

# June 2014

## Volume 2 Number 1

**Editors:** Kaye Wilson,  
Donna Jennings & Sarah Le Leu  
email: [schedule@pharmac.govt.nz](mailto:schedule@pharmac.govt.nz)  
Telephone +64 4 460 4990  
Facsimile +64 4 460 4995  
Level 9, 40 Mercer Street  
PO Box 10 254 Wellington 6143

**Freephone Information Line**  
**0800 66 00 50** (9am – 5pm weekdays)

### Circulation

Accessible in an electronic format at no cost from the Health Professionals section of the PHARMAC website [www.pharmac.govt.nz](http://www.pharmac.govt.nz)

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month. Alternatively there is a nominal charge for an annual subscription to the printed Schedule publications. To access either of these subscriptions visit our subscription website [www.schedule.co.nz](http://www.schedule.co.nz).

### Production

Typeset automatically from XML and T<sub>E</sub>X.  
XML version of the Schedule available from [www.pharmac.govt.nz/pub/schedule/archive/](http://www.pharmac.govt.nz/pub/schedule/archive/)

### Programmers

Anrik Drenth & John Geering  
email: [texschedule@pharmac.govt.nz](mailto:texschedule@pharmac.govt.nz)

©Pharmaceutical Management Agency



ISSN 1179-3708 pdf  
ISSN 1172-9694 print

This work is licensed under the Creative Commons Attribution 3.0 New Zealand licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to PHARMAC and abide by the other licence terms. To view a copy of this licence, visit:

[creativecommons.org/licenses/by/3.0/nz/](http://creativecommons.org/licenses/by/3.0/nz/).

Attribution to PHARMAC should be in written form and not by reproduction of the PHARMAC logo. While care has been taken in compiling this Schedule, PHARMAC takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

## Part I General Rules 4

<b>Part II</b>	Alimentary Tract and Metabolism	12
	Blood and Blood Forming Organs	25
	Cardiovascular System	35
	Dermatologicals	47
	Genito-Urinary System	53
	Hormone Preparations	57
	Infections	63
	Musculoskeletal System	86
	Nervous System	96
	Oncology Agents and Immunosuppressants	123
	Respiratory System and Allergies	161
	Sensory Organs	167
	Various	173
	Extemporaneous Compounds (ECPs)	181
	Special Foods	184
	Vaccines	198

## Part III Optional Pharmaceuticals 203

## Index 205

## Introducing PHARMAC

PHARMAC, the Pharmaceutical Management Agency, is a Crown entity established pursuant to the New Zealand Public Health and Disability Act 2000 (The Act). The primary objective of PHARMAC is to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided.

The PHARMAC Board consists of up to six members appointed by the Minister of Health. All decisions relating to PHARMAC's operation are made by or under the authority of the Board. More information on the Board can be found at [www.pharmac.govt.nz](http://www.pharmac.govt.nz). The functions of PHARMAC are set out in section 48 of the Act. PHARMAC is required to perform these functions within the amount of funding provided to it and in accordance with its statement of intent and any directions given by the Minister (Section 103 of the Crown Entities Act). The Government has agreed that PHARMAC will assume responsibility for the assessment, prioritisation and procurement of medical devices on behalf of DHBs. Medical devices come within the definition of Pharmaceuticals in the Act. PHARMAC is assuming responsibility for procurement of some medical devices categories immediately, as a first step to full PHARMAC management of these categories within the Pharmaceutical Schedule.

## Decision Criteria

PHARMAC takes into account the following criteria when considering amendments to the Schedule:

- a) the health needs of all eligible people within New Zealand;
- b) the particular health needs of Māori and Pacific peoples;
- c) the availability and suitability of existing medicines, therapeutic medical devices and related products and related things;
- d) the clinical benefits and risks of pharmaceuticals;
- e) the cost-effectiveness of meeting health needs by funding pharmaceuticals rather than using other publicly funded health and disability support services;
- f) the budgetary impact (in terms of the pharmaceutical budget and the Government's overall health budget) of any changes to the Schedule;
- g) the direct cost to health service users;
- h) the Government's priorities for health funding, as set out in any objectives notified by the Crown to PHARMAC, or in PHARMAC's Funding Agreement, or elsewhere; and
- i) such other criteria as PHARMAC thinks fit. PHARMAC will carry out appropriate consultation when it intends to take any such "other criteria" into account.

## PHARMAC's clinical advisors

### Pharmacology and Therapeutics Advisory Committee (PTAC)

PHARMAC works closely with the Pharmacology and Therapeutics Advisory Committee (PTAC), an expert medical committee which provides independent advice to PHARMAC on health needs and the clinical benefits of particular pharmaceuticals for use in the community and/or in DHB Hospitals. The chair of PTAC sits with the PHARMAC Board in an advisory capacity.

Contact PTAC C/-PTAC Secretary, Pharmaceutical Management Agency, PO Box 10 254, WELLINGTON 6143, Email: [PTAC@pharmac.govt.nz](mailto:PTAC@pharmac.govt.nz)

### PTAC Subcommittees

PTAC has subcommittees from which it can seek specialist advice in relation to funding applications. PTAC may seek advice from one or more subcommittees in relation to a funding application, or may make recommendations to PHARMAC without seeking the advice of a subcommittee:

Analgesic Subcommittee	Haematology Subcommittee	Reproductive and Sexual Health Subcommittee
Anti-Infective Subcommittee	Hospital Pharmaceuticals Subcommittee	Respiratory Subcommittee
Cancer Treatments Subcommittee	Immunisation Subcommittee	Rheumatology Subcommittee
Cardiovascular Subcommittee	Mental Health Subcommittee	Special Foods Subcommittee
Dermatology Subcommittee	Neurological Subcommittee	Transplant Immunosuppressants Subcommittee
Diabetes Subcommittee	Ophthalmology Subcommittee	
Endocrinology Subcommittee	Pulmonary Arterial Hypertension Subcommittee	
Gastrointestinal Subcommittee		

PTAC also has a Tender Medical Evaluation Subcommittee to provide advice on clinical matters relating to PHARMAC's annual multi-product tender and other purchasing strategies. Current membership of PTAC's subcommittees can be found on PHARMAC's website: <http://www.pharmac.health.nz/about/committees/ptac>

# Named Patient Pharmaceutical Assessment policy

Named Patient Pharmaceutical Assessment (NPPA) provides a mechanism for individual patients to receive funding for medicines not listed in the Pharmaceutical Schedule (either at all or for their clinical circumstances). PHARMAC will assess applications that meet the prerequisites according to its Decision Criteria before deciding whether to approve applications for funding. The Decision Criteria will be used to assess both the individual clinical circumstances of each NPPA applicant, and the implications of each NPPA funding decision on PHARMAC's ability to carry out its legislative functions.

For more information on NPPA, or to apply, visit the PHARMAC website at <http://www.pharmac.health.nz/tools- resources/forms/named-patient-pharmaceutical-assessment-nppa-forms>, or call the Panel Coordinators at (04) 9167553 or (04) 9167521.

## The Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by PHARMAC.

The purpose of the Schedule is not to show the final cost to Government of subsidising each Community Pharmaceutical or to DHBs in purchasing each Hospital Pharmaceutical or other Pharmaceuticals, including Medical Devices, used in DHB Hospitals, since that will depend on any rebate and other arrangements PHARMAC has with the supplier and, for some Hospital Pharmaceuticals, or other Pharmaceuticals, including Medical Devices, used in DHB Hospitals, on any logistics arrangements put in place by individual DHB Hospitals.

## Finding Information in Section H

Section H lists Pharmaceuticals that can be used in DHB Hospitals, and is split into the following parts:

- Part I lists the rules in relation to use of Pharmaceuticals by DHB Hospitals.
- Part II lists Hospital Pharmaceuticals that are funded for use in DHB Hospitals. These are classified based on the Anatomical Therapeutic Chemical (ATC) system used for Community Pharmaceuticals. It also provides information on any National Contracts that exist, and an indication of which products have Hospital Supply Status (HSS).
- Part III lists Optional Pharmaceuticals for which National Contracts exist, and DHB Hospitals may choose to fund. These are listed alphabetically by generic chemical entity name and line item, the relevant Price negotiated by PHARMAC and, if applicable, an indication of whether it has Hospital Supply Status (HSS) and any associated Discretionary Variance Limit (DV Limit).

The index located at the back of the Section H can be used to find page numbers for generic chemical entities and product brand names, for Hospital Pharmaceuticals. The listings are displayed alphabetically (where practical) within each level of the classification system. Each anatomical section contains a series of therapeutic headings, some of which may contain a further classification

## Glossary

### Units of Measure

gram .....	microgram.....	millimole.....
kilogram .....	milligram .....	unit.....
international unit .....	millilitre.....	

### Abbreviations

application .....	enteric coated .....	ointment.....
capsule .....	granules.....	solution.....
cream.....	injection .....	suppository.....
dispersible .....	linctus .....	tablet.....
effervescent .....	liquid.....	tincture.....
emulsion .....	lotion.....	

HSS    Hospital Supply Status (Refer to Rule 20)

### INTRODUCTION

Section H contains general rules that apply, and other information relating, to Hospital Pharmaceuticals and Optional Pharmaceuticals.

Where relevant, Section H shows the Price at which a Pharmaceutical can be purchased directly from the Pharmaceutical supplier by DHBs, providers of logistics services, wholesalers or other such distributors, or Contract Manufacturers.

The Price is determined via contractual arrangements between PHARMAC and the relevant Pharmaceutical supplier. Where a Pharmaceutical is listed in Part II of Section H, but no Price and/or brand of Pharmaceutical is indicated, each DHB may purchase any brand and/or pay the price that the DHB negotiates with the relevant Pharmaceutical supplier.

As required by section 23(7) of the Act, in performing any of its functions in relation to the supply of Pharmaceuticals, a DHB must not act inconsistently with the Pharmaceutical Schedule.

### INTERPRETATION AND DEFINITIONS

#### 1 Interpretation and Definitions

1.1 In this Schedule, unless the context otherwise requires:

**“Act”**, means the New Zealand Public Health and Disability Act 2000.

**“Combined Pharmaceutical Budget”**, means the pharmaceutical budget set for PHARMAC by the Crown for the subsidised supply of Community Pharmaceuticals and Pharmaceutical Cancer Treatments including for named patients in exceptional circumstances.

**“Community”**, means any setting outside of a DHB Hospital.

**“Community Pharmaceutical”**, means a Pharmaceutical listed in Sections A to G or I of the Pharmaceutical Schedule that is subsidised by the Funder from the Combined Pharmaceutical Budget and, for the purposes of this Section H, includes Pharmaceutical Cancer Treatments (PCTs).

**“Contract Manufacturer”**, means a manufacturer or a supplier that is a party to a contract with the relevant DHB Hospital to compound Pharmaceuticals, on request from that DHB Hospital.

**“Designated Delivery Point”**, means at a DHB Hospital's discretion:

- a) a delivery point agreed between a Pharmaceutical supplier and the relevant DHB Hospital, to which delivery point that Pharmaceutical supplier must supply a National Contract Pharmaceutical directly at the Price; and/or
- b) any delivery point designated by the relevant DHB Hospital or PHARMAC, such delivery point being within 30 km of the relevant Pharmaceutical supplier's national distribution centre.

**“DHB”**, means an organisation established as a District Health Board by or under Section 19 of the Act.

**“DHB Hospital”**, means a hospital (including community trust hospitals) and/or an associated health service that is funded by a DHB including (but not limited to) district nursing services and child dental services.

**“DV Limit”**, means, for a particular National Contract Pharmaceutical with HSS, the National DV Limit or the Individual DV Limit.

**“DV Pharmaceutical”**, means a discretionary variance Pharmaceutical that does not have HSS but is used in place of one that does. Usually this means it is the same chemical entity, at the same strength, and in the same or a similar presentation or form, as the relevant National Contract Pharmaceutical with HSS. Where this is not the case, a note will be included with the listing of the relevant Hospital Pharmaceutical.

**“Extemporaneously Compounded Product”**, means a Pharmaceutical that is compounded from two or more Pharmaceuticals, for the purposes of reconstitution, dilution or otherwise.

**“First Transition Period”**, means the period of time after notification that a Pharmaceutical has been awarded HSS and before HSS is implemented.

**“Funder”**, means the body or bodies responsible, pursuant to the Act, for the funding of Pharmaceuticals listed on the Schedule (which may be one or more DHBs and/or the Ministry of Health) and their successors.

**“Give”**, means to administer, provide or dispense (or, in the case of a Medical Device, use) a Pharmaceutical, or to arrange for the administration, provision or dispensing (or, in the case of a Medical Device, use) of a Pharmaceutical, and “Given” has a corresponding meaning.

**“Hospital Pharmaceuticals”**, means the list of Pharmaceuticals set out in Section H Part II of the Schedule which includes some National Contract Pharmaceuticals.

**“HSS”**, stands for hospital supply status, which means the status of being the brand of the relevant National Contract Pharmaceutical that DHBs are obliged to purchase, subject to any DV Limit, for the period of hospital supply,

as awarded under an agreement between PHARMAC and the relevant Pharmaceutical supplier. Pharmaceuticals with HSS are listed in Section H in bold text.

**“Indication Restriction”**, means a limitation placed by PHARMAC on the funding of a Hospital Pharmaceutical which restricts funding to treatment of particular clinical circumstances.

**“Individual DV Limit”**, means, for a particular National Contract Pharmaceutical with HSS and a particular DHB Hospital, the discretionary variance limit, being the specified percentage of that DHB Hospital's Total Market Volume up to which that DHB Hospital may purchase DV Pharmaceuticals of that National Contract Pharmaceutical.

**“Local Restriction”**, means a restriction on the use of a Pharmaceutical in specific DHB Hospitals on the basis of prescriber type that is implemented by the relevant DHB in accordance with rule 7.

**“Medical Device”**, has the meaning set out in the Medicines Act 1981.

**“Named Patient Pharmaceutical Assessment Advisory Panel”**, means the panel of clinicians, appointed by the PHARMAC Board, that is responsible for advising PHARMAC, in accordance with its Terms of Reference, on Named Patient Pharmaceutical Assessment applications and any Exceptional Circumstances renewal applications submitted after 1 March 2012.

**“National Contract”**, means a contractual arrangement between PHARMAC and a Pharmaceutical supplier which sets out the basis on which any Pharmaceutical may be purchased for use in a DHB Hospital, including an agreement as to a national price.

**“National Contract Pharmaceutical”**, means a brand of Pharmaceutical listed in Section H, where PHARMAC has entered into contractual arrangements with the relevant Pharmaceutical supplier that specify the terms and conditions of listing, including the Price. Such Pharmaceuticals are recognisable in Section H because the relevant listing identifies the brand and Price.

**“National DV Limit”**, means, for a particular National Contract Pharmaceutical with HSS, the discretionary variance limit, being the specified percentage of the Total Market Volume up to which all DHB Hospitals may collectively purchase DV Pharmaceuticals of that National Contract Pharmaceutical.

**“Optional Pharmaceuticals”**, means the list of National Contract Pharmaceuticals set out in Section H Part III of the Schedule.

**“PHARMAC”**, means the Pharmaceutical Management Agency established by Section 46 of the Act.

**“Pharmacode”**, means the six or seven digit identifier assigned to a Pharmaceutical by the Pharmacy Guild following application from a Pharmaceutical supplier.

**“Pharmaceutical”**, means a medicine, therapeutic medical device, or related product or related thing listed in Sections B to I of the Schedule.

**“Pharmaceutical Cancer Treatment”**, means Pharmaceuticals for the treatment of cancer, listed in Sections A to G of the Schedule and identified therein as a “PCT” or “PCT only” Pharmaceutical that DHBs must fund for use in their DHB hospitals, and/or in association with outpatient services provided by their DHB Hospitals, in relation to the treatment of cancers.

**“Prescriber Restriction”**, means a restriction placed by PHARMAC on the funding of a Pharmaceutical on the basis of prescriber type (and where relevant in these rules, includes a Local Restriction).

**“Price”**, means the standard national price for a National Contract Pharmaceutical, and, unless agreed otherwise between PHARMAC and the Pharmaceutical supplier, includes any costs associated with the supply of the National Contract Pharmaceutical to, at a DHB Hospital's discretion, any Designated Delivery Point, or to a Contract Manufacturer (expressly for the purpose of compounding), but does not include the effect of any rebates which may have been negotiated between PHARMAC and the Pharmaceutical supplier.

**“Restriction”**, means a limitation, put in place by PHARMAC or a DHB, restricting the funding of a Pharmaceutical and includes Indication Restrictions, Local Restrictions and Prescriber Restrictions (as defined in this Part I of Section H).

**“Schedule”**, means this Pharmaceutical Schedule and all its sections and appendices.

**“Special Authority Approval”**, means an approval for funding of a Community Pharmaceutical that is marked in Sections B-G of the Schedule as being subject to a Special Authority restriction.

**“Total Market Volume”**, means, for a particular Hospital Pharmaceutical with HSS in any given period, in accordance with the data available to PHARMAC, the sum of:

- the total number of Units of the relevant Hospital Pharmaceutical with HSS purchased by all DHB Hospitals, or by a particular DHB Hospital in the case of the Individual DV Limit; and
- the total number of Units of all the relevant DV Pharmaceuticals purchased by all DHB Hospitals, or by a particular DHB Hospital in the case of the Individual DV Limit.

**“Unapproved Indication”**, means, for a Pharmaceutical, an indication for which it is not approved under the Medicines Act 1981. Clinicians prescribing Pharmaceuticals for Unapproved Indications should be aware of, and comply with, their obligations under Section 25 and/or Section 29 of the Medicines Act 1981 and as set out in rule 23.

**“Unit”**, means an individual unit of a Pharmaceutical (e.g. a tablet, 1 ml of an oral liquid, an ampoule or a syringe).

**“Unlisted Pharmaceutical”**, means a Pharmaceutical that is within the scope of a Hospital Pharmaceutical, but is not listed in Section H Part II.

- 1.2 In addition to the above interpretations and definitions, unless the content requires otherwise, a reference in the Schedule to:
  - a) the singular includes the plural; and
  - b) any legislation includes a modification and re-enactment of, legislation enacted in substitution for, and a regulation, Order in Council, and other instrument from time to time issued or made under, that legislation.

## HOSPITAL SUPPLY OF PHARMACEUTICALS

### 2 Hospital Pharmaceuticals

- 2.1 Section H Part II contains the list of Hospital Pharmaceuticals that must be funded by DHB Hospitals. Section H Part II does not currently encompass the following categories of pharmaceuticals:

- a) Medical Devices;
- b) whole or fractionated blood products;
- c) diagnostic products which have an ex vivo use, such as pregnancy tests and reagents;
- d) disinfectants and sterilising products, except those that are to be used in or on a patient;
- e) foods and probiotics;
- f) radioactive materials;
- g) medical gases; and
- h) parenteral nutrition.

Subject to rule 2.2, the funding of pharmaceuticals identified in a)–h) above is a decision for individual DHB Hospitals.

- 2.2 Section H Part III lists Optional Pharmaceuticals that PHARMAC and the relevant Pharmaceutical supplier have entered into contractual arrangements for the purchase of, including an agreement on a national price and other obligations such as HSS. DHB Hospitals may choose whether or not to fund the Optional Pharmaceuticals listed in Part III of Section H, but if they do, they must comply with any National Contract requirements.

- 2.3 Section H Part II does not encompass the provision of pharmaceutical treatments for DHB Hospital staff as part of an occupational health and safety programme. DHB Hospitals may choose whether or not to fund pharmaceutical treatments for such use, but if they do, they must comply with any National Contract requirements.

### 3 DHB Supply Obligations

- 3.1 In accordance with section 23(7) of the Act, in performing any of its functions in relation to the supply of pharmaceuticals, a DHB must not act inconsistently with the Pharmaceutical Schedule, which includes these General Rules.
- 3.2 DHB Hospitals are not required to hold stock of every Hospital Pharmaceutical listed in Section H Part II, but they must Give it within a reasonable time if it is prescribed.
- 3.3 DHB Hospitals are able to hold stock of an Unlisted Pharmaceutical if doing so is considered necessary for the DHB Hospital to be able to Give the Unlisted Pharmaceutical in a timely manner under rules 11–17 inclusive.
- 3.4 Except where permitted in accordance with rule 11, DHBs must not Give:
  - a) an Unlisted Pharmaceutical; or
  - b) a Hospital Pharmaceutical outside of any relevant Restrictions.

### 4 Funding

- 4.1 The purchase costs of Hospital Pharmaceuticals or Optional Pharmaceuticals administered, provided or dispensed by DHB Hospitals must be funded by the relevant DHB Hospital from its own budget, with the exception of:
  - a) Pharmaceutical Cancer Treatments;
  - b) Community Pharmaceuticals that have been brought to the DHB hospital by the patient who is being treated by outpatient Services or who is admitted as an inpatient;
  - c) Community Pharmaceuticals that have been dispensed to a mental health day clinic under a Practitioner's Supply Order; and

- d) Unlisted Pharmaceutical that have been brought to the DHB Hospital by the patient who is admitted as an inpatient.
- 4.2 For the avoidance of doubt, Pharmaceutical Cancer Treatments and Community Pharmaceuticals are funded through the Combined Pharmaceutical Budget, and Unlisted Pharmaceuticals are funded by the patient.

## LIMITS ON SUPPLY

### 5 Prescriber Restrictions

- 5.1 A DHB Hospital may only Give a Hospital Pharmaceutical that has a Prescriber Restriction if it is prescribed:
  - a) by a clinician of the type specified in the restriction for that Pharmaceutical or, subject to rule 5.2, pursuant to a recommendation from such a clinician;
  - b) in accordance with a protocol or guideline that has been endorsed by the DHB Hospital; or
  - c) in an emergency situation, provided that the prescriber has made reasonable attempts to comply with rule 5.1(a) above. If on-going treatment is required (i.e. beyond 24 hours) subsequent prescribing must comply with rule 5.1(a).
- 5.2 Where a Hospital Pharmaceutical is prescribed pursuant to a recommendation from a clinician of the type specified in the restriction for that Pharmaceutical:
  - a) the prescriber must consult with a clinician of the type specified in the restriction for that Pharmaceutical; and
  - b) the consultation must relate to the patient for whom the prescription is written; and
  - c) the consultation may be in person, by telephone, letter, facsimile or email; and
  - d) appropriate records are kept of the consultation, including recording the name of the advising clinician on the prescription/chart.
- 5.3 Where a clinician is working under supervision of a consultant who is of the type specified in the restriction for that Pharmaceutical, the requirements of rule 5.2 can be deemed to have been met.

### 6 Indication Restrictions

- 6.1 A DHB Hospital may only Give a Hospital Pharmaceutical that has an Indication Restriction, if it is prescribed for treatment of a patient with the particular clinical circumstances set out in the Indication Restriction.
- 6.2 If a patient has a current Special Authority Approval for the Hospital Pharmaceutical that the DHB Hospital wishes to Give, then the Indication Restriction is deemed to have been met.
- 6.3 If a Hospital Pharmaceutical has an Indication Restriction that is "for continuation only" then the DHB Hospital should only Give the Hospital Pharmaceutical where:
  - a) the patient has been treated with the Pharmaceutical in the Community; or
  - b) the patient is unable to be treated with an alternative Hospital Pharmaceutical, and the prescriber has explained to the patient that the Pharmaceutical is not fully subsidised in the Community.

### 7 Local Restrictions

- 7.1 A DHB Hospital may implement a Local Restriction, provided that:
  - a) in doing so, it ensures that the Local Restriction does not unreasonably limit funded access to the Hospital Pharmaceutical or undermine PHARMAC's decision that the Hospital Pharmaceutical must be funded;
  - b) it provides PHARMAC with details of each Local Restriction that it implements; and
- 7.2 PHARMAC may, when it considers that a Local Restriction does not conform to rule 7.1 above, require a DHB to amend or remove that Local Restriction.

### 8 Community use of Hospital Pharmaceuticals

- 8.1 Except where otherwise specified in Section H, DHB Hospitals can Give any Hospital Pharmaceutical to a patient for use in the Community, provided that:
  - a) the quantity does not exceed that sufficient for up to 30 days' treatment, unless:
    - i) it would be inappropriate to provide less than the amount in an original pack; or
    - ii) the relevant DHB Hospital has a Dispensing for Discharge Policy and the quantity dispensed is in accordance with that policy; and
  - b) the Hospital Pharmaceutical is supplied consistent with any applicable Restrictions.

### 9 Community use of Medical Devices

- 9.1 Subject to rules 9.2 and 9.3, DHB Hospitals may Give a Medical Device for patients for use in the Community.
- 9.2 Where a Medical Device (or a similar Medical Device) is a Community Pharmaceutical, the DHB Hospital must supply:
  - a) the brand of Medical Device that is listed in Sections A-G of the Schedule; and

- b) only to patients who meet the funding eligibility criteria set out in Sections A-G of the Schedule.
- 9.3 Where a DHB Hospital has supplied a Medical Device to a patient; and
  - a) that Medical Device (or a similar Medical Device) is subsequently listed in Sections A-G of the Schedule; and
  - b) the patient would not meet any funding eligibility criteria for the Medical Device set out in Sections A-G of the Schedule; and
  - c) the Medical Device has consumable components that need to be replaced throughout its usable life; then DHB Hospitals may continue to fund consumable products for that patient until the end of the usable life of the Medical Device. At the end of the usable life of the device, funding for a replacement device must be consistent with the Pharmaceutical Schedule and/or in accordance with the Named Patient Pharmaceutical Assessment policy.
- 9.4 DHB Hospitals may also continue to fund consumable products, as in rule 9.3 above, in situations where the DHB has been funding consumable products but where the Medical Device was funded by the patient.
- 10 **Extemporaneous Compounding**
  - 10.1 A DHB Hospital may Give any Extemporaneously Compounded Product for a patient in its care, provided that:
    - a) all of the component Pharmaceuticals of the Extemporaneously Compounded Product are Hospital Pharmaceuticals; and
    - b) the Extemporaneously Compounded Product is supplied consistent with any applicable rules or Restrictions for its component Hospital Pharmaceuticals.
  - 10.2 For the avoidance of doubt, this rule 10.1 applies to any Extemporaneously Compounded Product, whether it is manufactured by the DHB Hospital or by a Contract Manufacturer.

## EXCEPTIONS

- 11 **Named Patient Pharmaceutical Assessment**
  - 11.1 A DHB Hospitals may only Give:
    - a) an Unlisted Pharmaceutical; or
    - b) a Hospital Pharmaceutical outside of any relevant Restrictions,in accordance with the Named Patient Pharmaceutical Assessment Policy or rules 12–17 inclusive.
- 12 **Continuation**
  - 12.1 Where a patient's clinical circumstances have been stabilised via treatment in the Community with a pharmaceutical that has not been funded by the Funder, and that patient is admitted to hospital as an inpatient, a DHB Hospital may fund that pharmaceutical for the duration of the patient's stay, where:
    - a) the patient has not brought (or cannot arrange to bring) the pharmaceuticals to the DHB Hospital, or pharmacy staff consider that the pharmaceuticals brought to the DHB Hospital by the patient cannot be used; and
    - b) interrupted or delayed treatment would have significant adverse clinical consequences; and
    - c) it is not considered appropriate to switch treatment to a Hospital Pharmaceutical.
- 13 **Pre-Existing Use**
  - 13.1 Subject to 13.2, where a DHB Hospital has Given a pharmaceutical for a patient prior to 1 July 2013, and the pharmaceutical:
    - a) is an Unlisted Pharmaceutical; or
    - b) treatment of the patient would not comply with any relevant Restrictions;the DHB Hospital may continue to Give that pharmaceutical if it is considered that there would be significant adverse clinical consequences from ceasing or switching treatment.
  - 13.2 Each DHB Hospital must, by no later than 1 October 2013, provide PHARMAC with a report on pharmaceuticals it has Given in accordance with this rule 13 where treatment has continued beyond 1 August 2013.
- 14 **Clinical Trials and Free Stock**
  - 14.1 DHB Hospitals may Give any pharmaceutical that is funded by a third party and is being used:
    - 14.1.1 as part of a clinical trial that has Ethics Committee approval; or
    - 14.1.2 for on-going treatment of patients following the end of such a clinical trial.
  - 14.2 DHB Hospitals may Give any pharmaceutical that is provided free of charge by a supplier, provided that the pharmaceutical is provided as part of a programme of which the DHB, or supplier, has notified PHARMAC.
- 15 **Pharmaceutical Cancer Treatments in Paediatrics**

DHB Hospitals may Give any pharmaceutical for use within a paediatric oncology/haematology service for the treatment of cancer.



**16 Other Government Funding**

DHB Hospitals may Give any pharmaceutical where funding for that pharmaceutical has been specifically provided by a Government entity other than PHARMAC or a DHB.

**17 Other Exceptions**

17.1 PHARMAC may also approve the funding of a pharmaceutical within a single DHB Hospital for information gathering purposes or otherwise related to PHARMAC's decision-making process for considering additions to or amendments to the Pharmaceutical Schedule.

17.2 Funding approvals granted under rule 17.1 will be subject to specific limitations on use as determined appropriate by PHARMAC in each circumstance, in consultation with the relevant DHB Hospital and/or DHB.

**NATIONAL CONTRACTING****18 Hospital Pharmaceutical Contracts**

18.1 A DHB Hospital may enter into a contract for the purchase of any Pharmaceutical, including any Medical Device, that it is entitled to fund in accordance with this Schedule H and that is not a National Contract Pharmaceutical, provided that such a contract:

- a) does not oblige the relevant DHB Hospital to purchase a volume of that Pharmaceutical, if that Pharmaceutical is a DV Pharmaceutical, that is greater than the relevant DV Limit;
- b) enables PHARMAC to access and use future price and volume data in respect of that Pharmaceutical; and
- c) enables the relevant DHB Hospital to terminate the contract or relevant parts of the contract in order to give full effect to the National Contract on no more than 3 months' written notice to the Pharmaceutical supplier.

18.2 From 1 July 2013, where a DHB Hospital has a pre-existing supply contract for a particular brand of chemical entity for which there is a National Contract Pharmaceutical, the DHB may continue purchasing the chemical entity in accordance with its pre-existing supply contract however:

- a) from the day its pre-existing supply contract expires, that DHB Hospital is to purchase the relevant National Contract Pharmaceutical listed in Section H at the Price, and is to comply with any DV Limits for the National Contract Pharmaceutical where it has HSS;
- b) if purchase of the relevant National Contract Pharmaceutical listed in Section H at the Price, where it has HSS, would not cause the relevant DHB Hospital to be in breach of its pre-existing supply contract for a particular brand of chemical entity; the DHB Hospital must purchase the National Contract Pharmaceutical.

18.3 Following written notification from PHARMAC that a Pharmaceutical is a National Contract Pharmaceutical, either through Section H updates or otherwise, DHB Hospitals must, unless PHARMAC expressly notifies otherwise:

- a) take any steps available to them to terminate pre-existing contracts or relevant parts of such a contract, and
- b) not enter any new contracts or extend the period of any current contracts, for the supply of that National Contract Pharmaceutical or the relevant chemical entity or Medical Device.

**19 National Contract Pharmaceuticals**

19.1 DHB Hospitals must take all necessary steps to enable any contracts between PHARMAC and a Pharmaceutical supplier in relation to National Contract Pharmaceuticals to be given full effect.

19.2 The contractual arrangement between PHARMAC and the relevant supplier of a National Contract Pharmaceutical requires it to be made available for purchase at the relevant Price by any or all of the following:

- a) DHB Hospitals at Designated Delivery Points; and/or
- b) Contract Manufacturers (expressly for the purpose of compounding).

In the case of Medical Devices, a National Contract may require the Medical Device to be purchased by, and/or supplied to, a third party logistics provider.

**20 Hospital Supply Status (HSS)**

20.1 The DV Limit for any National Contract Pharmaceutical which has HSS is set out in the listing of the relevant National Contract Pharmaceutical in Section H, and may be amended from time to time.

20.2 If a National Contract Pharmaceutical is listed in Section H as having HSS, DHB Hospitals:

- a) are expected to use up any existing stocks of DV Pharmaceuticals during the First Transition Period;
- b) must not purchase DV Pharmaceuticals in volumes exceeding their usual requirements, or in volumes exceeding those which they reasonably expect to use, within the First Transition Period;
- c) must ensure that Contract Manufacturers, when manufacturing an Extemporaneously Compounded Product on their behalf, use the National Contract Pharmaceutical with HSS; and
- d) must purchase the National Contract Pharmaceutical with HSS except:

- i) to the extent that the DHB Hospital may use its discretion to purchase a DV Pharmaceutical within the DV Limit, provided that (subject to rule 20.2(d)(iii) below) the DV Limit has not been exceeded nationally;
- ii) if the Pharmaceutical supplier fails to supply that National Contract Pharmaceutical, in which case the relevant DHB Hospital does not have to comply with the DV Limit for that National Contract Pharmaceutical during that period of non-supply (and any such month(s) included in a period of non-supply will be excluded in any review of the DV Limit in accordance with rule 20.3 below);
- iii) that where the DV Limit has been exceeded nationally, the DHB Hospital may negotiate with the Pharmaceutical supplier that supplies the National Contract Pharmaceutical with HSS for written permission to vary the application of that DHB Hospital's Individual DV Limit for any patient whose exceptional needs require a DV Pharmaceutical.

**20.3 PHARMAC may, in its discretion, for any period or part period:**

- a) review usage by DHB Hospitals of the National Contract Pharmaceutical and DV Pharmaceuticals to determine whether the DV Limit has been exceeded; and
- b) audit compliance by DHB Hospitals with the DV Limits and related requirements.

**20.4 PHARMAC will address any issues of non-compliance by any individual DHB or DHB Hospital with a DV Limit by:**

- a) obtaining the relevant DHB or DHB Hospital's assurance that it will comply with the DV Limit for that National Contract Pharmaceutical with HSS in the remainder of the applicable period and any subsequent periods; and
- b) informing the relevant supplier of the HSS Pharmaceutical of any individual DHB or DHB Hospital's non-compliance with the DV Limit for that HSS Pharmaceutical.

**20.5 In addition to the steps taken by PHARMAC under rule 20.4 above to address any issues of non-compliance by any individual DHB or DHB Hospital with a DV Limit, the relevant Pharmaceutical supplier may require, in its discretion, financial compensation from the relevant DHB or DHB Hospital:**

- a) an amount representing that DHB or DHB Hospital's contribution towards exceeding the DV Limit (where PHARMAC is able to quantify this based on the information available to it); or
- b) the sum of \$1,000 or \$5,000 (depending on the terms of the applicable national contract applying to the HSS Pharmaceutical),

whichever is the greater as between sub-paragraphs (a) and (b) within the number of business days specified in the notice from the Pharmaceutical supplier requiring such payment to be made.

**21 Collection of rebates and payment of financial compensation**

21.1 Following the receipt of any rebates from a Pharmaceutical supplier in respect of a particular National Contract Pharmaceutical, PHARMAC will notify each relevant DHB and DHB Hospital of the amount of the rebate owing to it, being a portion of the total rebate determined by PHARMAC on the basis of that DHB Hospital's usage of that National Contract Pharmaceutical, where this is able to be determined. Where data to determine individual DHB Hospitals' usage is not available, PHARMAC will apportion rebates on the basis of an alternative method agreed between the relevant DHBs and PHARMAC.

21.2 PHARMAC will pay each DHB Hospital the rebate amounts (if any) owing to it, no less frequently than once each calendar quarter in respect of rebates received quarterly (or more often).

**22 Price and Volume Data**

22.1 DHB Hospitals must provide to PHARMAC, on a monthly basis in accordance with PHARMAC's requirements, any volume data and, unless it would result in a breach of a pre-existing contract, price data held by those DHB Hospitals in respect of any Pharmaceutical (including any Medical Device) listed in Section H.

22.2 All price and volume data provided to PHARMAC under rule 22.1 above should identify the relevant Hospital Pharmaceutical by using a Pharmacode or some other unique numerical identifier, and the date (month and year) on which the DHB Hospital incurred a cost for the purchase of that Hospital Pharmaceutical. Volume is to be measured in units (that being the smallest possible whole Unit – e.g. a capsule, a vial, a millilitre etc).

## MISCELLANEOUS PROVISIONS

**23 Unapproved Pharmaceuticals**

Prescribers should, where possible, prescribe Hospital Pharmaceuticals that are approved under the Medicines Act 1981. However, the funding criteria (including Restrictions) under which a Hospital Pharmaceutical is listed in Section H of the Schedule may:

23.1 in some cases, explicitly permit a DHB to fund a Pharmaceutical that is not approved under the Medicines Act 1981 or for an Unapproved Indication; or

23.2 not explicitly prohibit a DHB from funding a Pharmaceutical for use for an Unapproved Indication;

Accordingly, if clinicians are planning on prescribing an unapproved Pharmaceutical or a Pharmaceutical for an Unapproved Indication, they should:

23.1 be aware of and comply with their obligations under sections 25 and/or 29 of the Medicines Act 1981, as applicable, and otherwise under that Act and the Medicines Regulations 1984;

23.2 be aware of and comply with their obligations under the Health and Disability Commissioner's Code of Consumer Rights, including the requirement to obtain informed consent from the patient (PHARMAC recommends that clinicians obtain written consent); and

23.3 exercise their own skill, judgment, expertise and discretion, and make their own prescribing decisions with respect to the use of an unapproved Pharmaceutical or a Pharmaceutical for an Unapproved Indication.

Clinicians should be aware that simply by listing a Pharmaceutical on the Pharmaceutical Schedule, PHARMAC makes no representations about whether that Pharmaceutical has any form of approval or consent under, or whether the supply or use of the Pharmaceutical otherwise complies with, the Medicines Act 1981. Further, the Pharmaceutical Schedule does not constitute an advertisement, advertising material or a medical advertisement as defined in the Medicines Act or otherwise.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Antacids and Antiflatulents</b>			
<b>Antacids and Reflux Barrier Agents</b>			
ALUMINIUM HYDROXIDE WITH MAGNESIUM HYDROXIDE AND SIMETHICONE			
Tab 200 mg with magnesium hydroxide 200 mg and simethicone 20 mg			<i>e.g. Mylanta</i>
Oral liq 200 mg with magnesium hydroxide 200 mg and simethicone 20 mg per 5 ml			<i>e.g. Mylanta</i>
Oral liq 400 mg with magnesium hydroxide 400 mg and simethicone 30 mg per 5 ml			<i>e.g. Mylanta Double Strength</i>
SIMETHICONE			
Oral drops 100 mg per ml			
SODIUM ALGINATE WITH MAGNESIUM ALGINATE			
Powder for oral soln 225 mg with magnesium alginate 87.5 mg, sachet			<i>e.g. Gaviscon Infant</i>
SODIUM ALGINATE WITH SODIUM BICARBONATE AND CALCIUM CARBONATE			
Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg			<i>e.g. Gaviscon Double Strength</i>
Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbon- ate 160 mg per 10 ml .....	4.95	500 ml	Acidex
SODIUM CITRATE			
Oral liq 8.8% (300 mmol/l)			
<b>Phosphate Binding Agents</b>			
ALUMINIUM HYDROXIDE			
Tab 600 mg			
CALCIUM CARBONATE – <b>Restricted</b> see terms below			
☞ Oral liq 250 mg per ml (100 mg elemental per ml) .....	39.00	500 ml	Roxane
☞ <b>Restricted</b>			
Only for use in children under 12 years of age for use as a phosphate binding agent			
<b>Antidiarrhoeals and Intestinal Anti-Inflammatory Agents</b>			
<b>Antipropulsives</b>			
DIPHENOXYLATE HYDROCHLORIDE WITH ATROPINE SULPHATE			
Tab 2.5 mg with atropine sulphate 25 mg			
LOPERAMIDE HYDROCHLORIDE			
Tab 2 mg			
Cap 2 mg – 1% DV Jul-14 to 2016 .....	7.84	400	Diamide Relief
<b>Rectal and Colonic Anti-Inflammatories</b>			
BUDESONIDE – <b>Restricted</b> see terms on the next page			
☞ Cap 3 mg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➔ Restricted</b>			
<b>Crohn's disease</b>			
Both:			
1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and			
2 Any of the following:			
2.1 Diabetes; or			
2.2 Cushingoid habitus; or			
2.3 Osteoporosis where there is significant risk of fracture; or			
2.4 Severe acne following treatment with conventional corticosteroid therapy; or			
2.5 History of severe psychiatric problems associated with corticosteroid treatment; or			
2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or			
2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).			
<b>Collagenous and lymphocytic colitis (microscopic colitis)</b>			
Patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies			
<b>Gut Graft versus Host disease</b>			
Patient has a gut Graft versus Host disease following allogenic bone marrow transplantation			
<b>HYDROCORTISONE ACETATE</b>			
Rectal foam 10% (14 applications) – <b>1% DV Jan-13 to 2015</b> .....	25.30	21.1 g	<b>Colifoam</b>
<b>MESALAZINE</b>			
Tab EC 400 mg .....	49.50	100	<b>Asacol</b>
Tab EC 500 mg .....	49.50	100	<b>Asamax</b>
Tab long-acting 500 mg .....	59.05	100	<b>Pentasa</b>
Modified release granules 1 g .....	141.72	120 g	<b>Pentasa</b>
Suppos 500 mg – <b>1% DV Sep-11 to 2014</b> .....	22.80	20	<b>Asacol</b>
Suppos 1 g .....	54.60	30	<b>Pentasa</b>
Enema 1 g per 100 ml – <b>1% DV Sep-12 to 2015</b> .....	44.12	7	<b>Pentasa</b>
<b>OLSALAZINE</b>			
Tab 500 mg			
Cap 250 mg			
<b>SODIUM CROMOGLYCATE</b>			
Cap 100 mg			
<b>SULPHASALAZINE</b>			
Tab 500 mg – <b>1% DV Oct-13 to 2016</b> .....	11.68	100	<b>Salazopyrin</b>
Tab EC 500 mg – <b>1% DV Oct-13 to 2016</b> .....	12.89	100	<b>Salazopyrin EN</b>
<b>Local Preparations for Anal and Rectal Disorders</b>			
<b>Antihæmorrhoidal Preparations</b>			
<b>CINCHOCAINE HYDROCHLORIDE WITH HYDROCORTISONE</b>			
Oint 5 mg with hydrocortisone 5 mg per g .....	15.00	30 g	<b>Proctosedyl</b>
Suppos 5 mg with hydrocortisone 5 mg per g .....	9.90	12	<b>Proctosedyl</b>
<b>FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE</b>			
Oint 950 mcg with fluocortolone pivalate 920 mcg and cinchocaine hydrochloride 5 mg per g .....	6.35	30 g	<b>Ultraproct</b>
Suppos 630 mcg with fluocortolone pivalate 610 mcg and cinchocaine hydrochloride 1 mg .....	2.66	12	<b>Ultraproct</b>
Products with Hospital Supply Status (HSS) are in <b>bold</b>			
Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Management of Anal Fissures</b>			
GLYCERYL TRINITRATE			
Oint 0.2% .....	22.00	30 g	Rectogesic
<b>Rectal Sclerosants</b>			
OILY PHENOL [PHENOL OILY]			
Inj 5%, 5 ml vial			
<b>Antispasmodics and Other Agents Altering Gut Motility</b>			
GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – 1% DV Oct-13 to 2016.....	28.56	10	Max Health
HYOSCINE BUTYLBROMIDE			
Tab 10 mg – 1% DV Sep-11 to 2014 .....	1.48	20	Gastrosoothe
Inj 20 mg, 1 ml ampoule – 1% DV Nov-11 to 2014.....	9.57	5	Buscopan
MEBEVERINE HYDROCHLORIDE			
Tab 135 mg – 1% DV Sep-11 to 2014 .....	18.00	90	Colofac
<b>Antiulcerants</b>			
<b>Antisecretory and Cytoprotective</b>			
MISOPROSTOL			
Tab 200 mcg			
<b>H2 Antagonists</b>			
CIMETIDINE			
Tab 200 mg			
Tab 400 mg			
RANITIDINE			
Tab 150 mg – 1% DV Sep-11 to 2014 .....	6.79	250	Arrow-Ranitidine
Tab 300 mg – 1% DV Sep-11 to 2014 .....	9.34	250	Arrow-Ranitidine
Oral liq 150 mg per 10 ml – 1% DV Sep-11 to 2014.....	5.92	300 ml	Peptisoothe
Inj 25 mg per ml, 2 ml ampoule .....	8.75	5	Zantac
<b>Proton Pump Inhibitors</b>			
LANSOPRAZOLE			
Cap 15 mg – 1% DV Jan-13 to 2015 .....	2.00	28	Solox
Cap 30 mg – 1% DV Jan-13 to 2015 .....	2.32	28	Solox
OMEPRAZOLE			
☞ Tab dispersible 20 mg			
☞Restricted			
Only for use in tube-fed patients			
Cap 10 mg – 1% DV Oct-11 to 2014 .....	2.91	90	Omezol Relief
Cap 20 mg – 1% DV Oct-11 to 2014 .....	3.78	90	Omezol Relief
Cap 40 mg – 1% DV Oct-11 to 2014 .....	5.57	90	Omezol Relief
Powder for oral liq – 1% DV Sep-11 to 2014.....	42.50	5 g	Midwest
Inj 40 mg ampoule – 1% DV Sep-11 to 2014.....	19.00	5	Dr Reddy's Omeprazole
Inj 40 mg ampoule with diluent – 1% DV Sep-11 to 2014.....	28.65	5	Dr Reddy's Omeprazole

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PANTOPRAZOLE</b>			
Tab EC 20 mg – 1% DV May-14 to 2016 .....	2.68	100	<b>Pantoprazole Actavis 20</b>
Tab EC 40 mg – 1% DV May-14 to 2016 .....	3.54	100	<b>Pantoprazole Actavis 40</b>
Inj 40 mg vial			

### Site Protective Agents

<b>BISMUTH TRIOXIDE</b>			
Tab 120 mg .....	32.50	112	De-Nol
<b>SUCRALFATE</b>			
Tab 1 g			

### Bile and Liver Therapy

L-ORNITHINE L-ASPARTATE – **Restricted** see terms below

⚡ Grans for oral liquid 3 g

➡ **Restricted**

For patients with chronic hepatic encephalopathy who have not responded to treatment with, or are intolerant to lactulose, or where lactulose is contraindicated.

### Diabetes

#### Alpha Glucosidase Inhibitors

<b>ACARBOSE</b>			
Tab 50 mg – 1% DV Dec-12 to 2015 .....	9.82	90	<b>Accarb</b>
Tab 100 mg – 1% DV Dec-12 to 2015 .....	15.83	90	<b>Accarb</b>

#### Hyperglycaemic Agents

**DIAZOXIDE** – **Restricted** see terms below

⚡ Cap 25 mg .....	110.00	100	Proglucem
⚡ Cap 100 mg .....	280.00	100	Proglucem

➡ **Restricted**

For patients with confirmed hypoglycaemia caused by hyperinsulinism.

<b>GLUCAGON HYDROCHLORIDE</b>			
Inj 1 mg syringe kit .....	32.00	1	Glucagen Hypokit

**GLUCOSE**

Tab 1.5 g  
Tab 3.1 g  
Gel 40%

**GLUCOSE WITH SUCROSE AND FRUCTOSE**

Gel 19.7% with sucrose 35% and fructose 19.7%, 18 g sachet

### Insulin - Intermediate-Acting Preparations

**INSULIN ASPART WITH INSULIN ASPART PROTAMINE**

Inj insulin aspart 30% with insulin aspart protamine 70%, 100 u per ml, 3 ml prefilled pen .....	52.15	5	NovoMix 30 FlexPen
---	-------	---	--------------------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>INSULIN ISOPHANE</b>			
Inj insulin human 100 u per ml, 10 ml vial			
Inj insulin human 100 u per ml, 3 ml cartridge			
<b>INSULIN LISPRO WITH INSULIN LISPRO PROTAMINE</b>			
Inj insulin lispro 25% with insulin lispro protamine 75%, 100 u per ml, 3 ml cartridge .....	42.66	5	Humalog Mix 25
Inj insulin lispro 50% with insulin lispro protamine 50%, 100 u per ml, 3 ml cartridge .....	42.66	5	Humalog Mix 50
<b>INSULIN NEUTRAL WITH INSULIN ISOPHANE</b>			
Inj insulin neutral 30% with insulin isophane 70%, 100 u per ml, 10 ml vial			
Inj insulin neutral 30% with insulin isophane 70%, 100 u per ml, 3 ml cartridge			
Inj insulin neutral 40% with insulin isophane 60%, 100 u per ml, 3 ml cartridge			
Inj insulin neutral 50% with insulin isophane 50%, 100 u per ml, 3 ml cartridge			
<b>Insulin - Long-Acting Preparations</b>			
<b>INSULIN GLARGINE</b>			
Inj 100 u per ml, 3 ml disposable pen .....	94.50	5	Lantus SoloStar
Inj 100 u per ml, 3 ml cartridge .....	94.50	5	Lantus
Inj 100 u per ml, 10 ml vial .....	63.00	1	Lantus
<b>Insulin - Rapid-Acting Preparations</b>			
<b>INSULIN ASPART</b>			
Inj 100 u per ml, 10 ml vial			
Inj 100 u per ml, 3 ml cartridge			
<b>INSULIN GLULISINE</b>			
Inj 100 u per ml, 10 ml vial .....	27.03	1	Apidra
Inj 100 u per ml, 3 ml cartridge .....	46.07	5	Apidra
Inj 100 u per ml, 3 ml disposable pen .....	46.07	5	Apidra Solostar
<b>INSULIN LISPRO</b>			
Inj 100 u per ml, 10 ml vial			
Inj 100 u per ml, 3 ml cartridge			
<b>Insulin - Short-Acting Preparations</b>			
<b>INSULIN NEUTRAL</b>			
Inj human 100 u per ml, 10 ml vial			
Inj human 100 u per ml, 3 ml cartridge			
<b>Oral Hypoglycaemic Agents</b>			
<b>GLIBENCLAMIDE</b>			
Tab 5 mg			
<b>GLICLAZIDE</b>			
Tab 80 mg – 1% DV Sep-11 to 2014 .....	17.60	500	Apo-Gliclazide



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>GLIPIZIDE</b>			
Tab 5 mg – 1% DV Dec-12 to 2015 .....	3.00	100	<b>Minidiab</b>
<b>METFORMIN</b>			
Tab immediate-release 500 mg – 1% DV Oct-12 to 2015.....	12.30	1,000	<b>Apotex</b>
Tab immediate-release 850 mg – 1% DV Oct-12 to 2015.....	10.10	500	<b>Apotex</b>
<b>PIOGLITAZONE</b>			
Tab 15 mg – 1% DV Sep-12 to 2015 .....	1.50	28	<b>Pizaccord</b>
Tab 30 mg – 1% DV Sep-12 to 2015 .....	2.50	28	<b>Pizaccord</b>
Tab 45 mg – 1% DV Sep-12 to 2015 .....	3.50	28	<b>Pizaccord</b>

## Digestives Including Enzymes

### PANCREATIC ENZYME

Cap EC 10,000 BP u lipase, 9,000 BP u amylase and 210 BP u protease

Cap EC 25,000 BP u lipase, 18,000 BP u amylase and 1,000 BP u protease

Cap EC 25,000 BP u lipase, 22,500 BP u amylase and 1,250 BP u protease

Powder 25,000 u lipase with 30,000 u amylase and 1,400 u protease per g

### URSODEOXYCHOLIC ACID – **Restricted** see terms below

☛ Cap 250 mg – 1% DV May-12 to 2014 ..... 71.50 100 **Ursosan**

#### ☛ **Restricted**

#### **Alagille syndrome or progressive familial intrahepatic cholestasis**

Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

#### **Chronic severe drug induced cholestatic liver injury**

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

#### **Cirrhosis**

Either:

- 1 Primary biliary cirrhosis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 µmol/l; decompensated cirrhosis).

#### **Pregnancy**

Patient diagnosed with cholestasis of pregnancy.

#### **Haematological transplant**

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

#### **Total parenteral nutrition induced cholestasis**

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by TPN; and
- 2 Liver function has not improved with modifying the TPN composition.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Laxatives</b>			
<b>Bowel-Cleansing Preparations</b>			
CITRIC ACID WITH MAGNESIUM OXIDE AND SODIUM PICOSULFATE			
Powder for oral soln 12 g with magnesium oxide 3.5 g and sodium picosulfate 10 mg per sachet			<i>e.g. PicoPrep</i>
MACROGOL 3350 WITH ASCORBIC ACID, POTASSIUM CHLORIDE AND SODIUM CHLORIDE			
Powder for oral soln 755.68 mg with ascorbic acid 85.16 mg, potassium chloride 10.55 mg, sodium chloride 37.33 mg and sodium sulphate 80.62 mg per g, 210 g sachet			<i>e.g. Glycoprep-C</i>
Powder for oral soln 755.68 mg with ascorbic acid 85.16 mg, potassium chloride 10.55 mg, sodium chloride 37.33 mg and sodium sulphate 80.62 mg per g, 70 g sachet			<i>e.g. Glycoprep-C</i>
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE, SODIUM CHLORIDE AND SODIUM SULPHATE			
Powder for oral soln 59 g with potassium chloride 0.7425 g, sodium bicarbonate 1.685 g, sodium chloride 1.465 g and sodium sulphate 5.685 g per sachet	14.31	4	Klean Prep
<b>Bulk-Forming Agents</b>			
ISPAGHULA (PSYLLIUM) HUSK			
Powder for oral soln – 1% DV Sep-13 to 2016	5.51	500 g	Konsyl-D
STERCULIA WITH FRANGULA – <b>Restricted:</b> For continuation only			
➔ Powder for oral soln			
<b>Faecal Softeners</b>			
DOCUSATE SODIUM			
Cap 50 mg – 1% DV Sep-11 to 2014	2.57	100	Laxofast 50
Cap 120 mg – 1% DV Sep-11 to 2014	3.48	100	Laxofast 120
DOCUSATE SODIUM WITH SENNOSIDES			
Tab 50 mg with sennosides 8 mg	6.38	200	Laxsol
PARAFFIN			
Oral liquid 1 mg per ml			
Enema 133 ml			
POLOXAMER			
Oral drops 10% – 1% DV Sep-11 to 2014	3.78	30 ml	Coloxyl
<b>Osmotic Laxatives</b>			
GLYCEROL			
Suppos 1.27 g			
Suppos 2.55 g			
Suppos 3.6 g – 1% DV Jan-13 to 2015	6.50	20	PSM
LACTULOSE			
Oral liq 10 g per 15 ml – 1% DV May-14 to 2014	3.84	500 ml	Laevolac

↑ Item restricted (see ➔ above); ↓ Item restricted (see ➔ below)

*e.g. Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE AND SODIUM CHLORIDE – <b>Restricted</b> see terms below			
⚡ Powder for oral soln 6.563 g with potassium chloride 23.3 mg, sodium bicarbonate 89.3 mg and sodium chloride 175.4 mg			
⚡ Powder for oral soln 13.125 g with potassium chloride 46.6 mg, sodium bicarbonate 178.5 mg and sodium chloride 350.7 mg – <b>1% DV</b>			
<b>Nov-13 to 2014</b> .....	10.00	30	<b>Lax-Sachets</b>

➡ **Restricted**

Either:

- 1 The patient has problematic constipation requiring intervention with a per rectal preparation despite an adequate trial of other oral pharmacotherapies including lactulose where lactulose is not contraindicated; or
- 2 For short-term use for faecal disimpaction.

## SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE

Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml – <b>1% DV Sep-13 to 2016</b> .....	19.95	50	<b>Micolette</b>
--	-------	----	------------------

## SODIUM PHOSPHATE WITH PHOSPHORIC ACID

Oral liq 16.4% with phosphoric acid 25.14% .....			
Enema 10% with phosphoric acid 6.58% .....	2.50	1	<b>Fleet Phosphate Enema</b>

**Stimulant Laxatives**

## BISACODYL

Tab 5 mg .....	4.99	200	<b>Lax-Tabs</b>
Suppos 5 mg .....	3.00	6	<b>Dulcolax</b>
Suppos 10 mg .....	3.00	6	<b>Dulcolax</b>

DANTHRON WITH POLOXAMER – **Restricted** see terms below

⚡ Oral liq 25 mg with poloxamer 200 mg per 5 ml .....	21.30	300 ml	<b>Pinorax</b>
⚡ Oral liq 75 mg with poloxamer 1 g per 5 ml .....	43.60	300 ml	<b>Pinorax Forte</b>

➡ **Restricted**

Only for the prevention or treatment of constipation in the terminally ill

## SENNOSIDES

Tab 7.5 mg

**Metabolic Disorder Agents**

## ARGININE

Powder  
Inj 600 mg per ml, 25 ml vial

BETAINE – **Restricted** see terms below

⚡ Powder

➡ **Restricted**

Metabolic disorders physician or metabolic disorders dietitian

BIOTIN – **Restricted** see terms below

⚡ Cap 50 mg  
⚡ Cap 100 mg  
⚡ Inj 10 mg per ml, 5 ml vial

➡ **Restricted**

Metabolic disorders physician or metabolic disorders dietitian.

## HAEM ARGINATE

Inj 25 mg per ml, 10 ml ampoule

# ALIMENTARY TRACT AND METABOLISM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>IMIGLUCERASE – Restricted</b> see terms below			
¶ Inj 40 iu per ml, 5 ml vial			
¶ Inj 40 iu per ml, 10 ml vial			
➡ <b>Restricted</b>			
Only for use in patients with approval by the Gaucher's Treatment Panel			
<b>LEVOCARNITINE – Restricted</b> see terms below			
¶ Cap 500 mg			
¶ Oral soln 500 mg per 15 ml			
¶ Inj 200 mg per ml, 5 ml vial			
➡ <b>Restricted</b>			
Metabolic disorders physician, metabolic disorders dietitian or neurologist			
<b>PYRIDOXAL-5-PHOSPHATE – Restricted</b> see terms below			
¶ Tab 50 mg			
➡ <b>Restricted</b>			
Metabolic disorders physician, metabolic disorders dietician or neurologist			
<b>SODIUM BENZOATE</b>			
Cap 500 mg			
Powder			
Soln 100 mg per ml			
Inj 20%, 10 ml ampoule			
<b>SODIUM PHENYLBUTYRATE</b>			
Tab 500 mg			
Oral liq 250 mg per ml			
Inj 200 mg per ml, 10 ml ampoule			
<b>TRIENTINE DIHYDROCHLORIDE</b>			
Cap 300 mg			

## Minerals

### Calcium

<b>CALCIUM CARBONATE</b>			
Tab 1.25 g (500 mg elemental) – 1% DV Feb-12 to 2014 .....	6.38	250	<b>Arrow-Calcium</b>
Tab 1.5 g (600 mg elemental)			
Tab eff 1.75 g (1 g elemental) – 1% DV Nov-11 to 2014 .....	6.21	30	<b>Calsource</b>

### Fluoride

<b>SODIUM FLUORIDE</b>			
Tab 1.1 mg (0.5 mg elemental)			

### Iodine

<b>POTASSIUM IODATE</b>			
Tab 256 mcg (150 mcg elemental iodine)			
<b>POTASSIUM IODATE WITH IODINE</b>			
Oral liq 10% with iodine 5%			

### Iron

<b>FERROUS FUMARATE</b>			
Tab 200 mg (65 mg elemental) .....	4.35	100	<b>Ferro-tab</b>

↑ Item restricted (see ➡ above); ¶ Item restricted (see ➡ below)  
e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>FERROUS FUMARATE WITH FOLIC ACID</b>			
Tab 310 mg (100 mg elemental) with folic acid 350 mcg .....	4.75	60	Ferro-F-Tabs
<b>FERROUS GLUCONATE WITH ASCORBIC ACID</b>			
Tab 170 mg (20 mg elemental) with ascorbic acid 40 mg			
<b>FERROUS SULPHATE</b>			
Tab long-acting 325 mg (105 mg elemental) .....	2.06	30	Ferrograd
Oral liq 30 mg (6 mg elemental) per ml – <b>1% DV Apr-14 to 2016</b> .....	10.28	500 ml	<b>Ferodan</b>
<b>FERROUS SULPHATE WITH ASCORBIC ACID</b>			
Tab long-acting 325 mg (105 mg elemental) with ascorbic acid 500 mg			
<b>FERROUS SULPHATE WITH FOLIC ACID</b>			
Tab long-acting 325 mg (105 mg elemental) with folic acid 350 mcg			
<b>IRON POLYMALTOSE</b>			
Inj 50 mg per ml, 2 ml ampoule – <b>1% DV Oct-11 to 2014</b> .....	19.90	5	<b>Ferrum H</b>
<b>IRON SUCROSE</b>			
Inj 20 mg per ml, 5 ml ampoule .....	100.00	5	Venofer
<b>Magnesium</b>			
<b>MAGNESIUM HYDROXIDE</b>			
Tab 311 mg (130 mg elemental)			
<b>MAGNESIUM OXIDE</b>			
Cap 663 mg (400 mg elemental)			
<b>MAGNESIUM SULPHATE</b>			
Inj 0.4 mmol per ml, 250 ml bag			
Inj 2 mmol per ml, 5 ml ampoule – <b>1% DV Feb-13 to 2014</b> .....	18.35	10	<b>Martindale</b>
<b>Zinc</b>			
<b>ZINC</b>			
Oral liq 5 mg per 5 drops			
<b>ZINC CHLORIDE</b>			
Inj 5.3 mg per ml (5.1 mg per ml elemental), 2 ml ampoule			
<b>ZINC SULPHATE</b>			
Cap 137.4 mg (50 mg elemental) – <b>1% DV Nov-11 to 2014</b> .....	11.00	100	<b>Zincaps</b>
<b>Mouth and Throat</b>			
<b>Agents Used in Mouth Ulceration</b>			
<b>BENZYDAMINE HYDROCHLORIDE</b>			
Soln 0.15%			
Spray 0.15%			
<b>BENZYDAMINE HYDROCHLORIDE WITH CETYLPYRIDINIUM CHLORIDE</b>			
Lozenge 3 mg with cetylpyridinium chloride			
<b>CARBOXYMETHYLCELLULOSE</b>			
Oral spray			
<b>CHLORHEXIDINE GLUCONATE</b>			
Mouthwash 0.2% – <b>1% DV Dec-12 to 2015</b> .....	2.68	200 ml	<b>healthE</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE Adhesive gel 8.7% with cetalkonium chloride 0.01%			
DICHLOROBENZYL ALCOHOL WITH AMYLMETACRESOL Lozenge 1.2 mg with amylmetacresol 0.6 mg			
SODIUM CARBOXYMETHYLCELLULOSE WITH PECTIN AND GELATINE Paste Powder			
TRIAMCINOLONE ACETONIDE Paste 0.1% – 1% DV Sep-11 to 2014 .....	4.34	5 g	Oracort
<b>Oropharyngeal Anti-Infectives</b>			
AMPHOTERICIN B Lozenge 10 mg .....	5.86	20	Fungilin
MICONAZOLE Oral gel 20 mg per g – 1% DV Feb-13 to 2015 .....	4.95	40 g	Decozol
NYSTATIN Oral liquid 100,000 u per ml – 1% DV Sep-11 to 2014 .....	3.19	24 ml	Nilstat
<b>Other Oral Agents</b>			
SODIUM HYALURONATE – <b>Restricted</b> see terms below ⚡ Inj 20 mg per ml, 1 ml syringe ➡ <b>Restricted</b> Otolaryngologist			
THYMOL GLYCERIN Compound, BPC			
<b>Vitamins</b>			
<b>Multivitamin Preparations</b>			
MULTIVITAMINS Tab (BPC cap strength)			e.g. <i>Mvite</i>
⚡ Cap vitamin A 2500 u, betacarotene 3 mg, cholecalciferol 11 mcg, al- pha tocopherol 150 u, phytomenadione 150 mcg, folic acid 0.2 mg, ascorbic acid 100 mg, thiamine 1.5 mg, pantothenic acid 12 mg, riboflavin 1.7 mg, niacin 20 mg, pyridoxine hydrochloride 1.9 mg, cyanocobalamin 3 mcg, zinc 7.5 mg and biotin 100 mcg			e.g. <i>Vitabdeck</i>
➡ <b>Restricted</b> Either:			
1 Patient has cystic fibrosis with pancreatic insufficiency; or 2 Patient is an infant or child with liver disease or short gut syndrome.			
⚡ Powder vitamin A 4200 mcg with vitamin D 155.5 mcg, vitamin E 21.4 mg, vitamin C 400 mg, vitamin K1 166 mcg thiamine 3.2 mg, riboflavin 4.4 mg, niacin 35 mg, vitamin B6 3.4 mg, folic acid 303 mcg, vitamin B12 8.6 mcg, biotin 214 mcg, pantothenic acid 17 mg, choline 350 mg and inositol 700 mg			e.g. <i>Paediatric Seravit</i>
➡ <b>Restricted</b> Patient has inborn errors of metabolism.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Inj thiamine hydrochloride 250 mg with riboflavin 4 mg and pyridoxine hydrochloride 50 mg, 5 ml ampoule (1) and inj ascorbic acid 500 mg with nicotinamide 160 mg and glucose 1000 mg, 5 ml ampoule (1)			<i>e.g. Pabrinex IV</i>
Inj thiamine hydrochloride 250 mg with riboflavin 4 mg and pyridoxine hydrochloride 50 mg, 5 ml ampoule (1) and inj ascorbic acid 500 mg with nicotinamide 160 mg, 2 ml ampoule (1)			<i>e.g. Pabrinex IM</i>
Inj thiamine hydrochloride 500 mg with riboflavin 8 mg and pyridoxine hydrochloride 100 mg, 10 ml ampoule (1) and inj ascorbic acid 1000 mg with nicotinamide 320 mg and glucose 2000 mg, 10 ml ampoule (1)			<i>e.g. Pabrinex IV</i>
<b>VITAMIN A WITH VITAMINS D AND C</b>			
Soln 1,000 u with vitamin D 400 u and ascorbic acid 30 mg per 10 drops			<i>e.g. Vitadol C</i>

### Vitamin A

#### RETINOL

Tab 10,000 iu  
Cap 25,000 iu  
Oral liq 150,000 iu per ml

### Vitamin B

#### HYDROXOCOBALAMIN ACETATE

Inj 1 mg per ml, 1 ml ampoule – 1% DV Sep-12 to 2015 .....	5.10	3	<b>ABM</b> <b>Hydroxocobalamin</b>
--	------	---	---------------------------------------

#### PYRIDOXINE HYDROCHLORIDE

Tab 25 mg – 1% DV Sep-11 to 2014 .....	2.20	90	<b>PyridoxADE</b>
Tab 50 mg – 1% DV Sep-11 to 2014 .....	12.16	500	<b>Apo-Pyridoxine</b>
Inj 100 mg per ml, 1 ml ampoule			

#### THIAMINE HYDROCHLORIDE

Tab 50 mg  
Tab 100 mg  
Inj 100 mg per ml, 2 ml vial

#### VITAMIN B COMPLEX

Tab strong, BPC

### Vitamin C

#### ASCORBIC ACID

Tab 100 mg – 1% DV Nov-13 to 2016 .....	7.00	500	<b>Cvite</b>
Tab chewable 250 mg			

### Vitamin D

#### ALFACALCIDOL

Cap 0.25 mcg .....	26.32	100	One-Alpha
Cap 1 mcg .....	87.98	100	One-Alpha
Oral drops 2 mcg per ml			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>CALCITRIOL</b>			
Cap 0.25 mcg .....	3.03	30	Airflow
	10.10	100	Calcitriol-AFT
Cap 0.5 mcg .....	5.62	30	Airflow
	18.73	100	Calcitriol-AFT
Oral liq 1 mcg per ml			
Inj 1 mcg per ml, 1 ml ampoule			
<b>CHOLECALCIFEROL</b>			
Tab 1.25 mg (50,000 iu) .....	7.76	12	Cal-d-Forte

## Vitamin E

ALPHA TOCOPHERYL ACETATE – **Restricted** see terms below

- ⚡ Cap 100 u
- ⚡ Cap 500 u
- ⚡ Oral liq 156 u per ml

➡ **Restricted**

### Cystic fibrosis

Both:

- 1 Cystic fibrosis patient; and
- 2 Either:
  - 2.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck); or
  - 2.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for the patient.

### Osteoradionecrosis

For the treatment of osteoradionecrosis

### Other indications

All of the following:

- 1 Infant or child with liver disease or short gut syndrome; and
- 2 Requires vitamin supplementation; and
- 3 Either:
  - 3.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplements (Vitabdeck); or
  - 3.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for patient.



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Antianaemics</b>			
<b>Hypoplastic and Haemolytic</b>			
ERYTHROPOIETIN ALPHA – <b>Restricted</b> see terms below			
⚡ Inj 1,000 iu in 0.5 ml syringe .....	48.68	6	Eprex
⚡ Inj 2,000 iu in 0.5 ml syringe .....	120.18	6	Eprex
⚡ Inj 3,000 iu in 0.3 ml syringe .....	166.87	6	Eprex
⚡ Inj 4,000 iu in 0.4 ml syringe .....	193.13	6	Eprex
⚡ Inj 5,000 iu in 0.5 ml syringe .....	243.26	6	Eprex
⚡ Inj 6,000 iu in 0.6 ml syringe .....	291.92	6	Eprex
⚡ Inj 10,000 iu in 1 ml syringe .....	395.18	6	Eprex
➡ <b>Restricted</b>			
Both:			
1 Both:			
1.1 Patient in chronic renal failure; and			
1.2 Haemoglobin $\leq$ 100g/L; and			
2 Any of the following:			
2.1 Both:			
2.1.1 Patient is not diabetic; and			
2.1.2 Glomerular filtration rate $\leq$ 30ml/min; or			
2.2 Both:			
2.2.1 Patient is diabetic; and			
2.2.2 Glomerular filtration rate $\leq$ 45ml/min; or			
2.3 Patient is on haemodialysis or peritoneal dialysis.			
ERYTHROPOIETIN BETA – <b>Restricted</b> see terms below			
⚡ Inj 2,000 iu in 0.3 ml syringe .....	120.18	6	NeoRecormon
⚡ Inj 3,000 iu in 0.3 ml syringe .....	166.87	6	NeoRecormon
⚡ Inj 4,000 iu in 0.3 ml syringe .....	193.13	6	NeoRecormon
⚡ Inj 5,000 iu in 0.3 ml syringe .....	243.26	6	NeoRecormon
⚡ Inj 6,000 iu in 0.3 ml syringe .....	291.92	6	NeoRecormon
⚡ Inj 10,000 iu in 0.6 ml syringe .....	395.18	6	NeoRecormon
➡ <b>Restricted</b>			
Both:			
1 Both:			
1.1 Patient in chronic renal failure; and			
1.2 Haemoglobin $\leq$ 100g/L; and			
2 Any of the following:			
2.1 Both:			
2.1.1 Patient is not diabetic; and			
2.1.2 Glomerular filtration rate $\leq$ 30ml/min; or			
2.2 Both:			
2.2.1 Patient is diabetic; and			
2.2.2 Glomerular filtration rate $\leq$ 45ml/min; or			
2.3 Patient is on haemodialysis or peritoneal dialysis.			
Products with Hospital Supply Status (HSS) are in <b>bold</b>			
Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Megaloblastic</b>			
<b>FOLIC ACID</b>			
Tab 0.8 mg			
Tab 5 mg			
Oral liq 50 mcg per ml .....	24.00	25 ml	Biomed
Inj 5 mg per ml, 10 ml vial			
<b>Antifibrinolytics, Haemostatics and Local Sclerosants</b>			
<b>APROTININ – Restricted</b> see terms below			
☞ Inj 10,000 kIU per ml (equivalent to 200 mg per ml), 50 ml vial			
<b>☞ Restricted</b>			
Cardiac anaesthetist			
Either:			
1 Paediatric patient undergoing cardiopulmonary bypass procedure; or			
2 Adult patient undergoing cardiac surgical procedure where the significant risk of massive bleeding outweighs the potential adverse effects of the drug.			
<b>ELTROMBOPAG – Restricted</b> see terms below			
☞ Tab 25 mg .....	1,771.00	28	Revolade
☞ Tab 50 mg .....	3,542.00	28	Revolade
<b>☞ Restricted</b>			
Haematologist			
<b>Initiation (idiopathic thrombocytopenic purpura - post-splenectomy)</b>			
<i>Re-assessment required after 6 weeks</i>			
All of the following:			
1 Patient has had a splenectomy; and			
2 Two immunosuppressive therapies have been trialed and failed after therapy of 3 months each (or 1 month for rituximab); and			
3 Any of the following:			
3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or			
3.2 Patient has a platelet count of ≤ 20,000 platelets per microlitre and has evidence of active bleeding; or			
3.3 Patient has a platelet count of ≤ 10,000 platelets per microlitre.			
<b>Initiation - (idiopathic thrombocytopenic purpura - preparation for splenectomy)</b>			
<i>Re-assessment required after 6 weeks</i>			
The patient requires eltrombopag treatment as preparation for splenectomy.			
<b>Continuation - (idiopathic thrombocytopenic purpura - post-splenectomy)</b>			
<i>Re-assessment required after 12 months</i>			
The patient has obtained a response (see note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.			
Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.			
<b>FERRIC SUBSULFATE</b>			
Gel 25.9%			
Soln 500 ml			
<b>POLIDOCANOL</b>			
Inj 0.5%, 30 ml vial			
<b>SODIUM TETRADECYL SULPHATE</b>			
Inj 3%, 2 ml ampoule			
<b>THROMBIN</b>			
Powder			

☞ Item restricted (see ☞ above); ☞ Item restricted (see ☞ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>TRANEXAMIC ACID</b>			
Tab 500 mg .....	32.92	100	Cyklokapron
Inj 100 mg per ml, 5 ml ampoule .....	124.73	10	Cyklokapron

## Blood Factors

### EPTACOG ALFA [RECOMBINANT FACTOR VIIA] – **Restricted** see terms below

⚡ Inj 1 mg syringe .....	1,163.75	1	NovoSeven RT
⚡ Inj 2 mg syringe .....	2,327.50	1	NovoSeven RT
⚡ Inj 5 mg syringe .....	5,818.75	1	NovoSeven RT
⚡ Inj 8 mg syringe .....	9,310.00	1	NovoSeven RT

#### →Restricted

When used in the treatment of haemophilia, treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

### FACTOR EIGHT INHIBITORS BYPASSING AGENT – **Restricted** see terms below

⚡ Inj 500 U .....	1,640.00	1	FEIBA
⚡ Inj 1,000 U .....	3,280.00	1	FEIBA

#### →Restricted

When used in the treatment of haemophilia, treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

### MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] – **Restricted** see terms below

⚡ Inj 250 iu vial .....	225.00	1	Xyntha
⚡ Inj 500 iu vial .....	450.00	1	Xyntha
⚡ Inj 1,000 iu vial .....	900.00	1	Xyntha
⚡ Inj 2,000 iu vial .....	1,800.00	1	Xyntha
⚡ Inj 3,000 iu vial .....	2,700.00	1	Xyntha

#### →Restricted

When used in the treatment of haemophilia, treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

### NONACOG ALFA [RECOMBINANT FACTOR IX] – **Restricted** see terms below

⚡ Inj 250 iu vial .....	310.00	1	BeneFIX
⚡ Inj 500 iu vial .....	620.00	1	BeneFIX
⚡ Inj 1,000 iu vial .....	1,240.00	1	BeneFIX
⚡ Inj 2,000 iu vial .....	2,480.00	1	BeneFIX

#### →Restricted

When used in the treatment of haemophilia, treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

### OCTOCOG ALFA [RECOMBINANT FACTOR VIII] – **Restricted** see terms on the next page

⚡ Inj 250 iu vial .....	237.50	1	Advate
	250.00		Kogenate FS
⚡ Inj 500 iu vial .....	475.00	1	Advate
	500.00		Kogenate FS
⚡ Inj 1,000 iu vial .....	950.00	1	Advate
	1,000.00		Kogenate FS
⚡ Inj 1,500 iu vial .....	1,425.00	1	Advate
⚡ Inj 2,000 iu vial .....	1,900.00	1	Advate
	2,000.00		Kogenate FS
⚡ Inj 3,000 iu vial .....	2,850.00	1	Advate
	3,000.00		Kogenate FS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## ➡Restricted

When used in the treatment of haemophilia, treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

## Vitamin K

### PHYTOMENADIONE

Inj 2 mg in 0.2 ml ampoule .....	8.00	5	Konaktion MM
Inj 10 mg per ml, 1 ml ampoule .....	9.21	5	Konaktion MM

## Anti-thrombotics

## Anticoagulants

BIVALIRUDIN – **Restricted** see terms below

⚡ Inj 250 mg vial

## ➡Restricted

Either:

- 1 For use in heparin-induced thrombocytopenia, heparin resistance or heparin intolerance; or
- 2 For use in patients undergoing endovascular procedures.

### DABIGATRAN

Cap 75 mg .....	148.00	60	Pradaxa
Cap 110 mg .....	148.00	60	Pradaxa
Cap 150 mg .....	148.00	60	Pradaxa

### DALTEPARIN

Inj 2,500 iu in 0.2 ml syringe .....	19.97	10	Fragmin
Inj 5,000 iu in 0.2 ml syringe .....	39.94	10	Fragmin
Inj 7,500 iu in 0.75 ml syringe .....	60.03	10	Fragmin
Inj 10,000 iu in 1 ml syringe .....	77.55	10	Fragmin
Inj 12,500 iu in 0.5 ml syringe .....	99.96	10	Fragmin
Inj 15,000 iu in 0.6 ml syringe .....	120.05	10	Fragmin
Inj 18,000 iu in 0.72 ml syringe .....	158.47	10	Fragmin

DANAPAROID – **Restricted** see terms below

⚡ Inj 750 u in 0.6 ml ampoule

## ➡Restricted

For use in heparin-induced thrombocytopenia, heparin resistance or heparin intolerance

DEFIBROTIDE – **Restricted** see terms below

⚡ Inj 80 mg per ml, 2.5 ml ampoule

## ➡Restricted

Haematologist

Patient has moderate or severe sinusoidal obstruction syndrome as a result of chemotherapy or regimen-related toxicities

DEXTROSE WITH SODIUM CITRATE AND CITRIC ACID [ACID CITRATE DEXTROSE A]

Inj 24.5 mg with sodium citrate 22 mg and citric acid 7.3 mg per ml,  
100 ml bag

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>ENOXAPARIN</b>			
Inj 20 mg in 0.2 ml syringe – <b>1% DV Sep-12 to 2015</b> .....	37.24	10	<b>Clexane</b>
Inj 40 mg in 0.4 ml ampoule			
Inj 40 mg in 0.4 ml syringe – <b>1% DV Sep-12 to 2015</b> .....	49.69	10	<b>Clexane</b>
Inj 60 mg in 0.6 ml syringe – <b>1% DV Sep-12 to 2015</b> .....	74.91	10	<b>Clexane</b>
Inj 80 mg in 0.8 ml syringe – <b>1% DV Sep-12 to 2015</b> .....	99.86	10	<b>Clexane</b>
Inj 100 mg in 1 ml syringe – <b>1% DV Sep-12 to 2015</b> .....	125.06	10	<b>Clexane</b>
Inj 120 mg in 0.8 ml syringe – <b>1% DV Sep-12 to 2015</b> .....	155.40	10	<b>Clexane</b>
Inj 150 mg in 1 ml syringe – <b>1% DV Sep-12 to 2015</b> .....	177.60	10	<b>Clexane</b>
<b>FONDAPARINUX SODIUM – Restricted</b> see terms below			
☞ Inj 2.5 mg in 0.5 ml syringe			
☞ Inj 7.5 mg in 0.6 ml syringe			
☞ <b>Restricted</b>			
For use in heparin-induced thrombocytopenia, heparin resistance or heparin intolerance			
<b>HEPARIN SODIUM</b>			
Inj 100 iu per ml, 250 ml bag			
Inj 1,000 iu per ml, 1 ml ampoule .....	66.80	50	Hospira
Inj 1,000 iu per ml, 35 ml ampoule			
Inj 1,000 iu per ml, 5 ml ampoule .....	11.44	10	Pfizer
	46.30	50	Pfizer
Inj 5,000 iu in 0.2 ml ampoule			
Inj 5,000 iu per ml, 1 ml ampoule .....	14.20	5	Hospira
Inj 5,000 iu per ml, 5 ml ampoule .....	182.00	50	Pfizer
<b>HEPARINISED SALINE</b>			
Inj 10 iu per ml, 5 ml ampoule .....	32.50	50	Pfizer
Inj 100 iu per ml, 2 ml ampoule			
Inj 100 iu per ml, 5 ml ampoule			
<b>PHENINDIONE</b>			
Tab 10 mg			
Tab 25 mg			
Tab 50 mg			
<b>PROTAMINE SULPHATE</b>			
Inj 10 mg per ml, 5 ml ampoule			
<b>RIVAROXABAN – Restricted</b> see terms below			
☞ Tab 10 mg .....	153.00	15	Xarelto
☞ <b>Restricted</b>			
Either:			
1 Limited to five weeks' treatment for the prophylaxis of venous thromboembolism following a total hip replacement; or			
2 Limited to two weeks' treatment for the prophylaxis of venous thromboembolism following a total knee replacement.			
<b>SODIUM CITRATE WITH SODIUM CHLORIDE AND POTASSIUM CHLORIDE</b>			
Inj 4.2 mg with sodium chloride 5.7 mg and potassium chloride			
74.6 mcg per ml, 5,000 ml bag			
<b>TRISODIUM CITRATE</b>			
Inj 4%, 5 ml ampoule			
Inj 46.7%, 3 ml syringe			
Inj 46.7%, 5 ml ampoule			

## BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>WARFARIN SODIUM</b>			
Tab 1 mg .....	6.86	100	Marevan
Tab 2 mg .....			
Tab 3 mg .....	9.70	100	Marevan
Tab 5 mg .....	11.75	100	Marevan
<b>Antiplatelets</b>			
<b>ASPIRIN</b>			
Tab 100 mg – 1% DV Mar-14 to 2016 .....	1.60	90	<b>Ethics Aspirin EC</b>
	10.50	990	<b>Ethics Aspirin EC</b>
Suppos 300 mg			
<b>CLOPIDOGREL</b>			
Tab 75 mg – 1% DV Dec-13 to 2016 .....	5.48	84	<b>Arrow - Clopid</b>
<b>DIPYRIDAMOLE</b>			
Tab 25 mg			
Tab long-acting 150 mg – 1% DV Oct-11 to 2014 .....	11.52	60	<b>Pytazen SR</b>
Inj 5 mg per ml, 2 ml ampoule			
<b>EPTIFIBATIDE – Restricted</b> see terms below			
⚡ Inj 2 mg per ml, 10 ml vial .....	111.00	1	Integrilin
⚡ Inj 750 mcg per ml, 100 ml vial .....	324.00	1	Integrilin
<b>➡Restricted</b>			
Either:			
1 For use in patients with acute coronary syndromes undergoing percutaneous coronary intervention; or			
2 For use in patients with definite or strongly suspected intra-coronary thrombus on coronary angiography.			
<b>PRASUGREL – Restricted</b> see terms below			
⚡ Tab 5 mg .....	108.00	28	Effient
⚡ Tab 10 mg .....	120.00	28	Effient
<b>➡Restricted</b>			
<b>Bare metal stents</b>			
Limited to 6 months' treatment			
Patient has undergone coronary angioplasty in the previous 4 weeks and is clopidogrel-allergic.			
<b>Drug-eluting stents</b>			
Limited to 12 months' treatment			
Patient has had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic.			
<b>Stent thrombosis</b>			
Patient has experienced cardiac stent thrombosis whilst on clopidogrel.			
<b>Myocardial infarction</b>			
Limited to 7 days' treatment			
For short term use while in hospital following ST-elevated myocardial infarction.			
Note: Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.			
<b>TICAGRELOR – Restricted</b> see terms below			
⚡ Tab 90 mg .....	90.00	56	Brilinta
<b>➡Restricted</b>			
Restricted to treatment of acute coronary syndromes specifically for patients who have recently been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome, and in whom fibrinolytic therapy has not been given in the last 24 hours and is not planned.			
<b>TICLOPIDINE</b>			
Tab 250 mg			

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

## Fibrinolytic Agents

### ALTEPLASE

Inj 10 mg vial  
Inj 50 mg vial

### TENECTEPLASE

Inj 50 mg vial

### UROKINASE

Inj 10,000 iu vial  
Inj 50,000 iu vial  
Inj 100,000 iu vial  
Inj 500,000 iu vial

## Colony-Stimulating Factors

### Granulocyte Colony-Stimulating Factors

FILGRASTIM – **Restricted** see terms below

⚡ Inj 300 mcg in 0.5 ml syringe – 1% DV Jan-13 to 31 Dec 2015 .....	540.00	5	<b>Zarzio</b>
⚡ Inj 300 mcg in 1 ml vial .....	650.00	5	Neupogen
⚡ Inj 480 mcg in 0.5 ml syringe – 1% DV Jan-13 to 31 Dec 2015 .....	864.00	5	<b>Zarzio</b>

#### →Restricted

Oncologist or haematologist

PEGFILGRASTIM – **Restricted** see terms below

⚡ Inj 6 mg per 0.6 ml syringe .....	1,080.00	1	Neulastim
-------------------------------------	----------	---	-----------

#### →Restricted

For prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk  $\geq$  20%\*).

\*Febrile neutropenia risk  $\geq$  20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

## Fluids and Electrolytes

### Intravenous Administration

#### CALCIUM CHLORIDE

Inj 100 mg per ml, 10 ml vial

#### CALCIUM GLUCONATE

Inj 10%, 10 ml ampoule .....	21.40	10	Hospira
------------------------------	-------	----	---------

#### COMPOUND ELECTROLYTES

Inj sodium 140 mmol/l with potassium 5 mmol/l, magnesium 1.5 mmol/l, chloride 98 mmol/l, acetate 27 mmol/l and gluconate 23 mmol/l, bag .....	5.00	500 ml	Baxter
	3.10	1,000 ml	Baxter

#### COMPOUND ELECTROLYTES WITH GLUCOSE

Inj glucose 50 g with 140 mmol/l sodium, 5 mmol/l potassium, 1.5 mmol/l magnesium, 98 mmol/l chloride, 27 mmol/l acetate and 23 mmol/l gluconate, bag .....	7.00	1,000 ml	Baxter
---	------	----------	--------

## BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>COMPOUND SODIUM LACTATE [HARTMANN'S SOLUTION]</b>			
Inj sodium 131 mmol/l with potassium 5 mmol/l, calcium 2 mmol/l, bi-carbonate 29 mmol/l, chloride 111 mmol/l, bag .....	1.77	500 ml	Baxter
	1.80	1,000 ml	Baxter
<b>COMPOUND SODIUM LACTATE WITH GLUCOSE</b>			
Inj sodium 131 mmol/l with potassium 5 mmol/l, calcium 2 mmol/l, bi-carbonate 29 mmol/l, chloride 111 mmol/l and glucose 5%, bag .....	5.38	1,000 ml	Baxter
<b>GLUCOSE</b>			
Inj 5%, bag .....	2.87	50 ml	Baxter
	2.84	100 ml	Baxter
	3.87	250 ml	Baxter
	1.77	500 ml	Baxter
	1.80	1,000 ml	Baxter
Inj 10%, bag .....	3.70	500 ml	Baxter
	5.29	1,000 ml	Baxter
Inj 50%, bag .....	6.84	500 ml	Baxter
Inj 50%, 10 ml ampoule – 1% DV Sep-11 to 2014 .....	19.50	5	<b>Biomed</b>
Inj 50%, 90 ml bottle – 1% DV Sep-11 to 2014 .....	11.25	1	<b>Biomed</b>
Inj 70%, 1,000 ml bag			
Inj 70%, 500 ml bag			
<b>GLUCOSE WITH POTASSIUM CHLORIDE</b>			
Inj 5% glucose with 20 mmol/l potassium chloride, bag .....	7.36	1,000 ml	Baxter
Inj 5% glucose with 30 mmol/l potassium chloride, 1,000 ml bag			
Inj 10% glucose with 10 mmol/l potassium chloride, 500 ml bag			
<b>GLUCOSE WITH POTASSIUM CHLORIDE AND SODIUM CHLORIDE</b>			
Inj 4% glucose with potassium chloride 20 mmol/l and sodium chloride 0.18%, bag .....	3.45	500 ml	Baxter
	4.30	1,000 ml	Baxter
Inj 4% glucose with potassium chloride 30 mmol/l and sodium chloride 0.18%, bag .....	3.62	1,000 ml	Baxter
Inj 2.5% glucose with potassium chloride 20 mmol/l and sodium chloride 0.45%, 3,000 ml bag			
Inj 10% glucose with potassium chloride 10 mmol/l and sodium chloride 15 mmol/l, 500 ml bag			
<b>GLUCOSE WITH SODIUM CHLORIDE</b>			
Inj glucose 2.5% with sodium chloride 0.45%, bag .....	4.95	500 ml	Baxter
Inj glucose 5% with sodium chloride 0.45%, bag .....	9.87	500 ml	Baxter
	5.80	1,000 ml	Baxter
Inj glucose 5% with sodium chloride 0.9%, bag .....	4.54	1,000 ml	Baxter
Inj glucose 5% with sodium chloride 0.2%, 500 ml bag			
<b>POTASSIUM CHLORIDE</b>			
Inj 75 mg (1 mmol) per ml, 10 ml ampoule			
Inj 225 mg (3 mmol) per ml, 20 ml ampoule			



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>POTASSIUM CHLORIDE WITH SODIUM CHLORIDE</b>			
Inj 20 mmol/l potassium chloride with 0.9% sodium chloride, bag .....	3.85	1,000 ml	Baxter
Inj 30 mmol/l potassium chloride with 0.9% sodium chloride, bag .....	2.59	1,000 ml	Baxter
Inj 40 mmol/l potassium chloride with 0.9% sodium chloride, bag .....	6.62	1,000 ml	Baxter
Inj 10 mmol potassium chloride with 0.29% sodium chloride, 100 ml bag			
Inj 40 mmol/l potassium chloride with 0.9% sodium chloride, 100 ml bag			
<b>POTASSIUM DIHYDROGEN PHOSPHATE</b>			
Inj 1 mmol per ml, 10 ml ampoule			
<b>RINGER'S SOLUTION</b>			
Inj sodium 147 mmol/l with potassium 4 mmol/l, calcium 2.2 mmol/l, chloride 156 mmol/l, bag .....	5.13	1,000 ml	Baxter
<b>SODIUM ACETATE</b>			
Inj 4 mmol per ml, 20 ml ampoule			
<b>SODIUM BICARBONATE</b>			
Inj 8.4%, 10 ml vial			
Inj 8.4%, 50 ml vial .....	19.95	1	Biomed
Inj 8.4%, 100 ml vial .....	20.50	1	Biomed
<b>SODIUM CHLORIDE</b>			
Inj 0.45%, bag .....	5.50	500 ml	Baxter
☯ Inj 0.9%, 3 ml syringe			
☛ <b>Restricted</b>			
For use in flushing of in-situ vascular access devices only.			
Inj 0.9%, bag .....	1.70	500 ml	Freeflex
	1.71	1,000 ml	Freeflex
	3.01	50 ml	Baxter
	2.28	100 ml	Baxter
	3.60	250 ml	Baxter
	1.77	500 ml	Baxter
	1.80	1,000 ml	Baxter
☯ Inj 0.9%, 5 ml syringe			
☛ <b>Restricted</b>			
For use in flushing of in-situ vascular access devices only.			
☯ Inj 0.9%, 10 ml syringe			
☛ <b>Restricted</b>			
For use in flushing of in-situ vascular access devices only.			
Inj 3%, bag .....	5.69	1,000 ml	Baxter
Inj 0.9%, 5 ml ampoule .....	10.85	50	Multichem
	15.50		Pfizer
Inj 0.9%, 10 ml ampoule .....	11.50	50	Multichem
	15.50		Pfizer
Inj 0.9%, 20 ml ampoule .....	8.41	20	Multichem
Inj 23.4% (4 mmol/ml), 20 ml – 1% DV Sep-13 to 2016 .....	31.25	5	<b>Biomed</b>
Inj 1.8%, 500 ml bottle			
<b>SODIUM DIHYDROGEN PHOSPHATE [SODIUM ACID PHOSPHATE]</b>			
Inj 1 mmol per ml, 20 ml ampoule			

## BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>WATER</b>			
Inj, bag .....	2.75	1,000 ml	Baxter
Inj 5 ml ampoule .....	10.25	50	Multichem
Inj 10 ml ampoule .....	11.25	50	Multichem
Inj 20 ml ampoule .....	6.50	20	Multichem
Inj 250 ml bag			
Inj 500 ml bag			
<b>Oral Administration</b>			
<b>CALCIUM POLYSTYRENE SULPHONATE</b>			
Powder .....	169.85	300 g	Calcium Resonium
<b>COMPOUND ELECTROLYTES</b>			
Powder for oral soln			
<b>COMPOUND ELECTROLYTES WITH GLUCOSE</b>			
Soln with electrolytes			
<b>PHOSPHORUS</b>			
Tab eff 500 mg (16 mmol)			
<b>POTASSIUM CHLORIDE</b>			
Tab eff 548 mg (14 mmol) with chloride 285 mg (8 mmol)			
Tab long-acting 600 mg (8 mmol) – 1% DV Oct-12 to 2015 .....	7.42	200	Span-K
Oral liq 2 mmol per ml			
<b>SODIUM BICARBONATE</b>			
Cap 840 mg .....	8.52	100	Sodibic
<b>SODIUM CHLORIDE</b>			
Tab 600 mg			
Oral liq 2 mmol/ml			
<b>SODIUM POLYSTYRENE SULPHONATE</b>			
Powder			
<b>Plasma Volume Expanders</b>			
<b>GELATINE, SUCCINYLATED</b>			
Inj 4%, 500 ml bag .....	92.50	10	Gelafusal
	108.00		Gelofusine
<b>HYDROXYETHYL STARCH 130/0.4 WITH MAGNESIUM CHLORIDE, POTASSIUM CHLORIDE, SODIUM ACETATE AND SODIUM CHLORIDE</b>			
Inj 6% with magnesium chloride 0.03%, potassium chloride 0.03%, sodium acetate 0.463% and sodium chloride 0.6%, 500 ml bag .....	198.00	20	Volulyte 6%
<b>HYDROXYETHYL STARCH 130/0.4 WITH SODIUM CHLORIDE</b>			
Inj 6% with sodium chloride 0.9%, 500 ml bag .....	198.00	20	Voluven

Price (ex man. excl. GST)	Brand or Generic Manufacturer
\$ Per	

## Agents Affecting the Renin-Angiotensin System

### ACE Inhibitors

#### CAPTOPRIL

☞ Oral liq 5 mg per ml .....94.99 95 ml Capoten

#### ☞ Restricted

Any of the following:

- 1 For use in children under 12 years of age; or
- 2 For use in tube-fed patients; or
- 3 For management of rebound transient hypertension following cardiac surgery.

#### CILAZAPRIL

Tab 0.5 mg – 1% DV Sep-13 to 2016 .....	2.00	90	<b>Zapril</b>
Tab 2.5 mg – 1% DV Sep-13 to 2016 .....	4.31	90	<b>Zapril</b>
Tab 5 mg – 1% DV Sep-13 to 2016 .....	6.98	90	<b>Zapril</b>

#### ENALAPRIL MALEATE

Tab 5 mg .....	1.19	100	Ethics Enalapril
Tab 10 mg .....	1.47	100	Ethics Enalapril
Tab 20 mg .....	1.91	100	Ethics Enalapril

#### LISINOPRIL

Tab 5 mg – 1% DV Jan-13 to 2015 .....	3.58	90	<b>Arrow-Lisinopril</b>
Tab 10 mg – 1% DV Jan-13 to 2015 .....	4.08	90	<b>Arrow-Lisinopril</b>
Tab 20 mg – 1% DV Jan-13 to 2015 .....	4.88	90	<b>Arrow-Lisinopril</b>

#### PERINDOPRIL

Tab 2 mg .....	3.75	30	Apo-Perindopril
Tab 4 mg .....	4.80	30	Apo-Perindopril

#### QUINAPRIL

Tab 5 mg – 1% DV Apr-13 to 2015 .....	3.44	90	<b>Arrow-Quinapril 5</b>
Tab 10 mg – 1% DV Apr-13 to 2015 .....	4.64	90	<b>Arrow-Quinapril 10</b>
Tab 20 mg – 1% DV Apr-13 to 2015 .....	6.34	90	<b>Arrow-Quinapril 20</b>

TRANOLAPRIL – **Restricted**: For continuation only

☞ Cap 1 mg

☞ Cap 2 mg

### ACE Inhibitors with Diuretics

#### CILAZAPRIL WITH HYDROCHLOROTHIAZIDE

Tab 5 mg with hydrochlorothiazide 12.5 mg – 1% DV Mar-14 to 2016 .....	10.72	100	<b>Apo-Cilazapril/ Hydrochlorothiazide</b>
--	-------	-----	--

ENALAPRIL MALEATE WITH HYDROCHLOROTHIAZIDE – **Restricted**: For continuation only

☞ Tab 20 mg with hydrochlorothiazide 12.5 mg

#### QUINAPRIL WITH HYDROCHLOROTHIAZIDE

Tab 10 mg with hydrochlorothiazide 12.5 mg – 1% DV Aug-12 to 2015 .....	3.37	30	<b>Accuretic 10</b>
Tab 20 mg with hydrochlorothiazide 12.5 mg – 1% DV Aug-12 to 2015 .....	4.57	30	<b>Accuretic 20</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Angiotensin II Antagonists</b>			
CANDESARTAN CILEXETIL – <b>Restricted</b> see terms below			
⚡ Tab 4 mg – 1% DV Nov-12 to 2015 .....	4.13	90	<b>Candestar</b>
⚡ Tab 8 mg – 1% DV Nov-12 to 2015 .....	6.10	90	<b>Candestar</b>
⚡ Tab 16 mg – 1% DV Nov-12 to 2015 .....	10.18	90	<b>Candestar</b>
⚡ Tab 32 mg – 1% DV Nov-12 to 2015 .....	17.66	90	<b>Candestar</b>
➡ <b>Restricted</b>			
<b>ACE inhibitor intolerance</b>			
Either:			
1 Patient has persistent ACE inhibitor induced cough that is not resolved by ACE inhibitor retrial (same or new ACE inhibitor); or			
2 Patient has a history of angioedema.			
<b>Unsatisfactory response to ACE inhibitor</b>			
Patient is not adequately controlled on maximum tolerated dose of an ACE inhibitor.			
LOSARTAN POTASSIUM			
Tab 12.5 mg – 1% DV Dec-11 to 2014 .....	2.88	90	<b>Lostaar</b>
Tab 25 mg – 1% DV Dec-11 to 2014 .....	3.20	90	<b>Lostaar</b>
Tab 50 mg – 1% DV Dec-11 to 2014 .....	5.22	90	<b>Lostaar</b>
Tab 100 mg – 1% DV Dec-11 to 2014 .....	8.68	90	<b>Lostaar</b>
<b>Angiotensin II Antagonists with Diuretics</b>			
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE			
Tab 50 mg with hydrochlorothiazide 12.5 mg – 1% DV Dec-11 to 2014 .....	4.89	30	<b>Arrow-Losartan &amp; Hydrochlorothiazide</b>
<b>Alpha-Adrenoceptor Blockers</b>			
DOXAZOSIN			
Tab 2 mg – 1% DV Jun-11 to 2014 .....	8.23	500	<b>Apo-Doxazosin</b>
Tab 4 mg – 1% DV Jun-11 to 2014 .....	12.40	500	<b>Apo-Doxazosin</b>
PHENOXYBENZAMINE HYDROCHLORIDE			
Cap 10 mg			
Inj 50 mg per ml, 2 ml ampoule			
PHENTOLAMINE MESYLATE			
Inj 10 mg per ml, 1 ml ampoule			
PRAZOSIN			
Tab 1 mg .....	5.53	100	<b>Apo-Prazo Apo-Prazosin</b>
Tab 2 mg .....	7.00	100	<b>Apo-Prazo Apo-Prazosin</b>
Tab 5 mg .....	11.70	100	<b>Apo-Prazo Apo-Prazosin</b>
TERAZOSIN			
Tab 1 mg – 1% DV Sep-13 to 2016 .....	0.50	28	<b>Arrow</b>
Tab 2 mg – 1% DV Sep-13 to 2016 .....	0.45	28	<b>Arrow</b>
Tab 5 mg – 1% DV Sep-13 to 2016 .....	0.68	28	<b>Arrow</b>

Price (ex man. excl. GST)	Brand or Generic
\$	Per Manufacturer

## Antiarrhythmics

### ADENOSINE

Inj 3 mg per ml, 2 ml vial

⚡ Inj 3 mg per ml, 10 ml vial

#### ➡Restricted

For use in cardiac catheterisation, electrophysiology and MRI.

AJMALINE – **Restricted** see terms below

⚡ Inj 5 mg per ml, 10 ml ampoule

#### ➡Restricted

Cardiologist

### AMIODARONE HYDROCHLORIDE

Tab 100 mg

Tab 200 mg

Inj 50 mg per ml, 3 ml ampoule – 1% DV Aug-13 to 2016 .....22.80      6      **Cordarone-X**

### ATROPINE SULPHATE

Inj 600 mcg per ml, 1 ml ampoule – 1% DV Jan-13 to 2015 .....71.00      50      **AstraZeneca**

### DIGOXIN

Tab 62.5 mcg

Tab 250 mcg

Oral liq 50 mcg per ml

Inj 250 mcg per ml, 2 ml vial

### DISOPYRAMIDE PHOSPHATE

Cap 100 mg

Cap 150 mg

### FLECAINIDE ACETATE

Tab 50 mg .....45.82      60      Tambocor

Tab 100 mg .....80.92      60      Tambocor

Cap long-acting 100 mg .....45.82      30      Tambocor CR

Cap long-acting 200 mg .....80.92      30      Tambocor CR

Inj 10 mg per ml, 15 ml ampoule .....52.45      5      Tambocor

### MEXILETINE HYDROCHLORIDE

Cap 150 mg .....65.00      100      Mexiletine Hydrochloride  
USP

Cap 250 mg .....102.00      100      Mexiletine Hydrochloride  
USP

### PROPAFENONE HYDROCHLORIDE

Tab 150 mg

## Antihypotensives

MIDODRINE – **Restricted** see terms on the next page

⚡ Tab 2.5 mg

⚡ Tab 5 mg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➡Restricted</b>			
All of the following:			
1 Disabling orthostatic hypotension not due to drugs; and			
2 Patient has tried fludrocortisone (unless contra-indicated) with unsatisfactory results; and			
3 Patient has tried non-pharmacological treatments such as support hose, increased salt intake, exercise, and elevation of head and trunk at night.			
<b>Beta-Adrenoceptor Blockers</b>			
<b>ATENOLOL</b>			
Tab 50 mg – 1% DV Oct-12 to 2015 .....	5.56	500	Mylan Atenolol
Tab 100 mg – 1% DV Oct-12 to 2015 .....	9.12	500	Mylan Atenolol
Oral liq 5 mg per ml .....	21.25	300 ml	Atenolol-AFT
<b>BISOPROLOL</b>			
Tab 2.5 mg .....	3.88	30	Bosvate
Tab 5 mg .....	4.74	30	Bosvate
Tab 10 mg .....	9.18	30	Bosvate
<b>CARVEDILOL</b>			
Tab 6.25 mg .....	21.00	30	Dilatrend
Tab 12.5 mg .....	27.00	30	Dilatrend
Tab 25 mg .....	33.75	30	Dilatrend
<b>CELIPROLOL</b>			
Tab 200 mg .....	19.00	180	Celol
<b>ESMOLOL HYDROCHLORIDE</b>			
Inj 10 mg per ml, 10 ml vial			
<b>LABETALOL</b>			
Tab 50 mg .....	8.23	100	Hybloc
Tab 100 mg .....	10.06	100	Hybloc
Tab 200 mg .....	17.55	100	Hybloc
Tab 400 mg			
Inj 5 mg per ml, 20 ml ampoule			
<b>METOPROLOL SUCCINATE</b>			
Tab long-acting 23.75 mg – 1% DV Sep-12 to 2015 .....	0.96	30	Metoprolol - AFT CR
Tab long-acting 47.5 mg – 1% DV Sep-12 to 2015 .....	1.41	30	Metoprolol - AFT CR
Tab long-acting 95 mg – 1% DV Sep-12 to 2015 .....	2.42	30	Metoprolol - AFT CR
Tab long-acting 190 mg – 1% DV Sep-12 to 2015 .....	4.66	30	Metoprolol - AFT CR
<b>METOPROLOL TARTRATE</b>			
Tab 50 mg – 1% DV Aug-12 to 2015 .....	16.00	100	Lopresor
Tab 100 mg – 1% DV Aug-12 to 2015 .....	21.00	60	Lopresor
Tab long-acting 200 mg – 1% DV Aug-12 to 2015 .....	18.00	28	Slow-Lopresor
Inj 1 mg per ml, 5 ml vial – 1% DV Dec-12 to 2015 .....	24.00	5	Lopresor
<b>NADOLOL</b>			
Tab 40 mg – 1% DV Apr-13 to 2015 .....	15.57	100	Apo-Nadolol
Tab 80 mg – 1% DV Apr-13 to 2015 .....	23.74	100	Apo-Nadolol
<b>PINDOLOL</b>			
Tab 5 mg – 1% DV Nov-13 to 2016 .....	9.72	100	Apo-Pindolol
Tab 10 mg – 1% DV Nov-13 to 2016 .....	15.62	100	Apo-Pindolol
Tab 15 mg – 1% DV Nov-13 to 2016 .....	23.46	100	Apo-Pindolol

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PROPRANOLOL</b>			
Tab 10 mg .....	3.65	100	Apo-Propranolol
Tab 40 mg .....	4.65	100	Apo-Propranolol
Cap long-acting 160 mg .....	16.06	100	Cardinol LA
Oral liq 4 mg per ml			
Inj 1 mg per ml, 1 ml ampoule			
<b>SOTALOL</b>			
Tab 80 mg .....	27.50	500	Mylan
Tab 160 mg .....	10.50	100	Mylan
Inj 10 mg per ml, 4 ml ampoule .....	65.39	5	Sotacor
<b>TIMOLOL MALEATE</b>			
Tab 10 mg			
<b>Calcium Channel Blockers</b>			
<b>Dihydropyridine Calcium Channel Blockers</b>			
<b>AMLODIPINE</b>			
Tab 2.5 mg – 1% DV Mar-12 to 2014 .....	2.45	100	<b>Apo-Amlodipine</b>
Tab 5 mg – 1% DV Oct-11 to 2014 .....	2.65	100	<b>Apo-Amlodipine</b>
Tab 10 mg – 1% DV Oct-11 to 2014 .....	4.15	100	<b>Apo-Amlodipine</b>
<b>FELODIPINE</b>			
Tab long-acting 2.5 mg – 1% DV Sep-12 to 2015 .....	2.90	30	<b>Plendil ER</b>
Tab long-acting 5 mg – 1% DV Nov-12 to 2015 .....	3.10	30	<b>Plendil ER</b>
Tab long-acting 10 mg – 1% DV Nov-12 to 2015 .....	4.60	30	<b>Plendil ER</b>
<b>ISRADIPINE</b>			
Tab 2.5 mg			
Cap long-acting 2.5 mg			
Cap long-acting 5 mg			
<b>NIFEDIPINE</b>			
Tab long-acting 10 mg			
Tab long-acting 20 mg .....	9.59	100	Nyefax Retard
Tab long-acting 30 mg .....	8.56	30	Adefin XL
			Arrow-Nifedipine XR
Tab long-acting 60 mg .....	12.28	30	Adefin XL
			Arrow-Nifedipine XR
Cap 5 mg			
<b>NIMODIPINE</b>			
Tab 30 mg			
Inj 200 mcg per ml, 50 ml vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Other Calcium Channel Blockers</b>			
<b>DILTIAZEM HYDROCHLORIDE</b>			
Tab 30 mg – 5% DV Sep-12 to 2015 .....	4.60	100	<b>Dilzem</b>
Tab 60 mg – 5% DV Sep-12 to 2015 .....	8.50	100	<b>Dilzem</b>
Cap long-acting 120 mg .....	1.91	30	Cardizem CD
	31.83	500	Apo-Diltiazem CD
Cap long-acting 180 mg .....	7.56	30	Cardizem CD
	47.67	500	Apo-Diltiazem CD
Cap long-acting 240 mg .....	10.22	30	Cardizem CD
	63.58	500	Apo-Diltiazem CD
Inj 5 mg per ml, 5 ml vial			
<b>PERHEXILINE MALEATE – Restricted</b> see terms below			
⚡ Tab 100 mg .....	62.90	100	Pexsig
<b>➡Restricted</b>			
Both:			
1 Patient has refractory angina; and			
2 Patient is on the maximal tolerated dose of a beta-blocker, a calcium channel blocker and a long-acting nitrate.			
<b>VERAPAMIL HYDROCHLORIDE</b>			
Tab 40 mg – 1% DV Sep-11 to 2014 .....	7.01	100	<b>Isoptin</b>
Tab 80 mg – 1% DV Sep-11 to 2014 .....	11.74	100	<b>Isoptin</b>
Tab long-acting 120 mg .....	15.20	250	Verpamil SR
Tab long-acting 240 mg .....	25.00	250	Verpamil SR
Inj 2.5 mg per ml, 2 ml ampoule .....	7.54	5	Isoptin
<b>Centrally-Acting Agents</b>			
<b>CLONIDINE</b>			
Patch 2.5 mg, 100 mcg per day – 1% DV Jul-14 to 2017 .....	12.80	4	<b>Catapres-TTS-1</b>
Patch 5 mg, 200 mcg per day – 1% DV Jul-14 to 2017 .....	18.04	4	<b>Catapres-TTS-2</b>
Patch 7.5 mg, 300 mcg per day – 1% DV Jul-14 to 2017 .....	22.68	4	<b>Catapres-TTS-3</b>
<b>CLONIDINE HYDROCHLORIDE</b>			
Tab 25 mcg – 1% DV Jul-13 to 2015 .....	15.09	112	<b>Clonidine BNM</b>
Tab 150 mcg – 1% DV Feb-13 to 2015 .....	34.32	100	<b>Catapres</b>
Inj 150 mcg per ml, 1 ml ampoule – 1% DV Nov-12 to 2015 .....	16.07	5	<b>Catapres</b>
<b>METHYLDOPA</b>			
Tab 125 mg .....	14.25	100	Prodopa
Tab 250 mg .....	15.10	100	Prodopa
Tab 500 mg .....	23.15	100	Prodopa
<b>Diuretics</b>			
<b>Loop Diuretics</b>			
<b>BUMETANIDE</b>			
Tab 1 mg .....	16.36	100	Burinex
Inj 500 mcg per ml, 4 ml vial			



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>FUROSEMIDE (FRUSEMIDE)</b>			
Tab 40 mg – 1% DV Sep-12 to 2015 .....	10.25	1,000	<b>Diurin 40</b>
Tab 500 mg – 1% DV Feb-13 to 2015.....	25.00	50	<b>Urex Forte</b>
Oral liq 10 mg per ml .....			
Inj 10 mg per ml, 2 ml ampoule .....	1.30	5	Frusemide-Claris
Inj 10 mg per ml, 25 ml ampoule .....			

## Osmotic Diuretics

<b>MANNITOL</b>			
Inj 10%, 1,000 ml bag .....	14.21	1,000 ml	Baxter
Inj 15%, 500 ml bag .....	9.84	500 ml	Baxter
Inj 20%, 500 ml bag .....	10.80	500 ml	Baxter

## Potassium Sparing Combination Diuretics

<b>AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE</b>			
Tab 5 mg with furosemide 40 mg .....			
<b>AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE</b>			
Tab 5 mg with hydrochlorothiazide 50 mg .....			

## Potassium Sparing Diuretics

<b>AMILORIDE HYDROCHLORIDE</b>			
Tab 5 mg .....	17.50	100	Apo-Amiloride
Oral liq 1 mg per ml .....	30.00	25 ml	Biomed
<b>SPIRONOLACTONE</b>			
Tab 25 mg – 1% DV Sep-13 to 2016 .....	3.65	100	<b>Spiractin</b>
Tab 100 mg – 1% DV Sep-13 to 2016 .....	11.80	100	<b>Spiractin</b>
Oral liq 5 mg per ml .....	30.00	25 ml	<b>Spirotone</b>
<i>(Spirotone Tab 100 mg to be delisted 1 August 2014)</i>			

## Thiazide and Related Diuretics

<b>BENDROFLUMETHAZIDE [BENDROFLUAZIDE]</b>			
Tab 2.5 mg – 1% DV Sep-11 to 2014 .....	6.48	500	<b>Arrow-Bendrofluazide</b>
Tab 5 mg – 1% DV Sep-11 to 2014 .....	9.95	500	<b>Arrow-Bendrofluazide</b>
<b>CHLOROTHIAZIDE</b>			
Oral liq 50 mg per ml .....	26.00	25 ml	Biomed
<b>CHLORTALIDONE [CHLORTHALIDONE]</b>			
Tab 25 mg .....	8.00	50	Hygroton
<b>INDAPAMIDE</b>			
Tab 2.5 mg – 1% DV Oct-13 to 2016 .....	2.25	90	<b>Dapa-Tabs</b>

**METOLAZONE – Restricted** see terms below

⬇ Tab 5 mg

➡ **Restricted**

Either:

- 1 Patient has refractory heart failure and is intolerant or has not responded to loop diuretics and/or loop-thiazide combination therapy; or
- 2 Patient has severe refractory nephrotic oedema unresponsive to high dose loop diuretics and concentrated albumin infusions

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Lipid-Modifying Agents</b>			
<b>Fibrates</b>			
<b>BEZAFIBRATE</b>			
Tab 200 mg – 1% DV Mar-13 to 2015 .....	9.70	90	<b>Bezalip</b>
Tab long-acting 400 mg – 1% DV Oct-12 to 2015 .....	5.70	30	<b>Bezalip Retard</b>
<b>GEMFIBROZIL</b>			
Tab 600 mg – 1% DV Nov-13 to 2016 .....	17.60	60	<b>Lipazil</b>
<b>HMG CoA Reductase Inhibitors (Statins)</b>			
<b>ATORVASTATIN</b>			
Tab 10 mg – 1% DV Oct-12 to 2015 .....	2.52	90	<b>Zarator</b>
Tab 20 mg – 1% DV Oct-12 to 2015 .....	4.17	90	<b>Zarator</b>
Tab 40 mg – 1% DV Oct-12 to 2015 .....	7.32	90	<b>Zarator</b>
Tab 80 mg – 1% DV Oct-12 to 2015 .....	16.23	90	<b>Zarator</b>
<b>PRAVASTATIN</b>			
Tab 10 mg			
Tab 20 mg – 1% DV Nov-11 to 2014 .....	5.44	30	<b>Cholvastin</b>
Tab 40 mg – 1% DV Nov-11 to 2014 .....	9.28	30	<b>Cholvastin</b>
<b>SIMVASTATIN</b>			
Tab 10 mg – 1% DV Sep-11 to 2014 .....	1.40	90	<b>Arrow-Simva</b>
Tab 20 mg – 1% DV Sep-11 to 2014 .....	1.95	90	<b>Arrow-Simva</b>
Tab 40 mg – 1% DV Sep-11 to 2014 .....	3.18	90	<b>Arrow-Simva</b>
Tab 80 mg – 1% DV Sep-11 to 2014 .....	9.31	90	<b>Arrow-Simva</b>
<b>Resins</b>			
<b>CHOLESTYRAMINE</b>			
Powder for oral liq 4 g			
<b>COLESTIPOL HYDROCHLORIDE</b>			
Grans for oral liq 5 g			
<b>Selective Cholesterol Absorption Inhibitors</b>			
<b>EZETIMIBE – Restricted</b> see terms below			
☞ Tab 10 mg			
☞ <b>Restricted</b>			
All of the following:			
1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and			
2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and			
3 Any of the following:			
3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 × normal) when treated with one statin; or			
3.2 The patient is intolerant to both simvastatin and atorvastatin; or			
3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.			

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

#### EZETIMIBE WITH SIMVASTATIN – **Restricted** see terms below

- ⚡ Tab 10 mg with simvastatin 10 mg
- ⚡ Tab 10 mg with simvastatin 20 mg
- ⚡ Tab 10 mg with simvastatin 40 mg
- ⚡ Tab 10 mg with simvastatin 80 mg

#### ➡ **Restricted**

All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

### Other Lipid-Modifying Agents

#### ACIPIMOX

Cap 250 mg

#### NICOTINIC ACID

Tab 50 mg

Tab 500 mg

### Nitrates

#### GLYCERYL TRINITRATE

Tab 600 mcg – 1% DV Sep-11 to 2014.....	8.00	100	<b>Lycinate</b>
Inj 1 mg per ml, 5 ml ampoule – 1% DV Dec-12 to 2015.....	22.70	10	<b>Nitronal</b>
Inj 1 mg per ml, 50 ml vial – 1% DV Dec-12 to 2015.....	86.60	10	<b>Nitronal</b>
Inj 5 mg per ml, 10 ml ampoule .....	40.00	5	Hospira
Oral spray, 400 mcg per dose – 1% DV Mar-12 to 2014.....	4.45	250 dose	<b>Glytrin</b>
Patch 25 mg, 5 mg per day – 1% DV Sep-11 to 2014 .....	16.56	30	<b>Nitroderm TTS 5</b>
Patch 50 mg, 10 mg per day – 1% DV Sep-11 to 2014 .....	19.50	30	<b>Nitroderm TTS 10</b>

#### ISOSORBIDE MONONITRATE

Tab 20 mg – 1% DV Jun-11 to 2014.....	17.10	100	<b>Ismo-20</b>
Tab long-acting 40 mg .....	7.50	30	Corangin
Tab long-acting 60 mg .....	3.94	90	Ismo 40 Retard

(Corangin Tab long-acting 40 mg to be delisted 1 August 2014)

### Other Cardiac Agents

#### LEVOSIMENDAN – **Restricted** see terms below

- ⚡ Inj 2.5 mg per ml, 5 ml vial
- ⚡ Inj 2.5 mg per ml, 10 ml vial

#### ➡ **Restricted**

#### Heart transplant

Either:

- 1 For use as a bridge to heart transplant, in patients who have been accepted for transplant; or
- 2 For the treatment of heart failure following heart transplant.

#### Heart failure - cardiologist or intensivist

For the treatment of severe acute decompensated heart failure that is non-responsive to dobutamine.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Sympathomimetics</b>			
<b>ADRENALINE</b>			
Inj 1 in 1,000, 1 ml ampoule .....	4.98	5	Aspen Adrenaline
	5.25		Hospira
Inj 1 in 1,000, 30 ml vial			
Inj 1 in 10,000, 10 ml ampoule .....	27.00	5	Hospira
	49.00	10	Aspen Adrenaline
Inj 1 in 10,000, 10 ml syringe			
<b>DOBUTAMINE HYDROCHLORIDE</b>			
Inj 12.5 mg per ml, 20 ml vial			
<b>DOPAMINE HYDROCHLORIDE</b>			
Inj 40 mg per ml, 5 ml ampoule – 1% DV Sep-12 to 2015 .....	69.77	10	Martindale
<b>EPHEDRINE</b>			
Inj 3 mg per ml, 10 ml syringe			
Inj 30 mg per ml, 1 ml ampoule – 1% DV Nov-12 to 2014 .....	66.00	10	Max Health
<b>ISOPRENALINE</b>			
Inj 200 mcg per ml, 1 ml ampoule			
Inj 200 mcg per ml, 5 ml ampoule			
<b>METARAMINOL</b>			
Inj 0.5 mg per ml, 20 ml syringe			
Inj 1 mg per ml, 1 ml ampoule			
Inj 1 mg per ml, 10 ml syringe			
Inj 10 mg per ml, 1 ml ampoule			
<b>NORADRENALINE</b>			
Inj 0.06 mg per ml, 100 ml bag			
Inj 0.06 mg per ml, 50 ml syringe			
Inj 0.1 mg per ml, 100 ml bag			
Inj 0.12 mg per ml, 100 ml bag			
Inj 0.12 mg per ml, 50 ml syringe			
Inj 0.16 mg per ml, 50 ml syringe			
Inj 1 mg per ml, 100 ml bag			
Inj 1 mg per ml, 2 ml ampoule .....	42.00	6	Levophed
<b>PHENYLEPHRINE HYDROCHLORIDE</b>			
Inj 10 mg per ml, 1 ml vial .....	115.50	25	Neosynephrine HCL
<b>Vasodilators</b>			
<b>ALPROSTADIL HYDROCHLORIDE</b>			
Inj 500 mcg per ml, 1 ml ampoule – 1% DV Oct-12 to 2015 .....	1,417.50	5	Prostin VR
<b>AMYL NITRITE</b>			
Liq 98% in 3 ml capsule			
<b>DIAZOXIDE</b>			
Inj 15 mg per ml, 20 ml ampoule			
<b>HYDRALAZINE HYDROCHLORIDE</b>			
⚡ Tab 25 mg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➔Restricted</b>			
Either:			
1 For the treatment of refractory hypertension; or			
2 For the treatment of heart failure, in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.			
Inj 20 mg ampoule .....	25.90	5	Apresoline
<b>MILRINONE</b>			
Inj 1 mg per ml, 10 ml ampoule			
<b>MINOXIDIL – Restricted</b> see terms below			
⚡ Tab 10 mg .....	70.00	100	Loniten
<b>➔Restricted</b>			
For patients with severe refractory hypertension who have failed to respond to extensive multiple therapies.			
<b>NICORANDIL – Restricted</b> see terms below			
⚡ Tab 10 mg .....	27.95	60	Ikorel
⚡ Tab 20 mg .....	33.28	60	Ikorel
<b>➔Restricted</b>			
Both:			
1 Patient has refractory angina; and			
2 Patient is on the maximal tolerated dose of a beta-blocker, a calcium channel blocker and a long-acting nitrate.			
<b>PAPAVERINE HYDROCHLORIDE</b>			
Inj 30 mg per ml, 1 ml vial			
Inj 12 mg per ml, 10 ml ampoule .....	73.12	5	Hospira
<b>PENTOXIFYLLINE [OXPENTIFYLLINE]</b>			
Tab 400 mg			
<b>SODIUM NITROPRUSSIDE</b>			
Inj 50 mg vial			

### Endothelin Receptor Antagonists

<b>AMBRISENTAN – Restricted</b> see terms below			
⚡ Tab 5 mg .....	4,585.00	30	Volibris
⚡ Tab 10 mg .....	4,585.00	30	Volibris
<b>➔Restricted</b>			
1 For use in patients with approval by the Pulmonary Arterial Hypertension Panel; or			
2 In hospital stabilisations in emergency situations.			
<b>BOSENTAN – Restricted</b> see terms below			
⚡ Tab 62.5 mg .....	1,500.00	60	pms-Bosentan
	4,585.00		Tracleer
⚡ Tab 125 mg .....	1,500.00	60	pms-Bosentan
	4,585.00		Tracleer
<b>➔Restricted</b>			
1 For use in patients with approval by the Pulmonary Arterial Hypertension Panel; or			
2 In hospital stabilisation in emergency situations.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Phosphodiesterase Type 5 Inhibitors</b>			
SILDENAFIL – <b>Restricted</b> see terms below			
☞ Tab 25 mg – 1% DV May-13 to 2014 .....	1.85	4	<b>Silagra</b>
☞ Tab 50 mg – 1% DV May-13 to 2014 .....	1.85	4	<b>Silagra</b>
☞ Tab 100 mg – 1% DV May-13 to 2014 .....	7.45	4	<b>Silagra</b>

**☞Restricted**

Any of the following:

- 1 For use in patients with approval by the Pulmonary Arterial Hypertension Panel; or
- 2 For use in neonatal units for persistent pulmonary hypertension of the newborn (PPHN); or
- 3 For use in weaning patients from inhaled nitric oxide; or
- 4 For perioperative use in cardiac surgery patients; or
- 5 For use in intensive care as an alternative to nitric oxide; or
- 6 In-hospital stabilisation in emergency situations; or
- 7 All of the following:
  - 7.1 Patient has Raynaud's phenomenon; and
  - 7.2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
  - 7.3 Patient is following lifestyle management (proper body insulation, avoidance of cold exposure, smoking cessation support, avoidance of sympathomimetic drugs); and
  - 7.4 Patient has persisting severe symptoms despite treatment with calcium channel blockers and nitrates (unless contraindicated or not tolerated).

**Prostacyclin Analogues**
**ILOPROST**

Inj 50 mcg in 0.5 ml ampoule – 1% DV Apr-14 to 2016 .....	925.00	5	<b>Ilomedin</b>
☞ Nebuliser soln 10 mcg per ml, 2 ml .....	1,185.00	30	<b>Ventavis</b>

**☞Restricted**

Any of the following:

- 1 For use in patients with approval by the Pulmonary Arterial Hypertension Panel; or
- 2 For diagnostic use in catheter laboratories; or
- 3 For use following mitral or tricuspid valve surgery; or
- 4 In-hospital stabilisation in emergency situations.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Anti-Infective Preparations</b>			
<b>Antibacterials</b>			
FUSIDATE SODIUM [FUSIDIC ACID]			
Crm 2% .....	3.25	15 g	Foban
Oint 2% – <b>1% DV Sep-13 to 2016</b> .....	3.45	15 g	<b>Foban</b>
HYDROGEN PEROXIDE			
Crm 1% .....	8.56	15 g	Crystaderm
Soln 3% (10 vol)			
MAFENIDE ACETATE – <b>Restricted</b> see terms below			
↓ Powder 50 g sachet			
➔ <b>Restricted</b>			
For the treatment of burns patients.			
MUPIROCIN			
Oint 2%			
SULPHADIAZINE SILVER			
Crm 1% .....	12.30	50 g	Flamazine
<b>Antifungals</b>			
AMOROLFINE – <b>Restricted:</b> For continuation only			
➔ Nail soln 5%			
CICLOPIROX OLAMINE			
Nail soln 8%			
➔ Soln 1% – <b>Restricted:</b> For continuation only			
CLOTTRIMAZOLE			
Crm 1% – <b>1% DV Nov-11 to 2014</b> .....	0.54	20 g	<b>Clomazol</b>
➔ Soln 1% – <b>Restricted:</b> For continuation only			
ECONAZOLE NITRATE			
➔ Crm 1% – <b>Restricted:</b> For continuation only			
Foaming soln 1%			
KETOCONAZOLE			
Shampoo 2% – <b>1% DV Sep-11 to 2014</b> .....	3.08	100 ml	<b>Sebizole</b>
METRONIDAZOLE			
Gel 0.75%			
MICONAZOLE NITRATE			
Crm 2% – <b>1% DV Nov-11 to 2014</b> .....	0.46	15 g	<b>Multichem</b>
➔ Lotn 2% – <b>Restricted:</b> For continuation only			
Tinc 2%			
NYSTATIN			
Crm 100,000 u per g			
<b>Antiparasitics</b>			
LINDANE [GAMMA BENZENE HEXACHLORIDE]			
Crm 1%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MALATHION [MALDISON] Lotn 0.5% Shampoo 1%			
MALATHION WITH PERMETHRIN AND PIPERONYL BUTOXIDE Spray 0.25% with permethrin 0.5% and piperonyl butoxide 2% Note: Temporary listing to cover out-of-stock.			
PERMETHRIN			
Crm 5% – 1% DV Sep-11 to 2014 .....	4.20	30 g	Lyderm
Lotn 5% – 1% DV Sep-11 to 2014 .....	3.24	30 ml	A-Scabies

## Antiacne Preparations

ADAPALENE			
Crm 0.1%			
Gel 0.1%			
BENZOYL PEROXIDE			
Soln 5%			
ISOTRETINOIN			
Cap 10 mg – 1% DV Jan-13 to 2015 .....	18.71	120	Oratane
Cap 20 mg – 1% DV Jan-13 to 2015 .....	28.91	120	Oratane
TRETINOIN			
Crm 0.05%			

## Antipruritic Preparations

CALAMINE			
Crm, aqueous, BP – 1% DV Mar-13 to 2015 .....	1.77	100 g	Pharmacy Health
Lotn, BP – 1% DV Nov-12 to 2015 .....	13.45	2,000 ml	PSM
CROTAMITON			
Crm 10% – 1% DV Sep-12 to 2015 .....	3.48	20 g	Itch-Soothe

## Barrier Creams and Emollients

### Barrier Creams

DIMETHICONE			
Crm 5% tube – 1% DV Apr-14 to 2016 .....	1.65	100 g	healthE Dimethicone 5%
Crm 5% pump bottle – 1% DV Apr-14 to 2016 .....	4.73	500 ml	healthE Dimethicone 5%
ZINC			
Crm			e.g. Zinc Cream (Orion); Zinc Cream (PSM)
Oint			e.g. Zinc oxide (PSM)
Paste			
ZINC AND CASTOR OIL			
Crm – 1% DV Apr-12 to 2014 .....	1.63	20 g	Orion
Oint, BP			



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ZINC WITH WOOL FAT			
Crm zinc 15.25% with wool fat 4%			<i>e.g. Sudocrem</i>
<b>Emollients</b>			
<b>AQUEOUS CREAM</b>			
Crm 100 g – 1% <b>DV Sep-11 to 2014</b> .....	1.23	100 g	<b>AFT</b>
Note: DV limit applies to the pack sizes of 100 g or less.			
Crm 500 g – 1% <b>DV Sep-11 to 2014</b> .....	1.96	500 g	<b>AFT</b>
Note: DV limit applies to the pack sizes of greater than 100 g.			
<b>CETOMACROGOL</b>			
Crm BP, 500 g .....	3.50	500 g	Pharmacy Health
Crm BP, 100 g .....	1.65	1	healthE
<b>CETOMACROGOL WITH GLYCEROL</b>			
Crm 90% with glycerol 10%, .....	2.10	100 g	Pharmacy Health
	2.00		Pharmacy Health
	3.20		healthE
Crm 90% with glycerol 10% .....	4.50	500 ml	Pharmacy Health
			Sorbolene with
			Glycerin
	6.50	1,000 ml	Pharmacy Health
			Sorbolene with
			Glycerin
Crm 90% with glycerol 10%, 500 ml, 1 bottle .....	5.46	1	healthE
<b>EMULSIFYING OINTMENT</b>			
Oint BP – 1% <b>DV Nov-11 to 2014</b> .....	1.95	100 g	<b>Jaychem</b>
Oint BP, 500 g – 1% <b>DV Sep-11 to 2014</b> .....	3.04	500 g	<b>AFT</b>
Note: DV limit applies to pack sizes of greater than 100 g.			
<b>GLYCEROL WITH PARAFFIN</b>			
Crm glycerol 10% with white soft paraffin 5% and liquid paraffin 10%			<i>e.g. QV cream</i>
<b>OIL IN WATER EMULSION</b>			
Crm – 1% <b>DV Dec-12 to 2015</b> .....	2.63	500 g	<b>healthE Fatty Cream</b>
Crm, 100 g .....	1.60	1	<b>healthE Fatty Cream</b>
<b>PARAFFIN</b>			
Oint liquid paraffin 50% with white soft paraffin 50% .....	3.10	100 g	healthE
White soft – 1% <b>DV Feb-13 to 2015</b> .....	0.92	10 g	<b>healthE</b>
Note: DV limit applies to pack sizes of 30 g or less, and to both white soft paraffin and yellow soft paraffin.			
Yellow soft			
<b>PARAFFIN WITH WOOL FAT</b>			
Lotn liquid paraffin 15.9% with wool fat 0.6%			<i>e.g. AlphaKeri;BK ;DP; Hydroderm Lotn</i>
Lotn liquid paraffin 91.7% with wool fat 3%			<i>e.g. Alpha Keri Bath Oil</i>
<b>UREA</b>			
Crm 10%			
<b>WOOL FAT</b>			
Crm			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Corticosteroids</b>			
<b>BETAMETHASONE DIPROPIONATE</b>			
Crm 0.05%			
Oint 0.05%			
<b>BETAMETHASONE VALERATE</b>			
Crm 0.1%			
Oint 0.1%			
Lotn 0.1%			
<b>CLOBETASOL PROPIONATE</b>			
Crm 0.05% .....	3.68	30 g	Dermol
Oint 0.05% .....	3.68	30 g	Dermol
<b>CLOBETASONE BUTYRATE</b>			
Crm 0.05%			
<b>DIFLUCORTOLONE VALERATE – Restricted:</b> For continuation only			
➡ Crm 0.1%			
➡ Fatty oint 0.1%			
<b>HYDROCORTISONE</b>			
Crm 1%, 100 g .....	3.75	100 g	Pharmacy Health
Crm 1%, 500 g – 1% DV Nov-11 to 2014 .....	14.00	500 g	Pharmacy Health
Note: DV limit applies to the pack sizes of greater than 100 g.			
<b>HYDROCORTISONE ACETATE</b>			
Crm 1% .....	2.48	14.2 g	AFT
<b>HYDROCORTISONE BUTYRATE</b>			
Crm 0.1% – 1% DV Mar-13 to 2015 .....	2.30	30 g	Locoid Lipocream
	6.85	100 g	Locoid Lipocream
Oint 0.1% – 1% DV Mar-13 to 2015 .....	6.85	100 g	Locoid
Milky emul 0.1% – 1% DV Mar-13 to 2015 .....	6.85	100 ml	Locoid Crelo
<b>HYDROCORTISONE WITH PARAFFIN AND WOOL FAT</b>			
Lotn 1% with paraffin liquid 15.9% and wool fat 0.6%			
<b>METHYLPREDNISOLONE ACEPONATE</b>			
Crm 0.1% .....	4.95	15 g	Advantan
Oint 0.1% .....	4.95	15 g	Advantan
<b>MOMETASONE FUROATE</b>			
Crm 0.1% – 1% DV Sep-12 to 2015 .....	1.78	15 g	m-Mometasone
	3.42	45 g	m-Mometasone
Oint 0.1% – 1% DV Sep-12 to 2015 .....	1.78	15 g	m-Mometasone
	3.42	45 g	m-Mometasone
Lotn 0.1%			
<b>TRIAMCINOLONE ACETONIDE</b>			
Crm 0.02% – 1% DV Sep-11 to 2014 .....	6.63	100 g	Aristocort
Oint 0.02% – 1% DV Sep-11 to 2014 .....	6.69	100 g	Aristocort

**Corticosteroids with Anti-Infective Agents**

**BETAMETHASONE VALERATE WITH CLIOQUINOL – Restricted** see terms on the next page

⚡ Crm 0.1% with clioquinol 3%

⚡ Oint 0.1% with clioquinol 3%

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➔ Restricted</b>			
Either:			
1 For the treatment of intertrigo; or			
2 For continuation use			
<b>BETAMETHASONE VALERATE WITH FUSIDIC ACID</b>			
Crm 0.1% with fusidic acid 2%			
<b>HYDROCORTISONE WITH MICONAZOLE</b>			
Crm 1% with miconazole nitrate 2% .....	2.20	15 g	Micreme H
<b>HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN</b>			
Crm 1% with natamycin 1% and neomycin sulphate 0.5% .....	2.79	15 g	Pimafucort
Oint 1% with natamycin 1% and neomycin sulphate 0.5% .....	2.79	15 g	Pimafucort
<b>TRIAMCINOLONE ACETONIDE WITH NEOMYCIN SULPHATE, GRAMICIDIN AND NYSTATIN</b>			
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g			

### Psoriasis and Eczema Preparations

<b>ACITRETIN</b>			
Cap 10 mg .....	35.95	100	Neotigason
	38.66	60	Novatretin
Cap 25 mg .....	83.11	60	Novatretin
	85.40	100	Neotigason
<b>BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL</b>			
Gel 500 mcg with calcipotriol 50 mcg per g .....	26.12	30 g	Daivobet
Oint 500 mcg with calcipotriol 50 mcg per g .....	26.12	30 g	Daivobet
<b>CALCIPOTRIOL</b>			
Crm 50 mcg per g .....	45.00	100 g	Daivonex
Oint 50 mcg per g .....	45.00	100 g	Daivonex
Soln 50 mcg per ml .....	16.00	30 ml	Daivonex
<b>COAL TAR WITH SALICYLIC ACID AND SULPHUR</b>			
Oint 12% with salicylic acid 2% and sulphur 4%			
<b>COAL TAR WITH TRIETHANOLAMINE LARYL SULPHATE AND FLUORESCEIN</b>			
Soln 2.3% with triethanolamine lauryl sulphate and fluorescein sodium – 1% DV Nov-11 to 2014 .....	3.05	500 ml	Pinetarsol
	5.82	1,000 ml	Pinetarsol
<b>METHOXSALEN [8-METHOXYPSORALEN]</b>			
Cap 10 mg			
Lotn 1.2%			
<b>POTASSIUM PERMANGANATE</b>			
Tab 400 mg			
Crystals			

### Scalp Preparations

<b>BETAMETHASONE VALERATE</b>			
Scalp app 0.1% .....	7.75	100 ml	Beta Scalp
<b>CLOBETASOL PROPIONATE</b>			
Scalp app 0.05% .....	6.96	30 ml	Dermol

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>HYDROCORTISONE BUTYRATE</b>			
Scalp lotn 0.1% – 1% DV Mar-13 to 2015 .....	3.65	100 ml	<b>Locoid</b>

## Wart Preparations

IMIQUIMOD – **Restricted** see terms below

⚡ Crm 5%, 250 mg sachet – 1% DV Nov-11 to 2014 ..... 62.00 12 **Aldara**

➡ **Restricted**

Any of the following:

- 1 The patient has external anogenital warts and podophyllotoxin has been tried and failed (or is contraindicated); or
- 2 The patient has external anogenital warts and podophyllotoxin is unable to be applied accurately to the site; or
- 3 The patient has confirmed superficial basal cell carcinoma where other standard treatments, including surgical excision, are contraindicated or inappropriate.

Notes:

Superficial basal cell carcinoma

- Surgical excision remains first-line treatment for superficial basal cell carcinoma as it has a higher cure rate than imiquimod and allows histological assessment of tumour clearance.
- Imiquimod has not been evaluated for the treatment of superficial basal cell carcinoma within 1 cm of the hairline, eyes, nose, mouth or ears.
- Imiquimod is not indicated for recurrent, invasive, infiltrating, or nodular basal cell carcinoma.
- Every effort should be made to biopsy the lesion to confirm that it is a superficial basal cell carcinoma.

External anogenital warts

- Imiquimod is only indicated for external genital and perianal warts (condyloma acuminata).

PODOPHYLLOTOXIN

Soln 0.5% ..... 33.60 3.5 ml **Condyline**

SILVER NITRATE

Sticks with applicator

## Other Skin Preparations

DIPHENYL METILSULFATE

Powder 2%

SUNSCREEN, PROPRIETARY

Crm

Lotn .....	3.30	100 g	Marine Blue Lotion SPF 50+
	5.10	200 g	Marine Blue Lotion SPF 50+

## Antineoplastics

FLUOROURACIL SODIUM

Crm 5% – 1% DV Feb-13 to 2015 ..... 25.16 20 g **Efudix**

METHYL AMINOLEVULINATE HYDROCHLORIDE – **Restricted** see terms below

⚡ Crm 16%

➡ **Restricted**

Dermatologist or plastic surgeon

## Wound Management Products

CALCIUM GLUCONATE

Gel 2.5% ..... 21.00 1 healthE

Price (ex man. excl. GST)	Brand or Generic
\$	Per Manufacturer

## Anti-Infective Agents

### ACETIC ACID

- Soln 3%
- Soln 5%

### ACETIC ACID WITH HYDROXYQUINOLINE, GLYCEROL AND RICINOLEIC ACID

- Jelly 0.94% with hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator

### CHLORHEXIDINE

- Crm 1% – **1% DV Oct-12 to 2015** ..... 1.24 50 g **healthE**

### CHLORHEXIDINE GLUCONATE

- Lotn 1%, 200 ml ..... 6.75 1 **healthE**

### CLOTRIMAZOLE

- Vaginal crm 1% with applicator – **1% DV Dec-13 to 2016** ..... 1.45 35 g **Clomazol**
- Vaginal crm 2% with applicator – **1% DV Dec-13 to 2016** ..... 2.20 20 g **Clomazol**

### MICONAZOLE NITRATE

- Vaginal crm 2% with applicator

### NYSTATIN

- Vaginal crm 100,000 u per 5 g with applicator(s)

## Contraceptives

### Antiandrogen Oral Contraceptives

### CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

- Tab 2 mg with ethinyloestradiol 35 mcg

### Combined Oral Contraceptives

### ETHINYLOESTRADIOL WITH DESOGESTREL

- Tab 20 mcg with desogestrel 150 mcg
- Tab 30 mcg with desogestrel 150 mcg

### ETHINYLOESTRADIOL WITH LEVONORGESTREL

- Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets ..... 2.65 84 **Ava 20 ED**
- Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets ..... 2.30 84 **Ava 30 ED**
- Tab 20 mcg with levonorgestrel 100 mcg
- Tab 30 mcg with levonorgestrel 150 mcg
- Tab 50 mcg with levonorgestrel 125 mcg ..... 9.45 84 **Microgynon 50 ED**

### ETHINYLOESTRADIOL WITH NORETHISTERONE

- Tab 35 mcg with norethisterone 1 mg
- Tab 35 mcg with norethisterone 500 mcg

### NORETHISTERONE WITH MESTRANOL

- Tab 1 mg with mestranol 50 mcg

## Contraceptive Devices

### INTRA-UTERINE DEVICE

- IUD

*e.g. Multiload Cu375,  
Multiload Cu375 SL*

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Emergency Contraception</b>			
LEVONORGESTREL			
Tab 1.5 mg – 1% DV Jul-13 to 2016 .....	3.50	1	Postinor-1
<b>Progestogen-Only Contraceptives</b>			
LEVONORGESTREL			
Tab 30 mcg			
Implant 75 mg .....	133.65	1	Jadelle
⚡ Intra-uterine system, 20 mcg per day			e.g. Mirena
➡ <b>Restricted</b>			
Obstetrician or gynaecologist			
<b>Initiation – heavy menstrual bleeding</b>			
All of the following:			
1 The patient has a clinical diagnosis of heavy menstrual bleeding; and			
2 The patient has failed to respond to or is unable to tolerate other appropriate pharmaceutical therapies as per the Heavy Menstrual Bleeding Guidelines; and			
3 Any of the following:			
3.1 Serum ferritin level < 16 mcg/l (within the last 12 months); or			
3.2 Haemoglobin level < 120 g/l; or			
3.3 The patient has had a uterine ultrasound and either a hysteroscopy or endometrial biopsy.			
<b>Continuation – heavy menstrual bleeding</b>			
Either:			
1 Patient demonstrated clinical improvement of heavy menstrual bleeding; or			
2 Previous insertion was removed or expelled within 3 months of insertion.			
<b>Initiation – endometriosis</b>			
The patient has a clinical diagnosis of endometriosis confirmed by laparoscopy.			
<b>Continuation – endometriosis</b>			
Either:			
1 Patient demonstrated satisfactory management of endometriosis; or			
2 Previous insertion was removed or expelled within 3 months of insertion.			
Note: endometriosis is an unregistered indication.			
MEDROXYPROGESTERONE ACETATE			
Inj 150 mg per ml, 1 ml syringe – 1% DV Sep-13 to 2016 .....	7.00	1	Depo-Provera
NORETHISTERONE			
Tab 350 mcg			
<b>Obstetric Preparations</b>			
<b>Antiprogestogens</b>			
MIFEPRISTONE			
Tab 200 mg			
<b>Oxytocics</b>			
CARBOPROST TROMETAMOL			
Inj 250 mcg per ml, 1 ml ampoule			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>DINOPROSTONE</b>			
Pessaries 10 mg			
Gel 1 mg in 2.5 ml .....	52.65	1	Prostin E2
Gel 2 mg in 2.5 ml .....	64.60	1	Prostin E2
<b>ERGOMETRINE MALEATE</b>			
Inj 500 mcg per ml, 1 ml ampoule – 1% DV Nov-11 to 2014 .....	31.00	5	<b>DBL Ergometrine</b>
<b>OXYTOCIN</b>			
Inj 5 iu per ml, 1 ml ampoule – 1% DV Feb-14 to 2015 .....	4.75	5	<b>Oxytocin BNM</b>
Inj 10 iu per ml, 1 ml ampoule – 1% DV Feb-14 to 2015 .....	5.98	5	<b>BNM</b>
<b>OXYTOCIN WITH ERGOMETRINE MALEATE</b>			
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampoule – 1% DV Oct-12 to 2015 .....	11.13	5	<b>Syntometrine</b>

## Tocolytics

**PROGESTERONE – Restricted** see terms below

☞ Cap 100 mg ..... 16.50 30 Utrogestan

☞ **Restricted**

Obstetrician or gynaecologist

Both:

- 1 For the prevention of pre-term labour\*; and
- 2 Either:
  - 2.1 The patient has a short cervix on ultrasound (defined as < 25mm at 16 to 28 weeks) or
  - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with \* are Unapproved Indications (refer to Section A: General Rules, Part I (Interpretations and Definitions) and Part IV (Miscellaneous Provisions) rule 23.1).

**TERBUTALINE – Restricted** see terms below

☞ Inj 500 mcg ampoule

☞ **Restricted**

Obstetrician

## Oestrogens

**OESTRIOL**

Crm 1 mg per g with applicator

Pessaries 500 mcg

## Urologicals

### 5-Alpha Reductase Inhibitors

**FINASTERIDE – Restricted** see terms below

☞ Tab 5 mg – 1% DV Nov-11 to 2014 ..... 5.10 30 **Rex Medical**

☞ **Restricted**

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
  - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
  - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Alpha-1A Adrenoceptor Blockers</b>			
TAMSULOSIN – <b>Restricted</b> see terms below			
☞ Cap 400 mcg – 1% DV Dec-13 to 2016 .....	13.51	100	Tamsulosin-Rex
☞ <b>Restricted</b>			
Both:			
1 Patient has symptomatic benign prostatic hyperplasia; and			
2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.			
<b>Urinary Alkalisers</b>			
POTASSIUM CITRATE – <b>Restricted</b> see terms below			
☞ Oral liq 3 mmol per ml .....	30.00	200 ml	Biomed
☞ <b>Restricted</b>			
Both:			
1 The patient has recurrent calcium oxalate urolithiasis; and			
2 The patient has had more than two renal calculi in the two years prior to the application.			
SODIUM CITRO-TARTRATE			
Grans eff 4 g sachets .....	3.93	28	Ural
<b>Urinary Antispasmodics</b>			
OXYBUTYNIN			
Tab 5 mg – 1% DV Jun-13 to 2016 .....	11.20	500	Apo-Oxybutynin
Oral liq 5 mg per 5 ml – 1% DV Jun-13 to 2016 .....	56.45	473 ml	Apo-Oxybutynin
SOLIFENACIN SUCCINATE – <b>Restricted</b> see terms below			
☞ Tab 5 mg .....	56.50	30	Vesicare
☞ Tab 10 mg .....	56.50	30	Vesicare
☞ <b>Restricted</b>			
Patient has overactive bladder and a documented intolerance of, or is non-responsive to, oxybutynin.			
TOLTERODINE TARTRATE – <b>Restricted</b> see terms below			
☞ Tab 1 mg .....	14.56	56	Arrow-Tolterodine
☞ Tab 2 mg .....	14.56	56	Arrow-Tolterodine
☞ <b>Restricted</b>			
Patient has overactive bladder and a documented intolerance of, or is non-responsive to, oxybutynin.			



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Anabolic Agents</b>			
OXANDROLINE			
↓ Tab 2.5 mg			
→ <b>Restricted</b>			
For the treatment of burns patients.			
<b>Androgen Agonists and Antagonists</b>			
CYPROTERONE ACETATE			
Tab 50 mg – 1% DV Oct-12 to 2015 .....	18.80	50	Siterone
Tab 100 mg – 1% DV Oct-12 to 2015 .....	34.25	50	Siterone
TESTOSTERONE			
Patch 2.5 mg per day .....	80.00	60	Androderm
TESTOSTERONE CYPIONATE			
Inj 100 mg per ml, 10 ml vial – 1% DV Feb-12 to 2014 .....	76.50	1	Depo-Testosterone
TESTOSTERONE ESTERS			
Inj testosterone decanoate 100 mg, testosterone isocarproate 60 mg, testosterone phenylpropionate 60 mg and testosterone propionate 30 mg per ml, 1 ml ampoule			
TESTOSTERONE UNDECANOATE			
Cap 40 mg – 1% DV Oct-12 to 2015 .....	31.17	60	Andriol Testocaps
Inj 250 mg per ml, 4 ml ampoule .....	86.00	1	Reandron 1000
<b>Calcium Homeostasis</b>			
CALCITONIN			
Inj 100 iu per ml, 1 ml ampoule – 1% DV Sep-11 to 2014 .....	110.00	5	Miacalcic
ZOLEDRONIC ACID			
↓ Inj 0.8 mg per ml, 5 ml vial .....	550.00	1	Zometa
→ <b>Restricted</b>			
For hypercalcaemia of malignancy			
<b>Corticosteroids</b>			
BETAMETHASONE			
Tab 500 mcg			
Inj 4 mg per ml, 1 ml ampoule			
BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE			
Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml ampoule			
DEXAMETHASONE			
Tab 1 mg – 1% DV Aug-12 to 2015 .....	5.87	100	Douglas
Tab 4 mg – 1% DV Aug-12 to 2015 .....	8.16	100	Douglas
Oral liq 1 mg per ml .....	45.00	25 ml	Biomed
DEXAMETHASONE PHOSPHATE			
Inj 4 mg per ml, 1 ml ampoule – 1% DV Apr-14 to 2016 .....	25.80	10	Dexamethasone-hameln
Inj 4 mg per ml, 2 ml ampoule – 1% DV Apr-14 to 2016 .....	17.98	5	Dexamethasone-hameln

## HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>FLUDROCORTISONE ACETATE</b>			
Tab 100 mcg .....	14.32	100	Florinef
<b>HYDROCORTISONE</b>			
Tab 5 mg – 1% DV Nov-12 to 2015 .....	8.10	100	<b>Douglas</b>
Tab 20 mg – 1% DV Nov-12 to 2015 .....	20.32	100	<b>Douglas</b>
Inj 100 mg vial – 1% DV Oct-13 to 2016 .....	4.99	1	<b>Solu-Cortef</b>
<b>METHYLPREDNISOLONE (AS SODIUM SUCCINATE)</b>			
Tab 4 mg – 1% DV Oct-12 to 2015 .....	60.00	100	<b>Medrol</b>
Tab 100 mg – 1% DV Oct-12 to 2015 .....	166.52	20	<b>Medrol</b>
Inj 40 mg vial – 1% DV Oct-12 to 2015 .....	7.50	1	<b>Solu-Medrol</b>
Inj 125 mg vial – 1% DV Oct-12 to 2015 .....	18.50	1	<b>Solu-Medrol</b>
Inj 500 mg vial – 1% DV Oct-12 to 2015 .....	18.00	1	<b>Solu-Medrol</b>
Inj 1 g vial – 1% DV Oct-12 to 2015 .....	37.50	1	<b>Solu-Medrol</b>
<b>METHYLPREDNISOLONE ACETATE</b>			
Inj 40 mg per ml, 1 ml vial – 1% DV Oct-12 to 2015 .....	6.70	1	<b>Depo-Medrol</b>
<b>METHYLPREDNISOLONE ACETATE WITH LIGNOCAINE</b>			
Inj 40 mg with lignocaine 10 mg per ml, 1 ml vial – 1% DV Oct-12 to 2015 .....	7.50	1	<b>Depo-Medrol with Lidocaine</b>
<b>PREDNISOLONE</b>			
Oral liq 5 mg per ml .....	10.45	30 ml	Redipred
Enema 200 mcg per ml, 100 ml			
<b>PREDNISONE</b>			
Tab 1 mg .....	2.13	100	Apo-Prednisone S29
	10.68	500	Apo-Prednisone
Tab 2.5 mg .....	12.09	500	Apo-Prednisone
Tab 5 mg .....	11.09	500	Apo-Prednisone
Tab 20 mg .....	29.03	500	Apo-Prednisone
<b>TRIAMCINOLONE ACETONIDE</b>			
Inj 10 mg per ml, 1 ml ampoule – 1% DV Jun-12 to 2014 .....	21.90	5	<b>Kenacort-A</b>
Inj 40 mg per ml, 1 ml ampoule – 1% DV Jun-12 to 2014 .....	53.79	5	<b>Kenacort-A40</b>
<b>TRIAMCINOLONE HEXACETONIDE</b>			
Inj 20 mg per ml, 1 ml vial			

## Hormone Replacement Therapy

### Oestrogens

#### OESTRADIOL

- Tab 1 mg
- Tab 2 mg
- Patch 25 mcg per day
- Patch 50 mcg per day
- Patch 100 mcg per day

#### OESTRADIOL VALERATE

- Tab 1 mg
- Tab 2 mg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
OESTROGENS (CONJUGATED EQUINE)			
Tab 300 mcg			
Tab 625 mcg			

### Progestogen and Oestrogen Combined Preparations

OESTRADIOL WITH NORETHISTERONE ACETATE			
Tab 1 mg with 0.5 mg norethisterone acetate			
Tab 2 mg with 1 mg norethisterone acetate			
Tab 2 mg with 1 mg norethisterone acetate (10), and tab 2 mg oestradiol (12) and tab 1 mg oestradiol (6)			
OESTROGENS WITH MEDROXYPROGESTERONE ACETATE			
Tab 625 mcg conjugated equine with 2.5 mg medroxyprogesterone acetate			
Tab 625 mcg conjugated equine with 5 mg medroxyprogesterone acetate			

### Progestogens

MEDROXYPROGESTERONE ACETATE			
Tab 2.5 mg – 1% DV Sep-13 to 2016 .....	3.09	30	<b>Provera</b>
Tab 5 mg – 1% DV Sep-13 to 2016 .....	13.06	100	<b>Provera</b>
Tab 10 mg – 1% DV Sep-13 to 2016 .....	6.85	30	<b>Provera</b>

### Other Endocrine Agents

CABERGOLINE – <b>Restricted</b> see terms below			
⚡ Tab 0.5 mg – 1% DV Sep-12 to 2015 .....	6.25	2	<b>Dostinex</b>
	25.00	8	<b>Dostinex</b>

#### ➡ **Restricted**

Any of the following:

- 1 Inhibition of lactation; or
- 2 Patient has pathological hyperprolactinemia; or
- 3 Patient has acromegaly.

CLOMIPHENE CITRATE			
Tab 50 mg – 1% DV Sep-13 to 2016 .....	29.84	10	<b>Serophene</b>
DANAZOL			
Cap 100 mg .....	68.33	100	<b>Azol</b>
Cap 200 mg .....	97.83	100	<b>Azol</b>

GESTRINONE  
Cap 2.5 mg

METYRAPONE  
Cap 250 mg

PENTAGASTRIN  
Inj 250 mcg per ml, 2 ml ampoule

### Other Oestrogen Preparations

ETHINYLOESTRADIOL  
Tab 10 mcg

## HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
OESTRADIOL Implant 50 mg			
OESTRIOL Tab 2 mg			

### Other Progestogen Preparations

MEDROXYPROGESTERONE			
Tab 100 mg – 1% DV Sep-13 to 2016 .....	96.50	100	<b>Provera</b>
Tab 200 mg .....	70.50	30	Provera
NORETHISTERONE			
Tab 5 mg – 1% DV Nov-11 to 2014 .....	26.50	100	<b>Primolut N</b>

### Pituitary and Hypothalamic Hormones and Analogues

CORTICOTRORELIN (OVINE) Inj 100 mcg vial
THYROTROPIN ALFA Inj 900 mcg vial

### Adrenocorticotrophic Hormones

TETRACOSACTIDE [TETRACOSACTRIN]			
Inj 250 mcg per ml, 1 ml ampoule – 1% DV Sep-11 to 2014 .....	177.18	10	<b>Synacthen</b>
Inj 1 mg per ml, 1 ml ampoule – 1% DV Sep-11 to 2014 .....	29.56	1	<b>Synacthen Depot</b>

### GnRH Agonists and Antagonists

BUSERELIN Inj 1 mg per ml, 5.5 ml vial			
GONADORELIN Inj 100 mcg vial			
GOSERELIN			
Implant 3.6 mg .....	166.20	1	Zoladex
Implant 10.8 mg .....	443.76	1	Zoladex
LEUPRORELIN ACETATE			
Inj 3.75 mg syringe .....	221.60	1	Lucrin Depot PDS
Inj 7.5 mg syringe .....	166.20	1	Eligard
Inj 11.25 mg syringe .....	591.68	1	Lucrin Depot PDS
Inj 22.5 mg syringe .....	443.76	1	Eligard
Inj 30 mg syringe .....	1,109.40	1	Lucrin Depot PDS
Inj 30 mg vial .....	591.68	1	Eligard
Inj 45 mg syringe .....	832.05	1	Eligard

### Gonadotrophins

CHORIOGONADOTROPIN ALFA Inj 250 mcg in 0.5 ml syringe
--

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

## Growth Hormone

SOMATROPIN – **Restricted** see terms below

⚡ Inj 16 iu (5.3 mg) vial

⚡ Inj 36 iu (12 mg) vial

### →Restricted

Only for use in patients with approval by the New Zealand Growth Hormone Committee or the Adult Growth Hormone Panel

## Thyroid and Antithyroid Preparations

CARBIMAZOLE

Tab 5 mg

IODINE

Soln BP 50 mg per ml

LEVOTHYROXINE

Tab 25 mcg

Tab 50 mcg

Tab 100 mcg

LIOTHYRONINE SODIUM

⚡ Tab 20 mcg

### →Restricted

For a maximum of 14 days' treatment in patients with thyroid cancer who are due to receive radioiodine therapy

Inj 20 mcg vial

POTASSIUM IODATE

Tab 170 mg

POTASSIUM PERCHLORATE

Cap 200 mg

PROPYLTHIOURACIL – **Restricted** see terms below

⚡ Tab 50 mg .....	35.00	100	PTU
-------------------	-------	-----	-----

### →Restricted

Both:

1 The patient has hyperthyroidism; and

2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Note: Propylthiouracil is not recommended for patients under the age of 18 years unless the patient is pregnant and other treatments are contraindicated.

PROTIRELIN

Inj 100 mcg per ml, 2 ml ampoule

## Vasopressin Agents

ARGIPRESSIN [VASOPRESSIN]

Inj 20 u per ml, 1 ml ampoule

DESMOPRESSIN ACETATE – **Some items restricted** see terms on the next page

⚡ Tab 100 mcg .....	36.40	30	Minirin
⚡ Tab 200 mcg .....	93.60	30	Minirin
Nasal spray 10 mcg per dose – 1% DV Sep-11 to 2014.....	27.48	6 ml	Desmopressin-PH&T
Inj 4 mcg per ml, 1 ml ampoule			
Inj 15 mcg per ml, 1 ml ampoule			
Nasal drops 100 mcg per ml			

## HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➡Restricted</b>			
<b>Nocturnal enuresis</b>			
Either:			
1 The nasal forms of desmopressin are contraindicated; or			
2 An enuresis alarm is contraindicated.			
Cranial diabetes insipidus and the nasal forms of desmopressin are contraindicated			
<b>TERLIPRESSIN</b>			
Inj 0.1 mg per ml, 8.5 ml ampoule .....	450.00	5	Glypressin
Inj 1 mg vial .....	450.00	5	Glypressin
<i>(Glypressin Inj 1 mg vial to be delisted 1 December 2014)</i>			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Antibacterials</b>			
<b>Aminoglycosides</b>			
AMIKACIN – <b>Restricted</b> see terms below			
⌚ Inj 5 mg per ml, 10 ml syringe			
⌚ Inj 5 mg per ml, 5 ml syringe – <b>1% DV Nov-12 to 2014</b> .....	176.00	10	<b>Biomed</b>
⌚ Inj 15 mg per ml, 5 ml syringe			
⌚ Inj 250 mg per ml, 2 ml vial			
➔ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist or respiratory physician			
GENTAMICIN SULPHATE			
Inj 10 mg per ml, 1 ml ampoule .....	8.56	5	Hospira
Inj 10 mg per ml, 2 ml ampoule .....	175.10	25	APP Pharmaceuticals
Inj 40 mg per ml, 2 ml ampoule – <b>1% DV Sep-12 to 2015</b> .....	6.50	10	<b>Pfizer</b>
PAROMOMYCIN – <b>Restricted</b> see terms below			
⌚ Cap 250 mg .....	126.00	16	Humatin
➔ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
STREPTOMYCIN SULPHATE – <b>Restricted</b> see terms below			
⌚ Inj 400 mg per ml, 2.5 ml ampoule			
➔ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist or respiratory physician			
TOBRAMYCIN – <b>Restricted</b> see terms below			
⌚ Inj 40 mg per ml, 2 ml vial – <b>1% DV Sep-11 to 2014</b> .....	29.32	5	<b>DBL Tobramycin</b>
⌚ Inj 100 mg per ml, 5 ml vial			
➔ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist or respiratory physician			
<b>Carbapenems</b>			
ERTAPENEM – <b>Restricted</b> see terms below			
⌚ Inj 1 g vial .....	70.00	1	Invanz
➔ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
IMIPENEM WITH CILASTATIN – <b>Restricted</b> see terms below			
⌚ Inj 500 mg with 500 mg cilastatin vial – <b>1% DV Dec-12 to 2014</b> .....	18.37	1	<b>Primaxin</b>
➔ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
MEROPENEM – <b>Restricted</b> see terms below			
⌚ Inj 500 mg vial – <b>1% DV Mar-12 to 2014</b> .....	10.50	1	<b>Penembact</b>
⌚ Inj 1 g vial – <b>1% DV Mar-12 to 2014</b> .....	21.00	1	<b>Penembact</b>
➔ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
<b>Cephalosporins and Cephamycins - 1st Generation</b>			
CEFALEXIN			
Cap 500 mg – <b>1% DV Oct-13 to 2016</b> .....	5.70	20	<b>Cephalexin ABM</b>
Grans for oral liq 25 mg per ml – <b>1% DV Oct-13 to 2016</b> .....	8.50	100 ml	<b>Cefalexin Sandoz</b>
Grans for oral liq 50 mg per ml – <b>1% DV Oct-13 to 2016</b> .....	11.50	100 ml	<b>Cefalexin Sandoz</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>CEFAZOLIN</b>			
Inj 500 mg vial – 1% DV Mar-12 to 2014 .....	3.99	5	<b>AFT</b>
Inj 1 g vial – 1% DV Mar-12 to 2014 .....	3.99	5	<b>AFT</b>

### Cephalosporins and Cephamycins - 2nd Generation

<b>CEFACLOR</b>			
Cap 250 mg – 1% DV Dec-13 to 2016 .....	26.00	100	<b>Ranbaxy-Cefaclor</b>
Grans for oral liq 25 mg per ml – 1% DV Dec-13 to 2016 .....	3.53	100 ml	<b>Ranbaxy-Cefaclor</b>
<b>CEFOXITIN</b>			
Inj 1 g vial .....	55.00	5	<b>Hospira</b>
<b>CEFUROXIME</b>			
Tab 250 mg .....	29.40	50	<b>Zinnat</b>
Inj 750 mg vial – 1% DV Mar-12 to 2014 .....	6.96	5	<b>m-Cefuroxime</b>
Inj 1.5 g vial – 1% DV Mar-12 to 2014 .....	2.65	1	<b>Mylan</b>

### Cephalosporins and Cephamycins - 3rd Generation

<b>CEFOTAXIME</b>			
Inj 500 mg vial – 1% DV Oct-11 to 2014 .....	1.90	1	<b>Cefotaxime Sandoz</b>
Inj 1 g vial – 1% DV Nov-11 to 2014 .....	15.58	10	<b>DBL Cefotaxime</b>
<b>CEFTAZADIME – Restricted</b> see terms below			
⚡ Inj 500 mg vial – 1% DV Oct-11 to 2014 .....	2.37	1	<b>Fortum</b>
⚡ Inj 1 g vial .....	3.25	1	<b>DBL Ceftazidime</b>
⚡ Inj 2 g vial .....	6.49	1	<b>DBL Ceftazidime</b>
➡ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist or respiratory physician			
<b>CEFTRIAXONE</b>			
Inj 500 mg vial – 1% DV Mar-14 to 2016 .....	1.50	1	<b>Ceftriaxone-AFT</b>
Inj 1 g vial – 1% DV Mar-14 to 2016 .....	5.22	5	<b>Ceftriaxone-AFT</b>
Inj 2 g vial – 1% DV Mar-14 to 2016 .....	2.75	1	<b>Ceftriaxone-AFT</b>

### Cephalosporins and Cephamycins - 4th Generation

<b>CEFEPIME – Restricted</b> see terms below			
⚡ Inj 1 g vial .....	8.80	1	<b>DBL Cefepime</b>
⚡ Inj 2 g vial .....	17.60	1	<b>DBL Cefepime</b>
➡ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			

### Macrolides

<b>AZITHROMYCIN – Restricted</b> see terms below			
⚡ Tab 250 mg .....	10.00	30	<b>Apo-Azithromycin</b>
⚡ Tab 500 mg – 1% DV Feb-13 to 2015 .....	1.25	2	<b>Apo-Azithromycin</b>
⚡ Oral liq 40 mg per ml .....	6.60	15 ml	<b>Zithromax</b>

#### ➡**Restricted**

Any of the following:

- 1 Patient has received a lung transplant and requires treatment or prophylaxis for bronchiolitis obliterans syndrome; or
- 2 Patient has cystic fibrosis and has chronic infection with *Pseudomonas aeruginosa* or *Pseudomonas* related gram negative organisms; or
- 3 For any other condition for five days' treatment, with review after five days.



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>CLARITHROMYCIN – Restricted</b> see terms below			
⚡ Tab 250 mg – <b>1% DV Jan-12 to 2014</b> .....	4.19	14	<b>Apo-Clarithromycin</b>
⚡ Tab 500 mg – <b>1% DV Apr-12 to 2014</b> .....	10.95	14	<b>Apo-Clarithromycin</b>
⚡ Grans for oral liq 25 mg per ml .....	23.12	70 ml	Klacid
⚡ Inj 500 mg vial – <b>1% DV Oct-11 to 2014</b> .....	30.00	1	<b>Klacid</b>
<b>➡ Restricted</b>			
<b>Tab 250 mg and oral liquid</b>			
Tab 250 mg and oral liquid			
1 Atypical mycobacterial infection; or			
2 Mycobacterium tuberculosis infection where there is drug resistance or intolerance to standard pharmaceutical agents.			
<b>Tab 500 mg</b>			
Helicobacter pylori eradication.			
<b>Infusion</b>			
Infusion			
1 Atypical mycobacterial infection; or			
2 Mycobacterium tuberculosis infection where there is drug resistance or intolerance to standard pharmaceutical agents; or			
3 Community-acquired pneumonia (clarithromycin is not to be used as the first-line macrolide).			
<b>ERYTHROMYCIN (AS ETHYLSUCCINATE)</b>			
Tab 400 mg .....	16.95	100	E-Mycin
Grans for oral liq 200 mg per 5 ml .....	4.35	100 ml	E-Mycin
Grans for oral liq 400 mg per 5 ml .....	5.85	100 ml	E-Mycin
<b>ERYTHROMYCIN (AS LACTOBIONATE)</b>			
Inj 1 g vial .....	16.00	1	Erythrocin IV
<b>ERYTHROMYCIN (AS STEARATE) – Restricted:</b> For continuation only			
➡ Tab 250 mg			
➡ Tab 500 mg			
<b>ROXITHROMYCIN</b>			
Tab 150 mg – <b>1% DV Sep-12 to 2015</b> .....	7.48	50	<b>Arrow-Roxithromycin</b>
Tab 300 mg – <b>1% DV Sep-12 to 2015</b> .....	14.40	50	<b>Arrow-Roxithromycin</b>
<b>Penicillins</b>			
<b>AMOXYCILLIN</b>			
Cap 250 mg – <b>1% DV Mar-14 to 2016</b> .....	16.18	500	<b>Apo-Amoxi</b>
Cap 500 mg – <b>1% DV Jul-14 to 2016</b> .....	20.94	500	<b>Apo-Amoxi</b>
	26.50		Alphamox
Grans for oral liq 25 mg per ml .....	1.55	100 ml	Ospamox
Grans for oral liq 50 mg per ml .....	1.10	100 ml	Ospamox
Inj 250 mg vial – <b>1% DV Nov-11 to 2014</b> .....	12.96	10	<b>Ibiamox</b>
Inj 500 mg vial – <b>1% DV Nov-11 to 2014</b> .....	15.08	10	<b>Ibiamox</b>
Inj 1 g vial – <b>1% DV Nov-11 to 2014</b> .....	21.94	10	<b>Ibiamox</b>
<i>(Alphamox Cap 500 mg to be delisted 1 July 2014)</i>			
<b>AMOXYCILLIN WITH CLAVULANIC ACID</b>			
Tab 500 mg with clavulanic acid 125 mg – <b>1% DV Aug-12 to 2014</b> .....	12.55	100	<b>Curam Duo</b>
Grans for oral liq 25 mg with clavulanic acid 6.25 mg per ml – <b>1% DV Nov-12 to 2015</b> .....	1.61	100 ml	<b>Augmentin</b>
Grans for oral liq 50 mg with clavulanic acid 12.5 mg per ml – <b>1% DV Nov-12 to 2015</b> .....	2.19	100 ml	<b>Augmentin</b>
Inj 500 mg with clavulanic acid 100 mg vial – <b>1% DV Jan-13 to 2015</b> .....	10.14	10	<b>m-Amoxiclav</b>
Inj 1,000 mg with clavulanic acid 200 mg vial – <b>1% DV Jan-13 to 2015</b> .....	14.03	10	<b>m-Amoxiclav</b>

Products with Hospital Supply Status (HSS) are in **bold**

Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>BENZATHINE BENZYL PENICILLIN</b>			
Inj 900 mg (1.2 million units) in 2.3 ml syringe – 1% DV Sep-12 to 2015 .....	315.00	10	<b>Bicillin LA</b>
<b>BENZYL PENICILLIN SODIUM [PENICILLIN G]</b>			
Inj 600 mg (1 million units) vial – 1% DV Nov-11 to 2014 .....	11.50	10	<b>Sandoz</b>
<b>FLUCLOXACILLIN</b>			
Cap 250 mg – 1% DV Oct-12 to 2015 .....	22.00	250	<b>Staphlex</b>
Cap 500 mg – 1% DV Oct-12 to 2015 .....	74.00	500	<b>Staphlex</b>
Grans for oral liq 25 mg per ml – 1% DV Sep-12 to 2015 .....	2.49	100 ml	<b>AFT</b>
Grans for oral liq 50 mg per ml – 1% DV Sep-12 to 2015 .....	3.25	100 ml	<b>AFT</b>
Inj 250 mg vial – 1% DV Nov-11 to 2014 .....	10.86	10	<b>Flucloxin</b>
Inj 500 mg vial – 1% DV Nov-11 to 2014 .....	11.32	10	<b>Flucloxin</b>
Inj 1 g vial – 1% DV Nov-11 to 2014 .....	14.28	10	<b>Flucloxin</b>
<b>PHENOXYMETHYL PENICILLIN [PENICILLIN V]</b>			
Cap 250 mg .....	11.99	50	<b>Cilicaine VK</b>
Cap 500 mg .....	14.45	50	<b>Cilicaine VK</b>
Grans for oral liq 125 mg per 5 ml – 1% DV Apr-14 to 2016 .....	1.64	100 ml	<b>AFT</b>
Grans for oral liq 250 mg per 5 ml – 1% DV Apr-14 to 2016 .....	1.74	100 ml	<b>AFT</b>
<b>PIPERACILLIN WITH TAZOBACTAM – Restricted</b> see terms below			
⚡ Inj 4 g with tazobactam 0.5 g vial – 1% DV Oct-13 to 2016 .....	5.84	1	<b>Tazocin EF</b>
➡ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist or respiratory physician			
<b>PROCAINE PENICILLIN</b>			
Inj 1.5 g in 3.4 ml syringe – 1% DV Nov-11 to 2014 .....	123.50	5	<b>Cilicaine</b>
<b>TICARCILLIN WITH CLAVULANIC ACID – Restricted</b> see terms below			
⚡ Inj 3 g with clavulanic acid 0.1 mg vial			
➡ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist or respiratory physician			
<b>Quinolones</b>			
<b>CIPROFLOXACIN – Restricted</b> see terms below			
⚡ Tab 250 mg – 1% DV Dec-11 to 2014 .....	2.20	28	<b>Cipflox</b>
⚡ Tab 500 mg – 1% DV Dec-11 to 2014 .....	3.00	28	<b>Cipflox</b>
⚡ Tab 750 mg – 1% DV Dec-11 to 2014 .....	5.15	28	<b>Cipflox</b>
⚡ Oral liq 50 mg per ml			
⚡ Oral liq 100 mg per ml			
⚡ Inj 2 mg per ml, 100 ml bag .....	41.00	10	<b>Aspen Ciprofloxacin</b>
➡ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
<b>MOXIFLOXACIN – Restricted</b> see terms on the next page			
⚡ Tab 400 mg .....	52.00	5	<b>Avelox</b>
⚡ Inj 1.6 mg per ml, 250 ml bag .....	70.00	1	<b>Avelox IV 400</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➔Restricted</b>			
<b>Mycobacterium infection</b>			
Infectious disease physician, clinical microbiologist or respiratory physician			
1 Active tuberculosis, with any of the following:			
1.1 Documented resistance to one or more first-line medications; or			
1.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or			
1.3 Impaired visual acuity (considered to preclude ethambutol use); or			
1.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or			
1.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications.			
2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated			
<b>Pneumonia</b>			
Infectious disease physician or clinical microbiologist			
1 Immunocompromised patient with pneumonia that is unresponsive to first-line treatment; or			
2 Pneumococcal pneumonia or other invasive pneumococcal disease highly resistant to other antibiotics.			
<b>Penetrating eye injury</b>			
Ophthalmologist			
Five days treatment for patients requiring prophylaxis following a penetrating eye injury			
<b>Mycoplasma genitalium</b>			
All of the following:			
1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium; and			
2 Has tried and failed to clear infection using azithromycin; and			
3 Treatment is only for 7 days.			
<b>NORFLOXACIN</b>			
Tab 400 mg – 1% DV Sep-11 to 2014 .....	15.45	100	<b>Arrow-Norfloxacin</b>
<b>Tetracyclines</b>			
<b>DEMECLOCYCLINE HYDROCHLORIDE</b>			
Cap 150 mg			
<b>DOXYCYCLINE</b>			
➔ Tab 50 mg – <b>Restricted:</b> For continuation only			
Tab 100 mg – 1% DV Sep-11 to 2014 .....	7.95	250	<b>Doxine</b>
Inj 5 mg per ml, 20 ml vial			
<b>MINOCYCLINE</b>			
Tab 50 mg			
➔ Cap 100 mg – <b>Restricted:</b> For continuation only			
<b>TETRACYCLINE</b>			
Tab 250 mg			
Cap 500 mg .....	46.00	30	<b>Tetracyclin Wolff</b>
<b>TIGECYCLINE</b> – <b>Restricted</b> see terms below			
⚡ Inj 50 mg vial			
<b>➔Restricted</b>			
Infectious disease physician or clinical microbiologist			
<b>Other Antibacterials</b>			
<b>AZTREONAM</b> – <b>Restricted</b> see terms below			
⚡ Inj 1 g vial – 1% DV Sep-11 to 2014 .....	131.00	5	<b>Azactam</b>
<b>➔Restricted</b>			
Infectious disease physician or clinical microbiologist			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>CHLORAMPHENICOL – Restricted</b> see terms below			
☞ Inj 1 g vial			
☞ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
<b>CLINDAMYCIN – Restricted</b> see terms below			
☞ Cap 150 mg – 1% DV Oct-13 to 2016 .....	5.80	16	<b>Clindamycin ABM</b>
☞ Oral liq 15 mg per ml			
☞ Inj 150 mg per ml, 4 ml ampoule – 1% DV Sep-13 to 2016 .....	100.00	10	<b>Dalacin C</b>
☞ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
<b>COLISTIN SULPHOMETHATE [COLESTIMETHATE] – Restricted</b> see terms below			
☞ Inj 150 mg per ml, 1 ml vial .....	65.00	1	<b>Colistin-Link</b>
☞ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist or respiratory physician			
<b>DAPTOMYCIN – Restricted</b> see terms below			
☞ Inj 350 mg vial			
☞ Inj 500 mg vial			
☞ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
<b>FOSFOMYCIN – Restricted</b> see terms below			
☞ Powder for oral solution, 3 g sachet			
☞ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
<b>FUSIDIC ACID – Restricted</b> see terms below			
☞ Tab 250 mg .....	34.50	12	<b>Fucidin</b>
☞ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
<b>HEXAMINE HIPPURATE</b>			
Tab 1 g			
<b>LINCOMYCIN – Restricted</b> see terms below			
☞ Inj 300 mg per ml, 2 ml vial			
☞ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
<b>LINEZOLID – Restricted</b> see terms below			
☞ Tab 600 mg			
☞ Oral liq 20 mg per ml			
☞ Inj 2 mg per ml, 300 ml bag			
☞ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
<b>NITROFURANTOIN</b>			
Tab 50 mg			
Tab 100 mg			
<b>PIVMECILLINAM – Restricted</b> see terms below			
☞ Tab 200 mg			
☞ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
<b>SULPHADIAZINE – Restricted</b> see terms on the next page			
☞ Tab 500 mg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➔Restricted</b>			
Infectious disease physician, clinical microbiologist or maternal-foetal medicine specialist			
TEICOPLANIN – <b>Restricted</b> see terms below			
☯ Inj 400 mg vial			
<b>➔Restricted</b>			
Infectious disease physician or clinical microbiologist			
TRIMETHOPRIM			
Tab 100 mg			
Tab 300 mg	9.28	50	TMP
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXAZOLE]			
Tab 80 mg with sulphamethoxazole 400 mg			
Oral liq 8 mg with sulphamethoxazole 40 mg per ml	2.15	100 ml	Deprim
Inj 16 mg with sulphamethoxazole 80 mg per ml, 5 ml ampoule			
VANCOMYCIN – <b>Restricted</b> see terms below			
☯ Inj 500 mg vial – 1% DV Sep-11 to 2014	3.58	1	Mylan

**➔Restricted**  
Infectious disease physician or clinical microbiologist

## Antifungals

### Imidazoles

KETOCONAZOLE

☯ Tab 200 mg

**➔Restricted**

Infectious disease physician, clinical microbiologist, dermatologist, endocrinologist or oncologist

### Polyene Antimycotics

AMPHOTERICIN B

☯ Inj (liposomal) 50 mg vial – 1% DV Oct-12 to 2015 3,450.00 10 **AmBisome**

**➔Restricted**

Infectious disease physician, clinical microbiologist, haematologist, oncologist, transplant specialist or respiratory physician

Either:

- 1 Proven or probable invasive fungal infection, to be prescribed under an established protocol; or
- 2 Both:
  - 2.1 Possible invasive fungal infection; and
  - 2.2 A multidisciplinary team (including an infectious disease physician or a clinical microbiologist) considers the treatment to be appropriate.

☯ Inj 50 mg vial

**➔Restricted**

Infectious disease physician, clinical microbiologist, haematologist, oncologist, transplant specialist or respiratory physician

NYSTATIN

Tab 500,000 u	17.09	50	Nilstat
Cap 500,000 u	15.47	50	Nilstat

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Triazoles</b>			
<b>FLUCONAZOLE – Restricted</b> see terms below			
☞ Cap 50 mg – 1% DV Jan-12 to 2014 .....	4.77	28	<b>Ozole</b>
☞ Cap 150 mg – 1% DV Jan-12 to 2014 .....	0.91	1	<b>Ozole</b>
☞ Cap 200 mg – 1% DV Jan-12 to 2014 .....	13.34	28	<b>Ozole</b>
☞ Oral liquid 50 mg per 5 ml .....	34.56	35 ml	Diflucan
☞ Inj 2 mg per ml, 50 ml vial – 1% DV Oct-13 to 2016 .....	4.95	1	<b>Fluconazole-Claris</b>
☞ Inj 2 mg per ml, 100 ml vial – 1% DV Oct-13 to 2016 .....	6.47	1	<b>Fluconazole-Claris</b>
☞ <b>Restricted</b>			
Consultant			
<b>ITRACONAZOLE – Restricted</b> see terms below			
☞ Cap 100 mg – 1% DV Oct-13 to 2016 .....	2.99	15	<b>Itrazole</b>
☞ Oral liquid 10 mg per ml .....			
☞ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist, clinical immunologist or dermatologist			
<b>POSACONAZOLE – Restricted</b> see terms below			
☞ Oral liq 40 mg per ml .....	761.13	105 ml	Noxafil
☞ <b>Restricted</b>			
Infectious disease physician or haematologist			
<b>Initiation</b>			
<i>Re-assessment required after 6 weeks</i>			
Both:			
1 Either:			
1.1 Patient has acute myeloid leukaemia; or			
1.2 Patient is planned to receive a stem cell transplant and is at high risk for aspergillus infection; and			
2 Patient is to be treated with high dose remission induction therapy or re-induction therapy			
<b>Continuation</b>			
<i>Re-assessment required after 6 weeks</i>			
Both:			
1 Patient has previously received posaconazole prophylaxis during remission induction therapy; and			
2 Any of the following:			
2.1 Patient is to be treated with high dose remission re-induction therapy; or			
2.2 Patient is to be treated with high dose consolidation therapy; or			
2.3 Patient is receiving a high risk stem cell transplant.			
<b>VORICONAZOLE – Restricted</b> see terms below			
☞ Tab 50 mg .....	730.00	56	Vfend
☞ Tab 200 mg .....	2,930.00	56	Vfend
☞ Oral liq 40 mg per ml .....	730.00	70 ml	Vfend
☞ Inj 200 mg vial .....	185.00	1	Vfend
☞ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist or haematologist			
<b>Proven or probable aspergillus infection</b>			
Both:			
1 Patient is immunocompromised; and			
2 Patient has proven or probable invasive aspergillus infection.			
<b>Possible aspergillus infection</b>			
All of the following:			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued. . .

- 1 Patient is immunocompromised; and
- 2 Patient has possible invasive aspergillus infection; and
- 3 A multidisciplinary team (including an infectious disease physician) considers the treatment to be appropriate.

#### Resistant candidiasis infections and other moulds

All of the following:

- 1 Patient is immunocompromised, and
- 2 Either:
  - 2.1 Patient has fluconazole resistant candidiasis; or
  - 2.2 Patient has mould strain such as *Fusarium* spp. and *Scedosporium* spp; and
- 3 A multidisciplinary team (including an infectious disease physician or clinical microbiologist) considers the treatment to be appropriate.

### Other Antifungals

CASPOFUNGIN – **Restricted** see terms below

⚡ Inj 50 mg vial – 1% DV Oct-12 to 2015	667.50	1	<b>Candidas</b>
⚡ Inj 70 mg vial – 1% DV Oct-12 to 2015	862.50	1	<b>Candidas</b>

#### ➡Restricted

Infectious disease physician, clinical microbiologist, haematologist, oncologist, transplant specialist or respiratory physician

Either:

- 1 Proven or probable invasive fungal infection, to be prescribed under an established protocol; or
- 2 Both:
  - 2.1 Possible invasive fungal infection; and
  - 2.2 A multidisciplinary team (including an infectious disease physician or a clinical microbiologist) considers the treatment to be appropriate.

FLUCYTOSINE – **Restricted** see terms below

⚡ Cap 500 mg

#### ➡Restricted

Infectious disease physician or clinical microbiologist.

TERBINAFINE

Tab 250 mg – 1% DV Nov-11 to 2014	1.78	14	<b>Dr Reddy's Terbinafine</b>
-----------------------------------	------	----	-------------------------------

### Antimycobacterials

#### Antileprotics

CLOFAZIMINE – **Restricted** see terms below

⚡ Cap 50 mg

#### ➡Restricted

Infectious disease physician, clinical microbiologist or dermatologist

DAPSONE – **Restricted** see terms below

⚡ Tab 25 mg

⚡ Tab 100 mg

#### ➡Restricted

Infectious disease physician, clinical microbiologist or dermatologist

### Antituberculotics

CYCLOSERINE – **Restricted** see terms below

⚡ Cap 250 mg

#### ➡Restricted

Infectious disease physician, clinical microbiologist or respiratory physician

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>ETHAMBUTOL HYDROCHLORIDE – Restricted</b> see terms below			
☞ Tab 100 mg .....	48.01	56	Myambutol
☞ Tab 400 mg .....	49.34	56	Myambutol
☞ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist or respiratory physician			
<b>ISONIAZID – Restricted</b> see terms below			
☞ Tab 100 mg – 1% DV Mar-13 to 2015 .....	20.00	100	PSM
☞ <b>Restricted</b>			
Internal medicine physician, paediatrician, clinical microbiologist, dermatologist or public health physician			
<b>ISONIAZID WITH RIFAMPICIN – Restricted</b> see terms below			
☞ Tab 100 mg with rifampicin 150 mg			
☞ Tab 150 mg with rifampicin 300 mg			
☞ <b>Restricted</b>			
Internal medicine physician, paediatrician, clinical microbiologist, dermatologist or public health physician			
<b>PARA-AMINOSALICYLIC ACID – Restricted</b> see terms below			
☞ Grans for oral liq 4 g .....	280.00	30	Paser
☞ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist or respiratory physician			
<b>PROTIONAMIDE – Restricted</b> see terms below			
☞ Tab 250 mg .....	305.00	100	Peteha
☞ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist or respiratory physician			
<b>PYRAZINAMIDE – Restricted</b> see terms below			
☞ Tab 500 mg			
☞ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist or respiratory physician			
<b>RIFABUTIN – Restricted</b> see terms below			
☞ Cap 150 mg – 1% DV Sep-13 to 2016 .....	213.19	30	Mycobutin
☞ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist, respiratory physician or gastroenterologist			
<b>RIFAMPICIN – Restricted</b> see terms below			
☞ Tab 600 mg			
☞ Cap 150 mg			
☞ Cap 300 mg			
☞ Oral liq 100 mg per 5 ml			
☞ Inj 600 mg vial			
☞ <b>Restricted</b>			
Internal medicine physician, clinical microbiologist, dermatologist, paediatrician or public health physician			

## Antiparasitics

### Anthelmintics

**ALBENDAZOLE – Restricted** see terms below

- ☞ Tab 200 mg
- ☞ Tab 400 mg

☞ **Restricted**

Infectious disease physician or clinical microbiologist



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>IVERMECTIN – Restricted</b> see terms below			
☞ Tab 3 mg .....	17.20	4	Stromectol
☛ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist or dermatologist.			
<b>MEBENDAZOLE</b>			
Tab 100 mg – <b>1% DV Nov-11 to 2014</b> .....	24.19	24	<b>De-Worm</b>
Oral liq 100 mg per 5 ml			
<b>PRAZIQUANTEL</b>			
Tab 600 mg			
<b>Antiprotzoals</b>			
<b>ARTEMETHER WITH LUMEFANTRINE – Restricted</b> see terms below			
☞ Tab 20 mg with lumefantrine 120 mg			
☛ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
<b>ARTESUNATE – Restricted</b> see terms below			
☞ Inj 60 mg vial			
☛ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
<b>ATOVAQUONE WITH PROGUANIL HYDROCHLORIDE – Restricted</b> see terms below			
☞ Tab 62.5 mg with proguanil hydrochloride 25 mg			
☞ Tab 250 mg with proguanil hydrochloride 100 mg			
☛ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
<b>CHLOROQUINE PHOSPHATE – Restricted</b> see terms below			
☞ Tab 250 mg			
☛ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist, dermatologist or rheumatologist			
<b>MEFLOQUINE HYDROCHLORIDE – Restricted</b> see terms below			
☞ Tab 250 mg			
☛ <b>Restricted</b>			
Infectious disease physician, clinical microbiologist, dermatologist or rheumatologist			
<b>METRONIDAZOLE</b>			
Tab 200 mg .....	10.45	100	Trichozole
Tab 400 mg .....	18.15	100	Trichozole
Oral liq benzoate 200 mg per 5 ml .....	25.00	100 ml	Flagyl-S
Inj 5 mg per ml, 100 ml bag .....	2.46	1	Baxter
	12.30	5	AFT
Suppos 500 mg .....	24.48	10	Flagyl
<b>NITAZOXANIDE – Restricted</b> see terms below			
☞ Tab 500 mg .....	1,680.00	30	Alinia
☞ Oral liq 100 mg per 5 ml			
☛ <b>Restricted</b>			
Infectious disease physician or clinical microbiologist			
<b>ORNIDAZOLE</b>			
Tab 500 mg .....	16.50	10	Arrow-Ornidazole
<b>PENTAMIDINE ISETHIONATE – Restricted</b> see terms on the next page			
☞ Inj 300 mg vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➡Restricted</b>			
Infectious disease physician or clinical microbiologist			
PRIMAQUINE PHOSPHATE – <b>Restricted</b> see terms below			
☞ Tab 7.5 mg			
<b>➡Restricted</b>			
Infectious disease physician or clinical microbiologist			
PYRIMETHAMINE – <b>Restricted</b> see terms below			
☞ Tab 25 mg			
<b>➡Restricted</b>			
Infectious disease physician, clinical microbiologist or maternal-foetal medicine specialist			
QUININE DIHYDROCHLORIDE – <b>Restricted</b> see terms below			
☞ Inj 60 mg per ml, 10 ml ampoule			
☞ Inj 300 mg per ml, 2 ml vial			
<b>➡Restricted</b>			
Infectious disease physician or clinical microbiologist			
QUININE SULPHATE			
Tab 300 mg .....	54.06	500	Q 300
SODIUM STIBOGLUCONATE – <b>Restricted</b> see terms below			
☞ Inj 100 mg per ml, 1 ml vial			
<b>➡Restricted</b>			
Infectious disease physician or clinical microbiologist			
SPIRAMYCIN – <b>Restricted</b> see terms below			
☞ Tab 500 mg			
<b>➡Restricted</b>			
Maternal-foetal medicine specialist			

## Antiretrovirals

### HIV Fusion Inhibitors

ENFUVIRTIDE – **Restricted** see terms below

☞ Inj 108 mg vial × 60 .....2,380.00 1 Fuzeon

**➡Restricted**

#### Initiation

*Re-assessment required after 12 months*

All of the following:

- 1 Confirmed HIV infection; and
- 2 Enfuvirtide to be given in combination with optimized background therapy (including at least 1 other antiretroviral drug that the patient has never previously been exposed to) for treatment failure; and
- 3 Either:
  - 3.1 Patient has evidence of HIV replication, despite ongoing therapy; or
  - 3.2 Patient has treatment-limiting toxicity to previous antiretroviral agents; and
- 4 Previous treatment with 3 different antiretroviral regimens has failed; and
- 5 All of the following:
  - 5.1 Previous treatment with a non-nucleoside reverse transcriptase inhibitor has failed; and
  - 5.2 Previous treatment with a nucleoside reverse transcriptase inhibitor has failed; and
  - 5.3 Previous treatment with a protease inhibitor has failed.

#### Continuation

Patient has had at least a 10-fold reduction in viral load at 12 months

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

## Non-Nucleoside Reverse Transcriptase Inhibitors

### ➔Restricted

#### Confirmed HIV

Both:

- 1 Confirmed HIV infection; and
- 2 Any of the following:
  - 2.1 Symptomatic patient; or
  - 2.2 Patient aged 12 months and under; or
  - 2.3 Both:
    - 2.3.1 Patient aged 1 to 5 years; and
    - 2.3.2 Any of the following:
      - 2.3.2.1 CD4 counts < 1000 cells/mm<sup>3</sup>; or
      - 2.3.2.2 CD4 counts < 0.25 × total lymphocyte count; or
      - 2.3.2.3 Viral load counts > 100000 copies per ml; or
  - 2.4 Both:
    - 2.4.1 Patient aged 6 years and over; and
    - 2.4.2 CD4 counts < 500 cells/mm<sup>3</sup>

#### Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

#### Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

#### Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

EFAVIRENZ – **Restricted** see terms above

⬆ Tab 50 mg .....	158.33	30	Stocrin
⬆ Tab 200 mg .....	474.99	90	Stocrin
⬆ Tab 600 mg .....	474.99	30	Stocrin
⬆ Oral liq 30 mg per ml			

ETRAVIRINE – **Restricted** see terms above

⬆ Tab 200 mg .....	770.00	60	Intelence
--------------------	--------	----	-----------

NEVIRAPINE – **Restricted** see terms above

⬆ Tab 200 mg – 1% DV Jan-13 to 2015 .....	95.94	60	<b>Nevirapine Alphapharm</b>
⬆ Oral suspension 10 mg per ml .....	134.55	240 ml	Viramune Suspension

## Nucleoside Reverse Transcriptase Inhibitors

### ➔Restricted

#### Confirmed HIV

Both:

- 1 Confirmed HIV infection; and
- 2 Any of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
2.1 Symptomatic patient; or			
2.2 Patient aged 12 months and under; or			
2.3 Both:			
2.3.1 Patient aged 1 to 5 years; and			
2.3.2 Any of the following:			
2.3.2.1 CD4 counts < 1000 cells/mm <sup>3</sup> ; or			
2.3.2.2 CD4 counts < 0.25 × total lymphocyte count; or			
2.3.2.3 Viral load counts > 100000 copies per ml; or			
2.4 Both:			
2.4.1 Patient aged 6 years and over; and			
2.4.2 CD4 counts < 500 cells/mm <sup>3</sup>			
<b>Prevention of maternal transmission</b>			
Either:			
1 Prevention of maternal foetal transmission; or			
2 Treatment of the newborn for up to eight weeks.			
<b>Post-exposure prophylaxis following non-occupational exposure to HIV</b>			
Both:			
1 Treatment course to be initiated within 72 hours post exposure; and			
2 Any of the following:			
2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or			
2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or			
2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.			
<b>Percutaneous exposure</b>			
Patient has percutaneous exposure to blood known to be HIV positive.			
ABACAVIR SULPHATE – <b>Restricted</b> see terms on the preceding page			
⌚ Tab 300 mg – 1% DV Jul-11 to 2014 .....	229.00	60	<b>Ziagen</b>
⌚ Oral liq 20 mg per ml – 1% DV Jul-11 to 2014 .....	50.00	240 ml	<b>Ziagen</b>
ABACAVIR SULPHATE WITH LAMIVUDINE – <b>Restricted</b> see terms on the preceding page			
⌚ Tab 600 mg with lamivudine 300 mg .....	630.00	30	Kivexa
DIDANOSINE [DDI] – <b>Restricted</b> see terms on the preceding page			
⌚ Cap 125 mg			
⌚ Cap 200 mg			
⌚ Cap 250 mg			
⌚ Cap 400 mg			
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPROXIL FUMARATE – <b>Restricted</b> see terms on the preceding page			
⌚ Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil fumarate 300 mg .....	1,313.19	30	Atripla
EMTRICITABINE – <b>Restricted</b> see terms on the preceding page			
⌚ Cap 200 mg .....	307.20	30	Emtriva
EMTRICITABINE WITH TENOFOVIR DISOPROXIL FUMARATE – <b>Restricted</b> see terms on the preceding page			
⌚ Tab 200 mg with tenofovir disoproxil fumarate 300 mg .....	838.20	30	Truvada
LAMIVUDINE – <b>Restricted</b> see terms on the preceding page			
⌚ Tab 150 mg			
⌚ Oral liq 10 mg per ml			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
STAVUDINE – <b>Restricted</b> see terms on page 75			
⬆ Cap 30 mg			
⬆ Cap 40 mg			
⬆ Powder for oral soln 1 mg per ml			
ZIDOVUDINE [AZT] – <b>Restricted</b> see terms on page 75			
⬆ Cap 100 mg – 1% DV Oct-13 to 2016	152.25	100	<b>Retrovir</b>
⬆ Oral liq 10 mg per ml – 1% DV Oct-13 to 2016	30.45	200 ml	<b>Retrovir</b>
⬆ Inj 10 mg per ml, 20 ml vial			
ZIDOVUDINE [AZT] WITH LAMIVUDINE – <b>Restricted</b> see terms on page 75			
⬆ Tab 300 mg with lamivudine 150 mg – 1% DV Dec-12 to 2014	63.50	60	<b>Alphapharm</b>

## Protease Inhibitors

### ➡ **Restricted**

#### Confirmed HIV

Both:

- 1 Confirmed HIV infection; and
- 2 Any of the following:
  - 2.1 Symptomatic patient; or
  - 2.2 Patient aged 12 months and under; or
  - 2.3 Both:
    - 2.3.1 Patient aged 1 to 5 years; and
    - 2.3.2 Any of the following:
      - 2.3.2.1 CD4 counts < 1000 cells/mm<sup>3</sup>; or
      - 2.3.2.2 CD4 counts < 0.25 × total lymphocyte count; or
      - 2.3.2.3 Viral load counts > 100000 copies per ml; or
  - 2.4 Both:
    - 2.4.1 Patient aged 6 years and over; and
    - 2.4.2 CD4 counts < 500 cells/mm<sup>3</sup>

#### Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

#### Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

#### Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

ATAZANAVIR SULPHATE – **Restricted** see terms above

⬆ Cap 150 mg	568.34	60	Reyataz
⬆ Cap 200 mg	757.79	60	Reyataz

DARUNAVIR – **Restricted** see terms above

⬆ Tab 400 mg	837.50	60	Prezista
⬆ Tab 600 mg	1,190.00	60	Prezista

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
INDINAVIR – <b>Restricted</b> see terms on the preceding page			
† Cap 200 mg			
† Cap 400 mg			
LOPINAVIR WITH RITONAVIR – <b>Restricted</b> see terms on the preceding page			
† Tab 100 mg with ritonavir 25 mg .....	183.75	60	Kaletra
† Tab 200 mg with ritonavir 50 mg .....	735.00	120	Kaletra
† Oral liq 80 mg with ritonavir 20 mg per ml .....	735.00	300 ml	Kaletra
RITONAVIR – <b>Restricted</b> see terms on the preceding page			
† Tab 100 mg – <b>1% DV Oct-12 to 2015</b> .....	43.31	30	Norvir
† Oral liq 80 mg per ml			

## Strand Transfer Inhibitors

### ➔Restricted

#### Confirmed HIV

Both:

- 1 Confirmed HIV infection; and
- 2 Any of the following:
  - 2.1 Symptomatic patient; or
  - 2.2 Patient aged 12 months and under; or
  - 2.3 Both:
    - 2.3.1 Patient aged 1 to 5 years; and
    - 2.3.2 Any of the following:
      - 2.3.2.1 CD4 counts < 1000 cells/mm<sup>3</sup>; or
      - 2.3.2.2 CD4 counts < 0.25 × total lymphocyte count; or
      - 2.3.2.3 Viral load counts > 100000 copies per ml; or
  - 2.4 Both:
    - 2.4.1 Patient aged 6 years and over; and
    - 2.4.2 CD4 counts < 500 cells/mm<sup>3</sup>

#### Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

#### Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

#### Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

RALTEGRAVIR POTASSIUM – **Restricted** see terms above

† Tab 400 mg .....	1,090.00	60	Isentress
--------------------	----------	----	-----------

## Antivirals

### Hepatitis B

ADEFOVIR DIPIVOXIL – **Restricted** see terms on the next page

† Tab 10 mg .....	670.00	30	Hepsera
-------------------	--------	----	---------

† Item restricted (see ➔ above); ‡ Item restricted (see ➔ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➡Restricted</b>			
Gastroenterologist or infectious disease physician			
All of the following:			
1 Patient has confirmed Hepatitis B infection (HBsAg+); and			
Documented resistance to lamivudine, defined as:			
1 Patient has raised serum ALT ( $> 1 \times \text{ULN}$ ); and			
2 Patient has HBV DNA greater than 100,000 copies per mL, or viral load $\geq 10$ -fold over nadir; and			
3 Detection of M204I or M204V mutation; and			
4 Either:			
4.1 Both:			
4.1.1 Patient is cirrhotic; and			
4.1.2 Adefovir dipivoxil to be used in combination with lamivudine; or			
4.2 Both:			
4.2.1 Patient is not cirrhotic; and			
4.2.2 Adefovir dipivoxil to be used as monotherapy.			
ENTECAVIR – <b>Restricted</b> see terms below			
⚡ Tab 0.5 mg .....	400.00	30	Baraclude
<b>➡Restricted</b>			
Gastroenterologist or infectious disease physician			
All of the following:			
1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and			
2 Patient is Hepatitis B nucleoside analogue treatment-naïve; and			
3 Entecavir dose 0.5 mg/day; and			
4 Either:			
4.1 ALT greater than upper limit of normal; or			
4.2 Bridging fibrosis or cirrhosis (Metavir stage 3 or greater or moderate fibrosis) on liver histology; and			
5 Either:			
5.1 HBeAg positive; or			
5.2 Patient has $\geq 2,000$ IU HBV DNA units per ml and fibrosis (Metavir stage 2 or greater) on liver histology; and			
6 No continuing alcohol abuse or intravenous drug use; and			
7 Not co-infected with HCV, HIV or HDV; and			
8 Neither ALT nor AST greater than 10 times upper limit of normal; and			
9 No history of hypersensitivity to entecavir; and			
10 No previous documented lamivudine resistance (either clinical or genotypic).			
LAMIVUDINE – <b>Restricted</b> see terms below			
⚡ Tab 100 mg – 1% DV Dec-12 to 2014 .....	32.50	28	Zetlam
⚡ Oral liq 5 mg per ml			
<b>➡Restricted</b>			
Gastroenterologist, infectious disease specialist, paediatrician or general physician			
<b>Initiation</b>			
<i>Re-assessment required after 12 months</i>			
Any of the following:			
1 HBV DNA positive cirrhosis prior to liver transplantation; or			
2 HBsAg positive and have had a liver, kidney, heart, lung or bone marrow transplant; or			
3 Hepatitis B virus naïve patient who has received a liver transplant from an anti-HBc (Hepatitis B core antibody) positive donor; or			
4 Hepatitis B surface antigen positive (HbsAg) patient who is receiving chemotherapy for a malignancy, or who has received such treatment within the previous two months; and			
5 Hepatitis B surface antigen positive patient who is receiving anti tumour necrosis factor treatment; or			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 6 Hepatitis B core antibody (anti-HBc) positive patient who is receiving rituximab plus high dose steroids (e.g. R-CHOP).

## **Continuation - patients who have maintained continuous treatment and response to lamivudine**

*Re-assessment required after 2 years*

All of the following:

- 1 Have maintained continuous treatment with lamivudine; and
- 2 Most recent test result shows continuing biochemical response (normal ALT); and
- 3 HBV DNA <100,00 copies per ml by quantitative PCR at a reference laboratory; or

## **Continuation - when given in combination with adefovir dipivoxil for patients with cirrhosis and resistance to lamivudine**

*Re-assessment required after 2 years*

All of the following:

- 1 Lamivudine to be used in combination with adefovir dipivoxil; and
- 2 Patient is cirrhotic; and

Documented resistance to lamivudine, defined as:

- 1 Patient has raised serum ALT ( $> 1 \times \text{ULN}$ ); and
- 2 Patient has HBV DNA greater than 100,000 copies per mL, or viral load  $\geq 10$ -fold over nadir; and
- 3 Detection of M204I or M204V mutation; or

## **Continuation - when given in combination with adefovir dipivoxil for patients with resistance to adefovir dipivoxil**

*Re-assessment required after 2 years*

All of the following:

- 1 Lamivudine to be used in combination with adefovir dipivoxil; and

Documented resistance to adefovir, defined as:

- 1 Patient has raised serum ALT ( $> 1 \times \text{ULN}$ ); and
- 2 Patient has HBV DNA greater than 100,000 copies per mL, or viral load  $\geq 10$ -fold over nadir; and
- 3 Detection of N236T or A181T/V mutation.

TENOFOVIR DISOPROXIL FUMARATE – **Restricted** see terms below

⚡ Tab 300 mg .....	531.00	30	Viread
--------------------	--------	----	--------

## **➡Restricted**

### **Confirmed hepatitis B**

Either:

- 1 All of the following:
  - 1.1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
  - 1.2 Patient has had previous lamivudine, adefovir or entecavir therapy; and
  - 1.3 HBV DNA greater than 20,000 IU/mL or increased  $\leq 10$ -fold over nadir; and
  - 1.4 Any of the following:
    - 1.4.1 Lamivudine resistance - detection of M204I/V mutation; or
    - 1.4.2 Adefovir resistance - detection of A181T/V or N236T mutation; or
    - 1.4.3 Entecavir resistance - detection of relevant mutations including I169T, L180M T184S/A/I/L/G/C/M, S202C/G/I, M204V or M250I/V mutation; or
- 2 Patient is either listed or has undergone liver transplantation for HBV; or
- 3 Patient has a decompensated cirrhosis with a Mayo score  $> 20$ .

### **Pregnant or Breastfeeding, Active hepatitis B**

Limited to twelve months' treatment

Both:

- 1 Patient is HBsAg positive and pregnant; and
- 2 HBV DNA  $> 20,000$  IU/mL and ALT  $> \text{ULN}$ .

### **Pregnant, prevention of vertical transmission**

Limited to six months' treatment

Both:

continued...



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient is HBsAg positive and pregnant; and
- 2 HBV DNA > 20 million IU/mL and ALT normal.

#### Confirmed HIV

Both:

- 1 Confirmed HIV infection; and
- 2 Any of the following:
  - 2.1 Symptomatic patient; or
  - 2.2 Patient aged 12 months and under; or
  - 2.3 Both:
    - 2.3.1 Patient aged 1 to 5 years; and
    - 2.3.2 Any of the following:
      - 2.3.2.1 CD4 counts < 1000 cells/mm<sup>3</sup>; or
      - 2.3.2.2 CD4 counts < 0.25 × total lymphocyte count; or
      - 2.3.2.3 Viral load counts > 100000 copies per ml; or
  - 2.4 Both:
    - 2.4.1 Patient aged 6 years and over; and
    - 2.4.2 CD4 counts < 500 cells/mm<sup>3</sup>

#### Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

#### Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

#### Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

### Hepatitis C

BOCEPREVIR – **Restricted** see terms below

☛ Cap 200 mg ..... 5,015.00 336 Victrelis

#### ☛ Restricted

#### Chronic hepatitis C - genotype 1, first-line from gastroenterologist, infectious disease physician or general physician

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has not received prior pegylated interferon treatment; and
- 3 Patient has IL-28B genotype CT or TT; and
- 4 Patient is to be treated in combination with pegylated interferon and ribavirin; and
- 5 Patient is hepatitis C protease inhibitor treatment-naïve; and
- 6 Maximum of 44 weeks therapy.

#### Chronic hepatitis C - genotype 1, second-line from gastroenterologist, infectious disease physician or general physician.

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has received pegylated interferon treatment; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

3 Any one of:

- 3.1 Patient was a responder relapser; or
- 3.2 Patient was a partial responder; or
- 3.3 Patient received pegylated interferon prior to 2004; and

4 Patient is to be treated in combination with pegylated interferon and ribavirin; and

5 Maximum of 44 weeks therapy.

Note: Due to risk of severe sepsis boceprevir should not be initiated if either Platelet count  $<100 \times 10^9 /l$  or Albumin  $<35 \text{ g/l}$ .

## Herpesviridae

### ACICLOVIR

Tab dispersible 200 mg – 1% DV Sep-13 to 2016 .....	1.78	25	Lovir
Tab dispersible 400 mg – 1% DV Sep-13 to 2016 .....	5.98	56	Lovir
Tab dispersible 800 mg – 1% DV Sep-13 to 2016 .....	6.64	35	Lovir
Inj 250 mg vial – 1% DV Mar-13 to 2015 .....	14.09	5	Zovirax IV

### CIDOFOVIR – **Restricted** see terms below

☞ Inj 75 mg per ml, 5 ml vial

#### ☞ **Restricted**

Infectious disease physician, clinical microbiologist, otolaryngologist or oral surgeon

### FOSCARNET SODIUM – **Restricted** see terms below

☞ Inj 24 mg per ml, 250 ml bottle

#### ☞ **Restricted**

Infectious disease physician or clinical microbiologist

### GANCICLOVIR – **Restricted** see terms below

☞ Inj 500 mg vial .....	380.00	5	Cymevene
-------------------------	--------	---	----------

#### ☞ **Restricted**

Infectious disease physician or clinical microbiologist

### VALACICLOVIR – **Restricted** see terms below

☞ Tab 500 mg .....	102.72	30	Valtrex
--------------------	--------	----	---------

#### ☞ **Restricted**

Any of the following:

- 1 Patient has genital herpes with 2 or more breakthrough episodes in any 6 month period while treated with aciclovir 400 mg twice daily.
- 2 Patient has previous history of ophthalmic zoster and the patient is at risk of vision impairment.
- 3 Patient has undergone organ transplantation.

### Immunocompromised patients

Limited to 7 days treatment

Both:

- 1 Patient is immunocompromised; and
- 2 Patient has herpes zoster.

### VALGANCICLOVIR – **Restricted** see terms on the next page

☞ Tab 450 mg .....	3,000.00	60	Valcyte
--------------------	----------	----	---------

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

➔ **Restricted**

**Transplant cytomegalovirus prophylaxis**

Limited to three months' treatment

Patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

**Lung transplant cytomegalovirus prophylaxis**

Limited to six months' treatment

Both:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
  - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
  - 2.2 The recipient is cytomegalovirus positive.

**Cytomegalovirus in immunocompromised patients**

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

**Influenza**

OSELTAMIVIR – **Restricted** see terms below

⚡ Tab 75 mg

⚡ Powder for oral suspension 6 mg per ml

➔ **Restricted**

Either:

- 1 Only for hospitalised patient with known or suspected influenza; or
- 2 For prophylaxis of influenza in hospitalised patients as part of a DHB hospital approved infections control plan.

ZANAMIVIR

⚡ Powder for inhalation 5 mg .....37.38 20 dose Relenza Rotadisk

➔ **Restricted**

Either:

- 1 Only for hospitalised patient with known or suspected influenza; or
- 2 For prophylaxis of influenza in hospitalised patients as part of a DHB hospital approved infections control plan.

**Immune Modulators**

INTERFERON ALFA-2A

- Inj 3 m iu prefilled syringe
- Inj 6 m iu prefilled syringe
- Inj 9 m iu prefilled syringe

INTERFERON ALFA-2B

- Inj 18 m iu, 1.2 ml multidose pen
- Inj 30 m iu, 1.2 ml multidose pen
- Inj 60 m iu, 1.2 ml multidose pen

INTERFERON GAMMA – **Restricted** see terms below

⚡ Inj 100 mcg in 0.5 ml vial

➔ **Restricted**

Patient has chronic granulomatous disease and requires interferon gamma.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PEGYLATED INTERFERON ALFA-2A – Restricted</b> see terms below			
☞ Inj 135 mcg prefilled syringe			
☞ Inj 135 mcg prefilled syringe (4) with ribavirin tab 200 mg (112)			
☞ Inj 135 mcg prefilled syringe (4) with ribavirin tab 200 mg (168)			
☞ Inj 180 mcg prefilled syringe .....	900.00	4	Pegasys
☞ Inj 180 mcg prefilled syringe (4) with ribavirin tab 200 mg (112) .....	1,159.84	1	Pegasus RBV Combination Pack
☞ Inj 180 mcg prefilled syringe (4) with ribavirin tab 200 mg (168) .....	1,290.00	1	Pegasus RBV Combination Pack

## ☞ Restricted

### Initiation – Chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant

Both:

- 1 Any of the following:
  - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
  - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
  - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant.
- 2 Maximum of 48 weeks therapy.

Notes:

Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.

Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml.

### Continuation – (Chronic hepatitis C - genotype 1 infection) - gastroenterologist, infectious disease physician or general physician

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

### Initiation (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) - Gastroenterologist, infectious disease physician or general physician

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; or
  - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

### Initiation – Chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

### Initiation – Hepatitis B

Gastroenterologist, infectious disease specialist or general physician

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naïve; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log<sub>10</sub> IU/ml; and
- 5 Either:
  - 5.1 HBeAg positive; or
  - 5.2 Serum HBV DNA ≥ 2,000 units/ml and significant fibrosis (≥ Metavir Stage F2 or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Notes:

Approved dose is 180 mcg once weekly.

The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.

In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon alfa-2a dose should be reduced to 135 mcg once weekly.

In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.

Pegylated Interferon alfa-2a is not approved for use in children.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Anticholinesterases</b>			
EDROPHONIUM CHLORIDE – <b>Restricted</b> see terms below			
☞ Inj 10 mg per ml, 15 ml vial			
☞ Inj 10 mg per ml, 1 ml ampoule			
☞ <b>Restricted</b>			
For the diagnosis of myasthenia gravis			
NEOSTIGMINE METILSULFATE			
Inj 2.5 mg per ml, 1 ml ampoule – <b>1% DV Sep-11 to 2014</b> .....	140.00	50	<b>AstraZeneca</b>
NEOSTIGMINE METILSULFATE WITH GLYCOPYRRONIUM BROMIDE			
Inj 2.5 mg with glycopyrronium bromide 0.5 mg per ml, 1 ml ampoule – <b>1% DV Nov-13 to 2016</b> .....	27.86	10	<b>Max Health</b>
PYRIDOSTIGMINE BROMIDE			
Tab 60 mg – <b>1% DV Sep-11 to 2014</b> .....	38.90	100	<b>Mestinon</b>
<b>Antirheumatoid Agents</b>			
AURANOFIN			
Tab 3 mg			
HYDROXYCHLOROQUINE			
Tab 200 mg – <b>1% DV Nov-12 to 2015</b> .....	18.00	100	<b>Plaquenil</b>
LEFLUNOMIDE			
Tab 10 mg .....	55.00	30	Arava
Tab 20 mg .....	76.00	30	Arava
Tab 100 mg .....	54.44	3	Arava
PENICILLAMINE			
Tab 125 mg .....	61.93	100	D-Penamine
Tab 250 mg .....	98.98	100	D-Penamine
SODIUM AUROTHIOMALATE			
Inj 10 mg in 0.5 ml ampoule			
Inj 20 mg in 0.5 ml ampoule			
Inj 50 mg in 0.5 ml ampoule			
<b>Drugs Affecting Bone Metabolism</b>			
<b>Bisphosphonates</b>			
ALENDRONATE SODIUM			
☞ Tab 40 mg .....	133.00	30	Fosamax
☞ <b>Restricted</b>			
Both:			
1 Paget's disease; and			
2 Any of the following:			
2.1 Bone or articular pain; or			
2.2 Bone deformity; or			
2.3 Bone, articular or neurological complications; or			
2.4 Asymptomatic disease, but risk of complications due to site (base of skull, spine, long bones of lower limbs); or			
2.5 Preparation for orthopaedic surgery.			
☞ Tab 70 mg .....	22.90	4	Fosamax

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➡ **Restricted**

**Osteoporosis**

Any of the following:

- History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD)  $\geq 2.5$  standard deviations below the mean normal value in young adults (i.e. T-Score  $\leq -2.5$ ) (see Note); or
- History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- History of two significant osteoporotic fractures demonstrated radiologically; or
- Documented T-Score  $\leq -3.0$  (see Note); or
- A 10-year risk of hip fracture  $\geq 3\%$ , calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- Patient has had a Special Authority approval for zoledronic acid (osteoporosis) or raloxifene.

**Initiation - glucocorticosteroid therapy**

*Re-assessment required after 12 months*

Both:

- The patient is receiving systemic glucocorticosteroid therapy ( $\geq 5$  mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- Any of the following:
  - The patient has documented BMD  $\geq 1.5$  standard deviations below the mean normal value in young adults (i.e. T-Score  $\leq -1.5$ ) (see Note); or
  - The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
  - The patient has had a Special Authority approval for zoledronic acid (glucocorticosteroid therapy) or raloxifene.

**Continuation - glucocorticosteroid therapy**

*Re-assessment required after 12 months*

The patient is continuing systemic glucocorticosteroid therapy ( $\geq 5$  mg per day prednisone equivalents)

Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score  $\leq -2.5$  and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

ALENDRONATE SODIUM WITH CHOLECALCIFEROL – **Restricted** see terms below

⚡ Tab 70 mg with cholecalciferol 5,600 iu .....22.90      4      Fosamax Plus

➡ **Restricted**

**Osteoporosis**

Any of the following:

- History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD)  $\geq 2.5$  standard deviations below the mean normal value in young adults (i.e. T-Score  $\leq -2.5$ ) (see Note); or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
<ol style="list-style-type: none"> <li>History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or</li> <li>History of two significant osteoporotic fractures demonstrated radiologically; or</li> <li>Documented T-Score <math>\leq -3.0</math> (see Note); or</li> <li>A 10-year risk of hip fracture <math>\geq 3\%</math>, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or</li> <li>Patient has had a Special Authority approval for zoledronic acid (osteoporosis) or raloxifene.</li> </ol>			
<b>Initiation - glucocorticosteroid therapy</b>			
<i>Re-assessment required after 12 months</i>			
Both:			
<ol style="list-style-type: none"> <li>The patient is receiving systemic glucocorticosteroid therapy (<math>\geq 5</math> mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and</li> <li>Any of the following: <ol style="list-style-type: none"> <li>The patient has documented BMD <math>\geq 1.5</math> standard deviations below the mean normal value in young adults (i.e. T-Score <math>\leq -1.5</math>) (see Note); or</li> <li>The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or</li> <li>The patient has had a Special Authority approval for zoledronic acid (glucocorticosteroid therapy) or raloxifene.</li> </ol> </li> </ol>			
<b>Continuation - glucocorticosteroid therapy</b>			
<i>Re-assessment required after 12 months</i>			
The patient is continuing systemic glucocorticosteroid therapy ( $\geq 5$ mg per day prednisone equivalents)			
Notes:			
<ol style="list-style-type: none"> <li>BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.</li> <li>Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score <math>\geq -2.5</math> and, therefore, do not require BMD measurement for treatment with bisphosphonates.</li> <li>Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.</li> <li>A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.</li> </ol>			
<b>ETIDRONATE DISODIUM</b>			
Tab 200 mg – 1% DV Sep-12 to 2015 .....	15.80	100	Arrow-Etidronate
<b>PAMIDRONATE DISODIUM</b>			
Inj 3 mg per ml, 5 ml vial .....	18.75	1	Pamisol
Inj 3 mg per ml, 10 ml vial – 1% DV Feb-13 to 2014 .....	16.00	1	Pamidronate BNM
Inj 6 mg per ml, 10 ml vial – 1% DV Feb-13 to 2014 .....	32.00	1	Pamidronate BNM
Inj 9 mg per ml, 10 ml vial – 1% DV Feb-13 to 2014 .....	48.00	1	Pamidronate BNM
<b>ZOLEDRONIC ACID – Restricted</b> see terms on the next page			
☯ Inj 0.05 mg per ml, 100 ml vial .....	600.00	100 ml	Aclasta



Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

➔ **Restricted**

**Osteogenesis imperfecta**

Patient has been diagnosed with clinical or genetic osteogenesis imperfecta.

**Osteoporosis**

Both:

- 1 Any of the following:
  - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD)  $\geq$  2.5 standard deviations below the mean normal value in young adults (i.e. T-Score  $\leq$  -2.5) (see Note); or
  - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
  - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
  - 1.4 Documented T-Score  $\geq$  -3.0 (see Note); or
  - 1.5 A 10-year risk of hip fracture  $\geq$  3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
  - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) or raloxifene; and
- 2 The patient will not be prescribed more than one infusion in a 12-month period.

**Initiation - glucocorticosteroid therapy**

*Re-assessment required after 12 months*

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy ( $\geq$  5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
  - 2.1 The patient has documented BMD  $\geq$  1.5 standard deviations below the mean normal value in young adults (i.e. T-Score  $\leq$  -1.5) (see Note); or
  - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
  - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause - glucocorticosteroid therapy) or raloxifene; and
- 3 The patient will not be prescribed more than one infusion in the 12-month approval period.

**Continuation - glucocorticosteroid therapy**

*Re-assessment required after 12 months*

Both:

- 1 The patient is continuing systemic glucocorticosteroid therapy ( $\geq$  5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than one infusion in the 12-month approval period.

**Initiation - Paget's disease**

*Re-assessment required after 12 months*

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
  - 2.1 Bone or articular pain; or
  - 2.2 Bone deformity; or
  - 2.3 Bone, articular or neurological complications; or
  - 2.4 Asymptomatic disease, but risk of complications; or
  - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than one infusion in the 12-month approval period.

**Continuation - Paget's disease**

*Re-assessment required after 12 months*

Both:

- 1 Any of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
- 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
- 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than one infusion in the 12-month approval period.

Notes:

- 1 BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- 2 Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score  $\leq -2.5$  and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- 3 Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- 4 A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

## Other Drugs Affecting Bone Metabolism

**RALOXIFENE – Restricted** see terms below

☞ Tab 60 mg .....53.76      28      Evista

☞ **Restricted**

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD)  $\geq 2.5$  standard deviations below the mean normal value in young adults (i.e. T-Score  $\leq -2.5$ ) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score  $\geq -3.0$  (see Notes); or
- 5 A 10-year risk of hip fracture  $\geq 3\%$ , calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a prior Special Authority approval for zoledronic acid (Underlying cause - Osteoporosis) or alendronate (Underlying cause - Osteoporosis).

Notes:

- 1 BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- 2 Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score  $\leq -2.5$  and, therefore, do not require BMD measurement for raloxifene funding.
- 3 Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- 4 A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

**RISEDRONATE SODIUM**

Tab 35 mg ..... 4.00 4 Risedronate Sandoz

**TERIPARATIDE – Restricted** see terms below

📌 Inj 250 mcg per ml, 2.4 ml cartridge ..... 490.00 1 Forteo

➡ **Restricted**

Limited to 18 months' treatment

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- 1 The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- 2 Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- 3 A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

**Enzymes**

**HYALURONIDASE**

Inj 1,500 iu ampoule

**Hyperuricaemia and Antigout**

**ALLOPURINOL**

Tab 100 mg – 1% DV Dec-11 to 2014 ..... 15.90 1,000 **Apo-Allopurinol**

Tab 300 mg – 1% DV Dec-11 to 2014 ..... 16.75 500 **Apo-Allopurinol**

**BENZBROMARONE – Restricted** see terms below

📌 Tab 100 mg ..... 45.00 100 Benzbromaron AL 100

➡ **Restricted**

Both:

- 1 Any of the following:
  - 1.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and appropriate doses of probenecid; or
  - 1.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite appropriate doses of probenecid; or
  - 1.3 Both:
    - 1.3.1 The patient has renal impairment and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); and
    - 1.3.2 The patient has a rate of creatinine clearance greater than or equal to 20 ml/min; or
  - 1.4 All of the following:
    - 1.4.1 The patient is taking azathioprine and requires urate-lowering therapy; and
    - 1.4.2 Allopurinol is contraindicated; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.4.3 Appropriate doses of probenecid are ineffective or probenecid cannot be used due to reduced renal function; and

- 2 The patient is receiving monthly liver function tests.

Notes: Benzbromarone has been associated with potentially fatal hepatotoxicity. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

The New Zealand Rheumatology Association has developed information for prescribers which can be accessed from its website at [http://www.rheumatology.org.nz/benzbromarone\\_prescriber\\_information.cfm](http://www.rheumatology.org.nz/benzbromarone_prescriber_information.cfm)

## COLCHICINE

Tab 500 mcg – **1% DV Oct-13 to 2016** ..... 10.08 100 **Colgout**

## FEBUXOSTAT – **Restricted** see terms below

⚡ Tab 80 mg ..... 39.50 28 Adenuric  
⚡ Tab 120 mg ..... 39.50 28 Adenuric

## ➡**Restricted**

Any of the following:

- The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and appropriate doses of probenecid; or
- The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite appropriate doses of probenecid; or
- Both:
  - The patient has renal impairment and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); and
  - The patient has a rate of creatinine clearance greater than or equal to 30 ml/min.

Note: Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

## PROBENECID

Tab 500 mg

## RASBURICASE – **Restricted** see terms below

⚡ Inj 1.5 mg vial

## ➡**Restricted**

Haematologist

## Muscle Relaxants and Related Agents

### ATRACURIUM BESYLATE

Inj 10 mg per ml, 2.5 ml ampoule – **1% DV Sep-12 to 2015** ..... 6.13 5 **Tracrium**  
Inj 10 mg per ml, 5 ml ampoule – **1% DV Sep-12 to 2015** ..... 9.19 5 **Tracrium**

### BACLOFEN

Tab 10 mg – **1% DV Jun-13 to 2016** ..... 3.85 100 **Pacifen**  
Oral liq 1 mg per ml  
Inj 0.05 mg per ml, 1 ml ampoule – **1% DV Oct-12 to 2015** ..... 11.55 1 **Lioresal Intrathecal**  
Inj 2 mg per ml, 5 ml ampoule – **1% DV Oct-12 to 2015** ..... 209.29 1 **Lioresal Intrathecal**

### CLOSTRIDIUM BOTULINUM TYPE A TOXIN

Inj 100 u vial ..... 467.50 1 Botox  
Inj 500 u vial ..... 1,295.00 2 Dysport

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>DANTROLENE</b>			
Cap 25 mg .....	65.00	100	Dantrium
Cap 50 mg .....	77.00	100	Dantrium
Inj 20 mg vial			<i>e.g. Dantrium IV</i>
<b>MIVACURIUM CHLORIDE</b>			
Inj 2 mg per ml, 5 ml ampoule .....	33.92	5	Mivacron
Inj 2 mg per ml, 10 ml ampoule .....	67.17	5	Mivacron
<b>ORPHENADRINE CITRATE</b>			
Tab 100 mg			
<b>PANCURONIUM BROMIDE</b>			
Inj 2 mg per ml, 2 ml ampoule – <b>1% DV Jan-13 to 2015</b> .....	260.00	50	<b>AstraZeneca</b>
<b>ROCURONIUM BROMIDE</b>			
Inj 10 mg per ml, 5 ml vial – <b>1% DV Sep-12 to 2015</b> .....	38.25	10	<b>DBL Rocuronium Bromide</b>
<b>SUXAMETHONIUM CHLORIDE</b>			
Inj 50 mg per ml, 2 ml ampoule – <b>1% DV Jun-14 to 2017</b> .....	78.00	50	<b>AstraZeneca</b>
<b>VECURONIUM BROMIDE</b>			
Inj 4 mg ampoule			
Inj 10 mg vial			

### Reversers of Neuromuscular Blockade

**SUGAMMADEX – Restricted** see terms below

⚡ Inj 100 mg per ml, 2 ml vial .....	1,200.00	10	Bridion
⚡ Inj 100 mg per ml, 5 ml vial .....	3,000.00	10	Bridion

#### ➡Restricted

Any of the following:

- 1 Patient requires reversal of profound neuromuscular blockade following rapid sequence induction that has been undertaken using rocuronium (i.e. suxamethonium is contraindicated or undesirable); or
- 2 Severe neuromuscular degenerative disease where the use of neuromuscular blockade is required; or
- 3 Patient has an unexpectedly difficult airway that cannot be intubated and requires a rapid reversal of anaesthesia and neuromuscular blockade; or
- 4 The duration of the patient's surgery is unexpectedly short; or
- 5 Neostigmine or a neostigmine/anticholinergic combination is contraindicated (for example the patient has ischaemic heart disease, morbid obesity or COPD); or
- 6 Patient has a partial residual block after conventional reversal.

### Non-Steroidal Anti-Inflammatory Drugs

**CELECOXIB – Restricted** see terms below

⚡ Cap 100 mg
⚡ Cap 200 mg
⚡ Cap 400 mg

#### ➡Restricted

For preoperative and/or postoperative use for a total of up to 8 days' use.

# MUSCULOSKELETAL SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>DICLOFENAC SODIUM</b>			
Tab EC 25 mg – 1% DV Mar-13 to 2015 .....	4.00	100	<b>Apo-Diclo</b>
Tab 50 mg dispersible .....			
Tab EC 50 mg – 1% DV Mar-13 to 2015 .....	16.00	500	<b>Apo-Diclo</b>
Tab long-acting 75 mg – 1% DV Dec-12 to 2015 .....	3.10	30	<b>Diclax SR</b>
	24.52	500	<b>Diclax SR</b>
Tab long-acting 100 mg – 1% DV Dec-12 to 2015 .....	42.25	500	<b>Diclax SR</b>
Inj 25 mg per ml, 3 ml ampoule – 1% DV Sep-11 to 2014 .....	12.00	5	<b>Voltaren</b>
Suppos 12.5 mg – 1% DV Sep-11 to 2014 .....	1.85	10	<b>Voltaren</b>
Suppos 25 mg – 1% DV Sep-11 to 2014 .....	2.22	10	<b>Voltaren</b>
Suppos 50 mg – 1% DV Sep-11 to 2014 .....	3.84	10	<b>Voltaren</b>
Suppos 100 mg – 1% DV Sep-11 to 2014 .....	6.36	10	<b>Voltaren</b>
<b>ETORICOXIB – Restricted</b> see terms below			
☞ Tab 30 mg			
☞ Tab 60 mg			
☞ Tab 90 mg			
☞ Tab 120 mg			
☞ <b>Restricted</b>			
For preoperative and/or postoperative use for a total of up to 8 days' use.			
<b>IBUPROFEN</b>			
Tab 200 mg			
☞ Tab 400 mg – <b>Restricted:</b> For continuation only			
☞ Tab 600 mg – <b>Restricted:</b> For continuation only			
Tab long-acting 800 mg – 1% DV Oct-11 to 2014 .....	8.12	30	<b>Brufen SR</b>
Oral liq 20 mg per ml – 1% DV Mar-14 to 2016 .....	1.89	200 ml	<b>Fenpaed</b>
Inj 5 mg per ml, 2 ml ampoule			
<b>INDOMETHACIN</b>			
Cap 25 mg			
Cap 50 mg			
Cap long-acting 75 mg			
Inj 1 mg vial			
Suppos 100 mg			
<b>KETOPROFEN</b>			
Cap long-acting 100 mg .....	21.56	100	<b>Oruvail SR</b>
Cap long-acting 200 mg .....	12.07	28	<b>Oruvail SR</b>
<i>(Oruvail SR Cap long-acting 100 mg to be delisted 1 September 2014)</i>			
<b>MEFENAMIC ACID – Restricted:</b> For continuation only			
☞ Cap 250 mg			
<b>MELOXICAM – Restricted</b> see terms below			
☞ Tab 7.5 mg			
☞ <b>Restricted</b>			
Either:			
1 Haemophilic arthropathy, with both of the following:			
1.1 The patient has moderate to severe haemophilia with less than or equal to 5% of normal circulating functional clotting factor; and			
1.2 Pain and inflammation associated with haemophilic arthropathy is inadequately controlled by alternative funded treatment options, or alternative funded treatment options are contraindicated; or			
2 For preoperative and/or postoperative use for a total of up to 8 days' use.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>NAPROXEN</b>			
Tab 250 mg – 1% DV Jan-13 to 2015 .....	21.25	500	<b>Noflam 250</b>
Tab 500 mg – 1% DV Jan-13 to 2015 .....	22.25	250	<b>Noflam 500</b>
Tab long-acting 750 mg			
Tab long-acting 1 g			
<b>PARECOXIB</b>			
Inj 40 mg vial .....	100.00	10	Dynastat
<b>SULINDAC – Restricted:</b> For continuation only			
➡ Tab 100 mg			
➡ Tab 200 mg			
<b>TENOXICAM</b>			
Tab 20 mg			
Inj 20 mg vial .....	9.95	1	AFT

### Topical Products for Joint and Muscular Pain

**CAPSAICIN – Restricted** see terms below

⚡ Crm 0.025% ..... 9.95 45 g Zostrix

➡ **Restricted**

Patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

## Agents for Parkinsonism and Related Disorders

### Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE – **Restricted** see terms below

⚡ Tab 50 mg .....	400.00	56	Rilutek
-------------------	--------	----	---------

#### ➡ **Restricted**

##### Initiation

Neurologist or respiratory specialist

*Re-assessment required after 6 months*

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
  - 5.1 The patient is ambulatory; or
  - 5.2 The patient is able to use upper limbs; or
  - 5.3 The patient is able to swallow.

##### Continuation

*Re-assessment required after 18 months*

All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
  - 3.1 The patient is ambulatory; or
  - 3.2 The patient is able to use upper limb; or
  - 3.3 The patient is able to swallow.

TETRABENAZINE

Tab 25 mg – 1% DV Sep-13 to 2016 .....	118.00	112	Motetis
--	--------	-----	---------

### Anticholinergics

BENZTROPINE MESYLATE

Tab 2 mg .....	7.99	60	Benztrop
Inj 1 mg per ml, 2 ml ampoule .....	95.00	5	Cogentin

ORPHENADRINE HYDROCHLORIDE

Tab 50 mg

PROCYCLIDINE HYDROCHLORIDE

Tab 5 mg

### Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE

Cap 100 mg – 1% DV Sep-11 to 2014 .....	38.24	60	Symmetrel
---	-------	----	-----------

APOMORPHINE HYDROCHLORIDE

Inj 10 mg per ml, 1 ml ampoule			
Inj 10 mg per ml, 2 ml ampoule .....	110.00	5	Apomine

BROMOCRIPTINE

Tab 2.5 mg

Cap 5 mg



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>ENTACAPONE</b>			
Tab 200 mg – <b>1% DV Dec-12 to 2015</b> .....	47.92	100	<b>Entapone</b>
<b>LEVODOPA WITH BENSERAZIDE</b>			
Tab dispersible 50 mg with benserazide 12.5 mg .....	10.00	100	Madopar Rapid
Cap 50 mg with benserazide 12.5 mg .....	8.00	100	Madopar 62.5
Cap 100 mg with benserazide 25 mg .....	12.50	100	Madopar 125
Cap long-acting 100 mg with benserazide 25 mg .....	17.00	100	Madopar HBS
Cap 200 mg with benserazide 50 mg .....	25.00	100	Madopar 250
<b>LEVODOPA WITH CARBIDOPA</b>			
Tab 100 mg with carbidopa 25 mg .....	20.00	100	Sinemet <i>e.g. Sindopa</i>
Tab long-acting 200 mg with carbidopa 50 mg .....	47.50	100	Sinemet CR
Tab 250 mg with carbidopa 25 mg .....	40.00	100	Sinemet <i>e.g. Sindopa</i>
<b>LISURIDE HYDROGEN MALEATE</b>			
Tab 200 mcg .....	25.00	30	Dopergin
<b>PRAMIPEXOLE HYDROCHLORIDE</b>			
Tab 0.125 mg .....	1.95	30	Dr Reddy's Pramipexole
Tab 0.25 mg .....	2.40	30	Dr Reddy's Pramipexole
	7.20	100	Ramipex
Tab 0.5 mg .....	4.20	30	Dr Reddy's Pramipexole
Tab 1 mg .....	7.20	30	Dr Reddy's Pramipexole
	24.39	100	Ramipex
<b>ROPINIROLE HYDROCHLORIDE</b>			
Tab 0.25 mg – <b>1% DV Mar-14 to 2016</b> .....	2.36	100	<b>Apo-Ropinirole</b>
Tab 1 mg – <b>1% DV Mar-14 to 2016</b> .....	5.32	100	<b>Apo-Ropinirole</b>
Tab 2 mg – <b>1% DV Mar-14 to 2016</b> .....	7.72	100	<b>Apo-Ropinirole</b>
Tab 5 mg – <b>1% DV Mar-14 to 2016</b> .....	14.48	100	<b>Apo-Ropinirole</b>
<b>SELEGILINE HYDROCHLORIDE</b>			
Tab 5 mg			
<b>TOLCAPONE</b>			
Tab 100 mg – <b>1% DV Sep-11 to 2014</b> .....	126.20	100	<b>Tasmar</b>

## Anaesthetics

### General Anaesthetics

<b>DESFLURANE</b>			
Soln for inhalation 100%, 240 ml bottle – <b>1% DV Dec-12 to 2015</b> .....	1,230.00	6	<b>Suprane</b>
<b>DEXMEDETOMIDINE HYDROCHLORIDE</b>			
Inj 100 mcg per ml, 2 ml vial			
<b>ETOMIDATE</b>			
Inj 2 mg per ml, 10 ml ampoule			
<b>ISOFLURANE</b>			
Soln for inhalation 100%, 250 ml bottle – <b>1% DV Dec-12 to 2015</b> .....	1,020.00	6	<b>Aerrane</b>

## NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>KETAMINE HYDROCHLORIDE</b>			
Inj 1 mg per ml, 100 ml bag			
Inj 4 mg per ml, 50 ml syringe			
Inj 10 mg per ml, 10 ml syringe			
Inj 100 mg per ml, 2 ml vial			
<b>METHOHEXITAL SODIUM</b>			
Inj 10 mg per ml, 50 ml vial			
<b>PROPOFOL</b>			
Inj 10 mg per ml, 20 ml ampoule	7.60	5	Fresofol 1%
Inj 10 mg per ml, 20 ml vial	7.60	5	Provoke MCT-LCT 1%
	42.00		Diprivan
Inj 10 mg per ml, 50 ml syringe	47.00	1	Diprivan
Inj 10 mg per ml, 50 ml vial	4.00	1	Fresofol 1%
			Provoke MCT-LCT 1%
	25.00		Diprivan
Inj 10 mg per ml, 100 ml vial	7.60	1	Fresofol 1%
			Provoke MCT-LCT 1%
	30.00		Diprivan
<b>SEVOFLURANE</b>			
Soln for inhalation 100%, 250 ml bottle – 1% DV Dec-12 to 2015	1,230.00	6	Baxter
<b>THIOPENTAL [THIOPENTONE] SODIUM</b>			
Inj 500 mg ampoule			

### Local Anaesthetics

#### ARTICAINE HYDROCHLORIDE WITH ADRENALINE

- Inj 4% with adrenaline 1:100,000, 1.7 ml dental cartridge
- Inj 4% with adrenaline 1:100,000, 2.2 ml dental cartridge
- Inj 4% with adrenaline 1:200,000, 1.7 ml dental cartridge
- Inj 4% with adrenaline 1:200,000, 2.2 ml dental cartridge

#### BENZOCAINE

- Gel 20%

#### BUPIVACAINE HYDROCHLORIDE

- Inj 5 mg per ml, 4 ml ampoule – 1% DV Jul-14 to 2017 50.00 5 Marcaïn Isobaric
- Inj 2.5 mg per ml, 20 ml ampoule
- Inj 2.5 mg per ml, 20 ml ampoule sterile pack – 1% DV Oct-12 to 2015 35.00 5 Marcaïn
- Inj 5 mg per ml, 10 ml ampoule 35.00 50 Marcaïn
- Inj 5 mg per ml, 10 ml ampoule sterile pack – 1% DV Oct-12 to 2015 28.00 5 Marcaïn
- Inj 5 mg per ml, 20 ml ampoule
- Inj 5 mg per ml, 20 ml ampoule sterile pack – 1% DV Oct-12 to 2015 28.00 5 Marcaïn
- Inj 1.25 mg per ml, 100 ml bag
- Inj 1.25 mg per ml, 200 ml bag
- Inj 2.5 mg per ml, 100 ml bag – 1% DV Jul-14 to 2017 150.00 5 Marcaïn
- Inj 2.5 mg per ml, 200 ml bag
- Inj 1.25 mg per ml, 500 ml bag

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>BUPIVACAINE HYDROCHLORIDE WITH ADRENALINE</b>			
Inj 2.5 mg per ml with adrenaline 1:400,000, 20 ml vial – <b>1% DV Nov-11 to 2014</b> .....	135.00	5	<b>Marcaïn with Adrenaline</b>
Inj 5 mg per ml with adrenaline 1:200,000, 20 ml vial – <b>1% DV Nov-11 to 2014</b> .....	115.00	5	<b>Marcaïn with Adrenaline</b>
<b>BUPIVACAINE HYDROCHLORIDE WITH FENTANYL</b>			
Inj 0.625 mg with fentanyl 2 mcg per ml, 100 ml bag			
Inj 1.25 mg with fentanyl 2 mcg per ml, 100 ml syringe			
Inj 1.25 mg with fentanyl 2 mcg per ml, 100 ml bag – <b>1% DV Nov-11 to 2014</b> .....	210.00	10	<b>Bupafen</b>
Inj 1.25 mg with fentanyl 2 mcg per ml, 200 ml bag – <b>1% DV Nov-11 to 2014</b> .....	210.00	10	<b>Bupafen</b>
Inj 1.25 mg with fentanyl 2 mcg per ml, 50 ml syringe			
Inj 1.25 mg with fentanyl 2 mcg per ml, 15 ml syringe – <b>1% DV Nov-11 to 2014</b> .....	72.00	10	<b>Biomed</b>
Inj 1.25 mg with fentanyl 2 mcg per ml, 20 ml syringe – <b>1% DV Nov-11 to 2014</b> .....	92.00	10	<b>Biomed</b>
<b>BUPIVACAINE HYDROCHLORIDE WITH GLUCOSE</b>			
Inj 0.5% with glucose 8%, 4 ml ampoule .....	38.00	5	<b>Marcaïn Heavy</b>
<b>COCAINE HYDROCHLORIDE</b>			
Paste 5%			
Soln 15%, 2 ml syringe			
Soln 4%, 2 ml syringe .....	25.46	1	<b>Biomed</b>
<b>COCAINE HYDROCHLORIDE WITH ADRENALINE</b>			
Paste 15% with adrenaline 0.06%			
Paste 25% with adrenaline 0.06%			
<b>ETHYL CHLORIDE</b>			
Spray 100%			
<b>LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE</b>			
Gel 2% – <b>1% DV Oct-12 to 2015</b> .....	3.40	20 ml	<b>Orion</b>
Soln 4%			
Spray 10% – <b>1% DV Sep-13 to 2016</b> .....	75.00	50 ml	<b>Xylocaine</b>
Oral (viscous) soln 2% – <b>1% DV Sep-11 to 2014</b> .....	55.00	200 ml	<b>Xylocaine Viscous</b>
Inj 1%, 20 ml ampoule, sterile pack			
Inj 2%, 20 ml ampoule, sterile pack			
Inj 1%, 5 ml ampoule – <b>1% DV Jul-13 to 2015</b> .....	8.75	25	<b>Lidocaine-Claris</b>
Inj 1%, 20 ml ampoule – <b>1% DV Jul-13 to 2015</b> .....	2.40	1	<b>Lidocaine-Claris</b>
Inj 2%, 5 ml ampoule – <b>1% DV Jul-13 to 2015</b> .....	6.90	25	<b>Lidocaine-Claris</b>
Inj 2%, 20 ml ampoule – <b>1% DV Jul-13 to 2015</b> .....	2.40	1	<b>Lidocaine-Claris</b>
Gel 2%, 10 ml urethral syringe .....	43.26	10	<b>Pfizer</b>

# NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH ADRENALINE</b>			
Inj 1% with adrenaline 1:100,000, 5 ml ampoule .....	27.00	10	Xylocaine
Inj 1% with adrenaline 1:200,000, 20 ml vial .....	50.00	5	Xylocaine
Inj 2% with adrenaline 1:80,000, 1.7 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 1.8 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 2.2 ml dental cartridge			
Inj 2% with adrenaline 1:200,000, 20 ml vial .....	60.00	5	Xylocaine
<b>LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH ADRENALINE AND TETRACAINE HYDROCHLORIDE</b>			
Soln 4% with adrenaline 0.1% and tetracaine hydrochloride 0.5%, 5 ml syringe			
<b>LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH CHLORHEXIDINE</b>			
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringe .....	43.26	10	Pfizer
<b>LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH PHENYLEPHRINE HYDROCHLORIDE</b>			
Nasal spray 5% with phenylephrine hydrochloride 0.5%			
<b>LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE</b>			
Crm 2.5% with prilocaine 2.5% .....	45.00	30 g	EMLA
Patch 25 mcg with prilocaine 25 mcg .....	115.00	20	EMLA
Crm 2.5% with prilocaine 2.5%, 5 g .....	45.00	5	EMLA
<b>MEPIVACAINE HYDROCHLORIDE</b>			
Inj 3%, 1.8 ml dental cartridge			
Inj 3%, 2.2 ml dental cartridge			
<b>PRILOCAINE HYDROCHLORIDE</b>			
Inj 0.5%, 50 ml vial .....	100.00	5	Citanest
Inj 2%, 5 ml ampoule .....	55.00	10	Citanest
<b>PRILOCAINE HYDROCHLORIDE WITH FELYPRESSIN</b>			
Inj 3% with felypressin 0.03 iu per ml, 1.8 ml dental cartridge			
Inj 3% with felypressin 0.03 iu per ml, 2.2 ml dental cartridge			
<b>ROPIVACAINE HYDROCHLORIDE</b>			
Inj 2 mg per ml, 10 ml ampoule			
Inj 2 mg per ml, 20 ml ampoule .....	75.00	5	Naropin
Inj 2 mg per ml, 100 ml bag .....	200.00	5	Naropin
Inj 2 mg per ml, 200 ml bag .....	265.00	5	Naropin
Inj 7.5 mg per ml, 10 ml ampoule .....	45.00	5	Naropin
Inj 7.5 mg per ml, 20 ml ampoule .....	84.00	5	Naropin
Inj 10 mg per ml, 10 ml ampoule .....	54.00	5	Naropin
Inj 10 mg per ml, 20 ml ampoule			
<b>ROPIVACAINE HYDROCHLORIDE WITH FENTANYL</b>			
Inj 2 mg with fentanyl 2 mcg per ml, 100 ml bag .....	198.50	5	Naropin
Inj 2 mg with fentanyl 2 mcg per ml, 200 ml bag .....	270.00	5	Naropin
<b>TETRACAINE [AMETHOCAINE] HYDROCHLORIDE</b>			
Gel 4%			

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

Price (ex man. excl. GST)	Brand or Generic
\$	Per Manufacturer

## Analgesics

### Non-Opioid Analgesics

#### ASPIRIN

Tab EC 300 mg

Tab dispersible 300 mg

#### CAPSAICIN – **Restricted** see terms below

¶ Crm 0.075% ..... 12.50 45 g Zostrix HP

#### ➔**Restricted**

For post-herpetic neuralgia or diabetic peripheral neuropathy

#### METHOXYFLURANE – **Restricted** see terms below

¶ Soln for inhalation 99.9%, 3 ml bottle

#### ➔**Restricted**

Both:

- 1 Patient is undergoing a painful procedure with an expected duration of less than one hour; and
- 2 Only to be used under supervision by a medical practitioner or nurse who is trained in the use of methoxyflurane.

#### NEFOPAM HYDROCHLORIDE

Tab 30 mg

#### PARACETAMOL – **Some items restricted** see terms below

Tab soluble 500 mg

Tab 500 mg

Oral liq 120 mg per 5 ml – 20% DV Dec-11 to 2014 .....	2.21	500 ml	<b>Ethics Paracetamol</b>
Oral liq 250 mg per 5 ml – 20% DV Sep-11 to 2014 .....	6.70	1,000 ml	<b>Paracare Double Strength</b>

¶ Inj 10 mg per ml, 50 ml vial – 1% DV Dec-13 to 2014 .....	22.50	10	<b>Paracetamol-AFT</b>
¶ Inj 10 mg per ml, 100 ml vial – 1% DV Apr-13 to 2014 .....	22.50	10	<b>Paracetamol-AFT</b>
Suppos 25 mg .....	56.35	20	<b>Biomed</b>
Suppos 50 mg .....	56.35	20	<b>Biomed</b>
Suppos 125 mg .....	7.49	20	<b>Panadol</b>
Suppos 250 mg .....	14.40	20	<b>Panadol</b>
Suppos 500 mg – 1% DV Jan-13 to 2015 .....	20.70	50	<b>Paracare</b>

#### ➔**Restricted**

Intravenous paracetamol is only to be used where other routes are unavailable or impractical, or where there is reduced absorption. The need for IV paracetamol must be re-assessed every 24 hours.

#### SUCROSE

Oral liq 25%

## Opioid Analgesics

#### ALFENTANIL HYDROCHLORIDE

Inj 0.5 mg per ml, 2 ml ampoule

#### CODEINE PHOSPHATE

Tab 15 mg – 1% DV Jul-13 to 2016 .....	4.75	100	<b>PSM</b>
Tab 30 mg – 1% DV Jul-13 to 2016 .....	5.80	100	<b>PSM</b>
Tab 60 mg – 1% DV Jul-13 to 2016 .....	12.50	100	<b>PSM</b>

#### DIHYDROCODEINE TARTRATE

Tab long-acting 60 mg – 1% DV Sep-13 to 2016 .....	13.64	60	<b>DHC Continus</b>
--	-------	----	---------------------

# NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>FENTANYL</b>			
Inj 10 mcg per ml, 10 ml syringe			
Inj 50 mcg per ml, 2 ml ampoule – 1% DV Sep-12 to 2015	4.50	10	<b>Boucher and Muir</b>
Inj 10 mcg per ml, 50 ml bag – 1% DV Dec-11 to 2014	210.00	10	<b>Biomed</b>
Inj 10 mcg per ml, 50 ml syringe – 1% DV Dec-11 to 2014	165.00	10	<b>Biomed</b>
Inj 50 mcg per ml, 10 ml ampoule – 1% DV Sep-12 to 2015	11.77	10	<b>Boucher and Muir</b>
Inj 10 mcg per ml, 100 ml bag – 1% DV Dec-11 to 2014	210.00	10	<b>Biomed</b>
Inj 20 mcg per ml, 50 ml syringe – 1% DV Dec-11 to 2014	185.00	10	<b>Biomed</b>
Inj 20 mcg per ml, 100 ml bag			
Patch 12.5 mcg per hour	8.90	5	Mylan Fentanyl Patch
Patch 25 mcg per hour	9.15	5	Mylan Fentanyl Patch
Patch 50 mcg per hour	11.50	5	Mylan Fentanyl Patch
Patch 75 mcg per hour	13.60	5	Mylan Fentanyl Patch
Patch 100 mcg per hour	14.50	5	Mylan Fentanyl Patch
<b>METHADONE HYDROCHLORIDE</b>			
Tab 5 mg	1.85	10	Methatabs
Oral liq 2 mg per ml – 1% DV Sep-12 to 2015	5.55	200 ml	<b>Biodone</b>
Oral liq 5 mg per ml – 1% DV Sep-12 to 2015	5.55	200 ml	<b>Biodone Forte</b>
Oral liq 10 mg per ml – 1% DV Sep-12 to 2015	6.55	200 ml	<b>Biodone Extra Forte</b>
Inj 10 mg per ml, 1 ml vial	61.00	10	AFT
<b>MORPHINE HYDROCHLORIDE</b>			
Oral liq 1 mg per ml – 1% DV Oct-12 to 2015	8.84	200 ml	<b>RA-Morph</b>
Oral liq 2 mg per ml – 1% DV Oct-12 to 2015	11.62	200 ml	<b>RA-Morph</b>
Oral liq 5 mg per ml – 1% DV Oct-12 to 2015	14.65	200 ml	<b>RA-Morph</b>
Oral liq 10 mg per ml – 1% DV Oct-12 to 2015	21.55	200 ml	<b>RA-Morph</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>MORPHINE SULPHATE</b>			
Tab long-acting 10 mg – 1% DV Sep-13 to 2016	1.95	10	<b>Arrow-Morphine LA</b>
Tab immediate-release 10 mg	2.80	10	Sevredol
Tab immediate-release 20 mg	5.52	10	Sevredol
Tab long-acting 30 mg – 1% DV Sep-13 to 2016	2.98	10	<b>Arrow-Morphine LA</b>
Tab long-acting 60 mg – 1% DV Sep-13 to 2016	5.75	10	<b>Arrow-Morphine LA</b>
Tab long-acting 100 mg – 1% DV Sep-13 to 2016	6.45	10	<b>Arrow-Morphine LA</b>
Cap long-acting 10 mg – 1% DV Feb-14 to 2016	1.70	10	<b>m-Eslon</b>
Cap long-acting 30 mg – 1% DV Feb-14 to 2016	2.50	10	<b>m-Eslon</b>
Cap long-acting 60 mg – 1% DV Feb-14 to 2016	5.40	10	<b>m-Eslon</b>
Cap long-acting 100 mg – 1% DV Feb-14 to 2016	6.38	10	<b>m-Eslon</b>
Inj 1 mg per ml, 100 ml bag – 1% DV Dec-11 to 2014	165.00	10	<b>Biomed</b>
Inj 1 mg per ml, 10 ml syringe – 1% DV Dec-11 to 2014	39.50	10	<b>Biomed</b>
Inj 1 mg per ml, 50 ml syringe – 1% DV Dec-11 to 2014	79.50	10	<b>Biomed</b>
Inj 1 mg per ml, 2 ml syringe			
Inj 2 mg per ml, 30 ml syringe – 1% DV Dec-11 to 2014	135.00	10	<b>Biomed</b>
Inj 5 mg per ml, 1 ml ampoule – 1% DV Nov-11 to 2014	5.51	5	<b>DBL Morphine Sulphate</b>
Inj 10 mg per ml, 1 ml ampoule – 1% DV Nov-11 to 2014	4.79	5	<b>DBL Morphine Sulphate</b>
Inj 10 mg per ml, 100 mg cassette			
Inj 10 mg per ml, 100 ml bag			
Inj 15 mg per ml, 1 ml ampoule – 1% DV Nov-11 to 2014	5.01	5	<b>DBL Morphine Sulphate</b>
Inj 30 mg per ml, 1 ml ampoule – 1% DV Nov-11 to 2014	5.30	5	<b>DBL Morphine Sulphate</b>
Inj 200 mcg in 0.4 ml syringe			
Inj 300 mcg in 0.3 ml syringe			
<b>MORPHINE TARTRATE</b>			
Inj 80 mg per ml, 1.5 ml ampoule – 1% DV Sep-13 to 2016	35.60	5	<b>Hospira</b>
Inj 80 mg per ml, 5 ml ampoule – 1% DV Sep-13 to 2016	107.67	5	<b>Hospira</b>
<b>OXYCODONE HYDROCHLORIDE</b>			
Tab controlled-release 5 mg	7.51	20	<b>OxyContin</b>
Tab controlled-release 10 mg – 1% DV Oct-13 to 2015	6.75	20	<b>BNM</b>
Tab controlled-release 20 mg – 1% DV Oct-13 to 2015	11.50	20	<b>Oxydone BNM</b>
Tab controlled-release 40 mg – 1% DV Oct-13 to 2015	18.50	20	<b>BNM</b>
Tab controlled-release 80 mg – 1% DV Oct-13 to 2015	34.00	20	<b>Oxydone BNM</b>
Cap immediate-release 5 mg	2.83	20	<b>Oxydone BNM</b>
Cap immediate-release 10 mg	5.58	20	<b>OxyNorm</b>
Cap immediate-release 20 mg	9.77	20	<b>OxyNorm</b>
Oral liq 5 mg per 5 ml	11.20	250 ml	<b>OxyNorm</b>
Inj 1 mg per ml, 100 ml bag			
Inj 10 mg per ml, 1 ml ampoule – 1% DV Dec-12 to 2015	10.08	5	<b>Oxycodone Orion</b>
Inj 10 mg per ml, 2 ml ampoule – 1% DV Dec-12 to 2015	19.87	5	<b>Oxycodone Orion</b>
Inj 50 mg per ml, 1 ml ampoule – 1% DV May-13 to 2015	60.00	5	<b>OxyNorm</b>
<i>(Oxydone BNM Tab controlled-release 10 mg to be delisted 1 August 2014)</i>			
<i>(Oxydone BNM Tab controlled-release 20 mg to be delisted 1 August 2014)</i>			

## NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PARACETAMOL WITH CODEINE</b>			
Tab paracetamol 500 mg with codeine phosphate 8 mg – 1% DV Nov-11 to 2014 .....	2.70	100	<b>Paracetamol + Codeine (Relieve)</b>
<b>PETHIDINE HYDROCHLORIDE</b>			
Tab 50 mg – 1% DV Mar-13 to 2015 .....	3.95	10	<b>PSM</b>
Tab 100 mg – 1% DV Mar-13 to 2015 .....	5.80	10	<b>PSM</b>
Inj 5 mg per ml, 10 ml syringe			
Inj 5 mg per ml, 100 ml bag			
Inj 10 mg per ml, 100 ml bag			
Inj 10 mg per ml, 50 ml syringe			
Inj 50 mg per ml, 1 ml ampoule – 1% DV Nov-11 to 2014 .....	5.51	5	<b>DBL Pethidine Hydrochloride</b>
Inj 50 mg per ml, 2 ml ampoule – 1% DV Nov-11 to 2014 .....	5.83	5	<b>DBL Pethidine Hydrochloride</b>
<b>REMIFENTANIL HYDROCHLORIDE</b>			
Inj 1 mg vial – 1% DV Feb-12 to 2014 .....	27.95	5	<b>Remifentanil-AFT</b>
Inj 2 mg vial – 1% DV Feb-12 to 2014 .....	41.80	5	<b>Remifentanil-AFT</b>
<b>TRAMADOL HYDROCHLORIDE</b>			
Tab sustained-release 100 mg .....	2.14	20	<b>Tramal SR 100</b>
Tab sustained-release 150 mg .....	3.21	20	<b>Tramal SR 150</b>
Tab sustained-release 200 mg .....	4.28	20	<b>Tramal SR 200</b>
Cap 50 mg – 1% DV Sep-11 to 2014 .....	4.95	100	<b>Arrow-Tramadol</b>
Oral drops 100 mg per ml			
Inj 10 mg per ml, 100 ml bag			
Inj 50 mg per ml, 1 ml ampoule .....	4.50	5	<b>Tramal 50</b>
Inj 50 mg per ml, 2 ml ampoule .....	4.50	5	<b>Tramal 100</b>
<b>Antidepressants</b>			
<b>Cyclic and Related Agents</b>			
<b>AMITRIPTYLINE</b>			
Tab 10 mg – 1% DV Jan-13 to 2014 .....	3.32	100	<b>Arrow-Amitriptyline</b>
Tab 25 mg – 1% DV Jun-11 to 2014 .....	1.85	100	<b>Amitrip</b>
Tab 50 mg – 1% DV Jun-11 to 2014 .....	3.60	100	<b>Amitrip</b>
<b>CLOMIPRAMINE HYDROCHLORIDE</b>			
Tab 10 mg – 1% DV Jan-13 to 2015 .....	12.60	100	<b>Apo-Clomipramine</b>
Tab 25 mg – 1% DV Jan-13 to 2015 .....	8.68	100	<b>Apo-Clomipramine</b>
<b>DOTHIEPIN HYDROCHLORIDE</b>			
Tab 75 mg .....	10.50	100	<b>Dopress</b>
Cap 25 mg .....	6.17	100	<b>Dopress</b>
<b>DOXEPIN HYDROCHLORIDE</b>			
Cap 10 mg			
Cap 25 mg			
Cap 50 mg			



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>IMIPRAMINE HYDROCHLORIDE</b>			
Tab 10 mg .....	5.48	50	Tofranil
	6.58	60	Tofranil
Tab 25 mg .....	8.80	50	Tofranil
<b>MAPROTILINE HYDROCHLORIDE</b>			
Tab 25 mg			
Tab 75 mg			
<b>MIANSERIN HYDROCHLORIDE</b>			
Tab 30 mg			
<b>NORTRIPTYLINE HYDROCHLORIDE</b>			
Tab 10 mg – 1% DV Jun-13 to 2016 .....	4.00	100	Norpress
Tab 25 mg – 1% DV Jun-13 to 2016 .....	9.00	180	Norpress

### Monoamine-Oxidase Inhibitors - Non-Selective

<b>PHENELZINE SULPHATE</b>			
Tab 15 mg			
<b>TRANLYCYPROMINE SULPHATE</b>			
Tab 10 mg			

### Monoamine-Oxidase Type A Inhibitors

<b>MOCLOBEMIDE</b>			
Tab 150 mg – 1% DV Apr-13 to 2015 .....	81.83	500	Apo-Moclobemide
Tab 300 mg – 1% DV Apr-13 to 2015 .....	29.51	100	Apo-Moclobemide

### Other Antidepressants

<b>MIRTAZAPINE – Restricted</b> see terms below			
⚡ Tab 30 mg – 1% DV Sep-12 to 2015 .....	8.78	30	Avanza
⚡ Tab 45 mg – 1% DV Sep-12 to 2015 .....	13.95	30	Avanza

#### ➡Restricted

#### Initiation

*Re-assessment required after two years*

- Both:
- 1 The patient has a severe major depressive episode; and
  - 2 Either:
    - 2.1 The patient must have had a trial of two different antidepressants and was unable to tolerate the treatments or failed to respond to an adequate dose over an adequate period of time (usually at least four weeks); or
    - 2.2 Both:
      - 2.2.1 The patient is currently a hospital in-patient as a result of an acute depressive episode; and
      - 2.2.2 The patient must have had a trial of one other antidepressant and either could not tolerate it or failed to respond to an adequate dose over an adequate period of time.

#### Continuation

*Re-assessment required after two years*

The patient has a high risk of relapse (prescriber determined)

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>VENLAFAXINE – Some items restricted</b> see terms below			
Tab modified release 37.5 mg .....	5.06	28	Arrow-Venlafaxine XR
Tab modified release 75 mg .....	6.44	28	Arrow-Venlafaxine XR
Tab modified release 150 mg .....	8.86	28	Arrow-Venlafaxine XR
Tab modified release 225 mg .....	14.34	28	Arrow-Venlafaxine XR
⚡ Cap modified release 37.5 mg .....	8.71	28	Efexor XR
⚡ Cap modified release 75 mg .....	17.42	28	Efexor XR
⚡ Cap modified release 150 mg .....	21.35	28	Efexor XR

## ➡ Restricted

### Initiation

*Re-assessment required after two years*

Both:

- 1 The patient has 'treatment-resistant' depression; and
- 2 Either:
  - 2.1 The patient must have had a trial of two different antidepressants and have had an inadequate response from an adequate dose over an adequate period of time (usually at least four weeks); or
  - 2.2 Both:
    - 2.2.1 The patient is currently a hospital in-patient as a result of an acute depressive episode; and
    - 2.2.2 The patient must have had a trial of one other antidepressant and have had an inadequate response from an adequate dose over an adequate period of time. Continuation.

### Continuation

*Re-assessment required after two years*

The patient has a high risk of relapse (prescriber determined)

## Selective Serotonin Reuptake Inhibitors

<b>CITALOPRAM HYDROBROMIDE</b>			
Tab 20 mg – 1% DV Sep-11 to 2014 .....	2.34	84	<b>Arrow-Citalopram</b>
<b>ESCITALOPRAM</b>			
Tab 10 mg .....	2.65	28	Loxalate
Tab 20 mg .....	4.20	28	Loxalate
<b>FLUOXETINE HYDROCHLORIDE</b>			
Tab dispersible 20 mg, scored – 1% DV Apr-14 to 2016 .....	2.50	30	<b>Arrow-Fluoxetine</b>
Cap 20 mg – 1% DV Apr-14 to 2016 .....	1.74	90	<b>Arrow-Fluoxetine</b>
<b>PAROXETINE HYDROCHLORIDE</b>			
Tab 20 mg .....	4.32	90	Loxamine
<b>SERTRALINE</b>			
Tab 50 mg – 1% DV Sep-13 to 2016 .....	3.64	90	<b>Arrow-Sertraline</b>
Tab 100 mg – 1% DV Sep-13 to 2016 .....	6.28	90	<b>Arrow-Sertraline</b>

## Antiepilepsy Drugs

### Agents for the Control of Status Epilepticus

<b>CLONAZEPAM</b>			
Inj 1 mg per ml, 1 ml ampoule .....	19.00	5	Rivotril
<b>DIAZEPAM</b>			
Inj 5 mg per ml, 2 ml ampoule .....	9.24	5	Hospira
Rectal tubes 5 mg .....	25.05	5	Stesolid
Rectal tubes 10 mg .....	30.50	5	Stesolid

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LORAZEPAM			
Inj 2 mg vial			
Inj 4 mg per ml, 1 ml vial			
PARALDEHYDE			
Inj 5 ml ampoule			
PHENYTOIN SODIUM			
Inj 50 mg per ml, 2 ml ampoule			
Inj 50 mg per ml, 5 ml ampoule			

Control of Epilepsy

CARBAMAZEPINE			
Tab 200 mg			
Tab long-acting 200 mg			
Tab 400 mg			
Tab long-acting 400 mg			
Oral liq 20 mg per ml			
CLOBAZAM			
Tab 10 mg			
CLONAZEPAM			
Oral drops 2.5 mg per ml			
ETHOSUXIMIDE			
Cap 250 mg			
Oral liq 50 mg per ml			
GABAPENTIN – <b>Restricted</b> see terms below			
⚡ Tab 600 mg			
⚡ Cap 100 mg .....	7.16	100	Arrow-Gabapentin Nupentin
⚡ Cap 300 mg .....	11.00	100	Arrow-Gabapentin Nupentin
⚡ Cap 400 mg .....	13.75	100	Arrow-Gabapentin Nupentin

➡ Restricted

- 1 For preoperative and/or postoperative use for up to a total of 8 days' use; or
- 2 For the pain management of burns patients with monthly review.

Initiation - epilepsy  
Re-assessment required after 15 months  
Either:

- 1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
- 2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Note: "Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

## Continuation - epilepsy

Patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life.

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

## Initiation - neuropathic pain

*Re-assessment required after 3 months*

Patient has tried and failed, or has been unable to tolerate, treatment with a tricyclic antidepressant.

## Continuation - neuropathic pain

Either:

- 1 The patient has demonstrated a marked improvement in their control of pain (prescriber determined); or
- 2 The patient has previously demonstrated clinical responsiveness to gabapentin and has now developed neuropathic pain in a new site.

LACOSAMIDE – **Restricted** see terms below

⚡ Tab 50 mg .....	25.04	14	Vimpat
⚡ Tab 100 mg .....	50.06	14	Vimpat
	200.24	56	Vimpat
⚡ Tab 150 mg .....	75.10	14	Vimpat
	300.40	56	Vimpat
⚡ Tab 200 mg .....	400.55	56	Vimpat
⚡ Inj 10 mg per ml, 20 ml vial			

## ➡Restricted

## Initiation

*Re-assessment required after 15 months*

Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Women of childbearing age are not required to have a trial of sodium valproate.

## Continuation

Patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>LAMOTRIGINE</b>			
Tab dispersible 2 mg .....	6.74	30	Lamictal
Tab dispersible 5 mg .....	9.64	30	Lamictal
	15.00	56	Arrow-Lamotrigine
Tab dispersible 25 mg .....	19.38	56	Logem
	20.40		Arrow-Lamotrigine
			Mogine
	29.09		Lamictal
Tab dispersible 50 mg .....	32.97	56	Logem
	34.70		Arrow-Lamotrigine
			Mogine
	47.89		Lamictal
Tab dispersible 100 mg .....	56.91	56	Logem
	59.90		Arrow-Lamotrigine
			Mogine
	79.16		Lamictal
<b>LEVETIRACETAM</b>			
Tab 250 mg .....	24.03	60	Levetiracetam-Rex
Tab 500 mg .....	28.71	60	Levetiracetam-Rex
Tab 750 mg .....	45.23	60	Levetiracetam-Rex
Inj 100 mg per ml, 5 ml vial			
<b>PHENOBARBITONE</b>			
Tab 15 mg – 1% DV Mar-13 to 2015 .....	28.00	500	<b>PSM</b>
Tab 30 mg – 1% DV Mar-13 to 2015 .....	29.00	500	<b>PSM</b>
<b>PHENYTOIN</b>			
Tab 50 mg			
<b>PHENYTOIN SODIUM</b>			
Cap 30 mg			
Cap 100 mg			
Oral liq 6 mg per ml			
<b>PRIMIDONE</b>			
Tab 250 mg			
<b>SODIUM VALPROATE</b>			
Tab 100 mg			
Tab EC 200 mg			
Tab EC 500 mg			
Oral liq 40 mg per ml			
Inj 100 mg per ml, 4 ml vial			
<b>STIRIPENTOL – <b>Restricted</b> see terms on the next page</b>			
⚡ Cap 250 mg .....	509.29	60	Diacomit
⚡ Powder for oral liq 250 mg sachet .....	509.29	60	Diacomit

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➡Restricted</b>			
Paediatric neurologist			
<b>Initiation</b>			
<i>Re-assessment required after 6 months</i>			
Both:			
1 Patient has confirmed diagnosis of Dravet syndrome; and			
2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.			
<b>Continuation</b>			
Patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.			
<b>TOPIRAMATE</b>			
Tab 25 mg .....	11.07	60	Arrow-Topiramate
	26.04		Topamax
Tab 50 mg .....	18.81	60	Arrow-Topiramate
	44.26		Topamax
Tab 100 mg .....	31.99	60	Arrow-Topiramate
	75.25		Topamax
Tab 200 mg .....	55.19	60	Arrow-Topiramate
	129.85		Topamax
Cap sprinkle 15 mg .....	20.84	60	Topamax
Cap sprinkle 25 mg .....	26.04	60	Topamax

**VIGABATRIN – Restricted** see terms below

⚡ Tab 500 mg

**➡Restricted**

- Both:
- Either:
    - 1.1 Patient has infantile spasms; or
    - 1.2 Both:
      - 1.2.1 Patient has epilepsy; and
      - 1.2.2 Either:
        - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
        - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and
  - 2 Either:
    - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
    - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Notes:

"Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

## Antimigraine Preparations

### Acute Migraine Treatment

**DIHYDROERGOTAMINE MESYLATE**

Inj 1 mg per ml, 1 ml ampoule

**ERGOTAMINE TARTRATE WITH CAFFEINE**

Tab 1 mg with caffeine 100 mg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>METOCLOPRAMIDE HYDROCHLORIDE WITH PARACETAMOL</b>			
Tab 5 mg with paracetamol 500 mg			
<b>RIZATRIPTAN BENZOATE</b>			
Tab orodispersible 10 mg – 1% DV May-12 to 2014 .....	18.00	30	<b>Rizamelt</b>
<b>SUMATRIPTAN</b>			
Tab 50 mg – 1% DV Sep-13 to 2016 .....	29.80	100	<b>Arrow-Sumatriptan</b>
Tab 100 mg – 1% DV Sep-13 to 2016 .....	54.80	100	<b>Arrow-Sumatriptan</b>
Inj 12 mg per ml, 0.5 ml cartridge – 1% DV Sep-13 to 2016 .....	13.80	2	<b>Arrow-Sumatriptan</b>

## Prophylaxis of Migraine

<b>PIZOTIFEN</b>			
Tab 500 mcg – 1% DV Mar-13 to 2015 .....	23.21	100	<b>Sandomigran</b>

## Antinausea and Vertigo Agents

APREPITANT – **Restricted** see terms below

⚡ Cap 2 × 80 mg and 1 × 125 mg .....	116.00	3	Emend Tri-Pack
--------------------------------------	--------	---	----------------

➡ **Restricted**

Patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

<b>BETAHISTINE DIHYDROCHLORIDE</b>			
Tab 16 mg – 1% DV Jun-14 to 2017 .....	4.95	84	<b>Vergo 16</b>

<b>CYCLIZINE HYDROCHLORIDE</b>			
Tab 50 mg – 1% DV Sep-12 to 2015 .....	0.59	10	<b>Nausicalm</b>

<b>CYCLIZINE LACTATE</b>			
Inj 50 mg per ml, 1 ml ampoule .....	14.95	5	Nausicalm

<b>DOMPERIDONE</b>			
Tab 10 mg – 1% DV Mar-13 to 2015 .....	3.25	100	<b>Prokinex</b>

<b>DROPERIDOL</b>			
Inj 2.5 mg per ml, 1 ml ampoule			

<b>HYOSCINE HYDROBROMIDE</b>			
Inj 400 mcg per ml, 1 ml ampoule .....	6.66	5	Hospira
⚡ Patch 1.5 mg – 1% DV Dec-13 to 2016 .....	11.95	2	<b>Scopoderm TTS</b>

➡ **Restricted**

Any of the following:

- Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective; or
- For treatment of post-operative nausea and vomiting where cyclizine, droperidol and a 5HT3 antagonist have proven ineffective, are not tolerated or are contraindicated.

<b>METOCLOPRAMIDE HYDROCHLORIDE</b>			
Tab 10 mg – 1% DV Jun-11 to 2014 .....	3.95	100	<b>Metamide</b>
Oral liq 5 mg per 5 ml			
Inj 5 mg per ml, 2 ml ampoule – 1% DV Sep-11 to 2014 .....	4.50	10	<b>Pfizer</b>

## NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>ONDANSETRON</b>			
Tab 4 mg – 1% DV Jan-14 to 2016 .....	5.51	50	<b>Onrex</b>
Tab dispersible 4 mg .....	1.70	10	Dr Reddy's Ondansetron
	17.18		Zofran Zydis
Tab 8 mg – 1% DV Jan-14 to 2016 .....	6.19	50	<b>Onrex</b>
Tab dispersible 8 mg .....	2.00	10	Dr Reddy's Ondansetron
Inj 2 mg per ml, 2 ml ampoule – 1% DV Sep-13 to 2016 .....	1.82	5	<b>Ondanaccord</b>
Inj 2 mg per ml, 4 ml ampoule – 1% DV Sep-13 to 2016 .....	2.18	5	<b>Ondanaccord</b>

### PROCHLORPERAZINE

Tab buccal 3 mg			
Tab 5 mg – 1% DV Jun-14 to 2017 .....	9.75	500	<b>Antinaus</b>
Inj 12.5 mg per ml, 1 ml ampoule			
Suppos 25 mg			

### PROMETHAZINE THEOCLATE – **Restricted:** For continuation only

➡ Tab 25 mg

### TROPISETRON

Cap 5 mg .....	77.41	5	Navoban
Inj 1 mg per ml, 2 ml ampoule – 1% DV May-14 to 2015 .....	8.95	1	<b>Tropisetron-AFT</b>
Inj 1 mg per ml, 5 ml ampoule – 1% DV May-14 to 2015 .....	13.95	1	<b>Tropisetron-AFT</b>

(Navoban Cap 5 mg to be delisted 1 August 2014)

## Antipsychotic Agents

### General

#### AMISULPRIDE

Tab 100 mg – 1% DV Jul-13 to 2016 .....	6.22	30	<b>Solian</b>
Tab 200 mg – 1% DV Jul-13 to 2016 .....	21.92	60	<b>Solian</b>
Tab 400 mg – 1% DV Jul-13 to 2016 .....	44.52	60	<b>Solian</b>
Oral liq 100 mg per ml – 1% DV Jul-13 to 2016 .....	52.50	60 ml	<b>Solian</b>

#### ARIPRAZOLE – **Restricted** see terms below

⚡ Tab 10 mg .....	123.54	30	Abilify
⚡ Tab 15 mg .....	175.28	30	Abilify
⚡ Tab 20 mg .....	213.42	30	Abilify
⚡ Tab 30 mg .....	260.07	30	Abilify

#### ➡ **Restricted**

Both:

- 1 Patient is suffering from schizophrenia or related psychoses; and
- 2 Either:
  - 2.1 An effective dose of risperidone or quetiapine has been trialled and has been discontinued, or is in the process of being discontinued, because of unacceptable side effects; or
  - 2.2 An effective dose of risperidone or quetiapine has been trialled and has been discontinued, or is in the process of being discontinued, because of inadequate clinical response.

#### CHLORPROMAZINE HYDROCHLORIDE

Tab 10 mg	
Tab 25 mg	
Tab 100 mg	
Oral liq 10 mg per ml	
Inj 25 mg per ml, 2 ml ampoule	

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>CLOZAPINE</b>			
Tab 25 mg .....	13.37	50	Clozaril
	26.74	100	Clozaril
	6.69	50	Clopine
	13.37	100	Clopine
Tab 50 mg .....	8.67	50	Clopine
	17.33	100	Clopine
Tab 100 mg .....	34.65	50	Clozaril
	69.30	100	Clozaril
	17.33	50	Clopine
	34.65	100	Clopine
Tab 200 mg .....	34.65	50	Clopine
	69.30	100	Clopine
Oral liq 50 mg per ml .....	17.33	100 ml	Clopine
<b>HALOPERIDOL</b>			
Tab 500 mcg – 1% DV Oct-13 to 2016 .....	6.23	100	<b>Serenace</b>
Tab 1.5 mg – 1% DV Oct-13 to 2016 .....	9.43	100	<b>Serenace</b>
Tab 5 mg – 1% DV Oct-13 to 2016 .....	29.72	100	<b>Serenace</b>
Oral liq 2 mg per ml – 1% DV Oct-13 to 2016 .....	23.84	100 ml	<b>Serenace</b>
Inj 5 mg per ml, 1ml ampoule – 1% DV Oct-13 to 2016 .....	21.55	10	<b>Serenace</b>
<b>LEVOMEPRMAZINE</b>			
Tab 25 mg			
Tab 100 mg			
Inj 25 mg per ml, 1 ml ampoule			
<b>LITHIUM CARBONATE</b>			
Tab long-acting 400 mg			
Tab 250 mg – 1% DV Sep-12 to 2015 .....	34.30	500	<b>Lithicarb FC</b>
Tab 400 mg – 1% DV Sep-12 to 2015 .....	12.83	100	<b>Lithicarb FC</b>
Cap 250 mg – 1% DV Nov-11 to 2014 .....	9.42	100	<b>Douglas</b>
<b>OLANZAPINE</b>			
Tab 2.5 mg .....	2.00	28	Zypine
Tab 5 mg .....	3.85	28	Olanzine
			Zypine
Tab orodispersible 5 mg .....	6.36	28	Olanzine-D
			Zypine ODT
Tab 10 mg .....	6.35	28	Olanzine
			Zypine
Tab orodispersible 10 mg .....	8.76	28	Olanzine-D
			Zypine ODT
Inj 10 mg vial			
<i>(Olanzine-D Tab orodispersible 5 mg to be delisted 1 August 2014)</i>			
<i>(Olanzine Tab 10 mg to be delisted 1 August 2014)</i>			
<i>(Olanzine-D Tab orodispersible 10 mg to be delisted 1 August 2014)</i>			
<b>PERICYZINE</b>			
Tab 2.5 mg			
Tab 10 mg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>QUETIAPINE</b>			
Tab 25 mg .....	7.00	60	Dr Reddy's Quetiapine
			Seroquel
	10.50	90	Quetapel
Tab 100 mg .....	14.00	60	Seroquel
	21.00	90	Dr Reddy's Quetiapine
			Quetapel
Tab 200 mg .....	24.00	60	Dr Reddy's Quetiapine
			Seroquel
	36.00	90	Quetapel
Tab 300 mg .....	40.00	60	Dr Reddy's Quetiapine
			Seroquel
	60.00	90	Quetapel
<b>RISPERIDONE – Some items restricted see terms below</b>			
Tab 0.5 mg .....	2.86	20	Risperdal
	3.51	60	Apo-Risperidone
			Dr Reddy's Risperidone
			Ridal
☞ Tab orodispersible 0.5 mg .....	21.42	28	Risperdal Quicklet
Tab 1 mg .....	6.00	60	Apo-Risperidone
			Dr Reddy's Risperidone
			Ridal
	16.92		Risperdal
☞ Tab orodispersible 1 mg .....	42.84	28	Risperdal Quicklet
Tab 2 mg .....	11.00	60	Apo-Risperidone
			Dr Reddy's Risperidone
			Ridal
	33.84		Risperdal
☞ Tab orodispersible 2 mg .....	85.71	28	Risperdal Quicklet
Tab 3 mg .....	15.00	60	Apo-Risperidone
			Dr Reddy's Risperidone
			Ridal
	50.78		Risperdal
Tab 4 mg .....	20.00	60	Apo-Risperidone
			Dr Reddy's Risperidone
			Ridal
	67.68		Risperdal
Oral liq 1 mg per ml .....	18.35	30 ml	Apo-Risperidone
			Risperon
	25.26		Risperdal

## ☞Restricted

### Acute situations

Both:

- 1 For a non-adherent patient on oral therapy with standard risperidone tablets or risperidone oral liquid; and
- 2 The patient is under direct supervision for administration of medicine.

### Chronic situations

Both:

- 1 The patient is unable to take standard risperidone tablets or oral liquid, or once stabilized refuses to take risperidone tablets or oral liquid; and
- 2 The patient is under direct supervision for administration of medicine.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>TRIFLUOPERAZINE HYDROCHLORIDE</b>			
Tab 1 mg			
Tab 2 mg			
Tab 5 mg			
<b>ZIPRASIDONE – Some items restricted see terms below</b>			
⚡ Cap 20 mg	87.88	60	Zeldox
⚡ Cap 40 mg	164.78	60	Zeldox
⚡ Cap 60 mg	247.17	60	Zeldox
⚡ Cap 80 mg	329.56	60	Zeldox
Inj 20 mg			
Inj 100 mg			
<b>➡ Restricted</b>			
1 Patient is suffering from schizophrenia or related psychoses; and			
2 Either:			
2.1 An effective dose of risperidone or quetiapine has been trialled and has been discontinued, or is in the process of being discontinued, because of unacceptable side effects; or			
2.2 An effective dose of risperidone or quetiapine has been trialled and has been discontinued, or is in the process of being discontinued, because of inadequate clinical response.			
<b>ZUCLOPENTHIXOL ACETATE</b>			
Inj 50 mg per ml, 1 ml ampoule			
Inj 50 mg per ml, 2 ml ampoule			
<b>ZUCLOPENTHIXOL HYDROCHLORIDE</b>			
Tab 10 mg	31.45	100	Clopixol
<b>Depot Injections</b>			
<b>FLUPENTHIXOL DECANOATE</b>			
Inj 20 mg per ml, 1 ml ampoule	13.14	5	Fluanxol
Inj 20 mg per ml, 2 ml ampoule	20.90	5	Fluanxol
Inj 100 mg per ml, 1 ml ampoule	40.87	5	Fluanxol
<b>FLUPHENAZINE DECANOATE</b>			
Inj 12.5 mg per 0.5 ml ampoule	17.60	5	Modecate
Inj 25 mg per ml, 1 ml ampoule	27.90	5	Modecate
Inj 100 mg per ml, 1 ml ampoule	154.50	5	Modecate
<b>HALOPERIDOL DECANOATE</b>			
Inj 50 mg per ml, 1 ml ampoule	28.39	5	Haldol
Inj 100 mg per ml, 1 ml ampoule	55.90	5	Haldol Concentrate
<b>OLANZAPINE – Restricted see terms on the next page</b>			
⚡ Inj 210 mg vial	280.00	1	Zyprexa Relprevv
⚡ Inj 300 mg vial	460.00	1	Zyprexa Relprevv
⚡ Inj 405 mg vial	560.00	1	Zyprexa Relprevv

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➡Restricted</b>			
<b>Initiation</b>			
<i>Re-assessment required after 12 months</i>			
Either:			
<ol style="list-style-type: none"> <li>The patient has had an initial Special Authority approval for risperidone depot injection or paliperidone depot injection; or</li> <li>All of the following: <ol style="list-style-type: none"> <li>The patient has schizophrenia; and</li> <li>The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and</li> <li>The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.</li> </ol> </li> </ol>			
<b>Continuation</b>			
<i>Re-assessment required after 12 months</i>			
The initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.			
<b>PALIPERIDONE – Restricted</b> see terms below			
⚡ Inj 25 mg syringe .....	194.25	1	Invega Sustenna
⚡ Inj 50 mg syringe .....	271.95	1	Invega Sustenna
⚡ Inj 75 mg syringe .....	357.42	1	Invega Sustenna
⚡ Inj 100 mg syringe .....	435.12	1	Invega Sustenna
⚡ Inj 150 mg syringe .....	435.12	1	Invega Sustenna
<b>➡Restricted</b>			
<b>Initiation</b>			
<i>Re-assessment required after 12 months</i>			
Either:			
<ol style="list-style-type: none"> <li>The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or</li> <li>All of the following: <ol style="list-style-type: none"> <li>The patient has schizophrenia or other psychotic disorder; and</li> <li>The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and</li> <li>The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.</li> </ol> </li> </ol>			
<b>Continuation</b>			
<i>Re-assessment required after 12 months</i>			
The initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.			
<b>PIPOTHAZINE PALMITATE</b>			
Inj 50 mg per ml, 1 ml ampoule			
Inj 50 mg per ml, 2 ml ampoule			
<b>RISPERIDONE – Restricted</b> see terms below			
⚡ Inj 25 mg vial .....	135.98	1	Risperdal Consta
⚡ Inj 37.5 mg vial .....	178.71	1	Risperdal Consta
⚡ Inj 50 mg vial .....	217.56	1	Risperdal Consta
<b>➡Restricted</b>			
<b>Initiation</b>			
<i>Re-assessment required after 12 months</i>			
Either:			
<ol style="list-style-type: none"> <li>The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or</li> <li>All of the following: <ol style="list-style-type: none"> <li>The patient has schizophrenia or other psychotic disorder; and</li> <li>The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and</li> </ol> </li> </ol>			
			continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

### Continuation

#### Re-assessment required after 12 months

The initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

#### ZUCLOPENTHIXOL DECANOATE

Inj 200 mg per ml, 1 ml ampoule	19.80	5	Clopixol
---------------------------------	-------	---	----------

### Anxiolytics

#### ALPRAZOLAM

Tab 1 mg  
Tab 250 mcg  
Tab 500 mcg

#### BUSPIRONE HYDROCHLORIDE

Tab 5 mg	28.00	100	Pacific Buspirone
Tab 10 mg	17.00	100	Pacific Buspirone

#### CLONAZEPAM

Tab 500 mcg	6.68	100	Paxam
Tab 2 mg	12.75	100	Paxam

#### DIAZEPAM

Tab 2 mg	11.44	500	Arrow-Diazepam
Tab 5 mg	13.71	500	Arrow-Diazepam

#### LORAZEPAM

Tab 1 mg	19.82	250	Ativan
Tab 2.5 mg	13.49	100	Ativan

#### OXAZEPAM

Tab 10 mg  
Tab 15 mg

### Multiple Sclerosis Treatments

GLATIRAMER ACETATE – **Restricted** see terms below

⚡ Inj 20 mg per ml, 1 ml syringe

#### ➡Restricted

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessments Committee

INTERFERON BETA-1-ALPHA – **Restricted** see terms below

⚡ Inj 6 million iu in 0.5 ml pen

⚡ Inj 6 million iu in 0.5 ml syringe

⚡ Inj 6 million iu vial

#### ➡Restricted

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessments Committee

INTERFERON BETA-1-BETA – **Restricted** see terms below

⚡ Inj 8 million iu per ml, 1 ml vial

#### ➡Restricted

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessments Committee

Price  
(ex man. excl. GST)  
\$ Per Brand or  
Generic  
Manufacturer

## Sedatives and Hypnotics

### CHLORAL HYDRATE

Oral liq 100 mg per ml  
Oral liq 200 mg per ml

### LORMETAZEPAM – **Restricted**: For continuation only

➡ Tab 1 mg

### MELATONIN – **Restricted** see terms below

⚡ Tab modified-release 2 mg

*e.g. Circadin*

⚡ Tab 1 mg

⚡ Tab 2 mg

⚡ Tab 3 mg

⚡ Cap 2 mg

⚡ Cap 3 mg

➡ **Restricted**

For in hospital use only. For the treatment of insomnia where benzodiazepines and zopiclone are contraindicated.

### MIDAZOLAM

Tab 7.5 mg .....	40.00	100	Hypnovel
Oral liq 2 mg per ml			
Inj 1 mg per ml, 5 ml ampoule .....	10.00	10	Pfizer
	10.75		Hypnovel
Inj 5 mg per ml, 3 ml ampoule .....	11.90	5	Hypnovel Pfizer

### NITRAZEPAM

Tab 5 mg

### PHENOBARBITONE

Inj 200 mg per ml, 1 ml ampoule

### TEMAZEPAM

Tab 10 mg – 1% DV Nov-11 to 2014 .....	1.27	25	<b>Normison</b>
--	------	----	-----------------

### TRIAZOLAM – **Restricted**: For continuation only

➡ Tab 125 mcg

➡ Tab 250 mcg

### ZOPICLONE

Tab 7.5 mg – 1% DV Jan-12 to 2014 .....	1.90	30	<b>Apo-Zopiclone</b>
---	------	----	----------------------

## Stimulants / ADHD Treatments

### ATOMOXETINE – **Restricted** see terms on the next page

⚡ Cap 10 mg .....	107.03	28	Strattera
⚡ Cap 18 mg .....	107.03	28	Strattera
⚡ Cap 25 mg .....	107.03	28	Strattera
⚡ Cap 40 mg .....	107.03	28	Strattera
⚡ Cap 60 mg .....	107.03	28	Strattera
⚡ Cap 80 mg .....	139.11	28	Strattera
⚡ Cap 100 mg .....	139.11	28	Strattera

Price  
(ex man. excl. GST)  
\$ Per Brand or  
Generic  
Manufacturer

➔ **Restricted**

All of the following:

- 1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder) diagnosed according to DSM-IV or ICD 10 criteria; and
- 2 Once-daily dosing; and
- 3 Any of the following:
  - 3.1 Treatment with a subsidised formulation of a stimulant has resulted in the development or worsening of serious adverse reactions or where the combination of subsidised stimulant treatment with another agent would pose an unacceptable medical risk; or
  - 3.2 Treatment with a subsidised formulation of a stimulant has resulted in worsening of co-morbid substance abuse or there is a significant risk of diversion with subsidised stimulant therapy; or
  - 3.3 An effective dose of a subsidised formulation of a stimulant has been trialled and has been discontinued because of inadequate clinical response; or
  - 3.4 Treatment with a subsidised formulation of a stimulant is considered inappropriate because the patient has a history of psychoses or has a first-degree relative with schizophrenia; and
- 4 The patient will not be receiving treatment with atomoxetine in combination with a subsidised formulation of a stimulant, except for the purposes of transitioning from subsidised stimulant therapy to atomoxetine.

Note: A "subsidised formulation of a stimulant" refers to currently listed methylphenidate hydrochloride tablet formulations (immediate-release, sustained-release and extended-release) or dexamphetamine sulphate tablets.

**CAFFEINE**

Tab 100 mg

DEXAMPHETAMINE SULPHATE – **Restricted** see terms below

⚡ Tab 5 mg – 1% DV Mar-13 to 2015 ..... 16.50 100 **PSM**

➔ **Restricted**

**ADHD**

Paediatrician or psychiatrist

Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria

**Narcolepsy**

Neurologist or respiratory specialist

Patient suffers from narcolepsy

METHYLPHENIDATE HYDROCHLORIDE – **Restricted** see terms on the next page

⚡ Tab extended-release 18 mg .....	58.96	30	Concerta
⚡ Tab extended-release 27 mg .....	65.44	30	Concerta
⚡ Tab extended-release 36 mg .....	71.93	30	Concerta
⚡ Tab extended-release 54 mg .....	86.24	30	Concerta
⚡ Tab immediate-release 5 mg .....	3.20	30	Rubifen
⚡ Tab immediate-release 10 mg .....	3.00	30	Ritalin
			Rubifen
⚡ Tab immediate-release 20 mg .....	7.85	30	Rubifen
⚡ Tab sustained-release 20 mg .....	10.95	30	Rubifen SR
	50.00	100	Ritalin SR
⚡ Cap modified-release 10 mg .....	19.50	30	Ritalin LA
⚡ Cap modified-release 20 mg .....	25.50	30	Ritalin LA
⚡ Cap modified-release 30 mg .....	31.90	30	Ritalin LA
⚡ Cap modified-release 40 mg .....	38.25	30	Ritalin LA

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## ➔Restricted

### ADHD (immediate-release and sustained-release formulations)

Paediatrician or psychiatrist

Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria

### Narcolepsy (immediate-release and sustained-release formulations)

Neurologist or respiratory specialist

Patient suffers from narcolepsy

### Extended-release and modified-release formulations

Paediatrician or psychiatrist

Both:

- 1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria; and
- 2 Either:
  - 2.1 Patient is taking a currently listed formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
  - 2.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

MODAFINIL – **Restricted** see terms below

⚡ Tab 100 mg

## ➔Restricted

Neurologist or respiratory specialist

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
  - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
  - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
  - 3.1 An effective dose of a listed formulation of methylphenidate or dexamphetamine has been trialled and discontinued because of intolerable side effects; or
  - 3.2 Methylphenidate and dexamphetamine are contraindicated.

## Treatments for Dementia

### DONEPEZIL HYDROCHLORIDE

Tab 5 mg .....	7.71	90	Donepezil-Rex
Tab 10 mg .....	14.06	90	Donepezil-Rex

## Treatments for Substance Dependence

### BUPRENORPHINE WITH NALOXONE – **Restricted** see terms below

⚡ Tab 2 mg with naloxone 0.5 mg .....	57.40	28	Suboxone
⚡ Tab 8 mg with naloxone 2 mg .....	166.00	28	Suboxone

## ➔Restricted

### Detoxification

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Prescriber works in an opioid treatment service approved by the Ministry of Health.

### Maintenance treatment

All of the following:

continued...



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
1 Patient is opioid dependent; and			
2 Patient will not be receiving methadone; and			
3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and			
4 Prescriber works in an opioid treatment service approved by the Ministry of Health.			
<b>BUPROPION HYDROCHLORIDE</b>			
Tab modified-release 150 mg – 1% DV Oct-13 to 2016.....	4.97	30	<b>Zyban</b>
<b>DISULFIRAM</b>			
Tab 200 mg .....	24.30	100	Antabuse
<b>NALTREXONE HYDROCHLORIDE – Restricted</b> see terms below			
⚡ Tab 50 mg – 1% DV Sep-13 to 2016 .....	76.00	30	<b>Naltraccord</b>
➡ <b>Restricted</b>			
<b>Alcohol dependence</b>			
Both:			
1 Patient is currently enrolled, or is planned to be enrolled, in a recognised comprehensive treatment programme for alcohol dependence; and			
2 Naltrexone is to be prescribed by, or on the recommendation of, a physician working in an Alcohol and Drug Service.			
<b>Constipation</b>			
For the treatment of opioid-induced constipation			
<b>NICOTINE – Some items restricted</b> see terms below			
Gum 2 mg – 1% DV Apr-14 to 2017 .....	36.47	384	<b>Habitrol (Classic)</b> <b>Habitrol (Fruit)</b> <b>Habitrol (Mint)</b>
Gum 4 mg – 1% DV Apr-14 to 2017 .....	42.04	384	<b>Habitrol (Classic)</b> <b>Habitrol (Fruit)</b> <b>Habitrol (Mint)</b>
Patch 7 mg per 24 hours – 1% DV Apr-14 to 2017 .....	18.13	28	<b>Habitrol</b>
Patch 14 mg per 24 hours – 1% DV Apr-14 to 2017 .....	18.81	28	<b>Habitrol</b>
Patch 21 mg per 24 hours – 1% DV Apr-14 to 2017 .....	19.14	28	<b>Habitrol</b>
Lozenge 1 mg – 1% DV Apr-14 to 2017 .....	19.94	216	<b>Habitrol</b>
Lozenge 2 mg – 1% DV Apr-14 to 2017 .....	24.27	216	<b>Habitrol</b>
⚡ Soln for inhalation 15 mg cartridge			<i>e.g. Nicorette Inhalator</i>
➡ <b>Restricted</b>			
Any of the following:			
1 For perioperative use in patients who have a 'nil by mouth' instruction; or			
2 For use within mental health inpatient units; or			
3 For acute use in agitated patients who are unable to leave the hospital facilities.			
<b>VARENICLINE – Restricted</b> see terms below			
⚡ Tab 0.5 mg × 11 and 1 mg × 14 .....	60.48	25	Champix
⚡ Tab 1 mg .....	67.74	28	Champix
	135.48	56	Champix
➡ <b>Restricted</b>			
All of the following:			
1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and			
2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and			
3 Either:			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
- 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not used funded varenicline in the last 12 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 3 months' funded varenicline in a 12 month period.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Chemotherapeutic Agents</b>			
<b>Alkylating Agents</b>			
BUSULFAN			
Tab 2 mg .....	59.50	100	Myleran
Inj 6 mg per ml, 10 ml ampoule			
CARMUSTINE			
Inj 100 mg vial			
CHLORAMBUCIL			
Tab 2 mg			
CYCLOPHOSPHAMIDE			
Tab 50 mg .....	79.00	50	Endoxan
	158.00	100	Procytox
Inj 1 g vial – 1% DV Nov-11 to 2014 .....	26.70	1	<b>Endoxan</b>
Inj 2 g vial – 1% DV Nov-11 to 2014 .....	56.90	1	<b>Endoxan</b>
IFOSFAMIDE			
Inj 1 g vial .....	96.00	1	Holoxan
Inj 2 g vial .....	180.00	1	Holoxan
LOMUSTINE			
Cap 10 mg – 1% DV Sep-11 to 2014 .....	132.59	20	<b>Ceenu</b>
Cap 40 mg – 1% DV Sep-11 to 2014 .....	399.15	20	<b>Ceenu</b>
MELPHALAN			
Tab 2 mg			
Inj 50 mg vial			
THIOTEPA			
Inj 15 mg vial			
<b>Anthracyclines and Other Cytotoxic Antibiotics</b>			
BLEOMYCIN SULPHATE			
Inj 15,000 iu (10 mg) vial			
DACTINOMYCIN [ACTINOMYCIN D]			
Inj 0.5 mg vial			
DAUNORUBICIN			
Inj 2 mg per ml, 10 ml vial – 1% DV Aug-13 to 2016 .....	118.72	1	<b>Pfizer</b>
DOXORUBICIN HYDROCHLORIDE			
Note: DV limit applies to all 50 mg presentations of doxorubicin hydrochloride.			
Inj 2 mg per ml, 5 ml vial			
Inj 2 mg per ml, 25 ml vial – 1% DV Mar-13 to 2015 .....	17.00	1	<b>Arrow-Doxorubicin</b>
Inj 50 mg vial			
Inj 2 mg per ml, 50 ml vial			
Inj 2 mg per ml, 100 ml vial – 1% DV Mar-13 to 2015 .....	65.00	1	<b>Arrow-Doxorubicin</b>

# ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>EPIRUBICIN HYDROCHLORIDE</b>			
Inj 2 mg per ml, 5 ml vial .....	25.00	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial – 1% DV Aug-12 to 2015 .....	39.38	1	<b>DBL Epirubicin Hydrochloride</b>
Inj 2 mg per ml, 50 ml vial – 1% DV Aug-12 to 2015 .....	58.20	1	<b>DBL Epirubicin Hydrochloride</b>
Inj 2 mg per ml, 100 ml vial – 1% DV Aug-12 to 2015 .....	94.50	1	<b>DBL Epirubicin Hydrochloride</b>
<b>IDARUBICIN HYDROCHLORIDE</b>			
Cap 5 mg .....	115.00	1	Zavedos
Cap 10 mg .....	144.50	1	Zavedos
Inj 5 mg vial – 1% DV Sep-12 to 2015 .....	100.00	1	<b>Zavedos</b>
Inj 10 mg vial – 1% DV Sep-12 to 2015 .....	200.00	1	<b>Zavedos</b>
<b>MITOMYCIN C</b>			
Inj 5 mg vial – 1% DV Oct-13 to 2016 .....	79.75	1	<b>Arrow</b>
<b>MITOZANTRONE</b>			
Inj 2 mg per ml, 5 ml vial .....	110.00	1	Mitozantrone Ebewe
Inj 2 mg per ml, 10 ml vial .....	100.00	1	Mitozantrone Ebewe
Inj 2 mg per ml, 12.5 ml vial .....	407.50	1	Onkotrone
<b>Antimetabolites</b>			
<b>CAPECITABINE</b>			
Tab 150 mg .....	115.00	60	Xeloda
Tab 500 mg .....	705.00	120	Xeloda
<b>CLADRIBINE</b>			
Inj 2 mg per ml, 5 ml vial .....			
Inj 1 mg per ml, 10 ml vial .....	5,249.72	7	Leustatin
<b>CYTARABINE</b>			
Inj 20 mg per ml, 5 ml vial – 1% DV Nov-13 to 2016 .....	55.00	5	<b>Pfizer</b>
Inj 20 mg per ml, 25 ml vial .....	18.15	1	Pfizer
Inj 100 mg per ml, 10 ml vial – 1% DV Nov-13 to 2016 .....	8.83	1	<b>Pfizer</b>
Inj 100 mg per ml, 20 ml vial – 1% DV Nov-13 to 2016 .....	17.65	1	<b>Pfizer</b>
<b>FLUDARABINE PHOSPHATE</b>			
Tab 10 mg – 1% DV Jun-12 to 2015 .....	433.50	20	<b>Fludara Oral</b>
Inj 50 mg vial – 1% DV Sep-11 to 2014 .....	525.00	5	<b>Fludarabine Ebewe</b>
<b>FLUOROURACIL</b>			
Inj 25 mg per ml, 100 ml vial .....	13.55	1	Hospira
Inj 50 mg per ml, 10 ml vial .....	26.25	5	Fluorouracil Ebewe
Inj 50 mg per ml, 20 ml vial .....	7.50	1	Fluorouracil Ebewe
Inj 50 mg per ml, 50 ml vial .....	18.00	1	Fluorouracil Ebewe
Inj 50 mg per ml, 100 ml vial .....	34.50	1	Fluorouracil Ebewe

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>GEMCITABINE</b>			
Inj 10 mg per ml, 100 ml vial .....	62.50	1	Gemcitabine Ebewe
Inj 10 mg per ml, 20 ml vial .....	12.50	1	Gemcitabine Ebewe
Inj 200 mg vial .....	12.50	1	Gemcitabine Actavis 200
Inj 1 g vial .....	62.50	1	DBL Gemcitabine Gemcitabine Actavis 1000
<i>(Gemcitabine Actavis 200 Inj 200 mg vial to be delisted 1 July 2014)</i>			
<i>(Gemcitabine Actavis 1000 Inj 1 g vial to be delisted 1 July 2014)</i>			
<b>MERCAPTOPURINE</b>			
Tab 50 mg – 1% DV Oct-13 to 2016 .....	49.41	25	Puri-nethol
<b>METHOTREXATE</b>			
Tab 2.5 mg – 1% DV Jun-14 to 2015 .....	3.82	30	Trexate
Tab 10 mg – 1% DV Jun-14 to 2015 .....	26.25	50	Trexate
Inj 2.5 mg per ml, 2 ml vial			
Inj 7.5 mg prefilled syringe – 1% DV Jan-14 to 2016 .....	17.19	1	Methotrexate Sandoz
Inj 10 mg prefilled syringe – 1% DV Jan-14 to 2016 .....	17.25	1	Methotrexate Sandoz
Inj 15 mg prefilled syringe – 1% DV Jan-14 to 2016 .....	17.38	1	Methotrexate Sandoz
Inj 20 mg prefilled syringe – 1% DV Jan-14 to 2016 .....	17.50	1	Methotrexate Sandoz
Inj 25 mg prefilled syringe – 1% DV Jan-14 to 2016 .....	17.63	1	Methotrexate Sandoz
Inj 30 mg prefilled syringe – 1% DV Jan-14 to 2016 .....	17.75	1	Methotrexate Sandoz
Inj 25 mg per ml, 2 ml vial – 1% DV Sep-13 to 2016 .....	20.20	5	Hospira
Inj 25 mg per ml, 20 ml vial – 1% DV Sep-13 to 2016 .....	27.78	1	Hospira
Inj 100 mg per ml, 10 ml vial – 1% DV Nov-08 to 2014 .....	25.00	1	Methotrexate Ebewe
Inj 100 mg per ml, 50 ml vial – 1% DV Nov-08 to 2014 .....	125.00	1	Methotrexate Ebewe
<b>THIOGUANINE</b>			
Tab 40 mg			

## Other Cytotoxic Agents

<b>AMSACRINE</b>			
Inj 50 mg per ml, 1.5 ml ampoule			
<b>ANAGRELIDE HYDROCHLORIDE</b>			
Cap 0.5 mg			
<b>ARSENIC TRIOXIDE</b>			
Inj 1 mg per ml, 10 ml vial .....	4,817.00	10	AFT
<b>BORTEZOMIB – Restricted</b> see terms below			
⚡ Inj 1 mg vial .....	540.70	1	Velcade
⚡ Inj 3.5 mg vial .....	1,892.50	1	Velcade

### ➡ Restricted

**Initiation - treatment naive multiple myeloma/amyloidosis**

Both:

- 1 Either:
  - 1.1 The patient has treatment-naive symptomatic multiple myeloma; or
  - 1.2 The patient has treatment-naive symptomatic systemic AL amyloidosis \*; and
- 2 Maximum of 9 treatment cycles.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Note: Indications marked with \* are Unapproved Indications.

## Initiation - relapsed/refractory multiple myeloma/amyloidosis

All of the following:

- 1 Either:
  - 1.1 The patient has relapsed or refractory multiple myeloma; or
  - 1.2 The patient has relapsed or refractory systemic AL amyloidosis \*; and
- 2 The patient has received only one prior front line chemotherapy for multiple myeloma or amyloidosis; and
- 3 The patient has not had prior publicly funded treatment with bortezomib; and
- 4 Maximum of 4 treatment cycles.

Note: Indications marked with \* are Unapproved Indications.

## Continuation - relapsed/refractory multiple myeloma/amyloidosis

Both:

- 1 The patient's disease obtained at least a partial response from treatment with bortezomib at the completion of cycle 4; and
- 2 Maximum of 4 further treatment cycles (making a total maximum of 8 consecutive treatment cycles).

Notes: Responding relapsed/refractory multiple myeloma patients should receive no more than 2 additional cycles of treatment beyond the cycle at which a confirmed complete response was first achieved. A line of therapy is considered to comprise either:

- 1 A known therapeutic chemotherapy regimen and supportive treatments; or
- 2 A transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments.

Refer to datasheet for recommended dosage and number of doses of bortezomib per treatment cycle.

### COLASPASE [L-ASPARAGINASE]

Inj 10,000 iu vial .....	102.32	1	Leunase
--------------------------	--------	---	---------

### DACARBAZINE

Inj 200 mg vial – 1% DV Oct-13 to 2016 .....	51.84	1	Hospira
--	-------	---	---------

### ETOPOSIDE

Cap 50 mg .....	340.73	20	Vepesid
Cap 100 mg .....	340.73	10	Vepesid
Inj 20 mg per ml, 5 ml vial .....	25.00	1	Hospira

### ETOPOSIDE (AS PHOSPHATE)

Inj 100 mg vial – 1% DV Sep-11 to 2014 .....	40.00	1	Etopophos
--	-------	---	-----------

### HYDROXYUREA

Cap 500 mg .....	31.76	100	Hydrea
------------------	-------	-----	--------

### IRINOTECAN HYDROCHLORIDE

Inj 20 mg per ml, 2 ml vial – 1% DV Nov-12 to 2015 .....	9.34	1	Irinotecan Actavis 40
Inj 20 mg per ml, 5 ml vial – 1% DV Nov-12 to 2015 .....	23.34	1	Irinotecan Actavis 100

### PEGASPARGASE – Restricted see terms below

☞ Inj 750 iu per ml, 5 ml vial .....	3,005.00	1	Oncaspar
--------------------------------------	----------	---	----------

### ☞ Restricted

#### Newly diagnosed ALL

Limited to 12 months' treatment

All of the following:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and
- 3 Treatment is with curative intent.

#### Relapsed ALL

Limited to 12 months' treatment

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued. . .

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and
- 3 Treatment is with curative intent.

#### PENTOSTATIN [DEOXYCOFORMYCIN]

Inj 10 mg vial

#### PROCARBAZINE HYDROCHLORIDE

Cap 50 mg .....	498.00	50	Natulan
-----------------	--------	----	---------

#### TEMOZOLOMIDE – **Restricted** see terms below

⚡ Cap 5 mg – 1% DV Sep-13 to 2016.....	8.00	5	<b>Temaccord</b>
⚡ Cap 20 mg – 1% DV Sep-13 to 2016.....	36.00	5	<b>Temaccord</b>
⚡ Cap 100 mg – 1% DV Sep-13 to 2016.....	175.00	5	<b>Temaccord</b>
⚡ Cap 250 mg – 1% DV Sep-13 to 2016.....	410.00	5	<b>Temaccord</b>

#### ➡ **Restricted**

All of the following:

- 1 Either:
  - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
  - 1.2 Patient has newly diagnosed anaplastic astrocytoma\*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of six cycles of 5 days treatment, at a maximum dose of 200 mg/m<sup>2</sup>.

Notes: Indication marked with a \* is an Unapproved Indication. Studies of temozolomide show that its benefit is predominantly in those patients with a good performance status (WHO grade 0 or 1 or Karnofsky score >80), and in patients who have had at least a partial resection of the tumour.

#### THALIDOMIDE – **Restricted** see terms below

⚡ Cap 50 mg .....	504.00	28	Thalomid
⚡ Cap 100 mg .....	1,008.00	28	Thalomid

#### ➡ **Restricted**

#### Initiation

Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis\*; or
- 3 The patient has erythema nodosum leprosum.

#### Continuation

Patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with \* is an Unapproved Indication

#### TRETINOIN

Cap 10 mg .....	435.90	100	Vesanoid
-----------------	--------	-----	----------

### Platinum Compounds

#### CARBOPLATIN

Inj 10 mg per ml, 5 ml vial .....	20.00	1	Carboplatin Ebewe
Inj 10 mg per ml, 15 ml vial – 1% DV Jan-13 to 2015.....	19.50	1	<b>Carbaccord</b>
Inj 10 mg per ml, 45 ml vial – 1% DV Jan-13 to 2015.....	48.50	1	<b>Carbaccord</b>
Inj 10 mg per ml, 100 ml vial .....	105.00	1	Carboplatin Ebewe

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>CISPLATIN</b>			
Inj 1 mg per ml, 50 ml vial .....	15.00	1	Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial .....	21.00	1	Cisplatin Ebewe
<b>OXALIPLATIN</b>			
Inj 50 mg vial – 1% DV Aug-12 to 2015 .....	15.32	1	<b>Oxaliplatin Actavis 50</b>
Inj 100 mg vial – 1% DV Aug-12 to 2015 .....	25.01	1	<b>Oxaliplatin Actavis 100</b>

### Protein-Tyrosine Kinase Inhibitors

**DASATINIB – Restricted** see terms below

☞ Tab 20 mg .....	3,774.06	60	Sprycel
☞ Tab 50 mg .....	6,214.20	60	Sprycel
☞ Tab 70 mg .....	7,692.58	60	Sprycel
☞ Tab 100 mg .....	6,214.20	30	Sprycel

☞ **Restricted**

For use in patients with approval from the CML/GIST Co-ordinator

**ERLOTINIB – Restricted** see terms below

☞ Tab 100 mg .....	1,133.00	30	Tarceva
☞ Tab 150 mg .....	1,700.00	30	Tarceva

☞ **Restricted**

#### Initiation

*Re-assessment required after 3 months*

Either:

- 1 All of the following:
  - 1.1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
  - 1.2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
  - 1.3 Either:
    - 1.3.1 Patient is treatment naive; or
    - 1.3.2 Both:
      - 1.3.2.1 Patient has documented disease progression following treatment with first line platinum based chemotherapy; and
      - 1.3.2.2 Patient has not received prior treatment with gefitinib; and
  - 1.4 Erlotinib is to be given for a maximum of 3 months, or
- 2 The patient received funded erlotinib prior to 31 December 2013 and radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

#### Continuation

*Re-assessment required after 6 months*

Radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

**GEFITINIB – Restricted** see terms below

☞ Tab 250 mg .....	1,700.00	30	Iressa
--------------------	----------	----	--------

☞ **Restricted**

#### Initiation

*Re-assessment required after 3 months*

Both

- 1 Patient has treatment naive locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase.

#### Continuation

*Re-assessment required after 6 months*

Radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
IMATINIB MESILATE			
⚡ Tab 100 mg .....	2,400.00	60	Glivec

➡ **Restricted**

For use in patients with approval from the CML/GIST Co-ordinator

Cap 100 mg – 1% DV Jul-14 to 2017 ..... 298.90 60 **Imatinib-AFT**

Note: Imatinib-AFT is not a registered for the treatment of Gastro Intestinal Stromal Tumours (GIST). The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA0643 in Section B of the Pharmaceutical Schedule.

LAPATINIB – **Restricted** see terms below

⚡ Tab 250 mg ..... 1,899.00 70 Tykerb

➡ **Restricted**

**Initiation**

*Re-assessment required after 12 months*

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The patient has not previously received trastuzumab treatment for HER 2 positive metastatic breast cancer; and
  - 1.3 Lapatinib not to be given in combination with trastuzumab; and
  - 1.4 Lapatinib to be discontinued at disease progression; or
- 2 All of the following:
  - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 2.2 The patient started trastuzumab for metastatic breast cancer but discontinued trastuzumab within 3 months of starting treatment due to intolerance; and
  - 2.3 The cancer did not progress whilst on trastuzumab; and
  - 2.4 Lapatinib not to be given in combination with trastuzumab; and
  - 2.5 Lapatinib to be discontinued at disease progression.

**Continuation**

*Re-assessment required after 12 months*

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

PAZOPANIB – **Restricted** see terms below

⚡ Tab 200 mg ..... 1,334.70 30 Votrient  
 ⚡ Tab 400 mg ..... 2,669.40 30 Votrient

➡ **Restricted**

**Initiation**

*Re-assessment required after 3 months*

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
- 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
- 5 The patient has intermediate or poor prognosis defined as any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal; or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of  $\leq$  70; or
  - 5.6  $\geq$  2 sites of organ metastasis.

## Continuation

*Re-assessment required after 3 months*

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

SUNITINIB – **Restricted** see terms below

⚡ Cap 12.5 mg .....	2,315.38	28	Sutent
⚡ Cap 25 mg .....	4,630.77	28	Sutent
⚡ Cap 50 mg .....	9,261.54	28	Sutent

## ➡Restricted

*Re-assessment required after 3 months*

Initiation - RCC

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
  - 2.4 Both:
    - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
    - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
- 5 The patient has intermediate or poor prognosis defined as any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal; or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of  $\leq$  70; or
  - 5.6  $\geq$  2 sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

## Continuation - RCC

*Re-assessment required after 3 months*

Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued. . .

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

#### Initiation - GIST

*Re-assessment required after 3 months*

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
  - 2.1 The patient's disease has progressed following treatment with imatinib; or
  - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

#### Continuation - GIST

*Re-assessment required after 6 months*

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
  - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
  - 1.2 The patient has had a partial response (a decrease in size of  $\geq 10\%$  or decrease in tumour density in Hounsfield Units (HU) of  $\geq 15\%$  on CT and no new lesions and no obvious progression of non-measurable disease); or
  - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: RCC - Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

GIST - It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of  $\geq 10\%$  and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

### Taxanes

#### DOCETAXEL

Inj 10 mg per ml, 2 ml vial – 1% DV May-13 to 2014 .....	48.75	1	<b>Docetaxel Sandoz</b>
Inj 10 mg per ml, 8 ml vial – 1% DV May-13 to 2014 .....	195.00	1	<b>Docetaxel Sandoz</b>

#### PACLITAXEL

Inj 6 mg per ml, 5 ml vial – 1% DV Oct-08 to 2014 .....	137.50	5	<b>Paclitaxel Ebewe</b>
Inj 6 mg per ml, 16.7 ml vial – 1% DV Oct-08 to 2014 .....	91.67	1	<b>Paclitaxel Actavis</b>
			<b>Paclitaxel Ebewe</b>
Inj 6 mg per ml, 25 ml vial – 1% DV Oct-08 to 2014 .....	137.50	1	<b>Anzatax</b>
			<b>Paclitaxel Actavis</b>
			<b>Paclitaxel Ebewe</b>
Inj 6 mg per ml, 50 ml vial – 1% DV Oct-08 to 2014 .....	275.00	1	<b>Anzatax</b>
			<b>Paclitaxel Actavis</b>
			<b>Paclitaxel Ebewe</b>
Inj 6 mg per ml, 100 ml vial – 1% DV Oct-08 to 2014 .....	550.00	1	<b>Paclitaxel Ebewe</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Treatment of Cytotoxic-Induced Side Effects</b>			
<b>CALCIUM FOLINATE</b>			
Tab 15 mg – 1% DV Nov-11 to 2014 .....	82.45	10	<b>DBL Leucovorin Calcium</b>
Inj 3 mg per ml, 1 ml ampoule			
Inj 10 mg per ml, 5 ml ampoule – 1% DV Sep-08 to 2014 .....	24.50	5	<b>Calcium Folate Ebewe</b>
Inj 10 mg per ml, 10 ml vial – 1% DV Sep-08 to 2014 .....	9.75	1	<b>Calcium Folate Ebewe</b>
Inj 10 mg per ml, 30 ml vial – 1% DV Sep-08 to 2014 .....	30.00	1	<b>Calcium Folate Ebewe</b>
Inj 10 mg per ml, 100 ml vial – 1% DV Sep-08 to 2014 .....	90.00	1	<b>Calcium Folate Ebewe</b>
<b>MESNA</b>			
Tab 400 mg – 1% DV Oct-13 to 2016 .....	227.50	50	<b>Uromitexan</b>
Tab 600 mg – 1% DV Oct-13 to 2016 .....	339.50	50	<b>Uromitexan</b>
Inj 100 mg per ml, 4 ml ampoule – 1% DV Oct-13 to 2016 .....	148.05	15	<b>Uromitexan</b>
Inj 100 mg per ml, 10 ml ampoule – 1% DV Oct-13 to 2016 .....	339.90	15	<b>Uromitexan</b>
<b>Vinca Alkaloids</b>			
<b>VINBLASTINE SULPHATE</b>			
Inj 1 mg per ml, 10 ml vial .....	137.50	5	<b>Hospira</b>
<b>VINCISTINE SULPHATE</b>			
Inj 1 mg per ml, 1 ml vial – 1% DV Sep-13 to 2016 .....	64.80	5	<b>Hospira</b>
Inj 1 mg per ml, 2 ml vial – 1% DV Sep-13 to 2016 .....	69.60	5	<b>Hospira</b>
<b>VINOELBINE</b>			
Inj 10 mg per ml, 1 ml vial – 1% DV Sep-12 to 2015 .....	12.85	1	<b>Navelbine</b>
Inj 10 mg per ml, 5 ml vial – 1% DV Sep-12 to 2015 .....	64.25	1	<b>Navelbine</b>
<b>Endocrine Therapy</b>			
<b>BICALUTAMIDE – Restricted</b> see terms below			
☞ Tab 50 mg – 1% DV Nov-11 to 2014 .....	10.00	28	<b>Bicalaccord</b>
☞ <b>Restricted</b>			
For the treatment of advanced prostate cancer			
<b>FLUTAMIDE</b>			
Tab 250 mg .....	55.00	100	<b>Flutamin</b>
<b>MEGESTROL ACETATE</b>			
Tab 160 mg – 1% DV Jan-13 to 2015 .....	51.55	30	<b>Apo-Megestrol</b>
<b>OCTREOTIDE – Some items restricted</b> see terms on the next page			
Inj 50 mcg per ml, 1 ml ampoule – 1% DV May-12 to 2014 .....	19.24	5	<b>Octreotide MaxRx</b>
Inj 100 mcg per ml, 1 ml ampoule – 1% DV May-12 to 2014 .....	36.38	5	<b>Octreotide MaxRx</b>
Inj 500 mcg per ml, 1 ml ampoule – 1% DV May-12 to 2014 .....	131.25	5	<b>Octreotide MaxRx</b>
☞ Inj 10 mg vial .....	1,772.50	1	<b>Sandostatin LAR</b>
☞ Inj 20 mg vial .....	2,358.75	1	<b>Sandostatin LAR</b>
☞ Inj 30 mg vial .....	2,951.25	1	<b>Sandostatin LAR</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➔Restricted</b>			
Note: restriction applies only to the long-acting formulations of octreotide			
<b>Malignant bowel obstruction</b>			
All of the following:			
1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and			
2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and			
3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.			
Note: Indications marked with * are Unapproved Indications			
<b>Initiation - acromegaly</b>			
<i>Re-assessment required after 3 months</i>			
Both:			
1 The patient has acromegaly; and			
2 Any of the following:			
2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or			
2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or			
2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.			
<b>Continuation - acromegaly</b>			
Both:			
1 IGF1 levels have decreased since starting octreotide; and			
2 The treatment remains appropriate and the patient is benefiting from treatment.			
Note: In patients with acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks.			
<b>Other indications</b>			
Any of the following:			
1 VIPomas and glucagonomas - for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or			
2 Both:			
2.1 Gastrinoma; and			
2.2 Either:			
2.2.1 Patient has failed surgery; or			
2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or			
3 Both:			
3.1 Insulinomas; and			
3.2 Surgery is contraindicated or has failed; or			
4 For pre-operative control of hypoglycaemia and for maintenance therapy; or			
5 Both:			
5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and			
5.2 Disabling symptoms not controlled by maximal medical therapy.			
<b>TAMOXIFEN CITRATE</b>			
Tab 10 mg .....	2.63	60	Genox
	17.50	100	Genox
Tab 20 mg – 1% DV Jun-11 to 2014.....	2.63	30	<b>Genox</b>
	8.75	100	<b>Genox</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Aromatase Inhibitors</b>			
ANASTROZOLE			
Tab 1 mg .....	26.55	30	Aremed DP-Anastrozole
EXEMESTANE			
Tab 25 mg – 1% DV Jun-11 to 2014.....	22.57	30	<b>Aromasin</b>
LETROZOLE			
Tab 2.5 mg – 1% DV Oct-12 to 2015 .....	4.85	30	<b>Letraccord</b>

## Immunosuppressants

### Calcineurin Inhibitors

CICLOSPORIN			
Cap 25 mg .....	44.63	50	Neoral
Cap 50 mg .....	88.91	50	Neoral
Cap 100 mg .....	177.81	50	Neoral
Oral liq 100 mg per ml – 1% DV Oct-12 to 2015.....	198.13	50 ml	<b>Neoral</b>
Inj 50 mg per ml, 5 ml ampoule – 1% DV Oct-12 to 2015 .....	276.30	10	<b>Sandimmun</b>
TACROLIMUS – <b>Restricted</b> see terms below			
☞ Cap 0.5 mg – 1% DV Nov-14 to 31 Oct 2018.....	85.60	100	<b>Tacrolimus Sandoz</b>
	214.00		Prograf
☞ Cap 1 mg – 1% DV Nov-14 to 31 Oct 2018.....	171.20	100	<b>Tacrolimus Sandoz</b>
	428.00		Prograf
☞ Cap 5 mg – 1% DV Nov-14 to 31 Oct 2018.....	428.00	50	<b>Tacrolimus Sandoz</b>
	1,070.00		Prograf
☞ Inj 5 mg per ml, 1 ml ampoule			
<i>(Prograf Cap 0.5 mg to be delisted 1 November 2014)</i>			
<i>(Prograf Cap 1 mg to be delisted 1 November 2014)</i>			
<i>(Prograf Cap 5 mg to be delisted 1 November 2014)</i>			

#### ☞ **Restricted**

For use in organ transplant recipients

### Fusion Proteins

ETANERCEPT – <b>Restricted</b> see terms below			
☞ Inj 25 mg vial .....	949.96	4	Enbrel
☞ Inj 50 mg autoinjector .....	1,899.92	4	Enbrel
☞ Inj 50 mg syringe .....	1,899.92	4	Enbrel

#### ☞ **Restricted**

#### Initiation - juvenile idiopathic arthritis

Rheumatologist or named specialist

*Re-assessment required after 4 months*

Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for adalimumab for juvenile idiopathic arthritis (JIA); and

1.2 Either:

1.2.1 The patient has experienced intolerable side effects from adalimumab; or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for JIA; or

2 All of the following:

- 2.1 Patient diagnosed with Juvenile Idiopathic Arthritis (JIA); and
- 2.2 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m<sup>2</sup> weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
- 2.5 Both:
  - 2.5.1 Either:
    - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
    - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
  - 2.5.2 Physician's global assessment indicating severe disease.

#### Continuation - juvenile idiopathic arthritis

Rheumatologist or named specialist

*Re-assessment required after 6 months*

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

#### Initiation - rheumatoid arthritis

Rheumatologist

*Re-assessment required after 6 months*

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe and active erosive rheumatoid arthritis for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulphasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
  - 2.5 Any of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of cyclosporin; or
- 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
  - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
  - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

## Continuation - rheumatoid arthritis

Rheumatologist

*Re-assessment required after 6 months*

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

## Initiation - ankylosing spondylitis

Rheumatologist

*Re-assessment required after 6 months*

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
  - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of an exercise regime supervised by a physiotherapist; and
- 2.5 Either:

continued...



Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or

2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and

2.6 Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of starting treatment.

Average normal chest expansion corrected for age and gender:

Age	Male	Female
18-24	7.0 cm	5.5 cm
25-34	7.5 cm	5.5 cm
35-44	6.5 cm	4.5 cm
45-54	6.0 cm	5.0 cm
55-64	5.5 cm	4.0 cm
65-74	4.0 cm	4.0 cm
75+	3.0 cm	2.5 cm

#### Continuation - ankylosing spondylitis

Rheumatologist

*Re-assessment required after 6 months*

All of the following:

- 1 Following 12 weeks of etanercept treatment, BASDAI has improved by 4 or more points from pre-treatment baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

#### Initiation - psoriatic arthritis

Rheumatologist

*Re-assessment required after 6 months*

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulphasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
- 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

## Continuation - psoriatic arthritis

Rheumatologist

*Re-assessment required after 6 months*

All of the following:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

## Initiation - plaque psoriasis, prior TNF use

Dermatologist

*Re-assessment required after 4 months*

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
  - 1.1 The patient has experienced intolerable side effects from adalimumab; or
  - 1.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; and
- 2 Patient must be reassessed for continuation after 3 doses.

## Initiation - plaque psoriasis, treatment-naïve

Dermatologist

*Re-assessment required after 4 months*

All of the following:

- 1 Either:
  - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, cyclosporin, or acitretin; and
- 3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

### Continuation - plaque psoriasis

Dermatologist

*Re-assessment required after 6 months*

- Both:
- 1 Either:
    - 1.1 Both:
      - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
      - 1.1.2 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-etanercept treatment baseline value; or
    - 1.2 Both:
      - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
      - 1.2.2 Either:
        - 1.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
        - 1.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-etanercept treatment baseline value; and
  - 2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

## Monoclonal Antibodies

ABCIXIMAB – **Restricted** see terms below

⚡ Inj 2 mg per ml, 5 ml vial ..... 579.53      1      ReoPro

➡ **Restricted**

Either:

- 1 For use in patients with acute coronary syndromes undergoing percutaneous coronary intervention; or
- 2 For use in patients undergoing intra-cranial intervention.

ADALIMUMAB – **Restricted** see terms below

⚡ Inj 20 mg per 0.4 ml syringe ..... 1,799.92      2      Humira

⚡ Inj 40 mg per 0.8 ml pen ..... 1,799.92      2      HumiraPen

⚡ Inj 40 mg per 0.8 ml syringe ..... 1,799.92      2      Humira

➡ **Restricted**

### Initiation - juvenile idiopathic arthritis

Rheumatologist or named specialist

*Re-assessment required after 4 months*

- Either:
- 1 Either:
    - 1.1 Both:
      - 1.1.1 The patient has had an initial Special Authority approval for etanercept for juvenile idiopathic arthritis (JIA); and
      - 1.1.2 Either:
        - 1.1.2.1 The patient has experienced intolerable side effects from etanercept; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

1.1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for JIA; or

2 All of the following:

2.1 Patient diagnosed with Juvenile Idiopathic Arthritis (JIA); and

2.2 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and

2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m<sup>2</sup> weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and

2.5 Both:

2.5.1 Either:

2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or

2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and

2.5.2 Physician's global assessment indicating severe disease.

## Continuation - juvenile idiopathic arthritis

Rheumatologist or named specialist

*Re-assessment required after 6 months*

All of the following:

1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

2 Either:

2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or

2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

## Initiation - fistulising Crohn's disease

Gastroenterologist

*Re-assessment required after 4 months*

All of the following

1 Patient has confirmed Crohn's disease; and

2 Either:

2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or

2.2 Patient has one or more rectovaginal fistula(e); and

3 A Baseline Fistula Assessment (a copy of which is available at [www.pharmac.govt.nz/latest/BaselineFistulaAssessment.pdf](http://www.pharmac.govt.nz/latest/BaselineFistulaAssessment.pdf)) has been completed and is no more than 1 month old at the time of application.

## Continuation - fistulising Crohn's disease

Gastroenterologist

*Re-assessment required after 6 months*

Either:

1 The number of open draining fistulae have decreased from baseline by at least 50%; or

2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

### Initiation - Crohn's disease

Gastroenterologist

*Re-assessment required after 3 months*

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

### Continuation - Crohn's disease

Gastroenterologist

*Re-assessment required after 3 months*

Both:

- 1 Either:
  - 1.1 Either:
    - 1.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
    - 1.1.2 CDAI score is 150 or less; or
  - 1.2 Both:
    - 1.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
    - 1.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

### Initiation - rheumatoid arthritis

Rheumatologist

*Re-assessment required after 6 months*

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe and active erosive rheumatoid arthritis for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulphasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
  - 2.5 Any of the following:
    - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of cyclosporin; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
  - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
  - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

## Continuation - rheumatoid arthritis

Rheumatologist

*Re-assessment required after 6 months*

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Adalimumab to be administered at doses no greater than 50 mg every 7 days.

## Initiation - ankylosing spondylitis

Rheumatologist

*Re-assessment required after 6 months*

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
  - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of an exercise regime supervised by a physiotherapist; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and

2.6 Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of starting treatment.

Average normal chest expansion corrected for age and gender:

Age	Male	Female
18-24	7.0 cm	5.5 cm
25-34	7.5 cm	5.5 cm
35-44	6.5 cm	4.5 cm
45-54	6.0 cm	5.0 cm
55-64	5.5 cm	4.0 cm
65-74	4.0 cm	4.0 cm
75+	3.0 cm	2.5 cm

### Continuation - ankylosing spondylitis

Rheumatologist

*Re-assessment required after 6 months*

All of the following:

- Following 12 weeks of adalimumab treatment, BASDAI has improved by 4 or more points from pre-treatment baseline on a 10 point scale, or by 50%, whichever is less; and
- Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- Adalimumab to be administered at doses no greater than 40 mg every 14 days.

### Initiation - psoriatic arthritis

Rheumatologist

*Re-assessment required after 6 months*

Either:

- Both:
  - The patient has had an initial Special Authority approval for etanercept for psoriatic arthritis; and
  - Either:
    - The patient has experienced intolerable side effects from etanercept; or
    - The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for psoriatic arthritis; or
- All of the following:
  - Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - Patient has tried and not responded to at least three months of sulphasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - Either:
    - Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
    - Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - Any of the following:
    - Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

## Continuation - psoriatic arthritis

Rheumatologist

*Re-assessment required after 6 months*

Both:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

## Initiation - plaque psoriasis, prior TNF use

Dermatologist

*Re-assessment required after 4 months*

Both:

- 1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from etanercept; or
  - 2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; and

## Initiation - plaque psoriasis, treatment-naïve

Dermatologist

*Re-assessment required after 4 months*

All of the following:

- 1 Either:
  - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, cyclosporin, or acitretin; and
- 3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

## Continuation - plaque psoriasis

Dermatologist

*Re-assessment required after 6 months*

Both:

- 1 Either:
  - 1.1 Both:

continued...



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
- 1.1.2 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
- 1.2 Both:
  - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
  - 1.2.2 Either:
    - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
    - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-etanercept treatment baseline value; and

2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**BASILIXIMAB – Restricted** see terms below

⬇ Inj 20 mg vial .....3,200.00 1 Simulect

➔**Restricted**

For use in solid organ transplants

**BEVACIZUMAB – Restricted** see terms below

⬇ Inj 25 mg per ml, 16 ml vial

⬇ Inj 25 mg per ml, 4 ml vial

➔**Restricted**

Either:

- 1 Ocular neovascularisation; or
- 2 Exudative ocular angiopathy.

**INFLIXIMAB – Restricted** see terms below

⬇ Inj 100 mg .....1,227.00 1 Remicade

➔**Restricted**

**Graft vs host disease**

Patient has steroid-refractory acute graft vs. host disease of the gut

**Initiation - rheumatoid arthritis**

Rheumatologist

*Re-assessment required after 3-4 months*

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance

**Continuation - rheumatoid arthritis**

Rheumatologist

*Re-assessment required after 6 months*

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

## **Initiation - ankylosing spondylitis**

Rheumatologist

*Re-assessment required after 3 months*

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

## **Continuation - ankylosing spondylitis**

Rheumatologist

*Re-assessment required after 6 months*

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

## **Initiation - psoriatic arthritis**

Rheumatologist

*Re-assessment required after 3-4 months*

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for psoriatic arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for psoriatic arthritis.

## **Continuation - psoriatic arthritis**

Rheumatologist

*Re-assessment required after 6 months*

Both:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

## **Initiation - severe ocular inflammation**

*Re-assessment required after 3 doses*

Both:

- 1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
- 2 Either:
  - 2.1 Patient has failed to achieve control of severe vision-threatening ocular inflammation following high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids; or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

2.2 Patient developed new inflammatory symptoms while receiving high dose steroids.

### Initiation - chronic ocular inflammation

*Re-assessment required after 3 doses*

Both:

- 1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2 Patient has tried at least two other immunomodulatory agents.

### Continuation - ocular inflammation

Both:

- 1 Patient had a good clinical response to initial treatment; and
- 2 Either:
  - 2.1 A withdrawal of infliximab has been trialled and patient has relapsed after trial withdrawal; or
  - 2.2 Patient has Behcet's disease.

### Pulmonary sarcoidosis

Both:

- 1 Patient has life-threatening pulmonary sarcoidosis that is refractory to other treatments; and
- 2 Treatment is to be prescribed by, or has been recommended by, a physician with expertise in the treatment of pulmonary sarcoidosis.

### Initiation - Crohn's disease (adults)

Gastroenterologist

*Re-assessment required after 3 months*

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

### Continuation - Crohn's disease (adults)

Gastroenterologist

*Re-assessment required after 6 months*

All of the following:

- 1 One of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
  - 1.2 CDAI score is 150 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle; and
- 3 Patient must be reassessed for continuation after further 6 months.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

## Initiation - Crohn's disease (children)

Gastroenterologist

*Re-assessment required after 3 months*

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

## Continuation - Crohn's disease (children)

Gastroenterologist

*Re-assessment required after 6 months*

All of the following:

- 1 One of the following:
  - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle; and
- 3 Patient must be reassessed for continuation after further 6 months.

## Initiation - fistulising Crohn's disease

Gastroenterologist

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Either:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 Patient must be reassessed for continuation after 4 months of therapy.

## Continuation - fistulising Crohn's disease

Gastroenterologist

All of the following:

- 1 Either:
  - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle; and
- 3 Patient must be reassessed for continuation after further 6 months.

## Initiation - acute severe fulminant ulcerative colitis

Gastroenterologist

All of the following:

- 1 Patient has acute, severe fulminant ulcerative colitis; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful; and
- 3 Patient must be reassessed for continuation after 6 weeks of therapy.

**Continuation - severe fulminant ulcerative colitis**

Gastroenterologist

All of the following:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle; and
- 3 Patient must be reassessed for continuation after further 6 months.

**Initiation - severe ulcerative colitis**

Gastroenterologist

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 The Simple Clinical Colitis Activity Index (SCCAI) is  $\geq 4$
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

**Continuation - severe ulcerative colitis**

Gastroenterologist

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 SCCAI score has reduced by  $\geq 2$  points from the SCCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

**Initiation - plaque psoriasis, prior TNF use**

Dermatologist

*Re-assessment required after 3 doses*

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plaque psoriasis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
  - 2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plaque psoriasis.

**Initiation - plaque psoriasis, treatment-naïve**

Dermatologist

*Re-assessment required after 3 doses*

All of the following:

- 1 Either:
  - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, thotrexate, cyclosporin, or acitretin; and
- 3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

## Continuation - plaque psoriasis

Dermatologist

*Re-assessment required after 3 doses*

Both:

- 1 Either:
  - 1.1 Both:
    - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
  - 1.2 Both:
    - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
    - 1.2.2 Either:
      - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
      - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

RANIBIZUMAB – **Restricted** see terms below

⚡ Inj 10 mg per ml, 0.23 ml vial

⚡ Inj 10 mg per ml, 0.3 ml vial

## ➡Restricted

### Initiation

*Re-assessment required after 3 doses*

Both:

- 1 Either
  - 1.1 Age-related macular degeneration; or
  - 1.2 Choroidal neovascular membrane; and
- 2 Any of the following:
  - 2.1 The patient has had a severe ophthalmic inflammatory response following bevacizumab; or
  - 2.2 The patient has had a myocardial infarction or stroke within the last three months; or
  - 2.3 The patient has failed to respond to bevacizumab following three intraocular injections; or
  - 2.4 The patient is of child-bearing potential and has not completed a family.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

### Continuation

Both:

- 1 Documented benefit after three doses must be demonstrated to continue; and
- 2 In the case of but previous non-response to bevacizumab, a retreat of bevacizumab is required to confirm non-response before continuing with ranibizumab.

RITUXIMAB – **Restricted** see terms below

⚡ Inj 10 mg per ml, 10 ml vial .....	1,075.50	2	Mabthera
⚡ Inj 10 mg per ml, 50 ml vial .....	2,688.30	1	Mabthera

### ➡ Restricted

#### Initiation - haemophilia with inhibitors

Haematologist

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

#### Continuation - haemophilia with inhibitors

Haematologist

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

#### Initiation - post-transplant

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with \* are Unapproved Indications.

#### Continuation - post-transplant

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with \* are Unapproved Indications

#### Initiation - indolent, low-grade lymphomas

Either:

- 1 Both:
  - 1.1 The patient has indolent low grade NHL with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
  - 1.3 Both:
    - 1.3.1 The patient has indolent, low grade lymphoma requiring first-line systemic chemotherapy; and
    - 1.3.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia.

#### Continuation - indolent, low-grade lymphomas

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia.

## Initiation - aggressive CD20 positive NHL

Either:

- 1 All of the following:
  - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
  - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
  - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
  - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia.

## Continuation - aggressive CD20 positive NHL

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia.

## Chronic lymphocytic leukaemia

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 The patient is rituximab treatment naive; and
- 3 Either:
  - 3.1 The patient is chemotherapy treatment naive; or
  - 3.2 Both:
    - 3.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
    - 3.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; and
- 4 The patient has good performance status; and
- 5 The patient has good renal function (creatinine clearance  $\geq$  30 ml/min); and
- 6 The patient does not have chromosome 17p deletion CLL; and
- 7 Rituximab to be administered in combination with fludarabine and cyclophosphamide for a maximum of 6 treatment cycles; and
- 8 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration).

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to <2.

## Initiation - rheumatoid arthritis - prior TNF inhibitor use

Rheumatologist

*Re-assessment required after 2 doses*

All of the following:

- 1 Both:
  - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
  - 1.2 Either:

continued...



Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
- 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and

2 Either:

- 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and

3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

#### **Initiation - rheumatoid arthritis - TNF inhibitors contraindicated**

Rheumatologist

*Re-assessment required after 2 doses*

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulphasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
  - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of cyclosporin; or
  - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
  - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
  - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
  - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
  - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

#### **Continuation - rheumatoid arthritis - re-treatment in 'partial responders' to rituximab**

Rheumatologist

*Re-assessment required after 2 doses*

All of the following:

- 1 Either:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

## **Continuation - rheumatoid arthritis - re-treatment in 'responders' to rituximab**

Rheumatologist

*Re-assessment required after 2 doses*

All of the following:

- 1 Either:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

## **Initiation – severe cold haemagglutinin disease (CHAD)**

Haematologist

*Limited to 4 weeks' treatment*

Both:

- 1 Patient has cold haemagglutinin disease\*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms.

Note: Indications marked with \* are Unapproved Indications.

## **Continuation – severe cold haemagglutinin disease (CHAD)**

Haematologist

*Limited to 4 weeks' treatment*

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m<sup>2</sup> weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are Unapproved Indications.

## **Initiation – warm autoimmune haemolytic anaemia (warm AIHA)**

Haematologist

*Limited to 4 weeks' treatment*

Both:

- 1 Patient has warm autoimmune haemolytic anaemia\*; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to >5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin.

Note: Indications marked with \* are Unapproved Indications.

### **Continuation – warm autoimmune haemolytic anaemia (warm AIHA)**

Haematologist

*Limited to 4 weeks' treatment*

Either:

- Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m<sup>2</sup> weekly for 4 weeks) is now planned; or
- All of the following:
  - Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia\*; and
  - An initial response lasting at least 12 months was demonstrated; and
  - Patient now requires repeat treatment.

Note: Indications marked with \* are Unapproved Indications.

### **Initiation – immune thrombocytopenic purpura (ITP)**

Haematologist

*Limited to 4 weeks' treatment*

Both:

- Either:
  - Patient has immune thrombocytopenic purpura\* with a platelet count of  $\leq$  20,000 platelets per microlitre; or
  - Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- Any of the following:
  - Treatment with steroids and splenectomy have been ineffective; or
  - Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
  - Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy).

Note: Indications marked with \* are Unapproved Indications.

### **Continuation – immune thrombocytopenic purpura (ITP)**

Haematologist

*Limited to 4 weeks' treatment*

Either:

- Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m<sup>2</sup> weekly for 4 weeks) is now planned; or
- All of the following:
  - Patient was previously treated with rituximab for immune thrombocytopenic purpura\*; and
  - An initial response lasting at least 12 months was demonstrated; and
  - Patient now requires repeat treatment.

Note: Indications marked with \* are Unapproved Indications.

### **Initiation – thrombotic thrombocytopenic purpura (TTP)**

Haematologist

*Limited to 4 weeks' treatment*

Either:

- Patient has thrombotic thrombocytopenic purpura\* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
- Patient has acute idiopathic thrombotic thrombocytopenic purpura\* with neurological or cardiovascular pathology.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Note: Indications marked with \* are Unapproved Indications.

## **Continuation – thrombotic thrombocytopenic purpura (TTP)**

Haematologist

*Limited to 4 weeks' treatment*

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura\*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Note: Indications marked with \* are Unapproved Indications.

## **Initiation – pure red cell aplasia (PRCA)**

Haematologist

*Limited to 6 weeks' treatment*

Patient has autoimmune pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with \* are Unapproved Indications.

## **Continuation – pure red cell aplasia (PRCA)**

Haematologist

*Limited to 6 weeks' treatment*

Patient was previously treated with rituximab for pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with \* are Unapproved Indications.

## **Initiation – ANCA associated vasculitis**

*Limited to 4 weeks' treatment*

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 Either:
  - 2.1 Patient does not have MPO-ANCA positive vasculitis\*; or
  - 2.2 Mycophenolate mofetil has not been effective in those patients who have MPO-ANCA positive vasculitis\*; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks; and
- 4 Any of the following:
  - 4.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve complete absence of disease after at least 3 months; or
  - 4.2 Patient has previously had a cumulative dose of cyclophosphamide >15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose >15 g; or
  - 4.3 Cyclophosphamide and methotrexate are contraindicated; or
  - 4.4 Patient is a female of child-bearing potential; or
  - 4.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with \* are Unapproved Indications.

## **Continuation – ANCA associated vasculitis**

*Limited to 4 weeks' treatment*

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks.

Note: Indications marked with \* are Unapproved Indications.

## **Initiation – treatment refractory systemic lupus erythematosus (SLE)**

Rheumatologist or nephrologist

All of the following:

continued...

Price (ex man. excl. GST)	Brand or Generic
\$ Per	Manufacturer

continued. . .

- 1 The patient has severe, immediately life- or organ-threatening SLE\*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with \* are Unapproved Indications.

#### **Continuation – treatment refractory systemic lupus erythematosus (SLE)**

Rheumatologist or nephrologist

All of the following:

- 1 Patient's SLE\* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with \* are Unapproved Indications.

#### **Antibody-mediated renal transplant rejection**

Nephrologist

Patient has been diagnosed with antibody-mediated renal transplant rejection\*.

Note: Indications marked with \* are Unapproved Indications.

#### **ABO-incompatible renal transplant**

Nephrologist

Patient is to undergo an ABO-incompatible renal transplant\*.

Note: Indications marked with \* are Unapproved Indications.

TOCILIZUMAB – **Restricted** see terms below

⚡ Inj 20 mg per ml, 4 ml vial .....	220.00	1	Actemra
⚡ Inj 20 mg per ml, 10 ml vial .....	550.00	1	Actemra
⚡ Inj 20 mg per ml, 20 ml vial .....	1,100.00	1	Actemra

#### **➡Restricted**

#### **Initiation - systemic juvenile idiopathic arthritis**

Paediatric rheumatologist

*Re-assessment required after 6 months*

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

#### **Continuation - systemic juvenile idiopathic arthritis**

Paediatric rheumatologist

*Re-assessment required after 6 months*

Either:

- 1 Following up to 6 months initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

TRASTUZUMAB – **Restricted** see terms on the next page

⚡ Inj 150 mg vial .....	1,350.00	1	Herceptin
⚡ Inj 440 mg vial .....	3,875.00	1	Herceptin

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## ➡Restricted

### Early breast cancer

Limited to 12 months' treatment

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
  - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
  - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
  - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
  - 3.4 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

### Initiation - metastatic breast cancer (trastuzumab-naïve patients)

*Re-assessment required after 12 months*

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The patient has not previously received lapatinib treatment for HER 2 positive metastatic breast cancer; and
  - 1.3 Trastuzumab not to be given in combination with lapatinib; and
  - 1.4 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
  - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 2.2 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
  - 2.3 The cancer did not progress whilst on lapatinib; and
  - 2.4 Trastuzumab not to be given in combination with lapatinib; and
  - 2.5 Trastuzumab to be discontinued at disease progression.

### Initiation - metastatic breast cancer (patients previously treated with trastuzumab)

*Re-assessment required after 12 months*

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
  - 3.1 All of the following:
    - 3.1.1 The patient has not previously received lapatinib treatment for metastatic breast cancer; and
    - 3.1.2 Trastuzumab not to be given in combination with lapatinib; and
    - 3.1.3 Trastuzumab to be discontinued at disease progression; or
  - 3.2 All of the following:
    - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
    - 3.2.2 The cancer did not progress whilst on lapatinib; and
    - 3.2.3 Trastuzumab not to be given in combination with lapatinib; and
    - 3.2.4 Trastuzumab to be discontinued at disease progression; or
  - 3.3 All of the following:
    - 3.3.1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
    - 3.3.2 Trastuzumab not to be given in combination with lapatinib; and
    - 3.3.3 Trastuzumab to be discontinued at disease progression.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

### Continuation - metastatic breast cancer

Re-assessment required after 12 months

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

### Other Immunosuppressants

#### ANTITHYMOCYTE GLOBULIN (EQUINE)

Inj 50 mg per ml, 5 ml ampoule .....	2,137.50	5	ATGAM
--------------------------------------	----------	---	-------

#### ANTITHYMOCYTE GLOBULIN (RABBIT)

Inj 25 mg vial

#### AZATHIOPRINE

Tab 50 mg – 1% DV Jun-14 to 2016 .....	13.22	100	<b>Azamun</b>
Inj 50 mg vial .....	126.00	1	Imuran

#### BACILLUS CALMETTE-GUERIN (BCG) – **Restricted** see terms below

⚡ Inj 2-8 × 10 <sup>8</sup> CFU vial – 1% DV Sep-13 to 2016 .....	149.37	1	<b>OncoTICE</b>
---	--------	---	-----------------

#### → **Restricted**

For use in bladder cancer

#### MYCOPHENOLATE MOFETIL – **Restricted** see terms below

⚡ Tab 500 mg – 1% DV Nov-13 to 2016 .....	25.00	50	<b>CellCept</b>
⚡ Cap 250 mg – 1% DV Nov-13 to 2016 .....	25.00	100	<b>CellCept</b>
⚡ Powder for oral liq 1 g per 5 ml – 1% DV Nov-13 to 2016 .....	187.25	165 ml	<b>CellCept</b>
⚡ Inj 500 mg vial – 1% DV Nov-13 to 2016 .....	133.33	4	<b>CellCept</b>

#### → **Restricted**

Either:

- 1 Transplant recipient; or
- 2 Patients with diseases where both:
  - 2.1 Steroids and azathioprine have been trialled and discontinued because of unacceptable side effects or inadequate clinical response; and
  - 2.2 Either:
    - 2.2.1 Cyclophosphamide has been trialled and discontinued because of unacceptable side effects or inadequate clinical response; or
    - 2.2.2 Cyclophosphamide treatment is contraindicated.

#### PICIBANIL

Inj 100 mg vial

#### SIROLIMUS – **Restricted** see terms on the next page

⚡ Tab 1 mg .....	813.00	100	Rapamune
⚡ Tab 2 mg .....	1,626.00	100	Rapamune
⚡ Oral liq 1 mg per ml .....	487.80	60 ml	Rapamune

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➡Restricted

For rescue therapy for an organ transplant recipient

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR < 30 ml/min; or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- HUS or TTP; or
- Leukoencephalopathy; or
- Significant malignant disease



Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

## Antiallergy Preparations

### Allergy Desensitisation

BEE VENOM – **Restricted** see terms below

⚡ Inj 120 mcg vial with diluent, 6 vial

⚡ Inj 550 mcg vial with diluent

➡ **Restricted**

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

PAPER WASP VENOM – **Restricted** see terms below

⚡ Inj 550 mcg vial with diluent

➡ **Restricted**

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

YELLOW JACKET WASP VENOM – **Restricted** see terms below

⚡ Inj 550 mcg vial with diluent

➡ **Restricted**

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

### Allergy Prophylactics

BECLOMETHASONE DIPROPIONATE

Nasal spray 50 mcg per dose .....	4.85	200 dose	Alanase
-----------------------------------	------	----------	---------

Nasal spray 100 mcg per dose .....	5.75	200 dose	Alanase
------------------------------------	------	----------	---------

BUDESONIDE

Nasal spray 50 mcg per dose .....	4.85	200 dose	Butacort Aqueous
-----------------------------------	------	----------	------------------

Nasal spray 100 mcg per dose .....	5.75	200 dose	Butacort Aqueous
------------------------------------	------	----------	------------------

FLUTICASONE PROPIONATE

Nasal spray 50 mcg per dose – 1% DV Apr-13 to 2015 .....	2.30	120 dose	<b>Flixonase Hayfever &amp; Allergy</b>
--	------	----------	---

IPRATROPIUM BROMIDE

Nasal spray 0.03%

SODIUM CROMOGLYCATE

Nasal spray 4%

### Antihistamines

CETIRIZINE HYDROCHLORIDE

Tab 10 mg – 1% DV Sep-11 to 2014 .....	1.59	100	<b>Zetop</b>
--	------	-----	--------------

Oral liq 1 mg per ml – 1% DV Nov-11 to 2014 .....	3.52	200 ml	<b>Cetirizine - AFT</b>
---	------	--------	-------------------------

CHLORPHENIRAMINE MALEATE

Oral liq 0.4 mg per ml

Inj 10 mg per ml, 1 ml ampoule

CYPROHEPTADINE HYDROCHLORIDE

Tab 4 mg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>FEXOFENADINE HYDROCHLORIDE</b>			
Tab 60 mg			
Tab 120 mg			
Tab 180 mg			
<b>LORATADINE</b>			
Tab 10 mg – 1% DV Dec-13 to 2016 .....	1.30	100	<b>Lorafix</b>
Oral liq 1 mg per ml .....	3.10	100 ml	Lorapaed
<b>PROMETHAZINE HYDROCHLORIDE</b>			
Tab 10 mg – 1% DV Sep-12 to 2015 .....	1.99	50	<b>Allersoothe</b>
Tab 25 mg – 1% DV Sep-12 to 2015 .....	2.99	50	<b>Allersoothe</b>
Oral liq 1 mg per ml – 1% DV Feb-13 to 2015 .....	2.79	100 ml	<b>Allersoothe</b>
Inj 25 mg per ml, 2 ml ampoule .....	11.00	5	Hospira
<b>TRIMEPAZINE TARTRATE</b>			
Oral liq 6 mg per ml			

## Anticholinergic Agents

<b>IPRATROPIUM BROMIDE</b>			
Aerosol inhaler 20 mcg per dose			
Nebuliser soln 250 mcg per ml, 1 ml ampoule – 1% DV Sep-13 to 2016 .....	3.26	20	<b>Univent</b>
Nebuliser soln 250 mcg per ml, 2 ml ampoule – 1% DV Sep-13 to 2016 .....	3.37	20	<b>Univent</b>
<b>TIOTROPIUM BROMIDE – Restricted</b> see terms below			
☞ Powder for inhalation 18 mcg per dose .....	70.00	30 dose	Spiriva

### ☞Restricted

All of the following:

- 1 To be used for the long-term maintenance treatment of bronchospasm and dyspnoea associated with COPD; and
- 2 In addition to standard treatment, the patient has trialled a short acting bronchodilator of at least 40 mcg ipratropium q.i.d for one month; and
- 3 The patient's breathlessness according to the Medical Research Council (UK) dyspnoea scale is either:
  - 3.1 Grade 4 (stops for breath after walking about 100 metres or after a few minutes on the level); or
  - 3.2 Grade 5 (too breathless to leave the house, or breathless when dressing or undressing); and
- 4 Actual FEV<sub>1</sub> as a % of predicted, must be below 60%.
- 5 Either:
  - 5.1 Patient is not a smoker; or
  - 5.2 Patient is a smoker and has been offered smoking cessation counselling; and
- 6 The patient has been offered annual influenza immunisation.

## Anticholinergic Agents with Beta-Adrenoceptor Agonists

<b>SALBUTAMOL WITH IPRATROPIUM BROMIDE</b>			
Aerosol inhaler 100 mcg with ipratropium bromide 20 mcg per dose			
Nebuliser soln 2.5 mg with ipratropium bromide 0.5 mg per 2.5 ml ampoule – 1% DV Nov-12 to 2015 .....	3.75	20	<b>Duolin</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Beta-Adrenoceptor Agonists</b>			
<b>SALBUTAMOL</b>			
Oral liq 400 mcg per ml – <b>1% DV Jan-14 to 2016</b> .....	2.06	150 ml	<b>Ventolin</b>
Inj 500 mcg per ml, 1 ml ampoule			
Inj 1 mg per ml, 5 ml ampoule			
Aerosol inhaler, 100 mcg per dose .....	4.00	200 dose	Salamol
	6.00		Ventolin
Nebuliser soln 1 mg per ml, 2.5 ml ampoule – <b>1% DV Nov-12 to 2015</b> .....	3.25	20	<b>Asthalin</b>
Nebuliser soln 2 mg per ml, 2.5 ml ampoule – <b>1% DV Nov-12 to 2015</b> .....	3.44	20	<b>Asthalin</b>
<b>TERBUTALINE SULPHATE</b>			
Powder for inhalation 250 mcg per dose			
Inj 0.5 mg per ml, 1 ml ampoule			
<b>Cough Suppressants</b>			
<b>PHOLCODINE</b>			
Oral liq 1 mg per ml			
<b>Decongestants</b>			
<b>OXYMETAZOLINE HYDROCHLORIDE</b>			
Aqueous nasal spray 0.25 mg per ml			
Aqueous nasal spray 0.5 mg per ml			
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b>			
Tab 60 mg			
<b>SODIUM CHLORIDE</b>			
Aqueous nasal spray 7.4 mg per ml			
<b>SODIUM CHLORIDE WITH SODIUM BICARBONATE</b>			
Soln for nasal irrigation			
<b>XYLOMETAZOLINE HYDROCHLORIDE</b>			
Aqueous nasal spray 0.05%			
Aqueous nasal spray 0.1%			
Nasal drops 0.05%			
Nasal drops 0.1%			
<b>Inhaled Corticosteroids</b>			
<b>BECLOMETHASONE DIPROPIONATE</b>			
Aerosol inhaler 50 mcg per dose .....	8.54	200 dose	Beclazone 50
Aerosol inhaler 100 mcg per dose .....	12.50	200 dose	Beclazone 100
Aerosol inhaler 250 mcg per dose .....	22.67	200 dose	Beclazone 250
<b>BUDESONIDE</b>			
Nebuliser soln 250 mcg per ml, 2 ml ampoule			
Nebuliser soln 500 mcg per ml, 2 ml ampoule			
Powder for inhalation 100 mcg per dose			
Powder for inhalation 200 mcg per dose			
Powder for inhalation 400 mcg per dose			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>FLUTICASONE</b>			
Aerosol inhaler 50 mcg per dose .....	7.50	120 dose	Flixotide
Powder for inhalation 50 mcg per dose .....	8.67	60 dose	Flixotide Accuhaler
Powder for inhalation 100 mcg per dose .....	13.87	60 dose	Flixotide Accuhaler
Aerosol inhaler 125 mcg per dose .....	13.60	120 dose	Flixotide
Aerosol inhaler 250 mcg per dose .....	27.20	120 dose	Flixotide
Powder for inhalation 250 mcg per dose .....	24.51	60 dose	Flixotide Accuhaler

## Leukotriene Receptor Antagonists

**MONTELUKAST – Restricted** see terms below

☞ Tab 4 mg .....	18.48	28	Singulair
☞ Tab 5 mg .....	18.48	28	Singulair
☞ Tab 10 mg .....	18.48	28	Singulair

☞ **Restricted**

**Pre-school wheeze**

Both:

- 1 To be used for the treatment of intermittent severe wheezing (possibly viral); and
- 2 The patient has had at least three episodes in the previous 12 months of acute wheeze severe enough to seek medical attention.

**Exercise-induced asthma**

Both:

- 1 Patient has been trialed with maximal asthma therapy, including inhaled corticosteroids and long-acting beta-adrenoceptor agonists; and
- 2 Patient continues to receive optimal inhaled corticosteroid therapy; and
- 3 Patient continues to experience frequent episodes of exercise-induced bronchoconstriction.

**Aspirin desensitisation**

Clinical immunologist or allergist

All of the following:

- 1 Patient is undergoing aspirin desensitisation therapy under the supervision of a clinical immunologist or allergist; and
- 2 Patient has moderate to severe aspirin-exacerbated respiratory disease or Samter's triad; and
- 3 Nasal polyposis, confirmed radiologically or surgically; and
- 4 Documented aspirin or NSAID allergy confirmed by aspirin challenge or a clinical history of severe reaction to aspirin or NSAID where challenge would be considered dangerous.

## Long-Acting Beta-Adrenoceptor Agonists

**EFORMOTEROL FUMARATE**

- Powder for inhalation 6 mcg per dose
- Powder for inhalation 12 mcg per dose

**SALMETEROL**

Aerosol inhaler 25 mcg per dose .....	26.46	120 dose	Serevent
Powder for inhalation 50 mcg per dose .....	26.46	60 dose	Serevent Accuhaler

## Inhaled Corticosteroids with Long-Acting Beta-Adrenoceptor Agonists

**BUDESONIDE WITH EFORMOTEROL – Restricted** see terms on the next page

- ☞ Powder for inhalation 100 mcg with eformoterol fumarate 6 mcg
- ☞ Powder for inhalation 200 mcg with eformoterol fumarate 6 mcg
- ☞ Powder for inhalation 400 mcg with eformoterol fumarate 12 mcg
- ☞ Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg
- ☞ Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## ➔Restricted

Either:

- 1 All of the following:
  - 1.1 Patient is a child under the age of 12; and
  - 1.2 Has been treated with inhaled corticosteroids of at least 400 mcg per day beclomethasone or budesonide, or 200 mcg per day fluticasone; and
  - 1.3 The prescriber considers that the patient would receive additional clinical benefit from switching to a combination product; or
- 2 All of the following:
  - 2.1 Patient is over the age of 12; and
  - 2.2 Has been treated with inhaled corticosteroids of at least 800 mcg per day beclomethasone or budesonide, or 500 mcg per day fluticasone; and
  - 2.3 The prescriber considers that the patient would receive additional clinical benefit from switching to a combination product.

### FLUTICASONE WITH SALMETEROL

Aerosol inhaler 50 mcg with salmeterol 25 mcg .....	37.48	120 dose	Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg .....	37.48	60 dose	Seretide Accuhaler
Aerosol inhaler 125 mcg with salmeterol 25 mcg .....	49.69	120 dose	Seretide
Powder for inhalation 250 mcg with salmeterol 50 mcg .....	49.69	60 dose	Seretide Accuhaler

## Mast Cell Stabilisers

### NEDOCROMIL

Aerosol inhaler 2 mg per dose

### SODIUM CROMOGLYCATE

Powder for inhalation 20 mg per dose

Aerosol inhaler 5 mg per dose

## Methylxanthines

### AMINOPHYLLINE

Inj 25 mg per ml, 10 ml ampoule – **1% DV Nov-11 to 2014**.....53.75      5      **DBL Aminophylline**

### CAFFEINE CITRATE

Oral liq 20 mg per ml (caffeine 10 mg per ml) .....	14.85	25 ml	Biomed
Inj 20 mg per ml (caffeine 10 mg per ml), 2.5 ml ampoule .....	55.75	5	Biomed

### THEOPHYLLINE

Tab long-acting 250 mg

Oral liq 80 mg per 15 ml

## Mucolytics and Expectorants

### DORNASE ALFA – **Restricted** see terms below

☞ Nebuliser soln 2.5 mg per 2.5 ml ampoule .....250.00      6      Pulmozyme

## ➔Restricted

Any of the following:

- 1 Cystic fibrosis and the patient has been approved by the Cystic Fibrosis Panel; and/or
- 2 Significant mucus production and meets the following criteria
- 3 Treatment for up to four weeks for patients meeting the following:
  - 3.1 Patient is an in-patient; and
  - 3.2 The mucus production cannot be cleared by first line chest techniques; or
- 4 Treatment for up to three days for patients diagnosed with empyema.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SODIUM CHLORIDE			
Nebuliser soln 7%, 90 ml bottle .....	23.50	90 ml	Biomed

### Pulmonary Surfactants

BERACTANT			
Soln 200 mg per 8 ml vial .....	550.00	1	Survanta
PORACTANT ALFA			
Soln 120 mg per 1.5 ml vial .....	425.00	1	Curosurf
Soln 240 mg per 3 ml vial .....	695.00	1	Curosurf

### Respiratory Stimulants

DOXAPRAM			
Inj 20 mg per ml, 5 ml vial			

### Sclerosing Agents

TALC			
Powder			
Soln (slurry) 100 mg per ml, 50 ml			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Anti-Infective Preparations</b>			
<b>Antibacterials</b>			
CHLORAMPHENICOL			
Eye oint 1% – <b>1% DV Jan-13 to 2015</b> .....	2.76	4 g	<b>Chlorsig</b>
Ear drops 0.5%			
Eye drops 0.5% – <b>1% DV Sep-12 to 2015</b> .....	1.20	10 ml	<b>Chlorafast</b>
Eye drops 0.5%, single dose			
CIPROFLOXACIN			
Eye drops 0.3%			
FRAMYCETIN SULPHATE			
Ear/eye drops 0.5%			
FUSIDIC ACID			
Eye drops 1% .....	4.50	5 g	Fucithalmic
GENTAMICIN SULPHATE			
Eye drops 0.3% .....	11.40	5 ml	Genoptic
PROPAMIDINE ISETHIONATE			
Eye drops 0.1%			
SULPHACETAMIDE SODIUM			
Eye drops 10%			
TOBRAMYCIN			
Eye oint 0.3% – <b>1% DV Sep-11 to 2014</b> .....	10.45	3.5 g	<b>Tobrex</b>
Eye drops 0.3% – <b>1% DV Sep-11 to 2014</b> .....	11.48	5 ml	<b>Tobrex</b>
<b>Antifungals</b>			
NATAMYCIN			
Eye drops 5%			
<b>Antivirals</b>			
ACICLOVIR			
Eye oint 3%			
<b>Combination Preparations</b>			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/eye drops 500 mcg with framycetin sulphate 5 mg and gramicidin 50 mcg per ml			
DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE			
Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin B sul- phate 6,000 u per g			
Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin B sul- phate 6,000 u per ml			
DEXAMETHASONE WITH TOBRAMYCIN			
Eye drops 0.1% with tobramycin 0.3%			
FLUMETASONE PIVALATE WITH CLIOQUINOL			
Ear drops 0.02% with clioquinol 1%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
HYDROCORTISONE WITH CIPROFLOXACIN Ear drops 1% with ciprofloxacin 0.2%			
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCIN AND NYSTATIN Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g .....	5.16	7.5 ml	Kenacomb

## Anti-Inflammatory Preparations

### Corticosteroids

DEXAMETHASONE Eye oint 0.1% – 1% DV Sep-11 to 2014 .....	5.86	3.5 g	Maxidex
Eye drops 0.1% .....	4.50	5 ml	Maxidex
FLUOROMETHOLONE Eye drops 0.1% – 1% DV Dec-12 to 2015 .....	3.80	5 ml	Flucon
PREDNISOLONE ACETATE Eye drops 0.12% Eye drops 1%			
PREDNISOLONE SODIUM PHOSPHATE Eye drops 0.5%, single dose			

### Non-Steroidal Anti-Inflammatory Drugs

DICLOFENAC SODIUM Eye drops 0.1% – 1% DV Sep-11 to 2014 .....	13.80	5 ml	Voltaren Ophtha
Eye drops 0.1%, single dose			
KETOROLAC TROMETAMOL Eye drops 0.5%			

## Decongestants and Antiallergics

### Antiallergic Preparations

LEVOCABASTINE Eye drops 0.05%			
LODOXAMIDE Eye drops 0.1%			
OLOPATADINE Eye drops 0.1%			
SODIUM CROMOGLYCATE Eye drops 2%			

### Decongestants

NAPHAZOLINE HYDROCHLORIDE Eye drops 0.1% – 1% DV Sep-11 to 2014 .....	4.15	15 ml	Naphcon Forte
--	------	-------	---------------



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Diagnostic and Surgical Preparations

Diagnostic Dyes

FLUORESC EIN SODIUM			
Eye drops 2%, single dose			
Inj 10%, 5 ml vial	125.00	12	Fluorescite
Ophthalmic strips 1 mg			
FLUORESC EIN SODIUM WITH LIGNOCAINE HYDROCHLORIDE			
Eye drops 0.25% with lignocaine hydrochloride 4%, single dose			
LISSAMINE GREEN			
Ophthalmic strips 1.5 mg			
ROSE BENGAL SODIUM			
Ophthalmic strips 1%			

Irrigation Solutions

CALCIUM CHLORIDE WITH MAGNESIUM CHLORIDE, POTASSIUM CHLORIDE, SODIUM ACETATE, SODIUM CHLORIDE AND SODIUM CITRATE			
Eye drops 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 15 ml			e.g. Balanced Salt Solution
Eye drops 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 250 ml			e.g. Balanced Salt Solution
Eye drops 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 500 ml			e.g. Balanced Salt Solution

Ocular Anaesthetics

OXYBUPROCAINE HYDROCHLORIDE	
Eye drops 0.4%, single dose	
PROXYMETACAINE HYDROCHLORIDE	
Eye drops 0.5%	
TETRACAINE [AMETHOCAINE] HYDROCHLORIDE	
Eye drops 0.5%, single dose	
Eye drops 1%, single dose	

Viscoelastic Substances

HYPROMELLOSE	
Inj 2%, 1 ml syringe	
Inj 2%, 2 ml syringe	

## SENSORY ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>SODIUM HYALURONATE</b>			
Inj 14 mg per ml, 0.85 ml syringe – 1% DV Oct-12 to 2015	50.00	1	Healon GV
Inj 14 mg per ml, 0.55 ml syringe – 1% DV Oct-12 to 2015	50.00	1	Healon GV
Inj 23 mg per ml, 0.6 ml syringe			
Inj 10 mg per ml, 0.85 ml syringe – 1% DV Oct-12 to 2015	30.00	1	Provisc
<b>SODIUM HYALURONATE WITH CHONDROITIN SULPHATE</b>			
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.35 ml syringe and inj 10 mg sodium hyaluronate per ml, 0.4 ml syringe	64.00	1	Duovisc
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.5 ml syringe and inj 10 mg sodium hyaluronate per ml, 0.55 ml syringe – 1% DV Sep-11 to 2014	74.00	1	Duovisc
Inj 30 mg with chondroitin sulphate 40 mg per ml, 0.75 ml syringe			
<b>Other</b>			
<b>RIBOFLAVIN 5-PHOSPHATE</b>			
Soln trans epithelial riboflavin			
Inj 0.1%			
Inj 0.1% plus 20% dextran T500			
<b>Glaucoma Preparations</b>			
<b>Beta Blockers</b>			
<b>BETAXOLOL</b>			
Eye drops 0.25%			
Eye drops 0.5%			
<b>LEVOBUNOLOL HYDROCHLORIDE</b>			
Eye drops 0.25%	7.00	5 ml	Betagan
Eye drops 0.5%	7.00	5 ml	Betagan
<b>TIMOLOL</b>			
Eye drops 0.25%			
Eye drops 0.25%, gel forming – 1% DV Mar-14 to 2016	3.30	2.5 ml	Timoptol XE
Eye drops 0.5%			
Eye drops 0.5%, gel forming – 1% DV Mar-14 to 2016	3.78	2.5 ml	Timoptol XE
<b>Carbonic Anhydrase Inhibitors</b>			
<b>ACETAZOLAMIDE</b>			
Tab 250 mg – 1% DV Nov-11 to 2014	17.03	100	Diamox
Inj 500 mg			
<b>BRINZOLAMIDE</b>			
Eye drops 1%			
<b>DORZOLAMIDE</b>			
Eye drops 2%			
<b>DORZOLAMIDE WITH TIMOLOL</b>			
Eye drops 2% with timolol 0.5%	15.50	5 ml	Cosopt
<b>Miotics</b>			
<b>ACETYLCHOLINE CHLORIDE</b>			
Inj 20 mg vial with diluent			

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PILOCARPINE HYDROCHLORIDE</b>			
Eye drops 1%			
Eye drops 2%			
Eye drops 2%, single dose			
Eye drops 4%			
<b>Prostaglandin Analogues</b>			
<b>BIMATOPROST</b>			
Eye drops 0.03%			
<b>LATANOPROST</b>			
Eye drops 0.005% – <b>1% DV Sep-12 to 2015</b> .....	1.99	2.5 ml	<b>Hysite</b>
<b>TRAVOPROST</b>			
Eye drops 0.004%			
<b>Sympathomimetics</b>			
<b>APRACLOPIDINE</b>			
Eye drops 0.5%			
<b>BRIMONIDINE TARTRATE</b>			
Eye drops 0.2% – <b>1% DV Jul-12 to 2014</b> .....	6.45	5 ml	<b>Arrow-Brimonidine</b>
<b>BRIMONIDINE TARTRATE WITH TIMOLOL</b>			
Eye drops 0.2% with timolol 0.5%			
<b>Mydriatics and Cycloplegics</b>			
<b>Anticholinergic Agents</b>			
<b>ATROPINE SULPHATE</b>			
Eye drops 0.5%			
Eye drops 1%, single dose			
Eye drops 1% – <b>1% DV Jul-14 to 2017</b> .....	17.36	15 ml	<b>Atropt</b>
<b>CYCLOPENTOLATE HYDROCHLORIDE</b>			
Eye drops 0.5%, single dose			
Eye drops 1%			
Eye drops 1%, single dose			
<b>TROPICAMIDE</b>			
Eye drops 0.5% – <b>1% DV Sep-11 to 2014</b> .....	7.15	15 ml	<b>Mydriacyl</b>
Eye drops 0.5%, single dose			
Eye drops 1% – <b>1% DV Sep-11 to 2014</b> .....	8.66	15 ml	<b>Mydriacyl</b>
Eye drops 1%, single dose			
<b>Sympathomimetics</b>			
<b>PHENYLEPHRINE HYDROCHLORIDE</b>			
Eye drops 2.5%, single dose			
Eye drops 10%, single dose			
<b>Ocular Lubricants</b>			
<b>CARBOMER</b>			
Ophthalmic gel 0.3%, single dose .....	8.25	30	Poly Gel
Ophthalmic gel 0.2%			

## SENSORY ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>CARMELLOSE SODIUM</b>			
Eye drops 0.5%			
Eye drops 0.5%, single dose			
Eye drops 1%			
Eye drops 1%, single dose			
<b>HYPROMELLOSE</b>			
Eye drops 0.5% .....	3.92	15 ml	Methopt
<b>HYPROMELLOSE WITH DEXTRAN</b>			
Eye drops 0.3% with dextran 0.1% .....	2.30	15 ml	Poly-Tears
Eye drops 0.3% with dextran 0.1%, single dose			
<b>MACROGOL 400 AND PROPYLENE GLYCOL</b>			
Eye drops 0.4% with propylene glycol 0.3% preservative free, single dose .....	4.30	24	Systane Unit Dose
<b>PARAFFIN LIQUID WITH SOFT WHITE PARAFFIN</b>			
Eye oint 42.5% with soft white paraffin 57.3%			
<b>PARAFFIN LIQUID WITH WOOL FAT</b>			
Eye oint 3% with wool fat 3% – 1% DV Jul-14 to 2017 .....	3.63	3.5 g	<b>Poly-Visc</b>
<b>POLYVINYL ALCOHOL</b>			
Eye drops 1.4% .....	2.95	15 ml	Vistil
	3.62		Liquifilm Tears
Eye drops 3% .....	3.80	15 ml	Vistil Forte
	3.88		Liquifilm Forte
<b>POLYVINYL ALCOHOL WITH POVIDONE</b>			
Eye drops 1.4% with povidone 0.6%, single dose			
<b>RETINOL PALMITATE</b>			
Oint 138 mcg per g .....	3.80	5 g	VitA-POS
<b>SODIUM HYALURONATE</b>			
Eye drops 1 mg per ml .....	22.00	10 ml	Hylo-Fresh

### Other Otolological Preparations

<b>ACETIC ACID WITH PROPYLENE GLYCOL</b>
Ear drops 2.3% with propylene glycol 2.8%
<b>DOCUSATE SODIUM</b>
Ear drops 0.5%

Price (ex man. excl. GST)	Brand or Generic
\$ Per	Manufacturer

## Agents Used in the Treatment of Poisonings

### Antidotes

#### ACETYLCYSTEINE

Tab eff 200 mg

Inj 200 mg per ml, 10 ml ampoule – **1% DV Jul-12 to 2015** ..... 178.00

10

**Martindale**

**Acetylcysteine**

Inj 200 mg per ml, 30 ml vial ..... 219.00

4

Acetadote

#### DIGOXIN IMMUNE FAB

Inj 38 mg vial

Inj 40 mg vial

#### ETHANOL

Liq 96%

#### ETHANOL WITH GLUCOSE

Inj 10% with glucose 5%, 500 ml bottle

#### ETHANOL, DEHYDRATED

Inj 100%, 5 ml ampoule

Inj 96%

#### FLUMAZENIL

Inj 0.1 mg per ml, 5 ml ampoule ..... 170.10

5

Anexate

#### HYDROXOCOBALAMIN

Inj 5 g vial

Inj 2.5 g vial

#### NALOXONE HYDROCHLORIDE

Inj 400 mcg per ml, 1 ml ampoule ..... 33.00

5

Hospira

#### PRALDOXIME IODIDE

Inj 25 mg per ml, 20 ml ampoule

#### SODIUM NITRITE

Inj 30 mg per ml, 10 ml ampoule

#### SODIUM THIOSULFATE

Inj 500 mg per ml, 20 ml ampoule

Inj 250 mg per ml, 10 ml vial

Inj 500 mg per ml, 10 ml vial

#### SOYA OIL

Inj 20%, 500 ml bag

Inj 20%, 500 ml bottle

### Antitoxins

#### BOTULISM ANTITOXIN

Inj 250 ml vial

#### DIPHTHERIA ANTITOXIN

Inj 10,000 iu vial

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Antivenoms</b>			
RED BACK SPIDER ANTIVENOM Inj 500 u vial			
SNAKE ANTIVENOM Inj 50 ml vial			
<b>Removal and Elimination</b>			
CHARCOAL Oral liq 200 mg per ml .....	43.50	250 ml	Carbasorb-X
DEFERIPRONE Tab 500 mg .....	533.17	100	Ferriprox
Oral liq 100 mg per ml .....	266.59	250 ml	Ferriprox
DESFERRIOXAMINE MESILATE Inj 500 mg vial .....	99.00	10	Hospira
DICOBALT EDETATE Inj 15 mg per ml, 20 ml ampoule			
DIMERCAPROL Inj 50 mg per ml, 2 ml ampoule			
DIMERCAPTOSUCCINIC ACID Cap 100 mg			
DISODIUM EDETATE Inj 150 mg per ml, 20 ml ampoule Inj 150 mg per ml, 20 ml vial Inj 150 mg per ml, 100 ml vial			
SODIUM CALCIUM EDETATE Inj 200 mg per ml, 2.5 ml ampoule Inj 200 mg per ml, 5 ml ampoule			
<b>Antiseptics and Disinfectants</b>			
CHLORHEXIDINE Soln 4% .....	1.86	50 ml	healthE
Soln 5% .....	15.50	500 ml	healthE
CHLORHEXIDINE WITH CETRIMIDE Crm 0.1% with cetrimide 0.5% Foaming soln 0.5% with cetrimide 0.5%			
CHLORHEXIDINE WITH ETHANOL Soln 0.5% with ethanol 70%, non-staining (pink) 100 ml .....	2.65	1	healthE
Soln 2% with ethanol 70%, non-staining (pink) 100 ml .....	3.54	1	healthE
Soln 0.5% with ethanol 70%, non-staining (pink) 25 ml .....	1.55	1	healthE
Soln 0.5% with ethanol 70%, staining (red) 100 ml .....	2.90	1	healthE
Soln 2% with ethanol 70%, staining (red) 100 ml .....	3.86	1	healthE
Soln 0.5% with ethanol 70%, non-staining (pink) 500 ml .....	5.45	1	healthE
Soln 0.5% with ethanol 70%, staining (red) 500 ml .....	5.90	1	healthE
Soln 2% with ethanol 70%, staining (red) 500 ml .....	9.56	1	healthE
IODINE WITH ETHANOL Soln 1% with ethanol 70%, 100 ml .....	9.30	1	healthE

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>ISOPROPYL ALCOHOL</b>			
Soln 70%, 500 ml .....	5.00	1	PSM
	5.65		healthE
<b>POVIDONE-IODINE</b>			
📌 Vaginal tab 200 mg			
<b>➡Restricted</b>			
Rectal administration pre-prostate biopsy.			
Oint 10% .....	3.27	25 g	Betadine
Soln 10% .....	2.95	100 ml	Riodine
	6.20	500 ml	Riodine
			Betadine
Soln 5%			
Soln 7.5%			
Pad 10%			
Swab set 10%			
<b>POVIDONE-IODINE WITH ETHANOL</b>			
Soln 10% with ethanol 30% .....	10.00	500 ml	Betadine Skin Prep
Soln 10% with ethanol 70%			
<b>SODIUM HYPOCHLORITE</b>			
Soln			

## Contrast Media

### Iodinated X-ray Contrast Media

<b>DIATRIZOATE MEGLUMINE WITH DIATRIZOATE SODIUM</b>			
Oral liq 660 mg per ml with diatrizoate sodium 100 mg per ml, 100 ml .....	21.00	100 ml	Gastrografin
Inj 370 mg with sodium amidotrizoate 100 mg per ml, 50 ml bottle			
Inj 146 mg with sodium amidotrizoate 40 mg per ml, 250 ml bottle .....	210.00	10	Gastrografin
<b>DIATRIZOATE SODIUM</b>			
Oral liq 370 mg per ml, 10 ml sachet .....	156.12	50	Ioscan
<b>IODISED OIL</b>			
Inj 480 mg per ml, 10 ml ampoule			
<b>IODIXANOL</b>			
Inj 270 mg per ml, 20 ml vial			
Inj 270 mg per ml, 50 ml bottle .....	223.50	10	Visipaque
Inj 270 mg per ml, 100 ml bottle .....	447.00	10	Visipaque
Inj 320 mg per ml, 20 ml vial			
Inj 320 mg per ml, 50 ml bottle .....	223.50	10	Visipaque
Inj 320 mg per ml, 100 ml bottle .....	447.00	10	Visipaque
Inj 320 mg per ml, 150 ml bottle .....	670.50	10	Visipaque
Inj 320 mg per ml, 200 ml bottle .....	565.56	6	Visipaque
	894.00	10	Visipaque

## VARIOUS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>IOHEXOL</b>			
Inj 240 mg per ml, 50 ml bottle .....	77.80	10	Omnipaque
Inj 300 mg per ml, 20 ml bottle .....	24.00	6	Omnipaque
Inj 300 mg per ml, 50 ml bottle .....	77.80	10	Omnipaque
Inj 300 mg per ml, 100 ml bottle .....	155.60	10	Omnipaque
Inj 300 mg per ml, 500 ml bottle .....	468.00	6	Omnipaque
Inj 350 mg per ml, 20 ml bottle .....	24.00	6	Omnipaque
Inj 350 mg per ml, 50 ml bottle .....	77.80	10	Omnipaque
Inj 350 mg per ml, 75 ml bottle .....	116.70	10	Omnipaque
Inj 350 mg per ml, 100 ml bottle .....	155.60	10	Omnipaque
Inj 350 mg per ml, 200 ml bottle .....	311.16	10	Omnipaque
<b>IOMEPROL</b>			
Inj 150 mg per ml, 50 ml bottle			
Inj 300 mg per ml, 20 ml vial			
Inj 300 mg per ml, 50 ml bottle			
Inj 300 mg per ml, 100 ml bottle			
Inj 350 mg per ml, 20 ml vial			
Inj 350 mg per ml, 50 ml bottle			
Inj 350 mg per ml, 75 ml bottle			
Inj 350 mg per ml, 100 ml bottle			
Inj 400 mg per ml, 50 ml bottle			
<b>IOPROMIDE</b>			
Inj 240 per ml, 50 ml bottle			
Inj 300 per ml, 20 ml vial			
Inj 300 per ml, 50 ml bottle			
Inj 370 per ml, 30 ml vial			
Inj 370 per ml, 50 ml bottle			
Inj 370 per ml, 100 ml bottle			
Inj 370 per ml, 200 ml bottle			
Inj 300 per ml, 100 ml bottle			
<b>IOTROLAN</b>			
Inj 240 mg per ml, 10 ml vial			



Price (ex man. excl. GST)		Brand or Generic
\$	Per	Manufacturer

## Non-iodinated X-ray Contrast Media

### BARIUM SULPHATE

Powder for enema 397 g			
Powder for oral liq 10,000 g			
Powder for oral liq 100 g			
Powder for oral liq 148 g			
Powder for oral liq 22.1 g			
Powder for oral liq 300 g			
Powder for oral liq 340 g			
Eosophogeal cream 30 mg per g			
Eosophogeal cream 600 mg per g			
Liq 1,000 mg per ml			
Oral liq 1 mg per ml			
Oral liq 1,250 mg per ml			
Oral liq 13 mg per ml			
Oral liq 130 mg per ml			
Oral liq 21 mg per ml			
Oral liq 400 mg per ml			
Eosophogeal paste 400 mg per ml			
Oral liq 22 mg per g, 250 ml .....	175.00	24	CT Plus+
Oral liq 22 mg per g, 450 ml .....	220.00	24	CT Plus+
Enema 1,250 mg per ml			

### CITRIC ACID WITH SODIUM BICARBONATE

Powder 382.2 mg per g with sodium bicarbonate 551.3 mg per g, 4 g sachet

*e.g. E-Z-GAS II*

## Paramagnetic Contrast Media

### GADOBENIC ACID

Inj 334 mg per ml, 10 ml vial .....	324.74	10	Multihance
Inj 334 mg per ml, 20 ml vial .....	636.28	10	Multihance

### GADOBUTROL

Inj 1 mmol per ml, 15 ml vial			
Inj 1 mmol per ml, 7.5 ml syringe .....	253.10	5	Gadovist

### GADODIAMIDE

Inj 287 mg per ml, 10 ml syringe .....	220.00	10	Omniscan
Inj 287 mg per ml, 10 ml vial .....	180.00	10	Omniscan
Inj 287 mg per ml, 5 ml vial			
Inj 287 mg per ml, 15 ml syringe .....	330.00	10	Omniscan
Inj 287 mg per ml, 15 ml vial .....	270.00	10	Omniscan
Inj 287 mg per ml, 20 ml syringe .....	440.00	10	Omniscan
Inj 287 mg per ml, 20 ml vial			

### GADOTERIC ACID

Inj 0.5 mmol per ml, 10 ml syringe
Inj 0.5 mmol per ml, 20 ml syringe
Inj 0.5 mmol per ml, 10 ml bottle
Inj 0.5 mmol per ml, 20 ml bottle
Inj 0.5 mmol per ml, 5 ml bottle

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>GADOXETATE DISODIUM</b>			
Inj 181 mg per ml, 10 ml syringe			
<b>MEGLUMINE GADOPENTETATE</b>			
Inj 469 mg per ml, 10 ml syringe .....	92.00	5	Magnevist
Inj 469 mg per ml, 10 ml vial .....	184.00	10	Magnevist
Inj 469 mg per ml, 15 ml vial			
Inj 469 mg per ml, 20 ml vial			

### Ultrasound Contrast Media

#### PERFLUTREN

Inj 1.1 mg per ml, 2 ml

### Diagnostic Agents

#### ARGININE

Inj 50 mg per ml, 500 ml bottle

Inj 100 mg per ml, 300 ml bottle

#### HISTAMINE ACID PHOSPHATE

Nebuliser soln 0.6%, 10 ml vial

Nebuliser soln 2.5%, 10 ml vial

Nebuliser soln 5%, 10 ml vial

#### METHACHOLINE CHLORIDE

Powder 100 mg

#### SECRETIN PENTAHYDROCHLORIDE

Inj 100 u ampoule

#### SINCALIDE

Inj 5 mcg per vial

#### TUBERCULIN, PURIFIED PROTEIN DERIVATIVE

Inj 5 TU per 0.1 ml, 1 ml vial

### Diagnostic Dyes

#### BONNEY'S BLUE DYE

Soln

#### INDIGO CARMINE

Inj 4 mg per ml, 5 ml ampoule

Inj 8 mg per ml, 5 ml ampoule

#### INDOCYANINE GREEN

Inj 25 mg vial

#### METHYLTHIONINIUM CHLORIDE [METHYLENE BLUE]

Inj 10 mg per ml, 10 ml ampoule

Inj 10 mg per ml, 5 ml ampoule

#### PATENT BLUE V

Inj 2.5%, 2 ml ampoule

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Irrigation Solutions</b>			
<b>CHLORHEXIDINE</b>			
Irrigation soln 0.02%, bottle .....	2.92	100 ml	Baxter
Irrigation soln 0.05%, bottle .....	3.02	100 ml	Baxter
	3.63	500 ml	Baxter
Irrigation soln 0.1%, bottle .....	3.10	100 ml	Baxter
Irrigation soln 0.5%, bottle .....	4.69	500 ml	Baxter
Irrigation soln 0.02%, 500 ml bottle			
Irrigation soln 0.1%, 30 ml ampoule			
<b>CHLORHEXIDINE WITH CETRIMIDE</b>			
Irrigation soln 0.015% with cetrimide 0.15%, 30 ml ampoule			
Irrigation soln 0.015% with cetrimide 0.15%, bottle .....	3.21	100 ml	Baxter
	3.47	500 ml	Baxter
	4.17	1,000 ml	Baxter
Irrigation soln 0.05% with cetrimide 0.5%, bottle .....	4.20	100 ml	Baxter
	3.87	500 ml	Baxter
Irrigation soln 0.1% with cetrimide 1%, bottle .....	4.38	100 ml	Baxter
	5.81	500 ml	Baxter
<b>GLYCINE</b>			
Irrigation soln 1.5%, bottle .....	11.38	2,000 ml	Baxter
	14.44	3,000 ml	Baxter
<b>SODIUM CHLORIDE</b>			
Irrigation soln 0.9%, 30 ml ampoule – 1% DV Nov-11 to 2014 .....	19.50	30 ml	<b>Pfizer</b>
Irrigation soln 0.9%, bottle .....	2.49	100 ml	Baxter
	2.88	500 ml	Baxter
	2.96	1,000 ml	Baxter
	10.00	2,000 ml	Baxter
	12.67	3,000 ml	Baxter
<b>WATER</b>			
Irrigation soln, bottle .....	2.68	100 ml	Baxter
	2.61	500 ml	Baxter
	2.75	1,000 ml	Baxter
	9.71	2,000 ml	Baxter
	15.80	3,000 ml	Baxter

## Surgical Preparations

### BISMUTH SUBNITRATE AND IODOFORM PARAFFIN

Paste

### DIMETHYL SULFOXIDE

Soln 50%

Soln 99%

### PHENOL

Inj 6%, 10 ml ampoule

### PHENOL WITH IOXAGLIC ACID

Inj 12%, 10 ml ampoule

### TROMETAMOL

Inj 36 mg per ml, 500 ml bottle

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Cardioplegia Solutions</b>			
<b>ELECTROLYTES</b>			
Inj aspartic acid 10.43 mg per ml, citric acid 0.22476 mg per ml, glutamic acid 11.53 mg per ml, sodium phosphate 0.1725 mg per ml, potassium chloride 2.15211 mg per ml, sodium citrate 1.80768 mg per ml, sodium hydroxide 6.31 mg per ml and trometamol 11.2369 mg per ml, 364 ml bag			<i>e.g. Cardioplegia Enriched Paed. Soln.</i>
Inj aspartic acid 8.481 mg per ml, citric acid 0.8188 mg per ml, glutamic acid 9.375 mg per ml, sodium phosphate 0.6285 mg per ml, potassium chloride 2.5 mg per ml, sodium citrate 6.585 mg per ml, sodium hydroxide 5.133 mg per ml and trometamol 9.097 mg per ml, 527 ml bag			<i>e.g. Cardioplegia Enriched Solution</i>
Inj citric acid 0.07973 mg per ml, sodium phosphate 0.06119 mg per ml, potassium chloride 2.181 mg per ml, sodium chloride 1.788 mg ml, sodium citrate 0.6412 mg per ml and trometamol 5.9 mg per ml, 523 ml bag			<i>e.g. Cardioplegia Base Solution</i>
Inj 110 mmol/l sodium, 16 mmol/l potassium, 1.2 mmol/l calcium, 16 mmol/l magnesium and 160 mmol/l chloride, 1,000 ml bag			<i>e.g. Cardioplegia Solution AHB7832</i>
Inj 143 mmol/l sodium, 16 mmol/l potassium, 16 mmol/l magnesium and 1.2 mmol/l calcium, 1,000 ml bag			<i>e.g. Cardioplegia Electrolyte Solution</i>
<b>MONOSODIUM GLUTAMATE WITH SODIUM ASPARTATE</b>			
Inj 42.68 mg with sodium aspartate 39.48 mg per ml, 250 ml bottle			
<b>MONOSODIUM L-ASPARTATE</b>			
Inj 14 mmol per 10 ml, 10 ml			
<b>Cold Storage Solutions</b>			
<b>SODIUM WITH POTASSIUM</b>			
Inj 29 mmol/l with potassium 125 mmol/l, 1,000 ml bag			

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
<b>Extemporaneously Compounded Preparations</b>		
ACETIC ACID Liq		
ALUM Powder BP		
ARACHIS OIL [PEANUT OIL] Liq		
ASCORBIC ACID Powder		
BENZOIN Tincture compound BP		
BISMUTH SUBGALLATE Powder		
BORIC ACID Powder		
CARBOXYMETHYLCELLULOSE Soln 1.5%		
CETRIMIDE Soln 40%		
CHLORHEXIDINE GLUCONATE Soln 20 %		
CHLOROFORM Liq BP		
CITRIC ACID Powder BP		
CLOVE OIL Liq		
COAL TAR Soln BP		
CODEINE PHOSPHATE Powder		
COLLODION FLEXIBLE Liq		
COMPOUND HYDROXYBENZOATE Soln		
CYSTEAMINE HYDROCHLORIDE Powder		
DISODIUM HYDROGEN PHOSPHATE WITH SODIUM DIHYDROGEN PHOSPHATE Inj 37.46 mg with sodium dihydrogen phosphate 47.7 mg in 1.5 ml ampoule		
DITHRANOL Powder		

# EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GLUCOSE			
Powder			
GLYCERIN WITH SODIUM SACCHARIN			
Suspension .....	35.50	473 ml	Ora-Sweet SF
GLYCERIN WITH SUCROSE			
Suspension .....	35.50	473 ml	Ora-Sweet
GLYCEROL			
Liq .....	19.80	2,000 ml	ABM
HYDROCORTISONE			
Powder – 1% DV Nov-11 to 2014 .....	44.00	25 g	ABM
LACTOSE			
Powder			
MAGNESIUM HYDROXIDE			
Paste			
MENTHOL			
Crystals			
METHADONE HYDROCHLORIDE			
Powder			
METHYL HYDROXYBENZOATE			
Powder			
METHYLCELLULOSE			
Powder			
Suspension .....	35.50	473 ml	Ora-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN			
Suspension .....	35.50	473 ml	Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE			
Suspension .....	35.50	473 ml	Ora-Blend
OLIVE OIL			
Liq			
PARAFFIN			
Liq			
PHENOBARBITONE SODIUM			
Powder			
PHENOL			
Liq			
PILOCARPINE NITRATE			
Powder			
POLYHEXAMETHYLENE BIGUANIDE			
Liq			
POVIDONE K30			
Powder			
PROPYLENE GLYCOL			
Liq .....	12.00	500 ml	ABM

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SALICYLIC ACID Powder			
SILVER NITRATE Crystals			
SODIUM BICARBONATE Powder BP			
SODIUM CITRATE Powder			
SODIUM METABISULFITE Powder			
STARCH Powder			
SULPHUR Precipitated Sublimed			
SYRUP Liq (pharmaceutical grade) .....	21.75	2,000 ml	Midwest
TRI-SODIUM CITRATE Crystals			
TRICHLORACETIC ACID Grans			
UREA Powder BP			
WOOL FAT Oint, anhydrous			
XANTHAN Gum 1%			
ZINC OXIDE Powder			

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

## Food Modules

### Carbohydrate

#### ➡ Restricted

##### Use as an additive

Any of the following:

- 1 Cystic fibrosis; or
- 2 Chronic kidney disease; or
- 3 Cancer in children; or
- 4 Cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 5 Faltering growth in an infant/child; or
- 6 Bronchopulmonary dysplasia; or
- 7 Premature and post premature infant; or
- 8 Inborn errors of metabolism.

##### Use as a module

For use as a component in a modular formula

CARBOHYDRATE SUPPLEMENT – **Restricted** see terms above

⬆ Powder 95 g carbohydrate per 100 g, 368 g can

⬆ Powder 96 g carbohydrate per 100 g, 400 g can

*e.g. Polycal*

### Fat

#### ➡ Restricted

##### Use as an additive

Any of the following:

- 1 Patient has inborn errors of metabolism; or
- 2 Faltering growth in an infant/child; or
- 3 Bronchopulmonary dysplasia; or
- 4 Fat malabsorption; or
- 5 Lymphangiectasia; or
- 6 Short bowel syndrome; or
- 7 Infants with necrotising enterocolitis; or
- 8 Biliary atresia; or
- 9 For use in a ketogenic diet; or
- 10 Chyle leak; or
- 11 Ascites; or
- 12 Patient has increased energy requirements, and for whom dietary measures have not been successful.

##### Use as a module

For use as a component in a modular formula

LONG-CHAIN TRIGLYCERIDE SUPPLEMENT – **Restricted** see terms above

⬆ Liquid 50 g fat per 100 ml, 200 ml bottle

*e.g. Calogen*

⬆ Liquid 50 g fat per 100 ml, 500 ml bottle

*e.g. Calogen*

MEDIUM-CHAIN TRIGLYCERIDE SUPPLEMENT – **Restricted** see terms above

⬆ Liquid 50 g fat per 100 ml, 250 ml bottle

*e.g. Liquigen*

⬆ Liquid 95 g fat per 100 ml, 500 ml bottle

*e.g. MCT Oil*

WALNUT OIL – **Restricted** see terms above

⬆ Liq



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Protein</b>			
➡ <b>Restricted</b>			
<b>Use as an additive</b>			
Either:			
1 Protein losing enteropathy; or			
2 High protein needs.			
<b>Use as a module</b>			
For use as a component in a modular formula			
PROTEIN SUPPLEMENT – <b>Restricted</b> see terms above			
⬆ Powder 5 g protein, 0.67 g carbohydrate and 0.6 g fat per 6.6 g, 275 g can			<i>e.g. Promod</i>
⬆ Powder 6 g protein per 7 g, can .....	8.95	227 g	Resource Beneprotein
⬆ Powder 89 g protein, <1.5 g carbohydrate and 2 g fat per 100 g, 225 g can			<i>e.g. Protifar</i>
<b>Other Supplements</b>			
BREAST MILK FORTIFIER			
Powder 0.2 g protein, 0.7 g carbohydrate and 0.02 g fat per 1 g sachet			<i>e.g. FM 85</i>
Powder 0.5 g protein, 1.2 g carbohydrate and 0.08 g fat per 2 g sachet			<i>e.g. S26 Human Milk Fortifier</i>
Powder 0.6 g protein and 1.4 g carbohydrate per 2.2 g sachet			<i>e.g. Nutricia Breast Milk Fortifier</i>
CARBOHYDRATE AND FAT SUPPLEMENT – <b>Restricted</b> see terms below			
⬆ Powder 72.7 g carbohydrate and 22.3 g fat per 100 g, 400 g can			<i>e.g. Super Soluble Duocal</i>
➡ <b>Restricted</b>			
Both:			
1 Infant or child aged four years or under; and			
2 Any of the following:			
2.1 Cystic fibrosis; or			
2.2 Cancer in children; or			
2.3 Faltering growth; or			
2.4 Bronchopulmonary dysplasia; or			
2.5 Premature and post premature infants.			
<b>Food/Fluid Thickeners</b>			
<b>NOTE:</b>			
While pre-thickened drinks and supplements have not been included in Section H, DHB hospitals may continue to use such products for patients with dysphagia, provided that:			
<ul style="list-style-type: none"> <li>• use was established prior to 1 July 2013; and</li> <li>• the product has not been specifically considered and excluded by PHARMAC; and</li> <li>• use of the product conforms to any applicable indication restrictions for similar products that are listed in Section H (for example, use of thickened high protein products should be in line with the restriction for high protein oral feed in Section H).</li> </ul>			
PHARMAC intends to make a further decision in relation to pre-thickened drinks and supplements in the future, and will notify of any change to this situation.			
CAROB BEAN GUM WITH MAIZE STARCH AND MALTODEXTRIN			
Powder			<i>e.g. Feed Thickener Karicare Aptamil</i>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GUAR GUM Powder			<i>e.g. Guarcol</i>
MAIZE STARCH Powder			<i>e.g. Resource Thicken Up; Nutilis</i>
MALTODEXTRIN WITH XANTHAN GUM Powder			<i>e.g. Instant Thick</i>
MALTODEXTRIN WITH XANTHAN GUM AND ASCORBIC ACID Powder			<i>e.g. Easy Thick</i>

## Metabolic Products

### ➡ Restricted

Any of the following:

- 1 For the dietary management of homocystinuria, maple syrup urine disease, phenylketonuria (PKU), glutaric aciduria, isovaleric acidemia, propionic acidemia, methylmalonic acidemia, tyrosinemia or urea cycle disorders; or
- 2 Patient has adrenoleukodystrophy; or
- 3 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

### Glutaric Aciduria Type 1 Products

AMINO ACID FORMULA (WITHOUT LYSINE AND LOW TRYPTOPHAN) – **Restricted** see terms above

- ⚡ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can *e.g. GA1 Anamix Infant*
- ⚡ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can *e.g. XLYS Low TRY  
Maxamaid*

### Homocystinuria Products

AMINO ACID FORMULA (WITHOUT METHIONINE) – **Restricted** see terms above

- ⚡ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can *e.g. HCU Anamix Infant*
- ⚡ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can *e.g. XMET Maxamaid*
- ⚡ Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can *e.g. XMET Maxamum*
- ⚡ Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle *e.g. HCU Anamix Junior  
LQ*

### Isovaleric Acidemia Products

AMINO ACID FORMULA (WITHOUT LEUCINE) – **Restricted** see terms above

- ⚡ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can *e.g. IVA Anamix Infant*
- ⚡ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can *e.g. XLEU Maxamaid*
- ⚡ Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can *e.g. XLEU Maxamum*

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

## Maple Syrup Urine Disease Products

AMINO ACID FORMULA (WITHOUT ISOLEUCINE, LEUCINE AND VALINE) – **Restricted** see terms on the preceding page

⬆ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can		<i>e.g. MSUD Anamix Infant</i>
⬆ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can		<i>e.g. MSUD Maxamaid</i>
⬆ Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can		<i>e.g. MSUD Maxamum</i>
⬆ Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle		<i>e.g. MSUD Anamix Junior LQ</i>

## Phenylketonuria Products

AMINO ACID FORMULA (WITHOUT PHENYLALANINE) – **Restricted** see terms on the preceding page

⬆ Tab 8.33 mg		<i>e.g. Phlexy-10</i>
⬆ Powder 29 g protein, 38 g carbohydrate and 13.5 g fibre per 100 g, 29 g sachet		<i>e.g. PKU Anamix Junior</i>
⬆ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can		<i>e.g. PKU Anamix Infant</i>
⬆ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can		<i>e.g. XP Maxamaid</i>
⬆ Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can		<i>e.g. XP Maxamum</i>
⬆ Powder 8.33 g protein and 8.8 g carbohydrate per 20 g sachet		<i>e.g. Phlexy-10</i>
⬆ Liquid 10 g protein, 4.4 g carbohydrate and 0.25 g fibre per 100 ml, 62.5 ml bottle		<i>e.g. PKU Lophlex LQ 10</i>
⬆ Liquid 20 g protein, 8.8 g carbohydrate and 0.34 g fibre per 100 ml, 125 ml bottle		<i>e.g. PKU Lophlex LQ 20</i>
⬆ Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, bottle .....	13.10	125 ml PKU Anamix Junior LQ (Berry)
		PKU Anamix Junior LQ (Orange)
		PKU Anamix Junior LQ (Unflavoured)
⬆ Liquid 16 g protein, 7 g carbohydrate and 0.27 g fibre per 100 ml, 125 ml bottle		<i>e.g. PKU Lophlex LQ 20</i>
⬆ Liquid 16 g protein, 7 g carbohydrate and 0.27 g fibre per 100 ml, 62.5 ml bottle		<i>e.g. PKU Lophlex LQ 10</i>
⬆ Liquid 16 g protein, 7 g carbohydrate and 0.4 g fibre per 100 ml, 125 ml bottle		<i>e.g. PKU Lophlex LQ 20</i>
⬆ Liquid 16 g protein, 7 g carbohydrate and 0.4 g fibre per 100 ml, 62.5 ml bottle		<i>e.g. PKU Lophlex LQ 10</i>
⬆ Liquid 6.7 g protein, 5.1 g carbohydrate and 2 g fat per 100 ml, 250 ml carton		<i>e.g. Easiphen</i>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Propionic Acidemia and Methylmalonic Acidemia Products</b>			
AMINO ACID FORMULA (WITHOUT ISOLEUCINE, METHIONINE, THREONINE AND VALINE) – <b>Restricted</b> see terms on page 186			
⚡ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can			e.g. MMA/PA Anamix Infant
⚡ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can			e.g. XMTVI Maxamaid
⚡ Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can			e.g. XMTVI Maxamum
<b>Protein Free Supplements</b>			
PROTEIN FREE SUPPLEMENT – <b>Restricted</b> see terms on page 186			
⚡ Powder nil added protein and 67 g carbohydrate per 100 g, 400 g can			e.g. Energivit
<b>Tyrosinaemia Products</b>			
AMINO ACID FORMULA (WITHOUT PHENYLALANINE AND TYROSINE) – <b>Restricted</b> see terms on page 186			
⚡ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can			e.g. TYR Anamix Infant
⚡ Powder 25 g protein and 51 g carbohydrate per 100 g, 400 g can			e.g. XPHEN, TYR Maxamaid
⚡ Powder 29 g protein, 38 g carbohydrate and 13.5 g fat per 100 g, 29 g sachet			e.g. TYR Anamix Junior
⚡ Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle			e.g. TYR Anamix Junior LQ
<b>Urea Cycle Disorders Products</b>			
AMINO ACID SUPPLEMENT – <b>Restricted</b> see terms on page 186			
⚡ Powder 25 g protein and 65 g carbohydrate per 100 g, 200 g can			e.g. Dialamine
⚡ Powder 79 g protein per 100 g, 200 g can			e.g. Essential Amino Acid Mix
<b>X-Linked Adrenoleukodystrophy Products</b>			
GLYCEROL TRIERUCATE – <b>Restricted</b> see terms on page 186			
⚡ Liquid, 1,000 ml bottle			
GLYCEROL TRIOLEATE – <b>Restricted</b> see terms on page 186			
⚡ Liquid, 500 ml bottle			
<b>Specialised Formulas</b>			
<b>Diabetic Products</b>			
➡ <b>Restricted</b>			
Any of the following:			
1 For patients with type I or type II diabetes suffering weight loss and malnutrition that requires nutritional support; or			
2 For patients with pancreatic insufficiency; or			
3 For patients who have, or are expected to, eat little or nothing for 5 days;			
4 For patients who have a poor absorptive capacity and/or high nutrient losses and/or increased nutritional needs from causes such as catabolism; or			
continued...			
⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below) e.g. Brand indicates brand example only. It is not a contracted product.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued. . .			
5 For use pre- and post-surgery; or			
6 For patients being tube-fed; or			
7 For tube-feeding as a transition from intravenous nutrition.			
LOW-GI ENTERAL FEED 1 KCAL/ML – <b>Restricted</b> see terms on the preceding page			
⬆ Liquid 5 g protein, 9.6 g carbohydrate and 5.4 g fat per 100 ml, 1,000 ml bottle .....	7.50	1,000 ml	Glucerna Select RTH (Vanilla)
⬆ Liquid 4.3 g protein, 11.3 g carbohydrate and 4.2 g fat per 100 ml, 1,000 ml bag			<i>e.g. Nutrison Advanced Diason</i>
LOW-GI ORAL FEED 1 KCAL/ML – <b>Restricted</b> see terms on the preceding page			
⬆ Liquid 4.5 g protein, 9.8 g carbohydrate, 4.4 g fat and 1.9 g fibre per 100 ml, can .....	2.10	237 ml	Sustagen Diabetic (Vanilla)
⬆ Liquid 5 g protein, 9.6 g carbohydrate and 5.4 g fat per 100 ml, 250 ml bottle .....	1.88	250 ml	Glucerna Select (Vanilla)
⬆ Liquid 6 g protein, 9.5 g carbohydrate, 4.7 g fat and 2.6 g fibre per 100 ml, can .....	2.10	237 ml	Resource Diabetic (Vanilla)
⬆ Liquid 4.9 g protein, 11.7 g carbohydrate, 3.8 g fat and 2 g fibre per 100 ml, 200 ml bottle			<i>e.g. Diasip</i>

## Elemental and Semi-Elemental Products

### ➡ **Restricted**

Any of the following:

- 1 Malabsorption; or
- 2 Short bowel syndrome; or
- 3 Enterocutaneous fistulas; or
- 4 Eosinophilic enteritis (including oesophagitis); or
- 5 Inflammatory bowel disease; or
- 6 Acute pancreatitis where standard feeds are not tolerated; or
- 7 Patients with multiple food allergies requiring enteral feeding.

AMINO ACID ORAL FEED – **Restricted** see terms above

⬆ Powder 11.5 g protein, 61.7 g carbohydrate and 0.8 g fat per sachet .....	4.50	80.4 g	Vivonex TEN
---	------	--------	-------------

AMINO ACID ORAL FEED 0.8 KCAL/ML – **Restricted** see terms above

⬆ Liquid 2.5 g protein, 11 g carbohydrate and 3.5 g fat per 100 ml, 250 ml carton			<i>e.g. Elemental 028 Extra</i>
---	--	--	---------------------------------

PEPTIDE-BASED ENTERAL FEED 1 KCAL/ML – **Restricted** see terms above

⬆ Liquid 4 g protein, 17.6 g carbohydrate and 1.7 g fat per 100 ml, 1,000 ml bag			<i>e.g. Nutrison Advanced Peptisorb</i>
--	--	--	---

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PEPTIDE-BASED ORAL FEED – Restricted</b> see terms on the preceding page			
⬆ Powder 12.5 g protein, 55.4 g carbohydrate and 3.25 g fat per sachet .....	4.40	79 g	Vital HN
⬆ Powder 13.7 g protein, 62.9 g carbohydrate and 17.5 g fat per 100 g, 400 g can			<i>e.g. Peptamen Junior</i>
⬆ Powder 13.8 g protein, 59 g carbohydrate and 18 g fat per 100 g, 400 g can			<i>e.g. MCT Peptide; MCT Peptide 1+</i>
⬆ Powder 15.8 g protein, 49.5 g carbohydrate and 4.65 g fat per 76 g sachet .....	7.50	76 g	Alitraq
<b>PEPTIDE-BASED ORAL FEED 1 KCAL/ML – Restricted</b> see terms on the preceding page			
⬆ Liquid 5 g protein, 16 g carbohydrate and 1.69 g fat per 100 ml, carton .....	4.95	237 ml	Peptamen OS 1.0 (Vanilla)

## Fat Modified Products

**FAT-MODIFIED FEED – Restricted** see terms below

⬆ Powder 11.4 g protein, 68 g carbohydrate and 11.8 g fat per 100 g, 400 g can			<i>e.g. Monogen</i>
---	--	--	---------------------

### ➡Restricted

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed for adults.

## Hepatic Products

### ➡Restricted

For children (up to 18 years) who require a liver transplant

**HEPATIC ORAL FEED – Restricted** see terms above

⬆ Powder 11 g protein, 64 g carbohydrate and 20 g fat per 100 g, can .....	78.97	400 g	Heparon Junior
--	-------	-------	----------------

## High Calorie Products

### ➡Restricted

Any of the following:

- 1 Patient is fluid volume or rate restricted; or
- 2 Patient requires low electrolyte; or
- 3 Both:
  - 3.1 Any of the following:
    - 3.1.1 Cystic fibrosis; or
    - 3.1.2 Any condition causing malabsorption; or
    - 3.1.3 Faltering growth in an infant/child; or
    - 3.1.4 Increased nutritional requirements; and
  - 3.2 Patient has substantially increased metabolic requirements.

**ENTERAL FEED 2 KCAL/ML – Restricted** see terms above

⬆ Liquid 7.5 g protein, 20 g carbohydrate and 10 g fat per 100 ml, bottle .....	5.50	500 ml	Nutrison Concentrated
⬆ Liquid 8.4 g protein, 21.9 g carbohydrate, 9.1 g fat and 0.5 g fibre per 100 ml, bottle .....	11.00	1,000 ml	TwoCal HN RTH (Vanilla)

**ORAL FEED 2 KCAL/ML – Restricted** see terms above

⬆ Liquid 8.4 g protein, 22.4 g carbohydrate, 8.9 g fat and 0.8 g fibre per 100 ml, bottle .....	1.90	200 ml	Two Cal HN
--	------	--------	------------

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

## High Protein Products

HIGH PROTEIN ENTERAL FEED 1.25 KCAL/ML – **Restricted** see terms below

☞ Liquid 6.3 g protein, 14.2 g carbohydrate and 4.9 g fat per 100 ml,  
1,000 ml bag

*e.g. Nutrison Protein  
Plus*

### ☞ Restricted

Both:

- 1 The patient has a high protein requirement; and
- 2 Any of the following:
  - 2.1 Patient has liver disease; or
  - 2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
  - 2.3 Patient is fluid restricted; or
  - 2.4 Patient's needs cannot be more appropriately met using high calorie product.

HIGH PROTEIN ENTERAL FEED 1.28 KCAL/ML – **Restricted** see terms below

☞ Liquid 6.3 g protein, 14.1 g carbohydrate, 4.9 g fat and 1.5 g fibre per  
100 ml, 1,000 ml bag

*e.g. Nutrison Protein  
Plus Multi Fibre*

### ☞ Restricted

Both:

- 1 The patient has a high protein requirement; and
- 2 Any of the following:
  - 2.1 Patient has liver disease; or
  - 2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
  - 2.3 Patient is fluid restricted; or
  - 2.4 Patient's needs cannot be more appropriately met using high calorie product.

HIGH PROTEIN ORAL FEED 1 KCAL/ML – **Restricted** see terms below

☞ Liquid 10 g protein, 10.3 g carbohydrate and 2.1 g fat per 100 ml,  
200 ml bottle

*e.g. Fortimel Regular*

### ☞ Restricted

Any of the following:

- 1 Decompensating liver disease without encephalopathy; or
- 2 Protein losing gastro-enteropathy; or
- 3 Patient has increased protein requirements without increased energy requirements.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Infant Formulas</b>			
AMINO ACID FORMULA – <b>Restricted</b> see terms below			
☞ Powder 1.95 g protein, 8.1 g carbohydrate and 3.5 g fat per 100 ml, 400 g can			<i>e.g. Neocate</i>
☞ Powder 13 g protein, 52.5 g carbohydrate and 24.5 g fat per 100 g, 400 g can			<i>e.g. Neocate LCP</i>
☞ Powder 13.5 g protein, 52 g carbohydrate and 24.5 g fat per 100 g, can .....53.00	400 g		Neocate Gold (Unflavoured)
☞ Powder 14 g protein, 50 g carbohydrate and 24.3 g fat per 100 g, 400 g can			<i>e.g. Neocate Advance</i>
☞ Powder 16 g protein, 51.4 g carbohydrate and 21 g fat per 100 g, can .....53.00	400 g		Neocate Advance (Vanilla)
☞ Powder 2.2 g protein, 7.8 g carbohydrate and 3.4 g fat per 100 ml, can .....53.00	400 g		Elecare LCP (Unflavoured)
☞ Powder 2.2 g protein, 7.8 g carbohydrate and 3.4 g fat per 100 ml, can .....53.00	400 g		Elecare (Unflavoured) Elecare (Vanilla)
☞ Powder 6 g protein, 31.5 g carbohydrate and 5.88 g fat per sachet .....6.00	48.5 g		Vivonex Paediatric

#### ☞**Restricted**

##### Initiation

Any of the following:

- 1 Extensively hydrolysed formula has been reasonably trialled and is inappropriate due to documented severe intolerance or allergy or malabsorption; or
- 2 History of anaphylaxis to cows' milk protein formula or dairy products; or
- 3 Eosinophilic oesophagitis.

##### Continuation

Both:

- 1 An assessment as to whether the infant can be transitioned to a cows' milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an amino acid infant formula.

EXTENSIVELY HYDROLYSED FORMULA – **Restricted** see terms below

- ☞ Powder 14 g protein, 53.4 g carbohydrate and 27.3 g fat per 100 g, 450 g can

*e.g. Gold Pepti Junior  
Karicare Aptamil*

#### ☞**Restricted**

##### Initiation - new patients

Any of the following:

- 1 Both:
  - 1.1 Cows' milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
  - 1.2 Either:
    - 1.2.1 Soy milk formula has been trialled without resolution of symptoms; or
    - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malabsorption; or
- 7 Cystic fibrosis; or

continued...



	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued. . .			
8 Proven fat malabsorption; or			
9 Severe intestinal motility disorders causing significant malabsorption; or			
10 Intestinal failure.			
<b>Initiation - step down from amino acid formula</b>			
Both:			
1 The infant is currently receiving funded amino acid formula; and			
2 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula.			
<b>Continuation</b>			
Both:			
1 An assessment as to whether the infant can be transitioned to a cows' milk protein or soy infant formula has been undertaken; and			
2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula.			
<b>FRUCTOSE-BASED FORMULA</b>			
Powder 14.6 g protein, 49.7 g carbohydrate and 30.8 g fat per 100 g, 400 g can			<i>e.g. Galactomin 19</i>
<b>LACTOSE-FREE FORMULA</b>			
Powder 1.3 g protein, 7.3 g carbohydrate and 3.5 g fat per 100 ml, 900 g can			<i>e.g. Karicare Aptamil Gold De-Lact</i>
Powder 1.5 g protein, 7.2 g carbohydrate and 3.6 g fat per 100 ml, 900 g can			<i>e.g. S26 Lactose Free</i>
<b>LOW-CALCIUM FORMULA</b>			
Powder 14.6 g protein, 53.7 g carbohydrate and 26.1 g fat per 100 g, 400 g can			<i>e.g. Locasol</i>
<b>PAEDIATRIC ORAL FEED 1 KCAL/ML – <b>Restricted</b> see terms below</b>			
⚡ Liquid 2.6 g protein, 10.3 g carbohydrate, 5.4 g fat and 0.6 g fibre per 100 ml, 100 ml bottle			<i>e.g. Infatrini</i>
<b>➡Restricted</b>			
Both:			
1 Either:			
1.1 The patient is fluid restricted; or			
1.2 The patient has increased nutritional requirements due to faltering growth; and			
2 Patient is under 18 months old and weighs less than 8kg.			
<b>PRETERM FORMULA – <b>Restricted</b> see terms below</b>			
⚡ Powder 1.9 g protein, 7.5 g carbohydrate and 3.9 g fat per 14 g, can .....15.25	400 g		S-26 Gold Premgro
⚡ Liquid 2.2 g protein, 8.4 g carbohydrate and 4.4 g fat per 100 ml, bottle .....0.75	100 ml		S26 LBW Gold RTF
⚡ Liquid 2.3 g protein, 8.6 g carbohydrate and 4.2 g fat per 100 ml, 90 ml bottle			<i>e.g. Pre Nan Gold RTF</i>
⚡ Liquid 2.6 g protein, 8.4 g carbohydrate and 3.9 g fat per 100 ml, 70 ml bottle			<i>e.g. Karicare Aptamil Gold+Preterm</i>
<b>➡Restricted</b>			
For infants born before 33 weeks' gestation or weighing less than 1.5 kg at birth.			
<b>THICKENED FORMULA</b>			
Powder 1.8 g protein, 8.1 g carbohydrate and 3.3 g fat per 100 ml, 900 g can			<i>e.g. Karicare Aptamil Thickened AR</i>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Ketogenic Diet Products</b>			
HIGH FAT FORMULA – <b>Restricted</b> see terms below			
☛ Powder 15.25 g protein, 3 g carbohydrate and 73 g fat per 100 g, can .....	35.50	300 g	Ketocal 4:1 (Unflavoured) Ketocal 4:1 (Vanilla)
☛ Powder 15.3 g protein, 7.2 g carbohydrate and 67.7 g fat per 100 g, can .....	35.50	300 g	Ketocal 3:1 (Unflavoured)
➡ <b>Restricted</b>			
For patients with intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.			
<b>Paediatric Products</b>			
➡ <b>Restricted</b>			
Both:			
1 Child is aged one to ten years; and			
2 Any of the following:			
2.1 The child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or			
2.2 Any condition causing malabsorption; or			
2.3 Faltering growth in an infant/child; or			
2.4 Increased nutritional requirements; or			
2.5 The child is being transitioned from TPN or tube feeding to oral feeding.			
PAEDIATRIC ORAL FEED – <b>Restricted</b> see terms above			
☛ Powder 14.9 g protein, 54.3 g carbohydrate and 24.7 g fat per 100 g, can .....	20.00	850 g	Pediasure (Vanilla)
PAEDIATRIC ENTERAL FEED 0.76 KCAL/ML – <b>Restricted</b> see terms above			
☛ Liquid 2.5 g protein, 12.5 g carbohydrate, 3.3 g fat and 0.7 g fibre per 100 ml, bag .....	4.00	500 ml	Nutrini Low Energy Multifibre RTH
PAEDIATRIC ENTERAL FEED 1 KCAL/ML – <b>Restricted</b> see terms above			
☛ Liquid 2.8 g protein, 11.2 g carbohydrate and 5 g fat per 100 ml, bag .....	2.68	500 ml	Pediasure RTH
☛ Liquid 2.8 g protein, 12.3 g carbohydrate and 4.4 g fat per 100 ml, 500 ml bag			<i>e.g. Nutrini RTH</i>
PAEDIATRIC ENTERAL FEED 1.5 KCAL/ML – <b>Restricted</b> see terms above			
☛ Liquid 4.1 g protein, 18.5 g carbohydrate, 6.7 g fat and 0.8 g fibre per 100 ml, bag .....	6.00	500 ml	Nutrini Energy Multi Fibre
☛ Liquid 4.1 g protein, 18.5 g carbohydrate and 6.7 g fat per 100 ml, 500 ml bag			<i>e.g. Nutrini Energy RTH</i>
PAEDIATRIC ORAL FEED 1 KCAL/ML – <b>Restricted</b> see terms above			
☛ Liquid 4.2 g protein, 16.7 g carbohydrate and 7.5 g fat per 100 ml, bottle .....	1.07	200 ml	Pediasure (Chocolate) Pediasure (Strawberry) Pediasure (Vanilla) Pediasure (Vanilla)
☛ Liquid 4.2 g protein, 16.7 g carbohydrate and 7.5 g fat per 100 ml, can .....	1.34	250 ml	
PAEDIATRIC ORAL FEED 1.5 KCAL/ML – <b>Restricted</b> see terms above			
☛ Liquid 3.4 g protein, 18.8 g carbohydrate and 6.8 g fat per 100 ml, 200 ml bottle			<i>e.g. Fortini</i>
☛ Liquid 4.0 g protein, 18.8 g carbohydrate, 6.8 g fat and 1.5 g fibre per 100 ml, 200 ml bottle			<i>e.g. Fortini Multifibre</i>

☛ Item restricted (see ➡ above); ☛ Item restricted (see ➡ below)

*e.g. Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Renal Products</b>			
LOW ELECTROLYTE ENTERAL FEED 1.8 KCAL/ML – <b>Restricted</b> see terms below			
☞ Liquid 8.1 g protein, 14.74 g carbohydrate, 9.77 g fat and 1.26 g fibre per 100 ml, bottle .....	6.08	500 ml	Nepro HP RTH
☛ <b>Restricted</b>			
For patients with acute or chronic kidney disease			
LOW ELECTROLYTE ENTERAL FEED 2 KCAL/ML – <b>Restricted</b> see terms below			
☞ Liquid 7 g protein, 20.6 g carbohydrate, 9.6 g fat and 1.56 g fibre per 100 ml, bottle .....	6.08	500 ml	Nepro RTH
<i>(Nepro RTH Liquid 7 g protein, 20.6 g carbohydrate, 9.6 g fat and 1.56 g fibre per 100 ml, bottle to be delisted 1 August 2014)</i>			
☛ <b>Restricted</b>			
For patients with acute or chronic kidney disease			
LOW ELECTROLYTE ORAL FEED – <b>Restricted</b> see terms below			
☞ Powder 7.5 g protein, 59 g carbohydrate and 26.3 g fat per 100 g, 400 g can .....			<i>e.g. Kindergen</i>
☛ <b>Restricted</b>			
For children (up to 18 years) with acute or chronic kidney disease			
LOW ELECTROLYTE ORAL FEED 1.8 KCAL/ML			
☞ Liquid 8 g protein, 14.74 g carbohydrate, 9.77 g fat and 1.26 g fibre per 100 ml, carton .....	2.67	220 ml	Nepro HP (Strawberry) Nepro HP (Vanilla)
☛ <b>Restricted</b>			
For patients with acute or chronic kidney disease.			
LOW ELECTROLYTE ORAL FEED 2 KCAL/ML – <b>Restricted</b> see terms below			
☞ Liquid 7 g protein, 20.6 g carbohydrate, 9.6 g fat and 1.56 g fibre per 100 ml, carton .....	2.43	200 ml	Nepro (Strawberry) Nepro (Vanilla)
☞ Liquid 9.1 g protein, 19 g carbohydrate and 10 g fat per 100 ml, carton .....	3.31	237 ml	Novasource Renal (Vanilla)
☞ Liquid 3 g protein, 25.5 g carbohydrate and 9.6 g fat per 100 ml, 237 ml bottle .....			<i>e.g. Suplena</i>
☞ Liquid 7.5 g protein, 20 g carbohydrate and 10 g fat per 100 ml, 125 ml carton .....			<i>e.g. Renilon 7.5</i>
<i>(Nepro (Strawberry) Liquid 7 g protein, 20.6 g carbohydrate, 9.6 g fat and 1.56 g fibre per 100 ml, carton to be delisted 1 August 2014)</i>			
<i>(Nepro (Vanilla) Liquid 7 g protein, 20.6 g carbohydrate, 9.6 g fat and 1.56 g fibre per 100 ml, carton to be delisted 1 August 2014)</i>			
☛ <b>Restricted</b>			
For patients with acute or chronic kidney disease.			
<b>Respiratory Products</b>			
LOW CARBOHYDRATE ORAL FEED 1.5 KCAL/ML – <b>Restricted</b> see terms below			
☞ Liquid 6.2 g protein, 10.5 g carbohydrate and 9.32 g fat per 100 ml, bottle .....	1.66	237 ml	Pulmocare (Vanilla)
☛ <b>Restricted</b>			
For patients with CORD and hypercapnia, defined as a CO <sub>2</sub> value exceeding 55 mmHg			
Products with Hospital Supply Status (HSS) are in <b>bold</b>			
Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Surgical Products</b>			
HIGH ARGININE ORAL FEED 1.4 KCAL/ML – <b>Restricted</b> see terms below			
☞ Liquid 7.6 g protein, 18.9 g carbohydrate, 3.9 g fat and 1.4 g fibre per 100 ml, carton .....	4.00	237 ml	Impact Advanced Recovery (Chocolate) Impact Advanced Recovery (Vanilla)

➡ **Restricted**

Three packs per day for 5 to 7 days prior to major gastrointestinal, head or neck surgery

**Standard Feeds**➡ **Restricted**

Any of the following:

- For patients with malnutrition, defined as any of the following:
  - BMI < 18.5; or
  - Greater than 10% weight loss in the last 3-6 months; or
  - BMI < 20 with greater than 5% weight loss in the last 3-6 months; or
- For patients who have, or are expected to, eat little or nothing for 5 days; or
- For patients who have a poor absorptive capacity and/or high nutrient losses and/or increased nutritional needs from causes such as catabolism; or
- For use pre- and post-surgery; or
- For patients being tube-fed; or
- For tube-feeding as a transition from intravenous nutrition; or
- For any other condition that meets the community Special Authority criteria.

ENTERAL FEED 1.5 KCAL/ML – **Restricted** see terms above

☞ Liquid 5.4 g protein, 13.6 g carbohydrate and 3.3 g fat per 100 ml, 1,000 ml bottle			<i>e.g. Isosource Standard RTH</i>
☞ Liquid 6 g protein, 18.3 g carbohydrate and 5.8 g fat per 100 ml, bag .....	7.00	1,000 ml	Nutrison Energy
☞ Liquid 6 g protein, 18.4 g carbohydrate, 5.8 g fat and 1.5 g fibre per 100 ml, 1,000 ml bag			<i>e.g. Nutrison Energy Multi Fibre</i>
☞ Liquid 6.25 g protein, 20 g carbohydrate and 5 g fat per 100 ml, can .....	1.75	250 ml	Ensure Plus HN
☞ Liquid 6.27 g protein, 20.4 g carbohydrate and 4.9 g fat per 100 ml, bag .....	7.00	1,000 ml	Ensure Plus HN RTH
☞ Liquid 6.38 g protein, 21.1 g carbohydrate, 4.9 g fat and 1.2 g fibre per 100 ml, bag .....	7.00	1,000 ml	Jevity HiCal RTH

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>ENTERAL FEED 1 KCAL/ML – Restricted</b> see terms on the preceding page			
☛ Liquid 4 g protein, 13.6 g carbohydrate and 3.4 g fat per 100 ml, bottle .....	2.65	500 ml	Osmolite RTH
	5.29	1,000 ml	Osmolite RTH
☛ Liquid 4 g protein, 13.6 g carbohydrate and 3.4 g fat per 100 ml, can .....	1.24	250 ml	Osmolite
☛ Liquid 4 g protein, 14.1 g carbohydrate, 3.47 g fat and 1.76 g fibre per 100 ml, bottle .....	2.65	500 ml	Jevity RTH
	5.29	1,000 ml	Jevity RTH
☛ Liquid 4 g protein, 14.1 g carbohydrate, 3.47 g fat and 1.76 g fibre per 100 ml, can .....	1.32	237 ml	Jevity
☛ Liquid 4 g protein, 12.3 g carbohydrate and 3.9 g fat per 100 ml, 1,000 ml bag			<i>e.g. NutrisonStdRTH; NutrisonLowSodium</i>
☛ Liquid 4 g protein, 12.3 g carbohydrate, 3.9 g fat and 1.5 g fibre per 100 ml, 1000 ml bag			<i>e.g. Nutrison Multi Fibre</i>
<b>ENTERAL FEED 1.2 KCAL/ML – Restricted</b> see terms on the preceding page			
☛ Liquid 5.55 g protein, 15.1 g carbohydrate, 3.93 g fat and 2 g fibre per 100 ml, 1,000 ml bag			<i>e.g. Jevity Plus RTH</i>
<b>ORAL FEED – Restricted</b> see terms on the preceding page			
☛ Powder 16 g protein, 59.8 g carbohydrate and 14 g fat per 100 g, can .....	13.00	850 g	Ensure (Chocolate) Ensure (Vanilla)
☛ Powder 18.7 g protein, 54.5 g carbohydrate and 18.9 g fat per 100 g, can .....	9.50	900 g	Fortisip (Vanilla)
☛ Powder 23 g protein, 65 g carbohydrate and 2.5 g fat per 100 g, can .....	10.22	900 g	Sustagen Hospital Formula (Chocolate) Sustagen Hospital Formula (Vanilla)
<b>ORAL FEED 1 KCAL/ML – Restricted</b> see terms on the preceding page			
☛ Liquid 3.8 g protein, 23 g carbohydrate and 12.7 g fibre per 100 ml, 237 ml carton			<i>e.g. Resource Fruit Beverage</i>
<b>ORAL FEED 1.5 KCAL/ML – Restricted</b> see terms on the preceding page			
☛ Liquid 5.5 g protein, 21.1 g carbohydrate and 4.81 g fat per 100 ml, can .....	1.33	237 ml	Ensure Plus (Chocolate) Ensure Plus (Vanilla)
☛ Liquid 6.25 g protein, 20.2 g carbohydrate and 4.92 g fat per 100 ml, carton .....	1.26	200 ml	Ensure Plus (Banana) Ensure Plus (Chocolate) Ensure Plus (Fruit of the Forest) Ensure Plus (Vanilla)
☛ Liquid 4 g protein and 33.5 g carbohydrate per 100 ml, 200 ml bottle			<i>e.g. Fortijuice</i>
☛ Liquid 6 g protein, 18.4 g carbohydrate and 5.8 g fat per 100 ml, 200 ml bottle			<i>e.g. Fortisip</i>
☛ Liquid 6 g protein, 18.4 g carbohydrate, 5.8 g fat and 2.3 g fibre per 100 ml, 200 ml bottle			<i>e.g. Fortisip Multi Fibre</i>

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

## Bacterial and Viral Vaccines

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE – **Restricted** see terms below

☞ Inj 30 IU diphtheria toxoid with 30 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units poliomyelitis virus in 0.5 ml syring

### ☞Restricted

For primary vaccination in children

DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE – **Restricted** see terms below

☞ Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin, 80 D-antigen units poliomyelitis virus, 10 mcg hepatitis B surface antigen in 0.5 ml syringe (1) and inj 10 mcg haemophilus influenzae type B vaccine vial

### ☞Restricted

Either:

- 1 For primary vaccination in children; or
- 2 For revaccination of children following immunosuppression.

## Bacterial Vaccines

BACILLUS CALMETTE-GUERIN VACCINE – **Restricted** see terms below

☞ Inj 1.5 mg vial with diluent

### ☞Restricted

For infants at increased risk of tuberculosis

Note: increased risk is defined as:

- 1 Living in a house or family with a person with current or past history of TB; or
- 2 Having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3 During their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000.

A list of countries with high rates of TB are available at [www.moh.govt.nz/immunisation](http://www.moh.govt.nz/immunisation) or [www.bcgatlas.org/index.php](http://www.bcgatlas.org/index.php).

DIPHTHERIA AND TETANUS VACCINE – **Restricted** see terms below

☞ Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid in 0.5 ml syringe

### ☞Restricted

Any of the following:

- 1 For vaccination of patients aged between 45 and 65 years old; or
- 2 For vaccination of previously unimmunised patients; or
- 3 For revaccination of children following immunosuppression; or
- 4 For revaccination for patients with tetanus-prone wounds; or
- 5 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE – **Restricted** see terms below

☞ Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis toxoid, 8 mcg pertussis filamentous haemagglutinin and 2.5 mcg pertactin in 0.5 ml syringe

### ☞Restricted

Either:

- 1 For primary vaccination in children aged 7-18 years; or
- 2 For pregnant women between gestational weeks 28 and 38 during epidemics.

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
<b>HAEMOPHILUS INFLUENZAE TYPE B VACCINE – Restricted</b> see terms below		
⚡ Inj 10 mcg vial with diluent syringe		
➔ <b>Restricted</b>		
Any of the following:		
1 For primary vaccination in children; or		
2 For revaccination of children following immunosuppression; or		
3 For children aged 0-18 years with functional asplenia; or		
4 For patients pre- and post-splenectomy; or		
5 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.		
<b>MENINGOCOCCAL (A, C, Y AND W-135) CONJUGATE VACCINE – Restricted</b> see terms below		
⚡ Inj 48 mcg in 0.5 ml vial		
➔ <b>Restricted</b>		
Any of the following:		
1 For patients pre- and post-splenectomy; or		
2 For children aged 0-18 years with functional asplenia; or		
3 For organisation and community based outbreaks; or		
4 For use in transplant patients; or		
5 For use following immunosuppression.		
<b>MENINGOCOCCAL (A, C, Y AND W-135) POLYSACCHARIDE VACCINE – Restricted</b> see terms below		
⚡ Inj 200 mcg vial with diluent		
➔ <b>Restricted</b>		
Any of the following:		
1 For patients pre- and post-splenectomy; or		
2 For children aged 2-18 years with functional asplenia; or		
3 For organisation and community based outbreaks.		
<b>MENINGOCOCCAL C CONJUGATE VACCINE – Restricted</b> see terms below		
⚡ Inj 10 mcg in 0.5 ml syringe		
➔ <b>Restricted</b>		
Any of the following:		
1 For patients pre- and post-splenectomy; or		
2 For children aged 0-18 years with functional asplenia; or		
3 For organisation and community based outbreaks; or		
4 For use in transplant patients aged under 2 years; or		
5 For use following immunosuppression in patients aged under 2 years.		
<b>PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE – Restricted</b> see terms below		
⚡ Inj 16 mcg in 0.5 ml syringe		
➔ <b>Restricted</b>		
For primary vaccination in children		
<b>PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE – Restricted</b> see terms below		
⚡ Inj 30.8 mcg in 0.5 ml syringe		
➔ <b>Restricted</b>		
Any of the following:		
1 For high risk children under the age of 5; or		
2 For patients aged less than 18 years pre- or post-splenectomy or with functional asplenia; or		
3 For revaccination of children following immunosuppression; or		
4 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.		

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE – Restricted</b> see terms below			
☞ Inj 575 mcg in 0.5 ml vial			
☞ <b>Restricted</b>			
Any of the following:			
1 For patients pre- and post-splenectomy; or			
2 For children aged 2-18 years with functional asplenia; or			
3 For revaccination of children following immunosuppression; or			
4 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.			
<b>SALMONELLA TYPHI VACCINE – Restricted</b> see terms below			
☞ Inj 25 mcg in 0.5 ml syringe			
☞ <b>Restricted</b>			
For use during typhoid fever outbreaks			
<b>Viral Vaccines</b>			
<b>HEPATITIS A VACCINE – Restricted</b> see terms below			
☞ Inj 720 ELISA units in 0.5 ml syringe			
☞ Inj 1440 ELISA units in 1 ml syringe			
☞ <b>Restricted</b>			
Any of the following:			
1 For use in transplant patients; or			
2 For use in children with chronic liver disease; or			
3 For close contacts of known hepatitis A carriers.			
<b>HEPATITIS B VACCINE – Restricted</b> see terms below			
☞ Inj 5 mcg in 0.5 ml vial			
☞ Inj 10 mcg in 1 ml vial			
☞ <b>Restricted</b>			
Any of the following:			
1 Household or sexual contacts of known hepatitis B carriers; or			
2 Children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or			
3 Dialysis patients; or			
4 HIV-positive patients; or			
5 Hepatitis C positive patients; or			
6 For use in transplant patients; or			
7 For use following immunosuppression; or			
8 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.			
<b>HUMAN PAPILLOMAVIRUS (6, 11, 16 AND 18) VACCINE [HPV] – Restricted</b> see terms below			
☞ Inj 120 mcg in 0.5 ml syringe			
☞ <b>Restricted</b>			
Any of the following:			
1 Women aged between 9 and 19 years old; or			
2 Male patients aged between 9 and 25 years old with confirmed HIV infection; or			
3 For use in transplant patients.			
<b>INFLUENZA VACCINE – Restricted</b> see terms on the next page			
☞ Inj 45 mcg in 0.5 ml syringe .....	90.00	10	Fluarix Influvac



Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

### ➔Restricted

Any of the following:

- 1 All people 65 years of age and over; or
- 2 People under 65 years of age who:
  - 2.1 Have any of the following cardiovascular diseases:
    - 2.1.1 Ischaemic heart disease; or
    - 2.1.2 Congestive heart disease; or
    - 2.1.3 Rheumatic heart disease; or
    - 2.1.4 Congenital heart disease; or
    - 2.1.5 Cerebro-vascular disease; or
  - 2.2 Have any of the following chronic respiratory diseases:
    - 2.2.1 Asthma, if on a regular preventative therapy; or
    - 2.2.2 Other chronic respiratory disease with impaired lung function; or
  - 2.3 Have diabetes;
  - 2.4 Have chronic renal disease;
  - 2.5 Have any cancer, excluding basal and squamous skin cancers if not invasive;
  - 2.6 Have any of the following other conditions:
    - 2.6.1 Autoimmune disease;
    - 2.6.2 Immune suppression;
    - 2.6.3 HIV;
    - 2.6.4 Transplant recipients;
    - 2.6.5 Neuromuscular and CNS diseases;
    - 2.6.6 Haemoglobinopathies;
    - 2.6.7 Are children on long term aspirin; or
  - 2.7 Are pregnant; or
  - 2.8 Are children aged four and under who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- 3 People under 18 years of age living within the boundaries of the Canterbury District Health Board.

Note: The following conditions are excluded from funding:

- asthma not requiring regular preventative therapy; and
- hypertension and/or dyslipidaemia without evidence of end-organ disease.

MEASLES, MUMPS AND RUBELLA VACCINE – **Restricted** see terms below

¶ Inj 1000 TCID50 measles, 12500 TCID50 mumps and 1000 TCID50 rubella vial with diluent

### ➔Restricted

Any of the following:

- 1 For primary vaccination in children; or
- 2 For revaccination following immunosuppression; or
- 3 For any individual susceptible to measles, mumps or rubella.

POLIOMYELITIS VACCINE – **Restricted** see terms below

¶ Inj 80 D-antigen units in 0.5 ml syringe

### ➔Restricted

Either:

- 1 For previously unvaccinated individuals; or
- 2 For revaccination following immunosuppression.

RABIES VACCINE

Inj 2.5 IU vial with diluent

VARICELLA ZOSTER VACCINE [CHICKEN POX VACCINE] – **Restricted** see terms on the next page

¶ Inj 1350 PFU vial with diluent

¶ Inj 2000 PFU vial with diluent

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## ➡Restricted

Any of the following:

- 1 For non-immune patients:
  - 1.1 with chronic liver disease who may in future be candidates for transplantation; or
  - 1.2 with deteriorating renal function before transplantation; or
  - 1.3 prior to solid organ transplant; or
  - 1.4 prior to any elective immunosuppression; or
  - 1.5 for post exposure prophylaxis who are immune competent inpatients.
- 2 For patients at least 2 years after bone marrow transplantation, on advice of their specialist; or
- 3 For patients at least 6 months after completion of chemotherapy, on advice of their specialist; or
- 4 For HIV positive non-immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
- 5 For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has:
  - 5.1 adult household contact - a negative serology result for varicella; or
  - 5.2 child household contact - no clinical history of varicella or negative varicella serology.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Optional Pharmaceuticals</b>			
<b>NOTE:</b>			
In addition to the products expressly listed here in Part III: Optional Pharmaceuticals, a number of additional Optional Pharmaceuticals, including some wound care products and disposable laparoscopic equipment, are listed in an addendum to Part III which is available at <a href="http://www.pharmac.govt.nz">www.pharmac.govt.nz</a> . The Optional Pharmaceuticals listed in the addendum are deemed to be listed in Part III, and the Rules of the Pharmaceutical Schedule applying to products listed in Part III apply to them.			
<b>BLOOD GLUCOSE DIAGNOSTIC TEST METER</b>			
1 meter with 50 lancets, a lancing device, and 10 diagnostic test strips .....	20.00	1	Caresens II
			Caresens N
			Caresens N POP
Meter .....	9.00	1	FreeStyle Lite
			On Call Advanced
	19.00		Accu-Chek Performa
<b>BLOOD GLUCOSE DIAGNOSTIC TEST STRIP</b>			
Blood glucose test strips .....	10.56	50 test	CareSens
			CareSens N
	21.65		FreeStyle Lite
	28.75		Accu-Chek Performa
			Freestyle Optium
Blood glucose test strips × 50 and lancets × 5 .....	19.10	50 test	On Call Advanced
<b>BLOOD KETONE DIAGNOSTIC TEST METER</b>			
Meter .....	40.00	1	Freestyle Optium
<b>INSULIN PEN NEEDLES</b>			
29 g × 12.7 mm .....	10.50	100	B-D Micro-Fine
31 g × 5 mm .....	11.75	100	B-D Micro-Fine
31 g × 6 mm .....	10.50	100	ABM
31 g × 8 mm .....	10.50	100	ABM
			B-D Micro-Fine
32 g × 4 mm .....	10.50	100	B-D Micro-Fine
<b>INSULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE</b>			
Syringe 0.3 ml with 29 g × 12.7 mm needle .....	13.00	100	B-D Ultra Fine
Syringe 0.3 ml with 31 g × 8 mm needle .....	13.00	100	B-D Ultra Fine II
Syringe 0.5 ml with 29 g × 12.7 mm needle .....	13.00	100	B-D Ultra Fine
Syringe 0.5 ml with 31 g × 8 mm needle .....	13.00	100	B-D Ultra Fine II
Syringe 1 ml with 29 g × 12.7 mm needle .....	13.00	100	ABM
			B-D Ultra Fine
Syringe 1 ml with 31 g × 8 mm needle .....	13.00	100	ABM
			B-D Ultra Fine II
<b>KETONE BLOOD BETA-KETONE ELECTRODES</b>			
Test strips .....	15.50	10 strip	Freestyle Optium Ketone
<b>MASK FOR SPACER DEVICE</b>			
Size 2 .....	2.99	1	EZ-fit Paediatric Mask
<b>PEAK FLOW METER</b>			
Low Range .....	11.44	1	Breath-Alert
Normal Range .....	11.44	1	Breath-Alert

## PART III - OPTIONAL PHARMACEUTICALS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PREGNANCY TEST - HCG URINE</b>			
Cassette .....	22.80	40 test	Innovacon hCG One Step Pregnancy Test
<b>SODIUM NITROPRUSSIDE</b>			
Test strip .....	6.00	50 strip	Accu-Chek Ketur-Test
<b>SPACER DEVICE</b>			
230 ml (single patient) .....	4.72	1	Space Chamber Plus
800 ml .....	8.50	1	Volumatic

<b>- Symbols -</b>		Agents Used in the Treatment of	Amyl nitrite .....	44
8-methoxypsoralen .....	51	Poisonings .....	Anabolic Agents .....	57
<b>- A -</b>		Ajmaline .....	Anaesthetics .....	97
A-Scabies .....	48	Alanase .....	Anagrelide hydrochloride .....	125
Abacavir sulphate .....	76	Albendazole .....	Analgesics .....	101
Abacavir sulphate with		Aldara .....	Anastrozole .....	134
lamivudine .....	76	Alendronate sodium .....	Andriol Testocaps .....	57
Abciximab .....	139	Alendronate sodium with	Androderm .....	57
Abilify .....	112	cholecalciferol .....	Androgen Agonists and	
ABM Hydroxocobalamin .....	23	Alfacalcidol .....	Antagonists .....	57
Acarbose .....	15	Alfentanil hydrochloride .....	Anexate .....	173
Accarb .....	15	Alinia .....	Antabuse .....	121
Accu-Chek Ketur-Test .....	204	Alitraq .....	Antacids and Antiflatulents .....	12
Accu-Chek Performa .....	203	Allersoothe .....	Anti-Infective Agents .....	53
Accuretic 10 .....	35	Allopurinol .....	Anti-Infective Preparations	
Accuretic 20 .....	35	Alpha tocopheryl acetate .....	Dermatological .....	47
Acetadote .....	173	Alpha-Adrenoceptor Blockers .....	Sensory .....	167
Acetazolamide .....	170	Alphamox .....	Anti-Inflammatory	
Acetic acid		Alprazolam .....	Preparations .....	168
Extemporaneous .....	181	Alprostadiol hydrochloride .....	Antiacne Preparations .....	48
Genito-Urinary .....	53	Alteplase .....	Antiallergy Preparations .....	161
Acetic acid with hydroxyquinoline,		Alum .....	Antianaemics .....	25
glycerol and ricinoleic acid .....	53	Aluminium hydroxide .....	Antiarrhythmics .....	37
Acetic acid with propylene		Aluminium hydroxide with	Antibacterials .....	63
glycol .....	172	magnesium hydroxide and	Anticholinergic Agents .....	162
Acetylcholine chloride .....	170	simethicone .....	Anticholinesterases .....	86
Acetylcysteine .....	173	Amantadine hydrochloride .....	Antidepressants .....	104
Aciclovir		AmBisome .....	Antidiarrhoeals and Intestinal	
Infection .....	82	Ambrisentan .....	Anti-Inflammatory Agents .....	12
Sensory .....	167	Amethocaine .....	Antiepilepsy Drugs .....	106
Acid Citrate Dextrose A .....	28	Nervous .....	Antifibrinolytics, Haemostatics	
Acidex .....	12	Sensory .....	and Local Sclerosants .....	26
Acipimox .....	43	Amikacin .....	Antifungals .....	69
Acitretin .....	51	Amiloride hydrochloride .....	Antihypotensives .....	37
Aclasta .....	88	Amiloride hydrochloride with	Antimigraine Preparations .....	110
Actemra .....	157	furosemide .....	Antimycobacterials .....	71
Actinomycin D .....	123	Amiloride hydrochloride with	Antinaus .....	112
Adalimumab .....	139	hydrochlorothiazide .....	Antinausea and Vertigo	
Adapalene .....	48	Aminophylline .....	Agents .....	111
Adefin XL .....	39	Amiodarone hydrochloride .....	Antiparasitics .....	72
Adefovir dipivoxil .....	78	Amisulpride .....	Antipruritic Preparations .....	48
Adenosine .....	37	Amitrip .....	Antipsychotic Agents .....	112
Adenuric .....	92	Amitriptyline .....	Antiretrovirals .....	74
Adrenaline .....	44	Amlodipine .....	Antirheumatoid Agents .....	86
Advantan .....	50	Amorolfine .....	Antiseptics and	
Advate .....	27	Amoxycillin .....	Disinfectants .....	174
Aerrane .....	97	Amoxycillin with clavulanic	Antispasmodics and Other	
Agents Affecting the		acid .....	Agents Altering Gut	
Renin-Angiotensin System .....	35	Amphotericin B	Motility .....	14
Agents for Parkinsonism and		Alimentary .....	Antithrombotics .....	28
Related Disorders .....	96	Infection .....	Antithymocyte globulin	
		Amsacrine .....	(equine) .....	159

## Generic Chemicals and Brands

Anti-thymocyte globulin		Arrow - Clopid	30	Atorvastatin	42
(rabbit)	159	Arrow-Amitriptyline	104	Atovaquone with proguanil	
Ant ulcerants	14	Arrow-Bendroflumazide	41	hydrochloride	73
Antivirals	78	Arrow-Brimonidine	171	Atracurium besylate	92
Anxiolytics	117	Arrow-Calcium	20	Atripila	76
Anzatax	131	Arrow-Citalopram	106	Atropine sulphate	
Apidra	16	Arrow-Diazepam	117	Cardiovascular	37
Apidra Solostar	16	Arrow-Doxorubicin	123	Sensory	171
Apo-Allopurinol	91	Arrow-Etidronate	88	Atropt	171
Apo-Amiloride	41	Arrow-Fluoxetine	106	Augmentin	65
Apo-Amlodipine	39	Arrow-Gabapentin	107	Auranofin	86
Apo-Amoxi	65	Arrow-Lamotrigine	109	Ava 20 ED	53
Apo-Azithromycin	64	Arrow-Lisinopril	35	Ava 30 ED	53
Apo-Cilazapril/ Hydrochlorothiazide	35	Arrow-Losartan & Hydrochlorothiazide	36	Avanza	105
Apo-Clarithromycin	65	Arrow-Morphine LA	103	Avelox	66
Apo-Clomipramine	104	Arrow-Nifedipine XR	39	Avelox IV 400	66
Apo-Diclo	94	Arrow-Norfloxacin	67	Azactam	67
Apo-Diltiazem CD	40	Arrow-Ornidazole	73	Azamun	159
Apo-Doxazosin	36	Arrow-Quinapril 10	35	Azathioprine	159
Apo-Gliclazide	16	Arrow-Quinapril 20	35	Azithromycin	64
Apo-Megestrol	132	Arrow-Quinapril 5	35	Azol	59
Apo-Moclobemide	105	Arrow-Ranitidine	14	AZT	77
Apo-Nadolol	38	Arrow-Roxithromycin	65	Aztreonam	67
Apo-Oxybutynin	56	Arrow-Sertraline	106	- B -	
Apo-Perindopril	35	Arrow-Simva	42	B-D Micro-Fine	203
Apo-Pindolol	38	Arrow-Sumatriptan	111	B-D Ultra Fine	203
Apo-Prazo	36	Arrow-Tolterodine	56	B-D Ultra Fine II	203
Apo-Prazosin	36	Arrow-Topiramate	110	Bacillus calmette-guerin (BCG)	159
Apo-Prednisone	58	Arrow-Tramadol	104	Bacillus calmette-guerin vaccine	198
Apo-Prednisone S29	58	Arrow-Venlafaxine XR	106	Baclofen	92
Apo-Propranolol	39	Arsenic trioxide	125	Bacterial and Viral Vaccines	198
Apo-Pyridoxine	23	Artemether with lumefantrine	73	Bacterial Vaccines	198
Apo-Risperidone	114	Artesanate	73	Baraclude	79
Apo-Ropinirole	97	Articaine hydrochloride with adrenaline	98	Barium sulphate	177
Apo-Zopiclone	118	Asacol	13	Barrier Creams and Emollients	48
Apomine	96	Asamax	13	Basiliximab	145
Apomorphine hydrochloride	96	Ascorbic acid		Beclazone 100	163
Apraclonidine	171	Alimentary	23	Beclazone 250	163
Aprepitant	111	Extemporaneous	181	Beclazone 50	163
Apresoline	45	Aspen Adrenaline	44	Beclomethasone	
Aprotinin	26	Aspen Ciprofloxacin	66	dipropionate	161, 163
Aqueous cream	49	Aspirin		Bee venom	161
Arachis oil [Peanut oil]	181	Blood	30	Bendroflumazide	41
Arava	86	Nervous	101	Bendroflumethazide	
Aremed	134	Asthalin	163	[Bendrofluazide]	41
Arginine		Atazanavir sulphate	77	BeneFIX	27
Alimentary	19	Atenolol	38	Benzathine benzylpenicillin	66
Various	178	Atenolol-AFT	38	Benzbromaron AL 100	91
Argipressin [Vasopressin]	61	ATGAM	159	Benzbromarone	91
Aripiprazole	112	Ativan	117	Benzocone	98
Aristocort	50	Atomoxetine	118		
Aromasin	134				

Benzoin .....	181	Blood glucose diagnostic test		Calcipotriol .....	51
Benzoyl peroxide .....	48	meter .....	203	Calcitonin .....	57
Benztrop .....	96	Blood glucose diagnostic test		Calcitriol .....	24
Benztropine mesylate .....	96	strip .....	203	Calcitriol-AFT .....	24
Benzylamine hydrochloride .....	21	Blood ketone diagnostic test		Calcium carbonate .....	12, 20
Benzylamine hydrochloride with		meter .....	203	Calcium Channel Blockers .....	39
cetylpyridinium chloride .....	21	BNM .....	55, 103	Calcium chloride .....	31
Benzylpenicillin sodium [Penicillin		Boceprevir .....	81	Calcium chloride with	
G] .....	66	Bonney's blue dye .....	178	magnesium chloride,	
Beractant .....	166	Boric acid .....	181	potassium chloride, sodium	
Beta Scalp .....	51	Bortezomib .....	125	acetate, sodium chloride and	
Beta-Adrenoceptor Agonists .....	163	Bosentan .....	45	sodium citrate .....	169
Beta-Adrenoceptor Blockers .....	38	Bosvate .....	38	Calcium folinate .....	132
Betadine .....	175	Botox .....	92	Calcium Folate Ebewe .....	132
Betadine Skin Prep .....	175	Botulism antitoxin .....	173	Calcium gluconate	
Betagan .....	170	Breath-Alert .....	203	Blood .....	31
Betahistine dihydrochloride .....	111	Bridion .....	93	Dermatological .....	52
Betaine .....	19	Brilinta .....	30	Calcium Homeostasis .....	57
Betamethasone .....	57	Brimonidine tartrate .....	171	Calcium polystyrene	
Betamethasone dipropionate .....	50	Brimonidine tartrate with		sulphonate .....	34
Betamethasone dipropionate		timolol .....	171	Calcium Resonium .....	34
with calcipotriol .....	51	Brinzolamide .....	170	Calsource .....	20
Betamethasone sodium		Bromocriptine .....	96	Candidas .....	71
phosphate with		Brufen SR .....	94	Candesartan cilexetil .....	36
betamethasone acetate .....	57	Budesonide		Candestar .....	36
Betamethasone		Alimentary .....	12	Capecitabine .....	124
valerate .....	50-51	Respiratory .....	161, 163	Capoten .....	35
Betamethasone valerate with		Budesonide with		Capsaicin	
clioquinol .....	50	eformoterol .....	164	Musculoskeletal System .....	95
Betamethasone valerate with		Bumetanide .....	40	Nervous .....	101
fusidic acid .....	51	Bupafen .....	99	Captopril .....	35
Betaxolol .....	170	Bupivacaine hydrochloride .....	98	Carbaccord .....	127
Bevacizumab .....	145	Bupivacaine hydrochloride with		Carbamazepine .....	107
Bezafibrate .....	42	adrenaline .....	99	Carbasorb-X .....	174
Bezalip .....	42	Bupivacaine hydrochloride with		Carbimazole .....	61
Bezalip Retard .....	42	fentanyl .....	99	Carbomer .....	171
Bicalaccord .....	132	Bupivacaine hydrochloride with		Carboplatin .....	127
Bicalutamide .....	132	glucose .....	99	Carboplatin Ebewe .....	127
Bicillin LA .....	66	Buprenorphine with		Carboprost trometamol .....	54
Bile and Liver Therapy .....	15	naloxone .....	120	Carboxymethylcellulose	
Bimatoprost .....	171	Bupropion hydrochloride .....	121	Alimentary .....	21
Biodone .....	102	Burinex .....	40	Extemporaneous .....	181
Biodone Extra Forte .....	102	Buscopan .....	14	Cardinol LA .....	39
Biodone Forte .....	102	Buserelin .....	60	Cardizem CD .....	40
Biotin .....	19	Buspirone hydrochloride .....	117	CareSens .....	203
Bisacodyl .....	19	Busulfan .....	123	Caresens II .....	203
Bismuth subgallate .....	181	Butacort Aqueous .....	161	CareSens N .....	203
Bismuth subnitrate and iodoform		- C -		Caresens N .....	203
paraffin .....	179	Cabergoline .....	59	Caresens N POP .....	203
Bismuth trioxide .....	15	Caffeine .....	119	Carmellose sodium .....	172
Bisoprolol .....	38	Caffeine citrate .....	165	Carmustine .....	123
Bivalirudin .....	28	Cal-d-Forte .....	24	Carvedilol .....	38
Bleomycin sulphate .....	123	Calamine .....	48	Casprofungin .....	71

Catapres .....	40	hydrochloride .....	112	Clopixol .....	115, 117
Catapres-TTS-1 .....	40	Chlorsig .....	167	Clostridium botulinum type A	
Catapres-TTS-2 .....	40	Chlorthalidone .....		toxin .....	92
Catapres-TTS-3 .....	40	[Chlorthalidone] .....	41	Clotrimazole	
Ceenu .....	123	Chlorthalidone .....	41	Dermatological .....	47
Cefaclor .....	64	Cholecalciferol .....	24	Genito-Urinary .....	53
Cefalexin .....	63	Cholestyramine .....	42	Clove oil .....	181
Cefalexin Sandoz .....	63	Choline salicylate with		Clozapine .....	113
Cefazolin .....	64	cetalkonium chloride .....	22	Clozaril .....	113
Cefepime .....	64	Cholvastin .....	42	Co-trimoxazole .....	69
Cefotaxime .....	64	Choriogonadotropin alfa .....	60	Coal tar .....	181
Cefotaxime Sandoz .....	64	Ciclopirox olamine .....	47	Coal tar with salicylic acid and	
Cefoxitin .....	64	Ciclosporin .....	134	sulphur .....	51
Ceftazadime .....	64	Cidofovir .....	82	Coal tar with triethanolamine laryl	
Ceftriaxone .....	64	Cilazapril .....	35	sulphate and fluorescein .....	51
Ceftriaxone-AFT .....	64	Cilazapril with		Cocaine hydrochloride .....	99
Cefuroxime .....	64	hydrochlorothiazide .....	35	Cocaine hydrochloride with	
Celecoxib .....	93	Cilicaine .....	66	adrenaline .....	99
Celiprolol .....	38	Cilicaine VK .....	66	Codeine phosphate	
CellCept .....	159	Cimetidine .....	14	Extemporaneous .....	181
Celol .....	38	Cinchocaine hydrochloride with		Nervous .....	101
Centrally-Acting Agents .....	40	hydrocortisone .....	13	Cogentin .....	96
Cephalexin ABM .....	63	Cipflox .....	66	Colaspase [L-asparaginase] .....	126
Cetirizine - AFT .....	161	Ciprofloxacin		Colchicine .....	92
Cetirizine hydrochloride .....	161	Infection .....	66	Colestimethate .....	68
Cetomacrogol .....	49	Sensory .....	167	Colestipol hydrochloride .....	42
Cetomacrogol with glycerol .....	49	Cisplatin .....	128	Colgot .....	92
Cetrimide .....	181	Cisplatin Ebewe .....	128	Colifoam .....	13
Champix .....	121	Citalopram hydrobromide .....	106	Colistin sulphomethate	
Charcoal .....	174	Citanest .....	100	[Colestimethate] .....	68
Chemotherapeutic Agents .....	123	Citric acid .....	181	Colistin-Link .....	68
Chemo-vax vaccine .....	201	Citric acid with magnesium oxide		Collodion flexible .....	181
Chlorafast .....	167	and sodium picosulfate .....	18	Colofac .....	14
Chloral hydrate .....	118	Citric acid with sodium		Colony-Stimulating Factors .....	31
Chlorambucil .....	123	bicarbonate .....	177	Coloxyl .....	18
Chloramphenicol		Cladribine .....	124	Compound electrolytes .....	31, 34
Infection .....	68	Clarithromycin .....	65	Compound electrolytes with	
Sensory .....	167	Clexane .....	29	glucose .....	31, 34
Chlorhexidine		Clindamycin .....	68	Compound	
Genito-Urinary .....	53	Clindamycin ABM .....	68	hydroxybenzoate .....	181
Various .....	174, 179	Clobazam .....	107	Compound sodium lactate	
Chlorhexidine gluconate		Clobetasol propionate .....	50–51	[Hartmann's solution] .....	32
Alimentary .....	21	Clobetasone butyrate .....	50	Compound sodium lactate with	
Extemporaneous .....	181	Clofazimine .....	71	glucose .....	32
Genito-Urinary .....	53	Clomazol .....	47, 53	Concerta .....	119
Chlorhexidine with		Clomiphene citrate .....	59	Condyline .....	52
cetrimide .....	174, 179	Clomipramine hydrochloride .....	104	Contraceptives .....	53
Chlorhexidine with ethanol .....	174	Clonazepam .....	106–107, 117	Contrast Media .....	175
Chloroform .....	181	Clonidine .....	40	Corangin .....	43
Chloroquine phosphate .....	73	Clonidine BNM .....	40	Cordarone-X .....	37
Chlorothiazide .....	41	Clonidine hydrochloride .....	40	Corticosteroids	
Chlorpheniramine maleate .....	161	Clopidogrel .....	30	Dermatological .....	50
Chlorpromazine		Clopine .....	113	Hormone .....	57



Corticotrorelin (ovine) .....	60	DBL Leucovorin Calcium .....	132	Diamox .....	170
Cosopt .....	170	DBL Morphine Sulphate .....	103	Diatrizoate meglumine with diatrizoate sodium .....	175
Cough Suppressants .....	163	DBL Pethidine Hydrochloride .....	104	Diatrizoate sodium .....	175
Crotamiton .....	48	DBL Rocuronium Bromide .....	93	Diazepam .....	106, 117
Crystaderm .....	47	DBL Tobramycin .....	63	Diazoxide Alimentary .....	15
CT Plus+ .....	177	DDI .....	76	Cardiovascular .....	44
Curam Duo .....	65	De-Nol .....	15	Dichlorobenzyl alcohol with amylmetacresol .....	22
Curosulf .....	166	De-Worm .....	73	Diclac SR .....	94
Cvite .....	23	Decongestants .....	163	Diclofenac sodium Musculoskeletal System .....	94
Cyclizine hydrochloride .....	111	Decongestants and Antiallergics .....	168	Sensory .....	168
Cyclizine lactate .....	111	Decozol .....	22	Dicobalt edetate .....	174
Cyclopentolate hydrochloride .....	171	Deferiprone .....	174	Didanosine [DDI] .....	76
Cyclophosphamide .....	123	Defibrotide .....	28	Diflucan .....	70
Cycloserine .....	71	Demeclocycline hydrochloride .....	67	Diflucortolone valerate .....	50
Cyklokapron .....	27	Deoxycoformycin .....	127	Digestives Including Enzymes .....	17
Cymevene .....	82	Depo-Medrol .....	58	Digoxin .....	37
Cyproheptadine hydrochloride .....	161	Depo-Medrol with Lidocaine .....	58	Digoxin immune Fab .....	173
Cyproterone acetate .....	57	Depo-Provera .....	54	Dihydrocodeine tartrate .....	101
Cyproterone acetate with ethinyloestradiol .....	53	Depo-Testosterone .....	57	Dihydroergotamine mesylate .....	110
Cysteamine hydrochloride .....	181	Deprim .....	69	Dilatrend .....	38
Cytarabine .....	124	Dermol .....	50, 51	Diltiazem hydrochloride .....	40
<b>- D -</b>		Desferrioxamine mesilate .....	174	Dilzem .....	40
D-Penamine .....	86	Desflurane .....	97	Dimercaprol .....	174
Dabigatran .....	28	Desmopressin acetate .....	61	Dimercaptosuccinic acid .....	174
Dacarbazine .....	126	Desmopressin-PH&T .....	61	Dimethicone .....	48
Dactinomycin [Actinomycin D] .....	123	Dexamethasone Hormone .....	57	Dimethyl sulfoxide .....	179
Daivobet .....	51	Sensory .....	168	Dinoprostone .....	55
Daivonex .....	51	Dexamethasone phosphate .....	57	Diphenamil metilsulfate .....	52
Dalacin C .....	68	Dexamethasone with framycetin and gramicidin .....	167	Diphenoxylate hydrochloride with atropine sulphate .....	12
Dalteparin .....	28	Dexamethasone with neomycin sulphate and polymyxin B sulphate .....	167	Diphtheria and tetanus vaccine .....	198
Danaparoid .....	28	Dexamethasone with tobramycin .....	167	Diphtheria antitoxin .....	173
Danazol .....	59	Dexamethasone-hameln .....	57	Diphtheria, tetanus and pertussis vaccine .....	198
Danthron with poloxamer .....	19	Dexamphetamine sulphate .....	119	Diphtheria, tetanus, pertussis and polio vaccine .....	198
Dantrium .....	93	Dexmedetomidine hydrochloride .....	97	Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine .....	198
Dantrolene .....	93	Dextrose with sodium citrate and citric acid [Acid Citrate Dextrose A] .....	28	Diprivan .....	98
Dapa-Tabs .....	41	DHC Continus .....	101	Dipyridamole .....	30
Dapsone .....	71	Diabetes .....	15	Disodium edetate .....	174
Daptomycin .....	68	Diacomit .....	109	Disodium hydrogen phosphate with sodium dihydrogen phosphate .....	181
Darunavir .....	77	Diagnostic Agents .....	178		
Dasatinib .....	128	Diagnostic and Surgical Preparations .....	169		
Daunorubicin .....	123	Diamide Relief .....	12		
DBL Aminophylline .....	165				
DBL Cefepime .....	64				
DBL Cefotaxime .....	64				
DBL Ceftazidime .....	64				
DBL Epirubicin Hydrochloride .....	124				
DBL Ergometrine .....	55				
DBL Gemcitabine .....	125				



Ferrograd .....	21	Flupenthixol decanoate .....	115	Gelatine, succinylated .....	34
Ferrous fumarate .....	20	Fluphenazine decanoate .....	115	Gelofusine .....	34
Ferrous fumarate with folic acid .....	21	Flutamide .....	132	Gemcitabine .....	125
Ferrous gluconate with ascorbic acid .....	21	Flutamin .....	132	Gemcitabine Actavis 1000 .....	125
Ferrous sulphate .....	21	Fluticasone .....	164	Gemcitabine Actavis 200 .....	125
Ferrous sulphate with ascorbic acid .....	21	Fluticasone propionate .....	161	Gemcitabine Ebewe .....	125
Ferrous sulphate with folic acid .....	21	Fluticasone with salmeterol .....	165	Gemfibrozil .....	42
Ferrum H .....	21	Foban .....	47	Genoptic .....	167
Fexofenadine hydrochloride .....	162	Folic acid .....	26	Genox .....	133
Filgrastim .....	31	Fondaparinux sodium .....	29	Gentamicin sulphate Infection .....	63
Finasteride .....	55	Food Modules .....	184	Sensory .....	167
Flagyl .....	73	Food/Fluid Thickeners .....	185	Gestrinone .....	59
Flagyl-S .....	73	Forteo .....	91	Glatiramer acetate .....	117
Flamazine .....	47	Fortisip (Vanilla) .....	197	Glaucoma Preparations .....	170
Flecainide acetate .....	37	Fortum .....	64	Glibenclamide .....	16
Fleet Phosphate Enema .....	19	Fosamax .....	86	Gliclazide .....	16
Flixonase Hayfever & Allergy .....	161	Fosamax Plus .....	87	Glipizide .....	17
Flixotide .....	164	Foscarnet sodium .....	82	Glivec .....	129
Flixotide Accuhaler .....	164	Fosfomycin .....	68	Glucagen Hypokit .....	15
Florinef .....	58	Fragmin .....	28	Glucagon hydrochloride .....	15
Fluanxol .....	115	Framycetin sulphate .....	167	Glucerna Select (Vanilla) .....	189
Fluarix .....	200	Freeflex .....	33	Glucerna Select RTH (Vanilla) .....	189
Flucloxacillin .....	66	FreeStyle Lite .....	203	Glucose .....	
Flucloxin .....	66	Freestyle Optium .....	203	Alimentary .....	15
Flucon .....	168	Freestyle Optium Ketone .....	203	Blood .....	32
Fluconazole .....	70	Fresofol 1% .....	98	Extemporaneous .....	182
Fluconazole-Claris .....	70	Frusemide-Claris .....	41	Glucose with potassium chloride .....	32
Flucytosine .....	71	Fucidin .....	68	Glucose with potassium chloride and sodium chloride .....	32
Fludara Oral .....	124	Fucithalmic .....	167	Glucose with sodium chloride .....	32
Fludarabine Ebewe .....	124	Fungilin .....	22	Glucose with sucrose and fructose .....	15
Fludarabine phosphate .....	124	Furosemide (frusemide) .....	41	Glycerin with sodium saccharin .....	182
Fludrocortisone acetate .....	58	Fusidate sodium [Fusidic acid] .....	47	Glycerin with sucrose .....	182
Fluids and Electrolytes .....	31	Fusidic acid .....	47	Glycerol .....	
Flumazenil .....	173	Dermatological .....	47	Alimentary .....	18
Flumetasone pivalate with clioquinol .....	167	Infection .....	68	Extemporaneous .....	182
Fluocortolone caproate with fluocortolone pivalate and cinchocaine .....	13	Sensory .....	167	Glycerol with paraffin .....	49
Fluorescein sodium .....	169	Fuzeon .....	74	Glyceryl trinitrate Alimentary .....	14
Fluorescein sodium with lignocaine hydrochloride .....	169	- G -		Cardiovascular .....	43
Fluorescite .....	169	Gabapentin .....	107	Glycine .....	179
Fluorometholone .....	168	Gadobenic acid .....	177	Glycopyrronium bromide .....	14
Fluorouracil .....	124	Gadobutrol .....	177	Glypressin .....	62
Fluorouracil Ebewe .....	124	Gadodiamide .....	177	Glytrin .....	43
Fluorouracil sodium .....	52	Gadoteric acid .....	177	Gonadorelin .....	60
Fluoxetine hydrochloride .....	106	Gadovist .....	177	Goserelin .....	60
		Gadoxetate disodium .....	178		
		Gamma benzene hexachloride .....	47		
		Ganciclovir .....	82		
		Gastrografin .....	175		
		Gastrosoothe .....	14		
		Gefitinib .....	128		
		Gelafusal .....	34		

## Generic Chemicals and Brands

Habitrol (Classic) .....	121	Hydrocortisone with paraffin and wool fat .....	50	Innovacon hCG One Step Pregnancy Test .....	204
Habitrol (Fruit) .....	121	Hydrogen peroxide .....	47	Insulin aspart .....	16
Habitrol (Mint) .....	121	Hydroxocobalamin .....	173	Insulin aspart with insulin aspart protamine .....	15
Haem arginate .....	19	Hydroxocobalamin acetate .....	23	Insulin glargine .....	16
Haemophilus influenzae type B vaccine .....	199	Hydroxychloroquine .....	86	Insulin glulisine .....	16
Haldol .....	115	Hydroxyethyl starch 130/0.4 with magnesium chloride, potassium chloride, sodium acetate and sodium chloride .....	34	Insulin isophane .....	16
Haldol Concentrate .....	115	Hydroxyethyl starch 130/0.4 with sodium chloride .....	34	Insulin lispro .....	16
Haloperidol .....	113	Hydroxyurea .....	126	Insulin lispro with insulin lispro protamine .....	16
Haloperidol decanoate .....	115	Hygroton .....	41	Insulin neutral .....	16
Hartmann's solution .....	31	Hylo-Fresh .....	172	Insulin neutral with insulin isophane .....	16
Healon GV .....	170	Hyoscine butylbromide .....	14	Insulin pen needles .....	203
healthE Dimethicone 5% .....	48	Hyoscine hydrobromide .....	111	Insulin syringes, disposable with attached needle .....	203
healthE Fatty Cream .....	49	Hyperuricaemia and Antigout .....	91	Integrilin .....	30
Heparin sodium .....	29	Hypnovel .....	118	Intelence .....	75
Heparinised saline .....	29	Hypromellose .....	169, 172	Interferon alfa-2a .....	83
Heparon Junior .....	190	Hypromellose with dextran .....	172	Interferon alfa-2b .....	83
Hepatitis A vaccine .....	200	Hysite .....	171	Interferon beta-1-alpha .....	117
Hepatitis B vaccine .....	200	- I -		Interferon beta-1-beta .....	117
Hepsera .....	78	Ibiamox .....	65	Interferon gamma .....	83
Herceptin .....	157	Ibuprofen .....	94	Intra-uterine device .....	53
Hexamine hippurate .....	68	Idarubicin hydrochloride .....	124	Invanz .....	63
Histamine acid phosphate .....	178	Ifosfamide .....	123	Invega Sustenna .....	116
Holoxan .....	123	Ikorel .....	45	Iodine .....	61
Hormone Replacement Therapy .....	58	Ilomedin .....	46	Iodine with ethanol .....	174
HPV .....	200	Iloprost .....	46	Iodised oil .....	175
Humalog Mix 25 .....	16	Imatinib mesilate .....	129	Iodixanol .....	175
Humalog Mix 50 .....	16	Imatinib-AFT .....	129	Iohexol .....	176
Human papillomavirus (6, 11, 16 and 18) vaccine [HPV] .....	200	Imiglucerase .....	20	Iomeprol .....	176
Humatin .....	63	Impipenem with cilastatin .....	63	Iopromide .....	176
Humira .....	139	Imipramine hydrochloride .....	105	Ioscan .....	175
HumiraPen .....	139	Imiquimod .....	52	Iotrolan .....	176
Hyaluronidase .....	91	Immune Modulators .....	83	Ipratropium bromide .....	161–162
Hybloc .....	38	Immunosuppressants .....	134	Iressa .....	128
Hydralazine hydrochloride .....	44–45	Impact Advanced Recovery (Chocolate) .....	196	Irinotecan Actavis 100 .....	126
Hydrea .....	126	Impact Advanced Recovery (Vanilla) .....	196	Irinotecan Actavis 40 .....	126
Hydrocortisone Dermatological .....	50	Imuran .....	159	Irinotecan hydrochloride .....	126
Hydrocortisone Extentoparaneous .....	182	Indapamide .....	41	Iron polymaltose .....	21
Hydrocortisone Hormone .....	58	Indigo carmine .....	178	Iron sucrose .....	21
Hydrocortisone acetate Alimentary .....	13	Indinavir .....	78	Irrigation Solutions .....	179
Hydrocortisone acetate Dermatological .....	50	Indocyanine green .....	178	Isentress .....	78
Hydrocortisone butyrate .....	50, 52	Indomethacin .....	94	Ismo 40 Retard .....	43
Hydrocortisone with ciprofloxacin .....	168	Infliximab .....	145	Ismo-20 .....	43
Hydrocortisone with miconazole .....	51	Influenza vaccine .....	200	Isoflurane .....	97
Hydrocortisone with natamycin and neomycin .....	51	Influvac .....	200	Isoniazid .....	72
		Inhaled Corticosteroids .....	163	Isoniazid with rifampicin .....	72
				Isoprenaline .....	44
				Isopropyl alcohol .....	175
				Isoptin .....	40

Isosorbide mononitrate .....	43	Lax-Tabs .....	19	Liquifilm Forte .....	172
Isotretinoin .....	48	Laxatives .....	18	Liquifilm Tears .....	172
Ispaghula (psyllium) husk .....	18	Laxofast 120 .....	18	Lisinopril .....	35
Isradipine .....	39	Laxofast 50 .....	18	Lissamine green .....	169
Itch-Soothe .....	48	Laxsol .....	18	Lisuride hydrogen maleate .....	97
Itraconazole .....	70	Leflunomide .....	86	Lithicarb FC .....	113
Itrazole .....	70	Letraccord .....	134	Lithium carbonate .....	113
Ivermectin .....	73	Letrozole .....	134	Local Preparations for Anal and Rectal Disorders .....	13
<b>- J -</b>		Leukotriene Receptor Antagonists .....	164	Locoid .....	50, 52
Jadelle .....	54	Leunase .....	126	Locoid Crelo .....	50
Jevity .....	197	Leuporelin acetate .....	60	Locoid Lipocream .....	50
Jevity HiCal RTH .....	196	Leustatin .....	124	Lodoxamide .....	168
Jevity RTH .....	197	Levetiracetam .....	109	Logem .....	109
<b>- K -</b>		Levetiracetam-Rex .....	109	Lomustine .....	123
Kaletra .....	78	Levobunolol hydrochloride .....	170	Long-Acting Beta-Adrenoceptor Agonists .....	164
Kenacomb .....	168	Levocabastine .....	168	Loniten .....	45
Kenacort-A .....	58	Levocarnitine .....	20	Loperamide hydrochloride .....	12
Kenacort-A40 .....	58	Levodopa with benserazide .....	97	Lopinavir with ritonavir .....	78
Ketamine hydrochloride .....	98	Levodopa with carbidopa .....	97	Lopresor .....	38
Ketocal 3:1 (Unflavoured) .....	194	Levomepromazine .....	113	Lorafix .....	162
Ketocal 4:1 (Unflavoured) .....	194	Levonorgestrel .....	54	Lorapaed .....	162
Ketocal 4:1 (Vanilla) .....	194	Levophed .....	44	Loratadine .....	162
Ketoconazole .....		Levosimendan .....	43	Lorazepam .....	107, 117
Dermatological .....	47	Levothyroxine .....	61	Lormetazepam .....	118
Infection .....	69	Lidocaine [Lignocaine] hydrochloride .....	99	Losartan potassium .....	36
Ketone blood beta-ketone electrodes .....	203	Lidocaine [Lignocaine] hydrochloride with adrenaline .....	100	Losartan potassium with hydrochlorothiazide .....	36
Ketoprofen .....	94	Lidocaine [Lignocaine] hydrochloride with adrenaline and tetracaine .....	100	Lostaar .....	36
Ketorolac trometamol .....	168	Lidocaine [Lignocaine] hydrochloride .....	100	Lovir .....	82
Kivexa .....	76	Lidocaine [Lignocaine] with chlorhexidine .....	100	Loxalate .....	106
Klacid .....	65	Lidocaine [Lignocaine] with phenylephrine hydrochloride .....	100	Loxamine .....	106
Klean Prep .....	18	Lidocaine [Lignocaine] with prilocaine .....	100	Lucrin Depot PDS .....	60
Kogenate FS .....	27	Lidocaine-Claris .....	99	Lycinate .....	43
Konakion MM .....	28	Lignocaine .....	99, 100	Lyderm .....	48
Konsyl-D .....	18	Lindane [Gamma benzene hexachloride] .....	47	<b>- M -</b>	
<b>- L -</b>		Linezolid .....	68	m-Amoxiclav .....	65
L-asparaginase .....	126	Lioresal Intrathecal .....	92	m-Cefuroxime .....	64
L-ornithine L-aspartate .....	15	Liothyronine sodium .....	61	m-Eslon .....	103
Labetalol .....	38	Lipazil .....	42	m-Mometasone .....	50
Lacosamide .....	108	Lipid-Modifying Agents .....	42	Mabthera .....	151
Lactose .....	182			Macrogl 3350 with ascorbic acid, potassium chloride and sodium chloride .....	18
Lactulose .....	18			Macrogl 3350 with potassium chloride, sodium bicarbonate and sodium chloride .....	19
Laevolac .....	18			Macrogl 3350 with potassium chloride, sodium bicarbonate, sodium chloride and sodium sulphate .....	18
Lamictal .....	109				
Lamivudine .....	76, 79				
Lamotrigine .....	109				
Lansoprazole .....	14				
Lantus .....	16				
Lantus SoloStar .....	16				
Lapatinib .....	129				
Latanoprost .....	171				
Lax-Sachets .....	19				

Macroglol 400 and propylene glycol .....	172	vaccine .....	199	Metoprolol - AFT CR .....	38
Madopar 125 .....	97	Menthol .....	182	Metoprolol succinate .....	38
Madopar 250 .....	97	Mepivacaine hydrochloride .....	100	Metoprolol tartrate .....	38
Madopar 62.5 .....	97	Mercaptopurine .....	125	Metronidazole	
Madopar HBS .....	97	Meropenem .....	63	Dermatological .....	47
Madopar Rapid .....	97	Mesalazine .....	13	Infection .....	73
Mafenide acetate .....	47	Mesna .....	132	Metyrapone .....	59
Magnesium hydroxide		Mestinon .....	86	Mexiletine hydrochloride .....	37
Alimentary .....	21	Metabolic Disorder Agents .....	19	Mexiletine Hydrochloride	
Extemporaneous .....	182	Metabolic Products .....	186	USP .....	37
Magnesium oxide .....	21	Metamide .....	111	Miacalcic .....	57
Magnesium sulphate .....	21	Metaraminol .....	44	Mianserin hydrochloride .....	105
Magnevist .....	178	Metformin .....	17	Micolette .....	19
Malathion [Maldison] .....	48	Methacholine chloride .....	178	Miconazole .....	22
Malathion with permethrin and piperonyl butoxide .....	48	Methadone hydrochloride		Miconazole nitrate	
Maldison .....	47	Extemporaneous .....	182	Dermatological .....	47
Mannitol .....	41	Nervous .....	102	Genito-Urinary .....	53
Maprotiline hydrochloride .....	105	Methatabs .....	102	Micreme H .....	51
Marcaïn .....	98	Methohexital sodium .....	98	Microgynon 50 ED .....	53
Marcaïn Heavy .....	99	Methopt .....	172	Midazolam .....	118
Marcaïn Isobaric .....	98	Methotrexate .....	125	Midodrine .....	37
Marcaïn with Adrenaline .....	99	Methotrexate Ebewe .....	125	Mifepristone .....	54
Marevan .....	30	Methotrexate Sandoz .....	125	Milrinone .....	45
Marine Blue Lotion SPF 50+ .....	52	Methoxsalen		Minerals .....	20
Martindale Acetylcysteine .....	173	[8-methoxypsoralen] .....	51	Minidiab .....	17
Mask for spacer device .....	203	Methoxyflurane .....	101	Minirin .....	61
Mast Cell Stabilisers .....	165	Methyl aminolevulinate hydrochloride .....	52	Minocycline .....	67
Maxidex .....	168	Methyl hydroxybenzoate .....	182	Minoxidil .....	45
Measles, mumps and rubella vaccine .....	201	Methylcellulose .....	182	Mirtazapine .....	105
Mebendazole .....	73	Methylcellulose with glycerin and sodium saccharin .....	182	Misoprostol .....	14
Mebeverine hydrochloride .....	14	Methylcellulose with glycerin and sucrose .....	182	Mitomycin C .....	124
Medrol .....	58	Methyldopa .....	40	Mitozantrone .....	124
Medroxyprogesterone .....	60	Methylene blue .....	178	Mitozantrone Ebewe .....	124
Medroxyprogesterone acetate		Methylphenidate		Mivacron .....	93
Genito-Urinary .....	54	hydrochloride .....	119	Mivacurium chloride .....	93
Hormone .....	59	Methylprednisolone (as sodium succinate) .....	58	Moclobemide .....	105
Mefenamic acid .....	94	Methylprednisolone		Modafinil .....	120
Mefloquine hydrochloride .....	73	aceponate .....	50	Moderate .....	115
Megestrol acetate .....	132	Methylprednisolone acetate .....	58	Mogine .....	109
Meglumine gadopentetate .....	178	Methylprednisolone acetate with lignocaine .....	58	Mometasone furoate .....	50
Melatonin .....	118	Methylthioninium chloride [Methylene blue] .....	178	Monosodium glutamate with sodium aspartate .....	180
Meloxicam .....	94	Methylxanthines .....	165	Monosodium l-aspartate .....	180
Melphalan .....	123	Metoclopramide		Montelukast .....	164
Meningococcal (A, C, Y and W-135) conjugate vaccine .....	199	hydrochloride .....	111	Morococog alfa [Recombinant factor VIII] .....	27
Meningococcal (A, C, Y and W-135) polysaccharide vaccine .....	199	Metoclopramide hydrochloride		Morphine hydrochloride .....	102
Meningococcal C conjugate		with paracetamol .....	111	Morphine sulphate .....	103
		Metolazone .....	41	Morphine tartrate .....	103
				Motetis .....	96
				Mouth and Throat .....	21
				Moxifloxacin .....	66
				Mucolytics and	

Expectorants .....	165	Nicotinic acid .....	43	Oestradiol valerate .....	58
Multihance .....	177	Nifedipine .....	39	Oestradiol with norethisterone	
Multiple Sclerosis		Nilstat .....	22, 69	acetate .....	59
Treatments .....	117	Nimodipine .....	39	Oestriol	
Multivitamins .....	22-23	Nitazoxanide .....	73	Genito-Urinary .....	55
Mupirocin .....	47	Nitrates .....	43	Hormone .....	60
Muscle Relaxants and Related		Nitrazepam .....	118	Oestrogens .....	55
Agents .....	92	Nitroderm TTS 10 .....	43	Oestrogens (conjugated	
Myambutol .....	72	Nitroderm TTS 5 .....	43	equine) .....	59
Mycobutin .....	72	Nitrofurantoin .....	68	Oestrogens with	
Mycophenolate mofetil .....	159	Nitronal .....	43	medroxyprogesterone	
Mydiacryl .....	171	Noflam 250 .....	95	acetate .....	59
Mydriatics and Cycloplegics .....	171	Noflam 500 .....	95	Oil in water emulsion .....	49
Mylan Atenolol .....	38	Non-Steroidal Anti-Inflammatory		Oily phenol [Phenol oily] .....	14
Mylan Fentanyl Patch .....	102	Drugs .....	93	Olanzapine .....	113, 115
Myleran .....	123	Nonacog alfa [Recombinant		Olanzine .....	113
<b>- N -</b>		factor IX] .....	27	Olanzine-D .....	113
Nadolol .....	38	Noradrenaline .....	44	Olive oil .....	182
Naloxone hydrochloride .....	173	Norethisterone		Olopatadine .....	168
Naltraccord .....	121	Genito-Urinary .....	54	Olisalazine .....	13
Naltrexone hydrochloride .....	121	Hormone .....	60	Omeprazole .....	14
Naphazoline hydrochloride .....	168	Norethisterone with		Omezol Relief .....	14
Naphcon Forte .....	168	mestranol .....	53	Omnipaque .....	176
Naproxen .....	95	Norfloxacin .....	67	Omniscan .....	177
Naropin .....	100	Normison .....	118	On Call Advanced .....	203
Natamycin .....	167	Norpress .....	105	Oncaspar .....	126
Natulan .....	127	Nortriptyline hydrochloride .....	105	OncoTICE .....	159
Nausicalm .....	111	Norvir .....	78	Ondanaccord .....	112
Navelbine .....	132	Novasource Renal (Vanilla) .....	195	Ondansetron .....	112
Navoban .....	112	Novatretin .....	51	One-Alpha .....	23
Nedocromil .....	165	NovoMix 30 FlexPen .....	15	Onkotrone .....	124
Nefopam hydrochloride .....	101	NovoSeven RT .....	27	Onrex .....	112
Neocate Advance (Vanilla) .....	192	Noxafil .....	70	Optional Pharmaceuticals .....	203
Neocate Gold (Unflavoured) .....	192	Nupentin .....	107	Ora-Blend .....	182
Neoral .....	134	Nutrini Energy Multi Fibre .....	194	Ora-Blend SF .....	182
NeoRecormon .....	25	Nutrini Low Energy Multifibre		Ora-Plus .....	182
Neostigmine metilsulfate .....	86	RTH .....	194	Ora-Sweet .....	182
Neostigmine metilsulfate with		Nutrison Concentrated .....	190	Ora-Sweet SF .....	182
glycopyrronium bromide .....	86	Nutrison Energy .....	196	Oracort .....	22
Neosynephrine HCL .....	44	Nyefax Retard .....	39	Oratane .....	48
Neotigason .....	51	Nystatin		Ornidazole .....	73
Nepro (Strawberry) .....	195	Alimentary .....	22	Orphenadrine citrate .....	93
Nepro (Vanilla) .....	195	Dermatological .....	47	Orphenadrine hydrochloride .....	96
Nepro HP (Strawberry) .....	195	Genito-Urinary .....	53	Oruvail SR .....	94
Nepro HP (Vanilla) .....	195	Infection .....	69	Oseltamivir .....	83
Nepro HP RTH .....	195	<b>- O -</b>		Osmolite .....	197
Nepro RTH .....	195	Obstetric Preparations .....	54	Osmolite RTH .....	197
Neulastim .....	31	Octocog alfa [Recombinant factor		Ospamox .....	65
Neupogen .....	31	VIII] .....	27	Other Cardiac Agents .....	43
Nevirapine .....	75	Octreotide .....	132	Other Endocrine Agents .....	59
Nevirapine Alphapharm .....	75	Octreotide MaxRx .....	132	Other Oestrogen	
Nicorandil .....	45	Ocular Lubricants .....	171	Preparations .....	59
Nicotine .....	121	Oestradiol .....	58, 60	Other Otological	

Preparations .....	172	Dermatological .....	49	Phenol oily .....	14
Other Progestogen .....		Extemporaneous .....	182	Phenol with ioxaglic acid .....	179
Preparations .....	60	Paraffin liquid with soft white .....		Phenoxymethylpenicillin .....	
Other Skin Preparations .....	52	paraffin .....	172	hydrochloride .....	36
Oxaliplatin .....	128	Paraffin liquid with wool fat .....	172	Phenoxybenzamine .....	
Oxaliplatin Actavis 100 .....	128	Paraffin with wool fat .....	49	[Penicillin V] .....	66
Oxaliplatin Actavis 50 .....	128	Paraldehyde .....	107	Phentolamine mesylate .....	36
Oxandrolone .....	57	Parecoxib .....	95	Phenylephrine hydrochloride .....	
Oxazepam .....	117	Paromomycin .....	63	Cardiovascular .....	44
Oxpentifylline .....	45	Paroxetine hydrochloride .....	106	Sensory .....	171
Oxybuprocaine .....		Paser .....	72	Phenytoin .....	109
hydrochloride .....	169	Patent blue V .....	178	Phenytoin sodium .....	107, 109
Oxybutynin .....	56	Paxam .....	117	Pholcodine .....	163
Oxycodone hydrochloride .....	103	Pazopanib .....	129	Phosphorus .....	34
Oxycodone Orion .....	103	Peak flow meter .....	203	Phytomenadione .....	28
OxyContin .....	103	Peanut oil .....	181	Picibanil .....	159
Oxydone BNM .....	103	Pediasure (Chocolate) .....	194	Pilocarpine hydrochloride .....	171
Oxymetazoline .....		Pediasure (Strawberry) .....	194	Pilocarpine nitrate .....	182
hydrochloride .....	163	Pediasure (Vanilla) .....	194	Pimafucort .....	51
OxyNorm .....	103	Pediasure RTH .....	194	Pindolol .....	38
Oxytocin .....	55	Pegaspargase .....	126	Pinetarsol .....	51
Oxytocin BNM .....	55	Pegasus RBV Combination .....		Pinorax .....	19
Oxytocin with ergometrine .....		Pack .....	84	Pinorax Forte .....	19
maleate .....	55	Pegasys .....	84	Pioglitazone .....	17
Ozole .....	70	Pegfilgrastim .....	31	Piperacillin with tazobactam .....	66
- P - .....		Pegylated interferon alfa-2a .....	84	Pipothiazine palmitate .....	116
Pacifen .....	92	Penembact .....	63	Pituitary and Hypothalamic .....	
Pacific Buspirone .....	117	Penicillamine .....	86	Hormones and Analogues .....	60
Paclitaxel .....	131	Penicillin G .....	66	Pivmecillinam .....	68
Paclitaxel Actavis .....	131	Penicillin V .....	66	Pizaccord .....	17
Paclitaxel Ebewe .....	131	Pentagastrin .....	59	Pizotifen .....	111
Paliperidone .....	116	Pentamidine isethionate .....	73	PKU Anamix Junior LQ .....	
Pamidronate BNM .....	88	Pentasa .....	13	(Berry) .....	187
Pamidronate disodium .....	88	Pentostatin .....		PKU Anamix Junior LQ .....	
Pamisol .....	88	[Deoxycytoformycin] .....	127	(Orange) .....	187
Panadol .....	101	Pentoxifylline [Oxpentifylline] .....	45	PKU Anamix Junior LQ .....	
Pancreatic enzyme .....	17	Peptamen OS 1.0 (Vanilla) .....	190	(Unflavoured) .....	187
Pancuronium bromide .....	93	Peptisoothe .....	14	Plaquenil .....	86
Pantoprazole .....	15	Perflutren .....	178	Plendil ER .....	39
Pantoprazole Actavis 20 .....	15	Perhexiline maleate .....	40	pms-Bosentan .....	45
Pantoprazole Actavis 40 .....	15	Pericyazine .....	113	Pneumococcal (PCV10) .....	
Papaverine hydrochloride .....	45	Perindopril .....	35	conjugate vaccine .....	199
Paper wasp venom .....	161	Permethrin .....	48	Pneumococcal (PCV13) .....	
Para-aminosalicylic Acid .....	72	Peteha .....	72	conjugate vaccine .....	199
Paracare .....	101	Pethidine hydrochloride .....	104	Pneumococcal (PPV23) .....	
Paracare Double Strength .....	101	Pexsig .....	40	polysaccharide vaccine .....	200
Paracetamol .....	101	Phenelzine sulphate .....	105	Podophyllotoxin .....	52
Paracetamol + Codeine .....		Phenindione .....	29	Polidocanol .....	26
(Relieve) .....	104	Phenobarbitone .....	109, 118	Poliomyelitis vaccine .....	201
Paracetamol with codeine .....	104	Phenobarbitone sodium .....	182	Poloxamer .....	18
Paracetamol-AFT .....	101	Phenol .....		Poly Gel .....	171
Paraffin .....		Extemporaneous .....	182	Poly Tears .....	172
Alimentary .....	18	Various .....	179	Poly-Visc .....	172



Polyhexamethylene biguanide .....	182	Progesterone .....	55	Ramipex .....	97
Polyvinyl alcohol .....	172	Proglycem .....	15	Ranbaxy-Cefaclor .....	64
Polyvinyl alcohol with povidone .....	172	Prograf .....	134	Ranibizumab .....	150
Poractant alfa .....	166	Prokinex .....	111	Ranitidine .....	14
Posaconazole .....	70	Promethazine hydrochloride .....	162	Rapamune .....	159
Postinor-1 .....	54	Promethazine theoclate .....	112	Rasburicase .....	92
Potassium chloride .....	32, 34	Propafenone hydrochloride .....	37	Reandron 1000 .....	57
Potassium chloride with sodium chloride .....	33	Propamide isethionate .....	167	Recombinant factor IX .....	27
Potassium citrate .....	56	Propofol .....	98	Recombinant factor VIIa .....	27
Potassium dihydrogen phosphate .....	33	Propranolol .....	39	Recombinant factor VIII .....	27
Potassium iodate		Propylene glycol .....	182	Rectogesic .....	14
Alimentary .....	20	Propylthiouracil .....	61	Red back spider antivenom .....	174
Hormone .....	61	Prostin E2 .....	55	Redipred .....	58
Potassium iodate with iodine .....	20	Prostin VR .....	44	Relenza Rotadisk .....	83
Potassium perchlorate .....	61	Protamine sulphate .....	29	Remicade .....	145
Potassium permanganate .....	51	Protionamide .....	72	Remifentanil hydrochloride .....	104
Povidone K30 .....	182	Protirelin .....	61	Remifentanil-AFT .....	104
Povidone-iodine .....	175	Provera .....	59, 60	ReoPro .....	139
Povidone-iodine with ethanol .....	175	Provisc .....	170	Resource Beneprotein .....	185
Pradaxa .....	28	Provive MCT-LCT 1% .....	98	Resource Diabetic (Vanilla) .....	189
Pralidoxime iodide .....	173	Proxymetacaine hydrochloride .....	169	Respiratory Stimulants .....	166
Pramipexole hydrochloride .....	97	Pseudoephedrine hydrochloride .....	163	Retinol .....	23
Prasugrel .....	30	Psoriasis and Eczema Preparations .....	51	Retinol Palmitate .....	172
Pravastatin .....	42	PTU .....	61	Retrovir .....	77
Praziquantel .....	73	Pulmocare (Vanilla) .....	195	Revolade .....	26
Prazosin .....	36	Pulmonary Surfactants .....	166	Reyataz .....	77
Prednisolone .....	58	Pulmozyme .....	165	Riboflavin 5-phosphate .....	170
Prednisolone acetate .....	168	Puri-nethol .....	125	Ridal .....	114
Prednisolone sodium phosphate .....	168	Pyrazinamide .....	72	Rifabutin .....	72
Prednisone .....	58	Pyridostigmine bromide .....	86	Rifampicin .....	72
Pregnancy test - hCG urine .....	204	PyridoxADE .....	23	Rilutek .....	96
Prezista .....	77	Pyridoxal-5-phosphate .....	20	Riluzole .....	96
Prilocaine hydrochloride .....	100	Pyridoxine hydrochloride .....	23	Ringer's solution .....	33
Prilocaine hydrochloride with felypressin .....	100	Pyrimethamine .....	74	Riodine .....	175
Primaquine phosphate .....	74	Pytazen SR .....	30	Risedronate Sandoz .....	91
Primaxin .....	63	- Q -		Risedronate sodium .....	91
Primidone .....	109	Q 300 .....	74	Risperdal .....	114
Primolut N .....	60	Quetapel .....	114	Risperdal Consta .....	116
Probenecid .....	92	Quetiapine .....	114	Risperdal Quicklet .....	114
Procaine penicillin .....	66	Quinapril .....	35	Risperidone .....	114, 116
Procarbazine hydrochloride .....	127	Quinapril with hydrochlorothiazide .....	35	Risperon .....	114
Prochlorperazine .....	112	Quinine dihydrochloride .....	74	Ritalin .....	119
Proctosedyl .....	13	Quinine sulphate .....	74	Ritalin LA .....	119
Procydiline hydrochloride .....	96	- R -		Ritalin SR .....	119
Procytox .....	123	RA-Morph .....	102	Ritonavir .....	78
Prodopa .....	40	Rabies vaccine .....	201	Rituximab .....	151
		Raloxifene .....	90	Rivaroxaban .....	29
		Raltegravir potassium .....	78	Rivotril .....	106
				Rizamelt .....	111
				Rizatriptan benzoate .....	111
				Rocuronium bromide .....	93
				Ropinrole hydrochloride .....	97
				Ropivacaine hydrochloride .....	100

Ropivacaine hydrochloride with fentanyl ..... 100	Siterone ..... 57	Sodium phenylbutyrate ..... 20
Rose bengal sodium ..... 169	Slow-Lopresor ..... 38	Sodium phosphate with phosphoric acid ..... 19
Roxane ..... 12	Snake antivenom ..... 174	Sodium polystyrene sulphonate ..... 34
Roxithromycin ..... 65	Sodibic ..... 34	Sodium stibogluconate ..... 74
Rubifen ..... 119	Sodium acetate ..... 33	Sodium tetradecyl sulphate ..... 26
Rubifen SR ..... 119	Sodium acid phosphate ..... 33	Sodium thiosulfate ..... 173
<b>- S -</b>		
S-26 Gold Premgro ..... 193	Sodium alginate with magnesium alginate ..... 12	Sodium valproate ..... 109
S26 LBW Gold RTF ..... 193	Sodium alginate with sodium bicarbonate and calcium carbonate ..... 12	Sodium with potassium ..... 180
Salamol ..... 163	Sodium aurothiomalate ..... 86	Solian ..... 112
Salazopyrin ..... 13	Sodium benzoate ..... 20	Solifenacin succinate ..... 56
Salazopyrin EN ..... 13	Sodium bicarbonate	Solox ..... 14
Salbutamol ..... 163	Blood ..... 33-34	Solu-Cortef ..... 58
Salbutamol with ipratropium bromide ..... 162	Extemporaneous ..... 183	Solu-Medrol ..... 58
Salicylic acid ..... 183	Sodium calcium edetate ..... 174	Somatropin ..... 61
Salmeterol ..... 164	Sodium carboxymethylcellulose with pectin and gelatine ..... 22	Sotacor ..... 39
Salmonella typhi vaccine ..... 200	Sodium chloride	Sotalol ..... 39
Sandimmun ..... 134	Blood ..... 33-34	Soya oil ..... 173
Sandomigran ..... 111	Respiratory ..... 163, 166	Space Chamber Plus ..... 204
Sandostatin LAR ..... 132	Various ..... 179	Spacer device ..... 204
Scalp Preparations ..... 51	Sodium chloride with sodium bicarbonate ..... 163	Span-K ..... 34
Sclerosing Agents ..... 166	Sodium citrate	Specialised Formulas ..... 188
Scopoderm TTS ..... 111	Alimentary ..... 12	Spiractin ..... 41
Sebizole ..... 47	Extemporaneous ..... 183	Spiramycin ..... 74
Secretin pentahydrochloride ..... 178	Sodium citrate with sodium chloride and potassium chloride ..... 29	Spiriva ..... 162
Sedatives and Hypnotics ..... 118	Sodium citrate with sodium lauryl sulphoacetate ..... 19	Spirolactone ..... 41
Selegiline hydrochloride ..... 97	Sodium citro-tartrate ..... 56	Spirotone ..... 41
Sennosides ..... 19	Sodium cromoglycate	Sprycel ..... 128
Serenace ..... 113	Alimentary ..... 13	Standard Feeds ..... 196
Seretide ..... 165	Respiratory ..... 161, 165	Staphlex ..... 66
Seretide Accuhaler ..... 165	Sensory ..... 168	Starch ..... 183
Serevent ..... 164	Sodium dihydrogen phosphate [Sodium acid phosphate] ..... 33	Stavudine ..... 77
Serevent Accuhaler ..... 164	Sodium fluoride ..... 20	Sterculia with frangula ..... 18
Serophene ..... 59	Sodium hyaluronate	Stesolid ..... 106
Seroquel ..... 114	Alimentary ..... 22	Stimulants / ADHD Treatments ..... 118
Sertraline ..... 106	Sensory ..... 170, 172	Stiripentol ..... 109
Sevoflurane ..... 98	Sodium hyaluronate with chondroitin sulphate ..... 170	Stocrin ..... 75
Sevredol ..... 103	Sodium hypochlorite ..... 175	Strattera ..... 118
Silagra ..... 46	Sodium metabisulfite ..... 183	Streptomycin sulphate ..... 63
Sildenafil ..... 46	Sodium nitrite ..... 173	Stromectol ..... 73
Silver nitrate	Sodium nitroprusside	Suboxone ..... 120
Dermatological ..... 52	Cardiovascular ..... 45	Sucalfate ..... 15
Extemporaneous ..... 183	Part III - OPTIONAL PHARMACEUTICALS ..... 204	Sucrose ..... 101
Simethicone ..... 12		Sugammadex ..... 93
Simulect ..... 145		Sulindac ..... 95
Simvastatin ..... 42		Sulphacetamide sodium ..... 167
Sincalide ..... 178		Sulphadiazine ..... 68
Sinemet ..... 97		Sulphadiazine silver ..... 47
Sinemet CR ..... 97		Sulphasalazine ..... 13
Singular ..... 164		Sulphur ..... 183
Sirolimus ..... 159		Sumatriptan ..... 111

Sunitinib .....	130	Sensory .....	169	Travoprost .....	171
Sunscreen, proprietary .....	52	Tetracosactide .....		Treatments for Dementia .....	120
Suprane .....	97	[Tetracosactrin] .....	60	Treatments for Substance	
Surgical Preparations .....	179	Tetracosactrin .....	60	Dependence .....	120
Survanta .....	166	Tetracyclin Wolff .....	67	Tretinoin .....	
Sustagen Diabetic (Vanilla) .....	189	Tetracycline .....	67	Dermatological .....	48
Sustagen Hospital Formula		Thalidomide .....	127	Oncology .....	127
(Chocolate) .....	197	Thalomid .....	127	Trexate .....	125
Sustagen Hospital Formula		Theophylline .....	165	Tri-sodium citrate .....	183
(Vanilla) .....	197	Thiamine hydrochloride .....	23	Triamcinolone acetoneide	
Sutent .....	130	Thioguanine .....	125	Alimentary .....	22
Suxamethonium chloride .....	93	Thiopental [Thiopentone]		Dermatological .....	50
Symmetrel .....	96	sodium .....	98	Hormone .....	58
Sympathomimetics .....	44	Thiopentone .....	98	Triamcinolone acetoneide with	
Synacthen .....	60	Thiotepa .....	123	gramicidin, neomycin and	
Synacthen Depot .....	60	Thrombin .....	26	nystatin .....	168
Syntometrine .....	55	Thymol glycerin .....	22	Triamcinolone acetoneide with	
Syrup .....	183	Thyroid and Antithyroid		neomycin sulphate, gramicidin	
Systane Unit Dose .....	172	Preparations .....	61	and nystatin .....	51
<b>- T -</b>		Thyrotropin alfa .....	60	Triamcinolone hexacetoneide .....	58
Tacrolimus .....	134	Ticagrelor .....	30	Triazolam .....	118
Tacrolimus Sandoz .....	134	Ticarcillin with clavulanic acid .....	66	Trichloracetic acid .....	183
Talc .....	166	Ticlopidine .....	30	Trichozole .....	73
Tambocor .....	37	Tigecycline .....	67	Trientine dihydrochloride .....	20
Tambocor CR .....	37	Timolol .....	170	Trifluoperazine	
Tamoxifen citrate .....	133	Timolol maleate .....	39	hydrochloride .....	115
Tamsulosin .....	56	Tioptol XE .....	170	Trimeprazine tartrate .....	162
Tamsulosin-Rex .....	56	Tiotropium bromide .....	162	Trimethoprim .....	69
Tarceva .....	128	TMP .....	69	Trimethoprim with	
Tasmar .....	97	Tobramycin		sulphamethoxazole	
Tazocin EF .....	66	Infection .....	63	[Co-trimoxazole] .....	69
Teicoplanin .....	69	Sensory .....	167	Trisodium citrate .....	29
Temaccord .....	127	Tobrex .....	167	Trometamol .....	179
Temazepam .....	118	Tocilizumab .....	157	Tropicamide .....	171
Temozolomide .....	127	Tofranil .....	105	Tropisetron .....	112
Tenecteplase .....	31	Tolcapone .....	97	Tropisetron-AFT .....	112
Tenofovir disoproxil fumarate .....	80	Tolterodine tartrate .....	56	Truvada .....	76
Tenoxicam .....	95	Topamax .....	110	Tuberculin, purified protein	
Terazosin .....	36	Topical Products for Joint and		derivative .....	178
Terbinafine .....	71	Muscular Pain .....	95	Two Cal HN .....	190
Terbutaline .....	55	Topiramate .....	110	TwoCal HN RTH (Vanilla) .....	190
Terbutaline sulphate .....	163	Tracleer .....	45	Tykerb .....	129
Teriparatide .....	91	Tracrrium .....	92	<b>- U -</b>	
Terlipressin .....	62	Tramadol hydrochloride .....	104	Ultraproct .....	13
Testosterone .....	57	Tramal 100 .....	104	Univent .....	162
Testosterone cypionate .....	57	Tramal 50 .....	104	Ural .....	56
Testosterone esters .....	57	Tramal SR 100 .....	104	Urea	
Testosterone undecanoate .....	57	Tramal SR 150 .....	104	Dermatological .....	49
Tetrabenazine .....	96	Tramal SR 200 .....	104	Extemporaneous .....	183
Tetracaine [Amethocaine]		Trandolapril .....	35	Urex Forte .....	41
hydrochloride		Tranexamic acid .....	27	Urokinase .....	31
Nervous .....	100	Tranylcypromine sulphate .....	105	Urologicals .....	55
		Trastuzumab .....	157	Uromitexan .....	132

Ursodeoxycholic acid .....	17	Vital HN .....	190	Zarzio .....	31
Ursosan .....	17	Vitamin A with vitamins D and C .....	23	Zavedos .....	124
Utrogestan .....	55	Vitamin B complex .....	23	Zeldox .....	115
<b>- V -</b>		Vitamins .....	22	Zetlam .....	79
Valaciclovir .....	82	Vivonex Paediatric .....	192	Zetop .....	161
Valcyte .....	82	Vivonex TEN .....	189	Ziagen .....	76
Valganciclovir .....	82	Volibris .....	45	Zidovudine [AZT] .....	77
Valtrex .....	82	Voltaren .....	94	Zidovudine [AZT] with lamivudine .....	77
Vancomycin .....	69	Voltaren Ophtha .....	168	Zinc .....	
Varenicline .....	121	Volulyte 6% .....	34	Alimentary .....	21
Varicella zoster vaccine [Chicken pox vaccine] .....	201	Volumatic .....	204	Dermatological .....	48
Vasodilators .....	44	Voluven .....	34	Zinc and castor oil .....	48
Vasopressin .....	61	Voriconazole .....	70	Zinc chloride .....	21
Vasopressin Agents .....	61	Votrient .....	129	Zinc oxide .....	183
Vecuronium bromide .....	93	<b>- W -</b>		Zinc sulphate .....	21
Velcade .....	125	Warfarin sodium .....	30	Zinc with wool fat .....	49
Venlafaxine .....	106	Wart Preparations .....	52	Zincaps .....	21
Venofer .....	21	Water .....		Zinnat .....	64
Ventavis .....	46	Blood .....	34	Ziprasidone .....	115
Ventolin .....	163	Various .....	179	Zithromax .....	64
Vepesid .....	126	Wool fat .....		Zofran Zydis .....	112
Verapamil hydrochloride .....	40	Dermatological .....	49	Zoladex .....	60
Vergo 16 .....	111	Extemporaneous .....	183	Zoledronic acid .....	
Verpamil SR .....	40	<b>- X -</b>		Hormone .....	57
Vesanoid .....	127	Xanthan .....	183	Musculoskeletal System .....	88
Vesicare .....	56	Xarelto .....	29	Zometa .....	57
Vfend .....	70	Xeloda .....	124	Zopiclone .....	118
Victrelis .....	81	Xylocaine .....	99, 100	Zostrix .....	95
Vigabatrin .....	110	Xylocaine Viscous .....	99	Zostrix HP .....	101
Vimpat .....	108	Xylometazoline hydrochloride .....	163	Zovirax IV .....	82
Vinblastine sulphate .....	132	Xyntha .....	27	Zuclopenthixol acetate .....	115
Vincristine sulphate .....	132	<b>- Y -</b>		Zuclopenthixol decanoate .....	117
Vinorelbine .....	132	Yellow jacket wasp venom .....	161	Zuclopenthixol hydrochloride .....	115
Viral Vaccines .....	200	<b>- Z -</b>		Zyban .....	121
Viramune Suspension .....	75	Zanamivir .....	83	Zypine .....	113
Viread .....	80	Zantac .....	14	Zypine ODT .....	113
Visipaque .....	175	Zapril .....	35	Zyprexa Relprevv .....	115
Vistil .....	172	Zarator .....	42		
Vistil Forte .....	172				
VitA-POS .....	172				







