Index 252

Introducing PHARMAC 2

August 2015

Volume 22 Number 2

Editor: Kaye Wilson email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am - 5pm weekdays)

Circulation

Published each April, August and December. Changes to the contents are published in monthly updates.

Accessible in an electronic format at no cost from the Health Professionals section of the PHARMAC website www.pharmac.govt.nz

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month. Alternatively there is a nominal charge for an annual subscription to the printed Schedule publications. To access either of these subscriptions visit our subscription website www.schedule.co.nz.

Production

Typeset automatically from XML and TEX. XML version of the Schedule available from www.pharmac.govt.nz/schedule/archive/

Programmers

Anrik Drenth & John Geering email: texschedule@pharmac.govt.nz

©Pharmaceutical Management Agency



ISSN 1179-3686 pdf ISSN 1172-9376 print

This work is licensed under the Creative Commons Attribution 3.0 New Zealand licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to PHARMAC and abide by the other licence terms. To view a copy of this licence, visit:

creativecommons.org/licenses/by/3.0/nz/.

Attribution to PHARMAC should be in written form and not by reproduction of the PHARMAC logo. While care has been taken in compiling this Schedule, PHARMAC takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

| Section / | 4 |
|-----------|---|
|-----------|---|

Section B

Alimentary Tract & Metabolism 20 Blood & Blood Forming Organs 40 Cardiovascular System 49 Dermatologicals 62 Genito Urinary System 73 Hormone Preparations – Systemic 79 Infections – Agents For Systemic Use 91 Musculoskeletal System 115

- Nervous System 124
- Oncology Agents & Immunosuppressants 157
 - Respiratory System & Allergies 193
 - Sensory Organs 200

Various 204

Section C Extemporaneous Compounds (ECPs) 206

Section DSpecial Foods213Section EPractitioner's Supply Orders235
Rural Areas239Section FDispensing Period Exemptions240Section GSafety Cap Medicines242Section INational Immunisation Schedule245

General Rules 6

Introducing PHARMAC

The Pharmaceutical Management Agency (PHARMAC) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. PHARMAC negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

PHARMAC's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about PHARMAC and the way we make funding decisions can be found on the PHARMAC website at http://www.pharmac.health.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by PHARMAC.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements PHARMAC has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A through to G and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

| gramg kilogramkg | milligrammg | m ur |
|----------------------|---------------|---------|
| international unitiu | millilitre ml | |

| millimole | mmol |
|-----------|------|
| unit | u |

Abbreviations Ampoule ...

| Ampoule | Amp | Gelatinous | Gel |
|----------------|------|------------|------|
| Capsule | Сар | Granules | Gran |
| Cream | | Infusion | Inf |
| Device | Dev | Injection | Inj |
| Dispersible | Disp | Liquid | Liq |
| Effervescent | Eff | | LA |
| Emulsion | Emul | Ointment | Oint |
| Enteric Coated | EC | Sachet | Sach |
| | | | |

| Solution | Soln |
|-----------------------|------|
| Suppository | Supp |
| Tablet | Tab |
| Tincture | Tinc |
| Trans Dermal Delivery | |
| System | TDDS |
| | |

BSO Bulk Supply Order.

CBS Cost Brand Source.

ECP Extemporaneously Compounded Preparation.

- Original Pack subsidy is rounded up to a multiple at whole packs. OP
- PSO Practitioner's Supply Order.

Sole Subsidised

Supplier Only brand of this medicine subsidised.

XPharm Pharmacies cannot claim subsidy because PHARMAC has made alternative distribution arrangements.

Three months supply may be dispensed at one time if the exempted medicine is endorsed 'certified exemption' by the practitioner or pharmacist.

- Three months dispensed all-at-once or, in the case of oral contraceptives, six months dispensed all-at-once, unless the * medicine meets the Dispensing Frequency Rule criteria.
- Safety cap required for oral liquid formulations, including extemporaneously compounded preparations. t
- Fully subsidised brand of a given medicine. Brands without the tick are not fully subsidised and may cost the patient a manufacturer's surcharge.
- S29 This medicine is an unapproved medication supplied under Section 29 of the Medicines Act 1981.
- HP3 Subsidised when dispensed from a pharmacy that has a contract to dispense Special Foods.
- HP4 Subsidised when dispensed from a pharmacy that has a contract to dispense from the Monitored Therapy Variation (for Clozapine Services).

Community Pharmaceutical costs met by the Government

Most of the cost of a subsidised prescription for a Community Pharmaceutical is met by the Government through the Combined Pharmaceutical Budget. The Government pays a subsidy for the Community Pharmaceutical to pharmacies, and a fee covering distribution and pharmacy dispensing services. The subsidy paid to pharmacies does not necessarily represent the final cost to Government of subsidising a particular Community Pharmaceutical. The final cost will depend on the nature of PHARMAC's contractual arrangements with the supplier. Fully subsidised medicines are identified with a V in the product's Schedule listing.

Patient costs

Everyone who is eligible for publicly funded health and disability services should in most circumstances pay only a \$5 co-payment for subsidised medicines, although co-payments can vary from \$0 to \$15. Where the price of a Pharmaceutical is higher than the subsidy, a patient may pay a manufacturer's surcharge in addition to the co-payment. A patient may also pay additional fees for services such as after-hours dispensing and special packaging.

Patients can check whether they are eligible for publicly funded health and disability services by referring to the Guide to eligibility on the Ministry of Health's website.

DHBs have a list of eligible providers in their respective regions. Any provider/prescriber not specifically listed by a DHB as an approved provider/prescriber should be regarded as not approved.

For more information on patient co-payments or eligibility please visit http://www.moh.govt.nz.

Special Authority Applications

Special Authority is an application process in which a prescriber requests government subsidy on a Community Pharmaceutical for a particular person.

Subsidy

Once approved, the applicant will be provided a Special Authority number which must appear on the prescription.

The authority number can provide access to subsidy, increased subsidy, or waive certain restrictions otherwise present on the Community Pharmaceutical.

Some approvals are dependent on the availability of funding from the Combined Pharmaceutical Budget.

Criteria

The criteria for approval of Special Authority applications are included below each Community Pharmaceutical listing, and on the application forms available on PHARMAC's website. For some Special Authority Community Pharmaceuticals, not all indications that have been approved by Medsafe are subsidised.

Making a Special Authority application

Application forms can be found at http://www.pharmac.govt.nz. Except where stated on the application form, applications are processed by the Ministry of Health, and are sent to:

Ministry of Health Sector Services, Fax: (06) 349 1983 or free fax 0800 100 131

Private Bag 3015, WANGANUI 4540

To register for submission of applications on-line - Contact the Ministry of Health on 0800 505 125 or email at

onlinehelpdesk@moh.govt.nz. For Special Authority approval numbers, applicants can phone the Ministry of Health Sector Services Call Centre, free phone 0800 243 666.

Named Patient Pharmaceutical Assessment policy

Named Patient Pharmaceutical Assessment (NPPA) provides a mechanism for individual patients to receive funding for medicines not listed in the Pharmaceutical Schedule (either at all or for their clinical circumstances). PHARMAC will assess applications that meet the prerequisites according to its Decision Criteria before deciding whether to approve applications for funding. The Decision Criteria will be used to assess both the individual clinical circumstances of each NPPA applicant, and the implications of each NPPA funding decision on PHARMAC's ability to carry out its legislative functions.

For more information on NPPA, or to apply, visit the PHARMAC website at

http://www.pharmac.health.nz/link/nppa, or call the Panel Coordinators at 0800 660 050 Option 2.

INTRODUCTION

Section A contains the restrictions and other general rules that apply to Subsidies on Community Pharmaceuticals. The amounts payable by the Funder to Contractors are currently determined by:

- the quantities, forms, and strengths, of subsidised Community Pharmaceuticals dispensed under valid prescription by each Contractor;
- the amount of the Subsidy on the Manufacturer's Price payable for each unit of the Community Pharmaceuticals dispensed by each Contractor and;
- the contractual arrangements between the Contractor and the Funder for the payment of the Contractor's dispensing services.

The Pharmaceutical Schedule shows the level of subsidy payable in respect of each Community Pharmaceutical so that the amount payable by the Government to Contractors, for each Community Pharmaceutical, can be calculated. The Pharmaceutical Schedule also shows the standard price (exclusive of GST) at which a Community Pharmaceutical is supplied ex-manufacturer to wholesalers if it differs from the subsidy. The manufacturer's surcharge to patients can be estimated using the subsidy and the standard manufacturer's price as set out in this Schedule.

The cost to Government of subsidising each Community Pharmaceutical and the manufacturer's prices may vary, in that suppliers may provide rebates to other stakeholders in the primary health care sector, including dispensers, wholesalers, and the Government. Rebates are not specified in the Pharmaceutical Schedule.

This Schedule is dated 1 August 2015 and is to be referred to as the Pharmaceutical Schedule Volume 22 Number 2, 2015. Distribution will be from 20 August 2015. This Schedule comes into force on 1 August 2015.

PART I INTERPRETATIONS AND DEFINITIONS

1.1 In this Schedule, unless the context otherwise requires:

"90 Day Lot", means the quantity of a Community Pharmaceutical required for the number of days' treatment covered by the Prescription, being up to 90 consecutive days' treatment;

"**180 Day Lot**", means the quantity of a Community Pharmaceutical required for the number of days' treatment covered by the Prescription, being up to 180 consecutive days' treatment;

"Access Exemption Criteria", means the criteria under which patients may receive greater than one Month's supply of a Community Pharmaceutical covered by Section F Part II (b) subsidised in one Lot. The specifics of these criteria are conveyed in the Ministry of Health guidelines, which are issued from time to time. The criteria the patient must meet are that they:

- a) have limited physical mobility;
- b) live and work more than 30 minutes from the nearest pharmacy by their normal form of transport;
- c) are relocating to another area;
- d) are travelling extensively and will be out of town when the repeat prescriptions are due.
- "Act", means the New Zealand Public Health and Disability Act 2000.

"Advisory Committee", means the Pharmaceutical Services Advisory Committee convened by the Ministry of Health under the terms of the Advice Notice issued to Contractors pursuant to Section 88 of the Act.

"Alternate Subsidy", means a higher level of subsidy that the Government will pay contractors for a particular community Pharmaceutical dispensed to a person who has either been granted a Special Authority for that pharmaceutical, or where the prescription is endorsed in accordance with the requirements of this Pharmaceutical Schedule.

"Annotation", means written annotation of a prescription by a dispensing pharmacist in the pharmacist's own handwriting following confirmation from the Prescriber if required, and "Annotated" has a corresponding meaning. The Annotation must include the details specified in the Schedule, including the date the prescriber was contacted (if applicable) and be initialled by the dispensing pharmacist.

"Authority to Substitute", means an authority for the dispensing pharmacist to change a prescribed medicine in accordance with regulation 42(4) of the Medicines Regulations 1984. An authority to substitute letter, which may be used by Practitioners, is available on the final page of the Schedule.

"Bulk Supply Order", means a written order, on a form supplied by the Ministry of Health, or approved by the Ministry of Health, made by the licensee or manager of an institution certified to provide hospital care under the Health and Disability Services (Safety) Act 2001 for the supply of such Community Pharmaceuticals as are expected to be required for the treatment of persons who are under the medical or dental supervision of such a Private Hospital or institution.

"Class B Controlled Drug", means a Class B controlled drug within the meaning of the Misuse of Drugs Act 1975. "Community Pharmaceutical", means a Pharmaceutical listed in Sections A to G and Section I of the Pharmaceutical Schedule that is subsidised by the Funder from the Pharmaceutical Budget for use in the community.

"Contractor", means a person who is entitled to receive a payment from the Crown or a DHB under a notice issued by the Crown or a DHB under Section 88 of the Act or under a contract with the Ministry of Health or a DHB for the supply of Community Pharmaceuticals.

"Controlled Drug", means a controlled drug within the meaning of the Misuse of Drugs Act 1975 (other than a controlled drug specified in Part VI of the Third Schedule to that Act).

"Cost, Brand, Source of Supply", means that the Community Pharmaceutical is eligible for Subsidy on the basis of the Contractor's annotated purchase price, brand, and source of supply. Alternatively a copy of the invoice for the purchase of the Pharmaceutical may be attached to the prescription, in the place of an annotation, in order to be eligible for Subsidy.

"Dentist", means a person registered with the Dental Council, and who holds a current annual practising certificate, under the HPCA Act 2003.

"Diabetes Nurse Prescriber", means a nurse who is a Designated Prescriber—Registered Nurses Practising in Diabetes Health as determined by the Nursing Council of New Zealand to practice in diabetes health and has authority to prescribe specified diabetes medicines in accordance with regulations made under the Medicines Act 1981.

"Dietitian", means a person registered as a dietitian with the Dietitians Board, and who holds a current annual practicing certificate under the HPCA Act 2003.

"DHB", means an organisation established as a District Health Board by or under Section 19 of the Act.

"DHB Hospital", means a DHB, including its hospital or associated provider unit that the DHB purchases Hospital Pharmaceuticals for.

"Dispensing Frequency Rule", means the rule in Part IV, Section A of the Pharmaceutical Schedule that defines patient groups or medicines eligible for more frequent dispensing periods.

"Doctor", means a medical Practitioner registered with the Medical Council of New Zealand and, who holds a current annual practising certificate under the HPCA Act 2003.

"DV Limit", means, for a particular Hospital Pharmaceutical with HSS, the National DV Limit or the Individual DV Limit.

"DV Pharmaceutical", means a discretionary variance Pharmaceutical, that does not have HSS and which:

- a) is either listed in Section H Part II of the Schedule as being a DV Pharmaceutical in association with the relevant Hospital Pharmaceutical with HSS; or
- b) is the same chemical entity, at the same strength, and in the same or a similar presentation or form, as the relevant Hospital Pharmaceutical with HSS, but which is not yet listed as being a DV Pharmaceutical.

"Endorsements", - unless otherwise specified, endorsements should be either handwritten or computer generated by the practitioner prescribing the medication. The endorsement can be written as "certified condition", or state the condition of the patient, where that condition is specified for the Community Pharmaceutical in Section B of the Pharmaceutical Schedule. Where the practitioner writes "certified condition" as the endorsement, he/she is making a declaration that the patient meets the criteria as set out in Section B of the Pharmaceutical Schedule.

"Funder", means the body or bodies responsible, pursuant to the Act, for the funding of pharmaceuticals listed on the Schedule (which may be one or more DHBs and/or the Ministry of Health) and their successors.

"GST", means goods and services tax under the Goods and Services Tax Act 1985.

"Hospital Care Operator", means a person for the time being in charge of providing hospital care, in accordance with the Health and Disability Services (Safety) Act 2001.

"Hospital Pharmaceuticals", means the list of pharmaceuticals set out in Section H part II of the Schedule which includes some National Contract Pharmaceuticals.

"Hospital Pharmacy", means that the Community Pharmaceutical is not eligible for Subsidy unless it is supplied by a hospital or pharmacy contracted to the Funder to dispense as a hospital pharmacy to an person on the Prescription of a Practitioner.

"Hospital Pharmacy-Specialist", means that the Community Pharmaceutical is not eligible for Subsidy unless it is supplied by a hospital or pharmacy contracted to the Funder to dispense as a hospital pharmacy to an Outpatient either:

- a) on a Prescription signed by a Specialist, or
- b) where the treatment with the Community Pharmaceutical has been recommended by a Specialist, on the Prescription of a practitioner which is either:
 - endorsed with the words "recommended by [name of specialist and year of authorisation]" and signed by the Practitioner, or
 - ii) endorsed with the word 'protocol' which means "initiated in accordance with DHB hospital approved protocol",
 - iii) annotated by the dispensing pharmacist, following verbal confirmation from the Practitioner of the name of the Specialist and date of recommendation, with the words "recommended by [name of specialist and date of authorisation], confirmed by [practitioner]". Where the Contractor has an electronic record of such an Endorsement or Annotation from a previous prescription for the same Community Pharmaceutical written by a prescriber for the same patient, they may annotate the prescription accordingly.

"As recommended by a Specialist" to be interpreted as either:

- i) follows a substantive consultation with an appropriate Specialist;
- ii) the consultation to relate to the Patient for whom the Prescription is written;
- iii) consultation to mean communication by referral, telephone, letter, facsimile or email;
- iv) except in emergencies consultation to precede annotation of the Prescription; and
- v) both the specialist and the General Practitioner must keep a written record of the consultation; or
- a) treatment with the Community Pharmaceutical has been initiated in accordance with a DHB hospital approved protocol.

For the purposes of the definition it makes no difference whether or not the Specialist is employed by a hospital.

"Hospital Pharmacy-Specialist Prescription", means that the Community Pharmaceutical is not eligible for Subsidy unless it is supplied by a hospital or pharmacy contracted to the Funder to dispense as a hospital pharmacy:

- a) to an Outpatient; and
- b) on a Prescription signed by a Specialist.

For the purposes of this definition, a "specialist" means a doctor who holds a current annual practicing certificate and who satisfies the criteria set out in paragraphs (a) or (b) or (c) of the definitions of Specialist below.

"HSS", means hospital supply status, the status of being the brand of the relevant Hospital Pharmaceutical listed in Section H Part II as HSS, that DHBs are obliged to purchase subject to any DV Limit for that Hospital Pharmaceutical for the period of hospital supply, as awarded under an agreement between PHARMAC and the relevant pharmaceutical supplier.

"In Combination", means that the Community Pharmaceutical is only subsidised when prescribed in combination with another subsidised pharmaceutical as specified in Section B or C of the Pharmaceutical Schedule.

"Individual DV Limit", means, for a particular Hospital Pharmaceutical with HSS and a particular DHB Hospital, the discretionary variance limit, being the specified percentage of that DHB Hospital's Total Market Volume up to which that DHB Hospital may purchase DV Pharmaceuticals of that Hospital Pharmaceutical.

"Licensed Hospital", means a place or institution that is certified to provide hospital care within the meaning of the Health and Disability Services (Safety) Act 2001.

"Lot", means a quantity of a Community Pharmaceutical supplied in one dispensing.

"Manufacturer's Price", means the standard price at which a Community Pharmaceutical is supplied to wholesalers (excluding GST), as notified to PHARMAC by the supplier.

"Maternity hospital", means that the Community Pharmaceutical is not eligible for Subsidy unless it is supplied pursuant to a Bulk Supply Order to a maternity hospital certified under the Health and Disability Services (Safety) Act 2001.

"Midwife", means a person registered as a midwife with the Midwifery Council, and who holds a current annual practising certificate under the HPCA Act 2003.

"Month", means a period of 30 consecutive days.

"Monthly Lot", means the quantity of a Community Pharmaceutical required for the number of days' treatment covered by the Prescription, being up to 30 consecutive days' treatment;

"Named Patient Pharmaceutical Assessment Advisory Panel", means the panel of clinicians, appointed by the PHARMAC Board, that is responsible for advising, within its Terms of Reference, on Named Patient Pharmaceutical Assessment applications and Exceptional Circumstances renewal applications submitted after 1 March 2012 (EC renewal application form located at http://www.pharmac.govt.nz/healthpros/EC/ECForms)

"National Contract Pharmaceutical", means a Hospital Pharmaceutical for which PHARMAC has negotiated a

national contract and the Price.

"National DV Limit", means, for a particular Hospital Pharmaceutical with HSS, the discretionary variance limit, being the specified percentage of the Total Market Volume up to which all DHB Hospitals may collectively purchase DV Pharmaceuticals of that Hospital Pharmaceutical.

"National Immunisation Schedule", means Section I of the Pharmaceutical Schedule, which is a schedule administered by PHARMAC, being a schedule specifying a programme of vaccinations to promote immunity against the diseases specified in the schedule.

"Not In Combination", means that no Subsidy is available for any Prescription containing the Community Pharmaceutical in combination with other ingredients unless the particular combination of ingredients is separately specified in Section B or C of the Schedule, and then only to the extent specified.

"Nurse Prescriber", means a person who is a nurse practitioner in terms of the Medicines Act 1981, or a Diabetes Nurse Prescriber

"Optional Pharmaceuticals", means the list of National Contract Pharmaceuticals set out in Section H Part II of the Schedule

"Optometrist", means a person registered with the Optometrists and Dispensing Opticians Board with a scope of practice that includes prescribing medicines (TPA endorsement)

"Outpatient", in relation to a Community Pharmaceutical, means a person who, as part of treatment at a hospital or other institution under the control of a DHB, is prescribed the Community Pharmaceutical for consumption or use in the person's home.

"PCT", means Pharmaceutical Cancer Treatment in respect of which DHB hospital pharmacies and other Contractors can claim Subsidies.

"PCT only", means Pharmaceutical Cancer Treatment in respect of which only DHB hospital pharmacies can claim Subsidies.

"Penal Institution", means a penal institution, as that term is defined in The Penal Institutions Act 1954;

"PHARMAC", means the Pharmaceutical Management Agency established by Section 46 of the Act (PHARMAC). "Pharmaceutical", means a medicine, therapeutic medical device, or related product or related thing listed in Sections B to I of the Schedule.

"Pharmaceutical Benefits", means the right of:

- a) a person; and
- b) any member under 16 years of age of that person's family, to have made by the Government on his or her behalf, subject to any conditions for the time being specified in the Schedule, such payment in respect of any Community Pharmaceutical supplied to that person or family member under the order of a Practitioner in the course of his or her practice.

"Pharmaceutical Budget", means the pharmaceutical budget set for PHARMAC by the Crown for the subsidised supply of Community Pharmaceuticals and Pharmaceutical Cancer Treatments including for named patients in exceptional circumstances.

"Pharmaceutical Cancer Treatment", means Pharmaceuticals for the treatment of cancer, listed in Sections A to G of the Schedule and identified therein as a "PCT" or "PCT only" Pharmaceutical that DHBs must provide access to, for use in their hospitals, and/or in association with Outpatient services provided in their DHB Hospitals, in relation to the treatment of cancers.

"Pharmacist Prescriber", means a person registered with the Pharmacy Council of New Zealand, who holds a current annual practising certificate under the HPCA Act 2003, and is approved by the Pharmacy Council of New Zealand to prescribe specified prescription medicines relating to his/her scope of practice.

"Pharmacist", means a person registered with the Pharmacy Council of New Zealand and who holds a current annual practicing certificate under the HPCA Act 2003.

"Practitioner", means a Doctor, a Dentist, a Dietitian, a Midwife, a Nurse Prescriber, an Optometrist, a Quitcard Provider, or a Pharmacist Prescriber as those terms are defined in the Pharmaceutical Schedule.

"Practitioner's Supply Order", means a written order made by a Practitioner on a form supplied by the Ministry of Health, or approved by the Ministry of Health, for the supply of Community Pharmaceuticals to the Practitioner, which the Practitioner requires to ensure medical supplies are available for emergency use, teaching and demonstration purposes, and for provision to certain patient groups where individual prescription is not practicable.

"Prescription", means a quantity of a Community Pharmaceutical prescribed for a named person on a document signed by a Practitioner.

"Prescription Medicine", means any Pharmaceutical listed in Part I of Schedule 1 of the Medicines Regulations

1984.

"Private Hospital", means a hospital certified under the Health and Disability Services (Safety) Act 2001 that is not owned or operated by a DHB.

"Quitcard Provider", means a person registered with the Ministry of Health as a Quitcard Provider.

"Residential Disability Care Institution", means premises used to provide residential disability care in accordance with the Health and Disability Services (Safety) Act 2001.

"Rest Home", means premises used to provide rest home care in accordance with the Health and Disability Services (Safety) Act 2001.

"Restricted Medicine", means any Pharmaceutical listed in Part II of Schedule 1 of the Medicines Regulations 1984.

"Retail Pharmacy-Specialist", means that the Community Pharmaceutical is only eligible for Subsidy if it is either:

- a) supplied on a Prescription or Practitioner's Supply Order signed by a Specialist, or,
- b) in the case of treatment recommended by a Specialist, supplied on a Prescription or Practitioner's Supply Order and either:
 - endorsed with the words "recommended by [name of Specialist and year of authorisation]" and signed by the Practitioner, or
 - ii) endorsed with the word 'protocol' which means "initiated in accordance with DHB hospital approved protocol", or
 - iii) Annotated by the dispensing pharmacist, following verbal confirmation from the Practitioner of the name of the Specialist and date of recommendation, with the words "recommended by [name of specialist and year of authorisation], confirmed by [practitioner]". Where the Contractor has an electronic record of such an Endorsement or Annotation from a previous prescription for the same Community Pharmaceutical written by a prescriber for the same patient, they may annotate the prescription accordingly.

"As recommended by a Specialist" to be interpreted as either:

- a) i) follows a substantive consultation with an appropriate Specialist;
 - ii) the consultation to relate to the Patient for whom the Prescription is written;
 - iii) consultation to mean communication by referral, telephone, letter, facsimile or email;
 - iv) except in emergencies consultation to precede annotation of the Prescription; and
 - v) both the Specialist and the General Practitioner must keep a written record of consultation; or
- b) treatment with the Community Pharmaceutical has been initiated in accordance with a DHB hospital approved protocol.

"Retail Pharmacy-Specialist Prescription", means that the Community Pharmaceutical is only eligible for Subsidy if it is supplied on a Prescription, or Practitioner's Supply Order, signed by a Specialist.

For the purposes of this definition, a "specialist" means a doctor who holds a current annual practicing certificate and who satisfies the criteria set out in paragraphs (a) or (b) or (c) of the definitions of Specialist below.

"Safety Medicine", means a Community Pharmaceutical defined in Section A, Part IV of the Pharmaceutical Schedule.

"Schedule", means this Pharmaceutical Schedule and all its sections and appendices.

"Special Authority", means that the Community Pharmaceutical or Pharmaceutical Cancer Treatment is only eligible for Subsidy or additional Subsidy for a particular person if an application meeting the criteria specified in the Schedule has been approved, and the valid Special Authority number is present on the prescription.

"Specialist", in relation to a Prescription, means a doctor who holds a current annual practising certificate and who satisfies the criteria set out in paragraphs (a) or (b) or (c) or (d) below:

- a) the doctor is vocationally registered in accordance with the criteria set out by the Medical Council of New Zealand and the HPCA Act 2003 and who has written the Prescription in the course of practising in that area of medicine; or
- b) the doctor is recognised by the Ministry of Health as a specialist for the purposes of this Schedule and receives remuneration from a DHB at a level which that DHB considers appropriate for specialists and who has written that prescription in the course of practising in that area of competency; or
- c) the doctor is recognised by the Ministry of Health as a specialist in relation to a particular area of medicine for the purpose of writing Prescriptions and who has written the Prescription in the course of practising in that area of competency; or
- d) the doctor writes the prescription on DHB stationery and is appropriately authorised by the relevant DHB to do so.

"Subsidy", means the maximum amount that the Government will pay Contractors for a Community Pharmaceutical dispensed to a person eligible for Pharmaceutical Benefits and is different from the cost to Government of subsidising that Community Pharmaceutical. For the purposes of a DHB hospital pharmacy claiming for Pharmaceutical Cancer Treatments, Subsidy refers to any payment made to the DHB hospital pharmacy or service provider to which that pharmacy serves, and does not relate to a specific payment that might be made on submission of a claim.

"Supply Order", means a Bulk Supply Order or a Practitioner's Supply Order.

"Unapproved Indication", means, for a Pharmaceutical, an indication for which it is not approved under the Medicines Act 1981. Practitioners prescribing Pharmaceuticals for Unapproved Indications should be aware of, and comply with, their obligations under Section 25 and/or Section 29 of the Medicines Act 1981 and as set out in Section A: General Rules, Part IV (Miscellaneous Provisions) rule 5.5.

"Unlisted Pharmaceutical", means a Pharmaceutical that is within the scope of a Hospital Pharmaceutical but is not listed in Section H part II

"Unusual Clinical Circumstances (UCC)", means the pathway under the Named Patient Pharmaceutical Assessment policy for funding consideration for named patients whose clinical circumstances are so unusual that PHARMAC is unlikely, for administrative reasons, to consider listing treatments for these circumstances on the Schedule.

"Urgent Assessment (UA)", means the pathway under the Named Patient Pharmaceutical Assessment policy for funding consideration for treatments for named patients where PHARMAC is also considering or is likely to consider the treatment for Schedule listing, but the patient's clinical circumstances justify urgent assessment, prior to a decision on Schedule listing.

- 1.2 In addition to the above interpretations and definitions, unless the content requires otherwise, a reference in the Schedule to:
 - a) the singular includes the plural; and
 - b) any legislation includes a modification and re-enactment of, legislation enacted in substitution for, and a regulation, Order in Council, and other instrument from time to time issued or made under that legislation, where that legislation, regulation, Order in Council or other instrument has an effect on the prescribing, dispensing or subsidising of Community Pharmaceuticals.

PART II COMMUNITY PHARMACEUTICALS SUBSIDY

- 2.1 Community Pharmaceuticals eligible for Subsidy include every medicine, therapeutic medical device or related product, or related thing listed in Sections B to G and I of the Schedule subject to:
 - 2.1.1 clauses 2.2 of the Schedule; and
 - 2.1.2 clauses 3.1 to 5.4 of the Schedule; and
 - 2.1.3 the conditions (if any) specified in Sections B to G and I of the Schedule;
- 2.2 No claim by a Contractor for payment in respect of the supply of Community Pharmaceuticals will be allowed unless the Community Pharmaceuticals so supplied:
 - 2.2.1 comply with the appropriate standards prescribed by regulations for the time being in force under the Medicines Act 1981; or
 - 2.2.2 in the absence of any such standards, comply with the appropriate standards for the time being prescribed by the British Pharmacopoeia; or
 - 2.2.3 in the absence of the standards prescribed in clauses 2.2.1 and 2.2.2, comply with the appropriate standards for the time being prescribed by the British Pharmaceutical Codex; or
 - 2.2.4 in the absence of the standards prescribed in clauses 2.2.1, 2.2.2 and 2.2.3 are of a grade and quality not lower than those usually applicable to Community Pharmaceuticals intended to be used for medical purposes.

PART III PERIOD AND QUANTITY OF SUPPLY

3.1 Doctors', Dentists', Dietitians', Midwives', Nurse Prescribers', Optometrists and Pharmacist Prescribers' Prescriptions (other than oral contraceptives) The following provisions apply to all Prescriptions, other than those for an oral contraceptive, written by a Doctor, Dentist, Dietitian, Midwife, Nurse Prescriber, an Optometrist, or a Pharmacist Prescriber unless specifically excluded:

- 3.1.1 For a Community Pharmaceutical other than a Class B Controlled Drug, only a quantity suffcient to provide treatment for a period not exceeding three Months will be subsidised.
- 3.1.2 For methylphenidate hydrochloride and dexamphetamine sulphate (except for Dentist prescriptions), only a quantity sufficient to provide treatment for a period not exceeding one Month will be subsidised.
- 3.1.3 For a Class B Controlled Drug:
 - a) other than Dentist prescriptions and methylphenidate hydrochloride and dexamphetamine sulphate, only a quantity:
 - i) sufficient to provide treatment for a period not exceeding 10 days; and
 - ii) which has been dispensed pursuant to a Prescription sufficient to provide treatment for a period not exceeding one Month, will be subsidised.
 - b) for a Dentist prescription only such quantity as is necessary to provide treatment for a period not exceeding five days will be subsidised.
- 3.1.4 Subject to clauses 3.1.3 and 3.1.7, for a Doctor, Dentist, Dietitian, Midwife or Nurse Prescriber and 3.1.7 for an Optometrist, where a practitioner has prescribed a quantity of a Community Pharmaceutical sufficient to provide treatment for:
 - A) one Month or less than one Month, but dispensed by the Contractor in quantities smaller than the quantity prescribed, the Community Pharmaceutical will only be subsidised as if that Community Pharmaceutical had been dispensed in a Monthly Lot;
 - B) more than one Month, the Community Pharmaceutical will be subsidised only if it is dispensed:
 - i) in a 90 Day Lot, where the Community Pharmaceutical is a Pharmaceutical covered by Section F Part I of the Pharmaceutical Schedule; or
 - ii) if the Community Pharmaceutical is not a Pharmaceutical referred to in Section F Part I of the Pharmaceutical Schedule, in Monthly Lots, unless:
 - a) the eligible person or his/her nominated representative endorses the back of the Prescription form with a statement identifying which Access Exemption Criterion (Criteria) applies and signs that statement to this effect; or
 - b) both:
 - the Practitioner endorses the Community Pharmaceutical on the Prescription with the words "certified exemption" written in the Practitioner's own handwriting, or signed or initialled by the Practitioner; and
 - every Community Pharmaceutical endorsed as "certified exemption" is covered by Section F Part II of the Pharmaceutical Schedule.
- 3.1.5 A Community Pharmaceutical is only eligible for Subsidy if the Prescription under which it has been dispensed was presented to the Contractor:
 - a) for a Class B Controlled Drug, within eight days of the date on which the Prescription was written; or
 - b) for any other Community Pharmaceutical, within three Months of the date on which the Prescription was written.
- 3.1.6 No subsidy will be paid for any Prescription, or part thereof, that is not fulfilled within:
 - a) in the case of a Prescription for a total supply of from one to three Months, three Months from the date the Community Pharmaceutical was first dispensed; or
 - b) in any other case, one Month from the date the Community Pharmaceutical was first dispensed. Only that part of any Prescription that is dispensed within the time frames specified above is eligible for Subsidy.
- 3.1.7 If a Community Pharmaceutical:
 - a) is stable for a limited period only, and the Practitioner has endorsed the Prescription with the words "unstable medicine" and has specified the maximum quantity that may be dispensed at any one time; or
 - b) is stable for a limited period only, and the Contractor has endorsed the Prescription with the words "unstable medicine" and has specified the maximum quantity that should be dispensed at any one time in all the circumstances of the particular case; or
 - c) is under the Dispensing Frequency Rule,

The actual quantity dispensed will be subsidised in accordance with any such specification.

3.2 Oral Contraceptives

The following provisions apply to all Prescriptions written by a Doctor, Midwife, Nurse Prescriber or a Pharmacist Prescriber for an oral contraceptive:

- 3.2.1 The prescribing Doctor, Midwife, Nurse Prescriber or a Pharmacist Prescriber must specify on the Prescription the period of treatment for which the Community Pharmaceutical is to be supplied. This period must not exceed six Months.
- 3.2.2 Where the period of treatment specified in the Prescription does not exceed six Months, the Community Pharmaceutical is to be dispensed:
 - a) in Lots as specified in the Prescription if the Community Pharmaceutical is under the Dispensing Frequency Rule; or
 - b) where no Lots are specified, in one Lot sufficient to provide treatment for the period prescribed.
- 3.2.3 An oral contraceptive is only eligible for Subsidy if the Prescription under which it has been dispensed was presented to the Contractor within three Months of the date on which it was written.
- 3.2.4 Where a Community Pharmaceutical on a Prescription is under the Dispensing Frequency Rule and a repeat on the Prescription remains unfulfilled after six Months from the date the Community Pharmaceutical was first dispensed only the actual quantity supplied by the Contractor within this time limit will be eligible for Subsidy.

3.3 Original Packs, Certain Antibiotics and Unapproved Medicines

- 3.3.1 Notwithstanding clauses 3.1 and 3.3 of the Schedule, if a Practitioner prescribes or orders a Community Pharmaceutical that is identified as an Original Pack (OP) on the Pharmaceutical Schedule and is packed in a container from which it is not practicable to dispense lesser amounts, every reference in those clauses to an amount or quantity eligible for Subsidy, is deemed to be a reference:
 - a) where an amount by weight or volume of the Community Pharmaceutical is specified in the Prescription, to the smallest container of the Community Pharmaceutical, or the smallest number of containers of the Community Pharmaceutical, sufficient to provide that amount; and
 - b) in every other case, to the amount contained in the smallest container of the Community Pharmaceutical that is manufactured in, or imported into, New Zealand.
- 3.3.2 If a Community Pharmaceutical is either:
 - a) the liquid oral form of an antibiotic to which a diluent must be added by the Contractor at the time of dispensing; or
 - b) an unapproved medicine supplied under Section 29 of the Medicines Act 1981, but excluding any medicine listed as Cost, Brand, Source of Supply, or
 - c) any other pharmaceutical that PHARMAC determines, from time to time and notes in the Pharmaceutical Schedule

and it is prescribed or ordered by a Practitioner in an amount that does not coincide with the amount contained in one or more standard packs of that Community Pharmaceutical, Subsidy will be paid for the amount prescribed or ordered by the Practitioner in accordance with either clause 3.1 or clause 3.3 of the Schedule, and for the balance of any pack or packs from which the Community Pharmaceutical has been dispensed. At the time of dispensing the Contractor must keep a record of the quantity discarded. To ensure wastage is reduced, the Contractor should reduce the amount dispensed to make it equal to the quantity contained in a whole pack where:

- a) the difference between the amount dispensed and the amount prescribed by the Practitioner is less than 10% (eg; if a prescription is for 105 mls then a 100ml pack would be dispensed); and
- b) in the reasonable opinion of the Contractor the difference would not affect the efficacy of the course of treatment prescribed by the Practitioner.

Note: For the purposes of audit and compliance it is an act of fraud to claim wastage and then use the wastage amount for any subsequent prescription.

3.4 Pharmacist Prescribers' Prescriptions

The following apply to every prescription written by a Pharmacist Prescriber

- 3.4.1 Prescriptions written by a Pharmacist Prescriber for a Community Pharmaceutical will only be subsidised where they are for either:
 - a) a Community Pharmaceutical classified as a Prescription Medicine and which a Pharmacist Prescriber is permitted under regulations to prescribe; or

- b) any other Community Pharmaceutical that is a Restricted Medicine (Pharmacist Only Medicine), a Pharmacy Only Medicine or a General Sales Medicine.
- 3.4.2 Any Pharmacist Prescribers' prescriptions for a medication requiring a Special Authority will only be subsidised if it is for a repeat prescription (ie after the initial prescription with Special Authority approval was dispensed).

3.5 Dietitians' Prescriptions

- The following provisions apply to every Prescription written by a Dietitian:
- 3.5.1 Prescriptions written by a Dietitian for a Community Pharmaceutical will only be subsidised where they are for either:
 - a) special foods, as listed in Section D; or
 - b) any other Pharmaceutical that has been identified in Section D of the Pharmaceutical Schedule as being able to be prescribed by a Dietitian, providing that the products being prescribed are not classified as Prescription Medicines or Restricted Medicines.
- 3.5.2 For the purposes of Dietitians prescribing pursuant to this clause 3.5, the prescribing and dispensing of these products is required to be in accordance with regulations 41 and 42 of the Medicines Regulations 1984.

3.6 Diabetes Nurse Prescribers' Prescriptions

The following provisions apply to every Prescription written by a Diabetes Nurse Prescriber:

- 3.6.1 Prescriptions written by a Diabetes Nurse Prescriber for a Community Pharmaceutical will only be subsidised where they are for either:
 - a) a Community Pharmaceutical classified as a Prescription Medicine or a Restricted Medicine and which a Diabetes Nurse Prescribers is permitted under regulations to prescribe; or
 - b) any other Community Pharmaceutical listed below:

aspirin, blood glucose diagnostic test meter, blood glucose diagnostic test strip, blood ketone diagnostic test meter, glucagon hydrochloride inj 1 mg syringe kit, insulin pen needles, insulin syringes disposable with attached needle, insulin pump accessories, insulin pump infusion set, insulin pump reservoir, ketone blood beta-ketone electrodes test strip, nicotine, sodium nitroprusside test strip,

3.6.2 Any Diabetes Nurse Prescribers' prescription for a medication requiring a Special Authority will only be subsidised if it is for a repeat prescription (ie after the initial prescription with Special Authority approval was dispensed).

3.7 Quitcard Providers' Prescriptions

Prescriptions written by a Quitcard Provider will only be subsidised where they are:

- a) for any of the following Community Pharmaceuticals: nicotine patches, nicotine lozenges or nicotine gum; and
- b) written on a Quitcard.

PART IV DISPENSING FREQUENCY RULE

Rule 3.1.4 of the Pharmaceutical Schedule specifies, for community patients, a default period of supply for each Community Pharmaceutical (a Monthly Lot, 90 Day Lot or for oral contraceptives 180 Day Lot). This Dispensing Frequency Rule defines patient groups or medicines eligible for more frequent dispensing periods for Community Pharmaceuticals; and the conditions that must be met to enable any pharmacy to claim for payment of handling fees for the additional dispensings made. This Dispensing Frequency Rule relates to the circumstances in which a subsidy is payable for the Community Pharmaceutical; it does not override alternative dispensing frequencies as expressly stated in the Medicines Act, Medicines Regulations, Pharmacy Services Agreement or Pharmaceutical Schedule.

For the purposes of this Dispensing Frequency Rule:

"Frequent Dispensing" means:

- i) for a Community Pharmaceutical referred to in Section F Part I, (the Stat exemption) dispensing in quantities less than one 90 Day Lot (or for oral contraceptives, less than one 180 Day Lot); or
- ii) for any other Community Pharmaceutical dispensing in quantities less than a Monthly Lot "Safety Medicine"
 - i) an antidepressant listed under the "Cyclic and Related Agents" subheading;
 - ii) an antipsychotic;

- iii) a benzodiazepine;
- iv) a Class B Controlled Drug;
- v) codeine (includes combination products);
- vi) buprenorphine with naloxone; or
- vii) zopiclone.

The Dispensing Frequency Rule covers 5 different circumstances where Frequent Dispensing for patients may be clinically or otherwise appropriate. These are:

- 1) Long Term Condition (LTC) patients and Core patients, or
- 2) Persons in residential care, or
- 3) Trial periods, or
- 4) Safety and co-prescribed medicines, or
- 5) Pharmaceutical Supply Management.

4.1 Frequent Dispensing for patients registered as Long Term Condition (LTC) or Core patients

If a Pharmacist considers Frequent Dispensing is required, then:

- 4.1.1 For LTC registered patients, Frequent Dispensing can occur as often as the dispensing Pharmacist deems appropriate to meet that patient's compliance and adherence needs;
- 4.1.2 For Core (non-LTC) patients, Frequent Dispensing should be no more often than a Monthly Lot. Pharmacists may authorise monthly dispensing on a Stat exemption Community Pharmaceutical without prescriber authority. If the Pharmacist considers more frequent (than monthly) dispensing is necessary, prescriber approval is required. Verbal approval from the prescriber is acceptable provided it is annotated by the Pharmacist on the Prescription and dated.

4.2 Frequent Dispensings for persons in residential care

- 4.2.1 Community Pharmaceuticals can be dispensed to:
 - any person whose placement in a Residential Disability Care Institution is funded by the Ministry of Health or a DHB; or
 - a person assessed as requiring long term residential care services and residing in an age related residential care facility;

on the request of the person, their agent or caregiver or community residential service provider via Frequent Dispensing, provided the following conditions are met:

- a) the quantity or period of supply to be dispensed at any one time is not less than:
 - i) 7 days' supply for a Class B Controlled Drug; or
 - ii) 7 days' supply for clozapine in accordance with a Clozapine Dispensing Protocol; or
 - iii) 28 days' supply for any other Community Pharmaceutical (except under conditions outlined in 4.3 (Trial periods) below; and
- b) the prescribing Practitioner or dispensing Pharmacist has
 - i) included the name of the patient's residential placement or facility on the Prescription; and
 - ii) included the patient's NHI number on the Prescription; and
 - iii) specified the maximum quantity or period of supply to be dispensed at any one time.
- 4.2.2 Any person meeting the criteria above who is being initiated onto a new medicine or having their dose changed is able to have their medicine dispensed in accordance with 4.3 (Trial periods) below.

4.3 Frequent Dispensings for Trial Periods

Frequent Dispensing can occur when a Community Pharmaceutical has been prescribed for a patient who requires close monitoring due to recent initiation onto, or dose change for, the Community Pharmaceutical (applicable to the patient's first changed Prescription only) and the prescribing Practitioner has:

- endorsed each Community Pharmaceutical on the Prescription clearly with the words "Trial Period", or "Trial"; and
- specified the maximum quantity or period of supply to be dispensed for each Community Pharmaceutical at any one time.

Patients who reside in Penal Institutions are not eligible for Trial Periods.

4.4 Frequent Dispensing for Safety and co-prescribed medicines

- 4.4.1 For a Safety Medicine to be dispensed via Frequent Dispensing, both of the following conditions must be met:
 - a) The patient is not a resident in a Penal Institution, or one of the residential placements or facilities referenced in 4.2 on the previous page; and
 - b) The prescribing Practitioner has:
 - i) Assessed clinical risk and determined the patient requires increased Frequent Dispensing; and
 - ii) Specified the maximum quantity or period of supply to be dispensed for each Safety Medicine at each dispensing.
- 4.4.2 A Community Pharmaceutical that is co-prescribed with a Safety Medicine, which can be dispensed in accordance with rule 4.4.1 above, may be dispensed at the same frequency as the Safety Medicine if the dispensing pharmacist has:
 - Assessed clinical risk and determined the patient requires Frequent Dispensing of their co-dispensed medicines; and
 - Annotated the Prescription with the amended dispensing quantity and frequency.

4.5 Frequent Dispensing for Pharmaceutical Supply Management

- 4.5.1 Frequent Dispensing may be required from time to time to manage stock supply issues or emergency situations. Pharmacists may dispense more frequently than the Schedule would otherwise allow when all of the following conditions are met:
 - a) PHARMAC has approved and notified pharmacists to annotate Prescriptions for a specified Community Pharmaceutical(s) "out of stock" without prescriber endorsement for a specified time; and
 - b) the dispensing pharmacist has:
 - clearly annotated each of the approved Community Pharmaceuticals that appear on the Prescription with the words "out of stock" or "OOS"; and
 - ii) initialled the annotation in their own handwriting; and
 - iii) has complied with maximum quantity or period of supply to be dispensed at any one time, as specified by PHARMAC at the time of notification.

Note – no claim shall be made to any DHB for subsidised dispensing under this rule where dispensing occurs more frequently than specified by PHARMAC to manage the supply management issue.

PART V MISCELLANEOUS PROVISIONS

5.1 Bulk Supply Orders

The following provisions apply to the supply of Community Pharmaceuticals under Bulk Supply Orders:

- 5.1.1 No Community Pharmaceutical supplied under a Bulk Supply Order will be subsidised unless all the requirements in Section B, C or D of the Schedule applicable to that pharmaceutical are met.
- 5.1.2 The person who placed the Bulk Supply Order may be called upon by the Ministry of Health to justify the amount ordered.
- 5.1.3 Class B Controlled Drugs will be subsidised only if supplied under Bulk Supply Orders placed by an institution certified to provide hospital care under the Health and Disability Services (Safety) Act 2001.
- 5.1.4 Any order for a Class B Controlled Drug or for buprenorphine hydrochloride must be written on a Special Bulk Supply Order Controlled Drug Form supplied by the Ministry of Health.
- 5.1.5 Community Pharmaceuticals listed in Part I of the First Schedule to the Medicines Regulations 1984 will be subsidised only if supplied under a Bulk Supply Order placed by an institution certified to provide hospital care under the Health and Disability Services (Safety) Act 2001 and:
 - a) that institution employs a registered general nurse, registered with the Nursing Council and who holds a current annual practicing certificate under the HPCA Act 2003; and
 - b) the Bulk Supply Order is supported by a written requisition signed by a Hospital Care Operator.
- 5.1.6 No Subsidy will be paid for any quantity of a Community Pharmaceutical supplied under a Bulk Supply Order in excess of what is a reasonable monthly allocation for the particular institution, after taking into account stock on hand.
- 5.1.7 The Ministry of Health may, at any time, by public notification, declare that any approved institution within its particular region, is not entitled to obtain supplies of Community Pharmaceuticals under Bulk Supply Orders

with effect from the date specified in that declaration. Any such notice may in like manner be revoked by the Ministry of Health at any time.

5.2 Practitioner's Supply Orders

The following provisions apply to the supply of Community Pharmaceuticals to Practitioners under a Practitioner's Supply Order:

- 5.2.1 Subject to clause 5.2.3 and 5.2.6, a Practitioner may only order under a Practitioner's Supply Order those Community Pharmaceuticals listed in Section E Part I and only in such quantities as set out in Section E Part I that the Practitioner requires to ensure medical supplies are available for emergency use, teaching and demonstration purposes, and for provision to certain patient groups where individual prescription is not practicable.
- 5.2.2 Any order for a Class B Controlled Drug or for buprenorphine hydrochloride must be written on a Special Practitioner's Supply Order Controlled Drug Form supplied by the Ministry of Health.
- 5.2.3 A Practitioner may order such Community Pharmaceuticals as he or she expects to be required for personal administration to patients under the Practitioner's care if:
 - a) the Practitioner's normal practice is in the specified areas listed in Section E Part II of the Schedule, or if the Practitioner is a locum for a Practitioner whose normal practice is in such an area.
 - b) the quantities ordered are reasonable for up to one Month's supply under the conditions normally existing in the practice. (The Practitioner may be called on by the Ministry of Health to justify the amounts of Community Pharmaceuticals ordered.)
- 5.2.4 No Community Pharmaceutical ordered under a Practitioner's Supply order will be eligible for Subsidy unless:
 - a) the Practitioner's Supply Order is made on a form supplied for that purpose by the Ministry of Health, or approved by the Ministry of Health and which:
 - i) is personally signed and dated by the Practitioner; and
 - ii) sets out the Practitioner's address; and
 - iii) sets out the Community Pharmaceuticals and quantities, and;
 - b) all the requirements of Sections B and C of the Schedule applicable to that pharmaceutical are met.
- 5.2.5 The Ministry of Health may, at any time, on the recommendation of an Advisory Committee appointed by the Ministry of Health for that purpose, by public notification, declare that a Practitioner specified in such a notice is not entitled to obtain supplies of Community Pharmaceuticals under Practitioner's Supply Orders until such time as the Ministry of Health notifies otherwise.
- 5.2.6 A Practitioner working in the Rheumatic Fever Prevention Programme (RFPP) may order under a Practitioner's Supply Order such Community Pharmaceuticals (identified below) as he or she requires to ensure medical supplies are available for patients with suspected or confirmed Group A Streptococcal throat infections for the purposes of the RFPP in the following circumstances:
 - a) the RFPP provider name is written on the Practitioner's Supply Order; and
 - b) the total quantity ordered does not exceed a multiple of:
 - ten times the Practitioner's Supply Order current maximum listed in Section E Part I for amoxicillin grans for oral liq 250 mg per 5 ml, amoxicillin cap 250 mg and amoxicillin cap 500 mg; or
 - ii) two times the Practitioner's Supply Order current maximum listed in Section E Part I for phenoxymethyl penicillin grans for oral liquid 250 mg per 5 ml, phenoxymethyl penicillin cap 500 mg, erythromycin ethyl succinate grans for oral liq 200 mg per 5 ml and erythromycin ethyl succinate tab 400 mg; and
 - c) the practitioner must specify the order quantity in course-specific amounts on the Practitioner's Supply Order (e.g. 10 x 300 ml amoxicillin grans for oral liq 250 mg per 5 ml). This will enable the pharmacy to dispense each course separately and claim multiple service fees as per the Community Pharmacy Services Agreement.

5.3 Retail Pharmacy and Hospital Pharmacy-Specialist Restriction

The following provisions apply to Prescriptions for Community Pharmaceuticals eligible to be subsidised as "Retail Pharmacy-Specialist" and "Hospital Pharmacy-Specialist":

5.3.1 Record Keeping

It is expected that a record will be kept by both the General Practitioner and the Specialist of the fact of consultation and enough of the clinical details to justify the recommendation. This means referral by telephone will need to be followed up by written consultation.

5.3.2 Expiry

The recommendation expires at the end of two years and can be renewed by a further consultation.

- 5.3.3 The circulation by Specialists of the circumstances under which they are prepared to recommend a particular Community Pharmaceutical is acceptable as a guide. It must however be followed up by the procedure in subclauses 5.3.1 and 5.3.2, for the individual Patient.
- 5.3.4 The use of preprinted forms and named lists of Specialists (as circulated by some pharmaceutical companies) is regarded as inappropriate.
- 5.3.5 The Rules for Retail Pharmacy-Specialist and Hospital Pharmacy-Specialist will be audited as part of the Ministry of Health's routine auditing procedures.

5.4 Pharmaceutical Cancer Treatments

- 5.4.1 DHBs must provide access to Pharmaceutical Cancer Treatments for the treatment of cancers in their DHB hospitals, and/or in association with Outpatient services provided in their DHB hospitals.
- 5.4.2 DHBs must only provide access to Pharmaceuticals for the treatment of cancer that are listed as Pharmaceutical Cancer Treatments in Sections A to G of the Schedule, provided that DHBs may provide access to an unlisted pharmaceutical for the treatment of cancer where that unlisted pharmaceutical:
 - a) has Named Patient Pharmaceutical Assessment (NPPA) approval;
 - b) is being used as part of a bona fide clinical trial which has Ethics Committee approval;
 - c) is being used and funded as part of a paediatric oncology service; or
 - d) was being used to treat the patient in question prior to 1 July 2005.
- 5.4.3 A DHB hospital pharmacy that holds a claiming agreement for Pharmaceutical Cancer Treatements with the Funder may claim a Subsidy for a Pharmaceutical Cancer Treatment marked as "PCT" or "PCT only" in Sections A to G of this Schedule subject to that Pharmaceutical Cancer Treatment being dispensed in accordance with:
 - a) Part 1;
 - b) clauses 2.1 to 2.2;
 - c) clauses 3.1 to 3.4; and
 - d) clause 5.4,
 - of Section A of the Schedule
- 5.4.4 A Contractor (other than a DHB hospital pharmacy) may only claim a Subsidy for a Pharmaceutical Cancer Treatment marked as "PCT" in Sections A to G of the Schedule subject to that Pharmaceutical Cancer Treatment being dispensed in accordance with the rules applying to Sections A to G of the Schedule.
- 5.4.5 Some indications for Pharmaceutical Cancer Treatments listed in the Schedule are Unapproved Indications. Some of these formed part of the October 2001 decision by the Minister of Health as to pharmaceuticals and indications for which DHBs must provide access. As far as reasonably practicable, these Unapproved Indications are marked in the Schedule. However, PHARMAC makes no representation and gives no guarantee as to the accuracy of this information. Practitioners prescribing Pharmaceutical Cancer Treatments for such Unapproved Indications should:
 - a) be aware of and comply with their obligations under sections 25 and 29 of the Medicines Act 1981, as applicable, and otherwise under that act and the Medicines Regulations 1984;
 - b) be aware of and comply with their obligations under the Health and Disability Commissioner's Code of Consumer Rights, including the requirement to obtain informed consent from the patient (PHARMAC recommends that Practitioners obtain written consent); and
 - c) exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of an unapproved Pharmaceutical Cancer Treatment or a Pharmaceutical Cancer Treatment for an Unapproved Indication.
- 5.4.6 Applications to add pharmaceuticals, and add or amend indications for Pharmaceutical Cancer Treatments, may be made in writing by pharmaceutical suppliers and/or clinicians to PHARMAC. Applications should follow the Guidelines for Funding Applications to PHARMAC 2010 and Recommended methods to derive clinical inputs for proposals to PHARMAC, copies of which are available from PHARMAC or PHARMAC's website.

5.5 Practitioners prescribing unapproved Pharmaceuticals

Practitioners should, where possible, prescribe Pharmaceuticals that are approved under the Medicines Act 1981. However, the access criteria under which a Pharmaceutical is listed on the Pharmaceutical Schedule may:

a) in some case, explicitly permit Government funded access to a Pharmaceutical that is not approved under

the Medicines Act 1981 or for an Unapproved Indication; or

b) not explicitly preclude Government funded access to a Pharmaceutical when it is used for an Unapproved Indication;

Accordingly, if Practitioners are planning on prescribing an unapproved Pharmaceutical or a Pharmaceutical for an Unapproved Indication, Practitioners should:

- a) be aware of and comply with their obligations under sections 25 and 29 of the Medicines Act 1981, as applicable, and otherwise under that Act and the Medicines Regulations 1984;
- b) be aware of and comply with their obligations under the Health and Disability Commissioner's Code of Consumer Rights, including the requirement to obtain informed consent from the patient (PHARMAC recommends that Practitioners obtain written consent); and
- c) exercise their own skill, judgment, expertise and discretion, and make their own prescribing decisions with respect to the use of an unapproved Pharmaceutical or a Pharmaceutical for an Unapproved Indication.

Practitioners should be aware that simply by listing a Pharmaceutical on the Pharmaceutical Schedule PHARMAC makes no representations about whether that Pharmaceutical has any form of approval or consent under, or whether the supply or use of the Pharmaceutical otherwise complies with, the Medicines Act 1981. Further, the Pharmaceutical Schedule does not constitute an advertisement, advertising material or a medical advertisement as defined in the Medicines Act or otherwise.

5.6 Substitution

Where a Practitioner has prescribed a brand of a Community Pharmaceutical that has no Subsidy or has a Manufacturer's Price that is greater than the Subsidy and there is an alternative fully subsidised Community Pharmaceutical available, a Contractor may dispense the fully subsidised Community Pharmaceutical, unless either or both of the following circumstances apply:

- a) there is a clinical reason why substitution should not occur; or
- b) the prescriber has marked the prescription with a statement such as 'no brand substitution permitted'

Such an Authority to Substitute is valid whether or not there is a financial implication for the Pharmaceutical Budget. When dispensing a subsidised alternative brand, the Contractor must annotate and sign the prescription and inform the patient of the brand change.

5.7 Alteration to Presentation of Pharmaceutical Dispensed

A Contractor, when dispensing a subsidised Community Pharmaceutical, may alter the presentation of a Pharmaceutical dispensed to another subsidised presentation but may not alter the dose, frequency and/or total daily dose. This may only occur when it is not practicable for the contractor to dispense the requested presentation. If the change will result in additional cost to the DHBs, then annotation of the prescription by the dispensing pharmacist must occur stating the reason for the change, and the Contractor must initial the change for the purposes of Audit.

5.8 Other DHB Funding

A DHB may fund a Community Pharmaceutical outside of the mechanisms established in the Pharmaceutical Schedule, provided that:

- a) specific prior agreement is obtained from PHARMAC for such funding;
- any funding restrictions set out in the Pharmaceutical Schedule for those Community Pharmaceuticals are applied; and
- c) a Contractor (including a DHB Hospital Pharmacy) may not claim a Subsidy for a Community Pharmaceutical dispensed and funded by the DHB via such an alternate mechanism.

5.9 Conflict in Provisions

If any rules in Sections B-G and Section I of this Schedule conflict with the rules in Section A, the rules in Sections B-G and Section I apply.

SECTION B: ALIMENTARY TRACT AND METABOLISM

| | Subsidy | | Fully | Brand or |
|--|-----------------------------|--------------------|------------|------------------------------------|
| | (Manufacturer's Price \$ | e) Per | Subsidised | |
| Antacids and Antiflatulants | | | | |
| Antacids and Reflux Barrier Agents | | | | |
| ALGINIC ACID | | | | |
| Sodium alginate 225 mg and magnesium alginate 87.5 mg per sachet | | 30 | • | Gaviscon Infant |
| SIMETHICONE | | | | |
| Oral liq aluminium hydroxide 200 mg with magnesium hydrox- ide 200 mg and activated simethicone 20 mg per 5 ml | | 500 ml | ſ | Mylanta P |
| SODIUM ALGINATE | | | | |
| * Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour | | 60 | (| Gaviscon Double Strength |
| Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml | | 500 ml | | Acidex |
| Phosphate Binding Agents | | | | |
| ALUMINIUM HYDROXIDE * Tab 600 mg | | 100 | ~ | Alu-Tab |
| CALCIUM CARBONATE | | | | |
| Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsement Only when prescribed for children under 12 years of ag endorsed accordingly. Antidiarrhoeals | | 500 ml sphate I | | Roxane ent and the prescription |
| Agents Which Reduce Motility | | | | |
| LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a | a PSO | | | |
| * Tab 2 mg * Cap 2 mg | | 400 400 | | Nodia Diamide Relief |
| Rectal and Colonic Anti-inflammatories | | | | |
| BUDESONIDE Cap 3 mg – Special Authority see SA1155 below – Retail pharmacy | | 90 | ~ 1 | Entocort CIR |
| SA1155 Special Authority for Subsidy nitial application — (Crohn's disease) from any relevant praction following criteria: Both: | | alid for (| 6 months f | or applications meeting th |
| Mild to moderate ileal, ileocaecal or proximal Crohn's dis Any of the following: | ease; and | | | |

20

continued...

| Subsidy (Manufacturer's Price) | | Fully Subsidised | Brand or Generic | |
|-----------------------------------|-----|---------------------|---------------------|--|
| (Manuacurer's Price) | | Subsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

- 2.1 Diabetes; or
- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or
- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an Unapproved Indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

| Rectal foam 10%, CFC-Free (14 applications) | 21.1 g OP | Colifoam |
|--|-----------|------------------------------|
| MESALAZINE | | |
| Tab 400 mg49.50 | 100 | Asacol |
| Tab EC 500 mg | 100 | Asamax |
| Tab long-acting 500 mg59.05 | 100 | Pentasa |
| Modified release granules, 1 g141.72 | 120 OP | Pentasa |
| Enema 1 g per 100 ml41.30 | 7 | Pentasa |
| Pentasa to be Sole Supply on 1 October 2015 | | |
| Suppos 500 mg | 20 | Asacol |
| Suppos 1 g | 30 | Pentasa |
| OLSALAZINE | | |
| Tab 500 mg | 100 | Dipentum |
| Cap 250 mg | 100 | Dipentum |
| SODIUM CROMOGLYCATE | | |
| | 100 | |
| Cap 100 mg89.21 | 100 | Nalcrom |
| SULPHASALAZINE | | |
| Tab 500 mg – For sulphasalazine oral liquid formulation refer, | | |
| page 207 | 100 | Salazopyrin |
| * Tab EC 500 mg12.89 | 100 | Salazopyrin EN |

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cin-

| chocaine hydrochloride 5 mg per g6.35 | 30 g OP | Ultraproct |
|--|---------|--------------------------------|
| Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and | | |
| cinchocaine hydrochloride 1 mg2.66 | 12 | Ultraproct |

| | Subsidy (Manufacturer's Pr \$ | ice) Sul Per | Fully Brand or osidised Generic Manufacturer |
|--|-------------------------------------|-----------------|---|
| HYDROCORTISONE WITH CINCHOCAINE Oint 5 mg with cinchocaine hydrochloride 5 mg per g Suppos 5 mg with cinchocaine hydrochloride 5 mg per g | | 30 g OP 12 | ✓ Proctosedyl ✓ Proctosedyl |
| Management of Anal Fissures | | | |
| GLYCERYL TRINITRATE – Special Authority see SA1329 below * Oint 0.2% SA1329 Special Authority for Subsidy | | 30 g OP | ✓ Rectogesic |
| Initial application from any relevant practitioner. Approvals vali chronic anal fissure that has persisted for longer than three week | | enewal unles | is notified where the patient has |
| Antispasmodics and Other Agents Altering Gut | Motility | | |
| GLYCOPYRRONIUM BROMIDE | | | |
| Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available or a PSO | | 10 | 🗸 Max Health |
| HYOSCINE N-BUTYLBROMIDE | | | - |
| * Tab 10 mg * Inj 20 mg, 1 ml – Up to 5 inj available on a PSO | | 20 5 | Gastrosoothe Buscopan |
| MEBEVERINE HYDROCHLORIDE | 9.57 | 5 | |
| * Tab 135 mg | | 90 | ✓ <u>Colofac</u> |
| Antiulcerants | | | |
| Antisecretory and Cytoprotective | | | |
| MISOPROSTOL * Tab 200 mcg | | 120 | ✔ Cytotec |
| Helicobacter Pylori Eradication | | | |
| CLARITHROMYCIN Tab 500 mg – Subsidy by endorsement a) Maximum of 14 tab per prescription b) Subsidiated ashuf prescripted for ballachester pulari are | | 14 | ✓ <u>Apo-Clarithromycin</u> |
| b) Subsidised only if prescribed for helicobacter pylori era Note: the prescription is considered endorsed if clarithromycin is amoxicillin or metronidazole. | | | |
| H2 Antagonists | | | |
| CIMETIDINE – Only on a prescription * Tab 200 mg | 5.00 (7.50) | 100 | Apo-Cimetidine |
| * Tab 400 mg | | 100 | |
| (Apo-Cimetidine Tab 200 mg to be delisted 1 February 2016) (Apo-Cimetidine Tab 400 mg to be delisted 1 February 2016) | (12.00) | | Apo-Cimetidine |
| RANITIDINE – Only on a prescription | | | 4 - - - - - |
| * Tab 150 mg * Tab 300 mg | | 500 500 | ✓ <u>Ranitidine Relief</u> ✓ Ranitidine Relief |
| • | | 300 ml | ✓ Peptisoothe |
| * Oral liq 150 mg per 10 ml | | 000 111 | Zantac |

| | Subsidy (Manufacturer's Price) \$ | Per | Full <u>y</u> Subsidised | |
|--|---|-----|-----------------------------|---------------|
| Proton Pump Inhibitors | | | | |
| LANSOPRAZOLE | | | | |
| * Cap 15 mg | 2.00 | 28 | | Solox |
| * Cap 30 mg | 2.32 | 28 | ~ | Solox |
| OMEPRAZOLE | | | | |
| For omeprazole suspension refer Standard Formulae, page 21 | 0 | | | |
| * Cap 10 mg | | 90 | | Omezol Relief |
| * Cap 20 mg | | 90 | | Omezol Relief |
| * Cap 40 mg | | 90 | | Omezol Relief |
| Powder – Only in combination | | 5 g | V | Midwest |
| Only in extemporaneously compounded omeprazole susper * Ini 40 mg | | 5 | | Dr Reddy's |
| ₭ Inj 40 mg | | 5 | v | Omeprazole |
| | | | | Omepiazole |
| PANTOPRAZOLE * Tab EC 20 mg | 2.69 | 100 | | Pantoprazole |
| * Tab EC 20 Trig | 2.00 | 100 | • | Actavis 20 |
| * Tab EC 40 mg | | 100 | ~ | Pantoprazole |
| | | | | Actavis 40 |
| Site Protective Agents | | | | |
| BISMUTH TRIOXIDE | | | | |
| Tab 120 mg | 32 50 | 112 | ~ | De Nol S29 |
| 0 | | 112 | • | |
| SUCRALFATE | 25 50 | 100 | | |
| Tab 1 g | (48.28) | 120 | | Carafate |
| | (40.20) | | | Jaraiale |
| Bile and Liver Therapy | | | | |
| RIFAXIMIN – Special Authority see SA1461 below – Retail pharm | acv | | | |
| Tab 550 mg | | 56 | ~ | Xifaxan |
| BACA1461 Crassial Authority for Subaidy | | | | |

➡SA1461 Special Authority for Subsidy

Initial application only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid for 6 months where the patient has hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

Renewal only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Hyperglycaemic Agents

DIAZOXIDE - Special Authority see SA1320 on the next page - Retail pharmacy

| Cap 25 mg110.00 | 100 |
|-----------------------------|----------|
| Cap 100 mg | 100 |
| Oral liq 50 mg per ml620.00 | 30 ml OP |

- Proglicem S29
- ✓ Proglicem S29
- ✔ Proglycem S29

ALIMENTARY TRACT AND METABOLISM

| | Subsidy (Manufacturer's P \$ | rice) Sub Per | Fully sidised | Brand or Generic Manufacturer |
|---|------------------------------------|------------------|------------------|--|
| ⇒SA1320 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid glycaemia caused by hyperinsulinism. Renewal from any relevant practitioner. Approvals valid without fur priate and the patient is benefiting from treatment. | | | | |
| GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit – Up to 5 kit available on a PSO | | 1 | 🖌 Glu | ucagen Hypokit |
| Insulin - Short-acting Preparations | | | | |
| NSULIN NEUTRAL ▲ Inj human 100 u per ml | 25.26 | 10 ml OP | ✔ Ac | trapid mulin R |
| ▲ Inj human 100 u per ml, 3 ml | 42.66 | 5 | 🖌 Ac | trapid Penfill mulin R |
| Insulin - Intermediate-acting Preparations | | | | |
| NSULIN ASPART WITH INSULIN ASPART PROTAMINE ▲ Inj 100 iu per ml, 3 ml prefilled pen | | 5 | 🗸 No | voMix 30 FlexPen |
| NSULIN ISOPHANE ▲ Inj human 100 u per ml | 17.68 | 10 ml OP | | mulin NPH otaphane |
| ▲ Inj human 100 u per ml, 3 ml | 29.86 | 5 | 🖌 Hu | mulin NPH otaphane Penfill |
| NSULIN ISOPHANE WITH INSULIN NEUTRAL Inj human with neutral insulin 100 u per ml | 25.26 | 10 ml OP | | mulin 30/70 ctard 30 |
| Inj human with neutral insulin 100 u per ml, 3 ml | 42.66 | 5 | ✓ Per ✓ Per | mulin 30/70 nMix 30 nMix 40 nMix 50 |
| NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE ▲ Inj lispro 25% with insulin lispro protamine 75% 100 u per ml, | | | | |
| 3 ml ▲ Inj lispro 50% with insulin lispro protamine 50% 100 u per ml, | | 5 | 🖌 Hu | malog Mix 25 |
| 3 ml | | 5 | 🖌 Hu | malog Mix 50 |
| Insulin - Long-acting Preparations | | | | |
| NSULIN GLARGINE ▲ Inj 100 u per ml, 10 ml ▲ Inj 100 u per ml, 3 ml | 94.50 | 1 5 | ✔ Lai ✔ Lai | ntus |
| Inj 100 u per ml, 3 ml disposable pen Insulin - Rapid Acting Preparations | 94.50 | 5 | 🖌 La | ntus SoloStar |
| | | | | |
| NSULIN ASPART ▲ Inj 100 u per ml, 3 ml syringe ▲ Inj 100 u per ml, 3 ml | 51.19 | 5 5 | 🖌 No | voRapid FlexPen voRapid Penfill |
| Inj 100 u per ml, 10 ml | | 1 | 🖌 No | voRapid |

| () | Subsidy ⁄Ianufacturer's Pri \$ | ice) Per | Full <u>y</u> Subsidised | d Generic |
|--|--------------------------------------|-------------|-----------------------------|-----------------|
| INSULIN GLULISINE | | | | |
| ▲ Inj 100 u per ml, 10 ml | | 1 | | Apidra |
| Inj 100 u per ml, 3 ml | | 5 | | Apidra |
| Inj 100 u per ml, 3 ml disposable pen | 46.07 | 5 | ~ | Apidra SoloStar |
| NSULIN LISPRO | | | | |
| Inj 100 u per ml, 10 ml | | 10 ml Ol | | Humalog |
| Inj 100 u per ml, 3 ml | 59.52 | 5 | ~ | Humalog |
| Alpha Glucosidase Inhibitors | | | | |
| CARBOSE | | | | |
| ₭ Tab 50 mg | | 90 | | Glucobay |
| | 9.82 | | - | Accarb |
| Tab 100 mg | | 90 | | Glucobay |
| | 15.83 | | V | Accarb |
| Oral Hypoglycaemic Agents | | | | |
| GLIBENCLAMIDE | | | | |
| k Tab 5 mg | 5.00 | 100 | ~ | Daonil |
| GLICLAZIDE | | | | |
| ₭ Tab 80 mg | 11.50 | 500 | ~ | Glizide |
| BLIPIZIDE | | | | |
| ₭ Tab 5 mg | 2.85 | 100 | ~ | Minidiab |
| Minidiab to be Sole Supply on 1 October 2015 | | | | |
| IETFORMIN HYDROCHLORIDE | | | | |
| Tab immediate-release 500 mg | 12.30 | 1,000 | ~ | Apotex |
| Tab immediate-release 850 mg | 10.10 | 500 | ~ | Apotex |
| PIOGLITAZONE | | | | |
| ₭ Tab 15 mg | 1.50 | 28 | ~ | Pizaccord |
| Tab 30 mg | 2.50 | 28 | ~ | Pizaccord |
| 🖌 Tab 45 mg | 3.50 | 28 | ~ | Pizaccord |

Diabetes Management

Ketone Testing

| BLOOD KETONE DIAGNOSTIC TEST METER – Up to 1 meter available on Meter funded for the purposes of blood ketone diagnostics only. Patient h at risk of future episodes or patient is on an insulin pump. Only one meter Meter | as had one or mor per patient will be | |
|--|--|---|
| KETONE BLOOD BETA-KETONE ELECTRODES a) Maximum of 20 strip per prescription b) Up to 10 strip available on a PSO | | |
| Test strip – Not on a BSO15.50 | 10 strip OP | Freestyle Optium Ketone |
| SODIUM NITROPRUSSIDE – Maximum of 50 strip per prescription | | |
| * Test strip - Not on a BSO6.00 | 50 strip OP | Accu-Chek Ketur-Test |
| 14.14 | | ✓ Ketostix |

ALIMENTARY TRACT AND METABOLISM

‡ safety cap *Three months or six months, as applicable, dispensed all-at-once

| Blood Glucose Testing BLOOD GLUCOSE DIAGNOSTIC TEST METER – Subsidy by end a) Maximum of 1 pack per prescription b) Up to 1 pack available on a PSO c) A diagnostic blood glucose test meter is subsidised for a patie 1) is receiving insulin or sulphonylurea therapy; or 2) is pregnant with diabetes; or 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia 4) has a genetic or an acquired disorder of glucose homeostas bnly one CareSens meter per patient. No further prescriptions will b | ent who: a; or | | | |
|---|--|---|--------------------------|---|
| a) Maximum of 1 pack per prescription b) Up to 1 pack available on a PSO c) A diagnostic blood glucose test meter is subsidised for a patient 1) is receiving insulin or sulphonylurea therapy; or 2) is pregnant with diabetes; or 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia 4) has a genetic or an acquired disorder of glucose homeostas | ent who: a; or | | | |
| or the avoidance of doubt patients who have previously received a f neter. The prescription must be endorsed accordingly. Pharmacists record of prior dispensing of insulin or sulphonylureas. Meter with 50 lancets, a lancing device and 10 diagnostic test | be subsidised funded meter, | for patients v other than C | /ho alread areSens, a | y have a CareSens mete are eligible for a CareSe |
| strips | 20.00 | 1 OP | V C | areSens II areSens N areSens N POP |
| Note: Only 1 meter available per PSO | | | | |
| as endorsed where there exists a record of prior dispensing 2) Prescribed on the same prescription as insulin or a sulphony or 3) Prescribed for a pregnant woman with diabetes and endors 4) Prescribed for a patient on home TPN at risk of hypoglycae 5) Prescribed for a patient with a genetic or an acquired disord and metabolic syndrome and endorsed accordingly. Blood glucose test strips – Note differing brand requirements | ylurea in which sed accordingly smia or hyperg | i case the pre y; or lycaemia and | escription i | d accordingly; or |
| below | 10.56 28.75 | 50 test OP | V C | areSens areSens N ccu-Chek |
| | 20.70 | | | Performa |
| a) Accu-Chek Performa brand: Special Authority see SA1294 | 1 holow Dot | il nhormoov | | reestyle Optium |
| b) Freestyle Optium brand: Special Authority see SA1291 be c) Note: Accu-Chek Performa and Freestyle Optium are not a SA1294 Special Authority for Subsidy otes: Application details may be obtained from PHARMAC's websi | elow – Retail p available on a | harmacy PSO | | an be sent to: |
| PHARMAC | | J | | |
| PO Box 10 254 Facsimile: (04) 974 4788 | | | | |
| Wellington Email: bgstrips@pharmac.govt.nz | | | | |

PO Box 10 254 Facsimile: (04) 974 4788 Wellington Email: bgstrips@pharmac.govt.nz

| | Subsidy (Manufacturer's F \$ | Price) Su Per | bsidised Ge | and or neric inufacturer |
|---|--|---------------------------------|------------------|--|
| OOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED) The number of test strips available on a prescription is r 1) Prescribed for a patient on insulin or a sulphonylurea as endorsed where there exists a record of prior dis | and endorsed accord | lingly. Pharma sulphonylurea | ; or | |
| Prescribed on the same prescription as insulin or a s or Prescribed for a pregnant woman with diabetes and Prescribed for a patient on home TPN at risk of hyp Prescribed for a patient with a genetic or an acquire and metabolic syndrome and endorsed accordingly. | l endorsed accordingl oglycaemia or hyperg d disorder of glucose | y; or lycaemia and | endorsed ac | cordingly; or |
| Blood glucose test strips | | 50 test OP | ✔ Sense | oCard |
| nsulin Syringes and Needles | | | | |
| ubsidy is available for disposable insulin syringes, needle r the supply of insulin or when prescribed for an insulin pr notate the prescription as endorsed where there exists a SULIN PEN NEEDLES – Maximum of 100 dev per prescri | atient and the prescri record of prior dispens | ption is endor | sed accordin | |
| 29 g $	imes$ 12.7 mm | | 30 | 🖌 B-D N | licro-Fine |
| | 10.50 | 100 | 🖌 B-D N | licro-Fine |
| 31 g $	imes$ 5 mm | 11.75 | 100 | 🖌 B-D N | licro-Fine |
| $31 \text{ g} \times 6 \text{ mm}$ | | 100 | 🖌 ABM | |
| 31 g × 8 mm | 3.15 | 30 | 🖌 B-D N | licro-Fine |
| - | 10.50 | 100 | ✔ B-D N ✔ ABM | licro-Fine |
| $32 \text{ g} \times 4 \text{ mm}$ | 10.50 | 100 | • • • • • • | licro-Fine |
| BM 31 g \times 8 mm to be delisted 1 September 2015) | | | | |
| SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEE | | 100 day par p | rocarintian | |
| Syringe 0.3 ml with 29 g × 12.7 mm needle | | 100 dev per p | rescription | |
| Synnge 0.5 mil with 29 g × 12.7 min needle | (1.99) | 10 | BDI | lltra Fine |
| | 13.00 | 100 | ✓ B-D U | |
| Syringe 0.3 ml with 31 g $	imes$ 8 mm needle | | 100 | P D-D 0 | |
| | (1.99) | 10 | B-D II | Itra Fine II |
| | 13.00 | 100 | - | litra Fine II |
| Syringe 0.5 ml with 29 g $	imes$ 12.7 mm needle | | 100 | • 0-00 | |
| | (1.99) | 10 | B-D I | Iltra Fine |
| | 13.00 | 100 | ✓ B-D U | |
| Syringe 0.5 ml with 31 g $	imes$ 8 mm needle | | 10 | • 550 | |
| | (1.99) | 10 | B-D I | Iltra Fine II |
| | 13.00 | 100 | | lltra Fine II |
| Syringe 1 ml with 29 g $	imes$ 12.7 mm needle | | 100 | ✓ ABM | |
| | 1.30 | 10 | | |
| | (1.99) | | B-D II | lltra Fine |
| | | 100 | ✓ B-D U | |
| | | | ✓ ABM | |
| Svringe 1 ml with 31 g \times 8 mm needle | 13.00 13.00 | 100 | | |
| Syringe 1 ml with 31 g \times 8 mm needle $\hfill \hfill \hf$ | 13.00 | 100 10 | | |
| Syringe 1 ml with 31 g \times 8 mm needle $\hfill \hfill \hf$ | 13.00 1.30 | 100 10 | | Iltra Fine II |
| Syringe 1 ml with 31 g \times 8 mm needle $\hfill \hfill \hf$ | 13.00 | | B-D U | Iltra Fine II I ltra Fine II |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|-------------|---------------------|--|
| Insulin Pumps | | | | |
| INSULIN PUMP – Special Authority see SA1237 below – Retail p a) Maximum of 1 dev per prescription b) Only on a prescription c) Maximum of 1 insulin pump per patient each four year perior Min basal rate 0.025 U/h; black colour Min basal rate 0.025 U/h; blue colour Min basal rate 0.025 U/h; green colour Min basal rate 0.025 U/h; pink colour | pd. 4,500.00 4,500.00 4,500.00 4,500.00 | 1 1 1 | | unimas Vibe unimas Vibe unimas Vibe unimas Vibe |
| Min basal rate 0.025 U/h; silver colour Min basal rate 0.05 U/h; blue colour | | 1 1 | V P | animas Vibe Paradigm 522 Paradigm 722 |
| Min basal rate 0.05 U/h; clear colour | 4,400.00 | 1 | 🖌 P | Paradigm 522 Paradigm 722 |
| Min basal rate 0.05 U/h; pink colour | | 1 | 🖌 P | Paradigm 522 Paradigm 722 |
| Min basal rate 0.05 U/h; purple colour | | 1 | 🖌 P | Paradigm 522 Paradigm 722 |
| Min basal rate 0.05 U/h; smoke colour | 4,400.00 | 1 | | Paradigm 522 Paradigm 722 |

SA1237 Special Authority for Subsidy

Notes: Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

| The IPP Co-ordinator | Phone: (04) 460 4990 |
|----------------------|----------------------------|
| PHARMAC | Facsimile: (04) 974 7806 |
| PO Box 10 254 | Email: ipp@pharmac.govt.nz |
| Wellington | |

Insulin Pump Consumables

➡SA1240 Special Authority for Subsidy

Notes: Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

| The IPP Co-ordinator | Phone: (04) 460 4990 | | | |
|-----------------------|--------------------------------------|---------------------|---------|----------------------|
| PHARMAC | Facsimile: (04) 974 7806 | | | |
| PO Box 10 254 | Email: ipp@pharmac.govt.nz | | | |
| Wellington | | | | |
| INSULIN PUMP ACCESS | ORIES - Special Authority see SA1240 |) above – Retail pl | harmacy | |
| a) Maximum of 1 cap | per prescription | | | |
| b) Only on a prescrip | tion | | | |
| c) Maximum of 1 pres | scription per 180 days. | | | |
| Battery cap | | | 1 | 🖌 Animas Battery Cap |
| | | | | |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|----------|---------------------|-------------------------------------|
| INSULIN PUMP INFUSION SET (STEEL CANNULA) – Special a) Maximum of 3 sets per prescription | Authority see SA1240 |) on the | e previous | page – Retail pharmacy |
| b) Only on a prescription | | | | |
| c) Maximum of 13 infusion sets will be funded per year. | | | | |
| 10 mm steel needle; 29 G; manual insertion; 60 cm tubing \times | | | | |
| 10 with 10 needles | 130.00 | 1 OP | 🖌 Pa | aradigm Sure-T MMT-884 |
| 10 mm steel needle; 29 G; manual insertion; 60 cm tubing $	imes$ | | | | |
| 10 with 10 needles; luer lock | 130.00 | 1 OP | 🗸 S | ure-T MMT-883 |
| 10 mm steel needle; 29 G; manual insertion; 80 cm tubing $	imes$ | | | | |
| 10 with 10 needles | 130.00 | 1 OP | 🖌 Pa | aradigm Sure-T MMT-886 |
| 10 mm steel needle; 29 G; manual insertion; 80 cm tubing $	imes$ | | | | |
| 10 with 10 needles; luer lock | 130.00 | 1 OP | 🗸 S | ure-T MMT-885 |
| 6 mm steel cannula; straight insertion; 60 cm grey line \times 10 | | | | |
| with 10 needles | 130.00 | 1 OP | 🖌 C | ontact-D |
| 6 mm steel needle; 29 G; manual insertion; 60 cm tubing $	imes$ | | | | |
| 10 with 10 needles | 130.00 | 1 OP | V P | aradigm Sure-T |
| | | | • • | MMT-864 |
| 6 mm steel needle; 29 G; manual insertion; 60 cm tubing $	imes$ | | | | |
| 10 with 10 needles; luer lock | 130.00 | 1 OP | 19 | ure-T MMT-863 |
| 6 mm steel needle; 29 G; manual insertion; 80 cm tubing \times | | | • 3 | |
| | | | | anadiana Cuna T |
| 10 with 10 needles | 130.00 | 1 OP | V Pa | aradigm Sure-T MMT-866 |
| | | | | IVIIVI 1-800 |
| 6 mm steel needle; 29 G; manual insertion; 80 cm tubing \times | 400.00 | | | |
| 10 with 10 needles; luer lock | | 1 OP | VS | ure-T MMT-865 |
| 8 mm steel cannula; straight insertion; 110 cm grey line $	imes$ 10 | | | | _ |
| with 10 needles | 130.00 | 1 OP | ✔ C | ontact-D |
| 8 mm steel cannula; straight insertion; 60 cm grey line \times 10 | | | | |
| with 10 needles | | 1 OP | 🖌 C | ontact-D |
| 8 mm steel needle; 29 G; manual insertion; 60 cm tubing $	imes$ | | | | |
| 10 with 10 needles | 130.00 | 1 OP | 🖌 Pa | aradigm Sure-T |
| | | | | MMT-874 |
| 8 mm steel needle; 29 G; manual insertion; 60 cm tubing $	imes$ | | | | |
| 10 with 10 needles; luer lock | 130.00 | 1 OP | 🗸 S | ure-T MMT-873 |
| 8 mm steel needle; 29 G; manual insertion; 80 cm tubing \times | | | | |
| 10 with 10 needles | 130.00 | 1 OP | 🖌 Pi | aradigm Sure-T |
| | | | • • | MMT-876 |
| 8 mm steel needle; 29 G; manual insertion; 80 cm tubing $	imes$ | | | | |
| 10 with 10 needles; luer lock | | 1 OP | ~ 9 | ure-T MMT-875 |
| | | | | |

| | Subsidy (Manufacturer's \$ | Price) Sub Per | Fully Brand or osidised Generic Manufactu | urer |
|--|----------------------------------|-------------------|---|------------------|
| ISULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN | | HINSERTION | DEVICE) - Spec | ial Authority se |
| A1240 on page 28 – Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription | | | opco | |
| c) Maximum of 13 infusion sets will be funded per year. 13 mm teflon cannula; angle insertion; insertion device; 110 cm grey line × 10 with 10 needles | | 1 OP | ✔ Inset 30 | |
| 13 mm teflon cannula; angle insertion; insertion device; 60 cm blue line × 10 with 10 needles | | 1 OP | ✓ Inset 30 | |
| 13 mm teflon cannula; angle insertion; insertion device; 60 cm grey line × 10 with 10 needles | | 1 OP | ✓ Inset 30 | |
| 13 mm teflon cannula; angle insertion; insertion device; 60 cm pink line × 10 with 10 needles | | 1 OP | ✓ Inset 30 | |
| ISULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN | | | | page 28 – Reta |
| narmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 13 mm teflon cannula; angel insertion; 60 cm grey line × 5 | | | | |
| with 10 needles | 120.00 | 1 OP | Comfort Sh | ort |
| 10 needles | | 1 OP | Paradigm S MMT-382 | ilhouette |
| 13 mm teflon cannula; angle insertion; 45 cm line \times 10 with 10 needles | 130.00 | 1 OP | ✓ Paradigm S MMT-368 | ilhouette |
| 13 mm teflon cannula; angle insertion; 60 cm line \times 10 with 10 needles | | 1 OP | ✓ Paradigm S MMT-381 | Silhouette |
| 13 mm teflon cannula; angle insertion; 80 cm line \times 10 with 10 needles | | 1 OP | ✓ Paradigm S MMT-383 | Silhouette |
| 17 mm teflon cannula; angle insertion; 110 cm grey line × 5 with 10 needles | | 1 OP | Comfort | |
| 17 mm teflon cannula; angle insertion; 110 cm line \times 10 with 10 needles | | 1 OP | ✓ Paradigm S MMT-377 | ilhouette |
| 17 mm teflon cannula; angle insertion; 110 cm line \times 10 with 10 needles; luer lock | | 1 OP | ✓ Silhouette | MMT-371 |
| 17 mm teflon cannula; angle insertion; 60 cm grey line × 5 with 10 needles | | 1 OP | ✓ Comfort | |
| 17 mm teflon cannula; angle insertion; 60 cm line \times 10 with 10 needles | | 1 OP | ✓ Paradigm S MMT-378 | Silhouette |
| 17 mm teflon cannula; angle insertion; 60 cm line \times 10 with 10 needles; luer lock | | 1 OP | Silhouette | MMT-373 |
| 17 mm teflon cannula; angle insertion; 80 cm line \times 10 with 10 needles | | 1 OP | Paradigm S | |
| | | | MMT-384 | |

| | Subsidy (Manufacturer's Price) \$ | Sub Per | Fully sidised | Brand or Generic Manufacturer |
|--|---|------------|------------------|-------------------------------------|
| INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH see SA1240 on page 28 – Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription | T INSERTION WITH | INSERTIO | ON DE | VICE) – Special Authority |
| c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; insertion device; 110 cm grey line × 10 with 10 needles | 140.00 1 | OP | 🗸 Ir | nset II |
| 6 mm teflon cannula; straight insertion; insertion device; 45 cm blue tubing \times 10 with 10 needles | 130.00 1 | OP | ✔ P | aradigm Mio MMT-941 |
| 6 mm teflon cannula; straight insertion; insertion device; 45 cm pink tubing \times 10 with 10 needles | 130.00 1 | OP | ✔ P | aradigm Mio MMT-921 |
| 6 mm teflon cannula; straight insertion; insertion device; 60 cm blue tubing × 10 with 10 needles | 130.00 1 | OP | ✔ P | aradigm Mio MMT-943 |
| 6 mm teflon cannula; straight insertion; insertion device; 60 cm pink tubing × 10 with 10 needles | 130.00 1 | OP | ✔ P | aradigm Mio MMT-923 |
| 6 mm teflon cannula; straight insertion; insertion device; 80 cm blue tubing \times 10 with 10 needles | 130.00 1 | OP | ✔ P | aradigm Mio MMT-945 |
| 6 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing \times 10 with 10 needles | 130.00 1 | OP | ✔ P | aradigm Mio MMT-965 |
| 6 mm teflon cannula; straight insertion; insertion device; 80 cm pink tubing × 10 with 10 needles | 130.00 1 | OP | ✔ P | aradigm Mio MMT-925 |
| 6 mm teflon cannula; straight insertionl insertion device; 60 cm blue line × 10 with 10 needles 6 mm teflon cannula; straight insertionl insertion device; 60 | 140.00 1 | OP | 🗸 in | iset II |
| cm grey line × 10 with 10 needles 6 mm teflon cannula; straight insertionl insertion device; 60 | | OP | | nset II |
| cm pink line × 10 with 10 needles 9 mm teflon cannula; straight insertion; insertion device; 60 cm blue line × 10 with 10 needles | | OP OP | | nset II nset II |
| 9 mm teflon cannula; straight insertion; insertion device; 60 cm grey line × 10 with 10 needles 9 mm teflon cannula; straight insertion; insertion device; 60 | 140.00 1 | OP | 🗸 Ir | nset II |
| 9 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing × 10 with 10 needles | | OP OP | | nset II |
| 9 mm teflon cannula; straight insertionl insertion device; 110 | | - | | aradigm Mio MMT-975 |
| cm grey line \times 10 with 10 needles | 140.00 1 | OP | 🖌 Ir | nset II |

| | Subsidy (Manufacturer's F \$ | Price) Su Per | Fully Brand or Ibsidised Generic Manufacturer |
|--|------------------------------------|------------------|---|
| INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGI | HT INSERTION) | - Special A | uthority see SA1240 on page 28 - |
| Retail pharmacy a) Maximum of 3 sets per prescription | | | |
| b) Only on a prescription | | | |
| c) Maximum of 13 infusion sets will be funded per year. | | | |
| 6 mm teflon cannula; straight insertion; 110 cm tubing \times 10 | | | |
| with 10 needles | | 1 OP | Paradigm Quick-Set MMT-398 |
| 6 mm teflon cannula; straight insertion; 110 cm tubing \times 10 | | | |
| with 10 needles; luer lock | | 1 OP | Quick-Set MMT-391 |
| 6 mm teflon cannula; straight insertion; 60 cm tubing \times 10 | | 4.00 | |
| with 10 needles | | 1 OP | Paradigm Quick-Set MMT-399 |
| 6 mm teflon cannula; straight insertion; 60 cm tubing $	imes$ 10 |) | | |
| with 10 needles; luer lock | | 1 OP | Quick-Set MMT-393 |
| 6 mm teflon cannula; straight insertion; 80 cm tubing \times 10 | | | |
| with 10 needles | | 1 OP | Paradigm Quick-Set MMT-387 |
| 9 mm teflon cannula; straight insertion; 106 cm tubing $	imes$ 10 | | | |
| with 10 needles | 130.00 | 1 OP | Paradigm Quick-Set MMT-396 |
| 9 mm teflon cannula; straight insertion; 110 cm tubing × 10 with 10 needles; luer lock | | 1 OP | ✔ Quick-Set MMT-390 |
| 9 mm teflon cannula; straight insertion; 60 cm tubing \times 10 | | | |
| with 10 needles | | 1 OP | Paradigm Quick-Set MMT-397 |
| 9 mm teflon cannula; straight insertion; 60 cm tubing $	imes$ 10 |) | | |
| with 10 needles; luer lock | | 1 OP | ✔ Quick-Set MMT-392 |
| 9 mm teflon cannula; straight insertion; 80 cm tubing $	imes$ 10 |) | | |
| with 10 needles | 130.00 | 1 OP | Paradigm Quick-Set MMT-386 |
| INSULIN PUMP RESERVOIR - Special Authority see SA1240 c | on page 28 – Ret | ail pharmacy | |
| a) Maximum of 3 sets per prescription b) Only on a prescription | | | |
| c) Maximum of 13 packs of reservoir sets will be funded per y | /oar | | |
| $10 \times \text{luer lock conversion cartridges 1.8 ml for Paradigm}$ | | | |
| pumps | | 1 OP | ADR Cartridge 1.8 |
| $10 \times$ luer lock conversion cartridges 3.0 ml for Paradigm | | | Ū |
| pumps | | 1 OP | ADR Cartridge 3.0 |
| Cartridge 200 U, luer lock × 10 | | 1 OP | Animas Cartridge |
| Cartridge for 5 and 7 series pump; 1.8 ml \times 10 | 50.00 | 1 OP | Paradigm 1.8 Reservoir |
| Cartridge for 7 series pump; 3.0 ml \times 10 | 50.00 | 1 OP | Paradigm 3.0 Reservoir |
| Syringe and cartridge for 50X pump, 3.0 ml $	imes$ 10 | | 1 OP | ✓ 50X 3.0 Reservoir |
| | | | |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|----------|---------------------|-------------------------------------|
| Digestives Including Enzymes | | | | |
| PANCREATIC ENZYME | | | | |
| Cap EC 10,000 BP u lipase, 9,000 BP u amylase and 210 BP u protease Creon 10000 to be Sole Supply on 1 November 2015 | | 100 | ✔ C | reon 10000 |
| Cap EC 25,000 BP u lipase, 18,000 BP u amylase, 1,000 BP u protease Creon 25000 to be Sole Supply on 1 November 2015 | 94.38 | 100 | ✔ C | reon 25000 |
| Cap EC 25,000 BP u lipase, 22,500 BP u amylase, 1,250 BP u protease | 94.40 | 100 | 🖌 P | anzytrat |
| URSODEOXYCHOLIC ACID – Special Authority see SA1383 bel Cap 250 mg – For ursodeoxycholic acid oral liquid formula- tion refer, page 207 | | / 100 | ✓ <u>U</u> | rsosan |

SA1383 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Cirrhosis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cirrhosis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

continued...

| Subsidy (Manufacturer's Price) | Subs | Fully sidised | Brand or Generic |
|-----------------------------------|------|------------------|---------------------|
| \$ | Per | ~ | Manufacturer |

continued...

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Cirrhosis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure – doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Laxatives

Bulk-forming Agents

| •• | | | |
|--|-----------------------------------|--|---|
| ISPAGHULA (PSYLLIUM) HUSK – Only on a prescription * Powder for oral soln MUCILAGINOUS LAXATIVES WITH STIMULANTS | 5.51 | 500 g OP | ✓ <u>Konsyl-D</u> |
| * Dry | 2.41 (8.72) 6.02 (17.32) | 200 g OP 500 g OP | Normacol Plus Normacol Plus |
| Faecal Softeners | | | |
| DOCUSATE SODIUM – Only on a prescription * Tab 50 mg * Tab 120 mg * Enema conc 18% DOCUSATE SODIUM WITH SENNOSIDES * Tab 50 mg with sennosides 8 mg POLOXAMER – Only on a prescription Not funded for use in the ear. * Oral drops 10% | 3.13 5.40 4.40 | 100 100 100 ml OP 200 30 ml OP | ✓ <u>Coloxyl</u> ✓ <u>Coloxyl</u> ✓ Coloxyl ✓ Laxsol ✓ <u>Coloxyl</u> |
| Osmotic Laxatives | | | |
| GLYCEROL * Suppos 3.6 g – Only on a prescription PSM to be Sole Supply on 1 October 2015 | 6.50 | 20 | 🗸 PSM |
| LACTULOSE – Only on a prescription * Oral liq 10 g per 15 ml | 3.84 | 500 ml | ✓ Laevolac |

MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE AND SODIUM CHLORIDE – Special Authority see SA1473 on the next page – Retail pharmacy Powder for oral soln 13.125 g with potassium chloride

| | Subsidy (Manufacturer's Price) \$ |) Si Per | Fully ubsidised | Brand or Generic Manufacturer |
|---|---|-------------|--------------------|-------------------------------------|
| SA1473 Special Authority for Subsidy | | | | |
| tial application from any relevant practitioner. Approvals va | lid for 6 months for app | lications | meeting t | he following criteria: |
| th: | | | | |
| The patient has problematic constipation despite an where lactulose is not contraindicated; and The patient would otherwise require a per rectal prepa | | oral pha | rmacothe | erapies including lactule |
| enewal from any relevant practitioner. Approvals valid for 1 | | otiont in d | ompliant | and is continuing to g |
| nefit from treatment. | | | Jomphani | and is continuing to g |
| DDIUM ACID PHOSPHATE – Only on a prescription | | | | |
| Enema 16% with sodium phosphate 8% | 2.50 | 1 | ✔ F | leet Phosphate Enema |
| DDIUM CITRATE WITH SODIUM LAURYL SULPHOACETAT | E – Only on a prescrip | tion | | |
| Enema 90 mg with sodium lauryl sulphoacetate 9 mg per | ml, | | | |
| 5 ml | | 50 | ✓ <u>M</u> | licolette |
| timulant Laxatives | | | | |
| SACODYL – Only on a prescription | | | | |
| Tab 5 mg Lax-Tab to be Sole Supply on 1 November 2015 | 5.99 | 200 | 🖌 Li | ax-Tab |
| Suppos 5 mg | 3.00 | 6 | 🖌 D | ulcolax |
| Suppos 10 mg | 3.00 | 6 | 🗸 D | ulcolax |
| ENNA – Only on a prescription | | | | |
| Tab, standardised | | 20 | | |
| | (1.72) | | S | enokot |
| | 2.17 | 100 | c | enokot |
| | (6.84) | | 3 | enokol |
| letabolic Disorder Agents | | | | |
| aucher's Disease | | | | |
| IGLUCERASE – Special Authority see SA0473 below – Ret | ail pharmacy | | | |
| Inj 40 iu per ml, 200 iu vial | 1,072.00 | 1 | 🖌 C | erezyme |
| Inj 40 iu per ml, 400 iu vial | 2,144.00 | 1 | 🗸 C | erezyme |
| SA0473 Special Authority for Subsidy | | | | |

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

| The Co-ordinator, Gaucher's Treatment Panel | Phone: (04) 460 4990 |
|---|-------------------------------------|
| PHARMAC, PO Box 10 254 | Facsimile: (04) 916 7571 |
| Wellington | Email: gaucherpanel@pharmac.govt.nz |

| | Subsidy (Manufacturer's | | Fully Brand or bidised Generic Manufacturer |
|---|------------------------------|-------------------|---|
| Mouth and Throat | \$ | Per | |
| Agents Used in Mouth Ulceration | | | |
| BENZYDAMINE HYDROCHLORIDE | | | |
| Soln 0.15% – Higher subsidy of up to \$17.01 per 500 ml with Endorsement | | 200 ml | |
| Lindischent | (8.50) | 200 111 | Difflam |
| | 9.00 | 500 ml | |
| Additional subsidy by endorsement for a patient who has o | (17.01) ral muccasitic ac | a recult of treat | Difflam |
| tion is endorsed accordingly. | rai mucosilis as | a result of freat | anent for cancer, and the prescrip- |
| CHLORHEXIDINE GLUCONATE | | | |
| Mouthwash 0.2% | 2.57 | 200 ml OP | ✓ healthE |
| healthE to be Sole Supply on 1 October 2015 | | | |
| CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE * Adhesive gel 8.7% with cetalkonium chloride 0.01% | 2.06 | 15 g OP | |
| Adhesive ger 0.7 /0 with cetaikonium chionde 0.01 /0 | (6.00) | 15 9 01 | Bonjela |
| SODIUM CARBOXYMETHYLCELLULOSE | , , | | |
| With pectin and gelatin paste | | 56 g OP | Stomahesive |
| | 1.52 (3.60) | 5 g OP | Orabase |
| | (3.60) 4.55 | 15 g OP | Ulabase |
| | (7.90) | 0 | Orabase |
| With pectin and gelatin powder | | 28 g OP | Stomahesive |
| TRIAMCINOLONE ACETONIDE | (10.95) | | Stomanesive |
| Paste 0.1% | 5.33 | 5 g OP | ✓ Kenalog in Orabase |
| Oropharyngeal Anti-infectives | | - 3 - | · <u></u> |
| | | | |
| AMPHOTERICIN B | F 00 | 00 | |
| Lozenges 10 mg | | 20 | Fungilin |
| MICONAZOLE Oral gel 20 mg per g | 4.79 | 40 g OP | ✓ Decozol |
| Decozol to be Sole Supply on 1 October 2015 | | | |
| NYSTATIN | | | |
| Oral liq 100,000 u per ml | 3.35 | 24 ml OP | ✓ Nilstat |
| Other Oral Agents | | | |
| For folinic mouthwash, pilocarpine oral liquid or saliva substitute for | ormula refer Sta | Indard Formula | e, page 210 |
| HYDROGEN PEROXIDE | | | |
| * Soln 10 vol – Maximum of 200 ml per prescription | 1.28 | 100 ml | ✔ PSM |
| THYMOL GLYCERIN | 0.15 | 500 ml | . / DOM |
| * Compound, BPC | 9.15 | 500 ml | ✔ PSM |

ALIMENTARY TRACT AND METABOLISM

| | Subsidy (Manufacturer's P | rice) Sub | Fully Brand sidised Gene | |
|--|------------------------------|------------|---|------------------|
| | \$ | Per | Manu | facturer |
| /itamins | | | | |
| /itamin A | | | | |
| ITAMIN A WITH VITAMINS D AND C Soln 1000 u with Vitamin D 400 u and ascorbic acid 30 mg per 10 drops | 4.50 | 10 ml OP | ✓ Vitadol | c |
| /itamin B | | | | |
| YDROXOCOBALAMIN | | | | |
| Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a PS | 02.31 5.10 | 3 | Neo-B12 ABM Hydro | 2 xocobalamin |
| YRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose b) Only on a prescription | | | | |
| Tab 25 mg – No patient co-payment payable | | 90 500 | Vitamin | |
| Tab 50 mg HIAMINE HYDROCHLORIDE – Only on a prescription | 11.00 | 500 | ✓ <u>Аро-Ру</u> | laoxine |
| Tab 50 mg | 5.62 | 100 | 🖌 Apo-Thi | amine |
| ITAMIN B COMPLEX | | | | |
| Tab, strong, BPC | 4.30 | 500 | ✓ Bplex | |
| Vitamin C | | | | |
| SCORBIC ACID a) No more than 100 mg per dose b) Only on a prescription | | | | |
| • Tab 100 mg | 7.00 | 500 | ✓ <u>Cvite</u> | |
| /itamin D | | | | |
| LFACALCIDOL • Cap 0.25 mcg • Cap 1 mcg | | 100 100 | ✔ One-Alp ✔ One-Alp | |
| Oral drops 2 mcg per ml | 60.68 | 20 ml OP | One-Alp | ha |
| ALCITRIOL · Cap 0.25 mcg | 3 03 | 30 | Airflow | |
| | 10.10 | 100 | Calcitrio | ol-AFT |
| Cap 0.5 mcg | 5.62 18.73 | 30 100 | Airflow Calcitric | AFT. |
| HOLECALCIFEROL | | | | |
| Tab 1.25 mg (50,000 iu) – Maximum of 12 tab per prescriptio | n 7.76 | 12 | Cal-d-Fo | orte |
| Multivitamin Preparations | | | | |
| ULTIVITAMINS - Special Authority see SA1036 on the next page | | | . | |
| Powder | 72.00 | 200 g OP | Paediati | ric Seravit |

ALIMENTARY TRACT AND METABOLISM

| | Subsidy (Manufacturer's Prio \$ | ce) Sul Per | Fully Brand or bsidised Generic Manufacturer |
|---|---------------------------------------|---------------------|--|
| ►SA1036 Special Authority for Subsidy | id without further | concernation of the | and notified where the notient he |
| Initial application from any relevant practitioner. Approvals va inborn errors of metabolism. | ia without further i | renewal unie | ess notified where the patient ha |
| Renewal from any relevant practitioner. Approvals valid without approval for multivitamins. | further renewal un | ess notified | where patient has had a previous |
| VITAMINS | | | |
| * Tab (BPC cap strength) * Cap (fat soluble vitamins A, D, E, K) – Special Authority see | | 1,000 | ✓ <u>Mvite</u> |
| SA1002 below - Retail pharmacy | 23.40 | 60 | Vitabdeck |
| Salong Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valie the following criteria: Either: | | newal unles | ss notified for applications meeting |
| Patient has cystic fibrosis with pancreatic insufficiency; c Patient is an infant or child with liver disease or short gut | | | |
| Minerals | | | |
| Calcium | | | |
| CALCIUM CARBONATE * Tab eff 1.75 g (1 g elemental) | 6.21 | 30 | ✓ Calsource |
| * Tab 1.25 g (500 mg elemental) | | 250 | ✓ <u>Arrow-Calcium</u> |
| CALCIUM GLUCONATE * Inj 10%, 10 ml ampoule | | 10 | ✓ Hospira |
| Fluoride | | | |
| SODIUM FLUORIDE * Tab 1.1 mg (0.5 mg elemental) | 5.00 | 100 | 🗸 PSM |
| lodine | | | |
| POTASSIUM IODATE * Tab 253 mcg (150 mcg elemental iodine) | 2.65 | 90 | ✓ NeuroTabs |
| | | 90 | • <u>Neurorabs</u> |
| | | | |
| FERROUS FUMARATE * Tab 200 mg (65 mg elemental) | 2.89 | 100 | ✓ <u>Ferro-tab</u> |
| FERROUS FUMARATE WITH FOLIC ACID Tab 310 mg (100 mg elemental) with folic acid 350 mcg | 4 75 | 60 | ✓ Ferro-F-Tabs |
| Iab 310 mg (100 mg elemental) with folic acid 350 mcg FERROUS SULPHATE | 4.75 | 00 | |
| * Tab long-acting 325 mg (105 mg elemental) | | 30 | ✓ Ferrograd |
| *‡ Oral liq 30 mg (6 mg elemental) per 1 ml | | 500 ml | Ferodan |
| FERROUS SULPHATE WITH FOLIC ACID | I | | |
| * Tab long-acting 325 mg (105 mg elemental) with folic acid 350 mcg | | 30 | |
| č | (4.29) | | Ferrograd F |
| | 15.00 | F | |
| * Inj 50 mg per ml, 2 ml ampoule | | 5 | ✓ <u>Ferrum H</u> |

ALIMENTARY TRACT AND METABOLISM

| | Subsidy (Manufacturer's Price) \$ | Sub Per | Fully osidised | Brand or Generic Manufacturer |
|--|---|------------|-------------------|-------------------------------------|
| Magnesium | | | | |
| For magnesium hydroxide mixture refer Standard Formulae, page MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule | | 10 | ✓ <u>DI</u> | <u>BL</u> |
| Zinc | | | | |
| ZINC SULPHATE * Cap 137.4 mg (50 mg elemental) | 11.00 | 100 | ✓ <u>Zi</u> | ncaps |

| Subsidy | Fu | Illy | Brand or |
|------------------------|----------|------|--------------|
| (Manufacturer's Price) | Subsidis | ed | Generic |
| \$ | Per | ~ | Manufacturer |

Antianaemics

Hypoplastic and Haemolytic

SA1469 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin ≤ 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate \leq 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate \leq 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Erythropoietin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin <100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum erythropoietin level of <500 IU/L; and
- 6 The minimum necessary dose of erythropoietin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an Unapproved Indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Erythropoietin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of erythropoietin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an Unapproved Indication

| | Subsidy | | Fully | Brand or |
|---|----------------------|------------|------------|----------------|
| | (Manufacturer's Pric | | ubsidised | Generic |
| | \$ | Per | ~ | Manufacturer |
| EPOETIN ALFA [ERYTHROPOIETIN ALFA] - Special Authority | see SA1469 on the | previous p | age – Re | etail pharmacy |
| Wastage claimable - see rule 3.3.2 on page 13 | | | • | |
| Inj 1,000 iu in 0.5 ml, syringe | | 6 | 🖌 <u>E</u> | prex |
| Inj 2,000 iu in 0.5 ml, syringe | 120.18 | 6 | ✓ E | prex |
| Inj 3,000 iu in 0.3 ml, syringe | 166.87 | 6 | 🖌 <u>E</u> | prex |
| Inj 4,000 iu in 0.4 ml, syringe | | 6 | 🖌 <u>E</u> | prex |
| Inj 5,000 iu in 0.5 ml, syringe | 243.26 | 6 | V E | prex |
| Inj 6,000 iu in 0.6 ml, syringe | | 6 | V E | prex |
| Inj 8,000 iu in 0.8 ml, syringe | 352.69 | 6 | 🖌 <u>E</u> | prex |
| Inj 10,000 iu in 1 ml, syringe | | 6 | 🖌 <u>E</u> | prex |
| Inj 40,000 iu in 1 ml, syringe | | 1 | ✓ E | prex |
| Megaloblastic | | | | |
| FOLIC ACID | | | | |
| * Tab 0.8 mg Apo-Folic Acid to be Sole Supply on 1 November 2015 | 20.60 | 1,000 | 🖌 A | po-Folic Acid |
| Tab 5 mg | 10.92 | 500 | 🗸 A | po-Folic Acid |
| Oral liq 50 mcg per ml | 24.00 | 25 ml OP | 🖌 В | iomed |
| Antifibrinolytics, Haemostatics and Local Sclere | osants | | | |
| ELTROMBOPAG – Special Authority see SA1418 below – Retai Wastage claimable – see rule 3.3.2 on page 13 | l pharmacy | | | |
| Tab 25 mg | 1.771.00 | 28 | 🖌 B | evolade |
| Tab 50 mg | | 28 | | evolade |
| | -, | | | |

SA1418 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of ≤ 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of \leq 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of >30,000 platelets per microlitre.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia, whose treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

| Inj 1 mg syringe | 1,163.75 | 1 | NovoSeven RT |
|------------------|----------|---|--------------|
| Inj 2 mg syringe | 2,327.50 | 1 | NovoSeven RT |
| Inj 5 mg syringe | 5,818.75 | 1 | NovoSeven RT |
| Inj 8 mg syringe | 9,310.00 | 1 | NovoSeven RT |

41

BLOOD AND BLOOD FORMING ORGANS

| | Subsidy (Manufacturer's Price |) Si | Fully ubsidised | Brand or Generic |
|--|---|---------------------------------|--|---|
| | \$ | Per | ~ | Manufacturer |
| ACTOR EIGHT INHIBITORS BYPASSING AGENT - [Xpharm | | | | |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. | l by the Haemophilia Tr | reaters Gr | roup in co | njunction with the Nation |
| Inj 500 U | 1,640.00 | 1 | 🖌 FE | EIBA |
| Inj 1,000 U | 3,280.00 | 1 | 🖌 FE | EIBA |
| OROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xph | arm] | | | |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. | l by the Haemophilia Tr | reaters Gr | roup in co | njunction with the Natior |
| Inj 250 iu vial | | 1 | 🖌 Xy | /ntha |
| Inj 500 iu vial | 450.00 | 1 | 🖌 X | /ntha |
| Inj 1,000 iu vial | 900.00 | 1 | 🖌 X | /ntha |
| Inj 2,000 iu vial | 1,800.00 | 1 | 🖌 X | /ntha |
| Inj 3,000 iu vial | 2,700.00 | 1 | 🖌 X | /ntha |
| IONACOG ALFA [RECOMBINANT FACTOR IX] – [Xpharm] | | | | |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. | l by the Haemophilia Tr | reaters Gr | roup in co | njunction with the Nation |
| Inj 250 iu vial | | 1 | V Be | eneFIX |
| Inj 500 iu vial | | 1 | V B | eneFIX |
| Inj 1,000 iu vial | | 1 | V B | eneFIX |
| Inj 2,000 iu vial | <i>'</i> | 1 | | eneFIX |
| For patients with haemophilia, whose treatment is managed | l by the Haemophilia Tr | reaters Gi | roup in co | njunction with the Natior |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. | | | | |
| For patients with haemophilia, whose treatment is managed | | reaters Gi 1 | ✓ A | dvate |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial | 237.50 250.00 | | ✓ A | dvate ogenate FS |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. | | 1 | ✓ Ac | dvate ogenate FS dvate |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial | | 1 1 | A K A K | dvate ogenate FS dvate ogenate FS |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial | | 1 | | dvate ogenate FS dvate ogenate FS dvate |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial | | 1 1 | | dvate ogenate FS dvate ogenate FS dvate ogenate FS |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial | | 1 1 1 | A K A K A K K K | dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 1,500 iu vial | | 1 1 1 1 | Ac Kc Kc Kc Kc Ac Kc Ac < | dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 1,500 iu vial Inj 2,000 iu vial | | 1 1 1 1 | Ac Kc Kc Kc Kc Ac Kc Ac < | dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate dvate dvate ogenate FS |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 1,500 iu vial | | 1 1 1 1 1 | | dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate dvate dvate ogenate FS |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 1,500 iu vial Inj 3,000 iu vial | | 1 1 1 1 1 | | dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate dvate dvate ogenate FS dvate |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 1,500 iu vial Inj 2,000 iu vial Inj 3,000 iu vial ODIUM TETRADECYL SULPHATE | | 1 1 1 1 1 | | dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate dvate dvate ogenate FS dvate |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 1,500 iu vial Inj 2,000 iu vial Inj 3,000 iu vial ODIUM TETRADECYL SULPHATE | | 1 1 1 1 1 | A K A K A K A A K A A K A K K K K K | dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate dvate ogenate FS dvate ogenate FS |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 1,500 iu vial Inj 3,000 iu vial Inj 3,000 iu vial SODIUM TETRADECYL SULPHATE ≰ Inj 3% 2 ml | | 1 1 1 1 1 | A K A K A K A A K A A K A K K K K K | dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate dvate dvate ogenate FS dvate |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 1,500 iu vial Inj 3,000 iu vial SODIUM TETRADECYL SULPHATE k Inj 3% 2 ml RANEXAMIC ACID | 237.50 250.00 475.00 500.00 950.00 1,000.00 1,425.00 1,900.00 2,000.00 2,850.00 3,000.00 28.50 (73.00) | 1 1 1 1 1 5 | A K A K A K A K A K K K Fi | dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate ogenate FS |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 1,500 iu vial Inj 2,000 iu vial Inj 3,000 iu vial CODIUM TETRADECYL SULPHATE Inj 3% 2 ml RANEXAMIC ACID Tab 500 mg | 237.50 250.00 475.00 500.00 950.00 1,000.00 1,425.00 1,900.00 2,000.00 2,850.00 3,000.00 28.50 (73.00) | 1 1 1 1 1 | A K A K A K A K A K K K Fi | dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate dvate ogenate FS dvate ogenate FS |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 1,500 iu vial Inj 3,000 iu vial Inj 3,000 iu vial SODIUM TETRADECYL SULPHATE | 237.50 250.00 475.00 500.00 950.00 1,000.00 1,425.00 1,900.00 2,000.00 2,850.00 3,000.00 28.50 (73.00) | 1 1 1 1 1 5 | A K A K A K A K A K K K Fi | dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate ogenate FS |
| For patients with haemophilia, whose treatment is managed Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 1,500 iu vial Inj 2,000 iu vial Inj 3,000 iu vial ODIUM TETRADECYL SULPHATE Inj 3% 2 ml RANEXAMIC ACID Tab 500 mg Vitamin K HYTOMENADIONE | 237.50 250.00 475.00 500.00 950.00 1,000.00 1,425.00 1,900.00 2,000.00 2,850.00 3,000.00 28.50 (73.00) 23.00 | 1 1 1 1 1 1 5 | AA Ka AA Ka AA Ka AA Ka Ka Ka Fi | dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate obvate ogenate FS dvate ogenate FS dvate bro-vein <u>/klokapron</u> |
| Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 1,500 iu vial Inj 3,000 iu vial SODIUM TETRADECYL SULPHATE k Inj 3% 2 ml RANEXAMIC ACID | | 1 1 1 1 1 5 | A | dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate ogenate FS dvate ogenate FS |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|--|-----------|---------------------|-------------------------------------|
| Antithrombotic Agents | | | | |
| Antiplatelet Agents | | | | |
| ASPIRIN * Tab 100 mg CLOPIDOGREL * Tab 75 mg – For clopidogrel oral liquid formulation refer, page | | 990 84 | _ | thics Aspirin EC |
| 207 DIPYRIDAMOLE * Tab 25 mg – For dipyridamole oral liquid formulation refer, page 207 | | 84 | _ | <u>rrow - Clopid</u> ersantin |
| * Tab long-acting 150 mg | | 60 | V P | ytazen SR |
| PRASUGREL – Special Authority see SA1201 below – Retail pha Tab 5 mg Tab 10 mg | 108.00 | 28 28 | ✓ Ei ✓ Ei | ffient ffient |
| ►SA1201 Special Authority for Subsidy Initial application — (coronary angioplasty and bare metal st where the patient has undergone coronary angioplasty in the prev Initial application — (drug eluting stent) from any relevant prac | ent) from any releva rious 4 weeks and is | clopido | ogrel-allergi | с*. |

a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Initial application — (stent thromobosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (coronary angioplasty and bare metal stent) from any relevant practitioner. Approvals valid for 6 months where the patient has undergone coronary angioplasty or had a bare metal cardiac stent inserted in the previous 4 weeks and is clopidogrelallergic*.

Renewal — (drug eluting stent) from any relevant practitioner. Approvals valid for 12 months where had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Note: * Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

TICAGRELOR – Special Authority see SA1382 below – Retail pharmacy

* Tab 90 mg90.00 56 🖌 Brilinta

➡SA1382 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Generic |
|--|---|-----|---------------------|---------|
| Heparin and Antagonist Preparations | | | | |
| DALTEPARIN SODIUM – Special Authority see SA1270 below – | Retail pharmacy | | | |
| Inj 2,500 iu per 0.2 ml prefilled syringe | | 10 | ~ | Fragmin |
| Inj 5,000 iu per 0.2 ml prefilled syringe | | 10 | ~ | Fragmin |
| Inj 7,500 iu per 0.75 ml graduated syringe | 60.03 | 10 | ~ | Fragmin |
| Inj 10,000 iu per 1 ml graduated syringe | 77.55 | 10 | ~ | Fragmin |
| Inj 12,500 iu per 0.5 ml prefilled syringe | | 10 | ~ | Fragmin |
| Inj 15,000 iu per 0.6 ml prefilled syringe | | 10 | ~ | Fragmin |
| Inj 18,000 iu per 0.72 ml prefilled syringe | 158.47 | 10 | ~ | Fragmin |

SA1270 Special Authority for Subsidy

Initial application — (Pregnancy or Malignancy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy or Malignancy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, Acute Coronary Syndrome, cardioversion, or prior to oral anti-coagulation).

ENOXAPARIN SODIUM – Special Authority see SA1174 on the next page – Retail pharmacy

| Inj 20 mg | | 10 | Clexane |
|------------|--------|----|-----------------------------|
| Inj 40 mg | | 10 | Clexane |
| Inj 60 mg | 74.91 | 10 | Clexane |
| Inj 80 mg | | 10 | Clexane |
| Inj 100 mg | | 10 | Clexane |
| Inj 120 mg | 155.40 | 10 | Clexane |
| Inj 150 mg | | 10 | Clexane |

| Subsidy (Manufacturer's Price) | Subsid | - ully ised | Brand or Generic | |
|-----------------------------------|--------|-------------------|---------------------|--|
| \$ | Per | ~ | Manufacturer | |

➡SA1174 Special Authority for Subsidy

Initial application — (Pregnancy or Malignancy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

- Any of the following:
 - 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
 - 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
 - 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
 - 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
 - 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy or Malignancy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

HEPARIN SODIUM

| Inj 1,000 iu per ml, 5 ml13.36 | 10 | Hospira | |
|---|----|-----------------------------|--|
| 66.80 | 50 | Hospira | |
| 61.04 | | Pfizer | |
| Inj 1,000 iu per ml, 35 ml16.00 | 1 | Hospira | |
| Inj 5,000 iu per ml, 1 ml14.20 | 5 | Hospira | |
| Inj 5,000 iu per ml, 5 ml236.60 | 50 | Pfizer | |
| Inj 25,000 iu per ml, 0.2 ml9.50 | 5 | Hospira | |
| HEPARINISED SALINE | | | |
| * Inj 10 iu per ml, 5 ml | 50 | Pfizer | |
| PROTAMINE SULPHATE | | | |
| * Inj 10 mg per ml, 5 ml | 10 | | |
| (119.23) | 10 | Artex | |
| | | 71100 | |
| Oral Anticoagulants | | | |
| DABIGATBAN | | | |
| Cap 75 mg - No more than 2 cap per day148.00 | 60 | Pradaxa | |
| Cap 110 mg | 60 | Pradaxa | |
| Cap 150 mg148.00 | 60 | ✓ Pradaxa | |
| | | | |
| RIVAROXABAN – Special Authority see SA1066 on the next page – Retail pharma | | A 34 U | |
| Tab 10 mg153.00 | 15 | Xarelto | |
| | | | |

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

SA1066 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 weeks for applications meeting the following criteria: Either:

- 1 For the prophylaxis of venous thromboembolism following a total hip replacement; or
- 2 For the prophylaxis of venous thromboembolism following a total knee replacement.

Note: Rivaroxaban is only currently indicated and subsidised for up to 5 weeks therapy for prophylaxis of venous thromboembolism following a total hip replacement and up to 2 weeks therapy for prophylaxis of venous thromboembolism following a total knee replacement.

Renewal from any relevant practitioner. Approvals valid for 5 weeks where prophylaxis for venous thromboembolism is required for patients following a subsequent total hip or knee replacement.

WARFARIN SODIUM

Note: Marevan and Coumadin are not interchangeable.

| * | Tab 1 mg | 50 | Coumadin |
|---|--------------|-----|----------|
| | 6.86 | 100 | Marevan |
| * | Tab 2 mg4.31 | 50 | Coumadin |
| * | Tab 3 mg9.70 | 100 | Marevan |
| * | Tab 5 mg5.93 | 50 | Coumadin |
| | 11.75 | 100 | Marevan |

Blood Colony-stimulating Factors

| FILGRASTIM – Special Authority see SA1259 below – Retail pl | harmacy | | |
|---|---------|---|----------------------------|
| Inj 300 mcg per 0.5 ml prefilled syringe | | 5 | Zarzio |
| Inj 480 mcg per 0.5 ml prefilled syringe | | 5 | Zarzio |

➡SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk ≥ 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5×10^9 /L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5×10^{9} /L).

Note: *Febrile neutropenia risk $\geq 20\%$ after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1384 below - Retail pharmacy

Inj 6 mg per 0.6 ml syringe 1,080.00 1 🖌 Veulastim

➡SA1384 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk $\geq 20\%^*$).

Note: *Febrile neutropenia risk \geq 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

| | Subsidy (Manufacturer's Price \$ | e) Per | Fully Subsidised | |
|--|--|-------------------|---------------------|------------------------------|
| Fluids and Electrolytes | | | | |
| Intravenous Administration | | | | |
| GLUCOSE [DEXTROSE] | | | | |
| Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO | | 5 1 | | <u>Biomed</u> Biomed |
| POTASSIUM CHLORIDE * Inj 75 mg per ml, 10 ml | | 50 | ~ | AstraZeneca |
| SODIUM BICARBONATE | | | | |
| Inj 8.4%, 50 ml a) Up to 5 inj available on a PSO b) Not in combination | 19.95 | 1 | ✔ E | Biomed |
| a) Up to 5 inj available on a PSO b) Not in combination | 20.50 | 1 | ✔ E | Biomed |
| SODIUM CHLORIDE Not funded for use as a nasal drop. Only funded for nebuliser use. | use when in conjur | nction w | ith an antil | piotic intended for nebulise |
| Inf 0.9% – Up to 2000 ml available on a PSO | | 500 ml 1.000 m | | Baxter Baxter |
| Only if prescribed on a prescription for renal dialysis, mate for emergency use. (500 ml and 1,000 ml packs) | | , | | |
| Inj 23.4%, 20 ml ampoule For Sodium chloride oral liquid formulation refer Standard I | | 5 | ✓ <u>E</u> | Biomed |
| Inj 0.9%, 5 ml – Up to 5 inj available on a PSO | | 50 | | Multichem Pfizer |
| Inj 0.9%, 10 ml – Up to 5 inj available on a PSO | | 50 | V | Multichem Pfizer |
| Inj 0.9%, 20 ml | | 6 | | Pharmacia |
| | 11.79 8.41 | 30 20 | | Pharmacia Multichem |
| TOTAL PARENTERAL NUTRITION (TPN) - Retail pharmacy-Spe | | | | |
| Infusion WATER | CBS | 1 OP | ✓ 1 | PN |
| On a prescription or Practitioner's Supply Order only whe Schedule requiring a solvent or diluent; or On a bulk supply order; or When used in the extemporaneous compounding of eye of | | n as an | injection li | isted in the Pharmaceutica |
| Purified for inj, 5 ml – Up to 5 inj available on a PSO | | 50 | | Multichem |
| Purified for inj, 10 ml – Up to 5 inj available on a PSO Purified for inj, 20 ml – Up to 5 inj available on a PSO | | 50 20 | | Multichem Multichem |
| Oral Administration | | | | |
| CALCIUM POLYSTYRENE SULPHONATE Powder | 160.95 | 100 g OI | | Calcium Resonium |
| COMPOUND ELECTROLYTES Powder for oral soln – Up to 10 sach available on a PSO | | 10 g Or | | Enerlyte |

| | Subsidy | | Fully | Brand or |
|---|-----------------|-------------|---------|--------------------------------|
| | (Manufacturer's | | sidised | Generic |
| | \$ | Per | ~ | Manufacturer |
| EXTROSE WITH ELECTROLYTES | | | | |
| Soln with electrolytes | 6.55 | 1,000 ml OP | ✓ P | <u>edialyte -</u> Bubblegum |
| HOSPHORUS | | | | |
| Tab eff 500 mg (16 mmol) | | 100 | 🖌 P | hosphate-Sandoz |
| OTASSIUM CHLORIDE | | | | |
| Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq) | 5.26 | 60 | | |
| | (11.85) | | С | hlorvescent |
| Tab long-acting 600 mg (8 mmol) Span-K to be Sole Supply on 1 October 2015 | 7.42 | 200 | ✔ S | pan-K |
| ODIUM BICARBONATE | | | | |
| Cap 840 mg | 8.52 | 100 | 🖌 S | odibic |
| ODIUM POLYSTYRENE SULPHONATE | | | | |
| Powder | | 454 g OP | 🖌 R | esonium-A |
| Resonium-A to be Sole Supply on 1 October 2015 | | 5 | | |

| | Subsidy | | Fully Dro | nd or |
|---|-------------------|-----------|----------------------------|----------------|
| | (Manufacturer's P | rice) Sub | | na or neric |
| | \$ | Per | | nufacturer |
| | | | | |
| Alpha Adrenoceptor Blockers | | | | |
| DOXAZOSIN | | | | |
| * Tab 2 mg | 6.75 | 500 | Apo-D | oxazosin |
| * Tab 4 mg | 9.67 | 500 | Apo-D | oxazosin |
| PHENOXYBENZAMINE HYDROCHLORIDE | | | | |
| * Cap 10 mg | 65.00 | 30 | 🖌 BNM® | 29 |
| PRAZOSIN | | | | |
| * Tab 1 mg | 5.53 | 100 | 🖌 Apo-P | razosin |
| * Tab 2 mg | 7.00 | 100 | 🖌 Apo-P | razosin |
| * Tab 5 mg | 11.70 | 100 | 🖌 Apo-P | razosin |
| TERAZOSIN | | | | |
| * Tab 1 mg | 0.50 | 28 | ✓ Arrow | |
| * Tab 2 mg | | 28 | ✓ <u>Arrow</u> | |
| * Tab 5 mg | 0.68 | 28 | ✓ <u>Arrow</u> | |
| Agents Affecting the Renin-Angiotensin System | | | | |
| | | | | |
| ACE Inhibitors | | | | |
| CAPTOPRIL | | | | |
| *‡ Oral liq 5 mg per ml | | 95 ml OP | 🖌 Capot | en |
| Oral liquid restricted to children under 12 years of age. | | | | |
| CILAZAPRIL | | | | |
| * Tab 0.5 mg | 2.00 | 90 | 🗸 Zapril | |
| * Tab 2.5 mg | 4.31 | 90 | Zapril | |
| * Tab 5 mg | 6.98 | 90 | Zapril | |
| ENALAPRIL MALEATE | | | | |
| * Tab 5 mg | 0.96 | 100 | Ethics | Enalapril |
| Ethics Enalapril to be Sole Supply on 1 October 2015 | | | 4 | |
| * Tab 10 mg | 1.24 | 100 | Ethics | Enalapril |
| Ethics Enalapril to be Sole Supply on 1 October 2015 * Tab 20 mg – For enalapril maleate oral liquid formulation re- | | | | |
| fer, page 207 | | 100 | Ethics | Enalapril |
| Ethics Enalapril to be Sole Supply on 1 October 2015 | | | | |
| LISINOPRIL | | | | |
| * Tab 5 mg | 3.58 | 90 | Arrow | -Lisinopril |
| * Tab 10 mg | 4.08 | 90 | Arrow | -Lisinopril |
| * Tab 20 mg | 4.88 | 90 | Arrow | -Lisinopril |
| PERINDOPRIL | | | | |
| * Tab 2 mg | 3.75 | 30 | | erindopril |
| * Tab 4 mg | 4.80 | 30 | ✓ <u>Apo-P</u> | erindopril |
| QUINAPRIL | | | | |
| * Tab 5 mg | 4.31 | 90 | Arrow | -Quinapril 5 |
| Arrow-Quinapril 5 to be Sole Supply on 1 October 2015 | 0.45 | 00 | | Outronall 10 |
| * Tab 10 mg Arrow-Quinapril 10 to be Sole Supply on 1 October 2015 | 3.15 | 90 | Arrow | -Quinapril 10 |
| Arrow-Quinaphi 10 to be Sole Supply on 1 October 2015 * Tab 20 mg | 5 97 | 90 | | -Quinapril 20 |
| Arrow-Quinapril 20 to be Sole Supply on 1 October 2015 | | 00 | * A110W | Samapin 20 |
| | | | | |

‡ safety cap *Three months or six months, as applicable, dispensed all-at-once

| Subsidy (Manufacturer's Price) | Subs | Fully sidised | Brand or Generic |
|-----------------------------------|------|------------------|---------------------|
| \$ | Per | ~ | Manufacturer |

TRANDOLAPRIL

| III | Higher subsidy by endorsement is available for patients who were the prior to 1 June 1998. The prescription must be endorsed accordin are "certified condition" or an appropriate description of the patt cardiac failure" or "CCF". For the purposes of this endorsement | ngly. We recommended tient such as "co t, congestive heat | end that the ongestive he art failure ind | e words used to indicate eligibility eart failure", "CHF", "congestive includes patients post myocardial |
|-----|---|--|---|--|
| | infarction with an ejection fraction of less than 40%. Patients who full subsidy by endorsement. | started on trand | Jolapril after | 1 June 1998 are not eligible for |
| * | Cap 1 mg - Higher subsidy of \$18.67 per 28 cap with En- | | | |
| - | dorsement | 3.06 | 28 | |
| | | (18.67) | | Gopten |
| * | Cap 2 mg - Higher subsidy of \$27.00 per 28 cap with En- | | | |
| | dorsement | | 28 | - |
| _ | | (27.00) | | Gopten |
| A | CE Inhibitors with Diuretics | | | |
| CIL | AZAPRIL WITH HYDROCHLOROTHIAZIDE | | | |
| * | Tab 5 mg with hydrochlorothiazide 12.5 mg | 10.72 | 100 | ✓ <u>Apo-</u> <u>Cilazapril/Hydrochlorothiazide</u> |
| EN/ | ALAPRIL MALEATE WITH HYDROCHLOROTHIAZIDE | | | |
| * | Tab 20 mg with hydrochlorothiazide 12.5 mg | 3.32 | 30 | |
| | | (8.70) | | Co-Renitec |
| QU | INAPRIL WITH HYDROCHLOROTHIAZIDE | | | |
| * | Tab 10 mg with hydrochlorothiazide 12.5 mg Accuretic 10 to be Sole Supply on 1 November 2015 | 3.65 | 30 | ✓ Accuretic 10 |
| * | Tab 20 mg with hydrochlorothiazide 12.5 mg | 4.78 | 30 | ✓ Accuretic 20 |
| | | | | |
| A | ngiotensin II Antagonists | | | |
| CA | NDESARTAN CILEXETIL – Special Authority see SA1223 below - | - Retail pharmac | :v | |
| | Tab 4 mg | | 90 | ✓ Candestar |
| | Candestar to be Sole Supply on 1 October 2015 | | | |
| * | Tab 8 mg | 3.68 | 90 | ✓ Candestar |
| | Candestar to be Sole Supply on 1 October 2015 | | | |
| * | Tab 16 mg | 6.12 | 90 | ✓ Candestar |
| .,, | Candestar to be Sole Supply on 1 October 2015 | 10.00 | ~~ | |
| * | Tab 32 mg | 10.66 | 90 | Candestar |
| | Candestar to be Sole Supply on 1 October 2015 | | | |

SA1223 Special Authority for Subsidy

Initial application — (ACE inhibitor intolerance) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has persistent ACE inhibitor induced cough that is not resolved by ACE inhibitor retrial (same or new ACE inhibitor); or
- 2 Patient has a history of angioedema.

Initial application — (Unsatisfactory response to ACE inhibitor) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is not adequately controlled on maximum tolerated dose of an ACE inhibitor.

| | Subsidy | | Fully | |
|---|------------------------------|-------|------------|---|
| | (Manufacturer's Price) \$ | Per | Subsidised | |
| | ¥ | . 01 | | |
| OSARTAN POTASSIUM | 1 55 | 04 | | ocorton Actoria |
| K Tab 12.5 mg | | 84 | | Losartan Actavis |
| ₭ Tab 25 mg | | 84 | | Losartan Actavis |
| K Tab 50 mg | | 84 | | Losartan Actavis |
| ≰ Tab 100 mg | 2.60 | 84 | V | Losartan Actavis |
| Angiotensin II Antagonists with Diuretics | | | | |
| OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE | | | | |
| Tab 50 mg with hydrochlorothiazide 12.5 mg | 2.18 | 30 | | Arrow-Losartan & Hydrochlorothiazide |
| Antiarrhythmics | | | | <u>Hydrochiorothiazide</u> |
| | | 101 | | |
| or lignocaine hydrochloride refer to NERVOUS SYSTEM, Ana MIODARONE HYDROCHLORIDE | esthetics, Local, page | 124 | | |
| Tab 100 mg – Retail pharmacy-Specialist | | 30 | • | Aratac |
| | | | | Cordarone-X |
| Tab 200 mg – Retail pharmacy-Specialist | | 30 | | Aratac |
| | | | ~ | Cordarone-X |
| Inj 50 mg per ml, 3 ml ampoule – Up to 6 inj available on | а | | | |
| PSO | | 6 | ~ | Cordarone-X |
| TROPINE SULPHATE | | | | |
| Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on | 2 | | | |
| PSO | | 50 | 1 | AstraZeneca |
| | | 50 | • | ASIIdZelleca |
| DIGOXIN | | | | |
| Tab 62.5 mcg – Up to 30 tab available on a PSO | | 240 | | Lanoxin PG |
| * Tab 250 mcg – Up to 30 tab available on a PSO | 14.52 | 240 | | Lanoxin |
| k‡ Oral liq 50 mcg per ml | | 60 ml | ~ | Lanoxin |
| DISOPYRAMIDE PHOSPHATE | | | | |
| Cap 100 mg | 15.00 | 100 | | |
| | (23.87) | 100 | | Rythmodan |
| Cap 150 mg | · · · · | 100 | | Rythmodan |
| | | | • | ., |
| EECAINIDE ACETATE – Retail pharmacy-Specialist | 00.05 | 00 | | - |
| Tab 50 mg | | 60 | ~ | Tambocor |
| Tab 100 mg – For flecainide acetate oral liquid formulation | | • - | | |
| refer, page 207 | | 60 | | Tambocor |
| Cap long-acting 100 mg | | 30 | | Tambocor CR |
| Cap long-acting 200 mg | | 30 | | Tambocor CR |
| Inj 10 mg per ml, 15 ml ampoule | | 5 | ~ | Tambocor |
| IEXILETINE HYDROCHLORIDE | | | | |
| Cap 150 mg | | 100 | ~ | Mexiletine |
| | | | | Hydrochloride |
| | | | | USP S29 |
| Cap 250 mg | | 100 | ~ | Mexiletine |
| | | | • | Hydrochloride USP §29 |
| ROPAFENONE HYDROCHLORIDE – Retail pharmacy-Speci | alist | | | |
| | | 50 | | Rytmonorm |

| | Subsidy (Manufacturer's Price) \$ | S Per | Fully ubsidised | Brand or Generic Manufacturer |
|---|---|----------|--------------------|-------------------------------------|
| Antihypotensives | | | | |
| MIDODRINE – Special Authority see SA1474 below – Retail pha | rmacy | | | |
| Tab 2.5 mg | 53.00 | 100 | 🖌 G | utron |
| Tab 5 mg | 79.00 | 100 | 🖌 G | utron |

►SA1474 Special Authority for Subsidy

Beta Adrenoceptor Blockers

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Note: Treatment should be started with small doses and titrated upwards as necessary. Hypertension should be avoided, and the usual target is a standing systolic blood pressure of 90 mm Hg.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

| ATE | ENOLOL | | | |
|------|--|---------|-----------|----------------|
| * | Tab 50 mg | 4.61 | 500 | Mylan Atenolol |
| | Mylan Atenolol to be Sole Supply on 1 October 2015 | | | |
| * | Tab 100 mg | 7.67 | 500 | Mylan Atenolol |
| | Mylan Atenolol to be Sole Supply on 1 October 2015 | | | |
| * | Oral liq 25 mg per 5 ml | 21.25 | 300 ml OP | Atenolol AFT |
| | Restricted to children under 12 years of age. | | | |
| BIS | OPROLOL FUMARATE | | | |
| | Tab 2.5 mg | 2.40 | 30 | ✓ Bosvate |
| | Tab 5 mg | 3.50 | 30 | Bosvate |
| | Tab 10 mg | 6.40 | 30 | Bosvate |
| CAI | RVEDILOL | | | |
| * | Tab 6.25 mg | | 30 | Dilatrend |
| | ······································ | 3.90 | 60 | ✓ Dicarz |
| | Dicarz to be Sole Supply on 1 September 2015 | | | |
| * | Tab 12.5 mg | 2.55 | 30 | Dilatrend |
| | - | 5.10 | 60 | Dicarz |
| | Dicarz to be Sole Supply on 1 September 2015 | | | |
| * | Tab 25 mg – For carvedilol oral liquid formulation refer, page | | | |
| | 207 | | 30 | Dilatrend |
| | | 6.30 | 60 | Dicarz |
| | Dicarz to be Sole Supply on 1 September 2015 | | | |
| • | atrend Tab 6.25 mg to be delisted 1 September 2015) | | | |
| · | atrend Tab 12.5 mg to be delisted 1 September 2015) | | | |
| (DII | atrend Tab 25 mg to be delisted 1 September 2015) | | | |
| CEI | LIPROLOL | | | |
| * | Tab 200 mg | 21.40 | 180 | Celol |
| LAE | BETALOL | | | |
| * | Tab 50 mg | 8.23 | 100 | Hybloc |
| * | Tab 100 mg – For labetalol oral liquid formulation refer, page | | | • |
| | 207 | 10.06 | 100 | Hybloc |
| * | Tab 200 mg | 17.55 | 100 | Hybloc |
| * | Inj 5 mg per ml, 20 ml ampoule | | 5 | - |
| | | (88.60) | | Trandate |
| | | | | |

| | | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|-----|---|---|--------|---------------------|-------------------------------------|
| ME | TOPROLOL SUCCINATE | | | | |
| * | Tab long-acting 23.75 mg | 0.96 | 30 | 🖌 N | letoprolol - AFT CR |
| * | Tab long-acting 47.5 mg | 1.41 | 30 | 🗸 N | letoprolol - AFT CR |
| * | Tab long-acting 95 mg | 2.42 | 30 | | letoprolol - AFT CR |
| * | Tab long-acting 190 mg | 4.66 | 30 | V N | letoprolol - AFT CR |
| ME | TOPROLOL TARTRATE | | | | |
| * | Tab 50 mg - For metoprolol tartrate oral liquid formulation | | | | |
| | refer, page 207 | | 100 | V L | opresor |
| * | Tab 100 mg | 21.00 | 60 | 🖌 L | opresor |
| * | Tab long-acting 200 mg | | 28 | 🖌 S | low-Lopresor |
| * | Inj 1 mg per ml, 5 ml vial | 24.00 | 5 | V L | opresor |
| NA | DOLOL | | | | |
| * | Tab 40 mg | | 100 | 🖌 A | po-Nadolol |
| | Apo-Nadolol to be Sole Supply on 1 November 2015 | | | | • |
| * | Tab 80 mg | 24.70 | 100 | 🗸 A | po-Nadolol |
| | Apo-Nadolol to be Sole Supply on 1 November 2015 | | | | • |
| PIN | IDOLOL | | | | |
| * | Tab 5 mg | 9.72 | 100 | 🖌 A | po-Pindolol |
| * | Tab 10 mg | | 100 | | po-Pindolol |
| * | Tab 15 mg | | 100 | VĀ | po-Pindolol |
| PR | OPRANOLOL | | | _ | |
| * | Tab 10 mg | | 100 | 🗸 A | -00 |
| | | | | | Propranolol S29 |
| | | | | | |
| * | Tab 40 mg | 4.65 | 100 | 🗸 A | po- |
| | | | | | Propranolol S29 |
| | Our lange esting 100 mg | 10.17 | 100 | | |
| * | Cap long-acting 160 mg | | 100 | v 0 | ardinol LA |
| * | Oral liq 4 mg per ml – Special Authority see SA1327 below – | | :00 ml | | overe can |
| _ | Retail pharmacy | | 600 ml | V H | oxane S29 |

SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTALOL

| * Tab 80 mg – For sotalol oral liquid formulation refer, page 207 * Tab 160 mg | | 500 100 | ✔ Mylan ✔ Mylan |
|---|-------|------------|--------------------|
| * Inj 10 mg per ml, 4 ml ampoule | | 5 | Sotacor |
| TIMOLOL * Tab 10 mg | 10.55 | 100 | 🖌 Apo-Timol |

CARDIOVASCULAR SYSTEM

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-----|---------------------|-------------------------------------|
| Calcium Channel Blockers | | | | |
| Dihydropyridine Calcium Channel Blockers | | | | |
| MLODIPINE | | | | |
| Tab 2.5 mg Tab 5 mg – For amlodipine oral liquid formulation refer, page | | 100 | V <u>I</u> | Apo-Amlodipine |
| 207 | | 250 | | Apo-Amlodipine |
| Fab 10 mg | 7.21 | 250 | <u> </u> | Apo-Amlodipine |
| ELODIPINE | | | | |
| Tab long-acting 2.5 mg | 1.45 | 30 | 🖌 F | Plendil ER |
| Plendil ER to be Sole Supply on 1 October 2015 | | | | |
| Tab long-acting 5 mg | 1.55 | 30 | 🖌 F | Plendil ER |
| Plendil ER to be Sole Supply on 1 October 2015 | 0.00 | ~~ | | |
| • Tab long-acting 10 mg | 2.30 | 30 | V | Plendil ER |
| Plendil ER to be Sole Supply on 1 October 2015 | | | | |
| RADIPINE | | | | |
| Cap long-acting 2.5 mg | | 30 | | Dynacirc-SRO |
| Cap long-acting 5 mg | | 30 | V | Dynacirc-SRO |
| FEDIPINE | | | | |
| Tab long-acting 10 mg | | 60 | | Adalat 10 |
| Tab long-acting 20 mg | | 100 | | Vyefax Retard |
| Tab long-acting 30 mg | | 30 | _ | Adefin XL |
| Tab long-acting 60 mg | 5./5 | 30 | • | Adefin XL |
| Other Calcium Channel Blockers | | | | |
| ILTIAZEM HYDROCHLORIDE | | | | |
| - Tab 30 mg | 4.60 | 100 | / [| Dilzem |
| Tab 60 mg - For diltiazem hydrochloride oral liquid formula- | | | | |
| tion refer, page 207 | | 100 | / [| Dilzem |
| Cap long-acting 120 mg | 1.91 | 30 | v (| Cardizem CD |
| | 31.83 | 500 | V | Apo-Diltiazem CD |
| Cap long-acting 180 mg | | 30 | | Cardizem CD |
| | 47.67 | 500 | | Apo-Diltiazem CD |
| Cap long-acting 240 mg | | 30 | | Cardizem CD |
| | 63.58 | 500 | V | Apo-Diltiazem CD |
| ERHEXILINE MALEATE | | | | |
| Tab 100 mg | 62.90 | 100 | 🖌 F | Pexsig |
| ERAPAMIL HYDROCHLORIDE | | | | |
| Tab 40 mg | 7.01 | 100 | ~ 1 | soptin |
| Tab 80 mg - For verapamil hydrochloride oral liquid formula- | | | | |
| tion refer, page 207 | | 100 | ~ | <u>soptin</u> |
| Tab long-acting 120 mg | 15.20 | 250 | v 1 | /erpamil SR |
| Tab long-acting 240 mg | 25.00 | 250 | • | /erpamil SR |
| Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a | | | | |
| PSO | 7.54 | 5 | ~ | soptin |

| | Subsidy (Manufacturer's Pric | (a) | Fully Subsidised | |
|--|---------------------------------|---------------|---------------------|--------------------|
| | (Manulacturer 31 no \$ | Per | | |
| Centrally-Acting Agents | | | | |
| CLONIDINE | | | | |
| Patch 2.5 mg, 100 mcg per day – Only on a prescription | 12.80 | 4 | ~ | Catapres-TTS-1 |
| Patch 5 mg, 200 mcg per day – Only on a prescription | | 4 | | Catapres-TTS-2 |
| Patch 7.5 mg, 300 mcg per day – Only on a prescription | 22.68 | 4 | | Catapres-TTS-3 |
| LONIDINE HYDROCHLORIDE | | | | |
| Fab 25 mcg | 10.53 | 112 | | Clonidine BNM |
| Clonidine BNM to be Sole Supply on 1 October 2015 | | 400 | | . . |
| Tab 150 mcg Ini 150 mcg per ml. 1 ml ampoule | | 100 | | Catapres |
|) · · · · · · · · · | | 5 | V | Catapres |
| ETHYLDOPA | 44.05 | 400 | | |
| • Tab 125 mg | | 100 | | Prodopa |
| Tab 250 mg Tab 500 mg | | 100 100 | | Prodopa Prodopa |
| Ŷ | 23.15 | 100 | | Ргодора |
| Diuretics | | | | |
| Loop Diuretics | | | | |
| UMETANIDE | | | | |
| 🗧 Tab 1 mg | 16.36 | 100 | ~ | Burinex |
| Inj 500 mcg per ml, 4 ml vial | 7.95 | 5 | ~ | Burinex |
| UROSEMIDE [FRUSEMIDE] | | | | |
| Tab 40 mg – Up to 30 tab available on a PSO | 8.00 | 1,000 | ~ | Diurin 40 |
| Diurin 40 to be Sole Supply on 1 October 2015 | | | | |
| Tab 500 mg | 25.00 | 50 | | Urex Forte |
| Urex Forte to be Sole Supply on 1 October 2015 | 10.00 | | | |
| toral liq 10 mg per ml | | 30 ml OF 5 | | Lasix Lasix |
| Inj 10 mg per ml, 25 ml ampoule Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a | | 5 | V | Lasix |
| Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO | | 5 | | Frusemide-Claris |
| | | 5 | • | |
| Potassium Sparing Diuretics | | | | |
| MILORIDE HYDROCHLORIDE | | | | |
| Tab 5 mg | | 100 | | Apo-Amiloride |
| Oral liq 1 mg per ml | | 25 ml OF | , / | Biomed |
| ETOLAZONE – Special Authority see SA1349 below – Retail | oharmacy | | | |
| Tab 5 mg | CBS | 1 | ~ | Metolazone S29 |
| | | 50 | v : | Zaroxolyn S29 |
| SA1349 Special Authority for Subsidy | | | | |

SA1349 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where used for the treatment of patients with refractory heart failure who are intolerant or have not responded to loop diuretics and/or loop-thiazide combination therapy.

SPIRONOLACTONE

| * | Tab 25 mg | 100 | Spiractin |
|---|----------------------|----------|-------------------------------|
| * | Tab 100 mg11.80 | 100 | Spiractin |
| ‡ | Oral liq 5 mg per ml | 25 ml OP | Biomed |

| | Subsidy (Manufacturer's Price \$ | e) Sub: Per | Fully Brand or sidised Generic Manufacturer |
|--|--|----------------|---|
| Potassium Sparing Combination Diuretics | Ų | | • Wanuacturer |
| | | | |
| AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE * Tab 5 mg with furosemide 40 mg | | 28 | 🗸 Frumil |
| AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZI | | | |
| * Tab 5 mg with hydrochlorothiazide 50 mg | | 50 | ✓ Moduretic |
| Thiazide and Related Diuretics | | | |
| BENDROFLUMETHIAZIDE [BENDROFLUAZIDE] | | | |
| * Tab 2.5 mg – Up to 150 tab available on a PSO | 5.48 | 500 | Arrow- Bendrofluazide |
| May be supplied on a PSO for reasons other than emerge | ency. | | |
| * Tab 5 mg | 8.95 | 500 | <u>Arrow-</u> <u>Bendrofluazide</u> |
| CHLOROTHIAZIDE | | | |
| Oral liq 50 mg per ml | | 25 ml OP | Biomed |
| CHLORTALIDONE [CHLORTHALIDONE] | 0.00 | | |
| * Tab 25 mg | 8.00 | 50 | Hygroton |
| INDAPAMIDE * Tab 2.5 mg | 2.25 | 90 | 🗸 Dapa-Tabs |
| Lipid-Modifying Agents | | | |
| Fibrates | | | |
| BEZAFIBRATE | | | |
| * Tab 200 mg | 9.05 | 90 | ✓ Bezalip |
| Bezalip to be Sole Supply on 1 November 2015 | 0.70 | 00 | A Denalia Deterri |
| Tab long-acting 400 mg Bezalip Retard to be Sole Supply on 1 November 2015 | 6.78 | 30 | Bezalip Retard |
| GEMFIBROZIL | | | |
| * Tab 600 mg | | 60 | ✓ Lipazil |
| Other Lipid-Modifying Agents | | | |
| ACIPIMOX | | | |
| * Cap 250 mg | | 30 | Olbetam |
| NICOTINIC ACID | | | |
| * Tab 50 mg | | 100 | ✓ <u>Apo-Nicotinic Acid</u> |
| * Tab 500 mg | | 100 | Apo-Nicotinic Acid |
| Resins | | | |
| CHOLESTYRAMINE | | | |
| Powder for oral liq 4 g | | 50 | |
| | (52.68) | | Questran-Lite |
| COLESTIPOL HYDROCHLORIDE | | | 4 a b b b b |
| Grans for oral liq 5 g | | 30 | Colestid |
| | | | |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|----------|---------------------|-------------------------------------|
| HMG CoA Reductase Inhibitors (Statins) | | | | |
| Prescribing Guidelines | | | | |
| Freatment with HMG CoA Reductase Inhibitors (statins) is recor cardiovascular risk of 15% or greater. | nmended for patients | with | dyslipidaen | nia and an absolute 5 y |
| ATORVASTATIN – See prescribing guideline above | | | | |
| * Tab 10 mg | 0.84 | 30 | V L | ipitor |
| | | | | fizer atorvastatin |
| | 2.52 | 90 | 🗸 Z | arator |
| * Tab 20 mg | 1.39 | 30 | V L | ipitor |
| | | | 🖌 F | fizer atorvastatin |
| | 4.17 | 90 | 🖌 Z | arator |
| * Tab 40 mg | 2.44 | 30 | | ipitor |
| | | | | fizer atorvastatin |
| | 7.32 | 90 | | arator |
| * Tab 80 mg | 5.41 | 30 | | ipitor |
| | 16.23 | 90 | | fizer atorvastatin Zarator |
| (Lipitor Tab 20 mg to be delisted 1 November 2015) (Pfizer atorvastatin Tab 20 mg to be delisted 1 November 2015) (Lipitor Tab 40 mg to be delisted 1 November 2015) (Pfizer atorvastatin Tab 40 mg to be delisted 1 November 2015) (Lipitor Tab 80 mg to be delisted 1 November 2015) (Pfizer atorvastatin Tab 80 mg to be delisted 1 November 2015) | | | | |
| PRAVASTATIN – See prescribing guideline above | | | | |
| * Tab 20 mg | 3.45 | 30 | . . | Cholvastin |
| * Tab 20 mg | | 30 | | Cholvastin |
| · ···· · · · · · · · · · · · · · · · · | | | • • | |
| SIMVASTATIN – See prescribing guideline above Tab 10 mg | 0.95 | 90 | <u> </u> | rrow-Simva 10mg |
| ★ Tab 10 mg | | 90 90 | _ | Arrow-Simva 20mg |
| ★ Tab 20 mg | | 90 90 | | Arrow-Simva 40mg |
| ★ Tab 40 mg | | 90 | | Arrow-Simva 80mg |
| • | | 00 | • • | arow onlive comy |
| Selective Cholesterol Absorption Inhibitors | | | | |
| EZETIMIBE – Special Authority see SA1045 below – Retail phar Tab 10 mg | | 30 | | zemibe zetrol |
| Ezemibe to be Sole Supply on 1 November 2015 (Ezetrol Tab 10 mg to be delisted 1 November 2015) | | | | |

(Ezetrol Tab 10 mg to be delisted 1 November 2015)

➡SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and

- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:

continued...

57

CARDIOVASCULAR SYSTEM

| Subsidy (Manufacturer's Price) | Suk | Fully | Brand or Generic | |
|-----------------------------------|-----|-------|---------------------|--|
| (ivialiulaciulei s Flice) \$ | Per | | Manufacturer | |

continued...

- 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 $\times\,$ normal) when treated with one statin; or
- 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
- 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

| EZETIMIBE WITH SIMVASTATIN - Special Authority see | SA1046 below - Retail pl | narmacy | |
|--|--------------------------|---------|----------------------------|
| Tab 10 mg with simvastatin 10 mg | 5.15 | 30 | Zimybe |
| | (36.68) | | Vytorin |
| Zimybe to be Sole Supply on 1 November 2015 | | | · |
| Tab 10 mg with simvastatin 20 mg | 6.15 | 30 | Zimybe |
| | (38.70) | | Vytorin |
| Zimybe to be Sole Supply on 1 November 2015 | | | · |
| Tab 10 mg with simvastatin 40 mg | 7.15 | 30 | Zimybe |
| | (41.40) | | Vytorin |
| Zimybe to be Sole Supply on 1 November 2015 | | | |
| Tab 10 mg with simvastatin 80 mg | 8.15 | 30 | Zimybe |
| | (45.45) | | Vytorin |
| Zimybe to be Sole Supply on 1 November 2015 | | | - |
| (Vytorin Tab 10 mg with simvastatin 10 mg to be delisted 1 | November 2015) | | |
| (Vytorin Tab 10 mg with simvastatin 20 mg to be delisted 1 | November 2015) | | |
| (Vytorin Tab 10 mg with simvastatin 40 mg to be delisted 1 | November 2015) | | |
| (Vytorin Tab 10 mg with simvastatin 80 mg to be delisted 1 | November 2015) | | |

➡SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to ≤ 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

| | Subsidy | | Fully Brand or |
|---|-----------------------|--------------------|---------------------------------------|
| | (Manufacturer's \$ | Price) Sub: Per | sidised Generic Manufacturer |
| | φ | Fei | |
| Nitrates | | | |
| LYCERYL TRINITRATE | | | |
| Tab 600 mcg – Up to 100 tab available on a PSO | 8.00 | 100 OP | Lycinate |
| Oral pump spray, 400 mcg per dose – Up to 250 dose avail- | | | |
| able on a PSO | 4.45 | 250 dose OP | Nitrolingual Pump |
| | | | Spray |
| • Oral spray, 400 mcg per dose – Up to 250 dose available on | | | |
| a PSO | | 250 dose OP | Glytrin |
| Patch 25 mg, 5 mg per day | | 30 | ✓ <u>Nitroderm TTS</u> |
| Patch 50 mg, 10 mg per day | 18.62 | 30 | Nitroderm TTS |
| SOSORBIDE MONONITRATE | | | |
| F Tab 20 mg | 17.10 | 100 | ✓ Ismo 20 |
| Tab long-acting 40 mg | 7.50 | 30 | Ismo 40 Retard |
| Tab long-acting 60 mg | | 90 | Duride |
| Sympathomimetics | | | |
| | | | |
| DRENALINE Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PSO | 4 98 | 5 | Aspen Adrenaline |
| | | 5 | ✓ Hospira |
| Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a | 0.20 | | • noopnu |
| PSO | 27.00 | 5 | ✓ Hospira |
| 1.00 | 49.00 | 10 | Aspen Adrenaline |
| | 40.00 | 10 | |
| SOPRENALINE Inj 200 mcg per ml, 1 ml ampoule | 00.00 | 25 | |
| f inj 200 mcg per mi, i mi ampoue | | 20 | Isuprel |
| | (104.20) | | Isuprei |
| Vasodilators | | | |
| MYL NITRITE | | | |
| E Liq 98% in 0.3 ml cap | 62.92 | 12 | |
| | (73.40) | | Baxter |
| YDRALAZINE HYDROCHLORIDE | | | |
| Tab 25 mg - Special Authority see SA1321 below - Retail | | | |
| pharmacy | CBS | 1 | Hydralazine |
| P | | 56 | ✓ Onelink S29 |
| Inj 20 mg ampoule | 25 90 | 5 | ✓ Apresoline |
| | 20.00 | 5 | • Apresolitie |
| ⇒SA1321 Special Authority for Subsidy aitial application from any relevant practitioner. Approvals valid the following criteria: ither: | without furthe | er renewal unless | s notified for applications mee |
| For the treatment of refractory hypertension; or For the treatment of heart failure in combination with a nitra inhibitors and/or angiotensin receptor blockers. | ate, in patients | s who are intolera | ant or have not responded to <i>i</i> |
| INOXIDIL – Special Authority see SA1271 below – Retail pharm. Tab 10 mg | | 100 | ✔ Loniten |
| • | | 100 | |
| SA1271 Special Authority for Subsidy itial application only from a relevant specialist. Approvals valid v fractory hypertension which has failed to respond to extensive multiple application of the second secon | | | notified where patient has se |

| | Subsidy (Manufacturer's Price) \$ | Per | Full Subsidise | d Generic |
|---|--|----------------------|---|--|
| NICORANDIL | | | | |
| Tab 10 mg | 27.95 | 60 | • | lkorel |
| ▲ Tab 20 mg | | 60 | ~ | lkorel |
| PAPAVERINE HYDROCHLORIDE | | | | |
| Inj 12 mg per ml, 10 ml ampoule | 217.90 | 5 | ~ | Hospira |
| PENTOXIFYLLINE [OXPENTIFYLLINE] | | | | |
| Tab 400 mg | | 50 | | |
| | (42.26) | | | Trental 400 |
| Endothelin Receptor Antagonists | | | | |
| SA0967 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertension | n Panel | | | |
| SA0967 Special Authority for Subsidy | site http://www.phar | mac.ç | <u>jovt.nz</u> or | |
| ⇒SA0967 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertension lotes: Application details may be obtained from PHARMAC's web The Coordinator, PAH Panel PHARMAC, PO Box 10-254, WELLINGTON | site <u>http://www.phar</u> | mac.ç | <u>jovt.nz</u> or | |
| SA0967 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertension Votes: Application details may be obtained from PHARMAC's web I'he Coordinator, PAH Panel PHARMAC, PO Box 10-254, WELLINGTON Fel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.go AMBRISENTAN – Special Authority see SA0967 above – Retail pl Tab 5 mg | site <u>http://www.phar</u> wt.nz harmacy 4,585.00 | 30 | V | Volibris |
| ⇒SA0967 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertension Votes: Application details may be obtained from PHARMAC's web I'he Coordinator, PAH Panel PHARMAC, PO Box 10-254, WELLINGTON Fel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.go AMBRISENTAN – Special Authority see SA0967 above – Retail pl | site <u>http://www.phar</u> wt.nz harmacy 4,585.00 | | V | |
| ⇒SA0967 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertension Notes: Application details may be obtained from PHARMAC's web The Coordinator, PAH Panel PHARMAC, PO Box 10-254, WELLINGTON Fel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.go AMBRISENTAN – Special Authority see SA0967 above – Retail pl Tab 5 mg Tab 10 mg | site <u>http://www.phar</u> <u>wt.nz</u> harmacy 4,585.00 4,585.00 nacy | 30 30 | ~ ~ | Volibris Volibris |
| ►SA0967 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertension Jotes: Application details may be obtained from PHARMAC's web The Coordinator, PAH Panel PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.go AMBRISENTAN – Special Authority see SA0967 above – Retail pl Tab 5 mg Tab 10 mg | site <u>http://www.phar</u> harmacy 4,585.00 4,585.00 nacy 375.00 | 30 30 56 | | Volibris Volibris Mylan-Bosentan |
| ►SA0967 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertension Jotes: Application details may be obtained from PHARMAC's web The Coordinator, PAH Panel PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.go AMBRISENTAN – Special Authority see SA0967 above – Retail pl Tab 5 mg Tab 10 mg | site <u>http://www.phar</u> harmacy 4,585.00 4,585.00 nacy 375.00 1,500.00 | 30 30 | | Volibris Volibris Mylan-Bosentan pms-Bosentan |
| SA0967 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertension lotes: Application details may be obtained from PHARMAC's web The Coordinator, PAH Panel PHARMAC, PO Box 10-254, WELLINGTON el: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.go MBRISENTAN – Special Authority see SA0967 above – Retail pl Tab 5 mg Tab 10 mg BOSENTAN – Special Authority see SA0967 above – Retail pharm Tab 62.5 mg | site <u>http://www.phar</u> harmacy 4,585.00 4,585.00 nacy 375.00 1,500.00 4,585.00 | 30 30 56 60 | ~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~ | Volibris Volibris Mylan-Bosentan pms-Bosentan Tracleer |
| SA0967 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertension lotes: Application details may be obtained from PHARMAC's web 'he Coordinator, PAH Panel PHARMAC, PO Box 10-254, WELLINGTON el: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.go MBRISENTAN – Special Authority see SA0967 above – Retail pl Tab 5 mg Tab 10 mg | site <u>http://www.phar</u> harmacy 4,585.00 4,585.00 nacy 375.00 1,500.00 4,585.00 | 30 30 56 | ~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~ | Volibris Volibris Mylan-Bosentan pms-Bosentan |

Phosphodiesterase Type 5 Inhibitors

SA1293 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon* - for Pulmonary Arterial Hypertension see note below) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Notes: Sildenafil is also funded for patients with Pulmonary Arterial Hypertension who are approved by the Pulmonary Arterial Hypertension Panel (an application must be made using form <u>SA1293-PAH</u>).

Application details may be obtained from:

The Coordinator, PAH Panel

PHARMAC, PO Box 10 254, Wellington

Phone: (04) 916 7561 Facsimile: (04) 974 4858 Email: PAH@pharmac.govt.nz

Indications marked with * are Unapproved Indications.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|-----|---------------------|-------------------------------------|
| | age – Retail pharmacy | | | |
| Tab 25 mg | 0.75 | 4 | 🖌 V | edafil |
| , | 1.85 | | V S | ilagra |
| Tab 50 mg | 0.75 | 4 | 🖌 V | edafil |
| - | 1.85 | | 🖌 S | ilagra |
| Tab 100 mg – For sildenafil oral liquid formulation refer, page | 9 | | | - |
| 207 | 2.75 | 4 | 🗸 V | edafil |
| | 7.45 | | 🖌 S | ilagra |
| | | | | - |
| Prostacyclin Analogues | | | | |

| SA0969 Special Authority for Subsidy | | |
|--|------------|------------------------------|
| Special Authority approved by the Pulmonary Arterial Hypertension Panel | | |
| Notes: Application details may be obtained from PHARMAC's website http://www.pha | rmac.govt. | nz or: |
| The Coordinator, PAH Panel | | |
| PHARMAC, PO Box 10-254, WELLINGTON | | |
| Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz | | |
| ILOPROST - Special Authority see SA0969 above - Retail pharmacy | | |
| Nebuliser soln 10 mcg per ml, 2 ml1,185.00 | 30 | Ventavis |

| (| Subsidy Manufacturer's Price | _ | Fully | Brand or Generic |
|--|---------------------------------|---------|------------|---------------------|
| | \$ | Per | <u> </u> | Manufacturer |
| Antiacne Preparations | | | | |
| For systemic antibacterials, refer to INFECTIONS, Antibacterials, pa | age 91 | | | |
| ADAPALENE | | | | |
| a) Maximum of 30 g per prescription | | | | |
| b) Only on a prescription | | | | |
| Ćrm 0.1% | 22.89 3 | 80 g OP | 🖌 Di | ifferin |
| Gel 0.1% | 22.89 3 | 80 g OP | 🖌 Di | ifferin |
| SOTRETINOIN - Special Authority see SA1475 below - Retail ph | armacv | | | |
| Cap 10 mg | , | 120 | v 0 | ratane |
| Cap 20 mg | | 120 | ✔0 | ratane |

➡SA1475 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
 - 3.2 Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
- 2 Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

TRETINOIN

| Crm 0.5 mg per g | Maximum of 50 g per prescription | 13.90 | 50 g OP | ReTrieve |
|------------------|--|-------|---------|----------|
|------------------|--|-------|---------|----------|

| | Subsidy (Manufacturer's \$ | Price) Sul Per | Fully Brand or osidised Generic ✓ Manufacturer |
|---|------------------------------------|-------------------|--|
| Antibacterials Topical | | | |
| For systemic antibacterials, refer to INFECTIONS, Antibacterial | s, page 91 | | |
| FUSIDIC ACID | | | |
| Crm 2% | 2.52 | 15 g OP | ✓ DP Fusidic Acid |
| a) Maximum of 15 g per prescription b) Only on a prescription c) Not in combination | | | <u>Cream</u> |
| Oint 2% a) Maximum of 15 g per prescription b) Only on a prescription c) Not in combination | 3.45 | 15 g OP | ✔ <u>Foban</u> |
| | 0.50 | 45 00 | |
| * Crm 1% | 8.56 | 15 g OP | Crystaderm |
| MUPIROCIN Oint 2% | 6.60 (9.26) | 15 g OP | Bactroban |
| a) Only on a prescriptionb) Not in combination | () | | |
| SILVER SULPHADIAZINE | | | |
| Crm 1%a) Up to 250 g available on a PSO b) Not in combination | 12.30 | 50 g OP | Flamazine |
| Antifungals Topical | | | |
| For systemic antifungals, refer to INFECTIONS, Antifungals, pa | ge 98 | | |
| AMOROLFINE | 0 | | |
| a) Only on a prescription | | | |
| b) Not in combination | | | |
| Nail soln 5% | 19.95 | 5 ml OP | ✓ MycoNail |
| CICLOPIROX OLAMINE a) Only on a prescription b) Not in combination | | | |
| Nail-soln 8% Apo-Ciclopirox to be Sole Supply on 1 October 2015 | 6.50 | 7 ml OP | Apo-Ciclopirox |
| CLOTRIMAZOLE | | | |
| Crm 1%a) Only on a prescription b) Not in combination | 0.52 | 20 g OP | ✓ <u>Clomazol</u> |
| * Soln 1% | 4.36 | 20 ml OP | |
| a) Only on a prescription b) Not in combination | (7.55) | | Canesten |

| | Subsidy (Manufacturer's F | Price) Sul | Fully Brand or bsidised Generic |
|---|------------------------------|-----------------|------------------------------------|
| | (Manulactule) S P | Per | Manufacturer |
| CONAZOLE NITRATE | | | |
| Crm 1% | 1.00 | 20 g OP | |
| | (7.48) | | Pevaryl |
| a) Only on a prescription | | | |
| b) Not in combination | | | |
| Foaming soln 1%, 10 ml sachets | | 3 | Devend |
| a) Only on a pressription | (17.23) | | Pevaryl |
| a) Only on a prescription | | | |
| b) Not in combination | | | |
| | 0.55 | 45 | |
| ₭ Crm 2% | 0.55 | 15 g OP | Multichem |
| a) Only on a prescription b) Not in combination | | | |
| € Lotn 2% | 1 36 | 30 ml OP | |
| | (10.03) | 50 111 01 | Daktarin |
| a) Only on a prescription | (10.00) | | Danarin |
| b) Not in combination | | | |
| ← Tinct 2% | 4.36 | 30 ml OP | |
| | (12.10) | | Daktarin |
| a) Only on a prescription | () | | |
| b) Not in combination | | | |
| IYSTATIN | | | |
| Crm 100,000 u per g | 1.00 | 15 g OP | |
| | (7.90) | • | Mycostatin |
| a) Only on a prescription | | | |
| b) Not in combination | | | |
| Antipruritic Preparations | | | |
| CALAMINE | | | |
| a) Only on a prescription | | | |
| b) Not in combination | | | |
| Crm, aqueous, BP | 1.77 | 100 g | Pharmacy Health |
| Lotn, BP | | 2,000 ml | ✔ PSM |
| ROTAMITON | | · | |
| a) Only on a prescription | | | |
| b) Not in combination | | | |
| Crm 10% | 3.37 | 20 g OP | Itch-Soothe |
| Itch-Soothe to be Sole Supply on 1 October 2015 | | 0 | |
| IENTHOL – Only in combination | | | |
| Only in combination with a dermatological base or pro | oprietary Topical C | orticosteriod - | - Plain, refer dermatological ba |
| page 206 | | | , |
| 2) With or without other dermatological galenicals. | | | |
| Crystals | 6.50 | 25 g | 🖌 PSM |
| , - | 6.92 | - 9 | ✓ MidWest |
| | | | |

| | Subsidy (Manufacturer's F \$ | rice) Sub Per | Fully Brand or osidised Generic ✓ Manufacturer |
|---|------------------------------------|------------------|--|
| | * | | |
| Corticosteroids Topical | | | |
| or systemic corticosteroids, refer to CORTICOSTEROIDS AND F | RELATED AGEN | TS, page 79 | |
| Corticosteroids - Plain | | | |
| BETAMETHASONE DIPROPIONATE | | | |
| Crm 0.05% | 2.96 | 15 g OP | Diprosone |
| | 8.97 | 50 g OP | ✓ Diprosone |
| Crm 0.05% in propylene glycol base | 4.33 | 30 g OP | Diprosone OV |
| Oint 0.05% | | 15 g OP | Diprosone |
| | 8.97 | 50 g OP | Diprosone |
| Oint 0.05% in propylene glycol base | 4.33 | 30 g OP | Diprosone OV |
| BETAMETHASONE VALERATE | | | |
| k Crm 0.1% | 3.15 | 50 g OP | 🖌 Beta Cream |
| ₭ Oint 0.1% | | 50 g OP | Beta Ointment |
| ₭ Lotn 0.1% | | 50 ml OP | Betnovate |
| CLOBETASOL PROPIONATE | | | |
| € Crm 0.05% | 3 20 | 30 g OP | Clobetasol BNM |
| CIII 0.0570 | | 30 g OI | ✓ Dermol |
| Clobetasol BNM to be Sole Supply on 1 October 2015 | | | • Dermor |
| Cont 0.05% | 3 20 | 30 g OP | Clobetasol BNM |
| | | 00 g 0. | ✓ Dermol |
| Clobetasol BNM to be Sole Supply on 1 October 2015 | | | |
| Dermol Crm 0.05% to be delisted 1 October 2015) | | | |
| Dermol Oint 0.05% to be delisted 1 October 2015) | | | |
| | | | |
| Crm 0.05% | 5 29 | 30 g OP | |
| CIII 0.05% | (7.09) | SU Y OF | Eumovate |
| | 16.13 | 100 g OP | Editiovale |
| | (22.00) | 100 g OI | Eumovate |
| | () | | Lanorato |
| DIFLUCORTOLONE VALERATE | 0.07 | 50 × 00 | |
| Crm 0.1% | | 50 g OP | Naviana |
| Fatty oint 0.1% | (15.86) | 50 ~ OD | Nerisone |
| rally uill U. 1% | | 50 g OP | Noricono |
| | (15.86) | | Nerisone |
| IYDROCORTISONE | | | 4 |
| Crm 1% – Only on a prescription | | 100 g | Pharmacy Health |
| | 14.00 | 500 g | Pharmacy Health |
| ✤ Powder – Only in combination | | 25 g | ✓ <u>ABM</u> |
| Up to 5% in a dermatological base (not proprietary Topi | cal Corticosterio | a – Plain) wit | n or without other dermatologic |
| galenicals. Refer, page 206 | | | |
| HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN | | | |
| Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only | | | |
| on a prescription | 10.57 | 250 ml | ✓ DP Lotn HC |

| | Subsidy (Manufacturer's | Prico) S | Full | |
|---|----------------------------|-----------------|----------------|------------------|
| | (Manulacturers) \$ | Price) S Per | ubsidised V | Manufacturer |
| IYDROCORTISONE BUTYRATE | | | | |
| Lipocream 0.1% | 2.30 | 30 g OP | V | Locoid Lipocream |
| | 6.85 | 100 g OP | | Locoid Lipocream |
| Oint 0.1% | 6.85 | 100 g OP | | Locoid |
| Milky emul 0.1% | 6.85 | 100 ml OP | V | Locoid Crelo |
| AETHYLPREDNISOLONE ACEPONATE | | | | |
| Crm 0.1% | 4.95 | 15 g OP | V | Advantan |
| Oint 0.1% | | 15 g OP | V | Advantan |
| IOMETASONE FUROATE | | Ū | | |
| Crm 0.1% | 1 78 | 15 g OP | ~ | m-Mometasone |
| | 3.42 | 45 g OP | | m-Mometasone |
| Oint 0.1% | ••••= | 15 g OP | | m-Mometasone |
| | 3.42 | 45 g OP | | m-Mometasone |
| Lotn 0.1% | 7.35 | 30 ml OP | V | Elocon |
| Elocon to be Sole Supply on 1 October 2015 | | | | |
| RIAMCINOLONE ACETONIDE | | | | |
| Crm 0.02% | 6.30 | 100 g OP | V | Aristocort |
| Oint 0.02% | | 100 g OP | | Aristocort |
| Corticosteroids - Combination | | | | |
| Controsteroius - Combination | | | | |
| BETAMETHASONE VALERATE WITH CLIOQUINOL - Only on a | prescription | | | |
| Crm 0.1% with clioquinol 3% | | 15 g OP | | |
| | (4.90) | Ũ | | Betnovate-C |
| ETAMETHASONE VALERATE WITH FUSIDIC ACID | | | | |
| Crm 0.1% with fusidic acid 2% | | 15 g OP | | |
| | (10.45) | | | Fucicort |
| a) Maximum of 15 g per prescription | (/ | | | |
| b) Only on a prescription | | | | |
| YDROCORTISONE WITH MICONAZOLE - Only on a prescription | on | | | |
| Crm 1% with miconazole nitrate 2% | | 15 g OP | ~ | Micreme H |
| Micreme H to be Sole Supply on 1 October 2015 | | - 5 - | | |
| YDROCORTISONE WITH NATAMYCIN AND NEOMYCIN - Onl | v on a prescrip | tion | | |
| Crm 1% with natamycin 1% and neomycin sulphate 0.5% | , , , | 15 g OP | V | Pimafucort |
| Oint 1% with natamycin 1% and neomycin sulphate 0.5% | | 15 g OP | ~ | Pimafucort |
| RIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCIN | AND NYSTAT | IN - | | |
| Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg | | | | |
| and gramicidin 250 mcg per g – Only on a prescription | 3 49 | 15 g OP | | |
| | (6.60) | 10 9 01 | | Viaderm KC |
| Disinfasting and Oleansing America | (0.00) | | | |
| Disinfecting and Cleansing Agents | | | | |
| HLORHEXIDINE GLUCONATE – Subsidy by endorsement | | | | |
| a) No more than 500 ml per month | | | | |
| b) Only if prescribed for a dialysis patient and the prescription | is endorsed ac | cordinaly. | | |
| Handrub 1% with ethanol 70% | | 500 ml | V | healthE |
| healthE to be Sole Supply on 1 October 2015 | | | • | |
| | | | | |
| ₭ Soln 4% wash | 3.98 | 500 ml | ~ | healthE |

| | Subsidy (Manufacturer's \$ | Price) Sub Per | Fully Brand or sidised Generic Manufacturer |
|--|----------------------------------|-------------------|--|
| RICLOSAN – Subsidy by endorsement a) Maximum of 500 ml per prescription b) | | | |
| a) Only if prescribed for a patient identified with Methicillin in hospital and the prescription is endorsed accordingly | ; or | | · // · |
| b) Only if prescribed for a patient with recurrent Staphyloco | | • | |
| Soln 1% | 4.50 5.90 | 500 ml OP | Pharmacy Health healthE |
| Barrier Creams and Emollients | | | |
| Barrier Creams | | | |
| | 4 70 | | |
| Crm 5% pump bottle | 4.73 | 500 ml OP | <u>healthE</u> <u>Dimethicone 5%</u> |
| INC AND CASTOR OIL ₭ Oint BP | | 500 g | ✓ Multichem |
| Emollients | | 0 | |
| QUEOUS CREAM | | | 4 · |
| ₭ Crm | 1.96 | 500 g | 🗸 AFT |
| ETOMACROGOL € Crm BP | 2 15 | 500 a | ✔ PSM |
| | | 500 g | ♥ F3W |
| ETOMACROGOL WITH GLYCEROL Crm 90% with glycerol 10% | 4.50 | 500 ml OP | Pharmacy Health |
| | | | Sorbolene with |
| | | | Glycerin |
| | 6.50 | 1,000 ml OP | Pharmacy Health Sorbolene with |
| | | | Glycerin |
| MULSIFYING OINTMENT | | | • |
| Gint BP | 2.73 | 500 g | ✓ <u>AFT</u> |
| IL IN WATER EMULSION | | | |
| € Crm | 2.63 | 500 g | healthE Fatty Cream |
| JREA | 4.05 | 400 00 | |
| Crm 10% | 1.65 | 100 g OP | healthE Urea Cream |
| /OOL FAT WITH MINERAL OIL − Only on a prescription Lotn hydrous 3% with mineral oil | 1.40 | 250 ml OP | |
| | (4.53) | 250 111 01 | DP Lotion |
| | 5.60 | 1,000 ml | |
| | (11.95) | | DP Lotion |
| | (20.53) 1.40 | 250 ml OP | Alpha-Keri Lotion |
| | (7.73) | 230 mi OF | BK Lotion |
| | 5.60 | 1,000 ml | |
| | (23.91) | | BK Lotion |

| | Subsidy (Manufacturer's P \$ | Price) Sul Per | Fully Brand or osidised Generic Manufacturer |
|--|------------------------------------|-------------------|--|
| Other Dermatological Bases | | | |
| PARAFFIN | | | |
| White soft – Only in combination | | 500 g | |
| | (7.78) | | IPW |
| | 20.20 | 2,500 g | 🖌 IPW |
| | 3.58 | 500 g | DOM |
| Only in combination with a demostal sized selection or | (8.69) | | PSM PSM |
| Only in combination with a dermatological galenical or as | s a diluent for a pro | prietary topic | al Corticosteroid – Plain. |
| Minor Skin Infections | | | |
| POVIDONE IODINE | | | |
| Oint 10% | 3.27 | 25 g OP | ✓ Betadine |
| a) Maximum of 100 g per prescription | | 0 | |
| b) Only on a prescription | | | |
| Antiseptic soln 10% | 0.19 | 15 ml | |
| | (4.45) | | Betadine |
| | 1.28 | 100 ml | |
| | (8.25) | | Betadine |
| | 6.20 | 500 ml | Betadine |
| | 1.28 | 100 ml | |
| | (4.20) | | Riodine |
| | 6.20 | 500 ml | Riodine |
| Skin preparation, povidone iodine 10% with 30% alcohol | 1.63 | 100 ml | |
| | (3.65) | | Betadine Skin Prep |
| | 10.00 | 500 ml | Betadine Skin Prep |
| Skin preparation, povidone iodine 10% with 70% alcohol | | 100 ml | |
| | (6.04) | | Orion |
| | 8.13 | 500 ml | |
| | (18.63) | | Orion |
| Parasiticidal Preparations | | | |
| AMMA BENZENE HEXACHLORIDE | | | |
| GAMMA BENZENE HEXACHLORIDE Crm 1% | 2 50 | | Benhex |
| | 3.50 | 50 g OP | V Dennex |
| Benhex Crm 1% to be delisted 1 January 2016) | | | |
| VERMECTIN – Special Authority see SA1225 on the next page | | | 4 |
| Tab 3 mg – Up to 100 tab available on a PSO | | 4 | ✓ Stromectol |
| 1) PSO for institutional use only. Must be endorsed with the | ne name of the ins | titution for whi | ch the PSO is required and a va |
| Special Authority for patient of that institution. | | | a the star of the star of the the st |

2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.

 For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or penal institutions.

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | S | ubsidised | Generic | |
| `\$´´ | Per | ~ | Manufacturer | |

➡SA1225 Special Authority for Subsidy

Initial application — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:

continued...

| | Fully Subsidised | Brand or Generic | |
|-----------|---------------------|---------------------|--|
| \$ Per | ~ | Manufacturer | |

continued...

- 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
- 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.
- Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria: Any of the following:

- 1 Filovicialess ev
- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

MALATHION WITH PERMETHRIN AND PIPERONYL BUTOXIDE

| Spray 0.25% with permethrin 0.5% and piperonyl butoxide 2%11.15 | 90 g OP | 🖌 Para Plus | |
|---|----------|--------------------|--|
| PERMETHRIN | | | |
| Crm 5%4.20 | 30 g OP | Lyderm | |
| Lotn 5% | 30 ml OP | ✓ <u>A-Scabies</u> | |
| Desvissis and Essense Drenarations | | | |

Psoriasis and Eczema Preparations

| ACITRETIN – Special Authority see SA1476 below – Retail pharmacy | |
|--|----|
| Cap 10 mg 17.86 | 60 |

| Cap 10 mg | 60 | Novatretin |
|-----------|--------|--------------|
| Cap 25 mg | 60 | ✓ Novatretin |

➡SA1476 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is female and has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of two years after the completion of the treatment; or
 - 3.2 Patient is male.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is female and has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of two years after the completion of the treatment; or
- 2 Patient is male.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

| Gel 500 mcg with calcipotriol 50 mcg per g26.12 | 30 g OP | Daivobet |
|---|---------|----------|
| Daivobet to be Sole Supply on 1 October 2015 | | |
| Oint 500 mcg with calcipotriol 50 mcg per g | 30 g OP | Daivobet |
| Daivobet to be Sole Supply on 1 October 2015 | - | |

| CALCIPOTRIOL Crm 50 mcg per g Soln 50 mcg per g COAL TAR Soln – Only in combination 1) Up to 10% only in combination with a dermatological base or pro- base, page 206 2) With or without other dermatological galenicals. COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5% COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint SALICYLIC ACID Powder – Only in combination 1) Only in combination with a dermatological base or proprietar dermatological base, page 206 2) With or without other dermatological galenicals. | 45.00 .45.00 .16.00 .12.55 oprietary Top 3.43 (4.35) 6.59 (8.00) 7.95 | Per 30 g OP 100 g OP 100 g OP 30 ml OP 200 ml | Dai Dai Dai Dai Dai Dai Dai Egr | |
|--|--|---|---|---|
| Crm 50 mcg per g Oint 50 mcg per g Soln 50 mcg per ml COAL TAR Soln – Only in combination 1) Up to 10% only in combination with a dermatological base or pro- base, page 206 2) With or without other dermatological galenicals. COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5% COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint SALICYLIC ACID Powder – Only in combination 1) Only in combination with a dermatological base or proprietar dermatological base, page 206 | 45.00 .45.00 .16.00 .12.55 oprietary Top 3.43 (4.35) 6.59 (8.00) 7.95 | 100 g OP 100 g OP 30 ml OP 200 ml ical Corticoste 30 g OP 75 g OP | Dai Dai Dai Miceriod – P | ivonex ivonex ivonex dwest Plain, refer dermatologic opsoryl TA opsoryl TA |
| Crm 50 mcg per g Oint 50 mcg per g Soln 50 mcg per ml COAL TAR Soln – Only in combination 1) Up to 10% only in combination with a dermatological base or pro- base, page 206 2) With or without other dermatological galenicals. COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5% COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint SALICYLIC ACID Powder – Only in combination 1) Only in combination with a dermatological base or proprietar dermatological base, page 206 | 45.00 .45.00 .16.00 .12.55 oprietary Top 3.43 (4.35) 6.59 (8.00) 7.95 | 100 g OP 100 g OP 30 ml OP 200 ml ical Corticoste 30 g OP 75 g OP | Dai Dai Dai Miceriod – P | ivonex ivonex ivonex dwest Plain, refer dermatologic opsoryl TA opsoryl TA |
| Oint 50 mcg per g Soln 50 mcg per ml COAL TAR Soln – Only in combination 1) Up to 10% only in combination with a dermatological base or probase, page 206 2) With or without other dermatological galenicals. COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5% COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint SALICYLIC ACID Powder – Only in combination 1) Only in combination with a dermatological base or proprietar dermatological base, page 206 | 45.00 .45.00 .16.00 .12.55 oprietary Top 3.43 (4.35) 6.59 (8.00) 7.95 | 100 g OP 100 g OP 30 ml OP 200 ml ical Corticoste 30 g OP 75 g OP | Dai Dai Dai Miceriod – P | ivonex ivonex ivonex dwest Plain, refer dermatologic opsoryl TA opsoryl TA |
| Soln 50 mcg per ml COAL TAR Soln – Only in combination 1) Up to 10% only in combination with a dermatological base or probase, page 206 2) With or without other dermatological galenicals. COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5% COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint SALICYLIC ACID Powder – Only in combination 1) Only in combination with a dermatological base or proprietar dermatological base, page 206 | .45.00 .16.00 .12.55 oprietary Top 3.43 (4.35) 6.59 (8.00) 7.95 | 100 g OP 30 ml OP 200 ml ical Corticoste 30 g OP 75 g OP | Dai Dai Dai Dai | ivonex ivonex d <u>west</u> dain, refer dermatologic opsoryl TA opsoryl TA |
| Soln 50 mcg per ml COAL TAR Soln – Only in combination 1) Up to 10% only in combination with a dermatological base or probase, page 206 2) With or without other dermatological galenicals. COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5% COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint SALICYLIC ACID Powder – Only in combination 1) Only in combination with a dermatological base or proprietar dermatological base, page 206 | . 16.00 . 12.55 oprietary Top 3.43 (4.35) 6.59 (8.00) 7.95 | 30 ml OP 200 ml ical Corticoste 30 g OP 75 g OP | ✓ Dai ✓ <u>Mic</u> erriod – P Egr Egr | ivonex dwest Plain, refer dermatologic opsoryl TA opsoryl TA |
| COAL TAR Soln – Only in combination 1) Up to 10% only in combination with a dermatological base or probase, page 206 2) With or without other dermatological galenicals. COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5% COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint | . 12.55 oprietary Top 3.43 (4.35) 6.59 (8.00) 7.95 | 200 ml iical Corticoste 30 g OP 75 g OP | <mark>✓ <u>Mic</u>eriod – P</mark> Ega Ega | dwest I'lain, refer dermatologic opsoryl TA opsoryl TA |
| Soln – Only in combination | 3.43 (4.35) 6.59 (8.00) 7.95 | ical Corticoste 30 g OP 75 g OP | eriod – P Ego Ego | lain, refer dermatologic opsoryl TA opsoryl TA |
| Up to 10% only in combination with a dermatological base or probase, page 206 With or without other dermatological galenicals. COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5% COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint COAL TAR UITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint | 3.43 (4.35) 6.59 (8.00) 7.95 | ical Corticoste 30 g OP 75 g OP | eriod – P Ego Ego | lain, refer dermatologic opsoryl TA opsoryl TA |
| base, page 206 2) With or without other dermatological galenicals. COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5% COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint COALICYLIC ACID Powder – Only in combination | 3.43 (4.35) 6.59 (8.00) 7.95 | 30 g OP 75 g OP | Ego | opsoryl TA opsoryl TA |
| COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5% COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint SALICYLIC ACID Powder – Only in combination | (4.35) 6.59 (8.00) 7.95 | 75 g OP | Ego | opsoryl TA |
| Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5% COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint SALICYLIC ACID Powder – Only in combination 1) Only in combination with a dermatological base or proprietar dermatological base, page 206 | (4.35) 6.59 (8.00) 7.95 | 75 g OP | Ego | opsoryl TA |
| allantoin crm 2.5% COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint SALICYLIC ACID Powder – Only in combination 1) Only in combination with a dermatological base or proprietar dermatological base, page 206 | (4.35) 6.59 (8.00) 7.95 | 75 g OP | Ego | opsoryl TA |
| COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint SALICYLIC ACID Powder – Only in combination | (4.35) 6.59 (8.00) 7.95 | 75 g OP | Ego | opsoryl TA |
| Soln 12% with salicylic acid 2% and sulphur 4% oint SALICYLIC ACID Powder - Only in combination | `6.59 [´] (8.00) 7.95 | Ū | Ego | opsoryl TA |
| Soln 12% with salicylic acid 2% and sulphur 4% oint SALICYLIC ACID Powder - Only in combination | (8.00) 7.95 | Ū | | |
| Soln 12% with salicylic acid 2% and sulphur 4% oint SALICYLIC ACID Powder - Only in combination | 7.95 | 40 g OP | | |
| Soln 12% with salicylic acid 2% and sulphur 4% oint SALICYLIC ACID Powder - Only in combination | | 40 g OP | 🖌 Co | co-Scalp |
| ALICYLIC ACID Powder – Only in combination | | 40 g OP | 🗸 Co | co-Scalp |
| Powder – Only in combination 1) Only in combination with a dermatological base or proprietar dermatological base, page 206 | 10.00 | | | |
| Powder – Only in combination 1) Only in combination with a dermatological base or proprietar dermatological base, page 206 | 10.00 | | | |
| Only in combination with a dermatological base or proprietar dermatological base, page 206 | . 18.88 | 250 a | 🖌 PS | М |
| , 3 6 | y Topical Co | orticosteroid – | Plain o | r collodion flexible, ref |
| SULPHUR | | | | |
| Precipitated – Only in combination | | 100 g | 🖌 Mic | |
| Only in combination with a dermatological base or proprietary page 206 | / Topical Cor | rticosteroid – | Plain, re | eter dermatological bas |
| 2) With or without other dermatological galenicals. | | | | |
| AR WITH TRIETHANOLAMINE LAURYL SULPHATE AND FLUORES | CEIN – Only | y on a prescrip | otion | |
| Soln 2.3% with triethanolamine lauryl sulphate and fluores- | | | | |
| cein sodium | 3.36 | 500 ml | 🖌 Pin | netarsol |
| Pinetarsol to be Sole Supply on 1 October 2015 | | | | |
| Scalp Preparations | | | | |
| | | | | |
| BETAMETHASONE VALERATE | | | | |
| ₭ Scalp app 0.1% | 7.75 | 100 ml OP | 🖌 Bei | ta Scalp |
| CLOBETASOL PROPIONATE | | | | |
| ₭ Scalp app 0.05% | 6.96 | 30 ml OP | 🖌 Dei | rmol |
| IYDROCORTISONE BUTYRATE | | | | |
| Scalp lotn 0.1% | 3.65 | 100 ml OP | 🖌 Lo | coid |
| • | | | ÷ 10 | ~~~~ |
| KETOCONAZOLE | 0.00 | 100 | | hi-ala |
| Shampoo 2% | 2.99 | 100 ml OP | ✓ Sel | DIZOIE |
| a) Maximum of 100 ml per prescription b) Only on a prescription | | | | |

| | Subsidy (Manufacturer's F | rice) Su | Fully bsidised | Brand or Generic |
|--|------------------------------|-----------------|-------------------|------------------------------|
| | \$ | Per | ~ | Manufacturer |
| Sunscreens | | | | |
| SUNSCREENS, PROPRIETARY – Subsidy by endorsement Only if prescribed for a patient with severe photosensitivity endorsed accordingly. | v secondary to a o | defined clinica | al conditi | on and the prescription i |
| Crm | 3 30 | 100 g OP | | |
| | (5.89) | 100 9 01 | н | amilton Sunscreen |
| Lotn, | (/ | 100 g OP | | arine Blue Lotion SPF 50+ |
| | 5.10 | 200 g OP | ✔ M | arine Blue Lotion SPF 50+ |
| Lotn | 4.13 | 125 ml OP | | |
| | (6.94) | | A | quasun 30+ |
| Wart Preparations | | | | |
| or salicylic acid preparations refer to PSORIASIS AND ECZEM | IA PREPARATION | S, page 70 | | |
| MIQUIMOD | | | | |
| Crm 5%, 250 mg sachet | 17.98 | 12 | ✓ <u>A</u> | po-Imiquimod Cream 5% |
| ODOPHYLLOTOXIN | | | | |
| Soln 0.5% a) Maximum of 3.50 ml per prescription b) Only on a prescription | 33.60 | 3.5 ml OP | ✔ C | ondyline |
| Other Skin Preparations | | | | |
| Antineoplastics | | | | |
| LUOROURACIL SODIUM Crm 5% | 8 95 | 20 g OP | | fudix |
| Efudix to be Sole Supply on 1 October 2015 | 0.00 | 20 9 01 | ΨL | |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|-----|---------------------|--|
| Contraceptives - Non-hormonal | | | | |
| Condoms | | | | |
| CONDOMS | | | | |
| ₭ 49 mm – Up to 144 dev available on a PSO | 13.36 | 144 | | larquisTantiliza hield 49 |
| € 52 mm – Up to 144 dev available on a PSO | 13.36 | 144 | V N | larquis Selecta Iarquis Sensolite |
| | | | | larquis Supalite |
| 52 mm extra strength – Up to 144 dev available on a P | | 144 | | larquis Protecta |
| € 53 mm – Up to 144 dev available on a PSO | | 12 | 🗸 S | iold Knight hield Blue |
| | 13.36 | 144 | V N | hield Blue Iarquis Black Iarquis Titillata |
| € 53 mm (chocolate) – Up to 144 dev available on a PSC |) 111 | 12 | | iold Knight |
| | 13.36 | 144 | | iold Knight |
| 53 mm (strawberry) – Up to 144 dev available on a PS | | 12 | | iold Knight |
| | 13.36 | 144 | | old Knight |
| 54 mm, shaped – Up to 144 dev available on a PSO | | 12 | • • | iona rungin |
| | (1.24) | 12 | 1 | ifestyles Flared |
| | 13.36 | 144 | _ | nootyloo i laioa |
| | (14.84) | | 1 | ifestyles Flared |
| 55 mm – Up to 144 dev available on a PSO | | 144 | | larguis Conforma |
| 56 mm – Up to 144 dev available on a PSO | | 12 | | old Knight |
| | 13.36 | 144 | | old Knight |
| | | | VD | ourex Extra Safe ourex Select Flavours |
| € 56 mm, shaped – Up to 144 dev available on a PSO | 1.11 | 12 | 🗸 D | urex Confidence |
| | 13.36 | 144 | 🖌 D | urex Confidence |
| 60 mm – Up to 144 dev available on a PSO | | 144 | 🖌 S | hield XL |
| Contraceptive Devices | | | | |
| DIAPHRAGM – Up to 1 dev available on a PSO One of each size is permitted on a PSO. | | | | |
| 65 mm | 10 00 | 1 | | ortho All-flex |
| 70 mm | | 1 | | ortho All-flex |
| 75 mm | | 1 | | ortho All-flex |
| 80 mm | | 1 | | ortho All-flex |
| | | • | | |
| ITRA-UTERINE DEVICE a) Up to 40 dev available on a PSO b) Only on a PSO | | | | |
| ► IUD 29.1 mm length × 23.2 mm width | | 1 | v 0 | hoice TT380 Short |
| UD 33.6 mm length × 29.9 mm width | | 1 | | choice TT380 Standard |

GENITO-URINARY SYSTEM

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

Contraceptives - Hormonal

Combined Oral Contraceptives

➡SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 Either:

- 1.1 Patient is on a Social Welfare benefit; or
- 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

| * | Tab 20 mcg with desogestrel 150 mcg and 7 inert tab | 6.62 (19.80) | 84 | Mercilon 28 |
|-----|--|-----------------|------|--------------------------------------|
| | a) Higher subsidy of \$13.80 per 84 tab with Special Authority | see SA0500 abov | e | |
| | b) Up to 84 tab available on a PSO | | | |
| * | Tab 30 mcg with desogestrel 150 mcg and 7 inert tab | 6.62 | 84 | |
| | | (19.80) | | Marvelon 28 |
| | a) Higher subsidy of \$13.80 per 84 tab with Special Authority | see SA0500 abov | e | |
| | b) Up to 84 tab available on a PSO | | | |
| ETI | HINYLOESTRADIOL WITH LEVONORGESTREL | | | |
| * | Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tab – Up | | | |
| | to 84 tab available on a PSO | 2.65 | 84 🖌 | 🖊 Ava 20 ED |
| * | Tab 50 mcg with levonorgestrel 125 mcg and 7 inert tab – Up | | | |
| | to 84 tab available on a PSO | 9.45 | 84 🖌 | Microgynon 50 ED |
| * | Tab 30 mcg with levonorgestrel 150 mcg | 6.62 | 63 | |
| | | (16.50) | | Microgynon 30 |
| | a) Higher subsidy of \$15.00 per 63 tab with Special Authority | see SA0500 abov | e | |
| | b) Up to 63 tab available on a PSO | | | |
| * | Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tab – Up | | | |
| | to 84 tab available on a PSO | 2.30 | 84 🖌 | 🖊 Ava 30 ED |
| | | | | |

GENITO-URINARY SYSTEM

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Generic |
|--|---|-----|---------------------|---------------|
| ETHINYLOESTRADIOL WITH NORETHISTERONE | | | | |
| * Tab 35 mcg with norethisterone 1 mg – Up to 63 tab available on a PSO | 6.62 | 63 | ~ | Brevinor 1/21 |
| Tab 35 mcg with norethisterone 1 mg and 7 inert tab – Up to 84 tab available on a PSO | 6.62 | 84 | ~ | Brevinor 1/28 |
| Tab 35 mcg with norethisterone 500 mcg – Up to 63 tab avail- able on a PSO | 6.62 | 63 | ~ | Brevinor 21 |
| * Tab 35 mcg with norethisterone 500 mcg and 7 inert tab – Up to 84 tab available on a PSO | 6.62 | 84 | ~ | Norimin |

Progestogen-only Contraceptives

➡SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 Either:

- 1.1 Patient is on a Social Welfare benefit; or
- 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

LEVONORGESTREL

| * Tab 30 mcg | 6.62 | 84 | |
|---|---------------------|-----|----------------|
| - | (16.50) | | Microlut |
| a) Higher subsidy of \$13.80 per 84 tab with Special Auth b) Up to 84 tab available on a PSO | ority see SA0500 at | ove | |
| * Subdermal implant (2 × 75 mg rods) | | 1 | ✓ Jadelle |
| MEDROXYPROGESTERONE ACETATE * Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a F | PSO7.00 | 1 | ✓ Depo-Provera |
| NORETHISTERONE * Tab 350 mcg – Up to 84 tab available on a PSO Noriday 28 to be Sole Supply on 1 November 2015 | 6.25 | 84 | 🗸 Noriday 28 |

| | Subsidy | | Fully Brand or | |
|--|--------------------------|--------------------|---|--|
| | (Manufacturer's Pr \$ | ice) Sub Per | osidised Generic Manufacturer | |
| Emergency Contraceptives | | | | |
| LEVONORGESTREL | | | | |
| * Tab 1.5 mga) Up to 5 tab available on a PSO b) Maximum of 2 tab per prescription | 3.50 | 1 | ✓ Postinor-1 | |
| Antiandrogen Oral Contraceptives | | | | |
| Prescribers may code prescriptions "contraceptive" (code "O") whe prescription charge will be as per other contraceptives, as follows: • \$5.00 prescription charge (patient co-payment) will apply. • prescription may be written for up to six months supply. Prescriptions coded in any other way are subject to the non contr of supply. ie. Prescriptions may be written for up to three months as CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL | aceptive prescrip | | | |
| * Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO | 5 36 | 168 | ✓ Ginet | |
| Gynaecological Anti-infectives | | 100 | • <u>dinet</u> | |
| | | | | |
| ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC A Jelly with glacial acetic acid 0.94%, hydroxyquinoline sul- | CID | | | |
| phate 0.025%, glycerol 5% and ricinoleic acid 0.75% with | | | | |
| applicator | 8.43 (24.00) | 100 g OP | Aci-Jel | |
| CLOTRIMAZOLE | | | | |
| * Vaginal crm 1% with applicators * Vaginal crm 2% with applicators | | 35 g OP 20 g OP | ✓ <u>Clomazol</u> ✓ Clomazol | |
| MICONAZOLE NITRATE | 2.20 | 20 y OF | | |
| Kontazole NITRATE Kontazole NITRATE Kontazole NITRATE | | 40 g OP | ✓ Micreme | |
| NYSTATIN | | - 5 - | · | |
| Vaginal crm 100,000 u per 5 g with applicator(s) | 4.71 | 75 g OP | ✓ Nilstat | |
| Myometrial and Vaginal Hormone Preparations | | | | |
| ERGOMETRINE MALEATE | | | | |
| Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on a | | | | |
| PSO | 94.70 | 5 | ✓ DBL Ergometrine | |
| OESTRIOL | | | | |
| * Crm 1 mg per g with applicator | | 15 g OP | ✓ Ovestin | |
| * Pessaries 500 mcg | 0.53 | 15 | Ovestin | |
| OXYTOCIN – Up to 5 inj available on a PSO Inj 5 iu per ml, 1 ml ampoule | 4 75 | 5 | Oxytocin BNM | |
| Inj 10 iu per ml, 1 ml ampoule | | 5 | | |
| OXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj availa | | | | |
| Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml Syntometrine to be Sole Supply on 1 October 2015 | | 5 | Syntometrine | |

| | | GENITO- | -URINARY SYSTEM |
|--|---|---------------|--|
| | Subsidy (Manufacturer's Price) \$ | | Fully Brand or dised Generic ✔ Manufacturer |
| Pregnancy Tests - hCG Urine | | | |
| PREGNANCY TESTS - HCG URINE a) Up to 200 test available on a PSO b) Only on a PSO Cassette | 17.60 40 22.80 | | EasyCheck Innovacon hCG One Step Pregnancy Test |
| Urinary Agents | | | |
| For urinary tract Infections refer to INFECTIONS, Antibacterials, p | bage 112 | | |
| 5-Alpha Reductase Inhibitors | | | |
| FINASTERIDE – Special Authority see SA0928 below – Retail pl * Tab 5 mg SA0928 Special Authority for Subsidy | | 28 | ✓ Finpro |
| Initial application from any relevant practitioner. Approvals valid the following criteria: Both: | d without further ren | ewal unless r | notified for applications meeting |
| Patient has symptomatic benign prostatic hyperplasia; ar Either: | nd | | |
| 2.1 The patient is intolerant of non-selective alpha blo2.2 Symptoms are not adequately controlled with nor | n-selective alpha bloo | ckers. | ed; or |
| Note: Patients with enlarged prostates are the appropriate candid Alpha-1A Adrenoreceptor Blockers | lates for therapy with | finasteride. | |
| TAMSULOSIN HYDROCHLORIDE – Special Authority see SA10 | 132 helow – Betail nh | armacy | |
| * Cap 400 mcg | | | ✓ Tamsulosin-Rex |
| ► SA1032 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals value the following criteria: Both: | d without further ren | ewal unless r | notified for applications meeting |
| Patient has symptomatic benign prostatic hyperplasia; ar The patient is intolerant of non-selective alpha blockers of | | dicated. | |
| Other Urinary Agents | | | |
| OXYBUTYNIN * Tab 5 mg * Oral liq 5 mg per 5 ml POTASSIUM CITRATE | | | <u>Apo-Oxybutynin</u> <u>Apo-Oxybutynin</u> |
| Oral liq 3 mmol per ml – Special Authority see SA1083 on the next page – Retail pharmacy | | 00 ml OP | ✓ Biomed |

| Subsidy | Fully | Brand or |
|------------------------|------------|----------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | |

➡SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE

| 2.93 | 28 | ✓ Ural |
|----------------|-------------------------|-------------------------------|
| – Retail pharm | acy | |
| 37.50 | 30 | Vesicare |
| 37.50 | 30 | Vesicare |
| | – Retail pharm 37.50 | – Retail pharmacy 37.50 30 |

SA0998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has overactive bladder and a documented intolerance of, or is non-responsive to oxybutynin.

| TOLTERODINE – Special Authority see SA1272 below – Retail pharmacy | | |
|--|----|-------------------|
| Tab 1 mg14.56 | 56 | Arrow-Tolterodine |
| Tab 2 mg14.56 | 56 | Arrow-Tolterodine |

SA1272 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where patient has overactive bladder and a documented intolerance of, or is non-responsive to oxybutynin.

| Detection of Substances in Urine | | | |
|----------------------------------|---------|-------------|----------|
| ORTHO-TOLIDINE | | | |
| * Compound diagnostic sticks | 7.50 | 50 test OP | |
| | (8.25) | | Hemastix |
| TETRABROMOPHENOL | | | |
| * Blue diagnostic strips | 7.02 | 100 test OP | |
| <u> </u> | (13.92) | | Albustix |

| | 0.1.11 | | |
|--|------------------------------|----------------|--|
| | Subsidy (Manufacturer's P | rice) Sub | Fully Brand or sidised Generic |
| | \$ | Per | Manufacturer |
| Calcium Homeostasis | | | |
| CALCITONIN | | | |
| * Inj 100 iu per ml, 1 ml ampoule | 121.00 | 5 | ✓ Miacalcic |
| ZOLEDRONIC ACID | | | |
| Inj 4 mg per 5 ml, vial – Special Authority see SA1512 below | | | |
| – Retail pharmacy | 550.00 | 1 | Zometa |
| ► SA1512 Special Authority for Subsidy | | | |
| Initial application only from an oncologist, haematologist or pa unless notified for applications meeting the following criteria: | lliative care spe | cialist. Appro | vals valid without further renewal |
| Any of the following: | | | |
| 1 Patient has hypercalcaemia of malignancy; or | | | |
| 2 Both: | | | |
| 2.1 Patient has bone metastases or involvement; and | l first line treatm | antai ar | |
| 2.2 Patient has severe bone pain resistant to standard3 Both: | i inst-ine treatm | ents; or | |
| 3.1 Patient has bone metastases or involvement; and | | | |
| 3.2 Patient is at risk of skeletal-related events patho | ological fracture, | spinal cord c | ompression, radiation to bone or |
| surgery to bone). | 0 , | , | |
| Corticosteroids and Related Agents for Systemic | c Use | | |
| BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHAS | | | |
| * Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml | | 5 | |
| · · · · · · · · · · · · · · · · · · · | (33.60) | - | Celestone |
| | | | Chronodose |
| DEXAMETHASONE | | | 4 - - |
| * Tab 1 mg – Retail pharmacy-Specialist Up to 30 tab available on a PSO | 5.87 | 100 | Douglas |
| Tab 4 mg – Retail pharmacy-Specialist | 8.16 | 100 | ✓ Douglas |
| Up to 30 tab available on a PSO | | | · |
| Oral liq 1 mg per ml – Retail pharmacy-Specialist | | 25 ml OP | Biomed |
| Oral liq prescriptions: 1) Must be written by a Paediatrician or Paediatric Cardiolog | iist: or | | |
| 2) On the recommendation of a Paediatrician or Paediatric C | | | |
| DEXAMETHASONE PHOSPHATE | 0 | | |
| Dexamethasone phosphate injection will not be funded for ora | | | 4 - |
| * Inj 4 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSC |)25.80 | 10 | <u>Dexamethasone-</u> hameln |
| * Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSC |)17.98 | 5 | ✓ Dexamethasone- |
| | | | hameln |
| FLUDROCORTISONE ACETATE | | | |
| * Tab 100 mcg | 14.32 | 100 | Florinef |
| | | | |

| | Subsidy (Manufacturer's Price \$ |) Per | Full Subsidise | |
|---|--|----------|-------------------|-------------------------------|
| IYDROCORTISONE | | | | |
| * Tab 5 mg | 8.10 | 100 | ~ | Douglas |
| Douglas to be Sole Supply on 1 October 2015 | | | | Ū |
| Tab 20 mg – For hydrocortisone oral liquid formulation refer, | | | | |
| page 207 | | 100 | ~ | Douglas |
| Douglas to be Sole Supply on 1 October 2015 | | | | Ū |
| € Inj 100 mg vial | | 1 | ~ | Solu-Cortef |
| a) Up to 5 inj available on a PSO | | | | |
| b) Only on a PSO | | | | |
| | | | | |
| ETHYLPREDNISOLONE – Retail pharmacy-Specialist | 80.00 | 100 | | Madral |
| • Tab 4 mg | | 100 | V | Medrol |
| Medrol to be Sole Supply on 1 November 2015 | 100.00 | 00 | | Madual |
| Tab 100 mg | | 20 | V | Medrol |
| Medrol to be Sole Supply on 1 November 2015 | | | | |
| ETHYLPREDNISOLONE (AS SODIUM SUCCINATE) – Retail p | harmacy-Specialist | | | |
| Inj 40 mg vial | | 1 | ~ | Solu-Medrol |
| Solu-Medrol to be Sole Supply on 1 November 2015 | | | | |
| Inj 125 mg vial | | 1 | ~ | Solu-Medrol |
| Solu-Medrol to be Sole Supply on 1 November 2015 | | | | |
| Inj 500 mg vial | 9.00 | 1 | ~ | Solu-Medrol |
| Solu-Medrol to be Sole Supply on 1 November 2015 | | | | |
| Inj 1 g vial | | 1 | ~ | Solu-Medrol |
| Solu-Medrol to be Sole Supply on 1 November 2015 | | | | |
| IETHYLPREDNISOLONE ACETATE | | | | |
| | 40.00 | F | | Done Madral |
| Inj 40 mg per ml, 1 ml vial | | 5 | V | Depo-Medrol |
| Depo-Medrol to be Sole Supply on 1 November 2015 | | | | |
| ETHYLPREDNISOLONE ACETATE WITH LIDOCAINE [LIGNOC | | | | |
| Inj 40 mg per ml with lidocaine [lignocaine] 1 ml vial | 9.25 | 1 | ~ | Depo-Medrol with Lidocaine |
| Depo-Medrol with Lidocaine to be Sole Supply on 1 Novem | nher 2015 | | | |
| | | | | |
| REDNISOLONE | 7.50 0 | 0 | n . 4 | Dedinand |
| Oral liq 5 mg per ml – Up to 30 ml available on a PSO | | 0 ml O | r V | Redipred |
| Restricted to children under 12 years of age. | | | | |
| REDNISONE | | | | |
| 🗧 Tab 1 mg | 2.13 | 100 | ~ | Apo-Prednisone |
| | | | | S29 S29 |
| | 10.68 | 500 | ~ | Apo-Prednisone |
| - Tab 2.5 mg | | 500 | | Apo-Prednisone |
| Tab 5 mg – Up to 30 tab available on a PSO | | 500 | | Apo-Prednisone |
| Tab 20 mg | | 500 | | Apo-Prednisone |
| 5 | | 500 | • | Aport realisone |
| ETRACOSACTRIN | | | | |
| Inj 250 mcg per ml, 1 ml ampoule | | 1 | | Synacthen |
| | 177.18 | 10 | | Synacthen |
| Inj 1 mg per ml, 1 ml | | 1 | ~ | Synacthen Depot |
| RIAMCINOLONE ACETONIDE | | | | |
| Inj 10 mg per ml, 1 ml ampoule | 20.80 | 5 | ~ | Kenacort-A 10 |
| Inj 40 mg per ml, 1 ml ampoule | | 5 | | Kenacort-A 40 |
| ייון דיט יווץ אבי וווו, ד ווו מווואטעופ | | 5 | | Nellacolt-A 40 |

| | Subsidy (Manufacturer's Price) \$ | Sul Per | Fully bsidised | Brand or Generic Manufacturer |
|---|---|------------|-------------------|-------------------------------------|
| Sex Hormones Non Contraceptive | | | | |
| Androgen Agonists and Antagonists | | | | |
| YPROTERONE ACETATE – Retail pharmacy-Specialist | | | | |
| Tab 50 mg | 15.87 18.80 | 50 | • • | rocur iterone |
| Tab 100 mg | | 50 | | rocur iterone |
| ESTOSTERONE | | | | |
| Transdermal patch, 2.5 mg per day | 80.00 | 60 | 🗸 A | ndroderm |
| ESTOSTERONE CYPIONATE – Retail pharmacy-Specialist Inj 100 mg per ml, 10 ml vial | 76.50 | 1 | ✓ <u>D</u> | epo-Testosterone |
| ESTOSTERONE ESTERS – Retail pharmacy-Specialist Inj 250 mg per ml, 1 ml | 12.98 | 1 | ✔ S | ustanon Ampoules |
| ESTOSTERONE UNDECANOATE – Retail pharmacy-Specialis Cap 40 mg | | 60 | 🗸 A | ndriol Testocaps |
| Andriol Testocaps to be Sole Supply on 1 October 2015 Inj 250 mg per ml, 4 ml vial | | 1 | 🗸 R | eandron 1000 |

SA1018 Special Authority for Alternate Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 years for applications meeting the following criteria: Any of the following:

- 1 acute or significant liver disease where oral oestrogens are contraindicated as determined by a gastroenterologist or general physician. The applicant must keep written confirmation from such a specialist with the patient's record; or
- 2 oestrogen induced hypertension requiring antihypertensive therapy documented evidence must be kept on file that raised blood pressure levels or inability to control blood pressure adequately occurred post oral oestrogens; or
- 3 hypertriglyceridaemia documented evidence must be kept on file that triglyceride levels increased to at least 2 × normal triglyceride levels post oral oestrogens; or
- 4 Somatropin co-therapy patient is being prescribed somatropin with subsidy provided under a valid approval issued under Special Authority.

Note: Prescriptions with a valid Special Authority (CHEM) number will be reimbursed at the level of the lowest priced TDDS product within the specified dose group.

Renewal from any relevant practitioner. Approvals valid for 5 years where the treatment remains appropriate and the patient is benefiting from treatment, or the patient remains on subsidised somatropin co-therapy.

Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

| | Subsidy (Manufacturer's Pr \$ | ice) Su Per | Fully Brand or ubsidised Generic Manufacturer |
|--|-------------------------------------|----------------|---|
| Oestrogens | | | |
| OESTRADIOL – See prescribing guideline on the previous pag | e | | |
| * Tab 1 mg | | 28 OP | |
| | (11.10) | | Estrofem |
| * Tab 2 mg | 4.12 | 28 OP | |
| | (11.10) | | Estrofem |
| * TDDS 25 mcg per day | 3.01 | 8 | |
| | (10.86) | | Estradot |
| a) Higher subsidy of \$10.86 per 8 patch with Special Aut b) No more than 2 patch per week c) Only on a prescription | hority see SA1018 | on the previo | ous page |
| TDDS 3.9 mg (releases 50 mcg of oestradiol per day) | 4 12 | 4 | |
| | (13.18) | Ŧ | Climara 50 |
| a) Higher subsidy of \$13.18 per 4 patch with Special Aut | ``` | on the previo | |
| b) No more than 1 patch per week c) Only on a prescription | nonty see SATUTO | | ous page |
| * TDDS 50 mcg per day | | 8 | |
| | (13.18) | · · | Estradot 50 mcg |
| a) Higher subsidy of \$13.18 per 8 patch with Special Aut b) No more than 2 patch per week c) Only on a prescription | - | on the previo | ous page |
| * TDDS 7.8 mg (releases 100 mcg of oestradiol per day) | 7.05 | 4 | |
| | (16.14) | | Climara 100 |
| a) Higher subsidy of \$16.14 per 4 patch with Special Aut b) No more than 1 patch per week c) Only on a prescription | hority see SA1018 | on the previo | ous page |
| * TDDS 100 mcg per day | 7.05 | 8 | |
| | (16.14) | · · | Estradot |
| a) Higher subsidy of \$16.14 per 8 patch with Special Aut b) No more than 2 patch per week c) Only on a prescription | · · · · | on the previo | |
| OESTRADIOL VALERATE – See prescribing guideline on the p | previous page | | |
| * Tab 1 mg | | 84 | Progynova |
| * Tab 2 mg | 12.36 | 84 | Progynova |
| OESTROGENS – See prescribing guideline on the previous pa | ae | | |
| * Conjugated, equine tab 300 mcg | • | 28 | |
| · · · · · · · · · · · · · · · · · · · | (11.48) | | Premarin |
| * Conjugated, equine tab 625 mcg | · · · · | 28 | |
| ······································ | (11.48) | | Premarin |
| Progestogens | (-) | | |
| MEDROXYPROGESTERONE ACETATE – See prescribing gui | deline on the previo | us page | |
| * Tab 2.5 mg | | 30 | Provera |
| * Tab 5 mg | | 100 | ✓ Provera |
| * Tab 3 mg | | 30 | ✓ Provera |
| * 1ab 10 mg | 0.05 | 50 | |

| | Subsidy (Manufacturer's Pric \$ | e) S Per | Fully Subsidised | |
|---|---------------------------------------|-------------|---------------------|------------------------------|
| Progestogen and Oestrogen Combined Prepara | tions | | | |
| OESTRADIOL WITH NORETHISTERONE – See prescribing gui * Tab 1 mg with 0.5 mg norethisterone acetate | | 28 OP | | Kliovance |
| * Tab 2 mg with 1 mg norethisterone acetate | 5.40 (18.10) | 28 OP | I | Kliogest |
| Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg oestradiol tab (12) and 1 mg oestradiol tab (6) | | 28 OP | | Trisequens |
| DESTROGENS WITH MEDROXYPROGESTERONE - See pres | 00 | n page 81 | | |
| Tab 625 mcg conjugated equine with 2.5 mg medroxyproges- terone acetate tab (28) | | 28 OP | I | Premia 2.5 Continuous |
| * Tab 625 mcg conjugated equine with 5 mg medroxyproges- terone acetate tab (28) | | 28 OP | I | Premia 5 Continuous |
| Other Oestrogen Preparations | | | | |
| ETHINYLOESTRADIOL * Tab 10 mcg NZ Medical and Scientific to be Sole Supply on 1 October | | 100 | • | NZ Medical and Scientific |
| OESTRIOL * Tab 2 mg | 7.00 | 30 | | Ovestin |
| Other Progestogen Preparations | | 50 | | ovestin |
| LEVONORGESTREL ★ Levonorgestrel - releasing intrauterine system 20 mcg/24 hr – Special Authority see SA0782 below – Retail pharmacy. | | 1 | ~ 1 | Mirena |
| SA0782 Special Authority for Subsidy nitial application — (No previous use) only from a relevant s applications meeting the following criteria: All of the following: | pecialist or general | practitio | ner. Appr | rovals valid for 6 months fo |
| The patient has a clinical diagnosis of heavy menstrual b The patient has failed to respond to or is unable to tolera Menstrual Bleeding Guidelines; and Either: | 0 | te pharm | aceutical | therapies as per the Heav |
| 3.1 serum ferritin level $<$ 16 mcg/l (within the last 12 s 3.2 haemoglobin level $<$ 120 g/l. | months); or | | | |
| Note: Applications are not to be made for use in patients as contra- nitial application — (Previous use before 1 October 2002) or valid for 6 months for applications meeting the following criteria: All of the following: | | | | |
| 1 The patient had a clinical diagnosis of heavy menstrual b | leeding; and | | | |

The patient had a clinical diagnosis of heavy menstrual bleeding; and
 Patient demonstrated clinical improvement of heavy menstrual bleeding; and

| Subsidy | Fully | Brand or | _ |
|------------------------|------------|--------------|---|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🖌 | Manufacturer | |

continued...

3 Applicant to state date of the previous insertion.

Note: Applications are not to be made for use in patients as contraception except where they meet the above criteria. **Renewal** only from a relevant specialist or general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient demonstrated clinical improvement of heavy menstrual bleeding; or
 - 1.2 Previous insertion was removed or expelled within 3 months of insertion; and
- 2 Applicant to state date of the previous insertion.

| * Tab 100 mg - Retail pharmacy-Specialist | 100 | ✓ <u>Provera</u> |
|--|-----|--------------------------------|
| NORETHISTERONE * Tab 5 mg – Up to 30 tab available on a PSO | 100 | ✓ Primolut N |
| PROGESTERONE | 100 | |
| Cap 100 mg - Special Authority see SA1392 below - Retail | | |
| pharmacy16.50 | 30 | Utrogestan |

➡SA1392 Special Authority for Subsidy

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 For the prevention of pre-term labour*; and

2 Either:

- 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
- 2.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are Unapproved Indications (refer to Interpretations and Definitions).

Thyroid and Antithyroid Agents

CARBIMAZOLE

| * Tab 5 mg | | 100 | Neo-Mercazole |
|---|---------------------------|-------|-------------------------------|
| LEVOTHYROXINE | | | |
| * Tab 25 mcg | | 90 | Synthroid |
| ‡ Safety cap for extemporaneously compounded of | oral liquid preparations. | | - |
| * Tab 50 mcg | | 90 | Synthroid |
| | 64.28 | 1,000 | Eltroxin |
| ‡ Safety cap for extemporaneously compounded of | oral liquid preparations. | | |
| * Tab 100 mcg | 4.21 | 90 | Synthroid |
| | 66.78 | 1,000 | Eltroxin |
| ‡ Safety cap for extemporaneously compounded of | oral liquid preparations. | | |
| LEVOTHYROXINE (MERCURY PHARMA) | | | |
| * Tab 50 mcg | 1.71 | 28 | Mercury Pharma |
| ‡ Safety cap for extemporaneously compounded of | oral liquid preparations. | | |
| * Tab 100 mcg | | 28 | Mercury Pharma |
| ‡ Safety cap for extemporaneously compounded of | oral liquid preparations. | | |

| | Subsidy (Manufacturer's Price) \$ | Subs Per | Fully idised | Brand or Generic Manufacturer |
|---|---|-------------|-----------------|-------------------------------------|
| PROPYLTHIOURACIL – Special Authority see SA1199 below – F | Retail pharmacy | | | |

Propylthiouracil is not recommended for patients under the age of 18 years unless the patient is pregnant and other treatments are contraindicated.

PTU \$29

SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

| SO | MATROPIN (OMNITROPE) – Special Authority see SA1451 below | / – Retail pharma | су | |
|----|---|-------------------|----|-------------------------------|
| * | Inj 5 mg cartridge | .109.50 | 1 | Omnitrope |
| * | Inj 10 mg cartridge | .219.00 | 1 | Omnitrope |
| * | Inj 15 mg cartridge | .328.50 | 1 | Omnitrope |

SA1451 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: Fither:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and</p>
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon followup laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is \leq 14 years (female patients) or \leq 16 years (male patients); and
- 2 Height velocity is ≥ 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is ≥ 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🖌 | Manufacturer | |

continued...

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity \geq 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is \geq 2 cm per year, calculated over six months; and
- 3 A current bone age is \leq 14 years; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is ≥ 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is ≥ 2 cm per year as calculated over six months; and
- 3 A current bone age is \leq 14 years (female patients) or \leq 16 years (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is \leq to 14 years (female patients) or \leq to 16 years (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

- 6.1 The patient has a GFR ≤ 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
- 6.2 The patient has received a renal transplant and has received < 5mg/m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is ≥ 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is $\geq 2 \text{ cm}$ per year as calculated over six months; and
- 3 A current bone age is \leq 14 years (female patients) or \leq 16 years (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient's height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 Either:
 - 3.1 The patient is under two years of age and height velocity has been assessed over a minimum six month period from the age of 12 months, with at least three supine length measurements over this period demonstrating clear and consistent evidence of linear growth failure (with height velocity < 25th percentile); or</p>
 - 3.2 The patient is aged two years or older; and
- 4 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 5 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 6 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by \geq 0.5 standard deviations in the preceding 12 months.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is ≥ 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is ≥ 2 cm per year as calculated over six months; and
- 3 A current bone age is \leq 14 years (female patients) or \leq 16 years (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and

continued...

| Subsidy (Manufacturer's Price) | | Fully Subsidised | Brand or Generic | |
|-----------------------------------|-----|---------------------|---------------------|--|
| \$ | Per | ~ | Manufacturer | |

continued...

6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by ≥ 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA[®]).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of $\leq 3 \text{ mcg}$ per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of $\leq 0.4 \text{ mcg per litre}$.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria: Either:

1 All of the following:

- 1.1 The patient has been treated with somatropin for < 12 months; and
- 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA^(B)) score from baseline; and
- 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
- 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA[®] score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients.

GnRH Analogues

| GOSERELIN A | CETATE |
|-------------|--------|
|-------------|--------|

| Inj 3.6 mg | 1 | Zoladex |
|-------------|---|---------|
| Inj 10.8 mg | 1 | Zoladex |

| | Subsidy | | Full | y Brand or |
|---|-----------------------|--------|------------|------------------|
| (| Manufacturer's Price) | | Subsidised | d Generic |
| | \$ | Per | ~ | Manufacturer |
| EUPRORELIN | | | | |
| Inj 3.75 mg prefilled syringe | 221.60 | 1 | ~ | Lucrin Depot PDS |
| Inj 7.5 mg | | 1 | ~ | Eligard |
| Inj 11.25 mg prefilled syringe | 591.68 | 1 | ~ | Lucrin Depot PDS |
| Inj 22.5 mg | | 1 | ~ | Eligard |
| Inj 30 mg | 591.68 | 1 | ~ | Eligard |
| Inj 30 mg prefilled syringe | | 1 | ~ | Lucrin Depot PDS |
| Inj 45 mg | 832.05 | 1 | ~ | Eligard |
| Vasopressin Agonists | | | | |
| ESMOPRESSIN ACETATE | | | | |
| Tab 100 mcg - Special Authority see SA1401 below - Retail | | | | |
| pharmacy | 36.40 | 30 | ~ | Minirin |
| Tab 200 mcg – Special Authority see SA1401 below – Retail | | | | |
| pharmacy | 93.60 | 30 | ~ | Minirin |
| Nasal drops 100 mcg per ml – Retail pharmacy-Specialist | | 5 ml O | - | Minirin |
| Nasal spray 10 mcg per dose – Retail pharmacy-Specialist | | ml OF | | Desmopressin- |
| | | | • | PH&T |
| Inj 4 mcg per ml, 1 ml – Special Authority see SA1401 below | | | | <u></u> |
| – Retail pharmacy | 67.18 | 10 | ~ | Minirin |
| | | | 5 | |

■SA1401 Special Authority for Subsidy

Initial application — (Desmopressin tablets for Nocturnal enuresis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has primary nocturnal enuresis; and
- 2 The nasal forms of desmopressin are contraindicated; and
- 3 An enuresis alarm is contraindicated.

Initial application — (Desmopressin tablets for Diabetes insipidus) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has cranial diabetes insipidus; and
- 2 The nasal forms of desmopressin are contraindicated.

Renewal — (Desmopressin tablets) from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from the treatment.

Initial application — (Desmopressin injection) only from a relevant specialist. Approvals valid for 2 years where the patient cannot use desmopressin nasal spray or nasal drops.

Renewal — (Desmopressin injection) only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Other Endocrine Agents

CABERGOLINE

| Tab 0.5 mg - Maximum of 2 tab per prescription; can be | | |
|---|---|----------|
| waived by Special Authority see SA1370 on the next page4.75 | 2 | Dostinex |
| 19.00 | 8 | Dostinex |
| Dostinex to be Sole Supply on 1 October 2015 | | |

| Subsidy | Fully | Brand or |
|-----------------|-------------------|--------------|
| (Manufacturer's | Price) Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

SA1370 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 pathological hyperprolactinemia; or
- 2 acromegaly*.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment. Note: Indication marked with * is an Unapproved indication.

CLOMIPHENE CITRATE

| Tab 50 mg | | 10 | ✓ Serophene |
|-------------------------------|------------------|-----|--------------------------------|
| DANAZOL | | | |
| Cap 100 mg | | 100 | 🖌 Azol |
| Cap 200 mg | | 100 | 🖌 Azol |
| METYRAPONE | | | |
| Cap 250 mg – Retail pharmacy- | Specialist520.00 | 50 | Metopirone |

| | Subsidy | | Fully Brand or | |
|--|---------------------|----------------|---|-------|
| | (Manufacturer's P | , | ibsidised Generic | |
| | \$ | Per | Manufacturer | |
| Anthelmintics | | | | |
| ALBENDAZOLE - Special Authority see SA1318 below - Retail | pharmacy | | | |
| Tab 400 mg | | 60 | Eskazole S29 | |
| ►SA1318 Special Authority for Subsidy | | | | |
| Initial application only from an infectious disease specialist or patient has hydatids. | r clinical microbio | logist. Appro | ovals valid for 6 months where | e the |
| Renewal only from an infectious disease specialist or clinical n remains appropriate and the patient is benefitting from the treatm | | provals valid | for 6 months where the treatr | ment |
| MEBENDAZOLE - Only on a prescription | | | | |
| Tab 100 mg | | 24 | De-Worm | |
| Oral liq 100 mg per 5 ml | | 15 ml | | |
| | (7.17) | | Vermox | |
| PRAZIQUANTEL | | | | |
| Tab 600 mg | | 8 | Biltricide | |
| Antibacterials | | | | |
| a) For topical antibacterials, refer to DERMATOLOGICALS, page | 63 | | | |
| b) For anti-infective eve preparations, refer to SENSORY ORGAN | | | | |
| | 10, pago 200 | | | |
| Cephalosporins and Cephamycins | | | | |
| CEFACLOR MONOHYDRATE | | | | |
| Cap 250 mg | | 100 | Ranbaxy-Cefaclor | |
| Grans for oral liq 125 mg per 5 ml – Wastage claimable – see | | | | |
| rule 3.3.2 on page 13 | 3.53 | 100 ml | Ranbaxy-Cefaclor | |
| CEFALEXIN | | | | |
| Cap 500 mg | | 20 | Cephalexin ABM | |
| Grans for oral liq 25 mg per ml - Wastage claimable - see | | | | |
| rule 3.3.2 on page 13 | | 100 ml | Cefalexin Sandoz | |
| a) Note: Cefalexin grans for oral liq will not be funded in a | | n 14 days trea | atment per dispensing. | |
| b) Cefalexin Sandoz to be Sole Supply on 1 October 2015 Grans for oral liq 50 mg per ml – Wastage claimable – see | | | | |
| rule 3.3.2 on page 13 | | 100 ml | Cefalexin Sandoz | |
| a) Note: Cefalexin grans for oral liq will not be funded in a | | | | |
| b) Cefalexin Sandoz to be Sole Supply on 1 October 2015 | | | | |
| CEFAZOLIN – Subsidy by endorsement | | | | |
| Only if prescribed for dialysis or cellulitis in accordance with a | a DHB approved p | protocol and t | he prescription is endorsed acc | cord |
| ingly. | | | | |
| Inj 500 mg vial | | 5 | ✓ <u>AFT</u> | |
| Inj 1 g vial | | 5 | ✓ <u>AFT</u> | |
| CEFTRIAXONE – Subsidy by endorsement | | | | |
| a) Up to 5 inj available on a PSO | | | . Construction of the state of | |
| b) Subsidised only if prescribed for a dialysis or cystic fibro | | | | |
| pelvic inflammatory disease, or the treatment of suspected the prescription or PSO is endorsed accordingly. | meningius in patie | ents who hav | e a known allergy to penicillin, | , and |
| Inj 500 mg vial | 1 50 | 1 | Ceftriaxone-AFT | |
| | | | | |
| Inj 1 g vial | | 5 | Ceftriaxone-AFT | |

| | Subsidy | | Fully Brand or |
|--|----------------------|-----------------|------------------------------------|
| | (Manufacturer's Pric | e) Sul Per | bsidised Generic Manufacturer |
| | \$ | Per | Manufacturer |
| CEFUROXIME AXETIL – Subsidy by endorsement Only if prescribed for prophylaxis of endocarditis and the presc | rintion is andorsa | d according | nlv. |
| Tab 250 mg | | 50 50 | Zinnat |
| Macrolides | | | |
| AZITHROMYCIN – Maximum of 5 days treatment per prescription For Endorsement, patient has either: | ; can be waived b | y endorsem | nent |
| Received a lung transplant and requires treatment or prop | hylaxis for bronch | iolitis obliter | rans syndrome*; or |
| Cystic fibrosis and has chronic infection with Pseudomor isms*. | | | |
| Indications marked with * are Unapproved Indications | | | |
| Tab 250 mg | 9.00 | 30 | Apo-Azithromycin |
| Apo-Azithromycin to be Sole Supply on 1 October 2015 Tab 500 mg – Up to 8 tab available on a PSO | 1.05 | 2 | Apo-Azithromycin |
| Apo-Azithromycin to be Sole Supply on 1 October 2015 | | 2 | • Apo-Azitinomycin |
| Grans for oral lig 200 mg per 5 ml (40 mg per ml) – Wastage | | | |
| claimable – see rule 3.3.2 on page 13 | 12.50 | 15 ml | Zithromax |
| Zithromax to be Sole Supply on 1 November 2015 | | | |
| CLARITHROMYCIN - Maximum of 500 mg per prescription; can b | be waived by Spec | cial Authority | y see SA1131 below |
| Tab 250 mg | 3.98 | 14 | Apo-Clarithromycin |
| Grans for oral liq 125 mg per 5 ml – Wastage claimable – see | | | |
| rule 3.3.2 on page 13 | | 70 ml | Klacid |
| SA1131 Special Authority for Waiver of Rule | | | |
| Initial application — (Mycobacterial infections) only from a res | | , infectious | disease specialist or paediatricia |
| Approvals valid for 2 years for applications meeting the following cr | iteria: | | |
| Either: | | | |
| 1 Atypical mycobacterial infection; or | registeres er intel | aranaa ta a | tandard pharmagautical agenta |
| 2 Mycobacterium tuberculosis infection where there is drug- | | | 1 |
| Renewal — (Mycobacterial infections) only from a respiratory sp valid for 2 years where the treatment remains appropriate and the | | | |
| | palient is benenui | iy nom dea | unen. |
| ERYTHROMYCIN ETHYL SUCCINATE Tab 400 mg | 16.05 | 100 | 🖌 E-Mycin |
| a) Up to 20 tab available on a PSO | | 100 | E-MyCIII |
| b) Up to 2 x the maximum PSO quantity for RFPP – see rul | e 5.2.6 on page 1 | 7 | |
| Grans for oral liq 200 mg per 5 ml | | 100 ml | E-Mycin |
| a) Up to 300 ml available on a PSO | | | • |
| b) Up to 2 x the maximum PSO quantity for RFPP - see rul | e 5.2.6 on page 1 | 7 | |
| c) Wastage claimable – see rule 3.3.2 on page 13 | | | |
| Grans for oral liq 400 mg per 5 ml | 6.77 | 100 ml | E-Mycin |
| a) Up to 200 ml available on a PSO | | | |
| b) Wastage claimable – see rule 3.3.2 on page 13 | | | |
| ERYTHROMYCIN LACTOBIONATE Inj 1 g | 16.00 | 1 | Erythrocin IV |
| | | I | |
| ERYTHROMYCIN STEARATE | 14.05 | 100 | |
| Tab 250 mg – Up to 30 tab available on a PSO | | 100 | ERA |
| Tab 500 mg | (22.29) 29.90 | 100 | |
| | (44.58) | 100 | ERA |
| | (| | _ |

| | Subsidy | | Subsidy Fully B | |
|---|---|---------|-----------------|---------------------|
| | | | Subsidised | |
| | (included of of of the second | Per | V | Manufacturer |
| | | | | |
| ROXITHROMYCIN | | | | |
| Tab 150 mg | | 50 | ~ | Arrow- |
| | | | | Roxithromycin |
| T-h 000 mm | 14.40 | 50 | | |
| Tab 300 mg | 14.40 | 50 | V | Arrow- |
| | | | | Roxithromycin |
| Penicillins | | | | |
| | | | | |
| AMOXICILLIN | | | | |
| Cap 250 mg | 16 18 | 500 | ~ | Apo-Amoxi |
| a) Up to 30 cap available on a PSO | | 500 | • | Apo Amoxi |
| , , , , | | | | |
| b) Up to 10 x the maximum PSO quantity for RFPP – see | | | | |
| Cap 500 mg | 20.94 | 500 | V | Apo-Amoxi |
| a) Up to 30 cap available on a PSO | | | | |
| b) Up to 10 x the maximum PSO quantity for RFPP - see | rule 5.2.6 on page | e 17 | | |
| Grans for oral lig 125 mg per 5 ml | | 100 ml | ~ | Alphamox |
| | | 100 111 | | Amoxicillin Actavis |
| | | | | |
| | | | V | Ranmoxy |
| a) Up to 200 ml available on a PSO | | | | |
| b) Wastage claimable – see rule 3.3.2 on page 13 | | | | |
| Grans for oral lig 250 mg per 5 ml | 0.97 | 100 ml | ~ | Alphamox |
| 1 31 | | | ~ | Amoxicillin Actavis |
| | | | | Ranmoxy |
| a) Up to 300 ml available on a PSO | | | • | nannoxy |
| , , | | 4- | | |
| b) Up to 10 x the maximum PSO quantity for RFPP – see | rule 5.2.6 on page | e 1/ | | |
| c) Wastage claimable - see rule 3.3.2 on page 13 | | | | |
| Inj 250 mg vial | | 10 | ~ | Ibiamox |
| Inj 500 mg vial | | 10 | V | Ibiamox |
| Inj 1 g vial – Up to 5 inj available on a PSO | | 10 | ~ | Ibiamox |
| (Ranmoxy Grans for oral lig 125 mg per 5 ml to be delisted 1 Octo | | | • | |
| | , | | | |
| (Ranmoxy Grans for oral liq 250 mg per 5 ml to be delisted 1 Octo | ober 2015) | | | |
| AMOXICILLIN WITH CLAVULANIC ACID | | | | |
| Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab avail- | | | | |
| able on a PSO | | 00 | | Augustin |
| able on a PSO | | 20 | | Augmentin |
| | 9.75 | 100 | V | Curam Duo |
| Grans for oral liq amoxicillin 125 mg with clavulanic acid | | | | |
| 31.25 mg per 5 ml | | 100 ml | ~ | Augmentin |
| oo | | | | Curam |
| a) Lin ta 000 mi available an a DCO | | | • | ourum |
| a) Up to 200 ml available on a PSO | | | | |
| b) Wastage claimable – see rule 3.3.2 on page 13 | | | | |
| Grans for oral liq amoxicillin 250 mg with clavulanic acid | | | | |
| 62.5 mg per 5 ml | 2.19 | 100 ml | v | Augmentin |
| | | | V | Curam |
| a) Up to 200 ml available on a PSO | | | | |
| b) Wastage claimable – see rule 3.3.2 on page 13 | | | | |
| | | | | |
| BENZATHINE BENZYLPENICILLIN | | | | |
| Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj | | | | |
| available on a PSO | | 10 | 4 | Bicillin LA |
| | | 10 | • | |
| Bicillin LA to be Sole Supply on 1 October 2015 | | | | |
| | | | | |

| | Subsidy | | Fully Brand or |
|--|----------------------------|--------------|-----------------------------------|
| | (Manufacturer's Pric \$ | xe) S Per | ubsidised Generic Manufacturer |
| | • | | |
| BENZYLPENICILLIN SODIUM (PENICILLIN G) Inj 600 mg (1 million units) vial – Up to 5 inj available on a | | | |
| PSO | 10.35 | 10 | Sandoz |
| FLUCLOXACILLIN | | 10 | <u>oundoz</u> |
| Cap 250 mg – Up to 30 cap available on a PSO | 18 70 | 250 | Staphlex |
| Staphlex to be Sole Supply on 1 October 2015 | | 200 | • oupliex |
| Cap 500 mg | | 500 | Staphlex |
| Staphlex to be Sole Supply on 1 October 2015 | | | |
| Grans for oral liq 25 mg per ml | 2.29 | 100 ml | 🖌 AFT |
| a) Up to 200 ml available on a PSO | | | |
| b) Wastage claimable – see rule 3.3.2 on page 13 | | | |
| c) AFT to be Sole Supply on 1 October 2015 | | | |
| Grans for oral lig 50 mg per ml | | 100 ml | 🖌 AFT |
| a) Up to 200 ml available on a PSO | | | |
| b) Wastage claimable – see rule 3.3.2 on page 13 | | | |
| c) AFT to be Sole Supply on 1 October 2015 | | | |
| Inj 250 mg vial | 8.80 | 10 | Flucloxin |
| Inj 500 mg vial | | 10 | ✓ Flucloxin |
| Inj 1 g vial – Up to 10 inj available on a PSO | 5.80 | 5 | ✓ DBL Flucloxacillin |
| | 11.60 | 10 | Flucloxin |
| PHENOXYMETHYLPENICILLIN (PENICILLIN V) | | | |
| Cap 250 mg – Up to 30 cap available on a PSO | 2.88 | 50 | Cilicaine VK |
| Cap 500 mg | | 50 | ✓ Cilicaine VK |
| a) Up to 20 cap available on a PSO | | 00 | |
| b) Up to 2 x the maximum PSO quantity for RFPP – see ru | le 5 2 6 on page 1 | 7 | |
| Grans for oral lig 125 mg per 5 ml | | 100 ml | 🖌 AFT |
| a) Up to 200 ml available on a PSO | | | - <u></u> |
| b) Wastage claimable – see rule 3.3.2 on page 13 | | | |
| Grans for oral lig 250 mg per 5 ml | | 100 ml | 🖌 AFT |
| a) Up to 300 ml available on a PSO | | | |
| b) Up to 2 x the maximum PSO quantity for RFPP - see ru | le 5.2.6 on page 1 | 7 | |
| c) Wastage claimable – see rule 3.3.2 on page 13 | 1.0 | | |
| PROCAINE PENICILLIN | | | |
| Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO | 123 50 | 5 | Cilicaine |
| | | Ŭ | • • |
| Tetracyclines | | | |
| DOXYCYCLINE | | | |
| * Tab 50 mg – Up to 30 tab available on a PSO | | 30 | |
| | (6.00) | | Doxy-50 |
| * Tab 100 mg – Up to 30 tab available on a PSO | 6.75 | 250 | ✓ Doxine |
| MINOCYCLINE HYDROCHLORIDE | | | |
| * Tab 50 mg - Additional subsidy by Special Authority see | | | |
| SA1355 below – Retail pharmacy | 5.79 | 60 | |
| | (12.05) | | Mino-tabs |
| * Cap 100 mg | | 100 | |
| | (52.04) | | Minomycin |

SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|-----|---------------------|-------------------------------------|
| TETRACYCLINE – Special Authority see SA1332 below – Retai Cap 500 mg | | 30 | ~ T | etracyclin Wolff §29 |

➡SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

| For topical antibiotics, refer to DERMATOLOGICALS, page 63 | | |
|---|----------|----------------------------|
| CIPROFLOXACIN | | |
| Recommended for patients with any of the following: i) microbiologically confirmed and clinically significant pseudomonas infec ii) prostatitis; or | tion; or | |
| iii) pyelonephritis; or | | |
| iv) gonorrhoea. | | |
| Tab 250 mg – Up to 5 tab available on a PSO1.75 | 28 | Cipflox |
| Tab 500 mg – Up to 5 tab available on a PSO2.00 | 28 | ✓ Cipflox |
| Tab 750 mg3.75 | 28 | ✓ <u>Cipflox</u> |
| CLINDAMYCIN | | |
| Cap hydrochloride 150 mg – Maximum of 4 cap per prescrip- tion; can be waived by endorsement - Retail pharmacy - | 40 | |
| Specialist | 16 | Clindamycin ABM |
| Inj phosphate 150 mg per ml, 4 ml – Retail pharmacy- Specialist100.00 | 10 | ✓ Dalacin C |
| CO-TRIMOXAZOLE | | |
| * Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - | | |
| Up to 30 tab available on a PSO | 500 | ✓ Trisul |
| * Oral lig trimethoprim 40 mg and sulphamethoxazole 200 mg | | |
| per 5 ml – Up to 200 ml available on a PSO | 100 ml | Deprim |
| COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist - Subsidy by endo | rsement | |
| Only if prescribed for dialysis or cystic fibrosis patient and the prescription is | | ordinalv. |
| Inj 150 mg65.00 | 1 | Colistin-Link |
| FUSIDIC ACID | | |
| Tab 250 mg – Retail pharmacy-Specialist | 12 | Fucidin |
| | | |

Prescriptions must be written by, or on the recommendation of, an infectious disease physician or a clinical microbiologist

| | Subsidy (Manufacturer's Price) \$ | Su Per | Fully bsidised | Brand or Generic Manufacturer |
|---|--|---|---------------------|--------------------------------------|
| GENTAMICIN SULPHATE | | | | |
| Inj 10 mg per ml, 1 ml – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient or or accordingly. | | 5 act infecti | | ospira e prescription is endorsed |
| Inj 10 mg per ml, 2 ml – Subsidy by endorsement | 175.10 | 25 | 🖌 AF | PP Pharmaceuticals S29 |
| Only if prescribed for a dialysis or cystic fibrosis patient or o accordingly. | complicated urinary tra | act infecti | on and th | e prescription is endorsed |
| Inj 40 mg per ml, 2 ml ampoule – Subsidy by endorsement a) Only if prescribed for a dialysis or cystic fibrosis patiendorsed accordingly. b) Pfizer to be Sole Supply on 1 October 2015 | | 10 rinary trae | ✓ Pf ct infectio | |
| MOXIFLOXACIN - Special Authority see SA1358 below - Retail | pharmacy | | | |
| No patient co-payment payable Tab 400 mg | 52.00 | 5 | 🖌 A) | elox |
| for applications meeting the following criteria: Either: 1 Both: 1.1 Active tuberculosis*; and 1.2 Any of the following: 1.2.1 Documented resistance to one or more firs 1.2.2 Suspected resistance to one or more firs area with known resistance), as part of reg 1.2.3 Impaired visual acuity (considered to preci 1.2.4 Significant pre-existing liver disease or her 1.2.5 Significant documented intolerance and/or or | t-line medications (tu gimen containing othe ude ethambutol use); patotoxicity from tuber | berculosi r second- or culosis m | line ager | nts; or ns; or |
| 2 Mycobacterium avium-intracellulare complex not respon | ding to other therapy | or where | such ther | apy is contraindicated.*. |
| Note: Indications marked with * are Unapproved Indications (refe Renewal only from a respiratory specialist or infectious diseases appropriate and the patient is benefiting from treatment. Initial application — (Mycoplasma genitalium) from any ref meeting the following criteria: All of the following: | r to Interpretations an specialist. Approvals | d Definiti valid for 1 | ons). year whe | ere the treatment remains |
| Has nucleic acid amplification test (NAAT) confirmed My Has tried and failed to clear infection using azithromycin; Treatment is only for 7 days. | | ; and | | |
| Initial application — (Penetrating eye injury) only from an requires prophylaxis following a penetrating eye injury and treatment Note: Indications marked with * are Unapproved Indications (reference) | ent is for 5 days only. | | | month where the patient |
| PAROMOMYCIN - Special Authority see SA1324 on the next pa | o 1 , | | | |
| Cap 250 mg | 126.00 | 16 | 🗸 Hu | umatin S29 |

| | Subsidy (Manufacturer's Price \$ | e) Sub Per | Fully sidised | Brand or Generic Manufacturer |
|---|--|-----------------|------------------|---------------------------------------|
| ►SA1324 Special Authority for Subsidy Initial application only from an infectious disease specialist or clin has confirmed cryptosporidium infection. Renewal only from an infectious disease specialist or clinical m confirmed cryptosporidium infection. | | | | |
| PYRIMETHAMINE – Special Authority see SA1328 below – Reta | il pharmacy | | | |
| Tab 25 mg | | 30 | 🖌 Da | araprim S29 |
| | 36.95 | 50 | | araprim S29 |
| ►SA1328 Special Authority for Subsidy | | | | |
| Initial application from any relevant practitioner. Approvals value the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months | or a period of 3 mon | | s notifie | d for applications meeting |
| SULFADIAZINE SODIUM - Special Authority see SA1331 below | - Retail pharmacy | | | |
| Tab 500 mg | | 56 | V W | ockhardt S29 |
| SA1331 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following: For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or | or a period of 3 mon | | s notifie | d for applications meeting |
| 3 For infants with congenital toxoplasmosis until 12 months | s of age. | | | |
| TOBRAMYCIN | | | | |
| Inj 40 mg per ml, 2 ml – Subsidy by endorsement Only if prescribed for dialysis or cystic fibrosis patient and Solution for inhalation 60 mg per ml, 5 ml – Subsidy by en- | the prescription is e | 5 ndorsed ac | | BL Tobramycin ^{y.} |
| dorsementa) Wastage claimable – see rule 3.3.2 on page 13 b) Only if prescribed for a cystic fibrosis patient and the pr | 2,200.00 | 56 dose | ✓ T | DBI |
| | | | giy. | |
| TRIMETHOPRIM * Tab 300 mg – Up to 30 tab available on a PSO TMP to be Sole Supply on 1 November 2015 | 15.00 | 50 | 🗸 TI | MP |
| VANCOMYCIN – Subsidy by endorsement | | | | |
| Only if prescribed for a dialysis or cystic fibrosis patient or for | | carditis or f | or treatr | nent of Clostridium difficile |
| following metronidazole failure and the prescription is endors | •• | | 4 | |
| Inj 500 mg | 2.64 | 1 | ✔ <u>М</u> | <u>ylan</u> |

| | Subsidy (Manufacturer's Price) \$ | S Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|----------|---------------------|---------------------------------------|
| Antifungals | | | | |
| a) For topical antifungals refer to DERMATOLOGICALS, page 63 | | | | |
| b) For topical antifungals refer to GENITO URINARY, page 76 | | | | |
| FLUCONAZOLE | | | | |
| Cap 50 mg – Retail pharmacy-Specialist | 3.49 | 28 | <u> </u> | <u>Dzole</u> |
| Cap 150 mg – Subsidy by endorsement | | 1 | | Dzole |
| a) Maximum of 1 cap per prescription; can be waived by e | | | <i>,</i> , | |
| b) Patient has vaginal candida albicans and the practition | | | | · · · · · · · · · · · · · · · · · · · |
| recommended and the prescription is endorsed according | | | | |
| Cap 200 mg – Retail pharmacy-Specialist | | 28 | <u> </u> | Dzole |
| Powder for oral suspension 10 mg per ml – Special Authority | | | | |
| see SA1359 below – Retail pharmacy | | 35 ml | ~ [| Diflucan S29 S29 |
| | 98.50 | | / [| Diflucan |
| Wastage claimable – see rule 3.3.2 on page 13 | | | | |

SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOLE

Funded for tinea vesicolor where topical treatment has not been successful and diagnosis has been confirmed by mycology, or for tinea unguium where terbinafine has not been successful in eradication or the patient is intolerant to terbinafine and diagnosis has been confirmed by mycology and the prescription is endorsed accordingly. Can be waived by endorsement - Retail pharmacy - Specialist Specialist must be an infectious disease physician, clinical microbiologist, clinical immunologist or dermatologist. Oral lig 10 mg per ml – Special Authority see SA1322 on the

next page - Retail pharmacy 141.80 150 ml OP V Sporanox

|--|

SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KFTOCONAZOI F

| Tab 200 mg - PCT - Retail pharmacy-Specialist - Subsidy by endorsement | CBS | 30 | Link Healthcare \$29 Nizoral \$29 |
|---|---------------|-----------|--|
| Prescriptions must be written by, or on the recommendation of | f an oncologi | st | |
| NYSTATIN | - | | |
| Tab 500,000 u | 14.16 | 50 | |
| | (17.09) | | Nilstat |
| Cap 500,000 u | 12.81 | 50 | |
| | (15.47) | | Nilstat |
| POSACONAZOLE - Special Authority see SA1285 below - Retail p | harmacv | | |
| Oral liq 40 mg per ml | | 105 ml OP | Noxafil |

SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy: or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Fither:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy: or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (\geq 1 mg per kilogram of body weight per day for patients with acute GVHD or \geq 0.8 mg per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE

| * Tab 250 mg – For terbinafine oral liquid formulation refer page 207 | , | 14 | ✓ <u>Dr Reddy's</u> <u>Terbinafine</u> |
|---|-------------------|-------|---|
| VORICONAZOLE - Special Authority see SA1273 on the next p | age – Retail phar | macy | |
| Tab 50 mg | | 56 | Vfend |
| Tab 200 mg | 2,930.00 | 56 | Vfend |
| Powder for oral suspension 40 mg per ml – Wastage claimable – see rule 3.3.2 on page 13 | | 70 ml | ✓ Vfend |

| Subsidy (Manufacturer's Price) | g | Fully Subsidised | Brand or Generic | |
|-----------------------------------|-----|---------------------|---------------------|--|
| \$ | Per | ~ | Manufacturer | |

➡SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

PRIMAQUINE PHOSPHATE – Special Authority see SA1326 below – Retail pharmacy

SA1326 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Antiparasitics

Antiprotozoals

| 2UININE SULPHATE ★ Tab 300 mg54. ‡ Safety cap for extemporaneously compounded oral liquid preparat | | 🗸 Q 300 |
|--|-----------|--------------------------------------|
| Antitrichomonal Agents | | |
| | | |
| IETRONIDAZOLE | | |
| Tab 200 mg – Up to 30 tab available on a PSO10. | 45 100 | Trichozole |
| Tab 400 mg18. | 15 100 | Trichozole |
| Oral lig benzoate 200 mg per 5 ml25. | 00 100 ml | FlagyI-S |
| Suppos 500 mg24. | | ✓ Flagyl |
| DRNIDAZOLE | | |
| Tab 500 mg16. | 50 10 | Arrow-Ornidazole |

| | Subsidy (Manufacturer's Price \$ | e) S Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|--|-----------------------|---------------------|---|
| Antituberculotics and Antileprotics | | | | |
| Note: There is no co-payment charge for all pharmaceuticals list immigration status. | ted in the Antituber | culotics a | nd Antilep | rotics group regardless of |
| CLOFAZIMINE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat | tion of, an infectious | s disease | physiciar | n, clinical microbiologist or |
| dermatologist. * Cap 50 mg | | 100 | 🖌 La | amprene S29 |
| CYCLOSERINE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician. Cap 250 mg | | s disease 100 | | n, clinical microbiologist or |
| DAPSONE – Retail pharmacy-Specialist | | | • | |
| a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat dermatologist | | | | - |
| Tab 25 mg Tab 100 mg | | 100 100 | | apsone apsone |
| ETHAMBUTOL HYDROCHLORIDE – Retail pharmacy-Specialis a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician Tab 100 mg Tab 400 mg | tion of, an infectious | s disease 56 56 | ✓ M | ı, clinical microbiologist or yambutol yambutol |
| ISONIAZID – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation biologist, dermatologist or public health physician | on of, an internal me | dicine ph | ysician, pa | aediatrician, clinical micro- |
| * Tab 100 mg PSM to be Sole Supply on 1 October 2015 | | 100 | ✔ P | SM |
| * Tab 100 mg with rifampicin 150 mg Rifinah to be Sole Supply on 1 October 2015 | | 100 | 🗸 R | ifinah |
| * Tab 150 mg with rifampicin 300 mg Rifinah to be Sole Supply on 1 October 2015 | | 100 | 🖌 R | ifinah |
| PARA-AMINO SALICYLIC ACID – Retail pharmacy-Specialist a) No patient co-payment payable b) Specialist must be an infectious disease specialist, clinical Grans for oral lig 4 g sachet | | spiratory 30 | | aser 529 |
| PROTIONAMIDE – Retail pharmacy-Specialist a) No patient co-payment payable | | | | |
| b) Specialist must be an infectious disease specialist, clinical Tab 250 mg | | spiratory 100 | | eteha S29 |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|--|--------------------|----------------------------|--|
| PYRAZINAMIDE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendati respiratory physician | ion of, an infectious | diseas | se physician | , clinical microbiologist or |
| Tab 500 mg – For pyrazinamide oral liquid formulation refer, page 207 | | 100 | 🗸 Al | FT-Pyrazinamide |
| RIFABUTIN – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendati gastroenterologist * Cap 150 mg – For rifabutin oral liquid formulation refer, page | | disea | se physiciar | n, respiratory physician or |
| 207 | | 30 | ✓ <u>M</u> | <u>ycobutin</u> |
| RIFAMPICIN – Subsidy by endorsement a) No patient co-payment payable b) For confirmed recurrent Staphylococcus aureus infection in based on susceptibilities and the prescription is endorsed as Specialist. Specialist must be an internal medicine physicial health physician. | ccordingly; can be w n, clinical microbiolc | vaived ogist, c | by endorse dermatologis | ment - Retail pharmacy - st, paediatrician, or public |
| * Tab 600 mg | | 30 100 | ✓ <u>Ri</u> | |
| * Cap 150 mg * Cap 300 mg | | 100 | ✓ <u>Ri</u> ✓ Ri | |
| * Oral liq 100 mg per 5 ml | | 60 ml | Ri | |
| Antivirals | | | | |
| For eye preparations refer to Eye Preparations, Anti-Infective Prep | parations, page 200 | | | |
| Hepatitis B Treatment | | | | |
| ADEFOVIR DIPIVOXIL – Special Authority see SA0829 below – F Tab 10 mg | | 30 | V He | epsera |

➡SA0829 Special Authority for Subsidy

Initial application only from a gastroenterologist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg+); and Documented resistance to lamivudine, defined as:
- 2 Patient has raised serum ALT (> 1 \times ULN); and
- 3 Patient has HBV DNA greater than 100,000 copies per mL, or viral load \geq 10 fold over nadir; and
- 4 Detection of M204I or M204V mutation; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 Patient is cirrhotic; and
 - 5.1.2 adefovir dipivoxil to be used in combination with lamivudine; or
 - 5.2 Both:
 - 5.2.1 Patient is not cirrhotic; and
 - 5.2.2 adefovir dipivoxil to be used as monotherapy.

| Subs | | Fully | Brand or | _ |
|-------------|----------------------|-------|--------------|---|
| (Manufactur | urer's Price) Subsid | lised | Generic | |
| \$ | \$ Per | ~ | Manufacturer | |

continued...

Renewal only from a gastroenterologist or infectious disease specialist. Approvals valid for 2 years where in the opinion of the treating physician, treatment remains appropriate and patient is benefiting from treatment.

Notes: Lamivudine should be added to adefovir dipivoxil if a patient develops documented resistance to adefovir dipivoxil, defined as:

- i) raised serum ALT (> 1 $\times\,$ ULN); and
- ii) HBV DNA greater than 100,000 copies per mL, or viral load \geq 10 fold over nadir; and
- iii) Detection of N236T or A181T/V mutation.

Adefovir dipivoxil should be stopped 6 months following HBeAg seroconversion for patients who were HBeAg+ prior to commencing adefovir dipivoxil.

The recommended dose of adefovir dipivoxil is no more than 10mg daily.

In patients with renal insufficiency adefovir dipivoxil dose should be reduced in accordance with the datasheet guidelines. Adefovir dipivoxil should be avoided in pregnant women and children.

ENTECAVIR - Special Authority see SA1361 below - Retail pharmacy

➡SA1361 Special Authority for Subsidy

Initial application only from a gastroenterologist or infectious disease specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B nucleoside analogue treatment-naive; and
- 3 Entecavir dose 0.5 mg/day; and
- 4 Either:
 - 4.1 ALT greater than upper limit of normal; or
 - 4.2 Bridging fibrosis (Metavir stage 3 or greater or moderate fibrosis) or cirrhosis on liver histology; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 patient has ≥ 2,000 IU HBV DNA units per mI and fibrosis (Metavir stage 2 or greater) on liver histology; and
- 6 No continuing alcohol abuse or intravenous drug use; and
- 7 Not co-infected with HCV, HIV or HDV; and
- 8 Neither ALT nor AST greater than 10 times upper limit of normal; and
- 9 No history of hypersensitivity to entecavir; and
- 10 No previous documented lamivudine resistance (either clinical or genotypic).

Notes:

- Entecavir should be continued for 6 months following documentation of complete HBeAg seroconversion (defined as loss
 of HBeAg plus appearance of anti-HBe plus loss of serum HBV DNA) for patients who were HBeAg positive prior to
 commencing this agent. This period of consolidation therapy should be extended to 12 months in patients with advanced
 fibrosis (Metavir Stage F3 or F4).
- Entecavir should be taken on an empty stomach to improve absorption.

LAMIVUDINE - Special Authority see SA1360 on the next page - Retail pharmacy

| Tab 100 mg6.00 | 28 | ✓ Zeffix |
|----------------------------|--------|----------|
| Oral liq 5 mg per ml270.00 | 240 ml | ✓ Zeffix |

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | | ubsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

SA1360 Special Authority for Subsidy

Initial application only from a gastroenterologist, infectious disease specialist, paediatrician, general physician or medical practitioner on the recommendation of a gastroenterologist, infectious disease specialist, paediatrician or general physician. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 HBV DNA positive cirrhosis prior to liver transplantation; or
- 2 HBsAg positive and have had a liver, kidney, heart, lung or bone marrow transplant; or
- 3 Hepatitis B virus naive patient who has received a liver transplant from an anti-HBc (Hepatitis B core antibody) positive donor; or
- 4 Hepatitis B surface antigen (HbsAg) positive patient who is receiving chemotherapy for a malignancy, or high dose steroids (at least 20mg/day for at least 7 days), or who has received such treatment within the previous two months; or
- 5 Hepatitis B surface antigen positive patient who is receiving anti tumour necrosis factor treatment; or
- 6 Hepatitis B core antibody (anti-HBc) positive patient who is receiving rituximab plus high dose steroids (e.g. R-CHOP).

Renewal only from a gastroenterologist, infectious disease specialist, paediatrician, general physician or medical practitioner on the recommendation of a gastroenterologist, infectious disease specialist, paediatrician or general physician. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

Renewal for patients who have maintained continuous treatment and response to lamivudine

- 1 All of the following:
 - 1.1 Have maintained continuous treatment with lamivudine; and
 - 1.2 Most recent test result shows continuing biochemical response (normal ALT); and
 - 1.3 HBV DNA <100,000 copies per ml by quantitative PCR at a reference laboratory; or

Renewal when given in combination with adefovir dipivoxil for patients with cirrhosis and resistance to lamivudine 2 All of the following:

- All of the following.
 - 2.1 Lamivudine to be used in combination with adefovir dipivoxil; and
 - 2.2 Patient is cirrhotic; and

Documented resistance to lamivudine, defined as:

- 2.3 Patient has raised serum ALT (> 1 \times ULN); and
- 2.4 Patient has HBV DNA greater than 100,000 copies per mL, or viral load = 10 fold over nadir; and
- 2.5 Detection of M204I or M204V mutation; or

Renewal when given in combination with adefovir dipivoxil for patients with resistance to adefovir dipivoxil All of the following:

- 3 All of the following:
 - 3.1 Lamivudine to be used in combination with adefovir dipivoxil; and Documented resistance to adefovir, defined as:
 - 3.2 Patient has raised serum ALT (> 1 \times ULN); and
 - 3.3 Patient has HBV DNA greater than 100,000 copies per mL, or viral load = 10 fold over nadir; and
 - 3.4 Detection of N236T or A181T/V mutation.

Herpesvirus Treatments

ACICLOVIR

| A CONTRACTOR OF A CONTRACTOR OFTA CONT | | |
|--|----|---------|
| * Tab dispersible 200 mg1.78 | 25 | Lovir |
| * Tab dispersible 400 mg5.98 | 56 | 🖌 Lovir |
| * Tab dispersible 800 mg6.64 | 35 | ✓ Lovir |
| | | • |
| VALACICLOVIR – Special Authority see SA1363 on the next page – Retail pharma | су | |
| Tab 500 mg102.72 | 30 | Valtrex |
| ····· • • • • • • • • • • • • • • • • • | | |

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | Su | lbsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

SA1363 Special Authority for Subsidy

Initial application — (recurrent genital herpes) from any medical practitioner. Approvals valid for 12 months where the patient has genital herpes with 2 or more breakthrough episodes in any 6 month period while treated with aciclovir 400 mg twice daily.

Renewal — (recurrent genital herpes) from any medical practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (ophthalmic zoster) from any medical practitioner. Approvals valid without further renewal unless notified where the patient has previous history of ophthalmic zoster and the patient is at risk of vision impairment.

Initial application — (CMV prophylaxis) from any medical practitioner. Approvals valid for 3 months where the patient has undergone organ transplantation.

Initial application — (immunocompromised patients) from any medical practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patients is immunocompromised; and
- 2 Patient has herpes zoster; and
- 3 Valaciclovir is to be given for a maximum of 7 days per course.

VALGANCICLOVIR - Special Authority see SA1404 below - Retail pharmacy

| Tab 450 mg | 60 | Valcyte |
|------------|--------|---------|

SA1404 Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
- 2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

- Both:
 - 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
 - 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

continued...

- 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
- 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
- 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis B/ HIV/AIDS Treatment

TENOFOVIR DISOPROXIL FUMARATE - Subsidy by endorsement; can be waived by Special Authority see SA1362 below

Endorsement for treatment of HIV: Prescription is deemed to be endorsed if tenofovir disoproxil fumarate is co-prescribed with another anti-retroviral subsidised under Special Authority SA1364 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Tenofovir disoproxil furnarate prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA1364, page 108

SA1362 Special Authority for Waiver of Rule

Initial application — (Chronic Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 All of the following:
 - 1.1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
 - 1.2 Patient has had previous lamivudine, adefovir or entecavir therapy; and
 - 1.3 HBV DNA greater than 20,000 IU/mL or increased \geq 10 fold over nadir; and
 - 1.4 Any of the following:
 - 1.4.1 Lamivudine resistance detection of M204I/V mutation; or
 - 1.4.2 Adefovir resistance detection of A181T/V or N236T mutation; or
 - 1.4.3 Entecavir resistance detection of relevant mutations including I169T, L180M T184S/A/I/L/G/C/M, S202C/G/I, M204V or M250I/V mutation; or
- 2 Patient is either listed or has undergone liver transplantation for HBV; or
- 3 Patient has decompensated cirrhosis with a Mayo score >20.

Initial application — (Pregnant, Active hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient is HBsAg positive and pregnant; and
- 2 HBV DNA > 20,000 IU/mL and ALT > ULN.

Renewal — (Confirmed Hepatitis B following funded tenofovir treatment for pregnancy within the previous two years) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

continued...

| Subsic (Manufacture | | |
|------------------------|-----|--------------|
| (Manuadae \$ | Per | Manufacturer |

continued...

- 1 All of the following:
 - 1.1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
 - 1.2 Patient has had previous lamivudine, adefovir or entecavir therapy; and
 - 1.3 HBV DNA greater than 20,000 IU/mL or increased $\geq~$ 10 fold over nadir; and
 - 1.4 Any of the following:
 - 1.4.1 Lamivudine resistance detection of M204I/V mutation; or
 - 1.4.2 Adefovir resistance detection of A181T/V or N236T mutation; or
 - 1.4.3 Entecavir resistance detection of relevant mutations including I169T, L180M T184S/A/I/L/G/C/M, S202C/G/I, M204V or M250I/V mutation; or
- 2 Patient is either listed or has undergone liver transplantation for HBV.

Renewal — (Subsequent pregnancy or Breastfeeding, Active hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient is HBsAg positive and pregnant or breastfeeding; and
- 2 HBV DNA > 20,000 IU/mL and ALT > ULN.

Initial application — (Pregnant, prevention of vertical transmission) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is HBsAg positive and pregnant; and
- 2 HBV DNA > 20 million IU/mL and ALT normal.

Renewal — (Subsequent pregnancy, prevention of vertical transmission) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is HBsAg positive and pregnant; and
- 2 HBV DNA > 20 million IU/mL and ALT normal.

Notes:

- Tenofovir disoproxil fumarate should be stopped 6 months following HBeAg seroconversion for patients who were HBeAg
 positive prior to commencing this agent and 6 months following HBsAg seroconversion for patients who were HBeAg
 negative prior to commencing this agent.
- The recommended dose of Tenofovir disoproxil fumarate for the treatment of all three indications is 300 mg once daily.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Tenofovir disoproxil fumarate dose should be reduced in accordance with the approved Medsafe datasheet guidelines.
- · Tenofovir disoproxil fumarate is not approved for use in children.

Hepatitis C Treatment

| BOCEPREVIR - Special Authority see SA1402 below - Retail pha | armacy | | |
|--|----------|-----|--------------------|
| Cap 200 mg - Wastage claimable - see rule 3.3.2 on page | | | |
| 13 | 5,015.00 | 336 | ✓ <u>Victrelis</u> |
| | | | |

SA1402 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, first-line) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has not received prior pegylated interferon treatment; and
- 3 Patient has IL-28B genotype CT or TT; and
- 4 Patient is to be treated in combination with pegylated interferon and ribavirin; and

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | S | ubsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

- 5 Patient is hepatitis C protease inhibitor treatment-naive; and
- 6 Maximum of 44 weeks therapy.

Initial application — (chronic hepatitis C - genotype 1, second-line) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has received pegylated interferon treatment; and
- 3 Any of the following:
 - 3.1 Patient was a responder relapser; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received pegylated interferon prior to 2004; and
- 4 Patient is to be treated in combination with pegylated interferon and ribavirin; and
- 5 Maximum of 44 weeks therapy.

Notes:

- Due to risk of severe sepsis boceprevir should not be initiated if either Platelet count < 100 x10⁹ /l or Albumin <35 g/l
- The wastage rule applies to boceprevir to allow dispensing to occur more frequently than monthly

Antiretrovirals

➡SA1364 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Confirmed HIV infection; and
- 2 Any of the following:
 - 2.1 Symptomatic patient; or
 - 2.2 Patient aged 12 months and under; or
 - 2.3 Both:
 - 2.3.1 Patient aged 1 to 5 years; and
 - 2.3.2 Any of the following:
 - 2.3.2.1 CD4 counts < 1000 cells/mm³; or
 - 2.3.2.2 CD4 counts $< 0.25 \times$ total lymphocyte count; or
 - 2.3.2.3 Viral load counts > 100000 copies per ml; or
 - 2.4 Both:
 - 2.4.1 Patient aged 6 years and over; and
 - 2.4.2 CD4 counts $< 500 \text{ cells/mm}^3$.

Notes: Tenofovir disoproxil fumarate prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals.

Subsidies for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

1 Prevention of maternal foetal transmission; or

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | S | ubsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil fumarate prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals.

Subsidies for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil fumarate prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals.

Subsidies for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria: Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil fumarate prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals.

Subsidies for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

| | Subsidy | | Fully Brand or |
|---|-------------------------|-------------------|--|
| | (Manufacturer's F \$ | Price) Sub Per | sidised Generic Manufacturer |
| Non-nucleosides Reverse Transcriptase Inhibito | rs | | |
| EFAVIRENZ – Special Authority see SA1364 on page 108 – Reta | ail pharmacy | | |
| Tab 50 mg Stocrin to be Sole Supply on 1 October 2015 | 63.38 | 30 | ✓ Stocrin S29 |
| Tab 200 mg Stocrin to be Sole Supply on 1 October 2015 | | 90 | ✓ Stocrin |
| Tab 600 mg Stocrin to be Sole Supply on 1 October 2015 | | 30 | ✓ Stocrin |
| Oral liq 30 mg per ml ETRAVIRINE – Special Authority see SA1364 on page 108 – Rel | | 180 ml OP | ✓ Stocrin S29 |
| Tab 200 mg | | 60 | ✓ Intelence |
| NEVIRAPINE – Special Authority see SA1364 on page 108 – Re Tab 200 mg – Brand switch fee payable (Pharmacode | | | |
| 2433265) - see page 204 for details | 95.94 | 60 | Nevirapine Alphapharm |
| Oral suspension 10 mg per ml | 134.55 | 240 ml | Viramune Suspension |
| Nucleosides Reverse Transcriptase Inhibitors | | | |
| ABACAVIR SULPHATE - Special Authority see SA1364 on page | | , | 4 |
| Tab 300 mg Oral lig 20 mg per ml | | 60 240 ml OP | ✓ <u>Ziagen</u> ✓ Ziagen |
| ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority Note: abacavir with lamivudine (combination tablets) counts retroviral Special Authority. | | | |
| Tab 600 mg with lamivudine 300 mg | 630.00 | 30 | Kivexa |
| DIDANOSINE [DDI] – Special Authority see SA1364 on page 108 Cap 125 mg | | acy 30 | ✔ Videx EC |
| Cap 125 mg Cap 200 mg | | 30 | Videx EC |
| Cap 250 mg | | 30 | ✓ Videx EC |
| Cap 400 mg | | 30 | ✓ Videx EC |
| EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPR – Retail pharmacy | OXIL FUMARAT | E – Special Au | uthority see SA1364 on page 108 |
| Note: Efavirenz with emtricitabine and tenofovir disoproxil fum of the anti-retroviral Special Authority | | three anti-retro | viral medications for the purposes |
| Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil fumarate 300 mg | | 30 | ✓ Atripla |
| EMTRICITABINE – Special Authority see SA1364 on page 108 – Cap 200 mg | | y 30 | ✓ Emtriva |
| EMTRICITABINE WITH TENOFOVIR DISOPROXIL FUMARATE Note: Emtricitabine with tenofovir disoproxil fumarate counts retroviral Special Authority | s as two anti-ret | roviral medicati | ions for the purposes of the anti- |
| Tab 200 mg with tenofovir disoproxil fumarate 300 mg | | 30 | Truvada |
| LAMIVUDINE – Special Authority see SA1364 on page 108 – Re Tab 150 mg | | 60 | ✓ Lamivudine |
| Oral liq 10 mg per ml | 102.50 | 240 ml OP | <u>Alphapharm</u> ✓ <u>3TC</u> |

| | Subsidy (Manufacturer's \$ | | Fully Brand or sidised Generic Manufacturer |
|--|----------------------------------|--------------------|---|
| | * | | |
| STAVUDINE [D4T] – Special Authority see SA1364 on page 10 | | • | |
| Cap 40 mg | | 60 | ✓ Zerit |
| Powder for oral soln 1 mg per ml | 100.76 | 200 ml OP | V Zerit S29 |
| ZIDOVUDINE [AZT] - Special Authority see SA1364 on page | 108 – Retail phar | macy | |
| Cap 100 mg | | 100 | Retrovir |
| Oral liq 10 mg per ml | | 200 ml OP | Retrovir |
| ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority s Note: zidovudine [AZT] with lamivudine (combination table anti-retroviral Special Authority. Tab 300 mg with lamivudine 150 mg | ets) counts as two | | |
| Protease Inhibitors | | | |
| TAZANAN/ID SLIL DHATE Special Authority and SA1264 on | naga 109 Data | il phormooy | |
| ATAZANAVIR SULPHATE – Special Authority see SA1364 on Cap 150 mg | | 60 | ✓ Reyataz |
| Cap 200 mg | | 60 | ✓ Reyataz |
| DARUNAVIR – Special Authority see SA1364 on page 108 – F | | | • |
| Tab 400 mg | | 60 | ✓ Prezista |
| Tab 600 mg | | 60 | ✓ Prezista |
| • | | 00 | • I ICLISIU |
| NDINAVIR – Special Authority see SA1364 on page 108 – Re Cap 200 mg | • • | 360 | Crixivan |
| Cap 200 mg | | 180 | Crixivan |
| | | | |
| OPINAVIR WITH RITONAVIR – Special Authority see SA136 | | | |
| Tab 100 mg with ritonavir 25 mg Tab 200 mg with ritonavir 50 mg | | 60 120 | ✓ Kaletra ✓ Kaletra |
| Oral liq 80 mg with ritonavir 20 mg per ml | | 300 ml OP | ✓ Kaletra |
| RITONAVIR – Special Authority see SA1364 on page 108 – Ri | | | • Halotta |
| Tab 100 mg | | 30 | ✓ Norvir |
| Oral liq 80 mg per ml | | 90 ml OP | ✓ Norvir |
| Strand Transfer Inhibitors | | | |
| RALTEGRAVIR POTASSIUM – Special Authority see SA1364 | on page 108 - R | etail pharmacy | |
| Tab 400 mg | | 60 | Isentress |
| Antiretrovirals - Additional Therapies | | | |
| HIV Fusion Inhibitors | | | |
| ENFUVIRTIDE – Special Authority see SA0845 below – Retail | l pharmacy | | |
| Powder for inj 90 mg per ml $	imes$ 60 | 2,380.00 | 1 | Fuzeon |
| SA0845 Special Authority for Subsidy nitial application only from a named specialist. Approvals val All of the following: | id for 3 months fo | or applications me | eeting the following criteria: |
| 1 Confirmed HIV infection; and | | | |
| 2 Enfuvirtide to be given in combination with optimized b the patient has never previously been exposed to) for t 3 Either: | | | east 1 other antiretroviral drug th |
| 3.1 Patient has evidence of HIV replication, despite | e ongoing therapy | /; or | |
| | | | continued. |

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

continued...

- 3.2 Patient has treatment-limiting toxicity to previous antiretroviral agents; and
- 4 Previous treatment with 3 different antiretroviral regimens has failed; and
- 5 All of the following:
 - 5.1 Previous treatment with a non-nucleoside reverse transcriptase inhibitor has failed; and
 - 5.2 Previous treatment with a nucleoside reverse transcriptase inhibitor has failed; and
 - 5.3 Previous treatment with a protease inhibitor has failed.

Renewal only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 Evidence of at least a 10 fold reduction in viral load at 12; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Immune Modulators

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

- a) Diagnosis
 - Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
 - · PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
 - Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Exclusion Criteria

- Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- b) Pregnancy.
- c) Neutropenia (<2.0 \times 10⁹) and/or thrombocytopenia.
- d) Continuing alcohol abuse and/or continuing intravenous drug users.

Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

INTERFERON ALFA-2A - PCT - Retail pharmacy-Specialist

a) See prescribing guideline above

| b) Prescriptions must be written by, or on the recommendation | of, an internal | medicine phys | sician or ophthalmologist |
|---|-----------------|---------------|---------------------------|
| Inj 3 m iu prefilled syringe | 31.32 | 1 | Roferon-A |

INTERFERON ALFA-2B - PCT - Retail pharmacy-Specialist

| a) See prescribing guideline above | | | |
|---|--------------------------|------------|----------------------------|
| b) Prescriptions must be written by, or on the recommendation | ation of, an internal me | edicine ph | ysician or ophthalmologist |
| Inj 18 m iu, 1.2 ml multidose pen | | 1 | ✓ Intron-A |
| Inj 30 m iu, 1.2 ml multidose pen | | 1 | Intron-A |
| Inj 60 m iu, 1.2 ml multidose pen | | 1 | Intron-A |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|---------|---------------------|-------------------------------------|
| PEGYLATED INTERFERON ALFA-2A – Special Authority see S/ | A1400 below – Retail | l pharm | acy | |
| See prescribing guideline on the previous page | | | | |
| Inj 135 mcg prefilled syringe | 1,448.00 | 4 | ✓ P | egasys |
| Inj 180 mcg prefilled syringe | 900.00 | 4 | ✓ P | egasys |
| Inj 135 mcg prefilled syringe $	imes$ 4 with ribavirin tab 200 mg $	imes$ | | | | |
| 112 | 1,799.68 | 1 OP | ✓ P | egasys RBV |
| | | | | Combination Pack |
| Inj 135 mcg prefilled syringe \times 4 with ribavirin tab 200 mg \times | | | 4 - | |
| 168 | 1,975.00 | 1 OP | | egasys RBV |
| lai 100 men avafillad avringer v 1 with site visia tak 000 men v | | | | Combination Pack |
| Inj 180 mcg prefilled syringe \times 4 with ribavirin tab 200 mg \times 112 | | 1 OP | | |
| 112 | 1,159.64 | TUP | | egasys RBV Combination Book |
| Inj 180 mcg prefilled syringe $	imes$ 4 with ribavirin tab 200 mg $	imes$ | | | | Combination Pack |
| 168 | | 1 OP | | eqasys RBV |
| 100 | 1,230.00 | I UF | | Combination Pack |
| | | | | Compination 1 dok |

SA1400 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:

| Subsidy Ful | | Fully | Brand or |
|----------------------------------|-----|--------|----------|
| (Manufacturer's Price) Subsidise | | idised | Generic |
| \$ | Per | ~ | |

continued...

- 3.1 Patient has responder relapsed; or
- 3.2 Patient was a partial responder; or
- 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA ≥ 2,000 units/ml and significant fibrosis (≥ Metavir Stage F2 or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Notes:

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

Urinary Tract Infections

| HEXAMINE HIPPURATE | | | |
|--|-------------------|----------------|--|
| * Tab 1 g | 18.40 | 100 | |
| J. J | (38.10) | | Hiprex |
| NITROFURANTOIN | | | |
| * Tab 50 mg - For nitrofurantoin oral liquid formulation refer, | | | |
| page 207 | 22.20 | 100 | Nifuran |
| * Tab 100 mg | 37.50 | 100 | Nifuran |
| NORFLOXACIN | | | |
| Tab 400 mg – Subsidy by endorsement | 13.50 | 100 | Arrow-Norfloxacin |
| Only if prescribed for a patient with an uncomplicated urinary | / tract infectior | n that is unre | sponsive to a first line agent or with |
| proven resistance to first line agents and the prescription is e | | | . 0 |

MUSCULOSKELETAL SYSTEM

| | Subsidy | | Fully Brand or |
|---|---------------------|--------|----------------------------------|
| | (Manufacturer's Pri | | bsidised Generic |
| | \$ | Per | Manufacturer |
| nticholinesterases | | | |
| | | | |
| OSTIGMINE METILSULFATE | | | <i>.</i> |
| Inj 2.5 mg per ml, 1 ml ampoule | | 50 | AstraZeneca |
| RIDOSTIGMINE BROMIDE | | | |
| Tab 60 mg | | 100 | Mestinon |
| on-Steroidal Anti-Inflammatory Drugs | | | |
| CLOFENAC SODIUM | | | |
| Tab EC 25 mg | 4.00 | 100 | Apo-Diclo |
| Tab 50 mg dispersible | | 20 | Voltaren D |
| Tab EC 50 mg | | 500 | ✓ Apo-Diclo |
| Tab long-acting 75 mg | | 500 | ✓ Diclax SR |
| Tab long-acting 100 mg | | 500 | ✓ Diclax SR |
| Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on | | 500 | |
| PSO | | 5 | ✓ Voltaren |
| Suppos 12.5 mg | | 10 | Voltaren |
| | | 10 | Voltaren |
| Suppos 25 mg | | | |
| Suppos 50 mg – Up to 10 supp available on a PSO | | 10 | Voltaren |
| Suppos 100 mg | | 10 | ✓ <u>Voltaren</u> |
| JPROFEN | | | |
| Tab 200 mg | 9.45 | 1,000 | Ibugesic |
| Tab long-acting 800 mg | 7.99 | 30 | Brufen SR |
| Oral liq 20 mg per ml | 1.89 | 200 ml | Fenpaed |
| TOPROFEN | | | |
| Cap long-acting 200 mg | 12.07 | 28 | Oruvail SR |
| | | | |
| | | | |
| Cap 250 mg | () | 20 | D . |
| | (5.60) | | Ponstan |
| | 1.25 | 50 | |
| | (9.16) | | Ponstan |
| PROXEN | | | 4 • • • • • • • • • |
| Tab 250 mg | | 500 | Noflam 250 |
| Noflam 250 to be Sole Supply on 1 October 2015 | | | 4 • • • • • • • • |
| Tab 500 mg | | 250 | Noflam 500 |
| Noflam 500 to be Sole Supply on 1 October 2015 | | | |
| Tab long-acting 750 mg | | 90 | Naprosyn SR 750 |
| Tab long-acting 1 g | 21.00 | 90 | Naprosyn SR 1000 |
| LINDAC | | | |
| Tab 100 mg | | 50 | Aclin |
| Tab 200 mg | | 50 | ✓ Aclin |
| • | | | - |
| | 0.05 | 20 | A Doutonov |
| Tab 20 mg | | 20 | ✓ <u>Reutenox</u> |
| Inj 20 mg vial | 9.95 | 1 | 🗸 AFT |
| SAIDs Other | | | |
| LOXICAM - Special Authority see SA1034 on the next page | – Retail pharmaou | | |
| Tab 7.5 mg | | 30 | Arrow-Meloxicam |
| | UG.II. | 30 | |

| Subsidy (Manufacturer's Price) | Subs | Fully | Brand or Generic |
|-----------------------------------|------|-------|---------------------|
| \$ | Per | ~ | Manufacturer |

SA1034 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has moderate to severe haemophilia with less than or equal to 5% of normal circulating functional clotting factor; and
- 2 The patient has haemophilic arthropathy; and
- 3 Pain and inflammation associated with haemophilic arthropathy is inadequately controlled by alternative funded treatment options, or alternative funded treatment options are contraindicated.

Topical Products for Joint and Muscular Pain

CAPSAICIN

| Crm 0.025% – Special Authority see SA1289 below – Retail | | |
|--|---------|---------|
| pharmacy | 25 g OP | Zostrix |
| 9.95 | 45 a OP | Zostrix |

SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

| Antirheumatoid Agents | | |
|---|-----|-----------------|
| AURANOFIN | | |
| Tab 3 mg68.99 | 60 | Ridaura s29 s29 |
| HYDROXYCHLOROQUINE | | |
| * Tab 200 mg10.50 | 100 | Plaquenil |
| Plaquenil to be Sole Supply on 1 October 2015 | | |
| LEFLUNOMIDE | | |
| Tab 10 mg55.00 | 30 | 🖌 Arava |
| Tab 20 mg76.00 | 30 | Arava |
| Tab 100 mg54.44 | 3 | Arava |
| PENICILLAMINE | | |
| Tab 125 mg61.93 | 100 | D-Penamine |
| Tab 250 mg | 100 | D-Penamine |
| SODIUM AUROTHIOMALATE | | |
| Inj 10 mg in 0.5 ml ampoule | 10 | ✓ Myocrisin |
| Inj 20 mg in 0.5 ml ampoule113.17 | 10 | ✓ Myocrisin |
| Inj 50 mg in 0.5 ml ampoule217.23 | 10 | ✓ Myocrisin |

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

SA1039 Special Authority for Subsidy

Initial application — (Underlying cause – Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD)
 - \geq 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score \leq -2.5) (see Note); or

MUSCULOSKELETAL SYSTEM

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score \leq -3.0 (see Note); or
- 5 A 10-year risk of hip fracture \geq 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or raloxifene.

Initial application — (Underlying cause – glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is receiving systemic glucocorticosteriod therapy (≥ 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD $\geq\,$ 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score $\leq\,$ -1.5) (see Note); or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for zoledronic acid (Underlying cause glucocorticosteroid therapy) or raloxifene.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year where the patient is continuing systemic glucocorticosteriod therapy (≥ 5 mg per day prednisone equivalents).

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) ≥ 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score ≤ -2.5) (see Note); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score \leq -3.0 (see Note); or
- 5 A 10-year risk of hip fracture \geq 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) or raloxifene.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score ≤ -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) In line with the Australian guidelines for funding alendronate, a vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

| | Subsidy (Manufacturer's Price) \$ | Sub: Per | Fully Brand or sidised Generic Manufacturer |
|--|--|-----------------|---|
| ALENDRONATE SODIUM – Special Authority see SA1039 on pa * Tab 70 mg | • | macy 4 | ✓ Fosamax |
| ALENDRONATE SODIUM WITH CHOLECALCIFEROL – Special * Tab 70 mg with cholecalciferol 5,600 iu | | 39 on page 4 | e 116 – Retail pharmacy V Fosamax Plus |
| Alendronate for Paget's Disease | | | |
| SA0949 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid is Both: Paget's disease; and Any of the following: Bone or articular pain; or Bone deformity; or Bone, articular or neurological complications; or Asymptomatic disease, but risk of complications or Preparation for orthopaedic surgery. Renewal from any relevant practitioner. Approvals valid for 6 moleonefiting from treatment. | due to site (base of s | kull, spine, | long bones of lower limbs); or |
| ALENDRONATE SODIUM – Special Authority see SA0949 above * Tab 40 mg | | 30 | ✓ Fosamax |
| Other Treatments | | | |
| ETIDRONATE DISODIUM – See prescribing guideline below * Tab 200 mg Arrow-Etidronate to be Sole Supply on 1 October 2015 Prescribing Guidelines | 13.50 | 100 | ✓ Arrow-Etidronate |
| Etidronate for osteoporosis should be prescribed for 14 days (400 not be taken at the same time of the day as any calcium supplement Etidronate should be taken at least 2 hours before or after any foo PAMIDRONATE DISODIUM | entation (minimum do od or fluid, except wa | ose – 500 r | |
| Inj 3 mg per ml, 10 ml vial Inj 6 mg per ml, 10 ml vial Inj 9 mg per ml, 10 ml vial | | 1 1 1 | ✓ Pamisol ✓ Pamisol ✓ Pamisol |
| RALOXIFENE HYDROCHLORIDE – Special Authority see SA113 * Tab 60 mg | | armacy 28 | ✓ Evista |

➡SA1138 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) \geq 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score \leq -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score \leq -3.0 (see Notes); or

MUSCULOSKELETAL SYSTEM

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

- 5 A 10-year risk of hip fracture \geq 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a prior Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or alendronate (Underlying cause - Osteoporosis).

Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score ≤ -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM

| Tab 35 mg | 4.00 | 4 | Risedronate Sandoz |
|---|-------|---|--|
| TERIPARATIDE - Special Authority see SA1139 below - Retail pharma | асу | | |
| Inj 250 mcg per ml, 2.4 ml4 | 90.00 | 1 | ✓ Forteo |

SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, vial − Special Authority see SA1187 on the next page − Retail pharmacy600.00 100 ml OP ✓ Aclasta

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

SA1187 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) ≥ 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score ≤ -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score \leq -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture ≥ 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) or raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (≥ 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD $\geq\,$ 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score $\leq\,$ -1.5) (see Note); or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) or raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined); and

MUSCULOSKELETAL SYSTEM

| , |
|---|
| |

continued...

2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
 - 1 The patient is continuing systemic glucocorticosteriod therapy (≥ 5 mg per day prednisone equivalents); and
 - 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) ≥ 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score ≤ -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score \leq -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture ≥ 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) or raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score ≤ -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

| ALL | OPURINOL | | |
|-----|---|-------|-----------------|
| * | Tab 100 mg15.11 | 1,000 | Apo-Allopurinol |
| * | Tab 300 mg – For allopurinol oral liquid formulation refer, | | |
| | page 20715.91 | 500 | Apo-Allopurinol |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|-----|---------------------|-------------------------------------|
| BENZBROMARONE – Special Authority see SA1537 below – Re Tab 100 mg | | 100 | 🗸 В | enzbromaron AL |

SA1537 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 Both:
 - 2.3.1 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Notes); and
 - 2.3.2 The patient has a rate of creatinine clearance greater than or equal to 20 ml/min; or
 - 2.4 All of the following:
 - 2.4.1 The patient is taking azathioprine and requires urate-lowering therapy; and
 - 2.4.2 Allopurinol is contraindicated; and
 - 2.4.3 Appropriate doses of probenecid are ineffective or probenecid cannot be used due to reduced renal function; and
- 3 The patient is receiving monthly liver function tests.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

Notes: Benzbromarone has been associated with potentially fatal hepatotoxicity.

In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

The New Zealand Rheumatology Association has developed information for prescribers which can be accessed from its website at www.rheumatology.org.nz/downloads/Benzbromarone-prescriber-information-NZRA-V2.pdf

| COLCHICINE * Tab 500 mcg | 10.08 | 100 | Colgout |
|--|-------|-----|------------------------------|
| FEBUXOSTAT – Special Authority see SA1538 on the next page | | | • <u>••••</u> |
| Tab 80 mg | , , , | 28 | Adenuric |
| Tab 120 mg | | 28 | Adenuric |

.

MUSCULOSKELETAL SYSTEM

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

SA1538 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note).

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

PROBENECID

| * Tab 500 mg | | 100 | ✓ Probenecid-AFT |
|------------------------------|--|-----|--|
| Muscle Relaxants | | | |
| BACLOFEN | | | |
| 5 | oral liquid formulation refer, page | 100 | ✓ Pacifen |
| Inj 0.05 mg per ml, 1 ml arr | npoule – Subsidy by endorsement | 1 | Lioresal Intrathecal |
| caused intolerable side | se in a programmable pump in patients where oral effects and the prescription is endorsed according be Sole Supply on 1 October 2015 | 1 0 | ents have been ineffective or have |
| Inj 2 mg per ml, 5 ml ampo | ule – Subsidy by endorsement209.29 | 1 | Lioresal Intrathecal |
| | in a programmable pump in patients where oral effects and the prescription is endorsed according | | ents have been ineffective or have |
| DANTROLENE | | | |
| | | 100 | Dantrium |
| * Cap 50 mg | | 100 | Dantrium |
| ORPHENADRINE CITRATE | 10 51 | 100 | A Novflow |
| iad 100 mg | | 100 | Norflex |

| | Subsidy (Manufacturer's Price) \$ | Fi Subsidis Per | ully Brand or eed Generic ✓ Manufacturer |
|---|---|-----------------------|--|
| Agents for Parkinsonism and Related Disorders | | | |
| Dopamine Agonists and Related Agents | | | |
| AMANTADINE HYDROCHLORIDE | 00.04 | | 40 |
| ▲ Cap 100 mg | | 60 • | Symmetrel |
| APOMORPHINE HYDROCHLORIDE ▲ Inj 10 mg per ml, 2 ml ampoule | 119.00 | 5 🖌 | Apomine |
| BROMOCRIPTINE MESYLATE | | • | Apointic |
| * Tab 2.5 mg | | 100 | Apo-Bromocriptine |
| ENTACAPONE | | | |
| ▲ Tab 200 mg | | 100 | Entapone |
| Entapone to be Sole Supply on 1 October 2015 | | | - |
| EVODOPA WITH BENSERAZIDE | | | |
| * Tab dispersible 50 mg with benserazide 12.5 mg | | | Madopar Rapid |
| Cap 50 mg with benserazide 12.5 mg | | | Madopar 62.5 |
| Cap 100 mg with benserazide 25 mg | | | Madopar 125 |
| Cap long-acting 100 mg with benserazide 25 mg | | | Madopar HBS |
| Cap 200 mg with benserazide 50 mg | 25.00 | 100 | Madopar 250 |
| LEVODOPA WITH CARBIDOPA | | | |
| * Tab 100 mg with carbidopa 25 mg – For levodopa with car- | | | 4 |
| bidopa oral liquid formulation refer, page 207 | | | Kinson |
| | | - | Sinemet |
| * Tab long-acting 200 mg with carbidopa 50 mg | | | Sinemet CR |
| * Tab 250 mg with carbidopa 25 mg | | 100 | Sinemet |
| LISURIDE HYDROGEN MALEATE | | | |
| ▲ Tab 200 mcg | 25.00 | 30 🖌 | Dopergin |
| PRAMIPEXOLE HYDROCHLORIDE | | | |
| ▲ Tab 0.25 mg | 7.20 | 100 🖌 | Ramipex |
| Tab 1 mg | 24.39 | 100 🖌 | Ramipex |
| ROPINIROLE HYDROCHLORIDE | | | |
| ▲ Tab 0.25 mg | 2.36 | 100 | Apo-Ropinirole |
| ▲ Tab 1 mg | | | Apo-Ropinirole |
| ▲ Tab 2 mg | | | Apo-Ropinirole |
| ▲ Tab 5 mg | 14.48 | 100 | Apo-Ropinirole |
| SELEGILINE HYDROCHLORIDE | | | |
| * Tab 5 mg | | 100 | Apo-Selegiline |
| | | | Apo-Selegiline |
| | | | S29 S29 |
| OLCAPONE | | | |
| ▲ Tab 100 mg | 126.20 | 100 | Tasmar |
| | | | 140110 |

| NERVOUS S | SYSTEM |
|-----------|--------|
|-----------|--------|

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-------------------|--------------------------|-------------------------------------|
| Anticholinergics | | | | |
| BENZTROPINE MESYLATE Tab 2 mg Inj 1 mg per ml, 2 ml a) Up to 5 inj available on a PSO b) Only on a PSO | | 60 5 | | enztrop ogentin |
| PROCYCLIDINE HYDROCHLORIDE Tab 5 mg | 7.40 | 100 | 🗸 Ke | emadrin |
| Agents for Essential Tremor, Chorea and Relate | ed Disorders | | | |
| RILUZOLE – Special Authority see SA1403 below – Retail phar Wastage claimable – see rule 3.3.2 on page 13 Tab 50 mg | | 56 | 🖌 Ri | lutek |
| nitial application only from a neurologist or respiratory speciollowing criteria: All of the following: The patient has amyotrophic lateral sclerosis with disea The patient has at least 60 percent of predicted forced v The patient has not undergone a tracheostomy; and The patient has not experienced respiratory failure; and Any of the following: The patient is ambulatory; or The patient is able to use upper limbs; or The patient is able to swallow. Renewal from any relevant practitioner. Approvals valid for 18 m All of the following: The patient has not undergone a tracheostomy; and | use duration of 5 years vital capacity within 2 r I | or less nonths | s; and s prior to the | initial application; and |
| 3.3 The patient is able to swallow. TETRABENAZINE Tab 25 mg Anaesthetics Local | 118.00 | 112 | ✓ <u>M</u> | otetis |
| IDOCAINE [LIGNOCAINE] Gel 2%, 10 ml urethral syringe – Subsidy by endorsement a) Up to 5 each available on a PSO b) Subsidised only if prescribed for urethral or cervical ac | | 10 rescrip | ✓ Pf ption is endored | |

| | Subsidy (Manufacturer's Price \$ | e) Per | Fully Subsidised | Generic |
|---|--|-----------|---------------------|--------------------|
| LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE | | | | |
| Oral (viscous) soln 2% | | 200 ml | V | Kylocaine Viscous |
| Inj 1%, 5 ml ampoule – Up to 25 inj available on a PSO | 8.75 | 25 | v 1 | idocaine-Claris |
| | 17.50 | 50 | | |
| | (35.00) | | 2 | Kylocaine |
| Inj 2%, 5 ml ampoule – Up to 5 inj available on a PSO | 6.90 | 25 | 1 | Lidocaine-Claris |
| Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO | 2.40 | 1 | 1 | Lidocaine-Claris |
| | 12.00 | 5 | | |
| | (20.00) | | 2 | Kylocaine |
| Inj 2%, 20 ml ampoule – Up to 5 inj available on a PSO | 2.40 | 1 | ~ | Lidocaine-Claris |
| LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE | | | | |
| Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes - | | | | |
| Subsidy by endorsement | | 10 | ~ | Pfizer |
| a) Up to 5 each available on a PSO | | | • | |
| b) Subsidised only if prescribed for urethral or cervical adm | ninistration and the r | orescrir | ntion is end | lorsed accordingly |

for urethral or cervical a ingiy.

Topical Local Anaesthetics

➡SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

| LIDOCAINE [LIGNOCAINE] - Special Authority see SA0906 above | - Retail phar | macy | |
|---|---------------|----------------|-------------|
| Crm 4% | 27.00 | 30 g OP | 🖌 LMX4 |
| Crm 4% (5 g tubes) | 27.00 | 5 | 🖌 LMX4 |
| LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Authorit | y see SA0906 | 6 above – Reta | il pharmacy |
| Crm 2.5% with prilocaine 2.5% | 45.00 | 30 g OP | 🖌 EMLA |
| Crm 2.5% with prilocaine 2.5% (5 g tubes) | 45.00 | 5 | 🖌 EMLA |

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 115

Non-opioid Analgesics

| For aspirin & chloroform application refer Standard Formulae, page ASPIRIN | 210 | | |
|--|-----------------|---------------|----------------------------------|
| * Tab EC 300 mg | 2.00 (8.50) | 100 | Aspec 300 |
| * Tab dispersible 300 mg - Up to 30 tab available on a PSO | () | 100 | ✓ <u>Ethics Aspirin</u> |
| CAPSAICIN – Subsidy by endorsement | | | |
| Subsidised only if prescribed for post-herpetic neuralgia or dia accordingly. | abetic peripher | al neuropathy | and the prescription is endorsed |
| Crm 0.075% | 12.50 | 45 g OP | ✓ Zostrix HP |
| NEFOPAM HYDROCHLORIDE Tab 30 mg | 23.40 | 90 | ✔ Acupan |

NERVOUS SYSTEM

| | Subsidy | | Fully Brand or |
|--|-------------------------|-------------------|----------------------------------|
| | (Manufacturer's F \$ | Price) Sul Per | osidised Generic Manufacturer |
| | Ŷ | | |
| PARACETAMOL | | | |
| * Tab 500 mg – Up to 30 tab available on a PSO | 8.47 | 1,000 | Pharmacare |
| *‡ Oral liq 120 mg per 5 ml | 4.15 | 1,000 ml | ✓ Paracare |
| a) Up to 200 ml available on a PSO | | | |
| b) Not in combination | | | |
| *‡ Oral liq 250 mg per 5 ml | 4.35 | 1,000 ml | Paracare Double |
| | | | Strength |
| a) Up to 100 ml available on a PSO | | | |
| b) Not in combination | | | |
| * Suppos 125 mg | | 20 | Panadol |
| * Suppos 250 mg | | 20 | Panadol |
| * Suppos 500 mg | 20.70 | 50 | Paracare |
| Opioid Analgesics | | | |
| CODEINE PHOSPHATE – Safety medicine; prescriber may d | etermine dispensing | g frequency | |
| Tab 15 mg | 4.75 | 100 | ✓ <u>PSM</u> |
| Tab 30 mg | 5.80 | 100 | ✓ <u>PSM</u> |
| Tab 60 mg | 12.50 | 100 | ✓ <u>PSM</u> |
| DIHYDROCODEINE TARTRATE | | | |
| Tab long-acting 60 mg | | 60 | DHC Continus |
| | | | · <u></u> |
| FENTANYL | | | |
| a) Only on a controlled drug form | | | |
| b) No patient co-payment payable | fraguanay | | |
| c) Safety medicine; prescriber may determine dispensing | | 10 | Boucher and Muir |
| Inj 50 mcg per ml, 2 ml ampoule | | 10 | |
| Boucher and Muir to be Sole Supply on 1 October 2015 Inj 50 mcg per ml, 10 ml ampoule | | 10 | Boucher and Muir |
| Boucher and Muir to be Sole Supply on 1 October 2015 | | 10 | |
| Patch 12.5 mcg per hour | | 5 | Fentanyl Sandoz |
| Patch 25 mcg per hour | | 5 | ✓ Fentanyl Sandoz |
| Patch 50 mcg per hour | | 5 | ✓ Fentanyl Sandoz |
| Patch 75 mcg per hour | | 5 | ✓ Fentanyl Sandoz |
| Patch 100 mcg per hour | | 5 | Fentanyl Sandoz |
| | | 5 | |
| METHADONE HYDROCHLORIDE | | | |
| a) Only on a controlled drug form | | | |
| b) No patient co-payment payable | | | |
| c) Safety medicine; prescriber may determine dispensing | | | |
| d) Extemporaneously compounded methadone will only b | e reimbursed at the | rate of the ch | eapest form available (methadone |
| powder, not methadone tablets). | | | |
| e) For methadone hydrochloride oral liquid refer Standard | | | |
| Tab 5 mg | 1.85 | 10 | Methatabs |
| Methatabs to be Sole Supply on 1 October 2015 | | 000 1 | |
| Oral liq 2 mg per ml Diadage to be Cale Currents on 1 October 2015 | 5.55 | 200 ml | Biodone |
| Biodone to be Sole Supply on 1 October 2015 | | 000 1 | |
| Oral liq 5 mg per ml Dia lang 5 mg per ml | 5.00 | 200 ml | Biodone Forte |
| Biodone Forte to be Sole Supply on 1 October 2015 | | 000 1 | |
| Oral liq 10 mg per ml Diadage Ender to be 20th 20th 20th 20th 20th 20th 20th 20th | | 200 ml | Biodone Extra Forte |
| Biodone Extra Forte to be Sole Supply on 1 October 20 | | 40 | |
| Inj 10 mg per ml, 1 ml | 61.00 | 10 | 🖌 AFT |
| | | | |

| | Subsidy (Manufacturer's Pi \$ | rice) Su Per | Fully Brand or bsidised Generic ✔ Manufacturer |
|--|-------------------------------------|-----------------|--|
| IORPHINE HYDROCHLORIDE | | | |
| a) Only on a controlled drug form | | | |
| b) No patient co-payment payable | | | |
| c) Safety medicine; prescriber may determine dispensing frequ | lency | | |
| Oral liq 1 mg per ml | 8.84 | 200 ml | RA-Morph |
| RA-Morph to be Sole Supply on 1 November 2015 | | | - |
| Oral liq 2 mg per ml | 14.00 | 200 ml | RA-Morph |
| RA-Morph to be Sole Supply on 1 November 2015 | | | |
| Oral liq 5 mg per ml | | 200 ml | RA-Morph |
| RA-Morph to be Sole Supply on 1 November 2015 | | | |
| Oral liq 10 mg per ml | | 200 ml | RA-Morph |
| RA-Morph to be Sole Supply on 1 November 2015 | | | |
| IORPHINE SULPHATE | | | |
| a) Only on a controlled drug form | | | |
| b) No patient co-payment payable | | | |
| c) Safety medicine; prescriber may determine dispensing frequ | lency | | |
| Tab immediate-release 10 mg | | 10 | Sevredol |
| Tab long-acting 10 mg | 1.95 | 10 | Arrow-Morphine LA |
| Tab immediate-release 20 mg | 5.52 | 10 | Sevredol |
| Tab long-acting 30 mg | 2.98 | 10 | ✓ Arrow-Morphine LA |
| Tab long-acting 60 mg | 5.75 | 10 | Arrow-Morphine LA |
| Tab long-acting 100 mg | 6.45 | 10 | Arrow-Morphine LA |
| Cap long-acting 10 mg | 1.70 | 10 | ✓ m-Eslon |
| Cap long-acting 30 mg | 2.50 | 10 | ✓ <u>m-Eslon</u> |
| Cap long-acting 60 mg | 5.40 | 10 | ✓ <u>m-Eslon</u> |
| Cap long-acting 100 mg | 6.38 | 10 | ✓ <u>m-Eslon</u> |
| Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO | 12.48 | 5 | DBL Morphine |
| | | | Sulphate |
| Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a | | | |
| PSO | 9.09 | 5 | DBL Morphine |
| | | | Sulphate |
| Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a | | | |
| PSO | 9.77 | 5 | DBL Morphine |
| | | | Sulphate |
| Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a | | | |
| PSO | 12.43 | 5 | DBL Morphine |
| | | | Sulphate |
| ORPHINE TARTRATE | | | |
| a) Only on a controlled drug form | | | |
| b) No patient co-payment payable | | | |
| c) Safety medicine; prescriber may determine dispensing frequ | iency | | |
| Inj 80 mg per ml, 1.5 ml | | 5 | ✓ Hospira |
| Inj 80 mg per ml, 5 ml | 107.67 | 5 | ✓ Hospira |

| | Subsidy (Manufacturer's Price | | Fully Subsidised | Generic |
|---|----------------------------------|--------|---------------------|--|
| | \$ | Per | ~ | Manufacturer |
| XYCODONE HYDROCHLORIDE | | | | |
| a) Only on a controlled drug form | | | | |
| b) No patient co-payment payable | | | | |
| c) Safety medicine; prescriber may determine dispensing | | | | |
| Tab controlled-release 5 mg | | 20 | | OxyContin |
| Tab controlled-release 10 mg | 6.75 | 20 | | Oxycodone ControlledRelease Tablets(BNM) |
| Tab controlled-release 20 mg | 11.50 | 20 | | Oxycodone ControlledRelease Tablets(BNM) |
| Tab controlled-release 40 mg | 18.50 | 20 | | Oxycodone ControlledRelease Tablets(BNM) |
| Tab controlled-release 80 mg | | 20 | | Oxycodone ControlledRelease Tablets(BNM) |
| Cap immediate-release 5 mg OxyNorm to be Sole Supply on 1 November 2015 | 1.98 | 20 | | OxyNorm |
| Cap immediate-release 10 mg OxyNorm to be Sole Supply on 1 November 2015 | 3.91 | 20 | | OxyNorm |
| Cap immediate-release 20 mg OxyNorm to be Sole Supply on 1 November 2015 | 6.84 | 20 | | OxyNorm |
| Oral liq 5 mg per 5 ml | 11.20 | 250 ml | ~ | OxyNorm |
| Inj 10 mg per ml, 1 ml | 10.08 | 5 | ~ | Oxycodone Orion |
| Inj 10 mg per ml, 2 ml | 19.87 | 5 | | Oxycodone Orion |
| Inj 50 mg per ml, 1 ml | 60.00 | 5 | | OxyNorm |
| ARACETAMOL WITH CODEINE – Safety medicine; prescrib | per may determine disp | ensina | freauencv | |
| Tab paracetamol 500 mg with codeine phosphate 8 mg | | 1,000 | | Paracetamol + Codeine (Relieve) |
| ETHIDINE HYDROCHLORIDE | | | | |
| a) Only on a controlled drug form | | | | |
| b) No patient co-payment payable | , | | | |
| c) Safety medicine; prescriber may determine dispensing | · · | 40 | | |
| Tab 50 mg | | 10 | | PSM |
| Tab 100 mg | | 10 | | PSM DBL Dathiding |
| Inj 50 mg per ml, 1 ml – Up to 5 inj available on a PSO | | 5 | - | DBL Pethidine Hydrochloride |
| Inj 50 mg per ml, 2 ml – Up to 5 inj available on a PSO | 5.83 | 5 | | DBL Pethidine Hydrochloride |
| RAMADOL HYDROCHLORIDE | | | | |
| Tab sustained-release 100 mg | | 20 | - | Tramal SR 100 |
| Tab sustained-release 150 mg | | 20 | - | Tramal SR 150 |
| Tab sustained-release 200 mg | | 20 | V _ | Tramal SR 200 |
| Cap 50 mg – For tramadol hydrochloride oral liquid form | | 100 | | A |
| tion refer, page 207 | 2.50 | 100 | V | Arrow-Tramadol |

‡ safety cap *Three months or six months, as applicable, dispensed all-at-once

NERVOUS SYSTEM

| | Subsidy | | Fully | |
|---|-----------------------------|----------|-------------|---------------------------------------|
| | (Manufacturer's Price \$ |) Per | Subsidised | I Generic Manufacturer |
| Antidepressants | | | | |
| Cyclic and Related Agents | | | | |
| MITRIPTYLINE – Safety medicine; prescriber may determine | dispensing frequency | | | |
| Tab 10 mg | | 100 | ~ | Arrow Amitriptyline |
| Tab 25 mg | | 100 | | Arrow-Amitriptyline |
| Tab 50 mg | 2.82 | 100 | | Arrow-Amitriptyline |
| CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; presc | riber may determine o | lispens | sina freaue | encv |
| Tab 10 mg | | 100 | 0 1 | Apo-Clomipramine |
| Apo-Clomipramine to be Sole Supply on 1 October 2015 | | | | · · · · · · · · · · · · · · · · · · · |
| Tab 25 mg | | 100 | ~ | Apo-Clomipramine |
| Apo-Clomipramine to be Sole Supply on 1 October 2015 | | | | |
| OTHIEPIN HYDROCHLORIDE - Safety medicine; prescriber | may determine dispe | nsing fr | requency | |
| Tab 75 mg | | 100 | | Dopress |
| Cap 25 mg | 6.17 | 100 | | Dopress |
| OXEPIN HYDROCHLORIDE - Safety medicine; prescriber ma | av determine dispens | na frec | nuency | |
| Cap 10 mg | • | 100 | | Anten |
| Cap 25 mg | | 100 | | Anten |
| Cap 50 mg | | 100 | V | Anten |
| MIPRAMINE HYDROCHLORIDE - Safety medicine; prescriber | r may determine disp | nsina | frequency | |
| Tab 10 mg | , , | 60 | | Tofranil s29 s29 |
| | 5.48 | 50 | - | Tofranil |
| | 10.96 | 100 | | Tofranil |
| Tab 25 mg | | 50 | - | Tofranil |
| IAPROTILINE HYDROCHLORIDE – Safety medicine; prescrib | | | | |
| Tab 25 mg | | 30 | | Ludiomil |
| 1ab 25 mg | 12.53 | 50 | | Ludiomil |
| | 25.06 | 100 | | Ludiomil |
| Tab 75 mg | | 20 | - | Ludiomil |
| ······································ | 21.01 | 30 | | Ludiomil |
| IIANSERIN HYDROCHLORIDE – Safety medicine; prescriber | may datarmina disna | ncina f | reguency | |
| Tab 30 mg – Subsidy by endorsement | | 30 | | Tolvon |
| Subsidised for patients who were taking mianserin hydroc | | | - | |
| ingly. Pharmacists may annotate the prescription as end hydrochloride. Note that supply of mianserin hydrochlori | orsed where there ex | ists a r | ecord of p | rior dispensing of mianser |
| there will be no stock of mianserin available beyond Nove | ember 2015. | | | |
| IORTRIPTYLINE HYDROCHLORIDE – Safety medicine; presc | | dispen | o 1 | |
| Tab 10 mg | | 100 | | Norpress |
| Tab 25 mg | 9.00 | 180 | ~ | Norpress |
| Monoamine-Oxidase Inhibitors (MAOIs) - Non S | elective | | | |
| PHENELZINE SULPHATE | | | | |
| ₭ Tab 15 mg | 95.00 | 100 | ~ | Nardil |
| RANYLCYPROMINE SULPHATE | | | | |
| Kan Teo Tr Homine Soler Hare | 22 94 | 50 | ~ | Parnate |
| | | | | |

| | Subsidy (Manufacturer's Price) \$ | S Per | Subsidised (| Brand or Generic Manufacturer |
|---|---|-----------|------------------|-------------------------------------|
| Monoamine-Oxidase Type A Inhibitors | | | | |
| IOCLOBEMIDE Note: There is a significant cost differential between moclob expensive). For depressive syndromes it is therefore more c ing prescribing moclobemide. | | | | |
| Tab 150 mg Apo-Moclobemide to be Sole Supply on 1 November 201 | | 500 | 🖌 Аро | -Moclobemide |
| K Tab 300 mg Apo-Moclobernide to be Sole Supply on 1 November 201. | | 100 | 🖌 Аро | -Moclobemide |
| Selective Serotonin Reuptake Inhibitors | | | | |
| ITALOPRAM HYDROBROMIDE | | | | |
| ₭ Tab 20 mg SCITALOPRAM | 2.34 | 84 | 🖌 Arro | ow-Citalopram |
| * Tab 10 mg | 1.40 | 28 | ✔ Air I ✔ Lox | Flow Products alate |
| Air Flow Products to be Sole Supply on 1 October 2015 k Tab 20 mg | 2.40 | 28 | ✔ Air I ✔ Lox | Flow Products |
| Air Flow Products to be Sole Supply on 1 October 2015 Loxalate Tab 10 mg to be delisted 1 October 2015) Loxalate Tab 20 mg to be delisted 1 October 2015) | | | U LOA | |
| LUOXETINE HYDROCHLORIDE Tab dispersible 20 mg, scored – Subsidy by endorsement Subsidised by endorsement | 2.50 | 30 | ✓ <u>Arro</u> | ow-Fluoxetine |
| When prescribed for a patient who cannot swallow whole or | e tablets or capsules a | ind the p | prescription i | s endorsed accordingl |
| 2) When prescribed in a daily dose that is not a multiple of Note: Tablets should be combined with capsules to facili | | | | leemed to be endorsed |
| ₭ Cap 20 mg | 1.74 | 90 | ✓ <u>Arro</u> | ow-Fluoxetine |
| AROXETINE HYDROCHLORIDE ₭ Tab 20 mg | 4.32 | 90 | 🖌 Lox | amine |
| ERTRALINE | | | | |
| K Tab 50 mg K Tab 100 mg | | 90 90 | | ow-Sertraline |
| Other Antidepressants | | | | |
| IIRTAZAPINE – Special Authority see SA0994 on the next page Tab 30 mg Tab 45 mg | 8.78 | 30 30 | ✔ Ava ✔ Ava | |

NERVOUS SYSTEM

| (| Subsidy (Manufacturer's Price) | | Fully ubsidised | Brand or Generic | |
|---|-----------------------------------|-----|--------------------|---------------------|--|
| | \$ | Per | ~ | Manufacturer | |

SA0994 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has a severe major depressive episode; and
- 2 Either:
 - 2.1 The patient must have had a trial of two different antidepressants and was unable to tolerate the treatments or failed to respond to an adequate dose over an adequate period of time (usually at least four weeks); or
 - 2.2 Both:
 - 2.2.1 The patient is currently a hospital in-patient as a result of an acute depressive episode; and
 - 2.2.2 The patient must have had a trial of one other antidepressant and either could not tolerate it or failed to respond to an adequate dose over an adequate period of time.

Renewal from any relevant practitioner. Approvals valid for 2 years where the patient has a high risk of relapse (prescriber determined).

VENLAFAXINE

| Tab 37.5 mg | 5.06 | 28 | Arrow-Venlafaxine XR |
|---|-------|----|---|
| Tab 75 mg | 6.44 | 28 | Arrow-Venlafaxine XR |
| Tab 150 mg | 8.86 | 28 | Arrow-Venlafaxine XR |
| Tab 225 mg | 14.34 | 28 | Arrow-Venlafaxine XR |
| Cap 37.5 mg - Special Authority see SA1061 below - Retail | | | |
| pharmacy | 5.69 | 28 | Efexor XR |
| Cap 75 mg – Special Authority see SA1061 below – Retail | | | |
| pharmacy | 11.40 | 28 | Efexor XR |
| Cap 150 mg - Special Authority see SA1061 below - Retail | | | |
| pharmacy | 13.98 | 28 | Efexor XR |
| | | | |

➡SA1061 Special Authority for Subsidy

Initial application only from a relevant specialist or vocationally registered general practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has 'treatment-resistant' depression; and
- 2 Either:
 - 2.1 The patient must have had a trial of two different antidepressants and have had an inadequate response from an adequate dose over an adequate period of time (usually at least four weeks); or
 - 2.2 Both:
 - 2.2.1 The patient is currently a hospital in-patient as a result of an acute depressive episode; and
 - 2.2.2 The patient must have had a trial of one other antidepressant and have had an inadequate response from an adequate dose over an adequate period of time.

Renewal from any medical practitioner. Approvals valid for 2 years where the patient has a high risk of relapse (prescriber determined).

Antiepilepsy Drugs

Agents for Control of Status Epilepticus

| CLONAZEPAM - Safety medicine; prescriber may determine disper | nsing frequency | | |
|---|-----------------|---|------------------------------|
| Inj 1 mg per ml, 1 ml | 19.00 | 5 | Rivotril |

NERVOUS SYSTEM

| | Subsidy (Manufacturer's Price) \$ |) Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|------------------------------------|----------------------|---|
| DIAZEPAM – Safety medicine; prescriber may determine dispens Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement a) Up to 5 inj available on a PSO b) Only on a PSO | | 5 | ✔ Н | ospira |
| c) PSO must be endorsed "not for anaesthetic procedures" Rectal tubes 5 mg – Up to 5 tube available on a PSO Rectal tubes 10 mg – Up to 5 tube available on a PSO | | 5 5 | | tesolid tesolid |
| PARALDEHYDE * Inj 5 ml | 1,500.00 | 5 | 🗸 A | FT |
| PHENYTOIN SODIUM * Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO Hospira to be Sole Supply on 1 November 2015 | | 5 | ✔ Н | ospira |
| Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a PSO Hospira to be Sole Supply on 1 November 2015 | | 5 | ✔ Н | ospira |
| Control of Epilepsy | | | | |
| CARBAMAZEPINE * Tab 200 mg | 16.98 34.58 39.17 | 100 100 100 100 250 ml | ✓ Te ✓ Te ✓ Te | egretol egretol CR egretol egretol CR egretol |
| CLOBAZAM – Safety medicine; prescriber may determine disper Tab 10 mg ‡ Safety cap for extemporaneously compounded oral liquid | 9.12 | 50 | 🖌 Fi | risium |
| CLONAZEPAM – Safety medicine; prescriber may determine disp ‡ Oral drops 2.5 mg per ml | | 0 ml Ol | ⊃ 🖌 R | ivotril |
| ETHOSUXIMIDE * Cap 250 mg | | 200 200 ml | | arontin arontin |
| GABAPENTIN – Special Authority see SA1477 below – Retail ph Cap 100 mg | | 100 | | rrow-Gabapentin upentin |
| Cap 300 mg – For gabapentin oral liquid formulation refer, page 207 | | 100 | | rrow-Gabapentin |
| ▲ Cap 400 mg | 13.75 | 100 | 🖌 A | upentin rrow-Gabapentin upentin |

➡SA1477 Special Authority for Subsidy

Initial application — (Epilepsy) from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:

Either:

- 1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
- 2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents.

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

continued...

Note: "Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Initial application — (Neuropathic pain or Chronic Kidney Disease associated pruritus) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

- Either:
 - 1 The patient has been diagnosed with neuropathic pain; or
 - 2 Both:
 - 2.1 The patient has Chronic Kidney Disease Stage 5-associated pruritus* where no other cause for pruritus can be identified (e.g. scabies, allergy); and
 - 2.2 The patient has persistent pruritus not relieved with a trial of emollient/moisturising creams alone.

Renewal — (Epilepsy) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life.

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Renewal — (Neuropathic pain or Chronic Kidney Disease associated pruritus) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- Either:
 - 1 The patient has demonstrated a marked improvement in their control of pain or itch (prescriber determined); or
 - 2 The patient has previously demonstrated clinical responsiveness to gabapentin and has now developed neuropathic pain in a new site.

Note: Indications marked with * are Unapproved Indications (see Interpretations and Definitions). Dosage adjustment of gabapentin is recommended for patients with renal impairment.

| GABAPENTIN (NEURONTIN) - Special Authority see SA0973 belo | w – Retail pha | irmacy | |
|--|----------------|--------|-------------------------------|
| ▲ Tab 600 mg | 67.50 | 100 | Neurontin |
| ▲ Cap 100 mg | 13.26 | 100 | Neurontin |
| ▲ Cap 300 mg – For gabapentin (neurontin) oral liquid formu- | | | |
| lation refer, page 207 | 39.76 | 100 | Neurontin |
| ▲ Cap 400 mg | 53.01 | 100 | Neurontin |

SA0973 Special Authority for Subsidy

Notes: Subsidy for patients pre-approved by PHARMAC on 1 August 2009. Approvals valid without further renewal unless notified. No new approvals will be granted from 1 August 2009.

LACOSAMIDE – Special Authority see SA1125 below – Retail pharmacy

| ▲ Tab 50 mg | | 14 | Vimpat |
|--------------|--------|----|----------------------------|
| ▲ Tab 100 mg | | 14 | Vimpat |
| - | 200.24 | 56 | Vimpat |
| ▲ Tab 150 mg | 75.10 | 14 | Vimpat |
| - | 300.40 | 56 | Vimpat |
| ▲ Tab 200 mg | | 56 | Vimpat |

➡SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

| Subsidy | | Fully | Brand or | |
|------------------------|-----|----------|--------------|--|
| (Manufacturer's Price) | Sub | bsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed guality of life from the patient's perspective.

LAMOTRIGINE

| LAMOTRIGINE | | |
|--|--------|---------------------------------------|
| ▲ Tab dispersible 2 mg6.74 | 30 | Lamictal |
| ▲ Tab dispersible 5 mg | 30 | Lamictal |
| 15.00 | 56 | Arrow-Lamotrigine |
| ▲ Tab dispersible 25 mg | 56 | Logem |
| 20.40 | | Arrow-Lamotrigine |
| | | V Mogine |
| 29.09 | | Lamictal |
| ▲ Tab dispersible 50 mg | 56 | Logem |
| 34.70 | | Arrow-Lamotrigine |
| | | ✓ Mogine |
| 47.89 | | ✓ Lamictal |
| ▲ Tab dispersible 100 mg | 56 | ✓ Logem |
| 59.90 | 00 | Arrow-Lamotrigine |
| 00.00 | | ✓ Mogine |
| 79.16 | | ✓ Lamictal |
| | | Lannetai |
| LEVETIRACETAM | | |
| Tab 250 mg24.03 | 60 | Levetiracetam-Rex |
| Tab 500 mg – For levetiracetam oral liquid formulation refer, | | |
| page 207 | 60 | Levetiracetam-Rex |
| Tab 750 mg45.23 | 60 | Levetiracetam-Rex |
| PHENOBARBITONE | | |
| For phenobarbitone oral liquid refer Standard Formulae, page 210 | | |
| * Tab 15 mg | 500 | ✔ PSM |
| * Tab 30 mg | 500 | ✓ PSM |
| - | 500 | ► FSW |
| PHENYTOIN SODIUM | | |
| * Tab 50 mg50.51 | 200 | Dilantin Infatab |
| * Cap 30 mg22.00 | 200 | Dilantin |
| * Cap 100 mg19.79 | 200 | Dilantin |
| *‡ Oral liq 30 mg per 5 ml22.03 | 500 ml | Dilantin |
| PRIMIDONE | | |
| * Tab 250 mg | 100 | Apo-Primidone |
| 5 | 100 | |
| SODIUM VALPROATE | | |
| * Tab 100 mg | 100 | Epilim Crushable |
| * Tab 200 mg EC27.44 | 100 | Epilim |
| * Tab 500 mg EC | 100 | Epilim |
| *‡ Oral liq 200 mg per 5 ml20.48 | 300 ml | Epilim S/F Liquid |
| | | Epilim Syrup |
| * Inj 100 mg per ml, 4 ml41.50 | 1 | 🖌 Epilim IV |
| | | |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-----|---------------------|-------------------------------------|
| STIRIPENTOL – Special Authority see SA1330 below – Retail pl | narmacy | | | |
| Cap 250 mg | | 60 | 🗸 D | iacomit S29 |
| Powder for oral liq 250 mg sachet | | 60 | 🗸 D | iacomit S29 |

SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

TOPIRAMATE

| ▲ Tab 25 mg | 60 | Arrow-Topiramate |
|---|-----|--------------------|
| • | | Topiramate Actavis |
| 26.04 | | Topamax |
| ▲ Tab 50 mg18.81 | 60 | Arrow-Topiramate |
| • | | Topiramate Actavis |
| 44.26 | | Topamax |
| ▲ Tab 100 mg31.99 | 60 | Arrow-Topiramate |
| | | Topiramate Actavis |
| 75.25 | | Topamax |
| ▲ Tab 200 mg55.19 | 60 | Arrow-Topiramate |
| • | | Topiramate Actavis |
| 129.85 | | Topamax |
| Sprinkle cap 15 mg20.84 | 60 | Topamax |
| Sprinkle cap 25 mg26.04 | 60 | Topamax |
| VIGABATRIN – Special Authority see SA1072 below – Retail pharmacy | | - |
| ▲ Tab 500 mg | 100 | ✓ Sabril |

►SA1072 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and

2 Either:

- 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6monthly basis thereafter); or
- 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | S | ubsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

Notes: "Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 115

Acute Migraine Treatment

| ERGOTAMINE TARTRATE WITH CAFFEINE | 100 | |
|---|---------------|------------------------------------|
| Tab 1 mg with caffeine 100 mg | 100 | Cafergot |
| RIZATRIPTAN | 00 | Discussion |
| Tab orodispersible 10 mg8.10 | 30 | ✓ <u>Rizamelt</u> |
| SUMATRIPTAN | 100 | |
| Tab 50 mg | 100 | Arrow-Sumatriptan |
| Tab 100 mg | 100 | Arrow-Sumatriptan |
| Inj 12 mg per ml, 0.5 ml cartridge – Maximum of 10 inj per prescription | 2 OP | Arrow-Sumatriptan |
| | 2 01 | |
| Prophylaxis of Migraine | | |
| For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 52 | | |
| PIZOTIFEN | | |
| * Tab 500 mcg23.21 | 100 | Sandomigran |
| Sandomigran to be Sole Supply on 1 October 2015 | | |
| Antinausea and Vertigo Agents | | |
| For Antispasmodics refer to ALIMENTARY TRACT, page 22 | | |
| APREPITANT – Special Authority see SA0987 below – Retail pharmacy | | |
| Cap 2×80 mg and 1×125 mg | 3 OP | Emend Tri-Pack |
| ►SA0987 Special Authority for Subsidy | | |
| Initial application from any relevant practitioner. Approvals valid for 12 months whe | ere the patie | nt is undergoing highly emetogenic |
| chemotherapy and/or anthracycline-based chemotherapy for the treatment of maligr | nancy. | |
| Renewal from any relevant practitioner. Approvals valid for 12 months where the patie | ent is underg | going highly emetogenic chemothe |
| apy and/or anthracycline-based chemotherapy for the treatment of malignancy. | | |
| BETAHISTINE DIHYDROCHLORIDE | | |
| * Tab 16 mg4.95 | 84 | Vergo 16 |
| | | |

‡ safety cap *Three months or six months, as applicable, dispensed all-at-once

137

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-----|---------------------|-------------------------------------|
| CYCLIZINE HYDROCHLORIDE | | | | |
| Tab 50 mg | 0.59 | 10 | 🖌 N | ausicalm |
| CYCLIZINE LACTATE Inj 50 mg per ml, 1 ml | 14.95 | 5 | 🗸 N | ausicalm |
| DOMPERIDONE | | | | |
| * Tab 10 mg – For domperidone oral liquid formulation refer, page 207 | | 100 | 🖌 P | rokinex |
| GRANISETRON | | | | |
| * Tab 1 mg | 5.98 | 50 | ✓ <u>G</u> | iranirex |
| HYOSCINE HYDROBROMIDE | | | | |
| * Inj 400 mcg per ml, 1 ml ampoule | 46.50 | 5 | 🗸 Н | ospira |
| | 93.00 | 10 | V N | lartindale S29 |
| Patch 1.5 mg – Special Authority see SA1387 below – Retail | | | | |
| pharmacy | 11.95 | 2 | ✓ <u>S</u> | copoderm TTS |

➡SA1387 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE

| * Tab 10 mg – For metoclopramide hydrochloride oral liquid | | |
|--|-----|-------------|
| formulation refer, page 2071.82 | 100 | Metamide |
| * Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO4.50 | 10 | ✓ Pfizer |
| ONDANSETRON | | |
| * Tab 4 mg5.51 | 50 | ✓ Onrex |
| * Tab disp 4 mg1.00 | 10 | Dr Reddy's |
| | | Ondansetron |
| * Tab 8 mg6.19 | 50 | ✓ Onrex |
| * Tab disp 8 mg1.50 | 10 | Ondansetron |
| | | ODT-DRLA |
| PROCHLORPERAZINE | | |
| * Tab 3 mg buccal | 50 | |
| (15.00) | | Buccastem |
| * Tab 5 mg – Up to 30 tab available on a PSO | 500 | Antinaus |
| * Inj 12.5 mg per ml, 1 ml – Up to 5 inj available on a PSO | 10 | ✓ Stemetil |
| * Suppos 25 mg23.87 | 5 | Stemetil |
| PROMETHAZINE THEOCLATE | | |
| * Tab 25 mg | 10 | |
| (6.24) | 10 | Avomine |
| (0.24) | | / |

| Subsidy | | Fully | Br |
|------------------------|-----|----------|----|
| (Manufacturer's Price) | Sul | osidised | G |
| \$ | Per | ~ | M |

Brand or Generic Manufacturer

Antipsychotics

Guidelines for the use of atypical antipsychotic agents

Diagnosis: Schizophrenia and related psychoses when positive symptoms (delusions, hallucinations and thought disorder) are prominent and/or disabling or when both positive symptoms and negative symptoms (flattened affect, emotional and social withdrawal and poverty of speech) are present. Treatment: Before initiating atypical antipsychotic therapy, physicians should consider whether the patient is likely to respond to and/or tolerate conventional antipsychotic therapy and, where appropriate, trial one or more conventional agent prior to use of an atypical agent.

General

| AMISULPRIDE - Safety medicine; prescriber may determine | e dispensing frequence | ;y | |
|---|------------------------|-------|-----------|
| Tab 100 mg | 6.22 | 30 | Solian |
| Tab 200 mg | 21.92 | 60 | Solian |
| Tab 400 mg | | 60 | Solian |
| Oral liq 100 mg per ml | | 60 ml | Solian |
| ARIPIPRAZOLE – Special Authority see SA1539 below – Re Safety medicine; prescriber may determine dispensing fr | | | |
| Tab 10 mg | | 30 | 🖌 Abilify |
| Tab 15 mg | | 30 | 🖌 Abilify |
| Tab 20 mg | 213.42 | 30 | 🖌 Abilify |
| Tab 30 mg | | 30 | Abilify |

➡SA1539 Special Authority for Subsidy

Initial application — (Schizophrenia or related psychoses) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient is suffering from schizophrenia or related psychoses; and
- 2 Either:
 - 2.1 An effective dose of risperidone or quetiapine has been trialled and has been discontinued, or is in the process of being discontinued, because of unacceptable side effects; or
 - 2.2 An effective dose of risperidone or quetiapine has been trialled and has been discontinued, or is in the process of being discontinued, because of inadequate clinical response.

Initial application — (Autism spectrum disorder*) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has been diagnosed with an autism spectrum disorder* and has symptoms of severe irritability; and
- 2 An effective dose of risperidone has been trialled and has been discontinued because of unacceptable side effects or inadequate response; and
- 3 The patient is aged less than 18 years.

Renewal — (Schizophrenia or related psychoses) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Autism spectrum disorder*) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indications marked with * are Unapproved Indications

CHLORPROMAZINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency

| Tab 10 mg - Up to 30 tab available on a PSO | | 100 | Largactil |
|---|-------|-----|-------------------------------|
| Tab 25 mg - Up to 30 tab available on a PSO | | 100 | Largactil |
| Tab 100 mg - Up to 30 tab available on a PSO | | 100 | Largactil |
| Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO | 25.66 | 10 | Largactil |

| | Subsidy (Manufacturer's Price | e) | Full Subsidise | |
|--|----------------------------------|----------|-------------------|--------------|
| | (Manulacturer's Frict | Per | Subsidise V | Manufacturer |
| OZAPINE – Hospital pharmacy [HP4] | | | | |
| Safety medicine; prescriber may determine dispensing frequ | Jency | | | |
| Tab 25 mg | | 50 | ~ | Clozaril |
| ····· | 11.36 | 100 | | Clozaril |
| | 6.69 | 50 | | Clopine |
| | 13.37 | 100 | | Clopine |
| Tab 50 mg | | 50 | | Clopine |
| ····· • • • • • • • | 17.33 | 100 | | Clopine |
| Tab 100 mg | | 50 | | Clozaril |
| | 29.45 | 100 | | Clozaril |
| | 17.33 | 50 | | Clopine |
| | 34.65 | 100 | | Clopine |
| Tab 200 mg | | 50 | | Clopine |
| 100 200 mg | | 100 | | Clopine |
| Suspension 50 mg per ml | | 100 ml | | Clopine |
| | | | • | olopille |
| LOPERIDOL – Safety medicine; prescriber may determine d | | | | |
| Tab 500 mcg – Up to 30 tab available on a PSO | 6.23 | 100 | ~ | Serenace |
| Tab 1.5 mg – Up to 30 tab available on a PSO | 9.43 | 100 | ~ | Serenace |
| Tab 5 mg – Up to 30 tab available on a PSO | | 100 | ~ | Serenace |
| Oral liq 2 mg per ml - Up to 200 ml available on a PSO | 23.84 | 100 ml | ~ | Serenace |
| Inj 5 mg per ml, 1 ml – Up to 5 inj available on a PSO | 21.55 | 10 | ~ | Serenace |
| VOMEPROMAZINE MALEATE – Safety medicine; prescriber | r may datarmina dian | oncina f | | |
| | | 100 | | Nozinan |
| Tab 25 mg | | 100 | | Nozinan |
| Tab 100 mg | | | | |
| Inj 25 mg per ml, 1 ml | | 10 | V | Nozinan |
| HIUM CARBONATE - Safety medicine; prescriber may dete | rmine dispensing fre | quency | | |
| Tab 250 mg | | 500 | ~ | Lithicarb FC |
| Lithicarb FC to be Sole Supply on 1 October 2015 | | | | |
| Tab 400 mg | | 100 | ~ | Lithicarb FC |
| Lithicarb FC to be Sole Supply on 1 October 2015 | | | | |
| Tab long-acting 400 mg | | 100 | ~ | Priadel |
| Cap 250 mg | | 100 | V | Douglas |
| | | | - | |
| ANZAPINE – Safety medicine; prescriber may determine dis | | | | _ . |
| Tab 2.5 mg | | 28 | | Zypine |
| Tab 5 mg | | 28 | | Zypine |
| Tab orodispersible 5 mg | | 28 | | Zypine ODT |
| Tab 10 mg | | 28 | | Zypine |
| Tab orodispersible 10 mg | 3.05 | 28 | ~ | Zypine ODT |
| RICYAZINE - Safety medicine; prescriber may determine dis | spensing frequency | | | |
| Tab 2.5 mg | | 100 | ~ | Neulactil |
| Tab 10 mg | | 100 | | Neulactil |
| • | | 100 | • | |
| JETIAPINE – Safety medicine; prescriber may determine disp | | | | |
| Tab 25 mg | 2.10 | 90 | ~ | Quetapel |
| Tab 100 mg | 4.20 | 90 | ~ | Quetapel |
| Tab 200 mg | 7.20 | 90 | ~ | Quetapel |
| Tab 300 mg | 12 00 | 90 | ~ | Quetapel |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Generic |
|--|---|-------|---------------------|--------------------|
| RISPERIDONE – Safety medicine; prescriber may determine disp | pensing frequency | | | |
| Tab orodispersible 0.5 mg - Special Authority see SA0927 | | | | |
| below – Retail pharmacy | | 28 | ~ | Risperdal Quicklet |
| Tab 0.5 mg | 1.90 | 60 | ~ | Actavis |
| Tab 1 mg | 2.10 | 60 | v | Actavis |
| Tab orodispersible 1 mg - Special Authority see SA0927 be- | | | | |
| low – Retail pharmacy | | 28 | ~ | Risperdal Quicklet |
| Tab 2 mg | 2.34 | 60 | v | Actavis |
| Tab orodispersible 2 mg - Special Authority see SA0927 be- | | | | |
| low – Retail pharmacy | | 28 | ~ | Risperdal Quicklet |
| Tab 3 mg | 2.55 | 60 | ~ | Actavis |
| Tab 4 mg | | 60 | v | Actavis |
| Oral liq 1 mg per ml | 9.75 | 30 ml | ~ [| Risperon |

➡SA0927 Special Authority for Subsidy

Initial application — (Acute situations) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 For a non-adherent patient on oral therapy with standard risperidone tablets or risperidone oral liquid; and
- 2 The patient is under direct supervision for administration of medicine.

Initial application — (Chronic situations) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is unable to take standard risperidone tablets or oral liquid, or once stabilized refuses to take risperidone tablets or oral liquid; and
- 2 The patient is under direct supervision for administration of medicine.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The patient is unable to take standard risperidone tablets or oral liquid, or once stabilized refuses to take risperidone tablets or oral liquid; and
- 2 The patient is under direct supervision for administration of medicine.

Note: Risperdal Quicklets cost significantly more than risperidone tablets and should only be used where necessary.

TRIFLUOPERAZINE HYDROCHLORIDE - Safety medicine; prescriber may determine dispensing frequency

| Tab 1 mg | 100 | Stelazine |
|----------|---------|-------------------------------|
| Tab 2 mg | 100 | Stelazine |
| Tab 5 mg | 100 | Stelazine |

ZIPRASIDONE - Subsidy by endorsement

a) Safety medicine; prescriber may determine dispensing frequency

b) Ziprasidone is subsidised for patients suffering from schizophrenia or related psychoses after a trial of an effective dose of risperidone or quetiapine that has been discontinued, or is in the process of being discontinued, because of unacceptable side effects or inadequate response, and the prescription is endorsed accordingly.

| Cap 20 mg Cap 40 mg | | 60 60 | Zeldox Zeldox |
|--|----------------------------|------------|--|
| Cap 60 mg Cap 80 mg | 247.17 | 60 60 | ✓ Zeldox ✓ Zeldox |
| ZUCLOPENTHIXOL HYDROCHLORIDE – Safety medic Tab 10 mg | ine; prescriber may determ | ine dispen | sing frequency |

| | Subsidy (Manufacturer's Price) \$ | S Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|----------|---------------------|-------------------------------------|
| Depot Injections | | | | |
| FLUPENTHIXOL DECANOATE – Safety medicine; prescriber ma | av determine dispensi | na frea | uencv | |
| Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PSO | / / | 5 | | uanxol |
| Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PSO | | 5 | 🖌 Fl | uanxol |
| Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO | 40.87 | 5 | 🖌 Fl | uanxol |
| FLUPHENAZINE DECANOATE - Safety medicine; prescriber ma | av determine dispensi | na frea | uency | |
| Inj 12.5 mg per 0.5 ml, 0.5 ml – Up to 5 inj available on a PSC | | 5 | | odecate |
| Inj 25 mg per ml, 1 ml - Up to 5 inj available on a PSO | | 5 | 🖌 M | odecate |
| Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO | | 5 | 🖌 M | odecate |
| HALOPERIDOL DECANOATE - Safety medicine; prescriber may | / determine dispensin | a freau | encv | |
| Inj 50 mg per ml, 1 ml – Up to 5 inj available on a PSO | | 5 | 🦳 и | aldol |
| Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO | | 5 | 🖌 Н | aldol Concentrate |
| OLANZAPINE – Special Authority see SA1428 below – Retail ph Safety medicine; prescriber may determine dispensing freque | | | | |
| Inj 210 mg vial | | 1 | V 7 | yprexa Relprevv |
| Inj 300 mg vial | | 1 | | vprexa Relprevv |
| Inj 405 mg vial | | 1 | | yprexa Relprevv |

SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

| Inj 25 mg syringe | 1 | Invega Sustenna |
|--------------------------|---|-----------------|
| Inj 50 mg syringe271.95 | 1 | Invega Sustenna |
| Inj 75 mg syringe | 1 | Invega Sustenna |
| Inj 100 mg syringe | 1 | Invega Sustenna |
| Inj 150 mg syringe435.12 | 1 | Invega Sustenna |

➡SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

| Subsidy (Manutacturer's I | Price) | Fully Subsidised | Brand or Generic | |
|------------------------------|--------|---------------------|---------------------|--|
| `\$ | Per | ~ | Manufacturer | |

continued...

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

PIPOTHIAZINE PALMITATE - Subsidy by endorsement

a) Safety medicine; prescriber may determine dispensing frequency

b) Subsidised for patients who were taking pipothiazine palmitate prior to 1 August 2014 and the prescription or PSO is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of pipothiazine palmitate.

| Inj 50 mg per ml, 1 ml – Up to 5 inj available on a PSO | 178.48 | 10 | Piportil |
|---|--------|----|------------------|
| Inj 50 mg per ml, 2 ml $-$ Up to 5 inj available on a PSO | | 10 | 🗸 Piportil |
| RISPERIDONE - Special Authority see SA1427 below - Retail p | | | |
| Safety medicine; prescriber may determine dispensing freque | ency | | |
| Inj 25 mg vial | | 1 | Risperdal Consta |
| Inj 37.5 mg vial | 178.71 | 1 | Risperdal Consta |
| Inj 50 mg vial | 217.56 | 1 | Risperdal Consta |
| | | | |

►SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

| Inj 200 mg per ml, 1 ml – Up to 5 inj available on a PSO | 5 | Clopixol |
|---|-----|-------------------|
| Anxiolytics | | |
| ALPRAZOLAM - Safety medicine; prescriber may determine dispensing frequency | | |
| Tab 250 mcg2.50 | 50 | ✓ Xanax |
| ‡ Safety cap for extemporaneously compounded oral liquid preparations. | | |
| Tab 500 mcg3.25 | 50 | ✓ Xanax |
| ‡ Safety cap for extemporaneously compounded oral liquid preparations. | | |
| Tab 1 mg5.00 | 50 | 🗸 Xanax |
| ‡ Safety cap for extemporaneously compounded oral liquid preparations. | | |
| BUSPIRONE HYDROCHLORIDE | | |
| * Tab 5 mg | 100 | Pacific Buspirone |
| * Tab 10 mg17.00 | 100 | Pacific Buspirone |
| - | | • |

| (1 | Subsidy Manufacturer's Price) \$ | Per | Full <u>y</u> Subsidised | |
|--|--|------------|-----------------------------|----------------|
| CLONAZEPAM - Safety medicine; prescriber may determine disper | 0 1 2 | | | |
| Tab 500 mcg Tab 2 mg | | 100 100 | - | Paxam Paxam |
| DIAZEPAM – Safety medicine; prescriber may determine dispensin Tab 2 mg ‡ Safety cap for extemporaneously compounded oral liquid p | 11.44 | 500 | ~ | Arrow-Diazepam |
| Tab 5 mg ‡ Safety cap for extemporaneously compounded oral liquid p | 13.71 | 500 | ~ | Arrow-Diazepam |
| LORAZEPAM – Safety medicine; prescriber may determine dispens Tab 1 mg ‡ Safety cap for extemporaneously compounded oral liquid p | 10.79 | 250 | V | <u>Ativan</u> |
| Tab 2.5 mg ‡ Safety cap for extemporaneously compounded oral liquid p | 13.88 | 100 | ~ | <u>Ativan</u> |
| DXAZEPAM - Safety medicine; prescriber may determine dispensir | 0 1 2 | | | |
| Tab 10 mg ‡ Safety cap for extemporaneously compounded oral liquid p | | 100 | V | <u>Ox-Pam</u> |
| Tab 15 mg ‡ Safety cap for extemporaneously compounded oral liquid p | 8.53 | 100 | ~ | <u>Ox-Pam</u> |
| Multiple Sclerosis Treatments | | | | |
| FINGOLIMOD – Special Authority see SA1487 below – Retail phari Wastage claimable – see rule 3.3.2 on page 13 | nacy | | | |
| Cap 0.5 mg | .2,650.00 | 28 | ~ | Gilenya |

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below). Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

| The coordinator | Phone: 04 460 4990 |
|---|---|
| Multiple Sclerosis Treatment Assessment Committee | Facsimile: 04 916 7571 |
| PHARMAC PO Box 10 254 | Email: mstaccoordinator@pharmac.govt.nz |
| | |

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- a) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- b) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- c) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of MRI activity on a scan within the past 24 months (either a contrast enhancing lesion or with new T2 lesions(s) compared with a previous scan); and
- d) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);

| Subsidy | Fu | lly Brand or | |
|-----------------|-----------------|----------------------------------|--|
| (Manufacturer's | Price) Subsidis | ed Generic | |
| \$ | Per | Manufacturer | |

- b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
- c) last at least one week;
- d) start at least one month after the onset of a previous relapse;
- e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
- f) be distinguishable from the effects of general fatigue; and
- g) not be associated with a fever (T>37.5°C); and
- e) applications must be made by the patient's neurologist or general physician; and
- f) patients must have no previous history of lack of response to fingolimod; and
- g) patients must have not previously had intolerance to fingolimod; and
- h) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; orh) 4.0 to 4.5.
- b) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- c) intolerance to fingolimod; or
- d) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab and fingolimod is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

NATALIZUMAB - Special Authority see SA1496 below - Retail pharmacy

Inj 20 mg per ml, 15 ml vial 1,750.00 1 🗸 Tysabri

SA1496 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below). Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

| The coordinator | Phone: 04 460 4990 |
|---|---|
| Multiple Sclerosis Treatment Assessment Committee | Facsimile: 04 916 7571 |
| PHARMAC PO Box 10 254 | Email: mstaccoordinator@pharmac.govt.nz |

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified). Entry Criteria

| Subsidy | Full | y Brand or | |
|------------------------|-----------|----------------------------------|--|
| (Manufacturer's Price) | Subsidise | d Generic | |
| \$ | Per 🖌 | Manufacturer | |

- a) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- b) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- c) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of MRI activity on a scan within the past 24 months (either a contrast enhancing lesion or with new T2 lesions(s) compared with a previous scan); and
- d) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T>37.5 $^{\circ}$ C); and
- e) applications must be made by the patient's neurologist or general physician; and
- f) treatment must be initiated and supervised by a neurologist who is registered in the Tysabri Australasian Prescribing Programme operated by the supplier; and
- g) patients must have no previous history of lack of response to natalizumab; and
- h) patients must have not previously had intolerance to natalizumab; and
- i) either
 - a) Patient is JC virus negative, or
 - b) Patient is JC virus positive and has given written informed consent acknowledging an understanding of the risk of progressive multifocal leucoencephalopathy (PML) associated with natalizumab
- j) patient will not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- a) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- b) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- c) intolerance to natalizumab; or
- d) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Switching between natalizumab and fingolimod is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the EDSS stopping criteria are not met.

If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

| Subsidy (Manufacturer's Price) © | Per | Fully Subsidised | Brand or Generic Manufacturer | |
|--|-----|---------------------|-------------------------------------|--|
| φ | Fei | • | Manulaciulei | |

Other Multiple Sclerosis Treatments

SA1484 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below). Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

| The coordinator | Phone: 04 460 4990 |
|---|---|
| Multiple Sclerosis Treatment Assessment Committee | Facsimile: 04 916 7571 |
| PHARMAC PO Box 10 254 | Email: mstaccoordinator@pharmac.govt.nz |

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

These agents will NOT be subsidised if dispensed from a community or hospital pharmacy. Regular supplies will be distributed to all approved patients or their clinicians by courier.

Prescribers must send quarterly prescriptions for approved patients to the MSTAC coordinator.

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, or 20 mg glatiramer acetate daily will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. The MSTAC coordinator should be notified of the change and a new prescription provided.

Entry Criteria

- a) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- b) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- c) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of MRI activity on a scan within the past 24 months (either a contrast enhancing lesion or with new T2 lesions(s) compared with a previous scan); and
- d) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T>37.5°C); and
- e) applications must be made by the patient's neurologist; and
- f) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- g) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or
 - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- h) patient will not be co-prescribed natalizumab or fingolimod.

| | Subsidy | | Fully | Brand or | |
|----|-----------------------|-------|-------|--------------|--|
| (M | lanufacturer's Price) | Subsi | dised | Generic | |
| | \$ | Per | ~ | Manufacturer | |

Stopping Criteria

Any of the following:

- a) Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment. Progression of disability is defined as progress by any of the following EDDSS Points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- b) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- c) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- d) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

Entry Criteria for patients with an EDSS of 4.5-5.5 who have not had an application for funding considered prior to 1 November 2014

- a) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis should as a rule include MRI confirmation. For patients diagnosed before MRI was widely utilised in New Zealand, confirmation of diagnosis via clinical assessment and laboratory/ancillary data must be provided; and
- b) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- c) patients must have:
 - a) EDSS score 4.5-5.5 with 2+ relapses:
 - Experienced at least 2 significant relapses of MS in the previous 12 months, and
 - An EDSS score of between 4.5-5.5; and
- d) Each relapse must:
 - a) be confirmed by a neurologist or general physician (the patient may not necessarily have been seen during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) follow a period of stability of at least one month;
 - be severe enough to change either the EDSS or at least one of the Kurtzke functional systems scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and

| Subsidy (Manufacturer's Price) | | Fully Subsidised | Brand or Generic | |
|-----------------------------------|-----|---------------------|---------------------|--|
| (Manulaclurer S Frice) | | Subsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

- g) not be associated with a fever (T>37.5°C); and
- e) applications must be made at least four weeks after the date of the onset of the last known relapse; and
- f) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate (see criteria for stopping).
- g) applications must be submitted to the Multiple Sclerosis Treatment Assessment Committee (MSTAC) by the patient's neurologist or a general physician; and
- h) patients must agree (via informed consent) to co-operate if as a result of their meeting the stopping criteria, funding is withdrawn. Patients must agree to the collection of clinical data relating to their MS and use of those data by PHARMAC; and
- patients must agree to allow clinical data to be collected and reviewed by MSTAC annually for each year in which they receive funding for beta-interferon or glatiramer acetate.

Stopping Criteria for patients with an EDSS of 4.5-5.5 who have not had an application for funding considered prior to 1 November 2014

Any of the following:

- a) Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment. Progression of disability is defined as progress by any of the following:
 - a) an increase of 1 EDSS point where starting EDSS 3.5 or greater; or
 - b) an increase in EDSS score to 6.0 or more; or
- b) stable or increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- c) pregnancy and/or lactation; or
- d) within the 12 month approval year, intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- e) non-compliance with treatment, including refusal to undergo annual assessment or refusal to allow the results of the assessment to be submitted to MSTAC; or
- f) patients may, subject to conclusions drawn from published evidence available at the time, be excluded if they develop a high titre of neutralising anti-bodies to beta-interferon or glatiramer acetate.

Note: Patients who have a stable or increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet any of the other Stopping Criteria at annual review may switch to a different class of funded treatment (i.e. patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa). Patients may switch classes of treatment for this reason only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to stable or increasing relapse rate over 12 months of treatment).

| GLATIRAMER ACETATE – Special Authority see SA1484 o Inj 20 mg prefilled syringe | | 28 | Copaxone | |
|--|------------------------|-------|-------------------------------|--|
| INTERFERON BETA-1-ALPHA - Special Authority see SA | | harm] | | |
| Inj 6 million iu prefilled syringe | 1,170.00 | 4 | Avonex | |
| Injection 6 million iu per 0.5 ml pen injector | 1,170.00 | 4 | Avonex Pen | |
| Inj 6 million iu per vial | | 4 | Avonex | |
| INTERFERON BETA-1-BETA – Special Authority see SA14 | | arm] | | |
| Inj 8 million iu per 1 ml | 1,322.89 | 15 | Betaferon | |
| Sedatives and Hypnotics | | | | |
| LORMETAZEPAM - Safety medicine; prescriber may deter | mine dispensing freque | псу | | |
| Tab 1 mg | 3.11 | 30 | | |
| | (23.50) | | Noctamid | |
| ‡ Safety cap for extemporaneously compounded ora | l liquid preparations. | | | |

| | Subsidy (Manufacturer's Price) \$ | Per | Full Subsidise | d Generic |
|--|---|-----|-------------------|----------------|
| MIDAZOLAM – Safety medicine; prescriber may determine disper | sing frequency | | | |
| Inj 1 mg per ml, 5 ml | | 10 | ~ | Pfizer |
| | 10.75 | | ~ | Hypnovel |
| Inj 5 mg per ml, 3 ml | 11.90 | 5 | ~ | Hypnovel |
| | | | ~ | Pfizer |
| NITRAZEPAM – Safety medicine; prescriber may determine dispe | nsing frequency | | | |
| Tab 5 mg | 0 1 2 | 100 | ~ | Nitrados |
| ‡ Safety cap for extemporaneously compounded oral liquid | preparations. | | | |
| PHENOBARBITONE SODIUM – Special Authority see SA1386 be | elow – Retail pharma | асу | | |
| Inj 200 mg per ml, 1 ml ampoule | | 10 | ~ | Martindale S29 |
| ■SA1386 Special Authority for Subsidy | | | | |

SA1386 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

| TEMAZEPAM – Safety medicine; prescriber may determine dispensing frequency Tab 10 mg | 25 | ✓ Normison | |
|---|-----|---------------|---|
| ‡ Safety cap for extemporaneously compounded oral liquid preparations. | | | |
| TRIAZOLAM – Safety medicine; prescriber may determine dispensing frequency | | | |
| Tab 125 mcg5.10 | 100 | | |
| (7.25) | | Hypam | |
| ‡ Safety cap for extemporaneously compounded oral liquid preparations. | | | |
| Tab 250 mcg4.10 | 100 | | |
| (8.70) | | Hypam | |
| ‡ Safety cap for extemporaneously compounded oral liquid preparations. | | | |
| ZOPICLONE – Safety medicine; prescriber may determine dispensing frequency | | | |
| Tab 7.5 mg11.90 | 500 | Apo-Zopiclone | |
| Stimulants/ADHD Treatments | | | |
| | | | |
| Stimulants/ADHD treatments | | | |
| ATOMOXETINE – Special Authority see SA1416 below – Retail pharmacy | | | 1 |
| Cap 10 mg107.03 | 28 | ✓ Strattera | |
| Cap 18 mg107.03 | 28 | ✓ Strattera | |
| Cap 25 mg | 28 | ✓ Strattera | |
| Cap 40 mg | 28 | ✓ Strattera | |
| Cap 60 mg | 28 | ✓ Strattera | |
| Cop 90 mg 120 11 | 00 | A Strattora | |

| ATOMOXETINE - Special Authority see SA1416 be | elow – Retail pharmacy | | |
|---|------------------------|----|-------------------------------|
| Cap 10 mg | | 28 | Strattera |
| Cap 18 mg | | 28 | Strattera |
| Cap 25 mg | | 28 | Strattera |
| Cap 40 mg | | 28 | Strattera |
| Cap 60 mg | | 28 | Strattera |
| Cap 80 mg | | 28 | Strattera |
| Cap 100 mg | | 28 | Strattera |
| | | | |

►SA1416 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder) diagnosed according to DSM-IV or ICD 10 criteria; and
- 2 Once-daily dosing; and
- 3 Any of the following:

| ξ | Subsidy | Fully | Brand or |
|---------|------------------|------------|--------------|
| (Manufa | acturer's Price) | Subsidised | Generic |
| | \$ Per | er 🖌 | Manufacturer |

- 3.1 Treatment with a subsidised formulation of a stimulant has resulted in the development or worsening of serious adverse reactions or where the combination of subsidised stimulant treatment with another agent would pose an unacceptable medical risk; or
- 3.2 Treatment with a subsidised formulation of a stimulant has resulted in worsening of co-morbid substance abuse or there is a significant risk of diversion with subsidised stimulant therapy; or
- 3.3 An effective dose of a subsidised formulation of a stimulant has been trialled and has been discontinued because of inadequate clinical response; or
- 3.4 Treatment with a subsidised formulation of a stimulant is considered inappropriate because the patient has a history of psychoses or has a first-degree relative with schizophrenia; and
- 4 The patient will not be receiving treatment with atomoxetine in combination with a subsidised formulation of a stimulant, except for the purposes of transitioning from subsidised stimulant therapy to atomoxetine.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: A "subsidised formulation of a stimulant" refers to currently subsidised methylphenidate hydrochloride tablet formulations (immediate-release, sustained-release and extended-release) or dexamphetamine sulphate tablets.

DEXAMFETAMINE SULFATE - Special Authority see SA1149 below - Retail pharmacy

a) Only on a controlled drug form

b) Safety medicine; prescriber may determine dispensing frequency

| Tab 5 mg16.50 100 | 🖌 PSM |
|-------------------|-------|

➡SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

| Subsidy (Manufacturer's Price) | Subs | Fully | Brand or Generic |
|-----------------------------------|------|-------|---------------------|
| \$ | Per | ~ | Manufacturer |

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1150 below - Retail pharmacy

a) Only on a controlled drug form

b) Safety medicine; prescriber may determine dispensing frequency

| 3.20 | 30 | Rubifen |
|-------|-----|--------------------------------|
| | 30 | Ritalin |
| | | Rubifen |
| 7.85 | 30 | Rubifen |
| 10.95 | 30 | Rubifen SR |
| 50.00 | 100 | Ritalin SR |
| | | |

➡SA1150 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

| | Subsidy (Manufacturer's Price) \$ | Subs Per | Fully sidised | Brand or Generic Manufacturer |
|--|---|-------------|------------------|-------------------------------------|
| METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEAS a) Only on a controlled drug form | | see SA11 | 51 belo | ow – Retail pharmacy |
| b) Safety medicine; prescriber may determine dispensing free | equency | | | |

| b) dalety medicine, presender may determine dispensi | ig inequency | | |
|--|--------------|----|------------------------------|
| Tab extended-release 18 mg | | 30 | Concerta |
| Tab extended-release 27 mg | 65.44 | 30 | Concerta |
| Tab extended-release 36 mg | 71.93 | 30 | Concerta |
| Tab extended-release 54 mg | | 30 | Concerta |
| Cap modified-release 10 mg | | 30 | Ritalin LA |
| Cap modified-release 20 mg | | 30 | Ritalin LA |
| Cap modified-release 30 mg | | 30 | Ritalin LA |
| Cap modified-release 40 mg | | 30 | Ritalin LA |
| | | | |

SA1151 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL – Special Authority see SA1126 below – Retail pharmacy

Tab 100 mg72.50 30 🗸 Modavigil

SA1126 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and

2 Either:

- 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
- 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and

| ubsidy turer's Price) Sub | Fully | Brand or Generic |
|------------------------------|-------|---------------------|
| \$ Per | ~ | Manufacturer |

3 Either:

- 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamphetamine has been trialled and discontinued because of intolerable side effects; or
- 3.2 Methylphenidate and dexamphetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

| DONEPEZIL HYDROCHLORIDE | | | |
|---|-------------------------------------|----|---------------|
| * Tab 5 mg | 5.48 | 90 | Donepezil-Rex |
| * Tab 10 mg | | 90 | Donepezil-Rex |
| RIVASTIGMINE - Special Authority see SA1488 below | Retail pharmacy | | |
| Patch 4.6 mg per 24 hour | | 30 | Exelon |
| Patch 9.5 mg per 24 hour | | 30 | Exelon |

SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

a) No patient co-payment payable

b) Safety medicine; prescriber may determine dispensing frequency

| Tab sublingual 2 mg with naloxone 0.5 mg | | 28 | Suboxone |
|--|--------|----|------------------------------|
| Tab sublingual 8 mg with naloxone 2 mg | 166.00 | 28 | Suboxone |

SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

| Subsidy | Fully | | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | I Generic | |
| \$ | Per 🖌 | Manufacturer | |

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

| Tab modified-release 150 mg | 4.97 | 30 | Zyban |
|---|------------------|-------------|---------------|
| DISULFIRAM | | | |
| Tab 200 mg | 24.30 | 100 | Antabuse |
| NALTREXONE HYDROCHLORIDE - Special Authority see SA14 | 08 below – Retai | il pharmacy | |
| Tab 50 mg | 76.00 | 30 | ✓ Naltraccord |

SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

| | Subsidy (Manufacturer's Price) \$ | S Per | Fully Subsidised | Brand or Generic Manufacturer | |
|----------|---|----------|---------------------|-------------------------------------|--|
| NICOTINE | | | | | |

NICOTINE

| Nicotine will not be funded under the Dispensing Frequency Rule in amounts I | ess than 4 w | eeks of treatment. |
|--|--------------|--------------------|
| Patch 7 mg – Up to 28 patch available on a PSO10.57 | 28 | Habitrol |
| Patch 14 mg – Up to 28 patch available on a PSO11.31 | 28 | Habitrol |
| Patch 21 mg – Up to 28 patch available on a PSO11.95 | 28 | Habitrol |
| Lozenge 1 mg - Up to 216 loz available on a PSO12.91 | 216 | ✓ Habitrol |
| Lozenge 2 mg - Up to 216 loz available on a PSO14.14 | 216 | Habitrol |
| Gum 2 mg (Classic) – Up to 384 piece available on a PSO | 384 | Habitrol |
| Gum 2 mg (Fruit) – Up to 384 piece available on a PSO22.26 | 384 | Habitrol |
| Gum 2 mg (Mint) – Up to 384 piece available on a PSO22.26 | 384 | Habitrol |
| Gum 4 mg (Classic) – Up to 384 piece available on a PSO25.67 | 384 | Habitrol |
| Gum 4 mg (Fruit) – Up to 384 piece available on a PSO25.67 | 384 | Habitrol |
| Gum 4 mg (Mint) – Up to 384 piece available on a PSO25.67 | 384 | Habitrol |

VARENICLINE TARTRATE - Special Authority see SA1161 below - Retail pharmacy

a) Varenicline will not be funded under the Dispensing Frequency Rule in amounts less than 2 weeks of treatment.

b) A maximum of 3 months' varenicline will be subsidised on each Special Authority approval.

| Tab 1 mg67.74 | 28 | Champix |
|--|-------|-----------------------------|
| 135.48 | 56 | Champix |
| Tab 0.5 mg \times 11 and 1 mg \times 1460.48 | 25 OP | Champix |

SA1161 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to guit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to guit smoking using bupropion or nortriptyline; and
- 4 The patient has not used funded varenicline in the last 12 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this: and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 3 months' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria:

All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 The patient has not used funded varenicline in the last 12 months: and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this: and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 3 months' funded varenicline (see note).

The patient must not have had an approval in the past 12 months.

Note: a maximum of 3 months' varenicline will be subsidised on each Special Authority approval.

| | Subsidy (Manufacturer's F | Price) | Fully Subsidised | Brand or Generic |
|---|------------------------------|----------|---------------------|---------------------|
| | \$ | Per | ~ | Manufacturer |
| Chemotherapeutic Agents | | | | |
| Alkylating Agents | | | | |
| BUSULPHAN – PCT – Retail pharmacy-Specialist | | | | |
| Tab 2 mg | | 100 | 🗸 M | yleran |
| CARBOPLATIN – PCT only – Specialist | | | | |
| Inj 10 mg per ml, 5 ml vial | | 1 | 🗸 D | BL Carboplatin |
| | 20.00 | | 🖌 C | arboplatin Ebewe |
| Inj 10 mg per ml, 15 ml vial | 14.05 | 1 | 🖌 D | BL Carboplatin |
| | 19.50 | | 🖌 C | arbaccord |
| | 22.50 | | | arboplatin Ebewe |
| Inj 10 mg per ml, 45 ml vial | | 1 | | BL Carboplatin |
| | 48.50 | | | arbaccord |
| | 50.00 | | | arboplatin Ebewe |
| Inj 10 mg per ml, 100 ml vial | | 1 | | arboplatin Ebewe |
| Inj 1 mg for ECP | | 1 mg | V B | axter |
| Carboplatin Ebewe Inj 10 mg per ml, 100 ml vial to be deliste | eu i September 2015 | 7 | | |
| CARMUSTINE – PCT only – Specialist | | | | |
| Inj 100 mg vial | | 1 | | iCNU |
| Inj 100 mg for ECP | 532.00 | 100 mg O | Р / В | axter |
| HLORAMBUCIL – PCT – Retail pharmacy-Specialist | | | | |
| Tab 2 mg | 29.06 | 25 | 🖌 L | eukeran FC |
| CISPLATIN – PCT only – Specialist | | | | |
| Inj 1 mg per ml, 50 ml | | 1 | V C | isplatin Ebewe |
| | | | | ospira |
| Inj 1 mg per ml, 100 ml | 21.00 | 1 | V C | isplatin Ebewe |
| | | | 🖌 Н | ospira |
| Inj 1 mg for ECP | 0.27 | 1 mg | 🖌 В | axter |
| YCLOPHOSPHAMIDE | | | | |
| Tab 50 mg – PCT – Retail pharmacy-Specialist | 79.00 | 50 | V F | ndoxan S29 |
| | 158.00 | 100 | | rocytox S29 |
| Wastage claimable - see rule 3.3.2 on page 13 | 130.00 | 100 | V F | IUCYIUX 320 |
| Inj 1 g vial – PCT – Retail pharmacy-Specialist | 35.03 | 1 | √ F | ndoxan |
| | 127.80 | 6 | | vtoxan |
| Inj 2 g vial – PCT only – Specialist | | 1 | | ndoxan |
| Inj 1 mg for ECP – PCT only – Specialist | | 1 mg | | axter |
| FOSFAMIDE – PCT only – Specialist | | 0 | | |
| Inj 1 g | 96.00 | 1 | ⊿ н | oloxan |
| Inj 2 g | | 1 | | oloxan |
| Inj 1 mg for ECP | | 1 mg | | axter |
| | | · …9 | ÷ 0 | |
| OMUSTINE – PCT – Retail pharmacy-Specialist | 120 50 | 20 | | eeNU |
| Cap 10 mg | | 20 20 | | eenu eeNU |
| Cap 40 mg | | 20 | v (| CENU |
| IELPHALAN | | | | |
| Tab 2 mg – PCT – Retail pharmacy-Specialist | | 25 | | lkeran |
| Inj 50 mg – PCT only – Specialist | 67.80 | 1 | V A | lkeran |

| (| Subsidy Manufacturer's Price) \$ | Per | Full Subsidise | d Generic |
|---|--|------|---|--|
| OXALIPLATIN – PCT only – Specialist Inj 50 mg | 15.32 | 1 | r | Oxaliplatin Actavis 50 |
| Inj 100 mg | 55.00 200.00 25.01 | 1 | ~ | Oxaliplatin Ebewe Eloxatin Oxaliplatin Actavis |
| | 110.00 400.00 | | ~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~ | 100 Oxaliplatin Ebewe Eloxatin |
| Inj 1 mg for ECP THIOTEPA – PCT only – Specialist | | 1 mg | | Baxter |
| Inj 15 mg | CBS | 1 | V | Bedford S29 THIO-TEPA S29 Tepadina S29 |
| Antimetabolites | | | | |
| AZACITIDINE – PCT only – Specialist – Special Authority see SA Inj 100 mg vial | | 1 | ~ | Vidaza |

| inj 100 mg viai | | 1 | Vidaza |
|------------------|-------|------|----------------------------|
| Inj 1 mg for ECF | 96.66 | 1 mg | Baxter |

SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- 4 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

| | Subsidy (Manufacturer's | Price) Sub | Fully Brand or osidised Generic |
|--|----------------------------|------------|---|
| | (Manulacturer 3 | Per | Manufacturer |
| ALCIUM FOLINATE | | | |
| Tab 15 mg – PCT – Retail pharmacy-Specialist | 82.45 | 10 | DBL Leucovorin Calcium |
| Inj 3 mg per ml, 1 ml – PCT – Retail pharmacy-Specialist | 17.10 | 5 | Hospira |
| Inj 50 mg – PCT – Retail pharmacy-Specialist | | 5 | <u>Calcium Folinate</u> <u>Ebewe</u> |
| Inj 100 mg – PCT only – Specialist | 7.33 | 1 | Calcium Folinate Ebewe |
| Inj 300 mg – PCT only – Specialist | 22.51 | 1 | Calcium Folinate Ebewe |
| Inj 1 g – PCT only – Specialist | 67.51 | 1 | Calcium Folinate Ebewe |
| Inj 1 mg for ECP – PCT only – Specialist | 0.06 | 1 mg | Baxter |
| APECITABINE – Retail pharmacy-Specialist | | | |
| Tab 150 mg | | 60 | ✓ <u>Capecitabine</u> Winthrop |
| Tab 500 mg | 120.00 | 120 | ✓ <u>Capecitabine</u> <u>Winthrop</u> |
| LADRIBINE – PCT only – Specialist | | | |
| Inj 1 mg per ml, 10 ml | | 7 | Leustatin |
| Inj 10 mg for ECP | 749.96 | 10 mg OP | Baxter |
| YTARABINE | | | |
| Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist | | 5 | Pfizer |
| | 80.00 | | Hospira |
| Inj 500 mg – PCT – Retail pharmacy-Specialist | | 1 | ✓ Pfizer |
| | 95.36 | 5 | Hospira |
| Inj 100 mg per ml, 10 ml vial – PCT – Retail pharmacy- | | | |
| Specialist | 8.83 | 1 | Pfizer |
| | 42.65 | | Hospira |
| Inj 100 mg per ml, 20 ml vial – PCT – Retail pharmacy- | | | |
| Specialist | 17.65 | 1 | Pfizer |
| | 34.47 | | Hospira |
| Inj 1 mg for ECP – PCT only – Specialist | | 10 mg | Baxter |
| Inj 100 mg intrathecal syringe for ECP – PCT only – Specialist | 11.00 | 100 mg OP | Baxter |
| LUDARABINE PHOSPHATE | | | |
| Tab 10 mg – PCT – Retail pharmacy-Specialist Fludara Oral to be Sole Supply on 1 October 2015 | 412.00 | 20 | Fludara Oral |
| Inj 50 mg – PCT only – Specialist | 525.00 | 5 | Fludarabine Ebewe |
| | 1,430.00 | | Fludara |
| Inj 50 mg for ECP – PCT only – Specialist | 105.00 | 50 mg OP | Baxter |
| LUOROURACIL | | | |
| Inj 50 mg per ml, 10 ml vial – PCT only – Specialist | | 5 | Fluorouracil Ebewe |
| Inj 50 mg per ml, 20 ml vial – PCT only – Specialist | | 1 | Fluorouracil Ebewe |
| Inj 25 mg per ml, 100 ml vial – PCT only – Specialist | | 1 | Hospira |
| Inj 50 mg per ml, 50 ml vial – PCT only – Specialist | | 1 | Fluorouracil Ebewe |
| Inj 50 mg per ml, 100 ml vial – PCT only – Specialist | | 1 | Fluorouracil Ebewe |
| Inj 1 mg for ECP – PCT only – Specialist Fluorouracil Ebewe Inj 50 mg per ml, 10 ml vial to be delisted 1 N | | 100 mg | Baxter |

| | Subsidy (Manufacturer's Pri | | Fully Brand or Subsidised Generic |
|--|--------------------------------|---------|---|
| | \$ | Per | Manufacturer |
| EMCITABINE HYDROCHLORIDE – PCT only – Specialist | | | |
| Inj 1 g | | 1 | Gemcitabine Ebewe |
| , 0 | 62.50 | | DBL Gemcitabine |
| | 349.20 | | Gemzar |
| lnj 200 mg | | 1 | Gemcitabine Ebewe |
| , 3 | 78.00 | | 🖌 Gemzar |
| Inj 1 mg for ECP | | 1 mg | ✓ Baxter |
| INOTECAN HYDROCHLORIDE – PCT only – Specialist | | 0 | |
| Inj 20 mg per ml, 2 ml vial | 11 50 | 1 | Irinotecan Actavis |
| 11 j 20 11 g per 11 i, 2 11 i viai | 11.50 | I | 40 |
| | 41.00 | | |
| | 41.00 | | Camptosar |
| | 17.00 | | Irinotecan-Rex |
| Inj 20 mg per ml, 5 ml vial | 17.80 | 1 | Irinotecan Actavis |
| | | | 100 |
| | 100.00 | | Camptosar |
| | | | Irinotecan-Rex |
| Inj 1 mg for ECP | 0.19 | 1 mg | Baxter |
| ERCAPTOPURINE – PCT – Retail pharmacy-Specialist | | | |
| Tab 50 mg | 49 41 | 25 | Puri-nethol |
| 0 | | 20 | |
| ETHOTREXATE | | | 4 - . |
| Tab 2.5 mg – PCT – Retail pharmacy-Specialist | 3.18 | 30 | Trexate |
| Trexate to be Sole Supply on 1 October 2015 | | | |
| Tab 10 mg – PCT – Retail pharmacy-Specialist | 21.00 | 50 | Trexate |
| Trexate to be Sole Supply on 1 October 2015 | | | |
| Inj 2.5 mg per ml, 2 ml – PCT – Retail pharmacy-Speciali | st23.65 | 5 | Hospira |
| Inj 7.5 mg prefilled syringe | 17.19 | 1 | Methotrexate |
| | | | Sandoz |
| Inj 10 mg prefilled syringe | 17.25 | 1 | Methotrexate |
| | | | <u>Sandoz</u> |
| Inj 15 mg prefilled syringe | 17.38 | 1 | Methotrexate |
| | | | Sandoz |
| Inj 20 mg prefilled syringe | 17.50 | 1 | Methotrexate |
| | | | Sandoz |
| Inj 25 mg prefilled syringe | 17.63 | 1 | Methotrexate |
| | | | Sandoz |
| Inj 30 mg prefilled syringe | 17.75 | 1 | Methotrexate |
| | | | Sandoz |
| Inj 25 mg per ml, 2 ml – PCT – Retail pharmacy-Specialis | st20.20 | 5 | Hospira |
| Inj 25 mg per ml, 20 ml - PCT - Retail pharmacy-Special | list27.78 | 1 | Hospira |
| Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specia | alist25.00 | 1 | ✓ Methotrexate Ebeween A Methotrexate Ebeween A Methotrexate Ebeween A Method A |
| Inj 100 mg per ml, 50 ml - PCT - Retail pharmacy-Specia | alist99.99 | 1 | ✓ Methotrexate Ebeween |
| Inj 1 mg for ECP - PCT only - Specialist | 0.10 | 1 mg | ✓ Baxter |
| Inj 5 mg intrathecal syringe for ECP - PCT only - Specia | | 5 mg ÖP | Baxter |
| | | 0 | |
| HOGUANINE – PCT – Retail pharmacy-Specialist | 106.01 | 05 | 1 I anvia |
| Tab 40 mg | | 25 | Lanvis |

| | Subsidy (Manufacturer's Price) \$ | Per | Ful Subsidise | d Generic |
|--|---|--------|------------------|---------------|
| Other Cytotoxic Agents | | | | |
| AMSACRINE – PCT only – Specialist | | | | |
| Inj 50 mg per ml, 1.5 ml ampoule | 1,500.00 | 6 | ~ | Amsidine S29 |
| Inj 75 mg | | 5 | ~ | AmsaLyo S29 |
| ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-Sp | pecialist | | | |
| Cap 0.5 mg | | 100 | ~ | Agrylin S29 |
| | | | | Teva S29 |
| ARSENIC TRIOXIDE – PCT only – Specialist | | | | |
| Inj 10 mg | 4.817.00 | 10 | ~ | AFT S29 |
| BLEOMYCIN SULPHATE – PCT only – Specialist | , | | | |
| Inj 15,000 iu (10 mg), vial | 150.48 | 1 | ~ | DBL Bleomycin |
| | | | | Sulfate |
| Inj 1,000 iu for ECP | | ,000 i | u 🗸 | Baxter |
| BORTEZOMIB - PCT only - Specialist - Special Authority see | SA1127 below | | | |
| Inj 1 mg | | 1 | ~ | Velcade |
| Inj 3.5 mg | 1,892.50 | 1 | ~ | Velcade |
| Inj 1 mg for ECP | 594.77 | 1 mg | ~ | Baxter |

➡SA1127 Special Authority for Subsidy

Initial application — (Treatment naive multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 The patient has treatment-naive symptomatic multiple myeloma; or
 - 1.2 The patient has treatment-naive symptomatic systemic AL amyloidosis *; and
- 2 Maximum of 9 treatment cycles.

Note: Indications marked with * are Unapproved Indications.

Initial application — (Relapsed/refractory multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 The patient has relapsed or refractory multiple myeloma; or
 - 1.2 The patient has relapsed or refractory systemic AL amyloidosis *; and
- 2 The patient has received only one prior front line chemotherapy for multiple myeloma or amyloidosis; and
- 3 The patient has not had prior publicly funded treatment with bortezomib; and
- 4 Maximum of 4 treatment cycles.

Note: Indications marked with * are Unapproved Indications.

Renewal — (Relapsed/refractory multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 The patient's disease obtained at least a partial response from treatment with bortezomib at the completion of cycle 4; and
- 2 Maximum of 4 further treatment cycles (making a total maximum of 8 consecutive treatment cycles).

Notes: Responding relapsed/refractory multiple myeloma patients should receive no more than 2 additional cycles of treatment beyond the cycle at which a confirmed complete response was first achieved. A line of therapy is considered to comprise either:

- a) a known therapeutic chemotherapy regimen and supportive treatments; or
- b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments.

Refer to datasheet for recommended dosage and number of doses of bortezomib per treatment cycle.

| | Subsidy (Manufacturer's | | Fully Brand or sidised Generic |
|--|----------------------------|----------------|---------------------------------------|
| | (Manalactarer 3 | Per | Manufacturer |
| COLASPASE [L-ASPARAGINASE] – PCT only – Specialist | | | |
| Inj 10,000 iu | | 1 | Leunase |
| Inj 10,000 iu for ECP | | 10,000 iu OP | Baxter |
| DACARBAZINE – PCT only – Specialist | | · | |
| Inj 200 mg vial | 51 84 | 1 | Hospira |
| Inj 200 mg for ECP | | 200 mg OP | ✓ Baxter |
| DACTINOMYCIN [ACTINOMYCIN D] – PCT only – Specialist | | | |
| | 145.00 | 1 | ✓ Cosmegen |
| Inj 0.5 mg vial Inj 0.5 mg for ECP | | ا 0.5 mg OP | Baxter |
| , . | 145.00 | 0.5 mg Oi | |
| DAUNORUBICIN – PCT only – Specialist | | | 4.54 |
| Inj 2 mg per ml, 10 ml | | 1 | ✓ Pfizer |
| Inj 20 mg for ECP | 118.72 | 20 mg OP | Baxter |
| DOCETAXEL – PCT only – Specialist | | | |
| Inj 20 mg | | 1 | DBL Docetaxel |
| | 48.75 | | Docetaxel Sandoz |
| Inj 20 mg per ml, 1 ml | | 1 | ✓ Taxotere |
| Inj 20 mg per ml, 4 ml | | 1 | ✓ Taxotere |
| Inj 80 mg | | 1 | ✓ DBL Docetaxel |
| laid and for ECD | 195.00 | 4 | Docetaxel Sandoz |
| Inj 1 mg for ECP | 0.01 | 1 mg | Baxter |
| DOXORUBICIN – PCT only – Specialist | | | |
| Inj 10 mg | | 1 | Doxorubicin Ebewe |
| Inj 50 mg | | 1 | Arrow-Doxorubicin |
| | 40.00 | | DBL Doxorubicin |
| | | | DBL Doxorubicin |
| | | | S29 S29 |
| | | | Doxorubicin Ebewe |
| lnj 100 mg | | 1 | Doxorubicin Ebewe |
| Inj 200 mg | | 1 | Arrow-Doxorubicin |
| | 150.00 | | Adriamycin |
| | 0.07 | 4 | Doxorubicin Ebewe |
| Inj 1 mg for ECP | 0.37 | 1 mg | Baxter |
| EPIRUBICIN – PCT only – Specialist | | | |
| lnj 2 mg per ml, 5 ml | | 1 | Epirubicin Ebewe |
| Inj 2 mg per ml, 25 ml | | 1 | DBL Epirubicin |
| | | | Hydrochloride |
| | 87.50 | | Epirubicin Ebewe |
| Inj 2 mg per ml, 50 ml | 58.20 | 1 | DBL Epirubicin |
| | | | Hydrochloride |
| | 125.00 | | Epirubicin Ebewe |
| Inj 2 mg per ml, 100 ml | 94.50 | 1 | DBL Epirubicin |
| | | | Hydrochloride |
| | 210.00 | | Epirubicin Ebewe |
| Inj 1 mg for ECP | 0.82 | 1 mg | Baxter |

| | Subsidy (Manufacturer's Price) | | Fully Subsidised | Generic |
|--|-----------------------------------|------|---------------------|--------------|
| | \$ | Per | ~ | Manufacturer |
| ETOPOSIDE | | | | |
| Cap 50 mg – PCT – Retail pharmacy-Specialist | | 20 | ~ | Vepesid |
| Cap 100 mg - PCT - Retail pharmacy-Specialist | | 10 | ~ | Vepesid |
| Inj 20 mg per ml, 5 ml - PCT - Retail pharmacy-Specialist | | 1 | ~ | Hospira |
| | 612.20 | 10 | ~ | Vepesid |
| Inj 1 mg for ECP – PCT only – Specialist | 0.30 | 1 mg | ~ | Baxter |
| ETOPOSIDE PHOSPHATE - PCT only - Specialist | | - | | |
| Inj 100 mg (of etoposide base) | 40.00 | 1 | ~ | Etopophos |
| Inj 1 mg (of etoposide base) for ECP | | 1 mg | | Baxter |
| | | | - | |
| HYDROXYUREA – PCT – Retail pharmacy-Specialist | 01 70 | 100 | | l hudua a |
| Cap 500 mg | | 100 | V | Hydrea |
| IDARUBICIN HYDROCHLORIDE | | | | |
| Inj 5 mg – PCT only – Specialist | 100.00 | 1 | ~ | Zavedos |
| Inj 10 mg – PCT only – Specialist | | 1 | ~ | Zavedos |
| Inj 1 mg for ECP – PCT only – Specialist | | 1 mg | ~ | Baxter |
| LENALIDOMIDE – Retail pharmacy-Specialist – Special Authorit | v see SA1468 below | , | | |
| Wastage claimable – see rule 3.3.2 on page 13 | , | | | |
| Cap 10 mg | 6.207.00 | 21 | ~ | Revlimid |
| Cap 25 mg | | 21 | V | Revlimid |

➡SA1468 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Either:
 - 2.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 2.2 Both:
 - 2.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 2.2.2 The patient has experienced severe (grade \geq 3), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 3 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an Unapproved Indication (refer to Interpretations and Definitions). A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MESNA

| Tab 400 mg – PCT – Retail pharmacy-Specialist | 50 | Uromitexan |
|--|--------|--------------------------------|
| Tab 600 mg – PCT – Retail pharmacy-Specialist | 50 | Uromitexan |
| Inj 100 mg per ml, 4 ml ampoule – PCT only – Specialist | 15 | Uromitexan |
| Inj 100 mg per ml, 10 ml ampoule – PCT only – Specialist | 15 | Uromitexan |
| Inj 1 mg for ECP – PCT only – Specialist | 100 mg | Baxter |

| | Subsidy (Manufacturer's Price |) | Fully Subsidised | |
|---|----------------------------------|------|---------------------|--------------------|
| | \$ | Per | | |
| MITOMYCIN C – PCT only – Specialist | | | | |
| Inj 5 mg vial | 79.75 | 1 | V | Arrow |
| Inj 1 mg for ECP | | 1 mg | ~ | Baxter |
| /ITOZANTRONE – PCT only – Specialist | | | | |
| Inj 2 mg per ml, 5 ml vial | 110.00 | 1 | ~ | Mitozantrone Ebewe |
| Inj 2 mg per ml, 10 ml vial | | 1 | ~ | Mitozantrone Ebewe |
| Inj 2 mg per ml, 12.5 ml vial | | 1 | | |
| | (413.21) | | | Onkotrone |
| Inj 1 mg for ECP | | 1 mg | ~ | Baxter |
| Inj 30 mg | 45.00 | 5 | ~ | Paclitaxel Ebewe |
| ACLITAXEL – PCT only – Specialist | | | | |
| Inj 100 mg | | 1 | • | Paclitaxel Ebewe |
| Inj 100 mg | 91.67 | 1 | - | Paclitaxel Actavis |
| lnj 150 mg | | 1 | - | Paclitaxel Ebewe |
| ing 100 mg | 137.50 | • | | Anzatax |
| | | | • | Paclitaxel Actavis |
| Inj 300 mg | | 1 | V | Paclitaxel Ebewe |
| , | 275.00 | | V | Anzatax |
| | | | V | Paclitaxel Actavis |
| Inj 600 mg | 73.06 | 1 | ~ | Paclitaxel Ebewe |
| Inj 1 mg for ECP | | 1 mg | ~ | Baxter |
| EGASPARGASE – PCT only – Special Authority see SA1325 b | | • | | |
| Inj 3,750 IU per 5 ml | | 1 | ~ | Oncaspar S29 |
| - O A 1005 Our stick Authority for Outholds | -, | | - | |

SA1325 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and
- 3 Treatment is with curative intent.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and
- 3 Treatment is with curative intent.

| PENTOSTATIN [DEOXYCOFORMYCIN] | PCT only – Specialist |
|-------------------------------|---|
|-------------------------------|---|

| Inj 10 mg | CBS | 1 | ✔ Nipent S29 |
|--|------------------|------|--------------|
| PROCARBAZINE HYDROCHLORIDE - PCT - Retail pharmacy- | Specialist | | |
| Cap 50 mg | 498.00 | 50 | Natulan S29 |
| TEMOZOLOMIDE - Special Authority see SA1063 on the next page | ge – Retail phar | macy | |
| Cap 5 mg | 8.00 | 5 | Temaccord |
| Cap 20 mg | 36.00 | 5 | Temaccord |
| Cap 100 mg | 175.00 | 5 | Temaccord |
| Cap 250 mg | 410.00 | 5 | Temaccord |

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

►SA1063 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 10 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of six cycles of 5 days treatment, at a maximum dose of 200 mg/m².

Notes: Indication marked with a * is an Unapproved Indication. Temozolomide is not subsidised for the treatment of relapsed glioblastoma multiforme. Reapplications will not be approved.

Studies of temozolomide show that its benefit is predominantly in those patients with a good performance status (WHO grade 0 or 1 or Karnofsky score >80), and in patients who have had at least a partial resection of the tumour.

| THALIDOMIDE | - PCT only - Specialist - Special Authority see SA1124 below | v | |
|-------------|--|----|------------------------------|
| Cap 50 mg | | 28 | Thalomid |
| Cap 100 mg | | 28 | Thalomid |

➡SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Fither:

1 The notiont has multipl

The patient has multiple myeloma; or
 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period. Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an Unapproved Indication.

TRETINOIN

| Cap 10 mg - PCT - Retail pharmacy-Specialist | 100 | Vesanoid |
|---|------|-----------------------------|
| VINBLASTINE SULPHATE | | |
| Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Specialist37.29 | 1 | Hospira |
| 186.46 | 5 | Hospira |
| Inj 1 mg for ECP – PCT only – Specialist | 1 mg | Baxter |
| VINCRISTINE SULPHATE | | |
| Inj 1 mg per ml, 1 ml – PCT – Retail pharmacy-Specialist64.80 | 5 | Hospira |
| Inj 1 mg per ml, 2 ml – PCT – Retail pharmacy-Specialist | 5 | Hospira |
| Inj 1 mg for ECP – PCT only – Specialist | 1 mg | Baxter |
| VINORELBINE – PCT only – Specialist | | |
| Inj 10 mg per ml, 1 ml vial8.00 | 1 | Navelbine |
| 42.00 | | Vinorelbine Ebewe |
| Inj 10 mg per ml, 5 ml vial40.00 | 1 | Navelbine |
| 210.00 | | Vinorelbine Ebewe |
| Inj 1 mg for ECP0.90 | 1 mg | Baxter |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|-----|---------------------|-------------------------------------|
| Protein-tyrosine Kinase Inhibitors | | | | |
| DASATINIB – Special Authority see SA0976 below – [Xpharm] | | | | |
| Tab 20 mg | 3,774.06 | 60 | 🗸 S | prycel |
| Tab 50 mg | 6,214.20 | 60 | 🗸 S | prycel |
| Tab 70 mg | 7,692.58 | 60 | 🗸 S | prycel |
| Tab 100 mg | 6 214 20 | 30 | V S | prvcel |

►SA0976 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from PHARMAC's website <u>http://www.pharmac.govt.nz</u>, and prescriptions should be sent to:

| The CML/GIST Co-ordinator | Phone: (04) 460 4990 |
|---------------------------|---|
| PHARMAC | Facsimile: (04) 916 7571 |
| PO Box 10 254 | Email: cmlgistcoordinator@pharmac.govt.nz |
| Wellington | |

Wellington

Special Authority criteria for CML - access by application

- a) Funded for patients with diagnosis (confirmed by a haematologist) of a chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase.
- b) Maximum dose of 140 mg/day for accelerated or blast phase, and 100 mg/day for chronic phase CML.
- c) Subsidised for use as monotherapy only.
- d) Initial approvals valid seven months.
- e) Subsequent approval(s) are granted on application and are valid for six months. The first reapplication (after seven months) should provide details of the haematological response. The third reapplication should provide details of the cytogenetic response after 14-18 months from initiating therapy. All other reapplications should provide details of haematological response, and cytogenetic response if such data is available. Applications to be made and subsequent prescriptions can be written by a haematologist or an oncologist.

Note: Dasatinib is indicated for the treatment of adults with chronic, accelerated or blast phase CML with resistance or intolerance to prior therapy including imatinib.

Guideline on discontinuation of treatment for patients with CML

- a) Prescribers should consider discontinuation of treatment if, after 6 months from initiating therapy, a patient did not obtain a haematological response as defined as any one of the following three levels of response:
 - a) complete haematologic response (as characterised by an absolute neutrophil count (ANC) > 1.5×10^{9} /L, platelets > 100 × 10⁹/L, absence of peripheral blood (PB) blasts, bone marrow (BM) blasts < 5% (or FISH Ph+ 0-35% metaphases), and absence of extramedullary disease); or
 - b) no evidence of leukaemia (as characterised by an absolute neutrophil count (ANC) > 1.0 × 10⁹/L, platelets > 20 × 10⁹/L, absence of peripheral blood (PB) blasts, bone marrow (BM) blasts < 5% (or FISH Ph+ 0-35% metaphases), and absence of extramedullary disease); or
 - c) return to chronic phase (as characterised by BM and PB blasts < 15%, BM and PB blasts and promyelocytes < 30%, PB basophils < 20% and absence of extramedullary disease other than spleen and liver).</p>
- Prescribers should consider discontinuation of treatment if, after 18 months from initiating therapy, a patient did not obtain a major cytogenetic response defined as 0-35% Ph+ metaphases.

ERLOTINIB - Retail pharmacy-Specialist - Special Authority see SA1519 on the next page

| Tab 100 mg | | ····· | 1 | ,000.00 | 30 | ✓ Tarceva |
|------------|------|-------|---|---------|----|-----------|
| Tab 150 mg | | | 1 | ,500.00 | 30 | ✓ Tarceva |

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

SA1519 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

- Either:
 - 1 All of the following:
 - 1.1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
 - 1.2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
 - 1.3 Any of the following:
 - 1.3.1 Patient is treatment naive; or
 - 1.3.2 Both:
 - 1.3.2.1 Patient has documented disease progression following treatment with first line platinum based chemotherapy; and
 - 1.3.2.2 Patient has not received prior treatment with gefitinib; or

1.3.3 Both:

- 1.3.3.1 The patient has discontinued gefitinib within 6 weeks of starting treatment due to intolerance; and
- 1.3.3.2 The cancer did not progress while on gefitinib; and
- 1.4 Erlotinib is to be given for a maximum of 3 months; or
- 2 The patient received funded erlotinib prior to 31 December 2013 and radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB - Retail pharmacy-Specialist

Tab 250 mg − Special Authority see SA1520 below......1,700.00 30 🖌 Iressa

➡SA1520 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
 - 2.1 Patient is treatment naive; or
 - 2.2 Both:

2.2.1 The patient has discontinued erlotinib within 6 weeks of starting treatment due to intolerance; and 2.2.2 The cancer did not progress whilst on erlotinib; and

- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

IMATINIB MESILATE

Note: Imatinib-AFT is not a registered for the treatment of Gastro Intestinal Stromal Tumours (GIST). The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule.

| | Tab Too mg - Special Automy see SA 1400 on the next page | | |
|---|--|----|----------------|
| | - [Xpharm]2,400.00 | 60 | Glivec |
| * | Cap 100 mg | 60 | Imatinib-AFT |
| * | Cap 400 mg | 30 | ✓ Imatinib-AFT |

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | 9 | Subsidised | Generic |
| \$ | Per | ~ | Manufacturer |

SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from PHARMAC's website <u>http://www.pharmac.govt.nz</u>, and prescriptions should be sent to:

| The CML/GIST Co-ordinator | Phone: (04) 460 4990 |
|---------------------------|---|
| PHARMAC | Facsimile: (04) 916 7571 |
| PO Box 10 254 | Email: cmlgistcoordinator@pharmac.govt.nz |
| | |

Wellington

Special Authority criteria for GIST - access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA1191 below - Retail pharmacy

Tab 250 mg 1,899.00 70 V Tykerb

SA1191 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Fither:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient has not previously received trastuzumab treatment for HER 2 positive metastatic breast cancer; and
 - 1.3 Lapatinib not to be given in combination with trastuzumab; and
 - 1.4 Lapatinib to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2.2 The patient started trastuzumab for metastatic breast cancer but discontinued trastuzumab within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on trastuzumab; and
 - 2.4 Lapatinib not to be given in combination with trastuzumab; and
 - 2.5 Lapatinib to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
 - 3 Lapatinib not to be given in combination with trastuzumab; and
 - 4 Lapatinib to be discontinued at disease progression.

NILOTINIB – Special Authority see SA1489 on the next page – Retail pharmacy

| Wastage claimable – see rule 5.5.2 on page 15 | | | |
|---|----------|-----|-----------------------------|
| Cap 150 mg | 4,680.00 | 120 | 🖌 Tasigna |
| Cap 200 mg | 6,532.00 | 120 | Tasigna |

| Subsidy (Manufacturer's Pric | 20) | Fully Subsidised | Brand or Generic | |
|---------------------------------|-----|---------------------|---------------------|--|
| (Waluadue S Fid | Per | | Manufacturer | |

➡SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Either:
 - 2.1 Patient has documented CML treatment failure* with imatinib; or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PAZOPANIB - Special Authority see SA1190 below - Retail pharmacy

| Tab 200 mg | 1,334.70 | 30 | Votrient |
|------------|----------|----|------------------------------|
| Tab 400 mg | 2,669.40 | 30 | Votrient |

SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
 - The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of \leq 70; or
 - 5.6 \geq 2 sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

| Subsidy | Fully | | Brand or |
|------------------------|----------|----|--------------|
| (Manufacturer's Price) | Subsidis | ed | Generic |
| \$ | Per | ~ | Manufacturer |

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

| SUNITINIB – Special Authority see SA1266 below – Retail pharmacy | | |
|--|------|----------------------------|
| Cap 12.5 mg2,315.38 | 3 28 | Sutent |
| Cap 25 mg4,630.77 | 7 28 | Sutent |
| Cap 50 mg9,261.54 | 4 28 | Sutent |

➡SA1266 Special Authority for Subsidy

Initial application - (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of \leq 70; or
 - 5.6 > 2 sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application - (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Fither:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

| Subsidy | | Fully | Brand or | |
|-----------------------|-----|------------|--------------|--|
| (Manufacturer's Price |) | Subsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of $\geq 10\%$ or decrease in tumour density in Hounsfield Units (HU) of $\geq 15\%$ on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of $\geq 10\%$ and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

►SA1515 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 5 months for applications meeting the following criteria:

All of the following:

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|---------------------|--|--|
| continued 1 Significant decrease in serum PSA from baseline; and 2 No evidence of clinical disease progression; and 3 No initiation of taxane chemotherapy with abiraterone; ar 4 The treatment remains appropriate and the patient is ber | | t. | | |
| BICALUTAMIDE Tab 50 mg | 4.90 | 28 | ✓ <u>B</u> | icalaccord |
| FLUTAMIDE – Retail pharmacy-Specialist | | | | |
| Tab 250 mg | | 30 100 | | lutamin S29 S29 lutamin |
| MEGESTROL ACETATE – Retail pharmacy-Specialist Tab 160 mg Apo-Megestrol to be Sole Supply on 1 November 2015 | 54.30 | 30 | 🗸 A | po-Megestrol |
| OCTREOTIDE Inj 50 mcg per ml, 1 ml vial Inj 100 mcg per ml, 1 ml vial Inj 500 mcg per ml, 1 ml vial | 22.40 | 5 5 5 | ✓ <u>D</u> ✓ <u>D</u> ✓ <u>D</u> | BL |
| OCTREOTIDE LAR (SOMATOSTATIN ANALOGUE) – Special Au Inj LAR 10 mg prefilled syringe Inj LAR 20 mg prefilled syringe Inj LAR 30 mg prefilled syringe | uthority see SA1016 b 1,772.50 2,358.75 | elow 1 1 1 | v∕s v∕s | armacy andostatin LAR andostatin LAR andostatin LAR |

➡SA1016 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are Unapproved Indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

| Subsidy | | Fully | Brand or | |
|------------------------|------|---------|--------------|--|
| (Manufacturer's Price) | Subs | sidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Any of the followina:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery: or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed: or
- 3 Both:
 - 3.1 Insulinomas: and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal - (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

| * Tab 10 mg * Tab 20 mg | 30 | ✓ Genox✓ Genox✓ Genox | |
|----------------------------|----|---|--|
| Aromatase Inhibitors | | | |

Aromatase innibitors

| ANASTROZOLE * Tab 1 mg26.55 | 30 | ✓ Aremed ✓ Arimidex ✓ DP-Anastrozole |
|--------------------------------|----|--|
| EXEMESTANE * Tab 25 mg14.50 | 30 | ✓ <u>Aromasin</u> |
| LETROZOLE * Tab 2.5 mg4.85 | 30 | ✓ Letraccord |

| | Subsidy (Manufacturer's Price \$ | e) Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|--|--|---------------------|--|
| Immunosuppressants | | | | |
| Cytotoxic Immunosuppressants | | | | |
| AZATHIOPRINE – Retail pharmacy-Specialist * Tab 25 mg * Tab 50 mg – For azathioprine oral liquid formulation refer, page 207 * Inj 50 mg MYCOPHENOLATE MOFETIL Tab 500 mg Cap 250 mg Powder for oral liq 1 g per 5 ml – Subsidy by endorsement Mycophenolate powder for oral liquid is subsidised only fo prescription is endorsed accordingly. | | 60 100 1 50 100 65 ml C swallo | | Azamun Azamun muran <u>Cellcept</u> <u>Cellcept</u> Cellcept nd capsules, and when the |
| Fusion Proteins | | | | |
| ETANERCEPT – Special Authority see SA1478 below – Retail ph Inj 25 mg Inj 50 mg autoinjector Inj 50 mg prefilled syringe | 949.96 1,899.92 | 4 4 4 | V E | Enbrel Enbrel Enbrel |

SA1478 Special Authority for Subsidy

Initial application — (juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

unon. 4 D

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient diagnosed with Juvenile Idiopathic Arthritis (JIA); and
 - 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
 - 2.5 Both:
 - 2.5.1 Either:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | ~ | Manufacturer |

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulphasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of cyclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

continued...

- 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
- 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, cyclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal antiinflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of an exercise regime supervised by a physiotherapist; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

continued...

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm

25-34 years - Male: 7.5 cm; Female: 5.5 cm

35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm

45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulphasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, cyclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 4 doses.

Note: Indications marked with * are Unapproved Indications (refer to Interpretations and Definitions).

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

1 Both:

- 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a named specialist or rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- - 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
 - 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
 - 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

| Subsidy | | Fully | Brand or |
|---------------------|------|------------|--------------|
| (Manufacturer's Pri | ice) | Subsidised | Generic |
| \$ | Per | ~ | Manufacturer |

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
 - 2 Following 12 weeks of etanercept treatment, BASDAI has improved by 4 or more points from pre-treatment baseline on a 10 point scale, or by 50%, whichever is less; and
 - 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
 - 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement: and
- 2 Patient continues to require treatment; and
- 3 A maximum of 4 doses.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Fither:

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Immune Modulators

| ANTITHYMOCYTE GLOBULIN (EQUINE) – PCT only – Specialist Inj 50 mg per ml, 5 ml2,351.25 | 5 | ✔ ATGAM |
|---|---|------------------|
| BACILLUS CALMETTE-GUERIN (BCG) VACCINE – PCT only – Specialist Subsidised only for bladder cancer. | | |
| Inj 2-8 \times 100 million CFU | 1 | V OncoTICE |
| • | | |
| Inj 40 mg per ml, vial149.37 | 3 | SII-Onco-BCG S29 |
| Monoclonal Antibodies | | |
| ADALIMUMAB – Special Authority see SA1479 below – Retail pharmacy | | |
| Inj 20 mg per 0.4 ml prefilled syringe1,799.92 | 2 | Humira |
| Inj 40 mg per 0.8 ml prefilled pen | 2 | HumiraPen |
| Inj 40 mg per 0.8 ml prefilled syringe1,799.92 | 2 | Humira |

SA1479 Special Authority for Subsidy

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis: or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | ~ | Manufacturer |

continued...

- 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulphasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of cyclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (Crohn's disease) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

continued...

- 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
- 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, cyclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. **Initial application — (ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal antiinflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of an exercise regime supervised by a physiotherapist; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

| | Subsidy | Fully | Brand or |
|--------|-------------------|------------|--------------|
| (Manuf | facturer's Price) | Subsidised | Generic |
| | \$ Per | r 🖌 | Manufacturer |

continued...

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application. Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm: Female: 5.5 cm

25-34 years - Male: 7.5 cm; Female: 5.5 cm

35-44 years - Male: 6.5 cm; Female: 4.5 cm

45-54 years - Male: 6.0 cm; Female: 5.0 cm

55-64 years - Male: 5.5 cm; Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulphasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for juvenile idiopathic arthritis; or
- 2 All of the following:

| Subsidy (Manufacturer's Price) | | Fully Subsidised | Brand or Generic | |
|-----------------------------------|-----|---------------------|---------------------|--|
| \$ | Per | ~ | Manufacturer | |

continued...

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient diagnosed with JIA; and
- 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
- 2.5 Both:
 - 2.5.1 Either:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's disease. **Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 4 doses.

Note: Note: Indications marked with * are Unapproved Indications (refer to (Interpretations and Definitions).

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or

2 All of the following:

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

- 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
- 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
- 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Either:

- 1.1 Applicant is a gastroenterologist; or
- 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 CDAI score is 150 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and

| Subsidy | Fu | lly Brand or | |
|------------------------|-----------|----------------------------------|--|
| (Manufacturer's Price) | Subsidise | ed Generic | |
| \$ | Per | Manufacturer | |

continued...

- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist: or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
 - 2 Following 12 weeks of adalimumab treatment, BASDAI has improved by 4 or more points from pre-adalimumab baseline on a 10 point scale, or by 50%, whichever is less; and
 - 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
 - 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Either:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

continued...

- 1.1 Applicant is a named specialist or rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Fither:

- 1.1 Applicant is a gastroenterologist; or
- 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 4 doses.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

oun:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

OMALIZUMAB - Special Authority see SA1490 below - Retail pharmacy

| Inj 150 mg vial | | 1 | 🖌 Xolair |
|-----------------|--|---|----------|
|-----------------|--|---|----------|

SA1490 Special Authority for Subsidy

Initial application only from a respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient is over the age of 6; and
- 2 Patient has a diagnosis of severe, life threatening asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and

| 5 | Fully Subsidised | Brand or Generic | |
|-----------|---------------------|---------------------|--|
| \$ Per | ~ | Manufacturer | |

continued...

- 5 Proven compliance with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1600 micrograms per day or fluticasone propionate 1000 micrograms per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 micrograms bd or eformoterol 12 micrograms bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; and
- 7 At least four admissions to hospital for a severe asthma exacerbation over the previous 24 months with at least one of those being in the previous 12 months; and
- 8 An Asthma Control Questionnaire (ACQ-5) score of at least 3.0 as assessed in the previous month .

Renewal only from a respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Hospital admissions have been reduced as a result of treatment; and
- 2 A reduction in the Asthma Control Questionnaire (ACQ-5) score of at least 1.0 from baseline; and
- 3 A reduction in the maintenance oral corticosteroid dose of at least 50% from baseline.

RITUXIMAB - PCT only - Specialist - Special Authority see SA1152 below

| Inj 100 mg per 10 ml vial | 2 | Mabthera |
|-----------------------------------|------|----------------------------|
| Inj 500 mg per 50 ml vial2,688.30 | 1 | Mabthera |
| Inj 1 mg for ECP5.64 | 1 mg | Baxter |

SA1152 Special Authority for Subsidy

Initial application — (Post-transplant) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are Unapproved Indications.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Either:

1 Both:

- 1.1 The patient has indolent low grade NHL with relapsed disease following prior chemotherapy; and
- 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia.

Initial application — (Aggressive CD20 positive NHL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or

2 Both:

2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and

| Subsidy | | Fully | Brand or | |
|------------------------|------|---------|--------------|--|
| (Manufacturer's Price) | Subs | sidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (Chronic Lymphocytic Leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 The patient is rituximab treatment naive; and
- 3 Either:
 - 3.1 The patient is chemotherapy treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
 - 3.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; and
- 4 The patient has good performance status; and
- 5 The patient has good renal function (creatinine clearance \geq 30 ml/min); and
- 6 The patient does not have chromosome 17p deletion CLL; and
- 7 Rituximab to be administered in combination with fludarabine and cyclophosphamide for a maximum of 6 treatment cycles; and
- 8 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration).

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to <2.

Renewal — (Post-transplant) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are Unapproved Indications.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia.

Renewal — (Aggressive CD20 positive NHL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

| | Subsidy (Manufacturer's Price) \$ | | | Brand or Generic Manufacturer | |
|--|---|----|------|-------------------------------------|--|
| TRASTUZUMAB – PCT only – Specialist – Specia | al Authority see SA1521 below | | | | |
| Inj 150 mg vial | | 1 | 🖌 He | erceptin | |
| Inj 440 mg vial | | 1 | 🖌 He | erceptin | |
| Ini 1 ma for ECP | | ma | 🖌 Ba | axter | |

SA1521 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient has not previously received lapatinib treatment for HER 2 positive metastatic breast cancer; and
 - 1.3 Trastuzumab not to be given in combination with lapatinib; and
 - 1.4 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2.2 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on lapatinib; and
 - 2.4 Trastuzumab not to be given in combination with lapatinib; and
 - 2.5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | ~ | Manufacturer |

continued...

- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 All of the following:
 - 3.1.1 The patient has not previously received lapatinib treatment for metastatic breast cancer; and
 - 3.1.2 Trastuzumab not to be given in combination with lapatinib; and
 - 3.1.3 Trastuzumab to be discontinued at disease progression; or
 - 3.2 All of the following:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; and
 - 3.2.3 Trastuzumab not to be given in combination with lapatinib; and
 - 3.2.4 Trastuzumab to be discontinued at disease progression; or
 - 3.3 All of the following:
 - 3.3.1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
 - 3.3.2 Trastuzumab not to be given in combination with lapatinib; and
 - 3.3.3 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

Other Immunosuppressants

CICLOSPORIN

| Cap 25 mg | | 50 | Neoral |
|---|----------------|----------|------------------------------|
| Cap 50 mg | | 50 | Neoral |
| Cap 100 mg | | 50 | Neoral |
| Oral liq 100 mg per ml | 198.13 | 50 ml OP | Neoral |
| EVEROLIMUS – Special Authority see SA1491 below – Re Wastage claimable – see rule 3.3.2 on page 13 | etail pharmacy | | |
| Tab 5 mg | 4,555.76 | 30 | Afinitor |
| Tab 10 mg | 6,512.29 | 30 | Afinitor |

►SA1491 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

SIROLIMUS - Special Authority see SA0866 on the next page - Retail pharmacy

| Tab 1 mg | 100 | Rapamune |
|----------------------|--------------|----------|
| 5 | 100 | Rapamune |
| Oral liq 1 mg per ml | 60 ml OP | Rapamune |

| Subsidy (Manufacturer's Price) Si | | lly | Brand or Generic |
|--------------------------------------|--------------------|-----|---------------------|
| (Manulactaler 3 1 nec) \$ | Subsidise Per o | ~ | Manufacturer |

SA0866 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR<30 ml/min; or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- HUS or TTP; or
- Leukoencepthalopathy; or
- Significant malignant disease

TACROLIMUS – Special Authority see SA1540 below – Retail pharmacy

| Cap 0.5 mg | | 100 | Tacrolimus Sandoz |
|---|--------|-----|---------------------|
| Cap 1 mg | 171.20 | 100 | ✓ Tacrolimus Sandoz |
| Cap 5 mg - For tacrolimus oral liquid formulation refer, page | | | |
| 207 | 428.00 | 50 | ✓ Tacrolimus Sandoz |

➡SA1540 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Initial application — (steroid-resistant nephrotic syndrome*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Fither:

- zither:
 - 1 The patient is a child with steroid-resistant nephrotic syndrome* (SRNS) where ciclosporin has been trialled in combination with prednisone and discontinued because of unacceptable side effects or inadequate clinical response; or
 - 2 All of the following:
 - 2.1 The patient is an adult with SRNS; and
 - 2.2 Ciclosporin has been trialled in combination with prednisone and discontinued because of unacceptable side effects or inadequate clinical response; and
 - 2.3 Cyclophosphamide or mycophenolate have been trialled and discontinued because of unacceptable side effects or inadequate clinical response, or these treatments are contraindicated.

Note: Indications marked with * are Unapproved Indications

Note: Subsidy applies for either primary or rescue therapy.

| | Subsidy (Manufacturer's P | rice) Sul | Full | |
|---|------------------------------|----------------|---------|-------------------------------|
| | \$ | Per | V | Manufacturer |
| Antiallergy Preparations | | | | |
| ►SA1367 Special Authority for Subsidy | | | | |
| Initial application only from a relevant specialist. Approvals valid Both: | for 2 years for a | oplications me | eting | the following criteria: |
| RAST or skin test positive; and Patient has had severe generalised reaction to the sensiti | sing agent. | | | |
| Renewal only from a relevant specialist. Approvals valid for 2 ye benefiting from treatment. | ears where the tr | eatment rema | ains ap | ppropriate and the patient is |
| BEE VENOM ALLERGY TREATMENT - Special Authority see SA | A1367 above – F | letail pharmad | су | |
| Maintenance kit - 6 vials 120 mcg freeze dried venom, 6 dilu- | | | | |
| ent 1.8 ml | | 1 OP | V | Albay |
| Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent 9 ml, 3 diluent 1.8 ml | | 1 OP | | Albey |
| (Albay Maintenance kit - 6 vials 120 mcg freeze dried venom, 6 di | | | | |
| WASP VENOM ALLERGY TREATMENT – Special Authority see | | | | , =0.0) |
| Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze | | | acy | |
| dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml | | 1 OP | V | Albey |
| Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze | | | | |
| dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml | | 1 OP | ~ | Albey |
| Antihistamines | | | | |
| CETIBIZINE HYDROCHLORIDE | | | | |
| * Tab 10 mg | 1.59 | 100 | ~ | Zetop |
| *‡ Oral liq 1 mg per ml | | 200 ml | | <u>Histaclear</u> |
| CHLORPHENIRAMINE MALEATE | | | | |
| *‡ Oral liq 2 mg per 5 ml | 8.06 | 500 ml | ~ | Histafen |
| DEXTROCHLORPHENIRAMINE MALEATE | | | | |
| * Tab 2 mg | 1.01 | 20 | | |
| | (5.99) | | | Polaramine |
| | 2.02 | 40 | | Delevenia |
| *‡ Oral liq 2 mg per 5 ml | (8.40) | 100 ml | | Polaramine |
| | | 100 111 | | Polaramine |
| FEXOFENADINE HYDROCHLORIDE | (10.20) | | | |
| * Tab 60 mg | 4.34 | 20 | | |
| | (11.53) | | | Telfast |
| * Tab 120 mg | | 10 | | |
| | (11.53) | •- | | Telfast |
| | 14.22 | 30 | | Talfaat |
| | (29.81) | | | Telfast |
| LORATADINE | 1 00 | 100 | | Lorofix |
| * Tab 10 mg * Oral lig 1 mg per ml | | 100 200 ml | | Lorafix LoraPaed |
| | | 200 111 | • | |

| | Subsidy | | Fully Brand or |
|--|-----------------------|---------------------------|--|
| | (Manufacturer's \$ | Price) Sur Per | osidised Generic Manufacturer |
| ROMETHAZINE HYDROCHLORIDE | | | |
| Fab 10 mg | 1.78 | 50 | Allersoothe |
| Allersoothe to be Sole Supply on 1 October 2015 | | | |
| Tab 25 mg | 1.99 | 50 | Allersoothe |
| Allersoothe to be Sole Supply on 1 October 2015 | | | 4 4 1 1 1 |
| t Oral liq 1 mg per 1 ml | 2.59 | 100 ml | Allersoothe |
| Allersoothe to be Sole Supply on 1 October 2015 Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a | | | |
| PSO | 11.99 | 5 | ✓ Hospira |
| | | Ũ | • noopnu |
| Oral lig 30 mg per 5 ml | 2 70 | 100 ml OP | |
| | (8.06) | | Vallergan Forte |
| | (0.00) | | valiergan i one |
| nhaled Corticosteroids | | | |
| ECLOMETHASONE DIPROPIONATE | | | |
| Aerosol inhaler, 50 mcg per dose | 9.30 | 200 dose OP | ✓ Qvar |
| Aerosol inhaler, 50 mcg per dose CFC-free | 8.54 | 200 dose OP | Beclazone 50 |
| Aerosol inhaler, 100 mcg per dose | | 200 dose OP | |
| Aerosol inhaler, 100 mcg per dose CFC-free | | 200 dose OP | |
| Aerosol inhaler, 250 mcg per dose CFC-free | 22.67 | 200 dose OP | Beclazone 250 |
| UDESONIDE | | | |
| Powder for inhalation, 100 mcg per dose | 17.00 | 200 dose OP | |
| | 10.00 | | Turbuhaler |
| Powder for inhalation, 200 mcg per dose | | 200 dose OP | ✓ Pulmicort |
| Develop (or inheliation, 400 menors do a | 00.00 | 000 days OD | Turbuhaler |
| Powder for inhalation, 400 mcg per dose | | 200 dose OP | Pulmicort Turbuhaler |
| | | | Turbunaler |
| LUTICASONE | 7 50 | 100 daga OD | |
| Aerosol inhaler, 50 mcg per dose CFC-free Powder for inhalation, 50 mcg per dose | | 120 dose OP 60 dose OP | Flixotide Flixotide Accuhaler |
| Powder for inhalation, 30 mcg per dose | | 60 dose OP | Flixotide Accuhaler Flixotide Accuhaler |
| Aerosol inhaler, 125 mcg per dose CFC-free | | 120 dose OP | |
| Aerosol inhaler, 250 mcg per dose CFC-free | | 120 dose OP | |
| Powder for inhalation, 250 mcg per dose | | 60 dose OP | Flixotide Accuhaler |
| nhaled Long-acting Beta-adrenoceptor Agonists | | | |
| Inialed Long-acting Deta-adrenoceptor Agonists | • | | |
| FORMOTEROL FUMARATE | | | |
| Powder for inhalation, 6 mcg per dose, breath activated | | 60 dose OP | |
| | (16.90) | | Oxis Turbuhaler |
| Powder for inhalation, 12 mcg per dose, and monodose de- | 00.04 | 00 I | |
| vice | (0 - 0 0) | 60 dose | Forodil |
| | (35.80) | | Foradil |
| IDACATEROL | 04.00 | | |
| Powder for inhalation 150 mcg | | 30 dose OP | Onbrez Breezhaler Onbrez Breezhaler |
| Powder for inhalation 300 mcg | | 30 dose OP | Onbrez Breezhaler |
| ALMETEROL | | 100 J 7 - | 4. |
| Aerosol inhaler CFC-free, 25 mcg per dose | | 120 dose OP | ✓ Serevent |
| Powder for inhalation, 50 mcg per dose, breath activated | | 60 dose OP | Serevent Accuhaler |

Turbuhaler 400/12

| | Subsidy (Manufacturer's F \$ | Price) Subs Per | Fully sidised | Brand or Generic Manufacturer |
|--|------------------------------------|-------------------------------|------------------|-------------------------------------|
| Inhaled Corticosteroids with Long-Acting Beta-A | drenocepto | or Agonists | | |
| BUDESONIDE WITH EFORMOTEROL – Special Authority see SA Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg Powder for inhalation 100 mcg with eformoterol fumarate | | Retail pharmac 120 dose OP | | annair |
| 6 mcg | 55.00 | 120 dose OP | | ymbicort Turbuhaler 100/6 |
| Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg Powder for inhalation 200 mcg with eformoterol fumarate | 31.25 | 120 dose OP | 🖌 Va | annair |
| 6 mcg | 60.00 | 120 dose OP | ✔ S | ymbicort Turbuhaler 200/6 |
| Powder for inhalation 400 mcg with eformoterol fumarate 12 mcg – No more than 2 dose per day | 60.00 | 60 dose OP | ✔ S | ymbicort |

SA1179 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Patient is a child under the age of 12; and
 - 1.2 Has been treated with inhaled corticosteroids of at least 400 mcg per day beclomethasone or budesonide, or 200 mcg per day fluticasone; and
 - 1.3 The prescriber considers that the patient would receive additional clinical benefit from switching to a combination product; or
- 2 All of the following:
 - 2.1 Patient is over the age of 12; and
 - 2.2 Has been treated with inhaled corticosteroids of at least 800 mcg per day beclomethasone or budesonide, or 500 mcg per day fluticasone; and
 - 2.3 The prescriber considers that the patient would receive additional clinical benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

FLUTICASONE WITH SALMETEROL

| | Aerosol inhaler 50 mcg with salmeterol 25 mcg | 120 dose OP | Seretide |
|----|---|-------------|------------------------------|
| | Aerosol inhaler 125 mcg with salmeterol 25 mcg49.69 | 120 dose OP | Seretide |
| | Powder for inhalation 100 mcg with salmeterol 50 mcg – No more than 2 dose per day | 60 dose OP | Seretide Accuhaler |
| | | 00 dose OF | Seletide Accultatel |
| | Powder for inhalation 250 mcg with salmeterol 50 mcg – No more than 2 dose per day | 60 dose OP | ✓ Seretide Accuhaler |
| В | eta-Adrenoceptor Agonists | | |
| SA | LBUTAMOL | | |
| ţ | Oral lig 400 mcg per ml2.06 | 150 ml | ✓ Ventolin |
| | Infusion 1 mg per ml, 5 ml | 10 | |
| | (130.21) | | Ventolin |
| | Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO | 5 | ✓ Ventolin |
| | | | |

| | Subsidy (Manufacturer's \$ | | Fully Brand or sidised Generic ✔ Manufacturer |
|--|----------------------------------|-------------|---|
| Inhaled Beta-Adrenoceptor Agonists | | | |
| SALBUTAMOL | | | |
| Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000 dose available on a PSO | | 200 dose OP | ✓ Respigen✓ Salamol |
| | (6.00) | | Ventolin |
| Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO | 3.19 | 20 | ✓ Asthalin |
| Asthalin to be Sole Supply on 1 October 2015 Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO Asthalin to be Sole Supply on 1 October 2015 | | 20 | ✓ Asthalin |
| TERBUTALINE SULPHATE Powder for inhalation, 250 mcg per dose, breath activated | 22.00 | 200 dose OP | Bricanyl Turbuhaler |
| Anticholinergic Agents | | | |
| PRATROPIUM BROMIDE | | | |
| Aerosol inhaler, 20 mcg per dose CFC-free Nebuliser soln, 250 mcg per ml, 1 ml – Up to 40 neb available | | 200 dose OP | ✓ Atrovent |
| on a PSO | | 20 | Univent |
| Nebuliser soln, 250 mcg per ml, 2 ml – Up to 40 neb available on a PSO | | 20 | ✓ Univent |
| Inhaled Beta-Adrenoceptor Agonists with Antich | nolinergic A | gents | |
| SALBUTAMOL WITH IPRATROPIUM BROMIDE | | | |
| Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg per dose CFC-free | | 200 dose OP | 🗸 Duolin HFA |
| Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule – Up to 20 neb available on a PSO Duolin to be Sole Supply on 1 October 2015 | | 20 | 🗸 Duolin |
| Long-Acting Muscarinic Antagonists | | | |

Long-Acting Muscarinic Antagonists

SA1485 Special Authority for Subsidy

Initial application only from a general practitioner or relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 To be used for the long-term maintenance treatment of bronchospasm and dyspnoea associated with COPD; and
- 2 In addition to standard treatment, the patient has trialled a short acting bronchodilator dose of at least 40 µg ipratropium g.i.d for one month; and
- 3 Either:
 - The patient's breathlessness according to the Medical Research Council (UK) dyspnoea scale is:
 - 3.1 Grade 4 (stops for breath after walking about 100 meters or after a few minutes on the level); or
 - 3.2 Grade 5 (too breathless to leave the house, or breathless when dressing or undressing); and
- 4 All of the following:

Applicant must state recent measurement of:

4.1 Actual FEV1 (litres); and

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

continued...

- 4.2 Predicted FEV1 (litres); and
- 4.3 Actual FEV₁ as a % of predicted (must be below 60%); and
- 5 Either:
 - 5.1 Patient is not a smoker (for reporting purposes only); or
 - 5.2 Patient is a smoker and has been offered smoking cessation counselling; and
- 6 The patient has been offered annual influenza immunisation.

Renewal only from a general practitioner or relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined); and
- 3 All of the following:

Applicant must state recent measurement of:

- 3.1 Actual FEV1 (litres); and
- 3.2 Predicted FEV₁ (litres); and
- 3.3 Actual FEV $_1$ as a % of predicted.

GLYCOPYRRONIUM – Special Authority see SA1485 on the previous page – Retail pharmacy

| Glycopyrronium treatment will not be subsidised if patient is also receiving tre | atment with sub | sidised tiotropium. |
|--|-----------------|---------------------------------------|
| Powder for inhalation 50 mcg per dose61.00 | 30 dose OP | Seebri Breezhaler |

TIOTROPIUM BROMIDE - Special Authority see SA1485 on the previous page - Retail pharmacy

Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised glycopyrronium.

Leukotriene Receptor Antagonists

MONTELUKAST - Special Authority see SA1421 below - Retail pharmacy

Prescribing Guideline: Clinical evidence indicates that the effectiveness of montelukast is strongest when montelukast is used in short treatment courses.

| Tab 4 mg | 28 | Singulair |
|-----------|----|-----------|
| Tab 5 mg | 28 | Singulair |
| Tab 10 mg | 28 | Singulair |

SA1421 Special Authority for Subsidy

Initial application — (Pre-school wheeze) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 To be used for the treatment of intermittent severe wheezing (possibly viral) in children under 5 years; and
- 2 The patient has had at least three episodes in the previous 12 months of acute wheeze severe enough to seek medical attention.

Renewal — (Pre-school wheeze) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (exercise-induced asthma) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been trialled with maximal asthma therapy, including inhaled corticosteroids and long-acting beta-adrenoceptor agonists; and
- 2 Patient continues to receive optimal inhaled corticosteroid therapy; and

| Subsidy | (a.a.) | Fully | Brand or | |
|--------------------------|--------|------------|-------------------------|--|
| (Manufacturer's Pr \$ | Per | Subsidised | Generic Manufacturer | |

continued...

3 Patient continues to experience frequent episodes of exercise-induced bronchoconstriction.

Initial application — (aspirin desensitisation) only from a clinical immunologist or allergist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient is undergoing aspirin desensitisation therapy under the supervision of a Clinical Immunologist or Allergist; and
- 2 Patient has moderate to severe aspirin-exacerbated respiratory disease or Samter's triad; and
- 3 Nasal polyposis, confirmed radiologically or surgically; and
- 4 Documented aspirin or NSAID allergy confirmed by aspirin challenge or a clinical history of severe reaction to aspirin or NSAID where challenge would be considered dangerous.

| NOAD WHELE CHAILENGE WOULD be CONSIDERE | u uangerous. | | |
|---|--|-------------------|------------------------------------|
| Mast Cell Stabilisers | | | |
| NEDOCROMIL | | | |
| Aerosol inhaler, 2 mg per dose CFC-free | | 112 dose OP | ✓ Tilade |
| SODIUM CROMOGLYCATE | | | |
| Powder for inhalation, 20 mg per dose | 17.94 | 50 dose | Intal Spincaps |
| Aerosol inhaler, 5 mg per dose CFC-free | | 112 dose OP | Intal Forte CFC Free |
| Methylxanthines | | | |
| AMINOPHYLLINE | | | |
| * Inj 25 mg per ml, 10 ml ampoule - Up to 5 inj a | vailable on a | | |
| PSO | | 5 | DBL Aminophylline |
| THEOPHYLLINE | | | |
| * Tab long-acting 250 mg | | 100 | ✓ Nuelin-SR |
| *‡ Oral liq 80 mg per 15 ml | 15.50 | 500 ml | Nuelin |
| Mucolytics | | | |
| DORNASE ALFA - Special Authority see SA0611 be | | | |
| Nebuliser soln, 2.5 mg per 2.5 ml ampoule | 250.00 | 6 | Pulmozyme |
| SA0611 Special Authority for Subsidy | | | |
| Special Authority approved by the Cystic Fibrosis Adv | , | | |
| Notes: Application details may be obtained from PHA | <u> </u> | w.pharmac.govt.i | nz or: |
| The Co-ordinator, Cystic Fibrosis Advisory Panel | | | |
| PHARMAC, PO Box 10 254 Wellington | Facsimile: (04) 916 7571 Email: CFPanel@pharm | ac govt nz | |
| Prescriptions for patients approved for treatment mus | | | ediatricians who have experience |
| and expertise in treating cystic fibrosis. | | physicians of par | |
| SODIUM CHLORIDE | | | |
| Not funded for use as a nasal drop. | | | |
| Soln 7% | | 90 ml OP | ✓ Biomed |
| | | | |

| | Subsidy (Manufacturer's | | Fully Brand or sidised Generic |
|--|----------------------------|-------------|--|
| | \$ | Per | Manufacturer |
| Nasal Preparations | | | |
| Allergy Prophylactics | | | |
| ECLOMETHASONE DIPROPIONATE | | | |
| Metered aqueous nasal spray, 50 mcg per dose | 2.35 | 200 dose OP | |
| | (4.85) | | Alanase |
| Metered aqueous nasal spray, 100 mcg per dose | 2.46 (5.75) | 200 dose OP | Alanase |
| UDESONIDE | (0.1.0) | | |
| Metered aqueous nasal spray, 50 mcg per dose | 2 35 | 200 dose OP | |
| Melered aqueous hasal spray, so meg per dose | (4.85) | 200 0036 01 | Butacort Aqueous |
| Metered aqueous nasal spray, 100 mcg per dose | | 200 dose OP | Bulacon Aqueous |
| | (5.75) | 200 0000 01 | Butacort Aqueous |
| LUTICASONE PROPIONATE | (0110) | | Datatori i i quotato |
| Metered aqueous nasal spray, 50 mcg per dose | 2.18 | 120 dose OP | Flixonase Hayfever & Allergy |
| Flixonase Hayfever & Allergy to be Sole Supply on 1 Octo | ber 2015 | | a Anorgy |
| PRATROPIUM BROMIDE | 5012010 | | |
| Aqueous nasal spray, 0.03% | 3.05 | 15 ml OP | ✓ Univent |
| | | 13 111 01 | |
| Respiratory Devices | | | |
| IASK FOR SPACER DEVICE | | | |
| a) Up to 20 dev available on a PSO | | | |
| b) Only on a PSO | | | |
| c) Only for children aged six years and under | | | |
| Size 2 | 2 99 | 1 | EZ-fit Paediatric |
| | | · | Mask |
| EAK FLOW METER | | | |
| a) Up to 10 dev available on a PSO | | | |
| | | | |
| b) Only on a PSO | | | |
| | 11.44 | 1 | ✓ Breath-Alert |
| b) Only on a PSO | | 1 1 | Breath-Alert Breath-Alert |
| b) Only on a PSO Low range Normal range | | | |
| b) Only on a PSO Low range Normal range | | | |
| b) Only on a PSO Low range Normal range PACER DEVICE | | | |
| b) Only on a PSO Low range Normal range PACER DEVICE a) Up to 20 dev available on a PSO b) Only on a PSO | 11.44 | | |
| b) Only on a PSO Low range Normal range PACER DEVICE a) Up to 20 dev available on a PSO | 11.44 | 1 | ✓ Breath-Alert |
| b) Only on a PSO Low range Normal range PACER DEVICE a) Up to 20 dev available on a PSO b) Only on a PSO | 11.44 | 1 | Breath-Alert Space Chamber |
| b) Only on a PSO Low range Normal range PACER DEVICE a) Up to 20 dev available on a PSO b) Only on a PSO 230 ml (single patient) 800 ml | 11.44 | 1 | Breath-Alert Space Chamber Plus |
| b) Only on a PSO Low range Normal range PACER DEVICE a) Up to 20 dev available on a PSO b) Only on a PSO 230 ml (single patient) 800 ml PACER DEVICE AUTOCLAVABLE | 11.44 | 1 | Breath-Alert Space Chamber Plus |
| b) Only on a PSO Low range Normal range PACER DEVICE a) Up to 20 dev available on a PSO b) Only on a PSO 230 ml (single patient) 800 ml PACER DEVICE AUTOCLAVABLE a) Up to 5 dev available on a PSO | 11.44 | 1 | Breath-Alert Space Chamber Plus |
| b) Only on a PSO Low range Normal range PACER DEVICE a) Up to 20 dev available on a PSO b) Only on a PSO 230 ml (single patient) 800 ml PACER DEVICE AUTOCLAVABLE | 11.44 4.72 8.50 | 1 | Breath-Alert Space Chamber Plus Volumatic |
| b) Only on a PSO Low range Normal range PACER DEVICE a) Up to 20 dev available on a PSO b) Only on a PSO 230 ml (single patient) 800 ml PACER DEVICE AUTOCLAVABLE a) Up to 5 dev available on a PSO b) Only on a PSO | | 1 1 1 | Breath-Alert Space Chamber Plus Volumatic Space Chamber |
| b) Only on a PSO Low range | | 1 1 1 | Breath-Alert Space Chamber Plus Volumatic Space Chamber |
| b) Only on a PSO Low range | | 1 1 1 | Breath-Alert Space Chamber Plus Volumatic Space Chamber |
| b) Only on a PSO Low range | | 1 1 1 | Breath-Alert Space Chamber Plus Volumatic Space Chamber |

199

| | Subsidy (Manufacturer's | Price) Sub | Fully Brand or osidised Generic |
|--|----------------------------|--------------------|--|
| | (Manulacturers | Price) Suc Per | Manufacturer |
| ar Preparations | | | |
| ETIC ACID WITH 1, 2- PROPANEDIOL DIACETATE AND BEN | ZETHONIUM | | |
| For Vosol ear drops with hydrocortisone powder refer Standar | | ge 210 | |
| Ear drops 2% with 1, 2-Propanediol diacetate 3% and | | • | |
| benzethonium chloride 0.02% | 6.97 | 35 ml OP | Vosol |
| | | | . |
| Ear drops 0.02% with clioquinol 1% | 4.46 | 7.5 ml OP | Locacorten-Viaform ED's |
| | | | Locorten-Vioform |
| RIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCIN | NAND NYSTAT | ΊN | |
| Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate | E 16 | 7.5 ml OP | ✓ Kenacomb |
| 2.5 mg and gramicidin 250 mcg per g | | 7.5 III OP | |
| ar/Eye Preparations | | | |
| EXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN | | | |
| Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and | | | |
| gramicidin 50 mcg per ml | | 8 ml OP | |
| | (9.27) | | Sofradex |
| | 4 10 | | |
| Ear/Eye drops 0.5% | 4.13 (8.65) | 8 ml OP | Soframycin |
| vo Proporationa | (0.00) | | Containgont |
| Eye Preparations | | | |
| ve preparations are only funded for use in the eye, unless explici | tly stated other | wise. | |
| Inti-Infective Preparations | | | |
| CICLOVIR | | | |
| Eye oint 3% | | 4.5 g OP | Zovirax |
| HLORAMPHENICOL | | | |
| Eye oint 1% | | 4 g OP 10 ml OP | Chlorsig Chlorafast |
| Eye drops 0.5% a) Funded for use in the ear*. Indications marked with * are | | | Chioratast |
| b) Chlorafast to be Sole Supply on 1 October 2015 | | 10100101 | |
| PROFLOXACIN | | | |
| Eye Drops 0.3% | | 5 ml OP | Ciloxan |
| For treatment of bacterial keratitis or severe bacterial conju | inctivitis resista | nt to chloramph | enicol. |
| | 4.50 | | |
| Eye drops 1% | 4.50 | 5 g OP | Fucithalmic |
| NCICLOVIR | | | |
| Eye gel 0.15% | | 5 g OP | Virgan S29 |
| | 44.46 | 5 ml 00 | |
| Eye drops 0.3% | 11.40 | 5 ml OP | Genoptic |
| ROPAMIDINE ISETHIONATE | | | |
| | | 10 | |
| Eye drops 0.1% | | 10 ml OP | Brolene |

| | Subsidy (Manufacturer's F | Price) Sub | Fully Brand or osidised Generic |
|--|------------------------------|------------|-------------------------------------|
| | \$ | Per | Manufacturer |
| OBRAMYCIN | | | |
| Eye oint 0.3% | 10.45 | 3.5 g OP | ✓ <u>Tobrex</u> |
| Eye drops 0.3% | 11.48 | 5 ml OP | ✓ <u>Tobrex</u> |
| Corticosteroids and Other Anti-Inflammatory P | reparations | | |
| EXAMETHASONE | | | |
| € Eye oint 0.1% | | 3.5 g OP | Maxidex |
| € Eye drops 0.1% | 4.50 | 5 ml OP | Maxidex |
| EXAMETHASONE WITH NEOMYCIN SULPHATE AND POLY | | ATE | |
| Eye oint 0.1% with neomycin sulphate 0.35% and polymyx | | | A |
| b sulphate 6,000 u per g | | 3.5 g OP | Maxitrol |
| Eye drops 0.1% with neomycin sulphate 0.35% and polym | | | / •• •• • |
| xin b sulphate 6,000 u per ml | 4.50 | 5 ml OP | Maxitrol |
| ICLOFENAC SODIUM | | | |
| Eye drops 0.1% | 13.80 | 5 ml OP | Voltaren Ophtha |
| LUOROMETHOLONE | | | |
| € Eye drops 0.1% | 3.09 | 5 ml OP | ✓ FML |
| | 3.80 | | Flucon |
| EVOCABASTINE | | | |
| Eye drops 0.5 mg per ml | | 4 ml OP | |
| | (10.34) | | Livostin |
| ODOXAMIDE | | | |
| Eye drops 0.1% | 8.71 | 10 ml OP | Lomide |
| REDNISOLONE ACETATE | | | |
| € Eye drops 0.12% | 4.50 | 5 ml OP | Pred Mild |
| Eye drops 1% | 4.50 | 5 ml OP | Pred Forte |
| ODIUM CROMOGLYCATE | | | |
| Eye drops 2% | 1.18 | 5 ml OP | Rexacrom |
| Glaucoma Preparations - Beta Blockers | | | |
| ETAXOLOL | | | |
| € Eye drops 0.25% | | 5 ml OP | Betoptic S |
| € Eye drops 0.5% | 7.50 | 5 ml OP | <u>Betoptic</u> |
| EVOBUNOLOL | | | 4 |
| Eye drops 0.5% | 7.00 | 5 ml OP | Betagan |
| IMOLOL | | | |
| € Eye drops 0.25% | | 5 ml OP | Arrow-Timolol |
| Eye drops 0.25%, gel forming | | 2.5 ml OP | Timoptol XE |
| Eye drops 0.5% | | 5 ml OP | Arrow-Timolol |
| Eye drops 0.5%, gel forming | | 2.5 ml OP | Timoptol XE |
| Glaucoma Preparations - Carbonic Anhydrase | Inhibitors | | |
| CETAZOLAMIDE | | | |
| | er, | | |
| Tab 250 mg – For acetazolamide oral liquid formulation refe | | | (D) |
| Tab 250 mg – For acetazolamide oral liquid formulation refe page 207 | 17.03 | 100 | ✓ <u>Diamox</u> |
| | 17.03 | 100 | ✓ <u>Diamox</u> |

‡ safety cap *Three months or six months, as applicable, dispensed all-at-once SENSORY ORGANS

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

SENSORY ORGANS

| | Subsidy (Manufacturer's P \$ | rice) Sub Per | Fully Brand or sidised Generic ✔ Manufacturer |
|--|------------------------------------|----------------------------------|--|
| DORZOLAMIDE HYDROCHLORIDE * Eye drops 2% | 9.77 (17.44) | 5 ml OP | Trusopt |
| DORZOLAMIDE HYDROCHLORIDE WITH TIMOLOL MALEATE * Eye drops 2% with timolol maleate 0.5% | | 5 ml OP | ✔ Cosopt |
| Glaucoma Preparations - Prostaglandin Analogu | ies | | |
| BIMATOPROST * Eye drops 0.03% | 18.50 | 3 ml OP | 🗸 Lumigan |
| * Eye drops 0.005% Hysite to be Sole Supply on 1 October 2015 | 1.50 | 2.5 ml OP | ✓ Hysite |
| TRAVOPROST * Eye drops 0.004% | | 2.5 ml OP | 🖌 Travatan |
| Glaucoma Preparations - Other | | | |
| BRIMONIDINE TARTRATE * Eye drops 0.2% BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE | 4.32 | 5 ml OP | ✓ <u>Arrow-Brimonidine</u> |
| * Eye drops 0.2% with timolol maleate 0.5% | | 5 ml OP | Combigan |
| * Eye drops 1% * Eye drops 2% * Eye drops 4% Subsidised for oral use pursuant to the Standard Formulae * Eye drops 2% single dose - Special Authority see SA0895 | 5.35 7.99 e. | 15 ml OP 15 ml OP 15 ml OP | ✓ Isopto Carpine ✓ Isopto Carpine ✓ Isopto Carpine |
| below – Retail pharmacy | | 20 dose | Minims |

►SA0895 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics

| ATROPINE SULPHATE * Eye drops 1% | 15 ml OP | ✓ <u>Atropt</u> |
|---|----------------------|--|
| CYCLOPENTOLATE HYDROCHLORIDE * Eye drops 1%8.76 | 15 ml OP | ✓ Cyclogyl |
| TROPICAMIDE * Eye drops 0.5% 7.15 * Eye drops 1% 8.66 | 15 ml OP 15 ml OP | ✓ <u>Mydriacyl</u> ✓ <u>Mydriacyl</u> |

| | Subsidy (Manufacturer's F \$ | Price) Sub Per | Fully Brand or osidised Generic ✔ Manufacturer |
|--|------------------------------------|----------------------------|--|
| Preparations for Tear Deficiency | | | |
| For acetylcysteine eye drops refer Standard Formulae, page 210 |) | | |
| HYPROMELLOSE * Eye drops 0.5% | 2.00 | 15 ml OP | |
| | (3.92) | 13 III OF | Methopt |
| HYPROMELLOSE WITH DEXTRAN | | | |
| * Eye drops 0.3% with dextran 0.1% | 2.30 | 15 ml OP | Poly-Tears |
| POLYVINYL ALCOHOL * Eve drops 1.4% | 0.60 | 15 ml OP | 🖌 Vistil |
| * Eye drops 3% | | 15 ml OP | Vistil Forte |
| Preservative Free Ocular Lubricants | | | |
| SA1388 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid Both: Confirmed diagnosis by slit lamp of severe secretory dry 2 Either: | / eye; and | | meeting the following criteria: |
| 2.1 Patient is using eye drops more than four times of2.2 Patient has had a confirmed allergic reaction to patient | , 0 | | |
| Renewal from any relevant practitioner. Approvals valid for 24 m and has benefited from treatment. CARBOMER – Special Authority see SA1388 above – Retail ph Ophthalmic gel 0.3%, 0.5 g | onths where the armacy | • | es to require lubricating eye drops |
| MACROGOL 400 AND PROPYLENE GLYCOL – Special Author | | •• | |
| Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml | | 24 | ✓ Systane Unit Dose |
| SODIUM HYALURONATE [HYALURONIC ACID] – Special Auth Eye drops 1 mg per ml Hylo-Fresh has a 6 month expiry after opening. The Phari is not relevant and therefore only the prescribed dosage t | | 10 ml OP Manual restric | ✓ <u>Hylo-Fresh</u> ction allowing one bottle per month |
| Other Eye Preparations | | | |
| NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1% | 4.15 | 15 ml OP | ✓ Naphcon Forte |
| OLOPATADINE Eye drops 0.1% | 17.00 | 5 ml OP | ✓ Patanol |
| PARAFFIN LIQUID WITH SOFT WHITE PARAFFIN * Eye oint with soft white paraffin | 3.63 | 3.5 g OP | 🗸 Refresh Night Time |
| PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3% | 0 60 | 35 c OP | |
| RETINOL PALMITATE | | 3.5 g OP | ✓ Poly-Visc |
| E HIVE FALMITALE Eye oint 138 mcg per g | 3.80 | 5 g OP | ✔ VitA-POS |

| | Subsidy (Manufacturer's Pri \$ | ce) Sub Per | Fully osidised | Brand or Generic Manufacturer |
|--|---|---|--------------------------------|--|
| Agents Used in the Treatment of Poisonings | | | | |
| Antidotes | | | | |
| ACETYLCYSTEINE – Retail pharmacy-Specialist Inj 200 mg per ml, 10 ml ampoule | | 10 | 🖌 Ma | BL Acetylcysteine artindale Acetylcysteine |
| Inj 200 mg per ml, 30 ml NALOXONE HYDROCHLORIDE a) Up to 5 inj available on a PSO b) Only on a PSO | 219.00 | 4 | 🖌 Ad | cetadote |
| * lnj 400 mcg per ml, 1 ml ampoule | | 5 | 🖌 Ho | ospira |
| Removal and Elimination | | | | |
| CHARCOAL * Oral liq 50 g per 250 mla) Up to 250 ml available on a PSO b) Only on a PSO | 43.50 | 250 ml OP | 🖌 Ca | arbosorb-X |
| DEFERASIROX – Special Authority see SA1492 below – Re Wastage claimable – see rule 3.3.2 on page 13 Tab 125 mg dispersible | | 28 | 🖌 Ex | |
| Tab 250 mg dispersible Tab 500 mg dispersible | | 28 28 | ✓ Ex ✓ Ex | |
| Salues Special Authority for Subsidy Initial application only from a haematologist. Approvals valid All of the following: The patient has been diagnosed with chronic iron ov 2 Deferasirox is to be given at a daily dose not exceed 3 Any of the following: | erload due to congenit | | - | - |
| 3.1 Treatment with maximum tolerated doses of nation therapy have proven ineffective as me 3.2 Treatment with deferiprone has resulted in se 3.3 Treatment with deferiprone has resulted in ar 3.4 Treatment with deferiprone is contraindicated count (ANC) of < 0.5 cells per µL) or recurre | asured by serum ferriti evere persistent vomitir thritis; or due to a history of agr | n levels, liver ng or diarrhoe anulocytosis | or cardi ea; or (defined | ac MRI T2*; or as an absolute neutroph |
| 0.5 - 1.0 cells per μ L). Renewal only from a haematologist. Approvals valid for 2 yea Either: | ars for applications me | eting the follo | wing crit | teria: |
| For the first renewal following 2 years of therapy, the in all three parameters namely serum ferritin, cardiac For subsequent renewals, the treatment has been to in all three parameters namely serum ferritin, cardiac | c MRI T2* and liver MR lerated and has resulte | I T2* levels; ed in clinical : | or | |
| DEFERIPRONE - Special Authority see SA1480 on the next | t page – Retail pharma | cy | | |

| DEFERIPRONE – Special Authority see SA1480 on the nex | xt page – Retail pharn | nacy | | |
|---|------------------------|-----------|-------------------------------|--|
| Tab 500 mg | | 100 | Ferriprox | |
| Oral liq 100 mg per 1 ml | | 250 ml OP | Ferriprox | |

| (N | Subsidy /anufacturer's Price) | Su | Fully bsidised | Brand or Generic |
|----|----------------------------------|-----|-------------------|---------------------|
| | \$ | Per | ~ | Manufacturer |

SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

| DESFERRIOXAMINE MESYLATE * Inj 500 mg vial | | 10 | ✓ Hospira |
|---|----------|----|-------------------------------|
| SODIUM CALCIUM EDETATE | | | |
| * Inj 200 mg per ml, 5 ml | | 6 | |
| | (156.71) | | Calcium Disodium Versenate |

INTRODUCTION

The following extemporaneously compounded products are eligible for subsidy:

- The "Standard Formulae".
- Oral liquid mixtures for patients unable to swallow subsidised solid dose oral formulations.
- The preparation of syringe drivers when prescribed by a general practitioner.
- Dermatological preparations
 - a) One or more subsidised dermatological galenical(s) in a subsidised dermatological base.
 - b) Dilution of proprietary Topical Corticosteroid-Plain preparations with a dermatological base (Retail pharmacyspecialist).
 - c) Menthol crystals only in the following bases:

Aqueous cream

Urea cream 10%

Wool fat with mineral oil lotion

Hydrocortisone 1% with wool fat and mineral oil lotion

Glycerol, paraffin and cetyl alcohol lotion.

Glossary

Dermatological base: The products listed in the Barrier creams and Emollients section and the Topical Corticosteroids-Plain section of the Pharmaceutical Schedule are classified as dermatological bases for the purposes of extemporaneous compounding and are the bases to which the dermatological galenicals can be added. Also the dermatological bases in the Barrier Creams and Emollients section of the Pharmaceutical Schedule can be used for diluting proprietary Topical Corticosteroid-Plain preparations. The following products are dermatological bases:

- Aqueous cream
- Cetomacrogol cream BP
- Collodion flexible
- Emulsifying ointment BP
- Hydrocortisone with wool fat and mineral oil lotion
- Oil in water emulsion
- Urea cream 10%
- White soft paraffin
- Wool fat with mineral oil lotion
- Zinc and castor oil ointment BP
- Proprietary Topical Corticosteroid-Plain preparations

Dermatological galenical: Dermatological galenicals will only be subsidised when added to a dermatological base. More than one dermatological galenical can be added to a dermatological base.

The following are dermatological galenicals:

- Coal tar solution up to 10%
- Hydrocortisone powder up to 5%
- Menthol crystals
- Salicylic acid powder
- Sulphur precipitated powder

Standard formulae: Standard formulae are a list of fomulae for ECPs that are subsidised. Their ingredients are listed under the appropriate therapeutic heading in Section B of the Pharmaceutical Schedule and also in Section C.

Explanatory notes

Oral liquid mixtures

Oral liquid mixtures are subsidised for patients unable to swallow subsidised solid oral dose forms where no suitable alternative proprietary formulation is subsidised. Suitable alternatives include dispersible and sublingual formulations, oral liquid formulations or rectal formulations. Before extemporaneously compounding an oral liquid mixture, other alternatives such as dispersing the solid dose form (if appropriate) or crushing the solid dose form in jam, honey or soft foods such as yoghurt should be explored.

The Emixt website **www.pharminfotech.co.nz** has evidence-based formulations which are intended to standardise compounded oral liquids within New Zealand.

Pharmaceuticals with standardised formula for compounding in Ora products

| Acetazolamide 25 mg/ml | Gabapentin 100 mg/ml | Sildenafil 2 mg/ml |
|----------------------------------|------------------------------------|----------------------------------|
| Allopurinol 20 mg/ml | Gabapentin (Neurontin) 100 mg/ml | Sotalol 5 mg/ml |
| Amlodipine 1 mg/ml | Hydrocortisone 1 mg/ml | Sulphasalazine 100 mg/ml |
| Azathioprine 50 mg/ml | Labetolol 10 mg/ml | Tacrolimus 1 mg/ml |
| Baclofen 10 mg/ml | Levetiracetam 100 mg/ml | Terbinafine 25 mg/ml |
| Carvedilol 1 mg/ml | Levodopa with carbidopa (5 mg lev- | Tramadol 10 mg/ml |
| Clopidogrel 5 mg/ml | odopa + 1.25 mg carbidopa)/ml | Ursodeoxycholic acid 50 mg/ml |
| Diltiazem hydrochloride 12 mg/ml | Metoclopramide 1 mg/ml | Valganciclovir 60 mg/ml* |
| Dipyridamole 10 mg/ml | Metoprolol tartrate 10 mg/ml | Verapamil hydrochloride 50 mg/ml |
| Domperidone 1 mg/ml | Nitrofurantoin 10 mg/ml | |
| Enalapril 1 mg/ml | Pyrazinamide 100 mg/ml | |
| Flecainide 20 mg/ml | Rifabutin 20 mg/ml | |

*Note this is a DCS formulation

PHARMAC endorses the recommendations of the Emixt website and encourages New Zealand pharmacists to use these formulations when compounding is appropriate. The Emixt website also provides stability and expiry data for compounded products. For the majority of products compounded with Ora-Blend, Ora-Blend SF, Ora-Plus, Ora-Sweet or Ora-Sweet SF a four week expiry is appropriate.

Please note that no oral liquid mixture will be eligible for Subsidy unless all the requirements of Section B and C of the Schedule applicable to that pharmaceutical are met.

Some community pharmacies may not have appropriate equipment to compound all of the listed products, please use appropriate clinical judgement.

Subsidy for extemporaneously compounded oral liquid mixtures is based on:

| Solid dose form | qs |
|------------------|---------|
| Preservative | qs |
| Suspending agent | qs |
| Water | to 100% |

or

Solid dose form

Ora-Blend, Ora-Blend SF, Ora-Plus, Ora-Sweet and/or Ora-Sweet SF

Prescribers may prescribe or pharmacists may add extra non-subsidised ingredients such as flavouring and colouring agents, but these extra ingredients will not be reimbursed. The subsidised ingredients in the formula will be reimbursed and a compounding fee paid.

qs to 100%

The majority of extemporaneously compounded oral liquid mixtures should contain a preservative and suspending agent.

- Ora-Blend, Ora-Blend SF, Ora-Plus, Ora-Sweet and Ora-Sweet SF when used correctly are an appropriate preservative
 and suspending agent.
- Methylcellulose 3% is considered a suitable suspending agent and compound hydroxybenzoate solution or methyl hydroxybenzoate 10% solution are considered to be suitable preservatives. Usually 1 ml of these preservative solutions is added to 100 ml of oral liquid mixture.

Some solid oral dose forms are not appropriate for compounding into oral liquid mixtures and should therefore not be used/considered for extemporaneously compounded oral liquid mixtures. This includes long-acting solid dose formulations, enteric coated tablets or capsules, sugar coated tablets, hard gelatin capsules and chemotherapeutic agents.

The following practices will not be subsidised:

- Where a Standard Formula exists in the Pharmaceutical Schedule for a solid dose form, compounding the solid dose form in Ora-Blend, Ora-Blend SF, Ora-Plus, Ora-Sweet and/or Ora-Sweet SF.
- Mixing one or more proprietary oral liquids (eg an antihistamine with pholcodine linctus).
- Extemporaneously compounding an oral liquid with more than one solid dose chemical.
- Mixing more than one extemporaneously compounded oral liquid mixture.
- Mixing one or more extemporaneously compounded oral liquid mixtures with one or more proprietary oral liquids.
- The addition of a chemical/powder/agent/solution to a proprietary oral liquid or extemporaneously compounded oral mixture.

Standard formulae

A list of standard formulae is contained in this section. All ingredients associated with a standard formula will be subsidised and an appropriate compounding fee paid.

Prescribers may prescribe or pharmacists may add extra non-subsidised ingredients, but these extra ingredients will not be reimbursed. The subsidised ingredients in the formula will be reimbursed and a compounding fee paid.


Dermatological Preparations

Proprietary topical corticosteroid preparations may be diluted with a dermatological base (see page 206) from the Barrier Creams and Emollients section of the Pharmaceutical Schedule (Retail pharmacy-Specialist). Dilution of proprietary topical corticosteroid preparations should only be prescribed for withdrawing patients off higher strength proprietary topical corticosteroid products where there is no suitable proprietary product of a lower strength available or an extemporaneously compounded product with up to 5% hydrocortisone is not appropriate. (In general proprietary topical corticosteroid preparations should not be diluted because dilution effects can be unpredictable and may not be linear, and usually there is no stability data available for diluted products).

One or more dermatological galenicals may be added to a dermatological base (including proprietary topical corticosteroid preparations). Prescribers may prescribe or pharmacists may add extra non-subsidised ingredients, but these extra ingredients will not be reimbursed. The subsidised ingredients in the formula will be reimbursed and a compounding fee paid.

The addition of dermatological galenicals to diluted proprietary Topical Corticosteroids-Plain will not be subsidised.

The flow diagram on the next page may assist you in deciding whether or not a dermatological ECP is subsidised.



EXTEMPORANEOUSLY COMPOUNDED PRODUCTS AND GALENICALS

to 100 ml

Standard Formulae

| ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base | qs qs |
|---|--|
| ASPIRIN AND CHLOROFORM APPLICAT Aspirin Soluble tabs 300 mg Chloroform | • |
| CODEINE LINCTUS PAEDIATRIC (3 mg pr Codeine phosphate Glycerol Preservative Water | er 5 ml) 60 mg 40 ml qs to 100 ml |
| CODEINE LINCTUS DIABETIC (15 mg per Codeine phosphate Glycerol Preservative Water | ⁵ ml) 300 mg 40 ml qs to 100 ml |
| FOLINIC MOUTHWASH Calcium folinate 15 mg tab Preservative Water (Preservative should be used if quantity sup more than 5 days. Maximum 500 ml per pro- | |
| MAGNESIUM HYDROXIDE 8% MIXTURE Magnesium hydroxide paste 29% Methyl hydroxybenzoate Water | 275 g 1.5 g to 1,000 ml |
| METHADONE MIXTURE Methadone powder Glycerol Water | qs qs to 100 ml |
| METHYL HYDROXYBENZOATE 10% SOL Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml o mixture) | 10 g to 100 ml |
| OMEPRAZOLE SUSPENSION Omeprazole capules or powder Sodium bicarbonate powder BP | qs 8.4 g |

| PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water | 1 g 70 ml to 100 ml |
|---|---------------------------|
| PHENOBARBITONE SODIUM PAEDIATR LIQUID (10 mg per ml) | IC ORAL |
| Phenobarbitone Sodium | 400 mg |
| Glycerol BP | 4 ml |
| Water | to 40 ml |
| PILOCARPINE ORAL LIQUID | |
| Pilocarpine 4% eye drops | qs |
| Preservative | qs |
| Water | to 500 ml |
| (Preservative should be used if quantity su more than 5 days.) | ipplied is for |
| more man 5 days.) | |
| SALIVA SUBSTITUTE FORMULA | |
| Methylcellulose | 5 g |
| Preservative | qs |
| Water | to 500 ml |
| (Preservative should be used if quantity su more than 5 days. Maximum 500 ml per pr | |
| more than 5 days. Maximum 500 mi per pi | coorpion.) |
| SODIUM CHLORIDE ORAL LIQUID | |
| Sodium chloride inj 23.4%, 20 ml | qs |
| Water | qs |
| (Only funded if prescribed for treatment of | hyponatraemia) |
| VANCOMYCIN ORAL SOLUTION (50 mg | ner ml) |
| Vancomycin 500 mg injection | 10 vials |
| Glycerol BP | 40 ml |
| Water | to 100 ml |
| (Only funded if prescribed for treatment of difficile following metronidazole failure) | Clostridium |
| - , | |
| VOSOL EAR DROPS | |
| WITH HYDROCORTISONE POWDER 1% | |
| Hydrocortisone powder Vosol Ear Drops | 1% to 35 ml |
| VUSUI EAI DIUPS | 10 33 111 |

Water

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

| | Subsidy | | Fully Brand or |
|--|---------------------|----------------|----------------------------------|
| | (Manufacturer's F | | bsidised Generic |
| | \$ | Per | Manufacturer |
| Extemporaneously Compounded Preparations | and Galenica | s | |
| BENZOIN | | | |
| Tincture compound BP | 2.44 | 50 ml | |
| | (5.10) | | Pharmacy Health |
| | 24.42 | 500 ml | |
| | (39.90) | 50 ml | Pharmacy Health |
| | 2.44 (5.93) | 50 mi | Home Essentials |
| Home Essentials Tincture compound BP to be delisted 1 Dece | ``` | | |
| CHLOROFORM – Only in combination | , | | |
| Only in aspirin and chloroform application. | | | |
| Chloroform BP | 25.50 | 500 ml | ✔ PSM |
| CODEINE PHOSPHATE – Safety medicine; prescriber may det | | frequency | |
| Powder – Only in combination | | 5 g | |
| | (25.46) | - 3 | Douglas |
| | 63.09 | 25 g | Ũ |
| | (90.09) | | Douglas |
| a) Only in extemporaneously compounded codeine linctu | | | ediatric. |
| b) ‡ Safety cap for extemporaneously compounded oral | liquid preparations | i. | |
| COLLODION FLEXIBLE | | | 4 |
| Collodion flexible | 19.30 | 100 ml | V PSM |
| COMPOUND HYDROXYBENZOATE – Only in combination | | | |
| Only in extemporaneously compounded oral mixtures. | | | |
| Soln | | 100 ml | ✓ Midwest |
| | 34.18 | | David Craig |
| GLYCERIN WITH SODIUM SACCHARIN – Only in combination | า | | |
| Only in combination with Ora-Plus. | 05 50 | 470 ml | |
| Suspension | | 473 ml | Ora-Sweet SF |
| GLYCERIN WITH SUCROSE – Only in combination | | | |
| Only in combination with Ora-Plus. | 05 50 | 470 ml | |
| Suspension | | 473 ml | Ora-Sweet |
| GLYCEROL | | | |
| Liquid – Only in combination | | 500 ml | healthE Glycerol BP |
| Only in extemporaneously compounded oral liquid prepa | irations. | | |
| /IAGNESIUM HYDROXIDE Paste 29% | 00.61 | 500 a | ✔ PSM |
| | 22.01 | 500 g | V PSM |
| METHADONE HYDROCHLORIDE | | | |
| a) Only on a controlled drug form | | | |
| b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing from the second seco | equency | | |
| d) Extemporaneously compounded methadone will only be | | rate of the ch | neapest form available (methador |
| powder, not methadone tablets). | . c.mbarood at the | | |
| Powder | 7.84 | 1 g | 🖌 AFT |
| ‡ Safety cap for extemporaneously compounded oral liquestication | uid preparations. | č | |
| METHYL HYDROXYBENZOATE | | | |
| Powder | 8.00 | 25 g | 🖌 PSM |
| | 8.98 | | ✓ Midwest |

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

| | Subsidy (Manufacturer's Pri \$ | ce) Per | Fully Subsidised | |
|--|--------------------------------------|---------------------|---------------------|---------------------|
| METHYLCELLULOSE | | | | |
| Powder Suspension – Only in combination | | 100 g 473 ml | - | MidWest Ora-Plus |
| METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHA Suspension | | mbination 473 ml | | Ora-Blend SF |
| METHYLCELLULOSE WITH GLYCERIN AND SUCROSE – Only Suspension | | 473 ml | ~ | Ora-Blend |
| PHENOBARBITONE SODIUM Powder – Only in combination | 52 50 | 10 a | | MidWest |
| | 325.00 | 10 g | - | MidWest |
| a) Only in children up to 12 years b) ‡ Safety cap for extemporaneously compounded oral liquing | uid preparations. | | | |
| PROPYLENE GLYCOL | | | | |
| Only in extemporaneously compounded methyl hydroxybenzo | | 500 ml | | PSM |
| Liq | | 500 m | - | Midwest |
| SODIUM BICARBONATE | | | | |
| Powder BP – Only in combination | | 500 g | ~ | Midwest |
| | 9.80 (29.50) | | | David Craig |
| Only in extemporaneously compounded omeprazole and la | nsoprazole suspe | ension. | | |
| SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparatior | IS. | | | |
| | 21.75 | 2,000 m | I 🗸 | Midwest |
| WATER Tap – Only in combination | 0.00 | 1 ml | ~ | Tap water |

EXPLANATORY NOTES

The list of special foods to which Subsidies apply is contained in this section. The list of available products, guidelines for use, subsidies and charges is reviewed as required. Applications for new listings and changes to subsidies and access criteria will be considered by the special foods sub-committee of PTAC which meets as and when required. In all cases, subsidies are available by Special Authority only. This means that, unless a patient has a valid Special Authority number for their special food requirements, they must pay the full cost of the products themselves.

Eligibility for Special Authority

Special Authorities will be approved for patients meeting conditions specified under the *Conditions and Guidelines* for each product. In some cases there are also limits to how products can be prescribed (for example quantity, use or duration). Only those brands, presentations and flavours of special foods listed in this section are subsidised.

Who can apply for Special Authority?

 Initial Applications:
 Only from a dietitian, relevant specialist or a vocationally registered general practitioner.

 Reapplications:
 Only from a dietitian, relevant specialist or a vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or a vocationally registered general practitioners.

 Very specialist or a vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or a vocationally registered general practitioner. Other general practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

All applications must be made on an official form available from the PHARMAC website www.pharmac.govt.nz. All applications must include specific details as requested on the form relating to the application. Applications must be forwarded to:

Ministry of Health Sector Services Private Bag 3015 WHANGANUI 4540 Freefax 0800 100 131

Subsidies and manufacturer's surcharges

The Subsidies for some special foods are based on the lowest priced product within each group. Where this is so, or where special foods are otherwise not fully subsidised, a manufacturer's surcharge may be payable by the patient. The manufacturer's surcharge is the difference between the price of the product and the subsidy attached to it and may be subject to mark-ups applied at a pharmacy level. As a result the manufacturer's surcharge may vary. Fully subsidised alternatives are available in most cases (as indicated by a tick in the left hand column). Patients should only have to pay a co-payment on these products.

Where are special foods available from?

Distribution arrangements for special foods vary from region to region. Special foods are available from hospital pharmacies providing an outpatient dispensing service as well as retail pharmacies in the Northern, Midland and Central (including Nelson and Blenheim) regions.

Definitions

 Failure to thrive
 An inability to gain or maintain weight resulting in physiological impairment.

 Growth deficiency
 Where the weight of the child is less than the fifth or possibly third percentile for their age, with evidence of malnutrition

Dietitian Prescribing

Prescriptions from Dietitians will be only valid for subsidy where they are for special foods, as listed in this section, or where they are for the following products:

ASCORBIC ACID Tab 100 mg

CALCIUM CARBONATE

- ✓ Tab eff 1.75 g (1 g elemental)
- ✓ Tab 1.25 g (500 mg elemental)
- COMPOUND ELECTROLYTES

Powder for oral soln

DEXTROSE WITH ELECTROLYTES Soln with electrolytes

FERROUS FUMARATE ✓ Tab 200 mg (65 mg elemental)

FERROUS FUMARATE WITH FOLIC ACID

✓ Tab 310 mg (100 mg elemental) with folic acid 350 mcg

FERROUS SULPHATE

- ✓ Tab long-acting 325 mg (105 mg elemental)
- ✓ Oral liq 30 mg (6 mg elemental) per 1 ml

FERROUS SULPHATE WITH FOLIC ACID Tab long-acting 325 mg (105 mg elemental) with folic acid 350 mcg

FOLIC ACID Tab 0.8 mg

MULTIVITAMINS

PANCREATIC ENZYME

✓ Cap EC 10,000 BP u lipase, 9,000 BP u amylase and 210 BP u protease PHOSPHORUS ✓ Tab eff 500 mg (16 mmol)

POTASSIUM CHLORIDE

Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)

Tab long-acting 600 mg (8 mmol)

POTASSIUM IODATE

✓ Tab 253 mcg (150 mcg elemental iodine)

PYRIDOXINE HYDROCHLORIDE

- ✔ Tab 25 mg
- ✔ Tab 50 mg

SODIUM CHLORIDE

✓ Inj 23.4%, 20 ml ampoule

SODIUM FLUORIDE

✓ Tab 1.1 mg (0.5 mg elemental)

THIAMINE HYDROCHLORIDE ✓ Tab 50 mg

VITAMIN A WITH VITAMINS D AND C

✓ Soln 1000 u with Vitamin D 400 u and ascorbic acid 30 mg per 10 drops

VITAMIN B COMPLEX

✓ Tab, strong, BPC

VITAMINS

- ✓ Tab (BPC cap strength)
- ✓ Cap (fat soluble vitamins A, D, E, K)

Subsidy (Manufacturer's Price) \$ Per Brand or Generic Manufacturer

Fully

Subsidised

Nutrient Modules

Carbohydrate

SA1522 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Either:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children; or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 inborn errors of metabolism; or
- 7 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Cystic fibrosis or renal failure)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT – Special Authority see SA1522 above – Hospital pharmacy [HP3]

| Powder5.29 | 400 g OP | Polycal |
|------------|----------|---------|
|------------|----------|---------|

Carbohydrate And Fat

SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

| Subsidy (Manufacturer's Price) | Fully ce) Subsidised | | Brand or Generic | |
|-----------------------------------|-------------------------|---|---------------------|--|
| \$ | Per | ~ | Manufacturer | |

continued...

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| CARBOHYDRATE AND FAT SUPPLEMENT | - Special Authority see SA1376 | on the previous p | age – Hospital pharmacy [HP3] |
|---------------------------------|--------------------------------|-------------------|-------------------------------|
| Powder (neutral) | | 400 g OP | ✓ Duocal Super |
| | | - | Soluble Powder |

Fat

➡SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome; or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or
- 10 acites; or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | S | ubsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Authority see SA1523 on the previous page - Hospital pharmacy [HP3]

| Emulsion (neutral) | 200 ml OP | ✓ Calogen |
|----------------------------|-----------|------------------------------|
| 30.75 | 500 ml OP | Calogen |
| Emulsion (strawberry)12.30 | 200 ml OP | Calogen |
| Oil | 500 ml OP | MCT oil (Nutricia) |
| Oil, 250 ml | 4 OP | Liquigen |

Protein

➡SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT - Special Authority see SA1524 above - Hospital pharmacy [HP3]

| | | ~····~~) [···· o] | |
|------------------|------|-------------------|-------------|
| Powder | | 225 g OP | Protifar |
| | 8.95 | 227 g OP | Resource |
| | | | Beneprotein |
| Powder (vanilla) | | 275 g OP | Promod |

| | | Subsidy (Manufacturer's Price) \$ | Fully Subsidised Per 🖌 | Generic |
|--|---|---|---|---|
| Oral | Supplements/Complete Diet (Nasogastric/ | Gastrostomy Tub | e Feed) | |
| Res | piratory Products | | | |
| Initial where Renew menda meetin Both: 1 | 1094 Special Authority for Subsidy application only from a dietitian, relevant specialist or voc the patient has CORD and hypercapnia, defined as a CO2 val only from a dietitian, relevant specialist, vocationally re tion of a dietitian, relevant specialist or vocationally registe g the following criteria: The treatment remains appropriate and the patient is be General Practitioners must include the name of the die | 2 value exceeding 55 rr gistered general practi ared general practitione nefiting from treatment | imHg. tioner or genera r. Approvals va ; and | al practitioner on the recom lid for 1 year for applications |
| | tioner and date contacted. ORAL FEED 1.5KCAL/ML – Special Authority see SA10 guid | | | Pulmocare |
| Diab | etic Products | | | |
| Initial where Renew menda | 1095 Special Authority for Subsidy application only from a dietitian, relevant specialist or voc the patient is a type I or and II diabetic who is suffering we val only from a dietitian, relevant specialist, vocationally re tion of a dietitian, relevant specialist or vocationally registe g the following criteria: | eight loss and malnutriti | on that require | s nutritional support. al practitioner on the recom |
| | The treatment remains appropriate and the patient is be General Practitioners must include the name of the die tioner and date contacted. | | | ly registered general practi |
| DIABE | TIC ENTERAL FEED 1KCAL/ML - Special Authority see | SA1095 above - Hosp | ital pharmacy [| HP3] |

| Liquid | 7.50 | 1,000 ml OP | Diason RTH Glucerna Select RTH |
|---|-----------------------|-----------------|---|
| DIABETIC ORAL FEED 1KCAL/ML - Special Authority | see SA1095 above - Ho | spital pharmacy | [HP3] |
| Liquid (strawberry) | 1.50 | 200 ml OP | Diasip |
| Liquid (vanilla) | 1.50 | 200 ml OP | Diasip |
| | 1.88 | 250 ml OP | Glucerna Select |
| | 1.78 | 237 ml OP | |
| | (2.10) | | Resource Diabetic |
| | (2.10) | | Sustagen Diabetic |

Fat Modified Products

SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or

continued...

| Subsidy (Manufacturer's Price) | Sub | Fully | Brand or Generic | |
|-----------------------------------|-----|-------|---------------------|--|
| \$ | Per | ~ | Manufacturer | |

continued...

3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED - Special Authority see SA1525 on the previous page - Hospital pharmacy [HP3]

| Powder | | | | 60.48 | 400 g OP | Monogen |
|--------|--|--|--|-------|----------|---------|
|--------|--|--|--|-------|----------|---------|

High Protein Products

➡SA1378 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 decompensating liver disease without encephalopathy; or
- 2 protein losing gastro-enteropathy.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

HIGH PROTEIN ORAL FEED 1KCAL/ML – Special Authority see SA1378 above – Hospital pharmacy [HP3]

Liquid1.90 200 ml OP 🖌 Fortimel Regular

(Fortimel Regular Liquid to be delisted 1 September 2015)

Paediatric Products For Children Awaiting Liver Transplant

➡SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| ENTERAL/ORAL FEED 1KCAL/ML - S | Special Authority see SA1098 above - | Hospital pharmacy | ' [HP3] |
|--------------------------------|--------------------------------------|-------------------|----------------|
| Powder (unflavoured) | | 400 g OP | Heparon Junior |

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

Paediatric Products For Children With Chronic Renal Failure

SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

Paediatric Products

SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| PAEDIATRIC ENTERAL FEED 1KCAL/ML – Special Authority see SA1379 above Liquid2.68 | | macy [HP3] V Nutrini RTH Pediasure RTH |
|---|--|---|
| PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority see Liquid6.00 | | Hospital pharmacy [HP3] Nutrini Energy Multi Fibre Nutrini Energy RTH |
| PAEDIATRIC ORAL FEED – Special Authority see SA1379 above – Hospital phan Powder (vanilla) | macy [HP3] 850 g OP | ✓ Pediasure |
| PAEDIATRIC ORAL FEED 1.5KCAL/ML – Special Authority see SA1379 above – Liquid (strawberry) | Hospital pharm 200 ml OP 200 ml OP | acy [HP3] ✔ Fortini ✔ Fortini |

Specialised And Elemental Products

➡SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 malabsorption: or
 - 2 short bowel syndrome; or
 - 3 enterocutaneous fistulas; or
 - 4 eosinophilic oesophagitis: or
 - 5 inflammatory bowel disease; or
 - 6 patients with multiple food allergies requiring enteral feeding.

continued...

SPECIAL FOODS

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

continued...

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| ENTERAL/ORAL ELEMENTAL FEED 1KCAL/ML | Special Authority see SA1377 | on the previous | s page – Hospital pharmacy [HP3] | |
|--------------------------------------|--|-----------------|----------------------------------|--|
| Powder | 4 40 | 79 a OP | Vital HN | |

| 7.50 | 76 g OP | ✓ Alitraq |
|---|-----------------|---|
| ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see SA1377 on the | previous page - | - Hospital pharmacy [HP3] |
| Liquid (grapefruit), 250 ml carton171.00 | 18 OP | Elemental 028 Extra |
| Liquid (pineapple & orange), 250 ml carton 171.00 | 18 OP | Elemental 028 Extra |
| Liquid (summer fruits), 250 ml carton171.00 | 18 OP | Elemental 028 Extra |
| ORAL ELEMENTAL FEED 1KCAL/ML – Special Authority see SA1377 on the pr Powder (unflavoured)4.50 | | Hospital pharmacy [HP3] |
| SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML – Special Authority see SA1377 Liquid12.04 | | s page – Hospital pharmacy [HP3] |

Paediatric Products For Children With Low Energy Requirements

➡SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| • | | | Multi Fibre |
|--|---------------------|----------------|-----------------------------|
| Liquid | 4.00 | 500 ml OP | Nutrini Low Energy |
| PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML - Sp | pecial Authority se | e SA1196 above | e – Hospital pharmacy [HP3] |

| (Manufacturer's Price) | Fully ice) Subsidised | | Brand or Generic | |
|------------------------|--------------------------|---|---------------------|--|
| \$ | Per | ~ | Manufacturer | |

Standard Supplements

SA1228 Special Authority for Subsidy

Initial application — (Children) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Adults) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - Patient is Malnourished
 - 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m2; or
 - 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
 - 1.3 Patient has a BMI of less than 20 kg/m2 and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:
 - Patient has not responded to first-line dietary measures over a 4 week period by:
 - 2.1 Increasing their food intake frequency (eg snacks between meals); or
 - 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
 - 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:
 - Patient is Malnourished
 - 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m2; or
 - 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
 - 2.3 Patient has a BMI of less than 20 kg/m2 and unintentional weight loss greater than 5% within the last 3-6 months.

continued...

| Subsidy (Manufacturer's Price | | Fully Subsidised | Brand or Generic | |
|----------------------------------|-----|---------------------|---------------------|--|
| \$ | Per | ~ | Manufacturer | |

continued...

Initial application - (Adults transitioning from hospital Discretionary Community Supply) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 The patient has had up to a 30 day supply of a 1.0 or a 1.5 kcal/ml Standard Oral Supplement; and
- 2 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 3 Any of the following:
 - Patient is Malnourished
 - 3.1 Patient has a body mass index (BMI) of less than 18.5 kg/m2; or
 - 3.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
 - 3.3 Patient has a BMI of less than 20 kg/m2 and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 Is being feed via a nasogastric tube or a nasogastric tube is to be inserted for feeding: or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant: and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (<13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum: or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result: or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant: and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (<13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum: or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

continued.

| Subsidy | | Fully | Brand or | |
|----------------------|-----|------------|--------------|--|
| (Manufacturer's Pric | e) | Subsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

Initial application - (Long-term medical condition) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria): or
- 2 Cystic Fibrosis; or
- 3 Liver disease: or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease: or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria): or
- 2 Cystic Fibrosis; or
- 3 Liver disease: or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease: or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

| ENTERAL FEED 1.5KCAL/ML – Special Authority see SA1228 on page 223 – Liquid | | y [HP3] ✔ Nutrison Energy |
|--|--------------------------|--|
| ENTERAL FEED 1KCAL/ML - Special Authority see SA1228 on page 223 - He | ospital pharmacy | [HP3] |
| Liquid | 250 ml OP | ✓ Isosource Standard ✓ Osmolite |
| 5.29 | 1,000 ml OP | Isosource Standard RTH |
| | | Nutrison Standard RTH |
| 2.65 5.29 | 500 ml OP 1,000 ml OP | Osmolite RTH Osmolite RTH |
| ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority see SA1228 on | 1 0 1 | |
| Liquid | 237 ml OP | ✓ Jevity |
| 2.65 | 500 ml OP | , |
| 5.29 | 1,000 ml OP | |
| | | Nutrison Multi Fibre |

| | Subsidy (Manufacturer's \$ | Price) Sub Per | Fully Brand or isidised Generic ✔ Manufacturer |
|---|----------------------------------|----------------------|--|
| ENTERAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority s | ee SA1228 on | page 223 – Ho | spital pharmacy [HP3] |
| Liquid | 1.75 | 250 ml OP | Ensure Plus HN |
| | 7.00 | 1,000 ml OP | Ensure Plus RTH Jevity HiCal RTH Nutrison Energy Multi Fibre |
| ORAL FEED (POWDER) - Special Authority see SA1228 on page | e 223 – Hospita | al pharmacy [Hi | 23] |
| Note: Higher subsidy for Sustagen Hospital Formula will onl number and an appropriately endorsed prescription. Powder (chocolate) – Higher subsidy of up to \$14.90 per | y be reimburse | ed for patients v | vith both a valid Special Authority |
| 900 g with Endorsement | 13.00 10.22 | 850 g OP 900 g OP | Ensure |
| | (14.90) | | Sustagen Hospital Formula |
| Additional subsidy by endorsement is available for patient | s with fat mala | bsorption, fat in | tolerance or chyle leak. The pre- |
| scription must be endorsed accordingly. | | | |
| Powder (vanilla) – Higher subsidy of up to \$14.90 per 900 g | | | |
| with Endorsement | | 350 g OP | ✓ Fortisip |
| | 13.00 | 850 g OP | Ensure |
| | 10.22 | 900 g OP | |
| | (14.90) | | Sustagen Hospital Formula |
| Additional subsidy by endorsement is available for patient scription must be endorsed accordingly. | s with fat mala | bsorption, fat in | tolerance or chyle leak. The pre |

| | Subsidy (Manufacturer's \$ | | Fully Brand or dised Generic ✔ Manufacturer |
|--|----------------------------------|------------------|---|
| ORAL FEED 1.5KCAL/ML – Special Authority see SA1228 on pa | ge 223 – Hospi | tal pharmacy [HP | 3] |
| Additional subsidy by endorsement is available for patients be | ing bolus fed th | | |
| molysis bullosa. The prescription must be endorsed according | | | |
| Liquid (banana) - Higher subsidy of \$1.26 per 200 ml with | | | |
| Endorsement | | 200 ml OP | |
| | (1.26) | | Ensure Plus |
| | (1.26) | | Fortisip |
| Liquid (chocolate) – Higher subsidy of up to \$1.33 per 237 ml | | 000 | |
| with Endorsement | | 200 ml OP | |
| | (1.26) | 007 | Ensure Plus |
| | 0.85 | 237 ml OP | |
| | (1.33) 0.72 | 200 ml OP | Ensure Plus |
| | (1.26) | 200 IIII OF | Forticip |
| Liquid (fruit of the fareat) Llipher subsidu of \$1.06 per 000 ml | () | | Fortisip |
| Liquid (fruit of the forest) – Higher subsidy of \$1.26 per 200 ml with Endorsement | | 200 ml OP | |
| | | 200 IIII OF | Ensure Plus |
| Liquid (strowborn) Higher subsidy of \$1.26 per 200 ml with | () | | LISUIE Flus |
| Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with Endorsement | | 200 ml OP | |
| Endorsement | | 200 IIII OF | Ensure Plus |
| | (1.26) | | Fortisip |
| Liquid (toffac) Higher subsidy of \$1.26 per 200 ml with | . , | | Fortisip |
| Liquid (toffee) – Higher subsidy of \$1.26 per 200 ml with Endorsement | | 200 ml OP | |
| | (1.26) | 200 IIII OF | Fortisip |
| Liquid (tropical fruit) - Higher subsidy of \$1.26 per 200 ml | · · · | | i oi usip |
| with Endorsement | | 200 ml OP | |
| | (1.26) | 200 IIII OF | Fortisip |
| Liquid (vanilla) - Higher subsidy of up to \$1.33 per 237 ml | () | | i oi usip |
| with Endorsement | | 200 ml OP | |
| | (1.26) | 200 IIII OF | Ensure Plus |
| | 0.85 | 237 ml OP | Elistic Flus |
| | (1.33) | 207 111 01 | Ensure Plus |
| | 0.72 | 200 ml OP | |
| | (1.26) | 200 0. | Fortisip |
| (Fortisip Liquid (toffee) to be delisted 1 September 2015) (Fortisip Liquid (tropical fruit) to be delisted 1 September 2015) | (| | |
| | 041000 | •• 000 User't 1 | n h a mar a su [l ID0] |
| ORAL FEED WITH FIBRE 1.5 KCAL/ML – Special Authority see Additional subsidy by endorsement is available for patients be molysis bullosa. The prescription must be endorsed accordin Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with | eing bolus fed th gly. | | |
| Endorsement | | 200 ml OP | |
| | (1.26) | | Fortisip Multi Fibre |
| Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with | (/ | | - F |
| Endorsement | | 200 ml OP | |
| | | | |

SPECIAL FOODS

| Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per 🖌 Manufacture | er |
|---|----|
|---|----|

High Calorie Products

►SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment: and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 above | e – Hospital | pharmacy [HP3] | |
|---|--------------|----------------|--|
| Liquid | 5.50 | 500 ml OP | Nutrison Concentrated |
| | 11.00 | 1,000 ml OP | 🖌 Two Cal HN RTH |
| ORAL FEED 2 KCAL/ML – Special Authority see SA1195 above – H Additional subsidy by endorsement is available for patients being molysis bullosa. The prescription must be endorsed accordingly. Liquid (vanilla) – Higher subsidy of \$1.90 per 200 ml with | bolus fed th | | tube, or who have severe epider- |
| Endorsement | 0.96 | 200 ml OP | |
| | (1.90) | | Two Cal HN |

| | Subsidy (Manufacturer's Pric \$ | e) Sul Per | Fully bsidised | Brand or Generic Manufacturer |
|--|---------------------------------------|---------------|-------------------|-------------------------------------|
| Food Thickeners | • | | | |
| ⇒SA1106 Special Authority for Subsidy itial application only from a dietitian, relevant specialist or voc here the patient has motor neurone disease with swallowing dis enewal only from a dietitian, relevant specialist, vocationally re endation of a dietitian, relevant specialist or vocationally register teeting the following criteria: oth: | sorder. gistered general pra | ctitioner or | general | practitioner on the recon |
| The treatment remains appropriate and the patient is be General Practitioners must include the name of the die tioner and date contacted. | • | | ationally | v registered general prac |
| OOD THICKENER - Special Authority see SA1106 above - H | ospital pharmacy [H | P3] | | |
| Powder | 6.53 | 300 g OP | 🖌 N | lutilis |
| | 7.25 | 380 g OP | 🗸 F | eed Thickener Karicare Aptamil |

The funding of gluten free foods is no longer being actively managed by PHARMAC from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

SA1107 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

1 Gluten enteropathy has been diagnosed by biopsy; or

2 Patient suffers from dermatitis herpetiformis.

| GLUTEN FREE BAKING MIX – Special Authority see SA1107 Powder | | pharmacy [HP3] 1,000 g OP | |
|---|-------------------|------------------------------|--|
| | (5.15) | - | Healtheries Simple Baking Mix |
| GLUTEN FREE BREAD MIX - Special Authority see SA1107 a | bove – Hospital p | harmacy [HP3] | |
| Powder | | 1,000 g OP | |
| | (7.32) | - | NZB Low Gluten Bread Mix |
| | 4.77 | | |
| | (8.71) | | Bakels Gluten Free Health Bread Mix |
| | 3.51 | | |
| | (10.87) | | Horleys Bread Mix |
| GLUTEN FREE FLOUR – Special Authority see SA1107 above | | | |
| Powder | 5.62 (18.10) | 2,000 g OP | Horleys Flour |

| | Subsidy (Manufacturer's \$ | | ully Brand or Sed Generic Manufacturer | | | | |
|---|----------------------------------|----------|--|--|--|--|--|
| GLUTEN FREE PASTA – Special Authority see SA1107 on the previous page – Hospital pharmacy [HP3] | | | | | | | |
| Buckwheat Spirals | 2.00 | 250 g OP | | | | | |
| | (3.11) | | Orgran | | | | |
| Corn and Vegetable Shells | | 250 g OP | | | | | |
| | (2.92) | | Orgran | | | | |
| Corn and Vegetable Spirals | | 250 g OP | | | | | |
| | (2.92) | | Orgran | | | | |
| Rice and Corn Lasagne Sheets | | 200 g OP | - | | | | |
| | (3.82) | | Orgran | | | | |
| Rice and Corn Macaroni | | 250 g OP | • | | | | |
| | (2.92) | | Orgran | | | | |
| Rice and Corn Penne | | 250 g OP | 0 | | | | |
| | (2.92) | | Orgran | | | | |
| Rice and Maize Pasta Spirals | | 250 g OP | 0 | | | | |
| Disc and Millet Onionia | (2.92) | 050 00 | Orgran | | | | |
| Rice and Millet Spirals | | 250 g OP | 0 | | | | |
| D'es and some successive Warranties | (3.11) | 075 . 00 | Orgran | | | | |
| Rice and corn spaghetti noodles | | 375 g OP | 0 | | | | |
| Verstehle and Rice Onivela | (2.92) | 050 - 00 | Orgran | | | | |
| Vegetable and Rice Spirals | | 250 g OP | Oraran | | | | |
| Italian lang at ta ang ta ti | (2.92) | 000 - 00 | Orgran | | | | |
| Italian long style spaghetti | | 220 g OP | Overver | | | | |
| | (3.11) | | Orgran | | | | |

Foods And Supplements For Inborn Errors Of Metabolism

SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

| AMINOACID FORMULA WITHOUT METHIONINE – Special Powder | • |)8 above – Hos 500 g OP | pital pharmacy [HP3] |
|---|---|------------------------------|-------------------------------|
| Supplements For MSUD | | | |
| AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND pharmacy [HP3] Powder | | Special Authorit 500 g OP | y see SA1108 above – Hospital |

| S | Subsidy | Fully Brai | nd or |
|----------|-------------------------|------------|------------|
| (Manufac | acturer's Price) Subsid | lised Ger | neric |
| | \$ Per | 🖌 Mar | nufacturer |

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

| Tabs | | 75 OP | Phlexy 10 |
|--|--------|-----------|-------------------|
| Powder (unflavoured) 29 g sachets | | 30 | PKU Anamix Junior |
| | | 400 g OP | PKU Anamix Infant |
| Powder (orange) | | 500 g OP | XP Maxamaid |
| | 320.00 | 0 | XP Maxamum |
| Powder (unflavoured) | | 500 g OP | XP Maxamaid |
| · · · · · · | 320.00 | 0 | XP Maxamum |
| Liquid (berry) | | 125 ml OP | PKU Anamix Junior |
| | | | LQ |
| Liquid (orange) | | 125 ml OP | PKU Anamix Junior |
| | | | LQ |
| Liquid (unflavoured) | | 125 ml OP | PKU Anamix Junior |
| | | | LQ |
| Liquid (forest berries), 250 ml carton | | 18 OP | 🖌 Easiphen Liquid |
| Liquid (juicy berries) 62.5 ml | | 60 OP | PKU Lophlex LQ 10 |
| | | 60 OP | PKU Lophlex LQ 10 |
| | | 60 OP | PKU Lophlex LQ 10 |
| | | 30 OP | PKU Lophlex LQ 20 |
| Liquid (juicy citrus) 125 ml | | 30 OP | PKU Lophlex LQ 20 |
| | | 30 OP | PKU Lophlex LQ 20 |
| · · · · · · · · · · · · · · · · · · · | | | |

Foods

| LOW PROTEIN BAKING MIX - Special Authority see SA1108 on the prev | 1 0 | | , , , |
|--|-----------|---------------|----------------------------------|
| Powder | 8.22 | 500 g OP | Loprofin Mix |
| LOW PROTEIN PASTA - Special Authority see SA1108 on the previous p | bage – Ho | spital pharma | cy [HP3] |
| Animal shapes11 | 1.91 | 500 g OP | Loprofin |
| Lasagne | 5.95 | 250 g OP | Loprofin |
| Low protein rice pasta11 | 1.91 | 500 g OP | Loprofin |
| Macaroni | 5.95 | 250 g OP | Loprofin |
| Penne11 | 1.91 | 500 g OP | Loprofin |
| Spaghetti11 | 1.91 | 500 g OP | Loprofin |
| Spirals1 | 1.91 | 500 g OP | Loprofin |

Infant Formulae

For Premature Infants

| PRETERM POST-DISCHARGE INFANT FORMULA - | Special Authority see SA1 | 198 on the next | page – Hospital pharmacy [HP3] |
|---|---------------------------|-----------------|--------------------------------|
| Powder | | 400 g OP | S-26 Gold Premgro |

| Subsidy | Fully | Brand or |
|-----------------|-------------------|--------------|
| (Manufacturer's | Price) Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

➡SA1198 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The infant was born before 33 weeks gestation or weighed less than 1.5 kg at birth; and
- 2 Either:
 - 2.1 The infant has faltering growth (downward crossing of percentiles); or
 - 2.2 The infant is not maintaining, or is considered unlikely to maintain, adequate growth on standard infant formula.

For Williams Syndrome

SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA – Special Authority see SA1110 above – Hospital pharmacy [HP3]

| Powder | | 4 | 400 g OP | ~ | Locasol |
|--------|------|---|----------|---|---------|

Gastrointestinal and Other Malabsorptive Problems

| Powder | 6.00 | 48.5 g OP | Vivonex Pediatric |
|----------------------|-------|-----------|-------------------|
| | 53.00 | 400 g OP | Neocate LCP |
| Powder (unflavoured) | 53.00 | 400 g OP | Elecare |
| | | - | Elecare LCP |
| | | | Neocate Advance |
| | | | Neocate Gold |
| Powder (vanilla) | 53.00 | 400 g OP | Elecare |
| · · · | | 5 | Neocate Advance |

➡SA1219 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Extensively hydrolysed formula has been reasonably trialled and is inappropriate due to documented severe intolerance or allergy or malabsorption; or
- 2 History of anaphylaxis to cows milk protein formula or dairy products; or
- 3 Eosinophilic oesophagitis.
- Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

continued...

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | Su | Ibsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein, soy, or extensively hydrolysed infant formula has been undertaken: and
- 2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

```
EXTENSIVELY HYDROLYSED FORMULA - Special Authority see SA1380 below - Hospital pharmacy [HP3]
```

| Powder15.21 | 450 g OP | Pepti Junior Gold |
|-------------|----------|-------------------|
| | | Karicare Aptamil |

➡SA1380 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria: Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption: or
- 3 Short bowel syndrome: or
- 4 Intractable diarrhea: or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption: or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken: and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Step Down from Amino Acid Formula) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 The infant is currently receiving funded amino acid formula: and
- 2 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

| Subsic (Manufacture \$ | | ubsidised | Brand or Generic Manufacturer |
|------------------------------|------|-----------|-------------------------------------|
| Ψ | 1.01 | • | Manuacturer |

Ketogenic Diet

SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

| Powder (unflavoured) | 300 g OP | KetoCal 4:1 Ketocal 3:1 |
|-----------------------|----------|--|
| Powder (vanilla)35.50 | 300 g OP | ✓ KetoCal 4:1 |

Pharmaceuticals and quantities that may be obtained on a Practitioner's Supply Order

| ADRENALINE ✓ Inj 1 in 1,000, 1 ml ampoule5 ✓ Inj 1 in 10,000, 10 ml ampoule5 |
|--|
| AMINOPHYLLINE ✓ Inj 25 mg per ml, 10 ml ampoule5 |
| AMIODARONE HYDROCHLORIDE Inj 50 mg per ml, 3 ml ampoule6 |
| AMOXICILLIN ✓ Cap 250 mg |
| AMOXICILLIN WITH CLAVULANIC ACID ✓ Tab 500 mg with clavulanic acid 125 mg |
| ✓ Grans for oral liq amoxicillin 250 mg with clavulanic acid 62.5 mg per 5 ml 200 ml |
| ASPIRIN V Tab dispersible 300 mg |
| ATROPINE SULPHATE Inj 600 mcg per ml, 1 ml ampoule5 |
| AZITHROMYCIN V Tab 500 mg – See note on page 928 |
| BENDROFLUMETHIAZIDE [BENDROFLUAZIDE] V Tab 2.5 mg – See note on page 56 |
| BENZATHINE BENZYLPENICILLIN Inj 900 mg (1.2 million units) in 2.3 ml syringe |
| BENZTROPINE MESYLATE ✔ Inj 1 mg per ml, 2 ml |
| BENZYLPENICILLIN SODIUM (PENICILLIN G) ✓ Inj 600 mg (1 million units) vial |
| BLOOD GLUCOSE DIAGNOSTIC TEST METER ✓ Meter with 50 lancets, a lancing device and 10 diagnostic test strips – Subsidy by endorsement – See note on page 26 |
| BLOOD GLUCOSE DIAGNOSTIC TEST STRIP ✓ Blood glucose test strips – See note on page 26 |
| BLOOD KETONE DIAGNOSTIC TEST METER Meter – See note on page 251 |

| CEFTRIAXONE | |
|---|----------|
| ✓ Inj 500 mg vial – Subsidy by endorsement See note on page 91 | - |
| Inj 1 g vial – Subsidy by endorsement – Se note on page 91 | е |
| CHARCOAL ✔ Oral liq 50 g per 250 ml | 250 ml |
| CHLORPROMAZINE HYDROCHLORIDE | |
| ✓ Tab 10 mg | 20 |
| ✓ Tab 25 mg | |
| ✓ Tab 25 mg | |
| ✓ Inj 25 mg per ml, 2 ml | |
| CIPROFLOXACIN | |
| ✓ Tab 250 mg – See note on page 95 | 5 |
| ✓ Tab 500 mg – See note on page 95 | |
| CO-TRIMOXAZOLE | |
| | |
| Tab trimethoprim 80 mg and sulphamethoxazole 400 mg | 20 |
| | |
| Oral liq trimethoprim 40 mg and autobarethousards 200 mg par | |
| sulphamethoxazole 200 mg per 5 ml | 200 ml |
| | |
| COMPOUND ELECTROLYTES | |
| Powder for oral soln | 10 |
| CONDOMS | |
| ✓ 49 mm | 144 |
| ✓ 52 mm | |
| ✓ 52 mm extra strength | |
| ✓ 53 mm | |
| ✓ 53 mm (chocolate) | |
| ✓ 53 mm (strawberry) | 144 |
| 54 mm, shaped | |
| ✓ 55 mm | |
| ✔ 56 mm | 144 |
| ✓ 56 mm, shaped | 144 |
| ✔ 60 mm | |
| CYPROTERONE ACETATE ETHINYLOESTRADIOL | WITH |
| ✓ Tab 2 mg with ethinyloestradiol 35 mcg and | ł |
| 7 inert tabs | |
| DEXAMETHASONE | |
| Tab 1 mg – Retail pharmacy-Specialist | 30 |
| Tab 4 mg – Retail pharmacy-Specialist | |
| | |
| DEXAMETHASONE PHOSPHATE | |
| ✓ Inj 4 mg per ml, 1 ml ampoule – See note o | |
| page 79 | 5 |
| c | ontinued |

(continued)

| Inj 4 mg per ml, 2 ml ampoule – See note on page 79 | 5 |
|--|---|
| DIAPHRAGM | |
| ✓ 65 mm – See note on page 73 | 1 |
| ✓ 70 mm – See note on page 73 | 1 |
| ✓ 75 mm – See note on page 73 | 1 |
| ✓ 80 mm – See note on page 73 | 1 |

DIAZEPAM

| ✓ Inj 5 mg per ml, 2 ml ampoule – Subsidy by | |
|--|---|
| endorsement - See note on page 133 | 5 |
| ✓ Rectal tubes 5 mg | 5 |
| ✓ Rectal tubes 10 mg | |

DICLOFENAC SODIUM

| ✓ Inj 25 mg per ml, 3 ml ampoule | 5 |
|----------------------------------|----|
| ✓ Suppos 50 mg | 10 |

DIGOXIN

| 🖌 Tab | o 62.5 mcg | 30 |
|-------|------------|----|
| 🗸 Tab | o 250 mcg | |

DOXYCYCLINE Tab 50 mg

| | Tab | 50 mg | 30 |
|---|-----|--------|----|
| 1 | Tab | 100 mg | 30 |

ERGOMETRINE MALEATE

| ~ | Inj | 500 mc | g per ml, | 1 n | nl ampoule5 |
|---|-----|--------|-----------|-----|-------------|
|---|-----|--------|-----------|-----|-------------|

ERYTHROMYCIN ETHYL SUCCINATE

| ✓ Tab 400 mg | 20 |
|--------------------------------------|--------|
| ✓ Grans for oral liq 200 mg per 5 ml | |
| ✓ Grans for oral liq 400 mg per 5 ml | 200 ml |

ERYTHROMYCIN STEARATE

| Tab 250 mg | 30 |
|------------|----|
|------------|----|

ETHINYLOESTRADIOL WITH DESOGESTREL

| Tab 20 mcg with desogestrel 150 mcg and 7 | |
|---|----|
| inert tab | 84 |
| Tab 30 mcg with desogestrel 150 mcg and 7 | |
| inert tab | |

ETHINYLOESTRADIOL WITH LEVONORGESTREL

| Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tab | 84 |
|--|----|
| ✓ Tab 50 mcg with levonorgestrel 125 mcg and 7 inert tab | 84 |
| Tab 30 mcg with levonorgestrel 150 mcg ✓ Tab 30 mcg with levonorgestrel 150 mcg and | |
| 7 inert tab | 84 |
| ETHINYLOESTRADIOL WITH NORETHISTERONE | |

| | r | Tab 35 | mcg with | norethisterone | 1 | mg63 |
|--|---|--------|----------|----------------|---|------|
|--|---|--------|----------|----------------|---|------|

| Tab 35 mcg with norethisterone 1 mg and 7 | |
|--|----------------------|
| inert tab ✓ Tab 35 mcg with norethisterone 500 mcg ✓ Tab 35 mcg with norethisterone 500 mcg | 63 |
| and 7 inert tab | 84 |
| FLUCLOXACILLIN ✓ Cap 250 mg ✓ Grans for oral liq 25 mg per ml ✓ Grans for oral liq 50 mg per ml ✓ Inj 1 g vial | . 200 ml . 200 ml |
| FLUPENTHIXOL DECANOATE ✓ Inj 20 mg per ml, 1 ml ✓ Inj 20 mg per ml, 2 ml ✓ Inj 100 mg per ml, 1 ml | 5 |
| FLUPHENAZINE DECANOATE ✔ Inj 12.5 mg per 0.5 ml, 0.5 ml ✔ Inj 25 mg per ml, 1 ml ✔ Inj 100 mg per ml, 1 ml | 5 |
| FUROSEMIDE [FRUSEMIDE] ✔ Tab 40 mg ✔ Inj 10 mg per ml, 2 ml ampoule | |
| GLUCAGON HYDROCHLORIDE ✔ Inj 1 mg syringe kit | 5 |
| GLUCOSE [DEXTROSE] ✔ Inj 50%, 10 ml ampoule ✔ Inj 50%, 90 ml bottle | |
| GLYCERYL TRINITRATE ✓ Tab 600 mcg ✓ Oral pump spray, 400 mcg per dose | 50 dose |
| GLYCOPYRRONIUM BROMIDE Inj 200 mcg per ml, 1 ml ampoule | 10 |
| HALOPERIDOL Tab 500 mcg Tab 1.5 mg Tab 5 mg Oral liq 2 mg per ml Inj 5 mg per ml, 1 ml | 30 30 . 200 ml |
| HALOPERIDOL DECANOATE ✔ Inj 50 mg per ml, 1 ml ✔ Inj 100 mg per ml, 1 ml | 5 5 |
| HYDROCORTISONE ✔ Inj 100 mg vial | 5 |
| HYDROXOCOBALAMIN ✔ Inj 1 mg per ml, 1 ml ampoule | |
| contin | |

PRACTITIONER'S SUPPLY ORDERS

| (continued) HYOSCINE N-BUTYLBROMIDE ✔ Inj 20 mg, 1 ml5 | ✔ Inj 15 con ✔ Inj 30 |
|--|---|
| INTRA-UTERINE DEVICE ✓ IUD 29.1 mm length × 23.2 mm width | con NALOXO ✔ Inj 400 |
| IPRATROPIUM BROMIDE ✓ Nebuliser soln, 250 mcg per ml, 1 ml40 ✓ Nebuliser soln, 250 mcg per ml, 2 ml40 | NICOTIN Patch Patch |
| IVERMECTIN ✓ Tab 3 mg – See note on page 68 100 | Patch Lozen Lozen |
| KETONE BLOOD BETA-KETONE ELECTRODES ✓ Test strip | ✓ Gum 2 ✓ Gum 2 ✓ Gum 2 |
| LEVONORGESTREL Tab 30 mcg84 ✔ Tab 1.5 mg5 | ✔ Gum 2 ✔ Gum 4 ✔ Gum 4 ✔ Gum 4 |
| LIDOCAINE [LIGNOCAINE] ✓ Gel 2%, 10 ml urethral syringe – Subsidy by endorsement – See note on page 1255 | NORETH Tab 35 Tab 5 |
| LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE ✓ Inj 1%, 5 ml ampoule | OXYTOC Inj 5 iu Inj 10 OXYTOC Inj 5 iu |
| LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE ✓ Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes – Subsidy by endorsement – See note on page 1265 | per PARACE ✓ Tab 50 ✓ Oral liv ✓ Oral liv |
| LOPERAMIDE HYDROCHLORIDE ✓ Tab 2 mg | PEAK FL |
| MASK FOR SPACER DEVICE ✓ Size 2 – See note on page 19920 | PETHIDI |
| MEDROXYPROGESTERONE ACETATE ✓ Inj 150 mg per ml, 1 ml syringe5 | ✓ Inj 50 dru ✓ Inj 50 |
| METOCLOPRAMIDE HYDROCHLORIDE ✔ Inj 5 mg per ml, 2 ml ampoule5 | dru PHENOX |
| METRONIDAZOLE ✔ Tab 200 mg | ✓ Cap 2 ✓ Cap 5 ✓ Grans |
| MORPHINE SULPHATE ✓ Inj 5 mg per ml, 1 ml ampoule – Only on a controlled drug form | ✓ Grans PHENYT ✓ Inj 50 ✓ Inj 50 |

| ALOXONE HYDROCHLORIDE 'Inj 400 mcg per ml, 1 ml ampoule | Inj 15 mg per ml, 1 ml ampoule – Only on a controlled drug form | |
|--|---|--|
| Patch 7 mg - See note on page 156 28 Patch 14 mg - See note on page 156 28 Patch 21 mg - See note on page 156 28 Lozenge 1 mg - See note on page 156 216 Courage 2 mg - See note on page 156 384 'Gum 2 mg (Classic) - See note on page 156 384 'Gum 2 mg (Kruit) - See note on page 156 384 'Gum 4 mg (Classic) - See note on page 156 384 'Gum 4 mg (Kruit) - See note on page 156 384 'Gum 4 mg (Kruit) - See note on page 156 384 'Gum 4 mg (Kruit) - See note on page 156 384 'Gum 4 mg (Kruit) - See note on page 156 384 'Gum 4 mg (Kruit) - See note on page 156 384 'Gum 4 mg (Mint) - See note on page 156 384 'Gum 4 mg (Kruit) - See note on page 156 384 'Gum 4 mg (Mint) - See note on page 156 384 'Tab 50 mcg 84 'Tab 50 mg 30 XYTOCIN 'Int BERGOMETRINE MALEATE 'Inj 5 iu with ergometrine maleate 500 mcg 9 'per ml, 1 ml 100 ml 'EAK FLOW METER 100 ml 'Oral liq 250 mg per 5 ml 100 ml 'Normal range | | |
| Y Tab 350 mcg | Patch 7 mg – See note on page 156 28 Patch 14 mg – See note on page 156 28 Patch 21 mg – See note on page 156 28 Lozenge 1 mg – See note on page 156 216 Lozenge 2 mg – See note on page 156 216 Gum 2 mg (Classic) – See note on page 156 384 Gum 2 mg (Fruit) – See note on page 156 384 Gum 2 mg (Mint) – See note on page 156 384 Gum 4 mg (Classic) – See note on page 156 384 Gum 4 mg (Fruit) – See note on page 156 384 Gum 4 mg (Fruit) – See note on page 156 384 Gum 4 mg (Fruit) – See note on page 156 384 | |
| Inj 5 iu per ml, 1 ml ampoule | ' Tab 350 mcg | |
| 'Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml | / Inj 5 iu per ml, 1 ml ampoule5 | |
| Yab 500 mg | Inj 5 iu with ergometrine maleate 500 mcg | |
| 'Low range 10 'Normal range 10 ETHIDINE HYDROCHLORIDE 10 'Inj 50 mg per ml, 1 ml – Only on a controlled drug form 5 'Inj 50 mg per ml, 2 ml – Only on a controlled drug form 5 HENOXYMETHYLPENICILLIN (PENICILLIN V) 20 'Cap 250 mg 20 'Grans for oral liq 125 mg per 5 ml 300 ml HENYTOIN SODIUM 300 ml HENYTOIN SODIUM 5 'Inj 50 mg per ml, 2 ml ampoule 5 | ' Tab 500 mg | |
| ^r Inj 50 mg per ml, 1 ml – Only on a controlled drug form | ' Low range 10 | |
| Cap 250 mg 30 Cap 500 mg 20 Grans for oral liq 125 mg per 5 ml 200 ml Grans for oral liq 250 mg per 5 ml 300 ml HENYTOIN SODIUM 10 j 50 mg per ml, 2 ml ampoule 1 j 50 mg per ml, 5 ml ampoule 5 | ' Inj 50 mg per ml, 1 ml – Only on a controlled drug form5 ' Inj 50 mg per ml, 2 ml – Only on a controlled | |
| ^r Inj 50 mg per ml, 2 ml ampoule5 ^r Inj 50 mg per ml, 5 ml ampoule5 | Cap 250 mg | |
| | ' Grans for oral liq 125 mg per 5 ml 200 ml ' Grans for oral liq 250 mg per 5 ml 300 ml | |

(continued)

SAI BUTAMOL WITH IPRATROPIUM BROMIDE ✓ Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule20 SILVER SULPHADIAZINE SODIUM BICARBONATE ✓ Inj 8.4%, 100 ml......5 SODIUM CHLORIDE ✓ Inf 0.9% – See note on page 47 2000 ml ✓ Inj 0.9%, 5 ml – See note on page 47......5 ✓ Inj 0.9%, 10 ml – See note on page 47......5 SPACER DEVICE SPACER DEVICE AUTOCLAVABLE ✓ 230 ml (autoclavable) – Subsidy by endorsement - See note on page 1995 TRIMETHOPRIM VERAPAMIL HYDROCHLORIDE ✓ Inj 2.5 mg per ml, 2 ml ampoule5 WATER ✓ Purified for inj, 5 ml – See note on page 47......5 ✓ Purified for inj, 10 ml – See note on page 47......5 ✓ Purified for inj, 20 ml – See note on page 47......5 ZUCLOPENTHIXOL DECANOATE

| ~ | Ini | 200 | ma | per | ml. | 1 | ml | 5 |
|---|-------|-----|----|-----|-----|---|------|---|
| • | ····y | 200 | mg | por | , | | 1111 | ~ |

Rural Areas for Practitioner's Supply Orders

NORTH ISLAND

Northland DHB Dargaville Hikurangi Kaeo Kaikohe Kaitaia Kawakawa Kerikeri Mangonui Maungaturoto Moerewa Naunauru Paihia Rawene Ruakaka Russell Tutukaka Waipu Whangaroa

Waitemata DHB

Helensville Huapai Kumeu Snells Beach Waimauku Warkworth Wellsford

Auckland DHB

Great Barrier Island Oneroa Ostend

Counties Manukau DHB Tuakau

Waiuku

Waikato DHB

Coromandel Huntly Kawhia Matamata Morrinsville Ngatea Otorohanga Paeroa Pauanui Beach Putaruru Raglan Tairua Taumarunui Te Aroha Te Kauwhata Te Kuiti Tokoroa Waihi Whangamata Whitianga

Bay of Plenty DHB

Edgecumbe Katikati Kawerau Murupara Opotiki Taneatua Te Kaha Waihi Beach Whakatane

Lakes DHB

Mangakino Turangi

Tairawhiti DHB

Ruatoria Te Araroa Te Karaka Te Puia Springs Tikitiki Tokomaru Bay Tolaga Bay

Taranaki DHB

Eltham Inglewood Manaia Oakura Okato Opunake Patea Stratford Waverley

Hawkes Bay DHB

Chatham Islands Waipawa Waipukurau Wairoa Whanganui DHB Bulls Marton Ohakune Raetihi Taihape Waiouru

MidCentral DHB

Dannevirke Foxton Levin Otaki Pahiatua Shannon Woodville

Wairarapa DHB

Carteron Featherston Greytown Martinborough

SOUTH ISLAND

Nelson/Marlborough DHB

Havelock Mapua Motueka Murchison Picton Takaka Wakefield

West Coast DHB

Dobson Greymouth Hokitika Karamea Reefton South Westland Westport Whataroa

Canterbury DHB

Akaroa Amberley Amuri Cheviot Darfield Diamond Harbour Hanmer Springs Kaikoura Leeston Lincoln Methven Oxford Rakaia Rolleston Rotherham Templeton Waikari

South Canterbury DHB

Fairlie Geraldine Pleasant Point Temuka Twizel Waimate

Southern DHB

Alexandra Balclutha Cromwell Gore Kurow I awrence Lumsden Mataura Milton Oamaru Oban Otautau Outram Owaka Palmerston Queenstown Ranfurly Riverton Roxburah Tapanui Te Anau Tokonui Tuatapere Wanaka Winton

SECTION F: PART I

A Community Pharmaceutical identified with a * within the other sections of the Pharmaceutical Schedule:

- a) is exempt from any requirement to dispense in Monthly Lots;
- b) will only be subsidised if it is dispensed in a 90 Day Lot unless it is under the Dispensing Frequency Rule.

A Community Pharmaceutical that is an oral contraceptive and that is identified with a * within the other sections of the Pharmaceutical Schedule:

- a) is exempt from any requirement to dispense in Monthly Lots;
- b) will only be subsidised if it is dispensed in a 180 Day Lot unless it is is under the Dispensing Frequency Rule.

SECTION F: PART II: CERTIFIED EXEMPTIONS AND ACCESS EXEMPTIONS TO MONTHLY DISPENSING

A Community Pharmaceutical, other than a Community Pharmaceutical identified with a * within the other sections of the Pharmaceutical Schedule, may be dispensed in a 90 Day Lot if:

a) the Community Pharmaceutical is identified with a ▲ within the other sections of the Pharmaceutical Schedule and the prescriber/pharmacist has endorsed/annotated the Prescription item(s) on the Prescription to which the exemption applies "certified exemption".

In endorsing/annotating the Prescription items for a certified exemption, the prescriber/pharmacist is certifying that:

- i) the patient wished to have the medicine dispensed in a quantity greater than a Monthly Lot; and
- ii) the patient has been stabilised on the same medicine for a reasonable period of time; and
- iii) the prescriber/pharmacist has reason to believe the patient will continue on the medicine and is compliant.
- b) a patient, who has difficulty getting to and from a pharmacy, signs the back of the Prescription to qualify for an Access Exemption. In signing the Prescription, the patient or his or her nominated representative must also certify which of the following criteria they meet:
 - i) have limited physical mobility;
 - ii) live and work more than 30 minutes from the nearest pharmacy by their normal form of transport;
 - iii) are relocating to another area;
 - iv) are travelling extensively and will be out of town when the repeat prescriptions are due.

SECTION F: PART III: FLEXIBLE AND VARIABLE DISPENSING PERIODS FOR PHARMACY

A Community Pharmaceutical, other than a Community Pharmaceutical identified with a ***** within the other sections of the Pharmaceutical Schedule, may be dispensed in variable dispensing periods under the following conditions:

- a) for stock management where the original pack(s) result in dispensing greater than 30 days supply,
- b) to synchronise a patients medication where multiple medicines result in uneven supply periods, note if dispensing a medicine other than a Pharmaceutical identified with a * please refer to Section F; Part II
- Note the total quantity and dispensing period can not exceed the total quantity and period prescribed on the prescription.

The following Community Pharmaceuticals are identified with a ▲ within the other sections of the Pharmaceutical Schedule and may be dispensed in a 90 Day Lot if endorsed as a certified exemption in accordance with paragraph (a) in Section F Part II above.

ALIMENTARY TRACT AND METABOLISM

INSULIN ASPART

INSULIN ASPART WITH INSULIN ASPART PROTAMINE

INSULIN GLARGINE

INSULIN GLULISINE

INSULIN ISOPHANE

INSULIN ISOPHANE WITH INSULIN NEUTRAL

INSULIN LISPRO

INSULIN LISPRO WITH INSULIN LISPRO PROTAMINE

INSULIN NEUTRAL

HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

DESMOPRESSIN ACETATE Nasal drops 100 mcg Minirin per ml Nasal spray 10 mcg per Desmopressin-PH&T

MUSCULOSKELETAL SYSTEM

PYRIDOSTIGMINE BROMIDE

NERVOUS SYSTEM

dose

AMANTADINE HYDROCHLORIDE

APOMORPHINE HYDROCHLORIDE

ENTACAPONE

GABAPENTIN

GABAPENTIN (NEURONTIN)

LACOSAMIDE

LAMOTRIGINE

LISURIDE HYDROGEN MALEATE

PRAMIPEXOLE HYDROCHLORIDE

ROPINIROLE HYDROCHLORIDE

TOLCAPONE

TOPIRAMATE

VIGABATRIN

CARDIOVASCULAR SYSTEM

| AMIODARONE HYDH | IOCHLORIDE |
|-----------------|-------------|
| Tab 100 mg | Cordarone-X |
| Tab 200 mg | Cordarone-X |

DISOPYRAMIDE PHOSPHATE

FLECAINIDE ACETATE

| lab 50 mg | lambocor |
|------------------------|-------------|
| Tab 100 mg | Tambocor |
| Cap long-acting 100 mg | Tambocor CR |
| Cap long-acting 200 mg | Tambocor CR |

MEXILETINE HYDROCHLORIDE

MINOXIDIL

NICORANDIL

PROPAFENONE HYDROCHLORIDE

SECTION G: SAFETY CAP MEDICINES

Pharmacists are required, under the Code of Ethics of the Pharmacy Council of New Zealand, to endeavour to use safety caps when dispensing any of the medicines listed in Section G in an oral liquid formulation pursuant to a prescription or Practitioner's Supply Order. This includes all proprietary and extemporaneously compounded oral liquid preparations of those pharmaceuticals listed in Section G of the Pharmaceutical Schedule. These medicines will be identified throughout Section B of the Pharmaceutical Schedule with the symbol '‡'.

Exemptions

Oral liquid preparations of the pharmaceuticals listed in Section G of the Pharmaceutical Schedule will be dispensed in a container with a safety cap unless:

- the practitioner has endorsed the Prescription or Practitioner's Supply Order, stating that, the Pharmaceutical is not to be dispensed in a container with a safety cap; or
- the Contractor has annotated the Prescription or Practitioner's Supply Order stating that, because of infirmity of the particular person, the Pharmaceutical to be used by that person should not be dispensed in a container with a safety cap; or
- the Pharmaceutical is packaged in an Original Pack so designed that on the professional judgement of the Contractor, transfer to a container with a safety cap would be inadvisable or a retrograde procedure.

Reimbursment

Pharmacists will be reimbursed according to their agreement. Where an additional fee is paid on safety caps it will be paid on all dispensings of oral liquid preparations for those pharmaceuticals listed in Section G of the Pharmaceutical Schedule unless the practitioner has endorsed or the contractor has annotated the Prescription or Practitioner's Supply Order that a safety cap has not been supplied.

Safety Caps (NZS 5825:1991)

| 20 mm | .Clic-Loc, United Closures & Plastics PLC, England |
|-------|--|
| | Kerr, Cormack Packaging, Sydney, under licence to Kerr USA |
| 24 mm | .Clic-Loc, United Closures & Plastics PLC, England |
| | Clic-Loc, ACI Closures under license to Owens-Illinois |
| | Kerr, Cormack Packaging, Sydney, under licence to Kerr USA |
| 28 mm | .Clic-Loc, United Closures & Plastics PLC, England |
| | Clic-Loc, ACI Closures under license to Owens-Illinois |
| | Kerr, Cormack Packaging, Sydney, under licence to Kerr USA |
| | PDL Squeezlok |
| | PDL FG |
| | |

ALIMENTARY TRACT AND METABOLISM

FERROUS SULPHATE Oral liq 30 mg (6 mg el- Ferodan emental) per 1 ml

CARDIOVASCULAR SYSTEM

AMILORIDE HYDROCHLORIDE Oral liq 1 mg per ml Biomed

CAPTOPRIL Oral lig 5 mg per ml Capoten

CHLOROTHIAZIDE Oral liq 50 mg per ml Biomed

DIGOXIN Oral liq 50 mcg per ml Lanoxin

FUROSEMIDE [FRUSEMIDE] Oral liq 10 mg per ml Lasix

SPIRONOLACTONE Oral liq 5 mg per ml Biomed

HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

LEVOTHYROXINE

| Tab 25 mcg | Synthroid | |
|-------------|-----------|--|
| Tab 50 mcg | Eltroxin | |
| - | Synthroid | |
| Tab 100 mcg | Eltroxin | |
| • | Synthroid | |
| . , | | |

(Extemporaneously compounded oral liquid preparations)

LEVOTHYROXINE (MERCURY PHARMA)

Tab 50 mcg Mercury Pharma Tab 100 mcg Mercury Pharma (Extemporaneously compounded oral liquid preparations)

INFECTIONS - AGENTS FOR SYSTEMIC USE

QUININE SULPHATE Tab 300 mg Q 300 (Extemporaneously compounded oral liquid preparations)

NERVOUS SYSTEM

ALPRAZOLAM Tab 250 mcg Xanax Tab 500 mcg Xanax Tab 1 mg Xanax (Extemporaneously compounded oral liquid preparations)

Tegretol

CARBAMAZEPINE

Oral liq 20 mg per ml

CLOBAZAM Tab 10 mg Frisium (Extemporaneously compounded oral liquid preparations)

CLONAZEPAM Oral drops 2.5 mg per Rivotril ml

DIAZEPAM

 Tab 2 mg
 Arrow-Diazepam

 Tab 5 mg
 Arrow-Diazepam

 (Extemporaneously compounded oral liquid preparations)

ETHOSUXIMIDE Oral liq 250 mg per 5 ml Zarontin

LORAZEPAM Tab 1 mg Ativan Tab 2.5 mg Ativan (Extemporaneously compounded oral liquid preparations)

LORMETAZEPAM

Tab 1 mg Noctamid (Extemporaneously compounded oral liquid preparations)

METHADONE HYDROCHLORIDE

Oral liq 2 mg per mlBiodoneOral liq 5 mg per mlBiodone ForteOral liq 10 mg per mlBiodone Extra Forte

MORPHINE HYDROCHLORIDE

Oral liq 1 mg per ml Oral liq 2 mg per ml Oral liq 5 mg per ml Oral liq 10 mg per ml

RA-Morph RA-Morph RA-Morph RA-Morph

NITRAZEPAM

Tab 5 mg Nitrados (Extemporaneously compounded oral liquid preparations)

OXAZEPAM

Tab 10 mg Ox-Pam Tab 15 mg Ox-Pam (Extemporaneously compounded oral liquid preparations)

OXYCODONE HYDROCHLORIDE Oral lig 5 mg per 5 ml OxyNorm

PARACETAMOL Oral liq 120 mg per 5 ml Paracare Oral liq 250 mg per 5 ml Paracare Double Strength

PHENYTOIN SODIUM Oral liq 30 mg per 5 ml Dilantin

243

SAFETY CAP MEDICINES

SODIUM VALPROATE Oral liq 200 mg per 5 ml

Epilim S/F Liquid Epilim Syrup

TEMAZEPAM Tab 10 mg Normison (Extemporaneously compounded oral liquid preparations)

TRIAZOLAM Tab 125 mcg Hypam Tab 250 mcg Hypam (Extemporaneously compounded oral liquid preparations)

RESPIRATORY SYSTEM AND ALLERGIES

CETIRIZINE HYDROCHLORIDE Oral liq 1 mg per ml Histaclear

CHLORPHENIRAMINE MALEATE Oral liq 2 mg per 5 ml Histafen

DEXTROCHLORPHENIRAMINE MALEATE Oral liq 2 mg per 5 ml Polaramine

PROMETHAZINE HYDROCHLORIDE Oral liq 1 mg per 1 ml Allersoothe SALBUTAMOL Oral liq 400 mcg per ml Ventolin

THEOPHYLLINE Oral liq 80 mg per 15 ml Nuelin

TRIMEPRAZINE TARTRATE Oral liq 30 mg per 5 ml Vallergan Forte

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

CODEINE PHOSPHATE Powder Douglas (Extemporaneously compounded oral liquid preparations)

METHADONE HYDROCHLORIDE Powder AFT (Extemporaneously compounded oral liquid preparations)

PHENOBARBITONE SODIUM Powder MidWest (Extemporaneously compounded oral liquid preparations)

| | Subsidy (Manufacturer's Price) \$ | Fi Subsidis Per | ised (| Brand or Generic Manufacturer |
|---|---|---|--|--|
| Vaccinations | | | | |
| ADULT DIPHTHERIA AND TETANUS VACCINE – [Xpharm] Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid in 0.5 ml | 0.00 | | | <u>r Booster</u> r Booster |
| Any of the following: For vaccination of patients aged 45 and 65 years old; or For vaccination of previously unimmunised or partially imm For revaccination following immunosuppression; or For boosting of patients with tetanus-prone wounds; or For use in testing for primary immunodeficiency disease paediatrician. Note: Please refer to the Immunisation Handbook for appropriate BACILLUS CALMETTE-GUERIN VACCINE – [Xpharm] For infants at increased risk of tuberculosis. Increased risk is living in a house or family with a person with current or paeding in the part of the | es, on the recommend schedule for catch up defined as: ast history of TB; or ithin the last 5 years in in a country with a rate ealth.govt.nz/tuberculo 0.00 | dation of an i p programmes ived in a coun te of TB > or e osis (search fo 1 ● | interna s. htry with equal to for down | al medicine physician or h a rate of TB > or equal o 40 per 100,000 |
| DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE – [Xpharm Funded for any of the following criteria: A single vaccine for pregnant woman between gestationa A course of up to four vaccines is funded for children from immunisation; or An additional four doses (as appropriate) are funded for transplantation or chemotherapy; pre or post splenector severely immunosuppressive regimens. Notes: Tdap is not registered for patients aged less than 10 yea schedule for catch up programmes. Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis toxoid, 8 mcg pertussis filamentous haemagluttinin and 2.5 mcg pertactin in 0.5 ml syringe | al weeks 28 and 38; or a age 7 up to the age o or (re-)immunisation f my; pre- or post solid ars. Please refer to th | of 18 years ind for patients p d organ trans he Immunisat 1 • | post ha splant, i | aematopoietic stem cell renal dialysis and other andbook for appropriate |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|--|---|---|
| DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE – Funded for any of the following: A single dose for children up to the age of 7 who have co A course of four vaccines is funded for catch up program immunisation; or An additional four doses (as appropriate) are funded for or post splenectomy; pre- or post solid organ transplant, or Five doses will be funded for children requiring solid organ Note: Please refer to the Immunisation Handbook for appropriate Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagluttinin, 8 mcg pertactin and 80 D-antigen units poliomyelitis virus in 0.5ml syringe | [Xpharm] mpleted primary imm mes for children (to th (re-)immunisation for renal dialysis and oth in transplantation. schedule for catch up | unisa le age patier ner se | tion; or e of 10 year nts post HSi everely immi rammes. | s) to complete full primary CT, or chemotherapy; pre- |
| DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AN | D HAEMOPHILUS IN | | | |
| Funded for patients meeting any of the following criteria: 1) Up to four doses for children up to and under the age of 1 2) An additional four doses (as appropriate) are funded for are patients post haematopoietic stem cell transplantatic organ transplant, renal dialysis and other severely immur 3) Up to five doses for children up to and under the age of 1 Note: A course of up-to four vaccines is funded for catch up p to complete full primary immunisation. Please refer to the Imm programmes. Inj 30IU diphtheriatoxoid with 40IU tetanustoxoid, 25mcg pertussistoxoid, 25mcg pertussisfilamentoushaemagluttinin, 8 mcgpertactin, 80 D-AgUpoliovirus, 10mcghepatitisB-surfaceantigen in 0.5ml syringe | (re-)immunisation for of on, or chemotherapy; losuppressive regimer 0 receiving solid orga rogrammes for childre hunisation Handbook | childr pre o ns; or n trar en (u | en up to and r post splen hsplantation p to and ur he appropria | ectomy; pre- or post solid nder the age of 10 years) |
| HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm] One dose for patients meeting any of the following: 1) For primary vaccination in children; or 2) An additional dose (as appropriate) is funded for (re-)imn tion, or chemotherapy; pre or post splenectomy; pre- or dialysis and other severely immunosuppressive regimens 3) For use in testing for primary immunodeficiency disease paediatrician. | post solid organ trans s; or | post | haematopo , pre- or pos | ietic stem cell transplanta- st cochlear implants, renal |
| Inj 10 mcg vial with diluent syringe | 0.00 | 1 | ✓ <u>A</u> | ct-HIB |
| HEPATITIS A VACCINE – [Xpharm] Funded for patients meeting any of the following criteria: 1) Two vaccinations for use in transplant patients; or 2) Two vaccinations for use in children with chronic liver diss 3) One dose of vaccine for close contacts of known hepatitis | s A cases. | | | |
| Inj 1440 ELISA units in 1 ml syringe | | 1 | | <u>avrix</u> avrix lupior |
| Inj 720 ELISA units in 0.5 ml syringe | 0.00 | I | ✓ <u>H</u> | avrix Junior |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|--|---------|---------------------|-------------------------------------|
| HEPATITIS B RECOMBINANT VACCINE – [Xpharm] Inj 5 mcg per 0.5 ml vial Funded for patients meeting any of the following criteria: | 0.00 | 1 | <u>✓ H</u> | BvaxPRO |
| for household or sexual contacts of known acute hepatitis for children born to mothers who are hepatitis B surface a for children up to and under the age of 18 years inclusive v require additional vaccination; or for HIV positive patients; or for hepatitis C positive patients; or for patients following non-consensual sexual intercourse; for patients following immunosuppression; or for transplant patients; or | ntigen (HBsAg) posit who are considered r | ive; o | r | ed a positive serology and |
| following needle stick injury. Inj 10 mcg per 1 ml vial Funded for patients meeting any of the following criteria: | 0.00 | 1 | ✓ <u>H</u> | <u>BvaxPRO</u> |
| for household or sexual contacts of known acute hepatitis for children born to mothers who are hepatitis B surface a for children up to and under the age of 18 years inclusive v require additional vaccination; or for HIV positive patients; or for hepatitis C positive patients; or for patients following non-consensual sexual intercourse; for transplant patients; or following needle stick injury. | ntigen (HBsAg) posit who are considered r | ive; o | r | ed a positive serology and |
| Inj 40 mcg per 1 ml vial Funded for any of the following criteria: 1) for dialysis patients; or 2) for liver or kidney transplant patient. | 0.00 | 1 | ✓ <u>H</u> | <u>BvaxPRO</u> |
| HUMAN PAPILLOMAVIRUS (6, 11, 16 AND 18) VACCINE [HPV] Maximum of three doses for patient meeting any of the followin 1) Females aged under 20 years old; or 2) Patients aged under 26 years old with confirmed HIV infer 3) For use in transplant (including stem cell) patients; or 4) An additional dose for patients under 26 years of age positions | ng criteria: ction; or | | | |
| Inj 120 mcg in 0.5 ml syringe | | 1 10 | | ardasil ardasil |

| Subsidy | Fully | Brand or |
|------------------------|------------|----------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | |

INFLUENZA VACCINE - [Xpharm]

- A) is available each year for patients who meet the following criteria, as set by PHARMAC:
 - a) all people 65 years of age and over; or
 - b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or
 - c) children aged four years and under who have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Individual DHBs may fund patients over and above the above criteria. The claiming process for these additional patients should be determined between the DHB and Contractor, or
- D) Stock of the seasonal influenza vaccine is typically available from February until late July with suppliers being required to ensure supply until at least 30 June. Exact start and end dates for each season will be notified each year.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|------------|---------------------|-------------------------------------|
| MEASLES, MUMPS AND RUBELLA VACCINE – [Xpharm] | | | | |
| A maximum of two doses for any patient meeting the following | criteria: | | | |
| 1) For primary vaccination in children; or | | | | |
| 2) For revaccination following immunosuppression; or | | | | |
| For any individual susceptible to measles, mumps or rube | | | | |
| A maximum of three doses for children who have had their | | | | |
| Note: Please refer to the Immunisation Handbook for appropriate s | schedule for catch up | prog | rammes. | |
| Inj 1000 TCID50 measles, 12500 TCID50 mumps and 1000 | | | | |
| TCID50 rubella vial with diluent 0.5 ml vial | 0.00 | 1 | | <u>-M-R II</u> |
| | | 10 | ✓ <u>M</u> | -M-R II |
| MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONGUGATI | E VACCINE – [Xpha | rm] | | |
| Any of the following: | | | | |
| 1) Up to three doses and a booster every five years for patient | nts pre- and post sple | enect | omy and for | patients with functional or |
| anatomic asplenia, HIV, complement deficiency (acquired | or inherited), or pre | or pos | st solid orga | n transplant; or |
| One dose for close contacts of meningococcal cases; or | | | | |
| A maximum of two doses for bone marrow transplant patie | | | | |
| A maximum of two doses for patients following immunosu | | | | |
| Note: children under seven years of age require two doses 8 week | s apart, a booster do | ose th | ree years at | ter the primary series and |
| then five yearly. | | | | |
| *Immunosuppression due to steroid or other immunosuppressive th | herapy must be for a | perio | d of greater | than 28 days. |
| Inj 4 mcg of each meningococcal polysaccharide conjugated | | | | |
| to a total of approximately 48 mcg of diphtheria toxoid | 0.00 | 4 | | |
| carrier per 0.5 ml vial | 0.00 | 1 | | enactra |
| MENINGOCOCCAL C CONGUGATED VACCINE - [Xpharm] | | | | |
| Any of the following: | | | | |
| 1) Up to three doses and a booster every five years for patient | | | | |
| anatomic asplenia, HIV, complement deficiency (acquired | or inherited), or pre | or pos | st solid orga | n transplant; or |
| 2) One dose for close contacts of meningococcal cases; or | | | | |
| A maximum of two doses for bone marrow transplant patie | , | | | |
| A maximum of two doses for patients following immunosup Nature shildren under source of any any instance of any second sec | | | | |
| Note: children under seven years of age require two doses 8 week | s apart, a booster do | ose tri | ree years a | ter the primary series and |
| then five yearly. *Immunosuppression due to steroid or other immunosuppressive to | norany must be for a | norio | d of arootor | than 29 days |
| Inj 10 mcg in 0.5 ml syringe | | perio 1 | | eisvac-C |
| | 0.00 | 10 | | eisvac-C |

| nj 10 mcg in 0.5 ml syringe0.00 | 1 | Neisvac-C |
|---------------------------------|----|-----------|
| | 10 | Neisvac-C |

| | Subsidy (Manufacturer's Price) \$ | Sub: Per | Fully sidised | Brand or Generic Manufacturer |
|--|---|-------------|------------------|-------------------------------------|
| PNEUMOCOCCAL (PCV13) VACCINE – [Xpharm] | | | | |
| Any of the following: | | | | |
| 1) A primary course of four doses for previously unvaccinate | | • | | |
| Up to three doses as appropriate to complete the primary who have received one to three doses of PCV10; or | course of immunisation | n for indiv | iduals u | inder the age of 59 months |
| 3) One dose is funded for high risk children (over the age of four doses of PCV10; or | 17 months and up to t | he age of | 18) wh | o have previously received |
| 4) Up to an additional four doses (as appropriate) are fun | ded for (re-)immunisa | tion of pa | atients | with HIV, for patients post |
| haematopoietic stem cell transplantation, or chemothera solid organ transplant, renal dialysis, complement deficie odeficiency; or | py; pre- or post spler | nectomy; | functio | nal asplenia, pre- or post- |
| For use in testing for primary immunodeficiency disease paediatrician. | es, on the recommend | dation of a | an inter | nal medicine physician or |
| Note: please refer to the Immunisation Handbook for the appropri- | ate schedule for catch | up progr | ammes | |
| Inj 30.8 mcg in 0.5 ml syringe | 0.00 | 1 | | revenar 13 |
| | | 10 | у Р | revenar 13 |
| PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE – [X Either of the following: | pharm] | | | |
| 1) Up to three doses for patients pre- or post-splenectomy of | | enia; or | | |
| 2) Up to two doses are funded for high risk children to the a | ge of 18. | | | |
| Inj 575 mcg in 0.5 ml vial (25 mcg of each 23 pneumococcal serotype) | 0.00 | 1 | • 7 D | neumovax 23 |
| Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 | 0.00 | I | VP | neumovax 23 |
| pneumococcal serotype) | 0.00 | 1 | V P | neumovax 23 |
| (Pneumovax 23 Inj 575 mcg in 0.5 ml vial (25 mcg of each 23 pne | | | | |
| POLIOMYELITIS VACCINE – [Xpharm] | 51 / | | | , |
| Up to three doses for patients meeting either of the following: | | | | |
| For partially vaccinated or previously unvaccinated individ | luals; or | | | |
| 2) For revaccination following immunosuppression. | | | | |
| Note: Please refer to the Immunisation Handbook for appropriate | | | | |
| Inj 80D antigen units in 0.5 ml syringe | 0.00 | 1 | ✓ IF | POL |
| ROTAVIRUS LIVE REASSORTANT ORAL VACCINE - [Xpharm] | | | | |
| Maximum of three doses for patients meeting the following: | | | | |
| 1) first dose to be administered in infants aged under 15 we | | | | |
| 2) no vaccination being administered to children aged 8 mol | nths or over. | | | |
| Oral susp G1, G2, G3, G4, P1(8)11.5 million CCID50 units per 2 ml, tube | 0.00 | 10 | ~ ¤ | otaTeq |
| Por 2 mil, lube | | 10 | • <u>n</u> | |

| Subsidy (Manufacturer's P | Price) | Fully Subsidised | Brand or Generic | |
|------------------------------|--------|---------------------|---------------------|--|
| \$ | Per | ~ | Manufacturer | |

VARICELLA VACCINE [CHICKEN POX VACCINE] - [Xpharm]

Maximum of two doses for any of the following:

- 1) For non-immune patients:
 - a) with chronic liver disease who may in future be candidates for transplantation; or
 - b) with deteriorating renal function before transplantation; or
 - c) prior to solid organ transplant; or
 - d) prior to any elective immunosuppression*.
- 2) For patients at least 2 years after bone marrow transplantation, on advice of their specialist.
- 3) For patients at least 6 months after completion of chemotherapy, on advice of their specialist.
- 4) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist.
- 5) For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella.
- 6) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.
- 7) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.

| * immunosuppression due to steroid or other immunosuppressiv | e therapy must be for | a treatment | t period of greater | than 28 days |
|--|-----------------------|-------------|------------------------------|--------------|
| Inj 2000 PFU vial with diluent | 0.00 | 1 | Varilrix | |

- Symbols -

| - A - A-Scabies 70 Abacavir sulphate 110 Abacavir sulphate with 1amivudine lamivudine 110 Abilify 139 Abiraterone acetate 171 ABM Hydroxocobalamin 37 Acarbose 25 Accarb 25 Accu-Chek Ketur-Test 25 Accu-Chek Performa 26 Accuretic 10 50 Accetadote 204 Acetazolamide 201 Acetadote 204 Acetazolamide 201 Acetic acid with 1, 2- propanediol diacetate and benzethonium 200 Acetic acid with hydroxyquinoline and ricinoleic acid and ricinoleic acid 76 Acetylcysteine 204 Aci-Jel 76 Acidex 20 Acipimox 56 Acitretin 70 Aclasta 119 Actinomycin D 162 Actavis | 3IC110 |
|--|-----------------------------------|
| A-Scabies 70 Abacavir sulphate 110 Abacavir sulphate with 110 Iamivudine 110 Abilify 139 Abiraterone acetate 171 ABM Hydroxocobalamin 37 Accarbose 25 Accarbose 25 Accarbose 25 Accu-Chek Ketur-Test 25 Accu-Chek Performa 26 Accuretic 10 50 Accuretic 20 50 Acetazolamide 201 Acetacote 204 Acetazolamide 201 Acetic acid with 1, 2- propanediol diacetate and benzethonium 200 Acetic acid with hydroxyquinoline and ricinoleic acid 76 Actolovir Infection 104 Sensory 200 Acidex 20 Acipimox 56 Acitretin 70 Aclasta 119 Actin 115 Act-HIB 246 Actavis 141 Actavis 141 Actapol | 50X 3.0 Reservoir32 |
| Abacavir sulphate 110 Abacavir sulphate with 139 Iamivudine 110 Abilify 139 Abiraterone acetate 171 ABM Hydroxocobalamin 37 Acarbose 25 AccurChek Ketur-Test 25 Accu-Chek Performa 26 Accuretic 10 50 Accetadote 204 Acetazolamide 201 Acetacolamide 201 Acetic acid with 1, 2- propanediol diacetate and benzethonium 200 Acetic acid with hydroxyquinoline 76 Acetylcysteine 204 Aci-Jel 76 Acetylcysteine 204 Aci-Jel 76 Acidovir Infection Infection 104 Sensory 200 Acidex 20 Acipimox 56 Acitretin 70 Aclasta 119 Actinomycin D 162 Actayis 141 Actoopin 141 <t< td=""><td></td></t<> | |
| Abacavir sulphate 110 Abacavir sulphate with 139 Iamivudine 110 Abilify 139 Abiraterone acetate 171 ABM Hydroxocobalamin 37 Acarbose 25 AccurChek Ketur-Test 25 Accu-Chek Performa 26 Accuretic 10 50 Accetadote 204 Acetazolamide 201 Acetacolamide 201 Acetic acid with 1, 2- propanediol diacetate and benzethonium 200 Acetic acid with hydroxyquinoline 76 Acetylcysteine 204 Aci-Jel 76 Acetylcysteine 204 Aci-Jel 76 Acidovir Infection Infection 104 Sensory 200 Acidex 20 Acipimox 56 Acitretin 70 Aclasta 119 Actinomycin D 162 Actayis 141 Actoopin 141 <t< td=""><td>A-Scabies70</td></t<> | A-Scabies70 |
| Abacavir sulphate with lamivudine 110 Abilify 139 Abiraterone acetate 171 ABM Hydroxocobalamin 37 Acarbose 25 Accarbose 25 Accarbose 25 AccurChek Ketur-Test 25 Accu-Chek Performa 26 Accuretic 10 50 Accuretic 20 50 Acetadote 204 Acetazolamide 201 Acetic acid with 1, 2- propanediol diacetate and benzethonium 200 Acetic acid with hydroxyquinoline and ricinoleic acid 76 Acetylcysteine 204 Acielovir Infection Infection 104 Sensory 200 Acidex 20 Acipimox 56 Acitretin 70 Aclasta 119 Actin 115 Act-HIB 246 Actavis 141 Actavis 141 Actavis 141 Actavis 141 Actavis 141 | |
| Iamivudine 110 Abilify 139 Abiraterone acetate 171 ABM Hydroxocobalamin 37 Acarbose 25 Accarbose 25 Accarbose 25 Accurechek Ketur-Test 25 Accurechek Performa 26 Accuretic 10 50 Accuretic 20 50 Acetadote 204 Acetazolamide 201 Acetacota with 1, 2- propanediol diacetate and benzethonium 200 Acetic acid with hydroxyquinoline and ricinoleic acid 76 Actylcysteine 204 Aci-Jel .76 Acielovir Infection Infection 104 Sensory 200 Acidex 20 Acipimox 56 Acitretin .70 Aclasta .119 Actin .141 Actavis .141 Actavis .141 Actavis .141 Actavis .141 Actapid Penfill .24 | Abacavir sulphate with |
| Abilify 139 Abiraterone acetate 171 ABM Hydroxocobalamin 37 Acarbose 25 Accarbose 25 Accur-Chek Ketur-Test 25 Accuretic 10 50 Accuretic 20 50 Accuretic 20 50 Accuretic 20 50 Acetazolamide 201 Acetacolamide 201 Acetic acid with 1, 2- propanediol diacetate and benzethonium 200 Acetic acid with hydroxyquinoline and ricinoleic acid 76 Actylcysteine 204 Aci-Jel 76 Aciclovir Infection Infection 104 Sensory 200 Acidex 20 Acipimox 56 Acitretin 70 Aclasta 119 Aclin 115 Actavis 141 Actinomycin D 162 Actavis 141 Actinomycin D 162 Actavis 141 Actrapid Penfill 24 <tr< td=""><td></td></tr<> | |
| Abiraterone acetate 171 ABM Hydroxocobalamin 37 Acarbose 25 Accarb 25 Accur-Chek Ketur-Test 25 Accur-Chek Performa 26 Accuretic 10 50 Accuretic 20 50 Acetadote 204 Acetazolamide 201 Acetacote aid with 1, 2- propanediol diacetate and benzethonium 200 Acetic acid with hydroxyquinoline and ricinoleic acid 76 Acetylcysteine 204 Aci-Jel .76 Aciclovir Infection 104 Sensory 200 Acidex 20 Acidax 104 Sensory 200 Acidex 200 Acidex 200 Acidex .20 Aciperimox .56 Actives .21 Acidex .20 Acitretin .70 Aciasta .119 Actinomycin D .62 Actavis .141 <td></td> | |
| ABM Hydroxocobalamin | |
| Acarbose | |
| Accarb | |
| Accu-Chek Ketur-Test | |
| Accu-Chek Performa | |
| Accuretic 10 | Accu-Chek Performa 26 |
| Accuretic 20 | |
| Acetadote 204 Acetazolamide 201 Acetic acid with 1, 2- propanediol diacetate and benzethonium 200 Acetic acid with hydroxyquinoline and ricinoleic acid 76 Acetylcysteine 204 Aci-Jel 76 Acitylcysteine 204 Aci-Jel 76 Aciclovir Infection Infection 104 Sensory 200 Acidex 20 Acipimox 56 Acitretin 70 Aclasta 119 Aclin 115 Act-HIB 246 Actavis 141 Actinomycin D 162 Actrapid 24 Actrapid 24 Actapid 24 Actapid 24 Actapid 24 Actapid Penfill 24 Acupan 126 Adalimumab 180 Adapalene 62 Adefin XL 54 Adeforir dipivoxil 102 | |
| Acetazolamide 201 Acetic acid with 1, 2- propanediol diacetate and benzethonium 200 Acetic acid with hydroxyquinoline and ricinoleic acid 76 Acetylcysteine 204 Aci-Jel .76 Aciclovir Infection 104 Sensory 200 Acidex .20 Acipimox .56 Acitretin .70 Aclasta .119 Actin .115 Act-HIB .246 Actavis .141 Actinomycin D .162 Actrapid .24 Actrapid .24 Actapid .24 Actavis .141 Actavis .141 Actavis .141 Actapid .24< | |
| Acetic acid with 1, 2- propanediol diacetate and benzethonium 200 Acetic acid with hydroxyquinoline and ricinoleic acid 76 Acetylcysteine 204 Aci-Jel 76 Acitovir 1nfection Infection 104 Sensory 200 Acidex 20 Acipimox 56 Acitretin 70 Aclasta 119 Actin 115 Actayis 141 Actrapid 24 Actrapid Penfill 24 Actrapid Penfill 24 Adalat 10 54 Adalimumab 180 Adapalene 62 Adefin XL 54 Aderovir dipivoxil 102 Aderovir dipivoxil 102 Aderovir dipivoxil 32 ADR Cartridge 1.8 32 ADR Cartridge 3.0 32 Adriamycin 162 Adriamycin 162 | Acetadole |
| diacetate and benzethonium | Acetazolamile |
| benzethonium 200 Acetic acid with hydroxyquinoline and ricinoleic acid .76 Acetylcysteine .204 Aci-Jel .76 Aciclovir | |
| Acetic acid with hydroxyquinoline and ricinoleic acid 76 Acetylcysteine 204 Aci-Jel 76 Aciclovir 104 Infection 104 Sensory 200 Acidex 20 Acipimox 56 Acitretin 70 Aclasta 119 Act-HIB 246 Actavis 141 Actinomycin D 162 Actrapid 24 Actavis 141 Actinomycin D 162 Actavis 141 Actinomycin D 162 Actavis 141 Actionomycin D 162 Actavis 141 Actionomycin D 162 Actavid Penfill 24 Actapid Penfill 24 Actapid NL 54 Adalimumab 180 Adapalene 62 Adefin XL 54 Adefovir dipivoxil 102 Aderovir dipivoxil </td <td>diacetate and</td> | diacetate and |
| and ricinoleic acid 76 Acetylcysteine 204 Aci-Jel 76 Aciclovir 104 Infection 104 Sensory 200 Acidex 20 Acipimox 56 Acitretin 70 Aclasta 119 Act-HIB 246 Actavis 141 Actinomycin D 162 Actrapid 24 Actapid Penfill 24 Acupan 126 Adalimumab 180 Adapalene 62 Adefin XL 54 Adefovir dipivoxil 102 Aderuric 122 ADR Cartridge 1.8 32 Adrenaline 59 Adriamycin 162 | benzetnonium 200 |
| Acetylcysteine 204 Aci-Jel | Acetic acid with hydroxyquinoline |
| Aci-Jel | and ricinoleic acid |
| Aciclovir 104 Infection 104 Sensory 200 Acidex 20 Acipimox 56 Actiretin 70 Aclasta 119 Actin 115 Act-HIB 246 Actavis 141 Actinomycin D 162 Actrapid 24 Actapid Penfill 24 Actapid Penfill 24 Actapalene 62 Adalimumab 180 Adapalene 62 Adefin XL 54 Adefovir dipivoxil 102 Aderouric 122 ADR Cartridge 1.8 32 ADR Cartridge 3.0 32 Adrenaline 59 Adriamycin 162 | |
| Infection 104 Sensory 200 Acidex 20 Acipimox 56 Acitretin 70 Aclasta 119 Aclin 115 Act-HIB 246 Actavis 141 Actavis 141 Actrapid 24 Actrapid Penfill 24 Actupan 126 Adalat 10 54 Adalimumab 180 Adapalene 62 Adefin XL 54 Adefovir dipivoxil 102 ADR Cartridge 1.8 32 ADR Cartridge 3.0 32 Adreanline 59 Adriamycin 162 | |
| Sensory 200 Acidex 20 Acipimox 56 Acitretin 70 Aclasta 119 Aclin 115 Act-HIB 246 Actavis 141 Actinomycin D 262 Actrapid 24 Actrapid Penfill 24 Acupan 126 Adalat 10 54 Adalimumab 180 Adapalene 62 Adefin XL 54 Adefovir dipivoxil 102 Aderoxir dipivoxil 102 ADR Cartridge 1.8 32 ADR Cartridge 3.0 32 Adrenaline 59 Adriamycin 162 ADT Booster 245 | |
| Acidex 20 Acipimox 56 Acitretin 70 Aclasta 119 Aclin 115 Act-HIB 246 Actayis 141 Actrapid 24 Actrapid Penfill 24 Actrapid Penfill 24 Actayan 126 Adalat 10 54 Adalimumab 180 Adapalene 62 Adefin XL 54 Adefovir dipivoxil 102 ADR Cartridge 1.8 32 ADR Cartridge 3.0 32 Adriamycin 162 Adriamycin 162 | |
| Acipimox | |
| Acitretin 70 Aclasta 119 Aclin 115 Act-HIB 246 Actavis 141 Actinomycin D 162 Actrapid 24 Actrapid Penfill 24 Actrapid Penfill 24 Adalat 10 54 Adalaimumab 180 Adapalene 62 Adefin XL 54 Adefovir dipivoxil 102 Aderuric 122 ADR Cartridge 1.8 32 ADR Cartridge 3.0 32 Adriamycin 162 ADT Booster 245 | |
| Aclasta 119 Aclin 115 Act-HIB 246 Actavis 141 Actinomycin D 162 Actrapid 24 Actrapid Penfill 24 Acupan 126 Adalat 10 54 Adalaimumab 180 Adapalene 62 Adefini XL 54 Adefovir dipivoxil 102 Adenuric 122 ADR Cartridge 1.8 32 ADR Cartridge 3.0 32 Adriamycin 162 ADT Booster 245 | |
| Aclin 115 Act-HIB 246 Actavis 141 Actinomycin D 162 Actrapid 24 Actrapid Penfill 24 Acupan 126 Adalat 10 54 Adalaimumab 180 Adapalene 62 Adefin XL 54 Adefovir dipivoxil 102 Aderuric 122 ADR Cartridge 1.8 32 ADR Cartridge 3.0 32 Adriamycin 162 ADT Booster 245 | |
| Act-HIB 246 Actavis 141 Actinomycin D 162 Actrapid 24 Actrapid Penfill 24 Acupan 126 Adalat 10 54 Adalimumab 180 Adapalene 62 Adefin XL 54 Adefovir dipivoxil 102 Aderuric 122 ADR Cartridge 1.8 32 Adrenaline 59 Adriamycin 162 ADT Booster 245 | |
| Actavis 141 Actinomycin D 162 Actrapid 24 Actrapid Penfill 24 Acupan 126 Adalat 10 54 Adalimumab 180 Adapalene 62 Adefin XL 54 Adefovir dipivoxil 102 Adenuric 122 ADR Cartridge 1.8 32 Adrenaline 59 Adriamycin 162 ADT Booster 245 | |
| Actinomycin D 162 Actrapid 24 Actrapid Penfill 24 Acupan 126 Adalat 10 54 Adalimumab 180 Adapalene 62 Adefovir dipivoxil 102 Aderviric 122 ADR Cartridge 1.8 32 Adrenaline 59 Adriamycin 162 ADT Booster 245 | Act-HIB246 |
| Actrapid | |
| Actrapid Penfill 24 Acupan 126 Adalat 10 .54 Adalimumab 180 Adapalene .62 Adefin XL .54 Adefovir dipivoxil .102 Adefovir dipivoxil .102 Adenuric .122 ADR Cartridge 1.8 .32 ADR Cartridge 3.0 .32 Adreanline .59 Adriamycin .162 ADT Booster | Actinomycin D162 |
| Acupan 126 Adalat 10 .54 Adalimumab 180 Adapalene .62 Adefin XL .54 Adefovir dipivoxil 102 Aderovir dipivoxil 102 Adenuric .122 ADR Cartridge 1.8 .32 ADR Cartridge 3.0 .32 Adrenaline .59 Adriamycin .162 ADT Booster .245 | Actrapid24 |
| Adalat 10 | |
| Adalimumab 180 Adapalene 62 Adefin XL 54 Adefovir dipivoxil 102 Adenuric 122 ADR Cartridge 1.8 32 ADR Cartridge 3.0 32 Adrenaline 59 Adriamycin 162 ADT Booster 245 | |
| Adapalene | Adalat 1054 |
| Adefin XL | Adalimumab180 |
| Adefin XL | Adapalene62 |
| Adefovir dipivoxil102Adenuric122ADR Cartridge 1.832ADR Cartridge 3.032Adrenaline59Adriamycin162ADT Booster245 | |
| Adenuric 122 ADR Cartridge 1.8 32 ADR Cartridge 3.0 32 Adrenaline 59 Adriamycin 162 ADT Booster 245 | Adefovir dipivoxil102 |
| ADR Cartridge 1.8 .32 ADR Cartridge 3.0 .32 Adrenaline .59 Adriamycin .162 ADT Booster .245 | |
| ADR Cartridge 3.0 | ADR Cartridge 1.8 |
| Adrenaline59 Adriamycin162 ADT Booster245 | ADR Cartridge 3.0 |
| Adriamycin162 ADT Booster245 | |
| ADT Booster245 | |
| | ADT Booster |
| | Adult diphtheria and tetanus |

| vaccine | . 245 |
|---------------------------------|-------|
| Advantan | |
| Advate | 42 |
| Afinitor | 191 |
| AFT-Pyrazinamide | 102 |
| Agents Affecting the | |
| Renin-Angiotensin System | 49 |
| Agents for Parkinsonism and | |
| Related Disorders | . 124 |
| Agents Used in the Treatment of | |
| Poisonings | . 204 |
| Agrylin | |
| Alanase | 199 |
| Albay | |
| Albendazole | 91 |
| Albey | 193 |
| Albustix | 78 |
| Alendronate sodium | 118 |
| Alendronate sodium with | |
| cholecalciferol | . 118 |
| Alfacalcidol | |
| Alginic acid | 20 |
| Alitraq | 222 |
| Alkeran | 157 |
| Allersoothe | 194 |
| Allopurinol | 121 |
| Alpha Adrenoceptor Blockers | |
| Alpha-Keri Lotion | 67 |
| Alphamox | 93 |
| Alprazolam | 143 |
| Alu-Tab | 20 |
| Aluminium hydroxide | 20 |
| Amantadine hydrochloride | 124 |
| Ambrisentan | 60 |
| Amiloride hydrochloride | 55 |
| Amiloride hydrochloride with | |
| furosemide | 56 |
| Amiloride hydrochloride with | |
| hydrochlorothiazide | 56 |
| Aminophylline | 198 |
| Amiodarone hydrochloride | 51 |
| Amisulpride | 139 |
| Amitriptyline | 130 |
| Amlodipine | |
| Amorolfine | |
| Amoxicillin | 93 |
| Amoxicillin Actavis | 93 |
| Amoxicillin with clavulanic | _ |
| acid | 93 |
| Amphotericin B | |
| Amsacrine | |
| AmsaLyo | 161 |

| Amsidine | 161 |
|--|--|
| Amyl nitrite | 59 |
| Anaesthetics | |
| Anagrelide hydrochloride | |
| Analgesics | 126 |
| Anastrozole | 173 |
| Andriol Testocaps | |
| Androderm | |
| Animas Battery Cap | |
| Animas Cartridge | 32 |
| Animas Vibe | 28 |
| Antabuse | |
| Antacids and Antiflatulants | 20 |
| Anten | |
| Anthelmintics | |
| Antiacne Preparations | 62 |
| Antiallergy Preparations | 102 |
| Antianaemics | |
| Antiandrogen Oral | 40 |
| Contraceptives | 76 |
| Antiarrhythmics | .70 |
| Antibacterials | |
| Antibacterials Topical | 91 |
| Antipacterials Topical | 100 |
| Anticholinergic Agents | 190 |
| Anticholinesterases | |
| Antidepressants | 130 |
| Antidiarrhoeals | |
| Antiepilepsy Drugs | 132 |
| Antifibrinolytics, Haemostatics | |
| and Local Sclerosants | |
| Antifungals | |
| Antifungals Topical | |
| Antihistamines | |
| Antihypotensives | 52 |
| Antimalarials | 100 |
| Antimigraine Preparations | |
| Antinaus | 138 |
| Antinausea and Vertigo | |
| · · · · · · · · · · · · · · · · · · · | |
| Agents | 137 |
| Agents Antiparasitics | 100 |
| Agents Antiparasitics Antipruritic Preparations | 100 64 |
| Agents Antiparasitics Antipruritic Preparations Antipsychotics | 100 64 139 |
| Agents Antiparasitics Antipruritic Preparations Antipsychotics | 100 64 139 |
| Agents Antiparasitics Antipruritic Preparations Antipsychotics Antiretrovirals - Additional | 100 64 139 108 |
| Agents Antiparasitics Antipruritic Preparations Antipsychotics Antiretrovirals - Additional | 100 64 139 108 |
| Agents Antiparasitics Antipruritic Preparations Antipsychotics Antiretrovirals | 100 64 139 108 111 |
| Agents Antiparasitics Antipruritic Preparations Antipsychotics Antiretrovirals Antiretrovirals - Additional Therapies Antirheumatoid Agents Antispasmodics and Other | 100 64 139 108 111 |
| Agents Antiparasitics Antipruritic Preparations Antipsychotics Antiretrovirals - Additional Therapies Antirheumatoid Agents Antispasmodics and Other Agents Altering Gut | 100 64 139 108 111 116 |
| Agents Antiparasitics Antipruritic Preparations Antipsychotics Antiretrovirals - Additional Therapies Antirheumatoid Agents Antispasmodics and Other Agents Altering Gut | 100 64 139 108 111 116 |
| Agents Antiparasitics Antipruritic Preparations Antipsychotics Antiretrovirals Antiretrovirals . Additional Therapies Antirheumatoid Agents Antispasmodics and Other Agents Altering Gut Motility Antithrombotic Agents | 100 64 139 108 111 116 .22 |
| Agents Antiparasitics Antipruritic Preparations Antipsychotics Antiretrovirals - Additional Therapies Antirheumatoid Agents Antispasmodics and Other Agents Altering Gut | 100 64 139 108 111 116 .22 43 |

| | 4.0.0 |
|--------------------------------|--------------------|
| Antitrichomonal Agents | 100 |
| Antituberculotics and | |
| Antileprotics | 101 |
| Antiulcerants | 22 |
| Antivirals | 102 |
| Anxiolytics | 143 |
| Anzatax | |
| Apidra | |
| Apidra SoloStar | |
| | |
| Apo-Allopurinol | |
| Apo-Amiloride | |
| Apo-Amlodipine | |
| Apo-Amoxi | |
| Apo-Azithromycin | |
| Apo-Bromocriptine | |
| Apo-Ciclopirox | 63 |
| Apo- | |
| Cilazapril/Hydrochlorothiazide | 50 |
| Apo-Cimetidine | |
| Apo-Clarithromycin | |
| Alimentary | 22 |
| Infection | |
| | |
| Apo-Clomipramine | |
| Apo-Diclo | |
| Apo-Diltiazem CD | |
| Apo-Doxazosin | |
| Apo-Folic Acid | 41 |
| Apo-Imiquimod Cream 5% | |
| Apo-Megestrol | 172 |
| Apo-Moclobemide | 131 |
| Apo-Nadolol | |
| Apo-Nicotinic Acid | |
| Apo-Oxybutynin | |
| Apo-Perindopril | |
| Apo-Pindolol | |
| | |
| Apo-Prazosin | 49 |
| Apo-Prednisone | |
| Apo-Prednisone S29 | |
| Apo-Primidone | |
| Apo-Propranolol | |
| Apo-Pyridoxine | 37 |
| Apo-Ropinirole | 124 |
| Apo-Selegiline | 124 |
| Apo-Selegiline S29 | 124 |
| Apo-Thiamine | |
| Apo-Timol | |
| Apo-Zopiclone | |
| Apomine | |
| Apomorphine hydrochloride | 124 |
| Aprepitant | |
| Apresoline | |
| Aquasun 30+ | 5 70 |
| | |
| Aqueous cream | ७/ |

| Aratac5 | 1 |
|------------------------|--------|
| Arava11 | 6 |
| Aremed175 | |
| Arimidex173 | |
| Aripiprazole13 | 9 |
| Aristocort6 | 6 |
| Aromasin17 | 3 |
| Arrow - Clopid4 | 3 |
| Arrow Amitriptyline13 | n |
| Arrow-Amitriptyline13 | n |
| Arrow-Bendrofluazide | 6 |
| Arrow-Brimonidine | 0 0 |
| Arrow-Calcium | |
| Arrow-Citalopram13 | |
| Arrow-Citalopram13 | 1 |
| Arrow-Diazepam14 | 4 |
| Arrow-Doxorubicin16 | 2 |
| Arrow-Etidronate11 | |
| Arrow-Fluoxetine13 | |
| Arrow-Gabapentin13 | 3 |
| Arrow-Lamotrigine13 | 5 |
| Arrow-Lisinopril4 | 9 |
| Arrow-Losartan & | |
| Hydrochlorothiazide5 | 1 |
| Arrow-Meloxicam11 | 5 |
| Arrow-Morphine LA12 | 8 |
| Arrow-Norfloxacin11 | 4 |
| Arrow-Ornidazole10 | |
| Arrow-Quinapril 104 | 9 |
| Arrow-Quinapril 204 | 9 |
| Arrow-Quinapril 54 | a |
| Arrow-Roxithromycin | 2 |
| Arrow-Sertraline13 | 1 |
| Arrow-Sertrainle | 7 |
| Arrow-Simva 20mg5 | 7 |
| Arrow Simua 2011g | 7 |
| Arrow-Simva 40mg5 | 7 |
| Arrow-Simva 80mg5 | / |
| Arrow-Sumatriptan13 | |
| Arrow-Timolol20 | 1 |
| Arrow-Tolterodine7 | 8 |
| Arrow-Topiramate13 | 6 |
| Arrow-Tramadol12 | 9 |
| Arrow-Venlafaxine XR13 | 2 |
| Arsenic trioxide16 | |
| Asacol2 | |
| Asamax2 | 1 |
| Ascorbic acid3 | |
| Aspec 30012 | 6 |
| Aspen Adrenaline5 | 9 |
| Aspirin | |
| Blood4 | 3 |
| Nervous12 | |
| Asthalin19 | |
| Atazanavir sulphate11 | 1 |
| | • |
| | |

| Atenolol | 52 |
|---|--|
| Atenolol AFT | 52 |
| ATGAM | 180 |
| Ativan | 144 |
| Atomoxetine | |
| Atorvastatin | 57 |
| Atripla | 110 |
| Atropine sulphate | |
| Cardiovascular | 51 |
| Sensory | 202 |
| Atropt | |
| Atrovent | |
| Augmentin | 93 |
| Auranofin | |
| Ava 20 ED | |
| Ava 30 ED | |
| Avanza | 131 |
| Avelox | |
| Avomine | |
| Avonex | |
| Avonex Pen | |
| Azacitidine | |
| Azamun | |
| Azathioprine | |
| Azithromycin | |
| Azol | 90 |
| | |
| Azopt | 201 |
| AZT | 201 |
| AZT | 201 111 |
| AZT - B - B-D Micro-Fine | 201 111 |
| AZT - B - B-D Micro-Fine B-D Ultra Fine | 201 111 27 27 |
| AZT - B - B-D Micro-Fine B-D Ultra Fine B-D Ultra Fine II | 201 111 27 27 |
| AZT B-D Micro-Fine B-D Ultra Fine B-D Ultra Fine II Bacillus Calmette-Guerin (BCG) | 201 111 27 27 27 |
| AZT | 201 111 27 27 27 |
| AZT B-D Micro-Fine B-D Ultra Fine B-D Ultra Fine II Bacillus Calmette-Guerin (BCG) vaccine Bacillus Calmette-Guerin | 201 111 27 27 27 180 |
| AZT B-D Micro-Fine B-D Ultra Fine B-D Ultra Fine II Bacillus Calmette-Guerin (BCG) vaccine Bacillus Calmette-Guerin vaccine | 201 111 27 27 27 180 245 |
| AZT B-D Micro-Fine B-D Ultra Fine B-D Ultra Fine II Bacillus Calmette-Guerin (BCG) vaccine Bacillus Calmette-Guerin vaccine Baclofen | 201 111 27 27 27 180 245 123 |
| AZT B-D Micro-Fine B-D Ultra Fine B-D Ultra Fine II Bacillus Calmette-Guerin (BCG) vaccine Bacillus Calmette-Guerin vaccine Baclofen Bactroban | 201 111 27 27 27 180 245 123 |
| AZT B-D Micro-Fine B-D Ultra Fine B-D Ultra Fine II Bacillus Calmette-Guerin (BCG) vaccine Bacillus Calmette-Guerin vaccine Baclofen Bactroban Bakels Gluten Free Health Bread | 201 111 27 27 27 180 245 123 63 |
| AZT B-D Micro-Fine B-D Ultra Fine B-D Ultra Fine II Bacillus Calmette-Guerin (BCG) vaccine Bacillus Calmette-Guerin vaccine Baclofen Bactroban Bakels Gluten Free Health Bread Mix | 201 1111 27 27 180 245 123 63 229 |
| AZT | 201 1111 27 27 180 245 123 63 229 |
| AZT | 201 1111 27 27 180 245 123 63 229 103 |
| AZT | 201 111 27 27 180 245 123 63 229 103 67 |
| AZT | 201 111 27 27 180 245 123 63 229 103 67 245 |
| AZT | 201 111 27 27 180 245 123 63 229 103 67 245 194 |
| AZT | 201 111 27 27 180 245 123 63 229 103 67 245 194 194 |
| AZT | 201 111 27 27 180 245 123 63 229 103 67 245 194 194 |
| AZT | 201 111 27 27 27 180 245 123 63 229 103 67 245 194 194 |
| AZT | 201 111 27 27 27 180 245 123 63 229 103 67 245 194 194 |
| AZT | 201 1111 27 27 180 245 123 63 229 103 67 245 194 194 194 |
| AZT | 201 1111 27 27 180 245 123 63 229 103 67 245 194 194 194 199 193 |
| AZT | 201 1111 27 27 180 245 123 63 229 103 67 245 194 194 194 199 193 |

| [Bendrofluazide] | |
|--|---|
| BeneFIX | 42 |
| Benhex | 68 |
| Benzathine benzylpenicillin | 93 |
| Benzbromaron AL 100 | 122 |
| Benzbromarone | |
| Benzoin | |
| Benztrop | |
| Benztropine mesylate | 125 |
| Benzydamine hydrochloride | |
| Benzylpenicillin sodium (penicillin | 00 |
| G) | |
| Beta Adrenoceptor Blockers | |
| Beta Cream | |
| Beta Ointment | |
| | |
| Beta Scalp | /1 |
| Beta-Adrenoceptor Agonists | |
| Betadine | |
| Betadine Skin Prep | |
| Betaferon | |
| Betagan | 201 |
| Betahistine dihydrochloride | 137 |
| Betamethasone dipropionate | 65 |
| Betamethasone dipropionate | |
| with calcipotriol | 70 |
| Betamethasone sodium | |
| phosphate with | |
| | |
| betamethasone acetate | 79 |
| betamethasone acetate6 | 79 5, 71 |
| Betamethasone valerate6 Betamethasone valerate with | 5, 71 |
| Betamethasone valerate6 Betamethasone valerate with | 5, 71 |
| Betamethasone valerate6 | 5, 71 |
| Betamethasone valerate | 5, 71 66 |
| Betamethasone valerate | 5, 71 66 66 |
| Betamethasone valerate | 5, 71 66 66 201 |
| Betamethasone valerate | 5, 71 66 66 201 65 |
| Betamethasone valerate | 5, 71 66 201 65 66 |
| Betamethasone valerate | 5, 71 66 201 65 66 201 |
| Betamethasone valerate | 5, 71 66 201 65 66 201 201 |
| Betamethasone valerate | 5, 71 66 201 65 66 201 201 56 |
| Betamethasone valerate | 5, 71 66 201 65 66 201 201 56 56 |
| Betamethasone valerate | 5, 71 66 201 65 66 201 201 56 56 |
| Betamethasone valerate | 5, 71 66 201 65 66 201 56 56 56 72 |
| Betamethasone valerate | 5, 71 66 201 65 66 201 56 56 56 172 172 |
| Betamethasone valerate | 5, 71 66 201 |
| Betamethasone valerate | 5, 71 66 201 201 201 201 201 56 56 56 172 172 93 157 |
| Betamethasone valerate | 5, 71 66 201 201 201 201 201 56 56 172 172 93 157 23 |
| Betamethasone valerate | 5, 71 66 201 201 201 201 56 56 172 172 172 93 157 23 91 |
| Betamethasone valerate | 5, 71 66 201 65 201 201 56 56 56 56 712 93 93 91 202 |
| Betamethasone valerate | 5, 71 66 201 65 201 201 56 56 56 56 93 93 93 91 202 91 |
| Betamethasone valerate | 5, 71 66 201 65 66 201 56 56 72 72 93 157 93 91 202 91 202 91 |
| Betamethasone valerate | 5, 71 66 201 65 201 201 201 56 56 72 72 93 91 202 27 2127 22 |
| Betamethasone valerate | 5, 71 66 201 65 66 201 56 56 56 56 56 93 93 91 |

| Bisoprolol fumarate52 | |
|-------------------------------------|---|
| BK Lotion67 | |
| Bleomycin sulphate161 | |
| Blood Colony-stimulating Factors | |
| Factors 46 | ; |
| Blood glucose diagnostic test | |
| meter | 5 |
| Blood glucose diagnostic test | |
| strip 26 | 5 |
| Blood glucose test strips (visually | |
| impaired) 27 | 7 |
| Blood ketone diagnostic test | |
| meter 25 | ; |
| BNM | |
| Cardiovascular49 | |
| Genito-Urinary76 | 5 |
| Boceprevir107 | 7 |
| Boceprevir107 Bonjela | 5 |
| Boostrix245 | |
| Bortezomib161 | í |
| Bosentan60 |) |
| Bosvate52 | |
| Bplex | |
| Breath-Alert199 | |
| Brevinor 1/2175 | |
| Brevinor 1/28 | |
| Brevinor 21 | |
| Bricanyl Turbuhaler | |
| Brilinta43 | 2 |
| Brimonidine tartrate | |
| Brimonidine tartrate with timolol | - |
| maleate | , |
| Brinzolamide201 | |
| Brolene200 | |
| Bromocriptine mesylate124 | |
| Brufen SR115 | • |
| | |
| Buccastem138 | 5 |
| Budesonide | |
| Alimentary20 | |
| Respiratory194, 199 |) |
| Budesonide with | |
| eformoterol 195 | |
| Bumetanide55 | ; |
| Buprenorphine with | |
| naloxone | ł |
| Bupropion hydrochloride155 | ; |
| Burinex55 | |
| Buscopan22 |) |
| Buspirone hydrochloride143 | |
| Busulphan157 | 7 |
| Butacort Aqueous199 |) |
| | |

- C -

| • |
|----------------------------|
| Cabergoline89 |
| Cafergot137 |
| Caffeine citrate199 |
| Cal-d-Forte37 |
| Calamine64 |
| Calcipotriol71 |
| Calcitonin79 |
| Calcitriol37 |
| Calcitriol-AFT37 |
| Calcium carbonate20, 38 |
| Calcium Channel Blockers54 |
| Calcium Disodium |
| Versenate |
| Calcium folinate159 |
| Calcium Folinate Ebewe159 |
| Calcium gluconate |
| Calcium Homeostasis79 |
| Calcium polystyrene |
| sulphonate |
| Calcium Resonium47 |
| Calogen217 |
| Calsource |
| Camptosar160 |
| Candesartan cilexetil50 |
| Candestar50 |
| Canesten63 |
| Capecitabine159 |
| Capecitabine Winthrop159 |
| Capoten49 |
| Capsaicin |
| Musculoskeletal System116 |
| Nervous126 |
| Captopril49 |
| Carafate23 |
| Carbaccord157 |
| Carbamazepine133 |
| Carbimazole |
| Carbomer203 |
| Carboplatin157 |
| Carboplatin Ebewe157 |
| Carbosorb-X204 |
| Cardinol LA53 |
| Cardizem CD54 |
| CareSens |
| CareSens II |
| CareSens N |
| CareSens N POP26 |
| Carmustine157 |
| Carvedilol52 |
| Catapres55 |
| Catapres-TTS-155 |
| |

| Catapres-TTS-255 |
|------------------------------|
| Catapres-TTS-355 |
| CeeNU |
| Cefaclor monohydrate |
| |
| Cefalexin91 |
| Cefalexin Sandoz91 |
| Cefazolin91 |
| Ceftriaxone91 |
| Ceftriaxone-AFT91 |
| Cefuroxime axetil92 |
| Celestone Chronodose |
| |
| Celiprolol |
| Cellcept174 |
| Celol |
| Centrally-Acting Agents55 |
| Cephalexin ABM91 |
| Cerezyme35 |
| Cetirizine hydrochloride193 |
| Cetomacrogol67 |
| Cetomacrogol with glycerol67 |
| Champix156 |
| Charcoal204 |
| |
| Chemotherapeutic Agents |
| Chicken pox vaccine250 |
| Chlorafast200 |
| Chlorambucil157 |
| Chloramphenicol200 |
| Chlorhexidine gluconate |
| Alimentary |
| Dermatological66 |
| Chloroform211 |
| Chlorothiazide |
| Chlorpheniramine maleate |
| • |
| Chlorpromazine |
| hydrochloride 139 |
| Chlorsig200 |
| Chlortalidone |
| [Chlorthalidone]56 |
| Chlorthalidone56 |
| Chlorvescent48 |
| Choice TT380 Short73 |
| Choice TT380 Standard73 |
| Cholecalciferol |
| |
| Cholestyramine |
| Choline salicylate with |
| cetalkonium chloride |
| Cholvastin57 |
| Ciclopirox olamine63 |
| Ciclosporin191 |
| Cilazapril49 |
| Cilazapril with |
| |
| hydrochlorothiazide |
| hydrochlorothiazide |

| Cilicaine VK94 | |
|-----------------------------------|---|
| Ciloxan200 | |
| Cimetidine22 | 2 |
| Cipflox95 | j |
| Ciprofloxacin | |
| Infection95 | 5 |
| Sensory200 |) |
| Cisplatin157 | |
| Cisplatin Ebewe157 | , |
| Citalopram hydrobromide137 | |
| | |
| Cladribine159 | , |
| Clarithromycin | |
| Alimentary22 | |
| Infection92 | |
| Clexane44 | |
| Climara 10082 | 2 |
| Climara 5082 | |
| Clindamycin95 | j |
| Clindamycin ABM95 | 5 |
| Clobazam133 | 3 |
| Clobetasol BNM65 | |
| Clobetasol propionate | |
| Clobetasone butyrate | |
| Clofazimine101 | |
| | |
| Clomazol | |
| Dermatological63 | |
| Genito-Urinary76 | |
| Clomiphene citrate90 |) |
| Clomipramine hydrochloride130 | |
| Clonazepam132-133, 144 | |
| Clonidine55 | ; |
| Clonidine BNM55 | |
| Clonidine hydrochloride55 | 5 |
| Clopidogrel43 | |
| Clopine |) |
| Clopixol141, 143 | 2 |
| Clotrimazole | ʻ |
| Dermatological63 | , |
| Genito-Urinary | ; |
| Clozapine140 | |
| | |
| Clozaril | |
| Co-Renitec50 |) |
| Co-trimoxazole95 | |
| Coal tar71 | |
| Coal tar with allantoin, menthol, | |
| phenol and sulphur71 | |
| Coal tar with salicylic acid and | |
| sulphur71 | |
| Coco-Scalp71 | |
| Codeine phosphate | |
| Extemporaneous | |
| Nervous127 | |
| Cogentin125 | 5 |
| 00g01011120 | · |
| | |

| Colaspase [L-asparaginase] | 162 |
|--|---|
| Colchicine | |
| Colestid | |
| Colestipol hydrochloride | |
| Colgout | |
| Colifoam | |
| Colistin sulphomethate | |
| Colistin-Link | 95 |
| Collodion flexible | |
| Colofac | |
| Coloxyl | |
| Combigan | |
| Comfort | |
| Comfort Short | 30 |
| Compound electrolytes | 47 |
| Compound | |
| hydroxybenzoate | 211 |
| Concerta | |
| Condoms | |
| Condyline | |
| Contact-D | 29 |
| Contraceptives - Hormonal | 74 |
| Contraceptives - | |
| Non-hormonal | |
| Copaxone | |
| Cordarone-X | 51 |
| Cartiagataraida and Dalatad | |
| Corticosteroids and Related | |
| Agents for Systemic Use | . 79 |
| Agents for Systemic Use | 65 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen | 65 162 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt | 65 162 202 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin | 65 162 202 46 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Cournadin Creon 10000 | 65 162 202 46 33 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Cournadin Creon 10000 Creon 25000 | 65 162 202 46 33 33 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin Creon 10000 Creon 25000 Crixivan | 65 162 202 46 33 33 111 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Cournadin Creon 10000 Creon 25000 Crixivan Crotamiton | 65 162 202 46 33 33 111 64 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Cournadin Creon 10000 Creon 25000 Crixivan Crotamiton Crystaderm | 65 162 202 46 33 33 111 64 63 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Cournadin Creon 10000 Creon 25000 Crixivan Crotamiton Crystaderm Curam | 65 162 202 46 33 33 111 64 63 93 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin Creon 10000 Creon 25000 Crixivan Crotamiton Crystaderm Curam Curam Duo | 65 162 202 46 33 33 111 64 93 93 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin Creon 10000 Creon 25000 Crixivan Crotamiton Crystaderm Curam Duo Cvite | 65 162 202 46 33 33 111 64 63 93 93 37 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin Creon 10000 Creon 25000 Crixivan Crotamiton Crystaderm Curam Duo Curam Duo Cvite Cyclizine hydrochloride | 65 162 202 46 33 33 111 64 93 93 93 37 138 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin Creon 10000 Creon 25000 Crixivan Crotamiton Crystaderm Curam Duo Curam Duo Cvite Cyclizine hydrochloride Cyclizine lactate | 65 162 202 46 33 33 111 64 93 93 37 138 138 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin Creon 10000 Creon 25000 Crixivan Crotamiton Crystaderm Curam Duo Curam Duo Cvite Cyclizine hydrochloride Cyclizine lactate Cyclogyl | 65 162 202 46 33 33 111 64 93 93 37 138 138 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin Creon 10000 Creon 25000 Crixivan Crotamiton Crystaderm Curam Duo Cvite Cyclizine hydrochloride Cyclizine hydrochloride Cyclizine lactate Cyclogyl Cyclopentolate | 65 162 202 46 33 33 111 64 93 93 37 138 138 202 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin Creon 10000 Creon 25000 Crixivan Crotamiton Crystaderm Curam Duo Cystaderm Curam Duo Cyclizine hydrochloride Cyclizine lactate Cyclogyl Cyclopentolate hydrochloride | 65 162 202 46 33 33 111 64 93 93 37 138 138 202 202 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin Creon 10000 Creon 25000 Crixivan Crotamiton Crotamiton Crystaderm Curam Duo Curam Duo Cyclizine hydrochloride Cyclizine lactate Cyclogyl Cyclopentolate hydrochloride Cyclophosphamide | 65 162 202 46 33 33 111 64 93 93 37 138 138 202 202 157 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin Creon 10000 Creon 25000 Crixivan Crotamiton Crotamiton Crystaderm Curam Duo Curam Duo Cyclizine hydrochloride Cyclizine lactate Cyclogyl Cyclopentolate hydrochloride Cyclophosphamide Cycloserine | 65 162 202 46 33 33 111 64 63 93 93 37 138 138 202 202 157 101 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin Creon 10000 Creon 25000 Crixivan Crotamiton Crystaderm Curam Duo Cystaderm Curam Duo Cyclizine hydrochloride Cyclizine hydrochloride Cyclizine lactate Cyclogyl Cyclopentolate hydrochloride Cyclophosphamide Cycloserine Cyklokapron | 65 162 202 46 33 33 111 64 93 93 37 138 138 202 202 157 101 42 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin Creon 10000 Creon 25000 Crixivan Crotamiton Crystaderm Curam Quara Curam Duo Cyclizine hydrochloride Cyclizine hydrochloride Cyclizine lactate Cyclogyl Cyclopentolate hydrochloride Cyclopentolate hydrochloride Cycloserine Cycloserine Cyklokapron Cyproterone acetate | 65 162 202 46 33 33 111 64 93 93 37 138 138 202 202 157 101 42 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin Creon 10000 Creon 25000 Crixivan Crotamiton Crystaderm Curam Duo Curam Duo Curam Duo Cyclizine hydrochloride Cyclizine hydrochloride Cyclizine lactate Cyclogyl Cyclopentolate hydrochloride Cyclopentolate hydrochloride Cycloperine Cycloserine Cyklokapron Cyproterone acetate Cyproterone acetate with | 65 162 202 46 33 33 111 64 93 93 37 138 202 202 157 101 42 81 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin Creon 10000 Creon 25000 Crixivan Crotamiton Crystaderm Curam Duo Curam Duo Curam Duo Cyclizine hydrochloride Cyclizine hydrochloride Cyclogyl Cyclopentolate hydrochloride Cyclophosphamide Cyclophosphamide Cycloserine Cyklokapron Cyptorerone acetate Cyproterone acetate with ethinyloestradiol | 65 162 202 46 33 33 111 64 93 93 37 138 138 202 202 157 101 42 81 |
| Agents for Systemic Use Corticosteroids Topical Cosmegen Cosopt Coumadin Creon 10000 Creon 25000 Crixivan Crotamiton Crystaderm Curam Duo Curam Duo Curam Duo Cyclizine hydrochloride Cyclizine hydrochloride Cyclizine lactate Cyclogyl Cyclopentolate hydrochloride Cyclopentolate hydrochloride Cycloperine Cycloserine Cyklokapron Cyproterone acetate Cyproterone acetate with | 65 162 202 46 33 33 111 64 93 93 37 138 202 202 157 101 42 81 76 159 |

| Cytoxan157 | ' |
|------------------------------|---|
| - D - | |
| D-Penamine116 | |
| d4T110 | |
| Dabigatran45 | |
| Dacarbazine | |
| Dactinomycin [Actinomycin | |
| D] 162 | • |
| Daivobet | |
| Daivonex71 | |
| Daktarin64 | |
| Dalacin C95 | |
| Dalteparin sodium44 | |
| Danazol | 1 |
| Dantrium | |
| Dantrolene | |
| Daonil25 | |
| Dapa-Tabs56 | |
| Dapsone101 | |
| Daraprim97 | , |
| Darunavir111 | |
| Dasatinib166 | 1 |
| Daunorubicin162 | |
| DBL Acetylcysteine204 | |
| DBL Aminophylline198 | |
| DBL Bleomycin Sulfate161 | |
| DBL Carboplatin157 | , |
| DBL Docetaxel162 | |
| DBL Doxorubicin162 | |
| DBL Doxorubicin S29162 | |
| DDI Entruckista | |
| Hydrochloride | |
| DBL Ergometrine76 | i |
| DBL Flucloxacillin94 | |
| DBL Gemcitabine160 | 1 |
| DBL Leucovorin Calcium159 | |
| DBL Morphine Sulphate128 | |
| DBL Pethidine | |
| Hydrochloride | |
| DBL Tobramycin97 | |
| DDI110 | |
| De Nol23 | |
| De-Worm91 | |
| Decozol | |
| Deferasirox204 | |
| Deferiprone204 | |
| Deoxycoformycin164 | |
| Depo-Medrol80 | ١ |
| Depo-Medrol with Lidocaine80 | ١ |
| Depo-Provera75 | |
| Depo-Testosterone81 | |
| Deprim95 | 1 |

| Dermol | 65 71 |
|---|-----------|
| Desferrioxamine mesylate | 205 |
| Desmopressin acetate | 205 QQ |
| | 00 |
| Desmopressin-PH&T Detection of Substances in | 09 |
| | 70 |
| Urine | |
| Dexamethasone | |
| Hormone | |
| Sensory | |
| Dexamethasone phosphate | |
| Dexamethasone with framyceti | |
| and gramicidin | |
| Dexamethasone with neomycir | ו |
| sulphate and polymyxin B | |
| sulphate | |
| Dexamethasone-hameln | |
| Dexamfetamine sulfate | 151 |
| Dextrochlorpheniramine | |
| maleate | 193 |
| Dextrose | 47 |
| Dextrose with electrolytes | 48 |
| DHC Continus | |
| Diabetes | |
| Diabetes Management | 20 |
| Diacomit | 136 |
| Diamide Relief | 20 |
| Diamox | |
| Diaphragm | |
| Diasip | |
| | |
| Diason RTH | |
| Diazepam1 | |
| Diazoxide | |
| Dicarz | |
| Diclax SR | 115 |
| Diclofenac sodium | |
| Musculoskeletal System | 115 |
| Sensory | 201 |
| Didanosine [DDI] | |
| Differin | |
| Difflam | |
| Diflucan | |
| Diflucan S29 | |
| Diflucortolone valerate | 65 |
| Digestives Including | |
| Enzymes | 33 |
| Digoxin | 51 |
| Dihydrocodeine tartrate | 127 |
| Dilantin | 135 |
| Dilantin Infatab | |
| Dilatrend | 52 |
| Diltiazem hydrochloride | |
| Dilzem | 54 |
| Dimethicone | |
| | |

| Dipentum | 21 |
|---|-----|
| Diphtheria, tetanus and pertussis | |
| vaccine Diphtheria, tetanus, pertussis | 245 |
| and polio vaccine | 246 |
| Diphtheria, tetanus, pertussis, | 240 |
| polio, hepatitis B and | |
| haemophilus influenzae type E | 3 |
| vaccine | |
| Diprosone | 65 |
| Diprosone OV | 65 |
| Dipyridamole | |
| Disinfecting and Cleansing | |
| Agents | 66 |
| Disopyramide phosphate | 51 |
| Disulfiram | |
| Diuretics | |
| Diurin 40 Docetaxel | |
| Docetaxel Sandoz | |
| Docusate sodium | |
| Docusate sodium with | 0+ |
| sennosides | |
| Domperidone | |
| Donepezil hydrochloride | |
| Donepezil-Rex | 154 |
| Dopergin | 124 |
| Dopress | 130 |
| Dornase alfa | 198 |
| Dorzolamide hydrochloride | 202 |
| Dorzolamide hydrochloride with | |
| timolol maleate | |
| Dostinex | 89 |
| Dothiepin hydrochloride | |
| Doxazosin Doxepin hydrochloride | 120 |
| Doxine | |
| Doxorubicin | |
| Doxorubicin Ebewe | 162 |
| Doxy-50 | |
| Doxycycline | 94 |
| DP Fusidic Acid Cream | 63 |
| DP Lotion | |
| DP Lotn HC | |
| DP-Anastrozole | |
| Dr Reddy's Omeprazole Dr Reddy's Ondansetron | 23 |
| Dr Heddy's Ondansetron | 138 |
| Dr Reddy's Terbinafine | 99 |
| Drugs Affecting Bone Metabolism | 110 |
| Dulcolax | |
| Ducciax Duocal Super Soluble | 55 |
| Powder | 216 |
| | |

| Duolin196 |
|----------------------------------|
| Duolin HFA196 |
| |
| Durex Confidence73 |
| Durex Extra Safe73 |
| Durex Select Flavours73 |
| Duride59 |
| Dynacirc-SRO |
| • |
| - E - |
| E-Mycin92 |
| Ear Preparations200 |
| |
| Ear/Eye Preparations200 |
| Easiphen Liquid231 |
| EasyCheck |
| Econazole nitrate64 |
| Efavirenz110 |
| Efavirenz with emtricitabine and |
| |
| tenofovir disoproxil |
| fumarate110 |
| Efexor XR132 |
| Effient43 |
| Eformoterol fumarate |
| |
| Efudix |
| Egopsoryl TA71 |
| Elecare232 |
| Elecare LCP232 |
| Elemental 028 Extra222 |
| Eligard |
| 5 |
| Elocon |
| Eloxatin158 |
| Eltrombopag41 |
| Eltroxin |
| Emend Tri-Pack137 |
| EMLA126 |
| |
| Emtricitabine110 |
| Emtricitabine with tenofovir |
| disoproxil fumarate 110 |
| Emtriva110 |
| Emulsifying ointment67 |
| Enalapril maleate49 |
| |
| Enalapril maleate with |
| hydrochlorothiazide |
| Enbrel174 |
| Endocrine Therapy171 |
| Endoxan157 |
| |
| Enerlyte |
| Enfuvirtide111 |
| Enoxaparin sodium44 |
| Ensure |
| Ensure Plus227 |
| Ensure Plus HN226 |
| Ensure Plus RTH226 |
| |
| Entacapone124 |
| |

| Entapone | 124 |
|--|-------|
| Entecavir | 103 |
| Entocort CIR | 20 |
| Epilim | |
| Epilim Crushable | |
| Epilim IV | |
| Epilim S/F Liquid | 135 |
| Epilim Syrup | 135 |
| Epirubicin | |
| Epirubicin Ebewe | |
| Epoetin alfa [Erythropoietin | 102 |
| | 44 |
| alfa] | |
| Eprex | |
| Eptacog alfa [Recombinant factor | |
| VIIa] | |
| ERA | |
| Ergometrine maleate | 76 |
| Ergotamine tartrate with | |
| caffeine | . 137 |
| Erlotinib | |
| Erythrocin IV | 92 |
| Erythromycin ethyl succinate | 92 |
| Erythromycin lactobionate | 92 |
| Erythromycin stearate | 92 |
| Erythromycin stearate Erythropoietin alfa | 40 |
| Escitalopram | 131 |
| Eskazole | |
| Estradot | |
| Estrofem | |
| Etanercept | |
| Ethambutol hydrochloride | 101 |
| Ethics Assiste | . 101 |
| Ethics Aspirin | 120 |
| Ethics Aspirin EC | 43 |
| Ethics Enalapril | 49 |
| Ethinyloestradiol | 83 |
| Ethinyloestradiol with desogestrel | |
| desogestrel | 74 |
| Ethinyloestradiol with | |
| levonorgestrel | 74 |
| Ethinyloestradiol with norethisterone | |
| norethisterone | 75 |
| Ethosuximide | |
| Etidronate disodium | 118 |
| Etopophos | 163 |
| Etoposide | 163 |
| Etoposide phosphate | |
| Etravirine | |
| Eumovate | |
| Everolimus | |
| Evista | |
| Exelon | |
| Exemestane | |
| Exjade | |
| слушие | 204 |

| Extemporaneously Compounded |
|-----------------------------------|
| Preparations and |
| Galenicals |
| Eve Preparations |
| EZ-fit Paediatric Mask199 |
| Ezemibe57 |
| Ezetimibe57 |
| Ezetimibe with simvastatin58 |
| Ezetrol57 |
| .F. |
| Factor eight inhibitors bypassing |
| agent |
| Febuxostat122 |
| Feed Thickener Karicare |
| Aptamil 229 |
| FEIBA42 |
| Felodipine54 |
| Fenpaed115 |
| Fentanyl |
| Fentanyl Sandoz127 |
| Ferodan |
| Ferriprox204 |
| Ferro-F-Tabs |
| Ferro-tab |
| Ferrograd |
| Ferrograd F |
| Ferrous fumarate |
| Ferrous fumarate with folic |
| acid |
| Ferrous sulphate |
| Ferrous sulphate with folic |
| acid |
| Ferrum H |
| Fexofenadine hydrochloride193 |
| Fibro-vein42 |
| Filgrastim46 |
| Finasteride77 |
| Fingolimod144 |
| Finpro77 |
| Flagyl100 |
| FlagyI-S100 |
| Flamazine63 |
| Flecainide acetate51 |
| Fleet Phosphate Enema35 |
| Flixonase Hayfever & |
| Allergy 199 |
| Flixotide194 |
| Flixotide Accuhaler194 |
| Florinef79 |
| Fluanxol142 |
| Fluarix248 |
| Flucloxacillin94 |
| Flucloxin94 |

| Flucon | |
|---|--|
| Fluconazole | |
| Fludara | |
| Fludara Oral | 159 |
| Fludarabine Ebewe | 159 |
| Fludarabine phosphate | 159 |
| Fludrocortisone acetate | 79 |
| Fluids and Electrolytes | 47 |
| Flumetasone pivalate | |
| Fluocortolone caproate with | |
| fluocortolone pivalate and | |
| cinchocaine | 21 |
| Fluorometholone | |
| Fluorouracil | |
| Fluorouracil Ebewe | 150 |
| Fluorouracil sodium | |
| Fluoxetine hydrochloride | |
| Flupenthixol decanoate | |
| Fluphenazine decanoate | 142 |
| | |
| Flutamide | |
| Flutamin | |
| Flutamin S29 | |
| Fluticasone | |
| Fluticasone propionate | 199 |
| Fluticasone with salmeterol | |
| FML | |
| Foban | 63 |
| | |
| Folic acid | |
| Food Thickeners | |
| Food Thickeners Foods And Supplements For | |
| Food Thickeners Foods And Supplements For Inborn Errors Of | 229 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism | 229 230 |
| Food Thickeners Foods And Supplements For Inborn Errors Of | 229 230 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Forteo | 229 230 194 119 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Forteo Fortimel Regular | 229 230 194 119 219 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Forteo Fortimel Regular Fortini | 229 230 194 119 219 220 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Forteo Fortimel Regular Fortini Fortini Multi Fibre | 229 230 194 119 219 220 221 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Forteo Fortimel Regular Fortini Multi Fibre Fortini Multi Fibre | 229 230 194 119 219 220 221 6, 227 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Forteo Fortimel Regular Fortini Multi Fibre Fortisip | 229 230 194 119 219 221 6, 227 227 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Forteo Fortimel Regular Fortini Multi Fibre Fortisip | 229 230 194 119 219 220 221 5,227 227 218 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Forteo Fortimel Regular Fortini Multi Fibre Fortisip | 229 230 194 119 219 220 221 5,227 227 218 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Forteo Fortimel Regular Fortini Multi Fibre Fortisip | 229 230 194 219 220 221 6,227 227 118 118 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Fortimel Regular Fortimel Regular Fortini Multi Fibre Fortisip Fortisip Multi Fibre Fosamax Plus Fragmin | 229 230 194 219 220 221 6,227 227 118 118 44 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Fortimel Regular Fortimel Regular Fortimi Multi Fibre Fortisip Fortisip Multi Fibre Fosamax Plus Fosamax Plus Fragmin Framycetin sulphate | 229 230 194 119 219 220 221 6,227 227 118 118 44 200 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Fortinel Regular Fortini Multi Fibre Fortisip Fortisip Multi Fibre Fortisip Multi Fibre Fosamax Plus Fragmin Framycetin sulphate Freestyle Optium | 229 230 194 119 219 221 6,227 227 227 118 118 44 200 25,26 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Fortinel Regular Fortini Multi Fibre Fortisip Multi Fibre Fortisip Multi Fibre Fosamax Plus Fragmin Fraestyle Optium Freestyle Optium Ketone | 229 230 194 119 219 221 6,227 227 118 118 118 44 200 25,26 25 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Fortinel Regular Fortinel Regular Fortini Multi Fibre Fortisip Multi Fibre Fortisip Multi Fibre Fosamax Plus Fragmin Fraestyle Optium Freestyle Optium Ketone Frisium | 229 230 194 219 220 221 227 227 118 118 118 210 25, 26 25 25 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Forteo Fortimel Regular Fortini Multi Fibre Fortisip Fortisip Multi Fibre Fortisip Multi Fibre Fosamax Fosamax Plus Fragmin Fraestyle Optium Freestyle Optium Ketone Frisium Frumil | 229 230 194 219 220 221 227 118 118 44 200 25, 26 25 25 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Forteo Fortimel Regular Fortimi Multi Fibre Fortisip Fortisip Multi Fibre Fortisip Multi Fibre Fosamax Fosamax Plus Fragmin Fragmin Freestyle Optium Ketone Frisium Frumil Frusemide | 229 230 194 119 219 220 221 227 118 118 200 25, 26 25 26 56 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Forteo Fortimel Regular Fortimi Multi Fibre Fortisip Fortisip Multi Fibre Fortisip Multi Fibre Fosamax Plus Fragmin Fragmin Freestyle Optium Freestyle Optium Ketone Frisium Frumil Frusemide Frusemide | 229 230 194 219 221 221 227 118 227 227 227 225 25 25 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Forteo Fortimel Regular Fortimel Regular Fortimi Multi Fibre Fortisip Multi Fibre Fortisip Multi Fibre Fosamax Plus Frosamax Plus Fragmin Fraestyle Optium Ketone Frisium Frumil Frusemide Frusemide Frusemide | 229 230 194 119 219 221 227 118 20 25, 26 25 25 55 55 |
| Food Thickeners | 229 230 194 119 219 221 3, 227 118 221 225 255 255 |
| Food Thickeners Foods And Supplements For Inborn Errors Of Metabolism Foradil Forteo Fortimel Regular Fortimel Regular Fortimi Multi Fibre Fortisip Multi Fibre Fortisip Multi Fibre Fosamax Plus Frosamax Plus Fragmin Fraestyle Optium Ketone Frisium Frumil Frusemide Frusemide Frusemide | 229 230 194 219 221 221 227 227 118 118 44 200 25, 26 255 555 555 555 955 955 955 |

| Furosemide [Frusemide] Fusidic acid | 55 |
|--|-----|
| Dermatological | ~~ |
| | |
| Infection | |
| Sensory | 200 |
| Fuzeon | 111 |
| - G - | |
| Gabapentin | 133 |
| Gabapentin (Neurontin) | 134 |
| Gamma benzene | |
| hexachloride | 68 |
| Ganciclovir | |
| | |
| Gardasil | |
| Gastrosoothe Gaviscon Double Strength | 22 |
| | |
| Gaviscon Infant | |
| Gefitinib | |
| Gemcitabine Ebewe | 160 |
| Gemcitabine hydrochloride | |
| Gemfibrozil | 56 |
| Gemzar | 160 |
| Genoptic | |
| Genox | |
| Gentamicin sulphate | |
| Infection | 96 |
| Infection Sensory | 200 |
| Gilenya | 1// |
| Ginet | 144 |
| | |
| Glatiramer acetate | |
| Glibenclamide | |
| Gliclazide | 25 |
| Glipizide | |
| Glivec | |
| Glizide | 25 |
| Glucagen Hypokit | 24 |
| Glucagon hydrochloride | 24 |
| Glucerna Select | 218 |
| Glucerna Select RTH | |
| Glucobay | 25 |
| Glucose [Dextrose] | 47 |
| Gluten Free Foods | |
| Glycerin with sodium | |
| saccharin | 211 |
| Glycerin with sucrose | 211 |
| Glycerol | |
| Alimentary | 04 |
| | |
| Extemporaneous | |
| Glyceryl trinitrate | ~ ~ |
| Alimentary | 22 |
| Cardiovascular | |
| Glycopyrronium | 197 |
| Glycopyrronium bromide | |
| Glytrin | 59 |
| | |

| Gold Knight Gopten Goserelin acetate Granisetron Gutron | 50 88 138 138 |
|---|------------------------|
| Gynaecological | |
| Anti-infectives | 76 |
| - H - | |
| Habitrol | 156 |
| Haemophilus influenzae type B | 0.40 |
| vaccine Haldol | |
| Haldol Concentrate | |
| Haloperidol | |
| Haloperidol decanoate | 142 |
| Hamilton Sunscreen | 72 |
| Havrix | |
| Havrix Junior | 246 |
| HBvaxPRO | 247 |
| healthE Dimethicone 5% | 67 |
| healthE Fatty Cream | 67 |
| healthE Glycerol BP | 211 |
| healthE Urea Cream | 67 |
| Healtheries Simple Baking | |
| Mix | |
| Hemastix | |
| Heparin sodium Heparinised saline | 45 |
| Heparon Junior | 210 |
| Hepatitis A vaccine | |
| Hepatitis B recombinant | |
| vaccine | |
| Hepsera | 102 |
| Herceptin | 190 |
| Hexamine hippurate | 114 |
| Hiprex | |
| Histaclear | |
| Histafen | |
| Holoxan Horleys Bread Mix | |
| Horleys Flour | 229 |
| Hormone Replacement Therapy - | |
| Systemic | |
| HPV | 247 |
| Humalog | 25 |
| Humalog Mix 25 | |
| Humalog Mix 50 | 24 |
| Human papillomavirus (6, 11, 16 and 18) vaccine [HPV] | |
| | |
| Humatin | |
| Humira | |
| HumiraPen | 180 |

| Humulin 30/7024 |
|-------------------------------|
| Humulin NPH24 |
| |
| Humulin R24 |
| Hyaluronic acid203 |
| Hybloc52 |
| Hydralazine |
| |
| Hydralazine hydrochloride59 |
| Hydrea163 |
| Hydrocortisone |
| Dermatological |
| |
| Hormone80 |
| Hydrocortisone acetate21 |
| Hydrocortisone and paraffin |
| liquid and lanolin65 |
| Hydrocortisone butyrate |
| |
| Hydrocortisone with |
| cinchocaine |
| Hydrocortisone with |
| miconazole66 |
| Hydrocortisone with natamycin |
| |
| and neomycin66 |
| Hydrogen peroxide |
| Alimentary |
| Dermatological63 |
| |
| Hydroxocobalamin37 |
| Hydroxychloroquine116 |
| Hydroxyurea163 |
| Hygroton56 |
| Hylo-Fresh203 |
| Hyoscine hydrobromide138 |
| |
| Hyoscine N-butylbromide22 |
| Hypam150 |
| Hyperuricaemia and |
| Antigout |
| Hypnovel150 |
| |
| Hypromellose203 |
| Hypromellose with Dextran203 |
| Hysite |
| • |
| -1- |
| Ibiamox93 |
| Ibugesic115 |
| Ibuprofen115 |
| Idarubicin hydrochloride163 |
| |
| Ifosfamide |
| Ikorel60 |
| lloprost61 |
| |
| Imatinib mesilate167 |

Immune Modulators112

| Indacaterol |
|------------------------------------|
| Indapamide56 |
| Indinavir111 |
| Infanrix IPV246 |
| Infanrix-hexa246 |
| Infant Formulae231 |
| Influenza vaccine248 |
| Influvac248 |
| Inhaled Corticosteroids194 |
| Inhaled Long-acting |
| Beta-adrenoceptor |
| Agonists 194 |
| Innovacon hCG One Step |
| Pregnancy Test |
| Inset 30 |
| Inset II |
| Insulin aspart24 |
| Insulin aspart with insulin aspart |
| protamine |
| protarnine |
| Insulin glargine |
| Insulin glulisine |
| Insulin isophane24 |
| Insulin isophane with insulin |
| neutral24 |
| Insulin lispro25 |
| Insulin lispro with insulin lispro |
| protamine24 |
| Insulin neutral24 |
| Insulin pen needles27 |
| Insulin pump28 |
| Insulin pump accessories28 |
| Insulin pump infusion set (steel |
| cannula)29 |
| Insulin pump infusion set (teflon |
| cannula, angle insertion with |
| insertion device) |
| Insulin pump infusion set (teflon |
| cannula, angle insertion) |
| Insulin pump infusion set (teflon |
| cannula, straight insertion with |
| insertion device) |
| Insulin pump infusion set (teflon |
| cannula, straight insertion) |
| Insulin pump reservoir |
| Insulin syringes, disposable with |
| attached needle |
| Intal Forte CFC Free |
| Intal Spincaps |
| Intelence |
| Interferon alfa-2a112 |
| Interferon alfa-2b112 |
| |
| |
| |

| Interferon beta-1-alpha149 |
|----------------------------------|
| Interferon beta-1-beta149 |
| Intra-uterine device |
| Intron-A112 |
| Invega Sustenna142 |
| IPOL |
| pratropium bromide |
| Iressa167 |
| Irinotecan Actavis 100160 |
| Irinotecan Actavis 40160 |
| Irinotecan hydrochloride160 |
| Irinotecan-Rex160 |
| Iron polymaltose |
| Isentress111 |
| Isenitess |
| Ismo 40 Retard59 |
| Isoniazid101 |
| soprenaline |
| Isoptin54 |
| Isopto Carpine202 |
| Isosorbide mononitrate |
| Isosource Standard225 |
| Isosource Standard RTH225 |
| Isotretinoin62 |
| Ispaghula (psyllium) husk |
| Isradipine |
| Isuprel |
| Itch-Soothe64 |
| Itraconazole98 |
| Itrazole |
| lvermectin68 |
| I- |
| - J - Jadelle |
| |
| Jevity225 Jevity HiCal RTH226 |
| Jevity RTH225 |
| |
| - К - |
| Kaletra111 |
| Kemadrin125 |
| Kenacomb200 |
| Kenacort-A 1080 |
| Kenacort-A 4080 |
| Kenalog in Orabase |
| Ketocal 3:1234 |
| KetoCal 4:1234 |
| Ketoconazole |
| Dermatological71 |
| Infection |
| Ketogenic Diet234 |
| Ketone blood beta-ketone |
| electrodes25 |

Ketoprofen115

| Ketostix25 |
|---|
| Kindergen220 |
| Kinson124 |
| Kivexa110 |
| Klacid92 |
| Kliogest83 |
| Kliovance83 |
| Kogenate FS42 |
| Konakion MM42 |
| Konsyl-D34 |
| -L- |
| -L- |
| L-asparaginase161 |
| Labetalol |
| Lacosamide |
| Lactulose |
| Laevolac |
| Lamictal135 |
| Lamivudine103, 110 |
| Lamivudine Alphapharm110 |
| Lamotrigine135 |
| Lamprene101 |
| Lanoxin51 |
| Lanoxin PG51 |
| Lansoprazole23 |
| Lantus |
| Lantus SoloStar24 |
| Lanvis160 |
| Lapatinib ditosylate168 |
| Largactil |
| Lasix |
| Latanoprost202 |
| Lax-Sachets |
| Lax-Tab |
| Lax tab |
| Laxol |
| Leflunomide |
| Lenalidomide |
| Letraccord |
| Letrazela 170 |
| Letrozole |
| Leukeran FC157 |
| Leukotriene Receptor Antagonists 197 |
| Antagonists 197 |
| Leunase162 |
| Leuprorelin89 |
| Leustatin159 |
| Levetiracetam135 |
| Levetiracetam-Rex135 |
| Levobunolol201 |
| Levocabastine201 |
| Levodopa with benserazide124 |
| Levodopa with carbidopa124 |
| l evomepromazine maleate |

| Levonorgestrel |
|-----------------------------------|
| Genito-Urinary75-76 |
| Hormone |
| Levothyroxine84 |
| 5 |
| Levothyroxine (mercury pharma) |
| Lidocaine |
| [Lignocaine] 125–126 |
| Lidocaine [Lignocaine] |
| hydrochloride 126 |
| Lidocaine [Lignocaine] with |
| chlorhexidine |
| Lidocaine [Lignocaine] with |
| Lidocaine [Lignocaine] with |
| prilocaine 126 |
| Lidocaine-Claris126 |
| Lifestyles Flared73 |
| Lignocaine80, 125, 126 |
| Hormone80 |
| Nervous125, 126 |
| Link Healthcare99 |
| Lioresal Intrathecal123 |
| Lipazil56 |
| Lipid-Modifying Agents56 |
| Lipitor |
| Liquigen217 |
| Lisinopril49 |
| Lisuride hydrogen maleate |
| Lithicarb FC140 |
| Lithium carbonate140 |
| Livostin |
| LIVOSUIT |
| |
| Locacorten-Viaform ED's |
| Local preparations for Anal and |
| Rectal Disorders21 |
| Locasol232 |
| Locoid66, 71 |
| Locoid Crelo66 |
| Locoid Lipocream66 |
| Locorten-Vioform200 |
| Lodoxamide201 |
| Logem135 |
| Lomide201 |
| Lomustine157 |
| Loniten59 |
| Loperamide hydrochloride20 |
| Lopinavir with ritonavir111 |
| Lopresor |
| Loprofin231 |
| Loprofin Mix231 |
| Lopionin with |
| Loralix |
| Loratadine193 |
| Luraraume |
| Lorazepam144 |
| |

| Lormetazepam149 |
|-------------------------------|
| Losartan Actavis51 |
| Losartan potassium51 |
| Losartan potassium with |
| hydrochlorothiazide51 |
| Lovir104 |
| Loxalate131 |
| Loxamine131 |
| Lucrin Depot PDS89 |
| Ludiomil130 |
| Lumigan202 |
| Lycinate59 |
| Lyderm70 |
| - M - |
| m-Eslon128 |
| M-M-R II249 |
| m-Mometasone66 |
| Mabthera188 |
| Macrogol 3350 with potassium |
| chloride, sodium bicarbonate |
| and sodium chloride |
| Macrogol 400 and propylene |
| glycol |
| Madopar 125124 |
| Madopar 250124 |
| Madopar 62.5124 |
| Madopar HBS124 |
| Madopar Rapid124 |
| Magnesium hydroxide211 |
| Magnesium sulphate |
| Malathion with permethrin and |
| piperonyl butoxide70 |
| Maprotiline hydrochloride130 |
| Marevan |
| Marine Blue Lotion SPF 50+72 |
| Marquis Black73 |
| Marquis Conforma73 |

Marquis Protecta73 Marquis Selecta73 Marquis Sensolite73 Marquis Supalite73 Marquis Titillata73 MarquisTantiliza73 Martindale Acetylcysteine204 Marvelon 2874 Mask for spacer device199 Mast Cell Stabilisers198 Maxitrol201 MCT oil (Nutricia)217 Measles, mumps and rubella Mebendazole91

| Mebeverine hydrochloride | |
|----------------------------------|-----|
| Medrol | 80 |
| Medroxyprogesterone acetate | |
| Genito-Urinary | |
| Hormone | |
| Mefenamic acid | |
| Megestrol acetate | |
| Meloxicam | 115 |
| Melphalan | 157 |
| Menactra | 249 |
| Meningococcal (groups A, C, Y | |
| and W-135) congugate | |
| vaccine | 249 |
| Meningococcal c congugated | |
| vaccine | 249 |
| Menthol | 64 |
| Mercaptopurine | 160 |
| Mercilon 28 | 74 |
| Mesalazine | 21 |
| Mesna | 163 |
| Mestinon | 115 |
| Metabolic Disorder Agents | 35 |
| Metamide | |
| Metformin hydrochloride | 25 |
| Methadone hydrochloride | |
| Extemporaneous | 211 |
| Nervous | |
| Methatabs | 127 |
| Methopt | 203 |
| Methotrexate | |
| Methotrexate Ebewe | 160 |
| Methotrexate Sandoz | |
| Methyl hydroxybenzoate | 211 |
| Methylcellulose | |
| Methylcellulose with glycerin an | |
| sodium saccharin | 212 |
| Methylcellulose with glycerin an | d |
| sucrose | 212 |
| Methyldopa | 55 |
| Methylphenidate | |
| hydrochloride | 152 |
| Methylphenidate hydrochloride | |
| extended-release | |
| Methylprednisolone | 80 |
| Methylprednisolone (as sodium | |
| succinate) | 80 |
| Methylprednisolone | |
| aceponate | 66 |
| Methylprednisolone acetate | 80 |
| Methylprednisolone acetate with | n |
| lidocaine [Lignocaine] | 80 |
| Methylxanthines | 198 |
| Metoclopramide | |

| hydrochloride | 138 |
|--|------|
| Metolazone | |
| Metopirone | 90 |
| Metoprolol - AFT CR | 53 |
| Metoprolol succinate | 53 |
| Metoprolol tartrate | |
| Metronidazole | 100 |
| Metyrapone | |
| Mexiletine hydrochloride | 51 |
| Mexiletine Hydrochloride | |
| USP | . 51 |
| Miacalcic | 79 |
| Mianserin hydrochloride | |
| Micolette | |
| Miconazole | 36 |
| Miconazole nitrate | |
| Dermatological | 64 |
| Genito-Urinary | |
| Micreme | |
| Micreme H | 66 |
| Microgynon 30 | 74 |
| Microgynon 50 ED | 74 |
| Microlut | |
| Midazolam | |
| Midodrine | |
| Minerals | |
| Minidiab | |
| Minirin | |
| Mino-tabs | |
| Minocycline hydrochloride | |
| Minomycin | |
| Minor Skin Infections | |
| Minoxidil | |
| Mirena | |
| Mirtazapine | |
| Misoprostol Mitomycin C | 22 |
| | |
| Mitozantrone | 164 |
| Mitozantrone Ebewe | |
| Mixtard 30 | |
| Moclobemide | |
| Modafinil | |
| Modavigil | |
| Modecate | |
| Moduretic | |
| Mogine | |
| Mometasone furoate | |
| Monogen | |
| Montelukast | 197 |
| Moroctocog alfa [Recombinant | |
| factor VIII] Morphine hydrochloride | . 42 |
| Norphine nyarochloride | 128 |
| Morphine sulphate | 128 |
| | |

| Morphine tartrate | 128 |
|--------------------------------|-----|
| Motetis | 125 |
| Mouth and Throat | 36 |
| Moxifloxacin | 96 |
| MSUD Maxamaid | 230 |
| MSUD Maxamum | |
| Mucilaginous laxatives with | |
| stimulants | 34 |
| Mucolytics | 198 |
| Multiple Sclerosis | |
| Treatments | 144 |
| Multivitamins | 37 |
| Mupirocin | 63 |
| Muscle Relaxants | 123 |
| Mvite | |
| Myambutol | 101 |
| Mycobutin | |
| MycoNail | 63 |
| Mycophenolate mofetil | 174 |
| Mycostatin | 64 |
| Mydriacyl | 202 |
| Mylan Atenolol | |
| Mylan-Bosentan | 60 |
| Mylanta P | 20 |
| Myleran | |
| Myocrisin | |
| Myometrial and Vaginal Hormone | |
| Preparations | 76 |
| - N - | |
| Nadolol | 53 |
| | |

| Nadolol | 53 |
|---------------------------|-----|
| Nalcrom | 21 |
| Naloxone hydrochloride | 204 |
| Naltraccord | 155 |
| Naltrexone hydrochloride | 155 |
| Naphazoline hydrochloride | 203 |
| Naphcon Forte | 203 |
| Naprosyn SR 1000 | |
| Naprosyn SR 750 | |
| Naproxen | 115 |
| Nardil | 130 |
| Nasal Preparations | 199 |
| Natalizumab | 145 |
| Natulan | 164 |
| Nausicalm | 138 |
| Navelbine | 165 |
| Nedocromil | 198 |
| Nefopam hydrochloride | |
| Neisvac-C | |
| Neo-B12 | 37 |
| Neo-Mercazole | 84 |
| Neocate Advance | |
| Neocate Gold | 232 |
| Neocate LCP | 232 |

| Neoral191 |
|---------------------------------|
| Neostigmine metilsulfate115 |
| Nepro HP (strawberry)221 |
| Nepro HP (vanilla)221 |
| Nepro HP RTH |
| 1 |
| Nerisone |
| Neulactil140 |
| Neulastim46 |
| Neurontin134 |
| NeuroTabs |
| Nevirapine110 |
| Nevirapine Alphapharm110 |
| Nicorandil |
| |
| Nicotine156 |
| Nicotinic acid56 |
| Nifedipine54 |
| Nifuran114 |
| Nilotinib168 |
| Nilstat |
| Alimentary |
| Allifiendary |
| Genito-Urinary76 |
| Infection99 |
| Nipent164 |
| Nitrados150 |
| Nitrates59 |
| Nitrazepam150 |
| Nitroderm TTS59 |
| Nitrofurantoin114 |
| Nitrolingual Pump Spray59 |
| Nizoral |
| |
| Noctamid149 |
| Nodia20 |
| Noflam 250115 |
| Noflam 500115 |
| Non-Steroidal Anti-Inflammatory |
| Drugs 115 |
| Nonacog alfa [Recombinant |
| factor IX] |
| Norethisterone |
| Genito-Urinary75 |
| Hormone |
| |
| Norflex |
| Norfloxacin114 |
| Noriday 2875 |
| Norimin75 |
| Normacol Plus |
| Normison150 |
| Norpress130 |
| Nortriptyline hydrochloride |
| Norvir111 |
| |
| NovaSource Renal |
| Novatretin70 |
| NovoMix 30 FlexPen24 |
| |

| NovoRapid24 | |
|---|--|
| NovoRapid FlexPen24 | |
| NovoRapid Penfill24 | |
| NovoSeven RT41 | |
| Noxafil99 | |
| Nozinan140 | |
| Nuelin | |
| Nuelin-SR198 | |
| Nupentin | |
| Nutilis | |
| | |
| Nutrient Modules | |
| Nutrini Energy Multi Fibre220 | |
| Nutrini Energy RTH220 | |
| Nutrini Low Energy Multi | |
| Fibre | |
| Nutrini RTH220 | |
| Nutrison Concentrated228 | |
| Nutrison Energy225 | |
| Nutrison Energy Multi Fibre | |
| Nutrison Multi Fibre | |
| Nutrison Standard RTH | |
| | |
| Nyefax Retard54 | |
| Nystatin | |
| Alimentary36 | |
| Dermatological64 | |
| Genito-Urinary76 | |
| Infection99 | |
| | |
| NZB Low Gluten Bread Mix229 | |
| NZB Low Gluten Bread Mix | |
| - 0 - | |
| - O - Octocog alfa [Recombinant factor | |
| - O - Octocog alfa [Recombinant factor VIII]42 | |
| - O - Octocog alfa [Recombinant factor VIII] | |
| - O - Octocog alfa [Recombinant factor VIII] | |
| - O - Octocog alfa [Recombinant factor VIII] | |
| - O - Octocog alfa [Recombinant factor VIII] | |
| - O - Octocog alfa [Recombinant factor VIII] | |
| - O - Octocog alfa [Recombinant factor VIII]42 Octreotide172 Octreotide LAR (somatostatin analogue)172 Oestradiol | |
| - O - Octocog alfa [Recombinant factor VIII]42 Octreotide172 Octreotide LAR (somatostatin analogue)172 Oestradiol82 Oestradiol valerate82 Oestradiol with | |
| - O - Octocog alfa [Recombinant factor VIII] | |
| - O - Octocog alfa [Recombinant factor VIII] | |
| - 0 - Octocog alfa [Recombinant factor VIII] | |
| - O - Octocog alfa [Recombinant factor VIII] | |
| - 0 - Octocog alfa [Recombinant factor VIII] | |
| - 0 - Octocog alfa [Recombinant factor VIII]42 Octreotide | |
| - 0 - Octocog alfa [Recombinant factor VIII] | |
| - 0 - Octocog alfa [Recombinant factor VIII] | |
| - O - Octocog alfa [Recombinant factor VIII] | |
| - 0 - Octocog alfa [Recombinant factor VIII] | |
| - 0 - Octocog alfa [Recombinant factor VIII] | |
| - O - Octocog alfa [Recombinant factor VIII] | |
| - 0 - Octocog alfa [Recombinant factor VIII] | |
| - 0 - Octocog alfa [Recombinant factor VIII] | |
| - 0 - Octocog alfa [Recombinant factor VIII] | |
| - 0 - Octocog alfa [Recombinant factor VIII] | |

| Onbrez Breezhaler | 194 |
|--------------------------------|-------|
| Oncaspar | 164 |
| OncoTICE | |
| Ondansetron | |
| Ondansetron ODT-DRLA | 138 |
| One-Alpha | 37 |
| Onelink | |
| Onkotrone | |
| Onrex | |
| | |
| Ora-Blend | |
| Ora-Blend SF | |
| Ora-Plus | |
| Ora-Sweet | .211 |
| Ora-Sweet SF | |
| Orabase | 36 |
| Oral Supplements/Complete Diet | |
| (Nasogastric/Gastrostomy | |
| Tube Feed) | 218 |
| Oratane | |
| Orgran | |
| Ornidazole | .100 |
| Orphenadrine citrate | |
| Ortho All-flex | 73 |
| Ortho-tolidine | 78 |
| Oruvail SR | .115 |
| Osmolite | .225 |
| Osmolite RTH | |
| Other Endocrine Agents | 89 |
| Other Oestrogen | |
| Preparations | . 83 |
| Other Progestogen | |
| Preparations | 83 |
| Other Skin Preparations | 72 |
| Ovestin | |
| Genito-Urinary | 76 |
| Hormone | |
| Ox-Pam | |
| Oxaliplatin | |
| Oxaliplatin Actavis 100 | |
| Oxaliplatin Actavis 50 | 150 |
| Oxaliplatin Ebewe | 150 |
| | |
| Oxazepam | .144 |
| Oxis Turbuhaler | . 194 |
| Oxpentifylline | |
| Oxybutynin | // |
| Oxycodone ControlledRelease | |
| Tablets(BNM) | 129 |
| Oxycodone hydrochloride | .129 |
| Oxycodone Orion | |
| OxyContin | |
| OxyNorm | |
| Oxytocin | |
| Oxytocin BNM | 76 |

INDEX Generic Chemicals and Brands

| Oxytocin with ergometrine |
|-------------------------------|
| maleate76 |
| Ozole |
| - P - |
| Pacifen123 |
| Pacific Buspirone143 |
| Paclitaxel |
| Paclitaxel Actavis |
| Paclitaxel Ebewe |
| Paediatric Seravit |
| Paliperidone |
| Pamidronate disodium118 |
| Pamisol118 |
| Panadol127 |
| |
| Pancreatic enzyme |
| Pantoprazole |
| Pantoprazole Actavis 2023 |
| Pantoprazole Actavis 4023 |
| Panzytrat |
| Papaverine hydrochloride60 |
| Para Plus70 |
| Para-amino salicylic acid101 |
| Paracare127 |
| Paracare Double Strength127 |
| Paracetamol127 |
| Paracetamol + Codeine |
| (Relieve)129 |
| Paracetamol with codeine129 |
| Paradigm 1.8 Reservoir32 |
| Paradigm 3.0 Reservoir32 |
| Paradigm 52228 |
| Paradigm 72228 |
| Paradigm Mio MMT-92131 |
| Paradigm Mio MMT-92331 |
| Paradigm Mio MMT-92531 |
| Paradigm Mio MMT-94131 |
| Paradigm Mio MMT-94331 |
| Paradigm Mio MMT-94531 |
| Paradigm Mio MMT-96531 |
| Paradigm Mio MMT-97531 |
| Paradigm Quick-Set |
| Paradigm Quick-Set MMT-386 |
| Paradiam Quick-Set |
| Paradigm Quick-Set MMT-387 |
| Paradigm Quick-Set |
| MMT-396 |
| Paradiam Quick Sat |
| Paradigm Quick-Set MMT-397 |
| Paradigm Quick-Set |
| MMT-398 |
| IVIIVI 1-330 |
| Paradigm Quick-Set MMT-399 |
| |
| Paradigm Silhouette |

| MMT-368 30 |
|-----------------------------------|
| Paradigm Silhouette MMT-377 |
| Paradigm Silhouette MMT-37830 |
| Paradigm Silhouette MMT-381 |
| Paradigm Silhouette MMT-382 |
| Paradigm Silhouette MMT-383 |
| Paradigm Silhouette MMT-384 |
| Paradigm Sure-T MMT-86429 |
| Paradigm Sure-T MMT-86629 |
| Paradigm Sure-T MMT-87429 |
| Paradigm Sure-T MMT-87629 |
| Paradigm Sure-T MMT-88429 |
| Paradigm Sure-T MMT-88629 |
| Paraffin |
| Paraffin liquid with soft white |
| paraffin |
| Paraffin liquid with wool fat |
| Paraldehyde |
| Parasiticidal Preparations |
| Parnate |
| Paromomycin96 |
| Paroxetine hydrochloride |
| Paser101 |
| Patanol203 |
| Paxam144 |
| Pazopanib169 |
| Peak flow meter |
| Pedialyte - Bubblegum48 |
| Pediasure |
| Pediasure RTH220 |
| Pegaspargase |
| Pegasys113 |
| Pegasys RBV Combination |
| Pack 113 |
| Pegfilgrastim |
| Pegylated interferon alfa-2a113 |
| Penicillamine |
| PenMix 3024 |
| PenMix 40 |
| PenMix 5024 |
| Pentasa21 |
| Pentostatin |
| [Deoxycoformycin]164 |
| Pentoxifylline [Oxpentifylline]60 |
| Pepti Junior Gold Karicare |
| |
| Aptamil |
| Peptisoothe22 |

| Peptisorb | 222 |
|---|-------------------|
| Perhexiline maleate | 54 |
| Pericyazine | |
| Perindopril | |
| Permethrin | 70 |
| Persantin | |
| Peteha | 101 |
| Pethidine hydrochloride | |
| Pevaryl | 64 |
| Pexsig | |
| Pharmacare | |
| Phenelzine sulphate | 130 |
| Phenobarbitone | |
| Phenobarbitone sodium | |
| Phenobarbitone sodium Extemporaneous | 212 |
| Nervous | |
| | |
| Phenoxybenzamine hydrochloride | 49 |
| Phonovy/motny/inonicillin | |
| (Penicillin V) | |
| Phenytoin sodium133 | 3. 135 |
| Phlexy 10 | 231 |
| Phosphate-Sandoz | 48 |
| Phosphorus | 48 |
| Phytomenadione | 42 |
| Pilocarpine hydrochloride | 202 |
| Pimafucort | 66 |
| Pindolol | 53 |
| Pinetarsol | |
| Pioglitazone | |
| Piportil | 143 |
| Pipothiazine palmitate | 143 |
| Pizaccord | 25 |
| Pizotifen | |
| PKU Anamix Infant | |
| PKU Anamix Junior | 231 |
| PKU Anamix Junior LQ | 231 |
| PKUL ophley I O 10 | 221 |
| PKU Lophlex LQ 10 PKU Lophlex LQ 20 | 221 |
| Plaquenil | 116 |
| Plendil ER | |
| pms-Bosentan | |
| Pneumococcal (PCV13) | 00 |
| Pneumococcal (PCV13) vaccine | 250 |
| Pneumococcal (PPV23) | 200 |
| polysaccharide vaccine | 250 |
| Pneumovax 23 | |
| Podophyllotoxin | <u>2</u> 00 70 |
| Polaramine | 102 |
| Poliomyelitis vaccine | 250 |
| Poliomyeinis vaccine Poloxamer | 200 24 |
| Poloxamer Poly-Gel | |
| Poly-Ger Poly-Tears | |
| ruiy-ieals | 203 |

| Poly-Visc203 |
|--|
| Polycal215 |
| Polyvinyl alcohol203 |
| Ponstan115 |
| Posaconazole |
| Postinor-176 |
| Potassium chloride47–48 |
| Potassium citrate77 |
| Potassium iodate |
| Povidone iodine68 |
| Pradaxa45 |
| Pramipexole hydrochloride |
| Prasugrel |
| Pravastatin |
| Praziquantel91 |
| Prazosin |
| Pred Forte201 |
| Pred Mild201 |
| Prednisolone80 |
| Prednisolone acetate |
| Prednisone |
| Pregnancy Tests - hCG Urine77 |
| Premarin |
| Premia 2.5 Continuous83 |
| Premia 5 Continuous |
| Prevenar 13250 |
| Prezista |
| |
| Priadel |
| Primacin100 Primaquine phosphate100 |
| Primidone135 |
| Primolut N84 |
| |
| Probenecid |
| Probenecid-AFT |
| Procaine penicillin |
| Procarbazine hydrochloride |
| Prochlorperazine |
| Proctosedyl |
| Procur |
| Procyclidine hydrochloride |
| Procytox157 |
| Prodopa55 |
| Progesterone |
| Proglicem |
| Proglycem23 |
| Progynova82 |
| Prokinex |
| Promethazine hydrochloride194 |
| Promethazine theoclate |
| Promod |
| Propafenone hydrochloride51 |
| Propamidine isethionate200 |
| Propranolol53 |

| Propylene glycol | 212 |
|--------------------------|---------|
| Propylthiouracil | |
| Protamine sulphate | |
| Protaphane | 24 |
| Protaphane Penfill | 24 |
| Protifar | 217 |
| Protionamide | |
| Provera | 82, 84 |
| PSO | 235–238 |
| Psoriasis and Eczema | |
| Preparations | 70 |
| PTU | 85 |
| Pulmicort Turbuhaler | 194 |
| Pulmocare | 218 |
| Pulmozyme | |
| Puri-nethol | |
| Pyrazinamide | 102 |
| Pyridostigmine bromide | 115 |
| Pyridoxine hydrochloride | 37 |
| Pyrimethamine | 97 |
| Pytazen SR | 43 |
| - Q - | |
| Q 300 | |

| Q 300 | 100 |
|---------------------|-----|
| Questran-Lite | 56 |
| Quetapel | 140 |
| Quetiapine | 140 |
| Quick-Set MMT-390 | 32 |
| Quick-Set MMT-391 | 32 |
| Quick-Set MMT-392 | 32 |
| Quick-Set MMT-393 | 32 |
| Quinapril | 49 |
| Quinapril with | |
| hydrochlorothiazide | 50 |
| Quinine sulphate | 100 |
| Qvar | 194 |
| | |

- R -

| - 11 - | |
|--------------------------|-----|
| RA-Morph | 128 |
| Raloxifene hydrochloride | 118 |
| Raltegravir potassium | 111 |
| Ramipex | 124 |
| Ranbaxy-Cefaclor | 91 |
| Ranitidine | 22 |
| Ranitidine Relief | 22 |
| Ranmoxy | 93 |
| Rapamune | 191 |
| Reandron 1000 | 81 |
| Recombinant factor IX | 42 |
| Recombinant factor VIIa | 41 |
| Recombinant factor VIII | 42 |
| Rectogesic | 22 |
| Redipred | 80 |
| Refresh Night Time | |
| | |

| Renilon 7.5 | 221 |
|---|-----|
| Resonium-A | 48 |
| Resource Beneprotein | 217 |
| Resource Diabetic | 218 |
| Respigen | 196 |
| Respiratory Devices | 199 |
| Respiratory Stimulants | 199 |
| Retinol palmitate | |
| ReTrieve | |
| Retrovir | |
| Reutenox | |
| Revlimid | |
| | |
| Revolade | |
| Rexacrom | 201 |
| Reyataz | |
| Ridaura s29 | |
| Rifabutin | |
| Rifadin | |
| Rifampicin | |
| Rifaximin | 23 |
| Rifinah | |
| Rilutek | 125 |
| Riluzole | 125 |
| Riodine | 68 |
| Risedronate Sandoz | |
| Risedronate sodium | |
| Risperdal Consta | |
| | |
| Risperdal Quicklet | 143 |
| Risperon | |
| Ritalin | |
| Ritalin LA | |
| Ritalin SR | |
| Ritonavir | |
| Ritonavir | 100 |
| Rituximab | |
| Rivaroxaban | 45 |
| Rivastigmine | |
| Rivotril132, | |
| Rizamelt | |
| Rizatriptan | |
| Roferon-A | 112 |
| Ropinirole hydrochloride | |
| RotaTeq | 250 |
| Rotavirus live reassortant oral | |
| vaccine | 250 |
| Roxane | 53 |
| Alimentary | 20 |
| Cardiovascular | 53 |
| Roxithromycin | 93 |
| Rubifen | 152 |
| Rubifen SR | 152 |
| Rythmodan | |
| Rytmonorm | |
| · · · <i>j</i> ·· · · v · · v · · · · · · · · · · · · · · · · · · · | |

| S-26 Gold Premgro231 |
|--|
| Sabril136 |
| Salamol196 |
| Salazopyrin21 |
| Salazopyrin EN21 |
| Salbutamol |
| Salbutamol with ipratropium |
| bromide |
| Salicylic acid71 |
| Salmeterol |
| Sandomigran |
| Sandostatin LAR |
| Scalp Preparations71 |
| Scopoderm TTS |
| Sebizole |
| Sedatives and Hypnotics |
| Seebri Breezhaler |
| Selegiline hydrochloride |
| Selegime Hydrochlonde |
| Senokot |
| SensoCard27 |
| Serenace |
| Seretide |
| Seretide Accuhaler |
| Serevent194 |
| Serevent Accuhaler |
| Serophene |
| Sertraline131 |
| Sevredol128 |
| Sev Hormones Non |
| Contraceptive |
| |
| Shield 4973 |
| Shield Blue |
| Shield XL73 SII-Onco-BCG180 |
| |
| Silagra61 Sildenafil61 |
| Silhouette MMT-37130 |
| |
| Silhouette MMT-37330 Silver sulphadiazine63 |
| Silver sulphadiazine |
| Simvastatin |
| |
| Sinemet |
| Sinemet CR |
| Singulair |
| Sirolimus191 |
| Siterone |
| Slow-Lopresor |
| Sodibic |
| Sodium acid phosphate35 |
| Sodium alginate20 |

- S -

| Sodium aurothiomalate116 | |
|-----------------------------------|--|
| Sodium bicarbonate | |
| Blood47–48 | |
| Extemporaneous212 | |
| Sodium calcium edetate205 | |
| Sodium | |
| carboxymethylcellulose | |
| Sodium chloride | |
| Blood47 | |
| Respiratory198 | |
| Sodium citrate with sodium lauryl | |
| sulphoacetate35 | |
| Sodium citro-tartrate78 | |
| Sodium cromoglycate | |
| Alimentary21 | |
| Respiratory198 | |
| Sensory201 | |
| Sodium fluoride38 | |
| Sodium hyaluronate [Hyaluronic | |
| acid] 203 | |
| Sodium nitroprusside25 | |
| Sodium polystyrene | |
| sulphonate | |
| Sodium tetradecyl sulphate42 | |
| Sodium valproate135 | |
| Sofradex | |
| Soframycin200 | |
| Solian | |
| Solifenacin succinate78 | |
| Solox23 | |
| Solu-Cortef80 | |
| Solu-Medrol 80 | |
| Somatropin (Omnitrope)85 | |
| Sotacor | |
| Sotalol53 | |
| Space Chamber199 | |
| Space Chamber Plus199 | |
| Spacer device199 | |
| Spacer device autoclavable | |
| Span-K48 | |
| Spiractin55 | |
| Spiriva197 | |
| Spironolactone55 | |
| Sporanox | |
| Sprycel166 | |
| Staphlex94 | |
| Stavudine [d4T]111 | |
| Stelazine141 | |
| Stemetil138 | |
| Stesolid133 | |
| Stimulants/ADHD | |
| Treatments 150 | |
| Stiripentol136 | |
| | |

| Stocrin110 |
|----------------------------|
| Stomahesive36 |
| Strattera150 |
| Stromectol68 |
| Suboxone154 |
| Sucralfate23 |
| Sulfadiazine sodium97 |
| Sulindac |
| Sulphasalazine21 |
| Sulphur71 |
| Sumatriptan137 |
| Sunitinib170 |
| Sunscreens72 |
| Sunscreens, proprietary72 |
| Sure-T MMT-863 |
| Sure-T MMT-86529 |
| Sure-T MMT-87329 |
| Sure-T MMT-87529 |
| Sure-T MMT-88329 |
| Sure-T MMT-88529 |
| Sustagen Diabetic218 |
| Sustagen Hospital Formula |
| Sustanon Ampoules81 |
| Sutent |
| Symbicort Turbuhaler 100/6 |
| Symbicort Turbuhaler 200/6 |
| Symbicort Turbuhaler |
| 400/12 |
| Symmetrel124 |
| Sympathomimetics59 |
| Synacthen80 |
| Synacthen Depot80 |
| Synthroid84 |
| Syntometrine76 |
| Syrup (pharmaceutical |
| grade) 212 |
| grade) |
| -T- |
| - I - Tacrolimus192 |
| Tacrolimus Sandoz192 |
| Tambocor51 |
| Tambocor CR51 |
| Tampocor CR51 |

| Tambocor CR | 51 |
|--------------------------|-----|
| Tamoxifen citrate | 173 |
| Tamsulosin hydrochloride | 77 |
| Tamsulosin-Rex | 77 |
| Tap water | 212 |
| | |
| sulphate and fluorescein | 71 |
| Tarceva | 166 |
| Tasigna | 168 |
| Tasmar | 124 |
| Taxotere | 162 |

| Tegretol133 |
|--------------------------|
| Tegretol CR133 |
| Telfast193 |
| Temaccord164 |
| Temazepam150 |
| |
| Temozolomide164 |
| Tenofovir disoproxil |
| fumarate 106 |
| Tenoxicam115 |
| Tepadina158 |
| Terazosin49 |
| Terbinafine99 |
| Terbutaline sulphate196 |
| Teriparatide |
| Testosterone81 |
| Testosterone cypionate81 |
| Testosterone esters |
| Testosterone undecanoate |
| |
| Tetrabenazine125 |
| Tetrabromophenol78 |
| Tetracosactrin80 |
| Tetracyclin Wolff95 |
| Tetracycline95 |
| Teva161 |
| Thalidomide165 |
| Thalomid165 |
| Theophylline198 |
| Thiamine hydrochloride |
| THIO-TEPA158 |
| Thioguanine |
| |
| Thiotepa158 |
| Thymol glycerin |
| Thyroid and Antithyroid |
| Agents 84 |
| Ticagrelor43 |
| Tilade |
| Timolol |
| Cardiovascular53 |
| Sensory201 |
| Timoptol XE201 |
| Tiotropium bromide |
| |
| TMP |
| TOBI |
| Tobramycin |
| Infection97 |
| Sensory201 |
| Tobrex201 |
| Tofranil130 |
| Tofranil s29130 |
| Tolcapone124 |
| Tolterodine |
| Tolvon |
| Topamax |
| юранал130 |

| Topical Products for Joint and | |
|--|----------|
| Muscular Pain1 | 16 |
| Topiramate1 | 36 |
| Topiramate Actavis1 | 36 |
| Total parenteral nutrition | |
| (TPN) | 47 |
| TPN | |
| Tracleer | 60 |
| Tramadol hydrochloride1 | 29 |
| Tramal SR 1001 | 29 |
| Tramal SR 1501 | 29 |
| Tramal SR 2001 | |
| Trandate | 52 |
| Trandolapril | 50 |
| Tranexamic acid | 42 |
| Tranylcypromine sulphate1 | |
| Trastuzumab1 | 90 |
| Travatan2 | |
| Travoprost2 | 02 |
| Treatments for Dementia1 | 54 |
| Treatments for Substance Dependence1 | - 4 |
| Dependence 1 | 54 |
| Trental 400 | 60 |
| Tretinoin | ~~ |
| Dermatological | 62 |
| Oncology1 | 65 |
| Trexate | 60 |
| Triamcinolone acetonide | ~~ |
| Alimentary | 36 |
| Dermatological | |
| Hormone | 80 |
| Triamcinolone acetonide with gramicidin, neomycin and nystati | _ |
| | |
| Dermatological | |
| Sensory | |
| Triazolam1 | 50 |
| Trichozole1 | |
| Triclosan Trifluoperazine | 07 |
| hydrochloride1 | 44 |
| Trimeprazine tartrate1 | 41 |
| Trimethoprim | |
| Trisequens | 91 02 |
| Trisul | 00 |
| Trophic Hormones | 90 |
| Tropicamide2 | |
| Trusopt2 | 02 |
| Truvada1 | 102 |
| Two Cal HN | 28 |
| Two Cal HN2 Two Cal HN RTH2 | 20 |
| Tykerb1 | 68 |
| Tysabri1 | |
| 1950.0111 | 40 |

- U -

| Ultraproct21 | |
|-----------------------------|---|
| Univent196, 199 | 9 |
| Ural78 | 3 |
| Urea | 7 |
| Urex Forte55 | ō |
| Urinary Agents77 | 7 |
| Urinary Tract Infections114 | 1 |
| Uromitexan163 | 3 |
| Ursodeoxycholic acid33 | 3 |
| Ursosan | 3 |
| Utrogestan84 | 1 |

v

| - V - | |
|--------------------------------|-----|
| Vaccinations | |
| Valaciclovir | 104 |
| Valcyte | |
| Valganciclovir | |
| Vallergan Forte | |
| Valtrex | |
| Vancomycin | |
| Vannair | 195 |
| Varenicline tartrate | 156 |
| Varicella vaccine [Chicken pox | |
| vaccine] | 251 |
| Varilrix | |
| Vasodilators | |
| Vasopressin Agonists | |
| Vedafil | |
| Velcade | |
| Venlafaxine | |
| Ventavis | |
| Ventolin195 | |
| Vepesid | 163 |
| Verapamil hydrochloride | 54 |
| Vergo 16 | |
| Vermox | |
| Verpamil SR | |
| Vesanoid | |
| Vesicare | |
| Vfend | |
| Viaderm KC | |
| Victrelis | |
| Vidaza | 158 |
| Videx EC | |
| Vigabatrin | |
| Vimpat | |
| Vinblastine sulphate | |
| Vincristine sulphate | |
| Vinorelbine | |
| Vinorelbine Ebewe | |
| Viramune Suspension | |
| Viread | 106 |

| Virgan 200 Vistil 203 Vistil Forte 203 VitA-POS 203 |
|---|
| Vitabdeck |
| Vitadol C37 |
| Vital HN222 |
| Vitamin A with vitamins D and |
| C |
| Vitamin B complex37 |
| Vitamin B6 25 |
| Vitamins37–38 |
| Vivonex Pediatric232 |
| Vivonex TEN222 |
| Volibris60 |
| Voltaren115 |
| Voltaren D115 |
| Voltaren Ophtha201 |
| Volumatic |
| Voriconazole99 |
| Vosol200 |
| Votrient169 |
| Vytorin58 |
| - W - |
| Warfarin sodium |
| Wart Preparations72 |
| Wasp venom allergy |
| treatment193 |

Water

| Blood47 |
|-----------------------------|
| Extemporaneous212 |
| Wool fat with mineral oil67 |

- X -

| Xanax | 143 |
|-------------------|-----|
| Xarelto | 45 |
| Xifaxan | 23 |
| XMET Maxamum | 230 |
| Xolair | 187 |
| XP Maxamaid | 231 |
| XP Maxamum | 231 |
| Xylocaine | 126 |
| Xylocaine Viscous | 126 |
| Xyntha | 42 |
| | |

- Z -

| Zantac | 22 |
|-----------|-----|
| Zapril | 49 |
| Zarator | 57 |
| Zarontin | 133 |
| Zaroxolyn | 55 |
| Zarzio | 46 |
| Zavedos | 163 |
| Zeffix | |
| Zeldox | 141 |
| Zerit | 111 |
| Zetop | 193 |
| Ziagen | 110 |
| | |

| Zidovudine [AZT] | 111 |
|--------------------------|-----|
| Zidovudine [AZT] with | |
| lamivudine | 111 |
| Zimybe | 58 |
| Zinc and castor oil | |
| Zinc sulphate | 39 |
| Zincaps | |
| Zinnat | |
| Ziprasidone | 141 |
| Zithromax | 92 |
| Zoladex | 88 |
| Zoledronic acid | |
| Hormone | 79 |
| Musculoskeletal System | 119 |
| Zometa | |
| Zopiclone | 150 |
| Zostrix | 116 |
| Zostrix HP | 126 |
| Zovirax | |
| Zuclopenthixol decanoate | 143 |
| Zuclopenthixol | |
| hydrochloride | 141 |
| Zyban | |
| Zypine | |
| Zypine ODT | 140 |
| Zyprexa Relprevv | |
| Zytiga | |
| , . | |