

April 2016

Volume 4 Number 2

Editor: Kaye Wilson

email: enquiry@pharmac.govt.nz

Telephone +64 4 460 4990

Facsimile +64 4 460 4995

Level 9, 40 Mercer Street

PO Box 10 254 Wellington 6143

Freephone Information Line

0800 66 00 50 (9am – 5pm weekdays)

Circulation

Accessible in an electronic format at no cost from the Health Professionals section of the PHARMAC website www.pharmac.govt.nz

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month. Alternatively there is a nominal charge for an annual subscription to the printed Schedule publications. To access either of these subscriptions visit our subscription website www.schedule.co.nz.

Production

Typeset automatically from XML and T_EX.

XML version of the Schedule available from www.pharmac.govt.nz/pub/schedule/archive/

Programmers

Anrik Drenth & John Geering

email: texschedule@pharmac.govt.nz

©Pharmaceutical Management Agency



ISSN 1179-3708 pdf
ISSN 1172-9694 print

This work is licensed under the Creative Commons Attribution 3.0 New Zealand licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to PHARMAC and abide by the other licence terms. To view a copy of this licence, visit:

creativecommons.org/licenses/by/3.0/nz/.

Attribution to PHARMAC should be in written form and not by reproduction of the PHARMAC logo. While care has been taken in compiling this Schedule, PHARMAC takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

Part I General Rules 5

Part II Alimentary Tract and Metabolism 13

Blood and Blood Forming Organs 27

Cardiovascular System 39

Dermatologicals 51

Genito-Urinary System 57

Hormone Preparations 61

Infections 71

Musculoskeletal System 95

Nervous System 105

Oncology Agents and Immunosuppressants 133

Respiratory System and Allergies 177

Sensory Organs 184

Various 190

Extemporaneous Compounds (ECPs) 198

Special Foods 201

Vaccines 215

Part III Optional Pharmaceuticals 222

Index 224

Introducing PHARMAC

The Pharmaceutical Management Agency (PHARMAC) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. PHARMAC negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

PHARMAC's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about PHARMAC and the way we make funding decisions can be found on the PHARMAC website at <http://www.pharmac.health.nz/about>.

Named Patient Pharmaceutical Assessment policy

Named Patient Pharmaceutical Assessment (NPPA) provides a mechanism for individual patients to receive funding for medicines not listed in the Pharmaceutical Schedule (either at all or for their clinical circumstances). PHARMAC will assess applications that meet the prerequisites according to its Decision Criteria before deciding whether to approve applications for funding. The Decision Criteria will be used to assess both the individual clinical circumstances of each NPPA applicant, and the implications of each NPPA funding decision on PHARMAC's ability to carry out its legislative functions.

For more information on NPPA, or to apply, visit the PHARMAC website at <http://www.pharmac.health.nz/link/nppa> or call the Panel Coordinators at 0800 660 050 Option 2.

The Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by PHARMAC.

The Schedule does not show the final cost to Government of subsidising each community pharmaceutical, nor to DHB hospitals in purchasing each hospital pharmaceutical or other pharmaceuticals, including medical devices. The final cost will depend on any rebate and other arrangements PHARMAC has with the supplier or on any logistics arrangements put in place.

Finding Information in Section H

This book contains Section H of the Pharmaceutical Schedule and lists pharmaceuticals that can be used in DHB hospitals:

- Part I lists the rules in relation to use of Pharmaceuticals by DHB hospitals.
- Part II lists hospital pharmaceuticals that are funded for use in DHB hospitals. These are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. It also provides information on any national contracts that exist, and indicates which products have Hospital Supply Status (HSS).
- Part III lists optional pharmaceuticals for which national contracts exist, and DHB hospitals may choose to fund. In addition to the products listed in this book, a number of additional Optional Pharmaceuticals are listed in an addendum to Part III available at <http://www.pharmac.govt.nz>.

The listings are displayed alphabetically under each heading. The index lists both chemical entities and product brand names.

Glossary

Units of Measure

gram	g	microgram.....	mcg	millimole.....	mmol
kilogram	kg	milligram	mg	unit.....	u
international unit	iu	millilitre.....	ml		

Abbreviations

application	app	enteric coated	EC	solution	soln
capsule	cap	granules	grans	suppository	suppos
cream.....	crm	injection	inj	tablet.....	tab
dispersible	disp	liquid	liq	tincture.....	tinc
effervescent	eff	lotion	lotn		
emulsion	emul	ointment.....	oint		

HSS Hospital Supply Status (Refer to Rule 20)

Guide to Section H listings

Example

ANATOMICAL HEADING			
	Price (ex man. Excl. GST) \$	Per	Brand or Generic Manufacturer
THERAPEUTIC HEADING			
Generic name listed by therapeutic group and subgroup	CHEMICAL A - Restricted see terms below ⬇ Presentation A.....10.00	100	Brand A
	⬇ Restricted Only for use in children under 12 years of age		Brand or manufacturer's name
Indicates only presentation B1 is Restricted	CHEMICAL B - Some items restricted see terms below ⬇ Presentation B1.....1,589.00 Presentation B2 ⬇ Restricted Oncologist or haematologist	1	Brand B1 e.g. Brand B2
From 1 January 2012 to 30 June 2014, at least 99% of the total volume of this item purchased must be Brand C	CHEMICAL C Presentation C - 1% DV Limit Jan-12 to 201415.00	28	Brand C
	CHEMICAL D - Restricted see terms below ⬇ Presentation D - 1% DV Limit Mar-13 to 201438.65	500	Brand D
Standard national price excluding GST	⬇ Restricted <i>Limited to five weeks' treatment</i> Either: 1 For the prophylaxis of venous thromboembolism following a total hip replacement; or 2 For the prophylaxis of venous thromboembolism following a total knee replacement.		Product with Hospital Supply Status (HSS) Quantity the Price applies to
Form and strength	CHEMICAL E Presentation E.....		e.g. Brand E Not a contracted product
⬆ Item restricted (see above); ⬇ Item restricted (see below) Products with Hospital Supply Status (HSS) are in bold			

INTRODUCTION

Section H contains general rules that apply, and other information relating, to Hospital Pharmaceuticals and Optional Pharmaceuticals.

Where relevant, Section H shows the Price at which a Pharmaceutical can be purchased directly from the Pharmaceutical supplier by DHBs, providers of logistics services, wholesalers or other such distributors, or Contract Manufacturers.

The Price is determined via contractual arrangements between PHARMAC and the relevant Pharmaceutical supplier. Where a Pharmaceutical is listed in Part II of Section H, but no Price and/or brand of Pharmaceutical is indicated, each DHB may purchase any brand and/or pay the price that the DHB negotiates with the relevant Pharmaceutical supplier.

As required by section 23(7) of the Act, in performing any of its functions in relation to the supply of Pharmaceuticals, a DHB must not act inconsistently with the Pharmaceutical Schedule.

INTERPRETATION AND DEFINITIONS

1 Interpretation and Definitions

1.1 In this Schedule, unless the context otherwise requires:

“Act”, means the New Zealand Public Health and Disability Act 2000.

“Combined Pharmaceutical Budget”, means the pharmaceutical budget set for PHARMAC by the Crown for the subsidised supply of Community Pharmaceuticals and Pharmaceutical Cancer Treatments including for named patients in exceptional circumstances.

“Community”, means any setting outside of a DHB Hospital.

“Community Pharmaceutical”, means a Pharmaceutical listed in Sections A to G or I of the Pharmaceutical Schedule that is subsidised by the Funder from the Combined Pharmaceutical Budget and, for the purposes of this Section H, includes Pharmaceutical Cancer Treatments (PCTs).

“Contract Manufacturer”, means a manufacturer or a supplier that is a party to a contract with the relevant DHB Hospital to compound Pharmaceuticals, on request from that DHB Hospital.

“Designated Delivery Point”, means at a DHB Hospital’s discretion:

- a) a delivery point agreed between a Pharmaceutical supplier and the relevant DHB Hospital, to which delivery point that Pharmaceutical supplier must supply a National Contract Pharmaceutical directly at the Price; and/or
- b) any delivery point designated by the relevant DHB Hospital or PHARMAC, such delivery point being within 30 km of the relevant Pharmaceutical supplier’s national distribution centre.

“DHB”, means an organisation established as a District Health Board by or under Section 19 of the Act.

“DHB Hospital”, means a hospital (including community trust hospitals) and/or an associated health service that is funded by a DHB including (but not limited to) district nursing services and child dental services.

“DV Limit”, means, for a particular National Contract Pharmaceutical with HSS, the National DV Limit or the Individual DV Limit.

“DV Pharmaceutical”, means a discretionary variance Pharmaceutical that does not have HSS but is used in place of one that does. Usually this means it is the same chemical entity, at the same strength, and in the same or a similar presentation or form, as the relevant National Contract Pharmaceutical with HSS. Where this is not the case, a note will be included with the listing of the relevant Hospital Pharmaceutical.

“Extemporaneously Compounded Product”, means a Pharmaceutical that is compounded from two or more Pharmaceuticals, for the purposes of reconstitution, dilution or otherwise.

“First Transition Period”, means the period of time after notification that a Pharmaceutical has been awarded HSS and before HSS is implemented.

“Funder”, means the body or bodies responsible, pursuant to the Act, for the funding of Pharmaceuticals listed on the Schedule (which may be one or more DHBs and/or the Ministry of Health) and their successors.

“Give”, means to administer, provide or dispense (or, in the case of a Medical Device, use) a Pharmaceutical, or to arrange for the administration, provision or dispensing (or, in the case of a Medical Device, use) of a Pharmaceutical, and “Given” has a corresponding meaning.

“Hospital Pharmaceuticals”, means the list of Pharmaceuticals set out in Section H Part II of the Schedule which includes some National Contract Pharmaceuticals.

“HSS”, stands for hospital supply status, which means the status of being the brand of the relevant National Contract Pharmaceutical that DHBs are obliged to purchase, subject to any DV Limit, for the period of hospital supply,

as awarded under an agreement between PHARMAC and the relevant Pharmaceutical supplier. Pharmaceuticals with HSS are listed in Section H in bold text.

“Indication Restriction”, means a limitation placed by PHARMAC on the funding of a Hospital Pharmaceutical which restricts funding to treatment of particular clinical circumstances.

“Individual DV Limit”, means, for a particular National Contract Pharmaceutical with HSS and a particular DHB Hospital, the discretionary variance limit, being the specified percentage of that DHB Hospital's Total Market Volume up to which that DHB Hospital may purchase DV Pharmaceuticals of that National Contract Pharmaceutical.

“Local Restriction”, means a restriction on the use of a Pharmaceutical in specific DHB Hospitals on the basis of prescriber type that is implemented by the relevant DHB in accordance with rule 7.

“Medical Device”, has the meaning set out in the Medicines Act 1981.

“Named Patient Pharmaceutical Assessment Advisory Panel”, means the panel of clinicians, appointed by the PHARMAC Board, that is responsible for advising PHARMAC, in accordance with its Terms of Reference, on Named Patient Pharmaceutical Assessment applications and any Exceptional Circumstances renewal applications submitted after 1 March 2012.

“National Contract”, means a contractual arrangement between PHARMAC and a Pharmaceutical supplier which sets out the basis on which any Pharmaceutical may be purchased for use in a DHB Hospital, including an agreement as to a national price.

“National Contract Pharmaceutical”, means a brand of Pharmaceutical listed in Section H, where PHARMAC has entered into contractual arrangements with the relevant Pharmaceutical supplier that specify the terms and conditions of listing, including the Price. Such Pharmaceuticals are recognisable in Section H because the relevant listing identifies the brand and Price.

“National DV Limit”, means, for a particular National Contract Pharmaceutical with HSS, the discretionary variance limit, being the specified percentage of the Total Market Volume up to which all DHB Hospitals may collectively purchase DV Pharmaceuticals of that National Contract Pharmaceutical.

“Optional Pharmaceuticals”, means the list of National Contract Pharmaceuticals set out in Section H Part III of the Schedule.

“PHARMAC”, means the Pharmaceutical Management Agency established by Section 46 of the Act.

“Pharmacode”, means the six or seven digit identifier assigned to a Pharmaceutical by the Pharmacy Guild following application from a Pharmaceutical supplier.

“Pharmaceutical”, means a medicine, therapeutic medical device, or related product or related thing listed in Sections B to I of the Schedule.

“Pharmaceutical Cancer Treatment”, means Pharmaceuticals for the treatment of cancer, listed in Sections A to G of the Schedule and identified therein as a “PCT” or “PCT only” Pharmaceutical that DHBs must fund for use in their DHB hospitals, and/or in association with outpatient services provided by their DHB Hospitals, in relation to the treatment of cancers.

“Prescriber Restriction”, means a restriction placed by PHARMAC on the funding of a Pharmaceutical on the basis of prescriber type (and where relevant in these rules, includes a Local Restriction).

“Price”, means the standard national price for a National Contract Pharmaceutical, and, unless agreed otherwise between PHARMAC and the Pharmaceutical supplier, includes any costs associated with the supply of the National Contract Pharmaceutical to, at a DHB Hospital's discretion, any Designated Delivery Point, or to a Contract Manufacturer (expressly for the purpose of compounding), but does not include the effect of any rebates which may have been negotiated between PHARMAC and the Pharmaceutical supplier.

“Restriction”, means a limitation, put in place by PHARMAC or a DHB, restricting the funding of a Pharmaceutical and includes Indication Restrictions, Local Restrictions and Prescriber Restrictions (as defined in this Part I of Section H).

“Schedule”, means this Pharmaceutical Schedule and all its sections and appendices.

“Special Authority Approval”, means an approval for funding of a Community Pharmaceutical that is marked in Sections B-G of the Schedule as being subject to a Special Authority restriction.

“Total Market Volume”, means, for a particular Hospital Pharmaceutical with HSS in any given period, in accordance with the data available to PHARMAC, the sum of:

- a) the total number of Units of the relevant Hospital Pharmaceutical with HSS purchased by all DHB Hospitals, or by a particular DHB Hospital in the case of the Individual DV Limit; and
- b) the total number of Units of all the relevant DV Pharmaceuticals purchased by all DHB Hospitals, or by a particular DHB Hospital in the case of the Individual DV Limit.

“Unapproved Indication”, means, for a Pharmaceutical, an indication for which it is not approved under the Medicines Act 1981. Clinicians prescribing Pharmaceuticals for Unapproved Indications should be aware of, and comply with, their obligations under Section 25 and/or Section 29 of the Medicines Act 1981 and as set out in rule 23.

“Unit”, means an individual unit of a Pharmaceutical (e.g. a tablet, 1 ml of an oral liquid, an ampoule or a syringe).

“Unlisted Pharmaceutical”, means a Pharmaceutical that is within the scope of a Hospital Pharmaceutical, but is not listed in Section H Part II.

- 1.2 In addition to the above interpretations and definitions, unless the content requires otherwise, a reference in the Schedule to:
 - a) the singular includes the plural; and
 - b) any legislation includes a modification and re-enactment of, legislation enacted in substitution for, and a regulation, Order in Council, and other instrument from time to time issued or made under, that legislation.

HOSPITAL SUPPLY OF PHARMACEUTICALS

2 Hospital Pharmaceuticals

- 2.1 Section H Part II contains the list of Hospital Pharmaceuticals that must be funded by DHB Hospitals. Section H Part II does not currently encompass the following categories of pharmaceuticals except for any items specifically listed in this Section H Part II:
 - a) Medical Devices;
 - b) whole or fractionated blood products;
 - c) diagnostic products which have an ex vivo use, such as pregnancy tests and reagents;
 - d) disinfectants and sterilising products, except those that are to be used in or on a patient;
 - e) foods and probiotics;
 - f) radioactive materials;
 - g) medical gases; and
 - h) parenteral nutrition.

Subject to rule 2.2, the funding of pharmaceuticals identified in a)–h) above is a decision for individual DHB Hospitals.

- 2.2 Section H Part III lists Optional Pharmaceuticals that PHARMAC and the relevant Pharmaceutical supplier have entered into contractual arrangements for the purchase of, including an agreement on a national price and other obligations such as HSS. DHB Hospitals may choose whether or not to fund the Optional Pharmaceuticals listed in Part III of Section H, but if they do, they must comply with any National Contract requirements.
- 2.3 Section H Part II does not encompass the provision of pharmaceutical treatments for DHB Hospital staff as part of an occupational health and safety programme. DHB Hospitals may choose whether or not to fund pharmaceutical treatments for such use, but if they do, they must comply with any National Contract requirements.

3 DHB Supply Obligations

- 3.1 In accordance with section 23(7) of the Act, in performing any of its functions in relation to the supply of pharmaceuticals, a DHB must not act inconsistently with the Pharmaceutical Schedule, which includes these General Rules.
- 3.2 DHB Hospitals are not required to hold stock of every Hospital Pharmaceutical listed in Section H Part II, but they must Give it within a reasonable time if it is prescribed.
- 3.3 DHB Hospitals are able to hold stock of an Unlisted Pharmaceutical if doing so is considered necessary for the DHB Hospital to be able to Give the Unlisted Pharmaceutical in a timely manner under rules 11–17 inclusive.
- 3.4 Except where permitted in accordance with rule 11, DHBs must not Give:
 - a) an Unlisted Pharmaceutical; or
 - b) a Hospital Pharmaceutical outside of any relevant Restrictions.

4 Funding

- 4.1 The purchase costs of Hospital Pharmaceuticals or Optional Pharmaceuticals administered, provided or dispensed by DHB Hospitals must be funded by the relevant DHB Hospital from its own budget, with the exception of:
 - a) Pharmaceutical Cancer Treatments;
 - b) Community Pharmaceuticals that have been brought to the DHB hospital by the patient who is being treated by outpatient Services or who is admitted as an inpatient;
 - c) Community Pharmaceuticals that have been dispensed to a mental health day clinic under a Practitioner's

- Supply Order; and
 - d) Unlisted Pharmaceutical that have been brought to the DHB Hospital by the patient who is admitted as an inpatient.
- 4.2 For the avoidance of doubt, Pharmaceutical Cancer Treatments and Community Pharmaceuticals are funded through the Combined Pharmaceutical Budget, and Unlisted Pharmaceuticals are funded by the patient.

LIMITS ON SUPPLY

5 Prescriber Restrictions

- 5.1 A DHB Hospital may only Give a Hospital Pharmaceutical that has a Prescriber Restriction if it is prescribed:
- a) by a clinician of the type specified in the restriction for that Pharmaceutical or, subject to rule 5.2, pursuant to a recommendation from such a clinician;
 - b) in accordance with a protocol or guideline that has been endorsed by the DHB Hospital; or
 - c) in an emergency situation, provided that the prescriber has made reasonable attempts to comply with rule 5.1(a) above. If on-going treatment is required (i.e. beyond 24 hours) subsequent prescribing must comply with rule 5.1(a).
- 5.2 Where a Hospital Pharmaceutical is prescribed pursuant to a recommendation from a clinician of the type specified in the restriction for that Pharmaceutical:
- a) the prescriber must consult with a clinician of the type specified in the restriction for that Pharmaceutical; and
 - b) the consultation must relate to the patient for whom the prescription is written; and
 - c) the consultation may be in person, by telephone, letter, facsimile or email; and
 - d) appropriate records are kept of the consultation, including recording the name of the advising clinician on the prescription/chart.
- 5.3 Where a clinician is working under supervision of a consultant who is of the type specified in the restriction for that Pharmaceutical, the requirements of rule 5.2 can be deemed to have been met.

6 Indication Restrictions

- 6.1 A DHB Hospital may only Give a Hospital Pharmaceutical that has an Indication Restriction, if it is prescribed for treatment of a patient with the particular clinical circumstances set out in the Indication Restriction.
- 6.2 If a patient has a current Special Authority Approval for the Hospital Pharmaceutical that the DHB Hospital wishes to Give, then the Indication Restriction is deemed to have been met.
- 6.3 If a Hospital Pharmaceutical has an Indication Restriction that is “for continuation only” then the DHB Hospital should only Give the Hospital Pharmaceutical where:
- a) the patient has been treated with the Pharmaceutical in the Community; or
 - b) the patient is unable to be treated with an alternative Hospital Pharmaceutical, and the prescriber has explained to the patient that the Pharmaceutical is not fully subsidised in the Community.

7 Local Restrictions

- 7.1 A DHB Hospital may implement a Local Restriction, provided that:
- a) in doing so, it ensures that the Local Restriction does not unreasonably limit funded access to the Hospital Pharmaceutical or undermine PHARMAC’s decision that the Hospital Pharmaceutical must be funded;
 - b) it provides PHARMAC with details of each Local Restriction that it implements; and
- 7.2 PHARMAC may, when it considers that a Local Restriction does not conform to rule 7.1 above, require a DHB to amend or remove that Local Restriction.

8 Community use of Hospital Pharmaceuticals

- 8.1 Except where otherwise specified in Section H, DHB Hospitals can Give any Hospital Pharmaceutical to a patient for use in the Community, provided that:
- a) the quantity does not exceed that sufficient for up to 30 days’ treatment, unless:
 - i) it would be inappropriate to provide less than the amount in an original pack; or
 - ii) the relevant DHB Hospital has a Dispensing for Discharge Policy and the quantity dispensed is in accordance with that policy; and
 - b) the Hospital Pharmaceutical is supplied consistent with any applicable Restrictions.

9 Community use of Medical Devices

- 9.1 Subject to rules 9.2 and 9.3, DHB Hospitals may Give a Medical Device for patients for use in the Community.
- 9.2 Where a Medical Device (or a similar Medical Device) is a Community Pharmaceutical, the DHB Hospital must supply:

- a) the brand of Medical Device that is listed in Sections A-G of the Schedule; and
 - b) only to patients who meet the funding eligibility criteria set out in Sections A-G of the Schedule.
- 9.3 Where a DHB Hospital has supplied a Medical Device to a patient; and
- a) that Medical Device (or a similar Medical Device) is subsequently listed in Sections A-G of the Schedule; and
 - b) the patient would not meet any funding eligibility criteria for the Medical Device set out in Sections A-G of the Schedule; and
 - c) the Medical Device has consumable components that need to be replaced throughout its usable life; then DHB Hospitals may continue to fund consumable products for that patient until the end of the usable life of the Medical Device. At the end of the usable life of the device, funding for a replacement device must be consistent with the Pharmaceutical Schedule and/or in accordance with the Named Patient Pharmaceutical Assessment policy.
- 9.4 DHB Hospitals may also continue to fund consumable products, as in rule 9.3 above, in situations where the DHB has been funding consumable products but where the Medical Device was funded by the patient.

10 Extemporaneous Compounding

- 10.1 A DHB Hospital may Give any Extemporaneously Compounded Product for a patient in its care, provided that:
- a) all of the component Pharmaceuticals of the Extemporaneously Compounded Product are Hospital Pharmaceuticals; and
 - b) the Extemporaneously Compounded Product is supplied consistent with any applicable rules or Restrictions for its component Hospital Pharmaceuticals.
- 10.2 For the avoidance of doubt, this rule 10.1 applies to any Extemporaneously Compounded Product, whether it is manufactured by the DHB Hospital or by a Contract Manufacturer.

EXCEPTIONS

11 Named Patient Pharmaceutical Assessment

- 11.1 A DHB Hospitals may only Give:
- a) an Unlisted Pharmaceutical; or
 - b) a Hospital Pharmaceutical outside of any relevant Restrictions,
- in accordance with the Named Patient Pharmaceutical Assessment Policy or rules 12–17 inclusive.

12 Continuation

- 12.1 Where a patient's clinical circumstances have been stabilised via treatment in the Community with a pharmaceutical that has not been funded by the Funder, and that patient is admitted to hospital as an inpatient, a DHB Hospital may fund that pharmaceutical for the duration of the patient's stay, where:
- a) the patient has not brought (or cannot arrange to bring) the pharmaceuticals to the DHB Hospital, or pharmacy staff consider that the pharmaceuticals brought to the DHB Hospital by the patient cannot be used; and
 - b) interrupted or delayed treatment would have significant adverse clinical consequences; and
 - c) it is not considered appropriate to switch treatment to a Hospital Pharmaceutical.

13 Pre-Existing Use

- 13.1 Subject to 13.2, where a DHB Hospital has Given a pharmaceutical for a patient prior to 1 July 2013, and the pharmaceutical:
- a) is an Unlisted Pharmaceutical; or
 - b) treatment of the patient would not comply with any relevant Restrictions;
- the DHB Hospital may continue to Give that pharmaceutical if it is considered that there would be significant adverse clinical consequences from ceasing or switching treatment.
- 13.2 Each DHB Hospital must, by no later than 1 October 2013, provide PHARMAC with a report on pharmaceuticals it has Given in accordance with this rule 13 where treatment has continued beyond 1 August 2013.

14 Clinical Trials and Free Stock

- 14.1 DHB Hospitals may Give any pharmaceutical that is funded by a third party and is being used:
- 14.1.1 as part of a clinical trial that has Ethics Committee approval; or
 - 14.1.2 for on-going treatment of patients following the end of such a clinical trial.
- 14.2 DHB Hospitals may Give any pharmaceutical that is provided free of charge by a supplier, provided that the pharmaceutical is provided as part of a programme of which the DHB, or supplier, has notified PHARMAC.

15 Pharmaceutical Cancer Treatments in Paediatrics

DHB Hospitals may Give any pharmaceutical for use within a paediatric oncology/haematology service for the treatment of

cancer.

16 Other Government Funding

DHB Hospitals may Give any pharmaceutical where funding for that pharmaceutical has been specifically provided by a Government entity other than PHARMAC or a DHB.

17 Other Exceptions

- 17.1 PHARMAC may also approve the funding of a pharmaceutical within a single DHB Hospital for information gathering purposes or otherwise related to PHARMAC's decision-making process for considering additions to or amendments to the Pharmaceutical Schedule.
- 17.2 Funding approvals granted under rule 17.1 will be subject to specific limitations on use as determined appropriate by PHARMAC in each circumstance, in consultation with the relevant DHB Hospital and/or DHB.

NATIONAL CONTRACTING

18 Hospital Pharmaceutical Contracts

- 18.1 A DHB Hospital may enter into a contract for the purchase of any Pharmaceutical, including any Medical Device, that it is entitled to fund in accordance with this Schedule H and that is not a National Contract Pharmaceutical, provided that such a contract:
 - a) does not oblige the relevant DHB Hospital to purchase a volume of that Pharmaceutical, if that Pharmaceutical is a DV Pharmaceutical, that is greater than the relevant DV Limit;
 - b) enables PHARMAC to access and use future price and volume data in respect of that Pharmaceutical; and
 - c) enables the relevant DHB Hospital to terminate the contract or relevant parts of the contract in order to give full effect to the National Contract on no more than 3 months' written notice to the Pharmaceutical supplier.
- 18.2 From 1 July 2013, where a DHB Hospital has a pre-existing supply contract for a particular brand of chemical entity for which there is a National Contract Pharmaceutical, the DHB may continue purchasing the chemical entity in accordance with its pre-existing supply contract however:
 - a) from the day its pre-existing supply contract expires, that DHB Hospital is to purchase the relevant National Contract Pharmaceutical listed in Section H at the Price, and is to comply with any DV Limits for the National Contract Pharmaceutical where it has HSS;
 - b) if purchase of the relevant National Contract Pharmaceutical listed in Section H at the Price, where it has HSS, would not cause the relevant DHB Hospital to be in breach of its pre-existing supply contract for a particular brand of chemical entity; the DHB Hospital must purchase the National Contract Pharmaceutical.
- 18.3 Following written notification from PHARMAC that a Pharmaceutical is a National Contract Pharmaceutical, either through Section H updates or otherwise, DHB Hospitals must, unless PHARMAC expressly notifies otherwise:
 - a) take any steps available to them to terminate pre-existing contracts or relevant parts of such a contract, and
 - b) not enter any new contracts or extend the period of any current contracts, for the supply of that National Contract Pharmaceutical or the relevant chemical entity or Medical Device.

19 National Contract Pharmaceuticals

- 19.1 DHB Hospitals must take all necessary steps to enable any contracts between PHARMAC and a Pharmaceutical supplier in relation to National Contract Pharmaceuticals to be given full effect.
- 19.2 The contractual arrangement between PHARMAC and the relevant supplier of a National Contract Pharmaceutical requires it to be made available for purchase at the relevant Price by any or all of the following:
 - a) DHB Hospitals at Designated Delivery Points; and/or
 - b) Contract Manufacturers (expressly for the purpose of compounding).

In the case of Medical Devices, a National Contract may require the Medical Device to be purchased by, and/or supplied to, a third party logistics provider.

20 Hospital Supply Status (HSS)

- 20.1 The DV Limit for any National Contract Pharmaceutical which has HSS is set out in the listing of the relevant National Contract Pharmaceutical in Section H, and may be amended from time to time.
- 20.2 If a National Contract Pharmaceutical is listed in Section H as having HSS, DHB Hospitals:
 - a) are expected to use up any existing stocks of DV Pharmaceuticals during the First Transition Period;
 - b) must not purchase DV Pharmaceuticals in volumes exceeding their usual requirements, or in volumes exceeding those which they reasonably expect to use, within the First Transition Period;
 - c) must ensure that Contract Manufacturers, when manufacturing an Extemporaneously Compounded Product on their behalf, use the National Contract Pharmaceutical with HSS; and

- d) must purchase the National Contract Pharmaceutical with HSS except:
 - i) to the extent that the DHB Hospital may use its discretion to purchase a DV Pharmaceutical within the DV Limit, provided that (subject to rule 20.2(d)(iii) below) the DV Limit has not been exceeded nationally;
 - ii) if the Pharmaceutical supplier fails to supply that National Contract Pharmaceutical, in which case the relevant DHB Hospital does not have to comply with the DV Limit for that National Contract Pharmaceutical during that period of non-supply (and any such month(s) included in a period of non-supply will be excluded in any review of the DV Limit in accordance with rule 20.3 below);
 - iii) that where the DV Limit has been exceeded nationally, the DHB Hospital may negotiate with the Pharmaceutical supplier that supplies the National Contract Pharmaceutical with HSS for written permission to vary the application of that DHB Hospital's Individual DV Limit for any patient whose exceptional needs require a DV Pharmaceutical.

20.3 PHARMAC may, in its discretion, for any period or part period:

- a) review usage by DHB Hospitals of the National Contract Pharmaceutical and DV Pharmaceuticals to determine whether the DV Limit has been exceeded; and
- b) audit compliance by DHB Hospitals with the DV Limits and related requirements.

20.4 PHARMAC will address any issues of non-compliance by any individual DHB or DHB Hospital with a DV Limit by:

- a) obtaining the relevant DHB or DHB Hospital's assurance that it will comply with the DV Limit for that National Contract Pharmaceutical with HSS in the remainder of the applicable period and any subsequent periods; and
- b) informing the relevant supplier of the HSS Pharmaceutical of any individual DHB or DHB Hospital's non-compliance with the DV Limit for that HSS Pharmaceutical.

20.5 In addition to the steps taken by PHARMAC under rule 20.4 above to address any issues of non-compliance by any individual DHB or DHB Hospital with a DV Limit, the relevant Pharmaceutical supplier may require, in its discretion, financial compensation from the relevant DHB or DHB Hospital:

- a) an amount representing that DHB or DHB Hospital's contribution towards exceeding the DV Limit (where PHARMAC is able to quantify this based on the information available to it); or
- b) the sum of \$1,000 or \$5,000 (depending on the terms of the applicable national contract applying to the HSS Pharmaceutical),

whichever is the greater as between sub-paragraphs (a) and (b) within the number of business days specified in the notice from the Pharmaceutical supplier requiring such payment to be made.

21 Collection of rebates and payment of financial compensation

21.1 Following the receipt of any rebates from a Pharmaceutical supplier in respect of a particular National Contract Pharmaceutical, PHARMAC will notify each relevant DHB and DHB Hospital of the amount of the rebate owing to it, being a portion of the total rebate determined by PHARMAC on the basis of that DHB Hospital's usage of that National Contract Pharmaceutical, where this is able to be determined. Where data to determine individual DHB Hospitals' usage is not available, PHARMAC will apportion rebates on the basis of an alternative method agreed between the relevant DHBs and PHARMAC.

21.2 PHARMAC will pay each DHB Hospital the rebate amounts (if any) owing to it, no less frequently than once each calendar quarter in respect of rebates received quarterly (or more often).

22 Price and Volume Data

22.1 DHB Hospitals must provide to PHARMAC, on a monthly basis in accordance with PHARMAC's requirements, any volume data and, unless it would result in a breach of a pre-existing contract, price data held by those DHB Hospitals in respect of any Pharmaceutical (including any Medical Device) listed in Section H.

22.2 All price and volume data provided to PHARMAC under rule 22.1 above should identify the relevant Hospital Pharmaceutical by using a Pharmacode or some other unique numerical identifier, and the date (month and year) on which the DHB Hospital incurred a cost for the purchase of that Hospital Pharmaceutical. Volume is to be measured in units (that being the smallest possible whole Unit – e.g. a capsule, a vial, a millilitre etc).

MISCELLANEOUS PROVISIONS

23 Unapproved Pharmaceuticals

Prescribers should, where possible, prescribe Hospital Pharmaceuticals that are approved under the Medicines Act 1981. However, the funding criteria (including Restrictions) under which a Hospital Pharmaceutical is listed in Section H of the

Schedule may:

- 23.1 in some cases, explicitly permit a DHB to fund a Pharmaceutical that is not approved under the Medicines Act 1981 or for an Unapproved Indication; or

- 23.2 not explicitly prohibit a DHB from funding a Pharmaceutical for use for an Unapproved Indication;

Accordingly, if clinicians are planning on prescribing an unapproved Pharmaceutical or a Pharmaceutical for an Unapproved Indication, they should:

- 23.1 be aware of and comply with their obligations under sections 25 and/or 29 of the Medicines Act 1981, as applicable, and otherwise under that Act and the Medicines Regulations 1984;

- 23.2 be aware of and comply with their obligations under the Health and Disability Commissioner's Code of Consumer Rights, including the requirement to obtain informed consent from the patient (PHARMAC recommends that clinicians obtain written consent); and

- 23.3 exercise their own skill, judgment, expertise and discretion, and make their own prescribing decisions with respect to the use of an unapproved Pharmaceutical or a Pharmaceutical for an Unapproved Indication.

Clinicians should be aware that simply by listing a Pharmaceutical on the Pharmaceutical Schedule, PHARMAC makes no representations about whether that Pharmaceutical has any form of approval or consent under, or whether the supply or use of the Pharmaceutical otherwise complies with, the Medicines Act 1981. Further, the Pharmaceutical Schedule does not constitute an advertisement, advertising material or a medical advertisement as defined in the Medicines Act or otherwise.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Antacids and Antiflatulents			
Antacids and Reflux Barrier Agents			
ALUMINIUM HYDROXIDE WITH MAGNESIUM HYDROXIDE AND SIMETHICONE			
Tab 200 mg with magnesium hydroxide 200 mg and simethicone 20 mg			<i>e.g. Mylanta</i>
Oral liq 200 mg with magnesium hydroxide 200 mg and simethicone 20 mg per 5 ml			<i>e.g. Mylanta</i>
Oral liq 400 mg with magnesium hydroxide 400 mg and simethicone 30 mg per 5 ml			<i>e.g. Mylanta Double Strength</i>
SIMETHICONE			
Oral drops 100 mg per ml			
SODIUM ALGINATE WITH MAGNESIUM ALGINATE			
Powder for oral soln 225 mg with magnesium alginate 87.5 mg, sachet			<i>e.g. Gaviscon Infant</i>
SODIUM ALGINATE WITH SODIUM BICARBONATE AND CALCIUM CARBONATE			
Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg			<i>e.g. Gaviscon Double Strength</i>
Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml	4.95	500 ml	Acidex
SODIUM CITRATE			
Oral liq 8.8% (300 mmol/l)			
Phosphate Binding Agents			
ALUMINIUM HYDROXIDE			
Tab 600 mg			
CALCIUM CARBONATE – Restricted see terms below			
☿ Oral liq 250 mg per ml (100 mg elemental per ml)	39.00	500 ml	Roxane
☞ Restricted			
Initiation			
Only for use in children under 12 years of age for use as a phosphate binding agent.			
Antidiarrhoeals and Intestinal Anti-Inflammatory Agents			
Antipropulsives			
DIPHENOXYLATE HYDROCHLORIDE WITH ATROPINE SULPHATE			
Tab 2.5 mg with atropine sulphate 25 mcg			
LOPERAMIDE HYDROCHLORIDE			
Tab 2 mg			
Cap 2 mg – 1% DV Jul-14 to 2016	7.84	400	Diamide Relief
Rectal and Colonic Anti-Inflammatories			
BUDESONIDE – Restricted see terms on the next page			
☿ Cap 3 mg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➡Restricted			
Initiation — Crohn's disease			
Both:			
1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and			
2 Any of the following:			
2.1 Diabetes; or			
2.2 Cushingoid habitus; or			
2.3 Osteoporosis where there is significant risk of fracture; or			
2.4 Severe acne following treatment with conventional corticosteroid therapy; or			
2.5 History of severe psychiatric problems associated with corticosteroid treatment; or			
2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or			
2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).			
Initiation — Collagenous and lymphocytic colitis (microscopic colitis)			
Patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.			
Initiation — Gut Graft versus Host disease			
Patient has gut Graft versus Host disease following allogenic bone marrow transplantation.			
HYDROCORTISONE ACETATE			
Rectal foam 10%, CFC free (14 applications) – 1% DV Oct-15 to 2018.....	26.55	21.1 g	Colifoam
MESALAZINE			
Tab EC 400 mg	49.50	100	Asacol
Tab EC 500 mg	49.50	100	Asamax
Tab long-acting 500 mg	59.05	100	Pentasa
Tab 800 mg	85.55	90	Asacol
Modified release granules 1 g	141.72	120 g	Pentasa
Suppos 500 mg	22.80	20	Asacol
Suppos 1 g – 1% DV Jun-15 to 2018	54.60	30	Pentasa
Enema 1 g per 100 ml – 1% DV Sep-15 to 2018	41.30	7	Pentasa
OLSALAZINE			
Tab 500 mg			
Cap 250 mg			
SODIUM CROMOGLYCATE			
Cap 100 mg			
SULPHASALAZINE			
Tab 500 mg – 1% DV Oct-13 to 2016	11.68	100	Salazopyrin
Tab EC 500 mg – 1% DV Oct-13 to 2016	12.89	100	Salazopyrin EN
Local Preparations for Anal and Rectal Disorders			
Antihæmorrhoidal Preparations			
CINCHOCAINE HYDROCHLORIDE WITH HYDROCORTISONE			
Oint 5 mg with hydrocortisone 5 mg per g	15.00	30 g	Proctosedyl
Suppos 5 mg with hydrocortisone 5 mg per g	9.90	12	Proctosedyl
FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE			
Oint 950 mcg with fluocortolone pivalate 920 mcg and cinchocaine hydrochloride 5 mg per g	6.35	30 g	Ultraproct
Suppos 630 mcg with fluocortolone pivalate 610 mcg and cinchocaine hydrochloride 1 mg	2.66	12	Ultraproct

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Management of Anal Fissures			
GLYCERYL TRINITRATE			
Oint 0.2%	22.00	30 g	Rectogesic
Rectal Sclerosants			
OILY PHENOL [PHENOL OILY]			
Inj 5%, 5 ml vial			
Antispasmodics and Other Agents Altering Gut Motility			
GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – 1% DV Oct-13 to 2016	17.14	10	Max Health
Max Health brand - HSS with 1% DV will apply 1 July 2016 to 30 June 2019.			
HYOSCINE BUTYLBROMIDE			
Tab 10 mg	2.18	20	Gastrosoothe
Inj 20 mg, 1 ml ampoule	9.57	5	Buscopan
MEBEVERINE HYDROCHLORIDE			
Tab 135 mg – 1% DV Sep-14 to 2017	18.00	90	Colofac
Antiulcerants			
Antisecretory and Cytoprotective			
MISOPROSTOL			
Tab 200 mcg – 1% DV Jun-16 to 2019	41.50	120	Cytotec
H2 Antagonists			
CIMETIDINE			
Tab 200 mg			
Tab 400 mg			
RANITIDINE			
Tab 150 mg – 1% DV Nov-14 to 2017	10.30	500	Ranitidine Relief
Tab 300 mg – 1% DV Nov-14 to 2017	14.73	500	Ranitidine Relief
Oral liq 150 mg per 10 ml – 1% DV Sep-14 to 2017	4.92	300 ml	Peptisoothe
Inj 25 mg per ml, 2 ml ampoule	8.75	5	Zantac
Proton Pump Inhibitors			
LANSOPRAZOLE			
Cap 15 mg – 1% DV Jan-16 to 2018	5.08	100	Lanzol Relief
Cap 30 mg – 1% DV Jan-16 to 2018	5.93	100	Lanzol Relief
OMEPRAZOLE			
⚡ Tab dispersible 20 mg			
➡ Restricted			
Initiation			
Only for use in tube-fed patients.			
Cap 10 mg – 1% DV Jan-15 to 2017	2.23	90	Omezol Relief
Cap 20 mg – 1% DV Jan-15 to 2017	2.91	90	Omezol Relief
Cap 40 mg – 1% DV Jan-15 to 2017	4.42	90	Omezol Relief
Powder for oral liq	42.50	5 g	Midwest

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Inj 40 mg ampoule	19.00	5	Dr Reddy's Omeprazole
Inj 40 mg ampoule with diluent	33.65	5	Dr Reddy's Omeprazole
PANTOPRAZOLE			
Tab EC 20 mg – 1% DV May-14 to 2016	2.68	100	Pantoprazole Actavis 20
Tab EC 40 mg – 1% DV May-14 to 2016	3.54	100	Pantoprazole Actavis 40
Inj 40 mg vial			

Site Protective Agents

BISMUTH TRIOXIDE			
Tab 120 mg	32.50	112	De-Nol
SUCRALFATE			
Tab 1 g			

Bile and Liver Therapy

L-ORNITHINE L-ASPARTATE – **Restricted** see terms below

⚡ Grans for oral liquid 3 g

➡ **Restricted**

Initiation

For patients with chronic hepatic encephalopathy who have not responded to treatment with, or are intolerant to lactulose, or where lactulose is contraindicated.

RIFAXIMIN – **Restricted** see terms below

⚡ Tab 550 mg – 1% DV Oct-14 to 2017 625.00 56 **Xifaxan**

➡ **Restricted**

Initiation

For patients with hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

Diabetes

Alpha Glucosidase Inhibitors

ACARBOSE			
Tab 50 mg – 1% DV Oct-15 to 2018	4.28	90	Glucobay
Tab 100 mg – 1% DV Oct-15 to 2018	7.78	90	Glucobay

Hyperglycaemic Agents

DIAZOXIDE – **Restricted** see terms below

⚡ Cap 25 mg	110.00	100	Proglidem
⚡ Cap 100 mg	280.00	100	Proglidem
⚡ Oral liq 50 mg per ml	620.00	30 ml	Proglycem

➡ **Restricted**

Initiation

For patients with confirmed hypoglycaemia caused by hyperinsulinism.

GLUCAGON HYDROCHLORIDE

Inj 1 mg syringe kit 32.00 1 Glucagen Hypokit

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GLUCOSE [DEXTROSE]			
Tab 1.5 g			
Tab 3.1 g			
Tab 4 g			
Gel 40%			
GLUCOSE WITH SUCROSE AND FRUCTOSE			
Gel 19.7% with sucrose 35% and fructose 19.7%, 18 g sachet			
Insulin - Intermediate-Acting Preparations			
INSULIN ASPART WITH INSULIN ASPART PROTAMINE			
Inj insulin aspart 30% with insulin aspart protamine 70%, 100 u per ml, 3 ml prefilled pen	52.15	5	NovoMix 30 FlexPen
INSULIN ISOPHANE			
Inj insulin human 100 u per ml, 10 ml vial			
Inj insulin human 100 u per ml, 3 ml cartridge			
INSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
Inj insulin lispro 25% with insulin lispro protamine 75%, 100 u per ml, 3 ml cartridge	42.66	5	Humalog Mix 25
Inj insulin lispro 50% with insulin lispro protamine 50%, 100 u per ml, 3 ml cartridge	42.66	5	Humalog Mix 50
INSULIN NEUTRAL WITH INSULIN ISOPHANE			
Inj insulin neutral 30% with insulin isophane 70%, 100 u per ml, 10 ml vial			
Inj insulin neutral 30% with insulin isophane 70%, 100 u per ml, 3 ml cartridge			
Inj insulin neutral 40% with insulin isophane 60%, 100 u per ml, 3 ml cartridge			
Inj insulin neutral 50% with insulin isophane 50%, 100 u per ml, 3 ml cartridge			
Insulin - Long-Acting Preparations			
INSULIN GLARGINE			
Inj 100 u per ml, 3 ml disposable pen	94.50	5	Lantus SoloStar
Inj 100 u per ml, 3 ml cartridge	94.50	5	Lantus
Inj 100 u per ml, 10 ml vial	63.00	1	Lantus
Insulin - Rapid-Acting Preparations			
INSULIN ASPART			
Inj 100 u per ml, 10 ml vial			
Inj 100 u per ml, 3 ml cartridge			
Inj 100 u per ml, 3 ml syringe	51.19	5	NovoRapid FlexPen
INSULIN GLULISINE			
Inj 100 u per ml, 10 ml vial	27.03	1	Apidra
Inj 100 u per ml, 3 ml cartridge	46.07	5	Apidra
Inj 100 u per ml, 3 ml disposable pen	46.07	5	Apidra Solostar
INSULIN LISPRO			
Inj 100 u per ml, 10 ml vial			
Inj 100 u per ml, 3 ml cartridge			

Products with Hospital Supply Status (HSS) are in **bold**

Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Insulin - Short-Acting Preparations			
INSULIN NEUTRAL			
Inj human 100 u per ml, 10 ml vial			
Inj human 100 u per ml, 3 ml cartridge			
Oral Hypoglycaemic Agents			
GLIBENCLAMIDE			
Tab 5 mg			
GLICLAZIDE			
Tab 80 mg – 1% DV Nov-14 to 2017	11.50	500	Glizide
GLIPIZIDE			
Tab 5 mg – 1% DV Sep-15 to 2018	2.85	100	Minidiab
METFORMIN HYDROCHLORIDE			
Tab immediate-release 500 mg – 1% DV Nov-15 to 2018	9.59	1,000	Metckek
Tab immediate-release 850 mg – 1% DV Dec-15 to 2018	7.82	500	Metformin Mylan
PIOGLITAZONE			
Tab 15 mg – 1% DV Dec-15 to 2018	3.47	90	Vexazone
Tab 30 mg – 1% DV Dec-15 to 2018	5.06	90	Vexazone
Tab 45 mg – 1% DV Dec-15 to 2018	7.10	90	Vexazone
Digestives Including Enzymes			
PANCREATIC ENZYME			
Cap EC 10,000 BP u lipase, 9,000 BP u amylase and 210 BP u pro- tease – 1% DV Oct-15 to 2018	34.93	100	Creon 10000
Cap EC 25,000 BP u lipase, 18,000 BP u amylase and 1,000 BP u protease – 1% DV Oct-15 to 2018	94.38	100	Creon 25000
Cap EC 25,000 BP u lipase, 22,500 BP u amylase and 1,250 BP u protease			
Powder 25,000 u lipase with 30,000 u amylase and 1,400 u protease per g			
URSODEOXYCHOLIC ACID – Restricted see terms below			
⚡ Cap 250 mg – 1% DV Sep-14 to 2017	53.40	100	Ursosan
➡ Restricted			
Initiation — Alagille syndrome or progressive familial intrahepatic cholestasis			
Either:			
1 Patient has been diagnosed with Alagille syndrome; or			
2 Patient has progressive familial intrahepatic cholestasis.			
Initiation — Chronic severe drug induced cholestatic liver injury			
All of the following:			
1 Patient has chronic severe drug induced cholestatic liver injury; and			
2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and			
3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.			
Initiation — Cirrhosis			
Both:			
1 Primary biliary cirrhosis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative by liver biopsy; and			

continued...

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

continued...

- 2 Patient not requiring a liver transplant (bilirubin > 100 $\mu\text{mol/l}$; decompensated cirrhosis).

Initiation — Pregnancy

Patient diagnosed with cholestasis of pregnancy.

Initiation — Haematological transplant

- Both:
- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogeneic stem cell or bone marrow transplantation; and
 - 2 Treatment for up to 13 weeks.

Initiation — Total parenteral nutrition induced cholestasis

- Both:
- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by TPN; and
 - 2 Liver function has not improved with modifying the TPN composition.

Laxatives

Bowel-Cleansing Preparations

CITRIC ACID WITH MAGNESIUM OXIDE AND SODIUM PICOSULFATE

Powder for oral soln 12 g with magnesium oxide 3.5 g and sodium picosulfate 10 mg per sachet *e.g. PicoPrep*

MACROGOL 3350 WITH ASCORBIC ACID, POTASSIUM CHLORIDE AND SODIUM CHLORIDE

Powder for oral soln 755.68 mg with ascorbic acid 85.16 mg, potassium chloride 10.55 mg, sodium chloride 37.33 mg and sodium sulphate 80.62 mg per g, 210 g sachet *e.g. Glycoprep-C*

Powder for oral soln 755.68 mg with ascorbic acid 85.16 mg, potassium chloride 10.55 mg, sodium chloride 37.33 mg and sodium sulphate 80.62 mg per g, 70 g sachet *e.g. Glycoprep-C*

MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE, SODIUM CHLORIDE AND SODIUM SULPHATE

Powder for oral soln 59 g with potassium chloride 0.7425 g, sodium bicarbonate 1.685 g, sodium chloride 1.465 g and sodium sulphate 5.685 g per sachet 14.31 4 Klean Prep

Bulk-Forming Agents

ISPAGHULA (PSYLLIUM) HUSK

Powder for oral soln – 1% DV **Sep-13 to 2016**.....5.51 500 g **Konsyl-D**

STERCULIA WITH FRANGULA – **Restricted:** For continuation only

➔ Powder for oral soln

Faecal Softeners

DOCUSATE SODIUM

Tab 50 mg – 1% DV **Jan-15 to 2017**.....2.31 100 **Coloxyl**
 Tab 120 mg – 1% DV **Jan-15 to 2017**.....3.13 100 **Coloxyl**

DOCUSATE SODIUM WITH SENNOSIDES

Tab 50 mg with sennosides 8 mg4.40 200 Laxsol

PARAFFIN

Oral liquid 1 mg per ml
 Enema 133 ml

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
POLOXAMER			
Oral drops 10% – 1% DV Sep-14 to 2017	3.78	30 ml	Coloxyl
Osmotic Laxatives			
GLYCEROL			
Suppos 1.27 g			
Suppos 2.55 g			
Suppos 3.6 g – 1% DV Sep-15 to 2018	6.50	20	PSM
LACTULOSE			
Oral liq 10 g per 15 ml – 1% DV May-14 to 2016	3.84	500 ml	Laevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE AND SODIUM CHLORIDE – Restricted see terms below			
⚡ Powder for oral soln 6.563 g with potassium chloride 23.3 mg, sodium bicarbonate 89.3 mg and sodium chloride 175.4 mg			
⚡ Powder for oral soln 13.125 g with potassium chloride 46.6 mg, sodium bicarbonate 178.5 mg and sodium chloride 350.7 mg – 1% DV Oct-14 to 2017	7.65	30	Lax-Sachets
➡ Restricted			
Initiation			
Either:			
1 Both:			
1.1 The patient has problematic constipation despite an adequate trial of other oral pharmacotherapies including lactulose where lactulose is not contraindicated; and			
1.2 The patient would otherwise require a per rectal preparation; or			
2 For short-term use for faecal disimpaction.			
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE			
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml – 1% DV Sep-13 to 2016	19.95	50	Micolette
SODIUM PHOSPHATE WITH PHOSPHORIC ACID			
Oral liq 16.4% with phosphoric acid 25.14%			
Enema 10% with phosphoric acid 6.58%	2.50	1	Fleet Phosphate Enema
Stimulant Laxatives			
BISACODYL			
Tab 5 mg – 1% DV Oct-15 to 2018	5.99	200	Lax-Tabs
Suppos 10 mg – 1% DV Jan-16 to 2018	3.78	10	Lax-Suppositories
SENNOSIDES			
Tab 7.5 mg			
Metabolic Disorder Agents			
ARGININE			
Powder			
Inj 600 mg per ml, 25 ml vial			
BETAINE – Restricted see terms below			
⚡ Powder			
➡ Restricted			
Metabolic physician or metabolic disorders dietitian			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BIOTIN – Restricted see terms below			
¶ Cap 50 mg			
¶ Cap 100 mg			
¶ Inj 10 mg per ml, 5 ml vial			
➔ Restricted			
Metabolic physician or metabolic disorders dietitian			
HAEM ARGINATE			
Inj 25 mg per ml, 10 ml ampoule			
IMIGLUCERASE – Restricted see terms below			
¶ Inj 40 iu per ml, 5 ml vial			
¶ Inj 40 iu per ml, 10 ml vial			
➔ Restricted			
Initiation			
Only for use in patients with approval by the Gaucher's Treatment Panel.			
LEVOCARNITINE – Restricted see terms below			
¶ Cap 500 mg			
¶ Oral soln 1,100 mg per 15 ml			
¶ Inj 200 mg per ml, 5 ml vial			
➔ Restricted			
Neurologist, metabolic physician or metabolic disorders dietitian			
PYRIDOXAL-5-PHOSPHATE – Restricted see terms below			
¶ Tab 50 mg			
➔ Restricted			
Neurologist, metabolic physician or metabolic disorders dietitian			
SODIUM BENZOATE			
Cap 500 mg			
Powder			
Soln 100 mg per ml			
Inj 20%, 10 ml ampoule			
SODIUM PHENYLBUTYRATE			
Tab 500 mg			
Oral liq 250 mg per ml			
Inj 200 mg per ml, 10 ml ampoule			
TRIENTINE DIHYDROCHLORIDE			
Cap 300 mg			

Minerals

Calcium

CALCIUM CARBONATE

Tab 1.25 g (500 mg elemental) – 1% DV Sep-14 to 2017	5.38	250	Arrow-Calcium
Tab eff 1.75 g (1 g elemental)	6.21	30	Calsource

Fluoride

SODIUM FLUORIDE

Tab 1.1 mg (0.5 mg elemental)

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Iodine			
POTASSIUM IODATE			
Tab 253 mcg (150 mcg elemental iodine) – 1% DV Dec-14 to 2017	3.65	90	NeuroTabs
POTASSIUM IODATE WITH IODINE			
Oral liq 10% with iodine 5%			
Iron			
FERRIC CARBOXYMALTOSE – Restricted see terms below			
☯ Inj 50 mg per ml, 10 ml vial	150.00	1	Ferinject
☛ Restricted			
Initiation			
Treatment with oral iron has proven ineffective or is clinically inappropriate.			
FERROUS FUMARATE			
Tab 200 mg (65 mg elemental) – 1% DV Jun-15 to 2018	2.89	100	Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID			
Tab 310 mg (100 mg elemental) with folic acid 350 mcg	4.75	60	Ferro-F-Tabs
FERROUS GLUCONATE WITH ASCORBIC ACID			
Tab 170 mg (20 mg elemental) with ascorbic acid 40 mg			
FERROUS SULPHATE			
Tab long-acting 325 mg (105 mg elemental)	2.06	30	Ferrograd
Oral liq 30 mg (6 mg elemental) per ml – 1% DV Apr-14 to 2016	10.28	500 ml	Ferodan
FERROUS SULPHATE WITH ASCORBIC ACID			
Tab long-acting 325 mg (105 mg elemental) with ascorbic acid 500 mg			
FERROUS SULPHATE WITH FOLIC ACID			
Tab long-acting 325 mg (105 mg elemental) with folic acid 350 mcg			
IRON POLYMALTOSE			
Inj 50 mg per ml, 2 ml ampoule – 1% DV Sep-14 to 2017	15.22	5	Ferrum H
IRON SUCROSE			
Inj 20 mg per ml, 5 ml ampoule	100.00	5	Venofer
Magnesium			
MAGNESIUM HYDROXIDE			
Tab 311 mg (130 mg elemental)			
MAGNESIUM OXIDE			
Cap 663 mg (400 mg elemental)			
MAGNESIUM SULPHATE			
Inj 0.4 mmol per ml, 250 ml bag			
Inj 2 mmol per ml, 5 ml ampoule – 1% DV Oct-14 to 2017	12.65	10	DBL
Zinc			
ZINC			
Oral liq 5 mg per 5 drops			
ZINC CHLORIDE			
Inj 5.3 mg per ml (5.1 mg per ml elemental), 2 ml ampoule			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ZINC SULPHATE			
Cap 137.4 mg (50 mg elemental) – 1% DV Mar-15 to 2017	11.00	100	Zincaps
Mouth and Throat			
Agents Used in Mouth Ulceration			
BENZYLAMINE HYDROCHLORIDE			
Soln 0.15%			
Spray 0.15%			
Spray 0.3%			
BENZYLAMINE HYDROCHLORIDE WITH CETYLPYRIDINIUM CHLORIDE			
Lozenge 3 mg with cetylpyridinium chloride			
CARBOXYMETHYLCELLULOSE			
Oral spray			
CHLORHEXIDINE GLUCONATE			
Mouthwash 0.2% – 1% DV Sep-15 to 2018	2.57	200 ml	healthE
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE			
Adhesive gel 8.7% with cetalkonium chloride 0.01%			
DICHLOROBENZYL ALCOHOL WITH AMYLMETACRESOL			
Lozenge 1.2 mg with amylmetacresol 0.6 mg			
SODIUM CARBOXYMETHYLCELLULOSE WITH PECTIN AND GELATINE			
Paste			
Powder			
TRIAMCINOLONE ACETONIDE			
Paste 0.1% – 1% DV Apr-15 to 2017	5.33	5 g	Kenalog in Orabase
Oropharyngeal Anti-Infectives			
AMPHOTERICIN B			
Lozenge 10 mg	5.86	20	Fungilin
MICONAZOLE			
Oral gel 20 mg per g – 1% DV Sep-15 to 2018	4.79	40 g	Decozol
NYSTATIN			
Oral liquid 100,000 u per ml – 1% DV Feb-16 to 2017	2.55	24 ml	m-Nystatin
Other Oral Agents			
SODIUM HYALURONATE [HYALURONIC ACID] – Restricted see terms below			
⚡ Inj 20 mg per ml, 1 ml syringe			
➡ Restricted			
Otolaryngologist			
THYMOL GLYCERIN			
Compound, BPC			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Vitamins			
Multivitamin Preparations			
MULTIVITAMIN AND MINERAL SUPPLEMENT – Restricted see terms below			
☞ Cap	23.35	180	Clinicians Multivit & Mineral Boost
☞ Restricted			
Initiation			
<i>Limited to 3 months treatment</i>			
Both:			
1 Patient was admitted to hospital with burns; and			
2 Any of the following:			
2.1 Burn size is greater than 15% of total body surface area (BSA) for all types of burns; or			
2.2 Burn size is greater than 10% of BSA for mid-dermal or deep dermal burns; or			
2.3 Nutritional status prior to admission or dietary intake is poor.			
MULTIVITAMIN RENAL – Restricted see terms below			
☞ Cap	8.39	30	Clinicians Renal Vit
☞ Restricted			
Initiation			
Either:			
1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or			
2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73m ² body surface area (BSA).			
MULTIVITAMINS			
Tab (BPC cap strength)			<i>e.g. Mvite</i>
☞ Cap vitamin A 2500 u, betacarotene 3 mg, cholecalciferol 11 mcg, alpha tocopherol 150 u, phytomenadione 150 mcg, folic acid 0.2 mg, ascorbic acid 100 mg, thiamine 1.5 mg, pantothenic acid 12 mg, riboflavin 1.7 mg, niacin 20 mg, pyridoxine hydrochloride 1.9 mg, cyanocobalamin 3 mcg, zinc 7.5 mg and biotin 100 mcg			<i>e.g. Vitabdeck</i>
☞ Restricted			
Initiation			
Either:			
1 Patient has cystic fibrosis with pancreatic insufficiency; or			
2 Patient is an infant or child with liver disease or short gut syndrome.			
☞ Powder vitamin A 4200 mcg with vitamin D 155.5 mcg, vitamin E 21.4 mg, vitamin C 400 mg, vitamin K1 166 mcg thiamine 3.2 mg, riboflavin 4.4 mg, niacin 35 mg, vitamin B6 3.4 mg, folic acid 303 mcg, vitamin B12 8.6 mcg, biotin 214 mcg, pantothenic acid 17 mg, choline 350 mg and inositol 700 mg			<i>e.g. Paediatric Seravit</i>
☞ Restricted			
Initiation			
Patient has inborn errors of metabolism.			
Inj thiamine hydrochloride 250 mg with riboflavin 4 mg and pyridoxine hydrochloride 50 mg, 5 ml ampoule (1) and inj ascorbic acid 500 mg with nicotinamide 160 mg and glucose 1000 mg, 5 ml ampoule (1)			<i>e.g. Pabrinex IV</i>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Inj thiamine hydrochloride 250 mg with riboflavin 4 mg and pyridoxine hydrochloride 50 mg, 5 ml ampoule (1) and inj ascorbic acid 500 mg with nicotinamide 160 mg, 2 ml ampoule (1)			<i>e.g. Pabrinex IM</i>
Inj thiamine hydrochloride 500 mg with riboflavin 8 mg and pyridoxine hydrochloride 100 mg, 10 ml ampoule (1) and inj ascorbic acid 1000 mg with nicotinamide 320 mg and glucose 2000 mg, 10 ml ampoule (1)			<i>e.g. Pabrinex IV</i>
VITAMIN A WITH VITAMINS D AND C			
Soln 1,000 u with vitamin D 400 u and ascorbic acid 30 mg per 10 drops			<i>e.g. Vitadol C</i>

Vitamin A

RETINOL

Tab 10,000 iu
Cap 25,000 iu
Oral liq 150,000 iu per ml

Vitamin B

HYDROXOCOBALAMIN

Inj 1 mg per ml, 1 ml ampoule – 1% DV Sep-15 to 20182.31 3 **Neo-B12**

PYRIDOXINE HYDROCHLORIDE

Tab 25 mg – 1% DV Apr-15 to 20172.15 90 **Vitamin B6 25**
Tab 50 mg – 1% DV Oct-14 to 201711.55 500 **Apo-Pyridoxine**
Inj 100 mg per ml, 1 ml ampoule

THIAMINE HYDROCHLORIDE

Tab 50 mg
Tab 100 mg
Inj 100 mg per ml, 1 ml vial
Inj 100 mg per ml, 2 ml vial *e.g. Benerva*

VITAMIN B COMPLEX

Tab strong, BPC

Vitamin C

ASCORBIC ACID

Tab 100 mg – 1% DV Nov-13 to 20167.00 500 **Cvite**
Tab chewable 250 mg

Vitamin D

ALFACALCIDOL

Cap 0.25 mcg26.32 100 One-Alpha
Cap 1 mcg87.98 100 One-Alpha
Oral drops 2 mcg per ml

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
CALCITRIOL			
Cap 0.25 mcg	3.03	30	Airflow
	10.10	100	Calcitriol-AFT
Cap 0.5 mcg	5.62	30	Airflow
	18.73	100	Calcitriol-AFT
Oral liq 1 mcg per ml			
Inj 1 mcg per ml, 1 ml ampoule			
CHOLECALCIFEROL			
Cap 1.25 mg (50,000 iu)	3.85	12	Vit.D3

Vitamin E

ALPHA TOCOPHERYL ACETATE – **Restricted** see terms below

- ¶ Cap 100 u
- ¶ Cap 500 u
- ¶ Oral liq 156 u per ml

➡ **Restricted**

Initiation — Cystic fibrosis

Both:

- 1 Cystic fibrosis patient; and
- 2 Either:
 - 2.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck); or
 - 2.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for the patient.

Initiation — Osteoradionecrosis

For the treatment of osteoradionecrosis.

Initiation — Other indications

All of the following:

- 1 Infant or child with liver disease or short gut syndrome; and
- 2 Requires vitamin supplementation; and
- 3 Either:
 - 3.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplements (Vitabdeck); or
 - 3.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for patient.

Price (ex man. excl. GST)	Brand or Generic
\$	Per Manufacturer

Antianaemics

Hypoplastic and Haemolytic

EPOETIN ALFA [ERYTHROPOIETIN ALFA] – **Restricted** see terms below

⚡ Inj 1,000 iu in 0.5 ml syringe – 5% DV Mar-15 to 28 Feb 2018	48.68	6	Eprex
⚡ Inj 2,000 iu in 0.5 ml syringe – 5% DV Mar-15 to 28 Feb 2018	120.18	6	Eprex
⚡ Inj 3,000 iu in 0.3 ml syringe – 5% DV Mar-15 to 28 Feb 2018	166.87	6	Eprex
⚡ Inj 4,000 iu in 0.4 ml syringe – 5% DV Mar-15 to 28 Feb 2018	193.13	6	Eprex
⚡ Inj 5,000 iu in 0.5 ml syringe – 5% DV Mar-15 to 28 Feb 2018	243.26	6	Eprex
⚡ Inj 6,000 iu in 0.6 ml syringe – 5% DV Mar-15 to 28 Feb 2018	291.92	6	Eprex
⚡ Inj 8,000 iu in 0.8 ml syringe – 5% DV May-15 to 28 Feb 2018	352.69	6	Eprex
⚡ Inj 10,000 iu in 1 ml syringe – 5% DV Mar-15 to 28 Feb 2018	395.18	6	Eprex
⚡ Inj 40,000 iu in 1 ml syringe – 5% DV May-15 to 28 Feb 2018	263.45	1	Eprex

→ Restricted

Initiation — chronic renal failure

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin \leq 100g/L; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate \leq 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate \leq 45ml/min; and
- 4 Patient is on haemodialysis or peritoneal dialysis.

Initiation — myelodysplasia*

Re-assessment required after 2 months

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS); and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification-based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum erythropoietin level of < 500 IU/L; and
- 6 The minimum necessary dose of erythropoietin would be used and will not exceed 80,000 iu per week.

Continuation — myelodysplasia*

Re-assessment required after 12 months

All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of erythropoietin would be used and will not exceed 80,000 iu per week.

Initiation — all other indications

Haematologist

For use in patients where blood transfusion is not a viable treatment alternative.

Note: Indications marked with * are Unapproved Indications

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

EPOETIN BETA [ERYTHROPOIETIN BETA] – **Restricted** see terms below

Note: Epoetin beta is considered a Discretionary Variance Pharmaceutical for epoetin alfa.

- ⚡ Inj 2,000 iu in 0.3 ml syringe
- ⚡ Inj 3,000 iu in 0.3 ml syringe
- ⚡ Inj 4,000 iu in 0.3 ml syringe
- ⚡ Inj 5,000 iu in 0.3 ml syringe
- ⚡ Inj 6,000 iu in 0.3 ml syringe
- ⚡ Inj 10,000 iu in 0.6 ml syringe

➡ **Restricted**

Initiation — chronic renal failure

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin \leq 100g/L; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate \leq 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate \leq 45ml/min; and
- 4 Patient is on haemodialysis or peritoneal dialysis.

Initiation — myelodysplasia*

Re-assessment required after 12 months

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS); and
- 2 Has had symptomatic anaemia with haemoglobin $<$ 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification-based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum erythropoietin level of $<$ 500 IU/L; and
- 6 The minimum necessary dose of erythropoietin would be used and will not exceed 80,000 iu per week.

Continuation — myelodysplasia*

Re-assessment required after 2 months

All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of erythropoietin would be used and will not exceed 80,000 iu per week.

Initiation — all other indications

Haematologist.

For use in patients where blood transfusion is not a viable treatment alternative.

*Note: Indications marked with * are Unapproved Indications.

Megaloblastic

FOLIC ACID

Tab 0.8 mg – 1% DV Oct-15 to 2018	20.60	1,000	Apo-Folic Acid
Tab 5 mg – 1% DV Oct-15 to 2018	10.92	500	Apo-Folic Acid
Oral liq 50 mcg per ml	24.00	25 ml	Biomed
Inj 5 mg per ml, 10 ml vial			

Price (ex man. excl. GST)	Brand or Generic
\$	Per Manufacturer

Antifibrinolytics, Haemostatics and Local Sclerosants

ALUMINIUM CHLORIDE – **Restricted** see terms below

☞ Topical soln 20% w/v

e.g. Driclor

➔Restricted

Initiation

For use as a haemostatis agent.

APROTININ – **Restricted** see terms below

☞ Inj 10,000 kIU per ml (equivalent to 200 mg per ml), 50 ml vial

➔Restricted

Initiation

Cardiac anaesthetist

Either:

- 1 Paediatric patient undergoing cardiopulmonary bypass procedure; or
- 2 Adult patient undergoing cardiac surgical procedure where the significant risk of massive bleeding outweighs the potential adverse effects of the drug.

ELTROMBOPAG – **Restricted** see terms below

☞ Tab 25 mg	1,771.00	28	Revolade
☞ Tab 50 mg	3,542.00	28	Revolade

➔Restricted

Initiation — idiopathic thrombocytopenic purpura - post-splenectomy

Haematologist

Limited to 6 weeks treatment

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of \leq 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of \leq 10,000 platelets per microlitre.

Initiation — (idiopathic thrombocytopenic purpura - preparation for splenectomy)

Haematologist

Limited to 6 weeks treatment

The patient requires eltrombopag treatment as preparation for splenectomy.

Continuation — (idiopathic thrombocytopenic purpura - post-splenectomy)

Haematologist

Re-assessment required after 12 months

The patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of $>$ 30,000 platelets per microlitre

FERRIC SUBSULFATE

Gel 25.9%

Soln 500 ml

POLIDOCANOL

Inj 0.5%, 30 ml vial

SODIUM TETRADECYL SULPHATE

Inj 3%, 2 ml ampoule

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
THROMBIN Powder			
TRANEXAMIC ACID			
Tab 500 mg – 1% DV Oct-14 to 2016	23.00	100	Cyklokapron
Inj 100 mg per ml, 5 ml ampoule – 1% DV Sep-15 to 2018	55.00	10	Cyklokapron

Blood Factors

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] – **Restricted** see terms below

⚡ Inj 1 mg syringe	1,163.75	1	NovoSeven RT
⚡ Inj 2 mg syringe	2,327.50	1	NovoSeven RT
⚡ Inj 5 mg syringe	5,818.75	1	NovoSeven RT
⚡ Inj 8 mg syringe	9,310.00	1	NovoSeven RT

➡ **Restricted**

Initiation

When used in the treatment of haemophilia, access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

FACTOR EIGHT INHIBITOR BYPASSING FRACTION – **Restricted** see terms below

⚡ Inj 500 U	1,450.00	1	FEIBA NF
⚡ Inj 1,000 U	2,900.00	1	FEIBA NF
⚡ Inj 2,500 U	7,250.00	1	FEIBA NF

➡ **Restricted**

Initiation

When used in the treatment of haemophilia, access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] – **Restricted** see terms below

⚡ Inj 250 iu prefilled syringe	210.00	1	Xyntha
⚡ Inj 500 iu prefilled syringe	420.00	1	Xyntha
⚡ Inj 1,000 iu prefilled syringe	840.00	1	Xyntha
⚡ Inj 2,000 iu prefilled syringe	1,680.00	1	Xyntha
⚡ Inj 3,000 iu prefilled syringe	2,520.00	1	Xyntha

➡ **Restricted**

Initiation

Note: Preferred Brand of recombinant factor VIII from 1 March 2016 until 28 February 2019. When used in the treatment of haemophilia, funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

NONACOG ALFA [RECOMBINANT FACTOR IX] – **Restricted** see terms below

⚡ Inj 250 iu vial	310.00	1	BeneFIX
⚡ Inj 500 iu vial	620.00	1	BeneFIX
⚡ Inj 1,000 iu vial	1,240.00	1	BeneFIX
⚡ Inj 2,000 iu vial	2,480.00	1	BeneFIX
⚡ Inj 3,000 iu vial	3,720.00	1	BeneFIX

➡ **Restricted**

Initiation

When used in the treatment of haemophilia, access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
NONACOG GAMMA, [RECOMBINANT FACTOR IX] – Restricted see terms below			
⌘ Inj 250 iu vial	287.50	1	RIXUBIS
⌘ Inj 500 iu vial	575.00	1	RIXUBIS
⌘ Inj 1,000 iu vial	1,150.00	1	RIXUBIS
⌘ Inj 2,000 iu vial	2,300.00	1	RIXUBIS
⌘ Inj 3,000 iu vial	3,450.00	1	RIXUBIS

➔ **Restricted**

Initiation

When used in the treatment of haemophilia, treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) – Restricted see terms below

⌘ Inj 250 iu vial	287.50	1	Advate
⌘ Inj 500 iu vial	575.00	1	Advate
⌘ Inj 1,000 iu vial	1,150.00	1	Advate
⌘ Inj 1,500 iu vial	1,725.00	1	Advate
⌘ Inj 2,000 iu vial	2,300.00	1	Advate
⌘ Inj 3,000 iu vial	3,450.00	1	Advate

➔ **Restricted**

Initiation

Notes: Rare Clinical Circumstances Brand of recombinant factor VIII from 1 March 2016 until 28 February 2019. When used in the treatment of haemophilia, access to funded treatment by application to the Haemophilia Treatments Panel. Application details may be obtained from PHARMAC.s website <http://www.pharmac.govt.nz> or:

The Co-ordinator, Haemophilia Treatments Panel	Phone: 0800 023 588 Option 2
PHARMAC PO Box 10 254	Facsimile: (04) 974 4881
Wellington	Email: haemophilia@pharmac.govt.nz

OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE FS) – Restricted see terms below

⌘ Inj 250 iu vial	237.50	1	Kogenate FS
⌘ Inj 500 iu vial	475.00	1	Kogenate FS
⌘ Inj 1,000 iu vial	950.00	1	Kogenate FS
⌘ Inj 2,000 iu vial	1,900.00	1	Kogenate FS
⌘ Inj 3,000 iu vial	2,850.00	1	Kogenate FS

➔ **Restricted**

Initiation

Notes: Second Brand of recombinant factor VIII from 1 March 2016 until 28 February 2019. When used in the treatment of haemophilia, access to funded treatment by application to the Haemophilia Treatments Panel. Application details may be obtained from PHARMAC.s website <http://www.pharmac.govt.nz> or:

The Co-ordinator, Haemophilia Treatments Panel	Phone: 0800 023 588 Option 2
PHARMAC PO Box 10 254	Facsimile: (04) 974 4881
Wellington	Email: haemophilia@pharmac.govt.nz

Vitamin K

PHYTOMENADIONE

Inj 2 mg in 0.2 ml ampoule	8.00	5	Konaktion MM
Inj 10 mg per ml, 1 ml ampoule	9.21	5	Konaktion MM

Antithrombotics

Anticoagulants

BIVALIRUDIN – **Restricted** see terms below

¶ Inj 250 mg vial

➡ **Restricted**

Initiation

Either:

- 1 For use in heparin-induced thrombocytopenia, heparin resistance or heparin intolerance; or
- 2 For use in patients undergoing endovascular procedures.

DABIGATRAN

Cap 75 mg	148.00	60	Pradaxa
Cap 110 mg	148.00	60	Pradaxa
Cap 150 mg	148.00	60	Pradaxa

DALTEPARIN

Inj 2,500 iu in 0.2 ml syringe	19.97	10	Fragmin
Inj 5,000 iu in 0.2 ml syringe	39.94	10	Fragmin
Inj 7,500 iu in 0.75 ml syringe	60.03	10	Fragmin
Inj 10,000 iu in 1 ml syringe	77.55	10	Fragmin
Inj 12,500 iu in 0.5 ml syringe	99.96	10	Fragmin
Inj 15,000 iu in 0.6 ml syringe	120.05	10	Fragmin
Inj 18,000 iu in 0.72 ml syringe	158.47	10	Fragmin

DANAPAROID – **Restricted** see terms below

¶ Inj 750 u in 0.6 ml ampoule

➡ **Restricted**

Initiation

For use in heparin-induced thrombocytopenia, heparin resistance or heparin intolerance.

DEFIBROTIDE – **Restricted** see terms below

¶ Inj 80 mg per ml, 2.5 ml ampoule

➡ **Restricted**

Initiation

Haematologist

Patient has moderate or severe sinusoidal obstruction syndrome as a result of chemotherapy or regimen-related toxicities.

DEXTROSE WITH SODIUM CITRATE AND CITRIC ACID [ACID CITRATE DEXTROSE A]

Inj 24.5 mg with sodium citrate 22 mg and citric acid 7.3 mg per ml,
100 ml bag

ENOXAPARIN

Inj 20 mg in 0.2 ml syringe	37.24	10	Clexane
Inj 40 mg in 0.4 ml ampoule			
Inj 40 mg in 0.4 ml syringe	49.69	10	Clexane
Inj 60 mg in 0.6 ml syringe	74.91	10	Clexane
Inj 80 mg in 0.8 ml syringe	99.86	10	Clexane
Inj 100 mg in 1 ml syringe	125.06	10	Clexane
Inj 120 mg in 0.8 ml syringe	155.40	10	Clexane
Inj 150 mg in 1 ml syringe	177.60	10	Clexane

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FONDAPARINUX SODIUM – Restricted see terms below			
¶ Inj 2.5 mg in 0.5 ml syringe			
¶ Inj 7.5 mg in 0.6 ml syringe			
➔Restricted			
Initiation			
For use in heparin-induced thrombocytopaenia, heparin resistance or heparin intolerance.			
HEPARIN SODIUM			
Inj 100 iu per ml, 250 ml bag			
Inj 1,000 iu per ml, 1 ml ampoule	66.80	50	Hospira
Inj 1,000 iu per ml, 35 ml vial			
Inj 1,000 iu per ml, 5 ml ampoule	61.04	50	Pfizer
Inj 5,000 iu in 0.2 ml ampoule			
Inj 5,000 iu per ml, 1 ml ampoule	14.20	5	Hospira
Inj 5,000 iu per ml, 5 ml ampoule	236.60	50	Pfizer
HEPARINISED SALINE			
Inj 10 iu per ml, 5 ml ampoule	39.00	50	Pfizer
Inj 100 iu per ml, 2 ml ampoule			
Inj 100 iu per ml, 5 ml ampoule			
PHENINDIONE			
Tab 10 mg			
Tab 25 mg			
Tab 50 mg			
PROTAMINE SULPHATE			
Inj 10 mg per ml, 5 ml ampoule			
RIVAROXABAN – Restricted see terms below			
¶ Tab 10 mg	153.00	15	Xarelto
➔Restricted			
Initiation — total hip replacement			
<i>Limited to 5 weeks treatment</i>			
For the prophylaxis of venous thromboembolism.			
Initiation — total knee replacement			
<i>Limited to 2 weeks treatment</i>			
For the prophylaxis of venous thromboembolism.			
SODIUM CITRATE WITH SODIUM CHLORIDE AND POTASSIUM CHLORIDE			
Inj 4.2 mg with sodium chloride 5.7 mg and potassium chloride 74.6 mcg per ml, 5,000 ml bag			
TRISODIUM CITRATE			
Inj 4%, 5 ml ampoule			
Inj 46.7%, 3 ml syringe			
Inj 46.7%, 5 ml ampoule			
WARFARIN SODIUM			
Tab 1 mg	6.86	100	Marevan
Tab 2 mg			
Tab 3 mg	9.70	100	Marevan
Tab 5 mg	11.75	100	Marevan

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Antiplatelets			
ASPIRIN			
Tab 100 mg – 1% DV Mar-14 to 2016	1.60	90	Ethics Aspirin EC
Suppos 300 mg	10.50	990	Ethics Aspirin EC
CLOPIDOGREL			
Tab 75 mg – 1% DV Dec-13 to 2016	5.48	84	Arrow - Clopid
DIPYRIDAMOLE			
Tab 25 mg			
Tab long-acting 150 mg	11.52	60	Pytazen SR
Inj 5 mg per ml, 2 ml ampoule			
EPTIFIBATIDE – Restricted see terms below			
☞ Inj 2 mg per ml, 10 ml vial	111.00	1	Integrilin
☞ Inj 750 mcg per ml, 100 ml vial	324.00	1	Integrilin
☞ Restricted			
Initiation			
Either:			
1 For use in patients with acute coronary syndromes undergoing percutaneous coronary intervention; or			
2 For use in patients with definite or strongly suspected intra-coronary thrombus on coronary angiography.			
PRASUGREL – Restricted see terms below			
☞ Tab 5 mg	108.00	28	Effient
☞ Tab 10 mg	120.00	28	Effient
☞ Restricted			
Initiation — Bare metal stents			
<i>Limited to 6 months treatment</i>			
Patient has undergone coronary angioplasty in the previous 4 weeks and is clopidogrel-allergic.			
Initiation — Drug-eluting stents			
<i>Limited to 12 months treatment</i>			
Patient has had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic.			
Initiation — Stent thrombosis			
Patient has experienced cardiac stent thrombosis whilst on clopidogrel.			
Initiation — Myocardial infarction			
<i>Limited to 1 week treatment</i>			
For short term use while in hospital following ST-elevated myocardial infarction.			
Note: Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment			
TICAGRELOR – Restricted see terms below			
☞ Tab 90 mg	90.00	56	Brilinta
☞ Restricted			
Initiation			
Restricted to treatment of acute coronary syndromes specifically for patients who have recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome, and in whom fibrinolytic therapy has not been given in the last 24 hours and is not planned.			
TICLOPIDINE			
Tab 250 mg			

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Fibrinolytic Agents

ALTEPLASE

Inj 2 mg vial
Inj 10 mg vial
Inj 50 mg vial

TENECTEPLASE

Inj 50 mg vial

UROKINASE

Inj 10,000 iu vial
Inj 50,000 iu vial
Inj 100,000 iu vial
Inj 500,000 iu vial

Colony-Stimulating Factors

Granulocyte Colony-Stimulating Factors

FILGRASTIM – **Restricted** see terms below

⚡ Inj 300 mcg in 0.5 ml prefilled syringe	270.00	5	Zarzio
⚡ Inj 300 mcg in 1 ml vial	650.00	5	Neupogen
⚡ Inj 480 mcg in 0.5 ml prefilled syringe	432.00	5	Zarzio

➔Restricted

Haematologist or oncologist

PEGFILGRASTIM – **Restricted** see terms below

⚡ Inj 6 mg per 0.6 ml syringe	1,080.00	1	Neulastim
-------------------------------------	----------	---	-----------

➔Restricted

Initiation

For prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk \geq 20%).

Note: *Febrile neutropenia risk \geq 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines

Fluids and Electrolytes

Intravenous Administration

CALCIUM CHLORIDE

Inj 100 mg per ml, 10 ml vial

CALCIUM GLUCONATE

Inj 10%, 10 ml ampoule	34.24	10	Hospira
------------------------------	-------	----	---------

COMPOUND ELECTROLYTES

Inj sodium 140 mmol/l with potassium 5 mmol/l, magnesium 1.5 mmol/l, chloride 98 mmol/l, acetate 27 mmol/l and gluconate 23 mmol/l, bag	3.10	1,000 ml	Baxter
	5.00	500 ml	Baxter

COMPOUND ELECTROLYTES WITH GLUCOSE

Inj glucose 50 g with 140 mmol/l sodium, 5 mmol/l potassium, 1.5 mmol/l magnesium, 98 mmol/l chloride, 27 mmol/l acetate and 23 mmol/l gluconate, bag	7.00	1,000 ml	Baxter
---	------	----------	--------

BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
COMPOUND SODIUM LACTATE [HARTMANN'S SOLUTION]			
Inj sodium 131 mmol/l with potassium 5 mmol/l, calcium 2 mmol/l, bi-carbonate 29 mmol/l, chloride 111 mmol/l, bag	1.77	500 ml	Baxter
	1.80	1,000 ml	Baxter
COMPOUND SODIUM LACTATE WITH GLUCOSE			
Inj sodium 131 mmol/l with potassium 5 mmol/l, calcium 2 mmol/l, bi-carbonate 29 mmol/l, chloride 111 mmol/l and glucose 5%, bag	5.38	1,000 ml	Baxter
GLUCOSE [DEXTROSE]			
Inj 5%, bag	1.77	500 ml	Baxter
	1.80	1,000 ml	Baxter
	2.84	100 ml	Baxter
	2.87	50 ml	Baxter
	3.87	250 ml	Baxter
Inj 10%, bag	3.70	500 ml	Baxter
	5.29	1,000 ml	Baxter
Inj 50%, bag	6.84	500 ml	Baxter
Inj 50%, 10 ml ampoule – 1% DV Oct-14 to 2017	27.50	5	Biomed
Inj 50%, 90 ml bottle – 1% DV Oct-14 to 2017	14.50	1	Biomed
Inj 70%, 1,000 ml bag			
Inj 70%, 500 ml bag			
GLUCOSE WITH POTASSIUM CHLORIDE			
Inj 5% glucose with 20 mmol/l potassium chloride, bag	7.36	1,000 ml	Baxter
Inj 5% glucose with 30 mmol/l potassium chloride, 1,000 ml bag			
Inj 10% glucose with 10 mmol/l potassium chloride, 500 ml bag			
GLUCOSE WITH POTASSIUM CHLORIDE AND SODIUM CHLORIDE			
Inj 4% glucose with potassium chloride 20 mmol/l and sodium chloride 0.18%, bag	3.45	500 ml	Baxter
	4.30	1,000 ml	Baxter
Inj 4% glucose with potassium chloride 30 mmol/l and sodium chloride 0.18%, bag	3.62	1,000 ml	Baxter
Inj 2.5% glucose with potassium chloride 20 mmol/l and sodium chloride 0.45%, 3,000 ml bag			
Inj 10% glucose with potassium chloride 10 mmol/l and sodium chloride 15 mmol/l, 500 ml bag			
GLUCOSE WITH SODIUM CHLORIDE			
Inj glucose 2.5% with sodium chloride 0.45%, bag	4.95	500 ml	Baxter
Inj glucose 5% with sodium chloride 0.45%, bag	5.80	1,000 ml	Baxter
	9.87	500 ml	Baxter
Inj glucose 5% with sodium chloride 0.9%, bag	4.54	1,000 ml	Baxter
Inj glucose 5% with sodium chloride 0.2%, 500 ml bag			
POTASSIUM CHLORIDE			
Inj 75 mg (1 mmol) per ml, 10 ml ampoule			
Inj 225 mg (3 mmol) per ml, 20 ml ampoule			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
POTASSIUM CHLORIDE WITH SODIUM CHLORIDE			
Inj 20 mmol/l potassium chloride with 0.9% sodium chloride, bag	3.85	1,000 ml	Baxter
Inj 30 mmol/l potassium chloride with 0.9% sodium chloride, bag	2.59	1,000 ml	Baxter
Inj 40 mmol/l potassium chloride with 0.9% sodium chloride, bag	6.62	1,000 ml	Baxter
Inj 10 mmol potassium chloride with 0.29% sodium chloride, 100 ml bag			
Inj 40 mmol potassium chloride with 0.9% sodium chloride, 100 ml bag			
POTASSIUM DIHYDROGEN PHOSPHATE			
Inj 1 mmol per ml, 10 ml ampoule – 1% DV Oct-15 to 2018	151.80	10	Hospira
RINGER'S SOLUTION			
Inj sodium 147 mmol/l with potassium 4 mmol/l, calcium 2.2 mmol/l, chloride 156 mmol/l, bag	5.13	1,000 ml	Baxter
SODIUM ACETATE			
Inj 4 mmol per ml, 20 ml ampoule			
SODIUM BICARBONATE			
Inj 8.4%, 10 ml vial			
Inj 8.4%, 50 ml vial	19.95	1	Biomed
Inj 8.4%, 100 ml vial	20.50	1	Biomed
SODIUM CHLORIDE			
Inj 0.45%, bag	5.50	500 ml	Baxter
Inj 0.9%, bag	1.77	500 ml	Baxter
	1.80	1,000 ml	Baxter
	2.28	100 ml	Baxter
	3.01	50 ml	Baxter
	3.60	250 ml	Baxter
	1.70	500 ml	Freeflex
	1.71	1,000 ml	Freeflex
Inj 3%, bag	5.69	1,000 ml	Baxter
Inj 0.9%, 5 ml ampoule	10.85	50	Multichem
	15.50		Pfizer
Inj 0.9%, 10 ml ampoule	11.50	50	Multichem
	15.50		Pfizer
⚡ Inj 0.9%, 3 ml syringe, non-sterile pack – 1% DV Jun-15 to 2018	10.65	30	BD PosiFlush
➡ Restricted			
Initiation			
For use in flushing of in-situ vascular access devices only.			
⚡ Inj 0.9%, 5 ml syringe, non-sterile pack – 1% DV Jun-15 to 2018	10.80	30	BD PosiFlush
➡ Restricted			
Initiation			
For use in flushing of in-situ vascular access devices only.			
⚡ Inj 0.9%, 10 ml syringe, non-sterile pack – 1% DV Jun-15 to 2018	11.25	30	BD PosiFlush
➡ Restricted			
Initiation			
For use in flushing of in-situ vascular access devices only.			
Inj 0.9%, 20 ml ampoule	8.41	20	Multichem
Inj 23.4% (4 mmol/ml), 20 ml – 1% DV Sep-13 to 2016	31.25	5	Biomed
Inj 1.8%, 500 ml bottle			
SODIUM DIHYDROGEN PHOSPHATE [SODIUM ACID PHOSPHATE]			
Inj 1 mmol per ml, 20 ml ampoule – 1% DV Oct-15 to 2018	47.50	5	Biomed

BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
WATER			
Inj, bag	2.75	1,000 ml	Baxter
Inj 5 ml ampoule	10.25	50	Multichem
Inj 10 ml ampoule	11.25	50	Multichem
Inj 20 ml ampoule	6.50	20	Multichem
Inj 250 ml bag			
Inj 500 ml bag			
Oral Administration			
CALCIUM POLYSTYRENE SULPHONATE			
Powder	169.85	300 g	Calcium Resonium
COMPOUND ELECTROLYTES			
Powder for oral soln			
COMPOUND ELECTROLYTES WITH GLUCOSE			
Soln with electrolytes			
PHOSPHORUS			
Tab eff 500 mg (16 mmol)			
POTASSIUM CHLORIDE			
Tab eff 548 mg (14 mmol) with chloride 285 mg (8 mmol)			
Tab long-acting 600 mg (8 mmol) – 1% DV Sep-15 to 2018	7.42	200	Span-K
Oral liq 2 mmol per ml			
SODIUM BICARBONATE			
Cap 840 mg	8.52	100	Sodibic
SODIUM CHLORIDE			
Tab 600 mg			
Oral liq 2 mmol/ml			
SODIUM POLYSTYRENE SULPHONATE			
Powder – 1% DV Sep-15 to 2018	84.65	454 g	Resonium A
Plasma Volume Expanders			
GELATINE, SUCCINYLATED			
Inj 4%, 500 ml bag	108.00	10	Gelofusine
HYDROXYETHYL STARCH 130/0.4 WITH MAGNESIUM CHLORIDE, POTASSIUM CHLORIDE, SODIUM ACETATE AND SODIUM CHLORIDE			
Inj 6% with magnesium chloride 0.03%, potassium chloride 0.03%, sodium acetate 0.463% and sodium chloride 0.6%, 500 ml bag	198.00	20	Volulyte 6%
HYDROXYETHYL STARCH 130/0.4 WITH SODIUM CHLORIDE			
Inj 6% with sodium chloride 0.9%, 500 ml bag	198.00	20	Voluven

Price (ex man. excl. GST)	Brand or Generic Manufacturer
\$ Per	

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

CAPTOPRIL

☞ Oral liq 5 mg per ml94.99 95 ml Capoten

☞ Restricted

Initiation

Any of the following:

- 1 For use in children under 12 years of age; or
- 2 For use in tube-fed patients; or
- 3 For management of rebound transient hypertension following cardiac surgery.

CILAZAPRIL

Tab 0.5 mg – 1% DV Sep-13 to 2016	2.00	90	Zapril
Tab 2.5 mg – 1% DV Sep-13 to 2016	4.31	90	Zapril
Tab 5 mg – 1% DV Sep-13 to 2016	6.98	90	Zapril

ENALAPRIL MALEATE

Tab 5 mg – 1% DV Sep-15 to 2018	0.96	100	Ethics Enalapril
Tab 10 mg – 1% DV Sep-15 to 2018	1.24	100	Ethics Enalapril
Tab 20 mg – 1% DV Sep-15 to 2018	1.78	100	Ethics Enalapril

LISINOPRIL

Tab 5 mg – 1% DV Jan-16 to 2018	1.80	90	Ethics Lisinopril
Tab 10 mg – 1% DV Jan-16 to 2018	2.05	90	Ethics Lisinopril
Tab 20 mg – 1% DV Jan-16 to 2018	2.76	90	Ethics Lisinopril

PERINDOPRIL

Tab 2 mg – 1% DV Oct-14 to 2017	3.75	30	Apo-Perindopril
Tab 4 mg – 1% DV Oct-14 to 2017	4.80	30	Apo-Perindopril

QUINAPRIL

Tab 5 mg – 1% DV Sep-15 to 2018	4.31	90	Arrow-Quinapril 5
Tab 10 mg – 1% DV Sep-15 to 2018	3.15	90	Arrow-Quinapril 10
Tab 20 mg – 1% DV Sep-15 to 2018	5.97	90	Arrow-Quinapril 20

TRANDOLAPRIL – **Restricted:** For continuation only

☞ Cap 1 mg

☞ Cap 2 mg

ACE Inhibitors with Diuretics

CILAZAPRIL WITH HYDROCHLOROTHIAZIDE

Tab 5 mg with hydrochlorothiazide 12.5 mg – 1% DV Mar-14 to 2016	10.72	100	Apo-Cilazapril/ Hydrochlorothiazide
--	-------	-----	--

ENALAPRIL MALEATE WITH HYDROCHLOROTHIAZIDE – **Restricted:** For continuation only

☞ Tab 20 mg with hydrochlorothiazide 12.5 mg

QUINAPRIL WITH HYDROCHLOROTHIAZIDE

Tab 10 mg with hydrochlorothiazide 12.5 mg – 1% DV Oct-15 to 2018	3.65	30	Accuretic 10
Tab 20 mg with hydrochlorothiazide 12.5 mg – 1% DV Oct-15 to 2018	4.78	30	Accuretic 20

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Angiotensin II Antagonists			
CANDESARTAN CILEXETIL – Restricted see terms below			
⚡ Tab 4 mg – 1% DV Sep-15 to 2018	2.50	90	Candestar
⚡ Tab 8 mg – 1% DV Sep-15 to 2018	3.68	90	Candestar
⚡ Tab 16 mg – 1% DV Sep-15 to 2018	6.12	90	Candestar
⚡ Tab 32 mg – 1% DV Sep-15 to 2018	10.66	90	Candestar
➡ Restricted			
Initiation — ACE inhibitor intolerance			
Either:			
1 Patient has persistent ACE inhibitor induced cough that is not resolved by ACE inhibitor retrial (same or new ACE inhibitor); or			
2 Patient has a history of angioedema.			
Initiation — Unsatisfactory response to ACE inhibitor			
Patient is not adequately controlled on maximum tolerated dose of an ACE inhibitor.			
LOSARTAN POTASSIUM			
Tab 12.5 mg – 1% DV Jan-15 to 2017	1.55	84	Losartan Actavis
Tab 25 mg – 1% DV Jan-15 to 2017	1.90	84	Losartan Actavis
Tab 50 mg – 1% DV Jan-15 to 2017	2.25	84	Losartan Actavis
Tab 100 mg – 1% DV Jan-15 to 2017	2.60	84	Losartan Actavis
Angiotensin II Antagonists with Diuretics			
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE			
Tab 50 mg with hydrochlorothiazide 12.5 mg – 1% DV Oct-14 to 2017	2.18	30	Arrow-Losartan & Hydrochlorothiazide
Alpha-Adrenoceptor Blockers			
DOXAZOSIN			
Tab 2 mg – 1% DV Sep-14 to 2017	6.75	500	Apo-Doxazosin
Tab 4 mg – 1% DV Sep-14 to 2017	9.67	500	Apo-Doxazosin
PHENOXYBENZAMINE HYDROCHLORIDE			
Cap 10 mg			
Inj 50 mg per ml, 2 ml ampoule			
PHENTOLAMINE MESYLATE			
Inj 10 mg per ml, 1 ml ampoule			
PRAZOSIN			
Tab 1 mg	5.53	100	Apo-Prazosin
Tab 2 mg	7.00	100	Apo-Prazosin
Tab 5 mg	11.70	100	Apo-Prazosin
TERAZOSIN			
Tab 1 mg – 1% DV Sep-13 to 2016	0.50	28	Arrow
Tab 2 mg – 1% DV Sep-13 to 2016	0.45	28	Arrow
Tab 5 mg – 1% DV Sep-13 to 2016	0.68	28	Arrow
Antiarrhythmics			
ADENOSINE			
Inj 3 mg per ml, 2 ml vial			
⚡ Inj 3 mg per ml, 10 ml vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Initiation			
For use in cardiac catheterisation, electrophysiology and MRI.			
AJMALINE – Restricted see terms below			
⚡ Inj 5 mg per ml, 10 ml ampoule			
➔Restricted			
Cardiologist			
AMIODARONE HYDROCHLORIDE			
Tab 100 mg			
Tab 200 mg			
Inj 50 mg per ml, 3 ml ampoule – 1% DV Aug-13 to 2016	22.80	6	Cordarone-X
ATROPINE SULPHATE			
Inj 600 mcg per ml, 1 ml ampoule	71.00	50	AstraZeneca
DIGOXIN			
Tab 62.5 mcg – 1% DV Jun-16 to 2019	6.67	240	Lanoxin PG
Tab 250 mcg – 1% DV Apr-16 to 2019	14.52	240	Lanoxin
Oral liq 50 mcg per ml			
Inj 250 mcg per ml, 2 ml vial			
DISOPYRAMIDE PHOSPHATE			
Cap 100 mg			
Cap 150 mg			
FLECAINIDE ACETATE			
Tab 50 mg	38.95	60	Tambocor
Cap long-acting 100 mg	38.95	30	Tambocor CR
Cap long-acting 200 mg	68.78	30	Tambocor CR
Inj 10 mg per ml, 15 ml ampoule	52.45	5	Tambocor
MEXILETINE HYDROCHLORIDE			
Cap 150 mg	162.00	100	Mexiletine Hydrochloride USP
Cap 250 mg	202.00	100	Mexiletine Hydrochloride USP
PROPAFENONE HYDROCHLORIDE			
Tab 150 mg			

Antihypotensives

MIDODRINE – **Restricted** see terms below

⚡ Tab 2.5 mg

⚡ Tab 5 mg

➔Restricted

Initiation

Patient has disabling orthostatic hypotension not due to drugs.

Beta-Adrenoceptor Blockers

ATENOLOL

Tab 50 mg – 1% DV Sep-15 to 2018	4.61	500	Mylan Atenolol
Tab 100 mg – 1% DV Sep-15 to 2018	7.67	500	Mylan Atenolol
Oral liq 5 mg per ml	21.25	300 ml	Atenolol-AFT

CARDIOVASCULAR SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BISOPROLOL FUMARATE			
Tab 2.5 mg – 1% DV Mar-15 to 2017	2.40	30	Bosvate
Tab 5 mg – 1% DV Mar-15 to 2017	3.50	30	Bosvate
Tab 10 mg – 1% DV Mar-15 to 2017	6.40	30	Bosvate
CARVEDILOL			
Tab 6.25 mg – 1% DV Jun-15 to 2017	3.90	60	Dicarz
Tab 12.5 mg – 1% DV Jun-15 to 2017	5.10	60	Dicarz
Tab 25 mg – 1% DV Jun-15 to 2017	6.30	60	Dicarz
CELIPROLOL			
Tab 200 mg	21.40	180	Celol
ESMOLOL HYDROCHLORIDE			
Inj 10 mg per ml, 10 ml vial			
LABETALOL			
Tab 50 mg	8.23	100	Hybloc
Tab 100 mg	10.06	100	Hybloc
Tab 200 mg	17.55	100	Hybloc
Tab 400 mg			
Inj 5 mg per ml, 20 ml ampoule			
METOPROLOL SUCCINATE			
Tab long-acting 23.75 mg – 1% DV Jun-16 to 2018	2.39	90	Metoprolol - AFT CR
Tab long-acting 47.5 mg – 1% DV Jun-16 to 2018	3.48	90	Metoprolol - AFT CR
Tab long-acting 95 mg – 1% DV Jun-16 to 2018	5.73	90	Metoprolol - AFT CR
Tab long-acting 190 mg – 1% DV Jun-16 to 2018	11.54	90	Metoprolol - AFT CR
METOPROLOL TARTRATE			
Tab 50 mg	16.00	100	Lopresor
Tab 100 mg	21.00	60	Lopresor
Tab long-acting 200 mg	18.00	28	Slow-Lopresor
Inj 1 mg per ml, 5 ml vial	24.00	5	Lopresor
NADOLOL			
Tab 40 mg – 1% DV Oct-15 to 2018	16.05	100	Apo-Nadolol
Tab 80 mg – 1% DV Oct-15 to 2018	24.70	100	Apo-Nadolol
PINDOLOL			
Tab 5 mg – 1% DV Nov-13 to 2016	9.72	100	Apo-Pindolol
Tab 10 mg – 1% DV Nov-13 to 2016	15.62	100	Apo-Pindolol
Tab 15 mg – 1% DV Nov-13 to 2016	23.46	100	Apo-Pindolol
PROPRANOLOL			
Tab 10 mg	3.65	100	Apo-Propranolol
Tab 40 mg	4.65	100	Apo-Propranolol
Cap long-acting 160 mg	18.17	100	Cardinol LA
Oral liq 4 mg per ml			
Inj 1 mg per ml, 1 ml ampoule			
SOTALOL			
Tab 80 mg	27.50	500	Mylan
Tab 160 mg	10.50	100	Mylan
Inj 10 mg per ml, 4 ml ampoule	65.39	5	Sotacor
TIMOLOL MALEATE			
Tab 10 mg			

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

Price (ex man. excl. GST)	Brand or Generic
\$	Per Manufacturer

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

AMLODIPINE

Tab 2.5 mg – 1% DV Feb-15 to 2017	2.21	100	Apo-Amlodipine
Tab 5 mg – 1% DV May-15 to 2017	5.04	250	Apo-Amlodipine
Tab 10 mg – 1% DV May-15 to 2017	7.21	250	Apo-Amlodipine

FELODIPINE

Tab long-acting 2.5 mg – 1% DV Sep-15 to 2018	1.45	30	Plendil ER
Tab long-acting 5 mg – 1% DV Sep-15 to 2018	1.55	30	Plendil ER
Tab long-acting 10 mg – 1% DV Sep-15 to 2018	2.30	30	Plendil ER

ISRADIPINE

Tab 2.5 mg
Cap 2.5 mg
Cap long-acting 2.5 mg
Cap long-acting 5 mg

NICARDIPINE HYDROCHLORIDE – **Restricted** see terms below

☞ Inj 2.5 mg per ml, 10 ml vial

☞ **Restricted**

Initiation

Anaesthetist, intensivist or paediatric cardiologist

Both:

- 1 Patient is a Paediatric Patient; and
- 2 Any of the following:
 - 2.1 Patient has hypertension requiring urgent treatment with an intravenous agent; or
 - 2.2 Patient has excessive ventricular afterload; or
 - 2.3 Patient is awaiting or undergoing cardiac surgery using cardiopulmonary bypass.

NIFEDIPINE

Tab long-acting 10 mg			
Tab long-acting 20 mg	9.59	100	Nyefax Retard
Tab long-acting 30 mg – 1% DV Sep-14 to 2017	3.75	30	Adefin XL
Tab long-acting 60 mg – 1% DV Sep-14 to 2017	5.75	30	Adefin XL
Cap 5 mg			

NIMODIPINE

Tab 30 mg
Inj 200 mcg per ml, 50 ml vial

Other Calcium Channel Blockers

DILTIAZEM HYDROCHLORIDE

Tab 30 mg	4.60	100	Dilzem
Tab 60 mg	8.50	100	Dilzem
Cap long-acting 120 mg	31.83	500	Apo-Diltiazem CD
	1.91	30	Cardizem CD
Cap long-acting 180 mg	47.67	500	Apo-Diltiazem CD
	7.56	30	Cardizem CD
Cap long-acting 240 mg	63.58	500	Apo-Diltiazem CD
	10.22	30	Cardizem CD
Inj 5 mg per ml, 5 ml vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PERHEXILINE MALEATE			
Tab 100 mg – 1% DV Jun-16 to 2019	62.90	100	Pexsig
VERAPAMIL HYDROCHLORIDE			
Tab 40 mg	7.01	100	Isoptin
Tab 80 mg – 1% DV Sep-14 to 2017	11.74	100	Isoptin
Tab long-acting 120 mg	15.20	250	Verpamil SR
Tab long-acting 240 mg	25.00	250	Verpamil SR
Inj 2.5 mg per ml, 2 ml ampoule	7.54	5	Isoptin

Centrally-Acting Agents

CLONIDINE			
Patch 2.5 mg, 100 mcg per day – 1% DV Jul-14 to 2017	12.80	4	Catapres-TTS-1
Patch 5 mg, 200 mcg per day – 1% DV Jul-14 to 2017	18.04	4	Catapres-TTS-2
Patch 7.5 mg, 300 mcg per day – 1% DV Jul-14 to 2017	22.68	4	Catapres-TTS-3
CLONIDINE HYDROCHLORIDE			
Tab 25 mcg – 1% DV Sep-15 to 2018	10.53	112	Clonidine BNM
Tab 150 mcg	34.32	100	Catapres
Inj 150 mcg per ml, 1 ml ampoule	16.07	5	Catapres
METHYLDOPA			
Tab 125 mg	14.25	100	Prodopa
Tab 250 mg	15.10	100	Prodopa
Tab 500 mg	23.15	100	Prodopa

Diuretics

Loop Diuretics

BUMETANIDE			
Tab 1 mg	16.36	100	Burinex
Inj 500 mcg per ml, 4 ml vial			
FUROSEMIDE [FRUSEMIDE]			
Tab 40 mg – 1% DV Sep-15 to 2018	8.00	1,000	Diurin 40
Tab 500 mg – 1% DV Sep-15 to 2018	25.00	50	Urex Forte
Oral liq 10 mg per ml			
Inj 10 mg per ml, 2 ml ampoule – 1% DV Jun-16 to 2019	1.20	5	Frusemide-Claris
Inj 10 mg per ml, 25 ml ampoule			

Osmotic Diuretics

MANNITOL			
Inj 10%, 1,000 ml bag	14.21	1,000 ml	Baxter
Inj 15%, 500 ml bag	9.84	500 ml	Baxter
Inj 20%, 500 ml bag	10.80	500 ml	Baxter

Potassium Sparing Combination Diuretics

AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE			
Tab 5 mg with furosemide 40 mg			
AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE			
Tab 5 mg with hydrochlorothiazide 50 mg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Potassium Sparing Diuretics			
AMILORIDE HYDROCHLORIDE			
Tab 5 mg	17.50	100	Apo-Amiloride
Oral liq 1 mg per ml	30.00	25 ml	Biomed
SPIRONOLACTONE			
Tab 25 mg – 1% DV Sep-13 to 2016	3.65	100	Spiractin
Tab 100 mg – 1% DV Sep-13 to 2016	11.80	100	Spiractin
Oral liq 5 mg per ml	30.00	25 ml	Biomed
Thiazide and Related Diuretics			
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]			
Tab 2.5 mg – 1% DV Sep-14 to 2017	5.48	500	Arrow-Bendrofluazide
Tab 5 mg – 1% DV Sep-14 to 2017	8.95	500	Arrow-Bendrofluazide
CHLOROTHIAZIDE			
Oral liq 50 mg per ml	26.00	25 ml	Biomed
CHLORTALIDONE [CHLORTHALIDONE]			
Tab 25 mg	8.00	50	Hygroton
INDAPAMIDE			
Tab 2.5 mg – 1% DV Oct-13 to 2016	2.25	90	Dapa-Tabs
METOLAZONE – Restricted see terms below			
⬇ Tab 5 mg			
Initiation			
Either:			
1 Patient has refractory heart failure and is intolerant or has not responded to loop diuretics and/or loop-thiazide combination therapy; or			
2 Patient has severe refractory nephrotic oedema unresponsive to high dose loop diuretics and concentrated albumin infusions.			
Lipid-Modifying Agents			
Fibrates			
BEZAFIBRATE			
Tab 200 mg – 1% DV Oct-15 to 2018	9.05	90	Bezalip
Tab long-acting 400 mg – 1% DV Oct-15 to 2018	6.78	30	Bezalip Retard
GEMFIBROZIL			
Tab 600 mg – 1% DV Nov-13 to 2016	17.60	60	Lipazil
HMG CoA Reductase Inhibitors (Statins)			
ATORVASTATIN			
Tab 10 mg	2.52	90	Zarator
Tab 20 mg	4.17	90	Zarator
Tab 40 mg	7.32	90	Zarator
Tab 80 mg	16.23	90	Zarator

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PRAVASTATIN			
Tab 10 mg			
Tab 20 mg – 1% DV Oct-14 to 2017	3.45	30	Cholvastin
Tab 40 mg – 1% DV Oct-14 to 2017	6.36	30	Cholvastin
SIMVASTATIN			
Tab 10 mg – 1% DV Sep-14 to 2017	0.95	90	Arrow-Simva
Tab 20 mg – 1% DV Sep-14 to 2017	1.61	90	Arrow-Simva
Tab 40 mg – 1% DV Sep-14 to 2017	2.83	90	Arrow-Simva
Tab 80 mg – 1% DV Sep-14 to 2017	7.91	90	Arrow-Simva

Resins

CHOLESTYRAMINE

Powder for oral liq 4 g

COLESTIPOL HYDROCHLORIDE

Grans for oral liq 5 g

Selective Cholesterol Absorption Inhibitors

EZETIMIBE – Restricted see terms below

⚡ Tab 10 mg – 1% DV Aug-15 to 2017 3.35 30 **Ezemibe**

➡ **Restricted**

Initiation

All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 × normal) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

EZETIMIBE WITH SIMVASTATIN – Restricted see terms below

⚡ Tab 10 mg with simvastatin 10 mg – 1% DV Aug-15 to 2017 5.15 30 **Zimybe**
 ⚡ Tab 10 mg with simvastatin 20 mg – 1% DV Aug-15 to 2017 6.15 30 **Zimybe**
 ⚡ Tab 10 mg with simvastatin 40 mg – 1% DV Aug-15 to 2017 7.15 30 **Zimybe**
 ⚡ Tab 10 mg with simvastatin 80 mg – 1% DV Aug-15 to 2017 8.15 30 **Zimybe**

➡ **Restricted**

Initiation

All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Other Lipid-Modifying Agents

ACIPIMOX

Cap 250 mg

NICOTINIC ACID

Tab 50 mg – 1% DV Oct-14 to 2017 3.96 100 **Apo-Nicotinic Acid**
 Tab 500 mg – 1% DV Oct-14 to 2017 17.37 100 **Apo-Nicotinic Acid**

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Nitrates			
GLYCERYL TRINITRATE			
Tab 600 mcg	8.00	100	Lycinate
Inj 1 mg per ml, 5 ml ampoule	22.70	10	Nitronal
Inj 1 mg per ml, 50 ml vial	86.60	10	Nitronal
Inj 5 mg per ml, 10 ml ampoule	100.00	5	Hospira
Oral pump spray, 400 mcg per dose	4.45	250 dose	Nitrolingual Pump Spray
Oral spray, 400 mcg per dose	4.45	250 dose	Glytrin
Patch 25 mg, 5 mg per day – 1% DV Sep-14 to 2017	15.73	30	Nitroderm TTS 5
Patch 50 mg, 10 mg per day – 1% DV Sep-14 to 2017	18.62	30	Nitroderm TTS 10
ISOSORBIDE MONONITRATE			
Tab 20 mg – 1% DV Sep-14 to 2017	17.10	100	Ismo-20
Tab long-acting 40 mg – 1% DV Jun-16 to 2019	7.50	30	Ismo 40 Retard
Tab long-acting 60 mg	3.94	90	Duride
Other Cardiac Agents			
LEVOSIMENDAN – Restricted see terms below			
⚡ Inj 2.5 mg per ml, 5 ml vial			
⚡ Inj 2.5 mg per ml, 10 ml vial			
➡Restricted			
Initiation — Heart transplant			
Either:			
1 For use as a bridge to heart transplant, in patients who have been accepted for transplant; or			
2 For the treatment of heart failure following heart transplant.			
Initiation — Heart failure			
Cardiologist or intensivist			
For the treatment of severe acute decompensated heart failure that is non-responsive to dobutamine.			
Sympathomimetics			
ADRENALINE			
Inj 1 in 1,000, 1 ml ampoule	4.98	5	Aspen Adrenaline
	5.25		Hospira
Inj 1 in 1,000, 30 ml vial			
Inj 1 in 10,000, 10 ml ampoule	49.00	10	Aspen Adrenaline
	27.00	5	Hospira
Inj 1 in 10,000, 10 ml syringe			
DOBUTAMINE HYDROCHLORIDE			
Inj 12.5 mg per ml, 20 ml ampoule – 1% DV Jan-16 to 2018	24.45	5	Dobutamine-Claris
DOPAMINE HYDROCHLORIDE			
Inj 40 mg per ml, 5 ml ampoule – 1% DV Sep-15 to 2018	16.89	5	DBL Sterile Dopamine Concentrate
EPHEDRINE			
Inj 3 mg per ml, 10 ml syringe			
Inj 30 mg per ml, 1 ml ampoule – 1% DV Mar-15 to 2017	51.48	10	Max Health
ISOPRENALINE			
Inj 200 mcg per ml, 1 ml ampoule			
Inj 200 mcg per ml, 5 ml ampoule			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
METARAMINOL			
Inj 0.5 mg per ml, 20 ml syringe			
Inj 1 mg per ml, 1 ml ampoule			
Inj 1 mg per ml, 10 ml syringe			
Inj 10 mg per ml, 1 ml ampoule			
NORADRENALINE			
Inj 0.06 mg per ml, 100 ml bag			
Inj 0.06 mg per ml, 50 ml syringe			
Inj 0.1 mg per ml, 100 ml bag			
Inj 0.12 mg per ml, 100 ml bag			
Inj 0.12 mg per ml, 50 ml syringe			
Inj 0.16 mg per ml, 50 ml syringe			
Inj 1 mg per ml, 100 ml bag			
Inj 1 mg per ml, 4 ml ampoule			
PHENYLEPHRINE HYDROCHLORIDE			
Inj 10 mg per ml, 1 ml vial	115.50	25	Neosynephrine HCL
Vasodilators			
ALPROSTADIL HYDROCHLORIDE			
Inj 500 mcg per ml, 1 ml ampoule – 1% DV Oct-15 to 2018	1,650.00	5	Prostin VR
AMYL NITRITE			
Liq 98% in 3 ml capsule			
DIAZOXIDE			
Inj 15 mg per ml, 20 ml ampoule			
HYDRALAZINE HYDROCHLORIDE			
☞ Tab 25 mg			
☞ Restricted			
Initiation			
Either:			
1 For the treatment of refractory hypertension; or			
2 For the treatment of heart failure, in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.			
Inj 20 mg ampoule	25.90	5	Apresoline
MILRINONE			
Inj 1 mg per ml, 10 ml ampoule			
MINOXIDIL – Restricted see terms below			
☞ Tab 10 mg	70.00	100	Loniten
☞ Restricted			
Initiation			
For patients with severe refractory hypertension who have failed to respond to extensive multiple therapies.			
NICORANDIL			
Tab 10 mg	27.95	60	Ikorel
Tab 20 mg	33.28	60	Ikorel
PAPAVERINE HYDROCHLORIDE			
Inj 30 mg per ml, 1 ml vial			
Inj 12 mg per ml, 10 ml ampoule	217.90	5	Hospira

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PENTOXIFYLLINE [OXPENTIFYLLINE] Tab 400 mg			
SODIUM NITROPRUSSIDE Inj 50 mg vial			

Endothelin Receptor Antagonists

AMBRISENTAN – **Restricted** see terms below

⚡ Tab 5 mg	4,585.00	30	Volibris
⚡ Tab 10 mg	4,585.00	30	Volibris

➔**Restricted**

Initiation

Either:

- 1 For use in patients with approval by the Pulmonary Arterial Hypertension Panel; or
- 2 In hospital stabilisations in emergency situations.

BOSENTAN – **Restricted** see terms below

⚡ Tab 62.5 mg – 1% DV Jan-16 to 2018	375.00	56	Mylan-Bosentan
⚡ Tab 125 mg – 1% DV Jan-16 to 2018	375.00	56	Mylan-Bosentan

➔**Restricted**

Initiation

Either:

- 1 For use in patients with approval by the Pulmonary Arterial Hypertension Panel; or
- 2 In hospital stabilisation in emergency situations.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – **Restricted** see terms below

⚡ Tab 25 mg – 1% DV Sep-15 to 2018	0.75	4	Vedafil
⚡ Tab 50 mg – 1% DV Sep-15 to 2018	0.75	4	Vedafil
⚡ Tab 100 mg – 1% DV Sep-15 to 2018	2.75	4	Vedafil

➔**Restricted**

Initiation

Any of the following:

- 1 For use in patients with approval by the Pulmonary Arterial Hypertension Panel; or
- 2 For use in neonatal units for persistent pulmonary hypertension of the newborn (PPHN); or
- 3 For use in weaning patients from inhaled nitric oxide; or
- 4 For perioperative use in cardiac surgery patients; or
- 5 For use in intensive care as an alternative to nitric oxide; or
- 6 In-hospital stabilisation in emergency situations; or
- 7 All of the following:
 - 7.1 Patient has Raynaud's phenomenon; and
 - 7.2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
 - 7.3 Patient is following lifestyle management (proper body insulation, avoidance of cold exposure, smoking cessation support, avoidance of sympathomimetic drugs); and
 - 7.4 Patient has persisting severe symptoms despite treatment with calcium channel blockers and nitrates (unless contraindicated or not tolerated).

Prostacyclin Analogues

EPOPROSTENOL – **Restricted** see terms on the next page

⚡ Inj 0.5 mg vial	36.61	1	Veletri
⚡ Inj 1.5 mg vial	73.21	1	Veletri

Products with Hospital Supply Status (HSS) are in **bold**

Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➡Restricted			
Initiation			
For use as a bridge to transplant for patients with Pulmonary Arterial Hypertension who are on the active waiting list for lung transplantation.			
ILOPROST			
Inj 50 mcg in 0.5 ml ampoule – 1% DV Sep-15 to 2016	89.50	1	Arrow-Iloprost
⚡ Nebuliser soln 10 mcg per ml, 2 ml	1,185.00	30	Ventavis

➡Restricted

Initiation

Any of the following:

- 1 For use in patients with approval by the Pulmonary Arterial Hypertension Panel; or
- 2 For diagnostic use in catheter laboratories; or
- 3 For use following mitral or tricuspid valve surgery; or
- 4 In hospital stabilisation in emergency situations.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Anti-Infective Preparations			
Antibacterials			
FUSIDIC ACID			
Crm 2% – 1% DV Jan-15 to 2016	2.52	15 g	DP Fusidic Acid Cream
Oint 2% – 1% DV Sep-13 to 2016	3.45	15 g	Foban
HYDROGEN PEROXIDE			
Crm 1%	8.56	15 g	Crystaderm
Soln 3% (10 vol) – 1% DV Nov-15 to 2018	1.40	100 ml	Pharmacy Health
MAFENIDE ACETATE – Restricted see terms below			
↓ Powder 50 g sachet			
➔ Restricted			
Initiation			
For the treatment of burns patients.			
MUPIROCIN			
Oint 2%			
SULPHADIAZINE SILVER			
Crm 1%	12.30	50 g	Flamazine
Antifungals			
AMOROLFINE			
Nail soln 5% – 1% DV Jan-15 to 2017	19.95	5 ml	MycoNail
CICLOPIROX OLAMINE			
Nail soln 8% – 1% DV Sep-15 to 2018	6.50	7 ml	Apo-Ciclopirox
➔ Soln 1% – Restricted: For continuation only			
CLOTTRIMAZOLE			
Crm 1% – 1% DV Sep-14 to 2017	0.52	20 g	Clomazol
➔ Soln 1% – Restricted: For continuation only			
ECONAZOLE NITRATE			
➔ Crm 1% – Restricted: For continuation only			
Foaming soln 1%			
KETOCONAZOLE			
Shampoo 2% – 1% DV Dec-14 to 2017	2.99	100 ml	Sebizole
METRONIDAZOLE			
Gel 0.75%			
MICONAZOLE NITRATE			
Crm 2% – 1% DV Mar-15 to 2017	0.55	15 g	Multichem
➔ Lotn 2% – Restricted: For continuation only			
Tinc 2%			
NYSTATIN			
Crm 100,000 u per g			
Antiparasitics			
MALATHION [MALDISON]			
Lotn 0.5%			
Shampoo 1%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MALATHION WITH PERMETHRIN AND PIPERONYL BUTOXIDE			
Spray 0.25% with permethrin 0.5% and piperonyl butoxide 2%			
PERMETHRIN			
Crm 5% – 1% DV Apr-15 to 2017	4.20	30 g	Lyderm
Lotn 5% – 1% DV Sep-14 to 2017	3.19	30 ml	A-Scabies

Antiacne Preparations

ADAPALENE			
Crm 0.1%			
Gel 0.1%			
BENZOYL PEROXIDE			
Soln 5%			
ISOTRETINOIN			
Cap 10 mg	12.47	100	Isotane 10
	14.96	120	Oratane
Cap 20 mg	19.27	100	Isotane 20
	23.12	120	Oratane
TRETINOIN			
Crm 0.05%			

Antipruritic Preparations

CALAMINE			
Crm, aqueous, BP – 1% DV Dec-15 to 2018	1.49	100 g	Pharmacy Health
Lotn, BP – 1% DV Dec-15 to 2018	12.94	2,000 ml	PSM
CROTAMITON			
Crm 10% – 1% DV Sep-15 to 2018	3.37	20 g	Itch-Soothe

Barrier Creams and Emollients

Barrier Creams

DIMETHICONE			
Crm 5% tube – 1% DV Apr-14 to 2016	1.65	100 g	healthE Dimethicone 5%
Crm 5% pump bottle – 1% DV Apr-14 to 2016	4.73	500 ml	healthE Dimethicone 5%
Crm 10% pump bottle – 1% DV Nov-15 to 2018	4.90	500 ml	healthE Dimethicone 10%
ZINC			
Crm			<i>e.g. Zinc Cream (Orion); Zinc Cream (PSM)</i>
Oint			<i>e.g. Zinc oxide (PSM)</i>
Paste			
ZINC AND CASTOR OIL			
Crm	1.63	20 g	Orion
Oint, BP – 1% DV Jul-15 to 2017	1.39	20 g	healthE

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ZINC WITH WOOL FAT			
Crn zinc 15.25% with wool fat 4%			<i>e.g. Sudocrem</i>
Emollients			
AQUEOUS CREAM			
Crn 100 g – 1% DV Jan-16 to 2018	1.00	100 g	Pharmacy Health SLS-free
Note: DV limit applies to the pack sizes of 100 g or less.			
Crn 500 g – 1% DV Mar-16 to 2018	1.99	500 g	AFT SLS-free
Note: DV limit applies to the pack sizes of greater than 100 g.			
CETOMACROGOL			
Crn BP, 500 g – 1% DV Nov-15 to 2018	2.74	500 g	healthE
Crn BP, 100 g – 1% DV Jan-16 to 2018	1.47	1	healthE
CETOMACROGOL WITH GLYCEROL			
Crn 90% with glycerol 10%,	2.00	100 g	Pharmacy Health
	2.10		Pharmacy Health
	3.20		healthE
Crn 90% with glycerol 10%	4.50	500 ml	Pharmacy Health
			Sorbolene with Glycerin
	6.50	1,000 ml	Pharmacy Health
			Sorbolene with Glycerin
Crn 90% with glycerol 10%, 500 ml, 1 bottle	5.46	1	healthE
EMULSIFYING OINTMENT			
Oint BP – 1% DV Apr-15 to 2017	1.84	100 g	Jaychem
Note: DV limit applies to pack sizes of less than 200 g.			
Oint BP, 500 g – 1% DV Jul-15 to 2017	2.73	500 g	AFT
Note: DV limit applies to pack sizes of greater than 200 g.			
GLYCEROL WITH PARAFFIN			
Crn glycerol 10% with white soft paraffin 5% and liquid paraffin 10%			<i>e.g. QV cream</i>
OIL IN WATER EMULSION			
Crn	2.63	500 g	healthE Fatty Cream
Crn, 100 g	1.60	1	healthE Fatty Cream
PARAFFIN			
Oint liquid paraffin 50% with white soft paraffin 50%	3.10	100 g	healthE
White soft – 1% DV Sep-15 to 2018	0.85	10 g	healthE
Note: DV limit applies to pack sizes of 30 g or less, and to both white soft paraffin and yellow soft paraffin.			
Yellow soft			
PARAFFIN WITH WOOL FAT			
Lotn liquid paraffin 15.9% with wool fat 0.6%			<i>e.g. AlphaKeri;BK ;DP; Hydroderm Lotn</i>
Lotn liquid paraffin 91.7% with wool fat 3%			<i>e.g. Alpha Keri Bath Oil</i>
UREA			
Crn 10%			
WOOL FAT			
Crn			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Corticosteroids			
BETAMETHASONE DIPROPIONATE			
Crm 0.05%			
Oint 0.05%			
BETAMETHASONE VALERATE			
Crm 0.1% – 1% DV Jun-15 to 2018	3.15	50 g	Beta Cream
Oint 0.1% – 1% DV Jun-15 to 2018	3.15	50 g	Beta Ointment
Lotn 0.1%			
CLOBETASOL PROPIONATE			
Crm 0.05% – 1% DV Jul-15 to 2016	3.20	30 g	Clobetasol BNM
Oint 0.05% – 1% DV Jul-15 to 2016	3.20	30 g	Clobetasol BNM
CLOBETASONE BUTYRATE			
Crm 0.05%			
DIFLUCORTOLONE VALERATE – Restricted: For continuation only			
➡ Crm 0.1%			
➡ Fatty oint 0.1%			
HYDROCORTISONE			
Crm 1%, 100 g	3.75	100 g	Pharmacy Health
Crm 1%, 500 g	14.00	500 g	Pharmacy Health
HYDROCORTISONE ACETATE			
Crm 1%	2.48	14.2 g	AFT
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN			
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – 1% DV Dec-14 to 2017	10.57	250 ml	DP Lotn HC
HYDROCORTISONE BUTYRATE			
Crm 0.1%	2.30	30 g	Locoid Lipocream
	6.85	100 g	Locoid Lipocream
Oint 0.1%	6.85	100 g	Locoid
Milky emul 0.1%	6.85	100 ml	Locoid Crelo
HYDROCORTISONE WITH PARAFFIN AND WOOL FAT			
Lotn 1% with paraffin liquid 15.9% and wool fat 0.6%			
METHYLPREDNISOLONE ACEPONATE			
Crm 0.1%	4.95	15 g	Advantan
Oint 0.1%	4.95	15 g	Advantan
MOMETASONE FUROATE			
Crm 0.1% – 1% DV Nov-15 to 2018	1.51	15 g	Elocon Alcohol Free
	2.90	50 g	Elocon Alcohol Free
Oint 0.1% – 1% DV Nov-15 to 2018	1.51	15 g	Elocon
	2.90	50 g	Elocon
Lotn 0.1% – 1% DV Sep-15 to 2018	7.35	30 ml	Elocon
TRIAMCINOLONE ACETONIDE			
Crm 0.02% – 1% DV Apr-15 to 2017	6.30	100 g	Aristocort
Oint 0.02% – 1% DV Apr-15 to 2017	6.35	100 g	Aristocort

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Corticosteroids with Anti-Infective Agents

BETAMETHASONE VALERATE WITH CLIOQUINOL – **Restricted** see terms below

¶ Crm 0.1% with clioquinol 3%

➔ Restricted

Initiation

Either:

- 1 For the treatment of intertrigo; or
- 2 For continuation use.

BETAMETHASONE VALERATE WITH FUSIDIC ACID

Crm 0.1% with fusidic acid 2%

HYDROCORTISONE WITH MICONAZOLE

Crm 1% with miconazole nitrate 2% – **1% DV Sep-15 to 2018**2.00 15 g **Micreme H**

HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN

Crm 1% with natamycin 1% and neomycin sulphate 0.5%2.79 15 g Pimafucort

Oint 1% with natamycin 1% and neomycin sulphate 0.5%2.79 15 g Pimafucort

TRIAMCINOLONE ACETONIDE WITH NEOMYCIN SULPHATE, GRAMICIDIN AND NYSTATIN

Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and
gramicidin 250 mcg per g

Psoriasis and Eczema Preparations

ACITRETIN

Cap 10 mg – **1% DV Nov-14 to 2017**17.86 60 **Novatretin**

Cap 25 mg – **1% DV Nov-14 to 2017**41.36 60 **Novatretin**

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

Gel 500 mcg with calcipotriol 50 mcg per g – **1% DV Sep-15 to 2018**26.12 30 g **Daivobet**

Oint 500 mcg with calcipotriol 50 mcg per g – **1% DV Sep-15 to 2018**26.12 30 g **Daivobet**

CALCIPOTRIOL

Crm 50 mcg per g45.00 100 g Daivonex

Oint 50 mcg per g45.00 100 g Daivonex

Soln 50 mcg per ml16.00 30 ml Daivonex

COAL TAR WITH SALICYLIC ACID AND SULPHUR

Oint 12% with salicylic acid 2% and sulphur 4%

METHOXSALEN [8-METHOXYPORALEN]

Tab 10 mg

Lotn 1.2%

PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN

Soln 2.3% with trolamine laurilsulfate and fluorescein sodium3.36 500 ml Pinetarsol

5.82 1,000 ml Pinetarsol

POTASSIUM PERMANGANATE

Tab 400 mg

Crystals

Scalp Preparations

BETAMETHASONE VALERATE

Scalp app 0.1%7.75 100 ml Beta Scalp

DERMATOLOGICALS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
CLOBETASOL PROPIONATE			
Scalp app 0.05%	6.96	30 ml	Dermol
HYDROCORTISONE BUTYRATE			
Scalp lotn 0.1%	3.65	100 ml	Locoid

Wart Preparations

IMIQUIMOD			
Crm 5%, 250 mg sachet – 1% DV Feb-15 to 2017	17.98	12	Apo-Imiquimod Cream 5%
PODOPHYLLOTOXIN			
Soln 0.5%	33.60	3.5 ml	Condyline
SILVER NITRATE			
Sticks with applicator			

Other Skin Preparations

DIPHEMANIL METILSULFATE			
Powder 2%			
SUNSCREEN, PROPRIETARY			
Crm			
Lotn	3.30	100 g	Marine Blue Lotion SPF 50+
	5.10	200 g	Marine Blue Lotion SPF 50+

Antineoplastics

FLUOROURACIL SODIUM			
Crm 5% – 1% DV Sep-15 to 2018	8.95	20 g	Efudix
METHYL AMINOLEVULINATE HYDROCHLORIDE – Restricted see terms below			
☞ Crm 16%			
☞ Restricted			
Dermatologist or plastic surgeon			

Wound Management Products

CALCIUM GLUCONATE			
Gel 2.5%	21.00	1	healthE

Price (ex man. excl. GST)	Brand or Generic
\$	Per Manufacturer

Anti-Infective Agents

ACETIC ACID

- Soln 3%
- Soln 5%

ACETIC ACID WITH HYDROXYQUINOLINE, GLYCEROL AND RICINOLEIC ACID

- Jelly 0.94% with hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator

CHLORHEXIDINE GLUCONATE

- | | | | |
|---|------|------|----------------|
| Crm 1% – 1% DV Sep-15 to 2018 | 1.21 | 50 g | healthE |
| Lotn 1%, 200 ml – 1% DV Sep-15 to 2018 | 2.98 | 1 | healthE |

CLOTRIMAZOLE

- | | | | |
|--|------|------|-----------------|
| Vaginal crm 1% with applicator – 1% DV Dec-13 to 2016 | 1.45 | 35 g | Clomazol |
| Vaginal crm 2% with applicator – 1% DV Dec-13 to 2016 | 2.20 | 20 g | Clomazol |

MICONAZOLE NITRATE

- | | | | |
|--|------|------|----------------|
| Vaginal crm 2% with applicator – 1% DV Oct-14 to 2017 | 3.95 | 40 g | Micreme |
|--|------|------|----------------|

NYSTATIN

- Vaginal crm 100,000 u per 5 g with applicator(s)

Contraceptives

Antiandrogen Oral Contraceptives

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

- | | | | |
|--|------|-----|--------------|
| Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tablets – 1% DV Dec-14 to 2017 | 5.36 | 168 | Ginet |
|--|------|-----|--------------|

Combined Oral Contraceptives

ETHINYLOESTRADIOL WITH DESOGESTREL

- Tab 20 mcg with desogestrel 150 mcg
- Tab 30 mcg with desogestrel 150 mcg

ETHINYLOESTRADIOL WITH LEVONORGESTREL

- | | | | |
|--|------|----|------------------|
| Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets | 2.65 | 84 | Ava 20 ED |
| Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets | 2.30 | 84 | Ava 30 ED |
| Tab 20 mcg with levonorgestrel 100 mcg | | | |
| Tab 30 mcg with levonorgestrel 150 mcg | | | |
| Tab 50 mcg with levonorgestrel 125 mcg | 9.45 | 84 | Microgynon 50 ED |

ETHINYLOESTRADIOL WITH NORETHISTERONE

- Tab 35 mcg with norethisterone 1 mg
- Tab 35 mcg with norethisterone 500 mcg

NORETHISTERONE WITH MESTRANOL

- Tab 1 mg with mestranol 50 mcg

Contraceptive Devices

INTRA-UTERINE DEVICE

- | | | | |
|--|-------|---|-----------------------|
| IUD 29.1 mm length × 23.2 mm width | 31.60 | 1 | Choice TT380 Short |
| IUD 33.6 mm length × 29.9 mm width | 31.60 | 1 | Choice TT380 Standard |

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Emergency Contraception			
LEVONORGESTREL			
Tab 1.5 mg – 1% DV Jul-13 to 2016	3.50	1	Postinor-1
Progestogen-Only Contraceptives			
LEVONORGESTREL			
Tab 30 mcg			
Subdermal implant (2 × 75 mg rods) – 5% DV Oct-14 to 31 Dec 2017	133.65	1	Jadelle
↳ Intra-uterine system, 20 mcg per day			e.g. Mirena
➡ Restricted			
Initiation — heavy menstrual bleeding			
Obstetrician or gynaecologist			
All of the following:			
1 The patient has a clinical diagnosis of heavy menstrual bleeding; and			
2 The patient has failed to respond to or is unable to tolerate other appropriate pharmaceutical therapies as per the Heavy Menstrual Bleeding Guidelines; and			
3 Any of the following:			
3.1 Serum ferritin level < 16 mcg/l (within the last 12 months); or			
3.2 Haemoglobin level < 120 g/l; or			
3.3 The patient has had a uterine ultrasound and either a hysteroscopy or endometrial biopsy.			
Continuation — heavy menstrual bleeding			
Obstetrician or gynaecologist			
Either:			
1 Patient demonstrated clinical improvement of heavy menstrual bleeding; or			
2 Previous insertion was removed or expelled within 3 months of insertion.			
Initiation — endometriosis			
Obstetrician or gynaecologist			
The patient has a clinical diagnosis of endometriosis confirmed by laparoscopy.			
Continuation — endometriosis			
Obstetrician or gynaecologist			
Either:			
1 Patient demonstrated satisfactory management of endometriosis; or			
2 Previous insertion was removed or expelled within 3 months of insertion.			
Note: endometriosis is an unregistered indication.			
MEDROXYPROGESTERONE ACETATE			
Inj 150 mg per ml, 1 ml syringe – 1% DV Sep-13 to 2016	7.00	1	Depo-Provera
NORETHISTERONE			
Tab 350 mcg – 1% DV Oct-15 to 2018	6.25	84	Noriday 28
Obstetric Preparations			
Antiprogestogens			
MIFEPRISTONE			
Tab 200 mg			
Oxytocics			
CARBOPROST TROMETAMOL			
Inj 250 mcg per ml, 1 ml ampoule			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DINOPROSTONE			
Pessaries 10 mg			
Vaginal gel 1 mg in 3 g	52.65	1	Prostin E2
Vaginal gel 2 mg in 3 g	64.60	1	Prostin E2
ERGOMETRINE MALEATE			
Inj 500 mcg per ml, 1 ml ampoule – 1% DV Oct-14 to 2017	94.70	5	DBL Ergometrine
OXYTOCIN			
Inj 5 iu per ml, 1 ml ampoule – 1% DV Nov-15 to 2018	4.03	5	Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule – 1% DV Nov-15 to 2018	5.03	5	Oxytocin BNM
OXYTOCIN WITH ERGOMETRINE MALEATE			
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampoule – 1% DV Sep-15 to 2018	11.13	5	Syntometrine

Tocolytics

PROGESTERONE – Restricted see terms below

☞ Cap 100 mg 16.50 30 Utrogestan

☞ **Restricted**

Initiation

Gynaecologist or obstetrician

Both:

- 1 For the prevention of pre-term labour*; and
- 2 Either:
 - 2.1 The patient has a short cervix on ultrasound (defined as < 25mm at 16 to 28 weeks); or
 - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are Unapproved Indications (refer to Section A: General Rules, Part I (Interpretations and Definitions) and Part IV (Miscellaneous Provisions) rule 23.1)

TERBUTALINE – Restricted see terms below

☞ Inj 500 mcg ampoule

☞ **Restricted**

Obstetrician

Oestrogens

OESTRIOL

Crm 1 mg per g with applicator

Pessaries 500 mcg

Urologicals

5-Alpha Reductase Inhibitors

FINASTERIDE – Restricted see terms below

☞ Tab 5 mg – **1% DV Dec-14 to 2017** 2.08 30 **Finpro**

☞ **Restricted**

Initiation

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Alpha-1A Adrenoceptor Blockers			
TAMSULOSIN – Restricted see terms below			
⚡ Cap 400 mcg – 1% DV Dec-13 to 2016	13.51	100	Tamsulosin-Rex
➡ Restricted			
Initiation			
Both:			
1 Patient has symptomatic benign prostatic hyperplasia; and			
2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.			
Urinary Alkalisers			
POTASSIUM CITRATE – Restricted see terms below			
⚡ Oral liq 3 mmol per ml	30.00	200 ml	Biomed
➡ Restricted			
Initiation			
Both:			
1 The patient has recurrent calcium oxalate urolithiasis; and			
2 The patient has had more than two renal calculi in the two years prior to the application.			
SODIUM CITRO-TARTRATE			
Grans eff 4 g sachets – 1% DV Feb-15 to 2017	2.93	28	Ural
Urinary Antispasmodics			
OXYBUTYNIN			
Tab 5 mg – 1% DV Jun-13 to 2016	11.20	500	Apo-Oxybutynin
Oral liq 5 mg per 5 ml – 1% DV Jun-13 to 2016	56.45	473 ml	Apo-Oxybutynin
SOLIFENACIN SUCCINATE – Restricted see terms below			
⚡ Tab 5 mg	37.50	30	Vesicare
⚡ Tab 10 mg	37.50	30	Vesicare
➡ Restricted			
Initiation			
Patient has overactive bladder and a documented intolerance of, or is non-responsive to, oxybutynin.			
TOLTERODINE TARTRATE – Restricted see terms below			
⚡ Tab 1 mg	14.56	56	Arrow-Tolterodine
⚡ Tab 2 mg	14.56	56	Arrow-Tolterodine
➡ Restricted			
Initiation			
Patient has overactive bladder and a documented intolerance of, or is non-responsive to, oxybutynin.			

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Anabolic Agents

OXANDROLONE

☯ Tab 2.5 mg

☞ **Restricted**

Initiation

For the treatment of burns patients.

Androgen Agonists and Antagonists

CYPROTERONE ACETATE

Tab 50 mg – 1% DV Oct-15 to 2018 15.87 50 **Procur**

Tab 100 mg – 1% DV Oct-15 to 2018 30.40 50 **Procur**

TESTOSTERONE

Patch 2.5 mg per day 80.00 60 Androderm

TESTOSTERONE CYPIONATE

Inj 100 mg per ml, 10 ml vial – 1% DV Sep-14 to 2017 76.50 1 **Depo-Testosterone**

TESTOSTERONE ESTERS

Inj testosterone decanoate 100 mg, testosterone isocarproate 60 mg,
testosterone phenylpropionate 60 mg and testosterone propionate
30 mg per ml, 1 ml ampoule

TESTOSTERONE UNDECANOATE

Cap 40 mg – 1% DV Sep-15 to 2018 16.80 60 **Andriol Testocaps**

Inj 250 mg per ml, 4 ml vial 86.00 1 **Reandron 1000**

Calcium Homeostasis

CALCITONIN

Inj 100 iu per ml, 1 ml ampoule – 1% DV Oct-14 to 2017 121.00 5 **Miacalcic**

ZOLEDRONIC ACID

☯ Inj 4 mg per 5 ml, vial 550.00 1 **Zometa**

☞ **Restricted**

Initiation

Oncologist, haematologist or palliative care specialist

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
 - 2.1 Patient has bone metastases or involvement; and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
 - 3.1 Patient has bone metastases or involvement; and
 - 3.2 Patient is at risk of skeletal-related events (pathological fracture, spinal cord compression, radiation to bone or surgery to bone).

Corticosteroids

BETAMETHASONE

Tab 500 mcg

Inj 4 mg per ml, 1 ml ampoule

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE

Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml ampoule

HORMONE PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DEXAMETHASONE			
Tab 0.5 mg – 1% DV Jan-16 to 2018.....	0.88	30	Dexamethsone
Tab 4 mg – 1% DV Jan-16 to 2018.....	1.84	30	Dexamethsone
Oral liq 1 mg per ml	45.00	25 ml	Biomed
DEXAMETHASONE PHOSPHATE			
Inj 4 mg per ml, 1 ml ampoule – 1% DV Apr-14 to 2016	14.19	10	Max Health
Max Health brand - HSS with 1% DV will apply 1 July 2016 to 30 June 2019.			
Inj 4 mg per ml, 2 ml ampoule – 1% DV Apr-14 to 2016	12.59	5	Max Health
Max Health brand - HSS with 1% DV will apply 1 July 2016 to 30 June 2019.			
FLUDROCORTISONE ACETATE			
Tab 100 mcg	14.32	100	Florinef
HYDROCORTISONE			
Tab 5 mg – 1% DV Sep-15 to 2018	8.10	100	Douglas
Tab 20 mg – 1% DV Sep-15 to 2018	20.32	100	Douglas
Inj 100 mg vial – 1% DV Oct-13 to 2016	4.99	1	Solu-Cortef
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)			
Tab 4 mg – 1% DV Oct-15 to 2018	80.00	100	Medrol
Tab 100 mg – 1% DV Oct-15 to 2018	180.00	20	Medrol
Inj 40 mg vial – 1% DV Oct-15 to 2018	10.50	1	Solu-Medrol
Inj 125 mg vial – 1% DV Oct-15 to 2018	22.25	1	Solu-Medrol
Inj 500 mg vial – 1% DV Oct-15 to 2018	9.00	1	Solu-Medrol
Inj 1 g vial – 1% DV Oct-15 to 2018	16.00	1	Solu-Medrol
METHYLPREDNISOLONE ACETATE			
Inj 40 mg per ml, 1 ml vial – 1% DV Oct-15 to 2018	40.00	5	Depo-Medrol
METHYLPREDNISOLONE ACETATE WITH LIDOCAINE [LIGNOCAINE]			
Inj 40 mg with lidocaine [lignocaine], 1 ml vial – 1% DV Oct-15 to 2018	9.25	1	Depo-Medrol with Lidocaine
PREDNISOLONE			
Oral liq 5 mg per ml	7.50	30 ml	Redipred
Enema 200 mcg per ml, 100 ml			
PREDNISONE			
Tab 1 mg	10.68	500	Apo-Prednisone
	2.13	100	Apo-Prednisone S29
Tab 2.5 mg	12.09	500	Apo-Prednisone
Tab 5 mg	11.09	500	Apo-Prednisone
Tab 20 mg	29.03	500	Apo-Prednisone
TRIAMCINOLONE ACETONIDE			
Inj 10 mg per ml, 1 ml ampoule – 1% DV Apr-15 to 2017	20.80	5	Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule – 1% DV Apr-15 to 2017	51.70	5	Kenacort-A 40
TRIAMCINOLONE HEXACETONIDE			
Inj 20 mg per ml, 1 ml vial			

Price (ex man. excl. GST)		Brand or Generic
\$	Per	Manufacturer

Hormone Replacement Therapy

Oestrogens

OESTRADIOL

- Tab 1 mg
- Tab 2 mg
- Patch 25 mcg per day
- Patch 50 mcg per day
- Patch 100 mcg per day

OESTRADIOL VALERATE

- | | | | |
|---------------------------------------|-------|----|------------------|
| Tab 1 mg – 1% DV Jun-15 to 2018 | 12.36 | 84 | Progynova |
| Tab 2 mg – 1% DV Jun-15 to 2018 | 12.36 | 84 | Progynova |

OESTROGENS (CONJUGATED EQUINE)

- Tab 300 mcg
- Tab 625 mcg

Progestogen and Oestrogen Combined Preparations

OESTRADIOL WITH NORETHISTERONE ACETATE

- Tab 1 mg with 0.5 mg norethisterone acetate
- Tab 2 mg with 1 mg norethisterone acetate
- Tab 2 mg with 1 mg norethisterone acetate (10), and tab 2 mg oestradiol (12) and tab 1 mg oestradiol (6)

OESTROGENS WITH MEDROXYPROGESTERONE ACETATE

- Tab 625 mcg conjugated equine with 2.5 mg medroxyprogesterone acetate
- Tab 625 mcg conjugated equine with 5 mg medroxyprogesterone acetate

Progestogens

MEDROXYPROGESTERONE ACETATE

- | | | | |
|---|-------|-----|----------------|
| Tab 2.5 mg – 1% DV Sep-13 to 2016 | 3.09 | 30 | Provera |
| Tab 5 mg – 1% DV Sep-13 to 2016 | 13.06 | 100 | Provera |
| Tab 10 mg – 1% DV Sep-13 to 2016 | 6.85 | 30 | Provera |

Other Endocrine Agents

CABERGOLINE – Restricted see terms below

- | | | | |
|---|-------|---|-----------------|
| ⚡ Tab 0.5 mg – 1% DV Sep-15 to 2018 | 4.75 | 2 | Dostinex |
| | 19.00 | 8 | Dostinex |

↪ Restricted

Initiation

Any of the following:

- 1 Inhibition of lactation; or
- 2 Patient has pathological hyperprolactinemia; or
- 3 Patient has acromegaly.

CLOMIPHENE CITRATE

- | | | | |
|--|-------|----|------------------|
| Tab 50 mg – 1% DV Sep-13 to 2016 | 29.84 | 10 | Serophene |
|--|-------|----|------------------|

HORMONE PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DANAZOL			
Cap 100 mg	68.33	100	Azol
Cap 200 mg	97.83	100	Azol
GESTRINONE			
Cap 2.5 mg			
METYRAPONE			
Cap 250 mg			
PENTAGASTRIN			
Inj 250 mcg per ml, 2 ml ampoule			

Other Oestrogen Preparations

ETHINYLOESTRADIOL			
Tab 10 mcg – 1% DV Sep-15 to 2018	17.60	100	NZ Medical & Scientific
OESTRADIOL			
Implant 50 mg			
OESTRIOL			
Tab 2 mg			

Other Progestogen Preparations

MEDROXYPROGESTERONE			
Tab 100 mg – 1% DV Sep-13 to 2016	96.50	100	Provera
NORETHISTERONE			
Tab 5 mg – 1% DV Jun-15 to 2018	18.29	100	Primolut N

Pituitary and Hypothalamic Hormones and Analogues

CORTICOTRORELIN (OVINE)			
Inj 100 mcg vial			
THYROTROPIN ALFA			
Inj 900 mcg vial			

Adrenocorticotrophic Hormones

TETRACOSACTIDE [TETRACOSACTRIN]			
Inj 250 mcg per ml, 1 ml ampoule	17.71	1	Synacthen
Inj 1 mg per ml, 1 ml ampoule	29.56	1	Synacthen Depot

GnRH Agonists and Antagonists

BUSERELIN			
Inj 1 mg per ml, 5.5 ml vial			
GONADORELIN			
Inj 100 mcg vial			
GOSERELIN			
Implant 3.6 mg	166.20	1	Zoladex
Implant 10.8 mg	443.76	1	Zoladex

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LEUPRORELIN ACETATE			
Inj 3.75 mg syringe	221.60	1	Lucrin Depot PDS
Inj 7.5 mg syringe	166.20	1	Eligard
Inj 11.25 mg syringe	591.68	1	Lucrin Depot PDS
Inj 22.5 mg syringe	443.76	1	Eligard
Inj 30 mg syringe	1,109.40	1	Lucrin Depot PDS
Inj 30 mg vial	591.68	1	Eligard
Inj 45 mg syringe	832.05	1	Eligard

Gonadotrophins

CHORIOGONADOTROPIN ALFA

Inj 250 mcg in 0.5 ml syringe

Growth Hormone

SOMATROPIN – **Restricted** see terms below

⚡ Inj 5 mg cartridge – 1% DV Jan-15 to 31 Dec 2017	109.50	1	Omnitrope
⚡ Inj 10 mg cartridge – 1% DV Jan-15 to 31 Dec 2017	219.00	1	Omnitrope
⚡ Inj 15 mg cartridge – 1% DV Jan-15 to 31 Dec 2017	328.50	1	Omnitrope

↪ Restricted

Initiation — growth hormone deficiency in children

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age; and adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Continuation — growth hormone deficiency in children

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 A current bone age is ≤ 14 years (female patients) or ≤ 16 years (male patients); and
- 2 Height velocity is ≥ 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is ≥ 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation — Turner syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Continuation — Turner syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 Height velocity \geq 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is \geq 2 cm per year, calculated over six months; and
- 3 A current bone age is \leq 14 years; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initiation — short stature without growth hormone deficiency

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Continuation — short stature without growth hormone deficiency

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 Height velocity is \geq 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is \geq 2 cm per year as calculated over six months; and
- 3 Current bone age is \leq 14 years (female patients) or \leq 16 years (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initiation — short stature due to chronic renal insufficiency

Endocrinologist, paediatric endocrinologist or renal physician on the recommendation of a endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is \leq to 14 years (female patients) or \leq to 16 years (male patients); and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a $GFR \leq 30 \text{ ml/min/1.73 m}^2$ as measured by the Schwartz method ($\text{Height(cm)/plasma creatinine (umol/l)} \times 40 = \text{corrected GFR (ml/min/1.73 m}^2)$) in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received $< 5 \text{ mg/m}^2$ /day of prednisone or equivalent for at least 6 months.

Continuation — short stature due to chronic renal insufficiency

Endocrinologist, paediatric endocrinologist or renal physician on the recommendation of an endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 Height velocity is ≥ 50 th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is $\geq 2 \text{ cm}$ per year as calculated over six months; and
- 3 A current bone age is ≤ 14 years (female patients) or ≤ 16 years (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initiation — Prader-Willi syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient's height velocity is < 25 th percentile for bone age adjusted for bone age/pubertal status if appropriate as calculated over 6 to 12 months using the standards of Tanner and Davies (1985) or pubertal status over 6 to 12 months; and
- 3 Either:
 - 3.1 The patient is under two years of age and height velocity has been assessed over a minimum six month period from the age of 12 months, with at least three supine length measurements over this period demonstrating clear and consistent evidence of linear growth failure (with height velocity < 25 th percentile); or
 - 3.2 The patient is aged two years or older; and
- 4 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 5 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 6 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by ≥ 0.5 standard deviations in the preceding 12 months.

Continuation — Prader-Willi syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Height velocity is \geq 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is \geq 2 cm per year as calculated over six months; and
- 3 A current bone age is \leq 14 years (female patients) or \leq 16 years (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by \geq 0.5 standard deviations in the preceding 12 months.

Initiation — adults and adolescents

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA[®]).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of \leq 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of \leq 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until it is within 1 standard deviation of the mean normal value for age and sex; and

The dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Continuation — adults and adolescents

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

Either:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in the Quality of Life Assessment defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA[®]) score from baseline; and
 - 1.3 Serum IGF-I levels have increased to within ± 1 SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin does not exceed 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA[®] score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ± 1 SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients.

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Thyroid and Antithyroid Preparations

CARBIMAZOLE

Tab 5 mg

IODINE

Soln BP 50 mg per ml

LEVOTHYROXINE

Tab 25 mcg

Tab 50 mcg

Tab 100 mcg

LIOTHYRONINE SODIUM

⚡ Tab 20 mcg

➡ **Restricted**

Initiation

For a maximum of 14 days' treatment in patients with thyroid cancer who are due to receive radioiodine therapy.

Inj 20 mcg vial

POTASSIUM IODATE

Tab 170 mg

POTASSIUM PERCHLORATE

Cap 200 mg

PROPYLTHIOURACIL – **Restricted** see terms below

⚡ Tab 50 mg 35.00 100 PTU

➡ **Restricted**

Initiation

Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Note: Propylthiouracil is not recommended for patients under the age of 18 years unless the patient is pregnant and other treatments are contraindicated.

PROTIRELIN

Inj 100 mcg per ml, 2 ml ampoule

Vasopressin Agents

ARGIPRESSIN [VASOPRESSIN]

Inj 20 u per ml, 1 ml ampoule

DESMOPRESSIN ACETATE – **Some items restricted** see terms below

⚡ Tab 100 mcg – 1% DV Jun-16 to 2019	25.00	30	Minirin
⚡ Tab 200 mcg – 1% DV Jun-16 to 2019	54.45	30	Minirin
Nasal spray 10 mcg per dose – 1% DV Sep-14 to 2017	22.95	6 ml	Desmopressin-PH&T
Inj 4 mcg per ml, 1 ml ampoule			
Inj 15 mcg per ml, 1 ml ampoule			
Nasal drops 100 mcg per ml			

➡ **Restricted**

Initiation — Nocturnal enuresis

Either:

- 1 The nasal forms of desmopressin are contraindicated; or
- 2 An enuresis alarm is contraindicated.

Note: Cranial diabetes insipidus and the nasal forms of desmopressin are contraindicated.

HORMONE PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
TERLIPRESSIN			
Inj 0.1 mg per ml, 8.5 ml ampoule	450.00	5	Glypressin
Inj 1 mg per 8.5 ml ampoule – 1% DV Jun-15 to 2018	215.00	5	Glypressin

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Antibacterials			
Aminoglycosides			
AMIKACIN – Restricted see terms below			
☞ Inj 5 mg per ml, 10 ml syringe			
☞ Inj 5 mg per ml, 5 ml syringe	176.00	10	Biomed
☞ Inj 15 mg per ml, 5 ml syringe			
☞ Inj 250 mg per ml, 2 ml vial – 1% DV Oct-14 to 2017	431.20	5	DBL Amikacin
☞ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
GENTAMICIN SULPHATE			
Inj 10 mg per ml, 1 ml ampoule	8.56	5	Hospira
Inj 10 mg per ml, 2 ml ampoule	175.10	25	APP Pharmaceuticals
Inj 40 mg per ml, 2 ml ampoule – 1% DV Sep-15 to 2018	6.00	10	Pfizer
PAROMOMYCIN – Restricted see terms below			
☞ Cap 250 mg	126.00	16	Humatin
☞ Restricted			
Clinical microbiologist or infectious disease specialist			
STREPTOMYCIN SULPHATE – Restricted see terms below			
☞ Inj 400 mg per ml, 2.5 ml ampoule			
☞ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
TOBRAMYCIN			
☞ Powder			
☞ Restricted			
Initiation			
For addition to orthopaedic bone cement.			
☞ Inj 40 mg per ml, 2 ml vial	38.00	5	DBL Tobramycin
☞ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
☞ Inj 100 mg per ml, 5 ml vial			
☞ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
☞ Solution for inhalation 60 mg per ml, 5 ml	2,200.00	56 dose	TOBI
☞ Restricted			
Initiation			
Patient has cystic fibrosis.			
Carbapenems			
ERTAPENEM – Restricted see terms below			
☞ Inj 1 g vial	73.50	1	Invanz
☞ Restricted			
Clinical microbiologist or infectious disease specialist			
IMIPENEM WITH CILASTATIN – Restricted see terms below			
☞ Inj 500 mg with 500 mg cilastatin vial – 1% DV Jun-15 to 2017	13.79	1	Imipenem+Cilastatin RBX
☞ Restricted			
Clinical microbiologist or infectious disease specialist			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MEROPENEM – Restricted see terms below			
‡ Inj 500 mg vial – 1% DV Oct-14 to 2017	35.22	10	DBL Meropenem
‡ Inj 1 g vial – 1% DV Oct-14 to 2017	65.21	10	DBL Meropenem

➡Restricted

Clinical microbiologist or infectious disease specialist

Cephalosporins and Cephamycins - 1st Generation
CEFALEXIN

Cap 500 mg – 1% DV Oct-13 to 2016	5.70	20	Cephalexin ABM
Grans for oral liq 25 mg per ml – 1% DV Sep-15 to 2018	8.00	100 ml	Cefalexin Sandoz
Grans for oral liq 50 mg per ml – 1% DV Sep-15 to 2018	11.00	100 ml	Cefalexin Sandoz

CEFAZOLIN

Inj 500 mg vial – 1% DV Sep-14 to 2017	3.99	5	AFT
Inj 1 g vial – 1% DV Sep-14 to 2017	3.38	5	AFT

Cephalosporins and Cephamycins - 2nd Generation
CEFACLOR

Cap 250 mg – 1% DV Dec-13 to 2016	26.00	100	Ranbaxy-Cefaclor
Grans for oral liq 25 mg per ml – 1% DV Dec-13 to 2016	3.53	100 ml	Ranbaxy-Cefaclor

CEFOXITIN

Inj 1 g vial – 1% DV Jan-16 to 2018	58.00	10	Cefoxitin Actavis
---	-------	----	--------------------------

CEFUROXIME

Tab 250 mg	29.40	50	Zinnat
Inj 750 mg vial	3.70	5	Zinacef
Inj 1.5 g vial	1.30	1	Zinacef

Cephalosporins and Cephamycins - 3rd Generation
CEFOTAXIME

Inj 500 mg vial	1.90	1	Cefotaxime Sandoz
Inj 1 g vial – 1% DV Oct-14 to 2017	17.10	10	DBL Cefotaxime

CEFTAZIDIME – Restricted see terms below

‡ Inj 500 mg vial – 1% DV Jan-15 to 2017	5.30	1	Fortum
‡ Inj 1 g vial – 1% DV Jan-15 to 2017	1.55	1	Fortum
‡ Inj 2 g vial – 1% DV Jan-15 to 2017	3.34	1	Fortum

➡Restricted

Clinical microbiologist, infectious disease specialist or respiratory specialist

CEFTRIAXONE

Inj 500 mg vial – 1% DV Mar-14 to 2016	1.50	1	Ceftriaxone-AFT
Inj 1 g vial – 1% DV Mar-14 to 2016	5.22	5	Ceftriaxone-AFT
Inj 2 g vial – 1% DV Mar-14 to 2016	2.75	1	Ceftriaxone-AFT

Cephalosporins and Cephamycins - 4th Generation
CEFEPIME – Restricted see terms below

‡ Inj 1 g vial – 1% DV Oct-15 to 2018	3.95	1	Cefepime-AFT
‡ Inj 2 g vial – 1% DV Oct-15 to 2018	6.92	1	Cefepime-AFT

➡Restricted

Clinical microbiologist or infectious disease specialist

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Cephalosporins and Cephamycins - 5th Generation			
CEFTAROLINE FOSAMIL – Restricted see terms below			
⚡ Inj 600 mg vial	1,450.00	10	Zinforo
➔ Restricted			
Initiation — multi-resistant organism salvage therapy			
Clinical microbiologist or infectious disease specialist			
Either:			
1 for patients where alternative therapies have failed; or			
2 for patients who have a contraindication or hypersensitivity to standard current therapies.			
Macrolides			
AZITHROMYCIN – Restricted see terms below			
⚡ Tab 250 mg – 1% DV Sep-15 to 2018	9.00	30	Apo-Azithromycin
⚡ Tab 500 mg – 1% DV Sep-15 to 2018	1.05	2	Apo-Azithromycin
⚡ Grans for oral liq 200 mg per 5 ml (40 mg per ml) – 1% DV Oct-15 to 2018	12.50	15 ml	Zithromax
➔ Restricted			
Initiation			
Any of the following:			
1 Patient has received a lung transplant and requires treatment or prophylaxis for bronchiolitis obliterans syndrome; or			
2 Patient has cystic fibrosis and has chronic infection with <i>Pseudomonas aeruginosa</i> or <i>Pseudomonas</i> related gram negative organisms; or			
3 For any other condition for five days' treatment, with review after five days.			
CLARITHROMYCIN – Restricted see terms below			
⚡ Tab 250 mg – 1% DV Sep-14 to 2017	3.98	14	Apo-Clarithromycin
⚡ Tab 500 mg – 1% DV Sep-14 to 2017	10.40	14	Apo-Clarithromycin
⚡ Grans for oral liq 50 mg per ml	23.12	50 ml	Klacid
⚡ Grans for oral liq 25 mg per ml	23.12	70 ml	Klacid
⚡ Inj 500 mg vial – 1% DV Mar-15 to 2017	20.40	1	Martindale
<i>(Klacid Grans for oral liq 25 mg per ml to be delisted 1 October 2016)</i>			
➔ Restricted			
Initiation — Tab 250 mg and oral liquid			
Either:			
1 Atypical mycobacterial infection; or			
2 <i>Mycobacterium tuberculosis</i> infection where there is drug resistance or intolerance to standard pharmaceutical agents.			
Initiation — Tab 500 mg			
<i>Helicobacter pylori</i> eradication.			
Initiation — Infusion			
Any of the following:			
1 Atypical mycobacterial infection; or			
2 <i>Mycobacterium tuberculosis</i> infection where there is drug resistance or intolerance to standard pharmaceutical agents; or			
3 Community-acquired pneumonia.			
ERYTHROMYCIN (AS ETHYLSUCCINATE)			
Tab 400 mg	16.95	100	E-Mycin
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	E-Mycin
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	E-Mycin
ERYTHROMYCIN (AS LACTOBIONATE)			
Inj 1 g vial	16.00	1	Erythrocin IV

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ERYTHROMYCIN (AS STEARATE) – Restricted: For continuation only			
➔ Tab 250 mg			
➔ Tab 500 mg			
ROXITHROMYCIN			
Tab 150 mg	7.48	50	Arrow-Roxithromycin
Tab 300 mg	14.40	50	Arrow-Roxithromycin
Penicillins			
AMOXICILLIN			
Cap 250 mg – 1% DV Mar-14 to 2016	16.18	500	Apo-Amoxi
Cap 500 mg – 1% DV Jul-14 to 2016	20.94	500	Apo-Amoxi
Grans for oral liq 125 mg per 5 ml	0.88	100 ml	Amoxicillin Actavis
.....	2.00		Ospamox
Grans for oral liq 250 mg per 5 ml	0.97	100 ml	Amoxicillin Actavis
.....	2.00		Ospamox
Inj 250 mg vial – 1% DV Oct-14 to 2017	10.67	10	Ibiamox
Inj 500 mg vial – 1% DV Oct-14 to 2017	12.41	10	Ibiamox
Inj 1 g vial – 1% DV Oct-14 to 2017	17.29	10	Ibiamox
AMOXICILLIN WITH CLAVULANIC ACID			
Tab 500 mg with clavulanic acid 125 mg	1.95	20	Augmentin
.....	9.75	100	Curam Duo
Grans for oral liq 25 mg with clavulanic acid 6.25 mg per ml	3.83	100 ml	Augmentin
Grans for oral liq 50 mg with clavulanic acid 12.5 mg per ml	4.97	100 ml	Augmentin
Inj 500 mg with clavulanic acid 100 mg vial – 1% DV Sep-15 to 2018	10.14	10	m-Amoxiclav
Inj 1,000 mg with clavulanic acid 200 mg vial – 1% DV Sep-15 to 2018	12.80	10	m-Amoxiclav
BENZATHINE BENZYL PENICILLIN			
Inj 900 mg (1.2 million units) in 2.3 ml syringe – 1% DV Sep-15 to 2018	315.00	10	Bicillin LA
BENZYL PENICILLIN SODIUM [PENICILLIN G]			
Inj 600 mg (1 million units) vial – 1% DV Sep-14 to 2017	10.35	10	Sandoz
FLUCLOXACILLIN			
Cap 250 mg – 1% DV Sep-15 to 2018	18.70	250	Staphlex
Cap 500 mg – 1% DV Sep-15 to 2018	62.90	500	Staphlex
Grans for oral liq 25 mg per ml – 1% DV Sep-15 to 2018	2.29	100 ml	AFT
Grans for oral liq 50 mg per ml – 1% DV Sep-15 to 2018	3.08	100 ml	AFT
Inj 250 mg vial – 1% DV Sep-14 to 2017	8.80	10	Flucloxin
Inj 500 mg vial – 1% DV Sep-14 to 2017	9.20	10	Flucloxin
Inj 1 g vial – 1% DV Jan-16 to 2017	11.60	10	Flucloxin
PHENOXYMETHYL PENICILLIN [PENICILLIN V]			
Cap 250 mg – 1% DV Jun-15 to 2018	2.88	50	Cilicaine VK
Cap 500 mg – 1% DV Jun-15 to 2018	4.73	50	Cilicaine VK
Grans for oral liq 125 mg per 5 ml – 1% DV Apr-14 to 2016	1.64	100 ml	AFT
Grans for oral liq 250 mg per 5 ml – 1% DV Apr-14 to 2016	1.74	100 ml	AFT
PIPERACILLIN WITH TAZOBACTAM – Restricted see terms below			
⚡ Inj 4 g with tazobactam 0.5 g vial	5.84	1	Hospira
➔ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
PROCAINE PENICILLIN			
Inj 1.5 g in 3.4 ml syringe – 1% DV Sep-14 to 2017	123.50	5	Cilicaine

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

TICARCILLIN WITH CLAVULANIC ACID – **Restricted** see terms below

⚡ Inj 3 g with clavulanic acid 0.1 mg vial

➔ **Restricted**

Clinical microbiologist, infectious disease specialist or respiratory specialist

Quinolones

CIPROFLOXACIN – **Restricted** see terms below

⚡ Tab 250 mg – 1% DV Sep-14 to 2017	1.75	28	Cipflo
⚡ Tab 500 mg – 1% DV Sep-14 to 2017	2.00	28	Cipflo
⚡ Tab 750 mg – 1% DV Sep-14 to 2017	3.75	28	Cipflo
⚡ Oral liq 50 mg per ml			
⚡ Oral liq 100 mg per ml			
⚡ Inj 2 mg per ml, 100 ml bag – 1% DV Mar-16 to 2018	30.58	10	Cipflo

➔ **Restricted**

Clinical microbiologist or infectious disease specialist

MOXIFLOXACIN – **Restricted** see terms below

⚡ Tab 400 mg	52.00	5	Avelox
⚡ Inj 1.6 mg per ml, 250 ml bottle	70.00	1	Avelox IV 400

➔ **Restricted**

Initiation — Mycobacterium infection

Infectious disease specialist, clinical microbiologist or respiratory specialist

Either:

- 1 Both:
 - 1.1 Active tuberculosis; and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.

Initiation — Pneumonia

Infectious disease specialist or clinical microbiologist

Either:

- 1 Immunocompromised patient with pneumonia that is unresponsive to first-line treatment; or
- 2 Pneumococcal pneumonia or other invasive pneumococcal disease highly resistant to other antibiotics.

Initiation — Penetrating eye injury

Ophthalmologist

Five days treatment for patients requiring prophylaxis following a penetrating eye injury.

Initiation — Mycoplasma genitalium

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium; and
- 2 Has tried and failed to clear infection using azithromycin; and
- 3 Treatment is only for 7 days.

NORFLOXACIN

Tab 400 mg – 1% DV Sep-14 to 2017	13.50	100	Arrow-Norfloxac
---	-------	-----	------------------------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Tetracyclines			
DEMECLOCYCLINE HYDROCHLORIDE			
Tab 150 mg			
Cap 150 mg			
Cap 300 mg			
DOXYCYCLINE			
➔ Tab 50 mg – Restricted: For continuation only			
Tab 100 mg – 1% DV Sep-14 to 2017	6.75	250	Doxine
Inj 5 mg per ml, 20 ml vial			
MINOCYCLINE			
Tab 50 mg			
➔ Cap 100 mg – Restricted: For continuation only			
TETRACYCLINE			
Tab 250 mg			
Cap 500 mg	46.00	30	Tetracyclin Wolff
TIGECYCLINE – Restricted see terms below			
⚡ Inj 50 mg vial			
➔ Restricted			
Clinical microbiologist or infectious disease specialist			
Other Antibacterials			
AZTREONAM – Restricted see terms below			
⚡ Inj 1 g vial	131.00	5	Azactam
➔ Restricted			
Clinical microbiologist or infectious disease specialist			
CHLORAMPHENICOL – Restricted see terms below			
⚡ Inj 1 g vial			
➔ Restricted			
Clinical microbiologist or infectious disease specialist			
CLINDAMYCIN – Restricted see terms below			
⚡ Cap 150 mg – 1% DV Oct-13 to 2016	5.80	16	Clindamycin ABM
⚡ Oral liq 15 mg per ml			
⚡ Inj 150 mg per ml, 4 ml ampoule – 1% DV Sep-13 to 2016	100.00	10	Dalacin C
➔ Restricted			
Clinical microbiologist or infectious disease specialist			
COLISTIN SULPHOMETHATE [COLESTIMETHATE] – Restricted see terms below			
⚡ Inj 150 mg per ml, 1 ml vial	65.00	1	Colistin-Link
➔ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
DAPTOMYCIN – Restricted see terms below			
⚡ Inj 350 mg vial – 1% DV Sep-15 to 2018	175.16	1	Cubicin
⚡ Inj 500 mg vial – 1% DV Sep-15 to 2018	243.52	1	Cubicin
➔ Restricted			
Clinical microbiologist or infectious disease specialist			
FOSFOMYCIN – Restricted see terms on the next page			
⚡ Powder for oral solution, 3 g sachet			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Clinical microbiologist or infectious disease specialist			
FUSIDIC ACID – Restricted see terms below			
⌚ Tab 250 mg	34.50	12	Fucidin
➔Restricted			
Clinical microbiologist or infectious disease specialist			
HEXAMINE HIPPURATE			
Tab 1 g			
LINCOMYCIN – Restricted see terms below			
⌚ Inj 300 mg per ml, 2 ml vial			
➔Restricted			
Clinical microbiologist or infectious disease specialist			
LINEZOLID – Restricted see terms below			
⌚ Tab 600 mg – 1% DV Sep-15 to 2018	800.00	10	Zyvox
⌚ Oral liq 20 mg per ml – 1% DV Sep-15 to 2018	775.00	150 ml	Zyvox
⌚ Inj 2 mg per ml, 300 ml bag – 1% DV Sep-15 to 2018	1,650.00	10	Zyvox
➔Restricted			
Clinical microbiologist or infectious disease specialist			
NITROFURANTOIN			
Tab 50 mg			
Tab 100 mg			
PIVMECILLINAM – Restricted see terms below			
⌚ Tab 200 mg			
➔Restricted			
Clinical microbiologist or infectious disease specialist			
SULPHADIAZINE – Restricted see terms below			
⌚ Tab 500 mg			
➔Restricted			
Clinical microbiologist, infectious disease specialist or maternal-foetal medicine specialist			
TEICOPLANIN – Restricted see terms below			
⌚ Inj 400 mg vial			
➔Restricted			
Clinical microbiologist or infectious disease specialist			
TRIMETHOPRIM			
Tab 100 mg			
Tab 300 mg – 1% DV Oct-15 to 2018	15.00	50	TMP
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXAZOLE]			
Tab 80 mg with sulphamethoxazole 400 mg			
Oral liq 8 mg with sulphamethoxazole 40 mg per ml	2.15	100 ml	Deprim
Inj 16 mg with sulphamethoxazole 80 mg per ml, 5 ml ampoule			
VANCOMYCIN – Restricted see terms below			
⌚ Inj 500 mg vial – 1% DV Oct-14 to 2017	2.64	1	Mylan
➔Restricted			
Clinical microbiologist or infectious disease specialist			

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Antifungals

Imidazoles

KETOCONAZOLE

☞ Tab 200 mg

☞ **Restricted**

Oncologist

Polyene Antimycotics

AMPHOTERICIN B

☞ Inj (liposomal) 50 mg vial – 1% DV Sep-15 to 20183,450.00 10 **AmBisome**

☞ **Restricted**

Initiation

Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist

Either:

- 1 Proven or probable invasive fungal infection, to be prescribed under an established protocol; or
- 2 Both:
 - 2.1 Possible invasive fungal infection; and
 - 2.2 A multidisciplinary team (including an infectious disease physician or a clinical microbiologist) considers the treatment to be appropriate.

☞ Inj 50 mg vial

☞ **Restricted**

Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist

NYSTATIN

Tab 500,000 u	17.09	50	Nilstat
Cap 500,000 u	15.47	50	Nilstat

Triazoles

FLUCONAZOLE – **Restricted** see terms below

☞ Cap 50 mg – 1% DV Nov-14 to 2017.....	3.49	28	Ozole
☞ Cap 150 mg – 1% DV Nov-14 to 2017	0.71	1	Ozole
☞ Cap 200 mg – 1% DV Nov-14 to 2017.....	9.69	28	Ozole
☞ Oral liquid 50 mg per 5 ml	98.50	35 ml	Diflucan
☞ Inj 2 mg per ml, 50 ml vial – 1% DV Oct-13 to 2016.....	4.95	1	Fluconazole-Clarix
☞ Inj 2 mg per ml, 100 ml vial – 1% DV Oct-13 to 2016	6.47	1	Fluconazole-Clarix

☞ **Restricted**

Consultant

ITRACONAZOLE – **Restricted** see terms below

☞ Cap 100 mg – 1% DV Oct-13 to 2016	2.99	15	Itrazole
☞ Oral liquid 10 mg per ml			

☞ **Restricted**

Clinical immunologist, clinical microbiologist, dermatologist or infectious disease specialist

POSACONAZOLE – **Restricted** see terms on the next page

☞ Oral liq 40 mg per ml	761.13	105 ml	Noxafil
-------------------------------	--------	--------	---------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Initiation			
Haematologist or infectious disease specialist			
<i>Re-assessment required after 6 weeks</i>			
Both:			
1 Either:			
1.1 Patient has acute myeloid leukaemia; or			
1.2 Patient is planned to receive a stem cell transplant and is at high risk for aspergillus infection; and			
2 Patient is to be treated with high dose remission induction therapy or re-induction therapy.			
Continuation			
Haematologist or infectious disease specialist			
<i>Re-assessment required after 6 weeks</i>			
Both:			
1 Patient has previously received posaconazole prophylaxis during remission induction therapy; and			
2 Any of the following:			
2.1 Patient is to be treated with high dose remission re-induction therapy; or			
2.2 Patient is to be treated with high dose consolidation therapy; or			
2.3 Patient is receiving a high risk stem cell transplant.			
VORICONAZOLE – Restricted see terms below			
⚡ Tab 50 mg – 1% DV Jan-16 to 2018	130.00	56	Vttack
⚡ Tab 200 mg – 1% DV Jan-16 to 2018	500.00	56	Vttack
⚡ Oral liq 40 mg per ml	730.00	70 ml	Vfend
⚡ Inj 200 mg vial	185.00	1	Vfend
➔Restricted			
Initiation — Proven or probable aspergillus infection			
Clinical microbiologist, haematologist or infectious disease specialist			
Both:			
1 Patient is immunocompromised; and			
2 Patient has proven or probable invasive aspergillus infection.			
Initiation — Possible aspergillus infection			
Clinical microbiologist, haematologist or infectious disease specialist			
All of the following:			
1 Patient is immunocompromised; and			
2 Patient has possible invasive aspergillus infection; and			
3 A multidisciplinary team (including an infectious disease physician) considers the treatment to be appropriate.			
Initiation — Resistant candidiasis infections and other moulds			
Clinical microbiologist, haematologist or infectious disease specialist			
All of the following:			
1 Patient is immunocompromised; and			
2 Either:			
2.1 Patient has fluconazole resistant candidiasis; or			
2.2 Patient has mould strain such as <i>Fusarium</i> spp. and <i>Scedosporium</i> spp; and			
3 A multidisciplinary team (including an infectious disease physician or clinical microbiologist) considers the treatment to be appropriate.			
Other Antifungals			
CASPOFUNGIN – Restricted see terms on the next page			
⚡ Inj 50 mg vial	667.50	1	Cancidas
⚡ Inj 70 mg vial	862.50	1	Cancidas

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➡Restricted			
Initiation			
Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist			
Either:			
1	Proven or probable invasive fungal infection, to be prescribed under an established protocol; or		
2	Both:		
2.1	Possible invasive fungal infection; and		
2.2	A multidisciplinary team (including an infectious disease physician or a clinical microbiologist) considers the treatment to be appropriate.		
FLUCYTOSINE – Restricted see terms below			
⚡	Cap 500 mg		
➡Restricted			
Clinical microbiologist or infectious disease specialist			
TERBINAFINE			
Tab 250 mg – 1% DV Sep-14 to 2017	1.50	14	Dr Reddy's Terbinafine
Antimycobacterials			
Antileprotics			
CLOFAZIMINE – Restricted see terms below			
⚡	Cap 50 mg		
➡Restricted			
Clinical microbiologist, dermatologist or infectious disease specialist			
DAPSONE – Restricted see terms below			
⚡ Tab 25 mg – 1% DV Sep-14 to 2017	95.00	100	Dapsone
⚡ Tab 100 mg – 1% DV Sep-14 to 2017	110.00	100	Dapsone
➡Restricted			
Clinical microbiologist, dermatologist or infectious disease specialist			
Antituberculotics			
CYCLOSERINE – Restricted see terms below			
⚡	Cap 250 mg		
➡Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
ETHAMBUTOL HYDROCHLORIDE – Restricted see terms below			
⚡ Tab 100 mg	48.01	56	Myambutol
⚡ Tab 400 mg	49.34	56	Myambutol
➡Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
ISONIAZID – Restricted see terms below			
⚡ Tab 100 mg – 1% DV Sep-15 to 2018	20.00	100	PSM
➡Restricted			
Clinical microbiologist, dermatologist, paediatrician, public health physician or internal medicine physician			
ISONIAZID WITH RIFAMPICIN – Restricted see terms below			
⚡ Tab 100 mg with rifampicin 150 mg – 1% DV Sep-15 to 2018	85.54	100	Rifinah
⚡ Tab 150 mg with rifampicin 300 mg – 1% DV Sep-15 to 2018	170.60	100	Rifinah
➡Restricted			
Clinical microbiologist, dermatologist, paediatrician, public health physician or internal medicine physician			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PARA-AMINOSALICYLIC ACID – Restricted see terms below			
¶ Grans for oral liq 4 g	280.00	30	Paser
➔ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
PROTIONAMIDE – Restricted see terms below			
¶ Tab 250 mg	305.00	100	Peteha
➔ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
PYRAZINAMIDE – Restricted see terms below			
¶ Tab 500 mg			
➔ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
RIFABUTIN – Restricted see terms below			
¶ Cap 150 mg – 1% DV Sep-13 to 2016.....	213.19	30	Mycobutin
➔ Restricted			
Clinical microbiologist, gastroenterologist, infectious disease specialist or respiratory specialist			
RIFAMPICIN – Restricted see terms below			
¶ Cap 150 mg – 1% DV Nov-14 to 2017.....	55.75	100	Rifadin
¶ Cap 300 mg – 1% DV Nov-14 to 2017.....	116.25	100	Rifadin
¶ Oral liq 100 mg per 5 ml – 1% DV Nov-14 to 2017.....	12.00	60 ml	Rifadin
¶ Inj 600 mg vial – 1% DV Nov-14 to 2017.....	128.85	1	Rifadin
➔ Restricted			
Clinical microbiologist, dermatologist, internal medicine physician, paediatrician or public health physician			

Antiparasitics

Anthelmintics

ALBENDAZOLE – Restricted see terms below			
¶ Tab 200 mg			
¶ Tab 400 mg			
➔ Restricted			
Clinical microbiologist or infectious disease specialist			
IVERMECTIN – Restricted see terms below			
¶ Tab 3 mg	17.20	4	Stromectol
➔ Restricted			
Clinical microbiologist, dermatologist or infectious disease specialist			
MEBENDAZOLE			
Tab 100 mg	24.19	24	De-Worm
Oral liq 100 mg per 5 ml			
PRAZQUANTEL			
Tab 600 mg			

Antiprotozoals

ARTEMETHER WITH LUMEFANTRINE – Restricted see terms below			
¶ Tab 20 mg with lumefantrine 120 mg			
➔ Restricted			
Clinical microbiologist or infectious disease specialist			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ARTESUNATE – Restricted see terms below			
☞ Inj 60 mg vial			
☞ Restricted			
Clinical microbiologist or infectious disease specialist			
ATOVAQUONE WITH PROGUANIL HYDROCHLORIDE – Restricted see terms below			
☞ Tab 62.5 mg with proguanil hydrochloride 25 mg – 1% DV Nov-14 to 2017	25.00	12	Malarone Junior
☞ Tab 250 mg with proguanil hydrochloride 100 mg – 1% DV Nov-14 to 2017	64.00	12	Malarone
☞ Restricted			
Clinical microbiologist or infectious disease specialist			
CHLOROQUINE PHOSPHATE – Restricted see terms below			
☞ Tab 250 mg			
☞ Restricted			
Clinical microbiologist, dermatologist, infectious disease specialist or rheumatologist			
MEFLOQUINE – Restricted see terms below			
☞ Tab 250 mg – 1% DV Dec-14 to 2017	33.48	8	Lariam
☞ Restricted			
Clinical microbiologist, dermatologist, infectious disease specialist or rheumatologist			
METRONIDAZOLE			
Tab 200 mg	10.45	100	Trichozole
Tab 400 mg	18.15	100	Trichozole
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	Flagyl-S
Inj 5 mg per ml, 100 ml bag – 1% DV Apr-15 to 2017	6.94	5	AFT
Suppos 500 mg	24.48	10	Flagyl
NITAZOXANIDE – Restricted see terms below			
☞ Tab 500 mg	1,680.00	30	Alinia
☞ Oral liq 100 mg per 5 ml			
☞ Restricted			
Clinical microbiologist or infectious disease specialist			
ORNIDAZOLE			
Tab 500 mg	16.50	10	Arrow-Ornidazole
PENTAMIDINE ISETHIONATE – Restricted see terms below			
☞ Inj 300 mg vial – 1% DV Mar-15 to 2017	180.00	5	Pentacarinat
☞ Restricted			
Clinical microbiologist or infectious disease specialist			
PRIMAQUINE PHOSPHATE – Restricted see terms below			
☞ Tab 7.5 mg			
☞ Restricted			
Clinical microbiologist or infectious disease specialist			
PYRIMETHAMINE – Restricted see terms below			
☞ Tab 25 mg			
☞ Restricted			
Clinical microbiologist, infectious disease specialist or maternal-foetal medicine specialist			
QUININE DIHYDROCHLORIDE – Restricted see terms on the next page			
☞ Inj 60 mg per ml, 10 ml ampoule			
☞ Inj 300 mg per ml, 2 ml vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Clinical microbiologist or infectious disease specialist			
QUININE SULPHATE			
Tab 300 mg	54.06	500	Q 300

SODIUM STIBOGLUCONATE – **Restricted** see terms below

⚡ Inj 100 mg per ml, 1 ml vial

➔Restricted

Clinical microbiologist or infectious disease specialist

SPIRAMYCIN – **Restricted** see terms below

⚡ Tab 500 mg

➔Restricted

Maternal-foetal medicine specialist

Antiretrovirals

HIV Fusion Inhibitors

ENFUVIRTIDE – **Restricted** see terms below

⚡ Inj 108 mg vial × 60 2,380.00 1 Fuzeon

➔Restricted

Initiation

Re-assessment required after 12 months

All of the following:

- 1 Confirmed HIV infection; and
- 2 Enfuvirtide to be given in combination with optimized background therapy (including at least 1 other antiretroviral drug that the patient has never previously been exposed to) for treatment failure; and
- 3 Either:
 - 3.1 Patient has evidence of HIV replication, despite ongoing therapy; or
 - 3.2 Patient has treatment-limiting toxicity to previous antiretroviral agents; and
- 4 Previous treatment with 3 different antiretroviral regimens has failed; and
- 5 All of the following:
 - 5.1 Previous treatment with a non-nucleoside reverse transcriptase inhibitor has failed; and
 - 5.2 Previous treatment with a nucleoside reverse transcriptase inhibitor has failed; and
 - 5.3 Previous treatment with a protease inhibitor has failed.

Continuation

Patient has had at least a 10-fold reduction in viral load at 12 months.

Non-Nucleoside Reverse Transcriptase Inhibitors

➔Restricted

Initiation — Confirmed HIV

Both:

- 1 Confirmed HIV infection; and
- 2 Any of the following:
 - 2.1 Symptomatic patient; or
 - 2.2 Patient aged 12 months and under; or
 - 2.3 Both:
 - 2.3.1 Patient aged 1 to 5 years; and
 - 2.3.2 Any of the following:
 - 2.3.2.1 CD4 counts < 1000 cells/mm³; or
 - 2.3.2.2 CD4 counts < 0.25 × total lymphocyte count; or

continued...

continued...

2.3.2.3 Viral load counts > 100000 copies per ml; or

2.4 Both:

2.4.1 Patient aged 6 years and over; and

2.4.2 CD4 counts < 500 cells/mm³.

Initiation — Prevention of maternal transmission

Either:

1 Prevention of maternal foetal transmission; or

2 Treatment of the newborn for up to eight weeks.

Initiation — Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

1 Treatment course to be initiated within 72 hours post exposure; and

2 Any of the following:

2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or

2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or

2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initiation — Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

EFAVIRENZ — **Restricted** see terms on the preceding page

⬆ Tab 50 mg – 1% DV Sep-15 to 2018	63.38	30	Stocrin
--	-------	----	----------------

⬆ Tab 200 mg – 1% DV Sep-15 to 2018	190.15	90	Stocrin
---	--------	----	----------------

⬆ Tab 600 mg – 1% DV Sep-15 to 2018	63.38	30	Stocrin
---	-------	----	----------------

⬆ Oral liq 30 mg per ml			
-------------------------------	--	--	--

ETRAVIRINE — **Restricted** see terms on the preceding page

⬆ Tab 200 mg	770.00	60	Intelence
--------------------	--------	----	-----------

NEVIRAPINE — **Restricted** see terms on the preceding page

⬆ Tab 200 mg – 1% DV Nov-15 to 2018	65.00	60	Nevirapine Alphapharm
---	-------	----	------------------------------

⬆ Oral suspension 10 mg per ml	134.55	240 ml	Viramune Suspension
--------------------------------------	--------	--------	---------------------

Nucleoside Reverse Transcriptase Inhibitors

➡Restricted

Initiation — Confirmed HIV

Both:

1 Confirmed HIV infection; and

2 Any of the following:

2.1 Symptomatic patient; or

2.2 Patient aged 12 months and under; or

2.3 Both:

2.3.1 Patient aged 1 to 5 years; and

2.3.2 Any of the following:

2.3.2.1 CD4 counts < 1000 cells/mm³; or

2.3.2.2 CD4 counts < 0.25 × total lymphocyte count; or

2.3.2.3 Viral load counts > 100000 copies per ml; or

2.4 Both:

2.4.1 Patient aged 6 years and over; and

2.4.2 CD4 counts < 500 cells/mm³.

Initiation — Prevention of maternal transmission

Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued. . .			
1 Prevention of maternal foetal transmission; or			
2 Treatment of the newborn for up to eight weeks.			
Initiation — Post-exposure prophylaxis following non-occupational exposure to HIV			
Both:			
1 Treatment course to be initiated within 72 hours post exposure; and			
2 Any of the following:			
2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or			
2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or			
2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.			
Initiation — Percutaneous exposure			
Patient has percutaneous exposure to blood known to be HIV positive.			
ABACAVIR SULPHATE – Restricted see terms on the preceding page			
⬆ Tab 300 mg – 1% DV Oct-14 to 2017	229.00	60	Ziagen
⬆ Oral liq 20 mg per ml – 1% DV Oct-14 to 2017.....	256.31	240 ml	Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE – Restricted see terms on the preceding page			
⬆ Tab 600 mg with lamivudine 300 mg	630.00	30	Kivexa
DIDANOSINE [DDI] – Restricted see terms on the preceding page			
⬆ Cap 125 mg			
⬆ Cap 200 mg			
⬆ Cap 250 mg			
⬆ Cap 400 mg			
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPROXIL FUMARATE – Restricted see terms on the preceding page			
⬆ Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil fumarate 300 mg	1,313.19	30	Atripla
EMTRICITABINE – Restricted see terms on the preceding page			
⬆ Cap 200 mg	307.20	30	Emtriva
EMTRICITABINE WITH TENOFOVIR DISOPROXIL FUMARATE – Restricted see terms on the preceding page			
⬆ Tab 200 mg with tenofovir disoproxil fumarate 300 mg	838.20	30	Truvada
LAMIVUDINE – Restricted see terms on the preceding page			
⬆ Oral liq 10 mg per ml			
STAVUDINE – Restricted see terms on the preceding page			
⬆ Cap 30 mg			
⬆ Cap 40 mg			
⬆ Powder for oral soln 1 mg per ml			
ZIDOVUDINE [AZT] – Restricted see terms on the preceding page			
⬆ Cap 100 mg – 1% DV Oct-13 to 2016	152.25	100	Retrovir
⬆ Oral liq 10 mg per ml – 1% DV Oct-13 to 2016.....	30.45	200 ml	Retrovir
⬆ Inj 10 mg per ml, 20 ml vial – 1% DV Oct-14 to 2017.....	750.00	5	Retrovir IV
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Restricted see terms on the preceding page			
⬆ Tab 300 mg with lamivudine 150 mg – 1% DV Sep-14 to 2017	44.00	60	Alphapharm

Protease Inhibitors

➔ Restricted

Initiation — Confirmed HIV

Both:

- 1 Confirmed HIV infection; and
- 2 Any of the following:
 - 2.1 Symptomatic patient; or
 - 2.2 Patient aged 12 months and under; or
 - 2.3 Both:
 - 2.3.1 Patient aged 1 to 5 years; and
 - 2.3.2 Any of the following:
 - 2.3.2.1 CD4 counts < 1000 cells/mm³; or
 - 2.3.2.2 CD4 counts < 0.25 × total lymphocyte count; or
 - 2.3.2.3 Viral load counts > 100000 copies per ml; or
 - 2.4 Both:
 - 2.4.1 Patient aged 6 years and over; and
 - 2.4.2 CD4 counts < 500 cells/mm³.

Initiation — Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Initiation — Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initiation — Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

ATAZANAVIR SULPHATE – Restricted see terms above

⚡ Cap 150 mg	568.34	60	Reyataz
⚡ Cap 200 mg	757.79	60	Reyataz

DARUNAVIR – Restricted see terms above

⚡ Tab 400 mg	837.50	60	Prezista
⚡ Tab 600 mg	1,190.00	60	Prezista

INDINAVIR – Restricted see terms above

- ⚡ Cap 200 mg
- ⚡ Cap 400 mg

LOPINAVIR WITH RITONAVIR – Restricted see terms above

⚡ Tab 100 mg with ritonavir 25 mg	183.75	60	Kaletra
⚡ Tab 200 mg with ritonavir 50 mg	735.00	120	Kaletra
⚡ Oral liq 80 mg with ritonavir 20 mg per ml	735.00	300 ml	Kaletra

RITONAVIR – Restricted see terms above

⚡ Tab 100 mg	43.31	30	Norvir
⚡ Oral liq 80 mg per ml			

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Strand Transfer Inhibitors

➔Restricted

Initiation — Confirmed HIV

Both:

- 1 Confirmed HIV infection; and
- 2 Any of the following:
 - 2.1 Symptomatic patient; or
 - 2.2 Patient aged 12 months and under; or
 - 2.3 Both:
 - 2.3.1 Patient aged 1 to 5 years; and
 - 2.3.2 Any of the following:
 - 2.3.2.1 CD4 counts < 1000 cells/mm³; or
 - 2.3.2.2 CD4 counts < 0.25 × total lymphocyte count; or
 - 2.3.2.3 Viral load counts > 100000 copies per ml; or
 - 2.4 Both:
 - 2.4.1 Patient aged 6 years and over; and
 - 2.4.2 CD4 counts < 500 cells/mm³.

Initiation — Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Initiation — Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initiation — Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

RALTEGRAVIR POTASSIUM – **Restricted** see terms above

⬆ Tab 400 mg	1,090.00	60	Isentress
--------------------	----------	----	-----------

Antivirals

Hepatitis B

ADEFOVIR DIPIVOXIL – **Restricted** see terms below

⬆ Tab 10 mg	670.00	30	Hepsera
-------------------	--------	----	---------

➔Restricted

Initiation

Gastroenterologist or infectious disease specialist

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg+); and Documented resistance to lamivudine defined as:
- 2 Patient has raised serum ALT (> 1 × ULN); and
- 3 Patient has HBV DNA greater than 100,000 copies per mL, or viral load ≥ 10-fold over nadir; and
- 4 Detection of M204I or M204V mutation; and
- 5 Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
5.1 Both:			
5.1.1 Patient is cirrhotic; and			
5.1.2 Adefovir dipivoxil to be used in combination with lamivudine; or			
5.2 Both:			
5.2.1 Patient is not cirrhotic; and			
5.2.2 Adefovir dipivoxil to be used as monotherapy.			
ENTECAVIR – Restricted see terms below			
⚡ Tab 0.5 mg	400.00	30	Baraclude
➡ Restricted			
Initiation			
Gastroenterologist or infectious disease specialist			
All of the following:			
1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and			
2 Patient is Hepatitis B nucleoside analogue treatment-naïve; and			
3 Entecavir dose 0.5 mg/day; and			
4 Either:			
4.1 ALT greater than upper limit of normal; or			
4.2 Bridging fibrosis or cirrhosis (Metavir stage 3 or greater or moderate fibrosis) on liver histology; and			
5 Either:			
5.1 HBeAg positive; or			
5.2 Patient has ≥ 2,000 IU HBV DNA units per ml and fibrosis (Metavir stage 2 or greater) on liver histology; and			
6 No continuing alcohol abuse or intravenous drug use; and			
7 Not co-infected with HCV, HIV or HDV; and			
8 Neither ALT nor AST greater than 10 times upper limit of normal; and			
9 No history of hypersensitivity to entecavir; and			
10 No previous documented lamivudine resistance (either clinical or genotypic).			
LAMIVUDINE – Restricted see terms below			
⚡ Tab 100 mg – 1% DV Nov-14 to 2017	6.00	28	Zeffix
⚡ Oral liq 5 mg per ml – 1% DV Nov-14 to 2017	270.00	240 ml	Zeffix
➡ Restricted			
Initiation			
Gastroenterologist, infectious disease specialist, paediatrician or general physician			
<i>Limited to 12 months treatment</i>			
Any of the following:			
1 HBV DNA positive cirrhosis prior to liver transplantation; or			
2 HBsAg positive and have had a liver, kidney, heart, lung or bone marrow transplant; or			
3 Hepatitis B virus naïve patient who has received a liver transplant from an anti-HBc (Hepatitis B core antibody) positive donor; or			
4 Hepatitis B surface antigen positive (HbsAg) patient who is receiving chemotherapy for a malignancy, or who has received such treatment within the previous two months; or			
5 Hepatitis B surface antigen positive patient who is receiving anti tumour necrosis factor treatment; or			
6 Hepatitis B core antibody (anti-HBc) positive patient who is receiving rituximab plus high dose steroids (e.g. R-CHOP).			
Continuation — patients who have maintained continuous treatment and response to lamivudine			
Gastroenterologist, infectious disease specialist, paediatrician or general physician			
<i>Re-assessment required after 2 years</i>			
All of the following:			
1 Have maintained continuous treatment with lamivudine; and			
2 Most recent test result shows continuing biochemical response (normal ALT); and			
continued...			
⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below) e.g. Brand indicates brand example only. It is not a contracted product.			

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 3 HBV DNA <100,000 copies per ml by quantitative PCR at a reference laboratory.

Continuation — when given in combination with adefovir dipivoxil for patients with cirrhosis and resistance to lamivudine

Gastroenterologist, infectious disease specialist, paediatrician or general physician

Re-assessment required after 2 years

All of the following:

- 1 Lamivudine to be used in combination with adefovir dipivoxil; and
- 2 Patient is cirrhotic; and
Documented resistance to lamivudine defined as:
- 3 All of the following:
 - 3.1 Patient has raised serum ALT ($> 1 \times \text{ULN}$); and
 - 3.2 Patient has HBV DNA greater than 100,000 copies per mL, or viral load ≥ 10 -fold over nadir; and
 - 3.3 Detection of M204I or M204V mutation.

Continuation — when given in combination with adefovir dipivoxil for patients with resistance to adefovir dipivoxil

Gastroenterologist, infectious disease specialist, paediatrician or general physician

Re-assessment required after 2 years

Both:

- 1 Lamivudine to be used in combination with adefovir dipivoxil; and
Documented resistance to lamivudine defined as:
- 2 All of the following:
 - 2.1 Patient has raised serum ALT ($> 1 \times \text{ULN}$); and
 - 2.2 Patient has HBV DNA greater than 100,000 copies per mL, or viral load ≥ 10 -fold over nadir; and
 - 2.3 Detection of N236T or A181T/V mutation.

TENOFOVIR DISOPROXIL FUMARATE – **Restricted** see terms below

⚡ Tab 300 mg	531.00	30	Viread
--------------------	--------	----	--------

➡ **Restricted**

Initiation — Confirmed hepatitis B

Any of the following:

- 1 All of the following:
 - 1.1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
 - 1.2 Patient has had previous lamivudine, adefovir or entecavir therapy; and
 - 1.3 HBV DNA greater than 20,000 IU/mL or increased ≤ 10 -fold over nadir; and
 - 1.4 Any of the following:
 - 1.4.1 Lamivudine resistance - detection of M204I/V mutation; or
 - 1.4.2 Adefovir resistance - detection of A181T/V or N236T mutation; or
 - 1.4.3 Entecavir resistance - detection of relevant mutations including I169T, L180M T184S/A/I/L/G/C/M, S202C/G/I, M204V or M250I/V mutation; or
- 2 Patient is either listed or has undergone liver transplantation for HBV; or
- 3 Patient has a decompensated cirrhosis with a Mayo score > 20 .

Initiation — Pregnant or Breastfeeding, Active hepatitis B

Limited to 12 months treatment

Both:

- 1 Patient is HBsAg positive and pregnant; and
- 2 HBV DNA $> 20,000$ IU/mL and ALT $> \text{ULN}$.

Initiation — Pregnant, prevention of vertical transmission

Limited to 6 months treatment

Both:

- 1 Patient is HBsAg positive and pregnant; and
- 2 HBV DNA > 20 million IU/mL and ALT normal.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation — Confirmed HIV

Both:

- 1 Confirmed HIV infection; and
- 2 Any of the following:
 - 2.1 Symptomatic patient; or
 - 2.2 Patient aged 12 months and under; or
 - 2.3 Both:
 - 2.3.1 Patient aged 1 to 5 years; and
 - 2.3.2 Any of the following:
 - 2.3.2.1 CD4 counts < 1000 cells/mm³; or
 - 2.3.2.2 CD4 counts < 0.25 × total lymphocyte count; or
 - 2.3.2.3 Viral load counts > 100000 copies per ml; or
 - 2.4 Both:
 - 2.4.1 Patient aged 6 years and over; and
 - 2.4.2 CD4 counts < 500 cells/mm³.

Initiation — Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Initiation — Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initiation — Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

Hepatitis C

BOCEPREVIR – **Restricted** see terms below

⚡ Cap 200 mg	5,015.00	336	Victrelis
--------------------	----------	-----	-----------

➡Restricted

Initiation — Chronic hepatitis C - genotype 1, first-line

Gastroenterologist, infectious disease specialist or general physician

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has not received prior pegylated interferon treatment; and
- 3 Patient has IL-28B genotype CT or TT; and
- 4 Patient is to be treated in combination with pegylated interferon and ribavirin; and
- 5 Patient is hepatitis C protease inhibitor treatment-naïve; and
- 6 Maximum of 44 weeks therapy.

Initiation — Chronic hepatitis C - genotype 1, second-line

Gastroenterologist, infectious disease specialist or general physician

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has received pegylated interferon treatment; and

continued...

Price (ex man. excl. GST)	Brand or Generic Manufacturer
\$	Per

continued...

- 3 Any of the following:
 - 3.1 Patient was a responder relapser; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received pegylated interferon prior to 2004; and
- 4 Patient is to be treated in combination with pegylated interferon and ribavirin; and
- 5 Maximum of 44 weeks therapy.

Note: Due to risk of severe sepsis boceprevir should not be initiated if either Platelet count $<100 \times 10^9/l$ or Albumin $<5 \text{ g/l}$.

Herpesviridae

ACICLOVIR

Tab dispersible 200 mg – 1% DV Sep-13 to 2016	1.78	25	Lovir
Tab dispersible 400 mg – 1% DV Sep-13 to 2016	5.98	56	Lovir
Tab dispersible 800 mg – 1% DV Sep-13 to 2016	6.64	35	Lovir
Inj 250 mg vial – 1% DV Jan-16 to 2018	10.10	5	Aciclovir-Clarix

CIDOFOVIR – **Restricted** see terms below

⚡ Inj 75 mg per ml, 5 ml vial

➡ **Restricted**

Clinical microbiologist, infectious disease specialist, otolaryngologist or oral surgeon

FOSCARNET SODIUM – **Restricted** see terms below

⚡ Inj 24 mg per ml, 250 ml bottle

➡ **Restricted**

Clinical microbiologist or infectious disease specialist

GANCICLOVIR – **Restricted** see terms below

⚡ Inj 500 mg vial 380.00 5 Cymevene

➡ **Restricted**

Clinical microbiologist or infectious disease specialist

VALACICLOVIR

Tab 500 mg – 1% DV Mar-16 to 2018	6.42	30	Vaclovir
Tab 1,000 mg – 1% DV Mar-16 to 2018	12.75	30	Vaclovir

VALGANCICLOVIR – **Restricted** see terms on the next page

⚡ Tab 450 mg – 1% DV Jun-15 to 2018 1,050.00 60 Valcyte

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➡Restricted

Initiation — Transplant cytomegalovirus prophylaxis

Limited to 3 months treatment

Patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Initiation — Lung transplant cytomegalovirus prophylaxis

Limited to 6 months treatment

Both:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive.

Initiation — Cytomegalovirus in immunocompromised patients

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Influenza

OSELTAMIVIR – **Restricted** see terms below

☞ Tab 75 mg

☞ Powder for oral suspension 6 mg per ml

➡Restricted

Initiation

Either:

- 1 Only for hospitalised patient with known or suspected influenza; or
- 2 For prophylaxis of influenza in hospitalised patients as part of a DHB hospital approved infections control plan.

ZANAMIVIR

☞ Powder for inhalation 5 mg37.38 20 dose Relenza Rotadisk

➡Restricted

Initiation

Either:

- 1 Only for hospitalised patient with known or suspected influenza; or
- 2 For prophylaxis of influenza in hospitalised patients as part of a DHB hospital approved infections control plan.

Immune Modulators

INTERFERON ALFA-2A

Inj 3 m iu prefilled syringe

Inj 6 m iu prefilled syringe

Inj 9 m iu prefilled syringe

INTERFERON ALFA-2B

Inj 18 m iu, 1.2 ml multidose pen

Inj 30 m iu, 1.2 ml multidose pen

Inj 60 m iu, 1.2 ml multidose pen

INTERFERON GAMMA – **Restricted** see terms below

☞ Inj 100 mcg in 0.5 ml vial

➡Restricted

Initiation

Patient has chronic granulomatous disease and requires interferon gamma.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PEGYLATED INTERFERON ALFA-2A – Restricted see terms below			
¶ Inj 135 mcg prefilled syringe			
¶ Inj 135 mcg prefilled syringe (4) with ribavirin tab 200 mg (112)			
¶ Inj 135 mcg prefilled syringe (4) with ribavirin tab 200 mg (168)			
¶ Inj 180 mcg prefilled syringe	900.00	4	Pegasys
¶ Inj 180 mcg prefilled syringe (4) with ribavirin tab 200 mg (112)	1,159.84	1	Pegasys RBV Combination Pack
¶ Inj 180 mcg prefilled syringe (4) with ribavirin tab 200 mg (168)	1,290.00	1	Pegasys RBV Combination Pack

➡ Restricted

Initiation — Chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant

Limited to 48 weeks treatment

Any of the following:

- 1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
- 2 Patient has chronic hepatitis C and is co-infected with HIV; or
- 3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant.

Notes: Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.

Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml.

Continuation — Chronic hepatitis C - genotype 1 infection

Gastroenterologist, infectious disease specialist or general physician

Re-assessment required after 48 weeks

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir.

Initiation — Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior

Gastroenterologist, infectious disease specialist or general physician

Limited to 48 weeks treatment

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir.

Initiation — Chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV

Limited to 6 months treatment

Patient has chronic hepatitis C, genotype 2 or 3 infection.

Initiation — Hepatitis B

Gastroenterologist, infectious disease specialist or general physician

Limited to 48 weeks treatment

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naïve; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log₁₀ IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 Serum HBV DNA ≥ 2,000 units/ml and significant fibrosis (≥ Metavir Stage F2 or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon.

Notes: Approved dose is 180 mcg once weekly.

The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.

In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon alfa-2a dose should be reduced to 135 mcg once weekly.

In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.

Pegylated Interferon alfa-2a is not approved for use in children.

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Anticholinesterases

EDROPHONIUM CHLORIDE – **Restricted** see terms below

- ⚡ Inj 10 mg per ml, 15 ml vial
- ⚡ Inj 10 mg per ml, 1 ml ampoule

➔ **Restricted**

Initiation

For the diagnosis of myasthenia gravis.

NEOSTIGMINE METILSULFATE

Inj 2.5 mg per ml, 1 ml ampoule – **1% DV Sep-14 to 2017** 98.00 50 **AstraZeneca**

NEOSTIGMINE METILSULFATE WITH GLYCOPYRRONIUM BROMIDE

Inj 2.5 mg with glycopyrronium bromide 0.5 mg per ml, 1 ml ampoule
– **1% DV Nov-13 to 2016** 20.90 10 **Max Health**
Max Health brand - HSS with 1% DV will apply 1 July 2016 to 30 June 2019.

PYRIDOSTIGMINE BROMIDE

Tab 60 mg 38.90 100 Mestinon

Antirheumatoid Agents

AURANOFIN

Tab 3 mg

HYDROXYCHLOROQUINE

Tab 200 mg – **1% DV Sep-15 to 2018** 10.50 100 **Plaquenil**

LEFLUNOMIDE

Tab 10 mg 55.00 30 Arava
Tab 20 mg 76.00 30 Arava

PENICILLAMINE

Tab 125 mg 61.93 100 D-Penamine
Tab 250 mg 98.98 100 D-Penamine

SODIUM AUROTHIOMALATE

Inj 10 mg in 0.5 ml ampoule
Inj 20 mg in 0.5 ml ampoule
Inj 50 mg in 0.5 ml ampoule

Drugs Affecting Bone Metabolism

Bisphosphonates

ALENDRONATE SODIUM

⚡ Tab 40 mg 133.00 30 Fosamax

➔ **Restricted**

Initiation — Paget's disease

Both:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications due to site (base of skull, spine, long bones of lower limbs); or
 - 2.5 Preparation for orthopaedic surgery.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
☞ Tab 70 mg	12.90	4	Fosamax

☞ Restricted

Initiation — Osteoporosis

Any of the following:

- History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) ≥ 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score ≤ -2.5) (see Note); or
- History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- History of two significant osteoporotic fractures demonstrated radiologically; or
- Documented T-Score ≤ -3.0 (see Note); or
- A 10-year risk of hip fracture $\geq 3\%$, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- Patient has had a Special Authority approval for zoledronic acid (underlying cause – osteoporosis) or raloxifene.

Initiation — glucocorticosteroid therapy

Re-assessment required after 12 months

Both:

- The patient is receiving systemic glucocorticosteroid therapy (≥ 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- Any of the following:
 - The patient has documented BMD ≥ 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score ≤ -1.5) (see Note); or
 - The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - The patient has had a Special Authority approval for zoledronic acid (glucocorticosteroid therapy) or raloxifene.

Continuation — glucocorticosteroid therapy

Re-assessment required after 12 months

The patient is continuing systemic glucocorticosteroid therapy (≥ 5 mg per day prednisone equivalents).

Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score ≤ -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

ALENDRONATE SODIUM WITH CHOLECALCIFEROL – **Restricted** see terms on the next page

☞ Tab 70 mg with cholecalciferol 5,600 iu	12.90	4	Fosamax Plus
---	-------	---	--------------

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

➔ **Restricted**

Initiation — Osteoporosis

Any of the following:

- History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) ≥ 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score ≤ -2.5) (see Note); or
- History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- History of two significant osteoporotic fractures demonstrated radiologically; or
- Documented T-Score ≤ -3.0 (see Note); or
- A 10-year risk of hip fracture $\geq 3\%$, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- Patient has had a Special Authority approval for zoledronic acid (underlying cause – osteoporosis) or raloxifene.

Initiation — glucocorticosteroid therapy

Re-assessment required after 12 months

Both:

- The patient is receiving systemic glucocorticosteroid therapy (≥ 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- Any of the following:
 - The patient has documented BMD ≥ 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score ≤ -1.5) (see Note); or
 - The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - The patient has had a Special Authority approval for zoledronic acid (glucocorticosteroid therapy) or raloxifene.

Continuation — glucocorticosteroid therapy

Re-assessment required after 12 months

The patient is continuing systemic glucocorticosteroid therapy (≥ 5 mg per day prednisone equivalents).

Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score ≥ -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

ETIDRONATE DISODIUM

Tab 200 mg – 1% DV Sep-15 to 2018	13.50	100	Arrow-Etidronate
---	-------	-----	-------------------------

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	6.80	1	Pamisol
Inj 6 mg per ml, 10 ml vial	13.20	1	Pamisol
Inj 9 mg per ml, 10 ml vial	19.20	1	Pamisol

RISEDRONATE SODIUM

Tab 35 mg	4.00	4	Risedronate Sandoz
-----------------	------	---	--------------------

ZOLEDRONIC ACID

⚡ Inj 5 mg per 100 ml, vial	600.00	100 ml	Aclasta
-----------------------------------	--------	--------	---------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Initiation — Inherited bone fragility disorders			
Any specialist			
Patient has been diagnosed with an inherited bone fragility disorder (e.g. osteogenesis imperfecta).			
Initiation — Osteoporosis			
Any specialist			
<i>Therapy limited to 3 doses</i>			
Both:			
1 Any of the following:			
1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) \geq 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score \leq -2.5) (see Note); or			
1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or			
1.3 History of two significant osteoporotic fractures demonstrated radiologically; or			
1.4 Documented T-Score \geq -3.0 (see Note); or			
1.5 A 10-year risk of hip fracture \geq 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or			
1.6 Patient has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) or raloxifene; and			
2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.			
Initiation — glucocorticosteroid therapy			
Any specialist			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 The patient is receiving systemic glucocorticosteroid therapy (\geq 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and			
2 Any of the following:			
2.1 The patient has documented BMD \geq 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score \leq -1.5) (see Note); or			
2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or			
2.3 The patient has had a Special Authority approval for alendronate (Underlying cause - glucocorticosteroid therapy) or raloxifene; and; and			
3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.			
Continuation — glucocorticosteroid therapy			
Any specialist			
<i>Re-assessment required after 12 months</i>			
Both:			
1 The patient is continuing systemic glucocorticosteroid therapy (\geq 5 mg per day prednisone equivalents); and			
2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.			
Initiation — Paget's disease			
Any specialist			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 Paget's disease; and			
2 Any of the following:			
2.1 Bone or articular pain; or			
2.2 Bone deformity; or			
2.3 Bone, articular or neurological complications; or			
2.4 Asymptomatic disease, but risk of complications; or			
continued...			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2.5 Preparation for orthopaedic surgery; and

3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Continuation — Paget's disease

Any specialist

Re-assessment required after 12 months

Both:


- Any of the following:
 - The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - Symptomatic disease (prescriber determined); and
- The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score ≤ -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Other Drugs Affecting Bone Metabolism

RALOXIFENE – **Restricted** see terms below

<div></div> Tab 60 mg	53.76	28	Evista
--	-------	----	--------

➡Restricted

Initiation

Any of the following:

- History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) ≥ 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score ≤ -2.5) (see Notes); or
- History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- History of two significant osteoporotic fractures demonstrated radiologically; or
- Documented T-Score ≥ -3.0 (see Notes); or
- A 10-year risk of hip fracture $\geq 3\%$, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- Patient has had a prior Special Authority approval for zoledronic acid (Underlying cause - Osteoporosis) or alendronate (Underlying cause - Osteoporosis).

Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score ≤ -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

TERIPARATIDE – **Restricted** see terms below

¶ Inj 250 mcg per ml, 2.4 ml cartridge490.00 1 Forteo

➡ **Restricted**

Initiation

Limited to 18 months treatment

All of the following:

- The patient has severe, established osteoporosis; and
- The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- The patient has had two or more fractures due to minimal trauma; and
- The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- Antiresorptive agents and their adequate doses for the purposes of this restriction are defined as: alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Enzymes

HYALURONIDASE

Inj 1,500 iu ampoule

Hyperuricaemia and Antigout

ALLOPURINOL

Tab 100 mg – 1% DV Mar-15 to 2017	15.11	1,000	Apo-Allopurinol
Tab 300 mg – 1% DV Mar-15 to 2017	15.91	500	Apo-Allopurinol

BENZBROMARONE – **Restricted** see terms on the next page

¶ Tab 100 mg45.00 100 Benzbromaron AL 100

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

➔ **Restricted**

Initiation

Any specialist

All of the following:

1 Patient has been diagnosed with gout; and

2 Any of the following:

2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or

2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or

2.3 Both:

2.3.1 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); and

2.3.2 The patient has a rate of creatinine clearance greater than or equal to 20 ml/min; or

2.4 All of the following:

2.4.1 The patient is taking azathioprine and requires urate-lowering therapy; and

2.4.2 Allopurinol is contraindicated; and

2.4.3 Appropriate doses of probenecid are ineffective or probenecid cannot be used due to reduced renal function; and

3 The patient is receiving monthly liver function tests.

Notes: Benzbromarone has been associated with potentially fatal hepatotoxicity. In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

The New Zealand Rheumatology Association has developed information for prescribers which can be accessed from its website at www.rheumatology.org.nz/downloads/Benzbromarone-prescriber-information-NZRA-V2.pdf

COLCHICINE

Tab 500 mcg – 1% DV Oct-13 to 2016	10.08	100	Colgout
--	-------	-----	----------------

FEBUXOSTAT – Restricted see terms below

⚡ Tab 80 mg	39.50	28	Adenuric
⚡ Tab 120 mg	39.50	28	Adenuric

➔ **Restricted**

Initiation

Any specialist

Both:

1 Patient has been diagnosed with gout; and

2 Any of the following:

2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or

2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or

2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note).

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

MUSCULOSKELETAL SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

PROBENECID

Tab 500 mg

RASBURICASE – **Restricted** see terms below

☞ Inj 1.5 mg vial

☞ **Restricted**

Haematologist

Muscle Relaxants and Related Agents

ATracURIUM BESYLATE

Inj 10 mg per ml, 2.5 ml ampoule – 1% DV Jan-16 to 2018	10.00	5	Tracrium
Inj 10 mg per ml, 5 ml ampoule – 1% DV Jan-16 to 2018	12.50	5	Tracrium

BACLOFEN

Tab 10 mg – 1% DV Jun-13 to 2016	3.85	100	Pacifen
Oral liq 1 mg per ml			
Inj 0.05 mg per ml, 1 ml ampoule – 1% DV Sep-15 to 2018	11.55	1	Lioresal Intrathecal
Inj 2 mg per ml, 5 ml ampoule	209.29	1	Lioresal Intrathecal

CLOSTRIDIUM BOTULINUM TYPE A TOXIN

Inj 100 u vial	467.50	1	Botox
Inj 300 u vial	388.50	1	Dysport
Inj 500 u vial	1,295.00	2	Dysport

DANTROLENE

Cap 25 mg	65.00	100	Dantrium
Cap 50 mg	77.00	100	Dantrium
Inj 20 mg vial	800.00	6	Dantrium IV

MIVACURIUM CHLORIDE

Inj 2 mg per ml, 5 ml ampoule	33.92	5	Mivacron
Inj 2 mg per ml, 10 ml ampoule	67.17	5	Mivacron

ORPHENADRINE CITRATE

Tab 100 mg

PANCURONIUM BROMIDE

Inj 2 mg per ml, 2 ml ampoule	260.00	50	AstraZeneca
-------------------------------------	--------	----	--------------------

ROCURONIUM BROMIDE

Inj 10 mg per ml, 5 ml vial	38.25	10	DBL Rocuronium Bromide
-----------------------------------	-------	----	-------------------------------

SUXAMETHONIUM CHLORIDE

Inj 50 mg per ml, 2 ml ampoule – 1% DV Jun-14 to 2017	78.00	50	AstraZeneca
---	-------	----	--------------------

VECURONIUM BROMIDE

Inj 4 mg ampoule
Inj 10 mg vial

Reversers of Neuromuscular Blockade

SUGAMMADEX – **Restricted** see terms on the next page

☞ Inj 100 mg per ml, 2 ml vial	1,200.00	10	Bridion
☞ Inj 100 mg per ml, 5 ml vial	3,000.00	10	Bridion

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

➔ **Restricted**

Initiation

Any of the following:

- 1 Patient requires reversal of profound neuromuscular blockade following rapid sequence induction that has been undertaken using rocuronium (i.e. suxamethonium is contraindicated or undesirable); or
- 2 Severe neuromuscular degenerative disease where the use of neuromuscular blockade is required; or
- 3 Patient has an unexpectedly difficult airway that cannot be intubated and requires a rapid reversal of anaesthesia and neuromuscular blockade; or
- 4 The duration of the patient's surgery is unexpectedly short; or
- 5 Neostigmine or a neostigmine/anticholinergic combination is contraindicated (for example the patient has ischaemic heart disease, morbid obesity or COPD); or
- 6 Patient has a partial residual block after conventional reversal.

Non-Steroidal Anti-Inflammatory Drugs

CELECOXIB – **Restricted** see terms below

- ⚡ Cap 100 mg
- ⚡ Cap 200 mg
- ⚡ Cap 400 mg

➔ **Restricted**

Initiation

For preoperative and/or postoperative use for a total of up to 8 days' use.

DICLOFENAC SODIUM

Tab EC 25 mg – 1% DV Dec-15 to 2018	1.30	50	Diclofenac Sandoz
Tab 50 mg dispersible	1.50	20	Voltaren D
Tab EC 50 mg – 1% DV Dec-15 to 2018	1.00	50	Diclofenac Sandoz
Tab long-acting 75 mg – 1% DV Dec-15 to 2018	15.20	500	Apo-Diclo SR
Tab long-acting 100 mg – 1% DV Dec-15 to 2018	26.20	500	Apo-Diclo SR
Inj 25 mg per ml, 3 ml ampoule – 1% DV Oct-14 to 2017	13.20	5	Voltaren
Suppos 12.5 mg – 1% DV Oct-14 to 2017	2.04	10	Voltaren
Suppos 25 mg – 1% DV Oct-14 to 2017	2.44	10	Voltaren
Suppos 50 mg – 1% DV Oct-14 to 2017	4.22	10	Voltaren
Suppos 100 mg – 1% DV Oct-14 to 2017	7.00	10	Voltaren

ETORICOXIB – **Restricted** see terms below

- ⚡ Tab 30 mg
- ⚡ Tab 60 mg
- ⚡ Tab 90 mg
- ⚡ Tab 120 mg

➔ **Restricted**

Initiation

For preoperative and/or postoperative use for a total of up to 8 days' use.

IBUPROFEN

Tab 200 mg			
➔ Tab 400 mg – Restricted : For continuation only			
➔ Tab 600 mg – Restricted : For continuation only			
Tab long-acting 800 mg – 1% DV Jul-15 to 2018	7.99	30	Brufen SR
Oral liq 20 mg per ml – 1% DV Mar-14 to 2016	1.89	200 ml	Fenpaed
Inj 5 mg per ml, 2 ml ampoule			
Inj 10 mg per ml, 2 ml vial			

MUSCULOSKELETAL SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
INDOMETHACIN			
Cap 25 mg			
Cap 50 mg			
Cap long-acting 75 mg			
Inj 1 mg vial			
Suppos 100 mg			
KETOPROFEN			
Cap long-acting 200 mg	12.07	28	Oruvail SR
MEFENAMIC ACID – Restricted: For continuation only			
➡ Cap 250 mg			
MELOXICAM – Restricted see terms below			
⚡ Tab 7.5 mg			
➡Restricted			
Initiation			
Either:			
1 All of the following:			
1.1 Haemophilic arthropathy; and			
1.2 The patient has moderate to severe haemophilia with less than or equal to 5% of normal circulating functional clotting factor; and			
1.3 Pain and inflammation associated with haemophilic arthropathy is inadequately controlled by alternative funded treatment options, or alternative funded treatment options are contraindicated; or			
2 For preoperative and/or postoperative use for a total of up to 8 days' use.			
NAPROXEN			
Tab 250 mg – 1% DV Sep-15 to 2018	18.06	500	Noflam 250
Tab 500 mg – 1% DV Sep-15 to 2018	18.91	250	Noflam 500
Tab long-acting 750 mg – 1% DV Jun-15 to 2018	18.00	90	Naprosyn SR 750
Tab long-acting 1 g – 1% DV Jun-15 to 2018	21.00	90	Naprosyn SR 1000
PARECOXIB			
Inj 40 mg vial	100.00	10	Dynastat
SULINDAC			
Tab 100 mg			
Tab 200 mg			
TENOXICAM			
Tab 20 mg – 1% DV Jan-15 to 2016	3.05	20	Reutenox
Inj 20 mg vial	9.95	1	AFT

Topical Products for Joint and Muscular Pain

CAPSAICIN – Restricted see terms below

⚡ Crm 0.025% 9.95 45 g Zostrix

➡Restricted

Initiation

Patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Agents for Parkinsonism and Related Disorders

Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE – **Restricted** see terms below

⚡ Tab 50 mg 400.00 56 Rilutek

➡ **Restricted**

Initiation

Neurologist or respiratory specialist

Re-assessment required after 6 months

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
 - 5.1 The patient is ambulatory; or
 - 5.2 The patient is able to use upper limbs; or
 - 5.3 The patient is able to swallow.

Continuation

Re-assessment required after 18 months

All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
 - 3.1 The patient is ambulatory; or
 - 3.2 The patient is able to use upper limbs; or
 - 3.3 The patient is able to swallow.

TETRABENAZINE

Tab 25 mg – **1% DV Sep-13 to 2016** 118.00 112 **Motetis**

Anticholinergics

BENZTROPINE MESYLATE

Tab 2 mg 7.99 60 Benztrop

Inj 1 mg per ml, 2 ml ampoule 95.00 5 Cogentin

PROCYCLIDINE HYDROCHLORIDE

Tab 5 mg

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE

Cap 100 mg – **1% DV Oct-14 to 2017** 38.24 60 **Symmetrel**

APOMORPHINE HYDROCHLORIDE

Inj 10 mg per ml, 1 ml ampoule

Inj 10 mg per ml, 2 ml ampoule 119.00 5 Apomine

BROMOCRIPTINE

Tab 2.5 mg

Cap 5 mg

ENTACAPONE

Tab 200 mg – **1% DV Sep-15 to 2018** 28.00 100 **Entapone**

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LEVODOPA WITH BENSERAZIDE			
Tab dispersible 50 mg with benserazide 12.5 mg	10.00	100	Madopar Rapid
Cap 50 mg with benserazide 12.5 mg	8.00	100	Madopar 62.5
Cap 100 mg with benserazide 25 mg	12.50	100	Madopar 125
Cap long-acting 100 mg with benserazide 25 mg	17.00	100	Madopar HBS
Cap 200 mg with benserazide 50 mg	25.00	100	Madopar 250
LEVODOPA WITH CARBIDOPA			
Tab 100 mg with carbidopa 25 mg	20.00	100	Sinemet <i>e.g. Kinson</i>
Tab long-acting 200 mg with carbidopa 50 mg	47.50	100	Sinemet CR
Tab 250 mg with carbidopa 25 mg	40.00	100	Sinemet <i>e.g. Sindopa</i>
LISURIDE HYDROGEN MALEATE			
Tab 200 mcg	25.00	30	Dopergin
PRAMIPEXOLE HYDROCHLORIDE			
Tab 0.25 mg – 1% DV Oct-14 to 2016	7.20	100	Ramipex
Tab 1 mg – 1% DV Oct-14 to 2016	24.39	100	Ramipex
ROPINIROLE HYDROCHLORIDE			
Tab 0.25 mg – 1% DV Mar-14 to 2016	2.36	100	Apo-Ropinirole
Tab 1 mg – 1% DV Mar-14 to 2016	5.32	100	Apo-Ropinirole
Tab 2 mg – 1% DV Mar-14 to 2016	7.72	100	Apo-Ropinirole
Tab 5 mg – 1% DV Mar-14 to 2016	14.48	100	Apo-Ropinirole
SELEGILINE HYDROCHLORIDE			
Tab 5 mg			
TOLCAPONE			
Tab 100 mg	126.20	100	Tasmar
Anaesthetics			
General Anaesthetics			
DESFLURANE			
Soln for inhalation 100%, 240 ml bottle	1,414.50	6	Suprane
DEXMEDETOMIDINE			
Inj 100 mcg per ml, 2 ml vial – 1% DV Oct-14 to 2017	479.85	5	Precedex
ETOMIDATE			
Inj 2 mg per ml, 10 ml ampoule			
ISOFLURANE			
Soln for inhalation 100%, 250 ml bottle	1,173.00	6	Aerrane
KETAMINE			
Inj 1 mg per ml, 100 ml bag – 1% DV Sep-14 to 2017	27.00	1	Biomed
Inj 4 mg per ml, 50 ml syringe – 1% DV Sep-14 to 2017	25.00	1	Biomed
Inj 10 mg per ml, 10 ml syringe – 1% DV Sep-14 to 2017	14.00	1	Biomed
Inj 100 mg per ml, 2 ml ampoule – 1% DV May-16 to 2018	47.05	5	Ketamine-Claris
METHOHEXITAL SODIUM			
Inj 10 mg per ml, 50 ml vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PROPOFOL			
Inj 10 mg per ml, 20 ml ampoule	7.60	5	Fresofol 1%
Inj 10 mg per ml, 20 ml vial – 10% DV Jun-16 to 2019	42.00	5	Diprivan
	5.27		Provide MCT-LCT 1%
Inj 10 mg per ml, 50 ml syringe	47.00	1	Diprivan
Inj 10 mg per ml, 50 ml vial – 10% DV Jun-16 to 2019	25.00	1	Diprivan
	4.00		Fresofol 1%
	24.50	10	Fresofol 1% MCT/LCT
	4.00	1	Provide MCT-LCT 1%
Inj 10 mg per ml, 100 ml vial – 10% DV Jun-16 to 2019	7.60	1	Fresofol 1%
	49.00	10	Fresofol 1% MCT/LCT
	7.60	1	Provide MCT-LCT 1%
<i>(Fresofol 1% Inj 10 mg per ml, 20 ml ampoule to be delisted 1 June 2016)</i>			
<i>(Diprivan Inj 10 mg per ml, 20 ml vial to be delisted 1 June 2016)</i>			
<i>(Diprivan Inj 10 mg per ml, 50 ml syringe to be delisted 1 June 2016)</i>			
<i>(Diprivan Inj 10 mg per ml, 50 ml vial to be delisted 1 June 2016)</i>			
<i>(Fresofol 1% Inj 10 mg per ml, 50 ml vial to be delisted 1 June 2016)</i>			
<i>(Provide MCT-LCT 1% Inj 10 mg per ml, 50 ml vial to be delisted 1 June 2016)</i>			
<i>(Fresofol 1% Inj 10 mg per ml, 100 ml vial to be delisted 1 June 2016)</i>			
<i>(Provide MCT-LCT 1% Inj 10 mg per ml, 100 ml vial to be delisted 1 June 2016)</i>			
SEVOFLURANE			
Soln for inhalation 100%, 250 ml bottle	1,365.00	6	Baxter
THIOPENTAL [THIOPENTONE] SODIUM			
Inj 500 mg ampoule			
Local Anaesthetics			
ARTICAINE HYDROCHLORIDE			
Inj 1%			
ARTICAINE HYDROCHLORIDE WITH ADRENALINE			
Inj 4% with adrenaline 1:100,000, 1.7 ml dental cartridge			
Inj 4% with adrenaline 1:100,000, 2.2 ml dental cartridge			
Inj 4% with adrenaline 1:200,000, 1.7 ml dental cartridge			
Inj 4% with adrenaline 1:200,000, 2.2 ml dental cartridge			
BENZOCAINE			
Gel 20%			
BUPIVACAINE HYDROCHLORIDE			
Inj 5 mg per ml, 4 ml ampoule – 1% DV Jul-14 to 2017	50.00	5	Marcaïn Isobaric
Inj 2.5 mg per ml, 20 ml ampoule			
Inj 2.5 mg per ml, 20 ml ampoule sterile pack – 1% DV Sep-15 to 2018	29.20	5	Marcaïn
Inj 5 mg per ml, 10 ml ampoule sterile pack – 1% DV Sep-15 to 2018	20.25	5	Marcaïn
Inj 5 mg per ml, 20 ml ampoule			
Inj 5 mg per ml, 20 ml ampoule sterile pack – 1% DV Sep-15 to 2018	20.70	5	Marcaïn
Inj 1.25 mg per ml, 100 ml bag			
Inj 1.25 mg per ml, 200 ml bag			
Inj 2.5 mg per ml, 100 ml bag – 1% DV Jul-14 to 2017	150.00	5	Marcaïn
Inj 2.5 mg per ml, 200 ml bag			
Inj 1.25 mg per ml, 500 ml bag			

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BUPIVACAINE HYDROCHLORIDE WITH ADRENALINE			
Inj 2.5 mg per ml with adrenaline 1:400,000, 20 ml vial – 1% DV Sep-14 to 2017	135.00	5	Marcaïn with Adrenaline
Inj 5 mg per ml with adrenaline 1:200,000, 20 ml vial – 1% DV Sep-14 to 2017	115.00	5	Marcaïn with Adrenaline
BUPIVACAINE HYDROCHLORIDE WITH FENTANYL			
Inj 0.625 mg with fentanyl 2 mcg per ml, 100 ml bag			
Inj 0.625 mg with fentanyl 2 mcg per ml, 200 ml bag			
Inj 1.25 mg with fentanyl 2 mcg per ml, 100 ml syringe			
Inj 1.25 mg with fentanyl 2 mcg per ml, 100 ml bag	210.00	10	Bupafen
Inj 1.25 mg with fentanyl 2 mcg per ml, 200 ml bag	210.00	10	Bupafen
Inj 1.25 mg with fentanyl 2 mcg per ml, 50 ml syringe			
Inj 1.25 mg with fentanyl 2 mcg per ml, 15 ml syringe	72.00	10	Biomed
Inj 1.25 mg with fentanyl 2 mcg per ml, 20 ml syringe	92.00	10	Biomed
BUPIVACAINE HYDROCHLORIDE WITH GLUCOSE			
Inj 0.5% with glucose 8%, 4 ml ampoule	38.00	5	Marcaïn Heavy
COCAINE HYDROCHLORIDE			
Paste 5%			
Soln 15%, 2 ml syringe			
Soln 4%, 2 ml syringe	25.46	1	Biomed
COCAINE HYDROCHLORIDE WITH ADRENALINE			
Paste 15% with adrenaline 0.06%			
Paste 25% with adrenaline 0.06%			
ETHYL CHLORIDE			
Spray 100%			
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE			
Gel 2% – 1% DV Sep-15 to 2018	3.40	20 ml	Orion
Soln 4%			
Spray 10% – 1% DV Sep-13 to 2016	75.00	50 ml	Xylocaine
Oral (viscous) soln 2% – 1% DV Sep-14 to 2017	55.00	200 ml	Xylocaine Viscous
Inj 1%, 20 ml ampoule, sterile pack			
Inj 2%, 20 ml ampoule, sterile pack			
Inj 1%, 5 ml ampoule	8.75	25	Lidocaine-Claris
Inj 1%, 20 ml ampoule	2.40	1	Lidocaine-Claris
Inj 2%, 5 ml ampoule	6.90	25	Lidocaine-Claris
Inj 2%, 20 ml ampoule	2.40	1	Lidocaine-Claris
Gel 2%, 10 ml urethral syringe	43.26	10	Pfizer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH ADRENALINE			
Inj 1% with adrenaline 1:100,000, 5 ml ampoule	27.00	10	Xylocaine
Inj 1% with adrenaline 1:200,000, 20 ml vial	50.00	5	Xylocaine
Inj 2% with adrenaline 1:80,000, 1.7 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 1.8 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 2.2 ml dental cartridge			
Inj 2% with adrenaline 1:200,000, 20 ml vial	60.00	5	Xylocaine

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH ADRENALINE AND TETRACAINE HYDROCHLORIDE			
Soln 4% with adrenaline 0.1% and tetracaine hydrochloride 0.5%, 5 ml syringe – 1% DV Oct-14 to 2017	17.50	1	Topicaine
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH CHLORHEXIDINE			
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringe	43.26	10	Pfizer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH PHENYLEPHRINE HYDROCHLORIDE			
Nasal spray 5% with phenylephrine hydrochloride 0.5%			
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE			
Crm 2.5% with prilocaine 2.5%	45.00	30 g	EMLA
Patch 25 mcg with prilocaine 25 mcg	115.00	20	EMLA
Crm 2.5% with prilocaine 2.5%, 5 g	45.00	5	EMLA
LIDOCAINE [LIGNOCAINE]			
Crm 4%	27.00	30 g	LMX4
Crm 4% (5 g tubes)	27.00	5	LMX4
MEPIVACAINE HYDROCHLORIDE			
Inj 3%, 1.8 ml dental cartridge – 1% DV Oct-14 to 2017	43.60	50	Scandonest 3%
Inj 3%, 2.2 ml dental cartridge – 1% DV Oct-14 to 2017	43.60	50	Scandonest 3%
PRILOCAINE HYDROCHLORIDE			
Inj 0.5%, 50 ml vial	100.00	5	Citanest
Inj 2%, 5 ml ampoule	55.00	10	Citanest
PRILOCAINE HYDROCHLORIDE WITH FELYPRESSIN			
Inj 3% with felypressin 0.03 iu per ml, 1.8 ml dental cartridge			
Inj 3% with felypressin 0.03 iu per ml, 2.2 ml dental cartridge			
ROPIVACAINE HYDROCHLORIDE			
Inj 2 mg per ml, 10 ml ampoule – 1% DV Aug-15 to 2017	9.05	5	Ropivacaine Kabi
Inj 2 mg per ml, 20 ml ampoule – 1% DV Aug-15 to 2017	9.50	5	Ropivacaine Kabi
Inj 2 mg per ml, 100 ml bag – 1% DV Jul-15 to 2017	60.00	5	Naropin
Inj 2 mg per ml, 200 ml bag – 1% DV Jul-15 to 2017	79.50	5	Naropin
Inj 7.5 mg per ml, 10 ml ampoule – 1% DV Aug-15 to 2017	10.20	5	Ropivacaine Kabi
Inj 7.5 mg per ml, 20 ml ampoule – 1% DV Aug-15 to 2017	12.50	5	Ropivacaine Kabi
Inj 10 mg per ml, 10 ml ampoule – 1% DV Aug-15 to 2017	10.90	5	Ropivacaine Kabi
Inj 10 mg per ml, 20 ml ampoule – 1% DV Aug-15 to 2017	16.30	5	Ropivacaine Kabi
ROPIVACAINE HYDROCHLORIDE WITH FENTANYL			
Inj 2 mg with fentanyl 2 mcg per ml, 100 ml bag	198.50	5	Naropin
Inj 2 mg with fentanyl 2 mcg per ml, 200 ml bag	270.00	5	Naropin
TETRACAINE [AMETHOCAINE] HYDROCHLORIDE			
Gel 4%			

Analgesics

Non-Opioid Analgesics

ASPIRIN

Tab dispersible 300 mg

CAPSAICIN – **Restricted** see terms on the next page

⚡ Crm 0.075% 12.50 45 g Zostrix HP

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➡Restricted			
Initiation			
For post-herpetic neuralgia or diabetic peripheral neuropathy.			
METHOXYFLURANE – Restricted see terms below			
☞ Soln for inhalation 99.9%, 3 ml bottle			
➡Restricted			
Initiation			
Both:			
1 Patient is undergoing a painful procedure with an expected duration of less than one hour; and			
2 Only to be used under supervision by a medical practitioner or nurse who is trained in the use of methoxyflurane.			
NEFOPAM HYDROCHLORIDE			
Tab 30 mg			
PARACETAMOL – Some items restricted see terms below			
Tab soluble 500 mg – 1% DV Oct-15 to 2017	1.60	20	Paragesic Soluble
Tab 500 mg			
Oral liq 120 mg per 5 ml – 20% DV Oct-14 to 2017	4.15	1,000 ml	Paracare
Oral liq 250 mg per 5 ml – 20% DV Sep-14 to 2017	4.35	1,000 ml	Paracare Double Strength
☞ Inj 10 mg per ml, 50 ml vial – 1% DV Sep-14 to 2017	12.90	12	Perfalgan
☞ Inj 10 mg per ml, 100 ml vial – 1% DV Sep-14 to 2017	12.90	12	Perfalgan
Suppos 25 mg	56.35	20	Biomed
Suppos 50 mg	56.35	20	Biomed
Suppos 125 mg – 1% DV Dec-15 to 2018	3.69	10	Gacet
Suppos 250 mg – 1% DV Dec-15 to 2018	3.79	10	Gacet
Suppos 500 mg – 1% DV Nov-15 to 2018	12.60	50	Paracare
➡Restricted			
Initiation			
Intravenous paracetamol is only to be used where other routes are unavailable or impractical, or where there is reduced absorption.			
The need for IV paracetamol must be re-assessed every 24 hours.			
SUCROSE			
Oral liq 25%			
Opioid Analgesics			
ALFENTANIL			
Inj 0.5 mg per ml, 2 ml ampoule – 1% DV Jan-15 to 2017	39.07	10	Hameln
CODEINE PHOSPHATE			
Tab 15 mg – 1% DV Jul-13 to 2016	4.75	100	PSM
Tab 30 mg – 1% DV Jul-13 to 2016	5.80	100	PSM
Tab 60 mg – 1% DV Jul-13 to 2016	12.50	100	PSM
DIHYDROCODEINE TARTRATE			
Tab long-acting 60 mg – 1% DV Sep-13 to 2016	13.64	60	DHC Continus

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FENTANYL			
Inj 10 mcg per ml, 10 ml syringe			
Inj 50 mcg per ml, 2 ml ampoule – 1% DV Sep-15 to 2018	3.95	10	Boucher and Muir
Inj 10 mcg per ml, 50 ml bag	210.00	10	Biomed
Inj 10 mcg per ml, 50 ml syringe	165.00	10	Biomed
Inj 50 mcg per ml, 10 ml ampoule – 1% DV Sep-15 to 2018	10.45	10	Boucher and Muir
Inj 10 mcg per ml, 100 ml bag	210.00	10	Biomed
Inj 20 mcg per ml, 50 ml syringe	185.00	10	Biomed
Inj 20 mcg per ml, 100 ml bag			
Patch 12.5 mcg per hour – 1% DV Aug-15 to 2016	2.92	5	Fentanyl Sandoz
Patch 25 mcg per hour – 1% DV Aug-15 to 2016	3.66	5	Fentanyl Sandoz
Patch 50 mcg per hour – 1% DV Aug-15 to 2016	6.64	5	Fentanyl Sandoz
Patch 75 mcg per hour – 1% DV Aug-15 to 2016	9.18	5	Fentanyl Sandoz
Patch 100 mcg per hour – 1% DV Aug-15 to 2016	11.29	5	Fentanyl Sandoz
METHADONE HYDROCHLORIDE			
Tab 5 mg – 1% DV Sep-15 to 2018	1.85	10	Methatabs
Oral liq 2 mg per ml – 1% DV Sep-15 to 2018	5.55	200 ml	Biodone
Oral liq 5 mg per ml – 1% DV Sep-15 to 2018	5.00	200 ml	Biodone Forte
Oral liq 10 mg per ml – 1% DV Sep-15 to 2018	6.55	200 ml	Biodone Extra Forte
Inj 10 mg per ml, 1 ml vial	61.00	10	AFT
MORPHINE HYDROCHLORIDE			
Oral liq 1 mg per ml – 1% DV Oct-15 to 2018	8.84	200 ml	RA-Morph
Oral liq 2 mg per ml – 1% DV Oct-15 to 2018	14.00	200 ml	RA-Morph
Oral liq 5 mg per ml – 1% DV Oct-15 to 2018	18.00	200 ml	RA-Morph
Oral liq 10 mg per ml – 1% DV Oct-15 to 2018	26.00	200 ml	RA-Morph

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MORPHINE SULPHATE			
Tab long-acting 10 mg – 1% DV Sep-13 to 2016	1.95	10	Arrow-Morphine LA
Tab immediate-release 10 mg – 1% DV Apr-15 to 2017	2.80	10	Sevredol
Tab immediate-release 20 mg – 1% DV Apr-15 to 2017	5.52	10	Sevredol
Tab long-acting 30 mg – 1% DV Sep-13 to 2016	2.98	10	Arrow-Morphine LA
Tab long-acting 60 mg – 1% DV Sep-13 to 2016	5.75	10	Arrow-Morphine LA
Tab long-acting 100 mg – 1% DV Sep-13 to 2016	6.45	10	Arrow-Morphine LA
Cap long-acting 10 mg – 1% DV Feb-14 to 2016	1.70	10	m-Eslon
Cap long-acting 30 mg – 1% DV Feb-14 to 2016	2.50	10	m-Eslon
Cap long-acting 60 mg – 1% DV Feb-14 to 2016	5.40	10	m-Eslon
Cap long-acting 100 mg – 1% DV Feb-14 to 2016	6.38	10	m-Eslon
Inj 1 mg per ml, 100 ml bag – 1% DV Oct-14 to 2017	185.00	10	Biomed
Inj 1 mg per ml, 10 ml syringe – 1% DV Oct-14 to 2017	45.00	10	Biomed
Inj 1 mg per ml, 50 ml syringe – 1% DV Oct-14 to 2017	87.50	10	Biomed
Inj 1 mg per ml, 2 ml syringe			
Inj 2 mg per ml, 30 ml syringe	135.00	10	Biomed
Inj 5 mg per ml, 1 ml ampoule – 1% DV Oct-14 to 2017	12.48	5	DBL Morphine Sulphate
Inj 10 mg per ml, 1 ml ampoule – 1% DV Oct-14 to 2017	9.09	5	DBL Morphine Sulphate
Inj 10 mg per ml, 100 mg cassette			
Inj 10 mg per ml, 100 ml bag			
Inj 15 mg per ml, 1 ml ampoule – 1% DV Oct-14 to 2017	9.77	5	DBL Morphine Sulphate
Inj 30 mg per ml, 1 ml ampoule – 1% DV Oct-14 to 2017	12.43	5	DBL Morphine Sulphate
Inj 200 mcg in 0.4 ml syringe			
Inj 300 mcg in 0.3 ml syringe			
MORPHINE TARTRATE			
Inj 80 mg per ml, 1.5 ml ampoule – 1% DV Sep-13 to 2016	35.60	5	Hospira
Inj 80 mg per ml, 5 ml ampoule – 1% DV Sep-13 to 2016	107.67	5	Hospira

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
OXYCODONE HYDROCHLORIDE			
Tab controlled-release 5 mg	7.51	20	OxyContin
Tab controlled-release 10 mg	6.75	20	Oxycodone ControlledRelease Tablets(BNM)
Tab controlled-release 20 mg	11.50	20	Oxycodone ControlledRelease Tablets(BNM)
Tab controlled-release 40 mg	18.50	20	Oxycodone ControlledRelease Tablets(BNM)
Tab controlled-release 80 mg	34.00	20	Oxycodone ControlledRelease Tablets(BNM)
Cap immediate-release 5 mg – 1% DV Oct-15 to 2018	1.98	20	OxyNorm
Cap immediate-release 10 mg – 1% DV Oct-15 to 2018	3.91	20	OxyNorm
Cap immediate-release 20 mg – 1% DV Oct-15 to 2018	6.84	20	OxyNorm
Oral liq 5 mg per 5 ml	11.20	250 ml	OxyNorm
Inj 1 mg per ml, 100 ml bag			
Inj 10 mg per ml, 1 ml ampoule – 1% DV Feb-16 to 2018	8.57	5	OxyNorm
Inj 10 mg per ml, 2 ml ampoule – 1% DV Feb-16 to 2018	16.89	5	OxyNorm
Inj 50 mg per ml, 1 ml ampoule – 1% DV Dec-15 to 2018	51.00	5	OxyNorm
PARACETAMOL WITH CODEINE			
Tab paracetamol 500 mg with codeine phosphate 8 mg	2.11	100	Paracetamol + Codeine (Relieve)
PETHIDINE HYDROCHLORIDE			
Tab 50 mg – 1% DV Nov-15 to 2018	4.46	10	PSM
Tab 100 mg – 1% DV Nov-15 to 2018	6.25	10	PSM
Inj 5 mg per ml, 10 ml syringe			
Inj 5 mg per ml, 100 ml bag			
Inj 10 mg per ml, 100 ml bag			
Inj 10 mg per ml, 50 ml syringe			
Inj 50 mg per ml, 1 ml ampoule – 1% DV Sep-14 to 2017	5.51	5	DBL Pethidine Hydrochloride
Inj 50 mg per ml, 2 ml ampoule – 1% DV Sep-14 to 2017	5.83	5	DBL Pethidine Hydrochloride
REMIFENTANIL HYDROCHLORIDE			
Inj 1 mg vial – 1% DV Nov-14 to 2017	10.00	5	Ultiva
Inj 2 mg vial – 1% DV Nov-14 to 2017	18.00	5	Ultiva
TRAMADOL HYDROCHLORIDE			
Tab sustained-release 100 mg – 1% DV Oct-14 to 2017	2.00	20	Tramal SR 100
Tab sustained-release 150 mg – 1% DV Oct-14 to 2017	3.00	20	Tramal SR 150
Tab sustained-release 200 mg – 1% DV Oct-14 to 2017	4.00	20	Tramal SR 200
Cap 50 mg – 1% DV Oct-14 to 2017	2.50	100	Arrow-Tramadol
Oral drops 100 mg per ml			
Inj 10 mg per ml, 100 ml bag			
Inj 50 mg per ml, 1 ml ampoule – 1% DV Oct-14 to 2017	4.50	5	Tramal 50
Inj 50 mg per ml, 2 ml ampoule – 1% DV Oct-14 to 2017	4.50	5	Tramal 100

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Antidepressants			
Cyclic and Related Agents			
AMITRIPTYLINE			
Tab 10 mg – 1% DV Sep-14 to 2017	1.68	100	Arrow-Amitriptyline
Tab 25 mg – 1% DV Jan-15 to 2017	1.68	100	Arrow-Amitriptyline
Tab 50 mg – 1% DV Jan-15 to 2017	2.82	100	Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE			
Tab 10 mg – 1% DV Sep-15 to 2018	12.60	100	Apo-Clomipramine
Tab 25 mg – 1% DV Sep-15 to 2018	8.68	100	Apo-Clomipramine
DOTIEPIN HYDROCHLORIDE			
Tab 75 mg	10.50	100	Dopress
Cap 25 mg	6.17	100	Dopress
DOXEPIH HYDROCHLORIDE			
Cap 10 mg			
Cap 25 mg			
Cap 50 mg			
IMIPRAMINE HYDROCHLORIDE			
Tab 10 mg	5.48	50	Tofranil
	6.58	60	Tofranil
Tab 25 mg	8.80	50	Tofranil
MAPROTILINE HYDROCHLORIDE			
Tab 25 mg			
Tab 75 mg			
MIANSERIN HYDROCHLORIDE – Restricted: For continuation only			
➡ Tab 30 mg			
NORTRIPTYLINE HYDROCHLORIDE			
Tab 10 mg – 1% DV Jun-13 to 2016	4.00	100	Norpress
Tab 25 mg – 1% DV Jun-13 to 2016	9.00	180	Norpress
Monoamine-Oxidase Inhibitors - Non-Selective			
PHENELZINE SULPHATE			
Tab 15 mg			
TRANLYCYPROMINE SULPHATE			
Tab 10 mg			
Monoamine-Oxidase Type A Inhibitors			
MOCLOBEMIDE			
Tab 150 mg – 1% DV Oct-15 to 2018	85.10	500	Apo-Moclobemide
Tab 300 mg – 1% DV Oct-15 to 2018	30.70	100	Apo-Moclobemide
Other Antidepressants			
MIRTAZAPINE			
Tab 30 mg – 1% DV Nov-15 to 2018	2.55	30	Apo-Mirtazapine
Tab 45 mg – 1% DV Nov-15 to 2018	3.25	30	Apo-Mirtazapine

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
VENLAFAXINE – Some items restricted see terms below			
Tab modified release 37.5 mg	5.06	28	Arrow-Venlafaxine XR
Tab modified release 75 mg	6.44	28	Arrow-Venlafaxine XR
Tab modified release 150 mg	8.86	28	Arrow-Venlafaxine XR
Tab modified release 225 mg	14.34	28	Arrow-Venlafaxine XR
⚡ Cap modified release 37.5 mg	5.69	28	Efexor XR
⚡ Cap modified release 75 mg	11.40	28	Efexor XR
⚡ Cap modified release 150 mg	13.98	28	Efexor XR

➡ Restricted

Initiation

Re-assessment required after 2 years

- Both:
- 1 The patient has 'treatment-resistant' depression; and
 - 2 Either:
 - 2.1 The patient must have had a trial of two different antidepressants and have had an inadequate response from an adequate dose over an adequate period of time (usually at least four weeks); or
 - 2.2 Both:
 - 2.2.1 The patient is currently a hospital in-patient as a result of an acute depressive episode; and
 - 2.2.2 The patient must have had a trial of one other antidepressant and have had an inadequate response from an adequate dose over an adequate period of time.

Continuation

Re-assessment required after 2 years

The patient has a high risk of relapse (prescriber determined).

Selective Serotonin Reuptake Inhibitors

CITALOPRAM HYDROBROMIDE			
Tab 20 mg – 1% DV Jan-16 to 2018	1.79	84	PSM Citalopram
ESCITALOPRAM			
Tab 10 mg – 1% DV Jul-15 to 2016	1.40	28	Air Flow Products
Tab 20 mg – 1% DV Jul-15 to 2016	2.40	28	Air Flow Products
FLUOXETINE HYDROCHLORIDE			
Tab dispersible 20 mg, scored – 1% DV Apr-14 to 2016	2.50	30	Arrow-Fluoxetine
Cap 20 mg – 1% DV Apr-14 to 2016	1.74	90	Arrow-Fluoxetine
PAROXETINE HYDROCHLORIDE			
Tab 20 mg	4.32	90	Loxamine
SERTRALINE			
Tab 50 mg	3.64	90	Arrow-Sertraline
Tab 100 mg – 1% DV Sep-13 to 2016	6.28	90	Arrow-Sertraline

Antiepilepsy Drugs

Agents for the Control of Status Epilepticus

CLONAZEPAM			
Inj 1 mg per ml, 1 ml ampoule	19.00	5	Rivotril
DIAZEPAM			
Inj 5 mg per ml, 2 ml ampoule	11.83	5	Hospira
Rectal tubes 5 mg	25.05	5	Stesolid
Rectal tubes 10 mg	30.50	5	Stesolid

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LORAZEPAM			
Inj 2 mg vial			
Inj 4 mg per ml, 1 ml vial			
PARALDEHYDE			
Inj 5 ml ampoule			
PHENYTOIN SODIUM			
Inj 50 mg per ml, 2 ml ampoule – 1% DV Oct-15 to 2018	88.63	5	Hospira
Inj 50 mg per ml, 5 ml ampoule – 1% DV Oct-15 to 2018	133.92	5	Hospira

Control of Epilepsy

CARBAMAZEPINE			
Tab 200 mg	14.53	100	Tegretol
Tab long-acting 200 mg	16.98	100	Tegretol CR
Tab 400 mg	34.58	100	Tegretol
Tab long-acting 400 mg	39.17	100	Tegretol CR
Oral liq 20 mg per ml	26.37	250 ml	Tegretol
CLOBAZAM			
Tab 10 mg			
CLONAZEPAM			
Oral drops 2.5 mg per ml			
ETHOSUXIMIDE			
Cap 250 mg			
Oral liq 50 mg per ml			
GABAPENTIN – Restricted see terms below			
⚡ Cap 100 mg	7.16	100	Arrow-Gabapentin Neurontin Nupentin
⚡ Cap 300 mg	11.00	100	Arrow-Gabapentin Neurontin Nupentin
⚡ Cap 400 mg	13.75	100	Arrow-Gabapentin Neurontin Nupentin

➡Restricted

Initiation — preoperative and/or postoperative use

Limited to 8 days treatment

Initiation — pain management of burns patients

Re-assessment required after 1 month

Continuation — pain management of burns patients

Re-assessment required after 1 month

The treatment remains appropriate and the patient is benefiting from treatment.

Initiation — epilepsy

Re-assessment required after 15 months

Either:

- 1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
- 2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued. . .

Note: "Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Continuation — epilepsy

Patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life.

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective

Initiation — Neuropathic pain or Chronic Kidney Disease-associated pruritus

Re-assessment required after 3 months

Either:

- 1 The patient has been diagnosed with neuropathic pain; or
- 2 Both:
 - 2.1 The patient has Chronic Kidney Disease Stage 5-associated pruritus* where no other cause for pruritus can be identified (e.g. scabies, allergy); and
 - 2.2 The patient has persistent pruritus not relieved with a trial of emollient/moisturising creams alone.

Continuation — Neuropathic pain or Chronic Kidney Disease-associated pruritus

Either:

- 1 The patient has demonstrated a marked improvement in their control of pain or itch (prescriber determined); or
- 2 The patient has previously demonstrated clinical responsiveness to gabapentin and has now developed neuropathic pain in a new site.

Note: Indications marked with * are Unapproved Indications. Dosage adjustment of gabapentin is recommended for patients with renal impairment.

LACOSAMIDE – **Restricted** see terms below

⚡ Tab 50 mg	25.04	14	Vimpat
⚡ Tab 100 mg	50.06	14	Vimpat
	200.24	56	Vimpat
⚡ Tab 150 mg	75.10	14	Vimpat
	300.40	56	Vimpat
⚡ Tab 200 mg	400.55	56	Vimpat
⚡ Inj 10 mg per ml, 20 ml vial			

→ Restricted

Initiation

Re-assessment required after 15 months

Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Women of childbearing age are not required to have a trial of sodium valproate.

Continuation

Patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LAMOTRIGINE			
Tab dispersible 2 mg	6.74	30	Lamictal
Tab dispersible 5 mg	15.00	56	Arrow-Lamotrigine
	9.64	30	Lamictal
Tab dispersible 25 mg	20.40	56	Arrow-Lamotrigine
	29.09		Lamictal
	19.38		Logem
Tab dispersible 50 mg	34.70	56	Arrow-Lamotrigine
	47.89		Lamictal
	32.97		Logem
Tab dispersible 100 mg	59.90	56	Arrow-Lamotrigine
	79.16		Lamictal
	56.91		Logem
LEVETIRACETAM			
Tab 250 mg	24.03	60	Everet
			Levetiracetam-Rex
Tab 500 mg	28.71	60	Everet
			Levetiracetam-Rex
Tab 750 mg	45.23	60	Everet
			Levetiracetam-Rex
Tab 1,000 mg	59.12	60	Everet
Inj 100 mg per ml, 5 ml vial			
<i>(Levetiracetam-Rex Tab 250 mg to be delisted 1 August 2016)</i>			
<i>(Levetiracetam-Rex Tab 500 mg to be delisted 1 August 2016)</i>			
<i>(Levetiracetam-Rex Tab 750 mg to be delisted 1 August 2016)</i>			
PHENOBARBITONE			
Tab 15 mg – 1% DV Dec-15 to 2018	30.00	500	PSM
Tab 30 mg – 1% DV Dec-15 to 2018	31.00	500	PSM
PHENYTOIN			
Tab 50 mg			
PHENYTOIN SODIUM			
Cap 30 mg			
Cap 100 mg			
Oral liq 6 mg per ml			
PRIMIDONE			
Tab 250 mg			
SODIUM VALPROATE			
Tab 100 mg			
Tab EC 200 mg			
Tab EC 500 mg			
Oral liq 40 mg per ml			
Inj 100 mg per ml, 4 ml vial – 1% DV Sep-15 to 2018	16.60	1	Epilim IV
STIRIPENTOL – Restricted see terms on the next page			
☞ Cap 250 mg	509.29	60	Diacomit
☞ Powder for oral liq 250 mg sachet	509.29	60	Diacomit

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Initiation			
Paediatric neurologist			
<i>Re-assessment required after 6 months</i>			
Both:			
1 Patient has confirmed diagnosis of Dravet syndrome; and			
2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.			
Continuation			
Paediatric neurologist			
Patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.			
TOPIRAMATE			
Tab 25 mg	11.07	60	Arrow-Topiramate
	26.04		Topamax
	11.07		Topiramate Actavis
Tab 50 mg	18.81	60	Arrow-Topiramate
	44.26		Topamax
	18.81		Topiramate Actavis
Tab 100 mg	31.99	60	Arrow-Topiramate
	75.25		Topamax
	31.99		Topiramate Actavis
Tab 200 mg	55.19	60	Arrow-Topiramate
	129.85		Topamax
	55.19		Topiramate Actavis
Cap sprinkle 15 mg	20.84	60	Topamax
Cap sprinkle 25 mg	26.04	60	Topamax

VIGABATRIN – **Restricted** see terms below

☿ Tab 500 mg

➔Restricted

Initiation

Re-assessment required after 15 months

Both:

- 1 Either:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Notes: "Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Continuation

Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Antimigraine Preparations

Acute Migraine Treatment

DIHYDROERGOTAMINE MESYLATE

Inj 1 mg per ml, 1 ml ampoule

ERGOTAMINE TARTRATE WITH CAFFEINE

Tab 1 mg with caffeine 100 mg

METOCLOPRAMIDE HYDROCHLORIDE WITH PARACETAMOL

Tab 5 mg with paracetamol 500 mg

RIZATRIPTAN

Tab orodispersible 10 mg – 1% DV Sep-14 to 2017	3.24	12	Rizamelt
	8.10	30	Rizamelt

SUMATRIPTAN

Tab 50 mg – 1% DV Sep-13 to 2016	29.80	100	Arrow-Sumatriptan
Tab 100 mg – 1% DV Sep-13 to 2016	54.80	100	Arrow-Sumatriptan
Inj 12 mg per ml, 0.5 ml cartridge	13.80	2	Arrow-Sumatriptan

Prophylaxis of Migraine

PIZOTIFEN

Tab 500 mcg – 1% DV Sep-15 to 2018	23.21	100	Sandomigran
--	-------	-----	--------------------

Antinausea and Vertigo Agents

APREPITANT – **Restricted** see terms below

☞ Cap 2 × 80 mg and 1 × 125 mg – 1% DV Sep-14 to 2017	100.00	3	Emend Tri-Pack
---	--------	---	-----------------------

☞ **Restricted**

Initiation

Patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

Tab 16 mg – 1% DV Jun-14 to 2017	4.95	84	Vergo 16
--	------	----	-----------------

CYCLIZINE HYDROCHLORIDE

Tab 50 mg – 1% DV Jan-16 to 2018	0.59	20	Nauzene
--	------	----	----------------

CYCLIZINE LACTATE

Inj 50 mg per ml, 1 ml ampoule	14.95	5	Nausicalm
--------------------------------------	-------	---	------------------

DOMPERIDONE

Tab 10 mg – 1% DV Dec-15 to 2018	3.20	100	Prokinex
--	------	-----	-----------------

DROPERIDOL

Inj 2.5 mg per ml, 1 ml ampoule

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GRANISETRON			
Tab 1 mg – 1% DV Jan-15 to 2017	5.98	50	Granirex
HYOSCINE HYDROBROMIDE			
Inj 400 mcg per ml, 1 ml ampoule	46.50	5	Hospira
⚡ Patch 1.5 mg – 1% DV Dec-13 to 2016	11.95	2	Scopoderm TTS
➡ Restricted			
Initiation			
Any of the following:			
1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or			
2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective; or			
3 For treatment of post-operative nausea and vomiting where cyclizine, droperidol and a 5HT3 antagonist have proven ineffective, are not tolerated or are contraindicated.			
METOCLOPRAMIDE HYDROCHLORIDE			
Tab 10 mg – 1% DV Sep-14 to 2017	1.82	100	Metamide
Oral liq 5 mg per 5 ml			
Inj 5 mg per ml, 2 ml ampoule – 1% DV Sep-14 to 2017	4.50	10	Pfizer
ONDANSETRON			
Tab 4 mg – 1% DV Jan-14 to 2016	5.51	50	Onrex
Tab dispersible 4 mg – 1% DV Oct-14 to 2017	1.00	10	Dr Reddy's Ondansetron
Tab 8 mg – 1% DV Jan-14 to 2016	6.19	50	Onrex
Tab dispersible 8 mg – 1% DV Oct-14 to 2017	1.50	10	Ondansetron ODT-DRLA
Inj 2 mg per ml, 2 ml ampoule – 1% DV Sep-13 to 2016	1.82	5	Ondanaccord
Inj 2 mg per ml, 4 ml ampoule – 1% DV Sep-13 to 2016	2.18	5	Ondanaccord
PROCHLORPERAZINE			
Tab buccal 3 mg			
Tab 5 mg – 1% DV Jun-14 to 2017	9.75	500	Antinaus
Inj 12.5 mg per ml, 1 ml ampoule			
Suppos 25 mg			
PROMETHAZINE THEOCLATE – Restricted: For continuation only			
➡ Tab 25 mg			
TROPISETRON			
Inj 1 mg per ml, 2 ml ampoule – 1% DV Sep-15 to 2018	8.95	1	Tropisetron-AFT
Inj 1 mg per ml, 5 ml ampoule – 1% DV Sep-15 to 2018	13.95	1	Tropisetron-AFT
Antipsychotic Agents			
General			
AMISULPRIDE			
Tab 100 mg – 1% DV Jul-13 to 2016	6.22	30	Solian
Tab 200 mg – 1% DV Jul-13 to 2016	21.92	60	Solian
Tab 400 mg – 1% DV Jul-13 to 2016	44.52	60	Solian
Oral liq 100 mg per ml – 1% DV Jul-13 to 2016	52.50	60 ml	Solian

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ARIPIPRAZOLE – Restricted see terms below			
⚡ Tab 5 mg	123.54	30	Abilify
⚡ Tab 10 mg	123.54	30	Abilify
⚡ Tab 15 mg	175.28	30	Abilify
⚡ Tab 20 mg	213.42	30	Abilify
⚡ Tab 30 mg	260.07	30	Abilify

➡ Restricted

Initiation — schizophrenia or related psychoses

Any specialist

Both:

- 1 Patient is suffering from schizophrenia or related psychoses; and
- 2 Either:
 - 2.1 An effective dose of risperidone or quetiapine has been trialled and has been discontinued, or is in the process of being discontinued, because of unacceptable side effect; or
 - 2.2 An effective dose of risperidone or quetiapine has been trialled and has been discontinued, or is in the process of being discontinued, because of inadequate clinical response.

Initiation — Autism spectrum disorder*

Psychiatrist or paediatrician

All of the following:

- 1 The patient has been diagnosed with an autism spectrum disorder* and has symptoms of severe irritability; and
- 2 An effective dose of risperidone has been trialled and has been discontinued because of unacceptable side effects or inadequate response; and
- 3 The patient is aged less than 18 years.

Note: Indications marked with * are Unapproved Indications

CHLORPROMAZINE HYDROCHLORIDE

Tab 10 mg
Tab 25 mg
Tab 100 mg
Oral liq 10 mg per ml
Inj 25 mg per ml, 2 ml ampoule

CLOZAPINE

Tab 25 mg	6.69	50	Clopine
	13.37	100	Clopine
	5.69	50	Clozaril
	11.36	100	Clozaril
Tab 50 mg	8.67	50	Clopine
	17.33	100	Clopine
Tab 100 mg	17.33	50	Clopine
	34.65	100	Clopine
	14.73	50	Clozaril
	29.45	100	Clozaril
Tab 200 mg	34.65	50	Clopine
	69.30	100	Clopine
Oral liq 50 mg per ml	17.33	100 ml	Clopine

HALOPERIDOL

Tab 500 mcg – 1% DV Oct-13 to 2016	6.23	100	Serenace
Tab 1.5 mg – 1% DV Oct-13 to 2016	9.43	100	Serenace
Tab 5 mg – 1% DV Oct-13 to 2016	29.72	100	Serenace
Oral liq 2 mg per ml – 1% DV Oct-13 to 2016	23.84	100 ml	Serenace
Inj 5 mg per ml, 1ml ampoule – 1% DV Oct-13 to 2016	21.55	10	Serenace

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LEVOMEPROMAZINE			
Tab 25 mg			
Tab 100 mg			
Inj 25 mg per ml, 1 ml ampoule			
LITHIUM CARBONATE			
Tab long-acting 400 mg			
Tab 250 mg – 1% DV Sep-15 to 2018	34.30	500	Lithicarb FC
Tab 400 mg – 1% DV Sep-15 to 2018	12.83	100	Lithicarb FC
Cap 250 mg – 1% DV Sep-14 to 2017	9.42	100	Douglas
OLANZAPINE			
Tab 2.5 mg – 1% DV Sep-14 to 2017	0.75	28	Zypine
Tab 5 mg – 1% DV Sep-14 to 2017	1.65	28	Zypine
Tab orodispersible 5 mg – 1% DV Sep-14 to 2017	1.75	28	Zypine ODT
Tab 10 mg – 1% DV Sep-14 to 2017	2.55	28	Zypine
Tab orodispersible 10 mg – 1% DV Sep-14 to 2017	3.05	28	Zypine ODT
Inj 10 mg vial			
PERICYAZINE			
Tab 2.5 mg			
Tab 10 mg			
QUETIAPINE			
Tab 25 mg – 1% DV Sep-14 to 2017	2.10	90	Quetapel
Tab 100 mg – 1% DV Sep-14 to 2017	4.20	90	Quetapel
Tab 200 mg – 1% DV Sep-14 to 2017	7.20	90	Quetapel
Tab 300 mg – 1% DV Sep-14 to 2017	12.00	90	Quetapel
RISPERIDONE – Some items restricted see terms below			
Tab 0.5 mg – 1% DV Feb-15 to 2017	1.90	60	Actavis
⚡ Tab orodispersible 0.5 mg	21.42	28	Risperdal Quicklet
Tab 1 mg – 1% DV Feb-15 to 30 Sep 2017	2.10	60	Actavis
⚡ Tab orodispersible 1 mg	42.84	28	Risperdal Quicklet
Tab 2 mg – 1% DV Feb-15 to 2017	2.34	60	Actavis
⚡ Tab orodispersible 2 mg	85.71	28	Risperdal Quicklet
Tab 3 mg – 1% DV Feb-15 to 2017	2.55	60	Actavis
Tab 4 mg – 1% DV Feb-15 to 2017	3.50	60	Actavis
Oral liq 1 mg per ml – 1% DV Sep-14 to 2017	9.75	30 ml	Risperon
➡Restricted			
Initiation — Acute situations			
Both:			
1 For a non-adherent patient on oral therapy with standard risperidone tablets or risperidone oral liquid; and			
2 The patient is under direct supervision for administration of medicine.			
Initiation — Chronic situations			
Both:			
1 The patient is unable to take standard risperidone tablets or oral liquid, or once stabilised refuses to take risperidone tablets or oral liquid; and			
2 The patient is under direct supervision for administration of medicine.			
TRIFLUOPERAZINE HYDROCHLORIDE			
Tab 1 mg			
Tab 2 mg			
Tab 5 mg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ZIPRASIDONE			
Cap 20 mg – 1% DV Jan-16 to 2018	14.56	60	Zusdone
Cap 40 mg – 1% DV Jan-16 to 2018	24.75	60	Zusdone
Cap 60 mg – 1% DV Jan-16 to 2018	33.87	60	Zusdone
Cap 80 mg – 1% DV Jan-16 to 2018	39.74	60	Zusdone
ZUCLOPENTHIXOL ACETATE			
Inj 50 mg per ml, 1 ml ampoule			
Inj 50 mg per ml, 2 ml ampoule			
ZUCLOPENTHIXOL HYDROCHLORIDE			
Tab 10 mg	31.45	100	Clopixol

Depot Injections

FLUPENTHIXOL DECANOATE			
Inj 20 mg per ml, 1 ml ampoule	13.14	5	Fluanxol
Inj 20 mg per ml, 2 ml ampoule	20.90	5	Fluanxol
Inj 100 mg per ml, 1 ml ampoule	40.87	5	Fluanxol
FLUPHENAZINE DECANOATE			
Inj 12.5 mg per 0.5 ml ampoule	17.60	5	Modecate
Inj 25 mg per ml, 1 ml ampoule	27.90	5	Modecate
Inj 25 mg per ml, 2 ml ampoule			e.g. Modecate
Inj 100 mg per ml, 1 ml ampoule	154.50	5	Modecate
HALOPERIDOL DECANOATE			
Inj 50 mg per ml, 1 ml ampoule	28.39	5	Haldol
Inj 100 mg per ml, 1 ml ampoule	55.90	5	Haldol Concentrate
OLANZAPINE – Restricted see terms below			
⚡ Inj 210 mg vial	280.00	1	Zyprexa Relprevv
⚡ Inj 300 mg vial	460.00	1	Zyprexa Relprevv
⚡ Inj 405 mg vial	560.00	1	Zyprexa Relprevv

➡ Restricted

Initiation

Re-assessment required after 12 months

Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or paliperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Continuation

Re-assessment required after 12 months

The initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE – Restricted see terms on the next page

⚡ Inj 25 mg syringe	194.25	1	Invega Sustenna
⚡ Inj 50 mg syringe	271.95	1	Invega Sustenna
⚡ Inj 75 mg syringe	357.42	1	Invega Sustenna
⚡ Inj 100 mg syringe	435.12	1	Invega Sustenna
⚡ Inj 150 mg syringe	435.12	1	Invega Sustenna

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

➔ **Restricted**

Initiation

Re-assessment required after 12 months

Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Continuation

Re-assessment required after 12 months

The initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PIPOTHIAZINE PALMITATE – **Restricted**: For continuation only

- ➔ Inj 50 mg per ml, 1 ml ampoule
- ➔ Inj 50 mg per ml, 2 ml ampoule

RISPERIDONE – **Restricted** see terms below

⚡ Inj 25 mg vial	135.98	1	Risperdal Consta
⚡ Inj 37.5 mg vial	178.71	1	Risperdal Consta
⚡ Inj 50 mg vial	217.56	1	Risperdal Consta

➔ **Restricted**

Initiation

Re-assessment required after 12 months

Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Continuation

Re-assessment required after 12 months

The initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE

Inj 200 mg per ml, 1 ml ampoule	19.80	5	Clopixol
Inj 500 mg per ml, 1 ml ampoule			<i>e.g. Clopixol Conc</i>

Anxiolytics

ALPRAZOLAM

- Tab 1 mg
- Tab 250 mcg
- Tab 500 mcg

BUSPIRONE HYDROCHLORIDE

Tab 5 mg	28.00	100	Pacific Buspirone
Tab 10 mg	17.00	100	Pacific Buspirone

CLONAZEPAM

Tab 500 mcg	7.53	100	Paxam
Tab 2 mg	14.37	100	Paxam

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DIAZEPAM			
Tab 2 mg	11.44	500	Arrow-Diazepam
Tab 5 mg	13.71	500	Arrow-Diazepam
LORAZEPAM			
Tab 1 mg – 1% DV Jun-15 to 2018	10.79	250	Ativan
Tab 2.5 mg – 1% DV Jun-15 to 2018	13.88	100	Ativan
OXAZEPAM			
Tab 10 mg – 1% DV Dec-14 to 2017	6.17	100	Ox-Pam
Tab 15 mg – 1% DV Dec-14 to 2017	8.53	100	Ox-Pam

Multiple Sclerosis Treatments

DIMETHYL FUMARATE – Restricted see terms below

☞ Cap 120 mg	520.00	14	Tecfidera
☞ Cap 240 mg	2,000.00	56	Tecfidera

☞ **Restricted**

Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

FINGOLIMOD – Restricted see terms below

☞ Cap 0.5 mg	2,650.00	28	Gilenya
--------------------	----------	----	---------

☞ **Restricted**

Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

NATALIZUMAB – Restricted see terms below

☞ Inj 20 mg per ml, 15 ml vial	1,750.00	1	Tysabri
--------------------------------------	----------	---	---------

☞ **Restricted**

Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

TERIFLUNOMIDE – Restricted see terms below

☞ Tab 14 mg	1,582.62	28	Aubagio
-------------------	----------	----	---------

☞ **Restricted**

Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

Other Multiple Sclerosis Treatments

☞ **Restricted**

Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

GLATIRAMER ACETATE – Restricted see terms above

☞ Inj 20 mg per ml, 1 ml syringe

☞ Item restricted (see ☞ above); ☞ Item restricted (see ☞ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
INTERFERON BETA-1-ALPHA – Restricted see terms on the preceding page			
⬆ Inj 6 million iu in 0.5 ml pen injector	1,170.00	4	Avonex Pen
⬆ Inj 6 million iu in 0.5 ml syringe	1,170.00	4	Avonex
⬆ Inj 6 million iu vial	1,170.00	4	Avonex
INTERFERON BETA-1-BETA – Restricted see terms on the preceding page			
⬆ Inj 8 million iu per ml, 1 ml vial			
Sedatives and Hypnotics			
CHLORAL HYDRATE			
Oral liq 100 mg per ml			
Oral liq 200 mg per ml			
LORMETAZEPAM – Restricted: For continuation only			
➡ Tab 1 mg			
MELATONIN – Restricted see terms below			
⬆ Tab modified-release 2 mg			<i>e.g. Circadin</i>
⬆ Tab 1 mg			
⬆ Tab 2 mg			
⬆ Tab 3 mg			
⬆ Cap 2 mg			
⬆ Cap 3 mg			
➡ Restricted			
Initiation			
For in hospital use only. For the treatment of insomnia where benzodiazepines and zopiclone are contraindicated.			
MIDAZOLAM			
Tab 7.5 mg	40.00	100	Hypnovel
Oral liq 2 mg per ml			
Inj 1 mg per ml, 5 ml ampoule	10.75	10	Hypnovel
	10.00		Pfizer
Inj 5 mg per ml, 3 ml ampoule	11.90	5	Hypnovel
			Pfizer
NITRAZEPAM			
Tab 5 mg – 1% DV Dec-14 to 2017	5.22	100	Nitrados
PHENOBARBITONE			
Inj 200 mg per ml, 1 ml ampoule			
TEMAZEPAM			
Tab 10 mg – 1% DV Sep-14 to 2017	1.27	25	Normison
TRIAZOLAM – Restricted: For continuation only			
➡ Tab 125 mcg			
➡ Tab 250 mcg			
ZOPICLONE			
Tab 7.5 mg – 1% DV Dec-15 to 2018	0.98	30	Zopiclone Actavis
	8.99	500	Zopiclone Actavis

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Stimulants / ADHD Treatments

ATOMOXETINE – **Restricted** see terms below

⚡ Cap 10 mg	107.03	28	Strattera
⚡ Cap 18 mg	107.03	28	Strattera
⚡ Cap 25 mg	107.03	28	Strattera
⚡ Cap 40 mg	107.03	28	Strattera
⚡ Cap 60 mg	107.03	28	Strattera
⚡ Cap 80 mg	139.11	28	Strattera
⚡ Cap 100 mg	139.11	28	Strattera

➡ **Restricted**

Initiation

All of the following:

- 1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder) diagnosed according to DSM-IV or ICD 10 criteria; and
- 2 Once-daily dosing; and
- 3 Any of the following:
 - 3.1 Treatment with a subsidised formulation of a stimulant has resulted in the development or worsening of serious adverse reactions or where the combination of subsidised stimulant treatment with another agent would pose an unacceptable medical risk; or
 - 3.2 Treatment with a subsidised formulation of a stimulant has resulted in worsening of co-morbid substance abuse or there is a significant risk of diversion with subsidised stimulant therapy; or
 - 3.3 An effective dose of a subsidised formulation of a stimulant has been trialled and has been discontinued because of inadequate clinical response; or
 - 3.4 Treatment with a subsidised formulation of a stimulant is considered inappropriate because the patient has a history of psychoses or has a first-degree relative with schizophrenia; and
- 4 The patient will not be receiving treatment with atomoxetine in combination with a subsidised formulation of a stimulant, except for the purposes of transitioning from subsidised stimulant therapy to atomoxetine.

Note: A "subsidised formulation of a stimulant" refers to currently listed methylphenidate hydrochloride tablet formulations (immediate-release, sustained-release and extended-release) or dexamphetamine sulphate tablets.

CAFFEINE

Tab 100 mg

DEXAMFETAMINE SULFATE – **Restricted** see terms below

⚡ Tab 5 mg – 1% DV Dec-15 to 2018	17.00	100	PSM
---	-------	-----	-----

➡ **Restricted**

Initiation — ADHD

Paediatrician or psychiatrist

Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria.

Initiation — Narcolepsy

Neurologist or respiratory specialist

Re-assessment required after 24 months

Patient suffers from narcolepsy.

Continuation — Narcolepsy

Neurologist or respiratory specialist

Re-assessment required after 24 months

The treatment remains appropriate and the patient is benefiting from treatment.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
METHYLPHENIDATE HYDROCHLORIDE – Restricted see terms below			
⚡ Tab extended-release 18 mg	58.96	30	Concerta
⚡ Tab extended-release 27 mg	65.44	30	Concerta
⚡ Tab extended-release 36 mg	71.93	30	Concerta
⚡ Tab extended-release 54 mg	86.24	30	Concerta
⚡ Tab immediate-release 5 mg	3.20	30	Rubifen
⚡ Tab immediate-release 10 mg	3.00	30	Ritalin
			Rubifen
⚡ Tab immediate-release 20 mg	7.85	30	Rubifen
⚡ Tab sustained-release 20 mg	50.00	100	Ritalin SR
	10.95	30	Rubifen SR
⚡ Cap modified-release 10 mg	15.60	30	Ritalin LA
⚡ Cap modified-release 20 mg	20.40	30	Ritalin LA
⚡ Cap modified-release 30 mg	25.52	30	Ritalin LA
⚡ Cap modified-release 40 mg	30.60	30	Ritalin LA

➡ **Restricted**

Initiation — ADHD (immediate-release and sustained-release formulations)

Paediatrician or psychiatrist

Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria.

Initiation — Narcolepsy (immediate-release and sustained-release formulations)

Neurologist or respiratory specialist

Re-assessment required after 24 months

Patient suffers from narcolepsy.

Continuation — Narcolepsy (immediate-release and sustained-release formulations)

Neurologist or respiratory specialist

Re-assessment required after 24 months

The treatment remains appropriate and the patient is benefiting from treatment.

Initiation — Extended-release and modified-release formulations

Paediatrician or psychiatrist

Both:

- 1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria; and
- 2 Either:
 - 2.1 Patient is taking a currently listed formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 2.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

MODAFINIL – Restricted see terms below

⚡ Tab 100 mg

➡ **Restricted**

Initiation — Narcolepsy

Neurologist or respiratory specialist

Re-assessment required after 24 months

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

3 Either:

3.1 An effective dose of a listed formulation of methylphenidate or dexamphetamine has been trialled and discontinued because of intolerable side effects; or

3.2 Methylphenidate and dexamphetamine are contraindicated.

Continuation — Narcolepsy

Neurologist or respiratory specialist

Re-assessment required after 24 months

The treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE

Tab 5 mg – 1% DV Feb-15 to 2017.....	5.48	90	Donepezil-Rex
Tab 10 mg – 1% DV Feb-15 to 2017.....	10.51	90	Donepezil-Rex

RIVASTIGMINE – Restricted see terms below

⚡ Patch 4.6 mg per 24 hour	90.00	30	Exelon
⚡ Patch 9.5 mg per 24 hour	90.00	30	Exelon

➡Restricted

Initiation

Re-assessment required after 6 months

Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Continuation

Re-assessment required after 12 months

Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE – Restricted see terms below

⚡ Tab 2 mg with naloxone 0.5 mg	57.40	28	Suboxone
⚡ Tab 8 mg with naloxone 2 mg	166.00	28	Suboxone

➡Restricted

Initiation — Detoxification

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Prescriber works in an opioid treatment service approved by the Ministry of Health.

Initiation — Maintenance treatment

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Prescriber works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

Tab modified-release 150 mg – 1% DV Oct-13 to 2016.....	4.97	30	Zyban
---	------	----	-------

DISULFIRAM

Tab 200 mg	24.30	100	Antabuse
------------------	-------	-----	----------

⚡Item restricted (see ➡ above); ⚡Item restricted (see ➡ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
NALTREXONE HYDROCHLORIDE – Restricted see terms below			
⚡ Tab 50 mg – 1% DV Sep-13 to 2016	76.00	30	Naltraccord
➡ Restricted			
Initiation — Alcohol dependence			
Both:			
1 Patient is currently enrolled, or is planned to be enrolled, in a recognised comprehensive treatment programme for alcohol dependence; and			
2 Naltrexone is to be prescribed by, or on the recommendation of, a physician working in an Alcohol and Drug Service.			
Initiation — Constipation			
For the treatment of opioid-induced constipation.			
NICOTINE – Some items restricted see terms below			
Patch 7 mg per 24 hours – 1% DV Apr-14 to 2017	10.57	28	Habitrol
Patch 14 mg per 24 hours – 1% DV Apr-14 to 2017	11.31	28	Habitrol
Patch 21 mg per 24 hours – 1% DV Apr-14 to 2017	11.95	28	Habitrol
⚡ Oral spray 1 mg per dose			<i>e.g. Nicorette QuickMist Mouth Spray</i>
Lozenge 1 mg – 1% DV Apr-14 to 2017	12.91	216	Habitrol
Lozenge 2 mg – 1% DV Apr-14 to 2017	14.14	216	Habitrol
⚡ Soln for inhalation 15 mg cartridge			<i>e.g. Nicorette Inhalator</i>
Gum 2 mg – 1% DV Apr-14 to 2017	22.26	384	Habitrol (Classic)
			Habitrol (Fruit)
			Habitrol (Mint)
Gum 4 mg – 1% DV Apr-14 to 2017	25.67	384	Habitrol (Classic)
			Habitrol (Fruit)
			Habitrol (Mint)
➡ Restricted			
Initiation			
Any of the following:			
1 For perioperative use in patients who have a 'nil by mouth' instruction; or			
2 For use within mental health inpatient units; or			
3 For acute use in agitated patients who are unable to leave the hospital facilities.			
VARENICLINE – Restricted see terms below			
⚡ Tab 0.5 mg × 11 and 1 mg × 14	60.48	25	Champix
⚡ Tab 1 mg	67.74	28	Champix
	135.48	56	Champix
➡ Restricted			
Initiation			
All of the following:			
1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and			
2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and			
3 Either:			
3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or			
3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and			
4 The patient has not used funded varenicline in the last 12 months; and			

continued...

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

continued...

- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline in a 12 month period.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Chemotherapeutic Agents			
Alkylating Agents			
BUSULFAN			
Tab 2 mg	89.25	100	Myleran
Inj 6 mg per ml, 10 ml ampoule			
CARMUSTINE			
Inj 100 mg vial – 1% DV Sep-15 to 2018	532.00	1	BiCNU
CHLORAMBUCIL			
Tab 2 mg			
CYCLOPHOSPHAMIDE			
Tab 50 mg	79.00	50	Endoxan
	158.00	100	Procytox
Inj 1 g vial – 1% DV Oct-15 to 2018	35.03	1	Endoxan
Inj 2 g vial – 1% DV Oct-15 to 2018	70.06	1	Endoxan
IFOSFAMIDE			
Inj 1 g vial	96.00	1	Holoxan
Inj 2 g vial	180.00	1	Holoxan
LOMUSTINE			
Cap 10 mg	132.59	20	Ceenu
Cap 40 mg	399.15	20	Ceenu
MELPHALAN			
Tab 2 mg			
Inj 50 mg vial			
THIOTEPA			
Inj 15 mg vial			
Inj 100 mg vial			
Anthracyclines and Other Cytotoxic Antibiotics			
BLEOMYCIN SULPHATE			
Inj 15,000 iu vial – 1% DV Oct-15 to 2018	150.48	1	DBL Bleomycin Sulfate
DACTINOMYCIN [ACTINOMYCIN D]			
Inj 0.5 mg vial	145.00	1	Cosmegen
DAUNORUBICIN			
Inj 2 mg per ml, 10 ml vial – 1% DV Aug-13 to 2016	118.72	1	Pfizer
DOXORUBICIN HYDROCHLORIDE			
Inj 2 mg per ml, 5 ml vial			
Inj 2 mg per ml, 25 ml vial – 1% DV Feb-16 to 2018	11.50	1	Doxorubicin Ebewe
Note: DV limit applies to all 50 mg presentations of doxorubicin hydrochloride.			
Inj 50 mg vial			
Inj 2 mg per ml, 50 ml vial – 1% DV Feb-16 to 2018	23.00	1	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial – 1% DV Feb-16 to 2018	46.00	1	Doxorubicin Ebewe

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
EPIRUBICIN HYDROCHLORIDE			
Inj 2 mg per ml, 5 ml vial	25.00	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial – 1% DV Nov-15 to 2018	30.00	1	Epirubicin Ebewe
Inj 2 mg per ml, 50 ml vial – 1% DV Nov-15 to 2018	32.50	1	Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial – 1% DV Nov-15 to 2018	65.00	1	Epirubicin Ebewe
IDARUBICIN HYDROCHLORIDE			
Inj 5 mg vial – 1% DV Nov-15 to 2018	125.00	1	Zavedos
Inj 10 mg vial – 1% DV Nov-15 to 2018	250.00	1	Zavedos
MITOMYCIN C			
Inj 5 mg vial – 1% DV Oct-13 to 2016	79.75	1	Arrow
MITOZANTRONE			
Inj 2 mg per ml, 10 ml vial – 1% DV Sep-15 to 2018	97.50	1	Mitozantrone Ebewe
Antimetabolites			
AZACITIDINE – Restricted see terms below			
☛ Inj 100 mg vial	605.00	1	Vidaza
☛ Restricted			
Initiation			
Haematologist			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 Any of the following:			
1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or			
1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or			
1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and			
2 The patient has performance status (WHO/ECOG) grade 0-2; and			
3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and			
4 The patient has an estimated life expectancy of at least 3 months.			
Continuation			
Haematologist			
<i>Re-assessment required after 12 months</i>			
Both:			
1 No evidence of disease progression, and; and			
2 The treatment remains appropriate and patient is benefitting from treatment.			
CAPECITABINE			
Tab 150 mg – 1% DV Sep-14 to 2016	30.00	60	Capecitabine Winthrop
Tab 500 mg – 1% DV Sep-14 to 2016	120.00	120	Capecitabine Winthrop
CLADRIBINE			
Inj 2 mg per ml, 5 ml vial			
Inj 1 mg per ml, 10 ml vial	5,249.72	7	Leustatin
CYTARABINE			
Inj 20 mg per ml, 5 ml vial – 1% DV Nov-13 to 2016	55.00	5	Pfizer
Inj 20 mg per ml, 25 ml vial	18.15	1	Pfizer
Inj 100 mg per ml, 10 ml vial – 1% DV Nov-13 to 2016	8.83	1	Pfizer
Inj 100 mg per ml, 20 ml vial – 1% DV Nov-13 to 2016	17.65	1	Pfizer

☛ Item restricted (see ☛ above); ☛ Item restricted (see ☛ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FLUDARABINE PHOSPHATE			
Tab 10 mg – 1% DV Sep-15 to 2018	412.00	20	Fludara Oral
Inj 50 mg vial	525.00	5	Fludarabine Ebewe
FLUOROURACIL			
Inj 50 mg per ml, 20 ml vial – 1% DV Oct-15 to 2018	10.00	1	Fluorouracil Ebewe
Inj 50 mg per ml, 50 ml vial – 1% DV Oct-15 to 2018	17.00	1	Fluorouracil Ebewe
Inj 50 mg per ml, 100 ml vial – 1% DV Oct-15 to 2018	30.00	1	Fluorouracil Ebewe
GEMCITABINE			
Inj 10 mg per ml, 20 ml vial – 1% DV Oct-14 to 2017	8.36	1	Gemcitabine Ebewe
Inj 10 mg per ml, 100 ml vial – 1% DV Oct-14 to 2017	15.89	1	Gemcitabine Ebewe
MERCAPTOPURINE			
Tab 50 mg – 1% DV Oct-13 to 2016	49.41	25	Puri-nethol
METHOTREXATE			
Tab 2.5 mg – 1% DV Sep-15 to 2018	3.18	30	Trexate
Tab 10 mg – 1% DV Sep-15 to 2018	21.00	50	Trexate
Inj 2.5 mg per ml, 2 ml vial			
Inj 7.5 mg prefilled syringe – 1% DV Jan-14 to 2016	17.19	1	Methotrexate Sandoz
Inj 10 mg prefilled syringe – 1% DV Jan-14 to 2016	17.25	1	Methotrexate Sandoz
Inj 15 mg prefilled syringe – 1% DV Jan-14 to 2016	17.38	1	Methotrexate Sandoz
Inj 20 mg prefilled syringe – 1% DV Jan-14 to 2016	17.50	1	Methotrexate Sandoz
Inj 25 mg prefilled syringe – 1% DV Jan-14 to 2016	17.63	1	Methotrexate Sandoz
Inj 30 mg prefilled syringe – 1% DV Jan-14 to 2016	17.75	1	Methotrexate Sandoz
Inj 25 mg per ml, 2 ml vial – 1% DV Sep-13 to 2016	20.20	5	Hospira
Inj 25 mg per ml, 20 ml vial – 1% DV Sep-13 to 2016	27.78	1	Hospira
Inj 100 mg per ml, 10 ml vial	25.00	1	Methotrexate Ebewe
Inj 100 mg per ml, 50 ml vial – 1% DV Oct-14 to 2017	99.99	1	Methotrexate Ebewe
THIOGUANINE			
Tab 40 mg			

Other Cytotoxic Agents

AMSACRINE

Inj 50 mg per ml, 1.5 ml ampoule
Inj 75 mg

ANAGRELIDE HYDROCHLORIDE

Cap 0.5 mg

ARSENIC TRIOXIDE

Inj 1 mg per ml, 10 ml vial 4,817.00 10 AFT

BORTEZOMIB – **Restricted** see terms on the next page

⚡ Inj 1 mg vial 540.70 1 Velcade
⚡ Inj 3.5 mg vial 1,892.50 1 Velcade

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➡Restricted			
Initiation — treatment naive multiple myeloma/amyloidosis			
<i>Limited to 15 months treatment</i>			
Both:			
1 Either:			
1.1 The patient has treatment-naive symptomatic multiple myeloma; or			
1.2 The patient has treatment-naive symptomatic systemic AL amyloidosis; and			
2 Maximum of 9 treatment cycles.			
Initiation — relapsed/refractory multiple myeloma/amyloidosis			
<i>Re-assessment required after 8 months</i>			
All of the following:			
1 Either:			
1.1 The patient has relapsed or refractory multiple myeloma; or			
1.2 The patient has relapsed or refractory systemic AL amyloidosis; and			
2 The patient has received only one prior front line chemotherapy for multiple myeloma or amyloidosis; and			
3 The patient has not had prior publicly funded treatment with bortezomib; and			
4 Maximum of 4 treatment cycles.			
Continuation — relapsed/refractory multiple myeloma/amyloidosis			
<i>Re-assessment required after 8 months</i>			
Both:			
1 The patient's disease obtained at least a partial response from treatment with bortezomib at the completion of cycle 4; and			
2 Maximum of 4 further treatment cycles (making a total maximum of 8 consecutive treatment cycles).			
Notes: Responding relapsed/refractory multiple myeloma patients should receive no more than 2 additional cycles of treatment beyond the cycle at which a confirmed complete response was first achieved. A line of therapy is considered to comprise either:			
1 A known therapeutic chemotherapy regimen and supportive treatments; or			
2 A transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments.			
Refer to datasheet for recommended dosage and number of doses of bortezomib per treatment cycle.			
COLASPASE [L-ASPARAGINASE]			
Inj 10,000 iu vial	102.32	1	Leunase
DACARBAZINE			
Inj 200 mg vial – 1% DV Oct-13 to 2016	51.84	1	Hospira
ETOPOSIDE			
Cap 50 mg	340.73	20	Vepesid
Cap 100 mg	340.73	10	Vepesid
Inj 20 mg per ml, 5 ml vial – 1% DV Apr-16 to 2018	7.90	1	Rex Medical
ETOPOSIDE (AS PHOSPHATE)			
Inj 100 mg vial	40.00	1	Etopophos
HYDROXYUREA			
Cap 500 mg	31.76	100	Hydrea
IRINOTECAN HYDROCHLORIDE			
Inj 20 mg per ml, 2 ml vial – 1% DV Sep-15 to 2018	11.50	1	Irinotecan Actavis 40
Inj 20 mg per ml, 5 ml vial – 1% DV Sep-15 to 2018	17.80	1	Irinotecan Actavis 100
LENALIDOMIDE – Restricted see terms on the next page			
⚡ Cap 10 mg	6,207.00	21	Revlimid
⚡ Cap 25 mg	7,627.00	21	Revlimid

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Initiation			
Haematologist			
<i>Re-assessment required after 6 months</i>			
All of the following:			
1 Patient has relapsed or refractory multiple myeloma with progressive disease; and			
2 Either:			
2.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or			
2.2 Both:			
2.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and			
2.2.2 The patient has experienced severe (grade ≥ 3), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and			
3 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.			
Continuation			
Haematologist			
<i>Re-assessment required after 6 months</i>			
Both:			
1 No evidence of disease progression; and			
2 The treatment remains appropriate and patient is benefitting from treatment.			
Note: Indication marked with * is an Unapproved Indication (refer to Interpretations and Definitions). A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.			
PEGASPARGASE – Restricted see terms below			
⚡ Inj 750 iu per ml, 5 ml vial	3,005.00	1	Oncaspar
➔Restricted			
Initiation — Newly diagnosed ALL			
<i>Limited to 12 months treatment</i>			
All of the following:			
1 The patient has newly diagnosed acute lymphoblastic leukaemia; and			
2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and			
3 Treatment is with curative intent.			
Initiation — Relapsed ALL			
<i>Limited to 12 months treatment</i>			
All of the following:			
1 The patient has relapsed acute lymphoblastic leukaemia; and			
2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and			
3 Treatment is with curative intent.			
PENTOSTATIN [DEOXYCOFORMYCIN]			
Inj 10 mg vial			
PROCARBAZINE HYDROCHLORIDE			
Cap 50 mg	498.00	50	Natulan
TEMOZOLOMIDE – Restricted see terms on the next page			
⚡ Cap 5 mg – 1% DV Sep-13 to 2016.....	8.00	5	Temaccord
⚡ Cap 20 mg – 1% DV Sep-13 to 2016.....	36.00	5	Temaccord
⚡ Cap 100 mg – 1% DV Sep-13 to 2016.....	175.00	5	Temaccord
⚡ Cap 250 mg – 1% DV Sep-13 to 2016.....	410.00	5	Temaccord

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Initiation			
All of the following:			
1 Either:			
1.1 Patient has newly diagnosed glioblastoma multiforme; or			
1.2 Patient has newly diagnosed anaplastic astrocytoma*; and			
2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and			
3 Following concomitant treatment temozolomide is to be used for a maximum of six cycles of 5 days treatment, at a maximum dose of 200 mg/m ² .			
Note: Indication marked with a * is an Unapproved Indication. Temozolomide is not funded for the treatment of relapsed glioblastoma multiforme. Reapplications will not be approved. Studies of temozolomide show that its benefit is predominantly in those patients with a good performance status (WHO grade 0 or 1 or Karnofsky score >80), and in patients who have had at least a partial resection of the tumour.			
THALIDOMIDE – Restricted see terms below			
⚡ Cap 50 mg	378.00	28	Thalomid
⚡ Cap 100 mg	756.00	28	Thalomid
➔Restricted			
Initiation			
<i>Re-assessment required after 12 months</i>			
Any of the following:			
1 The patient has multiple myeloma; or			
2 The patient has systemic AL amyloidosis*; or			
3 The patient has erythema nodosum leprosum.			
Continuation			
Patient has obtained a response from treatment during the initial approval period.			
Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier			
Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen			
Indication marked with * is an Unapproved Indication			
TRETINOIN			
Cap 10 mg	479.50	100	Vesanoid
Platinum Compounds			
CARBOPLATIN			
Inj 10 mg per ml, 5 ml vial – 1% DV Sep-15 to 2018	15.07	1	DBL Carboplatin
Inj 10 mg per ml, 15 ml vial – 1% DV Sep-15 to 2018	14.05	1	DBL Carboplatin
Inj 10 mg per ml, 45 ml vial – 1% DV Sep-15 to 2018	32.59	1	DBL Carboplatin
CISPLATIN			
Inj 1 mg per ml, 50 ml vial – 1% DV Nov-15 to 2018	12.29	1	DBL Cisplatin
Inj 1 mg per ml, 100 ml vial – 1% DV Nov-15 to 2018	22.46	1	DBL Cisplatin
OXALIPLATIN			
Inj 5 mg per ml, 10 ml vial	13.32	1	Oxaliccord
Inj 5 mg per ml, 20 ml vial	16.00	1	Oxaliccord
Protein-Tyrosine Kinase Inhibitors			
DASATINIB – Restricted see terms on the next page			
⚡ Tab 20 mg	3,774.06	60	Sprycel
⚡ Tab 50 mg	6,214.20	60	Sprycel
⚡ Tab 70 mg	7,692.58	60	Sprycel
⚡ Tab 100 mg	6,214.20	30	Sprycel

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Initiation			
For use in patients with approval from the CML/GIST Co-ordinator.			
ERLOTINIB – Restricted see terms below			
⚡ Tab 100 mg – 1% DV Jun-15 to 2018	1,000.00	30	Tarceva
⚡ Tab 150 mg – 1% DV Jun-15 to 2018	1,500.00	30	Tarceva
➔Restricted			
Initiation			
<i>Re-assessment required after 4 months</i>			
All of the following:			
1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and			
2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and			
3 Any of the following:			
3.1 Patient is treatment naive; or			
3.2 Both:			
3.2.1 Patient has documented disease progression following treatment with first line platinum based chemotherapy; and			
3.2.2 Patient has not received prior treatment with gefitinib; or			
3.3 Both:			
3.3.1 The patient has discontinued gefitinib within 12 weeks of starting treatment due to intolerance; and			
3.3.2 The cancer did not progress while on gefitinib; and			
4 Erlotinib is to be given for a maximum of 3 months.			
Continuation			
<i>Re-assessment required after 6 months</i>			
Both:			
1 Radiological assessment (preferably including CT scan) indicates NSCLC has not progressed; and			
2 Erlotinib is to be given for a maximum of 3 months.			
GEFITINIB – Restricted see terms below			
⚡ Tab 250 mg	1,700.00	30	Iressa
➔Restricted			
Initiation			
<i>Re-assessment required after 4 months</i>			
All of the following:			
1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and			
2 Either:			
2.1 Patient is treatment naive; or			
2.2 Both:			
2.2.1 The patient has discontinued erlotinib within 12 weeks of starting treatment due to intolerance; and			
2.2.2 The cancer did not progress whilst on erlotinib; and			
3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and			
4 Gefitinib is to be given for a maximum of 3 months.			
Continuation			
<i>Re-assessment required after 6 months</i>			
Both:			
1 Radiological assessment (preferably including CT scan) indicates NSCLC has not progressed; and			
2 Gefitinib is to be given for a maximum of 3 months.			
IMATINIB MESILATE			
Imatinib-AFT is not a registered for the treatment of Gastro Intestinal Stromal Tumours (GIST). The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule			
⚡ Tab 100 mg	2,400.00	60	Glivec

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➡Restricted			
Initiation			
<i>Re-assessment required after 12 months</i>			
Both:			
1 Patient has diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST); and			
2 Maximum dose of 400 mg/day.			
Continuation			
<i>Re-assessment required after 12 months</i>			
Adequate clinical response to treatment with imatinib (prescriber determined).			
Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule.			
Cap 100 mg – 1% DV Jul-14 to 2017	298.90	60	Imatinib-AFT
Cap 400 mg	597.80	30	Imatinib-AFT
LAPATINIB – Restricted see terms below			
⚡ Tab 250 mg	1,899.00	70	Tykerb
➡Restricted			
Initiation			
<i>Re-assessment required after 12 months</i>			
Either:			
1 All of the following:			
1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and			
1.2 The patient has not previously received trastuzumab treatment for HER 2 positive metastatic breast cancer; and			
1.3 Lapatinib not to be given in combination with trastuzumab; and			
1.4 Lapatinib to be discontinued at disease progression; or			
2 All of the following:			
2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and			
2.2 The patient started trastuzumab for metastatic breast cancer but discontinued trastuzumab within 3 months of starting treatment due to intolerance; and			
2.3 The cancer did not progress whilst on trastuzumab; and			
2.4 Lapatinib not to be given in combination with trastuzumab; and			
2.5 Lapatinib to be discontinued at disease progression.			
Continuation			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and			
2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and			
3 Lapatinib not to be given in combination with trastuzumab; and			
4 Lapatinib to be discontinued at disease progression.			
NILETINIB – Restricted see terms on the next page			
⚡ Cap 150 mg	4,680.00	120	Tasigna
⚡ Cap 200 mg	6,532.00	120	Tasigna

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ **Restricted**

Initiation

Haematologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Either:
 - 2.1 Patient has documented CML treatment failure* with imatinib; or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Continuation

Haematologist

Re-assessment required after 6 months

All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PAZOPANIB – **Restricted** see terms below

⚡ Tab 200 mg	1,334.70	30	Votrient
⚡ Tab 400 mg	2,669.40	30	Votrient

➔ **Restricted**

Initiation

Re-assessment required after 3 months

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
- 5 All of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; and
 - 5.2 Haemoglobin level < lower limit of normal; and
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); and
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; and
 - 5.5 Karnofsky performance score of ≤ 70; and
 - 5.6 ≥ 2 sites of organ metastasis.

Continuation

Re-assessment required after 3 months

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SUNITINIB – Restricted see terms below			
☞ Cap 12.5 mg	2,315.38	28	Sutent
☞ Cap 25 mg	4,630.77	28	Sutent
☞ Cap 50 mg	9,261.54	28	Sutent

☞Restricted
Initiation — RCC

Re-assessment required after 3 months

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
- 5 All of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; and
 - 5.2 Haemoglobin level < lower limit of normal; and
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); and
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; and
 - 5.5 Karnofsky performance score of ≤ 70; and
 - 5.6 ≥ 2 sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Notes: RCC - Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

Continuation — RCC

Re-assessment required after 3 months

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Initiation — GIST

Re-assessment required after 3 months

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Continuation — GIST

Re-assessment required after 6 months

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
1.2 The patient has had a partial response (a decrease in size of $\geq 10\%$ or decrease in tumour density in Hounsfield Units (HU) of $\geq 15\%$ on CT and no new lesions and no obvious progression of non-measurable disease); or			
1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and			
2 The treatment remains appropriate and the patient is benefiting from treatment.			
Note: GIST - It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of $\geq 10\%$ and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.			
Taxanes			
DOCETAXEL			
Inj 10 mg per ml, 2 ml vial – 1% DV Dec-14 to 2017	13.70	1	DBL Docetaxel
Inj 10 mg per ml, 8 ml vial – 1% DV Dec-14 to 2017	29.99	1	DBL Docetaxel
PACLITAXEL			
Inj 6 mg per ml, 5 ml vial – 1% DV Sep-14 to 2017	45.00	5	Paclitaxel Ebewe
Inj 6 mg per ml, 16.7 ml vial – 1% DV Sep-14 to 2017	19.02	1	Paclitaxel Ebewe
Inj 6 mg per ml, 25 ml vial – 1% DV Sep-14 to 2017	26.69	1	Paclitaxel Ebewe
Inj 6 mg per ml, 50 ml vial – 1% DV Sep-14 to 2017	36.53	1	Paclitaxel Ebewe
Inj 6 mg per ml, 100 ml vial – 1% DV Sep-14 to 2017	73.06	1	Paclitaxel Ebewe
Treatment of Cytotoxic-Induced Side Effects			
CALCIUM FOLINATE			
Tab 15 mg	104.26	10	DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml ampoule			
Inj 10 mg per ml, 5 ml ampoule – 1% DV Oct-14 to 2017	18.25	5	Calcium Folate Ebewe
Inj 10 mg per ml, 10 ml vial – 1% DV Oct-14 to 2017	7.33	1	Calcium Folate Ebewe
Inj 10 mg per ml, 30 ml vial – 1% DV Oct-14 to 2017	22.51	1	Calcium Folate Ebewe
Inj 10 mg per ml, 100 ml vial – 1% DV Oct-14 to 2017	67.51	1	Calcium Folate Ebewe
MESNA			
Tab 400 mg – 1% DV Oct-13 to 2016	227.50	50	Uromitexan
Tab 600 mg – 1% DV Oct-13 to 2016	339.50	50	Uromitexan
Inj 100 mg per ml, 4 ml ampoule – 1% DV Oct-13 to 2016	148.05	15	Uromitexan
Inj 100 mg per ml, 10 ml ampoule – 1% DV Oct-13 to 2016	339.90	15	Uromitexan
Vinca Alkaloids			
VINBLASTINE SULPHATE			
Inj 1 mg per ml, 10 ml vial	186.46	5	Hospira
VINCRISTINE SULPHATE			
Inj 1 mg per ml, 1 ml vial – 1% DV Sep-13 to 2016	64.80	5	Hospira
Inj 1 mg per ml, 2 ml vial – 1% DV Sep-13 to 2016	69.60	5	Hospira
VINORELBINE			
Inj 10 mg per ml, 1 ml vial – 1% DV Sep-15 to 2018	8.00	1	Navelbine
Inj 10 mg per ml, 5 ml vial – 1% DV Sep-15 to 2018	40.00	1	Navelbine

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Endocrine Therapy			
ABIRATERONE ACETATE – Restricted see terms below			
☞ Tab 250 mg	4,276.19	120	Zytiga
☞ Restricted			
Initiation			
Medical oncologist, radiation oncologist or urologist			
<i>Re-assessment required after 5 months</i>			
All of the following:			
1 Patient has prostate cancer; and			
2 Patient has metastases; and			
3 Patient's disease is castration resistant; and			
4 Either:			
4.1 All of the following:			
4.1.1 Patient is symptomatic; and			
4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and			
4.1.3 Patient has ECOG performance score of 0-1; and			
4.1.4 Patient has not had prior treatment with taxane chemotherapy; or			
4.2 All of the following:			
4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and			
4.2.2 Patient has ECOG performance score of 0-2; and			
4.2.3 Patient has not had prior treatment with abiraterone.			
Continuation			
Medical oncologist, radiation oncologist or urologist			
<i>Re-assessment required after 5 months</i>			
All of the following:			
1 Significant decrease in serum PSA from baseline; and			
2 No evidence of clinical disease progression; and			
3 No initiation of taxane chemotherapy with abiraterone; and			
4 The treatment remains appropriate and the patient is benefiting from treatment.			
BICALUTAMIDE			
Tab 50 mg – 1% DV Sep-14 to 2017	4.90	28	Bicalaccord
FLUTAMIDE			
Tab 250 mg	55.00	100	Flutamin
MEGESTROL ACETATE			
Tab 160 mg – 1% DV Oct-15 to 2018	54.30	30	Apo-Megestrol
OCTREOTIDE – Some items restricted see terms on the next page			
Inj 50 mcg per ml, 1 ml ampoule – 1% DV Sep-14 to 2017	13.50	5	DBL
Inj 100 mcg per ml, 1 ml ampoule – 1% DV Sep-14 to 2017	22.40	5	DBL
Inj 500 mcg per ml, 1 ml ampoule – 1% DV Sep-14 to 2017	89.40	5	DBL
☞ Inj 10 mg vial	1,772.50	1	Sandostatin LAR
☞ Inj 20 mg vial	2,358.75	1	Sandostatin LAR
☞ Inj 30 mg vial	2,951.25	1	Sandostatin LAR

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔ Restricted			
Initiation — Malignant bowel obstruction			
All of the following:			
1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and			
2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and			
3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.			
Note: Indications marked with * are Unapproved Indications			
Initiation — acromegaly			
<i>Re-assessment required after 3 months</i>			
Both:			
1 The patient has acromegaly; and			
2 Any of the following:			
2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or			
2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or			
2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.			
Continuation — acromegaly			
Both:			
1 IGF1 levels have decreased since starting octreotide; and			
2 The treatment remains appropriate and the patient is benefiting from treatment.			
Note: In patients with acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks.			
Initiation — Other indications			
Any of the following:			
1 VIPomas and glucagonomas - for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or			
2 Both:			
2.1 Gastrinoma; and			
2.2 Either:			
2.2.1 Patient has failed surgery; or			
2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or			
3 Both:			
3.1 Insulinomas; and			
3.2 Surgery is contraindicated or has failed; or			
4 For pre-operative control of hypoglycaemia and for maintenance therapy; or			
5 Both:			
5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and			
5.2 Disabling symptoms not controlled by maximal medical therapy.			
Note: restriction applies only to the long-acting formulations of octreotide			
TAMOXIFEN CITRATE			
Tab 10 mg	17.50	100	Genox
Tab 20 mg	2.63	30	Genox
	8.75	100	Genox
Aromatase Inhibitors			
ANASTROZOLE			
Tab 1 mg	26.55	30	Aremed DP-Anastrozole

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
EXEMESTANE			
Tab 25 mg – 1% DV Sep-14 to 2017	14.50	30	Aromasin
LETROZOLE			
Tab 2.5 mg – 1% DV Jan-16 to 2018	2.95	30	Letrole

Immunosuppressants

Calcineurin Inhibitors

CICLOSPORIN			
Cap 25 mg	44.63	50	Neoral
Cap 50 mg	88.91	50	Neoral
Cap 100 mg	177.81	50	Neoral
Oral liq 100 mg per ml	198.13	50 ml	Neoral
Inj 50 mg per ml, 5 ml ampoule – 1% DV Sep-15 to 2018	276.30	10	Sandimmun
TACROLIMUS – Restricted see terms below			
⚡ Cap 0.5 mg – 1% DV Nov-14 to 31 Oct 2018	85.60	100	Tacrolimus Sandoz
⚡ Cap 1 mg – 1% DV Nov-14 to 31 Oct 2018	171.20	100	Tacrolimus Sandoz
⚡ Cap 5 mg – 1% DV Nov-14 to 31 Oct 2018	428.00	50	Tacrolimus Sandoz
⚡ Inj 5 mg per ml, 1 ml ampoule			

➡ Restricted

Initiation — organ transplant recipients

Any specialist

For use in organ transplant recipients.

Initiation — Steroid-resistant nephrotic syndrome*

Any specialist

Either:

- The patient is a child with steroid-resistant nephrotic syndrome* (SRNS) where ciclosporin has been trialled in combination with prednisone and discontinued because of unacceptable side effects or inadequate clinical response; or
- All of the following:
 - The patient is an adult with SRNS; and
 - Ciclosporin has been trialled in combination with prednisone and discontinued because of unacceptable side effects or inadequate clinical response; and
 - Cyclophosphamide or mycophenolate have been trialled and discontinued because of unacceptable side effects or inadequate clinical response, or these treatments are contraindicated.

Note: Indications marked with * are Unapproved Indications

Fusion Proteins

ETANERCEPT – Restricted see terms below			
⚡ Inj 25 mg vial	799.96	4	Enbrel
⚡ Inj 50 mg autoinjector	1,599.96	4	Enbrel
⚡ Inj 50 mg syringe	1,599.96	4	Enbrel

➡ Restricted

Initiation — juvenile idiopathic arthritis

Rheumatologist or named specialist

Re-assessment required after 4 months

Either:

- Both:
 - The patient has had an initial Special Authority approval for adalimumab for juvenile idiopathic arthritis (JIA); and
continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for JIA; or
- 2 All of the following:
 - 2.1 Patient diagnosed with Juvenile Idiopathic Arthritis (JIA); and
 - 2.2 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
 - 2.5 Both:
 - 2.5.1 Either:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

Continuation — juvenile idiopathic arthritis

Rheumatologist or named specialist

Re-assessment required after 6 months

- Both:
- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation — rheumatoid arthritis

Rheumatologist

Re-assessment required after 6 months

- Either:
- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis; or
 - 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulphasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
- 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation — rheumatoid arthritis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initiation — ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of an exercise regime supervised by a physiotherapist; and
- 2.5 Either:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or

2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and

2.6 Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of starting treatment.

Average normal chest expansion corrected for age and gender:

Age	Male	Female
18-24	7.0 cm	5.5 cm
25-34	7.5 cm	5.5 cm
35-44	6.5 cm	4.5 cm
45-54	6.0 cm	5.0 cm
55-64	5.5 cm	4.0 cm
65-74	4.0 cm	4.0 cm
75+	3.0 cm	2.5 cm

Continuation — ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- Following 12 weeks of etanercept treatment, BASDAI has improved by 4 or more points from pre-treatment baseline on a 10 point scale, or by 50%, whichever is less; and
- Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initiation — psoriatic arthritis

Rheumatologist

Re-assessment required after 6 months

Either:

- Both:
 - The patient has had an initial Special Authority approval for adalimumab for psoriatic arthritis; and
 - Either:
 - The patient has experienced intolerable side effects from adalimumab; or
 - The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for psoriatic arthritis; or
- All of the following:
 - Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - Patient has tried and not responded to at least three months of sulphasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - Either:
 - Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- Any of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
- 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation — psoriatic arthritis

Rheumatologist

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initiation — plaque psoriasis, prior TNF use

Dermatologist

Limited to 4 months treatment

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; and
- 3 Patient must be reassessed for continuation after 3 doses.

Initiation — plaque psoriasis, treatment-naïve

Dermatologist

Limited to 4 months treatment

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Continuation — plaque psoriasis

Dermatologist

Re-assessment required after 6 months

- Both:
- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-etanercept treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-etanercept treatment baseline value; and
 - 2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initiation — pyoderma gangrenosum

Dermatologist

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 4 doses.

Note: Indications marked with * are Unapproved Indications (refer to Section A: General Rules, Part I (Interpretations and Definitions) and Part V (Miscellaneous Provisions) rule 5.5).

Continuation — pyoderma gangrenosum

Dermatologist

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 4 doses.

Initiation — adult-onset Still's disease

Rheumatologist

Re-assessment required after 6 months

- Either:
- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the Section H rules; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Continuation — adult-onset Still's disease

Rheumatologist

Re-assessment required after 6 months

The patient has a sustained improvement in inflammatory markers and functional status.

Monoclonal Antibodies

ABCIXIMAB – **Restricted** see terms below

⚡ Inj 2 mg per ml, 5 ml vial	579.53	1	ReoPro
------------------------------------	--------	---	--------

➡**Restricted**

Initiation

Either:

- 1 For use in patients with acute coronary syndromes undergoing percutaneous coronary intervention; or
- 2 For use in patients undergoing intra-cranial intervention.

ADALIMUMAB – **Restricted** see terms below

⚡ Inj 10 mg per 0.2 ml prefilled syringe	1,599.96	2	Humira
⚡ Inj 20 mg per 0.4 ml syringe	1,599.96	2	Humira
⚡ Inj 40 mg per 0.8 ml pen	1,599.96	2	HumiraPen
⚡ Inj 40 mg per 0.8 ml syringe	1,599.96	2	Humira

➡**Restricted**

Initiation — juvenile idiopathic arthritis

Rheumatologist or named specialist

Re-assessment required after 4 months

Either:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for juvenile idiopathic arthritis (JIA); and
 - 1.1.2 Either:
 - 1.1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for JIA; or
 - 2 All of the following:
 - 2.1 Patient diagnosed with Juvenile Idiopathic Arthritis (JIA); and
 - 2.2 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and

2.5 Both:

2.5.1 Either:

2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or

2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and

2.5.2 Physician's global assessment indicating severe disease.

Continuation — juvenile idiopathic arthritis

Rheumatologist or named specialist

Re-assessment required after 6 months

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation — fistulising Crohn's disease

Gastroenterologist

Re-assessment required after 4 months

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment (a copy of which is available at www.pharmac.govt.nz/latest/BaselineFistulaAssessment.pdf) has been completed and is no more than 1 month old at the time of application.

Continuation — fistulising Crohn's disease

Gastroenterologist

Re-assessment required after 6 months

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initiation — Crohn's disease

Gastroenterologist

Re-assessment required after 3 months

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Continuation — Crohn's disease

Gastroenterologist

Re-assessment required after 3 months

Both:

- 1 Either:
 - 1.1 Either:
 - 1.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 1.1.2 CDAI score is 150 or less; or
 - 1.2 Both:
 - 1.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 1.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation — rheumatoid arthritis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulphasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.7 Either:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation — rheumatoid arthritis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation — ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of an exercise regime supervised by a physiotherapist; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of starting treatment.

Average normal chest expansion corrected for age and gender:

Age	Male	Female
18-24	7.0 cm	5.5 cm
25-34	7.5 cm	5.5 cm
35-44	6.5 cm	4.5 cm
45-54	6.0 cm	5.0 cm
55-64	5.5 cm	4.0 cm
65-74	4.0 cm	4.0 cm
75+	3.0 cm	2.5 cm

Continuation — ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Following 12 weeks of adalimumab treatment, BASDAI has improved by 4 or more points from pre-treatment baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation — psoriatic arthritis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulphasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Continuation — psoriatic arthritis

Rheumatologist

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation — plaque psoriasis, prior TNF use

Dermatologist

Limited to 4 months treatment

Both:

- 1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from etanercept; or
 - 2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis.

Initiation — plaque psoriasis, treatment-naïve

Dermatologist

Limited to 4 months treatment

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Continuation — plaque psoriasis

Dermatologist

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.1.2 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
- 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-etanercept treatment baseline value; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation — pyoderma gangrenosum

Dermatologist

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 4 doses.

Note: Indications marked with * are Unapproved Indications (refer to Section A: General Rules, Part I (Interpretations and Definitions) and Part V (Miscellaneous Provisions) rule 5.5).

Continuation — pyoderma gangrenosum

Dermatologist

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 4 doses.

Initiation — adult-onset Still's disease

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the Section H rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Continuation — adult-onset Still's disease

Rheumatologist

Re-assessment required after 6 months

The patient has a sustained improvement in inflammatory markers and functional status.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BASILIXIMAB – Restricted see terms below			
⚡ Inj 20 mg vial	3,200.00	1	Simulect
➡ Restricted			
Initiation			
For use in solid organ transplants.			
BEVACIZUMAB – Restricted see terms below			
⚡ Inj 25 mg per ml, 4 ml vial			
⚡ Inj 25 mg per ml, 16 ml vial			
➡ Restricted			
Initiation			
Either:			
1 Ocular neovascularisation; or			
2 Exudative ocular angiopathy.			
INFLIXIMAB – Restricted see terms below			
⚡ Inj 100 mg – 10% DV Mar-15 to 29 Feb 2020	806.00	1	Remicade
➡ Restricted			
Initiation — Graft vs host disease			
Patient has steroid-refractory acute graft vs. host disease of the gut.			
Initiation — rheumatoid arthritis			
Rheumatologist			
<i>Re-assessment required after 4 months</i>			
All of the following:			
1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and			
2 Either:			
2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or			
2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and			
3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.			
Continuation — rheumatoid arthritis			
Rheumatologist			
<i>Re-assessment required after 6 months</i>			
All of the following:			
1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and			
2 Either:			
2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or			
2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and			
3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.			
Initiation — ankylosing spondylitis			
Rheumatologist			
<i>Re-assessment required after 3 months</i>			
Both:			
1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and			
2 Either:			
2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or			
continued...			

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Continuation — ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initiation — psoriatic arthritis

Rheumatologist

Re-assessment required after 4 months

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for psoriatic arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for psoriatic arthritis.

Continuation — psoriatic arthritis

Rheumatologist

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initiation — severe ocular inflammation

Therapy limited to 3 doses

Both:

- 1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
- 2 Either:
 - 2.1 Patient has failed to achieve control of severe vision-threatening ocular inflammation following high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids; or
 - 2.2 Patient developed new inflammatory symptoms while receiving high dose steroids.

Initiation — chronic ocular inflammation

Therapy limited to 3 doses

Both:

- 1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2 Patient has tried at least two other immunomodulatory agents.

Continuation — ocular inflammation

Both:

- 1 Patient had a good clinical response to initial treatment; and
- 2 Either:
 - 2.1 A withdrawal of infliximab has been trialled and patient has relapsed after trial withdrawal; or
 - 2.2 Patient has Behcet's disease.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Initiation — Pulmonary sarcoidosis

Both:

- 1 Patient has life-threatening pulmonary sarcoidosis that is refractory to other treatments; and
- 2 Treatment is to be prescribed by, or has been recommended by, a physician with expertise in the treatment of pulmonary sarcoidosis.

Initiation — Crohn's disease (adults)

Gastroenterologist

Re-assessment required after 3 months

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Continuation — Crohn's disease (adults)

Gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation — Crohn's disease (children)

Gastroenterologist

Re-assessment required after 3 months

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Continuation — Crohn's disease (children)

Gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Any of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
- 1.2 PCDAI score is 15 or less; or
- 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation — fistulising Crohn's disease

Gastroenterologist

Re-assessment required after 4 months

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Continuation — fistulising Crohn's disease

Gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation — acute severe fulminant ulcerative colitis

Gastroenterologist

Limited to 6 weeks treatment

Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Continuation — severe fulminant ulcerative colitis

Gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation — severe ulcerative colitis

Gastroenterologist

Re-assessment required after 3 months

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is ≥ 4 ; or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is ≥ 65 ; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Continuation — severe ulcerative colitis

Gastroenterologist

Re-assessment required after 6 months

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by ≥ 2 points from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by ≥ 30 points from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation — plaque psoriasis

Dermatologist

Re-assessment required after 3 doses

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, cyclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Continuation — plaque psoriasis

Dermatologist

Re-assessment required after 3 doses

Both:

1 Either:

1.1 Both:

- 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
- 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or

1.2 Both:

- 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and

1.2.2 Either:

- 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
- 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value; and

2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

OMALIZUMAB – **Restricted** see terms below

⚡ Inj 150 mg vial	500.00	1	Xolair
-------------------------	--------	---	--------

➡Restricted

Initiation

Respiratory specialist

Re-assessment required after 6 months

All of the following:

- 1 Patient is over the age of 6; and
- 2 Patient has a diagnosis of severe, life threatening asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven compliance with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1600 micrograms per day or fluticasone propionate 1000 micrograms per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 micrograms bd or formoterol 12 micrograms bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; and
- 7 At least four admissions to hospital for a severe asthma exacerbation over the previous 24 months with at least one of those being in the previous 12 months; and
- 8 An Asthma Control Questionnaire (ACQ-5) score of at least 3.0 as assessed in the previous month.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
Continuation			
Respiratory specialist			
<i>Re-assessment required after 6 months</i>			
All of the following:			
1 Hospital admissions have been reduced as a result of treatment; and			
2 A reduction in the Asthma Control Questionnaire (ACQ-5) score of at least 1.0 from baseline; and			
3 A reduction in the maintenance oral corticosteroid dose of at least 50% from baseline.			
RANIBIZUMAB – Restricted see terms below			
¶ Inj 10 mg per ml, 0.23 ml vial			
¶ Inj 10 mg per ml, 0.3 ml vial			
➔ Restricted			
Initiation			
<i>Re-assessment required after 3 doses</i>			
Both:			
1 Either:			
1.1 Age-related macular degeneration; or			
1.2 Choroidal neovascular membrane; and			
2 Any of the following:			
2.1 The patient has had a severe ophthalmic inflammatory response following bevacizumab; or			
2.2 The patient has had a myocardial infarction or stroke within the last three months; or			
2.3 The patient has failed to respond to bevacizumab following three intraocular injections; or			
2.4 The patient is of child-bearing potential and has not completed a family.			
Continuation			
Both:			
1 Documented benefit after three doses must be demonstrated to continue; and			
2 In the case of but previous non-response to bevacizumab, a retreat of bevacizumab is required to confirm non-response before continuing with ranibizumab.			
RITUXIMAB – Restricted see terms below			
¶ Inj 10 mg per ml, 10 ml vial	1,075.50	2	Mabthera
¶ Inj 10 mg per ml, 50 ml vial	2,688.30	1	Mabthera
➔ Restricted			
Initiation — haemophilia with inhibitors			
Haematologist			
Any of the following:			
1 Patient has mild congenital haemophilia complicated by inhibitors; or			
2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or			
3 Patient has acquired haemophilia.			
Continuation — haemophilia with inhibitors			
Haematologist			
All of the following:			
1 Patient was previously treated with rituximab for haemophilia with inhibitors; and			
2 An initial response lasting at least 12 months was demonstrated; and			
3 Patient now requires repeat treatment.			
Initiation — post-transplant			
Both:			
1 The patient has B-cell post-transplant lymphoproliferative disorder*; and			
2 To be used for a maximum of 8 treatment cycles.			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Note: Indications marked with * are Unapproved Indications.

Continuation — post-transplant

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are Unapproved Indications.

Initiation — indolent, low-grade lymphomas

Either:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia.

Continuation — indolent, low-grade lymphomas

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia.

Initiation — aggressive CD20 positive NHL

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia.

Continuation — aggressive CD20 positive NHL

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia.

Initiation — Chronic lymphocytic leukaemia

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 The patient is rituximab treatment naive; and
- 3 Either:
 - 3.1 The patient is chemotherapy treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

3.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; and

- 4 The patient has good performance status; and
- 5 The patient has good renal function (creatinine clearance \geq 30 ml/min); and
- 6 The patient does not have chromosome 17p deletion CLL; and
- 7 Rituximab to be administered in combination with fludarabine and cyclophosphamide for a maximum of 6 treatment cycles; and
- 8 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration).

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to <2.

Initiation — rheumatoid arthritis - prior TNF inhibitor use

Rheumatologist

Limited to 4 months treatment

All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initiation — rheumatoid arthritis - TNF inhibitors contraindicated

Rheumatologist

Limited to 4 months treatment

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulphasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of cyclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Continuation — rheumatoid arthritis - re-treatment in 'partial responders' to rituximab

Rheumatologist

Re-assessment required after 4 months

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Continuation — rheumatoid arthritis - re-treatment in 'responders' to rituximab

Rheumatologist

Re-assessment required after 4 months

All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initiation — severe cold haemagglutinin disease (CHAD)

Haematologist

Re-assessment required after 4 weeks

Both:

- 1 Patient has cold haemagglutinin disease*; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms.

Note: Indications marked with * are Unapproved Indications.

Continuation — severe cold haemagglutinin disease (CHAD)

Haematologist

Re-assessment required after 4 weeks

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are Unapproved Indications.

Initiation — warm autoimmune haemolytic anaemia (warm AIHA)

Haematologist

Re-assessment required after 4 weeks

Both:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to >5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin.

Note: Indications marked with * are Unapproved Indications.

Continuation — warm autoimmune haemolytic anaemia (warm AIHA)

Haematologist

Re-assessment required after 4 weeks

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are Unapproved Indications.

Initiation — immune thrombocytopenic purpura (ITP)

Haematologist

Re-assessment required after 4 weeks

Both:

- 1 Either:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of ≤ 20,000 platelets per microlitre; or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy).

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Note: Indications marked with * are Unapproved Indications.

Continuation — immune thrombocytopenic purpura (ITP)

Haematologist

Re-assessment required after 4 weeks

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are Unapproved Indications.

Initiation — thrombotic thrombocytopenic purpura (TTP)

Haematologist

Re-assessment required after 4 weeks

Either:

- 1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
- 2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are Unapproved Indications.

Continuation — thrombotic thrombocytopenic purpura (TTP)

Haematologist

Re-assessment required after 4 weeks

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Note: Indications marked with * are Unapproved Indications.

Initiation — pure red cell aplasia (PRCA)

Haematologist

Re-assessment required after 6 weeks

Patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are Unapproved Indications.

Continuation — pure red cell aplasia (PRCA)

Haematologist

Re-assessment required after 6 weeks

Patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are Unapproved Indications.

Initiation — ANCA associated vasculitis

Re-assessment required after 4 weeks

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Either:
 - 2.1 Patient does not have MPO-ANCA positive vasculitis*; or
 - 2.2 Mycophenolate mofetil has not been effective in those patients who have MPO-ANCA positive vasculitis*; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 4 Any of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 4.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve complete absence of disease after at least 3 months; or
- 4.2 Patient has previously had a cumulative dose of cyclophosphamide >15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose >15 g; or
- 4.3 Cyclophosphamide and methotrexate are contraindicated; or
- 4.4 Patient is a female of child-bearing potential; or
- 4.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are Unapproved Indications.

Continuation — ANCA associated vasculitis

Re-assessment required after 4 weeks

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are Unapproved Indications.

Initiation — treatment refractory systemic lupus erythematosus (SLE)

Rheumatologist or nephrologist

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are Unapproved Indications.

Continuation — treatment refractory systemic lupus erythematosus (SLE)

Rheumatologist or nephrologist

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are Unapproved Indications.

Initiation — Antibody-mediated renal transplant rejection

Nephrologist

Patient has been diagnosed with antibody-mediated renal transplant rejection*.

Note: Indications marked with * are Unapproved Indications.

Initiation — ABO-incompatible renal transplant

Nephrologist

Patient is to undergo an ABO-incompatible renal transplant*.

Note: Indications marked with * are Unapproved Indications.

TOCILIZUMAB – **Restricted** see terms on the next page

⚡ Inj 20 mg per ml, 4 ml vial	220.00	1	Actemra
⚡ Inj 20 mg per ml, 10 ml vial	550.00	1	Actemra
⚡ Inj 20 mg per ml, 20 ml vial	1,100.00	1	Actemra

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➡ Restricted

Initiation — Rheumatoid Arthritis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 All of the following:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
 - 1.3 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
 - 1.4 Either:
 - 1.4.1 The patient has experienced intolerable side effects from rituximab; or
 - 1.4.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis for six months duration or longer; and
 - 2.2 Tocilizumab is to be used as monotherapy; and
 - 2.3 Either:
 - 2.3.1 Treatment with methotrexate is contraindicated; or
 - 2.3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
 - 2.4 Either:
 - 2.4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of cyclosporin alone or in combination with another agent; or
 - 2.4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
 - 2.5 Either:
 - 2.5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 2.5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.6 Either:
 - 2.6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation — Rheumatoid Arthritis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Initiation — systemic juvenile idiopathic arthritis

Rheumatologist

Re-assessment required after 6 months

- Both:
- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
 - 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Continuation — systemic juvenile idiopathic arthritis

Rheumatologist

Re-assessment required after 6 months

- Either:
- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
 - 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Initiation — adult-onset Still's disease

Rheumatologist

Re-assessment required after 6 months

- Either:
- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
 - 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Continuation — adult-onset Still's disease

Rheumatologist

Re-assessment required after 6 months

The patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB – Restricted see terms below

⚡ Inj 150 mg vial	1,350.00	1	Herceptin
⚡ Inj 440 mg vial	3,875.00	1	Herceptin

➡Restricted

Initiation — Early breast cancer

Limited to 12 months treatment

- All of the following:
- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
 - 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
- 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
- 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
- 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Initiation — metastatic breast cancer (trastuzumab-naïve patients)

Limited to 12 months treatment

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient has not previously received lapatinib treatment for HER 2 positive metastatic breast cancer; and
 - 1.3 Trastuzumab not to be given in combination with lapatinib; and
 - 1.4 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2.2 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on lapatinib; and
 - 2.4 Trastuzumab not to be given in combination with lapatinib; and
 - 2.5 Trastuzumab to be discontinued at disease progression.

Initiation — metastatic breast cancer (patients previously treated with trastuzumab)

Limited to 12 months treatment

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 All of the following:
 - 3.1.1 The patient has not previously received lapatinib treatment for metastatic breast cancer; and
 - 3.1.2 Trastuzumab not to be given in combination with lapatinib; and
 - 3.1.3 Trastuzumab to be discontinued at disease progression; or
 - 3.2 All of the following:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; and
 - 3.2.3 Trastuzumab not to be given in combination with lapatinib; and
 - 3.2.4 Trastuzumab to be discontinued at disease progression; or
 - 3.3 All of the following:
 - 3.3.1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
 - 3.3.2 Trastuzumab not to be given in combination with lapatinib; and
 - 3.3.3 Trastuzumab to be discontinued at disease progression.

Continuation — metastatic breast cancer

Re-assessment required after 12 months

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Other Immunosuppressants			
ANTITHYMOCYTE GLOBULIN (EQUINE)			
Inj 50 mg per ml, 5 ml ampoule	2,351.25	5	ATGAM
ANTITHYMOCYTE GLOBULIN (RABBIT)			
Inj 25 mg vial			
AZATHIOPRINE			
Tab 25 mg	8.28	60	Azamun
Tab 50 mg – 1% DV Jun-14 to 2016	13.22	100	Azamun
Inj 50 mg vial	126.00	1	Imuran
BACILLUS CALMETTE-GUERIN (BCG) – Restricted see terms below			
⚡ Inj 2-8 × 10 ⁸ CFU vial – 1% DV Sep-13 to 2016	149.37	1	OncotICE
⚡ Inj 40 mg per ml, vial	149.37	3	SII-Onco-BCG
→Restricted			
Initiation			
For use in bladder cancer.			
EVEROLIMUS – Restricted see terms below			
⚡ Tab 5 mg	4,555.76	30	Afinitor
⚡ Tab 10 mg	6,512.29	30	Afinitor
→Restricted			
Initiation			
Neurologist or oncologist			
<i>Re-assessment required after 3 months</i>			
Both:			
1 Patient has tuberous sclerosis; and			
2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.			
Continuation			
Neurologist or oncologist			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and			
2 The treatment remains appropriate and the patient is benefiting from treatment; and			
3 Everolimus to be discontinued at progression of SEGAs.			
Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.			
MYCOPHENOLATE MOFETIL			
Tab 500 mg – 1% DV Nov-13 to 2016	25.00	50	CellCept
Cap 250 mg – 1% DV Nov-13 to 2016	25.00	100	CellCept
Powder for oral liq 1 g per 5 ml – 1% DV Nov-13 to 2016	187.25	165 ml	CellCept
Inj 500 mg vial – 1% DV Nov-13 to 2016	133.33	4	CellCept
PICIBANIL			
Inj 100 mg vial			
SIROLIMUS – Restricted see terms on the next page			
⚡ Tab 1 mg	813.00	100	Rapamune
⚡ Tab 2 mg	1,626.00	100	Rapamune
⚡ Oral liq 1 mg per ml	487.80	60 ml	Rapamune

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➡Restricted

Initiation

For rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR < 30 ml/min; or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- HUS or TTP; or
- Leukoencephalopathy; or
- Significant malignant disease

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Antiallergy Preparations

Allergic Emergencies

ICATIBANT – **Restricted** see terms below

⚡ Inj 10 mg per ml, 3 ml prefilled syringe2,668.00 1 Firazyr

➡ **Restricted**

Initiation

Clinical immunologist or relevant specialist

Re-assessment required after 12 months

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Continuation

Re-assessment required after 12 months

The treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

BEE VENOM – **Restricted** see terms below

⚡ Maintenance kit - 6 vials 120 mcg freeze dried venom, with diluent

⚡ Inj 550 mcg vial with diluent

➡ **Restricted**

Initiation

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

PAPER WASP VENOM – **Restricted** see terms below

⚡ Treatment kit - 6 vials 120 mcg freeze dried venom, with diluent

⚡ Inj 550 mcg vial with diluent

➡ **Restricted**

Initiation

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

YELLOW JACKET WASP VENOM – **Restricted** see terms below

⚡ Treatment kit - 6 vials 120 mcg freeze dried venom, with diluent

⚡ Inj 550 mcg vial with diluent

➡ **Restricted**

Initiation

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Allergy Prophylactics

BECLOMETHASONE DIPROPIONATE

Nasal spray 50 mcg per dose4.85 200 dose Alanase

Nasal spray 100 mcg per dose5.75 200 dose Alanase

RESPIRATORY SYSTEM AND ALLERGIES

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BUDESONIDE			
Nasal spray 50 mcg per dose	4.85	200 dose	Butacort Aqueous
Nasal spray 100 mcg per dose	5.75	200 dose	Butacort Aqueous
FLUTICASONE PROPIONATE			
Nasal spray 50 mcg per dose – 1% DV Sep-15 to 2018	2.18	120 dose	Flixonase Hayfever & Allergy
IPRATROPIUM BROMIDE			
Aqueous nasal spray 0.03% – 1% DV Jan-15 to 2017	3.95	15 ml	Univent
SODIUM CROMOGLYCATE			
Nasal spray 4%			

Antihistamines

CETIRIZINE HYDROCHLORIDE			
Tab 10 mg	1.59	100	Zetop
Oral liq 1 mg per ml – 1% DV Feb-15 to 2017	2.99	200 ml	Histaclear
CHLORPHENIRAMINE MALEATE			
Oral liq 0.4 mg per ml			
Inj 10 mg per ml, 1 ml ampoule			
CYPROHEPTADINE HYDROCHLORIDE			
Tab 4 mg			
FEXOFENADINE HYDROCHLORIDE			
Tab 60 mg			
Tab 120 mg			
Tab 180 mg			
LORATADINE			
Tab 10 mg – 1% DV Dec-13 to 2016	1.30	100	Lorafix
Oral liq 1 mg per ml – 1% DV Nov-14 to 2016	4.25	200 ml	LoraPaed
PROMETHAZINE HYDROCHLORIDE			
Tab 10 mg – 1% DV Sep-15 to 2018	1.78	50	Allersoothe
Tab 25 mg – 1% DV Sep-15 to 2018	1.99	50	Allersoothe
Oral liq 1 mg per ml – 1% DV Sep-15 to 2018	2.59	100 ml	Allersoothe
Inj 25 mg per ml, 2 ml ampoule	11.99	5	Hospira
TRIMEPRAZINE TARTRATE			
Oral liq 6 mg per ml			

Anticholinergic Agents

IPRATROPIUM BROMIDE			
Aerosol inhaler 20 mcg per dose			
Nebuliser soln 250 mcg per ml, 1 ml ampoule – 1% DV Sep-13 to 2016	3.26	20	Univent
Nebuliser soln 250 mcg per ml, 2 ml ampoule – 1% DV Sep-13 to 2016	3.37	20	Univent

Anticholinergic Agents with Beta-Adrenoceptor Agonists

SALBUTAMOL WITH IPRATROPIUM BROMIDE			
Aerosol inhaler 100 mcg with ipratropium bromide 20 mcg per dose			
Nebuliser soln 2.5 mg with ipratropium bromide 0.5 mg per 2.5 ml ampoule – 1% DV Sep-15 to 2018	3.59	20	Duolin

Price (ex man. excl. GST)	Brand or Generic Manufacturer
\$	Per

Long-Acting Muscarinic Agents

GLYCOPYRRONIUM

Note: inhaled glycopyrronium treatment must not be used if the patient is also receiving treatment with subsidised tiotropium or umeclidinium.

Powder for inhalation 50 mcg per dose61.00 30 dose Seebri Breezhaler

TIOTROPIUM BROMIDE – **Restricted** see terms below

Note: tiotropium treatment must not be used if the patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.

⬇ Soln for inhalation 2.5 mcg per dose70.00 60 dose Spiriva Respimat

⬇ Powder for inhalation 18 mcg per dose70.00 30 dose Spiriva

➡ **Restricted**

Initiation

All of the following:

- To be used for the long-term maintenance treatment of bronchospasm and dyspnoea associated with COPD; and
- In addition to standard treatment, the patient has trialled a short acting bronchodilator dose of at least 40 µg ipratropium q.i.d for one month; and
- Either:
 - the patient's breathlessness according to the Medical Research Council (UK) dyspnoea scale is:
 - Grade 4 (stops for breath after walking about 100 meters or after a few minutes on the level); or
 - Grade 5 (too breathless to leave the house, or breathless when dressing or undressing); and
- Actual FEV₁ as a % of predicted, must be below 60%; and
- Either:
 - Patient is not a smoker (for reporting purposes only); or
 - Patient is a smoker and has been offered smoking cessation counselling; and
- The patient has been offered annual influenza immunization.

UMECLIDINIUM

Note: Umeclidinium must not be used if the patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.

Powder for inhalation 62.5 mcg per dose61.50 30 dose Incruse Ellipta

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

➡ **Restricted**

Initiation

Re-assessment required after 2 years

Both:

- Patient has been stabilised on a long acting muscarinic antagonist; and
- The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Continuation

Re-assessment required after 2 years

Both:

- Patient is compliant with the medication; and
- Patient has experienced improved COPD symptom control (prescriber determined).

Note: Combination long acting muscarinic antagonist and long acting beta-2 agonist must not be used if the patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

GLYCOPYRRONIUM WITH INDACATEROL – **Restricted** see terms above

⬆ Powder for Inhalation 50 mcg with indacaterol 110 mcg81.00 30 dose Ultibro Breezhaler

TIOTROPIUM BROMIDE WITH OLODATEROL – **Restricted** see terms above

⬆ Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg81.00 60 dose Spiolto Respimat

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
UMECLIDINIUM WITH VILANTEROL – Restricted see terms on the preceding page			
† Powder for inhalation 62.5 mcg with vilanterol 25 mcg	77.00	30 dose	Anoro Ellipta

Beta-Adrenoceptor Agonists

SALBUTAMOL

Oral liq 400 mcg per ml – 1% DV Jan-14 to 2016	2.06	150 ml	Ventolin
Inj 500 mcg per ml, 1 ml ampoule			
Inj 1 mg per ml, 5 ml ampoule			
Aerosol inhaler, 100 mcg per dose	3.80	200 dose	SalAir
	4.00		Salamol
	6.00		Ventolin
Nebuliser soln 1 mg per ml, 2.5 ml ampoule – 1% DV Sep-15 to 2018	3.19	20	Asthalin
Nebuliser soln 2 mg per ml, 2.5 ml ampoule – 1% DV Sep-15 to 2018	3.29	20	Asthalin

TERBUTALINE SULPHATE

Powder for inhalation 250 mcg per dose			
Inj 0.5 mg per ml, 1 ml ampoule			

Cough Suppressants

PHOLCODINE

Oral liq 1 mg per ml			
----------------------	--	--	--

Decongestants

OXYMETAZOLINE HYDROCHLORIDE

Aqueous nasal spray 0.25 mg per ml			
Aqueous nasal spray 0.5 mg per ml			

PSEUDOEPHEDRINE HYDROCHLORIDE

Tab 60 mg			
-----------	--	--	--

SODIUM CHLORIDE

Aqueous nasal spray isotonic			
------------------------------	--	--	--

SODIUM CHLORIDE WITH SODIUM BICARBONATE

Soln for nasal irrigation			
---------------------------	--	--	--

XYLOMETAZOLINE HYDROCHLORIDE

Aqueous nasal spray 0.05%			
Aqueous nasal spray 0.1%			
Nasal drops 0.05%			
Nasal drops 0.1%			

Inhaled Corticosteroids

BECLOMETHASONE DIPROPIONATE

Aerosol inhaler 50 mcg per dose	8.54	200 dose	Beclazone 50
	9.30		Qvar
Aerosol inhaler 100 mcg per dose	12.50	200 dose	Beclazone 100
	15.50		Qvar
Aerosol inhaler 250 mcg per dose	22.67	200 dose	Beclazone 250

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BUDESONIDE			
Nebuliser soln 250 mcg per ml, 2 ml ampoule			
Nebuliser soln 500 mcg per ml, 2 ml ampoule			
Powder for inhalation 100 mcg per dose			
Powder for inhalation 200 mcg per dose			
Powder for inhalation 400 mcg per dose			
FLUTICASONE			
Aerosol inhaler 50 mcg per dose	7.50	120 dose	Flixotide Floair
Powder for inhalation 50 mcg per dose	8.67	60 dose	Flixotide Accuhaler
Powder for inhalation 100 mcg per dose	13.87	60 dose	Flixotide Accuhaler
Aerosol inhaler 125 mcg per dose	13.60	120 dose	Flixotide Floair
Aerosol inhaler 250 mcg per dose	27.20	120 dose	Flixotide Floair
Powder for inhalation 250 mcg per dose	24.51	60 dose	Flixotide Accuhaler
Leukotriene Receptor Antagonists			
MONTELUKAST – Restricted see terms below			
⚡ Tab 4 mg	18.48	28	Singulair
⚡ Tab 5 mg	18.48	28	Singulair
⚡ Tab 10 mg	18.48	28	Singulair
➡ Restricted			
Initiation — Pre-school wheeze			
Both:			
1 To be used for the treatment of intermittent severe wheezing (possibly viral) in children under 5 years; and			
2 The patient has had at least three episodes in the previous 12 months of acute wheeze severe enough to seek medical attention.			
Initiation — Exercise-induced asthma			
All of the following:			
1 Patient has been trialed with maximal asthma therapy, including inhaled corticosteroids and long-acting beta-adrenoceptor agonists; and			
2 Patient continues to receive optimal inhaled corticosteroid therapy; and			
3 Patient continues to experience frequent episodes of exercise-induced bronchoconstriction.			
Initiation — Aspirin desensitisation			
Clinical immunologist or allergist			
All of the following:			
1 Patient is undergoing aspirin desensitisation therapy under the supervision of a clinical immunologist or allergist; and			
2 Patient has moderate to severe aspirin-exacerbated respiratory disease or Samter's triad; and			
3 Nasal polyposis, confirmed radiologically or surgically; and			
4 Documented aspirin or NSAID allergy confirmed by aspirin challenge or a clinical history of severe reaction to aspirin or NSAID where challenge would be considered dangerous.			
Long-Acting Beta-Adrenoceptor Agonists			
EFORMOTEROL FUMARATE			
Powder for inhalation 6 mcg per dose			
Powder for inhalation 12 mcg per dose			
INDACATEROL			
Powder for inhalation 150 mcg per dose	61.00	30 dose	Onbrez Breezhaler
Powder for inhalation 300 mcg per dose	61.00	30 dose	Onbrez Breezhaler

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SALMETEROL			
Aerosol inhaler 25 mcg per dose	26.46	120 dose	Meterol
	25.00		Serevent
Powder for inhalation 50 mcg per dose	25.00	60 dose	Serevent Accuhaler

Inhaled Corticosteroids with Long-Acting Beta-Adrenoceptor Agonists

BUDESONIDE WITH EFORMOTEROL

- Powder for inhalation 100 mcg with eformoterol fumarate 6 mcg
- Powder for inhalation 200 mcg with eformoterol fumarate 6 mcg
- Powder for inhalation 400 mcg with eformoterol fumarate 12 mcg
- Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg
- Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg

FLUTICASONE FUROATE WITH VILANTEROL

- Powder for inhalation 100 mcg with vilanterol 25 mcg 44.08 30 dose Breo Ellipta

FLUTICASONE WITH SALMETEROL

- Aerosol inhaler 50 mcg with salmeterol 25 mcg 37.48 120 dose RexAir
-
-
- 33.74 Seretide
- Powder for inhalation 100 mcg with salmeterol 50 mcg 33.74 60 dose Seretide Accuhaler
- Aerosol inhaler 125 mcg with salmeterol 25 mcg 49.69 120 dose RexAir
-
-
- 44.08 Seretide
- Powder for inhalation 250 mcg with salmeterol 50 mcg 44.08 60 dose Seretide Accuhaler

Mast Cell Stabilisers

NEDOCROMIL

- Aerosol inhaler 2 mg per dose

SODIUM CROMOGLYCATE

- Powder for inhalation 20 mg per dose
- Aerosol inhaler 5 mg per dose

Methylxanthines

AMINOPHYLLINE

- Inj 25 mg per ml, 10 ml ampoule – 1% DV Oct-14 to 2017 118.25 5 DBL Aminophylline

CAFFEINE CITRATE

- Oral liq 20 mg per ml (caffeine 10 mg per ml) 14.85 25 ml Biomed
- Inj 20 mg per ml (caffeine 10 mg per ml), 2.5 ml ampoule 55.75 5 Biomed

THEOPHYLLINE

- Tab long-acting 250 mg
- Oral liq 80 mg per 15 ml

Mucolytics and Expectorants

DORNASE ALFA – **Restricted** see terms on the next page

- ☞ Nebuliser soln 2.5 mg per 2.5 ml ampoule 250.00 6 Pulmozyme

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted

Initiation — cystic fibrosis

The patient has cystic fibrosis and has been approved by the Cystic Fibrosis Panel.

Initiation — significant mucus production

Limited to 4 weeks treatment

Both:

- 1 Patient is an in-patient; and
- 2 The mucus production cannot be cleared by first line chest techniques.

Initiation — pleural emphyema

Limited to 3 days treatment

Both:

- 1 Patient is an in-patient; and
- 2 Patient diagnoses with pleural emphyema.

SODIUM CHLORIDE

Nebuliser soln 7%, 90 ml bottle	23.50	90 ml	Biomed
---------------------------------------	-------	-------	--------

Pulmonary Surfactants

BERACTANT

Soln 200 mg per 8 ml vial	550.00	1	Survanta
---------------------------------	--------	---	----------

PORACTANT ALFA

Soln 120 mg per 1.5 ml vial	425.00	1	Curosurf
Soln 240 mg per 3 ml vial	695.00	1	Curosurf

Respiratory Stimulants

DOXAPRAM

Inj 20 mg per ml, 5 ml vial

Sclerosing Agents

TALC

Powder

Soln (slurry) 100 mg per ml, 50 ml

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Anti-Infective Preparations			
Antibacterials			
CHLORAMPHENICOL			
Eye oint 1%	3.19	4 g	Chlorsig
Ear drops 0.5%			
Eye drops 0.5% – 1% DV Sep-15 to 2018	0.98	10 ml	Chlorafast
Eye drops 0.5%, single dose			
CIPROFLOXACIN			
Eye drops 0.3%			
FRAMYCETIN SULPHATE			
Ear/eye drops 0.5%			
FUSIDIC ACID			
Eye drops 1%	4.50	5 g	Fucithalmic
GENTAMICIN SULPHATE			
Eye drops 0.3%	11.40	5 ml	Genoptic
PROPAMIDINE ISETHIONATE			
Eye drops 0.1%			
SULPHACETAMIDE SODIUM			
Eye drops 10%			
TOBRAMYCIN			
Eye oint 0.3% – 1% DV Sep-14 to 2017	10.45	3.5 g	Tobrex
Eye drops 0.3% – 1% DV Sep-14 to 2017	11.48	5 ml	Tobrex
Antifungals			
NATAMYCIN			
Eye drops 5%			
Antivirals			
ACICLOVIR			
Eye oint 3%			
GANCICLOVIR			
Eye gel 0.15%			<i>e.g. Virgan</i>
Combination Preparations			
CIPROFLOXACIN WITH HYDROCORTISONE			
Ear drops ciprofloxacin 0.2% with 1% hydrocortisone – 1% DV Mar-15 to 2017	16.30	10 ml	Ciproxin HC Otic
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/eye drops 500 mcg with framycetin sulphate 5 mg and gramicidin 50 mcg per ml			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE			
Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per g – 1% DV Sep-14 to 2017	5.39	3.5 g	Maxitrol
Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml – 1% DV Sep-14 to 2017	4.50	5 ml	Maxitrol
DEXAMETHASONE WITH TOBRAMYCIN			
Eye drops 0.1% with tobramycin 0.3% – 1% DV Mar-15 to 2017	12.64	5 ml	Tobradex
FLUMETASONE PIVALATE WITH CLIOQUINOL			
Ear drops 0.02% with clioquinol 1%			
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCIN AND NYSTATIN			
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml	Kenacomb

Anti-Inflammatory Preparations

Corticosteroids

DEXAMETHASONE			
Eye oint 0.1% – 1% DV Oct-14 to 2017	5.86	3.5 g	Maxidex
Eye drops 0.1% – 1% DV Oct-14 to 2017	4.50	5 ml	Maxidex
FLUOROMETHOLONE			
Eye drops 0.1% – 1% DV Sep-15 to 2018	3.09	5 ml	FML
PREDNISOLONE ACETATE			
Eye drops 0.12%			
Eye drops 1%			
PREDNISOLONE SODIUM PHOSPHATE			
Eye drops 0.5%, single dose (preservative free)	38.50	20 dose	Minims Prednisolone

Non-Steroidal Anti-Inflammatory Drugs

DICLOFENAC SODIUM			
Eye drops 0.1% – 1% DV Sep-14 to 2017	13.80	5 ml	Voltaren Ophtha
KETOROLAC TROMETAMOL			
Eye drops 0.5%			

Decongestants and Antiallergics

Antiallergic Preparations

LEVOCABASTINE			
Eye drops 0.05%			
LODOXAMIDE			
Eye drops 0.1% – 1% DV Sep-14 to 2017	8.71	10 ml	Lomide
OLOPATADINE			
Eye drops 0.1%			
SODIUM CROMOGLYCATE			
Eye drops 2%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Decongestants			
NAPHAZOLINE HYDROCHLORIDE			
Eye drops 0.1% – 1% DV Sep-14 to 2017	4.15	15 ml	Naphcon Forte
Diagnostic and Surgical Preparations			
Diagnostic Dyes			
FLUORESCEIN SODIUM			
Eye drops 2%, single dose			
Inj 10%, 5 ml vial	125.00	12	Fluorescite
Ophthalmic strips 1 mg			
FLUORESCEIN SODIUM WITH LIGNOCAINE HYDROCHLORIDE			
Eye drops 0.25% with lignocaine hydrochloride 4%, single dose			
LISSAMINE GREEN			
Ophthalmic strips 1.5 mg			
ROSE BENGAL SODIUM			
Ophthalmic strips 1%			
Irrigation Solutions			
MIXED SALT SOLUTION FOR EYE IRRIGATION			
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 15 ml dropper bottle – 1% DV Jan-16 to 2018	5.00	15 ml	Balanced Salt Solution
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 250 ml			<i>e.g. Balanced Salt Solution</i>
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 500 ml bottle – 1% DV Jan-16 to 2018	10.50	500 ml	Balanced Salt Solution
Ocular Anaesthetics			
OXYBUPROCAINE HYDROCHLORIDE			
Eye drops 0.4%, single dose			
PROXYMETACAINE HYDROCHLORIDE			
Eye drops 0.5%			
TETRACAINE [AMETHOCAINE] HYDROCHLORIDE			
Eye drops 0.5%, single dose			
Eye drops 1%, single dose			
Viscoelastic Substances			
HYPROMELLOSE			
Inj 2%, 1 ml syringe			
Inj 2%, 2 ml syringe			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SODIUM HYALURONATE [HYALURONIC ACID]			
Inj 14 mg per ml, 0.85 ml syringe	50.00	1	Healon GV
Inj 14 mg per ml, 0.55 ml syringe	50.00	1	Healon GV
Inj 23 mg per ml, 0.6 ml syringe			
Inj 10 mg per ml, 0.85 ml syringe	30.00	1	Provisc
SODIUM HYALURONATE [HYALURONIC ACID] WITH CHONDROITIN SULPHATE			
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.35 ml syringe and inj 10 mg sodium hyaluronate [hyaluronic acid] per ml, 0.4 ml syringe	64.00	1	Duovisc
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.5 ml syringe and inj 10 mg sodium hyaluronate [hyaluronic acid] per ml, 0.55 ml syringe	74.00	1	Duovisc
Inj 30 mg with chondroitin sulphate 40 mg per ml, 0.75 ml syringe			

Other

DISODIUM EDETATE			
Inj 150 mg per ml, 20 ml ampoule			
Inj 150 mg per ml, 20 ml vial			
Inj 150 mg per ml, 100 ml vial			
RIBOFLAVIN 5-PHOSPHATE			
Soln trans epithelial riboflavin			
Inj 0.1%			
Inj 0.1% plus 20% dextran T500			

Glaucoma Preparations

Beta Blockers

BETAXOLOL			
Eye drops 0.25% – 1% DV Sep-14 to 2017	11.80	5 ml	Betoptic S
Eye drops 0.5% – 1% DV Sep-14 to 2017	7.50	5 ml	Betoptic
LEVOBUNOLOL HYDROCHLORIDE			
Eye drops 0.5%	7.00	5 ml	Betagan
TIMOLOL			
Eye drops 0.25% – 1% DV Sep-14 to 2017	1.45	5 ml	Arrow-Timolol
Eye drops 0.25%, gel forming – 1% DV Mar-14 to 2016	3.30	2.5 ml	Timoptol XE
Eye drops 0.5% – 1% DV Sep-14 to 2017	1.45	5 ml	Arrow-Timolol
Eye drops 0.5%, gel forming – 1% DV Mar-14 to 2016	3.78	2.5 ml	Timoptol XE

Carbonic Anhydrase Inhibitors

ACETAZOLAMIDE			
Tab 250 mg – 1% DV Sep-14 to 2017	17.03	100	Diamox
Inj 500 mg			
BRINZOLAMIDE			
Eye drops 1%			
DORZOLAMIDE			
Eye drops 2%			
DORZOLAMIDE WITH TIMOLOL			
Eye drops 2% with timolol 0.5% – 1% DV Dec-15 to 2018	3.45	5 ml	Arrow-Dortim

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Miotics			
ACETYLCHOLINE CHLORIDE			
Inj 20 mg vial with diluent			
PILOCARPINE HYDROCHLORIDE			
Eye drops 1% – 1% DV Sep-14 to 2017	4.26	15 ml	Isopto Carpine
Eye drops 2% – 1% DV Sep-14 to 2017	5.35	15 ml	Isopto Carpine
Eye drops 2%, single dose			
Eye drops 4% – 1% DV Sep-14 to 2017	7.99	15 ml	Isopto Carpine
Prostaglandin Analogues			
BIMATOPROST			
Eye drops 0.03%			
LATANOPROST			
Eye drops 0.005% – 1% DV Sep-15 to 2018	1.50	2.5 ml	Hysite
TRAVOPROST			
Eye drops 0.004%			
Sympathomimetics			
APRACLONIDINE			
Eye drops 0.5% – 1% DV Mar-15 to 2017	19.77	5 ml	Iopidine
BRIMONIDINE TARTRATE			
Eye drops 0.2% – 1% DV Sep-14 to 2017	4.32	5 ml	Arrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL			
Eye drops 0.2% with timolol 0.5%			
Mydriatics and Cycloplegics			
Anticholinergic Agents			
ATROPINE SULPHATE			
Eye drops 0.5%			
Eye drops 1%, single dose			
Eye drops 1% – 1% DV Jul-14 to 2017	17.36	15 ml	Atropt
CYCLOPENTOLATE HYDROCHLORIDE			
Eye drops 0.5%, single dose			
Eye drops 1% – 1% DV Sep-14 to 2017	8.76	15 ml	Cyclogyl
Eye drops 1%, single dose			
TROPICAMIDE			
Eye drops 0.5% – 1% DV Oct-14 to 2017	7.15	15 ml	Mydriacyl
Eye drops 0.5%, single dose			
Eye drops 1% – 1% DV Oct-14 to 2017	8.66	15 ml	Mydriacyl
Eye drops 1%, single dose			
Sympathomimetics			
PHENYLEPHRINE HYDROCHLORIDE			
Eye drops 2.5%, single dose			
Eye drops 10%, single dose			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Ocular Lubricants			
CARBOMER			
Ophthalmic gel 0.3%, single dose	8.25	30	Poly Gel
Ophthalmic gel 0.2%			
CARMELLOSE SODIUM			
Eye drops 0.5%			
Eye drops 0.5%, single dose			
Eye drops 1%			
Eye drops 1%, single dose			
HYPROMELLOSE			
Eye drops 0.5%	3.92	15 ml	Methopt
HYPROMELLOSE WITH DEXTRAN			
Eye drops 0.3% with dextran 0.1%	2.30	15 ml	Poly-Tears
Eye drops 0.3% with dextran 0.1%, single dose			
MACROGOL 400 AND PROPYLENE GLYCOL			
Eye drops 0.4% with propylene glycol 0.3% preservative free, single dose	4.30	24	Systane Unit Dose
PARAFFIN LIQUID WITH SOFT WHITE PARAFFIN			
Eye oint 42.5% with soft white paraffin 57.3%			
PARAFFIN LIQUID WITH WOOL FAT			
Eye oint 3% with wool fat 3% – 1% DV Jul-14 to 2017	3.63	3.5 g	Poly-Visc
POLYVINYL ALCOHOL			
Eye drops 1.4% – 1% DV Jun-16 to 2019	3.62	15 ml	Liquifilm Tears
	2.62		Vistil
Eye drops 3% – 1% DV Jun-16 to 2019	3.88	15 ml	Liquifilm Forte
	3.68		Vistil Forte
<i>(Liquifilm Tears Eye drops 1.4% to be delisted 1 June 2016)</i>			
<i>(Liquifilm Forte Eye drops 3% to be delisted 1 June 2016)</i>			
POLYVINYL ALCOHOL WITH POVIDONE			
Eye drops 1.4% with povidone 0.6%, single dose			
RETINOL PALMITATE			
Oint 138 mcg per g	3.80	5 g	VitA-POS
SODIUM HYALURONATE [HYALURONIC ACID]			
Eye drops 1 mg per ml	22.00	10 ml	Hyo-Fresh

Other Otological Preparations

ACETIC ACID WITH PROPYLENE GLYCOL			
Ear drops 2.3% with propylene glycol 2.8%			
DOCUSATE SODIUM			
Ear drops 0.5%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Agents Used in the Treatment of Poisonings			
Antidotes			
ACETYLCYSTEINE			
Tab eff 200 mg			
Inj 200 mg per ml, 10 ml ampoule – 1% DV Sep-15 to 2018	78.34	10	DBL Acetylcysteine
DIGOXIN IMMUNE FAB			
Inj 38 mg vial			
Inj 40 mg vial			
ETHANOL			
Liq 96%			
ETHANOL WITH GLUCOSE			
Inj 10% with glucose 5%, 500 ml bottle			
ETHANOL, DEHYDRATED			
Inj 100%, 5 ml ampoule			
Inj 96%			
FLUMAZENIL			
Inj 0.1 mg per ml, 5 ml ampoule – 1% DV Sep-15 to 2018	85.05	5	Anexate
HYDROXOCOBALAMIN			
Inj 5 g vial			
Inj 2.5 g vial			
NALOXONE HYDROCHLORIDE			
Inj 400 mcg per ml, 1 ml ampoule	48.84	5	Hospira
PRALIDOXIME IODIDE			
Inj 25 mg per ml, 20 ml ampoule			
SODIUM NITRITE			
Inj 30 mg per ml, 10 ml ampoule			
SODIUM THIOSULFATE			
Inj 500 mg per ml, 20 ml ampoule			
Inj 250 mg per ml, 10 ml vial			
Inj 500 mg per ml, 10 ml vial			
SOYA OIL			
Inj 20%, 500 ml bag			
Inj 20%, 500 ml bottle			
Antitoxins			
BOTULISM ANTITOXIN			
Inj 250 ml vial			
DIPHThERIA ANTITOXIN			
Inj 10,000 iu vial			
Antivenoms			
RED BACK SPIDER ANTIVENOM			
Inj 500 u vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SNAKE ANTIVENOM			
Inj 50 ml vial			

Removal and Elimination

CHARCOAL

Oral liq 200 mg per ml	43.50	250 ml	Carbasorb-X
------------------------------	-------	--------	-------------

DEFERASIROX – **Restricted** see terms below

⚡ Tab 125 mg dispersible	276.00	28	Exjade
⚡ Tab 250 mg dispersible	552.00	28	Exjade
⚡ Tab 500 mg dispersible	1,105.00	28	Exjade

➡ **Restricted**

Initiation

Haematologist

Re-assessment required after 2 years

All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 - 1.0 cells per μL).

Continuation

Haematologist

Re-assessment required after 2 years

Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels. .

DEFERIPRONE – **Restricted** see terms below

⚡ Tab 500 mg	533.17	100	Ferriprox
⚡ Oral liq 100 mg per ml	266.59	250 ml	Ferriprox

➡ **Restricted**

Initiation

Patient has been diagnosed with chronic iron overload due to congenital inherited anaemia or acquired red cell aplasia.

DEFERRIOXAMINE MESILATE

Inj 500 mg vial – 1% DV Feb-16 to 2018	51.52	10	Desferal
---	-------	----	-----------------

DICOBALT EDETATE

Inj 15 mg per ml, 20 ml ampoule

DIMERCAPROL

Inj 50 mg per ml, 2 ml ampoule

DIMERCAPTOSUCCINIC ACID

Cap 100 mg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SODIUM CALCIUM EDETATE			
Inj 200 mg per ml, 2.5 ml ampoule			
Inj 200 mg per ml, 5 ml ampoule			
Antiseptics and Disinfectants			
CHLORHEXIDINE			
Soln 4%	1.86	50 ml	healthE
Soln 5%	15.50	500 ml	healthE
CHLORHEXIDINE WITH CETRIMIDE			
Crm 0.1% with cetrimide 0.5%			
Foaming soln 0.5% with cetrimide 0.5%			
CHLORHEXIDINE WITH ETHANOL			
Soln 0.5% with ethanol 70%, non-staining (pink) 100 ml	2.65	1	healthE
Soln 2% with ethanol 70%, non-staining (pink) 100 ml	3.54	1	healthE
Soln 0.5% with ethanol 70%, non-staining (pink) 25 ml	1.55	1	healthE
Soln 0.5% with ethanol 70%, staining (red) 100 ml	2.90	1	healthE
Soln 2% with ethanol 70%, staining (red) 100 ml	3.86	1	healthE
Soln 0.5% with ethanol 70%, non-staining (pink) 500 ml	5.45	1	healthE
Soln 0.5% with ethanol 70%, staining (red) 500 ml	5.90	1	healthE
Soln 2% with ethanol 70%, staining (red) 500 ml	9.56	1	healthE
IODINE WITH ETHANOL			
Soln 1% with ethanol 70%, 100 ml	9.30	1	healthE
ISOPROPYL ALCOHOL			
Soln 70%, 500 ml	5.00	1	PSM
	5.65		healthE
POVIDONE-IODINE			
☼ Vaginal tab 200 mg			
➡Restricted			
Initiation			
Rectal administration pre-prostate biopsy.			
Oint 10%	3.27	25 g	Betadine
Soln 10%	6.20	500 ml	Betadine
	2.95	100 ml	Riodine
	6.20	500 ml	Riodine
Soln 5%			
Soln 7.5%			
Pad 10%			
Swab set 10%			
POVIDONE-IODINE WITH ETHANOL			
Soln 10% with ethanol 30%	10.00	500 ml	Betadine Skin Prep
Soln 10% with ethanol 70%			
SODIUM HYPOCHLORITE			
Soln			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Contrast Media			
Iodinated X-ray Contrast Media			
DIATRIZOATE MEGLUMINE WITH SODIUM AMIDOTRIZOATE			
Oral liq 660 mg per ml with sodium amidotrizoate 100 mg per ml, 100 ml bottle	22.50	100 ml	Gastrografin
Inj 260 mg with sodium amidotrizoate 40 mg per ml, 250 ml bottle	80.00	1	Urografin
DIATRIZOATE SODIUM			
Oral liq 370 mg per ml, 10 ml sachet	156.12	50	Ioscan
IODISED OIL			
Inj 38% w/w (480 mg per ml), 10 ml ampoule	191.00	1	Lipiodol Ultra Fluid
IODIXANOL			
Inj 270 mg per ml (iodine equivalent), 50 ml bottle – 5% DV Sep-14 to 2017	220.00	10	Visipaque
Inj 270 mg per ml (iodine equivalent), 100 ml bottle – 5% DV Sep-14 to 2017	430.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 50 ml bottle – 5% DV Sep-14 to 2017	220.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 100 ml bottle – 5% DV Sep-14 to 2017	430.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 200 ml bottle – 5% DV Sep-14 to 2017	850.00	10	Visipaque
IOHEXOL			
Inj 240 mg per ml (iodine equivalent), 50 ml bottle – 5% DV Sep-14 to 2017	75.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 20 ml bottle – 5% DV Sep-14 to 2017	57.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 50 ml bottle – 5% DV Sep-14 to 2017	75.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 100 ml bottle – 5% DV Sep-14 to 2017	150.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 20 ml bottle – 5% DV Sep-14 to 2017	59.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 50 ml bottle – 5% DV Sep-14 to 2017	75.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 75 ml bottle – 5% DV Sep-14 to 2017	114.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 100 ml bottle – 5% DV Sep-14 to 2017	150.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 200 ml bottle – 5% DV Sep-14 to 2017	290.00	10	Omnipaque

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Non-iodinated X-ray Contrast Media			
BARIUM SULPHATE			
Powder for oral liq 20 mg per g (2% w/w), 22.1 g sachet	507.50	50	E-Z-Cat Dry
Oral liq 400 mg per ml (40% w/v, 30% w/w), bottle	17.39	148 g	Varibar - Thin Liquid
Oral liq 600 mg per g (60% w/w), tube	36.51	454 g	E-Z-Paste
Oral liq 400 mg per ml (40% w/v), bottle	155.35	250 ml	Varibar - Honey
	38.40	240 ml	Varibar - Nectar
	145.04	230 ml	Varibar - Pudding
Enema 1,250 mg per ml (125% w/v), 500 ml bag	282.30	12	Liquibar
Oral liq 22 mg per g (2.2% w/w), 250 ml bottle	175.00	24	CT Plus+
Oral liq 22 mg per g (2.2% w/w), 450 ml bottle	220.00	24	CT Plus+
Oral liq 1 mg per ml (0.1% w/v, 0.1% w/w), 450 ml bottle	441.12	24	VoLumen
Oral liq 20.9 mg per ml (2.1% w/v, 2% w/w), 250 ml bottle	140.94	24	Readi-CAT 2
Powder for oral soln 97.65% w/w, 300 g bottle	237.76	24	X-Opaque-HD
Oral liq 400 mg per ml (40% w/v, 30% w/w), 20 ml bottle	52.35	3	Tagitol V
Oral liq 1,250 mg per ml (125% w/v), 2,000 ml bottle	91.77	1	Liquibar
BARIUM SULPHATE WITH SODIUM BICARBONATE			
Grans eff 382.2 mg per g with sodium bicarbonate 551.3 mg per g, 4 g sachet	102.93	50	E-Z-Gas II
CITRIC ACID WITH SODIUM BICARBONATE			
Powder 382.2 mg per g with sodium bicarbonate 551.3 mg per g, 4 g sachet			<i>e.g. E-Z-GAS II</i>
Paramagnetic Contrast Media			
GADOBENIC ACID			
Inj 334 mg per ml, 10 ml vial	324.74	10	Multihance
Inj 334 mg per ml, 20 ml vial	636.28	10	Multihance
GADOBUTROL			
Inj 1 mmol per ml, 15 ml vial			
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 7.5 ml prefilled syringe	180.00	5	Gadovist
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 15 ml prefilled syringe	700.00	10	Gadovist
GADODIAMIDE			
Inj 287 mg per ml, 10 ml prefilled syringe	200.00	10	Omniscan
Inj 287 mg per ml, 10 ml vial	170.00	10	Omniscan
Inj 287 mg per ml, 5 ml vial	120.00	10	Omniscan
Inj 287 mg per ml, 15 ml prefilled syringe	320.00	10	Omniscan
GADOTERIC ACID			
Inj 279.32 mg per ml (0.5 mmol per ml), 10 ml prefilled syringe	24.50	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 15 ml bottle	34.50	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 15 ml prefilled syringe	41.00	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 20 ml prefilled syringe	55.00	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 10 ml bottle	23.20	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 20 ml bottle	46.30	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 5 ml bottle	12.30	1	Dotarem

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GADOXETATE DISODIUM			
Inj 181.43 mg per ml (equivalent to 0.25 mmol per ml), 10 ml prefilled syringe	300.00	1	Primovist
MEGLUMINE GADOPENTETATE			
Inj 469 mg per ml, 10 ml prefilled syringe	95.00	5	Magnevist
Inj 469 mg per ml, 10 ml vial	185.00	10	Magnevist
MEGLUMINE IOTROXATE			
Inj 105 mg per ml, 100 ml bottle	150.00	100 ml	Biliscopin
Ultrasound Contrast Media			
PERFLUTREN			
Inj 1.1 mg per ml, 1.5 ml vial – 5% DV Sep-14 to 2017	180.00	1	Definity
	720.00	4	Definity
Diagnostic Agents			
ARGININE			
Inj 50 mg per ml, 500 ml bottle			
Inj 100 mg per ml, 300 ml bottle			
HISTAMINE ACID PHOSPHATE			
Nebuliser soln 0.6%, 10 ml vial			
Nebuliser soln 2.5%, 10 ml vial			
Nebuliser soln 5%, 10 ml vial			
MANNITOL			
Powder for inhalation			<i>e.g. Aridol</i>
METHACHOLINE CHLORIDE			
Powder 100 mg			
SECRETIN PENTAHYDROCHLORIDE			
Inj 100 u ampoule			
SINCALIDE			
Inj 5 mcg per vial			
TUBERCULIN, PURIFIED PROTEIN DERIVATIVE			
Inj 5 TU per 0.1 ml, 1 ml vial			
Diagnostic Dyes			
BONNEY'S BLUE DYE			
Soln			
INDIGO CARMINE			
Inj 4 mg per ml, 5 ml ampoule			
Inj 8 mg per ml, 5 ml ampoule			
INDOCYANINE GREEN			
Inj 25 mg vial			
METHYLTHIONINIUM CHLORIDE [METHYLENE BLUE]			
Inj 10 mg per ml, 10 ml ampoule			
Inj 10 mg per ml, 5 ml ampoule			
PATENT BLUE V			
Inj 2.5%, 2 ml ampoule	440.00	5	Obex Medical

Products with Hospital Supply Status (HSS) are in **bold**

Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Irrigation Solutions			
CHLORHEXIDINE			
Irrigation soln 0.02%, bottle	2.92	100 ml	Baxter
Irrigation soln 0.05%, bottle	3.02	100 ml	Baxter
	3.63	500 ml	Baxter
Irrigation soln 0.1%, bottle	3.10	100 ml	Baxter
Irrigation soln 0.5%, bottle	4.69	500 ml	Baxter
Irrigation soln 0.02%, 500 ml bottle			
Irrigation soln 0.1%, 30 ml ampoule			
CHLORHEXIDINE WITH CETRIMIDE			
Irrigation soln 0.015% with cetrimide 0.15%, 30 ml ampoule			
Irrigation soln 0.015% with cetrimide 0.15%, bottle	3.21	100 ml	Baxter
	3.47	500 ml	Baxter
	4.17	1,000 ml	Baxter
Irrigation soln 0.05% with cetrimide 0.5%, bottle	3.87	500 ml	Baxter
	4.20	100 ml	Baxter
Irrigation soln 0.1% with cetrimide 1%, bottle	4.38	100 ml	Baxter
	5.81	500 ml	Baxter
GLYCINE			
Irrigation soln 1.5%, bottle	11.38	2,000 ml	Baxter
	14.44	3,000 ml	Baxter
SODIUM CHLORIDE			
Irrigation soln 0.9%, 30 ml ampoule	19.50	30 ml	Pfizer
Irrigation soln 0.9%, bottle	2.49	100 ml	Baxter
	2.88	500 ml	Baxter
	2.96	1,000 ml	Baxter
	10.00	2,000 ml	Baxter
	12.67	3,000 ml	Baxter
WATER			
Irrigation soln, bottle	2.61	500 ml	Baxter
	2.68	100 ml	Baxter
	2.75	1,000 ml	Baxter
	9.71	2,000 ml	Baxter
	15.80	3,000 ml	Baxter

Surgical Preparations

BISMUTH SUBNITRATE AND IODOFORM PARAFFIN

Paste

DIMETHYL SULFOXIDE

Soln 50%

Soln 99%

PHENOL

Inj 6%, 10 ml ampoule

PHENOL WITH IOXAGLIC ACID

Inj 12%, 10 ml ampoule

TROMETAMOL

Inj 36 mg per ml, 500 ml bottle

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

Cardioplegia Solutions

ELECTROLYTES

Inj 15 mmol/l sodium chloride, 9 mmol/l potassium chloride, 1 mmol/l potassium hydrogen 2-ketoglutarate, 4 mmol/l magnesium chloride, 18 mmol/l histidine hydrochloride, 180 mmol/l histidine, 2 mmol/l tryptophan, 30 mmol/l mannitol, 0.015 mmol/l calcium chloride, 1,000 ml bag

e.g. Custodial-HTK

Inj aspartic acid 10.43 mg per ml, citric acid 0.22476 mg per ml, glutamic acid 11.53 mg per ml, sodium phosphate 0.1725 mg per ml, potassium chloride 2.15211 mg per ml, sodium citrate 1.80768 mg per ml, sodium hydroxide 6.31 mg per ml and trometamol 11.2369 mg per ml, 364 ml bag

*e.g. Cardioplegia
Enriched Paed.
Soln.*

Inj aspartic acid 8.481 mg per ml, citric acid 0.8188 mg per ml, glutamic acid 9.375 mg per ml, sodium phosphate 0.6285 mg per ml, potassium chloride 2.5 mg per ml, sodium citrate 6.585 mg per ml, sodium hydroxide 5.133 mg per ml and trometamol 9.097 mg per ml, 527 ml bag

*e.g. Cardioplegia
Enriched Solution*

Inj citric acid 0.07973 mg per ml, sodium phosphate 0.06119 mg per ml, potassium chloride 2.181 mg per ml, sodium chloride 1.788 mg ml, sodium citrate 0.6412 mg per ml and trometamol 5.9 mg per ml, 523 ml bag

*e.g. Cardioplegia Base
Solution*

Inj 110 mmol/l sodium, 16 mmol/l potassium, 1.2 mmol/l calcium, 16 mmol/l magnesium and 160 mmol/l chloride, 1,000 ml bag

*e.g. Cardioplegia
Solution AHB7832*

Inj 143 mmol/l sodium, 16 mmol/l potassium, 16 mmol/l magnesium and 1.2 mmol/l calcium, 1,000 ml bag

*e.g. Cardioplegia
Electrolyte Solution*

MONOSODIUM GLUTAMATE WITH SODIUM ASPARTATE

Inj 42.68 mg with sodium aspartate 39.48 mg per ml, 250 ml bottle

MONOSODIUM L-ASPARTATE

Inj 14 mmol per 10 ml, 10 ml

Cold Storage Solutions

SODIUM WITH POTASSIUM

Inj 29 mmol/l with potassium 125 mmol/l, 1,000 ml bag

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Extemporaneously Compounded Preparations		
ACETIC ACID Liq		
ALUM Powder BP		
ARACHIS OIL [PEANUT OIL] Liq		
ASCORBIC ACID Powder		
BENZOIN Tincture compound BP		
BISMUTH SUBGALLATE Powder		
BORIC ACID Powder		
CARBOXYMETHYLCELLULOSE Soln 1.5%		
CETRIMIDE Soln 40%		
CHLORHEXIDINE GLUCONATE Soln 20 %		
CHLOROFORM Liq BP		
CITRIC ACID Powder BP		
CLOVE OIL Liq		
COAL TAR Soln BP		
CODEINE PHOSPHATE Powder		
COLLODION FLEXIBLE Liq		
COMPOUND HYDROXYBENZOATE Soln		
CYSTEAMINE HYDROCHLORIDE Powder		
DISODIUM HYDROGEN PHOSPHATE WITH SODIUM DIHYDROGEN PHOSPHATE Inj 37.46 mg with sodium dihydrogen phosphate 47.7 mg in 1.5 ml ampoule		
DITHRANOL Powder		

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GLUCOSE [DEXTROSE] Powder			
GLYCERIN WITH SODIUM SACCHARIN Suspension	32.50	473 ml	Ora-Sweet SF
GLYCERIN WITH SUCROSE Suspension	32.50	473 ml	Ora-Sweet
GLYCEROL Liq	19.80	2,000 ml	ABM
HYDROCORTISONE Powder – 1% DV Dec-14 to 2017	59.50	25 g	ABM
LACTOSE Powder			
MAGNESIUM HYDROXIDE Paste			
MENTHOL Crystals			
METHADONE HYDROCHLORIDE Powder			
METHYL HYDROXYBENZOATE Powder			
METHYLCELLULOSE Powder			
Suspension	32.50	473 ml	Ora-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN Suspension	32.50	473 ml	Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE Suspension	32.50	473 ml	Ora-Blend
OLIVE OIL Liq			
PARAFFIN Liq			
PHENOBARBITONE SODIUM Powder			
PHENOL Liq			
PILOCARPINE NITRATE Powder			
POLYHEXAMETHYLENE BIGUANIDE Liq			
POVIDONE K30 Powder			
PROPYLENE GLYCOL Liq	12.00	500 ml	ABM

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SALICYLIC ACID Powder			
SILVER NITRATE Crystals			
SODIUM BICARBONATE Powder BP			
SODIUM CITRATE Powder			
SODIUM METABISULFITE Powder			
STARCH Powder			
SULPHUR Precipitated Sublimed			
SYRUP Liq (pharmaceutical grade)	21.75	2,000 ml	Midwest
THEOBROMA OIL Oint			
TRI-SODIUM CITRATE Crystals			
TRICHLORACETIC ACID Grans			
UREA Powder BP			
WOOL FAT Oint, anhydrous			
XANTHAN Gum 1%			
ZINC OXIDE Powder			

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Food Modules

Carbohydrate

➔Restricted

Initiation — Use as an additive

Any of the following:

- 1 Cystic fibrosis; or
- 2 Chronic kidney disease; or
- 3 Cancer in children; or
- 4 Cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 5 Faltering growth in an infant/child; or
- 6 Bronchopulmonary dysplasia; or
- 7 Premature and post premature infant; or
- 8 Inborn errors of metabolism.

Initiation — Use as a module

For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

CARBOHYDRATE SUPPLEMENT – **Restricted** see terms above

- ⬆ Powder 95 g carbohydrate per 100 g, 368 g can
- ⬆ Powder 96 g carbohydrate per 100 g, 400 g can

e.g. Polycal

Fat

➔Restricted

Initiation — Use as an additive

Any of the following:

- 1 Patient has inborn errors of metabolism; or
- 2 Faltering growth in an infant/child; or
- 3 Bronchopulmonary dysplasia; or
- 4 Fat malabsorption; or
- 5 Lymphangiectasia; or
- 6 Short bowel syndrome; or
- 7 Infants with necrotising enterocolitis; or
- 8 Biliary atresia; or
- 9 For use in a ketogenic diet; or
- 10 Chyle leak; or
- 11 Ascites; or
- 12 Patient has increased energy requirements, and for whom dietary measures have not been successful.

Initiation — Use as a module

For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk. .

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

LONG-CHAIN TRIGLYCERIDE SUPPLEMENT – **Restricted** see terms above

- ⬆ Liquid 50 g fat per 100 ml, 200 ml bottle
- ⬆ Liquid 50 g fat per 100 ml, 500 ml bottle

e.g. Calogen
e.g. Calogen

MEDIUM-CHAIN TRIGLYCERIDE SUPPLEMENT – **Restricted** see terms above

- ⬆ Liquid 50 g fat per 100 ml, 250 ml bottle
- ⬆ Liquid 95 g fat per 100 ml, 500 ml bottle

e.g. Liquigen
e.g. MCT Oil

WALNUT OIL – **Restricted** see terms above

- ⬆ Liq

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Protein			
➡ Restricted			
Initiation — Use as an additive			
Either:			
1 Protein losing enteropathy; or			
2 High protein needs.			
Initiation — Use as a module			
For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk. .			
Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.			
PROTEIN SUPPLEMENT – Restricted see terms above			
⬆ Powder 5 g protein, 0.67 g carbohydrate and 0.6 g fat per 6.6 g, 275 g can			<i>e.g. Promod</i>
⬆ Powder 6 g protein per 7 g, can8.95	227 g		Resource Beneprotein
⬆ Powder 89 g protein, <1.5 g carbohydrate and 2 g fat per 100 g, 225 g can			<i>e.g. Protifar</i>
Other Supplements			
BREAST MILK FORTIFIER			
Powder 0.2 g protein, 0.7 g carbohydrate and 0.02 g fat per 1 g sachet			<i>e.g. FM 85</i>
Powder 0.5 g protein, 1.2 g carbohydrate and 0.08 g fat per 2 g sachet			<i>e.g. S26 Human Milk Fortifier</i>
Powder 0.6 g protein and 1.4 g carbohydrate per 2.2 g sachet			<i>e.g. Nutricia Breast Milk Fortifier</i>
CARBOHYDRATE AND FAT SUPPLEMENT – Restricted see terms below			
⬆ Powder 72.7 g carbohydrate and 22.3 g fat per 100 g, 400 g can			<i>e.g. Super Soluble Duocal</i>
➡ Restricted			
Initiation			
Both:			
1 Infant or child aged four years or under; and			
2 Any of the following:			
2.1 Cystic fibrosis; or			
2.2 Cancer in children; or			
2.3 Faltering growth; or			
2.4 Bronchopulmonary dysplasia; or			
2.5 Premature and post premature infants.			
NOTE:			
While pre-thickened drinks and supplements have not been included in Section H, DHB hospitals may continue to use such products for patients with dysphagia, provided that:			
• use was established prior to 1 July 2013; and			
• the product has not been specifically considered and excluded by PHARMAC; and			
• use of the product conforms to any applicable indication restrictions for similar products that are listed in Section H (for example, use of thickened high protein products should be in line with the restriction for high protein oral feed in Section H).			
PHARMAC intends to make a further decision in relation to pre-thickened drinks and supplements in the future, and will notify of any change to this situation.			
⬆ Item restricted (see ➡ above); ⬆ Item restricted (see ➡ below)			
<i>e.g. Brand</i> indicates brand example only. It is not a contracted product.			

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Food/Fluid Thickeners		
CAROB BEAN GUM WITH MAIZE STARCH AND MALTODEXTRIN Powder		<i>e.g. Feed Thickener Karicare Aptamil</i>
GUAR GUM Powder		<i>e.g. Guarcol</i>
MAIZE STARCH Powder		<i>e.g. Resource Thicken Up; Nutilis</i>
MALTODEXTRIN WITH XANTHAN GUM Powder		<i>e.g. Instant Thick</i>
MALTODEXTRIN WITH XANTHAN GUM AND ASCORBIC ACID Powder		<i>e.g. Easy Thick</i>

Metabolic Products

➔ Restricted

Initiation

Any of the following:

- 1 For the dietary management of homocystinuria, maple syrup urine disease, phenylketonuria (PKU), glutaric aciduria, isovaleric acidemia, propionic acidemia, methylmalonic acidemia, tyrosinaemia or urea cycle disorders; or
- 2 Patient has adrenoleukodystrophy; or
- 3 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Glutaric Aciduria Type 1 Products

AMINO ACID FORMULA (WITHOUT LYSINE AND LOW TRYPTOPHAN) – **Restricted** see terms above

- ⬆ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre
per 100 g, 400 g can *e.g. GA1 Anamix Infant*
- ⬆ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can *e.g. XLYS Low TRY
Maxamaid*

Homocystinuria Products

AMINO ACID FORMULA (WITHOUT METHIONINE) – **Restricted** see terms above

- ⬆ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre
per 100 g, 400 g can *e.g. HCU Anamix Infant*
- ⬆ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can *e.g. XMET Maxamaid*
- ⬆ Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can *e.g. XMET Maxamum*
- ⬆ Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per
100 ml, 125 ml bottle *e.g. HCU Anamix Junior
LQ*

Isovaleric Acidemia Products

AMINO ACID FORMULA (WITHOUT LEUCINE) – **Restricted** see terms above

- ⬆ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre
per 100 g, 400 g can *e.g. IVA Anamix Infant*
- ⬆ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can *e.g. XLEU Maxamaid*
- ⬆ Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can *e.g. XLEU Maxamum*

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Maple Syrup Urine Disease Products			
AMINO ACID FORMULA (WITHOUT ISOLEUCINE, LEUCINE AND VALINE) – Restricted see terms on the preceding page			
☞ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can			e.g. MSUD Anamix Infant
☞ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can			e.g. MSUD Maxamaid
☞ Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can			e.g. MSUD Maxamum
☞ Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle			e.g. MSUD Anamix Junior LQ

Phenylketonuria Products

AMINO ACID FORMULA (WITHOUT PHENYLALANINE) – Restricted see terms on the preceding page			
☞ Tab 8.33 mg			e.g. Phlexy-10
☞ Powder 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet			e.g. PKU Anamix Junior
☞ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can			e.g. PKU Anamix Infant
☞ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can			e.g. XP Maxamaid
☞ Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can			e.g. XP Maxamum
☞ Powder 8.33 g protein and 8.8 g carbohydrate per 20 g sachet			e.g. Phlexy-10
☞ Liquid 10 g protein, 4.4 g carbohydrate and 0.25 g fibre per 100 ml, 62.5 ml bottle			e.g. PKU Lophlex LQ 10
☞ Liquid 20 g protein, 8.8 g carbohydrate and 0.34 g fibre per 100 ml, 125 ml bottle			e.g. PKU Lophlex LQ 20
☞ Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, bottle	13.10	125 ml	PKU Anamix Junior LQ (Berry)
			PKU Anamix Junior LQ (Orange)
			PKU Anamix Junior LQ (Unflavoured)
☞ Liquid 16 g protein, 7 g carbohydrate and 0.27 g fibre per 100 ml, 125 ml bottle			e.g. PKU Lophlex LQ 20
☞ Liquid 16 g protein, 7 g carbohydrate and 0.27 g fibre per 100 ml, 62.5 ml bottle			e.g. PKU Lophlex LQ 10
☞ Liquid 16 g protein, 7 g carbohydrate and 0.4 g fibre per 100 ml, 125 ml bottle			e.g. PKU Lophlex LQ 20
☞ Liquid 16 g protein, 7 g carbohydrate and 0.4 g fibre per 100 ml, 62.5 ml bottle			e.g. PKU Lophlex LQ 10
☞ Liquid 6.7 g protein, 5.1 g carbohydrate and 2 g fat per 100 ml, 250 ml carton			e.g. Easiphen

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

Propionic Acidaemia and Methylmalonic Acidaemia Products

AMINO ACID FORMULA (WITHOUT ISOLEUCINE, METHIONINE, THREONINE AND VALINE) – **Restricted** see terms on page 203

⬆ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can	<i>e.g. MMA/PA Anamix Infant</i>
⬆ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can	<i>e.g. XMTVI Maxamaid</i>
⬆ Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can	<i>e.g. XMTVI Maxamum</i>

Protein Free Supplements

PROTEIN FREE SUPPLEMENT – **Restricted** see terms on page 203

⬆ Powder nil added protein and 67 g carbohydrate per 100 g, 400 g can	<i>e.g. Energivit</i>
---	-----------------------

Tyrosinaemia Products

AMINO ACID FORMULA (WITHOUT PHENYLALANINE AND TYROSINE) – **Restricted** see terms on page 203

⬆ Powder 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet	<i>e.g. TYR Anamix Junior</i>
⬆ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can	<i>e.g. TYR Anamix Infant</i>
⬆ Powder 25 g protein and 51 g carbohydrate per 100 g, 400 g can	<i>e.g. XPHEN, TYR Maxamaid</i>
⬆ Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle	<i>e.g. TYR Anamix Junior LQ</i>

Urea Cycle Disorders Products

AMINO ACID SUPPLEMENT – **Restricted** see terms on page 203

⬆ Powder 25 g protein and 65 g carbohydrate per 100 g, 200 g can	<i>e.g. Dialamine</i>
⬆ Powder 79 g protein per 100 g, 200 g can	<i>e.g. Essential Amino Acid Mix</i>

X-Linked Adrenoleukodystrophy Products

GLYCEROL TRIERUCATE – **Restricted** see terms on page 203

⬆ Liquid, 1,000 ml bottle

GLYCEROL TRIOLEATE – **Restricted** see terms on page 203

⬆ Liquid, 500 ml bottle

Specialised Formulas

Diabetic Products

➡ **Restricted**

Initiation

Any of the following:

- 1 For patients with type I or type II diabetes suffering weight loss and malnutrition that requires nutritional support; or
- 2 For patients with pancreatic insufficiency; or
- 3 For patients who have, or are expected to, eat little or nothing for 5 days; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 4 For patients who have a poor absorptive capacity and/or high nutrient losses and/or increased nutritional needs from causes such as catabolism; or
- 5 For use pre- and post-surgery; or
- 6 For patients being tube-fed; or
- 7 For tube-feeding as a transition from intravenous nutrition.

LOW-GI ENTERAL FEED 1 KCAL/ML – **Restricted** see terms on the preceding page

⚡ Liquid 5 g protein, 9.6 g carbohydrate and 5.4 g fat per 100 ml, 1,000 ml bottle	7.50	1,000 ml	Glucerna Select RTH (Vanilla)
⚡ Liquid 4.3 g protein, 11.3 g carbohydrate and 4.2 g fat per 100 ml, 1,000 ml bag			<i>e.g. Nutrison Advanced Diason</i>

LOW-GI ORAL FEED 1 KCAL/ML – **Restricted** see terms on the preceding page

⚡ Liquid 4.5 g protein, 9.8 g carbohydrate, 4.4 g fat and 1.9 g fibre per 100 ml, can	2.10	237 ml	Sustagen Diabetic (Vanilla)
⚡ Liquid 5 g protein, 9.6 g carbohydrate and 5.4 g fat per 100 ml, 250 ml bottle	1.88	250 ml	Glucerna Select (Vanilla)
⚡ Liquid 6 g protein, 9.5 g carbohydrate, 4.7 g fat and 2.6 g fibre per 100 ml, can	2.10	237 ml	Resource Diabetic (Vanilla)
⚡ Liquid 4.9 g protein, 11.7 g carbohydrate, 3.8 g fat and 2 g fibre per 100 ml, 200 ml bottle			<i>e.g. Diasip</i>

Elemental and Semi-Elemental Products

➡ **Restricted**

Initiation

Any of the following:

- 1 Malabsorption; or
- 2 Short bowel syndrome; or
- 3 Enterocutaneous fistulas; or
- 4 Eosinophilic enteritis (including oesophagitis); or
- 5 Inflammatory bowel disease; or
- 6 Acute pancreatitis where standard feeds are not tolerated; or
- 7 Patients with multiple food allergies requiring enteral feeding.

AMINO ACID ORAL FEED – **Restricted** see terms above

⚡ Powder 11 g protein, 62 g carbohydrate and 1 g fat per sachet	4.50	80 g	Vivonex TEN
⚡ Powder 11.5 g protein, 61.7 g carbohydrate and 0.8 g fat per sachet	4.50	80.4 g	Vivonex TEN

(Vivonex TEN Powder 11.5 g protein, 61.7 g carbohydrate and 0.8 g fat per sachet to be delisted 1 June 2016)

AMINO ACID ORAL FEED 0.8 KCAL/ML – **Restricted** see terms above

⚡ Liquid 2.5 g protein, 11 g carbohydrate and 3.5 g fat per 100 ml, 250 ml carton			<i>e.g. Elemental 028 Extra</i>
---	--	--	---------------------------------

PEPTIDE-BASED ENTERAL FEED 1 KCAL/ML – **Restricted** see terms above

⚡ Liquid 4 g protein, 17.6 g carbohydrate and 1.7 g fat per 100 ml, 1,000 ml bag			<i>e.g. Nutrison Advanced Peptisorb</i>
--	--	--	---

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PEPTIDE-BASED ORAL FEED – Restricted see terms on the preceding page			
⬆ Powder 13.7 g protein, 62.9 g carbohydrate and 17.5 g fat per 100 g, 400 g can			<i>e.g. Peptamen Junior</i>
⬆ Powder 13.8 g protein, 59 g carbohydrate and 18 g fat per 100 g, 400 g can			<i>e.g. MCT Pepdite; MCT Pepdite 1+</i>
⬆ Powder 15.8 g protein, 49.5 g carbohydrate and 4.65 g fat per 76 g sachet	7.50	76 g	Alitraq
⬆ Liquid 6.75 g protein, 18.4 g carbohydrate and 5.5 g fat per 100 ml, bottle	18.06	1,000 ml	Vital
PEPTIDE-BASED ORAL FEED 1 KCAL/ML – Restricted see terms on the preceding page			
⬆ Liquid 5 g protein, 16 g carbohydrate and 1.69 g fat per 100 ml, carton	4.95	237 ml	Peptamen OS 1.0 (Vanilla)

Fat Modified Products

FAT-MODIFIED FEED – **Restricted** see terms below

⬆ Powder 11.4 g protein, 68 g carbohydrate and 11.8 g fat per 100 g, 400 g can			<i>e.g. Monogen</i>
---	--	--	---------------------

➡ Restricted

Initiation

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

Hepatic Products

➡ Restricted

Initiation

For children (up to 18 years) who require a liver transplant.

HEPATIC ORAL FEED – **Restricted** see terms above

⬆ Powder 11 g protein, 64 g carbohydrate and 20 g fat per 100 g, can	78.97	400 g	Heparon Junior
--	-------	-------	----------------

High Calorie Products

➡ Restricted

Initiation

Any of the following:

- 1 Patient is fluid volume or rate restricted; or
- 2 Patient requires low electrolyte; or
- 3 Both:
 - 3.1 Any of the following:
 - 3.1.1 Cystic fibrosis; or
 - 3.1.2 Any condition causing malabsorption; or
 - 3.1.3 Faltering growth in an infant/child; or
 - 3.1.4 Increased nutritional requirements; and
 - 3.2 Patient has substantially increased metabolic requirements.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ENTERAL FEED 2 KCAL/ML – Restricted see terms on the preceding page			
☞ Liquid 7.5 g protein, 20 g carbohydrate and 10 g fat per 100 ml, bottle	5.50	500 ml	Nutrison Concentrated
☞ Liquid 8.4 g protein, 21.9 g carbohydrate, 9.1 g fat and 0.5 g fibre per 100 ml, bottle	11.00	1,000 ml	TwoCal HN RTH (Vanilla)
ORAL FEED 2 KCAL/ML – Restricted see terms on the preceding page			
☞ Liquid 8.4 g protein, 22.4 g carbohydrate, 8.9 g fat and 0.8 g fibre per 100 ml, bottle	1.90	200 ml	Two Cal HN

High Protein Products

HIGH PROTEIN ENTERAL FEED 1.25 KCAL/ML – **Restricted** see terms below

☞ Liquid 6.3 g protein, 14.2 g carbohydrate and 4.9 g fat per 100 ml,
1,000 ml bag

*e.g. Nutrison Protein
Plus*

☞Restricted

Initiation

Both:

- 1 The patient has a high protein requirement; and
- 2 Any of the following:
 - 2.1 Patient has liver disease; or
 - 2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
 - 2.3 Patient is fluid restricted; or
 - 2.4 Patient's needs cannot be more appropriately met using high calorie product.

HIGH PROTEIN ENTERAL FEED 1.28 KCAL/ML – **Restricted** see terms below

☞ Liquid 6.3 g protein, 14.1 g carbohydrate, 4.9 g fat and 1.5 g fibre per
100 ml, 1,000 ml bag

*e.g. Nutrison Protein
Plus Multi Fibre*

☞Restricted

Initiation

Both:

- 1 The patient has a high protein requirement; and
- 2 Any of the following:
 - 2.1 Patient has liver disease; or
 - 2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
 - 2.3 Patient is fluid restricted; or
 - 2.4 Patient's needs cannot be more appropriately met using high calorie product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Infant Formulas			
AMINO ACID FORMULA – Restricted see terms below			
☞ Powder 1.95 g protein, 8.1 g carbohydrate and 3.5 g fat per 100 ml, 400 g can			<i>e.g. Neocate</i>
☞ Powder 13 g protein, 52.5 g carbohydrate and 24.5 g fat per 100 g, 400 g can			<i>e.g. Neocate LCP</i>
☞ Powder 13.5 g protein, 52 g carbohydrate and 24.5 g fat per 100 g, can53.00	400 g		Neocate Gold (Unflavoured)
☞ Powder 14 g protein, 50 g carbohydrate and 24.3 g fat per 100 g, 400 g can			<i>e.g. Neocate Advance</i>
☞ Powder 16 g protein, 51.4 g carbohydrate and 21 g fat per 100 g, can53.00	400 g		Neocate Advance (Vanilla)
☞ Powder 2.2 g protein, 7.8 g carbohydrate and 3.4 g fat per 100 ml, can53.00	400 g		Elecare LCP (Unflavoured)
☞ Powder 2.2 g protein, 7.8 g carbohydrate and 3.4 g fat per 100 ml, can53.00	400 g		Elecare (Unflavoured) Elecare (Vanilla)
☞ Powder 6 g protein, 31.5 g carbohydrate and 5.88 g fat per sachet6.00	48.5 g		Vivonex Paediatric

☞ Restricted

Initiation

Any of the following:

- 1 Extensively hydrolysed formula has been reasonably trialled and is inappropriate due to documented severe intolerance or allergy or malabsorption; or
- 2 History of anaphylaxis to cows' milk protein formula or dairy products; or
- 3 Eosinophilic oesophagitis.

Note: A reasonable trial is defined as a 2-4 week trial.

Continuation

Both:

- 1 An assessment as to whether the infant can be transitioned to a cows' milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an amino acid infant formula.

EXTENSIVELY HYDROLYSED FORMULA – **Restricted** see terms below

☞ Powder 14 g protein, 53.4 g carbohydrate and 27.3 g fat per 100 g, 450 g can	<i>e.g. Aptamil Gold+ Pepti Junior</i>
--	--

☞ Restricted

Initiation

Any of the following:

- 1 Both:
 - 1.1 Cows' milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malabsorption; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
7 Cystic fibrosis; or			
8 Proven fat malabsorption; or			
9 Severe intestinal motility disorders causing significant malabsorption; or			
10 Intestinal failure; or			
11 For step down from Amino Acid Formula.			
Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.			
Continuation			
Both:			
1 An assessment as to whether the infant can be transitioned to a cows' milk protein or soy infant formula has been undertaken; and			
2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula.			
FRUCTOSE-BASED FORMULA			
Powder 14.6 g protein, 49.7 g carbohydrate and 30.8 g fat per 100 g, 400 g can			<i>e.g. Galactomin 19</i>
LACTOSE-FREE FORMULA			
Powder 1.3 g protein, 7.3 g carbohydrate and 3.5 g fat per 100 ml, 900 g can			<i>e.g. Karicare Aptamil Gold De-Lact</i>
Powder 1.5 g protein, 7.2 g carbohydrate and 3.6 g fat per 100 ml, 900 g can			<i>e.g. S26 Lactose Free</i>
LOW-CALCIUM FORMULA			
Powder 14.6 g protein, 53.7 g carbohydrate and 26.1 g fat per 100 g, 400 g can			<i>e.g. Locasol</i>
PAEDIATRIC ORAL FEED 1 KCAL/ML – Restricted see terms below			
☞ Liquid 2.6 g protein, 10.3 g carbohydrate, 5.4 g fat and 0.6 g fibre per 100 ml, 100 ml bottle			<i>e.g. Infatrini</i>
☞Restricted			
Initiation			
Both:			
1 Either:			
1.1 The patient is fluid restricted; or			
1.2 The patient has increased nutritional requirements due to faltering growth; and			
2 Patient is under 18 months old and weighs less than 8kg.			
PRETERM FORMULA – Restricted see terms below			
☞ Powder 1.9 g protein, 7.5 g carbohydrate and 3.9 g fat per 14 g, can15.25	400 g		S-26 Gold Premgro
☞ Liquid 2.2 g protein, 8.4 g carbohydrate and 4.4 g fat per 100 ml, bottle0.75	100 ml		S26 LBW Gold RTF
☞ Liquid 2.3 g protein, 8.6 g carbohydrate and 4.2 g fat per 100 ml, 90 ml bottle			<i>e.g. Pre Nan Gold RTF</i>
☞ Liquid 2.6 g protein, 8.4 g carbohydrate and 3.9 g fat per 100 ml, 70 ml bottle			<i>e.g. Karicare Aptamil Gold+Preterm</i>
☞Restricted			
Initiation			
For infants born before 33 weeks' gestation or weighing less than 1.5 kg at birth.			
THICKENED FORMULA			
Powder 1.8 g protein, 8.1 g carbohydrate and 3.3 g fat per 100 ml, 900 g can			<i>e.g. Karicare Aptamil Thickened AR</i>

☞ Item restricted (see ☞ above); ☞ Item restricted (see ☞ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Ketogenic Diet Products			
HIGH FAT FORMULA – Restricted see terms below			
☛ Powder 15.25 g protein, 3 g carbohydrate and 73 g fat per 100 g, can	35.50	300 g	Ketocal 4:1 (Unflavoured) Ketocal 4:1 (Vanilla)
☛ Powder 15.3 g protein, 7.2 g carbohydrate and 67.7 g fat per 100 g, can	35.50	300 g	Ketocal 3:1 (Unflavoured)
➔ Restricted			
Initiation			
For patients with intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.			
Paediatric Products			
➔ Restricted			
Initiation			
Both:			
1 Child is aged one to ten years; and			
2 Any of the following:			
2.1 The child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or			
2.2 Any condition causing malabsorption; or			
2.3 Faltering growth in an infant/child; or			
2.4 Increased nutritional requirements; or			
2.5 The child is being transitioned from TPN or tube feeding to oral feeding; or			
2.6 The child has eaten, or is expected to eat, little or nothing for 3 days.			
PAEDIATRIC ORAL FEED – Restricted see terms above			
☛ Powder 14.9 g protein, 54.3 g carbohydrate and 24.7 g fat per 100 g, can	20.00	850 g	Pediasure (Vanilla)
PAEDIATRIC ENTERAL FEED 0.76 KCAL/ML – Restricted see terms above			
☛ Liquid 2.5 g protein, 12.5 g carbohydrate, 3.3 g fat and 0.7 g fibre per 100 ml, bag	4.00	500 ml	Nutrini Low Energy Multifibre RTH
PAEDIATRIC ENTERAL FEED 1 KCAL/ML – Restricted see terms above			
☛ Liquid 2.8 g protein, 11.2 g carbohydrate and 5 g fat per 100 ml, bag	2.68	500 ml	Pediasure RTH
☛ Liquid 2.8 g protein, 12.3 g carbohydrate and 4.4 g fat per 100 ml, 500 ml bag			<i>e.g. Nutrini RTH</i>
PAEDIATRIC ENTERAL FEED 1.5 KCAL/ML – Restricted see terms above			
☛ Liquid 4.1 g protein, 18.5 g carbohydrate, 6.7 g fat and 0.8 g fibre per 100 ml, bag	6.00	500 ml	Nutrini Energy Multi Fibre
☛ Liquid 4.1 g protein, 18.5 g carbohydrate and 6.7 g fat per 100 ml, 500 ml bag			<i>e.g. Nutrini Energy RTH</i>
PAEDIATRIC ORAL FEED 1 KCAL/ML – Restricted see terms above			
☛ Liquid 4.2 g protein, 16.7 g carbohydrate and 7.5 g fat per 100 ml, bottle	1.07	200 ml	Pediasure (Chocolate) Pediasure (Strawberry) Pediasure (Vanilla)
☛ Liquid 4.2 g protein, 16.7 g carbohydrate and 7.5 g fat per 100 ml, can	1.34	250 ml	Pediasure (Vanilla)

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PAEDIATRIC ORAL FEED 1.5 KCAL/ML – Restricted see terms on the preceding page			
☞ Liquid 3.4 g protein, 18.8 g carbohydrate and 6.8 g fat per 100 ml, 200 ml bottle			<i>e.g. Fortini</i>
☞ Liquid 4.0 g protein, 18.8 g carbohydrate, 6.8 g fat and 1.5 g fibre per 100 ml, 200 ml bottle			<i>e.g. Fortini Multifibre</i>

Renal Products

LOW ELECTROLYTE ENTERAL FEED 1.8 KCAL/ML – Restricted see terms below			
☞ Liquid 8.1 g protein, 14.74 g carbohydrate, 9.77 g fat and 1.26 g fibre per 100 ml, bottle	6.08	500 ml	Nepro HP RTH
☞ Restricted			
Initiation			
For patients with acute or chronic kidney disease.			
LOW ELECTROLYTE ORAL FEED – Restricted see terms below			
☞ Powder 7.5 g protein, 59 g carbohydrate and 26.3 g fat per 100 g, 400 g can			<i>e.g. Kindergen</i>
☞ Restricted			
Initiation			
For children (up to 18 years) with acute or chronic kidney disease.			
LOW ELECTROLYTE ORAL FEED 1.8 KCAL/ML			
☞ Liquid 8 g protein, 14.74 g carbohydrate, 9.77 g fat and 1.26 g fibre per 100 ml, carton	2.67	220 ml	Nepro HP (Strawberry) Nepro HP (Vanilla)
☞ Restricted			
Initiation			
For patients with acute or chronic kidney disease.			
LOW ELECTROLYTE ORAL FEED 2 KCAL/ML – Restricted see terms below			
☞ Liquid 9.1 g protein, 19 g carbohydrate and 10 g fat per 100 ml, carton	3.31	237 ml	Novasource Renal (Vanilla)
☞ Liquid 3 g protein, 25.5 g carbohydrate and 9.6 g fat per 100 ml, 237 ml bottle			
☞ Liquid 7.5 g protein, 20 g carbohydrate and 10 g fat per 100 ml, 125 ml carton			<i>e.g. Renilon 7.5</i>
☞ Restricted			
Initiation			
For patients with acute or chronic kidney disease.			

Respiratory Products

LOW CARBOHYDRATE ORAL FEED 1.5 KCAL/ML – Restricted see terms below			
☞ Liquid 6.2 g protein, 10.5 g carbohydrate and 9.32 g fat per 100 ml, bottle	1.66	237 ml	Pulmocare (Vanilla)
☞ Restricted			
Initiation			
For patients with CORD and hypercapnia, defined as a CO2 value exceeding 55 mmHg.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Surgical Products

HIGH ARGININE ORAL FEED 1.4 KCAL/ML – **Restricted** see terms below

⬇ Liquid 7.6 g protein, 18.9 g carbohydrate, 3.9 g fat and 1.4 g fibre per 100 ml, carton	4.00	237 ml	Impact Advanced Recovery (Chocolate) Impact Advanced Recovery (Vanilla)
--	------	--------	---

➡ **Restricted**
Initiation

Three packs per day for 5 to 7 days prior to major gastrointestinal, head or neck surgery.

PREOPERATIVE CARBOHYDRATE FEED 0.5 KCAL/ML – **Restricted** see terms below

⬇ Oral liq 0 g protein, 12.6 g carbohydrate and 0 g fat per 100 ml, 200 ml bottle	6.80	4	preOp
--	------	---	-------

➡ **Restricted**
Initiation

Maximum of 400 ml as part of an Enhanced Recovery After Surgery (ERAS) protocol 2 to 3 hours before major abdominal surgery.

Standard Feeds

➡ **Restricted**
Initiation

Any of the following:

For patients with malnutrition, defined as any of the following:

- 1 Any of the following:
 - 1.1 BMI < 18.5; or
 - 1.2 Greater than 10% weight loss in the last 3-6 months; or
 - 1.3 BMI < 20 with greater than 5% weight loss in the last 3-6 months; or
- 2 For patients who have, or are expected to, eat little or nothing for 5 days; or
- 3 For patients who have a poor absorptive capacity and/or high nutrient losses and/or increased nutritional needs from causes such as catabolism; or
- 4 For use pre- and post-surgery; or
- 5 For patients being tube-fed; or
- 6 For tube-feeding as a transition from intravenous nutrition; or
- 7 For any other condition that meets the community Special Authority criteria.

ENTERAL FEED 1.5 KCAL/ML – **Restricted** see terms above

⬆ Liquid 5.4 g protien, 13.6 g carbohydrate and 3.3 g fat per 100 ml, 1,000 ml bottle			<i>e.g. Isosource Standard RTH</i>
⬆ Liquid 6 g protein, 18.3 g carbohydrate and 5.8 g fat per 100 ml, bag	7.00	1,000 ml	Nutrison Energy
⬆ Liquid 6 g protein, 18.4 g carbohydrate, 5.8 g fat and 1.5 g fibre per 100 ml, 1,000 ml bag			<i>e.g. Nutrison Energy Multi Fibre</i>
⬆ Liquid 6.25 g protein, 20 g carbohydrate and 5 g fat per 100 ml, can	1.75	250 ml	Ensure Plus HN
⬆ Liquid 6.27 g protein, 20.4 g carbohydrate and 4.9 g fat per 100 ml, bag	7.00	1,000 ml	Ensure Plus HN RTH
⬆ Liquid 6.38 g protein, 21.1 g carbohydrate, 4.9 g fat and 1.2 g fibre per 100 ml, bag	7.00	1,000 ml	Jevity HiCal RTH

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ENTERAL FEED 1 KCAL/ML – Restricted see terms on the preceding page			
☞ Liquid 4 g protein, 13.6 g carbohydrate and 3.4 g fat per 100 ml, bottle	5.29	1,000 ml	Osmolite RTH
☞ Liquid 4 g protein, 13.6 g carbohydrate and 3.4 g fat per 100 ml, can	1.24	250 ml	Osmolite
☞ Liquid 4 g protein, 14.1 g carbohydrate, 3.47 g fat and 1.76 g fibre per 100 ml, bottle	2.65	500 ml	Jevity RTH
	5.29	1,000 ml	Jevity RTH
☞ Liquid 4 g protein, 14.1 g carbohydrate, 3.47 g fat and 1.76 g fibre per 100 ml, can	1.32	237 ml	Jevity
☞ Liquid 4 g protein, 12.3 g carbohydrate and 3.9 g fat per 100 ml, 1,000 ml bag			<i>e.g. NutrisonStdRTH; NutrisonLowSodium</i>
☞ Liquid 4 g protein, 12.3 g carbohydrate, 3.9 g fat and 1.5 g fibre per 100 ml, 1000 ml bag			<i>e.g. Nutrison Multi Fibre</i>
<i>(Osmolite Liquid 4 g protein, 13.6 g carbohydrate and 3.4 g fat per 100 ml, can to be delisted 1 June 2016)</i>			
ENTERAL FEED 1.2 KCAL/ML – Restricted see terms on the preceding page			
☞ Liquid 5.55 g protein, 15.1 g carbohydrate, 3.93 g fat and 2 g fibre per 100 ml, 1,000 ml bag			<i>e.g. Jevity Plus RTH</i>
ORAL FEED – Restricted see terms on the preceding page			
☞ Powder 16 g protein, 59.8 g carbohydrate and 14 g fat per 100 g, can	13.00	850 g	Ensure (Chocolate) Ensure (Vanilla)
☞ Powder 21.9 g protein, 53.5 g carbohydrate and 14.5 g fat per 100 g, can	3.67	350 g	Fortisip (Vanilla)
☞ Powder 23 g protein, 65 g carbohydrate and 2.5 g fat per 100 g, can	14.90	840 g	Sustagen Hospital Formula (Chocolate) Sustagen Hospital Formula (Vanilla)
Note: Community subsidy of Sustagen Hospital Formula is subject to both Special Authority criteria and a manufacturer's surcharge. Higher subsidy by endorsement is available for patients meeting the following endorsement criteria; fat malabsorption, fat intolerance or chyle leak.			
ORAL FEED 1 KCAL/ML – Restricted see terms on the preceding page			
☞ Liquid 3.8 g protein, 23 g carbohydrate and 12.7 g fibre per 100 ml, 237 ml carton			<i>e.g. Resource Fruit Beverage</i>
ORAL FEED 1.5 KCAL/ML – Restricted see terms on the preceding page			
☞ Liquid 5.5 g protein, 21.1 g carbohydrate and 4.81 g fat per 100 ml, can	1.33	237 ml	Ensure Plus (Chocolate) Ensure Plus (Vanilla)
☞ Liquid 6.25 g protein, 20.2 g carbohydrate and 4.92 g fat per 100 ml, carton	1.26	200 ml	Ensure Plus (Banana) Ensure Plus (Chocolate) Ensure Plus (Fruit of the Forest) Ensure Plus (Vanilla)
☞ Liquid 4 g protein and 33.5 g carbohydrate per 100 ml, 200 ml bottle			<i>e.g. Fortijuice</i>
☞ Liquid 6 g protein, 18.4 g carbohydrate and 5.8 g fat per 100 ml, 200 ml bottle			<i>e.g. Fortisip</i>
☞ Liquid 6 g protein, 18.4 g carbohydrate, 5.8 g fat and 2.3 g fibre per 100 ml, 200 ml bottle			<i>e.g. Fortisip Multi Fibre</i>

☞ Item restricted (see ➡ above); ☞ Item restricted (see ➡ below)

e.g. Brand indicates brand example only. It is not a contracted product.

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Bacterial and Viral Vaccines

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE – **Restricted** see terms below

<p>☞ Inj 30 IU diphtheria toxoid with 30IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units poliomyelitis virus in 0.5 ml syringe</p> <p>– 1% DV Jul-14 to 2017</p>	0.00	10	Infanrix IPV
---	------	----	--------------

➔ Restricted

Initiation

Any of the following:

- 1 A single dose for children up to the age of 7 who have completed primary immunisation; or
- 2 A course of up to four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
- 3 An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 4 Five doses will be funded for children requiring solid organ transplantation.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes

DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE – **Restricted** see terms below

<p>☞ Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin, 80 D-antigen units poliomyelitis virus, 10 mcg hepatitis B surface antigen in 0.5 ml syringe (1) and inj 10 mcg haemophilus influenzae type B vaccine vial – 1% DV Jul-14 to 2017.....</p>	0.00	10	Infanrix-hexa
---	------	----	---------------

➔ Restricted

Initiation

Any of the following:

- 1 Up to four doses for children up to and under the age of 10 for primary immunisation; or
- 2 An additional four doses (as appropriate) are funded for (re-)immunisation for children up to and under the age of 10 who are patients post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 3 Up to five doses for children up to and under the age of 10 receiving solid organ transplantation.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

Bacterial Vaccines

ADULT DIPHTHERIA AND TETANUS VACCINE

<p>☞ Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid in 0.5 ml syringe –</p> <p>1% DV Jul-14 to 2017</p>	0.00	5	ADT Booster
---	------	---	-------------

➔ Restricted

Initiation

Any of the following:

- 1 For vaccination of patients aged 45 and 65 years old; or
- 2 For vaccination of previously unimmunised or partially immunised patients; or
- 3 For revaccination following immunosuppression; or
- 4 For boosting of patients with tetanus-prone wounds; or
- 5 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BACILLUS CALMETTE-GUERIN VACCINE – Restricted see terms below			
⚡ Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin), Danish strain 1331, live attenuated, vial Danish strain 1331, live attenuated, vial with diluent – 1% DV Oct-14 to 2017	0.00	10	BCG Vaccine
➡Restricted			
Initiation			
All of the following:			
For infants at increased risk of tuberculosis defined as:			
1 Living in a house or family with a person with current or past history of TB; and			
2 Having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; and			
3 During their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000.			
Note: A list of countries with high rates of TB are available at http://www.health.govt.nz/tuberculosis (Search for Downloads) or www.bcgatlas.org/index.php			
DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE – Restricted see terms below			
⚡ Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis toxoid, 8 mcg pertussis filamentous haemagglutinin and 2.5 mcg pertactin in 0.5 ml syringe – 1% DV Jul-14 to 2017	0.00	1 10	Boostrix Boostrix
➡Restricted			
Initiation			
Any of the following:			
1 A single vaccine for pregnant woman between gestational weeks 28 and 38; or			
2 A course of up to four vaccines is funded for children from age 7 up the age of 18 years inclusive to complete full primary immunisation; or			
3 An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens.			
Note: Tdap is not registered for patients aged less than 10 years. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.			
HAEMOPHILUS INFLUENZAE TYPE B VACCINE – Restricted see terms below			
⚡ Inj 10 mcg vial with diluent syringe – 1% DV Jul-14 to 2017	0.00	1	Act-HIB
➡Restricted			
Initiation			
<i>Therapy limited to 1 dose</i>			
Any of the following:			
1 For primary vaccination in children; or			
2 An additional dose (as appropriate) is funded for (re-)immunisation for patients post haematopoietic stem cell transplantation, or chemotherapy; functional asplenic; pre or post splenectomy; pre- or post solid organ transplant, pre- or post cochlear implants, renal dialysis and other severely immunosuppressive regimens; or			
3 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.			
MENINGOCOCCAL (A, C, Y AND W-135) CONJUGATE VACCINE – Restricted see terms on the next page			
⚡ Inj 4 mcg or each meningococcal polysaccharide conjugated to a total of approximately 48 mcg of diphtheria toxoid carrier per 0.5 ml vial – 1% DV Jul-14 to 2017	0.00	1	Menactra

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔ Restricted			
Initiation			
Any of the following:			
1 Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with HIV, complement deficiency (acquired or inherited), functional or anatomic asplenia or pre or post solid organ transplant; or 2 One dose for close contacts of meningococcal cases; or 3 A maximum of two doses for bone marrow transplant patients; or 4 A maximum of two doses for patients following immunosuppression*.			
Notes: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.			
*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.			
MENINGOCOCCAL C CONJUGATE VACCINE – Restricted see terms below			
⚡ Inj 10 mcg in 0.5 ml syringe – 1% DV Jul-14 to 2017	0.00	1 10	Neisvac-C Neisvac-C
➔ Restricted			
Initiation			
Any of the following:			
1 Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with HIV, complement deficiency (acquired or inherited), functional or anatomic asplenia or pre or post solid organ transplant; or 2 One dose for close contacts of meningococcal cases; or 3 A maximum of two doses for bone marrow transplant patients; or 4 A maximum of two doses for patients following immunosuppression*.			
Notes: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.			
*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.			
PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE – Restricted see terms below			
⚡ Inj 30.8 mcg in 0.5 ml syringe – 1% DV Oct-14 to 2017	0.00	1 10	Prevenar 13 Prevenar 13
➔ Restricted			
Initiation			
Any of the following:			
1 A primary course of up to four doses for previously unvaccinated individuals up to the age of 59 months inclusive; or 2 Up to three doses as appropriate to complete the primary course of immunisation for individuals under the age of 59 months who have received one to three doses of PCV10; or 3 One dose is funded for high risk children (over the age of 17 months and up to the age of 18) who have previously received four doses of PCV10; or 4 Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients with HIV, for patients post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, primary immunodeficiency; or 5 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.			
Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes			
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE – Restricted see terms on the next page			
⚡ Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype) – 1% DV Jun-15 to 2017	0.00	1	Pneumovax 23
Products with Hospital Supply Status (HSS) are in bold			
Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➡Restricted

Initiation

Any of the following:

- 1 Up to three doses (as appropriate) for patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy; functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- 2 Up to two doses are funded for high risk children to the age of 18; or
- 3 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

SALMONELLA TYPHI VACCINE – Restricted see terms below

⚡ Inj 25 mcg in 0.5 ml syringe

➡Restricted

Initiation

For use during typhoid fever outbreaks.

Viral Vaccines

HEPATITIS A VACCINE – Restricted see terms below

⚡ Inj 720 ELISA units in 0.5 ml syringe – 1% DV Jul-14 to 2017	0.00	1	Havrix Junior
⚡ Inj 1440 ELISA units in 1 ml syringe – 1% DV Jul-14 to 2017	0.00	1	Havrix

➡Restricted

Initiation

All of the following:

- 1 Two vaccinations for use in transplant patients; and
- 2 Two vaccinations for use in children with chronic liver disease; and
- 3 One dose of vaccine for close contacts of known hepatitis A cases.

HEPATITIS B RECOMBINANT VACCINE

⚡ Inj 5 mcg in 0.5 ml vial – 1% DV Jul-14 to 2017	0.00	1	HBvaxPRO
---	------	---	-----------------

➡Restricted

Initiation

Any of the following:

- 1 For household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2 For children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3 For children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination; or
- 4 For HIV positive patients; or
- 5 For hepatitis C positive patients; or
- 6 for patients following non-consensual sexual intercourse; or
- 7 For patients following immunosuppression; or
- 8 For transplant patients; or
- 9 following needle stick injury.

⚡ Inj 10 mcg in 1 ml vial – 1% DV Jul-14 to 2017	0.00	1	HBvaxPRO
--	------	---	-----------------

➡Restricted

Initiation

Any of the following:

- 1 For household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2 For children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3 For children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued. . .			
4 For HIV positive patients; or			
5 For hepatitis C positive patients; or			
6 for patients following non-consensual sexual intercourse; or			
7 For patients following immunosuppression; or			
8 For transplant patients; or			
9 following needle stick injury.			
¶ Inj 40 mcg per 1 ml vial – 1% DV Jul-14 to 2017	0.00	1	HBvaxPRO
➔ Restricted			
Initiation			
Both:			
1 For dialysis patients; and			
2 For liver or kidney transplant patient.			
HUMAN PAPILLOMAVIRUS (6, 11, 16 AND 18) VACCINE [HPV] – Restricted see terms below			
¶ Inj 120 mcg in 0.5 ml syringe – 1% DV Jul-14 to 2017	0.00	10	Gardasil
➔ Restricted			
Initiation			
<i>Therapy limited to 3 doses</i>			
Any of the following:			
1 Females aged under 20 years old; or			
2 Patients aged under 26 years old with confirmed HIV infection; or			
3 For use in transplant (including stem cell) patients; or			
4 An additional dose for patients under 26 years of age post chemotherapy.			
INFLUENZA VACCINE – Restricted see terms below			
¶ Inj 45 mcg in 0.5 ml syringe	90.00	10	Fluarix Influvac
➔ Restricted			
Initiation — People over 65			
The patient is 65 years of age or over.			
Initiation — cardiovascular disease			
Any of the following:			
1 Ischaemic heart disease; or			
2 Congestive heart failure; or			
3 Rheumatic heart disease; or			
4 Longenital heart disease; or			
5 Cerebro-vascular disease.			
Note: hypertension and/or dyslipidaemia without evidence of end-organ disease is excluded from funding.			
Initiation — chronic respiratory disease			
Either:			
1 Asthma, if on a regular preventative therapy; or			
2 Other chronic respiratory disease with impaired lung function.			
Note: asthma not requiring regular preventative therapy is excluded from funding.			
Initiation — Other conditions			
Either:			
1 Any of the following:			
1.1 Diabetes; or			
1.2 chronic renal disease; or			
1.3 Any cancer, excluding basal and squamous skin cancers if not invasive; or			

continued. . .

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
1.4 Autoimmune disease; or			
1.5 Immune suppression or immune deficiency; or			
1.6 HIV; or			
1.7 Transplant recipient; or			
1.8 Neuromuscular and CNS diseases/ disorders; or			
1.9 Haemoglobinopathies; or			
1.10 Is a child on long term aspirin; or			
1.11 Has a cochlear implant; or			
1.12 Errors of metabolism at risk of major metabolic decompensation; or			
1.13 Pre and post splenectomy; or			
1.14 Down syndrome; or			
1.15 Is pregnant; or			
1.16 Is a child aged four and under who has been hospitalised for respiratory illness or has a history of significant respiratory illness; or			
2 Patients who are compulsorily detained long-term in a forensic unit within a DHB hospital.			
MEASLES, MUMPS AND RUBELLA VACCINE – Restricted see terms below			
¶ Inj 1000 TCID50 measles, 12500 TCID50 mumps and 1000 TCID50 rubella vial with diluent – 1% DV Jul-14 to 2017	0.00	10	M-M-R-II
➡ Restricted			
Initiation — first dose prior to 12 months			
<i>Therapy limited to 3 doses</i>			
Any of the following:			
1 For primary vaccination in children; or			
2 For revaccination following immunosuppression; or			
3 For any individual susceptible to measles, mumps or rubella.			
Initiation — first dose after to 12 months			
<i>Therapy limited to 2 doses</i>			
Any of the following:			
1 For primary vaccination in children; or			
2 For revaccination following immunosuppression; or			
3 For any individual susceptible to measles, mumps or rubella.			
Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.			
POLIOMYELITIS VACCINE – Restricted see terms below			
¶ Inj 80 D-antigen units in 0.5 ml syringe – 1% DV Jul-14 to 2017	0.00	1	IPOL
➡ Restricted			
Initiation			
<i>Therapy limited to 3 doses</i>			
Either:			
1 For partially vaccinated or previously unvaccinated individuals; or			
2 For revaccination following immunosuppression.			
Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.			
RABIES VACCINE			
Inj 2.5 IU vial with diluent			
ROTAVIRUS LIVE REASSORTANT ORAL VACCINE – Restricted see terms on the next page			
¶ Oral susp G1, G2, G3, G4, P1(8) 11.5 million CCID50 units per 2 ml, tube – 1% DV Jul-14 to 2017	0.00	10	RotaTeq

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Initiation			
<i>Therapy limited to 3 doses</i>			
Both:			
1 First dose to be administered in infants aged under 15 weeks of age; and			
2 No vaccination being administered to children aged 8 months or over.			
VARICELLA VACCINE [CHICKEN POX VACCINE] – Restricted see terms below			
¶ Inj 2,000 PFU vial with diluent – 1% DV Jul-14 to 2017			
	0.00	1	Varilrix

➔Restricted**Initiation***Therapy limited to 2 doses*

Any of the following:

- 1 Any of the following:
 - for non-immune patients
 - 1.1 With chronic liver disease who may in future be candidates for transplantation; or
 - 1.2 With deteriorating renal function before transplantation; or
 - 1.3 Prior to solid organ transplant; or
 - 1.4 Prior to any elective immunosuppression*; or
 - 1.5 For post exposure prophylaxis who are immune competent inpatients.; or
- 2 For patients at least 2 years after bone marrow transplantation, on advice of their specialist; or
- 3 For patients at least 6 months after completion of chemotherapy, on advice of their specialist; or
- 4 For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
- 5 For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or
- 6 For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or
- 7 For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.

Note: * immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

PART III - OPTIONAL PHARMACEUTICALS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
NOTE:			
In addition to the products expressly listed here in Part III: Optional Pharmaceuticals, a number of additional Optional Pharmaceuticals, including some wound care products and disposable laparoscopic equipment, are listed in an addendum to Part III which is available at www.pharmac.govt.nz . The Optional Pharmaceuticals listed in the addendum are deemed to be listed in Part III, and the Rules of the Pharmaceutical Schedule applying to products listed in Part III apply to them.			
Optional Pharmaceuticals			
BLOOD GLUCOSE DIAGNOSTIC TEST METER			
1 meter with 50 lancets, a lancing device, and 10 diagnostic test strips	20.00	1	Caresens II Caresens N Caresens N POP
Meter	19.00	1	Accu-Chek Performa FreeStyle Lite On Call Advanced
	9.00		
BLOOD GLUCOSE DIAGNOSTIC TEST STRIP			
Blood glucose test strips	28.75	50 test	Accu-Chek Performa CareSens CareSens N FreeStyle Lite
	10.56		FreeStyle Optium On Call Advanced
	21.65		
	28.75		
Blood glucose test strips × 50 and lancets × 5	19.10	50 test	
BLOOD KETONE DIAGNOSTIC TEST METER			
Meter	40.00	1	Freestyle Optium Freestyle Optium Neo
<i>(Freestyle Optium Meter to be delisted 1 May 2016)</i>			
INSULIN PEN NEEDLES			
29 g × 12.7 mm	10.50	100	B-D Micro-Fine
31 g × 5 mm	11.75	100	B-D Micro-Fine
31 g × 6 mm	10.50	100	ABM
31 g × 8 mm	10.50	100	B-D Micro-Fine
32 g × 4 mm	10.50	100	B-D Micro-Fine
INSULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE			
Syringe 0.3 ml with 29 g × 12.7 mm needle	13.00	100	B-D Ultra Fine
Syringe 0.3 ml with 31 g × 8 mm needle	13.00	100	B-D Ultra Fine II
Syringe 0.5 ml with 29 g × 12.7 mm needle	13.00	100	B-D Ultra Fine
Syringe 0.5 ml with 31 g × 8 mm needle	13.00	100	B-D Ultra Fine II
Syringe 1 ml with 29 g × 12.7 mm needle	13.00	100	B-D Ultra Fine
Syringe 1 ml with 31 g × 8 mm needle	13.00	100	B-D Ultra Fine II
KETONE BLOOD BETA-KETONE ELECTRODES			
Test strips	15.50	10 strip	Freestyle Optium Ketone
MASK FOR SPACER DEVICE			
Small	2.20	1	e-chamber Mask
PEAK FLOW METER			
Low Range	9.54	1	Mini-Wright AFS Low Range
Normal Range	9.54	1	Mini-Wright Standard
PREGNANCY TEST - HCG URINE			
Cassette – 1% DV Sep-15 to 2017	17.60	40 test	EasyCheck

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SODIUM NITROPRUSSIDE			
Test strip	6.00	50 strip	Accu-Chek Ketur-Test
SPACER DEVICE			
220 ml (single patient)	2.95	1	e-chamber Turbo
510 ml (single patient)	5.12	1	e-chamber La Grande
800 ml	6.50	1	Volumatic

- Symbols -		Agents Affecting the	Alimentary	23
8-methoxypsoralen	55	Renin-Angiotensin System	Infection	78
- A -		Agents for Parkinsonism and	Amsacrine	135
A-Scabies	52	Related Disorders	Amyl nitrite	48
Abacavir sulphate	85	Agents Used in the Treatment of	Anabolic Agents	61
Abacavir sulphate with		Poisonings	Anaesthetics	106
lamivudine	85	Air Flow Products	Anagrelide hydrochloride	135
Abciximab	152	Ajmaline	Analgesics	109
Abilify	122	Alanase	Anastrozole	145
Abiraterone acetate	144	Albendazole	Andriol Testocaps	61
Acarbose	16	Alendronate sodium	Androderm	61
Accu-Chek Ketur-Test	223	Alendronate sodium with	Androgen Agonists and	
Accu-Chek Performa	222	cholecalciferol	Antagonists	61
Accuretic 10	39	Alfacalcidol	Anexate	190
Accuretic 20	39	Alfentanil	Anoro Ellipta	180
Acetazolamide	187	Alinia	Antabuse	130
Acetic acid		Alitraq	Antacids and Antiflatulents	13
Extemporaneous	198	Allersoothe	Anti-Infective Agents	57
Genito-Urinary	57	Allopurinol	Anti-Infective Preparations	
Acetic acid with hydroxyquinoline,		Alpha tocopheryl acetate	Dermatological	51
glycerol and ricinoleic acid	57	Alpha-Adrenoceptor Blockers	Sensory	184
Acetic acid with propylene		Alprazolam	Anti-Inflammatory	
glycol	189	Alprostadiol hydrochloride	Preparations	185
Acetylcholine chloride	188	Alteplase	Antiacne Preparations	52
Acetylcysteine	190	Alum	Antiallergy Preparations	177
Aciclovir		Aluminium chloride	Antianaemics	27
Infection	91	Aluminium hydroxide	Antiarrhythmics	40
Sensory	184	Aluminium hydroxide with	Antibacterials	71
Aciclovir-Claris	91	magnesium hydroxide and	Anticholinergic Agents	178
Acid Citrate Dextrose A	32	simethicone	Anticholinesterases	95
Acidex	13	Amantadine hydrochloride	Antidepressants	114
Acipimox	46	AmBisome	Antidiarrhoeals and Intestinal	
Acitretin	55	Ambrisentan	Anti-Inflammatory Agents	13
Aclasta	97	Amethocaine	Antiepilepsy Drugs	115
Act-HIB	216	Nervous	Antifibrinolytics, Haemostatics	
Actemra	171	Sensory	and Local Sclerosants	29
Actinomycin D	133	Amikacin	Antifungals	78
Adalimumab	152	Amiloride hydrochloride	Antihypotensives	41
Adapalene	52	Amiloride hydrochloride with	Antimigraine Preparations	120
Adefin XL	43	furosemide	Antimycobacterials	80
Adefovir dipivoxil	87	Amiloride hydrochloride with	Antinaus	121
Adenosine	40-41	hydrochlorothiazide	Antinausea and Vertigo	
Adenuric	101	Aminophylline	Agents	120
Adrenaline	47	Amiodarone hydrochloride	Antiparasitics	81
ADT Booster	215	Amisulpride	Antipruritic Preparations	52
Adult diphtheria and tetanus		Amitriptyline	Antipsychotic Agents	121
vaccine	215	Amlodipine	Antiretrovirals	83
Advantan	54	Amorolfine	Antirheumatoid Agents	95
Advate	31	Amoxicillin	Antiseptics and	
Aerrane	106	Amoxicillin with clavulanic	Disinfectants	192
Afinitor	175	acid	Antispasmodics and Other	
AFT SLS-free	53	Amphotericin B	Agents Altering Gut	

Motility	15	Argipressin [Vasopressin]	69	Ativan	126
Antithrombotics	32	Aripiprazole	122	Atomoxetine	128
Antithymocyte globulin (equine)	175	Aristocort	54	Atorvastatin	45
Antithymocyte globulin (rabbit)	175	Aromasin	146	Atovaquone with proguanil hydrochloride	82
Antitumorants	15	Arrow - Clopid	34	Atracurium besylate	102
Antivirals	87	Arrow-Amitriptyline	114	Atripila	85
Anxiolytics	125	Arrow-Bendrofluazide	45	Atropine sulphate Cardiovascular	41
Apidra	17	Arrow-Brimonidine	188	Sensory	188
Apidra Solostar	17	Arrow-Calcium	21	Atropt	188
Apo-Allopurinol	100	Arrow-Diazepam	126	Aubagio	126
Apo-Amloride	45	Arrow-Dortim	187	Augmentin	74
Apo-Amlodipine	43	Arrow-Etidronate	97	Auranofin	95
Apo-Amoxi	74	Arrow-Fluoxetine	115	Avelox	75
Apo-Azithromycin	73	Arrow-Gabapentin	116	Avelox IV 400	75
Apo-Ciclopirox	51	Arrow-Iloprost	50	Avonex	127
Apo-Cilazapril/ Hydrochlorothiazide	39	Arrow-Lamotrigine	118	Avonex Pen	127
Apo-Clarithromycin	73	Arrow-Losartan & Hydrochlorothiazide	40	Azacitidine	134
Apo-Clomipramine	114	Arrow-Morphine LA	112	Azactam	76
Apo-Diclo SR	103	Arrow-Norfloxacin	75	Azamun	175
Apo-Diltiazem CD	43	Arrow-Ornidazole	82	Azathioprine	175
Apo-Doxazosin	40	Arrow-Quinapril 10	39	Azithromycin	73
Apo-Folic Acid	28	Arrow-Quinapril 20	39	Azol	64
Apo-Imiquimod Cream 5%	56	Arrow-Quinapril 5	39	AZT	85
Apo-Megestrol	144	Arrow-Roxithromycin	74	Aztreonam	76
Apo-Mirtazapine	114	Arrow-Sertraline	115		
Apo-Moclobemide	114	Arrow-Simva	46	- B -	
Apo-Nadolol	42	Arrow-Sumatriptan	120	B-D Micro-Fine	222
Apo-Nicotinic Acid	46	Arrow-Timolol	187	B-D Ultra Fine	222
Apo-Oxybutynin	60	Arrow-Tolterodine	60	B-D Ultra Fine II	222
Apo-Perindopril	39	Arrow-Topiramate	119	Bacillus calmette-guerin (BCG)	175
Apo-Pindolol	42	Arrow-Tramadol	113	Bacillus calmette-guerin vaccine	216
Apo-Prazosin	40	Arrow-Venlafaxine XR	115	Baclofen	102
Apo-Prednisone	62	Arsenic trioxide	135	Bacterial and Viral Vaccines	215
Apo-Prednisone S29	62	Artemether with lumefantrine	81	Bacterial Vaccines	215
Apo-Propranolol	42	Artesunate	82	Balanced Salt Solution	186
Apo-Pyridoxine	25	Articaine hydrochloride	107	Baraclude	88
Apo-Ropinirole	106	Articaine hydrochloride with adrenaline	107	Barium sulphate	194
Apomine	105	Asacol	14	Barium sulphate with sodium bicarbonate	194
Apomorphine hydrochloride	105	Asamax	14	Barrier Creams and Emollients	52
Apraclonidine	188	Ascorbic acid Alimentary	25	Basiliximab	159
Aprepitant	120	Extemporaneous	198	BCG Vaccine	216
Apresoline	48	Aspen Adrenaline	47	BD PosiFlush	37
Aprotinin	29	Aspirin Blood	34	Beclazone 100	180
Aqueous cream	53	Nervous	109	Beclazone 250	180
Arachis oil [Peanut oil]	198	Asthalin	180	Beclazone 50	180
Arava	95	Atazanavir sulphate	86	Beclomethasone dipropionate	177, 180
Aremed	145	Atenolol	41	Bee venom	177
Arginine Alimentary	20	Atenolol-AFT	41		
Various	195	ATGAM	175		

Bendrofluazide	45	Biliscopin	195	naloxone	130
Bendroflumethiazide [Bendrofluazide]	45	Bimatoprost	188	Bupropion hydrochloride	130
BeneFIX	30	Biodone	111	Burinex	44
Benzathine benzylpenicillin	74	Biodone Extra Forte	111	Buscopan	15
Benzbromaron AL 100	100	Biodone Forte	111	Buserelin	64
Benzbromarone	100	Biotin	21	Buspirone hydrochloride	125
Benzocaine	107	Bisacodyl	20	Busulfan	133
Benzoin	198	Bismuth subgallate	198	Butacort Aqueous	178
Benzoyl peroxide	52	Bismuth subnitrate and iodoform paraffin	196	- C -	
Benztrop	105	Bismuth trioxide	16		
Benztropine mesylate	105	Bisoprolol fumarate	42	Cabergoline	63
Benzydamine hydrochloride	23	Bivalirudin	32	Caffeine	128
Benzydamine hydrochloride with cetylpyridinium chloride	23	Bleomycin sulphate	133	Caffeine citrate	182
Benzylpenicillin sodium [Penicillin G]	74	Blood glucose diagnostic test meter	222	Calamine	52
Beractant	183	Blood glucose diagnostic test strip	222	Calcipotriol	55
Beta Cream	54	Blood ketone diagnostic test meter	222	Calcitonin	61
Beta Ointment	54	Boceprevir	90	Calcitriol	26
Beta Scalp	55	Bonney's blue dye	195	Calcitriol-AFT	26
Beta-Adrenoceptor Agonists	180	Boostrix	216	Calcium carbonate	13, 21
Beta-Adrenoceptor Blockers	41	Boric acid	198	Calcium Channel Blockers	43
Betadine	192	Bortezomib	135	Calcium chloride	35
Betadine Skin Prep	192	Bosentan	49	Calcium folinate	143
Betagan	187	Bosvate	42	Calcium Folate Ebewe	143
Betahistine dihydrochloride	120	Botox	102	Calcium gluconate Blood	35
Betaine	20	Botulism antitoxin	190	Dermatological	56
Betamethasone	61	Breo Ellipta	182	Calcium Homeostasis	61
Betamethasone dipropionate	54	Bridion	102	Calcium polystyrene sulphonate	38
Betamethasone dipropionate with calcipotriol	55	Brilinta	34	Calcium Resonium	38
Betamethasone sodium phosphate with betamethasone acetate	61	Brimonidine tartrate	188	Calsource	21
Betamethasone valerate	54–55	Brimonidine tartrate with timolol	188	Cancidas	79
Betamethasone valerate with clioquinol	55	Brinzolamide	187	Candesartan cilexetil	40
Betamethasone valerate with fusidic acid	55	Bromocriptine	105	Candestar	40
Betaxolol	187	Brufen SR	103	Capecitabine	134
Betoptic	187	Budesonide Alimentary	13	Capecitabine Winthrop	134
Betoptic S	187	Respiratory	178, 181	Capoten	39
Bevacizumab	159	Budesonide with eformoterol	182	Capsaicin Musculoskeletal	104
Bezafibrate	45	Bumetanide	44	Nervous	109
Bezalip	45	Bupafen	108	Captopril	39
Bezalip Retard	45	Bupivacaine hydrochloride	107	Carbamazepine	116
Bicalaccord	144	Bupivacaine hydrochloride with adrenaline	108	Carbasorb-X	191
Bicalutamide	144	Bupivacaine hydrochloride with fentanyl	108	Carbimazole	69
Bicillin LA	74	Bupivacaine hydrochloride with glucose	108	Carbomer	189
BicNU	133	Buprenorphine with		Carboplatin	138
Bile and Liver Therapy	16			Carboprost trometamol	58
				Carboxymethylcellulose Alimentary	23
				Extemporaneous	198
				Cardinol LA	42
				Cardizem CD	43
				CareSens	222
				Caresens II	222

CareSens N	222	Chlorhexidine with ethanol	192	Clobetasol propionate	54, 56
Caresens N	222	Chloroform	198	Clobetasone butyrate	54
Caresens N POP	222	Chloroquine phosphate	82	Clofazimine	80
Carmellose sodium	189	Chlorothiazide	45	Clomazol	51, 57
Carmustine	133	Chlorpheniramine maleate	178	Clomiphene citrate	63
Carvedilol	42	Chlorpromazine		Clomipramine hydrochloride	114
Caspofungin	79	hydrochloride	122	Clonazepam	115–116, 125
Catapres	44	Chlorsig	184	Clonidine	44
Catapres-TTS-1	44	Chlortalidone		Clonidine BNM	44
Catapres-TTS-2	44	[Chlorthalidone]	45	Clonidine hydrochloride	44
Catapres-TTS-3	44	Chlorthalidone	45	Clopidogrel	34
Ceenu	133	Cholecalciferol	26	Clopine	122
Cefaclor	72	Cholestyramine	46	Clopixol	124, 125
Cefalexin	72	Choline salicylate with		Clostridium botulinum type A	
Cefalexin Sandoz	72	cetalkonium chloride	23	toxin	102
Cefazolin	72	Cholvastin	46	Clotrimazole	
Cefepime	72	Choriogonadotropin alfa	65	Dermatological	51
Cefepime-AFT	72	Ciclopirox olamine	51	Genito-Urinary	57
Cefotaxime	72	Ciclosporin	146	Clove oil	198
Cefotaxime Sandoz	72	Cidofovir	91	Clozapine	122
Cefoxitin	72	Cilazapril	39	Clozaril	122
Ceftaroline fosamil	73	Cilazapril with		Co-trimoxazole	77
Ceftazidime	72	hydrochlorothiazide	39	Coal tar	198
Ceftriaxone	72	Cilicaine	74	Coal tar with salicylic acid and	
Ceftriaxone-AFT	72	Cilicaine VK	74	sulphur	55
Cefuroxime	72	Cimetidine	15	Cocaine hydrochloride	108
Celecoxib	103	Cinchocaine hydrochloride with		Cocaine hydrochloride with	
Celiprolol	42	hydrocortisone	14	adrenaline	108
CellCept	175	Cipflox	75	Codeine phosphate	
Celol	42	Ciprofloxacin		Extemporaneous	198
Centrally-Acting Agents	44	Infection	75	Nervous	110
Cephalexin ABM	72	Sensory	184	Cogentin	105
Cetirizine hydrochloride	178	Ciprofloxacin with		Colaspase [L-asparaginase]	136
Cetomacrogol	53	hydrocortisone	184	Colchicine	101
Cetomacrogol with glycerol	53	Ciproxin HC Otic	184	Colestimethate	76
Cetrimide	198	Cisplatin	138	Colestipol hydrochloride	46
Champix	131	Citalopram hydrobromide	115	Colgout	101
Charcoal	191	Citanest	109	Colifoam	14
Chemotherapeutic Agents	133	Citric acid	198	Colistin sulphomethate	
Chicken pox vaccine	221	Citric acid with magnesium oxide		[Colestimethate]	76
Chlorafast	184	and sodium picosulfate	19	Colistin-Link	76
Chloral hydrate	127	Citric acid with sodium		Collodion flexible	198
Chlorambucil	133	bicarbonate	194	Colofac	15
Chloramphenicol		Cladribine	134	Colony-Stimulating Factors	35
Infection	76	Clarithromycin	73	Coloxyl	19, 20
Sensory	184	Clexane	32	Compound electrolytes	35, 38
Chlorhexidine	192, 196	Clindamycin	76	Compound electrolytes with	
Chlorhexidine gluconate		Clindamycin ABM	76	glucose	35, 38
Alimentary	23	Clinicians Multivit & Mineral		Compound	
Extemporaneous	198	Boost	24	hydroxybenzoate	198
Genito-Urinary	57	Clinicians Renal Vit	24	Compound sodium lactate	
Chlorhexidine with		Clobazam	116	[Hartmann's solution]	36
cetrimide	192, 196	Clobetasol BNM	54	Compound sodium lactate with	

glucose	36	Dapa-Tabs	45	Sensory	185
Concerta	129	Dapsone		Dexamethasone phosphate	62
Condyline	56	Contracted	80	Dexamethasone with framycetin and gramicidin	184
Contraceptives	57	Infection	80	Dexamethasone with neomycin sulphate and polymyxin B sulphate	185
Contrast Media	193	Daptomycin	76	Dexamethasone with tobramycin	185
Cordarone-X	41	Darunavir	86	Dexamfetamine sulfate	128
Corticosteroids		Dasatinib	138	Dexmedetomidine	106
Dermatological	54	Daunorubicin	133	Dexmethsone	62
Hormone	61	DBL Acetylcysteine	190	Dextrose	
Corticotrorelin (ovine)	64	DBL Amikacin	71	Alimentary	16
Cosmegen	133	DBL Aminophylline	182	Blood	36
Cough Suppressants	180	DBL Bleomycin Sulfate	133	Extemporaneous	198
Creon 10000	18	DBL Carboplatin	138	Dextrose with sodium citrate and citric acid [Acid Citrate Dextrose A]	32
Creon 25000	18	DBL Cefotaxime	72	DHC Continus	110
Crotamiton	52	DBL Cisplatin	138	Diabetes	16
Crystaderm	51	DBL Docetaxel	143	Diacomit	118
CT Plus+	194	DBL Ergometrine	59	Diagnostic Agents	195
Cubicin	76	DBL Leucovorin Calcium	143	Diagnostic and Surgical Preparations	186
Curam Duo	74	DBL Meropenem	72	Diamide Relief	13
Curosurf	183	DBL Morphine Sulphate	112	Diamox	187
Cvite	25	DBL Pethidine		Diatrizoate meglumine with sodium amidotrizoate	193
Cyclizine hydrochloride	120	Hydrochloride	113	Diatrizoate sodium	193
Cyclizine lactate	120	DBL Rocuronium Bromide	102	Diazepam	115, 126
Cyclogyl	188	DBL Sterile Dopamine Concentrate	47	Diazoxide	
Cyclopentolate hydrochloride	188	DBL Tobramycin	71	Alimentary	16
Cyclophosphamide	133	DDI	85	Cardiovascular	48
Cycloserine	80	De-Nol	16	Dicarz	42
Cyklokapron	30	De-Worm	81	Dichlorobenzyl alcohol with amylmetacresol	23
Cymevene	91	Decongestants	180	Diclofenac Sandoz	103
Cyproheptadine hydrochloride	178	Decongestants and Antiallergics	185	Diclofenac sodium Musculoskeletal	103
Cyproterone acetate	61	Decozol	23	Sensory	185
Cyproterone acetate with ethinyloestradiol	57	Deferasirox	191	Dicobalt edetate	191
Cysteamine hydrochloride	198	Deferiprone	191	Didanosine [DDI]	85
Cytarabine	134	Defibrotide	32	Diflucan	78
Cytotec	15	Definity	195	Diflucortolone valerate	54
		Demeclocycline hydrochloride	76	Digestives Including Enzymes	18
- D -		Deoxycofornycin	137	Digoxin	41
D-Penamine	95	Depo-Medrol	62	Digoxin immune Fab	190
Dabigatran	32	Depo-Medrol with Lidocaine	62	Dihydrocodeine tartrate	110
Dacarbazine	136	Depo-Provera	58	Dihydroergotamine mesylate	120
Dactinomycin [Actinomycin D]	133	Depo-Testosterone	61	Diltiazem hydrochloride	43
Daivobet	55	Deprim	77		
Daivonex	55	Dermol	56		
Dalacin C	76	Desferal	191		
Dalteparin	32	Desferrioxamine mesilate	191		
Danaparoid	32	Desflurane	106		
Danazol	64	Desmopressin acetate	69		
Dantrium	102	Desmopressin-PH&T	69		
Dantrium IV	102	Dexamethasone Hormone	62		
Dantrolene	102				

Dilzem	43	Doxine	76	Enalapril maleate	39
Dimercaprol	191	Doxorubicin Ebewe	133	Enalapril maleate with hydrochlorothiazide	39
Dimercaptosuccinic acid	191	Doxorubicin hydrochloride	133	Enbrel	146
Dimethicone	52	Doxycycline	76	Endocrine Therapy	144
Dimethyl fumarate	126	DP Fusidic Acid Cream	51	Endoxan	133
Dimethyl sulfoxide	196	DP Lotn HC	54	Enfuvirtide	83
Dinoprostone	59	DP-Anastrozole	145	Enoxaparin	32
Diphemanil metilsulfate	56	Dr Reddy's Omeprazole	16	Ensure (Chocolate)	214
Diphenoxylate hydrochloride with atropine sulphate	13	Dr Reddy's Terbinafine	80	Ensure (Vanilla)	214
Diphtheria antitoxin	190	Droperidol	120	Ensure Plus (Banana)	214
Diphtheria, tetanus and pertussis vaccine	216	Drugs Affecting Bone Metabolism	95	Ensure Plus (Chocolate)	214
Diphtheria, tetanus, pertussis and polio vaccine	215	Duolin	178	Ensure Plus (Fruit of the Forest)	214
Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine	215	Duovisc	187	Ensure Plus (Vanilla)	214
Diprivan	107	Duride	47	Ensure Plus HN	213
Dipyridamole	34	Dynastat	104	Ensure Plus HN RTH	213
Disodium edetate	187	Dysport	102	Entacapone	105
Disodium hydrogen phosphate with sodium dihydrogen phosphate	198	- E -		Entapone	105
Disopyramide phosphate	41	e-chamber La Grande	223	Entecavir	88
Disulfiram	130	e-chamber Mask	222	Enzymes	100
Dithranol	198	e-chamber Turbo	223	Ephedrine	47
Diuretics	44	E-Mycin	73	Epilim IV	118
Diurin 40	44	E-Z-Cat Dry	194	Epirubicin Ebewe	134
Dobutamine hydrochloride	47	E-Z-Gas II	194	Epirubicin hydrochloride	134
Dobutamine-Claris	47	E-Z-Paste	194	Epoetin alfa [Erythropoietin alfa]	27
Docetaxel	143	EasyCheck	222	Epoetin beta [Erythropoietin beta]	28
Docusate sodium Alimentary	19	Econazole nitrate	51	Epoprostenol	49
Sensory	189	Edrophonium chloride	95	Eprex	27
Docusate sodium with sennosides	19	Efavirenz	84	Eptacog alfa [Recombinant factor Vlla]	30
Domperidone	120	Efavirenz with emtricitabine and tenofovir disoproxil fumarate	85	Eptifibatide	34
Donepezil hydrochloride	130	Efexor XR	115	Ergometrine maleate	59
Donepezil-Rex	130	Effient	34	Ergotamine tartrate with caffeine	120
Dopamine hydrochloride	47	Eformoterol fumarate	181	Erlotinib	139
Dopergin	106	Efudix	56	Ertapenem	71
Dopress	114	Elecare (Unflavoured)	209	Erythrocin IV	73
Dornase alfa	182	Elecare (Vanilla)	209	Erythromycin (as ethylsuccinate)	73
Dorzolamide	187	Elecare LCP (Unflavoured)	209	Erythromycin (as lactobionate)	73
Dorzolamide with timolol	187	Electrolytes	197	Erythromycin (as stearate)	74
Dostinex	63	Eligard	65	Erythropoietin alfa	27
Dotarem	194	Elocon	54	Erythropoietin beta	27
Dothiepin hydrochloride	114	Elocon Alcohol Free	54	Escitalopram	115
Doxapram	183	Eltrombopag	29	Esmolol hydrochloride	42
Doxazosin	40	Emend Tri-Pack	120	Etanercept	146
Doxepin hydrochloride	114	EMLA	109	Ethambutol hydrochloride	80
		Emtricitabine	85	Ethanol	190
		Emtricitabine with tenofovir disoproxil fumarate	85	Ethanol with glucose	190
		Emtriva	85		
		Emulsifying ointment	53		

Ethanol, dehydrated	190	Ferrous sulphate	22	Flutamin	144
Ethics Aspirin EC	34	Ferrous sulphate with ascorbic acid	22	Fluticasone	181
Ethics Enalapril	39	Ferrous sulphate with folic acid	22	Fluticasone furoate with vilanterol	182
Ethics Lisinopril	39	Ferrum H	22	Fluticasone propionate	178
Ethinylestradiol	64	Fexofenadine hydrochloride	178	Fluticasone with salmeterol	182
Ethinylestradiol with desogestrel	57	Filgrastim	35	FML	185
Ethinylestradiol with levonorgestrel	57	Finasteride	59	Foban	51
Ethinylestradiol with norethisterone	57	Fingolimod	126	Folic acid	28
Ethosuximide	116	Finpro	59	Fondaparinux sodium	33
Ethyl chloride	108	Firazyr	177	Food Modules	201
Etidronate disodium	97	Flagyl	82	Food/Fluid Thickeners	203
Etomidate	106	Flagyl-S	82	Forteo	100
Etopophos	136	Flamazine	51	Fortisip (Vanilla)	214
Etoposide	136	Flecainide acetate	41	Fortum	72
Etoposide (as phosphate)	136	Fleet Phosphate Enema	20	Fosamax	95, 96
Etoricoxib	103	Flixonase Hayfever & Allergy	178	Fosamax Plus	96
Etravirine	84	Flixotide	181	Foscarnet sodium	91
Everet	118	Flixotide Accuhaler	181	Fosfomycin	76
Everolimus	175	Floair	181	Fragmin	32
Evista	99	Florinef	62	Framycetin sulphate	184
Exelon	130	Fluanxol	124	Freeflex	37
Exemestane	146	Fluarix	219	FreeStyle Lite	222
Exjade	191	Flucloxacillin	74	Freestyle Optium	222
Extemporaneously Compounded Preparations	198	Flucloxin	74	Freestyle Optium Ketone	222
Ezemibe	46	Fluconazole	78	Freestyle Optium Neo	222
Ezetimibe	46	Fluconazole-Claris	78	Fresofol 1%	107
Ezetimibe with simvastatin	46	Flucytosine	80	Fresofol 1% MCT/LCT	107
- F -		Fludara Oral	135	frusemide	44
Factor eight inhibitor bypassing fraction	30	Fludarabine Ebewe	135	Frusemide-Claris	44
Febuxostat	101	Fludarabine phosphate	135	Fucidin	77
FEIBA NF	30	Fludrocortisone acetate	62	Fucithalmic	184
Felodipine	43	Fluids and Electrolytes	35	Fungilin	23
Fenpaed	103	Flumazenil	190	Furosemide [frusemide]	44
Fentanyl	111	Flumetasone pivalate with clioquinol	185	Fusidic acid	
Fentanyl Sandoz	111	Fluocortolone caproate with fluocortolone pivalate and cinchocaine	14	Dermatological	51
Ferinject	22	Fluorescein sodium	186	Infection	77
Ferodan	22	Fluorescein sodium with lignocaine hydrochloride	186	Sensory	184
Ferric carboxymaltose	22	Fluorescite	186	Fuzeon	83
Ferric subsulfate	29	Fluorometholone	185	- G -	
Ferriprox	191	Fluorouracil	135	Gabapentin	116
Ferro-F-Tabs	22	Fluorouracil Ebewe	135	Gacet	110
Ferro-tab	22	Fluorouracil sodium	56	Gadobenic acid	194
Ferrograd	22	Fluoxetine hydrochloride	115	Gadobutrol	194
Ferrous fumarate	22	Flupenthixol decanoate	124	Gadodiamide	194
Ferrous fumarate with folic acid	22	Fluphenazine decanoate	124	Gadoteric acid	194
Ferrous gluconate with ascorbic acid	22	Flutamide	144	Gadovist	194
				Gadoxetate disodium	195
				Ganciclovir	
				Infection	91
				Sensory	184
				Gardasil	219
				Gastrografin	193

Gastrosoothe	15	indacaterol	179	Hydralazine hydrochloride	48
Gefitinib	139	Glypressin	70	Hydrea	136
Gelatine, succinylated	38	Glytrin	47	Hydrocortisone	
Gelofusine	38	Gonadorelin	64	Dermatological	54
Gemcitabine	135	Goserelin	64	Extemporaneous	199
Gemcitabine Ebewe	135	Granirex	121	Hormone	62
Gemfibrozil	45	Granisetron	121	Hydrocortisone acetate	
Genoptic	184			Alimentary	14
Genox	145			Dermatological	54
Gentamicin sulphate				Hydrocortisone and paraffin	
Infection	71			liquid and lanolin	54
Sensory	184			Hydrocortisone butyrate	54, 56
Gestrinone	64			Hydrocortisone with	
Gilenya	126			miconazole	55
Ginet	57			Hydrocortisone with natamycin	
Glatiramer acetate	126			and neomycin	55
Glaucoma Preparations	187			Hydrocortisone with paraffin and	
Glibenclamide	18			wool fat	54
Gliclazide	18			Hydrogen peroxide	51
Glipizide	18			Hydroxocobalamin	
Glivec	139			Alimentary	25
Glizide	18			Various	190
Glucagen Hypokit	16			Hydroxychloroquine	95
Glucagon hydrochloride	16			Hydroxyurea	136
Glucerna Select (Vanilla)	206			Hygroton	45
Glucerna Select RTH				Hylo-Fresh	189
(Vanilla)	206			Hyoscine butylbromide	15
Glucobay	16			Hyoscine hydrobromide	121
Glucose [Dextrose]				Hyperuricaemia and	
Alimentary	17			Antigout	100
Blood	36			Hypnovel	127
Extemporaneous	199			Hypromellose	186, 189
Glucose with potassium				Hypromellose with dextran	189
chloride	36			Hysite	188
Glucose with potassium chloride					
and sodium chloride	36				
Glucose with sodium chloride	36				
Glucose with sucrose and					
fructose	17				
Glycerin with sodium					
saccharin	199				
Glycerin with sucrose	199				
Glycerol					
Alimentary	20				
Extemporaneous	199				
Glycerol with paraffin	53				
Glyceryl trinitrate					
Alimentary	15				
Cardiovascular	47				
Glycine	196				
Glycopyrronium	179				
Glycopyrronium bromide	15				
Glycopyrronium with					

- H -

- I -

(Chocolate)	213	Iron polymaltose	22	Labetalol	42
Impact Advanced Recovery (Vanilla)	213	Iron sucrose	22	Lacosamide	117
Imuran	175	Irrigation Solutions	196	Lactose	199
Incruse Ellipta	179	Isentress	87	Lactulose	20
Indacaterol	181	Ismo-20	47	Laevolac	20
Indapamide	45	Isoflurane	106	Lamictal	118
Indigo carmine	195	Isoniazid	80	Lamivudine	85, 88
Indinavir	86	Isoniazid with rifampicin	80	Lamotrigine	118
Indocyanine green	195	Isoprenaline	47	Lanoxin	41
Indomethacin	104	Isopropyl alcohol	192	Lanoxin PG	41
Infanrix IPV	215	Isoptin	44	Lansoprazole	15
Infanrix-hexa	215	Isopto Carpine	188	Lantus	17
Infliximab	159	Isosorbide mononitrate	47	Lantus SoloStar	17
Influenza vaccine	219	Isotane 10	52	Lanzol Relief	15
Influvac	219	Isotane 20	52	Lapatinib	140
Inhaled Corticosteroids	180	Isotretinoin	52	Lariam	82
Insulin aspart	17	Ispaghula (psyllium) husk	19	Latanoprost	188
Insulin aspart with insulin aspart protamine	17	Isradipine	43	Lax-Sachets	20
Insulin glargine	17	Itch-Soothe	52	Lax-Suppositories	20
Insulin glulisine	17	Itraconazole	78	Lax-Tabs	20
Insulin isophane	17	Itrazole	78	Laxatives	19
Insulin lispro	17	Ivermectin	81	Laxsol	19
Insulin lispro with insulin lispro protamine	17	- J -			95
Insulin neutral	18	Jadelle	58	Leclunomide	136
Insulin neutral with insulin isophane	17	Jaychem	53	Letrole	146
Insulin pen needles	222	Jevity	214	Letrozole	146
Insulin syringes, disposable with attached needle	222	Jevity HiCal RTH	213	Leukotriene Receptor Antagonists	181
Integrilin	34	Jevity RTH	214	Leunase	136
Intelence	84	- K -			65
Interferon alfa-2a	92	Kaletra	86	Leuprorelin acetate	134
Interferon alfa-2b	92	Kenacomb	185	Leustatin	118
Interferon beta-1-alpha	127	Kenacort-A 10	62	Levetiracetam	118
Interferon beta-1-beta	127	Kenacort-A 40	62	Levetiracetam-Rex	118
Interferon gamma	92	Kenalog in Orabase	23	Levobunolol hydrochloride	187
Intra-uterine device	57	Ketamine	106	Levocabastine	185
Invanz	71	Ketamine-Clarix	106	Levocarnitine	21
Invega Sustenna	124	Ketoconazole	51	Levodopa with benserazide	106
Iodine	69	Dermatological	51	Levodopa with carbidopa	106
Iodine with ethanol	192	Infection	78	Levomepromazine	123
Iodised oil	193	Ketone blood beta-ketone electrodes	222	Levonorgestrel	58
Iodixanol	193	Ketoprofen	104	Levosimendan	47
Iohexol	193	Ketorolac trometamol	185	Levothyroxine	69
Iopidine	188	Kivexa	85	Lidocaine [lignocaine]	109
Ioscan	193	Klacid	73	Lidocaine [Lignocaine] hydrochloride	108
IPOL	220	Klean Prep	19	Lidocaine [Lignocaine] hydrochloride with adrenaline	108
Ipratropium bromide	178	Kogenate FS	31	Lidocaine [Lignocaine] hydrochloride with adrenaline and tetracaine	109
Iressa	139	Konakion MM	31	Lidocaine [Lignocaine] hydrochloride	109
Irinotecan hydrochloride	136	Konsyl-D	19	Lidocaine [Lignocaine]	
		- L -			
		L-asparaginase	136		
		L-ornithine L-aspartate	16		

hydrochloride with chlorhexidine 109	Loxamine115	Mefloquine82
Lidocaine [Lignocaine] hydrochloride with phenylephrine hydrochloride 109	Lucrin Depot PDS65	Megestrol acetate144
Lidocaine [Lignocaine] with prilocaine 109	Lycinate47	Meglumine gadopentetate195
Lidocaine-Claris108	Lyderm52	Meglumine iotroxate195
Lignocaine108, 109	- M -	Melatonin127
lignocaine	m-Amoxiclav74	Meloxicam104
Hormone62	m-Eslon112	Melphalan133
Nervous109	M-M-R-II220	Menactra216
Lincomycin77	m-Nystatin23	Meningococcal (A, C, Y and W-135) conjugate vaccine 216
Linezolid77	Mabthera165	Meningococcal C conjugate vaccine 217
Lioresal Intrathecal102	Madopar 125106	Menthol199
Liothyronine sodium69	Madopar 250106	Mepivacaine hydrochloride109
Lipazil45	Madopar 62.5106	Mercaptopurine135
Lipid-Modifying Agents45	Madopar HBS106	Meropenem72
Lipodol Ultra Fluid193	Madopar Rapid106	Mesalazine14
Liquibar194	Maferide acetate51	Mesna143
Liquifilm Forte189	Magnesium hydroxide	Mestinon95
Liquifilm Tears189	Alimentary22	Metabolic Disorder Agents20
Lisinopril39	Extemporaneous199	Metabolic Products203
Lissamine green186	Magnesium oxide22	Metamide121
Lisuride hydrogen maleate106	Magnesium sulphate22	Metaraminol48
Lithicarb FC123	Magnevist195	Metchek18
Lithium carbonate123	Malarone82	Meterol182
LMX4109	Malarone Junior82	Metformin hydrochloride18
Local Preparations for Anal and Rectal Disorders 14	Malathion [Maldison]51	Methacholine chloride195
Locoid54, 56	Malathion with permethrin and piperonyl butoxide 52	Methadone hydrochloride Extemporaneous199
Locoid Crelo54	Maldison51	Nervous111
Locoid Lipocream54	Mannitol	Methatabs111
Lodoxamide185	Cardiovascular44	Methohexital sodium106
Logem118	Various195	Methopt189
Lomide185	Maprotiline hydrochloride114	Methotrexate135
Lomustine133	Marcaïn107	Methotrexate Ebewe135
Long-Acting Beta-Adrenoceptor Agonists 181	Marcaïn Heavy108	Methotrexate Sandoz135
Loniten48	Marcaïn Isobaric107	Methoxsalen [8-methoxypsoralen] 55
Loperamide hydrochloride13	Marcaïn with Adrenaline108	Methoxyflurane110
Lopinavir with ritonavir86	Marevan33	Methyl aminolevulinate hydrochloride 56
Lopresor42	Marine Blue Lotion SPF 50+56	Methyl hydroxybenzoate199
Lorafix178	Mask for spacer device222	Methylcellulose199
LoraPaed178	Mast Cell Stabilisers182	Methylcellulose with glycerin and sodium saccharin 199
Loratadine178	Max Health15, 47, 62, 95	Methylcellulose with glycerin and sucrose 199
Lorazepam116, 126	Maxidex185	Methyldopa44
Lormetazepam127	Maxitrol185	Methylene blue195
Losartan potassium40	Measles, mumps and rubella vaccine 220	Methylphenidate hydrochloride 129
Losartan potassium with hydrochlorothiazide 40	Mebendazole81	Methylprednisolone (as sodium
Lovir91	Mebeverine hydrochloride15	
	Medrol62	
	Medroxyprogesterone64	
	Medroxyprogesterone acetate Genito-Urinary58	
	Hormone63	
	Mefenamic acid104	

succinate) 62	Moclobemide 114	Navelbine 143
Methylprednisolone	Modafinil 129	Nedocromil 182
aceponate 54	Moderate 124	Nefopam hydrochloride 110
Methylprednisolone acetate 62	Mometasone furoate 54	Neisvac-C 217
Methylprednisolone acetate with	Monosodium glutamate with	Neo-B12 25
lidocaine [lignocaine] 62	sodium aspartate 197	Neocate Advance (Vanilla) 209
Methylthioninium chloride	Monosodium l-aspartate 197	Neocate Gold (Unflavoured) 209
[Methylene blue] 195	Montelukast 181	Neoral 146
Methylxanthines 182	Moroctocog alfa [Recombinant	Neostigmine metilsulfate 95
Metoclopramide	factor VIII] 30	Neostigmine metilsulfate with
hydrochloride 121	Morphine hydrochloride 111	glycopyrronium bromide 95
Metoclopramide hydrochloride	Morphine sulphate 112	Neosynephrine HCL 48
with paracetamol 120	Morphine tartrate 112	Nepro HP (Strawberry) 212
Metolazone 45	Motetis 105	Nepro HP (Vanilla) 212
Metoprolol - AFT CR 42	Mouth and Throat 23	Nepro HP RTH 212
Metoprolol succinate 42	Moxifloxacin 75	Neulastim 35
Metoprolol tartrate 42	Mucolytics and	Neupogen 35
Metronidazole	Expectorants 182	Neurotin 116
Dermatological 51	Multihance 194	NeuroTabs 22
Infection 82	Multiple Sclerosis	Nevirapine 84
Metyrapone 64	Treatments 126	Nevirapine Alphapharm 84
Mexiletine hydrochloride 41	Multivitamin and mineral	Nicardipine hydrochloride 43
Mexiletine Hydrochloride	supplement 24	Nicorandil 48
USP 41	Multivitamin renal 24	Nicotine 131
Miacalcic 61	Multivitamins 24–25	Nicotinic acid 46
Mianserin hydrochloride 114	Mupirocin 51	Nifedipine 43
Micolette 20	Muscle Relaxants and Related	Nilotinib 140
Miconazole 23	Agents 102	Nilstat 78
Miconazole nitrate	Myambutol 80	Nimodipine 43
Dermatological 51	Mycobutin 81	Nitazoxanide 82
Genito-Urinary 57	Mycosail 51	Nitrados 127
Micreme 57	Mycophenolate mofetil 175	Nitrates 47
Micreme H 55	Mydriacyl 188	Nitrazepam 127
Midazolam 127	Mydriatics and Cycloplegics 188	Nitroderm TTS 10 47
Midodrine 41	Myran Atenolol 41	Nitroderm TTS 5 47
Mifepristone 58	Myran-Bosentan 49	Nitrofurantoin 77
Milrinone 48	Myleran 133	Nitrolingual Pump Spray 47
Minerals 21		Nitronal 47
Mini-Wright AFS Low Range 222		Noflam 250 104
Mini-Wright Standard 222		Noflam 500 104
Minidiab 18		Non-Steroidal Anti-Inflammatory
Minirin 69		Drugs 103
Minocycline 76		Nonacog alfa [Recombinant
Minoxidil 48		factor IX] 30
Mirtazapine 114		Nonacog gamma, [Recombinant
Misoprostol 15		factor IX] 31
Mitomycin C 134		Noradrenaline 48
Mitozantrone 134		Norethisterone
Mitozantrone Ebewe 134		Genito-Urinary 58
Mivacron 102		Hormone 64
Mivacurium chloride 102		Norethisterone with
Mixed salt solution for eye		mestranol 57
irrigation 186		Norfloroxacin 75

- N -

Noriday 28	58	Omezol Relief	15	Oxytocin BNM	59
Normison	127	Omnipaque	193	Oxytocin with ergometrine maleate	59
Norpress	114	Omniscan	194	Ozole	78
Nortriptyline hydrochloride	114	Omnitrope	65		
Norvir	86	On Call Advanced	222	- P -	
Novasource Renal (Vanilla)	212	Onbrez Breezhaler	181	Pacifen	102
Novatretin	55	Oncaspar	137	Pacific Bupirone	125
NovoRapid FlexPen	17	OncoTICE	175	Paclitaxel	143
NovoSeven RT	30	Ondanaccord	121	Paclitaxel Ebewe	143
Noxafil	78	Ondansetron	121	Paliperidone	124
Nupentin	116	Ondansetron ODT-DRLA	121	Pamidronate disodium	97
Nutrini Energy Multi Fibre	211	One-Alpha	25	Pamisol	97
Nutrini Low Energy Multifibre RTH	211	Onrex	121	Pancreatic enzyme	18
Nutrison Concentrated	208	Optional Pharmaceuticals	222	Pancuronium bromide	102
Nutrison Energy	213	Ora-Blend	199	Pantoprazole	16
Nyefax Retard	43	Ora-Blend SF	199	Papaverine hydrochloride	48
Nystatin		Ora-Plus	199	Paper wasp venom	177
Alimentary	23	Ora-Sweet	199	Para-aminosalicylic Acid	81
Dermatological	51	Ora-Sweet SF	199	Paracare	110
Genito-Urinary	57	Oratane	52	Paracare Double Strength	110
Infection	78	Ornidazole	82	Paracetamol	110
NZ Medical & Scientific	64	Orphenadrine citrate	102	Paracetamol + Codeine (Relieve)	113
		Oruvail SR	104	Paracetamol with codeine	113
- O -		Oseltamivir	92	Paraffin	
Obex Medical	195	Osmolite	214	Alimentary	19
Obstetric Preparations	58	Osmolite RTH	214	Dermatological	53
Octocog alfa [Recombinant factor VIII] (Advate)	31	Ospamox	74	Extemporaneous	199
Octocog alfa [Recombinant factor VIII] (Kogenate FS)	31	Other Cardiac Agents	47	Paraffin liquid with soft white paraffin	189
Octreotide	144	Other Endocrine Agents	63	Paraffin liquid with wool fat	189
Ocular Lubricants	189	Other Oestrogen Preparations	64	Paraffin with wool fat	53
Oestradiol	63-64	Other Otological Preparations	189	Paragesic Soluble	110
Oestradiol valerate	63	Other Progestogen Preparations	64	Paraldehyde	116
Oestradiol with norethisterone acetate	63	Other Skin Preparations	56	Parecoxib	104
Oestriol		Ox-Pam	126	Paromomycin	71
Genito-Urinary	59	Oxallicord	138	Paroxetine hydrochloride	115
Hormone	64	Oxaliplatin	138	Paser	81
Oestrogens	59	Oxandrolone	61	Patent blue V	195
Oestrogens (conjugated equine)	63	Oxazepam	126	Paxam	125
Oestrogens with medroxyprogesterone acetate	63	Oxpentifylline	48	Pazopanib	141
Oil in water emulsion	53	Oxybuprocaine hydrochloride	186	Peak flow meter	222
Oily phenol [Phenol oily]	15	Oxybutynin	60	Peanut oil	198
Olanzapine	123-124	Oxycodone Controlled Release Tablets (BNM)	113	Pediasure (Chocolate)	211
Olive oil	199	Oxycodone hydrochloride	113	Pediasure (Strawberry)	211
Olopatadine	185	OxyContin	113	Pediasure (Vanilla)	211
Olsalazine	14	Oxymetazoline hydrochloride	180	Pediasure RTH	211
Omalizumab	164	OxyNorm	113	Pegaspargase	137
Omeprazole	15-16	Oxytocin	59	Pegasys	93
				Pegasys RBV Combination Pack	93
				Pegfilgrastim	35
				Pegylated interferon alfa-2a	93

Penicillamine	95	Pipothiazine palmitate	125	Pravastatin	46
Penicillin G	74	Pituitary and Hypothalamic Hormones and Analogues	64	Praziquantel	81
Penicillin V	74	Pivmecillinam	77	Prazosin	40
Pentacarinat	82	Pizotifen	120	Precedex	106
Pentagastrin	64	PKU Anamix Junior LQ (Berry)	204	Prednisolone	62
Pentamidine isethionate	82	PKU Anamix Junior LQ (Orange)	204	Prednisolone acetate	185
Pentasa	14	PKU Anamix Junior LQ (Unflavoured)	204	Prednisolone sodium phosphate	185
Pentostatin [Deoxycoformycin]	137	Plaquenil	95	Prednisone	62
Pentoxifylline [Oxpentifylline]	49	Plendil ER	43	Pregnancy test - hCG urine	222
Peptisoothe	15	Pneumococcal (PCV13) conjugate vaccine	217	preOp	213
Perfalgan	110	Pneumococcal (PPV23) polysaccharide vaccine	217	Prevenar 13	217
Perflutren	195	Pneumovax 23	217	Prezista	86
Perhexiline maleate	44	Podophyllotoxin	56	Prilocaine hydrochloride	109
Pericyazine	123	Polidocanol	29	Prilocaine hydrochloride with felypressin	109
Perindopril	39	Poliomyelitis vaccine	220	Primaquine phosphate	82
Permethrin	52	Poloxamer	20	Primidone	118
Peteha	81	Poly Gel	189	Primolut N	64
Pethidine hydrochloride	113	Poly-Tears	189	Primovist	195
Pexsig	44	Poly-Visc	189	Probenecid	102
Phenelzine sulphate	114	Polyhexamethylene biguanide	199	Procaine penicillin	74
Phenindione	33	Polyvinyl alcohol	189	Procarbazine hydrochloride	137
Phenobarbitone	118, 127	Polyvinyl alcohol with povidone	189	Prochlorperazine	121
Phenobarbitone sodium	199	Poractant alfa	183	Proctosedyl	14
Phenol		Posaconazole	78	Procur	61
Extemporaneous	199	Postinor-1	58	Procyclidine hydrochloride	105
Various	196	Potassium chloride	36, 38	Procytox	133
Phenol oily	15	Potassium chloride with sodium chloride	37	Prodopa	44
Phenol with ioxaglic acid	196	Potassium citrate	60	Progesterone	59
Phenoxybenzamine hydrochloride	40	Potassium dihydrogen phosphate	37	Proglucem	16
Phenoxyethylpenicillin [Penicillin V]	74	Potassium iodate Alimentary	22	Proglycem	16
Phentolamine mesylate	40	Potassium iodate with iodine	22	Progynova	63
Phenylephrine hydrochloride Cardiovascular	48	Potassium perchlorate	69	Prokinex	120
Sensory	188	Potassium permanganate	55	Promethazine hydrochloride	178
Phenytoin	118	Povidone K30	199	Promethazine theoclate	121
Phenytoin sodium	116, 118	Povidone-iodine	192	Propafenone hydrochloride	41
Pholcodine	180	Povidone-iodine with ethanol	192	Propamidine isethionate	184
Phosphorus	38	Pradaxa	32	Propofol	107
Phytomenadione	31	Pralidoxime iodide	190	Propranolol	42
Picibanil	175	Pramipexole hydrochloride	106	Propylene glycol	199
Pilocarpine hydrochloride	188	Prasugrel	34	Propylthiouracil	69
Pilocarpine nitrate	199			Prostin E2	59
Pimafucort	55			Prostin VR	48
Pindolol	42			Protamine sulphate	33
Pine tar with trolamine laurilsulfate and fluorescein	55			Protonamide	81
Pinetarsol	55			Protirelin	69
Pioglitazone	18			Provera	63, 64
Piperacillin with tazobactam	74			Provisc	187

hydrochloride 180	Retinol 25	Salazopyrin 14
Psoriasis and Eczema	Retinol Palmitate 189	Salazopyrin EN 14
Preparations 55	Retrovir 85	Salbutamol 180
PTU 69	Retrovir IV 85	Salbutamol with ipratropium
Pulmocare (Vanilla) 212	Reutenox 104	bromide 178
Pulmonary Surfactants 183	Revlimid 136	Salicylic acid 200
Pulmozyme 182	Revolade 29	Salmeterol 182
Puri-nethol 135	RexAir 182	Salmonella typhi vaccine 218
Pyrazinamide 81	Reyataz 86	Sandimmun 146
Pyridostigmine bromide 95	Riboflavin 5-phosphate 187	Sandomigran 120
Pyridoxal-5-phosphate 21	Rifabutin 81	Sandostatin LAR 144
Pyridoxine hydrochloride 25	Rifadin 81	Scalp Preparations 55
Pyrimethamine 82	Rifampicin 81	Scandonest 3% 109
Pytazen SR 34	Rifaximin 16	Sclerosing Agents 183
- Q -	Rifinah 80	Scopoderm TTS 121
Q 300 83	Rilutek 105	Sebizole 51
Quetapel 123	Riluzole 105	Secretin pentahydrochloride 195
Quetiapine 123	Ringer's solution 37	Sedatives and Hypnotics 127
Quinapril 39	Riodine 192	Seebri Breezhaler 179
Quinapril with	Risedronate Sandoz 97	Selegiline hydrochloride 106
hydrochlorothiazide 39	Risedronate sodium 97	Sennosides 20
Quinine dihydrochloride 82	Risperdal Consta 125	Serenace 122
Quinine sulphate 83	Risperdal Quicklet 123	Seretide 182
Qvar 180	Risperidone 123, 125	Seretide Accuhaler 182
- R -	Risperon 123	Serevent 182
RA-Morph 111	Ritalin 129	Serevent Accuhaler 182
Rabies vaccine 220	Ritalin LA 129	Serophene 63
Raloxifene 99	Ritalin SR 129	Sertraline 115
Raltegravir potassium 87	Ritonavir 86	Sevoflurane 107
Ramipex 106	Rituximab 165	Sevredol 112
Ranbaxy-Cefaclor 72	Rivaroxaban 33	SII-Onco-BCG 175
Ranibizumab 165	Rivastigmine 130	Sildenafil 49
Ranitidine 15	Rivotril 115	Silver nitrate
Ranitidine Relief 15	RIXUBIS 31	Dermatological 56
Rapamune 175	Rizamelt 120	Extemporaneous 200
Rasburicase 102	Rizatriptan 120	Simethicone 13
Readi-CAT 2 194	Rocuronium bromide 102	Simlect 159
Reandron 1000 61	Ropinirole hydrochloride 106	Simvastatin 46
Recombinant factor IX 30	Ropivacaine hydrochloride 109	Sincalide 195
Recombinant factor VIIa 30	Ropivacaine hydrochloride with	Sinemet 106
Recombinant factor VIII 30, 31	fentanyl 109	Sinemet CR 106
Rectogesic 15	Ropivacaine Kabi 109	Singular 181
Red back spider antivenom 190	Rose bengal sodium 186	Sirolimus 175
Redipred 62	RotaTeg 220	Slow-Lopresor 42
Relenza Rotadisk 92	Rotavirus live reassortant oral	Snake antivenom 191
Remicade 159	vaccine 220	Sodibic 38
Remifentanyl hydrochloride 113	Roxane 13	Sodium acetate 37
ReoPro 152	Roxithromycin 74	Sodium acid phosphate 37
Resonium A 38	Rubifen 129	Sodium alginate with magnesium
Resource Beneprotein 202	Rubifen SR 129	alginate 13
Resource Diabetic (Vanilla) 206	- S -	Sodium alginate with sodium
Respiratory Stimulants 183	SalAir 180	bicarbonate and calcium
	Salamol 180	carbonate 13

Sodium aurothiomalate	95	Solifenacin succinate	60	Symmetrel	105
Sodium benzoate	21	Solu-Cortef	62	Sympathomimetics	47
Sodium bicarbonate		Solu-Medrol	62	Synacthen	64
Blood	37-38	Somatropin	65	Synacthen Depot	64
Extemporaneous	200	Sotacor	42	Syntometrine	59
Sodium calcium edetate	192	Sotalol	42	Syrup	200
Sodium carboxymethylcellulose		Soya oil	190	Systane Unit Dose	189
with pectin and gelatine	23	Spacer device	223		
Sodium chloride		Span-K	38	- T -	
Blood	37-38	Specialised Formulas	205	Tacrolimus	146
Respiratory	180, 183	Spilto Respimat	179	Tacrolimus Sandoz	146
Various	196	Spiractin	45	Tagitol V	194
Sodium chloride with sodium		Spiramycin	83	Talc	183
bicarbonate	180	Spiriva	179	Tambocor	41
Sodium citrate		Spiriva Respimat	179	Tambocor CR	41
Alimentary	13	Spirolactone	45	Tamoxifen citrate	145
Extemporaneous	200	Sprycel	138	Tamsulosin	60
Sodium citrate with sodium		Standard Feeds	213	Tamsulosin-Rex	60
chloride and potassium		Staphlex	74	Tarceva	139
chloride	33	Starch	200	Tasigna	140
Sodium citrate with sodium lauryl		Stavudine	85	Tasmar	106
sulphoacetate	20	Sterculia with frangula	19	Tecfidera	126
Sodium citro-tartrate	60	Stesolid	115	Tegretol	116
Sodium cromoglycate		Stimulants / ADHD		Tegretol CR	116
Alimentary	14	Treatments	128	Teicoplanin	77
Respiratory	178, 182	Stiripentol	118	Temaccord	137
Sensory	185	Stocrin	84	Temazepam	127
Sodium dihydrogen phosphate		Strattera	128	Temozolomide	137
[Sodium acid phosphate]	37	Streptomycin sulphate	71	Tenecteplase	35
Sodium fluoride	21	Stromectol	81	Tenofovir disoproxil fumarate	89
Sodium hyaluronate [Hyaluronic acid]		Suboxone	130	Tenoxicam	104
Alimentary	23	Sucralfate	16	Terazosin	40
Sensory	187, 189	Sucrose	110	Terbinafine	80
Sodium hyaluronate [Hyaluronic		Sugammadex	102	Terbutaline	59
acid] with chondroitin		Sulindac	104	Terbutaline sulphate	180
sulphate	187	Sulphacetamide sodium	184	Teriflunomide	126
Sodium hypochlorite	192	Sulphadiazine	77	Teriparatide	100
Sodium metabisulfite	200	Sulphadiazine silver	51	Terlipressin	70
Sodium nitrite	190	Sulphasalazine	14	Testosterone	61
Sodium nitroprusside		Sulphur	200	Testosterone cypionate	61
Cardiovascular	49	Sumatriptan	120	Testosterone esters	61
.....	223	Sunitinib	142	Testosterone undecanoate	61
Sodium phenylbutyrate	21	Sunscreen, proprietary	56	Tetrabenazine	105
Sodium phosphate with		Suprane	106	Tetracaine [Amethocaine]	
phosphoric acid	20	Surgical Preparations	196	hydrochloride	
Sodium polystyrene		Survanta	183	Nervous	109
sulphonate	38	Sustagen Diabetic (Vanilla)	206	Sensory	186
Sodium stibogluconate	83	Sustagen Hospital Formula		Tetracosactide	
Sodium tetradecyl sulphate	29	(Chocolate)	214	[Tetracosactrin]	64
Sodium thiosulfate	190	Sustagen Hospital Formula		Tetracosactrin	64
Sodium valproate	118	(Vanilla)	214	Tetracyclin Wolff	76
Sodium with potassium	197	Sutant	142	Tetracycline	76
Solian	121	Suxamethonium chloride	102	Thalidomide	138
				Thalomid	138

Theobroma oil	200	Dependence	130	Urografin	193
Theophylline	182	Tretinoin		Urokinase	35
Thiamine hydrochloride	25	Dermatological	52	Urologicals	59
Thioguanine	135	Oncology	138	Uromitexan	143
Thiopental [Thiopentone]		Trexate	135	Ursodeoxycholic acid	18
sodium	107	Tri-sodium citrate	200	Ursosan	18
Thiopentone	107	Triamcinolone acetonide		Utrogestan	59
Thiotepa	133	Alimentary	23		
Thrombin	30	Dermatological	54		
Thymol glycerin	23	Hormone	62		
Thyroid and Antithyroid		Triamcinolone acetonide with			
Preparations	69	gramicidin, neomycin and			
Thyrotropin alfa	64	nystatin	185		
Ticagrelor	34	Triamcinolone acetonide with			
Ticarcillin with clavulanic acid	75	neomycin sulphate, gramicidin			
Ticlopidine	34	and nystatin	55		
Tigecycline	76	Triamcinolone hexacetonide	62		
Timolol	187	Triazolam	127		
Timolol maleate	42	Trichloroacetic acid	200		
Timoptol XE	187	Trichozole	82		
Tiotropium bromide	179	Trientine dihydrochloride	21		
Tiotropium bromide with		Trifluoperazine			
olodaterol	179	hydrochloride	123		
TMP	77	Trimeprazine tartrate	178		
TOBI	71	Trimethoprim	77		
Tobradex	185	Trimethoprim with			
Tobramycin		sulphamethoxazole			
Infection	71	[Co-trimoxazole]	77		
Sensory	184	Trisodium citrate	33		
Tobrex	184	Trometamol	196		
Tocilizumab	171	Tropicamide	188		
Tofranil	114	Tropisetron	121		
Tolcapone	106	Tropisetron-AFT	121		
Tolterodine tartrate	60	Truvada	85		
Topamax	119	Tuberculin, purified protein			
Topicalcaine	109	derivative	195		
Topical Products for Joint and		Two Cal HN	208		
Muscular Pain	104	TwoCal HN RTH (Vanilla)	208		
Topiramate	119	Tykerb	140		
Tracrium	102	Tysabri	126		
Tramadol hydrochloride	113				
Tramal 100	113				
Tramal 50	113				
Tramal SR 100	113				
Tramal SR 150	113				
Tramal SR 200	113				
Trandolapril	39				
Tranexamic acid	30				
Tranylcypromine sulphate	114				
Trastuzumab	173				
Travoprost	188				
Treatments for Dementia	130				
Treatments for Substance					

- V -

Vaclovir	91
Valaciclovir	91
Valcyte	91
Valganciclovir	91
Vancomycin	77
Varenicline	131
Varibar - Honey	194
Varibar - Nectar	194
Varibar - Pudding	194
Varibar - Thin Liquid	194
Varicella vaccine [Chicken pox	
vaccine]	221
Varilrix	221
Vasodilators	48
Vasopressin	69
Vasopressin Agents	69
Vecuronium bromide	102
Vedafil	49
Velcade	135
Veletri	49
Venlafaxine	115
Venofer	22
Ventavis	50
Ventolin	180
Vepesid	136
Verapamil hydrochloride	44
Vergo 16	120
Verpamil SR	44
Vesanoid	138
Vesicare	60
Vexazone	18
Vfend	79
Victralis	90
Vidaza	134
Vigabatrin	119
Vimpat	117
Vinblastine sulphate	143
Vincristine sulphate	143
Vinorelbine	143
Viral Vaccines	218
Viramune Suspension	84
Viread	89
Visipaque	193
Vistil	189
Vistil Forte	189

- U -

Ultibro Breezhaler	179
Ultiva	113
Ultraproct	14
Umeclidinium	179
Umeclidinium with vilanterol	180
Univent	178
Ural	60
Urea	
Dermatological	53
Extemporaneous	200
Urex Forte	44

Zyvox77



