

June 2016

Volume 4 Number 2

Editor: Kaye Wilsonemail: enquiry@pharmac.govt.nz

Telephone +64 4 460 4990

Facsimile +64 4 460 4995

Level 9, 40 Mercer Street

PO Box 10 254 Wellington 6143

Freephone Information Line**0800 66 00 50** (9am – 5pm weekdays)**Circulation**

Accessible in an electronic format at no cost from the Health Professionals section of the PHARMAC website www.pharmac.govt.nz

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month. Alternatively there is a nominal charge for an annual subscription to the printed Schedule publications. To access either of these subscriptions visit our subscription website www.schedule.co.nz.

ProductionTypeset automatically from XML and T_EX.

XML version of the Schedule available from www.pharmac.govt.nz/pub/schedule/archive/

Programmers

Anrik Drenth & John Geering

email: texschedule@pharmac.govt.nz

©Pharmaceutical Management Agency



ISSN 1179-3708 pdf
ISSN 1172-9694 print

This work is licensed under the Creative Commons Attribution 3.0 New Zealand licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to PHARMAC and abide by the other licence terms. To view a copy of this licence, visit:

creativecommons.org/licenses/by/3.0/nz/.

Attribution to PHARMAC should be in written form and not by reproduction of the PHARMAC logo. While care has been taken in compiling this Schedule, PHARMAC takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

Part I General Rules 5

Part II Alimentary Tract and Metabolism 13

Blood and Blood Forming Organs 27

Cardiovascular System 39

Dermatologicals 51

Genito-Urinary System 57

Hormone Preparations 61

Infections 71

Musculoskeletal System 95

Nervous System 105

Oncology Agents and Immunosuppressants 132

Respiratory System and Allergies 176

Sensory Organs 183

Various 189

Extemporaneous Compounds (ECPs) 197

Special Foods 200

Vaccines 214

Part III Optional Pharmaceuticals 221

Index 223

Introducing PHARMAC

The Pharmaceutical Management Agency (PHARMAC) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. PHARMAC negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

PHARMAC's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about PHARMAC and the way we make funding decisions can be found on the PHARMAC website at <http://www.pharmac.health.nz/about>.

Named Patient Pharmaceutical Assessment policy

Named Patient Pharmaceutical Assessment (NPPA) provides a mechanism for individual patients to receive funding for medicines not listed in the Pharmaceutical Schedule (either at all or for their clinical circumstances). PHARMAC will assess applications that meet the prerequisites according to its Decision Criteria before deciding whether to approve applications for funding. The Decision Criteria will be used to assess both the individual clinical circumstances of each NPPA applicant, and the implications of each NPPA funding decision on PHARMAC's ability to carry out its legislative functions.

For more information on NPPA, or to apply, visit the PHARMAC website at <http://www.pharmac.health.nz/link/nppa> or call the Panel Coordinators at 0800 660 050 Option 2.

The Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by PHARMAC.

The Schedule does not show the final cost to Government of subsidising each community pharmaceutical, nor to DHB hospitals in purchasing each hospital pharmaceutical or other pharmaceuticals, including medical devices. The final cost will depend on any rebate and other arrangements PHARMAC has with the supplier or on any logistics arrangements put in place.

Finding Information in Section H

This book contains Section H of the Pharmaceutical Schedule and lists pharmaceuticals that can be used in DHB hospitals:

- Part I lists the rules in relation to use of Pharmaceuticals by DHB hospitals.
- Part II lists hospital pharmaceuticals that are funded for use in DHB hospitals. These are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. It also provides information on any national contracts that exist, and indicates which products have Hospital Supply Status (HSS).
- Part III lists optional pharmaceuticals for which national contracts exist, and DHB hospitals may choose to fund. In addition to the products listed in this book, a number of additional Optional Pharmaceuticals are listed in an addendum to Part III available at <http://www.pharmac.govt.nz>.

The listings are displayed alphabetically under each heading. The index lists both chemical entities and product brand names.

Glossary

Units of Measure

gram	g	microgram.....	mcg	millimole.....	mmol
kilogram	kg	milligram	mg	unit.....	u
international unit	iu	millilitre.....	ml		

Abbreviations

application	app	enteric coated	EC	solution	soln
capsule	cap	granules	grans	suppository	suppos
cream.....	crm	injection	inj	tablet.....	tab
dispersible	disp	liquid	liq	tincture.....	tinc
effervescent	eff	lotion	lotn		
emulsion	emul	ointment.....	oint		

HSS Hospital Supply Status (Refer to Rule 20)

Guide to Section H listings

Example

ANATOMICAL HEADING			
	Price (ex man. Excl. GST) \$	Per	Brand or Generic Manufacturer
THERAPEUTIC HEADING			
Generic name listed by therapeutic group and subgroup	CHEMICAL A - Restricted see terms below ⬆ Presentation A.....10.00	100	Brand A
	⚡ Restricted Only for use in children under 12 years of age		Brand or manufacturer's name
Indicates only presentation B1 is Restricted	CHEMICAL B - Some items restricted see terms below ⬆ Presentation B1.....1,589.00 Presentation B2	1	Brand B1 e.g. Brand B2
	⚡ Restricted Oncologist or haematologist		
From 1 January 2012 to 30 June 2014, at least 99% of the total volume of this item purchased must be Brand C	CHEMICAL C Presentation C - 1% DV Limit Jan-12 to 201415.00	28	Brand C
	CHEMICAL D - Restricted see terms below ⬆ Presentation D - 1% DV Limit Mar-13 to 201438.65	500	Brand D
Standard national price excluding GST	⚡ Restricted <i>Limited to five weeks' treatment</i> Either: 1 For the prophylaxis of venous thromboembolism following a total hip replacement; or 2 For the prophylaxis of venous thromboembolism following a total knee replacement.		Product with Hospital Supply Status (HSS) Quantity the Price applies to
Form and strength	CHEMICAL E Presentation E.....		e.g. Brand E Not a contracted product
⬆ Item restricted (see above); ⬇ Item restricted (see below) Products with Hospital Supply Status (HSS) are in bold			

INTRODUCTION

Section H contains general rules that apply, and other information relating, to Hospital Pharmaceuticals and Optional Pharmaceuticals.

Where relevant, Section H shows the Price at which a Pharmaceutical can be purchased directly from the Pharmaceutical supplier by DHBs, providers of logistics services, wholesalers or other such distributors, or Contract Manufacturers.

The Price is determined via contractual arrangements between PHARMAC and the relevant Pharmaceutical supplier. Where a Pharmaceutical is listed in Part II of Section H, but no Price and/or brand of Pharmaceutical is indicated, each DHB may purchase any brand and/or pay the price that the DHB negotiates with the relevant Pharmaceutical supplier.

As required by section 23(7) of the Act, in performing any of its functions in relation to the supply of Pharmaceuticals, a DHB must not act inconsistently with the Pharmaceutical Schedule.

INTERPRETATION AND DEFINITIONS

1 Interpretation and Definitions

1.1 In this Schedule, unless the context otherwise requires:

“**Act**”, means the New Zealand Public Health and Disability Act 2000.

“**Combined Pharmaceutical Budget**”, means the pharmaceutical budget set for PHARMAC by the Crown for the subsidised supply of Community Pharmaceuticals and Pharmaceutical Cancer Treatments including for named patients in exceptional circumstances.

“**Community**”, means any setting outside of a DHB Hospital.

“**Community Pharmaceutical**”, means a Pharmaceutical listed in Sections A to G or I of the Pharmaceutical Schedule that is subsidised by the Funder from the Combined Pharmaceutical Budget and, for the purposes of this Section H, includes Pharmaceutical Cancer Treatments (PCTs).

“**Contract Manufacturer**”, means a manufacturer or a supplier that is a party to a contract with the relevant DHB Hospital to compound Pharmaceuticals, on request from that DHB Hospital.

“**Designated Delivery Point**”, means at a DHB Hospital’s discretion:

- a) a delivery point agreed between a Pharmaceutical supplier and the relevant DHB Hospital, to which delivery point that Pharmaceutical supplier must supply a National Contract Pharmaceutical directly at the Price; and/or
- b) any delivery point designated by the relevant DHB Hospital or PHARMAC, such delivery point being within 30 km of the relevant Pharmaceutical supplier’s national distribution centre.

“**DHB**”, means an organisation established as a District Health Board by or under Section 19 of the Act.

“**DHB Hospital**”, means a hospital (including community trust hospitals) and/or an associated health service that is funded by a DHB including (but not limited to) district nursing services and child dental services.

“**DV Limit**”, means, for a particular National Contract Pharmaceutical with HSS, the National DV Limit or the Individual DV Limit.

“**DV Pharmaceutical**”, means a discretionary variance Pharmaceutical that does not have HSS but is used in place of one that does. Usually this means it is the same chemical entity, at the same strength, and in the same or a similar presentation or form, as the relevant National Contract Pharmaceutical with HSS. Where this is not the case, a note will be included with the listing of the relevant Hospital Pharmaceutical.

“**Extemporaneously Compounded Product**”, means a Pharmaceutical that is compounded from two or more Pharmaceuticals, for the purposes of reconstitution, dilution or otherwise.

“**First Transition Period**”, means the period of time after notification that a Pharmaceutical has been awarded HSS and before HSS is implemented.

“**Funder**”, means the body or bodies responsible, pursuant to the Act, for the funding of Pharmaceuticals listed on the Schedule (which may be one or more DHBs and/or the Ministry of Health) and their successors.

“**Give**”, means to administer, provide or dispense (or, in the case of a Medical Device, use) a Pharmaceutical, or to arrange for the administration, provision or dispensing (or, in the case of a Medical Device, use) of a Pharmaceutical, and “**Given**” has a corresponding meaning.

“**Hospital Pharmaceuticals**”, means the list of Pharmaceuticals set out in Section H Part II of the Schedule which includes some National Contract Pharmaceuticals.

“**HSS**”, stands for hospital supply status, which means the status of being the brand of the relevant National Contract Pharmaceutical that DHBs are obliged to purchase, subject to any DV Limit, for the period of hospital supply,

as awarded under an agreement between PHARMAC and the relevant Pharmaceutical supplier. Pharmaceuticals with HSS are listed in Section H in bold text.

“Indication Restriction”, means a limitation placed by PHARMAC on the funding of a Hospital Pharmaceutical which restricts funding to treatment of particular clinical circumstances.

“Individual DV Limit”, means, for a particular National Contract Pharmaceutical with HSS and a particular DHB Hospital, the discretionary variance limit, being the specified percentage of that DHB Hospital's Total Market Volume up to which that DHB Hospital may purchase DV Pharmaceuticals of that National Contract Pharmaceutical.

“Local Restriction”, means a restriction on the use of a Pharmaceutical in specific DHB Hospitals on the basis of prescriber type that is implemented by the relevant DHB in accordance with rule 7.

“Medical Device”, has the meaning set out in the Medicines Act 1981.

“Named Patient Pharmaceutical Assessment Advisory Panel”, means the panel of clinicians, appointed by the PHARMAC Board, that is responsible for advising PHARMAC, in accordance with its Terms of Reference, on Named Patient Pharmaceutical Assessment applications and any Exceptional Circumstances renewal applications submitted after 1 March 2012.

“National Contract”, means a contractual arrangement between PHARMAC and a Pharmaceutical supplier which sets out the basis on which any Pharmaceutical may be purchased for use in a DHB Hospital, including an agreement as to a national price.

“National Contract Pharmaceutical”, means a brand of Pharmaceutical listed in Section H, where PHARMAC has entered into contractual arrangements with the relevant Pharmaceutical supplier that specify the terms and conditions of listing, including the Price. Such Pharmaceuticals are recognisable in Section H because the relevant listing identifies the brand and Price.

“National DV Limit”, means, for a particular National Contract Pharmaceutical with HSS, the discretionary variance limit, being the specified percentage of the Total Market Volume up to which all DHB Hospitals may collectively purchase DV Pharmaceuticals of that National Contract Pharmaceutical.

“Optional Pharmaceuticals”, means the list of National Contract Pharmaceuticals set out in Section H Part III of the Schedule.

“PHARMAC”, means the Pharmaceutical Management Agency established by Section 46 of the Act.

“Pharmacode”, means the six or seven digit identifier assigned to a Pharmaceutical by the Pharmacy Guild following application from a Pharmaceutical supplier.

“Pharmaceutical”, means a medicine, therapeutic medical device, or related product or related thing listed in Sections B to I of the Schedule.

“Pharmaceutical Cancer Treatment”, means Pharmaceuticals for the treatment of cancer, listed in Sections A to G of the Schedule and identified therein as a “PCT” or “PCT only” Pharmaceutical that DHBs must fund for use in their DHB hospitals, and/or in association with outpatient services provided by their DHB Hospitals, in relation to the treatment of cancers.

“Prescriber Restriction”, means a restriction placed by PHARMAC on the funding of a Pharmaceutical on the basis of prescriber type (and where relevant in these rules, includes a Local Restriction).

“Price”, means the standard national price for a National Contract Pharmaceutical, and, unless agreed otherwise between PHARMAC and the Pharmaceutical supplier, includes any costs associated with the supply of the National Contract Pharmaceutical to, at a DHB Hospital's discretion, any Designated Delivery Point, or to a Contract Manufacturer (expressly for the purpose of compounding), but does not include the effect of any rebates which may have been negotiated between PHARMAC and the Pharmaceutical supplier.

“Restriction”, means a limitation, put in place by PHARMAC or a DHB, restricting the funding of a Pharmaceutical and includes Indication Restrictions, Local Restrictions and Prescriber Restrictions (as defined in this Part I of Section H).

“Schedule”, means this Pharmaceutical Schedule and all its sections and appendices.

“Special Authority Approval”, means an approval for funding of a Community Pharmaceutical that is marked in Sections B-G of the Schedule as being subject to a Special Authority restriction.

“Total Market Volume”, means, for a particular Hospital Pharmaceutical with HSS in any given period, in accordance with the data available to PHARMAC, the sum of:

- a) the total number of Units of the relevant Hospital Pharmaceutical with HSS purchased by all DHB Hospitals, or by a particular DHB Hospital in the case of the Individual DV Limit; and
- b) the total number of Units of all the relevant DV Pharmaceuticals purchased by all DHB Hospitals, or by a particular DHB Hospital in the case of the Individual DV Limit.

“Unapproved Indication”, means, for a Pharmaceutical, an indication for which it is not approved under the Medicines Act 1981. Clinicians prescribing Pharmaceuticals for Unapproved Indications should be aware of, and comply with, their obligations under Section 25 and/or Section 29 of the Medicines Act 1981 and as set out in rule 23.

“Unit”, means an individual unit of a Pharmaceutical (e.g. a tablet, 1 ml of an oral liquid, an ampoule or a syringe).

“Unlisted Pharmaceutical”, means a Pharmaceutical that is within the scope of a Hospital Pharmaceutical, but is not listed in Section H Part II.

- 1.2 In addition to the above interpretations and definitions, unless the content requires otherwise, a reference in the Schedule to:
 - a) the singular includes the plural; and
 - b) any legislation includes a modification and re-enactment of, legislation enacted in substitution for, and a regulation, Order in Council, and other instrument from time to time issued or made under, that legislation.

HOSPITAL SUPPLY OF PHARMACEUTICALS

2 Hospital Pharmaceuticals

- 2.1 Section H Part II contains the list of Hospital Pharmaceuticals that must be funded by DHB Hospitals. Section H Part II does not currently encompass the following categories of pharmaceuticals except for any items specifically listed in this Section H Part II:
 - a) Medical Devices;
 - b) whole or fractionated blood products;
 - c) diagnostic products which have an ex vivo use, such as pregnancy tests and reagents;
 - d) disinfectants and sterilising products, except those that are to be used in or on a patient;
 - e) foods and probiotics;
 - f) radioactive materials;
 - g) medical gases; and
 - h) parenteral nutrition.

Subject to rule 2.2, the funding of pharmaceuticals identified in a)–h) above is a decision for individual DHB Hospitals.

- 2.2 Section H Part III lists Optional Pharmaceuticals that PHARMAC and the relevant Pharmaceutical supplier have entered into contractual arrangements for the purchase of, including an agreement on a national price and other obligations such as HSS. DHB Hospitals may choose whether or not to fund the Optional Pharmaceuticals listed in Part III of Section H, but if they do, they must comply with any National Contract requirements.
- 2.3 Section H Part II does not encompass the provision of pharmaceutical treatments for DHB Hospital staff as part of an occupational health and safety programme. DHB Hospitals may choose whether or not to fund pharmaceutical treatments for such use, but if they do, they must comply with any National Contract requirements.

3 DHB Supply Obligations

- 3.1 In accordance with section 23(7) of the Act, in performing any of its functions in relation to the supply of pharmaceuticals, a DHB must not act inconsistently with the Pharmaceutical Schedule, which includes these General Rules.
- 3.2 DHB Hospitals are not required to hold stock of every Hospital Pharmaceutical listed in Section H Part II, but they must Give it within a reasonable time if it is prescribed.
- 3.3 DHB Hospitals are able to hold stock of an Unlisted Pharmaceutical if doing so is considered necessary for the DHB Hospital to be able to Give the Unlisted Pharmaceutical in a timely manner under rules 11–17 inclusive.
- 3.4 Except where permitted in accordance with rule 11, DHBs must not Give:
 - a) an Unlisted Pharmaceutical; or
 - b) a Hospital Pharmaceutical outside of any relevant Restrictions.

4 Funding

- 4.1 The purchase costs of Hospital Pharmaceuticals or Optional Pharmaceuticals administered, provided or dispensed by DHB Hospitals must be funded by the relevant DHB Hospital from its own budget, with the exception of:
 - a) Pharmaceutical Cancer Treatments;
 - b) Community Pharmaceuticals that have been brought to the DHB hospital by the patient who is being treated by outpatient Services or who is admitted as an inpatient;
 - c) Community Pharmaceuticals that have been dispensed to a mental health day clinic under a Practitioner's

- Supply Order; and
 - d) Unlisted Pharmaceutical that have been brought to the DHB Hospital by the patient who is admitted as an inpatient.
- 4.2 For the avoidance of doubt, Pharmaceutical Cancer Treatments and Community Pharmaceuticals are funded through the Combined Pharmaceutical Budget, and Unlisted Pharmaceuticals are funded by the patient.

LIMITS ON SUPPLY

5 Prescriber Restrictions

- 5.1 A DHB Hospital may only Give a Hospital Pharmaceutical that has a Prescriber Restriction if it is prescribed:
- a) by a clinician of the type specified in the restriction for that Pharmaceutical or, subject to rule 5.2, pursuant to a recommendation from such a clinician;
 - b) in accordance with a protocol or guideline that has been endorsed by the DHB Hospital; or
 - c) in an emergency situation, provided that the prescriber has made reasonable attempts to comply with rule 5.1(a) above. If on-going treatment is required (i.e. beyond 24 hours) subsequent prescribing must comply with rule 5.1(a).
- 5.2 Where a Hospital Pharmaceutical is prescribed pursuant to a recommendation from a clinician of the type specified in the restriction for that Pharmaceutical:
- a) the prescriber must consult with a clinician of the type specified in the restriction for that Pharmaceutical; and
 - b) the consultation must relate to the patient for whom the prescription is written; and
 - c) the consultation may be in person, by telephone, letter, facsimile or email; and
 - d) appropriate records are kept of the consultation, including recording the name of the advising clinician on the prescription/chart.
- 5.3 Where a clinician is working under supervision of a consultant who is of the type specified in the restriction for that Pharmaceutical, the requirements of rule 5.2 can be deemed to have been met.

6 Indication Restrictions

- 6.1 A DHB Hospital may only Give a Hospital Pharmaceutical that has an Indication Restriction, if it is prescribed for treatment of a patient with the particular clinical circumstances set out in the Indication Restriction.
- 6.2 If a patient has a current Special Authority Approval for the Hospital Pharmaceutical that the DHB Hospital wishes to Give, then the Indication Restriction is deemed to have been met.
- 6.3 If a Hospital Pharmaceutical has an Indication Restriction that is “for continuation only” then the DHB Hospital should only Give the Hospital Pharmaceutical where:
- a) the patient has been treated with the Pharmaceutical in the Community; or
 - b) the patient is unable to be treated with an alternative Hospital Pharmaceutical, and the prescriber has explained to the patient that the Pharmaceutical is not fully subsidised in the Community.

7 Local Restrictions

- 7.1 A DHB Hospital may implement a Local Restriction, provided that:
- a) in doing so, it ensures that the Local Restriction does not unreasonably limit funded access to the Hospital Pharmaceutical or undermine PHARMAC’s decision that the Hospital Pharmaceutical must be funded;
 - b) it provides PHARMAC with details of each Local Restriction that it implements; and
- 7.2 PHARMAC may, when it considers that a Local Restriction does not conform to rule 7.1 above, require a DHB to amend or remove that Local Restriction.

8 Community use of Hospital Pharmaceuticals

- 8.1 Except where otherwise specified in Section H, DHB Hospitals can Give any Hospital Pharmaceutical to a patient for use in the Community, provided that:
- a) the quantity does not exceed that sufficient for up to 30 days’ treatment, unless:
 - i) it would be inappropriate to provide less than the amount in an original pack; or
 - ii) the relevant DHB Hospital has a Dispensing for Discharge Policy and the quantity dispensed is in accordance with that policy; and
 - b) the Hospital Pharmaceutical is supplied consistent with any applicable Restrictions.

9 Community use of Medical Devices

- 9.1 Subject to rules 9.2 and 9.3, DHB Hospitals may Give a Medical Device for patients for use in the Community.
- 9.2 Where a Medical Device (or a similar Medical Device) is a Community Pharmaceutical, the DHB Hospital must supply:

- a) the brand of Medical Device that is listed in Sections A-G of the Schedule; and
 - b) only to patients who meet the funding eligibility criteria set out in Sections A-G of the Schedule.
- 9.3 Where a DHB Hospital has supplied a Medical Device to a patient; and
- a) that Medical Device (or a similar Medical Device) is subsequently listed in Sections A-G of the Schedule; and
 - b) the patient would not meet any funding eligibility criteria for the Medical Device set out in Sections A-G of the Schedule; and
 - c) the Medical Device has consumable components that need to be replaced throughout its usable life; then DHB Hospitals may continue to fund consumable products for that patient until the end of the usable life of the Medical Device. At the end of the usable life of the device, funding for a replacement device must be consistent with the Pharmaceutical Schedule and/or in accordance with the Named Patient Pharmaceutical Assessment policy.
- 9.4 DHB Hospitals may also continue to fund consumable products, as in rule 9.3 above, in situations where the DHB has been funding consumable products but where the Medical Device was funded by the patient.

10 Extemporaneous Compounding

- 10.1 A DHB Hospital may Give any Extemporaneously Compounded Product for a patient in its care, provided that:
- a) all of the component Pharmaceuticals of the Extemporaneously Compounded Product are Hospital Pharmaceuticals; and
 - b) the Extemporaneously Compounded Product is supplied consistent with any applicable rules or Restrictions for its component Hospital Pharmaceuticals.
- 10.2 For the avoidance of doubt, this rule 10.1 applies to any Extemporaneously Compounded Product, whether it is manufactured by the DHB Hospital or by a Contract Manufacturer.

EXCEPTIONS

11 Named Patient Pharmaceutical Assessment

- 11.1 A DHB Hospitals may only Give:
- a) an Unlisted Pharmaceutical; or
 - b) a Hospital Pharmaceutical outside of any relevant Restrictions,
- in accordance with the Named Patient Pharmaceutical Assessment Policy or rules 12–17 inclusive.

12 Continuation

- 12.1 Where a patient's clinical circumstances have been stabilised via treatment in the Community with a pharmaceutical that has not been funded by the Funder, and that patient is admitted to hospital as an inpatient, a DHB Hospital may fund that pharmaceutical for the duration of the patient's stay, where:
- a) the patient has not brought (or cannot arrange to bring) the pharmaceuticals to the DHB Hospital, or pharmacy staff consider that the pharmaceuticals brought to the DHB Hospital by the patient cannot be used; and
 - b) interrupted or delayed treatment would have significant adverse clinical consequences; and
 - c) it is not considered appropriate to switch treatment to a Hospital Pharmaceutical.

13 Pre-Existing Use

- 13.1 Subject to 13.2, where a DHB Hospital has Given a pharmaceutical for a patient prior to 1 July 2013, and the pharmaceutical:
- a) is an Unlisted Pharmaceutical; or
 - b) treatment of the patient would not comply with any relevant Restrictions;
- the DHB Hospital may continue to Give that pharmaceutical if it is considered that there would be significant adverse clinical consequences from ceasing or switching treatment.
- 13.2 Each DHB Hospital must, by no later than 1 October 2013, provide PHARMAC with a report on pharmaceuticals it has Given in accordance with this rule 13 where treatment has continued beyond 1 August 2013.

14 Clinical Trials and Free Stock

- 14.1 DHB Hospitals may Give any pharmaceutical that is funded by a third party and is being used:
- 14.1.1 as part of a clinical trial that has Ethics Committee approval; or
 - 14.1.2 for on-going treatment of patients following the end of such a clinical trial.
- 14.2 DHB Hospitals may Give any pharmaceutical that is provided free of charge by a supplier, provided that the pharmaceutical is provided as part of a programme of which the DHB, or supplier, has notified PHARMAC.

15 Pharmaceutical Cancer Treatments in Paediatrics

DHB Hospitals may Give any pharmaceutical for use within a paediatric oncology/haematology service for the treatment of

cancer.

16 Other Government Funding

DHB Hospitals may Give any pharmaceutical where funding for that pharmaceutical has been specifically provided by a Government entity other than PHARMAC or a DHB.

17 Other Exceptions

- 17.1 PHARMAC may also approve the funding of a pharmaceutical within a single DHB Hospital for information gathering purposes or otherwise related to PHARMAC's decision-making process for considering additions to or amendments to the Pharmaceutical Schedule.
- 17.2 Funding approvals granted under rule 17.1 will be subject to specific limitations on use as determined appropriate by PHARMAC in each circumstance, in consultation with the relevant DHB Hospital and/or DHB.

NATIONAL CONTRACTING

18 Hospital Pharmaceutical Contracts

- 18.1 A DHB Hospital may enter into a contract for the purchase of any Pharmaceutical, including any Medical Device, that it is entitled to fund in accordance with this Schedule H and that is not a National Contract Pharmaceutical, provided that such a contract:
 - a) does not oblige the relevant DHB Hospital to purchase a volume of that Pharmaceutical, if that Pharmaceutical is a DV Pharmaceutical, that is greater than the relevant DV Limit;
 - b) enables PHARMAC to access and use future price and volume data in respect of that Pharmaceutical; and
 - c) enables the relevant DHB Hospital to terminate the contract or relevant parts of the contract in order to give full effect to the National Contract on no more than 3 months' written notice to the Pharmaceutical supplier.
- 18.2 From 1 July 2013, where a DHB Hospital has a pre-existing supply contract for a particular brand of chemical entity for which there is a National Contract Pharmaceutical, the DHB may continue purchasing the chemical entity in accordance with its pre-existing supply contract however:
 - a) from the day its pre-existing supply contract expires, that DHB Hospital is to purchase the relevant National Contract Pharmaceutical listed in Section H at the Price, and is to comply with any DV Limits for the National Contract Pharmaceutical where it has HSS;
 - b) if purchase of the relevant National Contract Pharmaceutical listed in Section H at the Price, where it has HSS, would not cause the relevant DHB Hospital to be in breach of its pre-existing supply contract for a particular brand of chemical entity; the DHB Hospital must purchase the National Contract Pharmaceutical.
- 18.3 Following written notification from PHARMAC that a Pharmaceutical is a National Contract Pharmaceutical, either through Section H updates or otherwise, DHB Hospitals must, unless PHARMAC expressly notifies otherwise:
 - a) take any steps available to them to terminate pre-existing contracts or relevant parts of such a contract, and
 - b) not enter any new contracts or extend the period of any current contracts, for the supply of that National Contract Pharmaceutical or the relevant chemical entity or Medical Device.

19 National Contract Pharmaceuticals

- 19.1 DHB Hospitals must take all necessary steps to enable any contracts between PHARMAC and a Pharmaceutical supplier in relation to National Contract Pharmaceuticals to be given full effect.
- 19.2 The contractual arrangement between PHARMAC and the relevant supplier of a National Contract Pharmaceutical requires it to be made available for purchase at the relevant Price by any or all of the following:
 - a) DHB Hospitals at Designated Delivery Points; and/or
 - b) Contract Manufacturers (expressly for the purpose of compounding).

In the case of Medical Devices, a National Contract may require the Medical Device to be purchased by, and/or supplied to, a third party logistics provider.

20 Hospital Supply Status (HSS)

- 20.1 The DV Limit for any National Contract Pharmaceutical which has HSS is set out in the listing of the relevant National Contract Pharmaceutical in Section H, and may be amended from time to time.
- 20.2 If a National Contract Pharmaceutical is listed in Section H as having HSS, DHB Hospitals:
 - a) are expected to use up any existing stocks of DV Pharmaceuticals during the First Transition Period;
 - b) must not purchase DV Pharmaceuticals in volumes exceeding their usual requirements, or in volumes exceeding those which they reasonably expect to use, within the First Transition Period;
 - c) must ensure that Contract Manufacturers, when manufacturing an Extemporaneously Compounded Product on their behalf, use the National Contract Pharmaceutical with HSS; and

- d) must purchase the National Contract Pharmaceutical with HSS except:
 - i) to the extent that the DHB Hospital may use its discretion to purchase a DV Pharmaceutical within the DV Limit, provided that (subject to rule 20.2(d)(iii) below) the DV Limit has not been exceeded nationally;
 - ii) if the Pharmaceutical supplier fails to supply that National Contract Pharmaceutical, in which case the relevant DHB Hospital does not have to comply with the DV Limit for that National Contract Pharmaceutical during that period of non-supply (and any such month(s) included in a period of non-supply will be excluded in any review of the DV Limit in accordance with rule 20.3 below);
 - iii) that where the DV Limit has been exceeded nationally, the DHB Hospital may negotiate with the Pharmaceutical supplier that supplies the National Contract Pharmaceutical with HSS for written permission to vary the application of that DHB Hospital's Individual DV Limit for any patient whose exceptional needs require a DV Pharmaceutical.

20.3 PHARMAC may, in its discretion, for any period or part period:

- a) review usage by DHB Hospitals of the National Contract Pharmaceutical and DV Pharmaceuticals to determine whether the DV Limit has been exceeded; and
- b) audit compliance by DHB Hospitals with the DV Limits and related requirements.

20.4 PHARMAC will address any issues of non-compliance by any individual DHB or DHB Hospital with a DV Limit by:

- a) obtaining the relevant DHB or DHB Hospital's assurance that it will comply with the DV Limit for that National Contract Pharmaceutical with HSS in the remainder of the applicable period and any subsequent periods; and
- b) informing the relevant supplier of the HSS Pharmaceutical of any individual DHB or DHB Hospital's non-compliance with the DV Limit for that HSS Pharmaceutical.

20.5 In addition to the steps taken by PHARMAC under rule 20.4 above to address any issues of non-compliance by any individual DHB or DHB Hospital with a DV Limit, the relevant Pharmaceutical supplier may require, in its discretion, financial compensation from the relevant DHB or DHB Hospital:

- a) an amount representing that DHB or DHB Hospital's contribution towards exceeding the DV Limit (where PHARMAC is able to quantify this based on the information available to it); or
- b) the sum of \$1,000 or \$5,000 (depending on the terms of the applicable national contract applying to the HSS Pharmaceutical),

whichever is the greater as between sub-paragraphs (a) and (b) within the number of business days specified in the notice from the Pharmaceutical supplier requiring such payment to be made.

21 Collection of rebates and payment of financial compensation

21.1 Following the receipt of any rebates from a Pharmaceutical supplier in respect of a particular National Contract Pharmaceutical, PHARMAC will notify each relevant DHB and DHB Hospital of the amount of the rebate owing to it, being a portion of the total rebate determined by PHARMAC on the basis of that DHB Hospital's usage of that National Contract Pharmaceutical, where this is able to be determined. Where data to determine individual DHB Hospitals' usage is not available, PHARMAC will apportion rebates on the basis of an alternative method agreed between the relevant DHBs and PHARMAC.

21.2 PHARMAC will pay each DHB Hospital the rebate amounts (if any) owing to it, no less frequently than once each calendar quarter in respect of rebates received quarterly (or more often).

22 Price and Volume Data

22.1 DHB Hospitals must provide to PHARMAC, on a monthly basis in accordance with PHARMAC's requirements, any volume data and, unless it would result in a breach of a pre-existing contract, price data held by those DHB Hospitals in respect of any Pharmaceutical (including any Medical Device) listed in Section H.

22.2 All price and volume data provided to PHARMAC under rule 22.1 above should identify the relevant Hospital Pharmaceutical by using a Pharmacode or some other unique numerical identifier, and the date (month and year) on which the DHB Hospital incurred a cost for the purchase of that Hospital Pharmaceutical. Volume is to be measured in units (that being the smallest possible whole Unit – e.g. a capsule, a vial, a millilitre etc).

MISCELLANEOUS PROVISIONS

23 Unapproved Pharmaceuticals

Prescribers should, where possible, prescribe Hospital Pharmaceuticals that are approved under the Medicines Act 1981. However, the funding criteria (including Restrictions) under which a Hospital Pharmaceutical is listed in Section H of the

Schedule may:

23.1 in some cases, explicitly permit a DHB to fund a Pharmaceutical that is not approved under the Medicines Act 1981 or for an Unapproved Indication; or

23.2 not explicitly prohibit a DHB from funding a Pharmaceutical for use for an Unapproved Indication;

Accordingly, if clinicians are planning on prescribing an unapproved Pharmaceutical or a Pharmaceutical for an Unapproved Indication, they should:

23.1 be aware of and comply with their obligations under sections 25 and/or 29 of the Medicines Act 1981, as applicable, and otherwise under that Act and the Medicines Regulations 1984;

23.2 be aware of and comply with their obligations under the Health and Disability Commissioner's Code of Consumer Rights, including the requirement to obtain informed consent from the patient (PHARMAC recommends that clinicians obtain written consent); and

23.3 exercise their own skill, judgment, expertise and discretion, and make their own prescribing decisions with respect to the use of an unapproved Pharmaceutical or a Pharmaceutical for an Unapproved Indication.

Clinicians should be aware that simply by listing a Pharmaceutical on the Pharmaceutical Schedule, PHARMAC makes no representations about whether that Pharmaceutical has any form of approval or consent under, or whether the supply or use of the Pharmaceutical otherwise complies with, the Medicines Act 1981. Further, the Pharmaceutical Schedule does not constitute an advertisement, advertising material or a medical advertisement as defined in the Medicines Act or otherwise.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Antacids and Antiflatulents			
Antacids and Reflux Barrier Agents			
ALUMINIUM HYDROXIDE WITH MAGNESIUM HYDROXIDE AND SIMETHICONE			
Tab 200 mg with magnesium hydroxide 200 mg and simethicone 20 mg			<i>e.g. Mylanta</i>
Oral liq 200 mg with magnesium hydroxide 200 mg and simethicone 20 mg per 5 ml			<i>e.g. Mylanta</i>
Oral liq 400 mg with magnesium hydroxide 400 mg and simethicone 30 mg per 5 ml			<i>e.g. Mylanta Double Strength</i>
<i>(e.g. Mylanta Oral liq 200 mg with magnesium hydroxide 200 mg and simethicone 20 mg per 5 ml to be delisted 1 August 2016)</i>			
SIMETHICONE			
Oral drops 100 mg per ml			
SODIUM ALGINATE WITH MAGNESIUM ALGINATE			
Powder for oral soln 225 mg with magnesium alginate 87.5 mg, sachet			<i>e.g. Gaviscon Infant</i>
SODIUM ALGINATE WITH SODIUM BICARBONATE AND CALCIUM CARBONATE			
Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg			<i>e.g. Gaviscon Double Strength</i>
Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml	4.95	500 ml	Acidex
SODIUM CITRATE			
Oral liq 8.8% (300 mmol/l)			
Phosphate Binding Agents			
ALUMINIUM HYDROXIDE			
Tab 600 mg			
CALCIUM CARBONATE – Restricted see terms below			
☞ Oral liq 250 mg per ml (100 mg elemental per ml)	39.00	500 ml	Roxane
☞ Restricted			
Initiation			
Only for use in children under 12 years of age for use as a phosphate binding agent.			
Antidiarrhoeals and Intestinal Anti-Inflammatory Agents			
Antipropulsives			
DIPHENOXYLATE HYDROCHLORIDE WITH ATROPINE SULPHATE			
Tab 2.5 mg with atropine sulphate 25 mcg			
LOPERAMIDE HYDROCHLORIDE			
Tab 2 mg			
Cap 2 mg – 1% DV Jul-14 to 2016	7.84	400	Diamide Relief
Rectal and Colonic Anti-Inflammatories			
BUDESONIDE – Restricted see terms on the next page			
☞ Cap 3 mg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➡Restricted			
Initiation — Crohn's disease			
Both:			
1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and			
2 Any of the following:			
2.1 Diabetes; or			
2.2 Cushingoid habitus; or			
2.3 Osteoporosis where there is significant risk of fracture; or			
2.4 Severe acne following treatment with conventional corticosteroid therapy; or			
2.5 History of severe psychiatric problems associated with corticosteroid treatment; or			
2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or			
2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).			
Initiation — Collagenous and lymphocytic colitis (microscopic colitis)			
Patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.			
Initiation — Gut Graft versus Host disease			
Patient has gut Graft versus Host disease following allogenic bone marrow transplantation.			
HYDROCORTISONE ACETATE			
Rectal foam 10%, CFC free (14 applications) – 1% DV Oct-15 to 2018.....	26.55	21.1 g	Colifoam
MESALAZINE			
Tab EC 400 mg	49.50	100	Asacol
Tab EC 500 mg	49.50	100	Asamax
Tab long-acting 500 mg	59.05	100	Pentasa
Tab 800 mg	85.55	90	Asacol
Modified release granules 1 g	141.72	120 g	Pentasa
Suppos 500 mg	22.80	20	Asacol
Suppos 1 g – 1% DV Jun-15 to 2018	54.60	30	Pentasa
Enema 1 g per 100 ml – 1% DV Sep-15 to 2018	41.30	7	Pentasa
OLSALAZINE			
Tab 500 mg			
Cap 250 mg			
SODIUM CROMOGLYCATE			
Cap 100 mg			
SULPHASALAZINE			
Tab 500 mg – 1% DV Oct-13 to 2016	11.68	100	Salazopyrin
Tab EC 500 mg – 1% DV Oct-13 to 2016	12.89	100	Salazopyrin EN
Local Preparations for Anal and Rectal Disorders			
Antihæmorrhoidal Preparations			
CINCHOCAINE HYDROCHLORIDE WITH HYDROCORTISONE			
Oint 5 mg with hydrocortisone 5 mg per g	15.00	30 g	Proctosedyl
Suppos 5 mg with hydrocortisone 5 mg per g	9.90	12	Proctosedyl
FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE			
Oint 950 mcg with fluocortolone pivalate 920 mcg and cinchocaine hydrochloride 5 mg per g	6.35	30 g	Ultraproct
Suppos 630 mcg with fluocortolone pivalate 610 mcg and cinchocaine hydrochloride 1 mg	2.66	12	Ultraproct

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Management of Anal Fissures			
GLYCERYL TRINITRATE			
Oint 0.2%	22.00	30 g	Rectogesic
Rectal Sclerosants			
OILY PHENOL [PHENOL OILY]			
Inj 5%, 5 ml vial			
Antispasmodics and Other Agents Altering Gut Motility			
GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – 1% DV Oct-13 to 2016	17.14	10	Max Health
Max Health brand - HSS with 1% DV will apply 1 July 2016 to 30 June 2019.			
HYOSCINE BUTYLBROMIDE			
Tab 10 mg	2.18	20	Gastrosoothe
Inj 20 mg, 1 ml ampoule	9.57	5	Buscopan
MEBEVERINE HYDROCHLORIDE			
Tab 135 mg – 1% DV Sep-14 to 2017	18.00	90	Colofac
Antiulcerants			
Antisecretory and Cytoprotective			
MISOPROSTOL			
Tab 200 mcg – 1% DV Jun-16 to 2019	41.50	120	Cytotec
H2 Antagonists			
CIMETIDINE			
Tab 200 mg			
Tab 400 mg			
RANITIDINE			
Tab 150 mg – 1% DV Nov-14 to 2017	10.30	500	Ranitidine Relief
Tab 300 mg – 1% DV Nov-14 to 2017	14.73	500	Ranitidine Relief
Oral liq 150 mg per 10 ml – 1% DV Sep-14 to 2017	4.92	300 ml	Peptisoothe
Inj 25 mg per ml, 2 ml ampoule	8.75	5	Zantac
Proton Pump Inhibitors			
LANSOPRAZOLE			
Cap 15 mg – 1% DV Jan-16 to 2018	5.08	100	Lanzol Relief
Cap 30 mg – 1% DV Jan-16 to 2018	5.93	100	Lanzol Relief
OMEPRAZOLE			
⚡ Tab dispersible 20 mg			
➡ Restricted			
Initiation			
Only for use in tube-fed patients.			
Cap 10 mg – 1% DV Jan-15 to 2017	2.23	90	Omezol Relief
Cap 20 mg – 1% DV Jan-15 to 2017	2.91	90	Omezol Relief
Cap 40 mg – 1% DV Jan-15 to 2017	4.42	90	Omezol Relief
Powder for oral liq	42.50	5 g	Midwest

ALIMENTARY TRACT AND METABOLISM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Inj 40 mg ampoule	19.00	5	Dr Reddy's Omeprazole
Inj 40 mg ampoule with diluent	33.65	5	Dr Reddy's Omeprazole
PANTOPRAZOLE			
Tab EC 20 mg – 1% DV May-14 to 2016	2.68	100	Pantoprazole Actavis 20
Tab EC 40 mg – 1% DV May-14 to 2016	3.54	100	Pantoprazole Actavis 40
Inj 40 mg vial			

Site Protective Agents

BISMUTH TRIOXIDE			
Tab 120 mg	32.50	112	De-Nol
<i>(De-Nol Tab 120 mg to be delisted 1 January 2017)</i>			
COLLOIDAL BISMUTH SUBCITRATE			
Tab 120 mg	14.51	50	Gastrodenol
SUCRALFATE			
Tab 1 g			

Bile and Liver Therapy

L-ORNITHINE L-ASPARTATE – **Restricted** see terms below

⚡ Grans for oral liquid 3 g

➡ **Restricted**

Initiation

For patients with chronic hepatic encephalopathy who have not responded to treatment with, or are intolerant to lactulose, or where lactulose is contraindicated.

RIFAXIMIN – **Restricted** see terms below

⚡ Tab 550 mg – 1% DV Oct-14 to 2017 625.00 56 **Xifaxan**

➡ **Restricted**

Initiation

For patients with hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

Diabetes

Alpha Glucosidase Inhibitors

ACARBOSE			
Tab 50 mg – 1% DV Oct-15 to 2018	4.28	90	Glucobay
Tab 100 mg – 1% DV Oct-15 to 2018	7.78	90	Glucobay

Hyperglycaemic Agents

DIAZOXIDE – **Restricted** see terms below

⚡ Cap 25 mg	110.00	100	Proglycem
⚡ Cap 100 mg	280.00	100	Proglycem
⚡ Oral liq 50 mg per ml	620.00	30 ml	Proglycem

➡ **Restricted**

Initiation

For patients with confirmed hypoglycaemia caused by hyperinsulinism.

GLUCAGON HYDROCHLORIDE

Inj 1 mg syringe kit 32.00 1 Glucagen Hypokit

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GLUCOSE [DEXTROSE]			
Tab 1.5 g			
Tab 3.1 g			
Tab 4 g			
Gel 40%			
GLUCOSE WITH SUCROSE AND FRUCTOSE			
Gel 19.7% with sucrose 35% and fructose 19.7%, 18 g sachet			
Insulin - Intermediate-Acting Preparations			
INSULIN ASPART WITH INSULIN ASPART PROTAMINE			
Inj insulin aspart 30% with insulin aspart protamine 70%, 100 u per ml, 3 ml prefilled pen	52.15	5	NovoMix 30 FlexPen
INSULIN ISOPHANE			
Inj insulin human 100 u per ml, 10 ml vial			
Inj insulin human 100 u per ml, 3 ml cartridge			
INSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
Inj insulin lispro 25% with insulin lispro protamine 75%, 100 u per ml, 3 ml cartridge	42.66	5	Humalog Mix 25
Inj insulin lispro 50% with insulin lispro protamine 50%, 100 u per ml, 3 ml cartridge	42.66	5	Humalog Mix 50
INSULIN NEUTRAL WITH INSULIN ISOPHANE			
Inj insulin neutral 30% with insulin isophane 70%, 100 u per ml, 10 ml vial			
Inj insulin neutral 30% with insulin isophane 70%, 100 u per ml, 3 ml cartridge			
Inj insulin neutral 40% with insulin isophane 60%, 100 u per ml, 3 ml cartridge			
Inj insulin neutral 50% with insulin isophane 50%, 100 u per ml, 3 ml cartridge			
Insulin - Long-Acting Preparations			
INSULIN GLARGINE			
Inj 100 u per ml, 3 ml disposable pen	94.50	5	Lantus SoloStar
Inj 100 u per ml, 3 ml cartridge	94.50	5	Lantus
Inj 100 u per ml, 10 ml vial	63.00	1	Lantus
Insulin - Rapid-Acting Preparations			
INSULIN ASPART			
Inj 100 u per ml, 10 ml vial			
Inj 100 u per ml, 3 ml cartridge			
Inj 100 u per ml, 3 ml syringe	51.19	5	NovoRapid FlexPen
INSULIN GLULISINE			
Inj 100 u per ml, 10 ml vial	27.03	1	Apidra
Inj 100 u per ml, 3 ml cartridge	46.07	5	Apidra
Inj 100 u per ml, 3 ml disposable pen	46.07	5	Apidra Solostar
INSULIN LISPRO			
Inj 100 u per ml, 10 ml vial			
Inj 100 u per ml, 3 ml cartridge			

Products with Hospital Supply Status (HSS) are in **bold**

Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Insulin - Short-Acting Preparations			
INSULIN NEUTRAL			
Inj human 100 u per ml, 10 ml vial			
Inj human 100 u per ml, 3 ml cartridge			
Oral Hypoglycaemic Agents			
GLIBENCLAMIDE			
Tab 5 mg			
GLICLAZIDE			
Tab 80 mg – 1% DV Nov-14 to 2017	11.50	500	Glizide
GLIPIZIDE			
Tab 5 mg – 1% DV Sep-15 to 2018	2.85	100	Minidiab
METFORMIN HYDROCHLORIDE			
Tab immediate-release 500 mg – 1% DV Nov-15 to 2018	9.59	1,000	Metckek
Tab immediate-release 850 mg – 1% DV Dec-15 to 2018	7.82	500	Metformin Mylan
PIOGLITAZONE			
Tab 15 mg – 1% DV Dec-15 to 2018	3.47	90	Vexazone
Tab 30 mg – 1% DV Dec-15 to 2018	5.06	90	Vexazone
Tab 45 mg – 1% DV Dec-15 to 2018	7.10	90	Vexazone
Digestives Including Enzymes			
PANCREATIC ENZYME			
Cap EC 10,000 BP u lipase, 9,000 BP u amylase and 210 BP u pro- tease – 1% DV Oct-15 to 2018	34.93	100	Creon 10000
Cap EC 25,000 BP u lipase, 18,000 BP u amylase and 1,000 BP u protease – 1% DV Oct-15 to 2018	94.38	100	Creon 25000
Cap EC 25,000 BP u lipase, 22,500 BP u amylase and 1,250 BP u protease			
Powder 25,000 u lipase with 30,000 u amylase and 1,400 u protease per g			
URSODEOXYCHOLIC ACID – Restricted see terms below			
⚡ Cap 250 mg – 1% DV Sep-14 to 2017	53.40	100	Ursosan
➡ Restricted			
Initiation — Alagille syndrome or progressive familial intrahepatic cholestasis			
Either:			
1 Patient has been diagnosed with Alagille syndrome; or			
2 Patient has progressive familial intrahepatic cholestasis.			
Initiation — Chronic severe drug induced cholestatic liver injury			
All of the following:			
1 Patient has chronic severe drug induced cholestatic liver injury; and			
2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and			
3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.			
Initiation — Cirrhosis			
Both:			
1 Primary biliary cirrhosis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative by liver biopsy; and			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
2 Patient not requiring a liver transplant (bilirubin > 100 μ mol/l; decompensated cirrhosis.			
Initiation — Pregnancy			
Patient diagnosed with cholestasis of pregnancy.			
Initiation — Haematological transplant			
Both:			
1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogeneic stem cell or bone marrow transplantation; and			
2 Treatment for up to 13 weeks.			
Initiation — Total parenteral nutrition induced cholestasis			
Both:			
1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by TPN; and			
2 Liver function has not improved with modifying the TPN composition.			

Laxatives

Bowel-Cleansing Preparations

CITRIC ACID WITH MAGNESIUM OXIDE AND SODIUM PICOSULFATE			
Powder for oral soln 12 g with magnesium oxide 3.5 g and sodium picosulfate 10 mg per sachet			<i>e.g. PicoPrep</i>
MACROGOL 3350 WITH ASCORBIC ACID, POTASSIUM CHLORIDE AND SODIUM CHLORIDE			
Powder for oral soln 755.68 mg with ascorbic acid 85.16 mg, potassium chloride 10.55 mg, sodium chloride 37.33 mg and sodium sulphate 80.62 mg per g, 210 g sachet			<i>e.g. Glycoprep-C</i>
Powder for oral soln 755.68 mg with ascorbic acid 85.16 mg, potassium chloride 10.55 mg, sodium chloride 37.33 mg and sodium sulphate 80.62 mg per g, 70 g sachet			<i>e.g. Glycoprep-C</i>
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE, SODIUM CHLORIDE AND SODIUM SULPHATE			
Powder for oral soln 59 g with potassium chloride 0.7425 g, sodium bicarbonate 1.685 g, sodium chloride 1.465 g and sodium sulphate 5.685 g per sachet	14.31	4	Klean Prep

Bulk-Forming Agents

ISPAGHULA (PSYLLIUM) HUSK			
Powder for oral soln – 1% DV Sep-13 to 2016	5.51	500 g	Konsyl-D
STERCULIA WITH FRANGULA – Restricted: For continuation only			
➔ Powder for oral soln			

Faecal Softeners

DOCUSATE SODIUM			
Tab 50 mg – 1% DV Jan-15 to 2017	2.31	100	Coloxyl
Tab 120 mg – 1% DV Jan-15 to 2017	3.13	100	Coloxyl
DOCUSATE SODIUM WITH SENNOSIDES			
Tab 50 mg with sennosides 8 mg	4.40	200	Laxsol
PARAFFIN			
Oral liquid 1 mg per ml			
Enema 133 ml			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
POLOXAMER			
Oral drops 10% – 1% DV Sep-14 to 2017	3.78	30 ml	Coloxyl
Osmotic Laxatives			
GLYCEROL			
Suppos 1.27 g			
Suppos 2.55 g			
Suppos 3.6 g – 1% DV Sep-15 to 2018	6.50	20	PSM
LACTULOSE			
Oral liq 10 g per 15 ml – 1% DV May-14 to 2016	3.84	500 ml	Laevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE AND SODIUM CHLORIDE – Restricted see terms below			
⚡ Powder for oral soln 6.563 g with potassium chloride 23.3 mg, sodium bicarbonate 89.3 mg and sodium chloride 175.4 mg			
⚡ Powder for oral soln 13.125 g with potassium chloride 46.6 mg, sodium bicarbonate 178.5 mg and sodium chloride 350.7 mg – 1% DV Oct-14 to 2017	7.65	30	Lax-Sachets
➡ Restricted			
Initiation			
Either:			
1 Both:			
1.1 The patient has problematic constipation despite an adequate trial of other oral pharmacotherapies including lactulose where lactulose is not contraindicated; and			
1.2 The patient would otherwise require a per rectal preparation; or			
2 For short-term use for faecal disimpaction.			
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE			
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml – 1% DV Sep-13 to 2016	19.95	50	Micolette
SODIUM PHOSPHATE WITH PHOSPHORIC ACID			
Oral liq 16.4% with phosphoric acid 25.14%			
Enema 10% with phosphoric acid 6.58%	2.50	1	Fleet Phosphate Enema
Stimulant Laxatives			
BISACODYL			
Tab 5 mg – 1% DV Oct-15 to 2018	5.99	200	Lax-Tabs
Suppos 10 mg – 1% DV Jan-16 to 2018	3.78	10	Lax-Suppositories
SENNOSIDES			
Tab 7.5 mg			
Metabolic Disorder Agents			
ARGININE			
Powder			
Inj 600 mg per ml, 25 ml vial			
BETAINE – Restricted see terms below			
⚡ Powder			
➡ Restricted			
Metabolic physician or metabolic disorders dietitian			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

BIOTIN – **Restricted** see terms below

- ⚡ Cap 50 mg
- ⚡ Cap 100 mg
- ⚡ Inj 10 mg per ml, 5 ml vial

➡ **Restricted**

Metabolic physician or metabolic disorders dietitian

GALSULFASE – **Restricted** see terms below

- | | | | |
|---|----------|---|------------------|
| ⚡ Inj 1 mg per ml, 5 ml vial – 1% DV May-16 to 2018 | 2,234.00 | 1 | Naglazyme |
|---|----------|---|------------------|

➡ **Restricted**

Initiation

Metabolic physician

Re-assessment required after 12 months

Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency confirmed by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI.

Continuation

Metabolic physician

Re-assessment required after 12 months

All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

HAEM ARGINATE

Inj 25 mg per ml, 10 ml ampoule

IMIGLUCERASE – **Restricted** see terms below

- ⚡ Inj 40 iu per ml, 5 ml vial
- ⚡ Inj 40 iu per ml, 10 ml vial

➡ **Restricted**

Initiation

Only for use in patients with approval by the Gaucher's Treatment Panel.

LEVOCARNITINE – **Restricted** see terms below

- ⚡ Cap 500 mg
- ⚡ Oral soln 1,100 mg per 15 ml
- ⚡ Inj 200 mg per ml, 5 ml vial

➡ **Restricted**

Neurologist, metabolic physician or metabolic disorders dietitian

PYRIDOXAL-5-PHOSPHATE – **Restricted** see terms below

- ⚡ Tab 50 mg

➡ **Restricted**

Neurologist, metabolic physician or metabolic disorders dietitian

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SODIUM BENZOATE			
Cap 500 mg			
Powder			
Soln 100 mg per ml			
Inj 20%, 10 ml ampoule			
SODIUM PHENYLBUTYRATE			
Tab 500 mg			
Oral liq 250 mg per ml			
Inj 200 mg per ml, 10 ml ampoule			
TRIENTINE DIHYDROCHLORIDE			
Cap 300 mg			
Minerals			
Calcium			
CALCIUM CARBONATE			
Tab 1.25 g (500 mg elemental) – 1% DV Sep-14 to 2017	5.38	250	Arrow-Calcium
Tab eff 1.75 g (1 g elemental)	6.21	30	Calsource
Fluoride			
SODIUM FLUORIDE			
Tab 1.1 mg (0.5 mg elemental)			
Iodine			
POTASSIUM IODATE			
Tab 253 mcg (150 mcg elemental iodine) – 1% DV Dec-14 to 2017	3.65	90	NeuroTabs
POTASSIUM IODATE WITH IODINE			
Oral liq 10% with iodine 5%			
Iron			
FERRIC CARBOXYMALTOSIDE – Restricted see terms below			
⚡ Inj 50 mg per ml, 10 ml vial	150.00	1	Ferinject
➡ Restricted			
Initiation			
Treatment with oral iron has proven ineffective or is clinically inappropriate.			
FERROUS FUMARATE			
Tab 200 mg (65 mg elemental) – 1% DV Jun-15 to 2018	2.89	100	Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID			
Tab 310 mg (100 mg elemental) with folic acid 350 mcg	4.75	60	Ferro-F-Tabs
FERROUS GLUCONATE WITH ASCORBIC ACID			
Tab 170 mg (20 mg elemental) with ascorbic acid 40 mg			
FERROUS SULPHATE			
Tab long-acting 325 mg (105 mg elemental)	2.06	30	Ferrograd
Oral liq 30 mg (6 mg elemental) per ml – 1% DV Apr-14 to 2016	10.28	500 ml	Ferodan
FERROUS SULPHATE WITH ASCORBIC ACID			
Tab long-acting 325 mg (105 mg elemental) with ascorbic acid 500 mg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FERROUS SULPHATE WITH FOLIC ACID			
Tab long-acting 325 mg (105 mg elemental) with folic acid 350 mcg			
IRON POLYMALTOSE			
Inj 50 mg per ml, 2 ml ampoule – 1% DV Sep-14 to 2017	15.22	5	Ferrum H
IRON SUCROSE			
Inj 20 mg per ml, 5 ml ampoule	100.00	5	Venofer
Magnesium			
MAGNESIUM HYDROXIDE			
Tab 311 mg (130 mg elemental)			
MAGNESIUM OXIDE			
Cap 663 mg (400 mg elemental)			
MAGNESIUM SULPHATE			
Inj 0.4 mmol per ml, 250 ml bag			
Inj 2 mmol per ml, 5 ml ampoule – 1% DV Oct-14 to 2017	12.65	10	DBL
Zinc			
ZINC			
Oral liq 5 mg per 5 drops			
ZINC CHLORIDE			
Inj 5.3 mg per ml (5.1 mg per ml elemental), 2 ml ampoule			
ZINC SULPHATE			
Cap 137.4 mg (50 mg elemental) – 1% DV Mar-15 to 2017	11.00	100	Zincaps
Mouth and Throat			
Agents Used in Mouth Ulceration			
BENZYDAMINE HYDROCHLORIDE			
Soln 0.15%			
Spray 0.15%			
Spray 0.3%			
BENZYDAMINE HYDROCHLORIDE WITH CETYLPYRIDINIUM CHLORIDE			
Lozenge 3 mg with cetylpyridinium chloride			
CARBOXYMETHYLCELLULOSE			
Oral spray			
CHLORHEXIDINE GLUCONATE			
Mouthwash 0.2% – 1% DV Sep-15 to 2018	2.57	200 ml	healthE
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE			
Adhesive gel 8.7% with cetalkonium chloride 0.01%			
DICHLOROBENZYL ALCOHOL WITH AMYLMETACRESOL			
Lozenge 1.2 mg with amylmetacresol 0.6 mg			
SODIUM CARBOXYMETHYLCELLULOSE WITH PECTIN AND GELATINE			
Paste			
Powder			
TRIAMCINOLONE ACETONIDE			
Paste 0.1% – 1% DV Apr-15 to 2017	5.33	5 g	Kenalog in Orabase

Products with Hospital Supply Status (HSS) are in **bold**

Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Oropharyngeal Anti-Infectives			
AMPHOTERICIN B			
Lozenge 10 mg	5.86	20	Fungilin
MICONAZOLE			
Oral gel 20 mg per g – 1% DV Sep-15 to 2018	4.79	40 g	Decozol
NYSTATIN			
Oral liquid 100,000 u per ml – 1% DV Feb-16 to 2017	2.55	24 ml	m-Nystatin
Other Oral Agents			
SODIUM HYALURONATE [HYALURONIC ACID] – Restricted see terms below			
☞ Inj 20 mg per ml, 1 ml syringe			
☞ Restricted			
Otolaryngologist			
THYMOL GLYCERIN			
Compound, BPC – 1% DV Aug-16 to 2019	9.15	500 ml	PSM
Vitamins			
Multivitamin Preparations			
MULTIVITAMIN AND MINERAL SUPPLEMENT – Restricted see terms below			
☞ Cap	23.35	180	Clinicians Multivit & Mineral Boost
☞ Restricted			
Initiation			
Limited to 3 months treatment			
Both:			
1 Patient was admitted to hospital with burns; and			
2 Any of the following:			
2.1 Burn size is greater than 15% of total body surface area (BSA) for all types of burns; or			
2.2 Burn size is greater than 10% of BSA for mid-dermal or deep dermal burns; or			
2.3 Nutritional status prior to admission or dietary intake is poor.			
MULTIVITAMIN RENAL – Restricted see terms below			
☞ Cap	8.39	30	Clinicians Renal Vit
☞ Restricted			
Initiation			
Either:			
1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or			
2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73m ² body surface area (BSA).			
MULTIVITAMINS			
Tab (BPC cap strength)			e.g. Mvite
☞ Cap vitamin A 2500 u, betacarotene 3 mg, cholecalciferol 11 mcg, al- pha tocopherol 150 u, phytomenadione 150 mcg, folic acid 0.2 mg, ascorbic acid 100 mg, thiamine 1.5 mg, pantothenic acid 12 mg, riboflavin 1.7 mg, niacin 20 mg, pyridoxine hydrochloride 1.9 mg, cyanocobalamin 3 mcg, zinc 7.5 mg and biotin 100 mcg			e.g. Vitabdeck

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔ Restricted			
Initiation			
Either:			
1 Patient has cystic fibrosis with pancreatic insufficiency; or			
2 Patient is an infant or child with liver disease or short gut syndrome.			
⚡ Powder vitamin A 4200 mcg with vitamin D 155.5 mcg, vitamin E 21.4 mg, vitamin C 400 mg, vitamin K1 166 mcg thiamine 3.2 mg, riboflavin 4.4 mg, niacin 35 mg, vitamin B6 3.4 mg, folic acid 303 mcg, vitamin B12 8.6 mcg, biotin 214 mcg, pantothenic acid 17 mg, choline 350 mg and inositol 700 mg			<i>e.g. Paediatric Seravit</i>
➔ Restricted			
Initiation			
Patient has inborn errors of metabolism.			
Inj thiamine hydrochloride 250 mg with riboflavin 4 mg and pyridoxine hydrochloride 50 mg, 5 ml ampoule (1) and inj ascorbic acid 500 mg with nicotinamide 160 mg and glucose 1000 mg, 5 ml ampoule (1)			<i>e.g. Pabrinex IV</i>
Inj thiamine hydrochloride 250 mg with riboflavin 4 mg and pyridoxine hydrochloride 50 mg, 5 ml ampoule (1) and inj ascorbic acid 500 mg with nicotinamide 160 mg, 2 ml ampoule (1)			<i>e.g. Pabrinex IM</i>
Inj thiamine hydrochloride 500 mg with riboflavin 8 mg and pyridoxine hydrochloride 100 mg, 10 ml ampoule (1) and inj ascorbic acid 1000 mg with nicotinamide 320 mg and glucose 2000 mg, 10 ml ampoule (1)			<i>e.g. Pabrinex IV</i>
VITAMIN A WITH VITAMINS D AND C			
Soln 1,000 u with vitamin D 400 u and ascorbic acid 30 mg per 10 drops			<i>e.g. Vitadol C</i>
Vitamin A			
RETINOL			
Tab 10,000 iu			
Cap 25,000 iu			
Oral liq 150,000 iu per ml			
Vitamin B			
HYDROXOCOBALAMIN			
Inj 1 mg per ml, 1 ml ampoule – 1% DV Sep-15 to 2018	2.31	3	Neo-B12
PYRIDOXINE HYDROCHLORIDE			
Tab 25 mg – 1% DV Apr-15 to 2017	2.15	90	Vitamin B6 25
Tab 50 mg – 1% DV Oct-14 to 2017	11.55	500	Apo-Pyridoxine
Inj 100 mg per ml, 1 ml ampoule			
THIAMINE HYDROCHLORIDE			
Tab 50 mg			
Tab 100 mg			
Inj 100 mg per ml, 1 ml vial			<i>e.g. Benerva</i>
Inj 100 mg per ml, 2 ml vial			
VITAMIN B COMPLEX			
Tab strong, BPC			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Vitamin C			
ASCORBIC ACID			
Tab 100 mg – 1% DV Nov-13 to 2016	7.00	500	Cvite
Tab chewable 250 mg			
Vitamin D			
ALFACALCIDOL			
Cap 0.25 mcg	26.32	100	One-Alpha
Cap 1 mcg	87.98	100	One-Alpha
Oral drops 2 mcg per ml			
CALCITRIOL			
Cap 0.25 mcg – 1% DV Aug-16 to 2019.....	3.03	30	Airflow
	9.95	100	Calcitriol-AFT
Cap 0.5 mcg – 1% DV Aug-16 to 2019.....	5.62	30	Airflow
	18.39	100	Calcitriol-AFT
Oral liq 1 mcg per ml			
Inj 1 mcg per ml, 1 ml ampoule			
<i>(Airflow Cap 0.25 mcg to be delisted 1 August 2016)</i>			
<i>(Airflow Cap 0.5 mcg to be delisted 1 August 2016)</i>			
CHOLECALCIFEROL			
Cap 1.25 mg (50,000 iu)	3.85	12	Vit.D3

Vitamin E

ALPHA TOCOPHERYL ACETATE – **Restricted** see terms below

- ☞ Cap 100 u
- ☞ Cap 500 u
- ☞ Oral liq 156 u per ml

☞ **Restricted**

Initiation — Cystic fibrosis

Both:

- 1 Cystic fibrosis patient; and
- 2 Either:
 - 2.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck); or
 - 2.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for the patient.

Initiation — Osteoradionecrosis

For the treatment of osteoradionecrosis.

Initiation — Other indications

All of the following:

- 1 Infant or child with liver disease or short gut syndrome; and
- 2 Requires vitamin supplementation; and
- 3 Either:
 - 3.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplements (Vitabdeck); or
 - 3.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for patient.

Price (ex man. excl. GST)	Brand or Generic
\$	Per Manufacturer

Antianaemics

Hypoplastic and Haemolytic

EPOETIN ALFA [ERYTHROPOIETIN ALFA] – **Restricted** see terms below

⚡ Inj 1,000 iu in 0.5 ml syringe – 5% DV Mar-15 to 28 Feb 2018	48.68	6	Eprex
⚡ Inj 2,000 iu in 0.5 ml syringe – 5% DV Mar-15 to 28 Feb 2018	120.18	6	Eprex
⚡ Inj 3,000 iu in 0.3 ml syringe – 5% DV Mar-15 to 28 Feb 2018	166.87	6	Eprex
⚡ Inj 4,000 iu in 0.4 ml syringe – 5% DV Mar-15 to 28 Feb 2018	193.13	6	Eprex
⚡ Inj 5,000 iu in 0.5 ml syringe – 5% DV Mar-15 to 28 Feb 2018	243.26	6	Eprex
⚡ Inj 6,000 iu in 0.6 ml syringe – 5% DV Mar-15 to 28 Feb 2018	291.92	6	Eprex
⚡ Inj 8,000 iu in 0.8 ml syringe – 5% DV May-15 to 28 Feb 2018	352.69	6	Eprex
⚡ Inj 10,000 iu in 1 ml syringe – 5% DV Mar-15 to 28 Feb 2018	395.18	6	Eprex
⚡ Inj 40,000 iu in 1 ml syringe – 5% DV May-15 to 28 Feb 2018	263.45	1	Eprex

→ Restricted

Initiation — chronic renal failure

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin \leq 100g/L; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate \leq 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate \leq 45ml/min; and
- 4 Patient is on haemodialysis or peritoneal dialysis.

Initiation — myelodysplasia*

Re-assessment required after 2 months

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS); and
- 2 Has had symptomatic anaemia with haemoglobin $<$ 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification-based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum erythropoietin level of $<$ 500 IU/L; and
- 6 The minimum necessary dose of erythropoietin would be used and will not exceed 80,000 iu per week.

Continuation — myelodysplasia*

Re-assessment required after 12 months

All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of erythropoietin would be used and will not exceed 80,000 iu per week.

Initiation — all other indications

Haematologist

For use in patients where blood transfusion is not a viable treatment alternative.

Note: Indications marked with * are Unapproved Indications

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

EPOETIN BETA [ERYTHROPOIETIN BETA] – **Restricted** see terms below

Note: Epoetin beta is considered a Discretionary Variance Pharmaceutical for epoetin alfa.

- ⚡ Inj 2,000 iu in 0.3 ml syringe
- ⚡ Inj 3,000 iu in 0.3 ml syringe
- ⚡ Inj 4,000 iu in 0.3 ml syringe
- ⚡ Inj 5,000 iu in 0.3 ml syringe
- ⚡ Inj 6,000 iu in 0.3 ml syringe
- ⚡ Inj 10,000 iu in 0.6 ml syringe

➡ **Restricted**

Initiation — chronic renal failure

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin \leq 100g/L; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate \leq 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate \leq 45ml/min; and
- 4 Patient is on haemodialysis or peritoneal dialysis.

Initiation — myelodysplasia*

Re-assessment required after 12 months

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS); and
- 2 Has had symptomatic anaemia with haemoglobin $<$ 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification-based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum erythropoietin level of $<$ 500 IU/L; and
- 6 The minimum necessary dose of erythropoietin would be used and will not exceed 80,000 iu per week.

Continuation — myelodysplasia*

Re-assessment required after 2 months

All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of erythropoietin would be used and will not exceed 80,000 iu per week.

Initiation — all other indications

Haematologist.

For use in patients where blood transfusion is not a viable treatment alternative.

*Note: Indications marked with * are Unapproved Indications.

Megaloblastic

FOLIC ACID

Tab 0.8 mg – 1% DV Oct-15 to 2018	20.60	1,000	Apo-Folic Acid
Tab 5 mg – 1% DV Oct-15 to 2018	10.92	500	Apo-Folic Acid
Oral liq 50 mcg per ml	24.00	25 ml	Biomed
Inj 5 mg per ml, 10 ml vial			

Price (ex man. excl. GST)	Brand or Generic
\$	Per Manufacturer

Antifibrinolytics, Haemostatics and Local Sclerosants

ALUMINIUM CHLORIDE – **Restricted** see terms below

☞ Topical soln 20% w/v

e.g. Driclor

☞ **Restricted**

Initiation

For use as a haemostatis agent.

APROTININ – **Restricted** see terms below

☞ Inj 10,000 kIU per ml (equivalent to 200 mg per ml), 50 ml vial

☞ **Restricted**

Initiation

Cardiac anaesthetist

Either:

- 1 Paediatric patient undergoing cardiopulmonary bypass procedure; or
- 2 Adult patient undergoing cardiac surgical procedure where the significant risk of massive bleeding outweighs the potential adverse effects of the drug.

ELTROMBOPAG – **Restricted** see terms below

☞ Tab 25 mg	1,771.00	28	Revolade
☞ Tab 50 mg	3,542.00	28	Revolade

☞ **Restricted**

Initiation — idiopathic thrombocytopenic purpura - post-splenectomy

Haematologist

Limited to 6 weeks treatment

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of \leq 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of \leq 10,000 platelets per microlitre.

Initiation — (idiopathic thrombocytopenic purpura - preparation for splenectomy)

Haematologist

Limited to 6 weeks treatment

The patient requires eltrombopag treatment as preparation for splenectomy.

Continuation — (idiopathic thrombocytopenic purpura - post-splenectomy)

Haematologist

Re-assessment required after 12 months

The patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of $>$ 30,000 platelets per microlitre

FERRIC SUBSULFATE

Gel 25.9%

Soln 500 ml

POLIDOCANOL

Inj 0.5%, 30 ml vial

SODIUM TETRADECYL SULPHATE

Inj 3%, 2 ml ampoule

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
THROMBIN Powder			
TRANEXAMIC ACID			
Tab 500 mg – 1% DV Oct-14 to 2016	23.00	100	Cyklokapron
Inj 100 mg per ml, 5 ml ampoule – 1% DV Sep-15 to 2018	55.00	10	Cyklokapron

Blood Factors

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] – **Restricted** see terms below

☞ Inj 1 mg syringe	1,163.75	1	NovoSeven RT
☞ Inj 2 mg syringe	2,327.50	1	NovoSeven RT
☞ Inj 5 mg syringe	5,818.75	1	NovoSeven RT
☞ Inj 8 mg syringe	9,310.00	1	NovoSeven RT

☞ **Restricted**

Initiation

When used in the treatment of haemophilia, access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

FACTOR EIGHT INHIBITOR BYPASSING FRACTION – **Restricted** see terms below

☞ Inj 500 U	1,450.00	1	FEIBA NF
☞ Inj 1,000 U	2,900.00	1	FEIBA NF
☞ Inj 2,500 U	7,250.00	1	FEIBA NF

☞ **Restricted**

Initiation

When used in the treatment of haemophilia, access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] – **Restricted** see terms below

☞ Inj 250 iu prefilled syringe	210.00	1	Xyntha
☞ Inj 500 iu prefilled syringe	420.00	1	Xyntha
☞ Inj 1,000 iu prefilled syringe	840.00	1	Xyntha
☞ Inj 2,000 iu prefilled syringe	1,680.00	1	Xyntha
☞ Inj 3,000 iu prefilled syringe	2,520.00	1	Xyntha

☞ **Restricted**

Initiation

Note: Preferred Brand of recombinant factor VIII from 1 March 2016 until 28 February 2019. When used in the treatment of haemophilia, funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

NONACOG ALFA [RECOMBINANT FACTOR IX] – **Restricted** see terms below

☞ Inj 250 iu vial	310.00	1	BeneFIX
☞ Inj 500 iu vial	620.00	1	BeneFIX
☞ Inj 1,000 iu vial	1,240.00	1	BeneFIX
☞ Inj 2,000 iu vial	2,480.00	1	BeneFIX
☞ Inj 3,000 iu vial	3,720.00	1	BeneFIX

☞ **Restricted**

Initiation

When used in the treatment of haemophilia, access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
NONACOG GAMMA, [RECOMBINANT FACTOR IX] – Restricted see terms below			
⌘ Inj 250 iu vial	287.50	1	RIXUBIS
⌘ Inj 500 iu vial	575.00	1	RIXUBIS
⌘ Inj 1,000 iu vial	1,150.00	1	RIXUBIS
⌘ Inj 2,000 iu vial	2,300.00	1	RIXUBIS
⌘ Inj 3,000 iu vial	3,450.00	1	RIXUBIS

➔ **Restricted**

Initiation

When used in the treatment of haemophilia, treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) – Restricted see terms below

⌘ Inj 250 iu vial	287.50	1	Advate
⌘ Inj 500 iu vial	575.00	1	Advate
⌘ Inj 1,000 iu vial	1,150.00	1	Advate
⌘ Inj 1,500 iu vial	1,725.00	1	Advate
⌘ Inj 2,000 iu vial	2,300.00	1	Advate
⌘ Inj 3,000 iu vial	3,450.00	1	Advate

➔ **Restricted**

Initiation

Notes: Rare Clinical Circumstances Brand of recombinant factor VIII from 1 March 2016 until 28 February 2019. When used in the treatment of haemophilia, access to funded treatment by application to the Haemophilia Treatments Panel. Application details may be obtained from PHARMAC.s website <http://www.pharmac.govt.nz> or:

The Co-ordinator, Haemophilia Treatments Panel	Phone: 0800 023 588 Option 2
PHARMAC PO Box 10 254	Facsimile: (04) 974 4881
Wellington	Email: haemophilia@pharmac.govt.nz

OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE FS) – Restricted see terms below

⌘ Inj 250 iu vial	237.50	1	Kogenate FS
⌘ Inj 500 iu vial	475.00	1	Kogenate FS
⌘ Inj 1,000 iu vial	950.00	1	Kogenate FS
⌘ Inj 2,000 iu vial	1,900.00	1	Kogenate FS
⌘ Inj 3,000 iu vial	2,850.00	1	Kogenate FS

➔ **Restricted**

Initiation

Notes: Second Brand of recombinant factor VIII from 1 March 2016 until 28 February 2019. When used in the treatment of haemophilia, access to funded treatment by application to the Haemophilia Treatments Panel. Application details may be obtained from PHARMAC.s website <http://www.pharmac.govt.nz> or:

The Co-ordinator, Haemophilia Treatments Panel	Phone: 0800 023 588 Option 2
PHARMAC PO Box 10 254	Facsimile: (04) 974 4881
Wellington	Email: haemophilia@pharmac.govt.nz

Vitamin K

PHYTOMENADIONE

Inj 2 mg in 0.2 ml ampoule	8.00	5	Konaktion MM
Inj 10 mg per ml, 1 ml ampoule	9.21	5	Konaktion MM

Price
(ex man. excl. GST)
\$ Per Brand or
Generic
Manufacturer

Antithrombotics

Anticoagulants

BIVALIRUDIN – **Restricted** see terms below

¶ Inj 250 mg vial

➡ **Restricted**

Initiation

Either:

- 1 For use in heparin-induced thrombocytopenia, heparin resistance or heparin intolerance; or
- 2 For use in patients undergoing endovascular procedures.

DABIGATRAN

Cap 75 mg	148.00	60	Pradaxa
Cap 110 mg	148.00	60	Pradaxa
Cap 150 mg	148.00	60	Pradaxa

DALTEPARIN

Inj 2,500 iu in 0.2 ml syringe	19.97	10	Fragmin
Inj 5,000 iu in 0.2 ml syringe	39.94	10	Fragmin
Inj 7,500 iu in 0.75 ml syringe	60.03	10	Fragmin
Inj 10,000 iu in 1 ml syringe	77.55	10	Fragmin
Inj 12,500 iu in 0.5 ml syringe	99.96	10	Fragmin
Inj 15,000 iu in 0.6 ml syringe	120.05	10	Fragmin
Inj 18,000 iu in 0.72 ml syringe	158.47	10	Fragmin

DANAPAROID – **Restricted** see terms below

¶ Inj 750 u in 0.6 ml ampoule

➡ **Restricted**

Initiation

For use in heparin-induced thrombocytopenia, heparin resistance or heparin intolerance.

DEFIBROTIDE – **Restricted** see terms below

¶ Inj 80 mg per ml, 2.5 ml ampoule

➡ **Restricted**

Initiation

Haematologist

Patient has moderate or severe sinusoidal obstruction syndrome as a result of chemotherapy or regimen-related toxicities.

DEXTROSE WITH SODIUM CITRATE AND CITRIC ACID [ACID CITRATE DEXTROSE A]

Inj 24.5 mg with sodium citrate 22 mg and citric acid 7.3 mg per ml,
100 ml bag

ENOXAPARIN

Inj 20 mg in 0.2 ml syringe	37.24	10	Clexane
Inj 40 mg in 0.4 ml ampoule			
Inj 40 mg in 0.4 ml syringe	49.69	10	Clexane
Inj 60 mg in 0.6 ml syringe	74.91	10	Clexane
Inj 80 mg in 0.8 ml syringe	99.86	10	Clexane
Inj 100 mg in 1 ml syringe	125.06	10	Clexane
Inj 120 mg in 0.8 ml syringe	155.40	10	Clexane
Inj 150 mg in 1 ml syringe	177.60	10	Clexane

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FONDAPARINUX SODIUM – Restricted see terms below			
¶ Inj 2.5 mg in 0.5 ml syringe			
¶ Inj 7.5 mg in 0.6 ml syringe			
➔ Restricted			
Initiation			
For use in heparin-induced thrombocytopaenia, heparin resistance or heparin intolerance.			
HEPARIN SODIUM			
Inj 100 iu per ml, 250 ml bag			
Inj 1,000 iu per ml, 1 ml ampoule	66.80	50	Hospira
Inj 1,000 iu per ml, 35 ml vial			
Inj 1,000 iu per ml, 5 ml ampoule	61.04	50	Pfizer
Inj 5,000 iu in 0.2 ml ampoule			
Inj 5,000 iu per ml, 1 ml ampoule	14.20	5	Hospira
Inj 5,000 iu per ml, 5 ml ampoule	236.60	50	Pfizer
HEPARINISED SALINE			
Inj 10 iu per ml, 5 ml ampoule	39.00	50	Pfizer
Inj 100 iu per ml, 2 ml ampoule			
Inj 100 iu per ml, 5 ml ampoule			
PHENINDIONE			
Tab 10 mg			
Tab 25 mg			
Tab 50 mg			
PROTAMINE SULPHATE			
Inj 10 mg per ml, 5 ml ampoule			
RIVAROXABAN – Restricted see terms below			
¶ Tab 10 mg	153.00	15	Xarelto
➔ Restricted			
Initiation — total hip replacement			
<i>Limited to 5 weeks treatment</i>			
For the prophylaxis of venous thromboembolism.			
Initiation — total knee replacement			
<i>Limited to 2 weeks treatment</i>			
For the prophylaxis of venous thromboembolism.			
SODIUM CITRATE WITH SODIUM CHLORIDE AND POTASSIUM CHLORIDE			
Inj 4.2 mg with sodium chloride 5.7 mg and potassium chloride 74.6 mcg per ml, 5,000 ml bag			
TRISODIUM CITRATE			
Inj 4%, 5 ml ampoule			
Inj 46.7%, 3 ml syringe			
Inj 46.7%, 5 ml ampoule			
WARFARIN SODIUM			
Tab 1 mg	6.86	100	Marevan
Tab 2 mg			
Tab 3 mg	9.70	100	Marevan
Tab 5 mg	11.75	100	Marevan

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Antiplatelets			
ASPIRIN			
Tab 100 mg – 1% DV Mar-14 to 2016	1.60	90	Ethics Aspirin EC
Suppos 300 mg	10.50	990	Ethics Aspirin EC
CLOPIDOGREL			
Tab 75 mg – 1% DV Dec-13 to 2016	5.48	84	Arrow - Clopid
DIPYRIDAMOLE			
Tab 25 mg			
Tab long-acting 150 mg	11.52	60	Pytazen SR
Inj 5 mg per ml, 2 ml ampoule			
EPTIFIBATIDE – Restricted see terms below			
☞ Inj 2 mg per ml, 10 ml vial	111.00	1	Integrilin
☞ Inj 750 mcg per ml, 100 ml vial	324.00	1	Integrilin
☞ Restricted			
Initiation			
Either:			
1 For use in patients with acute coronary syndromes undergoing percutaneous coronary intervention; or			
2 For use in patients with definite or strongly suspected intra-coronary thrombus on coronary angiography.			
PRASUGREL – Restricted see terms below			
☞ Tab 5 mg	108.00	28	Effient
☞ Tab 10 mg	120.00	28	Effient
☞ Restricted			
Initiation — Bare metal stents			
<i>Limited to 6 months treatment</i>			
Patient has undergone coronary angioplasty in the previous 4 weeks and is clopidogrel-allergic.			
Initiation — Drug-eluting stents			
<i>Limited to 12 months treatment</i>			
Patient has had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic.			
Initiation — Stent thrombosis			
Patient has experienced cardiac stent thrombosis whilst on clopidogrel.			
Initiation — Myocardial infarction			
<i>Limited to 1 week treatment</i>			
For short term use while in hospital following ST-elevated myocardial infarction.			
Note: Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment			
TICAGRELOR – Restricted see terms below			
☞ Tab 90 mg	90.00	56	Brilinta
☞ Restricted			
Initiation			
Restricted to treatment of acute coronary syndromes specifically for patients who have recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome, and in whom fibrinolytic therapy has not been given in the last 24 hours and is not planned.			
TICLOPIDINE			
Tab 250 mg			

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Fibrinolytic Agents

ALTEPLASE

- Inj 2 mg vial
- Inj 10 mg vial
- Inj 50 mg vial

TENECTEPLASE

- Inj 50 mg vial

UROKINASE

- Inj 10,000 iu vial
- Inj 50,000 iu vial
- Inj 100,000 iu vial
- Inj 500,000 iu vial

Colony-Stimulating Factors

Granulocyte Colony-Stimulating Factors

FILGRASTIM – **Restricted** see terms below

⚡ Inj 300 mcg in 0.5 ml prefilled syringe	270.00	5	Zarzio
⚡ Inj 300 mcg in 1 ml vial	650.00	5	Neupogen
⚡ Inj 480 mcg in 0.5 ml prefilled syringe	432.00	5	Zarzio

➡ **Restricted**

Haematologist or oncologist

PEGFILGRASTIM – **Restricted** see terms below

⚡ Inj 6 mg per 0.6 ml syringe	1,080.00	1	Neulastim
-------------------------------------	----------	---	-----------

➡ **Restricted**

Initiation

For prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk \geq 20%).

Note: *Febrile neutropenia risk \geq 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines

Fluids and Electrolytes

Intravenous Administration

CALCIUM CHLORIDE

- Inj 100 mg per ml, 10 ml vial

CALCIUM GLUCONATE

Inj 10%, 10 ml ampoule	34.24	10	Hospira
------------------------------	-------	----	---------

COMPOUND ELECTROLYTES

Inj sodium 140 mmol/l with potassium 5 mmol/l, magnesium 1.5 mmol/l, chloride 98 mmol/l, acetate 27 mmol/l and gluconate 23 mmol/l, bag	3.10	1,000 ml	Baxter
	5.00	500 ml	Baxter

COMPOUND ELECTROLYTES WITH GLUCOSE

Inj glucose 50 g with 140 mmol/l sodium, 5 mmol/l potassium, 1.5 mmol/l magnesium, 98 mmol/l chloride, 27 mmol/l acetate and 23 mmol/l gluconate, bag	7.00	1,000 ml	Baxter
---	------	----------	--------

BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
COMPOUND SODIUM LACTATE [HARTMANN'S SOLUTION]			
Inj sodium 131 mmol/l with potassium 5 mmol/l, calcium 2 mmol/l, bi-carbonate 29 mmol/l, chloride 111 mmol/l, bag	1.77	500 ml	Baxter
	1.80	1,000 ml	Baxter
COMPOUND SODIUM LACTATE WITH GLUCOSE			
Inj sodium 131 mmol/l with potassium 5 mmol/l, calcium 2 mmol/l, bi-carbonate 29 mmol/l, chloride 111 mmol/l and glucose 5%, bag	5.38	1,000 ml	Baxter
GLUCOSE [DEXTROSE]			
Inj 5%, bag	1.77	500 ml	Baxter
	1.80	1,000 ml	Baxter
	2.84	100 ml	Baxter
	2.87	50 ml	Baxter
	3.87	250 ml	Baxter
Inj 10%, bag	3.70	500 ml	Baxter
	5.29	1,000 ml	Baxter
Inj 50%, bag	6.84	500 ml	Baxter
Inj 50%, 10 ml ampoule – 1% DV Oct-14 to 2017	27.50	5	Biomed
Inj 50%, 90 ml bottle – 1% DV Oct-14 to 2017	14.50	1	Biomed
Inj 70%, 1,000 ml bag			
Inj 70%, 500 ml bag			
GLUCOSE WITH POTASSIUM CHLORIDE			
Inj 5% glucose with 20 mmol/l potassium chloride, bag	7.36	1,000 ml	Baxter
Inj 5% glucose with 30 mmol/l potassium chloride, 1,000 ml bag			
Inj 10% glucose with 10 mmol/l potassium chloride, 500 ml bag			
GLUCOSE WITH POTASSIUM CHLORIDE AND SODIUM CHLORIDE			
Inj 4% glucose with potassium chloride 20 mmol/l and sodium chloride 0.18%, bag	3.45	500 ml	Baxter
	4.30	1,000 ml	Baxter
Inj 4% glucose with potassium chloride 30 mmol/l and sodium chloride 0.18%, bag	3.62	1,000 ml	Baxter
Inj 2.5% glucose with potassium chloride 20 mmol/l and sodium chloride 0.45%, 3,000 ml bag			
Inj 10% glucose with potassium chloride 10 mmol/l and sodium chloride 15 mmol/l, 500 ml bag			
GLUCOSE WITH SODIUM CHLORIDE			
Inj glucose 2.5% with sodium chloride 0.45%, bag	4.95	500 ml	Baxter
Inj glucose 5% with sodium chloride 0.45%, bag	5.80	1,000 ml	Baxter
	9.87	500 ml	Baxter
Inj glucose 5% with sodium chloride 0.9%, bag	4.54	1,000 ml	Baxter
Inj glucose 5% with sodium chloride 0.2%, 500 ml bag			
POTASSIUM CHLORIDE			
Inj 75 mg (1 mmol) per ml, 10 ml ampoule			
Inj 225 mg (3 mmol) per ml, 20 ml ampoule			

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
POTASSIUM CHLORIDE WITH SODIUM CHLORIDE			
Inj 20 mmol/l potassium chloride with 0.9% sodium chloride, bag	3.85	1,000 ml	Baxter
Inj 30 mmol/l potassium chloride with 0.9% sodium chloride, bag	2.59	1,000 ml	Baxter
Inj 40 mmol/l potassium chloride with 0.9% sodium chloride, bag	6.62	1,000 ml	Baxter
Inj 10 mmol potassium chloride with 0.29% sodium chloride, 100 ml bag			
Inj 40 mmol potassium chloride with 0.9% sodium chloride, 100 ml bag			
POTASSIUM DIHYDROGEN PHOSPHATE			
Inj 1 mmol per ml, 10 ml ampoule – 1% DV Oct-15 to 2018	151.80	10	Hospira
RINGER'S SOLUTION			
Inj sodium 147 mmol/l with potassium 4 mmol/l, calcium 2.2 mmol/l, chloride 156 mmol/l, bag	5.13	1,000 ml	Baxter
SODIUM ACETATE			
Inj 4 mmol per ml, 20 ml ampoule			
SODIUM BICARBONATE			
Inj 8.4%, 10 ml vial			
Inj 8.4%, 50 ml vial	19.95	1	Biomed
Inj 8.4%, 100 ml vial	20.50	1	Biomed
SODIUM CHLORIDE			
Inj 0.45%, bag	5.50	500 ml	Baxter
Inj 0.9%, bag	1.77	500 ml	Baxter
	1.80	1,000 ml	Baxter
	2.28	100 ml	Baxter
	3.01	50 ml	Baxter
	3.60	250 ml	Baxter
	1.70	500 ml	Freeflex
	1.71	1,000 ml	Freeflex
Inj 3%, bag	5.69	1,000 ml	Baxter
Inj 0.9%, 5 ml ampoule	10.85	50	Multichem
	15.50		Pfizer
Inj 0.9%, 10 ml ampoule	11.50	50	Multichem
	15.50		Pfizer
⚡ Inj 0.9%, 3 ml syringe, non-sterile pack – 1% DV Jun-15 to 2018	10.65	30	BD PosiFlush
➡Restricted			
Initiation			
For use in flushing of in-situ vascular access devices only.			
⚡ Inj 0.9%, 5 ml syringe, non-sterile pack – 1% DV Jun-15 to 2018	10.80	30	BD PosiFlush
➡Restricted			
Initiation			
For use in flushing of in-situ vascular access devices only.			
⚡ Inj 0.9%, 10 ml syringe, non-sterile pack – 1% DV Jun-15 to 2018	11.25	30	BD PosiFlush
➡Restricted			
Initiation			
For use in flushing of in-situ vascular access devices only.			
Inj 0.9%, 20 ml ampoule	8.41	20	Multichem
Inj 23.4% (4 mmol/ml), 20 ml – 1% DV Sep-13 to 2016	31.25	5	Biomed
Inj 1.8%, 500 ml bottle			
SODIUM DIHYDROGEN PHOSPHATE [SODIUM ACID PHOSPHATE]			
Inj 1 mmol per ml, 20 ml ampoule – 1% DV Oct-15 to 2018	47.50	5	Biomed

BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
WATER			
Inj, bag	2.75	1,000 ml	Baxter
Inj 5 ml ampoule	10.25	50	Multichem
Inj 10 ml ampoule	11.25	50	Multichem
Inj 20 ml ampoule	6.50	20	Multichem
Inj 250 ml bag			
Inj 500 ml bag			
Oral Administration			
CALCIUM POLYSTYRENE SULPHONATE			
Powder	169.85	300 g	Calcium Resonium
COMPOUND ELECTROLYTES			
Powder for oral soln			
COMPOUND ELECTROLYTES WITH GLUCOSE			
Soln with electrolytes			
PHOSPHORUS			
Tab eff 500 mg (16 mmol)			
POTASSIUM CHLORIDE			
Tab eff 548 mg (14 mmol) with chloride 285 mg (8 mmol)			
Tab long-acting 600 mg (8 mmol) – 1% DV Sep-15 to 2018	7.42	200	Span-K
Oral liq 2 mmol per ml			
SODIUM BICARBONATE			
Cap 840 mg	8.52	100	Sodibic
SODIUM CHLORIDE			
Tab 600 mg			
Oral liq 2 mmol/ml			
SODIUM POLYSTYRENE SULPHONATE			
Powder – 1% DV Sep-15 to 2018	84.65	454 g	Resonium A
Plasma Volume Expanders			
GELATINE, SUCCINYLATED			
Inj 4%, 500 ml bag	108.00	10	Gelofusine
HYDROXYETHYL STARCH 130/0.4 WITH MAGNESIUM CHLORIDE, POTASSIUM CHLORIDE, SODIUM ACETATE AND SODIUM CHLORIDE			
Inj 6% with magnesium chloride 0.03%, potassium chloride 0.03%, sodium acetate 0.463% and sodium chloride 0.6%, 500 ml bag	198.00	20	Volulyte 6%
HYDROXYETHYL STARCH 130/0.4 WITH SODIUM CHLORIDE			
Inj 6% with sodium chloride 0.9%, 500 ml bag	198.00	20	Voluven

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

CAPTOPRIL

☞ Oral liq 5 mg per ml	94.99	95 ml	Capoten
------------------------------	-------	-------	---------

☞ Restricted

Initiation

Any of the following:

- 1 For use in children under 12 years of age; or
- 2 For use in tube-fed patients; or
- 3 For management of rebound transient hypertension following cardiac surgery.

CILAZAPRIL

Tab 0.5 mg – 1% DV Sep-13 to 2016	2.00	90	Zapril
Tab 2.5 mg – 1% DV Sep-13 to 2016	4.31	90	Zapril
Tab 5 mg – 1% DV Sep-13 to 2016	6.98	90	Zapril

ENALAPRIL MALEATE

Tab 5 mg – 1% DV Sep-15 to 2018	0.96	100	Ethics Enalapril
Tab 10 mg – 1% DV Sep-15 to 2018	1.24	100	Ethics Enalapril
Tab 20 mg – 1% DV Sep-15 to 2018	1.78	100	Ethics Enalapril

LISINOPRIL

Tab 5 mg – 1% DV Jan-16 to 2018	1.80	90	Ethics Lisinopril
Tab 10 mg – 1% DV Jan-16 to 2018	2.05	90	Ethics Lisinopril
Tab 20 mg – 1% DV Jan-16 to 2018	2.76	90	Ethics Lisinopril

PERINDOPRIL

Tab 2 mg – 1% DV Oct-14 to 2017	3.75	30	Apo-Perindopril
Tab 4 mg – 1% DV Oct-14 to 2017	4.80	30	Apo-Perindopril

QUINAPRIL

Tab 5 mg – 1% DV Sep-15 to 2018	4.31	90	Arrow-Quinapril 5
Tab 10 mg – 1% DV Sep-15 to 2018	3.15	90	Arrow-Quinapril 10
Tab 20 mg – 1% DV Sep-15 to 2018	5.97	90	Arrow-Quinapril 20

TRANDOLAPRIL – **Restricted:** For continuation only

☞ Cap 1 mg

☞ Cap 2 mg

ACE Inhibitors with Diuretics

CILAZAPRIL WITH HYDROCHLOROTHIAZIDE

Tab 5 mg with hydrochlorothiazide 12.5 mg – 1% DV Mar-14 to 2016	10.72	100	Apo-Cilazapril/ Hydrochlorothiazide
--	-------	-----	--

ENALAPRIL MALEATE WITH HYDROCHLOROTHIAZIDE – **Restricted:** For continuation only

☞ Tab 20 mg with hydrochlorothiazide 12.5 mg

QUINAPRIL WITH HYDROCHLOROTHIAZIDE

Tab 10 mg with hydrochlorothiazide 12.5 mg – 1% DV Oct-15 to 2018	3.65	30	Accuretic 10
Tab 20 mg with hydrochlorothiazide 12.5 mg – 1% DV Oct-15 to 2018	4.78	30	Accuretic 20

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Angiotensin II Antagonists			
CANDESARTAN CILEXETIL – Restricted see terms below			
⚡ Tab 4 mg – 1% DV Sep-15 to 2018	2.50	90	Candestar
⚡ Tab 8 mg – 1% DV Sep-15 to 2018	3.68	90	Candestar
⚡ Tab 16 mg – 1% DV Sep-15 to 2018	6.12	90	Candestar
⚡ Tab 32 mg – 1% DV Sep-15 to 2018	10.66	90	Candestar
➡ Restricted			
Initiation — ACE inhibitor intolerance			
Either:			
1 Patient has persistent ACE inhibitor induced cough that is not resolved by ACE inhibitor retrial (same or new ACE inhibitor); or			
2 Patient has a history of angioedema.			
Initiation — Unsatisfactory response to ACE inhibitor			
Patient is not adequately controlled on maximum tolerated dose of an ACE inhibitor.			
LOSARTAN POTASSIUM			
Tab 12.5 mg – 1% DV Jan-15 to 2017	1.55	84	Losartan Actavis
Tab 25 mg – 1% DV Jan-15 to 2017	1.90	84	Losartan Actavis
Tab 50 mg – 1% DV Jan-15 to 2017	2.25	84	Losartan Actavis
Tab 100 mg – 1% DV Jan-15 to 2017	2.60	84	Losartan Actavis
Angiotensin II Antagonists with Diuretics			
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE			
Tab 50 mg with hydrochlorothiazide 12.5 mg – 1% DV Oct-14 to 2017	2.18	30	Arrow-Losartan & Hydrochlorothiazide
Alpha-Adrenoceptor Blockers			
DOXAZOSIN			
Tab 2 mg – 1% DV Sep-14 to 2017	6.75	500	Apo-Doxazosin
Tab 4 mg – 1% DV Sep-14 to 2017	9.67	500	Apo-Doxazosin
PHENOXYBENZAMINE HYDROCHLORIDE			
Cap 10 mg			
Inj 50 mg per ml, 2 ml ampoule			
PHENTOLAMINE MESYLATE			
Inj 5 mg per ml, 1 ml ampoule			
Inj 10 mg per ml, 1 ml ampoule			
PRAZOSIN			
Tab 1 mg	5.53	100	Apo-Prazosin
Tab 2 mg	7.00	100	Apo-Prazosin
Tab 5 mg	11.70	100	Apo-Prazosin
TERAZOSIN			
Tab 1 mg – 1% DV Sep-13 to 2016	0.50	28	Arrow
Tab 2 mg – 1% DV Sep-13 to 2016	0.45	28	Arrow
Tab 5 mg – 1% DV Sep-13 to 2016	0.68	28	Arrow

Price (ex man. excl. GST)	Brand or Generic
\$ Per	Manufacturer

Antiarrhythmics

ADENOSINE

Inj 3 mg per ml, 2 ml vial

⚡ Inj 3 mg per ml, 10 ml vial

➡Restricted

Initiation

For use in cardiac catheterisation, electrophysiology and MRI.

AJMALINE – **Restricted** see terms below

⚡ Inj 5 mg per ml, 10 ml ampoule

➡Restricted

Cardiologist

AMIODARONE HYDROCHLORIDE

Tab 100 mg

Tab 200 mg

Inj 50 mg per ml, 3 ml ampoule – **1% DV Aug-13 to 2016** 22.80 6 **Cordarone-X**

ATROPINE SULPHATE

Inj 600 mcg per ml, 1 ml ampoule 71.00 50 AstraZeneca

DIGOXIN

Tab 62.5 mcg – **1% DV Jun-16 to 2019** 6.67 240 **Lanoxin PG**

Tab 250 mcg – **1% DV Apr-16 to 2019** 14.52 240 **Lanoxin**

Oral liq 50 mcg per ml

Inj 250 mcg per ml, 2 ml vial

DISOPYRAMIDE PHOSPHATE

Cap 100 mg

Cap 150 mg

FLECAINIDE ACETATE

Tab 50 mg 38.95 60 Tambocor

Cap long-acting 100 mg 38.95 30 Tambocor CR

Cap long-acting 200 mg 68.78 30 Tambocor CR

Inj 10 mg per ml, 15 ml ampoule 52.45 5 Tambocor

MEXILETINE HYDROCHLORIDE

Cap 150 mg 162.00 100 Mexiletine Hydrochloride
USP

Cap 250 mg 202.00 100 Mexiletine Hydrochloride
USP

PROPAFENONE HYDROCHLORIDE

Tab 150 mg

Antihypotensives

MIDODRINE – **Restricted** see terms below

⚡ Tab 2.5 mg

⚡ Tab 5 mg

➡Restricted

Initiation

Patient has disabling orthostatic hypotension not due to drugs.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Beta-Adrenoceptor Blockers			
ATENOLOL			
Tab 50 mg – 1% DV Sep-15 to 2018	4.61	500	Mylan Atenolol
Tab 100 mg – 1% DV Sep-15 to 2018	7.67	500	Mylan Atenolol
Oral liq 5 mg per ml	21.25	300 ml	Atenolol-AFT
BISOPROLOL FUMARATE			
Tab 2.5 mg – 1% DV Mar-15 to 2017	2.40	30	Bosvate
Tab 5 mg – 1% DV Mar-15 to 2017	3.50	30	Bosvate
Tab 10 mg – 1% DV Mar-15 to 2017	6.40	30	Bosvate
CARVEDILOL			
Tab 6.25 mg – 1% DV Jun-15 to 2017	3.90	60	Dicarz
Tab 12.5 mg – 1% DV Jun-15 to 2017	5.10	60	Dicarz
Tab 25 mg – 1% DV Jun-15 to 2017	6.30	60	Dicarz
CELIPROLOL			
Tab 200 mg	21.40	180	Celol
ESMOLOL HYDROCHLORIDE			
Inj 10 mg per ml, 10 ml vial			
LABETALOL			
Tab 50 mg	8.23	100	Hybloc
Tab 100 mg	10.06	100	Hybloc
Tab 200 mg	17.55	100	Hybloc
Tab 400 mg			
Inj 5 mg per ml, 20 ml ampoule			
METOPROLOL SUCCINATE			
Tab long-acting 23.75 mg – 1% DV Nov-16 to 2018	2.39	90	Metoprolol - AFT CR
Tab long-acting 47.5 mg – 1% DV Nov-16 to 2018	3.48	90	Metoprolol - AFT CR
Tab long-acting 95 mg – 1% DV Nov-16 to 2018	5.73	90	Metoprolol - AFT CR
Tab long-acting 190 mg – 1% DV Nov-16 to 2018	11.54	90	Metoprolol - AFT CR
METOPROLOL TARTRATE			
Tab 50 mg – 1% DV Aug-16 to 2018	4.64	100	Apo-Metoprolol
	16.00		Lopresor
Tab 100 mg – 1% DV Aug-16 to 2018	6.09	60	Apo-Metoprolol
	21.00		Lopresor
Tab long-acting 200 mg	18.00	28	Slow-Lopresor
Inj 1 mg per ml, 5 ml vial	24.00	5	Lopresor
<i>(Lopresor Tab 50 mg to be delisted 1 August 2016)</i>			
<i>(Lopresor Tab 100 mg to be delisted 1 August 2016)</i>			
NADOLOL			
Tab 40 mg – 1% DV Oct-15 to 2018	16.05	100	Apo-Nadolol
Tab 80 mg – 1% DV Oct-15 to 2018	24.70	100	Apo-Nadolol
PINDOLOL			
Tab 5 mg – 1% DV Nov-13 to 2016	9.72	100	Apo-Pindolol
Tab 10 mg – 1% DV Nov-13 to 2016	15.62	100	Apo-Pindolol
Tab 15 mg – 1% DV Nov-13 to 2016	23.46	100	Apo-Pindolol

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PROPRANOLOL			
Tab 10 mg	3.65	100	Apo-Propranolol
Tab 40 mg	4.65	100	Apo-Propranolol
Cap long-acting 160 mg	18.17	100	Cardinol LA
Oral liq 4 mg per ml			
Inj 1 mg per ml, 1 ml ampoule			
SOTALOL			
Tab 80 mg	27.50	500	Mylan
Tab 160 mg	10.50	100	Mylan
Inj 10 mg per ml, 4 ml ampoule	65.39	5	Sotacor
TIMOLOL MALEATE			
Tab 10 mg			
Calcium Channel Blockers			
Dihydropyridine Calcium Channel Blockers			
AMLODIPINE			
Tab 2.5 mg – 1% DV Feb-15 to 2017	2.21	100	Apo-Amlodipine
Tab 5 mg – 1% DV May-15 to 2017	5.04	250	Apo-Amlodipine
Tab 10 mg – 1% DV May-15 to 2017	7.21	250	Apo-Amlodipine
FELODIPINE			
Tab long-acting 2.5 mg – 1% DV Sep-15 to 2018	1.45	30	Plendil ER
Tab long-acting 5 mg – 1% DV Sep-15 to 2018	1.55	30	Plendil ER
Tab long-acting 10 mg – 1% DV Sep-15 to 2018	2.30	30	Plendil ER
ISRADIPINE			
Tab 2.5 mg			
Cap 2.5 mg			
Cap long-acting 2.5 mg			
Cap long-acting 5 mg			
NICARDIPINE HYDROCHLORIDE – Restricted see terms below			
⚡ Inj 2.5 mg per ml, 10 ml vial			
➡Restricted			
Initiation			
Anaesthetist, intensivist or paediatric cardiologist			
Both:			
1 Patient is a Paediatric Patient; and			
2 Any of the following:			
2.1 Patient has hypertension requiring urgent treatment with an intravenous agent; or			
2.2 Patient has excessive ventricular afterload; or			
2.3 Patient is awaiting or undergoing cardiac surgery using cardiopulmonary bypass.			
NIFEDIPINE			
Tab long-acting 10 mg			
Tab long-acting 20 mg	9.59	100	Nyefax Retard
Tab long-acting 30 mg – 1% DV Sep-14 to 2017	3.75	30	Adefin XL
Tab long-acting 60 mg – 1% DV Sep-14 to 2017	5.75	30	Adefin XL
Cap 5 mg			
NIMODIPINE			
Tab 30 mg			
Inj 200 mcg per ml, 50 ml vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Other Calcium Channel Blockers			
DILTIAZEM HYDROCHLORIDE			
Tab 30 mg	4.60	100	Dilzem
Tab 60 mg	8.50	100	Dilzem
Cap long-acting 120 mg	31.83	500	Apo-Diltiazem CD
Cap long-acting 180 mg	1.91	30	Cardizem CD
Cap long-acting 240 mg	47.67	500	Apo-Diltiazem CD
Cap long-acting 240 mg	7.56	30	Cardizem CD
Cap long-acting 240 mg	63.58	500	Apo-Diltiazem CD
Inj 5 mg per ml, 5 ml vial	10.22	30	Cardizem CD
PERHEXILINE MALEATE			
Tab 100 mg – 1% DV Jun-16 to 2019	62.90	100	Pexsig
VERAPAMIL HYDROCHLORIDE			
Tab 40 mg	7.01	100	Isoptin
Tab 80 mg – 1% DV Sep-14 to 2017	11.74	100	Isoptin
Tab long-acting 120 mg	15.20	250	Verpamil SR
Tab long-acting 240 mg	25.00	250	Verpamil SR
Inj 2.5 mg per ml, 2 ml ampoule	7.54	5	Isoptin
Centrally-Acting Agents			
CLONIDINE			
Patch 2.5 mg, 100 mcg per day – 1% DV Jul-14 to 2017	12.80	4	Catapres-TTS-1
Patch 5 mg, 200 mcg per day – 1% DV Jul-14 to 2017	18.04	4	Catapres-TTS-2
Patch 7.5 mg, 300 mcg per day – 1% DV Jul-14 to 2017	22.68	4	Catapres-TTS-3
CLONIDINE HYDROCHLORIDE			
Tab 25 mcg – 1% DV Sep-15 to 2018	10.53	112	Clonidine BNM
Tab 150 mcg	34.32	100	Catapres
Inj 150 mcg per ml, 1 ml ampoule	16.07	5	Catapres
METHYLDOPA			
Tab 125 mg	14.25	100	Prodopa
Tab 250 mg	15.10	100	Prodopa
Tab 500 mg	23.15	100	Prodopa
Diuretics			
Loop Diuretics			
BUMETANIDE			
Tab 1 mg	16.36	100	Burinex
Inj 500 mcg per ml, 4 ml vial			
FUROSEMIDE [FRUSEMIDE]			
Tab 40 mg – 1% DV Sep-15 to 2018	8.00	1,000	Diurin 40
Tab 500 mg – 1% DV Sep-15 to 2018	25.00	50	Urex Forte
Oral liq 10 mg per ml			
Inj 10 mg per ml, 2 ml ampoule – 1% DV Jun-16 to 2019	1.20	5	Frusemide-Claris
Inj 10 mg per ml, 25 ml ampoule			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Osmotic Diuretics			
MANNITOL			
Inj 10%, 1,000 ml bag	14.21	1,000 ml	Baxter
Inj 15%, 500 ml bag	9.84	500 ml	Baxter
Inj 20%, 500 ml bag	10.80	500 ml	Baxter
Potassium Sparing Combination Diuretics			
AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE			
Tab 5 mg with furosemide 40 mg			
AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE			
Tab 5 mg with hydrochlorothiazide 50 mg			
Potassium Sparing Diuretics			
AMILORIDE HYDROCHLORIDE			
Tab 5 mg	17.50	100	Apo-Amiloride
Oral liq 1 mg per ml	30.00	25 ml	Biomed
SPIRONOLACTONE			
Tab 25 mg – 1% DV Sep-13 to 2016	3.65	100	Spiractin
Tab 100 mg – 1% DV Sep-13 to 2016	11.80	100	Spiractin
Oral liq 5 mg per ml	30.00	25 ml	Biomed
Thiazide and Related Diuretics			
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]			
Tab 2.5 mg – 1% DV Sep-14 to 2017	5.48	500	Arrow-Bendrofluazide
Tab 5 mg – 1% DV Sep-14 to 2017	8.95	500	Arrow-Bendrofluazide
CHLOROTHIAZIDE			
Oral liq 50 mg per ml	26.00	25 ml	Biomed
CHLORTALIDONE [CHLORTHALIDONE]			
Tab 25 mg	8.00	50	Hygroton
INDAPAMIDE			
Tab 2.5 mg – 1% DV Oct-13 to 2016	2.25	90	Dapa-Tabs
METOLAZONE – Restricted see terms below			
⚡ Tab 5 mg			
➡ Restricted			
Initiation			
Either:			
1 Patient has refractory heart failure and is intolerant or has not responded to loop diuretics and/or loop-thiazide combination therapy; or			
2 Patient has severe refractory nephrotic oedema unresponsive to high dose loop diuretics and concentrated albumin infusions.			
Lipid-Modifying Agents			
Fibrates			
BEZAFIBRATE			
Tab 200 mg – 1% DV Oct-15 to 2018	9.05	90	Bezalip
Tab long-acting 400 mg – 1% DV Oct-15 to 2018	6.78	30	Bezalip Retard

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GEMFIBROZIL			
Tab 600 mg – 1% DV Nov-13 to 2016	17.60	60	Lipazil
HMG CoA Reductase Inhibitors (Statins)			
ATORVASTATIN			
Tab 10 mg	2.52	90	Zarator
Tab 20 mg	4.17	90	Zarator
Tab 40 mg	7.32	90	Zarator
Tab 80 mg	16.23	90	Zarator
PRAVASTATIN			
Tab 10 mg			
Tab 20 mg – 1% DV Oct-14 to 2017	3.45	30	Cholvastin
Tab 40 mg – 1% DV Oct-14 to 2017	6.36	30	Cholvastin
SIMVASTATIN			
Tab 10 mg – 1% DV Sep-14 to 2017	0.95	90	Arrow-Simva
Tab 20 mg – 1% DV Sep-14 to 2017	1.61	90	Arrow-Simva
Tab 40 mg – 1% DV Sep-14 to 2017	2.83	90	Arrow-Simva
Tab 80 mg – 1% DV Sep-14 to 2017	7.91	90	Arrow-Simva
Resins			
CHOLESTYRAMINE			
Powder for oral liq 4 g			
COLESTIPOL HYDROCHLORIDE			
Grans for oral liq 5 g			
Selective Cholesterol Absorption Inhibitors			
EZETIMIBE – Restricted see terms below			
⚡ Tab 10 mg – 1% DV Aug-15 to 2017	3.35	30	Ezemibe
➡ Restricted			
Initiation			
All of the following:			
1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and			
2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and			
3 Any of the following:			
3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 × normal) when treated with one statin; or			
3.2 The patient is intolerant to both simvastatin and atorvastatin; or			
3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.			
EZETIMIBE WITH SIMVASTATIN – Restricted see terms on the next page			
⚡ Tab 10 mg with simvastatin 10 mg – 1% DV Aug-15 to 2017	5.15	30	Zimybe
⚡ Tab 10 mg with simvastatin 20 mg – 1% DV Aug-15 to 2017	6.15	30	Zimybe
⚡ Tab 10 mg with simvastatin 40 mg – 1% DV Aug-15 to 2017	7.15	30	Zimybe
⚡ Tab 10 mg with simvastatin 80 mg – 1% DV Aug-15 to 2017	8.15	30	Zimybe

Price (ex man. excl. GST)	Brand or Generic Manufacturer
\$	Per

➔ **Restricted**

Initiation

All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Other Lipid-Modifying Agents

ACIPIMOX

Cap 250 mg

NICOTINIC ACID

Tab 50 mg – 1% DV Oct-14 to 2017	3.96	100	Apo-Nicotinic Acid
Tab 500 mg – 1% DV Oct-14 to 2017	17.37	100	Apo-Nicotinic Acid

Nitrates

GLYCERYL TRINITRATE

Tab 600 mcg	8.00	100	Lycinate
Inj 1 mg per ml, 5 ml ampoule	22.70	10	Nitronal
Inj 1 mg per ml, 50 ml vial	86.60	10	Nitronal
Inj 5 mg per ml, 10 ml ampoule	100.00	5	Hospira
Oral pump spray, 400 mcg per dose	4.45	250 dose	Nitrolingual Pump Spray
Oral spray, 400 mcg per dose	4.45	250 dose	Glytrin
Patch 25 mg, 5 mg per day – 1% DV Sep-14 to 2017	15.73	30	Nitroderm TTS 5
Patch 50 mg, 10 mg per day – 1% DV Sep-14 to 2017	18.62	30	Nitroderm TTS 10

ISOSORBIDE MONONITRATE

Tab 20 mg – 1% DV Sep-14 to 2017	17.10	100	Ismo-20
Tab long-acting 40 mg – 1% DV Jun-16 to 2019	7.50	30	Ismo 40 Retard
Tab long-acting 60 mg	3.94	90	Duride

Other Cardiac Agents

LEVOSIMENDAN – **Restricted** see terms below

- ⚡ Inj 2.5 mg per ml, 5 ml vial
- ⚡ Inj 2.5 mg per ml, 10 ml vial

➔ **Restricted**

Initiation — Heart transplant

Either:

- 1 For use as a bridge to heart transplant, in patients who have been accepted for transplant; or
- 2 For the treatment of heart failure following heart transplant.

Initiation — Heart failure

Cardiologist or intensivist

For the treatment of severe acute decompensated heart failure that is non-responsive to dobutamine.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Sympathomimetics			
ADRENALINE			
Inj 1 in 1,000, 1 ml ampoule	4.98	5	Aspen Adrenaline
	5.25		Hospira
Inj 1 in 1,000, 30 ml vial			
Inj 1 in 10,000, 10 ml ampoule	49.00	10	Aspen Adrenaline
	27.00	5	Hospira
Inj 1 in 10,000, 10 ml syringe			
DOBUTAMINE HYDROCHLORIDE			
Inj 12.5 mg per ml, 20 ml ampoule – 1% DV Jan-16 to 2018	24.45	5	Dobutamine-Claris
DOPAMINE HYDROCHLORIDE			
Inj 40 mg per ml, 5 ml ampoule – 1% DV Sep-15 to 2018	16.89	5	DBL Sterile Dopamine Concentrate
EPHEDRINE			
Inj 3 mg per ml, 10 ml syringe			
Inj 30 mg per ml, 1 ml ampoule – 1% DV Mar-15 to 2017	51.48	10	Max Health
ISOPRENALINE			
Inj 200 mcg per ml, 1 ml ampoule			
Inj 200 mcg per ml, 5 ml ampoule			
METARAMINOL			
Inj 0.5 mg per ml, 20 ml syringe			
Inj 1 mg per ml, 1 ml ampoule			
Inj 1 mg per ml, 10 ml syringe			
Inj 10 mg per ml, 1 ml ampoule			
NORADRENALINE			
Inj 0.06 mg per ml, 100 ml bag			
Inj 0.06 mg per ml, 50 ml syringe			
Inj 0.1 mg per ml, 100 ml bag			
Inj 0.12 mg per ml, 100 ml bag			
Inj 0.12 mg per ml, 50 ml syringe			
Inj 0.16 mg per ml, 50 ml syringe			
Inj 1 mg per ml, 100 ml bag			
Inj 1 mg per ml, 4 ml ampoule			
PHENYLEPHRINE HYDROCHLORIDE			
Inj 10 mg per ml, 1 ml vial	115.50	25	Neosynephrine HCL
Vasodilators			
ALPROSTADIL HYDROCHLORIDE			
Inj 500 mcg per ml, 1 ml ampoule – 1% DV Oct-15 to 2018	1,650.00	5	Prostin VR
AMYL NITRITE			
Liq 98% in 3 ml capsule			
DIAZOXIDE			
Inj 15 mg per ml, 20 ml ampoule			
HYDRALAZINE HYDROCHLORIDE			
⚡ Tab 25 mg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Initiation			
Either:			
1 For the treatment of refractory hypertension; or			
2 For the treatment of heart failure, in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.			
Inj 20 mg ampoule	25.90	5	Apresoline
MILRINONE			
Inj 1 mg per ml, 10 ml ampoule – 1% DV Jul-16 to 2018	300.30	10	Milrinone Generic Health
MINOXIDIL – Restricted see terms below			
⚡ Tab 10 mg	70.00	100	Loniten
➔Restricted			
Initiation			
For patients with severe refractory hypertension who have failed to respond to extensive multiple therapies.			
NICORANDIL			
Tab 10 mg	27.95	60	Ikorel
Tab 20 mg	33.28	60	Ikorel
PAPAVERINE HYDROCHLORIDE			
Inj 30 mg per ml, 1 ml vial			
Inj 12 mg per ml, 10 ml ampoule	217.90	5	Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]			
Tab 400 mg			
SODIUM NITROPRUSSIDE			
Inj 50 mg vial			

Endothelin Receptor Antagonists

AMBRISANTAN – Restricted see terms below			
⚡ Tab 5 mg	4,585.00	30	Volibris
⚡ Tab 10 mg	4,585.00	30	Volibris
➔Restricted			
Initiation			
Either:			
1 For use in patients with approval by the Pulmonary Arterial Hypertension Panel; or			
2 In hospital stabilisations in emergency situations.			
BOSENTAN – Restricted see terms below			
⚡ Tab 62.5 mg – 1% DV Jan-16 to 2018	375.00	56	Mylan-Bosentan
⚡ Tab 125 mg – 1% DV Jan-16 to 2018	375.00	56	Mylan-Bosentan
➔Restricted			
Initiation			
Either:			
1 For use in patients with approval by the Pulmonary Arterial Hypertension Panel; or			
2 In hospital stabilisation in emergency situations.			

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Restricted see terms on the next page			
⚡ Tab 25 mg – 1% DV Sep-15 to 2018	0.75	4	Vedafil
⚡ Tab 50 mg – 1% DV Sep-15 to 2018	0.75	4	Vedafil
⚡ Tab 100 mg – 1% DV Sep-15 to 2018	2.75	4	Vedafil

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔Restricted

Initiation

Any of the following:

- 1 For use in patients with approval by the Pulmonary Arterial Hypertension Panel; or
- 2 For use in neonatal units for persistent pulmonary hypertension of the newborn (PPHN); or
- 3 For use in weaning patients from inhaled nitric oxide; or
- 4 For perioperative use in cardiac surgery patients; or
- 5 For use in intensive care as an alternative to nitric oxide; or
- 6 In-hospital stabilisation in emergency situations; or
- 7 All of the following:
 - 7.1 Patient has Raynaud's phenomenon; and
 - 7.2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
 - 7.3 Patient is following lifestyle management (proper body insulation, avoidance of cold exposure, smoking cessation support, avoidance of sympathomimetic drugs); and
 - 7.4 Patient has persisting severe symptoms despite treatment with calcium channel blockers and nitrates (unless contraindicated or not tolerated).

Prostacyclin Analogues

EPOPROSTENOL – **Restricted** see terms below

⚡ Inj 0.5 mg vial	36.61	1	Veletri
⚡ Inj 1.5 mg vial	73.21	1	Veletri

➔Restricted

Initiation

For use as a bridge to transplant for patients with Pulmonary Arterial Hypertension who are on the active waiting list for lung transplantation.

ILOPROST

Inj 50 mcg in 0.5 ml ampoule – 1% DV Sep-15 to 2016	89.50	1	Arrow-Iloprost
⚡ Nebuliser soln 10 mcg per ml, 2 ml	1,185.00	30	Ventavis

➔Restricted

Initiation

Any of the following:

- 1 For use in patients with approval by the Pulmonary Arterial Hypertension Panel; or
- 2 For diagnostic use in catheter laboratories; or
- 3 For use following mitral or tricuspid valve surgery; or
- 4 In hospital stabilisation in emergency situations.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Anti-Infective Preparations			
Antibacterials			
FUSIDIC ACID			
Crm 2% – 1% DV Jan-15 to 2016	2.52	15 g	DP Fusidic Acid Cream
Oint 2% – 1% DV Sep-13 to 2016	3.45	15 g	Foban
HYDROGEN PEROXIDE			
Crm 1%	8.56	15 g	Crystaderm
Soln 3% (10 vol) – 1% DV Nov-15 to 2018	1.40	100 ml	Pharmacy Health
MAFENIDE ACETATE – Restricted see terms below			
↓ Powder 50 g sachet			
➔ Restricted			
Initiation			
For the treatment of burns patients.			
MUPIROCIN			
Oint 2%			
SULPHADIAZINE SILVER			
Crm 1%	12.30	50 g	Flamazine
Antifungals			
AMOROLFINE			
Nail soln 5% – 1% DV Jan-15 to 2017	19.95	5 ml	MycoNail
CICLOPIROX OLAMINE			
Nail soln 8% – 1% DV Sep-15 to 2018	6.50	7 ml	Apo-Ciclopirox
➔ Soln 1% – Restricted: For continuation only			
CLOTRIMAZOLE			
Crm 1% – 1% DV Sep-14 to 2017	0.52	20 g	Clomazol
➔ Soln 1% – Restricted: For continuation only			
ECONAZOLE NITRATE			
➔ Crm 1% – Restricted: For continuation only			
Foaming soln 1%			
KETOCONAZOLE			
Shampoo 2% – 1% DV Dec-14 to 2017	2.99	100 ml	Sebizole
METRONIDAZOLE			
Gel 0.75%			
MICONAZOLE NITRATE			
Crm 2% – 1% DV Mar-15 to 2017	0.55	15 g	Multichem
➔ Lotn 2% – Restricted: For continuation only			
Tinc 2%			
NYSTATIN			
Crm 100,000 u per g			
Antiparasitics			
MALATHION [MALDISON]			
Lotn 0.5%			
Shampoo 1%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MALATHION WITH PERMETHRIN AND PIPERONYL BUTOXIDE			
Spray 0.25% with permethrin 0.5% and piperonyl butoxide 2%			
PERMETHRIN			
Crm 5% – 1% DV Apr-15 to 2017	4.20	30 g	Lyderm
Lotn 5% – 1% DV Sep-14 to 2017	3.19	30 ml	A-Scabies

Antiacne Preparations

ADAPALENE			
Crm 0.1%			
Gel 0.1%			
BENZOYL PEROXIDE			
Soln 5%			
ISOTRETINOIN			
Cap 10 mg	12.47	100	Isotane 10
	14.96	120	Oratane
Cap 20 mg	19.27	100	Isotane 20
	23.12	120	Oratane
TRETINOIN			
Crm 0.05%			

Antipruritic Preparations

CALAMINE			
Crm, aqueous, BP – 1% DV Dec-15 to 2018	1.49	100 g	Pharmacy Health
Lotn, BP – 1% DV Dec-15 to 2018	12.94	2,000 ml	PSM
CROTAMITON			
Crm 10% – 1% DV Sep-15 to 2018	3.37	20 g	Itch-Soothe

Barrier Creams and Emollients

Barrier Creams

DIMETHICONE			
Crm 5% tube – 1% DV Apr-14 to 2016	1.65	100 g	healthE Dimethicone 5%
Crm 5% pump bottle – 1% DV Apr-14 to 2016	4.73	500 ml	healthE Dimethicone 5%
Crm 10% pump bottle – 1% DV Nov-15 to 2018	4.90	500 ml	healthE Dimethicone 10%
ZINC			
Crm			<i>e.g. Zinc Cream (Orion); Zinc Cream (PSM)</i>
Oint			<i>e.g. Zinc oxide (PSM)</i>
Paste			
ZINC AND CASTOR OIL			
Crm	1.63	20 g	Orion
Oint, BP – 1% DV Jul-15 to 2017	1.39	20 g	healthE

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ZINC WITH WOOL FAT			
Crm zinc 15.25% with wool fat 4%			<i>e.g. Sudocrem</i>
Emollients			
AQUEOUS CREAM			
Crm 100 g – 1% DV Jan-16 to 2018	1.00	100 g	Pharmacy Health SLS-free
Note: DV limit applies to the pack sizes of 100 g or less.			
Crm 500 g – 1% DV Mar-16 to 2018	1.99	500 g	AFT SLS-free
Note: DV limit applies to the pack sizes of greater than 100 g.			
CETOMACROGOL			
Crm BP, 500 g – 1% DV Nov-15 to 2018	2.74	500 g	healthE
Crm BP, 100 g – 1% DV Jan-16 to 2018	1.47	1	healthE
CETOMACROGOL WITH GLYCEROL			
Crm 90% with glycerol 10%,	2.00	100 g	Pharmacy Health
	2.10		Pharmacy Health
	3.20		healthE
Crm 90% with glycerol 10% – 1% DV Aug-16 to 2019	2.82	500 ml	Pharmacy Health Sorbolene with Glycerin
	3.87	1,000 ml	Pharmacy Health Sorbolene with Glycerin
Crm 90% with glycerol 10%, 500 ml, 1 bottle	5.46	1	healthE
<i>(healthE Crm 90% with glycerol 10%, 500 ml, 1 bottle to be delisted 1 August 2016)</i>			
EMULSIFYING OINTMENT			
Oint BP – 1% DV Apr-15 to 2017	1.84	100 g	Jaychem
Note: DV limit applies to pack sizes of less than 200 g.			
Oint BP, 500 g – 1% DV Jul-15 to 2017	2.73	500 g	AFT
Note: DV limit applies to pack sizes of greater than 200 g.			
GLYCEROL WITH PARAFFIN			
Crm glycerol 10% with white soft paraffin 5% and liquid paraffin 10%			<i>e.g. QV cream</i>
OIL IN WATER EMULSION			
Crm	2.63	500 g	healthE Fatty Cream
Crm, 100 g	1.60	1	healthE Fatty Cream
PARAFFIN			
Oint liquid paraffin 50% with white soft paraffin 50%	3.10	100 g	healthE
White soft – 1% DV Sep-15 to 2018	0.85	10 g	healthE
Note: DV limit applies to pack sizes of 30 g or less, and to both white soft paraffin and yellow soft paraffin			
Yellow soft			
PARAFFIN WITH WOOL FAT			
Lotn liquid paraffin 15.9% with wool fat 0.6%			<i>e.g. AlphaKeri;BK ;DP; Hydroderm Lotn</i>
Lotn liquid paraffin 91.7% with wool fat 3%			<i>e.g. Alpha Keri Bath Oil</i>
UREA			
Crm 10%			
WOOL FAT			
Crm			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Corticosteroids			
BETAMETHASONE DIPROPIONATE			
Crm 0.05%			
Oint 0.05%			
BETAMETHASONE VALERATE			
Crm 0.1% – 1% DV Jun-15 to 2018	3.15	50 g	Beta Cream
Oint 0.1% – 1% DV Jun-15 to 2018	3.15	50 g	Beta Ointment
Lotn 0.1%			
CLOBETASOL PROPIONATE			
Crm 0.05% – 1% DV Jul-15 to 2016	3.20	30 g	Clobetasol BNM
Oint 0.05% – 1% DV Jul-15 to 2016	3.20	30 g	Clobetasol BNM
CLOBETASONE BUTYRATE			
Crm 0.05%			
DIFLUCORTOLONE VALERATE – Restricted: For continuation only			
➡ Crm 0.1%			
➡ Fatty oint 0.1%			
HYDROCORTISONE			
Crm 1%, 100 g	3.75	100 g	Pharmacy Health
Crm 1%, 500 g	14.00	500 g	Pharmacy Health
HYDROCORTISONE ACETATE			
Crm 1%	2.48	14.2 g	AFT
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN			
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – 1% DV Dec-14 to 2017	10.57	250 ml	DP Lotn HC
HYDROCORTISONE BUTYRATE			
Crm 0.1%	2.30	30 g	Locoid Lipocream
	6.85	100 g	Locoid Lipocream
Oint 0.1%	6.85	100 g	Locoid
Milky emul 0.1%	6.85	100 ml	Locoid Crelo
HYDROCORTISONE WITH PARAFFIN AND WOOL FAT			
Lotn 1% with paraffin liquid 15.9% and wool fat 0.6%			
METHYLPREDNISOLONE ACEPONATE			
Crm 0.1%	4.95	15 g	Advantan
Oint 0.1%	4.95	15 g	Advantan
MOMETASONE FUROATE			
Crm 0.1% – 1% DV Nov-15 to 2018	1.51	15 g	Elocon Alcohol Free
	2.90	50 g	Elocon Alcohol Free
Oint 0.1% – 1% DV Nov-15 to 2018	1.51	15 g	Elocon
	2.90	50 g	Elocon
Lotn 0.1% – 1% DV Sep-15 to 2018	7.35	30 ml	Elocon
TRIAMCINOLONE ACETONIDE			
Crm 0.02% – 1% DV Apr-15 to 2017	6.30	100 g	Aristocort
Oint 0.02% – 1% DV Apr-15 to 2017	6.35	100 g	Aristocort

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Corticosteroids with Anti-Infective Agents

BETAMETHASONE VALERATE WITH CLIOQUINOL – **Restricted** see terms below

¶ Crm 0.1% with clioquinol 3%

➔Restricted

Initiation

Either:

- 1 For the treatment of intertrigo; or
- 2 For continuation use.

BETAMETHASONE VALERATE WITH FUSIDIC ACID

Crm 0.1% with fusidic acid 2%

HYDROCORTISONE WITH MICONAZOLE

Crm 1% with miconazole nitrate 2% – **1% DV Sep-15 to 2018**2.00 15 g **Micreme H**

HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN

Crm 1% with natamycin 1% and neomycin sulphate 0.5%2.79 15 g Pimafucort

Oint 1% with natamycin 1% and neomycin sulphate 0.5%2.79 15 g Pimafucort

TRIAMCINOLONE ACETONIDE WITH NEOMYCIN SULPHATE, GRAMICIDIN AND NYSTATIN

Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and
gramicidin 250 mcg per g

Psoriasis and Eczema Preparations

ACITRETIN

Cap 10 mg – **1% DV Nov-14 to 2017**17.86 60 **Novatretin**

Cap 25 mg – **1% DV Nov-14 to 2017**41.36 60 **Novatretin**

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

Gel 500 mcg with calcipotriol 50 mcg per g – **1% DV Sep-15 to 2018**26.12 30 g **Daivobet**

Oint 500 mcg with calcipotriol 50 mcg per g – **1% DV Sep-15 to 2018**26.12 30 g **Daivobet**

CALCIPOTRIOL

Crm 50 mcg per g45.00 100 g Daivonex

Oint 50 mcg per g45.00 100 g Daivonex

Soln 50 mcg per ml16.00 30 ml Daivonex

COAL TAR WITH SALICYLIC ACID AND SULPHUR

Oint 12% with salicylic acid 2% and sulphur 4%

METHOXSALEN [8-METHOXYPORALEN]

Tab 10 mg

Lotn 1.2%

PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN

Soln 2.3% with trolamine laurilsulfate and fluorescein sodium3.36 500 ml Pinetarsol

5.82 1,000 ml Pinetarsol

POTASSIUM PERMANGANATE

Tab 400 mg

Crystals

Scalp Preparations

BETAMETHASONE VALERATE

Scalp app 0.1%7.75 100 ml Beta Scalp

DERMATOLOGICALS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
CLOBETASOL PROPIONATE			
Scalp app 0.05%	6.96	30 ml	Dermol
HYDROCORTISONE BUTYRATE			
Scalp lotn 0.1%	3.65	100 ml	Locoid

Wart Preparations

IMIQUIMOD			
Crm 5%, 250 mg sachet – 1% DV Feb-15 to 2017	17.98	12	Apo-Imiquimod Cream 5%
PODOPHYLLOTOXIN			
Soln 0.5%	33.60	3.5 ml	Condyline
SILVER NITRATE			
Sticks with applicator			

Other Skin Preparations

DIPHEMANIL METILSULFATE			
Powder 2%			
SUNSCREEN, PROPRIETARY			
Crm			
Lotn	3.30	100 g	Marine Blue Lotion SPF 50+
	5.10	200 g	Marine Blue Lotion SPF 50+

Antineoplastics

FLUOROURACIL SODIUM			
Crm 5% – 1% DV Sep-15 to 2018	8.95	20 g	Efudix
METHYL AMINOLEVULINATE HYDROCHLORIDE – Restricted see terms below			
☞ Crm 16%			
☞ Restricted			
Dermatologist or plastic surgeon			

Wound Management Products

CALCIUM GLUCONATE			
Gel 2.5%	21.00	1	healthE

Price (ex man. excl. GST)	Brand or Generic
\$	Per Manufacturer

Anti-Infective Agents

ACETIC ACID

- Soln 3%
- Soln 5%

ACETIC ACID WITH HYDROXYQUINOLINE, GLYCEROL AND RICINOLEIC ACID

- Jelly 0.94% with hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator

CHLORHEXIDINE GLUCONATE

- | | | | |
|---|------|------|----------------|
| Crm 1% – 1% DV Sep-15 to 2018 | 1.21 | 50 g | healthE |
| Lotn 1%, 200 ml – 1% DV Sep-15 to 2018 | 2.98 | 1 | healthE |

CLOTRIMAZOLE

- | | | | |
|--|------|------|-----------------|
| Vaginal crm 1% with applicator – 1% DV Dec-13 to 2016 | 1.45 | 35 g | Clomazol |
| Vaginal crm 2% with applicator – 1% DV Dec-13 to 2016 | 2.20 | 20 g | Clomazol |

MICONAZOLE NITRATE

- | | | | |
|--|------|------|----------------|
| Vaginal crm 2% with applicator – 1% DV Oct-14 to 2017 | 3.95 | 40 g | Micreme |
|--|------|------|----------------|

NYSTATIN

- Vaginal crm 100,000 u per 5 g with applicator(s)

Contraceptives

Antiandrogen Oral Contraceptives

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

- | | | | |
|--|------|-----|--------------|
| Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tablets – 1% DV Dec-14 to 2017 | 5.36 | 168 | Ginet |
|--|------|-----|--------------|

Combined Oral Contraceptives

ETHINYLOESTRADIOL WITH DESOGESTREL

- Tab 20 mcg with desogestrel 150 mcg
- Tab 30 mcg with desogestrel 150 mcg

ETHINYLOESTRADIOL WITH LEVONORGESTREL

- | | | | |
|--|------|----|------------------|
| Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets | 2.65 | 84 | Ava 20 ED |
| Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets | 2.30 | 84 | Ava 30 ED |
| Tab 20 mcg with levonorgestrel 100 mcg | | | |
| Tab 30 mcg with levonorgestrel 150 mcg | | | |
| Tab 50 mcg with levonorgestrel 125 mcg | 9.45 | 84 | Microgynon 50 ED |

ETHINYLOESTRADIOL WITH NORETHISTERONE

- Tab 35 mcg with norethisterone 1 mg
- Tab 35 mcg with norethisterone 500 mcg

NORETHISTERONE WITH MESTRANOL

- Tab 1 mg with mestranol 50 mcg

Contraceptive Devices

INTRA-UTERINE DEVICE

- | | | | |
|--|-------|---|-----------------------|
| IUD 29.1 mm length × 23.2 mm width | 31.60 | 1 | Choice TT380 Short |
| IUD 33.6 mm length × 29.9 mm width | 31.60 | 1 | Choice TT380 Standard |
| IUD 35.5 mm length × 19.6 mm width | 31.60 | 1 | Choice Load 375 |

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Emergency Contraception			
LEVONORGESTREL			
Tab 1.5 mg – 1% DV Jul-13 to 2016	3.50	1	Postinor-1
Progestogen-Only Contraceptives			
LEVONORGESTREL			
Tab 30 mcg			
Subdermal implant (2 × 75 mg rods) – 5% DV Oct-14 to 31 Dec 2017	133.65	1	Jadelle
⚡ Intra-uterine system, 20 mcg per day			<i>e.g. Mirena</i>
➡ Restricted			
Initiation — heavy menstrual bleeding			
Obstetrician or gynaecologist			
All of the following:			
1 The patient has a clinical diagnosis of heavy menstrual bleeding; and			
2 The patient has failed to respond to or is unable to tolerate other appropriate pharmaceutical therapies as per the Heavy Menstrual Bleeding Guidelines; and			
3 Any of the following:			
3.1 Serum ferritin level < 16 mcg/l (within the last 12 months); or			
3.2 Haemoglobin level < 120 g/l; or			
3.3 The patient has had a uterine ultrasound and either a hysteroscopy or endometrial biopsy.			
Continuation — heavy menstrual bleeding			
Obstetrician or gynaecologist			
Either:			
1 Patient demonstrated clinical improvement of heavy menstrual bleeding; or			
2 Previous insertion was removed or expelled within 3 months of insertion.			
Initiation — endometriosis			
Obstetrician or gynaecologist			
The patient has a clinical diagnosis of endometriosis confirmed by laparoscopy.			
Continuation — endometriosis			
Obstetrician or gynaecologist			
Either:			
1 Patient demonstrated satisfactory management of endometriosis; or			
2 Previous insertion was removed or expelled within 3 months of insertion.			
Note: endometriosis is an unregistered indication.			
MEDROXYPROGESTERONE ACETATE			
Inj 150 mg per ml, 1 ml syringe – 1% DV Sep-13 to 2016	7.00	1	Depo-Provera
NORETHISTERONE			
Tab 350 mcg – 1% DV Oct-15 to 2018	6.25	84	Noriday 28
Obstetric Preparations			
Antiprogestogens			
MIFEPRISTONE			
Tab 200 mg			
Oxytocics			
CARBOPROST TROMETAMOL			
Inj 250 mcg per ml, 1 ml ampoule			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DINOPROSTONE			
Pessaries 10 mg			
Vaginal gel 1 mg in 3 g	52.65	1	Prostin E2
Vaginal gel 2 mg in 3 g	64.60	1	Prostin E2
ERGOMETRINE MALEATE			
Inj 500 mcg per ml, 1 ml ampoule – 1% DV Oct-14 to 2017	94.70	5	DBL Ergometrine
OXYTOCIN			
Inj 5 iu per ml, 1 ml ampoule – 1% DV Nov-15 to 2018	4.03	5	Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule – 1% DV Nov-15 to 2018	5.03	5	Oxytocin BNM
OXYTOCIN WITH ERGOMETRINE MALEATE			
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampoule – 1% DV Sep-15 to 2018	11.13	5	Syntometrine

Tocolytics

PROGESTERONE – Restricted see terms below

⚡ Cap 100 mg – **1% DV Aug-16 to 2019** 16.50 30 **Utrogestan**

➡ **Restricted**

Initiation

Gynaecologist or obstetrician

Both:

- 1 For the prevention of pre-term labour*; and
- 2 Either:
 - 2.1 The patient has a short cervix on ultrasound (defined as < 25mm at 16 to 28 weeks); or
 - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are Unapproved Indications (refer to Section A: General Rules, Part I (Interpretations and Definitions) and Part IV (Miscellaneous Provisions) rule 23.1)

TERBUTALINE – Restricted see terms below

⚡ Inj 500 mcg ampoule

➡ **Restricted**

Obstetrician

Oestrogens

OESTRIOL

Crm 1 mg per g with applicator

Pessaries 500 mcg

Urologicals

5-Alpha Reductase Inhibitors

FINASTERIDE – Restricted see terms below

⚡ Tab 5 mg – **1% DV Dec-14 to 2017** 2.08 30 **Finpro**

➡ **Restricted**

Initiation

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Alpha-1A Adrenoceptor Blockers			
TAMSULOSIN – Restricted see terms below			
⚡ Cap 400 mcg – 1% DV Dec-13 to 2016	13.51	100	Tamsulosin-Rex
➡ Restricted			
Initiation			
Both:			
1 Patient has symptomatic benign prostatic hyperplasia; and			
2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.			
Urinary Alkalisers			
POTASSIUM CITRATE – Restricted see terms below			
⚡ Oral liq 3 mmol per ml	30.00	200 ml	Biomed
➡ Restricted			
Initiation			
Both:			
1 The patient has recurrent calcium oxalate urolithiasis; and			
2 The patient has had more than two renal calculi in the two years prior to the application.			
SODIUM CITRO-TARTRATE			
Grans eff 4 g sachets – 1% DV Feb-15 to 2017	2.93	28	Ural
Urinary Antispasmodics			
OXYBUTYNIN			
Tab 5 mg – 1% DV Jun-13 to 2016	11.20	500	Apo-Oxybutynin
Oral liq 5 mg per 5 ml – 1% DV Jun-13 to 2016	56.45	473 ml	Apo-Oxybutynin
SOLIFENACIN SUCCINATE – Restricted see terms below			
⚡ Tab 5 mg	37.50	30	Vesicare
⚡ Tab 10 mg	37.50	30	Vesicare
➡ Restricted			
Initiation			
Patient has overactive bladder and a documented intolerance of, or is non-responsive to, oxybutynin.			
TOLTERODINE TARTRATE – Restricted see terms below			
⚡ Tab 1 mg	14.56	56	Arrow-Tolterodine
⚡ Tab 2 mg	14.56	56	Arrow-Tolterodine
➡ Restricted			
Initiation			
Patient has overactive bladder and a documented intolerance of, or is non-responsive to, oxybutynin.			

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Anabolic Agents

OXANDROLONE

⚡ Tab 2.5 mg

➡ **Restricted**

Initiation

For the treatment of burns patients.

Androgen Agonists and Antagonists

CYPROTERONE ACETATE

Tab 50 mg – 1% DV Oct-15 to 2018 15.87 50 **Procur**

Tab 100 mg – 1% DV Oct-15 to 2018 30.40 50 **Procur**

TESTOSTERONE

Patch 2.5 mg per day 80.00 60 Androderm

TESTOSTERONE CYPIONATE

Inj 100 mg per ml, 10 ml vial – 1% DV Sep-14 to 2017 76.50 1 **Depo-Testosterone**

TESTOSTERONE ESTERS

Inj testosterone decanoate 100 mg, testosterone isocaproate 60 mg,
testosterone phenylpropionate 60 mg and testosterone propionate
30 mg per ml, 1 ml ampoule

TESTOSTERONE UNDECANOATE

Cap 40 mg – 1% DV Sep-15 to 2018 16.80 60 **Andriol Testocaps**

Inj 250 mg per ml, 4 ml vial 86.00 1 **Reandron 1000**

Calcium Homeostasis

CALCITONIN

Inj 100 iu per ml, 1 ml ampoule – 1% DV Oct-14 to 2017 121.00 5 **Miacalcic**

CINACALCET – **Restricted** see terms below

⚡ Tab 30 mg 403.70 28 **Sensipar**

➡ **Restricted**

Initiation

Nephrologist or endocrinologist

Re-assessment required after 6 months

Either:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium \geq 3 mmol/L) despite previous first-line treatments including bisphosphonates and sodium thiosulfate; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciophylaxis (calcific uraemic arteriopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium \geq 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Continuation

Nephrologist or endocrinologist

Both:

continued...

HORMONE PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
1 The patient's serum calcium level has fallen to < 3mmol/L; and			
2 The patient has experienced clinically significant symptom improvement.			
Note: This does not include parathyroid adenomas unless these have become malignant.			
ZOLEDRONIC ACID			
☞ Inj 4 mg per 5 ml, vial	550.00	1	Zometa
☞ Restricted			
Initiation			
Oncologist, haematologist or palliative care specialist			
Any of the following:			
1 Patient has hypercalcaemia of malignancy; or			
2 Both:			
2.1 Patient has bone metastases or involvement; and			
2.2 Patient has severe bone pain resistant to standard first-line treatments; or			
3 Both:			
3.1 Patient has bone metastases or involvement; and			
3.2 Patient is at risk of skeletal-related events (pathological fracture, spinal cord compression, radiation to bone or surgery to bone).			
Corticosteroids			
BETAMETHASONE			
Tab 500 mcg			
Inj 4 mg per ml, 1 ml ampoule			
BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE			
Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml ampoule			
DEXAMETHASONE			
Tab 0.5 mg – 1% DV Jan-16 to 2018	0.88	30	Dexamethsone
Tab 4 mg – 1% DV Jan-16 to 2018	1.84	30	Dexamethsone
Oral liq 1 mg per ml	45.00	25 ml	Biomed
DEXAMETHASONE PHOSPHATE			
Inj 4 mg per ml, 1 ml ampoule – 1% DV Apr-14 to 2016	14.19	10	Max Health
Max Health brand - HSS with 1% DV will apply 1 July 2016 to 30 June 2019.			
Inj 4 mg per ml, 2 ml ampoule – 1% DV Apr-14 to 2016	12.59	5	Max Health
Max Health brand - HSS with 1% DV will apply 1 July 2016 to 30 June 2019.			
FLUDROCORTISONE ACETATE			
Tab 100 mcg	14.32	100	Florinef
HYDROCORTISONE			
Tab 5 mg – 1% DV Sep-15 to 2018	8.10	100	Douglas
Tab 20 mg – 1% DV Sep-15 to 2018	20.32	100	Douglas
Inj 100 mg vial – 1% DV Oct-13 to 2016	4.99	1	Solu-Cortef
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)			
Tab 4 mg – 1% DV Oct-15 to 2018	80.00	100	Medrol
Tab 100 mg – 1% DV Oct-15 to 2018	180.00	20	Medrol
Inj 40 mg vial – 1% DV Oct-15 to 2018	10.50	1	Solu-Medrol
Inj 125 mg vial – 1% DV Oct-15 to 2018	22.25	1	Solu-Medrol
Inj 500 mg vial – 1% DV Oct-15 to 2018	9.00	1	Solu-Medrol
Inj 1 g vial – 1% DV Oct-15 to 2018	16.00	1	Solu-Medrol

☞ Item restricted (see ☞ above); ☞ Item restricted (see ☞ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
METHYLPREDNISOLONE ACETATE			
Inj 40 mg per ml, 1 ml vial – 1% DV Oct-15 to 2018	40.00	5	Depo-Medrol
METHYLPREDNISOLONE ACETATE WITH LIDOCAINE [LIGNOCAINE]			
Inj 40 mg with lidocaine [lignocaine], 1 ml vial – 1% DV Oct-15 to 2018	9.25	1	Depo-Medrol with Lidocaine
PREDNISOLONE			
Oral liq 5 mg per ml	7.50	30 ml	Redipred
Enema 200 mcg per ml, 100 ml			
PREDNISONE			
Tab 1 mg	10.68	500	Apo-Prednisone
	2.13	100	Apo-Prednisone S29
Tab 2.5 mg	12.09	500	Apo-Prednisone
Tab 5 mg	11.09	500	Apo-Prednisone
Tab 20 mg	29.03	500	Apo-Prednisone
TRIAMCINOLONE ACETONIDE			
Inj 10 mg per ml, 1 ml ampoule – 1% DV Apr-15 to 2017	20.80	5	Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule – 1% DV Apr-15 to 2017	51.70	5	Kenacort-A 40
TRIAMCINOLONE HEXACETONIDE			
Inj 20 mg per ml, 1 ml vial			

Hormone Replacement Therapy

Oestrogens

OESTRADIOL

- Tab 1 mg
- Tab 2 mg
- Patch 25 mcg per day
- Patch 50 mcg per day
- Patch 100 mcg per day

OESTRADIOL VALERATE

- Tab 1 mg – **1% DV Jun-15 to 2018** 12.36 84 **Progynova**
- Tab 2 mg – **1% DV Jun-15 to 2018** 12.36 84 **Progynova**

OESTROGENS (CONJUGATED EQUINE)

- Tab 300 mcg
- Tab 625 mcg

Progestogen and Oestrogen Combined Preparations

OESTRADIOL WITH NORETHISTERONE ACETATE

- Tab 1 mg with 0.5 mg norethisterone acetate
- Tab 2 mg with 1 mg norethisterone acetate
- Tab 2 mg with 1 mg norethisterone acetate (10), and tab 2 mg oestradiol (12) and tab 1 mg oestradiol (6)

OESTROGENS WITH MEDROXYPROGESTERONE ACETATE

- Tab 625 mcg conjugated equine with 2.5 mg medroxyprogesterone acetate
- Tab 625 mcg conjugated equine with 5 mg medroxyprogesterone acetate

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Progestogens			
MEDROXYPROGESTERONE ACETATE			
Tab 2.5 mg – 1% DV Sep-13 to 2016	3.09	30	Provera
Tab 5 mg – 1% DV Sep-13 to 2016	13.06	100	Provera
Tab 10 mg – 1% DV Sep-13 to 2016	6.85	30	Provera

Other Endocrine Agents

CABERGOLINE – **Restricted** see terms below

⚡ Tab 0.5 mg – 1% DV Sep-15 to 2018	4.75	2	Dostinex
	19.00	8	Dostinex

➡ **Restricted**

Initiation

Any of the following:

- 1 Inhibition of lactation; or
- 2 Patient has pathological hyperprolactinemia; or
- 3 Patient has acromegaly.

CLOMIPHENE CITRATE

Tab 50 mg – 1% DV Sep-13 to 2016	29.84	10	Serophene
--	-------	----	------------------

DANAZOL

Cap 100 mg	68.33	100	Azol
Cap 200 mg	97.83	100	Azol

GESTRINONE

Cap 2.5 mg

METYRAPONE

Cap 250 mg

PENTAGASTRIN

Inj 250 mcg per ml, 2 ml ampoule

Other Oestrogen Preparations

ETHINYLOESTRADIOL

Tab 10 mcg – 1% DV Sep-15 to 2018	17.60	100	NZ Medical & Scientific
---	-------	-----	------------------------------------

OESTRADIOL

Implant 50 mg

OESTRIOL

Tab 2 mg

Other Progestogen Preparations

MEDROXYPROGESTERONE

Tab 100 mg – 1% DV Sep-13 to 2016	96.50	100	Provera
---	-------	-----	----------------

NORETHISTERONE

Tab 5 mg – 1% DV Jun-15 to 2018	18.29	100	Primolut N
---------------------------------------	-------	-----	-------------------

Pituitary and Hypothalamic Hormones and Analogues

CORTICOTRORELIN (OVINE)

Inj 100 mcg vial

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
THYROTROPIN ALFA			
Inj 900 mcg vial			
Adrenocorticotrophic Hormones			
TETRACOSACTIDE [TETRACOSACTRIN]			
Inj 250 mcg per ml, 1 ml ampoule	17.71	1	Synacthen
Inj 1 mg per ml, 1 ml ampoule	29.56	1	Synacthen Depot
GnRH Agonists and Antagonists			
BUSERELIN			
Inj 1 mg per ml, 5.5 ml vial			
GONADORELIN			
Inj 100 mcg vial			
GOSERELIN			
Implant 3.6 mg	166.20	1	Zoladex
Implant 10.8 mg	443.76	1	Zoladex
LEUPRORELIN ACETATE			
Inj 3.75 mg syringe	221.60	1	Lucrin Depot PDS
Inj 7.5 mg syringe	166.20	1	Eligard
Inj 11.25 mg syringe	591.68	1	Lucrin Depot PDS
Inj 22.5 mg syringe	443.76	1	Eligard
Inj 30 mg syringe	1,109.40	1	Lucrin Depot PDS
Inj 30 mg vial	591.68	1	Eligard
Inj 45 mg syringe	832.05	1	Eligard
Gonadotrophins			
CHORIOGONADOTROPIN ALFA			
Inj 250 mcg in 0.5 ml syringe			
Growth Hormone			
SOMATROPIN – Restricted see terms below			
⚡ Inj 5 mg cartridge – 1% DV Jan-15 to 31 Dec 2017	109.50	1	Omnitrope
⚡ Inj 10 mg cartridge – 1% DV Jan-15 to 31 Dec 2017	219.00	1	Omnitrope
⚡ Inj 15 mg cartridge – 1% DV Jan-15 to 31 Dec 2017	328.50	1	Omnitrope
➡Restricted			
Initiation — growth hormone deficiency in children			
Endocrinologist or paediatric endocrinologist			
<i>Re-assessment required after 12 months</i>			
Either:			
1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or			
2 All of the following:			
2.1 Height velocity < 25th percentile for age; and adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and			
2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
- 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
- 2.5 Appropriate imaging of the pituitary gland has been obtained.

Continuation — growth hormone deficiency in children

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 A current bone age is ≤ 14 years (female patients) or ≤ 16 years (male patients); and
- 2 Height velocity is ≥ 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is ≥ 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initiation — Turner syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Continuation — Turner syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 Height velocity ≥ 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is ≥ 2 cm per year, calculated over six months; and
- 3 A current bone age is ≤ 14 years; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initiation — short stature without growth hormone deficiency

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Continuation — short stature without growth hormone deficiency

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 Height velocity is \geq 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is \geq 2 cm per year as calculated over six months; and
- 3 Current bone age is \leq 14 years (female patients) or \leq 16 years (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initiation — short stature due to chronic renal insufficiency

Endocrinologist, paediatric endocrinologist or renal physician on the recommendation of a endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is $<$ 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is \leq to 14 years (female patients) or \leq to 16 years (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a $GFR \leq 30 \text{ ml/min/1.73 m}^2$ as measured by the Schwartz method ($\text{Height(cm)/plasma creatinine (umol/l)} \times 40 = \text{corrected GFR (ml/min/1.73 m}^2)$) in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received $< 5\text{mg/ m}^2$ /day of prednisone or equivalent for at least 6 months.

Continuation — short stature due to chronic renal insufficiency

Endocrinologist, paediatric endocrinologist or renal physician on the recommendation of a endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 Height velocity is \geq 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is \geq 2 cm per year as calculated over six months; and
- 3 A current bone age is \leq 14 years (female patients) or \leq 16 years (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initiation — Prader-Willi syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2 The patient's height velocity is < 25th percentile for bone age adjusted for bone age/pubertal status if appropriate as calculated over 6 to 12 months using the standards of Tanner and Davies (1985) or pubertal status over 6 to 12 months; and
- 3 Either:
 - 3.1 The patient is under two years of age and height velocity has been assessed over a minimum six month period from the age of 12 months, with at least three supine length measurements over this period demonstrating clear and consistent evidence of linear growth failure (with height velocity < 25th percentile); or
 - 3.2 The patient is aged two years or older; and
- 4 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 5 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 6 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by ≥ 0.5 standard deviations in the preceding 12 months.

Continuation — Prader-Willi syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 Height velocity is ≥ 50 th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is ≥ 2 cm per year as calculated over six months; and
- 3 A current bone age is ≤ 14 years (female patients) or ≤ 16 years (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by ≥ 0.5 standard deviations in the preceding 12 months.

Initiation — adults and adolescents

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA[®]).

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued. . .

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of ≤ 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of ≤ 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until it is within 1 standard deviation of the mean normal value for age and sex; and

The dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Continuation — adults and adolescents

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

Either:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in the Quality of Life Assessment defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA[®]) score from baseline; and
 - 1.3 Serum IGF-I levels have increased to within ± 1 SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin does not exceed 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA[®] score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ± 1 SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients.

Thyroid and Antithyroid Preparations

CARBIMAZOLE

Tab 5 mg

IODINE

Soln BP 50 mg per ml

LEVOTHYROXINE

Tab 25 mcg

Tab 50 mcg

Tab 100 mcg

LIOTHYRONINE SODIUM

↓ Tab 20 mcg

→ **Restricted**

Initiation

For a maximum of 14 days' treatment in patients with thyroid cancer who are due to receive radioiodine therapy.

Inj 20 mcg vial

POTASSIUM IODATE

Tab 170 mg

POTASSIUM PERCHLORATE

Cap 200 mg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PROPYLTHIOURACIL – Restricted see terms below			
⚡ Tab 50 mg	35.00	100	PTU

➡**Restricted**

Initiation

Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Note: Propylthiouracil is not recommended for patients under the age of 18 years unless the patient is pregnant and other treatments are contraindicated.

PROTIRELIN

Inj 100 mcg per ml, 2 ml ampoule

Vasopressin Agents

ARGIPRESSIN [VASOPRESSIN]

Inj 20 u per ml, 1 ml ampoule

DESMOPRESSIN ACETATE – **Some items restricted** see terms below

⚡ Tab 100 mcg – 1% DV Jun-16 to 2019	25.00	30	Minirin
⚡ Tab 200 mcg – 1% DV Jun-16 to 2019	54.45	30	Minirin
Nasal spray 10 mcg per dose – 1% DV Sep-14 to 2017	22.95	6 ml	Desmopressin-PH&T
Inj 4 mcg per ml, 1 ml ampoule			
Inj 15 mcg per ml, 1 ml ampoule			
Nasal drops 100 mcg per ml			

➡**Restricted**

Initiation — Nocturnal enuresis

Either:

- 1 The nasal forms of desmopressin are contraindicated; or
- 2 An enuresis alarm is contraindicated.

Note: Cranial diabetes insipidus and the nasal forms of desmopressin are contraindicated.

TERLIPRESSIN

Inj 0.1 mg per ml, 8.5 ml ampoule	450.00	5	Glypressin
Inj 1 mg per 8.5 ml ampoule – 1% DV Jun-15 to 2018	215.00	5	Glypressin

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Antibacterials			
Aminoglycosides			
AMIKACIN – Restricted see terms below			
☞ Inj 5 mg per ml, 10 ml syringe			
☞ Inj 5 mg per ml, 5 ml syringe	176.00	10	Biomed
☞ Inj 15 mg per ml, 5 ml syringe			
☞ Inj 250 mg per ml, 2 ml vial – 1% DV Oct-14 to 2017	431.20	5	DBL Amikacin
☞ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
GENTAMICIN SULPHATE			
Inj 10 mg per ml, 1 ml ampoule	8.56	5	Hospira
Inj 10 mg per ml, 2 ml ampoule	175.10	25	APP Pharmaceuticals
Inj 40 mg per ml, 2 ml ampoule – 1% DV Sep-15 to 2018	6.00	10	Pfizer
	30.00	50	Pfizer
PAROMOMYCIN – Restricted see terms below			
☞ Cap 250 mg	126.00	16	Humatin
☞ Restricted			
Clinical microbiologist or infectious disease specialist			
STREPTOMYCIN SULPHATE – Restricted see terms below			
☞ Inj 400 mg per ml, 2.5 ml ampoule			
☞ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
TOBRAMYCIN			
☞ Powder			
☞ Restricted			
Initiation			
For addition to orthopaedic bone cement.			
☞ Inj 40 mg per ml, 2 ml vial	38.00	5	DBL Tobramycin
☞ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
☞ Inj 100 mg per ml, 5 ml vial			
☞ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
☞ Solution for inhalation 60 mg per ml, 5 ml	2,200.00	56 dose	TOBI
☞ Restricted			
Initiation			
Patient has cystic fibrosis.			
Carbapenems			
ERTAPENEM – Restricted see terms below			
☞ Inj 1 g vial	73.50	1	Invanz
☞ Restricted			
Clinical microbiologist or infectious disease specialist			
IMIPENEM WITH CILASTATIN – Restricted see terms below			
☞ Inj 500 mg with 500 mg cilastatin vial – 1% DV Jun-15 to 2017	13.79	1	Imipenem+Cilastatin RBX
☞ Restricted			
Clinical microbiologist or infectious disease specialist			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MEROPENEM – Restricted see terms below			
‡ Inj 500 mg vial – 1% DV Oct-14 to 2017	35.22	10	DBL Meropenem
‡ Inj 1 g vial – 1% DV Oct-14 to 2017	65.21	10	DBL Meropenem

➡Restricted

Clinical microbiologist or infectious disease specialist

Cephalosporins and Cephamycins - 1st Generation
CEFALEXIN

Cap 500 mg – 1% DV Oct-13 to 2016	5.70	20	Cephalexin ABM
Grans for oral liq 25 mg per ml – 1% DV Sep-15 to 2018	8.00	100 ml	Cefalexin Sandoz
Grans for oral liq 50 mg per ml – 1% DV Sep-15 to 2018	11.00	100 ml	Cefalexin Sandoz

CEFAZOLIN

Inj 500 mg vial – 1% DV Sep-14 to 2017	3.99	5	AFT
Inj 1 g vial – 1% DV Sep-14 to 2017	3.38	5	AFT

Cephalosporins and Cephamycins - 2nd Generation
CEFACLOR

Cap 250 mg – 1% DV Dec-13 to 2016	26.00	100	Ranbaxy-Cefaclor
Grans for oral liq 25 mg per ml – 1% DV Dec-13 to 2016	3.53	100 ml	Ranbaxy-Cefaclor

CEFOXITIN

Inj 1 g vial – 1% DV Jan-16 to 2018	58.00	10	Cefoxitin Actavis
---	-------	----	--------------------------

CEFUROXIME

Tab 250 mg	29.40	50	Zinnat
Inj 750 mg vial	3.70	5	Zinacef
Inj 1.5 g vial	1.30	1	Zinacef

Cephalosporins and Cephamycins - 3rd Generation
CEFOTAXIME

Inj 500 mg vial	1.90	1	Cefotaxime Sandoz
Inj 1 g vial – 1% DV Oct-14 to 2017	17.10	10	DBL Cefotaxime

CEFTAZIDIME – Restricted see terms below

‡ Inj 500 mg vial – 1% DV Jan-15 to 2017	5.30	1	Fortum
‡ Inj 1 g vial – 1% DV Jan-15 to 2017	1.55	1	Fortum
‡ Inj 2 g vial – 1% DV Jan-15 to 2017	3.34	1	Fortum

➡Restricted

Clinical microbiologist, infectious disease specialist or respiratory specialist

CEFTRIAXONE

Inj 500 mg vial – 1% DV Mar-14 to 2016	1.50	1	Ceftriaxone-AFT
Inj 1 g vial – 1% DV Mar-14 to 2016	5.22	5	Ceftriaxone-AFT
Inj 2 g vial – 1% DV Mar-14 to 2016	2.75	1	Ceftriaxone-AFT

Cephalosporins and Cephamycins - 4th Generation
CEFEPIME – Restricted see terms below

‡ Inj 1 g vial – 1% DV Oct-15 to 2018	3.95	1	Cefepime-AFT
‡ Inj 2 g vial – 1% DV Oct-15 to 2018	6.92	1	Cefepime-AFT

➡Restricted

Clinical microbiologist or infectious disease specialist

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Cephalosporins and Cephamycins - 5th Generation			
CEFTAROLINE FOSAMIL – Restricted see terms below			
⚡ Inj 600 mg vial	1,450.00	10	Zinforo
➔ Restricted			
Initiation — multi-resistant organism salvage therapy			
Clinical microbiologist or infectious disease specialist			
Either:			
1 for patients where alternative therapies have failed; or			
2 for patients who have a contraindication or hypersensitivity to standard current therapies.			
Macrolides			
AZITHROMYCIN – Restricted see terms below			
⚡ Tab 250 mg – 1% DV Sep-15 to 2018	9.00	30	Apo-Azithromycin
⚡ Tab 500 mg – 1% DV Sep-15 to 2018	1.05	2	Apo-Azithromycin
⚡ Grans for oral liq 200 mg per 5 ml (40 mg per ml) – 1% DV Oct-15 to 2018	12.50	15 ml	Zithromax
➔ Restricted			
Initiation			
Any of the following:			
1 Patient has received a lung transplant and requires treatment or prophylaxis for bronchiolitis obliterans syndrome; or			
2 Patient has cystic fibrosis and has chronic infection with <i>Pseudomonas aeruginosa</i> or <i>Pseudomonas</i> related gram negative organisms; or			
3 For any other condition for five days' treatment, with review after five days.			
CLARITHROMYCIN – Restricted see terms below			
⚡ Tab 250 mg – 1% DV Sep-14 to 2017	3.98	14	Apo-Clarithromycin
⚡ Tab 500 mg – 1% DV Sep-14 to 2017	10.40	14	Apo-Clarithromycin
⚡ Grans for oral liq 50 mg per ml	23.12	50 ml	Klacid
⚡ Grans for oral liq 25 mg per ml	23.12	70 ml	Klacid
⚡ Inj 500 mg vial – 1% DV Mar-15 to 2017	20.40	1	Martindale
<i>(Klacid Grans for oral liq 25 mg per ml to be delisted 1 October 2016)</i>			
➔ Restricted			
Initiation — Tab 250 mg and oral liquid			
Either:			
1 Atypical mycobacterial infection; or			
2 <i>Mycobacterium tuberculosis</i> infection where there is drug resistance or intolerance to standard pharmaceutical agents.			
Initiation — Tab 500 mg			
<i>Helicobacter pylori</i> eradication.			
Initiation — Infusion			
Any of the following:			
1 Atypical mycobacterial infection; or			
2 <i>Mycobacterium tuberculosis</i> infection where there is drug resistance or intolerance to standard pharmaceutical agents; or			
3 Community-acquired pneumonia.			
ERYTHROMYCIN (AS ETHYLSUCCINATE)			
Tab 400 mg	16.95	100	E-Mycin
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	E-Mycin
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	E-Mycin
ERYTHROMYCIN (AS LACTOBIONATE)			
Inj 1 g vial	16.00	1	Erythrocin IV

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ERYTHROMYCIN (AS STEARATE) – Restricted: For continuation only			
➔ Tab 250 mg			
➔ Tab 500 mg			
ROXITHROMYCIN			
Tab 150 mg	7.48	50	Arrow-Roxithromycin
Tab 300 mg	14.40	50	Arrow-Roxithromycin
Penicillins			
AMOXICILLIN			
Cap 250 mg – 1% DV Mar-14 to 2016	16.18	500	Apo-Amoxi
Cap 500 mg – 1% DV Jul-14 to 2016	20.94	500	Apo-Amoxi
Grans for oral liq 125 mg per 5 ml	0.88	100 ml	Amoxicillin Actavis
.....	2.00		Ospamox
Grans for oral liq 250 mg per 5 ml	0.97	100 ml	Amoxicillin Actavis
.....	2.00		Ospamox
Inj 250 mg vial – 1% DV Oct-14 to 2017	10.67	10	Ibiamox
Inj 500 mg vial – 1% DV Oct-14 to 2017	12.41	10	Ibiamox
Inj 1 g vial – 1% DV Oct-14 to 2017	17.29	10	Ibiamox
AMOXICILLIN WITH CLAVULANIC ACID			
Tab 500 mg with clavulanic acid 125 mg – 1% DV Aug-16 to 2017	1.95	20	Augmentin
.....	9.75	100	Curam Duo
Grans for oral liq 25 mg with clavulanic acid 6.25 mg per ml	3.83	100 ml	Augmentin
Grans for oral liq 50 mg with clavulanic acid 12.5 mg per ml	4.97	100 ml	Augmentin
Inj 500 mg with clavulanic acid 100 mg vial – 1% DV Sep-15 to 2018	10.14	10	m-Amoxiclav
Inj 1,000 mg with clavulanic acid 200 mg vial – 1% DV Sep-15 to 2018	12.80	10	m-Amoxiclav
<i>(Curam Duo Tab 500 mg with clavulanic acid 125 mg to be delisted 1 August 2016)</i>			
BENZATHINE BENZYL PENICILLIN			
Inj 900 mg (1.2 million units) in 2.3 ml syringe – 1% DV Sep-15 to 2018	315.00	10	Bicillin LA
BENZYL PENICILLIN SODIUM [PENICILLIN G]			
Inj 600 mg (1 million units) vial – 1% DV Sep-14 to 2017	10.35	10	Sandoz
FLUCLOXACILLIN			
Cap 250 mg – 1% DV Sep-15 to 2018	18.70	250	Staphlex
Cap 500 mg – 1% DV Sep-15 to 2018	62.90	500	Staphlex
Grans for oral liq 25 mg per ml – 1% DV Sep-15 to 2018	2.29	100 ml	AFT
Grans for oral liq 50 mg per ml – 1% DV Sep-15 to 2018	3.08	100 ml	AFT
Inj 250 mg vial – 1% DV Sep-14 to 2017	8.80	10	Flucloxin
Inj 500 mg vial – 1% DV Sep-14 to 2017	9.20	10	Flucloxin
Inj 1 g vial – 1% DV Jan-16 to 2017	11.60	10	Flucloxin
PHENOXYMETHYL PENICILLIN [PENICILLIN V]			
Cap 250 mg – 1% DV Jun-15 to 2018	2.88	50	Cilicaine VK
Cap 500 mg – 1% DV Jun-15 to 2018	4.73	50	Cilicaine VK
Grans for oral liq 125 mg per 5 ml – 1% DV Apr-14 to 2016	1.64	100 ml	AFT
Grans for oral liq 250 mg per 5 ml – 1% DV Apr-14 to 2016	1.74	100 ml	AFT
PIPERACILLIN WITH TAZOBACTAM – Restricted see terms below			
⚡ Inj 4 g with tazobactam 0.5 g vial	5.84	1	Hospira
➔ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PROCAINE PENICILLIN			
Inj 1.5 g in 3.4 ml syringe – 1% DV Sep-14 to 2017	123.50	5	Cilicaïne
TICARCILLIN WITH CLAVULANIC ACID – Restricted see terms below			
⚡ Inj 3 g with clavulanic acid 0.1 mg vial			
➔ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
Quinolones			
CIPROFLOXACIN – Restricted see terms below			
⚡ Tab 250 mg – 1% DV Sep-14 to 2017	1.75	28	Cipflox
⚡ Tab 500 mg – 1% DV Sep-14 to 2017	2.00	28	Cipflox
⚡ Tab 750 mg – 1% DV Sep-14 to 2017	3.75	28	Cipflox
⚡ Oral liq 50 mg per ml			
⚡ Oral liq 100 mg per ml			
⚡ Inj 2 mg per ml, 100 ml bag – 1% DV Mar-16 to 2018	30.58	10	Cipflox
➔ Restricted			
Clinical microbiologist or infectious disease specialist			
MOXIFLOXACIN – Restricted see terms below			
⚡ Tab 400 mg	52.00	5	Avelox
⚡ Inj 1.6 mg per ml, 250 ml bottle	70.00	1	Avelox IV 400
➔ Restricted			
Initiation — Mycobacterium infection			
Infectious disease specialist, clinical microbiologist or respiratory specialist			
Either:			
1 Both:			
1.1 Active tuberculosis; and			
1.2 Any of the following:			
1.2.1 Documented resistance to one or more first-line medications; or			
1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or			
1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or			
1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or			
1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications;			
or			
2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.			
Initiation — Pneumonia			
Infectious disease specialist or clinical microbiologist			
Either:			
1 Immunocompromised patient with pneumonia that is unresponsive to first-line treatment; or			
2 Pneumococcal pneumonia or other invasive pneumococcal disease highly resistant to other antibiotics.			
Initiation — Penetrating eye injury			
Ophthalmologist			
Five days treatment for patients requiring prophylaxis following a penetrating eye injury.			
Initiation — Mycoplasma genitalium			
All of the following:			
1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium; and			
2 Has tried and failed to clear infection using azithromycin; and			
3 Treatment is only for 7 days.			
NORFLOXACIN			
Tab 400 mg – 1% DV Sep-14 to 2017	13.50	100	Arrow-Norfloxacin

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Tetracyclines			
DEMECLOCYCLINE HYDROCHLORIDE			
Tab 150 mg			
Cap 150 mg			
Cap 300 mg			
DOXYCYCLINE			
➔ Tab 50 mg – Restricted: For continuation only			
Tab 100 mg – 1% DV Sep-14 to 2017	6.75	250	Doxine
Inj 5 mg per ml, 20 ml vial			
MINOCYCLINE			
Tab 50 mg			
➔ Cap 100 mg – Restricted: For continuation only			
TETRACYCLINE			
Tab 250 mg			
Cap 500 mg	46.00	30	Tetracyclin Wolff
TIGECYCLINE – Restricted see terms below			
⚡ Inj 50 mg vial			
➔ Restricted			
Clinical microbiologist or infectious disease specialist			
Other Antibacterials			
AZTREONAM – Restricted see terms below			
⚡ Inj 1 g vial	131.00	5	Azactam
➔ Restricted			
Clinical microbiologist or infectious disease specialist			
CHLORAMPHENICOL – Restricted see terms below			
⚡ Inj 1 g vial			
➔ Restricted			
Clinical microbiologist or infectious disease specialist			
CLINDAMYCIN – Restricted see terms below			
⚡ Cap 150 mg – 1% DV Oct-13 to 2016	5.80	16	Clindamycin ABM
⚡ Oral liq 15 mg per ml			
⚡ Inj 150 mg per ml, 4 ml ampoule – 1% DV Sep-13 to 2016	100.00	10	Dalacin C
➔ Restricted			
Clinical microbiologist or infectious disease specialist			
COLISTIN SULPHOMETHATE [COLESTIMETHATE] – Restricted see terms below			
⚡ Inj 150 mg per ml, 1 ml vial	65.00	1	Colistin-Link
➔ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
DAPTOMYCIN – Restricted see terms below			
⚡ Inj 350 mg vial – 1% DV Sep-15 to 2018	175.16	1	Cubicin
⚡ Inj 500 mg vial – 1% DV Sep-15 to 2018	243.52	1	Cubicin
➔ Restricted			
Clinical microbiologist or infectious disease specialist			
FOSFOMYCIN – Restricted see terms on the next page			
⚡ Powder for oral solution, 3 g sachet			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Clinical microbiologist or infectious disease specialist			
FUSIDIC ACID – Restricted see terms below			
⌘ Tab 250 mg	34.50	12	Fucidin
➔Restricted			
Clinical microbiologist or infectious disease specialist			
HEXAMINE HIPPURATE			
Tab 1 g			
LINCOMYCIN – Restricted see terms below			
⌘ Inj 300 mg per ml, 2 ml vial			
➔Restricted			
Clinical microbiologist or infectious disease specialist			
LINEZOLID – Restricted see terms below			
⌘ Tab 600 mg – 1% DV Sep-15 to 2018	800.00	10	Zyvox
⌘ Oral liq 20 mg per ml – 1% DV Sep-15 to 2018	775.00	150 ml	Zyvox
⌘ Inj 2 mg per ml, 300 ml bag – 1% DV Sep-15 to 2018	1,650.00	10	Zyvox
➔Restricted			
Clinical microbiologist or infectious disease specialist			
NITROFURANTOIN			
Tab 50 mg			
Tab 100 mg			
PIVMECILLINAM – Restricted see terms below			
⌘ Tab 200 mg			
➔Restricted			
Clinical microbiologist or infectious disease specialist			
SULPHADIAZINE – Restricted see terms below			
⌘ Tab 500 mg			
➔Restricted			
Clinical microbiologist, infectious disease specialist or maternal-foetal medicine specialist			
TEICOPLANIN – Restricted see terms below			
⌘ Inj 400 mg vial			
➔Restricted			
Clinical microbiologist or infectious disease specialist			
TRIMETHOPRIM			
Tab 100 mg			
Tab 300 mg – 1% DV Oct-15 to 2018	15.00	50	TMP
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXAZOLE]			
Tab 80 mg with sulphamethoxazole 400 mg			
Oral liq 8 mg with sulphamethoxazole 40 mg per ml	2.15	100 ml	Deprim
Inj 16 mg with sulphamethoxazole 80 mg per ml, 5 ml ampoule			
VANCOMYCIN – Restricted see terms below			
⌘ Inj 500 mg vial – 1% DV Oct-14 to 2017	2.64	1	Mylan
➔Restricted			
Clinical microbiologist or infectious disease specialist			

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Antifungals

Imidazoles

KETOCONAZOLE

☞ Tab 200 mg

☞ **Restricted**

Oncologist

Polyene Antimycotics

AMPHOTERICIN B

☞ Inj (liposomal) 50 mg vial – 1% DV Sep-15 to 20183,450.00 10 **AmBisome**

☞ **Restricted**

Initiation

Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist
Either:

- 1 Proven or probable invasive fungal infection, to be prescribed under an established protocol; or
- 2 Both:
 - 2.1 Possible invasive fungal infection; and
 - 2.2 A multidisciplinary team (including an infectious disease physician or a clinical microbiologist) considers the treatment to be appropriate.

☞ Inj 50 mg vial

☞ **Restricted**

Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist

NYSTATIN

Tab 500,000 u	17.09	50	Nilstat
Cap 500,000 u	15.47	50	Nilstat

Triazoles

FLUCONAZOLE – **Restricted** see terms below

☞ Cap 50 mg – 1% DV Nov-14 to 2017.....	3.49	28	Ozole
☞ Cap 150 mg – 1% DV Nov-14 to 2017	0.71	1	Ozole
☞ Cap 200 mg – 1% DV Nov-14 to 2017.....	9.69	28	Ozole
☞ Oral liquid 50 mg per 5 ml	98.50	35 ml	Diflucan
☞ Inj 2 mg per ml, 50 ml vial – 1% DV Oct-13 to 2016.....	4.95	1	Fluconazole-Claris
☞ Inj 2 mg per ml, 100 ml vial – 1% DV Oct-13 to 2016	6.47	1	Fluconazole-Claris

☞ **Restricted**

Consultant

ITRACONAZOLE – **Restricted** see terms below

☞ Cap 100 mg – 1% DV Oct-13 to 2016	2.99	15	Itrazole
☞ Oral liquid 10 mg per ml			

☞ **Restricted**

Clinical immunologist, clinical microbiologist, dermatologist or infectious disease specialist

POSACONAZOLE – **Restricted** see terms on the next page

☞ Oral liq 40 mg per ml	761.13	105 ml	Noxafil
-------------------------------	--------	--------	---------

Price (ex man. excl. GST)	Brand or Generic
\$	Per Manufacturer

➔Restricted

Initiation

Haematologist or infectious disease specialist

Re-assessment required after 6 weeks

Both:

- 1 Either:
 - 1.1 Patient has acute myeloid leukaemia; or
 - 1.2 Patient is planned to receive a stem cell transplant and is at high risk for aspergillus infection; and
- 2 Patient is to be treated with high dose remission induction therapy or re-induction therapy.

Continuation

Haematologist or infectious disease specialist

Re-assessment required after 6 weeks

Both:

- 1 Patient has previously received posaconazole prophylaxis during remission induction therapy; and
- 2 Any of the following:
 - 2.1 Patient is to be treated with high dose remission re-induction therapy; or
 - 2.2 Patient is to be treated with high dose consolidation therapy; or
 - 2.3 Patient is receiving a high risk stem cell transplant.

VORICONAZOLE – **Restricted** see terms below

⚡ Tab 50 mg – 1% DV Jan-16 to 2018	130.00	56	Vttack
⚡ Tab 200 mg – 1% DV Jan-16 to 2018	500.00	56	Vttack
⚡ Oral liq 40 mg per ml	730.00	70 ml	Vfend
⚡ Inj 200 mg vial	185.00	1	Vfend

➔Restricted

Initiation — Proven or probable aspergillus infection

Clinical microbiologist, haematologist or infectious disease specialist

Both:

- 1 Patient is immunocompromised; and
- 2 Patient has proven or probable invasive aspergillus infection.

Initiation — Possible aspergillus infection

Clinical microbiologist, haematologist or infectious disease specialist

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient has possible invasive aspergillus infection; and
- 3 A multidisciplinary team (including an infectious disease physician) considers the treatment to be appropriate.

Initiation — Resistant candidiasis infections and other moulds

Clinical microbiologist, haematologist or infectious disease specialist

All of the following:

- 1 Patient is immunocompromised; and
- 2 Either:
 - 2.1 Patient has fluconazole resistant candidiasis; or
 - 2.2 Patient has mould strain such as *Fusarium* spp. and *Scedosporium* spp; and
- 3 A multidisciplinary team (including an infectious disease physician or clinical microbiologist) considers the treatment to be appropriate.

Other Antifungals

CASPOFUNGIN – **Restricted** see terms on the next page

⚡ Inj 50 mg vial	667.50	1	Cancidas
⚡ Inj 70 mg vial	862.50	1	Cancidas

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➡Restricted			
Initiation			
Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist			
Either:			
1	Proven or probable invasive fungal infection, to be prescribed under an established protocol; or		
2	Both:		
2.1	Possible invasive fungal infection; and		
2.2	A multidisciplinary team (including an infectious disease physician or a clinical microbiologist) considers the treatment to be appropriate.		
FLUCYTOSINE – Restricted see terms below			
⚡	Cap 500 mg		
➡Restricted			
Clinical microbiologist or infectious disease specialist			
TERBINAFINE			
Tab 250 mg – 1% DV Sep-14 to 2017	1.50	14	Dr Reddy's Terbinafine
Antimycobacterials			
Antileprotics			
CLOFAZIMINE – Restricted see terms below			
⚡	Cap 50 mg		
➡Restricted			
Clinical microbiologist, dermatologist or infectious disease specialist			
DAPSONE – Restricted see terms below			
⚡ Tab 25 mg – 1% DV Sep-14 to 2017	95.00	100	Dapsone
⚡ Tab 100 mg – 1% DV Sep-14 to 2017	110.00	100	Dapsone
➡Restricted			
Clinical microbiologist, dermatologist or infectious disease specialist			
Antituberculotics			
CYCLOSERINE – Restricted see terms below			
⚡	Cap 250 mg		
➡Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
ETHAMBUTOL HYDROCHLORIDE – Restricted see terms below			
⚡ Tab 100 mg	48.01	56	Myambutol
⚡ Tab 400 mg	49.34	56	Myambutol
➡Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
ISONIAZID – Restricted see terms below			
⚡ Tab 100 mg – 1% DV Sep-15 to 2018	20.00	100	PSM
➡Restricted			
Clinical microbiologist, dermatologist, paediatrician, public health physician or internal medicine physician			
ISONIAZID WITH RIFAMPICIN – Restricted see terms below			
⚡ Tab 100 mg with rifampicin 150 mg – 1% DV Sep-15 to 2018	85.54	100	Rifinah
⚡ Tab 150 mg with rifampicin 300 mg – 1% DV Sep-15 to 2018	170.60	100	Rifinah
➡Restricted			
Clinical microbiologist, dermatologist, paediatrician, public health physician or internal medicine physician			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PARA-AMINOSALICYLIC ACID – Restricted see terms below			
¶ Grans for oral liq 4 g	280.00	30	Paser
➔ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
PROTIONAMIDE – Restricted see terms below			
¶ Tab 250 mg	305.00	100	Peteha
➔ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
PYRAZINAMIDE – Restricted see terms below			
¶ Tab 500 mg			
➔ Restricted			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
RIFABUTIN – Restricted see terms below			
¶ Cap 150 mg – 1% DV Sep-13 to 2016.....	213.19	30	Mycobutin
➔ Restricted			
Clinical microbiologist, gastroenterologist, infectious disease specialist or respiratory specialist			
RIFAMPICIN – Restricted see terms below			
¶ Cap 150 mg – 1% DV Nov-14 to 2017.....	55.75	100	Rifadin
¶ Cap 300 mg – 1% DV Nov-14 to 2017.....	116.25	100	Rifadin
¶ Oral liq 100 mg per 5 ml – 1% DV Nov-14 to 2017.....	12.00	60 ml	Rifadin
¶ Inj 600 mg vial – 1% DV Nov-14 to 2017.....	128.85	1	Rifadin
➔ Restricted			
Clinical microbiologist, dermatologist, internal medicine physician, paediatrician or public health physician			

Antiparasitics

Anthelmintics

ALBENDAZOLE – Restricted see terms below			
¶ Tab 200 mg			
¶ Tab 400 mg			
➔ Restricted			
Clinical microbiologist or infectious disease specialist			
IVERMECTIN – Restricted see terms below			
¶ Tab 3 mg	17.20	4	Stromectol
➔ Restricted			
Clinical microbiologist, dermatologist or infectious disease specialist			
MEBENDAZOLE			
Tab 100 mg	24.19	24	De-Worm
Oral liq 100 mg per 5 ml			
PRAZQUANTEL			
Tab 600 mg			

Antiprotozoals

ARTEMETHER WITH LUMEFANTRINE – Restricted see terms below			
¶ Tab 20 mg with lumefantrine 120 mg			
➔ Restricted			
Clinical microbiologist or infectious disease specialist			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ARTESUNATE – Restricted see terms below			
☞ Inj 60 mg vial			
☞ Restricted			
Clinical microbiologist or infectious disease specialist			
ATOVAQUONE WITH PROGUANIL HYDROCHLORIDE – Restricted see terms below			
☞ Tab 62.5 mg with proguanil hydrochloride 25 mg – 1% DV Nov-14 to 2017	25.00	12	Malarone Junior
☞ Tab 250 mg with proguanil hydrochloride 100 mg – 1% DV Nov-14 to 2017	64.00	12	Malarone
☞ Restricted			
Clinical microbiologist or infectious disease specialist			
CHLOROQUINE PHOSPHATE – Restricted see terms below			
☞ Tab 250 mg			
☞ Restricted			
Clinical microbiologist, dermatologist, infectious disease specialist or rheumatologist			
MEFLOQUINE – Restricted see terms below			
☞ Tab 250 mg – 1% DV Dec-14 to 2017	33.48	8	Lariam
☞ Restricted			
Clinical microbiologist, dermatologist, infectious disease specialist or rheumatologist			
METRONIDAZOLE			
Tab 200 mg	10.45	100	Trichozole
Tab 400 mg	18.15	100	Trichozole
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	Flagyl-S
Inj 5 mg per ml, 100 ml bag – 1% DV Apr-15 to 2017	6.94	5	AFT
Suppos 500 mg	24.48	10	Flagyl
NITAZOXANIDE – Restricted see terms below			
☞ Tab 500 mg	1,680.00	30	Alinia
☞ Oral liq 100 mg per 5 ml			
☞ Restricted			
Clinical microbiologist or infectious disease specialist			
ORNIDAZOLE			
Tab 500 mg	16.50	10	Arrow-Ornidazole
PENTAMIDINE ISETHIONATE – Restricted see terms below			
☞ Inj 300 mg vial – 1% DV Mar-15 to 2017	180.00	5	Pentacarinat
☞ Restricted			
Clinical microbiologist or infectious disease specialist			
PRIMAQUINE PHOSPHATE – Restricted see terms below			
☞ Tab 7.5 mg			
☞ Restricted			
Clinical microbiologist or infectious disease specialist			
PYRIMETHAMINE – Restricted see terms below			
☞ Tab 25 mg			
☞ Restricted			
Clinical microbiologist, infectious disease specialist or maternal-foetal medicine specialist			
QUININE DIHYDROCHLORIDE – Restricted see terms on the next page			
☞ Inj 60 mg per ml, 10 ml ampoule			
☞ Inj 300 mg per ml, 2 ml vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Clinical microbiologist or infectious disease specialist			
QUININE SULPHATE			
Tab 300 mg	54.06	500	Q 300
SODIUM STIBOGLUCONATE – Restricted see terms below			
☞ Inj 100 mg per ml, 1 ml vial			
➔Restricted			
Clinical microbiologist or infectious disease specialist			
SPIRAMYCIN – Restricted see terms below			
☞ Tab 500 mg			
➔Restricted			
Maternal-foetal medicine specialist			

Antiretrovirals

HIV Fusion Inhibitors

ENFUVIRTIDE – **Restricted** see terms below

☞ Inj 108 mg vial × 60 2,380.00 1 Fuzeon

➔Restricted

Initiation

Re-assessment required after 12 months

All of the following:

- 1 Confirmed HIV infection; and
- 2 Enfuvirtide to be given in combination with optimized background therapy (including at least 1 other antiretroviral drug that the patient has never previously been exposed to) for treatment failure; and
- 3 Either:
 - 3.1 Patient has evidence of HIV replication, despite ongoing therapy; or
 - 3.2 Patient has treatment-limiting toxicity to previous antiretroviral agents; and
- 4 Previous treatment with 3 different antiretroviral regimens has failed; and
- 5 All of the following:
 - 5.1 Previous treatment with a non-nucleoside reverse transcriptase inhibitor has failed; and
 - 5.2 Previous treatment with a nucleoside reverse transcriptase inhibitor has failed; and
 - 5.3 Previous treatment with a protease inhibitor has failed.

Continuation

Patient has had at least a 10-fold reduction in viral load at 12 months.

Non-Nucleoside Reverse Transcriptase Inhibitors

➔Restricted

Initiation — Confirmed HIV

Both:

- 1 Confirmed HIV infection; and
- 2 Any of the following:
 - 2.1 Symptomatic patient; or
 - 2.2 Patient aged 12 months and under; or
 - 2.3 Both:
 - 2.3.1 Patient aged 1 to 5 years; and
 - 2.3.2 Any of the following:
 - 2.3.2.1 CD4 counts < 1000 cells/mm³; or
 - 2.3.2.2 CD4 counts < 0.25 × total lymphocyte count; or

continued...

continued...

2.3.2.3 Viral load counts > 100000 copies per ml; or

2.4 Both:

2.4.1 Patient aged 6 years and over; and

2.4.2 CD4 counts < 500 cells/mm³.

Initiation — Prevention of maternal transmission

Either:

1 Prevention of maternal foetal transmission; or

2 Treatment of the newborn for up to eight weeks.

Initiation — Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

1 Treatment course to be initiated within 72 hours post exposure; and

2 Any of the following:

2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or

2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or

2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initiation — Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

EFAVIRENZ — **Restricted** see terms on the preceding page

⬆ Tab 50 mg – 1% DV Sep-15 to 2018	63.38	30	Stocrin
--	-------	----	----------------

⬆ Tab 200 mg – 1% DV Sep-15 to 2018	190.15	90	Stocrin
---	--------	----	----------------

⬆ Tab 600 mg – 1% DV Sep-15 to 2018	63.38	30	Stocrin
---	-------	----	----------------

⬆ Oral liq 30 mg per ml			
-------------------------	--	--	--

ETRAVIRINE — **Restricted** see terms on the preceding page

⬆ Tab 200 mg	770.00	60	Intelence
--------------------	--------	----	-----------

NEVIRAPINE — **Restricted** see terms on the preceding page

⬆ Tab 200 mg – 1% DV Nov-15 to 2018	65.00	60	Nevirapine Alphapharm
---	-------	----	------------------------------

⬆ Oral suspension 10 mg per ml	134.55	240 ml	Viramune Suspension
--------------------------------------	--------	--------	---------------------

Nucleoside Reverse Transcriptase Inhibitors

➡Restricted

Initiation — Confirmed HIV

Both:

1 Confirmed HIV infection; and

2 Any of the following:

2.1 Symptomatic patient; or

2.2 Patient aged 12 months and under; or

2.3 Both:

2.3.1 Patient aged 1 to 5 years; and

2.3.2 Any of the following:

2.3.2.1 CD4 counts < 1000 cells/mm³; or

2.3.2.2 CD4 counts < 0.25 × total lymphocyte count; or

2.3.2.3 Viral load counts > 100000 copies per ml; or

2.4 Both:

2.4.1 Patient aged 6 years and over; and

2.4.2 CD4 counts < 500 cells/mm³.

Initiation — Prevention of maternal transmission

Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued. . .			
1 Prevention of maternal foetal transmission; or			
2 Treatment of the newborn for up to eight weeks.			
Initiation — Post-exposure prophylaxis following non-occupational exposure to HIV			
Both:			
1 Treatment course to be initiated within 72 hours post exposure; and			
2 Any of the following:			
2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or			
2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or			
2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.			
Initiation — Percutaneous exposure			
Patient has percutaneous exposure to blood known to be HIV positive.			
ABACAVIR SULPHATE – Restricted see terms on the preceding page			
⬆ Tab 300 mg – 1% DV Oct-14 to 2017	229.00	60	Ziagen
⬆ Oral liq 20 mg per ml – 1% DV Oct-14 to 2017.....	256.31	240 ml	Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE – Restricted see terms on the preceding page			
⬆ Tab 600 mg with lamivudine 300 mg	630.00	30	Kivexa
DIDANOSINE [DDI] – Restricted see terms on the preceding page			
⬆ Cap 125 mg			
⬆ Cap 200 mg			
⬆ Cap 250 mg			
⬆ Cap 400 mg			
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPROXIL FUMARATE – Restricted see terms on the preceding page			
⬆ Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil fumarate 300 mg	1,313.19	30	Atripla
EMTRICITABINE – Restricted see terms on the preceding page			
⬆ Cap 200 mg	307.20	30	Emtriva
EMTRICITABINE WITH TENOFOVIR DISOPROXIL FUMARATE – Restricted see terms on the preceding page			
⬆ Tab 200 mg with tenofovir disoproxil fumarate 300 mg	838.20	30	Truvada
LAMIVUDINE – Restricted see terms on the preceding page			
⬆ Oral liq 10 mg per ml			
STAVUDINE – Restricted see terms on the preceding page			
⬆ Cap 30 mg			
⬆ Cap 40 mg			
⬆ Powder for oral soln 1 mg per ml			
ZIDOVUDINE [AZT] – Restricted see terms on the preceding page			
⬆ Cap 100 mg – 1% DV Oct-13 to 2016	152.25	100	Retrovir
⬆ Oral liq 10 mg per ml – 1% DV Oct-13 to 2016.....	30.45	200 ml	Retrovir
⬆ Inj 10 mg per ml, 20 ml vial – 1% DV Oct-14 to 2017.....	750.00	5	Retrovir IV
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Restricted see terms on the preceding page			
⬆ Tab 300 mg with lamivudine 150 mg – 1% DV Sep-14 to 2017	44.00	60	Alphapharm

Protease Inhibitors

➔ Restricted

Initiation — Confirmed HIV

Both:

- 1 Confirmed HIV infection; and
- 2 Any of the following:
 - 2.1 Symptomatic patient; or
 - 2.2 Patient aged 12 months and under; or
 - 2.3 Both:
 - 2.3.1 Patient aged 1 to 5 years; and
 - 2.3.2 Any of the following:
 - 2.3.2.1 CD4 counts < 1000 cells/mm³; or
 - 2.3.2.2 CD4 counts < 0.25 × total lymphocyte count; or
 - 2.3.2.3 Viral load counts > 100000 copies per ml; or
 - 2.4 Both:
 - 2.4.1 Patient aged 6 years and over; and
 - 2.4.2 CD4 counts < 500 cells/mm³.

Initiation — Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Initiation — Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initiation — Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

ATAZANAVIR SULPHATE – Restricted see terms above

⚡ Cap 150 mg	568.34	60	Reyataz
⚡ Cap 200 mg	757.79	60	Reyataz

DARUNAVIR – Restricted see terms above

⚡ Tab 400 mg	837.50	60	Prezista
⚡ Tab 600 mg	1,190.00	60	Prezista

INDINAVIR – Restricted see terms above

- ⚡ Cap 200 mg
- ⚡ Cap 400 mg

LOPINAVIR WITH RITONAVIR – Restricted see terms above

⚡ Tab 100 mg with ritonavir 25 mg	183.75	60	Kaletra
⚡ Tab 200 mg with ritonavir 50 mg	735.00	120	Kaletra
⚡ Oral liq 80 mg with ritonavir 20 mg per ml	735.00	300 ml	Kaletra

RITONAVIR – Restricted see terms above

⚡ Tab 100 mg	43.31	30	Norvir
⚡ Oral liq 80 mg per ml			

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Strand Transfer Inhibitors

➔Restricted

Initiation — Confirmed HIV

Both:

- 1 Confirmed HIV infection; and
- 2 Any of the following:
 - 2.1 Symptomatic patient; or
 - 2.2 Patient aged 12 months and under; or
 - 2.3 Both:
 - 2.3.1 Patient aged 1 to 5 years; and
 - 2.3.2 Any of the following:
 - 2.3.2.1 CD4 counts < 1000 cells/mm³; or
 - 2.3.2.2 CD4 counts < 0.25 × total lymphocyte count; or
 - 2.3.2.3 Viral load counts > 100000 copies per ml; or
 - 2.4 Both:
 - 2.4.1 Patient aged 6 years and over; and
 - 2.4.2 CD4 counts < 500 cells/mm³.

Initiation — Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Initiation — Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initiation — Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

RALTEGRAVIR POTASSIUM – **Restricted** see terms above

⬆ Tab 400 mg	1,090.00	60	Isentress
--------------------	----------	----	-----------

Antivirals

Hepatitis B

ADEFOVIR DIPIVOXIL – **Restricted** see terms below

⬆ Tab 10 mg	670.00	30	Hepsera
-------------------	--------	----	---------

➔Restricted

Initiation

Gastroenterologist or infectious disease specialist

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg+); and Documented resistance to lamivudine defined as:
- 2 Patient has raised serum ALT (> 1 × ULN); and
- 3 Patient has HBV DNA greater than 100,000 copies per mL, or viral load ≥ 10-fold over nadir; and
- 4 Detection of M204I or M204V mutation; and
- 5 Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
5.1 Both:			
5.1.1 Patient is cirrhotic; and			
5.1.2 Adefovir dipivoxil to be used in combination with lamivudine; or			
5.2 Both:			
5.2.1 Patient is not cirrhotic; and			
5.2.2 Adefovir dipivoxil to be used as monotherapy.			
ENTECAVIR – Restricted see terms below			
⚡ Tab 0.5 mg	400.00	30	Baraclude
➡ Restricted			
Initiation			
Gastroenterologist or infectious disease specialist			
All of the following:			
1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and			
2 Patient is Hepatitis B nucleoside analogue treatment-naïve; and			
3 Entecavir dose 0.5 mg/day; and			
4 Either:			
4.1 ALT greater than upper limit of normal; or			
4.2 Bridging fibrosis or cirrhosis (Metavir stage 3 or greater or moderate fibrosis) on liver histology; and			
5 Either:			
5.1 HBeAg positive; or			
5.2 Patient has ≥ 2,000 IU HBV DNA units per ml and fibrosis (Metavir stage 2 or greater) on liver histology; and			
6 No continuing alcohol abuse or intravenous drug use; and			
7 Not co-infected with HCV, HIV or HDV; and			
8 Neither ALT nor AST greater than 10 times upper limit of normal; and			
9 No history of hypersensitivity to entecavir; and			
10 No previous documented lamivudine resistance (either clinical or genotypic).			
LAMIVUDINE – Restricted see terms below			
⚡ Tab 100 mg – 1% DV Nov-14 to 2017	6.00	28	Zeffix
⚡ Oral liq 5 mg per ml – 1% DV Nov-14 to 2017	270.00	240 ml	Zeffix
➡ Restricted			
Initiation			
Gastroenterologist, infectious disease specialist, paediatrician or general physician			
<i>Limited to 12 months treatment</i>			
Any of the following:			
1 HBV DNA positive cirrhosis prior to liver transplantation; or			
2 HBsAg positive and have had a liver, kidney, heart, lung or bone marrow transplant; or			
3 Hepatitis B virus naïve patient who has received a liver transplant from an anti-HBc (Hepatitis B core antibody) positive donor; or			
4 Hepatitis B surface antigen positive (HbsAg) patient who is receiving chemotherapy for a malignancy, or who has received such treatment within the previous two months; or			
5 Hepatitis B surface antigen positive patient who is receiving anti tumour necrosis factor treatment; or			
6 Hepatitis B core antibody (anti-HBc) positive patient who is receiving rituximab plus high dose steroids (e.g. R-CHOP).			
Continuation — patients who have maintained continuous treatment and response to lamivudine			
Gastroenterologist, infectious disease specialist, paediatrician or general physician			
<i>Re-assessment required after 2 years</i>			
All of the following:			
1 Have maintained continuous treatment with lamivudine; and			
2 Most recent test result shows continuing biochemical response (normal ALT); and			
continued...			
⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below) e.g. <i>Brand</i> indicates brand example only. It is not a contracted product.			

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- HBV DNA <100,000 copies per ml by quantitative PCR at a reference laboratory.

Continuation — when given in combination with adefovir dipivoxil for patients with cirrhosis and resistance to lamivudine

Gastroenterologist, infectious disease specialist, paediatrician or general physician

Re-assessment required after 2 years

All of the following:

- Lamivudine to be used in combination with adefovir dipivoxil; and
- Patient is cirrhotic; and
Documented resistance to lamivudine defined as:
- All of the following:
 - Patient has raised serum ALT ($> 1 \times \text{ULN}$); and
 - Patient has HBV DNA greater than 100,000 copies per mL, or viral load ≥ 10 -fold over nadir; and
 - Detection of M204I or M204V mutation.

Continuation — when given in combination with adefovir dipivoxil for patients with resistance to adefovir dipivoxil

Gastroenterologist, infectious disease specialist, paediatrician or general physician

Re-assessment required after 2 years

Both:

- Lamivudine to be used in combination with adefovir dipivoxil; and
Documented resistance to lamivudine defined as:
- All of the following:
 - Patient has raised serum ALT ($> 1 \times \text{ULN}$); and
 - Patient has HBV DNA greater than 100,000 copies per mL, or viral load ≥ 10 -fold over nadir; and
 - Detection of N236T or A181T/V mutation.

TENOFOVIR DISOPROXIL FUMARATE – **Restricted** see terms below

⚡ Tab 300 mg	531.00	30	Viread
--------------------	--------	----	--------

➡ **Restricted**

Initiation — Confirmed hepatitis B

Any of the following:

- All of the following:
 - Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
 - Patient has had previous lamivudine, adefovir or entecavir therapy; and
 - HBV DNA greater than 20,000 IU/mL or increased ≤ 10 -fold over nadir; and
 - Any of the following:
 - Lamivudine resistance - detection of M204I/V mutation; or
 - Adefovir resistance - detection of A181T/V or N236T mutation; or
 - Entecavir resistance - detection of relevant mutations including I169T, L180M T184S/A/I/L/G/C/M, S202C/G/I, M204V or M250I/V mutation; or
- Patient is either listed or has undergone liver transplantation for HBV; or
- Patient has a decompensated cirrhosis with a Mayo score > 20 .

Initiation — Pregnant or Breastfeeding, Active hepatitis B

Limited to 12 months treatment

Both:

- Patient is HBsAg positive and pregnant; and
- HBV DNA $> 20,000$ IU/mL and ALT $> \text{ULN}$.

Initiation — Pregnant, prevention of vertical transmission

Limited to 6 months treatment

Both:

- Patient is HBsAg positive and pregnant; and
- HBV DNA > 20 million IU/mL and ALT normal.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation — Confirmed HIV

Both:

- 1 Confirmed HIV infection; and
- 2 Any of the following:
 - 2.1 Symptomatic patient; or
 - 2.2 Patient aged 12 months and under; or
 - 2.3 Both:
 - 2.3.1 Patient aged 1 to 5 years; and
 - 2.3.2 Any of the following:
 - 2.3.2.1 CD4 counts < 1000 cells/mm³; or
 - 2.3.2.2 CD4 counts < 0.25 × total lymphocyte count; or
 - 2.3.2.3 Viral load counts > 100000 copies per ml; or
 - 2.4 Both:
 - 2.4.1 Patient aged 6 years and over; and
 - 2.4.2 CD4 counts < 500 cells/mm³.

Initiation — Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Initiation — Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initiation — Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

Hepatitis C

BOCEPREVIR – **Restricted** see terms below

⚡ Cap 200 mg	5,015.00	336	Victrelis
--------------------	----------	-----	-----------

➡Restricted

Initiation — Chronic hepatitis C - genotype 1, first-line

Gastroenterologist, infectious disease specialist or general physician

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has not received prior pegylated interferon treatment; and
- 3 Patient has IL-28B genotype CT or TT; and
- 4 Patient is to be treated in combination with pegylated interferon and ribavirin; and
- 5 Patient is hepatitis C protease inhibitor treatment-naïve; and
- 6 Maximum of 44 weeks therapy.

Initiation — Chronic hepatitis C - genotype 1, second-line

Gastroenterologist, infectious disease specialist or general physician

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has received pegylated interferon treatment; and

continued...

Price (ex man. excl. GST)	Brand or Generic Manufacturer
\$	Per

continued. . .

- 3 Any of the following:
 - 3.1 Patient was a responder relapser; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received pegylated interferon prior to 2004; and
- 4 Patient is to be treated in combination with pegylated interferon and ribavirin; and
- 5 Maximum of 44 weeks therapy.

Note: Due to risk of severe sepsis boceprevir should not be initiated if either Platelet count <100 x10⁹/l or Albumin <5 g/l.

Herpesviridae

ACICLOVIR

Tab dispersible 200 mg – 1% DV Sep-13 to 2016	1.78	25	Lovir
Tab dispersible 400 mg – 1% DV Sep-13 to 2016	5.98	56	Lovir
Tab dispersible 800 mg – 1% DV Sep-13 to 2016	6.64	35	Lovir
Inj 250 mg vial – 1% DV Jan-16 to 2018	10.10	5	Aciclovir-Clarix

CIDOFOVIR – **Restricted** see terms below

⚡ Inj 75 mg per ml, 5 ml vial

➡ **Restricted**

Clinical microbiologist, infectious disease specialist, otolaryngologist or oral surgeon

FOSCARNET SODIUM – **Restricted** see terms below

⚡ Inj 24 mg per ml, 250 ml bottle

➡ **Restricted**

Clinical microbiologist or infectious disease specialist

GANCICLOVIR – **Restricted** see terms below

⚡ Inj 500 mg vial 380.00 5 Cymevene

➡ **Restricted**

Clinical microbiologist or infectious disease specialist

VALACICLOVIR

Tab 500 mg – 1% DV Mar-16 to 2018	6.42	30	Vaclovir
Tab 1,000 mg – 1% DV Mar-16 to 2018	12.75	30	Vaclovir

VALGANCICLOVIR – **Restricted** see terms on the next page

⚡ Tab 450 mg – 1% DV Jun-15 to 2018 1,050.00 60 Valcyte

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➡Restricted

Initiation — Transplant cytomegalovirus prophylaxis

Limited to 3 months treatment

Patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Initiation — Lung transplant cytomegalovirus prophylaxis

Limited to 6 months treatment

Both:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive.

Initiation — Cytomegalovirus in immunocompromised patients

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Influenza

OSELTAMIVIR – **Restricted** see terms below

⚡ Tab 75 mg

⚡ Powder for oral suspension 6 mg per ml

➡Restricted

Initiation

Either:

- 1 Only for hospitalised patient with known or suspected influenza; or
- 2 For prophylaxis of influenza in hospitalised patients as part of a DHB hospital approved infections control plan.

ZANAMIVIR

⚡ Powder for inhalation 5 mg37.38 20 dose Relenza Rotadisk

➡Restricted

Initiation

Either:

- 1 Only for hospitalised patient with known or suspected influenza; or
- 2 For prophylaxis of influenza in hospitalised patients as part of a DHB hospital approved infections control plan.

Immune Modulators

INTERFERON ALFA-2A

Inj 3 m iu prefilled syringe

Inj 6 m iu prefilled syringe

Inj 9 m iu prefilled syringe

INTERFERON ALFA-2B

Inj 18 m iu, 1.2 ml multidose pen

Inj 30 m iu, 1.2 ml multidose pen

Inj 60 m iu, 1.2 ml multidose pen

INTERFERON GAMMA – **Restricted** see terms below

⚡ Inj 100 mcg in 0.5 ml vial

➡Restricted

Initiation

Patient has chronic granulomatous disease and requires interferon gamma.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PEGYLATED INTERFERON ALFA-2A – Restricted see terms below			
¶ Inj 135 mcg prefilled syringe			
¶ Inj 135 mcg prefilled syringe (4) with ribavirin tab 200 mg (112)			
¶ Inj 135 mcg prefilled syringe (4) with ribavirin tab 200 mg (168)			
¶ Inj 180 mcg prefilled syringe	900.00	4	Pegasys
¶ Inj 180 mcg prefilled syringe (4) with ribavirin tab 200 mg (112)	1,159.84	1	Pegasys RBV Combination Pack
¶ Inj 180 mcg prefilled syringe (4) with ribavirin tab 200 mg (168)	1,290.00	1	Pegasys RBV Combination Pack

(Any Inj 135 mcg prefilled syringe to be delisted 1 July 2016)

(Any Inj 135 mcg prefilled syringe (4) with ribavirin tab 200 mg (112) to be delisted 1 July 2016)

➡ Restricted

Initiation — Chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant

Limited to 48 weeks treatment

Any of the following:

- 1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
- 2 Patient has chronic hepatitis C and is co-infected with HIV; or
- 3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant.

Notes: Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.

Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml.

Continuation — Chronic hepatitis C - genotype 1 infection

Gastroenterologist, infectious disease specialist or general physician

Re-assessment required after 48 weeks

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir.

Initiation — Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior

Gastroenterologist, infectious disease specialist or general physician

Limited to 48 weeks treatment

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation — Chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV

Limited to 6 months treatment

Patient has chronic hepatitis C, genotype 2 or 3 infection.

Initiation — Hepatitis B

Gastroenterologist, infectious disease specialist or general physician

Limited to 48 weeks treatment

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naïve; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 Serum HBV DNA ≥ 2,000 units/ml and significant fibrosis (≥ Metavir Stage F2 or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon.

Notes: Approved dose is 180 mcg once weekly.

The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.

In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon alfa-2a dose should be reduced to 135 mcg once weekly.

In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.

Pegylated Interferon alfa-2a is not approved for use in children.

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Anticholinesterases

EDROPHONIUM CHLORIDE – **Restricted** see terms below

- ⚡ Inj 10 mg per ml, 15 ml vial
- ⚡ Inj 10 mg per ml, 1 ml ampoule

➔**Restricted**

Initiation

For the diagnosis of myasthenia gravis.

NEOSTIGMINE METILSULFATE

Inj 2.5 mg per ml, 1 ml ampoule – **1% DV Sep-14 to 2017** 98.00 50 **AstraZeneca**

NEOSTIGMINE METILSULFATE WITH GLYCOPYRRONIUM BROMIDE

Inj 2.5 mg with glycopyrronium bromide 0.5 mg per ml, 1 ml ampoule
– **1% DV Nov-13 to 2016** 20.90 10 **Max Health**
Max Health brand - HSS with 1% DV will apply 1 July 2016 to 30 June 2019.

PYRIDOSTIGMINE BROMIDE

Tab 60 mg 38.90 100 Mestinon

Antirheumatoid Agents

AURANOFIN

Tab 3 mg

HYDROXYCHLOROQUINE

Tab 200 mg – **1% DV Sep-15 to 2018** 10.50 100 **Plaquenil**

LEFLUNOMIDE

Tab 10 mg 55.00 30 Arava
Tab 20 mg 76.00 30 Arava

PENICILLAMINE

Tab 125 mg 61.93 100 D-Penamine
Tab 250 mg 98.98 100 D-Penamine

SODIUM AUROTHIOMALATE

Inj 10 mg in 0.5 ml ampoule
Inj 20 mg in 0.5 ml ampoule
Inj 50 mg in 0.5 ml ampoule

Drugs Affecting Bone Metabolism

Bisphosphonates

ALENDRONATE SODIUM

⚡ Tab 40 mg 133.00 30 Fosamax

➔**Restricted**

Initiation — Paget's disease

Both:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications due to site (base of skull, spine, long bones of lower limbs); or
 - 2.5 Preparation for orthopaedic surgery.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
⚡ Tab 70 mg	12.90	4	Fosamax

➡ Restricted

Initiation — Osteoporosis

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) ≥ 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score ≤ -2.5) (see Note); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score ≤ -3.0 (see Note); or
- 5 A 10-year risk of hip fracture $\geq 3\%$, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 6 Patient has had a Special Authority approval for zoledronic acid (underlying cause – osteoporosis) or raloxifene.

Initiation — glucocorticosteroid therapy

Re-assessment required after 12 months

Both:

- 1 The patient is receiving systemic glucocorticosteroid therapy (≥ 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD ≥ 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score ≤ -1.5) (see Note); or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for zoledronic acid (glucocorticosteroid therapy) or raloxifene.

Continuation — glucocorticosteroid therapy

Re-assessment required after 12 months

The patient is continuing systemic glucocorticosteroid therapy (≥ 5 mg per day prednisone equivalents).

Notes:

- 1 BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- 2 Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score ≤ -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- 3 Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- 4 A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

ALENDRONATE SODIUM WITH CHOLECALCIFEROL – **Restricted** see terms on the next page

⚡ Tab 70 mg with cholecalciferol 5,600 iu	12.90	4	Fosamax Plus
---	-------	---	--------------

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

➔ **Restricted**

Initiation — Osteoporosis

Any of the following:

- History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) ≥ 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score ≤ -2.5) (see Note); or
- History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- History of two significant osteoporotic fractures demonstrated radiologically; or
- Documented T-Score ≤ -3.0 (see Note); or
- A 10-year risk of hip fracture $\geq 3\%$, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- Patient has had a Special Authority approval for zoledronic acid (underlying cause – osteoporosis) or raloxifene.

Initiation — glucocorticosteroid therapy

Re-assessment required after 12 months

Both:

- The patient is receiving systemic glucocorticosteroid therapy (≥ 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- Any of the following:
 - The patient has documented BMD ≥ 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score ≤ -1.5) (see Note); or
 - The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - The patient has had a Special Authority approval for zoledronic acid (glucocorticosteroid therapy) or raloxifene.

Continuation — glucocorticosteroid therapy

Re-assessment required after 12 months

The patient is continuing systemic glucocorticosteroid therapy (≥ 5 mg per day prednisone equivalents).

Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score ≥ -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

ETIDRONATE DISODIUM

Tab 200 mg – 1% DV Sep-15 to 2018	13.50	100	Arrow-Etidronate
---	-------	-----	-------------------------

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	6.80	1	Pamisol
Inj 6 mg per ml, 10 ml vial	13.20	1	Pamisol
Inj 9 mg per ml, 10 ml vial	19.20	1	Pamisol

RISEDRONATE SODIUM

Tab 35 mg	4.00	4	Risedronate Sandoz
-----------------	------	---	--------------------

ZOLEDRONIC ACID

⚡ Inj 5 mg per 100 ml, vial	600.00	100 ml	Aclasta
-----------------------------------	--------	--------	---------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➡Restricted			
Initiation — Inherited bone fragility disorders			
Any specialist			
Patient has been diagnosed with an inherited bone fragility disorder (e.g. osteogenesis imperfecta).			
Initiation — Osteoporosis			
Any specialist			
<i>Therapy limited to 3 doses</i>			
Both:			
1 Any of the following:			
1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) \geq 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score \leq -2.5) (see Note); or			
1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or			
1.3 History of two significant osteoporotic fractures demonstrated radiologically; or			
1.4 Documented T-Score \geq -3.0 (see Note); or			
1.5 A 10-year risk of hip fracture \geq 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or			
1.6 Patient has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) or raloxifene; and			
2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.			
Initiation — glucocorticosteroid therapy			
Any specialist			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 The patient is receiving systemic glucocorticosteroid therapy (\geq 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and			
2 Any of the following:			
2.1 The patient has documented BMD \geq 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score \leq -1.5) (see Note); or			
2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or			
2.3 The patient has had a Special Authority approval for alendronate (Underlying cause - glucocorticosteroid therapy) or raloxifene; and; and			
3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.			
Continuation — glucocorticosteroid therapy			
Any specialist			
<i>Re-assessment required after 12 months</i>			
Both:			
1 The patient is continuing systemic glucocorticosteroid therapy (\geq 5 mg per day prednisone equivalents); and			
2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.			
Initiation — Paget's disease			
Any specialist			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 Paget's disease; and			
2 Any of the following:			
2.1 Bone or articular pain; or			
2.2 Bone deformity; or			
2.3 Bone, articular or neurological complications; or			
2.4 Asymptomatic disease, but risk of complications; or			
continued...			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2.5 Preparation for orthopaedic surgery; and

3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Continuation — Paget's disease

Any specialist

Re-assessment required after 12 months

- Both:
- Any of the following:
 - The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - Symptomatic disease (prescriber determined); and
 - The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

- Notes:
- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
 - Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score ≤ -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
 - Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
 - A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Other Drugs Affecting Bone Metabolism

RALOXIFENE – **Restricted** see terms below

Tab 60 mg	53.76	28	Evista
-----------------	-------	----	--------

➡Restricted

Initiation

- Any of the following:
- History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) ≥ 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score ≤ -2.5) (see Notes); or
 - History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - History of two significant osteoporotic fractures demonstrated radiologically; or
 - Documented T-Score ≥ -3.0 (see Notes); or
 - A 10-year risk of hip fracture $\geq 3\%$, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
 - Patient has had a prior Special Authority approval for zoledronic acid (Underlying cause - Osteoporosis) or alendronate (Underlying cause - Osteoporosis).

- Notes:
- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score ≤ -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

TERIPARATIDE – **Restricted** see terms below

¶ Inj 250 mcg per ml, 2.4 ml cartridge490.00 1 Forteo

➡ **Restricted**

Initiation

Limited to 18 months treatment

All of the following:

- The patient has severe, established osteoporosis; and
- The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- The patient has had two or more fractures due to minimal trauma; and
- The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- Antiresorptive agents and their adequate doses for the purposes of this restriction are defined as: alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Enzymes

HYALURONIDASE

Inj 1,500 iu ampoule

Hyperuricaemia and Antigout

ALLOPURINOL

Tab 100 mg – 1% DV Mar-15 to 2017	15.11	1,000	Apo-Allopurinol
Tab 300 mg – 1% DV Mar-15 to 2017	15.91	500	Apo-Allopurinol

BENZBROMARONE – **Restricted** see terms on the next page

¶ Tab 100 mg45.00 100 Benzbromaron AL 100

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

➔ **Restricted**

Initiation

Any specialist

All of the following:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 Both:
 - 2.3.1 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); and
 - 2.3.2 The patient has a rate of creatinine clearance greater than or equal to 20 ml/min; or
 - 2.4 All of the following:
 - 2.4.1 The patient is taking azathioprine and requires urate-lowering therapy; and
 - 2.4.2 Allopurinol is contraindicated; and
 - 2.4.3 Appropriate doses of probenecid are ineffective or probenecid cannot be used due to reduced renal function; and
- 3 The patient is receiving monthly liver function tests.

Notes: Benzbromarone has been associated with potentially fatal hepatotoxicity. In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

The New Zealand Rheumatology Association has developed information for prescribers which can be accessed from its website at www.rheumatology.org.nz/downloads/Benzbromarone-prescriber-information-NZRA-V2.pdf

COLCHICINE

Tab 500 mcg – 1% DV Oct-13 to 2016	10.08	100	Colgout
--	-------	-----	----------------

FEBUXOSTAT – Restricted see terms below

⚡ Tab 80 mg	39.50	28	Adenuric
⚡ Tab 120 mg	39.50	28	Adenuric

➔ **Restricted**

Initiation

Any specialist

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note).

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

MUSCULOSKELETAL SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PROBENECID Tab 500 mg			
RASBURICASE – Restricted see terms below ☞ Inj 1.5 mg vial ☞ Restricted Haematologist			
Muscle Relaxants and Related Agents			
ATRACURIUM BESYLATE Inj 10 mg per ml, 2.5 ml ampoule – 1% DV Jan-16 to 2018 Inj 10 mg per ml, 5 ml ampoule – 1% DV Jan-16 to 2018	10.00 12.50	5 5	Tracrium Tracrium
BACLOFEN Tab 10 mg – 1% DV Jun-13 to 2016 Oral liq 1 mg per ml Inj 0.05 mg per ml, 1 ml ampoule – 1% DV Sep-15 to 2018 Inj 2 mg per ml, 5 ml ampoule	3.85 11.55 209.29	100 1 1	Pacifen Lioresal Intrathecal Lioresal Intrathecal
CLOSTRIDIUM BOTULINUM TYPE A TOXIN Inj 100 u vial Inj 300 u vial Inj 500 u vial	467.50 388.50 1,295.00	1 1 2	Botox Dysport Dysport
DANTROLENE Cap 25 mg Cap 50 mg Inj 20 mg vial	65.00 77.00 800.00	100 100 6	Dantrium Dantrium Dantrium IV
MIVACURIUM CHLORIDE Inj 2 mg per ml, 5 ml ampoule Inj 2 mg per ml, 10 ml ampoule	33.92 67.17	5 5	Mivacron Mivacron
ORPHENADRINE CITRATE Tab 100 mg			
PANCURONIUM BROMIDE Inj 2 mg per ml, 2 ml ampoule	260.00	50	AstraZeneca
ROCURONIUM BROMIDE Inj 10 mg per ml, 5 ml vial – 1% DV Aug-16 to 2019	25.95	10	DBL Rocuronium Bromide
SUXAMETHONIUM CHLORIDE Inj 50 mg per ml, 2 ml ampoule – 1% DV Jun-14 to 2017	78.00	50	AstraZeneca
VECURONIUM BROMIDE Inj 4 mg ampoule Inj 10 mg vial			

Reversers of Neuromuscular Blockade

SUGAMMADEX – Restricted see terms on the next page ☞ Inj 100 mg per ml, 2 ml vial ☞ Inj 100 mg per ml, 5 ml vial	1,200.00 3,000.00	10 10	Bridion Bridion
---	----------------------	----------	--------------------

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

➔ **Restricted**

Initiation

Any of the following:

- 1 Patient requires reversal of profound neuromuscular blockade following rapid sequence induction that has been undertaken using rocuronium (i.e. suxamethonium is contraindicated or undesirable); or
- 2 Severe neuromuscular degenerative disease where the use of neuromuscular blockade is required; or
- 3 Patient has an unexpectedly difficult airway that cannot be intubated and requires a rapid reversal of anaesthesia and neuromuscular blockade; or
- 4 The duration of the patient's surgery is unexpectedly short; or
- 5 Neostigmine or a neostigmine/anticholinergic combination is contraindicated (for example the patient has ischaemic heart disease, morbid obesity or COPD); or
- 6 Patient has a partial residual block after conventional reversal.

Non-Steroidal Anti-Inflammatory Drugs

CELECOXIB – **Restricted** see terms below

- ⚡ Cap 100 mg
- ⚡ Cap 200 mg
- ⚡ Cap 400 mg

➔ **Restricted**

Initiation

For preoperative and/or postoperative use for a total of up to 8 days' use.

DICLOFENAC SODIUM

Tab EC 25 mg – 1% DV Dec-15 to 2018	1.30	50	Diclofenac Sandoz
Tab 50 mg dispersible	1.50	20	Voltaren D
Tab EC 50 mg – 1% DV Dec-15 to 2018	1.00	50	Diclofenac Sandoz
Tab long-acting 75 mg – 1% DV Dec-15 to 2018	15.20	500	Apo-Diclo SR
Tab long-acting 100 mg – 1% DV Dec-15 to 2018	26.20	500	Apo-Diclo SR
Inj 25 mg per ml, 3 ml ampoule – 1% DV Oct-14 to 2017	13.20	5	Voltaren
Suppos 12.5 mg – 1% DV Oct-14 to 2017	2.04	10	Voltaren
Suppos 25 mg – 1% DV Oct-14 to 2017	2.44	10	Voltaren
Suppos 50 mg – 1% DV Oct-14 to 2017	4.22	10	Voltaren
Suppos 100 mg – 1% DV Oct-14 to 2017	7.00	10	Voltaren

ETORICOXIB – **Restricted** see terms below

- ⚡ Tab 30 mg
- ⚡ Tab 60 mg
- ⚡ Tab 90 mg
- ⚡ Tab 120 mg

➔ **Restricted**

Initiation

For preoperative and/or postoperative use for a total of up to 8 days' use.

IBUPROFEN

Tab 200 mg			
➔ Tab 400 mg – Restricted : For continuation only			
➔ Tab 600 mg – Restricted : For continuation only			
Tab long-acting 800 mg – 1% DV Jul-15 to 2018	7.99	30	Brufen SR
Oral liq 20 mg per ml – 1% DV Mar-14 to 2016	1.89	200 ml	Fenpaed
Inj 5 mg per ml, 2 ml ampoule			
Inj 10 mg per ml, 2 ml vial			

MUSCULOSKELETAL SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
INDOMETHACIN			
Cap 25 mg			
Cap 50 mg			
Cap long-acting 75 mg			
Inj 1 mg vial			
Suppos 100 mg			
KETOPROFEN			
Cap long-acting 200 mg	12.07	28	Oruvail SR
MEFENAMIC ACID – Restricted: For continuation only			
➡ Cap 250 mg			
MELOXICAM – Restricted see terms below			
⚡ Tab 7.5 mg			
➡Restricted			
Initiation			
Either:			
1 All of the following:			
1.1 Haemophilic arthropathy; and			
1.2 The patient has moderate to severe haemophilia with less than or equal to 5% of normal circulating functional clotting factor; and			
1.3 Pain and inflammation associated with haemophilic arthropathy is inadequately controlled by alternative funded treatment options, or alternative funded treatment options are contraindicated; or			
2 For preoperative and/or postoperative use for a total of up to 8 days' use.			
NAPROXEN			
Tab 250 mg – 1% DV Sep-15 to 2018	18.06	500	Noflam 250
Tab 500 mg – 1% DV Sep-15 to 2018	18.91	250	Noflam 500
Tab long-acting 750 mg – 1% DV Jun-15 to 2018	18.00	90	Naprosyn SR 750
Tab long-acting 1 g – 1% DV Jun-15 to 2018	21.00	90	Naprosyn SR 1000
PARECOXIB			
Inj 40 mg vial	100.00	10	Dynastat
SULINDAC			
Tab 100 mg			
Tab 200 mg			
TENOXICAM			
Tab 20 mg – 1% DV Jan-15 to 2016	3.05	20	Reutenox
Inj 20 mg vial	9.95	1	AFT

Topical Products for Joint and Muscular Pain

CAPSAICIN – Restricted see terms below

⚡ Crm 0.025% 9.95 45 g Zostrix

➡Restricted

Initiation

Patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Price (ex man. excl. GST)	Brand or Generic
\$	Per Manufacturer

Agents for Parkinsonism and Related Disorders

Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE – **Restricted** see terms below

⚡ Tab 50 mg 400.00 56 Rilutek

➡ **Restricted**

Initiation

Neurologist or respiratory specialist

Re-assessment required after 6 months

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
 - 5.1 The patient is ambulatory; or
 - 5.2 The patient is able to use upper limbs; or
 - 5.3 The patient is able to swallow.

Continuation

Re-assessment required after 18 months

All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
 - 3.1 The patient is ambulatory; or
 - 3.2 The patient is able to use upper limbs; or
 - 3.3 The patient is able to swallow.

TETRABENAZINE

Tab 25 mg – **1% DV Sep-13 to 2016** 118.00 112 **Motetis**

Anticholinergics

BENZTROPINE MESYLATE

Tab 2 mg 7.99 60 Benztrop

Inj 1 mg per ml, 2 ml ampoule 95.00 5 Cogentin

PROCYCLIDINE HYDROCHLORIDE

Tab 5 mg

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE

Cap 100 mg – **1% DV Oct-14 to 2017** 38.24 60 **Symmetrel**

APOMORPHINE HYDROCHLORIDE

Inj 10 mg per ml, 1 ml ampoule

Inj 10 mg per ml, 2 ml ampoule 119.00 5 Apomine

Movapo

(Apomine Inj 10 mg per ml, 2 ml ampoule to be delisted 1 August 2016)

BROMOCRIPTINE

Tab 2.5 mg

Cap 5 mg

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ENTACAPONE			
Tab 200 mg – 1% DV Sep-15 to 2018	28.00	100	Entapone
LEVODOPA WITH BENSERAZIDE			
Tab dispersible 50 mg with benserazide 12.5 mg	10.00	100	Madopar Rapid
Cap 50 mg with benserazide 12.5 mg	8.00	100	Madopar 62.5
Cap 100 mg with benserazide 25 mg	12.50	100	Madopar 125
Cap long-acting 100 mg with benserazide 25 mg	17.00	100	Madopar HBS
Cap 200 mg with benserazide 50 mg	25.00	100	Madopar 250
LEVODOPA WITH CARBIDOPA			
Tab 100 mg with carbidopa 25 mg	20.00	100	Sinemet <i>e.g. Kinson</i>
Tab long-acting 200 mg with carbidopa 50 mg	47.50	100	Sinemet CR
Tab 250 mg with carbidopa 25 mg	40.00	100	Sinemet <i>e.g. Sindopa</i>
LISURIDE HYDROGEN MALEATE			
Tab 200 mcg	25.00	30	Dopergin
<i>(Dopergin Tab 200 mcg to be delisted 1 September 2016)</i>			
PRAMIPEXOLE HYDROCHLORIDE			
Tab 0.25 mg – 1% DV Oct-14 to 2016	7.20	100	Ramipex
Tab 1 mg – 1% DV Oct-14 to 2016	24.39	100	Ramipex
ROPINIROLE HYDROCHLORIDE			
Tab 0.25 mg – 1% DV Mar-14 to 2016	2.36	100	Apo-Ropinirole
Tab 1 mg – 1% DV Mar-14 to 2016	5.32	100	Apo-Ropinirole
Tab 2 mg – 1% DV Mar-14 to 2016	7.72	100	Apo-Ropinirole
Tab 5 mg – 1% DV Mar-14 to 2016	14.48	100	Apo-Ropinirole
SELEGILINE HYDROCHLORIDE			
Tab 5 mg			
TOLCAPONE			
Tab 100 mg	126.20	100	Tasmar
Anaesthetics			
General Anaesthetics			
DESFLURANE			
Soln for inhalation 100%, 240 ml bottle	1,414.50	6	Suprane
DEXMEDETOMIDINE			
Inj 100 mcg per ml, 2 ml vial – 1% DV Oct-14 to 2017	479.85	5	Precedex
ETOMIDATE			
Inj 2 mg per ml, 10 ml ampoule			
ISOFLURANE			
Soln for inhalation 100%, 250 ml bottle	1,173.00	6	Aerrane
KETAMINE			
Inj 1 mg per ml, 100 ml bag – 1% DV Sep-14 to 2017	27.00	1	Biomed
Inj 4 mg per ml, 50 ml syringe – 1% DV Sep-14 to 2017	25.00	1	Biomed
Inj 10 mg per ml, 10 ml syringe – 1% DV Sep-14 to 2017	14.00	1	Biomed
Inj 100 mg per ml, 2 ml ampoule – 1% DV May-16 to 2018	47.05	5	Ketamine-Claris

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
METHOHEXITAL SODIUM			
Inj 10 mg per ml, 50 ml vial			
PROPOFOL			
Inj 10 mg per ml, 20 ml vial – 10% DV Jun-16 to 2019	5.27	5	Provide MCT-LCT 1%
Inj 10 mg per ml, 50 ml vial – 10% DV Jun-16 to 2019	24.50	10	Fresofol 1% MCT/LCT
Inj 10 mg per ml, 100 ml vial – 10% DV Jun-16 to 2019	49.00	10	Fresofol 1% MCT/LCT
SEVOFLURANE			
Soln for inhalation 100%, 250 ml bottle	1,365.00	6	Baxter
THIOPENTAL [THIOPENTONE] SODIUM			
Inj 500 mg ampoule			

Local Anaesthetics

ARTICAINE HYDROCHLORIDE

Inj 1%

ARTICAINE HYDROCHLORIDE WITH ADRENALINE

Inj 4% with adrenaline 1:100,000, 1.7 ml dental cartridge

Inj 4% with adrenaline 1:100,000, 2.2 ml dental cartridge

Inj 4% with adrenaline 1:200,000, 1.7 ml dental cartridge

Inj 4% with adrenaline 1:200,000, 2.2 ml dental cartridge

BENZOCAINE

Gel 20%

BUPIVACAINE HYDROCHLORIDE

Inj 5 mg per ml, 4 ml ampoule – **1% DV Jul-14 to 2017**50.00

5

Marcaïn Isobaric

Inj 2.5 mg per ml, 20 ml ampoule

Inj 2.5 mg per ml, 20 ml ampoule sterile pack – **1% DV Sep-15 to 2018**29.20

5

Marcaïn

Inj 5 mg per ml, 10 ml ampoule sterile pack – **1% DV Sep-15 to 2018**.....20.25

5

Marcaïn

Inj 5 mg per ml, 20 ml ampoule

Inj 5 mg per ml, 20 ml ampoule sterile pack – **1% DV Sep-15 to 2018**.....20.70

5

Marcaïn

Inj 1.25 mg per ml, 100 ml bag

Inj 1.25 mg per ml, 200 ml bag

Inj 2.5 mg per ml, 100 ml bag – **1% DV Jul-14 to 2017** 150.00

5

Marcaïn

Inj 2.5 mg per ml, 200 ml bag

Inj 1.25 mg per ml, 500 ml bag

BUPIVACAINE HYDROCHLORIDE WITH ADRENALINE

Inj 2.5 mg per ml with adrenaline 1:400,000, 20 ml vial – **1% DV Sep-**

14 to 2017 135.00

5

**Marcaïn with
Adrenaline**

Inj 5 mg per ml with adrenaline 1:200,000, 20 ml vial – **1% DV Sep-14**

to 2017 115.00

5

**Marcaïn with
Adrenaline**

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BUPIVACAINE HYDROCHLORIDE WITH FENTANYL			
Inj 0.625 mg with fentanyl 2 mcg per ml, 100 ml bag			
Inj 0.625 mg with fentanyl 2 mcg per ml, 200 ml bag			
Inj 1.25 mg with fentanyl 2 mcg per ml, 100 ml syringe			
Inj 1.25 mg with fentanyl 2 mcg per ml, 100 ml bag	210.00	10	Bupafen
Inj 1.25 mg with fentanyl 2 mcg per ml, 200 ml bag	210.00	10	Bupafen
Inj 1.25 mg with fentanyl 2 mcg per ml, 50 ml syringe			
Inj 1.25 mg with fentanyl 2 mcg per ml, 15 ml syringe	72.00	10	Biomed
Inj 1.25 mg with fentanyl 2 mcg per ml, 20 ml syringe	92.00	10	Biomed
BUPIVACAINE HYDROCHLORIDE WITH GLUCOSE			
Inj 0.5% with glucose 8%, 4 ml ampoule	38.00	5	Marcaïn Heavy
COCAINE HYDROCHLORIDE			
Paste 5%			
Soln 15%, 2 ml syringe			
Soln 4%, 2 ml syringe	25.46	1	Biomed
COCAINE HYDROCHLORIDE WITH ADRENALINE			
Paste 15% with adrenaline 0.06%			
Paste 25% with adrenaline 0.06%			
ETHYL CHLORIDE			
Spray 100%			
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE			
Gel 2% – 1% DV Sep-15 to 2018	3.40	20 ml	Orion
Soln 4%			
Spray 10% – 1% DV Sep-13 to 2016	75.00	50 ml	Xylocaine
Oral (viscous) soln 2% – 1% DV Sep-14 to 2017	55.00	200 ml	Xylocaine Viscous
Inj 1%, 20 ml ampoule, sterile pack			
Inj 2%, 20 ml ampoule, sterile pack			
Inj 1%, 5 ml ampoule	8.75	25	Lidocaine-Claris
Inj 1%, 20 ml ampoule	2.40	1	Lidocaine-Claris
Inj 2%, 5 ml ampoule	6.90	25	Lidocaine-Claris
Inj 2%, 20 ml ampoule	2.40	1	Lidocaine-Claris
Gel 2%, 10 ml urethral syringe	43.26	10	Pfizer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH ADRENALINE			
Inj 1% with adrenaline 1:100,000, 5 ml ampoule	27.00	10	Xylocaine
Inj 1% with adrenaline 1:200,000, 20 ml vial	50.00	5	Xylocaine
Inj 2% with adrenaline 1:80,000, 1.7 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 1.8 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 2.2 ml dental cartridge			
Inj 2% with adrenaline 1:200,000, 20 ml vial	60.00	5	Xylocaine
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH ADRENALINE AND TETRACAINE HYDROCHLORIDE			
Soln 4% with adrenaline 0.1% and tetracaine hydrochloride 0.5%, 5 ml syringe – 1% DV Oct-14 to 2017	17.50	1	Topicaïne
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH CHLORHEXIDINE			
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringe	43.26	10	Pfizer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH PHENYLEPHRINE HYDROCHLORIDE			
Nasal spray 5% with phenylephrine hydrochloride 0.5%			

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE			
Crm 2.5% with prilocaine 2.5%	45.00	30 g	EMLA
Patch 25 mcg with prilocaine 25 mcg	115.00	20	EMLA
Crm 2.5% with prilocaine 2.5%, 5 g	45.00	5	EMLA
LIDOCAINE [LIGNOCAINE]			
Crm 4%	27.00	30 g	LMX4
Crm 4% (5 g tubes)	27.00	5	LMX4
MEPIVACAINE HYDROCHLORIDE			
Inj 3%, 1.8 ml dental cartridge – 1% DV Oct-14 to 2017	43.60	50	Scandonest 3%
Inj 3%, 2.2 ml dental cartridge – 1% DV Oct-14 to 2017	43.60	50	Scandonest 3%
PRILOCAINE HYDROCHLORIDE			
Inj 0.5%, 50 ml vial	100.00	5	Citanest
Inj 2%, 5 ml ampoule	55.00	10	Citanest
PRILOCAINE HYDROCHLORIDE WITH FELYPRESSIN			
Inj 3% with felypressin 0.03 iu per ml, 1.8 ml dental cartridge			
Inj 3% with felypressin 0.03 iu per ml, 2.2 ml dental cartridge			
ROPIVACAINE HYDROCHLORIDE			
Inj 2 mg per ml, 10 ml ampoule – 1% DV Aug-15 to 2017	9.05	5	Ropivacaine Kabi
Inj 2 mg per ml, 20 ml ampoule – 1% DV Aug-15 to 2017	9.50	5	Ropivacaine Kabi
Inj 2 mg per ml, 100 ml bag – 1% DV Jul-15 to 2017	60.00	5	Naropin
Inj 2 mg per ml, 200 ml bag – 1% DV Jul-15 to 2017	79.50	5	Naropin
Inj 7.5 mg per ml, 10 ml ampoule – 1% DV Aug-15 to 2017	10.20	5	Ropivacaine Kabi
Inj 7.5 mg per ml, 20 ml ampoule – 1% DV Aug-15 to 2017	12.50	5	Ropivacaine Kabi
Inj 10 mg per ml, 10 ml ampoule – 1% DV Aug-15 to 2017	10.90	5	Ropivacaine Kabi
Inj 10 mg per ml, 20 ml ampoule – 1% DV Aug-15 to 2017	16.30	5	Ropivacaine Kabi
ROPIVACAINE HYDROCHLORIDE WITH FENTANYL			
Inj 2 mg with fentanyl 2 mcg per ml, 100 ml bag	198.50	5	Naropin
Inj 2 mg with fentanyl 2 mcg per ml, 200 ml bag	270.00	5	Naropin
TETRACAINE [AMETHOCAINE] HYDROCHLORIDE			
Gel 4%			

Analgesics

Non-Opoid Analgesics

ASPIRIN

Tab dispersible 300 mg

CAPSAICIN – **Restricted** see terms below

⚡ Crm 0.075% 12.50 45 g Zostrix HP

➡**Restricted**

Initiation

For post-herpetic neuralgia or diabetic peripheral neuropathy.

METHOXYFLURANE – **Restricted** see terms below

⚡ Soln for inhalation 99.9%, 3 ml bottle

➡**Restricted**

Initiation

Both:

- 1 Patient is undergoing a painful procedure with an expected duration of less than one hour; and
- 2 Only to be used under supervision by a medical practitioner or nurse who is trained in the use of methoxyflurane.

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
NEFOPAM HYDROCHLORIDE			
Tab 30 mg			
PARACETAMOL – Some items restricted see terms below			
Tab soluble 500 mg – 1% DV Oct-15 to 2017	1.60	20	Paragesic Soluble
Tab 500 mg			
Oral liq 120 mg per 5 ml – 20% DV Oct-14 to 2017	4.15	1,000 ml	Paracare
Oral liq 250 mg per 5 ml – 20% DV Sep-14 to 2017	4.35	1,000 ml	Paracare Double Strength
¶ Inj 10 mg per ml, 50 ml vial – 1% DV Sep-14 to 2017	12.90	12	Perfalgan
¶ Inj 10 mg per ml, 100 ml vial – 1% DV Sep-14 to 2017	12.90	12	Perfalgan
Suppos 25 mg	56.35	20	Biomed
Suppos 50 mg	56.35	20	Biomed
Suppos 125 mg – 1% DV Dec-15 to 2018	3.69	10	Gacet
Suppos 250 mg – 1% DV Dec-15 to 2018	3.79	10	Gacet
Suppos 500 mg – 1% DV Nov-15 to 2018	12.60	50	Paracare
➔Restricted			
Initiation			
Intravenous paracetamol is only to be used where other routes are unavailable or impractical, or where there is reduced absorption.			
The need for IV paracetamol must be re-assessed every 24 hours.			
SUCROSE			
Oral liq 25%			
Opioid Analgesics			
ALFENTANIL			
Inj 0.5 mg per ml, 2 ml ampoule – 1% DV Jan-15 to 2017	39.07	10	Hameln
CODEINE PHOSPHATE			
Tab 15 mg – 1% DV Jul-13 to 2016	4.75	100	PSM
Tab 30 mg – 1% DV Jul-13 to 2016	5.80	100	PSM
Tab 60 mg – 1% DV Jul-13 to 2016	12.50	100	PSM
DIHYDROCODEINE TARTRATE			
Tab long-acting 60 mg – 1% DV Sep-13 to 2016	13.64	60	DHC Continus
FENTANYL			
Inj 10 mcg per ml, 10 ml syringe			
Inj 50 mcg per ml, 2 ml ampoule – 1% DV Sep-15 to 2018	3.95	10	Boucher and Muir
Inj 10 mcg per ml, 50 ml bag	210.00	10	Biomed
Inj 10 mcg per ml, 50 ml syringe	165.00	10	Biomed
Inj 50 mcg per ml, 10 ml ampoule – 1% DV Sep-15 to 2018	10.45	10	Boucher and Muir
Inj 10 mcg per ml, 100 ml bag	210.00	10	Biomed
Inj 20 mcg per ml, 50 ml syringe	185.00	10	Biomed
Inj 20 mcg per ml, 100 ml bag			
Patch 12.5 mcg per hour – 1% DV Aug-15 to 2016	2.92	5	Fentanyl Sandoz
Patch 25 mcg per hour – 1% DV Aug-15 to 2016	3.66	5	Fentanyl Sandoz
Patch 50 mcg per hour – 1% DV Aug-15 to 2016	6.64	5	Fentanyl Sandoz
Patch 75 mcg per hour – 1% DV Aug-15 to 2016	9.18	5	Fentanyl Sandoz
Patch 100 mcg per hour – 1% DV Aug-15 to 2016	11.29	5	Fentanyl Sandoz

↑ Item restricted (see ➔ above); ¶ Item restricted (see ➔ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
METHADONE HYDROCHLORIDE			
Tab 5 mg – 1% DV Sep-15 to 2018	1.85	10	Methatabs
Oral liq 2 mg per ml – 1% DV Sep-15 to 2018	5.55	200 ml	Biodone
Oral liq 5 mg per ml – 1% DV Sep-15 to 2018	5.00	200 ml	Biodone Forte
Oral liq 10 mg per ml – 1% DV Sep-15 to 2018	6.55	200 ml	Biodone Extra Forte
Inj 10 mg per ml, 1 ml vial	61.00	10	AFT
MORPHINE HYDROCHLORIDE			
Oral liq 1 mg per ml – 1% DV Oct-15 to 2018	8.84	200 ml	RA-Morph
Oral liq 2 mg per ml – 1% DV Oct-15 to 2018	14.00	200 ml	RA-Morph
Oral liq 5 mg per ml – 1% DV Oct-15 to 2018	18.00	200 ml	RA-Morph
Oral liq 10 mg per ml – 1% DV Oct-15 to 2018	26.00	200 ml	RA-Morph
MORPHINE SULPHATE			
Tab long-acting 10 mg – 1% DV Sep-13 to 2016	1.95	10	Arrow-Morphine LA
Tab immediate-release 10 mg – 1% DV Apr-15 to 2017	2.80	10	Sevredol
Tab immediate-release 20 mg – 1% DV Apr-15 to 2017	5.52	10	Sevredol
Tab long-acting 30 mg – 1% DV Sep-13 to 2016	2.98	10	Arrow-Morphine LA
Tab long-acting 60 mg – 1% DV Sep-13 to 2016	5.75	10	Arrow-Morphine LA
Tab long-acting 100 mg – 1% DV Sep-13 to 2016	6.45	10	Arrow-Morphine LA
Cap long-acting 10 mg – 1% DV Feb-14 to 2016	1.70	10	m-Eslon
Cap long-acting 30 mg – 1% DV Feb-14 to 2016	2.50	10	m-Eslon
Cap long-acting 60 mg – 1% DV Feb-14 to 2016	5.40	10	m-Eslon
Cap long-acting 100 mg – 1% DV Feb-14 to 2016	6.38	10	m-Eslon
Inj 1 mg per ml, 100 ml bag – 1% DV Oct-14 to 2017	185.00	10	Biomed
Inj 1 mg per ml, 10 ml syringe – 1% DV Oct-14 to 2017	45.00	10	Biomed
Inj 1 mg per ml, 50 ml syringe – 1% DV Oct-14 to 2017	87.50	10	Biomed
Inj 1 mg per ml, 2 ml syringe			
Inj 2 mg per ml, 30 ml syringe	135.00	10	Biomed
Inj 5 mg per ml, 1 ml ampoule – 1% DV Oct-14 to 2017	12.48	5	DBL Morphine Sulphate
Inj 10 mg per ml, 1 ml ampoule – 1% DV Oct-14 to 2017	9.09	5	DBL Morphine Sulphate
Inj 10 mg per ml, 100 mg cassette			
Inj 10 mg per ml, 100 ml bag			
Inj 15 mg per ml, 1 ml ampoule – 1% DV Oct-14 to 2017	9.77	5	DBL Morphine Sulphate
Inj 30 mg per ml, 1 ml ampoule – 1% DV Oct-14 to 2017	12.43	5	DBL Morphine Sulphate
Inj 200 mcg in 0.4 ml syringe			
Inj 300 mcg in 0.3 ml syringe			
MORPHINE TARTRATE			
Inj 80 mg per ml, 1.5 ml ampoule – 1% DV Sep-13 to 2016	35.60	5	Hospira
Inj 80 mg per ml, 5 ml ampoule – 1% DV Sep-13 to 2016	107.67	5	Hospira

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
OXYCODONE HYDROCHLORIDE			
Tab controlled-release 5 mg	7.51	20	OxyContin
Tab controlled-release 10 mg	6.75	20	Oxycodone ControlledRelease Tablets(BNM)
Tab controlled-release 20 mg	11.50	20	Oxycodone ControlledRelease Tablets(BNM)
Tab controlled-release 40 mg	18.50	20	Oxycodone ControlledRelease Tablets(BNM)
Tab controlled-release 80 mg	34.00	20	Oxycodone ControlledRelease Tablets(BNM)
Cap immediate-release 5 mg – 1% DV Oct-15 to 2018	1.98	20	OxyNorm
Cap immediate-release 10 mg – 1% DV Oct-15 to 2018	3.91	20	OxyNorm
Cap immediate-release 20 mg – 1% DV Oct-15 to 2018	6.84	20	OxyNorm
Oral liq 5 mg per 5 ml	11.20	250 ml	OxyNorm
Inj 1 mg per ml, 100 ml bag			
Inj 10 mg per ml, 1 ml ampoule – 1% DV Feb-16 to 2018	8.57	5	OxyNorm
Inj 10 mg per ml, 2 ml ampoule – 1% DV Feb-16 to 2018	16.89	5	OxyNorm
Inj 50 mg per ml, 1 ml ampoule – 1% DV Dec-15 to 2018	51.00	5	OxyNorm
PARACETAMOL WITH CODEINE			
Tab paracetamol 500 mg with codeine phosphate 8 mg	2.11	100	Paracetamol + Codeine (Relieve)
PETHIDINE HYDROCHLORIDE			
Tab 50 mg – 1% DV Nov-15 to 2018	4.46	10	PSM
Tab 100 mg – 1% DV Nov-15 to 2018	6.25	10	PSM
Inj 5 mg per ml, 10 ml syringe			
Inj 5 mg per ml, 100 ml bag			
Inj 10 mg per ml, 100 ml bag			
Inj 10 mg per ml, 50 ml syringe			
Inj 50 mg per ml, 1 ml ampoule – 1% DV Sep-14 to 2017	5.51	5	DBL Pethidine Hydrochloride
Inj 50 mg per ml, 2 ml ampoule – 1% DV Sep-14 to 2017	5.83	5	DBL Pethidine Hydrochloride
REMIFENTANIL HYDROCHLORIDE			
Inj 1 mg vial – 1% DV Nov-14 to 2017	10.00	5	Ultiva
Inj 2 mg vial – 1% DV Nov-14 to 2017	18.00	5	Ultiva
TRAMADOL HYDROCHLORIDE			
Tab sustained-release 100 mg – 1% DV Oct-14 to 2017	2.00	20	Tramal SR 100
Tab sustained-release 150 mg – 1% DV Oct-14 to 2017	3.00	20	Tramal SR 150
Tab sustained-release 200 mg – 1% DV Oct-14 to 2017	4.00	20	Tramal SR 200
Cap 50 mg – 1% DV Oct-14 to 2017	2.50	100	Arrow-Tramadol
Oral drops 100 mg per ml			
Inj 10 mg per ml, 100 ml bag			
Inj 50 mg per ml, 1 ml ampoule – 1% DV Oct-14 to 2017	4.50	5	Tramal 50
Inj 50 mg per ml, 2 ml ampoule – 1% DV Oct-14 to 2017	4.50	5	Tramal 100

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

Price (ex man. excl. GST)	Brand or Generic
\$	Per Manufacturer

Antidepressants

Cyclic and Related Agents

AMITRIPTYLINE

Tab 10 mg – 1% DV Sep-14 to 2017	1.68	100	Arrow-Amitriptyline
Tab 25 mg – 1% DV Jan-15 to 2017	1.68	100	Arrow-Amitriptyline
Tab 50 mg – 1% DV Jan-15 to 2017	2.82	100	Arrow-Amitriptyline

CLOMIPRAMINE HYDROCHLORIDE

Tab 10 mg – 1% DV Sep-15 to 2018	12.60	100	Apo-Clomipramine
Tab 25 mg – 1% DV Sep-15 to 2018	8.68	100	Apo-Clomipramine

DOTHIEPIN HYDROCHLORIDE

Tab 75 mg	10.50	100	Dopress
Cap 25 mg	6.17	100	Dopress

DOXEPIN HYDROCHLORIDE

Cap 10 mg
Cap 25 mg
Cap 50 mg

IMIPRAMINE HYDROCHLORIDE

Tab 10 mg	5.48	50	Tofranil
	6.58	60	Tofranil
Tab 25 mg	8.80	50	Tofranil

MAPROTILINE HYDROCHLORIDE

Tab 25 mg
Tab 75 mg

MIANSERIN HYDROCHLORIDE – **Restricted:** For continuation only

➔ Tab 30 mg

NORTRIPTYLINE HYDROCHLORIDE

Tab 10 mg – 1% DV Jun-13 to 2016	4.00	100	Norpress
Tab 25 mg – 1% DV Jun-13 to 2016	9.00	180	Norpress

Monoamine-Oxidase Inhibitors - Non-Selective

PHENELZINE SULPHATE

Tab 15 mg

TRANLYCYPROMINE SULPHATE

Tab 10 mg

Monoamine-Oxidase Type A Inhibitors

MOCLOBEMIDE

Tab 150 mg – 1% DV Oct-15 to 2018	85.10	500	Apo-Moclobemide
Tab 300 mg – 1% DV Oct-15 to 2018	30.70	100	Apo-Moclobemide

Other Antidepressants

MIRTAZAPINE

Tab 30 mg – 1% DV Nov-15 to 2018	2.55	30	Apo-Mirtazapine
Tab 45 mg – 1% DV Nov-15 to 2018	3.25	30	Apo-Mirtazapine

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
VENLAFAXINE – Some items restricted see terms below			
Tab modified release 37.5 mg	5.06	28	Arrow-Venlafaxine XR
Tab modified release 75 mg	6.44	28	Arrow-Venlafaxine XR
Tab modified release 150 mg	8.86	28	Arrow-Venlafaxine XR
Tab modified release 225 mg	14.34	28	Arrow-Venlafaxine XR
⚡ Cap modified release 37.5 mg	5.69	28	Efexor XR
⚡ Cap modified release 75 mg	11.40	28	Efexor XR
⚡ Cap modified release 150 mg	13.98	28	Efexor XR

➡ Restricted

Initiation

Re-assessment required after 2 years

Both:

- 1 The patient has 'treatment-resistant' depression; and
- 2 Either:
 - 2.1 The patient must have had a trial of two different antidepressants and have had an inadequate response from an adequate dose over an adequate period of time (usually at least four weeks); or
 - 2.2 Both:
 - 2.2.1 The patient is currently a hospital in-patient as a result of an acute depressive episode; and
 - 2.2.2 The patient must have had a trial of one other antidepressant and have had an inadequate response from an adequate dose over an adequate period of time.

Continuation

Re-assessment required after 2 years

The patient has a high risk of relapse (prescriber determined).

Selective Serotonin Reuptake Inhibitors

CITALOPRAM HYDROBROMIDE			
Tab 20 mg – 1% DV Jan-16 to 2018	1.79	84	PSM Citalopram
ESCITALOPRAM			
Tab 10 mg – 1% DV Jul-15 to 2016	1.40	28	Air Flow Products
Tab 20 mg – 1% DV Jul-15 to 2016	2.40	28	Air Flow Products
FLUOXETINE HYDROCHLORIDE			
Tab dispersible 20 mg, scored – 1% DV Apr-14 to 2016	2.50	30	Arrow-Fluoxetine
Cap 20 mg – 1% DV Apr-14 to 2016	1.74	90	Arrow-Fluoxetine
PAROXETINE HYDROCHLORIDE			
Tab 20 mg	4.32	90	Loxamine
SERTRALINE			
Tab 50 mg	3.64	90	Arrow-Sertraline
Tab 100 mg – 1% DV Sep-13 to 2016	6.28	90	Arrow-Sertraline

Antiepilepsy Drugs

Agents for the Control of Status Epilepticus

CLONAZEPAM			
Inj 1 mg per ml, 1 ml ampoule	19.00	5	Rivotril
DIAZEPAM			
Inj 5 mg per ml, 2 ml ampoule	11.83	5	Hospira
Rectal tubes 5 mg	25.05	5	Stesolid
Rectal tubes 10 mg	30.50	5	Stesolid

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LORAZEPAM			
Inj 2 mg vial			
Inj 4 mg per ml, 1 ml vial			
PARALDEHYDE			
Inj 5 ml ampoule			
PHENYTOIN SODIUM			
Inj 50 mg per ml, 2 ml ampoule – 1% DV Oct-15 to 2018	88.63	5	Hospira
Inj 50 mg per ml, 5 ml ampoule – 1% DV Oct-15 to 2018	133.92	5	Hospira

Control of Epilepsy

CARBAMAZEPINE			
Tab 200 mg	14.53	100	Tegretol
Tab long-acting 200 mg	16.98	100	Tegretol CR
Tab 400 mg	34.58	100	Tegretol
Tab long-acting 400 mg	39.17	100	Tegretol CR
Oral liq 20 mg per ml	26.37	250 ml	Tegretol
CLOBAZAM			
Tab 10 mg			
CLONAZEPAM			
Oral drops 2.5 mg per ml			
ETHOSUXIMIDE			
Cap 250 mg			
Oral liq 50 mg per ml			
GABAPENTIN – Restricted see terms below			
⚡ Cap 100 mg	7.16	100	Arrow-Gabapentin Neurontin Nupentin
⚡ Cap 300 mg	11.00	100	Arrow-Gabapentin Neurontin Nupentin
⚡ Cap 400 mg	13.75	100	Arrow-Gabapentin Neurontin Nupentin

➡ Restricted

Initiation — preoperative and/or postoperative use

Limited to 8 days treatment

Initiation — pain management of burns patients

Re-assessment required after 1 month

Continuation — pain management of burns patients

Re-assessment required after 1 month

The treatment remains appropriate and the patient is benefiting from treatment.

Initiation — epilepsy

Re-assessment required after 15 months

Either:

- Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
- Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Note: "Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Continuation — epilepsy

Patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life.

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective

Initiation — Neuropathic pain or Chronic Kidney Disease-associated pruritus

Re-assessment required after 3 months

Either:

- 1 The patient has been diagnosed with neuropathic pain; or
- 2 Both:
 - 2.1 The patient has Chronic Kidney Disease Stage 5-associated pruritus* where no other cause for pruritus can be identified (e.g. scabies, allergy); and
 - 2.2 The patient has persistent pruritus not relieved with a trial of emollient/moisturising creams alone.

Continuation — Neuropathic pain or Chronic Kidney Disease-associated pruritus

Either:

- 1 The patient has demonstrated a marked improvement in their control of pain or itch (prescriber determined); or
- 2 The patient has previously demonstrated clinical responsiveness to gabapentin and has now developed neuropathic pain in a new site.

Note: Indications marked with * are Unapproved Indications. Dosage adjustment of gabapentin is recommended for patients with renal impairment.

LACOSAMIDE – **Restricted** see terms below

⚡ Tab 50 mg	25.04	14	Vimpat
⚡ Tab 100 mg	50.06	14	Vimpat
	200.24	56	Vimpat
⚡ Tab 150 mg	75.10	14	Vimpat
	300.40	56	Vimpat
⚡ Tab 200 mg	400.55	56	Vimpat
⚡ Inj 10 mg per ml, 20 ml vial			

➡ **Restricted**

Initiation

Re-assessment required after 15 months

Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Continuation

Patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LAMOTRIGINE			
Tab dispersible 2 mg	6.74	30	Lamictal
Tab dispersible 5 mg	15.00	56	Arrow-Lamotrigine
	9.64	30	Lamictal
Tab dispersible 25 mg	20.40	56	Arrow-Lamotrigine
	29.09		Lamictal
	19.38		Logem
Tab dispersible 50 mg	34.70	56	Arrow-Lamotrigine
	47.89		Lamictal
	32.97		Logem
Tab dispersible 100 mg	59.90	56	Arrow-Lamotrigine
	79.16		Lamictal
	56.91		Logem
LEVETIRACETAM			
Tab 250 mg	24.03	60	Everet
			Levetiracetam-Rex
Tab 500 mg	28.71	60	Everet
			Levetiracetam-Rex
Tab 750 mg	45.23	60	Everet
			Levetiracetam-Rex
Tab 1,000 mg	59.12	60	Everet
Inj 100 mg per ml, 5 ml vial			
<i>(Levetiracetam-Rex Tab 250 mg to be delisted 1 August 2016)</i>			
<i>(Levetiracetam-Rex Tab 500 mg to be delisted 1 August 2016)</i>			
<i>(Levetiracetam-Rex Tab 750 mg to be delisted 1 August 2016)</i>			
PHENOBARBITONE			
Tab 15 mg – 1% DV Dec-15 to 2018	30.00	500	PSM
Tab 30 mg – 1% DV Dec-15 to 2018	31.00	500	PSM
PHENYTOIN			
Tab 50 mg			
PHENYTOIN SODIUM			
Cap 30 mg			
Cap 100 mg			
Oral liq 6 mg per ml			
PRIMIDONE			
Tab 250 mg			
SODIUM VALPROATE			
Tab 100 mg			
Tab EC 200 mg			
Tab EC 500 mg			
Oral liq 40 mg per ml			
Inj 100 mg per ml, 4 ml vial – 1% DV Sep-15 to 2018	16.60	1	Epilim IV
STIRIPENTOL – Restricted see terms on the next page			
⚡ Cap 250 mg	509.29	60	Diacomit
⚡ Powder for oral liq 250 mg sachet	509.29	60	Diacomit

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➡Restricted			
Initiation			
Paediatric neurologist			
<i>Re-assessment required after 6 months</i>			
Both:			
1 Patient has confirmed diagnosis of Dravet syndrome; and			
2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.			
Continuation			
Paediatric neurologist			
Patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.			
TOPIRAMATE			
Tab 25 mg	11.07	60	Arrow-Topiramate
	26.04		Topamax
	11.07		Topiramate Actavis
Tab 50 mg	18.81	60	Arrow-Topiramate
	44.26		Topamax
	18.81		Topiramate Actavis
Tab 100 mg	31.99	60	Arrow-Topiramate
	75.25		Topamax
	31.99		Topiramate Actavis
Tab 200 mg	55.19	60	Arrow-Topiramate
	129.85		Topamax
	55.19		Topiramate Actavis
Cap sprinkle 15 mg	20.84	60	Topamax
Cap sprinkle 25 mg	26.04	60	Topamax
VIGABATRIN – Restricted see terms below			
⚡ Tab 500 mg			
➡Restricted			
Initiation			
<i>Re-assessment required after 15 months</i>			
Both:			
1 Either:			
1.1 Patient has infantile spasms; or			
1.2 Both:			
1.2.1 Patient has epilepsy; and			
1.2.2 Either:			
1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or			
1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and			
2 Either:			
2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or			
2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.			
Notes: "Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.			
Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.			
Continuation			
Both:			
			continued...
⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below) e.g. Brand indicates brand example only. It is not a contracted product.			

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

continued...

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Antimigraine Preparations

Acute Migraine Treatment

DIHYDROERGOTAMINE MESYLATE

Inj 1 mg per ml, 1 ml ampoule

ERGOTAMINE TARTRATE WITH CAFFEINE

Tab 1 mg with caffeine 100 mg

METOCLOPRAMIDE HYDROCHLORIDE WITH PARACETAMOL

Tab 5 mg with paracetamol 500 mg

RIZATRIPTAN

Tab orodispersible 10 mg – 1% DV Sep-14 to 2017	3.24	12	Rizamelt
	8.10	30	Rizamelt

SUMATRIPTAN

Tab 50 mg – 1% DV Sep-13 to 2016	29.80	100	Arrow-Sumatriptan
Tab 100 mg – 1% DV Sep-13 to 2016	54.80	100	Arrow-Sumatriptan
Inj 12 mg per ml, 0.5 ml cartridge	13.80	2	Arrow-Sumatriptan

Prophylaxis of Migraine

PIZOTIFEN

Tab 500 mcg – 1% DV Sep-15 to 2018	23.21	100	Sandomigran
--	-------	-----	--------------------

Antinausea and Vertigo Agents

APREPITANT – **Restricted** see terms below

☒ Cap 2 × 80 mg and 1 × 125 mg – 1% DV Sep-14 to 2017	100.00	3	Emend Tri-Pack
---	--------	---	-----------------------

☒ **Restricted**

Initiation

Patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

Tab 16 mg – 1% DV Jun-14 to 2017	4.95	84	Vergo 16
--	------	----	-----------------

CYCLIZINE HYDROCHLORIDE

Tab 50 mg – 1% DV Jan-16 to 2018	0.59	20	Nauzene
--	------	----	----------------

CYCLIZINE LACTATE

Inj 50 mg per ml, 1 ml ampoule	14.95	5	Nausicalm
--------------------------------------	-------	---	------------------

DOMPERIDONE

Tab 10 mg – 1% DV Dec-15 to 2018	3.20	100	Prokinex
--	------	-----	-----------------

DROPERIDOL

Inj 2.5 mg per ml, 1 ml ampoule

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GRANISETRON			
Tab 1 mg – 1% DV Jan-15 to 2017	5.98	50	Granirex
HYOSCINE HYDROBROMIDE			
Inj 400 mcg per ml, 1 ml ampoule	46.50	5	Hospira
¶ Patch 1.5 mg – 1% DV Dec-13 to 2016	11.95	2	Scopoderm TTS
➡ Restricted			
Initiation			
Any of the following:			
1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or			
2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective; or			
3 For treatment of post-operative nausea and vomiting where cyclizine, droperidol and a 5HT3 antagonist have proven ineffective, are not tolerated or are contraindicated.			
METOCLOPRAMIDE HYDROCHLORIDE			
Tab 10 mg – 1% DV Sep-14 to 2017	1.82	100	Metamide
Oral liq 5 mg per 5 ml			
Inj 5 mg per ml, 2 ml ampoule – 1% DV Sep-14 to 2017	4.50	10	Pfizer
ONDANSETRON			
Tab 4 mg – 1% DV Jan-14 to 2016	5.51	50	Onrex
Tab dispersible 4 mg – 1% DV Oct-14 to 2017	1.00	10	Dr Reddy's ondansetron
Tab 8 mg – 1% DV Jan-14 to 2016	6.19	50	Onrex
Tab dispersible 8 mg – 1% DV Oct-14 to 2017	1.50	10	Ondansetron ODT-DRLA
Inj 2 mg per ml, 2 ml ampoule – 1% DV Sep-13 to 2016	1.82	5	Ondanaccord
Inj 2 mg per ml, 4 ml ampoule – 1% DV Sep-13 to 2016	2.18	5	Ondanaccord
PROCHLORPERAZINE			
Tab buccal 3 mg			
Tab 5 mg – 1% DV Jun-14 to 2017	9.75	500	Antinaus
Inj 12.5 mg per ml, 1 ml ampoule			
Suppos 25 mg			
PROMETHAZINE THEOCLATE – Restricted: For continuation only			
➡ Tab 25 mg			
TROPISETRON			
Inj 1 mg per ml, 2 ml ampoule – 1% DV Sep-15 to 2018	8.95	1	Tropisetron-AFT
Inj 1 mg per ml, 5 ml ampoule – 1% DV Sep-15 to 2018	13.95	1	Tropisetron-AFT

Antipsychotic Agents

General

AMISULPRIDE			
Tab 100 mg – 1% DV Jul-13 to 2016	6.22	30	Solian
Tab 200 mg – 1% DV Jul-13 to 2016	21.92	60	Solian
Tab 400 mg – 1% DV Jul-13 to 2016	44.52	60	Solian
Oral liq 100 mg per ml – 1% DV Jul-13 to 2016	52.50	60 ml	Solian

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ARIPIPRAZOLE – Restricted see terms below			
⚡ Tab 5 mg	123.54	30	Abilify
⚡ Tab 10 mg	123.54	30	Abilify
⚡ Tab 15 mg	175.28	30	Abilify
⚡ Tab 20 mg	213.42	30	Abilify
⚡ Tab 30 mg	260.07	30	Abilify

➡ **Restricted**

Initiation — schizophrenia or related psychoses

Any specialist

Both:

- 1 Patient is suffering from schizophrenia or related psychoses; and
- 2 Either:
 - 2.1 An effective dose of risperidone or quetiapine has been trialled and has been discontinued, or is in the process of being discontinued, because of unacceptable side effect; or
 - 2.2 An effective dose of risperidone or quetiapine has been trialled and has been discontinued, or is in the process of being discontinued, because of inadequate clinical response.

Initiation — Autism spectrum disorder*

Psychiatrist or paediatrician

All of the following:

- 1 The patient has been diagnosed with an autism spectrum disorder* and has symptoms of severe irritability; and
- 2 An effective dose of risperidone has been trialled and has been discontinued because of unacceptable side effects or inadequate response; and
- 3 The patient is aged less than 18 years.

Note: Indications marked with * are Unapproved Indications

CHLORPROMAZINE HYDROCHLORIDE

Tab 10 mg
Tab 25 mg
Tab 100 mg
Oral liq 10 mg per ml
Inj 25 mg per ml, 2 ml ampoule

CLOZAPINE

Tab 25 mg	6.69	50	Clopine
	13.37	100	Clopine
	5.69	50	Clozaril
	11.36	100	Clozaril
Tab 50 mg	8.67	50	Clopine
	17.33	100	Clopine
Tab 100 mg	17.33	50	Clopine
	34.65	100	Clopine
	14.73	50	Clozaril
	29.45	100	Clozaril
Tab 200 mg	34.65	50	Clopine
	69.30	100	Clopine
Oral liq 50 mg per ml	17.33	100 ml	Clopine

HALOPERIDOL

Tab 500 mcg – 1% DV Oct-13 to 2016	6.23	100	Serenace
Tab 1.5 mg – 1% DV Oct-13 to 2016	9.43	100	Serenace
Tab 5 mg – 1% DV Oct-13 to 2016	29.72	100	Serenace
Oral liq 2 mg per ml – 1% DV Oct-13 to 2016	23.84	100 ml	Serenace
Inj 5 mg per ml, 1ml ampoule – 1% DV Oct-13 to 2016	21.55	10	Serenace

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LEVOMEPRMAZINE			
Tab 25 mg			
Tab 100 mg			
Inj 25 mg per ml, 1 ml ampoule			
LITHIUM CARBONATE			
Tab long-acting 400 mg			
Tab 250 mg – 1% DV Sep-15 to 2018	34.30	500	Lithicarb FC
Tab 400 mg – 1% DV Sep-15 to 2018	12.83	100	Lithicarb FC
Cap 250 mg – 1% DV Sep-14 to 2017	9.42	100	Douglas
OLANZAPINE			
Tab 2.5 mg – 1% DV Sep-14 to 2017	0.75	28	Zypine
Tab 5 mg – 1% DV Sep-14 to 2017	1.65	28	Zypine
Tab orodispersible 5 mg – 1% DV Sep-14 to 2017	1.75	28	Zypine ODT
Tab 10 mg – 1% DV Sep-14 to 2017	2.55	28	Zypine
Tab orodispersible 10 mg – 1% DV Sep-14 to 2017	3.05	28	Zypine ODT
Inj 10 mg vial			
PERICYAZINE			
Tab 2.5 mg			
Tab 10 mg			
QUETIAPINE			
Tab 25 mg – 1% DV Sep-14 to 2017	2.10	90	Quetapel
Tab 100 mg – 1% DV Sep-14 to 2017	4.20	90	Quetapel
Tab 200 mg – 1% DV Sep-14 to 2017	7.20	90	Quetapel
Tab 300 mg – 1% DV Sep-14 to 2017	12.00	90	Quetapel
RISPERIDONE – Some items restricted see terms below			
Tab 0.5 mg – 1% DV Feb-15 to 2017	1.90	60	Actavis
⚡ Tab orodispersible 0.5 mg	21.42	28	Risperdal Quicklet
Tab 1 mg – 1% DV Feb-15 to 30 Sep 2017	2.10	60	Actavis
⚡ Tab orodispersible 1 mg	42.84	28	Risperdal Quicklet
Tab 2 mg – 1% DV Feb-15 to 2017	2.34	60	Actavis
⚡ Tab orodispersible 2 mg	85.71	28	Risperdal Quicklet
Tab 3 mg – 1% DV Feb-15 to 2017	2.55	60	Actavis
Tab 4 mg – 1% DV Feb-15 to 2017	3.50	60	Actavis
Oral liq 1 mg per ml – 1% DV Sep-14 to 2017	9.75	30 ml	Rispon
➡ Restricted			
Initiation — Acute situations			
Both:			
1 For a non-adherent patient on oral therapy with standard risperidone tablets or risperidone oral liquid; and			
2 The patient is under direct supervision for administration of medicine.			
Initiation — Chronic situations			
Both:			
1 The patient is unable to take standard risperidone tablets or oral liquid, or once stabilised refuses to take risperidone tablets or oral liquid; and			
2 The patient is under direct supervision for administration of medicine.			
TRIFLUOPERAZINE HYDROCHLORIDE			
Tab 1 mg			
Tab 2 mg			
Tab 5 mg			

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ZIPRASIDONE			
Cap 20 mg – 1% DV Jan-16 to 2018	14.56	60	Zusdone
Cap 40 mg – 1% DV Jan-16 to 2018	24.75	60	Zusdone
Cap 60 mg – 1% DV Jan-16 to 2018	33.87	60	Zusdone
Cap 80 mg – 1% DV Jan-16 to 2018	39.74	60	Zusdone
ZUCLOPENTHIXOL ACETATE			
Inj 50 mg per ml, 1 ml ampoule			
Inj 50 mg per ml, 2 ml ampoule			
ZUCLOPENTHIXOL HYDROCHLORIDE			
Tab 10 mg	31.45	100	Clopixol
Depot Injections			
FLUPENTHIXOL DECANOATE			
Inj 20 mg per ml, 1 ml ampoule	13.14	5	Fluanxol
Inj 20 mg per ml, 2 ml ampoule	20.90	5	Fluanxol
Inj 100 mg per ml, 1 ml ampoule	40.87	5	Fluanxol
FLUPHENAZINE DECANOATE			
Inj 12.5 mg per 0.5 ml ampoule	17.60	5	Modecate
Inj 25 mg per ml, 1 ml ampoule	27.90	5	Modecate
Inj 25 mg per ml, 2 ml ampoule			<i>e.g. Modecate</i>
Inj 100 mg per ml, 1 ml ampoule	154.50	5	Modecate
HALOPERIDOL DECANOATE			
Inj 50 mg per ml, 1 ml ampoule	28.39	5	Haldol
Inj 100 mg per ml, 1 ml ampoule	55.90	5	Haldol Concentrate
OLANZAPINE – Restricted see terms below			
⚡ Inj 210 mg vial	280.00	1	Zyprexa Relprev
⚡ Inj 300 mg vial	460.00	1	Zyprexa Relprev
⚡ Inj 405 mg vial	560.00	1	Zyprexa Relprev
➡ Restricted			
Initiation			
<i>Re-assessment required after 12 months</i>			
Either:			
1 The patient has had an initial Special Authority approval for risperidone depot injection or paliperidone depot injection; or			
2 All of the following:			
2.1 The patient has schizophrenia; and			
2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and			
2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.			
Continuation			
<i>Re-assessment required after 12 months</i>			
The initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.			
PALIPERIDONE – Restricted see terms on the next page			
⚡ Inj 25 mg syringe	194.25	1	Invega Sustenna
⚡ Inj 50 mg syringe	271.95	1	Invega Sustenna
⚡ Inj 75 mg syringe	357.42	1	Invega Sustenna
⚡ Inj 100 mg syringe	435.12	1	Invega Sustenna
⚡ Inj 150 mg syringe	435.12	1	Invega Sustenna

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔Restricted

Initiation

Re-assessment required after 12 months

Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Continuation

Re-assessment required after 12 months

The initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PIPOTHIAZINE PALMITATE – **Restricted:** For continuation only

- ➔ Inj 50 mg per ml, 1 ml ampoule
- ➔ Inj 50 mg per ml, 2 ml ampoule

RISPERIDONE – **Restricted** see terms below

⚡ Inj 25 mg vial	135.98	1	Risperdal Consta
⚡ Inj 37.5 mg vial	178.71	1	Risperdal Consta
⚡ Inj 50 mg vial	217.56	1	Risperdal Consta

➔Restricted

Initiation

Re-assessment required after 12 months

Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Continuation

Re-assessment required after 12 months

The initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE

Inj 200 mg per ml, 1 ml ampoule	19.80	5	Clopixol
Inj 500 mg per ml, 1 ml ampoule			e.g. Clopixol Conc

Anxiolytics

ALPRAZOLAM

- Tab 1 mg
- Tab 250 mcg
- Tab 500 mcg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BUSPIRONE HYDROCHLORIDE			
Tab 5 mg – 1% DV Jul-16 to 2018	23.80	100	Orion
	28.00		Pacific Buspirone
Tab 10 mg – 1% DV Jul-16 to 2018	14.96	100	Orion
	17.00		Pacific Buspirone
<i>(Pacific Buspirone Tab 5 mg to be delisted 1 July 2016)</i>			
<i>(Pacific Buspirone Tab 10 mg to be delisted 1 July 2016)</i>			
CLONAZEPAM			
Tab 500 mcg	7.53	100	Paxam
Tab 2 mg	14.37	100	Paxam
DIAZEPAM			
Tab 2 mg	11.44	500	Arrow-Diazepam
Tab 5 mg	13.71	500	Arrow-Diazepam
LORAZEPAM			
Tab 1 mg – 1% DV Jun-15 to 2018	10.79	250	Ativan
Tab 2.5 mg – 1% DV Jun-15 to 2018	13.88	100	Ativan
OXAZEPAM			
Tab 10 mg – 1% DV Dec-14 to 2017	6.17	100	Ox-Pam
Tab 15 mg – 1% DV Dec-14 to 2017	8.53	100	Ox-Pam

Multiple Sclerosis Treatments

DIMETHYL FUMARATE – Restricted see terms below

⚡ Cap 120 mg	520.00	14	Tecfidera
⚡ Cap 240 mg	2,000.00	56	Tecfidera

➡ **Restricted**

Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

FINGOLIMOD – Restricted see terms below

⚡ Cap 0.5 mg	2,650.00	28	Gilenya
--------------------	----------	----	---------

➡ **Restricted**

Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

NATALIZUMAB – Restricted see terms below

⚡ Inj 20 mg per ml, 15 ml vial	1,750.00	1	Tysabri
--------------------------------------	----------	---	---------

➡ **Restricted**

Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

TERIFLUNOMIDE – Restricted see terms on the next page

⚡ Tab 14 mg	1,582.62	28	Aubagio
-------------------	----------	----	---------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔Restricted

Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

Other Multiple Sclerosis Treatments

➔Restricted

Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

GLATIRAMER ACETATE – **Restricted** see terms above

⬆ Inj 20 mg per ml, 1 ml syringe

INTERFERON BETA-1-ALPHA – **Restricted** see terms above

⬆ Inj 6 million iu in 0.5 ml pen injector	1,170.00	4	Avonex Pen
⬆ Inj 6 million iu in 0.5 ml syringe	1,170.00	4	Avonex
⬆ Inj 6 million iu vial	1,170.00	4	Avonex

INTERFERON BETA-1-BETA – **Restricted** see terms above

⬆ Inj 8 million iu per ml, 1 ml vial

Sedatives and Hypnotics

CHLORAL HYDRATE

- Oral liq 100 mg per ml
- Oral liq 200 mg per ml

LORMETAZEPAM – **Restricted**: For continuation only

➔ Tab 1 mg

MELATONIN – **Restricted** see terms below

⬆ Tab modified-release 2 mg			<i>e.g. Circadin</i>
⬆ Tab 1 mg			
⬆ Tab 2 mg			
⬆ Tab 3 mg			
⬆ Cap 2 mg			
⬆ Cap 3 mg			

➔Restricted

Initiation

For in hospital use only. For the treatment of insomnia where benzodiazepines and zopiclone are contraindicated.

MIDAZOLAM

Tab 7.5 mg	40.00	100	Hypnovel
Oral liq 2 mg per ml			
Inj 1 mg per ml, 5 ml ampoule	10.75	10	Hypnovel
	10.00		Pfizer
Inj 5 mg per ml, 3 ml ampoule	11.90	5	Hypnovel
			Pfizer

NITRAZEPAM

Tab 5 mg – 1% DV Dec-14 to 2017	5.22	100	Nitrados
---------------------------------------	------	-----	----------

PHENOBARBITONE

Inj 200 mg per ml, 1 ml ampoule

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
TEMAZEPAM			
Tab 10 mg – 1% DV Sep-14 to 2017	1.27	25	Normison
TRIAZOLAM – Restricted: For continuation only			
➔ Tab 125 mcg			
➔ Tab 250 mcg			
ZOPICLONE			
Tab 7.5 mg – 1% DV Dec-15 to 2018	0.98	30	Zopiclone Actavis
	8.99	500	Zopiclone Actavis

Stimulants / ADHD Treatments

ATOMOXETINE – Restricted see terms below

⚡ Cap 10 mg	107.03	28	Strattera
⚡ Cap 18 mg	107.03	28	Strattera
⚡ Cap 25 mg	107.03	28	Strattera
⚡ Cap 40 mg	107.03	28	Strattera
⚡ Cap 60 mg	107.03	28	Strattera
⚡ Cap 80 mg	139.11	28	Strattera
⚡ Cap 100 mg	139.11	28	Strattera

➔ **Restricted**

Initiation

All of the following:

- 1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder) diagnosed according to DSM-IV or ICD 10 criteria; and
- 2 Once-daily dosing; and
- 3 Any of the following:
 - 3.1 Treatment with a subsidised formulation of a stimulant has resulted in the development or worsening of serious adverse reactions or where the combination of subsidised stimulant treatment with another agent would pose an unacceptable medical risk; or
 - 3.2 Treatment with a subsidised formulation of a stimulant has resulted in worsening of co-morbid substance abuse or there is a significant risk of diversion with subsidised stimulant therapy; or
 - 3.3 An effective dose of a subsidised formulation of a stimulant has been trialled and has been discontinued because of inadequate clinical response; or
 - 3.4 Treatment with a subsidised formulation of a stimulant is considered inappropriate because the patient has a history of psychoses or has a first-degree relative with schizophrenia; and
- 4 The patient will not be receiving treatment with atomoxetine in combination with a subsidised formulation of a stimulant, except for the purposes of transitioning from subsidised stimulant therapy to atomoxetine.

Note: A "subsidised formulation of a stimulant" refers to currently listed methylphenidate hydrochloride tablet formulations (immediate-release, sustained-release and extended-release) or dexamphetamine sulphate tablets.

CAFFEINE

Tab 100 mg

DEXAMFETAMINE SULFATE – Restricted see terms below

⚡ Tab 5 mg – 1% DV Dec-15 to 2018	17.00	100	PSM
---	-------	-----	------------

➔ **Restricted**

Initiation — ADHD

Paediatrician or psychiatrist

Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria.

Initiation — Narcolepsy

Neurologist or respiratory specialist

Re-assessment required after 24 months

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Patient suffers from narcolepsy.

Continuation — Narcolepsy

Neurologist or respiratory specialist

Re-assessment required after 24 months

The treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE – **Restricted** see terms below

⚡ Tab extended-release 18 mg	58.96	30	Concerta
⚡ Tab extended-release 27 mg	65.44	30	Concerta
⚡ Tab extended-release 36 mg	71.93	30	Concerta
⚡ Tab extended-release 54 mg	86.24	30	Concerta
⚡ Tab immediate-release 5 mg	3.20	30	Rubifen
⚡ Tab immediate-release 10 mg	3.00	30	Ritalin
			Rubifen
⚡ Tab immediate-release 20 mg	7.85	30	Rubifen
⚡ Tab sustained-release 20 mg	50.00	100	Ritalin SR
	10.95	30	Rubifen SR
⚡ Cap modified-release 10 mg	15.60	30	Ritalin LA
⚡ Cap modified-release 20 mg	20.40	30	Ritalin LA
⚡ Cap modified-release 30 mg	25.52	30	Ritalin LA
⚡ Cap modified-release 40 mg	30.60	30	Ritalin LA

➡ **Restricted**

Initiation — ADHD (immediate-release and sustained-release formulations)

Paediatrician or psychiatrist

Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria.

Initiation — Narcolepsy (immediate-release and sustained-release formulations)

Neurologist or respiratory specialist

Re-assessment required after 24 months

Patient suffers from narcolepsy.

Continuation — Narcolepsy (immediate-release and sustained-release formulations)

Neurologist or respiratory specialist

Re-assessment required after 24 months

The treatment remains appropriate and the patient is benefiting from treatment.

Initiation — Extended-release and modified-release formulations

Paediatrician or psychiatrist

Both:

- 1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria; and
- 2 Either:
 - 2.1 Patient is taking a currently listed formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 2.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

MODAFINIL – **Restricted** see terms on the next page

⚡ Tab 100 mg

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

➔Restricted

Initiation — Narcolepsy

Neurologist or respiratory specialist

Re-assessment required after 24 months

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a listed formulation of methylphenidate or dexamphetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamphetamine are contraindicated.

Continuation — Narcolepsy

Neurologist or respiratory specialist

Re-assessment required after 24 months

The treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE

Tab 5 mg – 1% DV Feb-15 to 2017	5.48	90	Donepezil-Rex
Tab 10 mg – 1% DV Feb-15 to 2017	10.51	90	Donepezil-Rex

RIVASTIGMINE – Restricted see terms below

⚡ Patch 4.6 mg per 24 hour	90.00	30	Exelon
⚡ Patch 9.5 mg per 24 hour	90.00	30	Exelon

➔Restricted

Initiation

Re-assessment required after 6 months

Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Continuation

Re-assessment required after 12 months

Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE – Restricted see terms on the next page

⚡ Tab 2 mg with naloxone 0.5 mg	57.40	28	Suboxone
⚡ Tab 8 mg with naloxone 2 mg	166.00	28	Suboxone

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➡Restricted			
Initiation — Detoxification			
All of the following:			
1 Patient is opioid dependent; and			
2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and			
3 Prescriber works in an opioid treatment service approved by the Ministry of Health.			
Initiation — Maintenance treatment			
All of the following:			
1 Patient is opioid dependent; and			
2 Patient will not be receiving methadone; and			
3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and			
4 Prescriber works in an opioid treatment service approved by the Ministry of Health.			
BUPROPION HYDROCHLORIDE			
Tab modified-release 150 mg – 1% DV Oct-13 to 2016	4.97	30	Zyban
DISULFIRAM			
Tab 200 mg	24.30	100	Antabuse
NALTREXONE HYDROCHLORIDE – Restricted see terms below			
⚡ Tab 50 mg – 1% DV Sep-13 to 2016	76.00	30	Nalttraccod
➡Restricted			
Initiation — Alcohol dependence			
Both:			
1 Patient is currently enrolled, or is planned to be enrolled, in a recognised comprehensive treatment programme for alcohol dependence; and			
2 Naltrexone is to be prescribed by, or on the recommendation of, a physician working in an Alcohol and Drug Service.			
Initiation — Constipation			
For the treatment of opioid-induced constipation.			
NICOTINE – Some items restricted see terms below			
Patch 7 mg per 24 hours – 1% DV Apr-14 to 2017	10.57	28	Habitrol
Patch 14 mg per 24 hours – 1% DV Apr-14 to 2017	11.31	28	Habitrol
Patch 21 mg per 24 hours – 1% DV Apr-14 to 2017	11.95	28	Habitrol
⚡ Oral spray 1 mg per dose			<i>e.g. Nicorette QuickMist Mouth Spray</i>
Lozenge 1 mg – 1% DV Apr-14 to 2017	12.91	216	Habitrol
Lozenge 2 mg – 1% DV Apr-14 to 2017	14.14	216	Habitrol
⚡ Soln for inhalation 15 mg cartridge			<i>e.g. Nicorette Inhalator</i>
Gum 2 mg – 1% DV Apr-14 to 2017	22.26	384	Habitrol (Classic) Habitrol (Fruit) Habitrol (Mint)
Gum 4 mg – 1% DV Apr-14 to 2017	25.67	384	Habitrol (Classic) Habitrol (Fruit) Habitrol (Mint)

➡Restricted

Initiation

Any of the following:

- 1 For perioperative use in patients who have a 'nil by mouth' instruction; or
- 2 For use within mental health inpatient units; or
- 3 For acute use in agitated patients who are unable to leave the hospital facilities.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
VARENICLINE – Restricted see terms below			
⚡ Tab 0.5 mg × 11 and 1 mg × 14	60.48	25	Champix
⚡ Tab 1 mg	67.74	28	Champix
	135.48	56	Champix

➡ **Restricted**

Initiation

All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not used funded varenicline in the last 12 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline in a 12 month period.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Chemotherapeutic Agents			
Alkylating Agents			
BUSULFAN			
Tab 2 mg	89.25	100	Myleran
Inj 6 mg per ml, 10 ml ampoule			
CARMUSTINE			
Inj 100 mg vial – 1% DV Sep-15 to 2018	532.00	1	BiCNU
CHLORAMBUCIL			
Tab 2 mg			
CYCLOPHOSPHAMIDE			
Tab 50 mg	79.00	50	Endoxan
	158.00	100	Procytox
Inj 1 g vial – 1% DV Oct-15 to 2018	35.03	1	Endoxan
Inj 2 g vial – 1% DV Oct-15 to 2018	70.06	1	Endoxan
IFOSFAMIDE			
Inj 1 g vial	96.00	1	Holoxan
Inj 2 g vial	180.00	1	Holoxan
LOMUSTINE			
Cap 10 mg	132.59	20	Ceenu
Cap 40 mg	399.15	20	Ceenu
MELPHALAN			
Tab 2 mg			
Inj 50 mg vial			
THIOTEPA			
Inj 15 mg vial			
Inj 100 mg vial			
Anthracyclines and Other Cytotoxic Antibiotics			
BLEOMYCIN SULPHATE			
Inj 15,000 iu vial – 1% DV Oct-15 to 2018	150.48	1	DBL Bleomycin Sulfate
DACTINOMYCIN [ACTINOMYCIN D]			
Inj 0.5 mg vial	145.00	1	Cosmegen
DAUNORUBICIN			
Inj 2 mg per ml, 10 ml vial – 1% DV Aug-13 to 2016	118.72	1	Pfizer
DOXORUBICIN HYDROCHLORIDE			
Inj 2 mg per ml, 5 ml vial			
Inj 2 mg per ml, 25 ml vial – 1% DV Feb-16 to 2018	11.50	1	Doxorubicin Ebewe
Note: DV limit applies to all 50 mg presentations of doxorubicin hydrochloride.			
Inj 50 mg vial			
Inj 2 mg per ml, 50 ml vial – 1% DV Feb-16 to 2018	23.00	1	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial – 1% DV Feb-16 to 2018	46.00	1	Doxorubicin Ebewe

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
EPIRUBICIN HYDROCHLORIDE			
Inj 2 mg per ml, 5 ml vial	25.00	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial – 1% DV Nov-15 to 2018	30.00	1	Epirubicin Ebewe
Inj 2 mg per ml, 50 ml vial – 1% DV Nov-15 to 2018	32.50	1	Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial – 1% DV Nov-15 to 2018	65.00	1	Epirubicin Ebewe
IDARUBICIN HYDROCHLORIDE			
Inj 5 mg vial – 1% DV Nov-15 to 2018	125.00	1	Zavedos
Inj 10 mg vial – 1% DV Nov-15 to 2018	250.00	1	Zavedos
MITOMYCIN C			
Inj 5 mg vial – 1% DV Oct-13 to 2016	79.75	1	Arrow
MITOZANTRONE			
Inj 2 mg per ml, 10 ml vial – 1% DV Sep-15 to 2018	97.50	1	Mitozantrone Ebewe
Antimetabolites			
AZACITIDINE – Restricted see terms below			
☞ Inj 100 mg vial	605.00	1	Vidaza
☞ Restricted			
Initiation			
Haematologist			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 Any of the following:			
1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or			
1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or			
1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and			
2 The patient has performance status (WHO/ECOG) grade 0-2; and			
3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and			
4 The patient has an estimated life expectancy of at least 3 months.			
Continuation			
Haematologist			
<i>Re-assessment required after 12 months</i>			
Both:			
1 No evidence of disease progression, and; and			
2 The treatment remains appropriate and patient is benefitting from treatment.			
CAPECITABINE			
Tab 150 mg – 1% DV Sep-14 to 2016	30.00	60	Capecitabine Winthrop
Tab 500 mg – 1% DV Sep-14 to 2016	120.00	120	Capecitabine Winthrop
CLADRIBINE			
Inj 2 mg per ml, 5 ml vial			
Inj 1 mg per ml, 10 ml vial	5,249.72	7	Leustatin
CYTARABINE			
Inj 20 mg per ml, 5 ml vial – 1% DV Nov-13 to 2016	55.00	5	Pfizer
Inj 20 mg per ml, 25 ml vial	18.15	1	Pfizer
Inj 100 mg per ml, 10 ml vial – 1% DV Nov-13 to 2016	8.83	1	Pfizer
Inj 100 mg per ml, 20 ml vial – 1% DV Nov-13 to 2016	17.65	1	Pfizer

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FLUDARABINE PHOSPHATE			
Tab 10 mg – 1% DV Sep-15 to 2018	412.00	20	Fludara Oral
Inj 50 mg vial	525.00	5	Fludarabine Ebewe
FLUOROURACIL			
Inj 50 mg per ml, 20 ml vial – 1% DV Oct-15 to 2018	10.00	1	Fluorouracil Ebewe
Inj 50 mg per ml, 50 ml vial – 1% DV Oct-15 to 2018	17.00	1	Fluorouracil Ebewe
Inj 50 mg per ml, 100 ml vial – 1% DV Oct-15 to 2018	30.00	1	Fluorouracil Ebewe
GEMCITABINE			
Inj 10 mg per ml, 20 ml vial – 1% DV Oct-14 to 2017	8.36	1	Gemcitabine Ebewe
Inj 10 mg per ml, 100 ml vial – 1% DV Oct-14 to 2017	15.89	1	Gemcitabine Ebewe
MERCAPTOPURINE			
Tab 50 mg – 1% DV Oct-13 to 2016	49.41	25	Puri-nethol
METHOTREXATE			
Tab 2.5 mg – 1% DV Sep-15 to 2018	3.18	30	Trexate
Tab 10 mg – 1% DV Sep-15 to 2018	21.00	50	Trexate
Inj 2.5 mg per ml, 2 ml vial			
Inj 7.5 mg prefilled syringe – 1% DV Jan-14 to 2016	17.19	1	Methotrexate Sandoz
Inj 10 mg prefilled syringe – 1% DV Jan-14 to 2016	17.25	1	Methotrexate Sandoz
Inj 15 mg prefilled syringe – 1% DV Jan-14 to 2016	17.38	1	Methotrexate Sandoz
Inj 20 mg prefilled syringe – 1% DV Jan-14 to 2016	17.50	1	Methotrexate Sandoz
Inj 25 mg prefilled syringe – 1% DV Jan-14 to 2016	17.63	1	Methotrexate Sandoz
Inj 30 mg prefilled syringe – 1% DV Jan-14 to 2016	17.75	1	Methotrexate Sandoz
Inj 25 mg per ml, 2 ml vial – 1% DV Sep-13 to 2016	20.20	5	Hospira
Inj 25 mg per ml, 20 ml vial – 1% DV Sep-13 to 2016	27.78	1	Hospira
Inj 100 mg per ml, 10 ml vial	25.00	1	Methotrexate Ebewe
Inj 100 mg per ml, 50 ml vial – 1% DV Oct-14 to 2017	99.99	1	Methotrexate Ebewe
THIOGUANINE			
Tab 40 mg			

Other Cytotoxic Agents

AMSACRINE			
Inj 50 mg per ml, 1.5 ml ampoule			
Inj 75 mg			
ANAGRELIDE HYDROCHLORIDE			
Cap 0.5 mg			
ARSENIC TRIOXIDE			
Inj 1 mg per ml, 10 ml vial	4,817.00	10	AFT
BORTEZOMIB – Restricted see terms on the next page			
⚡ Inj 1 mg vial	540.70	1	Velcade
⚡ Inj 3.5 mg vial	1,892.50	1	Velcade

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔ Restricted			
Initiation — treatment naive multiple myeloma/amyloidosis			
<i>Limited to 15 months treatment</i>			
Both:			
1 Either:			
1.1 The patient has treatment-naive symptomatic multiple myeloma; or			
1.2 The patient has treatment-naive symptomatic systemic AL amyloidosis; and			
2 Maximum of 9 treatment cycles.			
Initiation — relapsed/refractory multiple myeloma/amyloidosis			
<i>Re-assessment required after 8 months</i>			
All of the following:			
1 Either:			
1.1 The patient has relapsed or refractory multiple myeloma; or			
1.2 The patient has relapsed or refractory systemic AL amyloidosis; and			
2 The patient has received only one prior front line chemotherapy for multiple myeloma or amyloidosis; and			
3 The patient has not had prior publicly funded treatment with bortezomib; and			
4 Maximum of 4 treatment cycles.			
Continuation — relapsed/refractory multiple myeloma/amyloidosis			
<i>Re-assessment required after 8 months</i>			
Both:			
1 The patient's disease obtained at least a partial response from treatment with bortezomib at the completion of cycle 4; and			
2 Maximum of 4 further treatment cycles (making a total maximum of 8 consecutive treatment cycles).			
Notes: Responding relapsed/refractory multiple myeloma patients should receive no more than 2 additional cycles of treatment beyond the cycle at which a confirmed complete response was first achieved. A line of therapy is considered to comprise either:			
1 A known therapeutic chemotherapy regimen and supportive treatments; or			
2 A transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments.			
Refer to datasheet for recommended dosage and number of doses of bortezomib per treatment cycle.			
COLASPASE [L-ASPARAGINASE]			
Inj 10,000 iu vial	102.32	1	Leunase
DACARBAZINE			
Inj 200 mg vial – 1% DV Oct-13 to 2016	51.84	1	Hospira
ETOPOSIDE			
Cap 50 mg	340.73	20	Vepesid
Cap 100 mg	340.73	10	Vepesid
Inj 20 mg per ml, 5 ml vial – 1% DV Apr-16 to 2018	7.90	1	Rex Medical
ETOPOSIDE (AS PHOSPHATE)			
Inj 100 mg vial	40.00	1	Etopophos
HYDROXYUREA			
Cap 500 mg	31.76	100	Hydrea
IRINOTECAN HYDROCHLORIDE			
Inj 20 mg per ml, 2 ml vial – 1% DV Sep-15 to 2018	11.50	1	Irinotecan Actavis 40
Inj 20 mg per ml, 5 ml vial – 1% DV Sep-15 to 2018	17.80	1	Irinotecan Actavis 100
LENALIDOMIDE – Restricted see terms on the next page			
⚡ Cap 10 mg	6,207.00	21	Revlimid
⚡ Cap 25 mg	7,627.00	21	Revlimid

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➡Restricted			
Initiation			
Haematologist			
<i>Re-assessment required after 6 months</i>			
All of the following:			
1 Patient has relapsed or refractory multiple myeloma with progressive disease; and			
2 Either:			
2.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or			
2.2 Both:			
2.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and			
2.2.2 The patient has experienced severe (grade ≥ 3), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and			
3 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.			
Continuation			
Haematologist			
<i>Re-assessment required after 6 months</i>			
Both:			
1 No evidence of disease progression; and			
2 The treatment remains appropriate and patient is benefitting from treatment.			
Note: Indication marked with * is an Unapproved Indication (refer to Interpretations and Definitions). A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.			
PEGASPARGASE – Restricted see terms below			
⚡ Inj 750 iu per ml, 5 ml vial	3,005.00	1	Oncaspar
➡Restricted			
Initiation — Newly diagnosed ALL			
<i>Limited to 12 months treatment</i>			
All of the following:			
1 The patient has newly diagnosed acute lymphoblastic leukaemia; and			
2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and			
3 Treatment is with curative intent.			
Initiation — Relapsed ALL			
<i>Limited to 12 months treatment</i>			
All of the following:			
1 The patient has relapsed acute lymphoblastic leukaemia; and			
2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and			
3 Treatment is with curative intent.			
PENTOSTATIN [DEOXYCOFORMYCIN]			
Inj 10 mg vial			
PROCARBAZINE HYDROCHLORIDE			
Cap 50 mg	498.00	50	Natulan
TEMOZOLOMIDE – Restricted see terms on the next page			
⚡ Cap 5 mg – 1% DV Sep-13 to 2016.....	8.00	5	Temaccord
⚡ Cap 20 mg – 1% DV Sep-13 to 2016.....	36.00	5	Temaccord
⚡ Cap 100 mg – 1% DV Sep-13 to 2016.....	175.00	5	Temaccord
⚡ Cap 250 mg – 1% DV Sep-13 to 2016.....	410.00	5	Temaccord

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Initiation			
All of the following:			
1 Either:			
1.1 Patient has newly diagnosed glioblastoma multiforme; or			
1.2 Patient has newly diagnosed anaplastic astrocytoma*; and			
2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and			
3 Following concomitant treatment temozolomide is to be used for a maximum of six cycles of 5 days treatment, at a maximum dose of 200 mg/m ² .			
Note: Indication marked with a * is an Unapproved Indication. Temozolomide is not funded for the treatment of relapsed glioblastoma multiforme. Reapplications will not be approved. Studies of temozolomide show that its benefit is predominantly in those patients with a good performance status (WHO grade 0 or 1 or Karnofsky score >80), and in patients who have had at least a partial resection of the tumour.			
THALIDOMIDE – Restricted see terms below			
⚡ Cap 50 mg	378.00	28	Thalomid
⚡ Cap 100 mg	756.00	28	Thalomid
➔Restricted			
Initiation			
<i>Re-assessment required after 12 months</i>			
Any of the following:			
1 The patient has multiple myeloma; or			
2 The patient has systemic AL amyloidosis*; or			
3 The patient has erythema nodosum leprosum.			
Continuation			
Patient has obtained a response from treatment during the initial approval period.			
Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier			
Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen			
Indication marked with * is an Unapproved Indication			
TRETINOIN			
Cap 10 mg	479.50	100	Vesanoid
Platinum Compounds			
CARBOPLATIN			
Inj 10 mg per ml, 5 ml vial – 1% DV Sep-15 to 2018	15.07	1	DBL Carboplatin
Inj 10 mg per ml, 15 ml vial – 1% DV Sep-15 to 2018	14.05	1	DBL Carboplatin
Inj 10 mg per ml, 45 ml vial – 1% DV Sep-15 to 2018	32.59	1	DBL Carboplatin
CISPLATIN			
Inj 1 mg per ml, 50 ml vial – 1% DV Nov-15 to 2018	12.29	1	DBL Cisplatin
Inj 1 mg per ml, 100 ml vial – 1% DV Nov-15 to 2018	22.46	1	DBL Cisplatin
OXALIPLATIN			
Inj 5 mg per ml, 10 ml vial – 1% DV Jun-16 to 2018	13.32	1	Oxaliccord
Inj 5 mg per ml, 20 ml vial – 1% DV Jun-16 to 2018	16.00	1	Oxaliccord
Protein-Tyrosine Kinase Inhibitors			
DASATINIB – Restricted see terms on the next page			
⚡ Tab 20 mg	3,774.06	60	Sprycel
⚡ Tab 50 mg	6,214.20	60	Sprycel
⚡ Tab 70 mg	7,692.58	60	Sprycel
⚡ Tab 100 mg	6,214.20	30	Sprycel

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Initiation			
For use in patients with approval from the CML/GIST Co-ordinator.			
ERLOTINIB – Restricted see terms below			
☞ Tab 100 mg – 1% DV Jun-15 to 2018	1,000.00	30	Tarceva
☞ Tab 150 mg – 1% DV Jun-15 to 2018	1,500.00	30	Tarceva
➔Restricted			
Initiation			
<i>Re-assessment required after 4 months</i>			
All of the following:			
1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and			
2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and			
3 Any of the following:			
3.1 Patient is treatment naive; or			
3.2 Both:			
3.2.1 Patient has documented disease progression following treatment with first line platinum based chemotherapy; and			
3.2.2 Patient has not received prior treatment with gefitinib; or			
3.3 Both:			
3.3.1 The patient has discontinued gefitinib within 12 weeks of starting treatment due to intolerance; and			
3.3.2 The cancer did not progress while on gefitinib; and			
4 Erlotinib is to be given for a maximum of 3 months.			
Continuation			
<i>Re-assessment required after 6 months</i>			
Both:			
1 Radiological assessment (preferably including CT scan) indicates NSCLC has not progressed; and			
2 Erlotinib is to be given for a maximum of 3 months.			
GEFITINIB – Restricted see terms below			
☞ Tab 250 mg	1,700.00	30	Iressa
➔Restricted			
Initiation			
<i>Re-assessment required after 4 months</i>			
All of the following:			
1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and			
2 Either:			
2.1 Patient is treatment naive; or			
2.2 Both:			
2.2.1 The patient has discontinued erlotinib within 12 weeks of starting treatment due to intolerance; and			
2.2.2 The cancer did not progress whilst on erlotinib; and			
3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and			
4 Gefitinib is to be given for a maximum of 3 months.			
Continuation			
<i>Re-assessment required after 6 months</i>			
Both:			
1 Radiological assessment (preferably including CT scan) indicates NSCLC has not progressed; and			
2 Gefitinib is to be given for a maximum of 3 months.			
IMATINIB MESILATE			
Imatinib-AFT is not a registered for the treatment of Gastro Intestinal Stromal Tumours (GIST). The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule			
☞ Tab 100 mg	2,400.00	60	Glivec

☞ Item restricted (see ➔ above); ☞ Item restricted (see ➔ below)

e.g. Brand indicates brand example only. It is not a contracted product.

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

➔ **Restricted**

Initiation

Re-assessment required after 12 months

Both:

- 1 Patient has diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Maximum dose of 400 mg/day.

Continuation

Re-assessment required after 12 months

Adequate clinical response to treatment with imatinib (prescriber determined).

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule.

Cap 100 mg – 1% DV Jul-14 to 2017	298.90	60	Imatinib-AFT
Cap 400 mg	597.80	30	Imatinib-AFT

LAPATINIB – **Restricted** see terms below

⚡ Tab 250 mg	1,899.00	70	Tykerb
--------------------	----------	----	--------

➔ **Restricted**

Initiation

Re-assessment required after 12 months

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient has not previously received trastuzumab treatment for HER 2 positive metastatic breast cancer; and
 - 1.3 Lapatinib not to be given in combination with trastuzumab; and
 - 1.4 Lapatinib to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2.2 The patient started trastuzumab for metastatic breast cancer but discontinued trastuzumab within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on trastuzumab; and
 - 2.4 Lapatinib not to be given in combination with trastuzumab; and
 - 2.5 Lapatinib to be discontinued at disease progression.

Continuation

Re-assessment required after 12 months

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB – **Restricted** see terms on the next page

⚡ Cap 150 mg	4,680.00	120	Tasigna
⚡ Cap 200 mg	6,532.00	120	Tasigna

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Initiation			
Haematologist			
<i>Re-assessment required after 6 months</i>			
All of the following:			
1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and			
2 Either:			
2.1 Patient has documented CML treatment failure* with imatinib; or			
2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and			
3 Maximum nilotinib dose of 800 mg/day; and			
4 Subsidised for use as monotherapy only.			
Note: *treatment failure as defined by Leukaemia Net Guidelines.			
Continuation			
Haematologist			
<i>Re-assessment required after 6 months</i>			
All of the following:			
1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and			
2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and			
3 Maximum nilotinib dose of 800 mg/day; and			
4 Subsidised for use as monotherapy only.			
PAZOPANIB – Restricted see terms below			
☞ Tab 200 mg	1,334.70	30	Votrient
☞ Tab 400 mg	2,669.40	30	Votrient
➔Restricted			
Initiation			
<i>Re-assessment required after 3 months</i>			
All of the following:			
1 The patient has metastatic renal cell carcinoma; and			
2 Any of the following:			
2.1 The patient is treatment naive; or			
2.2 The patient has only received prior cytokine treatment; or			
2.3 Both:			
2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and			
2.3.2 The cancer did not progress whilst on sunitinib; and			
3 The patient has good performance status (WHO/ECOG grade 0-2); and			
4 The disease is of predominant clear cell histology; and			
5 All of the following:			
5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; and			
5.2 Haemoglobin level < lower limit of normal; and			
5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); and			
5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; and			
5.5 Karnofsky performance score of ≤ 70; and			
5.6 ≥ 2 sites of organ metastasis.			
Continuation			
<i>Re-assessment required after 3 months</i>			
Both:			
1 No evidence of disease progression; and			
2 The treatment remains appropriate and the patient is benefiting from treatment.			
Notes: Pazopanib treatment should be stopped if disease progresses.			
Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.			
⚡ Item restricted (see ➔ above); ☞ Item restricted (see ➔ below) e.g. Brand indicates brand example only. It is not a contracted product.			

140

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SUNITINIB – Restricted see terms below			
⚡ Cap 12.5 mg	2,315.38	28	Sutent
⚡ Cap 25 mg	4,630.77	28	Sutent
⚡ Cap 50 mg	9,261.54	28	Sutent

➡ **Restricted**

Initiation — RCC

Re-assessment required after 3 months

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
- 5 All of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; and
 - 5.2 Haemoglobin level < lower limit of normal; and
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); and
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; and
 - 5.5 Karnofsky performance score of ≤ 70; and
 - 5.6 ≥ 2 sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Notes: RCC - Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

Continuation — RCC

Re-assessment required after 3 months

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Initiation — GIST

Re-assessment required after 3 months

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Continuation — GIST

Re-assessment required after 6 months

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.2 The patient has had a partial response (a decrease in size of $\geq 10\%$ or decrease in tumour density in Hounsfield Units (HU) of $\geq 15\%$ on CT and no new lesions and no obvious progression of non-measurable disease); or
- 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and

2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: GIST - It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of $\geq 10\%$ and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Taxanes

DOCETAXEL

Inj 10 mg per ml, 2 ml vial – 1% DV Dec-14 to 2017	13.70	1	DBL Docetaxel
Inj 10 mg per ml, 8 ml vial – 1% DV Dec-14 to 2017	29.99	1	DBL Docetaxel

PACLITAXEL

Inj 6 mg per ml, 5 ml vial – 1% DV Sep-14 to 2017	45.00	5	Paclitaxel Ebewe
Inj 6 mg per ml, 16.7 ml vial – 1% DV Sep-14 to 2017	19.02	1	Paclitaxel Ebewe
Inj 6 mg per ml, 25 ml vial – 1% DV Sep-14 to 2017	26.69	1	Paclitaxel Ebewe
Inj 6 mg per ml, 50 ml vial – 1% DV Sep-14 to 2017	36.53	1	Paclitaxel Ebewe
Inj 6 mg per ml, 100 ml vial – 1% DV Sep-14 to 2017	73.06	1	Paclitaxel Ebewe

Treatment of Cytotoxic-Induced Side Effects

CALCIUM FOLINATE

Tab 15 mg	104.26	10	DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml ampoule			
Inj 10 mg per ml, 5 ml ampoule – 1% DV Oct-14 to 2017	18.25	5	Calcium Folate Ebewe
Inj 10 mg per ml, 10 ml vial – 1% DV Oct-14 to 2017	7.33	1	Calcium Folate Ebewe
Inj 10 mg per ml, 30 ml vial – 1% DV Oct-14 to 2017	22.51	1	Calcium Folate Ebewe
Inj 10 mg per ml, 100 ml vial – 1% DV Oct-14 to 2017	67.51	1	Calcium Folate Ebewe

MESNA

Tab 400 mg – 1% DV Oct-13 to 2016	227.50	50	Uromitexan
Tab 600 mg – 1% DV Oct-13 to 2016	339.50	50	Uromitexan
Inj 100 mg per ml, 4 ml ampoule – 1% DV Oct-13 to 2016	148.05	15	Uromitexan
Inj 100 mg per ml, 10 ml ampoule – 1% DV Oct-13 to 2016	339.90	15	Uromitexan

Vinca Alkaloids

VINBLASTINE SULPHATE

Inj 1 mg per ml, 10 ml vial	186.46	5	Hospira
-----------------------------------	--------	---	---------

VINCISTINE SULPHATE

Inj 1 mg per ml, 1 ml vial – 1% DV Sep-13 to 2016	64.80	5	Hospira
Inj 1 mg per ml, 2 ml vial – 1% DV Sep-13 to 2016	69.60	5	Hospira

VINORELBINE

Inj 10 mg per ml, 1 ml vial – 1% DV Sep-15 to 2018	8.00	1	Navelbine
Inj 10 mg per ml, 5 ml vial – 1% DV Sep-15 to 2018	40.00	1	Navelbine

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Endocrine Therapy			
ABIRATERONE ACETATE – Restricted see terms below			
⚡ Tab 250 mg	4,276.19	120	Zytiga
➡ Restricted			
Initiation			
Medical oncologist, radiation oncologist or urologist			
<i>Re-assessment required after 5 months</i>			
All of the following:			
1 Patient has prostate cancer; and			
2 Patient has metastases; and			
3 Patient's disease is castration resistant; and			
4 Either:			
4.1 All of the following:			
4.1.1 Patient is symptomatic; and			
4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and			
4.1.3 Patient has ECOG performance score of 0-1; and			
4.1.4 Patient has not had prior treatment with taxane chemotherapy; or			
4.2 All of the following:			
4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and			
4.2.2 Patient has ECOG performance score of 0-2; and			
4.2.3 Patient has not had prior treatment with abiraterone.			
Continuation			
Medical oncologist, radiation oncologist or urologist			
<i>Re-assessment required after 5 months</i>			
All of the following:			
1 Significant decrease in serum PSA from baseline; and			
2 No evidence of clinical disease progression; and			
3 No initiation of taxane chemotherapy with abiraterone; and			
4 The treatment remains appropriate and the patient is benefiting from treatment.			
BICALUTAMIDE			
Tab 50 mg – 1% DV Sep-14 to 2017	4.90	28	Bicalaccord
FLUTAMIDE			
Tab 250 mg	55.00	100	Flutamin
MEGESTROL ACETATE			
Tab 160 mg – 1% DV Oct-15 to 2018	54.30	30	Apo-Megestrol
OCTREOTIDE – Some items restricted see terms on the next page			
Inj 50 mcg per ml, 1 ml ampoule – 1% DV Sep-14 to 2017	13.50	5	DBL
Inj 100 mcg per ml, 1 ml ampoule – 1% DV Sep-14 to 2017	22.40	5	DBL
Inj 500 mcg per ml, 1 ml ampoule – 1% DV Sep-14 to 2017	89.40	5	DBL
⚡ Inj 10 mg vial	1,772.50	1	Sandostatin LAR
⚡ Inj 20 mg vial	2,358.75	1	Sandostatin LAR
⚡ Inj 30 mg vial	2,951.25	1	Sandostatin LAR

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Initiation — Malignant bowel obstruction			
All of the following:			
1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and			
2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and			
3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.			
Note: Indications marked with * are Unapproved Indications			
Initiation — acromegaly			
<i>Re-assessment required after 3 months</i>			
Both:			
1 The patient has acromegaly; and			
2 Any of the following:			
2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or			
2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or			
2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.			
Continuation — acromegaly			
Both:			
1 IGF1 levels have decreased since starting octreotide; and			
2 The treatment remains appropriate and the patient is benefiting from treatment.			
Note: In patients with acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks.			
Initiation — Other indications			
Any of the following:			
1 VIPomas and glucagonomas - for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or			
2 Both:			
2.1 Gastrinoma; and			
2.2 Either:			
2.2.1 Patient has failed surgery; or			
2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or			
3 Both:			
3.1 Insulinomas; and			
3.2 Surgery is contraindicated or has failed; or			
4 For pre-operative control of hypoglycaemia and for maintenance therapy; or			
5 Both:			
5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and			
5.2 Disabling symptoms not controlled by maximal medical therapy.			
Note: restriction applies only to the long-acting formulations of octreotide			
TAMOXIFEN CITRATE			
Tab 10 mg	17.50	100	Genox
Tab 20 mg	2.63	30	Genox
	8.75	100	Genox
Aromatase Inhibitors			
ANASTROZOLE			
Tab 1 mg	26.55	30	Aremed DP-Anastrozole

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
EXEMESTANE			
Tab 25 mg – 1% DV Sep-14 to 2017	14.50	30	Aromasin
LETROZOLE			
Tab 2.5 mg – 1% DV Jan-16 to 2018	2.95	30	Letrole

Immunosuppressants

Calcineurin Inhibitors

CICLOSPORIN			
Cap 25 mg	44.63	50	Neoral
Cap 50 mg	88.91	50	Neoral
Cap 100 mg	177.81	50	Neoral
Oral liq 100 mg per ml	198.13	50 ml	Neoral
Inj 50 mg per ml, 5 ml ampoule – 1% DV Sep-15 to 2018	276.30	10	Sandimmun
TACROLIMUS – Restricted see terms below			
⚡ Cap 0.5 mg – 1% DV Nov-14 to 31 Oct 2018	85.60	100	Tacrolimus Sandoz
⚡ Cap 1 mg – 1% DV Nov-14 to 31 Oct 2018	171.20	100	Tacrolimus Sandoz
⚡ Cap 5 mg – 1% DV Nov-14 to 31 Oct 2018	428.00	50	Tacrolimus Sandoz
⚡ Inj 5 mg per ml, 1 ml ampoule			

➡ Restricted

Initiation — organ transplant recipients

Any specialist

For use in organ transplant recipients.

Initiation — Steroid-resistant nephrotic syndrome*

Any specialist

Either:

- 1 The patient is a child with steroid-resistant nephrotic syndrome* (SRNS) where ciclosporin has been trialled in combination with prednisone and discontinued because of unacceptable side effects or inadequate clinical response; or
- 2 All of the following:
 - 2.1 The patient is an adult with SRNS; and
 - 2.2 Ciclosporin has been trialled in combination with prednisone and discontinued because of unacceptable side effects or inadequate clinical response; and
 - 2.3 Cyclophosphamide or mycophenolate have been trialled and discontinued because of unacceptable side effects or inadequate clinical response, or these treatments are contraindicated.

Note: Indications marked with * are Unapproved Indications

Fusion Proteins

ETANERCEPT – Restricted see terms below

⚡ Inj 25 mg vial	799.96	4	Enbrel
⚡ Inj 50 mg autoinjector	1,599.96	4	Enbrel
⚡ Inj 50 mg syringe	1,599.96	4	Enbrel

➡ Restricted

Initiation — juvenile idiopathic arthritis

Rheumatologist or named specialist

Re-assessment required after 4 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for juvenile idiopathic arthritis (JIA); and
- continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for JIA; or
- 2 All of the following:
 - 2.1 Patient diagnosed with Juvenile Idiopathic Arthritis (JIA); and
 - 2.2 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
 - 2.5 Both:
 - 2.5.1 Either:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

Continuation — juvenile idiopathic arthritis

Rheumatologist or named specialist

Re-assessment required after 6 months

- Both:
- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation — rheumatoid arthritis

Rheumatologist

Re-assessment required after 6 months

- Either:
- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis; or
 - 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulphasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:

continued...

continued...

- 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
- 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation — rheumatoid arthritis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initiation — ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of an exercise regime supervised by a physiotherapist; and
 - 2.5 Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or

2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and

2.6 Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of starting treatment.

Average normal chest expansion corrected for age and gender:

Age	Male	Female
18-24	7.0 cm	5.5 cm
25-34	7.5 cm	5.5 cm
35-44	6.5 cm	4.5 cm
45-54	6.0 cm	5.0 cm
55-64	5.5 cm	4.0 cm
65-74	4.0 cm	4.0 cm
75+	3.0 cm	2.5 cm

Continuation — ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Following 12 weeks of etanercept treatment, BASDAI has improved by 4 or more points from pre-treatment baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initiation — psoriatic arthritis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulphasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
- 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation — psoriatic arthritis

Rheumatologist

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initiation — plaque psoriasis, prior TNF use

Dermatologist

Limited to 4 months treatment

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; and
- 3 Patient must be reassessed for continuation after 3 doses.

Initiation — plaque psoriasis, treatment-naïve

Dermatologist

Limited to 4 months treatment

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Continuation — plaque psoriasis

Dermatologist

Re-assessment required after 6 months

Both:

1 Either:

1.1 Both:

1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and

1.1.2 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-etanercept treatment baseline value; or

1.2 Both:

1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and

1.2.2 Either:

1.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or

1.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-etanercept treatment baseline value; and

2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initiation — pyoderma gangrenosum

Dermatologist

All of the following:

1 Patient has pyoderma gangrenosum*; and

2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and

3 A maximum of 4 doses.

Note: Indications marked with * are Unapproved Indications (refer to Section A: General Rules, Part I (Interpretations and Definitions) and Part V (Miscellaneous Provisions) rule 5.5).

Continuation — pyoderma gangrenosum

Dermatologist

All of the following:

1 Patient has shown clinical improvement; and

2 Patient continues to require treatment; and

3 A maximum of 4 doses.

Initiation — adult-onset Still's disease

Rheumatologist

Re-assessment required after 6 months

Either:

1 Both:

1.1 Either:

1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the Section H rules; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Continuation — adult-onset Still's disease

Rheumatologist

Re-assessment required after 6 months

The patient has a sustained improvement in inflammatory markers and functional status.

Monoclonal Antibodies

ABCIXIMAB – **Restricted** see terms below

⚡ Inj 2 mg per ml, 5 ml vial 579.53 1 ReoPro

➡ **Restricted**

Initiation

Either:

- 1 For use in patients with acute coronary syndromes undergoing percutaneous coronary intervention; or
- 2 For use in patients undergoing intra-cranial intervention.

ADALIMUMAB – **Restricted** see terms below

⚡ Inj 10 mg per 0.2 ml prefilled syringe 1,599.96 2 Humira
 ⚡ Inj 20 mg per 0.4 ml syringe 1,599.96 2 Humira
 ⚡ Inj 40 mg per 0.8 ml pen 1,599.96 2 HumiraPen
 ⚡ Inj 40 mg per 0.8 ml syringe 1,599.96 2 Humira

➡ **Restricted**

Initiation — juvenile idiopathic arthritis

Rheumatologist or named specialist

Re-assessment required after 4 months

Either:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for juvenile idiopathic arthritis (JIA); and
 - 1.1.2 Either:
 - 1.1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for JIA; or
 - 2 All of the following:
 - 2.1 Patient diagnosed with Juvenile Idiopathic Arthritis (JIA); and
 - 2.2 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and

2.5 Both:

2.5.1 Either:

- 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or

- 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and

2.5.2 Physician's global assessment indicating severe disease.

Continuation — juvenile idiopathic arthritis

Rheumatologist or named specialist

Re-assessment required after 6 months

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation — fistulising Crohn's disease

Gastroenterologist

Re-assessment required after 4 months

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment (a copy of which is available at www.pharmac.govt.nz/latest/BaselineFistulaAssessment.pdf) has been completed and is no more than 1 month old at the time of application.

Continuation — fistulising Crohn's disease

Gastroenterologist

Re-assessment required after 6 months

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initiation — Crohn's disease

Gastroenterologist

Re-assessment required after 3 months

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Continuation — Crohn's disease

Gastroenterologist

Re-assessment required after 3 months

Both:

- 1 Either:
 - 1.1 Either:
 - 1.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 1.1.2 CDAI score is 150 or less; or
 - 1.2 Both:
 - 1.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 1.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation — rheumatoid arthritis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulphasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.7 Either:

continued...

continued...

- 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation — rheumatoid arthritis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation — ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of an exercise regime supervised by a physiotherapist; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of starting treatment.

Average normal chest expansion corrected for age and gender:

Age	Male	Female
18-24	7.0 cm	5.5 cm
25-34	7.5 cm	5.5 cm
35-44	6.5 cm	4.5 cm
45-54	6.0 cm	5.0 cm
55-64	5.5 cm	4.0 cm
65-74	4.0 cm	4.0 cm
75+	3.0 cm	2.5 cm

Continuation — ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Following 12 weeks of adalimumab treatment, BASDAI has improved by 4 or more points from pre-treatment baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation — psoriatic arthritis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulphasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation — psoriatic arthritis

Rheumatologist

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation — plaque psoriasis, prior TNF use

Dermatologist

Limited to 4 months treatment

Both:

- 1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from etanercept; or
 - 2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis.

Initiation — plaque psoriasis, treatment-naïve

Dermatologist

Limited to 4 months treatment

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Continuation — plaque psoriasis

Dermatologist

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 1.1.2 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
- 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-etanercept treatment baseline value; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation — pyoderma gangrenosum

Dermatologist

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 4 doses.

Note: Indications marked with * are Unapproved Indications (refer to Section A: General Rules, Part I (Interpretations and Definitions) and Part V (Miscellaneous Provisions) rule 5.5).

Continuation — pyoderma gangrenosum

Dermatologist

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 4 doses.

Initiation — adult-onset Still's disease

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the Section H rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Continuation — adult-onset Still's disease

Rheumatologist

Re-assessment required after 6 months

The patient has a sustained improvement in inflammatory markers and functional status.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BASILIXIMAB – Restricted see terms below			
☞ Inj 20 mg vial	3,200.00	1	Simulect
☞ Restricted			
Initiation			
For use in solid organ transplants.			
BEVACIZUMAB – Restricted see terms below			
☞ Inj 25 mg per ml, 4 ml vial			
☞ Inj 25 mg per ml, 16 ml vial			
☞ Restricted			
Initiation			
Either:			
1 Ocular neovascularisation; or			
2 Exudative ocular angiopathy.			
INFLIXIMAB – Restricted see terms below			
☞ Inj 100 mg – 10% DV Mar-15 to 29 Feb 2020	806.00	1	Remicade
☞ Restricted			
Initiation — Graft vs host disease			
Patient has steroid-refractory acute graft vs. host disease of the gut.			
Initiation — rheumatoid arthritis			
Rheumatologist			
<i>Re-assessment required after 4 months</i>			
All of the following:			
1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and			
2 Either:			
2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or			
2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and			
3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.			
Continuation — rheumatoid arthritis			
Rheumatologist			
<i>Re-assessment required after 6 months</i>			
All of the following:			
1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and			
2 Either:			
2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or			
2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and			
3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.			
Initiation — ankylosing spondylitis			
Rheumatologist			
<i>Re-assessment required after 3 months</i>			
Both:			
1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and			
2 Either:			
2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or			
continued...			
⚡ Item restricted (see ☞ above); ☞ Item restricted (see ☞ below) e.g. <i>Brand</i> indicates brand example only. It is not a contracted product.			

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Continuation — ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initiation — psoriatic arthritis

Rheumatologist

Re-assessment required after 4 months

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for psoriatic arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for psoriatic arthritis. .

Continuation — psoriatic arthritis

Rheumatologist

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initiation — severe ocular inflammation

Therapy limited to 3 doses

Both:

- 1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
- 2 Either:
 - 2.1 Patient has failed to achieve control of severe vision-threatening ocular inflammation following high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids; or
 - 2.2 Patient developed new inflammatory symptoms while receiving high dose steroids.

Initiation — chronic ocular inflammation

Therapy limited to 3 doses

Both:

- 1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2 Patient has tried at least two other immunomodulatory agents.

Continuation — ocular inflammation

Both:

- 1 Patient had a good clinical response to initial treatment; and
- 2 Either:
 - 2.1 A withdrawal of infliximab has been trialled and patient has relapsed after trial withdrawal; or
 - 2.2 Patient has Behcet's disease.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation — Pulmonary sarcoidosis

Both:

- 1 Patient has life-threatening pulmonary sarcoidosis that is refractory to other treatments; and
- 2 Treatment is to be prescribed by, or has been recommended by, a physician with expertise in the treatment of pulmonary sarcoidosis.

Initiation — Crohn's disease (adults)

Gastroenterologist

Re-assessment required after 3 months

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Continuation — Crohn's disease (adults)

Gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation — Crohn's disease (children)

Gastroenterologist

Re-assessment required after 3 months

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Continuation — Crohn's disease (children)

Gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Any of the following:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
- 1.2 PCDAI score is 15 or less; or
- 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation — fistulising Crohn's disease

Gastroenterologist

Re-assessment required after 4 months

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Continuation — fistulising Crohn's disease

Gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation — acute severe fulminant ulcerative colitis

Gastroenterologist

Limited to 6 weeks treatment

Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Continuation — severe fulminant ulcerative colitis

Gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation — severe ulcerative colitis

Gastroenterologist

Re-assessment required after 3 months

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is ≥ 4 ; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is ≥ 65 ; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Continuation — severe ulcerative colitis

Gastroenterologist

Re-assessment required after 6 months

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by ≥ 2 points from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by ≥ 30 points from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation — plaque psoriasis

Dermatologist

Re-assessment required after 3 doses

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, cyclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Continuation — plaque psoriasis

Dermatologist

Re-assessment required after 3 doses

Both:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

OMALIZUMAB – **Restricted** see terms below

⌚ Inj 150 mg vial	500.00	1	Xolair
-------------------------	--------	---	--------

➡ Restricted

Initiation

Respiratory specialist

Re-assessment required after 6 months

All of the following:

- 1 Patient is over the age of 6; and
- 2 Patient has a diagnosis of severe, life threatening asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven compliance with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1600 micrograms per day or fluticasone propionate 1000 micrograms per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 micrograms bd or eformoterol 12 micrograms bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; and
- 7 At least four admissions to hospital for a severe asthma exacerbation over the previous 24 months with at least one of those being in the previous 12 months; and
- 8 An Asthma Control Questionnaire (ACQ-5) score of at least 3.0 as assessed in the previous month.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation

Respiratory specialist

Re-assessment required after 6 months

All of the following:

- 1 Hospital admissions have been reduced as a result of treatment; and
- 2 A reduction in the Asthma Control Questionnaire (ACQ-5) score of at least 1.0 from baseline; and
- 3 A reduction in the maintenance oral corticosteroid dose of at least 50% from baseline.

RANIBIZUMAB – Restricted see terms below

⚡ Inj 10 mg per ml, 0.23 ml vial

⚡ Inj 10 mg per ml, 0.3 ml vial

➡ **Restricted**

Initiation

Re-assessment required after 3 doses

Both:

- 1 Either:
 - 1.1 Age-related macular degeneration; or
 - 1.2 Choroidal neovascular membrane; and
- 2 Any of the following:
 - 2.1 The patient has had a severe ophthalmic inflammatory response following bevacizumab; or
 - 2.2 The patient has had a myocardial infarction or stroke within the last three months; or
 - 2.3 The patient has failed to respond to bevacizumab following three intraocular injections; or
 - 2.4 The patient is of child-bearing potential and has not completed a family.

Continuation

Both:

- 1 Documented benefit after three doses must be demonstrated to continue; and
- 2 In the case of but previous non-response to bevacizumab, a retreat of bevacizumab is required to confirm non-response before continuing with ranibizumab.

RITUXIMAB – Restricted see terms below

⚡ Inj 10 mg per ml, 10 ml vial 1,075.50 2 Mabthera

⚡ Inj 10 mg per ml, 50 ml vial 2,688.30 1 Mabthera

➡ **Restricted**

Initiation — haemophilia with inhibitors

Haematologist

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Continuation — haemophilia with inhibitors

Haematologist

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initiation — post-transplant

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued. . .

Note: Indications marked with * are Unapproved Indications.

Continuation — post-transplant

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are Unapproved Indications.

Initiation — indolent, low-grade lymphomas

Either:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia.

Continuation — indolent, low-grade lymphomas

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia.

Initiation — aggressive CD20 positive NHL

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia.

Continuation — aggressive CD20 positive NHL

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia.

Initiation — Chronic lymphocytic leukaemia

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 The patient is rituximab treatment naive; and
- 3 Either:
 - 3.1 The patient is chemotherapy treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and

continued. . .

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

3.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; and

- 4 The patient has good performance status; and
- 5 The patient has good renal function (creatinine clearance \geq 30 ml/min); and
- 6 The patient does not have chromosome 17p deletion CLL; and
- 7 Rituximab to be administered in combination with fludarabine and cyclophosphamide for a maximum of 6 treatment cycles; and
- 8 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration).

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to <2 .

Initiation — rheumatoid arthritis - prior TNF inhibitor use

Rheumatologist

Limited to 4 months treatment

All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initiation — rheumatoid arthritis - TNF inhibitors contraindicated

Rheumatologist

Limited to 4 months treatment

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulphasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of cyclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Continuation — rheumatoid arthritis - re-treatment in 'partial responders' to rituximab

Rheumatologist

Re-assessment required after 4 months

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Continuation — rheumatoid arthritis - re-treatment in 'responders' to rituximab

Rheumatologist

Re-assessment required after 4 months

All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initiation — severe cold haemagglutinin disease (CHAD)

Haematologist

Re-assessment required after 4 weeks

Both:

- 1 Patient has cold haemagglutinin disease*; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms.

Note: Indications marked with * are Unapproved Indications.

Continuation — severe cold haemagglutinin disease (CHAD)

Haematologist

Re-assessment required after 4 weeks

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are Unapproved Indications.

Initiation — warm autoimmune haemolytic anaemia (warm AIHA)

Haematologist

Re-assessment required after 4 weeks

Both:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to >5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin.

Note: Indications marked with * are Unapproved Indications.

Continuation — warm autoimmune haemolytic anaemia (warm AIHA)

Haematologist

Re-assessment required after 4 weeks

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are Unapproved Indications.

Initiation — immune thrombocytopenic purpura (ITP)

Haematologist

Re-assessment required after 4 weeks

Both:

- 1 Either:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of \leq 20,000 platelets per microlitre; or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy).

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued. . .

Note: Indications marked with * are Unapproved Indications.

Continuation — immune thrombocytopenic purpura (ITP)

Haematologist

Re-assessment required after 4 weeks

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are Unapproved Indications.

Initiation — thrombotic thrombocytopenic purpura (TTP)

Haematologist

Re-assessment required after 4 weeks

Either:

- 1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
- 2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are Unapproved Indications.

Continuation — thrombotic thrombocytopenic purpura (TTP)

Haematologist

Re-assessment required after 4 weeks

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Note: Indications marked with * are Unapproved Indications.

Initiation — pure red cell aplasia (PRCA)

Haematologist

Re-assessment required after 6 weeks

Patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are Unapproved Indications.

Continuation — pure red cell aplasia (PRCA)

Haematologist

Re-assessment required after 6 weeks

Patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are Unapproved Indications.

Initiation — ANCA associated vasculitis

Re-assessment required after 4 weeks

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Either:
 - 2.1 Patient does not have MPO-ANCA positive vasculitis*; or
 - 2.2 Mycophenolate mofetil has not been effective in those patients who have MPO-ANCA positive vasculitis*; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 4 Any of the following:

continued. . .

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 4.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve complete absence of disease after at least 3 months; or
- 4.2 Patient has previously had a cumulative dose of cyclophosphamide >15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose >15 g; or
- 4.3 Cyclophosphamide and methotrexate are contraindicated; or
- 4.4 Patient is a female of child-bearing potential; or
- 4.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are Unapproved Indications.

Continuation — ANCA associated vasculitis

Re-assessment required after 4 weeks

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are Unapproved Indications.

Initiation — treatment refractory systemic lupus erythematosus (SLE)

Rheumatologist or nephrologist

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are Unapproved Indications.

Continuation — treatment refractory systemic lupus erythematosus (SLE)

Rheumatologist or nephrologist

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are Unapproved Indications.

Initiation — Antibody-mediated renal transplant rejection

Nephrologist

Patient has been diagnosed with antibody-mediated renal transplant rejection*.

Note: Indications marked with * are Unapproved Indications.

Initiation — ABO-incompatible renal transplant

Nephrologist

Patient is to undergo an ABO-incompatible renal transplant*.

Note: Indications marked with * are Unapproved Indications.

SILTUXIMAB – **Restricted** see terms on the next page

⚡ Inj 100 mg vial – 1% DV Jun-16 to 2018	770.57	1	Sylvant
⚡ Inj 400 mg vial – 1% DV Jun-16 to 2018	3,082.33	1	Sylvant

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔Restricted			
Initiation			
Haematologist or rheumatologist			
<i>Re-assessment required after 6 months</i>			
All of the following:			
1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and			
2 Treatment with an adequate trial of corticosteroids has proven ineffective; and			
3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.			
Continuation			
Haematologist or rheumatologist			
<i>Re-assessment required after 12 months</i>			
The treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.			
TOCILIZUMAB – Restricted see terms below			
⚡ Inj 20 mg per ml, 4 ml vial	220.00	1	Actemra
⚡ Inj 20 mg per ml, 10 ml vial	550.00	1	Actemra
⚡ Inj 20 mg per ml, 20 ml vial	1,100.00	1	Actemra
➔Restricted			
Initiation — Rheumatoid Arthritis			
Rheumatologist			
<i>Re-assessment required after 6 months</i>			
Either:			
1 All of the following:			
1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and			
1.2 Either:			
1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or			
1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and			
1.3 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and			
1.4 Either:			
1.4.1 The patient has experienced intolerable side effects from rituximab; or			
1.4.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis; or			
2 All of the following:			
2.1 Patient has had severe and active erosive rheumatoid arthritis for six months duration or longer; and			
2.2 Tocilizumab is to be used as monotherapy; and			
2.3 Either:			
2.3.1 Treatment with methotrexate is contraindicated; or			
2.3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and			
2.4 Either:			
2.4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of cyclosporin alone or in combination with another agent; or			
2.4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and			
2.5 Either:			
2.5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or			
2.5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2.6 Either:

- 2.6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 2.6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation — Rheumatoid Arthritis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initiation — systemic juvenile idiopathic arthritis

Rheumatologist

Re-assessment required after 6 months

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Continuation — systemic juvenile idiopathic arthritis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Initiation — adult-onset Still's disease

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Continuation — adult-onset Still's disease

Rheumatologist

Re-assessment required after 6 months

The patient has a sustained improvement in inflammatory markers and functional status.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
TRASTUZUMAB – Restricted see terms below			
¶ Inj 150 mg vial	1,350.00	1	Herceptin
¶ Inj 440 mg vial	3,875.00	1	Herceptin

➔ **Restricted**

Initiation — Early breast cancer

Limited to 12 months treatment

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Initiation — metastatic breast cancer (trastuzumab-naïve patients)

Limited to 12 months treatment

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient has not previously received lapatinib treatment for HER 2 positive metastatic breast cancer; and
 - 1.3 Trastuzumab not to be given in combination with lapatinib; and
 - 1.4 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2.2 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on lapatinib; and
 - 2.4 Trastuzumab not to be given in combination with lapatinib; and
 - 2.5 Trastuzumab to be discontinued at disease progression.

Initiation — metastatic breast cancer (patients previously treated with trastuzumab)

Limited to 12 months treatment

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 All of the following:
 - 3.1.1 The patient has not previously received lapatinib treatment for metastatic breast cancer; and
 - 3.1.2 Trastuzumab not to be given in combination with lapatinib; and
 - 3.1.3 Trastuzumab to be discontinued at disease progression; or
 - 3.2 All of the following:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; and
 - 3.2.3 Trastuzumab not to be given in combination with lapatinib; and
 - 3.2.4 Trastuzumab to be discontinued at disease progression; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

3.3 All of the following:

- 3.3.1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3.3.2 Trastuzumab not to be given in combination with lapatinib; and
- 3.3.3 Trastuzumab to be discontinued at disease progression.

Continuation — metastatic breast cancer

Re-assessment required after 12 months

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Other Immunosuppressants

ANTITHYMOCYTE GLOBULIN (EQUINE)

Inj 50 mg per ml, 5 ml ampoule	2,351.25	5	ATGAM
--------------------------------------	----------	---	-------

ANTITHYMOCYTE GLOBULIN (RABBIT)

Inj 25 mg vial

AZATHIOPRINE

Tab 25 mg	8.28	60	Azamun
Tab 50 mg – 1% DV Jun-14 to 2016	13.22	100	Azamun
Inj 50 mg vial	126.00	1	Imuran

BACILLUS CALMETTE-GUERIN (BCG) – **Restricted** see terms below

☞ Inj 2-8 × 10 ⁸ CFU vial – 1% DV Sep-13 to 2016	149.37	1	OncoTICE
☞ Inj 40 mg per ml, vial	149.37	3	SII-Onco-BCG

☞ **Restricted**

Initiation

For use in bladder cancer.

EVEROLIMUS – **Restricted** see terms below

☞ Tab 5 mg	4,555.76	30	Afinitor
☞ Tab 10 mg	6,512.29	30	Afinitor

☞ **Restricted**

Initiation

Neurologist or oncologist

Re-assessment required after 3 months

Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Continuation

Neurologist or oncologist

Re-assessment required after 12 months

All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MYCOPHENOLATE MOFETIL			
Tab 500 mg – 1% DV Nov-13 to 2016	25.00	50	CellCept
Cap 250 mg – 1% DV Nov-13 to 2016	25.00	100	CellCept
Powder for oral liq 1 g per 5 ml – 1% DV Nov-13 to 2016	187.25	165 ml	CellCept
Inj 500 mg vial – 1% DV Nov-13 to 2016	133.33	4	CellCept
PICIBANIL			
Inj 100 mg vial			
SIROLIMUS – Restricted see terms below			
⚡ Tab 1 mg	813.00	100	Rapamune
⚡ Tab 2 mg	1,626.00	100	Rapamune
⚡ Oral liq 1 mg per ml	487.80	60 ml	Rapamune

➡ **Restricted**

Initiation

For rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR < 30 ml/min; or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- HUS or TTP; or
- Leukoencephalopathy; or
- Significant malignant disease

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Antiallergy Preparations			
Allergic Emergencies			
ICATIBANT – Restricted see terms below			
⚡ Inj 10 mg per ml, 3 ml prefilled syringe	2,668.00	1	Firazyr
➡ Restricted			
Initiation			
Clinical immunologist or relevant specialist			
<i>Re-assessment required after 12 months</i>			
Both:			
1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and			
2 The patient has undergone product training and has agreed upon an action plan for self-administration.			
Continuation			
<i>Re-assessment required after 12 months</i>			
The treatment remains appropriate and the patient is benefiting from treatment.			
Allergy Desensitisation			
BEE VENOM – Restricted see terms below			
⚡ Maintenance kit - 6 vials 120 mcg freeze dried venom, with diluent			
⚡ Inj 550 mcg vial with diluent			
➡ Restricted			
Initiation			
Both:			
1 RAST or skin test positive; and			
2 Patient has had severe generalised reaction to the sensitising agent.			
PAPER WASP VENOM – Restricted see terms below			
⚡ Treatment kit - 6 vials 120 mcg freeze dried venom, with diluent			
⚡ Inj 550 mcg vial with diluent			
➡ Restricted			
Initiation			
Both:			
1 RAST or skin test positive; and			
2 Patient has had severe generalised reaction to the sensitising agent.			
YELLOW JACKET WASP VENOM – Restricted see terms below			
⚡ Treatment kit - 6 vials 120 mcg freeze dried venom, with diluent			
⚡ Inj 550 mcg vial with diluent			
➡ Restricted			
Initiation			
Both:			
1 RAST or skin test positive; and			
2 Patient has had severe generalised reaction to the sensitising agent.			
Allergy Prophylactics			
BECLOMETHASONE DIPROPIONATE			
Nasal spray 50 mcg per dose	4.85	200 dose	Alanase
Nasal spray 100 mcg per dose	5.75	200 dose	Alanase

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BUDESONIDE			
Nasal spray 50 mcg per dose	4.85	200 dose	Butacort Aqueous
Nasal spray 100 mcg per dose	5.75	200 dose	Butacort Aqueous
FLUTICASONE PROPIONATE			
Nasal spray 50 mcg per dose – 1% DV Sep-15 to 2018	2.18	120 dose	Flixonase Hayfever & Allergy
IPRATROPIUM BROMIDE			
Aqueous nasal spray 0.03% – 1% DV Jan-15 to 2017	3.95	15 ml	Univent
SODIUM CROMOGLYCATE			
Nasal spray 4%			

Antihistamines

CETIRIZINE HYDROCHLORIDE			
Tab 10 mg	1.59	100	Zetop
Oral liq 1 mg per ml – 1% DV Feb-15 to 2017	2.99	200 ml	Histaclear
CHLORPHENIRAMINE MALEATE			
Oral liq 0.4 mg per ml			
Inj 10 mg per ml, 1 ml ampoule			
CYPROHEPTADINE HYDROCHLORIDE			
Tab 4 mg			
FEXOFENADINE HYDROCHLORIDE			
Tab 60 mg			
Tab 120 mg			
Tab 180 mg			
LORATADINE			
Tab 10 mg – 1% DV Dec-13 to 2016	1.30	100	Lorafix
Oral liq 1 mg per ml – 1% DV Nov-14 to 2016	4.25	200 ml	LoraPaed
PROMETHAZINE HYDROCHLORIDE			
Tab 10 mg – 1% DV Sep-15 to 2018	1.78	50	Allersoothe
Tab 25 mg – 1% DV Sep-15 to 2018	1.99	50	Allersoothe
Oral liq 1 mg per ml – 1% DV Sep-15 to 2018	2.59	100 ml	Allersoothe
Inj 25 mg per ml, 2 ml ampoule	11.99	5	Hospira
TRIMEPRAZINE TARTRATE			
Oral liq 6 mg per ml			

Anticholinergic Agents

IPRATROPIUM BROMIDE			
Aerosol inhaler 20 mcg per dose			
Nebuliser soln 250 mcg per ml, 1 ml ampoule – 1% DV Sep-13 to 2016	3.26	20	Univent
Nebuliser soln 250 mcg per ml, 2 ml ampoule – 1% DV Sep-13 to 2016	3.37	20	Univent

Anticholinergic Agents with Beta-Adrenoceptor Agonists

SALBUTAMOL WITH IPRATROPIUM BROMIDE			
Aerosol inhaler 100 mcg with ipratropium bromide 20 mcg per dose			
Nebuliser soln 2.5 mg with ipratropium bromide 0.5 mg per 2.5 ml ampoule – 1% DV Sep-15 to 2018	3.59	20	Duolin

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Long-Acting Muscarinic Agents

GLYCOPYRRONIUM

Note: inhaled glycopyrronium treatment must not be used if the patient is also receiving treatment with subsidised tiotropium or umeclidinium.

Powder for inhalation 50 mcg per dose61.00 30 dose Seebri Breezhaler

TIOTROPIUM BROMIDE – **Restricted** see terms below

Note: tiotropium treatment must not be used if the patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.

⚡ Soln for inhalation 2.5 mcg per dose70.00 60 dose Spiriva Respimat

⚡ Powder for inhalation 18 mcg per dose70.00 30 dose Spiriva

➡**Restricted**

Initiation

All of the following:

- 1 To be used for the long-term maintenance treatment of bronchospasm and dyspnoea associated with COPD; and
- 2 In addition to standard treatment, the patient has trialled a short acting bronchodilator dose of at least 40 µg ipratropium q.i.d for one month; and
- 3 Either:
 - the patient's breathlessness according to the Medical Research Council (UK) dyspnoea scale is:
 - 3.1 Grade 3 (stops for breath after walking about 100 meters or after a few minutes on the level); or
 - 3.2 Grade 4 (too breathless to leave the house, or breathless when dressing or undressing); and
- 4 Actual FEV₁ as a % of predicted, must be below 60%; and
- 5 Either:
 - 5.1 Patient is not a smoker (for reporting purposes only); or
 - 5.2 Patient is a smoker and has been offered smoking cessation counselling; and
- 6 The patient has been offered annual influenza immunization.

UMECLIDINIUM

Note: Umeclidinium must not be used if the patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.

Powder for inhalation 62.5 mcg per dose61.50 30 dose Incruse Ellipta

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

➡**Restricted**

Initiation

Re-assessment required after 2 years

- Both:
- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
 - 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Continuation

Re-assessment required after 2 years

- Both:
- 1 Patient is compliant with the medication; and
 - 2 Patient has experienced improved COPD symptom control (prescriber determined).

Note: Combination long acting muscarinic antagonist and long acting beta-2 agonist must not be used if the patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

GLYCOPYRRONIUM WITH INDACATEROL – **Restricted** see terms above

⚡ Powder for Inhalation 50 mcg with indacaterol 110 mcg81.00 30 dose Ultibro Breezhaler

TIOTROPIUM BROMIDE WITH OLODATEROL – **Restricted** see terms above

⚡ Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg81.00 60 dose Spiolto Respimat

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
UMECLIDINIUM WITH VILANTEROL – Restricted see terms on the preceding page			
⬆ Powder for inhalation 62.5 mcg with vilanterol 25 mcg	77.00	30 dose	Anoro Ellipta

Beta-Adrenoceptor Agonists

SALBUTAMOL

Oral liq 400 mcg per ml – 1% DV Jan-14 to 2016.....	2.06	150 ml	Ventolin
Inj 500 mcg per ml, 1 ml ampoule			
Inj 1 mg per ml, 5 ml ampoule			
Aerosol inhaler, 100 mcg per dose	3.80	200 dose	SalAir
	4.00		Salamol
	6.00		Ventolin
Nebuliser soln 1 mg per ml, 2.5 ml ampoule – 1% DV Sep-15 to 2018	3.19	20	Asthalin
Nebuliser soln 2 mg per ml, 2.5 ml ampoule – 1% DV Sep-15 to 2018	3.29	20	Asthalin

TERBUTALINE SULPHATE

Powder for inhalation 250 mcg per dose			
Inj 0.5 mg per ml, 1 ml ampoule			

Cough Suppressants

PHOLCODINE

Oral liq 1 mg per ml			
----------------------	--	--	--

Decongestants

OXYMETAZOLINE HYDROCHLORIDE

Aqueous nasal spray 0.25 mg per ml			
Aqueous nasal spray 0.5 mg per ml			

PSEUDOEPHEDRINE HYDROCHLORIDE

Tab 60 mg			
-----------	--	--	--

SODIUM CHLORIDE

Aqueous nasal spray isotonic			
------------------------------	--	--	--

SODIUM CHLORIDE WITH SODIUM BICARBONATE

Soln for nasal irrigation			
---------------------------	--	--	--

XYLOMETAZOLINE HYDROCHLORIDE

Aqueous nasal spray 0.05%			
Aqueous nasal spray 0.1%			
Nasal drops 0.05%			
Nasal drops 0.1%			

Inhaled Corticosteroids

BECLOMETHASONE DIPROPIONATE

Aerosol inhaler 50 mcg per dose	8.54	200 dose	Beclazone 50
	9.30		Qvar
Aerosol inhaler 100 mcg per dose	12.50	200 dose	Beclazone 100
	15.50		Qvar
Aerosol inhaler 250 mcg per dose	22.67	200 dose	Beclazone 250

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BUDESONIDE			
Nebuliser soln 250 mcg per ml, 2 ml ampoule			
Nebuliser soln 500 mcg per ml, 2 ml ampoule			
Powder for inhalation 100 mcg per dose			
Powder for inhalation 200 mcg per dose			
Powder for inhalation 400 mcg per dose			
FLUTICASONE			
Aerosol inhaler 50 mcg per dose	7.50	120 dose	Flixotide Floair
Powder for inhalation 50 mcg per dose	8.67	60 dose	Flixotide Accuhaler
Powder for inhalation 100 mcg per dose	13.87	60 dose	Flixotide Accuhaler
Aerosol inhaler 125 mcg per dose	13.60	120 dose	Flixotide Floair
Aerosol inhaler 250 mcg per dose	27.20	120 dose	Flixotide Floair
Powder for inhalation 250 mcg per dose	24.51	60 dose	Flixotide Accuhaler
Leukotriene Receptor Antagonists			
MONTELUKAST – Restricted see terms below			
⚡ Tab 4 mg	18.48	28	Singulair
⚡ Tab 5 mg	18.48	28	Singulair
⚡ Tab 10 mg	18.48	28	Singulair
➡Restricted			
Initiation — Pre-school wheeze			
Both:			
1 To be used for the treatment of intermittent severe wheezing (possibly viral) in children under 5 years; and			
2 The patient has had at least three episodes in the previous 12 months of acute wheeze severe enough to seek medical attention.			
Initiation — Exercise-induced asthma			
All of the following:			
1 Patient has been trialed with maximal asthma therapy, including inhaled corticosteroids and long-acting beta-adrenoceptor agonists; and			
2 Patient continues to receive optimal inhaled corticosteroid therapy; and			
3 Patient continues to experience frequent episodes of exercise-induced bronchoconstriction.			
Initiation — Aspirin desensitisation			
Clinical immunologist or allergist			
All of the following:			
1 Patient is undergoing aspirin desensitisation therapy under the supervision of a clinical immunologist or allergist; and			
2 Patient has moderate to severe aspirin-exacerbated respiratory disease or Samter's triad; and			
3 Nasal polyposis, confirmed radiologically or surgically; and			
4 Documented aspirin or NSAID allergy confirmed by aspirin challenge or a clinical history of severe reaction to aspirin or NSAID where challenge would be considered dangerous.			
Long-Acting Beta-Adrenoceptor Agonists			
EFORMOTEROL FUMARATE			
Powder for inhalation 6 mcg per dose			
Powder for inhalation 12 mcg per dose			
INDACATEROL			
Powder for inhalation 150 mcg per dose	61.00	30 dose	Onbrez Breezhaler
Powder for inhalation 300 mcg per dose	61.00	30 dose	Onbrez Breezhaler

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

Inhaled Corticosteroids with Long-Acting Beta-Adrenoceptor Agonists

Powder for inhalation 100 mcg with eformoterol fumarate 6 mcg
Powder for inhalation 200 mcg with eformoterol fumarate 6 mcg
Powder for inhalation 400 mcg with eformoterol fumarate 12 mcg
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg

Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose	Breo Ellipta
--	-------	---------	--------------

Aerosol inhaler 50 mcg with salmeterol 25 mcg	37.48	120 dose	RexAir
	33.74		Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg	33.74	60 dose	Seretide Accuhaler
Aerosol inhaler 125 mcg with salmeterol 25 mcg	49.69	120 dose	RexAir
	44.08		Seretide
Powder for inhalation 250 mcg with salmeterol 50 mcg	44.08	60 dose	Seretide Accuhaler

Mast Cell Stabilisers

Aerosol inhaler 2 mg per dose

Powder for inhalation 20 mg per dose
Aerosol inhaler 5 mg per dose

Methylxanthines

Inj 25 mg per ml, 10 ml ampoule – 1% DV Oct-14 to 2017	118.25	5	DBL Aminophylline
--	--------	---	--------------------------

Oral liq 20 mg per ml (caffeine 10 mg per ml)	14.85	25 ml	Biomed
Inj 20 mg per ml (caffeine 10 mg per ml), 2.5 ml ampoule	55.75	5	Biomed

Tab long-acting 250 mg
Oral liq 80 mg per 15 ml

Mucolytics and Expectorants

↓ Nebuliser soln 2.5 mg per 2.5 ml ampoule	250.00	6	Pulmozyme
--	--------	---	-----------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➡Restricted

Initiation — cystic fibrosis

The patient has cystic fibrosis and has been approved by the Cystic Fibrosis Panel.

Initiation — significant mucus production

Limited to 4 weeks treatment

Both:

- 1 Patient is an in-patient; and
- 2 The mucus production cannot be cleared by first line chest techniques.

Initiation — pleural emphyema

Limited to 3 days treatment

Both:

- 1 Patient is an in-patient; and
- 2 Patient diagnoses with pleural emphyema.

SODIUM CHLORIDE

Nebuliser soln 7%, 90 ml bottle23.50 90 ml Biomed

Pulmonary Surfactants

BERACTANT

Soln 200 mg per 8 ml vial550.00 1 Survanta

PORACTANT ALFA

Soln 120 mg per 1.5 ml vial425.00 1 Curosurf

Soln 240 mg per 3 ml vial695.00 1 Curosurf

Respiratory Stimulants

DOXAPRAM

Inj 20 mg per ml, 5 ml vial

Sclerosing Agents

TALC

Powder

Soln (slurry) 100 mg per ml, 50 ml

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Anti-Infective Preparations			
Antibacterials			
CHLORAMPHENICOL			
Eye oint 1% – 1% DV Jul-16 to 2019	2.48	4 g	Chlorsig
Ear drops 0.5%			
Eye drops 0.5% – 1% DV Sep-15 to 2018	0.98	10 ml	Chlorafast
Eye drops 0.5%, single dose			
CIPROFLOXACIN			
Eye drops 0.3%			
FRAMYCETIN SULPHATE			
Ear/eye drops 0.5%			
FUSIDIC ACID			
Eye drops 1%	4.50	5 g	Fucithalmic
GENTAMICIN SULPHATE			
Eye drops 0.3%	11.40	5 ml	Genoptic
PROPAMIDINE ISETHIONATE			
Eye drops 0.1%			
SULPHACETAMIDE SODIUM			
Eye drops 10%			
TOBRAMYCIN			
Eye oint 0.3% – 1% DV Sep-14 to 2017	10.45	3.5 g	Tobrex
Eye drops 0.3% – 1% DV Sep-14 to 2017	11.48	5 ml	Tobrex
Antifungals			
NATAMYCIN			
Eye drops 5%			
Antivirals			
ACICLOVIR			
Eye oint 3%			
GANCICLOVIR			
Eye gel 0.15%			<i>e.g. Virgan</i>
Combination Preparations			
CIPROFLOXACIN WITH HYDROCORTISONE			
Ear drops ciprofloxacin 0.2% with 1% hydrocortisone – 1% DV Mar-15 to 2017	16.30	10 ml	Ciproxin HC Otic
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/eye drops 500 mcg with framycetin sulphate 5 mg and gramicidin 50 mcg per ml			
Products with Hospital Supply Status (HSS) are in bold Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE			
Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per g – 1% DV Sep-14 to 2017	5.39	3.5 g	Maxitrol
Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml – 1% DV Sep-14 to 2017	4.50	5 ml	Maxitrol
DEXAMETHASONE WITH TOBRAMYCIN			
Eye drops 0.1% with tobramycin 0.3% – 1% DV Mar-15 to 2017	12.64	5 ml	Tobradex
FLUMETASONE PIVALATE WITH CLIOQUINOL			
Ear drops 0.02% with clioquinol 1%			
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCIN AND NYSTATIN			
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml	Kenacomb

Anti-Inflammatory Preparations

Corticosteroids

DEXAMETHASONE			
Eye oint 0.1% – 1% DV Oct-14 to 2017	5.86	3.5 g	Maxidex
Eye drops 0.1% – 1% DV Oct-14 to 2017	4.50	5 ml	Maxidex
FLUOROMETHOLONE			
Eye drops 0.1% – 1% DV Sep-15 to 2018	3.09	5 ml	FML
PREDNISOLONE ACETATE			
Eye drops 0.12%			
Eye drops 1%			
PREDNISOLONE SODIUM PHOSPHATE			
Eye drops 0.5%, single dose (preservative free)	38.50	20 dose	Minims Prednisolone

Non-Steroidal Anti-Inflammatory Drugs

DICLOFENAC SODIUM			
Eye drops 0.1% – 1% DV Sep-14 to 2017	13.80	5 ml	Voltaren Ophtha
KETOROLAC TROMETAMOL			
Eye drops 0.5%			

Decongestants and Antiallergics

Antiallergic Preparations

LEVOCABASTINE			
Eye drops 0.05%			
LODOXAMIDE			
Eye drops 0.1% – 1% DV Sep-14 to 2017	8.71	10 ml	Lomide
OLOPATADINE			
Eye drops 0.1%	17.00	5 ml	Patanol
SODIUM CROMOGLYCATE			
Eye drops 2%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Decongestants			
NAPHAZOLINE HYDROCHLORIDE			
Eye drops 0.1% – 1% DV Sep-14 to 2017	4.15	15 ml	Naphcon Forte
Diagnostic and Surgical Preparations			
Diagnostic Dyes			
FLUORESCIEIN SODIUM			
Eye drops 2%, single dose			
Inj 10%, 5 ml vial	125.00	12	Fluorescite
Ophthalmic strips 1 mg			
FLUORESCIEIN SODIUM WITH LIGNOCAINE HYDROCHLORIDE			
Eye drops 0.25% with lignocaine hydrochloride 4%, single dose			
LISSAMINE GREEN			
Ophthalmic strips 1.5 mg			
ROSE BENGAL SODIUM			
Ophthalmic strips 1%			
Irrigation Solutions			
MIXED SALT SOLUTION FOR EYE IRRIGATION			
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 15 ml dropper bottle – 1% DV Jan-16 to 2018	5.00	15 ml	Balanced Salt Solution
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 250 ml			<i>e.g. Balanced Salt Solution</i>
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 500 ml bottle – 1% DV Jan-16 to 2018	10.50	500 ml	Balanced Salt Solution
Ocular Anaesthetics			
OXYBUPROCAINE HYDROCHLORIDE			
Eye drops 0.4%, single dose			
PROXYMETACAINE HYDROCHLORIDE			
Eye drops 0.5%			
TETRACAINE [AMETHOCAINE] HYDROCHLORIDE			
Eye drops 0.5%, single dose			
Eye drops 1%, single dose			
Viscoelastic Substances			
HYPROMELLOSE			
Inj 2%, 1 ml syringe			
Inj 2%, 2 ml syringe			

SENSORY ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SODIUM HYALURONATE [HYALURONIC ACID]			
Inj 14 mg per ml, 0.85 ml syringe	50.00	1	Healon GV
Inj 14 mg per ml, 0.55 ml syringe	50.00	1	Healon GV
Inj 23 mg per ml, 0.6 ml syringe			
Inj 10 mg per ml, 0.85 ml syringe	30.00	1	Provisc
SODIUM HYALURONATE [HYALURONIC ACID] WITH CHONDROITIN SULPHATE			
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.35 ml syringe and inj 10 mg sodium hyaluronate [hyaluronic acid] per ml, 0.4 ml syringe	64.00	1	Duovisc
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.5 ml syringe and inj 10 mg sodium hyaluronate [hyaluronic acid] per ml, 0.55 ml syringe	74.00	1	Duovisc
Inj 30 mg with chondroitin sulphate 40 mg per ml, 0.75 ml syringe			

Other

DISODIUM EDETATE			
Inj 150 mg per ml, 20 ml ampoule			
Inj 150 mg per ml, 20 ml vial			
Inj 150 mg per ml, 100 ml vial			
RIBOFLAVIN 5-PHOSPHATE			
Soln trans epithelial riboflavin			
Inj 0.1%			
Inj 0.1% plus 20% dextran T500			

Glaucoma Preparations

Beta Blockers

BETAXOLOL			
Eye drops 0.25% – 1% DV Sep-14 to 2017	11.80	5 ml	Betoptic S
Eye drops 0.5% – 1% DV Sep-14 to 2017	7.50	5 ml	Betoptic
LEVOBUNOLOL HYDROCHLORIDE			
Eye drops 0.5%	7.00	5 ml	Betagan
TIMOLOL			
Eye drops 0.25% – 1% DV Sep-14 to 2017	1.45	5 ml	Arrow-Timolol
Eye drops 0.25%, gel forming – 1% DV Mar-14 to 2016	3.30	2.5 ml	Timoptol XE
Eye drops 0.5% – 1% DV Sep-14 to 2017	1.45	5 ml	Arrow-Timolol
Eye drops 0.5%, gel forming – 1% DV Mar-14 to 2016	3.78	2.5 ml	Timoptol XE

Carbonic Anhydrase Inhibitors

ACETAZOLAMIDE			
Tab 250 mg – 1% DV Sep-14 to 2017	17.03	100	Diamox
Inj 500 mg			
BRINZOLAMIDE			
Eye drops 1%			
DORZOLAMIDE			
Eye drops 2%			
DORZOLAMIDE WITH TIMOLOL			
Eye drops 2% with timolol 0.5% – 1% DV Dec-15 to 2018	3.45	5 ml	Arrow-Dortim

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Miotics			
ACETYLCHOLINE CHLORIDE			
Inj 20 mg vial with diluent			
PILOCARPINE HYDROCHLORIDE			
Eye drops 1% – 1% DV Sep-14 to 2017	4.26	15 ml	Isopto Carpine
Eye drops 2% – 1% DV Sep-14 to 2017	5.35	15 ml	Isopto Carpine
Eye drops 2%, single dose			
Eye drops 4% – 1% DV Sep-14 to 2017	7.99	15 ml	Isopto Carpine
Prostaglandin Analogues			
BIMATOPROST			
Eye drops 0.03% – 1% DV Jul-16 to 2018	3.65	3 ml	Bimatoprost Actavis
LATANOPROST			
Eye drops 0.005% – 1% DV Sep-15 to 2018	1.50	2.5 ml	Hysite
TRAVOPROST			
Eye drops 0.004%			
Sympathomimetics			
APRACLOPIDINE			
Eye drops 0.5% – 1% DV Mar-15 to 2017	19.77	5 ml	Iopidine
BRIMONIDINE TARTRATE			
Eye drops 0.2% – 1% DV Sep-14 to 2017	4.32	5 ml	Arrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL			
Eye drops 0.2% with timolol 0.5%			
Mydriatics and Cycloplegics			
Anticholinergic Agents			
ATROPINE SULPHATE			
Eye drops 0.5%			
Eye drops 1%, single dose			
Eye drops 1% – 1% DV Jul-14 to 2017	17.36	15 ml	Atropt
CYCLOPENTOLATE HYDROCHLORIDE			
Eye drops 0.5%, single dose			
Eye drops 1% – 1% DV Sep-14 to 2017	8.76	15 ml	Cyclogyl
Eye drops 1%, single dose			
TROPICAMIDE			
Eye drops 0.5% – 1% DV Oct-14 to 2017	7.15	15 ml	Mydriacyl
Eye drops 0.5%, single dose			
Eye drops 1% – 1% DV Oct-14 to 2017	8.66	15 ml	Mydriacyl
Eye drops 1%, single dose			
Sympathomimetics			
PHENYLEPHRINE HYDROCHLORIDE			
Eye drops 2.5%, single dose			
Eye drops 10%, single dose			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Ocular Lubricants			
CARBOMER			
Ophthalmic gel 0.3%, single dose	8.25	30	Poly Gel
Ophthalmic gel 0.2%			
CARMELLOSE SODIUM			
Eye drops 0.5%			
Eye drops 0.5%, single dose			
Eye drops 1%			
Eye drops 1%, single dose			
HYPROMELLOSE			
Eye drops 0.5%	3.92	15 ml	Methopt
HYPROMELLOSE WITH DEXTRAN			
Eye drops 0.3% with dextran 0.1%	2.30	15 ml	Poly-Tears
Eye drops 0.3% with dextran 0.1%, single dose			
MACROGOL 400 AND PROPYLENE GLYCOL			
Eye drops 0.4% with propylene glycol 0.3% preservative free, single dose	4.30	24	Systane Unit Dose
PARAFFIN LIQUID WITH SOFT WHITE PARAFFIN			
Eye oint 42.5% with soft white paraffin 57.3%			
PARAFFIN LIQUID WITH WOOL FAT			
Eye oint 3% with wool fat 3% – 1% DV Jul-14 to 2017	3.63	3.5 g	Poly-Visc
POLYVINYL ALCOHOL			
Eye drops 1.4% – 1% DV Jun-16 to 2019	2.62	15 ml	Vistil
Eye drops 3% – 1% DV Jun-16 to 2019	3.68	15 ml	Vistil Forte
POLYVINYL ALCOHOL WITH POVIDONE			
Eye drops 1.4% with povidone 0.6%, single dose			
RETINOL PALMITATE			
Oint 138 mcg per g	3.80	5 g	VitA-POS
SODIUM HYALURONATE [HYALURONIC ACID]			
Eye drops 1 mg per ml	22.00	10 ml	Hylo-Fresh

Other Otological Preparations

ACETIC ACID WITH PROPYLENE GLYCOL			
Ear drops 2.3% with propylene glycol 2.8%			
DOCUSATE SODIUM			
Ear drops 0.5%			

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Agents Used in the Treatment of Poisonings

Antidotes

ACETYLCYSTEINE

Tab eff 200 mg

Inj 200 mg per ml, 10 ml ampoule – **1% DV Sep-15 to 2018** 78.34 10 **DBL Acetylcysteine**

DIGOXIN IMMUNE FAB

Inj 38 mg vial

Inj 40 mg vial

ETHANOL

Liq 96%

ETHANOL WITH GLUCOSE

Inj 10% with glucose 5%, 500 ml bottle

ETHANOL, DEHYDRATED

Inj 100%, 5 ml ampoule

Inj 96%

FLUMAZENIL

Inj 0.1 mg per ml, 5 ml ampoule – **1% DV Sep-15 to 2018** 85.05 5 **Anexate**

HYDROXOCOBALAMIN

Inj 5 g vial

Inj 2.5 g vial

NALOXONE HYDROCHLORIDE

Inj 400 mcg per ml, 1 ml ampoule 48.84 5 Hospira

PRALIDOXIME IODIDE

Inj 25 mg per ml, 20 ml ampoule

SODIUM NITRITE

Inj 30 mg per ml, 10 ml ampoule

SODIUM THIOSULFATE

Inj 500 mg per ml, 20 ml ampoule

Inj 250 mg per ml, 10 ml vial

Inj 500 mg per ml, 10 ml vial

SOYA OIL

Inj 20%, 500 ml bag

Inj 20%, 500 ml bottle

Antitoxins

BOTULISM ANTITOXIN

Inj 250 ml vial

DIPHThERIA ANTITOXIN

Inj 10,000 iu vial

Antivenoms

RED BACK SPIDER ANTIVENOM

Inj 500 u vial

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SNAKE ANTIVENOM			
Inj 50 ml vial			

Removal and Elimination

CHARCOAL

Oral liq 200 mg per ml	43.50	250 ml	Carbasorb-X
------------------------------	-------	--------	-------------

DEFERASIROX – **Restricted** see terms below

☞ Tab 125 mg dispersible	276.00	28	Exjade
☞ Tab 250 mg dispersible	552.00	28	Exjade
☞ Tab 500 mg dispersible	1,105.00	28	Exjade

☞ **Restricted**

Initiation

Haematologist

Re-assessment required after 2 years

All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 - 1.0 cells per μL).

Continuation

Haematologist

Re-assessment required after 2 years

Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels. .

DEFERIPRONE – **Restricted** see terms below

☞ Tab 500 mg	533.17	100	Ferriprox
☞ Oral liq 100 mg per ml	266.59	250 ml	Ferriprox

☞ **Restricted**

Initiation

Patient has been diagnosed with chronic iron overload due to congenital inherited anaemia or acquired red cell aplasia.

DESFERRIOXAMINE MESILATE

Inj 500 mg vial – 1% DV Feb-16 to 2018	51.52	10	Desferal
---	-------	----	-----------------

DICOBALT EDETATE

Inj 15 mg per ml, 20 ml ampoule

DIMERCAPROL

Inj 50 mg per ml, 2 ml ampoule

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DIMERCAPTOSUCCINIC ACID			
Cap 100 mg			<i>e.g. Pharmaceutical Compounding NZ</i>
Cap 200 mg			<i>e.g. Pharmaceutical Compounding NZ</i>
SODIUM CALCIUM EDETATE			
Inj 200 mg per ml, 2.5 ml ampoule			
Inj 200 mg per ml, 5 ml ampoule			
Antiseptics and Disinfectants			
CHLORHEXIDINE			
Soln 4%	1.86	50 ml	healthE
Soln 5%	15.50	500 ml	healthE
CHLORHEXIDINE WITH CETRIMIDE			
Crm 0.1% with cetrimide 0.5%			
Foaming soln 0.5% with cetrimide 0.5%			
CHLORHEXIDINE WITH ETHANOL			
Soln 0.5% with ethanol 70%, non-staining (pink) 100 ml	2.65	1	healthE
Soln 2% with ethanol 70%, non-staining (pink) 100 ml	3.54	1	healthE
Soln 0.5% with ethanol 70%, non-staining (pink) 25 ml	1.55	1	healthE
Soln 0.5% with ethanol 70%, staining (red) 100 ml	2.90	1	healthE
Soln 2% with ethanol 70%, staining (red) 100 ml	3.86	1	healthE
Soln 0.5% with ethanol 70%, non-staining (pink) 500 ml	5.45	1	healthE
Soln 0.5% with ethanol 70%, staining (red) 500 ml	5.90	1	healthE
Soln 2% with ethanol 70%, staining (red) 500 ml	9.56	1	healthE
IODINE WITH ETHANOL			
Soln 1% with ethanol 70%, 100 ml	9.30	1	healthE
ISOPROPYL ALCOHOL			
Soln 70%, 500 ml	5.00	1	PSM
	5.65		healthE
<i>(PSM Soln 70%, 500 ml to be delisted 1 July 2016)</i>			
POVIDONE-IODINE			
📌 Vaginal tab 200 mg			
➡ Restricted			
Initiation			
Rectal administration pre-prostate biopsy.			
Oint 10%	3.27	25 g	Betadine
Soln 10%	6.20	500 ml	Betadine
	2.95	100 ml	Riodine
	6.20	500 ml	Riodine
Soln 5%			
Soln 7.5%			
Pad 10%			
Swab set 10%			
POVIDONE-IODINE WITH ETHANOL			
Soln 10% with ethanol 30%	10.00	500 ml	Betadine Skin Prep
Soln 10% with ethanol 70%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SODIUM HYPOCHLORITE			
Soln			
Contrast Media			
Iodinated X-ray Contrast Media			
DIATRIZOATE MEGLUMINE WITH SODIUM AMIDOTRIZOATE			
Oral liq 660 mg per ml with sodium amidotrizoate 100 mg per ml, 100 ml bottle	22.50	100 ml	Gastrografin
Inj 260 mg with sodium amidotrizoate 40 mg per ml, 250 ml bottle	80.00	1	Urografin
DIATRIZOATE SODIUM			
Oral liq 370 mg per ml, 10 ml sachet	156.12	50	Ioscan
IODISED OIL			
Inj 38% w/w (480 mg per ml), 10 ml ampoule	191.00	1	Lipiodol Ultra Fluid
IODIXANOL			
Inj 270 mg per ml (iodine equivalent), 50 ml bottle – 5% DV Sep-14 to 2017	220.00	10	Visipaque
Inj 270 mg per ml (iodine equivalent), 100 ml bottle – 5% DV Sep-14 to 2017	430.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 50 ml bottle – 5% DV Sep-14 to 2017	220.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 100 ml bottle – 5% DV Sep-14 to 2017	430.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 200 ml bottle – 5% DV Sep-14 to 2017	850.00	10	Visipaque
IOHEXOL			
Inj 240 mg per ml (iodine equivalent), 50 ml bottle – 5% DV Sep-14 to 2017	75.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 20 ml bottle – 5% DV Sep-14 to 2017	57.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 50 ml bottle – 5% DV Sep-14 to 2017	75.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 100 ml bottle – 5% DV Sep-14 to 2017	150.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 20 ml bottle – 5% DV Sep-14 to 2017	59.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 50 ml bottle – 5% DV Sep-14 to 2017	75.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 75 ml bottle – 5% DV Sep-14 to 2017	114.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 100 ml bottle – 5% DV Sep-14 to 2017	150.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 200 ml bottle – 5% DV Sep-14 to 2017	290.00	10	Omnipaque

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Non-iodinated X-ray Contrast Media			
BARIUM SULPHATE			
Powder for oral liq 20 mg per g (2% w/w), 22.1 g sachet	507.50	50	E-Z-Cat Dry
Oral liq 400 mg per ml (40% w/v, 30% w/w), bottle	17.39	148 g	Varibar - Thin Liquid
Oral liq 600 mg per g (60% w/w), tube	36.51	454 g	E-Z-Paste
Oral liq 400 mg per ml (40% w/v), bottle	155.35	250 ml	Varibar - Honey
	38.40	240 ml	Varibar - Nectar
	145.04	230 ml	Varibar - Pudding
Enema 1,250 mg per ml (125% w/v), 500 ml bag	282.30	12	Liquibar
Oral liq 22 mg per g (2.2% w/w), 250 ml bottle	175.00	24	CT Plus+
Oral liq 22 mg per g (2.2% w/w), 450 ml bottle	220.00	24	CT Plus+
Oral liq 1 mg per ml (0.1% w/v, 0.1% w/w), 450 ml bottle	441.12	24	VoLumen
Oral liq 20.9 mg per ml (2.1% w/v, 2% w/w), 250 ml bottle	140.94	24	Readi-CAT 2
Powder for oral soln 97.65% w/w, 300 g bottle	237.76	24	X-Opaque-HD
Oral liq 400 mg per ml (40% w/v, 30% w/w), 20 ml bottle	52.35	3	Tagitol V
Oral liq 1,250 mg per ml (125% w/v), 2,000 ml bottle	91.77	1	Liquibar
BARIUM SULPHATE WITH SODIUM BICARBONATE			
Grans eff 382.2 mg per g with sodium bicarbonate 551.3 mg per g, 4 g sachet	102.93	50	E-Z-Gas II
CITRIC ACID WITH SODIUM BICARBONATE			
Powder 382.2 mg per g with sodium bicarbonate 551.3 mg per g, 4 g sachet			<i>e.g. E-Z-GAS II</i>
Paramagnetic Contrast Media			
GADOBENIC ACID			
Inj 334 mg per ml, 10 ml vial	324.74	10	Multihance
Inj 334 mg per ml, 20 ml vial	636.28	10	Multihance
GADOBUTROL			
Inj 1 mmol per ml, 15 ml vial			
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 7.5 ml prefilled syringe	180.00	5	Gadovist
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 15 ml prefilled syringe	700.00	10	Gadovist
GADODIAMIDE			
Inj 287 mg per ml, 10 ml prefilled syringe	200.00	10	Omniscan
Inj 287 mg per ml, 10 ml vial	170.00	10	Omniscan
Inj 287 mg per ml, 5 ml vial	120.00	10	Omniscan
Inj 287 mg per ml, 15 ml prefilled syringe	320.00	10	Omniscan
GADOTERIC ACID			
Inj 279.32 mg per ml (0.5 mmol per ml), 10 ml prefilled syringe	24.50	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 15 ml bottle	34.50	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 15 ml prefilled syringe	41.00	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 20 ml prefilled syringe	55.00	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 10 ml bottle	23.20	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 20 ml bottle	46.30	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 5 ml bottle	12.30	1	Dotarem

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GADOXETATE DISODIUM			
Inj 181.43 mg per ml (equivalent to 0.25 mmol per ml), 10 ml prefilled syringe	300.00	1	Primovist
MEGLUMINE GADOPENTETATE			
Inj 469 mg per ml, 10 ml prefilled syringe	95.00	5	Magnevist
Inj 469 mg per ml, 10 ml vial	185.00	10	Magnevist
MEGLUMINE IOTROXATE			
Inj 105 mg per ml, 100 ml bottle	150.00	100 ml	Biliscopin

Ultrasound Contrast Media

PERFLUTREN			
Inj 1.1 mg per ml, 1.5 ml vial – 5% DV Sep-14 to 2017	180.00	1	Definity
	720.00	4	Definity

Diagnostic Agents

ARGININE			
Inj 50 mg per ml, 500 ml bottle			
Inj 100 mg per ml, 300 ml bottle			
HISTAMINE ACID PHOSPHATE			
Nebuliser soln 0.6%, 10 ml vial			
Nebuliser soln 2.5%, 10 ml vial			
Nebuliser soln 5%, 10 ml vial			
MANNITOL			
Powder for inhalation			<i>e.g. Aridol</i>
METHACHOLINE CHLORIDE			
Powder 100 mg			
SECRETIN PENTAHYDROCHLORIDE			
Inj 100 u ampoule			
SINCALIDE			
Inj 5 mcg per vial			
TUBERCULIN, PURIFIED PROTEIN DERIVATIVE			
Inj 5 TU per 0.1 ml, 1 ml vial			

Diagnostic Dyes

BONNEY'S BLUE DYE			
Soln			
INDIGO CARMINE			
Inj 4 mg per ml, 5 ml ampoule			
Inj 8 mg per ml, 5 ml ampoule			
INDOCYANINE GREEN			
Inj 25 mg vial			
METHYLTHIONINIUM CHLORIDE [METHYLENE BLUE]			
Inj 10 mg per ml, 10 ml ampoule			
Inj 10 mg per ml, 5 ml ampoule			
PATENT BLUE V			
Inj 2.5%, 2 ml ampoule	440.00	5	Obex Medical

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Irrigation Solutions			
CHLORHEXIDINE			
Irrigation soln 0.02%, bottle	2.92	100 ml	Baxter
Irrigation soln 0.05%, bottle	3.02	100 ml	Baxter
	3.63	500 ml	Baxter
Irrigation soln 0.1%, bottle	3.10	100 ml	Baxter
Irrigation soln 0.5%, bottle	4.69	500 ml	Baxter
Irrigation soln 0.02%, 500 ml bottle			
Irrigation soln 0.1%, 30 ml ampoule			
CHLORHEXIDINE WITH CETRIMIDE			
Irrigation soln 0.015% with cetrimide 0.15%, 30 ml ampoule			
Irrigation soln 0.015% with cetrimide 0.15%, bottle	3.21	100 ml	Baxter
	3.47	500 ml	Baxter
	4.17	1,000 ml	Baxter
Irrigation soln 0.05% with cetrimide 0.5%, bottle	3.87	500 ml	Baxter
	4.20	100 ml	Baxter
Irrigation soln 0.1% with cetrimide 1%, bottle	4.38	100 ml	Baxter
	5.81	500 ml	Baxter
GLYCINE			
Irrigation soln 1.5%, bottle	11.38	2,000 ml	Baxter
	14.44	3,000 ml	Baxter
SODIUM CHLORIDE			
Irrigation soln 0.9%, 30 ml ampoule	19.50	30 ml	Pfizer
Irrigation soln 0.9%, bottle	2.49	100 ml	Baxter
	2.88	500 ml	Baxter
	2.96	1,000 ml	Baxter
	10.00	2,000 ml	Baxter
	12.67	3,000 ml	Baxter
WATER			
Irrigation soln, bottle	2.61	500 ml	Baxter
	2.68	100 ml	Baxter
	2.75	1,000 ml	Baxter
	9.71	2,000 ml	Baxter
	15.80	3,000 ml	Baxter

Surgical Preparations

BISMUTH SUBNITRATE AND IODOFORM PARAFFIN

Paste

DIMETHYL SULFOXIDE

Soln 50%

Soln 99%

PHENOL

Inj 6%, 10 ml ampoule

PHENOL WITH IOXAGLIC ACID

Inj 12%, 10 ml ampoule

TROMETAMOL

Inj 36 mg per ml, 500 ml bottle

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Cardioplegia Solutions			
ELECTROLYTES			
Inj 15 mmol/l sodium chloride, 9 mmol/l potassium chloride, 1 mmol/l potassium hydrogen 2-ketoglutarate, 4 mmol/l magnesium chloride, 18 mmol/l histidine hydrochloride, 180 mmol/l histidine, 2 mmol/l tryptophan, 30 mmol/l mannitol, 0.015 mmol/l calcium chloride, 1,000 ml bag			<i>e.g. Custodiol-HTK</i>
Inj aspartic acid 10.43 mg per ml, citric acid 0.22476 mg per ml, glutamic acid 11.53 mg per ml, sodium phosphate 0.1725 mg per ml, potassium chloride 2.15211 mg per ml, sodium citrate 1.80768 mg per ml, sodium hydroxide 6.31 mg per ml and trometamol 11.2369 mg per ml, 364 ml bag			<i>e.g. Cardioplegia Enriched Paed. Soln.</i>
Inj aspartic acid 8.481 mg per ml, citric acid 0.8188 mg per ml, glutamic acid 9.375 mg per ml, sodium phosphate 0.6285 mg per ml, potassium chloride 2.5 mg per ml, sodium citrate 6.585 mg per ml, sodium hydroxide 5.133 mg per ml and trometamol 9.097 mg per ml, 527 ml bag			<i>e.g. Cardioplegia Enriched Solution</i>
Inj citric acid 0.07973 mg per ml, sodium phosphate 0.06119 mg per ml, potassium chloride 2.181 mg per ml, sodium chloride 1.788 mg ml, sodium citrate 0.6412 mg per ml and trometamol 5.9 mg per ml, 523 ml bag			<i>e.g. Cardioplegia Base Solution</i>
Inj 110 mmol/l sodium, 16 mmol/l potassium, 1.2 mmol/l calcium, 16 mmol/l magnesium and 160 mmol/l chloride, 1,000 ml bag			<i>e.g. Cardioplegia Solution AHB7832</i>
Inj 143 mmol/l sodium, 16 mmol/l potassium, 16 mmol/l magnesium and 1.2 mmol/l calcium, 1,000 ml bag			<i>e.g. Cardioplegia Electrolyte Solution</i>

MONOSODIUM GLUTAMATE WITH SODIUM ASPARTATE

Inj 42.68 mg with sodium aspartate 39.48 mg per ml, 250 ml bottle

MONOSODIUM L-ASPARTATE

Inj 14 mmol per 10 ml, 10 ml

Cold Storage Solutions
SODIUM WITH POTASSIUM

Inj 29 mmol/l with potassium 125 mmol/l, 1,000 ml bag

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer Per
Extemporaneously Compounded Preparations		
ACETIC ACID Liq		
ALUM Powder BP		
ARACHIS OIL [PEANUT OIL] Liq		
ASCORBIC ACID Powder		
BENZOIN Tincture compound BP		
BISMUTH SUBGALLATE Powder		
BORIC ACID Powder		
CARBOXYMETHYLCELLULOSE Soln 1.5%		
CETRIMIDE Soln 40%		
CHLORHEXIDINE GLUCONATE Soln 20 %		
CHLOROFORM Liq BP		
CITRIC ACID Powder BP		
CLOVE OIL Liq		
COAL TAR Soln BP		
CODEINE PHOSPHATE Powder		
COLLODION FLEXIBLE Liq		
COMPOUND HYDROXYBENZOATE Soln		
CYSTEAMINE HYDROCHLORIDE Powder		
DISODIUM HYDROGEN PHOSPHATE WITH SODIUM DIHYDROGEN PHOSPHATE Inj 37.46 mg with sodium dihydrogen phosphate 47.7 mg in 1.5 ml ampoule		
DITHRANOL Powder		

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GLUCOSE [DEXTROSE] Powder			
GLYCERIN WITH SODIUM SACCHARIN Suspension	32.50	473 ml	Ora-Sweet SF
GLYCERIN WITH SUCROSE Suspension	32.50	473 ml	Ora-Sweet
GLYCEROL Liq	19.80	2,000 ml	ABM
HYDROCORTISONE Powder – 1% DV Dec-14 to 2017	59.50	25 g	ABM
LACTOSE Powder			
MAGNESIUM HYDROXIDE Paste			
MENTHOL Crystals			
METHADONE HYDROCHLORIDE Powder			
METHYL HYDROXYBENZOATE Powder			
METHYLCELLULOSE Powder			
Suspension	32.50	473 ml	Ora-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN Suspension	32.50	473 ml	Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE Suspension	32.50	473 ml	Ora-Blend
OLIVE OIL Liq			
PARAFFIN Liq			
PHENOBARBITONE SODIUM Powder			
PHENOL Liq			
PILOCARPINE NITRATE Powder			
POLYHEXAMETHYLENE BIGUANIDE Liq			
POVIDONE K30 Powder			
PROPYLENE GLYCOL Liq	12.00	500 ml	ABM

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SALICYLIC ACID Powder			
SILVER NITRATE Crystals			
SODIUM BICARBONATE Powder BP			
SODIUM CITRATE Powder			
SODIUM METABISULFITE Powder			
STARCH Powder			
SULPHUR Precipitated Sublimed			
SYRUP Liq (pharmaceutical grade)	21.75	2,000 ml	Midwest
THEOBROMA OIL Oint			
TRI-SODIUM CITRATE Crystals			
TRICHLORACETIC ACID Grans			
UREA Powder BP			
WOOL FAT Oint, anhydrous			
XANTHAN Gum 1%			
ZINC OXIDE Powder			

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Food Modules

Carbohydrate

➡ Restricted

Initiation — Use as an additive

Any of the following:

- 1 Cystic fibrosis; or
- 2 Chronic kidney disease; or
- 3 Cancer in children; or
- 4 Cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 5 Faltering growth in an infant/child; or
- 6 Bronchopulmonary dysplasia; or
- 7 Premature and post premature infant; or
- 8 Inborn errors of metabolism.

Initiation — Use as a module

For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

CARBOHYDRATE SUPPLEMENT – **Restricted** see terms above

⬆ Powder 95 g carbohydrate per 100 g, 368 g can

⬆ Powder 96 g carbohydrate per 100 g, 400 g can

e.g. Polycal

Fat

➡ Restricted

Initiation — Use as an additive

Any of the following:

- 1 Patient has inborn errors of metabolism; or
- 2 Faltering growth in an infant/child; or
- 3 Bronchopulmonary dysplasia; or
- 4 Fat malabsorption; or
- 5 Lymphangiectasia; or
- 6 Short bowel syndrome; or
- 7 Infants with necrotising enterocolitis; or
- 8 Biliary atresia; or
- 9 For use in a ketogenic diet; or
- 10 Chyle leak; or
- 11 Ascites; or
- 12 Patient has increased energy requirements, and for whom dietary measures have not been successful.

Initiation — Use as a module

For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

LONG-CHAIN TRIGLYCERIDE SUPPLEMENT – **Restricted** see terms above

⬆ Liquid 50 g fat per 100 ml, 200 ml bottle

e.g. Calogen

⬆ Liquid 50 g fat per 100 ml, 500 ml bottle

e.g. Calogen

MEDIUM-CHAIN TRIGLYCERIDE SUPPLEMENT – **Restricted** see terms above

⬆ Liquid 50 g fat per 100 ml, 250 ml bottle

e.g. Liquigen

⬆ Liquid 95 g fat per 100 ml, 500 ml bottle

e.g. MCT Oil

WALNUT OIL – **Restricted** see terms above

⬆ Liq

⬆ Item restricted (see ➡ above); ⬇ Item restricted (see ➡ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Protein

➔Restricted

Initiation — Use as an additive

Either:

- 1 Protein losing enteropathy; or
- 2 High protein needs.

Initiation — Use as a module

For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk. .

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

PROTEIN SUPPLEMENT – **Restricted** see terms above

⬆ Powder 5 g protein, 0.67 g carbohydrate and 0.6 g fat per 6.6 g, 275 g can			<i>e.g. Promod</i>
⬆ Powder 6 g protein per 7 g, can	8.95	227 g	Resource Beneprotein
⬆ Powder 89 g protein, <1.5 g carbohydrate and 2 g fat per 100 g, 225 g can			<i>e.g. Protifar</i>

Other Supplements

BREAST MILK FORTIFIER

Powder 0.2 g protein, 0.7 g carbohydrate and 0.02 g fat per 1 g sachet	<i>e.g. FM 85</i>
Powder 0.5 g protein, 1.2 g carbohydrate and 0.08 g fat per 2 g sachet	<i>e.g. S26 Human Milk Fortifier</i>
Powder 0.6 g protein and 1.4 g carbohydrate per 2.2 g sachet	<i>e.g. Nutricia Breast Milk Fortifier</i>

CARBOHYDRATE AND FAT SUPPLEMENT – **Restricted** see terms below

⬆ Powder 72.7 g carbohydrate and 22.3 g fat per 100 g, 400 g can	<i>e.g. Super Soluble Duocal</i>
--	----------------------------------

➔Restricted

Initiation

Both:

- 1 Infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 Cystic fibrosis; or
 - 2.2 Cancer in children; or
 - 2.3 Faltering growth; or
 - 2.4 Bronchopulmonary dysplasia; or
 - 2.5 Premature and post premature infants.

NOTE:

While pre-thickened drinks and supplements have not been included in Section H, DHB hospitals may continue to use such products for patients with dysphagia, provided that:

- use was established prior to 1 July 2013; and
- the product has not been specifically considered and excluded by PHARMAC; and
- use of the product conforms to any applicable indication restrictions for similar products that are listed in Section H (for example, use of thickened high protein products should be in line with the restriction for high protein oral feed in Section H).

PHARMAC intends to make a further decision in relation to pre-thickened drinks and supplements in the future, and will notify of any change to this situation.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Food/Fluid Thickeners			
CAROB BEAN GUM WITH MAIZE STARCH AND MALTODEXTRIN Powder			<i>e.g. Feed Thickener Karicare Aptamil</i>
GUAR GUM Powder			<i>e.g. Guarcol</i>
MAIZE STARCH Powder			<i>e.g. Resource Thicken Up; Nutilis</i>
MALTODEXTRIN WITH XANTHAN GUM Powder			<i>e.g. Instant Thick</i>
MALTODEXTRIN WITH XANTHAN GUM AND ASCORBIC ACID Powder			<i>e.g. Easy Thick</i>

Metabolic Products

➡ Restricted

Initiation

Any of the following:

- 1 For the dietary management of homocystinuria, maple syrup urine disease, phenylketonuria (PKU), glutaric aciduria, iso-valeric acidemia, propionic acidemia, methylmalonic acidemia, tyrosinaemia or urea cycle disorders; or
- 2 Patient has adrenoleukodystrophy; or
- 3 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Glutaric Aciduria Type 1 Products

AMINO ACID FORMULA (WITHOUT LYSINE AND LOW TRYPTOPHAN) – **Restricted** see terms above

- ⚡ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre
per 100 g, 400 g can *e.g. GA1 Anamix Infant*
- ⚡ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can *e.g. XLYS Low TRY
Maxamaid*

Homocystinuria Products

AMINO ACID FORMULA (WITHOUT METHIONINE) – **Restricted** see terms above

- ⚡ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre
per 100 g, 400 g can *e.g. HCU Anamix Infant*
- ⚡ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can *e.g. XMET Maxamaid*
- ⚡ Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can *e.g. XMET Maxamum*
- ⚡ Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per
100 ml, 125 ml bottle *e.g. HCU Anamix Junior
LQ*

Isovaleric Acidemia Products

AMINO ACID FORMULA (WITHOUT LEUCINE) – **Restricted** see terms above

- ⚡ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre
per 100 g, 400 g can *e.g. IVA Anamix Infant*
- ⚡ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can *e.g. XLEU Maxamaid*
- ⚡ Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can *e.g. XLEU Maxamum*

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Maple Syrup Urine Disease Products

AMINO ACID FORMULA (WITHOUT ISOLEUCINE, LEUCINE AND VALINE) – **Restricted** see terms on the preceding page

⬆ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can		<i>e.g. MSUD Anamix Infant</i>
⬆ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can		<i>e.g. MSUD Maxamaid</i>
⬆ Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can		<i>e.g. MSUD Maxamum</i>
⬆ Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle		<i>e.g. MSUD Anamix Junior LQ</i>

Phenylketonuria Products

AMINO ACID FORMULA (WITHOUT PHENYLALANINE) – **Restricted** see terms on the preceding page

⬆ Tab 8.33 mg		<i>e.g. Phlexy-10</i>
⬆ Powder 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet		<i>e.g. PKU Anamix Junior</i>
⬆ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can		<i>e.g. PKU Anamix Infant</i>
⬆ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can		<i>e.g. XP Maxamaid</i>
⬆ Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can		<i>e.g. XP Maxamum</i>
⬆ Powder 8.33 g protein and 8.8 g carbohydrate per 20 g sachet		<i>e.g. Phlexy-10</i>
⬆ Liquid 10 g protein, 4.4 g carbohydrate and 0.25 g fibre per 100 ml, 62.5 ml bottle		<i>e.g. PKU Lophlex LQ 10</i>
⬆ Liquid 20 g protein, 8.8 g carbohydrate and 0.34 g fibre per 100 ml, 125 ml bottle		<i>e.g. PKU Lophlex LQ 20</i>
⬆ Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, bottle	13.10	125 ml PKU Anamix Junior LQ (Berry)
		PKU Anamix Junior LQ (Orange)
		PKU Anamix Junior LQ (Unflavoured)
⬆ Liquid 16 g protein, 7 g carbohydrate and 0.27 g fibre per 100 ml, 125 ml bottle		<i>e.g. PKU Lophlex LQ 20</i>
⬆ Liquid 16 g protein, 7 g carbohydrate and 0.27 g fibre per 100 ml, 62.5 ml bottle		<i>e.g. PKU Lophlex LQ 10</i>
⬆ Liquid 16 g protein, 7 g carbohydrate and 0.4 g fibre per 100 ml, 125 ml bottle		<i>e.g. PKU Lophlex LQ 20</i>
⬆ Liquid 16 g protein, 7 g carbohydrate and 0.4 g fibre per 100 ml, 62.5 ml bottle		<i>e.g. PKU Lophlex LQ 10</i>
⬆ Liquid 6.7 g protein, 5.1 g carbohydrate and 2 g fat per 100 ml, 250 ml carton		<i>e.g. Easiphen</i>

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Propionic Acidemia and Methylmalonic Acidemia Products		
AMINO ACID FORMULA (WITHOUT ISOLEUCINE, METHIONINE, THREONINE AND VALINE) – Restricted see terms on page 202		
⚡ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can		e.g. MMA/PA Anamix Infant
⚡ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can		e.g. XMTVI Maxamaid
⚡ Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can		e.g. XMTVI Maxamum
Protein Free Supplements		
PROTEIN FREE SUPPLEMENT – Restricted see terms on page 202		
⚡ Powder nil added protein and 67 g carbohydrate per 100 g, 400 g can		e.g. Energivit
Tyrosinaemia Products		
AMINO ACID FORMULA (WITHOUT PHENYLALANINE AND TYROSINE) – Restricted see terms on page 202		
⚡ Powder 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet		e.g. TYR Anamix Junior
⚡ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can		e.g. TYR Anamix Infant
⚡ Powder 25 g protein and 51 g carbohydrate per 100 g, 400 g can		e.g. XPHEN, TYR Maxamaid
⚡ Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle		e.g. TYR Anamix Junior LQ
Urea Cycle Disorders Products		
AMINO ACID SUPPLEMENT – Restricted see terms on page 202		
⚡ Powder 25 g protein and 65 g carbohydrate per 100 g, 200 g can		e.g. Dialamine
⚡ Powder 79 g protein per 100 g, 200 g can		e.g. Essential Amino Acid Mix
X-Linked Adrenoleukodystrophy Products		
GLYCEROL TRIERUCATE – Restricted see terms on page 202		
⚡ Liquid, 1,000 ml bottle		
GLYCEROL TRIOLEATE – Restricted see terms on page 202		
⚡ Liquid, 500 ml bottle		
Specialised Formulas		
Diabetic Products		
➡ Restricted		
Initiation		
Any of the following:		
1 For patients with type I or type II diabetes suffering weight loss and malnutrition that requires nutritional support; or		
2 For patients with pancreatic insufficiency; or		
3 For patients who have, or are expected to, eat little or nothing for 5 days; or		
continued...		
⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below) e.g. Brand indicates brand example only. It is not a contracted product.		

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued. . .			
4 For patients who have a poor absorptive capacity and/or high nutrient losses and/or increased nutritional needs from causes such as catabolism; or			
5 For use pre- and post-surgery; or			
6 For patients being tube-fed; or			
7 For tube-feeding as a transition from intravenous nutrition.			
LOW-GI ENTERAL FEED 1 KCAL/ML – Restricted see terms on the preceding page			
⬆ Liquid 5 g protein, 9.6 g carbohydrate and 5.4 g fat per 100 ml, 1,000 ml bottle	7.50	1,000 ml	Glucerna Select RTH (Vanilla)
⬆ Liquid 4.3 g protein, 11.3 g carbohydrate and 4.2 g fat per 100 ml, 1,000 ml bag			<i>e.g. Nutrison Advanced Diason</i>
LOW-GI ORAL FEED 1 KCAL/ML – Restricted see terms on the preceding page			
⬆ Liquid 4.5 g protein, 9.8 g carbohydrate, 4.4 g fat and 1.9 g fibre per 100 ml, can	2.10	237 ml	Sustagen Diabetic (Vanilla)
⬆ Liquid 5 g protein, 9.6 g carbohydrate and 5.4 g fat per 100 ml, 250 ml bottle	1.88	250 ml	Glucerna Select (Vanilla)
⬆ Liquid 6 g protein, 9.5 g carbohydrate, 4.7 g fat and 2.6 g fibre per 100 ml, can	2.10	237 ml	Resource Diabetic (Vanilla)
⬆ Liquid 4.9 g protein, 11.7 g carbohydrate, 3.8 g fat and 2 g fibre per 100 ml, 200 ml bottle			<i>e.g. Diasip</i>

Elemental and Semi-Elemental Products

➔ **Restricted**

Initiation

Any of the following:

- 1 Malabsorption; or
- 2 Short bowel syndrome; or
- 3 Enterocutaneous fistulas; or
- 4 Eosinophilic enteritis (including oesophagitis); or
- 5 Inflammatory bowel disease; or
- 6 Acute pancreatitis where standard feeds are not tolerated; or
- 7 Patients with multiple food allergies requiring enteral feeding.

AMINO ACID ORAL FEED – **Restricted** see terms above

⬆ Powder 11 g protein, 62 g carbohydrate and 1 g fat per sachet	4.50	80 g	Vivonex TEN
---	------	------	-------------

AMINO ACID ORAL FEED 0.8 KCAL/ML – **Restricted** see terms above

⬆ Liquid 2.5 g protein, 11 g carbohydrate and 3.5 g fat per 100 ml, 250 ml carton			<i>e.g. Elemental 028 Extra</i>
---	--	--	---------------------------------

PEPTIDE-BASED ENTERAL FEED 1 KCAL/ML – **Restricted** see terms above

⬆ Liquid 4 g protein, 17.6 g carbohydrate and 1.7 g fat per 100 ml, 1,000 ml bag			<i>e.g. Nutrison Advanced Peptisorb</i>
--	--	--	---

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PEPTIDE-BASED ORAL FEED – Restricted see terms on the preceding page			
☞ Powder 13.7 g protein, 62.9 g carbohydrate and 17.5 g fat per 100 g, 400 g can			<i>e.g. Peptamen Junior</i>
☞ Powder 13.8 g protein, 59 g carbohydrate and 18 g fat per 100 g, 400 g can			<i>e.g. MCT Pepdite; MCT Pepdite 1+</i>
☞ Powder 15.8 g protein, 49.5 g carbohydrate and 4.65 g fat per 76 g sachet	7.50	76 g	Alitraq
☞ Liquid 6.75 g protein, 18.4 g carbohydrate and 5.5 g fat per 100 ml, bottle	18.06	1,000 ml	Vital
PEPTIDE-BASED ORAL FEED 1 KCAL/ML – Restricted see terms on the preceding page			
☞ Liquid 5 g protein, 16 g carbohydrate and 1.69 g fat per 100 ml, carton	4.95	237 ml	Peptamen OS 1.0 (Vanilla)

Fat Modified Products

FAT-MODIFIED FEED – **Restricted** see terms below

☞ Powder 11.4 g protein, 68 g carbohydrate and 11.8 g fat per 100 g, 400 g can			<i>e.g. Monogen</i>
☞ Powder 12.9 g protein, 69.1 g carbohydrate and 12.9 g fat per 100 g, 400 g can			<i>e.g. Monogen</i>

☞ **Restricted**

Initiation

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

Hepatic Products

☞ **Restricted**

Initiation

For children (up to 18 years) who require a liver transplant.

HEPATIC ORAL FEED – **Restricted** see terms above

☞ Powder 11 g protein, 64 g carbohydrate and 20 g fat per 100 g, can	78.97	400 g	Heparon Junior
--	-------	-------	----------------

High Calorie Products

☞ **Restricted**

Initiation

Any of the following:

- 1 Patient is fluid volume or rate restricted; or
- 2 Patient requires low electrolyte; or
- 3 Both:
 - 3.1 Any of the following:
 - 3.1.1 Cystic fibrosis; or
 - 3.1.2 Any condition causing malabsorption; or
 - 3.1.3 Faltering growth in an infant/child; or
 - 3.1.4 Increased nutritional requirements; and
 - 3.2 Patient has substantially increased metabolic requirements.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ENTERAL FEED 2 KCAL/ML – Restricted see terms on the preceding page			
⬆ Liquid 7.5 g protein, 20 g carbohydrate and 10 g fat per 100 ml, bottle	5.50	500 ml	Nutrison Concentrated
⬆ Liquid 8.4 g protein, 21.9 g carbohydrate, 9.1 g fat and 0.5 g fibre per 100 ml, bottle	11.00	1,000 ml	TwoCal HN RTH (Vanilla)
ORAL FEED 2 KCAL/ML – Restricted see terms on the preceding page			
⬆ Liquid 8.4 g protein, 22.4 g carbohydrate, 8.9 g fat and 0.8 g fibre per 100 ml, bottle	1.90	200 ml	Two Cal HN

High Protein Products

HIGH PROTEIN ENTERAL FEED 1.25 KCAL/ML – **Restricted** see terms below

⬆ Liquid 6.3 g protein, 14.2 g carbohydrate and 4.9 g fat per 100 ml, 1,000 ml bag	<i>e.g. Nutrison Protein Plus</i>
---	---------------------------------------

➡Restricted

Initiation

Both:

- 1 The patient has a high protein requirement; and
- 2 Any of the following:
 - 2.1 Patient has liver disease; or
 - 2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
 - 2.3 Patient is fluid restricted; or
 - 2.4 Patient's needs cannot be more appropriately met using high calorie product.

HIGH PROTEIN ENTERAL FEED 1.28 KCAL/ML – **Restricted** see terms below

⬆ Liquid 6.3 g protein, 14.1 g carbohydrate, 4.9 g fat and 1.5 g fibre per 100 ml, 1,000 ml bag	<i>e.g. Nutrison Protein Plus Multi Fibre</i>
--	---

➡Restricted

Initiation

Both:

- 1 The patient has a high protein requirement; and
- 2 Any of the following:
 - 2.1 Patient has liver disease; or
 - 2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
 - 2.3 Patient is fluid restricted; or
 - 2.4 Patient's needs cannot be more appropriately met using high calorie product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Infant Formulas			
AMINO ACID FORMULA – Restricted see terms below			
☞ Powder 1.95 g protein, 8.1 g carbohydrate and 3.5 g fat per 100 ml, 400 g can			<i>e.g. Neocate</i>
☞ Powder 13 g protein, 52.5 g carbohydrate and 24.5 g fat per 100 g, 400 g can			<i>e.g. Neocate LCP</i>
☞ Powder 13.5 g protein, 52 g carbohydrate and 24.5 g fat per 100 g, can53.00	400 g		Neocate Gold (Unflavoured)
☞ Powder 14 g protein, 50 g carbohydrate and 24.3 g fat per 100 g, 400 g can			<i>e.g. Neocate Advance</i>
☞ Powder 16 g protein, 51.4 g carbohydrate and 21 g fat per 100 g, can53.00	400 g		Neocate Advance (Vanilla)
☞ Powder 2.2 g protein, 7.8 g carbohydrate and 3.4 g fat per 100 ml, can53.00	400 g		Elecare LCP (Unflavoured)
☞ Powder 2.2 g protein, 7.8 g carbohydrate and 3.4 g fat per 100 ml, can53.00	400 g		Elecare (Unflavoured)
☞ Powder 2.2 g protein, 7.8 g carbohydrate and 3.4 g fat per 100 ml, can53.00	400 g		Elecare (Vanilla)
☞ Powder 6 g protein, 31.5 g carbohydrate and 5.88 g fat per sachet6.00	48.5 g		Vivonex Paediatric

☞**Restricted**

Initiation

Any of the following:

- 1 Extensively hydrolysed formula has been reasonably trialled and is inappropriate due to documented severe intolerance or allergy or malabsorption; or
- 2 History of anaphylaxis to cows' milk protein formula or dairy products; or
- 3 Eosinophilic oesophagitis.

Note: A reasonable trial is defined as a 2-4 week trial.

Continuation

Both:

- 1 An assessment as to whether the infant can be transitioned to a cows' milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an amino acid infant formula.

EXTENSIVELY HYDROLYSED FORMULA – **Restricted** see terms below

☞ Powder 14 g protein, 53.4 g carbohydrate and 27.3 g fat per 100 g, 450 g can	<i>e.g. Aptamil Gold+ Pepti Junior</i>
--	--

☞**Restricted**

Initiation

Any of the following:

- 1 Both:
 - 1.1 Cows' milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
7 Cystic fibrosis; or			
8 Proven fat malabsorption; or			
9 Severe intestinal motility disorders causing significant malabsorption; or			
10 Intestinal failure; or			
11 For step down from Amino Acid Formula.			
Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.			
Continuation			
Both:			
1 An assessment as to whether the infant can be transitioned to a cows' milk protein or soy infant formula has been undertaken; and			
2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula.			
FRUCTOSE-BASED FORMULA			
Powder 14.6 g protein, 49.7 g carbohydrate and 30.8 g fat per 100 g, 400 g can			<i>e.g. Galactomin 19</i>
LACTOSE-FREE FORMULA			
Powder 1.3 g protein, 7.3 g carbohydrate and 3.5 g fat per 100 ml, 900 g can			<i>e.g. Karicare Aptamil Gold De-Lact</i>
Powder 1.5 g protein, 7.2 g carbohydrate and 3.6 g fat per 100 ml, 900 g can			<i>e.g. S26 Lactose Free</i>
LOW-CALCIUM FORMULA			
Powder 14.6 g protein, 53.7 g carbohydrate and 26.1 g fat per 100 g, 400 g can			<i>e.g. Locasol</i>
PAEDIATRIC ORAL FEED 1 KCAL/ML – Restricted see terms below			
☞ Liquid 2.6 g protein, 10.3 g carbohydrate, 5.4 g fat and 0.6 g fibre per 100 ml, 100 ml bottle			<i>e.g. Infatrini</i>
☞ Restricted			
Initiation			
Both:			
1 Either:			
1.1 The patient is fluid restricted; or			
1.2 The patient has increased nutritional requirements due to faltering growth; and			
2 Patient is under 18 months old and weighs less than 8kg.			
PRETERM FORMULA – Restricted see terms below			
☞ Powder 1.9 g protein, 7.5 g carbohydrate and 3.9 g fat per 14 g, can 15.25	400 g		S-26 Gold Premgro
☞ Liquid 2.2 g protein, 8.4 g carbohydrate and 4.4 g fat per 100 ml, bottle 0.75	100 ml		S26 LBW Gold RTF
☞ Liquid 2.3 g protein, 8.6 g carbohydrate and 4.2 g fat per 100 ml, 90 ml bottle			<i>e.g. Pre Nan Gold RTF</i>
☞ Liquid 2.6 g protein, 8.4 g carbohydrate and 3.9 g fat per 100 ml, 70 ml bottle			<i>e.g. Karicare Aptamil Gold+Preterm</i>
☞ Restricted			
Initiation			
For infants born before 33 weeks' gestation or weighing less than 1.5 kg at birth.			
THICKENED FORMULA			
Powder 1.8 g protein, 8.1 g carbohydrate and 3.3 g fat per 100 ml, 900 g can			<i>e.g. Karicare Aptamil Thickened AR</i>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Ketogenic Diet Products			
HIGH FAT FORMULA – Restricted see terms below			
☞ Powder 14.4 g protein, 2.9 g carbohydrate and 69.2 g fat per 100 g, can	35.50	300 g	Ketocal 4:1 (Unflavoured) Ketocal 4:1 (Vanilla)
☞ Powder 15.3 g protein, 7.2 g carbohydrate and 67.7 g fat per 100 g, can	35.50	300 g	Ketocal 3:1 (Unflavoured)
☛ Restricted			
Initiation			
For patients with intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.			
Paediatric Products			
☛ Restricted			
Initiation			
Both:			
1 Child is aged one to ten years; and			
2 Any of the following:			
2.1 The child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or			
2.2 Any condition causing malabsorption; or			
2.3 Faltering growth in an infant/child; or			
2.4 Increased nutritional requirements; or			
2.5 The child is being transitioned from TPN or tube feeding to oral feeding; or			
2.6 The child has eaten, or is expected to eat, little or nothing for 3 days.			
PAEDIATRIC ORAL FEED – Restricted see terms above			
☞ Powder 14.9 g protein, 54.3 g carbohydrate and 24.7 g fat per 100 g, can	20.00	850 g	Pediasure (Vanilla)
PAEDIATRIC ENTERAL FEED 0.76 KCAL/ML – Restricted see terms above			
☞ Liquid 2.5 g protein, 12.5 g carbohydrate, 3.3 g fat and 0.7 g fibre per 100 ml, bag	4.00	500 ml	Nutrini Low Energy Multifibre RTH
PAEDIATRIC ENTERAL FEED 1 KCAL/ML – Restricted see terms above			
☞ Liquid 2.8 g protein, 11.2 g carbohydrate and 5 g fat per 100 ml, bag	2.68	500 ml	Pediasure RTH
☞ Liquid 2.8 g protein, 12.3 g carbohydrate and 4.4 g fat per 100 ml, 500 ml bag			e.g. <i>Nutrini RTH</i>
PAEDIATRIC ENTERAL FEED 1.5 KCAL/ML – Restricted see terms above			
☞ Liquid 4.1 g protein, 18.5 g carbohydrate, 6.7 g fat and 0.8 g fibre per 100 ml, bag	6.00	500 ml	Nutrini Energy Multi Fibre
☞ Liquid 4.1 g protein, 18.5 g carbohydrate and 6.7 g fat per 100 ml, 500 ml bag			e.g. <i>Nutrini Energy RTH</i>
PAEDIATRIC ORAL FEED 1 KCAL/ML – Restricted see terms above			
☞ Liquid 4.2 g protein, 16.7 g carbohydrate and 7.5 g fat per 100 ml, bottle	1.07	200 ml	Pediasure (Chocolate) Pediasure (Strawberry) Pediasure (Vanilla)
☞ Liquid 4.2 g protein, 16.7 g carbohydrate and 7.5 g fat per 100 ml, can	1.34	250 ml	Pediasure (Vanilla)

☞ Item restricted (see ☛ above); ☞ Item restricted (see ☛ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PAEDIATRIC ORAL FEED 1.5 KCAL/ML – Restricted see terms on the preceding page			
☞ Liquid 3.4 g protein, 18.8 g carbohydrate and 6.8 g fat per 100 ml, 200 ml bottle			<i>e.g. Fortini</i>
☞ Liquid 4.0 g protein, 18.8 g carbohydrate, 6.8 g fat and 1.5 g fibre per 100 ml, 200 ml bottle			<i>e.g. Fortini Multifibre</i>

Renal Products

LOW ELECTROLYTE ENTERAL FEED 1.8 KCAL/ML – **Restricted** see terms below

☞ Liquid 8.1 g protein, 14.74 g carbohydrate, 9.77 g fat and 1.26 g fibre per 100 ml, bottle	6.08	500 ml	Nepro HP RTH
--	------	--------	--------------

➔Restricted

Initiation

For patients with acute or chronic kidney disease.

LOW ELECTROLYTE ORAL FEED – **Restricted** see terms below

☞ Powder 7.5 g protein, 59 g carbohydrate and 26.3 g fat per 100 g, 400 g can			<i>e.g. Kindergen</i>
---	--	--	-----------------------

➔Restricted

Initiation

For children (up to 18 years) with acute or chronic kidney disease.

LOW ELECTROLYTE ORAL FEED 1.8 KCAL/ML

☞ Liquid 8 g protein, 14.74 g carbohydrate, 9.77 g fat and 1.26 g fibre per 100 ml, carton	2.67	220 ml	Nepro HP (Strawberry) Nepro HP (Vanilla)
--	------	--------	---

➔Restricted

Initiation

For patients with acute or chronic kidney disease.

LOW ELECTROLYTE ORAL FEED 2 KCAL/ML – **Restricted** see terms below

☞ Liquid 9.1 g protein, 19 g carbohydrate and 10 g fat per 100 ml, carton	3.31	237 ml	Novasource Renal (Vanilla)
---	------	--------	-------------------------------

☞ Liquid 3 g protein, 25.5 g carbohydrate and 9.6 g fat per 100 ml, 237 ml bottle			
---	--	--	--

☞ Liquid 7.5 g protein, 20 g carbohydrate and 10 g fat per 100 ml, 125 ml carton			<i>e.g. Renilon 7.5</i>
--	--	--	-------------------------

➔Restricted

Initiation

For patients with acute or chronic kidney disease.

Respiratory Products

LOW CARBOHYDRATE ORAL FEED 1.5 KCAL/ML – **Restricted** see terms below

☞ Liquid 6.2 g protein, 10.5 g carbohydrate and 9.32 g fat per 100 ml, bottle	1.66	237 ml	Pulmocare (Vanilla)
---	------	--------	---------------------

➔Restricted

Initiation

For patients with CORD and hypercapnia, defined as a CO₂ value exceeding 55 mmHg.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Surgical Products			
HIGH ARGININE ORAL FEED 1.4 KCAL/ML – Restricted see terms below			
☞ Liquid 7.6 g protein, 18.9 g carbohydrate, 3.9 g fat and 1.4 g fibre per 100 ml, carton	4.00	237 ml	Impact Advanced Recovery (Chocolate) Impact Advanced Recovery (Vanilla)

☞ **Restricted****Initiation**

Three packs per day for 5 to 7 days prior to major gastrointestinal, head or neck surgery.

PREOPERATIVE CARBOHYDRATE FEED 0.5 KCAL/ML – **Restricted** see terms below

☞ Oral liq 0 g protein, 12.6 g carbohydrate and 0 g fat per 100 ml, 200 ml bottle	6.80	4	preOp
--	------	---	-------

☞ **Restricted****Initiation**

Maximum of 400 ml as part of an Enhanced Recovery After Surgery (ERAS) protocol 2 to 3 hours before major abdominal surgery.

Standard Feeds☞ **Restricted****Initiation**

Any of the following:

For patients with malnutrition, defined as any of the following:

1 Any of the following:

1.1 BMI < 18.5; or

1.2 Greater than 10% weight loss in the last 3-6 months; or

1.3 BMI < 20 with greater than 5% weight loss in the last 3-6 months; or

2 For patients who have, or are expected to, eat little or nothing for 5 days; or

3 For patients who have a poor absorptive capacity and/or high nutrient losses and/or increased nutritional needs from causes such as catabolism; or

4 For use pre- and post-surgery; or

5 For patients being tube-fed; or

6 For tube-feeding as a transition from intravenous nutrition; or

7 For any other condition that meets the community Special Authority criteria.

ENTERAL FEED 1.5 KCAL/ML – **Restricted** see terms above

☞ Liquid 5.4 g protein, 13.6 g carbohydrate and 3.3 g fat per 100 ml, 1,000 ml bottle			e.g. <i>Isosource Standard RTH</i>
☞ Liquid 6 g protein, 18.3 g carbohydrate and 5.8 g fat per 100 ml, bag	7.00	1,000 ml	Nutrison Energy
☞ Liquid 6 g protein, 18.4 g carbohydrate, 5.8 g fat and 1.5 g fibre per 100 ml, 1,000 ml bag			e.g. <i>Nutrison Energy Multi Fibre</i>
☞ Liquid 6.25 g protein, 20 g carbohydrate and 5 g fat per 100 ml, can	1.75	250 ml	Ensure Plus HN
☞ Liquid 6.27 g protein, 20.4 g carbohydrate and 4.9 g fat per 100 ml, bag	7.00	1,000 ml	Ensure Plus HN RTH
☞ Liquid 6.38 g protein, 21.1 g carbohydrate, 4.9 g fat and 1.2 g fibre per 100 ml, bag	7.00	1,000 ml	Jevity HiCal RTH

☞ Item restricted (see ☞ above); ☞ Item restricted (see ☞ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ENTERAL FEED 1 KCAL/ML – Restricted see terms on the preceding page			
⬆ Liquid 4 g protein, 13.6 g carbohydrate and 3.4 g fat per 100 ml, bottle	5.29	1,000 ml	Osmolite RTH
⬆ Liquid 4 g protein, 14.1 g carbohydrate, 3.47 g fat and 1.76 g fibre per 100 ml, bottle	2.65	500 ml	Jevity RTH
	5.29	1,000 ml	Jevity RTH
⬆ Liquid 4 g protein, 14.1 g carbohydrate, 3.47 g fat and 1.76 g fibre per 100 ml, can	1.32	237 ml	Jevity
⬆ Liquid 4 g protein, 12.3 g carbohydrate and 3.9 g fat per 100 ml, 1,000 ml bag			<i>e.g. NutrisonStdRTH; NutrisonLowSodium</i>
⬆ Liquid 4 g protein, 12.3 g carbohydrate, 3.9 g fat and 1.5 g fibre per 100 ml, 1000 ml bag			<i>e.g. Nutrison Multi Fibre</i>
ENTERAL FEED 1.2 KCAL/ML – Restricted see terms on the preceding page			
⬆ Liquid 5.55 g protein, 15.1 g carbohydrate, 3.93 g fat and 2 g fibre per 100 ml, 1,000 ml bag			<i>e.g. Jevity Plus RTH</i>
ORAL FEED – Restricted see terms on the preceding page			
⬆ Powder 16 g protein, 59.8 g carbohydrate and 14 g fat per 100 g, can	13.00	850 g	Ensure (Chocolate) Ensure (Vanilla)
⬆ Powder 21.9 g protein, 53.5 g carbohydrate and 14.5 g fat per 100 g, can	3.67	350 g	Fortisip (Vanilla)
⬆ Powder 23 g protein, 65 g carbohydrate and 2.5 g fat per 100 g, can	14.90	840 g	Sustagen Hospital Formula (Chocolate) Sustagen Hospital Formula (Vanilla)
Note: Community subsidy of Sustagen Hospital Formula is subject to both Special Authority criteria and a manufacturer's surcharge. Higher subsidy by endorsement is available for patients meeting the following endorsement criteria; fat malabsorption, fat intolerance or chyle leak.			
ORAL FEED 1 KCAL/ML – Restricted see terms on the preceding page			
⬆ Liquid 3.8 g protein, 23 g carbohydrate and 12.7 g fibre per 100 ml, 237 ml carton			<i>e.g. Resource Fruit Beverage</i>
ORAL FEED 1.5 KCAL/ML – Restricted see terms on the preceding page			
⬆ Liquid 5.5 g protein, 21.1 g carbohydrate and 4.81 g fat per 100 ml, can	1.33	237 ml	Ensure Plus (Chocolate) Ensure Plus (Vanilla)
⬆ Liquid 6.25 g protein, 20.2 g carbohydrate and 4.92 g fat per 100 ml, carton	1.26	200 ml	Ensure Plus (Banana) Ensure Plus (Chocolate) Ensure Plus (Fruit of the Forest) Ensure Plus (Vanilla)
⬆ Liquid 4 g protein and 33.5 g carbohydrate per 100 ml, 200 ml bottle			<i>e.g. Fortijuce</i>
⬆ Liquid 6 g protein, 18.4 g carbohydrate and 5.8 g fat per 100 ml, 200 ml bottle			<i>e.g. Fortisip</i>
⬆ Liquid 6 g protein, 18.4 g carbohydrate, 5.8 g fat and 2.3 g fibre per 100 ml, 200 ml bottle			<i>e.g. Fortisip Multi Fibre</i>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Bacterial and Viral Vaccines

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE – **Restricted** see terms below

<p>☞ Inj 30 IU diphtheria toxoid with 30IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units poliomyelitis virus in 0.5 ml syringe – 1% DV Jul-14 to 2017</p>	0.00	10	Infanrix IPV
--	------	----	--------------

☞ **Restricted**

Initiation

Any of the following:

- 1 A single dose for children up to the age of 7 who have completed primary immunisation; or
- 2 A course of up to four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
- 3 An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 4 Five doses will be funded for children requiring solid organ transplantation.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes

DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE – **Restricted** see terms below

<p>☞ Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin, 80 D-antigen units poliomyelitis virus, 10 mcg hepatitis B surface antigen in 0.5 ml syringe (1) and inj 10 mcg haemophilus influenzae type B vaccine vial – 1% DV Jul-14 to 2017</p>	0.00	10	Infanrix-hexa
--	------	----	---------------

☞ **Restricted**

Initiation

Any of the following:

- 1 Up to four doses for children up to and under the age of 10 for primary immunisation; or
- 2 An additional four doses (as appropriate) are funded for (re-)immunisation for children up to and under the age of 10 who are patients post haematopoietic stem cell transplantation, or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 3 Up to five doses for children up to and under the age of 10 receiving solid organ transplantation.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

Bacterial Vaccines

ADULT DIPHTHERIA AND TETANUS VACCINE

<p>☞ Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid in 0.5 ml syringe – 1% DV Jul-14 to 2017</p>	0.00	5	ADT Booster
--	------	---	-------------

☞ **Restricted**

Initiation

Any of the following:

- 1 For vaccination of patients aged 45 and 65 years old; or
- 2 For vaccination of previously unimmunised or partially immunised patients; or
- 3 For revaccination following immunosuppression; or
- 4 For boosting of patients with tetanus-prone wounds; or
- 5 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BACILLUS CALMETTE-GUERIN VACCINE – Restricted see terms below			
☛ Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin), Danish strain 1331, live attenuated, vial Danish strain 1331, live attenuated, vial with diluent – 1% DV Oct-14 to 2017	0.00	10	BCG Vaccine
☛ Restricted			
Initiation			
All of the following:			
For infants at increased risk of tuberculosis defined as:			
1 Living in a house or family with a person with current or past history of TB; and			
2 Having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; and			
3 During their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000.			
Note: A list of countries with high rates of TB are available at http://www.health.govt.nz/tuberculosis (Search for Downloads) or www.bcgatlas.org/index.php			
DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE – Restricted see terms below			
☛ Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis toxoid, 8 mcg pertussis filamentous haemagglutinin and 2.5 mcg pertactin in 0.5 ml syringe – 1% DV Jul-14 to 2017	0.00	1 10	Boostrix Boostrix
☛ Restricted			
Initiation			
Any of the following:			
1 A single vaccine for pregnant woman between gestational weeks 28 and 38; or			
2 A course of up to four vaccines is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or			
3 An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens.			
Note: Tdap is not registered for patients aged less than 10 years. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.			
HAEMOPHILUS INFLUENZAE TYPE B VACCINE – Restricted see terms below			
☛ Inj 10 mcg vial with diluent syringe – 1% DV Jul-14 to 2017	0.00	1	Act-HIB
☛ Restricted			
Initiation			
<i>Therapy limited to 1 dose</i>			
Any of the following:			
1 For primary vaccination in children; or			
2 An additional dose (as appropriate) is funded for (re-)immunisation for patients post haematopoietic stem cell transplantation, or chemotherapy; functional asplenic; pre or post splenectomy; pre- or post solid organ transplant, pre- or post cochlear implants, renal dialysis and other severely immunosuppressive regimens; or			
3 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.			
MENINGOCOCCAL (A, C, Y AND W-135) CONJUGATE VACCINE – Restricted see terms on the next page			
☛ Inj 4 mcg or each meningococcal polysaccharide conjugated to a total of approximately 48 mcg of diphtheria toxoid carrier per 0.5 ml vial – 1% DV Jul-14 to 2017	0.00	1	Menactra

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➡Restricted			
Initiation			
Any of the following:			
1 Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with HIV, complement deficiency (acquired or inherited), functional or anatomic asplenia or pre or post solid organ transplant; or			
2 One dose for close contacts of meningococcal cases; or			
3 A maximum of two doses for bone marrow transplant patients; or			
4 A maximum of two doses for patients following immunosuppression*.			
Notes: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.			
*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.			
MENINGOCOCCAL C CONJUGATE VACCINE – Restricted see terms below			
☞ Inj 10 mcg in 0.5 ml syringe – 1% DV Jul-14 to 2017	0.00	1	Neisvac-C
		10	Neisvac-C
➡Restricted			
Initiation			
Any of the following:			
1 Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with HIV, complement deficiency (acquired or inherited), functional or anatomic asplenia or pre or post solid organ transplant; or			
2 One dose for close contacts of meningococcal cases; or			
3 A maximum of two doses for bone marrow transplant patients; or			
4 A maximum of two doses for patients following immunosuppression*.			
Notes: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.			
*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.			
PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE – Restricted see terms below			
☞ Inj 30.8 mcg in 0.5 ml syringe – 1% DV Oct-14 to 2017	0.00	1	Prevenar 13
		10	Prevenar 13
➡Restricted			
Initiation			
Any of the following:			
1 A primary course of up to four doses for previously unvaccinated individuals up to the age of 59 months inclusive; or			
2 Up to three doses as appropriate to complete the primary course of immunisation for individuals under the age of 59 months who have received one to three doses of PCV10; or			
3 One dose is funded for high risk children (over the age of 17 months and up to the age of 18) who have previously received four doses of PCV10; or			
4 Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients with HIV, for patients post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, primary immunodeficiency; or			
5 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.			
Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes			
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE – Restricted see terms on the next page			
☞ Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype) – 1% DV Jun-15 to 2017	0.00	1	Pneumovax 23

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ **Restricted****Initiation**

Any of the following:

- 1 Up to three doses (as appropriate) for patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy; functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- 2 Up to two doses are funded for high risk children to the age of 18; or
- 3 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

SALMONELLA TYPHI VACCINE – Restricted see terms below

⚡ Inj 25 mcg in 0.5 ml syringe

➔ **Restricted****Initiation**

For use during typhoid fever outbreaks.

Viral Vaccines

HEPATITIS A VACCINE – Restricted see terms below

⚡ Inj 720 ELISA units in 0.5 ml syringe – 1% DV Jul-14 to 2017	0.00	1	Havrix Junior
⚡ Inj 1440 ELISA units in 1 ml syringe – 1% DV Jul-14 to 2017	0.00	1	Havrix

➔ **Restricted****Initiation**

All of the following:

- 1 Two vaccinations for use in transplant patients; and
- 2 Two vaccinations for use in children with chronic liver disease; and
- 3 One dose of vaccine for close contacts of known hepatitis A cases.

HEPATITIS B RECOMBINANT VACCINE

⚡ Inj 5 mcg in 0.5 ml vial – 1% DV Jul-14 to 2017	0.00	1	HBvaxPRO
---	------	---	-----------------

➔ **Restricted****Initiation**

Any of the following:

- 1 For household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2 For children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3 For children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination; or
- 4 For HIV positive patients; or
- 5 For hepatitis C positive patients; or
- 6 for patients following non-consensual sexual intercourse; or
- 7 For patients following immunosuppression; or
- 8 For transplant patients; or
- 9 following needle stick injury.

⚡ Inj 10 mcg in 1 ml vial – 1% DV Jul-14 to 2017	0.00	1	HBvaxPRO
--	------	---	-----------------

➔ **Restricted****Initiation**

Any of the following:

- 1 For household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2 For children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3 For children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
4 For HIV positive patients; or			
5 For hepatitis C positive patients; or			
6 for patients following non-consensual sexual intercourse; or			
7 For patients following immunosuppression; or			
8 For transplant patients; or			
9 following needle stick injury.			
¶ Inj 40 mcg per 1 ml vial – 1% DV Jul-14 to 2017	0.00	1	HBvaxPRO
➡ Restricted			
Initiation			
Both:			
1 For dialysis patients; and			
2 For liver or kidney transplant patient.			
HUMAN PAPILLOMAVIRUS (6, 11, 16 AND 18) VACCINE [HPV] – Restricted see terms below			
¶ Inj 120 mcg in 0.5 ml syringe – 1% DV Jul-14 to 2017	0.00	10	Gardasil
➡ Restricted			
Initiation			
<i>Therapy limited to 3 doses</i>			
Any of the following:			
1 Females aged under 20 years old; or			
2 Patients aged under 26 years old with confirmed HIV infection; or			
3 For use in transplant (including stem cell) patients; or			
4 An additional dose for patients under 26 years of age post chemotherapy.			
INFLUENZA VACCINE – Restricted see terms below			
¶ Inj 45 mcg in 0.5 ml syringe	90.00	10	Fluarix Influvac
➡ Restricted			
Initiation — People over 65			
The patient is 65 years of age or over.			
Initiation — cardiovascular disease			
Any of the following:			
1 Ischaemic heart disease; or			
2 Congestive heart failure; or			
3 Rheumatic heart disease; or			
4 Longenital heart disease; or			
5 Cerebro-vascular disease.			
Note: hypertension and/or dyslipidaemia without evidence of end-organ disease is excluded from funding.			
Initiation — chronic respiratory disease			
Either:			
1 Asthma, if on a regular preventative therapy; or			
2 Other chronic respiratory disease with impaired lung function.			
Note: asthma not requiring regular preventative therapy is excluded from funding.			
Initiation — Other conditions			
Either:			
1 Any of the following:			
1.1 Diabetes; or			
1.2 chronic renal disease; or			
1.3 Any cancer, excluding basal and squamous skin cancers if not invasive; or			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
1.4 Autoimmune disease; or			
1.5 Immune suppression or immune deficiency; or			
1.6 HIV; or			
1.7 Transplant recipient; or			
1.8 Neuromuscular and CNS diseases/ disorders; or			
1.9 Haemoglobinopathies; or			
1.10 Is a child on long term aspirin; or			
1.11 Has a cochlear implant; or			
1.12 Errors of metabolism at risk of major metabolic decompensation; or			
1.13 Pre and post splenectomy; or			
1.14 Down syndrome; or			
1.15 Is pregnant; or			
1.16 Is a child aged four and under who has been hospitalised for respiratory illness or has a history of significant respiratory illness; or			
2 Patients who are compulsorily detained long-term in a forensic unit within a DHB hospital.			
MEASLES, MUMPS AND RUBELLA VACCINE – Restricted see terms below			
¶ Inj 1000 TCID50 measles, 12500 TCID50 mumps and 1000 TCID50 rubella vial with diluent – 1% DV Jul-14 to 2017	0.00	10	M-M-R-II
➔ Restricted			
Initiation — first dose prior to 12 months			
<i>Therapy limited to 3 doses</i>			
Any of the following:			
1 For primary vaccination in children; or			
2 For revaccination following immunosuppression; or			
3 For any individual susceptible to measles, mumps or rubella.			
Initiation — first dose after to 12 months			
<i>Therapy limited to 2 doses</i>			
Any of the following:			
1 For primary vaccination in children; or			
2 For revaccination following immunosuppression; or			
3 For any individual susceptible to measles, mumps or rubella.			
Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.			
POLIOMYELITIS VACCINE – Restricted see terms below			
¶ Inj 80 D-antigen units in 0.5 ml syringe – 1% DV Jul-14 to 2017	0.00	1	IPOL
➔ Restricted			
Initiation			
<i>Therapy limited to 3 doses</i>			
Either:			
1 For partially vaccinated or previously unvaccinated individuals; or			
2 For revaccination following immunosuppression.			
Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.			
RABIES VACCINE			
Inj 2.5 IU vial with diluent			
ROTAVIRUS LIVE REASSORTANT ORAL VACCINE – Restricted see terms on the next page			
¶ Oral susp G1, G2, G3, G4, P1(8) 11.5 million CCID50 units per 2 ml, tube – 1% DV Jul-14 to 2017	0.00	10	RotaTeg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➡Restricted			
Initiation			
<i>Therapy limited to 3 doses</i>			
Both:			
1 First dose to be administered in infants aged under 15 weeks of age; and			
2 No vaccination being administered to children aged 8 months or over.			
VARICELLA VACCINE [CHICKEN POX VACCINE] – Restricted see terms below			
¶ Inj 2,000 PFU vial with diluent – 1% DV Jul-14 to 2017	0.00	1	Varilrix

➡Restricted

Initiation

Therapy limited to 2 doses

Any of the following:

- 1 Any of the following:
 - for non-immune patients
 - 1.1 With chronic liver disease who may in future be candidates for transplantation; or
 - 1.2 With deteriorating renal function before transplantation; or
 - 1.3 Prior to solid organ transplant; or
 - 1.4 Prior to any elective immunosuppression*; or
 - 1.5 For post exposure prophylaxis who are immune competent inpatients.; or
 - 2 For patients at least 2 years after bone marrow transplantation, on advice of their specialist; or
 - 3 For patients at least 6 months after completion of chemotherapy, on advice of their specialist; or
 - 4 For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
 - 5 For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or
 - 6 For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or
 - 7 For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.

Note: * immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
NOTE:			
In addition to the products expressly listed here in Part III: Optional Pharmaceuticals, a number of additional Optional Pharmaceuticals, including some wound care products and disposable laparoscopic equipment, are listed in an addendum to Part III which is available at www.pharmac.govt.nz . The Optional Pharmaceuticals listed in the addendum are deemed to be listed in Part III, and the Rules of the Pharmaceutical Schedule applying to products listed in Part III apply to them.			
Optional Pharmaceuticals			
BLOOD GLUCOSE DIAGNOSTIC TEST METER			
1 meter with 50 lancets, a lancing device, and 10 diagnostic test strips	20.00	1	Caresens II
			Caresens N
			Caresens N POP
Meter	19.00	1	Accu-Chek Performa
	9.00		FreeStyle Lite
			On Call Advanced
BLOOD GLUCOSE DIAGNOSTIC TEST STRIP			
Blood glucose test strips	28.75	50 test	Accu-Chek Performa
	10.56		CareSens
			CareSens N
	21.65		FreeStyle Lite
	28.75		Freestyle Optium
Blood glucose test strips × 50 and lancets × 5	19.10	50 test	On Call Advanced
BLOOD KETONE DIAGNOSTIC TEST METER			
Meter	40.00	1	Freestyle Optium Neo
INSULIN PEN NEEDLES			
29 g × 12.7 mm	10.50	100	B-D Micro-Fine
31 g × 5 mm	11.75	100	B-D Micro-Fine
31 g × 6 mm	10.50	100	ABM
31 g × 8 mm	10.50	100	B-D Micro-Fine
32 g × 4 mm	10.50	100	B-D Micro-Fine
INSULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE			
Syringe 0.3 ml with 29 g × 12.7 mm needle	13.00	100	B-D Ultra Fine
Syringe 0.3 ml with 31 g × 8 mm needle	13.00	100	B-D Ultra Fine II
Syringe 0.5 ml with 29 g × 12.7 mm needle	13.00	100	B-D Ultra Fine
Syringe 0.5 ml with 31 g × 8 mm needle	13.00	100	B-D Ultra Fine II
Syringe 1 ml with 29 g × 12.7 mm needle	13.00	100	B-D Ultra Fine
Syringe 1 ml with 31 g × 8 mm needle	13.00	100	B-D Ultra Fine II
KETONE BLOOD BETA-KETONE ELECTRODES			
Test strips	15.50	10 strip	Freestyle Optium Ketone
MASK FOR SPACER DEVICE			
Small	2.20	1	e-chamber Mask
PEAK FLOW METER			
Low Range	9.54	1	Mini-Wright AFS Low Range
Normal Range	9.54	1	Mini-Wright Standard
PREGNANCY TEST - HCG URINE			
Cassette – 1% DV Sep-15 to 2017	17.60	40 test	EasyCheck
SODIUM NITROPRUSSIDE			
Test strip	6.00	50 strip	Accu-Chek Ketur-Test

Products with Hospital Supply Status (HSS) are in **bold**

Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.

PART III - OPTIONAL PHARMACEUTICALS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SPACER DEVICE			
220 ml (single patient)	2.95	1	e-chamber Turbo
510 ml (single patient)	5.12	1	e-chamber La Grande
800 ml	6.50	1	Volumatic

- Symbols -		Agents Affecting the	Alimentary	24
8-methoxypsoralen	55	Renin-Angiotensin System	Infection	78
- A -		Agents for Parkinsonism and	Amsacrine	134
A-Scabies	52	Related Disorders	Amyl nitrite	48
Abacavir sulphate	85	Agents Used in the Treatment of	Anabolic Agents	61
Abacavir sulphate with		Poisonings	Anaesthetics	106
lamivudine	85	Air Flow Products	Anagrelide hydrochloride	134
Abciximab	151	Ajmaline	Analgesics	109
Abilify	121	Alanase	Anastrozole	144
Abiraterone acetate	143	Albendazole	Andriol Testocaps	61
Acarbose	16	Alendronate sodium	Androderm	61
Accu-Chek Ketur-Test	221	Alendronate sodium with	Androgen Agonists and	
Accu-Chek Performa	221	cholecalciferol	Antagonists	61
Accuretic 10	39	Alfacalcidol	Anexate	189
Accuretic 20	39	Alfentanil	Anoro Ellipta	179
Acetazolamide	186	Alinia	Antabuse	130
Acetic acid		Alitraq	Antacids and Antiflatulents	13
Extemporaneous	197	Allersoothe	Anti-Infective Agents	57
Genito-Urinary	57	Allopurinol	Anti-Infective Preparations	
Acetic acid with hydroxyquinoline,		Alpha tocopheryl acetate	Dermatological	51
glycerol and ricinoleic acid	57	Alpha-Adrenoceptor Blockers	Sensory	183
Acetic acid with propylene		Alprazolam	Anti-Inflammatory	
glycol	188	Alprostadil hydrochloride	Preparations	184
Acetylcholine chloride	187	Alteplase	Antiacne Preparations	52
Acetylcysteine	189	Alum	Antiallergy Preparations	176
Aciclovir		Aluminium chloride	Antianaemics	27
Infection	91	Aluminium hydroxide	Antiarrhythmics	41
Sensory	183	Aluminium hydroxide with	Antibacterials	71
Aciclovir-Claris	91	magnesium hydroxide and	Anticholinergic Agents	177
Acid Citrate Dextrose A	32	simethicone	Anticholinesterases	95
Acidex	13	Amantadine hydrochloride	Antidepressants	113
Acipimox	47	AmBisome	Antidiarrhoeals and Intestinal	
Acitretin	55	Ambrisentan	Anti-Inflammatory Agents	13
Aclasta	97	Amethocaine	Antiepilepsy Drugs	114
Act-HIB	215	Nervous	Antifibrinolytics, Haemostatics	
Actemra	171	Sensory	and Local Sclerosants	29
Actinomycin D	132	Amikacin	Antifungals	78
Adalimumab	151	Amiloride hydrochloride	Antihypotensives	41
Adapalene	52	Amiloride hydrochloride with	Antimigraine Preparations	119
Adefin XL	43	furosemide	Antimycobacterials	80
Adefovir dipivoxil	87	Amiloride hydrochloride with	Antinaus	120
Adenosine	41	hydrochlorothiazide	Antinausea and Vertigo	
Adenuric	101	Aminophylline	Agents	119
Adrenaline	48	Amiodarone hydrochloride	Antiparasitics	81
ADT Booster	214	Amisulpride	Antipruritic Preparations	52
Adult diphtheria and tetanus		Amitriptyline	Antipsychotic Agents	120
vaccine	214	Amiodipine	Antiretrovirals	83
Advantan	54	Amorolfine	Antirheumatoid Agents	95
Advate	31	Amoxicillin	Antiseptics and	
Aerrane	106	Amoxicillin with clavulanic	Disinfectants	191
Afinitor	174	acid	Antispasmodics and Other	
AFT SLS-free	53	Amphotericin B	Agents Altering Gut	

Generic Chemicals and Brands

Motility	15	Various	194	ATGAM	174
Antithrombotics	32	Argipressin [Vasopressin]	70	Ativan	125
Antithymocyte globulin (equine)	174	Aripiprazole	121	Atomoxetine	127
Antithymocyte globulin (rabbit)	174	Aristocort	54	Atorvastatin	46
Antiulcerants	15	Aromasin	145	Atovaquone with proguanil hydrochloride	82
Antivirals	87	Arrow - Clopid	34	Atracurium besylate	102
Anxiolytics	124	Arrow-Amitriptyline	113	Atripila	85
Apidra	17	Arrow-Bendroflumazide	45	Atropine sulphate	
Apidra Solostar	17	Arrow-Brimonidine	187	Cardiovascular	41
Apo-Allopurinol	100	Arrow-Calcium	22	Sensory	187
Apo-Amiloride	45	Arrow-Diazepam	125	Atropt	187
Apo-Amlodipine	43	Arrow-Dortin	186	Aubagio	125
Apo-Amoxi	74	Arrow-Etidronate	97	Augmentin	74
Apo-Azithromycin	73	Arrow-Fluoxetine	114	Auranofin	95
Apo-Ciclopirox	51	Arrow-Gabapentin	115	Avelox	75
Apo-Cilazapril/ Hydrochlorothiazide	39	Arrow-Iloprost	50	Avelox IV 400	75
Apo-Clarithromycin	73	Arrow-Lamotrigine	117	Avonex	126
Apo-Clomipramine	113	Arrow-Losartan & Hydrochlorothiazide	40	Avonex Pen	126
Apo-Diclo SR	103	Arrow-Morphine LA	111	Azacitidine	133
Apo-Diltiazem CD	44	Arrow-Norfloxacin	75	Azactam	76
Apo-Doxazosin	40	Arrow-Ornidazole	82	Azamun	174
Apo-Folic Acid	28	Arrow-Quinapril 10	39	Azathioprine	174
Apo-Imiquimod Cream 5%	56	Arrow-Quinapril 20	39	Azithromycin	73
Apo-Megestrol	143	Arrow-Quinapril 5	39	Azol	64
Apo-Metoprolol	42	Arrow-Roxithromycin	74	AZT	85
Apo-Mirtazapine	113	Arrow-Sertraline	114	Aztreonam	76
Apo-Moclobemide	113	Arrow-Simba	46		
Apo-Nadolol	42	Arrow-Sumatriptan	119	- B -	
Apo-Nicotinic Acid	47	Arrow-Timolol	186	B-D Micro-Fine	221
Apo-Oxybutynin	60	Arrow-Tolterodine	60	B-D Ultra Fine	221
Apo-Perindopril	39	Arrow-Topiramate	118	B-D Ultra Fine II	221
Apo-Pindolol	42	Arrow-Tramadol	112	Bacillus calmette-guerin (BCG)	174
Apo-Prazosin	40	Arrow-Venlafaxine XR	114	Bacillus calmette-guerin vaccine	215
Apo-Prednisone	63	Arsenic trioxide	134	Baclofen	102
Apo-Prednisone S29	63	Artemether with lumefantrine	81	Bacterial and Viral Vaccines	214
Apo-Propranolol	43	Artunate	82	Bacterial Vaccines	214
Apo-Pyridoxine	25	Articaine hydrochloride	107	Balanced Salt Solution	185
Apo-Ropinirole	106	Articaine hydrochloride with adrenaline	107	Baraclude	88
Apomine	105	Asacol	14	Barium sulphate	193
Apomorphine hydrochloride	105	Asamax	14	Barium sulphate with sodium bicarbonate	193
Apraclonidine	187	Ascorbic acid		Barrier Creams and Emollients	52
Aprepitant	119	Alimentary	26	Basiliximab	158
Apresoline	49	Extemporaneous	197	BCG Vaccine	215
Aprotinin	29	Aspen Adrenaline	48	BD PosiFlush	37
Aqueous cream	53	Aspirin		Beclazone 100	179
Arachis oil [Peanut oil]	197	Blood	34	Beclazone 250	179
Arava	95	Nervous	109	Beclazone 50	179
Aremed	144	Asthalin	179	Beclomethasone dipropionate	176, 179
Arginine		Atazanavir sulphate	86		
Alimentary	20	Atenolol	42		
		Atenolol-AFT	42		

Bee venom176	Bile and Liver Therapy16	Buprenorphine with naloxone129
Bendroflumethiazide45	Biliscopin194	Bupropion hydrochloride130
Bendroflumethiazide [Bendroflumethiazide]45	Bimatoprost187	Burinex44
BeneFIX30	Biodone111	Buscopan15
Benzathine benzylpenicillin74	Biodone Extra Forte111	Buserelin65
Benzbromaron AL 100100	Biodone Forte111	Buspirone hydrochloride125
Benzbromarone100	Biotin21	Busulfan132
Benzocaine107	Bisacodyl20	Butacort Aqueous177
Benzoin197	Bismuth subgallate197	
Benzoyl peroxide52	Bismuth subnitrate and iodoform paraffin195	- C -
Benztrop105	Bismuth trioxide16	Cabergoline64
Benztropine mesylate105	Bisoprolol fumarate42	Caffeine127
Benzydamine hydrochloride23	Bivalirudin32	Caffeine citrate181
Benzydamine hydrochloride with cetylpyridinium chloride23	Bleomycin sulphate132	Calamine52
Benzylpenicillin sodium [Penicillin G]74	Blood glucose diagnostic test meter221	Calcipotriol55
Beractant182	Blood glucose diagnostic test strip221	Calcitonin61
Beta Cream54	Blood ketone diagnostic test meter221	Calcitriol26
Beta Ointment54	Boceprevir90	Calcitriol-AFT26
Beta Scalp55	Boenney's blue dye194	Calcium carbonate13, 22
Beta-Adrenoceptor Agonists179	Boostrix215	Calcium Channel Blockers43
Beta-Adrenoceptor Blockers42	Boric acid197	Calcium chloride35
Betadine191	Bortezomib134	Calcium folinate142
Betadine Skin Prep191	Bosentan49	Calcium Folate Ebewe142
Betagan186	Bosvate42	Calcium gluconate Blood35
Betahistine dihydrochloride119	Botox102	Dermatological56
Betaine20	Botulism antitoxin189	Calcium Homeostasis61
Betamethasone62	Breox Ellipta181	Calcium polystyrene sulphonate38
Betamethasone dipropionate54	Bridion102	Calcium Resonium38
Betamethasone dipropionate with calcipotriol55	Brilinta34	Calsource22
Betamethasone sodium phosphate with betamethasone acetate62	Brimonidine tartrate187	Cancidas79
Betamethasone valerate54-55	Brimonidine tartrate with timolol187	Candesartan cilexetil40
Betamethasone valerate with clioquinol55	Brinzolamide186	Candestar40
Betamethasone valerate with fusidic acid55	Bromocriptine105	Capecitabine133
Betaxolol186	Brufen SR103	Capecitabine Winthrop133
Betoptic186	Budesonide Alimentary13	Capoten39
Betoptic S186	Respiratory177, 180	Capsaicin Musculoskeletal104
Bevacizumab158	Budesonide with efomedoterol181	Nervous109
Bezafibrate45	Bumetanide44	Captopril39
Bezalip45	Bupafen108	Carbamazepine115
Bezalip Retard45	Bupivacaine hydrochloride107	Carbasorb-X190
Bicalaccord143	Bupivacaine hydrochloride with adrenaline107	Carbimazole69
Bicalutamide143	Bupivacaine hydrochloride with fentanyl108	Carbomer188
Bicillin LA74	Bupivacaine hydrochloride with glucose108	Carboplatin137
BiCNU132		Carboprost trometamol58
		Carboxymethylcellulose Alimentary23
		Extemporaneous197
		Cardinol LA43
		Cardizem CD44
		CareSens221

Caresens II	221	cetrimide	191, 195	Clinicians Renal Vit	24
CareSens N	221	Chlorhexidine with ethanol	191	Clobazam	115
Caresens N	221	Chloroform	197	Clobetasol BNM	54
Caresens N POP	221	Chloroquine phosphate	82	Clobetasol propionate	54, 56
Carmellose sodium	188	Chlorothiazide	45	Clobetasone butyrate	54
Carmustine	132	Chlorpheniramine maleate	177	Clofazimine	80
Carvedilol	42	Chlorpromazine		Clomazol	51, 57
Caspofungin	79	hydrochloride	121	Clomiphene citrate	64
Catapres	44	Chlorsig	183	Clomipramine hydrochloride	113
Catapres-TTS-1	44	Chlortalidone		Clonazepam	114–115, 125
Catapres-TTS-2	44	[Chlortalidone]	45	Clonidine	44
Catapres-TTS-3	44	Chlortalidone	45	Clonidine BNM	44
Ceenu	132	Choice Load 375	57	Clonidine hydrochloride	44
Cefaclor	72	Cholecalciferol	26	Clopidogrel	34
Cefalexin	72	Cholestyramine	46	Clopine	121
Cefalexin Sandoz	72	Choline salicylate with		Clopixol	123, 124
Cefazolin	72	cetalkonium chloride	23	Clostridium botulinum type A	
Cefepime	72	Cholvastin	46	toxin	102
Cefepime-AFT	72	Choriogonadotropin alfa	65	Clotrimazole	
Cefotaxime	72	Ciclopirox olamine	51	Dermatological	51
Cefotaxime Sandoz	72	Ciclosporin	145	Genito-Urinary	57
Cefoxitin	72	Cidofovir	91	Clove oil	197
Ceftaroline fosamil	73	Cilazapril	39	Clozapine	121
Ceftazidime	72	Cilazapril with		Clozaril	121
Ceftriaxone	72	hydrochlorothiazide	39	Co-trimoxazole	77
Ceftriaxone-AFT	72	Cilicaine	75	Coal tar	197
Cefuroxime	72	Cilicaine VK	74	Coal tar with salicylic acid and	
Celecoxib	103	Cimetidine	15	sulphur	55
Celiprolol	42	Cinacalcet	61	Cocaine hydrochloride	108
CelCept	175	Cinchocaine hydrochloride with		Cocaine hydrochloride with	
Celol	42	hydrocortisone	14	adrenaline	108
Centrally-Acting Agents	44	Cipflox	75	Codeine phosphate	
Cephalexin ABM	72	Ciprofloxacin		Extemporaneous	197
Cetirizine hydrochloride	177	Infection	75	Nervous	110
Cetomacrogol	53	Sensory	183	Cogentin	105
Cetomacrogol with glycerol	53	Ciprofloxacin with		Colaspase [L-asparaginase]	135
Cetrimide	197	hydrocortisone	183	Colchicine	101
Champix	131	Ciproxin HC Otic	183	Colestimethate	76
Charcoal	190	Cisplatin	137	Colestipol hydrochloride	46
Chemotherapeutic Agents	132	Citalopram hydrobromide	114	Colgout	101
Chicken pox vaccine	220	Citanest	109	Colifoam	14
Chlorafast	183	Citric acid	197	Colistin sulphomethate	
Chloral hydrate	126	Citric acid with magnesium oxide		[Colestimethate]	76
Chlorambucil	132	and sodium picosulfate	19	Colistin-Link	76
Chloramphenicol		Citric acid with sodium		Collodion flexible	197
Infection	76	bicarbonate	193	Colloidal bismuth subcitrate	16
Sensory	183	Cladribine	133	Colofac	15
Chlorhexidine	191, 195	Clarithromycin	73	Colony-Stimulating Factors	35
Chlorhexidine gluconate		Clexane	32	Coloxyl	19, 20
Alimentary	23	Clindamycin	76	Compound electrolytes	35, 38
Extemporaneous	197	Clindamycin ABM	76	Compound electrolytes with	
Genito-Urinary	57	Clinicians Multivit & Mineral		glucose	35, 38
Chlorhexidine with		Boost	24	Compound	

hydroxybenzoate	197	Danaparoid	32	Desflurane	106
Compound sodium lactate		Danazol	64	Desmopressin acetate	70
[Hartmann's solution]	36	Dantrium	102	Desmopressin-PH&T	70
Compound sodium lactate with		Dantrium IV	102	Dexamethasone	
glucose	36	Dantrolene	102	Hormone	62
Concerta	128	Dapa-Tabs	45	Sensory	184
Condyline	56	Dapsone		Dexamethasone phosphate	62
Contraceptives	57	Contracted	80	Dexamethasone with framycetin	
Contrast Media	192	Infection	80	and gramicidin	183
Cordarone-X	41	Daptomycin	76	Dexamethasone with neomycin	
Corticosteroids		Darunavir	86	sulphate and polymyxin B	
Dermatological	54	Dasatinib	137	sulphate	184
Hormone	62	Daunorubicin	132	Dexamethasone with	
Corticotorelin (ovine)	64	DBL Acetylcysteine	189	tobramycin	184
Cosmegen	132	DBL Amikacin	71	Dexamfetamine sulfate	127
Cough Suppressants	179	DBL Aminophylline	181	Dexmedetomidine	106
Creon 10000	18	DBL Bleomycin Sulfate	132	Dexmethsone	62
Creon 25000	18	DBL Carboplatin	137	Dextrose	
Crotamiton	52	DBL Cefotaxime	72	Alimentary	16
Crystaderm	51	DBL Cisplatin	137	Blood	36
CT Plus+	193	DBL Docetaxel	142	Extemporaneous	197
Cubicin	76	DBL Ergometrine	59	Dextrose with sodium citrate and	
Curam Duo	74	DBL Leucovorin Calcium	142	citric acid [Acid Citrate	
Curosurf	182	DBL Meropenem	72	Dextrose A]	32
Cvite	26	DBL Morphine Sulphate	111	DHC Continus	110
Cyclizine hydrochloride	119	DBL Pethidine		Diabetes	16
Cyclizine lactate	119	Hydrochloride	112	Diacomit	117
Cyclogyl	187	DBL Rocuronium Bromide	102	Diagnostic Agents	194
Cyclopentolate		DBL Sterile Dopamine		Diagnostic and Surgical	
hydrochloride	187	Concentrate	48	Preparations	185
Cyclophosphamide	132	DBL Tobramycin	71	Diamide Relief	13
Cycloserine	80	DDI	85	Diamox	186
Cyklokapron	30	De-Nol	16	Diatrizoate meglumine with	
Cymevene	91	De-Worm	81	sodium amidotrizoate	192
Cyproheptadine		Decongestants	179	Diatrizoate sodium	192
hydrochloride	177	Decongestants and		Diazepam	114, 125
Cyproterone acetate	61	Antiallergics	184	Diazoxide	
Cyproterone acetate with		Decozol	24	Alimentary	16
ethinyloestradiol	57	Deferasirox	190	Cardiovascular	48
Cysteamine hydrochloride	197	Deferiprone	190	Dicarz	42
Cytarabine	133	Defibrotide	32	Dichlorobenzyl alcohol with	
Cytotec	15	Definity	194	amylmetacresol	23
		Demeclocycline		Diclofenac Sandoz	103
- D -		hydrochloride	76	Diclofenac sodium	
D-Penamine	95	Deoxycoformycin	136	Musculoskeletal	103
Dabigatran	32	Depo-Medrol	63	Sensory	184
Dacarbazine	135	Depo-Medrol with Lidocaine	63	Dicobalt edetate	190
Dactinomycin [Actinomycin		Depo-Provera	58	Didanosine [DDI]	85
D]	132	Depo-Testosterone	61	Diflucan	78
Daivobet	55	Deprim	77	Diflucortolone valerate	54
Daivonex	55	Dermol	56	Digestives Including	
Dalacin C	76	Desferal	190	Enzymes	18
Dalteparin	32	Desferrioxamine mesilate	190	Digoxin	41

Generic Chemicals and Brands

Digoxin immune Fab	189	Dothiepin hydrochloride	113	disoproxil fumarate	85
Dihydrocodeine tartrate	110	Doxapram	182	Emtriva	85
Dihydroergotamine mesylate	119	Doxazosin	40	Emulsifying ointment	53
Diltiazem hydrochloride	44	Doxepin hydrochloride	113	Enalapril maleate	39
Dilzem	44	Doxine	76	Enalapril maleate with hydrochlorothiazide	39
Dimercaprol	190	Doxorubicin Ebewe	132	Enbrel	145
Dimercaptosuccinic acid	191	Doxorubicin hydrochloride	132	Endocrine Therapy	143
Dimethicone	52	Doxycycline	76	Endoxan	132
Dimethyl fumarate	125	DP Fusidic Acid Cream	51	Enfuvirtide	83
Dimethyl sulfoxide	195	DP Lotn HC	54	Enoxaparin	32
Dinoprostone	59	DP-Anastrozole	144	Ensure (Chocolate)	213
Diphemanil metilsulfate	56	Dr Reddy's Omeprazole	16	Ensure (Vanilla)	213
Diphenoxylate hydrochloride with atropine sulphate	13	Dr Reddy's Terbinafine	80	Ensure Plus (Banana)	213
Diphtheria antitoxin	189	Droperidol	119	Ensure Plus (Chocolate)	213
Diphtheria, tetanus and pertussis vaccine	215	Drugs Affecting Bone Metabolism	95	Ensure Plus (Fruit of the Forest)	213
Diphtheria, tetanus, pertussis and polio vaccine	214	Duolin	177	Ensure Plus (Vanilla)	213
Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine	214	Duovisc	186	Ensure Plus HN	212
Dipyridamole	34	Duride	47	Ensure Plus HN RTH	212
Disodium edetate	186	Dynastat	104	Entacapone	106
Disodium hydrogen phosphate with sodium dihydrogen phosphate	197	Dysport	102	Entapene	106
Disopyramide phosphate	41	- E -		Entecavir	88
Disulfiram	130	e-chamber La Grande	222	Enzymes	100
Dithranol	197	e-chamber Mask	221	Ephedrine	48
Diuretics	44	e-chamber Turbo	222	Epilim IV	117
Diurin 40	44	E-Mycin	73	Epirubicin Ebewe	133
Dobutamine hydrochloride	48	E-Z-Cat Dry	193	Epirubicin hydrochloride	133
Dobutamine-Clarix	48	E-Z-Gas II	193	Epoetin alfa [Erythropoietin alfa]	27
Docetaxel	142	E-Z-Paste	193	Epoetin beta [Erythropoietin beta]	28
Docusate sodium Alimentary	19	EasyCheck	221	Epoprostenol	50
Sensory	188	Econazole nitrate	51	Eprex	27
Docusate sodium with sennosides	19	Edrophonium chloride	95	Eptacog alfa [Recombinant factor VIIa]	30
Domperidon	119	Efavirenz	84	Eptifibatide	34
Donepezil hydrochloride	129	Efavirenz with emtricitabine and tenofovir disoproxil fumarate	85	Ergometrine maleate	59
Donepezil-Rex	129	Efexor XR	114	Ergotamine tartrate with caffeine	119
Dopamine hydrochloride	48	Effient	34	Erlotinib	138
Dopergin	106	Eformoterol fumarate	180	Ertapenem	71
Dopress	113	Efudix	56	Erythrocin IV	73
Dornase alfa	181	Elecare (Unflavoured)	208	Erythromycin (as ethylsuccinate)	73
Dorzolamide	186	Elecare (Vanilla)	208	Erythromycin (as lactobionate)	73
Dorzolamide with timolol	186	Elecare LCP (Unflavoured)	208	Erythromycin (as stearate)	74
Dostinex	64	Electrolytes	196	Erythropoietin alfa	27
Dotarem	193	Eligard	65	Erythropoietin beta	27
		Elocon	54	Escitalopram	114
		Elocon Alcohol Free	54	Esmolol hydrochloride	42
		Eltrombopag	29	Etanercept	145
		Emend Tri-Pack	119		
		EMLA	109		
		Emtricitabine	85		
		Emtricitabine with tenofovir			

Ethambutol hydrochloride	80	acid	22	Flupenthixol decanoate	123
Ethanol	189	Ferrous gluconate with ascorbic acid	22	Fluphenazine decanoate	123
Ethanol with glucose	189	Ferrous sulphate	22	Flutamide	143
Ethanol, dehydrated	189	Ferrous sulphate with ascorbic acid	22	Flutamin	143
Ethics Aspirin EC	34	Ferrous sulphate with folic acid	23	Fluticasone	180
Ethics Enalapril	39	Ferrum H	23	Fluticasone furoate with vilanterol	181
Ethics Lisinopril	39	Fexofenadine hydrochloride	177	Fluticasone propionate	177
Ethinylloestradiol	64	Filgrastim	35	Fluticasone with salmeterol	181
Ethinylloestradiol with desogestrel	57	Finasteride	59	FML	184
Ethinylloestradiol with levonorgestrel	57	Fingolimod	125	Foban	51
Ethinylloestradiol with norethisterone	57	Finpro	59	Folic acid	28
Ethosuximide	115	Firazyr	176	Fondaparinux sodium	33
Ethyl chloride	108	Flagyl	82	Food Modules	200
Etidronate disodium	97	Flagyl-S	82	Food/Fluid Thickeners	202
Etomidate	106	Flamazine	51	Forteo	100
Etopophos	135	Flecainide acetate	41	Fortisip (Vanilla)	213
Etoposide	135	Fleet Phosphate Enema	20	Fortum	72
Etoposide (as phosphate)	135	Flixonase Hayfever & Allergy	177	Fosamax	95, 96
Etoricoxib	103	Flixotide	180	Fosamax Plus	96
Etravirine	84	Flixotide Accuhaler	180	Foscarnet sodium	91
Everet	117	Floair	180	Fosfomycin	76
Everolimus	174	Florinef	62	Fragmin	32
Evista	99	Fluanxol	123	Framycetin sulphate	183
Exelon	129	Fluarix	218	Freeflex	37
Exemestane	145	Flucloxacillin	74	FreeStyle Lite	221
Exjade	190	Flucloxin	74	Freestyle Optium	221
Extemporaneously Compounded Preparations	197	Fluconazole	78	Freestyle Optium Ketone	221
Ezemibe	46	Fluconazole-Claris	78	Freestyle Optium Neo	221
Ezetimibe	46	Flucytosine	80	Fresofol 1% MCT/LCT	107
Ezetimibe with simvastatin	46	Fludara Oral	134	frusemide	44
- F -		Fludarabine Ebewe	134	Frusamide-Claris	44
Factor eight inhibitor bypassing fraction	30	Fludarabine phosphate	134	Fucidin	77
Febuxostat	101	Fludrocortisone acetate	62	Fucithalmic	183
FEIBA NF	30	Fluids and Electrolytes	35	Fungilin	24
Felodipine	43	Flumazenil	189	Furosemide [frusemide]	44
Fenpaed	103	Flumetasone pivalate with clioquinol	184	Fusidic acid	
Fentanyl	110	Fluocortolone caproate with fluocortolone pivalate and cinchocaine	14	Dermatological	51
Fentanyl Sandoz	110	Fluorescein sodium	185	Infection	77
Ferinject	22	Fluorescein sodium with lignocaine hydrochloride	185	Sensory	183
Ferodan	22	Fluorescite	185	Fuzeon	83
Ferric carboxymaltose	22	Fluorometholone	184	- G -	
Ferric subsulfate	29	Fluorouracil	134	Gabapentin	115
Ferriprox	190	Fluorouracil Ebewe	134	Gacet	110
Ferro-F-Tabs	22	Fluorouracil sodium	56	Gadobenid acid	193
Ferro-tab	22	Fluoxetine hydrochloride	114	Gadobutrol	193
Ferrograd	22			Gadodiamide	193
Ferrous fumarate	22			Gadoterid acid	193
Ferrous fumarate with folic acid	22			Gadovist	193
				Gadoxetate disodium	194
				Galsulfate	21
				Ganciclovir	
				Infection	91

Sensory	183	Glycine	195	Alimentary	24
Gardasil	218	Glycopyrronium	178	Sensory	185, 186, 188
Gastrodenol	16	Glycopyrronium bromide	15	Hyaluronidase	100
Gastrografin	192	Glycopyrronium with		Hybloc	42
Gastrosoothe	15	indacaterol	178	Hydralazine	
Gefitinib	138	Glypressin	70	hydrochloride	48–49
Gelatine, succinylated	38	Glytrin	47	Hydrea	135
Gelofusine	38	Gonadorelin	65	Hydrocortisone	
Gemcitabine	134	Goserelin	65	Dermatological	54
Gemcitabine Ebewe	134	Granirex	120	Extemporaneous	198
Gemfibrozil	46	Granisetron	120	Hormone	62
Genoptic	183			Hydrocortisone acetate	
Genox	144	- H -		Alimentary	14
Gentamicin sulphate		Habitrol	130	Dermatological	54
Infection	71	Habitrol (Classic)	130	Hydrocortisone and paraffin	
Sensory	183	Habitrol (Fruit)	130	liquid and lanolin	54
Gestrinone	64	Habitrol (Mint)	130	Hydrocortisone butyrate	54, 56
Gilenya	125	Haem arginate	21	Hydrocortisone with	
Ginet	57	Haemophilus influenzae type B		miconazole	55
Glatiramer acetate	126	vaccine	215	Hydrocortisone with natamycin	
Glaucoma Preparations	186	Haldol	123	and neomycin	55
Glibenclamide	18	Haldol Concentrate	123	Hydrocortisone with paraffin and	
Gliclazide	18	Haloperidol	121	wool fat	54
Glipizide	18	Haloperidol decanoate	123	Hydrogen peroxide	51
Glivec	138	Hameln	110	Hydroxocobalamin	
Glizide	18	Hartmann's solution	35	Alimentary	25
Glucagen Hypokit	16	Havrix	217	Various	189
Glucagon hydrochloride	16	Havrix Junior	217	Hydroxychloroquine	95
Glucerna Select (Vanilla)	205	HBvaxPRO	217, 218	Hydroxyurea	135
Glucerna Select RTH		Healon GV	186	Hygroton	45
(Vanilla)	205	healthE Dimethicone 10%	52	Hylo-Fresh	188
Glucobay	16	healthE Dimethicone 5%	52	Hyoscine butylbromide	15
Glucose [Dextrose]		healthE Fatty Cream	53	Hyoscine hydrobromide	120
Alimentary	17	Heparin sodium	33	Hyperuricaemia and	
Blood	36	Heparinised saline	33	Antigout	100
Extemporaneous	198	Heparon Junior	206	Hypnovel	126
Glucose with potassium		Hepatitis A vaccine	217	Hypromellose	185, 188
chloride	36	Hepatitis B recombinant		Hypromellose with dextran	188
Glucose with potassium chloride		vaccine	217–218	Hysite	187
and sodium chloride	36	Hepsera	87		
Glucose with sodium chloride	36	Herceptin	173	- I -	
Glucose with sucrose and		Hexamine hippurate	77	Ibiamox	74
fructose	17	Histaclear	177	Ibuprofen	103
Glycerin with sodium		Histamine acid phosphate	194	Icatibant	176
saccharin	198	Holoxan	132	Idarubicin hydrochloride	133
Glycerin with sucrose	198	Hormone Replacement		Ifosfamide	132
Glycerol		Therapy	63	Ikorel	49
Alimentary	20	HPV	218	Iloprost	50
Extemporaneous	198	Humalog Mix 25	17	Imatinib mesilate	138–139
Glycerol with paraffin	53	Humalog Mix 50	17	Imatinib-AFT	139
Glyceryl trinitrate		Humatin	71	Imiglucerase	21
Alimentary	15	Humira	151	Imipenem with cilastatin	71
Cardiovascular	47	HumiraPen	151	Imipenem+Cilastatin RBX	71
		Hyaluronic acid			

Imipramine hydrochloride	113	Ioscan	192	Konakion MM	31
Imiquimod	56	IPOL	219	Konsyl-D	19
Immune Modulators	92	Ipratropium bromide	177		
Immunosuppressants	145	Iressa	138	- L -	
Impact Advanced Recovery (Chocolate)	212	Irinotecan hydrochloride	135	L-asparaginase	135
Impact Advanced Recovery (Vanilla)	212	Iron polymaltose	23	L-ornithine L-aspartate	16
Imuran	174	Iron sucrose	23	Labetalol	42
Incruse Ellipta	178	Irrigation Solutions	195	Lacosamide	116
Indacaterol	180	Isentress	87	Lactose	198
Indapamide	45	Ismo-20	47	Lactulose	20
Indigo carmine	194	Isoflurane	106	Laevolac	20
Indinavir	86	Isoniazid	80	Lamictal	117
Indocyanine green	194	Isoniazid with rifampicin	80	Lamivudine	85, 88
Indomethacin	104	Isoprenaline	48	Lamotrigine	117
Infanrix IPV	214	Isopropyl alcohol	191	Lanoxin	41
Infanrix-hexa	214	Isoptin	44	Lanoxin PG	41
Infliximab	158	Isopto Carpine	187	Lansoprazole	15
Influenza vaccine	218	Isosorbide mononitrate	47	Lantus	17
Influvac	218	Isotane 10	52	Lantus SoloStar	17
Inhaled Corticosteroids	179	Isotane 20	52	Lanzol Relief	15
Insulin aspart	17	Isotretinoin	52	Lapatinib	139
Insulin aspart with insulin aspart protamine	17	Ispaghula (psyllium) husk	19	Lariam	82
Insulin glargine	17	Isradipine	43	Latanoprost	187
Insulin glulisine	17	Itch-Soothe	52	Lax-Sachets	20
Insulin isophane	17	Itraconazole	78	Lax-Suppositories	20
Insulin lispro	17	Itrazole	78	Lax-Tabs	20
Insulin lispro with insulin lispro protamine	17	Ivermectin	81	Laxatives	19
Insulin neutral	18	- J -		Laxsol	19
Insulin neutral with insulin isophane	17	Jadelle	58	Leflunomide	95
Insulin pen needles	221	Jaychem	53	Lenalidomide	135
Insulin syringes, disposable with attached needle	221	Jevity	213	Letrole	145
Integrilin	34	Jevity HiCal RTH	212	Letrozole	145
Intelence	84	Jevity RTH	213	Leukotriene Receptor Antagonists	180
Interferon alfa-2a	92	- K -		Leunase	135
Interferon alfa-2b	92	Kaletra	86	Leuprorelin acetate	65
Interferon beta-1-alpha	126	Kenacomb	184	Leustatin	133
Interferon beta-1-beta	126	Kenacort-A 10	63	Levetiracetam	117
Interferon gamma	92	Kenacort-A 40	63	Levetiracetam-Rex	117
Intra-uterine device	57	Kenalog in Orabase	23	Levobunolol hydrochloride	186
Invanz	71	Ketamine	106	Levocabastine	184
Invega Sustenna	123	Ketamine-Claris	106	Levocarnitine	21
Iodine	69	Ketoconazole	51	Levodopa with benserazide	106
Iodine with ethanol	191	Dermatological	51	Levodopa with carbidopa	106
Iodised oil	192	Infection	78	Levomepromazine	122
Iodixanol	192	Ketone blood beta-ketone electrodes	221	Levonorgestrel	58
Iohexol	192	Ketoprofen	104	Levosimendan	47
Iopidine	187	Ketorolac trometamol	184	Levothyroxine	69
		Kivexa	85	Lidocaine [lignocaine]	109
		Klacid	73	Lidocaine [Lignocaine] hydrochloride	108
		Klean Prep	19	Lidocaine [Lignocaine] hydrochloride with adrenaline	108
		Kogenate FS	31		

Generic Chemicals and Brands

Lidocaine [Lignocaine]	Losartan potassium with	Medroxyprogesterone acetate
hydrochloride with adrenaline	hydrochlorothiazide	Genito-Urinary
and tetracaine40	Hormone
hydrochloride	Lovir64
108	Loxamine	Mefenamic acid
Lidocaine [Lignocaine]114	104
hydrochloride with	Lucrin Depot PDS	Mefloquine
chlorhexidine65	82
108	Lycinate	Megestrol acetate
Lidocaine [Lignocaine]47	143
hydrochloride with	Lyderm	Meglumine gadopentetate
phenylephrine		194
hydrochloride	- M -	Meglumine iotroxate
108	m-Amoxiclav	194
Lidocaine [Lignocaine] with74	Melatonin
prilocaine	m-Eslon	126
109111	Meloxicam
Lidocaine-Clarix	M-M-R-II	104
108219	Melphalan
Lignocaine	m-Nystatin	132
108	Mabthera	215
lignocaine	Madopar 125	Menactra
Hormone106	Meningococcal (A, C, Y and
63	Madopar 250	W-135) conjugate
Nervous106	vaccine
109	Madopar 62.5	215
Lincomycin106	Meningococcal C conjugate
77	Madopar HBS	vaccine
Linezolid106	216
77	Madopar Rapid	Menthol
Lioresal Intrathecal	Mafenide acetate	198
10251	Mepivacaine hydrochloride
Liothyronine sodium	Magnesium hydroxide	109
69	Alimentary	Mercaptopurine
Lipazil23	134
46	Extemporaneous	Meropenem
Lipid-Modifying Agents198	72
45	Magnesium oxide	Mesalazine
Lipiodol Ultra Fluid23	14
192	Magnesium sulphate	Mesna
Liquibar23	142
193	Magnevist	Mestinon
Lisinopril194	95
39	Malarone	Metabolic Disorder Agents
Lissamine green82	20
185	Malarone Junior	Metabolic Products
Lisuride hydrogen maleate82	202
106	Malathion [Maldison]	Metamide
Lithicarb FC51	120
122	Malathion with permethrin and	Metaraminol
Lithium carbonate	piperonyl butoxide	48
12252	Metchek
LMX4	Maldison	18
10951	Meterol
Local Preparations for Anal and	Mannitol	181
Rectal Disorders	Cardiovascular	Methformin hydrochloride
1445	18
Locoid	Various	Methacholine chloride
54, 56194	194
Locoid Crelo	Maprotiline hydrochloride	Methadone hydrochloride
54113	Extemporaneous
Locoid Lipocream	Marcaïn	Nervous
54107	111
Lodoxamide	Marcaïn Heavy	Methatabs
184108	111
Logem	Marcaïn Isobaric	Methohexital sodium
117107	107
Lomide	Marcaïn with Adrenaline	Methopt
184107	188
Lomustine	Marevan	Methotrexate
13233	134
Long-Acting Beta-Adrenoceptor	Marine Blue Lotion SPF 50+	Methotrexate Ebewe
Agonists56	134
180	Mask for spacer device	Methotrexate Sandoz
Loniten221	134
49	Mast Cell Stabilisers	Methoxsalen
Loperamide hydrochloride181	[8-methoxypsoralen]
13	Max Health55
Lopinavir with ritonavir	15, 48, 62, 95	Methoxyflurane
86	Maxidex	109
Lopresor184	Methyl aminolevulinate
42	Maxitrol	hydrochloride
Lorafix184	56
LoraPaed	Measles, mumps and rubella	Methyl hydroxybenzoate
177	vaccine	198
Loratadine219	Methylcellulose
177	Mebendazole	198
Lorazepam81	Methylcellulose with glycerin and
115, 125	Mebeverine hydrochloride	sodium saccharin
12615	198
Losartan potassium	Medrol	Methylcellulose with glycerin and
4062	sucrose
	Medroxyprogesterone	198
64	Methyldopa
		44

Methylene blue	194	Mitozantrone Ebewe	133	Naprosyn SR 750	104
Methylphenidate hydrochloride	128	Mivacron	102	Naproxen	104
Methylprednisolone (as sodium succinate)	62	Mivacurium chloride	102	Naropin	109
Methylprednisolone aceponate	54	Mixed salt solution for eye irrigation	185	Natalizumab	125
Methylprednisolone acetate	63	Moclobemide	113	Natamycin	183
Methylprednisolone acetate with lidocaine [lignocaine]	63	Modafinil	128	Natulan	136
Methylthionium chloride [Methylene blue]	194	Moderate	123	Nausicalm	119
Methylxanthines	181	Mometasone furoate	54	Nauzene	119
Metoclopramide hydrochloride	120	Monosodium glutamate with sodium aspartate	196	Navelbine	142
Metoclopramide hydrochloride with paracetamol	119	Monosodium l-aspartate	196	Nedocromil	181
Metolazone	45	Montelukast	180	Nefopam hydrochloride	110
Metoprolol - AFT CR	42	Moroctocog alfa [Recombinant factor VIII]	30	Neisvac-C	216
Metoprolol succinate	42	Morphine hydrochloride	111	Neo-B12	25
Metoprolol tartrate	42	Morphine sulphate	111	Neocate Advance (Vanilla)	208
Metronidazole Dermatological	51	Morphine tartrate	111	Neocate Gold (Unflavoured)	208
Metronidazole Infection	82	Motetis	105	Neoral	145
Metyrapone	64	Mouth and Throat	23	Neostigmine metilsulfate	95
Mexiletine hydrochloride	41	Movapo	105	Neostigmine metilsulfate with glycopyrronium bromide	95
Mexiletine Hydrochloride USP	41	Moxifloxacin	75	Neosynephrine HCL	48
Miacalcic	61	Mucolytics and Expectorants	181	Nepro HP (Strawberry)	211
Mianserin hydrochloride	113	Multihance	193	Nepro HP (Vanilla)	211
Micolette	20	Multiple Sclerosis Treatments	125	Nepro HP RTH	211
Miconazole	24	Multivitamin and mineral supplement	24	Neulastim	35
Miconazole nitrate Dermatological	51	Multivitamin renal	24	Neupogen	35
Miconazole nitrate Genito-Urinary	57	Multivitamins	24-25	Neurontin	115
Micreme	57	Mupirocin	51	NeuroTabs	22
Micreme H	55	Muscle Relaxants and Related Agents	102	Nevirapine	84
Midazolam	126	Myambutol	80	Nevirapine Alphapharm	84
Midodrine	41	Mycobutin	81	Nicardipine hydrochloride	43
Mifepristone	58	MycoNail	51	Nicorandil	49
Milrinone	49	Mycophenolate mofetil	175	Nicotine	130
Milrinone Generic Health	49	Mydriacyl	187	Nicotinic acid	47
Minerals	22	Mydriatics and Cycloplegics	187	Nifedipine	43
Mini-Wright AFS Low Range	221	Mylan Atenolol	42	Nilotinib	139
Mini-Wright Standard	221	Mylan-Bosentan	49	Nilstat	78
Minidiab	18	Myleran	132	Nimodipine	43
Minirin	70	- N -		Nitazoxanide	82
Minocycline	76	Nadolol	42	Nitrados	126
Minoxidil	49	Naglazyme	21	Nitrates	47
Mirtazapine	113	Naloxone hydrochloride	189	Nitrazepam	126
Misoprostol	15	Naltracord	130	Nitroderm TTS 10	47
Mitomycin C	133	Naltrexone hydrochloride	130	Nitroderm TTS 5	47
Mitozantrone	133	Naphazoline hydrochloride	185	Nitrofurantoin	77
		Naphcon Forte	185	Nitrolingual Pump Spray	47
		Naprosyn SR 1000	104	Nitronal	47
				Noflam 250	104
				Noflam 500	104
				Non-Steroidal Anti-Inflammatory Drugs	103
				Nonacog alfa [Recombinant factor IX]	30
				Nonacog gamma, [Recombinant	

Pediasure (Vanilla)	210	Pimafucort	55	Povidone K30	198
Pediasure RTH	210	Pindolol	42	Povidone-iodine	191
Pegaspargase	136	Pine tar with tolramine laurilsulfate and fluorescein	55	Povidone-iodine with ethanol	191
Pegasys	93	Pinetarsol	55	Pradaxa	32
Pegasys RBV Combination Pack	93	Pioglitazone	18	Pralidoxime iodide	189
Pegfilgrastim	35	Piperacillin with tazobactam	74	Pramipexole hydrochloride	106
Pegylated interferon alfa-2a	93	Pipithiazine palmitate	124	Prasugrel	34
Penicillamine	95	Pituitary and Hypothalamic Hormones and Analogues	64	Pravastatin	46
Penicillin G	74	Pivmecillinam	77	Praziquantel	81
Penicillin V	74	Pizotifen	119	Prazosin	40
Pentacarinat	82	PKU Anamix Junior LQ (Berry)	203	Precedex	106
Pentagastrin	64	PKU Anamix Junior LQ (Orange)	203	Prednisolone	63
Pentamidine isethionate	82	PKU Anamix Junior LQ (Unflavoured)	203	Prednisolone acetate	184
Pentasa	14	Plaquenil	95	Prednisolone sodium phosphate	184
Pentostatin [Deoxycofomycin]	136	Plendil ER	43	Prednisona	63
Pentoxifylline [Oxpentifylline]	49	Pneumococcal (PCV13) conjugate vaccine	216	Pregnancy test - hCG urine	221
Peptisoothe	15	Pneumococcal (PPV23) polysaccharide vaccine	216	preOp	212
Perfalgan	110	Pneumovax 23	216	Prevenar 13	216
Perflutren	194	Podophyllotoxin	56	Prezista	86
Perhexiline maleate	44	Polidocanol	29	Prilocaine hydrochloride	109
Pericyazine	122	Poliomyelitis vaccine	219	Prilocaine hydrochloride with felypressin	109
Perindopril	39	Poloxamer	20	Primaquine phosphate	82
Permethrin	52	Poly Gel	188	Primidone	117
Peteha	81	Poly-Tears	188	Primolut N	64
Pethidine hydrochloride	112	Poly-Visc	188	Primovist	194
Pexsig	44	Polyhexamethylene biguanide	198	Probenecid	102
Phenelzine sulphate	113	Polyvinyl alcohol	188	Procaine penicillin	75
Phenindione	33	Polyvinyl alcohol with povidone	188	Procaine hydrochloride	136
Phenobarbitone	117, 126	Poractant alfa	182	Prochlorperazine	120
Phenobarbitone sodium	198	Posaconazole	78	Proctosedyl	14
Phenol		Postinor-1	58	Procur	61
Extemporaneous	198	Potassium chloride	36, 38	Procyclidine hydrochloride	105
Various	195	Potassium chloride with sodium chloride	37	Procytox	132
Phenol oily	15	Potassium citrate	60	Prodopa	44
Phenol with ioxaglic acid	195	Potassium dihydrogen phosphate	37	Progesterone	59
Phenoxybenzamine hydrochloride	40	Potassium iodate Alimentary	22	Proglider	16
Phenoxyethylpenicillin [Penicillin V]	74	Potassium iodate with iodine	22	Proglycem	16
Phentolamine mesylate	40	Potassium perchlorate	69	Prognova	63
Phenylephrine hydrochloride Cardiovascular	48	Potassium permanganate	55	Prokinex	119
Sensory	187			Promethazine hydrochloride	177
Phenytoin	117			Promethazine theoclate	120
Phenytoin sodium	115, 117			Propafenone hydrochloride	41
Pholcodine	179			Propamidine isethionate	183
Phosphorus	38			Propofol	107
Phytomenadione	31			Propranolol	43
Picibanil	175			Propylene glycol	198
Pilocarpine hydrochloride	187			Propylthiouracil	70
Pilocarpine nitrate	198			Prostin E2	59
				Prostin VR	48
				Protamine sulphate	33

Protionamide	81	Relenza Rotadisk	92	vaccine	219
Protirelin	70	Remicade	158	Roxane	13
Provera	64	Remifentanil hydrochloride	112	Roxithromycin	74
Provisc	186	ReoPro	151	Rubifen	128
Provine MCT-LCT 1%	107	Resonium A	38	Rubifen SR	128
Proxymetacaine		Resource Beneprotein	201		
hydrochloride	185	Resource Diabetic (Vanilla)	205	- S -	
Pseudoephedrine		Respiratory Stimulants	182	SalAir	179
hydrochloride	179	Retinol	25	Salamol	179
Psoriasis and Eczema		Retinol Palmitate	188	Salazopyrin	14
Preparations	55	Retrovir	85	Salazopyrin EN	14
PTU	70	Retrovir IV	85	Salbutamol	179
Pulmocare (Vanilla)	211	Reutenox	104	Salbutamol with ipratropium	
Pulmonary Surfactants	182	Revimid	135	bromide	177
Pulmozyme	181	Revolade	29	Salicylic acid	199
Puri-nethol	134	RexAir	181	Salmeterol	181
Pyrazinamide	81	Reyataz	86	Salmonella typhi vaccine	217
Pyridostigmine bromide	95	Riboflavin 5-phosphate	186	Sandimmun	145
Pyridoxal-5-phosphate	21	Rifabutin	81	Sandomigran	119
Pyridoxine hydrochloride	25	Rifadin	81	Sandostatin LAR	143
Pyrimethamine	82	Rifampicin	81	Scalp Preparations	55
Pytazen SR	34	Rifaximin	16	Scandone 3%	109
		Rifinah	80	Sclerosing Agents	182
- Q -		Rilutek	105	Scopoderm TTS	120
Q 300	83	Riluzole	105	Sebizole	51
Quetapel	122	Ringer's solution	37	Secretin pentahydrochloride	194
Quetiapine	122	Riodine	191	Sedatives and Hypnotics	126
Quinapril	39	Risedronate Sandoz	97	Seebri Breezhaler	178
Quinapril with		Risedronate sodium	97	Selegiline hydrochloride	106
hydrochlorothiazide	39	Risperdal Consta	124	Sennosides	20
Quinine dihydrochloride	82	Risperdal Quicklet	122	Sensipar	61
Quinine sulphate	83	Risperidone	122, 124	Serenace	121
Qvar	179	Risperon	122	Seretide	181
		Ritalin	128	Seretide Accuhaler	181
- R -		Ritalin LA	128	Serevent	181
RA-Morph	111	Ritalin SR	128	Serevent Accuhaler	181
Rabies vaccine	219	Ritonavir	86	Serophene	64
Raloxifene	99	Rituximab	164	Sertraline	114
Raltegravir potassium	87	Rivaroxaban	33	Sevoflurane	107
Ramipex	106	Rivastigmine	129	Sevredol	111
Ranbaxy-Cefaclor	72	Rivotril	114	SII-Onco-BCG	174
Ranibizumab	164	RIXUBIS	31	Sildenafil	49
Ranitidine	15	Rizamelt	119	Siltuximab	170
Ranitidine Relief	15	Rizatriptan	119	Silver nitrate	
Rapamune	175	Rocuronium bromide	102	Dermatological	56
Rasburicase	102	Ropinirole hydrochloride	106	Extemporaneous	199
Readi-CAT 2	193	Ropivacaine hydrochloride	109	Simethicone	13
Reandron 1000	61	Ropivacaine hydrochloride with		Simulect	158
Recombinant factor IX	30	fentanyl	109	Simvastatin	46
Recombinant factor VIIa	30	Ropivacaine Kabi	109	Sincalide	194
Recombinant factor VIII	30, 31	Rose bengal sodium	185	Sinemet	106
Rectogesic	15	RotaTeq	219	Sinemet CR	106
Red back spider antivenom	189	Rotavirus live reassortant oral		Singular	180
Redipred	63			Sirolimus	175

Slow-Lopresor	42	Sodium phosphate with phosphoric acid	20	Suprane	106
Snake antivenom	190	Sodium polystyrene		Surgical Preparations	195
Sodibic	38	sulphonate	38	Survanta	182
Sodium acetate	37	Sodium stibogluconate	83	Sustagen Diabetic (Vanilla)	205
Sodium acid phosphate	37	Sodium tetradecyl sulphate	29	Sustagen Hospital Formula (Chocolate)	213
Sodium alginate with magnesium alginate	13	Sodium thiosulfate	189	Sustagen Hospital Formula (Vanilla)	213
Sodium alginate with sodium bicarbonate and calcium carbonate	13	Sodium valproate	117	Sutent	141
Sodium aurothiomalate	95	Sodium with potassium	196	Suxamethonium chloride	102
Sodium benzoate	22	Solian	120	Sylvant	170
Sodium bicarbonate		Solifenacin succinate	60	Symmetrel	105
Blood	37-38	Solu-Cortef	62	Sympathomimetics	48
Extemporaneous	199	Solu-Medrol	62	Synacthen	65
Sodium calcium edetate	191	Somatropin	65	Synacthen Depot	65
Sodium carboxymethylcellulose with pectin and gelatine	23	Sotacor	43	Syntometrine	59
Sodium chloride		Sotalol	43	Syrup	199
Blood	37-38	Spacer device	222	Systane Unit Dose	188
Respiratory	179, 182	Span-K	38		
Various	195	Specialised Formulas	204	- T -	
Sodium chloride with sodium bicarbonate	179	Spiolto Respimat	178	Tacrolimus	145
Sodium citrate		Spiractin	45	Tacrolimus Sandoz	145
Alimentary	13	Spiramycin	83	Tagitol V	193
Extemporaneous	199	Spiriva	178	Talc	182
Sodium citrate with sodium chloride and potassium chloride	33	Spiriva Respimat	178	Tambocor	41
Sodium citrate with sodium lauryl sulphaacetate	20	Spiroinolactone	45	Tambocor CR	41
Sodium citro-tartrate	60	Sprycel	137	Tamoxifen citrate	144
Sodium cromoglycate		Standard Feeds	212	Tamsulosin	60
Alimentary	14	Staphlex	74	Tamsulosin-Rex	60
Respiratory	177, 181	Starch	199	Tarceva	138
Sensory	184	Stavudine	85	Tasigna	139
Sodium dihydrogen phosphate [Sodium acid phosphate]	37	Sterculia with frangula	19	Tasmar	106
Sodium fluoride	22	Stesolid	114	Tecfidera	125
Sodium hyaluronate [Hyaluronic acid]		Stimulants / ADHD		Tegretol	115
Alimentary	24	Treatments	127	Tegretol CR	115
Sensory	186, 188	Stiripentol	117	Teicoplanin	77
Sodium hyaluronate [Hyaluronic acid] with chondroitin sulphate	186	Stocrin	84	Temaccord	136
Sodium hypochlorite	192	Strattera	127	Temazepam	127
Sodium metabisulfite	199	Streptomycin sulphate	71	Temozolomide	136
Sodium nitrite	189	Stromectol	81	Tenecteplase	35
Sodium nitroprusside		Suboxone	129	Tenofovir disoproxil fumarate	89
Cardiovascular	49	Sucralfate	16	Tenoxicam	104
.....	221	Sucrose	110	Terazosin	40
Sodium phenylbutyrate	22	Sugammadex	102	Terbinafine	80
		Sulindac	104	Terbutaline	59
		Sulphacetamide sodium	183	Terbutaline sulphate	179
		Sulphadiazine	77	Teriflunomide	125
		Sulphadiazine silver	51	Teriparatide	100
		Sulphasalazine	14	Terlipressin	70
		Sulphur	199	Testosterone	61
		Sumatriptan	119	Testosterone cypionate	61
		Sunitinib	141	Testosterone esters	61
		Sunscreen, proprietary	56	Testosterone undecanoate	61
				Tetrabenazine	105

Tetracaine [Amethocaine]		Tramal 50	112	Ultiva	112
hydrochloride		Tramal SR 100	112	Ultraproct	14
Nervous	109	Tramal SR 150	112	Umeclidinium	178
Sensory	185	Tramal SR 200	112	Umeclidinium with vilanterol	179
Tetracosactide		Trandolapril	39	Univent	177
[Tetracosactrin]	65	Tranexamic acid	30	Ural	60
Tetracosactrin	65	Tranylcypromine sulphate	113	Urea	
Tetracyclin Wolff	76	Trastuzumab	173	Dermatological	53
Tetracycline	76	Travoprost	187	Extemporaneous	199
Thalidomide	137	Treatments for Dementia	129	Urex Forte	44
Thalomid	137	Treatments for Substance		Urografin	192
Theobroma oil	199	Dependence	129	Urokinase	35
Theophylline	181	Tretinoin		Urologicals	59
Thiamine hydrochloride	25	Dermatological	52	Uromitexan	142
Thioguanine	134	Oncology	137	Ursodeoxycholic acid	18
Thiopental [Thiopentone]		Trexate	134	Ursosan	18
sodium	107	Tri-sodium citrate	199	Utrogestan	59
Thiopentone	107	Triamcinolone acetonide			
Thiotepa	132	Alimentary	23	- V -	
Thrombin	30	Dermatological	54	Vaclovir	91
Thymol glycerin	24	Hormone	63	Valaciclovir	91
Thyroid and Antithyroid		Triamcinolone acetonide with		Valcyte	91
Preparations	69	gramicidin, neomycin and		Valganciclovir	91
Thyrotropin alfa	65	nystatin	184	Vancomycin	77
Ticagrelor	34	Triamcinolone acetonide with		Varenicline	131
Ticarcillin with clavulanic acid	75	neomycin sulphate, gramicidin		Varibar - Honey	193
Ticlopidine	34	and nystatin	55	Varibar - Nectar	193
Tigecycline	76	Triamcinolone hexacetonide	63	Varibar - Pudding	193
Timolol	186	Triazolam	127	Varibar - Thin Liquid	193
Timolol maleate	43	Trichloroacetic acid	199	Varicella vaccine [Chicken pox	
Timoptol XE	186	Trichozole	82	vaccine]	220
Tiotropium bromide	178	Trientine dihydrochloride	22	Varilrix	220
Tiotropium bromide with		Trifluoperazine		Vasodilators	48
olodaterol	178	hydrochloride	122	Vasopressin	70
TMP	77	Trimeprazine tartrate	177	Vasopressin Agents	70
TOBI	71	Trimethoprim	77	Vecuronium bromide	102
Tobradex	184	Trimethoprim with		Vedafil	49
Tobramycin		sulphamethoxazole		Velcade	134
Infection	71	[Co-trimoxazole]	77	Veletri	50
Sensory	183	Trisodium citrate	33	Venlafaxine	114
Tobrex	183	Trometamol	195	Venofer	23
Tocilizumab	171	Tropicamide	187	Ventavis	50
Tofranil	113	Tropisetron	120	Ventolin	179
Tolcapone	106	Tropisetron-AFT	120	Vepesid	135
Tolterodine tartrate	60	Truvada	85	Verapamil hydrochloride	44
Topamax	118	Tuberculin, purified protein		Vergo 16	119
Topicaïne	108	derivative	194	Verpamil SR	44
Topical Products for Joint and		Two Cal HN	207	Vesanoid	137
Muscular Pain	104	TwoCal HN RTH (Vanilla)	207	Vesicare	60
Topiramate	118	Tykerb	139	Vexazone	18
Tracrium	102	Tysabri	125	Vfend	79
Tramadol hydrochloride	112			Victrelis	90
Tramal 100	112	- U -		Vidaza	133
		Ultibro Breezhaler	178		

Vigabatrin	118	Water		Zinacef	72
Vimpat	116	Blood	38	Zinc	
Vinblastine sulphate	142	Various	195	Alimentary	23
Vincristine sulphate	142	Wool fat		Dermatological	52
Vinorelbine	142	Dermatological	53	Zinc and castor oil	52
Viral Vaccines	217	Extemporaneous	199	Zinc chloride	23
Viramune Suspension	84			Zinc oxide	199
Viread	89	- X -		Zinc sulphate	23
Visipaque	192	X-Opaque-HD	193	Zinc with wool fat	53
Vistil	188	Xanthan	199	Zincaps	23
Vistil Forte	188	Xarelto	33	Zinfofo	73
Vit.D3	26	Xifaxan	16	Zinnat	72
VitA-POS	188	Xolair	163	Ziprasidone	123
Vital	206	Xylocaine	108	Zithromax	73
Vitamin A with vitamins D and		Xylocaine Viscous	108	Zoladex	65
C	25	Xylometazoline		Zoledronic acid	
Vitamin B complex	25	hydrochloride	179	Hormone	62
Vitamins	24	Xyntha	30	Musculoskeletal	97-99
Vivonex Paediatric	208			Zometa	62
Vivonex TEN	205	- Y -		Zopiclone	127
Volibris	49	Yellow jacket wasp venom	176	Zostrix	104
Voltaren	103	- Z -		Zostrix HP	109
Voltaren D	103	Zanamivir	92	Zuclopenthixol acetate	123
Voltaren Ophtha	184	Zantac	15	Zuclopenthixol decanoate	124
Volulyte 6%	38	Zapril	39	Zuclopenthixol	
Volumatic	222	Zarator	46	hydrochloride	123
VoLumen	193	Zarzio	35	Zusdone	123
Voluven	38	Zavedos	133	Zyban	130
Voriconazole	79	Zeffix	88	Zypine	122
Votrient	140	Zetop	177	Zypine ODT	122
Vttack	79	Ziagen	85	Zyprexa Relprevv	123
- W -		Zidovudine [AZT]	85	Zytiga	143
Warfarin sodium	33	Zidovudine [AZT] with		Zyvox	77
Wart Preparations	56	lamivudine	85		
		Zimybe	46		