June	2018
Volume 25	Number [·]

Editors:

Kave Wilson & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am - 5pm weekdays)

Circulation

Published each April, August and December. Changes to the contents are published in monthly updates.

Accessible in an electronic format at no cost from the Health Professionals section of the PHARMAC website www.pharmac.govt.nz

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month. Alternatively there is a nominal charge for an annual subscription to the printed Schedule publications. To access either of these subscriptions visit our subscription website www.schedule.co.nz.


Production

Typeset automatically from XML and T_FX. XML version of the Schedule available from www.pharmac.govt.nz/pub/schedule/archive/

Programmers

Anrik Drenth & John Geering

email: texschedule@pharmac.govt.nz



ISSN 1179-3686 pdf ISSN 1172-9376 print

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to PHARMAC and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to PHARMAC should be in written form and not by reproduction of the PHARMAC logo. While care has been taken in compiling this Schedule, PHARMAC takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

Section A	General Rules	6
Section B	Alimentary Tract & Metabolism	20
	Blood & Blood Forming Organs	49
	Cardiovascular System	58
	Dermatologicals	71
	Genito Urinary System	82
	Hormone Preparations – Systemic	88
	Infections – Agents For Systemic Use	99
	Musculoskeletal System	121
	Nervous System	130
	Oncology Agents & Immunosuppressants	165
	Respiratory System & Allergies	208
	Sensory Organs	216
	Various	221

Section C	Extemporaneous Compounds (ECPs)	223
Section D	Special Foods	230
Section E	Practitioner's Supply Orders Rural Areas	250 254
Section F	Dispensing Period Exemptions	255
Section G	Safety Cap Medicines	257
Section I	National Immunisation Schedule	260

Index 271

Introducing PHARMAC

The Pharmaceutical Management Agency (PHARMAC) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. PHARMAC negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

PHARMAC's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about PHARMAC and the way we make funding decisions can be found on the PHARMAC website at http://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by PHARMAC.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements PHARMAC has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A through to G and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

•	g	microgram mcg	millimole mmol
	kg	milligram mg	unit u
international unit	iu	millilitreml	
Abbreviations			
Ampoule	Amp	GelatinousGel	SolutionSoln
Capsule	Сар	Granules Gran	SuppositorySupp
Cream	Crm	Infusion Inf	TabletTab
Device	Dev	Injection Inj	Tincture Tinc
Dispersible	Disp	LiquidLiq	Trans Dermal Delivery
	Ēff	Long ActingLA	SystemTDDS
Emulsion	Emul	OintmentOint	
Enteric Coated	EC	Sachet Sach	
BSO	Dulle Quante Order		
CBS	Bulk Supply Order. Cost Brand Source.		
ECP		Compounded Dreneration	
OP		Compounded Preparation.	
PSO	U U	idy is rounded up to a multiple at whole pack	.5.
Sole Subsidised	Practitioner's Supply	Oldel.	
	Only brand of this m	adiaina aubaidiaad	
<u>Supplier</u> XPharm	Only brand of this m	claim subsidy because PHARMAC has made	altornative distribution arrangements
		may be dispensed at one time if the exempter	
	by the practitioner or	, , , ,	u medicine is endorsed certilied exemption
*		nsed all-at-once or, in the case of oral contract	contives, six menths dispensed all at ence
*		meets the Dispensing Frequency Rule criteri	
+			
‡ ✓		for oral liquid formulations, including extemport nd of a given medicine. Brands without the tion	
•		5	ik are not rully subsidised and may cost the
S29	patient a manufactur		on 20 of the Medicines Act 1081
HP3		unapproved medication supplied under Section	
		pensed from a pharmacy that has a contract	
HP4		spensed from a pharmacy that has a contract	ci to dispense from the Monitored Therapy
	Variation (for Clozap	ine Services).	

Community Pharmaceutical costs met by the Government

Most of the cost of a subsidised prescription for a Community Pharmaceutical is met by the Government through the Combined Pharmaceutical Budget. The Government pays a subsidy for the Community Pharmaceutical to pharmacies, and a fee covering distribution and pharmacy dispensing services. The subsidy paid to pharmacies does not necessarily represent the final cost to Government of subsidising a particular Community Pharmaceutical. The final cost will depend on the nature of PHARMAC's contractual arrangements with the supplier. Fully subsidised medicines are identified with a \checkmark in the product's Schedule listing.

Patient costs

Everyone who is eligible for publicly funded health and disability services should in most circumstances pay only a \$5 co-payment for subsidised medicines, although co-payments can vary from \$0 to \$15. Where the price of a Pharmaceutical is higher than the subsidy, a patient may pay a manufacturer's surcharge in addition to the co-payment. A patient may also pay additional fees for services such as after-hours dispensing and special packaging.

Patients can check whether they are eligible for publicly funded health and disability services by referring to the Guide to eligibility on the Ministry of Health's website.

DHBs have a list of eligible providers in their respective regions. Any provider/prescriber not specifically listed by a DHB as an approved provider/prescriber should be regarded as not approved.

For more information on patient co-payments or eligibility please visit http://www.moh.govt.nz.

Special Authority Applications

Special Authority is an application process in which a prescriber requests government subsidy on a Community Pharmaceutical for a particular person.

Subsidy

Once approved, the applicant will be provided a Special Authority number which must appear on the prescription.

The authority number can provide access to subsidy, increased subsidy, or waive certain restrictions otherwise present on the Community Pharmaceutical.

Some approvals are dependent on the availability of funding from the Combined Pharmaceutical Budget.

Criteria

The criteria for approval of Special Authority applications are included below each Community Pharmaceutical listing, and on the application forms available on PHARMAC's website. For some Special Authority Community Pharmaceuticals, not all indications that have been approved by Medsafe are subsidised.

Making a Special Authority application

Application forms can be found at http://www.pharmac.govt.nz. Except where stated on the application form, applications are processed by the Ministry of Health, and are sent to:

Ministry of Health Sector Services, Private Bag 3015, WANGANUI 4540 Fax: (06) 349 1983 or free fax 0800 100 131

To register for submission of applications on-line - Contact the Ministry of Health on 0800 505 125 or email at onlinehelpdesk@moh.govt.nz. For Special Authority approval numbers, applicants can phone the Ministry of Health Sector Services Call Centre, free phone 0800 243 666.

Named Patient Pharmaceutical Assessment policy

Named Patient Pharmaceutical Assessment (NPPA) provides a mechanism for individual patients to receive funding for medicines not listed in the Pharmaceutical Schedule (either at all or for their clinical circumstances). PHARMAC will assess applications that meet the prerequisites according to its Factors for Consideration before deciding whether to approve applications for funding. The Factors for Consideration will be used to assess both the individual clinical circumstances of each NPPA applicant, and the implications of each NPPA funding decision on PHARMAC's ability to carry out its legislative functions.

For more information on NPPA, or to apply, visit the PHARMAC website at http://www.pharmac.govt.nz/nppa, or call the Panel Coordinators at 0800 660 050 Option 2.

INTRODUCTION

Section A contains the restrictions and other general rules that apply to Subsidies on Community Pharmaceuticals. The amounts payable by the Funder to Contractors are currently determined by:

- the quantities, forms, and strengths, of subsidised Community Pharmaceuticals dispensed under valid prescription by each Contractor;
- the amount of the Subsidy on the Manufacturer's Price payable for each unit of the Community Pharmaceuticals dispensed by each Contractor and;
- the contractual arrangements between the Contractor and the Funder for the payment of the Contractor's dispensing services.

The Pharmaceutical Schedule shows the level of subsidy payable in respect of each Community Pharmaceutical so that the amount payable by the Government to Contractors, for each Community Pharmaceutical, can be calculated. The Pharmaceutical Schedule also shows the standard price (exclusive of GST) at which a Community Pharmaceutical is supplied ex-manufacturer to whole-salers if it differs from the subsidy. The manufacturer's surcharge to patients can be estimated using the subsidy and the standard manufacturer's price as set out in this Schedule.

The cost to Government of subsidising each Community Pharmaceutical and the manufacturer's prices may vary, in that suppliers may provide rebates to other stakeholders in the primary health care sector, including dispensers, wholesalers, and the Government. Rebates are not specified in the Pharmaceutical Schedule.

This Schedule is dated 1 June 2018 and is to be referred to as the Pharmaceutical Schedule Volume 25 Number 1, 2018. Distribution will be from 20 June 2018. This Schedule comes into force on 1 June 2018.

PART I INTERPRETATIONS AND DEFINITIONS

1.1 In this Schedule, unless the context otherwise requires:

"90 Day Lot", means the quantity of a Community Pharmaceutical required for the number of days' treatment covered by the Prescription, being up to 90 consecutive days' treatment;

"180 Day Lot", means the quantity of a Community Pharmaceutical required for the number of days' treatment covered by the Prescription, being up to 180 consecutive days' treatment;

"Access Exemption Criteria", means the criteria under which patients may receive greater than one Month's supply of a Community Pharmaceutical covered by Section F Part II (b) subsidised in one Lot. The specifics of these criteria are conveyed in the Ministry of Health guidelines, which are issued from time to time. The criteria the patient must meet are that they:

- a) have limited physical mobility;
- b) live and work more than 30 minutes from the nearest pharmacy by their normal form of transport;
- c) are relocating to another area;
- d) are travelling extensively and will be out of town when the repeat prescriptions are due.

"Act", means the New Zealand Public Health and Disability Act 2000.

"Advisory Committee", means the Pharmaceutical Services Advisory Committee convened by the Ministry of Health under the terms of the Advice Notice issued to Contractors pursuant to Section 88 of the Act.

"Alternate Subsidy", means a higher level of subsidy that the Government will pay contractors for a particular community Pharmaceutical dispensed to a person who has either been granted a Special Authority for that pharmaceutical, or where the prescription is endorsed in accordance with the requirements of this Pharmaceutical Schedule.

"Annotation", means written annotation of a prescription by a dispensing pharmacist in the pharmacist's own handwriting following confirmation from the Prescriber if required, and "Annotated" has a corresponding meaning. The Annotation must include the details specified in the Schedule, including the date the prescriber was contacted (if applicable) and be initialled by the dispensing pharmacist.

"Authority to Substitute", means an authority for the dispensing pharmacist to change a prescribed medicine in accordance with regulation 42(4) of the Medicines Regulations 1984. An authority to substitute letter, which may be used by Practitioners, is available on the final page of the Schedule.

"Bulk Supply Order", means a written order, on a form supplied by the Ministry of Health, or approved by the Ministry of Health, made by the licensee or manager of an institution certified to provide hospital care under the Health and Disability Services (Safety) Act 2001 for the supply of such Community Pharmaceuticals as are expected to be required for the treatment of persons who are under the medical or dental supervision of such a Private Hospital or institution.

"Class B Controlled Drug", means a Class B controlled drug within the meaning of the Misuse of Drugs Act 1975.

"Community Pharmaceutical", means a Pharmaceutical listed in Sections A to G and Section I of the Pharmaceutical Schedule that is subsidised by the Funder from the Pharmaceutical Budget for use in the community.

"Contractor", means a person who is entitled to receive a payment from the Crown or a DHB under a notice issued by the Crown or a DHB under Section 88 of the Act or under a contract with the Ministry of Health or a DHB for the supply of Community Pharmaceuticals.

"Controlled Drug", means a controlled drug within the meaning of the Misuse of Drugs Act 1975 (other than a controlled drug specified in Part VI of the Third Schedule to that Act).

"Cost, Brand, Source of Supply", means that the Community Pharmaceutical is eligible for Subsidy on the basis of the Contractor's annotated purchase price, brand, and source of supply. Alternatively a copy of the invoice for the purchase of the Pharmaceutical may be attached to the prescription, in the place of an annotation, in order to be eligible for Subsidy.

"Dentist", means a person registered with the Dental Council, and who holds a current annual practising certificate, under the HPCA Act 2003.

"Dietitian", means a person registered as a dietitian with the Dietitians Board, and who holds a current annual practicing certificate under the HPCA Act 2003.

"DHB", means an organisation established as a District Health Board by or under Section 19 of the Act.

"DHB Hospital", means a DHB, including its hospital or associated provider unit that the DHB purchases Hospital Pharmaceuticals for.

"Dispensing Frequency Rule", means the rule in Part IV, Section A of the Pharmaceutical Schedule that defines patient groups or medicines eligible for more frequent dispensing periods.

"Doctor", means a medical Practitioner registered with the Medical Council of New Zealand and, who holds a current annual practising certificate under the HPCA Act 2003.

"DV Limit", means, for a particular Hospital Pharmaceutical with HSS, the National DV Limit or the Individual DV Limit. "DV Pharmaceutical", means a discretionary variance Pharmaceutical, that does not have HSS and which:

- a) is either listed in Section H Part II of the Schedule as being a DV Pharmaceutical in association with the relevant Hospital Pharmaceutical with HSS; or
- b) is the same chemical entity, at the same strength, and in the same or a similar presentation or form, as the relevant Hospital Pharmaceutical with HSS, but which is not yet listed as being a DV Pharmaceutical.

"Endorsements", unless otherwise specified, endorsements should be either handwritten or computer generated by the Prescriber. The endorsement can be written as "certified condition", or state the condition of the patient, where that condition is specified for the Community Pharmaceutical in Section B of the Pharmaceutical Schedule. Where the Prescriber writes "certified condition" as the endorsement, he/she is making a declaration that the patient meets the criteria as set out in Section B of the Pharmaceutical Schedule.

"Funder", means the body or bodies responsible, pursuant to the Act, for the funding of pharmaceuticals listed on the Schedule (which may be one or more DHBs and/or the Ministry of Health) and their successors.

"GST", means goods and services tax under the Goods and Services Tax Act 1985.

"Hospital Care Operator", means a person for the time being in charge of providing hospital care, in accordance with the Health and Disability Services (Safety) Act 2001.

"Hospital Pharmaceuticals", means the list of pharmaceuticals set out in Section H Part II of the Schedule which includes some National Contract Pharmaceuticals.

"Hospital Pharmacy", means that the Community Pharmaceutical is not eligible for Subsidy unless it is supplied by a hospital or pharmacy contracted to the Funder to dispense as a hospital pharmacy to an person on the Prescription of a Prescriber.

"Hospital Pharmacy-Specialist", means that the Community Pharmaceutical is not eligible for Subsidy unless it is supplied by a hospital or pharmacy contracted to the Funder to dispense as a hospital pharmacy to an Outpatient either:

- a) on a Prescription signed by a Specialist, or
- b) where the treatment with the Community Pharmaceutical has been recommended by a Specialist, on the Prescription of a Prescriber which is either:
 - i) endorsed with the words "recommended by [name of specialist and year of authorisation]" and signed by the Prescriber, or
 - ii) endorsed with the word 'protocol' which means "initiated in accordance with DHB hospital approved protocol",
 - iii) annotated by the dispensing pharmacist, following verbal confirmation from the Prescriber of the name of the Specialist and date of recommendation, with the words "recommended by [name of specialist and date of authorisation], confirmed by [Prescriber]". Where the Contractor has an electronic record of such an

Endorsement or Annotation from a previous prescription for the same Community Pharmaceutical written by a prescriber for the same patient, they may annotate the prescription accordingly.

"As recommended by a Specialist" to be interpreted as either:

- i) follows a substantive consultation with an appropriate Specialist;
- ii) the consultation to relate to the Patient for whom the Prescription is written;
- iii) consultation to mean communication by referral, telephone, letter, facsimile or email;
- iv) except in emergencies consultation to precede annotation of the Prescription; and
- v) both the specialist and the General Practitioner must keep a written record of the consultation; or
- a) treatment with the Community Pharmaceutical has been initiated in accordance with a DHB hospital approved protocol.

For the purposes of the definition it makes no difference whether or not the Specialist is employed by a hospital.

"Hospital Pharmacy-Specialist Prescription", means that the Community Pharmaceutical is not eligible for Subsidy unless it is supplied by a hospital or pharmacy contracted to the Funder to dispense as a hospital pharmacy:

- a) to an Outpatient; and
- b) on a Prescription signed by a Specialist.

For the purposes of this definition, a "specialist" means a doctor who holds a current annual practicing certificate and who satisfies the criteria set out in paragraphs (a) or (b) or (c) of the definitions of Specialist below.

"HSS", means hospital supply status, the status of being the brand of the relevant Hospital Pharmaceutical listed in Section H Part II as HSS, that DHBs are obliged to purchase subject to any DV Limit for that Hospital Pharmaceutical for the period of hospital supply, as awarded under an agreement between PHARMAC and the relevant pharmaceutical supplier.

"In Combination", means that the Community Pharmaceutical is only subsidised when prescribed in combination with another subsidised pharmaceutical as specified in Section B or C of the Pharmaceutical Schedule.

"Individual DV Limit", means, for a particular Hospital Pharmaceutical with HSS and a particular DHB Hospital, the discretionary variance limit, being the specified percentage of that DHB Hospital's Total Market Volume up to which that DHB Hospital may purchase DV Pharmaceuticals of that Hospital Pharmaceutical.

"Licensed Hospital", means a place or institution that is certified to provide hospital care within the meaning of the Health and Disability Services (Safety) Act 2001.

"Lot", means a quantity of a Community Pharmaceutical supplied in one dispensing.

"Manufacturer's Price", means the standard price at which a Community Pharmaceutical is supplied to wholesalers (excluding GST), as notified to PHARMAC by the supplier.

"Maternity hospital", means that the Community Pharmaceutical is not eligible for Subsidy unless it is supplied pursuant to a Bulk Supply Order to a maternity hospital certified under the Health and Disability Services (Safety) Act 2001.

"Midwife", means a person registered as a midwife with the Midwifery Council, and who holds a current annual practising certificate under the HPCA Act 2003.

"Month", means a period of 30 consecutive days.

"Monthly Lot", means the quantity of a Community Pharmaceutical required for the number of days' treatment covered by the Prescription, being up to 30 consecutive days' treatment;

"Named Patient Pharmaceutical Assessment Advisory Panel", means the panel of clinicians, appointed by the PHARMAC Board, that is responsible for advising, within its Terms of Reference, on Named Patient Pharmaceutical Assessment applications and Exceptional Circumstances renewal applications submitted after 1 March 2012 (EC renewal application form located at http://www.pharmac.govt.nz/nppa#oldec)

"National Contract Pharmaceutical", means a Hospital Pharmaceutical for which PHARMAC has negotiated a national contract and the Price.

"National DV Limit", means, for a particular Hospital Pharmaceutical with HSS, the discretionary variance limit, being the specified percentage of the Total Market Volume up to which all DHB Hospitals may collectively purchase DV Pharmaceuticals of that Hospital Pharmaceutical.

"National Immunisation Schedule", means Section I of the Pharmaceutical Schedule, which is a schedule administered by PHARMAC, being a schedule specifying a programme of vaccinations to promote immunity against the diseases specified in the schedule.

"Not In Combination", means that no Subsidy is available for any Prescription containing the Community Pharmaceutical in combination with other ingredients unless the particular combination of ingredients is separately specified in Section B or C of the Schedule, and then only to the extent specified. "Nurse Practitioner", means a nurse registered with Nursing Council of New Zealand, who holds a current annual practising certificate under the HPCA Act 2003 and for whom the Nursing Council has authorised a scope of practice that includes prescribing medicines

"Optional Pharmaceuticals", means the list of National Contract Pharmaceuticals set out in Section H Part II of the Schedule

"Optometrist", means a person registered with the Optometrists and Dispensing Opticians Board with a scope of practice that includes prescribing medicines (TPA endorsement)

"Outpatient", in relation to a Community Pharmaceutical, means a person who, as part of treatment at a hospital or other institution under the control of a DHB, is prescribed the Community Pharmaceutical for consumption or use in the person's home.

"PCT", means Pharmaceutical Cancer Treatment in respect of which DHB hospital pharmacies and other Contractors can claim Subsidies.

"PCT only", means Pharmaceutical Cancer Treatment in respect of which only DHB hospital pharmacies can claim Subsidies.

"Penal Institution", means a penal institution, as that term is defined in The Penal Institutions Act 1954;

"PHARMAC", means the Pharmaceutical Management Agency established by Section 46 of the Act (PHARMAC).

"Pharmaceutical", means a medicine, therapeutic medical device, or related product or related thing listed in Sections B to I of the Schedule.

"Pharmaceutical Benefits", means the right of:

- a) a person; and
- b) any member under 16 years of age of that person's family, to have made by the Government on his or her behalf, subject to any conditions for the time being specified in the Schedule, such payment in respect of any Community Pharmaceutical supplied to that person or family member under the order of a Practitioner in the course of his or her practice.

"Pharmaceutical Budget", means the pharmaceutical budget set for PHARMAC by the Crown for the subsidised supply of Community Pharmaceuticals and Pharmaceutical Cancer Treatments including for named patients in exceptional circumstances.

"Pharmaceutical Cancer Treatment", means Pharmaceuticals for the treatment of cancer, listed in Sections A to G of the Schedule and identified therein as a "PCT" or "PCT only" Pharmaceutical that DHBs must provide access to, for use in their hospitals, and/or in association with Outpatient services provided in their DHB Hospitals, in relation to the treatment of cancers.

"Pharmacist Prescriber", means a person registered with the Pharmacy Council of New Zealand, who holds a current annual practising certificate under the HPCA Act 2003, and is approved by the Pharmacy Council of New Zealand to prescribe specified prescription medicines relating to his/her scope of practice.

"Pharmacist", means a person registered with the Pharmacy Council of New Zealand and who holds a current annual practicing certificate under the HPCA Act 2003.

"Practitioner", means a Prescriber or any of the following: Quitcard Provider, a Pharmacist, or a Vaccinator as those terms are defined in the Pharmaceutical Schedule.

"Practitioner's Supply Order", means a written order made by a Prescriber on a form supplied by the Ministry of Health, or approved by the Ministry of Health, for the supply of Community Pharmaceuticals to the Prescriber, which the Prescriber requires to ensure medical supplies are available for emergency use, teaching and demonstration purposes, and for provision to certain patient groups where individual prescription is not practicable.

"Prescriber", means a Doctor, a Dentist, a Dietitian, a Midwife, a Nurse Practitioner, a Registered Nurse Prescriber, an Optometrist, or a Pharmacist Prescriber as those terms are defined in the Pharmaceutical Schedule.

"Prescription", means a quantity of a Community Pharmaceutical prescribed for a named person on a document signed by a Prescriber.

"Prescription Medicine", means any Pharmaceutical listed in Part I of Schedule 1 of the Medicines Regulations 1984. "Private Hospital", means a hospital certified under the Health and Disability Services (Safety) Act 2001 that is not owned or operated by a DHB.

"Quitcard Provider", means a person registered with the Ministry of Health as a Quitcard Provider.

"Registered Nurse Prescriber", means a registered nurse who meets specified requirements for qualifications, training and competence to be a designated prescriber for the purpose of prescribing specified prescription medicines under the Medicines (Designated Prescriber-Registered Nurses) Regulations 2016.

"Residential Disability Care Institution", means premises used to provide residential disability care in accordance with the Health and Disability Services (Safety) Act 2001.

a)

"Rest Home", means premises used to provide rest home care in accordance with the Health and Disability Services (Safety) Act 2001.

"Restricted Medicine", means any Pharmaceutical listed in Part II of Schedule 1 of the Medicines Regulations 1984.

- "Retail Pharmacy-Specialist", means that the Community Pharmaceutical is only eligible for Subsidy if it is either:
 - a) supplied on a Prescription or Practitioner's Supply Order signed by a Specialist, or,
 - b) in the case of treatment recommended by a Specialist, supplied on a Prescription or Practitioner's Supply Order and either:
 - i) endorsed with the words "recommended by [name of Specialist and year of authorisation]" and signed by the Prescriber, or
 - iii) endorsed with the word 'protocol' which means "initiated in accordance with DHB hospital approved protocol", or
 - iii) Annotated by the dispensing pharmacist, following verbal confirmation from the Prescriber of the name of the Specialist and date of recommendation, with the words "recommended by [name of specialist and year of authorisation], confirmed by [Prescriber]". Where the Contractor has an electronic record of such an Endorsement or Annotation from a previous prescription for the same Community Pharmaceutical written by a prescriber for the same patient, they may annotate the prescription accordingly.

"As recommended by a Specialist" to be interpreted as either:

- i) follows a substantive consultation with an appropriate Specialist;
 - ii) the consultation to relate to the Patient for whom the Prescription is written;
 - iii) consultation to mean communication by referral, telephone, letter, facsimile or email;
 - iv) except in emergencies consultation to precede annotation of the Prescription; and
 - v) both the Specialist and the General Practitioner must keep a written record of consultation; or
- b) treatment with the Community Pharmaceutical has been initiated in accordance with a DHB hospital approved protocol.

"Retail Pharmacy-Specialist Prescription", means that the Community Pharmaceutical is only eligible for Subsidy if it is supplied on a Prescription, or Practitioner's Supply Order, signed by a Specialist.

For the purposes of this definition, a "specialist" means a doctor who holds a current annual practicing certificate and who satisfies the criteria set out in paragraphs (a) or (b) or (c) of the definitions of Specialist below.

"Safety Medicine", means a Community Pharmaceutical defined in Section A, Part IV of the Pharmaceutical Schedule. "Schedule", means this Pharmaceutical Schedule and all its sections and appendices.

"Special Authority", means that the Community Pharmaceutical or Pharmaceutical Cancer Treatment is only eligible for Subsidy or additional Subsidy for a particular person if an application meeting the criteria specified in the Schedule has been approved, and the valid Special Authority number is present on the prescription.

"Specialist", in relation to a Prescription, means a doctor or nurse practitioner who holds a current annual practising certificate and who satisfies the criteria set out in paragraphs (a) or (b) or (c) or (d) below:

- a) the doctor is vocationally registered in accordance with the criteria set out by the Medical Council of New Zealand and the HPCA Act 2003 and who has written the Prescription in the course of practising in that area of medicine; or
- b) the doctor is recognised by the Ministry of Health as a specialist for the purposes of this Schedule and receives remuneration from a DHB at a level which that DHB considers appropriate for specialists and who has written that prescription in the course of practising in that area of competency; or
- c) the doctor is recognised by the Ministry of Health as a specialist in relation to a particular area of medicine for the purpose of writing Prescriptions and who has written the Prescription in the course of practising in that area of competency; or
- d) the doctor or nurse practitioner writes the prescription on DHB stationery and is appropriately authorised by the relevant DHB to do so.

"Subsidy", means the maximum amount that the Government will pay Contractors for a Community Pharmaceutical dispensed to a person eligible for Pharmaceutical Benefits and is different from the cost to Government of subsidising that Community Pharmaceutical. For the purposes of a DHB hospital pharmacy claiming for Pharmaceutical Cancer Treatments, Subsidy refers to any payment made to the DHB hospital pharmacy or service provider to which that pharmacy serves, and does not relate to a specific payment that might be made on submission of a claim. "Supply Order", means a Bulk Supply Order or a Practitioner's Supply Order.

"Unapproved Indication", means, for a Pharmaceutical, an indication for which it is not approved under the Medicines Act 1981. Prescribers of pharmaceuticals for Unapproved Indications should be aware of, and comply with, their obligations under Section 25 and/or Section 29 of the Medicines Act 1981 and as set out in Section A: General Rules, Part IV (Miscellaneous Provisions) rule 5.5.

"Unlisted Pharmaceutical", means a Pharmaceutical that is within the scope of a Hospital Pharmaceutical but is not listed in Section H Part II

"Unusual Clinical Circumstances (UCC)", means the pathway under the Named Patient Pharmaceutical Assessment policy for funding consideration for named patients whose clinical circumstances are so unusual that PHARMAC is unlikely, for administrative reasons, to consider listing treatments for these circumstances on the Schedule.

"Urgent Assessment (UA)", means the pathway under the Named Patient Pharmaceutical Assessment policy for funding consideration for treatments for named patients where PHARMAC is also considering or is likely to consider the treatment for Schedule listing, but the patient's clinical circumstances justify urgent assessment, prior to a decision on Schedule listing.

"Vaccinator", means either:

- a) a pharmacist who has successfully completed a vaccinator training course approved by the Ministry of Health and who is complying with the immunisation standards of the Ministry of Health; or
- b) any other person who is authorised by the Director-General of Health or a Medical Officer of Health to administer vaccines in accordance with this Section 44A of the Medicines Regulations 1984.
- 1.2 In addition to the above interpretations and definitions, unless the content requires otherwise, a reference in the Schedule to:
 - a) the singular includes the plural; and
 - b) any legislation includes a modification and re-enactment of, legislation enacted in substitution for, and a regulation, Order in Council, and other instrument from time to time issued or made under that legislation, where that legislation, regulation, Order in Council or other instrument has an effect on the prescribing, dispensing or subsidising of Community Pharmaceuticals.

PART II COMMUNITY PHARMACEUTICALS SUBSIDY

- 2.1 Community Pharmaceuticals eligible for Subsidy include every medicine, therapeutic medical device or related product, or related thing listed in Sections B to G and I of the Schedule subject to:
 - 2.1.1 clauses 2.2 of the Schedule; and
 - 2.1.2 clauses 3.1 to 5.4 of the Schedule; and
 - 2.1.3 the conditions (if any) specified in Sections B to G and I of the Schedule;
- 2.2 No claim by a Contractor for payment in respect of the supply of Community Pharmaceuticals will be allowed unless the Community Pharmaceuticals so supplied:
 - 2.2.1 comply with the appropriate standards prescribed by regulations for the time being in force under the Medicines Act 1981; or
 - 2.2.2 in the absence of any such standards, comply with the appropriate standards for the time being prescribed by the British Pharmacopoeia; or
 - 2.2.3 in the absence of the standards prescribed in clauses 2.2.1 and 2.2.2, comply with the appropriate standards for the time being prescribed by the British Pharmaceutical Codex; or
 - 2.2.4 in the absence of the standards prescribed in clauses 2.2.1, 2.2.2 and 2.2.3 are of a grade and quality not lower than those usually applicable to Community Pharmaceuticals intended to be used for medical purposes.

PART III PERIOD AND QUANTITY OF SUPPLY

3.1 Prescribers Prescriptions and provision of pharmaceuticals by other Practitioners (other than oral contraceptives)

The following provisions apply to all Prescriptions, other than those for an oral contraceptive, written by a Prescriber and provision of pharmaceuticals by other Practitioners unless specifically excluded:

- 3.1.1 For a Community Pharmaceutical other than a Class B Controlled Drug, only a quantity sufficient to provide treatment for a period not exceeding three Months will be subsidised.
- 3.1.2 For methylphenidate hydrochloride and dexamfetamine sulphate (except for Dentist prescriptions), only a quantity sufficient to provide treatment for a period not exceeding one Month will be subsidised.
- 3.1.3 For a Class B Controlled Drug:
 - a) other than Dentist prescriptions and methylphenidate hydrochloride and dexamfetamine sulphate, only a quantity:
 - i) sufficient to provide treatment for a period not exceeding 10 days; and
 - ii) which has been dispensed pursuant to a Prescription sufficient to provide treatment for a period not exceeding one Month, will be subsidised.
 - b) for a Dentist prescription only such quantity as is necessary to provide treatment for a period not exceeding five days will be subsidised.
- 3.1.4 Subject to clauses 3.1.3 and 3.1.7, for a Prescriber and 3.1.7 for an Optometrist, where a Prescriber has prescribed a quantity of a Community Pharmaceutical sufficient to provide treatment for:
 - A) one Month or less than one Month, but dispensed by the Contractor in quantities smaller than the quantity prescribed, the Community Pharmaceutical will only be subsidised as if that Community Pharmaceutical had been dispensed in a Monthly Lot;
 - B) more than one Month, the Community Pharmaceutical will be subsidised only if it is dispensed:
 - i) in a 90 Day Lot, where the Community Pharmaceutical is a Pharmaceutical covered by Section F Part I of the Pharmaceutical Schedule; or
 - ii) if the Community Pharmaceutical is not a Pharmaceutical referred to in Section F Part I of the Pharmaceutical Schedule, in Monthly Lots, unless:
 - a) the eligible person or his/her nominated representative endorses the back of the Prescription form with a statement identifying which Access Exemption Criterion (Criteria) applies and signs that statement to this effect; or
 - b) both:
 - the Prescriber endorses the Community Pharmaceutical on the Prescription with the words "certified exemption" written in the Prescriber's own handwriting, or signed or initialled by the Prescriber; and
 - 2) every Community Pharmaceutical endorsed as "certified exemption" is covered by Section F Part II of the Pharmaceutical Schedule.
- 3.1.5 A Community Pharmaceutical is only eligible for Subsidy if the Prescription under which it has been dispensed was presented to the Contractor:
 - a) for a Class B Controlled Drug, within eight days of the date on which the Prescription was written; or
 - b) for any other Community Pharmaceutical, within three Months of the date on which the Prescription was written.
- 3.1.6 No subsidy will be paid for any Prescription, or part thereof, that is not fulfilled within:
 - a) in the case of a Prescription for a total supply of from one to three Months, three Months from the date the Community Pharmaceutical was first dispensed; or
 - b) in any other case, one Month from the date the Community Pharmaceutical was first dispensed. Only that part of any Prescription that is dispensed within the time frames specified above is eligible for Subsidy.
- 3.1.7 If a Community Pharmaceutical:
 - a) is stable for a limited period only, and the Prescriber has endorsed the Prescription with the words "unstable medicine" and has specified the maximum quantity that may be dispensed at any one time; or
 - b) is stable for a limited period only, and the Contractor has endorsed the Prescription with the words "unstable medicine" and has specified the maximum quantity that should be dispensed at any one time in all the circumstances of the particular case; or
 - c) is under the Dispensing Frequency Rule,
 - The actual quantity dispensed will be subsidised in accordance with any such specification.

3.2 Oral Contraceptives

The following provisions apply to all Prescriptions written by a Prescriber for an oral contraceptive:

3.2.1 The Prescriber must specify on the Prescription the period of treatment for which the Community Pharmaceutical is to be supplied. This period must not exceed six Months.

- 3.2.2 Where the period of treatment specified in the Prescription does not exceed six Months, the Community Pharmaceutical is to be dispensed:
 - a) in Lots as specified in the Prescription if the Community Pharmaceutical is under the Dispensing Frequency Rule; or
 - b) where no Lots are specified, in one Lot sufficient to provide treatment for the period prescribed.
- 3.2.3 An oral contraceptive is only eligible for Subsidy if the Prescription under which it has been dispensed was presented to the Contractor within three Months of the date on which it was written.
- 3.2.4 Where a Community Pharmaceutical on a Prescription is under the Dispensing Frequency Rule and a repeat on the Prescription remains unfulfilled after six Months from the date the Community Pharmaceutical was first dispensed only the actual quantity supplied by the Contractor within this time limit will be eligible for Subsidy.

3.3 Original Packs, Certain Antibiotics and Unapproved Medicines

- 3.3.1 Notwithstanding clauses 3.1 and 3.3 of the Schedule, if a Prescriber prescribes or orders a Community Pharmaceutical that is identified as an Original Pack (OP) on the Pharmaceutical Schedule and is packed in a container from which it is not practicable to dispense lesser amounts, every reference in those clauses to an amount or quantity eligible for Subsidy, is deemed to be a reference:
 - a) where an amount by weight or volume of the Community Pharmaceutical is specified in the Prescription, to the smallest container of the Community Pharmaceutical, or the smallest number of containers of the Community Pharmaceutical, sufficient to provide that amount; and
 - b) in every other case, to the amount contained in the smallest container of the Community Pharmaceutical that is manufactured in, or imported into, New Zealand.
- 3.3.2 If a Community Pharmaceutical is either:
 - a) the liquid oral form of an antibiotic to which a diluent must be added by the Contractor at the time of dispensing; or
 - b) an unapproved medicine supplied under Section 29 of the Medicines Act 1981, but excluding any medicine listed as Cost, Brand, Source of Supply, or
 - c) any other pharmaceutical that PHARMAC determines, from time to time and notes in the Pharmaceutical Schedule

and it is prescribed or ordered by a Prescriber in an amount that does not coincide with the amount contained in one or more standard packs of that Community Pharmaceutical, Subsidy will be paid for the amount prescribed or ordered by the Prescriber in accordance with either clause 3.1 or clause 3.3 of the Schedule, and for the balance of any pack or packs from which the Community Pharmaceutical has been dispensed. At the time of dispensing the Contractor must keep a record of the quantity discarded. To ensure wastage is reduced, the Contractor should reduce the amount dispensed to make it equal to the quantity contained in a whole pack where:

- a) the difference between the amount dispensed and the amount prescribed or ordered by the Prescriber is less than 10% (eg; if a prescription is for 105 mls then a 100 ml pack would be dispensed); and
- b) in the reasonable opinion of the Contractor the difference would not affect the efficacy of the course of treatment prescribed or ordered by the Prescriber.

Note: For the purposes of audit and compliance it is an act of fraud to claim wastage and then use the wastage amount for any subsequent prescription.

3.4 Pharmacist Prescribers' Prescriptions

The following apply to every prescription written by a Pharmacist Prescriber

- 3.4.1 Prescriptions written by a Pharmacist Prescriber for a Community Pharmaceutical will only be subsidised where they are for either:
 - a) a Community Pharmaceutical classified as a Prescription Medicine and which a Pharmacist Prescriber is permitted under regulations to prescribe; or
 - b) any other Community Pharmaceutical that is a Restricted Medicine (Pharmacist Only Medicine), a Pharmacy Only Medicine or a General Sales Medicine.
- 3.4.2 Any Pharmacist Prescribers' prescriptions for a medication requiring a Special Authority will only be subsidised if it is for a repeat prescription (ie after the initial prescription with Special Authority approval was dispensed).

3.5 Registered Nurse Prescribers' Prescriptions

The following apply to every prescription written by a Registered Nurse Prescriber:

3.5.1 Prescriptions written by a Registered Nurse Prescriber for a Community Pharmaceutical will only be subsidised where they are for either:

- a Community Pharmaceutical classified as a Prescription Medicine and which a Registered Nurse Prescriber is permitted under regulations to prescribe; or
- b) any other Community Pharmaceutical that is a Restricted Medicine (Pharmacist Only Medicine), a Pharmacy Only Medicine or a General Sale Medicine.
- 3.5.2 Any Registered Nurse Prescribers' prescriptions for a medication requiring a Special Authority will only be subsidised if it is for a repeat prescription (ie after the initial prescription with Special Authority approval was dispensed). Registered Nurse Prescribers are not eligible to apply for Special Authority approvals (initial or renewal).

3.6 Non-prescribing Practitioners

- 3.6.1 Dispensing on the authority of a Quitcard will only be subsidised where it is:
 - a) for any of the following Community Pharmaceuticals: nicotine patches, nicotine lozenges or nicotine gum; and
 - b) written on a Quitcard.
- 3.6.2 Provision of vaccines by Vaccinators

Vaccines will only be valid for subsidy in accordance with an agreement between the Contractor and the DHB, and only for direct administration of a vaccine to a patient.

3.6.3 Provision of a Community Pharmaceutical by a Pharmacist Except where pursuant to a prescription, Quitcard or supply order, provision of a community pharmaceutical by a pharmacist will only be subsidised where specifically indicated in Section B of the Pharmaceutical Schedule.

PART IV DISPENSING FREQUENCY RULE

Rule 3.1.4 of the Pharmaceutical Schedule specifies, for community patients, a default period of supply for each Community Pharmaceutical (a Monthly Lot, 90 Day Lot or for oral contraceptives 180 Day Lot). This Dispensing Frequency Rule defines patient groups or medicines eligible for more frequent dispensing periods for Community Pharmaceuticals; and the conditions that must be met to enable any pharmacy to claim for payment of handling fees for the additional dispensings made. This Dispensing Frequency Rule relates to the circumstances in which a subsidy is payable for the Community Pharmaceutical; it does not override alternative dispensing frequencies as expressly stated in the Medicines Act, Medicines Regulations, Pharmacy Services Agreement or Pharmaceutical Schedule.

For the purposes of this Dispensing Frequency Rule:

"Frequent Dispensing" means:

- i) for a Community Pharmaceutical referred to in Section F Part I, (the Stat exemption) dispensing in quantities less than one 90 Day Lot (or for oral contraceptives, less than one 180 Day Lot); or
- ii) for any other Community Pharmaceutical dispensing in quantities less than a Monthly Lot

"Safety Medicine"

- i) an antidepressant listed under the "Cyclic and Related Agents" subheading;
- ii) an antipsychotic;
- iii) a benzodiazepine;
- iv) a Class B Controlled Drug;
- v) codeine (includes combination products);
- vi) buprenorphine with naloxone; or
- vii) zopiclone.

The Dispensing Frequency Rule covers 5 different circumstances where Frequent Dispensing for patients may be clinically or otherwise appropriate. These are:

- 1) Long Term Condition (LTC) patients and Core patients, or
- 2) Persons in residential care, or
- 3) Trial periods, or
- 4) Safety and co-prescribed medicines, or
- 5) Pharmaceutical Supply Management.
- 4.1 Frequent Dispensing for patients registered as Long Term Condition (LTC) or Core patients
 - If a Pharmacist considers Frequent Dispensing is required, then:
 - 4.1.1 For LTC registered patients, Frequent Dispensing can occur as often as the dispensing Pharmacist deems appropriate to meet that patient's compliance and adherence needs;

4.1.2 For Core (non-LTC) patients, Frequent Dispensing should be no more often than a Monthly Lot. Pharmacists may authorise monthly dispensing on a Stat exemption Community Pharmaceutical without prescriber authority. If the Pharmacist considers more frequent (than monthly) dispensing is necessary, prescriber approval is required. Verbal approval from the prescriber is acceptable provided it is annotated by the Pharmacist on the Prescription and dated.

4.2 Frequent Dispensings for persons in residential care

- 4.2.1 Community Pharmaceuticals can be dispensed to:
 - any person whose placement in a Residential Disability Care Institution is funded by the Ministry of Health or a DHB; or
 - a person assessed as requiring long term residential care services and residing in an age related residential care facility;

on the request of the person, their agent or caregiver or community residential service provider via Frequent Dispensing, provided the following conditions are met:

- a) the quantity or period of supply to be dispensed at any one time is not less than:
 - i) 7 days' supply for a Class B Controlled Drug; or
 - ii) 7 days' supply for clozapine in accordance with a Clozapine Dispensing Protocol; or
 - 28 days' supply for any other Community Pharmaceutical (except under conditions outlined in 4.3 (Trial periods) below; and
- b) the Prescriber or dispensing Pharmacist has
 - i) included the name of the patient's residential placement or facility on the Prescription; and
 - ii) included the patient's NHI number on the Prescription; and
 - iii) specified the maximum quantity or period of supply to be dispensed at any one time.
- 4.2.2 Any person meeting the criteria above who is being initiated onto a new medicine or having their dose changed is able to have their medicine dispensed in accordance with 4.3 (Trial periods) below.

4.3 Frequent Dispensings for Trial Periods

Frequent Dispensing can occur when a Community Pharmaceutical has been prescribed for a patient who requires close monitoring due to recent initiation onto, or dose change for, the Community Pharmaceutical (applicable to the patient's first changed Prescription only) and the Prescriber has:

- endorsed each Community Pharmaceutical on the Prescription clearly with the words "Trial Period", or "Trial"; and
- specified the maximum quantity or period of supply to be dispensed for each Community Pharmaceutical at any one time.
- Patients who reside in Penal Institutions are not eligible for Trial Periods.

4.4 Frequent Dispensing for Safety and co-prescribed medicines

- 4.4.1 For a Safety Medicine to be dispensed via Frequent Dispensing, both of the following conditions must be met:
 - a) The patient is not a resident in a Penal Institution, or one of the residential placements or facilities referenced in 4.2 on page 15; and
 - b) The prescribing Practitioner has:
 - i) Assessed clinical risk and determined the patient requires increased Frequent Dispensing; and
 - ii) Specified the maximum quantity or period of supply to be dispensed for each Safety Medicine at each dispensing.
- 4.4.2 A Community Pharmaceutical that is co-prescribed with a Safety Medicine, which can be dispensed in accordance with rule 4.4 above, may be dispensed at the same frequency as the Safety Medicine if the dispensing pharmacist has:
 - Assessed clinical risk and determined the patient requires Frequent Dispensing of their co-dispensed medicines; and
 - Annotated the Prescription with the amended dispensing quantity and frequency.

4.5 Frequent Dispensing for Pharmaceutical Supply Management

- 4.5.1 Frequent Dispensing may be required from time to time to manage stock supply issues or emergency situations. Pharmacists may dispense more frequently than the Schedule would otherwise allow when all of the following conditions are met:
 - PHARMAC has approved and notified pharmacists to annotate Prescriptions for a specified Community Pharmaceutical(s) "out of stock" without prescriber endorsement for a specified time; and
 - b) the dispensing pharmacist has:

- i) clearly annotated each of the approved Community Pharmaceuticals that appear on the Prescription with the words "out of stock" or "OOS"; and
- ii) initialled the annotation in their own handwriting; and
- iii) has complied with maximum quantity or period of supply to be dispensed at any one time, as specified by PHARMAC at the time of notification.

Note -- no claim shall be made to any DHB for subsidised dispensing under this rule where dispensing occurs more frequently than specified by PHARMAC to manage the supply management issue.

PART V MISCELLANEOUS PROVISIONS

5.1 Bulk Supply Orders

- The following provisions apply to the supply of Community Pharmaceuticals under Bulk Supply Orders:
- 5.1.1 No Community Pharmaceutical supplied under a Bulk Supply Order will be subsidised unless all the requirements in Section B, C or D of the Schedule applicable to that pharmaceutical are met.
- 5.1.2 The person who placed the Bulk Supply Order may be called upon by the Ministry of Health to justify the amount ordered.
- 5.1.3 Class B Controlled Drugs will be subsidised only if supplied under Bulk Supply Orders placed by an institution certified to provide hospital care under the Health and Disability Services (Safety) Act 2001.
- 5.1.4 Any order for a Class B Controlled Drug or for buprenorphine hydrochloride must be written on a Special Bulk Supply Order Controlled Drug Form supplied by the Ministry of Health.
- 5.1.5 Community Pharmaceuticals listed in Part I of the First Schedule to the Medicines Regulations 1984 will be subsidised only if supplied under a Bulk Supply Order placed by an institution certified to provide hospital care under the Health and Disability Services (Safety) Act 2001 and:
 - a) that institution employs a registered general nurse, registered with the Nursing Council and who holds a current annual practicing certificate under the HPCA Act 2003; and
 - b) the Bulk Supply Order is supported by a written requisition signed by a Hospital Care Operator.
- 5.1.6 No Subsidy will be paid for any quantity of a Community Pharmaceutical supplied under a Bulk Supply Order in excess of what is a reasonable monthly allocation for the particular institution, after taking into account stock on hand.
- 5.1.7 The Ministry of Health may, at any time, by public notification, declare that any approved institution within its particular region, is not entitled to obtain supplies of Community Pharmaceuticals under Bulk Supply Orders with effect from the date specified in that declaration. Any such notice may in like manner be revoked by the Ministry of Health at any time.

5.2 Practitioner's Supply Orders

The following provisions apply to the supply of Community Pharmaceuticals to Practitioners under a Practitioner's Supply Order:

- 5.2.1 Subject to clause 5.2.3 and 5.2.6, a Prescriber may only order under a Practitioner's Supply Order those Community Pharmaceuticals listed in Section E Part I and only in such quantities as set out in Section E Part I that the Practitioner requires to ensure medical supplies are available for emergency use, teaching and demonstration purposes, and for provision to certain patient groups where individual prescription is not practicable.
- 5.2.2 Any order for a Class B Controlled Drug or for buprenorphine hydrochloride must be written on a Special Practitioner's Supply Order Controlled Drug Form supplied by the Ministry of Health.
- 5.2.3 A Prescriber may order such Community Pharmaceuticals as he or she expects to be required for personal administration to patients under the Prescriber's care if:
 - a) the Prescriber's normal practice is in the specified areas listed in Section E Part II of the Schedule, or if the Prescriber is a locum for a Prescriber whose normal practice is in such an area.
 - b) the quantities ordered are reasonable for up to one Month's supply under the conditions normally existing in the practice. (The Prescriber may be called on by the Ministry of Health to justify the amounts of Community Pharmaceuticals ordered.)
- 5.2.4 No Community Pharmaceutical ordered under a Practitioner's Supply order will be eligible for Subsidy unless:
 - a) the Practitioner's Supply Order is made on a form supplied for that purpose by the Ministry of Health, or approved by the Ministry of Health and which:

- i) is personally signed and dated by the Prescriber; and
- ii) sets out the Prescriber's address; and
- iii) sets out the Community Pharmaceuticals and quantities, and;
- b) all the requirements of Sections B and C of the Schedule applicable to that pharmaceutical are met.
- 5.2.5 The Ministry of Health may, at any time, on the recommendation of an Advisory Committee appointed by the Ministry of Health for that purpose, by public notification, declare that a Prescriber specified in such a notice is not entitled to obtain supplies of Community Pharmaceuticals under Practitioner's Supply Orders until such time as the Ministry of Health notifies otherwise.
- 5.2.6 A Prescriber working in the Rheumatic Fever Prevention Programme (RFPP) may order under a Practitioner's Supply Order such Community Pharmaceuticals (identified below) as he or she requires to ensure medical supplies are available for patients with suspected or confirmed Group A Streptococcal throat infections for the purposes of the RFPP in the following circumstances:
 - a) the RFPP provider name is written on the Practitioner's Supply Order; and
 - b) the total quantity ordered does not exceed a multiple of:
 - i) ten times the Practitioner's Supply Order current maximum listed in Section E Part I for amoxicillin grans for oral liq 250 mg per 5 ml, amoxicillin cap 250 mg and amoxicillin cap 500 mg; or
 - two times the Practitioner's Supply Order current maximum listed in Section E Part I for phenoxymethyl penicillin grans for oral liquid 250 mg per 5 ml, phenoxymethyl penicillin cap 500 mg, erythromycin ethyl succinate grans for oral liq 200 mg per 5 ml and erythromycin ethyl succinate tab 400 mg; and
 - c) the Prescriber must specify the order quantity in course-specific amounts on the Practitioner's Supply Order (e.g. 10 x 300 ml amoxicillin grans for oral liq 250 mg per 5 ml). This will enable the pharmacy to dispense each course separately and claim multiple service fees as per the Community Pharmacy Services Agreement.

5.3 Retail Pharmacy and Hospital Pharmacy-Specialist Restriction

The following provisions apply to Prescriptions for Community Pharmaceuticals eligible to be subsidised as "Retail Pharmacy-Specialist" and "Hospital Pharmacy-Specialist":

5.3.1 Record Keeping

It is expected that a record will be kept by both the General Practitioner and the Specialist of the fact of consultation and enough of the clinical details to justify the recommendation. This means referral by telephone will need to be followed up by written consultation.

5.3.2 Expiry

The recommendation expires at the end of two years and can be renewed by a further consultation.

- 5.3.3 The circulation by Specialists of the circumstances under which they are prepared to recommend a particular Community Pharmaceutical is acceptable as a guide. It must however be followed up by the procedure in subclauses 5.3.1 and 5.3.2, for the individual Patient.
- 5.3.4 The use of preprinted forms and named lists of Specialists (as circulated by some pharmaceutical companies) is regarded as inappropriate.
- 5.3.5 The Rules for Retail Pharmacy-Specialist and Hospital Pharmacy-Specialist will be audited as part of the Ministry of Health's routine auditing procedures.

5.4 Pharmaceutical Cancer Treatments

- 5.4.1 DHBs must provide access to Pharmaceutical Cancer Treatments for the treatment of cancers in their DHB hospitals, and/or in association with Outpatient services provided in their DHB hospitals.
- 5.4.2 DHBs must only provide access to Pharmaceuticals for the treatment of cancer that are listed as Pharmaceutical Cancer Treatments in Sections A to G of the Schedule, provided that DHBs may provide access to an unlisted pharmaceutical for the treatment of cancer where that unlisted pharmaceutical:
 - a) has Named Patient Pharmaceutical Assessment (NPPA) approval;
 - b) is being used as part of a bona fide clinical trial which has Ethics Committee approval;
 - c) is being used and funded as part of a paediatric oncology service; or
 - d) was being used to treat the patient in question prior to 1 July 2005.
- 5.4.3 A DHB hospital pharmacy that holds a claiming agreement for Pharmaceutical Cancer Treatments with the Funder may claim a Subsidy for a Pharmaceutical Cancer Treatment marked as "PCT" or "PCT only" in Sections A to G of this Schedule subject to that Pharmaceutical Cancer Treatment being dispensed in accordance with:
 - a) Part 1;
 - b) clauses 2.1 to 2.2;

- c) clauses 3.1 to 3.4; and
- d) clause 5.4,

of Section A of the Schedule

- 5.4.4 A Contractor (other than a DHB hospital pharmacy) may only claim a Subsidy for a Pharmaceutical Cancer Treatment marked as "PCT" in Sections A to G of the Schedule subject to that Pharmaceutical Cancer Treatment being dispensed in accordance with the rules applying to Sections A to G of the Schedule.
- 5.4.5 Some indications for Pharmaceutical Cancer Treatments listed in the Schedule are Unapproved Indications. Some of these formed part of the October 2001 decision by the Minister of Health as to pharmaceuticals and indications for which DHBs must provide access. As far as reasonably practicable, these Unapproved Indications are marked in the Schedule. However, PHARMAC makes no representation and gives no guarantee as to the accuracy of this information. Prescribers of Pharmaceutical Cancer Treatments for such Unapproved Indications should:
 - a) be aware of and comply with their obligations under sections 25 and 29 of the Medicines Act 1981, as applicable, and otherwise under that act and the Medicines Regulations 1984;
 - b) be aware of and comply with their obligations under the Health and Disability Commissioner's Code of Consumer Rights, including the requirement to obtain informed consent from the patient (PHARMAC recommends that Prescribers obtain written consent); and
 - c) exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of an unapproved Pharmaceutical Cancer Treatment or a Pharmaceutical Cancer Treatment for an Unapproved Indication.
- 5.4.6 Applications to add pharmaceuticals, and add or amend indications for Pharmaceutical Cancer Treatments, may be made in writing by pharmaceutical suppliers and/or clinicians to PHARMAC. Applications should follow the Guidelines for Funding Applications to PHARMAC 2010 and Recommended methods to derive clinical inputs for proposals to PHARMAC, copies of which are available from PHARMAC or PHARMAC's website.

5.5 Prescribers of unapproved Pharmaceuticals

Prescribers should, where possible, prescribe Pharmaceuticals that are approved under the Medicines Act 1981. However, the access criteria under which a Pharmaceutical is listed on the Pharmaceutical Schedule may:

- a) in some case, explicitly permit Government funded access to a Pharmaceutical that is not approved under the Medicines Act 1981 or for an Unapproved Indication; or
- b) not explicitly preclude Government funded access to a Pharmaceutical when it is used for an Unapproved Indication;

Accordingly, if Prescribers are planning on prescribing an unapproved Pharmaceutical or a Pharmaceutical for an Unapproved Indication, Prescribers should:

- a) be aware of and comply with their obligations under sections 25 and 29 of the Medicines Act 1981, as applicable, and otherwise under that Act and the Medicines Regulations 1984;
- b) be aware of and comply with their obligations under the Health and Disability Commissioner's Code of Consumer Rights, including the requirement to obtain informed consent from the patient (PHARMAC recommends that Prescribers obtain written consent); and
- c) exercise their own skill, judgment, expertise and discretion, and make their own prescribing decisions with respect to the use of an unapproved Pharmaceutical or a Pharmaceutical for an Unapproved Indication.

Prescribers should be aware that simply by listing a Pharmaceutical on the Pharmaceutical Schedule PHARMAC makes no representations about whether that Pharmaceutical has any form of approval or consent under, or whether the supply or use of the Pharmaceutical otherwise complies with, the Medicines Act 1981. Further, the Pharmaceutical Schedule does not constitute an advertisement, advertising material or a medical advertisement as defined in the Medicines Act or otherwise.

5.6 Substitution

Where a Prescriber has prescribed a brand of a Community Pharmaceutical that has no Subsidy or has a Manufacturer's Price that is greater than the Subsidy and there is an alternative fully subsidised Community Pharmaceutical available, a Contractor may dispense the fully subsidised Community Pharmaceutical, unless either or both of the following circumstances apply:

- a) there is a clinical reason why substitution should not occur; or
- b) the prescriber has marked the prescription with a statement such as 'no brand substitution permitted'

Such an Authority to Substitute is valid whether or not there is a financial implication for the Pharmaceutical Budget. When dispensing a subsidised alternative brand, the Contractor must annotate and sign the prescription and inform the patient of the brand change.

5.7 Alteration to Presentation of Pharmaceutical Dispensed

A Contractor, when dispensing a subsidised Community Pharmaceutical, may alter the presentation of a Pharmaceutical dispensed to another subsidised presentation but may not alter the dose, frequency and/or total daily dose. This may only occur when it is not practicable for the contractor to dispense the requested presentation. If the change will result in additional cost to the DHBs, then annotation of the prescription by the dispensing pharmacist must occur stating the reason for the change, and the Contractor must initial the change for the purposes of Audit.

5.8 Other DHB Funding

A DHB may fund a Community Pharmaceutical outside of the mechanisms established in the Pharmaceutical Schedule, provided that:

- a) specific prior agreement is obtained from PHARMAC for such funding;
- b) any funding restrictions set out in the Pharmaceutical Schedule for those Community Pharmaceuticals are applied; and
- c) a Contractor (including a DHB Hospital Pharmacy) may not claim a Subsidy for a Community Pharmaceutical dispensed and funded by the DHB via such an alternate mechanism.

5.9 Conflict in Provisions

If any rules in Sections B-G and Section I of this Schedule conflict with the rules in Section A, the rules in Sections B-G and Section I apply.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Ful Subsidise Per	,
Antacids and Antiflatulants	Ŷ		Mandaotaroi
Antacids and Reflux Barrier Agents			
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg pe sachet		30 🗸	Gaviscon Infant
SODIUM ALGINATE * Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (8.60)	60	Gaviscon Double Strength
* Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 ml	Acidex
Phosphate Binding Agents			
ALUMINIUM HYDROXIDE * Tab 600 mg CALCIUM CARBONATE Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsement Only when prescribed for children under 12 years of age		500 ml 🗸	Alu-Tab Roxane ent and the prescription is
Antidiarrhoeals Agents Which Reduce Motility			
LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a * Tab 2 mg * Cap 2 mg	10.75		Nodia Diamide Relief
Rectal and Colonic Anti-inflammatories			
BUDESONIDE Cap 3 mg − Special Authority see SA1155 below − Retail pharmacy			Entocort CIR ns for applications meeting
Both: 1 Mild to moderate ileal, ileocaecal or proximal Crohn's dise 2 Any of the following: 2.1 Diabetes; or 2.2 Cushingoid habitus; or 2.3 Osteoporosis where there is significant risk of fract			
			continued.

20

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🖌	Manufacturer	

continued...

2.4 Severe acne following treatment with conventional corticosteroid therapy; or

2.5 History of severe psychiatric problems associated with corticosteroid treatment; or

2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or

2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an Unapproved Indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)	21.1 g OP	✓ Colifoam
MESALAZINE		
Tab 400 mg	100	Asacol
Tab EC 500 mg	100	Asamax
Tab long-acting 500 mg59.05	100	 Pentasa
Tab 800 mg	90	Asacol
Modified release granules, 1 g141.72	120 OP	 Pentasa
Enema 1 g per 100 ml41.30	7	 Pentasa
Suppos 500 mg22.80	20	Asacol
Suppos 1 g54.60	30	Pentasa
OLSALAZINE		
Tab 500 mg93.37	100	 Dipentum
Cap 250 mg53.00	100	 Dipentum
SODIUM CROMOGLICATE		
Cap 100 mg92.91	100	 Nalcrom
SULFASALAZINE		
* Tab 500 mg – For sulfasalazine oral liquid formulation refer,		
page 22414.00	100	 Salazopyrin
* Tab EC 500 mg13.50	100	 Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE

Oint 950 mcg, with fluocortolone pivalate 920 mcg, and			
cinchocaine hydrochloride 5 mg per g6.3	35 3	0gOP v	 Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and			
cinchocaine hydrochloride 1 mg2.6	66	12	 Ultraproct
HYDROCORTISONE WITH CINCHOCAINE			
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.0	00 3	0gOP v	Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.		12	Proctosedyl

‡ safety cap

	Subsidy		Fullv	Brand or
	(Manufacturer's Price)	Subsid	dised	Generic
	\$	Per	1	Manufacturer
Management of Anal Fissures				
GLYCERYL TRINITRATE – Special Authority see SA1329 below * Oint 0.2%		30 g OP	✔ R	ectogesic
➡SA1329 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals va chronic anal fissure that has persisted for longer than three wee		ewal unless	notified	d where the patient has a
chonic anal insolie that has persisted for longer than three wee	:KS.			
Antispasmodics and Other Agents Altering Gu	ıt Motility			
GLYCOPYRRONIUM BROMIDE				
Inj 200 mcg per ml, 1 ml ampoule - Up to 10 inj available o				
PSO	17.14	10	✓ M	ax Health
HYOSCINE BUTYLBROMIDE * Tab 10 mg	8 75	100	л в	uscopan
 * Inj 20 mg, 1 ml – Up to 5 inj available on a PSO 		5		uscopan
MEBEVERINE HYDROCHLORIDE				
* Tab 135 mg		90	✓ C	olofac
Antiulcerants				
Antisecretory and Cytoprotective				
MISOPROSTOL * Tab 200 mcg	41 50	120	✓ C	ytotec
			-	<u>,</u>
Helicobacter Pylori Eradication				
CLARITHROMYCIN	10.40			
Tab 500 mg – Subsidy by endorsement a) Maximum of 14 tab per prescription		14	• <u>A</u>	po-Clarithromycin
b) Subsidised only if prescribed for helicobacter pylori	eradication and press	ription is en	dorsed	accordingly.
Note: the prescription is considered endorsed if clarith and either amoxicillin or metronidazole.	romycin is prescribed	in conjunction	on with	a proton pump inhibitor
H2 Antagonists				
RANITIDINE – Only on a prescription				
* Tab 150 mg * Tab 300 mg		500 500	_	<u>anitidine Relief</u> anitidine Relief
* Oral lig 150 mg per 10 ml		300 ml		eptisoothe
* Inj 25 mg per ml, 2 ml	8.75	5	✓ Z	antac
Proton Pump Inhibitors				
LANSOPRAZOLE				
* Cap 15 mg		100	_	anzol Relief
* Cap 30 mg		100	• []	anzol Relief

Xifaxan

56

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
ОМ	EPRAZOLE				
	For omeprazole suspension refer Standard Formulae, page	227			
*	Cap 10 mg	1.98	90	1	Omeprazole actavis 10
*	Cap 20 mg	1.96	90	1	Omeprazole actavis 20
*	Cap 40 mg	3.12	90	1	Omeprazole actavis 40
*	Powder – Only in combination		5 g	1	Midwest
	Only in extemporaneously compounded omeprazole sus		Ũ		
*	Inj 40 mg ampoule with diluent		5	1	Dr Reddy's Omeprazole
PAN	NTOPRAZOLE				
*	Tab EC 20 mg	2.41	100	✓	Panzop Relief
	Tab EC 40 mg		100	1	Panzop Relief
Si	ite Protective Agents				
COI	LLOIDAL BISMUTH SUBCITRATE				
	Tab 120 mg		50	1	Gastrodenol S29
CI 1	CRALFATE				
300	Tab 1 g	35 50	120		
		(48.28)	120		Carafate
В	ile and Liver Therapy				

RIFAXIMIN – Special Authority see SA1461 below – Retail pharmacy	
Tab 550 mg	

➡SA1461 Special Authority for Subsidy

Initial application only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid for 6 months where the patient has hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

Renewal only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Diabetes

Hyperglycaemic Agents

Cap 25 mg	110.00	100	Proglicem S29
Cap 100 mg		100	Proglicem S29
Oral liq 50 mg per ml	620.00	30 ml OP	Proglycem S29
⇒SA1320 Special Authority for Subsidy			
nitial application from any relevant practitioner. Approval pypoglycaemia caused by hyperinsulinism. Renewal from any relevant practitioner. Approvals valid wit appropriate and the patient is benefiting from treatment. GLUCAGON HYDROCHLORIDE			

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Pr \$	ice) Subs Per	Fully sidised	Brand or Generic Manufacturer
Insulin - Short-acting Preparations				
NSULIN NEUTRAL ▲ Inj human 100 u per ml	25.26	10 ml OP		Actrapid
▲ Inj human 100 u per ml, 3 ml	42.66	5	✓ ,	Humulin R Actrapid Penfill Humulin R
Insulin - Intermediate-acting Preparations				
NSULIN ASPART WITH INSULIN ASPART PROTAMINE	52.15	5	~	NovoMix 30 FlexPen
NSULIN ISOPHANE ▲ Inj human 100 u per ml	17.68	10 ml OP		Humulin NPH
▲ Inj human 100 u per ml, 3 ml	29.86	5	✓	Protaphane Humulin NPH Protaphane Penfill
NSULIN ISOPHANE WITH INSULIN NEUTRAL Inj human with neutral insulin 100 u per ml	25.26	10 ml OP		Humulin 30/70 Mixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	\ \ \	Humulin 30/70 PenMix 30 PenMix 40 PenMix 50
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			•	rennix 50
3 ml Inj lispro 50% with insulin lispro protamine 50% 100 u per ml,		5		Humalog Mix 25
3 ml		5	•	Humalog Mix 50
Insulin - Long-acting Preparations				
NSULIN GLARGINE Inj 100 u per ml, 10 ml Inj 100 u per ml, 3 ml	63.00 94.50	1 5		Lantus Lantus
Inj 100 u per ml, 3 ml disposable pen		5	✓	Lantus SoloStar
Insulin - Rapid Acting Preparations				
NSULIN ASPART Inj 100 u per ml, 3 ml syringe Inj 100 u per ml, 3 ml Inj 100 u per ml, 10 ml	51.19	5 5 1	✓	NovoRapid FlexPen NovoRapid Penfill NovoRapid
NSULIN GLULISINE Inj 100 u per ml, 10 ml Inj 100 u per ml, 3 ml Inj 100 u per ml, 3 ml disposable pen	46.07	1 5 5	✓ .	Apidra Apidra Apidra SoloStar
NSULIN LISPRO ▲ Inj 100 u per ml, 10 ml ▲ Inj 100 u per ml, 3 ml		10 ml OP 5		Humalog Humalog

24

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	
Alpha Glucosidase Inhibitors				
ACARBOSE * Tab 50 mg * Tab 100 mg		90 90		<u>Glucobay</u> <u>Glucobay</u>
Oral Hypoglycaemic Agents				
GLIBENCLAMIDE * Tab 5 mg	5.00	100	1	Daonil
GLICLAZIDE * Tab 80 mg		500	1	Glizide
GLIPIZIDE * Tab 5 mg		100	1	Minidiab
METFORMIN HYDROCHLORIDE * Tab immediate-release 500 mg * Tab immediate-release 850 mg	9.59	1,000 500		<u>Metchek</u> Metformin Mylan
PIOGLITAZONE * Tab 15 mg * Tab 30 mg	3.47 5.06	90 90 90	<i>•</i> <i>•</i>	<u>Vexazone</u> Vexazone
 * Tab 45 mg Diabetes Management 		50	·	Vexazone
Ketone Testing				
 BLOOD KETONE DIAGNOSTIC TEST STRIP – Subsidy by er a) Not on a BSO b) Maximum of 20 strip per prescription c) Up to 10 strip available on a PSO d) Patient has any of the following: type 1 diabetes; or permanent neonatal diabetes; or undergone a pancreatectomy; or cystic fibrosis-related diabetes; or metabolic disease or epilepsy under the care of a The prescription must be endorsed accordingly. Test strips	paediatrician, neurolog	gist or n strip O		specialist. KetoSens
BLOOD KETONE DIAGNOSTIC TEST METER – Up to 1 meter Meter funded for the purposes of blood ketone diagnostics at risk of future episodes or patient is on an insulin pump. Meter	only. Patient has had Only one meter per pa		Il be subs	
 (Freestyle Optium Neo Meter to be delisted 1 August 2018) KETONE BLOOD BETA-KETONE ELECTRODES a) Maximum of 20 strip per prescription b) Up to 10 strip available on a PSO 				
Test strip – Not on a BSO		strip O	P 🗸	Freestyle Optium

(Freestyle Optium Ketone Test strip to be delisted 1 August 2018)

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Ketone

	Subsidy (Manufacturer's Price \$	e) Subs Per	Fully idised	Brand or Generic Manufacturer
SODIUM NITROPRUSSIDE – Maximum of 50 strip per prescrip * Test strip – Not on a BSO		50 strip OP	✔ К	etostix
Dual Blood Glucose and Blood Ketone Testing	I			
 DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC a) Maximum of 1 pack per prescription b) Up to 1 pack available on a PSO c) Note: may be provided by a pharmacist under the non-red d) A dual blood glucose and blood ketone diagnostic test m 1) type 1 diabetes; or 2) permanent neonatal diabetes; or 3) undergone a pancreatectomy; or 4) cystic fibrosis-related diabetes, or 5) metabolic disease or epilepsy under the care of a record of prior dispensing of insulin or sulphonylureas. Only 1 meter per patient will be subsidised (no repeat presc) For the avoidance of doubt patients who have previously refunded CareSens meter. From 1 February 2018 – 31 July 2018 patients who have us strips, as their only blood glucose diagnostic testing meter a meet the funding criteria. Meter with 50 lancets, a lancing device and 10 blood glucose diagnostic test strips. a) Brand switch fee payable (Pharmacode 2535890) - b) No patient co-payment payable c) CareSens Dual to be Sole Supply on 1 August 2018 	prescribing Practition neter is subsidised for paediatrician, neurol- sts may annotate the criptions). ceived a funded met sed a CareSens II bla and strips, are eligible se 	ogist or meta prescription er, other that bod glucose e for a new C 1 OP	ns in Pa rho has ubolic sp as end n Cares diagnos CareSer	art III of Section A. : pecialist. orsed where there exists Sens, are eligible for a stic meter and associated

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sul	bsidised	Generic
\$	Per	1	Manufacturer

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) Note: may be provided by a pharmacist under the non-prescribing Practitioners provisions in Part III of Section A.
- d) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylureas.

Only one CareSens meter per patient will be subsidised (no repeat prescriptions).

Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they meet the criteria for a dual blood glucose and blood ketone diagnostic test meter.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

From 1 February 2018 – 31 July 2018 patients who have used a CareSens II blood glucose diagnostic meter and associated strips, as their only blood glucose diagnostic testing meter and strips, are eligible for a new CareSens meter provided they meet the funding criteria.

20.00

Meter with 50 lancets, a lancing device and 10 diagnostic test



a) CareSens N brand: Brand switch fee payable (Pharmacode 2423138) - see page 221 for details

b) CareSens N POP brand: Brand switch fee payable (Pharmacode 2423154) - see page 221 for details

- c) CareSens N Premier brand: Brand switch fee payable (Pharmacode 2535882) see page 221 for details
- d) Note: Only 1 meter available per PSO

e) CareSens N to be Sole Supply on 1 August 2018

- f) CareSens N POP to be Sole Supply on 1 August 2018
- g) CareSens N Premier to be Sole Supply on 1 August 2018

Meter with 50 × lancets. 10 × diagnostic test strips and a

lancing device		1 OP	CareSens II
CareSens II Meter with 50 x lancets	10 x diagnostic test strips and a lancing device	to he del	listed 1 August 2018)

Subsidy		Fully	Brand or
(Manufacturer's Price)	Si	ubsidised	Generic
\$	Per	✓	Manufacturer

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- 2) Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed: or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Borforma	Test strips – Note differing brand rea		CareSens CareSens N CareSens PRO Accu-Chek
	a) Accur Chak Parforma brand:	 Image: A set of the set of the	Performa Freestyle Optium

- -Onek Penorma brand: Special Authority see SA1294 below Retall pharmacy
- b) Freestyle Optium brand: Special Authority see SA1291 below Retail pharmacy
- c) Note: Accu-Chek Performa and Freestyle Optium are not available on a PSO
- d) CareSens N to be Sole Supply on 1 August 2018
- e) CareSens PRO to be Sole Supply on 1 August 2018

(CareSens Test strips to be delisted 1 August 2018)

(Accu-Chek Performa Test strips to be delisted 1 August 2018)

(Freestyle Optium Test strips to be delisted 1 August 2018)

SA1294 Special Authority for Subsidy

Notes: Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz and can be sent to:

PHARMAC

PO Box 10 254 Facsimile: (04) 974 4788

Email: bgstrips@pharmac.govt.nz Wellington

SA1291 Special Authority for Subsidy

Notes: Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz and can be sent to:

PHARMAC

PO Box 10 254 Facsimile: (04) 974 4788

Wellington Email: bgstrips@pharmac.govt.nz

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- 2) Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed: or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

50 test OP ✓ SensoCard

		Subsidy (Manufacturer's Price) \$	Ful Subsidise Per	,
Ir	nsulin Syringes and Needles			
the	bsidy is available for disposable insulin syringes, needles, and supply of insulin or when prescribed for an insulin patient and notate the prescription as endorsed where there exists a recor	the prescription is er	ndorsed accord	
INS	SULIN PEN NEEDLES – Maximum of 100 dev per prescription	n		
*	29 g × 12.7 mm		100 🖌	B-D Micro-Fine
*	31 g × 5 mm	11.75	100 💌	B-D Micro-Fine
*	31 g × 6 mm		100 🖌	ABM
*	31 g × 8 mm		100 🖌	B-D Micro-Fine
*	32 g × 4 mm		100 💌	B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLI	E – Maximum of 100	dev per presc	ription
	Syringe 0.3 ml with 29 g × 12.7 mm needle			B-D Ultra Fine
	, , , , , , , , , , , , , , , , , , , ,	1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g × 8 mm needle		100 🖌	B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g × 12.7 mm needle		100 🖌	B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle		100 🖌	B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 1 ml with 29 g × 12.7 mm needle		100 🖌	B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle		100 🖌	B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 on the next page - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescriptionc) Maximum of 1 insulin pump per patient each four year period.

Min basal rate 0.025 U/h; black colour		1	 Animas Vibe
Min basal rate 0.025 U/h; blue colour	4,500.00	1	 Animas Vibe
Min basal rate 0.025 U/h; green colour	4,500.00	1	 Animas Vibe
Min basal rate 0.025 U/h; pink colour	4,500.00	1	 Animas Vibe
Min basal rate 0.025 U/h; silver colour		1	 Animas Vibe
Min basal rate 0.05 U/h; blue colour		1	Paradigm 522
	·		Paradigm 722
Min basal rate 0.05 U/h; clear colour		1	Paradigm 522
	·		Paradigm 722
Min basal rate 0.05 U/h; pink colour		1	Paradigm 522
	·		Paradigm 722
Min basal rate 0.05 U/h; purple colour		1	Paradigm 522
	·		Paradigm 722
Min basal rate 0.05 U/h; smoke colour		1	Paradigm 522
			Paradigm 722

‡ safety cap

▲ Three months supply may be dispensed at one time

*Three months or six months, as applicable, dispensed all-at-once

if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and

3 Either:

- 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
- 3.2 The pump is due for replacement; and

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

4 Either:

4.1 Applicant is a relevant specialist; or

4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and

*Three months or six months, as applicable, dispensed all-at-once

- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and

8 Either:

8.1 It has been at least 4 years since the last insulin pump was received by the patient; or

continued...

‡ safety cap

	Subsidy	F	ully	Brand or
(Ma	anufacturer's Price)	Subsidi	ised	Generic
	\$	Per	✓	Manufacturer

continued...

8.2 The pump is due for replacement; and

9 Either:

9.1 Applicant is a relevant specialist; or

9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Either:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Insulin Pump Consumables

⇒SA1604 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and

continued...

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	~	Manufacturer

continued...

- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and

3 Either:

3.1 Applicant is a relevant specialist; or

*Three months or six months, as applicable, dispensed all-at-once

3.2 Applicant is a nurse practitioner working within their vocational scope.

continued...

‡ safety cap

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

continued...

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

INSULIN PUMP ACCESSORIES - Special Authority see SA1604 on page 32 - Retail pharmacy

- a) Maximum of 1 cap per prescription
- b) Only on a prescription
- c) Maximum of 1 prescription per 180 days.

Battery cap				
-------------	--	--	--	--

1

Animas Battery Cap

	Subsidy (Manufacturer's Pric \$	ce) Per	Fully Subsidised	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (STEEL CANNULA) - Special	Authority see SA16	604 on pa	age 32 – F	Retail pharmacy
 a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 				
10 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles		1 OP	1	Paradigm Sure-T MMT-884
10 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock	130.00	1 OP	1	Sure-T MMT-883
10 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles	130.00	1 OP	1	Paradigm Sure-T MMT-886
10 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles; luer lock		1 OP	1	Sure-T MMT-885
6 mm steel cannula; straight insertion; 60 cm grey line × 10 v 10 needles		1 OP	1	Contact-D
6 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles	130.00	1 OP	1	Paradigm Sure-T MMT-864
6 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock 6 mm steel needle: 29 G; manual insertion; 80 cm tubing ×		1 OP	1	Sure-T MMT-863
10 with 10 needles	130.00	1 OP	1	Paradigm Sure-T MMT-866
6 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles; luer lock		1 OP	1	Sure-T MMT-865
8 mm steel cannula; straight insertion; 110 cm grey line × 10 with 10 needles		1 OP	1	Contact-D
8 mm steel cannula; straight insertion; 60 cm grey line × 10 v 10 needles		1 OP	1	Contact-D
8 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles	130.00	1 OP	1	Paradigm Sure-T MMT-874
8 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock		1 OP	1	Sure-T MMT-873
8 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles	130.00	1 OP	1	Paradigm Sure-T MMT-876
8 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles; luer lock		1 OP	1	Sure-T MMT-875
 INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE II SA1604 on page 32 – Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 13 mm teflon cannula; angle insertion; insertion device; 110 device 	cm			
grey line × 10 with 10 needles 13 mm teflon cannula; angle insertion; insertion device; 60 cr	m	1 OP		Inset 30
grey line × 10 with 10 needles	140.00	1 OP	1	Inset 30

\$\$ safety cap
\$\$ Three months or six months, as applicable, dispensed all-at-once

if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
ISULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE II	NSERTION) – Speci	al Au	thority see	SA1604 on page 32 -
etail pharmacy				
 Maximum of 3 sets per prescription 				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
13 mm teflon cannula; angle insertion; 120 cm line \times 10 with				
10 needles		1 OP		Paradigm Silhouette MMT-382
13 mm teflon cannula; angle insertion; 45 cm line × 10 with				
10 needles	130.00	1 OP		Paradigm Silhouette MMT-368
13 mm teflon cannula; angle insertion; 60 cm line \times 10 with				
10 needles		1 OP	-	Paradigm Silhouette MMT-381
13 mm teflon cannula; angle insertion; 80 cm line \times 10 with				
10 needles	130.00	1 OP	1	Paradigm Silhouette MMT-383
17 mm teflon cannula; angle insertion; 110 cm line \times 10 with				
10 needles		1 OP	-	Paradigm Silhouette MMT-377
17 mm teflon cannula; angle insertion; 110 cm line × 10 with				
10 needles; luer lock		1 OP	✓	Silhouette MMT-371
17 mm teflon cannula; angle insertion; 60 cm line × 10 with				
10 needles	130.00	1 OP		Paradigm Silhouette MMT-378
17 mm teflon cannula; angle insertion; 60 cm line \times 10 with				
10 needles; luer lock		1 OP	✓	Silhouette MMT-373
17 mm teflon cannula; angle insertion; 80 cm line × 10 with				
10 needles		1 OP	-	Paradigm Silhouette MMT-384

	Subsidy		Fully Brand or
	(Manufacturer's Pi \$	rice) Sub Per	osidised Generic Manufacturer
SULIN PUMP INFUSION SET (TEFLON CANNULA, STRAI	*	-	
e SA1604 on page 32 – Retail pharmacy	0		
a) Maximum of 3 sets per prescription			
b) Only on a prescription			
c) Maximum of 13 infusion sets will be funded per year.			
6 mm teflon cannula; straight insertion; insertion device;			
110 cm grey line × 10 with 10 needles		1 OP	 Inset II
6 mm teflon cannula; straight insertion; insertion device; 4	5 cm		
blue tubing × 10 with 10 needles		1 OP	Paradigm Mio
			MMT-941
6 mm teflon cannula; straight insertion; insertion device; 4			
pink tubing × 10 with 10 needles		1 OP	🗸 Paradigm Mio
			MMT-921
6 mm teflon cannula; straight insertion; insertion device; 6			
blue tubing × 10 with 10 needles		1 OP	 Paradigm Mio
			MMT-943
6 mm teflon cannula; straight insertion; insertion device; 6			• · · · ·
grey line × 10 with 10 needles		1 OP	 Inset II
6 mm teflon cannula; straight insertion; insertion device; 6			
pink tubing × 10 with 10 needles		1 OP	 Paradigm Mio
			MMT-923
6 mm teflon cannula; straight insertion; insertion device; 8		4.00	
blue tubing × 10 with 10 needles		1 OP	 Paradigm Mio
O must be first a second by a basis bet in a set in a single state in a second s	0		MMT-945
6 mm teflon cannula; straight insertion; insertion device; 8		1 00	
clear tubing × 10 with 10 needles		1 OP	 Paradigm Mio MMT-965
6 mm teflon cannula; straight insertion; insertion device; 8	0.0m		WIWI - 905
pink tubing × 10 with 10 needles		1 OP	 Paradigm Mio
		101	MMT-925
9 mm teflon cannula; straight insertion; insertion device;			
110 cm grey line × 10 with 10 needles		1 OP	✓ Inset II
9 mm teflon cannula; straight insertion; insertion device; 6			
grey line × 10 with 10 needles		1 OP	✓ Inset II
9 mm teflon cannula; straight insertion; insertion device; 8		-	
clear tubing × 10 with 10 needles		1 OP	 Paradigm Mio
č			MMT-975

	Subsidy (Manufacturer's Priv \$	ce) Per	Fully Subsidised	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH	T INSERTION) -	- Special	Authority s	ee SA1604 on page 32 -
Retail pharmacy a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; 110 cm tubing × 10 wi	th			
10 needles		1 OP	✓	Paradigm Quick-Set MMT-398
6 mm teflon cannula; straight insertion; 110 cm tubing × 10 wi	th			
10 needles; luer lock	130.00	1 OP		Quick-Set MMT-391
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with	า			
10 needles	130.00	1 OP	✓	Paradigm Quick-Set
				MMT-399
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with				
10 needles; luer lock		1 OP	•	Quick-Set MMT-393
6 mm teflon cannula; straight insertion; 80 cm tubing × 10 with		4.00		
10 needles		1 OP	✓	Paradigm Quick-Set
0 mm tellen som det staright insertions 100 om tubions 10 ui	44-			MMT-387
9 mm teflon cannula; straight insertion; 106 cm tubing × 10 wi 10 needles		1 OP		Paradigm Quick-Set
To needles	130.00	TOP	•	MMT-396
9 mm teflon cannula; straight insertion; 110 cm tubing $ imes$ 10 wi	th			WIWI 1-390
10 needles: luer lock		1 OP	1	Quick-Set MMT-390
9 mm teflon cannula; straight insertion; 60 cm tubing \times 10 with		101	•	Guick-Set Wiwi 1-550
10 needles		1 OP	~	Paradigm Quick-Set
		1 01		MMT-397
9 mm teflon cannula; straight insertion; 60 cm tubing \times 10 with	n			
10 needles; luer lock		1 OP	1	Quick-Set MMT-392
9 mm teflon cannula; straight insertion; 80 cm tubing × 10 with				
10 needles		1 OP	 Image: A second s	Paradigm Quick-Set
				MMT-386
INSULIN PUMP RESERVOIR - Special Authority see SA1604 or	page 32 – Retai	l pharma	cv	
a) Maximum of 3 sets per prescription	1.00	F	.,	
b) Only on a prescription				
c) Maximum of 13 packs of reservoir sets will be funded per				
10 × luer lock conversion cartridges 1.8 ml for Paradigm pum		1 OP		ADR Cartridge 1.8
Cartridge 200 U, luer lock × 10		1 OP		Animas Cartridge
Cartridge for 5 and 7 series pump; 1.8 ml \times 10		1 OP	✓	Paradigm
		_	_	1.8 Reservoir
Cartridge for 7 series pump; 3.0 ml × 10		1 OP	✓	Paradigm
				3.0 Reservoir
Syringe and cartridge for 50X pump, 3.0 ml × 10		1 OP	v !	50X 3.0 Reservoir

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per		Manufacturer
	100	✓ <u>(</u>	Creon 10000
e, 94.40	100	√ I	Panzytrat
94.38	100	✓ <u>(</u>	Creon 25000
<mark>elow</mark> – Retail pharmad	зy		
ion 37.95	100	✓ <u>I</u>	Jrsosan
	(Manufacturer's Price) \$ 	(Manufacturer's Price) \$ Per 	(Manufacturer's Price) Subsidised \$ Per •

⇒SA1383 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Cirrhosis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cirrhosis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

*Three months or six months, as applicable, dispensed all-at-once

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Cirrhosis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains

continued...

‡ safety cap

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Laxatives

Bulk-forming Agents

6.05	500 g OP	 ✓ Bonvit ✓ Konsyl-D
6.02 (17.32) 2.41 (8.72)	500 g OP 200 g OP	Normacol Plus Normacol Plus
2.31 3.13 5.40 3.10	100 100 100 ml OP 200 30 ml OP	 ✓ <u>Coloxyl</u> ✓ <u>Coloxyl</u> ✓ Coloxyl ✓ Laxsol ✓ <u>Coloxyl</u>
6 <mark>91 below –</mark> Re 36.00 246.00	tail pharmacy 1 7	✓ Relistor✓ Relistor
	6.02 (17.32) 2.41 (8.72) 3.13 5.40 3.10 3.78 3.78	

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

1 The patient is receiving palliative care; and

2 Either:

40

2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or

2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

(M	Subsidy anufacturer's Price \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Osmotic Laxatives				
GLYCEROL * Suppos 3.6 g – Only on a prescription	6.50	20	√ į	PSM
LACTULOSE – Only on a prescription X Oral liq 10 g per 15 ml	3.18	500 m	I ∕ I	Laevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICAR	RBONATE AND	SODIL	IM CHLOR	IDE
Powder for oral soln 13.125 g with potassium chloride 46.6 mg, sodium bicarbonate 178.5 mg and sodium chloride 350.7 m	g6.78	30	√ [Molaxole
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%	2.50	1	√	Fleet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE -	Only on a prescr	iption		
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml	26.72	50	√	Micolette
Stimulant Laxatives				
BISACODYL - Only on a prescription				
 * Tab 5 mg * Suppos 10 mg 		200 10	-	<u>Lax-Tab</u> Lax-Suppositories
SENNA – Only on a prescription			-	
* Tab, standardised	2.17 (6.84)	100	(Senokot
	0.43	20		
	(1.72)		9	Senokot

Metabolic Disorder Agents

ALGLUCOSIDASE ALFA - Special Authority see SA1622 below - Retail pharmacy 1

Myozyme

➡SA1622 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene: and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT): and

continued...

‡ safety cap

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

GALSULFASE - Special Authority see SA1593 below - Retail pharmacy

Inj 1 mg per ml, 5 ml vial......2,234.00 1 🖌 Naglazyme

⇒SA1593 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

IDURSULFASE – Special Authority see SA1623 below – Retail pharmacy

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:

- 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
- 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
continued 3 Patient is going to proceed with a haematopoietic stem c	oll transplant (USCT)	within	the payt 2 r	months and treatment with
 4 Patient has not required long-term invasive ventilation for (ERT); and 	, , ,			
 5 Idursulfase to be administered for a total of 24 weeks (ec greater than 0.5 mg/kg every week. 	uivalent to 12 weeks	pre- ar	nd 12 week	s post-HSCT) at doses no
LARONIDASE – Special Authority see SA1695 below – Retail p Inj 100 U per ml, 5 ml vial		1	✓ A	Idurazyme
■ SA1695 Special Authority for Subsidy Initial application only from a metabolic physician. Approvals we All of the following:	valid for 24 weeks for	applica	ations meet	ing the following criteria:
 The patient has been diagnosed with Hurler Syndrome (r Either: 	nucopolysacchardosi	s I-H);	and	
 Diagnosis confirmed by demonstration of alpha-L assay in cultured skin fibroblasts; or Detection of two disease causing mutations in the 				
to have Hurler syndrome; and 3 Patient is going to proceed with a haematopoietic stem c		•	·	·
laronidase would be bridging treatment to transplant; and 4 Patient has not required long-term invasive ventilation for	, , ,			
 (ERT); and 5 Laronidase to be administered for a total of 24 weeks (ec than 100 units/kg every week. 	uivalent to 12 weeks	pre- ar	nd 12 post-l	HSCT) at doses no greater
SODIUM BENZOATE - Special Authority see SA1599 below -	• •	100		mzoate S29
Soln 100 mg per ml SA1599 Special Authority for Subsidy Initial application only from a metabolic physician. Approvals v cycle disorder.		100 ml		
Renewal only from a metabolic physician. Approvals valid for 1 patient is benefiting from treatment.	2 months where the t	reatme	ent remains	appropriate and the
SODIUM PHENYLBUTYRATE – Special Authority see SA1598 Grans 483 mg per g		nacy 74 g O	P ⁄ F	Pheburane
► SA1598 Special Authority for Subsidy Initial application only from a metabolic physician. Approvals of cycle disorder involving a deficiency of carbamylphosphate synthetase.	netase, ornithine tran	scarba	mylase or a	rgininosuccinate
Renewal only from a metabolic physician. Approvals valid for 1 patient is benefiting from treatment.	2 months where the t	reatme	ent remains	appropriate and the
Gaucher's Disease				
IMIGLUCERASE - Special Authority see SA0473 on the next p	age – Retail pharmad	:V		

IMIGLUCERASE – Special Authority see SA0473 on the	next page – Retail pharma	су	
Inj 40 iu per ml, 200 iu vial		1	 Cerezyme
Inj 40 iu per ml, 400 iu vial	2,144.00	1	 Cerezyme

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's F \$	Price) S Per	Fully Subsidised	Brand or Generic Manufacturer
SA0473 Special Authority for Subsidy Special Authority approved by the Gaucher's Treat Notes: Subject to a budgetary cap. Applications of Application details may be obtained from PHARMA	will be considered and approve	,	0	vailability.
The Co-ordinator, Gaucher's Treatment Panel	Phone: (04) 460 4990			
PHARMAC, PO Box 10 254	Facsimile: (04) 916 7571			
Wellington	Email: gaucherpanel@pharr	nac.govt.nz	1	
Mouth and Throat				
Agents Used in Mouth Ulceration				
BENZYDAMINE HYDROCHLORIDE				
Soln 0.15% – Higher subsidy of up to \$17.01	1			
Endorsement		500 ml	_	
	(17.01)	000	[Difflam
	3.60 (8.50)	200 ml	г	Difflam
Additional subsidy by endorsement for a p prescription is endorsed accordingly.		as a result c	of treatmen	t for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PE Paste		56 g OF		Stomahesive
Faste	4.55	15 g OF		Stomanesive
	(7.90)	10 9 01		Drabase
	1.52	5 g OP		
	(3.60)	0	(Drabase
Powder	8.48	28 g OF		
	(10.95)		5	Stomahesive
CHLORHEXIDINE GLUCONATE				
Mouthwash 0.2%	2.57	200 ml O	P ∕	nealthE
CHOLINE SALICYLATE WITH CETALKONIUM C	HLORIDE			
Adhesive gel 8.7% with cetalkonium chloride (15 g OF		
	(6.00)		E	Bonjela
	5.33	5 g OP	✓ <u>I</u>	Kenalog in Orabase
Paste 0.1%				
TRIAMCINOLONE ACETONIDE Paste 0.1% Oropharyngeal Anti-infectives				
Paste 0.1% Oropharyngeal Anti-infectives AMPHOTERICIN B				
Paste 0.1% Oropharyngeal Anti-infectives		20	✓ F	Fungilin
Paste 0.1% Oropharyngeal Anti-infectives AMPHOTERICIN B Lozenges 10 mg	5.86			•
Paste 0.1% Oropharyngeal Anti-infectives AMPHOTERICIN B Lozenges 10 mg	5.86	20 40 g OF		⁻ ungilin Decozol
Paste 0.1% Oropharyngeal Anti-infectives AMPHOTERICIN B Lozenges 10 mg	5.86			•

	Subsidy		Fully Brand or	
	(Manufacturer's Pr \$	rice) Subsi Per	dised Generic Manufacturer	
	<u> </u>			
Other Oral Agents				
For folinic mouthwash, pilocarpine oral liquid or saliva substitute for	ormula refer Star	ndard Formulae	e, page 227	
HYDROGEN PEROXIDE				
Soln 3% (10 vol) – Maximum of 200 ml per prescription	1.40	100 ml	 Pharmacy Health 	
	0.15	500 ml		
* Compound, BPC	9.15	500 ml	✓ <u>PSM</u>	
Vitamins				
Vitamin A				
VITAMIN A WITH VITAMINS D AND C				
* Soln 1000 u with Vitamin D 400 u and ascorbic acid 30 mg pe	er			
10 drops		10 ml OP	 Vitadol C 	
Vitamin B				
HYDROXOCOBALAMIN			_	
* Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a PS	02.31	3	✓ <u>Neo-B12</u>	
PYRIDOXINE HYDROCHLORIDE				
a) No more than 100 mg per doseb) Only on a prescription				
 Tab 25 mg – No patient co-payment payable 	2.70	90	 Vitamin B6 25 	
* Tab 50 mg		500	 Apo-Pyridoxine 	
THIAMINE HYDROCHLORIDE – Only on a prescription			4 - - - - - -	
* Tab 50 mg	5.62	100	Apo-Thiamine	
VITAMIN B COMPLEX * Tab, strong, BPC	7 15	500	✓ Bplex	
		500		
Vitamin C				
ASCORBIC ACID				
a) No more than 100 mg per dose				
 b) Only on a prescription * Tab 100 mg 	9.10	500	✓ Cvite	
v		500	• <u>cvite</u>	
Vitamin D				
	06.00	100	. One Alaba	
* Cap 0.25 mcg * Cap 1 mcg		100 100	 ✓ <u>One-Alpha</u> ✓ One-Alpha 	
* Oral drops 2 mcg per ml		20 ml OP	✓ One-Alpha	
CALCITRIOL				
* Cap 0.25 mcg		100	 <u>Calcitriol-AFT</u> <u>Calcitriol AFT</u> 	
* Cap 0.5 mcg		100	 <u>Calcitriol-AFT</u> 	
COLECALCIFEROL * Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescription	on2.50	12	✓ <u>Vit.D3</u>	

‡ safety cap

 $\ensuremath{\boldsymbol{\ast}}$ Three months or six months, as applicable, dispensed all-at-once

	Subsidu		Fully Brand or
	Subsidy (Manufacturer's Price)		,
	\$	Per	 Manufacturer
Multivitamin Preparations			
MULTIVITAMIN RENAL – Special Authority see SA1546 below - * Cap	- Retail pharmacy 6.49	30	 Clinicians Renal Vit
SA1546 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Either:	d without further rene	ewal unless	notified for applications meeting
 The patient has chronic kidney disease and is receiving ei The patient has chronic kidney disease grade 5, defined a 15 ml/min/1.73 m² body surface area (BSA). 			
MULTIVITAMINS – Special Authority see SA1036 below – Retai		00 g OP	✓ Paediatric Seravit
■ SA1036 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid inborn errors of metabolism.			
Renewal from any relevant practitioner. Approvals valid without approval for multivitamins. VITAMINS	iuriner renewai unie:	ss nounied w	mere patient has had a previous
 * Tab (BPC cap strength) * Cap (fat soluble vitamins A, D, E, K) – Special Authority see 		1,000	✓ <u>Mvite</u>
 SA1720 below – Retail pharmacy SA1720 Special Authority for Subsidy 		60	 Vitabdeck
Initial application from any relevant practitioner. Approvals valion the following criteria:	d without further rene	ewal unless	notified for applications meeting
Any of the following:			
1 Patient has cystic fibrosis with pancreatic insufficiency; or			
 Patient is an infant or child with liver disease or short gut s Patient has severe malabsorption syndrome. 	syndrome; or		
Minerals			
Calcium			
CALCIUM CARBONATE			
* Tab eff 1.75 g (1 g elemental)		10 250	 ✓ Calsource ✓ Arrow-Calcium
* Tab 1.25 g (500 mg elemental)		250	Arrow-Calcium
CALCIUM GLUCONATE * Inj 10%, 10 ml ampoule		10	✓ Hospira
Fluoride			
SODIUM FLUORIDE * Tab 1.1 mg (0.5 mg elemental)	5.00	100	✓ PSM
lodine			
POTASSIUM IODATE * Tab 253 mcg (150 mcg elemental iodine)	4.69	90	✓ NeuroTabs

	Subsidy (Manufacturer's Pric \$	e) Su Per	Fully bsidised	Brand or Generic Manufacturer
Iron				
FERRIC CARBOXYMALTOSE - Special Authority see SA1675 Inj 50 mg per ml, 10 ml		macy 1	✓ F	erinject
■ SA1675 Special Authority for Subsidy Initial application — (serum ferritin less than or equal to 20 months for applications meeting the following criteria: Both:	mcg/L) from any m	edical prac	ctitioner.	Approvals valid for 3
 Patient has been diagnosed with iron-deficiency anaemia Any of the following: 				
2.1 Patient has been compliant with oral iron treatmer2.2 Treatment with oral iron has resulted in dose-limiti2.3 Rapid correction of anaemia is required.		s proven in	effective;	or
Renewal — (serum ferritin less than or equal to 20 mcg/L) fr applications meeting the following criteria: Both:	om any medical pra	actitioner.	Approval	s valid for 3 months for
 Patient continues to have iron-deficiency anaemia with a A re-trial with oral iron is clinically inappropriate. 				-
Initial application — (iron deficiency anaemia) only from an in anaesthetist or medical practitioner on the recommendation of a anaesthetist. Approvals valid for 3 months for applications meet Both:	internal medicine pl	hysician, o		
 Patient has been diagnosed with iron-deficiency anaemia Any of the following: 	; and			
 2.1 Patient has been compliant with oral iron treatmer 2.2 Treatment with oral iron has resulted in dose-limiti 2.3 Patient has symptomatic heart failure, chronic kidn and a trial of oral iron is unlikely to be effective; or 2.4 Rapid correction of anaemia is required. 	ng intolerance; or ney disease stage 3			
 Renewal — (iron deficiency anaemia) only from an internal m medical practitioner on the recommendation of a internal medicir Approvals valid for 3 months for applications meeting the followin Both: Patient continues to have iron-deficiency anaemia; and 2 A re-trial with oral iron is clinically inappropriate. 	ne physician, obstet			
FERROUS FUMARATE * Tab 200 mg (65 mg elemental)	2.89	100	✓ <u>F</u>	erro-tab
FERROUS FUMARATE WITH FOLIC ACID * Tab 310 mg (100 mg elemental) with folic acid 350 mcg Ferro-F-Tabs to be Sole Supply on 1 July 2018	4.68	60	✔ F	erro-F-Tabs
FERROUS SULPHATE * Tab long-acting 325 mg (105 mg elemental) Ferrograd to be Sole Supply on 1 July 2018	2.06	30	✔ F	errograd
*‡ Oral liq 30 mg (6 mg elemental) per 1 ml FERROUS SULPHATE WITH FOLIC ACID	10.80	500 ml	✓ <u>F</u>	<u>erodan</u>
* Tab long-acting 325 mg (105 mg elemental) with folic acid 350 mcg	1.80 (4.29)	30	F	errograd F
(Ferrograd F Tab long-acting 325 mg (105 mg elemental) with for		be delisted		

‡ safety cap

 $\ensuremath{\textbf{\#}}$ Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully osidised	Brand or Generic Manufacturer
IRON POLYMALTOSE * Inj 50 mg per ml, 2 ml ampoule		5	✔ F	errum H
Magnesium				
For magnesium hydroxide mixture refer Standard Formulae, page MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule		10	✓ <u>D</u>	BL
Zinc				
ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)	11.00	100	✓ z	incaps

Subsidised

Per

Fully

Subsidy (Manufacturer's Price) \$ Brand or Generic

Generic Manufacturer

Antianaemics

Hypoplastic and Haemolytic

► SA1469 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or

3.2 Both:

- 3.2.1 Patient has diabetes mellitus; and
- 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
- 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Erythropoietin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum erythropoietin level of < 500 IU/L; and
- 6 The minimum necessary dose of erythropoietin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an Unapproved Indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Erythropoietin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of erythropoietin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an Unapproved Indication

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
POETIN ALFA [ERYTHROPOIETIN ALFA] - Special Authorit	y see SA1469 on the	orevio	us page –	Retail pharmacy
Wastage claimable – see rule 3.3.2 on page 13	40.00	•		_
Inj 1,000 iu in 0.5 ml, syringe		6		Eprex
Inj 2,000 iu in 0.5 ml, syringe		6		Eprex
Inj 3,000 iu in 0.3 ml, syringe		6		Eprex
Inj 4,000 iu in 0.4 ml, syringe		6	~	Eprex
Inj 5,000 iu in 0.5 ml, syringe	243.26	6	✓	Eprex
Inj 6,000 iu in 0.6 ml, syringe		6	✓	Eprex
Inj 8,000 iu in 0.8 ml, syringe		6	✓	Eprex
Inj 10,000 iu in 1 ml, syringe		6	✓	Eprex
lnj 40,000 iu in 1 ml, syringe		1	✓	Eprex
Megaloblastic				
OLIC ACID				
Fab 0.8 mg		1.000	1	Apo-Folic Acid
K Tab 5 mg		500	-	Apo-Folic Acid
Oral liq 50 mcg per ml		5 ml C		Biomed

Antifibrinolytics, Haemostatics and Local Scierosants

ELTROMBOPAG – Special Authority se	e SA1418 below – Retail pharmacy
------------------------------------	----------------------------------

Wastage claimable – see rule 3.3.2 on page 13			
Tab 25 mg	1,771.00	28	 Revolade
Tab 50 mg	3,542.00	28	 Revolade

⇒SA1418 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia, whose funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 1 mg syringe1,178.30	1	NovoSeven RT
Inj 2 mg syringe2,356.60	1	NovoSeven RT
Inj 5 mg syringe	1	NovoSeven RT
Inj 8 mg syringe9,426.40	1	 NovoSeven RT

	Subsidy		Fully	
	(Manufacturer's Price) \$	S Per	ubsidised	Generic Manufacturer
FACTOR EIGHT INHIBITOR BYPASSING FRACTION -		1.01	•	Manulacturer
For patients with haemophilia, whose funded treatmer		philia ⁻	Freaters	Group in conjunction with
the National Haemophilia Management Group.	it is managed by the machine	prinia	incutoro	
Inj 500 U	1,450.00	1	✓	FEIBA NF
Inj 1,000 U		1		FEIBA NF
Inj 2,500 U	7,250.00	1	~	FEIBA NF
MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] -				
Preferred Brand of recombinant factor VIII for patients				
to funded treatment is managed by the Haemophilia T	reaters Group in conjunction	with t	he Natio	nal Haemophilia
Management Group.	010.00			Visitha
Inj 250 iu prefilled syringe		1 1		Xyntha Xyntha
Inj 500 iu prefilled syringe Inj 1.000 iu prefilled syringe		1		Xyntha
Inj 2,000 iu prefilled syringe		1		Xyntha
Inj 3,000 iu prefilled syringe	,	1		Xyntha
NONACOG ALFA [RECOMBINANT FACTOR IX] – [Xpha		•	-	<i>A</i> ynum
For patients with haemophilia, whose funded treatmer		nhilia ⁻	Treaters	Group in conjunction with
the National Haemophilia Management Group.	it is managed by the hadno	prilla	Tealers	
Inj 250 iu vial	310.00	1	1	BeneFIX
Inj 500 iu vial		1		BeneFIX
Inj 1,000 iu vial		1		BeneFIX
Inj 2,000 iu vial		1	-	BeneFIX
Inj 3,000 iu vial		1	~	BeneFIX
NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [X	pharm]			
For patients with haemophilia, whose funded treatmer	t is managed by the Haemo	philia ⁻	Freaters	Group in conjunction with
the National Haemophilia Management Group.				
Inj 250 iu vial		1		RIXUBIS
Inj 500 iu vial		1		RIXUBIS
Inj 1,000 iu vial	,	1		RIXUBIS
Inj 2,000 iu vial	,	1 1		RIXUBIS
Inj 3,000 iu vial	,	I	v	RIXUBIS
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVA				
Rare Clinical Circumstances Brand of recombinant fac				
28 February 2019. Access to funded treatment by ap be obtained from PHARMAC's website http://www.pha		reatri	ients Pal	nei. Application details may
The Co-ordinator, Haemophilia Treatments Panel	Phone: 0800 023 588 O	ption 2	2	
PHARMAC PO Box 10 254	Facsimile: (04) 974 4881			
Wellington	Email: <u>haemophilia@phar</u>	mac.g	ovt.nz	
Inj 250 iu vial	287 50	1	1	Advate
Inj 500 iu vial		1		Advate
Inj 1,000 iu vial		1		Advate
Inj 1,500 iu vial		1		Advate
Inj 2,000 iu vial	-	1	~	Advate
Inj 3,000 iu vial		1	1	Advate

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$) Su Per	Fully bsidised	Brand or Generic Manufacturer	
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGE Second Brand of recombinant factor VIII for patients w funded treatment by application to the Haemophilia Tr PHARMAC's website http://www.pharmac.govt.nz or:	vith haemophilia from 1 Mar				Access to
The Co-ordinator, Haemophilia Treatments Panel	Phone: 0800 023 588 (Option 2			
PHARMAC PO Box 10 254	Facsimile: (04) 974 488	1			
Wellington	Email: haemophilia@pha	armac.go	<u>vt.nz</u>		
Inj 250 iu vial		1		Kogenate FS	
Inj 500 iu vial		1		Kogenate FS	
Inj 1,000 iu vial		1 1		Cogenate FS	
Inj 2,000 iu vial Inj 3,000 iu vial	,	1		Cogenate FS	
SODIUM TETRADECYL SULPHATE				togenate i o	
* Inj 3% 2 ml	28.50	5			
	(73.00)	°,	F	ibro-vein	
TRANEXAMIC ACID					
Tab 500 mg	20.67	100	✓	Cyklokapron	
Vitamin K					
PHYTOMENADIONE Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSC		5 5	-	Konakion MM Konakion MM	
Antithrombotic Agents					
Antiplatelet Agents					
ASPIRIN					
* Tab 100 mg		990	✓ <u>E</u>	thics Aspirin I	EC
CLOPIDOGREL					
* Tab 75 mg – For clopidogrel oral liquid formulation rel			ŗ		
page 224	5.44	84	✓ <u>µ</u>	Arrow - Clopid	
DIPYRIDAMOLE	44.50				
* Tab long-acting 150 mg		60	✓ F	ytazen SR	
PRASUGREL - Special Authority see SA1201 below - Re					
Tab 5 mg Tab 10 mg		28 28		Effient Effient	
BS \$ 1201 Special Authority for Subsidy	120.00	20	• •		

⇒SA1201 Special Authority for Subsidy

Initial application — (coronary angioplasty and bare metal stent) from any relevant practitioner. Approvals valid for 6 months where the patient has undergone coronary angioplasty in the previous 4 weeks and is clopidogrel-allergic*.

Initial application — (drug eluting stent) from any relevant practitioner. Approvals valid for 12 months where the patient has had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Initial application — (stent thromobosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal - (coronary angioplasty and bare metal stent) from any relevant practitioner. Approvals valid for 6 months where

continued...

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
continued				
the patient has undergone coronary angioplasty or had a bare me clopidogrel-allergic*.	tal cardiac stent inse	rted i	n the previo	us 4 weeks and is
Renewal — (drug eluting stent) from any relevant practitioner. stent inserted in the previous 4 weeks and is clopidogrel-allergic*.				
Note: * Clopidogrel allergy is defined as a history of anaphylaxis, developing soon after clopidogrel is started and is considered unli				· · · · · · · · · · · · · · · · · · ·
TICAGRELOR – Special Authority see SA1382 below – Retail ph * Tab 90 mg		56	✓ B	rilinta
SA1382 Special Authority for Subsidy		00		innta
Initial application — (acute coronary syndrome) from any rele meeting the following criteria: Both:	vant practitioner. Ap	prova	Ils valid for ⁻	12 months for applications
 Patient has recently (within the last 60 days) been diagnos syndrome; and 	ed with an ST-elevat	ion or	a non-ST-e	elevation acute coronary
2 Fibrinolytic therapy has not been given in the last 24 hours	and is not planned.			
Renewal — (subsequent acute coronary syndrome) from any applications meeting the following criteria: Both:	relevant practitioner.	Арр	rovals valid	for 12 months for

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Heparin and Antagonist Preparations

TERARINA

DALTEPARIN SODIUM - Special Authority see SA1270 below	– Retail pharmacy		
Inj 2,500 iu per 0.2 ml prefilled syringe		10	 Fragmin
Inj 5,000 iu per 0.2 ml prefilled syringe		10	 Fragmin
Inj 7,500 iu per 0.75 ml graduated syringe	60.03	10	 Fragmin
Inj 10,000 iu per 1 ml graduated syringe	77.55	10	 Fragmin
Inj 12,500 iu per 0.5 ml prefilled syringe		10	 Fragmin
Inj 15,000 iu per 0.6 ml prefilled syringe		10	 Fragmin
Inj 18,000 iu per 0.72 ml prefilled syringe	158.47	10	🗸 Fragmin

⇒SA1270 Special Authority for Subsidy

Initial application — (Pregnancy or Malignancy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

- Any of the following:
 - 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
 - 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
 - 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
 - 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
 - 5 To be used in association with cardioversion of atrial fibrillation.

*Three months or six months, as applicable, dispensed all-at-once

continued...

‡ safety cap

Subsidy	F	ully Brand or	
(Manufacturer's	Price) Subsidis	sed Generic	
\$	Per	 Manufacturer 	

continued...

Renewal — (Pregnancy or Malignancy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, Acute Coronary Syndrome, cardioversion, or prior to oral anti-coagulation).

ENOXAPARIN SODIUM - Special Authority see SA1646 below - Retail pharmacy

Inj 20 mg in 0.2 ml syringe	 10	 Clexane
Inj 40 mg in 0.4 ml syringe	 10	 Clexane
Inj 60 mg in 0.6 ml syringe	10	 Clexane
Inj 80 mg in 0.8 ml syringe	10	 Clexane
Inj 100 mg in 1 ml syringe	10	 Clexane
Inj 120 mg in 0.8 ml syringe	10	 Clexane
Inj 150 mg in 1 ml syringe	10	 Clexane

⇒SA1646 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during home haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Any of the following:

- For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant
 - treatment; or
 - 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
 - 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
 - 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
 - 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during home haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

HEPARIN SODIUM

Inj 1,000 iu per ml, 35 ml vial	24.15	1	🖌 Hospira
Inj 1,000 iu per ml, 5 ml		10	 Hospira
	66.80	50	 Hospira
	99.50		 Pfizer
Inj 5,000 iu per ml, 1 ml	28.40	5	🗸 Hospira
Inj 5,000 iu per ml, 5 ml	341.89	50	 Pfizer
Inj 25,000 iu per ml, 0.2 ml	19.00	5	 Hospira

(1	Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	
HEPARINISED SALINE Inj 10 iu per ml, 5 ml	56.94	50	1	Pfizer
Oral Anticoagulants				
DABIGATRAN	70.00	~~		Pradaxa
Cap 75 mg – No more than 2 cap per day Cap 110 mg		60 60		Pradaxa Pradaxa
Cap 150 mg		60		Pradaxa
RIVAROXABAN – Special Authority see SA1066 below – Retail ph	armacy			
Tab 10 mg	153.00	15	1	Xarelto

⇒SA1066 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 weeks for applications meeting the following criteria: Either:

- 1 For the prophylaxis of venous thromboembolism following a total hip replacement; or
- 2 For the prophylaxis of venous thromboembolism following a total knee replacement.

Note: Rivaroxaban is only currently indicated and subsidised for up to 5 weeks therapy for prophylaxis of venous thromboembolism following a total hip replacement and up to 2 weeks therapy for prophylaxis of venous thromboembolism following a total knee replacement.

Renewal from any relevant practitioner. Approvals valid for 5 weeks where prophylaxis for venous thromboembolism is required for patients following a subsequent total hip or knee replacement.

WARFARIN SODIUM

Note: Marevan and Coumadin are not interchangeable.

*	Tab 1 mg	3.46	50	Coumadin
	5	6.86	100	 Marevan
*	Tab 2 mg	4.31	50	 Coumadin
*	Tab 3 mg	9.70	100	 Marevan
	Tab 5 mg		50	 Coumadin
	-	11.75	100	 Marevan

Blood Colony-stimulating Factors

FILGRASTIM - Special Authority see SA1259 below - Retail ph	armacy		
Inj 300 mcg per 0.5 ml prefilled syringe		5	 Zarzio
Inj 480 mcg per 0.5 ml prefilled syringe		5	🗸 Zarzio

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or

*Three months or six months, as applicable, dispensed all-at-once

5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5×10^{9} /L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Neulastim

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

[‡] safety cap

(Manufacturer 1 Hec) Subsidied Certeine \$ Per ✔ Manufacturer	Subsidy (Manufacturer's Price	9	Fully Subsidised	Brand or Generic	
	(Manulaction of Free				

■ SA1384 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*). Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration

GLUCOSE [DEXTROSE]			
* Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO		5	 Biomed
* Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO	14.50	1	 Biomed
POTASSIUM CHLORIDE			
* Inj 75 mg per ml, 10 ml	55.00	50	 AstraZeneca
SODIUM BICARBONATE			
Inj 8.4%, 50 ml	19.95	1	 Biomed
a) Up to 5 inj available on a PSO			
b) Not in combination	00 50	4	✓ Biomed
Inj 8.4%, 100 ml	20.50	1	Biomed
 a) Up to 5 inj available on a PSO b) Not in combination 			
,			
SODIUM CHLORIDE Not funded for use as a nasal drop. Only funded for nebuliser		oniunction with	an antibiotic intended for
nebuliser use.			
Inj 0.9%, bag – Up to 2000 ml available on a PSO		500 ml	✓ Baxter
	1.26	1,000 ml	✓ Baxter
Only if prescribed on a prescription for renal dialysis, mate	rnity or post-na	atal care in the	home of the patient, or on a PSO
for emergency use. (500 ml and 1,000 ml packs)			_
Inj 23.4% (4 mmol/ml), 20 ml ampoule		5	 Biomed
For Sodium chloride oral liquid formulation refer Standard			✓ InterPharma
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO	7.00	50	 Multichem
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO	6.63	50	✓ Pfizer
Inj 0.9%, 20 ml ampoule		20	✓ Multichem
1 · · · · · · · · · · · · · · · · · · ·	7.50	30	 InterPharma
TOTAL PARENTERAL NUTRITION (TPN) - Retail pharmacy-Spe	cialist		
Infusion		1 OP	🗸 TPN
WATER			
1) On a prescription or Practitioner's Supply Order only whe	n on the same	form as an ini	ection listed in the Pharmaceutical
Schedule requiring a solvent or diluent; or		,	
2) On a bulk supply order; or			
3) When used in the extemporaneous compounding of eye			
4) When used for the dilution of sodium chloride soln 7% for	r cystic fibrosis	patients only.	
Inj 5 ml ampoule – Up to 5 inj available on a PSO	7.00	50	 InterPharma
Inj 10 ml ampoule – Up to 5 inj available on a PSO		50 50	✓ <u>InterPharma</u> ✓ Pfizer
Inj 20 ml ampoule – Up to 5 inj available on a PSO		20	✓ Multichem
,	7.50	30	✓ InterPharma

56

	Subsidy (Manufacturer's F \$	Price) Subsi Per	Fully Brand or dised Generic ✓ Manufacturer
Oral Administration			
CALCIUM POLYSTYRENE SULPHONATE Powder		300 g OP	 Calcium Resonium
COMPOUND ELECTROLYTES Powder for oral soln – Up to 10 sach available on a PSO	2.30	10	✓ Enerlyte
DEXTROSE WITH ELECTROLYTES Soln with electrolytes (2 × 500 ml)	6.55	1,000 ml OP	 Pedialyte - Bubblegum
PHOSPHORUS Tab eff 500 mg (16 mmol)		100	✓ Phosphate-Sandoz
POTASSIUM CHLORIDE * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)		60	
* Tab long-acting 600 mg (8 mmol)	(11.85) 7.42	200	Chlorvescent ✓ Span-K
SODIUM BICARBONATE Cap 840 mg	8.52	100	 ✓ Sodibic ✓ Sodibic
SODIUM POLYSTYRENE SULPHONATE Powder		454 g OP	✓ <u>Resonium-A</u>

	Subsidy		Fully	
	(Manufacturer's Price \$	e) Per	Subsidised	Generic Manufacturer
Alpha Adrenoceptor Blockers				
DOXAZOSIN				
* Tab 2 mg	6.75	500	1	Apo-Doxazosin
* Tab 4 mg	9.09	500	1	Apo-Doxazosin
PHENOXYBENZAMINE HYDROCHLORIDE				
* Cap 10 mg	65.00	30	✓	BNM \$29
	216.67	100	1	Dibenzyline S29
PRAZOSIN				
* Tab 1 mg	5.53	100		Apo-Prazosin
* Tab 2 mg	7.00	100		Apo-Prazosin
* Tab 5 mg	11.70	100	~	Apo-Prazosin
TERAZOSIN				
* Tab 1 mg		28		Actavis
* Tab 2 mg		500		Apo-Terazosin
* Tab 5 mg		500	~	Apo-Terazosin
Agents Affecting the Renin-Angiotensin System	n			
Agents Anceting the Henni-Anglotensin bysten				
ACE Inhibitors				
CAPTOPRIL				
*‡ Oral liq 5 mg per ml		95 ml C	DP 🗸	Capoten
Oral liquid restricted to children under 12 years of age.				
CILAZAPRIL				
* Tab 0.5 mg	2.00	90	✓	Zapril
* Tab 2.5 mg	7.20	200		Apo-Cilazapril
* Tab 5 mg		200	1	Apo-Cilazapril
ENALAPRIL MALEATE				
* Tab 5 mg	0.96	100		Ethics Enalapril
* Tab 10 mg		100	~	Ethics Enalapril
* Tab 20 mg – For enalapril maleate oral liquid formulation ref				
page 224	1.78	100	~	Ethics Enalapril
LISINOPRIL			-	
* Tab 5 mg		90		Ethics Lisinopril
* Tab 10 mg		90		Ethics Lisinopril
* Tab 20 mg	2./b	90	~	Ethics Lisinopril
PERINDOPRIL	0.75	00		An a Davis da sull
* Tab 2 mg		30		Apo-Perindopril
* Tab 4 mg	4.80	30	~	Apo-Perindopril
QUINAPRIL	4.04	00		A
* Tab 5 mg		90		Arrow-Quinapril 5
* Tab 10 mg * Tab 20 mg		90 90		Arrow-Quinapril 10 Arrow-Quinapril 20
τ αυ 20 ΠΙΥ		90	•	

	Subsidy		Fully Brand or	r
	(Manufacturer's Price)		Subsidised Generic	
	\$	Per	 Manufac 	cturer
ACE Inhibitors with Diuretics				
CILAZAPRIL WITH HYDROCHLOROTHIAZIDE				
* Tab 5 mg with hydrochlorothiazide 12.5 mg	10.18	100	🗸 Apo-Cilaz	april/
		100		nlorothiazide
			nyuruci	norotmaziue
QUINAPRIL WITH HYDROCHLOROTHIAZIDE				
	0.05	00	(A a a sum of the	10
* Tab 10 mg with hydrochlorothiazide 12.5 mg		30	Accuretic	
* Tab 20 mg with hydrochlorothiazide 12.5 mg	4.78	30	✓ Accuretic	20
Angiotensin II Antagonists				
CANDESARTAN CILEXETIL – Special Authority see SA1223 bel	w – Retail pharma	ער		
			. Condesta	
* Tab 4 mg		90	✓ <u>Candesta</u>	-
* Tab 8 mg		90	 Candesta 	-
* Tab 16 mg	6.12	90	Candesta	<u>r</u>
* Tab 32 mg		90	Candesta	r
SA1223 Special Authority for Subsidy				-
 Patient has persistent ACE inhibitor induced cough that is r inhibitor); or Patient has a history of angioedema. Initial application — (Unsatisfactory response to ACE inhibito further renewal unless notified where patient is not adequately cor 	r) from any relevan	t prac	ctitioner. Approvals va	alid without
LOSARTAN POTASSIUM * Tab 12.5 mg * Tab 25 mg * Tab 50 mg * Tab 100 mg Angiotensin II Antagonists with Diuretics	1.63 2.00	84 84 84 84	✓ Losartan / ✓ Losartan / ✓ Losartan / ✓ Losartan /	Actavis Actavis Actavis
Tab 12.5 mg Tab 25 mg Tab 25 mg Tab 50 mg Tab 50 mg Tab 100 mg Divertion Angiotensin II Antagonists with Divertics LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE	1.63 2.00 2.31	84 84	✓ Losartan ✓ Losartan	Actavis Actavis Actavis
 Tab 12.5 mg Tab 25 mg Tab 25 mg Tab 50 mg Tab 100 mg Angiotensin II Antagonists with Diuretics	1.63 2.00 2.31	84 84	 ✓ Losartan / ✓ Losartan / ✓ Losartan / ✓ Losartan / ✓ Arrow-Los 	Actavis Actavis Actavis Actavis
Tab 12.5 mg Tab 25 mg Tab 25 mg Tab 50 mg Tab 50 mg Tab 100 mg Divertion Angiotensin II Antagonists with Divertics LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE	1.63 2.00 2.31	84 84 84	 ✓ Losartan / ✓ Losartan / ✓ Losartan / ✓ Losartan / ✓ Arrow-Los 	Actavis Actavis Actavis Actavis sartan &
 Tab 12.5 mg Tab 25 mg Tab 50 mg Tab 50 mg Tab 100 mg Angiotensin II Antagonists with Diuretics LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE Tab 50 mg with hydrochlorothiazide 12.5 mg Antiarrhythmics 		84 84 84 30	 ✓ Losartan / ✓ Losartan / ✓ Losartan / ✓ Losartan / ✓ Arrow-Los 	Actavis Actavis Actavis Actavis sartan &
 Tab 12.5 mg Tab 25 mg Tab 50 mg Tab 50 mg Tab 100 mg Angiotensin II Antagonists with Diuretics LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE Tab 50 mg with hydrochlorothiazide 12.5 mg Antiarrhythmics For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaest 		84 84 84 30	 ✓ Losartan / ✓ Losartan / ✓ Losartan / ✓ Losartan / ✓ Arrow-Los 	Actavis Actavis Actavis Actavis sartan &
 Tab 12.5 mg Tab 25 mg Tab 50 mg Tab 50 mg Tab 100 mg Angiotensin II Antagonists with Diuretics LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE Tab 50 mg with hydrochlorothiazide 12.5 mg Antiarrhythmics For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaest AMIODARONE HYDROCHLORIDE 		84 84 84 30	 ✓ Losartan / ✓ Losartan / ✓ Losartan / ✓ Arrow-Los Hydroch 	Actavis Actavis Actavis Actavis sartan & nlorothiazide
 Tab 12.5 mg Tab 25 mg Tab 50 mg Tab 50 mg Tab 100 mg Angiotensin II Antagonists with Diuretics LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE Tab 50 mg with hydrochlorothiazide 12.5 mg Antiarrhythmics For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaest 		84 84 84 30	 ✓ Losartan / ✓ Losartan / ✓ Losartan / ✓ Losartan / ✓ Arrow-Los 	Actavis Actavis Actavis Actavis sartan & nlorothiazide
 Tab 12.5 mg Tab 25 mg Tab 50 mg Tab 50 mg Tab 100 mg Angiotensin II Antagonists with Diuretics LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE Tab 50 mg with hydrochlorothiazide 12.5 mg Antiarrhythmics For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaest AMIODARONE HYDROCHLORIDE 		84 84 84 30	 ✓ Losartan / ✓ Losartan / ✓ Losartan / ✓ Arrow-Los Hydroch 	Actavis Actavis Actavis Actavis sartan & nlorothiazide
 Tab 12.5 mg Tab 25 mg Tab 50 mg Tab 50 mg Tab 100 mg Angiotensin II Antagonists with Diuretics LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE Tab 50 mg with hydrochlorothiazide 12.5 mg Antiarrhythmics For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaest AMIODARONE HYDROCHLORIDE Tab 100 mg – Retail pharmacy-Specialist Tab 200 mg – Retail pharmacy-Specialist 		84 84 84 30 130 30	 Losartan / Losartan / Losartan / Arrow-Los Hydroch 	Actavis Actavis Actavis Actavis sartan & nlorothiazide
 Tab 12.5 mg		84 84 84 30 130 30 30	 Losartan / Losartan / Losartan / Losartan / Arrow-Los Hydroch Cordaronu Cordaronu 	Actavis Actavis Actavis Actavis sartan & nlorothiazide
 Tab 12.5 mg Tab 25 mg Tab 50 mg Tab 50 mg Tab 100 mg Angiotensin II Antagonists with Diuretics LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE Tab 50 mg with hydrochlorothiazide 12.5 mg Antiarrhythmics For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaest AMIODARONE HYDROCHLORIDE Tab 100 mg – Retail pharmacy-Specialist Tab 200 mg – Retail pharmacy-Specialist 		84 84 84 30 130 30 30	 Losartan / Losartan / Losartan / Losartan / Arrow-Los Hydroch Cordaronu Cordaronu 	Actavis Actavis Actavis Actavis sartan & nlorothiazide
 Tab 12.5 mg		84 84 84 30 130 30 30	 Losartan / Losartan / Losartan / Losartan / Arrow-Los Hydroch Cordaronu Cordaronu 	Actavis Actavis Actavis Actavis sartan & nlorothiazide
 Tab 12.5 mg		84 84 84 30 130 30 30	 Losartan / Losartan / Losartan / Losartan / Arrow-Los Hydroch Cordaronu Cordaronu 	Actavis Actavis Actavis Actavis sartan & nlorothiazide

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully	
(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
DIGOXIN				
* Tab 62.5 mcg – Up to 30 tab available on a PSO	6.67	240	✓	Lanoxin PG
* Tab 250 mcg – Up to 30 tab available on a PSO		240	✓	Lanoxin
*‡ Oral liq 50 mcg per ml	16.60	60 m	✓	Lanoxin
			✓	Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE				
▲ Cap 100 mg	23.87	100	1	Rythmodan
FLECAINIDE ACETATE – Retail pharmacy-Specialist				
▲ Tab 50 mg		60	1	Tambocor
▲ Cap long-acting 100 mg		30	1	Tambocor CR
▲ Cap long-acting 200 mg		30	1	Tambocor CR
Inj 10 mg per ml, 15 ml ampoule	52.45	5	✓	Tambocor
MEXILETINE HYDROCHLORIDE				
▲ Cap 150 mg	162.00	100	1	Mexiletine
				Hydrochloride
				USP S29
▲ Cap 250 mg	202.00	100	1	Mexiletine
				Hydrochloride
				USP S29
PROPAFENONE HYDROCHLORIDE – Retail pharmacy-Specialis	t			
▲ Tab 150 mg		50	1	Rytmonorm
<u> </u>				
Antihypotensives				
MIDODRINE – Special Authority see SA1474 below – Retail pharr	201			
Tab 2.5 mg	,	100	1	Gutron
Tab 5 mg		100	-	Gutron
		100	•	Guuon

⇒SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Note: Treatment should be started with small doses and titrated upwards as necessary. Hypertension should be avoided, and the usual target is a standing systolic blood pressure of 90 mm Hg.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta Adrenoceptor Blockers

ATENOLOL			
* Tab 50 mg	4.61	500	 Mylan Atenolol
* Tab 100 mg	7.67	500	 Mylan Atenolol
* Oral lig 25 mg per 5 ml		300 ml OP	 Atenolol AFT
Restricted to children under 12 years of age.			
BISOPROLOL FUMARATE			
* Tab 2.5 mg	3.53	90	 Bosvate
* Tab 5 mg	5.15	90	 Bosvate
* Tab 10 mg	9.40	90	 Bosvate
CARVEDILOL			
* Tab 6.25 mg	2.24	60	 Carvedilol Sandoz
* Tab 12.5 mg		60	 Carvedilol Sandoz
* Tab 25 mg – For carvedilol oral liquid formulation refer,	page 224 2.95	60	 Carvedilol Sandoz

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
ELIPROLOL				
	21.40	180	1	Celol
ABETALOL				
Tab 50 mg	8 99	100	1	Hybloc
Tab 100 mg – For labetalol oral liquid formulation refer,		100		11,5.00
page 224	11.36	100	1	Hybloc
• Tab 200 mg		100		Hybloc
Inj 5 mg per ml, 20 ml ampoule		5		
	(88.60)	Ũ		Trandate
ETOPROLOL SUCCINATE	()			
Tab long-acting 23.75 mg	1.03	30	1	Betaloc CR
Tab long-acting 47.5 mg		30		Betaloc CR
Tab long-acting 95 mg		30	-	Betaloc CR
Tab long-acting 190 mg		30		Betaloc CR
ETOPROLOL TARTRATE				
Tab 50 mg – For metoprolol tartrate oral liquid formulation				
refer, page 224	4 64	100	1	Apo-Metoprolol
• Tab 100 mg		60		Apo-Metoprolol
Tab long-acting 200 mg		28		Slow-Lopresor
Inj 1 mg per ml, 5 ml vial		5		Lopresor
ADOLOL		5	•	Lopicsol
	16.05	100		Apo-Nadolol
 Tab 40 mg Tab 80 mg 		100		Apo-Nadolol
	24.70	100	•	Apo-Nauoloi
	0.70	400		An a Dividadad
Tab 5 mg		100		Apo-Pindolol
Tab 10 mg		100		Apo-Pindolol
• Tab 15 mg	23.46	100	•	Apo-Pindolol
ROPRANOLOL				
F Tab 10 mg	3.65	100		Apo-Propranolol
			~	Apo-Propranolol
				S29 S29
Tab 40 mg	4.65	100	1	Apo-Propranolol
			✓	Apo-Propranolol
				S29 S29
Cap long-acting 160 mg		100	1	Cardinol LA
 Oral liq 4 mg per ml – Special Authority see SA1327 below 				
Retail pharmacy		500 m	nl 🗸	Roxane S29
Apo-Propranolol S29 © Tab 10 mg to be delisted 1 July 2010				

(Apo-Propranolol S29 \$29 Tab 40 mg to be delisted 1 July 2018)

➡SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

continued...

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
	(Manufacturer's Frice) \$	Per		Manufacturer
pntinued				
ither:				
1 For the treatment of a child under 12 years with an haemar	igioma causing func	tional	impairment	: (not for cosmetic rea
only); or				
2 For the treatment of a child under 12 years with cardiac arr	inymias or congenita	al carc	liac aphorm	laitties.
OTALOL				
₭ Tab 80 mg – For sotalol oral liquid formulation refer, page 224		500		lylan
₭ Tab 160 mg		100		<u>lylan</u>
k Inj 10 mg per ml, 4 ml ampoule		5	✓ S	otacor
Sotacor Inj 10 mg per ml, 4 ml ampoule to be delisted 1 August 2	018)			
TIMOLOL				
* Tab 10 mg		100	🗸 🗸	po-Timol
Calcium Channel Blockers				
Dihydropyridine Calcium Channel Blockers				
MLODIPINE				
₭ Tab 2.5 mg		100	🗸 A	po-Amlodipine
₭ Tab 5 mg – For amlodipine oral liquid formulation refer, page		250		po-Amlodipine
₭ Tab 10 mg		250		po-Amlodipine
ELODIPINE			-	
★ Tab long-acting 2.5 mg	1 45	30	V P	lendil ER
 Tab long acting 5 mg. 		30	_	lendil ER
 Tab long-acting 10 mg 		30	_	lendil ER
		00	· -	
SRADIPINE ₭ Cap long-acting 2.5 mg	7.50	20		wasire CDO
		30 30)ynacirc-SRO)ynacirc-SRO
		30	• 1	ynacht-Sho
IIFEDIPINE	40.00	~~		
K Tab long-acting 10 mg		60		dalat 10
			-	defin S29
* Tab long-acting 20 mg		100		lyefax Retard
₭ Tab long-acting 30 mg		30		dalat Oros
 Tab long-acting 60 mg 	5.67	30	✓ <u>A</u>	dalat Oros
Other Calcium Channel Blockers				
ILTIAZEM HYDROCHLORIDE				
₭ Tab 30 mg	4.60	100	🗸 D	lizem
 Tab 60 mg – For diltiazem hydrochloride oral liquid formulation 				
refer, page 224		100	✓ D)ilzem
 Cap long-acting 120 mg 		500		po-Diltiazem CD
V Conlong opting 100 mg	47.07			

500 500

100

✓ Apo-Diltiazem CD

✓ Apo-Diltiazem CD

Pexsig

*

*

PERHEXILINE MALEATE

✓ fully subsidised

[HP4] refer page 4

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
VERAPAMIL HYDROCHLORIDE				
* Tab 40 mg	7.01	100	1	Isoptin
* Tab 80 mg – For verapamil hydrochloride oral liquid				
formulation refer, page 224	11.74	100	✓	Isoptin
* Tab long-acting 120 mg		250	✓	Verpamil SR
* Tab long-acting 240 mg		250	~	Verpamil SR
* Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a				
PSO	25.00	5	1	Isoptin
Centrally-Acting Agents				
CLONIDINE				
* Patch 2.5 mg, 100 mcg per day – Only on a prescription	7.40	4	1	<u>Mylan</u>
* Patch 5 mg, 200 mcg per day – Only on a prescription		4		Mylan
* Patch 7.5 mg, 300 mcg per day – Only on a prescription		4	1	Mylan
CLONIDINE HYDROCHLORIDE				
* Tab 25 mcg		112	1	Clonidine BNM
* Tab 150 mcg		100		Catapres
* Inj 150 mcg per ml, 1 ml ampoule		5	-	Catapres
	····· • • • •	-		
METHYLDOPA * Tab 250 mg	15 10	100		Methyldopa Mylan
Diuretics		100	•	methyldopa mylan
Loop Diuretics				
BUMETANIDE				
* Tab 1 mg		100	✓	Burinex
* Inj 500 mcg per ml, 4 ml vial	7.95	5	✓	Burinex
FUROSEMIDE [FRUSEMIDE]				
* Tab 40 mg – Up to 30 tab available on a PSO	8.00	1,000		Diurin 40
* Tab 500 mg		50		Urex Forte
* + Oral lig 10 mg per ml		0 ml C		Lasix
* Inj 10 mg per ml, 25 ml ampoule		6	1	Lasix
Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a P		5	1	Frusemide-Claris
Potassium Sparing Diuretics				
AMILORIDE HYDROCHLORIDE				
* Tab 5 mg	15.00	100	1	Apo-Amiloride
Oral liq 1 mg per ml		5 ml C	DP 🗸	Biomed
(Apo-Amiloride Tab 5 mg to be delisted 1 January 2019)				
METOLAZONE – Special Authority see SA1678 below – Retail p	harmacy			
Tab 5 mg	,	1	1	Metolazone S29
· · · · · · · · · · · · · · · · · · ·		50		Zaroxolyn S29
		50	•	Laioxoryn
SA1678 Special Authority for Subsidy				

SA1678 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

continued...

‡ safety cap

	Subsidy	,	Fully Brand or
	(Manufacturer's		sidised Generic
	\$	Per	Manufacturer
continued Either:			
 Patient has refractory heart failure and is therapy; or Paediatric patient has oedema secondar 			
SPIRONOLACTONE	,		
* Tab 25 mg		100	✓ Spiractin
 * Tab 100 mg ‡ Oral liq 5 mg per ml 		100 25 ml OP	 ✓ <u>Spiractin</u> ✓ Biomed
		23 III OF	• Diollieu
Potassium Sparing Combination D	liuretics		
AMILORIDE HYDROCHLORIDE WITH FUROS * Tab 5 mg with furosemide 40 mg		28	✓ Frumil
AMILORIDE HYDROCHLORIDE WITH HYDRO			
* Tab 5 mg with hydrochlorothiazide 50 mg	5.00	50	 Moduretic
Thiazide and Related Diuretics			
BENDROFLUMETHIAZIDE [BENDROFLUAZID			
* Tab 2.5 mg – Up to 150 tab available on a	PSO12.50	500	✓ <u>Arrow-</u> <u>Bendrofluazide</u>
May be supplied on a PSO for reasons	other than emergency.		
* Tab 5 mg		500	✓ <u>Arrow-</u> <u>Bendrofluazide</u>
CHLOROTHIAZIDE			
+ Oral liq 50 mg per ml		25 ml OP	 Biomed
CHLORTALIDONE [CHLORTHALIDONE] * Tab 25 mg	8.00	50	✓ Hygroton
INDAPAMIDE		50	• Hygroton
* Tab 2.5 mg		90	✓ Dapa-Tabs
Lipid-Modifying Agents			
Fibrates			
BEZAFIBRATE			
 * Tab 200 mg * Tab long-acting 400 mg 		90 30	 ✓ <u>Bezalip</u> ✓ Bezalip Retard
GEMFIBROZIL		50	
		60	✓ Lipazil
Other Lipid-Modifying Agents			
ACIPIMOX			
* Cap 250 mg		30	 Olbetam
NICOTINIC ACID * Tab 50 mg	<i>I</i> 10	100	 Apo-Nicotinic Acid
* Tab 50 mg		100	✓ <u>Apo-Nicotinic Acid</u>
-			

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic An Anufacturer
Resins			
CHOLESTYRAMINE			
Powder for oral liq 4 g	19.25 (52.68)	50	Questran-Lite
COLESTIPOL HYDROCHLORIDE			
Grans for oral liq 5 g		30	 Colestid
HMG CoA Reductase Inhibitors (Statins)			
Prescribing Guidelines Treatment with HMG CoA Reductase Inhibitors (statins) is reco cardiovascular risk of 15% or greater. ATORVASTATIN – See prescribing guideline above	·		
* Tab 10 mg		500	
✤ Tab 20 mg		500 500	
 ★ Tab 40 mg ★ Tab 80 mg 		500	
PRAVASTATIN – See prescribing guideline above		000	<u>Lorotat</u>
* Tab 20 mg		100	Apo-Pravastatin
* Tab 40 mg		100	
SIMVASTATIN – See prescribing guideline above			
* Tab 10 mg	0.95	90	 Simvastatin Mylan
* Tab 20 mg	1.52	90	 Simvastatin Mylan
* Tab 40 mg	2.63	90	 Simvastatin Mylan
* Tab 80 mg	6.00	90	 Simvastatin Mylan
Selective Cholesterol Absorption Inhibitors			

EZETIMIBE – Special Authority see SA1045 below – Reta	il pharmacy	
* Tab 10 mg		 Ezetimibe Sandoz

► SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and

*Three months or six months, as applicable, dispensed all-at-once

- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 × normal) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be

performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
EZETIMIBE WITH SIMVASTATIN - Special Authority see SA104	4 <mark>6 below –</mark> Retail pha	rmac	;y	
Tab 10 mg with simvastatin 10 mg	5.15	30	✓	Zimybe
Tab 10 mg with simvastatin 20 mg		30	✓	Zimybe
Tab 10 mg with simvastatin 40 mg		30	✓	Zimybe
Tab 10 mg with simvastatin 80 mg		30	✓	Zimybe

⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be

performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

GLYCERYL TRINITRATE		
* Tab 600 mcg – Up to 100 tab available on a PSO8.00	100 OP	 Lycinate
* Oral pump spray, 400 mcg per dose - Up to 250 dose		-
available on a PSO4.45	250 dose OP	 Nitrolingual Pump
	200 0000 01	Spray
* Oral spray, 400 mcg per dose – Up to 250 dose available on a		opiay
PSO	200 dose OP	✓ Glytrin
* Patch 25 mg, 5 mg per day15.73	30	 Nitroderm TTS
* Patch 50 mg, 10 mg per day18.62	30	 Nitroderm TTS
ISOSORBIDE MONONITRATE		
* Tab 20 mg	100	🖌 Ismo 20
* Tab long-acting 40 mg7.50	30	✓ Ismo 40 Retard
* Tab long-acting 60 mg	90	✓ Duride
	30	Dullde
Cumpathemimatica		
Sympathomimetics		
ADRENALINE		
Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PSO	5	Aspen Adrenaline
	5	•
5.25	_	 Hospira
Inj 1 in 10,000, 10 ml ampoule – Up to 5 inj available on a PSO27.00	5	 Hospira
49.00	10	 Aspen Adrenaline
ISOPRENALINE		
* Inj 200 mcg per ml, 1 ml ampoule	25	
(164.20)		Isuprel
(104.20)		loopioi

	Subsidy		Fully Brand or
	(Manufacturer's Price)	Subsi	dised Generic
	\$	Per	Manufacturer
Vasodilators			
AMYL NITRITE			
* Liq 98% in 0.3 ml cap		12	
	(73.40)		Baxter
HYDRALAZINE HYDROCHLORIDE			
* Tab 25 mg – Special Authority see SA1321 below – Retail			• · · · · ·
pharmacy	CBS	1	 Hydralazine
		56	Onelink S29
* Inj 20 mg ampoule	05.00	84 5	✓ AMDIPHARM S29
	25.90	5	Apresoline
■ SA1321 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valie	d without further repo		notified for applications mosting
the following criteria:		Nai uniess	nouned for applications meeting
Either:			
1 For the treatment of refractory hypertension; or			
2 For the treatment of heart failure in combination with a nit inhibitors and/or angiotensin receptor blockers.	rate, in patients who a	ire intolerai	nt or have not responded to ACE
MINOXIDIL			
▲ Tab 10 mg	70.00	100	 Loniten
NICORANDIL			
▲ Tab 10 mg		60	✓ Ikorel
▲ Tab 20 mg		60	✓ Ikorel
PAPAVERINE HYDROCHLORIDE			
* Inj 12 mg per ml, 10 ml ampoule		5	 Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]			
Tab 400 mg		50	 Trental 400
Endothelin Receptor Antagonists			
	nhormooy		
AMBRISENTAN – Special Authority see SA1702 below – Retail Tab 5 mg		30	✓ Volibris
Tab 10 mg	,	30 30	✓ Volibris
► SA1702 Special Authority for Subsidy	1,000.00	00	
Special Authority approved by the Pulmonary Arterial Hypertensi	on Panel		
Notes: Application details may be obtained from PHARMAC's w		rmac.govt.i	nz or:
The Coordinator, PAH Panel		-	
PHARMAC, PO Box 10-254, WELLINGTON			
Tel: (04) 916 7561, Fax: (04) 974 4858, Email: <u>PAH@pharmac</u>	· · · · · · · · · · · · · · · · · · ·		
BOSENTAN – Special Authority see SA1712 on the next page –		50	
Tab 62.5 mg		56 60	 ✓ <u>Mylan-Bosentan</u> ✓ Bosentan-Mylan
Tab 125 mg	401.79 375.00	60 56	 ✓ Bosentan-Mylan ✓ Mylan-Bosentan
	401.79	50 60	✓ Bosentan-Mylan
(Mylan-Bosentan Tab 62.5 mg to be delisted 1 July 2018)			
(Mylan-Bosentan Tab 125 mg to be delisted 1 July 2018)			
- , ,			

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy	Ful	ly Brand o	or
(Man	ufacturer's Price)	Subsidise	d Generio	0
	\$ F	Per •	 Manufa 	cturer

⇒SA1712 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil; or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria: Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:

- 3.1 Bosentan is to be used as PAH triple therapy; and
- 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully Brand or idised Generic ✓ Manufacturer	
Phosphodiesterase Type 5 Inhibitors				
SILDENAFIL – Special Authority see SA1704 below – Retail pha Tab 25 mg Tab 50 mg Tab 100 mg – For sildenafil oral liquid formulation refer, pag	0.75 0.75	4 4 4	✓ <u>Vedafil</u> ✓ <u>Vedafil</u> ✓ <u>Vedafil</u>	
■ SA1704 Special Authority for Subsidy Initial application — (Raynaud's Phenomenon*) from any rele notified for applications meeting the following criteria: All of the following:	evant practitioner. Ap	provals val	lid without further renewal unle	ess
 Patient has Raynaud's Phenomenon*; and Patient has severe digital ischaemia (defined as severe paulceration; digital ulcers; or gangrene); and Patient is following lifestyle management (avoidance of co avoidance of sympathomimetic drugs); and Patient is being treated with calcium channel blockers and 	ld exposure, sufficien	t protectior	n, smoking cessation support,	-
Initial application — (Pulmonary arterial hypertension*) only on the recommendation of a respiratory specialist or cardiologist. applications meeting the following criteria: All of the following:	from a respiratory spe	ecialist, car	rdiologist or medical practition	ıer
 Patient has pulmonary arterial hypertension (PAH)*; and Any of the following: PAH is in Group 1 of the WHO (Venice) clinical cla PAH is in Group 4 of the WHO (Venice) clinical cla PAH is in Group 5 of the WHO (Venice) clinical cla PAH is in Group 5 of the WHO (Venice) clinical cla PAH is in Group 5 of the WHO (Venice) clinical cla Any of the following:	ssifications; or ssifications; and ?) less than or equal to .Pm) > 25 mmHg; or		-	
Prostacyclin Analogues				
 EPOPROSTENOL – Special Authority see SA1696 below – Reta Inj 500 mcg vial Inj 1.5 mg vial ▶SA1696 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertension Notes: Application details may be obtained from PHARMAC's we The Coordinator, PAH Panel PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmacc ILOPROST – Special Authority see SA1705 on the next page – I 		1 1 rmac.govt.r	✓ Veletri ✓ Veletri nz or:	
Nebuliser soln 10 mcg per ml, 2 ml		30	 Ventavis 	

‡ safety cap

▲ Three months supply may be dispensed at one time

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	1	Manufacturer

SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website <u>http://www.pharmac.govt.nz</u> or: The Coordinator, PAH Panel PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

	Subsidy		Fully	Brand or
	(Manufacturer's Price) Sub	sidised	Generic
	\$	Per	1	Manufacturer
Antiacne Preparations				
For systemic antibacterials, refer to INFECTIONS, Antibacterial	s, page 99			
ADAPALENE				
a) Maximum of 30 g per prescription				
b) Only on a prescription				
Crm 0.1%	22.60	30 g OP	. -	Differin
		0	-	Differin
Gel 0.1%		30 g OP	• 0	merin
ISOTRETINOIN – Special Authority see SA1475 below – Retai	l pharmacy			
Cap 10 mg		100	🖌 İs	sotane 10
	14.96	120	✓ 0	Dratane
Cap 20 mg		100	🗸 s	sotane 20
	23.12	120	✓ 0	Dratane

⇒SA1475 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
 - 3.2 Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or

2 Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

TRETINOIN

Crm 0.5 mg per g - Maximum of 50 g per prescription	13.90	50 g OP	 ReTrieve
ReTrieve to be Sole Supply on 1 July 2018			

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 99

HYDROGEN PEROXIDE

Crystaderm

‡ safety cap

15 a OP

DERMATOLOGICALS

(Subsidy Manufacturer's Price		Ful Subsidise	d Generic
(UDID001)	\$	Pe	er •	Manufacturer
	0.00	15 - 1		
Oint 2%		15 g (OP	Bactroban
a) Only on a propagintian	(9.26)			Dactrobari
a) Only on a prescriptionb) Not in combination				
ODIUM FUSIDATE [FUSIDIC ACID]	0.50	45.00	0.0	
Crm 2%	2.52	15 g (UP •	DP Fusidic Acid Cream
a) Maximum of 15 a new properintian				Cream
a) Maximum of 15 g per prescriptionb) Only on a prescription				
c) Not in combination				
Oint 2%	3 45	15 g (OP 🖌	7 Foban
a) Maximum of 15 g per prescription		io g i	01 0	1 Obdil
b) Only on a prescription				
c) Not in combination				
JLFADIAZINE SILVER				
Crm 1%	10.90	50 g (Flamazine
a) Up to 250 g available on a PSO		50 y (UF •	FidilidZille
b) Not in combination				
b) Not in combination				
Antifungals Topical				
and topical				
or systemic antifungals, refer to INFECTIONS, Antifungals, page	106			
MOROLFINE				
a) Only on a prescription				
b) Not in combination				
Nail soln 5%	15.95	5 ml (OP 🖌	MycoNail
ICLOPIROX OLAMINE				
a) Only on a prescription				
b) Not in combination				
Nail-soln 8%	6.50	7 ml (OP 🖌	Apo-Ciclopirox
LOTRIMAZOLE				
Crm 1%	0 70	20 g (OP 🖌	Clomazol
a) Only on a prescription		20 g .	01 -	CICINIZEO
b) Not in combination				
Soln 1%	4.36	20 ml	OP	
	(7.55)		•	Canesten
a) Only on a prescription	(/			
b) Not in combination				
CONAZOLE NITRATE				
Crm 1%	1.00	20 g (OP	
	(7.48)	-09	0.	Pevaryl
a) Only on a prescription	(1.10)			· · · · · · · · ·
b) Not in combination				
Foaming soln 1%, 10 ml sachets	9.89	3		
	(17.23)	5		Pevaryl
a) Only on a prescription	(0)			
b) Not in combination				

	Subsidy		Fully	Brand or
	(Manufacturer's F \$	Price) Subs Per	sidised ✓	Generic Manufacturer
IICONAZOLE NITRATE	*	-		
₭ Crm 2%	0.74	15 g OP	🗸 W	ultichem
a) Only on a prescription		Ū	_	
b) Not in combination				
₭ Lotn 2%	4.36	30 ml OP		
	(10.03)		Da	aktarin
a) Only on a prescription				
b) Not in combination	4.00			
Finct 2%	4.36 (12.10)	30 ml OP	D	aktarin
a) Only on a prescription	(12.10)		D	aniaiiii
b) Not in combination				
IYSTATIN Crm 100,000 u per g	1.00	15 a OP		
	(7.90)	15 g OP	М	ycostatin
a) Only on a prescription	(7.50)		111	,
b) Not in combination				
Antipruritic Preparations				
CALAMINE				
a) Only on a prescriptionb) Not in combination				
Crm, aqueous, BP	1 /0	100 g	🖌 DI	narmacy Health
Lotn, BP		2,000 ml		
CROTAMITON		2,000		
a) Only on a prescription				
b) Not in combination				
Crm 10%	3.37	20 g OP	🖌 ita	ch-Soothe
IENTHOL – Only in combination		_0 g 0.	<u></u>	
-	or propriotory Topical C	Corticoptoriod	Diain r	for dormatalagical ba
 Only in combination with a dermatological base page 223 	or proprietary ropicar c	Jonicosteniou -	rialli, it	eler dermatological bas
 With or without other dermatological galenicals. 				
Crystals	6.50	25 g	✓ P\$	бм
- ,	6.92	3		idWest
	29.60	100 g		idWest
PSM Crystals to be delisted 1 November 2018)		-		
Corticosteroids Topical				
or systemic corticosteroids, refer to CORTICOSTEROID	S AND RELATED AGE	NTS, page 89		
Corticosteroids - Plain				
ETAMETHASONE DIPROPIONATE				
Crm 0.05%		15 g OP		prosone
	8.97	50 g OP		prosone
0 0 0 0 0 0 0 0 0 0	4.33	30 g OP		prosone OV
Crm 0.05% in propylene glycol base			.	prosone
Crm 0.05% in propylene glycol base Oint 0.05%	2.96	15 g OP		
	2.96 8.97	15 g OP 50 g OP 30 g OP	🗸 Di	prosone prosone OV

*Three months or six months, as applicable, dispensed all-at-once

if endorsed "certified exemption" by the prescriber or pharmacist.

s Per Manufacturer ETAMETHASONE VALERATE 3.15 50 g OP ✓ Beta Cream © Ont 0.1% 3.15 50 g OP ✓ Beta Cream © Ont 0.1% 3.15 50 g OP ✓ Beta Ointment © Ont 0.1% 10.05 50 m OP ✓ Beta Ointment © Ont 0.05% 2.20 30 g OP ✓ Dermol © Ont 0.05% 2.20 30 g OP ✓ Dermol © Ont 0.05% 2.20 30 g OP ✓ Dermol © Ont 0.05% 5.38 30 g OP ✓ Dermol UDBETASONE BUTYRATE (7.09) Eurovate IFLUCORTOLONE VALERATE (15.86) Nerisone Crm 0.1% (15.86) Nerisone YDPCOCRTISONE (15.86) Nerisone * Owder – Only in combination 11 30 g OP ✓ DermAssist * Powder – Only in combination 49.95 25 g ✓ ABM Up to 5% in a dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatologica galenicals. Refer, page 223 YDPCOCRTISONE BUTYRATE UporcORTISONE BUTYRATE<		Subsidy		Fully	
ETAMETHASONE VALERATE Cm 0.1% Chit 0.5% Chit 0.1% <				sidised V	
: Cm 0.1% 3.15 50 g OP ✓ Beta Cream : Oint 0.1% 3.15 50 g OP ✓ Beta Ontmatt : Lotin 0.1% 10.05 50 ml OP ✓ Beta Ontmatt : Lotin 0.1% 10.05 50 ml OP ✓ Beta Ontmatt : LOB 0.5% 2.20 30 g OP ✓ Dermol LOBETASONE BUTYPATE 7.09 Eumovate IUCORTOLONE VALERATE 7.09 Eumovate IFLUCORTOLONE VALERATE 8.97 50 g OP Cm 0.1% .8.97 50 g OP (15.86) Nerisone (15.86) YDROCORTISONE (15.86) Nerisone : Cm 1% - Only on a prescription 1.11 30 g OP ✓ DermAssist : Powder - Only in combination 49.95 25 g ABM Up to 5% in a dematological base (not proprietary Topical Corticosteriod - Plain) with or without other dematological galenicals. Refer, page 223 YDROCORTISONE & Locoid Lipocream : Powder - Only in combination 10.57 250 ml ✓ DP Lotn HC YDROCORTISONE BUTYRATE 2.30 30 g OP ✓ Locoid Lipocream		÷			manadataron
€ Ont 0.1% 3.15 50 g OP ✓ Beta Ointment € Lon 0.1% 10.05 50 m OP ✓ Beta Ointment € Crm 0.05% 2.20 30 g OP ✓ Dermol € Orn 0.05% 2.20 30 g OP ✓ Dermol LOBETASONE BUTYRATE Crm 0.05% 5.38 30 g OP ✓ Dermol LOBETASONE BUTYRATE (7.09) Eurovate Eurovate IFLUCORTOLONE VALERATE (7.09) Eurovate Crm 0.1% (15.86) Nerisone YDPOCORTISONE (15.86) Nerisone VDP OS% in a dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological galenicals. Refer, page 223 S0 g OP ✓ YDPOCORTISONE AND PARAFFIN LIQUID AND LANOLIN Lon 1% with parafin liquid 15.9% and lanolin 0.6% – Only on a prescription 10.57 250 ml ✓ DP Lotn HC YDPOCORTISONE BUTYRATE 2.30 30 g OP ✓ Locoid Lipocream Up to 5% in a dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological galenicals. Refer, page 223 S0 g OP ✓ </td <td></td> <td>2 15</td> <td></td> <td>1</td> <td>Bota Croam</td>		2 15		1	Bota Croam
			•		
LOBETASOL PROPIONATE 2.0 30 g OP ✓ Dermol ↓ Com 0.05% 2.20 30 g OP ✓ Dermol ↓ Com 0.05% 2.20 30 g OP ✓ Dermol ↓ Com 0.05% 5.38 30 g OP ✓ Dermol ↓ CDBETASONE BUTYRATE (7.09) Eurnovate ↓ CTUCORTOLONE VALERATE (7.09) Eurnovate ↓ CTUCORTOLONE VALERATE (15.86) Nerisone ↓ CTM \> Only on a prescription 1.11 30 g OP ✓ DermAssist ↓ Powder - Only in combination 49.95 25 g ✓ ABM ↓ Up to 5% in a dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatologica galericals. Refer, page 223 100 g OP ✓ Locoid Lipocream VDROCORTISONE BUTYRATE 2.30 30 g OP ✓ Locoid Lipocream Lipocream 0.1% 6.85 100 g OP ✓ Locoid Lipocream Oint 0.1% 6.85 100 g OP ✓ Locoid Lipocream Oint 0.1% 6.85 100 g OP ✓ Locoid Lipocream Oint 0.1% 6.85 100 g OP ✓ Locoid Lipocream Oint 0.1% 6.85 100 g OP ✓ Advantan			0		
 Cm 0.05%		10.05	50 mi OP	•	Betnovate
	CLOBETASOL PROPIONATE				
CODETASONE BUTYRATE 5.38 30 g OP (7.09) Eumovate VIFLUCORTOLONE VALERATE (15.86) Nerisone Crm 0.1% (15.86) Nerisone Fatty oint 0.1% (15.86) Nerisone (15.86) Nerisone (15.86) Nerisone VDROCORTISONE (15.86) Nerisone Nerisone * Ornly on a prescription 49.95 25 g ABM Up to 5% in a dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological base (not proprietary Topical Corticosteroly – Cocoid Lipocream – Locoid Clipocream –	₭ Crm 0.05%	2.20	30 g OP	-	Dermol
Crm 0.05% 5.38 30 g OP (7.09) Eumovate IFLUCORTOLONE VALERATE (7.09) Nerisone Crm 0.1% (15.86) Nerisone Fatty oint 0.1% (15.86) Nerisone YDROCORTISONE (15.86) Nerisone * Crm 1% - Only on a prescription 1.11 30 g OP ✓ DermAssist 16.25 500 g ✓ Pharmacy Health yp to 5% in a dematological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological galenicals. Refer, page 223 YDROCORTISONE AND PARAFTIN LIQUID AND LANOLIN Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only on a prescription 10.57 250 ml ✓ DP Lotn HC YDROCORTISONE BUTYRATE Lipcoream 0.1% 6.85 100 g OP ✓ Locoid Lipcoream Uint 0.1% 6.85 100 g OP ✓ Locoid Crelo Netremul 0.1% 4.95 15 g OP ✓ Advantan Oint 0.1% 1.51 15 g OP ✓ Advantan Oint 0.1% 1.51 15 g OP ✓ Elecon Alcohol Free Crm 0.1% 2.90 50 g OP ✓ Elecon Oint 0.1% 7.33 30 ml OP </td <td>← Oint 0.05%</td> <td>2.20</td> <td>30 g OP</td> <td>-</td> <td>Dermol</td>	← Oint 0.05%	2.20	30 g OP	-	Dermol
Crm 0.05% 5.38 30 g OP (7.09) Eumovate IFLUCORTOLONE VALERATE (7.09) Nerisone Crm 0.1% (15.86) Nerisone Fatty oint 0.1% (15.86) Nerisone VDROCORTISONE (15.86) Nerisone * Crm 1% - Only on a prescription 1.11 30 g OP ✓ DermAssist 16.25 500 g ✓ Pharmacy Health 49.95 25 g ✓ ABM Up to 5% in a dematological base (not proprietary Topical Corticosteriod – Plain) with or without other dematological galenicals. Refer, page 223 VDROCORTISONE AND PARAFTIN LIQUID AND LANOLIN Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only on a prescription 10.57 250 ml ✓ DP Lotn HC IVDROCORTISONE BUTYRATE Lipcoream 0.1% 6.85 100 g OP ✓ Locoid Lipcoream Uint 0.1% 6.85 100 g OP ✓ Locoid Crelo Crm 0.1% ✓ DP Lotn HC VDROCORTISONE ENDERDETARTE 2.30 30 g OP ✓ Locoid Lipcoream ✓ DP Lotn HC VDROCORTISONE BUTYRATE 10.57 250 ml ✓ DP Lotn HC ✓ DO Col Crelo NETHYLPREDNISOLONE ACEPONATE 6.85 100 g OP	OBETASONE BUTYBATE				
(7.09) Eumovate VIFLUCORTOLONE VALERATE 8.97 50 g OP Crm 0.1%		5.38	30 a OP		
IFLUCORTOLONE VALERATE 8.97 50 g OP Fatty oint 0.1%	0111 0.00 /0		00 9 01		Fumovate
Crm 0.1% 8.97 50 g OP Fatty oint 0.1% (15.86) Nerisone YDROCORTISONE (15.86) Nerisone * Crm 1% - Only on a prescription 1.11 30 g OP ✓ DermAssist * Crm 1% - Only in combination 49.95 25 g ✓ ABM Up to 5% in a dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological galenicals. Refer, page 223 YDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only on a prescription 10.57 250 ml ✓ DP Lotn HC YDROCORTISONE BUTYRATE 2.30 30 g OP ✓ Locoid Lipocream Lipocream 0.1% 6.85 100 g OP ✓ Locoid Lipocream Oint 0.1% 6.85 100 g OP ✓ Locoid Crelo IETHYLPREDNISOLONE ACEPONATE Crm 0.1% 4.95 15 g OP ✓ Advantan Oint 0.1% 2.90 50 g OP ✓ Elocon Alcohol Free 0 g OP ✓ Elocon Oint 0.1% 2.90 50 g OP ✓ Elocon Alcohol Free Crm 0.1% 2.90 50 g OP ✓ Elocon ICHTYLPREDNISOLONE ACEPONATE 1.51 15 g OP ✓ Advantan Oint 0.1% <td></td> <td>(7.00)</td> <td></td> <td></td> <td>Luniovale</td>		(7.00)			Luniovale
(15.86) Nerisone Fatty oint 0.1%					
Fatty oint 0.1% 8.97 50 g OP (15.86) Nerisone YDROCORTISONE (15.86) Verisone © Crm 1% - Only on a prescription 1.11 30 g OP ✓ DermAssist 16.25 500 g ✓ Pharmacy Health 16.25 500 g ✓ Pharmacy Health 16.25 500 g ✓ ABM Up to 5% in a dermatological base (not proprietary Topical Corticosteriod – Plain) with out other dermatological galenicals. Refer, page 223 YDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only on a prescription 10.57 250 ml ✓ DP Lotn HC YDROCORTISONE BUTYRATE Lipocream 0.1% 6.85 100 g OP ✓ Locoid Lipocream Milky emul 0.1% 6.85 100 g OP ✓ Locoid Crelo IETHYLPREDNISOLONE ACEPONATE Crm 0.1% 4.95 15 g OP ✓ Advantan Oint 0.1% 4.95 15 g OP ✓ Advantan IOMETASONE FUROATE Crm 0.1% 2.90 50 g OP ✓ Elocon Alcohol Free Oint 0.1% 2.90 50 g OP ✓ Elocon IOMETASONE FUROATE 7.35 30 ml OP	Crm 0.1%		50 g OP		
(15.86) Nerisone YDROCORTISONE 1.1.1 30 g OP DermAssist Crm 1% - Only on a prescription 16.25 500 g YParmacy Health 49.95 25 g XBM Yearmacy Health 10.1% Advantan 10.57 250 ml YDROCORTISONE BUTYRATE Lipocream 0.1% 2.30 30 g OP < Locoid Lipocream					Nerisone
YDROCORTISONE 1.11 30 g OP ✓ DermAssist 6 Crm 1% - Only on a prescription 16.25 500 g ✓ Pharmacy Health 49.95 25 g ✓ ABM Up to 5% in a dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological galenicals. Refer, page 223 YDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only on a prescription 10.57 250 ml ✓ DP Lotn HC YDROCORTISONE BUTYRATE Lipocream 0.1% 6.85 100 g OP ✓ Locoid Lipocream Oint 0.1% 6.85 100 g OP ✓ Locoid Crelo IETHYLPREDNISOLONE ACEPONATE 4.95 15 g OP ✓ Advantan Oint 0.1% 1.51 15 g OP ✓ Advantan IOMETASONE FUROATE 2.90 50 g OP ✓ Elocon Alcohol Free Crm 0.1% 1.51 15 g OP ✓ Elocon Oint 0.1% 6.35 100 g OP ✓ Elocon IOMETASONE FUROATE 2.90 50 g OP ✓ Elocon Crm 0.1% 6.35 100 g OP ✓ Elocon Iotn 0.1% 6.30 100 g OP ✓ Aristocor	Fatty oint 0.1%		50 g OP		
* Crm 1% - Only on a prescription 1.11 30 g OP ✓ DermAssist 16.25 500 g ✓ Pharmacy Health 49.95 25 g ✓ ABM Up to 5% in a dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological galenicals. Refer, page 223 IVDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only on a prescription 10.57 250 ml ✓ DP Lotn HC IVDROCORTISONE BUTYRATE 2.30 30 g OP ✓ Locoid Lipocream 0int 0.1% 6.85 100 g OP ✓ Locoid Lipocream 0int 0.1% 6.85 100 g OP ✓ Locoid Crelo NETHYPREDNISOLONE ACEPONATE 4.95 15 g OP ✓ Advantan Oint 0.1% 1.51 15 g OP ✓ Advantan NOMETASONE FUROATE 2.90 50 g OP ✓ Elocon Alcohol Free 0int 0.1% 2.90 50 g OP ✓ Elocon 0int 0.1% 7.35 30 ml OP ✓ Elocon 100 ml 0.1% 6.35 100 g OP ✓ Elocon 100 ml 0.1% 6.35 100 g OP ✓ Elocon 100 ml 0.1% 6.30 100 g OP ✓		(15.86)			Nerisone
16.25 500 g ✓ Pharmacy Health 49.95 25 g ✓ ABM Up to 5% in a dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatologica galenicals. Refer, page 223 YDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only on a prescription 10.57 250 ml ✓ DP Lotn HC YDROCORTISONE BUTYRATE Lipocream 0.1% 6.85 100 g OP ✓ Locoid Lipocream Oint 0.1% 6.85 100 g OP ✓ Locoid Lipocream Oint 0.1% 6.85 100 g OP ✓ Locoid Crelo NETHYLPREDNISOLONE ACEPONATE Crm 0.1% 4.95 15 g OP ✓ Advantan Oint 0.1% 4.95 15 g OP ✓ Advantan 0int 0.1% Elocon Alcohol Free Crm 0.1% 4.95 15 g OP ✓ Elocon Alcohol Free 2.90 50 g OP ✓ Elocon Oint 0.1% 1.51 15 g OP ✓ Elocon Alcohol Free 2.90 50 g OP ✓ Elocon Cond 0.1% 2.90 50 g OP ✓ Elocon Elocon Elocon Elocon IOMETASONE FUROATE Cond 0.0 g OP ✓ Advantan	YDROCORTISONE				
16.25 500 g ✓ Pharmacy Health 49.95 25 g ✓ ABM Up to 5% in a dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatologica galenicals. Refer, page 223 IYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only on a prescription 10.57 250 ml ✓ DP Lotn HC IYDROCORTISONE BUTYRATE Lipocream 0.1% 6.85 100 g OP ✓ Locoid Lipocream Oint 0.1% 6.85 100 g OP ✓ Locoid Lipocream Oint 0.1% 6.85 100 g OP ✓ Locoid Crelo NETHYLPREDNISOLONE ACEPONATE Crm 0.1% 4.95 15 g OP ✓ Advantan Oint 0.1% 4.95 15 g OP ✓ Advantan Oint 0.1% 4.95 15 g OP ✓ Elocon Alcohol Free Crm 0.1% 2.90 50 g OP ✓ Elocon Oint 0.1% 2.90 50 g OP ✓ Elocon Icon 1.1% 7.35 30 ml OP ✓ Elocon RIAMCINOLONE ACETONIDE 6.30 100 g OP ✓ Aristocort Oint 0.1% 6.35 100 g OP ✓ Aristocort	Crm 1% – Only on a prescription		30 a OP	1	DermAssist
 Powder - Only in combination			0		
Up to 5% in a dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological galenicals. Refer, page 223 YDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN Loth 1% with paraffin liquid 15.9% and lanolin 0.6% – Only on a prescription	Powder – Only in combination				
a prescription 10.57 250 ml ✓ DP Lotn HC IYDROCORTISONE BUTYRATE 30 g OP ✓ Locoid Lipocream Lipocream 0.1% 6.85 100 g OP ✓ Locoid Lipocream Oint 0.1% 6.85 100 g OP ✓ Locoid Lipocream Milky emul 0.1% 6.85 100 g OP ✓ Locoid Lipocream Milky emul 0.1% 6.85 100 ml OP ✓ Locoid Crelo NETHYLPREDNISOLONE ACEPONATE 15 g OP ✓ Advantan Oint 0.1% 4.95 15 g OP ✓ Advantan NOMETASONE FUROATE 1.51 15 g OP ✓ Elocon Alcohol Free Oint 0.1% 2.90 50 g OP ✓ Elocon Oint 0.1% 2.90 50 g OP ✓ Elocon Lotn 0.1% 7.35 30 ml OP ✓ Elocon RIAMCINOLONE ACETONIDE 6.35 100 g OP ✓ Aristocort Oint 0.02% 6.35 100 g OP ✓ Aristocort Oint 0.02% Combination 3.49 15 g OP	YDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN	n			
Lipocream 0.1%			250 ml	1	DP Lotn HC
Lipocream 0.1%					
6.85 100 g OP ✓ Locoid Lipocream Oint 0.1% 6.85 100 g OP ✓ Locoid Crelo Milky emul 0.1% 6.85 100 ml OP ✓ Locoid Crelo METHYLPREDNISOLONE ACEPONATE 4.95 15 g OP ✓ Advantan Oint 0.1% 4.95 15 g OP ✓ Advantan OINETASONE FUROATE 1.51 15 g OP ✓ Advantan NOMETASONE FUROATE 2.90 50 g OP ✓ Elocon Alcohol Free Oint 0.1% 1.51 15 g OP ✓ Elocon Alcohol Free Oint 0.1% 7.35 30 ml OP ✓ Elocon Lotn 0.1% 7.35 30 ml OP ✓ Elocon RIAMCINOLONE ACETONIDE 6.30 100 g OP ✓ Aristocort Oint 0.02% 6.35 100 g OP ✓ Aristocort Oint 0.02% Combination 3.49 15 g OP		2 30	30 a OP	1	Locoid Linocream
Oint 0.1% 6.85 100 g OP ✓ Locoid Milky emul 0.1% 6.85 100 ml OP ✓ Locoid Crelo NETHYLPREDNISOLONE ACEPONATE 4.95 15 g OP ✓ Advantan Oint 0.1% 4.95 15 g OP ✓ Advantan NOMETASONE FUROATE 4.95 15 g OP ✓ Advantan NOMETASONE FUROATE 1.51 15 g OP ✓ Elocon Alcohol Free Oint 0.1% 1.51 15 g OP ✓ Elocon Alcohol Free Oint 0.1% 1.51 15 g OP ✓ Elocon Alcohol Free Oint 0.1% 1.51 15 g OP ✓ Elocon Alcohol Free Oint 0.1% 7.35 30 ml OP ✓ Elocon Lotn 0.1% 7.35 30 ml OP ✓ Elocon RIAMCINOLONE ACETONIDE 6.30 100 g OP ✓ Aristocort Oint 0.02% 6.35 100 g OP ✓ Aristocort Oint 0.02% Combination 3.49 15 g OP			•		
Milky emul 0.1% 6.85 100 ml OP ✓ Locoid Crelo NETHYLPREDNISOLONE ACEPONATE 4.95 15 g OP ✓ Advantan Oint 0.1% 4.95 15 g OP ✓ Advantan NOMETASONE FUROATE 1.51 15 g OP ✓ Advantan NOMETASONE FUROATE 1.51 15 g OP ✓ Elocon Alcohol Free Oint 0.1% 1.51 15 g OP ✓ Elocon Alcohol Free Oint 0.1% 1.51 15 g OP ✓ Elocon Alcohol Free Oint 0.1% 1.51 15 g OP ✓ Elocon Lotn 0.1% 7.35 30 ml OP ✓ Elocon RIAMCINOLONE ACETONIDE 6.30 100 g OP ✓ Aristocort Oint 0.02% 6.35 100 g OP ✓ Aristocort Oint 0.02% Combination 100 g OP ✓ Aristocort VETAMETHASONE VALERATE WITH CLIOQUINOL – Only on a prescription Crm 0.1% with clioquinol 3% 3.49 15 g OP	Oint 0.1%		•		
NETHYLPREDNISOLONE ACEPONATE Crm 0.1% 4.95 15 g OP ✓ Advantan Oint 0.1% 4.95 15 g OP ✓ Advantan NOMETASONE FUROATE 1.51 15 g OP ✓ Advantan NOMETASONE FUROATE 2.90 50 g OP ✓ Elocon Alcohol Free Oint 0.1% 1.51 15 g OP ✓ Elocon Alcohol Free Oint 0.1% 1.51 15 g OP ✓ Elocon Alcohol Free Oint 0.1% 1.51 15 g OP ✓ Elocon Lotn 0.1% 7.35 30 ml OP ✓ Elocon RIAMCINOLONE ACETONIDE 6.30 100 g OP ✓ Aristocort Oint 0.02% 6.35 100 g OP ✓ Aristocort Oint 0.02% Combination ✓ Aristocort ✓ Aristocort VETAMETHASONE VALERATE WITH CLIOQUINOL – Only on a prescription Crm 0.1% with clioquinol 3% 3.49 15 g OP					
Crm 0.1% 4.95 15 g OP ✓ Advantan Oint 0.1% 4.95 15 g OP ✓ Advantan IOMETASONE FUROATE 1.51 15 g OP ✓ Elocon Alcohol Free Crm 0.1% 1.51 15 g OP ✓ Elocon Alcohol Free Oint 0.1% 2.90 50 g OP ✓ Elocon Alcohol Free Oint 0.1% 1.51 15 g OP ✓ Elocon Alcohol Free Oint 0.1% 2.90 50 g OP ✓ Elocon Lotn 0.1% 7.35 30 ml OP ✓ Elocon RIAMCINOLONE ACETONIDE 6.30 100 g OP ✓ Aristocort Oint 0.02% 6.35 100 g OP ✓ Aristocort Corticosteroids - Combination ETAMETHASONE VALERATE WITH CLIOQUINOL – Only on a prescription Crm 0.1% with clioquinol 3% 3.49 15 g OP		0.05		•	
Oint 0.1%					
IOMETASONE FUROATE 1.51 15 g OP ✓ Elocon Alcohol Free Crm 0.1% 2.90 50 g OP ✓ Elocon Alcohol Free Oint 0.1% 1.51 15 g OP ✓ Elocon Alcohol Free Lotn 0.1% 2.90 50 g OP ✓ Elocon Lotn 0.1% 7.35 30 ml OP ✓ Elocon RIAMCINOLONE ACETONIDE 6.30 100 g OP ✓ Aristocort Oint 0.02% 6.35 100 g OP ✓ Aristocort Corticosteroids - Combination 100 g OP ✓ Aristocort VETAMETHASONE VALERATE WITH CLIOQUINOL – Only on a prescription Crm 0.1% with clioquinol 3% 3.49 15 g OP			0		
Crm 0.1%	Oint 0.1%	4.95	15 g OP	-	Advantan
2.90 50 g OP ✓ Elocon Alcohol Free Oint 0.1% 1.51 15 g OP ✓ Elocon 2.90 50 g OP ✓ Elocon ✓ Elocon 2.90 50 g OP ✓ Elocon ✓ Elocon Lotn 0.1% .7.35 30 ml OP ✓ Elocon RIAMCINOLONE ACETONIDE .6.30 100 g OP ✓ Aristocort Oint 0.02% .6.35 100 g OP ✓ Aristocort Oint 0.02% .6.35 100 g OP ✓ Aristocort Corticosteroids - Combination	IOMETASONE FUROATE				
2.90 50 g OP ✓ Elocon Alcohol Free Oint 0.1% 1.51 15 g OP ✓ Elocon 2.90 50 g OP ✓ Elocon ✓ Elocon 2.90 50 g OP ✓ Elocon ✓ Elocon Lotn 0.1% .7.35 30 ml OP ✓ Elocon RIAMCINOLONE ACETONIDE .6.30 100 g OP ✓ Aristocort Oint 0.02% .6.35 100 g OP ✓ Aristocort Corticosteroids - Combination	Crm 0.1%	1.51	15 g OP	-	Elocon Alcohol Free
Oint 0.1% 1.51 15 g OP ✓ Elocon 2.90 50 g OP ✓ Elocon Atristocort 30 ml OP ✓ Elocon RIAMCINOLONE ACETONIDE 6.30 100 g OP ✓ Aristocort Oint 0.02% 6.35 100 g OP ✓ Aristocort Oint 0.02% 6.35 100 g OP ✓ Aristocort Corticosteroids - Combination 100 g OP ✓ Aristocort ETAMETHASONE VALERATE WITH CLIOQUINOL – Only on a prescription Crm 0.1% with clioquinol 3% 3.49 15 g OP			•		
2.90 50 g OP ✓ Elocon Lotn 0.1% 7.35 30 ml OP ✓ Elocon RIAMCINOLONE ACETONIDE 6.30 100 g OP ✓ Aristocort Oint 0.02% 6.35 100 g OP ✓ Aristocort Corticosteroids - Combination 6.35 100 g OP ✓ Aristocort ETAMETHASONE VALERATE WITH CLIOQUINOL – Only on a prescription Crm 0.1% with clioquinol 3% 3.49 15 g OP	Oint 0.1%		•		
Lotn 0.1%			•		
RIAMCINOLONE ACETONIDE 6.30 100 g OP Aristocort Crm 0.02% 6.35 100 g OP Aristocort Corticosteroids - Combination 6.35 100 g OP Aristocort ETAMETHASONE VALERATE WITH CLIOQUINOL – Only on a prescription Crm 0.1% with clioquinol 3% 15 g OP	Lotn 0.1%			-	
Crm 0.02% 6.30 100 g OP ✓ Aristocort Oint 0.02% 6.35 100 g OP ✓ Aristocort Corticosteroids - Combination ETAMETHASONE VALERATE WITH CLIOQUINOL – Only on a prescription Crm 0.1% with clioquinol 3% 15 g OP				-	
Oint 0.02%6.35 100 g OP ✓ Aristocort Corticosteroids - Combination ETAMETHASONE VALERATE WITH CLIOQUINOL – Only on a prescription Crm 0.1% with clioquinol 3%					
Corticosteroids - Combination ETAMETHASONE VALERATE WITH CLIOQUINOL – Only on a prescription Crm 0.1% with clioquinol 3%		0.00	100 - 00		A.J
ETAMETHASONE VALERATE WITH CLIOQUINOL – Only on a prescription Crm 0.1% with clioquinol 3%	Crm 0.02%		•		
Crm 0.1% with clioquinol 3%	Crm 0.02%		•		
Crm 0.1% with clioquinol 3%	Crm 0.02% Oint 0.02%		•		
	Crm 0.02% Oint 0.02% Corticosteroids - Combination	6.35	•		
	Crm 0.02% Oint 0.02% Corticosteroids - Combination BETAMETHASONE VALERATE WITH CLIOQUINOL – Only on a	a prescription	100 g OP		
	Crm 0.02% Oint 0.02% Corticosteroids - Combination ETAMETHASONE VALERATE WITH CLIOQUINOL – Only on a	a prescription	100 g OP		

			Fully Brand or
	Subsidy (Manufacturer's F		idised Generic
	\$	Per	 Manufacturer
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP	Fucicort
a) Maximum of 15 g per prescriptionb) Only on a prescription			
HYDROCORTISONE WITH MICONAZOLE - Only on a pres ← Crm 1% with miconazole nitrate 2%		15 g OP	✓ Micreme H
IYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN Crm 1% with natamycin 1% and neomycin sulphate 0.5% Oint 1% with natamycin 1% and neomycin sulphate 0.5%	62.79 62.79	15 g OP 15 g OP	✓ Pimafucort✓ Pimafucort
RIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOM Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2. and gramicidin 250 mcg per g – Only on a prescript	5 mg	15 g OP	Viaderm KC
Disinfecting and Cleansing Agents			
CHLORHEXIDINE GLUCONATE – Subsidy by endorsemen a) No more than 500 ml per month b) Only if prescribed for a dialysis patient and the prescri		cordingly	
 Handrub 1% with ethanol 70% 		500 ml	✓ <u>healthE</u> ✓ healthE
₭ Soln 4% wash	3.98	500 ml	
 Soln 4% wash RICLOSAN – Subsidy by endorsement Maximum of 500 ml per prescription b) 	3.98	500 mi	• <u>nearne</u>
RICLOSAN – Subsidy by endorsement a) Maximum of 500 ml per prescription	ethicillin-resistant Sta sed accordingly; or	aphylococcus a	ureus (MRSA) prior to electiv
 RICLOSAN – Subsidy by endorsement Maximum of 500 ml per prescription a) Only if prescribed for a patient identified with Mesurgery in hospital and the prescription is endor 	ethicillin-resistant Sta sed accordingly; or aphylococcus aureu	aphylococcus a	ureus (MRSA) prior to electiv
 RICLOSAN – Subsidy by endorsement Maximum of 500 ml per prescription Maximum of 500 ml per prescription Only if prescribed for a patient identified with Musurgery in hospital and the prescription is endor Only if prescribed for a patient with recurrent St accordingly Soln 1% 	ethicillin-resistant Sta sed accordingly; or aphylococcus aureu	aphylococcus a	ureus (MRSA) prior to electiv
 RICLOSAN – Subsidy by endorsement Maximum of 500 ml per prescription a) Only if prescribed for a patient identified with Ms surgery in hospital and the prescription is endor b) Only if prescribed for a patient with recurrent St accordingly Soln 1% 	ethicillin-resistant Sta sed accordingly; or aphylococcus aureu	aphylococcus a	ureus (MRSA) prior to electiv
RICLOSAN – Subsidy by endorsement a) Maximum of 500 ml per prescription b) a) Only if prescribed for a patient identified with Musurgery in hospital and the prescription is endor b) Only if prescribed for a patient with recurrent St accordingly Soln 1% Barrier Creams and Emollients BMETHICONE	ethicillin-resistant Sta sed accordingly; or aphylococcus aureu 5.90	aphylococcus a	ureus (MRSA) prior to elective the prescription is endorsed
 RICLOSAN – Subsidy by endorsement a) Maximum of 500 ml per prescription b) 	ethicillin-resistant Sta sed accordingly; or aphylococcus aureu 	aphylococcus a s infection and 500 ml OP	ureus (MRSA) prior to electivent the prescription is endorsed
 RICLOSAN – Subsidy by endorsement a) Maximum of 500 ml per prescription b) 	ethicillin-resistant Sta sed accordingly; or aphylococcus aureu 	aphylococcus a s infection and 500 ml OP 500 ml OP	ureus (MRSA) prior to elective the prescription is endorsed
RICLOSAN – Subsidy by endorsement a) Maximum of 500 ml per prescription b) a) Only if prescribed for a patient identified with M surgery in hospital and the prescription is endor b) Only if prescribed for a patient with recurrent St accordingly Soln 1% Barrier Creams and Emollients Barrier Creams DIMETHICONE k Crm 10% pump bottle	ethicillin-resistant Sta sed accordingly; or aphylococcus aureu 	aphylococcus a s infection and 3 500 ml OP 500 ml OP 500 ml OP	ureus (MRSA) prior to elective the prescription is endorsed
 RICLOSAN – Subsidy by endorsement a) Maximum of 500 ml per prescription b) 	ethicillin-resistant Sta sed accordingly; or aphylococcus aureu 	aphylococcus a s infection and 3 500 ml OP 500 ml OP 500 ml OP	ureus (MRSA) prior to elective the prescription is endorsed

*Three months or six months, as applicable, dispensed all-at-once

if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully Brand or
	(Manufacturer's F \$	Price) Subsi Per	dised Generic Manufacturer
CETOMACROGOL WITH GLYCEROL	· ·		
Crm 90% with glycerol 10%	2.82	500 ml OP	 Pharmacy Health
			Sorbolene with
			Glycerin
	3.87	1,000 ml OP	 Pharmacy Health
			Sorbolene with
			Glycerin
EMULSIFYING OINTMENT			_
* Oint BP	3.59	500 g	✓ <u>AFT</u>
DIL IN WATER EMULSION			
* Crm	2.25	500 g	✓ O/W Fatty Emulsion
			Cream
JREA			
* Crm 10%	1.37	100 g OP	 healthE Urea Cream
NOOL FAT WITH MINERAL OIL - Only on a prescription			
* Lotn hydrous 3% with mineral oil		1,000 ml	
	(11.95)		DP Lotion
	1.40	250 ml OP	DDLation
	(4.53) 5.60	1,000 ml	DP Lotion
	(20.53)	1,000 111	Alpha-Keri Lotion
	(23.91)		BK Lotion
	1.40	250 ml OP	
	(7.73)		BK Lotion
Other Dermatological Bases			
PARAFFIN			
White soft – Only in combination	20.20	2,500 g	🗸 IPW
	3.58	500 g	
	(7.78)	-	IPW
Only in combination with a dermatological galenical or	(8.69)		PSM

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid - Plain.

	Subsidy Manufacturer's Pric \$	e) Per	Fully Subsidised	
Minor Skin Infections				
POVIDONE IODINE				
Oint 10%	3.27	25 g O	Р 🗸	Betadine
a) Maximum of 100 g per prescription		Ũ		
b) Only on a prescription				
Antiseptic soln 10%	6.20	500 m	l 🗸	Betadine
			· 🗸	Riodine
	1.28	100 m	I	-
	(4.20)			Riodine
	(13.27)			Betadine
	0.19	15 ml		
	(7.41)			Betadine
Skin preparation, povidone iodine 10% with 30% alcohol		500 m	l 🗸	Betadine Skin Prep
	1.63	100 m		
	(3.48)			Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	8.13	500 m		
	(18.63)			Orion
	1.63	100 m		
	(6.04)			Orion
Parasiticidal Preparations				
METHICONE				
Lotn 4%	4.98	200 ml (א א	healthE
				Dimethicone 4%
ERMECTIN – Special Authority see SA1225 below – Retail pha			_	
Tab 3 mg – Up to 100 tab available on a PSO	17.20	4	1	Stromectol
 PSO for institutional use only. Must be endorsed wind special Authority for patient of that institution 		e institut	ion for wh	ich the PSO is required a

- a valid Special Authority for patient of that institution.2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or penal institutions.

⇒SA1225 Special Authority for Subsidy

Initial application — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:

2.1 Both:

- 2.1.1 The patient is in the community; and
- 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or

continued...

‡ safety cap

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sul	bsidised	Generic
\$	Per	1	Manufacturer

continued...

2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or

- 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently: and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scables hyperinfestation (Crusted/ Norwegian scables): or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy: or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application - (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

- Any of the following: 1 Filaricides: or

 - 2 Cutaneous larva migrans (creeping eruption); or
 - 3 Strongyloidiasis.

Renewal - (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist: and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy: or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution: and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy: or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal - (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

- Any of the following:
 - 1 Filaricides: or
 - 2 Cutaneous larva migrans (creeping eruption); or
 - 3 Strongvloidiasis.

PERMETHRIN

Crm 5%4	.95	30 g OP	 Lyderm
Lotn 5%		30 ml OP	 <u>A-Scabies</u>

			Fully Brand or
	Subsidy (Manufacturer's F	Price) Cub	,
	(IVIAIIUIACIUIEIS F	Per Sub	sidised Generic Manufacturer
HENOTHRIN	· · ·		
Shampoo 0.5%	11.36	200 ml OP	✓ Parasidose
•		200 01	
Psoriasis and Eczema Preparations			
CITRETIN – Special Authority see SA1476 below – Retail ph	narmacy		
Cap 10 mg		60	✓ Novatretin
Cap 25 mg	41.36	60	 Novatretin
SA1476 Special Authority for Subsidy			
itial application from any relevant practitioner. Approvals va I of the following:	alid for 1 year for a	pplications me	eting the following criteria:
1 Applicant is a vocationally registered dermatologist, voc	ationally registered	d general pract	itioner, or nurse practitioner
working in a relevant scope of practice; and	allorially registered	a gonorai praor	
 Applicant has an up to date knowledge of the safety iss Either: 	ues around acitreti	n and is compe	etent to prescribe acitretin; ar
3.1 Patient is female and has been counselled and u	inderstands the ris	k of teratogeni	city if acitretin is used during
pregnancy and the applicant has ensured that the			
commencement of the treatment and that the pa			
treatment and for a period of two years after the	completion of the t	treatment; or	
3.2 Patient is male.			
enewal from any relevant practitioner. Approvals valid for 1	wear for application	e mooting the	following criteria:
	year for application	is meeting the	0
ther:		-	-
ther: 1 Patient is female and has been counselled and understa	ands the risk of ter	atogenicity if a	citretin is used during pregna
ther:	ands the risk of ten gnancy has been e	atogenicity if a excluded prior t	citretin is used during pregna to the commencement of the
 Patient is female and has been counselled and understand the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 	ands the risk of ten gnancy has been e	atogenicity if a excluded prior t	citretin is used during pregna to the commencement of the
 ther: Patient is female and has been counselled and understand the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must 	ands the risk of ten gnancy has been e	atogenicity if a excluded prior t	citretin is used during pregna to the commencement of the
 Patient is female and has been counselled and understand the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 	ands the risk of ten gnancy has been e	atogenicity if a excluded prior t	citretin is used during pregna to the commencement of the
 ther: Patient is female and has been counselled and understand the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g 	ands the risk of ter gnancy has been e not become pregr	atogenicity if ac excluded prior t nant during trea 30 g OP	citretin is used during pregna to the commencement of the
 ther: Patient is female and has been counselled and understand the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or Patient is male. 	ands the risk of ter gnancy has been e not become pregr	atogenicity if ad excluded prior t eant during tree	citretin is used during pregna to the commencement of the ttment and for a period of two
 ther: Patient is female and has been counselled and understand the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g	ands the risk of ter gnancy has been e not become pregr 	atogenicity if ac excluded prior t nant during trea 30 g OP	citretin is used during pregna to the commencement of the atment and for a period of two <u>Daivobet</u>
 ther: Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pre treatment and that the patient is informed that she must years after the completion of the treatment; or Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g 	ands the risk of ter gnancy has been e not become pregr 	atogenicity if ac excluded prior t nant during trea 30 g OP	citretin is used during pregna to the commencement of the atment and for a period of two <u>Daivobet</u>
 ther: 1 Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 2 Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g ALCIPOTRIOL Oint 50 mcg per g 	ands the risk of ter gnancy has been e not become pregr 	atogenicity if a excluded prior t nant during trea 30 g OP 30 g OP	citretin is used during pregna to the commencement of the attment and for a period of two <u>Daivobet</u> <u>Daivobet</u>
 ther: Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g ALCIPOTRIOL Oint 50 mcg per g 	ands the risk of ten gnancy has been e not become pregr 	atogenicity if a excluded prior t nant during trea 30 g OP 30 g OP	citretin is used during pregna to the commencement of the attment and for a period of two <u>Daivobet</u> <u>Daivobet</u>
 ther: 1 Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 2 Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g ALCIPOTRIOL Oint 50 mcg per g DAL TAR 	ands the risk of ten gnancy has been e not become pregr 	atogenicity if a excluded prior t ant during trea 30 g OP 30 g OP 100 g OP 200 ml	citretin is used during pregna to the commencement of the strment and for a period of two <u>Daivobet</u> <u>Daivobet</u> <u>Daivonex</u> <u>Midwest</u>
 ther: 1 Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 2 Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g ALCIPOTRIOL Oint 50 mcg per g DAL TAR Soln BP – Only in combination 	ands the risk of ten gnancy has been e not become pregr 	atogenicity if a excluded prior t ant during trea 30 g OP 30 g OP 100 g OP 200 ml	citretin is used during pregna to the commencement of the ttment and for a period of two <u>Daivobet</u> <u>Daivobet</u> <u>Daivonex</u> <u>Midwest</u>
 ther: 1 Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 2 Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g ALCIPOTRIOL Oint 50 mcg per g DAL TAR Soln BP - Only in combination	ands the risk of ten gnancy has been e not become pregr 	atogenicity if a excluded prior t ant during trea 30 g OP 30 g OP 100 g OP 200 ml	citretin is used during pregna to the commencement of the ttment and for a period of two <u>Daivobet</u> <u>Daivobet</u> <u>Daivonex</u> <u>Midwest</u>
 ther: 1 Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 2 Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g ALCIPOTRIOL Oint 50 mcg per g DAL TAR Soln BP - Only in combination 1) Up to 10% only in combination with a dermatolog dermatological base, page 223 2) With or without other dermatological galenicals. 	ands the risk of ten gnancy has been e not become pregn 	atogenicity if a excluded prior t ant during trea 30 g OP 30 g OP 100 g OP 200 ml	citretin is used during pregna to the commencement of the ttment and for a period of two <u>Daivobet</u> <u>Daivobet</u> <u>Daivonex</u> <u>Midwest</u>
 ther: 1 Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 2 Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g ALCIPOTRIOL Oint 50 mcg per g DAL TAR Soln BP - Only in combination	ands the risk of tern gnancy has been e not become pregr 	atogenicity if a excluded prior t ant during trea 30 g OP 30 g OP 100 g OP 200 ml	citretin is used during pregna to the commencement of the ttment and for a period of two <u>Daivobet</u> <u>Daivobet</u> <u>Daivonex</u> <u>Midwest</u>
 ther: 1 Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 2 Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g ALCIPOTRIOL Oint 50 mcg per g DAL TAR Soln BP - Only in combination 1) Up to 10% only in combination with a dermatolog dermatological base, page 223 2) With or without other dermatological galenicals. 	ands the risk of ter gnancy has been e not become pregr 	atogenicity if a excluded prior t ant during trea 30 g OP 30 g OP 100 g OP 200 ml ietary Topical C	citretin is used during pregna to the commencement of the ttment and for a period of two <u>Daivobet</u> <u>Daivobet</u> <u>Daivonex</u> <u>Midwest</u>
 ther: 1 Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 2 Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g ALCIPOTRIOL Oint 50 mcg per g DAL TAR Soln BP - Only in combination	ands the risk of terr gnancy has been e not become pregr 	atogenicity if a excluded prior t ant during trea 30 g OP 30 g OP 100 g OP 200 ml	citretin is used during pregna to the commencement of the atment and for a period of two <u>Daivobet</u> <u>Daivobet</u> <u>Daivonex</u> <u>Midwest</u> Corticosteriod – Plain, refer
 ther: 1 Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 2 Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g ALCIPOTRIOL Oint 50 mcg per g DAL TAR Soln BP - Only in combination 1) Up to 10% only in combination with a dermatolog dermatological base, page 223 2) With or without other dermatological galenicals. 	ands the risk of ter gnancy has been e not become pregr 	atogenicity if a excluded prior t aant during trea 30 g OP 30 g OP 100 g OP 200 ml tetary Topical C	citretin is used during pregna to the commencement of the ttment and for a period of two <u>Daivobet</u> <u>Daivobet</u> <u>Daivonex</u> <u>Midwest</u>
 ther: 1 Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 2 Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g ALCIPOTRIOL Oint 50 mcg per g DAL TAR Soln BP - Only in combination 1) Up to 10% only in combination with a dermatolog dermatological base, page 223 2) With or without other dermatological galenicals. 	ands the risk of terr gnancy has been e not become pregr 	atogenicity if a excluded prior t ant during trea 30 g OP 30 g OP 100 g OP 200 ml ietary Topical C	citretin is used during pregna to the commencement of the attment and for a period of two <u>Daivobet</u> <u>Daivobet</u> <u>Daivonex</u> <u>Midwest</u> Corticosteriod – Plain, refer
 ther: 1 Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 2 Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g DALCIPOTRIOL Oint 50 mcg per g DAL TAR Soln BP – Only in combination 1) Up to 10% only in combination with a dermatolog dermatological base, page 223 2) With or without other dermatological galenicals. DAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SU Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% a allantoin crm 2.5%. 	ands the risk of ter gnancy has been e not become pregr 	atogenicity if a excluded prior t aant during trea 30 g OP 30 g OP 100 g OP 200 ml tetary Topical C	citretin is used during pregna to the commencement of the atment and for a period of two <u>Daivobet</u> <u>Daivobet</u> <u>Daivonex</u> <u>Midwest</u> Corticosteriod – Plain, refer
 ther: 1 Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 2 Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g CALCIPOTRIOL Oint 50 mcg per g DAL TAR Soln BP – Only in combination 1) Up to 10% only in combination with a dermatolog dermatological base, page 223 2) With or without other dermatological galenicals. DAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SU Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% a allantoin crm 2.5% 	ands the risk of ter gnancy has been e not become pregr 	atogenicity if a excluded prior t ant during trea 30 g OP 30 g OP 100 g OP 200 ml detary Topical C 75 g OP 30 g OP	citretin is used during pregna to the commencement of the attment and for a period of two <u>Daivobet</u> <u>Daivobet</u> <u>Daivonex</u> <u>Midwest</u> Corticosteriod – Plain, refer Egopsoryl TA Egopsoryl TA
 ther: 1 Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 2 Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g	ands the risk of ter gnancy has been e not become pregr 	atogenicity if ac excluded prior t ant during treat 30 g OP 30 g OP 100 g OP 200 ml tetary Topical O 75 g OP 30 g OP 40 g OP	citretin is used during pregna to the commencement of the stiment and for a period of two <u>Daivobet</u> <u>Daivobet</u> <u>Daivonex</u> <u>Midwest</u> Corticosteriod – Plain, refer Egopsoryl TA Egopsoryl TA <u>Egopsoryl TA</u>
 ther: 1 Patient is female and has been counselled and understa and the applicant has ensured that the possibility of pretreatment and that the patient is informed that she must years after the completion of the treatment; or 2 Patient is male. ETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g CALCIPOTRIOL Oint 50 mcg per g DAL TAR Soln BP – Only in combination 1) Up to 10% only in combination with a dermatolog dermatological base, page 223 2) With or without other dermatological galenicals. DAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SU Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% a allantoin crm 2.5% 	ands the risk of ter gnancy has been e not become pregr 	atogenicity if ac excluded prior t ant during treat 30 g OP 30 g OP 100 g OP 200 ml tetary Topical O 75 g OP 30 g OP 40 g OP	citretin is used during pregna to the commencement of the stiment and for a period of two <u>Daivobet</u> <u>Daivobet</u> <u>Daivonex</u> <u>Midwest</u> Corticosteriod – Plain, refer Egopsoryl TA Egopsoryl TA <u>Egopsoryl TA</u>

	Subsidy)rice) Cub	Fully	Brand or
	(Manufacturer's P \$	Per Sub	sidised ✓	Generic Manufacturer
SALICYLIC ACID				
Powder – Only in combination		250 g	🗸 PS	
1) Only in combination with a dermatological base or	proprietary Topic	cal Corticoster	oid – Plair	or collodion flexible,
refer dermatological base, page 223 2) With or without other dermatological galenicals.				
SULPHUR				
Precipitated – Only in combination	6.35	100 g	🗸 Mic	lwest
1) Only in combination with a dermatological base or	proprietary Topic	cal Corticoster	oid – Plair	n, refer dermatological
base, page 223				
2) With or without other dermatological galenicals.				
Scalp Preparations				
BETAMETHASONE VALERATE				
* Scalp app 0.1%	7.75	100 ml OP	🗸 Be	ta Scalp
CLOBETASOL PROPIONATE				
* Scalp app 0.05%	6.96	30 ml OP	🗸 De	rmol
HYDROCORTISONE BUTYRATE				
Scalp lotn 0.1%	3.65	100 ml OP	✓ Lo	coid
KETOCONAZOLE	2.00		✓ Sel	hizolo
a) Maximum of 100 ml per prescription	2.99	100 ml OP	• <u>5</u>	DIZOIE
b) Only on a prescription				
Sunscreens				
SUNSCREENS, PROPRIETARY - Subsidy by endorsement				
Only if prescribed for a patient with severe photosensitivity se	econdary to a de	fined clinical c	ondition a	nd the prescription is
endorsed accordingly. Crm	3 30	100 g OP		
Unit	(5.89)	100 9 01	Ha	milton Sunscreen
Lotn,	· · ·	100 g OP		rine Blue Lotion
	- /-			PF 50+
	5.10	200 g OP		rine Blue Lotion PF 50+
				NT 30+
Wart Preparations				
For salicylic acid preparations refer to PSORIASIS AND ECZEM	A PREPARATIO	NS, page 79		
		, page re		
Crm 5%, 250 mg sachet	17.98	12	🗸 Ap	o-Imiquimod
-			Ċ	cream 5%
	21.72	24	✓ Per	rrigo
PODOPHYLLOTOXIN	00.00		10	n du dine e
Soln 0.5%a) Maximum of 3.5 ml per prescription		3.5 ml OP	✔ Co	ndyline
b) Only on a prescription				
, , , , , , , , , , , , , , , , , , ,				

	Subsidy (Manufacturer's Price \$	e) Subsi Per	Fully idised	Brand or Generic Manufacturer
Other Skin Preparations				
Antineoplastics				
FLUOROURACIL SODIUM Crm 5%	8.95	20 g OP	✓ <u>E</u> f	fudix

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Contraceptives - Non-hormonal				
Condoms				
CONDOMS				
* 49 mm – Up to 144 dev available on a PSO		144	✓	Shield 49
* 53 mm – Up to 144 dev available on a PSO	1.11	12		Gold Knight Shield Blue
	13.36	144	✓	Shield Blue
* 53 mm (chocolate) - Up to 144 dev available on a PSO	1.11	12	✓	Gold Knight
	13.36	144	✓	Gold Knight
* 53 mm (strawberry) - Up to 144 dev available on a PSO	1.11	12	✓	Gold Knight
	13.36	144	✓	Gold Knight
* 56 mm – Up to 144 dev available on a PSO	1.11	12	✓	Gold Knight
	13.36	144	✓	Durex Extra Safe
			✓	Gold Knight
* 56 mm, shaped - Up to 144 dev available on a PSO	1.11	12	✓	Durex Confidence
	13.36	144	✓	Durex Confidence
* 60 mm – Up to 144 dev available on a PSO	13.36	144	1	Shield XL
Contraceptive Devices				
INTRA-UTERINE DEVICE				
a) Up to 40 dev available on a PSO b) Only on a PSO				
* IUD 29.1 mm length × 23.2 mm width		1	1	Choice TT380 Short
* IUD 33.6 mm length × 29.9 mm width		1		Choice TT380 Standard
* IUD 35.5 mm length × 19.6 mm width	31.60	1	✓	Choice Load 375
Contracontivos Hormonal				

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 Either:

- 1.1 Patient is on a Social Welfare benefit; or
- 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

continued...

	Subsidy (Manufacturer's Price) \$	Ful Subsidise Per •	
continued The additional subsidy will fund Mercilon and Marvelon up to the	manufacturer's price 1	for each of the	ese products as identified on
the Schedule at 1 November 1999. Special Authorities approved before 1 November 1999 remain va	lid until the expiry dat	e and can be	renewed providing that
women are still either:	and until the expiry dut		renewed providing that
 on a Social Welfare benefit; or 			
 have an income no greater than the benefit. 			
The approval numbers of Special Authorities approved before 1 N	November 1999 are in	terchangeable	e for products within the
combined oral contraceptives and progestogen-only contraceptiv	es groups, except Loe	ette and Micro	gynon 20 ED
ETHINYLOESTRADIOL WITH DESOGESTREL			
 * Tab 20 mcg with desogestrel 150 mcg and 7 inert tab 	6.62	84	
	(19.80)		Mercilon 28
a) Higher subsidy of \$13.80 per 84 tab with Special Aut	hority see SA0500 on	the previous	page
b) Up to 84 tab available on a PSO	6 60	0.4	
* Tab 30 mcg with desogestrel 150 mcg and 7 inert tab		84	Marvelon 28
a) Higher subsidy of \$13.80 per 84 tab with Special Aut	· · · ·	the previous	
b) Up to 84 tab available on a PSO			page
ETHINYLOESTRADIOL WITH LEVONORGESTREL			
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_		
Up to 84 tab available on a PSO		84 🖌	Microgynon 20 ED
* Tab 50 mcg with levonorgestrel 125 mcg and 7 inert tab - U			
to 84 tab available on a PSO		84 🖌	Microgynon 50 ED
* Tab 30 mcg with levonorgestrel 150 mcg	6.62	63	
	(16.50)		Microgynon 30
a) Higher subsidy of \$15.00 per 63 tab with Special Autb) Up to 63 tab available on a PSO	hority see SA0500 on	the previous	page
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets			• · · ·
Up to 84 tab available on a PSO	1.77	84 🖌	Levlen ED
ETHINYLOESTRADIOL WITH NORETHISTERONE			
* Tab 35 mcg with norethisterone 1 mg – Up to 63 tab availab on a PSO		63 🖌	Brevinor 1/21
* Tab 35 mcg with norethisterone 1 mg and 7 inert tab – Up to 84 tab available on a PSO		84 🖌	Brevinor 1/28
* Tab 35 mcg with norethisterone 500 mcg – Up to 63 tab available on a PSO	6.62	63 🖌	Brevinor 21
* Tab 35 mcg with norethisterone 500 mcg and 7 inert tab – U			
to 84 tab available on a PSO		84 🖌	Norimin

Progestogen-only Contraceptives

► SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 Either:

1.1 Patient is on a Social Welfare benefit; or

continued...

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once if

▲ Three months supply may be dispensed at one time

GENITO-URINARY SYSTEM

Subsidy	F	ully	Brand or
(Manufacturer's	Price) Subsid	ised	Generic
\$	Per	✓	Manufacturer

continued...

1.2 Patient has an income no greater than the benefit; and

2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED LEVONORGESTREL

ELIONONGEONNEE			
* Tab 30 mcg		84	Minuclut
	(16.50)		Microlut
 a) Higher subsidy of \$13.80 per 84 tab with Special Authority b) Up to 84 tab available on a PSO 	/ see SA0500 o	n the previ	ous page
✤ Subdermal implant (2 × 75 mg rods) – Up to 3 pack available			
on a PSO	.106.92	1	✓ Jadelle
MEDROXYPROGESTERONE ACETATE			
* Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PSO.	7.25	1	Depo-Provera
NORETHISTERONE		·	
* Tab 350 mcg – Up to 84 tab available on a PSO	6 25	84	Noriday 28
	0.20	04	V Nonday 20
Emergency Contraceptives			
LEVONORGESTREL			
* Tab 1.5 mg	4.95	1	Postinor-1
-) Manimum af O table and an initial			

a) Maximum of 2 tab per prescription

b) Up to 5 tab available on a PSO

c) Note: may be provided by a pharmacist under the non-prescribing Practitioners provisions in Part III of Section A.

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

*	Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up	
	to 168 tab available on a PSO4.67	168


Ginet

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's P	,	Fully Brand or idised Generic
Gynaecological Anti-infectives	\$	Per	Manufacturer
ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulpha			
0.025%, glycerol 5% and ricinoleic acid 0.75% with app	licator8.43 (24.00)	100 g OP	Aci-Jel
CLOTRIMAZOLE * Vaginal crm 1% with applicators * Vaginal crm 2% with applicators		35 g OP 20 g OP	✓ <u>Clomazol</u> ✓ Clomazol
MICONAZOLE NITRATE * Vaginal crm 2% with applicator		40 g OP	✓ Micreme
NYSTATIN Vaginal crm 100,000 u per 5 g with applicator(s)	4.45	75 g OP	✓ <u>Nilstat</u>
Myometrial and Vaginal Hormone Preparations			
ERGOMETRINE MALEATE			
Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on PSO		5	✓ DBL Ergometrine
OESTRIOL * Crm 1 mg per g with applicator Pessaries 500 mcg		15 g OP 15	 ✓ <u>Ovestin</u> ✓ Ovestin
OXYTOCIN – Up to 5 inj available on a PSO Inj 5 iu per ml, 1 ml ampoule Inj 10 iu per ml, 1 ml ampoule		5 5	✓ <u>Oxytocin BNM</u> ✓ Oxytocin Apotex
(Oxytocin Apotex Inj 10 iu per ml, 1 ml ampoule to be delisted 1	December 2018)		 Oxytocin BNM
OXYTOCIN WITH ERGOMETRINE MALEATE – Up to 5 inj ava Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml		5	✓ Syntometrine
Pregnancy Tests - hCG Urine			
PREGNANCY TESTS - HCG URINE a) Up to 200 test available on a PSO			
b) Only on a PSO Cassette	17.60	40 test OP	 EasyCheck
Urinary Agents			
For urinary tract Infections refer to INFECTIONS, Antibacterials,	page 117		
5-Alpha Reductase Inhibitors			
FINASTERIDE – Special Authority see SA0928 below – Retail p * Tab 5 mg		100	✓ <u>Ricit</u>
SA0928 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals vali the following criteria:	id without further	renewal unless	notified for applications meeting
			continued

‡ safety cap

 $\ensuremath{\boldsymbol{\ast}}$ Three months or six months, as applicable, dispensed all-at-once



GENITO-URINARY SYSTEM

	Subsidy Manufacturer's Price)	Subsid	- ully ised	Brand or Generic
·	\$	Per	1	Manufacturer

SA1272 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where patient has overactive bladder and a documented intolerance of, or is non-responsive to oxybutynin.

Detection of Substances in Urine			
ORTHO-TOLIDINE			
* Compound diagnostic sticks	7.50	50 test OP	
	(8.25)		Hemastix
TETRABROMOPHENOL			
* Blue diagnostic strips	7.02	100 test OP	
	(13.92)		Albustix

	Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
	\$	Per	✓	Manufacturer
Calcium Homeostasis				
CALCITONIN * Inj 100 iu per ml, 1 ml ampoule		5	✓ Mi	acalcic
CINACALCET – Special Authority see SA1618 below – Retail pr Tab 30 mg – Wastage claimable – see rule 3.3.2 on page 13		28	🗸 Se	ensipar
■ SA1618 Special Authority for Subsidy Initial application only from a nephrologist or endocrinologist. A following criteria: Either:	opprovals valid for 6 m	nonths for a	applicati	ons meeting the
1 All of the following:				
 1.1 The patient has been diagnosed with a parathyroid 1.2 The patient has persistent hypercalcaemia (serum first-line treatments including sodium thiosulfate (w 1.3 The patient is symptomatic; or 2 All of the following: 	calcium greater than	or equal to		
 2.1 The patient has been diagnosed with calciphylaxis 2.2 The patient has symptomatic (e.g. painful skin ulc mmol/L); and 2.3 The patient's condition has not responded to previous thiosulfate. 	ers) hypercalcaemia (serum calo	cium gre	
Renewal only from a nephrologist or endocrinologist. Approvals meeting the following criteria: Both:	valid without further r	enewal unl	ess not	ified for applications
 The patient's serum calcium level has fallen to < 3mmol/L The patient has experienced clinically significant symptom 	,			
Note: This does not include parathyroid adenomas unless these	have become malign	ant.		
ZOLEDRONIC ACID				
Inj 4 mg per 5 ml, vial – Special Authority see SA1687 below				
Retail pharmacy	84.50	1		oledronic acid Mylan
	550.00		✓ Zo	ometa
SA1687 Special Authority for Subsidy Initial application — (bone metastases) only from an oncologi without further renewal unless notified for applications meeting th Any of the following:		alliative ca	re spec	ialist. Approvals valid
 Patient has hypercalcaemia of malignancy; or Both: 				
2.1 Patient has bone metastases or involvement; and2.2 Patient has severe bone pain resistant to standard	first-line treatments;	or		
3 Both:				
3.1 Patient has bone metastases or involvement; and	rical fracture animal a	ord comer	nonior	radiation to hone or

3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer) only from an oncologist or medical practitioner on the recommendation of a oncologist. Approvals valid for 2 years for applications meeting the following criteria:

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	Manufacturer

continued...

All of the following:

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.

Corticosteroids and Related Agents for Systemic Use		
BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETA	TE	
* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	5	
(36.96)		Celestone
		Chronodose
DEXAMETHASONE		
* Tab 0.5 mg - Retail pharmacy-Specialist0.88	30	 Dexmethsone
Up to 60 tab available on a PSO		
* Tab 4 mg – Retail pharmacy-Specialist1.84	30	 Dexmethsone
Up to 30 tab available on a PSO		
Oral liq 1 mg per ml – Retail pharmacy-Specialist45.00	25 ml OP	 Biomed
Oral liq prescriptions:		
1) Must be written by a Paediatrician or Paediatric Cardiologist; or		
On the recommendation of a Paediatrician or Paediatric Cardiolog	jist.	
DEXAMETHASONE PHOSPHATE		
Dexamethasone phosphate injection will not be funded for oral use.		
* Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO 14.19	10	 Max Health
* Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO25.18	10	 Max Health
FLUDROCORTISONE ACETATE		
* Tab 100 mcg14.32	100	 Florinef
HYDROCORTISONE		
* Tab 5 mg	100	✓ Douglas
 * Tab 20 mg – For hydrocortisone oral liquid formulation refer, 	100	Dougluo
page 224	100	 Douglas
* Inj 100 mg vial	1	✓ Solu-Cortef
a) Up to 5 inj available on a PSO	·	
b) Only on a PSO		
METHYLPREDNISOLONE – Retail pharmacy-Specialist		
* Tab 4 mg	100	✓ Medrol
* Tab 100 mg	20	✓ <u>Medrol</u>
		Medior
METHYLPREDNISOLONE (AS SODIUM SUCCINATE) – Retail pharmacy-Spe		Colu Modual
Inj 40 mg vial	1	Solu-Medrol
Inj 125 mg vial	1	 ✓ <u>Solu-Medrol</u> ✓ <u>Solu-Medrol</u>
Inj 500 mg vial9.00 Inj 1 g vial16.00	1	 ✓ Solu-Medrol ✓ Solu-Medrol
, ,	I	
METHYLPREDNISOLONE ACETATE	-	Dama Madual
Inj 40 mg per ml, 1 ml vial40.00	5	Depo-Medrol
METHYLPREDNISOLONE ACETATE WITH LIDOCAINE [LIGNOCAINE]		
Inj 40 mg per ml with lidocaine [lignocaine] 1 ml vial	1	 Depo-Medrol with
		Lidocaine

‡ safety cap

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

PREDNISOLONE * Oral liq 5 mg per ml – Up to 30 ml available on a PSOa) a) Restricted to children under 12 years of age. b) Redipred to be Sole Supply on 1 July 2018 PREDNISONE * Tab 1 mg * Tab 2.5 mg * Tab 5 mg – Up to 30 tab available on a PSO * Tab 20 mg	10.68 12.09 11.09	30 ml OP 500 500 500	* *	Redipred <u>Apo-Prednisone</u> Apo-Prednisone
 a) Restricted to children under 12 years of age. b) Redipred to be Sole Supply on 1 July 2018 PREDNISONE * Tab 1 mg * Tab 2.5 mg * Tab 5 mg – Up to 30 tab available on a PSO 	10.68 12.09 11.09	500 500 500	* *	Apo-Prednisone
 * Tab 1 mg * Tab 2.5 mg * Tab 5 mg - Up to 30 tab available on a PSO 	12.09 11.09	500 500	✓	
 * Tab 2.5 mg * Tab 5 mg - Up to 30 tab available on a PSO 	12.09 11.09	500 500	✓	
* Tab 5 mg – Up to 30 tab available on a PSO	11.09	500		Ano-Prednisone
•			1	
* Tab 20 mg	29.03		•	Apo-Prednisone
		500	~	Apo-Prednisone
TETRACOSACTRIN				
* Inj 250 mcg per ml, 1 ml ampoule	75.00	1	1	Synacthen
* Inj 1 mg per ml, 1 ml ampoule		1	1	Synacthen Depot
TRIAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule	20.80	5	1	Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule		5		Kenacort-A 40
Sex Hormones Non Contraceptive				
Androgen Agonists and Antagonists				
CYPROTERONE ACETATE – Retail pharmacy-Specialist				
Tab 50 mg	15.87	50	1	Procur
Tab 100 mg		50		Procur
TESTOSTEBONE				
Patch 5 mg per day	80.00	30	1	Androderm
	00.00	50	•	Androuenni
TESTOSTERONE CIPIONATE – Retail pharmacy-Specialist				-
Inj 100 mg per ml, 10 ml vial	76.50	1	•	Depo-Testosterone
TESTOSTERONE ESTERS – Retail pharmacy-Specialist				
Inj 250 mg per ml, 1 ml	12.98	1	~	Sustanon Ampoules
TESTOSTERONE UNDECANOATE – Retail pharmacy-Specialist				
Cap 40 mg	16.80	60	✓	Andriol Testocaps
Inj 250 mg per ml, 4 ml vial	86.00	1	✓	Reandron 1000

Hormone Replacement Therapy - Systemic

Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

	Subsidy	Full	y Brand or
	(Manufacturer's Price)) Subsidise	d Generic
	\$	Per 🖌	Manufacturer
Oestrogens			
OESTRADIOL – See prescribing guideline on the previous page			
* Tab 1 mg		28 OP	
* Tab Ting		20 UF	Estrofem
* Tab Orea	(11.10)		Estiolem
* Tab 2 mg		28 OP	E dua fa m
	(11.10)		Estrofem
* Patch 25 mcg per day	6.12	8 🗸	Estradot
 a) No more than 2 patch per week 			
 b) Only on a prescription 			
* Patch 50 mcg per day	7.04	8 🖌	Estradot 50 mcg
a) No more than 2 patch per week			
b) Only on a prescription			
 Patch 75 mcg per day 	7 91	8 🗸	Estradot
•••••		0	
a) No more than 2 patch per week			
b) Only on a prescription			-
 Patch 100 mcg per day 	7.91	8 🗸	Estradot
 a) No more than 2 patch per week 			
 b) Only on a prescription 			
OESTRADIOL VALERATE - See prescribing guideline on the pr			
* Tab 1 mg		84 🗸	Progynova
5		• •	
* Tab 2 mg		84 🗸	Progynova
OESTROGENS – See prescribing guideline on the previous pag	e		
Conjugated, equine tab 300 mcg	3.01	28	
	(13.50)		Premarin
* Conjugated, equine tab 625 mcg		28	
- J. S	(13.50)		Premarin
	(10100)		
Progestogens			
MEDROXYPROGESTERONE ACETATE – See prescribing guid	leline on the previou	s page	
* Tab 2.5 mg			Provera
· · · · · · · · · · · · · · · · · · ·	7.00		Provera S29 S29
* Tab 5 mg			Provera
0			
* Tab 10 mg		30	Provera
(Provera S29 S29) Tab 2.5 mg to be delisted 1 September 2018))		
Progestogen and Oestrogen Combined Prepara	tions		
OESTRADIOL WITH NORETHISTERONE – See prescribing gui	ideline on the previo		
 Tab 1 mg with 0.5 mg norethisterone acetate 		28 OP	
י דמט ד וווץ שונו ט.ט וווץ ווטופנוווטופוטוופ מטפומופ		20 OF	Kliovance
* Tab 0 mg with 1 mg novathints range asstate	(18.10)		NIIOVATICE
* Tab 2 mg with 1 mg norethisterone acetate		28 OP	
	(18.10)		Kliogest
* Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg			
oestradiol tab (12) and 1 mg oestradiol tab (6)	5.40	28 OP	
	(18.10)		Trisequens
	· · /		·

‡ safety cap

	Subsidy (Manufacturer's Price)	c	Fully Subsidised	Brand or Generic
	(Manulacturer's Flice) \$	Per		Manufacturer
Other Oestrogen Preparations				
ETHINYLOESTRADIOL				
* Tab 10 mcg	17.60	100	✓ <u>N</u>	Z Medical and Scientific
DESTRIOL				
* Tab 2 mg	7.00	30	✓ 0	vestin
Other Progestogen Preparations				
EVONORGESTREL				
 Intra-uterine system 20 mcg per day – Special Authority see 				
SA1608 below – Retail pharmacy		1	✓ <u>N</u>	lirena
SA1608 Special Authority for Subsidy nitial application — (No previous use) only from a relevant sp	pecialist or general pr	actition	er. Approv	vals valid for 6 months for
applications meeting the following criteria:				
All of the following: 1 The patient has a clinical diagnosis of heavy menstrual bl	eeding: and			
2 The patient has failed to respond to or is unable to tolerati		harmad	ceutical the	rapies as per the Heavy
Menstrual Bleeding Guidelines; and				
3 Either:				
3.1 serum ferritin level $<$ 16 mcg/l (within the last 12 m 3.2 haemoglobin level $<$ 120 g/l.	nonths); or			
Note: Applications are not to be made for use in patients as cont				
Renewal only from a relevant specialist or general practitioner.	Approvals valid for 6	months	for applica	ations meeting the
ollowing criteria: Both:				
1 Either:				
1.1 Patient demonstrated clinical improvement of heav	y menstrual bleeding	j; or		
1.2 Previous insertion was removed or expelled within	3 months of insertion	n; and		
2 Applicant to state date of the previous insertion.				
MEDROXYPROGESTERONE ACETATE				
* Tab 100 mg – Retail pharmacy-Specialist	101.00	100	✓ <u>P</u>	rovera HD
NORETHISTERONE	10.00	400	10	
★ Tab 5 mg – Up to 30 tab available on a PSO		100	• •	rimolut N
PROGESTERONE				
Cap 100 mg – Special Authority see SA1609 below – Retail pharmacy		30	🗸 II	trogestan
SA1609 Special Authority for Subsidy		00	<u> </u>	
nitial application only from an obstetrician or gynaecologist. A	oprovals valid for 12	months	for applica	tions meeting the
ollowing criteria:				Ū
Both:				
1 For the prevention of pre-term labour*; and				
2 Either:				
2.1 The patient has a short cervix on ultrasound (defin 2.2 The patient has a history of pre-term birth at less the		to 28 v	veeks); or	

2.2 The patient has a history of pre-term birth at less than 28 weeks.

continued...

Subsidy (Manufacturer's Price)	5	Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 For the prevention of pre-term labour*; and

2 Treatment is required for second or subsequent pregnancy; and

3 Either:

3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or

3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are Unapproved Indications (refer to Interpretations and Definitions).

Thyroid and Antithyroid Agents CARBIMAZOI F 100 ✓ AFT Carbimazole S29 Neo-Mercazole **LEVOTHYROXINE** 90 Synthroid [±] Safety cap for extemporaneously compounded oral liquid preparations. Tab 50 mcg......1.71 28 Mercury Pharma 90 Svnthroid 4.05 64.28 1.000 ✓ Eltroxin [‡] Safety cap for extemporaneously compounded oral liquid preparations. 28 Mercury Pharma Synthroid 4.21 90 66.78 1.000 ✓ Fitroxin ‡ Safety cap for extemporaneously compounded oral liquid preparations. PROPYLTHIOURACIL - Special Authority see SA1199 below - Retail pharmacy Propylthiouracil is not recommended for patients under the age of 18 years unless the patient is pregnant and other treatments are contraindicated. 100 ✓ PTU S29 ■ SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 The patient has hyperthyroidism; and

2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) – Special Authority see S	SA1629 on the next page - I	Retail pha	rmacy
*	Inj 5 mg cartridge		1	 Omnitrope
*	Inj 10 mg cartridge		1	 Omnitrope
*	Inj 15 mg cartridge		1	 Omnitrope

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1629 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
- 2 All of the following:
 - Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under ; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

continued...

\$ safety cap

	Subsidy	F	ully	Brand or
(Mar	nufacturer's Price)	Subsidi	sed	Generic
	\$	Per	✓	Manufacturer

continued...

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight eximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

continued...

Subsidy	Fu	lly	Brand or	-
(Manufacturer's Price)	Subsidise	ed	Generic	
\$	Per	/	Manufacturer	

continued...

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients.

GnRH Analogues

GOSERELIN			
Implant 3.6 mg, syringe	66.48	1	✓ Zoladex
Implant 10.8 mg, syringe	177.50	1	✓ Zoladex
LEUPRORELIN			
Additional subsidy by endorsement where the patient is a child or goserelin and the prescription is endorsed accordingly.	or adolescen	t and is unable	to tolerate administration of
Inj 3.75 mg prefilled dual chamber syringe - Higher subsidy of			
\$221.60 per 1 inj with Endorsement	66.48	1	
	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe – Higher subsidy			
of \$591.68 per 1 inj with Endorsement	177.50	1	
	(591.68)		Lucrin Depot 3-month
Vasopressin Agonists DESMOPRESSIN ACETATE The descent of the desce			
Tab 100 mcg – Special Authority see SA1401 on the next page – Retail pharmacy	25.00	30	✓ Minirin
Tab 200 mcg – Special Authority see SA1401 on the next page			
 Retail pharmacy 		30	✓ Minirin
▲ Nasal drops 100 mcg per ml – Retail pharmacy-Specialist		2.5 ml OP	 Minirin
▲ Nasal spray 10 mcg per dose – Retail pharmacy-Specialist	23.95	6 ml OP	✓ <u>Desmopressin-</u> <u>PH&T</u>
Inj 4 mcg per ml, 1 ml – Special Authority see SA1401 on the			
next page – Retail pharmacy	67.18	10	 Minirin

‡ safety cap

lacksquare Three months supply may be dispensed at one time

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$ Per		Manufacturer

⇒SA1401 Special Authority for Subsidy

Initial application — (Desmopressin tablets for Nocturnal enuresis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has primary nocturnal enuresis; and
- 2 The nasal forms of desmopressin are contraindicated; and
- 3 An enuresis alarm is contraindicated.

Initial application — (Desmopressin tablets for Diabetes insipidus) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has cranial diabetes insipidus; and
- 2 The nasal forms of desmopressin are contraindicated.

Renewal — (Desmopressin tablets) from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from the treatment.

Initial application — (Desmopressin injection) only from a relevant specialist. Approvals valid for 2 years where the patient cannot use desmopressin nasal spray or nasal drops.

Renewal — (Desmopressin injection) only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Other Endocrine Agents

CABERGOLINE

		Tab 0.5 mg – Maximum of 2 tab per prescription; can be
 <u>Dostinex</u> 	2	waived by Special Authority see SA1370 below4.75
 Dostinex 	8	19.00

⇒SA1370 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

1 pathological hyperprolactinemia; or

2 acromegaly*.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment. Note: Indication marked with * is an Unapproved indication.

CLOMIFENE CITRATE

Tab 50 mg	29.84	10	 Mylan Clomiphen S29
			 Serophene
DANAZOL			
Cap 100 mg		100	🗸 Azol
Cap 200 mg	97.83	100	🗸 Azol
METYRAPONE			
Cap 250 mg – Retail pharmacy-Specialist	520.00	50	 Metopirone

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Per	Subsidised	Generic
	\$	Per		Manufacturer
Anthelmintics				
ALBENDAZOLE - Special Authority see SA1318 below - Retai	l pharmacy			
Tab 400 mg		60	1	Eskazole S29
➡SA1318 Special Authority for Subsidy				
Initial application only from an infectious disease specialist or o	clinical microbiologist.	Appr	ovals valic	for 6 months where the
patient has hydatids.	0			
Renewal only from an infectious disease specialist or clinical mi	crobiologist. Approva	als vali	d for 6 mo	nths where the treatment
remains appropriate and the patient is benefitting from the treatr	nent.			
MEBENDAZOLE – Only on a prescription				
Tab 100 mg		24	✓	De-Worm
Oral liq 100 mg per 5 ml	2.18	15 ml		
	(7.17)			Vermox
PRAZIQUANTEL				
Tab 600 mg		8	✓	Biltricide
Antibacterials				
a) For topical antibacterials, refer to DERMATOLOGICALS, page	71			
b) For anti-infective eve preparations, refer to SENSORY ORG.				
	-110, page 210			
Cephalosporins and Cephamycins				
CEFACLOR MONOHYDRATE				
Cap 250 mg	24.70	100	✓	Ranbaxy-Cefaclor
Grans for oral liq 125 mg per 5 ml – Wastage claimable – s				
rule 3.3.2 on page 13	3.53	100 m	✓	Ranbaxy-Cefaclor
CEFALEXIN				
Cap 250 mg		20		Cephalexin ABM
Cap 500 mg		20	~	Cephalexin ABM
Grans for oral liq 25 mg per ml – Wastage claimable – see				
3.3.2 on page 13		100 m		Cefalexin Sandoz
Note: Cefalexin grans for oral liq will not be funded in a		days	treatment	per dispensing.
Grans for oral liq 50 mg per ml – Wastage claimable – see				O falsada O sa da s
3.3.2 on page 13 Note: Cefalexin grans for oral lig will not be funded in a		100 m		Cefalexin Sandoz
o .		uays	liealment	per disperising.
CEFAZOLIN – Subsidy by endorsement			مريد مطلا امري	a substitute in an eleveral
Only if prescribed for dialysis or cellulitis in accordance with accordingly.	a DHB approved pro	locol a	ind the pre	scription is endorsed
Inj 500 mg vial	3 30	5	1	AFT
Inj 1 g vial		5		AFT
, .		5	-	<u></u>
CEFTRIAXONE – Subsidy by endorsement				
 a) Up to 5 inj available on a PSO b) Subsidised only if prescribed for a dialysis or cystic fibro 	sis nationt or the tree	tmont	of apport	ional or the treatment of
pelvic inflammatory disease, or the treatment of suspect				
and the prescription or PSO is endorsed accordingly.	ss morningnis in palle			aloan allorgy to periolilit,
Inj 500 mg vial		1	1	DEVA
Inj 1 g vial		1		DEVA
, ,	-			

‡ safety cap

 $\ensuremath{\boldsymbol{\ast}}$ Three months or six months, as applicable, dispensed all-at-once

(Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
CEFUROXIME AXETIL – Subsidy by endorsement Only if prescribed for prophylaxis of endocarditis and the presc Tab 250 mg		accoro 50		innat
Macrolides				
AZITHROMYCIN – Maximum of 5 days treatment per prescription; A maximum of 24 months of azithromycin treatment for non-cy- Authority.				
Tab 250 mg	9.00	30	🗸 🖌	po-Azithromycin
Tab 500 mg – Up to 8 tab available on a PSO Grans for oral lig 200 mg per 5 ml (40 mg per ml) – Wastage	1.05	2	✓ <u>A</u>	po-Azithromycin
claimable – see rule 3.3.2 on page 13	12.50	15 ml	✓ <u>Z</u>	ithromax

► SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are Unapproved Indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are Unapproved Indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are Unapproved Indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can	be waived by Sp	ecial Authori	ty see SA1131 on the next page
Tab 250 mg	3.98	14	 Apo-Clarithromycin
Grans for oral liq 250 mg per 5 ml – Wastage claimable – se	e		
rule 3.3.2 on page 13	23.12	50 ml	 Klacid

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

⇒SA1131 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria: Either:

1 Atypical mycobacterial infection; or

2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents. **Renewal — (Mycobacterial infections)** only from a respiratory specialist, infectious disease specialist or paediatrician.

Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN ETHYL SUCCINATE

Tab 400 mg	16.95	100	 E-Mycin
a) Up to 20 tab available on a PSO		100	
b) Up to 2 x the maximum PSO guantity for RFPP		no 17	
Grans for oral lig 200 mg per 5 ml		100 ml	 E-Mycin
		100 111	
a) Up to 300 ml available on a PSO		-	
b) Up to 2 x the maximum PSO quantity for RFPP	- see rule 5.2.6 on page	ge 17	
c) Wastage claimable – see rule 3.3.2 on page 13	o ==	100 1	/ - · ·
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	 E-Mycin
a) Up to 200 ml available on a PSO			
b) Wastage claimable – see rule 3.3.2 on page 13			
ERYTHROMYCIN LACTOBIONATE			
lnj 1 g		1	 Erythrocin IV
ERYTHROMYCIN STEARATE			-
Tab 250 mg – Up to 30 tab available on a PSO		100	
	(22.29)		ERA
Tab 500 mg		100	
	(44.58)		ERA
ROXITHROMYCIN			
Tab disp 50 mg		10	Rulide D
Restricted to children under 12 years of age.			
Tab 150 mg	7.48	50	Arrow-
5			Roxithromycin
Tab 300 mg	14.40	50	✓ Arrow-
			Roxithromycin

	Subsidy (Manufacturer's Pric \$	e) Per	Fully Subsidised	
Penicillins				
AMOXICILLIN				
Cap 250 mg	14.97	500	1	Apo-Amoxi
a) Up to 30 cap available on a PSO				-
b) Up to 10 x the maximum PSO quantity for RFPP – se	e rule 5.2.6 on pag	ge 17		
Cap 500 mg		500	✓	Apo-Amoxi
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP – se		ge 17		
Grans for oral liq 125 mg per 5 ml	1.20	100 ml	✓	Alphamox 125
 a) Up to 200 ml available on a PSO 				
b) Wastage claimable – see rule 3.3.2 on page 13				
Grans for oral liq 250 mg per 5 ml	1.31	100 ml	~	Alphamox 250
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP – se	e rule 5.2.6 on pag	ge 17		
c) Wastage claimable – see rule 3.3.2 on page 13	10.07	10		Ihlaman
Inj 250 mg vial Inj 500 mg vial		10 10		<u>Ibiamox</u> Ibiamox
Inj 1 g vial – Up to 5 inj available on a PSO		10		Ibiamox
		10	•	Indinox
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab	1.00	00		A
available on a PSO		20	•	Augmentin
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 r per ml		100 ml	1	Augmentin
a) Up to 200 ml available on a PSO		100 111	•	Augmentin
b) Wastage claimable – see rule 3.3.2 on page 13				
Grans for oral lig amoxicillin 50 mg with clavulanic acid 12.5 r	na			
per ml – Up to 200 ml available on a PSO		100 ml C	P 🗸	Curam
BENZATHINE BENZYLPENICILLIN			•	
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj				
available on a PSO	315.00	10	1	Bicillin LA
		10	•	
BENZYLPENICILLIN SODIUM [PENICILLIN G]	20 10.05	10		Conder
Inj 600 mg (1 million units) vial – Up to 5 inj available on a PS	50 10.55	10	•	Sandoz
FLUCLOXACILLIN	40.70	050	,	Otembler
Cap 250 mg – Up to 30 cap available on a PSO		250 500		Staphlex Staphlex
Cap 500 mg Grans for oral liq 25 mg per ml		100 ml		Staphlex AFT
a) Up to 200 ml available on a PSO		100 111	•	
b) Wastage claimable – see rule 3.3.2 on page 13				
Grans for oral lig 50 mg per ml		100 ml	1	AFT
a) Up to 200 ml available on a PSO			-	<u></u>
b) Wastage claimable – see rule 3.3.2 on page 13				
Inj 250 mg vial	9.00	10	1	Flucloxin
Inj 500 mg vial	9.40	10	-	Flucloxin
Inj 1 g vial – Up to 5 inj available on a PSO	5.22	5	✓	Flucil

	Subsidy (Manufacturer's Price \$) Sub Per	Fully sidised	Brand or Generic Manufacturer
PHENOXYMETHYLPENICILLIN (PENICILLIN V)				
Cap 250 mg – Up to 30 cap available on a PSO		50		ilicaine VK
Cap 500 mg	4.73	50	✓ <u>c</u>	ilicaine VK
 a) Up to 20 cap available on a PSO 				
b) Up to 2 x the maximum PSO quantity for RFPP – se				
Grans for oral liq 125 mg per 5 ml	1.48	100 ml	✓ <u>A</u>	FT
a) Up to 200 ml available on a PSO				
b) Wastage claimable – see rule 3.3.2 on page 13				
Grans for oral liq 250 mg per 5 ml	1.58	100 ml	✓ <u>A</u>	FT
a) Up to 300 ml available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP – se	e rule 5.2.6 on page	17		
c) Wastage claimable – see rule 3.3.2 on page 13				
ROCAINE PENICILLIN				
Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO.		5	✓ <u>C</u>	ilicaine
Tetracyclines				
OXYCYCLINE				
Fab 50 mg – Up to 30 tab available on a PSO	2.90	30		
	(6.00)		D	oxy-50
Tab 100 mg – Up to 30 tab available on a PSO	0.57	21		oxylin 100
	6.75	250	🗸 D	oxine
Doxylin 100 Tab 100 mg to be delisted 1 September 2018)				
IINOCYCLINE HYDROCHLORIDE				
Tab 50 mg – Additional subsidy by Special Authority see				
SA1355 below – Retail pharmacy	5.79	60		
· · ·	(12.05)		Μ	ino-tabs
€ Cap 100 mg		100		
	(52.04)		Μ	inomycin
»SA1355 Special Authority for Manufacturers Price				
itial application from any relevant practitioner. Approvals val	id without further ren	ewal unles	s notified	d where the patient ha
sacea.				•

TETRACYCLINE - Special Authority see SA1332 below - Re	tail pharmacy		
Cap 500 mg		30	 Tetracyclin
			Wolff S29

SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and

2 For use only in combination with bismuth as part of a quadruple therapy regimen.

‡ safety cap

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Other Antibiotics				
For topical antibiotics, refer to DERMATOLOGICALS, page 71 CIPROFLOXACIN Recommended for patients with any of the following: i) microbiologically confirmed and clinically significant ps ii) prostatitis; or iii) pyelonephritis; or iv) gonorrhoea.	eudomonas infection;	or		
Tab 250 mg – Up to 5 tab available on a PSO Tab 500 mg – Up to 5 tab available on a PSO Tab 750 mg CLINDAMYCIN Cap hydrochloride 150 mg – Maximum of 4 cap per	1.99	28 28 28	✓ (<u>Cipflox</u> <u>Cipflox</u> Cipflox
prescription; can be waived by endorsement - Retail pharmacy - Specialist Inj phosphate 150 mg per ml, 4 ml ampoule - Retail pharmacy-Specialist		16 10	-	Clindamycin ABM Dalacin C
COLISTIN SULPHOMETHATE – Retail pharmacy-Specialist – S Only if prescribed for dialysis or cystic fibrosis patient and th Inj 150 mg.	Subsidy by endorseme ne prescription is endor	ent	accordingly.	
GENTAMICIN SULPHATE Inj 10 mg per ml, 1 ml – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.		5 trac		lospira nd the prescription is
Inj 10 mg per ml, 2 ml – Subsidy by endorsement		25	-	APP Pharmaceuticals 529
 Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly. Inj 40 mg per ml, 2 ml ampoule – Subsidy by endorsement. Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly. 	6.00	10	✓ <u>I</u>	Pfizer
MOXIFLOXACIN – Special Authority see SA1358 below – Reta No patient co-payment payable Tab 400 mg		5	.	Avelox
SA1358 Special Authority for Subsidy Initial application — (Tuberculosis) only from a respiratory sp for applications meeting the following criteria: Either: 1 Both:		•	-	
 1.1 Active tuberculosis*; and 1.2 Any of the following: 1.2.1 Documented resistance to one or more first 1.2.2 Suspected resistance to one or more first-area with known resistance), as part of reg 	line medications (tuber			

	Subsidy	Fully	Brand or
(Man	ufacturer's Price)	Subsidised	Generic
	\$ Pe	er 🖌	Manufacturer

continued...

- 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
- 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
- 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*.

Note: Indications marked with * are Unapproved Indications (refer to Interpretations and Definitions).

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium*; and
- 2 Has tried and failed to clear infection using azithromycin; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are Unapproved Indications (refer to Interpretations and Definitions).

PAROMOMYCIN – Special Authority see SA1689 below – Retail pharmacy

Cap 250 mg...... 126.00 16 🖌 Humatin \$29

⇒SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

		al Authority see SA1328 below – Retail pharmacy	PYRIMETHAMINE – Special
 Daraprim S29 	30		Tab 25 mg
Daraprim S29	50	36.95	

⇒SA1328 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or

*Three months or six months, as applicable, dispensed all-at-once

3 For infants with congenital toxoplasmosis until 12 months of age.

SODIUM FUSIDATE [FUSIDIC ACID]

Prescriptions must be written by, or on the recommendation of, an infectious disease physician or a clinical microbiologist

SULFADIAZINE SODIUM	- Special Authority see SA1331 on the next page -	Retail pharmacy	
Tab 500 mg		56	✓ Wockhardt S29

	Subsidy (Manufacturer's Price	a) Si	Fully Ibsidised	Brand or Generic
	(Manulacialer 5 1 106 \$	Per		Manufacturer
SA1331 Special Authority for Subsidy tial application from any relevant practitioner. Approvals valid e following criteria: by of the following: 1 For the treatment of toxoplasmosis in patients with HIV for			ess notifie	d for applications meet
2 For pregnant patients for the term of the pregnancy; or3 For infants with congenital toxoplasmosis until 12 months of		,		
DBRAMYCIN				
Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement Only if prescribed for dialysis or cystic fibrosis patient and Solution for inhalation 60 mg per ml, 5 ml – Subsidy by		5 endorse		obramycin Mylan ıgly.
endorsement	2,200.00	56 dose	√ T	ОВІ
 b) Only if prescribed for a cystic fibrosis patient and the IMETHOPRIM 	prescription is endo	orsed acc	ordingly.	
Tab 300 mg - Up to 30 tab available on a PSO		50	🗸 Т	MP
RIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX/ Tab trimethoprim 80 mg and sulphamethoxazole 400 mg – U to 30 tab available on a PSO	lp	500	- л	risul
Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to 200 r available on a PSO	nl	100 ml	-	eprim
ANCOMYCIN – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient or for difficile following metronidazole failure and the prescription is Inj 500 mg vial	endorsed accordin			tment of Clostridium Iylan
Antifungals				
For topical antifungals refer to DERMATOLOGICALS, page 72 For topical antifungals refer to GENITO URINARY, page 85	2			
UCONAZOLE				
Cap 50 mg – Retail pharmacy-Specialist		28 1	_	lylan Iylan
 Cap 150 mg – Subsidy by endorsement a) Maximum of 1 cap per prescription; can be waived by b) Patient has vaginal candida albicans and the practition not recommended and the prescription is endorsed and Specialist. 	endorsement - Re ner considers that	tail pharm a topical i	nacy - Spe midazole	ecialist (used intra-vaginally) i
Cap 200 mg – Retail pharmacy-Specialist	5.08	28	🗸 N	lylan
Powder for oral suspension 10 mg per ml – Special Authority see SA1359 below – Retail pharmacy		35 ml	- ✓ D	iflucan S29 S29
	98.50		🗸 D	iflucan
Wastage claimable – see rule 3.3.2 on page 13				
•SA1359 Special Authority for Subsidy tial application — (Systemic candidiasis) from any relevant beeting the following criteria: oth:	practitioner. Appre	ovals valio	l for 6 wee	eks for applications

continued...

	Subsidy (Manufacturer's P)rice)	Cub	Fully sidised	Brand or Generic
	(Manulacturers F		Per	siuiseu ✓	Manufacturer
continued					
1 Patient requires prophylaxis for, or treatment of systemic ca	andidiasis; and				
2 Patient is unable to swallow capsules.					
Initial application - (Immunocompromised) from any relevant	practitioner. A	pprova	ls valid	for 6 mc	onths for applications
meeting the following criteria:					
All of the following:					
1 Patient is immunocompromised; and 2 Patient is at moderate to high risk of investive fungel infection	n; and				
 Patient is at moderate to high risk of invasive fungal infection Patient is unable to swallow capsules. 	n, anu				
Renewal — (Systemic candidiasis) from any relevant practitione	er Approvals v	alid fo	r 6 weel	s for an	plications meeting the
following criteria:			0 1000	to for up	plications meeting the
Both:					
1 Patient requires prophylaxis for, or treatment of systemic ca	andidiasis; and				
2 Patient is unable to swallow capsules.					
Renewal — (Immunocompromised) from any relevant practition	er. Approvals	valid fo	or 6 mor	ths for a	applications meeting the
following criteria:					
All of the following:					
 Patient remains immunocompromised; and Patient remains at moderate to high risk of invasive fungal 	infection: and				
3 Patient is unable to swallow capsules.					
ITRACONAZOLE					
Cap 100 mg – Subsidy by endorsement	2 79		15	✓ lt	razole
Funded for tinea vesicolor where topical treatment has no					<u> </u>
mycology, or for tinea unguium where terbinafine has not					
terbinafine and diagnosis has been confirmed by mycolog		cription	is endo	orsed ac	cordingly.
Can be waived by endorsement - Retail pharmacy - Spec					
Specialist must be an infectious disease physician, clinica Oral lig 10 mg per ml – Special Authority see SA1322 below -	•	, clinic	ai immu	nologist	or dermatologist.
Retail pharmacy		150	ml OP	✓ s	poranox
► SA1322 Special Authority for Subsidy		100		. 0	porunox
Initial application only from an infectious disease specialist, clinic	al microbiologis	st. clini	cal imm	unologis	st or any relevant
practitioner on the recommendation of a infectious disease physici					
valid for 6 months where the patient has a congenital immune defi			•		0 11
Renewal from any relevant practitioner. Approvals valid for 6 mor	ths where the t	treatme	ent rema	ains app	ropriate and the patient is
benefitting from the treatment.					
KETOCONAZOLE					
Tab 200 mg – PCT – Retail pharmacy-Specialist – Subsidy by endorsement					
endorsement			30		ink Healthcare S29
Prescriptions must be written by, or on the recommendation	on of an oncolo	aiet		♥ N	izoral S29
NYSTATIN		giot			
Tab 500.000 u	14 16		50		
	(17.09)			Ν	ilstat
Cap 500,000 u		!	50		
	(15.47)			Ν	ilstat
POSACONAZOLE - Special Authority see SA1285 on the next pa		armacy	'		
Tab modified-release 100 mg			24		oxafil
Oral liq 40 mg per ml	761.13	105	ml OP	✓ N	oxafil

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per		Manufacturer

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE

* Tab 250 mg – For terbinafine oral liquid formulation refer,		
page 2241.33	14	 Deolate
VORICONAZOLE - Special Authority see SA1273 below - Retail pharmacy		
Tab 50 mg	56	 Vttack
Tab 200 mg	56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage claimable		
- see rule 3.3.2 on page 131,156.32	70 ml	 Vfend

➡SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

Patient is immunocompromised; and

- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

	Subsidy (Manufacturer's Pr \$	ice) Sub Per	Fully Brand or sidised Generic ✓ Manufacturer	
Antimalarials				
PRIMAQUINE PHOSPHATE – Special Authority see SA1684		macy 56	Primacin S29	
Tab 7.5 mg				applicatior
1 The patient has vivax or ovale malaria; and 2 Primaquine is to be given for a maximum of 21 days. Renewal only from an infectious disease specialist or clinical the following criteria: toth:	microbiologist. Appr	ovals valid fo	r 1 month for applicatio	ns meetin
 The patient has relapsed vivax or ovale malaria; and Primaquine is to be given for a maximum of 21 days. 				
Antiparasitics				
Antiprotozoals				
UININE SULPHATE Tab 300 mg ‡ Safety cap for extemporaneously compounded oral li		500	✔ Q 300	
Antitrichomonal Agents				
IETRONIDAZOLE Tab 200 mg – Up to 30 tab available on a PSO Tab 400 mg Oral liq benzoate 200 mg per 5 ml Suppos 500 mg		100 100 100 ml 10	 ✓ Trichozole ✓ Trichozole ✓ Flagyl-S ✓ Flagyl 	
RNIDAZOLE Tab 500 mg	23.00	10	✓ Arrow-Ornidaze	ole
Antituberculotics and Antileprotics				
ote: There is no co-payment charge for all pharmaceuticals nmigration status. LOFAZIMINE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend darmatologist				
dermatologist. ⊱ Cap 50 mg		100	✓ Lamprene S29	
YCLOSERINE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend	dation of, an infectiou	us disease pł	nysician, clinical microb	iologist or
respiratory physician.				

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy	<u> </u>	Fully	Brand or
	(Manufacturer's Price) \$	Sul Per	osidised	Generic Manufacturer
	φ	Fei	•	Manulaciulei
DAPSONE – Retail pharmacy-Specialist				
 a) No patient co-payment payable 				
b) Prescriptions must be written by, or on the recommend	ation of, an infectious of	lisease p	hysician	, clinical microbiologist or
dermatologist				
Tab 25 mg		100		Dapsone
Tab 100 mg		100	✓	Dapsone
ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Specia	alist			
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommend	ation of, an infectious of	lisease p	hysician	, clinical microbiologist or
respiratory physician			,	,
Tab 100 mg		56	 Image: A second s	Myambutol S29
C C	85.73	100	 Image: A second s	EMB Fatol S29
Tab 400 mg		56		Myambutol S29
0		00		inyanibator 🛥
ISONIAZID – Retail pharmacy-Specialist				
a) No patient co-payment payable	allow of an internal mo			and the state of the first
b) Prescriptions must be written by, or on the recommend		dicine ph	ysıcıan,	paediatrician, clinical
microbiologist, dermatologist or public health physician		100		DOM
* Tab 100 mg		100	-	PSM Diffinish
* Tab 100 mg with rifampicin 150 mg		100		Rifinah Rifinah
* Tab 150 mg with rifampicin 300 mg		100	•	Rifinah
PARA-AMINO SALICYLIC ACID – Retail pharmacy-Specialist				
 a) No patient co-payment payable 				
b) Specialist must be an infectious disease specialist, clin	ical microbiologist or re	spiratory	speciali	ist.
Grans for oral liq 4 g sachet		30	 Image: A second s	Paser S29
PROTIONAMIDE – Retail pharmacy-Specialist				
a) No patient co-payment payable				
 b) Specialist must be an infectious disease specialist, clin 	ical microbiologist or re	spiratory	speciali	ist.
Tab 250 mg		100		Peteha S29
-		100	•	Cicila
PYRAZINAMIDE – Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommend	ation of, an infectious of	lisease p	hysician	, clinical microbiologist or
respiratory physician				
* Tab 500 mg – For pyrazinamide oral liquid formulation ref	,	400		
page 224		100		AFT-Pyrazinamide
			•	AFT-Pyrazinamide
				S29 S29
RIFABUTIN – Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommend	ation of, an infectious of	lisease p	hysician	, respiratory physician or
gastroenterologist		•		· · · · ·
* Cap 150 mg – For rifabutin oral liquid formulation refer,				
page 224		30	✓ [Mycobutin

	Subsidy (Manufacturer's Pri \$	ce) Sub Per	sidised G	rand or ieneric Ianufacturer
RIFAMPICIN – Subsidy by endorsement				
 a) No patient co-payment payable b) For confirmed recurrent Staphylococcus aureus infection antimicrobial based on susceptibilities and the prescription Retail pharmacy - Specialist. Specialist must be an in paediatrician, or public health physician. 	ption is endorsed acc nternal medicine physi	ordingly; car	h be waived	by endorsement -
卷 Сар 150 mg	55.75	100	✓ Rifa	
* Cap 300 mg		100	✓ <u>Rifa</u>	
 Vral liq 100 mg per 5 ml 		60 ml	✓ <u>Rifa</u>	din
Antivirals				
For eye preparations refer to Eye Preparations, Anti-Infective	Preparations, page 2	16		
Hepatitis B Treatment				
ADEFOVIR DIPIVOXIL – Special Authority see SA0829 belo Tab 10 mg		30	🗸 Нер	sera
SA0829 Special Authority for Subsidy Initial application only from a gastroenterologist or infectious meeting the following criteria: All of the following:	s disease specialist.	Approvals va	lid for 1 yea	r for applications
 Patient has confirmed Hepatitis B infection (HBsAg+); Documented resistance to lamivudine, defined as: Patient has raised serum ALT (> 1 × ULN); and Patient has HBV DNA greater than 100,000 copies per 		fold or highe	r over nadir	; and
4 Detection of M204I or M204V mutation; and 5 Either:				
5.1 Both:				
5.1.1 Patient is cirrhotic; and5.1.2 adefovir dipivoxil to be used in combina5.2 Both;	tion with lamivudine; o	or		
5.2.1 Patient is not cirrhotic; and 5.2.2 adefovir dipivoxil to be used as monoth	erany			
Renewal only from a gastroenterologist or infectious disease treating physician, treatment remains appropriate and patient Notes: Lamivudine should be added to adefovir dipivoxil if a	specialist. Approvals is benefiting from trea	atment.		
 i) raised serum ALT (> 1 × ULN); and ii) HBV DNA greater than 100,000 copies per mL, or vira iii) Detection of NOCCT or 42017 (contaction) 	l load 10 fold or highe	r over nadir;	and	
iii) Detection of N236T or A181T/V mutation. Adefovir dipivoxil should be stopped 6 months following HBe/ commencing adefovir dipivoxil.	Ag seroconversion for	patients who	o were HBe	Ag+ prior to
The recommended dose of adefovir dipivoxil is no more than n patients with renal insufficiency adefovir dipivoxil dose sho Adefovir dipivoxil should be avoided in pregnant women and	uld be reduced in acco	ordance with	the datash	eet guidelines.
ENTECAVIR * Tab 0.5 mg		30	🗸 Bara	

\$\$ safety cap
\$\$ Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Pric \$	e) Subs Per	Fully idised	Brand or Generic Manufacturer
	•	rei		Manulacturer
LAMIVUDINE – Special Authority see SA1685 below – Retail			<i>.</i> -	
Tab 100 mg		28	-	etlam
	6.00		✓ Z	•••••
Oral liq 5 mg per ml		240 ml OP	✓ Z	effix
SA1685 Special Authority for Subsidy				
nitial application only from a relevant specialist or medical pr	actitioner on the reco	mmendation	of a re	levant specialist.
Approvals valid for 1 year where used for the treatment or prev				
Renewal from any relevant practitioner. Approvals valid for 2 y	years where used for	the treatmer	nt or pre	evention of hepatitis B
TENOFOVIR DISOPROXIL				
Tenofovir disoproxil prescribed under endorsement for the		ncluded in the	e count	of up to 4 subsidised
antiretrovirals for the purposes of Special Authority SA165				
* Tab 245 mg (300 mg as a fumarate)		30	-	iread
* Tab 245 mg (300.6 mg as a succinate)		30	✓ T	enofovir Disoproxil
				Teva
Herpesvirus Treatments				
•				
ACICLOVIR				
* Tab dispersible 200 mg		25	✓ L	
* Tab dispersible 400 mg		56	✓ L	<u> </u>
* Tab dispersible 800 mg	5.98	35	✓ L	ovir
VALACICLOVIR				
Tab 500 mg	6.42	30	🗸 V	aclovir
Tab 1,000 mg		30	✓ V	aclovir
VALGANCICLOVIR - Special Authority see SA1404 below - F			_	
Tab 450 mg		60	🗸 V	alcyte
		00		aloyto

► SA1404 Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
- 2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Patient has undergone a lung transplant; and

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
 \$	Per	1	

continued...

2 Either:

- 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
- 2.2 The recipient is cytomegalovirus positive.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

LEDIPASVIR WITH SOFOSBUVIR – Special Authority see SA1605 below – [Xpharm] No patient co-payment payable
Tab 90 mg with sofosbuvir 400 mg 24,363.46 28 🖌 Harvoni
■SA1605 Special Authority for Subsidy
Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)
Notes: By application to the Hepatitis C Treatment Panel (HepCTP).
Applications will be considered by HepCTP and approved subject to confirmation of eligibility.
Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz/hepatitis-c-treatments or:
The Coordinator, Hepatitis C Treatment Panel
PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,
Email: hepcpanel@pharmac.govt.nz
PARITAPREVIR, RITONAVIR AND OMBITASVIR WITH DASABUVIR – [Xpharm]
a) No patient co-payment payable
b) Note – Supply of treatment is via PHARMAC's approved direct distribution supply. Application details for accessing
treatment may be obtained from PHARMAC's website http://www.pharmac.govt.nz/hepatitis-c-treatments
Tab 75 mg with ritonavir 50 mg, and ombitasvir 12.5 mg (56),
with dasabuvir tab 250 mg (56)
PARITAPREVIR, RITONAVIR AND OMBITASVIR WITH DASABUVIR AND RIBAVIRIN – [Xpharm]
a) No patient co-payment payable
b) Note – Supply of treatment is via PHARMAC's approved direct distribution supply. Application details for accessing
treatment may be obtained from PHARMAC's website http://www.pharmac.govt.nz/hepatitis-c-treatments
Tab 75 mg with ritonavir 50 mg, and ombitasvir 12.5 mg (56)
with dasabuvir tab 250 mg (56) and ribavirin tab 200 mg
(168)16,500.00 1 OP ✓ Viekira Pak-RBV

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	ubsidised	Generic
\$	Per	~	Manufacturer

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL FUMARATE - Subsidy by endorsement; can be waived by Special Authority see SA1714 below.

Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil fumarate is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note:

Emtricitabine with tenofovir disoproxil fumarate prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 115

There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the PHARMAC website.

Tab 200 mg with tenofovir disoproxil fumarate 300 mg......838.20 30 🗸 Truvada

➡SA1714 Special Authority for Waiver of Rule

Initial application only from a named specialist or medical practitioner on the recommendation of a named specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tested HIV negative; and
- 2 Either:
 - 2.1 All of the following:
 - 2.1.1 Patient is male or transgender; and
 - 2.1.2 Patient has sex with men; and
 - 2.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 2.1.4 Any of the following:
 - 2.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 2.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 2.1.4.3 Patient has used methamphetamine in the last three months; or
 - 2.2 All of the following:
 - 2.2.1 Patient has a regular partner who has HIV infection; and
 - 2.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 2.2.3 Condoms have not been consistently used.

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis; and
- 2 Patient has undergone testing for HIV, syphilis, and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative; and
- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	idised	Generic
\$	Per	1	Manufacturer

continued...

- 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
- 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
- 6.1.4.3 Patient has used methamphetamine in the last three months; or
- 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Antiretrovirals

⇒SA1651 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil fumarate prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals.

Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil fumarate prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals.

Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria: Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil fumarate prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals.

Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose

continued...

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	1	Manufacturer	

continued...

ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil fumarate prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals.

Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA1651 on the previous page - Retail pharmacy

Tab 50 mg		30	✓ Stocrin S29
Tab 200 mg		90	 Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
Oral liq 30 mg per ml	145.79	180 ml OP	 Stocrin S29
ETRAVIRINE - Special Authority see SA1651 on th	e previous page – Retail pha	rmacy	
Tab 200 mg	770.00	60	 Intelence
NEVIRAPINE - Special Authority see SA1651 on th	e previous page – Retail pha	rmacy	
Tab 200 mg		60	Nevirapine
			<u>Alphapharm</u>
Oral suspension 10 mg per ml		240 ml	 Viramune
			Suspension

Nucleosides Reverse Transcriptase Inhibitors

Tab	VIR SULPHATE – Special Authority see SA1651 on 300 mg I liq 20 mg per ml	229.00	letail pharma 60 240 ml OP	acy ✓ Ziagen ✓ Ziagen
ABACA Note	VIR SULPHATE WITH LAMIVUDINE – Special Auth e: abacavir with lamivudine (combination tablets) co -retroviral Special Authority.	ority see SA1651 on the		
Tab	600 mg with lamivudine 300 mg		30	 Kivexa
previous Note purp	ENZ WITH EMTRICITABINE AND TENOFOVIR DIS page – Retail pharmacy e: Efavirenz with emtricitabine and tenofovir disopro poses of the anti-retroviral Special Authority	xil fumarate counts as		,
	600 mg with emtricitabine 200 mg and tenofovir dise	•		4 • • • • •
	fumarate 300 mg	1,313.19	30	 Atripla
116	✓ fully subsidised	S29 Unappro	oved medicine	e supplied under Section 29
110	[HP4] refer page 4	Sole Subsidis	ed Supply	

	Subsidy (Manufacturer's F \$	Price) Subsi Per	Fully Brand or idised Generic ✓ Manufacturer
EMTRICITABINE – Special Authority see SA1651 on page 115 - Cap 200 mg		у 30	✓ Emtriva
LAMIVUDINE – Special Authority see SA1651 on page 115 – Re Tab 150 mg		60	 Lamivudine Alphapharm
Oral liq 10 mg per ml	102.50	240 ml OP	✓ 3TC
ZIDOVUDINE [AZT] – Special Authority see SA1651 on page 11 Cap 100 mg Oral liq 10 mg per ml		nacy 100 200 ml OP	 ✓ <u>Retrovir</u> ✓ <u>Retrovir</u>
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority see Note: zidovudine [AZT] with lamivudine (combination tablets the anti-retroviral Special Authority.	s) counts as two		edications for the purposes of
Tab 300 mg with lamivudine 150 mg		60	 Alphapharm
Protease Inhibitors			
ATAZANAVIR SULPHATE – Special Authority see SA1651 on p Cap 150 mg Cap 200 mg	568.34	pharmacy 60 60	 ✓ Reyataz ✓ Reyataz
DARUNAVIR – Special Authority see SA1651 on page 115 – Re Tab 400 mg Tab 600 mg		60 60	✓ <u>Prezista</u> ✓ Prezista
LOPINAVIR WITH RITONAVIR – Special Authority see SA1651 Tab 100 mg with ritonavir 25 mg Tab 200 mg with ritonavir 50 mg Oral lig 80 mg with ritonavir 20 mg per ml		Retail pharmacy 60 120 300 ml OP	 ✓ Kaletra ✓ Kaletra ✓ Kaletra
RITONAVIR – Special Authority see SA1651 on page 115 – Reta Tab 100 mg Oral liq 80 mg per ml	ail pharmacy 43.31	30 90 ml OP	✓ Norvir✓ Norvir
Strand Transfer Inhibitors			
DOLUTEGRAVIR – Special Authority see SA1651 on page 115 - Tab 50 mg	•	су 30	 Tivicay
RALTEGRAVIR POTASSIUM – Special Authority see SA1651 o Tab 400 mg		tail pharmacy 60	✓ Isentress

Immune Modulators

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects. Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

1) Diagnosis

Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a

continued...

‡ safety cap

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- supplementary RIBA test; or
- PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
- Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Exclusion Criteria

- 1) Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia (< 2.0 × 10⁹) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

Dosage

Ρ

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

INTERFERON ALFA-2A - PCT - Retail pharmacy-Specialist

- a) See prescribing guideline on the previous page
- b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician or ophthalmologist
- 1 ✓ Roferon-A

INTERFERON ALFA-2B - PCT - Retail pharmacy-Specialist

a) See prescribing guideline on the previous page

b)	Prescriptions must be written by	, or on the	recommendation of,	, an internal	medicine physician	or ophthalmologist
----	----------------------------------	-------------	--------------------	---------------	--------------------	--------------------

b) Prescriptions must be written by, or on the recommendation	ation of, an internal	medicine phy	sician or opninalmologist
Inj 18 m iu, 1.2 ml multidose pen	206.71	1	 Intron-A
Inj 30 m iu, 1.2 ml multidose pen		1	 Intron-A
Inj 60 m iu, 1.2 ml multidose pen	689.04	1	 Intron-A
PEGYLATED INTERFERON ALFA-2A – Special Authority see	SA1400 below - F	Retail pharma	су
See prescribing guideline on the previous page			
Inj 180 mcg prefilled syringe	500.00	4	 Pegasys
Inj 135 mcg prefilled syringe × 4 with ribavirin tab 200 mg >	(
168	1,975.00	1 OP	Pegasys RBV
			Combination Pack
Inj 180 mcg prefilled syringe × 4 with ribavirin tab 200 mg >	(
112	1,159.84	1 OP	Pegasys RBV
			Combination Pack
Inj 180 mcg prefilled syringe × 4 with ribavirin tab 200 mg >	<		
168	1,290.00	1 OP	Pegasys RBV

SA1400 Special Authority for Subsidy

Initial application - (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria: Both:

1 Any of the following:

- 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
- 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
- 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant: and

Combination Pack

Su	ubsidy	Fully	Brand or
(Manufac	cturer's Price) Subsid	lised	Generic
	\$ Per	1	Manufacturer

continued...

2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application - (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a

gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and

continued...

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once if endorsed "certified exemption" by the prescriber or pharmacist.

Three months supply may be dispensed at one time

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic
(Maintadaio) + 160)	Per	✓	Manufacturer

continued...

- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Notes:

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

Urinary Tract Infections			
HEXAMINE HIPPURATE			
* Tab 1 g1	18.40	100	
(4	40.01)		Hiprex
NITROFURANTOIN			
 Tab 50 mg – For nitrofurantoin oral liquid formulation refer, 			
page 224	22.20	100 🖌	 Nifuran
* Tab 100 mg	37.50	100 🖌	Nifuran
NORFLOXACIN			
Tab 400 mg - Subsidy by endorsement13	35.00	100 🖌	Arrow-Norfloxacin

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

(Manufacturer's Price) Subsidiated Per Generic Manufacturer Anticholinesterases EOSTIGMINE METILSULFATE 98.00 50 ✓ AstraZeneca VRIDOSTIGMINE BEROMIDE 98.00 50 ✓ Mestinon Tab 60 ng		Subsidy	Fi	Illy Brand or
s Per Manufacturer Anticholinesterases EOSTIGMINE METILSULFATE 98.00 50 ✓ AstraZenece VPIDOSTIGMINE BROMIDE 98.00 50 ✓ Mestinon Non-Steroidal Anti-Inflammatory Drugs Mestinon ICLOFENAC SODIUM 1.30 50 ✓ Voltaren D Tab 60 mg 1.50 20 ✓ Voltaren D Tab 50 mg dispersible 1.50 50 ✓ Voltaren D Tab 10 ng-acting 75 mg 1.52 500 ✓ App-Diclo SR Suppos 25 mg 2.44 10 ✓ Voltaren Suppos 25 mg 11.71 1.000 ✓ Relieve Tab long-acting 800 mg 7.99 30 ✓ Perpaed TDPROFEN Fenpaed Fording 20 mg ✓ Notaren Suppos 10 mg 1.2.07 28 Oruvail SR Fenpaed				
EOSTIGMINE METILSULFATE Inj 2.5 mg per ml, 1 ml ampoule 98.00 50 ✓ AstraZeneca VRIDOSTIGMINE BRONDE Tab 60 mg 42.79 100 ✓ Mestinon Von-Steroidal Anti-Inflammatory Drugs 1.30 50 ✓ Diclofenac Sandoz ICLOFENAC SODIUM 1.30 50 ✓ Voltaren D Tab E0 50 mg dispersible 1.50 20 ✓ Voltaren D Tab E0 50 mg 1.52 500 ✓ Appo-Diclo SR Tab long-acting 75 mg 2.44 10 ✓ Voltaren Suppos 21.5 mg 2.04 10 ✓ Voltaren Suppos 50 mg -Up to 5 inj available on a PSO 4.22 10 ✓ Voltaren Suppos 50 mg -Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 50 mg -Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 50 mg -Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 50 mg -Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 50 mg -Up to 10 supp available on a PSO 2.39				
EOSTIGMINE METILSULFATE Inj 2.5 mg per ml, 1 ml ampoule 98.00 50 ✓ AstraZeneca VRIDOSTIGMINE BRONDE Tab 60 mg 42.79 100 ✓ Mestinon Von-Steroidal Anti-Inflammatory Drugs 1.30 50 ✓ Diclofenac Sandoz ICLOFENAC SODIUM 1.30 50 ✓ Voltaren D Tab E0 50 mg dispersible 1.50 20 ✓ Voltaren D Tab E0 50 mg 1.52 500 ✓ Appo-Diclo SR Tab long-acting 75 mg 2.44 10 ✓ Voltaren Suppos 21.5 mg 2.04 10 ✓ Voltaren Suppos 50 mg -Up to 5 inj available on a PSO 4.22 10 ✓ Voltaren Suppos 50 mg -Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 50 mg -Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 50 mg -Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 50 mg -Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 50 mg -Up to 10 supp available on a PSO 2.39				
Inj 2.5 mg per ml, 1 ml ampoule	Anticholinesterases			
Inj 2.5 mg per ml, 1 ml ampoule	NEOSTIGMINE METII SUI FATE			
YRIDOSTIGMINE BROMIDE Tab 60 ng			50	AstraZeneca
Tab 60 mg				
Non-Steroidal Anti-Inflammatory Drugs ICLOFENAC SODIUM 130 50 ✓ Diclofenac Sandoz Tab EC 25 mg 1.50 20 ✓ Voltaren D Tab Go ng dispersible 1.50 20 ✓ Voltaren D Tab Iong-acting 75 mg 1.520 500 ✓ Apc-Diclo SR Tab Iong-acting 75 mg 26.20 500 ✓ Apc-Diclo SR Inj25 mg pernl, 3 ml ampoule – Up to 5 inj available on a PSO 4.22 10 ✓ Voltaren Suppos 25 mg _0 to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 100 mg _0 to 10 supp available on a PSO 4.22 10 ✓ Voltaren UPROFEN		42.79	100	Mestinon
CLOFENAC SODIUM 1.30 50 ✓ Diclofenac Sandoz Tab EC 50 mg 1.50 20 ✓ Voltaren D Tab EC 50 mg 1.520 500 ✓ Apo-Diclo SR Tab Long-acting 100 mg 26.20 500 ✓ Apo-Diclo SR Tab Long-acting 100 mg 26.20 500 ✓ Apo-Diclo SR Suppos 12.5 mg 2.04 10 ✓ Voltaren Suppos 25 mg 2.04 10 ✓ Voltaren Suppos 12.5 mg 2.04 10 ✓ Voltaren Suppos 20 mg -Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 100 mg -Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren UPROFEN -Tab long-acting 800 mg 7.99 30 ✓ Brufen SR ± Cap long-acting 200 mg 12.07 28 ✓ Oruvail SR EFENAMIC ACID - Cap Jong-acting 200 mg Ponstan ± Cap long-acting 750 mg 18.91 250 ✓ Norliam 500 * Tab Jong-acting 750 mg 18.91 250 ✓ Norliam 500 * Tab long-acting 750 mg 6.53 28 ✓ Maprosyn SR 750<	_ ·			
Tab EC 25 mg 1.30 50 ✓ Diclofenac Sandoz Tab EC 50 mg 1.50 20 ✓ Voltaren D Tab EC 50 mg 1.00 50 ✓ Diclofenac Sandoz Tab Long-acting 75 mg 15.20 500 ✓ Apo-Diclo SR Tab Long-acting 100 mg 26.20 500 ✓ Apo-Diclo SR Tab Long-acting 100 mg 26.20 500 ✓ Apo-Diclo SR Suppos 25 mg .204 10 ✓ Voltaren Suppos 25 mg .204 10 ✓ Voltaren Suppos 25 mg .204 10 ✓ Voltaren Suppos 100 mg .700 10 ✓ Voltaren Suppos 100 mg .700 10 ✓ Voltaren UPROFEN Tab Log acting 200 mg .7.99 30 ✓ Brufen SR ± Carl ling 20 mg per ml 2.39 200 ml ✓ Fenpaed ETOPROFEN .20 .00 Ponstan 0.50 20 (5.60) Ponstan .50 20 Ponstan 0.50 20 Noflam 550 * Tab 250 mg .12.57 .50 .40 .40 .40 <	Non-Steroidal Anti-Inflammatory Drugs			
Tab EC 25 mg 1.30 50 ✓ Diclofenac Sandoz Tab EC 50 mg 1.50 20 ✓ Voltaren D Tab EC 50 mg 1.00 50 ✓ Diclofenac Sandoz Tab Long-acting 75 mg 15.20 500 ✓ Apo-Diclo SR Tab Long-acting 100 mg 26.20 500 ✓ Apo-Diclo SR Tab Long-acting 100 mg 26.20 500 ✓ Apo-Diclo SR Suppos 25 mg .204 10 ✓ Voltaren Suppos 25 mg .204 10 ✓ Voltaren Suppos 25 mg .204 10 ✓ Voltaren Suppos 100 mg .700 10 ✓ Voltaren Suppos 100 mg .700 10 ✓ Voltaren UPROFEN Tab Log acting 200 mg .7.99 30 ✓ Brufen SR ± Carl ling 20 mg per ml 2.39 200 ml ✓ Fenpaed ETOPROFEN .20 .00 Ponstan 0.50 20 (5.60) Ponstan .50 20 Ponstan 0.50 20 Noflam 550 * Tab 250 mg .12.57 .50 .40 .40 .40 <				
Tab 50 mg dispersible 1.50 20 ✓ Voltaren D Tab LC 50 mg 1.00 50 Diclofenac Sandoz Tab long-acting 75 mg 15.20 500 ✓ Apo-Diclo SR Yab 50 mg dispersible 26.20 500 ✓ Apo-Diclo SR Yab 10 mg dispersible 15.20 500 ✓ Voltaren Suppos 12.5 mg 2.04 10 ✓ Voltaren Suppos 25 mg 2.04 10 ✓ Voltaren Suppos 12.5 mg 2.04 10 ✓ Voltaren Suppos 100 mg - Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 100 mg - Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 100 mg - Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren UPROFEN - Tab long-acting 800 mg 7.99 30 ✓ Brufen SR # Coral log 200 mg 12.07 28 ✓ Oruvail SR EFENAMIC ACID - Song 900 ml ✓ Fenpaed ETab long-acting 750 mg .60 28 ✓ Noftam 250 * Tab long-acting 750 mg .60 28		1.30	50	Diclofenac Sandoz
1: Tab EC 50 mg 1.00 50 ✓ Diclofenac Sandoz 1: Tab long-acting 75 mg 15.20 500 ✓ Apo-Diclo SR 1: Tab long-acting 100 mg 26.20 500 ✓ Apo-Diclo SR 1: Suppos 12.5 mg 2.04 10 ✓ Voltaren Suppos 52 mg 2.44 10 ✓ Voltaren Suppos 50 mg - Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 50 mg - Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 50 mg - Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 20 mg	5			
Tab long-acting 75 mg. 15.20 500 ✓ Apo-Diclo SR Tab long-acting 100 mg. 26.20 500 ✓ Apo-Diclo SR Tab long-acting 100 mg. 204 10 ✓ Voltaren Suppos 12.5 mg. 2.44 10 ✓ Voltaren Suppos 25 mg. 2.44 10 ✓ Voltaren Suppos 50 mg – Up to 10 supp available on a PSO. 4.22 10 ✓ Voltaren Suppos 100 mg. 7.00 10 ✓ Voltaren UPROFEN 11.71 1,000 ✓ Relieve Tab long-acting 800 mg. 7.99 30 ✓ Brufen SR #2 Oral liq 20 mg per ml. 2.39 200 ml ✓ Fenpaed ETOPROFEN E Cap long-acting 200 mg. 12.07 28 ✓ Oruvail SR EFENAMIC ACID 1.25 50 (9.16) Ponstan 0.50 20 (5.60) Ponstan 0.50 20 1 Tab 500 mg. 18.06 500 ✓ Noflam 550 Maprosyn SR 750 1 Tab long-acting 750 mg. 5.60 28 ✓ Naprosyn SR 1500 1 Tab long-acting 1 g. 6.53 28 ✓				_
Tab long-acting 100 mg. 26.20 500 ✓ Appo-Diclo SR Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a PSO 13.20 5 ✓ Voltaren Suppos 125 mg.	· _ · · ·			
Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a PSO13.20 5 ✓ Voltaren Suppos 12.5 mg 2.04 10 ✓ Voltaren Suppos 25 mg 2.44 10 ✓ Voltaren Suppos 50 mg – Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 100 mg - Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren BUPROFEN - Tab long-acting 800 mg 7.00 10 ✓ Voltaren Tab long-acting 800 mg 7.99 30 ✓ Brufen SR ± Carl lig 20 mg per ml 2.39 200 ml ✓ Fenpaed ETOPROFEN - Cap long-acting 200 mg 12.07 28 ✓ Oruvail SR EFENAMIC ACID - Fenpaed - Noflam 250 Ponstan 0.50 20 (5.60) Ponstan - Noflam 500 * Tab 500 mg - 18.06 500 ✓ Noflam 500 - Noflam 500 * Tab 10ng-acting 750 mg - 5.60 28 ✓ Naprosyn SR 750 - Tab 100 mg -				
Suppos 12.5 mg 2.04 10 ✓ Voltaren Suppos 25 mg 2.44 10 ✓ Voltaren Suppos 100 mg				
Suppos 25 mg 2.44 10 ✓ Voltaren Suppos 50 mg - Up to 10 supp available on a PSO 4.22 10 ✓ Voltaren Suppos 100 mg			10	✓ Voltaren
 Suppos 50 mg - Up to 10 supp available on a PSO			10	✓ Voltaren
HUPROFEN * Tab 200 mg 11.71 1,000 * Relieve * Tab long-acting 800 mg 7.99 30 * Brufen SR ** Oral liq 20 mg per ml 2.39 200 ml * Fenpaed ETOPROFEN 12.07 28 * Oruvail SR EFENAMIC ACID 12.05 20 ponstan 0.50 20 90 mstan 0.50 20 (9.16) Ponstan 0.50 20 ponstan APROXEN 18.91 250 * Noflam 250 * Noflam 500 * Tab 500 mg 18.91 250 * Naprosyn SR 750 * Tab long-acting 750 mg 6.53 28 * Naprosyn SR 1000 ULINDAC * * Tab 200 mg 8.55 50 * Aclin * Tab 200 mg 15.10 50 * Aclin * Aclin ENOXICAM * Tab 20 mg vial 9.95 1 * AFT NSAIDS Other <td< td=""><td></td><td></td><td>10</td><td> Voltaren </td></td<>			10	 Voltaren
Tab 200 mg 11.71 1,000 Relieve Brufen SR Brufen SR Brufen SR Brufen SR Fenpaed Fonstan 0.50 20 (5.60) Ponstan Ponstan 0.50 20 (5.60) Ponstan Moliam 250 Moliam 500 Moliam 500 Moliam 500 Moliam 500	* Suppos 100 mg	7.00	10	 Voltaren
Tab 200 mg 11.71 1,000 Relieve Brufen SR Brufen SR Brufen SR Brufen SR Fenpaed Fonstan 0.50 20 (5.60) Ponstan Ponstan 0.50 20 (5.60) Ponstan Moliam 250 Moliam 500 Moliam 500 Moliam 500 Moliam 500	BUPROFEN			
: Tab long-acting 800 mg. 7.99 30 ✓ Bruten SR : Cap long-acting 200 mg 2.39 200 ml ✓ Fenpaed ETOPROFEN : 2.39 200 ml ✓ Fenpaed ETOPROFEN : 2.39 200 ml ✓ Fenpaed ETENAMIC ACID : 2.25 50 (9.16) Ponstan 0.50 20 (5.60) Ponstan 0.50 20 : Tab 250 mg 18.06 500 ✓ Noflam 250 • Noflam 250 : Tab 500 mg 18.01 250 ✓ Naprosyn SR 750 : Tab long-acting 750 mg : 6.53 28 ✓ Naprosyn SR 1000 ULINDAC : : : 7ab 20 mg : 4.clin : Tab 200 mg : 15.10 50 ✓ Aclin ENOXICAM : : 7ab 20 mg vial : 9.95 1 ✓ AFT NSAIDS Other : : : 2.30 : 30 ✓ Celecoxib Pfizer : Cap 200 mg : 2.30 : 30 : Celecoxib Pfizer Celecoxib Pfizer			1.000	Relieve
#‡ Oral liq 20 mg per ml 2.39 200 ml ✓ Fenpaed ETOPROFEN : Cap long-acting 200 mg 12.07 28 ✓ Oruvail SR EFENAMIC ACID			,	
ETOPROFEN : Cap long-acting 200 mg 12.07 28 ✓ Oruvail SR EFENAMIC ACID :: 12.5 50 9000000000000000000000000000000000000				
:: Cap long-acting 200 mg 12.07 28 ✓ Oruvail SR EFENAMIC ACID 12.5 50 Ponstan :: Cap 250 mg 0.50 20 Ponstan 0.50 20 Ponstan Ponstan 0.50 20 Ponstan Ponstan : Tab 250 mg 18.06 500 ✓ Noflam 250 : Tab 500 mg 18.91 250 ✓ Noflam 500 : Tab long-acting 750 mg 5.60 28 ✓ Naprosyn SR 750 : Tab long-acting 1 g 6.53 28 ✓ Naprosyn SR 1000 ULINDAC - - Aclin - : Tab 200 mg 15.10 50 ✓ Aclin - : Tab 20 mg 10.95 100 ✓ Tilcotil - : Tab 20 mg vial 9.95 1 ✓ AFT NSAIDS Other - - - - ELECOXIB - - - - - Cap 100 mg 2.30 30 ✓ Celecoxib Pfizer - ELOXICAM - Special Authority see SA1034 on the next page – Retail pharmacy - <td< td=""><td></td><td></td><td></td><td></td></td<>				
EFENAMIC ACID • Cap 250 mg		12.07	28	Oruvail SP
# Cap 250 mg 1.25 50 (9.16) Ponstan 0.50 20 (5.60) Ponstan APROXEN 18.06 500 * Tab 250 mg 18.06 500 * Tab 500 mg 18.91 250 * Tab 500 mg 6.53 28 • Noflam 500 * Tab long-acting 750 mg 5.60 28 • Naprosyn SR 750 * Tab long-acting 1 g 6.53 28 • Naprosyn SR 1000 ULINDAC * * Naprosyn SR 1000 ULINDAC * Tab 200 mg 15.10 50 • Aclin ENOXICAM * * Tab 20 mg 10.95 100 • Tilcotil * Tab 20 mg vial 9.95 1 • AFT NSAIDs Other ELECOXIB 2.30 30 • Celecoxib Pfizer ELOXICAM - Special Authority see SA1034 on the next page - Retail pharmacy • Celecoxib Pfizer • Celecoxib Pfizer			20	
(9.16) Ponstan 0.50 20 (5.60) Ponstan APROXEN 18.06 500 ✓ Noflam 250 • Tab 250 mg 18.91 250 ✓ Noflam 500 • Tab long-acting 750 mg 5.60 28 ✓ Naprosyn SR 750 • Tab long-acting 1 g 6.53 28 ✓ Naprosyn SR 750 • Tab 100 mg 8.55 50 ✓ Aclin • Tab 200 mg 15.10 50 ✓ Aclin • Tab 20 mg 10.95 100 ✓ Tilcotil • Tab 20 mg vial 9.95 1 ✓ AFT NSAIDS Other ELECOXIB 3.63 60 ✓ Celecoxib Pfizer ELOXICAM Special Authority see SA1034 on the next page – Retail pharmacy 3.0 ✓ Celecoxib Pfizer		4.05	50	
0.50 20 (5.60) Ponstan APROXEN 18.06 500 ✓ Noflam 250 • Tab 250 mg 18.91 250 ✓ Noflam 500 • Tab long-acting 750 mg 5.60 28 ✓ Naprosyn SR 750 • Tab long-acting 1 g 6.53 28 ✓ Naprosyn SR 1000 ULINDAC - - Aclin • Tab 100 mg 8.55 50 ✓ Aclin • Tab 200 mg 15.10 50 ✓ Aclin ENOXICAM - - Tilcotil • Tab 20 mg vial 9.95 1 ✓ AFT NSAIDS Other - - - ELECOXIB - - - Cap 100 mg 3.63 60 ✓ Celecoxib Pfizer Cap 200 mg 2.30 30 ✓ Celecoxib Pfizer ELOXICAM - Special Authority see SA1034 on the next page - Retail pharmacy - -	* Cap 250 mg	()	50	Develop
(5.60) Ponstan APROXEN 18.06 500 ✓ Noflam 250 • Tab 250 mg 18.91 250 ✓ Noflam 500 • Tab long-acting 750 mg 5.60 28 ✓ Naprosyn SR 750 • Tab long-acting 1 g 6.53 28 ✓ Naprosyn SR 1000 ULINDAC - - Aclin • Tab 200 mg 15.10 50 ✓ Aclin ENOXICAM - - - • Tab 200 mg 10.95 100 ✓ Tilcotil • Inj 20 mg vial 9.95 1 ✓ AFT NSAIDS Other - - - ELECOXIB - - - Cap 100 mg 3.63 60 ✓ Celecoxib Pfizer Cap 200 mg 2.30 30 ✓ Celecoxib Pfizer ELOXICAM - Special Authority see SA1034 on the next page – Retail pharmacy ✓ Celecoxib Pfizer		()	00	Ponsian
APROXEN 18.06 500 ✓ Noflam 250 Tab 500 mg 18.91 250 ✓ Noflam 500 Tab long-acting 750 mg 5.60 28 ✓ Naprosyn SR 750 Tab long-acting 1 g 6.53 28 ✓ Naprosyn SR 1000 ULINDAC 6.53 28 ✓ Aclin Tab 100 mg 8.55 50 ✓ Aclin Tab 200 mg 15.10 50 ✓ Aclin ENOXICAM 10.95 100 ✓ Tilcotil Inj 20 mg vial 9.95 1 ✓ AFT NSAIDS Other 2.30 30 ✓ Celecoxib Pfizer ELECOXIB 2.30 30 ✓ Celecoxib Pfizer Cap 200 mg 2.30 30 ✓ Celecoxib Pfizer ELOXICAM – Special Authority see SA1034 on the next page – Retail pharmacy ✓ Celecoxib Pfizer			20	Donoton
# Tab 250 mg 18.06 500 ✓ Noflam 250 # Tab 500 mg 18.91 250 ✓ Noflam 500 # Tab long-acting 750 mg 5.60 28 ✓ Naprosyn SR 750 # Tab long-acting 1 g 6.53 28 ✓ Naprosyn SR 1000 ULINDAC 6.53 28 ✓ Maprosyn SR 1000 ULINDAC 8.55 50 ✓ Aclin # Tab 200 mg 15.10 50 ✓ Aclin ENOXICAM 10.95 100 ✓ Tilcotil # Tab 20 mg vial 9.95 1 ✓ AFT NSAIDS Other ELECOXIB 2.30 30 ✓ Celecoxib Pfizer ELOXICAM – Special Authority see SA1034 on the next page – Retail pharmacy 4.30 ✓ Celecoxib Pfizer		(5.60)		FUISIAII
: Tab 500 mg 18.91 250 ✓ Noflam 500 : Tab long-acting 750 mg 5.60 28 ✓ Naprosyn SR 750 : Tab long-acting 1 g 6.53 28 ✓ Naprosyn SR 1000 ULINDAC 6.53 28 ✓ Naprosyn SR 1000 ULINDAC 8.55 50 ✓ Aclin : Tab 200 mg 15.10 50 ✓ Aclin ENOXICAM 10.95 100 ✓ Tilcotil : Tab 20 mg vial 9.95 1 ✓ AFT NSAIDS Other ELECOXIB 3.63 60 ✓ Celecoxib Pfizer Cap 200 mg 2.30 30 ✓ Celecoxib Pfizer ELOXICAM – Special Authority see SA1034 on the next page – Retail pharmacy ✓ Naprosyn SR 1000 ✓ Celecoxib Pfizer	1	10.00		() // OF0
: Tab long-acting 750 mg				
# Tab long-acting 1 g				
ULINDAC 8.55 50 ✓ Aclin Tab 100 mg 15.10 50 ✓ Aclin Tab 200 mg 15.10 50 ✓ Aclin ENOXICAM 10.95 100 ✓ Tilcotil Inj 20 mg vial 9.95 1 ✓ AFT NSAIDS Other ELECOXIB 2.30 30 ✓ Celecoxib Pfizer ELOXICAM – Special Authority see SA1034 on the next page – Retail pharmacy Feature of the set of the se				
Tab 100 mg 8.55 50 ✓ Aclin Tab 200 mg 15.10 50 ✓ Aclin ENOXICAM 10.95 100 ✓ Tilcotil Inj 20 mg vial 9.95 1 ✓ AFT NSAIDS Other ELECOXIB 3.63 60 ✓ Celecoxib Pfizer Cap 100 mg 2.30 30 ✓ Celecoxib Pfizer ELOXICAM – Special Authority see SA1034 on the next page – Retail pharmacy ✓ Celecoxib Pfizer		0.03	20	Maprosyn SR 1000
 Tab 200 mg Tab 200 mg Tab 20 mg Tab 20 mg Tab 20 mg 10.95 100 Tilcotil Inj 20 mg vial 9.95 AFT NSAIDs Other ELECOXIB Cap 100 mg	SULINDAC			
ENOXICAM Tab 20 mg				
F Tab 20 mg 10.95 100 ✓ Tilcotil Inj 20 mg vial 9.95 1 ✓ AFT NSAIDs Other ELECOXIB 3.63 60 ✓ Celecoxib Pfizer Cap 100 mg 2.30 30 ✓ Celecoxib Pfizer ELOXICAM – Special Authority see SA1034 on the next page – Retail pharmacy Feature ✓ Celecoxib Pfizer	₭ Tab 200 mg		50	Aclin
 Inj 20 mg vial	TENOXICAM			
NSAIDs Other ELECOXIB Cap 100 mg	✤ Tab 20 mg		100	 <u>Tilcotil</u>
ELECOXIB 3.63 60 ✓ Celecoxib Pfizer Cap 100 mg 2.30 30 ✓ Celecoxib Pfizer Cap 200 mg 2.30 30 ✓ Celecoxib Pfizer ELOXICAM – Special Authority see SA1034 on the next page – Retail pharmacy Figure 100	* Inj 20 mg vial	9.95	1	🖌 AFT
ELECOXIB 3.63 60 ✓ Celecoxib Pfizer Cap 100 mg 2.30 30 ✓ Celecoxib Pfizer Cap 200 mg 2.30 30 ✓ Celecoxib Pfizer ELOXICAM – Special Authority see SA1034 on the next page – Retail pharmacy Figure 100	NSAIDs Other			
Cap 100 mg				
Cap 200 mg2.30 30 Celecoxib Pfizer ELOXICAM – Special Authority see SA1034 on the next page – Retail pharmacy	CELECOXIB			
ELOXICAM - Special Authority see SA1034 on the next page - Retail pharmacy	Cap 100 mg	3.63		
	Cap 200 mg	2.30	30	Celecoxib Pfizer
	MELOXICAM - Special Authority see SA1034 on the next page -	Retail pharmacv		
-			30	Arrow-Meloxicam
	-			

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once if end

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	1	Manufacturer	

⇒SA1034 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has moderate to severe haemophilia with less than or equal to 5% of normal circulating functional clotting factor; and
- 2 The patient has haemophilic arthropathy; and
- 3 Pain and inflammation associated with haemophilic arthropathy is inadequately controlled by alternative funded treatment options, or alternative funded treatment options are contraindicated.

Topical Products for Joint and Muscular Pain

CAPSAICIN

Crm 0.025% – Special Authority see SA1289 below – Retail			
pharmacy	6.95	25 g OP	 Zostrix
	9.95	45 a OP	 Zostrix

⇒SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Antirheumatoid Agents		
HYDROXYCHLOROQUINE * Tab 200 mg10.50 LEFLUNOMIDE	100	✓ <u>Plaquenil</u>
Tab 10 mg2.90 Tab 20 mg	30 30	 ✓ <u>Apo-Leflunomide</u> ✓ <u>Apo-Leflunomide</u>
PENICILLAMINE Tab 125 mg	100 100	 ✓ D-Penamine ✓ D-Penamine
SODIUM AUROTHIOMALATE Inj 10 mg in 0.5 ml ampoule76.87 Inj 20 mg in 0.5 ml ampoule113.17	10 10	 Myocrisin Myocrisin
Inj 50 mg in 0.5 ml ampoule217.23	10	 Myocrisin

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

⇒SA1039 Special Authority for Subsidy

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD)) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	✓	Manufacturer

continued...

densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or

- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Note); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or raloxifene.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is receiving systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for zoledronic acid (Underlying cause glucocorticosteroid therapy) or raloxifene.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year where the patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents).

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the `Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Note); or

*Three months or six months, as applicable, dispensed all-at-once

- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause was glucocorticosteroid therapy but patient now meets the `Underlying cause Osteoporosis' criteria) or raloxifene.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.

Subsidy		Fully	Brand or
(Manufacturer's Price)	_	Subsidised	Generic
\$	Per		Manufacturer

continued...

d) In line with the Australian guidelines for funding alendronate, a vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

AL	ENDRONATE SODIUM	- Special Authority see SA	1039 on page 122 - Ret	ail pharmacy	
*	Tab 70 mg		4.82	4	 Fosamax
AL	ENDRONATE SODIUM	WITH COLECALCIFEROL	- Special Authority see	SA1039 on page	122 – Retail pharmacy
*	Tab 70 mg with coleca	lciferol 5,600 iu	4.82	4	 Fosamax Plus

Alendronate for Paget's Disease

SA0949 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain: or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications due to site (base of skull, spine, long bones of lower limbs); or
 - 2.5 Preparation for orthopaedic surgery.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

ALENDRONATE SODIUM - Special Authority see SA0949 above	- Retail pharmacy		
* Tab 40 mg	133.00	30	 Fosamax

Other Treatments

ETIDRONATE DISODIUM – See prescribing guideline below		
* Tab 200 mg 13.50	100	Arrow-Etidronate
Prescribing Guidelines		

Prescribing Guidelines

Etidronate for osteoporosis should be prescribed for 14 days (400 mg in the morning) and repeated every three months. It should not be taken at the same time of the day as any calcium supplementation (minimum dose - 500 mg per day of elemental calcium). Etidronate should be taken at least 2 hours before or after any food or fluid, except water.

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	5.98	1	Pamisol
Inj 6 mg per ml, 10 ml vial		1	 Pamisol
Inj 9 mg per ml, 10 ml vial	17.05	1	 Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see SA11	38 below – Retail	pharmacy	
* Tab 60 mg	53.76	28	 Evista

■SA1138 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsic	lised	Generic	
\$	Per	1	Manufacturer	

continued...

equal to -2.5) (see Notes); or

- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a prior Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or alendronate (Underlying cause - Osteoporosis).

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and guantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM Tab 35 mg	3.80	4	✓ <u>Risedronate Sandoz</u>
TERIPARATIDE – Special Authority see SA1139 below – Retail p Inj 250 mcg per ml, 2.4 ml		1	✓ Forteo

SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

‡ safety cap

	Subsidy (Manufacturer's Prio \$	ce) Subsi Per	Fully dised	Brand or Generic Manufacturer
ZOLEDRONIC ACID				
Inj 0.05 mg per ml, 100 ml, vial – Special Authority see SA1187 below – Retail pharmacy	600.00	100 ml OP	🗸 Ac	lasta
► SA1187 Special Authority for Subsidy				
Initial application — (Paget's disease) from any relevant	practitioner. Approvals	valid for 1 ye	ar for ap	plications meeting the
following criteria:				
All of the following:				
1 Paget's disease; and				
2 Any of the following:				
2.1 Bone or articular pain; or				
2.2 Bone deformity; or2.3 Bone, articular or neurological complications;	or			
2.4 Asymptomatic disease, but risk of complications,				
2.5 Preparation for orthopaedic surgery; and	13, 01			
3 The patient will not be prescribed more than 5 mg of	zoledronic acid in the 1	2-month appr	oval per	iod.
Initial application — (Underlying cause - Osteoporosis)				
renewal unless notified for applications meeting the following				
Both:				
1 Any of the following:				
1.1 History of one significant osteoporotic fracture	e demonstrated radiolog	ically and do	cumente	d bone mineral density
(BMD) greater than or equal to 2.5 standard of	eviations below the me	an normal val	ue in yo	ung adults (i.e. T-Score
less than or equal to -2.5) (see Note); or				
1.2 History of one significant osteoporotic fracture				
densitometry scanning cannot be performed b				physiological reasons.
is unlikely that this provision would apply to m 1.3 History of two significant osteoporotic fracture			or	
1.4 Documented T-Score less than or equal to -3.		gically, of		
1.5 A 10-year risk of hip fracture greater than or e	()/	usina a nuhlis	had rick	assessment algorithm
(e.g. FRAX or Garvan) which incorporates Bl				abbeboment algorithm
1.6 Patient has had a Special Authority approval			Osteopoi	osis) or raloxifene; and
2 The patient will not be prescribed more than 5 mg of				, ,
Initial application — (Underlying cause - glucocorticoste	roid therapy) from an	y relevant pra	ctitioner	. Approvals valid for 1
year for applications meeting the following criteria:				
All of the following:				
1 The patient is receiving systemic glucocorticosteroid				
equivalents) and has already received or is expected	to receive therapy for a	at least three i	months;	and
2 Any of the following:				
2.1 The patient has documented BMD greater that		ard deviations	below t	he mean normal value i
young adults (i.e. T-Score less than or equal		onotrot-d us d		hu or
2.2 The patient has a history of one significant os2.3 The patient has had a Special Authority approx				
2.3 The patient has had a Special Authomy appro or raloxifene; and		denying caus	e - giuci	sonicosterola inerapy)
O The notions will not be preservibed more than E mg of		0		

3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	:	Subsidised	Generic	
 \$	Per	1	Manufacturer	

continued...

- 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
- 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
 - 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
 - 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) or raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

*Three months or six months, as applicable, dispensed all-at-once

DP-Allopurinol
DP-Allopurinol

‡ safety cap

▲ Three months supply may be dispensed at one time

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
BENZBROMARONE – Special Authority see SA1537 below – Re Tab 100 mg		100	1	Benzbromaron AL 100 629

⇒SA1537 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 Both:
 - 2.3.1 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Notes); and
 - 2.3.2 The patient has a rate of creatinine clearance greater than or equal to 20 ml/min; or
 - 2.4 All of the following:
 - 2.4.1 The patient is taking azathioprine and requires urate-lowering therapy; and
 - 2.4.2 Allopurinol is contraindicated; and
 - 2.4.3 Appropriate doses of probenecid are ineffective or probenecid cannot be used due to reduced renal function; and
- 3 The patient is receiving monthly liver function tests.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.
- Notes: Benzbromarone has been associated with potentially fatal hepatotoxicity.

In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

The New Zealand Rheumatology Association has developed information for prescribers which can be accessed from its website at www.rheumatology.org.nz/home/resources-2/

COLCHICINE

* Tab 500 mcg		100	 Colgout
FEBUXOSTAT - Special Authority see SA1538 below - Retail p	harmacy		
Tab 80 mg		28	 Adenuric
Tab 120 mg		28	 Adenuric

⇒SA1538 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or

Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
 (Manulactarer 3 1 hee) \$	Per	siuiseu ✓	Manufacturer

continued...

- 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
- 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note).

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

PROBENECID

* Tab 500 mg	55.00	100	Probenecid-AFT
Muscle Relaxants			
BACLOFEN			
* Tab 10 mg – For baclofen oral liquid formulation refer, page 2	<mark>24</mark> 3.85	100	 Pacifen
Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement.	11.55	1	 Lioresal Intrathecal
Subsidised only for use in a programmable pump in patien caused intolerable side effects and the prescription is end			nts have been ineffective or have
Inj 2 mg per ml, 5 ml ampoule – Subsidy by endorsement		1	 Lioresal Intrathecal
Subsidised only for use in a programmable pump in patier caused intolerable side effects and the prescription is end			nts have been ineffective or have
DANTROLENE			
Cap 25 mg	65.00	100	 Dantrium
			 Dantrium S29 S29
Cap 50 mg	77.00	100	 Dantrium
ORPHENADRINE CITRATE			
Tab 100 mg		100	 Norflex
Norflex to be Sole Supply on 1 July 2018			

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully Brand or
	(Manufacturer's Price)		Subsidised Generic
	\$	Per	 Manufacturer
Agents for Parkinsonism and Related Disorde	rs		
Dopamine Agonists and Related Agents			
AMANTADINE HYDROCHLORIDE			
▲ Cap 100 mg		60	 Symmetrel
APOMORPHINE HYDROCHLORIDE			-
Inj 10 mg per ml, 2 ml ampoule		5	 Movapo
BROMOCRIPTINE MESYLATE			-
₭ Tab 2.5 mg		100	Apo-Bromocriptine
ENTACAPONE			
Tab 200 mg		100	 Entapone
EVODOPA WITH BENSERAZIDE			
Tab dispersible 50 mg with benserazide 12.5 mg		100	 Madopar Rapid
₭ Cap 50 mg with benserazide 12.5 mg		100	✓ Madopar 62.5
K Cap 100 mg with benserazide 25 mg		100	 Madopar 125
Cap long-acting 100 mg with benserazide 25 mg		100	 Madopar HBS
Cap 200 mg with benserazide 50 mg	25.00	100	 Madopar 250
EVODOPA WITH CARBIDOPA			
Tab 100 mg with carbidopa 25 mg – For levodopa with			
carbidopa oral liquid formulation refer, page 224	17.97	100	 Kinson
			 Sinemet
Sinemet to be Sole Supply on 1 July 2018	07.45		
Tab long-acting 200 mg with carbidopa 50 mg		100	
Tab 250 mg with carbidopa 25 mg Kinson Tab 100 mg with carbidopa 25 mg to be delisted 1 July		100	✓ Sinemet
	(2010)		
PRAMIPEXOLE HYDROCHLORIDE Tab 0.25 mg	7.20	100	- Paminov
Tab 0.25 mg		100	 ✓ <u>Ramipex</u> ✓ Ramipex
-		100	• <u>Hampex</u>
ROPINIROLE HYDROCHLORIDE	0.79	100	Apo-Ropinirole
Tab 0.25 mg Tab 1 mg		100	 ✓ <u>Apo-Ropinirole</u> ✓ Apo-Ropinirole
Tab 2 mg		100	✓ Apo-Ropinirole
Tab 5 mg		100	✓ Apo-Ropinirole
★ Tab 5 mg		100	Apo-Selegiline
·			S29 S29
OLCAPONE			
Tab 100 mg	132.50	100	 Tasmar
·		100	<u>- ruomui</u>
Anticholinergics			
SENZATROPINE MESYLATE			_
Tab 2 mg		60	 Benztrop
Inj 1 mg per ml, 2 ml		5	✓ Cogentin
	190.00	10	 Omega
a) Up to 10 inj available on a PSO			
b) Only on a PSO			

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
PROCYCLIDINE HYDROCHLORIDE				
Tab 5 mg	7.40	100	✔ К	emadrin
Agents for Essential Tremor, Chorea and Relate	d Disorders			
RILUZOLE – Special Authority see SA1403 below – Retail pharm Wastage claimable – see rule 3.3.2 on page 13				
Tab 50 mg Rilutek to be Sole Supply on 1 September 2018	130.00	56	✓ R	ilutek
SA1403 Special Authority for Subsidy				
Initial application only from a neurologist or respiratory specialis following criteria:	t. Approvals valid fo	r 6 mont	hs for app	blications meeting the
All of the following: 1 The patient has amyotrophic lateral sclerosis with disease	duration of 5 years	or loca: a	nd	
 The patient has an yourophic tatefal scienciss with disease The patient has at least 60 percent of predicted forced vita The patient has not undergone a tracheostomy; and The patient has not experienced respiratory failure; and Any of the following: The patient is ambulatory; or The patient is able to use upper limbs; or The patient is able to swallow. 				initial application; and
Renewal from any relevant practitioner. Approvals valid for 18 m	onths for application	s meetin	g the follo	owing criteria:
All of the following:				
 The patient has not undergone a tracheostomy; and The patient has not experienced respiratory failure; and Any of the following: 				
3.1 The patient is ambulatory; or				
3.2 The patient is able to use upper limbs; or				
3.3 The patient is able to swallow.				
TETRABENAZINE Tab 25 mg	91.10	112	✓ <u>N</u>	lotetis
Anaesthetics				
Local				
LIDOCAINE [LIGNOCAINE] Gel 2%, tube – Subsidy by endorsement a) Up to 150 ml available on a PSO	14.50	30 ml	✓ X	ylocaine 2% Jelly
 b) Subsidised only if prescribed for urethral or cervical a Gel 2%, 10 ml urethral syringe – Subsidy by endorsement 		10	🗸 P	fizer
a) Up to 5 each available on a PSO	160.00	25		athejell
b) Subsidised only if prescribed for urethral or cervical a	aministration and the	e prescrip	ption is er	ndorsed accordingly.

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully	
	(Manufacturer's Price \$) Per	Subsidised	
	Ŷ	1.01		Manufacturor
IDOCAINE [LIGNOCAINE] HYDROCHLORIDE	~~~~			••
Oral (gel) soln 2%		200 m		Mucosoothe
Inj 1%, 5 ml ampoule – Up to 25 inj available on a PSO		25	v	Lidocaine-Claris
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule – Up to 5 inj available on a PSO		25	✓	Lidocaine-Claris
Inj 1%, 20 ml ampoule – Up to 5 inj available on a PSO	2.40	1	✓	Lidocaine-Claris
	12.00	5		
	(20.00)			Xylocaine
Inj 1%, 20 ml vial – Up to 5 inj available on a PSO		5	1	Lidocaine-Claris
Inj 2%, 20 ml ampoule - Up to 5 inj available on a PSO	2.40	1	✓	Lidocaine-Claris
Inj 2%, 20 ml vial – Up to 5 inj available on a PSO		5	✓	Lidocaine-Claris
IDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes -				
Subsidy by endorsement		10	1	Pfizer
a) Up to 5 each available on a PSO		10		
b) Subsidised only if prescribed for urethral or cervical	administration and th	o proc	orintion ic	andoread accordingly
b) Subsidised only if prescribed for dreamar of cervical		ie pies		endorsed accordingly.
Topical Local Anaesthetics				
SA0906 Special Authority for Subsidy				
nitial application from any relevant practitioner. Approvals val	id for 2 years where	the nati	iont is a cl	hild with a chronic medical
ondition requiring frequent injections or venepuncture.	in in 2 years where	ine pau		
Renewal from any relevant practitioner. Approvals valid for 2 ye	are whore the treate	ont ro	naine ann	ropriate and the patient is
enefiting from treatment.			nains app	nopriate and the patient is
5	Datallah			
IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 at	ove – Hetail pharma	су		

LIDOCAINE [LIGNOCAINE] - Special Authority see SA0906 above	– Hetali phar	macy	
Crm 4%	5.40	5 a OP	🖌 LMX4
	27.00	30 g OP	🖌 LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Authorit	y see SA0906	6 above – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	🖌 EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	🖌 EMLA

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 121

Non-opioid Analgesics

For aspirin & chloroform application refer Standard Formulae, page 2	27		
ASPIRIN			6 - .
* Tab dispersible 300 mg – Up to 30 tab available on a PSO	3.90	100	 Ethics Aspirin
CAPSAICIN – Subsidy by endorsement			
Subsidised only if prescribed for post-herpetic neuralgia or diabe accordingly.	tic peripheral	neuropathy an	d the prescription is endorsed
Crm 0.075%	12.50	45 a OP	 Zostrix HP
NEFOPAM HYDROCHLORIDE		- 5 -	
Tab 30 mg	23.40	90	 Acupan

	Subsidy		Fully	Brand or
	(Manufacturer's Pri		sidised	Generic
	\$	Per	/	Manufacturer
PARACETAMOL				
* Tab 500 mg - blister pack – Up to 30 tab available on a PSO	7.12	1,000		Pharmacare
* Tab 500 mg - bottle pack	6.32	1,000	-	Pharmacare
* ‡ Oral liq 120 mg per 5 ml	5.35	1,000 ml	-	Paracare
a) Up to 200 ml available on a PSO				
b) Not in combination				
*‡ Oral lig 250 mg per 5 ml	5.81	1,000 ml	-	Paracare Double
		,		Strength
a) Up to 100 ml available on a PSO				U U
b) Not in combination				
c) Paracare Double Strength to be Sole Supply on 1 Se	ntember 2018			
♣ Suppos 125 mg		10	1	Gacet
✤ Suppos 125 mg		10		Gacet
		50		Paracare
Suppos 500 mg	12.00	50	•	Falacale
Opioid Analgesics				
CODEINE PHOSPHATE – Safety medicine; prescriber may dete	rmine dispensing	frequency		
Tab 15 mg	5.75	100	✓	PSM
Tab 30 mg	6.80	100	-	PSM
Tab 60 mg		100	-	PSM
DIHYDROCODEINE TARTRATE				
Tab long-acting 60 mg	9 55	60	1	DHC Continus
		00	•	bilo continus
ENTANYL				
a) Only on a controlled drug form				
 b) No patient co-payment payable 				
 c) Safety medicine; prescriber may determine dispensing fre 				
Inj 50 mcg per ml, 2 ml ampoule	3.95	10	-	Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule	10.45	10	✓	Boucher and Muir
Patch 12.5 mcg per hour	2.95	5	-	Fentanyl Sandoz
Patch 25 mcg per hour	3.66	5	-	Fentanyl Sandoz
Patch 50 mcg per hour	6.65	5	✓	Fentanyl Sandoz
Patch 75 mcg per hour	9.25	5	-	Fentanyl Sandoz
Patch 100 mcg per hour		5		Fentanyl Sandoz
IETHADONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre				
 d) Extemporaneously compounded methadone will only be r 	eimpursed at the	rate of the ch	ieapes	a torm available
(methadone powder, not methadone tablets).		_		
e) For methadone hydrochloride oral liquid refer Standard Fo				
Tab 5 mg		10		Methatabs
Cral liq 2 mg per ml		200 ml		Biodone
		200 ml	-	Biodone Forte
Oral liq 5 mg per ml				
Oral liq 5 mg per ml Oral liq 10 mg per ml		200 ml	-	Biodone Extra Forte

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

		Subsidy	<u> </u>	Fully	Brand or
		(Manufacturer's Price) \$	Subsi Per	dised	Generic Manufacturer
ЛС	DRPHINE HYDROCHLORIDE	•			
///	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing free	anopov			
F	Oral lig 1 mg per ml		200 ml	1	RA-Morph
F	Oral lig 2 mg per ml		200 ml		RA-Morph
F	Oral lig 5 mg per ml		200 ml		RA-Morph
F	Oral lig 10 mg per ml		200 ml		RA-Morph
			200 111	•	
٩C	ORPHINE SULPHATE				
	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing free			-	
	Tab immediate-release 10 mg		10		Sevredol
	Tab long-acting 10 mg		10		Arrow-Morphine LA
	Tab immediate-release 20 mg		10		Sevredol
	Tab long-acting 30 mg	2.85	10		Arrow-Morphine LA
	Tab long-acting 60 mg		10		Arrow-Morphine LA
	Tab long-acting 100 mg	6.10	10		Arrow-Morphine LA
	Cap long-acting 10 mg	1.70	10		m-Eslon
	Cap long-acting 30 mg	2.50	10		m-Eslon
	Cap long-acting 60 mg	5.40	10		m-Eslon
	Cap long-acting 100 mg	6.38	10		m-Eslon
	Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS	O6.27	5		DBL Morphine
					Sulphate
	Inj 10 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS	SO4.47	5	✓	DBL Morphine
					Sulphate
	Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS	SO4.76	5	1	DBL Morphine
					Sulphate
	Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS	SO6.19	5	1	DBL Morphine
					Sulphate
л	DRPHINE TARTRATE				
	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing free	alleboy			
	Inj 80 mg per ml, 1.5 ml ampoule		5	1	DBL Morphine
			5	•	Tartrate
					Taillale

NERVOUS	SYSTEM
---------	--------

(1.	Subsidy Ianufacturer's Price)	Fully Brand or Subsidised Generic
(1)	s s rice	Per	Manufacturer
XYCODONE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
 c) Safety medicine; prescriber may determine dispensing frequ 	10001		
		00	
Tab controlled-release 5 mg		20	✓ <u>BNM</u>
Tab controlled-release 10 mg		20	✓ BNM
Tab controlled-release 20 mg		20	✓ <u>BNM</u>
Tab controlled-release 40 mg		20	✓ <u>BNM</u>
Tab controlled-release 80 mg		20	✓ <u>BNM</u>
Cap immediate-release 5 mg	1.98	20	 OxyNorm
Cap immediate-release 10 mg	3.91	20	✓ OxyNorm
Cap immediate-release 20 mg	6.84	20	 OxyNorm
Oral lig 5 mg per 5 ml	11.20	250 m	🖌 🗸 OxyNorm
Inj 10 mg per ml, 1 ml ampoule		5	✓ OxyNorm
Inj 10 mg per ml, 2 ml ampoule		5	✓ OxyNorm
Inj 50 mg per ml, 1 ml ampoule		5	✓ OxyNorm
		-	
ARACETAMOL WITH CODEINE - Safety medicine; prescriber ma			
K Tab paracetamol 500 mg with codeine phosphate 8 mg		1,000	
			Codeine (Relieve
ETHIDINE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing frequ	IODOV		
		10	✓ PSM
Tab 50 mg		10	✓ <u>PSM</u> ✓ PSM
Tab 100 mg			
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSC	J4.98	5	✓ <u>DBL Pethidine</u>
			Hydrochloride
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSC	D5.12	5	DBL Pethidine
			Hydrochloride
PSM Tab 100 mg to be delisted 1 July 2018)			
	1 55	00	Tramel CD 100
Tab sustained-release 100 mg		20	Tramal SR 100
Tab sustained-release 150 mg		20	Tramal SR 150
Tab sustained-release 200 mg		20	Tramal SR 200
Cap 50 mg - For tramadol hydrochloride oral liquid formulation			
refer, page 224	2.25	100	Arrow-Tramadol
Antidepressants			
Cyclic and Related Agents			
e jone and neuteu Agente			
MITRIPTYLINE - Safety medicine; prescriber may determine disp	ensing frequency	/	
Tab 10 mg		100	 Arrow-Amitriptyline
		100	✓ Arrow-Amitriptyline
5		100	
Tab 25 mg		100	Arrow Amitrintalia
Tab 25 mg Tab 50 mg	2.51	100	✓ Arrow-Amitriptyline
Tab 25 mg Tab 50 mg COMIPRAMINE HYDROCHLORIDE – Safety medicine; prescribe	2.51 r may determine	dispen	sing frequency
Tab 25 mg Tab 50 mg	2.51 r may determine		

DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequencyTab 75 mg11.19100✓ DopressCap 25 mg6.45100✓ Dopress

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once if

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	/ Brand or
	(Manufacturer's Price)	_	Subsidised	I Generic
	\$	Per		Manufacturer
DOXEPIN HYDROCHLORIDE – Safety medicine; prescriber ma	•	-		• •
Cap 10 mg		100		Anten
Cap 25 mg		100		Anten
Cap 50 mg	8.55	100	~	Anten
IMIPRAMINE HYDROCHLORIDE - Safety medicine; prescriber	may determine dispe	nsing	frequenc	у
Tab 10 mg	5.48	50	 ✓ 	Tofranil
	6.58	60	1	Tofranil s29 S29
	10.96	100	1	Tofranil
Tab 25 mg		50	1	Tofranil
MAPROTILINE HYDROCHLORIDE - Safety medicine; prescrib		nonci	na frequer	
Tab 25 mg	•	30		Ludiomil
Tab 23 mg	12.53	50		Ludiomil
	25.06	100		Ludiomil
Tab 75 mg		20		Ludiomil
Tab 75 mg	21.01	20 30		Ludiomil
NORTRIPTYLINE HYDROCHLORIDE - Safety medicine; presc				
Tab 10 mg		100	-	Norpress
Tab 25 mg	7.08	180	~	Norpress
Manageming Ouidage labilities (MAOle) New O	a la ativa			
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	elective			
PHENELZINE SULPHATE				
* Tab 15 mg	95.00	100	1	Nardil
FRANYLCYPROMINE SULPHATE				
	00.04	50		Parnate
* Tab 10 mg	22.94	50	•	Parnale
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE				
* Tab 150 mg		500	1	Apo-Moclobemide
* Tab 300 mg		100		Apo-Moclobemide
-				
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE				
* Tab 20 mg	1 70	84	1	PSM Citalopram
-	1.79	04	•	
ESCITALOPRAM				
* Tab 10 mg	1.11	28	<i>✓</i>	Escitalopram-
				Apotex
* Tab 20 mg	1 00	28		Escitalopram-
* Tab 20 mg	1.90	20	•	
				Apotex
FLUOXETINE HYDROCHLORIDE				
 Tab dispersible 20 mg, scored – Subsidy by endorsement 	2 47	30	1	Arrow-Fluoxetine
Subsidised by endorsement		00	•	
 When prescribed for a patient who cannot swallow 	whole tablete or con		and the e	recorintion is orderead
accordingly; or	whole lablets of caps	Sules	anu ine pi	
2) When prescribed in a daily dose that is not a multij	ale of 20 mg in which	0200	the prese	rintion is deemed to be
endorsed. Note: Tablets should be combined with				
chaorsea. Note. Tablets should be combined with	· supsuiss to identitate			u ing u0000.
* Cap 20 mg	1 99	90	1	Arrow-Fluoxetine
- Oup 20 mg		50	•	
fully subsidised	S29 Unapproved	d medi	cine sunnli	ed under Section 29
136 [HP4] refer page 4	Sole Subsidised			
נווו דן וטוטו אמשט ד	Cole Subsidised	Jahh	1	

	Subsidy		Fully	
	(Manufacturer's Price)		Subsidised	
	\$	Per	/	Manufacturer
PAROXETINE				
* Tab 20 mg	4.02	90	✓	Apo-Paroxetine
SERTRALINE				
* Tab 50 mg	3 05	90	1	Arrow-Sertraline
* Tab 100 mg		90	1	Arrow-Sertraline
		00	•	Arrow Certruinie
Other Antidepressants				
MIRTAZAPINE				
Tab 30 mg	2 55	30	1	Apo-Mirtazapine
Tab 45 mg		30		Apo-Mirtazapine
VENLAFAXINE	0.20			
	6.00	04		Enlefey VD
* Cap 37.5 mg		84 04		Enlafax XR Enlafax XR
* Cap 75 mg		84 94		Enlafax XR Enlafax XP
* Cap 150 mg		84	•	Enlafax XR
Antionilonov Drugo				
Antiepilepsy Drugs				
America for October 1 of Obstant Exilentians				
Agents for Control of Status Epilepticus				
CLONAZEPAM - Safety medicine; prescriber may determine d	ispansing fraguancy			
Inj 1 mg per ml, 1 ml		5	1	Rivotril
		5	•	nivoun
DIAZEPAM - Safety medicine; prescriber may determine dispe		_		
Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement	11.83	5	~	Hospira
a) Up to 5 inj available on a PSO				
b) Only on a PSO				
 c) PSO must be endorsed "not for anaesthetic procedu 				
Rectal tubes 5 mg - Up to 5 tube available on a PSO		5		Stesolid
Rectal tubes 10 mg – Up to 5 tube available on a PSO	40.87	5	~	Stesolid
PARALDEHYDE				
* Inj 5 ml		5	1	AFT S29
•				
PHENYTOIN SODIUM		F		Hearing
* Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a	PSU88.63	5	•	Hospira
* Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a	400.00	-		
PSO		5	~	Hospira
Control of Epilepsy				
CARBAMAZEPINE	44.50			-
* Tab 200 mg		100	-	Tegretol
* Tab long-acting 200 mg		100	-	Tegretol CR
* Tab 400 mg		100		Tegretol
* Tab long-acting 400 mg		100		Tegretol CR
*‡ Oral liq 20 mg per ml		250 m	· •	Tegretol
CLOBAZAM - Safety medicine; prescriber may determine disp	ensing frequency			
Tab 10 mg		50	✓	Frisium
‡ Safety cap for extemporaneously compounded oral lique	uid preparations.			
CLONAZEPAM - Safety medicine; prescriber may determine d	ispensing frequency			
Oral drops 2.5 mg per ml) ml O	P 🗸	Rivotril
, ,,				

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price	e)	Fully Subsidised	
	\$	Per	1	Manufacturer
ETHOSUXIMIDE				
Cap 250 mg		100	1	Zarontin
	32.90	200	✓	Zarontin
the second		200 n	nl 🖌	Zarontin
GABAPENTIN				
Note: Not subsidised in combination with subsidised preg	abalin			
Cap 100 mg – Note differing brand requirements below	2.65	100	✓	Apo-Gabapentin
	7.16			Arrow-Gabapentin
				Neurontin
			✓	Nupentin
 b) Arrow-Gabapentin brand: Special Authority see S c) Neurontin brand: Special Authority see SA1477 b ▲ Cap 300 mg – Note differing brand requirements below – gabapentin oral liquid formulation refer, page 224 a) Nupentin brand: Special Authority see SA1477 be 	elow – Retail pharmac For 4.07 11.00	y 100	1 1 1	Apo-Gabapentin Arrow-Gabapentin Neurontin Nupentin
 b) Arrow-Gabapentin brand: Special Authority see SA1477 b c) Neurontin brand: Special Authority see SA1477 b 	A1477 below - Retail	pharm	acy	
▲ Cap 400 mg - Note differing brand requirements below		<i></i> 100	1	Apo-Gabapentin
	13.75		1	Arrow-Gabapentin
			✓	Neurontin
			1	Nupentin
a) Nupentin brand: Special Authority see SA1477 be	elow – Retail pharmacy	/		

- b) Arrow-Gabapentin brand: Special Authority see SA1477 below Retail pharmacy
- c) Neurontin brand: Special Authority see SA1477 below Retail pharmacy

► SA1477 Special Authority for Subsidy

Initial application — (Epilepsy) from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:

Either:

- 1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
- 2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents.

Note: "Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Initial application — (Neuropathic pain or Chronic Kidney Disease associated pruritus) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with neuropathic pain; or
- 2 Both:
 - 2.1 The patient has Chronic Kidney Disease Stage 5-associated pruritus* where no other cause for pruritus can be identified (e.g. scabies, allergy); and
 - 2.2 The patient has persistent pruritus not relieved with a trial of emollient/moisturising creams alone.

Renewal — (Epilepsy) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life.

Subsidy (Manufacturer's Price)	s	Fully	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Renewal — (Neuropathic pain or Chronic Kidney Disease associated pruritus) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- Either:
 - 1 The patient has demonstrated a marked improvement in their control of pain or itch (prescriber determined); or
 - 2 The patient has previously demonstrated clinical responsiveness to gabapentin and has now developed neuropathic pain in a new site.

Note: Indications marked with * are Unapproved Indications (see Interpretations and Definitions). Dosage adjustment of gabapentin is recommended for patients with renal impairment.

LACOSAMIDE – Special Authority see SA1125 below – Retail pharmacy

▲ Tab 50 mg		14	 Vimpat
▲ Tab 100 mg		14	 Vimpat
C C	200.24	56	 Vimpat
▲ Tab 150 mg		14	 Vimpat
-	300.40	56	 Vimpat
▲ Tab 200 mg		56	 Vimpat

⇒SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

LAMOTRIGINE

Tab dispersible 2 mg	6.74	30	 Lamictal
▲ Tab dispersible 5 mg		30	 Lamictal
	15.00	56	Arrow-Lamotrigine
▲ Tab dispersible 25 mg		56	 Logem
	20.40		 Arrow-Lamotrigine
	29.09		 Lamictal
▲ Tab dispersible 50 mg		56	 Logem
	34.70		 Arrow-Lamotrigine
	47.89		 Lamictal
▲ Tab dispersible 100 mg		56	 Logem
1 3	59.90		 Arrow-Lamotrigine
	79.16		 Lamictal

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Pric	xe) S	Fully ubsidised	
	\$	Per	/	
EVETIRACETAM				
Tab 250 mg	24.03	60	1	Everet
Tab 500 mg		60	-	Everet
Tab 750 mg	45.23	60	1	Everet
Tab 1,000 mg		60	1	Everet
Toral liq 100 mg per ml		300 ml O	Р 🗸	Levetiracetam-AFT
Levetiracetam-AFT to be Sole Supply on 1 July 2018				
PHENOBARBITONE				
For phenobarbitone oral liquid refer Standard Formulae, pa	ge 227			
* Tab 15 mg		500	1	PSM
* Tab 30 mg		500		PSM
PHENYTOIN SODIUM				
* Tab 50 mg	50 51	200	1	Dilantin Infatab
Cap 30 mg		200		Dilantin
Cap 30 mg		200		Dilantin
*‡ Oral liq 30 mg per 5 ml		500 ml		Dilantin
		500 111	•	Dhanan
PREGABALIN				
Note: Not subsidised in combination with subsidised gabap		50		Deserve halin Dfinan
* Cap 25 mg	2.25	56	•	Pregabalin Pfizer
Pregabalin Pfizer to be Sole Supply on 1 July 2018	0.65	FC		Dreachalin Dfirer
* Cap 75 mg	2.00	56	•	Pregabalin Pfizer
Pregabalin Pfizer to be Sole Supply on 1 July 2018	4.01	FC		Dreachalin Dfirer
* Cap 150 mg	4.01	56	•	Pregabalin Pfizer
Pregabalin Pfizer to be Sole Supply on 1 July 2018	7 20	56		Drogobalin Dfizor
Cap 300 mg Pregabalin Pfizer to be Sole Supply on 1 July 2018		50	•	Pregabalin Pfizer
PRIMIDONE	17.05	100		
* Tab 250 mg	17.25	100	~	Apo-Primidone
SODIUM VALPROATE				
Tab 100 mg	13.65	100	~	Epilim Crushable
Tab 200 mg EC	27.44	100		Epilim
Tab 500 mg EC		100		Epilim
*‡ Oral liq 200 mg per 5 ml	20.48	300 ml	~	Epilim S/F Liquid
				Epilim Syrup
Inj 100 mg per ml, 4 ml	41.50	1	1	Epilim IV
STIRIPENTOL – Special Authority see SA1330 below – Retail	pharmacy			
Cap 250 mg	, ,	60	1	Diacomit S29
Powder for oral lig 250 mg sachet		60		Diacomit S29
Solution of a line 200 mg sachet		00	•	

⇒SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

Λ)	Subsidy /anufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
TOPIRAMATE				
Tab 25 mg	11.07	60	✓	Arrow-Topiramate
C C			-	Topiramate Actavis
	26.04		-	Topamax
▲ Tab 50 mg	18.81	60	~	Arrow-Topiramate
-			 Image: A start of the start of	Topiramate Actavis
	44.26		 Image: A start of the start of	Topamax
Tab 100 mg	31.99	60	✓	Arrow-Topiramate
			 Image: A start of the start of	Topiramate Actavis
	75.25		 Image: A start of the start of	Topamax
Tab 200 mg	55.19	60	✓	Arrow-Topiramate
			~	Topiramate Actavis
	129.85		~	Topamax
Sprinkle cap 15 mg	20.84	60	~	Topamax
Sprinkle cap 25 mg	26.04	60	✓	Topamax
/IGABATRIN - Special Authority see SA1072 below - Retail pharr	nacv			
▲ Tab 500 mg		100	1	Sabril

⇒SA1072 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

1 Either:

- 1.1 Patient has infantile spasms; or
- 1.2 Both:
 - 1.2.1 Patient has epilepsy; and

*Three months or six months, as applicable, dispensed all-at-once

- 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Notes: ``Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages. **Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

‡ safety cap

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	1	Manufacturer
Antimigraine Preparations				
or Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 1	age 121			
Acute Migraine Treatment				
ERGOTAMINE TARTRATE WITH CAFFEINE				
Tab 1 mg with caffeine 100 mg		100	🗸 Ca	fergot
			🗸 Ca	fergot S29 S29
IZATRIPTAN				
Tab orodispersible 10 mg	5.26	30	✓ <u>Riz</u>	amelt
UMATRIPTAN				
Tab 50 mg	24.44	100		o-Sumatriptan
Tab 100 mg		100	🗸 <u>Ap</u>	o-Sumatriptan
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj pe	er			
prescription		2 OP	🗸 Clu	ustran
			🗸 Su	n Pharma S29
Prophylaxis of Migraine				
or Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SY	STEM, page 60			
IZOTIFEN	o i zini, pago oo			
F Tab 500 mcg	23 21	100	🖌 Sa	ndomigran
		100	• 00	naomigran
Antinausea and Vertigo Agents				
or Antispasmodics refer to ALIMENTARY TRACT, page 22				
PREPITANT – Special Authority see SA0987 below – Retail ph	armacy			
Cap 2 × 80 mg and 1 × 125 mg		3 OP	🖌 En	end Tri-Pack
Emend Tri-Pack to be Sole Supply on 1 August 2018	74.40		<i>.</i> -	
Cap 40 mg	71.43	5 OP	🖌 En	nend
Emend Cap 40 mg to be delisted 1 August 2018)				
»SA0987 Special Authority for Subsidy	d fau 10 mandha udhau	- 4k -		evention binkly
itial application from any relevant practitioner. Approvals vali- metogenic chemotherapy and/or anthracycline-based chemothe			•	ergoing nignly
enewal from any relevant practitioner. Approvals valid for 12 m			0 ,	iahly emetogenic
nemotherapy and/or anthracycline-based chemotherapy for the			undergoing n	iginy entetogenie
ETAHISTINE DIHYDROCHLORIDE		·		
F Tab 16 mg	2.89	84	🗸 Ve	rgo 16
YCLIZINE HYDROCHLORIDE				-
Tab 50 mg	0.59	20	🗸 Na	uzene
YCLIZINE LACTATE		-		
Inj 50 mg per ml, 1 ml		5	🗸 Na	usicalm
OMPERIDONE		-		
Tab 10 mg – For domperidone oral liquid formulation refer,				
page 224		100	🗸 Pro	okinex
r-9 ·			<u></u>	

 Metoclopramide Actavis 10

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
HYOSCINE HYDROBROMIDE				
* Inj 400 mcg per ml, 1 ml ampoule		5	✓	Hospira
	93.00	10	 Image: A second s	Martindale S29
Patch 1.5 mg - Special Authority see SA1387 below - Retail				
pharmacy		2	v :	Scopoderm TTS

⇒SA1387 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE

*	Tab 10 mg – For metoclopramide hydrochloride oral liquid formulation refer, page 224	1.30	
*	Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PS	O4.50	

* Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO	4.50 10	 Pfizer
ONDANSETRON		
* Tab 4 mg	3.36 50	Apo-Ondansetron
* Tab disp 4 mg	0.95 10	 Ondansetron
		ODT-ORLA
* Tab 8 mg	4.77 50	Apo-Ondansetron
* Tab disp 8 mg	1.43 10	 Ondansetron
		ODT-DRLA
PROCHLORPERAZINE		
* Tab 3 mg buccal	5.97 50	
(1)	5.00)	Buccastem
* Tab 5 mg – Up to 30 tab available on a PSO	6.35 250	✓ <u>Nausafix</u>
* Inj 12.5 mg per ml, 1 ml – Up to 5 inj available on a PSO2	5.81 10	 Stemetil
PROMETHAZINE THEOCLATE		
* Tab 25 mg	1.20 10	
(8	5.59)	Avomine

Antipsychotics

General

AMISULPRIDE - Safety medicine; prescriber may determine dis	spensing frequence	;y	
Tab 100 mg		30	 Sulprix
Tab 200 mg	14.75	60	 Sulprix
Tab 400 mg		60	 Sulprix
Oral liq 100 mg per ml		60 ml	✓ Solian

*Three months or six months, as applicable, dispensed all-at-once

100

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
ARIPIPRAZOLE – Safety medicine; prescriber may determine dis Tab 5 mg		30	 Aripiprazole Sandoz
Tablet 5 mg – Special Authority (Abilify brand only) see SA1539 below – Retail pharmacy – No more than 1 tab per day	123.54	30	✓ Abilify
Tab 10 mg – Special Authority (Abilify brand only) see SA153 below – Retail pharmacy		30	 ✓ Aripiprazole Sandoz ✓ Abilify
Tab 15 mg – Special Authority (Abilify brand only) see SA153 below – Retail pharmacy		30	 ✓ Aripiprazole Sandoz ✓ Abilify
Tab 20 mg – Special Authority (Abilify brand only) see SA153 below – Retail pharmacy	9	30	 Aripiprazole Sandoz Abilify
Tab 30 mg – Special Authority (Abilify brand only) see SA153 below – Retail pharmacy	9	30	 Aripiprazole Sandoz Abilify

⇒SA1539 Special Authority for Subsidy

Initial application — (Schizophrenia or related psychoses) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient is suffering from schizophrenia or related psychoses; and
- 2 Either:
 - 2.1 An effective dose of risperidone or quetiapine has been trialled and has been discontinued, or is in the process of being discontinued, because of unacceptable side effects; or
 - 2.2 An effective dose of risperidone or quetiapine has been trialled and has been discontinued, or is in the process of being discontinued, because of inadequate clinical response.

Initial application — (Autism spectrum disorder*) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has been diagnosed with an autism spectrum disorder* and has symptoms of severe irritability; and
- 2 An effective dose of risperidone has been trialled and has been discontinued because of unacceptable side effects or inadequate response; and
- 3 The patient is aged less than 18 years.

Renewal — (Schizophrenia or related psychoses) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Autism spectrum disorder*) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indications marked with * are Unapproved Indications

CHLORPROMAZINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency

Tab 10 mg – Up to 30 tab available on a PSO	12.36	100	 Largactil
Tab 25 mg - Up to 30 tab available on a PSO	13.02	100	 Largactil
Tab 100 mg - Up to 30 tab available on a PSO	30.61	100	 Largactil
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO	25.66	10	 Largactil

NERVOUS SYSTEM

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
	Φ	rei	•	Manulaciurei
LOZAPINE – Hospital pharmacy [HP4]				
Safety medicine; prescriber may determine dispensing frequencies				
Tab 25 mg		50		Clozaril
	6.69			Clopine
	11.36	100		Clozaril
	13.37		-	Clopine
Tab 50 mg		50	-	Clopine
	17.33	100		Clopine
Tab 100 mg		50		Clozaril
	17.33			Clopine
	29.45	100		Clozaril
	34.65		✓	Clopine
Tab 200 mg		50		Clopine
	69.30	100		Clopine
Suspension 50 mg per ml		100 m	nl 🗸	Clopine
ALOPERIDOL – Safety medicine; prescriber may determine c	lispensing frequency			
Tab 500 mcg – Up to 30 tab available on a PSO		100	1	Serenace
Tab 1.5 mg – Up to 30 tab available on a PSO		100		Serenace
Tab 5 mg – Up to 30 tab available on a PSO		100		Serenace
Oral lig 2 mg per ml – Up to 200 ml available on a PSO		100 m		Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a F		10		Serenace
EVOMEPROMAZINE HYDROCHLORIDE – Safety medicine;				
Inj 25 mg per ml, 1 ml ampoule		10	~	Wockhardt
VOMEPROMAZINE MALEATE – Safety medicine; prescribe	r may determine dispe	ensing	frequenc	y
Tab 25 mg		100	1	Nozinan
Tab 100 mg	43.96	100	✓	Nozinan
				NOZINAN
THILIM (CARRONATE – Satety medicine: prescriber may dete		LIENCY	u l	NOZITATI
	ermine dispensing freq		,	
Tab 250 mg	ermine dispensing freq	500	 ✓ 	Lithicarb FC
Tab 250 mg Tab 400 mg	ermine dispensing freq 34.30 12.83	500 100	/ /	Lithicarb FC Lithicarb FC
Tab 250 mg Tab 400 mg Tab long-acting 400 mg	ermine dispensing freq 34.30 12.83 19.20	500 100 100	1 1 1	Lithicarb FC Lithicarb FC Priadel
Tab 250 mg Tab 400 mg Tab long-acting 400 mg Cap 250 mg	ermine dispensing freq 34.30 12.83 19.20 9.42	500 100	1 1 1	Lithicarb FC Lithicarb FC
Tab 250 mg Tab 400 mg Tab long-acting 400 mg Cap 250 mg LANZAPINE – Safety medicine; prescriber may determine dis	ermine dispensing freq 	500 100 100 100	1 1 1	<u>Lithicarb FC</u> <u>Lithicarb FC</u> Priadel Douglas
Tab 250 mg Tab 400 mg Tab long-acting 400 mg Cap 250 mg LANZAPINE – Safety medicine; prescriber may determine dis Tab 2.5 mg	ermine dispensing freq 	500 100 100 100 28	י י י י	Lithicarb FC Lithicarb FC Priadel Douglas Zypine
Tab 250 mg Tab 400 mg Tab long-acting 400 mg Cap 250 mg ANZAPINE – Safety medicine; prescriber may determine dis Tab 2.5 mg Tab 5 mg	ermine dispensing freq 	500 100 100 100 28 28 28	5 5 5 5 5	Lithicarb FC Lithicarb FC Priadel Douglas Zypine Zypine
Tab 250 mg Tab 400 mg Tab long-acting 400 mg Cap 250 mg LANZAPINE – Safety medicine; prescriber may determine dis Tab 2.5 mg Tab 5 mg Tab 5 mg Tab orodispersible 5 mg	ermine dispensing freq 	500 100 100 100 28 28 28 28	5 5 5 5 5	Lithicarb FC Lithicarb FC Priadel Douglas Zypine Zypine Zypine ODT
Tab 250 mg Tab 400 mg Tab long-acting 400 mg Cap 250 mg LANZAPINE – Safety medicine; prescriber may determine dis Tab 2.5 mg Tab 5 mg Tab orodispersible 5 mg Tab 10 mg	ermine dispensing freq 	500 100 100 100 28 28 28 28 28 28 28		Lithicarb FC Lithicarb FC Priadel Douglas Zypine Zypine Zypine ODT Zypine
Tab 250 mg Tab 400 mg Tab long-acting 400 mg Cap 250 mg ANZAPINE – Safety medicine; prescriber may determine dis Tab 2.5 mg Tab 5 mg Tab orodispersible 5 mg	ermine dispensing freq 	500 100 100 100 28 28 28 28		Lithicarb FC Lithicarb FC Priadel Douglas Zypine Zypine Zypine ODT
Tab 250 mg Tab 400 mg Tab long-acting 400 mg Cap 250 mg ANZAPINE – Safety medicine; prescriber may determine dis Tab 2.5 mg Tab 5 mg Tab orodispersible 5 mg Tab 10 mg Tab orodispersible 10 mg	ermine dispensing freq 	500 100 100 100 28 28 28 28 28 28 28		Lithicarb FC Lithicarb FC Priadel Douglas Zypine Zypine Zypine ODT Zypine
Tab 250 mg Tab 400 mg Tab long-acting 400 mg Cap 250 mg ANZAPINE – Safety medicine; prescriber may determine dis Tab 2.5 mg Tab 5 mg Tab orodispersible 5 mg Tab 10 mg Tab orodispersible 10 mg Tab orodispersible 10 mg ERICYAZINE – Safety medicine; prescriber may determine di	ermine dispensing freq 	500 100 100 100 28 28 28 28 28 28 28		Lithicarb FC Lithicarb FC Priadel Douglas Zypine Zypine Zypine ODT Zypine
Tab 250 mg Tab 400 mg Tab long-acting 400 mg Cap 250 mg ANZAPINE – Safety medicine; prescriber may determine dis Tab 2.5 mg Tab 5 mg Tab orodispersible 5 mg Tab 10 mg Tab orodispersible 10 mg Tab orodispersible 10 mg ERICYAZINE – Safety medicine; prescriber may determine di Tab 2.5 mg	ermine dispensing freq 	500 100 100 100 28 28 28 28 28 28 28 28 28 28		Lithicarb FC Lithicarb FC Priadel Douglas Zypine Zypine Zypine ODT Zypine ODT Zypine ODT Neulactil
Tab 250 mg Tab 400 mg Tab long-acting 400 mg Cap 250 mg ANZAPINE – Safety medicine; prescriber may determine dis Tab 2.5 mg Tab 5 mg Tab orodispersible 5 mg Tab orodispersible 5 mg Tab orodispersible 10 mg ERICYAZINE – Safety medicine; prescriber may determine di Tab 2.5 mg Tab 10 mg	ermine dispensing freq 	500 100 100 100 28 28 28 28 28 28 28 28		Lithicarb FC Lithicarb FC Priadel Douglas Zypine Zypine Zypine ODT Zypine ODT
Tab 250 mg Tab 400 mg Tab long-acting 400 mg Tab long-acting 400 mg Cap 250 mg Cap 250 mg ANZAPINE – Safety medicine; prescriber may determine dis Tab 2.5 mg Tab orodispersible 5 mg Tab orodispersible 5 mg Tab 10 mg Tab orodispersible 10 mg ERICYAZINE – Safety medicine; prescriber may determine di Tab 2.5 mg Tab 10 mg – Safety medicine; prescriber may determine di Jab 2.5 mg Tab 10 mg – JETIAPINE JETIAPINE – Safety medicine; prescriber may determine dis Jab 10 mg	ermine dispensing freq 	500 100 100 100 28 28 28 28 28 28 28 28 28 100 100	· · · · · · · · · · · · · · · · · · ·	Lithicarb FC Lithicarb FC Priadel Douglas Zypine Zypine Zypine ODT Zypine Zypine ODT Neulactil Neulactil
Tab 250 mg Tab 400 mg Tab long-acting 400 mg Tab long-acting 400 mg Cap 250 mg Cap 250 mg LANZAPINE – Safety medicine; prescriber may determine dis Tab 2.5 mg Tab orodispersible 5 mg Tab orodispersible 5 mg Tab 10 mg Tab orodispersible 10 mg ERICYAZINE – Safety medicine; prescriber may determine di Tab 2.5 mg Tab 10 mg – Safety medicine; prescriber may determine di Tab 2.5 mg JETIAPINE – Safety medicine; prescriber may determine dis Tab 25 mg	ermine dispensing freq 	500 100 100 100 100 28 28 28 28 28 28 28 28 100 100 90	· · · · · · · · · · · · · · · · · · ·	Lithicarb FC Lithicarb FC Priadel Douglas Zypine Zypine Zypine ODT Zypine ODT Zypine ODT Neulactil Neulactil Quetapel
Tab 250 mg Tab 400 mg Tab long-acting 400 mg Cap 250 mg LANZAPINE – Safety medicine; prescriber may determine dis Tab 5 mg Tab orodispersible 5 mg Tab orodispersible 10 mg Tab orodispersible 10 mg Tab orodispersible 10 mg Tab orodispersible 10 mg Tab 10 mg ERICYAZINE – Safety medicine; prescriber may determine dis Tab 10 mg UETIAPINE – Safety medicine; prescriber may determine dis Tab 25 mg Tab 100 mg	ermine dispensing freq 	500 100 100 100 28 28 28 28 28 28 100 100 90 90		Lithicarb FC Lithicarb FC Priadel Douglas Zypine Zypine Zypine ODT Zypine ODT Zypine ODT Neulactil Neulactil Quetapel Quetapel
Tab 400 mg Tab long-acting 400 mg Cap 250 mg LANZAPINE – Safety medicine; prescriber may determine dis Tab 2.5 mg Tab orodispersible 5 mg Tab orodispersible 5 mg Tab orodispersible 10 mg ERICYAZINE – Safety medicine; prescriber may determine di Tab 2.5 mg Tab 10 mg UETIAPINE – Safety medicine; prescriber may determine dis Tab 25 mg	ermine dispensing freq 	500 100 100 100 100 28 28 28 28 28 28 28 28 100 100 90		Lithicarb FC Lithicarb FC Priadel Douglas Zypine Zypine Zypine ODT Zypine ODT Zypine ODT Neulactil Neulactil Quetapel

	Subsidy		Fully Brand or
	(Manufacturer's Price)	Per	Subsidised Generic Manufacturer
	\$	Per	
RISPERIDONE – Safety medicine; prescriber may determine dis	spensing frequency		
Tab 0.5 mg	1.86	60	 Actavis
Tab 1 mg	2.06	60	 Actavis
Tab 2 mg	2.29	60	✓ Actavis
Tab 3 mg	2.50	60	✓ Actavis
Tab 4 mg	3.43	60	✓ Actavis
Oral liq 1 mg per ml	7.66	30 ml	Risperon
IPRASIDONE – Safety medicine; prescriber may determine dis	nensing frequency		
Cap 20 mg		60	Zeldox
0up 20 mg		00	✓ Zusdone
Cap 40 mg	24 75	60	✓ Zusdone
Cap 60 mg		60	✓ Zusdone
Cap 80 mg		60	✓ Zusdone
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pre	•		
Tab 10 mg		100	 Clopixol
Denet Inications			
Depot Injections			
LUPENTHIXOL DECANOATE - Safety medicine; prescriber m	av determine dispen	sina fr	requency
Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO		5	✓ Fluanxol
Inj 20 mg per ml, 2 ml – Up to 5 inj available on a PSO		5	✓ Fluanxol
Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO		5	✓ Fluanxol
		-	
ALOPERIDOL DECANOATE – Safety medicine; prescriber ma		•	
Inj 50 mg per ml, 1 ml – Up to 5 inj available on a PSO		5	✓ Haldol
Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO		5	 Haldol Concentrate
			 Haldol
			Decanoas S29
DLANZAPINE – Special Authority see SA1428 below – Retail ph	narmacy		
Safety medicine; prescriber may determine dispensing freque			
Inj 210 mg vial	•	1	 Zyprexa Relprevv
Inj 300 mg vial		1	 Zyprexa Relprevv
Inj 405 mg vial		1	✓ Zyprexa Relprevv
		•	-16.000

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer
PALIPERIDONE – Special Authority see SA1429 below – Retai Safety medicine: prescriber may determine dispensing frequ				Manufacturor
Inj 25 mg syringe	,	1	🖌 In	ivega Sustenna
Inj 50 mg syringe		1		ivega Sustenna
Inj 75 mg syringe		1	🖌 In	ivega Sustenna
Inj 100 mg syringe		1	🗸 In	ivega Sustenna
Inj 150 mg syringe		1	🗸 In	ivega Sustenna

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

PIPOTHIAZINE PALMITATE - Subsidy by endorsement

- a) Safety medicine; prescriber may determine dispensing frequency
- b) Subsidised for patients who were taking pipothiazine palmitate prior to 1 August 2014 and the prescription or PSO is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of pipothiazine palmitate.

Inj 50 mg per ml, 1 ml – Up to 5 inj available on a PSO 178.48	10	 Piportil
Inj 50 mg per ml, 2 ml – Up to 5 inj available on a PSO	10	 Piportil
(Piportil Inj 50 mg per ml, 1 ml to be delisted 1 June 2019)		
(Piportil Inj 50 mg per ml, 2 ml to be delisted 1 June 2019)		
RISPERIDONE – Special Authority see SA1427 below – Retail pharmacy		
Safety medicine; prescriber may determine dispensing frequency		
Inj 25 mg vial	1	Risperdal Consta
Inj 37.5 mg vial	1	 Risperdal Consta
Inj 50 mg vial217.56	1	 Risperdal Consta

■ SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

‡ safety cap

if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

(Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	
JCLOPENTHIXOL DECANOATE – Safety medicine; prescriber n Inj 200 mg per ml, 1 ml – Up to 5 inj available on a PSO		ensino 5		cy Clopixol
Anxiolytics				
JSPIRONE HYDROCHLORIDE				
Tab 5 mg		100		Orion
Tab 10 mg		100	~	Orion
ONAZEPAM - Safety medicine; prescriber may determine dispe				
Tab 500 mcg	5.64	100	~	Paxam
Paxam to be Sole Supply on 1 July 2018	10.79	100		Paxam
Tab 2 mg Paxam to be Sole Supply on 1 July 2018	10.76	100	•	Faxaiii
AZEPAM – Safety medicine; prescriber may determine dispensi	na frequency			
Tab 2 mg	0 1 2	500	1	Arrow-Diazepam
‡ Safety cap for extemporaneously compounded oral liquid				
Tab 5 mg		500	~	Arrow-Diazepam
‡ Safety cap for extemporaneously compounded oral liquid	preparations.			
DRAZEPAM – Safety medicine; prescriber may determine disper				
Tab 1 mg		250	~	Ativan
\$\$ \$\$ \$\$ \$\$ \$\$ \$\$ \$\$ \$\$ \$\$ \$\$ \$\$ \$\$		100		Ativon
Tab 2.5 mg ‡ Safety cap for extemporaneously compounded oral liquid		100	•	Ativan
XAZEPAM – Safety medicine; prescriber may determine dispens Tab 10 mg		100	1	Ox-Pam
\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$		100	•	
Tab 15 mg		100	✓	Ox-Pam
‡ Safety cap for extemporaneously compounded oral liquid				
Aultiple Sclerosis Treatments				
METHYL FUMARATE – Special Authority see SA1559 below – I	Retail pharmacy			
Wastage claimable – see rule 3.3.2 on page 13 Cap 120 mg	520.00	14	1	Tecfidera
Cap 240 mg		56		Tecfidera
SA1559 Special Authority for Subsidy	,			
becial Authority approved by the Multiple Sclerosis Treatment Co	nmittee			
otes: Special Authority approved by the Multiple Sclerosis Treatn		ommi	ttee (MST	AC). Applications will be

considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator	Phone: 04 460 4990
Multiple Sclerosis Treatment Assessment Committee	Facsimile: 04 916 7571
PHARMAC PO Box 10 254	Email: mstaccoordinator@pharmac.govt.nz
Wellington	

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	lised	Generic	
\$	Per	✓	Manufacturer	

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified). Entry Criteria

- a) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- b) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- c) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- d) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- e) applications must be made by the patient's neurologist or general physician; and
- f) patients must have no previous history of lack of response to dimethyl fumarate; and
- g) patients must have not previously had intolerance to dimethyl fumarate; and
- h) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to dimethyl fumarate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not

continued...

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Per Sub	sidised	Generic Manufacturer
continued	•			
met and both fingolimod and natalizumab are either not to	plerated or treatment with bo	th agents	would b	e clinically inappropriate.
Continued relapses on treatment would be expected to le		•		, ,, ,
a relapse has resulted in an increased EDSS score that p			of treatn	nent according to stopping
criteria, a period of 6 months is allowed from the start of t	, ,	cur.		
FINGOLIMOD – Special Authority see SA1562 below – F	Retail pharmacy			
Wastage claimable – see rule 3.3.2 on page 13 Cap 0.5 mg	2 650 00	28	10	ailenva
SA1562 Special Authority for Subsidy		20		anonyu
Special Authority approved by the Multiple Sclerosis Trea	tment Committee			
Notes: Special Authority approved by the Multiple Sclero		Committee	(MSTA	C). Applications will be
considered by MSTAC at its regular meetings and approv	ved subject to eligibility accord	rding to the	e Entry a	and Stopping criteria
(below).				
Application details may be obtained from PHARMAC's we		ovt.nz or:		
The coordinator	Phone: 04 460 4990			
Multiple Sclerosis Treatment Assessment Committee	Facsimile: 04 916 7571			
PHARMAC PO Box 10 254	Email: mstaccoordinator	pharmac	.govt.nz	
Wellington				
Completed application forms must be sent to the coordina	ator for MSTAC and will be c	onsidered	by MST	AC at the next practicable

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	1	Manufacturer	

- 6) patients must have no previous history of lack of response to fingolimod; and
- 7) patients must have not previously had intolerance to fingolimod; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to fingolimod; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

NATALIZUMAB – Special Authority see SA1563 below – Retail pharmacy

Inj 20 mg per ml, 15 ml vial...... 1,750.00 1 🗸 Tysabri

⇒SA1563 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator	Phone: 04 460 4990
Multiple Sclerosis Treatment Assessment Committee	Facsimile: 04 916 7571
PHARMAC PO Box 10 254	Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:

*Three months or six months, as applicable, dispensed all-at-once

 Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and

continued...

	Subsidy	Full	
(Mar	ufacturer's Price)	Subsidise	d Generic
	\$	Per 🖌	Manufacturer

- Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) treatment must be initiated and supervised by a neurologist who is registered in the Tysabri Australasian Prescribing Programme operated by the supplier; and
- 7) patients must have no previous history of lack of response to natalizumab; and
- 8) patients must have not previously had intolerance to natalizumab; and
- 9) a) Patient is JC virus negative, or
 - b) Patient is JC virus positive and has given written informed consent acknowledging an understanding of the risk of progressive multifocal leucoencephalopathy (PML) associated with natalizumab
- 10) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to natalizumab; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier.

Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate.

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
continued Continued relapses on treatment would be expected to le a relapse has resulted in an increased EDSS score that j criteria, a period of 6 months is allowed from the start of 1	potentially may lead to disconti the relapse for recovery to occ	inuation of treatr	
TERIFLUNOMIDE – Special Authority see SA1560 belo Wastage claimable – see rule 3.3.2 on page 13 Tab 14 mg		28 🗸 J	Aubagio
► SA1560 Special Authority for Subsidy Special Authority approved by the Multiple Sclerosis Trea Notes: Special Authority approved by the Multiple Sclero considered by MSTAC at its regular meetings and approv (below). Application details may be obtained from PHARMAC's w	osis Treatment Assessment Co ved subject to eligibility accord	ing to the Entry	,
The coordinator Multiple Sclerosis Treatment Assessment Committee PHARMAC PO Box 10 254 Wellington	Phone: 04 460 4990 Facsimile: 04 916 7571 Email: <u>mstaccoordinator@</u>	pharmac.govt.nz	<u> </u>
Completed application forms must be sent to the coordin opportunity. Notification of MSTAC's decision will be sent to the patient		,	

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5° C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) patients must have no previous history of lack of response to teriflunomide; and

continued...

NERVOUS SYSTEM

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price)	Fu Subsidise		
\$	Per	 Manufacturer 	

- 7) patients must have not previously had intolerance to teriflunomide; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to teriflunomide; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

Other Multiple Sclerosis Treatments

⇒SA1564 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator	Phone: 04 460 4990
Multiple Sclerosis Treatment Assessment Committee	Facsimile: 04 916 7571
PHARMAC PO Box 10 254	Email: mstaccoordinator@pharmac.govt.nz
Wellington	

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

These agents will NOT be subsidised if dispensed from a community or hospital pharmacy. Regular supplies will be distributed to all approved patients or their clinicians by courier.

Prescribers must send quarterly prescriptions for approved patients to the MSTAC coordinator.

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, or 20 mg glatiramer acetate daily will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. The MSTAC

Subsidy	Fully	/ Brand or	
(Manufacturer's Price)	Subsidised	d Generic	
\$	Per 🗸	Manufacturer	

coordinator should be notified of the change and a new prescription provided.

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or
 - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

*Three months or six months, as applicable, dispensed all-at-once

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment. Progression of disability is defined as progress by any of the following EDDSS Points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

continued...

\$ safety cap

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
\$	Per	1	Manufacturer

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

GLATIRAMER ACETATE - Special Authority see SA1564 on p	age 154 – [Xpharm	ן]	
Inj 20 mg prefilled syringe	1,089.25	28	 Copaxone
INTERFERON BETA-1-ALPHA - Special Authority see SA1564	4 on page 154 – [X	pharm]	
Inj 6 million iu prefilled syringe	1,170.00	4	 Avonex
Injection 6 million iu per 0.5 ml pen injector	1,170.00	4	 Avonex Pen
INTERFERON BETA-1-BETA - Special Authority see SA1564	on page 154 – [Xpł	harm]	
Inj 8 million iu per 1 ml	1,322.89	15	 Betaferon
O a da l'anna ann d dhanna a l'an			

Sedatives and Hypnotics

LORMETAZEPAM – Safety medicine; prescriber may de	etermine dispensing frequen	су		
Tab 1 mg		30		
	(23.50)		Noctamid	
‡ Safety cap for extemporaneously compounded of	oral liquid preparations.			
(Noctamid Tab 1 mg to be delisted 1 December 2018)				
MELATONIN - Special Authority see SA1666 below - R	etail pharmacy			
Tab modified-release 2 mg - No more than 5 tab pe	r day	30	 Circadin 	
	-			

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are Unapproved Indications.

NERVOUS SYSTEM

	Subsidy		Fully	
	(Manufacturer's Price)	_	Subsidised	
	\$	Per		Manufacturer
MIDAZOLAM - Safety medicine; prescriber may determine dispe	nsing frequency			
Inj 1 mg per ml, 5 ml ampoule	4.30	10	✓	Midazolam-Claris
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available				
on a PSO		10	1	Pfizer
On a PSO for status epilepticus use only. PSO must be	endorsed for status e	epiler	ticus use	only.
Inj 5 mg per ml, 3 ml ampoule	2.50	5		Midazolam-Claris
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj available				
a PSO		5	1	Pfizer
On a PSO for status epilepticus use only. PSO must be	endorsed for status e	epilep	ticus use	only.
NITRAZEPAM - Safety medicine; prescriber may determine disp	ensing frequency			
Tab 5 mg	0 1 7	100	1	Nitrados
‡ Safety cap for extemporaneously compounded oral liquid				
PHENOBARBITONE SODIUM – Special Authority see SA1386 b		0.01/		
				Mautin dala ana
Inj 200 mg per ml, 1 ml ampoule		10	•	Martindale S29
SA1386 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals valid	I without further rene	wal u	nless noti	fied for applications meeting
the following criteria:				
Both:				
1 For the treatment of terminal agitation that is unresponsive	to other agents; and	ł		
2 The applicant is part of a multidisciplinary team working in	palliative care.			
TEMAZEPAM - Safety medicine; prescriber may determine dispe	ansing frequency			
Tab 10 mg		25	1	Normison
\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$		20	•	NOTHISON
TRIAZOLAM – Safety medicine; prescriber may determine disper	• • •			
Tab 125 mcg		100		
	(9.85)			Hypam
‡ Safety cap for extemporaneously compounded oral liquid Tel: 050 manual		400		
Tab 250 mcg		100		
	(11.20)			Hypam
‡ Safety cap for extemporaneously compounded oral liquid				
ZOPICLONE - Safety medicine; prescriber may determine disper	nsing frequency			
Tab 7.5 mg	8.99	500	✓	Zopiclone Actavis
Stimulants/ADHD Treatments				
ATOMOVETINE Creatial Authority and CA1416 below Dataily				
ATOMOXETINE – Special Authority see SA1416 below – Retail g Cap 10 mg		28		Strattera
		20 28		Strattera
Cap 18 mg		28 28		
Cap 25 mg		28 28		Strattera Strattera
Cap 40 mg		28 28		Strattera
Cap 60 mg	107.03	20	•	Surallera

⇒SA1416 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder) diagnosed according to DSM-IV or ICD 10 criteria; and

continued...

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

28 28 Strattera

✓ Strattera

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	✓	Manufacturer

- 2 Once-daily dosing; and
- 3 Any of the following:
 - 3.1 Treatment with a subsidised formulation of a stimulant has resulted in the development or worsening of serious adverse reactions or where the combination of subsidised stimulant treatment with another agent would pose an unacceptable medical risk; or
 - 3.2 Treatment with a subsidised formulation of a stimulant has resulted in worsening of co-morbid substance abuse or there is a significant risk of diversion with subsidised stimulant therapy; or
 - 3.3 An effective dose of a subsidised formulation of a stimulant has been trialled and has been discontinued because of inadequate clinical response; or
 - 3.4 Treatment with a subsidised formulation of a stimulant is considered inappropriate because the patient has a history of psychoses or has a first-degree relative with schizophrenia; and
- 4 The patient will not be receiving treatment with atomoxetine in combination with a subsidised formulation of a stimulant, except for the purposes of transitioning from subsidised stimulant therapy to atomoxetine.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: A "subsidised formulation of a stimulant" refers to currently subsidised methylphenidate hydrochloride tablet formulations (immediate-release, sustained-release and extended-release) or dexamfetamine sulphate tablets.

DEXAMFETAMINE SULFATE - Special Authority see SA1149 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab 5 mg17.00 100 🗸 PSM

➡SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and

2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

1 The treatment remains appropriate and the patient is benefiting from treatment; and

2 Either:

- 2.1 Applicant is a paediatrician or psychiatrist; or
- 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

 Subsidy (Manufacturer's Price)		Fully lised	Brand or Generic
\$	Per	✓	Manufacturer

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1150 below - Retail pharmacy

a) Only on a controlled drug form

b) Safety medicine; prescriber may determine dispensing frequency

Tab immediate-release 5 mg		30	 Rubifen
Tab immediate-release 10 mg		30	 Ritalin
ç			 Rubifen
Tab immediate-release 20 mg	7.85	30	 Rubifen
Tab sustained-release 20 mg		30	 Rubifen SR
Ŭ	50.00	100	 Ritalin SR

⇒SA1150 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic r I Manufacturer
METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEAS	E – Special Authorit	y see	e SA1151 below – Retail pharmacy
a) Only on a controlled drug form			
b) Safety medicine; prescriber may determine dispensing free	equency		
Tab extended-release 18 mg		30	 Concerta
Tab extended-release 27 mg	65.44	30	 Concerta
Tab extended-release 36 mg	71.93	30	 Concerta
Tab extended-release 54 mg		30	 Concerta
Cap modified-release 10 mg		30	
Cap modified-release 20 mg		30	
Cap modified-release 30 mg		30	
Cap modified-release 40 mg		30	 Ritalin LA

⇒SA1151 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL – Special Authority see SA1126 below – Retail pharmacy

⇒SA1126 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and

2 Either:

- 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
- 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and

3 Either:

Subsidy (Manufacturer's Price)	SI	Fully ubsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

- 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
- 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia			
DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	4.34	90	 Donepezil-Rex
* Tab 10 mg	6.64	90	 Donepezil-Rex
RIVASTIGMINE - Special Authority see SA1488 below - Retail	pharmacy		
Patch 4.6 mg per 24 hour		30	 Exelon
Patch 9.5 mg per 24 hour	90.00	30	 Exelon

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 2 mg with naloxone 0.5 mg	57.40	28	 Suboxone
Tab sublingual 8 mg with naloxone 2 mg	166.00	28	 Suboxone

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal --- (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following

continued...

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fu	Illy Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufactu 	Jrer

criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE Tab modified-release 150 mg11.00 30	✓ Zyban
DISULFIRAM Tab 200 mg44.30 100	✓ Antabuse
NALTREXONE HYDROCHLORIDE – Special Authority see SA1408 below – Retail pharm Tab 50 mg112.55 30	

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
NICOTINE				
a) Nicotine will not be funded under the Dispensing Frequence	cy Rule in amounts le	ess tha	an 4 week	s of treatment.
b) Note: may be provided by a pharmacist under the non-pre-		's prov	risions in F	Part III of Section A.
Patch 7 mg – Up to 28 patch available on a PSO		28	✓	Habitrol
Patch 7 mg for direct distribution only – [Xpharm]	3.94	7	✓	Habitrol
Patch 14 mg – Up to 28 patch available on a PSO		28	✓	Habitrol
Patch 14 mg for direct distribution only - [Xpharm]	4.52	7	✓	Habitrol
Patch 21 mg – Up to 28 patch available on a PSO		28	✓	Habitrol
Patch 21 mg for direct distribution only - [Xpharm]	5.18	7	✓	Habitrol
Lozenge 1 mg - Up to 216 loz available on a PSO		216	✓	Habitrol
Lozenge 1 mg for direct distribution only - [Xpharm]	3.20	36	✓	Habitrol
Lozenge 2 mg - Up to 216 loz available on a PSO		216	✓	Habitrol
Lozenge 2 mg for direct distribution only - [Xpharm]	3.24	36	✓	Habitrol
Gum 2 mg (Fruit) - Up to 384 piece available on a PSO		384	✓	Habitrol
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]		96	✓	Habitrol
Gum 2 mg (Mint) - Up to 384 piece available on a PSO		384	✓	Habitrol
Gum 2 mg (Mint) for direct distribution only - [Xpharm]		96	✓	Habitrol
Gum 4 mg (Fruit) - Up to 384 piece available on a PSO		384	✓	Habitrol
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]		96	1	Habitrol
Gum 4 mg (Mint) – Up to 384 piece available on a PSO		384		Habitrol
Gum 4 mg (Mint) for direct distribution only – [Xpharm]		96		Habitrol

VARENICLINE TARTRATE - Special Authority see SA1575 below - Retail pharmacy

a) Varenicline will not be funded under the Dispensing Frequency Rule in amounts less than 2 weeks of treatment.

b) A maximum of 12 weeks' varenicline will be subsidised on	each Special Au	uthority appro	val, including the starter pack
Tab 1 mg	67.74	28	 Champix
	135.48	56	 Champix
Tab 0.5 mg × 11 and 1 mg × 14	60.48	25 OP	 Champix

➡SA1575 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not used funded varenicline in the last 12 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and

*Three months or six months, as applicable, dispensed all-at-once

7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	✓	Manufacturer

- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 The patient has not used funded varenicline in the last 12 months; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 12 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval. This includes the 2-week 'starter' pack.

	Subsidy	Fully	Brand or
	(Manufacturer's Price) \$	Subsidised Per 🖌	Generic Manufacturer
Chemotherapeutic Agents			
Alkylating Agents			
BENDAMUSTINE HYDROCHLORIDE – PCT only – Specialist Inj 25 mg vial Inj 100 mg vial Inj 1 mg for ECP		1 ✓ R 1 ✓ R	ibomustin ibomustin axter
► SA1667 Special Authority for Subsidy Initial application — (treatment naive CLL) only from a relevant relevant specialist. Approvals valid for 12 months for application All of the following:			he recommendation of a
 The patient has Binet stage B or C, or progressive stage The patient is chemotherapy treatment naive; and The patient is unable to tolerate toxicity of full-dose FCR Patient has ECOG performance status 0-2; and Patient has a Cumulative Illness Rating Scale (CIRS) sca Bendamustine is to be administered at a maximum dose 6 cycles. 	; and pre of < 6; and	·	
Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lym to comprise a known standard therapeutic chemotherapy regime Initial application — (Indolent, Low-grade lymphomas) only recommendation of a relevant specialist. Approvals valid for 9 r All of the following:	en and supportive treat from a relevant specia	ments. list or medical pra	actitioner on the
 The patient has indolent low grade NHL requiring treatment Patient has a WHO performance status of 0-2; and Either: 	ent; and		
 3.1 Both: 3.1.1 Patient is treatment naive; and 3.1.2 Bendamustine is to be administered for a CD20+); or 	maximum of 6 cycles (i	n combination wi	th rituximab when
 3.2 All of the following: 3.2.1 Patient has relapsed refractory disease fol 3.2.2 The patient has not received prior bendarr 3.2.3 Either: 3.2.3.1 Both: 		apy; and	
3.2.3.1 Both. 3.2.3.1.1 Bendamustine is to be admin combination with rituximab w 3.2.3.1.2 Patient has had a rituximab tu 3.2.3.2 Bendamustine is to be administered refractory patients.	hen CD20+); and reatment-free interval c	f 12 months or m	ore; or

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Either:
 - 2.1 Both:

*Three months or six months, as applicable, dispensed all-at-once

continued...

‡ safety cap

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy	\ ^	Fully Brand or
	(Manufacturer's Prie	ce) Subs Per	sidised Generic Manufacturer
ontinued			
2.1.1 Bendamustine is to be administered for a	a maximum of 6 cycle	es in relapsed	I patients (in combination with
rituximab when CD20+); and		·	
2.1.2 Patient has had a rituximab treatment-free	ee interval of 12 mont	ths or more; c	or
2.2 Bendamustine is to be administered as a mono	therapy for a maximu	m of 6 cycles	in rituximab refractory patien
ote: 'indolent, low-grade lymphomas' includes follicular, mar	ntle cell, marginal zon	e and lympho	oplasmacytic/ Waldenstrom's
nacroglobulinaemia.			
BUSULFAN – PCT – Retail pharmacy-Specialist			
Tab 2 mg		100	 Myleran
CARBOPLATIN – PCT only – Specialist			
Inj 10 mg per ml, 5 ml vial		1	 DBL Carboplatin
	20.00		 Carboplatin Ebewe
Inj 10 mg per ml, 15 ml vial		1	DBL Carboplatin
	19.50		 Carbaccord
	22.50		 Carboplatin Ebewe
Inj 10 mg per ml, 45 ml vial		1	 DBL Carboplatin
	48.50		 Carbaccord
	50.00		 Carboplatin Ebewe
Inj 1 mg for ECP	0.08	1 mg	 Baxter
ARMUSTINE – PCT only – Specialist			
Inj 100 mg vial	532.00	1	BiCNU
Inj 100 mg for ECP	532.00	100 mg OP	 Baxter
CHLORAMBUCIL – PCT – Retail pharmacy-Specialist			
Tab 2 mg		25	 Leukeran FC
CISPLATIN – PCT only – Specialist			
Inj 1 mg per ml, 50 ml vial	12 20	1	 DBL Cisplatin
	15.00	'	 Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial		1	 Cisplatin Ebewe
	22.46	·	✓ DBL Cisplatin
Inj 1 mg for ECP		1 mg	✓ Baxter
YCLOPHOSPHAMIDE		0	
Tab 50 mg – PCT – Retail pharmacy-Specialist	70.00	50	Endoxan S29
Tab 50 mg - PCT - Relair pharmacy-Specialist			
Western elsimptile and rule 2.2.0 on pore 12	158.00	100	Procytox S29
Wastage claimable – see rule 3.3.2 on page 13 Inj 1 g vial – PCT – Retail pharmacy-Specialist	25.02	1	Endoxan
inj i g viai – POT – Netali phannacy-Specialist		6	 ✓ Endoxan ✓ Cytoxan
Inj 2 g vial – PCT only – Specialist		1	
Inj 1 mg for ECP – PCT only – Specialist		1 mg	✓ Baxter
FOSFAMIDE – PCT only – Specialist		9	Bunton
Inj 1 g	06.00	1	✓ Holoxan
Inj 2 g		1	✓ Holoxan
Inj 2 g Inj 1 mg for ECP		-	✓ Baxter
, -		1 mg	
OMUSTINE – PCT – Retail pharmacy-Specialist	400 50	00	
Cap 10 mg		20	✓ CeeNU
Cap 40 mg		20	CeeNU
1ELPHALAN			
Tab 2 mg – PCT – Retail pharmacy-Specialist		25	 Alkeran
Inj 50 mg – PCT only – Specialist	67 80	1	 Alkeran

Subsidy acturer's Price \$) Su Per	Fully ubsidised	Generic
3.32	1	~	Oxaliccord
5.32	1	~	Oxaliplatin Actavis 50
5.00		1	Oxaliplatin Ebewe
5.01	1	~	Oxaliplatin Actavis 100
0.00		1	Oxaliplatin Ebewe
6.00	1		Oxaliccord
0.18	1 mg	1	Baxter
	•		
BS	1	1	Bedford S29
			THIO-TEPA S29
			Tepadina S29
DC	4		•
BS	1	•	Tepadina S29
elow 5.00	1		Vidaza Baxter
	0	0 1	0 1 🗸

⇒SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- 4 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 No evidence of disease progression; and

*Three months or six months, as applicable, dispensed all-at-once

2 The treatment remains appropriate and patient is benefitting from treatment.

	Subsidy (Manufacturer's Pric		Fully Brand or osidised Generic
	\$	Per	 Manufacturer
CALCIUM FOLINATE Tab 15 mg – PCT – Retail pharmacy-Specialist		10	 DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml – PCT – Retail pharmacy-Specialist		5	✓ Hospira
Inj 10 mg per ml, 5 ml vial – PCT – Retail pharmacy-Special		1	 Calcium Folinate Sandoz
Inj 50 mg - PCT - Retail pharmacy-Specialist		5	 Calcium Folinate Ebewe
Inj 10 mg per ml, 10 ml vial – PCT only – Specialist	7.30	1	 Calcium Folinate Sandoz
Inj 100 mg – PCT only – Specialist		1	 Calcium Folinate Ebewe
Inj 300 mg – PCT only – Specialist		1	 Calcium Folinate Ebewe
Inj 10 mg per ml, 35 ml vial – PCT only – Specialist		1	 Calcium Folinate Sandoz
Inj 1 g – PCT only – Specialist		1	 Calcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial – PCT only – Specialist		1	 Calcium Folinate Sandoz
Inj 1 mg for ECP – PCT only – Specialist	0.06	1 mg	 Baxter
CAPECITABINE – Retail pharmacy-Specialist			
Tab 150 mg		60 120	✓ <u>Brinov</u> ✓ Brinov
	02.20	120	
CLADRIBINE – PCT only – Specialist Inj 1 mg per ml, 10 ml	5 2/0 72	7	 Leustatin
Inj 10 mg for ECP		7 10 mg OP	✓ Baxter
YTARABINE	list 400.00	5	✓ Pfizer
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 10 ml vial – PCT – Retail pharmacy-Spec Inj 100 mg per ml, 20 ml vial – PCT – Retail		5 1	✓ Pfizer
pharmacy-Specialist		1	 Pfizer
Inj 1 mg for ECP – PCT only – Specialist		10 mg	✓ Baxter
Inj 100 mg intrathecal syringe for ECP – PCT only – Special Pfizer Inj 100 mg per ml, 10 ml vial to be delisted 1 October 201	list80.00 1	100 mg OP	✓ Baxter
LUDARABINE PHOSPHATE	412.00	20	Ludara Oral
Tab 10 mg – PCT – Retail pharmacy-Specialist Inj 50 mg vial – PCT only – Specialist		20 5	 ✓ <u>Fludara Oral</u> ✓ Fludarabine Ebewe
Inj 50 mg for ECP – PCT only – Specialist		50 mg OP	 ✓ Pludarabilie Ebewe ✓ Baxter
LUOROURACIL			
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist	10.00	1	 Fluorouracil Ebewe
Inj 50 mg per ml, 50 ml vial – PCT only – Specialist		1	 Fluorouracil Ebewe
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist		1	✓ Fluorouracil Ebewe
Inj 1 mg for ECP – PCT only – Specialist		100 mg	✓ Baxter

	Subsidy	-) 0	Fully Brand or	
	(Manufacturer's Pric \$	e) Sub Per	sidised	
GEMCITABINE HYDROCHLORIDE – PCT only – Specialist				
Inj 1 g, 26.3 ml vial	62.50	1	1	DBL Gemcitabine
Inj 1 g		1	✓	Gemcitabine Ebewe
	349.20		✓	Gemzar
Inj 200 mg	8.36	1	✓	Gemcitabine Ebewe
	78.00		✓	Gemzar
Inj 1 mg for ECP	0.02	1 mg	✓	Baxter
RINOTECAN HYDROCHLORIDE – PCT only – Specialist				
Inj 20 mg per ml, 2 ml vial	11.50	1	✓	Irinotecan Actavis
				40
	41.00			Camptosar
				Irinotecan-Rex
Inj 20 mg per ml, 5 ml vial	17.80	1	•	Irinotecan Actavis 100
	100.00		1	Camptosar
				Irinotecan-Rex
Inj 1 mg for ECP	0.19	1 mg	✓	Baxter
/ERCAPTOPURINE		č		
Tab 50 mg – PCT – Retail pharmacy-Specialist	49.41	25	1	Puri-nethol
Oral suspension 20 mg per ml – Retail pharmacy-Specialis		_5		
Special Authority see SA1725 below		100 ml OP	✓	Allmercap

➡SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

169

(Mar	Subsidy nufacturer's Price) \$	Per	Fully Subsidised	Generic
ETHOTREXATE				
Tab 2.5 mg – PCT – Retail pharmacy-Specialist	3.18	30	1	Trexate
Tab 10 mg - PCT - Retail pharmacy-Specialist	21.00	50	✓	Trexate
Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist	47.50	5	✓	Hospira
Inj 7.5 mg prefilled syringe	14.61	1	1	Methotrexate Sandoz
Inj 10 mg prefilled syringe	14.66	1	1	Methotrexate Sandoz
Inj 15 mg prefilled syringe	14.77	1	1	Methotrexate Sandoz
Inj 20 mg prefilled syringe	14.88	1	1	Methotrexate Sandoz
Inj 25 mg prefilled syringe	14.99	1	1	Methotrexate Sandoz
Inj 30 mg prefilled syringe	15.09	1	1	Methotrexate Sandoz
Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist	30.00	5	1	DBL Methotrexate Onco-Vial
Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist	45.00	1	1	DBL Methotrexate Onco-Vial
Inj 100 mg per ml, 10 ml – PCT – Retail pharmacy-Specialist Inj 100 mg per ml, 50 ml vial – PCT – Retail	25.00	1	1	Methotrexate Ebewe
pharmacy-Specialist	79.99	1	1	Methotrexate Ebewe
Inj 1 mg for ECP – PCT only – Specialist		1 mg	✓	Baxter
Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist	4.73 5	i mg Č	P 🗸	Baxter
METREXED - PCT only - Specialist - Special Authority see SA16		-		
Inj 100 mg vial		1	1	Juno Pemetrexed
Inj 500 mg vial		1	1	Juno Pemetrexed
Inj 1 mg for ECP		1 mg	1	Baxter

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and

2 Either:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	ubsidised	Generic
 \$	Per	1	Manufacturer

continued...

- 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
- 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

THIOGUANINE - PCT - Retail pharmacy-Specialist

Tab 40 mg	25	✓ Lanvis
Other Cytotoxic Agents		
AMSACRINE – PCT only – Specialist		
Inj 50 mg per ml, 1.5 ml ampoule1,500.00	6	 Amsidine S29
Inj 75 mg1,250.00	5	AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-Specialist		
Cap 0.5 mgCBS	100	 Agrylin S29
		Teva S29
ARSENIC TRIOXIDE – PCT only – Specialist		
Inj 10 mg4,817.00	10	✓ AFT \$29
BLEOMYCIN SULPHATE – PCT only – Specialist		
Inj 15,000 iu, vial	1	 DBL Bleomycin Sulfate
Inj 1,000 iu for ECP11.64	1,000 iu	 Baxter
BORTEZOMIB – PCT only – Specialist – Special Authority see SA1576 below		
Inj 3.5 mg vial1,892.50	1	 Velcade
Inj 1 mg for ECP594.77	1 mg	 Baxter

⇒SA1576 Special Authority for Subsidy

Initial application — (Treatment naive multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 Either:

- 1.1 The patient has treatment-naive symptomatic multiple myeloma; or
- 1.2 The patient has treatment-naive symptomatic systemic AL amyloidosis *; and
- 2 Maximum of 9 treatment cycles.

Note: Indications marked with * are Unapproved Indications.

*Three months or six months, as applicable, dispensed all-at-once

Initial application — (Relapsed/refractory multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following

continued...

Subsidy	Full	/ Brand or
(Manufacturer's Price)	Subsidised	d Generic
\$	Per 🗸	Manufacturer

criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has relapsed or refractory multiple myeloma; or
 - 1.2 The patient has relapsed or refractory systemic AL amyloidosis *; and
- 2 The patient has received only one prior front line chemotherapy for multiple myeloma or amyloidosis; and
- 3 The patient has not had prior publicly funded treatment with bortezomib; and
- 4 Maximum of 4 treatment cycles.
- Note: Indications marked with * are Unapproved Indications.

Renewal — (Relapsed/refractory multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 The patient's disease obtained at least a partial response from treatment with bortezomib at the completion of cycle 4; and
- 2 Maximum of 4 further treatment cycles (making a total maximum of 8 consecutive treatment cycles).

Notes: Responding relapsed/refractory multiple myeloma patients should receive no more than 2 additional cycles of treatment beyond the cycle at which a confirmed complete response was first achieved. A line of therapy is considered to comprise either:

a) a known therapeutic chemotherapy regimen and supportive treatments; or

b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments.

Refer to datasheet for recommended dosage and number of doses of bortezomib per treatment cycle.

COLASPASE [L-ASPARAGINASE] – PCT only – Specialist			
Inj 10,000 iu		1	 Leunase
Inj 10,000 iu for ECP		10,000 iu OP	 Baxter
DACARBAZINE – PCT only – Specialist			
Inj 200 mg vial	58.06	1	DBL Dacarbazine
	580.60	10	✓ Dacarbazine
			APP S29
Inj 200 mg for ECP	58.06	200 mg OP	✓ Baxter
		Loo nig or	Duxtor
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist	100 75	4	. Coomonon
Inj 0.5 mg vial		1	✓ Cosmegen
Inj 0.5 mg for ECP		0.5 mg OP	 Baxter
DAUNORUBICIN – PCT only – Specialist			
Inj 2 mg per ml, 10 ml		1	 Pfizer
Inj 20 mg for ECP	130.00	20 mg OP	 Baxter
DOCETAXEL – PCT only – Specialist			
Inj 10 mg per ml, 2 ml vial		1	 DBL Docetaxel
Inj 20 mg		1	 Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial		1	 DBL Docetaxel
Inj 80 mg		1	 Docetaxel Sandoz
Inj 1 mg for ECP	0.55	1 mg	 Baxter
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial		1	 Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	 Doxorubicin Ebewe
3 34- 7	17.00		 Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial		1	 Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	 Doxorubicin Ebewe
	65.00		 Arrow-Doxorubicin
Inj 1 mg for ECP	0.25	1 mg	✓ Baxter
		-	

	Subsidy		Fully	
	(Manufacturer's Price)	-	Subsidised	
	\$	Per	/	Manufacturer
EPIRUBICIN HYDROCHLORIDE – PCT only – Specialist				
Inj 2 mg per ml, 5 ml vial	25.00	1	✓	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	✓	Epirubicin Ebewe
Inj 2 mg per ml, 50 ml vial		1	✓	Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial	65.00	1	✓	Epirubicin Ebewe
Inj 1 mg for ECP	0.36	1 mg	✓	Baxter
ETOPOSIDE				
Cap 50 mg - PCT - Retail pharmacy-Specialist		20	1	Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10	✓	Vepesid
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specia		1	✓	Rex Medical
Inj 1 mg for ECP – PCT only – Specialist		1 mg	1	Baxter
ETOPOSIDE PHOSPHATE – PCT only – Specialist				
Inj 100 mg (of etoposide base)		1	1	Etopophos
Inj 1 mg (of etoposide base) for ECP		1 mg		Baxter
HYDROXYUREA – PCT – Retail pharmacy-Specialist				
Cap 500 mg	31.76	100	1	Hydrea
		100	•	nyulcu
IDARUBICIN HYDROCHLORIDE				_ .
Inj 5 mg vial – PCT only – Specialist		1		Zavedos
Inj 10 mg vial – PCT only – Specialist		1		Zavedos
Inj 1 mg for ECP – PCT only – Specialist		1 mg	v	Baxter
LENALIDOMIDE - Retail pharmacy-Specialist - Special Authori	ty see SA1468 below			
Wastage claimable – see rule 3.3.2 on page 13				
Cap 10 mg	6,207.00	21		Revlimid
Cap 15 mg	7,239.18	21		Revlimid
Cap 25 mg	7,627.00	21	✓	Revlimid

► SA1468 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Patient has relapsed or refractory multiple myeloma with progressive disease; and

2 Either:

2.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or

2.2 Both:

- 2.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
- 2.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 3 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an Unapproved Indication (refer to Interpretations and Definitions). A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

	Subsidy (Manufacturer's Price)	Fully Subsidised	
	(Manulacturer 3 Trice \$	Per		
IESNA				
Tab 400 mg – PCT – Retail pharmacy-Specialist	273.00	50	✓	Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist		50	✓	Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist	161.25	15	✓	Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist		15	✓	Uromitexan
Inj 1 mg for ECP – PCT only – Specialist	2.69	100 m	g 🗸	Baxter
IITOMYCIN C – PCT only – Specialist				
Inj 5 mg vial		1	1	Arrow
Inj 1 mg for ECP		1 mg	-	Baxter
ITOZANTRONE – PCT only – Specialist		5		
Inj 2 mg per ml, 10 ml vial	97 50	1	1	Mitozantrone Ebewe
Inj 1 mg for ECP		1 mg	-	Baxter
, ,		i ing	•	Daxier
ACLITAXEL – PCT only – Specialist	(7.00	-		
Inj 30 mg		5		Paclitaxel Ebewe
Inj 100 mg		1		Paclitaxel Ebewe
	91.67			Paclitaxel Actavis
Inj 150 mg		1		Paclitaxel Ebewe
	137.50			Anzatax
1 1 000				Paclitaxel Actavis
Inj 300 mg		1		Paclitaxel Ebewe
	275.00			Anzatax
	0.40		-	Paclitaxel Actavis
Inj 1 mg for ECP	0.19	1 mg	~	Baxter
EGASPARGASE – PCT only – Special Authority see SA1325	below			
Inj 3,750 IU per 5 ml	3,005.00	1	✓	Oncaspar S29

⇒SA1325 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 The patient has newly diagnosed acute lymphoblastic leukaemia; and

- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and
- 3 Treatment is with curative intent.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has relapsed acute lymphoblastic leukaemia; and

2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and

3 Treatment is with curative intent.

PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Specialist

Inj 10 mgCBS	1	 Nipent S29
PROCARBAZINE HYDROCHLORIDE – PCT – Retail pharmacy-Specialist		
Cap 50 mg498.00	50	 Natulan S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic r ✓ Manufacturer
TEMOZOLOMIDE – Special Authority see SA1616 below – Reta	ail pharmacy		
Cap 5 mg		5	✓ <u>Orion</u>
			Temozolomide
Cap 20 mg		5	✓ <u>Orion</u>
			Temozolomide
			Temizole 20 S29
Cap 100 mg	40.20	5	✓ Orion
			Temozolomide
Cap 140 mg		5	 Orion
			Temozolomide
Cap 250 mg	96.80	5	✓ Orion
			Temozolomide

■SA1616 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the followina:

- 1 Fither:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application - (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine: and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day: and
- 4 Temozolomide to be discontinued at disease progression.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient has glioblastoma multiforme; and
- 1.2 The treatment remains appropriate and the patient is benefitting from treatment: or
- 2 All of the following:

2.1 Patient has anaplastic astrocytoma*; and

- 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 No evidence of disease progression; and

*Three months or six months, as applicable, dispensed all-at-once

2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an Unapproved Indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special Authority see SA1124 on the next page	
Cap 50 mg	🗸 Т

Cap 50 mg	 ·		28	 Thalomid
Cap 100 mg	 	756.00	28	 Thalomid

Subsidy		Fully	Brand or	-
(Manufacturer's Price)	Su	Ibsidised	Generic	
\$	Per	✓	Manufacturer	

■ SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an Unapproved Indication. -----

TRETINOIN Cap 10 mg - PCT - Retail pharmacy-Specialist	100	✓ Vesanoid
VINBLASTINE SULPHATE	100	· vesanora
Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Specialist37.29	1	 Vinblastina
	I	Teva S29
186.46	5	✓ Hospira
Inj 1 mg for ECP – PCT only – Specialist	1 mg	✓ Baxter
(Vinblastina Teva ⁶²⁹ Inj 1 mg per ml, 10 ml vial to be delisted 1 August 2018) VINCRISTINE SULPHATE	, mg	- Bundi
Inj 1 mg per ml, 1 ml vial – PCT – Retail pharmacy-Specialist74.52	5	 DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist85.61	5	 DBL Vincristine Sulfate
Inj 1 mg for ECP – PCT only – Specialist 11.30	1 mg	 Baxter
VINORELBINE – PCT only – Specialist		
Inj 10 mg per ml, 1 ml vial8.00	1	 Navelbine
42.00		 Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial40.00	1	 Navelbine
210.00		 Vinorelbine Ebewe
Inj 1 mg for ECP0.90	1 mg	 Baxter
Protein-tyrosine Kinase Inhibitors		

Kinase innibito

1]		
	60	 Sprycel
6,214.20	60	 Sprycel
7,692.58	60	 Sprycel
6,214.20	30	 Sprycel
	1] 	

SA0976 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz, and prescriptions should be sent to:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
\$	Per	1	Manufacturer

continued...

The CML/GIST Co-ordinator	Phone: (04) 460 4990
PHARMAC	Facsimile: (04) 916 7571
PO Box 10 254	Email: cmlgistcoordinator@pharmac.govt.nz
Wellington	

Special Authority criteria for CML - access by application

- a) Funded for patients with diagnosis (confirmed by a haematologist) of a chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase.
- b) Maximum dose of 140 mg/day for accelerated or blast phase, and 100 mg/day for chronic phase CML.
- c) Subsidised for use as monotherapy only.
- d) Initial approvals valid seven months.
- e) Subsequent approval(s) are granted on application and are valid for six months. The first reapplication (after seven months) should provide details of the haematological response. The third reapplication should provide details of the cytogenetic response after 14-18 months from initiating therapy. All other reapplications should provide details of haematological response, and cytogenetic response if such data is available. Applications to be made and subsequent prescriptions can be written by a haematologist or an oncologist.

Note: Dasatinib is indicated for the treatment of adults with chronic, accelerated or blast phase CML with resistance or intolerance to prior therapy including imatinib.

Guideline on discontinuation of treatment for patients with CML

- a) Prescribers should consider discontinuation of treatment if, after 6 months from initiating therapy, a patient did not obtain a haematological response as defined as any one of the following three levels of response:
 - complete haematologic response (as characterised by an absolute neutrophil count (ANC) > 1.5 × 10⁹/L, platelets > 100 × 10⁹/L, absence of peripheral blood (PB) blasts, bone marrow (BM) blasts < 5% (or FISH Ph+ 0-35% metaphases), and absence of extramedullary disease); or
 - no evidence of leukaemia (as characterised by an absolute neutrophil count (ANC) > 1.0 × 10⁹/L, platelets > 20 × 10⁹/L, absence of peripheral blood (PB) blasts, bone marrow (BM) blasts < 5% (or FISH Ph+ 0-35% metaphases), and absence of extramedullary disease); or
 - return to chronic phase (as characterised by BM and PB blasts < 15%, BM and PB blasts and promyelocytes < 30%, PB basophils < 20% and absence of extramedullary disease other than spleen and liver).
- b) Prescribers should consider discontinuation of treatment if, after 18 months from initiating therapy, a patient did not obtain a major cytogenetic response defined as 0-35% Ph+ metaphases.

ERLOTINIB – Retail pharmacy-Specialist – Special A	uthority see SA1653 below		
Tab 100 mg		30	 Tarceva
Tab 150 mg	1,146.00	30	 Tarceva

► SA1653 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Either:
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

*Three months or six months, as applicable, dispensed all-at-once

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
GEFITINIB – Retail pharmacy-S					
v		1,700.00	30	✓ Ii	ressa
⇒SA1654 Special Authority for Initial application only from a re Approvals valid for 4 months for All of the following:	elevant specialist or medical pra		mend	ation of a re	levant specialist.
All of the following:	and as motostatia unsecontabl	a nan aguamaya Nan	Cmal		Concer (NICCL C), and
2 Either:	ced, or metastatic, unresectabl	e, non-squamous ivon	Smai		Jancer (NSCLC); and
2.1 Patient is treatment	nt naive: or				
2.2 Both:					
	t has discontinued erlotinib due r did not progress whilst on erlo				
3 There is documentation of4 Gefitinib is to be given for	onfirming that disease express a maximum of 3 months.	es activating mutations	s of E	GFR tyrosin	e kinase; and
Renewal only from a relevant sp for 6 months where radiological					
MATINIB MESILATE					
imatinib mesilate (supplied b	registered for the treatment of y Novartis) remains fully subsi- see SA1460 in Section B of the	dised under Special Au	Ithorit	· ·	,
Tab 100 mg - Special Auth	ority see SA1460 below -				
		,	60	-	Alivec
 Cap 100 mg Cap 400 mg 			60 30		<u>matinib-AFT</u> matinib-AFT
⇒SA1460 Special Authority f	or Subsidy				
Special Authority approved by th					
Notes: Application details may t sent to:	be obtained from PHARMAC's	website <u>http://www.pha</u>	rmac	<u>.govt.nz</u> , an	d prescriptions should be
The CML/GIST Co-ordinator	Phone: (04) 460 4990				
PHARMAC	Facsimile: (04) 916 7571				
PO Box 10 254	Email: cmlgistcoordinator@	pharmac.govt.nz			
Wellington					
Special Authority criteria for G Funded for patients:	IST – access by application				
a) With a diagnosis (confirm (GIST).	ed by an oncologist) of unrese	ctable and/or metastati	c mal	ignant gastr	ointestinal stromal tumou
b) Maximum dose of 400 mg					
d) Initial and subsequent ap	and subsequent prescriptions c plications are valid for one yea b (prescriber determined).				quate clinical response to
APATINIB DITOSYLATE – Sp	. , ,		70	ד 🗸	ykerb
■ SA1191 Special Authority f	or Subsidy				
nitial application — (metastat		relevant specialist or m	edica	al practitione	r on the recommendatior
of a relevant specialist. Approva	Is valid for 12 months for appli	cations meeting the foll	owing	g criteria:	

Either:

(Manut	Subsidy facturer's Price)	Ful Subsidise	<i>.</i>	Brand or Generic
· · · · · · · · · · · · · · · · · · ·	\$	Per •	/	Manufacturer

continued...

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient has not previously received trastuzumab treatment for HER 2 positive metastatic breast cancer; and
 - 1.3 Lapatinib not to be given in combination with trastuzumab; and
 - 1.4 Lapatinib to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2.2 The patient started trastuzumab for metastatic breast cancer but discontinued trastuzumab within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on trastuzumab; and
 - 2.4 Lapatinib not to be given in combination with trastuzumab; and
 - 2.5 Lapatinib to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

Wastage claimable – see rule 3.3.2 on page 13

Cap 150 mg	4,680.00	120	🗸 Tasigna
Cap 200 mg	6,532.00	120	🗸 Tasigna

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Either:
 - 2.1 Patient has documented CML treatment failure* with imatinib; or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and

*Three months or six months, as applicable, dispensed all-at-once

4 Subsidised for use as monotherapy only.

PAZOPANIB - Special Authority see SA1190 on the next pa	age – Retail pharmacy		
Tab 200 mg		30	 Votrient
Tab 400 mg	2,669.40	30	 Votrient

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
 - The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70: or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.
- Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

SUNITINIB - Special Authority see SA1266 below - Retail pharmacy

Cap 12.5 mg		28	 Sutent
Cap 25 mg		28	 Sutent
Cap 50 mg	9,261.54	28	 Sutent

SA1266 Special Authority for Subsidy

Initial application - (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval: or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

2.4.2 The cancer did not progress whilst on pazopanib; and

- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology: and
 - The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal: or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application - (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Fither:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal - (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.
- Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal - (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease): or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Subsi Per	dised	Generic Manufacturer
Endocrine Therapy				
For GnRH ANALOGUES – refer to HORMONE PREPARATION	S, Trophic Hormones	, page 93		
ABIRATERONE ACETATE - Retail pharmacy-Specialist - Specialist	cial Authority see SA	1515 below		
Wastage claimable – see rule 3.3.2 on page 13	4.070.40	100	17	h
Tab 250 mg	4,276.19	120	✓ Z	Żytiga
■ SA1515 Special Authority for Subsidy				an the measurement of
Initial application only from a medical oncologist, radiation once a medical oncologist, radiation oncologist or urologist. Approval				
All of the following:		or applicatio	/13 110	curing the following chiefta.
1 Patient has prostate cancer; and				
2 Patient has metastases; and				
3 Patient's disease is castration resistant; and				
4 Either:				
4.1 All of the following:				
4.1.1 Patient is symptomatic; and				
4.1.2 Patient has disease progression (rising ser		d line anti-a	indrog	en therapy; and
4.1.3 Patient has ECOG performance score of 04.1.4 Patient has not had prior treatment with tax		or		
4.1.4 Patient has not had phot treatment with tax 4.2 All of the following:	ane chemotherapy, (JI		
4.2.1 Patient's disease has progressed following	nrior chemotherany	containina :	a tavar	ne: and
4.2.2 Patient has ECOG performance score of 0		containing a	מ נמאמו	ie, allu
4.2.3 Patient has not had prior treatment with ab	,			
Renewal — (abiraterone acetate) only from a medical oncolog		ist, urologis	t or me	edical practitioner on the
recommendation of a medical oncologist, radiation oncologist or	urologist. Approvals	valid for 5 r	nonthe	s for applications meeting
the following criteria:				
All of the following:				
1 Significant decrease in serum PSA from baseline; and				
 No evidence of clinical disease progression; and No initiation of taxane chemotherapy with abiraterone; an 	d			
4 The treatment remains appropriate and the patient is ben		t.		
BICALUTAMIDE	g			
Tab 50 mg	3.80	28	✓ F	Binarex
FLUTAMIDE – Retail pharmacy-Specialist		20		
Tab 250 mg	16.50	30	✓ F	lutamide
				Mylan S29
	55.00	100	✓ F	lutamin
MEGESTROL ACETATE – Retail pharmacy-Specialist			-	
Tab 160 mg		30	✓ A	po-Megestrol
OCTREOTIDE			-	
Inj 50 mcg per ml, 1 ml vial		5	✓ C	BL Octreotide
Inj 100 mcg per ml, 1 ml vial		5	_	BL Octreotide
Inj 500 mcg per ml, 1 ml vial	72.50	5	✓ [BL Octreotide
OCTREOTIDE LAR (SOMATOSTATIN ANALOGUE) - Special	Authority see SA101	6 on the nex	kt page	e – Retail pharmacy
Inj LAR 10 mg prefilled syringe	1,772.50	1	✓ s	Sandostatin LAR
Inj LAR 20 mg prefilled syringe		1		andostatin LAR
Inj LAR 30 mg prefilled syringe	2 451 25	1	~ ~ ~	Sandostatin LAR

	Subsidy	Fully	Brand or
(Manuf		sidised	Generic
	\$ Per	1	Manufacturer

⇒SA1016 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.
- Note: Indications marked with * are Unapproved Indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma; and
 - 2.2 Either:

2.2.1 Patient has failed surgery; or

2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or

- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has failed; or

*Three months or six months, as applicable, dispensed all-at-once

- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
TAMOXIFEN CITRATE * Tab 10 mg * Tab 20 mg		100 30 100	1	Genox Genox Genox
Aromatase Inhibitors				
ANASTROZOLE – Brand switch fee payable (Pharmacode 25409 * Tab 1 mg EXEMESTANE	, , ,	for de 30	-	Rolin
 Tab 25 mg ETROZOLE 	14.50	30	1	Pfizer Exemestane
Tab 2.5 mg	2.95	30	1	<u>Letrole</u>
Immunosuppressants				
Cytotoxic Immunosuppressants				
IZATHIOPRINE – Retail pharmacy-Specialist ₭ Tab 25 mg	9.66	100	1	Imuran
page 224	10.58	100	1	Imuran
k Inj 50 mg vial	60.00	1	1	Imuran
	25.00	50		Colloant
Tab 500 mg Cap 250 mg		50 100		Cellcept Cellcept
Powder for oral liq 1 g per 5 ml – Subsidy by endorsement		5 ml (Cellcept
Mycophenolate powder for oral liquid is subsidised only for the prescription is endorsed accordingly.				

Fusion Proteins

ETANERCEPT – Special Authority see SA1620 below – Retail pharmacy		
Inj 25 mg	96 4	 Enbrel
Inj 50 mg autoinjector1,599.9	96 4	 Enbrel
Inj 50 mg prefilled syringe1,599.9	96 4	 Enbrel

► SA1620 Special Authority for Subsidy

Initial application — (juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for juvenile idiopathic arthritis (JIA); and 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for JIA; or

2 All of the following:

2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- or intolerance; and
- 2.2 Patient diagnosed with Juvenile Idiopathic Arthritis (JIA); and
- 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
- 2.5 Both:
 - 2.5.1 Either:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or

continued...

Subsidy	Fu	lly	Brand or	_
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	1	Manufacturer	

continued...

2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. **Initial application — (ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:

 Subsidy (Manufacturer's Price)	F Subsid	ully	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
- 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
- 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

- 18-24 years Male: 7.0 cm; Female: 5.5 cm
- 25-34 years Male: 7.5 cm; Female: 5.5 cm
- 35-44 years Male: 6.5 cm; Female: 4.5 cm

45-54 years - Male: 6.0 cm; Female: 5.0 cm

55-64 years - Male: 5.5 cm; Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Three months supply may be dispensed at one time

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Patient has pyoderma gangrenosum*; and

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 4 doses.

Note: Indications marked with * are Unapproved Indications (refer to Interpretations and Definitions).

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a named specialist or rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
 - 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 3 Either:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

continued...

- 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.
- Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
 - 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
 - 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
 - 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and

‡ safety cap

^{*}Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	1	Manufacturer

- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 4 doses.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) – PCT only – Specialist Inj 50 mg per ml, 5 ml2,351.25	5	🖌 ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE – PCT only – Specialist Subsidised only for bladder cancer.		
Inj 2-8 × 100 million CFU149.37	1	 OncoTICE
Monoclonal Antibodies		
ADALIMUMAB – Special Authority see SA1621 below – Retail pharmacy		
Inj 20 mg per 0.4 ml prefilled syringe1,599.96	2	🗸 Humira
Inj 40 mg per 0.8 ml prefilled pen1,599.96	2	 HumiraPen
Inj 40 mg per 0.8 ml prefilled syringe1,599.96	2	 Humira

⇒SA1621 Special Authority for Subsidy

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis; or
- 2 All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and

2.6 Either:

- 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints;
- 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Fither:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application - (Crohn's disease) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids: and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Initial application - (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. **Initial application — (ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm

Subsidy	Full	/ Brand or
(Manufacturer's Price)	Subsidise	d Generic
\$	Per 🖌	Manufacturer

continued...

25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Initial application - (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

```
1 Both:
```

- 1.1 The patient has had an initial Special Authority approval for etanercept for psoriatic arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept: or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Fither:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; ٥r
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application - (juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for juvenile idiopathic arthritis (JIA); and 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for juvenile idiopathic arthritis; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
 - 2.2 Patient diagnosed with JIA; and
 - 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	· · ·	Manufacturer	

- 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
- 2.5 Both:
 - 2.5.1 Either:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's disease.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 4 doses.

Note: Note: Indications marked with * are Unapproved Indications (refer to (Interpretations and Definitions).

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	Manufacturer

continued...

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 CDAI score is 150 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- All of the following
 - 1 Either:
 - 1.1 Applicant is a dermatologist; or

*Three months or six months, as applicable, dispensed all-at-once

- 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or

continued...

‡ safety cap

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	1	Manufacturer

- 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.
- Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
 - 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
 - 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a named specialist or rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and

2 Either:

- 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 4 doses.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT - Special Authority see SA1726 below - Retail pharmacy

➡SA1726 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or

*Three months or six months, as applicable, dispensed all-at-once

- 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
- 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
- 1.3 There is no structural damage to the central fovea of the treated eye; and

continued...

‡ safety cap

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

197

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Any of the following:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment; or
 - 2.3 Patient has current approval to use ranibizumab for treatment of wAMD; or
 - 2.4 Patient is currently receiving treatment with aflibercept and has documented previous poor response to bevacizumab.
- Note: Criteria 2.3 and 2.4 will be removed from 1 January 2019.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Patient has centre involving diabetic macular oedema (DMO); and
 - 1.2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
 - 1.3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
 - 1.4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
 - 1.5 There is no centre-involving sub-retinal fibrosis or foveal atrophy; or
- 2 Patient is currently receiving treatment with aflibercept and has documented previous poor response to bevacizumab.

Note: Criterion 2 will be removed from 1 January 2019.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with [2nd line anti-VEGF agent], patient has retrialled with at least one injection of bevacizumab and had no response.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 below

Inj 5 mg per ml, 20 ml vial		1	 Erbitux
Inj 5 mg per ml, 100 ml vial	1,820.00	1	 Erbitux
Inj 1 mg for ECP	3.82	1 mg	 Baxter

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

	Subsidy (Manufacturer's Price)	Qub	Fully sidised	Brand or Generic
	(Manulacturer's Frice)	Per	siuiseu ✓	Manufacturer
BINUTUZUMAB – PCT only – Specialist – Special Authority	see SA1627 below			
Inj 25 mg per ml, 40 ml vial		1		azyva
Inj 1 mg for ECP	6.21	1 mg	✓ В	axter
SA1627 Special Authority for Subsidy				
nitial application — (chronic lymphocytic leukaemia) only	r from a haematologist.	Approvals	s valid fo	or 12 months for
pplications meeting the following criteria: Il of the following:				
 The patient has progressive Binet stage A, B or C CD2 	0+ chronic lymphocytic	leukaemia	requirir	ng treatment: and
2 The patient is obinutuzumab treatment naive; and	or onionio lymphocyto	iounaonnio	roquin	ig troutinoni, and
3 The patient is not eligible for full dose FCR due to com	orbidities with a score >	6 on the C	Cumulat	ive Illness Rating Scale
(CIRS) or reduced renal function (creatinine clearance				
4 Patient has adequate neutrophil and platelet counts* un	nless the cytopenias are	a conseq	uence c	of marrow infiltration by
CLL; and 5 Patient has good performance status; and				
6 Obinutuzumab to be administered at a maximum cumu	lative dose of 8.000 mg	and in co	mbinatio	on with chlorambucil for
maximum of 6 cycles.				
otes: Chronic lymphocytic leukaemia includes small lympho	cytic lymphoma. Como	rbidity refe	rs only	to illness/impairment ot
an CLL induced illness/impairment in the patient. 'Good per				
mporarily debilitated by their CLL disease symptoms a higher		ptable wh	ere trea	tment with obinutuzuma
expected to improve symptoms and improve ECOG score to Neutrophil greater than or equal to 1.5×10^{9} /L and platelets g) < 2.	75 109/		
		75 X 10/L		
MALIZUMAB – Special Authority see SA1490 below – Reta Inj 150 mg vial		1	./ v	olair
▶SA1490 Special Authority for Subsidy		I	• ^	Uldii
itial application only from a respiratory specialist. Approva	Is valid for 6 months for	applicatio	ns meet	ing the following criteria
Il of the following:		applicatio		ang the fellowing officia
1 Patient is over the age of 6; and				
2 Patient has a diagnosis of severe, life threatening asthr				
3 Past or current evidence of atopy, documented by skin				
 4 Total serum human immunoglobulin E (IgE) between 7 5 Proven compliance with optimal inhaled therapy includi 				aanida 1600 miaraaran
per day or fluticasone propionate 1000 micrograms per				
salmeterol 50 micrograms bd or eformoterol 12 microgr				
tolerated; and	,			
6 Patient has received courses of systemic corticosteroid	Is equivalent to at least	28 days tr	eatment	in the past 12 months,
unless contraindicated or not tolerated; and				
7 At least four admissions to hospital for a severe asthma those being in the previous 12 months; and	a exacerbation over the	previous 2	24 mont	ns with at least one of
8 An Asthma Control Questionnaire (ACQ-5) score of at	least 3.0 as assessed ir	the previo	ous mor	nth
enewal only from a respiratory specialist. Approvals valid fo		•		
I of the following:	, app			<u>.</u>
1 Hospital admissions have been reduced as a result of t	reatment; and			
2 A reduction in the Asthma Control Questionnaire (ACQ	,		line; an	d
3 A reduction in the maintenance oral corticosteroid dose	e of at least 50% from ba	aseline.		
ERTUZUMAB – PCT only – Specialist – Special Authority se	CA1COC on the next i			
En l'Uzuwad – Fu l'uniy – Specialisi – Special Authonity se	e SA 1606 on the next	Jage		

Inj 30 mg per ml, 14 ml vial		1	🗸 Perjeta
Inj 1 mg for ECP	9.82	1 mg	 Baxter

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB - PCT only - Specialist - Special Authority see SA1686 below

Inj 100 mg per 10 ml vial		 Mabthera
Inj 500 mg per 50 ml vial		 Mabthera
Inj 1 mg for ECP	5.64 1 mg	 Baxter

⇒SA1686 Special Authority for Subsidy

Initial application — (Post-transplant) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are Unapproved Indications.

Initial application — (Indolent, Low-grade lymphomas or hairy cell leukaemia*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

1 Both:

- The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 1.2 To be used for a maximum of 6 treatment cycles; or

2 Both:

- 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
- 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Hairy cell leukaemia includes hairy cell leukaemia variant *Unapproved indication.

Initial application - (Aggressive CD20 positive NHL) only from a relevant specialist or medical practitioner on the

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia **Initial application — (Chronic Lymphocytic Leukaemia)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 The patient is rituximab treatment naive; and
- 3 Either:
 - 3.1 The patient is chemotherapy treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
 - 3.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; and
- 4 The patient has good performance status; and
- 5 The patient does not have chromosome 17p deletion CLL; and
- 6 Rituximab to be administered in combination with fludarabine and cyclophosphamide or bendamustine for a maximum of 6 treatment cycles; and
- 7 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2. **Renewal — (Post-transplant)** only from a relevant specialist or medical practitioner on the recommendation of a relevant

specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are Unapproved Indications.

Renewal — (Indolent, Low-grade lymphomas or hairy cell leukaemia*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

*Three months or six months, as applicable, dispensed all-at-once

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Hairy cell leukaemia includes hairy cell leukaemia variant *Unapproved indication.

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	bsidised	Generic	
\$	Per	~	Manufacturer	

Renewal — (Aggressive CD20 positive NHL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (Chronic Lymphocytic Leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
- 2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
- 3 The patient does not have chromosome 17p deletion CLL; and
- 4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide or bendamustine for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

Note: Siltuximab is to be administered at doses no great	ater than 11 mg/kg every	3 weeks.	
Inj 100 mg vial	770.57	1	 Sylvant
Inj 400 mg vial		1	 Sylvant

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB – PCT only – Specialist – Special Authority see SA1632 below

Inj 150 mg vial1,	350.00 1	 Herceptin
Inj 440 mg vial3,	875.00 1	 Herceptin
Inj 1 mg for ECP	9.36 1 m	g 🖌 Baxter

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and

2 Either:

2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or 2.2 Both:

Subsidy		Fully	Brand or	
(Manufacturer's Pr	,	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
- 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
- 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and 4 Either:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or

*Three months or six months, as applicable, dispensed all-at-once

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	✓	Manufacturer

- 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

Programmed Cell Death-1 (PD-1) Inhibitors

NIVOLUMAB - PCT only - Specialist - Special Authority see SA1656 below

Inj 10 mg per ml, 4 ml vial	1,051.98	1	 Opdivo
Inj 10 mg per ml, 10 ml vial	2,629.96	1	 Opdivo
Inj 1 mg for ECP		1 mg	 Baxter

⇒SA1656 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma stage III or IV; and
- 2 Patient has measurable disease as defined by the presence of at least one CT or MRI measurable lesion; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Nivolumab is to be used at a maximum dose of 3 mg/kg every 2 weeks for a maximum of 12 weeks (6 cycles); and
- 6 Baseline measurement of overall tumour burden is documented (see Note); and
- 7 Documentation confirming that the patient has been informed and acknowledges that the initial funded treatment period of nivolumab will not be continued beyond 12 weeks (6 cycles) if their disease progresses during this time.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note; or
 - 1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.3 Patient has stable disease according to RECIST criteria (see Note); and
- 2 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; and
- 3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; and
- 5 Nivolumab will be used at a maximum dose of 3 mg/kg every 2 weeks for a maximum of 12 weeks (6 cycles).

Notes: Disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version

Subsid	dy F	ully	Brand or
(Manufacture	er's Price) Subsidis	sed	Generic
\$	Per	✓	Manufacturer

continued...

1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Target lesion measurements should be assessed using CT or MRI imaging with the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB - PCT only - Specialist - Special Authority see SA1657 below

Inj 50 mg vial	 1	🗸 Keytruda
Inj 1 mg for ECP	 1 mg	 Baxter

► SA1657 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma stage III or IV; and
- 2 Patient has measurable disease as defined by the presence of at least one CT or MRI measurable lesion; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Pembrolizumab is to be used at a maximum dose of 2 mg/kg every 3 weeks for a maximum of 12 weeks (4 cycles); and
- 6 Baseline measurement of overall tumour burden is documented (see Note); and
- 7 Documentation confirming that the patient has been informed and acknowledges that the initial funded treatment period of pembrolizumab will not be continued beyond 12 weeks (4 cycles) if their disease progresses during this time.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.3 Patient has stable disease according to RECIST criteria (see Note); and
- 2 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; and
- 3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; and

continued...

\$ safety cap

*Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

5 Pembrolizumab will be used at a maximum dose of 2 mg/kg every 3 weeks for a maximum of 12 weeks (4 cycles). Notes: Disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Target lesion measurements should be assessed using CT or MRI imaging with the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

CICLOSPORIN

Cap 25 mg Cap 50 mg	50 50	✓ Neoral✓ Neoral
Cap 100 mg Oral liq 100 mg per ml	50 50 ml OP	 ✓ Neoral ✓ Neoral
EVEROLIMUS – Special Authority see SA1491 below – Retail Wastage claimable – see rule 3.3.2 on page 13		
Tab 10 mg Tab 5 mg	30 30	 ✓ Afinitor ✓ Afinitor

SA1491 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

SIROLIMUS - S	Special Authority see SA0866 on the next page - Retail pharma	су	
Tab 1 mg		100	 Rapamune
Tab 2 mg		100	✓ Rapamune
Oral lig 1 m	g per ml	60 ml OP	Rapamune

(Mar	Subsidy nufacturer's Price)	Ful Subsidise	
	\$	Per •	Manufacturer

► SA0866 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis: or
- HUS or TTP; or
- · Leukoencepthalopathy; or
- · Significant malignant disease

TACROLIMUS - Special Authority see SA1540 below - Retail pharmacy

Cap 0.5 mg	100	Tacrolimus Sandoz
Cap 1 mg	100	✓ Tacrolimus Sandoz
Cap 5 mg – For tacrolimus oral liquid formulation refer,		
page 224 428.00	50	Tacrolimus Sandoz

■ SA1540 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Initial application - (steroid-resistant nephrotic syndrome*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient is a child with steroid-resistant nephrotic syndrome* (SRNS) where ciclosporin has been trialled in combination with prednisone and discontinued because of unacceptable side effects or inadequate clinical response; or
- 2 All of the following:
 - 2.1 The patient is an adult with SRNS: and
 - 2.2 Ciclosporin has been trialled in combination with prednisone and discontinued because of unacceptable side effects or inadequate clinical response: and
 - 2.3 Cyclophosphamide or mycophenolate have been trialled and discontinued because of unacceptable side effects or inadequate clinical response, or these treatments are contraindicated.
- Note: Indications marked with * are Unapproved Indications
- Note: Subsidy applies for either primary or rescue therapy.

	Cubaidu		Fully	Brand or
	Subsidy (Manufacturer's Price)	Sub	Fully sidised	Generic
	\$	Per	✓	Manufacturer
Antiallergy Preparations				
Allergic Emergencies				
Allergic Elliergencies				
ICATIBANT – Special Authority see SA1558 below – Retail phan				
Inj 10 mg per ml, 3 ml prefilled syringe	2,668.00	1		irazyr
SA1558 Special Authority for Subsidy Initial application only from a clinical immunologist or relevant s the following criteria: Both:	pecialist. Approvals	valid for 1	2 month	s for applications meeting
 Supply for anticipated emergency treatment of laryngeal/ angioedema (HAE) for patients with confirmed diagnosis 	1 7 0			,
2 The patient has undergone product training and has agree	ed upon an action pl	an for self-	adminis	tration.
Renewal from any relevant practitioner. Approvals valid for 12 n is benefiting from treatment.	nonths where the trea	atment ren	nains ap	propriate and the patient
Allergy Desensitisation				
 SA1367 Special Authority for Subsidy Initial application only from a relevant specialist. Approvals val Both: RAST or skin test positive; and 	id for 2 years for app	lications m	neeting t	the following criteria:
2 Patient has had severe generalised reaction to the sensiti	sing agent.			
Renewal only from a relevant specialist. Approvals valid for 2 ye benefiting from treatment.	0 0	nent remai	ns appro	opriate and the patient is
BEE VENOM ALLERGY TREATMENT - Special Authority see	SA1367 above – Ret	ail pharma	CV	
Maintenance kit - 6 vials 120 mcg freeze dried venom, with diluent		' 1 OP	-	enomil \$29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent		1 OP		lbey
9 ml, 3 diluent 1.8 ml				libey
WASP VENOM ALLERGY TREATMENT – Special Authority see Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze	e SA 1367 above – H	etali pharn	nacy	
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml		1 OP	🗸 A	lbey
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze				•
dried venom, with diluent		1 OP	✓ V	enomil \$29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	. .	lbey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze		101	• •	libey
dried venom, with diluent		1 OP	🗸 V	enomil S29
Antihistamines				
CETIRIZINE HYDROCHLORIDE				
* Tab 10 mg		100	✓ Z	
*‡ Oral liq 1 mg per ml	2.99	200 ml	✓ H	listaclear
CHLORPHENIRAMINE MALEATE *+ Oral lig 2 mg per 5 ml	8 06	500 ml	/ u	listafen
*+ Oran ind 2 mg her 0 min	0.00	500 m	• 1	ווסנמוכוו

	Subsidy (Manufacturer's	Price) Subs	Fully Brand or idised Generic
	\$	Per	 Manufacturer
DEXTROCHLORPHENIRAMINE MALEATE			
* Tab 2 mg	2.02	40	
-	(8.40)		Polaramine
	1.01	20	
	(5.99)		Polaramine
*‡ Oral liq 2 mg per 5 ml	1.77	100 ml	
	(10.29)		Polaramine
FEXOFENADINE HYDROCHLORIDE			
* Tab 60 mg	4.34	20	
· · · · · · · · · · · · · · · · · · ·	(8.23)		Telfast
* Tab 120 mg		10	
· · · · · · · · · · · · · · · · · · ·	(8.23)		Telfast
	14.22	30	
	(26.44)		Telfast
LORATADINE	()		
· · · · · · · · · · · · · · · · · · ·	1 00	100	 Lorafix
5		120 ml	✓ Loranx ✓ Lorfast
· · · · · · · · · · · · · · · · · ·	2.10	120111	- LUHASI
PROMETHAZINE HYDROCHLORIDE			• • • • •
* Tab 10 mg		50	✓ <u>Allersoothe</u>
* Tab 25 mg		50	✓ <u>Allersoothe</u>
* + Oral liq 1 mg per 1 ml		100 ml	✓ <u>Allersoothe</u>
Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a	PSO 15.54	5	 Hospira
TRIMEPRAZINE TARTRATE			
Oral liq 30 mg per 5 ml	2.79	100 ml OP	
	(8.06)		Vallergan Forte
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler, 50 mcg per dose	0.20	200 dose OP	✓ Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP	✓ Gval ✓ Beclazone 50
Aerosol inhaler, 100 mcg per dose ch C-nee		200 dose OP	✓ Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP	✓ Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free		200 dose OP	 ✓ Beclazone 250
		200 0036 01	
BUDESONIDE			6 - • • •
Powder for inhalation, 100 mcg per dose	17.00	200 dose OP	✓ Pulmicort
			Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00	200 dose OP	 Pulmicort
			Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00	200 dose OP	 Pulmicort
			Turbuhaler
FLUTICASONE			
Aerosol inhaler, 50 mcg per dose	4.68	120 dose OP	✓ Floair
Aerosol inhaler, 50 mcg per dose CFC-free		120 dose OP	 Flixotide
Powder for inhalation, 50 mcg per dose	7.50	60 dose OP	 Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP	 Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose	7.22	120 dose OP	✓ Floair
Aerosol inhaler, 125 mcg per dose CFC-free	13.60	120 dose OP	 Flixotide
Aerosol inhaler, 250 mcg per dose		120 dose OP	✓ Floair
Aerosol inhaler, 250 mcg per dose CFC-free		120 dose OP	 Flixotide
Powder for inhalation, 250 mcg per dose		60 dose OP	 Flixotide Accuhaler

‡ safety cap

▲ Three months supply may be dispensed at one time

*Three months or six months, as applicable, dispensed all-at-once if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully Brand or
	(Manufacturer's \$	Price) Subsi Per	dised Generic Manufacturer
beled to see the Data set of the America			
nhaled Long-acting Beta-adrenoceptor Agonis	its		
FORMOTEROL FUMARATE			
Powder for inhalation, 6 mcg per dose, breath activated		60 dose OP	Oxis Turbuhaler
Powder for inhalation, 12 mcg per dose, and monodose dev	(16.90) ice 20.64	60 dose	
	(35.80)		Foradil
IDACATEROL	. ,		
Powder for inhalation 150 mcg	61.00	30 dose OP	 Onbrez Breezhaler
Powder for inhalation 300 mcg	61.00	30 dose OP	 Onbrez Breezhaler
ALMETEROL			_
Aerosol inhaler CFC-free, 25 mcg per dose		120 dose OP	 Serevent
Aerosol inhaler 25 mcg per dose Powder for inhalation, 50 mcg per dose, breath activated		120 dose OP 60 dose OP	 Meterol Serevent Accuhaler
Fowder for initialation, 50 mcg per dose, breath activated	25.00	60 dose OF	 Serevent Accunater
Inhaled Corticosteroids with Long-Acting Beta-	Adrenocept	tor Agonists	
	•	Ū	
UDESONIDE WITH EFORMOTEROL Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg	19.00	120 dose OP	✓ Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6 mcg		120 dose OP	✓ Symbicort
	109		Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg	21.40	120 dose OP	 Vannair
Powder for inhalation 200 mcg with eformoterol fumarate 6 r	ncg44.08	120 dose OP	 Symbicort
			Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate			
12 mcg – No more than 2 dose per day		60 dose OP	 Symbicort Turbuhaler 400/12
LUTICASONE FUROATE WITH VILANTEROL			
Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose OP	✓ Breo Ellipta
LUTICASONE WITH SALMETEROL		00 0030 01	
Aerosol inhaler 50 mcg with salmeterol 25 mcg	14 58	120 dose OP	✓ RexAir
	33.74		✓ Seretide
Aerosol inhaler 125 mcg with salmeterol 25 mcg	16.83	120 dose OP	 RexAir
	44.08		 Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg - No			4 a b b b b b b b b b b
more than 2 dose per day		60 dose OP	 Seretide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg – No		60 dose OP	 Seretide Accuhaler
more than 2 dose per day		60 dose OP	 Serelide Accunaler
Beta-Adrenoceptor Agonists			
ALBUTAMOL Oral liq 400 mcg per ml	11.00	150 ml	✓ Ventolin
Infusion 1 mg per ml, 5 ml		10	
	(130.21)		Ventolin
Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO		5	 Ventolin

	Subsidy (Manufacturer's		
	\$	Per	 Manufacturer
Inhaled Beta-Adrenoceptor Agonists			
SALBUTAMOL			
Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000 dose available on a PSO		200 dose OP	 Respigen Respigen
	(6.00)		 SalAir Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO		20	✓ <u>Asthalin</u>
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO		20	✓ <u>Asthalin</u>
TERBUTALINE SULPHATE Powder for inhalation, 250 mcg per dose, breath activated		200 dose OP	 Bricanyl Turbuhaler
Anticholinergic Agents			
IPRATROPIUM BROMIDE			
Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dos available on a PSO	16.20	200 dose OP	✓ Atrovent
Nebuliser soln, 250 mcg per ml, 1 ml ampoule – Up to 40 ne available on a PSO		20	✓ Univent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 ne available on a PSO	eb	20	✓ <u>Univent</u>
Inhaled Beta-Adrenoceptor Agonists with Antic	holinergic A	Agents	
SALBUTAMOL WITH IPRATROPIUM BROMIDE	•	5	
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p dose CFC-free		200 dose OP	✓ Duolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule – Up to 20 neb available on a PSO)3.59	20	✓ <u>Duolin</u>
Long-Acting Muscarinic Antagonists			
 GLYCOPYRRONIUM – Subsidy by endorsement a) Inhaled glycopyrronium treatment will not be subsidised i umeclidinium. 	f patient is also	receiving treatme	ent with subsidised tiotropium or
 b) Glycopyrronium powder for inhalation 50 mcg per dose is having COPD using spirometry, and the prescription is er Powder for inhalation 50 mcg per dose 	ndorsed accordi		o have been diagnosed as
TIOTROPIUM BROMIDE – Special Authority see SA1568 below Tiotropium treatment will not be subsidised if patient is also r	– Retail pharm	acy	
umeclidinium. Powder for inhalation, 18 mcg per dose Soln for inhalation 2.5 mcg per dose		30 dose 60 dose OP	✓ Spiriva✓ Spiriva Respimat
SA1568 Special Authority for Subsidy Initial application only from a general practitioner or relevant sp following criteria:	ecialist. Approv	vals valid for 2 ye	ars for applications meeting the

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

All of the following:

- 1 To be used for the long-term maintenance treatment of bronchospasm and dyspnoea associated with COPD; and
- 2 In addition to standard treatment, the patient has trialled a short acting bronchodilator dose of at least 40 µg ipratropium a.i.d for one month; and
- 3 Either:
 - The patient's breathlessness according to the Medical Research Council (UK) dyspnoea scale is:
 - 3.1 Grade 3 (stops for breath after walking about 100 meters or after a few minutes on the level); or
 - 3.2 Grade 4 (too breathless to leave the house, or breathless when dressing or undressing); and
- 4 All of the following:
 - Applicant must state recent measurement of:
 - 4.1 Actual FEV₁ (litres); and
 - 4.2 Predicted FEV₁ (litres); and
 - 4.3 Actual FEV_1 as a % of predicted (must be below 60%); and
- 5 Either:
 - 5.1 Patient is not a smoker (for reporting purposes only); or
 - 5.2 Patient is a smoker and has been offered smoking cessation counselling; and
- 6 The patient has been offered annual influenza immunisation.

Renewal only from a general practitioner or relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).
- GLYCOPYRRONIUM WITH INDACATEROL Special Authority see SA1584 above Retail pharmacy Powder for Inhalation 50 mcg with indacaterol 110 mcg......81.00 30 dose OP ✓ Ultibro Breezhaler

(1	Subsidy Manufacturer's P \$		Fully Brand or dised Generic ✓ Manufacturer
UMECLIDINIUM WITH VILANTEROL – Special Authority see SA1 Powder for inhalation 62.5 mcg with vilanterol 25 mcg		vious page – Re 30 dose OP	etail pharmacy ✓ Anoro Ellipta
Antifibrotics			
PIRFENIDONE - Retail pharmacy-Specialist - Special Authority se	ee SA1628 bel	wo	
Cap 267 mg – Wastage claimable – see rule 3.3.2 on page 13	3.645.00	270	✓ Esbriet
➡SA1628 Special Authority for Subsidy			
Initial application — (idiopathic pulmonary fibrosis) only from a applications meeting the following criteria: All of the following:	a respiratory sp	ecialist. Approv	vals valid for 12 months for
 Patient has been diagnosed with idiopathic pulmonary fibros Forced vital capacity is between 50% and 80% predicted; ar Pirfenidone is to be discontinued at disease progression (Se 	nd	d by histology, (CT or biopsy; and
Renewal — (idiopathic pulmonary fibrosis) only from a respirate meeting the following criteria: Both:		Approvals valid	for 12 months for applications
Treatment remains clinically appropriate and patient is bene Pirfenidone is to be discontinued at disease progression (Se		I tolerating treat	ment; and
Note: disease progression is defined as a decline in percent predic	,	% or more within	n any 12 month period.
Leukotriene Receptor Antagonists			
MONTELUKAST			
Prescribing Guideline: Clinical evidence indicates that the effe used in short treatment courses.	ctiveness of m	ontelukast is str	ongest when montelukast is
* Tab 4 mg * Tab 5 mg		28 28	 ✓ <u>Apo-Montelukast</u> ✓ Apo-Montelukast
* Tab 10 mg		28	✓ <u>Apo-Montelukast</u>
Mast Cell Stabilisers			
NEDOCROMIL			
Aerosol inhaler, 2 mg per dose CFC-free SODIUM CROMOGLICATE	28.07	112 dose OP	 Tilade
Aerosol inhaler, 5 mg per dose CFC-free		112 dose OP	✓ Intal Forte CFC Free
Methylxanthines			
AMINOPHYLLINE			
Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a PSO	124.37	5	✓ DBL Aminophylline
THEOPHYLLINE	04 54	100	
* Tab long-acting 250 mg *‡ Oral liq 80 mg per 15 ml		100 500 ml	✓ Nuelin-SR✓ Nuelin
Mucolytics			
DORNASE ALFA – Special Authority see SA0611 on the next page Nebuliser soln, 2.5 mg per 2.5 ml ampoule		macy 6	✓ Pulmozyme

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

RESPIRATORY SYSTEM AND ALLERGIES

<u> </u>	Subsidy (Manufacturer's \$	Price) Subs Per	Fully Brand or idised Generic ✓ Manufacturer
► SA0611 Special Authority for Subsidy Special Authority approved by the Cystic Fibrosis Adv Notes: Application details may be obtained from PHA	risory Panel ARMAC's website <u>http://ww</u>	w.pharmac.govt	. <u>nz</u> or:
The Co-ordinator, Cystic Fibrosis Advisory Panel PHARMAC, PO Box 10 254 Wellington	Phone: (04) 460 4990 Facsimile: (04) 916 757 Email: <u>CFPanel@pharm</u>		
Prescriptions for patients approved for treatment mus and expertise in treating cystic fibrosis. SODIUM CHLORIDE	t be written by respiratory p	physicians or pae	ediatricians who have experience
Not funded for use as a nasal drop. Soln 7%	23.50	90 ml OP	✓ Biomed
Nasal Preparations			
Allergy Prophylactics			
BECLOMETHASONE DIPROPIONATE Metered aqueous nasal spray, 50 mcg per dose .	2.35 (5.26)	200 dose OP	Alanase
Metered aqueous nasal spray, 100 mcg per dose		200 dose OP	Alanase
BUDESONIDE Metered aqueous nasal spray, 50 mcg per dose .	, , , , , , , , , , , , , , , , , , ,	200 dose OP	Butacort Aqueous
Metered aqueous nasal spray, 100 mcg per dose	2.61 (6.00)	200 dose OP	Butacort Aqueous
FLUTICASONE PROPIONATE Metered aqueous nasal spray, 50 mcg per dose .	2.18	120 dose OP	 Flixonase Hayfever & Allergy
IPRATROPIUM BROMIDE Aqueous nasal spray, 0.03%	4.61	15 ml OP	✓ <u>Univent</u>
Respiratory Devices			
MASK FOR SPACER DEVICE a) Up to 50 dev available on a PSO b) Only on a PSO c) Only for children aged six years and under Small.		1	✔ e-chamber Mask
PEAK FLOW METER a) Up to 25 dev available on a PSO b) Only on a PSO			
Low range	9.54	1	 Mini-Wright AFS Low Range
Normal range	9.54	1	✓ <u>Mini-Wright</u> Standard

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully osidised	Brand or Generic Manufacturer
SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
220 ml (single patient)	2.95	1	✓ <u>e</u>	-chamber Turbo
510 ml (single patient)	5.12	1	✔ e	-chamber La Grande
800 ml	6.50	1	🗸 V	olumatic
Respiratory Stimulants				
CAFFEINE CITRATE Oral liq 20 mg per ml (10 mg base per ml)		5 ml OP	✔ В	liomed

	Subsidy		Fully Brand or
	(Manufacturer's F \$	Price) Sub: Per	sidised Generic Manufacturer
Ear Preparations			
CETIC ACID WITH 1, 2- PROPANEDIOL DIACETATE AND BE	ENZETHONIUM		
For Vosol ear drops with hydrocortisone powder refer Standa	ard Formulae, <mark>pa</mark>	age 227	
Ear drops 2% with 1, 2-Propanediol diacetate 3% and	0.07		(Maral
benzethonium chloride 0.02%	6.97	35 ml OP	 Vosol
LUMETASONE PIVALATE Ear drops 0.02% with clioguinol 1%	4.46	7.5 ml OP	 Locacorten-Viaform
	4.40	7.5 111 01	ED's
			✓ Locorten-Vioform
RIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTA	ΓIN	
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate			
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	 Kenacomb
Ear/Eye Preparations			
EXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and	4.50		
gramicidin 50 mcg per ml	4.50 (9.27)	8 ml OP	Sofradex
RAMYCETIN SULPHATE	(0.27)		Conducx
Ear/Eye drops 0.5%	4.13	8 ml OP	
	(8.65)		Soframycin
Eye Preparations			
ye preparations are only funded for use in the eye, unless expli-	citly stated other	WISE.	
Anti-Infective Preparations			
CICLOVIR			
Eye oint 3%	14.92	4.5 g OP	ViruPOS
HLORAMPHENICOL Eye oint 1%	0.40		Chloroig
Eye drops 0.5%		4 g OP 10 ml OP	 ✓ <u>Chlorsig</u> ✓ Chlorafast
Funded for use in the ear*.			<u>emeralaci</u>
Indications marked with * are Unapproved Indications.			
IPROFLOXACIN			
Eye drops 0.3% – Subsidy by endorsement		5 ml OP	 Ciprofloxacin Teva
a) When prescribed for the treatment of bacterial kerati	(12.43) tis or severe bac	torial conjuncti	Ciloxan
or for the second line treatment of chronic suppurativ			
accordingly.		, ,	
b) Ciprofloxacin Teva to be Sole Supply on 1 September			
Note: Indication marked with a * is an Unapproved Indic	cation.		
Ciloxan Eye drops 0.3% to be delisted 1 September 2018)			
ENTAMICIN SULPHATE Eye drops 0.3%	11 40	5 ml OP	✓ Genoptic
Eye diops 0.3%	11.40	JIIIUF	
€ Eye drops 0.1%	2.97	10 ml OP	
	(14.55)		Brolene
16 fully subsidised		proved medicine	supplied under Section 29

S29 Unapproved medicine supplied under Section 29 Sole Subsidised Supply

SENSORY ORGANS

	Subsidy (Manufacturer's Pi \$	rice) Subs Per	Fully Bran idised Gene ✓ Man	
SODIUM FUSIDATE [FUSIDIC ACID] Eye drops 1% TOBRAMYCIN	4.50	5 g OP	 Fucitha 	Ilmic
Eye oint 0.3% Eye drops 0.3% Corticosteroids and Other Anti-Inflammatory P	11.48	3.5 g OP 5 ml OP	✓ Tobrex✓ Tobrex	
DEXAMETHASONE	. opulationo			
* Eye oint 0.1%		3.5 g OP	 Maxide 	x
* Eye drops 0.1% Ocular implant 700 mcg – Special Authority see SA1680 be		5 ml OP	 Maxide 	x
– Retail pharmacy	1,444.50	1	 Ozurde 	x
SA1680 Special Authority for Subsidy Initial application — (Diabetic macular oedema) only from ar	n ophthalmologist.	Approvals va	lid for 12 mon	ths for applications

meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Either:
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

*Three months or six months, as applicable, dispensed all-at-once

*	Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b			
	sulphate 6,000 u per g5.	39	3.5 g OP	 Maxitrol
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin		-	
	b sulphate 6,000 u per ml4.	50	5 ml OP	 Maxitrol

if endorsed "certified exemption" by the prescriber or pharmacist.

SENSORY ORGANS

	Subsidy	aa) Cubai	Fully Brand or
	(Manufacturer's Prio \$	ce) Subsi Per	dised Generic Manufacturer
DICLOFENAC SODIUM			
* Eye drops 0.1%		5 ml OP	 Voltaren Ophtha
FLUOROMETHOLONE			
* Eye drops 0.1%		5 ml OP	✓ <u>FML</u>
LEVOCABASTINE			
Eye drops 0.5 mg per ml	8.71	4 ml OP	
	(10.34)		Livostin
LODOXAMIDE			
Eye drops 0.1%	8.71	10 ml OP	 Lomide
PREDNISOLONE ACETATE			
Eye drops 1%	3.93	10 ml OP	 Prednisolone-AFT
	7.00	5 ml OP	 Pred Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority se		- Retail pharm	nacy
Eye drops 0.5%, single dose (preservative free)		20 dose	 Minims Prednisolone
► SA1715 Special Authority for Subsidy			
Initial application only from an ophthalmologist or optometrist. following criteria:	Approvals valid for	f 6 months for	applications meeting the
Both:			
 Patient has severe inflammation; and Patient has a confirmed allergic reaction to preservative in 	n eye drops.		
Renewal from any relevant practitioner. Approvals valid for 6 me benefiting from treatment.	onths where the tre	eatment remai	ns appropriate and the patient is
SODIUM CROMOGLICATE			
Eye drops 2%	0.85	5 ml OP	✓ <u>Rexacrom</u>
Glaucoma Preparations - Beta Blockers			
BETAXOLOL			
* Eye drops 0.25%	11.80	5 ml OP	 Betoptic S
* Eye drops 0.5%	7.50	5 ml OP	 Betoptic

LE	/OBUNOLOL		
*	Eye drops 0.5%7.00	5 ml OP	 Betagan
TIN	IOLOL		
*	Eye drops 0.25%	5 ml OP	Arrow-Timolol
*	Eye drops 0.25%, gel forming	2.5 ml OP	 Timoptol XE
*	Eye drops 0.5%	5 ml OP	Arrow-Timolol
*	Eye drops 0.5%, gel forming	2.5 ml OP	Timoptol XE

Glaucoma Preparations - Carbonic Anhydrase Inhibitors

ACETAZOLAMIDE		
* Tab 250 mg – For acetazolamide oral liquid formulation refer,		
page 22417.03	100	 Diamox
BRINZOLAMIDE		
* Eye drops 1%	5 ml OP	 Azopt
DORZOLAMIDE HYDROCHLORIDE		
* Eye drops 2%	5 ml OP	
(17.44)		Trusopt

SENSORY ORGANS

	Subsidy (Manufacturer's Pr \$	ice) Subs Per	Fully Brand or idised Generic ✓ Manufacturer
DORZOLAMIDE WITH TIMOLOL * Eye drops 2% with timolol 0.5%	3.45	5 ml OP	✓ <u>Arrow-Dortim</u>
Glaucoma Preparations - Prostaglandin Analog	ues		
BIMATOPROST * Eye drops 0.03%	3.65	3 ml OP	✓ Bimatoprost Actavis
* Eye drops 0.005%	1.50	2.5 ml OP	✓ <u>Hysite</u>
* Eye drops 0.004%	7.30 19.50	5 ml OP 2.5 ml OP	 ✓ <u>Travopt</u> ✓ Travatan
Glaucoma Preparations - Other			
BRIMONIDINE TARTRATE * Eye drops 0.2% BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE	4.29	5 ml OP	✓ <u>Arrow-Brimonidine</u>
* Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	 Combigan
 ¥ Eye drops 1% * Eye drops 2% * Eye drops 4% Subsidised for oral use pursuant to the Standard Formula 	5.35 7.99	15 ml OP 15 ml OP 15 ml OP	 ✓ Isopto Carpine ✓ Isopto Carpine ✓ Isopto Carpine
 ★ Eye drops 2% single dose – Special Authority see SA0895 below – Retail pharmacy		20 dose	 Minims Pilocarpine

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

1 Patient has to use an unpreserved solution due to an allergy to the preservative; or

2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics

ATROPINE SULPHATE * Eye drops 1%	15 ml OP	✓ Atropt
CYCLOPENTOLATE HYDROCHLORIDE		
* Eye drops 1%	15 ml OP	 Cyclogyl
* Eye drops 0.5%	15 ml OP	 ✓ Mydriacyl ✓ Mydriacyl
* Eye drops 1%	15 ml OP	 Mydriacyl

Preparations for Tear Deficiency

For acetylcysteine eye drops refer Standard Formulae, page 227			
HYPROMELLOSE			
* Eye drops 0.5%	2.00	15 ml OP	
	(3.92)		Methopt

‡ safety cap

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Pr			Brand or Generic
	\$	Per	-	Manufacturer
HYPROMELLOSE WITH DEXTRAN				
* Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	🗸 Po	oly-Tears
POLYVINYL ALCOHOL				
* Eye drops 1.4%	2.62	15 ml OP	🗸 <u>Vi</u>	stil
* Eye drops 3%	3.68	15 ml OP	✓ <u>Vi</u>	stil Forte

Preservative Free Ocular Lubricants

⇒SA1388 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

1 Confirmed diagnosis by slit lamp of severe secretory dry eye; and

- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

diops and has benefited from treatment.		
CARBOMER – Special Authority see SA1388 above – Retail pharmacy Ophthalmic gel 0.3%, 0.5 g	5 30	✓ Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL – Special Authority see SA1 Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml		narmacy Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] – Special Authority see SA Eye drops 1 mg per ml	0 10 ml OP cedures Manual restri	✓ Hylo-Fresh ction allowing one bottle per
Other Eye Preparations		
NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%	5 15 ml OP	✓ Naphcon Forte
OLOPATADINE Eye drops 0.1%	0 5 ml OP	✓ Patanol
PARAFFIN LIQUID WITH SOFT WHITE PARAFFIN * Eye oint with soft white paraffin	3 3.5 g OP	✓ Refresh Night Time
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%	3 3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE Eye oint 138 mcg per g	0 5 g OP	✔ VitA-POS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Various				
PHARMACY SERVICES				
May only be claimed once per patient. * Brand switch fee	4.50	1 fee	✓ B ✓ B	SF CareSens Dual SF CareSens N SF CareSens N POP SF CareSens N Premier
 a) The Pharmacode for BSF CareSens N is 2423138 - so b) The Pharmacode for BSF CareSens N POP is 242315 c) The Pharmacode for BSF CareSens N Premier is 253 d) The Pharmacode for BSF CareSens Dual is 2535890 e) The Pharmacode for BSF Rolin is 2540959 - see also (BSF CareSens Dual Brand switch fee to be delisted 1 August 2013) (BSF CareSens N POP Brand switch fee to be delisted 1 August 2018) (BSF CareSens N Premier Brand switch fee to be delisted 1 August 2018) (BSF CareSens N Premier Brand switch fee to be delisted 1 August 2018) 	54 - see also page 2 5882 - see also page - see also page 26 page 184 (8) 2018)		✔ B	SF Rolin
Agents Used in the Treatment of Poisonings				
Antidotes				
ACETYLCYSTEINE – Retail pharmacy-Specialist Inj 200 mg per ml, 10 ml ampoule NALOXONE HYDROCHLORIDE	78.34	10	✓ □	BL Acetylcysteine
 a) Up to 5 inj available on a PSO b) Only on a PSO * Inj 400 mcg per ml, 1 ml ampoule 		5	✓ D	BL Naloxone
				Hydrochloride
DBL Naloxone Hydrochloride to be Sole Supply on 1 Sep	tember 2018			
Removal and Elimination				
CHARCOAL * Oral liq 50 g per 250 ml a) Up to 250 ml available on a PSO b) Only on a PSO		0 ml (OP 🗸 C	arbosorb-X
DEFERASIROX – Special Authority see SA1492 on the next page Wastage claimable – see rule 3.3.2 on page 13				
Tab 125 mg dispersible Tab 250 mg dispersible Tab 500 mg dispersible	552.00	28 28 28	✓ E	xjade xjade xjade

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

VARIOUS

Subsidy (Manufacturer's Price)	Fu Subsidis		
\$	Per	 Manufacturer 	

SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea: or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per µL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 - 1.0 cells per µL).

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE - Special Authority see SA1480 below - Retail pharmacy

Tab 500 mg	 100	 Ferriprox
Oral liq 100 mg per 1 ml	 250 ml OP	 Ferriprox

SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESEEBBIOXAMINE MESILATE

* Inj 500 mg vial	51.52	10	 Desferal
SODIUM CALCIUM EDETATE			
* Inj 200 mg per ml, 5 ml	53.31	6	
	(156.71)		Calcium Disodium

m Versenate

INTRODUCTION

The following extemporaneously compounded products are eligible for subsidy:

- The "Standard Formulae".
- Oral liquid mixtures for patients unable to swallow subsidised solid dose oral formulations.
- The preparation of syringe drivers when prescribed by a general practitioner.
- Dermatological preparations
 - a) One or more subsidised dermatological galenical(s) in a subsidised dermatological base.
 - b) Dilution of proprietary Topical Corticosteroid-Plain preparations with a dermatological base (Retail pharmacy-Specialist).

Glossary

Dermatological base: The products listed in the Barrier creams and Emollients section and the Topical Corticosteroids-Plain section of the Pharmaceutical Schedule are classified as dermatological bases for the purposes of extemporaneous compounding and are the bases to which the dermatological galenicals can be added. Also the dermatological bases in the Barrier Creams and Emollients section of the Pharmaceutical Schedule can be used for diluting proprietary Topical Corticosteroid-Plain preparations. The following products are dermatological bases:

- Aqueous cream
- Cetomacrogol cream BP
- Collodion flexible
- · Emulsifying ointment BP
- · Hydrocortisone with wool fat and mineral oil lotion
- Oil in water emulsion
- Urea cream 10%
- White soft paraffin
- Wool fat with mineral oil lotion
- · Zinc and castor oil ointment BP
- Proprietary Topical Corticosteroid-Plain preparations

Dermatological galenical: Dermatological galenicals will only be subsidised when added to a dermatological base. More than one dermatological galenical can be added to a dermatological base.

The following are dermatological galenicals:

- Coal tar solution up to 10%
- Hydrocortisone powder up to 5%
- · Menthol crystals
- Salicylic acid powder
- Sulphur precipitated powder

Standard formulae: Standard formulae are a list of fomulae for ECPs that are subsidised. Their ingredients are listed under the appropriate therapeutic heading in Section B of the Pharmaceutical Schedule and also in Section C.

Explanatory notes

Oral liquid mixtures

Oral liquid mixtures are subsidised for patients unable to swallow subsidised solid oral dose forms where no suitable alternative proprietary formulation is subsidised. Suitable alternatives include dispersible and sublingual formulations, oral liquid formulations or rectal formulations. Before extemporaneously compounding an oral liquid mixture, other alternatives such as dispersing the solid dose form (if appropriate) or crushing the solid dose form in jam, honey or soft foods such as yoghurt should be explored. The Emixt website www.pharminfotech.co.nz has evidence-based formulations which are intended to standardise compounded oral liquids within New Zealand.

Pharmaceuticals with standardised formula for compounding in Ora products

- Acetazolamide 25 mg/ml Allopurinol 20 mg/ml Amlodipine 1 mg/ml Azathioprine 50 mg/ml Baclofen 10 mg/ml Carvedilol 1 mg/ml Clopidogrel 5 mg/ml Diltiazem hydrochloride 12 mg/ml Dipyridamole 10 mg/ml Domperidone 1 mg/ml Enalapril 1 mg/ml
- Flecainide 20 mg/ml Gabapentin 100 mg/ml Hydrocortisone 1 mg/ml Labetolol 10 mg/ml Levodopa with carbidopa (5 mg levodopa + 1.25 mg carbidopa)/ml Metoclopramide 1 mg/ml Metoprolol tartrate 10 mg/ml Nitrofurantoin 10 mg/ml Pyrazinamide 100 mg/ml Rifabutin 20 mg/ml
- Sildenafil 2 mg/ml Sotalol 5 mg/ml Sulfasalazine 100 mg/ml Tacrolimus 1 mg/ml Terbinafine 25 mg/ml Tramadol 10 mg/ml Ursodeoxycholic acid 50 mg/ml Valganciclovir 60 mg/ml* Verapamil hydrochloride 50 mg/ml

qs

*Note this is a DCS formulation

PHARMAC endorses the recommendations of the Emixt website and encourages New Zealand pharmacists to use these formulations when compounding is appropriate. The Emixt website also provides stability and expiry data for compounded products. For the majority of products compounded with Ora-Blend, Ora-Blend SF, Ora-Plus, Ora-Sweet or Ora-Sweet SF a four week expiry is appropriate.

Please note that no oral liquid mixture will be eligible for Subsidy unless all the requirements of Section B and C of the Schedule applicable to that pharmaceutical are met.

Some community pharmacies may not have appropriate equipment to compound all of the listed products, please use appropriate clinical iudgement.

Subsidy for extemporaneously compounded oral liquid mixtures is based on:

Solid dose form	qs
Preservative	qs
Suspending agent	qs
Water	to 100%

or

Solid dose form Ora-Blend, Ora-Blend SF, Ora-Plus, Ora-Sweet and/or Ora-Sweet SF to 100%

Prescribers may prescribe or pharmacists may add extra non-subsidised ingredients such as flavouring and colouring agents, but these extra ingredients will not be reimbursed. The subsidised ingredients in the formula will be reimbursed and a compounding fee paid.

The majority of extemporaneously compounded oral liquid mixtures should contain a preservative and suspending agent.

- Ora-Blend, Ora-Blend SF, Ora-Plus, Ora-Sweet and Ora-Sweet SF when used correctly are an appropriate preservative and suspending agent.
- Methylcellulose 3% is considered a suitable suspending agent and compound hydroxybenzoate solution or methyl hydroxybenzoate 10% solution are considered to be suitable preservatives. Usually 1 ml of these preservative solutions is added to 100 ml of oral liquid mixture.

Some solid oral dose forms are not appropriate for compounding into oral liquid mixtures and should therefore not be used/considered for extemporaneously compounded oral liquid mixtures. This includes long-acting solid dose formulations, enteric coated tablets or capsules, sugar coated tablets, hard gelatin capsules and chemotherapeutic agents.

The following practices will not be subsidised:

- Where a Standard Formula exists in the Pharmaceutical Schedule for a solid dose form, compounding the solid dose form in Ora-Blend, Ora-Blend SF, Ora-Plus, Ora-Sweet and/or Ora-Sweet SF.
- Mixing one or more proprietary oral liquids (eg an antihistamine with pholcodine linctus).
- Extemporaneously compounding an oral liquid with more than one solid dose chemical.
- Mixing more than one extemporaneously compounded oral liquid mixture.
- · Mixing one or more extemporaneously compounded oral liquid mixtures with one or more proprietary oral liquids.
- The addition of a chemical/powder/agent/solution to a proprietary oral liquid or extemporaneously compounded oral mixture.


Standard formulae

A list of standard formulae is contained in this section. All ingredients associated with a standard formula will be subsidised and an appropriate compounding fee paid.

Prescribers may prescribe or pharmacists may add extra non-subsidised ingredients, but these extra ingredients will not be reimbursed. The subsidised ingredients in the formula will be reimbursed and a compounding fee paid.

Dermatological Preparations

Proprietary topical corticosteroid preparations may be diluted with a dermatological base (see page 223) from the Barrier Creams and Emollients section of the Pharmaceutical Schedule (Retail pharmacy-Specialist). Dilution of proprietary topical corticosteroid preparations should only be prescribed for withdrawing patients off higher strength proprietary topical corticosteroid products where there is no suitable proprietary product of a lower strength available or an extemporaneously compounded product with up to 5% hydrocortisone is not appropriate. (In general proprietary topical corticosteroid preparations should not be diluted because dilution effects can be unpredictable and may not be linear, and usually there is no stability data available for diluted products). One or more dermatological galenicals may be added to a dermatological base (including proprietary, but these extra ingredients will not be reimbursed. The subsidised ingredients in the formula will be reimbursed and a compounding fee paid. The addition of dermatological galenicals to diluted proprietary Topical Corticosteroids-Plain will not be subsidised. The flow diagram on the next page may assist you in deciding whether or not a dermatological ECP is subsidised.



Standard Formulae

ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base	qs qs	PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water	1 g 70 ml to 100 ml
ASPIRIN AND CHLOROFORM APPLICATION Aspirin Soluble tabs 300 mg Chloroform	12 tabs to 100 ml	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml) Phenobarbitone Sodium	
CODEINE LINCTUS PAEDIATRIC (3 mg per 5 ml) Codeine phosphate Glycerol Preservative	60 mg 40 ml qs	Glycerol BP Water PILOCARPINE ORAL LIQUID	4 ml to 40 ml
Water CODEINE LINCTUS DIABETIC (15 mg per 5 ml)	to 100 ml	Pilocarpine 4% eye drops Preservative Water	qs qs to 500 ml
Codeine phosphate Glycerol Preservative	300 mg 40 ml qs	(Preservative should be used if quantity supplied is than 5 days.)	
Water	to 100 ml	SALIVA SUBSTITUTE FORMULA Methylcellulose	5 g
FOLINIC MOUTHWASH Calcium folinate 15 mg tab Preservative Water	1 tab qs to 500 ml	Preservative Water (Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	qs to 500 ml for more
(Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	for more	SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml	qs
MAGNESIUM HYDROXIDE 8% MIXTURE Magnesium hydroxide paste 29% Methyl hydroxybenzoate	275 g 1.5 g	Water (Only funded if prescribed for treatment of hyponatra	qs
Water	to 1,000 m	I VANCOMYCIN ORAL SOLUTION (50 mg per ml) Vancomycin 500 mg injection	10 vials
METHADONE MIXTURE Methadone powder Glycerol Water	qs qs to 100 ml	Glycerol BP Water (Only funded if prescribed for treatment of Clostridiu following metronidazole failure)	40 ml to 100 ml m difficile
METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liqu	10 g to 100 ml iid mixture)	VOSOL EAR DROPS WITH HYDROCORTISONE POWDER 1% Hydrocortisone powder Vosol Ear Drops	1% to 35 ml
OMEPRAZOLE SUSPENSION Omeprazole capules or powder Sodium bicarbonate powder BP Water	qs 8.4 g to 100 ml		

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy		Fully	Brand or
	(Manufacturer's Price	,	sidised	Generic
	\$	Per	1	Manufacturer
		-		
Extemporaneously Compounded Preparations a	and Galenical	S		
BENZOIN				
Tincture compound BP	24 42	500 ml		
	(39.90)	000 111		Pharmacy Health
	2.44	50 ml		r namaoy nounn
	(5.10)	00111		Pharmacy Health
OUL ODOFODM Only in combination	(0.10)			r namaoy noam
CHLOROFORM – Only in combination				
Only in aspirin and chloroform application.	05 50	500 ml		DOM
Chloroform BP		500 ml	•	PSM
CODEINE PHOSPHATE - Safety medicine; prescriber may dete		frequency		
Powder – Only in combination	63.09	25 g		
	(90.09)			Douglas
a) Only in extemporaneously compounded codeine linct	tus diabetic or cod	eine linctus	paediat	tric.
 b) Safety cap for extemporaneously compounded oral I 	iquid preparations.			
COLLODION FLEXIBLE				
Collodion flexible		100 ml	1	PSM
COMPOUND HYDROXYBENZOATE – Only in combination				
Only in extemporaneously compounded oral mixtures. Soln	20.00	100 ml		Midwest
S001		100 111		David Craig
	34.10		•	David Graig
GLYCERIN WITH SODIUM SACCHARIN - Only in combination				
Only in combination with Ora-Plus.				
Suspension		473 ml	-	Ora-Sweet SF
GLYCERIN WITH SUCROSE – Only in combination				
Only in combination with Ora-Plus.				
Suspension		473 ml	✓	Ora-Sweet
GLYCEROL				
* Liquid – Only in combination	3 28	500 ml	1	healthE Glycerol BP
Only in extemporaneously compounded oral liquid prepa		000 111	-	
MAGNESIUM HYDROXIDE				
Paste 29%	00.61	500 a		PSM
	22.01	500 g	•	FJIM
METHADONE HYDROCHLORIDE				
 a) Only on a controlled drug form 				
 b) No patient co-payment payable 				
c) Safety medicine; prescriber may determine dispensing free	equency			
d) Extemporaneously compounded methadone will only be i	reimbursed at the i	rate of the cl	neapes	t form available
(methadone powder, not methadone tablets).				
Powder		1 g	~	AFT
‡ Safety cap for extemporaneously compounded oral liqui	d preparations.			
METHYL HYDROXYBENZOATE				
Powder	8.00	25 g	✓	PSM
	8.98	-	✓	Midwest
METHYLCELLULOSE				
Powder	36.95	100 g	1	MidWest
Suspension – Only in combination		473 ml		Ora-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCH			-	
METHYLCELLOLOSE WITH GLYCERIN AND SODIOM SACCH. Suspension				Ora-Blend SF
JUSUE[[SI0[]	32.50	473 ml	v	Ula-Diellu SF

228

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy		Fully	Brand or
	(Manufacturer's Pric	e)	Subsidised	
	\$	Per	1	Manufacturer
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Onl	y in combination			
Suspension		473 m	I 🖌	Ora-Blend
PHENOBARBITONE SODIUM				
Powder – Only in combination		10 g	✓	MidWest
	325.00	100 g	✓	MidWest
a) Only in children up to 12 years		-		
b)‡ Safety cap for extemporaneously compounded oral I	iquid preparations.			
PROPYLENE GLYCOL				
Only in extemporaneously compounded methyl hydroxybenz	oate 10% solution.			
Liq		500 m	✓	Midwest
SODIUM BICARBONATE				
Powder BP – Only in combination		500 q	✓	Midwest
	9.80	0		
	(29.50)			David Craig
Only in extemporaneously compounded omeprazole and	l lansoprazole susp	ension.		-
SYRUP (PHARMACEUTICAL GRADE) - Only in combination				
Only in extemporaneously compounded oral liquid preparatic	ons.			
Liq		2,000 n	nl 🗸	Midwest
WATER				
Tap – Only in combination	0.00	1 ml	1	Tap water

EXPLANATORY NOTES

The list of special foods to which Subsidies apply is contained in this section. The list of available products, guidelines for use, subsidies and charges is reviewed as required. Applications for new listings and changes to subsidies and access criteria will be considered by the special foods sub-committee of PTAC which meets as and when required. In all cases, subsidies are available by Special Authority only. This means that, unless a patient has a valid Special Authority number for their special food requirements, they must pay the full cost of the products themselves.

Eligibility for Special Authority

Special Authorities will be approved for patients meeting conditions specified under the *Conditions and Guidelines* for each product. In some cases there are also limits to how products can be prescribed (for example quantity, use or duration). Only those brands, presentations and flavours of special foods listed in this section are subsidised.

Who can apply for Special Authority?

 Initial Applications:
 Only from a dietitian, relevant specialist or a vocationally registered general practitioner.

 Reapplications:
 Only from a dietitian, relevant specialist or a vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or a vocationally registered general practitioners.

 Weight of the second seco

All applications must be made on an official form available from the PHARMAC website www.pharmac.govt.nz. All applications must include specific details as requested on the form relating to the application. Applications must be forwarded to:

Ministry of Health Sector Services Private Bag 3015 WHANGANUI 4540 Freefax 0800 100 131

Subsidies and manufacturer's surcharges

The Subsidies for some special foods are based on the lowest priced product within each group. Where this is so, or where special foods are otherwise not fully subsidised, a manufacturer's surcharge may be payable by the patient. The manufacturer's surcharge is the difference between the price of the product and the subsidy attached to it and may be subject to mark-ups applied at a pharmacy level. As a result the manufacturer's surcharge may vary. Fully subsidised alternatives are available in most cases (as indicated by a tick in the left hand column). Patients should only have to pay a co-payment on these products.

Where are special foods available from?

Distribution arrangements for special foods vary from region to region. Special foods are available from hospital pharmacies providing an outpatient dispensing service as well as retail pharmacies in the Northern, Midland and Central (including Nelson and Blenheim) regions.

Definitions

 Failure to thrive
 An inability to gain or maintain weight resulting in physiological impairment.

 Growth deficiency
 Where the weight of the child is less than the fifth or possibly third percentile for their age, with evidence of malnutrition.

(Manufacturer's Price)

Per

Subsidy

\$

Fully Subsidised

Generic Manufacturer

Brand or

Nutrient Modules

Carbohydrate

■SA1522 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Fither:

- 1 cvstic fibrosis: or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia: or
- 5 premature and post premature infant; or
- 6 inborn errors of metabolism: or
- 7 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment: and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1522 above - Hospital pharmacy [HP3] Powder 5.29 400 a OP Polvcal

Carbohydrate And Fat

■SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

Subsidy		Fully	Brand or	
(Manufacturer's F	Price) S	ubsidised	Generic	
\$	Per	1	Manufacturer	

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE AND FAT SUI	PLEMENT - Special Author	ity see SA1376 on	the previous pa	ige -	Hospital pharmacy [HP3]
Powder (neutral)	-		400 g OP	1	Duocal Super
			-		Soluble Powder

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome; or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

- 10 ascites; or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** — (Inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Patho

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Authority see SA1523 on the previous page - Hospital pharmacy [HP3]

Emulsion (neutral)		200 ml OP	✓ Calogen
	30.75	500 ml OP	 Calogen
Emulsion (strawberry)		200 ml OP	 Calogen
Oil		500 ml OP	 MCT oil (Nutricia)
Oil, 250 ml		4 OP	 Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	armacy [HP3]	T – Special Authority see SA1524 above – Hospital pha	PROTEIN SUPPLEMENT
 Protifar 	225 g OP		Powder
 Resource 	227 g OP	8.95	
Banan	•		

Resource Beneprotein

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
\$	Per	1	Manufacture

Oral Supplements/Complete Diet (Nasogastric/Gastrostomy Tube Feed)

Respiratory Products

⇒SA1094 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has CORD and hypercapnia, defined as a CO2 value exceeding 55 mmHg.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CORD ORAL FEED 1.5KCAL/ML - Special Authority see SA	A1094 above – Hospi	tal pharmacy [H	HP3]
Liquid	1.66	237 ml OP	 Pulmocare

Diabetic Products

➡SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML – Special Authority see SA1095 above Liquid7.50	e – Hospital pharn 1,000 ml OP	nacy [HP3] ✓ Diason RTH ✓ Glucerna Select RTH
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 above - H	ospital pharmacy	[HP3]
Liquid (strawberry)1.50	200 ml OP	 Diasip
Liquid (vanilla)	200 ml OP	 Diasip
1.88	250 ml OP	 Glucerna Select
1.78	237 ml OP	
(2.10)		Resource Diabetic
(2.10)		Sustagen Diabetic

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED – Special Authority see SA1525 abov	e – Hospital pharma	cy [HP3]	
Powder	60.48	400 g OP	🗸 Monogen

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per ✓	Manufacturer
ENTERAL/ORAL FEED 1KCAL/ML – Special Authority see SA1 Liquid		• · ·	harmacy [HP3] Kindergen

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML – Special Author		ove – Hospital p	oharmacy [HP3]
Liquid		500 ml OP	✓ Nutrini Energy RTH
PAEDIATRIC ENTERAL FEED 1KCAL/ML – Special Authority		e – Hospital ph 500 ml OP	armacy [HP3] ✓ Nutrini RTH ✓ Pediasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML – S		e SA1379 abov	 re – Hospital pharmacy [HP3] ✓ Nutrini Energy Multi
Liquid		500 ml OP	Fibre
PAEDIATRIC ORAL FEED – Special Authority see SA1379 al Powder (vanilla)		armacy [HP3] 850 g OP	 Pediasure
PAEDIATRIC ORAL FEED 1.5KCAL/ML – Special Authority s Liquid (strawberry) Liquid (vanilla)	1.60	 Hospital pharr 200 ml OP 200 ml OP 	macy [HP3] ✓ Fortini ✓ Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML – Special Authority see Liquid (chocolate) Liquid (strawberry) Liquid (vanilla)	1.07 1.07	Hospital pharma 200 ml OP 200 ml OP 200 ml OP 250 ml OP	acy [HP3] Pediasure Pediasure Pediasure Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML – Spec		A1379 above –	Hospital pharmacy [HP3]
Liquid (chocolate)		200 ml OP	✓ Fortini Multi Fibre
Liquid (strawberry)		200 ml OP	✓ Fortini Multi Fibre
Liquid (vanilla)		200 ml OP	✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED – Special Authority see SA13		l pharmacy [HP	P3]
Powder		400 g OP	✓ Peptamen Junior

	Subsidy (Manufacturer's Pric \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
Renal Products				
 SA1101 Special Authority for Subsidy Initial application only from a dietitian, relevant specialist or voca years where the patient has acute or chronic kidney disease. Renewal only from a dietitian, relevant specialist, vocationally regrecommendation of a dietitian, relevant specialist or vocationally regrecommendations meeting the following criteria: Both: 1 The treatment remains appropriate and the patient is bene 2 General Practitioners must include the name of the dietitian practitioner and date contacted. 	gistered general pra registered general efiting from treatme	actitione practitio nt; and	r or general ner. Approv	practitioner on the als valid for 3 years for
RENAL ENTERAL FEED 1.8 KCAL/ML – Special Authority see S Liquid		ospital p 500 ml (P3] epro HP RTH
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see SA11 Liquid		al pharr 220 ml (OP 🖌 🖌 No	epro HP (strawberry) epro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML – Special Authority see SA110 Liquid Liquid (apricot) 125 ml Liquid (caramel) 125 ml	2.88 (3.31) 	pharma 237 ml (4 OP 4 OP) P ✓ R e	ovaSource Renal enilon 7.5 enilon 7.5

Specialised And Elemental Products

■ SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's P \$	Price) Subsi Per	Fully idised	Brand or Generic Manufacturer
ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML – Spe pharmacy [HP3] Liquid	,	e SA1377 on th 1,000 ml OP	e previ	
ORAL ELEMENTAL FEED 0.8KCAL/ML – Special Authority see Liquid (grapefruit), 250 ml carton Liquid (pineapple & orange), 250 ml carton Liquid (summer fruits), 250 ml carton		previous page - 18 OP 18 OP 18 OP 18 OP	✓ E ✓ E	tal pharmacy [HP3] lemental 028 Extra lemental 028 Extra lemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML – Special Authority see S Powder (unflavoured)		evious page – H 80 g OP		l pharmacy [HP3] i vonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML – Special Auth [HP3] Liquid		7 on the previou 1,000 ml OP		e – Hospital pharmacy eptisorb

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
 - 1 Child aged one to eight years; and
 - 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTE	RAL FEED WIT	H FIBRE 0.76 KCAL/	ML – Special A	Authority	see SA1196 a	bove -	- Hospital pharm	acy [HP3]
Liquid				4.00	500 ml OP	✓	Nutrini Low Er	ergy
							Multi Fibre	

Standard Supplements

⇒SA1554 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and

3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal - (Children - indications other than exclusive enteral nutrition for Crohn's disease) only from a dietitian, relevant

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist, dietitian on the recommendation of a gastroenterologist or vocationally registered general practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:
 - Patient has not responded to first-line dietary measures over a 4 week period by:
 - 2.1 Increasing their food intake frequency (eg snacks between meals); or
 - 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
 - 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Subsidy		ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	✓	Manufacturer

Initial application — (Short-term medical condition) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or</p>
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or</p>
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or

Subsidy	(Manufacturer's Price) Subsidised		Brand or Generic	
(Manufacturer's Price)				
\$	Per	✓	Manufacturer	

- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML – Special Authority see SA1554 on page 238 – Liquid	Hospital pharmacy [HP3] 1,000 ml OP ✓ Nutrison Energy	
ENTERAL FEED 1KCAL/ML – Special Authority see SA1554 on page 238 – H Liquid	lospital pharmacy [HP3] 250 ml OP ✓ Isosource Standar 1,000 ml OP ✓ Isosource Standar RTH ✓ Nutrison Standard RTH ✓ Osmolite RTH	ď
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML – Special Authority see SA1554 Liquid		
ENTERAL FEED WITH FIBRE 1 KCAL/ML – Special Authority see SA1554 on Liquid	n page 238 – Hospital pharmacy [HP3] 1,000 ml OP ✓ Jevity RTH ✓ Nutrison Multi Fib	re
ENTERAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority see SA1554 o Liquid1.75 7.00	on page 238 – Hospital pharmacy [HP3] 250 ml OP 1,000 ml OP V Ensure Plus RTH V Jevity HiCal RTH Nutrison Energy Multi Fibre	

	Subsidy		Fully	Brand or
	(Manufacturer's F	Price) Subs	idised	Generic
	`\$	Per	1	Manufacturer
ORAL FEED (POWDER) - Special Authority see SA1554 on page				
Note: Higher subsidy for Sustagen Hospital Formula will only	y be reimbursed	for patients wit	h both	a valid Special Authority
number and an appropriately endorsed prescription.				
Powder (chocolate) - Higher subsidy of up to \$26.00 per 85	0 a			
with Endorsement		050 ~ OD	./	Ensure
		850 g OP	v	Ensure
	9.54	840 g OP		
	(26.00)			Sustagen Hospital
				Formula
	(26.00)			Sustagen Hospital
	(20.00)			
				Formula Active
Additional subsidy by endorsement is available for patien	nts with fat mala	bsorption, fat in	tolera	nce or chyle leak. The
prescription must be endorsed accordingly.				
Powder (vanilla) - Higher subsidy of up to \$26.00 per 850 g				
	0.54	057 00		F
with Endorsement		857 g OP		Fortisip
	26.00	850 g OP	~	Ensure
	9.54	840 g OP		
	(26.00)	0		Sustagen Hospital
	(20100)			Formula
	(22.22)			
	(26.00)			Sustagen Hospital
				Formula Active
Additional subsidy by endorsement is available for patier	nts with fat mala	bsorption, fat in	tolera	nce or chyle leak. The
prescription must be endorsed accordingly.				,
(Sustagen Hospital Formula Powder (chocolate) to be delisted 1				
(Sustagen Hospital Formula Powder (vanilla) to be delisted 1 Oct	ober 2018)			
ORAL FEED 1.5KCAL/ML - Special Authority see SA1554 on pa		tal pharmaoy [L	וכסו	
Additional subsidy by endorsement is available for patients b				
epidermolysis bullosa, or as exclusive enteral nutrition in chil	dren under the a	age of 18 years	for the	e treatment of Crohn's
disease. The prescription must be endorsed accordingly.				
Liguid (banana) - Higher subsidy of \$1.26 per 200 ml with				
	0.70	200 ml OP		
Endorsement		200 mi OP		
	(1.26)			Ensure Plus
	(1.26)			Fortisip
Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with				
Endorsement		200 ml OP		
		200 IIII OF		
	(1.26)			Ensure Plus
	(1.26)			Fortisip
Liquid (fruit of the forest) – Higher subsidy of \$1.26 per 200	ml			
with Endorsement		200 ml OP		
		200 IIII OF		
	(1.26)			Ensure Plus
Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml wit	h			
Endorsement	0.72	200 ml OP		
	(1.26)			Ensure Plus
	· · ·			
· · · · · · · · · · · · · · · · · · ·	(1.26)			Fortisip
Liquid (vanilla) – Higher subsidy of up to \$1.33 per 237 ml w	ith			
Endorsement	0.85	237 ml OP		
	(1.33)			Ensure Plus
	0.72	200 ml OP		
	(1.26)			Ensure Plus
	(1.26)			Fortisip

	Subsidy		Fully	Brand or
	(Manufacturer's F	Price) Subsid	dised	Generic
	\$	Per	1	Manufacturer
ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see	SA1554 on pag	ne 238 – Hospital	l pharr	nacv [HP3]
Additional subsidy by endorsement is available for patients b				
epidermolysis bullosa. The prescription must be endorsed a	0			
Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with	0,			
Endorsement		200 ml OP		
		200 111 0F	-	artiain Multi Fibra
	(1.26)		Г	ortisip Multi Fibre
Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml wit				
Endorsement	0.72	200 ml OP		
	(1.26)		F	ortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with				
Endorsement	0 72	200 ml OP		
	(1.26)	200 01	F	ortisip Multi Fibre
	(1.20)		'	

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- All of the following:
 - 1 Cystic fibrosis; and
 - 2 other lower calorie products have been tried; and
 - 3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML – Special Authority see SA1195 abo	ove – Hospital j	oharmacy [HP3]	
Liquid	5.50	500 ml OP	 Nutrison
			Concentrated
	11.00	1,000 ml OP	Two Cal HN RTH

	Subsidy (Manufacturer's Pr \$	rice) Subs Per	Fully sidised	Brand or Generic Manufacturer
DRAL FEED 2 KCAL/ML – Special Authority see SA1195 on the Additional subsidy by endorsement is available for patients epidermolysis bullosa. The prescription must be endorsed Liquid (vanilla) – Higher subsidy of \$1.90 per 200 ml with	being bolus fed thr			
Endorsement	0.96 (1.90)	200 ml OP	T۱	wo Cal HN
Food Thickeners				
 Renewal only from a dietitian, relevant specialist, vocationally recommendation of a dietitian, relevant specialist or vocationally pplications meeting the following criteria: Both: The treatment remains appropriate and the patient is being 2 General Practitioners must include the name of the dietit practitioner and date contacted. 	y registered genera	Il practitioner. nent; and	Approv	als valid for 1 year for
OOD THICKENER – Special Authority see SA1106 above – H Powder		[HP3] 300 g OP 380 g OP		utilis eed Thickener Karicare Aptamil
Gluten Free Foods				
The funding of gluten free foods is no longer being actively man		C from 1 Anril	0011	T h.:

The funding of gluten free foods is no longer being actively managed by PHARMAC from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

➡SA1107 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

GLUTEN FREE BAKING MIX – Special Authority see SA1107 above – Hosp Powder	ital pharmacy [HP3] 1,000 g OP	
(5.15)	Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA1107 above - Hospi	al pharmacy [HP3]	
Powder	1,000 g OP	
(7.32)	NZB Low Gluten Bread Mix
3.51		
(10.87)	Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1107 above - Hospital ph	armacy [HP3]	
Powder	2,000 g OP	
(18.10)	Horleys Flour

	Subsidy (Manufacturer's P		Fully Brand or lised Generic
	(Manulacturers F	Per	Manufacturer
GLUTEN FREE PASTA – Special Authority see SA1107 on	the previous page -	Hospital pharma	icy [HP3]
Buckwheat Spirals	2.00	250 g OP	
	(3.11)		Orgran
Corn and Vegetable Shells	2.00	250 g OP	
	(2.92)		Orgran
Corn and Vegetable Spirals	2.00	250 g OP	
	(2.92)		Orgran
Rice and Corn Lasagne Sheets	1.60	200 g OP	
	(3.82)		Orgran
Rice and Corn Macaroni		250 g OP	
	(2.92)		Orgran
Rice and Corn Penne		250 g OP	-
	(2.92)	-	Orgran
Rice and Maize Pasta Spirals		250 g OP	-
	(2.92)	-	Orgran
Rice and Millet Spirals		250 g OP	-
	(3.11)	•	Orgran
Rice and corn spaghetti noodles		375 g OP	-
	(2.92)	•	Orgran
Vegetable and Rice Spirals		250 g OP	-
- ,	(2.92)	ů.	Orgran
Italian long style spaghetti	· · ·	220 g OP	2
	(3.11)	e e	Orgran

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

AMINOACID FORMULA WITHOUT METHIONINE	- Special Authority see SA1108	<mark>3 above</mark> – Hospi	tal pharmacy [HP3]
Powder		500 g OP	 XMET Maxamum

Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE	- Special	Authority see	SA1108 above – Hospital
pharmacy [HP3]		-	
Powder	22 50	0 a OP 🖌	MSUD Maxamum

	Subsidy (Manufacturer's \$	Price) Subs Per	Fully Brand or sidised Generic ✓ Manufacturer	
Supplements For PKU				
MINOACID FORMULA WITHOUT PHENYLALANINE – Spe harmacy [HP3]	ecial Authority see	SA1108 on the p	previous page – Hospita	al
Tabs		75 OP	Phiexy 10	
Powder (unflavoured) 27.8 g sachets	936.00	30	 PKU Lophlex Powder 	
Powder (unflavoured) 36 g sachets		30	🗸 PKU Anamix Ju	nior
Infant formula	174.72	400 g OP	PKU Anamix Infa	ant
Powder (orange)	221.00	500 g OP	🗸 XP Maxamaid	
	320.00	-	🗸 XP Maxamum	
Powder (unflavoured)	221.00	500 g OP	🗸 XP Maxamaid	
	320.00		🗸 XP Maxamum	
Liquid (berry)	13.10	125 ml OP	 PKU Anamix Jui LQ 	nior
Liquid (orange)	13.10	125 ml OP	 PKU Anamix Jui LQ 	nioı
Liquid (unflavoured)	13.10	125 ml OP	 PKU Anamix Jui LQ 	nioı
Liquid (forest berries), 250 ml carton		18 OP	 Easiphen Liquid 	
Liquid (juicy tropical) 125 ml		30 OP	PKU Lophlex LC	20
Oral semi-solid (berries) 109 g	1,123.20	36 OP	 PKU Lophlex Sensation 20 	
Liquid (juicy berries) 62.5 ml		60 OP	PKU Lophlex LG	10
Liquid (juicy citrus) 62.5 ml		60 OP	PKU Lophlex LG	
Liquid (juicy orange) 62.5 ml		60 OP	PKU Lophlex LG	
Liquid (juicy berries) 125 ml		30 OP	PKU Lophlex LG	20
Liquid (juicy citrus) 125 ml		30 OP	PKU Lophlex LG	
Liquid (juicy orange) 125 ml		30 OP	PKU Lophlex LC	20
PKU Lophlex LQ 20 Liquid (juicy citrus) 125 ml to be delisted			-	

Foods

LOW PROTEIN BAKING MIX – Special Authority see SA1108 on the previou Powder		pharmacy [HP3] ✓ Loprofin Mix
LOW PROTEIN PASTA - Special Authority see SA1108 on the previous page	e – Hospital pharm	acy [HP3]
Animal shapes11.91	500 g OP	 Loprofin
Lasagne	250 g OP	 Loprofin
Low protein rice pasta11.91	500 g OP	 Loprofin
Macaroni	250 g OP	 Loprofin
Penne	500 g OP	 Loprofin
Spaghetti	500 g OP	 Loprofin
Spirals	500 g OP	 Loprofin

Infant Formulae

For Premature Infants

PRETERM POST-DISCHARGE INFANT FORMULA - Spe	ecial Authority see SA1	198 on the nex	kt page – Hospital pharmacy
[HP3]			
Powder		400 g OP	 S-26 Gold Premgro
(S-26 Gold Premgro Powder to be delisted 1 July 2018)			

	Subsidy	Fu	ıllv	Brand or
	acturer's Price)	Subsidis	ed	Generic
Υ.	\$	Per	1	Manufacturer

⇒SA1198 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 The infant was born before 33 weeks gestation or weighed less than 1.5 kg at birth; and

2 Either:

- 2.1 The infant has faltering growth (downward crossing of percentiles); or
- 2.2 The infant is not maintaining, or is considered unlikely to maintain, adequate growth on standard infant formula.

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - H	lospital pharmac	y [HP3]
Powder	400 g OP	 Locasol

Gastrointestinal and Other Malabsorptive Problems

Powder4	3.60	400 g OP	 Alfamino Junior
5	3.00		Neocate LCP
Powder (unflavoured)5	3.00	400 g OP	 Elecare
		•	 Elecare LCP
			Neocate Advance
			Neocate Gold
			✓ Neocate Junior
			Unflavoured
Powder (vanilla)5	3.00	400 g OP	 Elecare
		0	Neocate Advance
			✓ Neocate Junior

(Neocate Advance Powder (unflavoured) to be delisted 1 September 2018) (Neocate Advance Powder (vanilla) to be delisted 1 September 2018)

⇒SA1219 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria: Any of the following:

1 Extensively hydrolysed formula has been reasonably trialled and is inappropriate due to documented severe intolerance or allergy or malabsorption; or

continued...

Vanilla

Subsidy (Manufacturer's Price)		Fully ubsidised	Brand or Generic	
\$	Per	~	Manufacturer	

- 2 History of anaphylaxis to cows milk protein formula or dairy products; or
- 3 Eosinophilic oesophagitis.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

			Junior
Powder	15.21	450 g OP	 Aptamil Gold+ Pepti
EXTENSIVELY HYDROLYSED FORMULA - Special Authority see \$	SA1557 belo	w – Hospital ph	armacy [HP3]

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- Any of the following:
 - 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
 - 2 Severe malabsorption; or
 - 3 Short bowel syndrome; or
 - 4 Intractable diarrhoea; or
 - 5 Biliary atresia; or
 - 6 Cholestatic liver diseases causing malsorption; or
 - 7 Cystic fibrosis; or
 - 8 Proven fat malabsorption; or
 - 9 Severe intestinal motility disorders causing significant malabsorption; or
 - 10 Intestinal failure; or
 - 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.
- Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

\$ Per 🖌 Manufacturer	Subsidy (Manufacturer's Price)	Su	Fully ubsidised	Brand or Generic	
	\$	Per	1	Manufacturer	

Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML - Special Authority see SA1698 below - Hospital pharmacy [HP3]

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Author	ority see SA1197	above – Retail	pharmacy
Powder (unflavoured)		300 g OP	KetoCal 4:1
		-	Ketocal 3:1
Powder (vanilla)	35.50	300 g OP	 KetoCal 4:1

Pharmaceuticals and quantities that may be obtained on a Practitioner's Supply Order

ADRENALINE	
✓ Inj 1 in 1,000, 1 ml ampoule	
✓ Inj 1 in 10,000, 10 ml ampoule	
AMINOPHYLLINE	
✓ Inj 25 mg per ml, 10 ml ampoule	
AMIODARONE HYDROCHLORIDE	
✓ Inj 50 mg per ml, 3 ml ampoule5	
AMOXICILLIN	
✓ Cap 250 mg	
✓ Cap 500 mg	
✓ Grans for oral liq 125 mg per 5 ml 200 ml	
 Grans for oral liq 250 mg per 5 ml	
✓ Inj 1 g vial5	
AMOXICILLIN WITH CLAVULANIC ACID	
✓ Tab 500 mg with clavulanic acid 125 mg	
 Grans for oral liq amoxicillin 25 mg with clavulanic 	
acid 6.25 mg per ml200 ml	
 Grans for oral liq amoxicillin 50 mg with clavulanic 	
acid 12.5 mg per ml 200 ml	
ASPIRIN	
✓ Tab dispersible 300 mg	
ATROPINE SULPHATE	
✓ Inj 600 mcg per ml, 1 ml ampoule	
AZITHROMYCIN	
✓ Tab 500 mg – See note on page 100	
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]	
✓ Tab 2.5 mg – See note on page 64150	
BENZATHINE BENZYLPENICILLIN	
✓ Inj 900 mg (1.2 million units) in 2.3 ml syringe5	
BENZATROPINE MESYLATE	
✓ Inj 1 mg per ml, 2 ml10	
BENZYLPENICILLIN SODIUM [PENICILLIN G]	
✓ Inj 600 mg (1 million units) vial	
BLOOD KETONE DIAGNOSTIC TEST STRIP	
BLOOD KETONE DIAGNOSTIC TEST STRIP	
 Test strips – Subsidy by endorsement – See note 	
 Test strips – Subsidy by endorsement – See note on page 2510 	
 Test strips – Subsidy by endorsement – See note on page 2510 BLOOD GLUCOSE DIAGNOSTIC TEST METER 	
 ✓ Test strips – Subsidy by endorsement – See note on page 25	
 Test strips – Subsidy by endorsement – See note on page 25	
 Test strips – Subsidy by endorsement – See note on page 25	
 Test strips – Subsidy by endorsement – See note on page 25	
 Test strips – Subsidy by endorsement – See note on page 25	
 Test strips – Subsidy by endorsement – See note on page 25	
 Test strips – Subsidy by endorsement – See note on page 25	
 Test strips – Subsidy by endorsement – See note on page 25	
 Test strips – Subsidy by endorsement – See note on page 25	

CEFTRIAXONE
 Inj 500 mg vial – Subsidy by endorsement – See
note on page 995
Inj 1 g vial – Subsidy by endorsement – See note
on page 995
CHARCOAL
 Oral liq 50 g per 250 ml
CHLORPROMAZINE HYDROCHLORIDE
✓ Tab 10 mg
✓ Tab 25 mg
✓ Tab 100 mg30
✓ Inj 25 mg per ml, 2 ml5
CIPROFLOXACIN
✓ Tab 250 mg – See note on page 1045
✓ Tab 500 mg – See note on page 1045
COMPOUND ELECTROLYTES
 Powder for oral soln10
CONDOMS
✓ 49 mm
✓ 53 mm
 53 mm (chocolate)
 53 mm (strawberry)
✓ 56 mm
✓ 56 mm, shaped
CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL
 Tab 2 mg with ethinyloestradiol 35 mcg and
7 inert tabs
DEXAMETHASONE
 Tab 0.5 mg – Retail pharmacy-Specialist
 Tab 0.5 mg – Retail pharmacy-Specialist
DEXAMETHASONE PHOSPHATE
✓ Inj 4 mg per ml, 1 ml ampoule – See note on page 895
 Inj 4 mg per ml, 2 ml ampoule – See note on page 895
DIAZEPAM
 Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement – See note on page 1375
 Rectal tubes 5 mg
 Rectal tubes 10 mg
DICLOFENAC SODIUM
 Inj 25 mg per ml, 3 ml ampoule
✓ Suppos 50 mg
DIGOXIN
✓ Tab 62.5 mcg
✓ Tab 250 mcg
DOXYCYCLINE
Tab 50 mg
✓ Tab 100 mg
continued

fully subsidised brand available

Please refer to Section A for a definition, and conditions of supply, of Practitioner's Supply Orders.

250

(continued)
DUAL BLOOD GLUCOSE AND BLOOD KETONE
DIAGNOSTIC TEST METER
 Meter with 50 lancets, a lancing device and
10 blood glucose diagnostic test strips –
Subsidy by endorsement – See note on page 261
ERGOMETRINE MALEATE
✓ Inj 500 mcg per ml, 1 ml ampoule
ERYTHROMYCIN ETHYL SUCCINATE
✓ Tab 400 mg
Grans for oral liq 200 mg per 5 ml
✓ Grans for oral liq 400 mg per 5 ml 200 ml ERYTHROMYCIN STEARATE
Tab 250 mg
ETHINYLOESTRADIOL WITH DESOGESTREL
Tab 20 mcg with desogestrel 150 mcg and 7 inert tab84
Tab 30 mcg with desogestrel 150 mcg and 7 inert tab 84
ETHINYLOESTRADIOL WITH LEVONORGESTREL
✓ Tab 20 mcg with levonorgestrel 100 mcg and
7 inert tablets
✓ Tab 50 mcg with levonorgestrel 125 mcg and
7 inert tab
Tab 30 mcg with levonorgestrel 150 mcg
✓ Tab 30 mcg with levonorgestrel 150 mcg and
7 inert tablets
ETHINYLOESTRADIOL WITH NORETHISTERONE
✓ Tab 35 mcg with norethisterone 1 mg63
✓ Tab 35 mcg with norethisterone 1 mg and 7 inert tab84
✓ Tab 35 mcg with norethisterone 500 mcg
 Tab 35 mcg with norethisterone 500 mcg and
7 inert tab84
FLUCLOXACILLIN
✓ Cap 250 mg
✓ Grans for oral liq 25 mg per ml 200 ml
✓ Grans for oral liq 50 mg per ml 200 ml
✓ Inj 1 g vial5
FLUPENTHIXOL DECANOATE
✓ Inj 20 mg per ml, 1 ml5
✓ Inj 20 mg per ml, 2 ml
✓ Inj 100 mg per ml, 1 ml
FUROSEMIDE [FRUSEMIDE]
✓ Tab 40 mg30
✓ Inj 10 mg per ml, 2 ml ampoule
GLUCAGON HYDROCHLORIDE
✓ Inj 1 mg syringe kit
GLUCOSE [DEXTROSE] ✓ Inj 50%, 10 ml ampoule
 ✓ Inj 50%, 10 mi ampoule
GLYCERYL TRINITRATE
✓ Tab 600 mcg
 Tab boo mcg
 Oral spray, 400 mcg per dose

✓ fully subsidised brand available

Please refer to Section A for a definition, and conditions of supply, of Practitioner's Supply Orders.

(continued)

LOPERAMIDE HYDROCHLORIDE
✓ Tab 2 mg
✓ Cap 2 mg
MASK FOR SPACER DEVICE
✓ Small – See note on page 214
MEDROXYPROGESTERONE ACETATE
✓ Inj 150 mg per ml, 1 ml syringe
METOCLOPRAMIDE HYDROCHLORIDE Inj 5 mg per ml, 2 ml ampoule
METRONIDAZOLE ✓ Tab 200 mg
MIDAZOLAM
Inj 1 mg per ml, 5 ml plastic ampoule – See note
on page 15710
✓ Inj 5 mg per ml, 3 ml plastic ampoule – See note
on page 1575
MORPHINE SULPHATE
 Inj 5 mg per ml, 1 ml ampoule – Only on a
controlled drug form5
 Inj 10 mg per ml, 1 ml ampoule – Only on a
controlled drug form5
✓ Inj 15 mg per ml, 1 ml ampoule – Only on a
controlled drug form
Inj 30 mg per mi, 1 mi ampoule – Only on a sentrelled drug forma
controlled drug form
NALOXONE HYDROCHLORIDE
NALOXONE HYDROCHLORIDE Inj 400 mcg per ml, 1 ml ampoule5
NALOXONE HYDROCHLORIDE Inj 400 mcg per ml, 1 ml ampoule5 NICOTINE
NALOXONE HYDROCHLORIDE ✓ Inj 400 mcg per ml, 1 ml ampoule5 NICOTINE ✓ Patch 7 mg – See note on page 16328
NALOXONE HYDROCHLORIDE ✓ Inj 400 mcg per ml, 1 ml ampoule .5 NICOTINE ✓ Patch 7 mg – See note on page 163
NALOXONE HYDROCHLORIDE ✓ Inj 400 mcg per ml, 1 ml ampoule .5 NICOTINE ✓ Patch 7 mg – See note on page 163
NALOXONE HYDROCHLORIDE ✓ Inj 400 mcg per ml, 1 ml ampoule .5 NICOTINE ✓ Patch 7 mg - See note on page 163
NALOXONE HYDROCHLORIDE ✓ Inj 400 mcg per ml, 1 ml ampoule .5 NICOTINE ✓ Patch 7 mg - See note on page 163
NALOXONE HYDROCHLORIDE ✓ Inj 400 mcg per ml, 1 ml ampoule .5 NICOTINE ✓ Patch 7 mg – See note on page 163 28 ✓ Patch 14 mg – See note on page 163 28 ✓ Patch 21 mg – See note on page 163 28 ✓ Lozenge 1 mg – See note on page 163 ✓ Lozenge 2 mg – See note on page 163 216 ✓ Gum 2 mg (Fruit) – See note on page 163 .384
NALOXONE HYDROCHLORIDE ✓ Inj 400 mcg per ml, 1 ml ampoule .5 NICOTINE ✓ Patch 7 mg – See note on page 163 .28 ✓ Patch 14 mg – See note on page 163 .28 ✓ Patch 21 mg – See note on page 163 .28 ✓ Lozenge 1 mg – See note on page 163 ✓ Lozenge 2 mg – See note on page 163 .216 ✓ Gum 2 mg (Fruit) – See note on page 163 .384 ✓ Gum 4 mg (Fruit) – See note on page 163
NALOXONE HYDROCHLORIDE ✓ Inj 400 mcg per ml, 1 ml ampoule .5 NICOTINE ✓ Patch 7 mg – See note on page 163
NALOXONE HYDROCHLORIDE ✓ Inj 400 mcg per ml, 1 ml ampoule .5 NICOTINE ✓ Patch 7 mg – See note on page 163 28 ✓ Patch 14 mg – See note on page 163 28 ✓ Patch 21 mg – See note on page 163 28 ✓ Lozenge 1 mg – See note on page 163 ✓ Lozenge 2 mg – See note on page 163 ✓ Gum 2 mg (Fruit) – See note on page 163 ✓ Gum 2 mg (Mint) – See note on page 163 ✓ Gum 4 mg (Fruit) – See note on page 163 384 ✓ Gum 4 mg (Fruit) – See note on page 163 384 ✓ Gum 4 mg (Mint) – See note on page 163 384 ✓ Gum 4 mg (Mint) – See note on page 163
NALOXONE HYDROCHLORIDE ✓ Inj 400 mcg per ml, 1 ml ampoule .5 NICOTINE ✓ Patch 7 mg – See note on page 163
NALOXONE HYDROCHLORIDE ✓ Inj 400 mcg per ml, 1 ml ampoule MICOTINE ✓ Patch 7 mg – See note on page 163. 28 ✓ Patch 14 mg – See note on page 163. 28 ✓ Patch 21 mg – See note on page 163. 28 ✓ Lozenge 1 mg – See note on page 163. 216 ✓ Lozenge 2 mg – See note on page 163. 216 ✓ Gum 2 mg (Fruit) – See note on page 163. 28 ✓ Gum 2 mg (Mint) – See note on page 163. 384 ✓ Gum 4 mg (Fruit) – See note on page 163. 384 ✓ Gum 4 mg (Mint) – See note on page 163. 384 ✓ Gum 4 mg (Mint) – See note on page 163. 384 ✓ Gum 4 mg (Mint) – See note on page 163. 384 ✓ Gum 4 mg (Mint) – See note on page 163. 384 ✓ Gum 4 mg (Mint) – See note on page 163. 384 ✓ Gum 4 mg (Mint) – See note on page 163. 384 ✓ Gum 4 mg (Mint) – See note on page 163. 384 ✓ Gum 4 mg (Mint) – See note on page 163. 384 ✓ Gum 4 mg (Mint) – See note on page 163.<
NALOXONE HYDROCHLORIDE ✓ Inj 400 mcg per ml, 1 ml ampoule MICOTINE ✓ Patch 7 mg - See note on page 163. 28 ✓ Patch 14 mg - See note on page 163. 28 ✓ Patch 21 mg - See note on page 163. 28 ✓ Lozenge 1 mg - See note on page 163. 216 ✓ Lozenge 2 mg - See note on page 163. 216 ✓ Gum 2 mg (Fruit) - See note on page 163. 284 ✓ Gum 4 mg (Fruit) - See note on page 163. 384 ✓ Gum 4 mg (Fruit) - See note on page 163. 384 ✓ Gum 4 mg (Mint) - See note on page 163. 384 ✓ Gum 4 mg (Mint) - See note on page 163. 384 ✓ Gum 4 mg (Mint) - See note on page 163. 384 ✓ Gum 4 mg (Mint) - See note on page 163. 384 ✓ NORETHISTERONE ✓ Tab 350 mcg. ¾ Tab 5 mg.
NALOXONE HYDROCHLORIDE ✓ Inj 400 mcg per ml, 1 ml ampoule MICOTINE ✓ Patch 7 mg – See note on page 163. 28 ✓ Patch 14 mg – See note on page 163. 28 ✓ Patch 21 mg – See note on page 163. 28 ✓ Lozenge 1 mg – See note on page 163. ✓ Lozenge 2 mg – See note on page 163. ✓ Lozenge 2 mg – See note on page 163. ✓ Gum 2 mg (Fruit) – See note on page 163. ✓ Gum 4 mg (Fruit) – See note on page 163. ✓ Gum 4 mg (Fruit) – See note on page 163. ✓ Gum 4 mg (Fruit) – See note on page 163. ✓ Gum 4 mg (Mint) – See note on page 163. ✓ Gum 4 mg (Mint) – See note on page 163. ✓ Gum 4 mg (Mint) – See note on page 163. ✓ Tab 350 mcg. ✓ Tab 350 mcg. Yab 5 mg. 300 OXYTOCIN
NALOXONE HYDROCHLORIDE Inj 400 mcg per ml, 1 ml ampoule SNICOTINE Patch 7 mg - See note on page 163 Patch 14 mg - See note on page 163 Patch 21 mg - See note on page 163 Lozenge 1 mg - See note on page 163 Lozenge 2 mg - See note on page 163 Lozenge 2 mg - See note on page 163 Sum 2 mg (Fruit) - See note on page 163 Gum 2 mg (Kint) - See note on page 163 Gum 4 mg (Fruit) - See note on page 163 Sum 4 mg (Fruit) - See note on page 163 MORETHISTERONE Tab 350 mcg Tab 5 mg 30 OXYTOCIN Inj 5 iu per ml, 1 ml ampoule Sup 10 iu per ml, 1 ml ampoule Sup 10 iu per ml, 1 ml ampoule
NALOXONE HYDROCHLORIDE ✓ Inj 400 mcg per ml, 1 ml ampoule .5 NICOTINE ✓ Patch 7 mg – See note on page 163. .28 ✓ Patch 14 mg – See note on page 163. .28 ✓ Patch 21 mg – See note on page 163. .28 ✓ Lozenge 1 mg – See note on page 163. .216 ✓ Lozenge 2 mg – See note on page 163. .216 ✓ Gum 2 mg (Fruit) – See note on page 163. .384 ✓ Gum 4 mg (Fruit) – See note on page 163. .384 ✓ Gum 4 mg (Mint) – See note on page 163. .384 ✓ Gum 4 mg (Mint) – See note on page 163. .384 ✓ Gum 4 mg (Mint) – See note on page 163. .384 ✓ Gum 4 mg (Mint) – See note on page 163. .384 ✓ Gum 4 mg (Mint) – See note on page 163. .384 ✓ Gum 4 mg (Mint) – See note on page 163. .384 ✓ Gum 4 mg (Mint) – See note on page 163. .384 ✓ Gum 4 mg (Mint) – See note on page 163. .384 ✓ Gum 4 mg (Mint) – See note on page 163. .384 ✓ Tab 350 mcg. .84 ✓ Tab 5 mg. .30 OXYTOCIN .5 ✓ Inj 5 iu per ml, 1 ml ampoule .5 ✓ Inj 5 iu with ergometrine malea
NALOXONE HYDROCHLORIDE Inj 400 mcg per ml, 1 ml ampoule
NALOXONE HYDROCHLORIDE Inj 400 mcg per ml, 1 ml ampoule
NALOXONE HYDROCHLORIDE Inj 400 mcg per ml, 1 ml ampoule

PEAK FLOW METER PETHIDINE HYDROCHLORIDE ✓ Inj 50 mg per ml, 1 ml ampoule – Only on a controlled drug form......5 ✓ Inj 50 mg per ml, 2 ml ampoule – Only on a controlled drug form.....5 PHENOXYMETHYLPENICILLIN (PENICILLIN V) PHENYTOIN SODIUM ✓ Inj 50 mg per ml, 2 ml ampoule......5 PHYTOMENADIONE PIPOTHIAZINE PAI MITATE Inj 50 mg per ml, 1 ml – Subsidy by endorsement - See note on page 1475 ✓ Ini 50 mg per ml. 2 ml – Subsidy by endorsement - See note on page 1475 PREDNISOLONE PREDNISONE PREGNANCY TESTS - HCG URINE PROCAINE PENICILLIN PROCHLORPERAZINE PROMETHAZINE HYDROCHLORIDE SALBUTAMOL ✓ Inj 500 mcg per ml, 1 ml......5 ✓ Aerosol inhaler, 100 mcg per dose CFC free 1000 dose SALBUTAMOL WITH IPRATROPIUM BROMIDE Nebuliser soln. 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule20 SODIUM BICARBONATE ✓ Inj 8.4%, 100 ml5 continued...

fully subsidised brand available

Please refer to Section A for a definition, and conditions of supply, of Practitioner's Supply Orders.

.. ..

(continued)	
SODIUM CHLORIDE	
✓ Inj 0.9%, bag – See note on page 56	
 Inj 0.9%, 5 ml ampoule – See note on page 56 	5
 Inj 0.9%, 10 ml ampoule – See note on page 56 	5
SPACER DEVICE ✓ 220 ml (single patient)	50
✓ 510 ml (single patient)	
✓ 800 ml	50

SULFADIAZINE SILVER

~	Crm 1%	 g
•	Crm 1%	 ć

TRIMETHOPRIM

✓	′ Tab 300 mg	
---	--------------	--

TRIMETHOPRIM WITH SULPHAMETHOXAZOLE	
[CO-TRIMOXAZOLE]	
 Tab trimethoprim 80 mg and sulphamethoxazole 	
400 mg	30
 Oral liq 8 mg sulphamethoxazole 40 mg per 	
ml	.200 ml
VERAPAMIL HYDROCHLORIDE	
 Inj 2.5 mg per ml, 2 ml ampoule 	5
WATER	
 Inj 5 ml ampoule – See note on page 56 	5
 Inj 10 ml ampoule – See note on page 56 	5
 Inj 20 ml ampoule – See note on page 56 	5
ZUCLOPENTHIXOL DECANOATE	
 Inj 200 mg per ml, 1 ml 	5

Rural Areas for Practitioner's Supply Orders

NORTH ISLAND

Northland DHB Dargaville Hikurangi Kaeo Kaikohe Kaitaia Kawakawa Kerikeri Mangonui Maungaturoto Moerewa Ngunguru Paihia Rawene Ruakaka Russell Tutukaka Waipu Whangaroa

Waitemata DHB

Helensville Huapai Kumeu Snells Beach Waimauku Warkworth Wellsford

Auckland DHB

Great Barrier Island Oneroa Ostend

Counties Manukau DHB

Tuakau Waiuku

Waikato DHB

Coromandel Huntly Kawhia Matamata Morrinsville Ngatea Otorohanga Paeroa Pauanui Beach Putaruru Raglan Tairua Taumarunui Te Aroha Te Kauwhata Te Kuiti Tokoroa Waihi Whangamata Whitianga

Bay of Plenty DHB

Edgecumbe Katikati Kawerau Murupara Opotiki Taneatua Te Kaha Waihi Beach Whakatane

Lakes DHB

Mangakino Turangi

Tairawhiti DHB

Ruatoria Te Araroa Te Karaka Te Puia Springs Tikitiki Tokomaru Bay Tolaga Bay

Taranaki DHB

Eltham Inglewood Manaia Oakura Okato Opunake Patea Stratford Waverley

Hawkes Bay DHB

Waipawa Waipukurau Wairoa Whanganui DHB Bulls

Marton Ohakune Raetihi Taihape Waiouru MidCentral DHB

MidCentral DHB

Dannevirke Foxton Levin Otaki Pahiatua Shannon Woodville

Wairarapa DHB

Carteron Featherston Greytown Martinborough

SOUTH ISLAND

Nelson/Marlborough DHB

Havelock Mapua Motueka Murchison Picton Takaka Wakefield

West Coast DHB

Dobson Greymouth Hokitika Karamea Reefton South Westland Westport Whataroa

Canterbury DHB

Akaroa Amberley Amuri Chatham Islands Cheviot Darfield

Diamond Harbour Hanmer Springs Kaikoura Leeston I incoln Methven Oxford Rakaia **Bolleston** Rotherham Templeton Waikari South Canterbury DHB Fairlie Geraldine Pleasant Point Temuka Twizel Waimate Southern DHB Alexandra Balclutha Cromwell Gore Kurow I awrence Lumsden Mataura Milton Oamaru Oban Otautau Outram Owaka Palmerston Queenstown Ranfurlv Riverton Roxburah Tapanui Te Anau Tokonui Tuatapere

Wanaka

Winton

SECTION F: PART I

A Community Pharmaceutical identified with a * within the other sections of the Pharmaceutical Schedule:

- a) is exempt from any requirement to dispense in Monthly Lots;
- b) will only be subsidised if it is dispensed in a 90 Day Lot unless it is under the Dispensing Frequency Rule.

A Community Pharmaceutical that is an oral contraceptive and that is identified with a * within the other sections of the Pharmaceutical Schedule:

- a) is exempt from any requirement to dispense in Monthly Lots;
- b) will only be subsidised if it is dispensed in a 180 Day Lot unless it is under the Dispensing Frequency Rule.

SECTION F: PART II: CERTIFIED EXEMPTIONS AND ACCESS EXEMPTIONS TO MONTHLY DISPENSING

A Community Pharmaceutical, other than a Community Pharmaceutical identified with a * within the other sections of the Pharmaceutical Schedule, may be dispensed in a 90 Day Lot if:

a) the Community Pharmaceutical is identified with a ▲ within the other sections of the Pharmaceutical Schedule and the prescriber/pharmacist has endorsed/annotated the Prescription item(s) on the Prescription to which the exemption applies "certified exemption".

In endorsing/annotating the Prescription items for a certified exemption, the prescriber/pharmacist is certifying that:

- i) the patient wished to have the medicine dispensed in a quantity greater than a Monthly Lot; and
- ii) the patient has been stabilised on the same medicine for a reasonable period of time; and
- iii) the prescriber/pharmacist has reason to believe the patient will continue on the medicine and is compliant.
- b) a patient, who has difficulty getting to and from a pharmacy, signs the back of the Prescription to qualify for an Access Exemption. In signing the Prescription, the patient or his or her nominated representative must also certify which of the following criteria they meet:
 - i) have limited physical mobility;
 - ii) live and work more than 30 minutes from the nearest pharmacy by their normal form of transport;
 - iii) are relocating to another area;
 - iv) are travelling extensively and will be out of town when the repeat prescriptions are due.

SECTION F: PART III: FLEXIBLE AND VARIABLE DISPENSING PERIODS FOR PHARMACY

A Community Pharmaceutical, other than a Community Pharmaceutical identified with a ***** within the other sections of the Pharmaceutical Schedule, may be dispensed in variable dispensing periods under the following conditions:

- a) for stock management where the original pack(s) result in dispensing greater than 30 days supply,
- b) to synchronise a patients medication where multiple medicines result in uneven supply periods, note if dispensing a medicine other than a Pharmaceutical identified with a * please refer to Section F; Part II
- Note the total quantity and dispensing period can not exceed the total quantity and period prescribed on the prescription.

COMMUNITY PHARMACEUTICALS DISPENSING PERIOD EXEMPTIONS

The following Community Pharmaceuticals are identified with a ▲ within the other sections of the Pharmaceutical Schedule and may be dispensed in a 90 Day Lot if endorsed as a certified exemption in accordance with paragraph (a) in Section F Part II above.

ALIMENTARY TRACT AND METABOLISM PROPAFENONE HYDROCHLORIDE INSULIN ASPART HORMONE PREPARATIONS - SYSTEMIC EXCLUDING INSULIN ASPART WITH INSULIN ASPART PROTAMINE CONTRACEPTIVE HORMONES DESMOPRESSIN ACETATE INSULIN GLARGINE Nasal drops 100 mcg Minirin per ml INSULIN GLULISINE Nasal spray 10 mcg Desmopressin-PH&T INSULIN ISOPHANE per dose INSULIN ISOPHANE WITH INSULIN NEUTRAL MUSCULOSKELETAL SYSTEM PYRIDOSTIGMINE BROMIDE INSULIN LISPRO NERVOUS SYSTEM INSULIN LISPRO WITH INSULIN LISPRO PROTAMINE AMANTADINE HYDROCHLORIDE INSULIN NEUTRAL APOMORPHINE HYDROCHLORIDE CARDIOVASCULAR SYSTEM ENTACAPONE AMIODARONE HYDROCHLORIDE Tab 100 mg Cordarone-X GABAPENTIN Tab 200 mg Cordarone-X I ACOSAMIDE DISOPYRAMIDE PHOSPHATE I AMOTRIGINE FI ECAINIDE ACETATE Tambocor PRAMIPEXOLE HYDROCHLORIDE Tab 50 mg Cap long-acting Tambocor CR **BOPINIBOLE HYDBOCHLOBIDE** 100 ma Cap long-acting Tambocor CR TOI CAPONE 200 ma TOPIRAMATE MEXILETINE HYDROCHLORIDE VIGABATRIN MINOXIDII

NICORANDIL

Pharmacists should endeavour to use safety caps when dispensing any of the medicines listed in Section G in an oral liquid formulation pursuant to a prescription or Practitioner's Supply Order. This includes all proprietary and extemporaneously compounded oral liquid preparations of those pharmaceuticals listed in Section G of the Pharmaceutical Schedule. These medicines will be identified throughout Section B of the Pharmaceutical Schedule with the symbol '‡'.

Exemptions

Oral liquid preparations of the pharmaceuticals listed in Section G of the Pharmaceutical Schedule will be dispensed in a container with a safety cap unless:

- the practitioner has endorsed the Prescription or Practitioner's Supply Order, stating that, the Pharmaceutical is not to be dispensed in a container with a safety cap; or
- the Contractor has annotated the Prescription or Practitioner's Supply Order stating that, because of infirmity of the
 particular person, the Pharmaceutical to be used by that person should not be dispensed in a container with a safety cap; or
- the Pharmaceutical is packaged in an Original Pack so designed that on the professional judgement of the Contractor, transfer to a container with a safety cap would be inadvisable or a retrograde procedure.

Reimbursement

Pharmacists will be reimbursed according to their agreement. Where an additional fee is paid on safety caps it will be paid on all dispensings of oral liquid preparations for those pharmaceuticals listed in Section G of the Pharmaceutical Schedule unless the practitioner has endorsed or the contractor has annotated the Prescription or Practitioner's Supply Order that a safety cap has not been supplied.

Safety Caps (NZS 5825:1991)

20 mm Clic-Loc, United Closures & Plastics PLC, England	
Kerr, Cormack Packaging, Sydney, under licence to Kerr US	А
24 mm Clic-Loc, United Closures & Plastics PLC, England	
Clic-Loc, ACI Closures under license to Owens-Illinois	
Kerr, Cormack Packaging, Sydney, under licence to Kerr US	A
28 mm Clic-Loc, United Closures & Plastics PLC, England	
Clic-Loc, ACI Closures under license to Owens-Illinois	
Kerr, Cormack Packaging, Sydney, under licence to Kerr US	А
PDL Squeezlok	
PDL FG	

SAFETY CAP MEDICI	NES		
ALIMENTARY TRACT AND META	ABOLISM	CLOBAZAM	
FERROUS SULPHATE		Tab 10 mg	Frisium
Oral liq 30 mg (6 mg elemental) per 1 ml	Ferodan	(Extemporaneously compounded of	oral liquid preparations)
, p		CLONAZEPAM	
		Oral drops 2.5 mg per ml	Rivotril
CARDIOVASCULAR SYSTEM			
AMILORIDE HYDROCHLORID	E	DIAZEPAM	
Oral liq 1 mg per ml	Biomed	Tab 2 mg	Arrow-Diazepam
		Tab 5 mg	Arrow-Diazepam
CAPTOPRIL		(Extemporaneously compounded of	oral liquid preparations)
Oral liq 5 mg per ml	Capoten		
		ETHOSUXIMIDE	
CHLOROTHIAZIDE		Oral liq 250 mg per 5 ml	Zarontin
Oral liq 50 mg per ml	Biomed		
		LEVETIRACETAM	
DIGOXIN		Oral liq 100 mg per ml	Levetiracetam-AFT
Oral liq 50 mcg per ml	Lanoxin		
	Lanoxin S29	LORAZEPAM	•
		Tab 1 mg	Ativan
FUROSEMIDE [FRUSEMIDE]	Leven	Tab 2.5 mg	Ativan
Oral liq 10 mg per ml	Lasix	(Extemporaneously compounded of	orai liquid preparations)
SPIRONOLACTONE		LORMETAZEPAM	
Oral liq 5 mg per ml	Biomed	Tab 1 mg	Noctamid
Oraning 5 mg per mi	Diomed	(Extemporaneously compounded of	
			ina inquita proparationoj
HORMONE PREPARATIONS - SY	STEMIC EXCLUDING	METHADONE HYDROCHLORI	DE
CONTRACEPTIVE HORMONES		Oral lig 2 mg per ml	Biodone
LEVOTHYROXINE		Oral lig 5 mg per ml	Biodone Forte
Tab 25 mcg	Synthroid	Oral liq 10 mg per ml	Biodone Extra Forte
Tab 50 mcg	Eltroxin		
-	Mercury Pharma	MORPHINE HYDROCHLORIDI	=
	Synthroid	Oral liq 1 mg per ml	RA-Morph
Tab 100 mcg	Eltroxin	Oral liq 2 mg per ml	RA-Morph
	Mercury Pharma	Oral liq 5 mg per ml	RA-Morph
	Synthroid	Oral liq 10 mg per ml	RA-Morph
(Extemporaneously compounded of	oral liquid preparations)		
		NITRAZEPAM	
		Tab 5 mg	Nitrados
INFECTIONS - AGENTS FOR SYS QUININE SULPHATE	STEMIC USE	(Extemporaneously compounded oral liquid prepa	
Tab 300 mg	Q 300	OXAZEPAM	
(Extemporaneously compounded of		Tab 10 mg	Ox-Pam
(Tab 15 mg	Ox-Pam
		(Extemporaneously compounded of	oral liquid preparations)
MUSCULOSKELETAL SYSTEM			
IBUPROFEN		OXYCODONE HYDROCHLOR	IDE
Oral liq 20 mg per ml	Fenpaed	Oral liq 5 mg per 5 ml	OxyNorm
		PARACETAMOL	D

NERVOUS SYSTEM

CARBAMAZEPINE Oral liq 20 mg per ml

Tegretol

Oral liq 120 mg per 5 ml Paracare Oral liq 250 mg per 5 ml Paracare Double Strength

SAFETY CAP MEDICINES

Ventolin

PHENYTOIN SODIUM Oral liq 30 mg per 5 ml Dilantin

Oral lig 200 mg per 5 ml

SODIUM VALPROATE

Epilim S/F Liquid Epilim Syrup

TEMAZEPAM Tab 10 mg Normison (Extemporaneously compounded oral liquid preparations)

TRIAZOLAM Tab 125 mcg Hypam Tab 250 mcg Hypam (Extemporaneously compounded oral liquid preparations)

RESPIRATORY SYSTEM AND ALLERGIES

CETIRIZINE HYDROCHLORIDE Oral liq 1 mg per ml Histaclear

CHLORPHENIRAMINE MALEATE Oral liq 2 mg per 5 ml Histafen

DEXTROCHLORPHENIRAMINE MALEATE Oral liq 2 mg per 5 ml Polaramine PROMETHAZINE HYDROCHLORIDE

Oral liq 1 mg per 1 ml Allersoothe

SALBUTAMOL

Oral liq 400 mcg per ml

THEOPHYLLINE Oral liq 80 mg per 15 ml Nuelin

TRIMEPRAZINE TARTRATE Oral liq 30 mg per 5 ml Vallergan Forte

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

CODEINE PHOSPHATE Powder Douglas (Extemporaneously compounded oral liquid preparations)

METHADONE HYDROCHLORIDE Powder AFT (Extemporaneously compounded oral liquid preparations)

PHENOBARBITONE SODIUM Powder MidWest (Extemporaneously compounded oral liquid preparations)

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		ubsidised	Generic
	\$	Per	1	Manufacturer
Vaccinations				
ADULT DIPHTHERIA AND TETANUS VACCINE – [Xpharm] Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid in 0.5 ml Any of the following: 1) For vaccination of patients aged 45 and 65 years o		5	✓ <u>A</u>	DT Booster
 For vaccination of previously unimmunised or partial For revaccination following immunosuppression; or For boosting of patients with tetanus-prone wounds For use in testing for primary immunodeficiency disor paediatrician. 	s; or	,	ion of an ii	nternal medicine physician
 Note: Please refer to the Immunisation Handbook for ap BACILLUS CALMETTE-GUERIN VACCINE – [Xpharm] For infants at increased risk of tuberculosis. Increased risk is 1) living in a house or family with a person with current or 2) having one or more household members or carers who equal to 40 per 100,000 for 6 months or longer; or 3) during their first 5 years will be living 3 months or longer 	s defined as: past history of TB; or within the last 5 years	s lived	in a countr	y with a rate of TB > or
Note a list of countries with high rates of TB are available at www.bcgatlas.org/index.php. Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin), Danish strain 1331, live attenuated, vial with diluent	-	bercul		ch for downloads) or
 DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE – [Xphar Funded for any of the following criteria: A single vaccine for pregnant woman between gestatio A course of up to four vaccines is funded for children freprimary immunisation; or An additional four doses (as appropriate) are funded for transplantation or chemotherapy; pre or post splenector severely immunosuppressive regimens. 	m] nal weeks 28 and 38; om age 7 up to the ag r (re-)immunisation foi	or e of 18 patier	years inc	clusive to complete full
 Notes: Tdap is not registered for patients aged less than 10 appropriate schedule for catch up programmes. Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis toxoid, 8 mcg pertussis filamentous haemagluttinin and 2.5 mcg pertactin in 0.5 ml syringe 		the Ir 10 1	✓ <u>B</u>	on Handbook for oostrix oostrix

	Subsidy (Manufacturer's Price) \$	F Subsidis Per	ully sed	Brand or Generic Manufacturer
PIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE	E – [Xpharm]			
Funded for any of the following:				
 A single dose for children up to the age of 7 who have A course of four vaccines is funded for catch up prog primary immunisation; or 				urs) to complete full
 An additional four doses (as appropriate) are funded pre- or post splenectomy; pre- or post solid organ transplant 	()			· · · · · · · · · · · · · · · · · · ·
regimens; or 4) Five doses will be funded for children requiring solid of	organ transplantation.			
Note: Please refer to the Immunisation Handbook for appr	•	tch up progra	amme	es.
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mc	g			
pertussis toxoid, 25 mcg pertussis filamentous haemagluttinin, 8 mcg pertactin and 80 D-antigen units	•			
poliomyelitis virus in 0.5ml syringe		10	🖌 In	fanrix IPV
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B				
Kpharm] Funded for patients meeting any of the following criteria:				
 Up to four doses for children up to and under the age 	of 10 for primary immu	inication: or		
2) An additional four doses (as appropriate) are funded			to ar	nd under the age of
10 who are patients post haematopoietic stem cell tra	()			0
post solid organ transplant, renal dialysis and other s		1.2.1		
3) Up to five doses for children up to and under the age	of 10 receiving solid or	gan transpla	ntatio	n.
Note: A course of up-to four vaccines is funded for catch u to complete full primary immunisation. Please refer to the		· ·		• •
programmes. Inj 30IU diphtheriatoxoid with 40IU tetanustoxoid, 25mcg				
pertussistoxoid, 25mcg pertussisfilamentoushaemagluttinin, 8 mcgpertactin,				
80 D-AgUpoliovirus, 10mcghepatitisBsurfaceantigen ir	n			
0.5ml syringe		10	🗸 In	fanrix-hexa
AEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm]				
One dose for patients meeting any of the following:				
1) For primary vaccination in children; or				
2) An additional dose (as appropriate) is funded for (re-)	immunisation for patier	nts post haen	natop	oietic stem cell
transplantation, or chemotherapy; functional asplenic	21 I I	271		lid organ transplant, p
or post cochlear implants, renal dialysis and other ser		•		
 For use in testing for primary immunodeficiency disea paediatrician. 	ases, on the recommen	dation of an	Intern	al medicine physician
Haemophilus Influenzae type B polysaccharide 10 mcg				
Haemophilus Influenzae type B polysaccharide 10 mcg conjugated to tetanus toxoid as carrier protein 20-40 m prefilled syringe plus vial 0.5 ml	0.	1		iberix

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

(Subsidy Manufacturer's Price) \$	F Subsid Per	Fully ised	Brand or Generic Manufacturer
HEPATITIS A VACCINE – [Xpharm]				
Funded for patients meeting any of the following criteria:				
1) Two vaccinations for use in transplant patients; or				
2) Two vaccinations for use in children with chronic liver dis	ease; or			
3) One dose of vaccine for close contacts of known hepatitie	s A cases.			
Inj 1440 ELISA units in 1 ml syringe	0.00	1	✓ <u>н</u>	avrix
Inj 720 ELISA units in 0.5 ml syringe	0.00	1	✓ Н	avrix Junior

				Manufacturer
PATITIS B RECOMBINANT VACCINE – [Xpharm]				
Inj 5 mcg per 0.5 ml vial	0.00	1	🗸 Н	BvaxPRO
Funded for patients meeting any of the following criteri	a:		_	
1) for household or sexual contacts of known acute	hepatitis B patients or h	nepatit	tis B carrier	s; or
2) for children born to mothers who are hepatitis B s	surface antigen (HBsAg	, posit	tive; or	
3) for children up to and under the age of 18 years i	inclusive who are consid	dered	not to have	achieved a positive
serology and require additional vaccination or rec	quire a primary course c	of vaco	cination; or	
for HIV positive patients; or				
5) for hepatitis C positive patients; or				
for patients following non-consensual sexual inter	rcourse; or			
for patients following immunosuppression; or				
for solid organ transplant patients; or				
for post-haematopoietic stem cell transplant (HSC)	CT) patients; or			
10) following needle stick injury.				
Inj 10 mcg per 1 ml vial		1	🗸 Н	BvaxPRO
Funded for patients meeting any of the following criteri				
1) for household or sexual contacts of known acute				s; or
2) for children born to mothers who are hepatitis B s				
3) for children up to and under the age of 18 years i				achieved a positive
serology and require additional vaccination or rec	quire a primary course o	of vaco	cination; or	
 for HIV positive patients; or for headtiling C positive patients; or 				
5) for hepatitis C positive patients; or	100U1001 01			
 for patients following non-consensual sexual inter for patients following immunocurpression: or 	rcourse; or			
7) for patients following immunosuppression; or8) for solid organ transplant patients; or				
 9) for post-haematopoietic stem cell transplant (HSC) 	CT) nationts: or			
10) following needle stick injury.	or) patiente, er			
Ini 20 mag new 1 ml profilled ovringe	0.00	4	./ 5	naniv B
Inj 20 mcg per 1 ml prefilled syringe Funded for patients meeting any of the following criteri		1	• -	ngerix-B
, , , ,		on otil		
 for household or sexual contacts of known acute for abildram harm to mathema who are heratilia B a 				s; or
 for children born to mothers who are hepatitis B s for children up to and under the age of 18 years i 				achieved a positive
serology and require additional vaccination or rec				achieveu a positive
4) for HIV positive patients; or	quile a plillary course c	n vacu		
5) for hepatitis C positive patients; or				
6) for patients following non-consensual sexual inter	rcourse: or			
7) for patients following immunosuppression; or	1000100, 01			
8) for solid organ transplant patients; or				
9) for post-haematopoietic stem cell transplant (HSC	CT) patients: or			
10) following needle stick injury.	e) paileine, ei			
Ini 40 meg ner 1 milviol	0.00	4	1 4	Puer PPO
Inj 40 mcg per 1 ml vial	0.00	1	• [BvaxPRO
Funded for any of the following criteria:				
 for dialysis patients; or for liver or kidney transplant patient 				
 for liver or kidney transplant patient. 				

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 54 Any of the following:	8) VACCINE [HPV] -	[Xpharm]	
 Maximum of two doses for children aged 14 years and the second sec			
 People aged 15 to 26 years inclusive; or Either: 			
People aged 9 to 26 years inclusive 1) Confirmed HIV infection; or			
 Transplant (including stem cell) patients: or Maximum of four doses for people aged 9 to 26 years in 	nclusive nost chemoth	erany	
Inj 270 mcg in 0.5 ml syringe	·		Gardasil 9

	Subsidy (Manufacturer's Price) \$	Subsic Per	Fully dised	Brand or Generic Manufacturer
INFLUENZA VACCINE				
Inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine) [Xpharm]		1	🖌 Fli	uarix Tetra
Inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine)		or or ot invasive; ion, or ignificant re n (within th bury District ounding reg	or or espirato ne Nelso t Health gion;	ry illness; on Marlborough District Board);
 a) asthma not requiring regular preventative ther b) hypertension and/or dyslipidaemia without evi B) Doctors are the only Contractors entitled to claim particular the only Contractors and the only contractor of t	apy, dence of end-organ o	disease.		
60 mcg in 0.5 ml syringe (paediatric quadrivalent va immunisation and they may only do so in respect of				

✓ Influvac Tetra 10

‡ safety cap

Subsidy	9	Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

- a) Only on a prescription
- b) No patient co-payment payable
- C)

A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by PHARMAC:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or
- c) children aged four years or less (but over three years) who have been hospitalised for respiratory illness or have a history of significant respiratory illness;
- d) people under 18 years of age living in the Seddon/Ward and rural Eastern Marlborough region (within the Nelson Marlborough District Health Board) and Kaikoura and Hurunui areas (within the Canterbury District Health Board);
- People under 18 years of age who have been displaced from their homes in Edgecumbe and the surrounding region;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

	Subsidy	0	Fully	Brand or
	(Manufacturer's Price) \$	Per	ubsidised ✓	Generic Manufacturer
MEASLES, MUMPS AND RUBELLA VACCINE – [Xpharm]	•	-		
A maximum of two doses for any patient meeting the followin	a criteria:			
1) For primary vaccination in children; or	g omona.			
2) For revaccination following immunosuppression; or				
 For any individual susceptible to measles, mumps or ru 	bella: or			
4) A maximum of three doses for children who have had the		12 mor	nths.	
Note: Please refer to the Immunisation Handbook for approp	priate schedule for cat	tch up p	programme	es.
Injection, measles virus 1,000 CCID50, mumps virus				
5,012 CCID50, Rubella virus 1,000 CCID50; prefilled				
syringe/ampoule of diluent 0.5 ml	0.00	10	✓ Pi	riorix
MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGAT Any of the following:	E VACCINE – [Xpha	arm]		
1) Up to three doses and a booster every five years for pa	tients pre- and post s	plenect	omy and f	or patients with functional
or anatomic asplenia, HIV, complement deficiency (acq		pre or	post solid (organ transplant; or
One dose for close contacts of meningococcal cases; o				
3) A maximum of two doses for bone marrow transplant pa				
A maximum of two doses for patients following immuno	suppression*.			
Note: children under coven vegra of ago require two doese C	waaka anart a haaa	tor doo	a three yes	are ofter the primers
Note: children under seven years of age require two doses & series and then five yearly.	weeks apart, a boos		e unee yea	als aller the phillary
*Immunosuppression due to steroid or other immunosuppres	sive therapy must be	for a pe	eriod of are	eater than 28 days
Inj 4 mcg of each meningococcal polysaccharide conjugated			strou et git	
a total of approximately 48 mcg of diphtheria toxoid carrie				
per 0.5 ml vial		1	🖌 <u>М</u>	enactra
MENINGOCOCCAL C CONJUGATE VACCINE – [Xpharm]				
Any of the following:				
1) Up to three doses and a booster every five years for pa	tients pre- and post s	plenect	omy and f	or patients with functional
or anatomic asplenia, HIV, complement deficiency (acq	uired or inherited), or	pre or	post solid (organ transplant; or
One dose for close contacts of meningococcal cases; o	r			
A maximum of two doses for bone marrow transplant particular				
 A maximum of two doses for patients following immuno 	suppression*.			
				6 H I
Note: children under seven years of age require two doses 8	weeks apart, a boos	ter dos	e three yea	ars after the primary
series and then five yearly. *Immunosuppression due to steroid or other immunosuppres	aive thereasy must be	for a p	oriod of ar	ator than 20 days
Inj 10 mcg in 0.5 ml syringe		1		eisvac-C
PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE – [Xpharm			• 11	
Either:	ıJ			
 A primary course of four doses for previously unvaccina 	ted individuals up to	the arc	of 50 mor	othe inclusive: or
2) Up to three doses as appropriate to complete the prima				
59 months who have received one to three doses of PC			i individud	lo undor the ugo of
Note: please refer to the Immunisation Handbook for the app	propriate schedule for	catch	up progran	nmes
Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6E	3,			
7F, 9V, 14 and 23F; 3 mcg of pneumococcal				
polysaccharide serotypes 4, 18C and 19F in 0.5 ml			-	
prefilled syringe	0.00	10	✓ <u>S</u>	<u>ynflorix</u>

‡ safety cap

*Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)	Si	ubsidised	Generic
\$	Per	1	Manufacturer

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- 1) One dose is funded for high risk children (over the age of 17 months and under 18 years) who have previously received four doses of PCV10; or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) with primary immune deficiencies; or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cyanosis or failure; or
 - I) with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes ini 30.8 mcg of pneumococcal polysaccharide services 1, 3, 4

30.6 mcg of pheumococcal polysacchande serotypes 1, 3, 4,	
5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml	
syringe0.00	10
	1

Prevenar 13
 Prevenar 13

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	

PNEUMOCOCCAL	(PPV23) POLYSACCHARIDE VACCINE – [Xpharm]		
Either:	. ,		
chemothe	e doses (as appropriate) for patients with HIV, for patients p rapy; pre- or post-splenectomy or with functional asplenia, p	ore- or post-solid c	organ transplant, renal dialysis,
	ent deficiency (acquired or inherited), cochlear implants, or p	rimary immunode	efficiency; or
2) All of the f	5		
	ent is a child under 18 years for (re-)immunisation; and then the standard strength the standard strength the		
	of the following:		
, ,	on immunosuppressive therapy or radiation therapy, vaccir	nata when there is	s expected to be a sufficient
9	immune response; or		
ii)	with primary immune deficiencies; or		
,	with HIV infection; or		
iv)	with renal failure, or nephrotic syndrome; or		
v)	who are immune-suppressed following organ transplantation	on (including haer	matopoietic stem cell transplant)
vi)	with cochlear implants or intracranial shunts; or		
	with cerebrospinal fluid leaks; or		
viii)	receiving corticosteroid therapy for more than two weeks, a		
	prednisone of 2 mg/kg per day or greater, or children who	weigh more than	10 kg on a total daily dosage of
	20 mg or greater; or		
	with chronic pulmonary disease (including asthma treated	with high-dose co	rticosteroid therapy); or
	pre term infants, born before 28 weeks gestation; or with cardiac disease, with cyanosis or failure; or		
	with diabetes; or		
,	with Down syndrome; or		
	who are pre-or post-splenectomy, or with functional asplen	nia.	
,	F F		
Inj 575 mcg in 0).5 ml prefilled syringe (25 mcg of each		
23 pneumo	coccal serotype)0.00	1	Pneumovax 23
POLIOMYELITIS VA	ACCINE – [Xpharm]		
Up to three dos	es for patients meeting either of the following:		
	lly vaccinated or previously unvaccinated individuals; or		
	cination following immunosuppression.		
	efer to the Immunisation Handbook for appropriate schedule	e for catch-up proc	grammes.
Inj 80D antigen	units in 0.5 ml syringe0.00	1	✓ <u>IPOL</u>
ROTAVIRUS ORAL	VACCINE – [Xpharm]		
Maximum of two	o doses for patients meeting the following:		
 first dose t 	to be administered in infants aged under 14 weeks of age; a	and	
no vaccina	ation being administered to children aged 24 weeks or over.		
	ttenuated human rotavirus	1.0	4 -
1,000,000 (CCID50 per dose, prefilled oral applicator0.00	10	✓ <u>Rotarix</u>

Subsidy	ç	Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

VARICELLA VACCINE [CHICKENPOX VACCINE] - [Xpharm]

Either:

- 1) Maximum of one dose for primary vaccination for either:
 - a) Any infant born on or after 1 April 2016; or
 - b) For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox), or
- 2) Maximum of two doses for any of the following:
 - a) Any of the following for non-immune patients:
 - i) with chronic liver disease who may in future be candidates for transplantation; or
 - ii) with deteriorating renal function before transplantation; or
 - iii) prior to solid organ transplant; or
 - iv) prior to any elective immunosuppression*, or
 - v) for post exposure prophylaxis who are immune competent inpatients.; or
 - b) For patients at least 2 years after bone marrow transplantation, on advice of their specialist, or
 - c) For patients at least 6 months after completion of chemotherapy, on advice of their specialist, or
 - d) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist, or
 - For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella, or
 - f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella, or
 - g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.

* immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

Inj 2000 PFU prefilled syringe plus vial	0.00	1	 Varilrix
		10	✓ Varilrix

VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED VACCINE [SHINGLES VACCINE] – [Xpharm] Funded for patients meeting either of the following criteria:

- 1) One dose for all people aged 65 years; or
- 2) One dose for all people aged between 66 and 80 years inclusive from 1 April 2018 and 31 March 2020.

Inj 19,400 PFU prefilled syringe plus vial0.00	1 10	✓ Zostavax✓ Zostavax	
Diagnostic Agents			
UBERCULIN PPD [MANTOUX] TEST – [Xpharm] Ini 5 TU per 0.1 ml, 1 ml vial	1	✓ Tubersol	

- Symbols -

-	Symbols -	
3TC	•	117
50X 3.0 Reserv	/oir	
	- A -	
A-Scabies		78
Abacavir sulpha	ate	116
Abacavir sulpha	ate with	
lamivudine		116
Abilify		144
	etate	
Accu-Chek Per	forma	
Accuretic 10		59
Acetazolamide		218
	1, 2- propanediol	
diacetate an		
henzethoniu	m	216
Acotic acid with	hydroxyquinoline	and
ricipoleic aci	d	
Acotylovetoino	u	00 001
Aci-Jer		00
		110
		112
Sensory		210
Acidex		20
Acipimox		64
Acitretin		
Aclasta		126
Aclin		121
Actinomycin D.		1/2
Actrapid		24
Actrapid Penfill		24
Acupan		132
Adalat Oros		62
Adalimumab		190
Adapalene		71
Adefin		62
Adefovir dipivo	xil	111
Adenuric		128
ADR Cartridge	1.8	38
Adrenaline		66
ADT Booster		260
Adult diphtheria	a and tetanus	
Advate		<mark>5</mark> 1
Afinitor		206
Aflibercept		197
AFT Carbimazo	ole	<mark>93</mark>
AFT-Pyrazinam	nide	110
AFT-Pyrazinam	nide S29	110

Agents Affecting the	
Renin-Angiotensin System	58
Agents for Parkinsonism and Related	
Disorders	130
Agents Used in the Treatment of	
Poisonings	221
Agrylin	171
Alanase	214
Albendazole	99
Albey	208
Albustix	87
Aldurazyme	43
Alendronate sodium	124
Alendronate sodium with	
colecalciferol	124
Alfacalcidol	45
Alfamino Junior	247
Alginic acid	20
Alglucosidase alfa	41
Alkeran	166
Allersoothe	209
Allmercap	
Allopurinol	127
Alpha Adrenoceptor Blockers	
Alpha-Keri Lotion	76
Alphamox 125	102
Alphamox 250	
Alu-Tab	
Aluminium hydroxide	20
Amantadine hydrochloride	
Ambrisentan	67
Amiloride hydrochloride	63
Amiloride hydrochloride with	
furosemide	64
Amiloride hydrochloride with	
hydrochlorothiazide	
Aminophylline Amiodarone hydrochloride	213
Amisulpride	
Amitriptyline	
Amlodipine	
Amorolfine	
Amoxicillin	
Amoxicillin with clavulanic acid	
Amphotericin B	
Amsacrine	171
AmsaLyo	
Amsidine	
Amyl nitrite	
Amzoate	
Anaesthetics	
Anagrelide hydrochloride	171
Analgesics	132
Anastrozole	

Andriol Testocaps	90
Androderm	90
Animas Battery Cap	34
Animas Cartridge	38
Animas Vibe	29
Anoro Ellipta	.213
Antabuse	
Antacids and Antiflatulants	20
Anten	
Anthelmintics	
Antiacne Preparations	71
Antiallergy Preparations	
Antianaemics	49
Antiandrogen Oral	
Contraceptives	84
Antiarrhythmics	
Antibacterials	00
Antibacterials Topical	33
Anticholinergic Agents	211
Anticholinesterases	
Antidepressants	
Antidiarrhoeals	
Antiepilepsy Drugs	107
Antifibrinolytics, Haemostatics and	. 137
Local Sclerosants	50
Antifibrotics	
Antifungals Antifungals Topical	. 100
Antihistamines	
Antihypotensives	
Antimalarials	.109
Antimigraine Preparations	. 142
Antinausea and Vertigo Agents	
Antiparasitics	.109
Antipruritic Preparations	
Antipsychotics	
Antiretrovirals	
Antirheumatoid Agents	. 122
Antispasmodics and Other Agents	
Altering Gut Motility	22
Antithrombotic Agents	52
Antithymocyte globulin	
(equine)	. 190
Antitrichomonal Agents	. 109
Antituberculotics and	
Antileprotics	. 109
Antiulcerants	
Antivirals	
Anxiolytics	
Anzatax	
Apidra	24
Apo-Amiloride	
Apo-Amlodipine	62

Apo-Amoxi
Apo-Amori 102
Apo-Azithromycin 100
Apo-Bromocriptine130
Apo-Ciclopirox72
Apo-Cilazapril
Apo-Cilazapril/
Apo-Cilazapin/
Hydrochlorothiazide59
Apo-Clarithromycin
Alimentary22
Infection 100
Apo-Clomipramine135
Apo Diolo CD 101
Apo-Diclo SR
Apo-Diltiazem CD62
Apo-Doxazosin58
Apo-Folic Acid
Apo-Gabapentin
Apo-Imiquimod Cream 5%80
Apo-Leflunomide 122
Apo-Megestrol 182
Apo-Metoprolol61
Apo-Mirtazapine137
Apo-Moclobemide136
Apo-Montelukast
Apo-Nadolol61
Apo-Nicotinic Acid64
Apo-Ondansetron 143
Apo-Oxybutynin86
Apo-Paroxetine
Apo-Perindopril
Apo-Pindolol
Apo-Pravastatin65
Apo-Prazosin58
Apo-Prednisone90
Apo-Primidone140
Apo-Propranolol
Apo-Propranolol S2961
Apo-Propranoioi 529
Apo-Pyridoxine
Apo-Ropinirole130
Apo-Selegiline S29130
Apo-Sumatriptan142
Apo-Terazosin58
Apo-Thiamine
Apo-Timol
Apomorphine hydrochloride130
Aprepitant 142
Apresoline67
Aptamil Gold+ Pepti Junior248
Aqueous cream
Aripiprazole
Aripiprazole Sandoz144
Aristocort74
Arrow - Clopid
Arrow-Amitriptyline 135
Arrow-Bendrofluazide
Arrow-Brimonidine
Arrow-Calcium46

Arrow-Diazepam148
Arrow-Dortim219
Arrow-Doxorubicin 172
Arrow-Etidronate124
Arrow-Fluoxetine136
Arrow-Gabapentin 138
Arrow-Lamotrigine 139
Arrow-Losartan &
Hydrochlorothiazide59
Arrow-Meloxicam121
Arrow-Morphine LA134
Arrow-Norfloxacin 120
Arrow-Ornidazole109
Arrow-Quinapril 1058
Arrow-Quinapril 2058
Arrow-Quinapril 5
Arrow-Roxithromycin101
Arrow-Sertraline137
Arrow-Timolol
Arrow-Tolterodine
Arrow-Topiramate141
Arrow-Tramadol135
Arsenic trioxide 171
Asacol21
Asamax21
Ascorbic acid 45
Aspen Adrenaline66
Aspirin
Blood52
Nervous132
Asthalin211
Atazanavir sulphate 117
Atenolol60
Atenolol AFT60
ATGAM190
Ativan148
Atomoxetine157
Atorvastatin65
Atripla 116
Atropine sulphate
Cardiovascular
Sensory219
Atropt219
Atrovent
Aubagio 153
Augmentin 102
Avelox104
Avomine143
Avonex
Avonex Pen156
Azacitidine
Azathioprine184
Azithromycin 100
Azol
Azopt
AZT117

- B -	
B-D Micro-Fine	29
B-D Ultra Fine	29
B-D Ultra Fine II	
Bacillus Calmette-Guerin (BCG)	
vaccine 1	90
Bacillus Calmette-Guerin	00
vaccine 2	020
Baclofen1	.00 20
Bactroban	72
Baraclude1	11
Barrier Creams and Emollients	75
BCG Vaccine	
Beclazone 100	
Beclazone 250	
Beclazone 502	
Beclomethasone	.09
dipropionate 209, 2	
Bee venom allergy treatment	
Bendamustine hydrochloride	.00
Bendrofluazide	00
Bendroflumethiazide	04
[Bendrofluazide]	61
BeneFIX	
Benzathine benzylpenicillin	01
Benzatropine mesylate 1	202
Benzbromaron AL 1001	20
Benzbromarone 1	
Benzoin	
Benztrop 1	
Benzydamine hydrochloride	11
Benzylpenicillin sodium [Penicillin	
G] 1	02
Beta Adrenoceptor Blockers	
Beta Cream	
Beta Ointment	74
Beta Scalp	
Beta-Adrenoceptor Agonists	210
Betadine	77
Betadine Skin Prep	77
Betaferon 1	56
Betagan2	18
Betahistine dihydrochloride1	
Betaloc CR	61
Betamethasone dipropionate	73
Betamethasone dipropionate with	
calcipotriol	79
Betamethasone sodium phosphate	
with betamethasone acetate	89
Betamethasone valerate74,	
Betamethasone valerate with	
clioquinol	74
Betamethasone valerate with sodium	
fusidate [fusidic acid]	75
Betaxolol2	18
Betnovate	74

Betnovate-C74
Betoptic218
Betoptic S218
Bezafibrate64
Bezalip64
Bezalip Retard 64
Bicalutamide
Bicillin LA 102
BiCNU166
Bile and Liver Therapy23
Biltricide
Bimatoprost
Bimatoprost Actavis
Binarex
Biodone
Biodone Extra Forte
Biodone Forte
Bisacodyl
Disacouyi
Bisoprolol fumarate
BK Lotion
Bleomycin sulphate
Blood Colony-stimulating
Factors
Blood glucose diagnostic test
meter
Blood glucose diagnostic test
strip 28
Blood glucose test strips (visually
impaired)28
Blood ketone diagnostic test
meter 25
Blood Ketone Diagnostic Test
Strip 25
Bonjela44
Bonvit40
Boostrix
Bortezomib171
Bosentan67
Bosentan-Mylan67
Bosvate60
Bplex
Breo Ellipta
Brevinor 1/21 83
Brevinor 1/21
Brevinor 1/21
Brevinor 1/21 83 Brevinor 1/28 83 Brevinor 21 83
Brevinor 1/21 83 Brevinor 1/28 83 Brevinor 21 83 Bricanyl Turbuhaler 211
Brevinor 1/21 83 Brevinor 1/28 83 Brevinor 21 83 Bricanyl Turbuhaler 211 Brilinta 53
Brevinor 1/21 83 Brevinor 1/28 83 Brevinor 21 83 Bricanyl Turbuhaler 211 Brilinta 53 Brimonidine tartrate 219
Brevinor 1/21 83 Brevinor 1/28 83 Brevinor 21 83 Bricanyl Turbuhaler 211 Brilinta 53 Brimonidine tartrate 219 Brimonidine tartrate with timolol
Brevinor 1/21 83 Brevinor 1/28 83 Brevinor 21 83 Bricanyl Turbuhaler 211 Brilinta 53 Brimonidine tartrate 219 Brimonidine tarte 219
Brevinor 1/21 83 Brevinor 1/28 83 Brevinor 21 83 Bricanyl Turbuhaler 211 Brilinta 53 Brimonidine tartrate 219 Brimonidine tartrate with timolol maleate 219 Brinov
Brevinor 1/21 83 Brevinor 1/28 83 Brevinor 21 83 Bricanyl Turbuhaler 211 Brilinta 53 Brimonidine tartrate 219 Brinov 168 Brinzolamide 218
Brevinor 1/21 83 Brevinor 1/28 83 Brevinor 21 83 Bricanyl Turbuhaler 211 Brilinta 53 Brimonidine tartrate 219 Brimonidine tartrate with timolol maleate Prinov 168 Brinzolamide 218 Brolene 216
Brevinor 1/21 83 Brevinor 1/28 83 Brevinor 21 83 Bricanyl Turbuhaler 211 Brilinta 53 Brimonidine tartrate 219 Brimonidine tartrate with timolol maleate Prinzolamide 219 Brinzolamide 219 Brinzolamide 219 Brolene 216 Bromocriptine mesylate 130
Brevinor 1/21 83 Brevinor 1/28 83 Brevinor 21 83 Bricanyl Turbuhaler 211 Brilinta 53 Brimonidine tartrate 219 Brimonidine tartrate with timolol maleate Prinov 168 Brinzolamide 218 Brolene 216

BSF CareSens N221
BSF CareSens N POP221
BSF CareSens N Premier221
BSF Rolin
Buccastem143
Budesonide
Alimentary
Respiratory209, 214
Budesonide with eformoterol
Bumetanide63
Buprenorphine with naloxone 161
Bupropion hydrochloride162
Burinex63
Buscopan22
Buspirone hydrochloride148
Busulfan
Butacort Aqueous
- C -
Cabergoline
Cafergot
Cafergot S29 142
Calefyor 529
Caffeine citrate
Calamine
Calcipotriol
Calcitonin
Calcitriol45
Calcitriol-AFT45
Calcium carbonate20, 46
Calcium Channel Blockers62
Calcium Disodium Versenate
Calcium folinate 168
Calcium Folinate Ebewe168
Calcium Folinate Sandoz168
Calcium gluconate46
Calcium Homeostasis
Calcium polystyrene sulphonate57
Calcium Resonium
Calogen
Calsource
Camptosar169
Candesartan cilexetil
Candestar
Canesten
Capecitabine
Capoten
Capsaicin
Musculoskeletal
Nervous
Captopril
Carafate
Carbaccord166
Carbamazepine
Carbimazole
Carbomer
Carboplatin
Carboplatin Ebewe 166

Carbosorb-X	1
Cardinol LA6	
CareSens	
CareSens Dual	6
CareSens II	7
CareSens N	, 0
CareSens N POP2	7
CareSens N Premier2	-
CareSens N Premier2	/
CareSens PRO2	8
Carmellose sodium with gelatin and	
pectin4	4
Carmustine16	
Carvedilol6	0
Carvedilol Sandoz6	0
Catapres6	3
Cathejell13	1
CeeNU16	
Cefaclor monohydrate9	9
Cefalexin9	
Cefalexin Sandoz	9
Cefazolin	
Ceftriaxone	
Cefuroxime axetil	ň
Celecoxib	
Celecoxib Pfizer	
Celestone Chronodose	1
Celestone Chronodose8	9
Celiprolol	
Cellcept18	
Celol6	
Centrally-Acting Agents 6	3
Cephalexin ABM9	9
Cerezyme4	3
Cetirizine hydrochloride 20	8
Cetomacrogol7	5
Cetomacrogol with glycerol7	6
Cetuximab 19	8
Champix16	
Charcoal	
Chemotherapeutic Agents 16	
Chickenpox vaccine	õ
Chlorafast	6
Chlorambucil	6
Chloramphenicol	
Chlorhexidine gluconate	0
Alimentary4	,
Dermatological	
Chloroform	
Chlorothiazide6	
Chlorpheniramine maleate20	8
Chlorpromazine hydrochloride14	4
Chlorsig21	6
Chlortalidone [Chlorthalidone]6	
Chlorthalidone6	
Chlorvescent5	7
Choice Load 3758	2
Choice TT380 Short8	2

Choice TT380 Standard82
Cholestyramine65
Choline salicylate with cetalkonium
chloride
Ciclopirox olamine72
Ciclosporin206
Cilazapril
Cilazapril with
hydrochlorothiazide
Cilicaine 103
Cilicaine VK 103
Ciloxan
Cinacalcet
Cipflox
Ciprofloxacin
Infection 104
Sensory
Ciprofloxacin Teva
Circadin
Cisplatin
Cisplatin Ebewe
Citalopram hydrobromide
Cladribine
Clarithromycin
Alimentary
Infection 100
Clexane
Clindamycin
Clindamycin ABM 104
Clinicians Renal Vit
Clobazam
Clobetasol propionate74, 80
Clobetasone butyrate74
Clofazimine109
Clomazol
Dermatological72
Genito-Urinary85
Clomifene citrate
Clomipramine hydrochloride135
Clonazepam 137, 148
Clonidine63
Clonidine BNM63
Clonidine hydrochloride63
Clopidogrel52
Clopine145
Clopixol146, 148
Clotrimazole
Dermatological72
Genito-Urinary85
Clozapine145
Clozaril145
Clustran142
Co-trimoxazole 106
Coal tar79
Coal tar with allantoin, menthol,
phenol and sulphur79

Coal tar with salicylic acid and	
sulphur7	' 9
Coco-Scalp7	' 9
Codeine phosphate	
Extemporaneous22	28
Nervous13	
Cogentin	
Colaspase [L-asparaginase]17	
Colchicine	
Colecalciferol	
Colestid6	
Colestipol hydrochloride6	5
Colgout	
Colifoam	21
Colistin sulphomethate	
Colistin-Link	
Collodion flexible	די, סמ
Colloidal bismuth subcitrate	.0 .0
Colofac	
Coloxyl	
Combigan	
Compound electrolytes5	3
Compound hydroxybenzoate	
Concerta	
Condoms	
Condyline	
Contact-D	55
Contraceptives - Hormonal	
Contraceptives - Non-hormonal	
Copaxone	56
Copaxone	56
Copaxone	56 59
Copaxone	56 59 39
Copaxone	56 59 39 73
Copaxone	56 59 39 73 72
Copaxone	56 59 39 73 72 55
Copaxone	56 59 39 73 72 55 39
Copaxone	56 59 73 72 55 39 39
Copaxone	56 59 73 72 55 59 73 73
Copaxone	56 59 73 72 55 59 73 73 71
Copaxone	56 59 73 72 55 59 73 72 59 59 73 71 20
Copaxone	56 59 72 55 59 73 72 59 59 73 71 22 55 73 71 22 55
Copaxone	56 59 57 55 59 59 59 59 59 59 59 59 59 59 59 59
Copaxone 15 Cordarone-X 5 Corticosteroids and Related Agents 6 for Systemic Use 8 Corticosteroids Topical 7 Cosmegen 17 Couradin 5 Creon 10000 6 Crotamiton 7 Crystaderm 7 Cystaderm 10 Cvite 4 Cyclizine hydrochloride 14 Cyclizine lactate 14	69 972599712522
Copaxone 15 Cordarone-X 5 Corticosteroids and Related Agents 6 for Systemic Use 6 Corticosteroids Topical 7 Cosmegen 17 Coumadin 5 Creon 10000 3 Crotamiton 7 Crystaderm 7 Curam 10 Cvite 4 Cyclizine hydrochloride 14 Cyclizine lactate 14 Cyclizyle 21	69 9325993125229 93125229 93125229 9312522 93125 93125 9312 93125 931
Copaxone 15 Cordarone-X 5 Corticosteroids and Related Agents 6 for Systemic Use 6 Corticosteroids Topical 7 Cosmegen 17 Coumadin 5 Creon 10000 3 Crotamiton 7 Crystaderm 7 Curam 10 Cvite 4 Cyclizine hydrochloride 14 Cyclizine lactate 14 Cyclogyl 21 Cyclopentolate hydrochloride 21	69 93259931252299
Copaxone 15 Cordarone-X 5 Corticosteroids and Related Agents 6 for Systemic Use 6 Corticosteroids Topical 7 Cosmegen 17 Coumadin 5 Creon 10000 3 Crotamiton 7 Crystaderm 7 Curam 10 Cvite 4 Cyclizine hydrochloride 14 Cyclizine lactate 14 Cyclogyl 21 Cyclopentolate hydrochloride 21 Cyclophosphamide 16	56 57 57 57 57 57 57 57 57 57 57
Copaxone 15 Cordarone-X 5 Corticosteroids and Related Agents 6 for Systemic Use 6 Corticosteroids Topical 7 Cosmegen 17 Coumadin 5 Creon 10000 3 Crotamiton 7 Cortamiton 7 Cyclagine hydrochloride 14 Cyclizine hydrochloride 14 Cyclogyl 21 Cyclophosphamide 16 Cycloserine 16	
Copaxone 15 Cordarone-X 5 Corticosteroids and Related Agents 6 for Systemic Use 6 Corticosteroids Topical 7 Cosmegen 17 Coumadin 5 Creon 10000 3 Crotamiton 7 Crystaderm 7 Curam 10 Cvite 4 Cyclizine hydrochloride 14 Cyclizine lactate 14 Cyclogyl 21 Cyclopentolate hydrochloride 21 Cyclophosphamide 16	
Copaxone 15 Cordarone-X 5 Corticosteroids and Related Agents 6 for Systemic Use 6 Corticosteroids Topical 7 Cosmegen 17 Coumadin 5 Creon 10000 3 Creon 25000 3 Crotamiton 7 Crystaderm 7 Curam 10 Cvite 4 Cyclizine hydrochloride 14 Cyclogyl 21 Cyclopentolate hydrochloride 16 Cycloserine 16	56 59 59 57 57 57 59 59 57 57 57 57 57 57 57 57 57 57 57 57 57
Copaxone 15 Cordarone-X 5 Corticosteroids and Related Agents 6 for Systemic Use 6 Corticosteroids Topical 7 Cosmegen 17 Coumadin 5 Creon 10000 6 Creon 25000 6 Crotamiton 7 Crystaderm 7 Curam 10 Cvite 4 Cyclizine hydrochloride 14 Cyclogyl 21 Cyclopentolate hydrochloride 16 Cyclopesphamide 16 Cycloserine 10 Cyklokapron 5 Cyproterone acetate 6 Cyproterone acetate 6	56 59 57 57 55 59 59 57 57 57 57 57 57 57 57 57 57 57 57 57
Copaxone 15 Cordarone-X 5 Corticosteroids and Related Agents 6 for Systemic Use 6 Corticosteroids Topical 7 Cosmegen 17 Coumadin 5 Creon 10000 6 Crotamiton 7 Crystaderm 7 Curam 10 Cvite 4 Cyclizine hydrochloride 14 Cyclogyl 21 Cyclopentolate hydrochloride 16 Cyclopohosphamide 16 Cycloserine 10 Cyklokapron 5 Cyptorerone acetate 6 Cyproterone acetate 6 Cyproterone acetate with 6 ethinyloestradiol 6	56 59 57 57 55 59 59 59 57 57 57 57 57 57 57 57 57 57 57 57 57
Copaxone 15 Cordarone-X 5 Corticosteroids and Related Agents 6 for Systemic Use 6 Corticosteroids Topical 7 Cosmegen 17 Coumadin 5 Creon 10000 6 Crotamiton 7 Crystaderm 7 Curam 10 Cvite 4 Cyclizine hydrochloride 14 Cyclogyl 21 Cyclopentolate hydrochloride 16 Cyclopohosphamide 16 Cycloserine 10 Cyklokapron 5 Cyptorerone acetate 6 Cyproterone acetate 6 Cyproterone acetate with 6 ethinyloestradiol 6	56 59 57 57 55 59 59 59 57 57 57 57 57 57 57 57 57 57 57 57 57
Copaxone 15 Cordarone-X 5 Corticosteroids and Related Agents 6 for Systemic Use 6 Corticosteroids Topical 7 Cosmegen 17 Coumadin 5 Creon 10000 6 Creon 25000 6 Crotamiton 7 Crystaderm 7 Curam 10 Cvite 4 Cyclizine hydrochloride 14 Cyclogyl 21 Cyclopentolate hydrochloride 16 Cyclopesphamide 16 Cycloserine 10 Cyklokapron 5 Cyproterone acetate 6 Cyproterone acetate 6	56 59 57 57 57 57 57 57 57 57 57 57 57 57 57
Copaxone 15 Cordarone-X 5 Corticosteroids and Related Agents 6 for Systemic Use 6 Corticosteroids Topical 7 Cosmegen 17 Coumadin 5 Creon 10000 6 Crotamiton 7 Crotamiton 7 Crystaderm 7 Curam 10 Cvite 4 Cyclizine hydrochloride 14 Cyclogyl 21 Cyclopentolate hydrochloride 16 Cyclopertolate hydrochloride 16 Cycloserine 10 Cycloserine 10 Cypterone acetate 5 Cyproterone acetate 5 Cyptarabine 16 Cytarabine 16	56 59 59 50 50 50 50 50 50 50 50 50 50

D-Penamine1	22
Dabigatran	55
Dacarbazine1	72
Dacarbazine APP1	72
Dactinomycin [Actinomycin D]1	72
Daivobet	79
Daivonex	79
Daktarin	73
Dalacin C	104
Dalteparin sodium	53
Danazol	00
Dantrium 1	00
Dantrium S29	120
Dantrolene	129
Daonil	.25
Dapa-Tabs	. 64
Dapsone1	110
Daraprim1	105
Darunavir1	117
Dasatinib1	176
Daunorubicin 1	172
DBL Acetylcysteine	221
DBL Aminophylline	213
DBL Bleomycin Sulfate1	171
DBL Carboplatin1	66
DBL Cisplatin1	66
DBL Dacarbazine1	172
DBL Docetaxel	
DBL Ergometrine	85
DBL Gemcitabine1	69
DBL Leucovorin Calcium	68
DBL Leucovorin Calcium	170
DBL Morphine Sulphate1	134
DBL Morphine Tartrate 1	121
DBL Naloxone Hydrochloride	201
DBL Octreotide1	100
DBL Pethidine Hydrochloride	102 195
DBL Vincristine Sulfate1	100
DBL VINCISUNE Sunate	00
De-worm	.99
Decozol	.44
Deferasirox	
Deferiprone	222
Deolate	08
Deoxycoformycin	174
Depo-Medrol Depo-Medrol with Lidocaine	.89
Depo-Medrol with Lidocaine	89
Depo-Provera	84
Depo-Testosterone	90
Deprim 1	
DermAssist	
Dermol74,	
Desferal	222
Desferrioxamine mesilate	
Desmopressin acetate	
Desmopressin-PH&T	
Detection of Substances in	

Urine
Dexamethasone
Hormone
Sensory217
Dexamethasone phosphate
Dexamethasone with framycetin and
gramicidin
Dexamethasone with neomycin
sulphate and polymyxin B
sulphate 217
Dexamfetamine sulfate
Dexmethsone
Dextrochlorpheniramine
maleate
Dextrose
Dextrose with electrolytes
DHC Continus
Diabetes
Diabetes Management
Diacomit
Diagnostic Agents
Diamide Relief
Diamox
Diasip
Diason RTH
Diazepam
Diazoxide
Diazonide
Diclofenac Sandoz
Diclofenac sodium
Musculoskeletal
Sensory
Differin
Difflam
Diflucan
Diflucan S29
Diflucortolone valerate
Digestives Including Enzymes
Digoxin
Dihydrocodeine tartrate
Dilantin
Dilantin Infatab
Diltiazem hydrochloride
Dilzem
Dimethicone75, 77 Dimethyl fumarate148
Dipentum
Diphtheria, tetanus and pertussis
vaccine
Diphtheria, tetanus, pertussis and
polio vaccine
Diphtheria, tetanus, pertussis, polio,
hepatitis B and haemophilus
influenzae type B vaccine
Diprosone
Diprosone OV73

Dipyridamole52
Disinfecting and Cleansing
Agents
Disopyramide phosphate
Disulfiram
Ditropan
Diuretics
Diurin 4063
Docetaxel172
Docetaxel Sandoz 172
Docusate sodium40
Docusate sodium with
sennosides 40
Dolutegravir 117
Domperidone 142
Donepezil hydrochloride 161
Donepezil-Rex161
Dopress
Dornase alfa
Dorzolamide hydrochloride
Dorzolamide with timolol
Dostinex
Dosulepin [Dothiepin]
hydrochloride
Dothiepin 135
Doxazosin
Doxepin hydrochloride
Doxine
Doxorubicin Ebewe
Doxorubicin hydrochloride
Doxy-50
Doxycycline 103
Doxylin 100103
DP Fusidic Acid Cream72
DP Lotion76
DP Lotn HC74
DP-Allopurinol127
Dr Reddy's Omeprazole23
Drugs Affecting Bone
Metabolism 122
Dual blood glucose and blood ketone
diagnostic test meter 26
Duocal Super Soluble Powder
Duolin
Duolin HFA
Durex Confidence82
Durex Extra Safe
Duride
Dynacirc-SRO62
- E -
e-chamber La Grande215
e-chamber Mask
e-chamber Turbo
E-Mycin 101
E-Mycin
E-Mycin

Easiphen Liquid	246
EasyCheck	85
Econazole nitrate	72
Efavirenz	
Efavirenz with emtricitabine and	
tenofovir disoproxil	
fumarate	116
Effient	
Eformoterol fumarate	
Efudix	
Egopsoryl TA	70
Elaprase	13
Elecare	
Elecare LCP	
Elemental 028 Extra	247
Elocon	74
Elocon Alcohol Free	
Eltrombopag	50
Eltroxin	
EMB Fatol	
Emend	142
Emend Tri-Pack	
EMLA	132
Emtricitabine	117
Emtricitabine with tenofovir disopr	
fumarate	114
Emtriva	117
Emulsifying ointment	76
Emulsifying ointment Enalapril maleate	76 58
Enalapril maleate	58
Enalapril maleate Enbrel	58 184
Enalapril maleate Enbrel Endocrine Therapy	58 184 182
Enalapril maleate Enbrel Endocrine Therapy Endoxan	58 184 182 166
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte	58 184 182 166 57
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B	58 184 182 166 57 263
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR	58 184 182 166 57 263 137
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR Enoxaparin sodium	58 184 182 166 57 263 137 54
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR Enoxaparin sodium Ensure	58 184 182 166 57 263 137 54 54 242
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR Enoxaparin sodium Ensure Ensure Plus	58 184 182 166 57 263 137 54 242 242
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR Enoxaparin sodium Ensure Plus Ensure Plus HN	58 184 182 166 57 263 137 54 242 242 241
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR Enoxaparin sodium Ensure Plus Ensure Plus Ensure Plus Ensure Plus RTH	58 184 182 166 57 263 137 54 242 242 241 241
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR Enoxaparin sodium Ensure Plus Ensure Plus Ensure Plus HN Ensure Plus RTH Ensure Plus RTH	58 184 182 166 57 263 137 54 242 242 241 241 130
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR Enatax XR Ensure Plus Ensure Plus HN Ensure Plus HN Ensure Plus RTH Entacapone Entapone	58 184 182 166 57 263 137 54 242 242 241 241 130 130
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR Ensure Blastar Sodium Ensure Plus Ensure Plus HN Ensure Plus HN Ensure Plus RTH Entacapone Entacapone Entecavir	58 184 182 166 57 263 137 54 242 241 241 241 241 130 130 131
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR. Enoxaparin sodium Ensure Plus Ensure Plus Ensure Plus HN Ensure Plus HN Ensure Plus RTH Entacapone Entacapone Entacavir. Entocort CIR	58 184 182 166 57 263 137 54 242 241 241 241 130 130 111
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR Enoxaparin sodium Ensure Plus Ensure Plus Ensure Plus HN Ensure Plus RTH Entacapone Entacapone Entacavir Entocort CIR Epilim	58 184 182 166 57 263 137 54 242 241 241 241 130 130 111 20 140
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR Enoxaparin sodium Ensure Plus Ensure Plus Ensure Plus HN Ensure Plus HN Ensure Plus RTH Entacapone Entacapone Entacore Entocort CIR Epilim Epilim Crushable	58 184 182 166 57 263 137 54 242 242 241 241 130 130 111 20 140 140
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B. Enlafax XR. Enoxaparin sodium. Ensure Plus Ensure Plus Ensure Plus HN. Ensure Plus HN. Ensure Plus HN. Ensure Plus HN. Entacapone Entacapone Entacort CIR. Epilim Epilim Crushable Epilim IV.	
Enalapril maleate Enbrel Endocrine Therapy Endoxan Eneryte Engerix-B Enlafax XR Enoxaparin sodium Ensure Plus Ensure Plus Ensure Plus HN Ensure Plus HN Ensure Plus RTH Entacapone Entacapone Entacapone Entacavir Entocort CIR Epilim Crushable Epilim IV Epilim S/F Liquid	
Enalapril maleate Enbrel Endocrine Therapy Endoxan Eneryte Engerix-B Enlafax XR Enoxaparin sodium Ensure Plus Ensure Plus Ensure Plus HN Ensure Plus HN Ensure Plus RTH Entacapone Entacapone Entacapone Entacavir Entocort CIR Epilim Crushable Epilim IV Epilim S/F Liquid Epilim Syrup	
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR Enoxaparin sodium Ensure Plus Ensure Plus Ensure Plus HN Ensure Plus HN Ensure Plus RTH Entacapone Entacapone Entacapone Entacapone Entacavir Entocort CIR Epilim Crushable Epilim S/F Liquid Epilim Syrup Epirubicin Ebewe	
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR Enoxaparin sodium Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Entacapone Epilim Crushable Epilim S/F Liquid Epilim Syrup Epirubicin Ebewe Epirubicin hydrochloride	
Enalapril maleate Enbrel Endocrine Therapy Endocrine Therapy Endoxan Enerlyte Engres Enalafax XR Enoxaparin sodium Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Entacapone Epilim Crushable Epilim S/F Liquid Epilim Syrup Epirubicin Ebewe Epirubicin hydrochloride Epoetin alfa [Erythropoietin alfa]	
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR Enoxaparin sodium Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Entacapone Epilim Crushable Epilim S/F Liquid Epilim Syrup Epirubicin Ebewe Epoetin alfa [Erythropoietin alfa] Epoprostenol	
Enalapril maleate Enbrel Endocrine Therapy Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR Enoxaparin sodium Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Entacapone Entacapone Entacapone Entacavir Entocort CIR Epilim Crushable Epilim S/F Liquid Epilim S/F Liquid Epilim Syrup Epirubicin Ebewe Eportin alfa [Erythropoietin alfa] Eportin alfa [Erythropoietin alfa] Eportin	
Enalapril maleate Enbrel Endocrine Therapy Endoxan Enerlyte Engerix-B Enlafax XR Enoxaparin sodium Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Ensure Plus Entacapone Epilim Crushable Epilim S/F Liquid Epilim Syrup Epirubicin Ebewe Epoetin alfa [Erythropoietin alfa] Epoprostenol	

ERA 101
Erbitux
Ergometrine maleate
Ergotamine tartrate with
caffeine 142
Erlotinib
Erythrocin IV101
Erythromycin ethyl succinate101
Erythromycin lactobionate 101
Erythromycin stearate101
Erythropoietin alfa50
Esbriet213
Escitalopram136
Escitalopram-Apotex 136
Eskazole
Estradot91
Estradot 50 mcg91
Estrofem
Etanercept 184
Ethambutol hydrochloride
Ethics Aspirin
Ethics Aspirin EC
Ethics Enalapril
Ethics Lisinopril
Ethinyloestradiol
Ethinyloestradiol with
desogestrel
Ethinyloestradiol with
levonorgestrel 83
Ethinyloestradiol with
norethisterone83
Ethosuximide 138
Etidronate disodium124
Etopophos 173
Etoposide173
Etoposide phosphate173
Etravirine
Eumovate74
Everet
Everolimus
Evista124
Evelon
Exemestane
Exjade
Extemporaneously Compounded
Preparations and
Galenicals
Eye Preparations
Eylea197
Ezetimibe65
Ezetimibe Sandoz65
Ezetimibe with simvastatin66
-F-
Factor eight inhibitor bypassing
fraction51
Febuxostat

Feed Thickener Karicare	
Aptamil	244
FEIBA NF	
Felodipine Fenpaed	
Fentanyl	133
Fentanyl Sandoz	
Ferinject	
Ferodan	47
Ferric carboxymaltose	47
Ferriprox	222
Ferro-F-Tabs	47
Ferro-tab	
Ferrograd	
Ferrograd F	47
Ferrous fumarate	47
Ferrous fumarate with folic acid	
Ferrous sulphate	47
Ferrous sulphate with folic acid	
Ferrum H	48
Fexofenadine hydrochloride	209
Fibro-vein	52
Filgrastim	55
Finasteride	85
Fingolimod	150
Firazyr	208
Flagyl	109
Flagyl-S	109
Flamazine	72
Flecainide acetate	
Fleet Phosphate Enema	
Flixonase Hayfever & Allergy	214
Flixotide	209
Flixotide Accuhaler	209
Floair	209
Florinef	
Fluanxol	
Fluarix Tetra	
Flucil	
Flucloxacillin	102
Flucloxin	102
Fluconazole	
Fludara Oral	
Fludarabine Ebewe	
Fludarabine phosphate	168
Fludrocortisone acetate	89
Fluids and Electrolytes	
Flumetasone pivalate	
Fluocortolone caproate with	
fluocortolone pivalate and	
cinchocaine	21
Fluorometholone	
Fluorouracil	
Fluorouracil Ebewe	. 168
Fluorouracil sodium	
Fluoxetine hydrochloride	

Flupenthixol decanoate	146
Flutamide	182
Flutamide Mylan	182
Flutamin	182
Fluticasone	209
Fluticasone furoate with	
vilanterol	210
Fluticasone propionate	214
Fluticasone with salmeterol	210
FML	
Foban	
Folic acid	
Food Thickeners	044
Foods And Supplements For Inborn	244
Errors Of Metabolism	045
Errors Or Metabolism	240
Foradil	210
Forteo	
Fortini	236
Fortini Multi Fibre	236
Fortisip	242
Fortisip Multi Fibre	243
Fosamax	
Fosamax Plus	
Fragmin	53
Framycetin sulphate	
Freestyle Optium	28
Freestyle Optium Ketone	25
Freestyle Optium Neo	25
Frisium	137
Frumil	. 64
Frusemide	63
Frusemide-Claris	63
Fucicort	75
Fucidin	
Fucithalmic	
Fungilin	_44
Furosemide [Frusemide]	63
fusidic acid	
Dermatological	75
Infection	
Sensory	
- G -	217
Gabapentin	138
Gacet	
Galsulfase	
Gardasil 9	204
Gastrodenol	
Gaviscon Double Strength	
Gaviscon Infant	
Gazyva	
Gefitinib	178
Gemcitabine Ebewe	169
Gemcitabine hydrochloride	
Gemfibrozil	
Gemzar	
Genoptic	216

	_
Genox1	84
Gentamicin sulphate	
Infection 1	04
Sensory2	
Gilenya1	50
Ginet	
Glatiramer acetate1	
Glibenclamide	
Gliclazide	
Glipizide	
Glivec1	
Glizide	
Glucagen Hypokit	
Glucagon hydrochloride	23
Glucerna Select	
Glucerna Select RTH2	34
Glucobay	25
Glucose [Dextrose]	56
Gluten Free Foods2	лл
Glycerin with sodium saccharin2	74 28
Glycerin with sucrose2	
Glycerol	20
Alimentary	11
Extemporaneous2	
Glyceryl trinitrate	20
Alimentary	იი
Cardiovascular	
Glycopyrronium	
Glycopyrronium bromide	22
Glycopyrronium with	
indacaterol2	
Glytrin	
Gold Knight	
Goserelin	
Gutron	60
Gynaecological Anti-infectives	85
-H-	~~
Habitrol1 Haemophilus influenzae type B	03
vaccine 2	61
Haldol1	
Haldol Concentrate1	
Haldol Decanoas1	
Haloperidol1	
Haloperidol decanoate1	
Hamilton Sunscreen	
Harvoni	
Havrix	
Havrix Junior2	
HBvaxPRO	03
healthE Dimethicone 10%	
healthE Dimethicone 4% Lotion	
healthE Dimethicone 5%	
healthE Glycerol BP2	
healthE Urea Cream	
Healtheries Simple Baking Mix2	44

Hyoscine butylbromide2	
Hyoscine hydrobromide14	3
Hypam 15	7
Hyperuricaemia and Antigout 12	7
Hypromellose21	
Hypromellose with dextran 220	0
Hysite21	9
-1-	
Ibiamox 102	
Ibuprofen 12	
Icatibant20	
Idarubicin hydrochloride 173	
Idursulfase4	
Ifosfamide16	
Ikorel6	
lloprost6	9
Imatinib mesilate 178	
Imatinib-AFT 178	
Imiglucerase4	
Imipramine hydrochloride13	
Imiquimod8	0
Immune Modulators11	
Immunosuppressants 184	4
Imuran	4
Incruse Ellipta212	2
Indacaterol21	
Indapamide64	
Infanrix IPV26	
Infanrix-hexa	
Infant Formulae	
Infatrini	
Influenza vaccine	
Influvac Tetra	
Inhaled Corticosteroids	9
Inhaled Long-acting Beta-adrenoceptor Agonists 210	^
Inset 30	
Inset II	
Insulin aspart	
Insulin aspart with insulin aspart	1
protamine	4
Insulin glargine	4
Insulin glulisine	
Insulin isophane	
Insulin isophane with insulin	ľ
neutral	4
Insulin lispro	4
Insulin lispro with insulin lispro	Ì
protamine	4
Insulin neutral	
Insulin pen needles	
Insulin pump	
Insulin pump accessories	
Insulin pump infusion set (steel	Ì
cannula)	5
Insulin pump infusion set (teflon	

cannula, angle insertion with
insertion device)
liseriion device)
Insulin pump infusion set (teflon
cannula, angle insertion) 36
Insulin pump infusion set (teflon
cannula, straight insertion with
insertion device)
Insulin pump infusion set (teflon
cannula, straight insertion)
Insulin pump reservoir
Insulin syringes, disposable with
attached needle
Intal Forte CFC Free213
Intelence
Interferon alfa-2a 118
Interferon alfa-2b 118
Interferon beta-1-alpha156
Interferon beta-1-beta156
Intra-uterine device82
Intron-A118
Invega Sustenna147
IPOĽ
Ipratropium bromide211, 214
Iressa
Irinotecan Actavis 100 169
Irinotecan Actavis 40
Irinotecan hydrochloride
Irinotecan-Rex
Iron polymaltose
Isentress
Ismo 20
Ismo 40 Retard66
Isoniazid 110
Isoprenaline66
Isoptin63
Isopto Carpine219
Isosorbide mononitrate
Isosource Standard241
Isosource Standard RTH241
Isotane 1071
Isotane 2071
Isotretinoin71
Ispaghula (psyllium) husk40
Isradipine
Isuprel
Itch-Soothe73
Itraconazole
Itrazole
Ivermectin
- J - Jadelle84
Jevity HiCal RTH
Jevity RTH241
Juno Pemetrexed 170
- K -

Kaletra 117

Kemadrin131
Kenacomb216
Kenacort-A 1090
Kenacort-A 4090
Kenalog in Orabase44
Ketocal 3:1249
KetoCal 4:1
Ketoconazole
Dermatological
Infection107
Ketogenic Diet249
Ketone blood beta-ketone
electrodes 25
Ketoprofen 121
KetoSens25
Ketostix
Keytruda
Kindergen
Kinson
Kivexa116
Klacid100
Kliogest
Kliovance
Kogenate FS52
Konakion MM
Konsyl-D40
-L-
L-asparaginase
Labetalol61
Lacosamide 139
Lactulose
Laevolac
Lamictal
Lamivudine 112, 117
Lamivudine Alphapharm117
Lamotrigine
Lamprene
Lanoxin
Lanoxin PG60
Lanoxin S2960
Lansoprazole22
Lantus24
Lantus SoloStar
Lanvis
Lanzol Relief22
Lapatinib ditosylate178
Largactil
Laronidase
Lasix
Latanoprost
Lax-Suppositories41
Lax-Tab
Laxatives
Laxsol
Ledipasvir with sofosbuvir 113
Leflunomide 122

Lenalidomide	173
Letrole	. 184
Letrozole	
Leukeran FC	166
Leukotriene Receptor	
Antagonists	. 213
Leunase	172
Leuprorelin	97
Leustatin	168
Levetiracetam	140
Levetiracetam-AFT	140
Levlen ED	
Levobunolol	218
Levocabastine	
Levodopa with benserazide	130
Levodopa with carbidopa	. 130
Levomepromazine	
hydrochloride	. 145
Levomepromazine maleate	145
Levonorgestrel	
Genito-Urinary	84
Hormone	92
Levothyroxine	93
Lidocaine [Lignocaine]131	-132
Lidocaine [Lignocaine]	
hydrochloride	. 132
Lidocaine [Lignocaine] with	
chlorhexidine	. 132
Lidocaine [Lignocaine] with	
prilocaine	. 132
Lidocaine-Claris	132
Lignocaine	
Hormone	89
Nervous131	-132
Lioresal Intrathecal	. 129
Lipazil	64
Lipid-Modifying Agents	64
Liquigen	233
Lisinopril	
Lithicarb FC	
Lithium carbonate	145
Livostin	
LMX4	132
Locacorten-Viaform ED's	216
Local preparations for Anal and	
Rectal Disorders	21
Locasol	
Locoid7	
Locoid Crelo	
Locoid Lipocream	
Locorten-Vioform	
Lodi	
Lodoxamide	
Logem	
Lomide	218
Lomustine	

Loniten	67
Loperamide hydrochloride	
Lopinavir with ritonavir	
Lopresor	
Loprofin	246
Loprofin Mix	
Lorafix	
Loratadine	
Lorazepam	
Lorfast	
Lormetazepam	
Lorstat	
Losartan Actavis	
Losartan potassium	
Losartan potassium with	
hydrochlorothiazide	59
Lovir	112
Lucrin Depot 1-month	
Lucrin Depot 3-month	
Ludiomil	
Lycinate	66
Lyderm	78
- M -	
m-Eslon	
Mabthera	
Macrogol 3350 with potassium	
chloride, sodium bicarbonate ar	nd
sodium chloride	
Macrogol 400 and propylene	
Macrogol 400 and propylene glycol	220
glycol	220 130
glycol Madopar 125	130
glycol	130 130
glycol Madopar 125 Madopar 250	130 130 130
glycol Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS	130 130 130 130
glycol Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide	130 130 130 130 130 130 228
glycol Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide	130 130 130 130 130 130 228
glycol Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide Magnesium sulphate	130 130 130 130 130 228 48
glycol Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide Magnesium sulphate Mantoux Maprotiline hydrochloride	130 130 130 130 130 228 228 48 270 136
glycol Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide Magnesium sulphate Mantoux Maprotiline hydrochloride Marevan	130 130 130 130 228 228 270 136 55
glycol Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide Magnesium sulphate Mantoux	130 130 130 130 228 228 270 136 55
glycol Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide Magnesium sulphate Mantoux Maprotiline hydrochloride Marvean Marve Blue Lotion SPF 50+ Marvelon 28	130 130 130 130 130 228 48 270 136 55 80 83
glycol Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide Magnesium sulphate Mantoux Marotiline hydrochloride Marevan Marine Blue Lotion SPF 50+ Marvelon 28 Mask for spacer device	130 130 130 130 130 228 48 270 136 55 80 83 214
glycol Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide Magnesium sulphate Mantoux Marotiline hydrochloride Marevan Marine Blue Lotion SPF 50+ Marvelon 28 Mask for spacer device Mast Cell Stabilisers	130 130 130 130 228 48 270 136 55 80 83 214 213
glycol Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide Magnesium sulphate Magnesium sulphate Marotiline hydrochloride Marotiline hydrochloride Marevan Marine Blue Lotion SPF 50+ Marvelon 28 Mask for spacer device Mast Cell Stabilisers Maxidex	130 130 130 130 130 228 48 270 136 55 80 83 214 213 217
glycol Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide Magnesium sulphate Mantoux Martouk Marotiline hydrochloride Marean Marine Blue Lotion SPF 50+ Marvelon 28 Mask for spacer device Mast Cell Stabilisers Maxidex Maxitrol	130 130 130 130 130 228 48 270 136 55 80 83 214 217 217
glycol Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide Magnesium sulphate Mantoux Martoux Martouk Marevan Marine Blue Lotion SPF 50+ Marvelon 28 Mask for spacer device Mask for spacer device Mast Cell Stabilisers Maxidex Maxidex Maxitrol MCT oil (Nutricia)	130 130 130 130 228 48 270 136 55 80 83 214 217 217 233
glycol Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide Magnesium sulphate Mantoux Martouk Marotiline hydrochloride Marean Marine Blue Lotion SPF 50+ Marvelon 28 Mask for spacer device Mast Cell Stabilisers Maxidex Maxitrol	130 130 130 130 228 48 270 136 55 80 83 214 217 217 233
glycol Madopar 125 Madopar 250 Madopar 250 Madopar Rapid Magnesium hydroxide Magnesium sulphate Mantoux Marvelon 28 Marvelon 28 Mask for spacer device Mask for spacer devi	
glycol Madopar 125 Madopar 250 Madopar 62.5 Madopar Rapid Magnesium sulphate Magnesium sulphate Maroux Marotiline hydrochloride Marotiline hydrochloride Marvelon 28 Marvelon 28 Mask for spacer device Mask for spacer device Mast Cell Stabilisers Maxitrol Maxitrol Maxitrol Mact oil (Nutricia) Measles, mumps and rubella vaccine Mebendazole	130 130 130 130 130 130 130 228 48 270 136 55 55 55 80 83 214 213 217 217 217 217 99
glycol Madopar 125 Madopar 250 Madopar 250 Madopar Agbid Magnesium sulphate Magnesium sulphate Martu and the subscription of the s	
glycol	
glycol	130 130 130 130 130 228 228 228 228 228 228 228 228 228 228 228 229 22
glycol	
glycol	

Megestrol acetate	182
Melatonin	156
Meloxicam	121
Melphalan	166
Menactra	267
Meningococcal (groups A, C, Y and	
W-135) conjugate vaccine	267
Meningococcal C conjugate	
vaccine	267
Menthol	
Mercaptopurine	
Mercilon 28	
Mesalazine	
Mesna	
Mestinon	
Metabolic Disorder Agents	
Metchek	
Meterol	
Metformin hydrochloride	
Metformin Mylan	
Methadone hydrochloride	20
Extemporaneous	000
Nervous Methatabs	
Methopt	
Methotrexate	
Methotrexate Ebewe	
Methotrexate Sandoz	
Methyl hydroxybenzoate	
Methylcellulose	228
Methylcellulose with glycerin and	
sodium saccharin	228
Methylcellulose with glycerin and	
sucrose	
Methyldopa	
Methyldopa Mylan	63
Methylnaltrexone bromide	
Methylphenidate hydrochloride	159
Methylphenidate hydrochloride	
extended-release	
Methylprednisolone	89
Methylprednisolone (as sodium	
succinate)	
Methylprednisolone aceponate	
Methylprednisolone acetate	89
Methylprednisolone acetate with	
lidocaine [Lignocaine]	89
Methylxanthines	
Metoclopramide Actavis 10	143
Metoclopramide hydrochloride	
Metolazone	
Metopirone	
Metoprolol succinate	
Metoprolol tartrate	
	61
Metronidazole	61 .109
	61 .109

Mexiletine hydrochloride	. 60
Mexiletine Hydrochloride USP	. 60
Miacalcic	.88
Micolette	.41
Miconazole	.44
Miconazole nitrate	
Dermatological	73
Genito-Urinary	
Micreme	
Micreme H	
Microgynon 20 ED	.70
Microgynon 30	.00 00
Microgynon 30	.83
Microgynon 50 ED	
Microlut	
Midazolam	157
Midazolam-Claris	
Midodrine	
Minerals	. 46
Mini-Wright AFS Low Range	214
Mini-Wright Standard	214
Minidiab	
Minims Pilocarpine	219
Minims Prednisolone	
Minirin	
Mino-tabs	
Minocycline hydrochloride	103
Minomycin	103
Minor Skin Infections	77
Minoxidil	
Mirena	
Mirtazapine	
Misoprostol	
Mitomycin C	
Mitozantrone	174
Mitozantrone Ebewe	
Mixtard 30	
Moclobemide	
Modafinil	
Modavigil	160
Moduretic	. 64
Molaxole	.41
Mometasone furoate	.74
Monogen	
Montelukast	213
Moroctocog alfa [Recombinant factor	
VIII]	
Morphine hydrochloride	134
Morphine sulphate	13/
Morphine tartrate	12/
Motetis	
Mouth and Throat	
Movapo	130
Moxifloxacin	
MSUD Maxamum	245
Mucilaginous laxatives with	
stimulants	40

Mucolytics	
Mucosoothe	
Multiple Sclerosis Treatments	148
Multivitamin renal	
Multivitamins	46
Mupirocin	
Muscle Relaxants	
Mvite	
Myambutol	
Mycobutin	110
Mycobulii	70
MycoNail	
Mycophenolate mofetil	
Mycostatin	73
Mydriacyl	
Mylan Atenolol	. 60
Mylan Clomiphen	
Mylan-Bosentan	. 67
Myleran	166
Myocrisin	122
Myometrial and Vaginal Hormone	
Preparations	. 85
Myozyme	41
- N -	
Nadolol	61
Naglazyme	
Nalcrom	01
Naloxone hydrochloride	001
Naltraccord	162
Naltrexone hydrochloride	
Naphazoline hydrochloride	220
Naphcon Forte	220
Naprosyn SR 1000	
Naprosyn SR 750	121
Naproxen	121
Nardil	136
Nasal Preparations	
Natalizumab	
Natulan	
Nausafix	143
Nausicalm	
Nauzene	
Navelbine	
Nedocromil	
Nefopam hydrochloride	
Neisvac-C	067
Neo-B12	
Neo-Mercazole	
Neocate Advance	
Neocate Gold	
Neocate Junior Unflavoured	
Neocate Junior Vanilla	247
Neocate LCP	
Neoral	
Neostigmine metilsulfate	
Nepro HP (strawberry)	237
Nepro HP (vanilla)	

Nepro HP RTH	.237
Nerisone	74
Neulactil	. 145
Neulastim	
Neurontin	
NeuroTabs	46
Vevirapine	
Nevirapine Alphapharm	116
Vicorandil	. 110
Vicotine	
Nicotinic acid	. 103
Nifedipine	
Nifuran	100
NIIUran	. 120
Nilotinib	. 179
Vilstat	
Alimentary	
Genito-Urinary	
Infection	
Nipent	. 174
Nitrados	
Nitrates	<mark>66</mark>
Nitrazepam	. 157
vitroderm TTS	<mark>66</mark>
Nitrofurantoin	. 120
Nitrolingual Pump Spray	<mark>66</mark>
Nivolumab	.204
Nizoral	. 107
Noctamid	
Nodia	20
Noflam 250	
Noflam 500	
Non-Steroidal Anti-Inflammatory	
Drugs	121
Nonacog alfa [Recombinant factor	
IX]	51
Nonacog gamma, [Recombinant	
Factor IX]	51
Vorethisterone	01
Genito-Urinary	8/
Hormone	
Norflex	
Norfloxacin	
Voriday 28	. 120
Norimin	83
Normacol Plus	
Normison	
Norpress	
Nortriptyline hydrochloride	
Norvir	
NovaSource Renal	
Novatretin	
NovoMix 30 FlexPen	
NovoRapid	
NovoRapid FlexPen	
NovoRapid Penfill	
NovoSeven RT	50

Noxafil1	
	07
Nozinan1	45
Nuelin2	13
Nuelin-SR 2	13
Nupentin1	38
Nutilis2	44
Nutrient Modules2	31
Nutrini Energy Multi Fibre 2	
Nutrini Energy RTH2	36
Nutrini Low Energy Multi Fibre 2	38
Nutrini RTH2	36
Nutrison 800 Complete Multi	
Fibre 2	41
Nutrison Concentrated2	43
Nutrison Energy2	41
Nutrison Energy Multi Fibre2	41
Nutrison Multi Fibre2	
Nutrison Standard RTH	41
Nyefax Retard	
Nystatin	02
Alimentary	<u>_</u>
Dermatological	79
Genito-Urinary	25
Infection 1	07
NZB Low Gluten Bread Mix	1
- 0 -	-
O/W Fatty Emulsion Cream	76
Obinutuzumab 1	ac
Octocog alfa [Recombinant factor	
VIII] (Advate)	51
Octocog alfa [Recombinant factor	Ŭ.
VIII] (Kogenate FS)	52
Octreotide	
Octreotide LAR (somatostatin	82
	82
analogue) 1	
analogue) 1	82
analogue)1 Oestradiol	82 91
analogue)1 Oestradiol Oestradiol valerate	82 91 91
analogue)1 Oestradiol Oestradiol valerate Oestradiol with norethisterone	82 91 91
analogue)1 Oestradiol Oestradiol valerate Oestradiol with norethisterone Oestriol	82 91 91 91
analogue)1 Oestradiol Oestradiol valerate Oestradiol with norethisterone Oestriol Genito-Urinary	82 91 91 91 91
analogue)1 Oestradiol Oestradiol valerate Oestradiol with norethisterone Oestriol Genito-Urinary Hormone	82 91 91 91 85 92
analogue)1 Oestradiol Oestradiol valerate Oestradiol with norethisterone Oestriol Genito-Urinary Hormone Oestrogens	82 91 91 91 85 92 91
analogue)1 Oestradiol Oestradiol valerate Oestradiol with norethisterone Oestriol Genito-Urinary Hormone Oestrogens Oil in water emulsion	82 91 91 91 85 92 91 76
analogue)	82 91 91 91 91 85 92 91 76 46
analogue)	82 91 91 91 85 92 91 76 46 64
analogue)	82 91 91 91 91 85 92 91 76 46 64
analogue)	82 91 91 91 91 85 92 91 76 46 20 21
analogue)	82 91 91 91 85 92 91 76 46 20 21 99
analogue)	82 91 91 91 91 85 92 91 76 64 20 21 99 23
analogue)	82 91 91 91 85 92 91 76 64 20 21 99 23 23
analogue)	82 91 91 91 85 92 91 76 64 20 23 23 23
analogue)	82 91 91 91 85 92 91 76 64 20 23 23 23 23
analogue)	82 91 91 91 91 85 92 91 64 20 23 23 23 23 93
analogue)	82 91 91 91 91 91 85 92 91 76 46 20 23 23 23 23 23 23 23 23 23 23 23 23 23
analogue)	82 91 91 91 85 92 91 76 64 20 23 23 23 23 23 23 23 23 23 23 23 23 23

Ondansetron143
Ondansetron ODT-DRLA 143
Ondansetron ODT-ORLA 143
One-Alpha45
Opdivo
Ora-Blend
Ora-Blend SF228
Ora-Plus
Ora-Sweet
Ora-Sweet SF228
Orabase
Oral Supplements/Complete Diet
(Nasogastric/Gastrostomy Tube
Feed) 234
Oratane
Orgran
Orion Temozolomide
Ornidazole 109
Orphenadrine citrate
Ortho-tolidine
Oruvail SR 121
Osmolite RTH
Other Endocrine Agents
Other Oestrogen Preparations
Other Progestogen
Preparations
Other Skin Preparations
Ovestin
Genito-Urinary
Hormone
Ox-Pam
Oxaliccord167
Oxaliplatin
Oxaliplatin Actavis 100
Oxaliplatin Actavis 50
Oxaliplatin Ebewe
Oxazepam
Oxis Turbuhaler
Oxpentifylline
Oxybutynin86
Oxybutynin
Oxycodone hydrochloride135
Oxycodone hydrochloride135 OxyNorm135
Oxycodone hydrochloride
Oxycodone hydrochloride 135 OxyNorm 135 Oxytocin 85 Oxytocin Apotex 85 Oxytocin BNM 85 Oxytocin with ergometrine maleate 85 Ozurdex 217 - P - Pacifen
Oxycodone hydrochloride 135 OxyNorm 135 Oxytocin 85 Oxytocin Apotex 85 Oxytocin BNM 85 Oxytocin with ergometrine maleate 85 Ozurdex 217 - P - Pacifien Paclitaxel 174
Oxycodone hydrochloride 135 OxyNorm 135 Oxytocin 85 Oxytocin Apotex 85 Oxytocin BNM 85 Oxytocin with ergometrine maleate 85 Ozurdex 217 - P - Pacifen Paclitaxel 174 Paclitaxel Actavis 174
Oxycodone hydrochloride 135 OxyNorm 135 Oxytocin 85 Oxytocin Apotex 85 Oxytocin BNM 85 Oxytocin with ergometrine maleate 85 Ozurdex 217 - P - Pacifen Paclitaxel 174 Paclitaxel Actavis 174 Paclitaxel Ebewe 174
Oxycodone hydrochloride 135 OxyNorm 135 Oxytocin 85 Oxytocin Apotex 85 Oxytocin BNM 85 Oxytocin with ergometrine 85 Ozurdex 217 - P - Pacifen Pacifen 129 Paclitaxel 174 Paclitaxel Actavis 174 Paclitaxel Ebewe 174 Paclitaxel Ebewe 174 Paclitaxel Ebewe 174
Oxycodone hydrochloride 135 OxyNorm 135 Oxytocin 85 Oxytocin Apotex 85 Oxytocin BNM 85 Oxytocin with ergometrine maleate 85 Ozurdex 217 - P - Pacifen Paclitaxel 174 Paclitaxel Actavis 174 Paclitaxel Ebewe 174

Pamisol124
Pancreatic enzyme
Pantoprazole23
Panzop Relief23
Panzytrat
Papaverine hydrochloride67
Para-amino salicylic acid 110
Paracare
Paracare Double Strength 133
Paracetamol 133
Paracetamol + Codeine
(Relieve) 135
Paracetamol with codeine 135
Paradigm 1.8 Reservoir
Paradigm 3.0 Reservoir
Paradigm 522 29
Paradigm 722
Paradigm Mio MMT-921
Paradigm Mio MMT-92337
Paradigm Mio MMT-92537
Paradigm Mio MMT-941 37
Paradigm Mio MMT-94337
Paradigm Mio MMT-94537
Paradigm Mio MMT-965
Paradigm Mio MMT-97537
Paradigm Quick-Set MMT-386
Paradigm Quick-Set MMT-387
Paradigm Quick-Set MMT-396
Paradigm Quick-Set MMT-397
Paradigm Quick-Set MMT-39838
Paradigm Quick-Set MMT-399
Paradigm Silhouette MMT-36836
Paradigm Silhouette MMT-37736
Paradigm Silhouette MMT-37836
Paradigm Silhouette MMT-381
Paradigm Silhouette MMT-38236
Paradigm Silhouette MMT-383
Paradigm Silhouette MMT-38436
Paradigm Sure-T MMT-86435
Paradigm Sure-T MMT-86635
Paradigm Sure-T MMT-87435
Paradigm Sure-T MMT-87635
Paradigm Sure-T MMT-88435
Paradigm Sure-T MMT-88635
Paraffin76
Paraffin liquid with soft white
paraffin220
Paraffin liquid with wool fat
Paraldehyde 137
Parasidose79
Parasiticidal Preparations77
Paritaprevir, ritonavir and ombitasvir
with dasabuvir113
Paritaprevir, ritonavir and ombitasvir
with dasabuvir and ribavirin 113
Parnate 136

Paromomycin	105
Paroxetine	. 137
Paser	110
Patanol	220
Paxam	. 148
Pazopanib	
Peak flow meter	
Pedialyte - Bubblegum	
Pediasure	.236
Pediasure RTH	
Pegaspargase	
Pegasys	118
Pegasys RBV Combination	
Pack	. 118
Pegfilgrastim	55
Pegylated interferon alfa-2a	
Pembrolizumab	
Pemetrexed	
Penicillamine	
Penicillin G	
PenMix 30	
PenMix 40	24 24
PenMix 50	
Pentasa	
Pentostatin [Deoxycoformycin]	
Pentoxifylline [Oxpentifylline]	
Peptamen Junior	
Peptisoothe	200
Peptisorb	228
Perhexiline maleate	
Pericyazine	1/5
Perindopril	
Perjeta	
Permethrin	
Perrigo Pertuzumab	100
Peteha	110
Pethidine hydrochloride	105
Pevaryl	
Pexsig	
Pfizer Exemestane Pharmacare	. 104
Pharmacy Health Sorbolene with	133
	70
Glycerin Pharmacy Services	
Pheburane Phenelzine sulphate	40 196
Phenobarbitone Phenobarbitone sodium	. 140
Extemporaneous	200
Nervous	
Phenothrin	
Phenothrin Phenoxybenzamine	79
hydrochloride	E0
Phenoxymethylpenicillin (Penicillin	38
V)	102
v / ·····	. 103

Phenytoin sodium137, 140
Phlexy 10246
Phosphate-Sandoz57
Phosphorus57
Phytomenadione52
Pilocarpine hydrochloride219
Pimafucort
Pindolol61
Pine tar with trolamine laurilsulfate
and fluorescein
Pinetarsol
Pioglitazone
Piportil
Pipothiazine palmitate
Pirfenidone
Pizotifen
PKU Anamix Infant
PKU Anamix Junior
PKU Anamix Junior LQ
PKU Lophlex LQ 10
PKU Lophlex LQ 20
PKU Lophlex Powder
PKU Lophlex Sensation 20246
Plaquenil
Plendil ER
Pneumococcal (PCV10) conjugate
vaccine
Pneumococcal (PCV13) conjugate
Theumococcal (TOVID) computate
vaccine
vaccine268Pneumococcal (PPV23)polysaccharide vaccinepolysaccharide vaccine269Pneumovax 23269Podophyllotoxin80Polaramine209Poliomyelitis vaccine269Poly-Gel220Poly-Gel220Poly-Tears220Poly-Visc220Poly-Visc220Polycal231Polyvinyl alcohol220Ponstan121
vaccine268Pneumococcal (PPV23)polysaccharide vaccinepolysaccharide vaccine269Pneumovax 23269Podophyllotoxin80Polaramine209Poliomyelitis vaccine269Poly-Gel220Poly-Gel220Poly-Tears220Poly-Visc220Poly-Visc220Poly-Uisc231Polyvinyl alcohol220Ponstan121Posaconazole107
vaccine
vaccine268Pneumococcal (PPV23)polysaccharide vaccinePneumovax 23269Podophyllotoxin80Polaramine209Poliomyelitis vaccine269Poloxamer40Poly-Gel220Poly-Tears220Polyval231Polyvinyl alcohol220Ponstan121Posaconazole107Postinor-184Potassium chloride56–57
vaccine268Pneumococcal (PPV23)polysaccharide vaccinePneumovax 23269Podophyllotoxin80Polaramine209Poliomyelitis vaccine269Poloxamer40Poly-Gel220Poly-Tears220Polyval231Polyvinyl alcohol220Ponstan121Posaconazole107Postinor-184Potassium chloride56–57Potassium citrate86
vaccine268Pneumococcal (PPV23)polysaccharide vaccinePneumovax 23269Podophyllotoxin80Polaramine209Poliomyelitis vaccine269Poloxamer40Poly-Gel220Poly-Tears220Polyval231Polyvinyl alcohol220Ponstan121Posaconazole107Postinor-184Potassium chloride56–57Potassium iodate46
vaccine268Pneumococcal (PPV23)polysaccharide vaccinePneumovax 23269Podophyllotoxin80Polaramine209Poliomyelitis vaccine269Poloxamer40Poly-Gel220Poly-Tears220Poly-Visc220Polyval231Polyvinyl alcohol220Ponstan121Posaconazole107Postinor-184Potassium chloride56–57Potassium citrate86Potassium iodate46Povidone iodine77
vaccine268Pneumococcal (PPV23)polysaccharide vaccinepolysaccharide vaccine269Pneumovax 23269Podophyllotoxin80Polaramine209Poliomyelitis vaccine269Poloxamer40Poly-Gel220Poly-Tears220Poly-Visc220Polycal231Polyvinyl alcohol220Ponstan121Posaconazole107Potassium citrate86Potassium iodate46Povidone iodine77Pradaxa55
vaccine268Pneumococcal (PPV23)polysaccharide vaccinepolysaccharide vaccine269Pneumovax 23269Podophyllotoxin80Polaramine209Poliomyelitis vaccine269Poloxamer40Poly-Gel220Poly-Gel220Poly-Gel220Polycal231Polycal231Polycal231Posaconazole107Postassium chloride56–57Potassium citrate86Potassium iodate46Povidone iodine77Pradaxa55Pramipexole hydrochloride130
vaccine268Pneumococcal (PPV23)polysaccharide vaccinepolysaccharide vaccine269Pneumovax 23269Podophyllotoxin80Polaramine209Polomyelitis vaccine269Poloxamer40Poly-Gel220Poly-Gel220Poly-Gel220Poly-Visc220Polycal231Polyuryl alcohol220Posaconazole107Postanor-184Potassium chloride56–57Potassium citrate86Potassium iodate46Povidone iodine775Pradaxa55Pramipexole hydrochloride130Prasugrel52
vaccine268Pneumococcal (PPV23)polysaccharide vaccinepolysaccharide vaccine269Pneumovax 23269Podophyllotoxin80Polaramine209Polomyelitis vaccine269Poloxamer40Poly-Gel220Poly-Gel220Poly-Gal231Polyvinyl alcohol220Ponstan121Posaconazole107Potassium chloride56–57Potassium citrate86Potassium cidate46Povidone iodine77Pradaxa55Pranipexole hydrochloride130Prasugrel52Pravastatin65
vaccine268Pneumococcal (PPV23)polysaccharide vaccine269Pneumovax 23269Podophyllotoxin80Polaramine209Poliomyelitis vaccine269Poloxamer40Poly-Gel220Poly-Gel220Poly-Tears220Poly-Visc220Poly-Visc220Poly-Visc220Polytal231Polytinyl alcohol220Postinor-184Potassium chloride56–57Potassium citrate86Potassium cidate46Povidone iodine77Pradaxa55Pramipexole hydrochloride130Prasugrel52Pravastatin65Praziquantel99
vaccine268Pneumococcal (PPV23)polysaccharide vaccine269Pneumovax 23269Podophyllotoxin80Polaramine209Poliomyelitis vaccine269Poloxamer40Poly-Gel220Poly-Gel220Poly-Tears220Poly-Visc220Polycal231Polycal231Polycal231Postinor-184Potassium chloride56–57Potassium citrate86Potassium citrate86Potadaxa55Pradaxa55Pramipexole hydrochloride130Prazyel52Pravastatin65Praziquantel99Prazosin58
vaccine268Pneumococcal (PPV23)polysaccharide vaccine269Pneumovax 23269Podophyllotoxin80Polaramine209Poliomyelitis vaccine269Poloxamer40Poly-Gel220Poly-Gel220Poly-Tears220Poly-Visc220Poly-Visc220Poly-Visc220Polytal231Polytinyl alcohol220Postinor-184Potassium chloride56–57Potassium citrate86Potassium cidate46Povidone iodine77Pradaxa55Pramipexole hydrochloride130Prasugrel52Pravastatin65Praziquantel99

Prednisolone acetate	010
	210
Prednisolone sodium	~ ~ ~
phosphate	218
Prednisolone-AFT	218
Prednisone	90
Pregabalin	140
Pregabalin Pfizer	140
Pregnancy Tests - hCG Urine	85
Premarin	
Prevenar 13	
Prezista	
Priadel	
Primacin	
Primaquine phosphate	
Primidone	
Primolut N	
Priorix	
Probenecid	129
Probenecid-AFT	129
Procaine penicillin	103
Procarbazine hydrochloride	174
Prochlorperazine	
Proctosedyl	
Procur	
Procyclidine hydrochloride	
Procytox	
Progesterone	92
Proglicem	
Proglycem	23
Progynova	91
Prokinex	142
Promethazine hydrochloride	209
Promethazine theoclate	143
Propafenone hydrochloride	
Propamidine isethionate	216
Propranolol	
Propylene glycol	229
Propylthiouracil	
Protaphane	
Protaphane Penfill	24
Protifar	233
Protionamide	
Provera	
Provera HD	
Provera S29	
PSM Citalopram	136
Psoriasis and Eczema	
Preparations	79
PTU	
Pulmicort Turbuhaler	209
Pulmocare	
Pulmozyme	
Puri-nethol	
Pyrazinamide	
Pyridostigmine bromide	
Pyridoxine hydrochloride	45

Pyrimethamine	.105
Pytazen SR	52
- Q -	
- Q - Q 300	. 109
Questran-Lite	65
Quetapel	. 145
Quetiapine	. 145
Quick-Set MMT-390 Quick-Set MMT-391	38
Quick-Set MMT-391	38
Quick-Set MMT-392	38
Quick-Set MMT-393	
Quinapril	58
Quinapril with hydrochlorothiazide	
hydrochlorothiazide	59
Quinine sulphate	. 109
Qvar	.209
RA-Morph	10/
Raloxifene hydrochloride	104
Raltegravir potassium	117
Ramipex	130
Ranbaxy-Cefaclor	001.
Ranitidine	33
Ranitidine Relief	
Rapamune	
Reandron 1000	
Recombinant factor IX	
Recombinant factor VIIa	
Recombinant factor VIII5	1-52
Rectogesic	22
Redipred	90
Refresh Night Time	.220
Relieve	.121
Relistor	40
Renilon 7.5	.237
Resonium-A	57
Resource Beneprotein	.233
Resource Diabetic	.234
Respigen	.211
Respiratory Devices	.214
Respiratory Stimulants	.215
Retinol palmitate	
ReTrieve	71
Retrovir	. 117
Revlimid	.173
Revolade	50
Rexacrom	
RexAir	
Reyataz	
Ribomustin	
Ricit	
Rifabutin	
Rifadin	
Rifampicin	
Rifaximin	
Rifinah	. 110

INDEX: Generic	Chemicals	and	Brands
----------------	-----------	-----	--------

Rilutek	131
Riluzole	
Riodine	.77
Risedronate Sandoz	125
Risedronate sodium	125
Risperdal Consta	147
Risperidone146–	147
Risperon	
Ritalin	
Ritalin LA	
Ritalin SR	
Ritonavir	
Rituximab	
Rivaroxaban	. 55
Rivastigmine	161
Rivotril	137
RIXUBIS	.51
Rizamelt	142
Rizatriptan	
Roferon-A	118
Rolin	
Ropinirole hydrochloride	
Rotarix	269
Rotavirus oral vaccine	269
Roxane	
Alimentary	.20
Cardiovascular	.61
Roxithromycin	
Rubifen	159
Rubifen SR	
Rulide D	
Rythmodan	.60
Rytmonorm	
- S -	
S-26 Gold Premgro	246
Sabril	
SalAir	
Salazopyrin	.21
Salazopyrin EN	21
Salbutamol210-	
Salbutamol with ipratropium	
bromide	211
Salicylic acid	
Salmeterol	
Sandomigran	
Sandostatin LAR	182
Scalp Preparations	
Scopoderm TTS	
Sebizole	
Sedatives and Hypnotics	
Seebri Breezhaler	
Selegiline hydrochloride	130
Senna	
Senokot	
Sensipar	
SensoCard	.28

=

Serenace	145
Seretide	210
Seretide Accuhaler	
Serevent	
Serevent Accuhaler	
Serophene	<mark>98</mark>
Sertraline	137
Sevredol	134
Sex Hormones Non	
Contraceptive	
Shield 49	
Shield Blue	
Shield XL	
shingles vaccine	
Sildenafil	<mark>69</mark>
Silhouette MMT-371	36
Silhouette MMT-373	
Siltuximab	
Simvastatin	
Simvastatin Mylan	
Sinemet	
Sinemet CR	
Sirolimus	
Slow-Lopresor	
Sodibic	
Sodium acid phosphate	41
Sodium alginate	20
Sodium aurothiomalate	122
Sodium benzoate	43
Sodium bicarbonate	
Blood	56–57
Extemporaneous	
Sodium calcium edetate	222
Sodium chloride	
Blood	<mark>56</mark>
Respiratory	214
Sodium citrate with sodium lauryl	
sulphoacetate	41
Sodium citro-tartrate	<mark>86</mark>
Sodium cromoglicate	
Alimentary	
Respiratory	213
Sensory	
Sodium fluoride	46
Sodium Fusidate [fusidic acid]	
Dermatological	72
Infection	
Sensory	217
Sodium hyaluronate [Hyaluronic	
acid]	
Sodium nitroprusside	
Sodium phenylbutyrate	
Sodium polystyrene sulphonate	57
Sodium tetradecyl sulphate	
Sodium valproate	
Sofradex	216

Soframycin	.216
Solian	.143
Solifenacin succinate	86
Solu-Cortef	
Solu-Medrol	89
Somatropin (Omnitrope)	93
Sotacor	62
Sotalol	
Spacer device	215
Span-K	
Spiolto Respimat	.212
Spiractin	64
Spiriva	211
Spiriva Respimat	
Spironolactone	64
Sporanox	107
Sprycel	176
Staphlex	102
Stemetil	
Stesolid	137
Stimulants/ADHD Treatments	157
Stiripentol	140
Stocrin	116
Stomahesive	44
Strattera	.157
Stromectol	
Suboxone	
Sucralfate	
Sulfadiazine Silver	72
Sulfadiazine sodium	. 105
Sulfasalazine	
Sulindac	
Sulphur	
Sulprix	.143
Sumatriptan	. 142
Sunitinib	. 180
Sunscreens	80
Sunscreens, proprietary	
Sure-T MMT-863	35
Sure-T MMT-865	35
Sure-T MMT-873	35
Sure-T MMT-875	35
Sure-T MMT-883	35
Sure-T MMT-885	35
Sustagen Diabetic	.234
Sustagen Hospital Formula	.242
Sustagen Hospital Formula	
Active	242
Sustanon Ampoules	
Sutent	
Sylvant	
Symbicort Turbuhaler 100/6	
Symbicort Turbuhaler 200/6	.210
Symbicort Turbuhaler 400/12	.210
Symmetrel	. 130
Sympathomimetics	66

Synacthen90
Synacthen Depot90
Synflorix
Synthroid
Syntometrine
Syrup (pharmaceutical grade)
Systane Unit Dose
- T -
Tacrolimus207
Tacrolimus Sandoz207
Tambocor60
Tambocor CR60
Tamoxifen citrate 184
Tamsulosin hydrochloride86
Tamsulosin-Rex86
Tap water
Tarceva177
Tasigna179
Tasmar
Tecfidera148
Tegretol 137
Tegretol CR 137
Telfast
Temazepam157
Temizole 20
Temozolomide
Tenofovir disoproxil
Tenofovir Disoproxil Teva112
Tenoxicam
Tepadina
Terazosin
Terbinafine
Terbutaline sulphate
Teriflunomide153
Teriparatide
Testosterone
Testosterone cipionate90
Testosterone esters90
Testosterone undecanoate90
Tetrabenazine131
Tetrabromophenol87
Tetracosactrin90
Tetracyclin Wolff103
Tetracycline 103
Teva171
Thalidomide 175
Thalomid175
Theophylline 213
Thiamine hydrochloride45
THIO-TEPA
Thioguanine171
Thiotepa
Thymol glycerin
Thyroid and Antithyroid Agents
Ticagrelor
Tilade

Tilcotil 121
Timolol
Cardiovascular 62
Sensory218
Timoptol XE
Tiotropium bromide
Tiotropium bromide with
olodaterol212
Tivicay117
TMP
TOBI106
Tobramycin
Infection
Sensory
Tobramycin Mylan 106
Tobrex217
Tofranil136
Tofranil s29136
Tolcapone130
Tolterodine
Topamax141
Topical Products for Joint and
Muscular Pain 122
Topiramate141
Topiramate Actavis141
Total parenteral nutrition (TPN)56
TPN
Tramadol hydrochloride
Tramal SR 100
Tramal SR 150 135
Tramal SR 200 135
Trandate61
Tranexamic acid52
Tranylcypromine sulphate136
Trastuzumab
Travatan
Travoprost
Travopt219
Treatments for Dementia161
Treatments for Substance
Dependence 161
Trental 40067
Tretinoin
Dermatological71
Oncology176
Trexate
Triamcinolone acetonide
Alimentary 44
Dermatological74
Hormone
Triamcinolone acetonide with
gramicidin, neomycin and nystatin
Dermatological
Sensory
Triazolam
Trichozole 109

Triclosan	75
Trimeprazine tartrate	.209
Trimethoprim	
Trimethoprim with	
sulphamethoxazole	
	100
[Co-trimoxazole]	
Trisequens	
Trisul	
Trophic Hormones	
Tropicamide	
Trusopt	
Truvada	.114
Tuberculin PPD [Mantoux] test	
Tubersol	
Two Cal HN	
Two Cal HN RTH	040
Tykerb	. 178
Tysabri	. 151
- U -	
Ultibro Breezhaler	.212
Ultraproct	21
Umeclidinium	.212
Umeclidinium with vilanterol	.213
Univent211	
Ural	
Urea	
Urex Forte Urinary Agents	03
Urinary Adents	
Urinary Tract Infections	.120
Urinary Tract Infections Uromitexan	.120 .174
Urinary Tract Infections Uromitexan Ursodeoxycholic acid	.120 .174 39
Urinary Tract Infections Uromitexan Ursodeoxycholic acid	.120 .174 39
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan	. 120 . 174 39 39
Urinary Tract Infections Uromitexan Ursodeoxycholic acid	. 120 . 174 39 39
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Utrogestan - V -	. 120 . 174 39 39 92
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Utrogestan - V - Vaccinations	. 120 . 174 39 39 92 92
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Utrogestan Vaccinations Vaccinations Vaclovir	. 120 . 174 39 39 92 . 260 . 112
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Utrogestan Vaccinations Vaccinations Valaciclovir	. 120 . 174 39 92 92 . 260 . 112 . 112
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Utrogestan Vaccinations Vacciovir Valaciclovir Valcyte	. 120 . 174 39 39 92 . 260 . 112 . 112 . 112
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Utrogestan Vaccinations Vacciovir Valaciclovir Valaganciclovir	. 120 . 174 39 92 . 260 . 112 . 112 . 112 . 112
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Utrogestan V - Vaccinations Valovir Valaciclovir Valoyte Valganciclovir Vallergan Forte	. 120 . 174 39 92 92 92 112 112 112 112 209
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Utrogestan V - Vaccinations Valaciclovir Valaciclovir Valaciclovir Valganciclovir Vallergan Forte Vancomycin	. 120 . 174 39 92 92 92 112 112 112 112 112 112 106
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Utrogestan V - Vaccinations Valaciclovir Valaciclovir Valaciclovir Vallergan Forte Vancomycin Vannair	. 120 . 174 39 92 . 260 . 112 . 112 . 112 . 112 . 209 . 106 . 210
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Utrogestan V - Vaccinations Valaciclovir Valaciclovir Valaciclovir Valganciclovir Vallergan Forte Vancomycin	. 120 . 174 39 92 . 260 . 112 . 112 . 112 . 112 . 209 . 106 . 210
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Utrogestan V - Vaccinations Valaciclovir Valaciclovir Valaciclovir Vallergan Forte Vancomycin Vannair	. 120 . 174 39 92 . 260 . 112 . 112 . 112 . 112 . 209 . 106 . 210
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Utrogestan Vacionations Valaciclovir Valaciclovir Valaciclovir Valaganciclovir Valganciclovir Vallergan Forte Vanlergan Forte Vancomycin Varenicline tartrate Varenicline tartrate Varicella vaccine [Chickenpox	. 120 . 174 39 92 . 260 . 112 . 112 . 112 . 112 . 209 . 106 . 210 . 163
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Utrogestan Vaccinations Vaccinations Valaciclovir Valaciclovir Valaciclovir Valaganciclovir Valaganciclovir Valeyte Valaganciclovir Valeyte Valaganciclovir Valeyte Valaganciclovir Valeyte Valaganciclovir Valeyte Valaganciclovir Valeyte Vanair Varenicline tartrate Varicella vaccine [Chickenpox vaccine]	. 120 . 174 39 92 . 260 . 112 . 112 . 112 . 112 . 209 . 106 . 210 . 163
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Utrogestan Vaccinations Vaclovir Valaciclovir Valaciclovir Valaganciclovir Varicella zoster virus (Oka strain) liv	. 120 . 174 39 92 . 260 . 112 . 112 . 112 . 112 . 209 . 106 . 210 . 163
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Vacosan Vaccinations Vaclovir Valaciclovir Varicella zoster virus (Oka strain) liv attenuated vaccine [shingles	. 120 . 174 39 39 92 . 260 . 112 . 112 . 112 . 112 . 210 . 163 . 270 /e
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Vaccinations Vaccinations Valaciclovir Vaccine] Varicella zoster virus (Oka strain) liv attenuated vaccine [shingles vaccine]	. 120 . 174 39 92 . 260 . 112 . 112 . 112 . 112 . 210 . 163 . 270 /e
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Vacosan Vaccinations Vaclovir Valaciclovir Valaciclovir Valaciclovir Valaciclovir Valaciclovir Valaciclovir Valacitovir	. 120 . 174 39 92 . 260 . 112 . 112 . 112 . 112 . 210 . 210 . 210 . 270 //e
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Vascinations Vaccinations Valcyte Valcyte Valganciclovir Valganciclovir Valganciclovir Valganciclovir Valganciclovir Valeyte Valganciclovir Valeyte Valganciclovir Valeyte Valeyte Valeyte Varenicline tartrate Varenicline tartrate Varenicline tartrate Varenicline tartrate Varcella vaccine [Chickenpox vaccine] Variella zoster virus (Oka strain) liv attenuated vaccine [shingles vaccine] Variux Various	. 120 . 174 39 92 260 112 112 112 112 209 210 210 270 /e 270 270 270
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Vacosan Vaccinations Valacitovir Valacitovir Valacitovir Valaganciclovir Valyte Valganciclovir Valganciclovir Valuergan Forte Valuergan Forte Valuergan Forte Varenicline tartrate Varenicline tartrate Varenicline tartrate Varenicline tartrate Varenicline tartrate Varicella vaccine [Chickenpox vaccine] Varicella zoster virus (Oka strain) liv attenuated vaccine [shingles vaccine] Varilix Various Vasodilators	. 120 . 174 39 92 260 112 112 112 112 209 210 210 210 270 270 270
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Vacosan Vacosan Vacosan Vacosan Vacosan Vacosan Vacosan Valter Varenic Vasodi Vasopressin Agonists	. 120 . 174 39 92 . 260 . 112 . 112 . 112 . 112 . 209 . 106 . 210 . 163 . 270 /e . 270 270 97 97
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Vacosan Vaccinations Valacitovir Valacitovir Valacitovir Valaganciclovir Valyte Valganciclovir Valganciclovir Valuergan Forte Valuergan Forte Valuergan Forte Varenicline tartrate Varenicline tartrate Varenicline tartrate Varenicline tartrate Varenicline tartrate Varicella vaccine [Chickenpox vaccine] Varicella zoster virus (Oka strain) liv attenuated vaccine [shingles vaccine] Varilix Various Vasodilators	. 120 . 174 39 92 . 260 . 112 . 112 . 112 . 112 . 209 . 106 . 210 . 163 . 270 /e . 270 270 97 97
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Vacosan Vacosan Vacosan Vacosan Vacosan Vacosan Vacosan Valter Varenic Vasodi Vasopressin Agonists	. 120 . 174 39 92 . 260 . 112 . 112 . 112 . 112 . 209 . 106 . 210 . 163 . 270 /e . 270 . 270 . 271 67 67 67
Urinary Tract Infections Uromitexan Ursodeoxycholic acid Ursosan Vacioacia Valoyie Valaciclovir Valaciclovir Valoyie Varicella vaccine [Chickenpox vaccine] Varicella zoster virus (Oka strain) liv attenuated vaccine [shingles vaccine] Various Vasodilators Vasopressin Agonists Vedafil	. 120 . 174 39 92 92 112 112 112 112 112 209 106 210 210 270 270 270 270 67 67 69

Venemil
Venomil
Ventavis
Ventolin210-211
Vepesid173
Verapamil hydrochloride63
Vergo 16142
Vermox
Verpamil SR63
Vesanoid176
Vesicare86
Vexazone25
Vfend 108
Viaderm KC75
Vidaza167
Viekira Pak 113
Viekira Pak-RBV113
Vigabatrin141
Vimpat
Vinblastina Teva
Vinblastine sulphate
Vincristine sulphate
Vincelbine
Vinorelbine Ebewe
Viramune Suspension
Viread
ViruPOS216
Vistil
Vistil Forte220
Vit.D345
VitA-POS
Vitabdeck46
Vitadol C45
Vital238
Vitamin A with vitamins D and C 45
Vitamin B complex45
Vitamin B6 25
Vitamins45–46
Vivonex TEN238
Volibris67
Voltaren 121
Voltaren D121
Voltaren Ophtha
Volumatic
Voriconazole108
Vonconazole
Vosol
Votrient
V IIdCK 100
- vv -
Warfarin sodium
Wart Preparations
Wasp venom allergy treatment208
Water
Blood56
Extemporaneous
Wool fat with mineral oil76
- X -

Xarelto
Xifaxan23
XMET Maxamum245
Xolair
XP Maxamaid246
XP Maxamum
Xylocaine
Xylocaine 2% Jelly131
Xyntha
- Z -
Zantac22
Zapril
Zarontin
Zaroxolyn63
Zarzio
Zavedos173
Zeffix112
Zeldox146
Zetlam
Ziagen
Zidovudine [AZT] 117
Zidovudine [AZT] with
lamivudine
Zimybe
Zinc and castor oil
Zinc sulphate
Zincaps
Zinnat
Ziprasidone
Zista
Zithromax
Zoladex
Zoledronic acid
Hormone
Musculoskeletal 126
Zoledronic acid Mylan
Zometa
Zopiclone
Zopicione Actavis
Zostavax
Zostrix
Zostrix HP132
Zuclopenthixol decanoate
Zuclopenthixol hydrochloride
Zusdone
Zyban
Zypine145
Zypine ODT
Zyprexa Relprevv146
Zytiga
Zyuya102



