February 2019 Volume 26 Number 0

Editors:

Kave Wilson & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9. 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am - 5pm weekdays)

Circulation

Published each April, August and December. Changes to the contents are published in monthly updates.

Accessible in an electronic format at no cost from the Health Professionals section of the PHARMAC website www.pharmac.govt.nz

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month. Alternatively there is a nominal charge for an annual subscription to the printed Schedule publications. To access either of these subscriptions visit our subscription website www.schedule.co.nz.

Production

Typeset automatically from XML and T_FX. XML version of the Schedule available from www.pharmac.govt.nz/pub/schedule/archive/

Programmers

Anrik Drenth & John Geering email: texschedule@pharmac.govt.nz ©Pharmaceutical Management Agency ISSN 1179-3686 pdf **(i)** Ccc ISSN 1172-9376 print This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to PHARMAC and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to PHARMAC should be in written form and not by reproduction of the PHARMAC logo. While care has been taken in compiling this Schedule, PHARMAC takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

	_	
Section A	General Rules	5
Section B	AI'	
Section b	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	37
	Cardiovascular System	47
	Dermatologicals	60
	Genito Urinary System	71
	Hormone Preparations – Systemic	77
	Infections – Agents For Systemic Use	88
	Musculoskeletal System	110
	Nervous System	119
	Oncology Agents & Immunosuppressants	152
	Respiratory System & Allergies	213
	Sensory Organs	221
	Various	226
Section C	Extemporaneous Compounds (ECPs)	228
	Extemporaneous Compounds (2013)	LLO
Section D	Special Foods	231
Section I	National Immunisation Schedule	250

Introducing PHARMAC

Index

Introducing PHARMAC

The Pharmaceutical Management Agency (PHARMAC) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. PHARMAC negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

PHARMAC's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about PHARMAC and the way we make funding decisions can be found on the PHARMAC website at http://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply:
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by PHARMAC.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements PHARMAC has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example




Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS



SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Sul Per	osidised G	rand or eneric anufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
LGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg p sachet		30	✓ Gavi	scon Infant
ODIUM ALGINATE Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (8.60)	60		scon Double ength
F Oral liq 500 mg with sodium bicarbonate 267 mg and calciul carbonate 160 mg per 10 ml		500 ml	Acide	ex
Phosphate Binding Agents				
LUMINIUM HYDROXIDE Tab 600 mg CALCIUM CARBONATE Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml)	12.56	100	✓ Alu-	Гab
Subsidy by endorsementOnly when prescribed for children under 12 years of agendorsed accordingly.		500 ml ate bindir	✓ Roxang agent and	
Antidiarrhoeals				
Agents Which Reduce Motility				
OPERAMIDE HYDROCHLORIDE – Up to 30 cap available on Fab 2 mg	10.75	400 400	✓ <u>Nodi</u> ✓ <u>Diam</u>	<u>a</u> nide Relief
Rectal and Colonic Anti-inflammatories				
UDESONIDE Cap 3 mg - Special Authority see SA1155 below - Retail pharmacy	166.50	90	✓ Ento	cort CIR

1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and

2 Any of the following:

- 2.1 Diabetes; or
- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
	Per	1	Manufacturer	

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)26.55	21.1 g OP	Colifoam
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab EC 500 mg49.50	100	✓ Asamax
Tab long-acting 500 mg59.05	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
Modified release granules, 1 g141.72	120 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g54.60	30	✓ Pentasa
OLSALAZINE		
Tab 500 mg93.37	100	✓ Dipentum
Cap 250 mg53.00	100	✓ Dipentum
SODIUM CROMOGLICATE		
Cap 100 mg92.91	100	✓ Nalcrom
SULFASALAZINE		
* Tab 500 mg14.00	100	✓ Salazopyrin
* Tab EC 500 mg	100	✓ Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

ELLIOCORTOLONE CAPROATE WITH ELLIOCORTOLONE PIVALATE AND CINCHOCAINE

INL	NOTIOUATIVE	ALA I L AIND OI	LUCCOTTOLONE OAI HOATE WITH LUCCOTTOLONE I W
P VIltraproct	30 g OP	6.35	Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cinchocaine hydrochloride 5 mg per g
✓ Ultraproct	12	2.66	Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and cinchocaine hydrochloride 1 mg
P Proctosedyl	30 g OP	15.00	HYDROCORTISONE WITH CINCHOCAINE Oint 5 mg with cinchocaine hydrochloride 5 mg per g
✓ Proctosedyl	12	9.90	Suppos 5 mg with cinchocaine hydrochloride 5 mg per g

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Management of Anal Fissures

GLYCERYL TRINITRATE - Special Authority see SA1329 below - Retail pharmacy

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

	/RRONII	

Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a
PSO.......17.14 10

HYOSCINE BUTYLBROMIDE

 ★ Tab 10 mg
 8.75
 100
 ✓ Buscopan

 ★ Inj 20 mg, 1 ml - Up to 5 inj available on a PSO
 9.57
 5
 ✓ Buscopan

MEBEVERINE HYDROCHLORIDE

Antiulcerants

Antisecretory and Cytoprotective

MISOPROSTOL

Helicobacter Pylori Eradication

CLARITHROMYCIN

Tab 500 mg − Subsidy by endorsement.......10.40 14 ✓ Apo-Clarithromycin

- a) Maximum of 14 tab per prescription
- Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.
 Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

H2 Antagonists

RANITIDINE - Only on a prescription

	Tab 150 mg	12.91	500	Ranitidine Relief
	Tab 300 mg		500	✓ Ranitidine Relief
*	Oral liq 150 mg per 10 ml	5.14	300 ml	✓ Peptisoothe
*	Inj 25 mg per ml, 2 ml	8.75	5	✓ Zantac

Proton Pump Inhibitors

LA	NS	OΡ	RA	١ZC)LE

*	Cap 15 mg4.58	100	Lanzoi Reliet
*	Cap 30 mg5.41	100	✓ Lanzol Relief

✓ Max Health

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
OMEPRAZOLE				
For omeprazole suspension refer Standard Formulae, page	228			
* Cap 10 mg	1.98	90	•	Omeprazole actavis 10
* Cap 20 mg	1.96	90	✓	Omeprazole actavis 20
* Cap 40 mg	3.12	90	•	Omeprazole actavis 40
Powder – Only in combination Only in extemporaneously compounded omegrazole sus		5 g	•	Midwest
* Inj 40 mg ampoule with diluent		5	•	<u>Dr Reddy's</u> <u>Omeprazole</u>
* Tab EC 20 mg	2 41	100	/	Panzop Relief
★ Tab EC 40 mg		100		Panzop Relief
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE				
Tab 120 mg	14.51	50	✓	Gastrodenol S29
SUCRALFATE				
Tab 1 g	35.50	120		
	(48.28)			Carafate
Bile and Liver Therapy				
RIFAXIMIN - Special Authority see SA1461 below - Retail phar Tab 550 mg	•	56	/	Xifaxan
■ SA1461 Special Authority for Subsidy nitial application only from a gastroenterologist, hepatologist or nepatologist. Approvals valid for 6 months where the patient has olerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or Practitio nepatologist. Approvals valid without further renewal unless noticenefiting from treatment.	hepatic encephalopa	thy d	lespite an n of a gast	adequate trial of maximun

Diabetes

Hyperglycaemic Agents

DIAZOXIDE - Special Authority see SA1320 below - Ref	ail pharmacy		
Cap 25 mg	110.00	100	✓ Proglicem S29
Cap 100 mg	280.00	100	✓ Proglicem S29
Oral liq 50 mg per ml	620.00	30 ml OP	✓ Proglycem §29

⇒SA1320 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where used for the treatment of confirmed hypoglycaemia caused by hyperinsulinism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

GLUCAGON HYDROCHLORIDE

Inj 1 mg syringe kit − Up to 5 kit available on a PSO......32.00 1 ✓ Glucagen Hypokit

	Subsidy		Fully	Brand or
	(Manufacturer's F		sidised	Generic
	\$	Per		Manufacturer
Inculin Chart acting Drangrations				
Insulin - Short-acting Preparations				
NSULIN NEUTRAL				
Inj human 100 u per ml	25.26	10 ml OP	✓	Actrapid
,			√ ⊦	lumulin R
▲ Inj human 100 u per ml, 3 ml	42.66	5	√ /	Actrapid Penfill
,			✓	lumulin R
Insulin - Intermediate-acting Preparations				
NSULIN ASPART WITH INSULIN ASPART PROTAMINE				
Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	✓ N	NovoMix 30 FlexPen
NSULIN ISOPHANE		-	-	
	17.60	10 ml OP	./ L	Humulin NPH
Inj human 100 u per ml	17.00	10 1111 OF		
Ini human 100 u nav ml. 0 ml	00.00	_		Protaphane
Inj human 100 u per ml, 3 ml	29.86	5		Humulin NPH
			•	Protaphane Penfill
NSULIN ISOPHANE WITH INSULIN NEUTRAL				
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP		lumulin 30/70
			-	Mixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5		lumulin 30/70
				PenMix 30
				PenMix 40
			✓ F	PenMix 50
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE				
Inj lispro 25% with insulin lispro protamine 75% 100 u per ml,				
3 ml		5	✓ ⊦	lumalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml,		ū		
3 ml		5	✓ F	lumalog Mix 50
V 111		<u> </u>	• •	idilialog illix oo
Insulin - Long-acting Preparations				
NSULIN GLARGINE				
▲ Inj 100 u per ml, 10 ml	63.00	1	√ I	_antus
Inj 100 u per ml, 3 ml	94 50	5		antus
Inj 100 u per ml, 3 ml disposable pen		5	_	antus SoloStar
L my 100 a por mi, o mi aloposablo por minimi minimi				
Insulin - Rapid Acting Preparations				
NSULIN ASPART				
▲ Inj 100 u per ml, 10 ml	30.03	1	√ N	NovoRapid
Inj 100 u per ml, 3 ml	51 10	5		NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe		5		NovoRapid FlexPen
		5	• 1	NOVONAPIU FIEKFEII
NSULIN GLULISINE				
Inj 100 u per ml, 10 ml		1	_	Apidra
Inj 100 u per ml, 3 ml		5		Apidra
Inj 100 u per ml, 3 ml disposable pen	46.07	5	✓	Apidra SoloStar
NSULIN LISPRO				
▲ Inj 100 u per ml, 10 ml	34.92	10 ml OP	✓	lumalog
▲ Inj 100 u per ml, 3 ml		5	_	lumalog
·				

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Alpha Glucosidase Inhibitors				
ACARBOSE				
Tab 50 mg		90		Glucobay
Tab 100 mg		90		Glucobay
	11.24	50	•	Acarbose Mylan S29
Oral Hypoglycaemic Agents				
GLIBENCLAMIDE				
* Tab 5 mg	6.00	100	✓	Daonil
GLICLAZIDE				
* Tab 80 mg	10.29	500	✓	Glizide
GLIPIZIDE				
* Tab 5 mg	3.27	100	✓	<u>Minidiab</u>
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg		1,000		Apotex
	(9.59)			Metchek
Apotex to be Sole Supply on 1 May 2019	7.04			
* Tab immediate-release 850 mg		500		Apotex Motformin Mulan
Apotex to be Sole Supply on 1 May 2019	(7.82)			Metformin Mylan
(Metchek Tab immediate-release 500 mg to be delisted 1 May 2	2019)			
(Metformin Mylan Tab immediate-release 850 mg to be delisted	,			
PIOGLITAZONE	ay =0.07			
* Tab 15 mg	3 47	90	1	Vexazone
* Tab 30 mg		90		Vexazone
* Tab 45 mg		90		Vexazone
VILDAGLIPTIN				
Tab 50 mg	40.00	60	✓	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride	40.00	60	1	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride		60	✓	Galvumet
- · ·				

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes: or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

1 OP CareSens Dual

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes; or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 diagnostic test

1 OP ✓ CareSens N

✓ CareSens N POP 20.00

✓ CareSens N Premier

Note: Only 1 meter available per PSO

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	•	CareSens N
		1	CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips	est OP 🗸 Se	nsoCard
---------------------------	-------------	---------

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

INSULIN PEN NEEDLES - Maximum of 100 dev per prescription

*	29 g × 12.7 mm	100	B-D Micro-Fine
	31 g × 5 mm11.75		✓ B-D Micro-Fine
	31 g × 6 mm		✓ ABM
*	31 g × 8 mm10.50	100	✓ B-D Micro-Fine
	32 g × 4 mm10.50		✓ B-D Micro-Fine

		Subsidy		Fully	Brand or
		(Manufacturer's Price)		Subsidised	Generic
_		\$	Per		Manufacturer
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE	E - Maximum of 100	dev p	er prescrip	tion
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.00	100	✓ [B-D Ultra Fine
		1.30	10		
		(1.99)		E	B-D Ultra Fine
*	Syringe 0.3 ml with 31 g × 8 mm needle	13.00	100	✓ [B-D Ultra Fine II
		1.30	10		
		(1.99)			B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g × 12.7 mm needle	13.00	100	✓ [B-D Ultra Fine
		1.30	10		
		(1.99)			B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	13.00	100	✓ [B-D Ultra Fine II
		1.30	10		
		(1.99)			B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.00	100	✓ [B-D Ultra Fine
		1.30	10		
		(1.99)			B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.00	100	✓ [B-D Ultra Fine II
		1.30	10		
		(1.99)		E	B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription
- c) Maximum of 1 insulin pump per patient each four year period.

Min basal rate 0.001 U/h	4,500.00	1	✓ Tandem t:slim X2
Min basal rate 0.025 U/h	8.800.00	1	✓ MiniMed 640G

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Fither:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
(Waitulatule 31 Noe)	Per	oubsidised ✓	Manufacturer	

continued...

- 4.1 Applicant is a relevant specialist; or
- 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — **(HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment: and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Fither:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Insulin Pump Consumables

⇒SA1604 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional): and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Fither:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist: or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 Applicant is a relevant specialist; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (**Previous use before 1 September 2012**) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less

1 OP

✓ Tandem Cartridge

	Subsidy (Manufacturer's Price) \$	Subsic Per	dised	Brand or Generic Manufacturer
continued				
than 80 mmol/mol; and				
2 The patient's HbA1c has not deteriorated more than 5 mr3 The patient has not had an increase in severe unexplaine4 Either:				; and
4.1 Applicant is a relevant specialist; or4.2 Applicant is a nurse practitioner working within the	ir vocational scope.			
INSULIN PUMP ACCESSORIES – Special Authority see SA160 a) Maximum of 1 cap per prescription b) Only on a prescription c) Maximum of 1 prescription per 180 days. Battery cap	32.00	pharmacy		imas Battery Cap
(Animas Battery Cap Battery cap to be delisted 1 October 2019)				
INSULIN PUMP CARTRIDGE – Special Authority see SA1604 (a) Maximum of 3 sets per prescription b) Only on a prescription	on page 17 – Retail ph	armacy		

c) Maximum of 13 packs of cartridge sets will be funded per year.

Cartridge 300 U, t:lock × 10......50.00

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA) - Special Authority see SA1604 on page 17 - Retail pharmacy

a١	Maximum	of 3 sets	ner nres	crintion

b) Only on a prescription

c)	Maximum	of 13	infusion sets will be funded per year	

	 Maximum of 13 infusion sets will be funded per year. 			
	10 mm steel needle; 29 G; manual insertion; 60 cm tubing x			
	10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T MMT-884
	10 mm steel needle; 29 G; manual insertion; 60 cm tubing ×			WIWI I -OO4
	10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-883
	10 mm steel needle; 29 G; manual insertion; 80 cm tubing x			
	10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T
				MMT-886
	10 mm steel needle; 29 G; manual insertion; 80 cm tubing x			
	10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-885
	6 mm steel cannula; straight insertion; 60 cm grey line \times 10 with			
	10 needles	130.00	1 OP	✓ Contact-D
	6 mm steel needle; 29 G; manual insertion; 60 cm tubing x			
	10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T MMT-864
	Construction CO on the contract of the contrac			IVIIVI I -864
	6 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock	120.00	1 OP	✓ Sure-T MMT-863
	6 mm steel needle; 29 G; manual insertion; 80 cm tubing ×	130.00	TOF	Jule-1 WIWI1-003
	10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T
				MMT-866
	6 mm steel needle; 29 G; manual insertion; 80 cm tubing ×			
	10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-865
	8 mm steel cannula; straight insertion; 110 cm grey line ×			
	10 with 10 needles	130.00	1 OP	✓ Contact-D
	8 mm steel cannula; straight insertion; 60 cm grey line \times 10 with			
	10 needles	130.00	1 OP	✓ Contact-D
	8 mm steel needle; 29 G; manual insertion; 60 cm tubing \times			
	10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T
				MMT-874
	8 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock	120.00	1 OP	✓ Sure-T MMT-873
	8 mm steel needle; 29 G; manual insertion; 80 cm tubing ×	130.00	TOP	Sure-1 WIWI1-073
	10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T
	10 With 10 Hoodies	100.00	1 01	MMT-876
	8 mm steel needle; 29 G; manual insertion; 80 cm tubing ×			
	10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-875
۰.,	ntact-D 6 mm steel cannula: straight insertion: 60 cm gray line > 1		lac to ha dalic	etad 1 October 2010)

(Contact-D 6 mm steel cannula; straight insertion; 60 cm grey line \times 10 with 10 needles to be delisted 1 October 2019) (Contact-D 8 mm steel cannula; straight insertion; 110 cm grey line \times 10 with 10 needles to be delisted 1 October 2019) (Contact-D 8 mm steel cannula; straight insertion; 60 cm grey line \times 10 with 10 needles to be delisted 1 October 2019)

Subsidy		Fully	Brand or
(Manufacturer's Price)		sidised	Generic
\$	Per	1	Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) – Special Authority see SA1604 on page 17 – Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

c) Maximum of to initiation sets will be funded per year.			
6 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles	130.00	1 OP	✓ TruSteel
6 mm steel cannula; straight insertion; 81 cm line × 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles	130.00	1 OP	✓ TruSteel

INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION WITH INSERTION DEVICE) - Special Authority see SA1604 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.
- 13 mm teflon cannula; angle insertion; insertion device; 110 cm 1 OP ✓ Inset 30 13 mm teflon cannula; angle insertion; insertion device; 60 cm ✓ Inset 30 1 OP 13 mm teflon cannula; angle insertion; insertion device; 110 cm 1 OP ✓ AutoSoft 30 13 mm teflon cannula; angle insertion; insertion device; 60 cm line x 10 with 10 needles......140.00 1 OP ✓ AutoSoft 30

(Inset 30 13 mm teflon cannula; angle insertion; insertion device; 110 cm grey line × 10 with 10 needles to be delisted 1 October 2019)

(Inset 30 13 mm teflon cannula; angle insertion; insertion device; 60 cm grey line × 10 with 10 needles to be delisted 1 October 2019)

Subsidy		Fully	Brand or	
(Manufacturer's Price)) Sul	osidised	Generic	
\$	Per	/	Manufacturer	

INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION) - Special Authority see SA1604 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

 c) Maximum of 13 infusion sets will be funded per year. 			
13 mm teflon cannula; angle insertion; 120 cm line \times 10 with			
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-382
13 mm teflon cannula; angle insertion; 45 cm line × 10 with			
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-368
13 mm teflon cannula; angle insertion; 60 cm line × 10 with			
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-381
13 mm teflon cannula; angle insertion; 80 cm line × 10 with			
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-383
17 mm teflon cannula; angle insertion; 110 cm line × 10 with			1111111 000
10 needles	120.00	1 OP	✓ Paradigm Silhouette
TO ficeules	130.00	TOF	MMT-377
17 mm teflon cannula; angle insertion; 110 cm line \times 10 with			
10 needles; luer lock	130.00	1 OP	✓ Silhouette MMT-371
17 mm teflon cannula; angle insertion; 60 cm line × 10 with			
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-378
17 mm teflon cannula; angle insertion; 60 cm line × 10 with			
10 needles; luer lock	130.00	1 OP	✓ Silhouette MMT-373
17 mm teflon cannula; angle insertion; 80 cm line × 10 with	100.00	1 01	- Simouette iiiii 1-070
	120.00	1 OP	✓ Paradigm Silhouette
10 needles	130.00	1 05	▼ Faraulylli Silliouelle

MMT-384

Subsidy Fully Brand or (Manufacturer's Price) Generic Subsidised \$ Per Manufacturer

1 OP

1 OP

✓ Inset II

Paradigm Mio

INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION WITH INSERTION DEVICE) - Special Authority see SA1604 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula: straight insertion: insertion device:

6 mm teflon cannula; straight insertion; insertion device; 80 cm

9 mm teflon cannula: straight insertion; insertion device:

9

clear tubing × 10 with 10 needles......130.00

6 mm teflon cannula; straight insertion; insertion device; 45 cm blue tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Mio MMT-941
6 mm teflon cannula; straight insertion; insertion device; 45 cm pink tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Mio MMT-921
6 mm teflon cannula; straight insertion; insertion device; 60 cm blue tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Mio MMT-943
6 mm teflon cannula; straight insertion; insertion device; 60 cm grey line × 10 with 10 needles		1 OP	✓ Inset II
pink tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Mio MMT-923
blue tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Mio

		MMT-965
6 mm teflon cannula; straight insertion; insertion device; 80 cm		
pink tubing × 10 with 10 needles130.00	1 OP	✓ Paradigm Mio
		MMT-925

•	110 cm grey line × 10 with 10 needles	.140.00	1 OP	✓ Inset II
,	9 mm teflon cannula; straight insertion; insertion device; 60 cm			
	gray line v 10 with 10 needles	140.00	1 OP	✓ Incat II

· · ·			
9 mm teflon cannula; straight insertion; insertion device; 80 c	m		
clear tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Mid MMT-975
			IVIIVI I -9/5

clear tubing × 10 with 10 needles130.00	1 OP	✓ Paradigm Mio MMT-975
6 mm teflon cannula; straight insertion; insertion device; 110 cm line × 10 with 10 needles140.00	1 OP	✓ AutoSoft 90
6 mm teflon cannula; straight insertion; insertion device; 60 cm		

line x 10 with 10 needles	140.00	1 OP	✓ AutoSoft 90
mm teflon cannula; straight insertion; insertion device;			
110 cm line x 10 with 10 needles	140 00	1 OP	✓ AutoSoft 90

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer	
9 mm teflon cannula; straight insertion; insertion device; 60 line × 10 with 10 needles		1 OP	✓ A	autoSoft 90	

(Inset II 6 mm teflon cannula; straight insertion; insertion device; 110 cm grey line x 10 with 10 needles to be delisted 1 October 2019)

(Inset II 6 mm teflon cannula; straight insertion; insertion device; 60 cm grey line x 10 with 10 needles to be delisted 1 October

(Inset II 9 mm teflon cannula; straight insertion; insertion device; 110 cm grey line x 10 with 10 needles to be delisted 1 October 2019)

(Inset II 9 mm teflon cannula; straight insertion; insertion device; 60 cm grey line x 10 with 10 needles to be delisted 1 October 2019)

INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION) - Special Authority see \$A1604 on page 17 -Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm teflon cannula; straight insertion; 110 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-398
6 mm teflon cannula; straight insertion; 110 cm tubing × 10 with 10 needles; luer lock	130.00	1 OP	✓ Quick-Set MMT-391
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Quick-Set
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with	100.00	4.00	MMT-399
10 needles; luer lock	130.00	1 OP	✓ Quick-Set MMT-393
10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-387
9 mm teflon cannula; straight insertion; 106 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-396
9 mm teflon cannula; straight insertion; 110 cm tubing x 10 with 10 needles; luer lock	130.00	1 OP	✓ Quick-Set MMT-390
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-397
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 10 needles; luer lock	130.00	1 OP	✓ Quick-Set MMT-392
9 mm teflon cannula; straight insertion; 80 cm tubing × 10 with	100.00	. 0.	- quick oot illim 1 ooz
10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-386

	Subsidy (Manufacturer's Price)		Fully	Brand or Generic
	\$	Per		Manufacturer
INSULIN PUMP RESERVOIR - Special Authority see SA1604	on page 17 – Retail pl	narmacy		
a) Maximum of 3 sets per prescription		•		
b) Only on a prescription				
c) Maximum of 13 packs of reservoir sets will be funded pe	r year.			
10 × luer lock conversion cartridges 1.8 ml for Paradigm pur	mps50.00	1 OP	1	ADR Cartridge 1.8
Cartridge 200 U, luer lock × 10	50.00	1 OP	1	Animas Cartridge
Cartridge for 5 and 7 series pump; 1.8 ml × 10	50.00	1 OP	✓	Paradigm
				1.8 Reservoir
Cartridge for 7 series pump; 3.0 ml x 10	50.00	1 OP	✓	Paradigm
		-		3.0 Reservoir
Syringe and cartridge for 50X pump, 3.0 ml × 10	50.00	1 OP	1	50X 3.0 Reservoir
(Animas Cartridge Cartridge 200 U, luer lock x 10 to be delisted				
(50X 3.0 Reservoir Syringe and cartridge for 50X pump, 3.0 ml)	,	October 20	019)	

Digestives Including Enzymes

PANCREATIC ENZYME

Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	.34.93	100	✓ Creon 10000
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase, 1,250 U protease))	.94.40	100	✓ Panzytrat
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	.94.38	100	✓ <u>Creon 25000</u>
URSODEOXYCHOLIC ACID – Special Authority see SA1739 below – Cap 250 mg		y 100	✓ <u>Ursosan</u>

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IqM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l: decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:
Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Laxatives

Bulk-forming Agents			
ISPAGHULA (PSYLLIUM) HUSK – Only on a prescription * Powder for oral soln	6.05	500 g OP	✓ Bonvit ✓ <u>Konsyl-D</u>
MUCILAGINOUS LAXATIVES WITH STIMULANTS			
* Dry	6.02 (17.32) 2.41	500 g OP 200 g OP	Normacol Plus
	(8.72)	Ü	Normacol Plus
Faecal Softeners			
DOCUSATE SODIUM — Only on a prescription # Tab 50 mg # Tab 120 mg # Enema conc 18% (Coloxyl Enema conc 18% to be delisted 1 April 2019)	3.13	100 100 100 ml OP	✓ <u>Coloxyl</u> ✓ <u>Coloxyl</u> ✓ Coloxyl
DOCUSATE SODIUM WITH SENNOSIDES * Tab 50 mg with sennosides 8 mg POLOXAMER – Only on a prescription Not funded for use in the ear.	3.10	200	✓ <u>Laxsol</u>

Oral drops 10%......3.78

✓ Coloxyl

30 ml OP

Subsidy		Fully	Brand or	
(Manufacturer's Price) 5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

Opioid Receptor Antagonists - Peripheral

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Either:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

Osmotic Laxatives

GLYCEROL			
* Suppos 3.6 g - Only on a prescription	9.25	20	✓ <u>PSM</u>
LACTULOSE - Only on a prescription			
* Oral liq 10 g per 15 ml	3.18	500 ml	✓ Laevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM B	ICARBONATE AN	ID SODIUM C	CHLORIDE
Powder for oral soln 13.125 g with potassium chloride 46.6			
sodium bicarbonate 178.5 mg and sodium chloride 350	0.	30	✓ Molaxole
SODIUM ACID PHOSPHATE – Only on a prescription	3		
Enema 16% with sodium phosphate 8%	2 50	1	✓ Fleet Phosphate
Enema 1076 with sodium phosphate 076	2.50	•	Enema
CODILIM CITRATE MITH CODILIM LAUDVI CHI DUOACETAT	C Only on a nya	aarintian	2.10.114
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETAT	, ,	scription	
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per m		50	/ Missiste
5 ml	26.72	50	✓ Micolette
Stimulant Laxatives			
Stilliulalit Laxatives			
BISACODYL - Only on a prescription			
* Tab 5 mg	5.99	200	✓ Lax-Tab
* Suppos 10 mg	3.74	10	✓ Lax-Suppositories
SENNA – Only on a prescription			
* Tab, standardised	2.17	100	
,	(6.84)		Senokot
	0.43	20	
	(1.72)		Senokot

Metabolic Disorder Agents

ALGLUCOSIDASE ALFA - Special Authority see SA1622 or	n the next page - Retail	pharmacy	
Inj 50 mg vial	1,142.60	1	✓ Myozyme

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

⇒SA1622 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT: and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

BETAINE - Special Authority see SA1727 below - Retail pharmacy

⇒SA1727 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE – Special Authority see SA1593 on the next page – Retail pharmacy
Inj 1 mg per ml, 5 ml vial.......2,234.00 1 ✓ Naglazyme

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1593 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI: and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
 \$	Per		Manufacturer	

continued...

laronidase would be bridging treatment to transplant; and

- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

SAPROPTERIN DIHYDROCHLORIDE – Special Authority see SA1757 below – Retail pharmacy
Tab soluble 100 mg.......1,452.70 30 OP
✓ Kuvan

⇒SA1757 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

(Manufacturer's Pri \$

⇒SA1598 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

IMIGLUCERASE - Special Authority see SA0473 below - Retail pharmacy

(Cerezyme Inj 40 iu per ml, 200 iu vial to be delisted 1 March 2019) (Cerezyme Inj 40 iu per ml, 400 iu vial to be delisted 1 March 2019)

⇒SA0473 Special Authority for Subsidy

Special Authority approved by the Gaucher's Treatment Panel

Notes: Subject to a budgetary cap. Applications will be considered and approved subject to funding availability.

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The Co-ordinator, Gaucher's Treatment Panel Phone: (04) 460 4990 PHARMAC. PO Box 10 254 Facsimile: (04) 916 7571

Wellington Email: gaucherpanel@pharmac.govt.nz

TALIGLUCERASE ALFA - Special Authority see SA1734 below - Retail pharmacy

Elelyso to be Sole Supply on 1 March 2019

⇒SA1734 Special Authority for Subsidy

Special Authority approved by the Gaucher's Treatment Panel

Notes: Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The Co-ordinator, Gaucher's Treatment Panel Phone: 04 460 4990 PHARMAC PO Box 10 254 Facsimile: 04 916 7571

Wellington Email: gaucherpanel@pharmac.govt.nz

Completed application forms must be sent to the coordinator for Gaucher's Treatment Panel and will be considered by Gaucher's Treatment Panel at the next practicable opportunity.

Notification of Gaucher's Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Access Criteria

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and
- 3) Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, serum glucosylsphingosine, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and

Subsidy		ully Brand or	
(Manufacturer's \$	Price) Subsidis Per	sed Generic Manufacturer	

continued...

- 5) Any of the following:
- 1) Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia, thrombocytopenia, at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or massive symptomatic splenomegaly; or
 - 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g. hips or shoulder); spontaneous fractures or vertebral collapse; chronic bone pain not controlled by other pharmaceuticals; or
 - 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 4) Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease: or
 - 5) Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1) Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated: and
- 2) Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size: and
- 3) Radiological (MRI) signs of bone activity performed at one year and two years since initiation of treatment begins, and two to three yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4) Serum glucosylsphingosine levels taken at least 6 to 12 monthly show a decrease compared with baseline; and
- 5) Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 6) Patient has not developed another medical condition that might reasonably be expected to compromise a response to
- 7) Patient is compliant with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- 8) Supporting clinical information including test reports, MRI whole body STIR, serum glucosylsphingosine, haematological data, and other relevant investigations are submitted to the Gaucher Panel for assessment as required.

Mouth and Throat

Agents Used in Mouth Ulceration

BENZYDAMINE HYDROCHLORIDE

Soln 0.15% - Higher subsidy of \$20.31 per 500 ml with			
Endorsement	9.00	500 ml	
	(20.31)		Difflam

Additional subsidy by endorsement for a patient who has oral mucositis as a result of treatment for cancer, and the prescription is endorsed accordingly.

CARMELLOSE SODIUM WITH GELATIN AND PECTIN

Paste	17.20	56 g OP	Stomahesive
	4.55	15 g OP	
	(7.90)		Orabase
	1.52	5 g OP	
	(3.60)	•	Orabase
Powder	8.48	28 g OP	
	(10.95)	•	Stomahesive

			<u> </u>
	Subsidy (Manufacturer's P \$	rice) Subsi Per	Fully Brand or dised Generic ✓ Manufacturer
HLORHEXIDINE GLUCONATE			
Mouthwash 0.2%	2.57	200 ml OP	✓ healthE
HOLINE SALICYLATE WITH CETALKONIUM CHLORIDE			
Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06 (6.00)	15 g OP	Bonjela
RIAMCINOLONE ACETONIDE	(0.00)		Donjoid
Paste 0.1%	5.33	5 g OP	✓ Kenalog in Orabase
Oropharyngeal Anti-infectives			
MPHOTERICIN B			
Lozenges 10 mg	5.86	20	✓ Fungilin
NICONAZOLE			
Oral gel 20 mg per g	4.74	40 g OP	✓ <u>Decozol</u>
IYSTATIN Oral liq 100,000 u per ml	1.05	24 ml OP	✓ Nilstat
Oral liq 100,000 u per mi	1.95	24 IIII OP	Niistat
Other Oral Agents			
or folinic mouthwash, pilocarpine oral liquid or saliva substitute	formula refer Sta	ndard Formulae	e, page 228
YDROGEN PEROXIDE			
Soln 3% (10 vol) – Maximum of 200 ml per prescription	1.40	100 ml	✓ Pharmacy Health
HYMOL GLYCERIN ← Compound, BPC	0.15	500 ml	✓ PSM
Compound, BFC	9.15	300 1111	▼ <u>FSWI</u>
Vitamins			
Vitamin A			
ITAMIN A WITH VITAMINS D AND C			
Soln 1000 u with Vitamin D 400 u and ascorbic acid 30 mg	per		
10 drops		10 ml OP	✓ Vitadol C
Vitadol C Soln 1000 u with Vitamin D 400 u and ascorbic acid 3	10 mg per 10 arop	s to be delisted	1 August 2019)
Vitamin B			
IYDROXOCOBALAMIN			
Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a P	SO1.89	3	✓ Neo-B12
YRIDOXINE HYDROCHLORIDE			
a) No more than 100 mg per dose			
 b) Only on a prescription Tab 25 mg – No patient co-payment payable 	2.70	90	✓ Vitamin B6 25
F Tab 50 mg		500	✓ Apo-Pyridoxine
HIAMINE HYDROCHLORIDE – Only on a prescription			
€ Tab 50 mg	4.89	100	✓ Max Health
TTAMIN B COMPLEX			
Fab, strong, BPC	7.15	500	✓ Bplex

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

Vitamin C

ASCORBIC ACID

- a) No more than 100 mg per dose
- b) Only on a prescription

* 1ab 100 mg8.10 500 • CVI	*	Tab 100 mg8.1	10 5	500	Cvite
----------------------------	---	---------------	------	-----	-------

Vitamin D

ALFACALCIDOL			
* Cap 0.25 mcg	26.32	100	✓ One-Alpha
* Cap 1 mcg	87.98	100	✓ One-Alpha
* Oral drops 2 mcg per ml	60.68	20 ml OP	✓ One-Alpha
CALCITRIOL			
* Cap 0.25 mcg	9.95	100	✓ Calcitriol-AFT
* Cap 0.5 mcg	18.39	100	✓ Calcitriol-AFT
COLECALCIFEROL			
* Cap 1.25 mg (50,000 iu) - Maximum of 12 cap per	prescription2.50	12	✓ Vit.D3
* Oral lig 188 mcg per ml (7,500 iu per ml)	9.00	4.8 ml OP	✓ Puria

Multivitamin Preparations

ΜL	JLTIVITAMIN RENAL – Special Authority see SA1546 below – Retail pharr	nacy		
*	Cap	30	•	Clinicians Renal Vit

⇒SA1546 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or
- 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73 m² body surface area (BSA).

MULTIVITAMINS - Special Authority see SA1036 below - Retail pharmacy

*	Powder	 	72.00	200	OP (1	Paediatric Seravit

⇒SA1036 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins.

VITAMINIS

VI	AWIIIO		
*	Tab (BPC cap strength)10.50	1,000	✓ Mvite
*	Cap (fat soluble vitamins A, D, E, K) - Special Authority see		
	SA1720 below – Retail pharmacy	60	✓ Vitabdeck

⇒SA1720 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency; or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

Fully

Brand or

Subsidy

	(Manufacturer's Price) \$	Su Per		eneric anufacturer
Minerals				
Calcium				
CALCIUM CARBONATE * Tab eff 1.75 g (1 g elemental) * Tab 1.25 g (500 mg elemental)	7.52	10 250	✓ Calso ✓ <u>Arro</u>	ource w-Calcium
* Inj 10%, 10 ml ampoule to be delisted 1 July 2019)	34.24 64.00	10 20	✓ Hosp ✓ Max	oira Health ^{S29}
Fluoride				
SODIUM FLUORIDE * Tab 1.1 mg (0.5 mg elemental)	5.75	100	✓ PSM	
lodine				
POTASSIUM IODATE * Tab 253 mcg (150 mcg elemental iodine) NeuroTabs to be Sole Supply on 1 April 2019	4.69	90	✓ Neur	oTabs
Iron				
FERRIC CARBOXYMALTOSE - Special Authority see SA1675 b Inj 50 mg per ml, 10 ml		acy 1	✓ Ferir	iject

⇒SA1675 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any medical practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any medical practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	
continued			

EEDDOLIO ELIMADATE

- 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
- 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
- 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
- 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

FERROUS FUMARATE			
* Tab 200 mg (65 mg elemental)	3.09	100	✓ Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID			
* Tab 310 mg (100 mg elemental) with folic acid 350 mcg	4.68	60	✓ Ferro-F-Tabs
FERROUS SULPHATE			
* Tab long-acting 325 mg (105 mg elemental)	2.06	30	✓ Ferrograd
* Oral liq 30 mg (6 mg elemental) per 1 ml	.10.80	500 ml	✓ Ferodan
IRON POLYMALTOSE			
* Inj 50 mg per ml, 2 ml ampoule	.15.22	5	✓ Ferrum H
	34.50		✓ Ferrosig
(Ferrum H Inj 50 mg per ml, 2 ml ampoule to be delisted 1 July 2019)			-
Magnesium			
magnesium			
For magnesium hydroxide mixture refer Standard Formulae, page 228			
MAGNESIUM SULPHATE			
* Ini 2 mmol per ml. 5 ml ampoule	.10.21	10	✓ DBL

Zinc		
ZINC SULPHATE		
* Cap 137.4 mg (50 mg elemental) 11.00	100	✓ Zincaps

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus: and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus: and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*: and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Generic
EPOETIN ALFA - Special Authority see SA1775 on the previous	s page – Retail pharm	асу		
Wastage claimable			_	_
Inj 1,000 iu in 0.5 ml, syringe		6		Eprex
	250.00			Binocrit
Inj 2,000 iu in 0.5 ml, syringe		6		Eprex
Inj 2,000 iu in 1 ml, syringe		6		Binocrit
Inj 3,000 iu in 0.3 ml, syringe		6		Binocrit
	166.87			Eprex
Inj 4,000 iu in 0.4 ml, syringe	96.50	6		Binocrit
	193.13		1	Eprex
Inj 5,000 iu in 0.5 ml, syringe	125.00	6	✓	Binocrit
	243.26		1	Eprex
Inj 6,000 iu in 0.6 ml, syringe	145.00	6		Binocrit
	291.92		✓	Eprex
Inj 8,000 iu in 0.8 ml, syringe	175.00	6	✓	Binocrit
	352.69		✓	Eprex
Inj 10,000 iu in 1 ml, syringe	197.50	6	✓	Binocrit
	395.18		✓	Eprex
Inj 40,000 iu in 1 ml, syringe	250.00	1	✓	Binocrit
	263.45		✓	Eprex
(Eprex Inj 1,000 iu in 0.5 ml, syringe to be delisted 1 April 2019)				
(Eprex Inj 2,000 iu in 0.5 ml, syringe to be delisted 1 April 2019)				
(Eprex Inj 3,000 iu in 0.3 ml, syringe to be delisted 1 April 2019)				
(Eprex Inj 4,000 iu in 0.4 ml, syringe to be delisted 1 April 2019)				
(Eprex Inj 5,000 iu in 0.5 ml, syringe to be delisted 1 April 2019)				
(Eprex Inj 6,000 iu in 0.6 ml, syringe to be delisted 1 April 2019)				
(Eprex Inj 8,000 iu in 0.8 ml, syringe to be delisted 1 April 2019)				
(Eprex Inj 10,000 iu in 1 ml, syringe to be delisted 1 April 2019)				
(Eprex Inj 40,000 iu in 1 ml, syringe to be delisted 1 April 2019)				

Megaloblastic

FΟ	LIC	ACI	D

*	Tab 0.8 mg21.84	1,000	1	Apo-Folic Acid
*	Tab 5 mg	500	1	Apo-Folic Acid
	Oral lig 50 mcg per ml	25 ml OP	1	Biomed

Antifibrinolytics, Haemostatics and Local Sclerosants

ELTROMBOPAG - Special Authority see SA1743	below – Retail pharmacy		
Wastage claimable			
Tab 25 mg	1,550.00	28	✓ Revolade
Tab 50 mg	3,100.00	28	Revolade

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and

Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic
 \$	Per 🗸	Manufacturer

continued...

- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding: or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (**idiopathic thrombocytopenic purpura - preparation for splenectomy**) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopage treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Fither:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter: or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

4 7...

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
EPTACOG ALFA [RECOMBINANT FACTOR VIIA]	- [Xpharm]			
For patients with haemophilia, whose funded tr	eatment is managed by the Haemo	philia	a Treaters (Group in conjunction with
the National Haemophilia Management Group.	• •			
Inj 1 mg syringe	1,178.30	1	✓	NovoSeven RT
Inj 2 mg syringe	2,356.60	1	✓	NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓	NovoSeven RT
Inj 8 mg syringe	9,426.40	1	✓	NovoSeven RT
ACTOR EIGHT INHIBITOR BYPASSING FRACTI	ON – [Xnharm]			
For patients with haemophilia, whose funded tr		nhilis	Troatore (Group in conjunction wit
the National Haemophilia Management Group.	eathern is managed by the rideme	prime	i i i oatoio i	aroup in conjunction wit
Inj 500 U	1 450 00	1	1	FEIBA NF
Inj 1,000 U	•	1		FEIBA NF
Inj 2,500 U	•	1		FEIBA NF
IOROCTOCOG ALFA [RECOMBINANT FACTOR	•	•	•	I LIDA III
Preferred Brand of recombinant factor VIII for p to funded treatment is managed by the Haemo				
Management Group.	prillia Treaters Group in Conjunction	ı wılı	i ine manoi	іаі паетіорііііа
Inj 250 iu prefilled syringe	210.00	1	1	Xyntha
Inj 500 iu prefilled syringe		1		Xyntha
Inj 1,000 iu prefilled syringe		1		Xyntha
Inj 2,000 iu prefilled syringe		1		Xyntha
		1		Xyntha
Inj 3,000 iu prefilled syringe		ı	•	Ayılıla
ONACOG ALFA [RECOMBINANT FACTOR IX] -				
For patients with haemophilia, whose funded tr		philia	a Treaters (Group in conjunction wit
the National Haemophilia Management Group.			_	
Inj 250 iu vial		1		BeneFIX
Inj 500 iu vial		1		BeneFIX
Inj 1,000 iu vial	•	1		BeneFIX
Inj 2,000 iu vial	2,480.00	1		BeneFIX
Inj 3,000 iu vial	3,720.00	1	✓	BeneFIX
ONACOG GAMMA, [RECOMBINANT FACTOR IX	K] – [Xpharm]			
For patients with haemophilia, whose funded tr		philia	a Treaters (Group in conjunction with
the National Haemophilia Management Group.		p		p
Inj 250 iu vial		1	/	RIXUBIS
			٠,	DIVIDIO

Inj 500 iu vial......575.00

Inj 2,000 iu vial......2,300.00

Inj 3,000 iu vial......3,450.00

1

1

1

✓ RIXUBIS

✓ RIXUBIS

✓ RIXUBIS

✓ RIXUBIS

	BLOOD AND	BLC	OD FOR	MING ORGANS
	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVA Rare Clinical Circumstances Brand of recombinant fa 28 February 2019. Access to funded treatment by ap be obtained from PHARMAC's website http://www.ph	ctor VIII for patients with happlication to the Haemophil	aemoph ia Treat	nilia from 1 N	March 2016 until el. Application details may
The Co-ordinator, Haemophilia Treatments Panel	Phone: 0800 023 588	Ontion	2	
PHARMAC PO Box 10 254	Facsimile: (04) 974 488		_	
Wellington	Email: haemophilia@ph		govt.nz	
·				
Inj 250 iu vial	287.50	1	✓	Advate
Inj 500 iu vial		1	✓	Advate
Inj 1,000 iu vial	1,150.00	1	✓	Advate
Inj 1,500 iu vial	1,725.00	1	✓	Advate
Inj 2,000 iu vial	2,300.00	1	√	Advate
Inj 3,000 iu vial	3,450.00	1	✓	Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGI	ENATE FS) - [Xpharm]			
Second Brand of recombinant factor VIII for patients v		arch 20	16 until 28 F	ebruary 2019. Access to
funded treatment by application to the Haemophilia T PHARMAC's website http://www.pharmac.govt.nz or:	reatments Panel. Applicat			
The Co-ordinator, Haemophilia Treatments Panel	Phone: 0800 023 588	Option	2	
PHARMAC PO Box 10 254	Facsimile: (04) 974 488		_	
Wellington	Email: haemophilia@ph		govt nz	
Weilington	шаш. <u>паеттортина егрг</u>	aiiiiau.	govi.nz	
Inj 250 iu vial	227 50	1	./ 1	Cogenate FS
Inj 500 iu vial		1		Kogenate FS
Inj 1,000 iu vial		1		Kogenate FS
Inj 2,000 iu vial		1		Kogenate FS
Inj 3,000 iu vial		1		Cogenate FS
SODIUM TETRADECYL SULPHATE			• •	togenate i o
	00.50	_		
* Inj 3% 2 ml		5	-	Thro voin
	(73.00)		Г	Fibro-vein
TRANEXAMIC ACID	20.07	400		
Tab 500 mg	20.67	100	y <u>(</u>	<u>Cyklokapron</u>
Vitamin K				
PHYTOMENADIONE				
Inj 2 mg per 0.2 ml - Up to 5 inj available on a PSO	8.00	5	✓ k	Conakion MM
Inj 10 mg per ml, 1 ml - Up to 5 inj available on a PS		5	✓ k	Conakion MM
Antithrombotic Agents				
Antiplatelet Agents				
ASPIRIN				
* Tab 100 mg	12.50	990	✓ <u>E</u>	thics Aspirin EC
CLOPIDOGREL				
* Tab 75 mg	5.44	84	✓	Arrow - Clopid
DIDVDID ALIQUE			_	<u> </u>

60

✓ Pytazen SR

* Tab long-acting 150 mg......11.52

DIPYRIDAMOLE

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
PRASUGREL - Special Authority see SA1201 below - Retail ph	armacy				
Tab 5 mg	108.00	28	√ E	ffient	
Tab 10 mg	120.00	28	√ E	ffient	

⇒SA1201 Special Authority for Subsidy

Initial application — (coronary angioplasty and bare metal stent) from any relevant practitioner. Approvals valid for 6 months where the patient has undergone coronary angioplasty in the previous 4 weeks and is clopidogrel-allergic*.

Initial application — (drug eluting stent) from any relevant practitioner. Approvals valid for 12 months where the patient has had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Initial application — (stent thromobosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (coronary angioplasty and bare metal stent) from any relevant practitioner. Approvals valid for 6 months where the patient has undergone coronary angioplasty or had a bare metal cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Renewal — (drug eluting stent) from any relevant practitioner. Approvals valid for 12 months where had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Note: * Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

TICAGRELOR - Special Authority see SA1382 below - Retail pharmacy

⇒SA1382 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Heparin and Antagonist Preparations

DALTEPARIN SODIUM - Special Authority see SA1270 belo	w - Retail pharmacy		
Inj 2,500 iu per 0.2 ml prefilled syringe	19.97	10	✓ Fragmin
Inj 5,000 iu per 0.2 ml prefilled syringe	39.94	10	✓ Fragmin
Inj 7,500 iu per 0.75 ml graduated syringe		10	✓ Fragmin
Inj 10,000 iu per 1 ml graduated syringe	77.55	10	✓ Fragmin
Inj 12,500 iu per 0.5 ml prefilled syringe	99.96	10	✓ Fragmin
Inj 15,000 iu per 0.6 ml prefilled syringe	120.05	10	✓ Fragmin
Inj 18,000 iu per 0.72 ml prefilled syringe		10	✓ Fragmin
			-

⇒SA1270 Special Authority for Subsidy

Initial application — (Pregnancy or Malignancy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

1 Low molecular weight heparin treatment is required during a patient's pregnancy; or

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

2 For the treatment of venous thromboembolism where the patient has a malignancy.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (**Pregnancy or Malignancy**) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, Acute Coronary Syndrome, cardioversion, or prior to oral anti-coagulation).

ENOXAPARIN SODIUM - Special Authority see SA1646 below - Retail pharmacy

Inj 20 mg in 0.2 ml syringe	27.93	10	Clexane
Inj 40 mg in 0.4 ml syringe		10	Clexane
Inj 60 mg in 0.6 ml syringe		10	✓ Clexane
Inj 80 mg in 0.8 ml syringe		10	✓ Clexane
Inj 100 mg in 1 ml syringe		10	✓ Clexane
Inj 120 mg in 0.8 ml syringe		10	✓ Clexane
Inj 150 mg in 1 ml syringe		10	✓ Clexane

⇒SA1646 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy: or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

1 Low molecular weight heparin treatment is required during a patient's pregnancy; or

	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per	Manufacturer
continued			

2 For the treatment of venous thromboembolism where the patient has a malignancy; or

3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

HEPARIN SODIUM			
Inj 1,000 iu per ml, 5 ml ampoule	58.57	50	✓ Pfizer
Inj 5,000 iu per ml, 1 ml		5	✓ Hospira
Inj 5,000 iu per ml, 5 ml ampoule	203.68	50	✓ Pfizer
Inj 25,000 iu per ml, 0.2 ml	19.00	5	✓ Hospira
HEPARINISED SALINE			
Inj 10 iu per ml, 5 ml	53.40	30	✓ BD PosiFlush S29
	56.94	50	✓ Pfizer
(BD PosiFlush S29) Inj 10 iu per ml, 5 ml to be delisted 1 N	farch 2019)		

Oral Anticoagulants

DABIGATRAN

Cap 110 mg	76.36	60	✓ Pradaxa
Cap 150 mg	76.36	60	Pradaxa
RIVAROXABAN			
Tab 10 mg - No more than 1 tab per day	83.10	30	✓ Xarelto
Tab 15 mg	77.56	28	✓ Xarelto
Tab 20 mg	77.56	28	✓ Xarelto
WARFARIN SODIUM			
Note: Marevan and Coumadin are not interchangeable.			
* Tab 1 mg	3.46	50	Coumadin
•	6.86	100	Marevan
* Tab 2 mg	4.31	50	✓ Coumadin
* Tab 3 mg	9.70	100	Marevan
* Tab 5 mg	5.93	50	Coumadin
· ·	11.75	100	✓ Marevan

Blood Colony-stimulating Factors

FILGRASTIM – Special Authority see SA1259 below – Retail p	oharmacy		
Inj 300 mcg per 0.5 ml prefilled syringe	270.00	5	✓ Zarzio
Inj 480 mcg per 0.5 ml prefilled syringe	432.00	5	✓ Zarzio

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or

continued...

Pradaxa

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$ P	Per 🗸	Manufacturer

continued...

- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10⁹/L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1384 below - Retail pharmacy

⇒SA1384 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*). Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

GLUCOSE (DEXTROSE)

Intravenous Administration

GLUCUSE [DEXTRUSE]		
* Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO29.50	5	✓ Biomed
* Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO14.50	1	✓ Biomed
POTASSIUM CHLORIDE		
* Inj 75 mg per ml, 10 ml55.00	50	✓ AstraZeneca
SODIUM BICARBONATE		
Inj 8.4%, 50 ml19.95	1	✓ Biomed
a) Up to 5 inj available on a PSO		
b) Not in combination		
Inj 8.4%, 100 ml20.50	1	Biomed
a) Up to 5 inj available on a PSO		
b) Not in combination		

SODIUM CHLORIDE

Not funded for use as a nasal drop. Only funded for nebuliser use when in conjunction with an antibiotic intended for nebuliser use.

Inj 0.9%, bag - Up to 2000 ml available on a PSO	1.23	500 ml	✓ Baxter
	1.26	1.000 ml	✓ Baxter

Only if prescribed on a prescription for renal dialysis, maternity or post-natal care in the home of the patient, or on a PSO for emergency use. (500 ml and 1 000 ml packs)

Inj 23.4% (4 mmol/ml), 20 ml ampoule	33.00	5	✓ Biomed
For Sodium chloride oral liquid formulation refer Sta	ndard Formulae, page	228	
Inj 0.9%, 5 ml ampoule - Up to 5 inj available on a PSC	7.00	50	✓ InterPharma
			✓ Multichem
Inj 0.9%, 10 ml ampoule - Up to 5 inj available on a PS	O6.63	50	✓ Pfizer
Inj 0.9%, 20 ml ampoule	5.00	20	✓ Multichem
,	7.50	30	✓ InterPharma
TOTAL PARENTERAL NUTRITION (TPN) - Retail pharmac	cy-Specialist		
Infusion	CBS	1 OP	✓ TPN

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

WATER

- 1) On a prescription or Practitioner's Supply Order only when on the same form as an injection listed in the Pharmaceutical Schedule requiring a solvent or diluent; or
- 2) On a bulk supply order; or
- 3) When used in the extemporaneous compounding of eye drops; or
- 4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.

Inj 5 ml ampoule - Up to 5 inj available on a PSO7.00	50	✓ InterPharma
Inj 10 ml ampoule – Up to 5 inj available on a PSO6.63	50	✓ Pfizer
Inj 20 ml ampoule - Up to 5 inj available on a PSO5.00	20	Multichem
7.50	30	✓ InterPharma

Oral Administration		
CALCIUM POLYSTYRENE SULPHONATE	300 g OP	✓ Calcium Resonium
Powder for oral soln — Up to 10 sach available on a PSO2.30	10	✓ Enerlyte
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml)6.55	1,000 ml OP	✓ Pedialyte - Bubblegum
PHOSPHORUS Tab eff 500 mg (16 mmol)82.50	100	✓ Phosphate Phebra ✓ Phosphate-Sandoz
(Phosphate-Sandoz Tab eff 500 mg (16 mmol) to be delisted 1 May 2019) POTASSIUM CHLORIDE		
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)5.26 (11.85)	60	Chlorvescent
* Tab long-acting 600 mg (8 mmol)8.90	200	✓ Span-K
SODIUM BICARBONATE Cap 840 mg8.52	100	✓ Sodibic ✓ Sodibic
SODIUM POLYSTYRENE SULPHONATE		
Powder	454 g OP	✓ Resonium-A

Subsidy
(Manufacturer's Price) Subs

Fully Subsidised Brand or Generic Manufacturer

Alpha-Adrenoceptor Blockers

Alpha Adrenoceptor Blockers

DOXAZOSIN	500	✓ Apo-Doxazosin
* Tab 4 mg	500	✓ Apo-Doxazosin
PHENOXYBENZAMINE HYDROCHLORIDE		
* Cap 10 mg65.00	30	✓ BNM S29
216.67	100	✓ Dibenzyline S29
PRAZOSIN		
* Tab 1 mg5.53	100	✓ Apo-Prazosin
* Tab 2 mg7.00	100	✓ Apo-Prazosin
* Tab 5 mg11.70	100	✓ Apo-Prazosin
TERAZOSIN		
* Tab 1 mg0.59	28	✓ Actavis
* Tab 2 mg7.50	500	✓ Apo-Terazosin
* Tab 5 mg10.90	500	✓ Apo-Terazosin

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

CARTORRII	
CAPTOPRIL	

the Court of the C		
* Oral liq 5 mg per ml		Capoten
Oral liquid restricted to children under 12 years of ac	je.	
CILAZAPRIL		
* Tab 0.5 mg	2.00 90	✓ Zapril
* Tab 2.5 mg		✓ Apo-Cilazapril
* Tab 5 mg		✓ Apo-Cilazapril
ENALAPRIL MALEATE		
	0.96 100	✓ Ethics Enalapril
: _ - · · · 9		•
* Tab 10 mg		✓ Ethics Enalapril
* Tab 20 mg	1.78 100	Ethics Enalapril
LISINOPRIL		
* Tab 5 mg	2.07 90	Ethics Lisinopril
* Tab 10 mg	2.36 90	✓ Ethics Lisinopril
* Tab 20 mg		✓ Ethics Lisinopril
PERINDOPRIL		
	3.75 30	✓ Apo-Perindopril
· · · · · · · · · · · · · · · · · · ·		
* Tab 4 mg	4.80 30	✓ Apo-Perindopril
QUINAPRIL		
* Tab 5 mg	6.01 90	 Arrow-Quinapril 5
* Tab 10 mg	3.16 90	✓ Arrow-Quinapril 10
* Tab 20 mg	4.89 90	✓ Arrow-Quinapril 20

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
ACE Inhibitors with Diuretics				
CILAZAPRIL WITH HYDROCHLOROTHIAZIDE * Tab 5 mg with hydrochlorothiazide 12.5 mg	10.18	100	✓ ,	Apo-Cilazapril/ Hydrochlorothiazide
QUINAPRIL WITH HYDROCHLOROTHIAZIDE * Tab 10 mg with hydrochlorothiazide 12.5 mg * Tab 20 mg with hydrochlorothiazide 12.5 mg		30 30	-	Accuretic 10 Accuretic 20
Angiotensin II Antagonists				
CANDESARTAN CILEXETIL * Tab 4 mg	2.28 3.67 6.39 1.39 1.63	90 90 90 90 84 84 84 84	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Candestar Candestar Candestar Candestar Losartan Actavis Losartan Actavis Losartan Actavis Losartan Actavis
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE Tab 50 mg with hydrochlorothiazide 12.5 mg	1.88	30	✓ ,	Arrow-Losartan & Hydrochlorothiazide

Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN - Special Authority see SA1751 below - Retail pharmacy

Note: Due to the angiotensin II receptor blocking activity of sacubitril with valsartan it should not be co-administered with an ACE inhibitor or another ARB.

Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/103

⇒SA1751 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II: or
 - 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Brand or

Generic

Manufacturer

Fully

Subsidised

Per

r lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, p	age 119	
MIODARONE HYDROCHLORIDE		4.4
Tab 100 mg - Retail pharmacy-Specialist	30	✓ Cordarone-X
Tab 200 mg - Retail pharmacy-Specialist	30 5	✓ Cordarone-X
Inj 50 mg per ml, 3 ml ampoule – Up to 5 inj available on a PSO9.98	5	✓ <u>Lodi</u>
ROPINE SULPHATE		
Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO12.07	10	✓ Martindale
	10	<u>Martindale</u>
GOXIN	0.40	(Lawrente BO
Tab 62.5 mcg – Up to 30 tab available on a PSO	240	✓ <u>Lanoxin PG</u>
Tab 250 mcg - Up to 30 tab available on a PSO	240 60 ml	✓ <u>Lanoxin</u>✓ Lanoxin
Oral liq 50 flicg per fill10.00	00 1111	✓ Lanoxin S29 S29
		V Lanoxiii 329 329
SOPYRAMIDE PHOSPHATE	400	/ Duthan alon
Cap 100 mg23.87	100	✓ Rythmodan
ECAINIDE ACETATE – Retail pharmacy-Specialist		. .
Tab 50 mg	60	✓ Tambocor
Cap long-acting 100 mg	30	✓ Tambocor CR
Cap long-acting 200 mg	30 5	✓ Tambocor CR✓ Tambocor
Inj 10 mg per ml, 15 ml ampoule	5	▼ Tallibocol
EXILETINE HYDROCHLORIDE	400	/ Marritation
Cap 150 mg162.00	100	Mexiletine Hydrochloride
		USP S29
Cap 250 mg202.00	100	✓ Mexiletine
		Hydrochloride USP \$29
OODAEENONE HYDDOCHI ODIDE Batail pharmagu Chaoialiat		001
ROPAFENONE HYDROCHLORIDE - Retail pharmacy-Specialist Tab 150 mg40.90	50	✓ Rytmonorm
140 100 mg	50	- riyunonomi
Antihypotensives		
DODRINE - Special Authority see SA1474 below - Retail pharmacy		
Tab 2.5 mg53.00	100	✓ Gutron
Tab 5 mg	100	✓ Gutron
SA1474 Special Authority for Subsidy		

Note: Treatment should be started with small doses and titrated upwards as necessary. Hypertension should be avoided, and

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is

Subsidy

(Manufacturer's Price)

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

49

the usual target is a standing systolic blood pressure of 90 mm Hg.

benefiting from treatment.

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

ATENOLOL			
* Tab 50 mg	4.26	500	✓ Mylan Atenolol
* Tab 100 mg	7.30	500	Mylan Atenolol
* Oral liq 25 mg per 5 ml	21.25	300 ml OP	Atenolol AFT
Restricted to children under 12 years of age.			
BISOPROLOL FUMARATE			
* Tab 2.5 mg	3.53	90	✓ Bosvate
* Tab 5 mg	5.15	90	✓ Bosvate
* Tab 10 mg	9.40	90	✓ Bosvate
CARVEDILOL			
* Tab 6.25 mg	2.24	60	✓ Carvedilol Sandoz
* Tab 12.5 mg		60	✓ Carvedilol Sandoz
* Tab 25 mg		60	✓ Carvedilol Sandoz
CELIPROLOL			
* Tab 200 mg	21.40	180	✓ Celol
LABETALOL		100	- 00101
* Tab 50 mg	9.00	100	✓ Hybloc
* Tab 50 mg		100	✓ Hybloc
		100	✓ Hybloc
· · · · · · · · · · · · · · · · · · ·		5	• пушос
* Inj 5 mg per ml, 20 ml ampoule	(88.60)	5	Trandate
(Hybloc Tab 50 mg to be delisted 1 August 2019) (Hybloc Tab 100 mg to be delisted 1 December 2019) (Hybloc Tab 200 mg to be delisted 1 February 2020)			
METOPROLOL SUCCINATE			
* Tab long-acting 23.75 mg	1.03	30	✓ Betaloc CR
* Tab long-acting 47.5 mg	1.25	30	✓ Betaloc CR
* Tab long-acting 95 mg		30	✓ Betaloc CR
* Tab long-acting 190 mg	3.00	30	✓ Betaloc CR
METOPROLOL TARTRATE			
* Tab 50 mg	5.66	100	✓ Apo-Metoprolol
* Tab 100 mg	7.55	60	✓ Apo-Metoprolol
* Tab long-acting 200 mg	23.40	28	✓ Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial	29.50	5	✓ Metroprolol IV
			<u>Mylan</u>
NADOLOL			
* Tab 40 mg	16.69	100	✓ Apo-Nadolol
* Tab 80 mg	26.43	100	✓ Apo-Nadolol
PINDOLOL			
* Tab 5 mg	13.22	100	✓ Apo-Pindolol
* Tab 10 mg		100	✓ Apo-Pindolol
* Tab 15 mg		100	✓ Apo-Pindolol
•			

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PROPRANOLOL				
* Tab 10 mg	4.64	100	✓	Apo-Propranolol
* Tab 40 mg	5.72	100	✓	Apo-Propranolol
* Cap long-acting 160 mg	18.17	100	1	Cardinol LA
* Oral lig 4 mg per ml - Special Authority see SA1327 below -				
Retail pharmacy		500 m	nl 🗸	Roxane \$29

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only): or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SO	TALOL		
*	Tab 80 mg39.53	500	✓ Mylan
	Tab 160 mg		✓ Mylan
TIM	OLOL		
*	Tab 10 mg10.55	100	✓ Apo-Timol

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

AMLODIPINE		
* Tab 2.5 mg	100	✓ Apo-Amlodipine
* Tab 5 mg	250	✓ Apo-Amlodipine
* Tab 10 mg4.40	250	✓ Apo-Amlodipine
FELODIPINE		
* Tab long-acting 2.5 mg	30	✓ Plendil ER
* Tab long-acting 5 mg	90	✓ Felo 5 ER
1.31	30	
(1.55)		Plendil ER
Felo 5 ER to be Sole Supply on 1 March 2019		
* Tab long-acting 10 mg4.32	90	✓ Felo 10 ER
1.44	30	
(2.30)		Plendil ER
Felo 10 ER to be Sole Supply on 1 March 2019		
(Plendil ER Tab long-acting 5 mg to be delisted 1 March 2019)		
(Plendil ER Tab long-acting 10 mg to be delisted 1 March 2019)		
NIFEDIPINE		
* Tab long-acting 10 mg10.63	60	✓ Adalat 10
		✓ Adefin S29
* Tab long-acting 20 mg9.59	100	✓ Nyefax Retard
* Tab long-acting 20 mg	30	✓ Adalat Oros
Tab long-acting 50 mg	30	✓ Adefin XL
* Tab long-acting 60 mg5.67	30	✓ Adeliii AL
Tab long adding od mg	30	✓ Adefin XL
		- AUCIIII AL

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

^{*}Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	
Other Calcium Channel Blockers	Ψ	1 61		Wandacturer
ILTIAZEM HYDROCHLORIDE				
F Tab 30 mg	4.60	100	1	Dilzem
F Tab 60 mg	8.50	100	✓	Dilzem
Cap long-acting 120 mg	33.42	500	✓	Apo-Diltiazem CD
Cap long-acting 180 mg		500	✓	Apo-Diltiazem CD
Cap long-acting 240 mg	66.76	500	✓	Apo-Diltiazem CD
ERHEXILINE MALEATE	62 00	100	1	Pexsig
Ç	02.90	100	•	rexsig
ERAPAMIL HYDROCHLORIDE				
F Tab 40 mg		100	_	Isoptin
F Tab 80 mg		100	_	Isoptin
Tab long-acting 120 mg		250	_	Verpamil SR
Tab long-acting 240 mg		250	/	Verpamil SR
Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO		5	/	Isoptin
				•
Centrally-Acting Agents				
LONIDINE				
Patch 2.5 mg, 100 mcg per day - Only on a prescription	7.40	4	/	<u>Mylan</u>
Patch 5 mg, 200 mcg per day – Only on a prescription		4		Mylan
Patch 7.5 mg, 300 mcg per day – Only on a prescription		4	_	Mylan
LONIDINE HYDROCHLORIDE		·		<u>,</u>
	0.75	112		Clanidina PMM
Tab 25 mcg		100		Clonidine BNM Catapres
: Tab 150 mcg : Inj 150 mcg per ml, 1 ml ampoule		100	_	Medsurge
	25.90	10	•	<u>weusurge</u>
ETHYLDOPA				
Tab 250 mg	15.10	100		Methyldopa Mylan
Diuretics				
Loop Diuretics				
UMETANIDE				
÷ Tab 1 mg	16.36	100	✓	Burinex
Inj 500 mcg per ml, 4 ml vial	7.95	5	✓	Burinex
JROSEMIDE [FRUSEMIDE]				
Tab 40 mg - Up to 30 tab available on a PSO	8.00	1,000	/	Diurin 40
Tab 500 mg		50		Urex Forte
Urex Forte to be Sole Supply on 1 April 2019				
Oral liq 10 mg per ml	10.66	30 ml C)P 🗸	Lasix
Inj 10 mg per ml, 25 ml ampoule		6		Lasix
Finj 10 mg per ml, 2 ml ampoule - Up to 5 inj available on a		5		Frusemide-Claris
Potassium Sparing Diuretics				
MILORIDE HYDROCHLORIDE				
Oral lig 1 mg per ml	30.00	25 ml C)P 🗸	Biomed
1 91				

		CARDIC	VASC	ULAR SYSTEM
	Subsidy (Manufacturer's Price \$) Sub	Fully sidised	Brand or Generic Manufacturer
EPLERENONE - Special Authority see SA1728 below - Retail pt Tab 50 mg Tab 25 mg >>SA1728 Special Authority for Subsidy	17.00 11.87	30 30	✓ <u>Ir</u>	nspra nspra
Initial application from any relevant practitioner. Approvals valid the following criteria: Both:		ewai unies	ss notitie	a for applications meeting
 1 Patient has heart failure with ejection fraction less than 40% 2 Either: 2.1 Patient is intolerant to optimal dosing of spironolacte 				
2.2 Patient has experienced a clinically significant adve METOLAZONE	rse effect while on	optimal do	sing of s	spironolactone.
Tab 5 mg	CBS	1		letolazone S29
SPIRONOLACTONE		50	✓ Z	aroxolyn S29
* Tab 25 mg * Tab 100 mg		100 100		piractin piractin
Oral liq 5 mg per ml		25 ml OP		iomed
Potassium Sparing Combination Diuretics				
AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE * Tab 5 mg with furosemide 40 mg		28	√ F	rumil
* Tab 5 mg with hydrochlorothiazide 50 mg		50	✓ N	loduretic
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE] * Tab 2.5 mg – Up to 150 tab available on a PSO	12.50	500	✓ <u>A</u>	rrow- Bendrofluazide
May be supplied on a PSO for reasons other than emerge * Tab 5 mg	•	500	✓ <u>A</u>	<u>rrow-</u> <u>Bendrofluazide</u>
CHLOROTHIAZIDE Oral liq 50 mg per ml	26.00	25 ml OP	✓ B	iomed
* Tab 25 mgINDAPAMIDE	8.00	50	✓ H	lygroton
* Tab 2.5 mg	2.60	90	✓ <u>D</u>	apa-Tabs
Lipid-Modifying Agents				
Fibrates				
BEZAFIBRATE * Tab 200 mg * Tab long-acting 400 mg		90 30		ezalip ezalip Retard

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
GEMFIBROZIL * Tab 600 mg	19.56	60	√ <u>I</u>	_ipazil
Other Lipid-Modifying Agents				
ACIPIMOX ★ Cap 250 mg	18.75	30	✓ (Dibetam
★ Tab 50 mg ★ Tab 500 mg		100 100	_	Apo-Nicotinic Acid Apo-Nicotinic Acid
Resins				
CHOLESTYRAMINE Powder for oral liq 4 g	19.25 (52.68) (52.68)	50		Questran-Lite Questran-Lite S29 ⁸²⁹
Questran-Lite Powder for oral liq 4 g to be delisted 1 June 20 Questran-Lite S29 S29 Powder for oral liq 4 g to be delisted COLESTIPOL HYDROCHLORIDE	,			
Grans for oral liq 5 g	28.60	30	✓ (Colestid

Prescribing Guidelines

Treatment with HMG CoA Reductase Inhibitors (statins) is recommended for patients with dyslipidaemia and an absolute 5 year cardiovascular risk of 15% or greater.

ATORVASTATINI - Sec	prescribing guideline above

* Tab 10 mg	6.96	500	✓ Lorstat
* Tab 20 mg	9.99	500	✓ Lorstat
* Tab 40 mg	15.93	500	✓ Lorstat
* Tab 80 mg		500	✓ Lorstat
PRAVASTATIN - See prescribing guideline above			
* Tab 20 mg	4.72	100	✓ Apo-Pravastatin
* Tab 40 mg		100	✓ Apo-Pravastatin
SIMVASTATIN - See prescribing guideline above			
* Tab 10 mg	0.95	90	 Simvastatin Mylan
* Tab 20 mg	1.52	90	✓ Simvastatin Mylan
* Tab 40 mg	2.63	90	✓ Simvastatin Mylan
* Tab 80 mg	6.00	90	✓ Simvastatin Mylan

Selective Cholesterol Absorption Inhibitors

	: HivibE – Special Authority see SA1045 below – Retail pharmacy			
*	Tab 10 mg	2.00	30	 Ezetimibe Sandoz

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
\$	Per	✓	Manufacturer

continued

All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 x normal) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 below - Retail pharmacy

Tab 10 mg with simvastatin 10 mg5.15	30	Zimybe
Tab 10 mg with simvastatin 20 mg	30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg7.15	30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg8.15	30	✓ Zimybe

⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

GL	YCERYL TRINITRATE		
*	Tab 600 mcg - Up to 100 tab available on a PSO8.00	100 OP	Lycinate
*	Oral pump spray, 400 mcg per dose - Up to 250 dose		
	available on a PSO4.45	250 dose OP	✓ Nitrolingual Pump Spray
*	Oral spray, 400 mcg per dose - Up to 200 dose available on a		
	PSO4.45	200 dose OP	✓ Glytrin
*	Patch 25 mg, 5 mg per day15.73	30	✓ Nitroderm TTS
*	Patch 50 mg, 10 mg per day	30	✓ Nitroderm TTS
(L)	cinate Tab 600 mcg to be delisted 1 March 2019)		

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	•	Manufacturer
SOSORBIDE MONONITRATE				
₭ Tab 20 mg	18.80	100	1	Ismo 20
★ Tab long-acting 40 mg	7.50	30	1	Ismo 40 Retard
Tab long-acting 60 mg	8.29	90	•	<u>Duride</u>
Sympathomimetics				
DRENALINE				
Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PSO	4 98	5	1	Aspen Adrenaline
ing this 1,000, this ampould op to only available on a too	5.25	Ü		Hospira
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PS		5		Hospira
, .,,	49.00	10		Aspen Adrenaline
OPRENALINE [ISOPROTERENOL]				•
Inj 200 mcg per ml, 1 ml ampoule	36.80	25		
.,g p, p	(164.20)			Isuprel
	,			'
/asodilators				
/DRALAZINE HYDROCHLORIDE				
Tab 25 mg - Special Authority see SA1321 below - Retail			_	
pharmacy	CBS	1	/	Hydralazine
		56	•	Onelink S29
		84	1	AMDIPHARM \$29
		100	1	Onelink S29
Inj 20 mg ampoule	25.90	5	✓	Apresoline
SA1321 Special Authority for Subsidy				
itial application from any relevant practitioner. Approvals valid	without further rene	wal u	nless notif	ied for applications mee
e following criteria:				
ither:				
1 For the treatment of refractory hypertension; or				
For the treatment of refractory hypertension; or For the treatment of heart failure in combination with a nitral	ate, in patients who a	are in	tolerant or	have not responded to
1 For the treatment of refractory hypertension; or	ate, in patients who a	are in	tolerant or	have not responded to
For the treatment of refractory hypertension; or For the treatment of heart failure in combination with a nitral inhibitors and/or angiotensin receptor blockers. INOXIDIL				·
For the treatment of refractory hypertension; or For the treatment of heart failure in combination with a nitral inhibitors and/or angiotensin receptor blockers. INOXIDIL		are in		have not responded to
For the treatment of refractory hypertension; or For the treatment of heart failure in combination with a nitral inhibitors and/or angiotensin receptor blockers. INOXIDIL				·
For the treatment of refractory hypertension; or For the treatment of heart failure in combination with a nitral inhibitors and/or angiotensin receptor blockers. INOXIDIL Tab 10 mg	70.00		<i>y</i>	Loniten
For the treatment of refractory hypertension; or For the treatment of heart failure in combination with a nitral inhibitors and/or angiotensin receptor blockers. INOXIDIL Tab 10 mg	70.00 27.95	100	<i>y</i>	Loniten
1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitral inhibitors and/or angiotensin receptor blockers. IINOXIDIL Tab 10 mg Tab 10 mg Tab 10 mg Tab 20 mg	70.00 27.95	100	<i>y</i>	Loniten
1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitral inhibitors and/or angiotensin receptor blockers. IINOXIDIL 1 Tab 10 mg 1 Tab 10 mg 1 Tab 20 mg APAVERINE HYDROCHLORIDE	70.00 27.95 33.28	100	<i>y y y</i>	Loniten
1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitral inhibitors and/or angiotensin receptor blockers. INOXIDIL Tab 10 mg Tab 10 mg Tab 20 mg APAVERINE HYDROCHLORIDE In 12 mg per ml, 10 ml ampoule	70.00 27.95 33.28	100 60 60	<i>y y y</i>	Loniten Ikorel Ikorel
1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitral inhibitors and/or angiotensin receptor blockers. INOXIDIL Tab 10 mg Tab 10 mg Tab 20 mg APAVERINE HYDROCHLORIDE In 12 mg per ml, 10 ml ampoule	70.00 27.95 33.28 217.90	100 60 60	\rightarrow \right	Loniten Ikorel Ikorel
1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitra inhibitors and/or angiotensin receptor blockers. INOXIDIL 1 Tab 10 mg 1 Tab 10 mg 1 Tab 20 mg 1 Tab 30 mg 1 Tab 400 mg 1 Tab 400 mg 1 Tab 400 mg 1 Tab 400 mg	70.00 27.95 33.28 217.90	100 60 60 5	\rightarrow \right	Loniten Ikorel Ikorel Hospira
1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitral inhibitors and/or angiotensin receptor blockers. INOXIDIL 1 Tab 10 mg 1 Tab 10 mg 1 Tab 20 mg 1 Tab 3 mg Tab 20 mg 1 Tab 400 mg	70.00 27.95 33.28 217.90 42.26	100 60 60 5	\rightarrow \right	Loniten Ikorel Ikorel Hospira
2 For the treatment of heart failure in combination with a nitral inhibitors and/or angiotensin receptor blockers. IINOXIDIL Tab 10 mg Tab 10 mg Tab 20 mg APAVERINE HYDROCHLORIDE Inj 12 mg per ml, 10 ml ampoule ENTOXIFYLLINE [OXPENTIFYLLINE]	70.0027.9533.28217.9042.26 e – Retail pharmacy	100 60 60 5	<i>y y y y</i>	Loniten Ikorel Ikorel Hospira

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1702 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The Coordinator, PAH Panel

PHARMAC. PO Box 10-254. WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz
BOSENTAN – Special Authority see SA1712 below – Retail pharmacy

Reddy's ✓ Bosentan-Mylan

Bosentan Dr Reddy's to be Sole Supply on 1 March 2019

Tab 125 mg141.00 60

✓ Bosentan Dr Reddy's Bosentan-Mylan

Bosentan Dr Reddy's to be Sole Supply on 1 March 2019

(Bosentan-Mylan Tab 62.5 mg to be delisted 1 March 2019) (Bosentan-Mylan Tab 125 mg to be delisted 1 March 2019)

⇒SA1712 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

(401.79)

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II. III. or IV: and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Fither:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil: or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV: or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHAWHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised: or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHAWHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see SA1738 below – Retail pharn	nacy		
Tab 25 mg	0.64	4	✓ Vedafil
Tab 50 mg	0.64	4	✓ Vedafil
Tab 100 mg		12	✓ Vedafil

⇒SA1738 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II; or
 - 3.2 PAH is in NYHA/WHO functional class III; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)		osidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 3.3 PAH is in NYHA/WHO functional class IV; and
 - 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Fither
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Prostacyclin Analogues

EPOPROSTENOL - Special Authority see SA1696 below	- Retail pharmacy		
Inj 500 mcg vial	36.61	1	✓ Veletri
Inj 1.5 mg vial	73.21	1	✓ Veletri

⇒SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ILOPROST - Special Authority see SA1705 below - Retail pharmacy

Nebuliser soln 10 mcg per ml, 2 ml1,185.00 30 **✓ Ventavis**

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 88

ADAPALENE

- a) Maximum of 30 g per prescription
- h) Only on a prescription

b) Only on a prescription			
Crm 0.1%	22.89	30 g OP	Differin
Gel 0.1%	22.89	30 g OP	Differin
ISOTRETINOIN - Special Authority see SA1475 below - Retail p	harmacy		
Cap 5 mg	8.14	60	Oratane
Cap 10 mg	13.34	120	✓ Oratane
Can 20 mg	20.49	120	✓ Oratane

⇒SA1475 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
 - 3.2 Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
- Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

TRFTINOIN

Crm 0.5 mg per g - Maximum of 50 g per prescription13.90 50 q OP ✓ ReTrieve

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 88

HYDROGEN PEROXIDI

HYDROGEN PEROXIDE			
* Crm 1%	8.56	15 g OP	 Crystaderm
MUPIROCIN		J	,
Oint 2%	6.60	15 g OP	
	(9.26)	Ū	Bactroban

- a) Only on a prescription
- b) Not in combination

		L	DERIMATOLOGICALS
	Subsidy (Manufacturer's I \$	Price) Subs Per	Fully Brand or sidised Generic ✓ Manufacturer
SODIUM FUSIDATE [FUSIDIC ACID] Crm 2%	2.52	15 g OP	✓ DP Fusidic Acid Cream
a) Maximum of 15 g per prescription b) Only on a prescription c) Not in combination Oint 2% a) Maximum of 15 g per prescription b) Only on a prescription	3.45	15 g OP	✔ Foban
c) Not in combination SULFADIAZINE SILVER Crm 1%	10.80	50 g OP	✓ <u>Flamazine</u>
Antifungals Topical			
For systemic antifungals, refer to INFECTIONS, Antifungals, pa AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		5 ml OP	✓ <u>MycoNail</u>
a) Only on a prescription b) Not in combination Nail-soln 8%	5.72	7 ml OP	✓ Apo-Ciclopirox
CLOTRIMAZOLE ** Crm 1% a) Only on a prescription b) Not in combination	0.70	20 g OP	✓ <u>Clomazol</u>
Soln 1% a) Only on a prescription b) Not in combination	4.36 (7.55)	20 ml OP	Canesten
ECONAZOLE NITRATE Crm 1%	1.00 (7.48)	20 g OP	Pevaryl
b) Not in combination Foaming soln 1%, 10 ml sachets	9.89 (17.23)	3	Pevaryl

a) Only on a prescriptionb) Not in combination

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

DERMATOLOGICALS

	Subsidy (Manufacturer's Price)		Fully Brand or sidised Generic
	\$	Per	✓ Manufacturer
MICONAZOLE NITRATE			
* Crm 2%	0.74	15 g OP	✓ <u>Multichem</u>
a) Only on a prescription			
b) Not in combination * Lotn 2%	4 36	30 ml OP	
	(10.03)	00 1111 01	Daktarin
a) Only on a prescription	()		
b) Not in combination			
* Tinct 2%		30 ml OP	
	(12.10)		Daktarin
a) Only on a prescription			
b) Not in combination			
NYSTATIN Crm 100,000 u per g	1.00	15 g OP	
Citii 100,000 u pei g	(7.90)	13 y OF	Mycostatin
a) Only on a prescription	(7.00)		myoodaan
b) Not in combination			
Antipruritic Preparations			
CALAMINE			
a) Only on a prescription			
b) Not in combination			
Crm, aqueous, BP	1.26	100 g	✓ <u>healthE Calamine</u>
			Aqueous Cream
			<u>BP</u>
Lotn, BP	12.94	2,000 ml	✓ PSM
CROTAMITON			
a) Only on a prescription			
b) Not in combination Crm 10%	3 20	20 g OP	✓ Itch-Soothe
		20 y OF	· Iteli-200tile
MENTHOL – Only in combination	mulatam . Tarabaal O	41	Diain
 Only in combination with a dermatological base or pro With or without other dermatological galenicals. 	prietary Topical Cor	ticosteriod –	Plain
2) Will of willout other definationogical galeficals.			
Crystals	6.92	25 g	✓ MidWest
,	29.60	100 g	✓ MidWest
Corticosteroids Topical			
For systemic corticosteroids, refer to CORTICOSTEROIDS AND	RELATED AGENT	ΓS, page 78	
Corticosteroids - Plain			
BETAMETHASONE DIPROPIONATE			
Crm 0.05%	2.96	15 g OP	✓ Diprosone
2.44 /		.0 9 0.	

BETAMETHASONE DIPROPIONATE			
Crm 0.05%	2.96	15 g OP	Diprosone
	8.97	50 g OP	Diprosone
Crm 0.05% in propylene glycol base	4.33	30 g OP	Diprosone OV
Oint 0.05%	2.96	15 g OP	Diprosone
	8.97	50 g OP	Diprosone
Oint 0.05% in propylene glycol base	4.33	30 g OP	✓ Diprosone OV

	Subsidy		Fully	
	(Manufacturer's P	rice) Subs Per	idised •	
ETAMETIJA CONE VALEDATE	Ψ	1 01		Mandiadaror
BETAMETHASONE VALERATE	0.45	50 ~ OD	./	Bata Craam
★ Crm 0.1% ★ Oint 0.1%		50 g OP		Beta Cream
		50 g OP 50 ml OP		Beta Ointment Betnovate
k Lotn 0.1%	18.00	50 IIII OP	•	<u>betnovate</u>
CLOBETASOL PROPIONATE			_	
₭ Crm 0.05%		30 g OP		<u>Dermol</u>
★ Oint 0.05%	2.20	30 g OP	1	<u>Dermol</u>
CLOBETASONE BUTYRATE				
Crm 0.05%	5.38	30 g OP		
	(7.09)	•		Eumovate
DIFLUCORTOLONE VALERATE	, ,			
Crm 0.1%	8 97	50 g OP		
0111 0.1 /0	(15.86)	30 g Oi		Nerisone
Fatty oint 0.1%		50 g OP		1101100110
Tany on the ort /o	(15.86)	00 g O1		Nerisone
IVPROCORTICONE	(10.00)			140/100/10
HYDROCORTISONE	4 4 4	00 ~ OD		Darm Assist
★ Crm 1% – Only on a prescription		30 g OP		DermAssist
K December October continuities	16.25	500 g		Pharmacy Health
Powder – Only in combination		25 g		<u>ABM</u>
galenicals HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN		,		-
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only of	on			
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only of a prescription		250 ml	/	DP Lotn HC
a prescription		250 ml	1	DP Lotn HC
a prescription HYDROCORTISONE BUTYRATE	10.57			
a prescription	10.57	30 g OP	/	Locoid Lipocream
a prescription		30 g OP 100 g OP	√	Locoid Lipocream
a prescription		30 g OP	√	Locoid Lipocream
a prescription		30 g OP 100 g OP 100 g OP	111	Locoid Lipocream Locoid Lipocream Locoid
a prescription		30 g OP 100 g OP	111	Locoid Lipocream
a prescription		30 g OP 100 g OP 100 g OP	111	Locoid Lipocream Locoid Lipocream Locoid
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP	111	Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP	111	Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP 15 g OP 15 g OP	/// / / //	Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo Advantan Advantan
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP 15 g OP 15 g OP	/// / // /	Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo Advantan Advantan Elocon Alcohol Free
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP 15 g OP 15 g OP 50 g OP	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo Advantan Advantan Elocon Alcohol Free Elocon Alcohol Free
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP 15 g OP 15 g OP 50 g OP 15 g OP		Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo Advantan Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP 15 g OP 15 g OP 50 g OP 15 g OP 50 g OP 50 g OP		Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo Advantan Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon Elocon
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP 15 g OP 15 g OP 50 g OP 15 g OP		Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo Advantan Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP 15 g OP 15 g OP 50 g OP 15 g OP 50 g OP 50 g OP		Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo Advantan Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon Elocon
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP 15 g OP 15 g OP 50 g OP 15 g OP 50 g OP 30 ml OP		Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo Advantan Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon Elocon Elocon
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP 15 g OP 15 g OP 50 g OP 15 g OP 50 g OP 50 g OP		Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo Advantan Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon Elocon Elocon Elocon Aristocort
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP 15 g OP 15 g OP 15 g OP 50 g OP 30 ml OP		Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo Advantan Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon Elocon Elocon
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP 15 g OP 15 g OP 15 g OP 50 g OP 30 ml OP		Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo Advantan Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon Elocon Elocon Elocon Aristocort
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP 15 g OP 15 g OP 15 g OP 50 g OP 30 ml OP		Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo Advantan Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon Elocon Elocon Elocon Aristocort
a prescription		30 g OP 100 g OP 100 g OP 100 ml OP 15 g OP 15 g OP 15 g OP 50 g OP 30 ml OP		Locoid Lipocream Locoid Lipocream Locoid Locoid Crelo Advantan Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon Elocon Elocon Elocon Aristocort

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

DERMATOLOGICALS

	Subsidy (Manufacturaria I	Orion) Cubo	Fully	Brand or Generic
	(Manufacturer's F \$	Per Per	idised •	Manufacturer
ETAMETHASONE VALERATE WITH SODIUM FUSIDATE [Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP	F	ucicort
a) Maximum of 15 g per prescriptionb) Only on a prescription				
/DROCORTISONE WITH MICONAZOLE - Only on a prese Crm 1% with miconazole nitrate 2%	•	15 g OP	✓ N	licreme H
DROCORTISONE WITH NATAMYCIN AND NEOMYCIN - Crm 1% with natamycin 1% and neomycin sulphate 0.5%	3.35	ption 15 g OP	✓ P	imafucort
Oint 1% with natamycin 1% and neomycin sulphate 0.5% RIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOM		15 g OP TIN	✓ P	imafucort
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 and gramicidin 250 mcg per g - Only on a prescription		15 g OP	V	iaderm KC
Disinfecting and Cleansing Agents	, ,			
HLORHEXIDINE GLUCONATE – Subsidy by endorsement				
a) No more than 500 ml per month b) Only if prescribed for a dialysis patient and the prescri Handrub 1% with ethanol 70% Soln 4% wash RICLOSAN – Subsidy by endorsement	4.29	ccordingly. 500 ml 500 ml		ealthE ealthE
a) Maximum of 500 ml per prescription				
a) Maximum of 500 ml per prescription b) a) Only if prescribed for a patient identified with Me surgery in hospital and the prescription is endors b) Only if prescribed for a patient with recurrent Sta	sed accordingly; or		•	
a) Only if prescribed for a patient identified with Me surgery in hospital and the prescription is endors	sed accordingly; or phylococcus aureu		the pres	
b) a) Only if prescribed for a patient identified with Me surgery in hospital and the prescription is endors b) Only if prescribed for a patient with recurrent Sta accordingly Soln 1%	sed accordingly; or phylococcus aureu	s infection and	the pres	scription is endorsed
b) a) Only if prescribed for a patient identified with Me surgery in hospital and the prescription is endors b) Only if prescribed for a patient with recurrent Sta accordingly Soln 1%	sed accordingly; or phylococcus aureu	s infection and	the pres	scription is endorsed
a) Only if prescribed for a patient identified with Me surgery in hospital and the prescription is endors b) Only if prescribed for a patient with recurrent Sta accordingly Soln 1%	sed accordingly; or aphylococcus aureu5.90	s infection and	the pres	ealthE
a) Only if prescribed for a patient identified with Me surgery in hospital and the prescription is endors b) Only if prescribed for a patient with recurrent Sta accordingly Soln 1%	sed accordingly; or uphylococcus aureu	s infection and	the pres	ealthE Dimethicone 5% ealthE
b) a) Only if prescribed for a patient identified with Me surgery in hospital and the prescription is endors b) Only if prescribed for a patient with recurrent Sta accordingly Soln 1%	sed accordingly; or uphylococcus aureu 5.90	s infection and 500 ml OP	the pres	ealthE Dimethicone 5%
a) Only if prescribed for a patient identified with Me surgery in hospital and the prescription is endors b) Only if prescribed for a patient with recurrent Sta accordingly Soln 1% Barrier Creams and Emollients Barrier Creams METHICONE Crm 5% pump bottle	sed accordingly; or uphylococcus aureu 5.90	s infection and 500 ml OP 500 ml OP 500 ml OP	the pres	ealthE Dimethicone 10%
a) Only if prescribed for a patient identified with Me surgery in hospital and the prescription is endors b) Only if prescribed for a patient with recurrent Sta accordingly Soln 1% Barrier Creams and Emollients Barrier Creams METHICONE Crm 5% pump bottle Crm 10% pump bottle NC AND CASTOR OIL Oint	sed accordingly; or aphylococcus aureu	s infection and 500 ml OP 500 ml OP 500 ml OP	the present the pr	ealthE Dimethicone 10%

			LIMATOLOGICALO
	Subsidy		Fully Brand or
	(Manufacturer's \$	Price) Subsi	idised Generic Manufacturer
OFTOMACDOCOL	Ψ	1 01	- Wandadard
CETOMACROGOL * Crm BP	2.48	500 g	✓ healthE
CETOMACROGOL WITH GLYCEROL	2.70	300 g	· IICAILIIL
Crm 90% with glycerol 10%	2 82	500 ml OP	✓ Pharmacy Health
Offit 00 /0 with gryboror 10 /0		000 1111 01	Sorbolene with
			Glycerin
	3.87	1,000 ml OP	✓ Pharmacy Health
			Sorbolene with
			<u>Glycerin</u>
EMULSIFYING OINTMENT			_
* Oint BP	3.59	500 g	✓ <u>AFT</u>
OIL IN WATER EMULSION			_
* Crm	2.19	500 g	✓ <u>O/W Fatty Emulsion</u>
			<u>Cream</u>
PARAFFIN	5.05	500 I OD	/ backle
Oint liquid paraffin 50% with white soft paraffin 50%	5.35	500 ml OP	✓ <u>healthE</u>
UREA	4.07	100 00	A beautiful time of the con-
* Crm 10%	1.37	100 g OP	✓ <u>healthE Urea Cream</u>
WOOL FAT WITH MINERAL OIL — Only on a prescription	F 00	1 000	
* Lotn hydrous 3% with mineral oil	(11.95)	1,000 ml	DP Lotion
	1.40	250 ml OP	DI LOUOII
	(4.53)		DP Lotion
	5.60	1,000 ml	
	(20.53)		Alpha-Keri Lotion
	(23.91)	050 1 05	BK Lotion
	1.40	250 ml OP	BK Lotion
	(7.73)		בית בטווטוו
Other Dermatological Bases			
PARAFFIN			
/ U V U I U I			

White soft - Only in combination	20.20	2,500 g	✓ IPW
•	3.58	500 g	
	(7.78)	_	IPW
	(8.69)		PSM

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid – Plain.

DERMATOLOGICALS

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
\$	Per	1	Manufacturer

Minor Skin Infections

/IDONE IODINE	0.07	05 00	4 B + 11
Oint 10%	3.27	25 g OP	Betadine
a) Maximum of 100 g per prescription			
b) Only on a prescription			
Antiseptic soln 10%	6.20	500 ml	Betadine
			✓ Riodine
	1.28	100 ml	
	(4.20)		Riodine
	(13.27)		Betadine
	0.19	15 ml	
	(7.41)		Betadine
Skin preparation, povidone iodine 10% with 30% alcohol	10.00 [′]	500 ml	✓ Betadine Skin Pre
	1.63	100 ml	'
	(3.48)		Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	` ,	100 ml	
	(6.04)		Orion
	(6.64)		Pfizer

Parasiticidal Preparations

DIMETHICONE

200 ml OP ✓ healthE Dimethicone 4% Lotion

IVERMECTIN - Special Authority see SA1225 below - Retail pharmacy

Tab 3 mg - Up to 100 tab available on a PSO.......17.20 ✓ Stromectol

- 1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- 3) For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or penal institutions.

⇒SA1225 Special Authority for Subsidy

Initial application — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy: or

Subsidy	Fu	lly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
\$	Per	/	Manufacturer

continued...

- 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
- 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently;
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy: or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- Strongyloidiasis.

	HRI	

Crm 5%4.95	30 g OP	✓ Lyderm
Lotn 5%	30 ml OP	✓ A-Scables

	Subsidy (Manufacturer's Pr \$	ice) Subs Per	Fully idised	Brand or Generic Manufacturer	
PHENOTHRIN Shampoo 0.5%	11.36	200 ml OP	√ P	arasidose	

Psoriasis and Eczema Preparations

ACITRETIN - Special Authority see SA1476 below - Retail pharm	acy		
Cap 10 mg	17.86	60	✓ Novatretin
Cap 25 mg	41.36	60	✓ Novatretin

⇒SA1476 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is female and has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of two years after the completion of the treatment; or
 - 3.2 Patient is male.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is female and has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of two years after the completion of the treatment; or
- 2 Patient is male.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL			
Gel 500 mcg with calcipotriol 50 mcg per g52	2.24	60 g OP	Daivobet
Daivobet to be Sole Supply on 1 March 2019		_	
Oint 500 mcg with calcipotriol 50 mcg per g19	9.95	30 g OP	 Daivobet
CALCIPOTRIOL			
Oint 50 mcg per g45	5.00 1	00 g OP	✓ Daivonex
COAL TAR		· ·	
Soln BP – Only in combination32	2.95	200 ml	✓ Midwest

- 1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR

Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and			
allantoin crm 2.5%	6.59	75 g OP	
	(8.00)	-	Egopsoryl TA
	3.43	30 g OP	•
	(4.35)		Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR			
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97	25 g OP	✓ Coco-Scalp
	7.95	40 g OP	✓ Coco-Scalp
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN	N - Only	on a prescription	
* Soln 2.3% with trolamine laurilsulfate and fluorescein sodium	3.86	500 ml	✓ Pinetarsol

SPF 50+

	Subsidy	F	Fully Brand or
	(Manufacturer's Price)	Subsidi	ised Generic
	\$	Per	✓ Manufacturer
SALICYLIC ACID			
Powder - Only in combination	18.88	250 g	✓ PSM
 Only in combination with a dermatological base of With or without other dermatological galenicals. 	or proprietary Topical C	orticosteroid	- Plain or collodion flexible
SULPHUR			
Precipitated - Only in combination	6.35	100 g	✓ Midwest
1) Only in combination with a dermatological base of	or proprietary Topical C	orticosteroid	– Plain
With or without other dermatological galenicals.			

Scalp Preparations			
BETAMETHASONE VALERATE * Scalp app 0.1%	7.75	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE * Scalp app 0.05%	6.96	30 ml OP	✓ Dermol
HYDROCORTISONE BUTYRATE Scalp lotn 0.1% Locoid to be Sole Supply on 1 April 2019	7.30	100 ml OP	✓ Locoid
KETOCONAZOLE Shampoo 2% a) Maximum of 100 ml per prescription b) Only on a prescription	2.99	100 ml OP	✓ <u>Sebizole</u>
-,,			

Sunscreens

SUNSCREENS, PROPE	ETARY <i>–</i> Subsid	٧b٧	/ endorsement
-------------------	-----------------------	-----	---------------

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

Crm	3.30 100 g Ol	
	(5.89)	Hamilton Sunscreen
Lotn,	3.30 100 g Ol	Marine Blue Lotion
		SPF 50+
	5.10 200 g Ol	✓ Marine Blue Lotion

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 68

IMIQUIMOD

Crm 5%, 250 mg sachet......21.72 24 **✓ Perrigo**

PODOPHYLLOTOXIN

- a) Maximum of 3.5 ml per prescription
- b) Only on a prescription

DERMATOLOGICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

✓ Efudix

Other Skin Preparations

Antineoplastics

FLUOROURACIL SODIUM

✓ Durex Confidence

✓ Shield XI

144

144

	Subsidy (Manufacturer's Price) \$	F Subsidis Per	ully Brand or Sed Generic Manufacturer
Contraceptives - Non-hormonal			
Condoms			
CONDOMS * 49 mm – Up to 144 dev available on a PSO * 53 mm – Up to 144 dev available on a PSO		12	✓ Shield 49 ✓ Gold Knight ✓ Shield Blue
	13.36		✓ Shield Blue

	13.36	144	✓ Shield Blue
*	53 mm (chocolate) - Up to 144 dev available on a PSO1.11	12	✓ Gold Knight
	13.36	144	✓ Gold Knight
*	53 mm (strawberry) – Up to 144 dev available on a PSO1.11	12	✓ Gold Knight
	13.36	144	✓ Gold Knight
*	56 mm - Up to 144 dev available on a PSO	12	✓ Gold Knight
	13.36	144	✓ Durex Extra Safe
			✓ Gold Knight
*	56 mm, shaped – Up to 144 dev available on a PSO1.11	12	✓ Durex Confidence

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

*	IUD 29.1 mm length × 23.2 mm width31.60	1	✓ Choice TT380 Short
*	IUD 33.6 mm length × 29.9 mm width31.60	1	✓ Choice
			TT380 Standard
*	IUD 35.5 mm length × 19.6 mm width31.60	1	Choice Load 375

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and

2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

GENITO-URINARY SYSTEM								
	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer				
continued								
The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on								
the Schedule at 1 November 1999. Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that								
women are still either:	iiu uniii ine expiry da	te and ca	n be rem	ewed providing that				
on a Social Welfare benefit; or								
 have an income no greater than the benefit. 								
The approval numbers of Special Authorities approved before 1 N	November 1999 are ir	nterchang	eable fo	r products within the				
combined oral contraceptives and progestogen-only contraceptive	es groups, except Lo	ette and I	Microgyn	on 20 ED				
ETHINYLOESTRADIOL WITH DESOGESTREL								
* Tab 20 mcg with desogestrel 150 mcg and 7 inert tab		84	_					
	(19.80)			lercilon 28				
a) Higher subsidy of \$13.80 per 84 tab with Special Aut	hority see SA0500 or	n the prev	ious pag	е				
b) Up to 84 tab available on a PSO * Tab 30 mcg with desogestrel 150 mcg and 7 inert tab	6 62	84						
Tab of may with accognition from may and 7 more tab	(19.80)	04	M	larvelon 28				
a) Higher subsidy of \$13.80 per 84 tab with Special Aut	` '	n the prev	ious pag	е				
b) Up to 84 tab available on a PSO	•	•						
ETHINYLOESTRADIOL WITH LEVONORGESTREL								
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_							
Up to 84 tab available on a PSO		84	✓ M	licrogynon 20 ED				
* Tab 50 mcg with levonorgestrel 125 mcg and 7 inert tab – U								
to 84 tab available on a PSO		84	✓ M	licrogynon 50 ED				
* Tab 30 mcg with levonorgestrel 150 mcg	(16.50)	63	M	licrogynon 30				
a) Higher subsidy of \$15.00 per 63 tab with Special Aut	` '	the prev		0,				
b) Up to 63 tab available on a PSO	nonly dod dridded dr	i alo piov	loud pag					
Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets	_							
Up to 28 tab available on a PSO		84	_	evlen ED				
Note: Ethinyloestradiol with levonorgestrel (Levlen ED) tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets is to								
be dispensed in 28 day lots only.								
ETHINYLOESTRADIOL WITH NORETHISTERONE								
* Tab 35 mcg with norethisterone 1 mg – Up to 63 tab availab		60	./ D	revinor 1/21				
on a PSO	0.0∠	63	▼ B	TEVITION 1/21				

*	on a PSO on a PSO	6.62	63	✓ Brevinor 1/21
*	Tab 35 mcg with norethisterone 1 mg and 7 inert tab – Up to 84 tab available on a PSO	6.62	84	✓ Brevinor 1/28
*	Tab 35 mcg with norethisterone 500 mcg - Up to 63 tab available on a PSO	6.62	63	✓ Brevinor 21
*	Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - Up to 84 tab available on a PSO		84	✓ Norimin
	available on a PSO	6.62		

(Brevinor 1/21 Tab 35 mcg with norethisterone 1 mg to be delisted 1 January 2020) (Brevinor 21 Tab 35 mcg with norethisterone 500 mcg to be delisted 1 July 2019)

GENITO-URINARY SYSTEM

Subsidy (Manufactured Price)		Fully	Brand or	=
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit: and

2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

84

LEVONORGESTREL

((16.50)		Microlut
 a) Higher subsidy of \$13.80 per 84 tab with Special Authority s b) Up to 84 tab available on a PSO 	see SA0500 ab	ove	
* Subdermal implant (2 × 75 mg rods) – Up to 3 pack available			
on a PSO1	06.92	1	✓ <u>Jadelle</u>
MEDROXYPROGESTERONE ACETATE			
Inj 150 mg per ml, 1 ml syringe - Up to 5 inj available on a PSO	7.25	1	✓ Depo-Provera
NORETHISTERONE			
* Tab 350 mcg – Up to 84 tab available on a PSO	6.25	84	✓ Noriday 28

Emergency Contraceptives

LEVONORGESTREL

*	Tab 1.5 mg	4.95	1	✓ Postinor-1

- a) Maximum of 2 tab per prescription
- b) Up to 5 tab available on a PSO
- c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

GENITO-URINARY SYSTEM			
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
Antiandrogen Oral Contraceptives			
Prescribers may code prescriptions "contraceptive" (code "O") wh and prescription charge will be as per other contraceptives, as foll • \$5.00 prescription charge (patient co-payment) will apply. • prescription may be written for up to six months supply. Prescriptions coded in any other way are subject to the non control of supply. ie. Prescriptions may be written for up to three months CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL * Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO	lows: aceptive prescription s supply.	charges, and the	non-contraceptive period
	4.67	168 V <u>G</u>	<u>inet</u>
Gynaecological Anti-infectives ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC A Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applications.	е	00 g OP	ci-Jel
CLOTRIMAZOLE * Vaginal crm 1% with applicators * Vaginal crm 2% with applicators MICONAZOLE NITRATE	1.60 3		lomazol lomazol
* Vaginal crm 2% with applicator NYSTATIN Vaginal crm 100,000 u per 5 g with applicator(s)		0 g OP	icreme ilstat
Myometrial and Vaginal Hormone Preparations			
ERGOMETRINE MALEATE Inj 250 mcg per ml, 1 ml ampoule – Up to 5 inj available on a		5 ✓ E	rgonovine \$29

Inj 250 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO454	.00 5	✓ Ergonovine \$29
Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO105 (Ergonovine S29 Inj 250 mcg per ml, 1 ml ampoule to be delisted 1 July 2		✓ <u>DBL Ergometrine</u>
OESTRIOL * Crm 1 mg per g with applicator	.62 15 g OP	✓ <u>Ovestin</u> ✓ <u>Ovestin</u>
OXYTOCIN — Up to 5 inj available on a PSO Inj 5 iu per ml, 1 ml ampoule		✓ Oxytocin BNM ✓ Oxytocin BNM
OXYTOCIN WITH ERGOMETRINE MALEATE – Up to 5 inj available on a Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml		✓ Syntometrine

Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE
-\

a) Up to 200 test available on a PSO

b) Only on a PSO

GENITO-URINARY SYSTEM

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 106

5-Alpha Reductase Inhibitors

FINASTERIDE - Special Authority see SA0928 below - Retail pharmacy

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benian prostatic hyperplasia: and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

OXYBUTYNIN

***** Tab 5 mg8.85 500 **✓ Apo-**

Oxybutynin S29

POTASSIUM CITRATE

Oral liq 3 mmol per ml - Special Authority see SA1083 below -

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE

GENITO-URINARY SYSTEM

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	•	Manufacturer
SOLIFENACIN SUCCINATE				
Tab 5 mg	3.00	30	✓	Solifenacin Mylan
Solifenacin Mylan to be Sole Supply on 1 March 2019				•
Tablet 5 mg	3.00	30		
•	(37.50)			Vesicare
Tab 10 mg	5.50	30	✓	Solifenacin Mylan
Solifenacin Mylan to be Sole Supply on 1 March 2019				•
Tablet 10 mg	5.50	30		
·	(37.50)			Vesicare
(Vesicare Tablet 5 mg to be delisted 1 March 2019)	, ,			
(Vesicare Tablet 10 mg to be delisted 1 March 2019)				
TOLTERODINE - Special Authority see SA1272 below - Retail	pharmacy			
Tab 1 mg	14.56	56	✓	Arrow-Tolterodine
Tab 2 mg	14.56	56	✓	Arrow-Tolterodine
OA4070 On a shall A salt a salt of an Oash a lab.				

⇒SA1272 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where patient has overactive bladder and a documented intolerance of, or is non-responsive to oxybutynin.

Detection of Substances in Urine

OF	THO-TOLIDINE			
*	Compound diagnostic sticks	7.50	50 test OP	
		(8.25)		Hemastix
ΤE	TRABROMOPHENOL			
*	Blue diagnostic strips	7.02	100 test OP	
	•	(13.92)		Albustix

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Por	1	Manufacturer	

Calcium Homeostasis

0 4 1	_			
CA	()	1()	N	IΝ

CINACALCET - Special Authority see SA1618 below - Retail pharmacy

Tab 30 mg − Wastage claimable210.30 28 ✓ Sensipar

⇒SA1618 Special Authority for Subsidy

Initial application only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

ZOLEDRONIC ACID

⇒SA1687 Special Authority for Subsidy

Initial application — **(bone metastases)** only from an oncologist, haematologist or palliative care specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy: or
- 2 Both:
 - 2.1 Patient has bone metastases or involvement; and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
 - 3.1 Patient has bone metastases or involvement: and
 - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — **(early breast cancer)** only from an oncologist or medical practitioner on the recommendation of a oncologist. Approvals valid for 2 years for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.

Corticostero	ids and Related	d Agents for S	Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMET	HASONE ACETAT	Έ	
* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	19.20	5	
	(36.96)		Celestone
			Chronodose
DEXAMETHASONE			
* Tab 0.5 mg - Retail pharmacy-Specialist	0.99	30	✓ <u>Dexmethsone</u>
Up to 60 tab available on a PSO			
* Tab 4 mg - Retail pharmacy-Specialist	1.90	30	✓ <u>Dexmethsone</u>
Up to 30 tab available on a PSO	45.00	05 00	/ Diament
Oral liq 1 mg per ml – Retail pharmacy-Specialist Oral liq prescriptions:	45.00	25 ml OP	✓ Biomed
······	Cardialagiat, ar		
 Must be written by a Paediatrician or Paediatric On the recommendation of a Paediatrician or Pa 		ct	
2) On the recommendation of a raediatrician of ra	ediatric Gardiologi	51.	
DEXAMETHASONE PHOSPHATE			
Dexamethasone phosphate injection will not be funded for # Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a		10	✓ Max Health
* Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a		10	✓ Max Health
, , ,	F3025.16	10	• Wax Health
FLUDROCORTISONE ACETATE	14.00	100	/ Flavings
* Tab 100 mcg	14.32	100	✓ Florinef
HYDROCORTISONE			4.5
* Tab 5 mg		100	✓ <u>Douglas</u>
* Tab 20 mg		100	✓ <u>Douglas</u>
* Inj 100 mg vial	5.30	1	✓ <u>Solu-Cortef</u>
a) Up to 5 inj available on a PSO			
b) Only on a PSO			
METHYLPREDNISOLONE – Retail pharmacy-Specialist	110.00	400	/ Mardon I
* Tab 4 mg		100 20	✓ <u>Medrol</u>
* Tab 100 mg			✓ <u>Medrol</u>
METHYLPREDNISOLONE (AS SODIUM SUCCINATE) - Ret			Code Maded Ast
Inj 40 mg vial	18.90	1	✓ <u>Solu-Medrol-Act-</u>
			<u>O-Vial</u>
Inj 125 mg vial	28.90	1	✓ Solu-Medrol-Act-
., . <u></u>		•	O-Vial
			
Inj 500 mg vial	22.78	1	✓ Solu-Medrol-Act-
			<u>O-Vial</u>
Inj 1 g vial	27 02	1	✓ Solu-Medrol
	21.03	ı	- Join-Menioi
METHYLPREDNISOLONE ACETATE	44.40	-	✓ Dana Madual
Inj 40 mg per ml, 1 ml vial	44.40	5	✓ <u>Depo-Medrol</u>

	Subsidy (Manufacturer's Pri	ca) Subs	Fully Brand or sidised Generic
	\$	Per	✓ Manufacturer
ETHYLPREDNISOLONE ACETATE WITH LIDOCAINE [LIG	NOCAINE]		
Inj 40 mg per ml with lidocaine [lignocaine] 1 ml vial	9.25	1	✓ Depo-Medrol with Lidocaine
epo-Medrol with Lidocaine Inj 40 mg per ml with lidocaine [lig	gnocaine] 1 ml vial t	be delisted	1 April 2019)
REDNISOLONE			
Oral liq 5 mg per ml — Up to 30 ml available on a PSO Restricted to children under 12 years of age.	6.00	30 ml OP	✓ Redipred
REDNISONE			
Tab 1 mg		500	✓ Apo-Prednisone
Tab 2.5 mg		500	✓ Apo-Prednisone
Tab 5 mg - Up to 30 tab available on a PSO		500	✓ Apo-Prednisone
Tab 20 mg	29.03	500	✓ Apo-Prednisone
TRACOSACTRIN			_
Inj 250 mcg per ml, 1 ml ampoule		1	✓ Synacthen
Inj 1 mg per ml, 1 ml ampoule	690.00	1	✓ Synacthen Depot
			✓ Synacthene Retard \$29
RIAMCINOLONE ACETONIDE			
Inj 10 mg per ml, 1 ml ampoule		5	✓ Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule	51.10	5	✓ Kenacort-A 40
Sex Hormones Non Contraceptive			
Androgen Agonists and Antagonists			
PROTERONE ACETATE - Retail pharmacy-Specialist			
Tab 50 mg	13.17	50	✓ Siterone
•	(15.87)		Procur
Siterone to be Sole Supply on 1 March 2019	` '		
Tab 100 mg	26.75	50	✓ Siterone
	(30.40)		Procur
Siterone to be Sole Supply on 1 March 2019			
rocur Tab 50 mg to be delisted 1 March 2019)			
rocur Tab 100 mg to be delisted 1 March 2019)			

Hormone Replacement Therapy - Systemic

TESTOSTERONE CIPIONATE - Retail pharmacy-Specialist

TESTOSTERONE ESTERS - Retail pharmacy-Specialist

TESTOSTERONE UNDECANOATE - Retail pharmacy-Specialist

Patch 5 mg per day90.00

Inj 250 mg per ml, 4 ml vial......86.00

Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

30

1

1

60

✓ Androderm

✓ Depo-Testosterone

✓ Sustanon Ampoules

✓ Andriol Testocaps

✓ Reandron 1000

	Subsidy		Fully	Brand or
	(Manufacturer's Pr	rice) S Per	ubsidised	Generic Manufacturer
	\$	Per		Manufacturer
Oestrogens				
DESTRADIOL - See prescribing guideline on the previous page				
* Tab 1 mg	4.12	28 OP		
	(11.10)			Estrofem
₭ Tab 2 mg	4.12	28 OP		
	(11.10)		E	Estrofem
* Patch 25 mcg per day	6.12	8	✓ <u>[</u>	<u>Estradot</u>
 a) No more than 2 patch per week 				
b) Only on a prescription				
★ Patch 50 mcg per day	7.04	8	✓ [Estradot 50 mcg
a) No more than 2 patch per week				
b) Only on a prescription				
▶ Patch 75 mcg per day	7.91	8	✓ [Estradot
a) No more than 2 patch per week			-	
b) Only on a prescription				
▶ Patch 100 mcg per day	7.91	8	✓ 1	Estradot
a) No more than 2 patch per week		·	-	
b) Only on a prescription				
, , , , ,				
DESTRADIOL VALERATE – See prescribing guideline on the pre		0.4	, .	_
₭ Tab 1 mg		84	_	Progynova
★ Tab 2 mg	12.36	84	✓ <u>I</u>	Progynova
DESTROGENS - See prescribing guideline on the previous page	9			
Conjugated, equine tab 300 mcg	3.01	28		
	(13.50)		F	Premarin
★ Conjugated, equine tab 625 mcg	4.12	28		
	(13.50)		F	Premarin
Progestogens				
MEDROXYPROGESTERONE ACETATE - See prescribing guide	eline on the prev	ious page		
★ Tab 2.5 mg	3.75	30	✓ [Provera
-	7.00	56	✓ [Provera S29 S29
₭ Tab 5 mg		56		Provera S29 S29
r 100 0 mg	14.00	100		Provera
₭ Tab 10 mg		30	_	Provera
Progestogen and Oestrogen Combined Prepara			-	
DESTRADIOL WITH NORETHISTERONE – See prescribing guid				
★ Tab 1 mg with 0.5 mg norethisterone acetate		28 OP		
	(18.10)		ŀ	Kliovance
* Tab 2 mg with 1 mg norethisterone acetate	5.40	28 OP		
	(18.10)		ŀ	Kliogest
★ Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg				
oestradiol tab (12) and 1 mg oestradiol tab (6)	5.40	28 OP		
. , , , , , , , , , , , , , ,	(18.10)		7	Trisequens
oestradiol tab (12) and 1 mg oestradiol tab (6)		28 OP	1	Trisequens

Subsidy

(Manufacturer's Price)

Fully

Subsidised

Brand or

Generic

	\$	Per	✓ Manufacturer
Other Oestrogen Preparations			
ETHINYLOESTRADIOL * Tab 10 mcg	17.60	100	✓ NZ Medical and Scientific
OESTRIOL * Tab 2 mg	7.00	30	✓ Ovestin

Other Progestogen Preparations

LEVONORGESTREL

* Intra-uterine system 20 mcg per day - Special Authority see Mirena

⇒SA1608 Special Authority for Subsidy

Initial application — (No previous use) only from a relevant specialist or general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 The patient has a clinical diagnosis of heavy menstrual bleeding; and

- 2 The patient has failed to respond to or is unable to tolerate other appropriate pharmaceutical therapies as per the Heavy Menstrual Bleeding Guidelines; and
- 3 Either:
 - 3.1 serum ferritin level < 16 mcg/l (within the last 12 months); or
 - 3.2 haemoglobin level < 120 g/l.

Note: Applications are not to be made for use in patients as contraception except where they meet the above criteria. Renewal only from a relevant specialist or general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient demonstrated clinical improvement of heavy menstrual bleeding; or
 - 1.2 Previous insertion was removed or expelled within 3 months of insertion; and
- 2 Applicant to state date of the previous insertion.

MEDROXYPROGESTERONE ACETATE Tab 100 mg - Retail pharmacy-Specialist......101.00 100 Provera HD NORETHISTERONE * Tab 5 mg - Up to 30 tab available on a PSO.......18.29 Primolut N 100 **PROGESTERONE** Cap 100 mg - Special Authority see SA1609 below - Retail pharmacy.......16.50 Utrogestan

⇒SA1609 Special Authority for Subsidy

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 For the prevention of pre-term labour*; and
- 2 Fither:
 - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 For the prevention of pre-term labour*; and
- 2 Treatment is required for second or subsequent pregnancy; and
- 3 Either
 - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are unapproved indications.

Thyroid and Antithyroid Agents

CARBIMAZOLE			
* Tab 5 mg	10.80	100	✓ AFT
			Carbimazole S29
			✓ Neo-Mercazole
LEVOTHYROXINE			
* Tab 25 mcg	3.89	90	✓ Synthroid
* Tab 50 mcg	1.71	28	✓ Mercury Pharma
•	4.05	90	✓ Synthroid
	64.28	1,000	✓ Eltroxin
* Tab 100 mcg	1.78	28	Mercury Pharma
Ç	4.21	90	✓ Synthroid
	66.78	1,000	✓ Eltroxin
PROPYLTHIOURACIL - Special Aut	hority see SA1199 below – Retail pharmacy		
•	ded for patients under the age of 18 years un	less the patie	ent is pregnant and other
Tab 50 mg	35.00	100	✓ PTU S29

⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) - Special Authority see SA1629 below	- Retail pharma	су	
*	Inj 5 mg cartridge	34.88	1	✓ Omnitrope
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
*	Inj 15 mg cartridge	104.63	1	✓ Omnitrope

⇒SA1629 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		ubsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and</p>
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and

(Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer		Subsidy		Fully	Brand or
\$ Per ✓ Manufacturer	(M	anufacturer's Price)		sidised	
		\$	Per		Manufacturer

continued...

- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred

Initial application — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months...

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred;
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight eximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Subsidy (Manufacturer's Price)	,	Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients.

GnRH Analogues

1	✓ Zoladex
1	✓ Zoladex
	1

LEUPRORELIN

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

Inj 3.75 mg prefilled dual chamber syringe – Higher subsidy of			
\$221.60 per 1 inj with Endorsement	66.48	1	
•	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe - Higher subsidy			
of \$591.68 per 1 inj with Endorsement	177.50	1	
	(591.68)		Lucrin Depot 3-month

Vasopressin Agonists

DESMOPRESSIN ACETATE

Tab 100 mcg - Special Authority see SA1401 on the next page - Retail pharmacy	25.00	30	✓ <u>Minirin</u>
Tab 200 mcg — Special Authority see SA1401 on the next page — Retail pharmacy Nasal drops 100 mcg per ml — Retail pharmacy-Specialist Nasal spray 10 mcg per dose — Retail pharmacy-Specialist	39.03	30 2.5 ml OP 6 ml OP	✓ Minirin ✓ Minirin ✓ Desmopressin- PH&T
Inj 4 mcg per ml, 1 ml – Special Authority see SA1401 on the next page – Retail pharmacy	67.18	10	✓ Minirin

Subsidy		Fully	Brand or
(Manufacturer's Price)	ırer's Price) Subsidise		Generic
 \$	Per	✓	Manufacturer

⇒SA1401 Special Authority for Subsidy

Initial application — (Desmopressin tablets for Nocturnal enuresis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has primary nocturnal enuresis; and
- 2 The nasal forms of desmopressin are contraindicated; and
- 3 An enuresis alarm is contraindicated.

Initial application — (Desmopressin tablets for Diabetes insipidus) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has cranial diabetes insipidus; and
- 2 The nasal forms of desmopressin are contraindicated.

Renewal — (Desmopressin tablets) from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from the treatment.

Initial application — (Desmopressin injection) only from a relevant specialist. Approvals valid for 2 years where the patient cannot use desmopressin nasal spray or nasal drops.

Renewal — (Desmopressin injection) only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Other Endocrine Agents

CABERGOLINE

		Tab 0.5 mg - Maximum of 2 tab per prescription; can be
Dostinex	2	waived by Special Authority see SA1370 below3.75
✓ Dostinex	8	15 20

⇒SA1370 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 pathological hyperprolactinemia; or
- 2 acromegaly*.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

•				^:	
	()N/	๚⊢⊢Ւ	JH.	CHI	RATE

Tab 50 mg	29.84	10	✓ Mylan Clomiphen S29 ✓ Serophene
(Serophene Tab 50 mg to be delisted 1 March 2019)			·
DANAZOL			
Cap 100 mg	68.33	100	✓ Azol
Cap 200 mg	97.83	100	✓ Azol
METYRAPONE			
Cap 250 mg - Retail pharmacy-Specialist	520.00	50	✓ Metopirone

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Anthelmintics

ALBENDAZOLE - Special Authority see	SA1318 below – Retail pharmacy
= 1 .00	100.00

60 ✓ Eskazole S29

⇒SA1318 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

MEBENDAZOLE - Only on a prescription

Tab 100 mg	24.19	24	De-Worm
Oral liq 100 mg per 5 ml		15 ml	
	(7.17)		Vermox
PRAZIOLIANTEI			

Biltricide

Antibacterials

- a) For topical antibacterials, refer to DERMATOLOGICALS, page 60
- b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 221

Cephalosporins and Cephamycins

CEFACLOR MONOHYDRATE		
Cap 250 mg24.70	100	✓ Ranbaxy-Cefaclor
Grans for oral liq 125 mg per 5 ml - Wastage claimable	100 ml	✓ Ranbaxy-Cefaclor

CEFALEXIN

Cap 250 mg3.50	20	✔ Cepnalexin ABIVI
Cap 500 mg	20	✓ Cephalexin ABM
Grans for oral liq 25 mg per ml - Wastage claimable8.75	100 ml	✓ Cefalexin Sandoz

Note: Cefalexin grans for oral liq will not be funded in amounts more than 14 days treatment per dispensing.

✓ Cefalexin Sandoz Grans for oral liq 50 mg per ml – Wastage claimable......11.75 100 ml

Note: Cefalexin grans for oral liq will not be funded in amounts more than 14 days treatment per dispensing.

CEFAZOLIN - Subsidy by endorsement

Only if prescribed for dialysis or cellulitis in accordance with a DHB approved protocol and the prescription is endorsed accordingly.

Inj 500 mg vial	3.39	5	✓ <u>AFT</u>
Inj 1 g vial	3.29	5	✓ AFT

CEFTRIAXONE - Subsidy by endorsement

- a) Up to 10 inj available on a PSO
- b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of pelvic inflammatory disease, or the treatment of suspected meningococcal disease, and the prescription or PSO is endorsed accordingly.

Inj 500 mg vial	1.20	1	DEVA
Inj 1 g vial	0.84	1	✓ DEVA

CEFUROXIME AXETIL - Subsidy by endorsement

Only if prescribed for prophylaxis of endocarditis and the prescription is endorsed accordingly

Only if prescribed for propriyaxis of endocarditis and the	prescription is endorsed	accordin	igiy.
Tab 250 mg	29.40	50	Zinnat

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ıbsidised	Generic	
\$	Per	✓	Manufacturer	

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	8.19	30	✓ Apo-Azithromycin
•	8.50	6	✓ Zithromax
Tab 500 mg - Up to 8 tab available on a PSO	0.93	2	✓ Apo-Azithromycin
Grans for oral lig 200 mg per 5 ml (40 mg per ml) - Wastage			
claimable	14.38	15 ml	✓ Zithromax

(Zithromax Tab 250 mg to be delisted 1 June 2019)

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN

Tab 250 mg − Maximum of 28 tab per prescription; can be waived by Special Authority see SA1131 on the next page3.98

Grans for oral liq 250 mg per 5 ml23.12

✓ Apo-Clarithromycin ✓ Klacid

- a) Maximum of 500 mg per prescription; can be waived by Special Authority see SA1131 on the next page
- b) Wastage claimable

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1131 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Renewal — **(Mycobacterial infections)** only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1131 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Renewal — **(Mycobacterial infections)** only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	E-Mycin
a) Up to 20 tab available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	E-Mycin
a) Up to 300 ml available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	E-Mycin
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
ERYTHROMYCIN LACTOBIONATE			
Inj 1 g	16.00	1	✓ Erythrocin IV
ERYTHROMYCIN STEARATE		-	
Tab 250 mg – Up to 30 tab available on a PSO	14.05	100	
rab 250 mg – Op to 50 tab available on a P50	(22.29)	100	ERA
Tab 500 mg		100	ENA
Tab 500 Hig	(44.58)	100	ERA
	(44.50)		LITA
ROXITHROMYCIN			4 - 11 -
Tab disp 50 mg	7.19	10	✓ Rulide D
Restricted to children under 12 years of age.			
Tab 150 mg	7.48	50	✓ Arrow-
			Roxithromycin
Tab 300 mg	14 40	50	✓ Arrow-
100 000 mg		00	Roxithromycin
			ctii oiiiyoiii

	Subsidy (Manufacturer's Pr \$	rice) Subs Per	Fully Brand or idised Generic Manufacturer
Penicillins			
AMOXICILLIN			
Cap 250 mg	14.97	500	✓ Apo-Amoxi
a) Up to 30 cap available on a PSO			
b) Up to 10 x the maximum PSO quantity for RFPP			
Cap 500 mg	16.75	500	✓ Apo-Amoxi
a) Up to 30 cap available on a PSO			
b) Up to 10 x the maximum PSO quantity for RFPP			4 4 4 4 4 4 4 4 4
Grans for oral liq 125 mg per 5 ml	1.20	100 ml	✓ Alphamox 125
a) Up to 200 ml available on a PSO			
b) Wastage claimable	4.04	400 1	4 41 1 250
Grans for oral liq 250 mg per 5 ml	1.31	100 ml	✓ Alphamox 250
a) Up to 300 ml available on a PSO			
b) Up to 10 x the maximum PSO quantity for RFPPc) Wastage claimable			
Inj 250 mg vial	10.67	10	✓ Ibiamox
Inj 500 mg vial		10	✓ Ibiamox
Inj 1 g vial – Up to 5 inj available on a PSO	17.29	10	✓ Ibiamox
AMOXICILLIN WITH CLAVULANIC ACID			
Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab			
available on a PSO	1 99	20	✓ Augmentin
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25		20	Augmentin
per ml		100 ml	✓ Augmentin
a) Up to 200 ml available on a PSO		100 1111	- Augmontin
b) Wastage claimable			
Grans for oral lig amoxicillin 50 mg with clavulanic acid 12.5	ma		
per ml – Up to 200 ml available on a PSO		100 ml OP	✓ Curam
BENZATHINE BENZYLPENICILLIN			
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj			
available on a PSO	344 93	10	✓ Bicillin LA
		10	5 BIOIIIII EA
BENZYLPENICILLIN SODIUM [PENICILLIN G] Inj 600 mg (1 million units) vial – Up to 5 inj available on a P.	SO 10.25	10	✓ Sandoz
, , , , ,	30 10.33	10	Januoz
FLUCLOXACILLIN	16.00	050	./ Ctambley
Cap 250 mg - Up to 30 cap available on a PSO Cap 500 mg		250 500	✓ <u>Staphlex</u>✓ Staphlex
Grans for oral liq 25 mg per ml		100 ml	✓ <u>Staprilex</u> ✓ AFT
a) Up to 200 ml available on a PSO		100 1111	· <u>BL I</u>
b) Wastage claimable			
Grans for oral liq 50 mg per ml	3.68	100 ml	✓ AFT
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
Inj 250 mg vial	9.00	10	✓ Flucloxin
Inj 500 mg vial	9.40	10	✓ Flucloxin
Inj 1 g vial – Up to 5 inj available on a PSO	5.22	5	✓ Flucil

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
PHENOXYMETHYLPENICILLIN (PENICILLIN V)				
Cap 250 mg - Up to 30 cap available on a PSO	2.59	50	✓	Cilicaine VK
Cap 500 mg	4.26	50	✓	Cilicaine VK
a) Up to 20 cap available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	1.48	100 m		<u>AFT</u>
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	1.58	100 m		<u>AFT</u>
a) Up to 300 ml available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
PROCAINE PENICILLIN				
Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO	123.50	5	/	Cilicaine
Tetracyclines				
DOXYCYCLINE				
* Tab 50 mg - Up to 30 tab available on a PSO	2.90	30		
•	(6.00)			Doxy-50
* Tab 100 mg - Up to 30 tab available on a PSO	6.75	250	1	Doxine
MINOCYCLINE HYDROCHLORIDE				
* Tab 50 mg - Additional subsidy by Special Authority see				
SA1355 below – Retail pharmacy	5.79	60		
,	(12.05)			Mino-tabs
* Cap 100 mg		100		
· •	(52.04)			Minomycin
⇒SA1355 Special Authority for Manufacturers Price				
Initial application from any relevant practitioner. Approvals valid	id without further rene	wal u	nless notif	ied where the patient has
rosacea.				
TETRACYCLINE - Special Authority see SA1332 below - Retain	il pharmacy			
Cap 500 mg	46.00	30	✓	Tetracyclin

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Wolff S29

INFECTIONS - AGENTS FOR SYSTEMIC USE				
	Subsidy (Manufacturer's Price) \$; Per	Fully Subsidised	Brand or Generic Manufacturer
Other Antibiotics				
For topical antibiotics, refer to DERMATOLOGICALS, page 60				
CIPROFLOXACIN				
Recommended for patients with any of the following: i) microbiologically confirmed and clinically significant ps ii) prostatitis; or	eudomonas infection;	or		
iii) pyelonephritis; oriv) gonorrhoea.				
Tab 250 mg - Up to 5 tab available on a PSO	1.45	28	•	Cipflox
Tab 500 mg - Up to 5 tab available on a PSO	1.99	28	✓	Cipflox
Tab 750 mg	3.15	28	✓	Cipflox
CLINDAMYCIN				
Cap hydrochloride 150 mg - Maximum of 4 cap per				
prescription; can be waived by endorsement - Retail				
pharmacy - Specialist	4.10	16	1	Clindamycin ABM
Inj phosphate 150 mg per ml, 4 ml ampoule - Retail				
pharmacy-Specialist	65.00	10	✓	Dalacin C
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist -	Subsidy by endorseme	nt		
Only if prescribed for dialysis or cystic fibrosis patient and the			ccordinaly	
Inj 150 mg		1	٠,	Colistin-Link
GENTAMICIN SULPHATE				
Inj 10 mg per ml, 1 ml ampoule – Subsidy by endorsement.	25 00	5	1	DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient		-		
endorsed accordingly.	or complicated armary	liuot	iiiicolloii c	and the presemption is
Inj 10 mg per ml, 2 ml – Subsidy by endorsement	62.00	5	1	Wockhardt S29
,	175.10	25		APP
				Pharmaceuticals S29
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	/ tract	infection a	and the prescription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement.	6.00	10	✓	Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.		/ tract	infection a	and the prescription is
(Wockhardt S29 Inj 10 mg per ml, 2 ml to be delisted 1 April 20	119)			
(APP Pharmaceuticals S29 Inj 10 mg per ml, 2 ml to be deliste	d 1 April 2019)			
MOXIFLOXACIN - Special Authority see SA1740 below - Reta	il pharmacy			
No patient co-payment payable	•			
Tab 400 mg	52.00	5	✓.	Avelox
OA4740 Outside Auditority for Outside				

⇒SA1740 Special Authority for Subsidy

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

1 Both:

1.1 Active tuberculosis*; and

	ubsidy cturer's Price) Subs	Fully	Brand or Generic
·	\$ Per	•	Manufacturer

continued...

- 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications;
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
 - 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy

Cap 250 mg.......126.00 16 ✓ Humatin S29

SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Fither:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharmacy

30 ✓ Daraprim S29 50 ✓ Daraprim S29 36.95

⇒SA1328 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

	II LOTIONS - A			TOTOTE WITO OOL
	Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	
SODIUM FUSIDATE [FUSIDIC ACID]				
Tab 250 mg — Retail pharmacy-Specialist Prescriptions must be written by, or on the recommendat		12 diseas		Fucidin an or a clinical microbiologist
SULFADIAZINE SODIUM – Special Authority see SA1331 below Tab 500 mg		56	/	Wockhardt \$29
⇒SA1331 Special Authority for Subsidy		30	•	Wockhardt
Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following:	without further ren	ewal ur	nless notifi	ied for applications meeting
1 For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months of the pregnancy.	·	ns; or		
TOBRAMYCIN	-			
Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement Only if prescribed for dialysis or cystic fibrosis patient and		5 endors		Tobramycin Mylan lingly.
Solution for inhalation 60 mg per ml, 5 ml — Subsidy by endorsement	2 200 00	56 dos	· /	ТОВІ
a) Wastage claimable b) Only if prescribed for a cystic fibrosis patient and the	,			
TRIMETHOPRIM			3,	
* Tab 300 mg - Up to 30 tab available on a PSO	16.50	50	•	<u>TMP</u>
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX/ * Tab trimethoprim 80 mg and sulphamethoxazole 400 mg – U	lp	500	,	Tribud
to 30 tab available on a PSO * Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to 200 r available on a PSO	nl	500 100 m		Trisul
	2.91	100 111	•	<u>Deprim</u>
VANCOMYCIN – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient or for difficile following metronidazole failure and the prescription is	prophylaxis of endo	ocarditi:	s or for tre	eatment of Clostridium
Inj 500 mg vial		1	✓	<u>Mylan</u>
Antifungals				
a) For topical antifungals refer to DERMATOLOGICALS, page 6° b) For topical antifungals refer to GENITO URINARY, page 74				
FLUCONAZOLE				
Cap 50 mg - Retail pharmacy-Specialist Cap 150 mg - Subsidy by endorsement	0.33	28 1	1	<u>Mylan</u> Mylan
 a) Maximum of 1 cap per prescription; can be waived by b) Patient has vaginal candida albicans and the practition of recommended and the prescription is endorsed as Specialist. 	ner considers that a	a topica	l imidazol	e (used intra-vaginally) is
Cap 200 mg - Retail pharmacy-Specialist	5.08	28	1	<u>Mylan</u>
Powder for oral suspension 10 mg per ml - Special Authority			_	
see SA1359 on the next page – Retail pharmacy	34.56 98.50	35 ml		Diflucan S29 S29 Diflucan
Wastage claimable	90.00		•	Dillucali
Š				

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
<u> </u>	Per	✓	Manufacturer

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOLE

Cap 100 mg − Subsidy by endorsement2.79 15 ✓ Itrazole

Funded for tinea vesicolor where topical treatment has not been successful and diagnosis has been confirmed by mycology, or for tinea unguium where terbinafine has not been successful in eradication or the patient is intolerant to terbinafine and diagnosis has been confirmed by mycology and the prescription is endorsed accordingly. Can be waived by endorsement - Retail pharmacy - Specialist Specialist must be an infectious disease physician, clinical microbiologist, clinical immunologist or dermatologist.

Oral liq 10 mg per ml − Special Authority see SA1322 below −
Retail pharmacy......141.80 150 ml OP ✓ Sporanox

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE

Tab 200 mg - PCT - Retail pharmacy-Specia endorsement	, ,	30	✓ Link Healthcare \$29 ✓ Nizoral \$29
Prescriptions must be written by, or on the	e recommendation of an oncologist		- MEGICI
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.00)		Nilotot

	Subsidy (Manufacturer's Price	,	Fully ubsidised	Brand or Generic
	\$	Per		Manufacturer
POSACONAZOLE - Special Authority see SA1285 below - Reta	ail pharmacy			
Tab modified-release 100 mg	869.86	24	✓ N	oxafil
Oral liq 40 mg per ml	761.13 1	05 ml OF	• √ N	oxafil
SA1285 Special Authority for Subsidy				

SA1285 | Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy: or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Fither:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE

* Tab 250 mg	14	✓ Deolate
VORICONAZOLE - Special Authority see SA1273 below - Retail pharmacy		
Tab 50 mg91.00	56	✓ Vttack
Tab 200 mg350.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage		
claimable1,437.00	70 ml	✓ Vfend

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis: or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
- 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
- 3.3 Patient has fluconazole resistant candidiasis; or
- 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

PRIMAQUINE PHOSPHATE - Special Authority see SA1684 below - Retail pharmacy

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Antiparasitics

Antiprotozoals

QUININE SULPHATE

Antitrichomonal Agents

METRONIDAZOI E

Tab 200 mg – Up to 30 tab available on a PSO	10.45	100	Trichozole
Tab 400 mg - Up to 15 tab available on a PSO	18.15	100	✓ Trichozole
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg	24.48	10	✓ Flagyl
ORNIDAZOLE			
Tab 500 mg	23.00	10	✓ Arrow-Ornidazole

Antituberculotics and Antileprotics

Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status.

CLOFAZIMINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
CYCLOSERINE - Retail pharmacy-Specialist				
a) No patient co-payment payableb) Prescriptions must be written by, or on respiratory physician.	ne recommendation of, an infectious o	liseas	e physicia	n, clinical microbiologist
Cap 250 mg	1,294.50	100	/	King S29
APSONE - Retail pharmacy-Specialist				
a) No patient co-payment payableb) Prescriptions must be written by, or on dermatologist	ne recommendation of, an infectious of	liseas	e physicia	n, clinical microbiologist
Tab 25 mg		100		Dapsone
Tab 100 mg		100	•	Dapsone
THAMBUTOL HYDROCHLORIDE – Retail p	armacy-Specialist			
a) No patient co-payment payableb) Prescriptions must be written by, or on respiratory physician		liseas	se physicia	n, clinical microbiologist
Tab 100 mg	85.73	100	/	EMB Fatol \$29
Tab 400 mg	49.34	56	/	Myambutol S29
ONIAZID – Retail pharmacy-Specialist				
a) No patient co-payment payable b) Prescriptions must be written by, or on microbiologist, dermatologist or public to the state of the	ealth physician	dicine		, paediatrician, clinical
ONIAZID WITH RIFAMPICIN – Retail pharm		100		<u>r Jiii</u>
a) No patient co-payment payable b) Prescriptions must be written by, or on microbiologist, dermatologist or public h	ne recommendation of, an internal me ealth physician	dicine	e physician	, paediatrician, clinical
Tab 100 mg with rifampicin 150 mg		100		Rifinah
Tab 150 mg with rifampicin 300 mg		100	/	<u>Rifinah</u>
ARA-AMINO SALICYLIC ACID – Retail phan	acy-Specialist			
a) No patient co-payment payable				
b) Specialist must be an infectious disease		•		
Grans for oral liq 4 g sachet	280.00	30	•	Paser S29
ROTIONAMIDE – Retail pharmacy-Specialis				
a) No patient co-payment payableb) Specialist must be an infectious diseas	enecialist clinical microhiologist or re	enirat	on, enocia	liet
Tab 250 mg	. •	100		Peteha S29
· ·		100	•	i ctciia 💴
YRAZINAMIDE – Retail pharmacy-Specialist a) No patient co-payment payable				
b) Prescriptions must be written by, or on respiratory physician	ne recommendation of, an infectious of	liseas		•
Tab 500 mg	59.00	100	/	AFT-Pyrazinamide
FABUTIN - Retail pharmacy-Specialist				
a) No patient co-payment payable b) Prescriptions must be written by, or on gastroenterologist	ne recommendation of, an infectious of	liseas	e physicia	n, respiratory physician o
	275.00	30		Mycobutin

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

RIFAMPICIN - Subsidy by endorsement

- a) No patient co-payment payable
- b) For confirmed recurrent Staphylococcus aureus infection in combination with other effective anti-staphylococcal antimicrobial based on susceptibilities and the prescription is endorsed accordingly; can be waived by endorsement -Retail pharmacy - Specialist. Specialist must be an internal medicine physician, clinical microbiologist, dermatologist, paediatrician, or public health physician.

*	Cap 150 mg55.75	100	1	Rifadin
*	Cap 300 mg116.25	100	1	Rifadin
*	Oral liq 100 mg per 5 ml12.00	60 ml	1	Rifadin

Antivirals

For eye preparations refer to Eye Preparations, Anti-Infective Preparations, page 221

Hepatitis B Treatment

⇒SA0829 Special Authority for Subsidy

Initial application only from a gastroenterologist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg+); and Documented resistance to lamiyudine, defined as:
- 2 Patient has raised serum ALT (> 1 x ULN); and
- 3 Patient has HBV DNA greater than 100,000 copies per mL, or viral load 10 fold or higher over nadir; and
- 4 Detection of M204I or M204V mutation: and
- 5 Fither:
 - 5.1 Both
 - 5.1.1 Patient is cirrhotic; and
 - 5.1.2 adefovir dipivoxil to be used in combination with lamivudine: or
 - 5.2 Both:
 - 5.2.1 Patient is not cirrhotic; and
 - 5.2.2 adefovir dipivoxil to be used as monotherapy.

Renewal only from a gastroenterologist or infectious disease specialist. Approvals valid for 2 years where in the opinion of the treating physician, treatment remains appropriate and patient is benefiting from treatment.

Notes: Lamivudine should be added to adefovir dipivoxil if a patient develops documented resistance to adefovir dipivoxil, defined as:

- i) raised serum ALT (> 1 × ULN); and
- ii) HBV DNA greater than 100,000 copies per mL, or viral load 10 fold or higher over nadir; and
- iii) Detection of N236T or A181T/V mutation.

Adefovir dipivoxil should be stopped 6 months following HBeAg seroconversion for patients who were HBeAg+ prior to commencing adefovir dipivoxil.

The recommended dose of adefovir dipivoxil is no more than 10mg daily.

In patients with renal insufficiency adefovir dipivoxil dose should be reduced in accordance with the datasheet guidelines. Adefovir dipivoxil should be avoided in pregnant women and children.

ENTECAVIR – Brand switch fee payable (Pharmacode 2559420) - see page 22	o for details				
* Tab 0.5 mg	52.00	30	 Entecavir Sandoz 			
LAMIVUDINE - Special Authority see SA1685 on the next page - Retail pharmacy						
Tab 100 mg	4.20	28	✓ Zetlam			
Oral liq 5 mg per ml	270.00	240 ml OP	✓ Zeffix			

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1685 Special Authority for Subsidy

Harnaavirus Traatmanta

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B. TENOFOVIR DISOPROXIL

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA1651., page 104

nerpesvirus rreatilients			
ACICLOVIR			
* Tab dispersible 200 mg	1.60	25	✓ Lovir
* Tab dispersible 400 mg		56	✓ <u>Lovir</u> ✓ Lovir
* Tab dispersible 800 mg		35	✓ Lovir
VALACICLOVIR			
Tab 500 mg	5.75	30	✓ Vaclovir
Tah 1 000 mg		30	✓ Vaclovir

SA1404 Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis: and
- 2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Roth:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via PHARMAC's approved direct distribution supply. Further details can be found on

PHARMAC's website https://www.pharmac.govt.nz/hepatitis-c-treatments

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz/hepatitis-c-treatments or:

The Coordinator, Hepatitis C Treatment Panel

PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL FUMARATE – Subsidy by endorsement; can be waived by Special Authority see SA1714 on the next page

Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil fumarate is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil fumarate prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 104 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the PHARMAC website.

Tab 200 mg with tenofovir disoproxil fumarate 300 mg......190.02 30 ✓ Truvada

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1714 Special Authority for Waiver of Rule

Initial application only from a named specialist or medical practitioner on the recommendation of a named specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- Patient has tested HIV negative; and
 - 2 Fither:
 - 2.1 All of the following:
 - 2.1.1 Patient is male or transgender; and
 - 2.1.2 Patient has sex with men; and
 - 2.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 2.1.4 Any of the following:
 - 2.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 2.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 2.1.4.3 Patient has used methamphetamine in the last three months; or
 - 2.2 All of the following:
 - 2.2.1 Patient has a regular partner who has HIV infection; and
 - 2.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 2.2.3 Condoms have not been consistently used.

All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis; and
- 2 Patient has undergone testing for HIV, syphilis, and a full STI screen in the previous two weeks; and

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative; and
- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men: and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Antiretrovirals

⇒SA1651 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil fumarate prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals.

Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil fumarate prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals.

Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil fumarate prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals.

Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates

Subsidy		Fully	Brand or
(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
Ψ	rei		Manuacturei

continued...

prophylaxis is required.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil fumarate prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals.

Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA1651 on the pre	evious page – Retail phar	macy	
Tab 50 mg	63.38	30	✓ Stocrin S29
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
Oral liq 30 mg per ml	145.79	180 ml OP	✓ Stocrin S29
ETRAVIRINE - Special Authority see SA1651 on the pr	revious page – Retail pha	ırmacy	
Tab 200 mg	770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA1651 on the p	revious page – Retail pha	ırmacy	
Tab 200 mg	60.00	60	✓ Nevirapine
			<u>Alphapharm</u>
Oral suspension 10 mg per ml	203.55	240 ml	✓ Viramune
			Suspension

Nucleosides Reverse Transcriptase Inhibitors

ABACAVIR SULPHATE – Special Authority see SA1651 on the Tab 300 mg		Retail pharmad	cy ✔ Ziagen
Oral liq 20 mg per ml		240 ml OP	✓ Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE — Special Author Note: abacavir with lamivudine (combination tablets) cou anti-retroviral Special Authority.	,		• ,
Tab 600 mg with lamivudine 300 mg	427.29	30	✓ Kivexa
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISC previous page – Retail pharmacy Note: Efavirenz with emtricitabine and tenofovir disoprox purposes of the anti-retroviral Special Authority		·	•
Tab 600 mg with emtricitabine 200 mg and tenofovir disor fumarate 300 mg		30	✓ Atripla
EMTRICITABINE - Special Authority see SA1651 on the pre-		pharmacy	·
Cap 200 mg		30	✓ Emtriva
LAMIVUDINE - Special Authority see SA1651 on the previou	s page – Retail pha	armacy	
Tab 150 mg	52.50	60	LamivudineAlphapharm
Oral liq 10 mg per ml	102.50	240 ml OP	✓ 3TC
ZIDOVUDINE [AZT] - Special Authority see SA1651 on the p	revious page - Ret	ail pharmacy	
Cap 100 mg	152.25	100	✓ Retrovir

200 ml OP

Retrovir

	Subsidy		Fully Brand or
	(Manufacturer's Price) \$	Subsidi Per	lised Generic ✓ Manufacturer
	•		
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority see Note: zidovudine [AZT] with lamivudine (combination tablets) the anti-retroviral Special Authority.	counts as two anti-re		
Tab 300 mg with lamivudine 150 mg	33.00	60	✓ <u>Alphapharm</u>
Protease Inhibitors			
ATAZANAVIR SULPHATE - Special Authority see SA1651 on pa	ige 104 – Retail phar	macy	
Cap 150 mg	568.34	60	✓ Reyataz
Cap 200 mg	757.79	60	✓ Reyataz
DARUNAVIR - Special Authority see SA1651 on page 104 - Ret	ail pharmacy		
Tab 400 mg	335.00	60	✓ Prezista
Tab 600 mg	476.00	60	✓ Prezista
LOPINAVIR WITH RITONAVIR - Special Authority see SA1651 of	on page 104 – Retail	pharmacy	
Tab 100 mg with ritonavir 25 mg		['] 60	✓ Kaletra
Tab 200 mg with ritonavir 50 mg	463.00	120	✓ Kaletra
Oral liq 80 mg with ritonavir 20 mg per ml	735.00 300	0 ml OP	✓ Kaletra
RITONAVIR - Special Authority see SA1651 on page 104 - Reta	il pharmacy		
Tab 100 mg		30	✓ Norvir
Strand Transfer Inhibitors			
DOLUTEGRAVIR - Special Authority see SA1651 on page 104 -	- Retail pharmacy		
Tab 50 mg	, ,	30	✓ Tivicay
RALTEGRAVIR POTASSIUM – Special Authority see SA1651 or	•		•

Immune Modulators

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

- 1) Diagnosis
 - Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test: or
 - PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
 - Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Exclusion Criteria

- 1) Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia (< 2.0 × 10⁹) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

Dosage

continued...

✓ Isentress

60

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	1	

continued...

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

INTERFERON ALFA-2A - PCT - Retail pharmacy-Specialist

- a) See prescribing guideline on the previous page
- b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician or ophthalmologist
- ✓ Roferon-A 1

INTERFERON ALFA-2B - PCT - Retail pharmacy-Specialist

- a) See prescribing guideline on the previous page
- b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician or ophthalmologist
- ✓ Intron-A 1 ✓ Intron-A ✓ Intron-A

(Intron-A Inj 18 m iu, 1.2 ml multidose pen to be delisted 1 May 2019)

(Intron-A Inj 30 m iu, 1.2 ml multidose pen to be delisted 1 May 2019)

(Intron-A Inj 60 m iu, 1.2 ml multidose pen to be delisted 1 May 2019)

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA1400 below - Retail pharmacy

- a) See prescribing guideline on the previous page
- b) Note: PHARMAC will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at PHARMAC on 0800-023-588 option 4.
- ✓ Pegasys

⇒SA1400 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C. genotype 1: and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Fither:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and

(Ma	Subsidy	Fully		Brand or
	nufacturer's Price)	Subsidised		Generic
	\$	Per	1	Manufacturer

continued...

5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — **(Hepatitis B)** only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Notes:

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet quidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

Urinary Tract Infections

HEXAMINE HIPPURATE			
* Tab 1 g	18.40	100	
·	(40.01)		Hiprex

INFECTIONS - AGENTS FOR SYSTEMIC USE

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
NITROFURANTOIN				
* Tab 50 mg	22.20	100	✓	Nifuran
* Tab 100 mg	37.50	100	1	Nifuran
NORFLOXACIN				
Tab 400 mg - Subsidy by endorsement	135.00	100	✓	Arrow-Norfloxacin
Only if prescribed for a patient with an uncomplicated urin with proven resistance to first line agents and the prescrip				ive to a first line agent of

	Cubaidu		Eully	Brand or
	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	✓	Manufacturer
Anticholinesterases				
NEOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	98.00	50	/	AstraZeneca
PYRIDOSTIGMINE BROMIDE				
▲ Tab 60 mg	42.79	100	1	Mestinon
Non-Steroidal Anti-Inflammatory Drugs				
DICLOFENAC SODIUM				
* Tab EC 25 mg	1.23	50	1	Diclofenac Sandoz
* Tab 50 mg dispersible		20		Voltaren D
* Tab EC 50 mg	1.23	50	✓	Diclofenac Sandoz
Tab long-acting 75 mg	22.80	500	1	Apo-Diclo SR
* Tab long-acting 100 mg	25.15	500		Apo-Diclo SR
Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a	a PSO 13.20	5	✓	Voltaren
* Suppos 12.5 mg		10	✓	Voltaren
* Suppos 25 mg		10	_	Voltaren
* Suppos 50 mg - Up to 10 supp available on a PSO		10	_	Voltaren
* Suppos 100 mg	7.00	10	•	Voltaren
IBUPROFEN				
* Tab 200 mg	11.71	1,000	/	Relieve
* Tab long-acting 800 mg	7.99	30	✓	Brufen SR
* Oral liq 20 mg per ml	2.39	200 m	ı 🗸	Fenpaed
KETOPROFEN				
* Cap long-acting 200 mg	12.07	28	✓	Oruvail SR
MEFENAMIC ACID				
* Cap 250 mg	1 25	50		
- σαρ 200 mg	(9.16)	00		Ponstan
	0.50	20		Tonotan
	(5.60)			Ponstan
NAPROXEN	(0.00)			
* Tab 250 mg	32.60	500	1	Noflam 250
* Tab 500 mg		250	_	Noflam 500
* Tab long-acting 750 mg		28		Naprosyn SR 750
* Tab long-acting 1 g		28		Naprosyn SR 1000
SULINDAC				
* Tab 100 mg	9.55	50	_	Aclin
* Tab 100 mg*		50		Aclin
•	13.10	30	•	Aciiii
TENOXICAM	10.05	100		Tileadil
* Tab 20 mg		100	_	Tilcotil
* Inj 20 mg vial	9.95	1	•	AFT
NSAIDs Other				
CELECOXIB				
Cap 100 mg	3.63	60		Celebrex
				Celecoxib Pfizer
Cap 200 mg	2.30	30	1	Celecoxib Pfizer

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	sidised	Generic	
\$	Per	1	Manufacturer	

Topical Products for Joint and Muscular Pain

CAPSAICIN

Crm 0.025% - Special Authority see SA1289 below - Retail		
pharmacy	25 g OP	✓ Zostrix
9.95		✓ Zostrix

⇒SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Antirheumatoid Agents

HYDROXYCHLOROQUINE			
* Tab 200 mg	7.98	100	✓ <u>Plaquenil</u>
LEFLUNOMIDE			
Tab 10 mg	2.90	30	✓ Apo-Leflunomide
Tab 20 mg		30	✓ Apo-Leflunomide
PENICILLAMINE			
Tab 125 mg	67.23	100	✓ D-Penamine
Tab 250 mg	110.12	100	D-Penamine
SODIUM AUROTHIOMALATE			
Inj 10 mg in 0.5 ml ampoule	76.87	10	✓ Myocrisin
Inj 20 mg in 0.5 ml ampoule	113.17	10	✓ Myocrisin
Inj 50 mg in 0.5 ml ampoule		10	✓ Myocrisin

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

ALENDRONATE CODILIM

* Tab 70 mg2.4	4	4	✓ Fosamax
Fosamax to be Sole Supply on 1 May 2019			
ALENDRONATE SODIUM WITH COLECALCIFEROL			
* Tab 70 mg with colecalciferol 5,600 iu1.5	1	4	✓ Fosamax Plus
Fosamax Plus to be Sole Supply on 1 May 2019			

Alendronate for Paget's Disease

⇒SA0949 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity: or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications due to site (base of skull, spine, long bones of lower limbs); or
 - 2.5 Preparation for orthopaedic surgery.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

ALENDRONATE SODIUM - Special Authority see SA0949 on the previous page - Retail pharmacy

(Fosamax Tab 40 mg to be delisted 1 May 2019)

Other Treatments

DENOSUMAB – Special Authority see SA1777 below – Retail pharmacy
Inj 60 mg prefilled syringe......326.00 1 ✓ Prolia

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Either:
 - 2.1 The patient is female and postmenopausal; or
 - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
PAMIDRONATE DISODIUM				
Inj 3 mg per ml, 10 ml vial	5.98	1	✓	Pamisol
Inj 6 mg per ml, 10 ml vial		1	1	Pamisol
Inj 9 mg per ml, 10 ml vial	17.05	1	✓	Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see SA17	779 below – Retail ph	armac	:V	
* Tab 60 mg		28	′ 🗸	Evista

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Tab 35 mg	3.80 4	✓ Risedronate Sandoz
TERIPARATIDE - Special Authority see SA1139 below - Retail p	oharmacy	
Inj 250 mcg per ml, 2.4 ml	490.00 1	✓ Forteo

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once dailv: zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, vial - Special Authority see

SA1780 below – Retail pharmacy600.00 100 ml OP ✓ Aclasta

⇒SA1780 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note): or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined): and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2010 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Notes:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fracility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has guantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

ALLOPURINOL			
* Tab 100 mg	4.54	500	✓ DP-Allopurinol
* Tab 300 mg	10.35	500	✓ DP-Allopurinol
BENZBROMARONE - S	ecial Authority see SA1537 below – Retail pharmacy		
Tab 100 mg	45.00	100	✓ Benzbromaron AL
			100 S29

⇒SA1537 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose: or
 - 2.3 Both:
 - 2.3.1 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Notes); and
 - 2.3.2 The patient has a rate of creatinine clearance greater than or equal to 20 ml/min; or
 - 2.4 All of the following:
 - 2.4.1 The patient is taking azathioprine and requires urate-lowering therapy; and
 - 2.4.2 Allopurinol is contraindicated; and
 - 2.4.3 Appropriate doses of probenecid are ineffective or probenecid cannot be used due to reduced renal function: and
- 3 The patient is receiving monthly liver function tests.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	•	Manufacturer

continued...

Notes: Benzbromarone has been associated with potentially fatal hepatotoxicity.

In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

The New Zealand Rheumatology Association has developed information for prescribers which can be accessed from its website at www.rheumatology.org.nz/home/resources-2/

COLCHICINE

* Tab 500 mcg	9.58	100	✓ Colgout
FEBUXOSTAT - Special Authority see SA1538 below - Reta	il pharmacy		
Tab 80 mg	39.50	28	✓ Adenuric
Tab 120 mg	39.50	28	✓ Adenuric

⇒SA1538 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose: or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note).

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

PROBENECID

*	Tab 500 mg	100	Probenecid-AFT
---	------------	-----	----------------

Muscle Relaxants			
BACLOFEN			
* Tab 10 mg	4.20	100	✓ Pacifen
Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement	11.55	1	✓ Lioresal Intrathecal
Subsidised only for use in a programmable pump in patier caused intolerable side effects and the prescription is end-		, ,	ents have been ineffective or have
Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement	209.29	1	✓ Lioresal Intrathecal
	372.98	5	✓ Medsurge
Subsidised only for use in a programmable pump in patier			ents have been ineffective or have

DANTROLENE

Cap 25 mg	65.00	100	✓ Dantrium ✓ Dantrium S29 S29
Cap 50 mg	77.00	100	✓ Dantrium

	Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
	\$	Per	✓	Manufacturer
ORPHENADRINE CITRATE Tab 100 mg	18.54	100	✓ N	orflex

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Agents for Parkinsonism and Related Disorders

Dopamine Agonists and Related Agents			
AMANTADINE HYDROCHLORIDE	00.04	60	√ Symmetre!
▲ Cap 100 mg	38.24	60	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE ▲ Inj 10 mg per ml, 2 ml ampoule	110.00	5	✓ Movapo
	119.00	3	▼ wovapo
ROMOCRIPTINE MESYLATE	22.00	100	✓ Ano Bromocrintino
₹ Tab 2.5 mg	32.00	100	✓ Apo-Bromocriptine
NTACAPONE ■ Tab 200 mg	22.00	100	✓ Entapone
v	22.00	100	▼ <u>спіаропе</u>
EVODOPA WITH BENSERAZIDE	10.05	100	/ Madamar Danid
Tab dispersible 50 mg with benserazide 12.5 mg		100 100	✓ Madopar Rapid✓ Madopar 62.5
Cap 30 mg with benserazide 12.3 mg		100	✓ Madopar 62.5 ✓ Madopar 125
Cap long-acting 100 mg with benserazide 25 mg		100	✓ Madopar HBS
Cap 200 mg with benserazide 50 mg		100	✓ Madopar 250
EVODOPA WITH CARBIDOPA			
★ Tab 100 mg with carbidopa 25 mg	17 07	100	✓ Kinson
rab 100 mg with carbidopa 20 mg		100	✓ Sinemet
★ Tab long-acting 200 mg with carbidopa 50 mg	37.15	100	✓ Sinemet CR
★ Tab 250 mg with carbidopa 25 mg		100	✓ Sinemet
Kinson Tab 100 mg with carbidopa 25 mg to be delisted 1 Jui			
PRAMIPEXOLE HYDROCHLORIDE			
▲ Tab 0.25 mg	7.20	100	✓ Ramipex
▲ Tab 1 mg		100	✓ Ramipex
ROPINIROLE HYDROCHLORIDE			
▲ Tab 0.25 mg	2.78	100	✓ Apo-Ropinirole
▲ Tab 1 mg	5.00	100	✓ Apo-Ropinirole
▲ Tab 2 mg	7.72	100	✓ Apo-Ropinirole
▲ Tab 5 mg	16.51	100	✓ Apo-Ropinirole
SELEGILINE HYDROCHLORIDE			
₭ Tab 5 mg	22.00	100	✓ Apo-Selegiline
			S29 S29
OLCAPONE			
▲ Tab 100 mg	132.50	100	✓ <u>Tasmar</u>
-			
Anticholinergics			
SENZATROPINE MESYLATE			
Tab 2 mg		60	✓ Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	✓ Cogentin
	190.00	10	✓ Omega
a) Up to 10 inj available on a PSOb) Only on a PSO			

100

✓ Kemadrin

Tab 5 mg7.40

PROCYCLIDINE HYDROCHLORIDE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE - Special Authority see SA1403 below - Retail pharmacy

Wastage claimable

⇒SA1403 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
 - 5.1 The patient is ambulatory; or
 - 5.2 The patient is able to use upper limbs; or
 - 5.3 The patient is able to swallow.

Renewal from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
 - 3.1 The patient is ambulatory; or
 - 3.2 The patient is able to use upper limbs; or
 - 3.3 The patient is able to swallow.

TETRABENAZINE

Tab 25 mg91.10 112 ✓ <u>Motetis</u>

Anaesthetics

LIDOCAINE [LIGNOCAINE]

Local

Gel 2%, tube - Subsidy by endorsement14	.50 30 ml	✓ Xylocaine 2% Jelly
a) Up to 150 ml available on a PSO		
b) Subsidised only if prescribed for urethral or cervical administra	tion and the prescription	is endorsed accordingly.
Gel 2%, 10 ml urethral syringe - Subsidy by endorsement81	.50 10	✓ Pfizer
160	.00 25	✓ Cathejell
a) Up to 5 each available on a PSO		
b) Subsidised only if prescribed for urethral or cervical administra	tion and the prescription	is endorsed accordingly.
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE		
Oral (gel) soln 2%38	.00 200 ml	✓ Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO8	.75 25	✓ Lidocaine-Claris
17	.50 50	
(35	.00)	Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO6	.75 25	✓ Lidocaine-Claris
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO12	.00 5	
(20	.00)	Xylocaine
Inj 1%, 20 ml vial – Up to 5 inj available on a PSO12	.00 5	Lidocaine-Claris
Inj 2%, 20 ml vial – Up to 5 inj available on a PSO12	.00 5	Lidocaine-Claris

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes -				
Subsidy by endorsement	81.50	10	✓	Pfizer
a) Up to 5 each available on a PSO				
b) Subsidised only if prescribed for urethral or cervical a	administration and the	pres	cription is e	endorsed accordingly.

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

Non-opioid Analgesics

For aspirin & chloroform application refer Standard Formulae, page 228

ASI	ΊK	IN
ASI	ΊK	IIN

*	Tab dispersible 300 mg - Up to 30 tab available on a PSO3.90	100	✓ Ethics Aspirin
CA	PSAICIN - Subsidy by endorsement		
	Subsidized only if prescribed for post-harnetic neuralgia or diabetic periph	aral nauronathy	and the prescription is a

Subsidised only if prescribed for post-herpetic neuralgia or diabetic peripheral neuropathy and the prescription is endorsed accordingly.

		Subsidy (Manufacturer's Pri	ce) Subs	Fully Brand or sidised Generic
		\$	Per	✓ Manufacturer
PA	RACETAMOL			_
	Tab 500 mg - blister pack	0.71 7.12	100 1,000	 ✓ Priceline ✓ Paracetamol Pharmacare ✓ Pharmacare ✓ Pharmacy Health
	 a) Maximum of 300 tab per prescription; can be waived b) Up to 30 tab available on a PSO 	by endorsement		
	Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater w annotated accordingly. Pharmacists may anno supports a long-term condition. Maximum of 100 tab per dispensing for non-endorsements.	no do not use com tate the prescription dorsed patients. I	npliance pack on as endorse f quantities p	aging, and the prescription is ed where dispensing history rescribed for more than 100 tab
	(for non-endorsed patients), then dispense in re			
*	Tab 500 mg - bottle pack		1,000	✓ Pharmacare
*	Oral liq 120 mg per 5 mla) Up to 200 ml available on a PSO b) Not in combination	5.35	1,000 ml	✓ <u>Paracare</u>
*	Oral liq 250 mg per 5 ml	5.81	1,000 ml	✓ <u>Paracare Double</u> <u>Strength</u>
	a) Up to 100 ml available on a PSOb) Not in combination			• • •
*	Suppos 125 mg		10	✓ Gacet
*	Suppos 250 mg		10	✓ <u>Gacet</u> ✓ Gacet
* (Pa	Suppos 500 mgracare Suppos 500 mg to be delisted 1 May 2019)	(12.60)	50	Paracare
`	. ,			
	pioid Analgesics			
CO	DEINE PHOSPHATE – Safety medicine; prescriber may dete	, ,		4
	Tab 15 mg		100	✓ PSM
	Tab 30 mg Tab 60 mg		100 100	✓ <u>PSM</u> ✓ PSM
	v	13.30	100	▼ <u>FSIWI</u>
	IYDROCODEINE TARTRATE	0.55	00	A DUO Continue
DIF	Tab long-acting 60 mg	9.55	60	✓ <u>DHC Continus</u>
	NTANYL			
	a) Only on a controlled drug form			
	a) Only on a controlled drug form b) No patient co-payment payable	aguanov		
	a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing free.		10	✓ Boucher and Muir
	a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing fre Inj 50 mcg per ml, 2 ml ampoule	3.56	10 10	✓ Boucher and Muir ✓ Boucher and Muir
	a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing fro lnj 50 mcg per ml, 2 ml ampoule	3.56 9.41		✓ Boucher and Muir
	a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing fre Inj 50 mcg per ml, 2 ml ampoule	3.56 9.41 2.95	10	
	a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing fro lnj 50 mcg per ml, 2 ml ampoule	3.56 9.41 2.95 3.66	10 5	✓ Boucher and Muir✓ Fentanyl Sandoz
	a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing fro lnj 50 mcg per ml, 2 ml ampoule	3.56 9.41 2.95 3.66 6.65	10 5 5	✓ Boucher and Muir ✓ Fentanyl Sandoz ✓ Fentanyl Sandoz

			NER\	OUS SYSTEM
	Subsidy (Manufacturer's Price)) Sub	Fully osidised	Brand or Generic Manufacturer
METHADONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre				- mar - 11 - 11 - 1
d) Extemporaneously compounded methadone will only be i	reimbursed at the ra	te of the c	neapest f	orm available
(methadone powder, not methadone tablets).e) For methadone hydrochloride oral liquid refer Standard F.	ormulae nage 229			
Tab 5 mg		10	✓ M	ethatabs
Oral liq 2 mg per ml		200 ml		iodone
Oral liq 5 mg per ml		200 ml		iodone Forte
Oral liq 10 mg per ml		200 ml	_	iodone Extra Forte
Inj 10 mg per ml, 1 ml		10	✓ Al	FT
MORPHINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre				
Oral liq 1 mg per ml	9.28	200 ml	_	A-Morph
Oral liq 2 mg per ml	16.24	200 ml		A-Morph
Oral liq 5 mg per ml		200 ml		A-Morph
Oral liq 10 mg per ml	27.74	200 ml	✓ <u>R</u> /	A-Morph
MORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre		40		
Tab immediate-release 10 mg		10		evredol rrow-Morphine LA
Tab long-acting 10 mg Tab immediate-release 20 mg		10 10		rrow-Morphine LA evredol
Tab long-acting 30 mg		10		rrow-Morphine LA
Tab long-acting 50 mg		10		rrow-Morphine LA
Tab long-acting 100 mg		10	_	rrow-Morphine LA
Cap long-acting 10 mg		10	_	-Eslon
Cap long-acting 30 mg		10	✓ m	-Eslon
Cap long-acting 60 mg		10		-Eslon
Cap long-acting 100 mg		10		-Eslon
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS	SO6.27	5		BL Morphine
http://www.nan.id.id.)OO 4 :=	_		Sulphate
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a F	∕SU4.47	5	_	BL Morphine
haide management and comments of the control of the	100 470	-		Sulphate
Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a F	′ 5∪4.76	5		BL Morphine
Ini 20 ma normi, 1 ml amnaula . Lie to 5 ini available and 5)CO 640	F		Sulphate
Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a F	-300.19	5		BL Morphine Sulphate
MORPHINE TARTRATE				-aipiiate
INCALL LINE LADIDATE				

MORPHINE TARTRATE

- a) Only on a controlled drug form
- b) No patient co-payment payable
- c) Safety medicine; prescriber may determine dispensing frequency
- Inj 80 mg per ml, 1.5 ml ampoule42.72

✓ DBL Morphine **Tartrate**

5

	Subsidy (Manufacturer's Price)	S	Fully Brand or ubsidised Generic
	\$	Per	✓ Manufacturer
OXYCODONE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing fr	equency		
Tab controlled-release 5 mg	2.63	20	✓ BNM
Tab controlled-release 10 mg	2.76	20	✓ BNM
Tab controlled-release 20 mg		20	✓ BNM
Tab controlled-release 40 mg		20	✓ BNM
Tab controlled-release 80 mg		20	✓ BNM
Cap immediate-release 5 mg		20	✓ <u>OxyNorm</u>
Cap immediate-release 10 mg		20	✓ <u>OxyNorm</u>
Cap immediate-release 20 mg		20	✓ <u>OxyNorm</u>
Oral liq 5 mg per 5 ml		250 ml	✓ OxyNorm
Inj 10 mg per ml, 1 ml ampoule		5	✓ <u>OxyNorm</u>
Inj 10 mg per ml, 2 ml ampoule		5	✓ <u>OxyNorm</u>
Inj 50 mg per ml, 1 ml ampoule		5	✓ <u>OxyNorm</u>
PARACETAMOL WITH CODEINE - Safety medicine; prescribe		-	
* Tab paracetamol 500 mg with codeine phosphate 8 mg	18.21	1,000	✓ Paracetamol +
			Codeine (Relieve)
PETHIDINE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing from	equency		
Tab 50 mg	4.46	10	✓ <u>PSM</u>
Inj 50 mg per ml, 1 ml ampoule - Up to 5 inj available on a I	PSO4.98	5	✓ DBL Pethidine
			<u>Hydrochloride</u>
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a I	PSO5.12	5	✓ DBL Pethidine
			<u>Hydrochloride</u>
TRAMADOL HYDROCHLORIDE			
Tab sustained-release 100 mg	1.55	20	✓ Tramal SR 100
Tab sustained-release 150 mg	2.10	20	✓ Tramal SR 150
Tab sustained-release 200 mg	2.75	20	✓ Tramal SR 200
Cap 50 mg	2.25	100	✓ Arrow-Tramadol
Antidepressants			
0 " IB I I I I I I			
Cyclic and Related Agents			
AMITRIPTYLINE - Safety medicine; prescriber may determine of	dispensina frequency		
Tab 10 mg	1.96	100	✓ Arrow-Amitriptyline
Tab 25 mg	1.52	100	✓ Arrow-Amitriptyline
Tab 50 mg	2.51	100	✓ Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; presci	riber may determine o	dispensi	na frequency
Tab 10 mg	,	100	✓ Apo-Clomipramine
Tab 25 mg		100	✓ Apo-Clomipramine
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Safety medici			
Tab 75 mg		100	✓ Dopress
Cap 25 mg		100	✓ Dopress
Oup 20 mg		100	Dopiess

	Subsidy	Full	,
	(Manufacturer's Price)	Subsidise	
	\$	Per 🗸	Manufacturer
DOXEPIN HYDROCHLORIDE - Subsidy by endorsement			
a) Safety medicine; prescriber may determine dispensing fi	requency		
b) Subsidy by endorsement - Subsidised for patients who			
prescription is endorsed accordingly. Pharmacists may	annotate the prescripti	on as endorse	d where there exists a record
of prior dispensing of doxepin hydrochloride.			
Cap 10 mg	6.30	100	' Anten
Cap 25 mg	6.86		' Anten
Cap 50 mg	8.55	100	['] Anten
IMIPRAMINE HYDROCHLORIDE - Safety medicine; prescribe	r may determine dispe	nsing frequenc	cy
Tab 10 mg			['] Tofranil
· ·	10.96	100	Tofranil
Tab 25 mg	8.80	50	Tofranil
MAPROTILINE HYDROCHLORIDE - Safety medicine; prescrib	er may determine disc	ensina freaue	ncv
Tab 25 mg			Ludiomil
	12.53		Ludiomil
	25.06		Ludiomil
Tab 75 mg	14.01	20	Ludiomil
· ·	21.01	30	Ludiomil
NORTRIPTYLINE HYDROCHLORIDE - Safety medicine; presi	criher may determine o	lisnensina frea	HENCY
Tab 10 mg			Norpress
Tab 25 mg			Norpress
. 45 _5g			р.сос
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	Selective		
PHENELZINE SULPHATE			
* Tab 15 mg	95.00	100	Nardil
TRANYLCYPROMINE SULPHATE			
* Tab 10 mg	22 04	50	Parnate
* Tab To Hig	22.34	30	railiale
Monoamine-Oxidase Type A Inhibitors			
MOCLOBEMIDE			
* Tab 150 mg	6.40	60	Aurorix
	85.10		Apo-Moclobemide
* Tab 300 mg	9.80		Aurorix
	30.70	100	Apo-Moclobemide
			.,
Selective Serotonin Reuptake Inhibitors			
CITALOPRAM HYDROBROMIDE			
* Tab 20 mg	1.52	84	PSM Citalopram
ESCITALOPRAM			
* Tab 10 mg	1.11	28	Escitalopram-
			Apotex
			1
* Tab 20 mg	1.90	28	Escitalopram-
			<u>Apotex</u>

	Subsidy		Fully	
(Manufacturer's Price)	Subsi	idised •	Generic Manufacturer
	\$	Per		Manufacturer
FLUOXETINE HYDROCHLORIDE				
* Tab dispersible 20 mg, scored – Subsidy by endorsement	2.47	30	•	Arrow-Fluoxetine
Subsidised by endorsement				
When prescribed for a patient who cannot swallow we cannot swallow with the control of t	nole tablets or caps	ules and t	ne pr	escription is endorsed
accordingly; or 2) When prescribed in a daily dose that is not a multiple	of 20 ma in which a	aca tha n	racor	intion is deemed to be
endorsed. Note: Tablets should be combined with a				
	sapoulos to lasimato			,g doodo.
* Cap 20 mg	1.99	90	1	Arrow-Fluoxetine
PAROXETINE				
* Tab 20 mg	4.02	90	1	Apo-Paroxetine
SERTRALINE				<u></u>
* Tab 50 mg	3.05	90	1	Arrow-Sertraline
* Tab 100 mg		90		Arrow-Sertraline
Other Antidepressants				
MIRTAZAPINE				
Tab 30 mg	2.63	30	J	Apo-Mirtazapine
Tab 45 mg		30		Apo-Mirtazapine
VENLAFAXINE		00	-	Tipo inii tazapino
* Cap 37.5 mg	6 38	84	1	Enlafax XR
* Cap 75 mg		84		Enlafax XR
* Cap 150 mg		84		Enlafax XR
Antiepilepsy Drugs				
Agents for Control of Status Epilepticus				
CLONAZEPAM - Safety medicine; prescriber may determine disp	ensing frequency			
Inj 1 mg per ml, 1 ml		5	1	Rivotril
DIAZEPAM – Safety medicine; prescriber may determine dispensi				
Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement		5	1	Hospira
a) Up to 5 inj available on a PSO				
b) Only on a PSO				
c) PSO must be endorsed "not for anaesthetic procedure				
Rectal tubes 5 mg - Up to 5 tube available on a PSO		5		Stesolid
Rectal tubes 10 mg - Up to 5 tube available on a PSO	40.87	5	1	Stesolid
PARALDEHYDE				
* Inj 5 ml	1,500.00	5	1	AFT S29
PHENYTOIN SODIUM				
* Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a PS	60 88.63	5	✓	Hospira
* Inj 50 mg per ml, 5 ml ampoule - Up to 5 inj available on a				
PSO	133.92	5	1	Hospira

Brand or

Fully

	Subsidy	` 0.1	Fully	Brand or
	(Manufacturer's Pric	e) Sub Per	sidised •	Generic Manufacturer
	<u> </u>			
Control of Epilepsy				
CARBAMAZEPINE				
* Tab 200 mg	14.53	100	✓ T	egretol
* Tab long-acting 200 mg	16.98	100	✓ T	egretol CR
* Tab 400 mg	34.58	100	✓ T	egretol
* Tab long-acting 400 mg	39.17	100	✓ T	egretol CR
* Oral liq 20 mg per ml	26.37	250 ml	✓ T	egretol
CLOBAZAM - Safety medicine; prescriber may determine dispe				
Tab 10 mg	. ,	50	√ F	risium
CLONAZEPAM - Safety medicine; prescriber may determine di	snensina freatiency			
Oral drops 2.5 mg per ml		10 ml OP	✓ B	livotril
ETHOSUXIMIDE		10 1111 01		
	001.75	200	./7	arontin
Cap 250 mg		200 200 ml		aronun arontin
Oral liq 250 mg per 5 ml		200 1111	• 2	aronun
GABAPENTIN				
Note: Not subsidised in combination with subsidised pregal				
* Cap 100 mg		100	_	po-Gabapentin
* Cap 300 mg		100	_	po-Gabapentin
* Cap 400 mg	5.64	100	✓ <u>A</u>	po-Gabapentin
LACOSAMIDE - Special Authority see SA1125 below - Retail p	oharmacy			
▲ Tab 50 mg	25.04	14	✓ V	impat
▲ Tab 100 mg	50.06	14	✓ V	impat
	200.24	56	✓ V	impat
▲ Tab 150 mg	75.10	14	✓ V	impat
-	300.40	56	✓ V	impat
▲ Tab 200 mg	400.55	56	✓ V	impat

Subeidy

⇒SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

	Subsidy (Manufacturar's Price	20/ 0:-1	Fully	
	(Manufacturer's Pric	ce) Sur Per	sidised •	I Generic Manufacturer
MOTRIGINE	· · · · · · · · · · · · · · · · · · ·			
Tab dispersible 2 mg	6.74	30	1	Lamictal
Tab dispersible 5 mg		30		Lamictal
rab dispersible 5 mg	15.00	56		Arrow-Lamotrigine
Tab dispersible 25 mg		56	_	Logem
rab dispersible 25 mg		30		Arrow-Lamotrigine
	20.40			•
Tab discountible 50 mm	29.09	50		Lamictal
Tab dispersible 50 mg		56		Logem
	34.70			Arrow-Lamotrigine
	47.89			Lamictal
Tab dispersible 100 mg		56		Logem
	59.90			Arrow-Lamotrigine
	79.16		/	Lamictal
VETIRACETAM				
Tab 250 mg	24.03	60	1	Everet
Tab 500 mg		60		Everet
Tab 750 mg		60		Everet
Tab 1,000 mg		60		Everet
Oral lig 100 mg per ml		300 ml OP		Levetiracetam-AFT
, ,,	44.70	300 IIII OF	•	Levelii acelaiii-AF i
IENOBARBITONE				
For phenobarbitone oral liquid refer Standard Formulae, p	age 228			
Tab 15 mg	40.00	500		<u>PSM</u>
Tab 30 mg	40.00	500	/	PSM
HENYTOIN SODIUM				
Tab 50 mg	50 51	200	1	Dilantin Infatab
Cap 30 mg		200		Dilantin
Cap 100 mg		200		Dilantin
, ,				Dilantin
Oral liq 30 mg per 5 ml	22.03	500 ml	•	Diianun
REGABALIN				
Note: Not subsidised in combination with subsidised gaba	pentin			
Cap 25 mg	2.25	56	/	Pregabalin Pfizer
Cap 75 mg	2.65	56	1	Pregabalin Pfizer
Cap 150 mg	4.01	56	1	Pregabalin Pfizer
Cap 300 mg	7.38	56	1	Pregabalin Pfizer
RIMIDONE				
	17.05	100	1	Apo-Primidone
Tab 250 mg				•
	62.00	200	•	Mysoline S29 S29
DDIUM VALPROATE				
Tab 100 mg	13.65	100	1	Epilim Crushable
Tab 200 mg EC	27.44	100	1	Epilim
Tab 500 mg EC		100	_	Epilim
Oral liq 200 mg per 5 ml		300 ml		Epilim S/F Liquid
1 Ur				Epilim Syrup
Inj 100 mg per ml, 4 ml	41.50	1		Epilim IV
, ,			-	-r
'IRIPENTOL – Special Authority see SA1330 on the next pa	•	•	_	
Cap 250 mg	509.29	60	/	Diacomit S29
Powder for oral lig 250 mg sachet	509.29	60	1	Diacomit S29

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

TOPIRAMATE

\blacktriangle	Tab 25 mg	11.07	60	Arrow-Topiramate
	•			✓ Topiramate Actavis
		26.04		✓ Topamax
\blacktriangle	Tab 50 mg	18.81	60	✓ Arrow-Topiramate
	•			✓ Topiramate Actavis
		44.26		✓ Topamax
\blacktriangle	Tab 100 mg	31.99	60	✓ Arrow-Topiramate
	•			✓ Topiramate Actavis
		75.25		✓ Topamax
\blacktriangle	Tab 200 mg	55.19	60	✓ Arrow-Topiramate
	Ÿ			✓ Topiramate Actavis
		129.85		✓ Topamax
\blacktriangle	Sprinkle cap 15 mg	20.84	60	✓ Topamax
\blacktriangle	Sprinkle cap 25 mg	26.04	60	✓ Topamax
VIC	GABATRIN – Special Authority see SA1072 below – Retail	pharmacy		
\blacktriangle	Tab 500 mg		100	✓ Sabril

SA1072 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Fither:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Notes: ``Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages. **Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
continued Both:			

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

Acute Migraine Treatment

ERGOTAMINE TARTRATE WITH CAFFEINE		
Tab 1 mg with caffeine 100 mg31.0	0 100	Cafergot
		✓ Cafergot S29 S29
RIZATRIPTAN		
Tab orodispersible 10 mg5.2	6 30	✓ <u>Rizamelt</u>
SUMATRIPTAN		
Tab 50 mg24.4-	4 100	✓ Apo-Sumatriptan
Tab 100 mg46.2	3 100	✓ Apo-Sumatriptan
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj per		
prescription42.6	7 2 OP	✓ Clustran
		✓ Sun Pharma S29

Prophylaxis of Migraine

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 50 PIZOTIFEN

* Tab 500 mcg......23.21 100

✓ Sandomigran

✓ Sandomigran S29 S29

Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT – Special Authority see SA0987 below – Retail pharmacy

Cap 2 × 80 mg and 1 × 125 mg.....84.00

3 OP

✓ Emend Tri-Pack

⇒SA0987 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

	Subsidy		Fully	Brand or
	(Manufacturer's Price	e)	Subsidised	I Generic
	\$	Per	•	Manufacturer
CYCLIZINE HYDROCHLORIDE				
Tab 50 mg	0.55	10	1	Nausicalm
ŭ	0.59	20	1	Nauzene
Nausicalm to be Sole Supply on 1 April 2019 (Nauzene Tab 50 mg to be delisted 1 April 2019)				
CYCLIZINE LACTATE				
Inj 50 mg per ml, 1 ml	14.95	5	1	Nausicalm
DOMPERIDONE				
* Tab 10 mg	2.25	100	1	Pharmacy Health
ŭ	3.20		1	Prokinex
HYOSCINE HYDROBROMIDE				
* Inj 400 mcg per ml, 1 ml ampoule	46.50	5	1	Hospira
,	93.00	10	✓	Martindale S29
Patch 1.5 mg - Special Authority see SA1387 below - Retail				
pharmacy		2	•	Scopoderm TTS

⇒SA1387 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE		
* Tab 10 mg	100	✓ Metoclopramide Actavis 10
* Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO4.50 ONDANSETRON	10	✓ Pfizer
* Tab 4 mg	50	✓ Apo-Ondansetron
* Tab disp 4 mg0.95	10	✓ Ondansetron ODT-ORLA
* Tab 8 mg4.77	50	✓ Apo-Ondansetron
* Tab disp 8 mg	10	✓ Ondansetron ODT-DRLA
PROCHLORPERAZINE		
* Tab 3 mg buccal5.97	50	
(15.00)		Buccastem
* Tab 5 mg - Up to 30 tab available on a PSO	250	✓ Nausafix
* Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO25.81	10	✓ Stemetil
PROMETHAZINE THEOCLATE		
* Tab 25 mg1.20	10	
(5.59)		Avomine
(Avomine Tab 25 mg to be delisted 1 March 2019)		

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Antipsychotics

General

AMISULPRIDE - Safety medicine; prescriber may determine dis	spensing frequen	су	
Tab 100 mg	4.56	30	✓ Sulprix
Tab 200 mg	14.75	60	✓ Sulprix
Tab 400 mg	27.70	60	✓ Sulprix
Oral liq 100 mg per ml	65.53	60 ml	✓ Solian
ARIPIPRAZOLE - Safety medicine; prescriber may determine of	lispensina freauei	ncv	
Tab 5 mg		30	✓ Aripiprazole Sandoz
Tab 10 mg	17.50	30	✓ Aripiprazole Sandoz
Tab 15 mg	17.50	30	✓ Aripiprazole Sandoz
Tab 20 mg	17.50	30	✓ Aripiprazole Sandoz
Tab 30 mg	17.50	30	✓ Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; pr	escriber may dete	ermine dispen	sing frequency
Tab 10 mg – Up to 30 tab available on a PSO		100	✓ Largactil
Tab 25 mg - Up to 30 tab available on a PSO		100	✓ Largactil
Tab 100 mg - Up to 30 tab available on a PSO		100	✓ Largactil
Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO		10	✓ Largactil
CLOZAPINE – Hospital pharmacy [HP4]			3
Safety medicine; prescriber may determine dispensing frequ	ionov		
Tab 25 mg	•	50	✓ Clozaril
1 ab 25 mg	6.69	30	✓ Clopine
	11.36	100	✓ Clozaril
	13.37	100	✓ Clopine
Tab 50 mg		50	✓ Clopine
1 ab 30 mg	17.33	100	✓ Clopine
Tab 100 mg		50	✓ Clozaril
1 db 100 mg	17.33	00	✓ Clopine
	29.45	100	✓ Clozaril
	34.65	100	✓ Clopine
Tab 200 mg		50	✓ Clopine
140 200 mg	69.30	100	✓ Clopine
Suspension 50 mg per ml		100 ml	✓ Clopine
HALOPERIDOL – Safety medicine; prescriber may determine d		201	
Tab 500 mcg – Up to 30 tab available on a PSO		100	✓ Serenace
Tab 1.5 mg - Up to 30 tab available on a PSO		100	✓ Serenace
Tab 5 mg - Up to 30 tab available on a PSO		100	✓ Serenace
Oral lig 2 mg per ml – Up to 200 ml available on a PSO		100 ml	✓ Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a P		100 1111	✓ Serenace
LEVOMEPROMAZINE HYDROCHLORIDE – Safety medicine;			
Inj 25 mg per ml, 1 ml ampoule		10	✓ Wockhardt
LEVOMEPROMAZINE MALEATE – Safety medicine; prescriber	•		
Tab 25 mg	16.93	100	✓ Nozinan
Tab 100 mg	43.96	100	✓ Nozinan

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	(Manufacturer's Frice)	Per	✓	Manufacturer
ITHIUM CARBONATE - Safety medicine; prescriber may deteri	mine dispensina frea	Henc	, , , , , , , , , , , , , , , , , , ,	
Tab 250 mg		500	_	Lithicarb FC
Tab 400 mg		100	_	Lithicarb FC
Tab long-acting 400 mg		100	_	Priadel
Cap 250 mg		100	_	Douglas
Lithicarb FC Tab 400 mg to be delisted 1 March 2019)	9.42	100	•	Douglas
DLANZAPINE – Safety medicine; prescriber may determine disp				
Tab 2.5 mg		28	_	Zypine
Tab 5 mg		28	_	Zypine
Tab orodispersible 5 mg		28	_	Zypine ODT
Tab 10 mg	1.65	28	_	Zypine
Tab orodispersible 10 mg	2.05	28	•	Zypine ODT
PERICYAZINE - Safety medicine; prescriber may determine disp	pensing frequency			
Tab 2.5 mg		84	✓	Neulactil
ŭ	12.49	100	1	Neulactil
Tab 10 mg	37.34	84	1	Neulactil
J	44.45	100		Neulactil
NIETIADINE Sofaty modicino: proporihar may datarmina diana				- ==-
QUETIAPINE – Safety medicine; prescriber may determine dispe	. ,	90	.1	Quetanal
Tab 25 mg				Quetapel
Tab 100 mg		90	_	Quetapel
Tab 200 mg		90	_	Quetapel
Tab 300 mg	9.60	90	•	<u>Quetapel</u>
RISPERIDONE - Safety medicine; prescriber may determine dis	pensing frequency			
Tab 0.5 mg	1.86	60	✓	<u>Actavis</u>
Tab 1 mg	2.06	60	1	Actavis
Tab 2 mg	2.29	60	1	Actavis
Tab 3 mg	2.50	60	1	Actavis
Tab 4 mg	3.43	60	1	Actavis
Oral liq 1 mg per ml		30 m	1	Risperon
IPRASIDONE - Safety medicine; prescriber may determine disp				
Cap 20 mg		60	1	Zusdone
Oap 20 mg	14.56	00		Zeldox
Zuadana ta ha Cala Sunniy an 1 March 2010	14.50		•	Zeidox
Zusdone to be Sole Supply on 1 March 2019	04.70	60	./	7uodono
Cap 40 mg		60	_	Zusdone
Cap 60 mg		60	_	Zusdone
Cap 80 mg	39.70	60	•	Zusdone
Zeldox Cap 20 mg to be delisted 1 March 2019)				
${\tt UCLOPENTHIXOL\ HYDROCHLORIDE\ - Safety\ medicine;\ pres}$	scriber may determin	e disp	ensing fre	equency
Tab 10 mg	31.45	100	1	Clopixol
Depot Injections				
THEFTHING PECANOATE Of the west to the	and the section of the			
LUPENTHIXOL DECANOATE – Safety medicine; prescriber ma		-		Fl
Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		Fluanxol
Inj 20 mg per ml, 2 ml – Up to 5 inj available on a PSO		5		Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	40.87	5	•	Fluanxol
IALOPERIDOL DECANOATE - Safety medicine; prescriber may	y determine dispensi	ng fre	equency	
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO		5		Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO		5	✓	Haldol Concentrate
			_	Haldol

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price)		Fully ubsidised	Brand or Generic
	\$	Per	✓	Manufacturer
OLANZAPINE - Special Authority see SA1428 below - Retail p				
Safety medicine; prescriber may determine dispensing frequency	,			
Inj 210 mg vial	252.00	1	✓ Z ₁	/prexa Relprevv
Inj 300 mg vial	414.00	1	✓ Z ₁	prexa Relprevv
Inj 405 mg vial	504.00	1	✓ Zy	prexa Relprevv

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing to	frequency		
Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe	435.12	1	✓ Invega Sustenna

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

PIPOTHIAZINE PALMITATE - Subsidy by endorsement

- a) Safety medicine; prescriber may determine dispensing frequency
- b) Subsidised for patients who were taking pipothiazine palmitate prior to 1 August 2014 and the prescription or PSO is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of pipothiazine palmitate.

Inj 50 mg per ml, 1 ml	- Up to 5 inj available on a PSO	.178.48	10	Piportil
Inj 50 mg per ml, 2 ml	- Up to 5 inj available on a PSO	.353.32	10	Piportil

(Piportil Inj 50 mg per ml, 1 ml to be delisted 1 June 2019)

(Piportil Inj 50 mg per ml, 2 ml to be delisted 1 June 2019)

	Subsidy (Manufacturer's Price) \$	Su Per	Fully ubsidised	Brand or Generic Manufacturer
RISPERIDONE – Special Authority see SA1427 below – Retail Safety medicine; prescriber may determine dispensing frequ				
Inj 25 mg vial	,	1	✓ Ri	sperdal Consta
Inj 37.5 mg vial	178.71	1	✓ Ri	sperdal Consta
Inj 50 mg vial	217.56	1	✓ Ri	sperdal Consta
TO CA1427 Special Authority for Subsidy				

SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atvoical antipsychotic agents: and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine: prescriber may determine dispensing frequency

✓ Clopixol

An	Ж	אוכ	lics
-	-		

BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	20.23	100	✓ Orion
* Tab 10 mg	13.16	100	✓ Orion
CLONAZEPAM - Safety medicine; prescriber may determine dis	pensing frequency		
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ <u>Paxam</u>
DIAZEPAM - Safety medicine; prescriber may determine dispen	sing frequency		
Tab 2 mg	15.05	500	✓ Arrow-Diazepam
Tab 5 mg	16.18	500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine dispersional dis	ensing frequency		
Tab 1 mg	9.72	250	✓ <u>Ativan</u>
Tab 2.5 mg	12.50	100	✓ <u>Ativan</u>
OXAZEPAM - Safety medicine; prescriber may determine disper	nsing frequency		
Tab 10 mg	6.17	100	✓ Ox-Pam
Tab 15 mg	8.53	100	✓ Ox-Pam

Multiple Sclerosis Treatments

DIMETHYL FUMARATE - Sp	pecial Authority see SA1559 on the next page – F	Retail pharmacy	
Wastage claimable			
Cap 120 mg	520.00	14	Tecfidera
Cap 240 mg	2,000.00	56	Tecfidera



Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

⇒SA1559 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- a) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- b) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- c) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- d) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week:
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- e) applications must be made by the patient's neurologist or general physician; and
- f) patients must have no previous history of lack of response to dimethyl fumarate; and
- g) patients must have not previously had intolerance to dimethyl fumarate; and
- h) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDSS points:

Subsidy Fully (Manufacturer's Price) Subsidised Per

Brand or Generic Manufacturer

continued...

- a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
- b) 1.0 to 3.0; or
- c) 1.5 to 3.5; or
- d) 2.0 to 4.0: or
- e) 2.5 to 4.5; or
- f) 3.0 to 4.5; or g) 3.5 to 4.5; or
- h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to dimethyl fumarate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

FINGOLIMOD - Special Authority see SA1562 below - Retail pharmacy

Wastage claimable

Cap 0.5 mg......2,200.00 28 **✓ Gilenya**

⇒SA1562 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (helow).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and



Subsidy		Fully	Brand or	
(Manufacturer's Price)	;	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week:
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) patients must have no previous history of lack of response to fingolimod; and
- 7) patients must have not previously had intolerance to fingolimod; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5: or
 - d) 2.0 to 4.0: or
 - e) 2.5 to 4.5: or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to fingolimod; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

NATALIZUMAB - Special Authority see SA1563 below - Retail pharmacy

⇒SA1563 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

NERVOUS SYSTEM

Subsidy (Manufacturer's Price) \$

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

continued...

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria):
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) treatment must be initiated and supervised by a neurologist who is registered in the Tysabri Australasian Prescribing Programme operated by the supplier; and
- 7) patients must have no previous history of lack of response to natalizumab; and
- 8) patients must have not previously had intolerance to natalizumab; and
- 9) a) Patient is JC virus negative, or
 - Patient is JC virus positive and has given written informed consent acknowledging an understanding of the risk of progressive multifocal leucoencephalopathy (PML) associated with natalizumab
- 10) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or

Subsidy (Manufacturer's Price) Fully Subsidised

Per

Brand or Generic Manufacturer

continued...

b) 1.0 to 3.0; or

c) 1.5 to 3.5; or

d) 2.0 to 4.0; or

e) 2.5 to 4.5; or

f) 3.0 to 4.5; or g) 3.5 to 4.5; or

h) 4.0 to 4.5.

- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to natalizumab: or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier.

Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate.

Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

TERIFLUNOMIDE - Special Authority see SA1560 below - Retail pharmacy

Wastage claimable

⇒SA1560 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi		Generic
	Per	•	Manufacturer

continued...

- v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week:
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) patients must have no previous history of lack of response to teriflunomide; and
- 7) patients must have not previously had intolerance to teriflunomide; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0: or
 - c) 1.5 to 3.5: or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5: or
 - f) 3.0 to 4.5; or
 - a) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to teriflunomide: or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

Other Multiple Sclerosis Treatments

⇒SA1564 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).



Subsidy (Manufacturer's Price)

Fully Subsidised Brand or Generic Manufacturer

continued...

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator Phone: 04 460 4990 Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only glatiramer acetate inj 40 mg prefilled syringe will be subsidised if dispensed in a community pharmacy. The other agents will NOT be subsidised if dispensed from a community or hospital pharmacy. Regular supplies will be distributed to all approved patients or their clinicians by courier.

Prescribers must send quarterly prescriptions for approved patients to the MSTAC coordinator.

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 20 mg glatiramer acetate daily or 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. The MSTAC coordinator should be notified of the change and a new prescription provided.

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria):
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week:
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or
 - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

8) patient will not be co-prescribed natalizumab or fingolimod.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment.
 Progression of disability is defined as progress by any of the following EDDSS Points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

GLATIRAMER ACETATE - Special Authority see SA1564	on page 141 - Retail p	harmacy	
Inj 20 mg prefilled syringe - [Xpharm]	2,250.00	28	Copaxone
Inj 40 mg prefilled syringe - No patient co-payment pa	yable2,275.00	12	Copaxone
(Copaxone Inj 20 mg prefilled syringe to be delisted 1 July 2			•
INTERFERON BETA-1-ALPHA - [Xpharm] - Special Author	ority see SA1564 on pa	ge 141	
Inj 6 million iu prefilled syringe	1,170.00	4	✓ Avonex
Injection 6 million iu per 0.5 ml pen injector	1,170.00	4	Avonex Pen
INTERFERON BETA-1-BETA - [Xpharm] - Special Author	ity see SA1564 on pag	e 141	
Inj 8 million iu per 1 ml	1.322.89	15	✓ Betaferon

Sedatives and Hypnotics

LORMETAZEPAM – Safety medicine; prescriber may deter	mine dispensing frequen	СУ	
Tab 1 mg	3.11	30	
· ·	(23.50)		Noctamid
(Noctamid Tab 1 mg to be delisted 1 March 2019)			
MELATONIN - Special Authority see SA1666 on the next p	age - Retail pharmacy		
Tab modified-release 2 mg - No more than 5 tab per da	ay28.22	30	Circadin



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

MIDAZOLAM – Safety medicine; prescriber may determine dispens	sing frequency		
Inj 1 mg per ml, 5 ml ampoule	4.30	10	✓ Midazolam-Claris
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available			
on a PSO	14.90	10	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be er	dorsed for statu	s epilepticu	s use only.
Inj 5 mg per ml, 3 ml ampoule	2.50	5	✓ Midazolam-Claris
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj available or	1		
a PSO	11.90	5	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be er	ndorsed for statu	s epilepticu	s use only.
NITRAZEPAM - Safety medicine; prescriber may determine disper	nsing frequency		
Tab 5 mg	0 1 7	100	✓ Nitrados
PHENOBARBITONE SODIUM - Special Authority see SA1386 bel	ow – Retail pha	macy	
Inj 200 mg per ml, 1 ml ampoule		5	✓ Aspen S29
, Jr. , p	46.20	10	✓ Martindale S29
(Martindale \$29 Inj 200 mg per ml, 1 ml ampoule to be delisted 1 c	lune 2019)		

⇒SA1386 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

TEMAZEPAM – Safety medicine; prescriber may determine	dispensing frequency		
Tab 10 mg	1.27	25	✓ Normison
TRIAZOLAM - Safety medicine; prescriber may determine	dispensing frequency		
Tab 125 mcg	5.10	100	
•	(9.85)		Hypam
Tab 250 mcg	4.10	100	
	(11.20)		Hypam

				-
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
ZOPICLONE – Safety medicine; prescriber may determine dispertable 7.5 mg		500	•	Zopiclone Actavis
Stimulants/ADHD Treatments				
ATOMOXETINE - Special Authority see SA1416 below - Retail	pharmacy			
Cap 10 mg	107.03	28	1	Strattera
Cap 18 mg		28	✓.	Strattera
Cap 25 mg		28	✓.	Strattera
Cap 40 mg		28	1	Strattera
Cap 60 mg		28	1	Strattera
Cap 80 mg		28	1	Strattera
Cap 100 mg		28	✓	Strattera

⇒SA1416 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder) diagnosed according to DSM-IV or ICD 10 criteria; and
- 2 Once-daily dosing; and
- 3 Any of the following:
 - 3.1 Treatment with a subsidised formulation of a stimulant has resulted in the development or worsening of serious adverse reactions or where the combination of subsidised stimulant treatment with another agent would pose an unacceptable medical risk; or
 - 3.2 Treatment with a subsidised formulation of a stimulant has resulted in worsening of co-morbid substance abuse or there is a significant risk of diversion with subsidised stimulant therapy; or
 - 3.3 An effective dose of a subsidised formulation of a stimulant has been trialled and has been discontinued because of inadequate clinical response; or
 - 3.4 Treatment with a subsidised formulation of a stimulant is considered inappropriate because the patient has a history of psychoses or has a first-degree relative with schizophrenia; and
- 4 The patient will not be receiving treatment with atomoxetine in combination with a subsidised formulation of a stimulant, except for the purposes of transitioning from subsidised stimulant therapy to atomoxetine.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: A "subsidised formulation of a stimulant" refers to currently subsidised methylphenidate hydrochloride tablet formulations (immediate-release, sustained-release and extended-release) or dexamfetamine sulphate tablets.

DEXAMFETAMINE SULFATE - Special Authority see SA1149 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab 5 mg20.00 100 ✓ <u>PSM</u>

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist: or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the



(Manu	Subsidy	Fully	Brand or
	facturer's Price)	Subsidised	Generic
	\$ P	Per 🗸	Manufacturer

continued...

last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1150 below - Retail pharmacy

- a) Only on a controlled drug form
- Safety medicine: prescriber may determine dispensing frequency

b) Safety medicine; prescriber may determine dispensing	j irequency		
Tab immediate-release 5 mg	3.20	30	Rubifen
Tab immediate-release 10 mg		30	✓ Ritalin
-			Rubifen
Tab immediate-release 20 mg	7.85	30	Rubifen
Tab sustained-release 20 mg	10.95	30	Rubifen SR
·	50.00	100	Ritalin SR

⇒SA1150 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the

NERVOUS SYSTEM

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1151 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine: prescriber may determine dispensing frequency

b) carety meaning, processing may actermine dispersion	.9		
Tab extended-release 18 mg	58.96	30	Concerta
Tab extended-release 27 mg	65.44	30	Concerta
Tab extended-release 36 mg	71.93	30	Concerta
Tab extended-release 54 mg	86.24	30	Concerta
Cap modified-release 10 mg		30	Ritalin LA
Cap modified-release 20 mg		30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg		30	✓ Ritalin LA
'			

⇒SA1151 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

NERVOUS SYSTEM

	,	ully Brand or	
(Manufact	turer's Price) Subsidi	sed Generic	
	\$ Per	 Manufac 	turer

⇒SA1126 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more: and
- 2 Either:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eve movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects: or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

* Tab 5 mg	4.34	90	✓ Donepezil-Rex
* Tab 10 mg		90	✓ Donepezil-Rex
RIVASTIGMINE - Special Authority see SA1488 bel	ow - Retail pharmacy		
Patch 4.6 mg per 24 hour	90.00	30	✓ Exelon

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Roth:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable

		dispensing nequency	b) Salety medicine, prescriber may determine disp
Suboxone	28	57.40	Tab sublingual 2 mg with naloxone 0.5 mg
✓ Suboxone	28	166.00	Tab sublingual 8 mg with naloxone 2 mg

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

1 Patient is opioid dependent; and

Subsidy (Manufacturer's Pric	e)	Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient received but failed detoxification with buprenorphine with naloxone; and
 - 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
 - 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
 - 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE Tab modified-release 150 mg	11.00	30	✓ <u>Zyban</u>
DISULFIRAM Tab 200 mg	75.57	100	✓ Antabuse
NALTREXONE HYDROCHLORIDE — S Tab 50 mg	Special Authority see SA1408 below – F		✓ Naltraccord

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.



Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.

b) Note: Direct Provision by a pharmacist permitted under the provisions	in Part I of Section A.	
Patch 7 mg - Up to 28 patch available on a PSO16.00	28	<u>Habitrol</u>
Patch 7 mg for direct distribution only - [Xpharm]3.94	7	<u>Habitrol</u>
Patch 14 mg - Up to 28 patch available on a PSO17.59	28	<u>Habitrol</u>
Patch 14 mg for direct distribution only - [Xpharm]4.52	7	<u>Habitrol</u>
Patch 21 mg - Up to 28 patch available on a PSO20.16	28	<u>Habitrol</u>
Patch 21 mg for direct distribution only - [Xpharm]5.18	7	<u>Habitrol</u>
Lozenge 1 mg - Up to 216 loz available on a PSO16.61	216	<u>Habitrol</u>
Lozenge 1 mg for direct distribution only - [Xpharm]	36	<u>Habitrol</u>
Lozenge 2 mg - Up to 216 loz available on a PSO18.20	216	<u>Habitrol</u>
Lozenge 2 mg for direct distribution only - [Xpharm]	36	<u>Habitrol</u>
Gum 2 mg (Fruit) - Up to 384 piece available on a PSO33.69	384	<u>Habitrol</u>
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]8.64	96	<u>Habitrol</u>
Gum 2 mg (Mint) - Up to 384 piece available on a PSO33.69	384	<u>Habitrol</u>
Gum 2 mg (Mint) for direct distribution only - [Xpharm]8.64	96	<u>Habitrol</u>
Gum 4 mg (Fruit) - Up to 384 piece available on a PSO38.95	384	<u>Habitrol</u>
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]10.01	96	<u>Habitrol</u>
Gum 4 mg (Mint) - Up to 384 piece available on a PSO38.95	384	<u>Habitrol</u>
Gum 4 mg (Mint) for direct distribution only - [Xpharm]10.01	96	<u>Habitrol</u>

VARENICLINE TARTRATE - Special Authority see SA1771 below - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.

Tab 0.5 mg × 11 and 1 mg × 42	25.64	53 OP	✓ Varenicline Pfizer
Tab 1 mg		56	✓ Varenicline Pfizer
v	67.74	28	✓ Champix
	135.48	56	✓ Champix
Tab 0.5 mg × 11 and 1 mg × 14	60.48	25 OP	✓ Champix

SA1771 | Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking;
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or

NERVOUS SYSTEM

continued...

- 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not used funded varenicline in the last 12 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 The patient has not used funded varenicline in the last 12 months; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 12 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see SA1667 below

Inj 25 mg vial	271.35 1	✓ Ribomustin
Inj 100 mg vial	1,085.38 1	✓ Ribomustin
Inj 1 mg for ECP	11.40 1 mg	✓ Baxter

⇒SA1667 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 All of the following:
 - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
 - 3.2.2 The patient has not received prior bendamustine therapy; and
 - 3.2.3 Fither:
 - 3.2.3.1 Both:
 - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Either:
 - 2.1 Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	;	Subsidised	Generic
\$	Per	1	Manufacturer

- 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.
 Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

macrogiobulinaemia.			
BUSULFAN - PCT - Retail pharmacy-Specialist			
Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			
Inj 10 mg per ml, 5 ml vial	15.07	1	 DBL Carboplatin
, -,	20.00		✓ Carboplatin Ebewe
Inj 10 mg per ml, 15 ml vial	14.05	1	✓ DBL Carboplatin
	19.50		✓ Carbaccord
	22.50		 Carboplatin Ebewe
Inj 10 mg per ml, 45 ml vial	32.59	1	DBL Carboplatin
	48.50		 Carbaccord
	50.00		 Carboplatin Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
(DBL Carboplatin Inj 10 mg per ml, 5 ml vial to be delisted 1 Ma (Carboplatin Ebewe Inj 10 mg per ml, 5 ml vial to be delisted 1 (DBL Carboplatin Inj 10 mg per ml, 15 ml vial to be delisted 1 M (Carbaccord Inj 10 mg per ml, 15 ml vial to be delisted 1 March (Carboplatin Ebewe Inj 10 mg per ml, 15 ml vial to be delisted CARMUSTINE – PCT only – Specialist	March 2019) March 2019) n 2019)		
Inj 100 mg vial	522.00	1	✓ BiCNU
Inj 100 mg for ECP		100 mg OP	✓ Baxter
		100 mg Oi	Dantei
CHLORAMBUCIL – PCT – Retail pharmacy-Specialist	00.00	05	()
Tab 2 mg	29.06	25	Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial		1	DBL Cisplatin
	15.00		Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial		1	✓ DBL Cisplatin
	21.00		Cisplatin Ebewe
Inj 1 mg for ECP	0.25	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE			
Tab 50 mg - PCT - Retail pharmacy-Specialist	79.00	50	✓ Endoxan S29
	158.00	100	✓ Procytox S29
Wastage claimable			
Inj 1 g vial - PCT - Retail pharmacy-Specialist	35.65	1	✓ Endoxan
, ,	127.80	6	✓ Cytoxan
Inj 2 g vial - PCT only - Specialist	71.25	1	✓ Endoxan
Inj 1 mg for ECP - PCT only - Specialist		1 mg	✓ Baxter
IFOSFAMIDE - PCT only - Specialist		•	
Inj 1 g	96.00	1	✓ Holoxan
Inj 2 g		1	✓ Holoxan
Inj 1 mg for ECP		1 mg	✓ Baxter
LOMUSTINE – PCT – Retail pharmacy-Specialist		و	
Cap 10 mg	132.50	20	✓ CeeNU
Cap 40 mg		20	✓ CeeNU
σαρ το πιχ		20	- 000110

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy anufacturer's Price) \$	Sul Per	Fully Brand or bsidised Generic ✓ Manufacturer
40.70	25	✓ Alkeran
67.80	1	✓ Alkeran
25.01	1	Oxaliplatin Actavis 100
110.00		✓ Oxaliplatin Ebewe
46.32	1	✓ Oxaliccord
0.48	1 mg	✓ Baxter
CBS	1	✓ Bedford S29
		✓ THIO-TEPA S29
		✓ Tepadina S29
CBS	1	✓ Tepadina S29
CBS		1

		ialist – Special Authority see SA1467 below	AZACITIDINE - PCT only - Special
Azacitidine Dr	1	139.00	Inj 100 mg vial
Reddy's			, 3
✓ Vidaza		605.00	
✓ Baxter	1 ma	1.53	Ini 1 ma for ECP

⇒SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome: or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- 4 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

	Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
	\$	Per	•	Manufacturer
CALCIUM FOLINATE				
Tab 15 mg - PCT - Retail pharmacy-Specialist	104 26	10	1	DBL Leucovorin
Tab 10 mg 1 01 Trotal pharmacy operation	10 1.20			Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17.10	5	1	Hospira
Inj 10 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialis:		1		Calcium Folinate
ing to mg per mi, 5 mi viai – POT – Hetali pharmacy-Specialis	14.55	'	•	Sandoz
lei 50 mm. BOT. Beteil abanna an Occasiolist	40.05	-	,	
Inj 50 mg - PCT - Retail pharmacy-Specialist	18.25	5	•	Calcium Folinate
				Ebewe
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	7.30	1	•	Calcium Folinate
				Sandoz
Inj 100 mg - PCT only - Specialist	7.33	1	✓	Calcium Folinate
				Ebewe
Inj 300 mg - PCT only - Specialist	22.51	1	1	Calcium Folinate
, , , ,				Ebewe
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	20.95	1	1	Calcium Folinate
ing to mg por mi, so mi viai i o r only opposition	20.00	•		Sandoz
Inj 1 g - PCT only - Specialist	67 51	1	1	Calcium Folinate
IIIJ 1 g = 1 01 offiy = opecialist	07.51	'	•	Ebewe
Ini 10 man man and 100 mal viral DOT and a Consciolist	00.00			
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	60.00	1	•	Calcium Folinate
			_	Sandoz
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	•	Baxter
CAPECITABINE - Retail pharmacy-Specialist				
Tab 150 mg	11.15	60	1	Brinov
Tab 500 mg	62.28	120	1	Brinov
CLADRIBINE - PCT only - Specialist				
Inj 1 mg per ml, 10 ml	5 240 72	7	1	Leustatin
Inj 10 mg for ECP		, 10 mg C		Baxter
, 0	/ 49.90	io ilig c)F •	Daxiei
CYTARABINE				
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specialis	t400.00	5	•	Pfizer
Inj 100 mg per ml, 20 ml vial – PCT – Retail				
pharmacy-Specialist	41.36	1	✓	Pfizer
Inj 1 mg for ECP - PCT only - Specialist	0.25	10 mg	/	Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Specialis	t80.00 1	00 mg	OP 🗸	Baxter
LUDARABINE PHOSPHATE				
Tab 10 mg - PCT - Retail pharmacy-Specialist	412.00	20	1	Fludara Oral
Inj 50 mg vial – PCT only – Specialist		5		Fludarabine Ebewe
Inj 50 mg for ECP – PCT only – Specialist		50 mg (Baxter
	105.00	ou my c)F •	Daxiei
LUOROURACIL				
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist		1		Fluorouracil Ebewe
Inj 50 mg per ml, 50 ml vial – PCT only – Specialist		1		Fluorouracil Ebewe
Inj 50 mg per ml, 100 ml vial - PCT only - Specialist		1	•	Fluorouracil Ebewe
Inj 1 mg for ECP - PCT only - Specialist	0.66	100 m	g 🗸	Baxter
(Fluorouracil Ebewe Inj 50 mg per ml, 50 ml vial to be delisted 1 Ma	arch 2019)			
, , ,	,			

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
GEMCITABINE HYDROCHLORIDE – PCT only – Specialist				
Inj 1 g, 26.3 ml vial		1	✓	DBL Gemcitabine
lnj 1 g	15.89	1	✓	Gemcitabine Ebewe
	349.20		1	Gemzar
Inj 200 mg	8.36	1	✓	Gemcitabine Ebewe
	78.00		✓	Gemzar
Inj 1 mg for ECP	0.02	1 mg	✓	Baxter
(Gemcitabine Ebewe Inj 200 mg to be delisted 1 June 2019)				
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist				
Inj 20 mg per ml, 5 ml vial	71.44	1	•	Irinotecan Actavis 100
	100.00		1	Irinotecan-Rex
Inj 1 mg for ECP	0.19	1 mg	1	Baxter
MERCAPTOPURINE		·		
Tab 50 mg - PCT - Retail pharmacy-Specialist	49 41	25	1	Puri-nethol
Oral suspension 20 mg per ml – Retail pharmacy-Specialist		_0	•	i wii iiowioi
Special Authority see SA1725 below		00 ml (OP 🗸	Allmercap

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

METHOTREXATE

IVIL	MOTIENATE		
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist8.05 Trexate to be Sole Supply on 1 April 2019	90	✓ Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist31.75 Trexate to be Sole Supply on 1 April 2019	90	✓ Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist47.50	5	✓ Hospira
*	Inj 7.5 mg prefilled syringe14.61	1	✓ Methotrexate Sandoz
*	Inj 10 mg prefilled syringe14.66	1	Methotrexate Sandoz
*	Inj 15 mg prefilled syringe14.77	1	Methotrexate Sandoz
*	Inj 20 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 25 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 30 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist30.00	5	✓ <u>DBL Methotrexate</u> Onco-Vial
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist45.00	1	✓ DBL Methotrexate Onco-Vial
*	Inj 100 mg per ml, 10 ml – PCT – Retail pharmacy-Specialist25.00 Inj 100 mg per ml, 50 ml vial – PCT – Retail	1	✓ Methotrexate Ebewe
-,-	pharmacy-Specialist	1	✓ Methotrexate Ebewe
*		1 mg	✓ Baxter
	Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist4.73	5 mg OP	✓ Baxter
	and the state of t	5 mg 51	-4/101

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully osidised	Brand or Generic Manufacturer	
PEMETREXED - PCT only - Specialist - Special Authority see	SA1679 below				_
Inj 100 mg vial	60.89	1	√ Jı	uno Pemetrexed	
Inj 500 mg vial	217.77	1	√ Jı	uno Pemetrexed	
Inj 1 mg for ECP	0.55	1 mg	✓ B	axter	

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Roth:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

- All of the following:
 - 1 No evidence of disease progression; and

THIOGUANINE - PCT - Retail pharmacy-Specialist

- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

Tab 40 mg	126.31	25	✓ Lanvis	
Other Cytotoxic Agents				
AMSACRINE - PCT only - Specialist				
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	✓ Amsidine S29	
Inj 75 mg	1,250.00	5	✓ AmsaLyo S29	
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharm	nacy-Specialist			
Cap 0.5 mg	CBS	100	✓ Agrylin S29	
· · · ·			✓ Teva S29	
ARSENIC TRIOXIDE - PCT only - Specialist				
Inj 10 mg	4,817.00	10	✓ AFT S29	

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price	e) Sub	Fully sidised	Brand or Generic	
	\$	Per	✓	Manufacturer	
BLEOMYCIN SULPHATE - PCT only - Specialist					
Inj 15,000 iu, vial	161.01	1	✓ D	BL Bleomycin	
				Sulfate	
Inj 1,000 iu for ECP	12.45	1,000 iu	✓ B	Baxter	
BORTEZOMIB - PCT only - Specialist - Special Authority see \$	SA1576 below				
Inj 3.5 mg vial	1,892.50	1	✓ V	/elcade	
Inj 1 mg for ECP	594.77	1 mg	✓ B	Baxter	

⇒SA1576 Special Authority for Subsidy

Initial application — (Treatment naive multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient has treatment-naive symptomatic multiple myeloma; or
 - 1.2 The patient has treatment-naive symptomatic systemic AL amyloidosis *; and
- 2 Maximum of 9 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Relapsed/refractory multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has relapsed or refractory multiple myeloma; or
 - 1.2 The patient has relapsed or refractory systemic AL amyloidosis *; and
- 2 The patient has received only one prior front line chemotherapy for multiple myeloma or amyloidosis; and
- 3 The patient has not had prior publicly funded treatment with bortezomib; and
- 4 Maximum of 4 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Relapsed/refractory multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 The patient's disease obtained at least a partial response from treatment with bortezomib at the completion of cycle 4; and
- 2 Maximum of 4 further treatment cycles (making a total maximum of 8 consecutive treatment cycles).

Notes: Responding relapsed/refractory multiple myeloma patients should receive no more than 2 additional cycles of treatment beyond the cycle at which a confirmed complete response was first achieved. A line of therapy is considered to comprise either:

- a) a known therapeutic chemotherapy regimen and supportive treatments; or
- b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments.

Refer to datasheet for recommended dosage and number of doses of bortezomib per treatment cycle.

COLASPASE [L-ASPARAGINASE] - POT OTHY - Specialist			
Inj 10,000 iu	102.32	1	✓ Leunase
Inj 10,000 iu for ECP		10,000 iu OP	✓ Baxter
DACARBAZINE - PCT only - Specialist			
Inj 200 mg vial	58.06	1	DBL Dacarbazine
, ,	580.60	10	 Dacarbazine
			APP S29
Inj 200 mg for ECP	58.06	200 mg OP	✓ Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist			
Inj 0.5 mg vial	166.75	1	✓ Cosmegen
Inj 0.5 mg for ECP	166.75	0.5 mg OP	✓ Baxter

	Subsidy		Fully	
	(Manufacturer's Pri \$	ce) Subs Per	sidised	I Generic Manufacturer
	Ψ	FEI		Manuacturei
DAUNORUBICIN - PCT only - Specialist				
Inj 2 mg per ml, 10 ml		1		Pfizer
Inj 20 mg for ECP	130.00	20 mg OP	•	Baxter
DOCETAXEL - PCT only - Specialist				
Inj 10 mg per ml, 2 ml vial	12.40	1	/	DBL Docetaxel
Inj 20 mg	48.75	1		Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial		1		DBL Docetaxel
Inj 80 mg	195.00	1	1	Docetaxel Sandoz
Inj 1 mg for ECP	0.55	1 mg	/	Baxter
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist				
Inj 2 mg per ml, 5 ml vial		1	1	Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial	11.50	1	1	Doxorubicin Ebewe
, •,	17.00		1	Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial	23.00	1	1	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	1	Doxorubicin Ebewe
	65.00		1	Arrow-Doxorubicin
Inj 1 mg for ECP	0.29	1 mg	1	Baxter
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist		•		
Inj 2 mg per ml, 5 ml vial	25.00	1	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1		Epirubicin Ebewe
Inj 2 mg per ml, 50 ml vial		1		Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1		Epirubicin Ebewe
Inj 1 mg for ECP		1 mg		Baxter
(Epirubicin Ebewe Inj 2 mg per ml, 50 ml vial to be delisted 1		9		
ETOPOSIDE				
Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	1	Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10		Vepesid
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Spe		1		Rex Medical
Inj 1 mg for ECP – PCT only – Specialist		1 mg		Baxter
, , ,		ring	-	Duxio
ETOPOSIDE PHOSPHATE – PCT only – Specialist	40.00	1	,	F4
Inj 100 mg (of etoposide base)		•		Etopophos
Inj 1 mg (of etoposide base) for ECP	0.47	1 mg	•	Baxter
HYDROXYUREA - PCT - Retail pharmacy-Specialist			_	
Cap 500 mg	31.76	100		Hydrea
IDARUBICIN HYDROCHLORIDE				
Inj 5 mg vial - PCT only - Specialist	93.00	1	1	Zavedos
Inj 10 mg vial - PCT only - Specialist	198.00	1	1	Zavedos
Inj 1 mg for ECP - PCT only - Specialist	21.84	1 mg	1	Baxter
LENALIDOMIDE – Retail pharmacy-Specialist – Special Auti Wastage claimable	hority see SA1468 bel	low		
Cap 10 mg	6.207.00	21	1	Revlimid
Cap 15 mg		21		Revlimid
Cap 25 mg		21		Revlimid
	,0=00			
⇒SA1468 Special Authority for Subsidy				

Initial application — (Relapsed/refractory disease) only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)) 5	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Either
 - 2.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 2.2 Both:
 - 2.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 2.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 3 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

М	FS	NΔ

Tab 400 mg - PCT - Retail pharmacy-Specialist	273.00	50	Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist	407.50	50	Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist.	161.25	15	Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialis	t370.35	15	Uromitexan
Inj 1 mg for ECP - PCT only - Specialist	2.69	100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist		-	
Inj 5 mg vial	204.08	1	✓ Arrow
Inj 1 mg for ECP	42.04	1 mg	✓ Baxter
		9	24
MITOZANTRONE – PCT only – Specialist	07.50	4	/ Mitagantyona Ehaura
Inj 2 mg per ml, 10 ml vial		4	✓ Mitozantrone Ebewe
Inj 1 mg for ECP	5.51	1 mg	✓ Baxter
PACLITAXEL - PCT only - Specialist			
Inj 30 mg	47.30	5	Paclitaxel Ebewe
Inj 100 mg	20.00	1	Paclitaxel Ebewe
	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
, 0	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg	35.35	1	✓ Paclitaxel Ebewe
,	275.00		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.19	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see SA132		-	
Inj 3,750 IU per 5 ml		1	✓ Oncaspar S29
			•

⇒SA1325 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Sub (Manufactu		Fully	Brand or Generic	
`	\$ Per	✓	Manufacturer	

continued...

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and
- 3 Treatment is with curative intent.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and
- 3 Treatment is with curative intent.

PENTOSTATIN [DEOXYCOFORMYCIN] – PCT only – Specialist Inj 10 mgCBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE – PCT – Retail pharmacy-Specialist		4 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
Cap 50 mg980.00	50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA1741 below - Retail pharmacy		
Cap 5 mg10.20	5	✓ Orion
		<u>Temozolomide</u>
Cap 20 mg18.30	5	✓ Orion
, •		Temozolomide
		✓ Temizole 20 S29
Cap 100 mg40.20	5	✓ Orion
		Temozolomide
Cap 140 mg56.00	5	✓ Orion
3		Temozolomide
Cap 250 mg96.80	5	✓ Orion
		Temozolomide

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special	Authority see SA1124 below		
Cap 50 mg	378.00	28	✓ Thalomid
Cap 100 mg	756.00	28	✓ Thalomid

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN		
Cap 10 mg - PCT - Retail pharmacy-Specialist479.50	100	✓ Vesanoid
VINBLASTINE SULPHATE		
Inj 1 mg per ml, 10 ml vial - PCT - Retail pharmacy-Specialist 186.46	5	✓ Hospira
Inj 1 mg for ECP - PCT only - Specialist4.14	1 mg	✓ Baxter
VINCRISTINE SULPHATE		
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	✓ DBL Vincristine
		Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist85.61	5	DBL Vincristine
		Sulfate
Inj 1 mg for ECP - PCT only - Specialist11.30	1 mg	✓ Baxter

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised •	
VINORELBINE - PCT only - Specialist	· · · · · · · · · · · · · · · · · · ·			
Inj 10 mg per ml, 1 ml vial	8.00	1	1	Navelbine
ing to mg per mi, i mi viai	42.00	'	_	Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial		1	_	Navelbine
,	210.00	•	_	Vinorelbine Ebewe
Inj 1 mg for ECP		1 mg	1	Baxter
Protein-tyrosine Kinase Inhibitors DASATINIB – [Xpharm] – Special Authority see SA0976 below				
Tab 20 mg	3 774 06	60	1	Sprycel
Tab 50 mg		60		Sprycel
Tab 70 mg		60		Sprycel
Tab 100 mg		30		Sprycel
⇒SA0976 Special Authority for Subsidy	0,214.20	00	•	opi yeei

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 **PHARMAC** Facsimile: (04) 916 7571

Email: cmlgistcoordinator@pharmac.govt.nz PO Box 10 254

Wellington

Special Authority criteria for CML - access by application

- a) Funded for patients with diagnosis (confirmed by a haematologist) of a chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase.
- b) Maximum dose of 140 mg/day for accelerated or blast phase, and 100 mg/day for chronic phase CML.
- c) Subsidised for use as monotherapy only.
- d) Initial approvals valid seven months.
- e) Subsequent approval(s) are granted on application and are valid for six months. The first reapplication (after seven months) should provide details of the haematological response. The third reapplication should provide details of the cytogenetic response after 14-18 months from initiating therapy. All other reapplications should provide details of haematological response, and cytogenetic response if such data is available. Applications to be made and subsequent prescriptions can be written by a haematologist or an oncologist.

Note: Dasatinib is indicated for the treatment of adults with chronic, accelerated or blast phase CML with resistance or intolerance to prior therapy including imatinib.

Guideline on discontinuation of treatment for patients with CML

- a) Prescribers should consider discontinuation of treatment if, after 6 months from initiating therapy, a patient did not obtain a haematological response as defined as any one of the following three levels of response:
 - 1) complete haematologic response (as characterised by an absolute neutrophil count (ANC) > 1.5 × 10⁹/L, platelets > 100 × 109/L, absence of peripheral blood (PB) blasts, bone marrow (BM) blasts < 5% (or FISH Ph+ 0-35%) metaphases), and absence of extramedullary disease); or
 - 2) no evidence of leukaemia (as characterised by an absolute neutrophil count (ANC) > 1.0×10^9 /L, platelets > 20×10^9 /L, 10⁹/L, absence of peripheral blood (PB) blasts, bone marrow (BM) blasts < 5% (or FISH Ph+ 0-35% metaphases), and absence of extramedullary disease); or
 - 3) return to chronic phase (as characterised by BM and PB blasts < 15%, BM and PB blasts and promyelocytes < 30%, PB basophils < 20% and absence of extramedullary disease other than spleen and liver).
- b) Prescribers should consider discontinuation of treatment if, after 18 months from initiating therapy, a patient did not obtain a major cytogenetic response defined as 0-35% Ph+ metaphases.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
ERLOTINIB - Retail pharmacy-Specialist - Special Authority see	e SA1653 below			
Tab 100 mg	764.00	30	✓.	Tarceva
Tab 150 mg	1,146.00	30	✓.	Tarceva
- CA4CEO Consist Authority for Cubaldy				

SA1653 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB - Retail pharmacy-Specialist - Special Authority see SA1654 below Tab 250 mg1,700.00 ✓ Iressa

⇒SA1654 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- - 2.1 Patient is treatment naive; or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
 - 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
 - 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

IMATINIB MESII ATE

Note: Imatinib-AFT is not a registered for the treatment of Gastro Intestinal Stromal Tumours (GIST). The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule.

Tab 100 mg - [Xpharm] - Special Authority see SA1460

	below2,40	0.00 6	60 🗸	Glivec
*	Cap 100 mg		60	Imatinib-AFT
*	Cap 400 mg19	7.50 3	80	Imatinib-AFT

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz, and prescriptions should be sent to:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

The CML/GIST Co-ordinator Phone: (04) 460 4990 PHARMAC Facsimile: (04) 916 7571

PO Box 10, 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

Special Authority criteria for GIST – access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA1191 below - Retail pharmacy 70 Tykerb

⇒SA1191 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient has not previously received trastuzumab treatment for HER 2 positive metastatic breast cancer; and
 - 1.3 Lapatinib not to be given in combination with trastuzumab; and
 - 1.4 Lapatinib to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2.2 The patient started trastuzumab for metastatic breast cancer but discontinued trastuzumab within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on trastuzumab; and
 - 2.4 Lapatinib not to be given in combination with trastuzumab; and
 - 2.5 Lapatinib to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology);
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 on the next page - Retail pharmacy

Wastage claimable 120 ✓ Tasigna 120 ✓ Tasigna

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Either:
 - 2.1 Patient has documented CML treatment failure* with imatinib; or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day: and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PAZOPANIB - Special Authority see SA1190 below - Retail pharmacy

Tab 200 mg	1,334.70	30	✓ Votrient
Tab 400 mg	2,669.40	30	✓ Votrient

⇒SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
RUXOLITINIB – Special Authority see SA1753 below – Retail pt Wastage claimable	narmacy			
Tab 5 mg	2 500 00	56	√ .i	lakavi
Tab 15 mg	,	56	_	akavi
Tab 20 mg	,	56	√ J	akavi

⇒SA1753 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA1266 below - Retail pharmacy

Cap 12.5 mg	38 28	✓ Sutent
Cap 25 mg		✓ Sutent
Cap 50 mg		✓ Sutent

⇒SA1266 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

specialist. Approvals valid for 3 months for applications meeting the following criteria:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and 2 Fither:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 82

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA1767 below

Wastage claimable

Tab 250 mg4,276.19 120 ✓ Zytiga

⇒SA1767 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Fither:

Subsidy	Fully		Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
- 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

BICALUTAMIDE Tab 50 mg	3.80	28	✓ Binarex
FLUTAMIDE - Retail pharmacy-Specialist			
Tab 250 mg	16.50	30	✓ Flutamide
			Mylan S29
	55.00	100	✓ Flutamin
MEGESTROL ACETATE - Retail pharmacy-Specialist			
Tab 160 mg	63.53	30	✓ Apo-Megestrol
OCTREOTIDE			
Inj 50 mcg per ml, 1 ml vial		5	✓ DBL Octreotide
Inj 100 mcg per ml, 1 ml vial	18.69	5	✓ DBL Octreotide
Inj 500 mcg per ml, 1 ml vial	72.50	5	✓ DBL Octreotide
OCTREOTIDE LAR (SOMATOSTATIN ANALOGUE) - S		16 below -	Retail pharmacy
Inj LAR 10 mg prefilled syringe	1,772.50	1	 Sandostatin LAR
Inj LAR 20 mg prefilled syringe		1	Sandostatin LAR
Inj LAR 30 mg prefilled syringe	2,951.25	1	Sandostatin LAR

⇒SA1016 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	✓	

continued...

specialist. Approvals valid for 3 months for applications meeting the following criteria:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed: or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

TAMOXIFEN CITRATE

	Tab 10 mg	11.75	60	✓ Tamoxifen Sandoz
	·	19.50	100	✓ Genox
	Tamoxifen Sandoz to be Sole Supply on 1 April 2019			
*	Tab 20 mg	5.60	60	✓ Tamoxifen Sandoz
	-	9.33	100	✓ Genox

Tamoxifen Sandoz to be Sole Supply on 1 April 2019

(Genox Tab 10 mg to be delisted 1 April 2019)

(Genox Tab 20 mg to be delisted 1 April 2019)

	Subsidy (Manufacturer's Price)	Subs Per	Fully sidised	Brand or Generic Manufacturer
Aromatase Inhibitors				
ANASTROZOLE * Tab 1 mg	5.04	30	✓ <u>R</u>	<u>olin</u>
* Tab 25 mg	14.50	30	✓ <u>P</u>	fizer Exemestane
LETROZOLE * Tab 2.5 mg	4.68	30	✓ <u>L</u>	<u>etrole</u>
Immunosuppressants				
Cytotoxic Immunosuppressants				
AZATHIOPRINE – Retail pharmacy-Specialist # Tab 25 mg # Tab 50 mg # Inj 50 mg vial	10.58 60.00 25.00 25.00 187.25 16	100 100 1 50 100 5 ml OP swallow ta	✓ <u>in</u> ✓ <u>in</u> ✓ <u>c</u>	nuran nuran nuran elicept elicept elicept nd capsules, and when

Fusion Proteins

ETANERCEPT – Special Authority see SA1620 below	 Retail pharmacy 		
Inj 25 mg	799.96	4	Enbrel
Inj 50 mg autoinjector	1,599.96	4	Enbrel
Inj 50 mg prefilled syringe	1,599.96	4	Enbrel

⇒SA1620 Special Authority for Subsidy

Initial application — (juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient diagnosed with Juvenile Idiopathic Arthritis (JIA); and
 - 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufactu 	rer

continued...

- 2.5 Both:
 - 2.5.1 Either:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis: or

2 All of the following:

- 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
- 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold: or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Fither:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints;
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Fither:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plague psoriasis; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Fither:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

continued...

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm: Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for psoriatic arthritis; and
- 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for psoriatic arthritis: or

2 All of the following:

- 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
- 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
- 2.4 Fither:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 4 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a named specialist or rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	,	Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment: and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Fither:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Renewal — **(psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 4 doses.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or

ANTITUVMOCVTE CLOPI II IN /EQUINE) DOT only Specialist

- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Immune Modulators

ANTITITIMOCTTE GLOBOLIN (EQUINE) - FOT ONLY -	Specialist		
Inj 50 mg per ml, 5 ml	2,351.25	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PC	T only – Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	OncoTICE

Monoclonal Antibodies

ADALIMUMAB - Special Authority see SA1742 below - Ref	tail pharmacy		
Inj 20 mg per 0.4 ml prefilled syringe	'	2	✓ Humira
Inj 40 mg per 0.8 ml prefilled pen	•	2	✓ HumiraPen
Inj 40 mg per 0.8 ml prefilled syringe	1,599.96	2	Humira

⇒SA1742 Special Authority for Subsidy

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis: or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:

Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
- 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold: or
- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection;
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Fither
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for psoriatic arthritis; and
 - 1.2 Either
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

4 0

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for juvenile idiopathic arthritis; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient diagnosed with JIA; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
- 2.5 Both:
 - 2.5.1 Either:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Fither:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); and
- $3\,$ A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's disease.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 4 doses.

Note: Note: Indications marked with * are unapproved indications.

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and

Subsidy	Fu	ly Brand or	
(Manufacturer's Price)	Subsidise	ed Generic	
\$	Per	Manufacturer	

continued...

2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (**rheumatoid arthritis**) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Fither:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 Fither:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab;

O

- 2.1.2 CDAI score is 150 or less; or
- 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 Either:
 - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
	Per 🗸	Manufacturer

continued...

- 2.1.2 PCDAI score is 15 or less; or
- 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value: and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — **(psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Fither:

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a named specialist or rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 4 doses.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2 The patient has a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT – Special Authority see SA1772 below – Retail pharmacy

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
CETUXIMAB - PCT only - Specialist - Special Authority see SA	1697 below			
Inj 5 mg per ml, 20 ml vial	364.00	1	✓	Erbitux
Inj 5 mg per ml, 100 ml vial	1,820.00	1	✓	Erbitux
Inj 1 mg for ECP	3.82	1 mg	/	Baxter

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB - PCT only	 Special Authority s 	see SA1778 below
-----------------------	---	------------------

Inj 100 mg	806.00	1	Remicade
Inj 1 mg for ECP	8.29	1 mg	✓ Baxter

⇒SA1778 Special Authority for Subsidy

Initial application — (Graft vs host disease) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis: or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis: or
 - 2.4 Severe ocular inflammation: or
 - 2.5 Chronic ocular inflammation: or
 - 2.6 Crohn's disease (adults): or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Chron's disease: or
 - 2.9 Severe fulminant ulcerative colitis; or
 - 2.10 Severe ulcerative colitis; or
 - 2.11 Plague psoriasis; or
 - 2.12 Neurosarcoidosis: or
 - 2.13 Severe Behcet's disease.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
	Per	1	Manufacturer

continued...

toxicity or intolerance.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for psoriatic arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for psoriatic arthritis.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
- 2 Any of the following:
 - 2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptomst; or
 - 2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2 Any of the following:
 - 2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or

Subsidy (Manufacturer's Price)	Fu Subsidis	,	
\$	Per	 Manufacturer 	

continued...

- 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection;
- 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:
Both:

- 1 Patient has confirmed Crohn's disease: and
- 2 Fither:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria: Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (severe ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:
All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
 - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

1 Roth

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	lised	Generic
\$	Per	1	Manufacturer

continued...

- 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plague psoriasis: and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plague psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis: or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plague or plagues have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from. at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement: and
- 3 Patient has steroid-refractory disease; and
- 4 Fither:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990:335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004:31:931-7.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less: and
 - 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
 - 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 The patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema), following 12 months' treatment; or</p>
- 3 The patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old, following 12 months' treatment.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 The patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < 1/2+

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema), following 12 months' treatment; or
- 3 The patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old, following 12 months' treatment.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be

Subsidy	Fu	ly Brand or	
(Manufacturer's Price)	Subsidise	ed Generic	
\$	Per	Manufacturer	

continued...

considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Renewal — (severe ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
 - 1.2.2 Fither:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values: or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Fither:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Either:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

OBINUTUZUMAB - PCT only - Specialist - Special Authority see SA1627 below

Inj 25 mg per ml, 40 ml vial	5,910.00	1	✓ Gazyva
Inj 1 mg for ECP	6.21	1 mg	✓ Baxter

⇒SA1627 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

Inj 150 mg prefilled syringe	450.00	1	Xolair
Inj 150 mg vial	450.00	1	✓ Xolair

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IqE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Fither:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Subsidy		Fully	Brand or	
(Manufacturer's Price) :	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Fither:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses: or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PERTUZUMAB - PCT only - Specialist - Special Authority see SA1606 below

Inj 30 mg per ml, 14 ml vial	3,927.00	1	Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	Baxter

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Fither:
 - 2.1 Patient is chemotherapy treatment naïve: or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least

Subsidy		Fully	Brand or	
(Manufacturer's Price		Subsidised	Generic	
<u> </u>	Per	✓	Manufacturer	

continued...

12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and

- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB - PCT only - Specialist - Special Authority see SA178	83 below	
---	----------	--

✓ Mabthera	2	Inj 100 mg per 10 ml vial
✓ Mabthera	1	Inj 500 mg per 50 ml vial
✓ Baxter	1 mg	Inj 1 mg for ECP

⇒SA1783 Special Authority for Subsidy

Initial application — (Antibody-mediated renal transplant rejection) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated renal transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (ABO-incompatible renal transplant) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible renal transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with rituximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 haemophilia with inhibitors: or
 - 2.2 rheumatoid arthritis; or
 - 2.3 severe cold haemagglutinin disease (CHAD); or
 - 2.4 warm autoimmune haemolytic anaemia (warm AIHA); or
 - 2.5 immune thrombocytopenic purpura (ITP); or
 - 2.6 thrombotic thrombocytopenic purpura (TTP); or
 - 2.7 pure red cell aplasia (PRCA); or
 - 2.8 ANCA associated vasculitis; or
 - 2.9 treatment refractory systemic lupus erythematosus (SLE); or
 - 2.10 steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS).

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia **Initial application — (Chronic lymphocytic leukaemia)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 The patient is rituximab treatment naive; and
- 3 Either:
 - 3.1 The patient is chemotherapy treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
 - 3.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; and
- 4 The patient has good performance status; and
- 5 The patient does not have chromosome 17p deletion CLL; and
- 6 Rituximab to be administered in combination with fludarabine and cyclophosphamide or bendamustine for a maximum of 6 treatment cycles; and
- 7 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2. Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

1 Both:

- 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis: and

2 Either:

- 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin: or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin.

Note: Indications marked with * are unapproved indications.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria: Both:

1 Fither:

- 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre: or
- 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy).

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria: Either:

- 1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
- 2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:

Subsidy (Manufacturer's Price)	Fully Subsidised		
(ivialitulacture) 5 Filoe)	Per 🗸	Manufacturer	
Φ	rei •	Manuacturei	

continued...

- 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
- 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
- 3.3 Cyclophosphamide and methotrexate are contraindicated; or
- 3.4 Patient is a female of child-bearing potential; or
- 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
- 2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
- 3 The patient does not have chromosome 17p deletion CLL; and
- 4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide or bendamustine for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
	Per 🗸	Manufacturer

continued...

- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1.000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physiciann; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	1	

continued...

2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

- All of the following:
 - 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
 - 2 An initial response lasting at least 12 months was demonstrated; and
 - 3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Note: Indications marked with * are unapproved indications.

SECUKINUMAB - Special Authority see SA1754 below - Retail pharmacy

Inj 150 mg per ml, 1 ml prefilled syringe...................................1,599.00 2 **✓ Cosentyx**

⇒SA1754 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plague psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab: or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

	Subsidy (Manufacturer's Price)	Sul	Fully osidised	Brand or Generic Manufacturer	
	φ	rei		Manuacturer	
SILTUXIMAB - Special Authority see SA1596 below - Retail ph	armacy				
Note: Siltuximab is to be administered at doses no greater t	han 11 mg/kg every 3	weeks.			
Inj 100 mg vial	770.57	1	✓ S	ylvant	
Inj 400 mg vial	3,082.33	1	✓ S	ylvant	

SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB - PCT only - Special Authority see SA1781 below

Inj 20 mg per ml, 4 ml vial220.00	1	Actemra
Inj 20 mg per ml, 10 ml vial550.00	1	✓ Actemra
Inj 20 mg per ml, 20 ml vial	1	✓ Actemra
Inj 1 mg for ECP	1 mg	✓ Baxter

⇒SA1781 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1331 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses; or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

Subsidy (Manufacturer's Price)	Fully Subsidised		
\$	Per 🗸	Manufacturer	

continued...

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Bules of the Pharmaceutical Schedule: and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
 - 2.4 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.5 Both:
 - 2.5.1 Either:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

(Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	Subsidy		Fully	Brand or
\$ Per ✓ Manufacturer	(Manufacturer's Price		Subsidised	Generic
	\$	Per	✓	Manufacturer

continued...

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline: or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB - PCT only - Specialist - Special Authority see SA1632 below

Inj 150 mg vial	1,350.00	1	Herceptin
Inj 440 mg vial	3,875.00	1	✓ Herceptin
Inj 1 mg for ECP	9.36	1 mg	✓ Baxter

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Fither:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and

Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
- 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria: All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadiuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology);
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and

ibsidy	Fully	Brand or
turer's Price) Sub	osidised	Generic
 \$ Per	/	

continued...

6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

Programmed Cell Death-1 (PD-1) Inhibitors

		NIVOLUMAB - PCT only - Specialist - Special Authority see SA1656 below
Opdivo	1	Inj 10 mg per ml, 4 ml vial
✓ Opdivo	1	Inj 10 mg per ml, 10 ml vial2,629.96
✓ Baxter	1 mg	Inj 1 mg for ECP27.62

⇒SA1656 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma stage III or IV; and
- 2 Patient has measurable disease as defined by the presence of at least one CT or MRI measurable lesion; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Nivolumab is to be used at a maximum dose of 3 mg/kg every 2 weeks for a maximum of 12 weeks (6 cycles); and
- 6 Baseline measurement of overall tumour burden is documented (see Note); and
- 7 Documentation confirming that the patient has been informed and acknowledges that the initial funded treatment period of nivolumab will not be continued beyond 12 weeks (6 cycles) if their disease progresses during this time.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note; or
 - 1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.3 Patient has stable disease according to RECIST criteria (see Note); and
- 2 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; and
- 3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; and
- 5 Nivolumab will be used at a maximum dose of 3 mg/kg every 2 weeks for a maximum of 12 weeks (6 cycles).

Notes: Disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Target lesion measurements should be assessed using CT or MRI imaging with the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

• Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
	Per	1	Manufacturer

continued...

must have reduction in short axis to < 10 mm.

- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB - PCT only - Specialist - Special Authority see SA1657 below

Inj 50 mg vial	2,340.00	1	Keytruda
Inj 1 mg for ECP	49.14	1 mg	Baxter

⇒SA1657 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma stage III or IV: and
- 2 Patient has measurable disease as defined by the presence of at least one CT or MRI measurable lesion; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Pembrolizumab is to be used at a maximum dose of 2 mg/kg every 3 weeks for a maximum of 12 weeks (4 cycles); and
- 6 Baseline measurement of overall tumour burden is documented (see Note); and
- 7 Documentation confirming that the patient has been informed and acknowledges that the initial funded treatment period of pembrolizumab will not be continued beyond 12 weeks (4 cycles) if their disease progresses during this time.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.3 Patient has stable disease according to RECIST criteria (see Note); and
- 2 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; and
- 3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; and
- 5 Pembrolizumab will be used at a maximum dose of 2 mg/kg every 3 weeks for a maximum of 12 weeks (4 cycles).

Notes: Disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Target lesion measurements should be assessed using CT or MRI imaging with the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline

Subsidy		Fully	Brand or
(Manufacturer's Price	,	Subsidised	Generic
<u> </u>	Per		Manufacturer

continued...

and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

leoral
leoral
leoral
leoral
finitor
finitor

⇒SA1491 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

SIROLIMUS - Special Authority see SA0866 below - Retail pharmacy

Tab 1 mg	749.99	100	Rapamune
Tab 2 mg	1,499.99	100	✓ Rapamune
Oral liq 1 mg per ml	449.99	60 ml OP	✓ Rapamune

⇒SA0866 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- · Rapidly progressive transplant vasculopathy; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- · Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP; or
- · Leukoencepthalopathy; or
- · Significant malignant disease

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

Cap 0.5 mg	55.64	100 •	✓ Tacrolimus Sandoz
Cap 1 mg1	11.28	100 •	Tacrolimus Sandoz
Cap 5 mg	78.20	50	Tacrolimus Sandoz

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antiallaray	Droparations	

Allergic Emergencies

ICATIBANT – Special Authority see SA1558 below – Retail pharmacy
Inj 10 mg per ml, 3 ml prefilled syringe.......2,668.00 1 ✓ Firazyr

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Maintenance kit - 6 vials 120 mon freeze dried venom with

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above - Retail pharmacy

diluent	285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent			
9 ml, 3 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent .	305.00	1 OP	✓ Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority see S.	A1367 above – F	Retail pharma	асу
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze			
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze			·
dried venom, with diluent	305.00	1 OP	✓ Venomil \$29

	Subsidy		Fully Brand or
	(Manufacturer's Price	e) Subs	idised Generic
	\$	Per	✓ Manufacturer
Autibistanius			
Antihistamines			
CETIRIZINE HYDROCHLORIDE			
* Tab 10 mg	1.01	100	✓ Zista
* Oral liq 1 mg per ml		200 ml	✓ Histaclear
CHLORPHENIRAMINE MALEATE	2.00	200	
	0.06	500 ml	✓ Histafen
* Oral liq 2 mg per 5 ml	0.00	300 1111	▼ nistaleli
DEXTROCHLORPHENIRAMINE MALEATE			
* Tab 2 mg		40	
	(8.40)		Polaramine
	1.01	20	
	(5.99)		Polaramine
* Oral liq 2 mg per 5 ml		100 ml	5
	(10.29)		Polaramine
FEXOFENADINE HYDROCHLORIDE			
* Tab 60 mg	4.34	20	
	(8.23)		Telfast
* Tab 120 mg	4.74	10	
	(8.23)		Telfast
	14.22	30	
	(26.44)		Telfast
LORATADINE			
* Tab 10 mg	1.28	100	✓ Lorafix
* Oral liq 1 mg per ml		120 ml	✓ Lorfast
PROMETHAZINE HYDROCHLORIDE			
* Tab 10 mg	1 68	50	✓ Allersoothe
* Tab 25 mg		50	✓ Allersoothe
* Oral liq 1 mg per 1 ml		100 ml	✓ Allersoothe
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a		5	✓ Hospira
The first control of the control of	1 00 10.0+	ŭ	• поорни
Inhaled Corticosteroids			
initial de de la control de la			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler, 50 mcg per dose	9.30 2	00 dose OP	✓ Qvar
Aerosol inhaler, 50 mcg per dose CFC-free	8.54 2	00 dose OP	✓ Beclazone 50
Aerosol inhaler, 100 mcg per dose	15.50 2	00 dose OP	✓ Qvar
Aerosol inhaler, 100 mcg per dose CFC-free	12.50 2	00 dose OP	✓ Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free	22.67 2	00 dose OP	✓ Beclazone 250
BUDESONIDE			
Powder for inhalation, 100 mcg per dose	17.00 2	00 dose OP	✓ Pulmicort
, 01			Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00 2	00 dose OP	✓ Pulmicort
5 - 5 - 5 - 5 - 5 - 5 - 5 - 5 - 5 - 5 -			Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00 2	00 dose OP	✓ Pulmicort
			Turbuhaler

	Subsidy		Fully	Brand or
	(Manufacturer's		idised	Generic
	\$	Per		Manufacturer
FLUTICASONE				
Aerosol inhaler, 50 mcg per dose		120 dose OP	-	Floair
Aerosol inhaler, 50 mcg per dose CFC-free		120 dose OP	_	lixotide
Powder for inhalation, 50 mcg per dose		60 dose OP		Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP 120 dose OP		Flixotide Accuhaler Floair
Aerosol inhaler, 125 mcg per dose Aerosol inhaler, 125 mcg per dose CFC-free		120 dose OP	_	Flixotide
Aerosol inhaler, 125 mcg per dose of 0-free		120 dose OP		Floair
Aerosol inhaler, 250 mcg per dose CFC-free		120 dose OP	_	Flixotide
Powder for inhalation, 250 mcg per dose		60 dose OP	_	Flixotide Accuhaler
Towast for initiation, 200 mag per dood	10.00	00 0000 01	• •	iixotiae Aooanalei
Inhaled Long-acting Beta-adrenoceptor Agonis	ts			
EFORMOTEROL FUMARATE				
Powder for inhalation, 6 mcg per dose, breath activated	10.32	60 dose OP		
	(16.90)		(Oxis Turbuhaler
Powder for inhalation, 12 mcg per dose, and monodose devi	ce20.64	60 dose		
	(35.80)			oradil
(Oxis Turbuhaler Powder for inhalation, 6 mcg per dose, breath a	activated to be c	delisted 1 April 20	019)	
EFORMOTEROL FUMARATE DIHYDRATE				
Powder for inhalation 4.5 mcg per dose, breath activated				
(equivalent to eformoterol fumarate 6 mcg metered dose) 10.32	60 dose OP		
	(16.90)		(Oxis Turbuhaler
INDACATEROL				
Powder for inhalation 150 mcg	61.00	30 dose OP	1	Onbrez Breezhaler
Powder for inhalation 300 mcg	61.00	30 dose OP	1	Onbrez Breezhaler
SALMETEROL				
Aerosol inhaler CFC-free, 25 mcg per dose	25.00	120 dose OP	√ 9	Serevent
Aerosol inhaler 25 mcg per dose		120 dose OP	1	Meterol
Powder for inhalation, 50 mcg per dose, breath activated		60 dose OP	√ 9	Serevent Accuhaler
Inhaled Corticosteroids with Long-Acting Beta-	Adrenocept	or Agonists		
BUDESONIDE WITH EFORMOTEROL				
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg	10.00	120 dose OP	1	/annair
Powder for inhalation 100 mcg with eformoterol furnarate 6 mcg		120 dose OP		Symbicort
1 owder for initial ation 100 miles with elormoteror furnarate of it	icg33.74	120 0036 01	•	Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg	21 40	120 dose OP	/ \	/annair
Powder for inhalation 200 mcg with eformoterol furnarate 6 m		120 dose OP		Symbicort
. 555 for initialization 255 may with distinistion fulfidate 6 ff		120 0000 01	- (Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate				
12 mcg - No more than 2 dose per day	44.08	60 dose OP	19	Symbicort
· J				Turbuhaler 400/12
FLUTICASONE FUROATE WITH VILANTEROL				
Powder for inhalation 100 mcg with vilanterol 25 mcg	44 08	30 dose OP	√ F	Breo Ellipta
. Strate for initialization 100 mbg with vilanteror 20 mbg		00 0000 01	٠.	oo Emplu

	Subsidy (Manufacturer's	Price) Subs	Fully idised	Brand or Generic
	\$	Per	1	Manufacturer
LUTICASONE WITH SALMETEROL				
Aerosol inhaler 50 mcg with salmeterol 25 mcg	14.58	120 dose OP	✓ [RexAir
•	33.74		1	Seretide
Aerosol inhaler 125 mcg with salmeterol 25 mcg	16.83	120 dose OP	✓	RexAir
•	44.08		1	Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg - No)			
more than 2 dose per day	33.74	60 dose OP	1	Seretide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg - No)			
more than 2 dose per day	44.08	60 dose OP	1	Seretide Accuhaler
. ,				
Beta-Adrenoceptor Agonists				
ALBUTAMOL				
Oral liq 400 mcg per ml	20.00	150 ml	1	/entolin
Infusion 1 mg per ml, 5 ml		10	-	
····•	(130.21)	. •	١	Ventolin
Inj 500 mcg per ml, 1 ml - Up to 5 inj available on a PSO	` ,	5		/entolin
, , ,				
Inhaled Beta-Adrenoceptor Agonists				
ALBUTAMOL				
Aerosol inhaler, 100 mcg per dose CFC free - Up to 1000				
dose available on a PSO	3.80	200 dose OP	✓ [Respigen
				SalAir
	(6.00)		١	Ventolin Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule - Up to 30 neb)			
available on a PSO	3.93	20	1	<u>Asthalin</u>
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule - Up to 30 neb)			
available on a PSO	4.03	20	1	<u>Asthalin</u>
ERBUTALINE SULPHATE				
Powder for inhalation, 250 mcg per dose, breath activated	27.30	200 dose OP	✓ F	Bricanyl Turbuhaler
Toward for mindiduoti, 200 mag por dood, broadi dourated		200 0000 01		onounyi ranbanalor
Anticholinergic Agents				
PRATROPIUM BROMIDE				
Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dos	:Α			
available on a PSO		200 dose OP	1	Atrovent
Nebuliser soln, 250 mcg per ml, 1 ml ampoule – Up to 40 ne		_00 0000 OI	- /	JYOIIL
available on a PSO		20	√ 1	Univent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 ne		20	• [<u>Jinvelit</u>
available on a PSO		20	/ I	Univent
available off a f oo		20	٠ ١	<u>viii4£iif</u>
nhaled Beta-Adrenoceptor Agonists with Antic	holinergic A	Agents		
ALBUTAMOL WITH IPRATROPIUM BROMIDE				
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg	ner			
dose CFC-free		200 dose OP	√ 1	Duolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per	12.13	200 0036 OF	٠,١	ZWIII III A
vial, 2.5 ml ampoule – Up to 20 neb available on a PSC	5 20	20	√ 1	Duolin

RESPIRATORY SYSTEM AND ALLERGIES

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium.
- Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL – Special Authority see SA1584 above – Retail pharmacy

TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 above - Retail pharmacy

Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg81.00 60 dose OP ✓ Spiolto Respimat

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

Antifibrotics

NINTEDANIB - Special Authority see SA1755 on the next page - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

 Cap 100 mg
 2,554.00
 60 OP
 ✓ Ofev

 Cap 150 mg
 3,870.00
 60 OP
 ✓ Ofev

RESPIRATORY SYSTEM AND ALLERGIES

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1755 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA1748 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Cap 267 mg − Wastage claimable......3,645.00 270 ✓ Esbriet

⇒SA1748 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 80% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

	RESPIRATO	RY SYSTE	EM AND ALLERGIES
	Subsidy (Manufacturer's Pri \$	ice) Subsi Per	Fully Brand or idised Generic Manufacturer
Leukotriene Receptor Antagonists			
MONTELUKAST * Tab 4 mg * Tab 5 mg * Tab 10 mg	5.50	28 28 28	✓ Apo-Montelukast ✓ Apo-Montelukast ✓ Accord \$29 ✓ Apo-Montelukast
Mast Cell Stabilisers			
NEDOCROMIL Aerosol inhaler, 2 mg per dose CFC-freeSODIUM CROMOGLICATE Aerosol inhaler, 5 mg per dose CFC-free		112 dose OP	✓ Tilade ✓ Intal Forte CFC Free
Methylxanthines	20.07	112 dose OF	• Illiai Forte CFC Free
AMINOPHYLLINE Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj av PSO THEOPHYLLINE Tab long-acting 250 mg Oral liq 80 mg per 15 ml	124.37	5 100 500 ml	✓ <u>DBL Aminophylline</u> ✓ Nuelin-SR ✓ Nuelin
Mucolytics DORNASE ALFA — Special Authority see SA0611 be Nebuliser soln, 2.5 mg per 2.5 ml ampoule ➤ SA0611 Special Authority for Subsidy Special Authority approved by the Cystic Fibrosis Adv	250.00	6	✓ Pulmozyme
Notes: Application details may be obtained from PH/ The Co-ordinator, Cystic Fibrosis Advisory Panel PHARMAC, PO Box 10 254 Wellington	ARMAC's website http://www. Phone: (04) 460 4990 Facsimile: (04) 916 7571 Email: CFPanel@pharmac		<u>nz</u> or:
Prescriptions for patients approved for treatment mus and expertise in treating cystic fibrosis. SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7%	t be written by respiratory phy		ediatricians who have experience
Nasal Preparations			
Allergy Prophylactics			

Allergy Prophylactics

RECLOM	FTHASONE	DIDBO	DIONATE
	EIDAOUNE	וארהני	FILMMIT

2.35	200 dose OP	
(5.26)		Alanase
2.46	200 dose OP	
(6.00)		Alanase
	(5.26) 2.46	(5.26) 2.46 200 dose OP

(Alanase Metered aqueous nasal spray, 50 mcg per dose to be delisted 1 January 2020) (Alanase Metered aqueous nasal spray, 100 mcg per dose to be delisted 1 January 2020)

RESPIRATORY SYSTEM AND ALLERGIES

	Subsidy (Manufacturer's \$	Price) Subsi	Fully Brand or idised Generic Manufacturer
BUDESONIDE			
Metered aqueous nasal spray, 50 mcg per dose Metered aqueous nasal spray, 100 mcg per dose		200 dose OP 200 dose OP	✓ <u>SteroClear</u> ✓ <u>SteroClear</u>
LUTICASONE PROPIONATE			
Metered aqueous nasal spray, 50 mcg per dose	1.98	120 dose OP	✓ Flixonase Hayfever & Allergy
PRATROPIUM BROMIDE			
Aqueous nasal spray, 0.03%	4.61	15 ml OP	✓ <u>Univent</u>
Respiratory Devices			
IASK FOR SPACER DEVICE			
a) Up to 50 dev available on a PSO			
b) Only on a PSO			
 c) Only for children aged six years and under 			
Small	2.20	1	e-chamber Mask
EAK FLOW METER			
a) Up to 25 dev available on a PSO			
b) Only on a PSO			_
Low range	9.54	1	Mini-Wright AFS Low Range
Normal range	9.54	1	✓ Mini-Wright Standard
PACER DEVICE			Guiradia
a) Up to 50 dev available on a PSO			
b) Only on a PSO			
220 ml (single patient)	2.95	1	e-chamber Turbo
510 ml (single patient)	5.12	1	e-chamber La
			Grande
800 ml	6.50	1	✓ Volumatic
Respiratory Stimulants			
AFFEINE CITRATE			
0 111 00 1/10 1			4

Oral liq 20 mg per ml (10 mg base per ml)......14.85 25 ml OP

✓ Biomed

	Subsidy (Manufacturer's I \$	Price) Subs Per	Fully Brand or sidised Generic Manufacturer
Ear Preparations			
ACETIC ACID WITH 1, 2- PROPANEDIOL DIACETATE AND B		000	
For Vosol ear drops with hydrocortisone powder refer Stand Ear drops 2% with 1, 2-Propanediol diacetate 3% and	ard Formulae, pa	age 228	
benzethonium chloride 0.02%	6.97	35 ml OP	✓ Vosol
FLUMETASONE PIVALATE		00 1111 01	10001
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform ED's
			✓ Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMY	CIN AND NYSTA	TIN	
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate			
2.5 mg and gramicidin 250 mcg per g		7.5 ml OP	✓ Kenacomb
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and			
gramicidin 50 mcg per ml	4.50	8 ml OP	
	(9.27)		Sofradex
FRAMYCETIN SULPHATE			
Ear/Eye drops 0.5%		8 ml OP	Ontonomia
	(8.65)		Soframycin
Eye Preparations			
-yo i roparationo			
Eye preparations are only funded for use in the eye, unless exp	licitly stated other	wise.	
Anti-Infective Preparations			
ACICLOVIR			
# Eye oint 3%	14 92	4.5 g OP	✓ ViruPOS
CHLORAMPHENICOL	17.02	7.0 y O1	- <u>VIIUI 00</u>
CHLORAMPHENICOL Eye oint 1%	2 48	4 g OP	✓ Chlorsig
Eye drops 0.5%		10 ml OP	✓ Chlorafast
Funded for use in the ear*. Indications marked with * a			
CIPROFLOXACIN			
Eye drops 0.3% - Subsidy by endorsement	9.99	5 ml OP	✓ Ciprofloxacin Teva
When prescribed for the treatment of bacterial keratitis	or severe bacteri	al conjunctivitis	resistant to chloramphenico

for the second line treatment of chronic suppurative otitis media (CSOM)*; and the prescription is endorsed accordingly.

(14.55)

✓ Genoptic

Brolene

✓ Fucithalmic

5 ml OP

10 ml OP

5 g OP

GENTAMICIN SULPHATE

PROPAMIDINE ISETHIONATE

SODIUM FUSIDATE [FUSIDIC ACID]

Note: Indication marked with a * is an unapproved indication.

	Subsidy		Fully	Brand or
	(Manufacturer's P \$	Price) Subs Per	sidised •	Generic Manufacturer
TOBRAMYCIN				
Eye oint 0.3% Eye drops 0.3%		3.5 g OP 5 ml OP	-	obrex obrex
Corticosteroids and Other Anti-Inflammatory Pro	eparations			
DEXAMETHASONE				
** Eye oint 0.1% *Eye drops 0.1% Ocular implant 700 mcg – Special Authority see SA1680 beld	4.50	3.5 g OP 5 ml OP		Maxidex Maxidex

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient has diabetic macular oedema with pseudophakic lens; and

- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- Fither
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not vet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

*	Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b			
	sulphate 6,000 u per g	5.39	3.5 g OP	✓ Maxitrol
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin			
	b sulphate 6,000 u per ml	4.50	5 ml OP	✓ Maxitrol
DIC	CLOFENAC SODIUM			
	Eye drops 0.1%	13.80	5 ml OP	✓ Voltaren Ophtha

✓ Ozurdex

✓ Rexacrom

5 ml OP

	Subsidy		Fully	Brand or
	(Manufacturer's Pr	rice) Subs	idised	Generic
	\$	Per	1	Manufacturer
FLUOROMETHOLONE				
* Eye drops 0.1%	3.09	5 ml OP	√ F	ML
LEVOCABASTINE				
Eye drops 0.5 mg per ml	8.71	4 ml OP		
	(10.34)		L	ivostin
LODOXAMIDE				
Eye drops 0.1%	8.71	10 ml OP	√ L	.omide
PREDNISOLONE ACETATE				
Eye drops 1%	3.93	10 ml OP	✓ P	rednisolone-AFT
	7.00	5 ml OP	✓ P	red Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority so	ee SA1715 below	- Retail pharr	nacy	
Eye drops 0.5%, single dose (preservative free)	38.50	20 dose	N	Minims
				Prednisolone

⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Eye drops 2%

SODIUM CROMOGLICATE

Glaucoma Preparations - Beta Blockers		
BETAXOLOL # Eye drops 0.25%	5 ml OP 5 ml OP	✓ Betoptic S✓ Betoptic
* Eye drops 0.5%	5 ml OP	✓ Betagan
TIMOLOL * Eye drops 0.25% 1.43 * Eye drops 0.25%, gel forming 3.30 * Eye drops 0.5% 1.43 * Eye drops 0.5%, gel forming 3.78	5 ml OP 2.5 ml OP 5 ml OP 2.5 ml OP	✓ Arrow-Timolol ✓ Timoptol XE ✓ Arrow-Timolol ✓ Timoptol XE
Glaucoma Preparations - Carbonic Anhydrase Inhibitors		
ACETAZOLAMIDE * Tab 250 mg	100	✓ <u>Diamox</u>
* Eye drops 1%	5 ml OP	✓ Azopt
DORZOLAMIDE HYDROCHLORIDE * Eye drops 2%	5 ml OP	Trusopt

	(Manufacturer's Pri \$	ce) Sub Per	sidised	Brand or Generic Manufacturer
DORZOLAMIDE WITH TIMOLOL * Eye drops 2% with timolol 0.5%	2.87 (3.45)	5 ml OP		Dortimopt Arrow-Dortim
Arrow-Dortim Eye drops 2% with timolol 0.5% to be delisted 1 Ap	oril 2019)			
Glaucoma Preparations - Prostaglandin Analogu	ues			
BIMATOPROST ★ Eye drops 0.03%	3.30	3 ml OP	•	Bimatoprost Multichem
	(3.65)			Bimatoprost Actavis
Bimatoprost Actavis Eye drops 0.03% to be delisted 1 May 2019) ATANOPROST			٠.	_
* Eye drops 0.005%	1.57 1.50 (1.84)	2.5 ml OP		Teva Hysite
Hysite Eye drops 0.005% to be delisted 1 July 2019) FRAVOPROST	, ,			.,,
* Eye drops 0.004%	7.30 19.50	5 ml OP 2.5 ml OP		Travopt Travatan
Glaucoma Preparations - Other				
BRIMONIDINE TARTRATE * Eye drops 0.2%	4.29	5 ml OP	✓	Arrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE ★ Eye drops 0.2% with timolol maleate 0.5% PILOCARPINE HYDROCHLORIDE	18.50	5 ml OP	•	Combigan
* Eye drops 1%		15 ml OP		Isopto Carpine
Eye drops 2% Eye drops 4% Subsidised for oral use pursuant to the Standard Formula	7.99	15 ml OP 15 ml OP		Isopto Carpine Isopto Carpine
Eye drops 2% single dose – Special Authority see SA0895 below – Retail pharmacy	31.95	20 dose	✓	Minims Pilocarpine
➤SA0895 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals valid Either: 1 Patient has to use an unpreserved solution due to an allergent of the province of the			eeting	the following criteria:

2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics

ATROPINE SULPHATE		
* Eye drops 1%	15 ml OP	✓ Atropt
CYCLOPENTOLATE HYDROCHLORIDE		
* Eye drops 1%	15 ml OP	Cyclogyl

	Subsidy		Fully	Brand or
	(Manufacturer's F	Price) Subsi	idised	Generic
	\$	Per	1	Manufacturer
TROPICAMIDE				
* Eye drops 0.5%	7.15	15 ml OP	✓ N	lydriacyl
* Eye drops 1%	8.66	15 ml OP		lydriacyl
Preparations for Tear Deficiency				
For acetylcysteine eye drops refer Standard Formulae, page 228				
HYPROMELLOSE				
* Eye drops 0.5%	2.00	15 ml OP		
	(3.92)		N	lethopt
HYPROMELLOSE WITH DEXTRAN				
* Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	✓ P	oly-Tears
POLYVINYL ALCOHOL				•
* Eye drops 1.4%	2 62	15 ml OP	✓ ∨	'ietil
* Eye drops 3%	2.02	15 ml OP		istii Forte
* Eye urups 3 /0	3.00	15 IIII OF	<u>v</u>	ISHI FUILE

Preservative Free Ocular Lubricants

⇒SA1388 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER – Special Authority see SA1388 above – Retail pha	armacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL - Special Author	ity see SA1388 a	above – Retail	pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	✓ Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special Auth	ority see SA1388	3 above – Reta	ail pharmacy
Eye drops 1 mg per ml			
Hylo-Fresh has a 6 month expiry after opening. The Pha	armacy Procedur	es Manual res	triction allowing one bottle per
month is not relevant and therefore only the prescribed d	losage to the nea	arest OP may b	be claimed.

Other Eye Preparations

NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%4.15	15 ml OP	✓ Naphcon Forte
OLOPATADINE Eye drops 0.1%10.00	5 ml OP	✓ Patanol
PARAFFIN LIQUID WITH SOFT WHITE PARAFFIN * Eye oint with soft white paraffin	3.5 g OP	✓ Refresh Night Time
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE Eye oint 138 mcg per g	5 g OP	✓ VitA-POS

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.



Subsidy (Manufacturer's Price) Fully Subsidised Brand or Generic Manufacturer

Sandoz

Various

PHARMACY SERVICES

May only be claimed once per patient.

* Brand switch fee4.50

1 fee **✓ BSF Entecavir**

The Pharmacode for BSF Entecavir Sandoz is 2559420 - see also page 100 (BSF Entecavir Sandoz Brand switch fee to be delisted 1 April 2019)

Agents Used in the Treatment of Poisonings

Antidotes

ACETYLCYSTEINE - Retail pharmacy-Specialist

Inj 200 mg per ml, 10 ml ampoule58.76

10

✓ DBL Acetylcysteine

NALOXONE HYDROCHLORIDE

a) Up to 5 ini available on a PSO

b) Only on a PSO

* Inj 400 mcg per ml, 1 ml ampoule22.60

5

✓ DBL Naloxone Hydrochloride

Removal and Elimination

CHARCOAL

a) Up to 250 ml available on a PSO

b) Only on a PSO

DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy

Wastage claimable

Tab 125 mg dispersible	276.00	28	Exjade
Tab 250 mg dispersible		28	Exjade
Tab 500 mg dispersible	1,105.00	28	Exjade

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μ L) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μ L).

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

	Subsidy		Fully	Brand or
	(Manufacturer's Pri	ice) Sub	sidised	Generic
	\$	Per	1	Manufacturer
DEFERIPRONE - Special Authority see SA1480 below - Retail	pharmacy			
Tab 500 mg	533.17	100	√ F	erriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	√ F	erriprox
⇒SA1480 Special Authority for Subsidy				
Initial application only from a haematologist. Approvals valid w	ithout further rene	wal unless no	otified fo	or applications meeting the
following criteria:				
Either:				

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE * Inj 500 mg vial	51.52	10	✓ Desferal
, ,	84.53		✓ DBL Desferrioxamine Mesylate for Inj BP
(Desferal Inj 500 mg vial to be delisted 1 June 2019)			
SODIUM CALCIUM EDETATE			
* Inj 200 mg per ml, 5 ml	53.31	6	
	(156.71)		Calcium Disodium Versenate



Standard Formulae

Standard Formulae			
ACETYLCYSTEINE EYE DROPS		PHENOBARBITONE ORAL LIQUID	
Acetylcysteine inj 200 mg per ml, 10 ml	qs	Phenobarbitone Sodium	1 g
Suitable eye drop base	qs	Glycerol BP	70 ml
	7-	Water	to 100 ml
ASPIRIN AND CHLOROFORM APPLICATION			
Aspirin Soluble tabs 300 mg	12 tabs	PHENOBARBITONE SODIUM PAEDIATRIC ORAL	LIQUID (10
Chloroform	to 100 ml	mg per ml)	
000500500500000000000000000000000000000		Phenobarbitone Sodium	400 mg
CODEINE LINCTUS PAEDIATRIC (3 mg per 5 ml)	00	Glycerol BP	4 ml
Codeine phosphate	60 mg	Water	to 40 ml
Glycerol Preservative	40 ml	PILOCARPINE ORAL LIQUID	
Water	qs to 100 ml		00
water	10 100 1111	Pilocarpine 4% eye drops Preservative	qs
CODEINE LINCTUS DIABETIC (15 mg per 5 ml)		Water	qs to 500 ml
Codeine phosphate	300 mg	(Preservative should be used if quantity supplied is f	
Glycerol	40 ml	than 5 days.)	or more
Preservative	qs	than 5 days.)	
Water	to 100 ml	SALIVA SUBSTITUTE FORMULA	
		Methylcellulose	5 g
FOLINIC MOUTHWASH		Preservative	qs
Calcium folinate 15 mg tab	1 tab	Water	to 500 ml
Preservative	qs	(Preservative should be used if quantity supplied is f	or more
Water	to 500 ml	than 5 days. Maximum 500 ml per prescription.)	
(Preservative should be used if quantity supplied is f	or more	SODIUM CHLORIDE ORAL LIQUID	
than 5 days. Maximum 500 ml per prescription.)		Sodium chloride inj 23.4%, 20 ml	qs
MAGNESIUM HYDROXIDE 8% MIXTURE		Water	qs
Magnesium hydroxide paste 29%	275 g	(Only funded if prescribed for treatment of hyponatra	
Methyl hydroxybenzoate	1.5 g	` , , , , , , , , , , , , , , , , , , ,	,
Water	to 1,000 m	VANCOMYCIN ORAL SOLUTION (50 mg per ml)	
		Vancomycin 500 mg injection	10 vials
METHADONE MIXTURE		Glycerol BP	40 ml
Methadone powder	qs	Water	to 100 ml
Glycerol	qs	(Only funded if prescribed for treatment of Clostridiu	m difficile
Water	to 100 ml	following metronidazole failure)	
METHYL HYDROXYBENZOATE 10% SOLUTION		VOSOL EAR DROPS	
Methyl hydroxybenzoate	10 g	WITH HYDROCORTISONE POWDER 1%	
Propylene glycol	to 100 ml	Hydrocortisone powder	1%
(Use 1 ml of the 10% solution per 100 ml of oral liqu	id mixture)	Vosol Ear Drops	to 35 ml
	-,	- r -	
OMEPRAZOLE SUSPENSION			
Omeprazole capules or powder	qs		
Sodium bicarbonate powder BP	8.4 g		

to 100 ml

Water

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy

(Manufacturer's Price)

Fully

Subsidised

Brand or

Generic

Per Manufacturer Extemporaneously Compounded Preparations and Galenicals BENZOIN Tincture compound BP......24.42 500 ml (39.90)Pharmacy Health 2.44 50 ml (5.10)Pharmacy Health CHI OROFORM a) Only in combination b) Maximum of 100 ml per prescription c) Only in aspirin and chloroform application. Chloroform BP......25.50 500 ml ✓ PSM CODEINE PHOSPHATE - Safety medicine; prescriber may determine dispensing frequency Powder - Only in combination......63.09 (90.09)Douglas Only in extemporaneously compounded codeine linctus diabetic or codeine linctus paediatric. COLLODION FLEXIBI F ✓ PSM 100 ml COMPOUND HYDROXYBENZOATE - Only in combination Only in extemporaneously compounded oral mixtures. ✓ Midwest 100 ml ✓ David Craig 34.18 GLYCERIN WITH SODIUM SACCHARIN - Only in combination Only in combination with Ora-Plus. Suspension......32.50 ✓ Ora-Sweet SF 473 ml GLYCERIN WITH SUCROSE - Only in combination Only in combination with Ora-Plus. Suspension......32.50 473 ml ✓ Ora-Sweet **GLYCEROL** ✓ healthE Glycerol BP 500 ml Only in extemporaneously compounded oral liquid preparations. MAGNESIUM HYDROXIDE 500 g ✓ PSM METHADONE HYDROCHLORIDE a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing frequency d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets). ✓ AFT 1 a METHYL HYDROXYBENZOATE 25 q ✓ Midwest METHYLCELLULOSE 100 g MidWest 473 ml ✓ Ora-Plus METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN - Only in combination ✓ Ora-Blend SF 473 ml

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Pric	e) 5	Fully Subsidised		
	` \$	Per	•	Manufacturer	
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Onl	ly in combination				
Suspension	32.50	473 ml	1	Ora-Blend	
PHENOBARBITONE SODIUM					
Powder - Only in combination	52.50	10 g	1	MidWest	
•	325.00	100 g	1	MidWest	
Only in children up to 12 years					
PROPYLENE GLYCOL					
Only in extemporaneously compounded methyl hydroxybenz	oate 10% solution.				
Liq	11.25	500 ml	1	Midwest	
SODIUM BICARBONATE					
Powder BP - Only in combination	8.95	500 g	1	Midwest	
	9.80				
	(29.50)			David Craig	
Only in extemporaneously compounded omeprazole and	d lansoprazole susp	ension.			
SYRUP (PHARMACEUTICAL GRADE) - Only in combination					
Only in extemporaneously compounded oral liquid preparation					
Liq	21.75	2,000 m		Midwest	
WATER					
Tap - Only in combination	0.00	1 ml	✓	Tap water	

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1522 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 inborn errors of metabolism: or
- 7 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Cystic fibrosis or renal failure)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

continued...



Subsidy (Manufacture de Brise)		Fully	Brand or	
(Manufacturer's Price)	5	ubsidised	Generic	
\$	Per	✓	Manufacturer	

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

Subsidy (Manufacturer's Price)	Sul	Fully	Brand or Generic
 \$	Per	✓	Manufacturer

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Authority see SA1523 on the previous page - Hospital pharmacy [HP3]

Emulsion (neutral)	12.30	200 ml OP	✓ Calogen
	30.75	500 ml OP	✓ Calogen
Emulsion (strawberry)	12.30	200 ml OP	✓ Calogen
Oil	30.00	500 ml OP	✓ MCT oil (Nutricia)
Oil, 250 ml	114.92	4 OP	✓ Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	ırmacy [HP3]	SUPPLEMENT – Special Authority see SA1524 above – Hospital pha	PROTEIN SUPPLEMENT
✓ Protifar	225 g OP	r7.90	Powder
✓ Resource		8.95	
Beneprotein	•		

Subsidy (Manufacturer's Price) Fully Subsidised Brand or Generic Manufacturer

Oral Supplements/Complete Diet (Nasogastric/Gastrostomy Tube Feed)

Respiratory Products

⇒SA1094 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has CORD and hypercapnia, defined as a CO2 value exceeding 55 mmHg.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3] Liquid7.50 1.000 ml OP ✓ Diason RTH ✓ Glucerna Select **RTH** DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3] ✓ Diasip 200 ml OP Liquid (strawberry)......1.50 200 ml OP ✓ Diasip 250 ml OP ✓ Glucerna Select 1 88 237 ml OP 1.78 (2.10)Resource Diabetic (2.10)Sustagen Diabetic

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism: or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED - Special Authority see SA1525 above - Hospital pharmacy [HP3]

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

r	
PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1379 ab Liquid6.00	ove – Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini Energy RTH
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see SA1379 about Liquid	ve – Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini RTH ✓ Pediasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML — Special Authority se Liquid	ee SA1379 above – Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML – Special Authority see SA1379 above Liquid (strawberry)	 Hospital pharmacy [HP3] 200 ml OP ✓ Fortini 200 ml OP ✓ Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA1379 above - Liquid (chocolate)	Hospital pharmacy [HP3] 200 ml OP
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority see S Liquid (unflavoured)	200 ml OP 200 m
PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 above – Hospita	ai pnarmacy [HP3]

Peptamen Junior

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML - Special Authority s		Hospital pharm	,
ции	0.00	300 1111 01	• Nepro III IIIII
RENAL ORAL FEED 1.8 KCAL/ML - Special Authority see S	A1101 above - Hos	spital pharmacy	[HP3]
Liquid	2.67	220 ml OP	✓ Nepro HP (strawberry)
			✓ Nepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA	1101 above – Hosp	ital pharmacy [H	HP3]
Liquid	2.88	237 ml OP	
	(3.31)		NovaSource Renal
Liquid (apricot) 125 ml	11.52	4 OP	✓ Renilon 7.5
Liquid (caramel) 125 ml	11.52	4 OP	✓ Renilon 7.5

Specialised And Elemental Products

SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	(Manufacturer's	Price) Subs	sidised Generic Manufacturer
ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML – Sp pharmacy [HP3] Liquid	,	ee SA1377 on th	ne previous page – Hospital Vital
ORAL ELEMENTAL FEED 0.8KCAL/ML – Special Authority se Liquid (grapefruit), 250 ml carton Liquid (pineapple & orange), 250 ml carton Liquid (summer fruits), 250 ml carton	171.00 171.00	e previous page 18 OP 18 OP 18 OP	 Hospital pharmacy [HP3] ✓ Elemental 028 Extra ✓ Elemental 028 Extra ✓ Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see Powder (unflavoured)		orevious page – 80 g OP	Hospital pharmacy [HP3] ✓ Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Aut [HP3] Liquid	•	77 on the previo 1,000 ml OP	us page – Hospital pharmacy ✓ Peptisorb

Subsidy

Fully

Brand or

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Standard Supplements

⇒SA1554 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age: and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) only from a dietitian, relevant

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist, dietitian on the recommendation of a gastroenterologist or vocationally registered general practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

continued...



Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

Initial application — (Short-term medical condition) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result: or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (**Long-term medical condition**) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis: or
- 3 Liver disease: or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	•	Manufacturer	

- 8 Bowel fistula; or
 - 9 Severe chronic neurological conditions; or
 - 10 Epidermolysis bullosa; or
 - 11 AIDS (CD4 count < 200 cells/mm3); or
 - 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease: or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML – Special Authority see SA1554 on pa	•		/ [HP3] ✓ Nutrison Energy
ENTERAL FEED 1KCAL/ML - Special Authority see SA1554 on page Liquid		spital pharmacy [l 250 ml OP 1,000 ml OP	
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authority see Liquid		n page 238 – Ho 1,000 ml OP	spital pharmacy [HP3] Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority see S. Liquid		age 238 - Hospi 1,000 ml OP	tal pharmacy [HP3] Jevity RTH Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority see S Liquid		page 238 – Hosp 250 ml OP 1,000 ml OP	oital pharmacy [HP3] ✓ Ensure Plus HN ✓ Ensure Plus RTH ✓ Jevity HiCal RTH ✓ Nutrison Energy Multi Fibre

Subsidy		Fully	Brand or
(Manufacturer's P	rice)	Subsidised	Generic
\$	Per	✓	Manufacturer

ORAL FEED (POWDER) - Special Authority see SA1554 on page 238 - Hospital pharmacy [HP3]

Note: Higher subsidy for Sustagen Hospital Formula will only be reimbursed for patients with both a valid Special Authority number and an appropriately endorsed prescription.

Additional subsidy by endorsement is available for patients with fat malabsorption, fat intolerance or chyle leak. The prescription must be endorsed accordingly.

Additional subsidy by endorsement is available for patients with fat malabsorption, fat intolerance or chyle leak. The prescription must be endorsed accordingly.

Formula Active

ORAL FEED 1.5KCAL/ML - Special Authority see SA1554 on page 238 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease. The prescription must be endorsed accordingly.

Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with	0.72	000 ml OD	
Endorsement	(1.26) (1.26)	200 ml OP	Ensure Plus Fortisip
Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	Casura Diva
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (fruit of the forest) – Higher subsidy of \$1.26 per 200 ml	(1.20)		Fortisip
with Endorsement	0.72	200 ml OP	
The Englishment of the English of th	(1.26)	200 1111 01	Ensure Plus
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with	` ,		
Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
Liquid (vanilla) – Higher subsidy of up to \$1.33 per 237 ml with			
Endorsement		237 ml OP	
	(1.33)	000 OD	Ensure Plus
	0.72	200 ml OP	Ensure Plus
	(1.26) (1.26)		Fortisip
	(1.20)		i ortioip

Fortisip Multi Fibre

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1554 on page 238 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with

Elquia (onocolate) Triginer subsity of \$1.20 per 200 mil with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	

(1.26)

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 ab	ove – Hospital į	pharmacy [HP3]	
Liquid	5.50	500 ml OP	✓ Nutrison
			Concentrated
	11 00	1 000 ml OP	✓ Two Cal HN RTH

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$1.90 per 200 ml with

(1.90) Two Cal HN

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FOOD THICKENER - Special Authority see SA1106 above - Hospital pharmacy [HP3]

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by PHARMAC from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX - Special Authority see SA1729 above - Hospital pharmacy [HP3]

Powder	2.81 1,000 g OP	•
	(5.15)	Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see	e SA1729 above – Hospital pharmacy [HP	3]
Powder	3.93 1,000 g OP	
	(7.32)	NZB Low Gluten
		Bread Mix

3.51 (10.87) Horleys Bread Mix

	Subsidy	F	Fully Brand or	
	(Manufacturer's I		lised Generic	
	\$	Per	✓ Manufacturer	
GLUTEN FREE FLOUR - Special Authority see SA1729 on th	e previous page -	- Hospital pharma	icv [HP3]	
Powder		2.000 g OP	, []	
	(18.10)	_,000 g 0.	Horleys Flour	
GLUTEN FREE PASTA - Special Authority see SA1729 on th	e previous page -	- Hospital pharma	cv [HP3]	
Buckwheat Spirals		250 g OP	, [-]	
	(3.11)	3 -	Orgran	
Corn and Vegetable Shells	, ,	250 g OP	- 3 ··	
ŭ	(2.92)	J	Orgran	
Corn and Vegetable Spirals	` ,	250 g OP	•	
ŭ i	(2.92)	J	Orgran	
Rice and Corn Lasagne Sheets	1.60 [′]	200 g OP	•	
U	(3.82)	J	Orgran	
Rice and Corn Macaroni	2.00 [°]	250 g OP	· ·	
	(2.92)	Ü	Orgran	
Rice and Corn Penne	2.00	250 g OP	•	
	(2.92)	•	Orgran	
Rice and Maize Pasta Spirals	2.00	250 g OP	•	
	(2.92)	•	Orgran	
Rice and Millet Spirals	2.00	250 g OP	-	
	(3.11)	•	Orgran	
Rice and corn spaghetti noodles	2.00	375 g OP	-	
	(2.92)	_	Orgran	
Vegetable and Rice Spirals	2.00	250 g OP	-	
	(2.92)	_	Orgran	
Italian long style spaghetti	2.00	220 g OP	-	
• • •	(3.11)	-	Orgran	

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

Supplements For MSUD

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

Tabs		75 OP	✓ Phlexy 10
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior Chocolate
Powder (unflavoured) 27.8 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 36 g sachets	393.00	30	PKU Anamix Junior
Powder (vanilla) 36 g sachet	393.00	30	PKU Anamix Junior Vanilla
Infant formula	174.72	400 g OP	PKU Anamix Infant
Powder (orange)	221.00	500 g OP	XP Maxamaid
	320.00		XP Maxamum
Powder (unflavoured)	221.00	500 g OP	XP Maxamaid
	320.00		XP Maxamum
Liquid (berry)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	 Easiphen Liquid
Liquid (juicy tropical) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g		36 OP	✓ PKU Lophlex Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20

(XP Maxamaid Powder (orange) to be delisted 1 April 2019)

(XP Maxamaid Powder (unflavoured) to be delisted 1 April 2019)

Foods

LOW PROTEIN BAKING MIX - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3] 500 a OP ✓ Loprofin Mix LOW PROTEIN PASTA - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

Animal shapes	11.91	500 g OP	✓ Loprofin
Lasagne		250 g OP	✓ Loprofin
Low protein rice pasta		500 g OP	✓ Loprofin
Macaroni		250 g OP	Loprofin
Penne	11.91	500 g OP	✓ Loprofin
Spaghetti	11.91	500 g OP	✓ Loprofin
Spirals	11.91	500 g OP	✓ Loprofin

Subsidy (Manufacturer's Price) \$ Pe

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Gastrointestinal and Other Malabsorptive Problems

AMINO ACID FORMULA - Special Authority see SA1219 b	elow - Hospital phar	macy [HP3]	
Powder	43.60	400 g OP	 Alfamino Junior
	53.00		✓ Neocate LCP
Powder (unflavoured)	53.00	400 g OP	✓ Elecare
		_	✓ Elecare LCP
			Neocate Gold
			✓ Neocate Junior Unflavoured
			✓ Neocate SYNEO
Powder (vanilla)	53.00	400 g OP	✓ Elecare
•		Ū	Neocate Junior Vanilla

(Neocate LCP Powder to be delisted 1 May 2019)

⇒SA1219 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Extensively hydrolysed formula has been reasonably trialled and is inappropriate due to documented severe intolerance or allergy or malabsorption; or
- 2 History of anaphylaxis to cows milk protein formula or dairy products; or
- 3 Eosinophilic oesophagitis.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see SA1557 below - Hospital pharmacy [HP3]

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome: or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia: or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis: or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure: or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula: and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula;

continued...

Subs	sidy Full	/ Brand or
(Manufactur	rer's Price) Subsidise	d Generic
\$	Per 🕶	Manufacturer

and

3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority	see SA1197	above - Retail	pharmacy
Powder (unflavoured)	35.50	300 g OP	✓ KetoCal 4:1
		-	✓ Ketocal 3:1
Powder (vanilla)	35.50	300 g OP	✓ KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer \$

Vaccinations

ADULT DIPHTHERIA AND TETANUS VACCINE - [Xpharm]

Ini 2 IU diphtheria toxoid with 20 IU tetanus toxoid in 0.5 ml..................0.00 ✓ ADT Booster

Any of the following:

- 1) For vaccination of patients aged 45 and 65 years old; or
- 2) For vaccination of previously unimmunised or partially immunised patients: or
- 3) For revaccination following immunosuppression; or
- 4) For boosting of patients with tetanus-prone wounds; or
- 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or

Ini Mycobacterium boyis BCG (Bacillus Calmette-Guerin). Danish strain 1331, live attenuated, vial with diluent.................0.00

10 ✓ BCG Vaccine

DIPHTHERIA. TETANUS AND PERTUSSIS VACCINE - [Xpharm]

Funded for any of the following criteria:

www.bcgatlas.org/index.php.

- 1) A single vaccine for pregnant woman between gestational weeks 28 and 38; or
- 2) A course of up to four vaccines is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
- 3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens.

Notes: Tdap is not registered for patients aged less than 10 years. Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid, 8 mcg pertussis filamentous

haemagluttinin and 2.5 mcg pertactin in 0.5 ml syringe0.00 10 **Boostrix** Boostrix

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE - [Xpharm]

Funded for any of the following:

- 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
- 2) A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
- 3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive
- 4) Five doses will be funded for children requiring solid organ transplantation.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

pertussis toxoid, 25 mcg pertussis filamentous

haemagluttinin, 8 mcg pertactin and 80 D-antigen units

✓ fully subsidised

✓ Infanrix IPV

	NATIONAL I	MMUNIS	ATIC	ON SCHEDULE
(Ma	Subsidy nufacturer's Price) \$	F Subsidi Per	iully sed	Brand or Generic Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND I	HAEMOPHILUS II	NFLUENZA	E TYP	PE B VACCINE -
[Xpharm]				
Funded for patients meeting any of the following criteria:				
1) Up to four doses for children up to and under the age of 10				d
2) An additional four doses (as appropriate) are funded for (re-				
10 who are patients post haematopoietic stem cell transplar post solid organ transplant, renal dialysis and other severely				st spienectomy; pre- or
3) Up to five doses for children up to and under the age of 10 i				1
Note: A course of up-to four vaccines is funded for catch up prog				
to complete full primary immunisation. Please refer to the Immur				
programmes.				
Inj 30IU diphtheriatoxoid with 40IU tetanustoxoid, 25mcg				
pertussistoxoid, 25mcg				
pertussisfilamentoushaemagluttinin, 8 mcgpertactin,				
80 D-AgUpoliovirus, 10mcghepatitisBsurfaceantigen in				
0.5ml syringe	0.00	10	✓ <u>Inf</u>	anrix-hexa
HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm] One dose for patients meeting any of the following:				
1) For primary vaccination in children; or				
2) An additional dose (as appropriate) is funded for (re-)immur				
transplantation, or chemotherapy; functional asplenic; pre o				id organ transplant, pre-
or post cochlear implants, renal dialysis and other severely				
3) For use in testing for primary immunodeficiency diseases, o	n the recommend	ation of an	ınterna	al medicine physician or
paediatrician.				
Haemophilus Influenzae type B polysaccharide 10 mcg				
conjugated to tetanus toxoid as carrier protein 20-40 mcg;				
prefilled syringe plus vial 0.5 ml	0.00	1	✓ Hil	berix
HEPATITIS A VACCINE – [Xpharm]				
Funded for patients meeting any of the following criteria:				
Two vaccinations for use in transplant patients; or				
2) Two vaccinations for use in children with chronic liver disea	se; or			
3) One dose of vaccine for close contacts of known hepatitis A	cases.			
Inj 1440 ELISA units in 1 ml syringe	0.00	1	✓ Ha	vrix
Inj 720 ELISA units in 0.5 ml syringe		1	_	vrix Junior

NATIONAL IMMUNISATION SCHEDULE

		Subsidy		Fully	Brand or
		(Manufacturer's Price)		Subsidised	Generic
		\$	Per		Manufacturer
HEPATITIS E	RECOMBINANT VACCINE - [Xpharm]				
Inj 5 mcg	per 0.5 ml vial	0.00	1	√ H	BvaxPRO
	led for patients meeting any of the following criteria:			_	
	for household or sexual contacts of known acute he	enatitis B natients or h	enatiti	is B carriers	s: or
,	for children born to mothers who are hepatitis B sui				o, o.
	for children up to and under the age of 18 years inc				achieved a positive
0,	serology and require additional vaccination or requi				aomovoa a poomvo
4)	for HIV positive patients; or	io a piiiiai y couloo c		manori, or	
	for hepatitis C positive patients; or				
,	for patients following non-consensual sexual interco	ourse: or			
,	for patients following immunosuppression; or	Jul. 55, 51			
,	for solid organ transplant patients; or				
,	for post-haematopoietic stem cell transplant (HSCT) natients: or			
,	following needle stick injury.) patiente, ei			
10)	Tollowing needle stick injury.				
Ini 10 ma	g per 1 ml vial	0.00	1	./ ⊔	BvaxPRO
•	led for patients meeting any of the following criteria:	0.00	'	• 11	DVAXENO
				:- Di	
,	for household or sexual contacts of known acute he				s; or
	for children born to mothers who are hepatitis B sur				achieved a positive
3)	for children up to and under the age of 18 years inc				acriieved a positive
4)	serology and require additional vaccination or requi	re a primary course of	vacc	mation, or	
,	for HIV positive patients; or				
,	for hepatitis C positive patients; or				
	for patients following non-consensual sexual interce for patients following immunosuppression; or	Juise, oi			
,					
,	for solid organ transplant patients; or	') notionto: or			
,	for post-haematopoietic stem cell transplant (HSCT) patients; or			
10)	following needle stick injury.				
In: 00 m	a man 4 mal manafilla di accidenta	0.00			manada D
	g per 1 ml prefilled syringe	0.00	1	V E	ngerix-B
	led for patients meeting any of the following criteria:				
	for household or sexual contacts of known acute he				s; or
,	for children born to mothers who are hepatitis B sur	0 1			1.1 1 92
3)	for children up to and under the age of 18 years inc				achieved a positive
	serology and require additional vaccination or requi	ire a primary course o	t vacc	ination; or	
,	for HIV positive patients; or				
,	for hepatitis C positive patients; or				
,	for patients following non-consensual sexual interco	ourse; or			
	for patients following immunosuppression; or				
,	for solid organ transplant patients; or				
,	for post-haematopoietic stem cell transplant (HSCT) patients; or			
10)	following needle stick injury.				
•	g per 1 ml vial	0.00	1	✓ <u>H</u>	BvaxPRO
	led for any of the following criteria:				
	for dialysis patients; or				
2)	for liver or kidney transplant patient.				

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	/	Manufacturer	

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV] - [Xpharm]

Any of the following:

- 1) Maximum of two doses for children aged 14 years and under; or
- 2) Maximum of three doses for patients meeting any of the following criteria:
 - 1) People aged 15 to 26 years inclusive; or
 - 2) Either:

People aged 9 to 26 years inclusive

- 1) Confirmed HIV infection; or
- 2) Transplant (including stem cell) patients: or
- 3) Maximum of four doses for people aged 9 to 26 years inclusive post chemotherapy

	Subsidy (Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer
INFLUENZA VACCINE Inj 45 mcg in 0.5 ml syringe (trivalent vaccine)	90.00	10	√ In	fluvac

- a) Only on a prescription
- b) No patient co-payment payable
- С
- A) is available each year for patients who meet the following criteria, as set by PHARMAC, for use if a funded quadrivalent influenza vaccine is not available:
 - a) all people 65 years of age and over; or
 - b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease: or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV. or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or

Ini 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine) -

- c) children aged four years and under who have been hospitalised for respiratory illness or have a history of significant respiratory illness;
- d) people under 18 years of age living in the Seddon/Ward and rural Eastern Marlborough region (within the Nelson Marlborough District Health Board) and Kaikoura and Hurunui areas (within the Canterbury District Health Board);
- e) People under 18 years of age who have been displaced from their homes in Edgecumbe and the surrounding region;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

,	3 (1		
[Xpharm]	9.00	1	Fluarix Tetra

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by PHARMAC:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes; or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;
- viii) are living in the Seddon/Ward and rural Eastern Marlborough region (within the Nelson Marlborough District Health Board) and Kaikoura and Hurunui areas (within the Canterbury District Health Board);
 - ix) have been displaced from their homes in Edgecumbe and the surrounding region;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

ni 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	90.00	10	✓ Influyac Tetra

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

- a) Only on a prescription
- b) No patient co-payment payable

С

A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by PHARMAC:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders. or
 - f) haemoglobinopathies, or
 - a) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or
- c) children aged four years or less (but over three years) who have been hospitalised for respiratory illness or have a history of significant respiratory illness;
- d) people under 18 years of age living in the Seddon/Ward and rural Eastern Marlborough region (within the Nelson Marlborough District Health Board) and Kaikoura and Hurunui areas (within the Canterbury District Health Board):
- People under 18 years of age who have been displaced from their homes in Edgecumbe and the surrounding region;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

	Subsidy (Manufacturer's Price)	F Subsidi	ully Brand or sed Generic
	\$	Per	✓ Manufacturer
MEASLES, MUMPS AND RUBELLA VACCINE - [Xpharm]			
A maximum of two doses for any patient meeting the following	g criteria:		
 For primary vaccination in children; or 			
2) For revaccination following immunosuppression; or			
For any individual susceptible to measles, mumps or rul	,		
 A maximum of three doses for children who have had the 	eir first dose prior to	12 months.	
Note: Please refer to the Immunisation Handbook for approp	riate schedule for ca	tch up progra	ammes.
Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50			
Rubella virus 1,000 CCID50; prefilled syringe/ampoule of			
diluent 0.5 ml		10	✓ <u>Priorix</u>
MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGAT	E VACCINE - [Xph	arm]	
Any of the following:			
 Up to three doses and a booster every five years for part 			
or anatomic asplenia, HIV, complement deficiency (acq	,,	pre or post	solid organ transplant; or
One dose for close contacts of meningococcal cases; o			
A maximum of two doses for bone marrow transplant particular for the doses.	•		
4) A maximum of two doses for patients following immunos	• •		
Note: children under seven years of age require two doses 8	weeks apart, a boos	ster dose thre	e years after the primary
series and then five yearly. *Immunosuppression due to steroid or other immunosuppression.	sive therapy must be	for a pariod	of areator than 20 days
Inj 4 mcg of each meningococcal polysaccharide conjugated		ioi a peliou	of greater than 20 days.
a total of approximately 48 mcg of diphtheria toxoid carrie			
per 0.5 ml vial		1	✓ Menactra
MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm]			
Any of the following:			
Up to three doses and a booster every five years for particle.	ients pre- and post s	plenectomy	and for patients with functional
or anatomic asplenia, HIV, complement deficiency (acqu			•
2) One dose for close contacts of meningococcal cases; o	,.	F F	3
3) A maximum of two doses for bone marrow transplant pa	atients; or		
4) A maximum of two doses for patients following immunos	suppression*.		
Note: children under seven years of age require two doses 8	weeks apart, a boos	ster dose thre	ee years after the primary
series and then five yearly.			
*Immunosuppression due to steroid or other immunosuppression			,
Inj 10 mcg in 0.5 ml syringe	0.00	1	✓ Neisvac-C
PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm]		
Either:			
 A primary course of four doses for previously unvaccina 			
2) Up to three doses as appropriate to complete the prima	•	ation for indi	viduals under the age of
59 months who have received one to three doses of PC			
Note: please refer to the Immunisation Handbook for the app	•	r catch up pr	ogrammes
Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6E	,		
7F, 9V, 14 and 23F; 3 mcg of pneumococcal			
polysaccharide serotypes 4, 18C and 19F in 0.5 ml prefilled syringe	0.00	10	✓ Synflorix
promised syminge		10	- <u>Symiona</u>

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	1	Manufacturer	

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- One dose is funded for high risk children (over the age of 17 months and under 18 years) who have previously received four doses of PCV10: or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) with primary immune deficiencies; or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cyanosis or failure; or
 - I) with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- 4) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

	NATIONAL	IMMUNISA	TION SCHEDULE
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	d Generic
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE -[Xpharm]		
Either:	/ for notions and ha		
 Up to three doses (as appropriate) for patients with HIV chemotherapy; pre- or post-splenectomy or with function complement deficiency (acquired or inherited), cochlea All of the following: 	onal asplenia, pre- or p	post-solid orga	n transplant, renal dialysis,
a) Patient is a child under 18 years for (re-)immunisi	ation; and		
b) Treatment is for a maximum of two doses; andc) Any of the following:			
 i) on immunosuppressive therapy or radiation immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or 	therapy, vaccinate wl	hen there is ex	pected to be a sufficient
iv) with renal failure, or nephrotic syndrome; or			
v) who are immune-suppressed following orga		luding haemato	ppoietic stem cell transplant);
vi) with cochlear implants or intracranial shunts	s; or		
vii) with cerebrospinal fluid leaks; or			
 viii) receiving corticosteroid therapy for more that prednisone of 2 mg/kg per day or greater, o 20 mg or greater; or 			
ix) with chronic pulmonary disease (including a	sthma treated with his	ah-dose cortico	osteroid therapy); or
x) pre term infants, born before 28 weeks gest		9	7,7,
xi) with cardiac disease, with cyanosis or failure	e; or		
xii) with diabetes; or xiii) with Down syndrome; or			
xiii) with Down syndrome, or xiv) who are pre-or post-splenectomy, or with fu	nctional asplenia.		
an) mis are pro-criptor opinionally, or min a	asp.oa.		
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	0.00	1	Pneumovax 23
POLIOMYELITIS VACCINE - [Xpharm]			
Up to three doses for patients meeting either of the following			
 For partially vaccinated or previously unvaccinated indi For revaccination following immunosuppression. 	viduals; or		
Note: Please refer to the Immunisation Handbook for approling 80D antigen units in 0.5 ml syringe		1 1 0	nmes. ´ <mark>IPOL</mark>
ROTAVIRUS ORAL VACCINE - [Xpharm]			
Maximum of two doses for patients meeting the following:			
 first dose to be administered in infants aged under 14 v no vaccination being administered to children aged 24 			
Oral susp live attenuated human rotavirus			
1,000,000 CCID50 per dose, prefilled oral applicator	0.00	10	Rotarix

	e candidates for tranantation; or e competent inpatie splantation, on adviemotherapy, on ad r moderate immuncof major metabolic re immunocompronias no clinical history of ading to immune contact in transport in the contact in the c	ents.; or ice of their vice of the osuppress decompenised, or ury of varice varicella	on; or r specia eir spec sion on a nsation undergo ella, or and who	alist, or cialist, or advice of HIV specialist, or , with no clinical history of coing a procedure leading to o are severely	
1) Maximum of one dose for primary vaccination for either: a) Any infant born on or after 1 April 2016; or b) For previously unvaccinated children turning 11 year varicella infection (chickenpox), or 2) Maximum of two doses for any of the following: a) Any of the following for non-immune patients: i) with chronic liver disease who may in future bii) with deteriorating renal function before transpliii) prior to solid organ transplant; or iv) prior to any elective immunosuppression*, or v) for post exposure prophylaxis who are immun b) For patients at least 2 years after bone marrow tranticly for HIV positive non immune to varicella with mild of the HIV positive non immune to varicella with mild of the For patients with inborn errors of metabolism at risk varicella, or f) For household contacts of paediatric patients who a immune compromise where the household contact g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure le has no clinical history of varicella. * immunosuppression due to steroid or other immunosuppress 28 days Inj 2000 PFU prefilled syringe plus vial VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED	e candidates for tranantation; or e competent inpatie splantation, on adviemotherapy, on ad r moderate immuncof major metabolic re immunocompronias no clinical history of ading to immune contact in transport in the contact in the c	ents.; or ice of their vice of the osuppress decompenised, or ury of varice varicella	on; or r specia eir spec sion on a nsation undergo ella, or and who	alist, or ialist, or advice of HIV specialist, or , with no clinical history of sing a procedure leading to o are severely	
a) Any infant born on or after 1 April 2016; or b) For previously unvaccinated children turning 11 year varicella infection (chickenpox), or 2) Maximum of two doses for any of the following: a) Any of the following for non-immune patients: i) with chronic liver disease who may in future bii) with deteriorating renal function before transplii) prior to solid organ transplant; or iv) prior to any elective immunosuppression*, or v) for post exposure prophylaxis who are immun b) For patients at least 2 years after bone marrow tran c) For patients at least 6 months after completion of cl d) For HIV positive non immune to varicella with mild of e) For patients with inborn errors of metabolism at risk varicella, or f) For household contacts of paediatric patients who a immune compromise where the household contact g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure le has no clinical history of varicella. * immunosuppression due to steroid or other immunosuppress 28 days Inj 2000 PFU prefilled syringe plus vial WARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED	e candidates for tranantation; or e competent inpatie splantation, on adviemotherapy, on ad r moderate immuncof major metabolic re immunocompronias no clinical history of ading to immune contact in transport in the contact in the c	ents.; or ice of their vice of the osuppress decompenised, or ury of varice varicella	on; or r specia eir spec sion on a nsation undergo ella, or and who	alist, or cialist, or advice of HIV specialist, or , with no clinical history of coing a procedure leading to o are severely	
b) For previously unvaccinated children turning 11 year varicella infection (chickenpox), or 2) Maximum of two doses for any of the following: a) Any of the following for non-immune patients: i) with chronic liver disease who may in future bii) with deteriorating renal function before transpliii) prior to solid organ transplant; or iv) prior to any elective immunosuppression*, or v) for post exposure prophylaxis who are immun b) For patients at least 2 years after bone marrow tran c) For patients at least 6 months after completion of cl d) For HIV positive non immune to varicella with mild of e) For patients with inborn errors of metabolism at risk varicella, or f) For household contacts of paediatric patients who a immune compromise where the household contact g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure le has no clinical history of varicella. * immunosuppression due to steroid or other immunosuppress 28 days Inj 2000 PFU prefilled syringe plus vial	e candidates for tranantation; or e competent inpatie splantation, on adviemotherapy, on ad r moderate immuncof major metabolic re immunocompronias no clinical history of ading to immune contact in transport in the contact in the c	ents.; or ice of their vice of the osuppress decompenised, or ury of varice varicella	on; or r specia eir spec sion on a nsation undergo ella, or and who	alist, or cialist, or advice of HIV specialist, or , with no clinical history of coing a procedure leading to o are severely	
a) Any of the following for non-immune patients: i) with chronic liver disease who may in future bii) with deteriorating renal function before transplii) prior to solid organ transplant; or iv) prior to any elective immunosuppression*, or v) for post exposure prophylaxis who are immun b) For patients at least 2 years after bone marrow tran c) For patients at least 6 months after completion of cl d) For HIV positive non immune to varicella with mild of e) For patients with inborn errors of metabolism at risk varicella, or f) For household contacts of paediatric patients who a immune compromise where the household contact g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure le has no clinical history of varicella. * immunosuppression due to steroid or other immunosuppress 28 days Inj 2000 PFU prefilled syringe plus vial VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED	e competent inpatie splantation, on advi emotherapy, on ad r moderate immunc of major metabolic re immunocompron has no clinical history of ading to immune contact of the	ents.; or ice of their vice of the osuppress decompe nised, or ury of varice	r specia eir spec sion on a ensation undergo cella, or and who	ialist, or advice of HIV specialist, or , with no clinical history of bing a procedure leading to o are severely	
i) with chronic liver disease who may in future b ii) with deteriorating renal function before transpl iii) prior to solid organ transplant; or iv) prior to any elective immunosuppression*, or v) for post exposure prophylaxis who are immun b) For patients at least 2 years after bone marrow tran c) For patients at least 6 months after completion of cl d) For HIV positive non immune to varicella with mild of e) For patients with inborn errors of metabolism at risk varicella, or f) For household contacts of paediatric patients who a immune compromise where the household contact g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure le has no clinical history of varicella. * immunosuppression due to steroid or other immunosuppress 28 days Inj 2000 PFU prefilled syringe plus vial VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED	e competent inpatie splantation, on advi emotherapy, on ad r moderate immunc of major metabolic re immunocompron has no clinical history of ading to immune contact of the	ents.; or ice of their vice of the osuppress decompe nised, or ury of varice	r specia eir spec sion on a ensation undergo cella, or and who	ialist, or advice of HIV specialist, or , with no clinical history of sing a procedure leading to o are severely	
ii) with deteriorating renal function before transpliii) prior to solid organ transplant; or iv) prior to any elective immunosuppression*, or v) for post exposure prophylaxis who are immun b) For patients at least 2 years after bone marrow tran c) For patients at least 6 months after completion of cl d) For HIV positive non immune to varicella with mild of e) For patients with inborn errors of metabolism at risk varicella, or f) For household contacts of paediatric patients who a immune compromise where the household contact g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure le has no clinical history of varicella. * immunosuppression due to steroid or other immunosuppress 28 days Inj 2000 PFU prefilled syringe plus vial VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED	e competent inpatie splantation, on advi emotherapy, on ad r moderate immunc of major metabolic re immunocompron has no clinical history of ading to immune contact of the	ents.; or ice of their vice of the osuppress decompe nised, or ury of varice	r specia eir spec sion on a ensation undergo cella, or and who	ialist, or advice of HIV specialist, or , with no clinical history of sing a procedure leading to o are severely	
iv) prior to any elective immunosuppression*, or v) for post exposure prophylaxis who are immun b) For patients at least 2 years after bone marrow tran c) For patients at least 6 months after completion of cl d) For HIV positive non immune to varicella with mild of e) For patients with inborn errors of metabolism at risk varicella, or f) For household contacts of paediatric patients who a immune compromise where the household contact g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure le has no clinical history of varicella. * immunosuppression due to steroid or other immunosuppress 28 days Inj 2000 PFU prefilled syringe plus vial VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED	splantation, on advi emotherapy, on ad r moderate immund of major metabolic re immunocompron as no clinical history of clinical history of ading to immune co	ce of their lyice of the osuppress decompe nised, or u ry of varice varicella	eir spec sion on a ensation undergo cella, or and who	cialist, or advice of HIV specialist, or , with no clinical history of sing a procedure leading to o are severely	
b) For patients at least 2 years after bone marrow tran c) For patients at least 6 months after completion of cl d) For HIV positive non immune to varicella with mild of the positive non immune to varicella with mild of the positive non immune to varicella with mild of the positive non immune to varicella, or f) For household contacts of paediatric patients who a immune compromise where the household contact g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure least no clinical history of varicella. * immunosuppression due to steroid or other immunosuppress 28 days Inj 2000 PFU prefilled syringe plus vial	splantation, on advi emotherapy, on ad r moderate immund of major metabolic re immunocompron as no clinical history of clinical history of ading to immune co	ce of their lyice of the osuppress decompe nised, or u ry of varice varicella	eir spec sion on a ensation undergo cella, or and who	cialist, or advice of HIV specialist, or , with no clinical history of sing a procedure leading to o are severely	
c) For patients at least 6 months after completion of cl d) For HIV positive non immune to varicella with mild of e) For patients with inborn errors of metabolism at risk varicella, or f) For household contacts of paediatric patients who a immune compromise where the household contact g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure le has no clinical history of varicella. * immunosuppression due to steroid or other immunosuppress 28 days Inj 2000 PFU prefilled syringe plus vial	emotherapy, on ad r moderate immuno of major metabolic re immunocompron has no clinical histo no clinical history of ading to immune co	vice of the osuppress decompe nised, or u ry of varic varicella	eir spec sion on a ensation undergo cella, or and who	ialist, or advice of HIV specialist, or , with no clinical history of sing a procedure leading to o are severely	
d) For HIV positive non immune to varicella with mild of e) For patients with inborn errors of metabolism at risk varicella, or f) For household contacts of paediatric patients who a immune compromise where the household contact g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure le has no clinical history of varicella. * immunosuppression due to steroid or other immunosuppress 28 days Inj 2000 PFU prefilled syringe plus vial	r moderate immuno of major metabolic re immunocompron has no clinical histo no clinical history of ading to immune co	osuppress decompe nised, or u ry of varic varicella	sion on a ensation undergo cella, or and who	advice of HIV specialist, or , with no clinical history of oing a procedure leading to o are severely	
e) For patients with inborn errors of metabolism at risk varicella, or f) For household contacts of paediatric patients who a immune compromise where the household contact g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure le has no clinical history of varicella. * immunosuppression due to steroid or other immunosuppress 28 days Inj 2000 PFU prefilled syringe plus vial	of major metabolic re immunocompron las no clinical histo lo clinical history of ading to immune co	decompe nised, or u ry of varic varicella	ensation undergo cella, or and who	, with no clinical history of oing a procedure leading to o are severely	
f) For household contacts of paediatric patients who a immune compromise where the household contact g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure le has no clinical history of varicella. * immunosuppression due to steroid or other immunosuppress 28 days Inj 2000 PFU prefilled syringe plus vial	nas no clinical histo no clinical history of ading to immune co	ry of varic varicella	ella, or and wh	o are severely	
g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure le has no clinical history of varicella. * immunosuppression due to steroid or other immunosuppress 28 days Inj 2000 PFU prefilled syringe plus vial	o clinical history of ading to immune co	varicella	and wh	o are severely	
immunocompromised, or undergoing a procedure le has no clinical history of varicella. * immunosuppression due to steroid or other immunosuppress 28 days Inj 2000 PFU prefilled syringe plus vial	ading to immune co				
28 days Inj 2000 PFU prefilled syringe plus vial VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED	o thorony must be				
VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED		for a trea	itment p	period of greater than	
	0.00	1 10		<u>'arilrix</u> 'arilrix	
VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED VACCINE [SHINGLES VACCINE] – [Xpharm] Funded for patients meeting either of the following criteria:					
1) One dose for all people aged 65 years; or					
2) One dose for all people aged between 66 and 80 years in	clusive from 1 April	2018 and	d 31 Ma	rch 2020.	
Inj 19,400 PFU prefilled syringe plus vial	0.00	1 10	_	ostavax ostavax	
Diagnostic Agents					
TUBERCULIN PPD [MANTOUX] TEST - [Xpharm] Inj 5 TU per 0.1 ml, 1 ml vial			_		

- Symbols -		Agents Affecting the		Androderm	79
3TC	105	Renin-Angiotensin System	47	Animas Battery Cap	19
50X 3.0 Reservoir	25	Agents for Parkinsonism and Rela	ted	Animas Cartridge	25
- A -		Disorders	119	Anoro Ellipta	217
A-Scabies	67	Agents Used in the Treatment of		Antabuse	149
Abacavir sulphate	105	Poisonings	226	Antacids and Antiflatulents	6
Abacavir sulphate with		Agrylin	157	Anten	125
lamivudine	105	Alanase	219	Anthelmintics	88
Abiraterone acetate	168	Albendazole	88	Antiacne Preparations	60
Acarbose	11	Albey	213	Antiallergy Preparations	213
Acarbose Mylan	11	Albustix	76	Antianaemics	37
Accuretic 10	48	Aldurazyme	29	Antiandrogen Oral	
Accuretic 20	48	Alendronate sodium11	1-112	Contraceptives	74
Acetazolamide	223	Alendronate sodium with		Antiarrhythmics	
Acetic acid with 1, 2- propaned		colecalciferol	111	Antibacterials	
diacetate and		Alfacalcidol		Antibacterials Topical	
benzethonium	221	Alfamino Junior		Anticholinergic Agents	
Acetic acid with hydroxyquinol		Alginic acid		Anticholinesterases	
ricinoleic acid		Alglucosidase alfa		Antidepressants	
Acetylcysteine		Alkeran		Antidiarrhoeals	
Aci-Jel		Allersoothe		Antiepilepsy Drugs	
Aciclovir	74	Allmercap		Antifibrinolytics, Haemostatics and	
Infection	101	Allopurinol		Local Sclerosants	
Sensory		Alpha-Adrenoceptor Blockers		Antifibrotics	
Acidex		Alpha-Keri Lotion		Antifungals	
Acipimox		Alphamox 125		Antifungals Topical	
Acitretin		Alphamox 250		Antihistamines	
Aclasta		Alu-Tab		Antihypotensives	
Aclin		Aluminium hydroxide		Antimalarials	
Actemra		•		Antimigraine Preparations	
Actinomycin D		Amantadine hydrochloride Ambrisentan		ů .	
,				Antinausea and Vertigo Agents	
Actrapid Actrapid Penfill		Amiloride hydrochloride Amiloride hydrochloride with	52	Antiparasitics Antipruritic Preparations	
		furosemide	E0	Antipsychotics	
AcupanAdalat 10			55	Antiretrovirals	
		Amiloride hydrochloride with	E0		
Adalat Oros		hydrochlorothiazide		Antirheumatoid Agents	1111
Adamstone		Aminophylline	219	Antispasmodics and Other Agents	0
Adapalene		Amiodarone hydrochloride		Altering Gut Motility	
Adefin		Amisulpride		Antithrombotic Agents	41
Adefin XL		Amitriptyline		Antithymocyte globulin	177
Adefovir dipivoxil		Amlodipine		(equine)	
Adenuric		Amorolfine		Antitrichomonal Agents	98
ADR Cartridge 1.8		Amoxicillin		Antituberculotics and	00
Adrenaline		Amoxicillin with clavulanic acid		Antileprotics	
ADT Booster	250	Amphotericin B		Antiulcerants	
Adult diphtheria and tetanus	050	Amsacrine		Antivirals	
vaccine		AmsaLyo		Anxiolytics	
Advantan		Amsidine		Anzatax	
Advate		Amzoate		Apidra	
Afinitor		Anaesthetics		Apidra SoloStar	
Aflibercept		Anagrelide hydrochloride		Apo-Amlodipine	
AFT OLO (Analgesics		Apo-Amoxi	
AFT SLS-free		Anastrozole		Apo-Azithromycin	
AFT-Pyrazinamide	99	Andriol Testocaps	79	Apo-Bromocriptine	119

Apo-Ciclopirox	61	Hydrochlorothiazide	48	Bacillus Calmette-Guerin (BCG)	
Apo-Cilazapril	47	Arrow-Morphine LA	123	vaccine	17
Apo-Cilazapril/		Arrow-Norfloxacin	109	Bacillus Calmette-Guerin	
Hydrochlorothiazide	48	Arrow-Ornidazole	98	vaccine	250
Apo-Clarithromycin		Arrow-Quinapril 10	47	Baclofen	117
Alimentary	8	Arrow-Quinapril 20	47	Bactroban	60
Infection	89	Arrow-Quinapril 5	47	Barrier Creams and Emollients	64
Apo-Clomipramine	124	Arrow-Roxithromycin		BCG Vaccine	250
Apo-Diclo SR		Arrow-Sertraline		BD PosiFlush	4
Apo-Diltiazem CD		Arrow-Timolol	223	Beclazone 100	
Apo-Doxazosin		Arrow-Tolterodine	76	Beclazone 250	
Apo-Folic Acid		Arrow-Topiramate		Beclazone 50	
Apo-Gabapentin		Arrow-Tramadol		Beclomethasone	
Apo-Leflunomide		Arsenic trioxide		dipropionate2	14, 219
Apo-Megestrol		Asacol		Bee venom allergy treatment	
Apo-Metoprolol		Asamax		Bendamustine hydrochloride	
Apo-Mirtazapine		Ascorbic acid		Bendrofluazide	
Apo-Moclobemide		Aspen		Bendroflumethiazide	
Apo-Montelukast		Aspen Adrenaline	56	[Bendrofluazide]	5
Apo-Nadolol		Aspirin		BeneFIX	40
Apo-Nicotinic Acid		Blood	41	Benzathine benzylpenicillin	
Apo-Ondansetron		Nervous		, ,	
		Asthalin		Benzatropine mesylate Benzbromaron AL 100	
Apo-Oxybutynin		Atazanavir sulphate		Benzbromarone	
Apo-Paroxetine					
Apo-Perindopril		Atenolol		Benzoin	
Apo-Pindolol		Atenolol AFT		Benztrop	
Apo-Pravastatin		ATGAM		Benzydamine hydrochloride	
Apo-Prazosin		Ativan		Benzylpenicillin sodium [Penicillin)
Apo-Prednisone		Atomoxetine		G]	
Apo-Primidone	128	Atorvastatin		Beta Cream	
Apo-Propranolol		Atripla	105	Beta Ointment	
Apo-Pyridoxine		Atropine sulphate	40	Beta Scalp	
Apo-Ropinirole		Cardiovascular		Beta-Adrenoceptor Agonists	
Apo-Selegiline S29	119	Sensory		Beta-Adrenoceptor Blockers	
Apo-Sumatriptan		Atropt		Betadine	
Apo-Terazosin		Atrovent		Betadine Skin Prep	
Apo-Timol		Aubagio		Betaferon	
Apomorphine hydrochloride		Augmentin		Betagan	
Aprepitant		Aurorix		Betahistine dihydrochloride	
Apresoline		AutoSoft 30		Betaine	
Aptamil Gold+ Pepti Junior		AutoSoft 90		Betaloc CR	
Aqueous cream		Avelox		Betamethasone dipropionate	
Aripiprazole		Avomine		Betamethasone dipropionate with	
Aripiprazole Sandoz		Avonex		calcipotriol	
Aristocort		Avonex Pen		Betamethasone sodium phospha	
Arrow - Clopid	41	Azacitidine	154	with betamethasone acetate	78
Arrow-Amitriptyline	124	Azacitidine Dr Reddy's	154	Betamethasone valerate	63, 69
Arrow-Bendrofluazide	53	Azathioprine	171	Betamethasone valerate with	
Arrow-Brimonidine	224	Azithromycin		clioquinol	63
Arrow-Calcium	35	Azol	87	Betamethasone valerate with soc	dium
Arrow-Diazepam		Azopt	223	fusidate [fusidic acid]	
Arrow-Dortim		AZT [']	105–106	Betaxolol	
Arrow-Doxorubicin		- B -		Betnovate	6
Arrow-Fluoxetine		B-D Micro-Fine	13	Betnovate-C	
Arrow-Lamotrigine		B-D Ultra Fine	14	Betoptic	22
Arrow-Losartan &		B-D Ultra Fine II	14	Betoptic S	

Bezafibrate53	Budesonide	CareSens PRO1
Bezalip53	Alimentary6	Carmellose sodium with gelatin and
Bezalip Retard53	Respiratory214, 220	pectin3
Bicalutamide169	Budesonide with eformoterol215	Carmustine15
Bicillin LA91	Bumetanide52	Carvedilol5
BiCNU153	Buprenorphine with naloxone148	Carvedilol Sandoz5
Bile and Liver Therapy9	Bupropion hydrochloride149	Catapres5
Biltricide88	Burinex52	Cathejell12
Bimatoprost224	Buscopan8	CeeNÚ15
Bimatoprost Actavis224	Buspirone hydrochloride135	Cefaclor monohydrate8
Bimatoprost Multichem224	Busulfan153	Cefalexin8
Binarex169	- C -	Cefalexin Sandoz8
Binocrit38	Cabergoline87	Cefazolin8
Biodone123	Cafergot130	Ceftriaxone8
Biodone Extra Forte123	Cafergot S29130	Cefuroxime axetil8
Biodone Forte123	Caffeine citrate220	Celebrex11
Bisacodyl27	Calamine62	Celecoxib11
Bisoprolol fumarate50	Calcipotriol68	Celecoxib Pfizer11
BK Lotion65	Calcitonin77	Celestone Chronodose7
Bleomycin sulphate158	Calcitriol34	Celiprolol5
Blood Colony-stimulating	Calcitriol-AFT34	Cellcept17
Factors44	Calcium carbonate6, 35	Celol5
Blood glucose diagnostic test	Calcium Channel Blockers51	Centrally-Acting Agents5
meter 12	Calcium Disodium Versenate227	Cephalexin ABM8
Blood glucose diagnostic test	Calcium folinate155	Cerezyme3
strip13	Calcium Folinate Ebewe155	Cetirizine hydrochloride21
Blood glucose test strips (visually	Calcium Folinate Sandoz155	Cetomacrogol6
impaired)13	Calcium gluconate35	Cetomacrogol with glycerol6
Blood Ketone Diagnostic Test	Calcium Homeostasis77	Cetuximab18
Strip 11	Calcium polystyrene sulphonate46	Champix15
Bonjela33	Calcium Resonium46	Charcoal22
Bonvit26	Calogen233	Chemotherapeutic Agents15
Boostrix250	Calsource35	Chickenpox vaccine26
Bortezomib	Candesartan cilexetil48	Chlorafast22
Bosentan57	Candestar48	Chlorambucil15
Bosentan Dr Reddy's57	Canesten61	Chloramphenicol22
Bosentan-Mylan57	Capecitabine155	Chlorhexidine gluconate
Bosvate50	Capoten47	Alimentary3
Bplex33	Capsaicin	Dermatological6
Breo Ellipta215	Musculoskeletal111	Chloroform22
Brevinor 1/2172	Nervous121	Chlorothiazide5
Brevinor 1/2872	Captopril47	Chlorpheniramine maleate21
Brevinor 2172	Carafate9	Chlorpromazine hydrochloride13
Bricanyl Turbuhaler216	Carbaccord	Chlorsig22
Brilinta42	Carbamazepine 127	Chlortalidone [Chlorthalidone]5
Brimonidine tartrate224	Carbimazole82	Chlorthalidone5
Brimonidine tartrate with timolol	Carbomer225	Chlorvescent4
maleate224	Carboplatin 153	Choice Load 3757
Brinov	Carboplatin Ebewe153	Choice TT380 Short7
Brinzolamide223	Carbosorb-X226	Choice TT380 Standard7
Brolene221	Cardinol LA51	Cholestyramine5
Bromocriptine mesylate119	CareSens Dual	Choline salicylate with cetalkonium
Brufen SR110	CareSens N	chloride3
BSF Entecavir Sandoz226	CareSens N POP	Ciclopirox olamine6
Buccastem131	CareSens N Premier12	Ciclosporin21
		Cilazapril4

Cilazapril with		Colchicine	117	Dactinomycin [Actinomycin D]	158
hydrochlorothiazide	48	Colecalciferol	34	Daivobet	68
Cilicaine	92	Colestid	54	Daivonex	68
Cilicaine VK	92	Colestipol hydrochloride	54	Daktarin	
Cinacalcet	77	Colgout	117	Dalacin C	9
Cipflox	93	Colifoam		Dalteparin sodium	42
Ciprofloxacin		Colistin sulphomethate	93	Danazol	
Infection	93	Colistin-Link	93	Dantrium	117
Sensory		Collodion flexible		Dantrium S29	117
Ciprofloxacin Teva		Colloidal bismuth subcitrate		Dantrolene	
Circadin		Colofac	8	Daonil	
Cisplatin	153	Coloxyl	26	Dapa-Tabs	5
Cisplatin Ebewe		Combigan		Dapsone	
Citalopram hydrobromide		Compound electrolytes		Daraprim	
Cladribine		Compound electrolytes with glu-		Darunavir	
Clarithromycin		[Dextrose]		Dasatinib	
Alimentary	8	Compound hydroxybenzoate		Daunorubicin	
Infection		Concerta		DBL Acetylcysteine	
Clexane		Condoms		DBL Aminophylline	
Clindamycin		Condyline		DBL Bleomycin Sulfate	
Clindamycin ABM		Contact-D		DBL Carboplatin	
Clinicians Renal Vit		Contraceptives - Hormonal		DBL Cisplatin	15
Clobazam		Contraceptives - Non-hormonal		DBL Dacarbazine	
Clobetasol propionate		Copaxone		DBL Desferrioxamine Mesylate for	
Clobetasone butyrate		Cordarone-X		BP	
Clofazimine		Corticosteroids and Related Age		DBL Docetaxel	
Clomazol		for Systemic Use		DBL Ergometrine	
Dermatological	61	Corticosteroids Topical		DBL Gemcitabine	156
Genito-Urinary		Cosentyx		DBL Gentamicin	
Clomifene citrate		Cosmegen		DBL Leucovorin Calcium	
Clomipramine hydrochloride		Coumadin		DBL Methotrexate Onco-Vial	
Clonazepam126–12		Creon 10000		DBL Morphine Sulphate	
Clonidine		Creon 25000		DBL Morphine Tartrate	
Clonidine BNM		Crotamiton		DBL Naloxone Hydrochloride	
Clonidine hydrochloride		Crystaderm		DBL Octreotide	
		•		DBL Pethidine Hydrochloride	
Clopidogrel		Curam Cvite		DBL Vincristine Sulfate	
Clopine				De-Worm	
Clopixol13	00, 100	Cyclizine hydrochloride			
Clotrimazole	61	Cyclizine lactate		Decozol	
Dermatological		Cyclogyl		Deferasirox	
Genito-Urinary		Cyclopentolate hydrochloride		Deferiprone Denosumab	
		Cyclophosphamide			
Clustran		Cycloserine		Deolate	
Clustran		Cyklokapron		Deoxycoformycin	
Co-trimoxazole		Cyproterone acetate	/9	Depo-Medrol	
Coal tar	68	Cyproterone acetate with	7.4	Depo-Medrol with Lidocaine	
Coal tar with allantoin, menthol,		ethinyloestradiol		Depo-Provera	
phenol and sulphur	68	Cystadane		Depo-Testosterone	/\
Coal tar with salicylic acid and		Cytarabine		Deprim	9
sulphur		Cytotec		DermAssist	
Coco-Scalp	68	Cytoxan	153	Dermol	
Codeine phosphate		-D-		Desferal	
Extemporaneous		D-Penamine		Desferrioxamine mesilate	
Nervous		Dabigatran	44	Desmopressin acetate	8
Cogentin		Dacarbazine		Desmopressin-PH&T	8
Colaspase [L-asparaginase]	158	Dacarbazine APP	158	Detection of Substances in	

Urine	Disinfecting and Cleansing	Efavirenz with emtricitabine and
Dexamethasone	Agents 64	tenofovir disoproxil
Hormone78	Disopyramide phosphate49	fumarate105
Sensory222	Disulfiram149	Effient42
Dexamethasone phosphate78	Diuretics52	Eformoterol fumarate218
Dexamethasone with framycetin and	Diurin 4052	Eformoterol fumarate dihydrate218
gramicidin221	Docetaxel159	Efudix70
Dexamethasone with neomycin	Docetaxel Sandoz159	Egopsoryl TA68
sulphate and polymyxin B	Docusate sodium26	Elaprase29
sulphate222	Docusate sodium with	Elecare24
Dexamfetamine sulfate145	sennosides26	Elecare LCP247
Dexmethsone78	Dolutegravir106	Elelyso3
Dextrochlorpheniramine	Domperidone 131	Elemental 028 Extra238
maleate214	Donepezil hydrochloride148	Elocon63
Dextrose45–46	Donepezil-Rex148	Elocon Alcohol Free63
DHC Continus122	Dopress	Eltrombopag38
Diabetes9	Dornase alfa219	Eltroxin82
Diabetes Management11	Dortimopt224	EMB Fatol99
Diacomit	Dorzolamide hydrochloride223	Emend Tri-Pack
Diagnostic Agents260	Dorzolamide with timolol224	EMLA12
Diamide Relief6	Dostinex	Emtricitabine 105
Diamox	Dosulepin [Dothiepin]	Emtricitabilite with tenofovir disoproxil
Diasip	hydrochloride124	fumarate102
Diason RTH234		Emtriva102
Diazepam	Dothiepin 124 Doxazosin 47	Emulsifying ointment68
	Doxepin hydrochloride125	Enalapril maleate4
Diazoxide 9 Dibenzyline 47		
•	Doxine	Enbrel 17 ² Endocrine Therapy
Diclofenac Sandoz110		
Diclofenac sodium	Doxorubicin hydrochloride159	Endoxan
Musculoskeletal110	Doxy-50	Enerlyte
Sensory222	Doxycycline	Engerix-B252
Differin	DP Fusidic Acid Cream61	Enlafax XR126
Difflam32	DP Lotion65	Enoxaparin sodium43
Diflucan95	DP Lotn HC63	Ensure242
Diflucan S2995	DP-Allopurinol116	Ensure Plus242
Diflucortolone valerate63	Dr Reddy's Omeprazole9	Ensure Plus HN24
Digestives Including Enzymes25	Drugs Affecting Bone	Ensure Plus RTH24
Digoxin49	Metabolism 111	Entacapone119
Dihydrocodeine tartrate122	Dual blood glucose and blood ketone	Entapone119
Dilantin	diagnostic test meter 12	Entecavir100
Dilantin Infatab128	Duocal Super Soluble Powder232	Entecavir Sandoz100
Diltiazem hydrochloride52	Duolin216	Entocort CIR
Dilzem52	Duolin HFA216	Entresto 24/2648
Dimethicone64, 66	Durex Confidence71	Entresto 49/5148
Dimethyl fumarate135	Durex Extra Safe71	Entresto 97/10348
Dipentum7	Duride56	Epilim128
Diphtheria, tetanus and pertussis	- E -	Epilim Crushable128
vaccine 250	e-chamber La Grande220	Epilim IV128
Diphtheria, tetanus, pertussis and	e-chamber Mask220	Epilim S/F Liquid128
polio vaccine250	e-chamber Turbo220	Epilim Syrup128
Diphtheria, tetanus, pertussis, polio,	E-Mycin90	Epirubicin Ebewe159
hepatitis B and haemophilus	Ear Preparations221	Epirubicin hydrochloride159
influenzae type B vaccine 251	Ear/Eye Preparations221	Eplerenone53
Diprosone62	Easiphen Liquid246	Epoetin alfa38
Diprosone OV62	Econazole nitrate61	Epoprostenol59
Dipyridamole41	Efavirenz105	Eprex
r /		F

Eptacog alfa [Recombinant factor	Febuxostat	117	Fluorouracil sodium	70
VIIa]40	Feed Thickener Karicare		Fluoxetine hydrochloride	126
ERA90	Aptamil	244	Flupenthixol decanoate	133
Erbitux186	FEIBA NF		Flutamide	
Ergometrine maleate74	Felo 10 ER	51	Flutamide Mylan	
Ergonovine74	Felo 5 ER	51	Flutamin	169
Ergotamine tartrate with	Felodipine	51	Fluticasone	21
caffeine130	Fenpaed		Fluticasone furoate with	
Erlotinib164	Fentanyl		vilanterol	21
Erythrocin IV90	Fentanyl Sandoz		Fluticasone propionate	
Erythromycin ethyl succinate90	Ferinject		Fluticasone with salmeterol	
Erythromycin lactobionate90	Ferodan		FML	
Erythromycin stearate90	Ferric carboxymaltose		Foban	
Esbriet218	Ferriprox		Folic acid	
Escitalopram125	Ferro-F-Tabs		Food Thickeners	
Escitalopram-Apotex125	Ferro-tab		Foods And Supplements For Inbo	
Eskazole	Ferrograd		Errors Of Metabolism	
Estradot80	Ferrosig		Foradil	
			Forteo	
Estradot 50 mcg80	Ferrous fumarate			
Estrofem	Ferrous fumarate with folic acid		Fortini	
Etanercept	Ferrous sulphate		Fortini Multi Fibre	
Ethambutol hydrochloride99	Ferrum H		Fortisip	
Ethics Aspirin	Fexofenadine hydrochloride		Fortisip Multi Fibre	
Ethics Aspirin EC41	Fibro-vein		Fosamax1	
Ethics Enalapril47	Filgrastim		Fosamax Plus	
Ethics Lisinopril47	Finasteride		Fragmin	
Ethinyloestradiol81	Fingolimod		Framycetin sulphate	
Ethinyloestradiol with	Firazyr		Frisium	
desogestrel72	Flagyl		Frumil	
Ethinyloestradiol with	Flagyl-S		Frusemide	
levonorgestrel72	Flamazine	61	Frusemide-Claris	
Ethinyloestradiol with	Flecainide acetate	49	Fucicort	
norethisterone72	Fleet Phosphate Enema	27	Fucidin	
Ethosuximide127	Flixonase Hayfever & Allergy	220	Fucithalmic	
Etopophos159	Flixotide	215	Fungilin	3
Etoposide	Flixotide Accuhaler	215	Furosemide [Frusemide]	52
Etoposide phosphate159	Floair	215	fusidic acid	
Etravirine105	Florinef	78	Dermatological	61, 64
Eumovate63	Fluanxol	133	Infection	9
Everet	Fluarix Tetra	254	Sensory	
Everolimus211	Flucil	91	- G -	
Evista113	Flucloxacillin		Gabapentin	12
Exelon148	Flucloxin	91	Gacet	12
Exemestane171	Fluconazole		Galsulfase	
Exjade226	Fludara Oral		Galvumet	
Extemporaneously Compounded	Fludarabine Ebewe		Galvus	
Preparations and	Fludarabine phosphate		Gardasil 9	
Galenicals229	Fludrocortisone acetate		Gastrodenol	
Eye Preparations221	Fluids and Electrolytes		Gaviscon Double Strength	
Eylea	Flumetasone pivalate		Gaviscon Infant	
Ezetimibe 54	Fluocortolone caproate with	44 1	Gazyva	10
Ezetimibe Sandoz 54	fluocortolone pivalate and		Gefitinib	
Ezetimibe Sandoz	cinchocaine	7	Gemcitabine Ebewe	
- F -	Fluorometholone			
			Gemcitabine hydrochloride	
Factor eight inhibitor bypassing	Fluorouracil		Gemfibrozil	
fraction40	Fluorouracil Ebewe	155	Gemzar	156

Genoptic22		Hydroxychloroquine	111
Genox17	healthE Glycerol BP229	Hydroxyurea	
Gentamicin sulphate	healthE Urea Cream65	Hygroton	
Infection9	Healtheries Simple Baking Mix244	Hylo-Fresh	
Sensory22		Hymenoptera	
Gilenya13	7 Heparin sodium44	Hyoscine butylbromide	
Ginet7	4 Heparinised saline44	Hyoscine hydrobromide	
Glatiramer acetate14		Hypam	144
Glecaprevir with pibrentasvir10	2 Hepatitis A vaccine251	Hyperuricaemia and Antigout	116
Glibenclamide1	1 Hepatitis B recombinant	Hypromellose	225
Gliclazide1	1 vaccine	Hypromellose with dextran	225
Glipizide1		Hysite	224
Glivec16	4 Herceptin207	-1-	
Glizide1	1 Hexamine hippurate108	Ibiamox	91
Glucagen Hypokit	9 Hiberix251	Ibuprofen	110
Glucagon hydrochloride	9 Hiprex108	Icatibant	213
Glucerna Select23	4 Histaclear214	Idarubicin hydrochloride	159
Glucerna Select RTH23	4 Histafen214	Idursulfase	
Glucobay1	1 Holoxan153	Ifosfamide	
Glucose [Dextrose]4		Ikorel	56
Gluten Free Foods24		lloprost	
Glycerin with sodium saccharin22		Imatinib mesilate	
Glycerin with sucrose22		Imatinib-AFT	164
Glycerol	HPV253	Imiglucerase	
Alimentary2	7 Humalog10	Imipramine hydrochloride	
Extemporaneous22		Imiquimod	
Glyceryl trinitrate	Humalog Mix 5010	Immune Modulators	
Alimentary		Immunosuppressants	
Cardiovascular5		Imuran	171
Glycopyrronium21	· · · · · · · · · · · · · · · · · · ·	Incruse Ellipta	
Glycopyrronium bromide		Indacaterol	
Glycopyrronium with	Humira177	Indapamide	
indacaterol21		Infanrix IPV	
Glytrin		Infanrix-hexa	
Gold Knight7		Infant Formulae	
Goserelin		Infatrini	
Gutron4		Infliximab	
Gynaecological Anti-infectives		Influenza vaccine	
- H -	Hydralazine56	Influvac	
Habitrol15		Influvac Tetra	
Haemophilus influenzae type B	Hydrea159	Inhaled Corticosteroids	
vaccine25	•	Inhaled Long-acting	
Haldol13		Beta-adrenoceptor Agonists	. 215
Haldol Concentrate13	3	Inset 30	21
Haldol Decanoas		Inset II	
Haloperidol13		Inspra	
Haloperidol decanoate13		Insulin aspart	
Hamilton Sunscreen		Insulin aspart with insulin aspart	
Harvoni10		protamine	10
Havrix	•	Insulin glargine	10
Havrix Junior25	,	Insulin glulisine	10
HBvaxPRO		Insulin isophane	
healthE Calamine Aqueous Cream	Hydrogen peroxide	Insulin isophane with insulin	10
BP6		neutral	10
healthE Dimethicone 10%	4 Dermatological 60	Insulin lispro	
healthE Dimethicone 4% Lotion6		Insulin lispro with insulin lispro	10
	, a.		

protamine	10	Ivermectin	66	Latanoprost	22
Insulin neutral	10	- J -		Lax-Suppositories	2
Insulin pen needles	13	Jadelle	73	Lax-Tab	
Insulin pump	14	Jakavi	167	Laxatives	2
Insulin pump accessories	19	Jevity HiCal RTH	241	Laxsol	2
Insulin pump cartridge	19	Jevity RTH	241	Ledipasvir with sofosbuvir	10
Insulin pump infusion set (steel		Juno Pemetrexed		Leflunomide	
cannula)	20	- K -		Lenalidomide	
Insulin pump infusion set (steel		Kaletra	106	Letrole	17
cannula, straight insertion)	21	Kemadrin	119	Letrozole	17
Insulin pump infusion set (teflon		Kenacomb	221	Leukeran FC	15
cannula, angle insertion with		Kenacort-A 10		Leukotriene Receptor	
insertion device)	21	Kenacort-A 40	79	Antagonists	21
Insulin pump infusion set (teflon		Kenalog in Orabase	33	Leunase	
cannula, angle insertion)	22	Ketocal 3:1		Leuprorelin	8
Insulin pump infusion set (teflon		KetoCal 4:1	249	Leustatin	
cannula, straight insertion with		Ketoconazole		Levetiracetam	12
insertion device)	23	Dermatological	69	Levetiracetam-AFT	12
Insulin pump infusion set (teflon		Infection		Levlen ED	
cannula, straight insertion)	24	Ketogenic Diet		Levobunolol	
Insulin pump reservoir		Ketoprofen		Levocabastine	22
Insulin syringes, disposable with		KetoSens		Levodopa with benserazide	11
attached needle	14	Keytruda		Levodopa with carbidopa	
Intal Forte CFC Free		Kindergen		Levomepromazine	
Intelence	105	Kinson		hydrochloride	13
Interferon alfa-2a		Kivexa	105	Levomepromazine maleate	
Interferon alfa-2b		Klacid	89	Levonorgestrel	
Interferon beta-1-alpha		Kliogest		Genito-Urinary	<mark>7</mark>
Interferon beta-1-beta		Kliovance		Hormone	
Intra-uterine device		Kogenate FS	41	Levothyroxine	
Intron-A	107	Konakion MM		Lidocaine [Lignocaine]	
Invega Sustenna		Konsyl-D		Lidocaine [Lignocaine]	
IPOĽ		Kuvan		hydrochloride	12
Ipratropium bromide21		-L-		Lidocaine [Lignocaine] with	
Iressa		L-asparaginase	158	chlorhexidine	12
Irinotecan Actavis 100		Labetalol		Lidocaine [Lignocaine] with	
Irinotecan hydrochloride		Lacosamide		prilocaine	12
Irinotecan-Rex		Lactulose		Lidocaine-Claris	
Iron polymaltose		Laevolac		Lignocaine	
Isentress		Lamictal		Hormone	7
Ismo 20		Lamivudine		Nervous	
Ismo 40 Retard		Lamivudine Alphapharm		Lioresal Intrathecal	11
Isoniazid		Lamotrigine		Lipazil	
Isoniazid with rifampicin		Lamprene		Lipid-Modifying Agents	
Isoprenaline [Isoproterenol]		Lanoxin		Liquigen	
Isoproterenol		Lanoxin PG		Lisinopril	4
Isoptin		Lanoxin S29		Lithicarb FC	
Isopto Carpine		Lansoprazole	8	Lithium carbonate	
Isosorbide mononitrate		Lantus		Livostin	
Isosource Standard		Lantus SoloStar		LMX4	
Isotretinoin		Lanvis		Locacorten-Viaform ED's	
Ispaghula (psyllium) husk		Lanzol Relief		Local preparations for Anal and	
Isuprel		Lapatinib ditosylate		Rectal Disorders	
Itch-Soothe		Largactil		Locasol	
Itraconazole		Laronidase		Locoid	63.6
Itrazole		Lasix		Locoid Crelo	6

Locoid Lipocream	63	Mebendazole88 Methylxanthines	
Locorten-Vioform		Mebeverine hydrochloride8 Metoclopramide Actavis 10	
Lodi		Medrol78 Metoclopramide hydrochloride	
Lodoxamide		Medroxyprogesterone acetate Metolazone	
Logem		Genito-Urinary73 Metopirone	
Lomide	223	Hormone80–81 Metoprolol succinate	
Lomustine	153	Medsurge Metoprolol tartrate	
Loniten		Cardiovascular52 Metronidazole	
Loperamide hydrochloride		Musculoskeletal117 Metroprolol IV Mylan	
Lopinavir with ritonavir		Mefenamic acid110 Metyrapone	
Loprofin		Megestrol acetate169 Mexiletine hydrochloride	
Loprofin Mix	246	Melatonin	4
Lorafix	214	Melphalan154 Miacalcic	
Loratadine	214	Menactra257 Micolette	<mark>2</mark>
Lorazepam		Meningococcal (groups A, C, Y and Miconazole	3
Lorfast		W-135) conjugate vaccine 257 Miconazole nitrate	
Lormetazepam		Meningococcal C conjugate Dermatological	
Lorstat	54	vaccine257 Genito-Urinary	
Losartan Actavis	48	Menthol62 Micreme	
Losartan potassium	48	Mercaptopurine156 Micreme H	6
Losartan potassium with		Mercilon 28	
hydrochlorothiazide	48	Mesalazine7 Microgynon 30	7
Lovir	101	Mesna160 Microgynon 50 ED	7
Lucrin Depot 1-month	86	Mestinon110 Microlut	<mark>7</mark> :
Lucrin Depot 3-month	86	Metabolic Disorder Agents27 Midazolam	14
Ludiomil	125	Metchek11 Midazolam-Claris	14
Lycinate	55	Meterol215 Midodrine	4
Lyderm	67	Metformin hydrochloride11 Minerals	
- M -		Metformin Mylan11 Mini-Wright AFS Low Range	220
m-Eslon		Methadone hydrochloride Mini-Wright Standard	220
Mabthera		Extemporaneous229 Minidiab	
Macrogol 3350 with potassium		Nervous	
chloride, sodium bicarbonat	e and	Methatabs123 Minims Pilocarpine	22
sodium chloride	<mark>27</mark>	Methopt225 Minims Prednisolone	22
Macrogol 400 and propylene		Methotrexate	
glycol	225	Methotrexate Ebewe156 Mino-tabs	
Madopar 125	119	Methotrexate Sandoz156 Minocycline hydrochloride	9
Madopar 250		Methyl hydroxybenzoate229 Minomycin	9
Madopar 250 Madopar 62.5	119	Methyl hydroxybenzoate	
Madopar 62.5Madopar HBS	119 119 119	Methylcellulose	60 50
Madopar 62.5	119 119 119 119	Methylcellulose	
Madopar 62.5	119 119 119 119 229	Methylcellulose	60 8 ¹ 120
Madopar 62.5	119 119 119 119 229 36	Methylcellulose	60 8 ¹ 120
Madopar 62.5	119 119 119 119 229 36	Methylcellulose	
Madopar 62.5	119 119 119 119 229 36 260	Methylcellulose	60 8 120
Madopar 62.5	11911911911922936260125	Methylcellulose 229 Minor Skin Infections Methylcellulose with glycerin and sodium saccharin 229 Mirena Methylcellulose with glycerin and sucrose 230 Mirazapine Methyldopa 52 Mitomycin C Methyldopa Mylan 52 Mitozantrone Methylnaltrexone bromide 27 Mitozantrone Ebewe	668120160
Madopar 62.5	119119119119119229362602601254469	Methylcellulose	
Madopar 62.5	119119119119119229362602601254469	Methylcellulose	
Madopar 62.5		Methylcellulose with glycerin and sodium saccharin 229 Minor Skin Infections Minoxidil Mirena Mirtazapine Misoprostol Misoprostol Methyldopa Mylan 52 Mitoyantrone Methylnaltrexone bromide 27 Methylphenidate hydrochloride extended-release 147 Minoz Skin Infections Minoxidil Minoxidil Mirena Mirtazapine Mirena Mirtazapine Mirtazapine Mirtazapine Mirtazapine Misoprostol Misoprostol Misoprostol Mitowacin C Mitowacin C Mitozantrone Ebewe Mitozantrone Ebewe Moclobemide Moclobemide Moclobemide Moclobemide Modafinil Modafinil Minoxidil Minoxidil Mirena Mirtazapine Misoprostol Misoprostol Misoprostol Misoprostol Mitozantrone Ebewe Moclobemide Moclobemide Moclobemide Modafinil Modafinil Minoxidil Misoprostol Misopr	
Madopar 62.5		Methylcellulose with glycerin and sodium saccharin	
Madopar 62.5		Methylcellulose with glycerin and sodium saccharin	
Madopar 62.5		Methylcellulose with glycerin and sodium saccharin	
Madopar 62.5		Methylcellulose with glycerin and sodium saccharin. 229 Methylcellulose with glycerin and sucrose. 230 Methyldopa	66 68 8 8 12 12 16 16 16 16 17 14 14 14 14 15 5 6 6 6 6
Madopar 62.5		Methylcellulose with glycerin and sodium saccharin. 229 Methylcellulose with glycerin and sucrose. 230 Methyldopa	66 68 8 8 12 12 16 16 16 16 17 14 14 14 14 15 5 6 6 6 6
Madopar 62.5		Methylcellulose with glycerin and sodium saccharin. 229 Methylcellulose with glycerin and sucrose. 230 Methyldopa	

VIII]	40	Neo-B12	33	Normison	144
Morphine hydrochloride	123	Neo-Mercazole	82	Norpress	125
Morphine sulphate	123	Neocate Gold	247	Nortriptyline hydrochloride	125
Morphine tartrate		Neocate Junior Unflavoured	247	Norvir	
Motetis		Neocate Junior Vanilla		NovaSource Renal	237
Mouth and Throat	32	Neocate LCP	247	Novatretin	68
Movapo	119	Neocate SYNEO	247	NovoMix 30 FlexPen	10
Moxifloxacin		Neoral	211	NovoRapid	10
MSUD Maxamum	245	Neostigmine metilsulfate	110	NovoRapid FlexPen	10
Mucilaginous laxatives with		Nepro HP (strawberry)	237	NovoRapid Penfill	10
stimulants	26	Nepro HP (vanilla)		NovoSeven RT	40
Mucolytics	219	Nepro HP RTH		Noxafil	97
Mucosoothe	120	Nerisone	63	Nozinan	132
Multiple Sclerosis Treatments	135	Neulactil	133	Nuelin	219
Multivitamin renal	34	Neulastim	45	Nuelin-SR	219
Multivitamins	34	NeuroTabs	35	Nutilis	244
Mupirocin	60	Nevirapine	105	Nutrient Modules	231
Muscle Relaxants	117	Nevirapine Alphapharm	105	Nutrini Energy Multi Fibre	236
Mvite	34	Nicorandil		Nutrini Energy RTH	
Myambutol	99	Nicotine	150	Nutrini Low Energy Multi Fibre	238
Mycobutin		Nicotinic acid	54	Nutrini RTH	236
MycoNail		Nifedipine		Nutrison 800 Complete Multi	
Mycophenolate mofetil	171	Nifuran	109	Fibre	241
Mycostatin	62	Nilotinib	165	Nutrison Concentrated	243
Mydriacyl	225	Nilstat		Nutrison Energy	241
Mylan Atenolol	50	Alimentary	33	Nutrison Energy Multi Fibre	241
Mylan Clomiphen		Genito-Urinary	74	Nutrison Multi Fibre	241
Myleran	153	Infection	96	Nutrison Standard RTH	241
Myocrisin		Nintedanib	217	Nyefax Retard	51
Myometrial and Vaginal Hormo		Nipent	161	Nystatin	
Preparations	74	Nitrados		Alimentary	33
Myozyme		Nitrates	55	Dermatological	62
Mysoline S29		Nitrazepam		Genito-Urinary	
- N -		Nitroderm TTS		Infection	
Nadolol	50	Nitrofurantoin	109	NZB Low Gluten Bread Mix	244
Naglazyme	28	Nitrolingual Pump Spray	55	-0-	
Nalcrom	7	Nivolumab		O/W Fatty Emulsion Cream	65
Naloxone hydrochloride	226	Nizoral	96	Obinutuzumab	193
Naltraccord		Noctamid	143	Octocog alfa [Recombinant factor	
Naltrexone hydrochloride	149	Nodia	6	VIII] (Advate)	
Naphazoline hydrochloride	225	Noflam 250	110	Octocog alfa [Recombinant factor	
Naphcon Forte		Noflam 500	110	VIII] (Kogenate FS)	
Naprosyn SR 1000		Non-Steroidal Anti-Inflammatory		Octreotide	
Naprosyn SR 750		Drugs	110	Octreotide LAR (somatostatin	
Naproxen		Nonacog alfa [Recombinant facto	r	analogue)	169
Nardil	125	IX]		Oestradiol	
Nasal Preparations	219	Nonacog gamma, [Recombinant		Oestradiol valerate	80
Natalizumab		Factor IX]	40	Oestradiol with norethisterone	80
Natulan	161	Norethisterone		Oestriol	
Nausafix		Genito-Urinary	73	Genito-Urinary	74
Nausicalm		Hormone	81	Hormone	
Nauzene		Norflex		Oestrogens	
Navelbine		Norfloxacin		Ofev	
Nedocromil		Noriday 28		Oil in water emulsion	
Nefopam hydrochloride		Norimin		Olanzapine13	
Neisvac-C		Normacol Plus		Olbetam	

Olopatadine	.225	Ozurdex	222	Paraldehyde	
Olsalazine	7	- P -		Parasidose	6
Omalizumab	. 193	Pacifen	117	Parasiticidal Preparations	
Omeprazole	9	Paclitaxel	160	Parnate	12
Omeprazole actavis 10	9	Paclitaxel Actavis	160	Paromomycin	94
Omeprazole actavis 20	9	Paclitaxel Ebewe	160	Paroxetine	12
Omeprazole actavis 40	9	Paediatric Seravit	34	Paser	9
Omnitrope	82	Paliperidone	134	Patanol	22
Onbrez Breezhaler	.215	Pamidronate disodium		Paxam	13
Oncaspar	.160	Pamisol	113	Pazopanib	16
OncoTICE	.177	Pancreatic enzyme	25	Peak flow meter	
Ondansetron	.131	Pantoprazole	9	Pedialyte - Bubblegum	4
Ondansetron ODT-DRLA	.131	Panzop Relief		Pediasure	23
Ondansetron ODT-ORLA	.131	Panzytrat		Pediasure RTH	
One-Alpha	34	Papaverine hydrochloride	56	Pegaspargase	16
Opdivo	.209	Para-amino salicylic acid	99	Pegasys	
Ora-Blend	.230	Paracare	122	Pegfilgrastim	
Ora-Blend SF	.229	Paracare Double Strength	122	Pegylated interferon alfa-2a	
Ora-Plus		Paracetamol		Pembrolizumab	
Ora-Sweet		Paracetamol + Codeine		Pemetrexed	
Ora-Sweet SF	.229	(Relieve)	124	Penicillamine	
Orabase		Paracetamol Pharmacare		Penicillin G	
Oral Supplements/Complete Diet		Paracetamol with codeine		PenMix 30	
(Nasogastric/Gastrostomy Tube		Paradigm 1.8 Reservoir		PenMix 40	
Feed)	234	Paradigm 3.0 Reservoir		PenMix 50	
Oratane		Paradigm Mio MMT-921		Pentasa	
Orgran		Paradigm Mio MMT-923		Pentostatin [Deoxycoformycin]	
Orion Temozolomide		Paradigm Mio MMT-925		Pentoxifylline [Oxpentifylline]	
Ornidazole		Paradigm Mio MMT-941		Peptamen Junior	
Orphenadrine citrate		Paradigm Mio MMT-943	23	Peptisoothe	
Ortho-tolidine		Paradigm Mio MMT-945		Peptisorb	. 23
Oruvail SR		Paradigm Mio MMT-965		Perhexiline maleate	
Osmolite RTH		Paradigm Mio MMT-975		Pericyazine	
Other Endocrine Agents		Paradigm Quick-Set MMT-386		Perindopril	4
Other Oestrogen Preparations		Paradigm Quick-Set MMT-387		Perjeta	
Other Progestogen		Paradigm Quick-Set MMT-396		Permethrin	
Preparations	81	Paradigm Quick-Set MMT-397		Perrigo	6
Other Skin Preparations		Paradigm Quick-Set MMT-398		Pertuzumab	
Ovestin		Paradigm Quick-Set MMT-399		Peteha	
Genito-Urinary	74	Paradigm Silhouette MMT-368		Pethidine hydrochloride	
Hormone		Paradigm Silhouette MMT-377		Pevaryl	6
Ox-Pam		Paradigm Silhouette MMT-378		Pexsig	
Oxaliccord		Paradigm Silhouette MMT-381		Pfizer Exemestane	
Oxaliplatin		Paradigm Silhouette MMT-382		Pharmacy Health Sorbolene with	
Oxaliplatin Actavis 100		Paradigm Silhouette MMT-383		Glycerin	6
Oxaliplatin Ebewe		Paradigm Silhouette MMT-384		Pharmacy Services	
Oxazepam		Paradigm Sure-T MMT-864		Pheburane	
Oxis Turbuhaler		Paradigm Sure-T MMT-866		Phenelzine sulphate	
Oxpentifylline		Paradigm Sure-T MMT-874		Phenobarbitone	. 12
Oxybutynin		Paradigm Sure-T MMT-876		Phenobarbitone sodium	
Oxycodone hydrochloride	.124	Paradigm Sure-T MMT-884		Extemporaneous	231
OxyNorm		Paradigm Sure-T MMT-886		Nervous	
Oxytocin		Paraffin		Phenothrin	
Oxytocin BNM		Paraffin liquid with soft white	00	Phenoxybenzamine	
Oxytocin with ergometrine	, +	paraffin	225	hydrochloride	4
maleate	74	Paraffin liquid with wool fat		Phenoxymethylpenicillin (Penicillin	*

V)	92	Praziquantel	88	Pulmocare	23
Phenytoin sodium		Prazosin		Pulmozyme	
Phlexy 10	246	Pred Forte	223	Puri-nethol	15
Phosphate Phebra		Prednisolone	79	Puria	3
Phosphate-Sandoz		Prednisolone acetate	223	Pyrazinamide	9
Phosphorus		Prednisolone sodium		Pyridostigmine bromide	
Phytomenadione		phosphate	223	Pyridoxine hydrochloride	
Pilocarpine hydrochloride		Prednisolone-AFT		Pyrimethamine	
Pimafucort		Prednisone	79	Pytazen SR	4
Pindolol	50	Pregabalin	128	- Q -	
Pine tar with trolamine laurilsul	fate	Pregabalin Pfizer		Q 300	9
and fluorescein	68	Pregnancy Tests - hCG Urine .		Questran-Lite	5
Pinetarsol	68	Premarin	80	Questran-Lite S29	5
Pioglitazone	11	Prevenar 13	258	Quetapel	13
Piportil		Prezista	106	Quetiapine	
Pipothiazine palmitate		Priadel	133	Quick-Set MMT-390	
Pirfenidone		Priceline		Quick-Set MMT-391	
Pizotifen		Primacin		Quick-Set MMT-392	
PKU Anamix Infant		Primaquine phosphate		Quick-Set MMT-393	
PKU Anamix Junior		Primidone		Quinapril	
PKU Anamix Junior Chocolate		Primolut N		Quinapril with	
PKU Anamix Junior LQ		Priorix		hydrochlorothiazide	4
PKU Anamix Junior Vanilla		Probenecid		Quinine sulphate	
PKU Lophlex LQ 10		Probenecid-AFT		Qvar	
PKU Lophlex LQ 20		Procaine penicillin		-R-	
PKU Lophlex Powder		Procarbazine hydrochloride		RA-Morph	19
PKU Lophlex Sensation 20		Prochlorperazine		Raloxifene hydrochloride	
Plaquenil		Proctosedyl		Raltegravir potassium	
Plendil ER		Procur			
				RamipexRanbaxy-Cefaclor	
Pneumococcal (PCV10) conjug		Procyclidine hydrochloride			
vaccine		Procytox		Ranitidine	
Pneumococcal (PCV13) conjug	-	Progesterone		Ranitidine Relief	
vaccine	258	Proglicem		Rapamune	21
Pneumococcal (PPV23)	050	Proglycem	9	Reandron 1000	/
polysaccharide vaccine		Progynova		Recombinant factor IX	
Pneumovax 23		Prokinex		Recombinant factor VIIa	
Podophyllotoxin		Prolia		Recombinant factor VIII	
Polaramine		Promethazine hydrochloride		Rectogesic	
Poliomyelitis vaccine		Promethazine theoclate		Redipred	
Poloxamer		Propafenone hydrochloride		Refresh Night Time	
Poly-Gel		Propamidine isethionate		Relieve	
Poly-Tears	225	Propranolol		Relistor	
Poly-Visc		Propylene glycol		Remicade	
Polycal	231	Propylthiouracil	82	Renilon 7.5	
Polyvinyl alcohol		Protaphane		Resonium-A	
Ponstan	110	Protaphane Penfill	10	Resource Beneprotein	23
Posaconazole	97	Protifar	233	Resource Diabetic	23
Postinor-1	73	Protionamide	99	Respigen	21
Potassium chloride	45-46	Provera	80	Respiratory Devices	
Potassium citrate		Provera HD	81	Respiratory Stimulants	22
Potassium iodate	35	Provera S29	80	Retinol palmitate	22
Povidone iodine	66	PSM Citalopram		ReTrieve	
Pradaxa		Psoriasis and Eczema		Retrovir	
Pramipexole hydrochloride		Preparations	68	Revlimid	
Prasugrel		PTU		Revolade	
Pravastatin		Pulmicort Turbuhaler		Rexacrom	

RexAir	216	Sandostatin LAR	169	Sodium cromoglicate	
Reyataz	106	Sapropterin dihydrochloride		Alimentary	
Ribomustin	152	Scalp Preparations	69	Respiratory	
Ricit		Scopoderm TTS	131	Sensory	22
Rifabutin	99	Sebizole	69	Sodium fluoride	3
Rifadin	100	Secukinumab	203	Sodium Fusidate [fusidic acid]	
Rifampicin	100	Sedatives and Hypnotics	143	Dermatological	6
Rifaximin		Seebri Breezhaler	217	Infection	9
Rifinah	99	Selegiline hydrochloride	119	Sensory	22
Rilutek	120	Senna	<mark>27</mark>	Sodium hyaluronate [Hyaluronic	
Riluzole	120	Senokot	<mark>27</mark>	acid]	22
Riodine	66	Sensipar		Sodium phenylbutyrate	
Risedronate Sandoz		SensoCard	13	Sodium polystyrene sulphonate	
Risedronate sodium		Serenace		Sodium tetradecyl sulphate	4
Risperdal Consta	135	Seretide	216	Sodium valproate	
Risperidone	133, 135	Seretide Accuhaler		Sofradex	
Risperon	133	Serevent	215	Soframycin	22
Ritalin	146	Serevent Accuhaler	215	Solian	
Ritalin LA	147	Serophene		Solifenacin Mylan	7
Ritalin SR	146	Sertraline	126	Solifenacin succinate	70
Ritonavir	106	Sevredol	123	Solu-Cortef	
Rituximab	195	Sex Hormones Non		Solu-Medrol	7
Rivaroxaban		Contraceptive	79	Solu-Medrol-Act-O-Vial	
Rivastigmine	148	Shield 49	71	Somatropin (Omnitrope)	8
Rivotril	126–127	Shield Blue	71	Sotalol	5
RIXUBIS	40	Shield XL	71	Spacer device	
Rizamelt	130	shingles vaccine	260	Span-K	4
Rizatriptan	130	Sildenafil		Spiolto Respimat	21
Roferon-A	107	Silhouette MMT-371	<mark>22</mark>	Spiractin	5
Rolin	171	Silhouette MMT-373	<mark>22</mark>	Spiriva	21
Ropinirole hydrochloride	119	Siltuximab	204	Spiriva Respimat	21
Rotarix	259	Simvastatin	54	Spironolactone	5
Rotavirus oral vaccine	259	Simvastatin Mylan	54	Sporanox	9
Roxane		Sinemet		Sprycel	16
Alimentary	6	Sinemet CR	119	Staphlex	9
Cardiovascular	51	Sirolimus	211	Stemetil	
Roxithromycin	90	Siterone	79	SteroClear	22
Rubifen		Slow-Lopresor	50	Stesolid	120
Rubifen SR	146	Smith BioMed Rapid Pregnand	СУ	Stimulants/ADHD Treatments	14
Rulide D	90	Test	74	Stiripentol	12
Ruxolitinib	167	Sodibic	46	Stocrin	10
Rythmodan	49	Sodium acid phosphate	<mark>27</mark>	Stomahesive	3
Rytmonorm		Sodium alginate	6	Strattera	
-\$-		Sodium aurothiomalate	111	Stromectol	
Sabril	129	Sodium benzoate	30	Suboxone	14
Sacubitril with valsartan	48	Sodium bicarbonate		Sucralfate	
SalAir	216	Blood	45-46	Sulfadiazine Silver	
Salazopyrin	7	Extemporaneous	230	Sulfadiazine sodium	9
Salazopyrin EN	7	Sodium calcium edetate		Sulfasalazine	
Salbutamol	216	Sodium chloride		Sulindac	110
Salbutamol with ipratropium		Blood	45	Sulphur	
bromide		Respiratory		Sulprix	
Salicylic acid		Sodium citrate with sodium lau		Sumatriptan	
Salmeterol		sulphoacetate		Sunitinib	
Sandomigran		Sodium citro-tartrate		Sunscreens	
Sandomigran S29	130				

Sunscreens, proprietary	69	Teriflunomide	140	Trastuzumab	20
Sure-T MMT-863	20	Teriparatide	113	Travatan	224
Sure-T MMT-865	20	Testosterone	79	Travoprost	224
Sure-T MMT-873	20	Testosterone cipionate	79	Travopt	224
Sure-T MMT-875	20	Testosterone esters	79	Treatments for Dementia	148
Sure-T MMT-883	20	Testosterone undecanoate	79	Treatments for Substance	
Sure-T MMT-885	20	Tetrabenazine	120	Dependence	148
Sustagen Diabetic	234	Tetrabromophenol	76	Trental 400	56
Sustagen Hospital Formula		Tetracosactrin	79	Tretinoin	
Active	242	Tetracyclin Wolff	92	Dermatological	60
Sustanon Ampoules	79	Tetracycline	92	Oncology	162
Sutent	167	Thalidomide	162	Trexate	156
Sylvant	204	Thalomid	162	Triamcinolone acetonide	
Symbicort Turbuhaler 100/6	215	Theophylline	219	Alimentary	3
Symbicort Turbuhaler 200/6	215	Thiamine hydrochloride	33	Dermatological	6
Symbicort Turbuhaler 400/12	215	THIO-TEPA	154	Hormone	79
Symmetrel	119	Thioguanine	157	Triamcinolone acetonide with	
Sympathomimetics	<mark>56</mark>	Thiotepa	154	gramicidin, neomycin and nys	statin
Synacthen	79	Thymol glycerin		Dermatological	64
Synacthen Depot	79	Thyroid and Antithyroid Agents		Sensory	22
Synacthene Retard	79	Ticagrelor	42	Triazolam	144
Synflorix	257	Tilade	219	Trichozole	98
Synthroid	82	Tilcotil	110	Triclosan	
Syntometrine	74	Timolol		Trimethoprim	9
Syrup (pharmaceutical grade)	230	Cardiovascular		Trimethoprim with	
Systane Unit Dose	225	Sensory	223	sulphamethoxazole	
- T -		Timoptol XE		[Co-trimoxazole]	9
Tacrolimus	212	Tiotropium bromide	217	Trisequens	
Tacrolimus Sandoz	212	Tiotropium bromide with		Trisul	
Taliglucerase alfa	31	olodaterol	217	Trophic Hormones	
Tambocor	49	Tivicay	106	Tropicamide	22
Tambocor CR	49	TMP	95	Trusopt	223
Tamoxifen citrate	170	TOBI	95	TruSteel	2
Tamoxifen Sandoz		Tobramycin		Truvada	102
Tamsulosin hydrochloride	75	Infection	95	Tuberculin PPD [Mantoux] test	260
Tamsulosin-Rex	75	Sensory	222	Tubersol	260
Tandem Cartridge	19	Tobramycin Mylan	95	Two Cal HN	24
Tandem t:slim X2	14	Tobrex	222	Two Cal HN RTH	
Tap water	230	Tocilizumab	204	Tykerb	16
Tarceva		Tofranil	125	Tysabri	138
Tasigna	165	Tolcapone	119	- U -	
Tasmar	119	Tolterodine	76	Ultibro Breezhaler	217
Tecfidera		Topamax	129	Ultraproct	
Tegretol	127	Topical Products for Joint and		Umeclidinium	
Tegretol CR	127	Muscular Pain	111	Umeclidinium with vilanterol	
Telfast	214	Topiramate	129	Univent2	216, 220
Temazepam	144	Topiramate Actavis	129	Ural	
Temizole 20	161	Total parenteral nutrition (TPN)	45	Urea	6
Temozolomide		TPN		Urex Forte	
Tenofovir disoproxil		Tramadol hydrochloride	124	Urinary Agents	
Tenofovir Disoproxil Teva	101	Tramal SR 100	124	Urinary Tract Infections	
Tenoxicam		Tramal SR 150		Uromitexan	160
Tepadina		Tramal SR 200		Ursodeoxycholic acid	
Terazosin	47	Trandate		Ursosan	2
Terbinafine		Tranexamic acid		Utrogestan	8
Terbutaline sulphate	216	Tranylcypromine sulphate	125	- V -	

Vaccinations250	Vitamin B6 2533
Vaclovir101	Vitamins33–34
Valaciclovir101	Vivonex TEN238
Valcyte101	Volibris56
Valganciclovir101	Voltaren110
Vancomycin95	Voltaren D110
Vannair215	Voltaren Ophtha222
Varenicline Pfizer150	Volumatic220
Varenicline tartrate150	Voriconazole97
Varicella vaccine [Chickenpox	Vosol221
vaccine]260	Votrient
Varicella zoster virus (Oka strain) live	Vttack
attenuated vaccine [shingles	- W -
vaccine] 260	Warfarin sodium44
Varilrix260	Wart Preparations69
Various	Wasp venom allergy treatment213
Vasodilators	Water
Vasopressin Agonists	Blood46
Vedafil	Extemporaneous230
Velcade	Wool fat with mineral oil65
Veletri	- X -
Venlafaxine 126	Xarelto44
Venomil	Xifaxan 9
Ventavis 59	XMET Maxamum245
Ventolin	
	Xolair
Vepesid	XP Maxamum246
Verapamil hydrochloride52 Vergo 16130	
	Xylocaine 120 Xylocaine 2% Jelly 120
Vermox	
Verpamil SR	Xyntha40
Vesanoid	Zantac8
Vesicare	
Vexazone11	Zapril
Vfend	
Viaderm KC64	Zaroxolyn53
Vidaza	Zarzio44
Vigabatrin	Zavedos
Vildagliptin11	Zeffix
Vildagliptin with metformin	Zeldox
hydrochloride11	Zetlam100
Vimpat	Ziagen
Vinblastine sulphate162	Zidovudine [AZT]105
Vincristine sulphate162	Zidovudine [AZT] with
Vinorelbine163	lamivudine106
Vinorelbine Ebewe163	Zimybe55
Viramune Suspension105	Zinc and castor oil64
ViruPOS221	Zinc sulphate36
Vistil225	Zincaps36
Vistil Forte225	Zinnat88
Vit.D334	Ziprasidone133
VitA-POS225	Zista214
Vitabdeck34	Zithromax89
Vitadol C33	Zoladex86
Vital238	Zoledronic acid
Vitamin A with vitamins D and C 33	Hormone77
Vitamin B complex33	Musculoskeletal114

Zoledronic acid Mylan	7
Zometa	77
Zopiclone	
Zopiclone Actavis	145
Zostavax	260
Zostrix	111
Zostrix HP	121
Zuclopenthixol decanoate	135
Zuclopenthixol hydrochloride	133
Zusdone	133
Zyban	149
Zypine	
Zypine ODT	133
Zyprexa Relprevv	
Zvtina	