5

6

June	2019
Volume 26	Number 1

Editors:

Kave Wilson & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am - 5pm weekdays)

Circulation

Published each April, August and December. Changes to the contents are published in monthly updates.

Accessible in an electronic format at no cost from the Health Professionals section of the PHARMAC website www.pharmac.govt.nz

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month. Alternatively there is a nominal charge for an annual subscription to the printed Schedule publications. To access either of these subscriptions visit our subscription website www.schedule.co.nz.

Production

Typeset automatically from XML and T_FX. XML version of the Schedule available from www.pharmac.govt.nz/pub/schedule/archive/

Programmers

Anrik Drenth & John Geering

email: texschedule@pharmac.govt.nz



ISSN 1179-3686 pdf ISSN 1172-9376 print

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to PHARMAC and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to PHARMAC should be in written form and not by reproduction of the PHARMAC logo. While care has been taken in compiling this Schedule, PHARMAC takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

Section A	General Rules
Section B	Alimentary Tract & Metabolism
	Blood & Blood Forming Organs
	Cardiovascular System

	Ourdiovasodiar Oystorn
60	Dermatologicals
71	Genito Urinary System
77	Hormone Preparations – Systemic
88	Infections – Agents For Systemic Use
110	Musculoskeletal System
118	Nervous System
151	Oncology Agents & Immunosuppressants
212	Respiratory System & Allergies
220	Sensory Organs
225	Various

Section C	Extemporaneous Compounds (ECPs)	227
Section D	Special Foods	230
Section I	National Immunisation Schedule	249
	Index	259

Introducing PHARMAC

The Pharmaceutical Management Agency (PHARMAC) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. PHARMAC negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

PHARMAC's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about PHARMAC and the way we make funding decisions can be found on the PHARMAC website at http://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by PHARMAC.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements PHARMAC has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gram g	
kilogram kg	
international unit iu	

Abbreviations

Capsule Cream Device Dispersible Effervescent Emulsion	Amp Cap Crm Dev Disp Eff Emul EC
Enteric Coated	EC

microgram me	cg
milligramn	ng
millilitreı	ml

millimole	mmol
unit	u

Gelatinous	Gel	SolutionSoln
Granules	Gran	SuppositorySupp
Infusion	Inf	TabletTab
Injection	Inj	Tincture Tinc
Liquid	Liq	Trans Dermal Delivery
Long Acting	LA	SystemTDDS
Ointment	Oint	-
Sachet	Sach	

General Rules for the Pharmaceutical Schedule are located on the PHARMAC website.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$) Per	Fully Subsidised	
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg p sachet SODIUM ALGINATE		30	v	Gaviscon Infant
* Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (8.60)	60		Gaviscon Double Strength
* Oral liq 500 mg with sodium bicarbonate 267 mg and calciur carbonate 160 mg per 10 ml		500 m	I	Acidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE * Tab 600 mg CALCIUM CARBONATE Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsement Only when prescribed for children under 12 years of age endorsed accordingly.		100 500 m nate bir	🗸	Alu-Tab Roxane nt and the prescription is
Antidiarrhoeals Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on	a PSO			
* Tab 2 mg * Cap 2 mg	10.75	400 400		<u>Nodia</u> Diamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE Cap 3 mg – Special Authority see SA1155 below – Retail pharmacy		90 valid for		Entocort CIR
the following criteria: Both:				
 Mild to moderate ileal, ileocaecal or proximal Crohn's dise Any of the following: 2.1 Diabetes; or 2.2 Cushingoid habitus; or 	ease; and			
2.3 Osteoporosis where there is significant risk of frac	ture; or			
				continued.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

continued...

2.4 Severe acne following treatment with conventional corticosteroid therapy; or

2.5 History of severe psychiatric problems associated with corticosteroid treatment; or

2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or

2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)2	6.55 21	.1 g OP 🖌	Colifoam
MESALAZINE			
Tab 400 mg4	9.50	100 🖌	Asacol
Tab EC 500 mg4	9.50	100 🗸	Asamax
Tab long-acting 500 mg5	9.05	100 🖌	Pentasa
Tab 800 mg	5.50	90 🖌	Asacol
Modified release granules, 1 g14	1.72 1	20 OP 🖌 🖌	Pentasa
Enema 1 g per 100 ml4	1.30	7 🖌	Pentasa
Suppos 500 mg2		20 🖌	Asacol
Suppos 1 g5	4.60	30 🖌	Pentasa
OLSALAZINE			
Tab 500 mg9	3.37	100 🖌	Dipentum
Cap 250 mg5		100 🗸	Dipentum
SODIUM CROMOGLICATE			
Cap 100 mg	2.91	100 🖌	Nalcrom
SULFASALAZINE			
* Tab 500 mg	4.00	100 🗸	Salazopyrin
* Tab EC 500 mg1			Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE

Oint 950 mcg, with fluocortolone pivalate 920 mcg, and		
cinchocaine hydrochloride 5 mg per g6.35	30 g OP	 Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and		
cinchocaine hydrochloride 1 mg2.66	12	 Ultraproct
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	 Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g	12	 Proctosedyl

	Subsidy	F	ully Brand or
	(Manufacturer's Price) \$	Subsidis Per	,
Management of Anal Fissures			
GLYCERYL TRINITRATE – Special Authority see SA1329 below * Oint 0.2%		0 g OP	✓ Rectogesic
► SA1329 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid chronic anal fissure that has persisted for longer than three week		wal unless no	ptified where the patient has a
Antispasmodics and Other Agents Altering Gut	Motility		
GLYCOPYRRONIUM BROMIDE Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on PSO		10	✓ Max Health
HYOSCINE BUTYLBROMIDE		10	
Tab 10 mg Tab 10 mg			✓ <u>Buscopan</u> ✓ Buscopan
MEBEVERINE HYDROCHLORIDE * Tab 135 mg		90	✓ Colofac
Antiulcerants			
Antisecretory and Cytoprotective			
MISOPROSTOL * Tab 200 mcg	41.50	120	✓ Cytotec
Helicobacter Pylori Eradication			
CLARITHROMYCIN Tab 500 mg – Subsidy by endorsement a) Maximum of 14 tab per prescription	10.40	14	✓ <u>Apo-Clarithromycin</u>
 b) Subsidised only if prescribed for helicobacter pylori e Note: the prescription is considered endorsed if clari inhibitor and either amoxicillin or metronidazole. 			0,
H2 Antagonists			
RANITIDINE – Only on a prescription * Tab 150 mg		500 300 ml	 <u>Ranitidine Relief</u> <u>Ranitidine Relief</u> <u>Peptisoothe</u> Zantac
Proton Pump Inhibitors			
LANSOPRAZOLE * Cap 15 mg * Cap 30 mg			 ✓ Lanzol Relief ✓ Lanzol Relief

Xifaxan

56

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
OMEPRAZOLE				
For omeprazole suspension refer Standard Formulae, pag	e 227			
* Cap 10 mg	1.98	90	 Image: A set of the set of the	Omeprazole actavis 10
* Cap 20 mg	1.96	90	1	Omeprazole actavis 20
* Cap 40 mg	3.12	90	1	Omeprazole actavis 40
* Powder – Only in combination		5 g	1	Midwest
Only in extemporaneously compounded omeprazole s	uspension.			
* Inj 40 mg ampoule with diluent		5	v	<u>Dr Reddy's</u> Omeprazole
PANTOPRAZOLE				
* Tab EC 20 mg	2.41	100	✓	Panzop Relief
* Tab EC 40 mg	3.35	100	✓	Panzop Relief
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE				
Tab 120 mg	14.51	50	1	Gastrodenol S29
SUCRALFATE				
Tab 1 g	35 50	120		
100 1 9	(48.28)	120		Carafate
Bile and Liver Therapy				

RIFAXIMIN - Special Authority see SA146	1 below – Retail pharmacy
Tab 550 mg	625.00

➡SA1461 Special Authority for Subsidy

Initial application only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid for 6 months where the patient has hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

Renewal only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Diabetes

Hyperglycaemic Agents

DIAZOXIDE - Special Authority see SA1320 below - Retail pha	armacy		
Cap 25 mg.	-	100	Proglicem S29
Cap 100 mg		100	 Proglicem S29
Oral lig 50 mg per ml		30 ml OP	✓ Proglycem S29
► SA1320 Special Authority for Subsidy			-37
Initial application from any relevant practitioner. Approvals va	lid for 12 months	where used for	the treatment of confirmed
hypoglycaemia caused by hyperinsulinism.			
Renewal from any relevant practitioner. Approvals valid without appropriate and the patient is benefiting from treatment.	t further renewal	unless notified	where the treatment remains
GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit – Up to 5 kit available on a PSO	32.00	1	 Glucagen Hypokit

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Pr \$	ice) Subs Per	Fully idised	Brand or Generic Manufacturer
Insulin - Short-acting Preparations				
NSULIN NEUTRAL ▲ Inj human 100 u per ml	25.26	10 ml OP		ctrapid
▲ Inj human 100 u per ml, 3 ml	42.66	5	🗸 A	umulin R ctrapid Penfill umulin R
Insulin - Intermediate-acting Preparations				
NSULIN ASPART WITH INSULIN ASPART PROTAMINE		5	🗸 N	ovoMix 30 FlexPen
NSULIN ISOPHANE ▲ Inj human 100 u per ml		10 ml OP		umulin NPH
Inj human 100 u per ml, 3 ml	29.86	5	🗸 Н	rotaphane umulin NPH rotaphane Penfill
NSULIN ISOPHANE WITH INSULIN NEUTRAL Inj human with neutral insulin 100 u per ml	25.26	10 ml OP		umulin 30/70 lixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ H ✓ P ✓ P	umulin 30/70 enMix 30 enMix 40 enMix 50
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE Inj lispro 25% with insulin lispro protamine 75% 100 u per ml,			• •	enività 50
3 ml Inj lispro 50% with insulin lispro protamine 50% 100 u per ml, 3 ml		5 5		umalog Mix 25
		5	• п	umalog Mix 50
Insulin - Long-acting Preparations				
Inj 100 u per ml, 10 ml	63.00 94.50	1 5		antus antus
Inj 100 u per ml, 3 ml disposable pen		5		antus SoloStar
Insulin - Rapid Acting Preparations				
NSULIN ASPART Inj 100 u per ml, 10 ml Inj 100 u per ml, 3 ml Inj 100 u per ml, 3 ml syringe	51.19	1 5 5	🗸 N	ovoRapid ovoRapid Penfill ovoRapid FlexPen
NSULIN GLULISINE Inj 100 u per ml, 10 ml Inj 100 u per ml, 3 ml		1 5		pidra pidra
Inj 100 u per ml, 3 ml disposable pen NSULIN LISPRO		5		pidra SoloStar
 Inj 100 u per ml, 10 ml Inj 100 u per ml, 3 ml 		10 ml OP 5		umalog umalog

10

ALIMENTARY	TRACT	AND N	METABOLIS	М
------------	-------	-------	------------------	---

	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully dised	Brand or Generic Manufacturer
Alpha Glucosidase Inhibitors				
ACARBOSE			_	
* Tab 50 mg		90		<u>Glucobay</u> Accarb
* Tab 100 mg	10.47	90		Accarb Glucobay
* Tab Too mg	11.24	50 50	-	Acarbose Mylan S29
	11.24	00	• •	
	20.23	90	✓	Accarb
(Acarbose Mylan S29) Tab 100 mg to be delisted 1 October 20	019)			
Oral Hypoglycaemic Agents				
GLIBENCLAMIDE				
* Tab 5 mg	6.00	100	✓ [Daonil
GLICLAZIDE				
* Tab 80 mg	10.29	500	√	Glizide
GLIPIZIDE				
* Tab 5 mg	3.27	100	✓ I	<u> Minidiab</u>
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg		1,000		Apotex
* Tab immediate-release 850 mg	7.04	500	 I 	Apotex
PIOGLITAZONE				
* Tab 15 mg		90	-	/exazone
* Tab 30 mg * Tab 45 mg		90 90	-	<u>/exazone</u> /exazone
5		90	• •	VEXAZUITE
VILDAGLIPTIN Tab 50 mg	40.00	60	1	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE		00	- (uaivud
Tab 50 mg with 1,000 mg metformin hydrochloride	40.00	60	1	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride		60 60		Galvumet
			- (our annot

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or

5)	metabolic disease	or epilepsy un	ider the care	of a paediatrician	, neurologist or m	etabolic specialis
The	prescription must be	endorsed ac	cordinaly			

i ne prescription must be endorsed accordingly.	
Test strips15.5	0

.... 15.50 10 strip OP

KetoSens

Dual Blood Glucose and Blood Ketone Testing DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER – Subsidy by endorsement a) Maximum of 1 pack per prescription b) Up to 1 pack available on a PSO c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has: 1) type 1 diabetes; or 2) permanent neonatal diabetes; or 3) undergone a pancreatectomy; or 4) cystic fibrosis-related diabetes; or 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist. The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). The avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter. Meter with 50 lancets, a lancing device and 10 blood glucose diagnostic test strips BLOOD GLUCOSE DIAGNOSTIC TEST METER – Subsidy by endorsement a) Maximum of 1 pack per prescription b) Up to 1 pack available on a PSO c) A diagnostic blood glucose test meter is subsidised for a patient who: 1) is receiving insulin or sulphonylurea therapy; or 2) is pregnant with diabetes; or 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diab		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
a) Maximum of 1 pack per prescription b) Up to 1 pack available on a PSO c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has: 1) type 1 diabetes; or 2) permanent neonatal diabetes; or 3) undergone a pancreatectomy; or 4) cystic fibrosis-related diabetes; or 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist. The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). I the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter. Meter with 50 lancets, a lancing device and 10 blood glucose diagnostic test strips	Dual Blood Glucose and Blood Ketone Testing				
Meter with 50 lancets, a lancing device and 10 blood glucose diagnostic test strips	 DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC a) Maximum of 1 pack per prescription b) Up to 1 pack available on a PSO c) A dual blood glucose and blood ketone diagnostic test m 1) type 1 diabetes; or 2) permanent neonatal diabetes; or 3) undergone a pancreatectomy; or 4) cystic fibrosis-related diabetes; or 5) metabolic disease or epilepsy under the care of a participation must be endorsed accordingly. Only 1 	C TEST METER – Su neter is subsidised for paediatrician, neurolog meter per patient will	a pati gist or be su	ient who has metabolic sj bsidised (no	: pecialist. repeat prescriptions). Fc
 diagnostic test strips	funded CareSens meter. Meter with 50 lancets, a lancing device and 10 blood glucos	Se .			
 BLOOD GLUCOSE DIAGNOSTIC TEST METER – Subsidy by endorsement Maximum of 1 pack per prescription Up to 1 pack available on a PSO A diagnostic blood glucose test meter is subsidised for a patient who: is receiving insulin or sulphonylurea therapy; or is pregnant with diabetes; or is on home TPN at risk of hypoglycaemia or hyperglycaemia; or has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome. The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have: type 1 diabetes; or undergone a pancreatectomy; or cystic fibrosis-related diabetes. For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for funded CareSens meter. Meter with 50 lancets, a lancing device and 10 diagnostic test strips			1 OP	✓ <u>c</u>	areSens Dual
 a) Maximum of 1 pack per prescription b) Up to 1 pack available on a PSO c) A diagnostic blood glucose test meter is subsidised for a patient who: is receiving insulin or sulphonylurea therapy; or is pregnant with diabetes; or is on home TPN at risk of hypoglycaemia or hyperglycaemia; or has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome. The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have: type 1 diabetes; or permanent neonatal diabetes; or undergone a pancreatectomy; or cystic fibrosis-related diabetes. For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for funded CareSens meter. Meter with 50 lancets, a lancing device and 10 diagnostic test strips	Blood Glucose Testing				
	 a) Maximum of 1 pack per prescription b) Up to 1 pack available on a PSO c) A diagnostic blood glucose test meter is subsidised for a is receiving insulin or sulphonylurea therapy; or is pregnant with diabetes; or is on home TPN at risk of hypoglycaemia or hyperg has a genetic or an acquired disorder of glucose he syndrome. The prescription must be endorsed accordingly. Only or prescriptions). Patients already using the CareSens N F meter, unless they have: type 1 diabetes; or permanent neonatal diabetes; or undergone a pancreatectomy; or cystic fibrosis-related diabetes. For the avoidance of doubt patients who have previously funded CareSens meter. 	a patient who: glycaemia; or omeostasis, excluding ne CareSens meter pe POP meter and CareS y received a funded m	er pati ens N eter, c	ent will be su I meter are n other than Ca	ubsidised (no repeat lot eligible for a new areSens, are eligible for a lareSens N
	Note: Only 1 meter available per PSO	20.00			

	Subsidy (Manufacturer's Pr \$	rice) Sub Per	Fully sidised	Brand or Generic Manufacturer
BLOOD GLUCOSE DIAGNOSTIC TEST STRIP – Up to 50 test	available on a PS	60		
The number of test strips available on a prescription is restri	icted to 50 unless:			
1) Prescribed for a patient on insulin or a sulphonylurea a				
prescription as endorsed where there exists a record of	1 1 0			,
 Prescribed on the same prescription as insulin or a sul endorsed; or 	phonylurea in whi	ch case the p	rescriptio	on is deemed to be
Prescribed for a pregnant woman with diabetes and er				
4) Prescribed for a patient on home TPN at risk of hypog				
 Prescribed for a patient with a genetic or an acquired of 2 diabetes and metabolic syndrome and endorsed acc 		e nomeostasis	s excludi	ng type T or type
Test strips	10.56	50 test OP		areSens N areSens PRO
BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)				
The number of test strips available on a prescription is restri	cted to 50 unless:			
1) Prescribed for a patient on insulin or a sulphonylurea a				
prescription as endorsed where there exists a record of 2) Prescribed on the same prescription as insulin or a sul				
endorsed; or	ipitoriyiurea in will	ch case the p	rescription	
 Prescribed for a pregnant woman with diabetes and er 	ndorsed according	ly; or		
4) Prescribed for a patient on home TPN at risk of hypog				
 Prescribed for a patient with a genetic or an acquired of 2 diabetes and metabolic syndrome and endorsed acc 	0	e homeostasis	s excludi	ng type 1 or type
Blood glucose test strips	26.20	50 test OP	✓ Se	ensoCard
Insulin Syringes and Needles				
Subsidy is available for disposable insulin syringes, needles, and he supply of insulin or when prescribed for an insulin patient and annotate the prescription as endorsed where there exists a reco	d the prescription	is endorsed a	ccording	
NSULIN PEN NEEDLES – Maximum of 100 dev per prescription				
 ✗ 29 g × 12.7 mm ⅔ 31 g × 5 mm 		100 100	_	-D Micro-Fine -D Micro-Fine

*	31 g × 5 mm	11./5
*	31 g × 6 mm	9.50
	31 g × 8 mm	
	32 g × 4 mm	
	0	

✓	B-D Micro-Fine
-	B-D Micro-Fine
-	' Berpu

✓ B-D Micro-Fine

100

100 100

✓ B-D Micro-Fine

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE	E – Maximum of 100	dev p	per prescri	ption
*	Syringe 0.3 ml with 29 g × 12.7 mm needle		100	· 🗸	B-D Ultra Fine
		1.30	10		
		(1.99)			B-D Ultra Fine
*	Syringe 0.3 ml with 31 g × 8 mm needle	13.00	100	✓	B-D Ultra Fine II
		1.30	10		
		(1.99)			B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g × 12.7 mm needle	13.00	100	✓	B-D Ultra Fine
		1.30	10		
		(1.99)			B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	13.00	100	✓	B-D Ultra Fine II
		1.30	10		
		(1.99)			B-D Ultra Fine II
*	Syringe 1 ml with 29 g × 12.7 mm needle	13.00	100	✓	B-D Ultra Fine
		1.30	10		
		(1.99)			B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.00	100	✓	B-D Ultra Fine II
		1.30	10		
		(1.99)			B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription
- c) Maximum of 1 insulin pump per patient each four year period.

Min basal rate 0.025 U/h		1	 MiniMed 640G
Min basal rate 0.1 U/h	4,500.00	1	 Tandem t:slim X2

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:

14

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	1	Manufacturer	

continued...

- 4.1 Applicant is a relevant specialist; or
- 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and

8 Either:

- 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
- 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Either:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufact

ic Manufacturer

Insulin Pump Consumables

⇒SA1604 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes: and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Fither:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the followina:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application - (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person): and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist: or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 vears for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events: and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 Applicant is a relevant specialist; or

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	1	Manufacturer	

continued...

3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

18

1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less

	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully idised	Brand or Generic Manufacturer
continued than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mm 3 The patient has not had an increase in severe unexplaine 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within the	d hypoglycaemic epis			ne; and
INSULIN PUMP ACCESSORIES – Special Authority see SA160 a) Maximum of 1 cap per prescription b) Only on a prescription c) Maximum of 1 prescription per 180 days. Battery cap		l pharmacy 1		nimas Battery Cap
 INSULIN PUMP CARTRIDGE – Special Authority see SA1604 c a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per Cartridge 300 U, t:lock × 10 	year.	narmacy 1 OP	✓ т	andem Cartridge

	Subsidy (Manufacturer's Pi \$	rice) Sub Per	Fully Brand or osidised Generic Manufacturer
JLIN PUMP INFUSION SET (STEEL CANNULA) – Special A a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year.	uthority see SA	1604 on page	17 – Retail pharmacy
10 mt steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T MMT-884
10 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-883
10 mm steel needle; 29 G; manual insertion; 80 cm tubing x 10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T MMT-886
10 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles; luer lock		1 OP	✓ Sure-T MMT-885
6 mm steel cannula; straight insertion; 60 cm grey line × 10 with 10 needles		1 OP	 Contact-D
6 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T MMT-864
6 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-863
6 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T MMT-866
6 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-865
8 mm steel cannula; straight insertion; 110 cm grey line × 10 with 10 needles		1 OP	✓ Contact-D
8 mm steel cannula; straight insertion; 60 cm grey line × 10 with 10 needles		1 OP	 Contact-D
8 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T MMT-874
8 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-873
8 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T MMT-876
8 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-875

(Contact-D 6 mm steel cannula; straight insertion; 60 cm grey line × 10 with 10 needles to be delisted 1 October 2019) (Contact-D 8 mm steel cannula; straight insertion; 110 cm grey line × 10 with 10 needles to be delisted 1 October 2019) (Contact-D 8 mm steel cannula; straight insertion; 60 cm grey line × 10 with 10 needles to be delisted 1 October 2019)

	Subsidy (Manufacturer's Pr \$	ice) Sub Per	Fully sidised	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT	INSERTION) -	Special Author	ority see	SA1604 on page 17 -
Retail pharmacy			•	
 Maximum of 3 sets per prescription 				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm steel cannula; straight insertion; 60 cm line \times 10 with			<i>.</i> –	
10 needles	130.00	1 OP	✓ Tr	uSteel
6 mm steel cannula; straight insertion; 81 cm line × 10 with			<i>.</i> -	. .
10 needles	130.00	1 OP	✓ Tr	uSteel
8 mm steel cannula; straight insertion; 60 cm line \times 10 with			<i>.</i> -	. .
10 needles	130.00	1 OP	✓ Tr	uSteel
8 mm steel cannula; straight insertion; 81 cm line × 10 with	100.00	4.05	/ -	o
10 needles		1 OP		uSteel
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	SERTION WITH	I INSERTION	DEVICE) – Special Authority see
SA1604 on page 17 – Retail pharmacy				
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
13 mm teflon cannula; angle insertion; insertion device; 110 c		1.00	In	set 30
grey line × 10 with 10 needles		1 OP	♥ Ins	set 30
13 mm teflon cannula; angle insertion; insertion device; 60 cm		1.00	. In	
grey line × 10 with 10 needles		1 OP	♥ Ins	set 30
13 mm teflon cannula; angle insertion; insertion device; 110 c		1.00		4.0.4
line × 10 with 10 needles		1 OP	♥ AL	itoSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cm line × 10 with 10 needles		1 OP	۸.	utoSoft 30
(Inset 30 13 mm teflon cannula; angle insertion; insertion device; 1 2019)	i i o cili grey line	x IU WITH IU	needles	o de delisted i October
2013/				

(Inset 30 13 mm teflon cannula; angle insertion; insertion device; 60 cm grey line \times 10 with 10 needles to be delisted 1 October 2019)

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
SULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE I	NSERTION) – Speci	al Au	hority see	SA1604 on page 17 -
etail pharmacy				
a) Maximum of 3 sets per prescriptionb) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
13 mm teflon cannula; angle insertion; 120 cm line \times 10 with				
10 needles	130.00	1 OP	1	Paradigm Silhouette MMT-382
13 mm teflon cannula; angle insertion; 45 cm line × 10 with				
10 needles	130.00	1 OP	1	Paradigm Silhouette MMT-368
13 mm teflon cannula; angle insertion; 60 cm line × 10 with				
10 needles	130.00	1 OP	1	Paradigm Silhouette MMT-381
13 mm teflon cannula; angle insertion; 80 cm line \times 10 with		_		
10 needles		1 OP	-	Paradigm Silhouette MMT-383
17 mm teflon cannula; angle insertion; 110 cm line × 10 with				-
10 needles		1 OP	•	Paradigm Silhouette MMT-377
17 mm teflon cannula; angle insertion; 110 cm line × 10 with	100.00			
10 needles; luer lock		1 OP	~	Silhouette MMT-371
17 mm teflon cannula; angle insertion; 60 cm line × 10 with 10 needles	130.00	1 OP	1	Paradigm Silhouette
17 mm toffen computer angle incention, co and the statistic				MMT-378
17 mm teflon cannula; angle insertion; 60 cm line × 10 with 10 needles; luer lock	130.00	1 OP	1	Silhouette MMT-373
17 mm teflon cannula; angle insertion; 80 cm line × 10 with		I UF	•	Simouelle wiwi -3/3
10 needles	130.00	1 OP	1	Paradigm Silhouette MMT-384

	Subsidy (Manufacturer's Prio	ce) Sub	Fully	Brand or Generic
	\$	Per	✓	Manufacturer
SULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH	HT INSERTION W	TH INSERT	ION DE	VICE) – Special Authorit
e SA1604 on page 17 – Retail pharmacy				, ,
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; insertion device;				
110 cm grey line × 10 with 10 needles	140.00	1 OP	🖌 In	iset II
6 mm teflon cannula; straight insertion; insertion device; 45 c	m			
blue tubing × 10 with 10 needles		1 OP	🗸 P	aradigm Mio
0				MMT-941
6 mm teflon cannula; straight insertion; insertion device; 45 c	m			
pink tubing × 10 with 10 needles		1 OP	✓ P	aradigm Mio
P				MMT-921
6 mm teflon cannula; straight insertion; insertion device; 60 c	m			
blue tubing × 10 with 10 needles		1 OP	✓ P	aradigm Mio
, , , , , , , , , , , , , , , , , , ,		-		MMT-943
6 mm teflon cannula; straight insertion; insertion device; 60 c	em			
grey line × 10 with 10 needles		1 OP	🗸 In	iset II
6 mm teflon cannula; straight insertion; insertion device; 60 c				
pink tubing × 10 with 10 needles		1 OP	✓ P	aradigm Mio
		1.01		MMT-923
6 mm teflon cannula; straight insertion; insertion device; 80 c	m			
blue tubing × 10 with 10 needles		1 OP	✓ P	aradigm Mio
		1 01		MMT-945
6 mm teflon cannula; straight insertion; insertion device; 80 c	m			
clear tubing × 10 with 10 needles		1 OP	✓ P	aradigm Mio
			-	MMT-965
6 mm teflon cannula; straight insertion; insertion device; 80 c	m			
pink tubing × 10 with 10 needles		1 OP	✓ P	aradigm Mio
			-	MMT-925
9 mm teflon cannula; straight insertion; insertion device;				
110 cm grey line × 10 with 10 needles		1 OP	🗸 In	Í Inset II
9 mm teflon cannula; straight insertion; insertion device; 60 c				
grey line × 10 with 10 needles		1 OP	🖌 In	iset II
9 mm teflon cannula; straight insertion; insertion device; 80 c				
clear tubing × 10 with 10 needles		1 OP	V P	aradigm Mio
			-	MMT-975
6 mm teflon cannula; straight insertion; insertion device;				
110 cm line × 10 with 10 needles		1 OP	🗸 A	utoSoft 90
6 mm teflon cannula; straight insertion; insertion device; 60 c				
line × 10 with 10 needles		1 OP	۸ 🗸	utoSoft 90
9 mm teflon cannula; straight insertion; insertion device;			- 1	
110 cm line × 10 with 10 needles	140.00	1 OP	۸ 🗸	utoSoft 90
			- 4	

	Subsidy (Manufacturer's P \$	rice) Sub Per	Fully osidised	Brand or Generic Manufacturer
9 mm teflon cannula; straight insertion; insertion device; 60				
line × 10 with 10 needles		1 OP		itoSoft 90
set II 6 mm teflon cannula; straight insertion; insertion device; 19)				
set II 6 mm teflon cannula; straight insertion; insertion device; 19)	0 7			
set II 9 mm teflon cannula; straight insertion; insertion device; 19)	110 cm grey line	× 10 with 10	needles to	be delisted 1 October
set II 9 mm teflon cannula; straight insertion; insertion device; 19)	60 cm grey line ×	: 10 with 10 n	eedles to	be delisted 1 October
SULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIG tail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription	HT INSERTION)	 Special Aut 	thority see	SA1604 on page 17 -
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; 110 cm tubing x 10 m 10 needles		1 OP		radigm Quick-Set MMT-398
6 mm teflon cannula; straight insertion; 110 cm tubing × 10 10 needles; luer lock	130.00	1 OP	🗸 Qı	iick-Set MMT-391
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 w 10 needles		1 OP		radigm Quick-Set MMT-399
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 w 10 needles; luer lock		1 OP		iick-Set MMT-393
6 mm teflon cannula; straight insertion; 80 cm tubing × 10 w 10 needles		1 OP		radigm Quick-Set
O man before a second a stariabling of the door second bins and o				MMT-387
9 mm teflon cannula; straight insertion; 106 cm tubing x 10 10 needles		1 OP		radigm Quick-Set MMT-396
9 mm teflon cannula; straight insertion; 110 cm tubing × 10 10 needles; luer lock		1 OP		lick-Set MMT-390
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 w	ith			
10 needles		1 OP		radigm Quick-Set MMT-397
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 w 10 needles; luer lock		1 OP	🗸 Qı	ick-Set MMT-392
9 mm teflon cannula; straight insertion; 80 cm tubing × 10 w				
10 needles		1 OP		radigm Quick-Set MMT-386

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
NSULIN PUMP RESERVOIR - Special Authority see SA1604 on	page 17 – Retail p	harma	су	
a) Maximum of 3 sets per prescription				
 b) Only on a prescription 				
c) Maximum of 13 packs of reservoir sets will be funded per y	/ear.			
10 × luer lock conversion cartridges 1.8 ml for Paradigm pump		1 OP		ADR Cartridge 1.8
Cartridge 200 U, luer lock × 10	50.00	1 OP	✓ .	Animas Cartridge
Cartridge for 5 and 7 series pump; 1.8 ml × 10	50.00	1 OP	✓	Paradigm
				1.8 Reservoir
Cartridge for 7 series pump; 3.0 ml × 10		1 OP	✓	Paradigm
				3.0 Reservoir
Syringe and cartridge for 50X pump, 3.0 ml × 10		1 OP		50X 3.0 Reservoir
Digestives Including Enzymes				
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase				
10,000 Ph Eur U, total protease 600 Ph Eur U)	34 93	100	1	Creon 10000
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase,		100		
1,250 U protease))		100	~	Panzytrat
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase		100	•	i anzyriar
25,000 Ph Eur U, total protease 1,000 Ph Eur U)	04.29	100	1	Creon 25000
			•	<u>CIEUII 23000</u>
RSODEOXYCHOLIC ACID – Special Authority see SA1739 bek				
Cap 250 mg		100	v	Ursosan
SA1739 Special Authority for Subsidy				
itial application — (Alagille syndrome or progressive familia	I intrahepatic cho	lestasi	is) from ar	ny relevant practitioner.
pprovals valid without further renewal unless notified for applicati				

Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Laxatives

Bulk-forming Agents

SPAGHULA (PSYLLIUM) HUSK – Only on a prescription * Powder for oral soln	6.05	500 g OP	✓ Konsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS			
* Dry	6.02	500 g OP	
	(17.32)	•	Normacol Plus
	2.41	200 g OP	
	(8.72)	•	Normacol Plus

Faecal Softeners

DOCUSATE SODIUM – Only on a prescription		
* Tab 50 mg2.31	100	 Coloxyl
* Tab 120 mg	100	 Coloxyl
DOCUSATE SODIUM WITH SENNOSIDES * Tab 50 mg with sennosides 8 mg	200	✓ Laxsol
POLOXAMER – Only on a prescription Not funded for use in the ear.		
* Oral drops 10%	30 ml OP	 Coloxyl

Opioid Receptor Antagonists - Peripheral

METHYLNALTREXONE BROMIDE - Special Authority see	SA1691 on the next pa	<mark>ge</mark> – Retai	l pharmacy
Inj 12 mg per 0.6 ml vial		1	 Relistor
	246.00	7	 Relistor

(Subsidy Manufacturer's Price \$) (Per	Fully Subsidised	Brand or Generic Manufacturer
SA1691 Special Authority for Subsidy initial application — (Opioid induced constipation) from any re unless notified for applications meeting the following criteria: Both:	levant practitioner	. Appro	ovals valid v	without further renewal
 The patient is receiving palliative care; and Either: 2.1 Oral and rectal treatments for opioid induced constip 				
2.2 Oral and rectal treatments for opioid induced constip	ation are unable to	o be tol	erated.	
Osmotic Laxatives				
GLYCEROL * Suppos 3.6 g – Only on a prescription	9.25	20	✓ <u>P</u>	SM
_ACTULOSE – Only on a prescription 券 Oral liq 10 g per 15 ml	3.18	500 ml	· ✓L	aevolac
ACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BIC/	ARBONATE AND	SODIU	M CHLORI	DE
Powder for oral soln 13.125 g with potassium chloride 46.6 mg sodium bicarbonate 178.5 mg and sodium chloride 350.7 r		30	✓ <u>N</u>	lolaxole
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%	2.50	1	✔ F	leet Phosphate Enema
ODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE - Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml,	 Only on a prescr 	iption		
5 ml		50	✓ N	licolette
Stimulant Laxatives				
BISACODYL - Only on a prescription				
₭ Tab 5 mg ₭ Suppos 10 mg		200 10		<u>ax-Tab</u> ax-Suppositories
		10	• ⊑	ax-suppositories
ENNA – Only on a prescription K Tab, standardised		100		
·,	(6.84)		S	enokot
	0.43	20		
	(1.72)		S	senokot
Metabolic Disorder Agents				
ALGLUCOSIDASE ALFA – Special Authority see SA1622 below –	Retail pharmacy			
Inj 50 mg vial		1	🗸 N	lyozyme
SA1622 Special Authority for Subsidy				
nitial application only from a metabolic physician. Approvals vali All of the following:	d for 12 months fo	r applic	ations mee	ting the following criter

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

villus biopsies and/or cultured amniotic cells; or

- 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
- 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
- 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

BETAINE – Special Authority see SA1727 below – Retail pharmacy

➡SA1727 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and

3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation. **Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE - Special Authority see SA1593 below - Retail pharmacy

Inj 1 mg per ml, 5 ml vial2,234.00	1	 Naglazyme
------------------------------------	---	-------------------------------

⇒SA1593 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:

28

2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either

	Subsidy	Fu	lly Brand or	
(Ma	nufacturer's Price)	Subsidise	ed Generic	
	\$	Per	 Manufacturer 	

continued...

- enzyme activity assay in leukocytes or skin fibroblasts; or
- 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

IDURSULFASE - Special Authority see SA1623 below - Retail pharmacy

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

LARONIDASE – Special Authority see SA1695 below – Retail pharmacy

Inj 100 0 per mi, 5 mi viai 1,335.16 1 Aldurazyme

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

SAPROPTERIN DIHYDROCHLORIDE – Special Authority see SA	A1757 on the next	page - Retai	pharmacy
Tab soluble 100 mg	1,452.70	30 OP	 Kuvan

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	 ✓ 	Manufacturer

⇒SA1757 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

Approvals valid for 12 months for applications meeting the following criteri All of the following:

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE - Special Authority see SA1599 below - Retail pharmacy

Soln 100 mg per mlCBS 100 ml 🖌 Amzoate \$29

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE - Special Authority see SA1598 below - Retail pharmacy

⇒SA1598 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

30

IMIGLUCERASE - Special Authority see SA0473 of	n the next page – Retail pharmacy
Inj 40 iu per ml, 400 iu vial	2,144.00 1
(Cerezyme Inj 40 iu per ml, 400 iu vial to be delisted	1 September 2019)

✓ Cerezvme

(Subsidy (Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer
SA0473 Special Authority for Subsidy Special Authority approved by the Gaucher Treatment Panel Notes: Subject to a budgetary cap. Applications will be considered Application details may be obtained from PHARMAC's website http			•	ailability.
The Co-ordinator, Gaucher Treatment PanelPhone: (04) 460PHARMAC, PO Box 10 254Facsimile: (04)WellingtonEmail: gaucher		<u>vt.nz</u>		
TALIGLUCERASE ALFA – Special Authority see SA1734 below – Inj 200 unit vial	1,072.00	1 rmac.g	_	<u>lelyso</u>
The Co-ordinator, Gaucher Treatment PanelPhone: 04 460PHARMAC PO Box 10 254Facsimile: 04 9WellingtonEmail: gaucher		<u>vt.nz</u>		

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Access Criteria

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2) Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and
- Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, serum glucosylsphingosine, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
- Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia, thrombocytopenia; at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or massive symptomatic splenomegaly; or
 - 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g. hips or shoulder); spontaneous fractures or vertebral collapse; chronic bone pain not controlled by other pharmaceuticals; or
 - 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
 - 5) Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	lbsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1) Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated; and
- 2) Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3) Radiological (MRI) signs of bone activity performed at one year and two years since initiation of treatment begins, and two to three yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4) Serum glucosylsphingosine levels taken at least 6 to 12 monthly show a decrease compared with baseline; and
- Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 7) Patient is compliant with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- Supporting clinical information including test reports, MRI whole body STIR, serum glucosylsphingosine, haematological data, and other relevant investigations are submitted to the Gaucher Panel for assessment as required.

Mouth and Throat

Agents Used in Mouth Ulceration

SENZYDAMINE HYDROCHLORIDE			
Soln 0.15% – Higher subsidy of \$20.31 per 500 ml with			
Endorsement	9.00	500 ml	
	(20.31)		Difflam
Additional subsidy by endorsement for a patient who has prescription is endorsed accordingly.	s oral mucositis	as a result of tre	eatment for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			
Paste		56 g OP	 Stomahesive
	4.55	15 g OP	
	(7.90)		Orabase
	1.52	5 g OP	
	(3.60)	- 3	Orabase
Powder	· · ·	28 g OP	
	(10.95)		Stomahesive
CHLORHEXIDINE GLUCONATE	· · · ·		
Mouthwash 0.2%	2 57	200 ml OP	✓ healthE
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE		200 01	
	0.00		
Adhesive gel 8.7% with cetalkonium chloride 0.01%		15 g OP	Deviale
	(6.00)		Bonjela
FRIAMCINOLONE ACETONIDE			
Paste 0.1%	5.33	5 g OP	 Kenalog in Orabase
Oropharyngeal Anti-infectives			
AMPHOTERICIN B			
Lozenges 10 mg	5.86	20	✓ Fungilin
5 5		20	• i uligiili
MICONAZOLE			
Oral gel 20 mg per g	4	40 g OP	 Decozol

32

	ALIMENTA	RY TRACT	AND	D METABOLISM
۸)	Subsidy /anufacturer's Pr \$	ice) Subsi Per	Fully dised	Brand or Generic Manufacturer
VYSTATIN Oral liq 100,000 u per ml	1.95	24 ml OP	✓ <u>I</u>	lilstat
Other Oral Agents				
For folinic mouthwash, pilocarpine oral liquid or saliva substitute for	nula refer Star	ndard Formulae	e, page	e 227
Soln 3% (10 vol) – Maximum of 200 ml per prescription	1.40	100 ml	√ F	Pharmacy Health
THYMOL GLYCERIN 卷 Compound, BPC	9.15	500 ml	✓ <u>F</u>	PSM
Vitamins				
Vitamin A				
VITAMIN A WITH VITAMINS D AND C * Soln 1000 u with Vitamin D 400 u and ascorbic acid 30 mg per 10 drops	4.50	10 ml OP	• 1	/itadol C
(Vitadol C Soln 1000 u with Vitamin D 400 u and ascorbic acid 30 m	ng per 10 drops	s to be delisted	1 Aug	ust 2019)
Vitamin B				
 HYDROXOCOBALAMIN Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a PSO PYRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose b) Only on a prescription 	1.89	3	✓ <u>I</u>	<u>leo-B12</u>
 Tab 25 mg – No patient co-payment payable Tab 50 mg 	2.70	90 500		/itamin B6 25 Apo-Pyridoxine
THIAMINE HYDROCHLORIDE – Only on a prescription				
* Tab 50 mg VITAMIN B COMPLEX		100	_	<u>lax Health</u>
* Tab, strong, BPC	7.15	500	✓Ē	<u> Splex</u>
Vitamin C				
ASCORBIC ACID a) No more than 100 mg per dose b) Only on a prescription				
* Tab 100 mg	8.10	500	✓ <u>(</u>	<u>Svite</u>
Vitamin D				
ALFACALCIDOL * Cap 0.25 mcg * Cap 1 mcg * Oral drops 2 mcg per ml	87.98	100 100 20 ml OP	✓ (Dne-Alpha Dne-Alpha Dne-Alpha
CALCITRIOL * Cap 0.25 mcg * Cap 0.5 mcg		100 100	_	Calcitriol-AFT Calcitriol-AFT

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Pr \$		Fully Brand or dised Generic ✓ Manufacturer
COLECALCIFEROL * Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescrip * Oral liq 188 mcg per ml (7,500 iu per ml)		12 4.8 ml OP	✓ <u>Vit.D3</u> ✓ Puria
Multivitamin Preparations			
MULTIVITAMIN RENAL Special Authority see SA1546 below * Cap		/ 30	✓ Clinicians Renal Vit
SA1546 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals va he following criteria: Either:	id without further r	enewal unless	notified for applications meeting
 The patient has chronic kidney disease and is receiving a The patient has chronic kidney disease grade 5, defined 15 ml/min/1.73 m² body surface area (BSA). 			
MULTIVITAMINS – Special Authority see SA1036 below – Retain the second		200 g OP	✓ Paediatric Seravit
 SA1036 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals va nborn errors of metabolism. Renewal from any relevant practitioner. Approvals valid withou approval for multivitamins. 			
/ITAMINS ₭ Tab (BPC cap strength)		1,000	✓ Mvite
Cap (fat soluble vitamins A, D, E, K) – Special Authority se SA1720 below – Retail pharmacy	е	60	✓ Vitabdeck
 SA1720 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals val he following criteria: Any of the following: 1 Patient has cystic fibrosis with pancreatic insufficiency; o 2 Patient is an infant or child with liver disease or short gut 3 Patient has severe malabsorption syndrome. 	r	renewal unless	notified for applications meeting
Minerals			
Calcium			
CALCIUM CARBONATE ≰ Tab eff 1.75 g (1 g elemental) ≰ Tab 1.25 g (500 mg elemental) Calsource Tab eff 1.75 g (1 g elemental) to be delisted 1 Septe	7.52	10 250	 ✓ Calsource ✓ <u>Arrow-Calcium</u>
CALCIUM GLUCONATE		40	
Inj 10%, 10 ml ampoule		10 20	 ✓ Hospira ✓ Max Health S29
Hospira Inj 10%, 10 ml ampoule to be delisted 1 July 2019)	2		
Fluoride			
ODIUM FLUORIDE ₭ Tab 1.1 mg (0.5 mg elemental)	5.75	100	✔ PSM
34 ✓ fully subsidised	S29 Unappi	roved medicine su	upplied under Section 29

34 Sole Subsidised Supply

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	Brand or Generic Manufacturer
lodine				
POTASSIUM IODATE * Tab 253 mcg (150 mcg elemental iodine)	4.69	90	✓ <u>Ne</u>	euroTabs
Iron				
FERRIC CARBOXYMALTOSE – Special Authority see SA1675 be Inj 50 mg per ml, 10 ml		acy 1	✔ Fe	erinject

⇒SA1675 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any medical practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and

- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any medical practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and

2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

FERROUS FUMARATE

* Tab 200 mg (65 mg elemental)	.3.09	100	✓ Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID * Tab 310 mg (100 mg elemental) with folic acid 350 mcg	.4.68	60	✓ Ferro-F-Tabs
FERROUS SULPHATE * Tab long-acting 325 mg (105 mg elemental) * Oral liq 30 mg (6 mg elemental) per 1 ml		30 500 ml	 ✓ Ferrograd ✓ Ferodan

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
IRON POLYMALTOSE * Inj 50 mg per ml, 2 ml ampoule (Ferrum H Inj 50 mg per ml, 2 ml ampoule to be delisted 1 July 2	34.50	5	-	errum H errosig
Magnesium				
For magnesium hydroxide mixture refer Standard Formulae, pag MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule		10	✓ <u>D</u> ✓ D	<u>BL</u> BL S29 529
Zinc				
ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)	11.00	100	✓ Z	incaps

Subsidised

Per

Fully

Subsidy (Manufacturer's Price) \$ Brand or Generic Manufacturer

Antianaemics

Hypoplastic and Haemolytic

➡SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or

3.2 Both:

- 3.2.1 Patient has diabetes mellitus; and
- 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
- 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy. Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following

criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
EPOETIN ALFA - Special Authority see SA1775 on the previous	s page – Retail pharr	nacy		
Wastage claimable				
Inj 1,000 iu in 0.5 ml, syringe		6	~	Binocrit
Binocrit to be Sole Supply on 1 July 2019				
Inj 2,000 iu in 1 ml, syringe		6	v	Binocrit
Binocrit to be Sole Supply on 1 July 2019				
Inj 3,000 iu in 0.3 ml, syringe	150.00	6	v	Binocrit
Binocrit to be Sole Supply on 1 July 2019				
Inj 4,000 iu in 0.4 ml, syringe	96.50	6	~	Binocrit
Binocrit to be Sole Supply on 1 July 2019				
Inj 5,000 iu in 0.5 ml, syringe		6	~	Binocrit
Binocrit to be Sole Supply on 1 July 2019				
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	~	Binocrit
Binocrit to be Sole Supply on 1 July 2019				
Inj 8,000 iu in 0.8 ml, syringe	175.00	6	~	Binocrit
Binocrit to be Sole Supply on 1 July 2019				
Inj 10,000 iu in 1 ml, syringe		6	~	Binocrit
Binocrit to be Sole Supply on 1 July 2019				
Inj 40,000 iu in 1 ml, syringe		1	-	Binocrit
Binocrit to be Sole Supply on 1 July 2019				
Megaloblastic				
FOLIC ACID				
* Tab 0.8 mg	21.84	1,000) 🗸	Apo-Folic Acid
* Tab 5 mg		500	✓	Apo-Folic Acid
Oral lig 50 mcg per ml		5 ml (DP 🗸	Biomed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Inj 250 iu vial	612.50	. 1	Alprolix
Inj 500 iu vial		1	 Alprolix
Inj 1,000 iu vial		1	 Alprolix
Inj 2,000 iu vial	4,900.00	1	 Alprolix
Inj 3,000 iu vial	7,350.00	1	 Alprolix
ELTROMBOPAG - Special Authority see SA1743 below - Re	tail pharmacy		
Wastage claimable			
Tab 25 mg		28	Revolade
Tab 50 mg		28	Revolade

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

38

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	sidised	Generic Manufacturer
PTACOG ALFA [RECOMBINANT FACTOR VIIA] – [Xpharm	1			
For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group.		emophilia	a Treat	ers Group in conjunctio
Inj 1 mg syringe	1,178.30	1	-	NovoSeven RT
lnj 2 mg syringe		1	✓	NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓	NovoSeven RT
Inj 8 mg syringe	9,426.40	1	~	NovoSeven RT
CTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xph	arm]			
For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group.	it is managed by the Hae	emophilia	a Treat	ers Group in conjunctio
Inj 500 U		1	1	FEIBA NF
Inj 1,000 U		1	1	FEIBA NF
Inj 2,500 U		1	✓	FEIBA NF
ROCTOCOG ALFA [RECOMBINANT FACTOR VIII] – [Xpl	harml			
For patients with haemophilia. Access to funded treatmen		emophilia	a Treat	ers Group in coniunctio
with the National Haemophilia Management Group.	,			· · · · · · · · · · · ·
Inj 250 iu prefilled syringe	210.00	1	✓	Xyntha
Inj 500 iu prefilled syringe		1		Xyntha
Inj 1,000 iu prefilled syringe		1	✓	Xyntha
Inj 2,000 iu prefilled syringe	1,680.00	1		Xyntha
Inj 3,000 iu prefilled syringe	2,520.00	1	~	Xyntha
NACOG ALFA [RECOMBINANT FACTOR IX] – [Xpharm]				
For patients with haemophilia, whose funded treatment is r	managed by the Haemo	philia Tre	eaters	Group in conjunction wi
the National Haemophilia Management Group.				
Inj 250 iu vial		1		BeneFIX
Inj 500 iu vial		1		BeneFIX
Inj 1,000 iu vial		1		BeneFIX
Inj 2,000 iu vial		1		BeneFIX
Inj 3,000 iu vial		1	~	BeneFIX
eneFIX Inj 250 iu vial to be delisted 1 November 2019)				
eneFIX Inj 500 iu vial to be delisted 1 November 2019)				
eneFIX Inj 1,000 iu vial to be delisted 1 November 2019)				
naElV Ini 2,000 in vial to be delicted 1 Nevember 2010)				
neFIX Inj 3,000 iu vial to be delisted 1 November 2019)	,			
neFIX Inj 3,000 iu vial to be delisted 1 November 2019) NACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xphar			. T	ere Orenne in een innetie
neFIX Inj 3,000 iu vial to be delisted 1 November 2019) NACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xphar For patients with haemophilia. Access to funded treatmen		emophilia	a Treat	ers Group in conjunctio
neFIX Inj 3,000 iu vial to be delisted 1 November 2019) NACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xphar For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group.	t is managed by the Hae		_	
eneFIX Inj 3,000 iu vial to be delisted 1 November 2019) NACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xphar For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group. Inj 500 iu vial	it is managed by the Hae	1	1	RIXUBIS
eneFIX Inf 3,000 iu vial to be delisted 1 November 2019) NACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xphar For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group. Inj 500 iu vial Inj 1,000 iu vial	it is managed by the Hae 435.00 	1	1	RIXUBIS RIXUBIS
aneFIX Inf 3,000 iu vial to be delisted 1 November 2019) NACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xphar For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group. Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial	it is managed by the Hae 435.00 	1	1 1 1	RIXUBIS RIXUBIS RIXUBIS
eneFIX Inj 3,000 iu vial to be delisted 1 November 2019) NACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xphar For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group. Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial Inj 3,000 iu vial	t is managed by the Hae 435.00 	1 1 1	1 1 1	RIXUBIS RIXUBIS
eneFIX Inj 3,000 iu vial to be delisted 1 November 2019) NACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xphar For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group. Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial Inj 3,000 iu vial TOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE)	t is managed by the Hae 	1 1 1 1	\$ \$ \$ \$ \$	RIXUBIS RIXUBIS RIXUBIS RIXUBIS
eneFIX Inj 3,000 iu vial to be delisted 1 November 2019) NACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xphar For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group. Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial Inj 3,000 iu vial TOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) For patients with haemophilia. Access to funded treatmen	t is managed by the Hae 	1 1 1 1	\$ \$ \$ \$ \$	RIXUBIS RIXUBIS RIXUBIS RIXUBIS
eneFIX Inj 3,000 iu vial to be delisted 1 November 2019) NACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xphar For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group. Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial Inj 3,000 iu vial TOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group.	t is managed by the Had 	1 1 1 1 emophilia	✓ ✓ ✓ ▲ Treat	RIXUBIS RIXUBIS RIXUBIS RIXUBIS ers Group in conjunctio
eneFIX Inj 3,000 iu vial to be delisted 1 November 2019) NACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xphar For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group. Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial TOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group. Inj 250 iu vial	t is managed by the Had 	1 1 1 1	✓ ✓ ✓ A Treat	RIXUBIS RIXUBIS RIXUBIS RIXUBIS ers Group in conjunctio Advate
eneFIX Inj 3,000 iu vial to be delisted 1 November 2019) NACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xphar For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group. Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial TOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group. Inj 250 iu vial	t is managed by the Had 	1 1 1 1 emophilia	✓ ✓ ✓ ✓ A Treat	RIXUBIS RIXUBIS RIXUBIS RIXUBIS ers Group in conjunctio
eneFIX Inj 3,000 iu vial to be delisted 1 November 2019) NACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xphar For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group. Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial TOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group. Inj 250 iu vial	t is managed by the Had 	1 1 1 emophilia 1	a Treat	RIXUBIS RIXUBIS RIXUBIS RIXUBIS ers Group in conjunctio Advate Advate
with the National Haemophilia Management Group. Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial Inj 3,000 iu vial CTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) For patients with haemophilia. Access to funded treatmen with the National Haemophilia Management Group. Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial	t is managed by the Had 	1 1 1 emophilia 1 1	a Treat	RIXUBIS RIXUBIS RIXUBIS RIXUBIS ers Group in conjunctio Advate Advate Advate

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE For patients with haemophilia. Access to funded treatment with the National Haemophilia Management Group.		emop	hilia Treat	ers Group in conjunction
Inj 250 iu vial		1	1	Kogenate FS
Inj 500 iu vial		1	✓	Kogenate FS
Inj 1,000 iu vial	950.00	1		Kogenate FS
Inj 2,000 iu vial	'	1		Kogenate FS
Inj 3,000 iu vial	2,850.00	1	✓	Kogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII For patients with haemophilia A receiving prophylaxis treatn Treaters Group in conjunction with the National Haemophilia Inj 250 iu vial	nent. Access to funder a Management group.	d trea 1		nanaged by the Haemophili Adynovate
Ini 500 iu vial		1		Adynovate
Inj 1,000 iu vial		1		Adynovate
Inj 2,000 iu vial		1		Adynovate
SODIUM TETRADECYL SULPHATE	,			
* Inj 3% 2 ml	28.50 (73.00)	5		Fibro-vein
TRANEXAMIC ACID Tab 500 mg	20.67	100	1	<u>Cyklokapron</u>
Vitamin K				
PHYTOMENADIONE Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO		5 5		Konakion MM Konakion MM
Antithrombotic Agents				
Antiplatelet Agents				
ASPIRIN * Tab 100 mg CLOPIDOGREL	12.50	990	1	Ethics Aspirin EC
* Tab 75 mg	5.44	84	1	Arrow - Clopid
DIPYRIDAMOLE * Tab long-acting 150 mg		60	1	Pytazen SR
PRASUGREL – Special Authority see SA1201 below – Retail p	•		-	
Tab 5 mg Tab 10 mg		28		Effient
	100.00	28		Effient

➡SA1201 Special Authority for Subsidy

Initial application — (coronary angioplasty and bare metal stent) from any relevant practitioner. Approvals valid for 6 months where the patient has undergone coronary angioplasty in the previous 4 weeks and is clopidogrel-allergic*. **Initial application** — (drug eluting stent) from any relevant practitioner. Approvals valid for 12 months where the patient has

had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Initial application — (stent thromobosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal --- (coronary angioplasty and bare metal stent) from any relevant practitioner. Approvals valid for 6 months where

Subsidy (Manufacturer's Price)	Ful Subsidise	,	
 (Manulacturer s r nee) \$	Per •	Manufacturer	

continued...

the patient has undergone coronary angioplasty or had a bare metal cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Renewal — (drug eluting stent) from any relevant practitioner. Approvals valid for 12 months where had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Note: * Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

TICAGRELOR - Special Authority see SA1382 below - Retail pharmacy

⇒SA1382 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Heparin and Antagonist Preparations

DALTEPARIN SODIUM - Special Authority see SA1270 below - Retail pharmacy

Inj 2,500 iu per 0.2 ml prefilled syringe1	9.97 1	0 🗸	Fragmin
Inj 5,000 iu per 0.2 ml prefilled syringe		0 🖌	Fragmin
Inj 7,500 iu per 0.75 ml graduated syringe60	0.03 1	0 🖌	Fragmin
Inj 10,000 iu per 1 ml graduated syringe7		0 🖌	Fragmin
Inj 12,500 iu per 0.5 ml prefilled syringe	9.96 1	0 🖌	Fragmin
Inj 15,000 iu per 0.6 ml prefilled syringe120		0 🖌	Fragmin
Inj 18,000 iu per 0.72 ml prefilled syringe158	8.47 1	0 🖌	Fragmin

⇒SA1270 Special Authority for Subsidy

Initial application — (Pregnancy or Malignancy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

42

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

continued...

Renewal — (Pregnancy or Malignancy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, Acute Coronary Syndrome, cardioversion, or prior to oral anti-coagulation).

ENOXAPARIN SODIUM - Special Authority see SA1646 below - Retail pharmacy

Inj 20 mg in 0.2 ml syringe	 10	 Clexane
Inj 40 mg in 0.4 ml syringe	 10	 Clexane
Inj 60 mg in 0.6 ml syringe	10	 Clexane
Inj 80 mg in 0.8 ml syringe	10	 Clexane
Inj 100 mg in 1 ml syringe	 10	 Clexane
Inj 120 mg in 0.8 ml syringe	10	 Clexane
Inj 150 mg in 1 ml syringe	 10	 Clexane

⇒SA1646 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
 - 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
 - 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

HEPARIN SODIUM

Inj 1,000 iu per ml, 5 ml ampoule		50	 Pfizer
Inj 5,000 iu per ml, 1 ml		5	✓ Hospira
			 Pfizer
Inj 5,000 iu per ml, 5 ml ampoule		50	 Pfizer
Inj 25,000 iu per ml, 0.2 ml		5	✓ Hospira
	190.00	50	 Pfizer S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
HEPARINISED SALINE				
Inj 10 iu per ml, 5 ml	56.94	50	1	Pfizer
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg – No more than 2 cap per day		60	✓	Pradaxa
Cap 110 mg		60	✓	Pradaxa
Cap 150 mg		60	✓	Pradaxa
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day		30	1	Xarelto
Tab 15 mg - Up to 14 tab available on a PSO	77.56	28	1	Xarelto
Tab 20 mg	77.56	28	~	Xarelto
WARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	3.46	50	1	Coumadin
•	6.86	100	✓	Marevan
* Tab 2 mg	4.31	50	✓	Coumadin
* Tab 3 mg		100	✓	Marevan
* Tab 5 mg	5.93	50	~	Coumadin
	11.75	100	✓	Marevan

Blood Colony-stimulating Factors

FILGRASTIM - Special Authority see SA1259 below - Retail	pharmacy		
Inj 300 mcg per 0.5 ml prefilled syringe		10	 Nivestim
	48.11	5	
	(270.00)		Zarzio
Nivestim to be Sole Supply on 1 August 2019			
Inj 480 mcg per 0.5 ml prefilled syringe		10	 Nivestim
	80.75	5	
	(432.00)		Zarzio

Nivestim to be Sole Supply on 1 August 2019 (Zarzio Inj 300 mcg per 0.5 ml prefilled syringe to be delisted 1 August 2019) (Zarzio Inj 480 mcg per 0.5 ml prefilled syringe to be delisted 1 August 2019)

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5×10^9 /L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM – Special Authority see SA1384 on the next page – Retail pharmacy

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer
--

➡SA1384 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*). Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration

GLUCOSE [DEXTROSE]			
Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO		5	Biomed
Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO	14.50	1	Biomed
POTASSIUM CHLORIDE			
k Inj 75 mg per ml, 10 ml	55.00	50	AstraZeneca
ODIUM BICARBONATE			
Inj 8.4%, 50 ml	19.95	1	 Biomed
a) Up to 5 inj available on a PSO			
b) Not in combination			
Inj 8.4%, 100 ml	20.50	1	 Biomed
a) Up to 5 inj available on a PSO			
b) Not in combination			
ODIUM CHLORIDE			
Not funded for use as a nasal drop. Only funded for nebuliser us	e when in c	onjunction with	an antibiotic intended for
nebuliser use. Inj 0.9%, bag – Up to 2000 ml available on a PSO	4.00		
		E00 ml	- Dovtor
		500 ml 1 000 ml	✓ <u>Baxter</u> ✓ Baxter
	1.26	1,000 ml	✓ Baxter
Only if prescribed on a prescription for renal dialysis, materni	1.26	1,000 ml	✓ Baxter
Only if prescribed on a prescription for renal dialysis, materni for emergency use. (500 ml and 1,000 ml packs)	1.26 ity or post-na	1,000 ml	✓ Baxter
Only if prescribed on a prescription for renal dialysis, materni	1.26 ity or post-n: 33.00	1,000 ml atal care in the 5	✓ Baxter home of the patient, or on a PS
Only if prescribed on a prescription for renal dialysis, materni for emergency use. (500 ml and 1,000 ml packs) Inj 23.4% (4 mmol/ml), 20 ml ampoule	1.26 ity or post-na 33.00 rrmulae, pag	1,000 ml atal care in the 5	✓ Baxter home of the patient, or on a PS
Only if prescribed on a prescription for renal dialysis, materni for emergency use. (500 ml and 1,000 ml packs) Inj 23.4% (4 mmol/ml), 20 ml ampoule For Sodium chloride oral liquid formulation refer Standard Fo Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO	1.26 ity or post-na 33.00 irmulae, pag 7.00	1,000 ml atal care in the 5 ge 227 50	Baxter bome of the patient, or on a PS Biomed InterPharma Multichem
Only if prescribed on a prescription for renal dialysis, materni for emergency use. (500 ml and 1,000 ml packs) Inj 23.4% (4 mmol/ml), 20 ml ampoule For Sodium chloride oral liquid formulation refer Standard Fo Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO	1.26 ity or post-n: 33.00 irmulae, pag 7.00 6.63	1,000 ml atal care in the ge 227 50 50	 <u>Baxter</u> home of the patient, or on a PS <u>Biomed</u> InterPharma Multichem <u>Pfizer</u>
Only if prescribed on a prescription for renal dialysis, materni for emergency use. (500 ml and 1,000 ml packs) Inj 23.4% (4 mmol/ml), 20 ml ampoule For Sodium chloride oral liquid formulation refer Standard Fo Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO	1.26 ity or post-n: 33.00 rrmulae, pag 7.00 6.63 5.00	1,000 ml atal care in the 5 ge 227 50 50 20	 <u>Baxter</u> home of the patient, or on a PS <u>Biomed</u> InterPharma Multichem <u>Pfizer</u> Multichem
Only if prescribed on a prescription for renal dialysis, materni for emergency use. (500 ml and 1,000 ml packs) Inj 23.4% (4 mmol/ml), 20 ml ampoule For Sodium chloride oral liquid formulation refer Standard Fo Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO	1.26 ity or post-n: 33.00 irmulae, pag 7.00 6.63	1,000 ml atal care in the ge 227 50 50	 <u>Baxter</u> home of the patient, or on a PS <u>Biomed</u> InterPharma Multichem <u>Pfizer</u>
Only if prescribed on a prescription for renal dialysis, materni for emergency use. (500 ml and 1,000 ml packs) Inj 23.4% (4 mmol/ml), 20 ml ampoule For Sodium chloride oral liquid formulation refer Standard Fo Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO Inj 0.9%, 20 ml ampoule	1.26 ity or post-n. 33.00 rrmulae, pag 7.00 6.63 5.00 7.50 alist	1,000 ml atal care in the 5 50 50 50 20 30	 <u>Baxter</u> home of the patient, or on a PS <u>Biomed</u> InterPharma Multichem <u>Pfizer</u> Multichem InterPharma
Only if prescribed on a prescription for renal dialysis, materni for emergency use. (500 ml and 1,000 ml packs) Inj 23.4% (4 mmol/ml), 20 ml ampoule For Sodium chloride oral liquid formulation refer Standard Fo Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO Inj 0.9%, 20 ml ampoule	1.26 ity or post-n. 33.00 rrmulae, pag 7.00 6.63 5.00 7.50 alist	1,000 ml atal care in the 5 ge 227 50 50 20	 <u>Baxter</u> home of the patient, or on a PS <u>Biomed</u> InterPharma Multichem <u>Pfizer</u> Multichem
Only if prescribed on a prescription for renal dialysis, materni for emergency use. (500 ml and 1,000 ml packs) Inj 23.4% (4 mmol/ml), 20 ml ampoule For Sodium chloride oral liquid formulation refer Standard Fo Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO Inj 0.9%, 20 ml ampoule	1.26 ity or post-n. 33.00 rrmulae, pag 7.00 6.63 5.00 7.50 alist	1,000 ml atal care in the 5 50 50 50 20 30	 <u>Baxter</u> home of the patient, or on a PS <u>Biomed</u> InterPharma Multichem <u>Pfizer</u> Multichem InterPharma

- Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or
- When used in the extemporaneous compounding of eye drops; or
- 4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.

Inj 5 ml ampoule – Up to 5 inj available on a PSO	7.00	50	 InterPharma
Inj 10 ml ampoule - Up to 5 inj available on a PSO	6.63	50	 Pfizer
Inj 20 ml ampoule - Up to 5 inj available on a PSO	5.00	20	 Multichem
	7.50	30	 InterPharma

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's F \$	Price) Subsi Per	Fully dised	Brand or Generic Manufacturer
Oral Administration				
CALCIUM POLYSTYRENE SULPHONATE Powder	169.85	300 g OP	🗸 Ca	Ilcium Resonium
COMPOUND ELECTROLYTES Powder for oral soln – Up to 10 sach available on a PSO	2.30	10	✓ <u>En</u>	erlyte
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml)	6.55	1,000 ml OP		<u>dialyte -</u> Bubblegum
PHOSPHORUS Tab eff 500 mg (16 mmol)		100	🗸 Ph	osphate Phebra
POTASSIUM CHLORIDE				
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	5.26 (11.85)	60	Ch	lorvescent
* Tab long-acting 600 mg (8 mmol) SODIUM BICARBONATE	()	200	✓ <u>Sp</u>	<u>oan-K</u>
Cap 840 mg	8.52	100	✓ So ✓ So	
SODIUM POLYSTYRENE SULPHONATE				
Powder		454 g OP	✓ <u>Re</u>	esonium-A

		Subsidy		Fully	Brand or
		(Manufacturer's Price)		Subsidised	I Generic
		\$	Per		Manufacturer
A	Ipha-Adrenoceptor Blockers				
A	Ipha Adrenoceptor Blockers				
DO	XAZOSIN				
*	Tab 2 mg	6.75	500	1	Apo-Doxazosin
*	Tab 4 mg	9.09	500	1	Apo-Doxazosin
PH	ENOXYBENZAMINE HYDROCHLORIDE				_
*	Cap 10 mg	65.00	30	1	BNM S29
		216.67	100	✓	Dibenzyline S29
PR	AZOSIN				
*	Tab 1 mg	5.53	100	1	Apo-Prazosin
*	Tab 2 mg		100	1	Apo-Prazosin
*	Tab 5 mg		100	1	Apo-Prazosin
TE	RAZOSIN				
*	Tab 1 mg	0.59	28	1	Actavis
*	Tab 2 mg		500		Apo-Terazosin
*	Tab 5 mg		500	1	Apo-Terazosin

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

CAPTOPRIL		
* Oral liq 5 mg per ml94.99	95 ml OP	 Capoten
Oral liquid restricted to children under 12 years of age.		
CILAZAPRIL		
* Tab 0.5 mg2.00	90	 Zapril
* Tab 2.5 mg7.20	200	Apo-Cilazapril
* Tab 5 mg	200	✓ Apo-Cilazapril
ENALAPRIL MALEATE		
* Tab 5 mg	100	 Ethics Enalapril
* Tab 10 mg	100	 Ethics Enalapril
* Tab 20 mg7.12	100	 Ethics Enalapril
LISINOPRIL		•
* Tab 5 mg2.07	90	 Ethics Lisinopril
* Tab 10 mg2.36	90	 Ethics Lisinopril
* Tab 20 mg	90	 Ethics Lisinopril
PERINDOPRIL		i
* Tab 2 mg	30	Apo-Perindopril
* Tab 4 mg	30	✓ Apo-Perindopril
QUINAPRIL		- <u></u>
* Tab 5 mg6.01	90	Arrow-Quinapril 5
* Tab 10 mg	90	✓ Arrow-Quinapril 10
1	90 90	✓ Arrow-Quinapril 20
* Tab 20 mg4.89	30	

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic r Manufacturer
ACE Inhibitors with Diuretics			
CILAZAPRIL WITH HYDROCHLOROTHIAZIDE * Tab 5 mg with hydrochlorothiazide 12.5 mg	10.18	100	✓ <u>Apo-Cilazapril/</u> Hydrochlorothiazide
QUINAPRIL WITH HYDROCHLOROTHIAZIDE * Tab 10 mg with hydrochlorothiazide 12.5 mg * Tab 20 mg with hydrochlorothiazide 12.5 mg		30 30	✓ <u>Accuretic 10</u> ✓ <u>Accuretic 20</u>
Angiotensin II Antagonists			
CANDESARTAN CILEXETIL * Tab 4 mg * Tab 8 mg * Tab 16 mg * Tab 32 mg LOSARTAN POTASSIUM * Tab 12.5 mg * Tab 25 mg * Tab 50 mg * Tab 50 mg * Tab 100 mg * Tab 100 mg	2.28 	90 90 90 90 84 84 84 84	 Candestar Candestar Candestar Candestar Candestar Candestar Losartan Actavis Losartan Actavis Losartan Actavis Losartan Actavis Losartan Actavis
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE Tab 50 mg with hydrochlorothiazide 12.5 mg	1.88	30	✓ <u>Arrow-Losartan &</u> Hydrochlorothiazide

Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN - Special Authority see SA1751 below - Retail pharmacy

Note: Due to the angiotensin II receptor blocking activity of sacubitril with valsartan it should not be co-administered with an ACE inhibitor or another ARB.

Tab 24.3 mg with valsartan 25.7 mg	.190.00	56 •	Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	.190.00	56 •	Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	.190.00	56 •	Entresto 97/103

⇒SA1751 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or
 - 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per	• • • • • • • • • • • • • • • • • • •	Manufacturer
Antiarrhythmics				
For lignocaine hydrochloride refer to NERVOUS SYSTEM, Ana	esthetics. Local. page	118		
AMIODARONE HYDROCHLORIDE				
▲ Tab 100 mg - Retail pharmacy-Specialist	1 66	30	1	Cordarone-X
 Tab 100 mg – Retail pharmacy-Specialist Tab 200 mg – Retail pharmacy-Specialist 		30		Cordarone-X
Inj 50 mg per ml, 3 ml ampoule – Up to 6 inj available on a		5		Lodi
	11.98	6		Cordarone-X
	11.90	0	•	Coluarone-A
ATROPINE SULPHATE				
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available or				
PSO		10	~	Martindale
DIGOXIN				
* Tab 62.5 mcg - Up to 30 tab available on a PSO	6.67	240	1	Lanoxin PG
* Tab 250 mcg - Up to 30 tab available on a PSO		240	1	Lanoxin
* Oral lig 50 mcg per ml		60 ml		Lanoxin
			1	Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE				
	00.07	100		Duthmodon
▲ Cap 100 mg	23.87	100	v	Rythmodan
FLECAINIDE ACETATE – Retail pharmacy-Specialist				
▲ Tab 50 mg		60		Tambocor
▲ Cap long-acting 100 mg		30		Tambocor CR
Cap long-acting 200 mg		30		Tambocor CR
Inj 10 mg per ml, 15 ml ampoule	52.45	5	✓	Tambocor
MEXILETINE HYDROCHLORIDE				
▲ Cap 150 mg		100	1	Mexiletine
1 3				Hydrochloride
				USP S29
▲ Cap 250 mg	202.00	100	1	Mexiletine
				Hydrochloride
				USP S29
PROPAFENONE HYDROCHLORIDE - Retail pharmacy-Spec	ialict			
Tab 150 mg		50	1	Rytmonorm
		50	•	nyullollollii
Antihypotensives				
MIDODRINE – Special Authority see SA1474 below – Retail ph		100		Outron
Tab 2.5 mg		100		Gutron
Tab 5 mg		100	~	Gutron

■SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Note: Treatment should be started with small doses and titrated upwards as necessary. Hypertension should be avoided, and the usual target is a standing systolic blood pressure of 90 mm Hg.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

_
<u>)</u> Z
) <u>Z</u>)Z
12
)

		Subsidy (Manufacturer's Price) \$	Per		
PR	OPRANOLOL				
*	Tab 10 mg	4.64	100	✓	Apo-Propranolol
*	Tab 40 mg	5.72	100	1	Apo-Propranolol
*	Cap long-acting 160 mg		100	1	Cardinol LA
*	Oral liq 4 mg per ml – Special Authority see SA1327 below – Retail pharmacy		500 m	nl 🗸	Roxane S29

SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- Either:
 - 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
 - 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTALOL

* Tab 80 mg	500	✓ Mylan
* Tab 160 mg	 100	✓ Mylan
TIMOLOL		
* Tab 10 mg	 100	🗸 Apo-Timol

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

AMLODIPINE			
* Tab 2.5 mg	1.72	100	Apo-Amlodipine
* Tab 5 mg		250	Apo-Amlodipine
* Tab 10 mg		250	✓ Apo-Amlodipine
FELODIPINE			
* Tab long-acting 2.5 mg		30	Plendil ER
* Tab long-acting 5 mg		90	✓ Felo 5 ER
* Tab long-acting 10 mg		90	 Felo 10 ER
NIFEDIPINE			
* Tab long-acting 10 mg		60	 Adalat 10
			Adefin S29
* Tab long-acting 20 mg		100	Nyefax Retard
* Tab long-acting 30 mg	3.14	30	 Adalat Oros
			 Adefin XL
* Tab long-acting 60 mg	5.67	30	Adalat Oros
			 Adefin XL

	Subsidy (Manufacturer's Price		Fully Subsidised	
	\$	Per		Manufacturer
Other Calcium Channel Blockers				
DILTIAZEM HYDROCHLORIDE				
* Tab 30 mg	4.60	100		Dilzem
* Tab 60 mg		100		Dilzem
* Cap long-acting 120 mg		500		Apo-Diltiazem CD
* Cap long-acting 180 mg		500	-	Apo-Diltiazem CD
* Cap long-acting 240 mg		500	v	Apo-Diltiazem CD
PERHEXILINE MALEATE			-	
* Tab 100 mg	62.90	100	v	Pexsig
VERAPAMIL HYDROCHLORIDE				
* Tab 40 mg	7.01	100	-	Isoptin
* Tab 80 mg		100	-	Isoptin
* Tab long-acting 120 mg		250		Verpamil SR
* Tab long-acting 240 mg		250	v	Verpamil SR
* Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a		-		
PS0	25.00	5	v	Isoptin
Centrally-Acting Agents				
Centrally-Acting Agents				
CLONIDINE				
* Patch 2.5 mg, 100 mcg per day – Only on a prescription	7.40	4	✓	Mylan
 Patch 5 mg, 200 mcg per day – Only on a prescription 		4		Mylan
 Patch 7.5 mg, 300 mcg per day – Only on a prescription 	12.34	4	1	<u>Mylan</u>
CLONIDINE HYDROCHLORIDE				
* Tab 25 mcg		112	✓	Clonidine BNM
* Tab 150 mcg		100	✓	Catapres
 Inj 150 mcg per ml, 1 ml ampoule 	25.96	10	✓	Medsurge
METHYLDOPA				
* Tab 250 mg	15.10	100	✓	Methyldopa Mylan
	52.85	500	✓	Methyldopa Mylan
				S29 S29
Diuretics				
Loop Diuretics				
BUMETANIDE				
BOMETANIDE * Tab 1 mg	16.26	100	1	Burinex
* Tab Tring		5		Burinex
		5	•	Durniex
FUROSEMIDE [FRUSEMIDE] * Tab 40 mg – Up to 30 tab available on a PSO	0.00	1 000		Diurin 40
* Tab 40 mg – Op to 30 tab available on a PSO		1,000 50	-	Diurin 40 Urex Forte
* Oral liq 10 mg per ml		30 ml C		Lasix
 * Inj 10 mg per ml, 25 ml ampoule 		6		Lasix
 * Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a 		5		Frusemide-Claris
Potassium Sparing Diuretics				
AMILORIDE HYDROCHLORIDE	20.00	05 ml C		Riomod
Oral liq 1 mg per ml		25 ml C	7r° ▼	Biomed

fully subsidised
 Sole Subsidised Supply

52

(\$29) Unapproved medicine supplied under Section 29

	0.1.11			
	Subsidy		Fully	
	(Manufacturer's Price \$	e) 3 Per	Subsidised	Generic Manufacturer
	*	1.01	•	Manufacturer
EPLERENONE – Special Authority see SA1728 below – Re				
Tab 50 mg		30		Inspra
Tab 25 mg	11.87	30	-	Inspra
➡SA1728 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals	s valid without further rer	newal unl	less notifi	ed for applications meeting
the following criteria:				
Both:				
1 Patient has heart failure with ejection fraction less that	an 40%; and			
2 Either:	an io,o, and			
2.1 Patient is intolerant to optimal dosing of spiror	alactora: or			
		ontimal	docina of	cnironolactono
2.2 Patient has experienced a clinically significan	adverse enect while on	optimat	uosing oi	spironolacione.
METOLAZONE				
Tab 5 mg	CBS	1	✓	Metolazone S29
		50	1	Zaroxolyn S29
		00	•	Laroxolyn
SPIRONOLACTONE	4.00	400		.
* Tab 25 mg		100		Spiractin
* Tab 100 mg		100	-	Spiractin
Oral liq 5 mg per ml		25 ml OF	√	Biomed
Potassium Sparing Combination Diuretics				
AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE				
 * Tab 5 mg with furosemide 40 mg 	8 63	28	1	Frumil
6 6		20	•	- Tunin
AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTH				
* Tab 5 mg with hydrochlorothiazide 50 mg	5.00	50	~	Moduretic
Thispide and Deleted Diverties				
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]				
* Tab 2.5 mg – Up to 150 tab available on a PSO	12 50	500	1	Arrow-
	12.00	000	•	Bendrofluazide
				Denaronuaziae
May be supplied on a PSO for reasons other than e	mergency.			
* Tab 5 mg		500	✓	Arrow-
C C C C C C C C C C C C C C C C C C C				Bendrofluazide
CHLOROTHIAZIDE				
Oral liq 50 mg per ml		25 ml OF	> √	Biomed
CHLORTALIDONE [CHLORTHALIDONE]				
* Tab 25 mg	8.00	50	1	Hygroton
-	0.00	50	•	
INDAPAMIDE	0.00	~~		David Taka
* Tab 2.5 mg	2.60	90	~	Dapa-Tabs
Lipid-Modifying Agents				
Lipid-modifying Agents				
Fibrates				
BEZAFIBRATE				
* Tab 200 mg	19.01	90		Bezalip
* Tab long-acting 400 mg		30	✓	Bezalip Retard

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
EMFIBROZIL Tab 600 mg		60	1	Lipazil
Other Lipid-Modifying Agents				
CIPIMOX				•
Cap 250 mg		30	~	Olbetam
COTINIC ACID Tab 50 mg	4 12	100	1	Apo-Nicotinic Acid
Tab 500 mg		100		Apo-Nicotinic Acid
lesins				
DLESTIPOL HYDROCHLORIDE				
Grans for oral liq 5 g		30	v	Colestid
IMG CoA Reductase Inhibitors (Statins)				
rdiovascular risk of 15% or greater. FORVASTATIN – See prescribing guideline above Tab 10 mg Tab 20 mg Tab 40 mg Tab 80 mg	9.99 15.93	500 500 500 500	1 1	Lorstat Lorstat Lorstat Lorstat
AVASTATIN – See prescribing guideline above	4.70	100		Ana Dravastatin
Tab 20 mg Tab 40 mg		100 100		Apo-Pravastatin Apo-Pravastatin
VVASTATIN – See prescribing guideline above		100	-	<u> </u>
Tab 10 mg	0.95	90	1	Simvastatin Mylan
Tab 20 mg		90		Simvastatin Mylan
Tab 40 mg		90		Simvastatin Mylan
Tab 80 mg	6.00	90	•	Simvastatin Mylan
elective Cholesterol Absorption Inhibitors				
ETIMIBE – Special Authority see SA1045 below – Retail ph Tab 10 mg		30	1	Ezetimibe Sandoz
SA1045 Special Authority for Subsidy		00	•	
tial application from any relevant practitioner. Approvals va of the following:	lid for 2 years for appli	catio	ns meeting	the following criteria:
 Patient has a calculated absolute risk of cardiovascular Patient's LDL cholesterol is 2.0 mmol/litre or greater; an Any of the following: 		ove	r 5 years; a	and
3.1 The patient has rhabdomyolysis (defined as mus treated with one statin; or		kinas	se more th	an 10 × normal) when

- 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
- 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	;	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy. If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 below - Retail pharmacy

Tab 10 mg with simvastatin 10 mg5.15	30	 Zimybe
Tab 10 mg with simvastatin 20 mg6.15	30	 Zimybe
Tab 10 mg with simvastatin 40 mg7.15	30	 Zimybe
Tab 10 mg with simvastatin 80 mg8.15	30	 Zimybe

⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be

performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

GLYCERYL TRINITRATE

*	Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO4.45	250 dose OP	 Nitrolingual Pump Spray
*	Oral spray, 400 mcg per dose – Up to 200 dose available on a		
	PSO	200 dose OP	 Glytrin
*	Patch 25 mg, 5 mg per day15.73	30	 Nitroderm TTS
*	Patch 50 mg, 10 mg per day	30	 Nitroderm TTS
ISC	SORBIDE MONONITRATE		
*	Tab 20 mg	100	🖌 Ismo 20
	Tab long-acting 40 mg7.50	30	 Ismo 40 Retard
*	Tab long-acting 60 mg8.29	90	✓ Duride

	Subsidy		Fully Brand or
	(Manufacturer's Price)		Subsidised Generic
	\$	Per	 Manufacturer
Sympathomimetics			
ADRENALINE			
Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PS	SO4.98 5.25	5	 ✓ Aspen Adrenaline ✓ Hospira
Inj 1 in 10,000, 10 ml ampoule – Up to 5 inj available on a	PSO27.00	5	 Hospira
	49.00	10	 Aspen Adrenaline
ISOPRENALINE [ISOPROTERENOL]			
* Inj 200 mcg per ml, 1 ml ampoule		25	
	(164.20)		Isuprel
N/ III I			
Vasodilators			
HYDRALAZINE HYDROCHLORIDE			
* Tab 25 mg – Special Authority see SA1321 below – Retail			
pharmacy		1	 Hydralazine
		56	 Onelink S29
		84	✓ AMDIPHARM S29
		100	 Onelink S29
* Inj 20 mg ampoule		5	✓ Apresoline
 For the treatment of refractory hypertension; or For the treatment of heart failure in combination with a n inhibitors and/or angiotensin receptor blockers. 	itrate, in patients who a	are into	lerant or have not responded to ACE
MINOXIDIL			
Tab 10 mg	70.00	100	 Loniten
NICORANDIL			
Tab 10 mg		60	✓ Ikorel
Tab 20 mg		60	 Ikorel
PAPAVERINE HYDROCHLORIDE			• • •
* Inj 12 mg per ml, 10 ml ampoule		5	 Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]			_
Tab 400 mg	42.26	50	 Trental 400
Endothelin Receptor Antagonists			
AMBRISENTAN – Special Authority see SA1702 below – Reta	il pharmacy		
Tab 5 mg		30	✓ Volibris
Tab 10 mg	4,585.00	30	 Volibris
SA1702 Special Authority for Subsidy			
Special Authority approved by the Pulmonary Arterial Hyperten			
Notes: Application details may be obtained from PHARMAC's	website http://www.pha	rmac.g	j <u>ovt.nz</u> or:
The Coordinator, PAH Panel			
PHARMAC, PO Box 10-254, WELLINGTON	an anut nz		
Tel: (04) 916 7561, Fax: (04) 974 4858, Email: <u>PAH@pharma</u>	ac.yuvi.nz		

✓ fully subsidised Sole Subsidised Supply

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
BOSENTAN - Special Authority see SA1712 below - Retail pha	rmacy			
Tab 62.5 mg	141.00	60	✓	Bosentan Dr
Tab 125 mg	141.00	60	1	<u>Reddy's</u> Bosentan Dr <u>Reddy's</u>

► SA1712 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil; or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria: Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic	
\$	Per	~	Manufacturer	

continued...

- 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
- 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see SA1738 below – Reta	ail pharmacy		
Tab 25 mg	0.64	4	 Vedafil
Tab 50 mg	0.64	4	✓ Vedafil
Tab 100 mg	6.60	12	✓ Vedafil

⇒SA1738 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II; or
 - 3.2 PAH is in NYHA/WHO functional class III; or
 - 3.3 PAH is in NYHA/WHO functional class IV; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Prostacyclin Analogues

EPOPROSTENOL - Special Authority see SA1696 on the	next page – Retail pharm	nacy	
Inj 500 mcg vial		1	 Veletri
Inj 1.5 mg vial	73.21	1	 Veletri

	Subsidy (Manufacturer's Price) \$	Su Per	Fully Ibsidised	Brand or Generic Manufacturer
SA1696 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertensi Notes: Application details may be obtained from PHARMAC's w The Coordinator, PAH Panel PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 916 7561, Fax: (04) 974 4858, Email: <u>PAH@pharmac</u>	ebsite <u>http://www.pha</u>	rmac.gc	ovt.nz or:	
ILOPROST – Special Authority see SA1705 below – Retail phar Nebuliser soln 10 mcg per ml, 2 ml		30	✓ V	entavis
► SA1705 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertensi Notes: Application details may be obtained from PHARMAC's w The Coordinator, PAH Panel PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 916 7561, Fax: (04) 974 4858, Email: <u>PAH@pharmac</u>	ebsite <u>http://www.pha</u>	rmac.gc	ovt.nz or:	

Subsidy (Manufacturer's Price) \$) Sub Per	Fully sidised	Brand or Generic Manufacturer
, page 88			
	0	-	Differin
	30 g OP	✓ L	Differin
pharmacy			
8.14	60	✓	Dratane
	120	✓	Dratane
20.49	120	✓	Dratane
	(Manufacturer's Price) \$, page 88 22.89	(Manufacturer's Price) Sub \$ Per , page 88 	(Manufacturer's Price) Subsidised \$ Per ✓ , page 88

⇒SA1475 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
 - 3.2 Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
- 2 Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

TRETINOIN Crm 0.5 mg per g – Maximum of 50 g per prescription13.90	50 g OP	✓ <u>ReTrieve</u>	
Antibacterials Topical			
For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 88			
HYDROGEN PEROXIDE			
* Crm 1%	10 g OP 15 a OP	 Crystaderm Crystaderm 	

	Subsidy (Manufacturer's Pric		Fully Brand or bidised Generic
	(Manulacturer's Pric	e) Suc Per	Manufacturer
MUPIROCIN			
Oint 2%	6.60 (9.26)	15 g OP	Bactroban
a) Only on a prescription	(0.20)		Baotroball
b) Not in combination			
SODIUM FUSIDATE [FUSIDIC ACID]			
Crm 2%	1.59 2.52	5 g OP 15 g OP	 ✓ Foban ✓ DP Fusidic Acid Cream
a) Maximum of 15 g per prescription			•••••
b) Only on a prescription			
c) Not in combination			
d) Foban to be Sole Supply on 1 August 2019	4 50		/ - ·
Oint 2%	1.59	5 g OP	 Foban
 a) Maximum of 15 g per prescription b) Only on a prescription 			
c) Not in combination			
d) Foban to be Sole Supply on 1 August 2019			
DP Fusidic Acid Cream Crm 2% to be delisted 1 August 2019)			
SULFADIAZINE SILVER			
Crm 1%	10.80	50 g OP	✓ Flamazine
a) Up to 250 g available on a PSO			
b) Not in combination			
Antifungals Topical			
For systemic antifungals, refer to INFECTIONS, Antifungals, pa	ge 95		
AMOROLFINE	-		
a) Only on a prescription			
b) Not in combination			
Nail soln 5%	15.95	5 ml OP	✓ MycoNail
CICLOPIROX OLAMINE			
a) Only on a prescription			
b) Not in combination			
Nail-soln 8%	5.72	7 ml OP	✓ <u>Apo-Ciclopirox</u>
	0.70	00 05	
* Crm 1%	0.70	20 g OP	✓ Clomazol
a) Only on a prescriptionb) Not in combination			
* Soln 1%		20 ml OP	
	(7.55)		Canesten
a) Only on a prescription	· · /		
b) Not in combination			

	Subsidy				
	(Manufacturer's F \$	Price) Sub: Per	sidised Generic Manufacturer		
CONAZOLE NITRATE	Ŷ		manataotaron		
Crm 1%	1.00	20 g OP			
0111 1/0	(7.48)	20 9 01	Pevaryl		
a) Only on a prescription	(1.10)		i ovaryi		
b) Not in combination					
Foaming soln 1%, 10 ml sachets	9.89	3			
	(17.23)	C C	Pevaryl		
a) Only on a prescription	(-)		, , , , , , , , , , , , , , , , , , ,		
b) Not in combination					
₭ Crm 2%	0.74	15 g OP	✓ Multichem		
a) Only on a prescription		10 9 01			
b) Not in combination					
€ Lotn 2%	4.36	30 ml OP			
	(10.03)		Daktarin		
a) Only on a prescription	()				
b) Not in combination					
₭ Tinct 2%		30 ml OP			
	(12.10)		Daktarin		
a) Only on a prescription	· · · · ·				
b) Not in combination					
VYSTATIN					
Crm 100,000 u per g	1.00	15 g OP			
e	(7.90)		Mycostatin		
a) Only on a prescription	()				
b) Not in combination					
,					
Antipruritic Preparations					
CALAMINE					
a) Only on a prescription					
b) Not in combination					
Crm, aqueous, BP		100 g	 healthE Calamine 		
		5	Aqueous Cream		
			BP		
Lotn, BP		2,000 ml	✓ PSM		
CROTAMITON					
a) Only on a prescription					
b) Not in combination					
Crm 10%	3.29	20 g OP	✓ Itch-Soothe		
IENTHOL – Only in combination		0			
1) Only in combination with a dermatological base	or proprietary Topical (Corticosteriod -	Plain		
2) With or without other dermatological galenicals.	or propriotary Topical C				
Crystals	6 00	25 a	✓ MidWest		
Crystals		25 g 100 g	✓ Midwest ✓ MidWest		
	29.00	100 y	- WILLWEST		

	Subsidy (Manufacturer's P \$	rice) Subs Per	Fully Brand or sidised Generic ✓ Manufacturer	
Corticosteroids Topical				
For systemic corticosteroids, refer to CORTICOSTEROIDS AN	D RELATED AGE	NTS, page 78		
Corticosteroids - Plain				
BETAMETHASONE DIPROPIONATE				
Crm 0.05%		15 g OP	 Diprosone 	
	8.97	50 g OP	✓ Diprosone	
Crm 0.05% in propylene glycol base		30 g OP	 Diprosone OV 	
Oint 0.05%		15 g OP	✓ Diprosone	
	8.97	50 g OP	 Diprosone 	
Oint 0.05% in propylene glycol base	4.33	30 g OP	 Diprosone OV 	
BETAMETHASONE VALERATE		5	P	
* Crm 0.1%	2.45	50 g OP	 Beta Cream 	
♣ Oint 0.1%		50 g OP 50 g OP	✓ Beta Ointment	
★ Onit 0.1 %		50 g OF 50 ml OP	✓ Betnovate	
	10.00	50 mi OF	• Delilovale	
CLOBETASOL PROPIONATE			<i>.</i> .	
₭ Crm 0.05%		30 g OP	✓ <u>Dermol</u>	
₭ Oint 0.05%	2.20	30 g OP	Dermol	
CLOBETASONE BUTYRATE				
Crm 0.05%	5.38	30 g OP		
	(7.09)	-	Eumovate	
DIFLUCORTOLONE VALERATE	. ,			
Crm 0.1%	8 97	50 g OP		
	(15.86)	00 9 01	Nerisone	
Fatty oint 0.1%		50 g OP	Nonoono	
	(15.86)	00 9 01	Nerisone	
IVERGOODTICONE	(10.00)		Honoono	
		00 - 00		
Crm 1% – Only on a prescription		30 g OP	✓ <u>DermAssist</u>	
Powder – Only in combination	16.25	500 g	Pharmacy Health	
		25 g	✓ <u>ABM</u>	
Up to 5% in a dermatological base (not proprietary Top galenicals	Dical Corticosteriod	– Plain) with c	or without other dermatologica	ai
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only	/ on			
a prescription		250 ml	DP Lotn HC	
Lipocream 0.1%	3 / 2	30 g OP	Locoid Lipocream	
Liputicani U. I /0		30 g OP 100 g OP	 Locoid Lipocream 	
Oint 0.1%		100 g OP 100 g OP	 Locoid Lipocream Locoid 	
Oint 0.1% Milky emul 0.1%		100 g OP 100 ml OP	✓ Locoid Crelo	
,	10.70			
METHYLPREDNISOLONE ACEPONATE				
Crm 0.1%		15 g OP	 Advantan 	
Oint 0.1%		15 g OP	Advantan	

	Subsidy		Fully Brand or
	(Manufacturer's Pr \$	ice) Subs Per	sidised Generic Manufacturer
OMETASONE FUROATE	Ŷ		manaratara
Crm 0.1%	1 51	15 g OP	Elocon Alcohol Free
	2.50	50 g OP	✓ Elocon Alcohol Free
Oint 0.1%		15 g OP	✓ Elocon
	2.90	50 g OP	✓ Elocon
Lotn 0.1%		30 ml OP	✓ Elocon
RIAMCINOLONE ACETONIDE			
Crm 0.02%	6.30	100 g OP	 Aristocort
Oint 0.02%		100 g OP	✓ Aristocort
		100 g 01	
Corticosteroids - Combination			
ETAMETHASONE VALERATE WITH CLIOQUINOL - Only of	on a prescription		
Crm 0.1% with clioquinol 3%	3.49	15 g OP	
	(4.90)		Betnovate-C
ETAMETHASONE VALERATE WITH SODIUM FUSIDATE [F	USIDIC ACID		
Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP	
	(10.45)	Ũ	Fucicort
a) Maximum of 15 g per prescription			
b) Only on a prescription			
YDROCORTISONE WITH MICONAZOLE - Only on a presc	ription		
Crm 1% with miconazole nitrate 2%		15 g OP	 Micreme H
YDROCORTISONE WITH NATAMYCIN AND NEOMYCIN -		0	
Crm 1% with natamycin 1% and neomycin sulphate 0.5%.		15 g OP	Pimafucort
Oint 1% with natamycin 1% and neomycin sulphate 0.5%.		15 g OP	 Pimafucort Pimafucort
		•	
RIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMY		N	
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5	•	15 × OD	
and gramicidin 250 mcg per g – Only on a prescription		15 g OP	
	(6.60)		Viaderm KC
Disinfecting and Cleansing Agents			
HLORHEXIDINE GLUCONATE – Subsidy by endorsement			
a) No more than 500 ml per month			
b) Only if prescribed for a dialysis patient and the prescrip		cordingly.	
Handrub 1% with ethanol 70%		500 ml	 healthE
Soln 4% wash	3.98	500 ml	 healthE
RICLOSAN – Subsidy by endorsement			
a) Maximum of 500 ml per prescription			
b)			
a) Only if prescribed for a patient identified with Meth	nicillin-resistant Star	phylococcus a	ureus (MRSA) prior to electiv
surgery in hospital and the prescription is endorse			· · · · ·
		infaction and	the preserintion is endereed
b) Only if prescribed for a patient with recurrent Stap	hylococcus aureus	intection and	the prescription is endorsed
	hylococcus aureus	intection and	the prescription is endorsed

	Subsidy (Manufacturer's F		Fully Brand or idised Generic
	\$	Per	 Manufacturer
Barrier Creams and Emollients			
Barrier Creams			
DIMETHICONE			_
* Crm 5% pump bottle	4.59	500 ml OP	✓ <u>healthE</u> Dimethicone 5%
* Crm 10% pump bottle	4.52	500 ml OP	✓ <u>healthE</u> Dimethicone 10%
ZINC AND CASTOR OIL			
* Oint	4.25	500 g	 Boucher
Emollients			
AQUEOUS CREAM			
* Crm	1.92	500 g	 Boucher
CETOMACROGOL * Crm BP	2 48	500 g	✓ healthE
CETOMACROGOL WITH GLYCEROL		000 g	Indulina
Crm 90% with glycerol 10%	2.82	500 ml OP	 <u>Pharmacy Health</u> <u>Sorbolene with</u> Glycerin
	3.87	1,000 ml OP	Pharmacy Health Sorbolene with Glycerin
EMULSIFYING OINTMENT			<i></i>
	3.59	500 g	✓ <u>AFT</u>
OIL IN WATER EMULSION * Crm	2.19	500 g	✓ O/W Fatty Emulsion
		Ū	Cream
PARAFFIN Oint liquid paraffin 50% with white soft paraffin 50%	5.35	500 ml OP	✓ <u>healthE</u>
UREA * Crm 10%	1.37	100 g OP	✓ healthE Urea Cream
WOOL FAT WITH MINERAL OIL – Only on a prescription		.00 9 01	- Inculting of our of our
* Lotn hydrous 3% with mineral oil		1,000 ml	
	(11.95) 1.40	250 ml OP	DP Lotion
	(4.53)	200 IIII OF	DP Lotion
	5.60	1,000 ml	Alaba Karilatian
	(20.53) (23.91)		Alpha-Keri Lotion BK Lotion
	1.40	250 ml OP	
	(7.73)		BK Lotion

	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully Brand or sidised Generic Manufacturer
Other Dermatological Bases			
ARAFFIN White soft – Only in combination	20.20 3.58	2,500 g 500 g	✔ IPW
	(7.78) (8.69)	Ū	IPW PSM
Only in combination with a dermatological galenical or a	s a diluent for a p	proprietary Top	ical Corticosteroid – Plain.
Minor Skin Infections			
POVIDONE IODINE			
Oint 10%	3.27	25 g OP	 Betadine
 a) Maximum of 100 g per prescription b) Only on a prescription 			
Antiseptic soln 10%	6.20	500 ml	✓ Betadine✓ Riodine
	1.28	100 ml	
	(4.20)		Riodine
	(13.27) 0.19	15 ml	Betadine
	(7.41)	10 111	Betadine
Skin preparation, povidone iodine 10% with 30% alcohol		500 ml 100 ml	 Betadine Skin Prep
Ohio presention, posidence is direct 00% with 700% placehol	(3.48)	100	Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	1.63 (6.64)	100 ml	Pfizer
Parasiticidal Preparations			
IMETHICONE			
₭ Lotn 4%		200 ml OP	✓ <u>healthE</u> <u>Dimethicone 4%</u> <u>Lotion</u>
/ERMECTIN – Special Authority see SA1225 below – Retail pl		4	✓ Stromectol
Tab 3 mg – Up to 100 tab available on a PSO 1) PSO for institutional use only. Must be endorsed a a valid Special Authority for patient of that institution	with the name of		

- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or penal institutions.

⇒SA1225 Special Authority for Subsidy

Initial application — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

66

1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	~	Manufacturer	

continued...

2 Either:

- 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
- 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.
- Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's F \$	^P rice) Per	Fully Subsidised	Brand or Generic Manufacturer
 continued Renewal — (Other parasitic infections) only from an infections valid for 1 month for applications meeting the folloc Any of the following: Filaricides; or Cutaneous larva migrans (creeping eruption); or Strongyloidiasis. 		list, clinica	l microbiolo	gist or dermatologist.
PERMETHRIN Crm 5% Lotn 5%		30 g C 30 ml C		Lyderm A-Scabies
PHENOTHRIN Shampoo 0.5%	11.36	200 ml	OP 🗸 I	Parasidose
Psoriasis and Eczema Preparations				
ACITRETIN – Special Authority see SA1476 below – Retail Cap 10 mg Cap 25 mg		d general p in and is co sk of terato gnancy ha nat she mu treatment;	s meeting the practitioner, competent to genicity if a s been exc st not become or	or nurse practitioner prescribe acitretin; and ucitretin is used during luded prior to the me pregnant during
Either: 1 Patient is female and has been counselled and under and the applicant has ensured that the possibility of p treatment and that the patient is informed that she mu years after the completion of the treatment; or 2 Patient is male. BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIO Cal E00 mag with advincting E0 mag parts	regnancy has been e ust not become pregr L	excluded p	rior to the c treatment	commencement of the
Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g		30 g C		Daivobet
CALCIPOTRIOL Oint 50 mcg per g	45.00	100 g (DP √	Daivonex

COAL TAR

68

1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteriod – Plain

2) With or without other dermatological galenicals.

	Subsidy (Manufacturer's F	Price) Subs	Fully Brand or idised Generic
	(Manulacturers F	Per	Manufacturer
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SUL	PHUR		
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% at			
allantoin crm 2.5%		75 g OP	
	(8.00)	70 g 01	Egopsoryl TA
	3.43	30 g OP	
	(4.35)	00 9 01	Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR	(1.00)		_gopool):
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.07	25 g OP	 Coco-Scalp
	7.95	25 g OP 40 g OP	✓ Coco-Scalp
		0	•
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORI			
₭ Soln 2.3% with trolamine laurilsulfate and fluorescein sodium	113.86	500 ml	Pinetarsol
SALICYLIC ACID			
Powder – Only in combination		250 g	✓ PSM
 Only in combination with a dermatological base o With or without other dermatological galenicals. 	r proprietary Topic	cal Corticostero	id – Plain or collodion flexible
SULPHUR			
Precipitated – Only in combination	6.35	100 g	 Midwest
 With or without other dermatological galenicals. Scalp Preparations 			
BETAMETHASONE VALERATE ₭ Scalp app 0.1%	7 75	100 ml OD	/ Pata Caalm
		100 ml OP	 Beta Scalp
CLOBETASOL PROPIONATE			
₭ Scalp app 0.05%	6.96	30 ml OP	 Dermol
IYDROCORTISONE BUTYRATE			
Scalp lotn 0.1%	7.30	100 ml OP	✓ Locoid
ETOCONAZOLE			
Shampoo 2%	2.99	100 ml OP	 Sebizole
a) Maximum of 100 ml per prescription			
b) Only on a prescription			
·/ - / · ··· F···			
Sunscreens			
SUNSCREENS, PROPRIETARY – Subsidy by endorsement			
Only if prescribed for a patient with severe photosensitivity	secondary to a de	fined clinical co	ondition and the prescription is
endorsed accordingly.			
Crm	3.30	100 g OP	
	(5.89)		Hamilton Sunscreen
Lotn,	3.30	100 g OP	 Marine Blue Lotion
			SPF 50+
	5.10	200 g OP	 Marine Blue Lotion

SPF 50+

	Subsidy (Manufacturer's Price \$) Sub Per	Fully sidised	Brand or Generic Manufacturer
Wart Preparations				
For salicylic acid preparations refer to PSORIASIS AND ECZEM	A PREPARATIONS,	page 68		
IMIQUIMOD Crm 5%, 250 mg sachet	21.72	24	✓ <u>P</u>	errigo
PODOPHYLLOTOXIN Soln 0.5%		8.5 ml OP	√ c	condyline
a) Maximum of 3.5 ml per prescriptionb) Only on a prescription				
Other Skin Preparations				
Antineoplastics				
FLUOROURACIL SODIUM Crm 5%	7.95	20 g OP	✓ <u>E</u>	fudix

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Contraceptives - Non-hormonal				
Condoms				
CONDOMS				
* 49 mm – Up to 144 dev available on a PSO		144	✓	Shield 49
* 53 mm – Up to 144 dev available on a PSO	1.11	12		Gold Knight Shield Blue
	13.36	144	✓	Shield Blue
* 53 mm (chocolate) – Up to 144 dev available on a PSO	1.11	12	✓	Gold Knight
	13.36	144	✓	Gold Knight
* 53 mm (strawberry) – Up to 144 dev available on a PSO	1.11	12	✓	Gold Knight
	13.36	144		Gold Knight
* 56 mm – Up to 144 dev available on a PSO	1.11	12		Gold Knight
	13.36	144		Durex Extra Safe Gold Knight
* 56 mm, shaped – Up to 144 dev available on a PSO	1.11	12	✓	Durex Confidence
	13.36	144	✓	Durex Confidence
* 60 mm – Up to 144 dev available on a PSO	13.36	144	1	Shield XL
Contraceptive Devices				
INTRA-UTERINE DEVICE a) Up to 40 dev available on a PSO b) Only on a PSO				
* IUD 29.1 mm length × 23.2 mm width		1	✓	Choice TT380 Short
# IUD 33.6 mm length × 29.9 mm width		1	1	Choice TT380 Standard
* IUD 35.5 mm length × 19.6 mm width	31.60	1	1	Choice Load 375
Contraceptives - Hormonal				

GENITO-URINARY SYSTEM

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 Either:

- 1.1 Patient is on a Social Welfare benefit; or
- 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Subsic Per	Fully lised	Brand or Generic Manufacturer
continued				
The additional subsidy will fund Mercilon and Marvelon up to the	manufacturer's price	for each of	these	products as identified on
the Schedule at 1 November 1999.				
Special Authorities approved before 1 November 1999 remain va	lid until the expiry dat	e and can I	be ren	ewed providing that
 women are still either: on a Social Welfare benefit; or 				
 bit a Social Weilare benefit, of have an income no greater than the benefit. 				
The approval numbers of Special Authorities approved before 1 N	November 1999 are in	terchangea	ble fo	r products within the
combined oral contraceptives and progestogen-only contraceptiv		•		
ETHINYLOESTRADIOL WITH DESOGESTREL	0 1 7 1		0,	
* Tab 20 mcg with desogestrel 150 mcg and 7 inert tab	6.62	84		
	(19.80)		N	lercilon 28
 a) Higher subsidy of \$13.80 per 84 tab with Special Aut b) Up to 84 tab available on a PSO 	hority see SA0500 on	the previou	us pag	je
* Tab 30 mcg with desogestrel 150 mcg and 7 inert tab		84		
	(19.80)			larvelon 28
 a) Higher subsidy of \$13.80 per 84 tab with Special Aut b) Up to 84 tab available on a PSO 	hority see SA0500 on	the previou	us pag	Je
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets				
Up to 84 tab available on a PSO		84	✓ <u>N</u>	licrogynon 20 ED
* Tab 50 mcg with levonorgestrel 125 mcg and 7 inert tab – U				
to 84 tab available on a PSO		84	✓ N	licrogynon 50 ED
* Tab 30 mcg with levonorgestrel 150 mcg	(16.50)	63	N	licrogynon 30
a) Higher subsidy of \$15.00 per 63 tab with Special Aut	()	the previou		
b) Up to 63 tab available on a PSO			io pug	
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets	_			
Up to 84 tab available on a PSO		84	✓ L	evlen ED
ETHINYLOESTRADIOL WITH NORETHISTERONE				
* Tab 35 mcg with norethisterone 1 mg - Up to 63 tab availab	le			
on a PSO	6.62	63	🗸 В	revinor 1/21
* Tab 35 mcg with norethisterone 1 mg and 7 inert tab – Up to				
84 tab available on a PSO	6.62	84	✓ В	Frevinor 1/28
* Tab 35 mcg with norethisterone 500 mcg – Up to 63 tab	6.60	60		and the second
available on a PSO * Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - U		63	• B	revinor 21
to 84 tab available on a PSO		84	🖌 N	lorimin
(Brevinor 1/21 Tab 35 mcg with norethisterone 1 mg to be deliste		J -	- 0	····
(Brevinor 21 Tab 35 mcg with norethisterone 500 mcg to be delise	, ,			

Progestogen-only Contraceptives

► SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

GENITO-URINARY SYSTEM

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

1 Either:

- 1.1 Patient is on a Social Welfare benefit; or
- 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

* Tab 30 mcg	6.62	84	
-	(16.50)		Microlut
 a) Higher subsidy of \$13.80 per 84 tab with Special Authority b) Up to 84 tab available on a PSO 	/ see SA0500) on the prev	<i>r</i> ious page
 Subdermal implant (2 x 75 mg rods) – Up to 3 pack available on a PSO 	. 106.92	1	✓ Jadelle
MEDROXYPROGESTERONE ACETATE Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PSO	7.25	1	✓ Depo-Provera
NORETHISTERONE			
* Tab 350 mcg – Up to 84 tab available on a PSO	6.25	84	 Noriday 28
Emergency Contraceptives			
LEVONORGESTREL			
* Tab 1.5 mg	4.95	1	Postinor-1

a) Maximum of 2 tab per prescription

b) Up to 5 tab available on a PSO

c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

 Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO......4.67 168 ✓ <u>Ginet</u>

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully Brand or
	(Manufacturer's P \$	rice) Subs Per	idised Generic Manufacturer
Gynaecological Anti-infectives			
ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC A	CID		
Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate		100 - 00	
0.025%, glycerol 5% and ricinoleic acid 0.75% with applic	ator8.43 (24.00)	100 g OP	Aci-Jel
CLOTRIMAZOLE	(24.00)		
* Vaginal crm 1% with applicators		35 g OP	✓ Clomazol
* Vaginal crm 2% with applicators	2.10	20 g OP	✓ <u>Clomazol</u>
MICONAZOLE NITRATE * Vaginal crm 2% with applicator	2 00	40 g OP	✓ Micreme
NYSTATIN		40 y OF	• <u>Micreille</u>
Vaginal crm 100,000 u per 5 g with applicator(s)	4.45	75 g OP	✓ <u>Nilstat</u>
Myometrial and Vaginal Hormone Preparations			
ERGOMETRINE MALEATE			
Inj 250 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO	454.00	5	Ergonovine S29
Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on a		Ũ	g
PSO		5	 DBL Ergometrine
(Ergonovine ^{\$29} Inj 250 mcg per ml, 1 ml ampoule to be delisted	1 July 2019)		
OESTRIOL * Crm 1 mg per g with applicator	6 62	15 g OP	✓ Ovestin
 Pessaries 500 mcg 		15	✓ <u>Ovestin</u>
OXYTOCIN – Up to 5 inj available on a PSO			
Inj 5 iu per ml, 1 ml ampoule		5	Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule OXYTOCIN WITH ERGOMETRINE MALEATE – Up to 5 inj availa		5	 Oxytocin BNM
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml		5	 Syntometrine
			_
Pregnancy Tests - hCG Urine			
PREGNANCY TESTS - HCG URINE			
a) Up to 200 test available on a PSOb) Only on a PSO			
Cassette		40 test OP	 Smith BioMed Rapid
			Pregnancy Test
Urinary Agents			
	107		
For urinary tract Infections refer to INFECTIONS, Antibacterials, pa			
5-Alpha Reductase Inhibitors			
FINASTERIDE - Special Authority see SA0928 on the next page			
* Tab 5 mg	4.81	100	✓ <u>Ricit</u>

	GENITO-URI	NARY SYSTEM	
Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
¢ .	Por 🧹	Manufacturor	

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both.

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Fither:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy Tamsulosin-Rex

100

► SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

OXYBUTYNIN * Tab 5 mg	500	 Apo-Oxybutynin
* Oral liq 5 mg per 5 ml	473 ml	✓ Apo-Oxybutynin
POTASSIUM CITRATE		
Oral liq 3 mmol per ml – Special Authority see SA1083 below –		
Retail pharmacy31.80	200 ml OP	 Biomed

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

1 The patient has recurrent calcium oxalate urolithiasis: and

2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TABTBATE

* Grans eff 4 g sachets	2.34 2	28 🗸	Ural
SOLIFENACIN SUCCINATE			
Tab 5 mg	3.00 3	30 🗸	Solifenacin Mylan
Tab 10 mg	5.50 3	30 🗸	Solifenacin Mylan
TOLTERODINE - Special Authority see SA1272 below - Retail pha	rmacy		
Tab 1 mg	14.56	56 🖌	Arrow-Tolterodine
Tab 2 mg	14.56	56 🗸	Arrow-Tolterodine

► SA1272 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where patient has overactive bladder and a documented intolerance of, or is non-responsive to oxybutynin.

	Subsidy (Manufacturer's Pr \$	· _	1
Detection of Substances in Urine			
ORTHO-TOLIDINE			
* Compound diagnostic sticks	7.50	50 test OP	
	(8.25)		Hemastix
TETRABROMOPHENOL			
* Blue diagnostic strips	7.02	100 test OP	
- ·	(13.92)		Albustix

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully sidised	Brand or Generic Manufacturer
Calcium Homeostasis				
CALCITONIN				
* Inj 100 iu per ml, 1 ml ampoule		5	✓ M	iacalcic
CINACALCET – Special Authority see SA1618 below – Retail p Tab 30 mg – Wastage claimable	bharmacy	28	✓ S	ensipar
► SA1618 Special Authority for Subsidy Initial application only from a nephrologist or endocrinologist. following criteria: Either:	Approvals valid for 6 m	onths for	applicat	ions meeting the
1 All of the following:				
 1.1 The patient has been diagnosed with a parathyro 1.2 The patient has persistent hypercalcaemia (serur first-line treatments including sodium thiosulfate (1.3 The patient is symptomatic; or 2 All of the following: 	n calcium greater than	or equal t		
 2.1 The patient has been diagnosed with calciphylaxi 2.2 The patient has symptomatic (e.g. painful skin ul 3 mmol/L); and 	•		, ·	eater than or equal to
2.3 The patient's condition has not responded to prev thiosulfate.	rious first-line treatment	ts includir	ıg bisph	osphonates and sodium
Renewal only from a nephrologist or endocrinologist. Approval meeting the following criteria: Both:	s valid without further re	enewal ur	less no	tified for applications
 The patient's serum calcium level has fallen to < 3mmol/ The patient has experienced clinically significant sympto 				
Note: This does not include parathyroid adenomas unless these	e have become maligna	ant.		
ZOLEDRONIC ACID				
Inj 4 mg per 5 ml, vial – Special Authority see SA1687 belo Retail pharmacy		1	✓ Z	oledronic acid Mylan
	(550.00)		Z	ometa
Zoledronic acid Mylan to be Sole Supply on 1 August 2 (Zometa Inj 4 mg per 5 ml, vial to be delisted 1 August 2019)	()			
■ SA1687 Special Authority for Subsidy Initial application — (bone metastases) only from an oncolog without further renewal unless notified for applications meeting to Any of the following:		alliative ca	are spec	sialist. Approvals valid
 Patient has hypercalcaemia of malignancy; or Both: 				
2.1 Patient has bone metastases or involvement; and2.2 Patient has severe bone pain resistant to standar		or		
3 Both:				
3.1 Patient has bone metastases or involvement; and3.2 Patient is at risk of skeletal-related events pathole surgery to bone.		ord comp	ression,	radiation to bone or

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

(Ma	Subsidy nufacturer's Price)	Subsid	Fully dised	Brand or Generic
· · · · · · · · · · · · · · · · · · ·	\$	Per	1	Manufacturer

continued...

Initial application — (early breast cancer) only from an oncologist or medical practitioner on the recommendation of a oncologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

1 Treatment to be used as adjuvant therapy for early breast cancer; and

- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.

Corticosteroids and Related Agents for Systemic Use		
BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE	=	
* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	- 5	
(36.96)		Celestone
		Chronodose
DEXAMETHASONE		
* Tab 0.5 mg – Retail pharmacy-Specialist0.99	30	 Dexmethsone
Up to 60 tab available on a PSO * Tab 4 mg – Retail pharmacy-Specialist1.90	30	✓ Dexmethsone
Up to 30 tab available on a PSO	30	Dexmetrisone
Oral lig 1 mg per ml – Retail pharmacy-Specialist	25 ml OP	 Biomed
Oral lig prescriptions:	20 01	
1) Must be written by a Paediatrician or Paediatric Cardiologist; or		
2) On the recommendation of a Paediatrician or Paediatric Cardiologis	t.	
DEXAMETHASONE PHOSPHATE		
Dexamethasone phosphate injection will not be funded for oral use.		
* Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO 14.19	10	 Max Health
* Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO25.18	10	 Max Health
FLUDROCORTISONE ACETATE		_
* Tab 100 mcg14.32	100	 Florinef
HYDROCORTISONE		
* Tab 5 mg	100	✓ <u>Douglas</u>
* Tab 20 mg	100 1	 ✓ <u>Douglas</u> ✓ Solu-Cortef
 * Inj 100 mg vial5.30 a) Up to 5 inj available on a PSO 	I	• Sold-Corter
b) Only on a PSO		
METHYLPREDNISOLONE – Retail pharmacy-Specialist		
* Tab 4 mg	100	✓ Medrol
* Tab 100 mg	20	✓ Medrol
METHYLPREDNISOLONE (AS SODIUM SUCCINATE) – Retail pharmacy-Specia	alist	
Inj 40 mg vial	1	Solu-Medrol-Act-
, ,		O-Vial
Inj 125 mg vial28.90	1	✓ <u>Solu-Medrol-Act-</u>
		<u>O-Vial</u>
Inj 500 mg vial22.78	1	Solu-Medrol-Act-
		O-Vial
1-14		Costa Madual
Inj 1 g vial27.83	1	✓ <u>Solu-Medrol</u>

78

(\$29) Unapproved medicine supplied under Section 29

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
METHYLPREDNISOLONE ACETATE	•			
Inj 40 mg per ml, 1 ml vial	44 40	5	1	Depo-Medrol
		5	•	
PREDNISOLONE * Oral liq 5 mg per ml – Up to 30 ml available on a PSO	6.00 2	0 ml O		Redipred
Restricted to children under 12 years of age.			/r •	neulpieu
PREDNISONE				
* Tab 1 mg	10.69	500	1	Apo-Prednisone
* Tab 2.5 mg		500		Apo-Prednisone
 * Tab 5 mg – Up to 30 tab available on a PSO 		500		Apo-Prednisone
* Tab 20 mg		500		Apo-Prednisone
TETRACOSACTRIN				
* Inj 250 mcg per ml, 1 ml ampoule	75.00	1	1	AU Synacthen
			-	S29 S29
			1	Synacthen
				Synacthen S29 S29
* Inj 1 mg per ml, 1 ml ampoule	600.00	1		Synacthen Depot
				Synacthene
			-	Retard S29
(Synacthen S29 s29 Inj 250 mcg per ml, 1 ml ampoule to be del	listed 1 January 202	0)		
TRIAMCINOLONE ACETONIDE	isicu i bandary 202	0)		
Inj 10 mg per ml, 1 ml ampoule	20.90	F	1	Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule		5 5		Kenacort-A 40
		5	•	Kenacon-A 40
Sex Hormones Non Contraceptive				
Androgen Agonists and Antagonists				
CYPROTERONE ACETATE – Retail pharmacy-Specialist				
Tab 50 mg		50	1	Siterone
Tab 100 mg		50	✓	Siterone
TESTOSTERONE				
Patch 5 mg per day		30	1	Androderm
TESTOSTERONE CIPIONATE – Retail pharmacy-Specialist				
Inj 100 mg per ml, 10 ml vial	76.50	1	1	Depo-Testosterone
				Sopo restorerone
TESTOSTERONE ESTERS – Retail pharmacy-Specialist Inj 250 mg per ml, 1 ml	12.08	1	1	Sustanon Ampoules
		I	•	Sustantin Ampoules
TESTOSTERONE UNDECANOATE - Retail pharmacy-Specialis		00		An defet Testerson
Cap 40 mg		60		Andriol Testocaps
Inj 250 mg per ml, 4 ml vial		1	•	Reandron 1000

Hormone Replacement Therapy - Systemic

Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

	Subsidy (Manufacturer's Pr \$	rice) Sub Per	Fully sidised	Brand or Generic Manufacturer
Oestrogens				
DESTRADIOL – See prescribing guideline on the previous pag	9			
₭ Tab 1 mg		28 OP		
-	(11.10)		Est	rofem
₭ Tab 2 mg		28 OP	_	
K. Datak 05 manual and	(11.10)	•		rofem
✤ Patch 25 mcg per day	6.12	8	✓ <u>Es</u>	tradot
 a) No more than 2 patch per week b) Only on a programination 				
b) Only on a prescription ₭ Patch 50 mcg per day	7 04	8	🖌 Fe	tradot 50 mcg
a) No more than 2 patch per week		0	• []	induction mog
b) Only on a prescription				
It Patch 75 mcg per day	7.91	8	🖌 Es	tradot
a) No more than 2 patch per week				
b) Only on a prescription				
Patch 100 mcg per day	7.91	8	✓ <u>Es</u>	tradot
a) No more than 2 patch per week				
 b) Only on a prescription 				
DESTRADIOL VALERATE – See prescribing guideline on the p	previous page			
k Tab 1 mg		84		ogynova
₭ Tab 2 mg	12.36	84	✓ <u>Pro</u>	ogynova
DESTROGENS – See prescribing guideline on the previous pa				
Conjugated, equine tab 300 mcg	3.01	28		
	(13.50)		Pre	emarin
Conjugated, equine tab 625 mcg	(· · · · · · · ·	28	Dre	morin
	(13.50)		Pre	emarin
Progestogens				
IEDROXYPROGESTERONE ACETATE - See prescribing gui	deline on the prev	vious page		
🖌 Tab 2.5 mg	3.75	30	✓ <u>Pro</u>	overa
₭ Tab 5 mg		100	✓ <u>Pro</u>	
₭ Tab 10 mg	7.15	30	✓ <u>Pro</u>	overa
Progestogen and Oestrogen Combined Prepar	ations			
DESTRADIOL WITH NORETHISTERONE – See prescribing gr	uideline on the pre	vious page		
K Tab 1 mg with 0.5 mg norethisterone acetate		28 OP		
	(18.10)		Klie	ovance
K Tab 2 mg with 1 mg norethisterone acetate		28 OP		
	(18.10)		Klie	ogest
₭ Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg				
oestradiol tab (12) and 1 mg oestradiol tab (6)		28 OP	T	
	(18.10)		I ris	sequens
Other Oestrogen Preparations				
THINYLOESTRADIOL				
Tab 10 mcg	17.60	100	🖌 N7	Medical and
- 140 TO HOY		100		Scientific
			-	

_					
		Subsidy	Cul	Fully	Brand or
		(Manufacturer's Price) \$	Per	sidised ✓	Generic Manufacturer
OES	STRIOL				
	Tab 2 mg	7.00	30	✓ 0	ovestin
01	ther Progestogen Preparations				
LEV	ONORGESTREL				
*	Intra-uterine system 20 mcg per day - Special Authority see				
	SA1608 below – Retail pharmacy		1	✓ <u>N</u>	lirena
	A1608 Special Authority for Subsidy				
app	al application — (No previous use) only from a relevant sp lications meeting the following criteria: f the following:	pecialist or general pra	actitioner	. Approv	vals valid for 6 months for
	 The patient has a clinical diagnosis of heavy menstrual bl The patient has failed to respond to or is unable to tolerat Menstrual Bleeding Guidelines; and Either: 		narmaceu	utical the	rapies as per the Heavy
	3.1 serum ferritin level $<$ 16 mcg/l (within the last 12 r 3.2 haemoglobin level $<$ 120 g/l.	nonths); or			
Ren	e: Applications are not to be made for use in patients as con newal only from a relevant specialist or general practitioner. A wing criteria: n:				
200	1 Either:				
	1.1 Patient demonstrated clinical improvement of heav1.2 Previous insertion was removed or expelled within2 Applicant to state date of the previous insertion.	, ,	-		
	DROXYPROGESTERONE ACETATE				
	Tab 100 mg – Retail pharmacy-Specialist	101.00	100	✓ P	rovera HD
	RETHISTERONE			-	
-	Tab 5 mg – Up to 30 tab available on a PSO		100	✓ P	rimolut N
	DGESTERONE				
	Cap 100 mg – Special Authority see SA1609 below – Retail pharmacy		30	~ 11	trogestan
	A1609 Special Authority for Subsidy		00		a ogotan
Initi	al application only from an obstetrician or gynaecologist. A wing criteria:	pprovals valid for 12 n	nonths fo	r applica	tions meeting the
	 For the prevention of pre-term labour*; and Either: 				
	2.1 The patient has a short cervix on ultrasound (defin2.2 The patient has a history of pre-term birth at less t		to 28 wee	eks); or	
	newal only from an obstetrician or gynaecologist. Approvals of the following:	valid for 12 months for	r applicat	ions me	eting the following criteria:
	1 For the prevention of pre-term labour*; and				
	2 Treatment is required for second or subsequent pregnand3 Either:	cy; and			
	3.1 The patient has a short cervix on ultrasound (defin3.2 The patient has a history of pre-term birth at less t		to 28 wee	eks); or	

Note: Indications marked with * are unapproved indications.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Thyroid and Antithyroid Agents				
CARBIMAZOLE * Tab 5 mg	10.80	100		FT Carbimazole S29 eo-Mercazole
LEVOTHYROXINE				
 * Tab 25 mcg * Tab 50 mcg * Tab 100 mcg PROPYLTHIOURACIL – Special Authority see SA1199 below – Propylthiouracil is not recommended for patients under the a 		90 28 90 1,000 28 90 1,000	✓ M ✓ S ✓ E ✓ M ✓ S ✓ E	ynthroid Iercury Pharma ynthroid Itroxin Iercury Pharma ynthroid Itroxin
treatments are contraindicated.	25.00	100		TU 529
Tab 50 mg SA1199 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals vali Both: 1 The patient has hyperthyroidism; and 2 The patient is intolerant of carbimazole or carbimazole is	d for 2 years for appli	100 cation	-	

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) – Special Authority see SA	1629 below – Retail phar	nacy	
*	Inj 5 mg cartridge		1	 Omnitrope
*	Inj 10 mg cartridge	69.75	1	 Omnitrope
*	Inj 15 mg cartridge	104.63	1	 Omnitrope

► SA1629 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: Either:

1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or

2 All of the following:

- Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
- 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

children who are 5 years or older, GH testing with sex steroid priming is required; and

- 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
- 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under ; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application - (short stature without growth hormone deficiency) only from a paediatric endocrinologist or

endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
`\$	Per	✓	Manufacturer

continued...

- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and

6 Either:

- 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
- 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight eximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:

5.1 Both:

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise		
\$	Per •	Manufacturer	

continued...

- 5.1.1 The patient is aged two years or older; and
- 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
- 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ± 1 SD of the mean of the normal range for age and sex; and

Subsidy	Fully		Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or 2 All of the following:

- 2.1 The patient has been treated with somatropin for more than 12 months; and
- 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
- 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
- 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients.

GnRH Analogues

GOSERELIN			
Implant 3.6 mg, syringe	66.48	1	 Zoladex
Implant 10.8 mg, syringe	177.50	1	 Zoladex

LEUPRORELIN

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

Inj 3.75 mg prefilled dual chamber syringe – Higher subsid	ly of		
\$221.60 per 1 inj with Endorsement		1	
	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe - Higher subs	idy		
of \$591.68 per 1 inj with Endorsement		1	
	(591.68)		Lucrin Depot 3-month

Vasopressin Agonists

Tab 100 mcg – Special Authority see SA1401 below – Retail pharmacy Tab 200 mcg – Special Authority see SA1401 below – Retail	25.00	30	✓ <u>Minirin</u>
pharmacy Nasal drops 100 mcg per ml – Retail pharmacy-Specialist Nasal spray 10 mcg per dose – Retail pharmacy-Specialist	39.03	30 2.5 ml OP 6 ml OP	 ✓ <u>Minirin</u> ✓ Minirin ✓ <u>Desmopressin</u> <u>PH&T</u>
Inj 4 mcg per ml, 1 ml – Special Authority see SA1401 below – Retail pharmacy	67.18	10	✓ Minirin

► SA1401 Special Authority for Subsidy

Initial application — (Desmopressin tablets for Nocturnal enuresis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has primary nocturnal enuresis; and
- 2 The nasal forms of desmopressin are contraindicated; and
- 3 An enuresis alarm is contraindicated.

Initial application — (Desmopressin tablets for Diabetes insipidus) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

 Subsidy	Fully Subsidised		Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	V	Manufacturer

continued...

1 The patient has cranial diabetes insipidus; and

2 The nasal forms of desmopressin are contraindicated.

Renewal — (Desmopressin tablets) from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from the treatment.

Initial application — (Desmopressin injection) only from a relevant specialist. Approvals valid for 2 years where the patient cannot use desmopressin nasal spray or nasal drops.

Renewal — (Desmopressin injection) only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Other Endocrine Agents

CABERGOLINE

		Tab 0.5 mg – Maximum of 2 tab per prescription; can be
 Dostinex 	2	waived by Special Authority see SA1370 below
 Dostinex 	8	15.20

⇒SA1370 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

1 pathological hyperprolactinemia; or

2 acromegaly*.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment. Note: Indication marked with * is an unapproved indication.

CLOMIFENE CITRATE

Tab 50 mg	84 1	0 🗸	Mylan Clomiphen S29
DANAZOL			
Cap 100 mg	.33 10	00 🗸	Azol
Cap 200 mg	.83 10	00 🗸	Azol
METYRAPONE			
Cap 250 mg – Retail pharmacy-Specialist	.00 5	0 🖌	Metopirone

	Subsidy (Manufacturer's Price) \$	Subsic Per	Fully lised	Brand or Generic Manufacturer
Anthelmintics				
ALBENDAZOLE - Special Authority see SA1318 below - Retai	l pharmacy			
Tab 400 mg		60	✓ E	skazole S29
► SA1318 Special Authority for Subsidy Initial application only from an infectious disease specialist or of patient has hydatids.	Ũ			
Renewal only from an infectious disease specialist or clinical mi remains appropriate and the patient is benefitting from the treatm		ls valid for 6	6 mont	ths where the treatment
MEBENDAZOLE – Only on a prescription				
Tab 100 mg Oral lig 100 mg per 5 ml		24 15 ml	✓ D	0e-Worm
	(7.17)	15 111	۷	/ermox
PRAZIQUANTEL Tab 600 mg		8	✓ В	Biltricide
Antibacterials				
a) For topical antibacterials, refer to DERMATOLOGICALS, page b) For anti-infective eye preparations, refer to SENSORY ORG.				
Cephalosporins and Cephamycins				
CEFACLOR MONOHYDRATE				
Cap 250 mg Grans for oral lig 125 mg per 5 ml – Wastage claimable		100 100 ml		<u>lanbaxy-Cefaclor</u> lanbaxy-Cefaclor
CEFALEXIN			• <u>n</u>	andaky-Celacion
Cap 250 mg		20	√ c	ephalexin ABM
Cap 500 mg		20	_	Cephalexin ABM
Grans for oral liq 25 mg per ml - Wastage claimable	8.75 ·	100 ml	 C 	efalexin Sandoz
Note: Cefalexin grans for oral liq will not be funded in a				
Grans for oral liq 50 mg per ml – Wastage claimable Note: Cefalexin grans for oral lig will not be funded in a		100 ml		Cefalexin Sandoz
CEFAZOLIN – Subsidy by endorsement	anounts more than 14	uays ireairi	ient p	er dispensing.
Only if prescribed for dialysis or cellulitis in accordance with accordingly.	a DHB approved prot	ocol and the	e pres	cription is endorsed
Inj 500 mg vial		5	🗸 A	FT
Inj 1 g vial		5	✓ <u>A</u>	
CEFTRIAXONE – Subsidy by endorsement				
 a) Up to 10 inj available on a PSO b) Subsidised only if prescribed for a dialysis or cystic fibro pelvic inflammatory disease, or the treatment of suspect endorsed accordingly. 	ed meningococcal dise			
Inj 500 mg vial		1	_	DEVA
Inj 1 g vial	0.84	1	✓ <u>D</u>	DEVA
CEFUROXIME AXETIL – Subsidy by endorsement	accription is andoread	accordingly		
Only if prescribed for prophylaxis of endocarditis and the pro Tab 250 mg		accordingly 50		innat
			-	

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Macrolides				
AZITHROMYCIN – Maximum of 5 days treatment per prescription A maximum of 24 months of azithromycin treatment for non- Authority.	· ·		,	

Tab 250 mg8.19	30	Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO0.93		 Apo-Azithromycin
Grans for oral liq 200 mg per 5 ml (40 mg per ml) – Wastage		
claimable14.38	15 ml	 Zithromax

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.
- Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).
- The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN -	- Maximum of 500 mg per prescription; can be w	aived by Special	Authority s	ee SA1131 below
Tab 250 mg		3.98	14	 Apo-Clarithromvcin

Tab 250 mg	3.98	14	Apo-Cla
Grans for oral lig 250 mg per 5 ml - Wastage claimable	23.12	50 ml	Klacid

► SA1131 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria: Either:

	Subsidy (Manufacturer's Price) \$) ; Per	Fully Subsidised	Brand or Generic Manufacturer
pntinued				
1 Atypical mycobacterial infection; or				
2 Mycobacterium tuberculosis infection where there is drug	g-resistance or intoler	ance to	standard	pharmaceutical agents.
enewal — (Mycobacterial infections) only from a respirator				
pprovals valid for 2 years where the treatment remains approp	priate and the patient i	s benef	iting from	treatment.
RYTHROMYCIN ETHYL SUCCINATE				
Tab 400 mg		100	✓	E-Mycin
 a) Up to 20 tab available on a PSO 				
b) Up to 2 x the maximum PSO quantity for RFPP				
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	✓	E-Mycin
 a) Up to 300 ml available on a PSO 				
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	✓	E-Mycin
 a) Up to 200 ml available on a PSO 				
b) Wastage claimable				
RYTHROMYCIN LACTOBIONATE				
Inj 1 g	16.00	1	✓	Erythrocin IV
RYTHROMYCIN STEARATE				
Tab 250 mg - Up to 30 tab available on a PSO		100		
	(22.29)		I	ERA
Tab 500 mg		100		
	(44.58)		I	ERA
OXITHROMYCIN				
Tab disp 50 mg	7.19	10	✓ 1	Rulide D
Restricted to children under 12 years of age.				
Tab 150 mg	8.28	50	 Image: A second s	Arrow-
				Roxithromycin
Arrow-Roxithromycin to be Sole Supply on 1 September	ar 2019			
Tab 300 mg		50	1	Arrow-
140 000 mg		50		Roxithromycin

Arrow-Roxithromycin to be Sole Supply on 1 September 2019

	Subsidy (Manufacturer's Price	e) 5	Fully Subsidised	
	\$	Per	1	Manufacturer
Penicillins				
AMOXICILLIN				
Cap 250 mg	14.97	500	✓	Apo-Amoxi
 a) Up to 30 cap available on a PSO 				
b) Up to 10 x the maximum PSO quantity for RFPP				
Cap 500 mg	16.75	500	~	Apo-Amoxi
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	1.20	100 ml	~	Alphamox 125
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	1.31	100 ml	~	Alphamox 250
a) Up to 300 ml available on a PSO				
 b) Up to 10 x the maximum PSO quantity for RFPP b) Western advised to 				
c) Wastage claimable	10.67	10		lhiemey
Inj 250 mg vial Inj 500 mg vial		10 10		Ibiamox Ibiamox
Inj 1 g vial – Up to 5 inj available on a PSO		10		Ibiamox
		10	•	IDIAIIIOX
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab	1 00	00		Auguantia
available on a PSO		20	•	Augmentin
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 r per ml		100 ml	1	Augmentin
a) Up to 200 ml available on a PSO		100 111	•	Auginentin
b) Wastage claimable				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 r	na			
per ml – Up to 200 ml available on a PSO		00 ml O	Р 🖌	Curam
BENZATHINE BENZYLPENICILLIN		00 111 0		ourum
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj available on a PSO	244.02	10		Diaillin I A
		10	•	Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]				• •
Inj 600 mg (1 million units) vial – Up to 5 inj available on a PS	50 10.35	10	~	Sandoz
FLUCLOXACILLIN				
Cap 250 mg – Up to 30 cap available on a PSO		250		Staphlex
Cap 500 mg		500	-	Staphlex
Grans for oral liq 25 mg per ml	2.29	100 ml	•	<u>AFT</u>
a) Up to 200 ml available on a PSO				
b) Wastage claimable	2 60	100 ml		AET
Grans for oral liq 50 mg per ml	3.00			<u>AFT</u>
 a) Up to 200 ml available on a PSO b) Wastage claimable 				
Inj 250 mg vial	9 00	10	1	Flucloxin
Inj 500 mg vial		10		Flucloxin
Inj 1 g vial – Up to 5 inj available on a PSO		5		Flucil
,		-		

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price \$		Fully Brand or idised Generic ✓ Manufacturer
PHENOXYMETHYLPENICILLIN (PENICILLIN V) Cap 250 mg – Up to 30 cap available on a PSO Cap 500 mg a) Up to 20 cap available on a PSO		50 50	 ✓ <u>Cilicaine VK</u> ✓ <u>Cilicaine VK</u>
 b) Up to 2 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml a) Up to 200 ml available on a PSO 	1.48	100 ml	✓ <u>AFT</u>
 b) Wastage claimable Grans for oral liq 250 mg per 5 ml a) Up to 300 ml available on a PSO b) Up to 2 x the maximum PSO quantity for RFPP c) Wastage claimable 	1.58	100 ml	✓ <u>AFT</u>
PROCAINE PENICILLIN Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO		5	✓ <u>Cilicaine</u>
Tetracyclines			
DOXYCYCLINE * Tab 50 mg – Up to 30 tab available on a PSO	2.90 (6.00)	30	Doxy-50
* Tab 100 mg – Up to 30 tab available on a PSO (Doxy-50 Tab 50 mg to be delisted 1 January 2020)		500	✓ Doxine
MINOCYCLINE HYDROCHLORIDE * Tab 50 mg – Additional subsidy by Special Authority see			
SA1355 below – Retail pharmacy	(12.05)	60 100	Mino-tabs
· · ·	(52.04)	100	Minomycin
■ SA1355 Special Authority for Manufacturers Price Initial application from any relevant practitioner. Approvals valid rosacea.	d without further ren	ewal unless	notified where the patient has
TETRACYCLINE – Special Authority see SA1332 below – Retail Cap 500 mg		30	 Tetracyclin

► SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and

2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Wolff S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Other Antibiotics				
For topical antibiotics, refer to DERMATOLOGICALS, page 60				
CIPROFLOXACIN				
Recommended for patients with any of the following:				
 i) microbiologically confirmed and clinically significant pse ii) prostatitis; or iii) pyelonephritis; or 	eudomonas infection;	or		
iv) gonorrhoea.				
Tab 250 mg – Up to 5 tab available on a PSO		28		ipflox
Tab 500 mg – Up to 5 tab available on a PSO		28		ipflox
Tab 750 mg	3.15	28	✓ <u>U</u>	ipflox
CLINDAMYCIN Cap hydrochloride 150 mg – Maximum of 4 cap per				
prescription; can be waived by endorsement - Retail				
pharmacy - Specialist	4.10	16	✓ C	lindamycin ABM
Inj phosphate 150 mg per ml, 4 ml ampoule – Retail pharmacy-Specialist	65.00	10	✓ <u>D</u>	alacin C
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist - S				
Only if prescribed for dialysis or cystic fibrosis patient and the Inj 150 mg		rsed a		olistin-Link
GENTAMICIN SULPHATE				
Inj 10 mg per ml, 1 ml ampoule – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient c endorsed accordingly.		5 / trac		BL Gentamicin d the prescription is
Inj 40 mg per ml, 2 ml ampoule – Subsidy by endorsement	6.00	10	🗸 P	fizer
	30.00	50	🗸 P	
Only if prescribed for a dialysis or cystic fibrosis patient c endorsed accordingly.	or complicated urinary	/ trac	t infection an	d the prescription is
MOXIFLOXACIN - Special Authority see SA1740 below - Retail	pharmacy			
No patient co-payment payable Tab 400 mg	52.00	5	۷ ۸	velox
SA1740 Special Authority for Subsidy		5	• •	VEIUX
Initial application — (Tuberculosis) only from a respiratory spe for applications meeting the following criteria:	ecialist or infectious d	iseas	e specialist.	Approvals valid for 1 year
Any of the following: 1 Both:				
1.1 Active tuberculosis*; and				
1.2 Any of the following:				
1.2.1 Documented resistance to one or more first				
1.2.2 Suspected resistance to one or more first-lin	,			
area with known resistance), as part of regi 1.2.3 Impaired visual acuity (considered to preclu			nu-ine agen	15, 01
1.2.4 Significant pre-existing liver disease or hepa	atotoxicity from tubero	culos		
1.2.5 Significant documented intolerance and/or s	side effects following	a rea	sonable trial	of first-line medications;
or				

continued...

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully	Brand or
	(Manufacturer's Price \$) Si Per	ubsidised	Brand or Generic Manufacturer
continued				
2 Mycobacterium avium-intracellulare complex not respondin3 Patient is under five years of age and has had close contact				
Note: Indications marked with * are unapproved indications. Renewal only from a respiratory specialist or infectious disease sp	pecialist. Approval	s valid fo	r 1 year w	here the treatment
remains appropriate and the patient is benefiting from treatment. Initial application — (Mycoplasma genitalium) only from a sexu sexual health specialist. Approvals valid for 1 month for application All of the following:				the recommendation of a
All of the following: 1 Has nucleic acid amplification test (NAAT) confirmed Myco 2 Either:	plasma genitalium'	* and is s	ymptoma	tic; and
2.1 Has tried and failed to clear infection using azithrom2.2 Has laboratory confirmed azithromycin resistance; a				
3 Treatment is only for 7 days.				
Initial application — (Penetrating eye injury) only from an opht requires prophylaxis following a penetrating eye injury and treatme Note: Indications marked with * are unapproved indications.			d for 1 mo	onth where the patient
PAROMOMYCIN - Special Authority see SA1689 below - Retail	pharmacy			
Cap 250 mg	126.00	16	✓ F	lumatin S29
SA1689 Special Authority for Subsidy Initial application only from an infectious disease specialist, clinic month for applications meeting the following criteria: Either:	cal microbiologist o	r gastroe	nterologis	t. Approvals valid for 1
1 Patient has confirmed cryptosporidium infection; or 2 For the eradication of Entamoeba histolyica carriage.				
Renewal only from an infectious disease specialist, clinical microb applications meeting the following criteria: Either:	iologist or gastroer	nterologis	st. Approv	vals valid for 1 month for
 Patient has confirmed cryptosporidium infection; or For the eradication of Entamoeba histolyica carriage. 				
PYRIMETHAMINE - Special Authority see SA1328 below - Retain	il pharmacy			
Tab 25 mg	26.14	30	🗸 [Daraprim S29
	36.95	50	✓ [Daraprim S29
SA1328 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following:	without further ren	ewal unle	ess notifie	d for applications meeting
 For the treatment of toxoplasmosis in patients with HIV for For pregnant patients for the term of the pregnancy; or For infants with congenital toxoplasmosis until 12 months contact to the term of the pregnancy. 	•	ns; or		
SODIUM FUSIDATE [FUSIDIC ACID] Tab 250 mg – Retail pharmacy-Specialist Prescriptions must be written by, or on the recommendati		12 disease	_	ucidin or a clinical microbiologist
SULFADIAZINE SODIUM – Special Authority see SA1331 on the				
Tab 500 mg	543.20	56	✓ V	Vockhardt S29

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
SA1331 Special Authority for Subsidy itial application from any relevant practitioner. Approvals valid	d without further rer	newal ur	nless notifi	ed for applications meetir
e following criteria: ny of the following:				
 For the treatment of toxoplasmosis in patients with HIV for 	r a period of 3 mont	hs: or		
2 For pregnant patients for the term of the pregnancy; or3 For infants with congenital toxoplasmosis until 12 months				
OBRAMYCIN				
Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement	15.00	5	1	Tobramycin Mylan
Only if prescribed for dialysis or cystic fibrosis patient an	d the prescription is	endors	ed accord	ingly.
Solution for inhalation 60 mg per ml, 5 ml – Subsidy by				
endorsement	2,200.00	56 dos	e 🗸	ТОВІ
a) Wastage claimableb) Only if prescribed for a cystic fibrosis patient and the	prescription is endo	orsed ad	cordingly.	
RIMETHOPRIM				
Tab 300 mg – Up to 30 tab available on a PSO		50	~	TMP
RIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX				
Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - L	Jp	500		Tuland
to 30 tab available on a PSO Oral lig 8 mg sulphamethoxazole 40 mg per ml – Up to 200		500	~	Trisul
 Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to 200 available on a PSO 		100 m	· •	Deprim
ANCOMYCIN – Subsidy by endorsement		100 111	•	Deprim
Only if prescribed for a dialysis or cystic fibrosis patient or for	r prophylaxis of end	ocarditi	s or for tre	atment of Clostridium
difficile following metronidazole failure and the prescription is	endorsed accordin	igly.		
Inj 500 mg vial		1	✓	Mylan
A 11/2 1				
Antifungals				
For topical antifungals refer to DERMATOLOGICALS, page 6	1			
For topical antifungals refer to GENITO URINARY, page 74				
LUCONAZOLE				
Cap 50 mg – Retail pharmacy-Specialist	2.09	28	1	Mylan
Cap 150 mg - Subsidy by endorsement		1	✓	Mylan
a) Maximum of 1 cap per prescription; can be waived by				
b) Patient has vaginal candida albicans and the practition				
not recommended and the prescription is endorsed a	accordingly; can be	waived	by endorse	ement - Retail pharmacy
Specialist. Cap 200 mg – Retail pharmacy-Specialist	5 08	28	1	Mylan
Powder for oral suspension 10 mg per ml – Special Authorit		20	,	in y lait
a smash isi shu suspension ne mg per mi opecial Authonit		35 ml	1	Diflucan S29 S29
see SA1359 below - Retail pharmacy				
see SA1359 below - Retail pharmacy	98.50		~	Diflucan
see SA1359 below – Retail pharmacy Wastage claimable •SA1359 Special Authority for Subsidy			~	Diflucan

meeting the following criteria: Both:

continued...

	Subsidy (Manufacturer's F \$	Price) Per	Fully Subsidised	Brand or Generic Manufacturer
continued				
 Patient requires prophylaxis for, or treatment of system Patient is unable to swallow capsules. 	mic candidiasis; and			
Initial application — (Immunocompromised) from any rele meeting the following criteria: All of the following:	evant practitioner. A	pprovals va	alid for 6 m	onths for applications
 Patient is immunocompromised; and Patient is at moderate to high risk of invasive fungal ir Patient is unable to swallow capsules. 	nfection; and			
Renewal — (Systemic candidiasis) from any relevant prac following criteria: Both:	titioner. Approvals v	alid for 6 w	veeks for ap	oplications meeting the
 Patient requires prophylaxis for, or treatment of system Patient is unable to swallow capsules. 	mic candidiasis; and			
Renewal — (Immunocompromised) from any relevant praci following criteria: All of the following:	ctitioner. Approvals	valid for 6 r	months for	applications meeting the
 Patient remains immunocompromised; and Patient remains at moderate to high risk of invasive fu Patient is unable to swallow capsules. 	ingal infection; and			
TRACONAZOLE				
TRACONAZOLE Cap 100 mg – Subsidy by endorsement Funded for tinea vesicolor where topical treatment h mycology, or for tinea unguium where terbinafine ha terbinafine and diagnosis has been confirmed by my by endorsement - Retail pharmacy - Specialist Spec clinical immunologist or dermatologist	as not been success s not been successfu cology and the prese	ul in eradica cription is e	gnosis has ation or the ndorsed ac	patient is intolerant to cordingly. Can be waive
Cap 100 mg – Subsidy by endorsement Funded for tinea vesicolor where topical treatment h mycology, or for tinea unguium where terbinafine ha terbinafine and diagnosis has been confirmed by my by endorsement - Retail pharmacy - Specialist Spec clinical immunologist or dermatologist. Oral liq 10 mg per ml – Special Authority see SA1322 be	as not been success s not been successfu cology and the preso ialist must be an infe elow –	ful and dia ul in eradic cription is e	gnosis has ation or the indorsed ac ase physici	been confirmed by patient is intolerant to cordingly. Can be waive an, clinical microbiologist
Cap 100 mg – Subsidy by endorsement Funded for tinea vesicolor where topical treatment h mycology, or for tinea unguium where terbinafine ha terbinafine and diagnosis has been confirmed by my by endorsement - Retail pharmacy - Specialist Spec clinical immunologist or dermatologist. Oral liq 10 mg per ml – Special Authority see SA1322 be Retail pharmacy	as not been success s not been successfi roology and the preso ialist must be an infe elow – 	ful and dia ul in eradica cription is e ctious dise 150 ml C st, clinical i robiologist	gnosis has ation or the ndorsed ac ase physici OP ✓ S mmunologi or clinical ir	been confirmed by patient is intolerant to coordingly. Can be waive ian, clinical microbiologist Sporanox st or any relevant nmunologist. Approvals
Cap 100 mg – Subsidy by endorsement Funded for tinea vesicolor where topical treatment h mycology, or for tinea unguium where terbinafine ha terbinafine and diagnosis has been confirmed by my by endorsement - Retail pharmacy - Specialist Spec clinical immunologist or dermatologist. Oral liq 10 mg per ml – Special Authority see SA1322 be Retail pharmacy	as not been success s not been success icology and the preso ialist must be an infe elow – 	ful and dia ul in eradica cription is e ctious dise 150 ml C st, clinical i robiologist	gnosis has ation or the ndorsed ac ase physici OP ✓ S mmunologi or clinical ir	been confirmed by patient is intolerant to coordingly. Can be waive ian, clinical microbiologist Sporanox st or any relevant nmunologist. Approvals
Cap 100 mg – Subsidy by endorsement Funded for tinea vesicolor where topical treatment h mycology, or for tinea unguium where terbinafine ha terbinafine and diagnosis has been confirmed by my by endorsement - Retail pharmacy - Specialist Spec clinical immunologist or dermatologist. Oral liq 10 mg per ml – Special Authority see SA1322 be Retail pharmacy	as not been success s not been success icology and the prese ialist must be an infe elow – 	ful and dia ul in eradica cription is e ctious dise 150 ml C st, clinical i robiologist	gnosis has ation or the indorsed ac ase physici OP S mmunologi or clinical ir emains app L	been confirmed by patient is intolerant to cordingly. Can be waive ian, clinical microbiologist Sporanox st or any relevant nmunologist. Approvals propriate and the patient is .ink Healthcare 529
Cap 100 mg – Subsidy by endorsement Funded for tinea vesicolor where topical treatment h mycology, or for tinea unguium where terbinafine ha terbinafine and diagnosis has been confirmed by my by endorsement - Retail pharmacy - Specialist Spec- clinical immunologist or dermatologist. Oral liq 10 mg per ml – Special Authority see SA1322 be Retail pharmacy	as not been success s not been success ialist must be an infe elow – 	ful and dia ul in eradic: cription is e ctious dise 150 ml C st, clinical i robiologist treatment re 30	gnosis has ation or the indorsed ac ase physici OP S mmunologi or clinical ir emains app L	been confirmed by patient is intolerant to cordingly. Can be waive ian, clinical microbiologist Sporanox st or any relevant nmunologist. Approvals propriate and the patient is
Cap 100 mg – Subsidy by endorsement Funded for tinea vesicolor where topical treatment h mycology, or for tinea unguium where terbinafine ha terbinafine and diagnosis has been confirmed by my by endorsement - Retail pharmacy - Specialist Spec clinical immunologist or dermatologist. Oral liq 10 mg per ml – Special Authority see SA1322 be Retail pharmacy	as not been success s not been success ialist must be an infe elow – 	ful and dia ul in eradic: cription is e ctious dise 150 ml C st, clinical i robiologist treatment n 30 gist	gnosis has ation or the indorsed ac ase physici OP S mmunologi or clinical ir emains app L	been confirmed by patient is intolerant to cordingly. Can be waive ian, clinical microbiologist Sporanox st or any relevant nmunologist. Approvals propriate and the patient is .ink Healthcare 529
Cap 100 mg – Subsidy by endorsement Funded for tinea vesicolor where topical treatment h mycology, or for tinea unguium where terbinafine ha terbinafine and diagnosis has been confirmed by my by endorsement - Retail pharmacy - Specialist Spec- clinical immunologist or dermatologist. Oral liq 10 mg per ml – Special Authority see SA1322 be Retail pharmacy	as not been success s not been success icology and the preso ialist must be an infe elow – 	ful and dia ul in eradic: cription is e ctious dise 150 ml C st, clinical i robiologist treatment re 30	gnosis has ation or the ndorsed ac ase physici OP	been confirmed by patient is intolerant to cordingly. Can be waive ian, clinical microbiologist Sporanox st or any relevant nmunologist. Approvals propriate and the patient is .ink Healthcare 529
Cap 100 mg – Subsidy by endorsement Funded for tinea vesicolor where topical treatment h mycology, or for tinea unguium where terbinafine ha terbinafine and diagnosis has been confirmed by my by endorsement - Retail pharmacy - Specialist Spec clinical immunologist or dermatologist. Oral liq 10 mg per ml – Special Authority see SA1322 be Retail pharmacy	as not been success s not been success icology and the preso ialist must be an infe elow – 	ful and dia ul in eradic: cription is e ctious dise 150 ml C st, clinical i robiologist treatment n 30 gist	gnosis has ation or the ndorsed ac ase physici OP	been confirmed by patient is intolerant to cordingly. Can be waive ian, clinical microbiologist sporanox st or any relevant nmunologist. Approvals propriate and the patient is .ink Healthcare s29 lizoral s29
Cap 100 mg – Subsidy by endorsement Funded for tinea vesicolor where topical treatment h mycology, or for tinea unguium where terbinafine ha terbinafine and diagnosis has been confirmed by my by endorsement - Retail pharmacy - Specialist Spec- clinical immunologist or dermatologist. Oral liq 10 mg per ml – Special Authority see SA1322 be Retail pharmacy	as not been success s not been success vology and the prese ialist must be an infe elow – 	ful and dia ul in eradic: cription is e ctious dise 150 ml C st, clinical i robiologist treatment n 30 gist 50 50 armacy	gnosis has ation or the indorsed ac ase physici OP	been confirmed by patient is intolerant to coordingly. Can be waive ian, clinical microbiologist Sporanox st or any relevant mmunologist. Approvals propriate and the patient is ink Healthcare 529 lizoral 529
Cap 100 mg – Subsidy by endorsement Funded for tinea vesicolor where topical treatment h mycology, or for tinea unguium where terbinafine ha terbinafine and diagnosis has been confirmed by my by endorsement - Retail pharmacy - Specialist Spec- clinical immunologist or dermatologist. Oral liq 10 mg per ml – Special Authority see SA1322 be Retail pharmacy	as not been success s not been success roology and the prese ialist must be an infe elow – 	ful and dia ul in eradic: cription is e ctious dise 150 ml C st, clinical i robiologist treatment n 30 gist 50 50	gnosis has ation or the indorsed ac ase physici OP	been confirmed by patient is intolerant to cordingly. Can be waive ian, clinical microbiologis Sporanox st or any relevant mmunologist. Approvals propriate and the patient i ink Healthcare S29 lizoral S29

Subsidy (Manufacture's	Price) Subsi Per	Fully dised	Brand or Generic Manufacturer	
φ	Fei		Warlulaclurei	

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE

* Tab 250 mg	1.33	14	✓ Deolate
VORICONAZOLE - Special Authority see SA1273 below - Retail pharma	acy		
Tab 50 mg9	1.00	56	 Vttack
Tab 200 mg	0.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage			
claimable	7.00	70 ml	 Vfend

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

	Subsidy		Fully Brand or
	(Manufacturer's Price)	Subsid	
	\$	Per	 Manufacturer
Antimalarials			
PRIMAQUINE PHOSPHATE - Special Authority see SA1684 be	low – Retail pharmad	CV .	
Tab 7.5 mg	•	56	 Primacin S29
► SA1684 Special Authority for Subsidy			
Initial application only from an infectious disease specialist or cli meeting the following criteria: Both:	inical microbiologist.	Approvals	valid for 1 month for applications
1 The patient has vivax or ovale malaria; and			
2 Primaquine is to be given for a maximum of 21 days.			
Renewal only from an infectious disease specialist or clinical mice the following criteria:	robiologist. Approva	Is valid for 1	1 month for applications meeting
Both:			
 The patient has relapsed vivax or ovale malaria; and Primaquine is to be given for a maximum of 21 days. 			
Antiparasitics			
Antiprotozoals			
QUININE SULPHATE			
* Tab 300 mg	61.91	500	✓ Q 300
Antitrichomonal Agents			
METRONIDAZOLE			
Tab 200 mg – Up to 30 tab available on a PSO		100	 Trichozole
Tab 400 mg – Up to 15 tab available on a PSO		100	 Trichozole
Oral liq benzoate 200 mg per 5 ml Suppos 500 mg		100 ml 10	 ✓ FlagyI-S ✓ FlagyI
ORNIDAZOLE		10	• Tidgyi
Tab 500 mg		10	Arrow-Ornidazole
· • • • • • • •			
Antituberculotics and Antileprotics			
Note: There is no co-payment charge for all pharmaceuticals liste immigration status.	ed in the Antitubercul	lotics and A	ntileprotics group regardless of
CLOFAZIMINE – Retail pharmacy-Specialist			
a) No patient co-payment payable			
b) Prescriptions must be written by, or on the recommendation dermatologist.		isease phys	sician, clinical microbiologist or
* Cap 50 mg	442.00	100	 Lamprene S29
CYCLOSERINE – Retail pharmacy-Specialist			
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation 	on of, an infectious d	isease phys	sician, clinical microbiologist or
respiratory physician. Cap 250 mg	344.00	60	 Cyclorin S29
	1,294.50	100	✓ King S29
(King S29) Cap 250 mg to be delisted 1 November 2019)	.,_000		

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per Brand or Generic Manufacturer DAPSONE - Retail pharmacy-Specialist a) No patient co-payment payable b) b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist - Tab 25 mg
\$ Per ✓ Manufacturer DAPSONE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist Tab 25 mg 268.50 100 ✓ Dapsone Tab 100 mg 268.50 100 ✓ Dapsone ETHAMBUTOL HYDROCHLORIDE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician Tab 100 mg 85.73 100 ✓ EMB Fatol 100 Tab 400 mg 49.34 56 ✓ Myambutol 100 ISONIAZID – Retail pharmacy-Specialist a) No patient co-payment payable 49.34 56
DAPSONE - Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist Tab 25 mg 268.50 100 ✓ Dapsone Tab 100 mg 329.50 100 ✓ Dapsone ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician Tab 100 mg 85.73 100 ✓ EMB Fatol 529 Tab 400 mg 49.34 56 ✓ Myambutol 529 ISONIAZID - Retail pharmacy-Specialist a) No patient co-payment payable a) No patient co-payment payable
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist Tab 25 mg
 b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist Tab 25 mg
dermatologist Tab 25 mg Tab 100 mg Tab 100 mg Bapsone ETHAMBUTOL HYDROCHLORIDE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician Tab 100 mg Tab 100 mg Bapsone Start of the second
Tab 25 mg 268.50 100 ✓ Dapsone Tab 100 mg 329.50 100 ✓ Dapsone ETHAMBUTOL HYDROCHLORIDE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician Tab 100 mg 85.73 100 ✓ EMB Fatol S29 Tab 400 mg 49.34 56 ✓ Myambutol S29 ISONIAZID – Retail pharmacy-Specialist a) No patient co-payment payable An opatient co-payment payable
Tab 100 mg
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician Tab 100 mg
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician Tab 100 mg
 b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician Tab 100 mg B5.73 100 ✓ EMB Fatol Tab 400 mg Horman 49.34 49.34 49.34 49.34 49.34 400 mg Myambutol 100 ISONIAZID – Retail pharmacy-Specialist a) No patient co-payment payable
respiratory physician Tab 100 mg
Tab 100 mg
ISONIAZID – Retail pharmacy-Specialist a) No patient co-payment payable
a) No patient co-payment payable
a) No patient co-payment payable
a,
microbiologist, dermatologist or public health physician
★ Tab 100 mg
ISONIAZID WITH RIFAMPICIN – Retail pharmacy-Specialist
a) No patient co-payment payable
b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician, paediatrician, clinical
microbiologist, dermatologist or public health physician
★ Tab 100 mg with rifampicin 150 mg
★ Tab 150 mg with rifampicin 300 mg
PARA-AMINO SALICYLIC ACID – Retail pharmacy-Specialist
a) No patient co-payment payable
b) Prescriptions must be written by, or on the recommendation of, an infectious disease specialist, clinical microbiologist or
respiratory physician Grans for oral lig 4 g sachet
PROTIONAMIDE – Retail pharmacy-Specialist
a) No patient co-payment payable
b) Prescriptions must be written by, or on the recommendation of, an infectious disease specialist, clinical microbiologist or respiratory physician
Tab 250 mg
•
PYRAZINAMIDE – Retail pharmacy-Specialist
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or
respiratory physician
* Tab 500 mg
RIFABUTIN – Retail pharmacy-Specialist
a) No patient co-payment payable
b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, respiratory physician or
gastroenterologist
* Cap 150 mg

	Subsidy		Fully	Brand or
	(Manufacturer's Price \$) Sub Per	sidised ✓	Generic Manufacturer
IFAMPICIN – Subsidy by endorsement				
 a) No patient co-payment payable b) For confirmed recurrent Staphylococcus aureus infection antimicrobial based on susceptibilities and the prescriptio Retail pharmacy - Specialist. Specialist must be an interr paediatrician, or public health physician. Cap 150 mg	n is endorsed accor nal medicine physici 55.75 116.25	dingly; can	be waiv microbi	/ed by endorsement -
Antivirals				
or eye preparations refer to Eye Preparations, Anti-Infective Pre	parations, page 220)		
Hepatitis B Treatment				
DEFOVIR DIPIVOXIL – Special Authority see SA0829 below – Tab 10 mg ■ SA0829 Special Authority for Subsidy nitial application only from a gastroenterologist or infectious dis		30		lepsera
 neeting the following criteria: Il of the following: 1 Patient has confirmed Hepatitis B infection (HBsAg+); and Documented resistance to lamivudine, defined as: 2 Patient has raised serum ALT (> 1 × ULN); and 3 Patient has HBV DNA greater than 100,000 copies per ml 4 Detection of M204I or M204V mutation; and 5 Either: 5.1 Both: 	I			
 5.1.1 Patient is cirrhotic; and 5.1.2 adefovir dipivoxil to be used in combination 5.2 Both: 5.2.1 Patient is not cirrhotic; and 5.2.2 adefovir dipivoxil to be used as monotherap 				
 lenewal only from a gastroenterologist or infectious disease speeating physician, treatment remains appropriate and patient is blotes: Lamivudine should be added to adefovir dipivoxil if a patiefined as: i) raised serum ALT (> 1 × ULN); and ii) HBV DNA greater than 100,000 copies per mL, or viral loading to the serum and th	enefiting from treatr ent develops docum	nent. ented resis	stance to	
iii) Detection of N236T or A181T/V mutation. defovir dipivoxil should be stopped 6 months following HBeAg s ommencing adefovir dipivoxil. he recommended dose of adefovir dipivoxil is no more than 10m o patients with renal insufficiency adefovir dipivoxil dose should I defovir dipivoxil should be avoided in pregnant women and child	eroconversion for pang daily.	atients who	were H	
NTECAVIR	<i>וו</i> כוו.			
← Tab 0.5 mg		30	✓ E	intecavir Sandoz
AMIVUDINE – Special Authority see SA1685 on the next page Tab 100 mg Oral lig 5 mg per ml	4.20	28 40 ml OP	_	<u>etlam</u> effix

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		idised	Generic
	\$	Per		Manufacturer
SA1685 Special Authority for Subsidy				
initial application only from a relevant specialist or medical prac	titioner on the recom	mendation	of a re	levant specialist.
Approvals valid for 1 year where used for the treatment or prever				
Renewal from any relevant practitioner. Approvals valid for 2 ye		e treatmen	t or pre	evention of hepatitis B.
TENOFOVIR DISOPROXIL				
Tenofovir disoproxil prescribed under endorsement for the tr	eatment of HIV is incl	udad in the	count	of up to 4 subsidised
antiretrovirals for the purposes of Special Authority SA1651.			Count	or up to 4 subsidised
 * Tab 245 mg (300.6 mg as a succinate) 		30	🖌 Т	enofovir Disoproxil
		00	• 1	Teva
				1014
Herpesvirus Treatments				
ACICLOVIR				
* Tab dispersible 200 mg	1.60	25	✓ L	ovir
* Tab dispersible 400 mg	5.38	56	✓ L	ovir
* Tab dispersible 800 mg	5.98	35	✓ L	ovir
VALACICLOVIR				
Tab 500 mg		30	🗸 V	aclovir
Tab 1,000 mg		30	_	aclovir
VALGANCICLOVIR – Special Authority see SA1404 below – Re			-	<u> </u>
Tab 450 mg		60	√ v	alganciclovir
1 au 700 1119		00	• v	Mylan
	(1.050.00)		v	•
Valganaiolovir Mulan to be Sale Supply on 1 August 001	(1,050.00)		v	alcyte
Valganciclovir Mylan to be Sole Supply on 1 August 201	9			

(Valcyte Tab 450 mg to be delisted 1 August 2019)

➡SA1404 Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
- 2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a lung transplant; and
- 2 Either:

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
- 2.2 The recipient is cytomegalovirus positive.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised: and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease: or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via PHARMAC's approve	ed direct distribution s	upply. Furthe	r details can be found on
PHARMAC's website https://www.pharmac.govt.nz/hepa	atitis-c-treatments		
Tab 100 mg with pibrentasvir 40 mg	24,750.00	84 OP	 Maviret
LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Au	uthority see SA1605 b	elow	
No patient co-payment payable			
Tab 90 mg with sofosbuvir 400 mg	24,363.46	28	 Harvoni
SA1605 Special Authority for Subsidy			

SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP) Notes: By application to the Hepatitis C Treatment Panel (HepCTP). Applications will be considered by HepCTP and approved subject to confirmation of eligibility. Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz/hepatitis-c-treatments or: The Coordinator, Hepatitis C Treatment Panel PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 460 4990. Email: hepcpanel@pharmac.govt.nz

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL – Subsidy by endorsement; can be waived by Special Authority see SA1714 below

below Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 104 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the PHARMAC website.

Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a

fumarate)	61.15	30	
,	(190.02)		Truvada
Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a			
succinate)	61.15	30	🗸 Teva
Teva to be Sole Supply on 1 September 2019			

(Truvada Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a fumarate) to be delisted 1 September 2019)

➡SA1714 Special Authority for Waiver of Rule

Initial application only from a named specialist or medical practitioner on the recommendation of a named specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tested HIV negative; and
- 2 Either:
 - 2.1 All of the following:
 - 2.1.1 Patient is male or transgender; and
 - 2.1.2 Patient has sex with men; and
 - 2.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 2.1.4 Any of the following:
 - 2.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 2.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 2.1.4.3 Patient has used methamphetamine in the last three months; or
 - 2.2 All of the following:
 - 2.2.1 Patient has a regular partner who has HIV infection; and
 - 2.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 2.2.3 Condoms have not been consistently used.

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis; and
- 2 Patient has undergone testing for HIV, syphilis, and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative; and
- 6 Either:
 - 6.1 All of the following:

continued...

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price)	Subs	Fully idised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 6.1.1 Patient is male or transgender; and
- 6.1.2 Patient has sex with men; and
- 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
- 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
- 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Antiretrovirals

⇒SA1651 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals.

Su	ubsidy	Fully	Brand or
(Manufac	turer's Price) Subsid	lised	Generic
	\$ Per	1	Manufacturer

continued...

Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA1651 on the p	revious page – Retail phar	rmacy	
Tab 50 mg	63.38	30	 Stocrin S29
Tab 200 mg		90	 Stocrin
Tab 600 mg	63.38	30	 Stocrin
Oral liq 30 mg per ml	145.79	180 ml OP	 Stocrin S29
ETRAVIRINE - Special Authority see SA1651 on the	previous page – Retail pha	armacy	
Tab 200 mg	770.00	60	 Intelence
NEVIRAPINE - Special Authority see SA1651 on the	previous page – Retail pha	armacy	
Tab 200 mg	60.00	60	 Nevirapine
			<u>Alphapharm</u>
Oral suspension 10 mg per ml		240 ml	 Viramune
			Suspension

Nucleosides Reverse Transcriptase Inhibitors

ABACAVIR SULPHATE – Special Authority see SA1651 on the Tab 300 mg		Retail pharmac 60	y ✔ Ziagen
Ziagen to be Sole Supply on 1 July 2019 Oral lig 20 mg per ml		240 ml OP	✓ Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority Note: abacavir with lamivudine (combination tablets) counts	/ see SA1651 on	the previous pa	age – Retail pharmacy
anti-retroviral Special Authority.			
Tab 600 mg with lamivudine 300 mg Kivexa to be Sole Supply on 1 July 2019		30	 Kivexa

	Subsidy		Fully	Brand or
	,	anufacturer's Price) Subsic		Generic
	\$	Per	1	Manufacturer
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOP	ROXIL – Specia	I Authority see	SA165 ⁻	1 on page 104 – Retail
pharmacy		,		
Note: Efavirenz with emtricitabine and tenofovir disoproxil c	ounts as three a	nti-retroviral me	dicatio	ns for the purposes of the
anti-retroviral Special Authority				
Tab 600 mg with emtricitabine 200 mg and tenofovir disopro				
245 mg (300 mg as a fumarate)		30		to be to
	(237.52)		Α	tripla
Tab 600 mg with emtricitabine 200 mg and tenofovir disopro		20	. ()	hulon
245 mg (300 mg as a maleate) Mylan to be Sole Supply on 1 September 2019		30	• 1	lylan
(Atripla Tab 600 mg with emtricitabine 200 mg and tenofovir disc	provil 245 ma (2	00 ma ac a fum	arata)	to be delicted 1 September
2019)	ipi0xii 245 iliy (5	oo my as a rum	aiale)	
EMTRICITABINE – Special Authority see SA1651 on page 104			_	
Cap 200 mg		30	✓ E	mtriva
LAMIVUDINE - Special Authority see SA1651 on page 104 - R	etail pharmacy			
Tab 150 mg	52.50	60	✓ L	amivudine
			_	Alphapharm
Oral liq 10 mg per ml		240 ml OP	✓ 3	TC
ZIDOVUDINE [AZT] - Special Authority see SA1651 on page 10	04 – Retail pharn	nacy		
Cap 100 mg		100		letrovir
Oral liq 10 mg per ml		200 ml OP	✓ <u>F</u>	letrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority se Note: zidovudine [AZT] with lamivudine (combination tablets the anti-retroviral Special Authority. Tab 300 mg with lamivudine 150 mg	s) counts as two		edicati	
Protease Inhibitors				
ATAZANAVIR SULPHATE - Special Authority see SA1651 on p	ago 104 – Rotai	Inharmaov		
Cap 150 mg		60	√ T	eva
	(568.34)			Reyataz
Teva to be Sole Supply on 1 September 2019	()			· · · · ·
Cap 200 mg		60	🗸 I	eva
	(757.79)		F	leyataz
Teva to be Sole Supply on 1 September 2019				
(Reyataz Cap 150 mg to be delisted 1 September 2019)				
(Reyataz Cap 200 mg to be delisted 1 September 2019)				
DARUNAVIR - Special Authority see SA1651 on page 104 - Re				
Tab 400 mg		60		rezista
Tab 600 mg	476.00	60	✓ P	Prezista
LOPINAVIR WITH RITONAVIR - Special Authority see SA1651				
Tab 100 mg with ritonavir 25 mg		60	-	aletra
Tab 200 mg with ritonavir 50 mg		120	_	<u>Caletra</u>
Oral liq 80 mg with ritonavir 20 mg per ml		300 ml OP	✓ K	Caletra
RITONAVIR - Special Authority see SA1651 on page 104 - Ret				
Tab 100 mg	43.31	30	🗸 N	lorvir
Norvir to be Sole Supply on 1 July 2019				

	Subsidy (Manufacturer's Price) \$	Subsic Per	Fully lised	Brand or Generic Manufacturer
Strand Transfer Inhibitors				
DOLUTEGRAVIR – Special Authority see SA1651 on page 104 - Tab 50 mg		30	🗸 Ti	vicay
RALTEGRAVIR POTASSIUM – Special Authority see SA1651 o Tab 400 mg		harmacy 60	✔ Is	entress
Immune Modulators				

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

- 1) Diagnosis
 - Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
 - · PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
 - Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Exclusion Criteria

- 1) Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia (< 2.0×10^9) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

INTERFERON ALFA-2A - PCT - Retail pharmacy-Specialist

- a) See prescribing guideline above
- b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician or ophthalmologist

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA1400 below - Retail pharmacy

- a) See prescribing guideline above

⇒SA1400 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post **liver transplant)** from any specialist. Approvals valid for 18 months for applications meeting the following criteria: Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	~	Manufacturer	

continued...

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or

INFECTIONS - AGENTS FOR SYSTEMIC USE

Subsidy (Manufacturer's Price)	Fully Subsidised	
\$	Per 🗸	Manufacturer

continued...

- 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Notes:

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.
- · Pegylated Interferon-alfa 2a is not approved for use in children.

Urinary Tract Infections

HEXAMINE HIPPURATE

	10.40	100	
* Tab 1 g		100	
	(40.01)		Hiprex
NITROFURANTOIN			
* Tab 50 mg		100	 Nifuran
* Tab 100 mg		100	 Nifuran
NORFLOXACIN			
Tab 400 mg – Subsidy by endorsement		100	Arrow-Norfloxacin

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

	a t · · ·			
	Subsidy (Manufacturer's Pr	rico) Sub	Fully Brand sidised Gene	
	(Manulactuler's Fi	Per Sub		facturer
	•			
Anticholinesterases				
EOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule		50	✓ AstraZe	neca
YRIDOSTIGMINE BROMIDE				
Tab 60 mg	42 79	100	✓ Mesting	'n
	12.70	100	mootine	<u></u>
Non-Steroidal Anti-Inflammatory Drugs				
ICLOFENAC SODIUM				
 Tab EC 25 mg 	1 23	50	🖌 Diclofer	nac Sandoz
 Tab E0 25 mg Tab 50 mg dispersible 		20	✓ Voltare	
 Tab 50 mg dispersible Tab EC 50 mg 		50		nac Sandoz
 Tab long-acting 75 mg 		500	✓ Apo-Dic	
 Tab long acting 70 mg Tab long-acting 100 mg 		500	✓ Apo-Dic	
 Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available or 		5	✓ Voltare	
 Suppos 12.5 mg 		10	✓ Voltarei	-
 Suppos 12.5 mg		10	✓ Voltarei	-
 Suppos 20 mg – Up to 10 supp available on a PSO 		10	✓ Voltarei	
Suppos 100 mg		10	✓ Voltarei	
		10	• voltarei	
BUPROFEN		4 000		
Tab 200 mg		1,000	Relieve	
Tab long-acting 800 mg		30	 Brufen 	SR
Oral liq 20 mg per ml	1.88	200 ml	 Ethics 	
			 Fenpae 	d
Ethics to be Sole Supply on 1 August 2019				
Fenpaed Oral liq 20 mg per ml to be delisted 1 August 2019	9			
ETOPROFEN				
 Cap long-acting 200 mg 	12.07	28	🗸 Oruvail	SR
	12.07	28	🗸 Oruvail	SR
IEFENAMIC ACID			 Oruvail 	SR
 Cap long-acting 200 mg IEFENAMIC ACID € Cap 250 mg 	1.25	28 50		-
IEFENAMIC ACID	1.25 (9.16)	50	 Oruvail Ponstan 	-
IEFENAMIC ACID	1.25 (9.16) 0.50		Ponstan	-
IEFENAMIC ACID Cap 250 mg	1.25 (9.16)	50		-
IEFENAMIC ACID Cap 250 mg		50 20	Ponstan Ponstan	-
IEFENAMIC ACID Cap 250 mg		50 20 500	Ponstan Ponstan ✔ <u>Noflam</u>	250
IEFENAMIC ACID Cap 250 mg IAPROXEN Tab 250 mg Tab 500 mg		50 20 500 250	Ponstan Ponstan ✔ <u>Noflam</u> ✔ <u>Noflam</u>	<u>250</u> 500
IEFENAMIC ACID ← Cap 250 mg IAPROXEN ← Tab 250 mg		50 20 500 250 28	Ponstan Ponstan ✓ <u>Noflam</u> ✓ <u>Noflam</u> ✓ <u>Napros</u>	250 500 yn SR 750
IEFENAMIC ACID		50 20 500 250	Ponstan Ponstan ✓ <u>Noflam</u> ✓ <u>Noflam</u> ✓ <u>Napros</u>	<u>250</u> 500
IEFENAMIC ACID		50 20 500 250 28 28 28	Ponstan Ponstan ✓ <u>Noflam</u> ✓ <u>Noflam</u> ✓ <u>Napros</u>	250 500 yn SR 750
IEFENAMIC ACID		50 20 500 250 28 28 28 50	Ponstan Ponstan <u>Noflam</u> <u>Noflam</u> <u>Napros</u> <u>Aclin</u>	250 500 yn SR 750
IEFENAMIC ACID		50 20 500 250 28 28 28	Ponstan Ponstan ✓ <u>Noflam</u> ✓ <u>Noflam</u> ✓ <u>Napros</u>	250 500 yn SR 750
IEFENAMIC ACID		50 20 500 250 28 28 28 50	Ponstan Ponstan <u>Noflam</u> <u>Noflam</u> <u>Napros</u> <u>Aclin</u>	250 500 yn SR 750
IEFENAMIC ACID		50 20 500 250 28 28 28 50	Ponstan Ponstan <u>Noflam</u> <u>Noflam</u> <u>Napros</u> <u>Aclin</u>	250 500 yn SR 750

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per Brand or Generic Manufacture Manufacture NSAIDs Other Generic Manufacture's Price) Manufacture's Price) <th>- </th>	-
\$ Per ✓ Manufactur NSAIDs Other SELECOXIB 3.63 60 ✓ Celebrex Cap 100 mg	-
ELECOXIB Cap 100 mg	fizer
Cap 100 mg	fizer
Cap 100 mg	fizer
Cap 200 mg2.30 30 Celebrex Celebrex Cap 100 mg to be delisted 1 September 2019) Topical Products for Joint and Muscular Pain CAPSAICIN Crm 0.025% – Special Authority see SA1289 below – Retail	fizer
Celebrex Cap 100 mg to be delisted 1 September 2019) Topical Products for Joint and Muscular Pain CAPSAICIN Crm 0.025% – Special Authority see SA1289 below – Retail	
Celebrex Cap 100 mg to be delisted 1 September 2019) Topical Products for Joint and Muscular Pain APSAICIN Crm 0.025% – Special Authority see SA1289 below – Retail	6
APSAICIN Crm 0.025% – Special Authority see SA1289 below – Retail	lizer
Crm 0.025% - Special Authority see SA1289 below - Retail	
pharmacy6.95 25 g OP 🖌 Zostrix	
9.95 45 g OP 🗸 Zostrix	
Special Authority for Subsidy itial application from any relevant practitioner. Approvals valid without further renewal unless notified where the	patient ha
steoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.	
Antirheumatoid Agents	
YDROXYCHLOROQUINE	
€ Tab 200 mg	
EFLUNOMIDE	
Tab 10 mg	
Tab 20 mg2.90 30 🖌 Apo-Lefluno	mide
ENICILLAMINE	
Tab 125 mg	
Tab 250 mg 110.12 100 ✓ D-Penamine	
ODIUM AUROTHIOMALATE	
Inj 10 mg in 0.5 ml ampoule ✓ Myocrisin Inj 20 mg in 0.5 ml ampoule 113.17 10 ✓ Myocrisin	
Inj 50 mg in 0.5 ml ampoule	
Myocrisin Inj 10 mg in 0.5 ml ampoule to be delisted 1 March 2020)	
Myocrisin Inj 20 mg in 0.5 ml ampoule to be delisted 1 March 2020)	
Ívocrisin Inj 50 mg in 0.5 ml ampoule to be delisted 1 March 2020)	
Drugs Affecting Bone Metabolism	
Alendronate for Osteoporosis	
LENDRONATE SODIUM	
€ Tab 70 mg	
LENDRONATE SODIUM WITH COLECALCIFEROL	
← Tab 70 mg with colecalciferol 5,600 iu	

Other Treatments

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy Fully Brand	lor
(Manufacturer's Price) Subsidised Gener	ric
\$ Per 🖌 Manuf	facturer

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Either:
 - 2.1 The patient is female and postmenopausal; or
 - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

	Inj 3 mg per ml, 10 ml vial	5.98	1	 Pamisol
	Inj 6 mg per ml, 10 ml vial	15.02	1	 Pamisol
	Inj 9 mg per ml, 10 ml vial		1	 Pamisol
RAL	OXIFENE HYDROCHLORIDE – Special Authority see SA1779 or	n the next page -	- Retail pha	armacy
*	Tab 60 mg	53.76	28	 Evista
	-			

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ubsidised	Generic	
\$	Per	1	Manufacturer	

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM

Tab 35 mg	4	 Risedronate Sandoz
TERIPARATIDE – Special Authority see SA1139 below – Retail pharmacy		
Inj 250 mcg per ml, 2.4 ml	1	 Forteo

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops

	Subsidy	Ful		
Λ)	Anufacturer's Price)	Subsidise	ed Generic	
	\$	Per •	 Manufacturer 	

continued...

during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

00.00 100 ml OP

Aclasta

SA1780 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- All of the following:
 - 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
 - 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	~	Manufacturer	

continued...

- 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
- 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

-2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.

d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

ALLOPURINOL

* Tab 100 mg * Tab 300 mg		DP-Allopurinol DP-Allopurinol
BENZBROMARONE - Special Authority see SA1537 below		
Tab 100 mg	 0 🗸	Benzbromaron AL
		100 S29

⇒SA1537 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 Both:
 - 2.3.1 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Notes); and
 - 2.3.2 The patient has a rate of creatinine clearance greater than or equal to 20 ml/min; or
 - 2.4 All of the following:
 - 2.4.1 The patient is taking azathioprine and requires urate-lowering therapy; and
 - 2.4.2 Allopurinol is contraindicated; and
 - 2.4.3 Appropriate doses of probenecid are ineffective or probenecid cannot be used due to reduced renal function; and
- 3 The patient is receiving monthly liver function tests.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.
- Notes: Benzbromarone has been associated with potentially fatal hepatotoxicity.

In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

The New Zealand Rheumatology Association has developed information for prescribers which can be accessed from its website at www.rheumatology.org.nz/home/resources-2/

COLCHICINE

*	Tab 500 mcg9.	.58 100	 Colgout
---	---------------	---------	-----------------------------

	Subsidy (Manufacturer's Price) \$	Su Per	Fully ubsidised	Brand or Generic Manufacturer	
FEBUXOSTAT – Special Authority see SA1538 below – Retail pl	harmacy				_
Tab 80 mg		28	🗸 A	denuric	
Tab 120 mg		28	🗸 🗸	denuric	

► SA1538 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note).

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

PROBENECID

* Tab 500 mg	55.00	100	✓ Probenecid-AFT
Muscle Relaxants			
BACLOFEN			
* Tab 10 mg		100	✓ Pacifen
Inj 0.05 mg per ml, 1 ml ampoule – Subsidy by endorsement.	11.55	1	 Lioresal Intrathecal
Subsidised only for use in a programmable pump in patie caused intolerable side effects and the prescription is end			ents have been ineffective or have
Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement	372.98	5	 Medsurge
	74.60	1	
	(209.29)		Lioresal Intrathecal
 a) Subsidised only for use in a programmable pump in practice by have caused intolerable side effects and the prescription by Medsurge to be Sole Supply on 1 July 2019 (Lioresal Intrathecal Inj 2 mg per ml, 5 ml ampoule to be delisted to be de	on is endorsed ad		agents have been ineffective or
DANTROLENE			
Cap 25 mg	65.00	100	 Dantrium
Cap 50 mg	77.00	100	 ✓ Dantrium S29 S29 ✓ Dantrium
ORPHENADRINE CITRATE Tab 100 mg	18.54	100	✓ <u>Norflex</u>

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	1	Manufacturer
Agents for Parkinsonism and Related Disord	ers			
Dopamine Agonists and Related Agents				
AMANTADINE HYDROCHLORIDE				
▲ Cap 100 mg		60	1	Symmetrel
APOMORPHINE HYDROCHLORIDE				•
Inj 10 mg per ml, 2 ml ampoule		5	1	Movapo
BROMOCRIPTINE MESYLATE				-
₭ Tab 2.5 mg		100	1	Apo-Bromocriptine
ENTACAPONE				
Tab 200 mg		100	1	Entapone
EVODOPA WITH BENSERAZIDE				.
Tab dispersible 50 mg with benserazide 12.5 mg		100	1	Madopar Rapid
₭ Cap 50 mg with benserazide 12.5 mg		100	-	Madopar 62.5
Cap 100 mg with benserazide 25 mg	15.80	100	✓	Madopar 125
Cap long-acting 100 mg with benserazide 25 mg		100	-	Madopar HBS
Cap 200 mg with benserazide 50 mg		100	~	Madopar 250
EVODOPA WITH CARBIDOPA				
K Tab 100 mg with carbidopa 25 mg	17.97	100		Kinson
				Sinemet
k Tab long-acting 100 mg with carbidopa 25 mg		100		Mylan S29
Tab long-acting 200 mg with carbidopa 50 mg		100		Sinemet CR
★ Tab 250 mg with carbidopa 25 mg		100	•	Sinemet
	7.00	400		D
Tab 0.25 mg		100 100		<u>Ramipex</u> Ramipex
	24.09	100	•	namipex
	0.70	100		Ana Daninirala
Tab 0.25 mg Tab 1 mg		100 100		<u>Apo-Ropinirole</u> Apo-Ropinirole
Tab 2 mg		100		Apo-Ropinirole
Tab 5 mg		100		Apo-Ropinirole
ELEGILINE HYDROCHLORIDE				
★ Tab 5 mg		100	1	Apo-Selegiline
·				S29 S29
OLCAPONE				
Tab 100 mg		100	1	Tasmar
Anticholinergics				
-				
BENZATROPINE MESYLATE	7.00	60		Pontron
Tab 2 mg Inj 1 mg per ml, 2 ml		60 5		Benztrop Cogentin
יוון י וווץ אָכו וווו, ב וווי	190.00	5 10		Omega
a) Up to 10 inj available on a PSO	100.00	10	•	e
b) Only on a PSO				
PROCYCLIDINE HYDROCHLORIDE				
Tab 5 mg	7 40	100	1	Kemadrin
· · · · · · · · · · · · · · · · · · ·	VT. 7		-	

			NER	VOUS SYSTEM
	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
Agents for Essential Tremor, Chorea and Relate	d Disorders			
RILUZOLE – Special Authority see SA1403 below – Retail pharm Wastage claimable Tab 50 mg		56 r 6 month	_	ilutek
 All of the following: 1 The patient has amyotrophic lateral sclerosis with disease 2 The patient has at least 60 percent of predicted forced vita 3 The patient has not undergone a tracheostomy; and 4 The patient has not experienced respiratory failure; and 5 Any of the following: 5.1 The patient is ambulatory; or 5.2 The patient is able to use upper limbs; or 5.3 The patient is able to swallow. 				initial application; and
 Renewal from any relevant practitioner. Approvals valid for 18 m All of the following: The patient has not undergone a tracheostomy; and The patient has not experienced respiratory failure; and Any of the following: The patient is ambulatory; or The patient is able to use upper limbs; or The patient is able to swallow. 	onths for application	s meeting	the follo	wing criteria:
TETRABENAZINE Tab 25 mg	91.10	112	✓ <u>N</u>	lotetis
Anaesthetics				
Local				
LIDOCAINE [LIGNOCAINE] Gel 2%, tube – Subsidy by endorsement a) Up to 150 ml available on a PSO	14.50	30 ml	✓ X	ylocaine 2% Jelly
 b) Subsidised only if prescribed for urethral or cervical a Gel 2%, 10 ml urethral syringe – Subsidy by endorsement 		prescript 10 25	✓ P	ndorsed accordingly. fizer atheiell
a) Up to 5 each available on a PSO	100.00	20		utilejen

a) Up to 5 each available on a PSO
b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly. (*Pfizer Gel 2%, 10 ml urethral syringe to be delisted 1 November 2019*)

	Subsidy		Fully Brand or	
	(Manufacturer's Pri	ce) Sub Per	sidised Generic Manufacturer	
	\$	rei		
	00.00	000	(Marca and the	
Oral (gel) soln 2%		200 ml	✓ <u>Mucosoothe</u>	
Inj 1%, 5 ml ampoule – Up to 25 inj available on a PSO		25	 Lidocaine-Claris 	
	17.50	50	Vulaccina	
Inj 2%, 5 ml ampoule – Up to 5 inj available on a PSO	(35.00)	25	Xylocaine Lidocaine-Claris	
Inj 1%, 20 ml ampoule – Up to 5 inj available on a PSO		25 5		
	(20.00)	5	Xylocaine	
Inj 1%, 20 ml vial – Up to 5 inj available on a PSO	()	5	✓ Lidocaine-Claris	
Lidocaine-Claris to be Sole Supply on 1 July 2019		0		
Inj 2%, 20 ml vial – Up to 5 inj available on a PSO	6.45	5	 Lidocaine-Claris 	
Lidocaine-Claris to be Sole Supply on 1 July 2019		Ũ		
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes – Subsidy by endorsement		10	✓ Pfizer	
		10	▼ Plizer	
 a) Up to 5 each available on a PSO b) Subsidised only if prescribed for urethral or cervical 	administration and	the preseriet	ion is anderead accordingly	
b) Subsidised only it prescribed for dretifial of cervical	auministration and	the prescript	ion is endorsed accordingly	•
Topical Local Anaesthetics				
Topiour Loour Anacotricitos				
Renewal from any relevant practitioner. Approvals valid for 2 ye benefiting from treatment. LIDOCAINE [LIGNOCAINE] – Special Authority see SA0906 ab Crm 4%	ove – Retail pharm		LMX4	111 15
	27.00	30 g OP	🗸 LMX4	
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Auth	nority see SA0906	Ũ	il nharmacy	
Crm 2.5% with prilocaine 2.5%		30 g OP		
Crm 2.5% with prilocaine 2.5% (5 g tubes)		5	✓ EMLA	
		-		
Analgesics				
For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, p	000 110			
	age 110			
Non-opioid Analgesics				
For aspirin & chloroform application refer Standard Formulae, pa	age 227			
ASPIRIN	-			
* Tab dispersible 300 mg – Up to 30 tab available on a PSO.	3 90	100	 Ethics Aspirin 	
		100		
CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or c accordingly.	liabetic peripheral	neuropathy a	nd the prescription is endor	rsed
Crm 0.075%		45 g OP	 Zostrix HP 	
NEFOPAM HYDROCHLORIDE		- 3		
Tab 30 mg	02 10	90	 Acupan 	
		90	ACIONIC	

Subsidy Fully Brand or Generic ARACETAMOL \$ Per ✓ Manufacturer ARACETAMOL 7.12 1,000 ✓ Priceline 7.12 1,000 ✓ Paracetamol Pharmacare ✓ Pharmacare ✓ Pharmacare ✓ Pharmacare ✓ Pharmacare ✓ Oral liq 120 mg per 5 ml 5.35 1,000 ml ✓ Paracare a) Up to 200 ml available on a PSO b) Not in combination 5.81 1,000 ml ✓ Paracare Double Strength 3.29 10 ✓ Gacet Suppos 250 mg 3.73 10 ✓ Gacet Suppos 250 mg .3.73 10 ✓ Gacet Suppos 500 mg .12.40 50 ✓ Gacet Priceline .5.75 100 ✓ PSM Tab 15 mg .5.75 100 ✓ PSM Tab 15 mg .5.75 100 ✓ PSM Tab 15 mg .5.75 100 ✓ PSM Tab 15 mg .13.50 100 ✓
\$ Per Manufacturer ARACETAMOL 7.12 100 Priceline 7.12 1,000 7.12 1,000 Pharmacare 7.12 1,000 Pharmacare Pharmacare Yearacetamol Pharmacare Pharmacare Yearacetamol Pharmacare Yearacetamol Yearacetamol Pharmacare Yearacetamol Yearacetamol Pharmacare Yearacetamol Yearacare Yearacare Yearacare Yearacare Yearacare Year
ARACETAMOL Tab 500 mg - blister pack – Up to 30 tab available on a PSO
 Tab 500 mg - blister pack – Up to 30 tab available on a PSO0.71 7.12 Tab 500 mg - bottle pack
7.12 1,000 ✓ Paracetamol Pharmacare ✓ Pharmacare ✓ Pharmacare ✓ Pharmacy Health É Tab 500 mg - bottle pack
Pharmacare Pharmacare Pharmacare Pharmacy Health Pharmacy Health Pharmacy Health Pharmacare Pharmacare Pharmacare Pharmacare Pharacare Pharacare
 Tab 500 mg - bottle pack
 Tab 500 mg - bottle pack
 Tab 500 mg - bottle pack
 Gral liq 120 mg per 5 ml
 a) Up to 200 ml available on a PSO b) Not in combination c) Oral liq 250 mg per 5 ml a) Up to 100 ml available on a PSO b) Not in combination c) Suppos 125 mg c) Suppos 250 mg c) Suppos 250 mg c) Suppos 500 mg <lic) 500="" li="" mg<="" suppos=""> c) Suppos 500 mg <lic) sup<="" td=""></lic)></lic)>
b) Not in combination Coral liq 250 mg per 5 ml
 Gral liq 250 mg per 5 ml
a) Up to 100 ml available on a PSO b) Not in combination
a) Up to 100 ml available on a PSO b) Not in combination
a) Up to 100 ml available on a PSO b) Not in combination Suppos 125 mg
b) Not in combination # Suppos 125 mg # Suppos 250 mg # Suppos 250 mg # Suppos 500 mg # Supos 500 mg # Suppos 5
Suppos 125 mg 3.29 10 ✓ Gacet Suppos 250 mg 3.79 10 ✓ Gacet Suppos 500 mg 12.40 50 ✓ Gacet Priceline Tab 500 mg - blister pack to be delisted 1 August 2019) 12.40 50 ✓ Gacet ODEINE PHOSPHATE - Safety medicine; prescriber may determine dispensing frequency Tab 15 mg 5.75 100 ✓ PSM Tab 30 mg 6.80 100 ✓ PSM Tab 60 mg 13.50 100 ✓ PSM IHYDROCODEINE TARTRATE 100 ✓ PSM
 Suppos 250 mg
Suppos 500 mg
Opioid Analgesics ODEINE PHOSPHATE - Safety medicine; prescriber may determine dispensing frequency Tab 15 mg 5.75 100 <u>PSM</u> Tab 30 mg 6.80 100 <u>PSM</u> Tab 60 mg 13.50 100 <u>PSM</u> IHYDROCODEINE TARTRATE
Opioid Analgesics ODEINE PHOSPHATE - Safety medicine; prescriber may determine dispensing frequency Tab 15 mg 5.75 100
ODEINE PHOSPHATE – Safety medicine; prescriber may determine dispensing frequency Tab 15 mg
ODEINE PHOSPHATE – Safety medicine; prescriber may determine dispensing frequency Tab 15 mg
Tab 15 mg 5.75 100 ✓ PSM Tab 30 mg 6.80 100 ✓ PSM Tab 60 mg 13.50 100 ✓ PSM IHYDROCODEINE TARTRATE 100 ✓ PSM
Tab 15 mg 5.75 100 ✓ PSM Tab 30 mg 6.80 100 ✓ PSM Tab 60 mg 13.50 100 ✓ PSM IHYDROCODEINE TARTRATE 100 ✓ PSM
Tab 30 mg 6.80 100 PSM Tab 60 mg 13.50 100 PSM IHYDROCODEINE TARTRATE
Tab 60 mg
IHYDROCODEINE TARTRATE
Tab long-acting 60 mg
ENTANYL
a) Only on a controlled drug form
b) No patient co-payment payable
c) Safety medicine; prescriber may determine dispensing frequency
Inj 50 mcg per ml, 2 ml ampoule
Inj 50 mcg per ml, 10 ml ampoule
Patch 12.5 mcg per hour
Patch 25 mcg per hour
51 · · · · · · · · · · · · · · · · · · ·
<u> </u>
Patch 75 mcg per hour
Patch 100 mcg per hour 11.40 5 Fentanyl Sandoz

	Subsidy		Fully	
	(Manufacturer's Price \$) Per	Subsidised	Generic Manufacturer
IETHADONE HYDROCHLORIDE	•			manaration
a) Only on a controlled drug form				
b) No patient co-payment payable				
 c) Safety medicine; prescriber may determine dispensing free 				
 d) Extemporaneously compounded methadone will only be r 		to of th	o oboono	at form available
, , , , , , ,	ennouiseu al lite la		e cheape	St IUIIII avallable
(methadone powder, not methadone tablets). e) For methadone hydrochloride oral liquid refer Standard Fo				
Tab 5 mg		10		Methatabs
5	1.40	10	•	Weinalaus
Methatabs to be Sole Supply on 1 September 2019	1.40	10		Methatabs
Tab 5 mg - bottle pack				
Oral liq 2 mg per ml		200 m		Biodone Biodone Forte
Oral liq 5 mg per ml		200 m	-	Biodone Forte
Oral liq 10 mg per ml		200 m		Biodone Extra Forte
Inj 10 mg per ml, 1 ml		10	•	AFT
Methatabs Tab 5 mg - bottle pack to be delisted 1 December 20	19)			
IORPHINE HYDROCHLORIDE				
 a) Only on a controlled drug form 				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	quency			
Oral liq 1 mg per ml		200 m	l 🗸	RA-Morph
Oral liq 2 mg per ml		200 m	l 🗸	RA-Morph
Oral liq 5 mg per ml		200 m	l 🗸	Ordine S29
			1	RA-Morph
Oral liq 10 mg per ml	27 74	200 m		Ordine S29
		200 11		RA-Morph
			•	
NORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre				• • •
Tab immediate-release 10 mg		10		Sevredol
Tab long-acting 10 mg		10		Arrow-Morphine LA
Tab immediate-release 20 mg		10		Sevredol
Tab long-acting 30 mg		10		Arrow-Morphine LA
Tab long-acting 60 mg		10		Arrow-Morphine LA
Tab long-acting 100 mg		10		Arrow-Morphine LA
Cap long-acting 10 mg		10		m-Eslon
Cap long-acting 30 mg		10		m-Eslon
Cap long-acting 60 mg		10		m-Eslon
Cap long-acting 100 mg		10		m-Eslon
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS	6.27	5	1	DBL Morphine
				Sulphate
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a F	SO 4.47	5	✓	DBL Morphine
				Sulphate
Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a F	SO 4.76	5	✓	DBL Morphine
				Sulphate
Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a F	SO6.19	5	1	DBL Morphine
, or ,		-		Sulphate

	Subsidy (Manufacturer's Price)	s	Fully ubsidised	
	\$	Per	1	Manufacturer
MORPHINE TARTRATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	equency			
Inj 80 mg per ml, 1.5 ml ampoule		5	✓	DBL Morphine
				Tartrate
OXYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
 c) Safety medicine; prescriber may determine dispensing free 	auencv			
Tab controlled-release 5 mg		20	1	Oxycodone Sandoz
	(2.63)			BNM
Oxycodone Sandoz to be Sole Supply on 1 August 2019	()			
Tab controlled-release 10 mg		20	1	Oxycodone Sandoz
	(2.76)			BNM
Oxycodone Sandoz to be Sole Supply on 1 August 2019	()			
Tab controlled-release 20 mg		20	1	Oxycodone Sandoz
0	(4.72)			BNM
Oxycodone Sandoz to be Sole Supply on 1 August 2019	()			
Tab controlled-release 40 mg		20	-	Oxycodone Sandoz
-	(7.69)			BNM
Oxycodone Sandoz to be Sole Supply on 1 August 2019				
Tab controlled-release 80 mg		20	✓	Oxycodone Sandoz
	(14.11)			BNM
Oxycodone Sandoz to be Sole Supply on 1 August 2019				
Cap immediate-release 5 mg		20		OxyNorm
Cap immediate-release 10 mg		20		OxyNorm
Cap immediate-release 20 mg		20		OxyNorm
Oral liq 5 mg per 5 ml		250 ml		OxyNorm
Inj 10 mg per ml, 1 ml ampoule		5		OxyNorm
Inj 10 mg per ml, 2 ml ampoule		5		OxyNorm
Inj 50 mg per ml, 1 ml ampoule		5	~	OxyNorm
(BNM Tab controlled-release 5 mg to be delisted 1 August 2019)				
(BNM Tab controlled-release 10 mg to be delisted 1 August 2019	,			
(BNM Tab controlled-release 20 mg to be delisted 1 August 2019	,			
(BNM Tab controlled-release 40 mg to be delisted 1 August 2019	,			
(BNM Tab controlled-release 80 mg to be delisted 1 August 2019)			
PARACETAMOL WITH CODEINE - Safety medicine; prescriber		ensing f	requenc	у
* Tab paracetamol 500 mg with codeine phosphate 8 mg		1,000	✓	Paracetamol +
				Codeine (Relieve)
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing free	equency			
Tab 50 mg		10	1	PSM
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a P		5		DBL Pethidine
				Hydrochloride
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a P	SO5.12	5	1	DBL Pethidine
				Hydrochloride

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Full	
	(Manufacturer's Price) \$	Per	Subsidise	
RAMADOL HYDROCHLORIDE	*			
Tab sustained-release 100 mg	1.55	20	-	Tramal SR 100
Tab sustained-release 150 mg		20		Tramal SR 150
Tab sustained-release 200 mg		20		Tramal SR 200
Cap 50 mg		100		Arrow-Tramadol
Antidepressants				
Cyclic and Related Agents				
MITRIPTYLINE – Safety medicine; prescriber may determin				
Tab 10 mg		100		Arrow-Amitriptyline
Tab 25 mg		100		Arrow-Amitriptyline
Tab 50 mg	2.51	100	~	Arrow-Amitriptyline
LOMIPRAMINE HYDROCHLORIDE – Safety medicine; pre	scriber may determine	disper	ising frequ	lency
Tab 10 mg		100		Apo-Clomipramine
Tab 25 mg		50		Apo-Clomipramine
	9.46	100		Apo-Clomipramine
OOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy by	endorsement			
a) Safety medicine; prescriber may determine dispensing				
				in as endorsed where there
2019 and the prescription is endorsed accordingly. Pl exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg	pin] hydrochloride.	100 100		Dopress Dopress
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg Dopress Tab 75 mg to be delisted 1 August 2020)	pin] hydrochloride.	100		Dopress
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg Dopress Tab 75 mg to be delisted 1 August 2020) Dopress Cap 25 mg to be delisted 1 January 2020)	pin] hydrochloride.	100		Dopress
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg Dopress Tab 75 mg to be delisted 1 August 2020) Dopress Cap 25 mg to be delisted 1 January 2020) DOXEPIN HYDROCHLORIDE – Subsidy by endorsement	pin] hydrochloride. 11.19 6.45	100		Dopress
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg Dopress Tab 75 mg to be delisted 1 August 2020) Dopress Cap 25 mg to be delisted 1 January 2020)	pin] hydrochloride. 	100 100 ydroc	hloride pr	Dopress Dopress
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg Dopress Tab 75 mg to be delisted 1 August 2020) Dopress Cap 25 mg to be delisted 1 January 2020) DOXEPIN HYDROCHLORIDE – Subsidy by endorsement a) Safety medicine; prescriber may determine dispensing b) Subsidy by endorsement – Subsidised for patients wh prescription is endorsed accordingly. Pharmacists ma	pin] hydrochloride. 	100 100 ydroc	hloride pris endorse	Dopress Dopress
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg Dopress Tab 75 mg to be delisted 1 August 2020) Dopress Cap 25 mg to be delisted 1 January 2020) IOXEPIN HYDROCHLORIDE – Subsidy by endorsement a) Safety medicine; prescriber may determine dispensing b) Subsidy by endorsement – Subsidised for patients wh prescription is endorsed accordingly. Pharmacists may of prior dispensing of doxepin hydrochloride.	pin] hydrochloride. 	100 100 ydroc	hloride pris s endorse	Dopress Dopress for to 1 March 2019 and the d where there exists a reco
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg Dopress Tab 75 mg to be delisted 1 August 2020) Dopress Cap 25 mg to be delisted 1 January 2020) IOXEPIN HYDROCHLORIDE – Subsidy by endorsement a) Safety medicine; prescriber may determine dispensing b) Subsidy by endorsement – Subsidised for patients wh prescription is endorsed accordingly. Pharmacists ma of prior dispensing of doxepin hydrochloride. Cap 10 mg Cap 25 mg	pin] hydrochloride. 	100 100 ydroc tion as 100	hloride prise endorse	Dopress Dopress for to 1 March 2019 and the d where there exists a reco
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg Dopress Tab 75 mg to be delisted 1 August 2020) Dopress Cap 25 mg to be delisted 1 January 2020) DOXEPIN HYDROCHLORIDE – Subsidy by endorsement a) Safety medicine; prescriber may determine dispensing b) Subsidy by endorsement – Subsidised for patients wh prescription is endorsed accordingly. Pharmacists ma of prior dispensing of doxepin hydrochloride. Cap 10 mg Cap 25 mg Cap 50 mg Cap 50 mg Can 10 mg to be delisted 1 January 2020) Anten Cap 25 mg to be delisted 1 Anuary 2020)	pin] hydrochloride. 	100 100 ydroc tion as 100 100	hloride prise endorse	Dopress Dopress for to 1 March 2019 and the d where there exists a reco
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg Dopress Tab 75 mg to be delisted 1 August 2020) Dopress Cap 25 mg to be delisted 1 January 2020) DOXEPIN HYDROCHLORIDE – Subsidy by endorsement a) Safety medicine; prescriber may determine dispensing b) Subsidy by endorsement – Subsidised for patients wh prescription is endorsed accordingly. Pharmacists ma of prior dispensing of doxepin hydrochloride. Cap 10 mg Cap 25 mg Cap 50 mg Anten Cap 10 mg to be delisted 1 January 2020) Anten Cap 50 mg to be delisted 1 April 2020) Anten Cap 50 mg to be delisted 1 May 2020)	pin] hydrochloride. 	100 100 ydroc tion as 100 100	hloride pr s endorse	Dopress Dopress for to 1 March 2019 and the d where there exists a reco Anten Anten Anten
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg Dopress Tab 75 mg to be delisted 1 August 2020) Dopress Cap 25 mg to be delisted 1 January 2020) DOXEPIN HYDROCHLORIDE – Subsidy by endorsement a) Safety medicine; prescriber may determine dispensing b) Subsidy by endorsement – Subsidised for patients wh prescription is endorsed accordingly. Pharmacists ma of prior dispensing of doxepin hydrochloride. Cap 10 mg Cap 25 mg	pin] hydrochloride. 	100 100 ydroc tion as 100 100	hloride prisendorse	Dopress Dopress for to 1 March 2019 and the d where there exists a reco Anten Anten Anten
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg Dopress Tab 75 mg to be delisted 1 August 2020) Dopress Cap 25 mg to be delisted 1 January 2020) DOXEPIN HYDROCHLORIDE – Subsidy by endorsement a) Safety medicine; prescriber may determine dispensing b) Subsidy by endorsement – Subsidised for patients wh prescription is endorsed accordingly. Pharmacists ma of prior dispensing of doxepin hydrochloride. Cap 10 mg Cap 25 mg Cap 50 mg Anten Cap 10 mg to be delisted 1 January 2020) Anten Cap 50 mg to be delisted 1 April 2020) Anten Cap 50 mg to be delisted 1 May 2020) MIPRAMINE HYDROCHLORIDE – Safety medicine; prescril	pin] hydrochloride. 	100 100 ydroc tion as 100 100 100	hloride pr s endorse	Dopress Dopress for to 1 March 2019 and the d where there exists a reco Anten Anten Anten
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg Dopress Tab 75 mg to be delisted 1 August 2020) Dopress Cap 25 mg to be delisted 1 January 2020) IOXEPIN HYDROCHLORIDE – Subsidy by endorsement a) Safety medicine; prescriber may determine dispensing b) Subsidy by endorsement – Subsidised for patients wh prescription is endorsed accordingly. Pharmacists ma of prior dispensing of doxepin hydrochloride. Cap 10 mg Cap 25 mg Cap 50 mg Cap 50 mg to be delisted 1 January 2020) Anten Cap 25 mg to be delisted 1 April 2020) Anten Cap 50 mg to be delisted 1 May 2020) MIPRAMINE HYDROCHLORIDE – Safety medicine; prescril	pin] hydrochloride. 	100 100 vdroc tion as 100 100 100	hloride pr s endorse	Dopress Dopress for to 1 March 2019 and the d where there exists a reco Anten Anten Anten Y Tofranil
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg	pin] hydrochloride. 	100 100 ydroc tion as 100 100 50 50 100 50	hloride pr s endorse	Dopress Dopress or to 1 March 2019 and the d where there exists a reco Anten Anten Anten Tofranil Tofranil Tofranil
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg Dopress Tab 75 mg to be delisted 1 August 2020) Dopress Cap 25 mg to be delisted 1 January 2020) OXEPIN HYDROCHLORIDE – Subsidy by endorsement a) Safety medicine; prescriber may determine dispensing b) Subsidy by endorsement – Subsidised for patients wh prescription is endorsed accordingly. Pharmacists ma of prior dispensing of doxepin hydrochloride. Cap 10 mg Cap 25 mg Cap 50 mg Anten Cap 25 mg to be delisted 1 January 2020) Anten Cap 50 mg to be delisted 1 January 2020) Anten Cap 50 mg to be delisted 1 May 2020) MIPRAMINE HYDROCHLORIDE – Safety medicine; prescri Tab 25 mg MAPROTILINE HYDROCHLORIDE – Safety medicine; prescri	pin] hydrochloride. 	100 100 ydrocc tion as 100 100 50 100 50 pensi	hloride prisendorse	Dopress Dopress or to 1 March 2019 and the d where there exists a reco Anten Anten Anten Tofranil Tofranil Tofranil Tofranil
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg	pin] hydrochloride. 	100 100 ydrocc tion as 100 100 50 100 50 ypensi 30	hloride pris s endorse	Dopress Dopress or to 1 March 2019 and the d where there exists a reco Anten Anten Anten Tofranil Tofranil Tofranil Tofranil Tofranil
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg	pin] hydrochloride. 	100 100 ydroc tion as 100 100 50 100 50 pensi 30 50	hloride pris s endorse	Dopress Dopress or to 1 March 2019 and the d where there exists a reco Anten Anten Anten Tofranil Tofranil Tofranil Tofranil ncy Ludiomil
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg	pin] hydrochloride. 	100 100 ydroc tion as 100 100 50 50 50 50 100	hloride pr s endorse	Dopress Dopress or to 1 March 2019 and the d where there exists a reco Anten Anten Anten Tofranil Tofranil Tofranil Tofranil ney Ludiomil Ludiomil
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg	pin] hydrochloride. 	100 100 ydroc tion as 100 100 100 50 pensi 30 50 100 20	hloride pr s endorse	Dopress Dopress or to 1 March 2019 and the d where there exists a reco Anten Anten Anten Tofranil Tofranil Tofranil Tofranil Ludiomil Ludiomil Ludiomil
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg	pin] hydrochloride. 	100 100 ydroc tion as 100 100 100 50 pensi 30 50 100 20 30	hloride pris s endorse	y Tofranil Tofranil Tofranil Tofranil Tofranil Tofranil Tofranil Ludiomil Ludiomil Ludiomil
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg	pin] hydrochloride. 11.19	100 100 ydroc tion as 100 100 100 50 pensi 30 50 100 20 30 disper	hloride pr s endorse	y Tofranil Tofranil Tofranil Tofranil Tofranil Ludiomil L
exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg	pin] hydrochloride.	100 100 ydroc tion as 100 100 100 50 pensi 30 50 100 20 30	hloride pr s endorse	y Tofranil Tofranil Tofranil Tofranil Tofranil Tofranil Tofranil Tudiomil Ludiomil Ludiomil

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	Selective			
PHENELZINE SULPHATE				
* Tab 15 mg	70.80 118.00	60 100		Nardil S29 S29 Nardil
TRANYLCYPROMINE SULPHATE * Tab 10 mg	22.94	50	1	Parnate
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE				
* Tab 150 mg	53.33	60 500	1	Aurorix
Auroriu to ha Cala Curachi an 1 July 0010	(85.10)			Apo-Moclobemide
Aurorix to be Sole Supply on 1 July 2019 * Tab 300 mg	9.80 16.33	60 100	1	Aurorix
Aurorix to be Sole Supply on 1 July 2019 (Apo-Moclobemide Tab 150 mg to be delisted 1 July 2019) (Apo-Moclobemide Tab 300 mg to be delisted 1 July 2019)	(30.70)			Apo-Moclobemide
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE * Tab 20 mg		84	1	PSM Citalopram
ESCITALOPRAM				
* Tab 10 mg	1.11	28	1	Escitalopram- Apotex
* Tab 20 mg	1.90	28	1	Escitalopram- Apotex
FLUOXETINE HYDROCHLORIDE				
* Tab dispersible 20 mg, scored – Subsidy by endorsement Subsidised by endorsement	2.47	30	1	Arrow-Fluoxetine
 When prescribed for a patient who cannot swallow accordingly; or 	whole tablets or caps	sules	and the pr	escription is endorsed
 When prescribed in a daily dose that is not a multi endorsed. Note: Tablets should be combined wit 	, v		•	
* Cap 20 mg	1.99	90	1	Arrow-Fluoxetine
PAROXETINE * Tab 20 mg	4.02	90	1	Apo-Paroxetine
SERTRALINE				
 * Tab 50 mg * Tab 100 mg 		90 90		Arrow-Sertraline Arrow-Sertraline

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Cubaidu		Fully Drand or	
	Subsidy (Manufacturer's Price		Fully Brand or Subsidised Generic	
	(Manulaciulei S Frice \$	Per	Manufacturer	
Other Antidepressants				
MIRTAZAPINE				
Tab 30 mg	2.63	30	 <u>Apo-Mirtazapine</u> 	
Tab 45 mg	3.48	30	Apo-Mirtazapine	
VENLAFAXINE				
* Cap 37.5 mg	6.38	84	Enlafax XR	
* Cap 75 mg	8.11	84	 Enlafax XR 	
* Cap 150 mg	11.16	84	 Enlafax XR 	
Antiepilepsy Drugs				
Agents for Control of Status Epilepticus				
CLONAZEPAM – Safety medicine; prescriber may determine dis	pensing frequency			
Inj 1 mg per ml, 1 ml		5	Rivotril	
DIAZEPAM – Safety medicine; prescriber may determine dispension		•		
Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement		5	 Hospira 	
	11.03	5		
a) Up to 5 inj available on a PSO				
b) Only on a PSO				
c) PSO must be endorsed "not for anaesthetic procedur		E	Ctopolid	
Rectal tubes 5 mg – Up to 5 tube available on a PSO		5	 Stesolid 	
Rectal tubes 10 mg – Up to 5 tube available on a PSO		5	 Stesolid 	
PARALDEHYDE				
* Inj 5 ml	1,500.00	5	 AFT \$29 	
PHENYTOIN SODIUM				
* Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a P	SO 88.63	5	 Hospira 	
* Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a				
PSO		5	 Hospira 	
Control of Epilepsy				
CARBAMAZEPINE			/ -	
* Tab 200 mg		100	 Tegretol 	
* Tab long-acting 200 mg		100	 Tegretol CR 	
* Tab 400 mg		100	 Tegretol 	
* Tab long-acting 400 mg		100	 Tegretol CR 	
* Oral liq 20 mg per ml		250 ml	 Tegretol 	
CLOBAZAM - Safety medicine; prescriber may determine disper	nsing frequency			
Tab 10 mg	9.12	50	 Frisium 	
CLONAZEPAM - Safety medicine; prescriber may determine dis	pensing frequency			
Oral drops 2.5 mg per ml		10 ml OF	Rivotril	
ETHOSUXIMIDE		-		
Cap 250 mg	281 75	200	 Zarontin 	
Oral lig 250 mg per 5 ml		200 ml	✓ Zarontin	
		200 111		
GABAPENTIN	alin			
Note: Not subsidised in combination with subsidised pregaba		100	Ano Cohanantin	
* Cap 100 mg		100	✓ <u>Apo-Gabapentin</u>	
* Cap 300 mg * Cap 400 mg		100	 ✓ <u>Apo-Gabapentin</u> ✓ Apo-Gabapentin 	
* Cap 400 mg		100	 Apo-Gapapentin 	

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
LACOSAMIDE - Special Authority see SA1125 below - Retail pl	harmacy			
▲ Tab 50 mg		14	✓	Vimpat
▲ Tab 100 mg		14	✓	Vimpat
	200.24	56	1	Vimpat
▲ Tab 150 mg	75.10	14	1	Vimpat
-	300.40	56	1	Vimpat
▲ Tab 200 mg		56	1	Vimpat

➡SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

LAMOTRIGINE

Editorritante			
▲ Tab dispersible 2 mg	6.74	30	 Lamictal
▲ Tab dispersible 5 mg	9.64	30	 Lamictal
	15.00	56	 Arrow-Lamotrigine
▲ Tab dispersible 25 mg	2.76	56	 Logem
	20.40		 Arrow-Lamotrigine
	29.09		 Lamictal
▲ Tab dispersible 50 mg	3.31	56	 Logem
	34.70		 Arrow-Lamotrigine
	47.89		 Lamictal
▲ Tab dispersible 100 mg	4.40	56	 Logem
	59.90		 Arrow-Lamotrigine
	79.16		 Lamictal
(Arrow-Lamotrigine Tab dispersible 50 mg to be delisted 1 Oc (Lamictal Tab dispersible 50 mg to be delisted 1 October 201 (Arrow-Lamotrigine Tab dispersible 100 mg to be delisted 1 October 20 (Lamictal Tab dispersible 100 mg to be delisted 1 October 20	9) October 2019)		
LEVETIRACETAM			
Tab 250 mg	4.99	60	 Everet
Everet to be Sole Supply on 1 August 2019			
Tab 500 mg	8.79	60	 Everet
Everet to be Sole Supply on 1 August 2019			
Tab 750 mg	14.39	60	 Everet
Everet to be Sole Supply on 1 August 2019			
Tab 1,000 mg		60	 Everet
Everet to be Sole Supply on 1 August 2019			
Oral liq 100 mg per ml	44.78	300 ml OP	 Levetiracetam-AFT

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	(Manulacialer 3 1 100) \$	Per		Manufacturer
HENOBARBITONE				
For phenobarbitone oral liquid refer Standard Formulae, pa	age 227			
F Tab 15 mg		500	1	PSM
F Tab 30 mg		500	✓	PSM
HENYTOIN SODIUM				
• Tab 50 mg		200	1	Dilantin Infatab
Cap 30 mg		200	1	Dilantin
Cap 100 mg		200	1	Dilantin
Oral liq 30 mg per 5 ml		500 m	l 🗸	Dilantin
REGABALIN				
Note: Not subsidised in combination with subsidised gaba	pentin			
🗧 Cap 25 mg	2.25	56	1	Pregabalin Pfizer
Cap 75 mg	2.65	56	✓	Pregabalin Pfizer
Cap 150 mg	4.01	56	1	Pregabalin Pfizer
Cap 300 mg	7.38	56	~	Pregabalin Pfizer
RIMIDONE				
• Tab 250 mg		100	1	Apo-Primidone
	62.00	200		Mysoline S29 S29
ODIUM VALPROATE				•
Tab 100 mg		100	1	Epilim Crushable
Tab 200 mg EC		100		Epilim
Tab 500 mg EC		100	1	Epilim
• Oral liq 200 mg per 5 ml		300 m	l 🗸	Epilim S/F Liquid
			~	Epilim Syrup
• Inj 100 mg per ml, 4 ml	41.50	1	1	Epilim IV
TIRIPENTOL – Special Authority see SA1330 below – Retail	pharmacy			
Cap 250 mg		60	1	Diacomit S29
Powder for oral lig 250 mg sachet		60	1	Diacomit S29

► SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

11.07			
11.07			
	60	↓</td <td>Arrow-Topiramate</td>	Arrow-Topiramate
			Fopiramate Actavis
26.04		🗸 1	Fopamax
18.81	60	✓	Arrow-Topiramate
		 1 	Topiramate Actavis
44.26		 1 	Горатах
31.99	60	✓	Arrow-Topiramate
		 1 	Copiramate Actavis
75.25		 1 	Горатах
55.19	60	✓	Arrow-Topiramate
		1 🗸	Fopiramate Actavis
129.85		1 🗸	Горатах
20.84	60	1 🗸	Горатах
26.04	60	1 🗸	Горатах
ν.			
	100	√ §	Sabril
•••	26.04 acy 119.30	26.04 60 acy	26.04 60 🖌 1 acy

⇒SA1072 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

1 Either:

- 1.1 Patient has infantile spasms; or
- 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Notes: ``Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages. **Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	_	Subsidised	Generic
	\$	Per	~	Manufacturer
Antimigraine Preparations				
For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 1	age 110			
Acute Migraine Treatment				
ERGOTAMINE TARTRATE WITH CAFFEINE				
Tab 1 mg with caffeine 100 mg	31.00	100	1	Cafergot
			1	Cafergot S29 S29
RIZATRIPTAN				
Tab orodispersible 10 mg	5.26	30	1	Rizamelt
SUMATRIPTAN				
Tab 50 mg		100		Apo-Sumatriptan
Tab 100 mg		100	~	Apo-Sumatriptan
Inj 12 mg per ml, 0.5 ml prefilled pen – Maximum of 10 inj pe prescription		2 OP	1	Clustran
prescription		2 01		Sun Pharma S29
			•	oun marma des
Prophylaxis of Migraine				
For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SY	STEM, page 50			
PIZOTIFEN				
₭ Tab 500 mcg	23.21	100	✓	Sandomigran
Antinous and Vorting Amonto				
Antinausea and Vertigo Agents				
For Antispasmodics refer to ALIMENTARY TRACT, page 8				
APREPITANT – Special Authority see SA0987 below – Retail ph	armacy			
Cap 2 × 80 mg and 1 × 125 mg		3 OP	✓	Emend Tri-Pack
SA0987 Special Authority for Subsidy				
,			•	
metogenic chemotherapy and/or anthracycline-based chemothe	rapy for the treatmer	nt of m	nalignancy	
emetogenic chemotherapy and/or anthracycline-based chemothe Renewal from any relevant practitioner. Approvals valid for 12 m	rapy for the treatmer onths where the pati	nt of m ent is	nalignancy	
emetogenic chemotherapy and/or anthracycline-based chemothe Renewal from any relevant practitioner. Approvals valid for 12 m themotherapy and/or anthracycline-based chemotherapy for the	rapy for the treatmer onths where the pati	nt of m ent is	nalignancy	
emetogenic chemotherapy and/or anthracycline-based chemothe Renewal from any relevant practitioner. Approvals valid for 12 m themotherapy and/or anthracycline-based chemotherapy for the BETAHISTINE DIHYDROCHLORIDE	rapy for the treatmer nonths where the pati treatment of maligna	nt of m ent is	alignancy undergoir	g highly emetogenic
emetogenic chemotherapy and/or anthracycline-based chemothe Renewal from any relevant practitioner. Approvals valid for 12 m chemotherapy and/or anthracycline-based chemotherapy for the BETAHISTINE DIHYDROCHLORIDE K Tab 16 mg	rapy for the treatmer nonths where the pati treatment of maligna	nt of m ent is ncy.	alignancy undergoir	
emetogenic chemotherapy and/or anthracycline-based chemothe Renewal from any relevant practitioner. Approvals valid for 12 m chemotherapy and/or anthracycline-based chemotherapy for the BETAHISTINE DIHYDROCHLORIDE * Tab 16 mg CYCLIZINE HYDROCHLORIDE	rapy for the treatmer onths where the pati treatment of maligna 2.89	nt of m ent is ncy.	nalignancy undergoir	g highly emetogenic <u>Vergo 16</u>
emetogenic chemotherapy and/or anthracycline-based chemothe Renewal from any relevant practitioner. Approvals valid for 12 m chemotherapy and/or anthracycline-based chemotherapy for the BETAHISTINE DIHYDROCHLORIDE ★ Tab 16 mg CYCLIZINE HYDROCHLORIDE Tab 50 mg	rapy for the treatmer onths where the pati treatment of maligna 2.89	nt of m ent is ncy. 84	nalignancy undergoir	g highly emetogenic
emetogenic chemotherapy and/or anthracycline-based chemothe Renewal from any relevant practitioner. Approvals valid for 12 m shemotherapy and/or anthracycline-based chemotherapy for the BETAHISTINE DIHYDROCHLORIDE * Tab 16 mg CYCLIZINE HYDROCHLORIDE Tab 50 mg CYCLIZINE LACTATE	rapy for the treatmer ionths where the pati treatment of maligna 2.89 0.55	nt of m ent is ncy. 84	ialignancy undergoir	g highly emetogenic <u>Vergo 16</u>
emetogenic chemotherapy and/or anthracycline-based chemothe Renewal from any relevant practitioner. Approvals valid for 12 m shemotherapy and/or anthracycline-based chemotherapy for the BETAHISTINE DIHYDROCHLORIDE * Tab 16 mg CYCLIZINE HYDROCHLORIDE Tab 50 mg CYCLIZINE LACTATE Inj 50 mg per ml, 1 ml	rapy for the treatmer ionths where the pati treatment of maligna 2.89 0.55	nt of m ent is ncy. 84 10	ialignancy undergoir	g highly emetogenic <u>Vergo 16</u> <u>Nausicalm</u>
emetogenic chemotherapy and/or anthracycline-based chemothe Renewal from any relevant practitioner. Approvals valid for 12 m chemotherapy and/or anthracycline-based chemotherapy for the BETAHISTINE DIHYDROCHLORIDE * Tab 16 mg CYCLIZINE HYDROCHLORIDE Tab 50 mg CYCLIZINE LACTATE Inj 50 mg per ml, 1 ml DOMPERIDONE	rapy for the treatmer ionths where the pati treatment of maligna 	nt of m ent is ncy. 84 10	alignancy undergoir	g highly emetogenic <u>Vergo 16</u> <u>Nausicalm</u>
emetogenic chemotherapy and/or anthracycline-based chemothe Renewal from any relevant practitioner. Approvals valid for 12 m chemotherapy and/or anthracycline-based chemotherapy for the BETAHISTINE DIHYDROCHLORIDE * Tab 16 mg CYCLIZINE HYDROCHLORIDE Tab 50 mg CYCLIZINE LACTATE Inj 50 mg per ml, 1 ml DOMPERIDONE * Tab 10 mg	rapy for the treatmer ionths where the pati treatment of maligna 	nt of m ent is ncy. 84 10 5	alignancy undergoir	g highly emetogenic <u>Vergo 16</u> <u>Nausicalm</u> Nausicalm
emetogenic chemotherapy and/or anthracycline-based chemothe Renewal from any relevant practitioner. Approvals valid for 12 m chemotherapy and/or anthracycline-based chemotherapy for the BETAHISTINE DIHYDROCHLORIDE * Tab 16 mg CYCLIZINE HYDROCHLORIDE Tab 50 mg CYCLIZINE LACTATE Inj 50 mg per ml, 1 ml DOMPERIDONE * Tab 10 mg HYOSCINE HYDROBROMIDE	rapy for the treatmer ionths where the pati treatment of maligna 	nt of m ent is ncy. 84 10 5	alignancy undergoir • •	g highly emetogenic <u>Vergo 16</u> <u>Nausicalm</u> Nausicalm
emetogenic chemotherapy and/or anthracycline-based chemothe Renewal from any relevant practitioner. Approvals valid for 12 m chemotherapy and/or anthracycline-based chemotherapy for the BETAHISTINE DIHYDROCHLORIDE * Tab 16 mg CYCLIZINE HYDROCHLORIDE Tab 50 mg CYCLIZINE LACTATE Inj 50 mg per ml, 1 ml DOMPERIDONE * Tab 10 mg HYOSCINE HYDROBROMIDE	rapy for the treatmer ionths where the pati treatment of maligna 	nt of m ent is ncy. 84 10 5 100	ialignancy undergoir	g highly emetogenic <u>Vergo 16</u> <u>Nausicalm</u> Nausicalm <u>Pharmacy Health</u>
CYCLIZINE LACTATE	rapy for the treatmer ionths where the pati treatment of maligna 	nt of m ent is ncy. 84 10 5 100 5	ialignancy undergoir	g highly emetogenic <u>Vergo 16</u> <u>Nausicalm</u> Nausicalm <u>Pharmacy Health</u> Hospira

Subsidy		Fully	Brand or	
(Manufacturer's Price)	;	Subsidised	Generic	
\$	Per	✓	Manufacturer	

➡SA1387 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE

* Tab 10 mg1.30	100	 Metoclopramide Actavis 10
Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO 13.56	10	 Link Healthcare S29 Pfizer
ONDANSETRON		
* Tab 4 mg	50	 Apo-Ondansetron
* Tab disp 4 mg0.95	10	✓ <u>Ondansetron</u> ODT-ORLA
* Tab 8 mg4.77	50	 Apo-Ondansetron
* Tab disp 8 mg1.43	10	✓ <u>Ondansetron</u> ODT-DRLA
PROCHLORPERAZINE		
* Tab 3 mg buccal	50	
(15.00)		Buccastem
* Tab 5 mg – Up to 30 tab available on a PSO	250	✓ Nausafix
* Inj 12.5 mg per ml, 1 ml – Up to 5 inj available on a PSO25.81	10	✓ Stemetil

Antipsychotics

General

AMISULPRIDE - Safety medicine; prescriber may determine d	ispensing frequenc	;y	
Tab 100 mg		30	✓ Sulprix
Tab 200 mg	14.75	60	 Sulprix
Tab 400 mg	27.70	60	✓ Sulprix
Oral liq 100 mg per ml	65.53	60 ml	✓ Solian
ARIPIPRAZOLE - Safety medicine; prescriber may determine	dispensing frequen	су	
Tab 5 mg		30	 Aripiprazole Sandoz
Tab 10 mg	17.50	30	 Aripiprazole Sandoz
Tab 15 mg	17.50	30	 Aripiprazole Sandoz
Tab 20 mg	17.50	30	 Aripiprazole Sandoz
Tab 30 mg	17.50	30	 Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; p	rescriber may dete	rmine disper	nsing frequency
Tab 10 mg – Up to 30 tab available on a PSO		100	 Largactil
Tab 25 mg – Up to 30 tab available on a PSO		100	 Largactil
Tab 100 mg – Up to 30 tab available on a PSO		100	 Largactil
Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO	25.66	10	 Largactil

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per	•	Manufacturer
CLOZAPINE – Hospital pharmacy [HP4]				
Safety medicine; prescriber may determine dispensing frequ	•	50		Ola-aril
Tab 25 mg		50		Clozaril
	6.69 11.36	100		Clopine Clozaril
	13.37	100		Clopine
Tab 50 mg		50		Clopine
Tab 50 mg	17.33	100		Clopine
Tab 100 mg		50		Clozaril
	17.33	50		Clopine
	29.45	100		Clozaril
	34.65	100		Clopine
Tab 200 mg		50		Clopine
140 200 mg	69.30	100		Clopine
Suspension 50 mg per ml		100 m		Clopine
		100 11		elephile
HALOPERIDOL – Safety medicine; prescriber may determine d Tab 500 mcg – Up to 30 tab available on a PSO		100	1	Serenace
Tab 1.5 mg – Up to 30 tab available on a PSO		100		Serenace
Tab 5 mg – Up to 30 tab available on a PSO		100		Serenace
Oral lig 2 mg per ml – Up to 200 ml available on a PSO		100 m		Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a P		100 11		Serenace
LEVOMEPROMAZINE HYDROCHLORIDE - Safety medicine;				
Inj 25 mg per ml, 1 ml ampoule		10		Wockhardt
LEVOMEPROMAZINE MALEATE - Safety medicine; prescriber	r may determine disp	ensing		
Tab 25 mg	16.10	100	1	Nozinan
Nozinan to be Sole Supply on 1 September 2019				
Tab 100 mg	41.75	100	1	Nozinan
Nozinan to be Sole Supply on 1 September 2019				
LITHIUM CARBONATE - Safety medicine; prescriber may dete	rmine dispensing free	quency	/	
Tab 250 mg		500	1	Lithicarb FC
Tab long-acting 400 mg		100	✓	Priadel
Cap 250 mg	9.42	100	1	Douglas
OLANZAPINE - Safety medicine; prescriber may determine dis	oensina freauency			
Tab 2.5 mg		28	1	Zypine
Tab 5 mg		28		Zypine
Tab orodispersible 5 mg		28		Zypine ODT
Tab 10 mg		28	-	Zypine
Tab orodispersible 10 mg		28		Zypine ODT
PERICYAZINE – Safety medicine; prescriber may determine dis				
Tab 2.5 mg		84	1	Neulactil
Tab 2.5 mg	12.49	100		Neulactil
Tab 10 mg		84		Neulactil
	44.45	100		Neulactil
OLIETIADINE Sofaty medicing, properitor may determine dia		100	•	
QUETIAPINE – Safety medicine; prescriber may determine disp		90		Quatanal
Tab 25 mg				Quetapel
Tab 100 mg		90 90		Quetapel Quetapel
Tab 200 mg		90 90		Quetapel Quetapel
Tab 300 mg		90	•	auelapel

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Sub Per	sidised	Generic
	\$	Per	1	Manufacturer
RISPERIDONE - Safety medicine; prescriber may determine	dispensing frequency			
Tab 0.5 mg		60	✓ <u>A</u>	ctavis
Tab 1 mg	2.06	60	🗸 A	ctavis
Tab 2 mg	2.29	60	🗸 🗸	ctavis
Tab 3 mg	2.50	60	🗸 🗸	ctavis
Tab 4 mg	3.43	60	🗸 🗸	ctavis
Oral lig 1 mg per ml	7.66	30 ml	🗸 🖌	lisperon
IPRASIDONE – Safety medicine; prescriber may determine	dispensing frequency		_	
Cap 20 mg		60	√ 7	usdone
Cap 40 mg		60	_	usdone
Cap 60 mg		60		usdone
Cap 80 mg		60	_	usdone
			_	
CUCLOPENTHIXOL HYDROCHLORIDE – Safety medicine; p	,		0 1	,
Tab 10 mg		100	• 0	lopixol
Denet Injections				
Depot Injections				
LUPENTHIXOL DECANOATE - Safety medicine; prescriber	r may determine dispen	sina freau	encv	
Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO		5 5		luanxol
Inj 20 mg per ml, 2 ml – Up to 5 inj available on a PSO		5		luanxol
Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO		5	-	luanxol
		-	-	launixon
ALOPERIDOL DECANOATE – Safety medicine; prescriber		•		
Inj 50 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		aldol
Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		aldol Concentrate
			• H	aldol
				Decanoas S29
LANZAPINE – Special Authority see SA1428 below – Retail	pharmacy			
Safety medicine; prescriber may determine dispensing fre	quency			
Inj 210 mg vial		1	🗸 Z	yprexa Relprevv
Inj 300 mg vial		1	✓ Z	yprexa Relprevv
				vprexa Relprevv

➡SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE - Special Authority see SA1429 on the next page - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Inj 25 mg syringe	1	Invega Sustenna
Inj 50 mg syringe	1	Invega Sustenna
Inj 75 mg syringe	1	Invega Sustenna
Inj 100 mg syringe	1	Invega Sustenna
Inj 150 mg syringe	1	 Invega Sustenna

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE - Special Authority see SA1427 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Inj 25 mg vial	1	Risperdal Consta
Inj 37.5 mg vial	1	 Risperdal Consta
lnj 50 mg vial217.56	1	 Risperdal Consta

➡SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE – Safety medicine; prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml – Up to 5 inj available on a	PSO19.80	5	 Clopixol
Anxiolytics			
BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	20.23	100	✓ <u>Orion</u>
* Tab 10 mg		100	✓ <u>Orion</u>
CLONAZEPAM - Safety medicine; prescriber may dete	rmine dispensing frequency		
Tab 500 mcg	5.64	100	✓ <u>Paxam</u>
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determin	ne dispensing frequency		
Tab 2 mg		500	 <u>Arrow-Diazepam</u>
Tab 5 mg		500	 Arrow-Diazepam

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
LORAZEPAM – Safety medicine; prescriber may determine dispertine	0 1 2	250		Ativan
Tab 1 mg Tab 2.5 mg		250 100		Ativan
OXAZEPAM - Safety medicine; prescriber may determine disper	0 1 2			
Tab 10 mg Tab 15 mg		100 100		Ox-Pam Ox-Pam

Multiple Sclerosis Treatments

DIMETHYL FUMARATE - Special Authority see SA1559 below - Retail pharmacy

Wastage claimable

Cap 120 mg		14	 Tecfidera
Cap 240 mg	2,000.00	56	 Tecfidera

⇒SA1559 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator	Phone: 04 460 4990
Multiple Sclerosis Treatment Assessment Committee	Facsimile: 04 916 7571
PHARMAC PO Box 10 254	Email: <u>mstaccoordinator@pharmac.govt.nz</u>

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified). Entry Criteria

- a) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- b) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- c) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- d) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;

Subsidy	0.1	Fully	Brand or	
(Manufacturer's Price) \$	Per	osidised ✓	Generic Manufacturer	

- e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
- f) be distinguishable from the effects of general fatigue; and
- g) not be associated with a fever (T> 37.5° C); and
- e) applications must be made by the patient's neurologist or general physician; and
- f) patients must have no previous history of lack of response to dimethyl fumarate; and
- g) patients must have not previously had intolerance to dimethyl fumarate; and
- h) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to dimethyl fumarate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

FINGOLIMOD - Special Authority see SA1562 below - Retail pharmacy

Wastage claimable Cap 0.5 mg

ap 0.5 mg	2,200.00	28	🗸 Gilenya
-----------	----------	----	-----------

► SA1562 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator	Phone: 04 460 4990
Multiple Sclerosis Treatment Assessment Committee	Facsimile: 04 916 7571
PHARMAC PO Box 10 254	Email: mstaccoordinator@pharmac.govt.nz
Wellington	

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified). Entry Criteria

S	Subsidy	Fully	Brand or
(Manufa	acturer's Price) Subsi	dised	Generic
	\$ Per	1	Manufacturer

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) patients must have no previous history of lack of response to fingolimod; and
- 7) patients must have not previously had intolerance to fingolimod; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to fingolimod; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🖌	Manufacturer	

a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

NATALIZUMAB – Special Authority see SA1563 below – Retail pharmacy

Inj 20 mg per ml, 15 ml vial...... 1,750.00 1 🗸 Tysabri

⇒SA1563 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator	Phone: 04 460 4990
Multiple Sclerosis Treatment Assessment Committee	Facsimile: 04 916 7571
PHARMAC PO Box 10 254	Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) treatment must be initiated and supervised by a neurologist who is registered in the Tysabri Australasian Prescribing Programme operated by the supplier; and
- 7) patients must have no previous history of lack of response to natalizumab; and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

- 8) patients must have not previously had intolerance to natalizumab; and
- 9) a) Patient is JC virus negative, or
 - b) Patient is JC virus positive and has given written informed consent acknowledging an understanding of the risk of progressive multifocal leucoencephalopathy (PML) associated with natalizumab
- 10) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to natalizumab; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier.

Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate.

Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

TERIFLUNOMIDE – Special Authority see SA1560 below – Retail pharmacy

Wastage claimable	•		
Tah 14 mg		1 582 62	28

lab	14 mg	1,582.62	28	Aubagio

⇒SA1560 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator	Phone: 04 460 4990
Multiple Sclerosis Treatment Assessment Committee	Facsimile: 04 916 7571
PHARMAC PO Box 10 254	Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) patients must have no previous history of lack of response to teriflunomide; and
- 7) patients must have not previously had intolerance to teriflunomide; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to teriflunomide; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping

Subsidy		Fully	Brand or	
(Manufacturer's Price	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

Other Multiple Sclerosis Treatments

⇒SA1564 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator	Phone: 04 460 4990
Multiple Sclerosis Treatment Assessment Committee	Facsimile: 04 916 7571
PHARMAC PO Box 10 254	Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only glatiramer acetate inj 40 mg prefilled syringe will be subsidised if dispensed in a community pharmacy. The other agents will NOT be subsidised if dispensed from a community or hospital pharmacy. Regular supplies will be distributed to all approved patients or their clinicians by courier.

Prescribers must send quarterly prescriptions for approved patients to the MSTAC coordinator.

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 20 mg glatiramer acetate daily or 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. The MSTAC coordinator should be notified of the change and a new prescription provided.

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	✓	Manufacturer

- d) start at least one month after the onset of a previous relapse;
- e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
- f) be distinguishable from the effects of general fatigue; and
- g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or
 - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment. Progression of disability is defined as progress by any of the following EDDSS Points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

GLATIRAMER ACETATE - Special Authority see SA1564 on the previous page - Retail pharmacy

2,250.00	28	 Copaxone
2,275.00	12	 Copaxone
SA1564 on the pr	evious page	
1,170.00	4	 Avonex
1,170.00	4	 Avonex Pen
SA1564 on the prev	vious page	
1,322.89	15	 Betaferon
	1,170.00 1,170.00	2,275.00 12 SA1564 on the previous page 1,170.00 4 1,170.00 4 SA1564 on the previous page

Subsidy (Manufacturer's Price		Fully Subsidised	
(ivianuiacturer s Frice \$	Per		Manufacturer
Sedatives and Hypnotics			
MELATONIN – Special Authority see SA1666 below – Retail pharmacy	00	(Oireadin
Tab modified-release 2 mg – No more than 5 tab per day	30	•	Circadin
SA1666 Special Authority for Subsidy Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist. applications meeting the following criteria: All of the following:			
 Patient has been diagnosed with persistent and distressing insomnia seconda (including, but not limited to, autism spectrum disorder or attention deficit hyp Behavioural and environmental approaches have been tried and were unsucc Funded modified-release melatonin is to be given at doses no greater than 10 Patient is aged 18 years or under*. 	eractivi cessful,) mg pe	ity disorde , or are ina er day; and	r)*; and ppropriate; and I
Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid following criteria:			
All of the following:			
 Patient is aged 18 years or under*; and Patient has demonstrated clinically meaningful benefit from funded modified-r Patient has had a trial of funded modified-release melatonin discontinuation w 			
recurrence of persistent and distressing insomnia; and)	ar day	
4 Funded modified-release melatonin is to be given at doses no greater than 10 Note: Indications marked with * are unapproved indications.	ng pe	er uay.	
MIDAZOLAM – Safety medicine; prescriber may determine dispensing frequency			
Inj 1 mg per ml, 5 ml ampoule	10	1	Midazolam-Claris
Inj 1 mg per ml, 5 ml plastic ampoule – Up to 10 inj available on a PSO14.90	10	1	Pfizer
On a PSO for status epilepticus use only. PSO must be endorsed for status	s epilep		
Inj 5 mg per ml, 3 ml ampoule2.50	5	~	Midazolam-Claris
Inj 5 mg per ml, 3 ml plastic ampoule – Up to 5 inj available on a PSO11.90	5	1	Pfizer
On a PSO for status epilepticus use only. PSO must be endorsed for status	-		
NITRAZEPAM – Safety medicine; prescriber may determine dispensing frequency			,
Tab 5 mg	100	1	Nitrados
PHENOBARBITONE SODIUM - Special Authority see SA1386 below - Retail phare	nacy		
Inj 200 mg per ml, 1 ml ampoule	5	✓	Aspen S29
■SA1386 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further ren the following criteria: Both:	iewal u	nless notif	ied for applications meeting
 For the treatment of terminal agitation that is unresponsive to other agents; at The applicant is part of a multidisciplinary team working in palliative care. 	nd		
TEMAZEPAM – Safety medicine; prescriber may determine dispensing frequency Tab 10 mg1.27	25	1	Normison

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
TRIAZOLAM - Safety medicine; prescriber may determine disp	ensing frequency			
Tab 125 mcg		100		
	(9.85)		H	Hypam
Tab 250 mcg	4.10	100		
	(11.20)		ŀ	Hypam
ZOPICLONE - Safety medicine; prescriber may determine disp	pensing frequency			
Tab 7.5 mg	0 1 2	500	✓ 2	Zopiclone Actavis
Stimulants/ADHD Treatments ATOMOXETINE – Special Authority see SA1416 below – Reta	il pharmacy			
Cap 10 mg		28		Strattera
Cap 18 mg		28	√ 9	Strattera
Cap 25 mg		28	√ 9	Strattera
Cap 40 mg	107.03	28	√ 9	Strattera
Cap 60 mg		28	√ 9	Strattera
Cap 80 mg		28	✓ 9	Strattera
Cap 100 mg		28	√ 9	Strattera

⇒SA1416 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder) diagnosed according to DSM-IV or ICD 10 criteria; and
- 2 Once-daily dosing; and
- 3 Any of the following:
 - 3.1 Treatment with a subsidised formulation of a stimulant has resulted in the development or worsening of serious adverse reactions or where the combination of subsidised stimulant treatment with another agent would pose an unacceptable medical risk; or
 - 3.2 Treatment with a subsidised formulation of a stimulant has resulted in worsening of co-morbid substance abuse or there is a significant risk of diversion with subsidised stimulant therapy; or
 - 3.3 An effective dose of a subsidised formulation of a stimulant has been trialled and has been discontinued because of inadequate clinical response; or
 - 3.4 Treatment with a subsidised formulation of a stimulant is considered inappropriate because the patient has a history of psychoses or has a first-degree relative with schizophrenia; and
- 4 The patient will not be receiving treatment with atomoxetine in combination with a subsidised formulation of a stimulant, except for the purposes of transitioning from subsidised stimulant therapy to atomoxetine.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: A "subsidised formulation of a stimulant" refers to currently subsidised methylphenidate hydrochloride tablet formulations (immediate-release, sustained-release and extended-release) or dexamfetamine sulphate tablets.

DEXAMFETAMINE SULFATE - Special Authority see SA1149 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency
- Tab 5 mg20.00 100 🗸 PSM

➡SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1150 below - Retail pharmacy

a) Only on a controlled drug form

 b) Safety medicine; prescriber may determine dispensing frequer 	b) Safe	tv medicine: p	rescriber may	/ determine di	ispensina freauency	
---	---------	----------------	---------------	----------------	---------------------	--

Tab immediate-release 5 mg	30	 Rubifen
Tab immediate-release 10 mg	30	 Ritalin
°		 Rubifen
Tab immediate-release 20 mg7.85	30	 Rubifen
Tab sustained-release 20 mg	30	Rubifen SR
50.00	100	 Ritalin SR

⇒SA1150 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application - (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for

Subsidy		Fully	Brand or	Ī
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	~	Manufacturer	

continued...

applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1151 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab extended-release 18 mg58.9	96 30 Concerta
Tab extended-release 27 mg65.4	44 30 ✓ Concerta
Tab extended-release 36 mg71.9	93 30 ✓ Concerta
Tab extended-release 54 mg	
Cap modified-release 10 mg 15.6	
Cap modified-release 20 mg	
Cap modified-release 30 mg25.8	
Cap modified-release 40 mg	

⇒SA1151 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

	Subsidy	0.1	Fully	Brand or
	(Manufacturer's Price) \$	Per	sidised ✓	Generic Manufacturer
continued				
1 The treatment remains appropriate and the patient is ben 2 Either:	efiting from treatment;	and		
2.1 Applicant is a paediatrician or psychiatrist; or2.2 Applicant is a medical practitioner and confirms th last 2 years and has recommended treatment for the		sychiatris	st has be	en consulted within the
MODAFINIL – Special Authority see SA1126 below – Retail pha Tab 100 mg		60	✓ N	lodaviqil
■ SA1126 Special Authority for Subsidy Initial application only from a neurologist or respiratory specialis following criteria: All of the following:	st. Approvals valid for	24 mon	ths for ap	oplications meeting the
 The patient has a diagnosis of narcolepsy and has exces almost daily for three months or more; and Either: 	sive daytime sleepines	ss assoc	iated wit	n narcolepsy occurring
2.1 The patient has a multiple sleep latency test with a more sleep onset rapid eye movement periods; or2.2 The patient has at least one of: cataplexy, sleep p				
 3 Either: 3.1 An effective dose of a subsidised formulation of m discontinued because of intolerable side effects; o 3.2 Methylphenidate and dexamfetamine are contrain 	r dicated.			
Renewal only from a neurologist or respiratory specialist. Appro and the patient is benefiting from treatment.	vais valiu ioi 24 monti	is where	e une urea	ument remains appropriate
Treatments for Dementia				

* Tab 5 mg * Tab 10 mg		90 90	 ✓ <u>Donepezil-Rex</u> ✓ <u>Donepezil-Rex</u>
RIVASTIGMINE - Special Authority see SA1488 below - Reta	il pharmacy		
Patch 4.6 mg per 24 hour		30	 Exelon
Patch 9.5 mg per 24 hour	90.00	30	 Exelon

➡SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 on the next page - Retail pharmacy

a) No patient co-payment payable

b) Safety medicine; prescriber may determine dispensing frequency		
Tab sublingual 2 mg with naloxone 0.5 mg57.40	28	 Suboxone
Tab sublingual 8 mg with naloxone 2 mg 166.00	28	 Suboxone

*Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from

any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

Tab modified-release 150 mg	11.00	30	 Zyban
DISULFIRAM			
Tab 200 mg	75.57	100	 Antabuse
NALTREXONE HYDROCHLORIDE - Special Authority see SA1	408 below – Reta	il pharmacy	
Tab 50 mg	112.55	30	✓ Naltraccord
- CA1409 Enocial Authority for Subsidy			

SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

- continued...
 - 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
 - 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

a) Nicotine will not be funded in amounts less than 4 weeks of treatment.

b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

b) Note. Direct Provision by a pharmacist permitted under the provisions in	Fart For Section	JII A.
Patch 7 mg – Up to 28 patch available on a PSO	28	 Habitrol
Patch 7 mg for direct distribution only - [Xpharm]	7	 Habitrol
Patch 14 mg – Up to 28 patch available on a PSO 17.59	28	 Habitrol
Patch 14 mg for direct distribution only - [Xpharm]4.52	7	 Habitrol
Patch 21 mg – Up to 28 patch available on a PSO20.16	28	 Habitrol
Patch 21 mg for direct distribution only - [Xpharm]5.18	7	 Habitrol
Lozenge 1 mg - Up to 216 loz available on a PSO16.61	216	 Habitrol
Lozenge 1 mg for direct distribution only - [Xpharm]	36	 Habitrol
Lozenge 2 mg - Up to 216 loz available on a PSO	216	✓ Habitrol
Lozenge 2 mg for direct distribution only - [Xpharm]	36	 Habitrol
Gum 2 mg (Fruit) – Up to 384 piece available on a PSO	384	 Habitrol
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]8.64	96	 Habitrol
Gum 2 mg (Mint) – Up to 384 piece available on a PSO	384	 Habitrol
Gum 2 mg (Mint) for direct distribution only - [Xpharm]8.64	96	 Habitrol
Gum 4 mg (Fruit) – Up to 384 piece available on a PSO	384	 Habitrol
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]10.01	96	 Habitrol
Gum 4 mg (Mint) – Up to 384 piece available on a PSO	384	 Habitrol
Gum 4 mg (Mint) for direct distribution only - [Xpharm]10.01	96	✓ Habitrol

VARENICLINE TARTRATE – Special Authority see SA1771 below – Retail pharmacy

a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
 b) Varenicline will not be funded in amounts less than 4 weeks of treatment.

Tab 0.5 mg × 11 and 1 mg × 42	25.64	53 OP	 Varenicline Pfizer
Tab 1 mg	27.10	56	 Varenicline Pfizer

➡SA1771 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and

3 Either:

3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at

(Subsidy Manufacturer's Price)	Subsid	Fully lised	Brand or Generic
	\$	Per	1	Manufacturer

continued...

- least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
- 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not used funded varenicline in the last 12 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 The patient has not used funded varenicline in the last 12 months; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 12 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer
Chemotherapeutic Agents			
Alkylating Agents			
BENDAMUSTINE HYDROCHLORIDE – PCT only – Specialist – Inj 25 mg vial Inj 100 mg vial Inj 1 mg for ECP		1 ✓ 1 ✓	Ribomustin Ribomustin Baxter
► SA1667 Special Authority for Subsidy		•	
Initial application - (treatment naive CLL) only from a relevant			the recommendation of a
relevant specialist. Approvals valid for 12 months for applications All of the following:	s meeting the following	g criteria:	
1 The patient has Binet stage B or C, or progressive stage A	chronic lymphocytic	loukaamia raa	iring treatment: and
2 The patient has bliet stage b of C, or progressive stage 7		ieukaeiilla ieyt	inning treatment, and
3 The patient is unable to tolerate toxicity of full-dose FCR;	and		
4 Patient has ECOG performance status 0-2; and			
5 Patient has a Cumulative Illness Rating Scale (CIRS) score			
6 Bendamustine is to be administered at a maximum dose of	of 100 mg/m ² on days	1 and 2 every 4	weeks for a maximum of
6 cycles. Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymp	booutio lumphoma (SI	I) Chomothou	any treatment is considered
to comprise a known standard therapeutic chemotherapy regimer			apy treatment is considered
Initial application — (Indolent, Low-grade lymphomas) only f			ractitioner on the
recommendation of a relevant specialist. Approvals valid for 9 m	onths for applications	meeting the fol	lowing criteria:
All of the following:			
1 The patient has indolent low grade NHL requiring treatment	nt; and		
2 Patient has a WHO performance status of 0-2; and			
3 Either:			
3.1 Both:			
3.1.1 Patient is treatment naive; and3.1.2 Bendamustine is to be administered for a m	avimum of 6 cycles (ii	n combination v	with rituximah when
CD20+); or		reombination	
3.2 All of the following:			
3.2.1 Patient has relapsed refractory disease follo	wing prior chemother	apy; and	
3.2.2 The patient has not received prior bendamu	istine therapy; and		
3.2.3 Either:			
3.2.3.1 Both:			
3.2.3.1.1 Bendamustine is to be adminis combination with rituximab wh		of 6 cycles in re	elapsed patients (in
3.2.3.1.2 Patient has had a rituximab tre	atment-free interval o	f 12 months or	more; or
3.2.3.2 Bendamustine is to be administered refractory patients.	as a monotherapy for	a maximum of	6 cycles in rituximab

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Either:
 - 2.1 Both:

continued...

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Pric		Subsidised	Brand or Generic
	\$	Per		Manufacturer
ontinued				
2.1.1 Bendamustine is to be administered for	a maximum of 6 cycle	s in relap	sed patients	(in combination with
rituximab when CD20+); and				
2.1.2 Patient has had a rituximab treatment-fro			-	
2.2 Bendamustine is to be administered as a mono				
lote: 'indolent, low-grade lymphomas' includes follicular, man nacroglobulinaemia.	ntle cell, marginal zon	e and lym	phoplasmac	ytic/ Waldenstrom's
BUSULFAN – PCT – Retail pharmacy-Specialist				
Tab 2 mg		100	🗸 My	leran
ARBOPLATIN – PCT only – Specialist				
Inj 10 mg per ml, 45 ml vial		1	🗸 DB	L Carboplatin
	45.20			rboplatin Ebewe
	48.50			rbaccord
Inj 1 mg for ECP	0.10	1 mg	🗸 Ba	xter
ARMUSTINE – PCT only – Specialist				
Inj 100 mg vial	532.00	1	🗸 BiO	NU
	1,380.00		🖌 Em	cure S29
Inj 100 mg for ECP	1,380.00	100 mg O	P 🖌 🖌 Ba	xter
BiCNU Inj 100 mg vial to be delisted 1 July 2019)				
HLORAMBUCIL – PCT – Retail pharmacy-Specialist				
Tab 2 mg		25	🖌 Lei	ukeran FC
ISPLATIN – PCT only – Specialist				
Inj 1 mg per ml, 50 ml vial		1	🗸 DB	L Cisplatin
	15.00		🗸 Cis	platin Ebewe
Inj 1 mg per ml, 100 ml vial	19.70	1		L Cisplatin
	21.00			platin Ebewe
Inj 1 mg for ECP	0.25	1 mg	🗸 Ba	xter
YCLOPHOSPHAMIDE				
Tab 50 mg – PCT – Retail pharmacy-Specialist	79.00	50	🖌 En	doxan S29
- · · ·	158.00	100	🗸 Pro	ocytox S29
Wastage claimable				-
Inj 1 g vial – PCT – Retail pharmacy-Specialist	35.65	1	🖌 En	doxan
	127.80	6	🗸 Cy	
Inj 2 g vial – PCT only – Specialist		1	✓ En	
Inj 1 mg for ECP – PCT only – Specialist	0.04	1 mg	🗸 Ba	xter
OSFAMIDE – PCT only – Specialist				
lnj 1 g		1	🗸 Ho	
lnj 2 g		1	✓ Ho	
Inj 1 mg for ECP	0.10	1 mg	🗸 Ba	xter
OMUSTINE – PCT – Retail pharmacy-Specialist				
Cap 10 mg	132.59	20	🗸 Ce	
Cap 40 mg		20	🗸 Ce	eNU
IELPHALAN				
Tab 2 mg – PCT – Retail pharmacy-Specialist	40.70	25	🗸 Alk	
Inj 50 mg – PCT only – Specialist	67.80	1	🖌 Alk	eran

()	Subsidy /anufacturer's Price) \$	Per	Fully Subsidised	
OXALIPLATIN – PCT only – Specialist				
Inj 100 mg vial	25.01	1	~	Oxaliplatin Actavis 100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	46.32	1	~	Oxaliccord
Inj 1 mg for ECP	0.48	1 mg	∕ √	Baxter
THIOTEPA – PCT only – Specialist				
Inj 15 mg vial	CBS	1	1	Bedford S29
			1	THIO-TEPA S29
			1	Tepadina S29
Inj 100 mg vial	CBS	1		Tepadina S29
Antimetabolites				
AZACITIDINE - PCT only - Specialist - Special Authority see SA1	467 below			
Inj 100 mg vial		1	~	Azacitidine Dr Reddy's
	605.00		1	Vidaza
Inj 1 mg for ECP	1.53	1 mg	· ✓	Baxter

► SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
 - 2 The patient has performance status (WHO/ECOG) grade 0-2; and
 - 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
 - 4 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

- Both:
 - 1 No evidence of disease progression; and
 - 2 The treatment remains appropriate and patient is benefitting from treatment.

	Subsidy	\ <u>•</u>	Fully Brand or
(Manufacturer's Pri \$	ce) Subs Per	idised Generic Manufacturer
LCIUM FOLINATE	•		
Tab 15 mg – PCT – Retail pharmacy-Specialist	104.26	10	 DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml – PCT – Retail pharmacy-Specialist		5	✓ Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specialis		1	 Calcium Folinate Sandoz
Inj 50 mg - PCT - Retail pharmacy-Specialist		5	 Calcium Folinate Ebewe
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	7.30	1	 Calcium Folinate Sandoz
Inj 100 mg - PCT only - Specialist	7.33	1	 Calcium Folinate Ebewe
Inj 300 mg - PCT only - Specialist	22.51	1	 Calcium Folinate Ebewe
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	20.95	1	 Calcium Folinate Sandoz
Inj 1 g – PCT only – Specialist	67.51	1	 Calcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	60.00	1	 Calcium Folinate Sandoz
Inj 1 mg for ECP – PCT only – Specialist	0.06	1 mg	✓ Baxter
PECITABINE – Retail pharmacy-Specialist		-	
Tab 150 mg	11.15	60	 Brinov
Tab 500 mg	62.28	120	 Brinov
ADRIBINE – PCT only – Specialist			
Inj 1 mg per ml, 10 ml		7	 Leustatin
Inj 10 mg for ECP	749.96	10 mg OP	 Baxter
TARABINE			
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialis	t400.00	5	 Pfizer
Inj 100 mg per ml, 20 ml vial – PCT – Retail			
pharmacy-Specialist		1	✓ Pfizer
Inj 1 mg for ECP – PCT only – Specialist		10 mg	✓ Baxter
Inj 100 mg intrathecal syringe for ECP – PCT only – Specialis	t80.00	100 mg OP	 Baxter
UDARABINE PHOSPHATE			
Tab 10 mg – PCT – Retail pharmacy-Specialist		20	Fludara Oral
Inj 50 mg vial – PCT only – Specialist		5	✓ Fludarabine Ebewe
Inj 50 mg for ECP – PCT only – Specialist	105.00	50 mg OP	 Baxter
UOROURACIL			_
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist		1	 Fluorouracil Ebewe
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist		1	 Fluorouracil Ebewe
Inj 1 mg for ECP – PCT only – Specialist	0.66	100 mg	 Baxter
MCITABINE HYDROCHLORIDE – PCT only – Specialist			
Inj 1 g, 26.3 ml vial		1	 DBL Gemcitabine
lnj 1 g		1	 Gemcitabine Ebewe
	349.20		 Gemzar
Inj 200 mg		1	 Gemzar
Inj 1 mg for ECP	0.02	1 mg	 Baxter

_					
		Subsidy		Fully	Brand or
		(Manufacturer's Pri		sidised	Generic
		\$	Per		Manufacturer
RI	NOTECAN HYDROCHLORIDE – PCT only – Specialist				
	Inj 20 mg per ml, 5 ml vial	71.44	1	✓	Irinotecan
					Accord S29
				I	Irinotecan Actavis
					100
		100.00		1	Irinotecan-Rex
	Inj 1 mg for ECP		1 mg		Baxter
	RCAPTOPURINE				
		07.00	25		Puri-nethol
	Tab 50 mg – PCT – Retail pharmacy-Specialist		25	•	Puri-neuroi
	Puri-nethol to be Sole Supply on 1 July 2019				
	Oral suspension 20 mg per ml – Retail pharmacy-Specialist				
	Special Authority see SA1725 below		100 ml OP	•	Allmercap
	SA1725 Special Authority for Subsidy				
ni	tial application only from a paediatric haematologist or paedia	tric oncologist. A	pprovals valid	d for 12	2 months where the patien
ec	uires a total dose of less than one full 50 mg tablet per day.				
Re	newal only from a paediatric haematologist or paediatric oncol	ogist. Approvals	valid for 12 m	nonths	where patient still requires
a to	otal dose of less than one full 50 mg tablet per day.				
ЛE	THOTREXATE				
	Tab 2.5 mg – PCT – Retail pharmacy-Specialist		90	1	Trexate
	Tab 10 mg - PCT - Retail pharmacy-Specialist		90		Trexate
	Inj 2.5 mg per ml, 2 ml – PCT – Retail pharmacy-Specialist		5		Hospira
	Inj 7.5 mg prefilled syringe		1		Methotrexate
•			·		Sandoz
*	Inj 10 mg prefilled syringe	14 66	1	1	Methotrexate
					Sandoz
*	Inj 15 mg prefilled syringe	14 77	1	1	Methotrexate
T	Ing 15 mg premied synnige		I	•	Sandoz
	lai 00 ma avefilled avriane	14.00			
*	Inj 20 mg prefilled syringe	14.88	1	•	Methotrexate
					Sandoz
*	Inj 25 mg prefilled syringe	14.99	1	✓	Methotrexate
					Sandoz
*	Inj 30 mg prefilled syringe	15.09	1	✓	Methotrexate
					Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Speciali	st30.00	5	✓]	DBL Methotrexate
					Onco-Vial
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specia	llist45.00	1	✓	DBL Methotrexate
					Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist	t25.00	1	I	Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail				
	pharmacy-Specialist	79.99	1	1	Methotrexate Ebewe
*	Inj 1 mg for ECP – PCT only – Specialist		1 mg		Baxter
*	Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist.		5 mg OP		Baxter
			0		
۲E	METREXED - PCT only - Specialist - Special Authority see				lune Demotranal
	Inj 100 mg vial		1		Juno Pemetrexed
	Inj 500 mg vial		1		Juno Pemetrexed
	Inj 1 mg for ECP	0.55	1 mg	•	Baxter

Subsidy (Manufacturer's Price)	Full Subsidise	d Generic
\$	Per 🖌	Manufacturer

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and

2 Either:

- 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
- 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

- All of the following:
 - 1 No evidence of disease progression; and
 - 2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

THIOGUANINE - PCT - Retail pharmacy-Specialist

Tab 40 mg126.31	25	 Lanvis
Other Cytotoxic Agents		
AMSACRINE – PCT only – Specialist		
Inj 50 mg per ml, 1.5 ml ampoule1,500.00) 6	 Amsidine S29
Inj 75 mg1,250.00) 5	AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE – PCT – Retail pharmacy-Specialist		
Cap 0.5 mgCBS	100	🗸 Agrylin S29
		 Teva S29
ARSENIC TRIOXIDE – PCT only – Specialist		
Inj 1 mg per ml, 10 ml vial) 10	 Phenasen
Inj 10 mg4,817.00) 10	✓ AFT \$29
Inj 10 mg for ECP481.70) 10 mg OP	 Baxter
(AFT \$29 Inj 10 mg to be delisted 1 September 2019)		

	Subsidy (Manufacturer's Price \$) Sub Per	Fully sidised	Brand or Generic Manufacturer
BLEOMYCIN SULPHATE – PCT only – Specialist Inj 15,000 iu, vial	161.01	1	✓ D	BL Bleomycin Sulfate
Inj 1,000 iu for ECP	12.45	1,000 iu	🗸 В	axter
BORTEZOMIB – PCT only – Specialist – Special Authority see S/ Inj 3.5 mg vial Inj 1 mg for ECP	1,892.50	1 1 mg	•	elcade eaxter

⇒SA1576 Special Authority for Subsidy

Initial application — (Treatment naive multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 Either:

- 1.1 The patient has treatment-naive symptomatic multiple myeloma; or
- 1.2 The patient has treatment-naive symptomatic systemic AL amyloidosis *; and
- 2 Maximum of 9 treatment cycles.
- Note: Indications marked with * are unapproved indications.

Initial application — (Relapsed/refractory multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has relapsed or refractory multiple myeloma; or
 - 1.2 The patient has relapsed or refractory systemic AL amyloidosis *; and
- 2 The patient has received only one prior front line chemotherapy for multiple myeloma or amyloidosis; and
- 3 The patient has not had prior publicly funded treatment with bortezomib; and
- 4 Maximum of 4 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Relapsed/refractory multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

1 The patient's disease obtained at least a partial response from treatment with bortezomib at the completion of cycle 4; and

2 Maximum of 4 further treatment cycles (making a total maximum of 8 consecutive treatment cycles).

Notes: Responding relapsed/refractory multiple myeloma patients should receive no more than 2 additional cycles of treatment beyond the cycle at which a confirmed complete response was first achieved. A line of therapy is considered to comprise either:

- a) a known therapeutic chemotherapy regimen and supportive treatments; or
- b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments.

Refer to datasheet for recommended dosage and number of doses of bortezomib per treatment cycle.

COLASPASE [L-ASPARAGINASE] - PCT only - Specialist

Inj 10,000 iu	102.32	1	 Leunase
Inj 10,000 iu for ECP	102.32	10,000 iu OP	 Baxter
DACARBAZINE – PCT only – Specialist			
Inj 200 mg vial	58.06	1	 DBL Dacarbazine
	580.60	10	 Dacarbazine
			APP S29
Inj 200 mg for ECP	58.06	200 mg OP	 Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist			
Inj 0.5 mg vial	166.75	1	 Cosmegen
Inj 0.5 mg for ECP	166.75	0.5 mg OP	 Baxter

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturar's Price	o) Cirk	Fully	
	(Manufacturer's Pric \$	e) Sub Per	sidised ✓	I Generic Manufacturer
AUNORUBICIN – PCT only – Specialist				
Inj 2 mg per ml, 10 ml	130.00	1	✓	Pfizer
Inj 20 mg for ECP	130.00	20 mg OP	✓	Baxter
OCETAXEL - PCT only - Specialist		-		
Inj 10 mg per ml, 2 ml vial		1	✓	DBL Docetaxel
Inj 20 mg		1	1	Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial		1	✓	DBL Docetaxel
Inj 80 mg		1	1	Docetaxel Sandoz
Inj 1 mg for ECP	0.55	1 mg	✓	Baxter
OXORUBICIN HYDROCHLORIDE – PCT only – Specialist				
Inj 2 mg per ml, 5 ml vial		1	1	Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial	11.50	1	✓	Doxorubicin Ebewe
	17.00		1	Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial		1	1	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial	56.15	1	1	Doxorubicin Ebewe
	65.00		✓	Arrow-Doxorubicin
Inj 1 mg for ECP	0.29	1 mg	1	Baxter
PIRUBICIN HYDROCHLORIDE – PCT only – Specialist				
Inj 2 mg per ml, 5 ml vial		1	✓	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1		Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1		Epirubicin Ebewe
Inj 1 mg for ECP	0.37	1 mg	1	Baxter
TOPOSIDE				
Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	1	Vepesid
Vepesid to be Sole Supply on 1 July 2019				
Cap 100 mg - PCT - Retail pharmacy-Specialist		10	1	Vepesid
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specialis		1		Rex Medical
Inj 1 mg for ECP – PCT only – Specialist	0.09	1 mg	1	Baxter
OPOSIDE PHOSPHATE - PCT only - Specialist				
Inj 100 mg (of etoposide base)	40.00	1	1	Etopophos
Inj 1 mg (of etoposide base) for ECP	0.47	1 mg	✓	Baxter
DROXYUREA – PCT – Retail pharmacy-Specialist				
Cap 500 mg		100	1	Hydrea
ARUBICIN HYDROCHLORIDE				•
Inj 5 mg vial – PCT only – Specialist	93.00	1	1	Zavedos
Inj 10 mg vial – PCT only – Specialist		1		Zavedos
Inj 1 mg for ECP – PCT only – Specialist		1 mg		Baxter
ENALIDOMIDE – Retail pharmacy-Specialist – Special Authority		Ũ	-	
Wastage claimable	y SEE SA 1400 DEIC	JVV		
Cap 10 mg	6 207 00	21	1	Revlimid
Cap 15 mg		21		Revlimid
Cap 25 mg		21		Revlimid
SA1468 Special Authority for Subsidy	1,021.00	21	•	

Initial application — (Relapsed/refractory disease) only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Patient has relapsed or refractory multiple myeloma with progressive disease; and

Subsidy	Fu	lly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
\$	Per	~	Manufacturer

continued...

2 Either:

2.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or

2.2 Both:

2.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and

- 2.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 3 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 No evidence of disease progression; and

2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MESNA

Tab 400 mg – PCT – Retail pharmacy-Specialist	50	 Uromitexan
Tab 600 mg – PCT – Retail pharmacy-Specialist	50	 Uromitexan
Inj 100 mg per ml, 4 ml ampoule – PCT only – Specialist	15	 Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist	15	 Uromitexan
Inj 1 mg for ECP – PCT only – Specialist2.69	100 mg	 Baxter
MITOMYCIN C – PCT only – Specialist		
Inj 5 mg vial204.08	1	 Arrow
Inj 1 mg for ECP	1 mg	✓ Baxter
MITOZANTRONE – PCT only – Specialist		
Inj 2 mg per ml, 10 ml vial	1	 Mitozantrone Ebewe
Inj 1 mg for ECP5.51	1 mg	 Baxter
PACLITAXEL – PCT only – Specialist	-	
Inj 30 mg	5	Paclitaxel Ebewe
Inj 100 mg20.00	1	Paclitaxel Ebewe
91.67		Paclitaxel Actavis
Inj 150 mg26.69	1	 Paclitaxel Ebewe
137.50		Anzatax
		 Paclitaxel Actavis
Inj 300 mg35.35	1	 Paclitaxel Ebewe
275.00		 Anzatax
		 Paclitaxel Actavis
Inj 1 mg for ECP0.19	1 mg	 Baxter
PEGASPARGASE – PCT only – Special Authority see SA1325 below		
Inj 3,750 IU per 5 ml	1	 Oncaspar S29

SA1325 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Subsi Per	idised	Generic Manufacturer
	Ŧ			
continued	and the second			
 The patient has newly diagnosed acute lymphoblastic leuk Pegaspargase to be used with a contemporary intensive new patient of the second second	,	rapy treatn	nent pr	otocol; and
3 Treatment is with curative intent.	0		·	
Renewal only from a relevant specialist or medical practitioner or for 12 months for applications meeting the following criteria: All of the following:	n the recommendation	n of a relev	ant sp	ecialist. Approvals valid
 The patient has relapsed acute lymphoblastic leukaemia; Pegaspargase to be used with a contemporary intensive n Treatment is with curative intent. 		rapy treatn	nent pr	otocol; and
PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Specialis	t			
Inj 10 mg		1	🗸 N	lipent S29
PROCARBAZINE HYDROCHLORIDE – PCT – Retail pharmacy	-Specialist			•
Cap 50 mg	•	50	🗸 N	latulan S29
TEMOZOLOMIDE - Special Authority see SA1741 below - Reta	il pharmacy			
Cap 5 mg		5	✓ <u>c</u>	Prion
				Temozolomide
Cap 20 mg		5	✓ <u>c</u>	<u>Prion</u>
				Temozolomide
Con 100 mg	40.00	5		emizole 20 ^(\$29) Drion
Cap 100 mg		5	• <u>u</u>	Temozolomide
Cap 140 mg		5	√ 0	rion
		-	-	Temozolomide
Cap 250 mg	96.80	5	✓ <u>c</u>	Prion
				Temozolomide

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient has glioblastoma multiforme; and
- 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special Authority	see SA1124 belo	w	
Cap 50 mg		28	 Thalomid
Cap 100 mg	756.00	28	 Thalomid

■SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Either:

1 The patient has multiple myeloma; or

2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

Cap 10 mg – PCT – Retail pharmacy-Specialist	100	 Vesanoid
VINBLASTINE SULPHATE Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Specialist 186.46 Inj 1 mg for ECP – PCT only – Specialist	5 1 mg	✓ Hospira✓ Baxter
VINCRISTINE SULPHATE Inj 1 mg per ml, 1 ml vial – PCT – Retail pharmacy-Specialist74.52	5	✓ DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist85.61	5	 DBL Vincristine Sulfate
Inj 1 mg for ECP – PCT only – Specialist11.30	1 mg	✓ Baxter

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

(Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	Generic
VINORELBINE – PCT only – Specialist				
Inj 10 mg per ml, 1 ml vial	12.00 42.00	1		Navelbine Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial	56.00 210.00	1	-	Navelbine Vinorelbine Ebewe
Inj 1 mg for ECP	1.25	1 mg	1	Baxter
Protein-tyrosine Kinase Inhibitors				
DASATINIB – Special Authority see SA1805 below – Retail pharm Wastage claimable	acy			
Tab 20 mg	3,774.06	60	1	Sprycel
Tab 50 mg	6,214.20	60	1	Sprycel
Tab 70 mg		60	1	Sprycel
SA1805 Special Authority for Subsidy				

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
 - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
 - 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB - Retail pharmacy-Specialist - Special Authority see SA1653 on the next page

Tab 100 mg	 30	 Tarceva
Tab 150 mg	 30	 Tarceva

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
SA1653 Special Authority for Subsidy				
nitial application only from a relevant specialist or medical p	practitioner on the recom	mend	ation of a re	levant specialist.
Approvals valid for 4 months for applications meeting the follo				
Ill of the following:	-			
1 Patient has locally advanced or metastatic, unresectal				
2 There is documentation confirming that the disease ex	presses activating muta	tions o	of EGFR tyre	osine kinase; and
3 Either:				
3.1 Patient is treatment naive; or3.2 Both:				
3.2.1 The patient has discontinued gefitinib d	up to intoloronoo; and			
3.2.2 The cancer did not progress while on ge				
4 Erlotinib is to be given for a maximum of 3 months.				
Renewal only from a relevant specialist or medical practitione	er on the recommendation	on of a	relevant sp	ecialist. Approvals valid
or 6 months where radiological assessment (preferably inclu				
GEFITINIB - Retail pharmacy-Specialist - Special Authority	see SA1654 below			
Tab 250 mg	1,700.00	30	🗸 li	ressa
SA1654 Special Authority for Subsidy				
nitial application only from a relevant specialist or medical		nmend	ation of a re	elevant specialist.
Approvals valid for 4 months for applications meeting the follo	wing criteria:			
All of the following:	hla waa amuunaa Naw	Omel		
 Patient has locally advanced, or metastatic, unresecta Either: 	ble, non-squamous Non	Smail	I Cell Lung (Jancer (NSCLC); and
2.1 Patient is treatment naive; or				
2.2 Both:				
2.2.1 The patient has discontinued erlotinib d	ue to intolerance; and			
2.2.2 The cancer did not progress whilst on e				
3 There is documentation confirming that disease expre	sses activating mutation	s of E	GFR tyrosin	e kinase; and
4 Gefitinib is to be given for a maximum of 3 months.				
Renewal only from a relevant specialist or medical practitione				
or 6 months where radiological assessment (preferably inclu-	aing CT scan) indicates	NSCL	C nas not p	rogressea.
MATINIB MESILATE Note: Imatinib-AFT is not a registered for the treatment of	of Coatro Intentinal Strar			T) The Clives brand of
imatinib mesilate (supplied by Novartis) remains fully sub				
metastatic malignant GIST, see SA1460 in Section B of t			y for pation	
Tab 100 mg - [Xpharm] - Special Authority see SA1460				
below	,	60		Blivec
₭ Cap 100 mg		60		matinib-AFT
₭ Cap 400 mg		30	✓ <u>li</u>	matinib-AFT
■ SA1460 Special Authority for Subsidy				
Special Authority approved by the CML/GIST Co-ordinator lotes: Application details may be obtained from PHARMAC	eele elte letter //ele			al muse substitutes also utal la

Notes: Application details may be obtained from PHARMAC's website <u>http://www.pharmac.govt.nz</u>, and prescriptions should be sent to:

Subsidy (Manufacturer's Price)	Fi Subsidis	ully	Brand or Generic
\$	Per	✓	Manufacturer

continued...

The CML/GIST Co-ordinator	Phone: (04) 460 4990
PHARMAC	Facsimile: (04) 916 7571
PO Box 10 254	Email: cmlgistcoordinator@pharmac.govt.nz
Wellington	

Special Authority criteria for GIST - access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA1191 below - Retail pharmacy

Tab 250 mg 1,899.00 70 🗸 Tykerb

⇒SA1191 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient has not previously received trastuzumab treatment for HER 2 positive metastatic breast cancer; and
 - 1.3 Lapatinib not to be given in combination with trastuzumab; and
 - 1.4 Lapatinib to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2.2 The patient started trastuzumab for metastatic breast cancer but discontinued trastuzumab within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on trastuzumab; and
 - 2.4 Lapatinib not to be given in combination with trastuzumab; and
 - 2.5 Lapatinib to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 on the next page - Retail pharmacy

Wastage claimable			
Cap 150 mg4	,680.00	120 🖌	Tasigna
Cap 200 mg6	,532.00	120 🖌	Tasigna

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and 2 Either:

2.1 Patient has documented CML treatment failure* with imatinib; or

2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and

- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PAZOPANIB - Special Authority see SA1190 below - Retail pharmacy

Tab 200 mg1,334.70) 30	Votrient
Tab 400 mg2,669.40) 30	 Votrient

⇒SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
 - The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

	Subsidy (Manufacturer's Price) \$	Per		
RUXOLITINIB - Special Authority see SA1753 below - Retail ph	armacy			
Wastage claimable			_	
Tab 5 mg	2,500.00	56	~	Jakavi
Tab 15 mg	5,000.00	56	✓	Jakavi
Tab 20 mg	5,000.00	56	✓	Jakavi
- CA1752 Created Authority for Subaidy				

⇒SA1753 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA1266 below - Retail pharmacy

Cap 12.5 mg	2,315.38	28	 Sutent
Cap 25 mg	4,630.77	28	 Sutent
Cap 50 mg	9,261.54	28	 Sutent

➡SA1266 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
 - The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application - (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

specialist. Approvals valid for 3 months for applications meeting the following criteria:

- Both:
 - 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
 - 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 82

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA1767 below

Wastage claimable

Tab 250 mg4,276.19 120 🗸 Zytiga

► SA1767 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
- 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

BICALUTAMIDE

Tab 50 mg		28	 Binarex
FLUTAMIDE – Retail pharmacy-Specialist			
Tab 250 mg		30	 Flutamide
			Mylan S29
	46.20	84	 Flutamide
			Mylan S29
	55.00	100	 Flutamin
MEGESTROL ACETATE – Retail pharmacy-Specialist			
Tab 160 mg	63.53	30	Apo-Megestrol
OCTREOTIDE			
Inj 50 mcg per ml, 1 ml vial		5	DBL Octreotide
Inj 100 mcg per ml, 1 ml vial		5	✓ DBL Octreotide
Inj 500 mcg per ml, 1 ml vial	72.50	5	✓ DBL Octreotide
OCTREOTIDE LAR (SOMATOSTATIN ANALOGUE) - S		16 below –	Retail pharmacy
Inj LAR 10 mg prefilled syringe	1,772.50	1	 Sandostatin LAR
Inj LAR 20 mg prefilled syringe		1	 Sandostatin LAR
Inj LAR 30 mg prefilled syringe	2,951.25	1	 Sandostatin LAR

➡SA1016 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal - (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment

	Subsidy	Fu	ully	Brand or
(M	anufacturer's Price)	Subsidis	ed	Generic
	\$	Per	✓	Manufacturer

continued...

remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma; and
 - 2.2 Either:

2.2.1 Patient has failed surgery; or

2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or

- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

* Tab 10 mg 11.75 * Tab 20 mg 5.60	60 60	 ✓ Tamoxifen Sandoz ✓ Tamoxifen Sandoz
Aromatase Inhibitors		
ANASTROZOLE * Tab 1 mg5.04	30	✓ <u>Rolin</u>
EXEMESTANE * Tab 25 mg14.50	30	✓ Pfizer Exemestane

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

4.68	30	1	Letrole
	100 5 ml C	✓ ✓ ✓ ✓	Imuran Imuran Celicept Celicept Celicept Celicept and capsules, and when
rmacy 799.96	4		Enbrel Enbrel
	25.00 25.00 187.25 16 patients unable to rmacy	25.00 50 25.00 100 187.25 165 ml C patients unable to swallo rmacy 799.96 4	25.00 50 • 25.00 100 • 187.25 165 ml OP • patients unable to swallow tablets rmacy 799.96 4 •

Initial application — (juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for juvenile idiopathic arthritis (JIA): and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for JIA: or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
 - 2.2 Patient diagnosed with Juvenile Idiopathic Arthritis (JIA); and
 - 2.3 Patient has had severe active polvarticular course JIA for 6 months duration or longer; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
 - 2.5 Both:

2.5.1 Either:

- 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender ioints: or
- 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

2.5.2 Physician's global assessment indicating severe disease.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

continued...

- 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
- 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. **Initial application — (ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm

25-34 years - Male: 7.5 cm; Female: 5.5 cm

35-44 years - Male: 6.5 cm; Female: 4.5 cm

	ONCOLOGY AGENTS AN		JPPRESSANIS
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✔	Brand or Generic Manufacturer
continued			
45-54 years - Male: 6.0 cm; Female: 5.0 cm			
55-64 years - Male: 5.5 cm; Female: 4.0 cm			
65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm			
Initial application — (psoriatic arthritis) only	v from a rheumatologist Approvals vali	d for 6 months fo	r applications meeting the
following criteria:	internationalitic spirit and the		
Either:			
1 Both:			
1.1 The patient has had an initial Sp1.2 Either:	ecial Authority approval for adalimumab	o for psoriatic arth	ritis; and
	ced intolerable side effects from adalimit insufficient benefit from adalimumab to		l criteria for adalimumab
2 All of the following:			
2.1 Patient has had severe active pa	soriatic arthritis for six months duration o	or longer; and	
	ded to at least three months of oral or pa	arenteral methotr	exate at a dose of at leas
20 mg weekly or a maximum tole			
	ded to at least three months of sulfasala		at least 2 g per day or
2.4 Either:) mg daily (or maximum tolerated doses); and	
	mptoms of poorly controlled and active of	disaasa in at laas	t 15 swollon tonderigints
Or	inploms of poony controlled and active c		
2.4.2 Patient has persistent sy	mptoms of poorly controlled and active of	disease in at leas	t four joints from the
following: wrist, elbow, k	nee, ankle, and either shoulder or hip; a	Ind	
2.5 Any of the following:			
2.5.1 Patient has a C-reactive date of this application; o	protein level greater than 15 mg/L meas r	sured no more that	an one month prior to the
2.5.3 ESR and CRP not measu	erythrocyte sedimentation rate (ESR) greater as patient is currently receiving pre- ne so for more than three months.		
Initial application — (pyoderma gangrenosu meeting the following criteria: All of the following:	Im) only from a dermatologist. Approva	als valid for 4 mo	nths for applications

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 4 doses.

Note: Indications marked with * are unapproved indications.

Initial application - (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

1 Both:

- 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and

Subsidy	F	ully Br	and or
(Manufacturer's F	Price) Subsidi	sed Ge	eneric
\$	Per	 Ma 	anufacturer

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a named specialist or rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
 - 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
 - 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Both:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

continued...

- 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
- 2.1.2 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
- 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.
- Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
 - 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
 - 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
 - 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
 - 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
 - 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 4 doses.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

Both:

1 Either:

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) – PCT only – Specialist Inj 50 mg per ml, 5 ml2,351.25	5	✔ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE – PCT only – Specialist Subsidised only for bladder cancer.		
Inj 2-8 × 100 million CFU	1	✓ OncoTICE
Inj 40 mg per ml, vial162.70	3	 SII-Onco-BCG §29
(SII-Onco-BCG 529 Inj 40 mg per ml, vial to be delisted 1 January 2020)		
Monoclonal Antibodies		
ADALIMUMAB – Special Authority see SA1742 below – Retail pharmacy		
Inj 20 mg per 0.4 ml prefilled syringe1,599.96	2	 Humira
Inj 40 mg per 0.8 ml prefilled pen1,599.96	2	 HumiraPen
Inj 40 mg per 0.8 ml prefilled syringe1,599.96	2	 Humira

⇒SA1742 Special Authority for Subsidy

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis; or

2 All of the following:

- 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

continued...

2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and

2.6 Either:

- 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
- 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or

2 All of the following:

2.1 Either:

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

continued...

- 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
- 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. **Initial application — (ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm

25-34 years - Male: 7.5 cm; Female: 5.5 cm

35-44 years - Male: 6.5 cm; Female: 4.5 cm

		Subsidy (Manufacturer's Price) \$	Fully Subsidised Per 🖌	Brand or Generic Manufacturer
ontinued				
,	5.0 cm; Female: 5.0 cm			
	5.5 cm; Female: 4.0 cm			
	4.0 cm; Female: 4.0 cm 0 cm; Female: 2.5 cm			
	(psoriatic arthritis) only from a rhe	umatologist Approvals vali	d for 6 months fo	r applications meeting the
ollowing criteria:				
Either:				
1 Both:				
1.1 The pate 1.2 Either:	tient has had an initial Special Autho	rity approval for etanercept f	or psoriatic arthri	tis; and
1.2.2	The patient has experienced intolera The patient has received insufficient psoriatic arthritis; or			criteria for etanercept for
2 All of the follow	ving:			
2.1 Patient	has had severe active psoriatic arthr	ritis for six months duration c	or longer; and	
	has tried and not responded to at least		arenteral methotr	exate at a dose of at leas
	weekly or a maximum tolerated dose			
	has tried and not responded to at lean nide at a dose of up to 20 mg daily (at least 2 g per day or
2.4 Either:	nide at a dose of up to 20 mg daily (<i>)</i> , and	
2.4.1	Patient has persistent symptoms of p	poorly controlled and active of	disease in at leas	t 15 swollen, tender joints
	Patient has persistent symptoms of p following: wrist, elbow, knee, ankle,			t four joints from the
2.5 Any of	the following:			
	Patient has a C-reactive protein leve date of this application; or	I greater than 15 mg/L meas	sured no more that	an one month prior to the
2.5.2 2.5.3	Patient has an elevated erythrocyte ESR and CRP not measured as pati 5 mg per day and has done so for m	ent is currently receiving pre		
	(juvenile idiopathic arthritis) only is meeting the following criteria:	from a named specialist or r	heumatologist. A	Approvals valid for 6

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for juvenile idiopathic arthritis (JIA); and 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for juvenile idiopathic arthritis; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient diagnosed with JIA; and
 - 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

2.5 Both:

2.5.1 Either:

- 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
- 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
- 2.5.2 Physician's global assessment indicating severe disease.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has confirmed Crohn's disease; and

2 Either:

- 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
- 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's disease.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 4 doses.

Note: Note: Indications marked with * are unapproved indications.

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either: 1 Both:

1.1 Fither:

- 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
- 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

	Subsidy	Fully	Brand or
(Manufa	acturer's Price) Subs	idised	Generic
	\$ Per	1	Manufacturer

continued...

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and

4 Either:

- 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
- 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a

gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 CDAI score is 150 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
 - 2 Either:
 - 2.1 Either:
 - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 PCDAI score is 15 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fu	ully	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
\$	Per	✓	Manufacturer

continued...

- 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
 - 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
 - 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
 - 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a named specialist or rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
 - 2 Either:
 - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 4 doses.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.
- AFLIBERCEPT Special Authority see SA1772 on the next page Retail pharmacy

Inj 40 mg per ml, 0.1 ml vial	1,250.00	1	🗸 Eylea
-------------------------------	----------	---	---------

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

► SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or

2 Either:

- 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
- 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 on the next page

Inj 5 mg per ml, 20 ml vial		1	 Erbitux
Inj 5 mg per ml, 100 ml vial		1	🗸 Erbitux
Inj 1 mg for ECP	3.82	1 mg	 Baxter

	Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic
	\$	Per	✓	Manufacturer
SA1697 Special Authority for Subsidy Initial application only from a medical oncologist or medical	al practitioner on the recom	mendation	ofam	edical oncologist
11 , 5				ourour orroorogion
Approvals valid for 6 months for applications meeting the fo All of the following:				ourour onconogion

- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB - PCT only - Special Authority see SA1778 below

Inj 100 mg	 1	 Remicade
Inj 1 mg for ECP	 1 mg	 Baxter

► SA1778 Special Authority for Subsidy

Initial application — (Graft vs host disease) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis; or
 - 2.4 Severe ocular inflammation; or
 - 2.5 Chronic ocular inflammation; or
 - 2.6 Crohn's disease (adults); or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease; or
 - 2.9 Severe fulminant ulcerative colitis; or
 - 2.10 Severe ulcerative colitis; or
 - 2.11 Plaque psoriasis; or
 - 2.12 Neurosarcoidosis; or
 - 2.13 Severe Behcet's disease.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- in:
- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for psoriatic arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for psoriatic arthritis.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
- 2 Any of the following:
 - 2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptomst; or
 - 2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2 Any of the following:
 - 2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

continued...

- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

- All of the following:
 - 1 Paediatric patient has severe active Crohn's disease; and
 - 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
 - 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
 - 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
 - 5 Patient must be reassessed for continuation after 3 months of therapy.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 Detient has soufi
 - 1 Patient has confirmed Crohn's disease; and
 - 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria: Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (severe ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- All of the following:
 - 1 Patient has histologically confirmed ulcerative colitis; and
 - 2 Either:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
 - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65; and
 - 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
 - 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria: Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plaque psoriasis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or

Subsidy	Fi	ully	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
\$	Per	✓	Manufacturer

continued...

- 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist.

Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and

2 Either:

- 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
- 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Subsidy		Fully	Brand or
(Manufacturer's Pric	ce) Su	ubsidised	Generic
\$	Per	1	Manufacturer

continued...

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 The patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema), following 12 months' treatment; or</p>
- 3 The patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old, following 12 months' treatment.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 The patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema), following 12 months' treatment; or</p>
- 3 The patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old, following 12 months' treatment.

Subsidy	Fu	ully Brand or	
(Manufacturer's P	rice) Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

- Both:
 - 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
 - 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Renewal — (severe ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy	Fully Subsidised	Brand or
 (Manufacturer's Price) \$	Per ✓	Generic Manufacturer

continued...

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Either:

- 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
- 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria: Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Either:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

OBINUTUZUMAB - PCT only - Specialist - Special Autho	rity see SA1627 on the	next page	
Inj 25 mg per ml, 40 ml vial		1	🖌 Gazyva
Inj 1 mg for ECP	6.21	1 mg	 Baxter

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1627 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^{9} /L and platelets greater than or equal to 75×10^{9} /L.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

Inj 150 mg prefilled syringe	1	🗸 Xolair
Inj 150 mg vial	1	🗸 Xolair

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and

6 Either:

- 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
- 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
- 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
- 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and

4 Either:

- 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
- 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Patient has previously adequately responded* to 6 doses of omalizumab; or

2 Both:

- 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
- 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PERTUZUMAB – PCT only – Specialist – Special Authority see SA1606 below

Inj 30 mg per ml, 14 ml vial	 1	 Perjeta
Inj 420 mg for ECP	 420 mg OP	 Baxter

➡SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal --- (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB - PCT only - Specialist - Special Authority see SA1783 below

Inj 100 mg per 10 ml vial		 Mabthera
Inj 500 mg per 50 ml vial		 Mabthera
Inj 1 mg for ECP	5.64 1 mg	Baxter

⇒SA1783 Special Authority for Subsidy

Initial application — (Antibody-mediated renal transplant rejection) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated renal transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (ABO-incompatible renal transplant) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible renal transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with rituximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 haemophilia with inhibitors; or
 - 2.2 rheumatoid arthritis; or
 - 2.3 severe cold haemagglutinin disease (CHAD); or
 - 2.4 warm autoimmune haemolytic anaemia (warm AIHA); or
 - 2.5 immune thrombocytopenic purpura (ITP); or
 - 2.6 thrombotic thrombocytopenic purpura (TTP); or
 - 2.7 pure red cell aplasia (PRCA); or
 - 2.8 ANCA associated vasculitis; or
 - 2.9 treatment refractory systemic lupus erythematosus (SLE); or
 - 2.10 steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS).

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Either:

1 Both:

- 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant. **Initial application — (aggressive CD20 positive NHL)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.
- Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 The patient is rituximab treatment naive; and
- 3 Either:
 - 3.1 The patient is chemotherapy treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
 - 3.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; and
- 4 The patient has good performance status; and
- 5 The patient does not have chromosome 17p deletion CLL; and
- 6 Rituximab to be administered in combination with fludarabine and cyclophosphamide or bendamustine for a maximum of 6 treatment cycles; and
- 7 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2. **Initial application — (rheumatoid arthritis - prior TNF inhibitor use)** only from a rheumatologist or Practitioner on the

recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

1 Both:

Su	ubsidy	Fully	Brand or
(Manufac	cturer's Price) Subsi	dised	Generic
	\$ Per	1	Manufacturer

continued...

- 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and

2 Either:

- 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.
- **Initial application (rheumatoid arthritis TNF inhibitors contraindicated)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria: Both:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms.

Subsidy	Ful	ly Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	 Manufacturer 	

continued...

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria: Both:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin.
- Note: Indications marked with * are unapproved indications.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre; or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy).

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria: Either:

- 1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
- 2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or

continued...

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	Ful	y Brand or
(Manufacturer's F	Price) Subsidise	d Generic
\$	Per	Manufacturer

continued...

3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy. Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	 ✓ 	Manufacturer	

continued...

Note: Indications marked with * are unapproved indications.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
- 2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
- 3 The patient does not have chromosome 17p deletion CLL; and
- 4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide or bendamustine for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and

3 Either:

- 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and 4 Maximum of two 1.000 mg infusions of rituximab given two weeks apart.
- 4 maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	1	Manufacturer	

continued...

the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physiciann; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and

3 Either:

- 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria: All of the following:

1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only

from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB - Special Authority see SA1754 on the next page - Retail pharmacy

Inj 150 mg per ml, 1 ml prefilled syringe......1,599.00

Cosentyx

2

Sub		Fully	Brand or
(Manufactu		Ibsidised	Generic
	6 Per	✓	Manufacturer

⇒SA1754 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Either:

- 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
- 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
- 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 on the next page - Retail pharmacy

Note: Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Inj 100 mg vial	 1	 Sylvant
Inj 400 mg vial	 1	 Sylvant

Subsidy		Fully	Brand or
 (Manufacturer's Price)	Subsi	dised	Generic
\$	Per	✓	Manufacturer

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB - PCT only - Special Authority see SA1781 below

Inj 20 mg per ml, 4 ml vial	 Actemra
Inj 20 mg per ml, 10 ml vial	 Actemra
Inj 20 mg per ml, 20 ml vial	 Actemra
Inj 1 mg for ECP	 Baxter

⇒SA1781 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1331 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses; or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or

Subsidy		Fully	Brand or
(Manufacturer's P	rice)	Subsidised	Generic
\$	Pe	r 🖌	Manufacturer

continued...

- 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- Both:
 - 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
 - 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither

- 1 Both:
 - 1.1 Either:

Subsidy		Fully	Brand or
(Manufacturer's Price)) 5	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
- 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule; and

1.2 Either:

- 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
- 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for juvenile idiopathic arthritis (JIA); and
- 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
 - 2.4 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.5 Both:
 - 2.5.1 Either:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

Inj 150 mg vial		1	 Herceptin
Inj 440 mg vial		1	 Herceptin
Inj 1 mg for ECP	9.36	1 mg	 Baxter

➡SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of

 e)	Fully Subsidised	Brand or Generic	
\$ Per	1	Manufacturer	

continued...

at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and

- 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria: All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

continued...

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price)	ę	Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

Programmed Cell Death-1 (PD-1) Inhibitors

NIVOLUMAB – PCT only – Specialist – Special Authority see SA1656 below		
Inj 10 mg per ml, 4 ml vial1,051.98	1	 Opdivo
Inj 10 mg per ml, 10 ml vial2,629.96	1	 Opdivo
Inj 1 mg for ECP27.62	1 mg	 Baxter

⇒SA1656 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma stage III or IV; and
- 2 Patient has measurable disease as defined by the presence of at least one CT or MRI measurable lesion; and
- 3 The patient has ECOG performance score of 0-2; and

4 Either:

- 4.1 Patient has not received funded pembrolizumab; or
- 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Nivolumab is to be used at a maximum dose of 3 mg/kg every 2 weeks for a maximum of 12 weeks (6 cycles); and
- 6 Baseline measurement of overall tumour burden is documented (see Note); and
- 7 Documentation confirming that the patient has been informed and acknowledges that the initial funded treatment period of nivolumab will not be continued beyond 12 weeks (6 cycles) if their disease progresses during this time.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note; or
 - 1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.3 Patient has stable disease according to RECIST criteria (see Note); and
- 2 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; and
- 3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; and
- 5 Nivolumab will be used at a maximum dose of 3 mg/kg every 2 weeks for a maximum of 12 weeks (6 cycles).

Notes: Disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Target lesion measurements should be assessed using CT or MRI imaging with the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

• Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB – PCT only – Specialist – Special Authority see SA1657 below

Inj 50 mg vial	 	 2,340.00	1	 Keytruda
Inj 1 mg for ECP	 	 	1 mg	 Baxter

➡SA1657 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma stage III or IV; and
- 2 Patient has measurable disease as defined by the presence of at least one CT or MRI measurable lesion; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Pembrolizumab is to be used at a maximum dose of 2 mg/kg every 3 weeks for a maximum of 12 weeks (4 cycles); and
- 6 Baseline measurement of overall tumour burden is documented (see Note); and
- 7 Documentation confirming that the patient has been informed and acknowledges that the initial funded treatment period of pembrolizumab will not be continued beyond 12 weeks (4 cycles) if their disease progresses during this time.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.3 Patient has stable disease according to RECIST criteria (see Note); and
- 2 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; and
- 3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; and
- 5 Pembrolizumab will be used at a maximum dose of 2 mg/kg every 3 weeks for a maximum of 12 weeks (4 cycles).

Notes: Disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Target lesion measurements should be assessed using CT or MRI imaging with the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

CICLOSPORIN

Cap 25 mg		50	Neoral
Cap 50 mg		50	 Neoral
Cap 100 mg		50	 Neoral
Oral liq 100 mg per ml		50 ml OP	 Neoral
VEROLIMUS – Special Authority see SA1491 below – Reta Wastage claimable	il pharmacy		
Tab 10 mg	6,512.29	30	 Afinitor
Tab 5 mg		30	 Afinitor

⇒SA1491 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

F١

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

SIROLIMUS - Special Authority see SA0866 below - Retail pharmacy

Tab 1 mg	749.99	100	 Rapamune
Tab 2 mg	1,499.99	100	 Rapamune
Oral liq 1 mg per ml	449.99	60 ml OP	 Rapamune

⇒SA0866 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min; or
- · Rapidly progressive transplant vasculopathy; or
- · Rapidly progressive obstructive bronchiolitis; or
- HUS or TTP; or

210

- Leukoencepthalopathy; or
- Significant malignant disease

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
TACROLIMUS – Special Authority see SA1745 belo	ow – Retail pharmacy			
Cap 0.5 mg		100	🖌 T	acrolimus Sandoz
Cap 1 mg		100	🗸 T	acrolimus Sandoz
Cap 5 mg	070.00	50	<i>.</i> / T	acrolimus Sandoz

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

(M	Subsidy anufacturer's Price)		Fully sidised	Brand or Generic
	\$	Per	1	Manufacturer
Antiallergy Preparations				
Allergic Emergencies				
ICATIBANT – Special Authority see SA1558 below – Retail pharmad Inj 10 mg per ml, 3 ml prefilled syringe	2,668.00	1 valid for 12		i razyr s for applications meeting
 Supply for anticipated emergency treatment of laryngeal/oro- angioedema (HAE) for patients with confirmed diagnosis of C The patient has undergone product training and has agreed u Renewal from any relevant practitioner. Approvals valid for 12 mont is benefiting from treatment. 	1-esterase inhibiti pon an action pla	or deficien n for self-a	icy; and administ	ration.
Allergy Desensitisation				
Initial application only from a relevant specialist. Approvals valid for Both: 1 RAST or skin test positive; and 2 Patient has had severe generalised reaction to the sensitising Renewal only from a relevant specialist. Approvals valid for 2 years benefiting from treatment.	agent.		-	-
BEE VENOM ALLERGY TREATMENT – Special Authority see SA1	367 above – Beta	il nharma	21/	
Maintenance kit - 6 vials 120 mcg freeze dried venom, with		ii phanna	у	
diluent	285.00	1 OP	🗸 V	enomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent			_	
9 ml, 3 diluent 1.8 ml		1 OP	✓ A	•
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent		1 OP		ymenoptera S29
WASP VENOM ALLERGY TREATMENT – Special Authority see SA	A1367 above – Re	etail pharm	nacy	
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	🗸 A	lbey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze dried venom, with diluent	305.00	1 OP	✔ Н	ymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze dried venom, with diluent	305.00	1 OP	🗸 V	enomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze dried venom, with diluent	305.00	1 OP	✔ Н	ymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	🗸 A	lbey

Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze

Venomil S29

	Subsidy		Fully	Brand or
	(Manufacturer's Pri	ice) Sub	osidised	
	\$	Per	 ✓ 	Manufacturer
Antihistamines				
CETIRIZINE HYDROCHLORIDE				
* Tab 10 mg	1.01	100	1	Zista
* Oral liq 1 mg per ml	2.99	200 ml	1	Histaclear
CHLORPHENIRAMINE MALEATE				
* Oral lig 2 mg per 5 ml		500 ml	1	Histafen
DEXTROCHLORPHENIRAMINE MALEATE				
* Tab 2 mg	2 02	40		
	(8.40)	-10		Polaramine
	1.01	20		
	(5.99)	_0		Polaramine
* Oral liq 2 mg per 5 ml		100 ml		
	(10.29)			Polaramine
FEXOFENADINE HYDROCHLORIDE	(<i>'</i>			
* Tab 60 mg	4 34	20		
* Tab 00 mg	(8.23)	20		Telfast
* Tab 120 mg		10		rondot
· · · · · · · · · · · · · · · · · · ·	(8.23)			Telfast
	14.22	30		
	(26.44)			Telfast
LORATADINE	(<i>'</i>			
* Tab 10 mg	1 28	100	1	Lorafix
* Oral liq 1 mg per ml		120 ml		Lorfast
PROMETHAZINE HYDROCHLORIDE				
* Tab 10 mg	1 69	50	1	Allersoothe
* Tab 10 mg		50 50		Allersoothe
* Oral lig 1 mg per 1 ml		100 ml		Allersoothe
 Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a F 		5		Hospira
Inhaled Corticosteroids		-		<u> </u>
Innaled Corticosteroids				
BECLOMETHASONE DIPROPIONATE				
Aerosol inhaler, 50 mcg per dose		200 dose OF) 🗸	Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OF) 🗸	Beclazone 50
Aerosol inhaler, 100 mcg per dose		200 dose OF		Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OF		Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free		200 dose OF	, ,	Beclazone 250
BUDESONIDE				
Powder for inhalation, 100 mcg per dose		200 dose OF	, /	Pulmicort
				Turbuhaler
Powder for inhalation, 200 mcg per dose		200 dose OF) 🗸	Pulmicort
				Turbuhaler
Powder for inhalation, 400 mcg per dose		200 dose OF) 🗸	Pulmicort
				Turbuhaler

Subsidy	Drice) Out-1	Fully Brand or
		dised Generic ✓ Manufacturer
Ψ	1.01	
4.00	100 1000 000	
		✓ Floair
		✓ Flixotide
		 Flixotide Accuhaler
		 Flixotide Accuhaler Flasir
		✓ Floair
		 ✓ Flixotide ✓ Floair
		 ✓ Floar ✓ Flixotide
	60 dose OP	 Flixotide Accuhaler
sts		
evice20.64	60 dose	
(35.80)		Foradil
. ,		
se) 10.32	60 dose OP	
,		Oxis Turbuhaler
(10.00)		
C1 00		/ Onlyne Dyserboley
		 Onbrez Breezhaler Onbrez Breezhaler
	30 dose OP	 Onbrez Breezhaler
		4 -
		 Serevent
		✓ Meterol
25.00	60 dose OP	 Serevent Accuhaler
a-Adrenocept	or Agonists	
	120 dose OP	🗸 Vannair
	120 dose OP 120 dose OP	 ✓ Vannair ✓ Symbicort
18.23 6 mcg33.74		 ✓ Vannair ✓ Symbicort Turbuhaler 100/6
6 mcg33.74		 Symbicort
6 mcg33.74	120 dose OP 120 dose OP	 ✓ Symbicort Turbuhaler 100/6 ✓ Vannair
6 mcg33.74	120 dose OP	 Symbicort Turbuhaler 100/6
6 mcg33.74	120 dose OP 120 dose OP	 ✓ Symbicort Turbuhaler 100/6 ✓ Vannair ✓ Symbicort
3 mcg33.74 21.40 3 mcg44.08	120 dose OP 120 dose OP 120 dose OP	 Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6
6 mcg33.74	120 dose OP 120 dose OP	 Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort
3 mcg33.74 21.40 3 mcg44.08	120 dose OP 120 dose OP 120 dose OP	 Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6
3 mcg33.74 	120 dose OP 120 dose OP 120 dose OP 60 dose OP	 Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12
3 mcg33.74 21.40 3 mcg44.08	120 dose OP 120 dose OP 120 dose OP	 Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort
3 mcg33.74 	120 dose OP 120 dose OP 120 dose OP 60 dose OP 30 dose OP	 Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta
3 mcg33.74 	120 dose OP 120 dose OP 120 dose OP 60 dose OP	 Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta RexAir
3 mcg33.74 	120 dose OP 120 dose OP 120 dose OP 60 dose OP 30 dose OP 120 dose OP	 Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta RexAir Seretide
3 mcg33.74 	120 dose OP 120 dose OP 120 dose OP 60 dose OP 30 dose OP	 Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta RexAir Seretide RexAir
3 mcg33.74 	120 dose OP 120 dose OP 120 dose OP 60 dose OP 30 dose OP 120 dose OP	 Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta RexAir Seretide
3 mcg33.74 	120 dose OP 120 dose OP 120 dose OP 60 dose OP 30 dose OP 120 dose OP 120 dose OP	 Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta RexAir Seretide RexAir Seretide
3 mcg33.74 	120 dose OP 120 dose OP 120 dose OP 60 dose OP 30 dose OP 120 dose OP	 Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta RexAir Seretide RexAir
3 mcg33.74 	120 dose OP 120 dose OP 120 dose OP 60 dose OP 30 dose OP 120 dose OP 120 dose OP	 Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta RexAir Seretide RexAir Seretide
	\$	4.68 120 dose OP 7.50 120 dose OP 7.50 60 dose OP 7.50 60 dose OP 7.50 60 dose OP 7.50 60 dose OP 7.50 120 dose OP 7.22 120 dose OP 13.60 120 dose OP 10.18 120 dose OP 13.60 60 dose OP ists 60 dose OP ists 60 dose OP (35.80) 60 dose OP

	Subsidy (Manufacturer's Pri \$		Fully Brand or dised Generic ✓ Manufacturer
Beta-Adrenoceptor Agonists			
SALBUTAMOL			
Oral liq 400 mcg per ml Infusion 1 mg per ml, 5 ml		150 ml 10	✓ <u>Ventolin</u>
Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO	(130.21) 12.90	5	Ventolin Ventolin
Inhaled Beta-Adrenoceptor Agonists			
SALBUTAMOL			
Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000 dose available on a PSO	3.80 2	200 dose OP	✓ Respigen
			✓ SalAir
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb	(6.00)		Ventolin
available on a PSO Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb	3.93	20	✓ <u>Asthalin</u>
available on a PSO		20	✓ Asthalin
ERBUTALINE SULPHATE Powder for inhalation, 250 mcg per dose, breath activated	27.30 2	200 dose OP	 Bricanyl Turbuhaler
Anticholinergic Agents			
PRATROPIUM BROMIDE			
Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dos available on a PSO		200 dose OP	✓ Atrovent
Nebuliser soln, 250 mcg per ml, 1 ml ampoule - Up to 40 ne	eb		
available on a PSO Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 ne		20	✓ <u>Univent</u>
available on a PSO		20	✓ <u>Univent</u>
Inhaled Beta-Adrenoceptor Agonists with Antic	holinergic Ag	ents	
ALBUTAMOL WITH IPRATROPIUM BROMIDE			
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p dose CFC-free		200 dose OP	✓ Duolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule – Up to 20 neb available on a PSO		20	✓ <u>Duolin</u>
Long-Acting Muscarinic Antagonists			
GLYCOPYRRONIUM – Subsidy by endorsement			
 a) Inhaled glycopyrronium treatment will not be subsidised i umeclidinium. 	f patient is also rec	ceiving treatme	ent with subsidised tiotropium of
 b) Glycopyrronium powder for inhalation 50 mcg per dose is having COPD using spirometry, and the prescription is er 			o have been diagnosed as

	Subsidy		Fully	Brand or
	(Manufacturer's Pri		Subsidised	Generic
	\$	Per	1	Manufacturer
TIOTROPIUM BROMIDE – Subsidy by endorsement				
 a) Tiotropium treatment will not be subsidised if patient is umeclidinium. 	also receiving treatr	nent with s	subsidised	inhaled glycopyrronium or
b) Tiotropium bromide is subsidised only for patients who prescription is endorsed accordingly. Patients who had Authority are deemed endorsed.				
Powder for inhalation, 18 mcg per dose	50.37	30 dose		Spiriva
Soln for inhalation 2.5 mcg per dose		60 dose C)P 🖌 S	Spiriva Respimat
UMECLIDINIUM – Subsidy by endorsement				
 a) Umeclidinium will not be subsidised if patient is also red tiotropium bromide. 	ceiving treatment wit	h subsidis	ed inhaled	d glycopyrronium or
b) Umeclidinium powder for inhalation 62.5 mcg per dose COPD using spirometry, and the prescription is endorse		or patients	who have	been diagnosed as having
Powder for inhalation 62.5 mcg per dose	61.50	30 dose C)P 🖌 I	ncruse Ellipta

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

■ SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 Patient has been stabilised on a long acting muscarinic antagonist; and

2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication: and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA1584	above – Retail pharmacy
Powder for Inhalation 50 mcg with indacaterol 110 mcg81.00	30 dose OP 🖌 Ultibro Breezhaler
TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA15	84 above – Retail pharmacy
Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg81.00	60 dose OP Spiolto Respimat
UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above	 Retail pharmacy

Powder for inhalation 62.5 mcg with vilanterol 25 mcg77.00 30 dose OP Anoro Ellipta

Antifibrotics

NINTEDANIB - Special Authority see SA1755 below - R	letail pharmacy		
Note: Nintedanib not subsidised in combination with	subsidised pirfenidone.		
Cap 100 mg		60 OP	 Ofev
Cap 150 mg		60 OP	 Ofev

■SA1755 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and

RESPIRATORY SYSTEM AND ALLERGIES

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA1748 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

► SA1748 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 80% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Leukotriene Receptor Antagonists

MONTELUKAST

*	Tab 4 mg5.25	28	Apo-Montelukast
	Tab 5 mg5.50	28	✓ Apo-Montelukast
	Tab 10 mg5.65	28	✓ Accord S29
	5		Apo-Montelukast

RESPIRATORY SYSTEM AND ALLERGIES

	0.1.11		
	Subsidy (Manufacturer's		Fully Brand or dised Generic
	\$	Per	 Manufacturer
Mast Cell Stabilisers			
NEDOCROMIL			
Aerosol inhaler, 2 mg per dose CFC-free		112 dose OP	 Tilade
SODIUM CROMOGLICATE			
Aerosol inhaler, 5 mg per dose CFC-free		112 dose OP	 Intal Forte CFC Free
Methylxanthines			
AMINOPHYLLINE			
 Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available PSO 		5	✓ DBL Aminophylline
THEOPHYLLINE			
* Tab long-acting 250 mg	21.51	100	✓ Nuelin-SR
* Oral liq 80 mg per 15 ml		500 ml	Nuelin
Mucolytics			
DORNASE ALFA - Special Authority see SA0611 below - I			
Nebuliser soln, 2.5 mg per 2.5 ml ampoule	250.00	6	Pulmozyme
▶ SA0611 Special Authority for Subsidy Special Authority approved by the Cystic Fibrosis Advisory F Notes: Application details may be obtained from PHARMAC		w.pharmac.govt.	nz or:
	e: (04) 460 4990		
	imile: (04) 916 757	1	
	I: CFPanel@pharm		
Prescriptions for patients approved for treatment must be wr and expertise in treating cystic fibrosis.	itten by respiratory p	physicians or pae	diatricians who have experience
SODIUM CHLORIDE			
Not funded for use as a nasal drop.			_
Soln 7%	23.50	90 ml OP	 Biomed
Nasal Preparations			
Allergy Prophylactics			
BECLOMETHASONE DIPROPIONATE			
Metered aqueous nasal spray, 50 mcg per dose		200 dose OP	
Metered equeeue peoplement 100 mes per door	(5.26)	000 daga OP	Alanase
Metered aqueous nasal spray, 100 mcg per dose	2.46 (6.00)	200 dose OP	Alanase
(Alanase Metered aqueous nasal spray, 50 mcg per dose to (Alanase Metered aqueous nasal spray, 100 mcg per dose t	be delisted 1 Janua		
BUDESONIDE			
Metered aqueous nasal spray, 50 mcg per dose		200 dose OP	✓ <u>SteroClear</u>
Metered aqueous nasal spray, 100 mcg per dose	2.8/	200 dose OP	✓ SteroClear
FLUTICASONE PROPIONATE Metered aqueous nasal spray, 50 mcg per dose	1.98	120 dose OP	 Flixonase Hayfever <u>& Allergy</u>

	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully Brand or sidised Generic ✔ Manufacturer
PRATROPIUM BROMIDE Aqueous nasal spray, 0.03%	4.61	15 ml OP	✓ <u>Univent</u>
Respiratory Devices			
MASK FOR SPACER DEVICE			
a) Up to 50 dev available on a PSO			
b) Only on a PSO			
c) Only for children aged six years and under Small.	2 20	1	✓ e-chamber Mask
PEAK FLOW METER			
a) Up to 25 dev available on a PSO			
b) Only on a PSO			
Low range	9.54	1	 Mini-Wright AFS Low Range
Normal range	9.54	1	 Mini-Wright Standard
SPACER DEVICE			• • • • • • • • • • • • • • • • • • • •
a) Up to 50 dev available on a PSO			
b) Only on a PSO			
220 ml (single patient)		1	 e-chamber Turbo
510 ml (single patient)	5.12	1	 e-chamber La Grande
800 ml	6.50	1	✓ Volumatic
Respiratory Stimulants			
CAFFEINE CITRATE			
Oral liq 20 mg per ml (10 mg base per ml)	14.85	25 ml OP	 Biomed

	Subsidy		Fully Brand or
	(Manufacturer's Pi \$	rice) Subs Per	sidised Generic Manufacturer
Ear Preparations			
ACETIC ACID WITH 1, 2- PROPANEDIOL DIACETATE AND E	BENZETHONIUM		
For Vosol ear drops with hydrocortisone powder refer Stan		ge 227	
Ear drops 2% with 1, 2-Propanediol diacetate 3% and	C 07		. Nacal
benzethonium chloride 0.02%	6.97	35 ml OP	✓ Vosol
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	 Locacorten-Viaform ED's
			 Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMY		IN	
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g		7.5 ml OP	✓ Kenacomb
2.5 mg and gramicidin 250 mcg per g		7.5 111 01	• Reliacomb
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and			
gramicidin 50 mcg per ml		8 ml OP	Cafradau
	(9.27)		Sofradex
FRAMYCETIN SULPHATE Ear/Eye drops 0.5%	4.13	8 ml OP	
	(8.65)	0.111 0.	Soframycin
Eye Preparations			
Eye preparations are only funded for use in the eye, unless exp	licitly stated otherw	viso	
Anti-Infective Preparations	nonly olated entern	100.	
•			
ACICLOVIR * Eye oint 3%	14 92	4.5 g OP	✓ ViruPOS
CHLORAMPHENICOL	14.02	4.0 g OI	
Eye oint 1%	2.48	4 g OP	✓ Chlorsig
Eye drops 0.5%		10 ml OP	 Chlorafast
Funded for use in the ear*. Indications marked with * a	are unapproved ind	lications.	
CIPROFLOXACIN Eye drops 0.3% – Subsidy by endorsement	9 99	5 ml OP	 Ciprofloxacin Teva
When prescribed for the treatment of bacterial keratitis			
for the second line treatment of chronic suppurative oti Note: Indication marked with a * is an unapproved ind	, ,	; and the pres	cription is endorsed accordingly.
GENTAMICIN SULPHATE		5	
Eye drops 0.3%	11.40	5 ml OP	 Genoptic
PROPAMIDINE ISETHIONATE * Eye drops 0.1%	2 07	10 ml OP	
	(14.55)		Brolene
	·/		
SODIUM FUSIDATE [FUSIDIC ACID]			

()	Subsidy Vanufacturer's F	Price) Subs	Fully sidised	Brand or Generic
·	\$	Per	1	Manufacturer
TOBRAMYCIN				
Eye oint 0.3%	10.45	3.5 g OP	🗸 I	obrex
Eye drops 0.3%	11.48	5 ml OP	√ T	obrex
Corticosteroids and Other Anti-Inflammatory Pre	parations			
DEXAMETHASONE				
* Eye oint 0.1%	5.86	3.5 g OP	✓ N	laxidex
* Eye drops 0.1%	4.50	5 ml OP	✓ N	laxidex
Ocular implant 700 mcg - Special Authority see SA1680 below	v			
- Retail pharmacy		1	√ (zurdex

SENSORY ORGANS

■ SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Either:
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

*	Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b			· · · · ·
	sulphate 6,000 u per g5.3	39 3	8.5 g OP	 Maxitrol
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin			
	b sulphate 6,000 u per ml4.5	50 5	5 ml OP	 Maxitrol
DIC	CLOFENAC SODIUM			
	Eye drops 0.1%	30 5	5 ml OP	 Voltaren Ophtha
				-

*Three months or six months, as applicable, dispensed all-at-once

	Subaide		Fully Prond or
	Subsidy (Manufacturer's Pr	ice) Subs	Fully Brand or idised Generic
	\$	Per	 Manufacturer
FLUOROMETHOLONE			
* Eye drops 0.1%	3.09	5 ml OP	✓ FML
	5.20		 Flucon
LEVOCABASTINE			
Eye drops 0.5 mg per ml		4 ml OP	Lineatie
	(10.34)		Livostin
LODOXAMIDE Eye drops 0.1%	0.71	10 ml OP	✓ Lomide
	0./1	10 IIII OF	✓ Lonnue
PREDNISOLONE ACETATE Eye drops 1%	2.02	10 ml OP	Prednisolone-AFT
	7.00	5 ml OP	 ✓ Pred lisoione-APT ✓ Pred Forte
PREDNISOLONE SODIUM PHOSPHATE – Special Authority se			
Eye drops 0.5%, single dose (preservative free)		20 dose	✓ Minims
_) = = = = = = = = = = = = = = = = = = =			Prednisolone
► SA1715 Special Authority for Subsidy			
Initial application only from an ophthalmologist or optometrist.	Approvals valid fo	r 6 months for	applications meeting the
following criteria:			
Both:			
1 Patient has severe inflammation; and			
2 Patient has a confirmed allergic reaction to preservative in	• •		
Renewal from any relevant practitioner. Approvals valid for 6 mo	onths where the tr	eatment rema	ins appropriate and the patient is
benefiting from treatment.			
SODIUM CROMOGLICATE Eye drops 2%	0.85	5 ml OP	✓ Rexacrom
	0.05	51111 01	
Glaucoma Preparations - Beta Blockers			
BETAXOLOL			
Eve drops 0.25%	11.80	5 ml OP	 Betoptic S
* Eye drops 0.5%		5 ml OP	 ✓ Betoptic ✓ Betoptic
TIMOLOL			
* Eve drops 0.25%	1.43	5 ml OP	 Arrow-Timolol
* Eye drops 0.25%, gel forming	3.30	2.5 ml OP	✓ Timoptol XE
* Eye drops 0.5%	1.43	5 ml OP	 Arrow-Timolol
* Eye drops 0.5%, gel forming	3.78	2.5 ml OP	 <u>Timoptol XE</u>
Glaucoma Preparations - Carbonic Anhydrase I	nhihitoro		
Giaucoma Freparations - Carbonic Annyarase I			
ACETAZOLAMIDE			
* Tab 250 mg	17.03	100	 <u>Diamox</u>
BRINZOLAMIDE			_
* Eye drops 1%	9.77	5 ml OP	 Azopt
DORZOLAMIDE HYDROCHLORIDE			
* Eye drops 2%		5 ml OP	
	(17.44)		Trusopt
DORZOLAMIDE WITH TIMOLOL			
* Eye drops 2% with timolol 0.5%	2.87	5 ml OP	 Dortimopt

	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully Brand or idised Generic ✓ Manufacturer
Glaucoma Preparations - Prostaglandin Analog	ues		
BIMATOPROST * Eye drops 0.03%	3.30	3 ml OP	 Bimatoprost Multichem
LATANOPROST * Eye drops 0.005% Teva to be Sole Supply on 1 July 2019 (Hysite Eye drops 0.005% to be delisted 1 July 2019)	1.50 1.57	2.5 ml OP	✓ Hysite✓ Teva
TRAVOPROST * Eye drops 0.004%	7.30 19.50	5 ml OP 2.5 ml OP	✓ Travopt✓ Travatan
Glaucoma Preparations - Other			
BRIMONIDINE TARTRATE * Eye drops 0.2% BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE * Eye drops 0.2% with timolol maleate 0.5%		5 ml OP 5 ml OP	 Arrow-Brimonidine Combigan
PILOCARPINE HYDROCHLORIDE	4.26 5.35 7.99	15 ml OP 15 ml OP 15 ml OP	 ✓ Isopto Carpine ✓ Isopto Carpine ✓ Isopto Carpine
 Eye drops 2% single dose – Special Authority see SA0895 below – Retail pharmacy 		20 dose	✓ Minims Pilocarpine

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

1 Patient has to use an unpreserved solution due to an allergy to the preservative; or

2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics

ATROPINE SULPHATE * Eye drops 1%	15 ml OP	✓ <u>Atropt</u>
CYCLOPENTOLATE HYDROCHLORIDE * Eye drops 1%8.76	15 ml OP	 Cyclogyl
TROPICAMIDE * Eye drops 0.5%	15 ml OP 15 ml OP	✓ Mydriacyl ✓ Mydriacyl

SENSORY ORGANS

	Subsidy (Manufacturer's Pric \$	e) Subsi Per	Fully dised	Brand or Generic Manufacturer
Preparations for Tear Deficiency				
For acetylcysteine eye drops refer Standard Formulae, page 227				
HYPROMELLOSE				
* Eye drops 0.5%	2.00 (3.92)	15 ml OP	Μ	lethopt
HYPROMELLOSE WITH DEXTRAN				
* Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	🗸 P	oly-Tears
POLYVINYL ALCOHOL				
* Eye drops 1.4%		15 ml OP	✓ V	istil
* Eye drops 3%	3.68	15 ml OP	✓ <u>v</u>	istil Forte
Preservative Free Ocular Lubricants				

► SA1388 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER - Special Authority see SA1388 above - Retail ph	armacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	 Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL - Special Autho	rity see SA1388 at	ove – Retai	l pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml		24	 Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special Aut	hority see SA1388	above - Ret	tail pharmacy
Eye drops 1 mg per ml		10 ml OP	 Hylo-Fresh
Hylo-Fresh has a 6 month expiry after opening. The Ph			
month is not relevant and therefore only the prescribed	dosage to the near	est OP may	be claimed.

Other Eye Preparations

	15 ml OP	 Naphcon Forte
OLOPATADINE Eye drops 0.1%	5 ml OP	 Patanol
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE Eye oint 138 mcg per g	5 g OP	✓ VitA-POS

VARIOUS

	Subsidy (Manufacturer's Pric \$	e) Subs Per	Fully idised	Brand or Generic Manufacturer
Agents Used in the Treatment of Poisonings				
Antidotes				
ACETYLCYSTEINE – Retail pharmacy-Specialist Inj 200 mg per ml, 10 ml ampoule NALOXONE HYDROCHLORIDE a) Up to 5 inj available on a PSO b) Only on a PSO	58.76	10	✓ <u>D</u>	BL Acetylcysteine
 * Inj 400 mcg per ml, 1 ml ampoule 	22.60	5	✓ <u>D</u>	BL Naloxone Hydrochloride
Removal and Elimination				
CHARCOAL * Oral liq 50 g per 250 ml a) Up to 250 ml available on a PSO b) Only on a PSO	43.50	250 ml OP	✔ C	arbosorb-X
DEFERASIROX – Special Authority see SA1492 below – Retail Wastage claimable	l pharmacy			
Tab 125 mg dispersible		28	✓ E	xjade
Tab 250 mg dispersible		28		xjade
Tab 500 mg dispersible	1,105.00	28	✓ E	xjade
SA1492 Special Authority for Subsidy Initial application only from a haematologist. Approvals valid for All of the following: 1 The natient has been diagnosed with chronic iron overlage			-	-

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE - Special Authority see SA1480 on the next page - Retail pharmacy

Tab 500 mg	·	 533.17	100	 Ferriprox
Oral liq 100 mg p	oer 1 ml	 266.59	250 ml OP	 Ferriprox

\$ Per 🗸 Manufacturer	 Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic
	 \$	Per 🗸	Manufacturer

► SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE

* Inj 500 mg vial	84.53	10	✓ <u>DBL</u> Desferrioxamine Mesylate for Inj <u>BP</u>
SODIUM CALCIUM EDETATE			
* Inj 200 mg per ml, 5 ml		6	
	(156.71)		Calcium Disodium Versenate

Standard Formulae

ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base	qs qs	PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water	1 g 70 ml to 100 ml
ASPIRIN AND CHLOROFORM APPLICATION Aspirin Soluble tabs 300 mg Chloroform	12 tabs to 100 ml	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml) Phenobarbitone Sodium	LIQUID (10
CODEINE LINCTUS PAEDIATRIC (3 mg per 5 ml) Codeine phosphate Glycerol Preservative	60 mg 40 ml qs	Glycerol BP Water PILOCARPINE ORAL LIQUID	400 mg 4 ml to 40 ml
Water CODEINE LINCTUS DIABETIC (15 mg per 5 ml) Codeine phosphate	to 100 ml	Pilocarpine 4% eye drops Preservative Water (Preservative should be used if quantity supplied is	qs qs to 500 ml for more
Glycerol Preservative Water	40 ml qs to 100 ml	than 5 days.) SALIVA SUBSTITUTE FORMULA	
FOLINIC MOUTHWASH Calcium folinate 15 mg tab Preservative Water	1 tab qs to 500 ml	Methylcellulose Preservative Water (Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	5 g qs to 500 ml for more
(Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	for more	SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml	qs
MAGNESIUM HYDROXIDE 8% MIXTURE Magnesium hydroxide paste 29% Methyl hydroxybenzoate Water	275 g 1.5 g to 1 000 m	Water (Only funded if prescribed for treatment of hyponatr _{II} VANCOMYCIN ORAL SOLUTION (50 mg per ml)	qs aemia)
METHADONE MIXTURE Methadone powder Glycerol	qs qs	Vancomycin 500 mg injection Glycerol BP Water (Only funded if prescribed for treatment of Clostridiu	10 vials 40 ml to 100 ml ım difficile
Water METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol	to 100 ml 10 g to 100 ml	following metronidazole failure) VOSOL EAR DROPS WITH HYDROCORTISONE POWDER 1% Hydrocortisone powder	1%
(Use 1 ml of the 10% solution per 100 ml of oral liqu OMEPRAZOLE SUSPENSION Omeprazole capules or powder	,	Vosol Ear Drops	to 35 ml
Sodium bicarbonate powder BP Water	qs 8.4 g to 100 ml		

	Outside		E. III	Durand au
	Subsidy (Manufacturer's Pric	0) C.L	Fully sidised	Brand or Generic
	(Manulacturer's Pric	Per Sub	siaisea ✓	Manufacturer
	Ŧ			
Extemporaneously Compounded Preparations	and Galenicals	S		
BENZOIN				
Tincture compound BP	24.42	500 ml		
	(39.90)		F	harmacy Health
	2.44	50 ml		
	(5.10)		F	harmacy Health
CHLOROFORM				
a) Only in combination				
b) Maximum of 100 ml per prescription				
c) Only in aspirin and chloroform application.				
d) Note: This product is no longer being manufactured by the	he supplier and will	be delisted	from the	e Schedule at a date to be
determined.				
Chloroform BP		500 ml	✓ F	PSM
CODEINE PHOSPHATE - Safety medicine; prescriber may dete				
Powder – Only in combination		25 g		
	(90.09)			Douglas
Only in extemporaneously compounded codeine linctus	diabetic or codeine	e linctus pae	diatric.	
COLLODION FLEXIBLE				
Note: This product is no longer being manufactured by the s	supplier and will be	delisted from	n the S	chedule at a date to be
determined.				
Collodion flexible	19.30	100 ml	✓ F	PSM
COMPOUND HYDROXYBENZOATE - Only in combination				
Only in extemporaneously compounded oral mixtures.				
Soln		100 ml		lidwest
Midure the her Oals Overslever 4 Average 0040	34.18		✓ [David Craig
Midwest to be Sole Supply on 1 August 2019				
(David Craig Soln to be delisted 1 August 2019)				
GLYCERIN WITH SODIUM SACCHARIN – Only in combination				
Only in combination with Ora-Plus. Suspension	20.05	473 ml		Dra-Sweet SF
		473 m	• (Jra-Sweel Sr
Ora-Sweet SF to be Sole Supply on 1 July 2019				
GLYCERIN WITH SUCROSE – Only in combination				
Only in combination with Ora-Plus. Suspension	20.05	473 ml		Dra-Sweet
Ora-Sweet to be Sole Supply on 1 July 2019		4/3 111	• (Jia-Sweet
GLYCEROL * Liquid – Only in combination	2.00	500 ml		anithE Changeral BD
Only in extemporaneously compounded oral liquid prepa		500 ml	• <u>n</u>	ealthE Glycerol BP
	aralions.			
MAGNESIUM HYDROXIDE	00.01	F00 m	✓ F	2014
Paste 29%	22.01	500 g	• •	· 51M
METHADONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
 c) Safety medicine; prescriber may determine dispensing from d) System are accurately compounded methodene will ach be 		oto of the - I		form available
d) Extemporaneously compounded methadone will only be (methadone powder, not methadone tablete)	reimpursed at the r	ale of the Ch	ieapest	IOTTI available
(methadone powder, not methadone tablets). Powder	7 8/	1 g	✓ A) FT
		' y	• •	

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy			and or
	(Manufacturer's F \$	Price) Subs Per		eneric anufacturer
METHYL HYDROXYBENZOATE Powder Midwest to be Sole Supply on 1 July 2019		25 g	✓ Midw	
METHYLCELLULOSE Powder		100 g	🖌 MidW	/est
MidWest to be Sole Supply on 1 July 2019 Suspension – Only in combination Ora-Plus to be Sole Supply on 1 July 2019		473 ml	✔ Ora-F	Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCH Suspension Ora-Blend SF to be Sole Supply on 1 July 2019	,	combination 473 ml	🗸 Ora-I	Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE – Or Suspension Ora-Blend to be Sole Supply on 1 July 2019		473 ml	🗸 Ora-I	Blend
PHENOBARBITONE SODIUM Powder – Only in combination	52.50 325.00	10 g 100 g	✓ MidW ✓ MidW	
Only in children up to 12 years PROPYLENE GLYCOL Only in extemporaneously compounded methyl hydroxyben				
	11.25	500 ml	 Midw 	est
SODIUM BICARBONATE Powder BP – Only in combination	8.95 9.80	500 g	🗸 Midw	est
Only in extemporaneously compounded omeprazole an	(29.50) d lansoprazole si	ispension	David	l Craig
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparati				
Liq		2,000 ml	🖌 Midw	est
WATER Tap – Only in combination	0.00	1 ml	🖌 Тар у	vater

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1522 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children; or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 inborn errors of metabolism; or
- 7 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Cystic fibrosis or renal failure)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE AND FAT SU	JPPLEMENT – Special Author	ity see SA1376 on t	he previous pag	e – Hosp	oital pharmacy [HP3]
Powder (neutral)	-		400 g OP	Duod	al Super
			-	So	uble Powder

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome; or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic
 \$	Per	1	Manufacturer

- 10 ascites; or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Patho

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Autho	ity see SA1523 on the previous	page – Hospital pharmacy [HP3]
--------------------------------	--------------------------------	--------------------------------

Emulsion (neutral)		200 ml OP	✓ Calogen
	30.75	500 ml OP	 Calogen
Emulsion (strawberry)		200 ml OP	 Calogen
Oil		500 ml OP	 MCT oil (Nutricia)
Oil, 250 ml		4 OP	 Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Resource Beneprotein

	armacy [HP3]	PROTEIN SUPPLEMENT – Special Authority see SA1524 above – Hospital ph	PR
 Protifar 	225 g OP	Powder	
 Resource 	227 g OP	8.95	
Donon	•		

Subsidy (Manufacturer's Price)

¢

Fully Subsidised

Per

Generic Manufacturer

Brand or

Oral Supplements/Complete Diet (Nasogastric/Gastrostomy Tube Feed)

Respiratory Products

⇒SA1094 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has CORD and hypercapnia, defined as a CO2 value exceeding 55 mmHg.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CORD ORAL FEED 1.5KCAL/ML - Special Author	rity see SA1094 above – Hosp	oital pharmacy [I	HP3]
Liquid	1.66	237 ml OP	 Pulmocare

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML – Special Authority see SA1095 above – Liquid	- Hospital pharn 1,000 ml OP	acy [HP3] ✓ Diason RTH ✓ Glucerna Select RTH
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hot	spital pharmacv	[HP3]
Liquid (strawberry)1.50	200 ml OP	✓ Diasip
Liquid (vanilla)	200 ml OP	✓ Diasip
1.88	250 ml OP	 Glucerna Select
1.78	237 ml OP	
(2.10)		Resource Diabetic
(2.10)		Sustagen Diabetic

Subsid	y Full	Brand or
(Manufacturer	s Price) Subsidise	I Generic
\$	Per 🗸	Manufacturer

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED	- Special Authority see SA152	5 above – Hospital phar	macy [HP3]		
Powder		60.48	400 g OP	~	Monogen

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
·	\$	Per 🗸	Manufacturer
ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA10		•	
Liquid		00 g OP 🛛 🗸	Kindergen

SPECIAL FOODS

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML – Special Authority see SA1379 Liquid	
PAEDIATRIC ENTERAL FEED 1KCAL/ML – Special Authority see SA1379 al Liquid2.68	bove – Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini RTH ✓ Pediasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority Liquid	y see SA1379 above – Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML – Special Authority see SA1379 abo Liquid (strawberry)1.60 Liquid (vanilla)1.60	ve – Hospital pharmacy [HP3] 200 ml OP ✓ Fortini 200 ml OP ✓ Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML – Special Authority see SA1379 above Liquid (chocolate)	e – Hospital pharmacy [HP3] 200 ml OP ✓ Pediasure 200 ml OP ✓ Pediasure 200 ml OP ✓ Pediasure 250 ml OP ✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority see Liquid (unflavoured) 1.60 Liquid (chocolate) 1.60 Liquid (strawberry) 1.60 Liquid (vanilla) 1.60	SA1379 above – Hospital pharmacy [HP3] 200 ml OP ✓ Fortini Multi Fibre 200 ml OP ✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 above – Hos Powder	pital pharmacy [HP3] 400 g OP ✓ Peptamen Junior

	Subsidy (Manufacturer's Pric \$	Fully e) Subsidised Per ✓	Brand or Generic Manufacturer
Renal Products			
 SA1101 Special Authority for Subsidy itial application only from a dietitian, relevant specialist or ears where the patient has acute or chronic kidney disease Renewal only from a dietitian, relevant specialist, vocational ecommendation of a dietitian, relevant specialist or vocatior pplications meeting the following criteria: The treatment remains appropriate and the patient is General Practitioners must include the name of the di practitioner and date contacted. 	y registered general pr ally registered general benefiting from treatme	actitioner or general practitioner. Approv	practitioner on the vals valid for 3 years for
RENAL ENTERAL FEED 1.8 KCAL/ML – Special Authority			IP31
Liquid	6.08	500 ml OP 🛛 🖌 🖊	lepro HP RTH
· · · · · ·	SA1101 above – Hospi	al pharmacy [HP3] 220 ml OP 🖌 N	•

Specialised And Elemental Products

SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

SPECIAL FOODS

	Subsidy (Manufacturer's I \$	Price) Subsi Per	Fully idised	Brand or Generic Manufacturer
ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML – Spi pharmacy [HP3] Liquid	,	e SA1377 on th 1,000 ml OP	e previ	
ORAL ELEMENTAL FEED 0.8KCAL/ML – Special Authority see Liquid (grapefruit), 250 ml carton Liquid (pineapple & orange), 250 ml carton Liquid (summer fruits), 250 ml carton	171.00 171.00	previous page - 18 OP 18 OP 18 OP 18 OP	✓ E	ital pharmacy [HP3] Elemental 028 Extra Elemental 028 Extra Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML – Special Authority see S Powder (unflavoured)		r <mark>evious page</mark> – H 80 g OP		al pharmacy [HP3] /ivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML – Special Aut [HP3] Liquid	,	7 on the previou 1,000 ml OP		e – Hospital pharmacy Peptisorb

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
 - 1 Child aged one to eight years; and
 - 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML	- Special Authority	see SA1196	above -	- Hospital pharmacy [HP3]
Liquid	4.00	500 ml OP	-	Nutrini Low Energy
				Multi Fibre

Standard Supplements

⇒SA1554 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and

3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) only from a dietitian, relevant

continued...

Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic	
\$	Per 🗸	Manufacturer	

specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist, dietitian on the recommendation of a gastroenterologist or vocationally registered general practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:
 - Patient has not responded to first-line dietary measures over a 4 week period by:
 - 2.1 Increasing their food intake frequency (eg snacks between meals); or
 - 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
 - 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:
 - Patient is Malnourished
 - 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
 - 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
 - 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	✓	Manufacturer	

Initial application — (Short-term medical condition) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding: or

- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or</p>
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or</p>
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or

continued...

Subsidy (Manufacturer's Price)	5	Fully Subsidised	Brand or Generic
\$	Per	1	Manufacturer

- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML – Special Authority see SA1554 on page 237 – I Liquid	Hospital pharmacy [HP3] 1,000 ml OP ✓ Nutrison Energy
ENTERAL FEED 1KCAL/ML – Special Authority see SA1554 on page 237 – Ho Liquid	spital pharmacy [HP3] 250 ml OP 1,000 ml OP Vutrison Standard RTH Standard RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML – Special Authority see SA1554 Liquid	on page 237 – Hospital pharmacy [HP3] 1,000 ml OP ✓ Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML – Special Authority see SA1554 on Liquid	page 237 – Hospital pharmacy [HP3] 1,000 ml OP ✓ Jevity RTH ✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority see SA1554 or Liquid	a page 237 – Hospital pharmacy [HP3] 250 ml OP ✓ Ensure Plus HN 1,000 ml OP ✓ Ensure Plus RTH ✓ Jevity HiCal RTH ✓ Nutrison Energy Multi Fibre

	Subsidy			nd or
	(Manufacturer's P \$	Price) Subs Per		ieric nufacturer
ORAL FEED (POWDER) – Special Authority see SA1554 on page Note: Higher subsidy for Sustagen Hospital Formula will only number and an appropriately endorsed prescription.	be reimbursed			d Special Authority
Powder (chocolate) – Higher subsidy of up to \$26.00 per 850		050 - 00		-
with Endorsement		850 g OP 840 g OP	 Ensure 	e
	(26.00)	040 g 01		en Hospital Jula Active
Additional subsidy by endorsement is available for patien prescription must be endorsed accordingly.	ts with fat malak	osorption, fat in		
Powder (vanilla) – Higher subsidy of up to \$26.00 per 850 g	0.54	057 - 00	/ Faultai	-
with Endorsement	8.54 26.00	857 g OP 850 g OP	 Fortisi Ensure 	
	20.00 9.54	840 g OP		5
	(26.00)	040 g 01		jen Hospital nula Active
Additional subsidy by endorsement is available for patien prescription must be endorsed accordingly.	ts with fat malak	osorption, fat in		
epidermolysis bullosa, or as exclusive enteral nutrition in child disease. The prescription must be endorsed accordingly. Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	ige of 18 years 200 ml OP		
	(1.26) (1.26)		Ensure Fortisip	
Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with Endorsement		200 ml OP		
Endorsement	0.72 (1.26)	200 MI OP	Ensure	Plue
	(1.26)		Fortisip	
Liquid (fruit of the forest) – Higher subsidy of \$1.26 per 200 r				
with Endorsement		200 ml OP	_	
	(1.26)		Ensure	e Plus
Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with		000 ml OD		
Endorsement	0.72 (1.26) (1.26)	200 ml OP	Ensure Fortisig	
Liquid (vanilla) – Higher subsidy of up to \$1.33 per 237 ml wi			ronisi)
Endorsement		237 ml OP		
	(1.33)		Ensure	e Plus
	0.72	200 ml OP		
	(1.26) (1.26)		Ensure Fortisip	

SPECIAL FOODS

	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully idised	Brand or Generic Manufacturer
ORAL FEED WITH FIBRE 1.5 KCAL/ML – Special Authority see Additional subsidy by endorsement is available for patients b epidermolysis bullosa. The prescription must be endorsed a Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with	eing bolus fed th ccordingly.			
Endorsement		200 ml OP	F	ortisip Multi Fibre
Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with Endorsement		200 ml OP	F	ortisip Multi Fibre
Liquid (vanilla) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72 (1.26)	200 ml OP	F	ortisip Multi Fibre

High Calorie Products

➡SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML – Special Authority see SA1195 abo	ve – Hospital	pharmacy [HP3]	
Liquid	5.50	500 ml OP	 Nutrison
			Concentrated
	11.00	1,000 ml OP	🗸 Two Cal HN RTH
		,	

	Subsidy (Manufacturer's Pri \$	ce) Subs Per	Fully idised	Brand or Generic Manufacturer
ORAL FEED 2 KCAL/ML – Special Authority see SA1195 on the Additional subsidy by endorsement is available for patients b epidermolysis bullosa. The prescription must be endorsed a	eing bolus fed three			
Liquid (vanilla) – Higher subsidy of \$1.90 per 200 ml with Endorsement	0.96 (1.90)	200 ml OP	T۱	wo Cal HN
Food Thickeners				
 SA1106 Special Authority for Subsidy Initial application only from a dietitian, relevant specialist or voc year where the patient has motor neurone disease with swallowin Renewal only from a dietitian, relevant specialist, vocationally represented in the second state of the following criteria: Both: The treatment remains appropriate and the patient is benue General Practitioners must include the name of the dietitian practitioner and date contacted. 	ng disorder. gistered general p registered general efiting from treatm an, relevant specia	ractitioner or o practitioner. ent; and list or vocatio	general Approv	practitioner on the als valid for 1 year for
FOOD THICKENER – Special Authority see SA1106 above – He Powder		HP3] 300 g OP 380 g OP		utilis eed Thickener Karicare Aptamil
Gluten Free Foods				
The funding of gluten free foods is no longer being actively mana no longer considering the listing of new products, or making subs anticipate that the range of funded items will reduce over time.	idy, or other chan	ges to the exis	sting list	tings. As a result we

► SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

necessary for good outcomes. A range of gluten free options are available through retail outlets.

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX – Special Authority see SA17	729 above – Hospital pharmacy [HP3]	
Powder	2.81 1,000 g OP	
	(5.15)	Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA17	29 above – Hospital pharmacy [HP3]	
Powder		
	(7.32)	NZB Low Gluten Bread Mix
	3.51	
	(10.87)	Horleys Bread Mix

SPECIAL FOODS

	Subsidy (Manufacturer's Pr \$	ice) S Per	Fully ubsidised	Brand or Generic Manufacturer
GLUTEN FREE FLOUR - Special Authority see SA1729 on the				IP3]
Powder		2,000 g O		
	(18.10)		ł	Horleys Flour
GLUTEN FREE PASTA - Special Authority see SA1729 on the	previous page - H	lospital pha	armacy [H	IP3]
Buckwheat Spirals	2.00	250 g OP)	
	(3.11)		(Orgran
Corn and Vegetable Shells	2.00	250 g OP)	
	(2.92)			Orgran
Corn and Vegetable Spirals		250 g OP		
	(2.92)			Orgran
Rice and Corn Lasagne Sheets		200 g OP		_
	(3.82)			Orgran
Rice and Corn Macaroni		250 g OP		~
	(2.92)			Orgran
Rice and Corn Penne		250 g OP		^
Disc and Maine Deate Opinale	(2.92)	050		Orgran
Rice and Maize Pasta Spirals		250 g OP		O wenne w
Dice and Millet Chirole	(2.92)	050 ~ 00		Orgran
Rice and Millet Spirals		250 g OP)
Pice and corp apaghetti peedlee	(3.11)	275 a OB		Orgran
Rice and corn spaghetti noodles		375 g OP		Orgran
Vegetable and Rice Spirals	(2.92)	250 g OP		Jigian
vegetable and title opliato	(2.92)	200 Y OF		Orgran
Italian long style spaghetti		220 g OP		orgran
	(3.11)	220 y OI		Orgran

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

AMINOACID FORMULA WITHOUT METHIONINE - Special Autho	rity see SA110	8 above – Hos	pital pharmacy [HP3]
Powder	461.94	500 g OP	 XMET Maxamum

Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Sp	pecial Authority s	see SA1108 above – Hospital
pharmacy [HP3]		
Powder	500 g OP	MSUD Maxamum

	Subsidy (Manufacturer's Price) \$		Fully Brand or dised Generic Manufacturer
Supplements For PKU			
AMINOACID FORMULA WITHOUT PHENYLALANINE – Special pharmacy [HP3]	Authority see SA11	108 on the p	revious page – Hospital
Tabs		75 OP	 Phlexy 10
Powder (chocolate) 36 g sachet	393.00	30	 PKU Anamix Junior Chocolate
Powder (unflavoured) 27.8 g sachets	936.00	30	 PKU Lophlex Powder
Powder (unflavoured) 36 g sachets		30	PKU Anamix Junior
Powder (vanilla) 36 g sachet		30	 PKU Anamix Junior Vanilla
Infant formula	174.72 4	00 g OP	 PKU Anamix Infant
Powder (orange)		00 g OP	✓ XP Maxamum
Powder (unflavoured)		00 g OP	✓ XP Maxamum
Liquid (berry)		25 ml OP	 PKU Anamix Junior LQ
Liquid (orange)		25 ml OP	 PKU Anamix Junior LQ
Liquid (unflavoured)		25 ml OP	 PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	 Easiphen Liquid
Liquid (juicy tropical) 125 ml		30 OP	PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g		36 OP	 PKU Lophlex Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml		60 OP	PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml		60 OP	PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml.		30 OP	PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml		30 OP	 PKU Lophlex LQ 20

Foods

LOW PROTEIN BAKING MIX - Special Authority see SA1108 on	the previous pa	<mark>age</mark> – Hospital p	oharmacy [HP3]
Powder	8.22	500 g OP	 Loprofin Mix
LOW PROTEIN PASTA - Special Authority see SA1108 on the pr	evious page –	Hospital pharm	acy [HP3]
Animal shapes	11.91	500 g OP	 Loprofin
Lasagne	5.95	250 g OP	 Loprofin
Low protein rice pasta		500 g OP	 Loprofin
Macaroni	5.95	250 g OP	 Loprofin
Penne		500 g OP	 Loprofin
Spaghetti		500 g OP	 Loprofin
Spirals	11.91	500 g OP	 Loprofin

SPECIAL FOODS

	Subsidy (Manufacturer's Pric \$	e) Sub Per	Fully sidised	Brand or Generic Manufacturer
Infant Formulae				
For Williams Syndrome				
 SA1110 Special Authority for Subsidy nitial application only from a dietitian, relevant specialist or v vear where the patient is an infant suffering from Williams Syn Renewal only from a dietitian, relevant specialist, vocationally vecarmendation of a dietitian, relevant specialist or vocational applications meeting the following criteria: 3oth: 1 The treatment remains appropriate and the patient is bi 2 General Practitioners must include the name of the diel practitioner and date contacted. WO CALCIUM INFANT FORMULA – Special Authority see S Powder	drome and associated registered general pra lly registered general pra enefiting from treatme litian, relevant special SA1110 above – Hosp	I hypercalca actitioner or oractitioner. nt; and st or vocati	aemia. general Approv onally re cy [HP3	practitioner on the vals valid for 1 year for
Gastrointestinal and Other Malabsorptive Pro	blems			
AMINO ACID FORMULA – Special Authority see SA1219 bel Powder		cy [HP3] 400 g OP	-	Ifamino Junior leocate LCP
Powder (unflavoured)		400 g OP	✓ E ✓ E ✓ N	lecare lecare LCP leocate Gold leocate Junior Unflavoured leocate SYNEO
Powder (vanilla)	53.00	400 g OP		leocate Junior

(Neocate LCP Powder to be delisted 1 August 2019)

⇒SA1219 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

1 Extensively hydrolysed formula has been reasonably trialled and is inappropriate due to documented severe intolerance or allergy or malabsorption; or

Vanilla

- 2 History of anaphylaxis to cows milk protein formula or dairy products; or
- 3 Eosinophilic oesophagitis.
- Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's Price) \$Pe	Fully Subsidised er	Brand or Generic Manufacturer
EXTENSIVELY HYDROLYSED FORMULA – Special Authority Powder			y [HP3] µptamil Gold+ Pepti Junior
■ SA1557 Special Authority for Subsidy Initial application only from a dietitian, relevant specialist or very months for applications meeting the following criteria: Any of the following: 1 Both:	ocationally registered gene	ral practitione	r. Approvals valid for 6
1.1 Cows milk formula is inappropriate due to severe1.2 Either:			ent; and
1.2.1 Soy milk formula has been reasonably tri1.2.2 Soy milk formula is considered clinically i			
 2 Severe malabsorption; or 3 Short bowel syndrome; or 4 Intractable diarrhoea; or 5 Biliary atresia; or 6 Cholestatic liver diseases causing malsorption; or 7 Cystic fibrosis; or 8 Proven fat malabsorption; or 9 Severe intestinal motility disorders causing significant m 10 Intestinal failure; or 11 All of the following: 11.1 For step down from Amino Acid Formula; and 11.2 The infant is currently receiving funded amino acid 11.3 The infant is to be trialled on, or transitioned to, a 11.4 General Practitioners must include the name of the following in the infant is to be trialled on and the following in the infant is to be trialled on and the following in the following in the following in the infant is the provide the following in the provide the following in the infant is the provide the following in the provide the provide the provide the provide the following in the provide the pro	nalabsorption; or cid formula; and an extensively hydrolysed f	ormula; and	nally registered general
practitioner and the date contacted. Note: A reasonable trial is defined as a 2-4 week trial, or signs Renewal only from a dietitian, relevant specialist, vocationally recommendation of a dietitian, relevant specialist or vocationall applications meeting the following criteria: All of the following:	registered general practition y registered general practition	ner or general tioner. Approv	practitioner on the vals valid for 6 months for
 An assessment as to whether the infant can be transitio undertaken; and The outcome of the assessment is that the infant contin 			

3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML – Special Authority see SA1698 below – Hospital pharmacy [HP3] Liquid......2.35 125 ml OP ✓ Infatrini

➡SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula;

continued...

SPECIAL FOODS

Subsidy (Manufacturer's Price)	Su	Fully Ibsidised	Brand or Generic
 \$	Per	1	Manufacturer

and

3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special	Authority see SA1197 above	 Retail pharmacy
Powder (unflavoured)		g OP 🖌 KetoCal 4:1
		 Ketocal 3:1
Powder (vanilla)		g OP 🖌 KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

	Subsidy (Manufacturer's Price)		Fully	Brand or Generic
	\$	Per	1	Manufacturer
Vaccinations				
ADULT DIPHTHERIA AND TETANUS VACCINE – [Xpharm]				
Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid in 0.5 ml. Any of the following:	0.00	5	✓ <u>A</u>	DT Booster
1) For vaccination of patients aged 45 and 65 years	,			
 For vaccination of previously unimmunised or part For vaccination following immunosurpressions 		nts; or		
 For revaccination following immunosuppression; c For boosting of patients with tetanus-prone wound 				
5) For use in testing for primary immunodeficiency di or paediatrician.		mendatior	n of an i	nternal medicine physicia
Note: Please refer to the Immunisation Handbook for a	ppropriate schedule fo	or catch up	o progra	mmes.
BACILLUS CALMETTE-GUERIN VACCINE – [Xpharm]				
For infants at increased risk of tuberculosis. Increased risk				
 living in a house or family with a person with current or having one or more household members or carers who 			a counti	ry with a rate of TB > or
equal to 40 per 100,000 for 6 months or longer; or 3) during their first 5 years will be living 3 months or longer	er in a country with a r	rate of TR	> or ea	ual to 40 per 100 000
Note a list of countries with high rates of TB are available at	•		•	
www.bcgatlas.org/index.php.			0 (000.0	
Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),				
Danish strain 1331, live attenuated, vial with diluent	0.00	10	✓ В	CG Vaccine
DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE – [Xpha Funded for any of the following criteria:				
 A single vaccine for pregnant woman between gestation A course of up to four vaccines is funded for children for 			ears inc	clusive to complete full
primary immunisation; or3) An additional four doses (as appropriate) are funded for transplantation or chemotherapy; pre or post splenector				
severely immunosuppressive regimens.		-		-
Notes: Tdap is not registered for patients aged less than 10	years. Please refer t	to the Imm	nunisatio	on Handbook for
appropriate schedule for catch up programmes.				
Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis toxoid, 8 mcg pertussis filamentous				
haemagluttinin and 2.5 mcg pertactin in 0.5 ml syringe .	0.00	10 1	_	oostrix oostrix
DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE Funded for any of the following:	– [Xpharm]		_	
 A single dose for children up to the age of 7 who have A course of four vaccines is funded for catch up program 				ars) to complete full
primary immunisation; or 3) An additional four doses (as appropriate) are funded for	or (ro.)immunication fo	or potionto	noct Ll	CT or chamatharapy:
pre- or post splenectomy; pre- or post solid organ trans regimens; or				
4) Five doses will be funded for children requiring solid or				
Note: Please refer to the Immunisation Handbook for appro	priate schedule for ca	itch up pro	gramm	es.
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg				
pertussis toxoid, 25 mcg pertussis filamentous				
haemagluttinin, 8 mcg pertactin and 80 D-antigen units poliomyelitis virus in 0.5ml syringe	0.00	10	🖌 lı	nfanrix IPV
		10	· · II	

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully dised	Brand or Generic Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B	AND HAEMOPHILUS	INFLUENZ	AE TY	PE B VACCINE -
[Xpharm] Funded for patients meeting any of the following criteria:				
1) Up to four doses for children up to and under the age	of 10 for primary immu	nisation: o	r	
2) An additional four doses (as appropriate) are funded f				nd under the age of
10 who are patients post haematopoietic stem cell tra				
post solid organ transplant, renal dialysis and other se		1 2 / 1		1 271
3) Up to five doses for children up to and under the age	of 10 receiving solid or	gan transpl	antatio	on.
Note: A course of up-to four vaccines is funded for catch u	p programmes for child	Iren (up to	and un	ider the age of 10 years)
to complete full primary immunisation. Please refer to the I	mmunisation Handboo	k for the ap	propri	ate schedule for catch up
programmes.				
Inj 30IU diphtheriatoxoid with 40IU tetanustoxoid, 25mcg				
pertussistoxoid, 25mcg				
pertussisfilamentoushaemagluttinin, 8 mcgpertactin,				
80 D-AgUpoliovirus, 10mcghepatitisBsurfaceantigen in		40		6
0.5ml syringe	0.00	10	♥ <u>II</u>	nfanrix-hexa
HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm]				
One dose for patients meeting any of the following:				
 For primary vaccination in children; or An additional dose (as appropriate) is funded for (re-)i 	mmunication for notion	to post had	motor	voiatia atom coll
transplantation, or chemotherapy; functional asplenic;				
or post cochlear implants, renal dialysis and other sev				bild organ transplant, pre-
 For use in testing for primary immunodeficiency disea 				nal medicine physician or
paediatrician.				
Haemophilus Influenzae type B polysaccharide 10 mcg				
conjugated to tetanus toxoid as carrier protein 20-40 m				
prefilled syringe plus vial 0.5 ml	0.00	1	✓ <u>н</u>	liberix
HEPATITIS A VACCINE – [Xpharm]				
Funded for patients meeting any of the following criteria:				
1) Two vaccinations for use in transplant patients; or				
2) Two vaccinations for use in children with chronic liver				
 One dose of vaccine for close contacts of known hepa 	atitis A cases.			
Inj 1440 ELISA units in 1 ml syringe	0.00	1	✔ н	avrix
Inj 720 ELISA units in 0.5 ml syringe	0.00	1	_	lavrix Junior

NATIONAL IMMUNISATION SCHEDULE

		Subsidy		Fully	Brand or
		(Manufacturer's Price)	Subsi	dised	Generic
		\$	Per	✓	Manufacturer
	RECOMBINANT VACCINE – [Xpharm]				
	per 0.5 ml vial	0.00	1	🖌 ні	BvaxPRO
	led for patients meeting any of the following criteria:		1	• <u> </u>	JVAAFIIO
,	for household or sexual contacts of known acute h				; or
,	for children born to mothers who are hepatitis B su	0 (0,			
3)	for children up to and under the age of 18 years in				achieved a positive
	serology and require additional vaccination or requ	iire a primary course o	f vaccinati	on; or	
4)	for HIV positive patients; or				
5)	for hepatitis C positive patients; or				
6)	for patients following non-consensual sexual interc	ourse; or			
7)	for patients following immunosuppression; or				
8)	for solid organ transplant patients; or				
9)	for post-haematopoietic stem cell transplant (HSC	F) patients; or			
,	following needle stick injury.	/1 /			
,					
Inj 10 mc	g per 1 ml vial	0.00	1	🗸 Hi	BvaxPRO
	led for patients meeting any of the following criteria:				
	for household or sexual contacts of known acute h		enatitis B	carriers	s: or
	for children born to mothers who are hepatitis B su				, 01
,	for children up to and under the age of 18 years in				achieved a nositive
0)	serology and require additional vaccination or requ				
4)	for HIV positive patients; or	ine a primary bourse e	1 vaooniau	011, 01	
,	for hepatitis C positive patients; or				
,	for patients following non-consensual sexual interc	ourse: or			
	for patients following immunosuppression; or	00136, 01			
,	for solid organ transplant patients; or				
,		T) notionto: or			
,	for post-haematopoietic stem cell transplant (HSC	r) patients, or			
10)	following needle stick injury.				
Ini 00 ma	a part minrefilled ouringe	0.00	1	./ с.	aariy D
	g per 1 ml prefilled syringe		I	• =	ngerix-B
	led for patients meeting any of the following criteria:				
,	for household or sexual contacts of known acute h		•		; or
,	for children born to mothers who are hepatitis B su				
3)	for children up to and under the age of 18 years in				achieved a positive
	serology and require additional vaccination or requ	ire a primary course o	f vaccinati	on; or	
,	for HIV positive patients; or				
	for hepatitis C positive patients; or				
6)	for patients following non-consensual sexual interc	ourse; or			
7)	for patients following immunosuppression; or				
8)	for solid organ transplant patients; or				
9)	for post-haematopoietic stem cell transplant (HSC	F) patients; or			
10)	following needle stick injury; or				
11)	for dialysis patients; or				
12)	for liver or kidney transplant patients.				
Inj 40 mc	g per 1 ml vial	0.00	1	✓ <u>H</u>	BvaxPRO
Func	led for any of the following criteria:				
1)	for dialysis patients; or				
,	for liver or kidney transplant patient.				
,					

NATIONAL IMMUNISATION SCHEDULE

	Subsidy (Manufacturer's Price) \$	Fu Subsidise Per	,
HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND Any of the following:	58) VACCINE [HPV] -	- [Xpharm]	
 Maximum of two doses for children aged 14 years and Maximum of three doses for patients meeting any of the 	,		
 People aged 15 to 26 years inclusive; or Either: 			
People aged 9 to 26 years inclusive 1) Confirmed HIV infection; or			
 Transplant (including stem cell) patients: o Maximum of four doses for people aged 9 to 26 years 		nerapy	
Inj 270 mcg in 0.5 ml syringe	·	10 •	Gardasil 9

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Su Per	Ibsidised	Generic Manufacturer
INFLUENZA VACCINE	Ψ	1.01		Mandiactarci
Inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine)			
[Xpharm]		1		luarix Tetra
A) INFLUENZA VACCINE – child aged 6 months to		1	• 1	
is available each year for patients aged 6 months t		et the fo	llowing cr	iteria, as set by
PHARMAC:				
 have any of the following cardiovascular dise 	ases			
a) ischaemic heart disease, or				
b) congestive heart failure, or				
c) rheumatic heart disease, or				
 congenital heart disease, or cerebo-vascular disease; or 				
ii) have either of the following chronic respirator	n dispasos.			
a) asthma, if on a regular preventative the				
b) other chronic respiratory disease with in		or		
iii) have diabetes; or	······································	•		
iv) have chronic renal disease; or				
v) have any cancer, excluding basal and squar	nous skin cancers if no	ot invasi	ve; or	
vi) have any of the following other conditions:				
a) autoimmune disease, or				
 b) immune suppression or immune deficie 	ency, or			
c) HIV, or				
d) transplant recipients, or	udana au			
 e) neuromuscular and CNS diseases/diso f) haemoglobinopathies, or 	rders, or			
g) on long term aspirin, or				
h) have a cochlear implant, or				
i) errors of metabolism at risk of major me	etabolic decompensat	ion, or		
j) pre and post splenectomy, or		,		
k) down syndrome, or				
vii) have been hospitalised for respiratory illness	or have a history of s	ignificar	nt respirat	ory illness;
Unless meeting the criteria set out above, the follo	wing conditions are ex	kcluded	from fund	ling:
 asthma not requiring regular preventative the 				
 b) hypertension and/or dyslipidaemia without ev 	Ũ			
B) Doctors are the only Contractors entitled to claim p 60 mcg in 0.5 ml syringe (paediatric quadrivalent v immunisation and they may only do so in respect of	accine) to patients elig	gible un	der the al	bove criteria for subsidised

Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)......90.00 10 🖌 Influvac Tetra

Subsidy	5	Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

- a) Only on a prescription
- b) No patient co-payment payable
- C)

A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by PHARMAC:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or
- children aged four years or less (but over three years) who have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

MEASLES, MUMPS AND RUBELLA VACCINE - [Xpharm]

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

diluent 0.5 ml0.00 10

(Mar	Subsidy ufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE V/	ACCINE - [Xph	arm]		
Any of the following:				
 Up to three doses and a booster every five years for patients or anatomic asplenia, HIV, complement deficiency (acquired 2) One dose for close contacts of meningococcal cases; or A maximum of two doses for bone marrow transplant patient A maximum of two doses for patients following immunosupp Note: children under seven years of age require two doses 8 wee series and then five yearly. 	or inherited), or s; or ression*.	pre o	r post solid	organ transplant; or
*Immunosuppression due to steroid or other immunosuppressive	horany must ho	for a	neriod of ar	pater than 28 days
Inj 4 mcg of each meningococcal polysaccharide conjugated to a total of approximately 48 mcg of diphtheria toxoid carrier per 0.5 ml vial		1		lenactra
MENINGOCOCCAL C CONJUGATE VACCINE – [Xpharm] Any of the following:			-	
 Up to three doses and a booster every five years for patients or anatomic asplenia, HIV, complement deficiency (acquired One dose for close contacts of meningococcal cases; or A maximum of two doses for bone marrow transplant patient A maximum of two doses for patients following immunosupp 	or inherited), or s; or			
Note: children under seven years of age require two doses 8 wee series and then five yearly.	ks apart, a boos	ster do	ose three ye	ars after the primary
*Immunosuppression due to steroid or other immunosuppressive Inj 10 mcg in 0.5 ml syringe		for a 1		eater than 28 days. leisvac-C
PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE – [Xpharm] Either:				
 A primary course of four doses for previously unvaccinated i Up to three doses as appropriate to complete the primary co 59 months who have received one to three doses of PCV13 	urse of immunis			
Note: please refer to the Immunisation Handbook for the appropr	ate schedule fo	r catch	n up prograi	mmes
Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B, 7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml				
prefilled syringe	0.00	10	✓ <u>s</u>	ynflorix

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Si	ubsidised	Generic	
\$	Per	1	Manufacturer	

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- 1) One dose is funded for high risk children (over the age of 17 months and under 18 years) who have previously received four doses of PCV10; or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) with primary immune deficiencies; or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cyanosis or failure; or
 - with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.
- Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,		
5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml		
syringe0.00	10	Prevenar 13

1

Prevenar 13

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✔	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE – Either:	[Xpharm]		
 Up to three doses (as appropriate) for patients with H chemotherapy; pre- or post-splenectomy or with funct complement deficiency (acquired or inherited), cochle All of the following: 	tional asplenia, pre- or p	oost-solid organ t	ransplant, renal dialysis,
a) Patient is a child under 18 years for (re-)immunib) Treatment is for a maximum of two doses; andc) Any of the following:	sation; and		
 i) on immunosuppressive therapy or radiatio immune response; or ii) with primary immune deficiencies; or 	n therapy, vaccinate wh	nen there is expe	cted to be a sufficient

- iii) with HIV infection; or
- iv) with renal failure, or nephrotic syndrome; or
- v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
- vi) with cochlear implants or intracranial shunts; or
- vii) with cerebrospinal fluid leaks; or
- viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
- ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
- x) pre term infants, born before 28 weeks gestation; or
- xi) with cardiac disease, with cyanosis or failure; or
- xii) with diabetes; or
- xiii) with Down syndrome; or
- xiv) who are pre-or post-splenectomy, or with functional asplenia.

Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each

23 pneumococcal serotype)	0.00	1	Pneumovax	<u>23</u>
POLIOMYELITIS VACCINE – [Xpharm]				
Up to three doses for patients meeting either of the following:				
1) For partially vaccinated or previously unvaccinated indiv	iduals; or			
For revaccination following immunosuppression.				
Note: Please refer to the Immunisation Handbook for approp	riate schedule for	catch-up pr	ogrammes.	
Inj 80D antigen units in 0.5 ml syringe	0.00	1	✓ IPOL	
ROTAVIRUS ORAL VACCINE – [Xpharm]				
Maximum of two doses for patients meeting the following:				
1) first dose to be administered in infants aged under 14 w	eeks of age; and			
no vaccination being administered to children aged 24 w	eeks or over.			
Oral susp live attenuated human rotavirus				
1,000,000 CCID50 per dose, prefilled oral applicator	0.00	10	Rotarix	

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] – [Xpharm] Either:				
1) Maximum of one dose for primary vaccination for eithe	r:			
 a) Any infant born on or after 1 April 2016; or b) For previously unvaccinated children turning 11 y varicella infection (chickenpox), or 	vears old on or after 1	July	2017, who l	nave not previously had a
2) Maximum of two doses for any of the following:				
 Any of the following for non-immune patients: 				
 i) with chronic liver disease who may in future ii) with deteriorating renal function before tran iii) prior to solid organ transplant; or 		nspla	intation; or	
iv) prior to any elective immunosuppression*, of	or			
 v) for post exposure prophylaxis who are imm 				
b) For patients at least 2 years after bone marrow tr				
c) For patients at least 6 months after completion of				
 d) For HIV positive non immune to varicella with mil a) For policity with inhome errors of matcheolism at a 				
 e) For patients with inborn errors of metabolism at r varicella, or 	isk of major metabolic	uecc	Inpensatio	
 f) For household contacts of paediatric patients wh immune compromise where the household conta g) For household contacts of adult patients who hav immunocompromised, or undergoing a procedur 	ct has no clinical histo ve no clinical history of	ory of f vario	varicella, o	no are severely
has no clinical history of varicella.		p.		
* immunosuppression due to steroid or other immunosuppre 28 days		e for a	a treatment	period of greater than
Inj 2000 PFU prefilled syringe plus vial	0.00	1 10		<u>Varilrix</u> Varilrix
VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATI Funded for patients meeting either of the following criteria:	ED VACCINE [SHING	LES	VACCINE]	– [Xpharm]
1) One dose for all people aged 65 years; or				
2) One dose for all people aged between 66 and 80 years	s inclusive from 1 Apri	l 201	8 and 31 M	arch 2020.
Inj 19,400 PFU prefilled syringe plus vial	0.00	1 10		Zostavax Zostavax
Diagnostic Agents				
TUBERCULIN PPD [MANTOUX] TEST – [Xpharm] Inj 5 TU per 0.1 ml, 1 ml vial	0.00	1		Tubersol
	0.00	I.		

- Symbols -

3TC	. 106
50X 3.0 Reservoir	25
- A -	
A-Scabies	68
Abacavir sulphate	105
Abacavir sulphate with	
lamivudine	105
Abiraterone acetate	167
Acarbose	
Acarbose Mylan	۱۱ ۱۹
Accarb	۱۱ ۱۹
Accuretic 10	
Accuretic 20	48
Acetazolamide	222
Acetic acid with 1, 2- propanediol	
diacetate and	
benzethonium	220
Acetic acid with hydroxyquinoline a	
ricinoleic acid	74
Acetylcysteine	
Aci-Jel	74
Aciclovir	
Infection	
Sensory	220
Acidex	6
Acipimox	54
Acitretin	68
Aclasta	
Aclin	
Actemra	203
Actinomycin D	157
Actrapid	10
Actrapid Penfill	10
Acupan	120
Adalat 10	
Adalat Oros	
Adalimumab	
Adapalene	60
Adefin	
Adefin XL	
Adefovir dipivoxil	100
Adenuric	117
ADR Cartridge 1.8	
Adrenaline	
ADT Booster	040
Adult diphtheria and tetanus	249
vaccine	240
Advantan	
Advate	40
Adynovate	
Afinitor	
Aflibercept	
AFT Carpimazole	82

AFT-Pyrazinamide99
Agents Affecting the
Renin-Angiotensin System 47
Agents for Parkinsonism and Related
Disorders 118
Agents Used in the Treatment of
Poisonings 225
Agrylin
Alanase
Albendazole
Albey
Albustix
Alendronate sodium
Alendronate sodium with
colecalciferol
Alfacalcidol
Alfamino Junior
Alginic acid
Alglucosidase alfa
Alkeran
Allersoothe
Allmercap155
Allopurinol116
Alpha-Adrenoceptor Blockers47
Alpha-Keri Lotion65
Alphamox 12591
Alphamox 25091
Alprolix
Alu-Tab6
Aluminium hydroxide6
Amantadine hydrochloride118
Ambrisentan
Amiloride hydrochloride
Amiloride hydrochloride with
furosemide
hydrochlorothiazide
Aminophylline
Amiodarone hydrochloride
Amisulpride131
Amitriptyline 124
Amlodipine51
Amorolfine61
Amoxicillin
Amoxicillin with clavulanic acid91
Amphotericin B 32
Amsacrine156
AmsaLyo156
Amsidine156
Amzoate
Anaesthetics
Anagrelide hydrochloride 156
Analgesics 120

Anastrozole	169
Andriol Testocaps	79
Androderm	79
Animas Battery Cap	
Animas Cartridge	25
Anoro Ellipta	
Antabuse	148
Antacids and Antiflatulents	6
Anten	. 124
Anthelmintics	88
Antiacne Preparations	60
Antiallergy Preparations	
Antianaemics	
Antiandrogen Oral	
Contraceptives	73
Antiarrhythmics	49
Antibacterials	
Antibacterials Topical	
Anticholinergic Agents	215
Anticholinesterases	110
Antidepressants	12/
Antidiarrhoeals	+ <u>۲</u> ۱
Antiepilepsy Drugs	
Antifibrinolytics, Haemostatics and	120
Local Sclerosants	20
Antifibrotics	
Antifungals	
Antifungals Topical	90 61
Antihistamines	
Antihypotensives	
Antimalarials	98
Antimigraine Preparations Antinausea and Vertigo Agents	. 130
Antiparasitics	
Antipruritic Preparations	
Antipsychotics	
Antiretrovirals	
Antirheumatoid Agents	111
Antispasmodics and Other Agents	~
Altering Gut Motility	8
Antithrombotic Agents	41
Antithymocyte globulin	470
(equine)	
Antitrichomonal Agents	98
Antituberculotics and	
Antileprotics	98
Antiulcerants	
	100
Antivirals	
Antivirals Anxiolytics	. 134
Antivirals Anxiolytics Anzatax	. 134 . 159
Antivirals Anxiolytics Anzatax Apidra	134 159 10
Antivirals Anxiolytics Anzatax Apidra Apidra SoloStar	134 159 10 10
Antivirals Anxiolytics Anzatax Apidra	134 159 10 10 51

Apo-Azithromycin
Apo-Bromocriptine118
Apo-Ciclopirox61
Apo-Cilazapril
Apo-Cilazapril/
Hydrochlorothiazide 48
Apo-Clarithromycin
Alimentary
Infection
Apo-Clomipramine124
Apo-Diclo SR110
Apo-Diltiazem CD
Apo-Doxazosin
Apo-Folic Acid
Apo-Gabapentin
Apo-Leflunomide
Apo-Megestrol
Apo-Meteorolal
Apo-Metoprolol
Apo-Mirtazapine
Apo-Moclobemide
Apo-Montelukast
Apo-Nadolol
Apo-Nicotinic Acid
Apo-Ondansetron 131
Apo-Oxybutynin
Apo-Paroxetine
Apo-Perindopril47
Apo-Pindolol
Apo-Pravastatin54
Apo-Prazosin47
Apo-Prednisone79
Apo-Primidone128
Apo-Propranolol51
Apo-Pyridoxine
Apo-Ropinirole118
Apo-Selegiline S29118
Apo-Sumatriptan130
Apo-Terazosin47
Apo-Timol51
Apomorphine hydrochloride118
Aprepitant 130
Apresoline56
Aptamil Gold+ Pepti Junior247
Aqueous cream65
Aripiprazole131
Aripiprazole Sandoz131
Aristocort64
Arrow - Clopid41
Arrow-Amitriptyline 124
Arrow-Bendrofluazide53
Arrow-Brimonidine 223
Arrow-Calcium
Arrow-Diazepam134
Arrow-Doxorubicin 158
Arrow-Fluoxetine 125
Arrow-Lamotrigine 127

Arrow-Losartan &
Hydrochlorothiazide48
Arrow-Morphine LA122
Arrow-Norfloxacin 109
Arrow-Ornidazole98
Arrow-Quinapril 1047
Arrow-Quinapril 2047
Arrow-Quinapril 547
Arrow-Roxithromycin90
Arrow-Sertraline
Arrow-Timolol
Arrow-Tolterodine
Arrow-Topiramate129
Arrow-Tramadol124
Arsenic trioxide
Asacol
Asamax7
Ascorbic acid
Aspen Adrenaline
Aspirin Blood41
Nervous120
Asthalin
Atazanavir sulphate 106
Atenolol
ATGAM
Ativan
Atomoxetine
Atorvastatin
Atropine sulphate
Cardiovascular
Sensory
Atropt
Atrovent
All Synacthen S29
Aubagio
Augmentin
Aurorix
AutoSoft 30
AutoSoft 9023–24
Avelox
Avonex
Avonex Pen
Azacitidine
Azacitidine Dr Reddy's
Azathioprine170
Azithromycin
Azol
Azopt
AZT
-B-
B-D Micro-Fine
B-D Ultra Fine
B-D Ultra Fine II14

Bacillus Calmette-Guerin (BCG)
vaccine 176
Bacillus Calmette-Guerin
vaccine 249
Baclofen117
Bactroban61
Barrier Creams and Emollients65
BCG Vaccine
Beclazone 100213
Beclazone 250213
Beclazone 50213
Beclomethasone
dipropionate 213, 218
Bee venom allergy treatment212
Bendamustine hydrochloride
Bendrofluazide
Bendroflumethiazide
[Bendrofluazide]
BeneFIX
Benzathine benzylpenicillin
Benzatropine mesylate 118
Benzbromaron AL 100
Benzbromarone
Benzoin
Benztrop
Benzydamine hydrochloride
Benzylpenicillin sodium [Penicillin
G]
Berpu13
Beta Cream63
Beta Ointment63
Beta Scalp 69
Beta-Adrenoceptor Agonists215
Beta-Adrenoceptor Blockers50
Betadine66
Betadine Skin Prep66
Betaferon 142
Betahistine dihydrochloride130
Betaine28
Betaloc CR 50
Betamethasone dipropionate63
Betamethasone dipropionate with
calcipotriol 68
Betamethasone sodium phosphate
with betamethasone acetate
Betamethasone valerate63, 69
Betamethasone valerate with
clioquinol64
Betamethasone valerate with sodium
fusidate [fusidic acid]
Betaxolol
Betnovate
Betnovate-C
Betoptic
Betoptic S
Bezafibrate 53

Bezalip53
Bezalip Retard
Bicalutamide
Bicillin LA
BiCNU
Bile and Liver Therapy
Biltricide
Bimatoprost
Bimatoprost Multichem
Binatoprost Multichem
Bindex
Biodone
Biodone Extra Forte
Biodone Forte
Bisacodyl
Bisoprolol fumarate
BK Lotion
Bleomycin sulphate 157
Blood Colony-stimulating Factors
Factors 44
Blood glucose diagnostic test
meter 12
Blood glucose diagnostic test
strip 13
Blood glucose test strips (visually
impaired)
Blood Ketone Diagnostic Test
Strip 11
Bonjela32
Bonjela

Bupropion hydrochloride	
Buscopan	ð
Buspirone hydrochloride	
Busulfan	152
- C -	
Cabergoline	. 87
Cafergot	130
Cafergot S29	130
Caffeine citrate	
Calamine	
Calcipotriol	
Calcitonin	77
Calcitriol	
Calcitriol-AFT	33
Calcium carbonate6,	3/
Calcium Channel Blockers	51
Calcium Disodium Versenate	
Calcium folinate Calcium Folinate Ebewe	154
Calcium Folinate Sandoz	154
Calcium gluconate	. 34
Calcium Homeostasis	
Calcium polystyrene sulphonate	. 46
Calcium Resonium	.46
Calogen	232
Calsource	
Candesartan cilexetil	.48
Candestar	.48
Canesten	.61
Capecitabine	
Capoten	
Capsaicin	
Musculoskeletal	111
Nervous	
Captopril	
Carafate	
Carbaccord	
Carbamazepine	
Carbimazole	021
Carbomer	
Carboplatin	152
Carboplatin Ebewe	152
Carbosorb-X	
Cardinol LA	
CareSens Dual	
CareSens N 12-	
CareSens N POP	
CareSens N Premier	
CareSens PRO	. 13
Carmellose sodium with gelatin and	
pectin	32
Carmustine	152
Carvedilol	
Carvedilol Sandoz	. 50
Catapres	.52

	119
CeeNU	152
Cefaclor monohydrate	. 88
Cefalexin	. 88
Cefalexin Sandoz	88
Cefazolin	
Ceftriaxone	88
Cefuroxime axetil	88
Celebrex	111
Celecoxib	111
Celecoxib Pfizer	111
Celestone Chronodose	
Celiprolol	50
Cellcept	170
Celol	
Centrally-Acting Agents	52
Cephalexin ABM	. 88
Cerezyme	30
Cetirizine hydrochloride	213
Cetomacrogol	
Cetomacrogol with glycerol	65
Cetuximab	184
Charcoal	
Chemotherapeutic Agents	
Chickenpox vaccine	
Chlorafast	220
Chlorambucil	152
Chloramphenicol	
Chlorhexidine aluconate	
Alimentary	32
Dermatological	64
Chloroform	228
011101010101111	
Chlorothiazide	53
Chlorothiazide	. 53
Chlorothiazide Chlorpheniramine maleate	. 53 213
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig	. 53 213 131 220
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig	. 53 213 131 220
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone]	53 213 131 220 53
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone] Chlorthalidone	53 213 131 220 53 53
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone] Chlorthalidone Chlorvescent	53 213 131 220 53 53 46
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone] Chlorthalidone Chlorvescent Choice Load 375 Choice TT380 Short	53 213 131 220 53 53 46 71 71
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone] Chlorthalidone Chlorvescent Choice Load 375 Choice TT380 Short Choice TT380 Standard	53 213 131 220 53 53 46 71 71
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone] Chlorthalidone Chlorthalidone Chlorvescent Choice Load 375 Choice TT380 Standard Choice TT380 Standard Choline salicylate with cetalkonium	53 213 131 220 53 53 53 46 71 71
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone] Chlorthalidone Chlorvescent Choice Load 375 Choice TT380 Short Choice TT380 Standard Choline salicylate with cetalkonium chloride	53 213 131 220 53 46 71 71 71
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone] Chlorthalidone Chlore Load 375 Choice TT380 Short Choice TT380 Standard Choline salicylate with cetalkonium chloride Ciclopirox olamine	53 213 131 220 53 46 71 71 71 32 61
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone] Chlorthalidone Chlore Load 375 Choice TT380 Short Choice TT380 Standard Choline salicylate with cetalkonium chloride Ciclopirox olamine	53 213 131 220 53 46 71 71 71 32 61
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone] Chlorthalidone Chlorvescent Choice Load 375 Choice TT380 Short Choice TT380 Standard Choline salicylate with cetalkonium chloride	53 213 131 220 53 53 46 71 71 71 32 61 210
Chlorothiazide Chlorpheniramine maleate Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlorthalidone [Chlorthalidone] Chlorthalidone Chlorvescent Choice Load 375 Choice TT380 Short Choice TT380 Short Choice TT380 Standard Choine salicylate with cetalkonium chloride Cicloppirox olamine Ciclosporin Ciclazapril Cilazapril with	53 213 131 220 53 46 71 71 71 32 61 210 47
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone] Chlorthalidone Chlorthalidone Choice Load 375 Choice TT380 Short Choice TT380 Standard Choline salicylate with cetalkonium chloride Ciclopirox olamine Ciclosporin	53 213 131 220 53 46 71 71 71 32 61 210 47
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone] Chlorthalidone Chlorthalidone Choice Load 375 Choice TT380 Short Choice TT380 Short Choice TT380 Standard Choline salicylate with cetalkonium chloride Ciclopirox olamine Cicloprox olamine Ciclosporin Cilazapril Cilazapril with hydrochlorothiazide	53 213 131 220 53 46 71 71 71 210 47 210 48
Chlorothiazide Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone] Chlorthalidone Chlorte Load 375 Choice TT380 Short Choice TT380 Short Choice TT380 Standard Choice TT380 Standard Choire alicylate with cetalkonium chloride Ciclopprox olamine Ciclosporin Ciclazapril Cilazapril with hydrochlorothiazide Cilicaine	53 213 131 220 53 46 71 71 71 210 47 210 48 92
Chlorothiazide Chloropheniramine maleate Chlorppromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone] Chlorthalidone Chlortescent. Choice Load 375 Choice TT380 Short Choice TT380 Standard Choice TT380 Standard Choire aslicylate with cetalkonium chloride Ciclopporin Ciclosporin Cilazapril Cilazapril with hydrochlorothiazide Cilicaine	53 213 131 220 53 53 46 71 71 71 32 61 210 47 48 92 92
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone] Chlorthalidone Chlore Load 375 Choice TT380 Short Choice TT380 Short Choice TT380 Standard Choline salicylate with cetalkonium chloride Ciclopirox olamine Ciclosporin Cilazapril Cilazapril with hydrochlorothiazide Cilicaine VK Cinacalcet	53 213 131 220 53 53 46 71 71 71 210 47 48 92 92 77
Chlorothiazide Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlortalidone [Chlorthalidone] Chlorthalidone Chlore Load 375 Choice TT380 Short Choice TT380 Standard Choice TT380 Standard Choline salicylate with cetalkonium chloride Ciclopirox olamine Ciclosporin Cilosporin Cilazapril with hydrochlorothiazide Cilicaine VK Cinacalcet Cipflox	53 213 131 220 53 53 46 71 71 71 210 47 48 92 92 77
Chlorothiazide Chlorpheniramine maleate Chlorpheniramine maleate Chlorpromazine hydrochloride Chlorsig Chlorthalidone [Chlorthalidone] Chlorthalidone Chlorvescent Choice Load 375 Choice TT380 Short Choice TT380 Short Choice TT380 Standard Choine salicylate with cetalkonium chloride Cicloppirox olamine Ciclosporin Ciclazapril Cilazapril with	53 213 131 220 53 53 46 71 71 71 32 61 210 47 48 92 93

Ciprofloxacin Teva220
Circadin
Cisplatin152
Cisplatin Ebewe152
Citalopram hydrobromide 125
Cladribine
Clarithromycin
Alimentary
Infection
Clexane
Clindamycin
Clindamycin ABM
Clinicians Renal Vit
Clobazam
Clobetasol propionate
Clobetasone butyrate63
Clofazimine
Clomazol
Dermatological61
Genito-Urinary74
Clomifene citrate87
Clomipramine hydrochloride124
Clonazepam 126, 134
Clonidine
Clonidine BNM52
Clonidine hydrochloride52
Clopidogrel41
Clopine 132
Clopixol
Clotrimazole
Dermatological61
Genito-Urinary74
Clozapine
Clozaril
Clustran
Co-trimoxazole
Coal tar
Coal tar with allantoin, menthol,
phenol and sulphur
Coal tar with salicylic acid and
sulphur
Coco-Scalp
Codeine phosphate
Extemporaneous
Nervous
Cogentin
Colaspase [L-asparaginase]
Colchicine
Colecalciferol
Colestid
Colestipol hydrochloride
Colgout
Colifoam
Colistin sulphomethate
Colistin-Link
Collodion flexible

Colloidal bismuth subcitrate	9
Colofac	8
Coloxyl	26
Combigan	
Compound electrolytes	
Compound electrolytes with glucose	
[Dextrose]	46
Compound hydroxybenzoate	000
Concerta	
Condoms	
Condyline	
Contact-D	
Contraceptives - Hormonal	71
Contraceptives - Non-hormonal	
Copaxone	142
Cordarone-X	. 49
Corticosteroids and Related Agents	
for Systemic Use	. 78
Corticosteroids Topical	63
Cosentyx	201
Cosmegen	157
Coumadin	
Creon 10000	
Creon 25000	
Crotamiton	
Crystaderm	
Curam	
Cvite	
Cyclizine hydrochloride	130
Cyclizine lactate	130
Cyclogyl	
Cyclopentolate hydrochloride	
Cyclophosphamide	
Cyclorin	
Cyklokapron	41
Cyproterone acetate	79
Cyproterone acetate with	70
ethinyloestradiol	73
Cystadane	
Cytarabine	154
Cytotec	8
Cytoxan	152
- D -	
D-Penamine	111
Dabigatran	
Dacarbazine	
Dacarbazine APP	
Dactinomycin [Actinomycin D]	
Daivobet	
Daivonex	68
Daktarin	
Dalacin C	
Dalteparin sodium	42
Danazol	87
Dantrium	117

	_
Dantrium S29	
Dantrolene	
Daonil	
Dapa-Tabs	
Dapsone	
Daraprim	
Darunavir	
Dasatinib	
Daunorubicin	
DBL Acetylcysteine DBL Aminophylline	220
DBL Bleomycin Sulfate	
DBL Carboplatin	157
DBL Cisplatin	
DBL Dacarbazine	157
DBL Desferrioxamine Mesylate for In	i 107
BP	
DBL Docetaxel	
DBL Ergometrine	
DBL Gemcitabine	
DBL Gentamicin	.93
DBL Leucovorin Calcium	154
DBL Methotrexate Onco-Vial	
DBL Morphine Sulphate	122
DBL Morphine Tartrate	123
DBL Naloxone Hydrochloride	225
DBL Octreotide	168
DBL Pethidine Hydrochloride	123
DBL Vincristine Sulfate	
De-Worm	
Decozol	
Deferasirox	
Deferiprone	
Denosumab	
Deolate	
Deoxycoformycin	160
Depo-Medrol	
Depo-Provera	. 73
Depo-Testosterone	
Deprim DermAssist	. 95
Dermol	
Desferrioxamine mesilate	
Desmopressin acetate	
Desmopressin-PH&T	.00 86
Detection of Substances in	.00
Urine	76
Dexamethasone	
Hormone	78
Sensory	
Dexamethasone phosphate	
Dexamethasone with framycetin and	
gramicidin	220
Dexamethasone with neomycin	
sulphate and polymyxin B	
sulphate	221

INDEX: Generic Chemicals	and	Brands
--------------------------	-----	--------

Dexamfetamine sulfate144
Dexmethsone78
Dextrochlorpheniramine
maleate
Dextrose
DHC Continus121
Diabetes9
Diabetes Management11
Diacomit128
Diagnostic Agents258
Diamide Relief6
Diamox
Diasip233
Diason RTH
Diazepam 126, 134
Diazoxide9
Dibenzyline
Diclofenac Sandoz110
Diclofenac sodium
Musculoskeletal 110
Sensory221
Differin
Difflam
Diflucan
Diflucan S29
Diflucortolone valerate
Digestives Including Enzymes25
Digoxin
Dihydrocodeine tartrate
Dilantin
Dilantin Infatab128
Diltiazem hydrochloride
Dilzem
Dimethicone
Dimethyl fumarate 135
Dipentum
Diphtheria, tetanus and pertussis
vaccine
Diphtheria, tetanus, pertussis and
polio vaccine
Diphtheria, tetanus, pertussis, polio,
hepatitis B and haemophilus
influenzae type B vaccine
Diprosone
Diprosone OV63
Dipyridamole
Disinfecting and Cleansing
Agents
Disopyramide phosphate
Disulfiram
Diuretics
Diurin 40
Docetaxel
Docetaxel Sandoz
Docusate sodium
Docusate sodium with

sennosides	
Dolutegravir	107
Domperidone	130
Donepezil hydrochloride	147
Donepezil-Rex	147
Dopress	124
Dornase alfa	218
Dortimopt	222
Dorzolamide hydrochloride	
Dorzolamide with timolol	
Dostinex	87
Dosulepin [Dothiepin]	
hydrochloride	124
Dothiepin	124
Doxazosin	47
Doxepin hydrochloride	
Doxine	92
Doxorubicin Ebewe	158
Doxorubicin hydrochloride	
Doxy-50	
Doxycycline	92
DP Fusidic Acid Cream	
DP Lotion	65
DP Lotn HC	63
DP-Allopurinol	
Dr Reddy's Omeprazole	9
Druge Affecting Pone	
Metabolism	
Dual blood glucose and blood ket	
diagnostic test meter	12
diagnostic test meter Duocal Super Soluble Powder	12 231
diagnostic test meter Duocal Super Soluble Powder Duolin	12 231 215
diagnostic test meter Duocal Super Soluble Powder Duolin Duolin HFA	12 231 215 215
diagnostic test meter Duocal Super Soluble Powder Duolin Duolin HFA Durex Confidence	12 231 215 215 71
diagnostic test meter Duocal Super Soluble Powder Duolin Duolin HFA Durex Confidence Durex Extra Safe	12 231 215 215 71 71
diagnostic test meter Duocal Super Soluble Powder Duolin Duolin HFA Durex Confidence	12 231 215 215 71 71
diagnostic test meter Duocal Super Soluble Powder Duolin Duolin HFA Durex Confidence Durex Extra Safe Duride - E -	12 215 215 71 71 55
diagnostic test meter Duocal Super Soluble Powder Duolin Duolin HFA Durex Confidence Durex Extra Safe Duride - E - e-chamber La Grande	12 215 215 71 71 55 219
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Duride	12 215 215 71 71 55 219 219
diagnostic test meter Duocal Super Soluble Powder Duolin Duolin HFA Durex Confidence Durex Extra Safe Duride e-chamber La Grande e-chamber Mask	12 215 215 71 71 75 219 219 219
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Duride - E - e-chamber La Grande e-chamber Mask. e-chamber Turbo E-Mycin.	
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Duride - E - e-chamber La Grande e-chamber Mask e-chamber Turbo E-Mycin Ear Preparations	
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Duride e-chamber La Grande e-chamber Mask e-chamber Turbo E-Mycin Ear Preparations	
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Duride - E - e-chamber La Grande e-chamber Mask e-chamber Turbo E-Mycin Ear Preparations Ear/Eye Preparations Easiphen Liguid	
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Duride e-chamber La Grande e-chamber Mask e-chamber Turbo E-Mycin Ear Preparations Ear/Eye Preparations Easiphen Liquid Econazole nitrate	
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Duride - E - e-chamber La Grande e-chamber Mask e-chamber Turbo E-Mycin Ear Preparations Ear/Eye Preparations Easiphen Liguid	
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Duride - E - e-chamber La Grande e-chamber Mask e-chamber Turbo E-Mycin Ear Preparations Ear/Eye Preparations Easiphen Liquid Econazole nitrate Efavirenz	
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Durex Extra Safe Duride - E - e-chamber La Grande e-chamber Mask e-chamber Turbo E-Mycin Ear Preparations Ear/Eye Preparations Easiphen Liquid Econazole nitrate Efavirenz.	
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Durex Extra Safe Duride - E - e-chamber La Grande e-chamber Mask e-chamber Mask e-chamber Turbo E-Mycin Ear Preparations Ear/Eye Preparations Easiphen Liquid Econazole nitrate Efavirenz Efavirenz with emtricitabine and tenofovir disoproxil	12 231 215 215 215 215 219 219 219 219 219 220 220 220 220 220 220 210 5 220 210 5 245 622 5 215 215 215 215 215 215 215 215 215
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Durex Extra Safe Duride - E - e-chamber La Grande e-chamber Mask e-chamber Mask e-chamber Turbo E-Mycin Ear Preparations Ear/Eye Preparations Easiphen Liquid Econazole nitrate Efavirenz with emtricitabine and tenofovir disoproxil	12 2311 215 215 215 215 219 219 219 220 220 220 220 220 220 220 220 220 22
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Duride - E - e-chamber La Grande e-chamber Mask e-chamber Turbo E-Mycin Ear Preparations Ear/Eye Preparations Ear/Eye Preparations Easiphen Liquid Econazole nitrate Efavirenz Efavirenz Efavirenz with emtricitabine and tenofovir disoproxil Effient Eformoterol fumarate	12 2311 215 215 215 215 219 219 219 220 220 220 220 220 220 220 220 220 22
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Duride - E - e-chamber La Grande e-chamber Turbo E-dwcin Ear Preparations Ear/Eye Preparations Ear/Eye Preparations Easiphen Liquid Econazole nitrate Efavirenz Efavirenz with emtricitabine and tenofovir disoproxil Effient Eformoterol fumarate Eformoterol fumarate dihydrate	12 231 215 215 215 215 219 219 219 200 220 220 220 220 220 220 220 245 62 105 106 41 214
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Duride - E - e-chamber La Grande e-chamber Mask e-chamber Turbo Ear Preparations Ear/Eye Preparations Ear/Eye Preparations Easiphen Liquid Econazole nitrate Efavirenz with emtricitabine and tenofovir disoproxil Effient Eformoterol fumarate Eformoterol fumarate Eftrenonacog alfa [Recombinant factor IX]	12 231 215 215 215 215 215 219 219 219 200 220 220 220 220 220 220 220 221 5 62 245 62 245 62 245 62 245 71 71 219 71 71 71 71 71 71 71 71 71 71 71 71 71
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Duride - E - e-chamber La Grande e-chamber Mask e-chamber Turbo Ear Preparations Ear/Eye Preparations Ear/Eye Preparations Ear/Eye Preparations Easiphen Liquid Econazole nitrate Efavirenz with emtricitabine and tenofovir disoproxil Effient Efformoterol fumarate Efformoterol fumarate Eftrenonacog alfa [Recombinant factor IX] Efudix Efudix	12 231 215 215 215 215 215 219 219 219 200 220 220 220 220 220 220 220 221 5 62 245 62 245 62 245 62 245 62 245 62 62 62 62 62 62 62 66 9 69 9 69 9 69
diagnostic test meter Duocal Super Soluble Powder Duolin HFA Durex Confidence Durex Extra Safe Duride - E - e-chamber La Grande e-chamber Mask e-chamber Turbo Ear Preparations Ear/Eye Preparations Ear/Eye Preparations Easiphen Liquid Econazole nitrate Efavirenz with emtricitabine and tenofovir disoproxil Effient Eformoterol fumarate Eformoterol fumarate Eftrenonacog alfa [Recombinant factor IX]	12 231 215 215 215 215 215 219 219 219 200 220 220 220 220 220 220 220 221 5 62 245 62 245 62 245 62 245 62 245 62 62 62 62 62 62 62 66 9 69 9 69 9 69

Elecare	
Elecare LCP	246
Elelyso	
Elemental 028 Extra	237
Elocon	
Elocon Alcohol Free	
Eltrombopag	
Eltroxin	
EMB Fatol	
Emcure	152
Emend Tri-Pack	
EMLA	120
Emtricitabine	106
Emtricitabine with tenofovir	
disoproxil	103
Emtriva	106
Emulsifying ointment	100
Enalapril maleate	
Enbrel	170
Endocrine Therapy	167
Endoxan	
Enerlyte	46
Engerix-B	251
Enlafax XR	126
Enoxaparin sodium	
Ensure	
Ensure Plus	241
Ensure Plus HN	
Ensure Plus RTH	
Entacapone	
Entapone	
Entecavir	
Entecavir Sandoz	
Entocort CIR	
Entresto 24/26	48
Entresto 49/51	48
Entresto 97/103	48
Epilim	
Epilim Crushable	128
Epilim IV	
Epilim S/F Liquid	128
Epilim Syrup	100
Epirubicin Ebewe	120
Epirubicin Ebewe	100
Eplerenone	
Epoetin alfa	
Epoprostenol	58
Eptacog alfa [Recombinant factor	
VIIa]	40
ERA	
Erbitux	
Ergometrine maleate	
Ergonovine	
Ergotamine tartrate with	
caffeine	120
Erlotinib	102

Erythrocin IV90
Erythromycin ethyl succinate90
Erythromycin lactobionate90
Erythromycin stearate
Esbriet
Escitalopram
Escilatopiani
Escitalopram-Apotex
Eskazole
Estradot
Estradot 50 mcg80
Estrofem
Etanercept 170
Ethambutol hydrochloride99
Ethics Aspirin120
Ethics Aspirin EC41
Ethics Enalapril47
Ethics Lisinopril47
Ethinyloestradiol
Ethinyloestradiol with
Ethinyloestradiol with desogestrel72
Ethinyloestradiol with
levonorgestrel
Evolution for the sector of th
Ethinyloestradiol with norethisterone72
Ethosuximide
Etopophos 158
Etoposide158
Etoposide phosphate158
Etravirine 105
Eumovate63
Everet
Everolimus210
Evista112
Exelon
Exemestane
Exjade
Extemporaneously Compounded
Preparations and
Galenicals 228
Eye Preparations
Eylea
Ezetimibe54
Ezetimibe Sandoz54
Ezetimibe with simvastatin55
- F -
Factor eight inhibitor bypassing
fraction
Febuxostat117
Feed Thickener Karicare
Aptamil
FEIBA NF
Felo 10 ER
Felo 5 ER
Felodipine
Fenpaed
Fentanyl121

	_
Fentanyl Sandoz12	
Ferinject	35
Ferodan	
Ferric carboxymaltose	
Ferriprox	25
Ferro-F-Tabs	35
Ferro-tab	
Ferrograd	35
Ferrosig	36
Ferrous fumarate	35
Ferrous fumarate with folic acid	
Ferrous sulphate	35
Ferrum H	
Fexofenadine hydrochloride21	3
Fibro-vein4	11
Filgrastim4	
Finasteride7	74
Fingolimod 13	36
Firazyr21	12
FlagyI	98
FlagyI-S	98
Flamazine6	
Flecainide acetate4	
Fleet Phosphate Enema	27
Flixonase Hayfever & Allergy21	8
Flixotide	14
Flixotide Accuhaler21	14
Floair21	
Florinef	
Fluanxol	
Fluarix Tetra25	
Flucil	91
Flucloxacillin	
Flucloxin	
Flucon	
Fluconazole	
Fludara Oral15	
Fludarabine Ebewe15	
Fludarabine phosphate	54
Fludrocortisone acetate	
Fluids and Electrolytes4	
Flumetasone pivalate	
Fluocortolone caproate with	
fluocortolone pivalate and	
cinchocaine	7
Fluorometholone	>?
Fluorouracil15	54
Fluorouracil Ebewe	54
Fluorouracil sodium	
Fluoxetine hydrochloride	
Flupenthixol decanoate	
Flutamide	
Flutamide Mylan	
Flutamin	
Fluticasone	
Fluticasone furoate with	. 1

vilanterol	
Fluticasone propionate	
Fluticasone with salmeterol	.214
FML	.222
Foban	61
Folic acid	
Food Thickeners	040
Food Trickeners	.243
Foods And Supplements For Inborn	
Errors Of Metabolism	. 244
Foradil	
Forteo	.113
Fortini	.235
Fortini Multi Fibre	.235
Fortisip	241
Fortisip Multi Fibre	242
Fosamax	111
Fosamax Plus	
Fragmin	42
Framycetin sulphate	
Frisium	. 126
Frumil	53
Frusemide	52
Frusemide-Claris	52
Fucicort	64
Fucidin	
Fucithalmic	
Fungilin	20
Furosemide [Frusemide]	32
	52
fusidic acid	
Dermatological6	1, 64
Infection	
Sensory	. 220
- G -	
Gabapentin	.126
Gacet	121
Galsulfase	28
Galvumet	
Galvus	1 1
Gardasil 9	11
Gastrodenol	9
Gaviscon Double Strength	
Gaviscon Infant	6
Gazyva	. 191
Gefitinib	.163
Gemcitabine Ebewe	.154
Gemcitabine hydrochloride	.154
Gemfibrozil	54
Gemzar	15/
Genoptic	
	. 220
Gentamicin sulphate	
Infection	
Sensory	. 220
Gilenya	. 136
Ginet	73
Glatiramer acetate	.142
Glecaprevir with pibrentasvir	.102

Glibenclamide11
Gliclazide 11
Glipizide11
Glivec163
Glizide11
Glucagen Hypokit9
Glucagon hydrochloride9
Glucerna Select 233
Glucerna Select RTH233
Glucobay11
Glucose [Dextrose]45
Gluten Free Foods243
Glycerin with sodium saccharin 228
Glycerin with sucrose228
Glycerol
Alimentary27
Extemporaneous228
Glyceryl trinitrate
Alimentary8
Cardiovascular55
Glycopyrronium
Glycopyrronium bromide8
Glycopyrronium with
indacaterol
Glytrin
Gold Knight71
Goserelin
Gutron49
Gynaecological Anti-infectives74
Gynaecological Anti-infectives74 - H -
Gynaecological Anti-infectives
Gynaecological Anti-infectives74 - H - Habitrol
Gynaecological Anti-infectives
Gynaecological Anti-infectives74 - H - Habitrol
Gynaecological Anti-infectives
Gynaecological Anti-infectives74 - H - Habitrol
Gynaecological Anti-infectives74 - H - Habitrol
Gynaecological Anti-infectives

Hepatitis B recombinant	
vaccine 2	251
Hepsera1	00
Herceptin2	
Hexamine hippurate1	
Hiberix	250
Hiprex 1	09
Histaclear2	
Histafen2	
Holoxan1	52
Horleys Bread Mix2	243
Horleys Flour2	
Hormone Replacement Therapy -	
Systemic	79
HPV2	252
Humalog	10
Humalog Mix 25	10
Humalog Mix 50	
Human papillomavirus (6, 11, 16, 18,	
31, 33, 45, 52 and 58) vaccine	
[HPV]2	252
Humatin	
Humira1	
HumiraPen1	
Humulin 30/70	
Humulin NPH	10
Humulin R	10
Hyaluronic acid2	
Hybloc	
Hydralazine	
Hydralazine hydrochloride	56
Hydrea1	
Hvdrocortisone	
Dermatological	63
Hormone	
Hydrocortisone acetate	
Hydrocortisone and paraffin liquid	
and lanolin	63
Hydrocortisone butyrate63,	
Hydrocortisone with cinchocaine	
Hydrocortisone with miconazole	64
Hydrocortisone with natamycin and	
neomycin	64
Hvdrogen peroxide	
Alimentary	33
Dermatological	
Hydroxocobalamin	33
Hydroxychloroquine1	
Hydroxyurea1	58
Hygroton	53
Hylo-Fresh2	24
Hymenoptera2	
Hyoscine butylbromide	
Hyoscine hydrobromide1	30
Hypam 1	44
Hyperuricaemia and Antigout 1	16

Hypromellose	224
Hypromellose with dextran	224
Hysite	223
-1-	
Ibiamox	
Ibuprofen	110
Icatibant	212
Idarubicin hydrochloride	158
Idursulfase	29
Ifosfamide	152
Ikorel	
lloprost	
Imatinib mesilate	
Imatinib-AFT	
Imiglucerase	
Imipramine hydrochloride	124
Imiquimod	70
Immune Modulators	
Immunosuppressants	
Imuran	
Incruse Ellipta	216
Indacaterol	
Indapamide	
Infanrix IPV	
Infanrix-hexa	
Infant Formulae	246
Infatrini	
Infliximab	
Influenza vaccine	
Influvac Tetra	
Inhaled Corticosteroids	
Inhaled Long-acting	213
Beta-adrenoceptor Agonists	014
bela-aurenoceptor Agonists	214
Inset 30	21
Inset II	
Inspra	
Insulin aspart	10
Insulin aspart with insulin aspart	
protamine	10
Insulin glargine	
Insulin glulisine	10
Insulin isophane	10
Insulin isophane with insulin	
neutral	
Insulin lispro	10
Insulin lispro with insulin lispro	
protamine	
Insulin neutral	10
Insulin pen needles	13
Insulin pump	14
Insulin pump accessories	
Insulin pump cartridge	19
Insulin pump infusion set (steel	
cannula)	20
Insulin pump infusion set (steel	
cannula, straight insertion)	21

Insulin pump infusion set (teflon
cannula, angle insertion with
insertion device) 21
Insulin pump infusion set (teflon
cannula, angle insertion) 22
Insulin pump infusion set (teflon
cannula, straight insertion with
insertion device)
Insulin pump infusion set (teflon
cannula, straight insertion)
Insulin pump reservoir
Insulin syringes, disposable with
attached needle
Intal Forte CFC Free
Intelence
Interferon alfa-2a 107
Interferon beta-1-alpha 142
Interferon beta-1-beta142
Intra-uterine device71
Invega Sustenna133
IPOL
Ipratropium bromide 215, 219
Iressa163
Irinotecan Accord155
Irinotecan Actavis 100 155
Irinotecan hydrochloride155
Irinotecan-Rex155
Iron polymaltose
Isentress 107
Ismo 2055
Ismo 40 Retard55
Isoniazid
Isoniazid with rifampicin
Isoprenaline [Isoproterenol]
Isoproterenol
Isoptin
Isopto Carpine
Isosorbide mononitrate
Isosource Standard
Isotretinoin
Ispaghula (psyllium) husk
Isuprel
Itch-Soothe
Itraconazole
Itrazole
Ivermectin
- J -
Jadelle73
Jakavi 166
Jevity HiCal RTH240
Jevity RTH240
Juno Pemetrexed 155
- K -
Kaletra 106
Kemadrin 118
Kenacomb220

Kenacort-A 1079
Kenacort-A 4079
Kenalog in Orabase32
Ketocal 3:1
KetoCal 4:1248
Ketoconazole
Dermatological
Infection
Ketogenic Diet
Ketoprofen 110
KetoSens11
Keytruda
Kindergen
Kinson118
Kivexa
Klacid
Kliogest80
Kliovance
Kogenate FS41
Konakion MM41
Konsyl-D
Kuvan
- L -
L-asparaginase
Labetalol
Lacosamide
Lactulose
Laevolac
Lamictal
Lamivudine 100, 106
Lamivudine Alphapharm
Lamotrigine
Lamprene
Lanoxin
Lanoxin PG49
Lanoxin S29
Lansoprazole
Lantus10
Lantus SoloStar10
Lanvis
Lanzol Relief
Lapatinib ditosylate164
Largactil
Laronidase
Lasix
Latanoprost
Lax-Suppositories
Lax-Tab
Laxatives
Laxsol
Ledipasvir with sofosbuvir 102
Leflunomide 111
Lenalidomide
Letrole170
Letrozole170
Leukeran FC152

Leukotriene Receptor	
Antagonists	217
Leunase	157
Leuprorelin	86
Leustatin	154
Levetiracetam	127
Levetiracetam-AFT	127
Levlen ED	72
Levocabastine	
Levodopa with benserazide	118
Levodopa with carbidopa	118
Levomepromazine	
hydrochloride	132
Levomepromazine maleate	132
Levonorgestrel	
Genito-Urinary	
Hormone	
Levothyroxine	82
Lidocaine [Lignocaine]11	
Lidocaine [Lignocaine]	
hydrochloride	120
Lidocaine [Lignocaine] with	
chlorhexidine	120
Lidocaine [Lignocaine] with	
prilocaine	120
Lidocaine-Claris	120
Lignocaine11	9–120
Lioresal Intrathecal	117
Lipazil	54
Lipid-Modifying Agents	53
Liquigen	232
Lisinopril	47
Lithicarb FC	132
Lithium carbonate	132
Livostin	222
LMX4	120
Locacorten-Viaform ED's	220
Local preparations for Anal and	
Rectal Disorders	7
Locasol	246
Locoid	63, 69
Locoid Crelo	63
Locoid Lipocream	63
Locorten-Vioform	220
Lodi	
Lodoxamide	
Logem	127
Lomide	222
Lomustine	152
Loniten	56
Loperamide hydrochloride	6
Lopinavir with ritonavir	106
Loprofin	
Loprofin Mix	
Lorafix	213
Loratadine	213

Lorazepam	135
Lorfast	213
Lorstat	. 54
Losartan Actavis	48
Losartan potassium	. 48
Losartan potassium with	
hydrochlorothiazide	. 48
Lovir	
Lucrin Depot 1-month	
Lucrin Depot 3-month	. 86
Ludiomil	124
Lyderm	. 68
- M -	
m-Eslon	
Mabthera	194
Macrogol 3350 with potassium	
chloride, sodium bicarbonate and	
sodium chloride	. 27
Macrogol 400 and propylene	
glycol	224
Madopar 125	118
Madopar 250	118
Madopar 62.5	118
Madopar HBS	118
Madopar Rapid	
Magnesium hydroxide	
Magnesium sulphate	
Mantoux	
Maprotiline hydrochloride	124
Marevan	. 44
Marine Blue Lotion SPF 50+	. 69
Marvelon 28	
Mask for spacer device	219
Mast Cell Stabilisers	
Maviret	
Maxidex	
Maxitrol	221
MCT oil (Nutricia)	
Measles, mumps and rubella	
vaccine	254
Mebendazole	88
Mebeverine hydrochloride	8
Medrol	
Medroxyprogesterone acetate	
Genito-Urinary	73
Hormone	-81
Mefenamic acid	
Megestrol acetate	168
Melatonin	
Melphalan	
Menactra	
Meningococcal (groups A, C, Y and	
W-135) conjugate vaccine	255
Meningococcal C conjugate	
vaccine	255
Menthol	

Mercaptopurine15	5
Mercilon 287	5
Mesalazine	
Mesna15	
Mestinon 11	
Metabolic Disorder Agents2	
Meterol21	4
Metformin hydrochloride1	1
Methadone hydrochloride	
Extemporaneous22	8
Nervous12	2
Methatabs12	
Methopt	
Methotrexate	
Methotrexate Ebewe15	
Methotrexate Sandoz15	
Methyl hydroxybenzoate22	
Methylcellulose 22	9
Methylcellulose with glycerin and	
sodium saccharin	9
Methylcellulose with glycerin and	
sucrose	q
Methyldopa5	
Methyldopa Mylan	2
Methyldopa Mylan S295	
Methylnaltrexone bromide2	
Methylphenidate hydrochloride14	5
Methylphenidate hydrochloride	
extended-release 14	6
Methylprednisolone7	8
Methylprednisolone (as sodium	
succinate)7	8
Methylprednisolone aceponate6	
Methylprednisolone acetate	
Methylxanthines	
Metoclopramide Actavis 10 13	
Metoclopramide hydrochloride 13	
Metolazone	
Metopirone8	
Metoprolol succinate5	
Metoprolol tartrate5	0
Metronidazole9	8
Metroprolol IV Mylan5	0
Metyrapone8	
Mexiletine hydrochloride	9
Mexiletine Hydrochloride USP4	,q
Miacalcic	7
Miconazole3	2
Miconazole nitrate	
Dermatological6	
Genito-Urinary7	4
Micreme7	4
Micreme H6	4
Microgynon 20 ED7	2
Microgynon 307	
	-

Microgynon 50 ED	.72
Microlut	.73
Midazolam	143
Midazolam-Claris	143
Midodrine	.49
Minerals	. 34
Mini-Wright AFS Low Range	219
Mini-Wright Standard	219
Minidiab	11
MiniMed 640G	14
Minims Pilocarpine	223
Minims Prednisolone	220
Minimis i realissione	
Mino-tabs	
Minocycline hydrochloride	. 92
Minomycin Minor Skin Infections	.92
Minoxidil	
Mirena	
Mirtazapine	
Misoprostol	8
Mitomycin C	159
Mitozantrone	
Mitozantrone Ebewe	
Mixtard 30	
Moclobemide	125
Modafinil	
Modavigil	147
Moduretic	E 0
	. 00
Molaxole	.27
Molaxole Mometasone furoate	.27 .64
Molaxole Mometasone furoate	.27 .64
Molaxole Mometasone furoate Monogen Montelukast	.27 .64 234 217
Molaxole Mometasone furoate Monogen Montelukast	.27 .64 234 217
Molaxole Mometasone furoate Monogen Montelukast Moroctocog alfa [Recombinant factor	. 27 . 64 234 217
Molaxole Mometasone furoate Monogen Montelukast Moroctocog alfa [Recombinant factor VIII]	. 27 . 64 234 217 . 40
Molaxole Mometasone furoate Monogen Montelukast Moroctocog alfa [Recombinant factor VIII] Morphine hydrochloride	.27 .64 234 217 .40 122
Molaxole Mometasone furoate Monogen Morotlukast Moroctocog alfa [Recombinant factor VIII] Morphine hydrochloride Morphine sulphate	.27 .64 234 217 .40 122 122
Molaxole Mometasone furoate Monogen Morotlukast Moroctocog alfa [Recombinant factor VIII] Morphine hydrochloride Morphine sulphate Morphine tartrate	.27 .64 234 217 .40 122 122 123
Molaxole Mometasone furoate Monogen Moroctocog alfa [Recombinant factor VIII] Morphine hydrochloride Morphine sulphate Morphine tartrate Motetis	.27 .64 234 217 .40 122 122 123
Molaxole	.27 .64 234 217 .40 122 122 123 119 .32
Molaxole	.27 .64 234 217 .40 122 123 123 119 .32
Molaxole	.27 .64 234 217 .40 122 123 112 .32 .32 .118 .93
Molaxole	.27 .64 234 217 .40 122 123 112 .32 .32 .118 .93
Molaxole	.27 .64 234 217 .40 122 123 112 .32 118 .93 244
Molaxole	.27 .64 234 217 .40 122 123 123 123 119 .32 118 .93 244
Molaxole	.27 .64 234 217 .40 122 123 123 119 .32 118 .93 244 .26 218
Molaxole	.27 .64 234 217 .40 122 123 112 .32 118 .93 244 .26 218 120
Molaxole	.27 .64 234 217 .40 122 123 112 .32 118 .93 244 .26 218 120 135
Molaxole	.27 .64 234 217 .40 122 123 122 123 119 .32 118 .93 244 .26 218 120 135 .34
Molaxole	.27 .64 234 217 .40 122 123 112 .32 118 .93 244 .26 218 120 135 .34 .34
Molaxole	.27 .64 234 217 .40 122 123 112 .32 118 .93 244 .26 218 120 135 .34 .34 .61
Molaxole	.27 .64 234 217 .40 122 123 123 123 123 .32 244 .26 218 120 135 .34 .34 .61 117
Molaxole	.27 .64 234 217 .40 122 122 123 119 .32 119 .32 244 .26 218 120 .34 .34 .34 .34 .34 .34 .34
Molaxole	.27 .64 234 217 .40 122 123 119 .32 118 .93 244 .26 218 120 135 .34 .34 .61 117 .34 .99
Molaxole	.27 .64 234 217 .40 122 123 119 .32 118 .93 244 .26 218 120 135 .34 .34 .61 117 .34 .99 .99

Mycophenolate mofetil170
Mycostatin62
Mydriacyl 223
Mylan Atenolol
Mylan Clomiphen87
Myleran152
Myocrisin111
Myometrial and Vaginal Hormone
Preparations
Myozyme
Mysoline S29128
- N -
Nadolol
Naglazyme
Nalcrom
Naloxone hydrochloride
Naltraccord
Naltrexone hydrochloride148
Naphazoline hydrochloride
Naphcon Forte
Naprosyn SR 1000 110
Naprosyn SR 750 110
Naproxen 110
Nardil
Nardil S29
Nasal Preparations
Natalizumab
Natulan
Nausafix
Nausicalm
Navelbine
Nedocromil
Nefopam hydrochloride
Neisvac-C
Neo-B12
Neo-Mercazole
Neocate Gold
Neocate Junior Unflavoured
Neocate Junior Vanilla
Neocate LCP
Neocate SYNEO
Neoral
Neostigmine metilsulfate
Nepro HP (strawberry)
Nepro HP (vanilla)
Nepro HP RTH
Nerisone
Neulactil
NeuroTabs
Nevirapine
Nicorondil
Nicorandil
Nicotinic acid
Nifedipine51

Nifuran
Nilotinib164
Nilstat
Alimentary 33
Genito-Urinary74
Infection96
Nintedanib216
Nipent
Nitrados143
Nitrates
Nitrazepam143
Nitroderm TTS55
Nitrofurantoin 109
Nitrolingual Pump Spray55
Nivestim
Nivolumab
Nizoral
Nodia
Noflam 250 110
Noflam 500 110 Non-Steroidal Anti-Inflammatory
Non-Steroidal Anti-Inflammatory
Drugs
Nonacog alfa [Recombinant factor
IX] 40
Nonacog gamma, [Recombinant
Factor IX] 40
Norethisterone
Genito-Urinary73
Hormone81
Norflex 117
Norfloxacin109
Noriday 28
Norimin
Normacol Plus
Normison143
Norpress 124
Nortriptyline hydrochloride124
Norvir
NovaSource Renal
Novatretin
NovoMix 30 FlexPen 10
NovoRapid10
NovoRapid FlexPen10
NovoRapid Penfill
NovoSeven RT40
Noxafil
Nozinan
Nuelin-SR
Nutilis
Nutrient Modules
Nutrini Energy Multi Fibre
Nutrini Energy RTH235
Nutrini Low Energy Multi Fibre237
Nutrini RTH235
Nutrison 800 Complete Multi

Fibre	040
Nutrison Concentrated	
Nutrison Energy	240
Nutrison Energy Multi Fibre	240
Nutrison Multi Fibre	240
Nutrison Standard RTH	240
Nyefax Retard	51
Nystatin	
Alimentary	33
Dermatological	62
Genito-Urinary	74
Infection	
NZB Low Gluten Bread Mix	040
- 0 -	240
	05
O/W Fatty Emulsion Cream	
Obinutuzumab	191
Octocog alfa [Recombinant factor	
VIII] (Advate)	40
Octocog alfa [Recombinant factor	
VIII] (Kogenate FS)	41
Octreotide	168
Octreotide LAR (somatostatin	
analogue)	168
Oestradiol	
Oestradiol valerate	
Oestradiol with norethisterone	
Oestriol	
Genito-Urinary	74
Hormone	81
Oestrogens	80
Ofev	
Oil in water emulsion	210
	00
Olanzapine	2-133
Olbetam	
Olopatadine	
Olsalazine	7
Omalizumab	192
Omeprazole	9
Omeprazole actavis 10	9
Omeprazole actavis 20	
Omeprazole actavis 40	9
Omnitrope	
Onbrez Breezhaler	214
Oncaspar	159
OncoTICE	176
Ondansetron	131
Ondansetron ODT-DRLA	131
Ondansetron ODT-ORLA	131
One-Alpha	
Opdivo	
Ora-Blend	
Ora-Blend SF	
Ora-Plus	
Ora-Sweet	
Ora-Sweet SF	
Orabase	

Oral Supplements/Complete Diet
(Nasogastric/Gastrostomy Tube
Feed) 233
Oratane
Ordine122
Orgran
Orion Temozolomide 160
Ornidazole
Orphenadrine citrate117
Ortho-tolidine
Oruvail SR 110
Osmolite RTH240
Other Endocrine Agents87
Other Oestrogen Preparations80
Other Progestogen
Preparations 81
Other Skin Preparations70
Ovestin
Genito-Urinary74
Hormone
Ox-Pam
Oxaliccord153
Oxaliplatin
Oxaliplatin Actavis 100
Oxaliplatin Ebewe
Oxazepam
Oxis Turbuhaler
Oxpentifylline
Oxybutynin
Oxycodone hydrochloride
Oxycodone Sandoz
OxyNorm
Oxytocin
Oxytocin BNM
Oxytocin with ergometrine
maleate
Ozurdex
- P -
Pacifen117
Paclitaxel
Paclitaxel Actavis
Paclitaxel Ebewe
Paediatric Seravit
Paliperidone
Pamidronate disodium
Pamisol
Pancreatic enzyme
Pantoprazole
Panzop Relief
Panzytrat
Panzytrat
Para-amino salicylic acid
Paracare
Paracare Double Strength 121
Paracetamol
Paracetamol + Codeine

(Relieve) 123
Paracetamol Pharmacare 121
Paracetamol with codeine123
Paradigm 1.8 Reservoir25
Paradigm 3.0 Reservoir
Paradigm Mio MMT-921
Paradigm Mio MMT-92323
Paradigm Mio MMT-92523
Paradigm Mio MMT-94123
Paradigm Mio MMT-94323
Paradigm Mio MMT-94523
Paradigm Mio MMT-96523
Paradigm Mio MMT-97523
Paradigm Mio MMT-97523 Paradigm Quick-Set MMT-38624
Paradigm Quick-Set MMT-38724
Paradigm Quick-Set MMT-396
Paradigm Quick-Set MMT-39724
Paradigm Quick-Set MMT 000
Paradigm Quick-Set MMT-39824
Paradigm Quick-Set MMT-39924
Paradigm Silhouette MMT-36822
Paradigm Silhouette MMT-37722
Paradigm Silhouette MMT-37822
Paradigm Silhouette MMT-38122
Paradigm Silhouette MMT-38222
Paradigm Silhouette MMT-38322
Paradigm Silhouette MMT-38422
Paradigm Sure-T MMT-86420
Paradigm Sure-T MMT-86620
Paradigm Sure-T MMT-80020
Paradigm Sure-T MMT-87620
Paradigm Sure-1 MMT-87620
Paradigm Sure-T MMT-88420
Paradigm Sure-T MMT-88620
Paraffin
Paraffin liquid with wool fat 224
Paraldehyde 126
Parasidose68
Parasiticidal Preparations
Parnate125
Paromomycin
Paroxetine
Paser
Patanol
Palarioi
Paxam
Pazopanib165
Peak flow meter219
Pedialyte - Bubblegum46
Pediasure
Pediasure RTH235
Pegaspargase159
Pegasys107
Pegfilgrastim
Pegylated interferon alfa-2a 107
Pembrolizumab
Pemetrexed
Penicillamine
Penicillin G91

PenMix 30	10
PenMix 40	10
PenMix 50	10
Pentasa	7
Pentostatin [Deoxycoformycin]	.160
Pentoxifylline [Oxpentifylline]	
Peptamen Junior	
Peptisoothe	8
Peptisorb	.237
Perhexiline maleate	
Pericyazine	.132
Perindopril	
Perjeta	. 193
Permethrin	
Perrigo	
Pertuzumab	
Peteha	
Pethidine hydrochloride	
Pevaryl	
Pexsig	
Pfizer Exemestane	169
Pharmacy Health Sorbolene with	
Glycerin	65
Pheburane	
Phenasen	
Phenelzine sulphate	125
Phenobarbitone	128
Phenobarbitone sodium	
Extemporaneous	229
Nervous	143
Phenothrin	
Phenoxybenzamine	
hydrochloride	47
Phenoxymethylpenicillin (Penicillin	
V)	92
Phenytoin sodium	128
Phlexy 10	
Phosphate Phebra	
Phosphorus	
Phytomenadione	
Pilocarpine hydrochloride	202
Pimafucort	.220 64
Pindolol	
Pine tar with trolamine laurilsulfate	50
Pine tar with trolamine laurilsulfate and fluorescein	50 69
Pine tar with trolamine laurilsulfate and fluorescein Pinetarsol	50 69 69
Pine tar with trolamine laurilsulfate and fluorescein Pinetarsol Pioglitazone	50 69 69 11
Pine tar with trolamine laurilsulfate and fluorescein Pinetarsol Pioglitazone Pirfenidone	50 69 69 11 . 217
Pine tar with trolamine laurilsulfate and fluorescein Pinetarsol Pioglitazone Pirfenidone Pizotifen	50 69 11 . 217 . 130
Pine tar with trolamine laurilsulfate and fluorescein Pinetarsol Pioglitazone Pirfenidone Pizotifen PKU Anamix Infant	50 69 11 .217 .130 .245
Pine tar with trolamine laurilsulfate and fluorescein Pinetarsol Pioglitazone Pirfenidone Pizotifen PKU Anamix Infant PKU Anamix Junior	50 69 11 .217 .130 .245 .245
Pine tar with trolamine laurilsulfate and fluorescein Pinetarsol Piglitazone Pirfenidone Pizotifen PKU Anamix Infant PKU Anamix Junior PKU Anamix Junior Chocolate	50 69 11 .217 .130 .245 .245 .245
Pine tar with trolamine laurilsulfate and fluorescein Pioglitazone Pirfenidone Pizotifen PKU Anamix Infant PKU Anamix Junior PKU Anamix Junior Chocolate PKU Anamix Junior LQ	50 69 11 .217 .130 .245 .245 .245 .245
Pine tar with trolamine laurilsulfate and fluorescein Pioglitazone Pirfenidone Pitrenidone PKU Anamix Infant PKU Anamix Junior PKU Anamix Junior Chocolate PKU Anamix Junior LQ PKU Anamix Junior Vanilla	50 69 11 .217 .130 .245 .245 .245 .245 .245
Pine tar with trolamine laurilsulfate and fluorescein Pioglitazone Pirfenidone Pizotifen PKU Anamix Infant PKU Anamix Junior PKU Anamix Junior Chocolate PKU Anamix Junior LQ	50 69 11 .217 .130 .245 .245 .245 .245 .245 .245

PKU Lophlex Powder245
PKU Lophlex Sensation 20245
Plaquenil111
Plendil ER51
Pneumococcal (PCV10) conjugate
vaccine
vaccine
Pneumococcal (PPV23)
polysaccharide vaccine
Pneumovax 23
Podophyllotoxin
Polaramine
Poliomyelitis vaccine
Poloxamer
Poly-Gel
Poly-Ger 224 Poly-Tears 224
Poly-Visc
Poly-VISC
Polycal
Polyvinyl alcohol
Ponstan
Posaconazole
Postinor-1
Potassium chloride
Potassium citrate
Potassium iodate
Povidone iodine
Pradaxa
Pramipexole hydrochloride 118
Prasugrel
Pravastatin
Praziquantel
Prazosin
Pred Forte
Prednisolone
Prednisolone acetate
Prednisolone sodium
phosphate
Prednisolone-AFT222
Prednisone
Pregabalin
Pregabalin Pfizer
Pregnancy Tests - hCG Urine74
Premarin
Presolol
Prevenar 13256
Prezista106
Priadel
Priceline121
Primacin
Primaquine phosphate
Primidone
Primolut N81
Priorix
Probenecid
Probenecid-AFT

	_
Procaine penicillin9)2
Procarbazine hydrochloride16	60
Prochlorperazine13	31
Proctosedyl	
Procyclidine hydrochloride11	8
Procytox15	
Progesterone	21
Proglicem	
	9
Proglycem	9
Progynova	
Prolia11	1
Promethazine hydrochloride21	
Propafenone hydrochloride4	9
Propamidine isethionate 22	20
Propranolol5	i 1
Propylene glycol22	29
Propylthiouracil	32
Protaphane 1	0
Protaphane Penfill1	0
Protifar	ž
Protionamide	20
Provera	20
Provera HD	
PSM Citalopram12	
	:0
Psoriasis and Eczema Preparations6	
Preparations	٥٥ م
PTU8	
Pulmicort Turbuhaler21	
Pulmocare	
Pulmozyme21	
Puri-nethol 15	5
Puria3	
Pyrazinamide9)9
Pyridostigmine bromide11	
Pyridoxine hydrochloride	33
Pyrimethamine	94
Pytazen SR4	1
- Q -	
Q 300	98
Quetapel13	32
Quetiapine 13	
Quick-Set MMT-3902	
Quick-Set MMT-3912	24
Quick-Set MMT-3922	24
Quick-Set MMT-393	 24
Quinapril4	
Quinapril with	
•	8
Quinine sulphate	
Qvar21 - R -	3
	0
RA-Morph	2
Raloxifene hydrochloride 11	
Raltegravir potassium10	
Ramipex11	8
Ranbaxy-Cefaclor8	18

Ranitidine	8
Ranitidine Relief	8
Rapamune	
Reandron 1000	79
Recombinant factor IX	8 40
Recombinant factor VIIa	
Recombinant factor VIII	0_41
Rectogesic	л – О Я
Redipred	
Relieve	
Relistor	
Remicade	
Renilon 7.5	236
Resonium-A	
Resource Beneprotein	
Resource Diabetic	000
Respigen	015
Respiratory Devices	
Respiratory Stimulants	
Retinol palmitate	
ReTrieve	
Retrovir	. 106
Revlimid	. 158
Revolade	
Rexacrom	
RexAir	214
Reyataz	. 106
Ribomustin	. 151
Ricit	
Rifabutin	
Rifadin Rifampicin	
Rifaximin	
Rifinah	
Rilutek	
Riluzole	
Riodine Risedronate Sandoz	
Risedronate sodium	. 113
Risperdal Consta	104
Risperidone133	104
Risperon	104
Ritalin	145
Ritalin LA	
Ritalin SR	
Ritonavir	
Rituximab	100
Rivaroxaban	. 194
Rivastigmine	
Rivasugmine	
RIXUBIS	
Rizamelt	
Rizatriptan	
Roferon-A	
Rolin	
Ropinirole hydrochloride	

INDEX: Ge	eneric	Chemicals	and	Brands
-----------	--------	-----------	-----	--------

Rotarix
Rotavirus oral vaccine257
Roxane
Alimentary6
Cardiovascular
Roxithromycin
Rubifen
Rubifen SR 145
Rulide D90
Rurioctocog alfa pegol [Recombinant
factor VIII] 41
Ruxolitinib
Rythmodan
Rytmonorm
- S -
-3-
Sabril
Sacubitril with valsartan48
SalAir215
Salazopyrin7
Salazopyrin EN7
Salbutamol
Salbutamol with ipratropium
bromide
Salicylic acid
Salmeterol
Sandomigran 130
Sandostatin LAR 168
Sapropterin dihydrochloride29
Scalp Preparations 69
Scopoderm TTS 130
Sebizole
Secukinumab
Sedatives and Hypnotics 143
Seebri Breezhaler
Selegiline hydrochloride 118
Senna27
Senokot27
Sensipar77
SensoCard13
Serenace
Seretide
Seretide Accuhaler
Serevent
Serevent Accuhaler
Sertraline 125
Sevredol 122
Sex Hormones Non
Contraceptive79
Shield 4971
Shield Blue
Shield XL
chingles vessing
shingles vaccine
SII-Onco-BCG
Sildenafil
Silhouette MMT-371 22
Silhouette MMT-37322

Siltuximab	202
Simvastatin	
Simvastatin Mylan	<mark>5</mark> 4
Sinemet	118
Sinemet CR	118
Sirolimus	210
Siterone	
Slow-Lopresor	
Smith BioMed Rapid Pregnancy	
Test	74
Sodibic	/ 4
Souipic	40
Sodium acid phosphate	21
Sodium alginate	6
Sodium aurothiomalate	
Sodium benzoate	30
Sodium bicarbonate	
Blood	45–46
Extemporaneous	
Sodium calcium edetate	
Sodium chloride	
Blood	45
Respiratory	
Sodium citrate with sodium lauryl	
sulphoacetate	07
Sodium citro-tartrate	21
	75
Sodium cromoglicate	_
Alimentary	
Respiratory	218
Sensory	
Sodium fluoride	34
Sodium Fusidate [fusidic acid]	
Dermatological	61
Infection	94
Sensory	220
Sodium hyaluronate [Hyaluronic	
acid]	224
Sodium phenylbutyrate	
Sodium polystyrene sulphonate	
Sodium tetradecyl sulphate	41
Sodium valproate	
Sofradex	
Soframycin	
Solian	
Solifenacin Mylan	
Solifenacin succinate	
Solu-Cortef	
Solu-Medrol	78
Solu-Medrol-Act-O-Vial	
Somatropin (Omnitrope)	
Sotalol	
Spacer device	219
Span-K	46
	216
Spiractin	53
Spiriva	
Spiriva Respimat	

Spironolactone53
Sporanox96
Sprycel162
Staphlex
Stemetil131
SteroClear
Stesolid126
Stimulants/ADHD Treatments 144
Stiripentol128
Stocrin 105
Stomahesive
Strattera144
Stromectol
Suboxone 147
Sucralfate
Sulfadiazine Silver61
Sulfadiazine sodium
Sulfasalazine
Sulindac
Sulphur
Sulprix
Sumatriptan
Sunitinib
Sunscreens
Sunscreens, proprietary 69
Sure-T MMT-863
Sure-T MMT-865
Sure-T MMT-873
Sure-T MMT-875
Sure-T MMT-873
Sure-T MMT-885
Sustagen Diabetic
Sustagen Hospital Formula
Active
Sustanon Ampoules
Sutent
Sylvant
Symbicort Turbuhaler 100/6
Symbicort Turbuhaler 200/6
Symbicort Turbuhaler 200/0
Symmetrel
Sympathomimetics
Synacthen
Synacthen Depot
Synacthen S29
Synacthene Retard
Synflorix
Synthroid
Syntometrine
Syrup (pharmaceutical grade)
Systane Unit Dose224
Tacrolimus
Tacrolimus Sandoz
Tambooor
Tambocor49

Tambocor CR
Tamoxifen citrate 169
Tamoxifen Sandoz169
Tamsulosin hydrochloride75
Tamsulosin-Rex75
Tandem Cartridge19
Tandem t:slim X214
Tap water
Tarceva162
Tasigna164
Tasmar
Tecfidera135
Tegretol
Tegretol CR 126
Telfast
Temazepam143
Temizole 20 160
Temozolomide 160
Tenofovir disoproxil 101
Tenofovir Disoproxil Teva101
Tenoxicam
Tepadina153
Terazosin47
Terbinafine97
Terbutaline sulphate215
Teriflunomide139
Teriparatide 113
Testosterone79
Testosterone cipionate79
Testosterone esters
Testosterone undecanoate79
Tetrabenazine119
Tetrabromophenol76
Tetracosactrin79
Tetracyclin Wolff92
Tetracycline
Thalidomide161
Thalomid161
Theophylline
Thiamine hydrochloride
THIO-TEPA 153
Thioguanine156
Thiotepa153
Thymol glycerin
Thyroid and Antithyroid Agents82
Ticagrelor
Tilade
Tilcotil
Timolol
Cardiovascular
Sensory
Timoptol XE
Tiotropium bromide
Tiotropium bromide with
olodaterol
Tivicay
,

TMP	05
TOBI	95
Tobramycin	
Infection	
Sensory	
Tobramycin Mylan	95
Tobrex	221
Tocilizumab	203
Tofranil	
Tolcapone	
Tolterodine	
Topamax	129
Topical Products for Joint and	
Muscular Pain	
Topiramate	
Topiramate Actavis	129
Total parenteral nutrition (TPN)	45
TPN	
Tramadol hydrochloride	124
Tramal SR 100	
Tramal SR 150	
Tramal SR 200	
Trandate	
Tranexamic acid	
Tranylcypromine sulphate	
Trastuzumab	206
Travatan	
Travoprost	
Travopt	
Treatments for Dementia	
	147
Treatments for Substance	
Dependence	. 147
Trental 400	56
Tretinoin	
Dermatological	60
Oncology	
Trexate	
Triamcinolone acetonide	
Alimentary	30
Dermatological	02 64
Dermatological	04
Hormone	79
Triamcinolone acetonide with	
gramicidin, neomycin and nystat	in
Dermatological	64
Sensory	
Triazolam	. 144
Trichozole	98
	64
Trimethoprim	
	50
Trimethoprim with	
sulphamethoxazole	
[Co-trimoxazole]	95
Trisequens	
Trisul	
Trophic Hormones	
Tropicamide	
1	

Trusopt	222
TruSteel	
Truvada	103
Tuberculin PPD [Mantoux] test	258
Tubersol	
Two Cal HN	
Two Cal HN RTH	
Tykerb	
Tysabri	138
- U -	
Ultibro Breezhaler	
Ultraproct	
Umeclidinium	216
Umeclidinium with vilanterol	216
Univent215,	219
Ural	
Urea	
Urex Forte	
Urinary Agents	
Urinary Tract Infections	100
Uromitexan	
Ursodeoxycholic acid	. 159
Ursosan	
Utrogestan	81
- V -	
Vaccinations	
Vaclovir	
Valaciclovir	
Valcyte	.101
Valganciclovir	101
Valganciclovir Mylan	.101
Vancomycin	95
Vannair	
Varenicline Pfizer	
Varenicline tartrate	
Varicella vaccine [Chickenpox	. 140
vaccine]	250
Varicella zoster virus (Oka strain) liv	200
attenuated vaccine [shingles	e
	050
vaccine]	
Varilrix	
Vasodilators	
Vasopressin Agonists	
Vedafil	
Velcade	
Veletri	58
Venlafaxine	126
Venomil	212
Ventavis	
Ventolin	
Vepesid	
Verapamil hydrochloride	52
Vergo 16	
Vermox	
Verpamil SR	
Vesanoid	. 161

Vexazone11
Vfend
Viaderm KC64
Vidaza
Vigabatrin 129
Vildagliptin11
Vildagliptin with metformin hydrochloride11
hydrochloride 11
Vimpat 127
Vinblastine sulphate161
Vincristine sulphate 161
Vinorelbine
Vinorelbine Ebewe162
Viramune Suspension 105
ViruPOS
Vistil
Vistil Forte
Vit.D3
VitA-POS
Vitabdeck
Vitadol C33
Vital
Vitamin A with vitamins D and C 33
Vitamin B complex
Vitamin B6 25
Vitamins
Vivonex TEN
Volibris
Voltaren 110
Voltaren D110
Voltaren Ophtha
Volumatic
Voriconazole
Vosol
Votrient
Vttack
- W -
Warfarin sodium
Wart Preparations
Wasp venom allergy treatment
Wator
Blood
Extemporaneous229
Wool fat with mineral oil
- X -
Xarelto
Xifaxan
XMET Maxamum
Xolair
XP Maxamum
Xylocaine
Xylocaine 2% Jelly
Xyntha
- Z -
Zantac8
Zapril

Zarantin 10	0
Zarontin	
Zaroxolyn	
Zarzio4	
Zavedos15	
Zeffix10	
Zetlam10	
Ziagen10	
Zidovudine [AZT] 10	6
Zidovudine [AZT] with	
lamivudine 10	
Zimybe5	
Zinc and castor oil6	
Zinc sulphate3	6
Zincaps3	6
Zinnat8	8
Ziprasidone13	3
Zista21	3
Zithromax8	9
Zoladex8	6
Zoledronic acid	
Hormone7	7
Musculoskeletal 11	4
Zoledronic acid Mylan7	
Zometa7	
Zopiclone14	
Zopiclone Actavis14	
Zostavax25	
Zostrix11	
Zostrix HP12	
Zuclopenthixol decanoate13	
Zuclopenthixol hydrochloride	
Zusdone13	
 Zyban14	
Zypine13	
Zypine ODT13	
Zyprexa Relprevv13	
-jpiona i loipiott	U
Zytiga16	















