

**October 2019**  
**Volume 26 Number 2**

**Editors:**

Kaye Wilson & Doris Chong  
email: [enquiry@pharmac.govt.nz](mailto:enquiry@pharmac.govt.nz)  
Telephone +64 4 460 4990  
Facsimile +64 4 460 4995  
Level 9, 40 Mercer Street  
PO Box 10 254 Wellington

**Freephone Information Line**

**0800 66 00 50 (9am – 5pm weekdays)**

**Circulation**

Published each April, August and December.  
Changes to the contents are published in  
monthly updates.

Accessible in an electronic format at  
no cost from the PHARMAC website  
[www.pharmac.govt.nz](http://www.pharmac.govt.nz).

You can register to have an electronic  
version of the Pharmaceutical Schedule (link  
to PDF copy) emailed to your nominated  
email address each month by subscribing at  
[www.pharmac.govt.nz/subscriptions](http://www.pharmac.govt.nz/subscriptions).

**Production**

Typeset automatically from XML and T<sub>E</sub>X.  
XML version of the Schedule available from  
[www.pharmac.govt.nz/pub/schedule/archive/](http://www.pharmac.govt.nz/pub/schedule/archive/)

**Programmers**

Anrik Drenth & John Geering

email: [texschedule@pharmac.govt.nz](mailto:texschedule@pharmac.govt.nz)

©Pharmaceutical Management Agency



ISSN 1179-3686 pdf  
ISSN 1172-9376 print

This work is licensed under the Creative  
Commons Attribution 4.0 International licence.  
In essence, you are free to copy, distribute  
and adapt it, as long as you attribute the work  
to PHARMAC and abide by the other licence  
terms. To view a copy of this licence, visit:  
[creativecommons.org/licenses/by/4.0/](http://creativecommons.org/licenses/by/4.0/).

Attribution to PHARMAC should be in  
written form and not by reproduction of the  
PHARMAC logo. While care has been taken  
in compiling this Schedule, PHARMAC takes  
no responsibility for any errors or omissions,  
and shall not be liable for any consequences  
arising there from.

<b>Section A</b>	General Rules	<b>5</b>
------------------	---------------	----------

<b>Section B</b>	Alimentary Tract & Metabolism	<b>6</b>
	Blood & Blood Forming Organs	<b>35</b>
	Cardiovascular System	<b>45</b>
	Dermatologicals	<b>59</b>
	Genito Urinary System	<b>70</b>
	Hormone Preparations – Systemic	<b>78</b>
	Infections – Agents For Systemic Use	<b>89</b>
	Musculoskeletal System	<b>110</b>
	Nervous System	<b>118</b>
	Oncology Agents & Immunosuppressants	<b>155</b>
	Respiratory System & Allergies	<b>220</b>
	Sensory Organs	<b>228</b>
	Various	<b>233</b>

<b>Section C</b>	Extemporaneous Compounds (ECPs)	<b>235</b>
------------------	---------------------------------	------------

<b>Section D</b>	Special Foods	<b>238</b>
------------------	---------------	------------

<b>Section I</b>	National Immunisation Schedule	<b>257</b>
------------------	--------------------------------	------------

Index	<b>267</b>
-------	------------

# Introducing PHARMAC

The Pharmaceutical Management Agency (PHARMAC) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. PHARMAC negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

## PHARMAC's role:

***“Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided.”***

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about PHARMAC and the way we make funding decisions can be found on the PHARMAC website at <http://www.pharmac.govt.nz/about>.

## Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by PHARMAC.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements PHARMAC has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that can be used in DHB hospitals and is a separate publication.

The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

# Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

## Example

ANATOMICAL HEADING			
THERAPEUTIC HEADING			
CHEMICAL			
Presentation, form and strength			
10.00	100	Brand A	Brand B
Presentation - Available on a PSO	15.00	50	Brand C
Presentation - Retail pharmacy-specialist	18.00	250 ml OP	Brand D
a) Prescriptions must be written by a paediatrician or paediatric cardiologist; or b) on the recommendation of a paediatrician or a paediatric cardiologist			
CHEMICAL			
Presentation, form and strength			
26.53	100	Brand E	
(35.27)			
Sole Supply			
Fully Subsidised			
Three months supply may be dispensed at one time if endorsed 'certified exemption' by the prescriber or pharmacist.			

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Practitioner's Supply Order

Conditions of and restrictions on prescribing (including Special Authority where it applies)

Three months or six months, as applicable, dispensed all-at-once

Brand or manufacturer's name

Sole subsidised supply product

Fully subsidised product

Original Pack - Subsidy is rounded up to a multiple of whole packs

Quantity the Subsidy applies to

Subsidy paid on a product before mark-ups and GST

Manufacturer's Price if different from Subsidy

# Glossary

## Units of Measure

gram .....	g	microgram.....	mcg	millimole.....	mmol
kilogram.....	kg	milligram .....	mg	unit.....	u
international unit.....	iu	millilitre.....	ml		

## Abbreviations

Ampoule .....	Amp	Gelatinous .....	Gel	Solution.....	Soln
Capsule .....	Cap	Granules .....	Gran	Suppository.....	Supp
Cream.....	Crm	Infusion .....	Inf	Tablet.....	Tab
Device.....	Dev	Injection .....	Inj	Tincture.....	Tinc
Dispersible.....	Disp	Liquid.....	Liq	Trans Dermal Delivery	
Effervescent.....	Eff	Long Acting.....	LA	System.....	TDDS
Emulsion.....	Emul	Ointment.....	Oint		
Enteric Coated.....	EC	Sachet .....	Sach		

General Rules for the Pharmaceutical Schedule are located on the PHARMAC website.

## SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

### Antacids and Antiflatulents

#### Antacids and Reflux Barrier Agents

##### ALGINIC ACID

Sodium alginate 225 mg and magnesium alginate 87.5 mg per sachet.....	5.31	30	✓	Gaviscon Infant
---	------	----	---	-----------------

##### SODIUM ALGINATE

* Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour.....	1.80 (8.60)	60		Gaviscon Double Strength
* Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml.....	1.50 (4.95)	500 ml		Acidex

#### Phosphate Binding Agents

##### ALUMINIUM HYDROXIDE

* Tab 600 mg .....	12.56	100	✓	Alu-Tab
--------------------	-------	-----	---	---------

##### CALCIUM CARBONATE

Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsement .....	39.00	500 ml	✓	Roxane
Only when prescribed for patients unable to swallow calcium carbonate tablets or where calcium carbonate tablets are inappropriate and the prescription is endorsed accordingly.				

### Antidiarrhoeals

#### Agents Which Reduce Motility

##### LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a PSO

* Tab 2 mg .....	10.75	400	✓	Nodia
* Cap 2 mg .....	6.25	400	✓	Diamide Relief

#### Rectal and Colonic Anti-inflammatories

##### BUDESONIDE

Cap 3 mg – Special Authority see <a href="#">SA1155 below</a> – Retail pharmacy.....	166.50	90	✓	Entocort CIR
--	--------	----	---	--------------

##### ► [SA1155](#) Special Authority for Subsidy

**Initial application — (Crohn's disease)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
  - 2.1 Diabetes; or
  - 2.2 Cushingoid habitus; or
  - 2.3 Osteoporosis where there is significant risk of fracture; or

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

**Initial application — (collagenous and lymphocytic colitis (microscopic colitis))** from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

**Initial application — (gut Graft versus Host disease)** from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation\*.

Note: Indication marked with \* is an unapproved indication.

**Renewal** from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

#### HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications).....	26.55	21.1 g OP	✓ Colifoam
--	-------	-----------	------------

#### HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE

Topical aerosol foam, 1% with pramoxine hydrochloride 1%.....	26.55	10 g OP	✓ Proctofoam S29
---	-------	---------	------------------

#### MESALAZINE

Tab 400 mg .....	49.50	100	✓ Asacol
Tab EC 500 mg .....	49.50	100	✓ Asamax
Tab long-acting 500 mg.....	59.05	100	✓ Pentasa
Tab 800 mg .....	85.50	90	✓ Asacol
Modified release granules, 1 g .....	141.72	120 OP	✓ Pentasa
Enema 1 g per 100 ml .....	41.30	7	✓ Pentasa
Suppos 500 mg .....	22.80	20	✓ Asacol
Suppos 1 g .....	54.60	30	✓ Pentasa

#### OLSALAZINE

Tab 500 mg .....	93.37	100	✓ Dipentum
Cap 250 mg .....	53.00	100	✓ Dipentum

#### SODIUM CROMOGLICATE

Cap 100 mg .....	92.91	100	✓ Nalcrom
------------------	-------	-----	-----------

#### SULFASALAZINE

* Tab 500 mg .....	14.00	100	✓ Salazopyrin
* Tab EC 500 mg .....	15.53	100	✓ Salazopyrin EN

Salazopyrin EN to be Sole Supply on 1 December 2019

## Local preparations for Anal and Rectal Disorders

### Antihæmorrhoidal Preparations

#### FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE

Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cinchocaine hydrochloride 5 mg per g.....	6.35	30 g OP	✓ Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and cinchocaine hydrochloride 1 mg .....	2.66	12	✓ Ultraproct

#### HYDROCORTISONE WITH CINCHOCAINE

Oint 5 mg with cinchocaine hydrochloride 5 mg per g.....	15.00	30 g OP	✓ Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g .....	9.90	12	✓ Proctosedyl

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

## Management of Anal Fissures

GLYCERYL TRINITRATE – Special Authority see [SA1329 below](#) – Retail pharmacy

\* Oint 0.2% ..... 22.00 30 g OP ✓ **Rectogesic**

► [SA1329](#) Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

## Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE

Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a  
PSO ..... 17.14 10 ✓ **Max Health**

HYOSCINE BUTYLBROMIDE

\* Tab 10 mg ..... 8.75 100 ✓ **Buscopan**

\* Inj 20 mg, 1 ml – Up to 5 inj available on a PSO ..... 9.57 5 ✓ **Buscopan**

MEBEVERINE HYDROCHLORIDE

\* Tab 135 mg ..... 18.00 90 ✓ **Colofac**

## Antiulcerants

### Antisecretory and Cytoprotective

MISOPROSTOL

\* Tab 200 mcg ..... 41.50 120 ✓ **Cytotec**

### Helicobacter Pylori Eradication

CLARITHROMYCIN

Tab 500 mg – Subsidy by endorsement ..... 10.40 14 ✓ **Apo-Clarithromycin**

a) Maximum of 14 tab per prescription

b) Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.

Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

## H2 Antagonists

RANITIDINE – Only on a prescription

\* Tab 150 mg ..... 12.91 500 ✓ **Ranitidine Relief**

\* Tab 300 mg ..... 18.21 500 ✓ **Ranitidine Relief**

\* Oral liq 150 mg per 10 ml ..... 5.14 300 ml ✓ **Peptisoothe**

\* Inj 25 mg per ml, 2 ml ..... 13.40 5 ✓ **Zantac**

## Proton Pump Inhibitors

LANSOPRAZOLE

\* Cap 15 mg ..... 4.58 100 ✓ **Lanzol Relief**

\* Cap 30 mg ..... 5.41 100 ✓ **Lanzol Relief**



	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>OMEPRAZOLE</b>				
For omeprazole suspension refer Standard Formulae, <a href="#">page 235</a>				
* Cap 10 mg.....	1.98	90	✓	<b>Omeprazole actavis 10</b>
* Cap 20 mg.....	1.96	90	✓	<b>Omeprazole actavis 20</b>
* Cap 40 mg.....	3.12	90	✓	<b>Omeprazole actavis 40</b>
* Powder – Only in combination.....	42.50	5 g	✓	<b>Midwest</b>
Only in extemporaneously compounded omeprazole suspension.				
* Inj 40 mg ampoule with diluent.....	33.98	5	✓	<b>Dr Reddy's Omeprazole</b>
<b>PANTOPRAZOLE</b>				
* Tab EC 20 mg.....	2.02	100	✓	<b>Panzop Relief</b>
* Tab EC 40 mg.....	2.85	100	✓	<b>Panzop Relief</b>

### Site Protective Agents

<b>COLLOIDAL BISMUTH SUBCITRATE</b>				
Tab 120 mg.....	14.51	50	✓	<b>Gastrodenol</b> <small>\$29</small>
<b>SUCRALFATE</b>				
Tab 1 g.....	35.50 (48.28)	120		Carafate

### Bile and Liver Therapy

<b>RIFAXIMIN</b> – Special Authority see <a href="#">SA1461 below</a> – Retail pharmacy				
Tab 550 mg.....	625.00	56	✓	<b>Xifaxan</b>

#### ►SA1461 Special Authority for Subsidy

**Initial application** only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid for 6 months where the patient has hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

**Renewal** only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

### Diabetes

#### Hyperglycaemic Agents

<b>DIAZOXIDE</b> – Special Authority see <a href="#">SA1320 below</a> – Retail pharmacy				
Cap 25 mg.....	110.00	100	✓	<b>Proglycem</b> <small>\$29</small>
Cap 100 mg.....	280.00	100	✓	<b>Proglycem</b> <small>\$29</small>
Oral liq 50 mg per ml.....	620.00	30 ml OP	✓	<b>Proglycem</b> <small>\$29</small>

#### ►SA1320 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months where used for the treatment of confirmed hypoglycaemia caused by hyperinsulinism.

**Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

<b>GLUCAGON HYDROCHLORIDE</b>				
Inj 1 mg syringe kit – Up to 5 kit available on a PSO.....	32.00	1	✓	<b>Glucagen Hypokit</b>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>Insulin - Short-acting Preparations</b>				
INSULIN NEUTRAL				
▲ Inj human 100 u per ml.....	25.26	10 ml OP	✓	Actrapid
			✓	Humulin R
▲ Inj human 100 u per ml, 3 ml.....	42.66	5	✓	Actrapid Penfill
			✓	Humulin R
<b>Insulin - Intermediate-acting Preparations</b>				
INSULIN ASPART WITH INSULIN ASPART PROTAMINE				
▲ Inj 100 iu per ml, 3 ml prefilled pen.....	52.15	5	✓	NovoMix 30 FlexPen
INSULIN ISOPHANE				
▲ Inj human 100 u per ml.....	17.68	10 ml OP	✓	Humulin NPH
			✓	Protaphane
▲ Inj human 100 u per ml, 3 ml.....	29.86	5	✓	Humulin NPH
			✓	Protaphane Penfill
INSULIN ISOPHANE WITH INSULIN NEUTRAL				
▲ Inj human with neutral insulin 100 u per ml.....	25.26	10 ml OP	✓	Humulin 30/70
			✓	Mixtard 30
▲ Inj human with neutral insulin 100 u per ml, 3 ml.....	42.66	5	✓	Humulin 30/70
			✓	PenMix 30
			✓	PenMix 40
			✓	PenMix 50
INSULIN LISPRO WITH INSULIN LISPRO PROTAMINE				
▲ Inj lispro 25% with insulin lispro protamine 75% 100 u per ml, 3 ml.....	42.66	5	✓	Humalog Mix 25
▲ Inj lispro 50% with insulin lispro protamine 50% 100 u per ml, 3 ml.....	42.66	5	✓	Humalog Mix 50
<b>Insulin - Long-acting Preparations</b>				
INSULIN GLARGINE				
▲ Inj 100 u per ml, 10 ml.....	63.00	1	✓	Lantus
▲ Inj 100 u per ml, 3 ml.....	94.50	5	✓	Lantus
▲ Inj 100 u per ml, 3 ml disposable pen.....	94.50	5	✓	Lantus SoloStar
<b>Insulin - Rapid Acting Preparations</b>				
INSULIN ASPART				
▲ Inj 100 u per ml, 10 ml.....	30.03	1	✓	NovoRapid
▲ Inj 100 u per ml, 3 ml.....	51.19	5	✓	NovoRapid Penfill
▲ Inj 100 u per ml, 3 ml syringe.....	51.19	5	✓	NovoRapid FlexPen
INSULIN GLULISINE				
▲ Inj 100 u per ml, 10 ml.....	27.03	1	✓	Apidra
▲ Inj 100 u per ml, 3 ml.....	46.07	5	✓	Apidra
▲ Inj 100 u per ml, 3 ml disposable pen.....	46.07	5	✓	Apidra SoloStar
INSULIN LISPRO				
▲ Inj 100 u per ml, 10 ml.....	34.92	10 ml OP	✓	Humalog
▲ Inj 100 u per ml, 3 ml.....	59.52	5	✓	Humalog

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>Alpha Glucosidase Inhibitors</b>				
ACARBOSE				
* Tab 50 mg .....	3.50	90	✓	<u>Glucobay</u>
	10.47		✓	<u>Accarb</u>
* Tab 100 mg .....	6.40	90	✓	<u>Glucobay</u>
	11.24	50	✓	<u>Acarbose Mylan</u> <small>\$29</small>
	20.23	90	✓	<u>Accarb</u>

(Acarbose Mylan \$29 Tab 100 mg to be delisted 1 January 2020)

<b>Oral Hypoglycaemic Agents</b>				
GLIBENCLAMIDE				
* Tab 5 mg .....	6.00	100	✓	<u>Daonil</u>
GLICLAZIDE				
* Tab 80 mg .....	10.29	500	✓	<u>Glizide</u>
GLIPIZIDE				
* Tab 5 mg .....	3.27	100	✓	<u>Minidiab</u>
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg .....	8.63	1,000	✓	<u>Apotex</u>
* Tab immediate-release 850 mg .....	7.04	500	✓	<u>Apotex</u>
PIOGLITAZONE				
* Tab 15 mg .....	3.47	90	✓	<u>Vexazone</u>
* Tab 30 mg .....	5.06	90	✓	<u>Vexazone</u>
* Tab 45 mg .....	7.10	90	✓	<u>Vexazone</u>
VILDAGLIPTIN				
Tab 50 mg .....	40.00	60	✓	<u>Galvus</u>
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride .....	40.00	60	✓	<u>Galvumet</u>
Tab 50 mg with 850 mg metformin hydrochloride .....	40.00	60	✓	<u>Galvumet</u>

## Diabetes Management

### Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP – Subsidy by endorsement

- Not on a BSO
- Maximum of 20 strip per prescription
- Up to 10 strip available on a PSO
- Patient has any of the following:
  - type 1 diabetes; or
  - permanent neonatal diabetes; or
  - undergone a pancreatectomy; or
  - cystic fibrosis-related diabetes; or
  - metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Test strips ..... 15.50      10 strip OP      ✓ KetoSens

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

## Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER – Subsidy by endorsement

- Maximum of 1 pack per prescription
- Up to 1 pack available on a PSO
- A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
  - type 1 diabetes; or
  - permanent neonatal diabetes; or
  - undergone a pancreatectomy; or
  - cystic fibrosis-related diabetes; or
  - metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

diagnostic test strips.....	20.00	1 OP	✓ <u>CareSens Dual</u>
-----------------------------	-------	------	------------------------

## Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER – Subsidy by endorsement

- Maximum of 1 pack per prescription
- Up to 1 pack available on a PSO
- A diagnostic blood glucose test meter is subsidised for a patient who:
  - is receiving insulin or sulphonylurea therapy; or
  - is pregnant with diabetes; or
  - is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
  - has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- type 1 diabetes; or
- permanent neonatal diabetes; or
- undergone a pancreatectomy; or
- cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 diagnostic test

strips.....	10.00	1 OP	✓ <u>CareSens N</u>
			✓ <u>CareSens N POP</u>
	20.00		✓ <u>CareSens N Premier</u>

Note: Only 1 meter available per PSO

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

**BLOOD GLUCOSE DIAGNOSTIC TEST STRIP – Up to 50 test available on a PSO**

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- 2) Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips .....	10.56	50 test OP	✓ <b>CareSens N</b> ✓ <b>CareSens PRO</b>
-------------------	-------	------------	--

**BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)**

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- 2) Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips.....	26.20	50 test OP	✓ <b>SensoCard</b>
--------------------------------	-------	------------	--------------------

**Insulin Syringes and Needles**

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

**INSULIN PEN NEEDLES – Maximum of 200 dev per prescription**

* 29 g x 12.7 mm .....	10.50	100	✓ <b>B-D Micro-Fine</b>
* 31 g x 5 mm .....	11.75	100	✓ <b>B-D Micro-Fine</b>
* 31 g x 6 mm .....	9.50	100	✓ <b>Berpu</b>
* 31 g x 8 mm .....	10.50	100	✓ <b>B-D Micro-Fine</b>
* 32 g x 4 mm .....	10.50	100	✓ <b>B-D Micro-Fine</b>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
INSULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE – Maximum of 200 dev per prescription				
* Syringe 0.3 ml with 29 g x 12.7 mm needle .....	13.00	100	✓	<b>B-D Ultra Fine</b>
	1.30	10		
	(1.99)			B-D Ultra Fine
* Syringe 0.3 ml with 31 g x 8 mm needle .....	13.00	100	✓	<b>B-D Ultra Fine II</b>
	1.30	10		
	(1.99)			B-D Ultra Fine II
* Syringe 0.5 ml with 29 g x 12.7 mm needle .....	13.00	100	✓	<b>B-D Ultra Fine</b>
	1.30	10		
	(1.99)			B-D Ultra Fine
* Syringe 0.5 ml with 31 g x 8 mm needle .....	13.00	100	✓	<b>B-D Ultra Fine II</b>
	1.30	10		
	(1.99)			B-D Ultra Fine II
* Syringe 1 ml with 29 g x 12.7 mm needle .....	13.00	100	✓	<b>B-D Ultra Fine</b>
	1.30	10		
	(1.99)			B-D Ultra Fine
* Syringe 1 ml with 31 g x 8 mm needle .....	13.00	100	✓	<b>B-D Ultra Fine II</b>
	1.30	10		
	(1.99)			B-D Ultra Fine II

## Insulin Pumps

INSULIN PUMP – Special Authority see [SA1603 below](#) – Retail pharmacy

- Maximum of 1 dev per prescription
- Only on a prescription
- Maximum of 1 insulin pump per patient each four year period.

Min basal rate 0.025 U/h .....	8,800.00	1	✓	<b>MiniMed 640G</b>
Min basal rate 0.1 U/h .....	4,500.00	1	✓	<b>Tandem t:slim X2</b>

### ➔SA1603 Special Authority for Subsidy

**Initial application — (permanent neonatal diabetes)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- Patient has permanent neonatal diabetes; and
- A MDI regimen trial is inappropriate; and
- Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- Either:
  - Applicant is a relevant specialist; or
  - Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (permanent neonatal diabetes)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- Either:

continued...

Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic Manufacturer
\$	Per	✓

continued...

- 4.1 Applicant is a relevant specialist; or
- 4.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application — (severe unexplained hypoglycaemia)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
  - 8.1 Applicant is a relevant specialist; or
  - 8.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (severe unexplained hypoglycaemia)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
  - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 3.2 The pump is due for replacement; and
- 4 Either:
  - 4.1 Applicant is a relevant specialist; or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application — (HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1c; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

**Renewal — (HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
  - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 3.2 The pump is due for replacement; and
- 4 Either:
  - 4.1 Applicant is a relevant specialist; or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application — (Previous use before 1 September 2012)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
  - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 8.2 The pump is due for replacement; and
- 9 Either:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (Previous use before 1 September 2012)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
  - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 4.2 The pump is due for replacement; and
- 5 Either:
  - 5.1 Applicant is a relevant specialist; or
  - 5.2 Applicant is a nurse practitioner working within their vocational scope.



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## Insulin Pump Consumables

### ►SA1604 Special Authority for Subsidy

**Initial application — (permanent neonatal diabetes)** only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
  - 6.1 Applicant is a relevant specialist; or
  - 6.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (permanent neonatal diabetes)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Either:
  - 3.1 Applicant is a relevant specialist; or
  - 3.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application — (severe unexplained hypoglycaemia)** only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
  - 8.1 Applicant is a relevant specialist; or
  - 8.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (severe unexplained hypoglycaemia)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
  - 3.1 Applicant is a relevant specialist; or

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

3.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application — (HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1c; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
  - 3.1 Applicant is a relevant specialist; or
  - 3.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application — (Previous use before 1 September 2012)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
  - 8.1 Applicant is a relevant specialist; or
  - 8.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (Previous use before 1 September 2012)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
continued...			
than 80 mmol/mol; and			
2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and			
3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and			
4 Either:			
4.1 Applicant is a relevant specialist; or			
4.2 Applicant is a nurse practitioner working within their vocational scope.			
INSULIN PUMP CARTRIDGE – Special Authority see <a href="#">SA1604 on page 17</a> – Retail pharmacy			
a) Maximum of 3 sets per prescription			
b) Only on a prescription			
c) Maximum of 13 packs of cartridge sets will be funded per year.			
Cartridge 300 U, t:lock x 10.....	50.00	1 OP	✓ Tandem Cartridge
INSULIN PUMP INFUSION SET (STEEL CANNULA) – Special Authority see <a href="#">SA1604 on page 17</a> – Retail pharmacy			
a) Maximum of 3 sets per prescription			
b) Only on a prescription			
c) Maximum of 13 infusion sets will be funded per year.			
10 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles.....	130.00	1 OP	✓ Paradigm Sure-T MMT-884
10 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles; luer lock .....	130.00	1 OP	✓ Sure-T MMT-883
10 mm steel needle; 29 G; manual insertion; 80 cm tubing x 10 with 10 needles.....	130.00	1 OP	✓ Paradigm Sure-T MMT-886
10 mm steel needle; 29 G; manual insertion; 80 cm tubing x 10 with 10 needles; luer lock .....	130.00	1 OP	✓ Sure-T MMT-885
6 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles.....	130.00	1 OP	✓ Paradigm Sure-T MMT-864
6 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles; luer lock .....	130.00	1 OP	✓ Sure-T MMT-863
6 mm steel needle; 29 G; manual insertion; 80 cm tubing x 10 with 10 needles.....	130.00	1 OP	✓ Paradigm Sure-T MMT-866
6 mm steel needle; 29 G; manual insertion; 80 cm tubing x 10 with 10 needles; luer lock .....	130.00	1 OP	✓ Sure-T MMT-865
8 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles.....	130.00	1 OP	✓ Paradigm Sure-T MMT-874
8 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles; luer lock .....	130.00	1 OP	✓ Sure-T MMT-873
8 mm steel needle; 29 G; manual insertion; 80 cm tubing x 10 with 10 needles.....	130.00	1 OP	✓ Paradigm Sure-T MMT-876
8 mm steel needle; 29 G; manual insertion; 80 cm tubing x 10 with 10 needles; luer lock .....	130.00	1 OP	✓ Sure-T MMT-875

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) – Special Authority see <a href="#">SA1604 on page 17</a> – Retail pharmacy				
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles.....	130.00	1 OP	✓	TruSteel
6 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles.....	130.00	1 OP	✓	TruSteel
8 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles.....	130.00	1 OP	✓	TruSteel
8 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles.....	130.00	1 OP	✓	TruSteel
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION WITH INSERTION DEVICE) – Special Authority see <a href="#">SA1604 on page 17</a> – Retail pharmacy				
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
13 mm teflon cannula; angle insertion; insertion device; 110 cm line x 10 with 10 needles.....	140.00	1 OP	✓	AutoSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cm line x 10 with 10 needles.....	140.00	1 OP	✓	AutoSoft 30

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION) – Special Authority see <a href="#">SA1604 on page 17</a> – Retail pharmacy			
a) Maximum of 3 sets per prescription			
b) Only on a prescription			
c) Maximum of 13 infusion sets will be funded per year.			
13 mm teflon cannula; angle insertion; 120 cm line x 10 with 10 needles.....	130.00	1 OP	✓ <b>Paradigm Silhouette MMT-382</b>
13 mm teflon cannula; angle insertion; 45 cm line x 10 with 10 needles.....	130.00	1 OP	✓ <b>Paradigm Silhouette MMT-368</b>
13 mm teflon cannula; angle insertion; 60 cm line x 10 with 10 needles.....	130.00	1 OP	✓ <b>Paradigm Silhouette MMT-381</b>
13 mm teflon cannula; angle insertion; 80 cm line x 10 with 10 needles.....	130.00	1 OP	✓ <b>Paradigm Silhouette MMT-383</b>
17 mm teflon cannula; angle insertion; 110 cm line x 10 with 10 needles.....	130.00	1 OP	✓ <b>Paradigm Silhouette MMT-377</b>
17 mm teflon cannula; angle insertion; 110 cm line x 10 with 10 needles; luer lock.....	130.00	1 OP	✓ <b>Silhouette MMT-371</b>
17 mm teflon cannula; angle insertion; 60 cm line x 10 with 10 needles.....	130.00	1 OP	✓ <b>Paradigm Silhouette MMT-378</b>
17 mm teflon cannula; angle insertion; 60 cm line x 10 with 10 needles; luer lock.....	130.00	1 OP	✓ <b>Silhouette MMT-373</b>
17 mm teflon cannula; angle insertion; 80 cm line x 10 with 10 needles.....	130.00	1 OP	✓ <b>Paradigm Silhouette MMT-384</b>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION WITH INSERTION DEVICE) – Special Authority see <a href="#">SA1604 on page 17</a> – Retail pharmacy				
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; insertion device; 45 cm blue tubing x 10 with 10 needles.....	130.00	1 OP	✓	Paradigm Mio MMT-941
6 mm teflon cannula; straight insertion; insertion device; 45 cm pink tubing x 10 with 10 needles.....	130.00	1 OP	✓	Paradigm Mio MMT-921
6 mm teflon cannula; straight insertion; insertion device; 60 cm blue tubing x 10 with 10 needles.....	130.00	1 OP	✓	Paradigm Mio MMT-943
6 mm teflon cannula; straight insertion; insertion device; 60 cm pink tubing x 10 with 10 needles.....	130.00	1 OP	✓	Paradigm Mio MMT-923
6 mm teflon cannula; straight insertion; insertion device; 80 cm blue tubing x 10 with 10 needles.....	130.00	1 OP	✓	Paradigm Mio MMT-945
6 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing x 10 with 10 needles.....	130.00	1 OP	✓	Paradigm Mio MMT-965
6 mm teflon cannula; straight insertion; insertion device; 80 cm pink tubing x 10 with 10 needles.....	130.00	1 OP	✓	Paradigm Mio MMT-925
9 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing x 10 with 10 needles.....	130.00	1 OP	✓	Paradigm Mio MMT-975
6 mm teflon cannula; straight insertion; insertion device; 110 cm line x 10 with 10 needles.....	140.00	1 OP	✓	AutoSoft 90
6 mm teflon cannula; straight insertion; insertion device; 60 cm line x 10 with 10 needles.....	140.00	1 OP	✓	AutoSoft 90
9 mm teflon cannula; straight insertion; insertion device; 110 cm line x 10 with 10 needles.....	140.00	1 OP	✓	AutoSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 cm line x 10 with 10 needles.....	140.00	1 OP	✓	AutoSoft 90

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION) – Special Authority see <a href="#">SA1604 on page 17</a> – Retail pharmacy			
a) Maximum of 3 sets per prescription			
b) Only on a prescription			
c) Maximum of 13 infusion sets will be funded per year.			
6 mm teflon cannula; straight insertion; 110 cm tubing × 10 with 10 needles.....	130.00	1 OP	✓ <b>Paradigm Quick-Set MMT-398</b>
6 mm teflon cannula; straight insertion; 110 cm tubing × 10 with 10 needles; luer lock.....	130.00	1 OP	✓ <b>Quick-Set MMT-391</b>
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 10 needles.....	130.00	1 OP	✓ <b>Paradigm Quick-Set MMT-399</b>
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 10 needles; luer lock.....	130.00	1 OP	✓ <b>Quick-Set MMT-393</b>
6 mm teflon cannula; straight insertion; 80 cm tubing × 10 with 10 needles.....	130.00	1 OP	✓ <b>Paradigm Quick-Set MMT-387</b>
9 mm teflon cannula; straight insertion; 106 cm tubing × 10 with 10 needles.....	130.00	1 OP	✓ <b>Paradigm Quick-Set MMT-396</b>
9 mm teflon cannula; straight insertion; 110 cm tubing × 10 with 10 needles; luer lock.....	130.00	1 OP	✓ <b>Quick-Set MMT-390</b>
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 10 needles.....	130.00	1 OP	✓ <b>Paradigm Quick-Set MMT-397</b>
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 10 needles; luer lock.....	130.00	1 OP	✓ <b>Quick-Set MMT-392</b>
9 mm teflon cannula; straight insertion; 80 cm tubing × 10 with 10 needles.....	130.00	1 OP	✓ <b>Paradigm Quick-Set MMT-386</b>

INSULIN PUMP RESERVOIR – Special Authority see [SA1604 on page 17](#) – Retail pharmacy

a) Maximum of 3 sets per prescription			
b) Only on a prescription			
c) Maximum of 13 packs of reservoir sets will be funded per year.			
10 × luer lock conversion cartridges 1.8 ml for Paradigm pumps.....	50.00	1 OP	✓ <b>ADR Cartridge 1.8</b>
Cartridge for 5 and 7 series pump; 1.8 ml × 10 .....	50.00	1 OP	✓ <b>Paradigm 1.8 Reservoir</b>
Cartridge for 7 series pump; 3.0 ml × 10 .....	50.00	1 OP	✓ <b>Paradigm 3.0 Reservoir</b>

**Digestives Including Enzymes**

## PANCREATIC ENZYME

Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U) .....	34.93	100	✓ <b><u>Creon 10000</u></b>
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase, 1,250 U protease)) .....	94.40	100	✓ <b><u>Panzytrat</u></b>
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U) .....	94.38	100	✓ <b><u>Creon 25000</u></b>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
URSODEOXYCHOLIC ACID – Special Authority see <a href="#">SA1739 below</a> – Retail pharmacy				
Cap 250 mg.....	37.95	100	✓	<b>Ursosan</b>

## ► **SA1739** Special Authority for Subsidy

**Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis)** from any relevant practitioner.

Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

**Initial application — (Chronic severe drug induced cholestatic liver injury)** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

**Initial application — (Primary biliary cholangitis)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

**Initial application — (Pregnancy)** from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

**Initial application — (Haematological Transplant)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogeneic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

**Initial application — (Total parenteral nutrition induced cholestasis)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

**Renewal — (Chronic severe drug induced cholestatic liver injury)** from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

**Renewal — (Pregnancy/Primary biliary cholangitis)** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

**Renewal — (Total parenteral nutrition induced cholestasis)** from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## Laxatives

### Bulk-forming Agents

ISPAGHULA (PSYLLIUM) HUSK – Only on a prescription

\* Powder for oral soln.....6.05      500 g OP      ✓ **Konsyl-D**

MUCILAGINOUS LAXATIVES WITH STIMULANTS

\* Dry.....6.02      500 g OP  
(17.32)  
2.41      200 g OP  
(8.72)      Normacol Plus  
Normacol Plus

### Faecal Softeners

DOCUSATE SODIUM – Only on a prescription

\* Tab 50 mg .....2.31      100      ✓ **Coloxyl**

\* Tab 120 mg .....3.13      100      ✓ **Coloxyl**

DOCUSATE SODIUM WITH SENNOSIDES

\* Tab 50 mg with sennosides 8 mg.....3.10      200      ✓ **Laxsol**

POLOXAMER – Only on a prescription

Not funded for use in the ear.

\* Oral drops 10%.....3.78      30 ml OP      ✓ **Coloxyl**

### Opioid Receptor Antagonists - Peripheral

METHYLNALTREXONE BROMIDE – Special Authority see [SA1691 below](#) – Retail pharmacy

Inj 12 mg per 0.6 ml vial .....36.00      1      ✓ **Relistor**

246.00      7      ✓ **Relistor**

#### ►SA1691 Special Authority for Subsidy

**Initial application — (Opioid induced constipation)** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Either:
  - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
  - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

### Osmotic Laxatives

GLYCEROL

\* Suppos 3.6 g – Only on a prescription .....9.25      20      ✓ **PSM**

LACTULOSE – Only on a prescription

\* Oral liq 10 g per 15 ml .....3.33      500 ml      ✓ **Laevolac**

Laevolac to be Sole Supply on 1 November 2019

MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE AND SODIUM CHLORIDE

Powder for oral soln 13.125 g with potassium chloride 46.6 mg,  
sodium bicarbonate 178.5 mg and sodium chloride 350.7 mg .....6.78      30      ✓ **Molaxole**

SODIUM ACID PHOSPHATE – Only on a prescription

Enema 16% with sodium phosphate 8% .....2.50      1      ✓ **Fleet Phosphate Enema**

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE – Only on a prescription				
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml.....	29.98	50	✓	Micolette
Micolette to be Sole Supply on 1 November 2019				

## Stimulant Laxatives

BISACODYL – Only on a prescription				
* Tab 5 mg .....	5.99	200	✓	Lax-Tab
* Suppos 10 mg .....	3.74	10	✓	Lax-Suppositories
SENNA – Only on a prescription				
* Tab, standardised.....	2.17	100		
	(6.84)			Senokot
	0.43	20		
	(1.72)			Senokot

## Metabolic Disorder Agents

ALGLUCOSIDASE ALFA – Special Authority see <a href="#">SA1622 below</a> – Retail pharmacy				
Inj 50 mg vial .....	1,142.60	1	✓	Myozyme

### ➡SA1622 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:  
All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
  - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
  - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
  - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
  - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

**Renewal** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>BETAINE</b> – Special Authority see <a href="#">SA1727 below</a> – Retail pharmacy				
Powder for oral soln.....	575.00	180 g OP	✓	<b>Cystadane</b>
<b>➤SA1727 Special Authority for Subsidy</b>				
<b>Initial application</b> only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:				
All of the following:				
1 The patient has a confirmed diagnosis of homocystinuria; and				
2 Any of the following:				
2.1 A cystathionine beta-synthase (CBS) deficiency; or				
2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or				
2.3 A disorder of intracellular cobalamin metabolism; and				
3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.				
<b>Renewal</b> only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.				
<b>GALSULFASE</b> – Special Authority see <a href="#">SA1593 below</a> – Retail pharmacy				
Inj 1 mg per ml, 5 ml vial.....	2,234.00	1	✓	<b>Naglazyme</b>
<b>➤SA1593 Special Authority for Subsidy</b>				
<b>Initial application</b> only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:				
Both:				
1 The patient has been diagnosed with mucopolysaccharidosis VI; and				
2 Either:				
2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or				
2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI.				
<b>Renewal</b> only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:				
All of the following:				
1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and				
2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and				
3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and				
4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.				
<b>IDURSULFASE</b> – Special Authority see <a href="#">SA1623 below</a> – Retail pharmacy				
Inj 2 mg per ml, 3 ml vial.....	4,608.30	1	✓	<b>Elaprase</b>
<b>➤SA1623 Special Authority for Subsidy</b>				
<b>Initial application</b> only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria:				
All of the following:				
1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and				
2 Either:				
2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or				
2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and				
3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and				
4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and				
5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.				

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

LARONIDASE – Special Authority see [SA1695 below](#) – Retail pharmacy

Inj 100 U per ml, 5 ml vial .....	1,335.16	1	✓	Aldurazyme
-----------------------------------	----------	---	---	------------

► **SA1695** Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria:  
All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysaccharidosis I-H); and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

SAPROPTERIN DIHYDROCHLORIDE – Special Authority see [SA1757 below](#) – Retail pharmacy

Tab soluble 100 mg .....	1,452.70	30 OP	✓	Kuvan
--------------------------	----------	-------	---	-------

► **SA1757** Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria:  
All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

**Renewal** only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
  - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
  - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
  - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
  - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE – Special Authority see [SA1599 on the next page](#) – Retail pharmacy

Soln 100 mg per ml .....	CBS	100 ml	✓	Amzoate <sup>S29</sup>
--------------------------	-----	--------	---	------------------------

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

### ►SA1599 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE – Special Authority see SA1598 below – Retail pharmacy

Grans 483 mg per g..... 1,920.00 174 g OP ✓ Pheburane

### ►SA1598 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

## Gaucher's Disease

TALIGLUCERASE ALFA – Special Authority see SA1734 below – Retail pharmacy

Inj 200 unit vial..... 1,072.00 1 ✓ Eleyso

### ►SA1734 Special Authority for Subsidy

Special Authority approved by the Gaucher Treatment Panel

Notes: Application details may be obtained from PHARMAC's website <http://www.pharmac.govt.nz> or:

The Co-ordinator, Gaucher Treatment Panel Phone: 04 460 4990

PHARMAC PO Box 10 254

Facsimile: 04 916 7571

Wellington

Email: [gaucherpanel@pharmac.govt.nz](mailto:gaucherpanel@pharmac.govt.nz)

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

### Access Criteria

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1) The patient has a diagnosis of symptomatic type 1 or type 3\* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2) Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and
- 3) Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, serum glucosylsphingosine, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
  - 1) Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia, thrombocytopenia; at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or massive symptomatic splenomegaly; or
  - 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g. hips or shoulder); spontaneous fractures or vertebral collapse; chronic bone pain not controlled by other pharmaceuticals; or
  - 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 4) Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
- 5) Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

\*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1) Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated; and
- 2) Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3) Radiological (MRI) signs of bone activity performed at one year and two years since initiation of treatment begins, and two to three yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4) Serum glucosylsphingosine levels taken at least 6 to 12 monthly show a decrease compared with baseline; and
- 5) Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 6) Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 7) Patient is compliant with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- 8) Supporting clinical information including test reports, MRI whole body STIR, serum glucosylsphingosine, haematological data, and other relevant investigations are submitted to the Gaucher Panel for assessment as required.

## Mouth and Throat

### Agents Used in Mouth Ulceration

#### BENZYDAMINE HYDROCHLORIDE

Soln 0.15% – Higher subsidy of \$20.31 per 500 ml with

Endorsement .....	9.00	500 ml	
	(20.31)		Difflam

Additional subsidy by endorsement for a patient who has oral mucositis as a result of treatment for cancer, and the prescription is endorsed accordingly.

#### CARMELLOSE SODIUM WITH GELATIN AND PECTIN

Paste .....	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)		Orabase
	1.52	5 g OP	
	(3.60)		Orabase
Powder .....	8.48	28 g OP	
	(10.95)		Stomahesive

#### CHLORHEXIDINE GLUCONATE

Mouthwash 0.2% .....	2.57	200 ml OP	✓ healthE
----------------------	------	-----------	-----------

#### CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE

* Adhesive gel 8.7% with cetalkonium chloride 0.01% .....	2.06	15 g OP	
	(6.00)		Bonjela

#### TRIAMCINOLONE ACETONIDE

Paste 0.1% .....	5.33	5 g OP	✓ Kenalog in Orabase
------------------	------	--------	----------------------

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
<b>Oropharyngeal Anti-infectives</b>			
AMPHOTERICIN B			
Lozenges 10 mg .....	5.86	20	✓ <b>Fungilin</b>
MICONAZOLE			
Oral gel 20 mg per g .....	4.74	40 g OP	✓ <b>Decozol</b>
NYSTATIN			
Oral liq 100,000 u per ml .....	1.95	24 ml OP	✓ <b>Nilstat</b>
<b>Other Oral Agents</b>			
For folinic mouthwash, pilocarpine oral liquid or saliva substitute formula refer Standard Formulae, <a href="#">page 235</a>			
HYDROGEN PEROXIDE			
* Soln 3% (10 vol) – Maximum of 200 ml per prescription .....	1.40	100 ml	✓ <b>Pharmacy Health</b>
<i>(Pharmacy Health Soln 3% (10 vol) to be delisted 1 July 2020)</i>			
THYMOL GLYCERIN			
* Compound, BPC .....	9.15	500 ml	✓ <b>PSM</b>
<b>Vitamins</b>			
<b>Vitamin A</b>			
VITAMIN A WITH VITAMINS D AND C			
* Soln 1000 u with Vitamin D 400 u and ascorbic acid 30 mg per 10 drops .....	4.50	10 ml OP	✓ <b>Vitadol C</b>
<i>(Vitadol C Soln 1000 u with Vitamin D 400 u and ascorbic acid 30 mg per 10 drops to be delisted 1 December 2019)</i>			
<b>Vitamin B</b>			
HYDROXOCOBALAMIN			
* Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a PSO .....	1.89	3	✓ <b>Neo-B12</b>
PYRIDOXINE HYDROCHLORIDE			
a) No more than 100 mg per dose			
b) Only on a prescription			
* Tab 25 mg – No patient co-payment payable .....	2.70	90	✓ <b>Vitamin B6 25</b>
* Tab 50 mg .....	13.63	500	✓ <b>Apo-Pyridoxine</b>
THIAMINE HYDROCHLORIDE – Only on a prescription			
* Tab 50 mg .....	4.89	100	✓ <b>Max Health</b>
VITAMIN B COMPLEX			
* Tab, strong, BPC .....	7.15	500	✓ <b>Bplex</b>
<b>Vitamin C</b>			
ASCORBIC ACID			
a) No more than 100 mg per dose			
b) Only on a prescription			
* Tab 100 mg .....	8.10	500	✓ <b>Cvite</b>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>Vitamin D</b>				
ALFACALCIDOL				
* Cap 0.25 mcg .....	26.32	100	✓	<u>One-Alpha</u>
* Cap 1 mcg .....	87.98	100	✓	<u>One-Alpha</u>
* Oral drops 2 mcg per ml .....	60.68	20 ml OP	✓	<u>One-Alpha</u>
CALCITRIOL				
* Cap 0.25 mcg .....	7.95	100	✓	<u>Calcitriol-AFT</u>
* Cap 0.5 mcg .....	13.75	100	✓	<u>Calcitriol-AFT</u>
COLECALCIFEROL				
* Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescription.....	2.50	12	✓	<u>Vit.D3</u>
* Oral liq 188 mcg per ml (7,500 iu per ml) .....	9.00	4.8 ml OP	✓	<u>Puria</u>

## Multivitamin Preparations

MULTIVITAMIN RENAL – Special Authority see [SA1546 below](#) – Retail pharmacy

* Cap .....	6.49	30	✓	Clinicians Renal Vit
-------------	------	----	---	----------------------

► [SA1546](#) **Special Authority for Subsidy**

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or
- 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73 m<sup>2</sup> body surface area (BSA).

MULTIVITAMINS – Special Authority see [SA1036 below](#) – Retail pharmacy

* Powder .....	72.00	200 g OP	✓	Paediatric Seravit
----------------	-------	----------	---	--------------------

► [SA1036](#) **Special Authority for Subsidy**

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism.

**Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins.

VITAMINS

* Tab (BPC cap strength) .....	10.50	1,000	✓	Mvite
* Cap (fat soluble vitamins A, D, E, K) – Special Authority see <a href="#">SA1720 below</a> – Retail pharmacy .....	23.40	60	✓	Vitabdeck

► [SA1720](#) **Special Authority for Subsidy**

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency; or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.



	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>Minerals</b>				
<b>Calcium</b>				
CALCIUM CARBONATE				
* Tab eff 1.75 g (1 g elemental) .....	28.40	20	✓	Calcium Sandoz <sup>S29</sup>
* Tab 1.25 g (500 mg elemental) .....	7.52	250	✓	Arrow-Calcium
CALCIUM GLUCONATE				
* Inj 10%, 10 ml ampoule .....	64.00	20	✓	Max Health <sup>S29</sup>
<b>Fluoride</b>				
SODIUM FLUORIDE				
* Tab 1.1 mg (0.5 mg elemental) .....	5.75	100	✓	PSM
<b>Iodine</b>				
POTASSIUM IODATE				
* Tab 253 mcg (150 mcg elemental iodine) .....	4.69	90	✓	NeuroTabs
<b>Iron</b>				
FERRIC CARBOXYMALTOSE – Special Authority see <a href="#">SA1840 below</a> – Retail pharmacy				
Inj 50 mg per ml, 10 ml .....	150.00	1	✓	Ferinject
» <b>SA1840</b> Special Authority for Subsidy				
<b>Initial application — (serum ferritin less than or equal to 20 mcg/L)</b> from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:				
Both:				
1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and				
2 Any of the following:				
2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or				
2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or				
2.3 Rapid correction of anaemia is required.				
<b>Renewal — (serum ferritin less than or equal to 20 mcg/L)</b> from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:				
Both:				
1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and				
2 A re-trial with oral iron is clinically inappropriate.				
<b>Initial application — (iron deficiency anaemia)</b> only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:				
Both:				
1 Patient has been diagnosed with iron-deficiency anaemia; and				
2 Any of the following:				
2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or				
2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or				
2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease				

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

# ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

and a trial of oral iron is unlikely to be effective; or

2.4 Rapid correction of anaemia is required.

**Renewal — (iron deficiency anaemia)** only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist.

Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

## FERROUS FUMARATE

\* Tab 200 mg (65 mg elemental) .....3.09 100 ✓ **Ferro-tab**

## FERROUS FUMARATE WITH FOLIC ACID

\* Tab 310 mg (100 mg elemental) with folic acid 350 mcg .....4.68 60 ✓ **Ferro-F-Tabs**

## FERROUS SULFATE

\* Oral liq 30 mg (6 mg elemental) per 1 ml .....12.08 500 ml ✓ **Ferodan**  
Ferodan to be Sole Supply on 1 November 2019

## FERROUS SULPHATE

\* Tab long-acting 325 mg (105 mg elemental) .....2.06 30 ✓ **Ferrograd**

## IRON POLYMALTOSE

\* Inj 50 mg per ml, 2 ml ampoule .....15.22 5 ✓ **Ferrum H**  
34.50 ✓ **Ferrosig**

## Magnesium

For magnesium hydroxide mixture refer Standard Formulae, [page 235](#)

## MAGNESIUM HYDROXIDE

Suspension 8% .....72.20 500 ml ✓ **T&R** S29

## MAGNESIUM SULPHATE

\* Inj 2 mmol per ml, 5 ml ampoule .....10.21 10 ✓ **DBL**  
✓ **DBL** S29 S29

## Zinc

## ZINC SULPHATE

\* Cap 137.4 mg (50 mg elemental) .....11.00 100 ✓ **Zincaps**  
Zincaps to be Sole Supply on 1 December 2019

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## Antianaemics

### Hypoplastic and Haemolytic

#### ►SA1775 Special Authority for Subsidy

**Initial application — (chronic renal failure)** from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
  - 3.1 Both:
    - 3.1.1 Patient does not have diabetes mellitus; and
    - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
  - 3.2 Both:
    - 3.2.1 Patient has diabetes mellitus; and
    - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
  - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

**Initial application — (myelodysplasia)** from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)\*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with \* is an unapproved indication

**Renewal — (chronic renal failure)** from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

**Renewal — (myelodysplasia)** from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with \* is an unapproved indication

## BLOOD AND BLOOD FORMING ORGANS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
EPOETIN ALFA – Special Authority see <a href="#">SA1775 on the previous page</a> – Retail pharmacy				
Wastage claimable				
Inj 1,000 iu in 0.5 ml, syringe.....	250.00	6	✓	<b>Binocrit</b>
Inj 2,000 iu in 1 ml, syringe.....	100.00	6	✓	<b>Binocrit</b>
Inj 3,000 iu in 0.3 ml, syringe.....	150.00	6	✓	<b>Binocrit</b>
Inj 4,000 iu in 0.4 ml, syringe.....	96.50	6	✓	<b>Binocrit</b>
Inj 5,000 iu in 0.5 ml, syringe.....	125.00	6	✓	<b>Binocrit</b>
Inj 6,000 iu in 0.6 ml, syringe.....	145.00	6	✓	<b>Binocrit</b>
Inj 8,000 iu in 0.8 ml, syringe.....	175.00	6	✓	<b>Binocrit</b>
Inj 10,000 iu in 1 ml, syringe.....	197.50	6	✓	<b>Binocrit</b>
Inj 40,000 iu in 1 ml, syringe.....	250.00	1	✓	<b>Binocrit</b>

### Megaloblastic

FOLIC ACID				
* Tab 0.8 mg .....	21.84	1,000	✓	<b>Apo-Folic Acid</b>
* Tab 5 mg .....	12.12	500	✓	<b>Apo-Folic Acid</b>
Oral liq 50 mcg per ml .....	26.00	25 ml OP	✓	<b>Biomed</b>

### Antifibrinolytics, Haemostatics and Local Sclerosants

#### EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Inj 250 iu vial.....	612.50	1	✓	<b>Alprolix</b>
Inj 500 iu vial.....	1,225.00	1	✓	<b>Alprolix</b>
Inj 1,000 iu vial.....	2,450.00	1	✓	<b>Alprolix</b>
Inj 2,000 iu vial.....	4,900.00	1	✓	<b>Alprolix</b>
Inj 3,000 iu vial.....	7,350.00	1	✓	<b>Alprolix</b>

#### ELTROMBOPAG – Special Authority see [SA1743 below](#) – Retail pharmacy

Wastage claimable				
Tab 25 mg .....	1,550.00	28	✓	<b>Revolade</b>
Tab 50 mg .....	3,100.00	28	✓	<b>Revolade</b>

#### ► [SA1743](#) Special Authority for Subsidy

**Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy)** only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
  - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
  - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
  - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

**Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy)** only from a haematologist.

Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

**Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy)** only from a haematologist.

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
  - 3.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per microliter; or
  - 3.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

**Initial application — (severe aplastic anaemia)** only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
  - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
  - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

**Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy)** only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

**Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy)** only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

**Renewal — (severe aplastic anaemia)** only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

## EPTACOG ALFA [RECOMBINANT FACTOR VIIA] – [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 1 mg syringe .....	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe .....	2,356.60	1	✓ NovoSeven RT
Inj 5 mg syringe .....	5,891.50	1	✓ NovoSeven RT
Inj 8 mg syringe .....	9,426.40	1	✓ NovoSeven RT

## FACTOR EIGHT INHIBITOR BYPASSING FRACTION – [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U .....	1,315.00	1	✓ FEIBA NF
Inj 1,000 U .....	2,630.00	1	✓ FEIBA NF
Inj 2,500 U .....	6,575.00	1	✓ FEIBA NF

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

# BLOOD AND BLOOD FORMING ORGANS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] – [Xpharm]</b>				
For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.				
Inj 250 iu prefilled syringe.....	210.00	1	✓	<b>Xyntha</b>
Inj 500 iu prefilled syringe.....	420.00	1	✓	<b>Xyntha</b>
Inj 1,000 iu prefilled syringe.....	840.00	1	✓	<b>Xyntha</b>
Inj 2,000 iu prefilled syringe.....	1,680.00	1	✓	<b>Xyntha</b>
Inj 3,000 iu prefilled syringe.....	2,520.00	1	✓	<b>Xyntha</b>
<b>NONACOG ALFA [RECOMBINANT FACTOR IX] – [Xpharm]</b>				
For patients with haemophilia, whose funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.				
Inj 250 iu vial.....	310.00	1	✓	<b>BeneFIX</b>
Inj 500 iu vial.....	620.00	1	✓	<b>BeneFIX</b>
Inj 1,000 iu vial.....	1,240.00	1	✓	<b>BeneFIX</b>
Inj 2,000 iu vial.....	2,480.00	1	✓	<b>BeneFIX</b>
Inj 3,000 iu vial.....	3,720.00	1	✓	<b>BeneFIX</b>
<i>(BeneFIX Inj 250 iu vial to be delisted 1 November 2019)</i>				
<i>(BeneFIX Inj 500 iu vial to be delisted 1 November 2019)</i>				
<i>(BeneFIX Inj 1,000 iu vial to be delisted 1 November 2019)</i>				
<i>(BeneFIX Inj 2,000 iu vial to be delisted 1 November 2019)</i>				
<i>(BeneFIX Inj 3,000 iu vial to be delisted 1 November 2019)</i>				
<b>NONACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xpharm]</b>				
For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.				
Inj 500 iu vial.....	435.00	1	✓	<b>RIXUBIS</b>
Inj 1,000 iu vial.....	870.00	1	✓	<b>RIXUBIS</b>
Inj 2,000 iu vial.....	1,740.00	1	✓	<b>RIXUBIS</b>
Inj 3,000 iu vial.....	2,610.00	1	✓	<b>RIXUBIS</b>
<b>OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) – [Xpharm]</b>				
For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.				
Inj 250 iu vial.....	210.00	1	✓	<b>Advate</b>
Inj 500 iu vial.....	420.00	1	✓	<b>Advate</b>
Inj 1,000 iu vial.....	840.00	1	✓	<b>Advate</b>
Inj 1,500 iu vial.....	1,260.00	1	✓	<b>Advate</b>
Inj 2,000 iu vial.....	1,680.00	1	✓	<b>Advate</b>
Inj 3,000 iu vial.....	2,520.00	1	✓	<b>Advate</b>
<b>OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE FS) – [Xpharm]</b>				
For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.				
Inj 250 iu vial.....	237.50	1	✓	<b>Kogenate FS</b>
Inj 500 iu vial.....	475.00	1	✓	<b>Kogenate FS</b>
Inj 1,000 iu vial.....	950.00	1	✓	<b>Kogenate FS</b>
Inj 2,000 iu vial.....	1,900.00	1	✓	<b>Kogenate FS</b>
Inj 3,000 iu vial.....	2,850.00	1	✓	<b>Kogenate FS</b>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII] – [Xpharm]</b>				
For patients with haemophilia A receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.				
Inj 250 iu vial.....	300.00	1	✓	Adynovate
Inj 500 iu vial.....	600.00	1	✓	Adynovate
Inj 1,000 iu vial.....	1,200.00	1	✓	Adynovate
Inj 2,000 iu vial.....	2,400.00	1	✓	Adynovate
<b>SODIUM TETRADECYL SULPHATE</b>				
* Inj 3% 2 ml.....	28.50 (73.00)	5		Fibro-vein
<b>TRANEXAMIC ACID</b>				
Tab 500 mg .....	20.67	100	✓	Cyklokapron
<b>Vitamin K</b>				
<b>PHYTOMENADIONE</b>				
Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO.....	8.00	5	✓	Konaktion MM
Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO.....	9.21	5	✓	Konaktion MM
<b>Antithrombotic Agents</b>				
<b>Antiplatelet Agents</b>				
<b>ASPIRIN</b>				
* Tab 100 mg .....	10.80	990	✓	Ethics Aspirin EC
Ethics Aspirin EC to be Sole Supply on 1 November 2019				
<b>CLOPIDOGREL</b>				
* Tab 75 mg .....	5.44	84	✓	Arrow - Clopid
<b>DIPYRIDAMOLE</b>				
* Tab long-acting 150 mg.....	10.90	60	✓	Pytazen SR
<b>PRASUGREL – Special Authority see SA1201 below – Retail pharmacy</b>				
Tab 5 mg .....	108.00	28	✓	Effient
Tab 10 mg .....	120.00	28	✓	Effient
<b>»SA1201 Special Authority for Subsidy</b>				
<b>Initial application — (coronary angioplasty and bare metal stent)</b> from any relevant practitioner. Approvals valid for 6 months where the patient has undergone coronary angioplasty in the previous 4 weeks and is clopidogrel-allergic*.				
<b>Initial application — (drug eluting stent)</b> from any relevant practitioner. Approvals valid for 12 months where the patient has had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.				
<b>Initial application — (stent thrombosis)</b> from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.				
<b>Renewal — (coronary angioplasty and bare metal stent)</b> from any relevant practitioner. Approvals valid for 6 months where the patient has undergone coronary angioplasty or had a bare metal cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.				
<b>Renewal — (drug eluting stent)</b> from any relevant practitioner. Approvals valid for 12 months where had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.				
Note: * Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.				
<b>TICAGRELOR – Special Authority see SA1382 on the next page – Retail pharmacy</b>				
* Tab 90 mg .....	90.00	56	✓	Brilinta

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

## ►SA1382 Special Authority for Subsidy

**Initial application — (acute coronary syndrome)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

**Renewal — (subsequent acute coronary syndrome)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

## Heparin and Antagonist Preparations

DALTEPARIN SODIUM – Special Authority see [SA1270 below](#) – Retail pharmacy

Inj 2,500 iu per 0.2 ml prefilled syringe.....	19.97	10	✓ Fragmin
Inj 5,000 iu per 0.2 ml prefilled syringe.....	39.94	10	✓ Fragmin
Inj 7,500 iu per 0.75 ml graduated syringe .....	60.03	10	✓ Fragmin
Inj 10,000 iu per 1 ml graduated syringe .....	77.55	10	✓ Fragmin
Inj 12,500 iu per 0.5 ml prefilled syringe.....	99.96	10	✓ Fragmin
Inj 15,000 iu per 0.6 ml prefilled syringe.....	120.05	10	✓ Fragmin
Inj 18,000 iu per 0.72 ml prefilled syringe.....	158.47	10	✓ Fragmin

(Fragmin Inj 2,500 iu per 0.2 ml prefilled syringe to be delisted 1 April 2020)

(Fragmin Inj 5,000 iu per 0.2 ml prefilled syringe to be delisted 1 April 2020)

(Fragmin Inj 7,500 iu per 0.75 ml graduated syringe to be delisted 1 April 2020)

(Fragmin Inj 10,000 iu per 1 ml graduated syringe to be delisted 1 April 2020)

(Fragmin Inj 12,500 iu per 0.5 ml prefilled syringe to be delisted 1 January 2020)

(Fragmin Inj 15,000 iu per 0.6 ml prefilled syringe to be delisted 1 January 2020)

(Fragmin Inj 18,000 iu per 0.72 ml prefilled syringe to be delisted 1 January 2020)

## ►SA1270 Special Authority for Subsidy

**Initial application — (Pregnancy or Malignancy)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy.

**Initial application — (Venous thromboembolism other than in pregnancy or malignancy)** from any relevant practitioner.

Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

**Renewal — (Pregnancy or Malignancy)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the

continued...



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

continued...

following criteria:

Either:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy.

**Renewal — (Venous thromboembolism other than in pregnancy or malignancy)** from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, Acute Coronary Syndrome, cardioversion, or prior to oral anti-coagulation).

ENOXAPARIN SODIUM – Special Authority see [SA1646 below](#) – Retail pharmacy

Inj 20 mg in 0.2 ml syringe.....	27.93	10	✓ <b>Clexane</b>
Inj 40 mg in 0.4 ml syringe.....	37.27	10	✓ <b>Clexane</b>
Inj 60 mg in 0.6 ml syringe.....	56.18	10	✓ <b>Clexane</b>
Inj 80 mg in 0.8 ml syringe.....	74.90	10	✓ <b>Clexane</b>
Inj 100 mg in 1 ml syringe.....	93.80	10	✓ <b>Clexane</b>
Inj 120 mg in 0.8 ml syringe.....	116.55	10	✓ <b>Clexane</b>
Inj 150 mg in 1 ml syringe.....	133.20	10	✓ <b>Clexane</b>

► [SA1646](#) Special Authority for Subsidy

**Initial application — (Pregnancy, Malignancy or Haemodialysis)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

**Initial application — (Venous thromboembolism other than in pregnancy or malignancy)** from any relevant practitioner.

Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

**Renewal — (Pregnancy, Malignancy or Haemodialysis)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

**Renewal — (Venous thromboembolism other than in pregnancy or malignancy)** from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

HEPARIN SODIUM

Inj 1,000 iu per ml, 5 ml ampoule .....	58.57	50	✓ <b>Pfizer</b>
Inj 5,000 iu per ml, 1 ml .....	28.40	5	✓ <b>Hospira</b>
			✓ <b>Pfizer</b>
Inj 5,000 iu per ml, 5 ml ampoule .....	203.68	50	✓ <b>Pfizer</b>
Inj 25,000 iu per ml, 0.2 ml .....	19.00	5	✓ <b>Hospira</b>
	122.00	10	✓ <b>Wockhardt</b> <sup>S29</sup>
	190.00	50	✓ <b>Pfizer</b> <sup>S29</sup>

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\* Three months or six months, as applicable, dispensed all-at-once

## BLOOD AND BLOOD FORMING ORGANS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
HEPARINISED SALINE				
Inj 10 iu per ml, 5 ml .....	56.94	50	✓	Pfizer

### Oral Anticoagulants

DABIGATRAN				
Cap 75 mg – No more than 2 cap per day .....	76.36	60	✓	Pradaxa
Cap 110 mg .....	76.36	60	✓	Pradaxa
Cap 150 mg .....	76.36	60	✓	Pradaxa
RIVAROXABAN				
Tab 10 mg – No more than 1 tab per day .....	83.10	30	✓	Xarelto
Tab 15 mg – Up to 14 tab available on a PSO .....	77.56	28	✓	Xarelto
Tab 20 mg .....	77.56	28	✓	Xarelto
WARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg .....	3.46	50	✓	Coumadin
	7.60	100	✓	Marevan
* Tab 2 mg .....	4.31	50	✓	Coumadin
* Tab 3 mg .....	11.80	100	✓	Marevan
* Tab 5 mg .....	5.93	50	✓	Coumadin
	13.50	100	✓	Marevan

### Blood Colony-stimulating Factors

FILGRASTIM – Special Authority see <a href="#">SA1259 below</a> – Retail pharmacy				
Inj 300 mcg per 0.5 ml prefilled syringe .....	96.22	10	✓	Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe .....	161.50	10	✓	Nivestim

#### ► [SA1259](#) Special Authority for Subsidy

**Initial application** only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%\*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia ( $ANC < 0.5 \times 10^9/L$ ); or
- 5 Treatment of drug-induced prolonged neutropenia ( $ANC < 0.5 \times 10^9/L$ ).

Note: \*Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM – Special Authority see <a href="#">SA1384 below</a> – Retail pharmacy				
Inj 6 mg per 0.6 ml syringe .....	1,080.00	1	✓	Neulastim

#### ► [SA1384](#) Special Authority for Subsidy

**Initial application** only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%\*).

Note: \*Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>Fluids and Electrolytes</b>				
<b>Intravenous Administration</b>				
<b>GLUCOSE [DEXTROSE]</b>				
* Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO .....	29.50	5	✓	<b>Biomed</b>
* Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO .....	14.50	1	✓	<b>Biomed</b>
<b>POTASSIUM CHLORIDE</b>				
* Inj 75 mg per ml, 10 ml .....	55.00	50	✓	<b>AstraZeneca</b>
<b>SODIUM BICARBONATE</b>				
Inj 8.4%, 50 ml .....	19.95	1	✓	<b>Biomed</b>
a) Up to 5 inj available on a PSO				
b) Not in combination				
Inj 8.4%, 100 ml .....	20.50	1	✓	<b>Biomed</b>
a) Up to 5 inj available on a PSO				
b) Not in combination				
<b>SODIUM CHLORIDE</b>				
Not funded for use as a nasal drop. Not funded for nebuliser use except when used in conjunction with an antibiotic intended for nebuliser use.				
Inj 0.9%, bag – Up to 2000 ml available on a PSO .....	1.23	500 ml	✓	<b>Baxter</b>
	1.26	1,000 ml	✓	<b>Baxter</b>
Only if prescribed on a prescription for renal dialysis, maternity or post-natal care in the home of the patient, or on a PSO for emergency use. (500 ml and 1,000 ml packs)				
Inj 23.4% (4 mmol/ml), 20 ml ampoule .....	33.00	5	✓	<b>Biomed</b>
For Sodium chloride oral liquid formulation refer Standard Formulae, <a href="#">page 235</a>				
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO .....	2.80	20	✓	<b>Fresenius Kabi</b>
	7.00	50	✓	<b>InterPharma</b>
			✓	<b>Multichem</b>
Fresenius Kabi to be Sole Supply on 1 December 2019				
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO .....	5.40	50	✓	<b>Fresenius Kabi</b>
	6.63		✓	<b>Pfizer</b>
Fresenius Kabi to be Sole Supply on 1 December 2019				
Inj 0.9%, 20 ml ampoule .....	5.00	20	✓	<b>Fresenius Kabi</b>
			✓	<b>Multichem</b>
	7.50	30	✓	<b>InterPharma</b>
Fresenius Kabi to be Sole Supply on 1 December 2019				
<i>(InterPharma Inj 0.9%, 5 ml ampoule to be delisted 1 December 2019)</i>				
<i>(Multichem Inj 0.9%, 5 ml ampoule to be delisted 1 December 2019)</i>				
<i>(Pfizer Inj 0.9%, 10 ml ampoule to be delisted 1 December 2019)</i>				
<i>(Multichem Inj 0.9%, 20 ml ampoule to be delisted 1 December 2019)</i>				
<i>(InterPharma Inj 0.9%, 20 ml ampoule to be delisted 1 December 2019)</i>				
<b>TOTAL PARENTERAL NUTRITION (TPN) – Retail pharmacy-Specialist</b>				
Infusion .....	CBS	1 OP	✓	<b>TPN</b>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>WATER</b>				
1) On a prescription or Practitioner's Supply Order only when on the same form as an injection listed in the Pharmaceutical Schedule requiring a solvent or diluent; or				
2) On a bulk supply order; or				
3) When used in the extemporaneous compounding of eye drops; or				
4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.				
Inj 5 ml ampoule – Up to 5 inj available on a PSO .....	7.00	50	✓	<b>InterPharma</b>
Inj 10 ml ampoule – Up to 5 inj available on a PSO .....	6.63	50	✓	<b>Pfizer</b>
Inj 20 ml ampoule – Up to 5 inj available on a PSO .....	5.00	20	✓	<b>Fresenius Kabi</b>
			✓	<b>Multichem</b>
	7.50	30	✓	<b>InterPharma</b>

## Oral Administration

<b>CALCIUM POLYSTYRENE SULPHONATE</b>				
Powder .....	169.85	300 g OP	✓	<b>Calcium Resonium</b>
<b>COMPOUND ELECTROLYTES</b>				
Powder for oral soln – Up to 10 sach available on a PSO .....	2.30	10	✓	<b>Enerlyte</b>
<b>COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]</b>				
Soln with electrolytes (2 x 500 ml) .....	6.55	1,000 ml OP	✓	<b>Pedialyte - Bubblegum</b>
<b>PHOSPHORUS</b>				
Tab eff 500 mg (16 mmol) .....	82.50	100	✓	<b>Phosphate Phebra</b>
<b>POTASSIUM CHLORIDE</b>				
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq) .....	5.26 (11.85)	60		Chlorvescent
* Tab long-acting 600 mg (8 mmol) .....	8.90	200	✓	<b>Span-K</b>
<b>SODIUM BICARBONATE</b>				
Cap 840 mg .....	8.52	100	✓	<b>Sodibic</b>
			✓	<b>Sodibic</b>
<b>SODIUM POLYSTYRENE SULPHONATE</b>				
Powder .....	84.65	454 g OP	✓	<b>Resonium-A</b>

Subsidy Fully  
(Manufacturer's Price) Subsidised Brand or  
\$ Per ✓ Generic  
Manufacturer

## Alpha-Adrenoceptor Blockers

### Alpha Adrenoceptor Blockers

#### DOXAZOSIN

* Tab 2 mg .....	6.75	500	✓ Apo-Doxazosin
* Tab 4 mg .....	9.09	500	✓ Apo-Doxazosin

#### PHENOXYBENZAMINE HYDROCHLORIDE

* Cap 10 mg .....	65.00	30	✓ BNM <sup>\$29</sup>
	216.67	100	✓ Dibenzyline <sup>\$29</sup>

#### PRAZOSIN

* Tab 1 mg .....	5.53	100	✓ Apo-Prazosin
* Tab 2 mg .....	7.00	100	✓ Apo-Prazosin
* Tab 5 mg .....	11.70	100	✓ Apo-Prazosin

#### TERAZOSIN

* Tab 1 mg .....	0.59	28	✓ Actavis
* Tab 2 mg .....	7.50	500	✓ Apo-Terazosin
* Tab 5 mg .....	10.90	500	✓ Apo-Terazosin

## Agents Affecting the Renin-Angiotensin System

### ACE Inhibitors

#### CAPTOPRIL

* Oral liq 5 mg per ml .....	94.99	95 ml OP	✓ Capoten
Oral liquid restricted to children under 12 years of age.			

#### CILAZAPRIL

* Tab 0.5 mg .....	2.09	90	✓ Zapril
* Tab 2.5 mg .....	4.80	90	✓ Zapril
	7.20	200	✓ Apo-Cilazapril
* Tab 5 mg .....	8.35	90	✓ Zapril
	12.00	200	✓ Apo-Cilazapril

(Apo-Cilazapril Tab 2.5 mg to be delisted 1 February 2020)

(Apo-Cilazapril Tab 5 mg to be delisted 1 February 2020)

#### ENALAPRIL MALEATE

* Tab 5 mg .....	3.84	100	✓ Ethics Enalapril
* Tab 10 mg .....	4.96	100	✓ Ethics Enalapril
* Tab 20 mg .....	7.12	100	✓ Ethics Enalapril

#### LISINOPRIL

* Tab 5 mg .....	2.07	90	✓ Ethics Lisinopril
* Tab 10 mg .....	2.36	90	✓ Ethics Lisinopril
* Tab 20 mg .....	3.17	90	✓ Ethics Lisinopril

#### PERINDOPRIL

* Tab 2 mg .....	3.75	30	✓ Apo-Perindopril
* Tab 4 mg .....	4.80	30	✓ Apo-Perindopril

#### QUINAPRIL

* Tab 5 mg .....	6.01	90	✓ Arrow-Quinapril 5
* Tab 10 mg .....	3.16	90	✓ Arrow-Quinapril 10
* Tab 20 mg .....	4.89	90	✓ Arrow-Quinapril 20

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

## ACE Inhibitors with Diuretics

### CILAZAPRIL WITH HYDROCHLOROTHIAZIDE

* Tab 5 mg with hydrochlorothiazide 12.5 mg.....	10.18	100	✓	<b>Apo-Cilazapril/ Hydrochlorothiazide</b>
--	-------	-----	---	--

### QUINAPRIL WITH HYDROCHLOROTHIAZIDE

* Tab 10 mg with hydrochlorothiazide 12.5 mg.....	3.83	30	✓	<b>Accuretic 10</b>
* Tab 20 mg with hydrochlorothiazide 12.5 mg.....	4.92	30	✓	<b>Accuretic 20</b>

## Angiotensin II Antagonists

### CANDESARTAN CILEXETIL

* Tab 4 mg .....	1.90	90	✓	<b>Candestar</b>
* Tab 8 mg .....	2.28	90	✓	<b>Candestar</b>
* Tab 16 mg .....	3.67	90	✓	<b>Candestar</b>
* Tab 32 mg .....	6.39	90	✓	<b>Candestar</b>

### LOSARTAN POTASSIUM

* Tab 12.5 mg .....	1.39	84	✓	<b>Losartan Actavis</b>
* Tab 25 mg .....	1.63	84	✓	<b>Losartan Actavis</b>
* Tab 50 mg .....	2.00	84	✓	<b>Losartan Actavis</b>
* Tab 100 mg .....	2.31	84	✓	<b>Losartan Actavis</b>

## Angiotensin II Antagonists with Diuretics

### LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE

Tab 50 mg with hydrochlorothiazide 12.5 mg.....	1.88	30	✓	<b>Arrow-Losartan &amp; Hydrochlorothiazide</b>
---	------	----	---	---

## Angiotensin II Antagonists with Neprilysin Inhibitors

### SACUBITRIL WITH VALSARTAN – Special Authority see [SA1751 below](#) – Retail pharmacy

Note: Due to the angiotensin II receptor blocking activity of sacubitril with valsartan it should not be co-administered with an ACE inhibitor or another ARB.

Tab 24.3 mg with valsartan 25.7 mg .....	190.00	56	✓	<b>Entresto 24/26</b>
Tab 48.6 mg with valsartan 51.4 mg .....	190.00	56	✓	<b>Entresto 49/51</b>
Tab 97.2 mg with valsartan 102.8 mg .....	190.00	56	✓	<b>Entresto 97/103</b>

### ► **SA1751** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
  - 2.1 Patient is in NYHA/WHO functional class II; or
  - 2.2 Patient is in NYHA/WHO functional class III; or
  - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>Antiarrhythmics</b>				
For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, <a href="#">page 118</a>				
<b>AMIODARONE HYDROCHLORIDE</b>				
▲ Tab 100 mg – Retail pharmacy-Specialist.....	3.80	30	✓	Aratac
	4.66		✓	Cordarone-X
Aratac to be Sole Supply on 1 December 2019				
▲ Tab 200 mg – Retail pharmacy-Specialist.....	5.25	30	✓	Aratac
	7.63		✓	Cordarone-X
Aratac to be Sole Supply on 1 December 2019				
Inj 50 mg per ml, 3 ml ampoule – Up to 6 inj available on a PSO .....	9.98	5	✓	Lodi
	11.98	6	✓	Cordarone-X
	16.37	10	✓	Max Health
<i>(Cordarone-X Tab 100 mg to be delisted 1 December 2019)</i>				
<i>(Cordarone-X Tab 200 mg to be delisted 1 December 2019)</i>				
<i>(Lodi Inj 50 mg per ml, 3 ml ampoule to be delisted 1 February 2020)</i>				
<i>(Cordarone-X Inj 50 mg per ml, 3 ml ampoule to be delisted 1 February 2020)</i>				
<b>ATROPINE SULPHATE</b>				
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO .....	12.07	10	✓	Martindale
<b>DIGOXIN</b>				
* Tab 62.5 mcg – Up to 30 tab available on a PSO .....	7.00	240	✓	Lanoxin PG
Lanoxin PG to be Sole Supply on 1 November 2019				
* Tab 250 mcg – Up to 30 tab available on a PSO .....	15.20	240	✓	Lanoxin
Lanoxin to be Sole Supply on 1 November 2019				
* Oral liq 50 mcg per ml .....	16.60	60 ml	✓	Lanoxin
			✓	Lanoxin S29 <small>\$29</small>
<b>DISOPYRAMIDE PHOSPHATE</b>				
▲ Cap 100 mg .....	23.87	100	✓	Rythmodan
<b>FLECAINIDE ACETATE – Retail pharmacy-Specialist</b>				
▲ Tab 50 mg .....	19.95	60	✓	Flecainide BNM
	38.95		✓	Tambocor
▲ Cap long-acting 100 mg .....	38.95	30	✓	Tambocor CR
	39.51	90	✓	Flecainide Controlled Release Teva
Flecainide Controlled Release Teva to be Sole Supply on 1 December 2019				
▲ Cap long-acting 200 mg .....	61.06	90	✓	Flecainide Controlled Release Teva
	68.78	30	✓	Tambocor CR
Flecainide Controlled Release Teva to be Sole Supply on 1 December 2019				
Inj 10 mg per ml, 15 ml ampoule .....	52.45	5	✓	Tambocor
<i>(Tambocor Tab 50 mg to be delisted 1 February 2020)</i>				
<i>(Tambocor CR Cap long-acting 100 mg to be delisted 1 December 2019)</i>				
<i>(Tambocor CR Cap long-acting 200 mg to be delisted 1 December 2019)</i>				

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\* Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>MEXILETINE HYDROCHLORIDE</b>				
▲ Cap 150 mg.....	162.00	100	✓	<b>Mexiletine Hydrochloride USP</b> <sup>\$29</sup>
▲ Cap 250 mg.....	202.00	100	✓	<b>Mexiletine Hydrochloride USP</b> <sup>\$29</sup>
<b>PROPAFENONE HYDROCHLORIDE – Retail pharmacy-Specialist</b>				
▲ Tab 150 mg .....	40.90	50	✓	<b>Rytmonorm</b>

## Antihypotensives

<b>MIDODRINE – Special Authority see SA1474 below – Retail pharmacy</b>				
Tab 2.5 mg .....	53.00	100	✓	<b>Gutron</b>
Tab 5 mg .....	79.00	100	✓	<b>Gutron</b>

### ►SA1474 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Note: Treatment should be started with small doses and titrated upwards as necessary. Hypertension should be avoided, and the usual target is a standing systolic blood pressure of 90 mm Hg.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

## Beta-Adrenoceptor Blockers

### Beta Adrenoceptor Blockers

<b>ATENOLOL</b>				
* Tab 50 mg .....	4.26	500	✓	<b>Mylan Atenolol</b>
* Tab 100 mg .....	7.30	500	✓	<b>Mylan Atenolol</b>
* Oral liq 25 mg per 5 ml .....	21.25	300 ml OP	✓	<b>Atenolol AFT</b>
Restricted to children under 12 years of age.				
<b>BISOPROLOL FUMARATE</b>				
* Tab 2.5 mg .....	3.53	90	✓	<b>Bosvate</b>
* Tab 5 mg .....	5.15	90	✓	<b>Bosvate</b>
* Tab 10 mg .....	9.40	90	✓	<b>Bosvate</b>
<b>CARVEDILOL</b>				
* Tab 6.25 mg .....	2.24	60	✓	<b>Carvedilol Sandoz</b>
* Tab 12.5 mg .....	2.30	60	✓	<b>Carvedilol Sandoz</b>
* Tab 25 mg .....	2.95	60	✓	<b>Carvedilol Sandoz</b>
<b>CELIPROLOL</b>				
* Tab 200 mg .....	21.40	180	✓	<b>Celol</b>
<b>LABETALOL</b>				
Tab 100 mg .....	11.36	100	✓	<b>Hybloc</b>
			✓	<b>Presolol</b> <sup>\$29</sup>
Tab 200 mg .....	29.74	100	✓	<b>Hybloc</b>
			✓	<b>Presolol</b> <sup>\$29</sup>
* Inj 5 mg per ml, 20 ml ampoule .....	59.06 (88.60)	5		Trandate

(Hybloc Tab 100 mg to be delisted 1 December 2019)

(Hybloc Tab 200 mg to be delisted 1 February 2020)



	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>METOPROLOL SUCCINATE</b>				
* Tab long-acting 23.75 mg.....	1.03	30	✓	<b>Betaloc CR</b>
* Tab long-acting 47.5 mg.....	1.25	30	✓	<b>Betaloc CR</b>
* Tab long-acting 95 mg.....	1.99	30	✓	<b>Betaloc CR</b>
* Tab long-acting 190 mg.....	3.00	30	✓	<b>Betaloc CR</b>
<b>METOPROLOL TARTRATE</b>				
* Tab 50 mg.....	5.66	100	✓	<b>Apo-Metoprolol</b>
* Tab 100 mg.....	7.55	60	✓	<b>Apo-Metoprolol</b>
* Tab long-acting 200 mg.....	23.40	28	✓	<b>Slow-Lopresor</b>
* Inj 1 mg per ml, 5 ml vial.....	29.50	5	✓	<b>Metoprolol IV</b> <b>Mylan</b>
<b>NADOLOL</b>				
* Tab 40 mg.....	16.69	100	✓	<b>Apo-Nadolol</b>
* Tab 80 mg.....	26.43	100	✓	<b>Apo-Nadolol</b>
<b>PINDOLOL</b>				
* Tab 5 mg.....	13.22	100	✓	<b>Apo-Pindolol</b>
* Tab 10 mg.....	23.12	100	✓	<b>Apo-Pindolol</b>
* Tab 15 mg.....	33.31	100	✓	<b>Apo-Pindolol</b>
<b>PROPRANOLOL</b>				
* Tab 10 mg.....	4.64	100	✓	<b>Apo-Propranolol</b>
* Tab 40 mg.....	5.72	100	✓	<b>Apo-Propranolol</b>
* Cap long-acting 160 mg.....	18.17	100	✓	<b>Cardinol LA</b>
* Oral liq 4 mg per ml – Special Authority see SA1327 below – Retail pharmacy.....	CBS	500 ml	✓	<b>Roxane S29</b>
<b>►SA1327 Special Authority for Subsidy</b>				
<b>Initial application</b> from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:				
Either:				
1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or				
2 For the treatment of a child under 12 years with cardiac arrhythmias or congenital cardiac abnormalities.				
<b>Renewal</b> from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:				
Either:				
1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or				
2 For the treatment of a child under 12 years with cardiac arrhythmias or congenital cardiac abnormalities.				
<b>SOTALOL</b>				
* Tab 80 mg.....	32.58	500	✓	<b>Mylan</b>
* Tab 160 mg.....	10.98	100	✓	<b>Mylan</b>
<b>TIMOLOL</b>				
* Tab 10 mg.....	10.55	100	✓	<b>Apo-Timol</b>

## Calcium Channel Blockers

### Dihydropyridine Calcium Channel Blockers

#### AMLODIPINE

* Tab 2.5 mg.....	1.72	100	✓	<b>Apo-Amlodipine</b>
* Tab 5 mg.....	3.33	250	✓	<b>Apo-Amlodipine</b>
* Tab 10 mg.....	4.40	250	✓	<b>Apo-Amlodipine</b>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

# CARDIOVASCULAR SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>FELODIPINE</b>				
* Tab long-acting 2.5 mg.....	1.45	30	✓	<b>Plendil ER</b>
* Tab long-acting 5 mg.....	3.93	90	✓	<b>Felo 5 ER</b>
* Tab long-acting 10 mg.....	4.32	90	✓	<b>Felo 10 ER</b>
<b>NIFEDIPINE</b>				
* Tab long-acting 10 mg.....	10.63	60	✓	<b>Adalat 10</b>
			✓	<b>Adefin</b> <sup>S29</sup>
* Tab long-acting 20 mg.....	9.59	100	✓	<b>Nyefax Retard</b>
* Tab long-acting 30 mg.....	3.14	30	✓	<b>Adalat Oros</b>
			✓	<b>Adefin XL</b>
* Tab long-acting 60 mg.....	5.67	30	✓	<b>Adalat Oros</b>
			✓	<b>Adefin XL</b>

(Adefin XL Tab long-acting 30 mg to be delisted 1 March 2020)

## Other Calcium Channel Blockers

<b>DILTIAZEM HYDROCHLORIDE</b>				
* Tab 30 mg .....	4.60	100	✓	<b>Dilzem</b>
* Tab 60 mg .....	8.50	100	✓	<b>Dilzem</b>
* Cap long-acting 120 mg .....	33.42	500	✓	<b>Apo-Diltiazem CD</b>
* Cap long-acting 180 mg .....	50.05	500	✓	<b>Apo-Diltiazem CD</b>
* Cap long-acting 240 mg .....	66.76	500	✓	<b>Apo-Diltiazem CD</b>
<b>PERHEXILINE MALEATE</b>				
* Tab 100 mg .....	62.90	100	✓	<b>Pexsig</b>
<b>VERAPAMIL HYDROCHLORIDE</b>				
* Tab 40 mg .....	7.01	100	✓	<b>Isoptin</b>
* Tab 80 mg .....	11.74	100	✓	<b>Isoptin</b>
* Tab long-acting 120 mg.....	15.20	250	✓	<b>Verpamil SR</b>
	36.02	100	✓	<b>Isoptin SR</b>
* Tab long-acting 240 mg.....	25.00	250	✓	<b>Verpamil SR</b>
* Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO .....	25.00	5	✓	<b>Isoptin</b>

(Verpamil SR Tab long-acting 120 mg to be delisted 1 May 2020)

## Centrally-Acting Agents

<b>CLONIDINE</b>				
* Patch 2.5 mg, 100 mcg per day – Only on a prescription.....	7.40	4	✓	<b>Mylan</b>
* Patch 5 mg, 200 mcg per day – Only on a prescription.....	10.04	4	✓	<b>Mylan</b>
* Patch 7.5 mg, 300 mcg per day – Only on a prescription.....	12.34	4	✓	<b>Mylan</b>
<b>CLONIDINE HYDROCHLORIDE</b>				
* Tab 25 mcg.....	8.75	112	✓	<b>Clonidine BNM</b>
* Tab 150 mcg.....	34.32	100	✓	<b>Catapres</b>
* Inj 150 mcg per ml, 1 ml ampoule .....	25.96	10	✓	<b>Medsurge</b>
<b>METHYLDOPA</b>				
* Tab 250 mg .....	15.10	100	✓	<b>Methyldopa Mylan</b>
	52.85	500	✓	<b>Methyldopa Mylan</b>
				<b>S29</b> <sup>S29</sup>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>Diuretics</b>				
<b>Loop Diuretics</b>				
<b>BUMETANIDE</b>				
* Tab 1 mg .....	16.36	100	✓	Burinex
* Inj 500 mcg per ml, 4 ml vial .....	7.95	5	✓	Burinex
<b>FUROSEMIDE [FRUSEMIDE]</b>				
Tab 40 mg – Up to 30 tab available on a PSO .....	7.24	1,000	✓	Apo-Furosemide
	8.00		✓	Diurin 40
	20.40		✓	Milan
Laboratories <sup>S29</sup>				
Note: Wastage may only be claimed once on Milan Laboratories.				
* Tab 500 mg .....	25.00	50	✓	Urex Forte
* Oral liq 10 mg per ml .....	11.20	30 ml OP	✓	Lasix
Lasix to be Sole Supply on 1 January 2020				
* Inj 10 mg per ml, 25 ml ampoule .....	60.65	6	✓	Lasix
Lasix to be Sole Supply on 1 January 2020				
* Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO .....	1.15	5	✓	Frusemide-Clarix
(Milan Laboratories <sup>S29</sup> Tab 40 mg to be delisted 1 November 2019)				
<b>Potassium Sparing Diuretics</b>				
<b>AMILORIDE HYDROCHLORIDE</b>				
Oral liq 1 mg per ml .....	30.00	25 ml OP	✓	Biomed
<b>EPLERENONE – Special Authority see SA1728 below – Retail pharmacy</b>				
Tab 50 mg .....	17.00	30	✓	Inspra
Tab 25 mg .....	11.87	30	✓	Inspra
<b>SA1728 Special Authority for Subsidy</b>				
<b>Initial application</b> from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:				
Both:				
1 Patient has heart failure with ejection fraction less than 40%; and				
2 Either:				
2.1 Patient is intolerant to optimal dosing of spironolactone; or				
2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone.				
<b>METOLAZONE</b>				
Tab 5 mg .....	CBS	1	✓	Metolazone <sup>S29</sup>
		50	✓	Zaroxolyn <sup>S29</sup>
<b>SPIRONOLACTONE</b>				
* Tab 25 mg .....	4.38	100	✓	Spiractin
* Tab 100 mg .....	11.80	100	✓	Spiractin
Oral liq 5 mg per ml .....	30.60	25 ml OP	✓	Biomed
Biomed to be Sole Supply on 1 November 2019				
<b>Potassium Sparing Combination Diuretics</b>				
<b>AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE</b>				
* Tab 5 mg with furosemide 40 mg .....	8.63	28	✓	Frumil

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE</b>				
* Tab 5 mg with hydrochlorothiazide 50 mg.....	5.00	50	✓	<b>Moduretic</b>

## Thiazide and Related Diuretics

<b>BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]</b>				
* Tab 2.5 mg – Up to 150 tab available on a PSO .....	12.50	500	✓	<b>Arrow- Bendroflumethiazide</b>
May be supplied on a PSO for reasons other than emergency.				
* Tab 5 mg .....	20.42	500	✓	<b>Arrow- Bendroflumethiazide</b>

<b>CHLOROTHIAZIDE</b>				
Oral liq 50 mg per ml .....	26.00	25 ml OP	✓	<b>Biomed</b>
<b>CHLORTALIDONE [CHLORTHALIDONE]</b>				
* Tab 25 mg .....	6.50	50	✓	<b>Hygroton</b>
Hygroton to be Sole Supply on 1 December 2019				
<b>INDAPAMIDE</b>				
* Tab 2.5 mg .....	2.60	90	✓	<b>Dapa-Tabs</b>

## Lipid-Modifying Agents

### Fibrates

<b>BEZAFIBRATE</b>				
* Tab 200 mg .....	19.01	90	✓	<b>Bezalip</b>
* Tab long-acting 400 mg.....	12.89	30	✓	<b>Bezalip Retard</b>
<b>GEMFIBROZIL</b>				
* Tab 600 mg .....	19.56	60	✓	<b>Lipazil</b>

### Other Lipid-Modifying Agents

<b>ACIPIMOX</b>				
* Cap 250 mg.....	18.75	30	✓	<b>Olbetam</b>
<b>NICOTINIC ACID</b>				
* Tab 50 mg .....	4.12	100	✓	<b>Apo-Nicotinic Acid</b>
* Tab 500 mg .....	17.89	100	✓	<b>Apo-Nicotinic Acid</b>

### Resins

<b>COLESTIPOL HYDROCHLORIDE</b>				
Grans for oral liq 5 g .....	28.60	30	✓	<b>Colestid</b>

## HMG CoA Reductase Inhibitors (Statins)

### Prescribing Guidelines

Treatment with HMG CoA Reductase Inhibitors (statins) is recommended for patients with dyslipidaemia and an absolute 5 year cardiovascular risk of 15% or greater.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
ATORVASTATIN – See prescribing guideline <a href="#">on the previous page</a>				
* Tab 10 mg .....	6.96	500	✓	<b>Lorstat</b>
* Tab 20 mg .....	9.99	500	✓	<b>Lorstat</b>
* Tab 40 mg .....	15.93	500	✓	<b>Lorstat</b>
* Tab 80 mg .....	27.19	500	✓	<b>Lorstat</b>
PRAVASTATIN – See prescribing guideline <a href="#">on the previous page</a>				
* Tab 20 mg .....	4.72	100	✓	<b>Apo-Pravastatin</b>
* Tab 40 mg .....	8.06	100	✓	<b>Apo-Pravastatin</b>
SIMVASTATIN – See prescribing guideline <a href="#">on the previous page</a>				
* Tab 10 mg .....	0.95	90	✓	<b>Simvastatin Mylan</b>
* Tab 20 mg .....	1.52	90	✓	<b>Simvastatin Mylan</b>
* Tab 40 mg .....	2.63	90	✓	<b>Simvastatin Mylan</b>
* Tab 80 mg .....	6.00	90	✓	<b>Simvastatin Mylan</b>

## Selective Cholesterol Absorption Inhibitors

EZETIMIBE – Special Authority see [SA1045 below](#) – Retail pharmacy

* Tab 10 mg .....	2.00	30	✓	<b>Ezetimibe Sandoz</b>
-------------------	------	----	---	-------------------------

### ►SA1045 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
  - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 × normal) when treated with one statin; or
  - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
  - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN – Special Authority see [SA1046 below](#) – Retail pharmacy

Tab 10 mg with simvastatin 10 mg .....	5.15	30	✓	<b>Zimybe</b>
Tab 10 mg with simvastatin 20 mg .....	6.15	30	✓	<b>Zimybe</b>
Tab 10 mg with simvastatin 40 mg .....	7.15	30	✓	<b>Zimybe</b>
Tab 10 mg with simvastatin 80 mg .....	8.15	30	✓	<b>Zimybe</b>

### ►SA1046 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

## Nitrates

### GLYCERYL TRINITRATE

* Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO .....	4.45	250 dose OP	✓ Nitrolingual Pump Spray
* Oral spray, 400 mcg per dose – Up to 200 dose available on a PSO .....	4.45	200 dose OP	✓ Glytrin
* Patch 25 mg, 5 mg per day .....	15.73	30	✓ Nitroderm TTS
* Patch 50 mg, 10 mg per day .....	18.62	30	✓ Nitroderm TTS

(Glytrin Oral spray, 400 mcg per dose to be delisted 1 May 2020)

### ISOSORBIDE MONONITRATE

* Tab 20 mg .....	18.80	100	✓ Ismo 20
* Tab long-acting 40 mg.....	7.50	30	✓ Ismo 40 Retard
* Tab long-acting 60 mg.....	8.29	90	✓ Duride

## Sympathomimetics

### ADRENALINE

Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PSO .....	4.98	5	✓ Aspen Adrenaline
	5.25		✓ Hospira
Inj 1 in 10,000, 10 ml ampoule – Up to 5 inj available on a PSO .....	27.00	5	✓ Hospira
	49.00	10	✓ Aspen Adrenaline

### ISOPRENALINE [ISOPROTERENOL]

* Inj 200 mcg per ml, 1 ml ampoule .....	36.80	25	
	(164.20)		Isuprel

## Vasodilators

### HYDRALAZINE HYDROCHLORIDE

* Tab 25 mg – Special Authority see <a href="#">SA1321 on the next page</a> – Retail pharmacy.....	CBS	1	✓ Hydralazine
		56	✓ Onelink <sup>S29</sup>
		84	✓ AMDIPHARM <sup>S29</sup>
		100	✓ Onelink <sup>S29</sup>
* Inj 20 mg ampoule.....	25.90	5	✓ Apresoline

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

►SA1321 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 For the treatment of refractory hypertension; or
- 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

MINOXIDIL

▲ Tab 10 mg ..... 70.00 100 ✓ **Loniten**

NICORANDIL

▲ Tab 10 mg ..... 25.57 60 ✓ **Ikorel**

Ikorel to be Sole Supply on 1 December 2019

▲ Tab 20 mg ..... 32.28 60 ✓ **Ikorel**

Ikorel to be Sole Supply on 1 December 2019

PAPAVERINE HYDROCHLORIDE

\* Inj 12 mg per ml, 10 ml ampoule ..... 217.90 5 ✓ **Hospira**

PENTOXIFYLLINE [OXPENTIFYLLINE]

Tab 400 mg ..... 42.26 50 ✓ **Trental 400**

**Endothelin Receptor Antagonists**

AMBRISENTAN – Special Authority see SA1702 below – Retail pharmacy

Tab 5 mg ..... 4,585.00 30 ✓ **Volibris**

Tab 10 mg ..... 4,585.00 30 ✓ **Volibris**

►SA1702 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website <http://www.pharmac.govt.nz> or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: [PAH@pharmac.govt.nz](mailto:PAH@pharmac.govt.nz)

BOSENTAN – Special Authority see SA1712 below – Retail pharmacy

Tab 62.5 mg ..... 141.00 60 ✓ **Bosentan Dr**

**Reddy's**

Tab 125 mg ..... 141.00 60 ✓ **Bosentan Dr**

**Reddy's**

►SA1712 Special Authority for Subsidy

**Initial application** only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:

4.1 Both:

4.1.1 Bosentan is to be used as PAH monotherapy; and

4.1.2 Either:

continued...

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\* Three months or six months, as applicable, dispensed all-at-once

continued...

- 4.1.2.1 Patient is intolerant or contraindicated to sildenafil; or
  - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
  - 4.2 Both:
    - 4.2.1 Bosentan is to be used as PAH dual therapy; and
    - 4.2.2 Either:
      - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
      - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
  - 4.3 Both:
    - 4.3.1 Bosentan is to be used as PAH triple therapy; and
    - 4.3.2 Any of the following:
      - 4.3.2.1 Patient is on the lung transplant list; or
      - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
      - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
      - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.
- Renewal** only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:
- Any of the following:
- 1 Both:
    - 1.1 Bosentan is to be used as PAH monotherapy; and
    - 1.2 Patient is stable or has improved while on bosentan; or
  - 2 Both:
    - 2.1 Bosentan is to be used as PAH dual therapy; and
    - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
  - 3 Both:
    - 3.1 Bosentan is to be used as PAH triple therapy; and
    - 3.2 Any of the following:
      - 3.2.1 Patient is on the lung transplant list; or
      - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
      - 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
      - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

## Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see [SA1825 below](#) – Retail pharmacy

Tab 25 mg .....	0.64	4	✓ <u>Vedafil</u>
Tab 50 mg .....	0.64	4	✓ <u>Vedafil</u>
Tab 100 mg .....	6.60	12	✓ <u>Vedafil</u>

➔ [SA1825](#) **Special Authority for Subsidy**

**Initial application — (Raynaud's Phenomenon\*)** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

continued...



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

continued...

All of the following:

- 1 Patient has Raynaud's Phenomenon\*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

**Initial application — (Pulmonary arterial hypertension\*)** only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 Any of the following:
  - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
  - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
  - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
  - 3.1 PAH is in NYHA/WHO functional class II; or
  - 3.2 PAH is in NYHA/WHO functional class III; or
  - 3.3 PAH is in NYHA/WHO functional class IV; and
- 4 Either:
  - 4.1 All of the following:
    - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.2 Either:
      - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
      - 4.1.2.2 Patient is peri Fontan repair; and
    - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm<sup>-5</sup>); or
  - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with \* are unapproved indications.

**Initial application — (erectile dysfunction due to spinal cord injury)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

**Renewal — (erectile dysfunction due to spinal cord injury)** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

## Prostacyclin Analogues

EPOPROSTENOL – Special Authority see [SA1696 below](#) – Retail pharmacy

Inj 500 mcg vial.....	36.61	1	✓ Veletri
Inj 1.5 mg vial .....	73.21	1	✓ Veletri

### ►SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website <http://www.pharmac.govt.nz> or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: [PAH@pharmac.govt.nz](mailto:PAH@pharmac.govt.nz)

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
ILOPROST – Special Authority see <a href="#">SA1705 below</a> – Retail pharmacy				
Nebuliser soln 10 mcg per ml, 2 ml .....	740.10	30	✓	Ventavis
Ventavis to be Sole Supply on 1 January 2020				

## ► [SA1705](#) Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website <http://www.pharmac.govt.nz> or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: [PAH@pharmac.govt.nz](mailto:PAH@pharmac.govt.nz)

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, [page 89](#)

### ADAPALENE

- Maximum of 30 g per prescription
- Only on a prescription

Crm 0.1%.....	22.89	30 g OP	✓ Differin
Gel 0.1%.....	22.89	30 g OP	✓ Differin

### ISOTRETINOIN – Special Authority see [SA1475 below](#) – Retail pharmacy

Cap 5 mg.....	8.14	60	✓ Oratane
Cap 10 mg.....	13.34	120	✓ Oratane
Cap 20 mg.....	20.49	120	✓ Oratane

### ►SA1475 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- Either:
  - Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
  - Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

**Renewal** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
- Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

### TRETINOIN

Crm 0.5 mg per g – Maximum of 50 g per prescription .....	13.90	50 g OP	✓ ReTrieve
---	-------	---------	------------

## Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, [page 89](#)

### HYDROGEN PEROXIDE

* Crm 1%.....	8.56	10 g OP	✓ Crystaderm
		15 g OP	✓ Crystaderm

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>MUPIROCI</b>				
Oint 2%.....	6.60 (9.26)	15 g OP		Bactroban
a) Only on a prescription				
b) Not in combination				
<b>SODIUM FUSIDATE [FUSIDIC ACID]</b>				
Crm 2%.....	1.59	5 g OP	✓ <b>Foban</b>	
a) Maximum of 5 g per prescription				
b) Only on a prescription				
c) Not in combination				
Oint 2%.....	1.59	5 g OP	✓ <b>Foban</b>	
a) Maximum of 5 g per prescription				
b) Only on a prescription				
c) Not in combination				
<b>SULFADIAZINE SILVER</b>				
Crm 1%.....	10.80	50 g OP	✓ <b>Flamazine</b>	
a) Up to 250 g available on a PSO				
b) Not in combination				

## Antifungals Topical

For systemic antifungals, refer to INFECTIONS, Antifungals, [page 96](#)

### AMOROLFINE

- a) Only on a prescription
- b) Not in combination

Nail soln 5%.....	15.95	5 ml OP	✓ <b>MycoNail</b>
-------------------	-------	---------	-------------------

### CICLOPIROX OLAMINE

- a) Only on a prescription
- b) Not in combination

Nail-soln 8%.....	5.72	7 ml OP	✓ <b>Apo-Ciclopirox</b>
-------------------	------	---------	-------------------------

### CLOTRIMAZOLE

* Crm 1%.....	0.70	20 g OP	✓ <b>Clomazol</b>
---------------	------	---------	-------------------

- a) Only on a prescription
- b) Not in combination

* Soln 1%.....	4.36 (7.55)	20 ml OP	Canesten
----------------	----------------	----------	----------

- a) Only on a prescription
- b) Not in combination

### ECONAZOLE NITRATE

Crm 1%.....	1.00 (7.48)	20 g OP	Pevaryl
-------------	----------------	---------	---------

- a) Only on a prescription
- b) Not in combination

Foaming soln 1%, 10 ml sachets.....	9.89 (17.23)	3	Pevaryl
-------------------------------------	-----------------	---	---------

- a) Only on a prescription
- b) Not in combination

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>MICONAZOLE NITRATE</b>				
* Crm 2%.....	0.74	15 g OP	✓	<b>Multichem</b>
a) Only on a prescription				
b) Not in combination				
* Lotn 2% .....	4.36 (10.03)	30 ml OP		Daktarin
a) Only on a prescription				
b) Not in combination				
* Tinct 2%.....	4.36 (12.10)	30 ml OP		Daktarin
a) Only on a prescription				
b) Not in combination				
<b>NYSTATIN</b>				
Crm 100,000 u per g.....	1.00 (7.90)	15 g OP		Mycostatin
a) Only on a prescription				
b) Not in combination				

### Antipruritic Preparations

#### CALAMINE

a) Only on a prescription				
b) Not in combination				
Crm, aqueous, BP .....	1.26	100 g	✓	<b>healthE Calamine Aqueous Cream BP</b>
Lotn, BP .....	12.94	2,000 ml	✓	<b>PSM</b>
<i>(PSM Lotn, BP to be delisted 1 July 2020)</i>				

#### CROTAMITON

a) Only on a prescription				
b) Not in combination				
Crm 10%.....	3.29	20 g OP	✓	<b>Itch-Soothe</b>

#### MENTHOL – Only in combination

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteroid – Plain
- 2) With or without other dermatological galenicals.

Crystals.....	6.92 29.60	25 g 100 g	✓	<b>MidWest</b> <b>MidWest</b>
---------------	---------------	---------------	---	----------------------------------

## Corticosteroids Topical

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 79

### Corticosteroids - Plain

#### BETAMETHASONE DIPROPIONATE

Crm 0.05%.....	2.96	15 g OP	✓ <b>Diprosone</b>
	8.97	50 g OP	✓ <b>Diprosone</b>
Crm 0.05% in propylene glycol base .....	4.33	30 g OP	✓ <b>Diprosone OV</b>
Oint 0.05% .....	2.96	15 g OP	✓ <b>Diprosone</b>
	8.97	50 g OP	✓ <b>Diprosone</b>
Oint 0.05% in propylene glycol base .....	4.33	30 g OP	✓ <b>Diprosone OV</b>

*(Diprosone OV Crm 0.05% in propylene glycol base to be delisted 1 May 2020)*

#### BETAMETHASONE VALERATE

* Crm 0.1%.....	3.45	50 g OP	✓ <b>Beta Cream</b>
* Oint 0.1%.....	3.45	50 g OP	✓ <b>Beta Ointment</b>
* Lotn 0.1% .....	18.00	50 ml OP	✓ <b>Betnovate</b>

#### CLOBETASOL PROPIONATE

* Crm 0.05%.....	2.18	30 g OP	✓ <b>Dermol</b>
Dermol to be Sole Supply on 1 November 2019			
* Oint 0.05%.....	2.12	30 g OP	✓ <b>Dermol</b>
Dermol to be Sole Supply on 1 November 2019			

#### CLOBETASONE BUTYRATE

Crm 0.05%.....	5.38	30 g OP	
	(7.09)		Eumovate

#### DIFLUCORTOLONE VALERATE

Crm 0.1%.....	8.97	50 g OP	
	(15.86)		Nerisone
Fatty oint 0.1%.....	8.97	50 g OP	
	(15.86)		Nerisone

#### HYDROCORTISONE

* Crm 1% – Only on a prescription.....	1.11	30 g OP	✓ <b>DermAssist</b>
	16.25	500 g	✓ <b>Pharmacy Health</b>
* Powder – Only in combination.....	49.95	25 g	✓ <b>ABM</b>
Up to 5% in a dermatological base (not proprietary Topical Corticosteroid – Plain) with or without other dermatological galenicals			

#### HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN

Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only on a prescription .....	10.57	250 ml	✓ <b>DP Lotn HC</b>
--	-------	--------	---------------------

#### HYDROCORTISONE BUTYRATE

Lipocream 0.1%.....	3.42	30 g OP	✓ <b>Locoid Lipocream</b>
	6.85	100 g OP	✓ <b>Locoid Lipocream</b>
Oint 0.1%.....	13.70	100 g OP	✓ <b>Locoid</b>
Milky emul 0.1% .....	13.70	100 ml OP	✓ <b>Locoid Crelo</b>

#### METHYLPREDNISOLONE ACEPONATE

Crm 0.1%.....	4.95	15 g OP	✓ <b>Advantan</b>
Oint 0.1%.....	4.95	15 g OP	✓ <b>Advantan</b>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>MOMETASONE FUROATE</b>				
Crm 0.1%.....	1.51	15 g OP	✓	<b>Elocon Alcohol Free</b>
	2.50	50 g OP	✓	<b>Elocon Alcohol Free</b>
Oint 0.1%.....	1.51	15 g OP	✓	<b>Elocon</b>
	2.90	50 g OP	✓	<b>Elocon</b>
Lotn 0.1% .....	6.30	30 ml OP	✓	<b>Elocon</b>
<b>TRIAMCINOLONE ACETONIDE</b>				
Crm 0.02%.....	6.30	100 g OP	✓	<b>Aristocort</b>
Oint 0.02%.....	6.35	100 g OP	✓	<b>Aristocort</b>

### Corticosteroids - Combination

<b>BETAMETHASONE VALERATE WITH CLIOQUINOL – Only on a prescription</b>				
Crm 0.1% with clioquinol 3%.....	3.49 (4.90)	15 g OP		Betnovate-C
<b>BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUSIDIC ACID]</b>				
Crm 0.1% with sodium fusidate (fusidic acid) 2%.....	3.49 (10.45)	15 g OP		Fucicort
a) Maximum of 15 g per prescription				
b) Only on a prescription				
<b>HYDROCORTISONE WITH MICONAZOLE – Only on a prescription</b>				
* Crm 1% with miconazole nitrate 2%.....	2.00	15 g OP	✓	<b>Micreme H</b>
<b>HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN – Only on a prescription</b>				
Crm 1% with natamycin 1% and neomycin sulphate 0.5% .....	3.35	15 g OP	✓	<b>Pimafucort</b>
Oint 1% with natamycin 1% and neomycin sulphate 0.5%.....	3.35	15 g OP	✓	<b>Pimafucort</b>
<b>TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCIN AND NYSTATIN</b>				
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g – Only on a prescription .....	3.49 (6.60)	15 g OP		Viaderm KC

### Disinfecting and Cleansing Agents

<b>CHLORHEXIDINE GLUCONATE – Subsidy by endorsement</b>				
a) No more than 500 ml per month				
b) Only if prescribed for a dialysis patient and the prescription is endorsed accordingly.				
* Handrub 1% with ethanol 70% .....	4.29	500 ml	✓	<b>healthE</b>
* Soln 4% wash.....	3.98	500 ml	✓	<b>healthE</b>
<b>TRICLOSAN – Subsidy by endorsement</b>				
a) Maximum of 500 ml per prescription				
b)				
a) Only if prescribed for a patient identified with Methicillin-resistant Staphylococcus aureus (MRSA) prior to elective surgery in hospital and the prescription is endorsed accordingly; or				
b) Only if prescribed for a patient with recurrent Staphylococcus aureus infection and the prescription is endorsed accordingly				
Soln 1% .....	5.90	500 ml OP	✓	<b>healthE</b>

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\* Three months or six months, as applicable, dispensed all-at-once

## Barrier Creams and Emollients

### Barrier Creams

#### DIMETHICONE

* Crm 5% pump bottle.....	4.48	500 ml OP	✓ <u>healthE</u> <u>Dimethicone 5%</u>
* Crm 10% pump bottle.....	4.52	500 ml OP	✓ <u>healthE</u> <u>Dimethicone 10%</u>

#### ZINC AND CASTOR OIL

* Oint.....	4.25	500 g	✓ <u>Boucher</u>
-------------	------	-------	------------------

### Emollients

#### AQUEOUS CREAM

* Crm.....	1.92	500 g	✓ <u>Boucher</u>
------------	------	-------	------------------

#### CETOMACROGOL

* Crm BP.....	2.48	500 g	✓ <u>healthE</u>
---------------	------	-------	------------------

#### CETOMACROGOL WITH GLYCEROL

Crm 90% with glycerol 10%.....	2.35	500 ml OP	✓ <u>Boucher</u>
	2.82		✓ <u>Pharmacy Health</u> <u>Sorbolene with</u> <u>Glycerin</u>
	3.10	1,000 ml OP	✓ <u>Boucher</u>
	3.87		✓ <u>Pharmacy Health</u> <u>Sorbolene with</u> <u>Glycerin</u>

(Pharmacy Health Sorbolene with Glycerin Crm 90% with glycerol 10% to be delisted 1 March 2020)

(Pharmacy Health Sorbolene with Glycerin Crm 90% with glycerol 10% to be delisted 1 March 2020)

#### EMULSIFYING OINTMENT

* Oint BP.....	3.59	500 g	✓ <u>AFT</u>
----------------	------	-------	--------------

#### OIL IN WATER EMULSION

* Crm.....	2.19	500 g	✓ <u>O/W Fatty Emulsion</u> <u>Cream</u>
------------	------	-------	---

#### PARAFFIN

Oint liquid paraffin 50% with white soft paraffin 50%.....	5.35	500 ml OP	✓ <u>healthE</u>
--	------	-----------	------------------

#### UREA

* Crm 10%.....	1.37	100 g OP	✓ <u>healthE Urea Cream</u>
----------------	------	----------	-----------------------------

#### WOOL FAT WITH MINERAL OIL – Only on a prescription

* Lotn hydrous 3% with mineral oil.....	5.60	1,000 ml	
	(11.95)		DP Lotion
	1.40	250 ml OP	
	(4.53)		DP Lotion
	5.60	1,000 ml	
	(20.53)		Alpha-Keri Lotion
	(23.91)		BK Lotion
	1.40	250 ml OP	
	(7.73)		BK Lotion



Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

## Other Dermatological Bases

### PARAFFIN

White soft – Only in combination .....	20.20	2,500 g	✓ IPW
	3.58	500 g	
	(7.78)		IPW
	(8.69)		PSM

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid – Plain.  
(PSM White soft to be delisted 1 May 2020)

## Minor Skin Infections

### POVIDONE IODINE

Oint 10% .....	3.27	25 g OP	✓ Betadine
a) Maximum of 100 g per prescription			
b) Only on a prescription			
Antiseptic soln 10% .....	2.55	100 ml	✓ Riodine
	3.83	15 ml	✓ Riodine
	5.40	500 ml	✓ Riodine
	6.20		✓ Betadine
	1.28	100 ml	
	(13.27)		Betadine
	0.19	15 ml	
	(7.41)		Betadine
Skin preparation, povidone iodine 10% with 30% alcohol .....	10.00	500 ml	✓ Betadine Skin Prep
	1.63	100 ml	
	(3.48)		Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol .....	1.63	100 ml	
	(6.64)		Pfizer

(Betadine Antiseptic soln 10% to be delisted 1 February 2020)

(Betadine Antiseptic soln 10% to be delisted 1 February 2020)

(Betadine Antiseptic soln 10% to be delisted 1 February 2020)

## Parasiticial Preparations

### DIMETHICONE

* Lotn 4% .....	4.98	200 ml OP	✓ <u>healthE</u> <u>Dimethicone 4%</u> <u>Lotion</u>
-----------------	------	-----------	--

IVERMECTIN – Special Authority see [SA1225 on the next page](#) – Retail pharmacy

Tab 3 mg – Up to 100 tab available on a PSO .....	17.20	4	✓ Stromectol
---	-------	---	--------------

- 1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- 3) For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## ►SA1225 Special Authority for Subsidy

**Initial application — (Scabies)** from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 The patient is in the community; and
    - 2.1.2 Any of the following:
      - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
      - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
      - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
  - 2.2 All of the following:
    - 2.2.1 The Patient is a resident in an institution; and
    - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
    - 2.2.3 Any of the following:
      - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
      - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
      - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

**Initial application — (Other parasitic infections)** only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

**Renewal — (Scabies)** from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 The patient is in the community; and
    - 2.1.2 Any of the following:
      - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
      - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
      - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
  - 2.2 All of the following:
    - 2.2.1 The Patient is a resident in an institution; and
    - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
    - 2.2.3 Any of the following:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
- 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

**Renewal — (Other parasitic infections)** only from an infectious disease specialist, clinical microbiologist or dermatologist.

Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

#### PERMETHRIN

Crm 5% .....	4.95	30 g OP	✓ <b>Lyderm</b>
Lotn 5% .....	3.69	30 ml OP	✓ <b>A-Scabies</b>

#### PHENOTHRIN

Shampoo 0.5% .....	11.36	200 ml OP	✓ <b>Parasidose</b>
--------------------	-------	-----------	---------------------

### Psoriasis and Eczema Preparations

ACITRETIN – Special Authority see [SA1476 below](#) – Retail pharmacy

Cap 10 mg .....	17.86	60	✓ <b>Novatretin</b>
Cap 25 mg .....	41.36	60	✓ <b>Novatretin</b>

#### ► [SA1476](#) Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
  - 3.1 Patient is female and has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of two years after the completion of the treatment; or
  - 3.2 Patient is male.

**Renewal** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Patient is female and has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of two years after the completion of the treatment; or
- 2 Patient is male.

#### BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

Gel 500 mcg with calcipotriol 50 mcg per g .....	52.24	60 g OP	✓ <b>Daivobet</b>
Oint 500 mcg with calcipotriol 50 mcg per g .....	19.95	30 g OP	✓ <b>Daivobet</b>

#### CALCIPOTRIOL

Oint 50 mcg per g .....	45.00	100 g OP	✓ <b>Daivonex</b>
-------------------------	-------	----------	-------------------

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\* Three months or six months, as applicable, dispensed all-at-once

## DERMATOLOGICALS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>COAL TAR</b>				
Soln BP – Only in combination.....	36.25	200 ml	✓	<b>Midwest</b>
a)				
1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteroid – Plain				
2) With or without other dermatological galenicals.				
b) Midwest to be Sole Supply on 1 November 2019				
<b>COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR</b>				
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5%.....	6.59 (8.00) 3.43 (4.35)	75 g OP  30 g OP		Egopsoryl TA  Egopsoryl TA
<b>COAL TAR WITH SALICYLIC ACID AND SULPHUR</b>				
Soln 12% with salicylic acid 2% and sulphur 4% oint.....	4.97 7.95	25 g OP 40 g OP	✓ ✓	<b>Coco-Scalp</b> <b>Coco-Scalp</b>
<b>PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN – Only on a prescription</b>				
* Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.....	3.86	500 ml	✓	<b>Pinetarsol</b>
<b>SALICYLIC ACID</b>				
Powder – Only in combination.....	18.88	250 g	✓ ✓	<b>Midwest</b> <b>PSM</b>
1) Only in combination with a dermatological base or proprietary Topical Corticosteroid – Plain or collodion flexible				
2) With or without other dermatological galenicals.				
<b>SULPHUR</b>				
Precipitated – Only in combination.....	6.35	100 g	✓	<b>Midwest</b>
1) Only in combination with a dermatological base or proprietary Topical Corticosteroid – Plain				
2) With or without other dermatological galenicals.				

## Scalp Preparations

<b>BETAMETHASONE VALERATE</b>				
* Scalp app 0.1% .....	7.75	100 ml OP	✓	<b>Beta Scalp</b>
<b>CLOBETASOL PROPIONATE</b>				
* Scalp app 0.05% .....	5.69	30 ml OP	✓	<b>Dermol</b>
Dermol to be Sole Supply on 1 November 2019				
<b>HYDROCORTISONE BUTYRATE</b>				
Scalp lotn 0.1%.....	7.30	100 ml OP	✓	<b>Locoid</b>
<b>KETOCONAZOLE</b>				
Shampoo 2% .....	2.99	100 ml OP	✓	<b>Sebizole</b>
a) Maximum of 100 ml per prescription				
b) Only on a prescription				

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

## Sunscreens

SUNSCREENS, PROPRIETARY – Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

Crm.....	3.30 (5.89)	100 g OP	
Lotn.....	5.10	200 g OP	✓ Hamilton Sunscreen <b>Marine Blue Lotion SPF 50+</b>

(Hamilton Sunscreen Crm to be delisted 1 March 2020)

## Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, [page 67](#)

IMIQUIMOD

Crm 5%, 250 mg sachet.....	21.72	24	✓ <b>Perrigo</b>
----------------------------	-------	----	------------------

PODOPHYLLOTOXIN

Soln 0.5% .....	33.60	3.5 ml OP	✓ <b>Condyline</b>
a) Maximum of 3.5 ml per prescription			
b) Only on a prescription			

## Other Skin Preparations

### Antineoplastics

FLUOROURACIL SODIUM

Crm 5%.....	7.95	20 g OP	✓ <b>Efudix</b>
-------------	------	---------	-----------------

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\* Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

Subsidy  
(Manufacturer's Price)  
\$ Per Fully  
Subsidised  
✓ Brand or  
Generic  
Manufacturer

## Contraceptives - Non-hormonal

### Condoms

#### CONDOMS

* 49 mm – Up to 144 dev available on a PSO	11.42	144	✓ Moments
	13.36		✓ Shield 49
* 53 mm	0.95	10	✓ Moments
	1.11	12	✓ Gold Knight
	11.64	144	✓ Moments
	13.36		✓ Shield Blue
a) Maximum of 60 dev per prescription			
b) Up to 60 dev available on a PSO			
* 53 mm (chocolate)	13.36	144	✓ Gold Knight
a) Maximum of 60 dev per prescription			
b) Up to 60 dev available on a PSO			
* 53 mm (strawberry)	13.36	144	✓ Gold Knight
a) Maximum of 60 dev per prescription			
b) Up to 60 dev available on a PSO			
* 53 mm, 0.05 mm thickness	0.95	10	✓ Moments
	11.42	144	✓ Moments
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
* 53 mm, chocolate, brown	0.95	10	✓ Moments
	11.64	144	✓ Moments
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
* 53 mm, strawberry, red	0.95	10	✓ Moments
	11.64	144	✓ Moments
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
* 56 mm	0.97	10	✓ Moments
	11.64	144	✓ Moments
	13.36		✓ Durex Extra Safe
			✓ Gold Knight
a) Maximum of 60 dev per prescription			
b) Up to 60 dev available on a PSO			
* 56 mm, 0.05 mm thickness	1.30	12	✓ Gold Knight
	15.57	144	✓ Gold Knight
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
* 56 mm, 0.08 mm thickness	0.97	10	✓ Moments
	11.64	144	✓ Moments
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
* 56 mm, 0.08 mm thickness, red	0.97	10	✓ Moments
	11.64	144	✓ Moments
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
* 56 mm, chocolate	1.30	12	✓ Gold Knight
	15.57	144	✓ Gold Knight
a) Up to 60 dev available on a PSO			

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\* Three months or six months, as applicable, dispensed all-at-once

## GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
* 56 mm, shaped.....	13.36 (16.08)	144		Durex Confidence
a) Maximum of 60 dev per prescription				
b) Up to 60 dev available on a PSO				
* 56 mm, strawberry.....	1.30 15.57	12 144	✓ ✓	Gold Knight Gold Knight
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription				
* 60 mm – Up to 144 dev available on a PSO .....	13.36	144	✓	Shield XL
<i>(Shield 49 49 mm to be delisted 1 March 2020)</i>				
<i>(Gold Knight 53 mm to be delisted 1 March 2020)</i>				
<i>(Shield Blue 53 mm to be delisted 1 March 2020)</i>				
<i>(Gold Knight 53 mm (chocolate) to be delisted 1 March 2020)</i>				
<i>(Gold Knight 53 mm (strawberry) to be delisted 1 March 2020)</i>				
<i>(Durex Extra Safe 56 mm to be delisted 1 March 2020)</i>				
<i>(Gold Knight 56 mm to be delisted 1 March 2020)</i>				
<i>(Durex Confidence 56 mm, shaped to be delisted 1 March 2020)</i>				

### Contraceptive Devices

#### INTRA-UTERINE DEVICE

a) Up to 40 dev available on a PSO				
b) Only on a PSO				
* IUD 29.1 mm length × 23.2 mm width .....	18.45	1	✓	Choice TT380 Short
Choice TT380 Short to be Sole Supply on 1 November 2019				
* IUD 33.6 mm length × 29.9 mm width .....	18.45	1	✓	Choice TT380 Standard
Choice TT380 Standard to be Sole Supply on 1 November 2019				
* IUD 35.5 mm length × 19.6 mm width .....	15.50	1	✓	Choice Load 375
Choice Load 375 to be Sole Supply on 1 November 2019				

### Contraceptives - Hormonal

#### Combined Oral Contraceptives

#### ►SA0500 Special Authority for Alternate Subsidy

**Initial application** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Patient is on a Social Welfare benefit; or
- 1.2 Patient has an income no greater than the benefit; and

2 Has tried at least one of the fully funded options and has been unable to tolerate it.

**Renewal** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

continued...



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

#### ETHINYLOESTRADIOL WITH DESOGESTREL

* Tab 20 mcg with desogestrel 150 mcg and 7 inert tab.....	6.62	84	
	(19.80)		Mercilon 28
a) Higher subsidy of \$13.80 per 84 tab with Special Authority see <a href="#">SA0500 on the previous page</a>			
b) Up to 84 tab available on a PSO			
* Tab 30 mcg with desogestrel 150 mcg and 7 inert tab.....	6.62	84	
	(19.80)		Marvelon 28
a) Higher subsidy of \$13.80 per 84 tab with Special Authority see <a href="#">SA0500 on the previous page</a>			
b) Up to 84 tab available on a PSO			

#### ETHINYLOESTRADIOL WITH LEVONORGESTREL

* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets – Up to 112 tab available on a PSO .....	2.18	84	✓ <b>Microgynon 20 ED</b>
	6.45	112	✓ <b>Femme-Tab ED</b>
* Tab 50 mcg with levonorgestrel 125 mcg and 7 inert tab – Up to 84 tab available on a PSO .....	9.45	84	✓ <b>Microgynon 50 ED</b>
* Tab 30 mcg with levonorgestrel 150 mcg .....	6.62	63	
	(16.50)		Microgynon 30
a) Higher subsidy of \$15.00 per 63 tab with Special Authority see <a href="#">SA0500 on the previous page</a>			
b) Up to 63 tab available on a PSO			
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets – Up to 112 tab available on a PSO .....	1.77	84	✓ <b>Levien ED</b>
	6.45	112	✓ <b>Femme-Tab ED</b>

#### ETHINYLOESTRADIOL WITH NORETHISTERONE

* Tab 35 mcg with norethisterone 1 mg – Up to 63 tab available on a PSO .....	6.62	63	✓ <b>Brevinor 1/21</b>
* Tab 35 mcg with norethisterone 1 mg and 7 inert tab – Up to 84 tab available on a PSO .....	6.95	84	✓ <b>Brevinor 1/28</b>
* Tab 35 mcg with norethisterone 500 mcg – Up to 63 tab available on a PSO .....	6.62	63	✓ <b>Brevinor 21</b>
* Tab 35 mcg with norethisterone 500 mcg and 7 inert tab – Up to 84 tab available on a PSO .....	6.62	84	✓ <b>Norimin</b>

(Brevinor 1/21 Tab 35 mcg with norethisterone 1 mg to be delisted 1 July 2020)  
(Brevinor 21 Tab 35 mcg with norethisterone 500 mcg to be delisted 1 July 2020)

## Progestogen-only Contraceptives

### ►SA0500 Special Authority for Alternate Subsidy

**Initial application** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Patient is on a Social Welfare benefit; or
  - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

**Renewal** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

### LEVONORGESTREL

* Tab 30 mcg.....	6.62 (16.50)	84	
			Microlut
a) Higher subsidy of \$13.80 per 84 tab with Special Authority see SA0500 above			
b) Up to 84 tab available on a PSO			
* Subdermal implant (2 x 75 mg rods) – Up to 3 pack available on a PSO .....	106.92	1	✓ <b>Jadelle</b>
<b>MEDROXYPROGESTERONE ACETATE</b>			
Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PSO .....	7.98	1	✓ <b>Depo-Provera</b>
Depo-Provera to be Sole Supply on 1 December 2019			

### NORETHISTERONE

* Tab 350 mcg – Up to 84 tab available on a PSO .....	6.25	84	✓ <b>Noriday 28</b>
---	------	----	---------------------

## Emergency Contraceptives

### LEVONORGESTREL

* Tab 1.5 mg .....	4.95	1	✓ <b>Postinor-1</b>
a) Maximum of 2 tab per prescription			
b) Up to 5 tab available on a PSO			
c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.			

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

## Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

### CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

* Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO.....	4.67	168	✓ <u>Ginet</u>
---	------	-----	----------------

## Gynaecological Anti-infectives

### ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID

Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator....	8.43 (24.00)	100 g OP	Aci-Jel
---	-----------------	----------	---------

### CLOTRIMAZOLE

* Vaginal crm 1% with applicators.....	2.50	35 g OP	✓ <u>Clomazol</u>
Clomazol to be Sole Supply on 1 January 2020			
* Vaginal crm 2% with applicators.....	3.00	20 g OP	✓ <u>Clomazol</u>
Clomazol to be Sole Supply on 1 January 2020			

### MICONAZOLE NITRATE

* Vaginal crm 2% with applicator .....	3.88	40 g OP	✓ <u>Micreme</u>
--	------	---------	------------------

### NYSTATIN

Vaginal crm 100,000 u per 5 g with applicator(s) .....	4.45	75 g OP	✓ <u>Nilstat</u>
--	------	---------	------------------

## Myometrial and Vaginal Hormone Preparations

### ERGOMETRINE MALEATE

Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO .....	105.00	5	✓ <u>DBL Ergometrine</u>
---	--------	---	--------------------------

### OESTRIOL

* Crm 1 mg per g with applicator.....	6.62	15 g OP	✓ <u>Ovestin</u>
* Pessaries 500 mcg .....	6.86	15	✓ <u>Ovestin</u>

### OXYTOCIN – Up to 5 inj available on a PSO

Inj 5 iu per ml, 1 ml ampoule .....	3.98	5	✓ <u>Oxytocin BNM</u>
Inj 10 iu per ml, 1 ml ampoule .....	4.98	5	✓ <u>Oxytocin BNM</u>

### OXYTOCIN WITH ERGOMETRINE MALEATE – Up to 5 inj available on a PSO

Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml .....	15.00	5	✓ <u>Syntometrine</u>
--	-------	---	-----------------------

## Pregnancy Tests - hCG Urine

### PREGNANCY TESTS - HCG URINE

a) Up to 200 test available on a PSO			
b) Only on a PSO			
Cassette .....	12.00	40 test OP	✓ <u>Smith BioMed Rapid Pregnancy Test</u>

## Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, [page 107](#)

### 5-Alpha Reductase Inhibitors

FINASTERIDE – Special Authority see [SA0928 below](#) – Retail pharmacy

* Tab 5 mg .....	4.81	100	✓ <b>Ricit</b>
------------------	------	-----	----------------

➔ [SA0928](#) **Special Authority for Subsidy**

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
  - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
  - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

### Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE – Special Authority see [SA1032 below](#) – Retail pharmacy

* Cap 400 mcg .....	17.73	100	✓ <b>Tamsulosin-Rex</b>
---------------------	-------	-----	-------------------------

Tamsulosin-Rex to be Sole Supply on 1 January 2020

➔ [SA1032](#) **Special Authority for Subsidy**

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

### Other Urinary Agents

OXYBUTYNIN

* Tab 5 mg .....	8.85	500	✓ <b>Apo-Oxybutynin</b>
* Oral liq 5 mg per 5 ml .....	60.40	473 ml	✓ <b>Apo-Oxybutynin</b>

POTASSIUM CITRATE

Oral liq 3 mmol per ml – Special Authority see [SA1083 below](#) –

Retail pharmacy.....	31.80	200 ml OP	✓ <b>Biomed</b>
----------------------	-------	-----------	-----------------

➔ [SA1083](#) **Special Authority for Subsidy**

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE

* Grans eff 4 g sachets .....	2.34	28	✓ <b>Ural</b>
-------------------------------	------	----	---------------

SOLIFENACIN SUCCINATE

Tab 5 mg .....	3.00	30	✓ <b>Solifenacin Mylan</b>
Tab 10 mg .....	5.50	30	✓ <b>Solifenacin Mylan</b>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
TOLTERODINE – Special Authority see <a href="#">SA1272 below</a> – Retail pharmacy				
Tab 1 mg .....	14.56	56	✓	Arrow-Tolterodine
Tab 2 mg .....	14.56	56	✓	Arrow-Tolterodine

(Arrow-Tolterodine Tab 1 mg to be delisted 1 March 2020)

► [SA1272](#) **Special Authority for Subsidy**

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has overactive bladder and a documented intolerance of, or is non-responsive to oxybutynin.

## Detection of Substances in Urine

### ORTHO-TOLIDINE

* Compound diagnostic sticks.....	7.50 (8.25)	50 test OP	Hemastix
-----------------------------------	----------------	------------	----------

### TETRABROMOPHENOL

* Blue diagnostic strips.....	7.02 (13.92)	100 test OP	Albustix
-------------------------------	-----------------	-------------	----------

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

## Calcium Homeostasis

### CALCITONIN

\* Inj 100 iu per ml, 1 ml ampoule ..... 121.00      5      ✓ **Miacalcic**

CINACALCET – Special Authority see [SA1618 below](#) – Retail pharmacy

Tab 30 mg – Wastage claimable ..... 210.30      28      ✓ **Sensipar**

### ►SA1618 Special Authority for Subsidy

**Initial application** only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
  - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
  - 1.3 The patient is symptomatic; or
- 2 All of the following:
  - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriopathy); and
  - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
  - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

**Renewal** only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

### ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial – Special Authority see [SA1687 below](#) –

Retail pharmacy ..... 38.03      1      ✓ **Zoledronic acid**  
**Mylan**

### ►SA1687 Special Authority for Subsidy

**Initial application — (bone metastases)** only from an oncologist, haematologist or palliative care specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
  - 2.1 Patient has bone metastases or involvement; and
  - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
  - 3.1 Patient has bone metastases or involvement; and
  - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

**Initial application — (early breast cancer)** only from an oncologist or medical practitioner on the recommendation of a oncologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

continued...

# HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.

## Corticosteroids and Related Agents for Systemic Use

### BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE

* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml .....	19.20	5	
	(36.96)		
			Celestone Chronodose

### DEXAMETHASONE

* Tab 0.5 mg – Retail pharmacy-Specialist.....	0.99	30	✓ <u>Dexamethasone</u>
Up to 60 tab available on a PSO			
* Tab 4 mg – Retail pharmacy-Specialist.....	1.90	30	✓ <u>Dexamethasone</u>
Up to 30 tab available on a PSO			
Oral liq 1 mg per ml – Retail pharmacy-Specialist.....	45.00	25 ml OP	✓ <u>Biomed</u>
Oral liq prescriptions:			
1) Must be written by a Paediatrician or Paediatric Cardiologist; or			
2) On the recommendation of a Paediatrician or Paediatric Cardiologist.			

### DEXAMETHASONE PHOSPHATE

Dexamethasone phosphate injection will not be funded for oral use.

* Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO .....	14.19	10	✓ <u>Max Health</u>
* Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO .....	25.18	10	✓ <u>Max Health</u>

### FLUDROCORTISONE ACETATE

* Tab 100 mcg.....	14.32	100	✓ <u>Florinef</u>
--------------------	-------	-----	-------------------

### HYDROCORTISONE

* Tab 5 mg .....	8.10	100	✓ <u>Douglas</u>
* Tab 20 mg .....	20.32	100	✓ <u>Douglas</u>
* Inj 100 mg vial .....	5.30	1	✓ <u>Solu-Cortef</u>
a) Up to 5 inj available on a PSO			
b) Only on a PSO			

### METHYLPREDNISOLONE – Retail pharmacy-Specialist

* Tab 4 mg .....	112.00	100	✓ <u>Medrol</u>
* Tab 100 mg .....	194.00	20	✓ <u>Medrol</u>

### METHYLPREDNISOLONE (AS SODIUM SUCCINATE) – Retail pharmacy-Specialist

Inj 40 mg vial .....	18.90	1	✓ <u>Solu-Medrol-Act-O-Vial</u>
Inj 125 mg vial .....	28.90	1	✓ <u>Solu-Medrol-Act-O-Vial</u>
Inj 500 mg vial .....	22.78	1	✓ <u>Solu-Medrol-Act-O-Vial</u>
Inj 1 g vial .....	27.83	1	✓ <u>Solu-Medrol</u>
METHYLPREDNISOLONE ACETATE			
Inj 40 mg per ml, 1 ml vial.....	44.40	5	✓ <u>Depo-Medrol</u>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

## HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>PREDNISOLONE</b>				
* Oral liq 5 mg per ml – Up to 30 ml available on a PSO .....6.00		30 ml OP	✓	<b>Redipred</b>
Restricted to children under 12 years of age.				
<b>PREDNISONE</b>				
* Tab 1 mg .....10.68		500	✓	<b>Apo-Prednisone</b>
* Tab 2.5 mg .....12.09		500	✓	<b>Apo-Prednisone</b>
* Tab 5 mg – Up to 30 tab available on a PSO .....11.09		500	✓	<b>Apo-Prednisone</b>
* Tab 20 mg .....29.03		500	✓	<b>Apo-Prednisone</b>
<b>TETRACOSACTRIN</b>				
* Inj 250 mcg per ml, 1 ml ampoule .....75.00		1	✓	<b>AU Synacthen</b>
			✓	<b>Synacthen</b>
			✓	<b>Synacthen S29</b> <sup>S29</sup>
* Inj 1 mg per ml, 1 ml ampoule .....690.00		1	✓	<b>Synacthen Depot</b>
			✓	<b>Synacthene</b>
				<b>Retard</b> <sup>S29</sup>
<i>(Synacthen S29 <sup>S29</sup> Inj 250 mcg per ml, 1 ml ampoule to be delisted 1 January 2020)</i>				
<b>TRIAMCINOLONE ACETONIDE</b>				
Inj 10 mg per ml, 1 ml ampoule .....20.80		5	✓	<b>Kenacort-A 10</b>
Inj 40 mg per ml, 1 ml ampoule .....51.10		5	✓	<b>Kenacort-A 40</b>

### Sex Hormones Non Contraceptive

#### Androgen Agonists and Antagonists

<b>CYPROTERONE ACETATE – Retail pharmacy-Specialist</b>				
Tab 50 mg .....13.17		50	✓	<b>Siterone</b>
Tab 100 mg .....26.75		50	✓	<b>Siterone</b>
<b>TESTOSTERONE</b>				
Patch 5 mg per day .....90.00		30	✓	<b>Androderm</b>
<b>TESTOSTERONE CIPIONATE – Retail pharmacy-Specialist</b>				
Inj 100 mg per ml, 10 ml vial.....76.50		1	✓	<b>Depo-Testosterone</b>
<b>TESTOSTERONE ESTERS – Retail pharmacy-Specialist</b>				
Inj 250 mg per ml, 1 ml .....12.98		1	✓	<b>Sustanon Ampoules</b>
<b>TESTOSTERONE UNDECANOATE – Retail pharmacy-Specialist</b>				
Cap 40 mg .....21.00		60	✓	<b>Andriol Testocaps</b>
Inj 250 mg per ml, 4 ml vial.....86.00		1	✓	<b>Reandron 1000</b>

### Hormone Replacement Therapy - Systemic

#### Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG “Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004”.



# HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>Oestrogens</b>				
OESTRADIOL – See prescribing guideline <a href="#">on the previous page</a>				
* Tab 1 mg .....	4.12 (11.10)	28 OP		Estrofem
* Tab 2 mg .....	4.12 (11.10)	28 OP		Estrofem
* Patch 25 mcg per day.....	6.12	8	✓	<b>Estradot</b>
a) No more than 2 patch per week				
b) Only on a prescription				
* Patch 50 mcg per day.....	7.04	8	✓	<b>Estradot 50 mcg</b>
a) No more than 2 patch per week				
b) Only on a prescription				
* Patch 75 mcg per day.....	7.91	8	✓	<b>Estradot</b>
a) No more than 2 patch per week				
b) Only on a prescription				
* Patch 100 mcg per day.....	7.91	8	✓	<b>Estradot</b>
a) No more than 2 patch per week				
b) Only on a prescription				
OESTRADIOL VALERATE – See prescribing guideline <a href="#">on the previous page</a>				
* Tab 1 mg .....	12.36	84	✓	<b>Progynova</b>
* Tab 2 mg .....	12.36	84	✓	<b>Progynova</b>
OESTROGENS – See prescribing guideline <a href="#">on the previous page</a>				
* Conjugated, equine tab 300 mcg.....	3.01 (13.50)	28		Premarin
* Conjugated, equine tab 625 mcg.....	4.12 (13.50)	28		Premarin
<b>Progestogens</b>				
MEDROXYPROGESTERONE ACETATE – See prescribing guideline <a href="#">on the previous page</a>				
* Tab 2.5 mg .....	3.75	30	✓	<b>Provera</b>
* Tab 5 mg .....	14.00	100	✓	<b>Provera</b>
* Tab 10 mg .....	7.15	30	✓	<b>Provera</b>
<b>Progestogen and Oestrogen Combined Preparations</b>				
OESTRADIOL WITH NORETHISTERONE – See prescribing guideline <a href="#">on the previous page</a>				
* Tab 1 mg with 0.5 mg norethisterone acetate .....	5.40 (18.10)	28 OP		Kliovance
* Tab 2 mg with 1 mg norethisterone acetate .....	5.40 (18.10)	28 OP		Kliogest
* Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg oestradiol tab (12) and 1 mg oestradiol tab (6).....	5.40 (18.10)	28 OP		Trisequens
<b>Other Oestrogen Preparations</b>				
ETHINYLOESTRADIOL				
* Tab 10 mcg.....	17.60	100	✓	<b>NZ Medical and Scientific</b>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

# HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
OESTRIOL				
* Tab 2 mg .....	7.00	30	✓	Ovestin

## Other Progestogen Preparations

### LEVONORGESTREL

* Intra-uterine system 20 mcg per day – Special Authority see SA1608 below – Retail pharmacy .....	269.50	1	✓	Mirena
---	--------	---	---	--------

#### ►SA1608 Special Authority for Subsidy

**Initial application — (No previous use)** only from a relevant specialist or general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a clinical diagnosis of heavy menstrual bleeding; and
- 2 The patient has failed to respond to or is unable to tolerate other appropriate pharmaceutical therapies as per the Heavy Menstrual Bleeding Guidelines; and
- 3 Either:
  - 3.1 serum ferritin level < 16 mcg/l (within the last 12 months); or
  - 3.2 haemoglobin level < 120 g/l.

Note: Applications are not to be made for use in patients as contraception except where they meet the above criteria.

**Renewal** only from a relevant specialist or general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Patient demonstrated clinical improvement of heavy menstrual bleeding; or
  - 1.2 Previous insertion was removed or expelled within 3 months of insertion; and
- 2 Applicant to state date of the previous insertion.

### MEDROXYPROGESTERONE ACETATE

Tab 100 mg – Retail pharmacy-Specialist.....	101.00	100	✓	Provera HD
--	--------	-----	---	------------

### NORETHISTERONE

* Tab 5 mg – Up to 30 tab available on a PSO .....	18.29	100	✓	Primolut N
Primolut N to be Sole Supply on 1 January 2020				

### PROGESTERONE

Cap 100 mg – Special Authority see SA1609 below – Retail pharmacy.....	16.50	30	✓	Utrogestan
---	-------	----	---	------------

#### ►SA1609 Special Authority for Subsidy

**Initial application** only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 For the prevention of pre-term labour\*; and
- 2 Either:
  - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
  - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

**Renewal** only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For the prevention of pre-term labour\*; and
- 2 Treatment is required for second or subsequent pregnancy; and
- 3 Either:
  - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
  - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with \* are unapproved indications.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>Thyroid and Antithyroid Agents</b>				
<b>CARBIMAZOLE</b>				
* Tab 5 mg .....	10.80	100	✓ AFT	
			✓ Carbimazole <sup>S29</sup>	
			✓ Neo-Mercazole	
<b>LEVOTHYROXINE</b>				
* Tab 25 mcg.....	3.89	90	✓ Synthroid	
* Tab 50 mcg.....	1.71	28	✓ Mercury Pharma	
	4.05	90	✓ Synthroid	
	64.28	1,000	✓ Eltroxin	
* Tab 100 mcg.....	1.78	28	✓ Mercury Pharma	
	4.21	90	✓ Synthroid	
	66.78	1,000	✓ Eltroxin	
<b>PROPYLTHIOURACIL</b> – Special Authority see <a href="#">SA1199 below</a> – Retail pharmacy				
Propylthiouracil is not recommended for patients under the age of 18 years unless the patient is pregnant and other treatments are contraindicated.				
Tab 50 mg .....	35.00	100	✓ PTU <sup>S29</sup>	

#### ►SA1199 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:  
Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

## Trophic Hormones

### Growth Hormones

**SOMATROPIN (OMNITROPE)** – Special Authority see [SA1629 below](#) – Retail pharmacy

* Inj 5 mg cartridge.....	34.88	1	✓ Omnitrope
* Inj 10 mg cartridge.....	69.75	1	✓ Omnitrope
* Inj 15 mg cartridge.....	104.63	1	✓ Omnitrope

#### ►SA1629 Special Authority for Subsidy

**Initial application** — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist.

Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
- 2 All of the following:
  - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
  - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
  - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

continued...

- children who are 5 years or older, GH testing with sex steroid priming is required; and
- 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
- 2.5 Appropriate imaging of the pituitary gland has been obtained.

**Renewal — (growth hormone deficiency in children)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

**Initial application — (Turner syndrome)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

**Renewal — (Turner syndrome)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under ; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

**Initial application — (short stature without growth hormone deficiency)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

**Renewal — (short stature without growth hormone deficiency)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

**Initial application — (short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
  - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m<sup>2</sup> as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) x 40 = corrected GFR (ml/min/1.73m<sup>2</sup> in a child who may or may not be receiving dialysis; or
  - 6.2 The patient has received a renal transplant and has received < 5mg/ m<sup>2</sup>/day of prednisone or equivalent for at least 6 months..

**Renewal — (short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

**Initial application — (Prader-Willi syndrome)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
  - 5.1 Both:

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

continued...

- 5.1.1 The patient is aged two years or older; and
- 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
- 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

**Renewal — (Prader-Willi syndrome)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

**Initial application — (adults and adolescents)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

**Notes:** For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

**Renewal — (adults and adolescents)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has been treated with somatropin for < 12 months; and
  - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
  - 1.3 Serum IGF-I levels have been increased within  $\pm 1SD$  of the mean of the normal range for age and sex; and

continued...

# HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
  - 2.1 The patient has been treated with somatropin for more than 12 months; and
  - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
  - 2.3 Serum IGF-I levels have continued to be maintained within  $\pm 1SD$  of the mean of the normal range for age and sex (other than for obvious external factors); and
  - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients.

## GnRH Analogues

### GOSERELIN

Implant 3.6 mg, syringe .....	66.48	1	✓ Zoladex
Implant 10.8 mg, syringe .....	177.50	1	✓ Zoladex

### LEUPRORELIN

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

Inj 3.75 mg prefilled dual chamber syringe – Higher subsidy of \$221.60 per 1 inj with Endorsement .....	66.48 (221.60)	1	Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe – Higher subsidy of \$591.68 per 1 inj with Endorsement .....	177.50 (591.68)	1	Lucrin Depot 3-month

## Vasopressin Agonists

### DESMOPRESSIN ACETATE

Tab 100 mcg – Special Authority see <a href="#">SA1401 below</a> – Retail pharmacy .....	25.00	30	✓ Minirin
Tab 200 mcg – Special Authority see <a href="#">SA1401 below</a> – Retail pharmacy .....	54.45	30	✓ Minirin
▲ Nasal drops 100 mcg per ml – Retail pharmacy-Specialist .....	39.03	2.5 ml OP	✓ Minirin
▲ Nasal spray 10 mcg per dose – Retail pharmacy-Specialist .....	23.95	6 ml OP	✓ <u>Desmopressin- PH&amp;T</u>
Inj 4 mcg per ml, 1 ml – Special Authority see <a href="#">SA1401 below</a> – Retail pharmacy .....	67.18	10	✓ Minirin

#### ► [SA1401](#) Special Authority for Subsidy

**Initial application — (Desmopressin tablets for Nocturnal enuresis)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has primary nocturnal enuresis; and
- 2 The nasal forms of desmopressin are contraindicated; and
- 3 An enuresis alarm is contraindicated.

**Initial application — (Desmopressin tablets for Diabetes insipidus)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

continued...

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\* Three months or six months, as applicable, dispensed all-at-once

# HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 1 The patient has cranial diabetes insipidus; and
- 2 The nasal forms of desmopressin are contraindicated.

**Renewal — (Desmopressin tablets)** from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from the treatment.

**Initial application — (Desmopressin injection)** only from a relevant specialist. Approvals valid for 2 years where the patient cannot use desmopressin nasal spray or nasal drops.

**Renewal — (Desmopressin injection)** only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

## Other Endocrine Agents

### CABERGOLINE

Tab 0.5 mg – Maximum of 2 tab per prescription; can be

waived by Special Authority see <a href="#">SA1370 below</a> .....	3.75	2	✓ <b>Dostinex</b>
	15.20	8	✓ <b>Dostinex</b>

### ► [SA1370](#) Special Authority for Waiver of Rule

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 pathological hyperprolactinemia; or
- 2 acromegaly\*.

**Renewal — (for patients who have previously been funded under Special Authority form SA1031)** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with \* is an unapproved indication.

### CLOMIFENE CITRATE

Tab 50 mg .....	29.84	10	✓ <b>Mylan</b> <b>Clomiphen</b> <small>\$29</small>
-----------------	-------	----	--

### DANAZOL

Cap 100 mg .....	68.33	100	✓ <b>Azol</b>
Cap 200 mg .....	97.83	100	✓ <b>Azol</b>

### METYRAPONE

Cap 250 mg – Retail pharmacy-Specialist .....	520.00	50	✓ <b>Metopirone</b>
---	--------	----	---------------------



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

## Anthelmintics

ALBENDAZOLE – Special Authority see [SA1318 below](#) – Retail pharmacy

Tab 400 mg .....	469.20	60	✓ Eskazole <sup>S29</sup>
------------------	--------	----	---------------------------

### ►SA1318 Special Authority for Subsidy

**Initial application** only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids.

**Renewal** only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

MEBENDAZOLE – Only on a prescription

Tab 100 mg .....	24.19	24	✓ De-Worm
Oral liq 100 mg per 5 ml .....	2.18	15 ml	Vermox
	(7.17)		

PRAZIQUANTEL

Tab 600 mg .....	68.00	8	✓ Biltricide
------------------	-------	---	--------------

## Antibacterials

- a) For topical antibacterials, refer to DERMATOLOGICALS, [page 59](#)  
b) For anti-infective eye preparations, refer to SENSORY ORGANS, [page 228](#)

## Cephalosporins and Cephamycins

CEFACLOR MONOHYDRATE

Cap 250 mg .....	24.70	100	✓ <u>Ranbaxy-Cefaclor</u>
Grans for oral liq 125 mg per 5 ml – Wastage claimable.....	3.53	100 ml	✓ <u>Ranbaxy-Cefaclor</u>
	4.33		✓ Keflor

CEFALEXIN

Cap 250 mg .....	3.33	20	✓ Cephalexin ABM
Cephalexin ABM to be Sole Supply on 1 November 2019			
Cap 500 mg .....	3.95	20	✓ Cephalexin ABM
Grans for oral liq 25 mg per ml – Wastage claimable.....	8.75	100 ml	✓ <u>Cefalexin Sandoz</u>
Note: Cefalexin grans for oral liq will not be funded in amounts more than 14 days treatment per dispensing.			
Grans for oral liq 50 mg per ml – Wastage claimable.....	11.75	100 ml	✓ <u>Cefalexin Sandoz</u>
Note: Cefalexin grans for oral liq will not be funded in amounts more than 14 days treatment per dispensing.			

CEFAZOLIN – Subsidy by endorsement

Only if prescribed for dialysis or cellulitis in accordance with a DHB approved protocol and the prescription is endorsed accordingly.

Inj 500 mg vial .....	3.39	5	✓ <u>AFT</u>
Inj 1 g vial .....	3.29	5	✓ <u>AFT</u>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>CEFTRIAXONE – Subsidy by endorsement</b>				
a) Up to 10 inj available on a PSO				
b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of pelvic inflammatory disease, or the treatment of suspected meningococcal disease, and the prescription or PSO is endorsed accordingly.				
Inj 500 mg vial .....	0.89	1	✓	Ceftriaxone-AFT
	1.20		✓	DEVA
Ceftriaxone-AFT to be Sole Supply on 1 January 2020				
Inj 1 g vial .....	0.84	1	✓	DEVA
	3.99	5	✓	Ceftriaxone-AFT
Ceftriaxone-AFT to be Sole Supply on 1 January 2020				
<i>(DEVA Inj 500 mg vial to be delisted 1 January 2020)</i>				
<i>(DEVA Inj 1 g vial to be delisted 1 January 2020)</i>				
<b>CEFUROXIME AXETIL – Subsidy by endorsement</b>				
Only if prescribed for prophylaxis of endocarditis and the prescription is endorsed accordingly.				
Tab 250 mg .....	45.93	50	✓	Zinnat

## Macrolides

<b>AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below</b>				
A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.				
Tab 250 mg .....	8.19	30	✓	Apo-Azithromycin
Tab 500 mg – Up to 8 tab available on a PSO .....	0.93	2	✓	Apo-Azithromycin
Grans for oral liq 200 mg per 5 ml (40 mg per ml) – Wastage claimable .....				
	14.38	15 ml	✓	Zithromax

### ►SA1683 Special Authority for Waiver of Rule

**Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections)** only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome\*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome\*; or
- 3 Patient has cystic fibrosis and has chronic infection with *Pseudomonas aeruginosa* or *Pseudomonas*-related gram negative organisms\*; or
- 4 Patient has an atypical *Mycobacterium* infection.

Note: Indications marked with \* are unapproved indications.

**Initial application — (non-cystic fibrosis bronchiectasis\*)** only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis\*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
  - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
  - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with \* are unapproved indications.

**Renewal — (non-cystic fibrosis bronchiectasis\*)** only from a respiratory specialist or paediatrician. Approvals valid for 12

continued...



	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>Penicillins</b>				
<b>AMOXICILLIN</b>				
Cap 250 mg.....	14.97	500	✓	<b>Apo-Amoxi</b>
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Cap 500 mg.....	16.75	500	✓	<b>Apo-Amoxi</b>
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml.....	1.20	100 ml	✓	<b>Alphamox 125</b>
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml.....	1.31	100 ml	✓	<b>Alphamox 250</b>
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
Inj 250 mg vial .....	10.67	10	✓	<b>Ibiamox</b>
Inj 500 mg vial .....	12.41	10	✓	<b>Ibiamox</b>
Inj 1 g vial – Up to 5 inj available on a PSO .....	17.29	10	✓	<b>Ibiamox</b>
<b>AMOXICILLIN WITH CLAVULANIC ACID</b>				
Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab available on a PSO.....	1.88	20	✓	<b>Augmentin</b>
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 mg per ml.....	3.83	100 ml	✓	<b>Augmentin</b>
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 mg per ml – Up to 200 ml available on a PSO .....	2.20	100 ml OP	✓	<b>Curam</b>
<b>BENZATHINE BENZYL PENICILLIN</b>				
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj available on a PSO.....	344.93	10	✓	<b>Bicillin LA</b>
<b>BENZYL PENICILLIN SODIUM [PENICILLIN G]</b>				
Inj 600 mg (1 million units) vial – Up to 5 inj available on a PSO .....	10.35	10	✓	<b>Sandoz</b>
<b>FLUCLOXACILLIN</b>				
Cap 250 mg – Up to 30 cap available on a PSO.....	16.83	250	✓	<b>Staphlex</b>
Cap 500 mg.....	56.61	500	✓	<b>Staphlex</b>
Grans for oral liq 25 mg per ml.....	2.29	100 ml	✓	<b>AFT</b>
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 50 mg per ml.....	3.68	100 ml	✓	<b>AFT</b>
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Inj 250 mg vial .....	9.00	10	✓	<b>Flucloxin</b>
Inj 500 mg vial .....	9.40	10	✓	<b>Flucloxin</b>
Inj 1 g vial – Up to 5 inj available on a PSO .....	5.22	5	✓	<b>Flucil</b>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>PHENOXYMETHYLPENICILLIN (PENICILLIN V)</b>				
Cap 250 mg – Up to 30 cap available on a PSO .....	2.59	50	✓	<b>Cilicaine VK</b>
Cap 500 mg .....	4.26	50	✓	<b>Cilicaine VK</b>
a) Up to 20 cap available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml .....	2.99	100 ml	✓	<b>AFT</b>
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
c) AFT to be Sole Supply on 1 January 2020				
Grans for oral liq 250 mg per 5 ml .....	3.99	100 ml	✓	<b>AFT</b>
a) Up to 300 ml available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
d) AFT to be Sole Supply on 1 January 2020				
<b>PROCAINE PENICILLIN</b>				
Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO .....	123.50	5	✓	<b>Cilicaine</b>

### Tetracyclines

<b>DOXYCYCLINE</b>				
* Tab 50 mg – Up to 30 tab available on a PSO .....	2.90 (6.00)	30		
* Tab 100 mg – Up to 30 tab available on a PSO .....	64.43	500	✓	<b>Doxine</b>
<i>(Doxo-50 Tab 50 mg to be delisted 1 January 2020)</i>				
<b>MINOCYCLINE HYDROCHLORIDE</b>				
* Tab 50 mg – Additional subsidy by Special Authority see SA1355 below – Retail pharmacy .....	5.79 (12.05)	60		Mino-tabs
* Cap 100 mg .....	19.32 (52.04)	100		Minomycin

#### ►SA1355 Special Authority for Manufacturers Price

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

**TETRACYCLINE** – Special Authority see SA1332 below – Retail pharmacy

Cap 500 mg .....	46.00	30	✓	<b>Tetracyclin Wolff S29</b>
------------------	-------	----	---	----------------------------------

#### ►SA1332 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

## Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, [page 59](#)

### CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg – Up to 5 tab available on a PSO .....	1.45	28	✓ <b>Cipflox</b>
Tab 500 mg – Up to 5 tab available on a PSO .....	1.99	28	✓ <b>Cipflox</b>
Tab 750 mg .....	3.15	28	✓ <b>Cipflox</b>

### CLINDAMYCIN

Cap hydrochloride 150 mg – Maximum of 4 cap per prescription; can be waived by endorsement - Retail pharmacy - Specialist .....

4.10	16	✓ <b>Clindamycin ABM</b>
4.61	24	✓ <b>Dalacin C</b>

Inj phosphate 150 mg per ml, 4 ml ampoule – Retail pharmacy-Specialist .....

39.00	10	✓ <b>Dalacin C</b>
-------	----	--------------------

### COLISTIN SULPHOMETHATE – Retail pharmacy-Specialist – Subsidy by endorsement

Only if prescribed for dialysis or cystic fibrosis patient and the prescription is endorsed accordingly.

Inj 150 mg .....	65.00	1	✓ <b>Colistin-Link</b>
------------------	-------	---	------------------------

### GENTAMICIN SULPHATE

Inj 10 mg per ml, 1 ml ampoule – Subsidy by endorsement.....

25.00	5	✓ <b>DBL Gentamicin</b>
-------	---	-------------------------

Only if prescribed for a dialysis or cystic fibrosis patient or complicated urinary tract infection and the prescription is endorsed accordingly.

Inj 40 mg per ml, 2 ml ampoule – Subsidy by endorsement.....	17.50	10	✓ <b>Pfizer</b>
	30.00	50	✓ <b>Pfizer</b>

Only if prescribed for a dialysis or cystic fibrosis patient or complicated urinary tract infection and the prescription is endorsed accordingly.

### MOXIFLOXACIN – Special Authority see [SA1740 below](#) – Retail pharmacy

No patient co-payment payable

Tab 400 mg .....	52.00	5	✓ <b>Avelox</b>
------------------	-------	---	-----------------

### ►SA1740 Special Authority for Subsidy

**Initial application — (Tuberculosis)** only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

1 Both:

- 1.1 Active tuberculosis\*; and
- 1.2 Any of the following:

- 1.2.1 Documented resistance to one or more first-line medications; or
- 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
- 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
- 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

continued...

1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications;  
or

- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.\*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with \* are unapproved indications.

**Renewal** only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

**Initial application — (Mycoplasma genitalium)** only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium\* and is symptomatic; and
- 2 Either:
  - 2.1 Has tried and failed to clear infection using azithromycin; or
  - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

**Initial application — (Penetrating eye injury)** only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with \* are unapproved indications.

PAROMOMYCIN – Special Authority see [SA1689 below](#) – Retail pharmacy

Cap 250 mg .....	126.00	16	✓ Humatin <small>\$29</small>
------------------	--------	----	-------------------------------

► **SA1689** Special Authority for Subsidy

**Initial application** only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolytica carriage.

**Renewal** only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolytica carriage.

PYRIMETHAMINE – Special Authority see [SA1328 below](#) – Retail pharmacy

Tab 25 mg .....	48.00	30	✓ Daraprim <small>\$29</small>
-----------------	-------	----	--------------------------------

► **SA1328** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

SODIUM FUSIDATE [FUSIDIC ACID]

Tab 250 mg – Retail pharmacy-Specialist.....	34.50	12	✓ Fucidin
--	-------	----	-----------

Prescriptions must be written by, or on the recommendation of, an infectious disease physician or a clinical microbiologist

SULFADIAZINE SODIUM – Special Authority see [SA1331 on the next page](#) – Retail pharmacy

Tab 500 mg .....	543.20	56	✓ Wockhardt <small>\$29</small>
------------------	--------	----	---------------------------------

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

## ►SA1331 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

### TOBRAMYCIN

Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement.....	15.00	5	✓ <b>Tobramycin Mylan</b>
Only if prescribed for dialysis or cystic fibrosis patient and the prescription is endorsed accordingly.			
Solution for inhalation 60 mg per ml, 5 ml – Subsidy by endorsement.....	2,200.00	56 dose	✓ <b>TOBI</b>
a) Wastage claimable			
b) Only if prescribed for a cystic fibrosis patient and the prescription is endorsed accordingly.			

### TRIMETHOPRIM

* Tab 300 mg – Up to 30 tab available on a PSO.....	16.50	50	✓ <b>TMP</b>
---	-------	----	--------------

### TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXAZOLE]

* Tab trimethoprim 80 mg and sulphamethoxazole 400 mg – Up to 30 tab available on a PSO.....	53.96	500	✓ <b>Trisul</b>
* Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to 200 ml available on a PSO.....	2.97	100 ml	✓ <b>Deprim</b>

### VANCOMYCIN – Subsidy by endorsement

Only if prescribed for a dialysis or cystic fibrosis patient or for prophylaxis of endocarditis or for treatment of Clostridium difficile following metronidazole failure and the prescription is endorsed accordingly.

Inj 500 mg vial .....	2.37	1	✓ <b>Mylan</b>
-----------------------	------	---	----------------

## Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, [page 60](#)
- b) For topical antifungals refer to GENITO URINARY, [page 75](#)

### FLUCONAZOLE

Cap 50 mg – Retail pharmacy-Specialist .....	2.09	28	✓ <b>Mylan</b>
Cap 150 mg – Subsidy by endorsement .....	0.33	1	✓ <b>Mylan</b>
a) Maximum of 1 cap per prescription; can be waived by endorsement - Retail pharmacy - Specialist			
b) Patient has vaginal candida albicans and the practitioner considers that a topical imidazole (used intra-vaginally) is not recommended and the prescription is endorsed accordingly; can be waived by endorsement - Retail pharmacy - Specialist.			
Cap 200 mg – Retail pharmacy-Specialist .....	5.08	28	✓ <b>Mylan</b>
Powder for oral suspension 10 mg per ml – Special Authority see <a href="#">SA1359 below</a> – Retail pharmacy.....	34.56	35 ml	✓ <b>Diflucan S29</b> <small>S29</small>
	98.50		✓ <b>Diflucan</b>
Wastage claimable			

## ►SA1359 Special Authority for Subsidy

**Initial application — (Systemic candidiasis)** from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

continued...



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

**Initial application — (Immunocompromised)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

**Renewal — (Systemic candidiasis)** from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

**Renewal — (Immunocompromised)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

## ITRACONAZOLE

Cap 100 mg – Subsidy by endorsement .....	4.27	15	✓ Itrazole
a) Funded for tinea versicolor where topical treatment has not been successful and diagnosis has been confirmed by mycology, or for tinea unguium where terbinafine has not been successful in eradication or the patient is intolerant to terbinafine and diagnosis has been confirmed by mycology and the prescription is endorsed accordingly. Can be waived by endorsement - Retail pharmacy - Specialist Specialist must be an infectious disease physician, clinical microbiologist, clinical immunologist or dermatologist.			
b) Itrazole to be Sole Supply on 1 November 2019			
Oral liq 10 mg per ml – Special Authority see <a href="#">SA1322 below</a> –			
Retail pharmacy.....	141.80	150 ml OP	✓ Sporanox

### ►SA1322 Special Authority for Subsidy

**Initial application** only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

**Renewal** from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

## KETOCONAZOLE

Tab 200 mg – PCT – Retail pharmacy-Specialist – Subsidy by endorsement.....	CBS	30	✓ Link Healthcare <sup>S29</sup> ✓ Nizoral <sup>S29</sup>
---	-----	----	--

Prescriptions must be written by, or on the recommendation of an oncologist

## NYSTATIN

Tab 500,000 u .....	14.16 (17.09)	50	Nilstat
Cap 500,000 u .....	12.81 (15.47)	50	Nilstat

POSACONAZOLE – Special Authority see [SA1285 on the next page](#) – Retail pharmacy

Tab modified-release 100 mg.....	869.86	24	✓ Noxafil
Oral liq 40 mg per ml .....	761.13	105 ml OP	✓ Noxafil

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

## ►SA1285 Special Authority for Subsidy

**Initial application** only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy\*.

**Renewal** only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression\* and requires on going posaconazole treatment.

Note: \* Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

### TERBINAFINE

\* Tab 250 mg ..... 1.33 14 ✓ **Deolate**

### VORICONAZOLE – Special Authority see SA1273 below – Retail pharmacy

Tab 50 mg ..... 91.00 56 ✓ **Vttack**

Tab 200 mg ..... 350.00 56 ✓ **Vttack**

Powder for oral suspension 40 mg per ml – Wastage

claimable ..... 1,437.00 70 ml ✓ **Vfend**

## ►SA1273 Special Authority for Subsidy

**Initial application — (invasive fungal infection)** only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient has proven or probable invasive aspergillus infection; or
  - 3.2 Patient has possible invasive aspergillus infection; or
  - 3.3 Patient has fluconazole resistant candidiasis; or
  - 3.4 Patient has mould strain such as *Fusarium* spp. and *Scedosporium* spp.

**Renewal — (invasive fungal infection)** only from a haematologist, infectious disease specialist or clinical microbiologist.

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
  - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
  - 3.3 Patient has fluconazole resistant candidiasis; or
  - 3.4 Patient has mould strain such as *Fusarium* spp. and *Scedosporium* spp.

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

## Antimalarials

PRIMAQUINE PHOSPHATE – Special Authority see [SA1684 below](#) – Retail pharmacy

Tab 7.5 mg .....	117.00	56	✓ Primacin <sup>\$29</sup>
------------------	--------	----	----------------------------

### ►SA1684 Special Authority for Subsidy

**Initial application** only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

**Renewal** only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

## Antiparasitics

### Antiprotozoals

QUININE SULPHATE

* Tab 300 mg .....	61.91	500	✓ Q 300
--------------------	-------	-----	---------

## Antitrichomonal Agents

METRONIDAZOLE

Tab 200 mg – Up to 30 tab available on a PSO .....	10.45	100	✓ Trichozole
Tab 400 mg – Up to 15 tab available on a PSO .....	18.15	100	✓ Trichozole
Oral liq benzoate 200 mg per 5 ml .....	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg .....	24.48	10	✓ Flagyl

ORNIDAZOLE

Tab 500 mg .....	23.00	10	✓ Arrow-Ornidazole
------------------	-------	----	--------------------

## Antituberculotics and Antileprotics

Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status.

CLOFAZIMINE – Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist.

* Cap 50 mg .....	442.00	100	✓ Lamprene <sup>\$29</sup>
-------------------	--------	-----	----------------------------

CYCLOSERINE – Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician.

Cap 250 mg .....	344.00	60	✓ Cycloclin <sup>\$29</sup>
	1,294.50	100	✓ King <sup>\$29</sup>

(King <sup>\$29</sup> Cap 250 mg to be delisted 1 November 2019)

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

# INFECTIONS - AGENTS FOR SYSTEMIC USE

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>DAPSONE – Retail pharmacy-Specialist</b>				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist				
Tab 25 mg .....	268.50	100	✓	<b>Dapsone</b>
Tab 100 mg .....	329.50	100	✓	<b>Dapsone</b>
<b>ETHAMBUTOL HYDROCHLORIDE – Retail pharmacy-Specialist</b>				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician				
Tab 100 mg .....	85.73	100	✓	<b>EMB Fato1</b> <small>\$29</small>
Tab 400 mg .....	49.34	56	✓	<b>Myambutol</b> <small>\$29</small>
<b>ISONIAZID – Retail pharmacy-Specialist</b>				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician, paediatrician, clinical microbiologist, dermatologist or public health physician				
* Tab 100 mg .....	22.00	100	✓	<b>PSM</b>
<b>ISONIAZID WITH RIFAMPICIN – Retail pharmacy-Specialist</b>				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician, paediatrician, clinical microbiologist, dermatologist or public health physician				
* Tab 100 mg with rifampicin 150 mg.....	85.54	100	✓	<b>Rifinah</b>
* Tab 150 mg with rifampicin 300 mg.....	170.60	100	✓	<b>Rifinah</b>
<b>PARA-AMINO SALICYLIC ACID – Retail pharmacy-Specialist</b>				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendation of, an infectious disease specialist, clinical microbiologist or respiratory physician				
Grans for oral liq 4 g sachet .....	280.00	30	✓	<b>Paser</b> <small>\$29</small>
<b>PROTIONAMIDE – Retail pharmacy-Specialist</b>				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendation of, an infectious disease specialist, clinical microbiologist or respiratory physician				
Tab 250 mg .....	305.00	100	✓	<b>Peteha</b> <small>\$29</small>
<b>PYRAZINAMIDE – Retail pharmacy-Specialist</b>				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician				
* Tab 500 mg .....	59.00	100	✓	<b>AFT-Pyrazinamide</b>
<b>RIFABUTIN – Retail pharmacy-Specialist</b>				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, respiratory physician or gastroenterologist				
* Cap 150 mg .....	275.00	30	✓	<b>Mycobutin</b>

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
<b>RIFAMPICIN – Subsidy by endorsement</b>			
a) No patient co-payment payable			
b) For confirmed recurrent <i>Staphylococcus aureus</i> infection in combination with other effective anti-staphylococcal antimicrobial based on susceptibilities and the prescription is endorsed accordingly; can be waived by endorsement - Retail pharmacy - Specialist. Specialist must be an internal medicine physician, clinical microbiologist, dermatologist, paediatrician, or public health physician.			
* Cap 150 mg .....	55.75	100	✓ <b>Rifadin</b>
* Cap 300 mg .....	116.25	100	✓ <b>Rifadin</b>
* Oral liq 100 mg per 5 ml .....	12.00	60 ml	✓ <b>Rifadin</b>

## Antivirals

For eye preparations refer to Eye Preparations, Anti-Infective Preparations, [page 228](#)

### Hepatitis B Treatment

**ADEFOVIR DIPIVOXIL – Special Authority** see [SA0829 below](#) – Retail pharmacy

Tab 10 mg .....	670.00	30	✓ <b>Hepsera</b>
-----------------	--------	----	------------------

#### ► [SA0829](#) Special Authority for Subsidy

**Initial application** only from a gastroenterologist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg+); and  
Documented resistance to lamivudine, defined as:
- 2 Patient has raised serum ALT ( $> 1 \times \text{ULN}$ ); and
- 3 Patient has HBV DNA greater than 100,000 copies per mL, or viral load 10 fold or higher over nadir; and
- 4 Detection of M204I or M204V mutation; and
- 5 Either:
  - 5.1 Both:
    - 5.1.1 Patient is cirrhotic; and
    - 5.1.2 adefovir dipivoxil to be used in combination with lamivudine; or
  - 5.2 Both:
    - 5.2.1 Patient is not cirrhotic; and
    - 5.2.2 adefovir dipivoxil to be used as monotherapy.

**Renewal** only from a gastroenterologist or infectious disease specialist. Approvals valid for 2 years where in the opinion of the treating physician, treatment remains appropriate and patient is benefiting from treatment.

Notes: Lamivudine should be added to adefovir dipivoxil if a patient develops documented resistance to adefovir dipivoxil, defined as:

- i) raised serum ALT ( $> 1 \times \text{ULN}$ ); and
- ii) HBV DNA greater than 100,000 copies per mL, or viral load 10 fold or higher over nadir; and
- iii) Detection of N236T or A181T/V mutation.

Adefovir dipivoxil should be stopped 6 months following HBsAg seroconversion for patients who were HBsAg+ prior to commencing adefovir dipivoxil.

The recommended dose of adefovir dipivoxil is no more than 10mg daily.

In patients with renal insufficiency adefovir dipivoxil dose should be reduced in accordance with the datasheet guidelines.

Adefovir dipivoxil should be avoided in pregnant women and children.

#### ENTECAVIR

* Tab 0.5 mg .....	52.00	30	✓ <b>Entecavir Sandoz</b>
--------------------	-------	----	---------------------------

**LAMIVUDINE – Special Authority** see [SA1685 on the next page](#) – Retail pharmacy

Tab 100 mg .....	4.20	28	✓ <b>Zetiam</b>
Oral liq 5 mg per ml .....	270.00	240 ml OP	✓ <b>Zeffix</b>

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\* Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

**►SA1685 Special Authority for Subsidy**

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B.

**TENOFOVIR DISOPROXIL**

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA1651., [page 105](#)

* Tab 245 mg (300.6 mg as a succinate) .....	38.10	30	✓	<b>Tenofovir Disoproxil</b> <b>Teva</b>
--	-------	----	---	--

**Herpesvirus Treatments**
**ACICLOVIR**

* Tab dispersible 200 mg .....	1.60	25	✓	<b>Lovir</b>
* Tab dispersible 400 mg .....	5.38	56	✓	<b>Lovir</b>
* Tab dispersible 800 mg .....	5.98	35	✓	<b>Lovir</b>

**VALACICLOVIR**

Tab 500 mg .....	5.75	30	✓	<b>Vaclovir</b>
Tab 1,000 mg .....	11.35	30	✓	<b>Vaclovir</b>

**VALGANCICLOVIR – Special Authority see SA1404 below – Retail pharmacy**

Tab 450 mg .....	225.00	60	✓	<b>Valganciclovir</b> <b>Mylan</b>
------------------	--------	----	---	---------------------------------------

**►SA1404 Special Authority for Subsidy**

**Initial application — (transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

**Renewal — (transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
- 2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin.

**Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin)** only from a relevant specialist.

Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

**Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin)** only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

**Initial application — (Lung transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
  - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
  - 2.2 The recipient is cytomegalovirus positive.

**Initial application — (Cytomegalovirus in immunocompromised patients)** only from a relevant specialist. Approvals valid for

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

**Renewal — (Cytomegalovirus in immunocompromised patients)** only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

## Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR – [Xpharm]

Note the supply of treatment is via PHARMAC's approved direct distribution supply. Further details can be found on PHARMAC's website <https://www.pharmac.govt.nz/hepatitis-c-treatments>

Tab 100 mg with pibrentasvir 40 mg .....	24,750.00	84 OP	✓ Maviret
--	-----------	-------	-----------

LEDIPASVIR WITH SOFOSBUVIR – [Xpharm] – Special Authority see [SA1605 below](#)

No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg .....	24,363.46	28	✓ Harvoni
--	-----------	----	-----------

### ➔SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from PHARMAC's website <http://www.pharmac.govt.nz/hepatitis-c-treatments> or:

The Coordinator, Hepatitis C Treatment Panel

PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: [hepcpanel@pharmac.govt.nz](mailto:hepcpanel@pharmac.govt.nz)

## HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL – Subsidy by endorsement; can be waived by Special Authority see [SA1842 on the next page](#)

- a) Brand switch fee payable (Pharmacode 2573865) - see [page 233](#) for details
- b) Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, [page 105](#) There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the PHARMAC website.

Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a succinate) .....	61.15	30	✓ Teva
---	-------	----	--------

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## ➔SA1842 Special Authority for Waiver of Rule

**Initial application** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or <https://ashm.org.au/HIV/PrEP/> for training materials); and
- 2 Patient has undergone testing for HIV, syphilis, Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
  - 6.1 All of the following:
    - 6.1.1 Patient is male or transgender; and
    - 6.1.2 Patient has sex with men; and
    - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
    - 6.1.4 Any of the following:
      - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
      - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
      - 6.1.4.3 Patient has used methamphetamine in the last three months; or
  - 6.2 All of the following:
    - 6.2.1 Patient has a regular partner who has HIV infection; and
    - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
    - 6.2.3 Condoms have not been consistently used.

**Renewal** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or <https://ashm.org.au/HIV/PrEP/> for training materials); and
- 2 Patient has undergone testing for HIV, syphilis, Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
  - 6.1 All of the following:
    - 6.1.1 Patient is male or transgender; and
    - 6.1.2 Patient has sex with men; and
    - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
    - 6.1.4 Any of the following:
      - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
      - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
      - 6.1.4.3 Patient has used methamphetamine in the last three months; or
  - 6.2 All of the following:
    - 6.2.1 Patient has a regular partner who has HIV infection; and
    - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
    - 6.2.3 Condoms have not been consistently used.



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## Antiretrovirals

### ►SA1651 Special Authority for Subsidy

**Initial application — (Confirmed HIV)** only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

**Renewal — (Confirmed HIV)** only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

**Initial application — (Prevention of maternal transmission)** only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

**Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV)** only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

**Renewal — (second or subsequent post-exposure prophylaxis)** only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

**Initial application — (Percutaneous exposure)** only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

**Renewal** — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

## Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ – Special Authority see [SA1651 on the previous page](#) – Retail pharmacy

Tab 50 mg .....	63.38	30	✓ Stocrin <sup>\$29</sup>
Tab 200 mg .....	190.15	90	✓ Stocrin
Tab 600 mg .....	63.38	30	✓ Stocrin
Oral liq 30 mg per ml .....	145.79	180 ml OP	✓ Stocrin <sup>\$29</sup>

(Stocrin <sup>\$29</sup> Tab 50 mg to be delisted 1 April 2020)

(Stocrin <sup>\$29</sup> Oral liq 30 mg per ml to be delisted 1 August 2020)

ETRAVIRINE – Special Authority see [SA1651 on the previous page](#) – Retail pharmacy

Tab 200 mg .....	770.00	60	✓ Intencele
------------------	--------	----	-------------

NEVIRAPINE – Special Authority see [SA1651 on the previous page](#) – Retail pharmacy

Tab 200 mg .....	60.00	60	✓ Nevirapine Alphapharm
Oral suspension 10 mg per ml .....	203.55	240 ml	✓ Viramune Suspension

## Nucleosides Reverse Transcriptase Inhibitors

ABACAVIR SULPHATE – Special Authority see [SA1651 on the previous page](#) – Retail pharmacy

Tab 300 mg .....	180.00	60	✓ Ziagen
Oral liq 20 mg per ml .....	256.31	240 ml OP	✓ Ziagen

ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority see [SA1651 on the previous page](#) – Retail pharmacy

Note: abacavir with lamivudine (combination tablets) counts as two anti-retroviral medications for the purposes of the anti-retroviral Special Authority.

Tab 600 mg with lamivudine 300 mg .....	63.00	30	✓ Kivexa
---	-------	----	----------

EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPROXIL – Special Authority see [SA1651 on the previous page](#) – Retail pharmacy

- Brand switch fee payable (Pharmacode 2573873) - see [page 233](#) for details
- Note: Efavirenz with emtricitabine and tenofovir disoproxil counts as three anti-retroviral medications for the purposes of the anti-retroviral Special Authority

Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil 245 mg (300 mg as a maleate) .....	106.88	30	✓ Mylan
---	--------	----	---------

EMTRICITABINE – Special Authority see [SA1651 on the previous page](#) – Retail pharmacy

Cap 200 mg .....	307.20	30	✓ Emtriva
------------------	--------	----	-----------

LAMIVUDINE – Special Authority see [SA1651 on the previous page](#) – Retail pharmacy

Tab 150 mg .....	52.50	60	✓ Lamivudine Alphapharm
Oral liq 10 mg per ml .....	102.50	240 ml OP	✓ 3TC

ZIDOVUDINE [AZT] – Special Authority see [SA1651 on the previous page](#) – Retail pharmacy

Cap 100 mg .....	152.25	100	✓ Retrovir
Oral liq 10 mg per ml .....	30.45	200 ml OP	✓ Retrovir

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority see <a href="#">SA1651 on page 105</a> – Retail pharmacy			
Note: zidovudine [AZT] with lamivudine (combination tablets) counts as two anti-retroviral medications for the purposes of the anti-retroviral Special Authority.			
Tab 300 mg with lamivudine 150 mg.....	33.00	60	✓ <b>Alphapharm</b>

## Protease Inhibitors

ATAZANAVIR SULPHATE – Special Authority see <a href="#">SA1651 on page 105</a> – Retail pharmacy			
Brand switch fee payable (Pharmacode 2573857) - see <a href="#">page 233</a> for details			
Cap 150 mg.....	141.68	60	✓ <b>Teva</b>
Cap 200 mg.....	188.91	60	✓ <b>Teva</b>
DARUNAVIR – Special Authority see <a href="#">SA1651 on page 105</a> – Retail pharmacy			
Tab 400 mg.....	335.00	60	✓ <b>Prezista</b>
Tab 600 mg.....	476.00	60	✓ <b>Prezista</b>
LOPINAVIR WITH RITONAVIR – Special Authority see <a href="#">SA1651 on page 105</a> – Retail pharmacy			
Tab 100 mg with ritonavir 25 mg.....	183.75	60	✓ <b>Kaletra</b>
Tab 200 mg with ritonavir 50 mg.....	463.00	120	✓ <b>Kaletra</b>
Oral liq 80 mg with ritonavir 20 mg per ml.....	735.00	300 ml OP	✓ <b>Kaletra</b>
RITONAVIR – Special Authority see <a href="#">SA1651 on page 105</a> – Retail pharmacy			
Tab 100 mg.....	43.31	30	✓ <b>Norvir</b>

## Strand Transfer Inhibitors

DOLUTEGRAVIR – Special Authority see <a href="#">SA1651 on page 105</a> – Retail pharmacy			
Tab 50 mg.....	1,090.00	30	✓ <b>Tivicay</b>
RALTEGRAVIR POTASSIUM – Special Authority see <a href="#">SA1651 on page 105</a> – Retail pharmacy			
Tab 400 mg.....	1,090.00	60	✓ <b>Isentress</b>
Tab 600 mg.....	1,090.00	60	✓ <b>Isentress HD</b>

## Immune Modulators

### Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

### Criteria for Treatment

#### 1) Diagnosis

- Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
- PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
- Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

### Exclusion Criteria

- 1) Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia ( $< 2.0 \times 10^9$ ) and/or thrombocytopenia.

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

## Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

## Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

INTERFERON ALFA-2A – PCT – Retail pharmacy-Specialist

- a) See prescribing guideline [on the previous page](#)
- b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician or ophthalmologist

Inj 3 m iu prefilled syringe ..... 38.00      1      ✓ **Roferon-A**

PEGYLATED INTERFERON ALFA-2A – Special Authority see [SA1400 below](#) – Retail pharmacy

- a) See prescribing guideline [on the previous page](#)
- b) Note: PHARMAC will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at PHARMAC on 0800-023-588 option 4.

Inj 180 mcg prefilled syringe ..... 500.00      4      ✓ **Pegasys**

## ►SA1400 Special Authority for Subsidy

**Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant)** from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
  - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
  - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

**Renewal — (Chronic hepatitis C - genotype 1 infection)** only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

**Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior)** only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

- 3 Any of the following:
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; or
  - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

**Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV)** from any specialist.

Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

**Initial application — (Hepatitis B)** only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naïve; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log<sub>10</sub> IU/ml; and
- 5 Either:
  - 5.1 HBeAg positive; or
  - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Notes:

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

## Urinary Tract Infections

### HEXAMINE HIPPURATE

* Tab 1 g .....	18.40	100	
	(40.01)		Hiprex

### NITROFURANTOIN

* Tab 50 mg .....	22.20	100	✓ Nifuran
* Tab 100 mg .....	37.50	100	✓ Nifuran

### NORFLOXACIN

Tab 400 mg – Subsidy by endorsement.....	135.00	100	✓ Arrow-Norfloxacin
--	--------	-----	---------------------

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

# MUSCULOSKELETAL SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>Anticholinesterases</b>				
<b>NEOSTIGMINE METILSULFATE</b>				
Inj 2.5 mg per ml, 1 ml ampoule .....	98.00	50	✓	<b>AstraZeneca</b>
<b>PYRIDOSTIGMINE BROMIDE</b>				
▲ Tab 60 mg .....	45.79	100	✓	<b>Mestinon</b>
Mestinon to be Sole Supply on 1 November 2019				
<b>Non-Steroidal Anti-Inflammatory Drugs</b>				
<b>DICLOFENAC SODIUM</b>				
* Tab EC 25 mg .....	1.23	50	✓	<b>Diclofenac Sandoz</b>
* Tab 50 mg dispersible .....	1.50	20	✓	<b>Voltaren D</b>
* Tab EC 50 mg .....	1.23	50	✓	<b>Diclofenac Sandoz</b>
* Tab long-acting 75 mg .....	22.80	500	✓	<b>Apo-Diclo SR</b>
* Tab long-acting 100 mg .....	25.15	500	✓	<b>Apo-Diclo SR</b>
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a PSO .....	13.20	5	✓	<b>Voltaren</b>
* Suppos 12.5 mg .....	2.04	10	✓	<b>Voltaren</b>
* Suppos 25 mg .....	2.44	10	✓	<b>Voltaren</b>
* Suppos 50 mg – Up to 10 supp available on a PSO .....	4.22	10	✓	<b>Voltaren</b>
* Suppos 100 mg .....	7.00	10	✓	<b>Voltaren</b>
<b>IBUPROFEN</b>				
* Tab 200 mg .....	11.71	1,000	✓	<b>Relieve</b>
* Tab long-acting 800 mg .....	7.99	30	✓	<b>Brufen SR</b>
* Oral liq 20 mg per ml .....	1.88	200 ml	✓	<b>Ethics</b>
<b>KETOPROFEN</b>				
* Cap long-acting 200 mg .....	12.07	28	✓	<b>Oruvail SR</b>
<b>MEFENAMIC ACID</b>				
* Cap 250 mg .....	1.25	50		
	(9.16)			Ponstan
	0.50	20		
	(5.60)			Ponstan
<b>NAPROXEN</b>				
* Tab 250 mg .....	32.69	500	✓	<b>Noflam 250</b>
* Tab 500 mg .....	22.19	250	✓	<b>Noflam 500</b>
* Tab long-acting 750 mg .....	6.16	28	✓	<b>Naprosyn SR 750</b>
* Tab long-acting 1 g .....	8.21	28	✓	<b>Naprosyn SR 1000</b>
<b>SULINDAC</b>				
* Tab 100 mg .....	8.55	50	✓	<b>Aclin</b>
* Tab 200 mg .....	15.10	50	✓	<b>Aclin</b>
<b>TENOXICAM</b>				
* Tab 20 mg .....	9.15	100	✓	<b>Tilcotil</b>
* Inj 20 mg vial .....	9.95	1	✓	<b>AFT</b>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>NSAIDs Other</b>				
CELECOXIB				
Cap 100 mg .....	3.63	60	✓	Celebrex
			✓	Celecoxib Pfizer
Cap 200 mg .....	2.30	30	✓	Celebrex
			✓	Celecoxib Pfizer
<i>(Celebrex Cap 100 mg to be delisted 1 January 2020)</i>				
<b>Topical Products for Joint and Muscular Pain</b>				
CAPSAICIN				
Crm 0.025% – Special Authority see <a href="#">SA1289 below</a> – Retail pharmacy .....	6.95	25 g OP	✓	Zostrix
	9.95	45 g OP	✓	Zostrix
➔ <a href="#">SA1289</a> Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.				
<b>Antirheumatoid Agents</b>				
HYDROXYCHLOROQUINE				
* Tab 200 mg .....	7.98	100	✓	Plaquenil
LEFLUNOMIDE				
Tab 10 mg .....	2.90	30	✓	Apo-Leflunomide
Tab 20 mg .....	2.90	30	✓	Apo-Leflunomide
PENICILLAMINE				
Tab 125 mg .....	67.23	100	✓	D-Penamine
Tab 250 mg .....	110.12	100	✓	D-Penamine
SODIUM AUROTHIOMALATE				
Inj 10 mg in 0.5 ml ampoule .....	76.87	10	✓	Myocrisin
Inj 20 mg in 0.5 ml ampoule .....	113.17	10	✓	Myocrisin
Inj 50 mg in 0.5 ml ampoule .....	217.23	10	✓	Myocrisin
<i>(Myocrisin Inj 10 mg in 0.5 ml ampoule to be delisted 1 March 2020)</i>				
<i>(Myocrisin Inj 20 mg in 0.5 ml ampoule to be delisted 1 March 2020)</i>				
<i>(Myocrisin Inj 50 mg in 0.5 ml ampoule to be delisted 1 March 2020)</i>				
<b>Drugs Affecting Bone Metabolism</b>				
<b>Alendronate for Osteoporosis</b>				
ALENDRONATE SODIUM				
* Tab 70 mg .....	2.44	4	✓	Fosamax
ALENDRONATE SODIUM WITH COLECALCIFEROL				
* Tab 70 mg with colecalciferol 5,600 iu .....	1.51	4	✓	Fosamax Plus
<b>Other Treatments</b>				
DENOSUMAB – Special Authority see <a href="#">SA1777 on the next page</a> – Retail pharmacy				
Inj 60 mg prefilled syringe .....	326.00	1	✓	Prolia

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\* Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## ►SA1777 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Either:
  - 2.1 The patient is female and postmenopausal; or
  - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
  - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
  - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
  - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
  - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
  - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
  - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

### PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial.....	5.98	1	✓ <b>Pamisol</b>
Inj 6 mg per ml, 10 ml vial.....	15.02	1	✓ <b>Pamisol</b>
Inj 9 mg per ml, 10 ml vial.....	17.05	1	✓ <b>Pamisol</b>

**RALOXIFENE HYDROCHLORIDE** – Special Authority see [SA1779 on the next page](#) – Retail pharmacy

* Tab 60 mg .....	53.76	28	✓ <b>Evista</b>
-------------------	-------	----	-----------------



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

►SA1779 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- History of two significant osteoporotic fractures demonstrated radiologically; or
- Documented T-Score less than or equal to -3.0 (see Notes); or
- A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- Patient has had a Special Authority approval for zoledronic acid (Underlying cause – Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

**RISEDRONATE SODIUM**

Tab 35 mg .....3.10 4 ✓ **Risedronate Sandoz**

**TERIPARATIDE – Special Authority see SA1139 below – Retail pharmacy**

Inj 250 mcg per ml, 2.4 ml .....490.00 1 ✓ **Forteo**

►SA1139 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- The patient has severe, established osteoporosis; and
- The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- The patient has had two or more fractures due to minimal trauma; and
- The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

## ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, vial – Special Authority see

SA1780 below – Retail pharmacy ..... 60.00      100 ml OP      ✓ **Aclasta**

### ►SA1780 Special Authority for Subsidy

**Initial application — (Paget's disease)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
  - 2.1 Bone or articular pain; or
  - 2.2 Bone deformity; or
  - 2.3 Bone, articular or neurological complications; or
  - 2.4 Asymptomatic disease, but risk of complications; or
  - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

**Initial application — (Underlying cause - Osteoporosis)** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
  - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
  - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
  - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
  - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
  - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

**Initial application — (Underlying cause - glucocorticosteroid therapy)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
  - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
- 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause - glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

**Renewal — (Paget's disease)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
  - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
  - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

**Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

**Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria)** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
  - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
  - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
  - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
  - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
  - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause - Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

-2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.

- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

## Hyperuricaemia and Antigout

### ALLOPURINOL

* Tab 100 mg .....	4.54	500	✓ <b>DP-Allopurinol</b>
* Tab 300 mg .....	10.35	500	✓ <b>DP-Allopurinol</b>

### BENZBROMARONE – Special Authority see [SA1537 below](#) – Retail pharmacy

Tab 100 mg .....	45.00	100	✓ <b>Benzbromaron AL</b> 100 <small>\$29</small>
------------------	-------	-----	---

### ► [SA1537](#) Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
  - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
  - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
  - 2.3 Both:
    - 2.3.1 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Notes); and
    - 2.3.2 The patient has a rate of creatinine clearance greater than or equal to 20 ml/min; or
  - 2.4 All of the following:
    - 2.4.1 The patient is taking azathioprine and requires urate-lowering therapy; and
    - 2.4.2 Allopurinol is contraindicated; and
    - 2.4.3 Appropriate doses of probenecid are ineffective or probenecid cannot be used due to reduced renal function; and
- 3 The patient is receiving monthly liver function tests.

**Renewal** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

Notes: Benzbromarone has been associated with potentially fatal hepatotoxicity.

In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

The New Zealand Rheumatology Association has developed information for prescribers which can be accessed from its website at [www.rheumatology.org.nz/home/resources-2/](http://www.rheumatology.org.nz/home/resources-2/)

### COLCHICINE

* Tab 500 mcg .....	9.58	100	✓ <b>Colgout</b>
---------------------	------	-----	------------------

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
FEBUXOSTAT – Special Authority see SA1538 below – Retail pharmacy				
Tab 80 mg .....	39.50	28	✓	Adenuric
Tab 120 mg .....	39.50	28	✓	Adenuric

►SA1538 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
  - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
  - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
  - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note).

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

PROBENECID

* Tab 500 mg .....	55.00	100	✓	Probenecid-AFT
--------------------	-------	-----	---	----------------

## Muscle Relaxants

BACLOFEN

* Tab 10 mg .....	4.20	100	✓	Pacifen
Inj 0.05 mg per ml, 1 ml ampoule – Subsidy by endorsement.....	11.55	1	✓	Lioresal Intrathecal
Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have caused intolerable side effects and the prescription is endorsed accordingly.				
Inj 2 mg per ml, 5 ml ampoule – Subsidy by endorsement.....	372.98	5	✓	Medsurge
Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have caused intolerable side effects and the prescription is endorsed accordingly.				

DANTROLENE

Cap 25 mg .....	65.00	100	✓	Dantrium
Cap 50 mg .....	77.00	100	✓	Dantrium S29 <sup>S29</sup>
			✓	Dantrium

ORPHENADRINE CITRATE

Tab 100 mg .....	18.54	100	✓	Norflex
------------------	-------	-----	---	---------

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

**Agents for Parkinsonism and Related Disorders**
**Dopamine Agonists and Related Agents**
**AMANTADINE HYDROCHLORIDE**

▲ Cap 100 mg ..... 38.24      60      ✓ **Symmetrel**

**APOMORPHINE HYDROCHLORIDE**

▲ Inj 10 mg per ml, 2 ml ampoule ..... 119.00      5      ✓ **Movapo**

**BROMOCRIPTINE MESYLATE**

\* Tab 2.5 mg ..... 32.08      100      ✓ **Apo-Bromocriptine**

**ENTACAPONE**

▲ Tab 200 mg ..... 22.00      100      ✓ **Entapone**

**LEVODOPA WITH BENSERAZIDE**

\* Tab dispersible 50 mg with benserazide 12.5 mg ..... 13.25      100      ✓ **Madopar Rapid**

\* Cap 50 mg with benserazide 12.5 mg ..... 13.75      100      ✓ **Madopar 62.5**

\* Cap 100 mg with benserazide 25 mg ..... 15.80      100      ✓ **Madopar 125**

\* Cap long-acting 100 mg with benserazide 25 mg ..... 22.85      100      ✓ **Madopar HBS**

\* Cap 200 mg with benserazide 50 mg ..... 26.25      100      ✓ **Madopar 250**

**LEVODOPA WITH CARBIDOPA**

\* Tab 100 mg with carbidopa 25 mg ..... 17.97      100      ✓ **Kinson**

✓ **Sinemet**

\* Tab long-acting 100 mg with carbidopa 25 mg ..... 23.84      100      ✓ **Mylan <sup>S29</sup>**

\* Tab long-acting 200 mg with carbidopa 50 mg ..... 37.15      100      ✓ **Sinemet CR**

46.73      ✓ **Mylan <sup>S29</sup>**

\* Tab 250 mg with carbidopa 25 mg ..... 32.67      100      ✓ **Sinemet**

**PRAMIPEXOLE HYDROCHLORIDE**

▲ Tab 0.25 mg ..... 6.12      100      ✓ **Ramipex**

▲ Tab 1 mg ..... 20.73      100      ✓ **Ramipex**

**ROPINIROLE HYDROCHLORIDE**

▲ Tab 0.25 mg ..... 2.78      100      ✓ **Apo-Ropinirole**

2.85      84      ✓ **Ropin**

▲ Tab 1 mg ..... 3.95      84      ✓ **Ropin**

5.00      100      ✓ **Apo-Ropinirole**

▲ Tab 2 mg ..... 5.48      84      ✓ **Ropin**

7.72      100      ✓ **Apo-Ropinirole**

▲ Tab 5 mg ..... 12.50      84      ✓ **Ropin**

16.51      100      ✓ **Apo-Ropinirole**

*(Apo-Ropinirole Tab 0.25 mg to be delisted 1 March 2020)*

*(Apo-Ropinirole Tab 1 mg to be delisted 1 March 2020)*

*(Apo-Ropinirole Tab 2 mg to be delisted 1 March 2020)*

*(Apo-Ropinirole Tab 5 mg to be delisted 1 March 2020)*

**SELEGILINE HYDROCHLORIDE**

\* Tab 5 mg ..... 22.00      100      ✓ **Apo-Selegiline**  
**S29 <sup>S29</sup>**

**TOLCAPONE**

▲ Tab 100 mg ..... 132.50      100      ✓ **Tasmar**

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>Anticholinergics</b>				
<b>BENZATROPINE MESYLATE</b>				
Tab 2 mg .....	7.99	60	✓	<b>Benztrop</b>
Inj 1 mg per ml, 2 ml .....	95.00	5	✓	<b>Cogentin</b>
	190.00	10	✓	<b>Omega</b>
a) Up to 10 inj available on a PSO				
b) Only on a PSO				
<b>PROCYCLIDINE HYDROCHLORIDE</b>				
Tab 5 mg .....	7.40	100	✓	<b>Kemadrin</b>
<b>Agents for Essential Tremor, Chorea and Related Disorders</b>				
<b>RILUZOLE</b> – Special Authority see <a href="#">SA1403 below</a> – Retail pharmacy				
Wastage claimable				
Tab 50 mg .....	130.00	56	✓	<b>Rilutek</b>
<b>►SA1403 Special Authority for Subsidy</b>				
<b>Initial application</b> only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria:				
All of the following:				
1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and				
2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and				
3 The patient has not undergone a tracheostomy; and				
4 The patient has not experienced respiratory failure; and				
5 Any of the following:				
5.1 The patient is ambulatory; or				
5.2 The patient is able to use upper limbs; or				
5.3 The patient is able to swallow.				
<b>Renewal</b> from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria:				
All of the following:				
1 The patient has not undergone a tracheostomy; and				
2 The patient has not experienced respiratory failure; and				
3 Any of the following:				
3.1 The patient is ambulatory; or				
3.2 The patient is able to use upper limbs; or				
3.3 The patient is able to swallow.				
<b>TETRABENAZINE</b>				
Tab 25 mg .....	91.10	112	✓	<b>Motetis</b>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

## Anaesthetics

### Local

#### LIDOCAINE [LIGNOCAINE]

Gel 2%, tube – Subsidy by endorsement.....	14.50	30 ml	✓ <b>Xylocaine 2% Jelly</b>
a) Up to 150 ml available on a PSO			
b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.			
Gel 2%, 10 ml urethral syringe – Subsidy by endorsement.....	81.50	10	✓ <b>Pfizer</b>
	105.00	25	✓ <b>Cathejell</b>
a) Up to 5 each available on a PSO			
b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.			
c) Cathejell to be Sole Supply on 1 November 2019			

(Pfizer Gel 2%, 10 ml urethral syringe to be delisted 1 November 2019)

#### LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE

Oral (gel) soln 2%.....	38.00	200 ml	✓ <b>Mucosoothe</b>
Inj 1%, 5 ml ampoule – Up to 25 inj available on a PSO.....	8.75	25	✓ <b>Lidocaine-Clarix</b>
	17.50	50	
	(35.00)		Xylocaine
Inj 2%, 5 ml ampoule – Up to 5 inj available on a PSO.....	8.25	25	✓ <b>Lidocaine-Clarix</b>
Lidocaine-Clarix to be Sole Supply on 1 November 2019			
Inj 1%, 20 ml ampoule – Up to 5 inj available on a PSO.....	12.00	5	
	(20.00)		Xylocaine
Inj 1%, 20 ml vial – Up to 5 inj available on a PSO.....	6.20	5	✓ <b>Lidocaine-Clarix</b>
Inj 2%, 20 ml vial – Up to 5 inj available on a PSO.....	6.45	5	✓ <b>Lidocaine-Clarix</b>

#### LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE

Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –			
Subsidy by endorsement.....	81.50	10	✓ <b>Pfizer</b>
a) Up to 5 each available on a PSO			
b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.			

### Topical Local Anaesthetics

#### ►SA0906 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] – Special Authority see SA0906 above – Retail pharmacy

Crm 4%.....	5.40	5 g OP	✓ <b>LMX4</b>
	27.00	30 g OP	✓ <b>LMX4</b>

LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 above – Retail pharmacy

Crm 2.5% with prilocaine 2.5%.....	45.00	30 g OP	✓ <b>EMLA</b>
Crm 2.5% with prilocaine 2.5% (5 g tubes) .....	45.00	5	✓ <b>EMLA</b>



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

## Analgesics

For Anti-inflammatory NSAIDs refer to MUSCULOSKELETAL, [page 110](#)

### Non-opioid Analgesics

For aspirin & chloroform application refer Standard Formulae, [page 235](#)

#### ASPIRIN

\* Tab dispersible 300 mg – Up to 30 tab available on a PSO ..... 4.50 100 ✓ Ethics Aspirin

#### CAPSAICIN – Subsidy by endorsement

Subsidised only if prescribed for post-herpetic neuralgia or diabetic peripheral neuropathy and the prescription is endorsed accordingly.

Crm 0.075% ..... 12.50 45 g OP ✓ Zostrix HP

#### NEFOPAM HYDROCHLORIDE

Tab 30 mg ..... 23.40 90 ✓ Acupan

#### PARACETAMOL

\* Tab 500 mg - blister pack – Up to 30 tab available on a PSO ..... 7.12 1,000 ✓ Paracetamol Pharmacare  
✓ Pharmacare  
✓ Pharmacy Health  
✓ Pharmacare  
✓ Paracare

\* Tab 500 mg - bottle pack ..... 6.32 1,000

\* Oral liq 120 mg per 5 ml ..... 5.35 1,000 ml

a) Up to 200 ml available on a PSO

b) Not in combination

\* Oral liq 250 mg per 5 ml ..... 5.81 1,000 ml ✓ Paracare Double Strength

a) Up to 100 ml available on a PSO

b) Not in combination

\* Suppos 125 mg ..... 3.29 10 ✓ Gacet

\* Suppos 250 mg ..... 3.79 10 ✓ Gacet

\* Suppos 500 mg ..... 12.40 50 ✓ Gacet

(Pharmacy Health Tab 500 mg - blister pack to be delisted 1 January 2020)

### Opioid Analgesics

CODEINE PHOSPHATE – Safety medicine; prescriber may determine dispensing frequency

Tab 15 mg ..... 5.75 100 ✓ PSM

Tab 30 mg ..... 6.80 100 ✓ PSM

Tab 60 mg ..... 13.50 100 ✓ PSM

#### DIHYDROCODEINE TARTRATE

Tab long-acting 60 mg ..... 8.60 60 ✓ DHC Continus

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>FENTANYL</b>				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency				
Inj 50 mcg per ml, 2 ml ampoule .....	3.56	10	✓	<b>Boucher and Muir</b>
Inj 50 mcg per ml, 10 ml ampoule .....	9.41	10	✓	<b>Boucher and Muir</b>
Patch 12.5 mcg per hour .....	2.95	5	✓	<b>Fentanyl Sandoz</b>
Patch 25 mcg per hour .....	3.66	5	✓	<b>Fentanyl Sandoz</b>
Patch 50 mcg per hour .....	6.65	5	✓	<b>Fentanyl Sandoz</b>
Patch 75 mcg per hour .....	9.25	5	✓	<b>Fentanyl Sandoz</b>
Patch 100 mcg per hour .....	11.40	5	✓	<b>Fentanyl Sandoz</b>
<b>METHADONE HYDROCHLORIDE</b>				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency				
d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets).				
e) For methadone hydrochloride oral liquid refer Standard Formulae, <a href="#">page 235</a>				
Tab 5 mg .....	1.40	10	✓	<b>Methatabs</b>
Tab 5 mg - bottle pack .....	1.40	10	✓	<b>Methatabs</b>
Oral liq 2 mg per ml .....	5.79	200 ml	✓	<b>Biodone</b>
Oral liq 5 mg per ml .....	5.79	200 ml	✓	<b>Biodone Forte</b>
Oral liq 10 mg per ml .....	6.79	200 ml	✓	<b>Biodone Extra Forte</b>
Inj 10 mg per ml, 1 ml .....	61.00	10	✓	<b>AFT</b>
<i>(Methatabs Tab 5 mg - bottle pack to be delisted 1 December 2019)</i>				
<b>MORPHINE HYDROCHLORIDE</b>				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency				
Oral liq 1 mg per ml .....	9.28	200 ml	✓	<b>RA-Morph</b>
Oral liq 2 mg per ml .....	16.24	200 ml	✓	<b>RA-Morph</b>
Oral liq 5 mg per ml .....	19.44	200 ml	✓	<b>Ordine <sup>S29</sup></b>
			✓	<b>RA-Morph</b>
Oral liq 10 mg per ml .....	27.74	200 ml	✓	<b>Ordine <sup>S29</sup></b>
			✓	<b>RA-Morph</b>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>MORPHINE SULPHATE</b>				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency				
Tab immediate-release 10 mg.....	2.80	10	✓	<u>Sevredol</u>
Tab long-acting 10 mg.....	1.93	10	✓	<u>Arrow-Morphine LA</u>
Tab immediate-release 20 mg.....	5.52	10	✓	<u>Sevredol</u>
Tab long-acting 30 mg.....	2.85	10	✓	<u>Arrow-Morphine LA</u>
Tab long-acting 60 mg.....	5.60	10	✓	<u>Arrow-Morphine LA</u>
Tab long-acting 100 mg.....	6.10	10	✓	<u>Arrow-Morphine LA</u>
Cap long-acting 10 mg.....	2.05	10	✓	<u>m-Eslon</u>
m-Eslon to be Sole Supply on 1 January 2020				
Cap long-acting 30 mg.....	3.00	10	✓	<u>m-Eslon</u>
m-Eslon to be Sole Supply on 1 January 2020				
Cap long-acting 60 mg.....	6.12	10	✓	<u>m-Eslon</u>
m-Eslon to be Sole Supply on 1 January 2020				
Cap long-acting 100 mg.....	7.13	10	✓	<u>m-Eslon</u>
m-Eslon to be Sole Supply on 1 January 2020				
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO .....	6.27	5	✓	<u>DBL Morphine Sulphate</u>
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO .....	4.47	5	✓	<u>DBL Morphine Sulphate</u>
Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO .....	4.76	5	✓	<u>DBL Morphine Sulphate</u>
Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO .....	6.19	5	✓	<u>DBL Morphine Sulphate</u>
<b>MORPHINE TARTRATE</b>				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency				
Inj 80 mg per ml, 1.5 ml ampoule .....	42.72	5	✓	<u>DBL Morphine Tartrate</u>
<b>OXYCODONE HYDROCHLORIDE</b>				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency				
Tab controlled-release 5 mg.....	2.15	20	✓	<u>Oxycodone Sandoz</u>
Tab controlled-release 10 mg.....	2.15	20	✓	<u>Oxycodone Sandoz</u>
Tab controlled-release 20 mg.....	2.15	20	✓	<u>Oxycodone Sandoz</u>
Tab controlled-release 40 mg.....	3.20	20	✓	<u>Oxycodone Sandoz</u>
Tab controlled-release 80 mg.....	10.98	20	✓	<u>Oxycodone Sandoz</u>
Cap immediate-release 5 mg.....	1.88	20	✓	<u>OxyNorm</u>
Cap immediate-release 10 mg.....	3.32	20	✓	<u>OxyNorm</u>
Cap immediate-release 20 mg.....	5.81	20	✓	<u>OxyNorm</u>
Oral liq 5 mg per 5 ml.....	11.20	250 ml	✓	<u>OxyNorm</u>
Inj 10 mg per ml, 1 ml ampoule .....	7.28	5	✓	<u>OxyNorm</u>
Inj 10 mg per ml, 2 ml ampoule .....	14.36	5	✓	<u>OxyNorm</u>
Inj 50 mg per ml, 1 ml ampoule .....	30.60	5	✓	<u>OxyNorm</u>
<b>PARACETAMOL WITH CODEINE – Safety medicine; prescriber may determine dispensing frequency</b>				
* Tab paracetamol 500 mg with codeine phosphate 8 mg.....	18.21	1,000	✓	<u>Paracetamol + Codeine (Relieve)</u>

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\* Three months or six months, as applicable, dispensed all-at-once

## NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>PETHIDINE HYDROCHLORIDE</b>				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency				
Tab 50 mg .....	4.46	10	✓	<b>PSM</b>
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO .....	4.98	5	✓	<b>DBL Pethidine Hydrochloride</b>
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO .....	5.12	5	✓	<b>DBL Pethidine Hydrochloride</b>
<b>TRAMADOL HYDROCHLORIDE</b>				
Tab sustained-release 100 mg .....	1.55	20	✓	<b>Tramal SR 100</b>
Tab sustained-release 150 mg .....	2.10	20	✓	<b>Tramal SR 150</b>
Tab sustained-release 200 mg .....	2.75	20	✓	<b>Tramal SR 200</b>
Cap 50 mg .....	2.25	100	✓	<b>Arrow-Tramadol</b>

## Antidepressants

### Cyclic and Related Agents

<b>AMITRIPTYLINE</b> – Safety medicine; prescriber may determine dispensing frequency				
Tab 10 mg .....	1.96	100	✓	<b>Arrow-Amitriptyline</b>
Tab 25 mg .....	1.52	100	✓	<b>Arrow-Amitriptyline</b>
Tab 50 mg .....	2.51	100	✓	<b>Arrow-Amitriptyline</b>
<b>CLOMIPRAMINE HYDROCHLORIDE</b> – Safety medicine; prescriber may determine dispensing frequency				
Tab 10 mg .....	13.99	100	✓	<b>Apo-Clomipramine</b>
Tab 25 mg .....	4.73	50	✓	<b>Apo-Clomipramine</b>
	9.46	100	✓	<b>Apo-Clomipramine</b>
<b>DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE</b> – Subsidy by endorsement				
a) Safety medicine; prescriber may determine dispensing frequency				
b) Subsidy by endorsement – Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 June 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.				
Tab 75 mg .....	11.19	100	✓	<b>Dopress</b>
Cap 25 mg .....	6.45	100	✓	<b>Dopress</b>
<i>(Dopress Tab 75 mg to be delisted 1 August 2020)</i>				
<i>(Dopress Cap 25 mg to be delisted 1 January 2020)</i>				
<b>DOXEPIN HYDROCHLORIDE</b> – Subsidy by endorsement				
a) Safety medicine; prescriber may determine dispensing frequency				
b) Subsidy by endorsement – Subsidised for patients who were taking doxepin hydrochloride prior to 1 March 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of doxepin hydrochloride.				
Cap 10 mg .....	6.30	100	✓	<b>Anten</b>
Cap 25 mg .....	6.86	100	✓	<b>Anten</b>
Cap 50 mg .....	8.55	100	✓	<b>Anten</b>
<i>(Anten Cap 10 mg to be delisted 1 January 2020)</i>				
<i>(Anten Cap 25 mg to be delisted 1 April 2020)</i>				
<i>(Anten Cap 50 mg to be delisted 1 May 2020)</i>				
<b>IMIPRAMINE HYDROCHLORIDE</b> – Safety medicine; prescriber may determine dispensing frequency				
Tab 10 mg .....	5.48	50	✓	<b>Tofranil</b>
	10.96	100	✓	<b>Tofranil</b>
Tab 25 mg .....	8.80	50	✓	<b>Tofranil</b>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
MAPROTIline HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency				
Tab 25 mg .....	7.52	30	✓	Ludiomil
	12.53	50	✓	Ludiomil
	25.06	100	✓	Ludiomil
Tab 75 mg .....	14.01	20	✓	Ludiomil
	21.01	30	✓	Ludiomil
NORTRIPTYLINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency				
Tab 10 mg .....	2.44	100	✓	Norpress
Tab 25 mg .....	5.98	180	✓	Norpress

### Monoamine-Oxidase Inhibitors (MAOIs) - Non Selective

PHENELZINE SULPHATE				
* Tab 15 mg .....	70.80	60	✓	Nardil S29
	118.00	100	✓	Nardil
TRANLYCYPROMINE SULPHATE				
* Tab 10 mg .....	12.85	28	✓	Parnate S29
	22.94	50	✓	Parnate
	96.00	100	✓	Parnate S29

### Monoamine-Oxidase Type A Inhibitors

MOCLOBEMIDE				
* Tab 150 mg .....	6.40	60	✓	Aurorix
* Tab 300 mg .....	9.80	60	✓	Aurorix

### Selective Serotonin Reuptake Inhibitors

CITALOPRAM HYDROBROMIDE				
* Tab 20 mg .....	1.52	84	✓	PSM Citalopram
ESCITALOPRAM				
* Tab 10 mg .....	1.11	28	✓	Escitalopram- Apotex
* Tab 20 mg .....	1.90	28	✓	Escitalopram- Apotex
FLUOXETINE HYDROCHLORIDE				
* Tab dispersible 20 mg, scored – Subsidy by endorsement .....	2.47	30	✓	Arrow-Fluoxetine
Subsidised by endorsement				
1) When prescribed for a patient who cannot swallow whole tablets or capsules and the prescription is endorsed accordingly; or				
2) When prescribed in a daily dose that is not a multiple of 20 mg in which case the prescription is deemed to be endorsed. Note: Tablets should be combined with capsules to facilitate incremental 10 mg doses.				
* Cap 20 mg .....	1.99	90	✓	Arrow-Fluoxetine
PAROXETINE				
* Tab 20 mg .....	3.61	90	✓	Loxamine
	4.02		✓	Apo-Paroxetine

(Apo-Paroxetine Tab 20 mg to be delisted 1 March 2020)

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\* Three months or six months, as applicable, dispensed all-at-once

## NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
SERTRALINE				
* Tab 50 mg .....	0.92	30	✓	Setrona
	3.05	90	✓	Arrow-Sertraline
* Tab 100 mg .....	1.61	30	✓	Setrona
	5.25	90	✓	Arrow-Sertraline

(Arrow-Sertraline Tab 50 mg to be delisted 1 March 2020)

(Arrow-Sertraline Tab 100 mg to be delisted 1 March 2020)

### Other Antidepressants

MIRTAZAPINE				
Tab 30 mg .....	2.63	30	✓	Apo-Mirtazapine
Tab 45 mg .....	3.48	30	✓	Apo-Mirtazapine
VENLAFAXINE				
* Cap 37.5 mg .....	6.38	84	✓	Enlafax XR
* Cap 75 mg .....	8.11	84	✓	Enlafax XR
* Cap 150 mg .....	11.16	84	✓	Enlafax XR

### Antiepilepsy Drugs

#### Agents for Control of Status Epilepticus

CLONAZEPAM – Safety medicine; prescriber may determine dispensing frequency				
Inj 1 mg per ml, 1 ml .....	21.00	5	✓	Rivotril
DIAZEPAM – Safety medicine; prescriber may determine dispensing frequency				
Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement .....	11.83	5	✓	Hospira
a) Up to 5 inj available on a PSO				
b) Only on a PSO				
c) PSO must be endorsed "not for anaesthetic procedures".				
Rectal tubes 5 mg – Up to 5 tube available on a PSO .....	40.87	5	✓	Stesolid
Rectal tubes 10 mg – Up to 5 tube available on a PSO .....	40.87	5	✓	Stesolid
PARALDEHYDE				
* Inj 5 ml .....	1,500.00	5	✓	AFT <sup>\$29</sup>
PHENYTOIN SODIUM				
* Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO ....	88.63	5	✓	Hospira
* Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a PSO .....	133.92	5	✓	Hospira

### Control of Epilepsy

CARBAMAZEPINE				
* Tab 200 mg .....	14.53	100	✓	Tegretol
* Tab long-acting 200 mg .....	16.98	100	✓	Tegretol CR
* Tab 400 mg .....	34.58	100	✓	Tegretol
* Tab long-acting 400 mg .....	39.17	100	✓	Tegretol CR
* Oral liq 20 mg per ml .....	26.37	250 ml	✓	Tegretol
CLOBAZAM – Safety medicine; prescriber may determine dispensing frequency				
Tab 10 mg .....	9.12	50	✓	Frisium
CLONAZEPAM – Safety medicine; prescriber may determine dispensing frequency				
Oral drops 2.5 mg per ml .....	7.38	10 ml OP	✓	Rivotril

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>ETHOSUXIMIDE</b>				
Cap 250 mg .....	140.88	100	✓	<b>Zarontin</b>
Oral liq 250 mg per 5 ml .....	56.35	200 ml	✓	<b>Zarontin</b>
<b>GABAPENTIN</b>				
Note: Not subsidised in combination with subsidised pregabalin				
* Cap 100 mg .....	2.65	100	✓	<b>Apo-Gabapentin</b>
* Cap 300 mg .....	4.07	100	✓	<b>Apo-Gabapentin</b>
* Cap 400 mg .....	5.64	100	✓	<b>Apo-Gabapentin</b>
<b>LACOSAMIDE</b> – Special Authority see <a href="#">SA1125 below</a> – Retail pharmacy				
▲ Tab 50 mg .....	25.04	14	✓	<b>Vimpat</b>
▲ Tab 100 mg .....	50.06	14	✓	<b>Vimpat</b>
	200.24	56	✓	<b>Vimpat</b>
▲ Tab 150 mg .....	75.10	14	✓	<b>Vimpat</b>
	300.40	56	✓	<b>Vimpat</b>
▲ Tab 200 mg .....	400.55	56	✓	<b>Vimpat</b>
<b>►SA1125 Special Authority for Subsidy</b>				
<b>Initial application</b> from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:				
Both:				
1 Patient has partial-onset epilepsy; and				
2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).				
Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.				
<b>Renewal</b> from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).				
Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.				
<b>LAMOTRIGINE</b>				
▲ Tab dispersible 2 mg .....	6.74	30	✓	<b>Lamictal</b>
▲ Tab dispersible 5 mg .....	9.64	30	✓	<b>Lamictal</b>
	15.00	56	✓	<b>Arrow-Lamotrigine</b>
▲ Tab dispersible 25 mg – Brand switch fee payable (Pharmacode 2575949) - see <a href="#">page 233</a> for details .....	2.76	56	✓	<b>Logem</b>
▲ Tab dispersible 50 mg – Brand switch fee payable (Pharmacode 2575949) - see <a href="#">page 233</a> for details .....	3.31	56	✓	<b>Logem</b>
▲ Tab dispersible 100 mg – Brand switch fee payable (Pharmacode 2575949) - see <a href="#">page 233</a> for details .....	4.40	56	✓	<b>Logem</b>
<b>LEVETIRACETAM</b>				
Tab 250 mg .....	4.99	60	✓	<b>Everet</b>
Tab 500 mg .....	8.79	60	✓	<b>Everet</b>
Tab 750 mg .....	14.39	60	✓	<b>Everet</b>
Tab 1,000 mg .....	18.59	60	✓	<b>Everet</b>
Oral liq 100 mg per ml .....	44.78	300 ml OP	✓	<b>Levetiracetam-AFT</b>
<b>PHENOBARBITONE</b>				
For phenobarbitone oral liquid refer Standard Formulae, <a href="#">page 235</a>				
* Tab 15 mg .....	40.00	500	✓	<b>PSM</b>
* Tab 30 mg .....	40.00	500	✓	<b>PSM</b>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>PHENYTOIN SODIUM</b>				
* Tab 50 mg .....	75.00	200	✓	Dilantin Infatab
Cap 30 mg .....	74.00	200	✓	Dilantin
Cap 100 mg .....	37.00	200	✓	Dilantin
* Oral liq 30 mg per 5 ml .....	22.03	500 ml	✓	Dilantin
<b>PREGABALIN</b>				
Note: Not subsidised in combination with subsidised gabapentin				
* Cap 25 mg .....	2.25	56	✓	Pregabalin Pfizer
* Cap 75 mg .....	2.65	56	✓	Pregabalin Pfizer
* Cap 150 mg .....	4.01	56	✓	Pregabalin Pfizer
* Cap 300 mg .....	7.38	56	✓	Pregabalin Pfizer
<b>PRIMIDONE</b>				
* Tab 250 mg .....	17.25	100	✓	Apo-Primidone
	62.00	200	✓	Mysoline S29
<b>SODIUM VALPROATE</b>				
Tab 100 mg .....	13.65	100	✓	Epilim Crushable
Tab 200 mg EC .....	27.44	100	✓	Epilim
Tab 500 mg EC .....	52.24	100	✓	Epilim
* Oral liq 200 mg per 5 ml .....	20.48	300 ml	✓	Epilim S/F Liquid
			✓	Epilim Syrup
			✓	Epilim IV
* Inj 100 mg per ml, 4 ml .....	41.50	1		
<b>STIRIPENTOL</b> – Special Authority see <a href="#">SA1330 below</a> – Retail pharmacy				
Cap 250 mg .....	509.29	60	✓	Diacomit S29
Powder for oral liq 250 mg sachet .....	509.29	60	✓	Diacomit S29
➔ <b>SA1330 Special Authority for Subsidy</b>				
<b>Initial application</b> only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist.				
Approvals valid for 6 months for applications meeting the following criteria:				
Both:				
1 Patient has confirmed diagnosis of Dravet syndrome; and				
2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.				
<b>Renewal</b> from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.				
<b>TOPIRAMATE</b>				
▲ Tab 25 mg .....	11.07	60	✓	Arrow-Topiramate
	26.04		✓	Topiramate Actavis
			✓	Topamax
▲ Tab 50 mg .....	18.81	60	✓	Arrow-Topiramate
	44.26		✓	Topiramate Actavis
			✓	Topamax
▲ Tab 100 mg .....	31.99	60	✓	Arrow-Topiramate
	75.25		✓	Topiramate Actavis
			✓	Topamax
▲ Tab 200 mg .....	55.19	60	✓	Arrow-Topiramate
	129.85		✓	Topiramate Actavis
			✓	Topamax
▲ Sprinkle cap 15 mg .....	20.84	60	✓	Topamax
▲ Sprinkle cap 25 mg .....	26.04	60	✓	Topamax
<b>VIGABATRIN</b> – Special Authority see <a href="#">SA1072 on the next page</a> – Retail pharmacy				
▲ Tab 500 mg .....	119.30	100	✓	Sabril



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

### ►SA1072 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Patient has infantile spasms; or
  - 1.2 Both:
    - 1.2.1 Patient has epilepsy; and
    - 1.2.2 Either:
      - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
      - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and
- 2 Either:
  - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Notes: "Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

**Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
  - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

## Antimigraine Preparations

For Anti-inflammatory NSAIDs refer to MUSCULOSKELETAL, [page 110](#)

### Acute Migraine Treatment

#### ERGOTAMINE TARTRATE WITH CAFFEINE

Tab 1 mg with caffeine 100 mg .....	31.00	100	✓ <b>Cafergot</b>
			✓ <b>Cafergot S29</b> <small>S29</small>

#### RIZATRIPTAN

Tab orodispersible 10 mg .....	5.26	30	✓ <b>Rizamelt</b>
--------------------------------	------	----	-------------------

#### SUMATRIPTAN

Tab 50 mg .....	24.44	100	✓ <b>Apo-Sumatriptan</b>
Tab 100 mg .....	46.23	100	✓ <b>Apo-Sumatriptan</b>
Inj 12 mg per ml, 0.5 ml prefilled pen – Maximum of 10 inj per prescription .....	42.67	2 OP	✓ <b>Sun Pharma</b> <small>S29</small>
	81.15		✓ <b>Clustran</b>

## Prophylaxis of Migraine

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, [page 48](#)

PIZOTIFEN

* Tab 500 mcg.....	23.21	100	✓ <b>Sandomigran</b>
--------------------	-------	-----	----------------------

## Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, [page 8](#)

APREPITANT – Special Authority see [SA0987 below](#) – Retail pharmacy

Cap 2 x 80 mg and 1 x 125 mg.....	84.00	3 OP	✓ <b>Emend Tri-Pack</b>
-----------------------------------	-------	------	-------------------------

► [SA0987](#) **Special Authority for Subsidy**

**Initial application** from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

* Tab 16 mg .....	2.89	84	✓ <b>Vergo 16</b>
-------------------	------	----	-------------------

CYCLIZINE HYDROCHLORIDE

Tab 50 mg .....	0.55	10	✓ <b>Nausicalm</b>
-----------------	------	----	--------------------

CYCLIZINE LACTATE

Inj 50 mg per ml, 1 ml .....	14.95	5	✓ <b>Nausicalm</b>
------------------------------	-------	---	--------------------

DOMPERIDONE

* Tab 10 mg .....	2.25	100	✓ <b>Pharmacy Health</b>
-------------------	------	-----	--------------------------

HYOSCINE HYDROBROMIDE

* Inj 400 mcg per ml, 1 ml ampoule .....	46.50	5	✓ <b>Hospira</b>
	93.00	10	✓ <b>Martindale</b> <small>§29</small>

Patch 1.5 mg – Special Authority see [SA1387 below](#) – Retail pharmacy.....

14.11	2	✓ <b>Scopoderm TTS</b>
-------	---	------------------------

► [SA1387](#) **Special Authority for Subsidy**

**Initial application** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

**Renewal** from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE

* Tab 10 mg .....	1.30	100	✓ <b>Metoclopramide Actavis 10</b>
-------------------	------	-----	------------------------------------

* Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO .....	9.50	10	✓ <b>Pfizer</b>
	13.56		✓ <b>Link Healthcare</b> <small>§29</small>

Pfizer to be Sole Supply on 1 January 2020

(Link Healthcare §29 Inj 5 mg per ml, 2 ml ampoule to be delisted 1 January 2020)

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>ONDANSETRON</b>				
* Tab 4 mg .....	3.36	50	✓	<b>Apo-Ondansetron</b>
* Tab disp 4 mg – Up to 10 tab available on a PSO .....	0.95	10	✓	<b><u>Ondansetron</u></b> <b><u>ODT-ORLA</u></b>
* Tab 8 mg .....	4.77	50	✓	<b>Apo-Ondansetron</b>
* Tab disp 8 mg – Up to 10 tab available on a PSO .....	1.43	10	✓	<b><u>Ondansetron</u></b> <b><u>ODT-DRLA</u></b>
<b>PROCHLORPERAZINE</b>				
* Tab 3 mg buccal .....	5.97 (15.00)	50		Buccastem
* Tab 5 mg – Up to 30 tab available on a PSO .....	6.35	250	✓	<b><u>Nausafix</u></b>
* Inj 12.5 mg per ml, 1 ml – Up to 5 inj available on a PSO .....	25.81	10	✓	<b><u>Stemetil</u></b>

## Antipsychotics

### General

AMISULPRIDE – Safety medicine; prescriber may determine dispensing frequency

Tab 100 mg .....	5.15	30	✓	<b>Sulprix</b>
Sulprix to be Sole Supply on 1 November 2019				
Tab 200 mg .....	14.96	60	✓	<b>Sulprix</b>
Sulprix to be Sole Supply on 1 November 2019				
Tab 400 mg .....	27.70	60	✓	<b>Sulprix</b>
Oral liq 100 mg per ml .....	65.53	60 ml	✓	<b>Solian</b>

(Solian Oral liq 100 mg per ml to be delisted 1 July 2020)

ARIPIPRAZOLE – Safety medicine; prescriber may determine dispensing frequency

Tab 5 mg .....	17.50	30	✓	<b><u>Aripiprazole Sandoz</u></b>
Tab 10 mg .....	17.50	30	✓	<b><u>Aripiprazole Sandoz</u></b>
Tab 15 mg .....	17.50	30	✓	<b><u>Aripiprazole Sandoz</u></b>
Tab 20 mg .....	17.50	30	✓	<b><u>Aripiprazole Sandoz</u></b>
Tab 30 mg .....	17.50	30	✓	<b><u>Aripiprazole Sandoz</u></b>

CHLORPROMAZINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency

Tab 10 mg – Up to 30 tab available on a PSO .....	14.83	100	✓	<b>Largactil</b>
Largactil to be Sole Supply on 1 January 2020				
Tab 25 mg – Up to 30 tab available on a PSO .....	15.62	100	✓	<b>Largactil</b>
Largactil to be Sole Supply on 1 January 2020				
Tab 100 mg – Up to 30 tab available on a PSO .....	36.73	100	✓	<b>Largactil</b>
Largactil to be Sole Supply on 1 January 2020				
Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO .....	30.79	10	✓	<b>Largactil</b>
Largactil to be Sole Supply on 1 January 2020				

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

# NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>CLOZAPINE – Hospital pharmacy [HP4]</b>				
Safety medicine; prescriber may determine dispensing frequency				
Tab 25 mg .....	5.69	50	✓ Clozaril	
	6.69		✓ Clopine	
	11.36	100	✓ Clozaril	
	13.37		✓ Clopine	
Tab 50 mg .....	8.67	50	✓ Clopine	
	17.33	100	✓ Clopine	
Tab 100 mg .....	14.73	50	✓ Clozaril	
	17.33		✓ Clopine	
	29.45	100	✓ Clozaril	
	34.65		✓ Clopine	
Tab 200 mg .....	34.65	50	✓ Clopine	
	69.30	100	✓ Clopine	
Suspension 50 mg per ml .....	17.33	100 ml	✓ Clopine	
<b>HALOPERIDOL – Safety medicine; prescriber may determine dispensing frequency</b>				
Tab 500 mcg – Up to 30 tab available on a PSO .....	6.23	100	✓ Serenace	
Tab 1.5 mg – Up to 30 tab available on a PSO .....	9.43	100	✓ Serenace	
Tab 5 mg – Up to 30 tab available on a PSO .....	29.72	100	✓ Serenace	
Oral liq 2 mg per ml – Up to 200 ml available on a PSO .....	23.84	100 ml	✓ Serenace	
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO .....	21.55	10	✓ Serenace	
<b>LEVOMEPROMAZINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency</b>				
Inj 25 mg per ml, 1 ml ampoule .....	47.89	10	✓ Wockhard	
<b>LEVOMEPROMAZINE MALEATE – Safety medicine; prescriber may determine dispensing frequency</b>				
Tab 25 mg .....	16.10	100	✓ Nozinan	
Tab 100 mg .....	41.75	100	✓ Nozinan	
<b>LITHIUM CARBONATE – Safety medicine; prescriber may determine dispensing frequency</b>				
Tab 250 mg .....	34.30	500	✓ Lithicarb FC	
Tab long-acting 400 mg .....	19.20	100	✓ Priadel	
Cap 250 mg .....	9.42	100	✓ Douglas	
<b>OLANZAPINE – Safety medicine; prescriber may determine dispensing frequency</b>				
Tab 2.5 mg .....	0.64	28	✓ Zypine	
Tab 5 mg .....	1.15	28	✓ Zypine	
Tab orodispersible 5 mg .....	1.25	28	✓ Zypine ODT	
Tab 10 mg .....	1.65	28	✓ Zypine	
Tab orodispersible 10 mg .....	2.05	28	✓ Zypine ODT	
<b>PERICYAZINE – Safety medicine; prescriber may determine dispensing frequency</b>				
Tab 2.5 mg .....	10.49	84	✓ Neulactil	
	12.49	100	✓ Neulactil	
Tab 10 mg .....	37.34	84	✓ Neulactil	
	44.45	100	✓ Neulactil	
<b>QUETIAPINE – Safety medicine; prescriber may determine dispensing frequency</b>				
Tab 25 mg .....	1.79	90	✓ Quetapel	
Tab 100 mg .....	3.45	90	✓ Quetapel	
Tab 200 mg .....	5.75	90	✓ Quetapel	
Tab 300 mg .....	9.60	90	✓ Quetapel	

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>RISPERIDONE</b> – Safety medicine; prescriber may determine dispensing frequency				
Tab 0.5 mg .....	1.86	60	✓	<b>Actavis</b>
Tab 1 mg .....	2.06	60	✓	<b>Actavis</b>
Tab 2 mg .....	2.29	60	✓	<b>Actavis</b>
Tab 3 mg .....	2.50	60	✓	<b>Actavis</b>
Tab 4 mg .....	3.43	60	✓	<b>Actavis</b>
Oral liq 1 mg per ml .....	7.66	30 ml	✓	<b>Risperon</b>
<b>ZIPRASIDONE</b> – Safety medicine; prescriber may determine dispensing frequency				
Cap 20 mg .....	14.50	60	✓	<b>Zusdone</b>
Cap 40 mg .....	24.70	60	✓	<b>Zusdone</b>
Cap 60 mg .....	33.80	60	✓	<b>Zusdone</b>
Cap 80 mg .....	39.70	60	✓	<b>Zusdone</b>
<b>ZUCLOPENTHIXOL HYDROCHLORIDE</b> – Safety medicine; prescriber may determine dispensing frequency				
Tab 10 mg .....	31.45	100	✓	<b>Clopixol</b>

## Depot Injections

<b>FLUPENTHIXOL DECANOATE</b> – Safety medicine; prescriber may determine dispensing frequency				
Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO .....	13.14	5	✓	<b>Fluanxol</b>
Inj 20 mg per ml, 2 ml – Up to 5 inj available on a PSO .....	20.90	5	✓	<b>Fluanxol</b>
Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO .....	40.87	5	✓	<b>Fluanxol</b>
<b>HALOPERIDOL DECANOATE</b> – Safety medicine; prescriber may determine dispensing frequency				
Inj 50 mg per ml, 1 ml – Up to 5 inj available on a PSO .....	28.39	5	✓	<b>Haldol</b>
Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO .....	55.90	5	✓	<b>Haldol Concentrate</b>
			✓	<b>Haldol</b>
				<b>Decanoate</b> S29
<b>OLANZAPINE</b> – Special Authority see <a href="#">SA1428 below</a> – Retail pharmacy				
Safety medicine; prescriber may determine dispensing frequency				
Inj 210 mg vial .....	252.00	1	✓	<b>Zyprexa Relprevv</b>
Inj 300 mg vial .....	414.00	1	✓	<b>Zyprexa Relprevv</b>
Inj 405 mg vial .....	504.00	1	✓	<b>Zyprexa Relprevv</b>

### ➤SA1428 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:  
Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia; and
  - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

**Note:** The patient should be monitored for post-injection syndrome for at least two hours after each injection.

**PALIPERIDONE** – Special Authority see [SA1429 on the next page](#) – Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency				
Inj 25 mg syringe .....	194.25	1	✓	<b>Invega Sustenna</b>
Inj 50 mg syringe .....	271.95	1	✓	<b>Invega Sustenna</b>
Inj 75 mg syringe .....	357.42	1	✓	<b>Invega Sustenna</b>
Inj 100 mg syringe .....	435.12	1	✓	<b>Invega Sustenna</b>
Inj 150 mg syringe .....	435.12	1	✓	<b>Invega Sustenna</b>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

## ►SA1429 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:  
Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE – Special Authority see SA1427 below – Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Inj 25 mg vial .....	135.98	1	✓ Risperdal Consta
Inj 37.5 mg vial .....	178.71	1	✓ Risperdal Consta
Inj 50 mg vial .....	217.56	1	✓ Risperdal Consta

## ►SA1427 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:  
Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE – Safety medicine; prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml – Up to 5 inj available on a PSO .....	19.80	5	✓ Clopixol
--	-------	---	------------

## Anxiolytics

BUSPIRONE HYDROCHLORIDE

* Tab 5 mg .....	20.23	100	✓ Orion
* Tab 10 mg .....	13.16	100	✓ Orion

CLONAZEPAM – Safety medicine; prescriber may determine dispensing frequency

Tab 500 mcg .....	5.64	100	✓ Paxam
Tab 2 mg .....	10.78	100	✓ Paxam

DIAZEPAM – Safety medicine; prescriber may determine dispensing frequency

Tab 2 mg .....	15.05	500	✓ Arrow-Diazepam
Tab 5 mg .....	16.18	500	✓ Arrow-Diazepam

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>LORAZEPAM – Safety medicine; prescriber may determine dispensing frequency</b>				
Tab 1 mg .....	9.72	250	✓	<b>Ativan</b>
Tab 2.5 mg .....	12.50	100	✓	<b>Ativan</b>
<b>OXAZEPAM – Safety medicine; prescriber may determine dispensing frequency</b>				
Tab 10 mg .....	6.17	100	✓	<b>Ox-Pam</b>
Tab 15 mg .....	8.53	100	✓	<b>Ox-Pam</b>

## Multiple Sclerosis Treatments

**DIMETHYL FUMARATE – Special Authority see SA1559 below – Retail pharmacy**

Wastage claimable

Cap 120 mg .....	520.00	14	✓	<b>Tecfidera</b>
Cap 240 mg .....	2,000.00	56	✓	<b>Tecfidera</b>

### ►SA1559 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website <http://www.pharmac.govt.nz> or:

The coordinator	Phone: 04 460 4990
Multiple Sclerosis Treatment Assessment Committee	Facsimile: 04 916 7571
PHARMAC PO Box 10 254	Email: <a href="mailto:mstaccoordinator@pharmac.govt.nz">mstaccoordinator@pharmac.govt.nz</a>
Wellington	

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

### Entry Criteria

- Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- patients must have:
  - EDSS score 0 - 4.0 and:
    - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
    - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
      - a gadolinium enhancing lesion; or
      - a Diffusion Weighted Imaging positive lesion; or
      - a T2 lesion with associated local swelling; or
      - a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
      - new T2 lesions compared with a previous MR scan; and
- A significant relapse must:
  - be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
  - be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
  - last at least one week;
  - start at least one month after the onset of a previous relapse;

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

continued...

- e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
- f) be distinguishable from the effects of general fatigue; and
- g) not be associated with a fever ( $T > 37.5^{\circ}\text{C}$ ); and
- e) applications must be made by the patient's neurologist or general physician; and
- f) patients must have no previous history of lack of response to dimethyl fumarate; and
- g) patients must have not previously had intolerance to dimethyl fumarate; and
- h) patient must not be co-prescribed beta interferon or glatiramer acetate.

## Stopping Criteria

### Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDS points:
  - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
  - b) 1.0 to 3.0; or
  - c) 1.5 to 3.5; or
  - d) 2.0 to 4.0; or
  - e) 2.5 to 4.5; or
  - f) 3.0 to 4.5; or
  - g) 3.5 to 4.5; or
  - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to dimethyl fumarate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

FINGOLIMOD – Special Authority see [SA1562 below](#) – Retail pharmacy

Wastage claimable

Cap 0.5 mg.....	2,200.00	28	✓ Gilenya
-----------------	----------	----	-----------

### ►SA1562 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website <http://www.pharmac.govt.nz> or:

The coordinator	Phone: 04 460 4990
Multiple Sclerosis Treatment Assessment Committee	Facsimile: 04 916 7571
PHARMAC PO Box 10 254	Email: <a href="mailto:mstaccoordinator@pharmac.govt.nz">mstaccoordinator@pharmac.govt.nz</a>
Wellington	

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

## Entry Criteria

continued...



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
  - a) EDSS score 0 - 4.0 and:
    - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
    - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
      - i) a gadolinium enhancing lesion; or
      - ii) a Diffusion Weighted Imaging positive lesion; or
      - iii) a T2 lesion with associated local swelling; or
      - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
      - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
  - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
  - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
  - c) last at least one week;
  - d) start at least one month after the onset of a previous relapse;
  - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
  - f) be distinguishable from the effects of general fatigue; and
  - g) not be associated with a fever ( $T > 37.5^{\circ}\text{C}$ ); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) patients must have no previous history of lack of response to fingolimod; and
- 7) patients must have not previously had intolerance to fingolimod; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

### Stopping Criteria

#### Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDS points:
  - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
  - b) 1.0 to 3.0; or
  - c) 1.5 to 3.5; or
  - d) 2.0 to 4.0; or
  - e) 2.5 to 4.5; or
  - f) 3.0 to 4.5; or
  - g) 3.5 to 4.5; or
  - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to fingolimod; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
--	---	----------------------------	-------------------------------------

continued...

a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

NATALIZUMAB – Special Authority see [SA1563 below](#) – Retail pharmacy

Inj 20 mg per ml, 15 ml vial.....1,750.00 1 ✓ Tysabri

## ►SA1563 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website <http://www.pharmac.govt.nz> or:

The coordinator Phone: 04 460 4990  
Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571  
PHARMAC PO Box 10 254 Email: [mstaccoordinator@pharmac.govt.nz](mailto:mstaccoordinator@pharmac.govt.nz)  
Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

## Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
  - a) EDSS score 0 - 4.0 and:
    - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
    - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
      - i) a gadolinium enhancing lesion; or
      - ii) a Diffusion Weighted Imaging positive lesion; or
      - iii) a T2 lesion with associated local swelling; or
      - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
      - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
  - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
  - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
  - c) last at least one week;
  - d) start at least one month after the onset of a previous relapse;
  - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
  - f) be distinguishable from the effects of general fatigue; and
  - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) treatment must be initiated and supervised by a neurologist who is registered in the Tysabri Australasian Prescribing Programme operated by the supplier; and
- 7) patients must have no previous history of lack of response to natalizumab; and

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 8) patients must have not previously had intolerance to natalizumab; and
- 9)
  - a) Patient is JC virus negative, or
  - b) Patient is JC virus positive and has given written informed consent acknowledging an understanding of the risk of progressive multifocal leucoencephalopathy (PML) associated with natalizumab
- 10) patient must not be co-prescribed beta interferon or glatiramer acetate.

### Stopping Criteria

#### Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDS points:
  - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
  - b) 1.0 to 3.0; or
  - c) 1.5 to 3.5; or
  - d) 2.0 to 4.0; or
  - e) 2.5 to 4.5; or
  - f) 3.0 to 4.5; or
  - g) 3.5 to 4.5; or
  - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to natalizumab; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier.

Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate.

Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

TERIFLUNOMIDE – Special Authority see [SA1560 below](#) – Retail pharmacy

Wastage claimable

Tab 14 mg .....	1,582.62	28	✓ Aubagio
-----------------	----------	----	-----------

### ►SA1560 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website <http://www.pharmac.govt.nz> or:

The coordinator	Phone: 04 460 4990
Multiple Sclerosis Treatment Assessment Committee	Facsimile: 04 916 7571
PHARMAC PO Box 10 254	Email: <a href="mailto:mstacordinator@pharmac.govt.nz">mstacordinator@pharmac.govt.nz</a>
Wellington	

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

### Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and

continued...

continued...

- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
  - a) EDSS score 0 - 4.0 and:
    - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
    - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
      - i) a gadolinium enhancing lesion; or
      - ii) a Diffusion Weighted Imaging positive lesion; or
      - iii) a T2 lesion with associated local swelling; or
      - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
      - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
  - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
  - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
  - c) last at least one week;
  - d) start at least one month after the onset of a previous relapse;
  - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
  - f) be distinguishable from the effects of general fatigue; and
  - g) not be associated with a fever ( $T > 37.5^{\circ}\text{C}$ ); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) patients must have no previous history of lack of response to teriflunomide; and
- 7) patients must have not previously had intolerance to teriflunomide; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

## Stopping Criteria

Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDS points:
  - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
  - b) 1.0 to 3.0; or
  - c) 1.5 to 3.5; or
  - d) 2.0 to 4.0; or
  - e) 2.5 to 4.5; or
  - f) 3.0 to 4.5; or
  - g) 3.5 to 4.5; or
  - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to teriflunomide; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

## Other Multiple Sclerosis Treatments

GLATIRAMER ACETATE – Special Authority see [SA1808 below](#) – Retail pharmacy

Inj 40 mg prefilled syringe – No patient co-payment payable.....2,275.00 12 ✓ Copaxone

### ►SA1808 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website <http://www.pharmac.govt.nz> or:

The coordinator	Phone: 04 460 4990
Multiple Sclerosis Treatment Assessment Committee	Facsimile: 04 916 7571
PHARMAC PO Box 10 254	Email: <a href="mailto:mstaccordinator@pharmac.govt.nz">mstaccordinator@pharmac.govt.nz</a>
Wellington	

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. As a new Special Authority approval number is required the MSTAC coordinator should be notified of the change and a new prescription provided.

### Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
  - a) EDSS score 0 - 4.0 and:
    - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
    - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
      - i) a gadolinium enhancing lesion; or
      - ii) a Diffusion Weighted Imaging positive lesion; or
      - iii) a T2 lesion with associated local swelling; or
      - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
      - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
  - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
  - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
  - c) last at least one week;
  - d) start at least one month after the onset of a previous relapse;
  - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- f) be distinguishable from the effects of general fatigue; and
- g) not be associated with a fever ( $T > 37.5^{\circ}\text{C}$ ); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
  - a) intolerance to both natalizumab and fingolimod; or
  - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

## Stopping Criteria

### Any of the following:

- 1) Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment.  
Progression of disability is defined as progress by any of the following EDDS Points:
  - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
  - b) 1.0 to 3.0; or
  - c) 1.5 to 3.5; or
  - d) 2.0 to 4.0; or
  - e) 2.5 to 4.5; or
  - f) 3.0 to 4.5; or
  - g) 3.5 to 4.5; or
  - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

INTERFERON BETA-1-ALPHA – Special Authority see [SA1809 below](#) – Retail pharmacy

No patient co-payment payable

Inj 6 million iu prefilled syringe.....	1,170.00	4	✓ Avonex
Injection 6 million iu per 0.5 ml pen injector.....	1,170.00	4	✓ Avonex Pen

### ►SA1809 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website <http://www.pharmac.govt.nz> or:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

The coordinator

Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee

Facsimile: 04 916 7571

PHARMAC PO Box 10 254

Email: [mstaccoordinator@pharmac.govt.nz](mailto:mstaccoordinator@pharmac.govt.nz)

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. As a new Special Authority approval number is required the MSTAC coordinator should be notified of the change and a new prescription provided.

### Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
  - a) EDSS score 0 - 4.0 and:
    - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
    - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
      - i) a gadolinium enhancing lesion; or
      - ii) a Diffusion Weighted Imaging positive lesion; or
      - iii) a T2 lesion with associated local swelling; or
      - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
      - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
  - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
  - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
  - c) last at least one week;
  - d) start at least one month after the onset of a previous relapse;
  - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
  - f) be distinguishable from the effects of general fatigue; and
  - g) not be associated with a fever ( $T > 37.5^{\circ}\text{C}$ ); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
  - a) intolerance to both natalizumab and fingolimod; or
  - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

### Stopping Criteria

#### Any of the following:

- 1) Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment. Progression of disability is defined as progress by any of the following EDDSS Points:

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
  - 1.0 to 3.0; or
  - 1.5 to 3.5; or
  - 2.0 to 4.0; or
  - 2.5 to 4.5; or
  - 3.0 to 4.5; or
  - 3.5 to 4.5; or
  - 4.0 to 4.5.
- increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
  - intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
  - non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

INTERFERON BETA-1-BETA – Special Authority see [SA1810 below](#) – Retail pharmacy

No patient co-payment payable

Inj 8 million iu per 1 ml.....	1,322.89	15	✓ <b>Betaferon</b>
--------------------------------	----------	----	--------------------

## ► [SA1810](#) Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website <http://www.pharmac.govt.nz> or:

The coordinator	Phone: 04 460 4990
Multiple Sclerosis Treatment Assessment Committee	Facsimile: 04 916 7571
PHARMAC PO Box 10 254	Email: <a href="mailto:mtaccordinator@pharmac.govt.nz">mtaccordinator@pharmac.govt.nz</a>
Wellington	

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. As a new Special Authority approval number is required the MSTAC coordinator should be notified of the change and a new prescription provided.

## Entry Criteria

- Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and

continued...



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 3) patients must have:
  - a) EDSS score 0 - 4.0 and:
    - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
    - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
      - i) a gadolinium enhancing lesion; or
      - ii) a Diffusion Weighted Imaging positive lesion; or
      - iii) a T2 lesion with associated local swelling; or
      - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
      - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
  - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
  - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
  - c) last at least one week;
  - d) start at least one month after the onset of a previous relapse;
  - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
  - f) be distinguishable from the effects of general fatigue; and
  - g) not be associated with a fever ( $T > 37.5^{\circ}\text{C}$ ); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
  - a) intolerance to both natalizumab and fingolimod; or
  - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

### Stopping Criteria

#### Any of the following:

- 1) Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment.  
Progression of disability is defined as progress by any of the following EDDS Points:
  - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
  - b) 1.0 to 3.0; or
  - c) 1.5 to 3.5; or
  - d) 2.0 to 4.0; or
  - e) 2.5 to 4.5; or
  - f) 3.0 to 4.5; or
  - g) 3.5 to 4.5; or
  - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on

continued...

continued...

starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

## Sedatives and Hypnotics

MELATONIN – Special Authority see [SA1666 below](#) – Retail pharmacy

Tab modified-release 2 mg – No more than 5 tab per day .....	28.22	30	✓ Circadin
--	-------	----	------------

### ► [SA1666](#) Special Authority for Subsidy

**Initial application** only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)\*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under\*.

**Renewal** only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under\*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with \* are unapproved indications.

MIDAZOLAM – Safety medicine; prescriber may determine dispensing frequency

Inj 1 mg per ml, 5 ml ampoule .....	4.30	10	✓ Midazolam-Clarix
Inj 1 mg per ml, 5 ml plastic ampoule – Up to 10 inj available on a PSO .....	14.90	10	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be endorsed for status epilepticus use only.			
Inj 5 mg per ml, 3 ml ampoule .....	2.50	5	✓ Midazolam-Clarix
Inj 5 mg per ml, 3 ml plastic ampoule – Up to 5 inj available on a PSO .....	11.90	5	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be endorsed for status epilepticus use only.			

NITRAZEPAM – Subsidy by endorsement

- a) Safety medicine; prescriber may determine dispensing frequency
- b) Subsidy by endorsement – subsidised for patients who were taking nitrazepam prior to 1 August 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of nitrazepam in the preceding 12 months.

Tab 5 mg .....	5.22	100	✓ Nitrados
----------------	------	-----	------------

(Nitrados Tab 5 mg to be delisted 1 January 2021)

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
PHENOBARBITONE SODIUM – Special Authority see <a href="#">SA1386</a> below – Retail pharmacy			
Inj 200 mg per ml, 1 ml ampoule.....	30.00	5	✓ Aspen <small>\$29</small>
➔ <a href="#">SA1386</a> Special Authority for Subsidy			
Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:			
Both:			
1 For the treatment of terminal agitation that is unresponsive to other agents; and			
2 The applicant is part of a multidisciplinary team working in palliative care.			
TEMAZEPAM – Safety medicine; prescriber may determine dispensing frequency			
Tab 10 mg .....	1.27	25	✓ Normison
TRIAZOLAM – Safety medicine; prescriber may determine dispensing frequency			
Tab 125 mcg.....	5.10	100	
	(9.85)		Hypam
Tab 250 mcg.....	4.10	100	
	(11.20)		Hypam
ZOPICLONE – Safety medicine; prescriber may determine dispensing frequency			
Tab 7.5 mg .....	9.56	500	✓ Zopiclone Actavis

## Stimulants/ADHD Treatments

ATOMOXETINE – Special Authority see <a href="#">SA1416</a> below – Retail pharmacy			
Cap 10 mg.....	107.03	28	✓ Strattera
Cap 18 mg.....	107.03	28	✓ Strattera
Cap 25 mg.....	107.03	28	✓ Strattera
Cap 40 mg.....	107.03	28	✓ Strattera
Cap 60 mg.....	107.03	28	✓ Strattera
Cap 80 mg.....	139.11	28	✓ Strattera
Cap 100 mg.....	139.11	28	✓ Strattera

### ➔[SA1416](#) Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder) diagnosed according to DSM-IV or ICD 10 criteria; and
- 2 Once-daily dosing; and
- 3 Any of the following:
  - 3.1 Treatment with a subsidised formulation of a stimulant has resulted in the development or worsening of serious adverse reactions or where the combination of subsidised stimulant treatment with another agent would pose an unacceptable medical risk; or
  - 3.2 Treatment with a subsidised formulation of a stimulant has resulted in worsening of co-morbid substance abuse or there is a significant risk of diversion with subsidised stimulant therapy; or
  - 3.3 An effective dose of a subsidised formulation of a stimulant has been trialled and has been discontinued because of inadequate clinical response; or
  - 3.4 Treatment with a subsidised formulation of a stimulant is considered inappropriate because the patient has a history of psychoses or has a first-degree relative with schizophrenia; and
- 4 The patient will not be receiving treatment with atomoxetine in combination with a subsidised formulation of a stimulant, except for the purposes of transitioning from subsidised stimulant therapy to atomoxetine.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: A "subsidised formulation of a stimulant" refers to currently subsidised methylphenidate hydrochloride tablet formulations (immediate-release, sustained-release and extended-release) or dexamfetamine sulphate tablets.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

DEXAMFETAMINE SULFATE – Special Authority see [SA1149 below](#) – Retail pharmacy

a) Only on a controlled drug form				
b) Safety medicine; prescriber may determine dispensing frequency				
Tab 5 mg .....	20.00	100	✓	PSM

## ►►SA1149 Special Authority for Subsidy

**Initial application — (ADHD in patients 5 or over)** only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

**Initial application — (ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

**Initial application — (Narcolepsy)** only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

**Renewal — (ADHD in patients 5 or over)** only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
  - 2.1 Applicant is a paediatrician or psychiatrist; or
  - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

**Renewal — (ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

**Renewal — (Narcolepsy)** only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE – Special Authority see [SA1150 below](#) – Retail pharmacy

a) Only on a controlled drug form				
b) Safety medicine; prescriber may determine dispensing frequency				
Tab immediate-release 5 mg.....	3.20	30	✓	Rubifen
Tab immediate-release 10 mg.....	3.00	30	✓	Ritalin
			✓	Rubifen
Tab immediate-release 20 mg.....	7.85	30	✓	Rubifen
Tab sustained-release 20 mg.....	10.95	30	✓	Rubifen SR
	50.00	100	✓	Ritalin SR

## ►►SA1150 Special Authority for Subsidy

**Initial application — (ADHD in patients 5 or over)** only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

**Initial application — (ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

**Initial application — (Narcolepsy)** only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

**Renewal — (ADHD in patients 5 or over)** only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
  - 2.1 Applicant is a paediatrician or psychiatrist; or
  - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

**Renewal — (ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

**Renewal — (Narcolepsy)** only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

**METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE** – Special Authority see [SA1151 on the next page](#) – Retail pharmacy

a) Only on a controlled drug form

b) Safety medicine; prescriber may determine dispensing frequency

Tab extended-release 18 mg.....	18.20	30	✓ Methylphenidate ER - Teva
	58.96		✓ Concerta
Tab extended-release 27 mg.....	22.00	30	✓ Methylphenidate ER - Teva
	65.44		✓ Concerta
Tab extended-release 36 mg.....	22.40	30	✓ Methylphenidate ER - Teva
	71.93		✓ Concerta
Tab extended-release 54 mg.....	26.40	30	✓ Methylphenidate ER - Teva
	86.24		✓ Concerta
Cap modified-release 10 mg .....	15.60	30	✓ Ritalin LA
Cap modified-release 20 mg .....	20.40	30	✓ Ritalin LA
Cap modified-release 30 mg .....	25.52	30	✓ Ritalin LA
Cap modified-release 40 mg .....	30.60	30	✓ Ritalin LA

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

## ►SA1151 Special Authority for Subsidy

**Initial application** only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
  - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
  - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

**Renewal** only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
  - 2.1 Applicant is a paediatrician or psychiatrist; or
  - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL – Special Authority see SA1126 below – Retail pharmacy

Tab 100 mg .....	64.00	60	✓ Modavigil
------------------	-------	----	-------------

## ►SA1126 Special Authority for Subsidy

**Initial application** only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
  - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
  - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
  - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
  - 3.2 Methylphenidate and dexamfetamine are contraindicated.

**Renewal** only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

## Treatments for Dementia

DONEPEZIL HYDROCHLORIDE

* Tab 5 mg .....	4.34	90	✓ Donepezil-Rex
* Tab 10 mg .....	6.64	90	✓ Donepezil-Rex

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
RIVASTIGMINE – Special Authority see <a href="#">SA1488 below</a> – Retail pharmacy				
Patch 4.6 mg per 24 hour .....	90.00	30	✓	Exelon
Patch 9.5 mg per 24 hour .....	90.00	30	✓	Exelon

► **SA1488** **Special Authority for Subsidy**

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

**Renewal** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

## Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE – Special Authority see [SA1203 below](#) – Retail pharmacy

- a) No patient co-payment payable
  - b) Safety medicine; prescriber may determine dispensing frequency
- |  |        |    |   |          |
|--|--------|----|---|----------|
| Tab sublingual 2 mg with naloxone 0.5 mg ..... | 57.40  | 28 | ✓ | Suboxone |
| Tab sublingual 8 mg with naloxone 2 mg .....   | 166.00 | 28 | ✓ | Suboxone |

► **SA1203** **Special Authority for Subsidy**

**Initial application — (Detoxification)** from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health..

**Initial application — (Maintenance treatment)** from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

**Renewal — (Detoxification)** from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

**Renewal — (Maintenance treatment)** from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and

continued...

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\* Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

**Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification)** from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

## BUPROPION HYDROCHLORIDE

Tab modified-release 150 mg.....	11.00	30	✓ <b>Zyban</b>
----------------------------------	-------	----	----------------

## DISULFIRAM

Tab 200 mg .....	75.57	100	✓ <b>Antabuse</b>
------------------	-------	-----	-------------------

## NALTREXONE HYDROCHLORIDE – Special Authority see [SA1408 below](#) – Retail pharmacy

Tab 50 mg .....	112.55	30	✓ <b>Naltrexone</b>
-----------------	--------	----	---------------------

## ► [SA1408](#) Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

**Renewal** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
  - 2.1 Patient is still unstable and requires further treatment; or
  - 2.2 Patient achieved significant improvement but requires further treatment; or
  - 2.3 Patient is well controlled but requires maintenance therapy.



	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>NICOTINE</b>				
a) Nicotine will not be funded in amounts less than 4 weeks of treatment.				
b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.				
Patch 7 mg – Up to 28 patch available on a PSO .....	17.28	28	✓	<b>Habitrol</b>
Patch 7 mg for direct distribution only – [Xpharm] .....	3.94	7	✓	<b>Habitrol</b>
Patch 14 mg – Up to 28 patch available on a PSO .....	19.00	28	✓	<b>Habitrol</b>
Patch 14 mg for direct distribution only – [Xpharm] .....	4.52	7	✓	<b>Habitrol</b>
Patch 21 mg – Up to 28 patch available on a PSO .....	21.77	28	✓	<b>Habitrol</b>
Patch 21 mg for direct distribution only – [Xpharm] .....	5.18	7	✓	<b>Habitrol</b>
Lozenge 1 mg – Up to 216 loz available on a PSO .....	18.27	216	✓	<b>Habitrol</b>
Lozenge 1 mg for direct distribution only – [Xpharm] .....	3.20	36	✓	<b>Habitrol</b>
Lozenge 2 mg – Up to 216 loz available on a PSO .....	20.02	216	✓	<b>Habitrol</b>
Lozenge 2 mg for direct distribution only – [Xpharm] .....	3.24	36	✓	<b>Habitrol</b>
Gum 2 mg (Fruit) – Up to 384 piece available on a PSO .....	36.39	384	✓	<b>Habitrol</b>
Gum 2 mg (Fruit) for direct distribution only – [Xpharm] .....	8.64	96	✓	<b>Habitrol</b>
Gum 2 mg (Mint) – Up to 384 piece available on a PSO .....	36.39	384	✓	<b>Habitrol</b>
Gum 2 mg (Mint) for direct distribution only – [Xpharm] .....	8.64	96	✓	<b>Habitrol</b>
Gum 4 mg (Fruit) – Up to 384 piece available on a PSO .....	42.07	384	✓	<b>Habitrol</b>
Gum 4 mg (Fruit) for direct distribution only – [Xpharm] .....	10.01	96	✓	<b>Habitrol</b>
Gum 4 mg (Mint) – Up to 384 piece available on a PSO .....	42.07	384	✓	<b>Habitrol</b>
Gum 4 mg (Mint) for direct distribution only – [Xpharm] .....	10.01	96	✓	<b>Habitrol</b>

**VARENICLINE TARTRATE** – Special Authority see [SA1845 below](#) – Retail pharmacy

- A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- Varenicline will not be funded in amounts less than 4 weeks of treatment.
- The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg x 11 and 1 mg x 42 .....	25.64	53 OP	✓	<b>Varenicline Pfizer</b>
Tab 1 mg .....	27.10	56	✓	<b>Varenicline Pfizer</b>

➡ **SA1845** **Special Authority for Subsidy**

**Initial application** from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria:

All of the following:

- Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- Either:
  - The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
  - The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- The patient is not pregnant; and
- The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

**Renewal** from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria:

All of the following:

- Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking;

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- and
- 2

The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3

It has been 6 months since the patient's previous Special Authority was approved; and
- 4

Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5

The patient is not pregnant; and
- 6

The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.  
Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.  
This includes the 4-week 'starter' pack.

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## Chemotherapeutic Agents

### Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE – PCT only – Specialist – Special Authority see [SA1667](#) below

Inj 25 mg vial .....	271.35	1	✓ Ribomustin
Inj 100 mg vial .....	1,085.38	1	✓ Ribomustin
Inj 1 mg for ECP .....	11.40	1 mg	✓ Baxter

#### ►SA1667 Special Authority for Subsidy

**Initial application — (treatment naive CLL)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m<sup>2</sup> on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

**Initial application — (Indolent, Low-grade lymphomas)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
  - 3.1 Both:
    - 3.1.1 Patient is treatment naive; and
    - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
  - 3.2 All of the following:
    - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
    - 3.2.2 The patient has not received prior bendamustine therapy; and
    - 3.2.3 Either:
      - 3.2.3.1 Both:
        - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
        - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
      - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

**Renewal — (Indolent, Low-grade lymphomas)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Either:
  - 2.1 Both:

continued...

# ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
continued...				
2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and				
2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or				
2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.				
Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.				
BUSULFAN – PCT – Retail pharmacy-Specialist				
Tab 2 mg .....	89.25	100	✓	Myleran
CARBOPLATIN – PCT only – Specialist				
Inj 10 mg per ml, 45 ml vial.....	32.59	1	✓	DBL Carboplatin
	45.20		✓	Carboplatin Ebewe
	48.50		✓	Carbaccord
Inj 1 mg for ECP .....	0.10	1 mg	✓	Baxter
CARMUSTINE – PCT only – Specialist				
Inj 100 mg vial .....	1,387.00	1	✓	BiCNU
			✓	Bicnu Heritage <sup>\$29</sup>
Inj 100 mg for ECP .....	1,387.00	100 mg OP	✓	Baxter
CHLORAMBUCIL – PCT – Retail pharmacy-Specialist				
Tab 2 mg .....	29.06	25	✓	Leukeran FC
CISPLATIN – PCT only – Specialist				
Inj 1 mg per ml, 50 ml vial.....	12.29	1	✓	DBL Cisplatin
	15.00		✓	Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial.....	19.70	1	✓	DBL Cisplatin
	21.00		✓	Cisplatin Ebewe
Inj 1 mg for ECP .....	0.25	1 mg	✓	Baxter
CYCLOPHOSPHAMIDE				
Tab 50 mg – PCT – Retail pharmacy-Specialist.....	79.00	50	✓	Endoxan <sup>\$29</sup>
	158.00	100	✓	Procytox <sup>\$29</sup>
Wastage claimable				
Inj 1 g vial – PCT – Retail pharmacy-Specialist.....	35.65	1	✓	Endoxan
	127.80	6	✓	Cytoxan
Inj 2 g vial – PCT only – Specialist.....	71.25	1	✓	Endoxan
Inj 1 mg for ECP – PCT only – Specialist.....	0.04	1 mg	✓	Baxter
IFOSFAMIDE – PCT only – Specialist				
Inj 1 g.....	96.00	1	✓	Holoxan
Inj 2 g.....	180.00	1	✓	Holoxan
Inj 1 mg for ECP .....	0.10	1 mg	✓	Baxter
LOMUSTINE – PCT – Retail pharmacy-Specialist				
Cap 10 mg .....	132.59	20	✓	CeeNU
Cap 40 mg .....	399.15	20	✓	CeeNU
MELPHALAN				
Tab 2 mg – PCT – Retail pharmacy-Specialist.....	40.70	25	✓	Alkeran
Inj 50 mg – PCT only – Specialist.....	67.80	1	✓	Alkeran

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>OXALIPLATIN – PCT only – Specialist</b>				
Inj 100 mg vial .....	25.01	1	✓	<b>Oxaliplatin Actavis 100</b>
	110.00		✓	<b>Oxaliplatin Ebewe</b>
Inj 5 mg per ml, 20 ml vial.....	46.32	1	✓	<b>Oxaliccord</b>
			✓	<b>Oxaliplatin Accord</b>
Inj 1 mg for ECP .....	0.48	1 mg	✓	<b>Baxter</b>
<i>(Oxaliccord Inj 5 mg per ml, 20 ml vial to be delisted 1 February 2020)</i>				
<b>THIOTEPA – PCT only – Specialist</b>				
Inj 15 mg vial .....	CBS	1	✓	<b>Bedford</b> <small>\$29</small>
			✓	<b>THIO-TEPA</b> <small>\$29</small>
			✓	<b>Tepadina</b> <small>\$29</small>
Inj 100 mg vial .....	CBS	1	✓	<b>Tepadina</b> <small>\$29</small>

### Antimetabolites

**AZACITIDINE – PCT only – Specialist – Special Authority see SA1467 below**

Inj 100 mg vial .....	139.00	1	✓	<b>Azacitidine Dr Reddy's</b>
	605.00		✓	<b>Vidaza</b>
Inj 1 mg for ECP .....	1.53	1 mg	✓	<b>Baxter</b>

#### ►SA1467 Special Authority for Subsidy

**Initial application** only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- Any of the following:
  - The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
  - The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
  - The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- The patient has performance status (WHO/ECOG) grade 0-2; and
- The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- The patient has an estimated life expectancy of at least 3 months.

**Renewal** only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- No evidence of disease progression; and
- The treatment remains appropriate and patient is benefitting from treatment.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

# ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>CALCIUM FOLINATE</b>				
Tab 15 mg – PCT – Retail pharmacy-Specialist.....	104.26	10	✓	<b>DBL Leucovorin Calcium</b>
Inj 3 mg per ml, 1 ml – PCT – Retail pharmacy-Specialist .....	17.10	5	✓	<b>Hospira</b>
Inj 10 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist.....	7.28	1	✓	<b>Calcium Folate Sandoz</b>
Inj 50 mg – PCT – Retail pharmacy-Specialist .....	18.25	5	✓	<b>Calcium Folate Ebewe</b>
Inj 10 mg per ml, 10 ml vial – PCT only – Specialist .....	9.49	1	✓	<b>Calcium Folate Sandoz</b>
Inj 100 mg – PCT only – Specialist.....	7.33	1	✓	<b>Calcium Folate Ebewe</b>
Inj 300 mg – PCT only – Specialist.....	22.51	1	✓	<b>Calcium Folate Ebewe</b>
Inj 10 mg per ml, 35 ml vial – PCT only – Specialist .....	25.14	1	✓	<b>Calcium Folate Sandoz</b>
Inj 1 g – PCT only – Specialist.....	67.51	1	✓	<b>Calcium Folate Ebewe</b>
Inj 10 mg per ml, 100 ml vial – PCT only – Specialist .....	72.00	1	✓	<b>Calcium Folate Sandoz</b>
Inj 1 mg for ECP – PCT only – Specialist.....	0.06	1 mg	✓	<b>Baxter</b>
<i>(Calcium Folate Ebewe Inj 50 mg to be delisted 1 March 2020)</i>				
<b>CAPECITABINE – Retail pharmacy-Specialist</b>				
Tab 150 mg .....	11.15	60	✓	<b>Brinov</b>
Tab 500 mg .....	62.28	120	✓	<b>Brinov</b>
<b>CLADRIBINE – PCT only – Specialist</b>				
Inj 1 mg per ml, 10 ml .....	5,249.72	7	✓	<b>Leustatin</b>
Inj 10 mg for ECP .....	749.96	10 mg OP	✓	<b>Baxter</b>
<b>CYTARABINE</b>				
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist.....	400.00	5	✓	<b>Pfizer</b>
Inj 100 mg per ml, 20 ml vial – PCT – Retail pharmacy-Specialist .....	41.36	1	✓	<b>Pfizer</b>
Inj 1 mg for ECP – PCT only – Specialist.....	0.25	10 mg	✓	<b>Baxter</b>
Inj 100 mg intrathecal syringe for ECP – PCT only – Specialist.....	80.00	100 mg OP	✓	<b>Baxter</b>
<b>FLUDARABINE PHOSPHATE</b>				
Tab 10 mg – PCT – Retail pharmacy-Specialist.....	412.00	20	✓	<b>Fludara Oral</b>
Inj 50 mg vial – PCT only – Specialist .....	576.45	5	✓	<b>Fludarabine Ebewe</b>
Inj 50 mg for ECP – PCT only – Specialist.....	115.29	50 mg OP	✓	<b>Baxter</b>
<b>FLUOROURACIL</b>				
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist .....	12.00	1	✓	<b>Fluorouracil Ebewe</b>
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist .....	30.00	1	✓	<b>Fluorouracil Ebewe</b>
Inj 1 mg for ECP – PCT only – Specialist.....	0.66	100 mg	✓	<b>Baxter</b>
<b>GEMCITABINE HYDROCHLORIDE – PCT only – Specialist</b>				
Inj 1 g, 26.3 ml vial.....	62.50	1	✓	<b>DBL Gemcitabine</b>
Inj 1 g.....	15.89	1	✓	<b>Gemcitabine Ebewe</b>
	349.20		✓	<b>Gemzar</b>
Inj 1 mg for ECP .....	0.02	1 mg	✓	<b>Baxter</b>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>IRINOTECAN HYDROCHLORIDE – PCT only – Specialist</b>				
Inj 20 mg per ml, 5 ml vial.....	71.44	1	✓	<b>Irinotecan</b> <b>Accord S29</b>
			✓	<b>Irinotecan Actavis</b> <b>100</b>
	100.00		✓	<b>Irinotecan-Rex</b>
Inj 1 mg for ECP .....	0.75	1 mg	✓	<b>Baxter</b>
<b>MERCAPTOPURINE</b>				
Tab 50 mg – PCT – Retail pharmacy-Specialist.....	37.00	25	✓	<b>Puri-nethol</b>
Oral suspension 20 mg per ml – Retail pharmacy-Specialist – Special Authority see <a href="#">SA1725 below</a> .....	428.00	100 ml OP	✓	<b>Allmercap</b>
<b>►SA1725 Special Authority for Subsidy</b>				
<b>Initial application</b> only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.				
<b>Renewal</b> only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.				
<b>METHOTREXATE</b>				
* Tab 2.5 mg – PCT – Retail pharmacy-Specialist.....	8.05	90	✓	<b>Trexate</b>
* Tab 10 mg – PCT – Retail pharmacy-Specialist.....	31.75	90	✓	<b>Trexate</b>
* Inj 2.5 mg per ml, 2 ml – PCT – Retail pharmacy-Specialist .....	47.50	5	✓	<b>Hospira</b>
* Inj 7.5 mg prefilled syringe.....	14.61	1	✓	<b>Methotrexate</b> <b>Sandoz</b>
* Inj 10 mg prefilled syringe.....	14.66	1	✓	<b>Methotrexate</b> <b>Sandoz</b>
* Inj 15 mg prefilled syringe.....	14.77	1	✓	<b>Methotrexate</b> <b>Sandoz</b>
* Inj 20 mg prefilled syringe.....	14.88	1	✓	<b>Methotrexate</b> <b>Sandoz</b>
* Inj 25 mg prefilled syringe.....	14.99	1	✓	<b>Methotrexate</b> <b>Sandoz</b>
* Inj 30 mg prefilled syringe.....	15.09	1	✓	<b>Methotrexate</b> <b>Sandoz</b>
* Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist.....	30.00	5	✓	<b>DBL Methotrexate</b> <b>Onco-Vial</b>
* Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Specialist.....	45.00	1	✓	<b>DBL Methotrexate</b> <b>Onco-Vial</b>
* Inj 100 mg per ml, 10 ml – PCT – Retail pharmacy-Specialist .....	25.00	1	✓	<b>Methotrexate Ebewe</b>
* Inj 100 mg per ml, 50 ml vial – PCT – Retail pharmacy-Specialist .....	79.99	1	✓	<b>Methotrexate Ebewe</b>
* Inj 1 mg for ECP – PCT only – Specialist .....	0.06	1 mg	✓	<b>Baxter</b>
* Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist.....	4.73	5 mg OP	✓	<b>Baxter</b>
<b>PEMETREXED – PCT only – Specialist – Special Authority see <a href="#">SA1679 on the next page</a></b>				
Inj 100 mg vial .....	60.89	1	✓	<b>Juno Pemetrexed</b>
Inj 500 mg vial .....	217.77	1	✓	<b>Juno Pemetrexed</b>
Inj 1 mg for ECP .....	0.55	1 mg	✓	<b>Baxter</b>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## ►SA1679 Special Authority for Subsidy

**Initial application — (mesothelioma)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m<sup>2</sup> every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

**Renewal — (mesothelioma)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m<sup>2</sup> every 21 days for a maximum of 6 cycles.

**Initial application — (non-small cell lung carcinoma)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient has chemotherapy-naïve disease; and
    - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m<sup>2</sup> every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
  - 2.2 All of the following:
    - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
    - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
    - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m<sup>2</sup> every 21 days for a maximum of 6 cycles.

**Renewal — (non-small cell lung carcinoma)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m<sup>2</sup> every 21 days.

THIOGUANINE – PCT – Retail pharmacy-Specialist

Tab 40 mg .....	126.31	25	✓ Lanvis
-----------------	--------	----	----------

## Other Cytotoxic Agents

AMSACRINE – PCT only – Specialist

Inj 50 mg per ml, 1.5 ml ampoule .....	1,500.00	6	✓ Amsidine <sup>S29</sup>
Inj 75 mg .....	1,250.00	5	✓ AmsaLyo <sup>S29</sup>

ANAGRELIDE HYDROCHLORIDE – PCT – Retail pharmacy-Specialist

Cap 0.5 mg .....	CBS	100	✓ Agrylin <sup>S29</sup> ✓ Teva <sup>S29</sup>
------------------	-----	-----	---

ARSENIC TRIOXIDE – PCT only – Specialist

Inj 1 mg per ml, 10 ml vial .....	4,817.00	10	✓ Phenasen
Inj 10 mg for ECP .....	481.70	10 mg OP	✓ Baxter



	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>BLEOMYCIN SULPHATE – PCT only – Specialist</b>				
Inj 15,000 iu, vial.....	161.01	1	✓	<b>DBL Bleomycin Sulfate</b>
Inj 1,000 iu for ECP .....	12.45	1,000 iu	✓	<b>Baxter</b>
<b>BORTEZOMIB – PCT only – Specialist – Special Authority see SA1576 below</b>				
Inj 3.5 mg vial .....	1,892.50	1	✓	<b>Velcade</b>
Inj 1 mg for ECP .....	594.77	1 mg	✓	<b>Baxter</b>
<b>►SA1576 Special Authority for Subsidy</b>				
<b>Initial application — (Treatment naïve multiple myeloma/amyloidosis)</b> only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:				
Both:				
1 Either:				
1.1 The patient has treatment-naïve symptomatic multiple myeloma; or				
1.2 The patient has treatment-naïve symptomatic systemic AL amyloidosis *; and				
2 Maximum of 9 treatment cycles.				
Note: Indications marked with * are unapproved indications.				
<b>Initial application — (Relapsed/refractory multiple myeloma/amyloidosis)</b> only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:				
All of the following:				
1 Either:				
1.1 The patient has relapsed or refractory multiple myeloma; or				
1.2 The patient has relapsed or refractory systemic AL amyloidosis *; and				
2 The patient has received only one prior front line chemotherapy for multiple myeloma or amyloidosis; and				
3 The patient has not had prior publicly funded treatment with bortezomib; and				
4 Maximum of 4 treatment cycles.				
Note: Indications marked with * are unapproved indications.				
<b>Renewal — (Relapsed/refractory multiple myeloma/amyloidosis)</b> only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:				
Both:				
1 The patient's disease obtained at least a partial response from treatment with bortezomib at the completion of cycle 4; and				
2 Maximum of 4 further treatment cycles (making a total maximum of 8 consecutive treatment cycles).				
Notes: Responding relapsed/refractory multiple myeloma patients should receive no more than 2 additional cycles of treatment beyond the cycle at which a confirmed complete response was first achieved. A line of therapy is considered to comprise either:				
a) a known therapeutic chemotherapy regimen and supportive treatments; or				
b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments.				
Refer to datasheet for recommended dosage and number of doses of bortezomib per treatment cycle.				
<b>COLASPASE [L-ASPARAGINASE] – PCT only – Specialist</b>				
Inj 10,000 iu .....	102.32	1	✓	<b>Leunase</b>
Inj 10,000 iu for ECP .....	102.32	10,000 iu OP	✓	<b>Baxter</b>
<b>DACARBAZINE – PCT only – Specialist</b>				
Inj 200 mg vial .....	58.06	1	✓	<b>DBL Dacarbazine</b>
	580.60	10	✓	<b>Dacarbazine</b>
				<b>APP S29</b>
Inj 200 mg for ECP .....	58.06	200 mg OP	✓	<b>Baxter</b>
<b>DACTINOMYCIN [ACTINOMYCIN D] – PCT only – Specialist</b>				
Inj 0.5 mg vial .....	166.75	1	✓	<b>Cosmegen</b>
Inj 0.5 mg for ECP .....	166.75	0.5 mg OP	✓	<b>Baxter</b>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

# ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>DAUNORUBICIN – PCT only – Specialist</b>				
Inj 2 mg per ml, 10 ml .....	130.00	1	✓	<b>Pfizer</b>
Inj 20 mg for ECP .....	130.00	20 mg OP	✓	<b>Baxter</b>
<b>DOCETAXEL – PCT only – Specialist</b>				
Inj 10 mg per ml, 2 ml vial .....	12.40	1	✓	<b>DBL Docetaxel</b>
Inj 20 mg .....	48.75	1	✓	<b>Docetaxel Sandoz</b>
Inj 10 mg per ml, 8 ml vial .....	26.95	1	✓	<b>DBL Docetaxel</b>
Inj 20 mg per ml, 4 ml vial .....	26.95	1	✓	<b>Docetaxel</b>
				<b>Accord <sup>S29</sup></b>
Inj 80 mg .....	195.00	1	✓	<b>Docetaxel Sandoz</b>
Inj 1 mg for ECP .....	0.55	1 mg	✓	<b>Baxter</b>
<b>DOXORUBICIN HYDROCHLORIDE – PCT only – Specialist</b>				
Inj 2 mg per ml, 5 ml vial .....	10.00	1	✓	<b>Doxorubicin Ebewe</b>
Inj 2 mg per ml, 25 ml vial .....	11.50	1	✓	<b>Doxorubicin Ebewe</b>
	17.00		✓	<b>Arrow-Doxorubicin</b>
Inj 2 mg per ml, 50 ml vial .....	23.00	1	✓	<b>Doxorubicin Ebewe</b>
Inj 2 mg per ml, 100 ml vial .....	56.15	1	✓	<b>Doxorubicin Ebewe</b>
	65.00		✓	<b>Arrow-Doxorubicin</b>
Inj 1 mg for ECP .....	0.29	1 mg	✓	<b>Baxter</b>
<b>EPIRUBICIN HYDROCHLORIDE – PCT only – Specialist</b>				
Inj 2 mg per ml, 5 ml vial .....	25.00	1	✓	<b>Epirubicin Ebewe</b>
Inj 2 mg per ml, 25 ml vial .....	30.00	1	✓	<b>Epirubicin Ebewe</b>
Inj 2 mg per ml, 100 ml vial .....	85.00	1	✓	<b>Epirubicin Ebewe</b>
Inj 1 mg for ECP .....	0.37	1 mg	✓	<b>Baxter</b>
<b>ETOPOSIDE</b>				
Cap 50 mg – PCT – Retail pharmacy-Specialist .....	340.73	20	✓	<b>Vepesid</b>
Cap 100 mg – PCT – Retail pharmacy-Specialist .....	340.73	10	✓	<b>Vepesid</b>
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist .....	7.90	1	✓	<b>Rex Medical</b>
Inj 1 mg for ECP – PCT only – Specialist .....	0.09	1 mg	✓	<b>Baxter</b>
<b>ETOPOSIDE PHOSPHATE – PCT only – Specialist</b>				
Inj 100 mg (of etoposide base) .....	40.00	1	✓	<b>Etopophos</b>
Inj 1 mg (of etoposide base) for ECP .....	0.47	1 mg	✓	<b>Baxter</b>
<b>HYDROXYUREA – PCT – Retail pharmacy-Specialist</b>				
Cap 500 mg .....	31.76	100	✓	<b>Hydrea</b>
<b>IDARUBICIN HYDROCHLORIDE</b>				
Inj 5 mg vial – PCT only – Specialist .....	93.00	1	✓	<b>Zavedos</b>
Inj 10 mg vial – PCT only – Specialist .....	198.00	1	✓	<b>Zavedos</b>
Inj 1 mg for ECP – PCT only – Specialist .....	21.84	1 mg	✓	<b>Baxter</b>
<b>LENALIDOMIDE – Retail pharmacy-Specialist – Special Authority see SA1468 below</b>				
Wastage claimable				
Cap 10 mg .....	6,207.00	21	✓	<b>Revlimid</b>
Cap 15 mg .....	7,239.18	21	✓	<b>Revlimid</b>
Cap 25 mg .....	7,627.00	21	✓	<b>Revlimid</b>

## ►SA1468 Special Authority for Subsidy

**Initial application — (Relapsed/refractory disease)** only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
continued...			
1 Patient has relapsed or refractory multiple myeloma with progressive disease; and			
2 Either:			
2.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or			
2.2 Both:			
2.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and			
2.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and			
3 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.			
<b>Renewal</b> only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:			
Both:			
1 No evidence of disease progression; and			
2 The treatment remains appropriate and patient is benefitting from treatment.			
Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.			
<b>MESNA</b>			
Tab 400 mg – PCT – Retail pharmacy-Specialist.....	314.00	50	✓ Uromitexan
Uromitexan to be Sole Supply on 1 November 2019			
Tab 600 mg – PCT – Retail pharmacy-Specialist.....	448.50	50	✓ Uromitexan
Uromitexan to be Sole Supply on 1 November 2019			
Inj 100 mg per ml, 4 ml ampoule – PCT only – Specialist.....	177.45	15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule – PCT only – Specialist.....	407.40	15	✓ Uromitexan
Inj 1 mg for ECP – PCT only – Specialist.....	2.96	100 mg	✓ Baxter
<b>MITOMYCIN C – PCT only – Specialist</b>			
Inj 5 mg vial .....	204.08	1	✓ Arrow
Inj 1 mg for ECP .....	42.04	1 mg	✓ Baxter
<b>MITOZANTRONE – PCT only – Specialist</b>			
Inj 2 mg per ml, 10 ml vial.....	97.50	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP .....	5.51	1 mg	✓ Baxter
<b>PACLITAXEL – PCT only – Specialist</b>			
Inj 30 mg.....	47.30	5	✓ Paclitaxel Ebewe
Inj 100 mg.....	20.00	1	✓ Paclitaxel Ebewe
	91.67		✓ Paclitaxel Actavis
Inj 150 mg.....	26.69	1	✓ Paclitaxel Ebewe
	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg.....	35.35	1	✓ Paclitaxel Ebewe
	275.00		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 1 mg for ECP .....	0.19	1 mg	✓ Baxter
<b>PEGASPARGASE – PCT only – Special Authority see SA1325 on the next page</b>			
Inj 3,750 IU per 5 ml .....	3,005.00	1	✓ Oncaspar S29

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

# ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

## ►SA1325 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and
- 3 Treatment is with curative intent.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and
- 3 Treatment is with curative intent.

PENTOSTATIN [DEOXYCOFORMYCIN] – PCT only – Specialist

Inj 10 mg.....	CBS	1	✓ Nipent <sup>\$29</sup>
----------------	-----	---	--------------------------

PROCARBAZINE HYDROCHLORIDE – PCT – Retail pharmacy-Specialist

Cap 50 mg.....	980.00	50	✓ Natulan <sup>\$29</sup>
----------------	--------	----	---------------------------

TEMOZOLOMIDE – Special Authority see SA1741 below – Retail pharmacy

Cap 5 mg.....	10.20	5	✓ Orion Temozolomide
Cap 20 mg.....	16.38	5	✓ Temaccord
	18.30		✓ Apo-Temozolomide
			✓ Orion Temozolomide
Cap 100 mg.....	35.98	5	✓ Temizole 20 <sup>\$29</sup>
	40.20		✓ Temaccord
			✓ Apo-Temozolomide
			✓ Orion Temozolomide
Cap 140 mg.....	50.12	5	✓ Temaccord
	56.00		✓ Orion Temozolomide
Cap 250 mg.....	86.34	5	✓ Temaccord
	96.80		✓ Orion Temozolomide

## ►SA1741 Special Authority for Subsidy

**Initial application — (high grade gliomas)** only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
  - 1.2 Patient has newly diagnosed anaplastic astrocytoma\*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m<sup>2</sup> per day.

**Initial application — (neuroendocrine tumours)** only from a relevant specialist. Approvals valid for 9 months for applications

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour\*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m<sup>2</sup> per day; and
- 4 Temozolomide to be discontinued at disease progression.

**Initial application — (ewing's sarcoma)** only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

**Renewal — (high grade gliomas)** only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has glioblastoma multiforme; and
  - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
  - 2.1 Patient has anaplastic astrocytoma\*; and
  - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
  - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

**Renewal — (neuroendocrine tumours)** only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

**Renewal — (ewing's sarcoma)** only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a \* is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE – Retail pharmacy-Specialist – Special Authority see [SA1124 below](#)

Cap 50 mg .....	378.00	28	✓ <b>Thalomid</b>
Cap 100 mg .....	756.00	28	✓ <b>Thalomid</b>

## ►SA1124 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis\*.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with \* is an unapproved indication.

TRETINOIN

Cap 10 mg – PCT – Retail pharmacy-Specialist .....	479.50	100	✓ <b>Vesanoid</b>
--	--------	-----	-------------------

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>VINBLASTINE SULPHATE</b>				
Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Specialist.....	186.46	5	✓	<b>Hospira</b>
Inj 1 mg for ECP – PCT only – Specialist.....	4.14	1 mg	✓	<b>Baxter</b>
<b>VINCISTINE SULPHATE</b>				
Inj 1 mg per ml, 1 ml vial – PCT – Retail pharmacy-Specialist.....	74.52	5	✓	<b>DBL Vincristine Sulfate</b>
Inj 1 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist.....	85.61	5	✓	<b>DBL Vincristine Sulfate</b>
Inj 1 mg for ECP – PCT only – Specialist.....	11.30	1 mg	✓	<b>Baxter</b>
<b>VINOELBINE – PCT only – Specialist</b>				
Inj 10 mg per ml, 1 ml vial.....	12.00	1	✓	<b>Navelbine</b>
	42.00		✓	<b>Vinorelbine Ebewe</b>
Inj 10 mg per ml, 5 ml vial.....	56.00	1	✓	<b>Navelbine</b>
	210.00		✓	<b>Vinorelbine Ebewe</b>
Inj 1 mg for ECP.....	1.25	1 mg	✓	<b>Baxter</b>

## Protein-tyrosine Kinase Inhibitors

**DASATINIB** – Special Authority see [SA1805 below](#) – Retail pharmacy

Wastage claimable

Tab 20 mg .....	3,774.06	60	✓	<b>Sprycel</b>
Tab 50 mg .....	6,214.20	60	✓	<b>Sprycel</b>
Tab 70 mg .....	7,692.58	60	✓	<b>Sprycel</b>

### ►SA1805 Special Authority for Subsidy

**Initial application** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

1 Both:

- 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
- 1.2 Maximum dose of 140 mg/day; or

2 Both:

- 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
- 2.2 Maximum dose of 140 mg/day; or

3 All of the following:

- 3.1 The patient has a diagnosis of CML in chronic phase; and
- 3.2 Maximum dose of 100 mg/day; and
- 3.3 Any of the following:
  - 3.3.1 Patient has documented treatment failure\* with imatinib; or
  - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
  - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
  - 3.3.4 Patients is enrolled in the KISS study\*\* and requires dasatinib treatment according to the study protocol.

**Renewal** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib\*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: \*treatment failure for CML as defined by Leukaemia Net Guidelines. \*\*Kinase-Inhibition Study with Sprycel Start-up <https://www.cancertrialsnz.ac.nz/kiss/>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>ERLOTINIB – Retail pharmacy-Specialist – Special Authority see <a href="#">SA1653 below</a></b>				
Tab 100 mg .....	764.00	30	✓	<b>Tarceva</b>
Tab 150 mg .....	1,146.00	30	✓	<b>Tarceva</b>

► **SA1653 Special Authority for Subsidy**

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Either:
  - 3.1 Patient is treatment naive; or
  - 3.2 Both:
    - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
    - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

**GEFITINIB – Retail pharmacy-Specialist – Special Authority see [SA1654 below](#)**

Tab 250 mg .....	1,700.00	30	✓	<b>Iressa</b>
------------------	----------	----	---	---------------

► **SA1654 Special Authority for Subsidy**

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
  - 2.1 Patient is treatment naive; or
  - 2.2 Both:
    - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
    - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

**IMATINIB MESILATE**

Note: Imatinib-AFT is not a registered for the treatment of Gastro Intestinal Stromal Tumours (GIST). The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule.

Tab 100 mg – [Xpharm] – Special Authority see [SA1460 below](#)

below .....	2,400.00	60	✓	<b>Glivec</b>
* Cap 100 mg .....	98.00	60	✓	<b>Imatinib-AFT</b>
* Cap 400 mg .....	197.50	30	✓	<b>Imatinib-AFT</b>

► **SA1460 Special Authority for Subsidy**

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from PHARMAC's website <http://www.pharmac.govt.nz>, and prescriptions should be sent to:

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

The CML/GIST Co-ordinator Phone: (04) 460 4990  
 PHARMAC Facsimile: (04) 916 7571  
 PO Box 10 254 Email: [cmlgistcoordinator@pharmac.govt.nz](mailto:cmlgistcoordinator@pharmac.govt.nz)  
 Wellington

## Special Authority criteria for GIST – access by application

Funded for patients:

- With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- Maximum dose of 400 mg/day.
- Applications to be made and subsequent prescriptions can be written by an oncologist.
- Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE – Special Authority see [SA1191 below](#) – Retail pharmacy

Tab 250 mg .....	1,899.00	70	✓ Tykerb
------------------	----------	----	----------

## ►SA1191 Special Authority for Subsidy

**Initial application — (metastatic breast cancer)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- All of the following:
  - The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - The patient has not previously received trastuzumab treatment for HER 2 positive metastatic breast cancer; and
  - Lapatinib not to be given in combination with trastuzumab; and
  - Lapatinib to be discontinued at disease progression; or
- All of the following:
  - The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - The patient started trastuzumab for metastatic breast cancer but discontinued trastuzumab within 3 months of starting treatment due to intolerance; and
  - The cancer did not progress whilst on trastuzumab; and
  - Lapatinib not to be given in combination with trastuzumab; and
  - Lapatinib to be discontinued at disease progression.

**Renewal — (metastatic breast cancer)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- Lapatinib not to be given in combination with trastuzumab; and
- Lapatinib to be discontinued at disease progression.

NILOTINIB – Special Authority see [SA1489 on the next page](#) – Retail pharmacy

Wastage claimable

Cap 150 mg.....	4,680.00	120	✓ Tasigna
Cap 200 mg.....	6,532.00	120	✓ Tasigna



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## ►SA1489 Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Either:
  - 2.1 Patient has documented CML treatment failure\* with imatinib; or
  - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: \*treatment failure as defined by Leukaemia Net Guidelines.

**Renewal** only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PAZOPANIB – Special Authority see SA1190 below – Retail pharmacy

Tab 200 mg .....	1,334.70	30	✓ Votrient
Tab 400 mg .....	2,669.40	30	✓ Votrient

## ►SA1190 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 Both:
    - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
    - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and  
The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal; or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of less than or equal to 70; or
  - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>RUXOLITINIB</b> – Special Authority see <a href="#">SA1753 below</a> – Retail pharmacy				
Wastage claimable				
Tab 5 mg .....	2,500.00	56	✓ <b>Jakavi</b>	
Tab 15 mg .....	5,000.00	56	✓ <b>Jakavi</b>	
Tab 20 mg .....	5,000.00	56	✓ <b>Jakavi</b>	

## ► **SA1753** Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
- 3 A maximum dose of 20 mg twice daily is to be given.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

**SUNITINIB** – Special Authority see [SA1266 below](#) – Retail pharmacy

Cap 12.5 mg .....	2,315.38	28	✓ <b>Sutent</b>
Cap 25 mg .....	4,630.77	28	✓ <b>Sutent</b>
Cap 50 mg .....	9,261.54	28	✓ <b>Sutent</b>

## ► **SA1266** Special Authority for Subsidy

**Initial application** — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
  - 2.4 Both:
    - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
    - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and  
The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal; or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of less than or equal to 70; or
  - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

**Initial application** — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

continued...

specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
  - 2.1 The patient's disease has progressed following treatment with imatinib; or
  - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

**Renewal — (RCC)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

**Renewal — (GIST)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
  - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
  - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
  - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

## Endocrine Therapy

For GnRH ANALOGUES – refer to HORMONE PREPARATIONS, Tropic Hormones, [page 83](#)

ABIRATERONE ACETATE – Retail pharmacy-Specialist – Special Authority see [SA1767 below](#)

Wastage claimable

Tab 250 mg .....	4,276.19	120	✓ Zytiga
------------------	----------	-----	----------

► **SA1767** Special Authority for Subsidy

**Initial application** only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:

continued...

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\* Three months or six months, as applicable, dispensed all-at-once

# ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 4.1 All of the following:
  - 4.1.1 Patient is symptomatic; and
  - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
  - 4.1.3 Patient has ECOG performance score of 0-1; and
  - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
- 4.2 All of the following:
  - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
  - 4.2.2 Patient has ECOG performance score of 0-2; and
  - 4.2.3 Patient has not had prior treatment with abiraterone.

**Renewal — (abiraterone acetate)** only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

## BICALUTAMIDE

Tab 50 mg .....	3.80	28	✓ <b>Binarex</b>
-----------------	------	----	------------------

## FLUTAMIDE – Retail pharmacy-Specialist

Tab 250 mg .....	100.38	84	✓ <b>Flutamide</b>
			<b>Mylan</b> <small>\$29</small>
	119.50	100	✓ <b>Flutamin</b>

## MEGESTROL ACETATE – Retail pharmacy-Specialist

Tab 160 mg .....	63.53	30	✓ <b>Apo-Megestrol</b>
------------------	-------	----	------------------------

## OCTREOTIDE

Inj 50 mcg per ml, 1 ml vial.....	30.64	5	✓ <b>DBL Octreotide</b>
Inj 100 mcg per ml, 1 ml vial.....	18.69	5	✓ <b>DBL Octreotide</b>
Inj 500 mcg per ml, 1 ml vial.....	72.50	5	✓ <b>DBL Octreotide</b>

## OCTREOTIDE LAR (SOMATOSTATIN ANALOGUE) – Special Authority see [SA1016 below](#) – Retail pharmacy

Inj LAR 10 mg prefilled syringe.....	1,772.50	1	✓ <b>Sandostatin LAR</b>
Inj LAR 20 mg prefilled syringe.....	2,358.75	1	✓ <b>Sandostatin LAR</b>
Inj LAR 30 mg prefilled syringe.....	2,951.25	1	✓ <b>Sandostatin LAR</b>

### ► [SA1016](#) Special Authority for Subsidy

**Initial application — (Malignant Bowel Obstruction)** from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea\* and vomiting\* due to malignant bowel obstruction\*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Renewal — (Malignant Bowel Obstruction)** from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

**Initial application — (Acromegaly)** only from a relevant specialist or medical practitioner on the recommendation of a relevant

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
  - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
  - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
  - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

**Renewal — (Acromegaly)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

**Initial application — (Other Indications)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas - for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
  - 2.1 Gastrinoma; and
  - 2.2 Either:
    - 2.2.1 Patient has failed surgery; or
    - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
  - 3.1 Insulinomas; and
  - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
  - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
  - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

**Renewal — (Other Indications)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

## TAMOXIFEN CITRATE

* Tab 10 mg .....	11.75	60	✓ <b>Tamoxifen Sandoz</b>
* Tab 20 mg .....	5.60	60	✓ <b>Tamoxifen Sandoz</b>

## Aromatase Inhibitors

### ANASTROZOLE

* Tab 1 mg .....	5.04	30	✓ <b>Rolin</b>
------------------	------	----	----------------

### EXEMESTANE

* Tab 25 mg .....	14.50	30	✓ <b>Pfizer Exemestane</b>
-------------------	-------	----	----------------------------

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
LETROZOLE				
* Tab 2.5 mg .....	4.68	30	✓	<b>Letrole</b>

## Immunosuppressants

### Cytotoxic Immunosuppressants

AZATHIOPRINE – Retail pharmacy-Specialist

* Tab 25 mg .....	7.35	60	✓	<b>Azamun</b>
	9.66	100	✓	<b>Imuran</b>
Azamun to be Sole Supply on 1 January 2020				
* Tab 50 mg .....	7.60	100	✓	<b>Azamun</b>
	10.58		✓	<b>Imuran</b>
Azamun to be Sole Supply on 1 January 2020				
* Inj 50 mg vial .....	199.00	1	✓	<b>Imuran</b>
<i>(Imuran Tab 25 mg to be delisted 1 January 2020)</i>				
<i>(Imuran Tab 50 mg to be delisted 1 January 2020)</i>				

MYCOPHENOLATE MOFETIL

Tab 500 mg .....	25.00	50	✓	<b>Cellcept</b>
Cap 250 mg .....	25.00	100	✓	<b>Cellcept</b>
Powder for oral liq 1 g per 5 ml – Subsidy by endorsement .....	187.25	165 ml OP	✓	<b>Cellcept</b>

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

## Fusion Proteins

ETANERCEPT – Special Authority see [SA1812 below](#) – Retail pharmacy

Inj 25 mg .....	799.96	4	✓	<b>Enbrel</b>
Inj 50 mg autoinjector .....	1,599.96	4	✓	<b>Enbrel</b>
Inj 50 mg prefilled syringe .....	1,599.96	4	✓	<b>Enbrel</b>

### ►SA1812 Special Authority for Subsidy

**Initial application — (juvenile idiopathic arthritis)** only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
  - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for JIA; or

2 All of the following:

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient diagnosed with Juvenile Idiopathic Arthritis (JIA); and
- 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m<sup>2</sup> weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
- 2.5 Both:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

2.5.1 Either:

2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or

2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and

2.5.2 Physician's global assessment indicating severe disease.

**Initial application — (rheumatoid arthritis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and

1.2 Either:

1.2.1 The patient has experienced intolerable side effects from adalimumab; or

1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis; or

2 All of the following:

2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and

2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and

2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and

2.5 Any of the following:

2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or

2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or

2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and

2.6 Either:

2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or

2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

2.7 Either:

2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or

2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Initial application — (severe chronic plaque psoriasis)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and

1.2 Either:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or

2 All of the following:

2.1 Either:

2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or

2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and

2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and

2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and

2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

**Initial application — (ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and

1.2 Either:

1.2.1 The patient has experienced intolerable side effects from adalimumab; or

1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or

2 All of the following:

2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and

2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and

2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and

2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and

2.5 Either:

2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or

2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and

2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID

continued...



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm

25-34 years - Male: 7.5 cm; Female: 5.5 cm

35-44 years - Male: 6.5 cm; Female: 4.5 cm

45-54 years - Male: 6.0 cm; Female: 5.0 cm

55-64 years - Male: 5.5 cm; Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

**Initial application — (psoriatic arthritis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for adalimumab for psoriatic arthritis; and

1.2 Either:

1.2.1 The patient has experienced intolerable side effects from adalimumab; or

1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for psoriatic arthritis; or

2 All of the following:

2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and

2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and

2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and

2.4 Either:

2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or

2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

2.5 Any of the following:

2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or

2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or

2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Patient has pyoderma gangrenosum\*; and

2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and

3 A maximum of 4 doses.

Note: Indications marked with \* are unapproved indications.

**Initial application — (adult-onset Still's disease)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

- 1.1 Either:
  - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
  - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
  - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

**Renewal — (juvenile idiopathic arthritis)** only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a named specialist or rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
  - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

**Renewal — (rheumatoid arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
  - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Renewal — (severe chronic plaque psoriasis)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 1.1 Applicant is a dermatologist; or
- 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 2.1.2 Either:
      - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
      - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
  - 2.2 Both:
    - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
    - 2.2.2 Either:
      - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
      - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Renewal — (psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist.

Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 4 doses.

**Renewal — (adult-onset Still's disease)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

## Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) – PCT only – Specialist

Inj 50 mg per ml, 5 ml.....	2,351.25	5	✓ ATGAM
-----------------------------	----------	---	---------

BACILLUS CALMETTE-GUERIN (BCG) VACCINE – PCT only – Specialist

Subsidised only for bladder cancer.

Inj 2-8 x 100 million CFU.....	149.37	1	✓ OncoTICE
--------------------------------	--------	---	------------

Inj 40 mg per ml, vial .....	162.70	3	✓ SII-Onco-BCG <sup>\$29</sup>
------------------------------	--------	---	--------------------------------

(SII-Onco-BCG <sup>\$29</sup> Inj 40 mg per ml, vial to be delisted 1 January 2020)

## Monoclonal Antibodies

ADALIMUMAB – Special Authority see [SA1847 below](#) – Retail pharmacy

Inj 20 mg per 0.4 ml prefilled syringe.....	1,599.96	2	✓ Humira
Inj 40 mg per 0.8 ml prefilled pen .....	1,599.96	2	✓ HumiraPen
Inj 40 mg per 0.8 ml prefilled syringe.....	1,599.96	2	✓ Humira

► [SA1847](#) Special Authority for Subsidy

**Initial application — (Crohn's disease - adults)** only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

**Renewal — (Crohn's disease - adults)** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

All of the following:

- 1 Either:
  - 1.1 Applicant is a gastroenterologist; or
  - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
  - 2.1 Either:
    - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
    - 2.1.2 CDAI score is 150 or less; or
  - 2.2 Both:
    - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
    - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (Crohn's disease - children)** only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

**Renewal — (Crohn's disease - children)** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a gastroenterologist; or
  - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
  - 2.1 Either:
    - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
    - 2.1.2 PCDAI score is 15 or less; or
  - 2.2 Both:
    - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
    - 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (adult-onset Still's disease)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Either:

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
- 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
  - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

**Renewal — (adult-onset Still's disease)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

**Initial application — (ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
  - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes); and
  - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm

25-34 years - Male: 7.5 cm; Female: 5.5 cm

35-44 years - Male: 6.5 cm; Female: 4.5 cm

45-54 years - Male: 6.0 cm; Female: 5.0 cm

55-64 years - Male: 5.5 cm; Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
    - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for chronic ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
    - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
    - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

**Renewal — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or

- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and

- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

**Initial application — (fistulising Crohn's disease)** only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Either:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's disease.

**Renewal — (fistulising Crohn's disease)** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Applicant is a gastroenterologist; or
  - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
  - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

**Initial application — (juvenile idiopathic arthritis)** only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for juvenile idiopathic arthritis; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient diagnosed with JIA; and
  - 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
  - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m<sup>2</sup> weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
  - 2.5 Both:
    - 2.5.1 Either:

continued...



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
- 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
- 2.5.2 Physician's global assessment indicating severe disease.

**Renewal — (juvenile idiopathic arthritis)** only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a named specialist or rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
  - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

**Initial application — (psoriatic arthritis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

All of the following:

1 Either:

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and

2 Either:

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and

3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 4 doses.

Note: Note: Indications marked with \* are unapproved indications.

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist.

Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 4 doses.

**Initial application — (rheumatoid arthritis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
  - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis; or

2 All of the following:

- 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 2.5 Any of the following:
  - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
  - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
  - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (rheumatoid arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
  - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Either:
  - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
  - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

**Initial application — (severe Behcet's disease)** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease that is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
  - 2.1 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has not responded adequately to treatment with infliximab (see Notes); or
  - 2.2 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has experienced intolerable side effects from treatment with infliximab; and
- 3 The patient is experiencing significant loss of quality of life; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: Behcet's disease diagnosed according to the International Study Group for Behcet's disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al, J Rheumatol. 2004;31:931-7.

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

**Renewal — (severe Behcet's disease)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (severe chronic plaque psoriasis)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
  - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
  - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

**Renewal — (severe chronic plaque psoriasis)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a dermatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 2.1.2 Either:
      - 2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or

2.2 Both:

2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and

2.2.2 Either:

2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or

2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and

3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

**Initial application — (severe ocular inflammation)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; and

1.2 Either:

1.2.1 The patient has experienced intolerable side effects from infliximab; or

1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for severe ocular inflammation; or

2 Both:

2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and

2.2 Any of the following:

2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or

2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or

2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

**Renewal — (severe ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Any of the following:

1.1 The patient has had a good clinical response following 3 initial doses; or

1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or

1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and

2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

**Initial application — (hidradenitis suppurativa)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 The patient has 3 or more active lesions (e.g. inflammatory nodules, abscesses, draining fistulae); and
- 4 The patient has a Dermatology Quality of Life Index of 10 or more and the assessment is no more than 1 month old at time of application; and
- 5 Following the initial loading doses, adalimumab is to be administered at doses no greater than 40mg every 7 days.

**Renewal — (hidradenitis suppurativa)** only from a dermatologist or Practitioner on the recommendation of a dermatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

**AFLIBERCEPT** – Special Authority see [SA1772 below](#) – Retail pharmacy

Inj 40 mg per ml, 0.1 ml vial.....	1,250.00	1	✓ Eylea
------------------------------------	----------	---	---------

► [SA1772](#) **Special Authority for Subsidy**

**Initial application — (wet age related macular degeneration)** only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Wet age-related macular degeneration (wet AMD); or
    - 1.1.2 Polypoidal choroidal vasculopathy; or
    - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
  - 1.2 Either:
    - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
    - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
- 1.3 There is no structural damage to the central fovea of the treated eye; and
- 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
  - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
  - 2.2 Patient has previously\* (\*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

**Initial application — (diabetic macular oedema)** only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 – 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

**Renewal — (wet age related macular degeneration)** only from an ophthalmologist. Approvals valid for 12 months for

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

continued...

applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

**Renewal — (diabetic macular oedema)** only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

CETUXIMAB – PCT only – Specialist – Special Authority see [SA1697 below](#)

Inj 5 mg per ml, 20 ml vial.....	364.00	1	✓ <b>Erbix</b>
Inj 5 mg per ml, 100 ml vial.....	1,820.00	1	✓ <b>Erbix</b>
Inj 1 mg for ECP .....	3.82	1 mg	✓ <b>Baxter</b>

► **SA1697** **Special Authority for Subsidy**

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB – PCT only – Special Authority see [SA1831 below](#)

Inj 100 mg.....	806.00	1	✓ <b>Remicade</b>
Inj 1 mg for ECP .....	8.29	1 mg	✓ <b>Baxter</b>

► **SA1831** **Special Authority for Subsidy**

**Initial application — (Crohn's disease (adults))** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

**Renewal — (Crohn's disease (adults))** only from a gastroenterologist or Practitioner on the recommendation of a

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
  - 1.2 CDAI score is 150 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application — (Crohn's disease (children))** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

**Renewal — (Crohn's disease (children))** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application — (Graft vs host disease)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

**Initial application — (Pulmonary sarcoidosis)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

**Initial application — (acute severe fulminant ulcerative colitis)** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

**Initial application — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:

continued...



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
- 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

**Initial application — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
    - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
    - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

**Renewal — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

**Initial application — (fistulising Crohn's disease)** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Either:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

2.2 Patient has one or more rectovaginal fistula(e).

**Renewal — (fistulising Crohn's disease)** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and

2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application — (neurosarcoidosis)** only from a neurologist or Practitioner on the recommendation of a neurologist.

Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoidosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
  - 4.1 IV cyclophosphamide has been tried; or
  - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

**Renewal — (neurosarcoidosis)** only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
  - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
  - 2.2 There has been a marked reduction in prednisone dose; and
  - 2.3 Either:
    - 2.3.1 There has been an improvement in MRI appearances; or
    - 2.3.2 Marked improvement in other symptomatology.

**Initial application — (plaque psoriasis)** only from a dermatologist or Practitioner on the recommendation of a dermatologist.

Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

**Renewal — (plaque psoriasis)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

1.1 Both:

- 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
- 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or

1.2 Both:

- 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
- 1.2.2 Either:
  - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
  - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value; and

2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

**Initial application — (previous use)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Patient was being treated with infliximab prior to 1 February 2019; and

2 Any of the following:

- 2.1 Rheumatoid arthritis; or
- 2.2 Ankylosing spondylitis; or
- 2.3 Psoriatic arthritis; or
- 2.4 Severe ocular inflammation; or
- 2.5 Chronic ocular inflammation; or
- 2.6 Crohn's disease (adults); or
- 2.7 Crohn's disease (children); or
- 2.8 Fistulising Crohn's disease; or
- 2.9 Severe fulminant ulcerative colitis; or

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 2.10 Severe ulcerative colitis; or
- 2.11 Plaque psoriasis; or
- 2.12 Neurosarcoidosis; or
- 2.13 Severe Behcet's disease.

**Initial application — (psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for psoriatic arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for psoriatic arthritis.

**Renewal — (psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

**Initial application — (rheumatoid arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

**Renewal — (rheumatoid arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

**Initial application — (severe Behcet's disease)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
- 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

**Renewal — (severe Behcet's disease)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

**Renewal — (severe fulminant ulcerative colitis)** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application — (severe ocular inflammation)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  - 2.2 Any of the following:
    - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
    - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
    - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

**Renewal — (severe ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

continued...

- Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

**Initial application — (severe ulcerative colitis)** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- Patient has histologically confirmed ulcerative colitis; and
- Either:
  - Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
  - Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65; and
- Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
- Surgery (or further surgery) is considered to be clinically inappropriate.

**Renewal — (severe ulcerative colitis)** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- Either:
  - Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
  - Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

OBINUTUZUMAB – PCT only – Specialist – Special Authority see [SA1627 below](#)

Inj 25 mg per ml, 40 ml vial.....	5,910.00	1	✓ Gazyva
Inj 1 mg for ECP .....	6.21	1 mg	✓ Baxter

➔ [SA1627](#) Special Authority for Subsidy

**Initial application — (chronic lymphocytic leukaemia)** only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- The patient is obinutuzumab treatment naive; and
- The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- Patient has adequate neutrophil and platelet counts\* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- Patient has good performance status; and

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

continued...

- Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

\* Neutrophil greater than or equal to  $1.5 \times 10^9/L$  and platelets greater than or equal to  $75 \times 10^9/L$ .

OMALIZUMAB – Special Authority see [SA1744 below](#) – Retail pharmacy

Inj 150 mg prefilled syringe.....	450.00	1	✓ <b>Xolair</b>
Inj 150 mg vial .....	450.00	1	✓ <b>Xolair</b>

► **SA1744 Special Authority for Subsidy**

**Initial application — (severe asthma)** only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
  - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
  - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

**Initial application — (severe chronic spontaneous urticaria)** only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
    - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
  - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
  - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
  - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
  - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:

continued...

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\* Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 4.1 Treatment to be stopped if inadequate response\* following 4 doses; or
- 4.2 Complete response\* to 6 doses of omalizumab.

**Renewal — (severe asthma)** only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

- Both:
- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
  - 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

**Renewal — (severe chronic spontaneous urticaria)** only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- Either:
- 1 Patient has previously adequately responded\* to 6 doses of omalizumab; or
  - 2 Both:
    - 2.1 Patient has previously had a complete response\* to 6 doses of omalizumab; and
    - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: \*Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/cyclosporin does not justify the funding of omalizumab.

**PERTUZUMAB – PCT only – Specialist – Special Authority see SA1606 below**

Inj 30 mg per ml, 14 ml vial.....	3,927.00	1	✓ <b>Perjeta</b>
Inj 420 mg for ECP .....	3,927.00	420 mg OP	✓ <b>Baxter</b>

► **SA1606** **Special Authority for Subsidy**

**Initial application — (metastatic breast cancer)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
  - 2.1 Patient is chemotherapy treatment naïve; or
  - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

**Renewal — (metastatic breast cancer)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

- Both:
- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

**RITUXIMAB – PCT only – Specialist – Special Authority see SA1818 on the next page**

Inj 100 mg per 10 ml vial .....	1,075.50	2	✓ <b>Mabthera</b>
Inj 500 mg per 50 ml vial .....	2,688.30	1	✓ <b>Mabthera</b>
Inj 1 mg for ECP .....	5.64	1 mg	✓ <b>Baxter</b>



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## ►SA1818 Special Authority for Subsidy

**Initial application — (ABO-incompatible renal transplant)** only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible renal transplant\*.

Note: Indications marked with \* are unapproved indications.

**Initial application — (ANCA associated vasculitis)** from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
  - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
  - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
  - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
  - 3.4 Patient is a female of child-bearing potential; or
  - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with \* are unapproved indications.

**Renewal — (ANCA associated vasculitis)** from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Initial application — (Antibody-mediated renal transplant rejection)** only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated renal transplant rejection\*.

Note: Indications marked with \* are unapproved indications.

**Initial application — (Chronic lymphocytic leukaemia)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 The patient is rituximab treatment naive; and
- 3 Either:
  - 3.1 The patient is chemotherapy treatment naive; or
  - 3.2 Both:
    - 3.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
    - 3.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; and
- 4 The patient has good performance status; and
- 5 The patient does not have chromosome 17p deletion CLL; and
- 6 Rituximab to be administered in combination with fludarabine and cyclophosphamide or bendamustine for a maximum of 6 treatment cycles; and

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

continued...

- 7 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

**Renewal — (Chronic lymphocytic leukaemia)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
- 2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
- 3 The patient does not have chromosome 17p deletion CLL; and
- 4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide or bendamustine for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

**Renewal — (Neuromyelitis Optica Spectrum Disorder)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m<sup>2</sup> administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

**Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD))** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m<sup>2</sup> administered weekly for four weeks; and
- 2 Either:
  - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
  - 2.2 All of the following:
    - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
    - 2.2.2 The patient is receiving treatment with mycophenolate; and
    - 2.2.3 The patients is receiving treatment with corticosteroids.

**Initial application — (Post-transplant)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with \* are unapproved indications.

**Renewal — (Post-transplant)** from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with \* are unapproved indications.

**Initial application — (Severe Refractory Myasthenia Gravis)** only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m<sup>2</sup> of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
  - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
  - 2.2 Both:
    - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
    - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

**Renewal — (Severe Refractory Myasthenia Gravis)** only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m<sup>2</sup> of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
  - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
  - 3.2 Both:
    - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
    - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

**Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS))** only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS\* or FRNS\*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS))** only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Initial application — (Steroid resistant nephrotic syndrome (SRNS))** only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SRNS\* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Renewal — (Steroid resistant nephrotic syndrome (SRNS))** only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Initial application — (aggressive CD20 positive NHL)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
  - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
  - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
  - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

**Renewal — (aggressive CD20 positive NHL)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

**Initial application — (haemophilia with inhibitors)** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

**Renewal — (haemophilia with inhibitors)** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

**Initial application — (immune thrombocytopenic purpura (ITP))** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per microlitre; or
  - 1.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
  - 2.1 Treatment with steroids and splenectomy have been ineffective; or
  - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
  - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy).

Note: Indications marked with \* are unapproved indications.

**Renewal — (immune thrombocytopenic purpura (ITP))** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m<sup>2</sup> weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

**Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia\*)** from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
  - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia\* requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenström macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

continued...

**Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia\*)** from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

**Initial application — (previous use)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with rituximab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 haemophilia with inhibitors; or
  - 2.2 rheumatoid arthritis; or
  - 2.3 severe cold haemagglutinin disease (CHAD); or
  - 2.4 warm autoimmune haemolytic anaemia (warm AIHA); or
  - 2.5 immune thrombocytopenic purpura (ITP); or
  - 2.6 thrombotic thrombocytopenic purpura (TTP); or
  - 2.7 pure red cell aplasia (PRCA); or
  - 2.8 ANCA associated vasculitis; or
  - 2.9 treatment refractory systemic lupus erythematosus (SLE); or
  - 2.10 steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS).

**Initial application — (pure red cell aplasia (PRCA))** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with \* are unapproved indications.

**Renewal — (pure red cell aplasia (PRCA))** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with \* are unapproved indications.

**Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
  - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
  - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
  - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

leflunomide alone or in combination with oral or parenteral methotrexate; and

6 Either:

- 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
- 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

7 Either:

- 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and

8 Either:

- 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and

9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

**Initial application — (rheumatoid arthritis - prior TNF inhibitor use)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Both:

- 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and

1.2 Either:

- 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
- 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and

2 Either:

- 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and

3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

**Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

- 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and

2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and

3 Either:

- 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and

4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

**Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

**Initial application — (severe cold haemagglutinin disease (CHAD))** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Patient has cold haemagglutinin disease\*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms.

Note: Indications marked with \* are unapproved indications.

**Renewal — (severe cold haemagglutinin disease (CHAD))** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m<sup>2</sup> weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

**Initial application — (thrombotic thrombocytopenic purpura (TTP))** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

Either:

- 1 Patient has thrombotic thrombocytopenic purpura\* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
- 2 Patient has acute idiopathic thrombotic thrombocytopenic purpura\* with neurological or cardiovascular pathology.

Note: Indications marked with \* are unapproved indications.

**Renewal — (thrombotic thrombocytopenic purpura (TTP))** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura\*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

**Initial application — (treatment refractory systemic lupus erythematosus (SLE))** only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting

continued...



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE\*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

**Renewal — (treatment refractory systemic lupus erythematosus (SLE))** only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE\* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

**Initial application — (warm autoimmune haemolytic anaemia (warm AIHA))** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Patient has warm autoimmune haemolytic anaemia\*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin.

Note: Indications marked with \* are unapproved indications.

**Renewal — (warm autoimmune haemolytic anaemia (warm AIHA))** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m<sup>2</sup> weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

SECUKINUMAB – Special Authority see [SA1754 below](#) – Retail pharmacy

Inj 150 mg per ml, 1 ml prefilled syringe.....	2	✓ Cosentyx
--	---	------------

➔ **SA1754** Special Authority for Subsidy

**Initial application — (severe chronic plaque psoriasis – second-line biologic)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
  - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and

4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

**Initial application — (severe chronic plaque psoriasis – first-line biologic)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Either:

- 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
- 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and

2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and

3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and

4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

**Renewal — (severe chronic plaque psoriasis – first and second-line biologic)** only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
- 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and

2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

**SILTUXIMAB** – Special Authority see [SA1596 below](#) – Retail pharmacy

Note: Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Inj 100 mg vial .....	770.57	1	✓ Sylvant
Inj 400 mg vial .....	3,082.33	1	✓ Sylvant

## ►SA1596 Special Authority for Subsidy

**Initial application** only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

**Renewal** only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
TOCILIZUMAB – PCT only – Special Authority see <a href="#">SA1781</a> below				
Inj 20 mg per ml, 4 ml vial.....	220.00	1	✓	Actemra
Inj 20 mg per ml, 10 ml vial.....	550.00	1	✓	Actemra
Inj 20 mg per ml, 20 ml vial.....	1,100.00	1	✓	Actemra
Inj 1 mg for ECP .....	2.85	1 mg	✓	Baxter

► **SA1781** Special Authority for Subsidy

**Initial application — (cytokine release syndrome)** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient is enrolled in the Children's Oncology Group AALL1331 trial; and
  - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
  - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses; or
- 2 All of the following:
  - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
  - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
  - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

**Initial application — (previous use)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 rheumatoid arthritis; or
  - 2.2 systemic juvenile idiopathic arthritis; or
  - 2.3 adult-onset Still's disease; or
  - 2.4 polyarticular juvenile idiopathic arthritis; or
  - 2.5 idiopathic multicentric Castleman's disease.

**Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept))** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
  - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
  - 3.2 Both:
    - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
    - 3.2.2 Either:
      - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
      - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

**Initial application — (Rheumatoid Arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
  - 3.1 Treatment with methotrexate is contraindicated; or
  - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
  - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
  - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
  - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
  - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
  - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Initial application — (systemic juvenile idiopathic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

**Initial application — (adult-onset Still's disease)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Either:
    - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
    - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
    - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

**Initial application — (polyarticular juvenile idiopathic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for juvenile idiopathic arthritis (JIA); and
- 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or

2 All of the following:

- 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
- 2.2 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
- 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m<sup>2</sup> weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
- 2.4 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.5 Both:

2.5.1 Either:

- 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
- 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and

2.5.2 Physician's global assessment indicating severe disease.

**Initial application — (idiopathic multicentric Castleman's disease)** only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

**Renewal — (Rheumatoid Arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

**Renewal — (systemic juvenile idiopathic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

**Renewal — (adult-onset Still's disease)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status.

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

**Renewal — (polyarticular juvenile idiopathic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

**Renewal — (idiopathic multicentric Castleman's disease)** only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB – PCT only – Specialist – Special Authority see [SA1632 below](#)

Inj 150 mg vial .....	1,350.00	1	✓ Herceptin
Inj 440 mg vial .....	3,875.00	1	✓ Herceptin
Inj 1 mg for ECP .....	9.36	1 mg	✓ Baxter

## ►SA1632 Special Authority for Subsidy

**Initial application — (metastatic breast cancer)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
  - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
  - 2.2 Both:
    - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
    - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
  - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
  - 3.2 All of the following:
    - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
    - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
    - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

**Renewal — (metastatic breast cancer)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 4 Trastuzumab to be discontinued at disease progression.

**Initial application — (early breast cancer)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
  - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
  - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
  - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
  - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
  - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

**Renewal — (early breast cancer\*)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
  - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
  - 3.2 Both:
    - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
    - 3.2.2 The cancer did not progress whilst on lapatinib; or
  - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
  - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
  - 4.2 All of the following:
    - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
    - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
    - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: \* For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

## Programmed Cell Death-1 (PD-1) Inhibitors

NIVOLUMAB – PCT only – Specialist – Special Authority see [SA1656 below](#)

Inj 10 mg per ml, 4 ml vial.....	1,051.98	1	✓ Opdivo
Inj 10 mg per ml, 10 ml vial.....	2,629.96	1	✓ Opdivo
Inj 1 mg for ECP .....	27.62	1 mg	✓ Baxter

► [SA1656](#) Special Authority for Subsidy

**Initial application — (unresectable or metastatic melanoma)** only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

All of the following:

- 1 Patient has metastatic or unresectable melanoma stage III or IV; and
- 2 Patient has measurable disease as defined by the presence of at least one CT or MRI measurable lesion; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
  - 4.1 Patient has not received funded pembrolizumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Nivolumab is to be used at a maximum dose of 3 mg/kg every 2 weeks for a maximum of 12 weeks (6 cycles); and
- 6 Baseline measurement of overall tumour burden is documented (see Note); and
- 7 Documentation confirming that the patient has been informed and acknowledges that the initial funded treatment period of nivolumab will not be continued beyond 12 weeks (6 cycles) if their disease progresses during this time.

**Renewal — (unresectable or metastatic melanoma)** only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
  - 1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
  - 1.3 Patient has stable disease according to RECIST criteria (see Note); and
- 2 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; and
- 3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; and
- 5 Nivolumab will be used at a maximum dose of 3 mg/kg every 2 weeks for a maximum of 12 weeks (6 cycles).

Notes: Disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Target lesion measurements should be assessed using CT or MRI imaging with the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB – PCT only – Specialist – Special Authority see [SA1657 on the next page](#)

Inj 25 mg per ml, 4 ml vial.....	4,680.00	1	✓ Keytruda
Inj 1 mg for ECP .....	49.14	1 mg	✓ Baxter



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## ►SA1657 Special Authority for Subsidy

**Initial application — (unresectable or metastatic melanoma)** only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma stage III or IV; and
- 2 Patient has measurable disease as defined by the presence of at least one CT or MRI measurable lesion; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
  - 4.1 Patient has not received funded nivolumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Pembrolizumab is to be used at a maximum dose of 2 mg/kg every 3 weeks for a maximum of 12 weeks (4 cycles); and
- 6 Baseline measurement of overall tumour burden is documented (see Note); and
- 7 Documentation confirming that the patient has been informed and acknowledges that the initial funded treatment period of pembrolizumab will not be continued beyond 12 weeks (4 cycles) if their disease progresses during this time.

**Renewal — (unresectable or metastatic melanoma)** only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
  - 1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
  - 1.3 Patient has stable disease according to RECIST criteria (see Note); and
- 2 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; and
- 3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; and
- 5 Pembrolizumab will be used at a maximum dose of 2 mg/kg every 3 weeks for a maximum of 12 weeks (4 cycles).

Notes: Disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Target lesion measurements should be assessed using CT or MRI imaging with the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>Other Immunosuppressants</b>				
<b>CICLOSPORIN</b>				
Cap 25 mg .....	44.63	50	✓	Neoral
Cap 50 mg .....	88.91	50	✓	Neoral
Cap 100 mg .....	177.81	50	✓	Neoral
Oral liq 100 mg per ml .....	198.13	50 ml OP	✓	Neoral
<b>EVEROLIMUS – Special Authority see SA1491 below – Retail pharmacy</b>				
Wastage claimable				
Tab 10 mg .....	6,512.29	30	✓	Afinitor
Tab 5 mg .....	4,555.76	30	✓	Afinitor
<b>►SA1491 Special Authority for Subsidy</b>				
<b>Initial application</b> only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:				
Both:				
1 Patient has tuberous sclerosis; and				
2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.				
<b>Renewal</b> only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria:				
All of the following:				
1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and				
2 The treatment remains appropriate and the patient is benefiting from treatment; and				
3 Everolimus to be discontinued at progression of SEGAs.				
Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.				
<b>SIROLIMUS – Special Authority see SA0866 below – Retail pharmacy</b>				
Tab 1 mg .....	749.99	100	✓	Rapamune
Tab 2 mg .....	1,499.99	100	✓	Rapamune
Oral liq 1 mg per ml .....	449.99	60 ml OP	✓	Rapamune
<b>►SA0866 Special Authority for Subsidy</b>				
<b>Initial application</b> from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.				
Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:				
• GFR < 30 ml/min; or				
• Rapidly progressive transplant vasculopathy; or				
• Rapidly progressive obstructive bronchiolitis; or				
• HUS or TTP; or				
• Leukoencephalopathy; or				
• Significant malignant disease				
<b>TACROLIMUS – Special Authority see SA1745 below – Retail pharmacy</b>				
Cap 0.5 mg .....	49.60	100	✓	Tacrolimus Sandoz
Cap 0.75 mg .....	99.30	100	✓	Tacrolimus Sandoz
Cap 1 mg .....	84.30	100	✓	Tacrolimus Sandoz
Cap 5 mg .....	248.20	50	✓	Tacrolimus Sandoz
<b>►SA1745 Special Authority for Subsidy</b>				
<b>Initial application — (organ transplant)</b> only from a relevant specialist. Approvals valid without further renewal unless notified				

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

**Initial application — (non-transplant indications\*)** only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with \* are unapproved indications

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

## Antiallergy Preparations

### Allergic Emergencies

ICATIBANT – Special Authority see [SA1558 below](#) – Retail pharmacy

Inj 10 mg per ml, 3 ml prefilled syringe .....	2,668.00	1	✓ <b>Firazyr</b>
--	----------	---	------------------

► **SA1558** **Special Authority for Subsidy**

**Initial application** only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

### Allergy Desensitisation

► **SA1367** **Special Authority for Subsidy**

**Initial application** only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

**Renewal** only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT – Special Authority see [SA1367 above](#) – Retail pharmacy

Maintenance kit - 6 vials 120 mcg freeze dried venom, with diluent .....	285.00	1 OP	✓ <b>Venomil</b> <sup>S29</sup>
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent 9 ml, 3 diluent 1.8 ml .....	305.00	1 OP	✓ <b>Albey</b>
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent .....	305.00	1 OP	✓ <b>Hymenoptera</b> <sup>S29</sup>

WASP VENOM ALLERGY TREATMENT – Special Authority see [SA1367 above](#) – Retail pharmacy

Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml .....	305.00	1 OP	✓ <b>Albey</b>
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze dried venom, with diluent .....	305.00	1 OP	✓ <b>Hymenoptera</b> <sup>S29</sup>
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze dried venom, with diluent .....	305.00	1 OP	✓ <b>Venomil</b> <sup>S29</sup>
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze dried venom, with diluent .....	305.00	1 OP	✓ <b>Hymenoptera</b> <sup>S29</sup>
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml .....	305.00	1 OP	✓ <b>Albey</b>
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze dried venom, with diluent .....	305.00	1 OP	✓ <b>Venomil</b> <sup>S29</sup>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>Antihistamines</b>				
<b>CETIRIZINE HYDROCHLORIDE</b>				
* Tab 10 mg .....	1.12	100	✓	<b>Zista</b>
Zista to be Sole Supply on 1 November 2019				
* Oral liq 1 mg per ml .....	2.99	200 ml	✓	<b>Histaclear</b>
<b>CHLORPHENIRAMINE MALEATE</b>				
* Oral liq 2 mg per 5 ml .....	9.37	500 ml	✓	<b>Histafen</b>
<b>DEXTROCHLORPHENIRAMINE MALEATE</b>				
* Tab 2 mg .....	2.02	40		
	(8.40)			Polaramine
	1.01	20		
	(5.99)			Polaramine
* Oral liq 2 mg per 5 ml .....	1.77	100 ml		Polaramine
	(10.29)			Polaramine
<b>FEXOFENADINE HYDROCHLORIDE</b>				
* Tab 60 mg .....	4.34	20		
	(8.23)			Telfast
* Tab 120 mg .....	4.74	10		
	(8.23)			Telfast
	14.22	30		
	(26.44)			Telfast
<b>LORATADINE</b>				
* Tab 10 mg .....	1.28	100	✓	<b>Lorafix</b>
* Oral liq 1 mg per ml .....	2.15	120 ml	✓	<b>Lorfast</b>
<b>PROMETHAZINE HYDROCHLORIDE</b>				
* Tab 10 mg .....	1.68	50	✓	<b>Allersoothe</b>
* Tab 25 mg .....	1.89	50	✓	<b>Allersoothe</b>
* Oral liq 1 mg per 1 ml .....	2.69	100 ml	✓	<b>Allersoothe</b>
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO ....	15.54	5	✓	<b>Hospira</b>
<b>Inhaled Corticosteroids</b>				
<b>BECLOMETHASONE DIPROPIONATE</b>				
Aerosol inhaler, 50 mcg per dose .....	9.30	200 dose OP	✓	<b>Qvar</b>
Aerosol inhaler, 50 mcg per dose CFC-free .....	8.54	200 dose OP	✓	<b>Beclazone 50</b>
Aerosol inhaler, 100 mcg per dose .....	15.50	200 dose OP	✓	<b>Qvar</b>
Aerosol inhaler, 100 mcg per dose CFC-free .....	12.50	200 dose OP	✓	<b>Beclazone 100</b>
Aerosol inhaler, 250 mcg per dose CFC-free .....	22.67	200 dose OP	✓	<b>Beclazone 250</b>
<b>BUDESONIDE</b>				
Powder for inhalation, 100 mcg per dose .....	17.00	200 dose OP	✓	<b>Pulmicort Turbuhaler</b>
Powder for inhalation, 200 mcg per dose .....	19.00	200 dose OP	✓	<b>Pulmicort Turbuhaler</b>
Powder for inhalation, 400 mcg per dose .....	32.00	200 dose OP	✓	<b>Pulmicort Turbuhaler</b>

# RESPIRATORY SYSTEM AND ALLERGIES

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>FLUTICASONE</b>				
Aerosol inhaler, 50 mcg per dose.....	4.68	120 dose OP	✓	Floair
Aerosol inhaler, 50 mcg per dose CFC-free .....	7.50	120 dose OP	✓	Flixotide
Powder for inhalation, 50 mcg per dose.....	7.50	60 dose OP	✓	Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose.....	7.50	60 dose OP	✓	Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose.....	7.22	120 dose OP	✓	Floair
Aerosol inhaler, 125 mcg per dose CFC-free .....	13.60	120 dose OP	✓	Flixotide
Aerosol inhaler, 250 mcg per dose.....	10.18	120 dose OP	✓	Floair
Aerosol inhaler, 250 mcg per dose CFC-free .....	27.20	120 dose OP	✓	Flixotide
Powder for inhalation, 250 mcg per dose .....	13.60	60 dose OP	✓	Flixotide Accuhaler

## Inhaled Long-acting Beta-adrenoceptor Agonists

<b>EFORMOTEROL FUMARATE</b>				
Powder for inhalation, 12 mcg per dose, and monodose device.....	20.64 (35.80)	60 dose		Foradil
<b>EFORMOTEROL FUMARATE DIHYDRATE</b>				
Powder for inhalation 4.5 mcg per dose, breath activated (equivalent to eformoterol fumarate 6 mcg metered dose).....	10.32 (16.90)	60 dose OP		Oxis Turbuhaler
<b>INDACATEROL</b>				
Powder for inhalation 150 mcg.....	61.00	30 dose OP	✓	Onbrez Breezhaler
Powder for inhalation 300 mcg.....	61.00	30 dose OP	✓	Onbrez Breezhaler
<b>SALMETEROL</b>				
Aerosol inhaler CFC-free, 25 mcg per dose.....	25.00	120 dose OP	✓	Serevent
Aerosol inhaler 25 mcg per dose.....	9.90	120 dose OP	✓	Meterol
Powder for inhalation, 50 mcg per dose, breath activated .....	25.00	60 dose OP	✓	Serevent Accuhaler

## Inhaled Corticosteroids with Long-Acting Beta-Adrenoceptor Agonists

<b>BUDESONIDE WITH EFORMOTEROL</b>				
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg .....	18.23	120 dose OP	✓	Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6 mcg .....	33.74	120 dose OP	✓	Symbicort Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg .....	21.40	120 dose OP	✓	Vannair
Powder for inhalation 200 mcg with eformoterol fumarate 6 mcg .....	44.08	120 dose OP	✓	Symbicort Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate 12 mcg – No more than 2 dose per day .....	44.08	60 dose OP	✓	Symbicort Turbuhaler 400/12
<b>FLUTICASONE FUROATE WITH VILANTEROL</b>				
Powder for inhalation 100 mcg with vilanterol 25 mcg .....	44.08	30 dose OP	✓	Breo Ellipta
<b>FLUTICASONE WITH SALMETEROL</b>				
Aerosol inhaler 50 mcg with salmeterol 25 mcg .....	14.58	120 dose OP	✓	RexAir
	33.74		✓	Seretide
Aerosol inhaler 125 mcg with salmeterol 25 mcg .....	16.83	120 dose OP	✓	RexAir
	44.08		✓	Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg – No more than 2 dose per day.....	33.74	60 dose OP	✓	Seretide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg – No more than 2 dose per day.....	44.08	60 dose OP	✓	Seretide Accuhaler

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

## Beta-Adrenoceptor Agonists

### SALBUTAMOL

Oral liq 400 mcg per ml .....	20.00	150 ml	✓ <b>Ventolin</b>
Infusion 1 mg per ml, 5 ml .....	118.38	10	✓ <b>Ventolin</b>
Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO .....	53.00	5	✓ <b>Ventolin</b>

## Inhaled Beta-Adrenoceptor Agonists

### SALBUTAMOL

Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000 dose available on a PSO .....	3.80	200 dose OP	✓ <b>Respigen</b> ✓ <b>SalAir</b> Ventolin
(6.00)			
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO .....	3.93	20	✓ <b>Asthalin</b>
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO .....	4.03	20	✓ <b>Asthalin</b>

### TERBUTALINE SULPHATE

Powder for inhalation, 250 mcg per dose, breath activated .....	27.30	200 dose OP	✓ <b>Bricanyl Turbuhaler</b>
---	-------	-------------	------------------------------

## Anticholinergic Agents

### IPRATROPIUM BROMIDE

Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dose available on a PSO .....	16.20	200 dose OP	✓ <b>Atrovent</b>
Nebuliser soln, 250 mcg per ml, 1 ml ampoule – Up to 40 neb available on a PSO .....	3.35	20	✓ <b>Univent</b>
Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 neb available on a PSO .....	11.73	20	✓ <b>Univent</b>
Univent to be Sole Supply on 1 January 2020			

## Inhaled Beta-Adrenoceptor Agonists with Anticholinergic Agents

### SALBUTAMOL WITH IPRATROPIUM BROMIDE

Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg per dose CFC-free .....	12.19	200 dose OP	✓ <b>Duolin HFA</b>
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule – Up to 20 neb available on a PSO .....	5.20	20	✓ <b>Duolin</b>

## Long-Acting Muscarinic Antagonists

### GLYCOPYRRONIUM – Subsidy by endorsement

- Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium.
- Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

Powder for inhalation 50 mcg per dose .....	61.00	30 dose OP	✓ <b>Seebri Breezhaler</b>
---	-------	------------	----------------------------

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

## TIOTROPIUM BROMIDE – Subsidy by endorsement

- Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

Powder for inhalation, 18 mcg per dose .....	50.37	30 dose	✓	<b>Spiriva</b>
Soln for inhalation 2.5 mcg per dose .....	50.37	60 dose OP	✓	<b>Spiriva Respimat</b>

## UMECLIDINIUM – Subsidy by endorsement

- Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

Powder for inhalation 62.5 mcg per dose .....	61.50	30 dose OP	✓	<b>Incruse Ellipta</b>
---	-------	------------	---	------------------------

## Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

### ►SA1584 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- Both:
- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
  - 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

**Renewal** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- Both:
- 1 Patient is compliant with the medication; and
  - 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL – Special Authority see [SA1584 above](#) – Retail pharmacy

Powder for Inhalation 50 mcg with indacaterol 110 mcg .....	81.00	30 dose OP	✓	<b>Ultibro Breezhaler</b>
---	-------	------------	---	---------------------------

TIOTROPIUM BROMIDE WITH OLODATEROL – Special Authority see [SA1584 above](#) – Retail pharmacy

Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg .....	81.00	60 dose OP	✓	<b>Spiolto Respimat</b>
---	-------	------------	---	-------------------------

UMECLIDINIUM WITH VILANTEROL – Special Authority see [SA1584 above](#) – Retail pharmacy

Powder for inhalation 62.5 mcg with vilanterol 25 mcg .....	77.00	30 dose OP	✓	<b>Anoro Ellipta</b>
---	-------	------------	---	----------------------

## Antifibrotics

NINTEDANIB – Special Authority see [SA1755 below](#) – Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

Cap 100 mg .....	2,554.00	60 OP	✓	<b>Ofev</b>
Cap 150 mg .....	3,870.00	60 OP	✓	<b>Ofev</b>

### ►SA1755 Special Authority for Subsidy

**Initial application** — (**idiopathic pulmonary fibrosis**) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and

continued...



	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
		✓	

continued...

- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with pirfenidone; or
  - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

**Renewal — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

**PIRFENIDONE** – Retail pharmacy-Specialist – Special Authority see [SA1748 below](#)

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Cap 267 mg – Wastage claimable.....3,645.00      270      ✓ **Esbriet**

➔ **SA1748** **Special Authority for Subsidy**

**Initial application — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 80% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with nintedanib; or
  - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

**Renewal — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

## Leukotriene Receptor Antagonists

### MONTELUKAST

* Tab 4 mg .....	4.25	28	✓	Montelukast Mylan
	5.25		✓	Apo-Montelukast
Montelukast Mylan to be Sole Supply on 1 January 2020				
* Tab 5 mg .....	4.25	28	✓	Montelukast Mylan
	5.50		✓	Apo-Montelukast
Montelukast Mylan to be Sole Supply on 1 January 2020				
* Tab 10 mg .....	3.95	28	✓	Montelukast Mylan
	5.65		✓	Accord <sup>S29</sup>
			✓	Apo-Montelukast
Montelukast Mylan to be Sole Supply on 1 January 2020				
(Apo-Montelukast Tab 4 mg to be delisted 1 January 2020)				
(Apo-Montelukast Tab 5 mg to be delisted 1 January 2020)				
(Accord <sup>S29</sup> Tab 10 mg to be delisted 1 January 2020)				
(Apo-Montelukast Tab 10 mg to be delisted 1 January 2020)				

## Mast Cell Stabilisers

### NEDOCROMIL

Aerosol inhaler, 2 mg per dose CFC-free.....28.07 112 dose OP ✓ Tilade

### SODIUM CROMOGLICATE

Aerosol inhaler, 5 mg per dose CFC-free.....28.07 112 dose OP ✓ Intal Forte CFC Free

## Methylxanthines

### AMINOPHYLLINE

\* Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a  
PSO .....124.37 5 ✓ DBL Aminophylline

### THEOPHYLLINE

\* Tab long-acting 250 mg.....23.02 100 ✓ Nuelin-SR  
Nuelin-SR to be Sole Supply on 1 January 2020  
\* Oral liq 80 mg per 15 ml .....16.60 500 ml ✓ Nuelin  
Nuelin to be Sole Supply on 1 January 2020

## Mucolytics

### DORNASE ALFA – Special Authority see SA0611 below – Retail pharmacy

Nebuliser soln, 2.5 mg per 2.5 ml ampoule .....250.00 6 ✓ Pulmozyme

### ►SA0611 Special Authority for Subsidy

Special Authority approved by the Cystic Fibrosis Advisory Panel

Notes: Application details may be obtained from PHARMAC's website <http://www.pharmac.govt.nz> or:

The Co-ordinator, Cystic Fibrosis Advisory Panel Phone: (04) 460 4990  
PHARMAC, PO Box 10 254 Facsimile: (04) 916 7571  
Wellington Email: [CFPanel@pharmac.govt.nz](mailto:CFPanel@pharmac.govt.nz)

Prescriptions for patients approved for treatment must be written by respiratory physicians or paediatricians who have experience and expertise in treating cystic fibrosis.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>SODIUM CHLORIDE</b>				
Not funded for use as a nasal drop.				
Soln 7% .....	24.50	90 ml OP	✓	<b>Biomed</b>
Biomed to be Sole Supply on 1 November 2019				

## Nasal Preparations

### Allergy Prophylactics

<b>BECLOMETHASONE DIPROPIONATE</b>				
Metered aqueous nasal spray, 50 mcg per dose .....	2.35 (5.26)	200 dose OP		Alanase
Metered aqueous nasal spray, 100 mcg per dose .....	2.46 (6.00)	200 dose OP		Alanase
<i>(Alanase Metered aqueous nasal spray, 50 mcg per dose to be delisted 1 January 2020)</i>				
<i>(Alanase Metered aqueous nasal spray, 100 mcg per dose to be delisted 1 January 2020)</i>				
<b>BUDESONIDE</b>				
Metered aqueous nasal spray, 50 mcg per dose .....	2.59	200 dose OP	✓	<b>SteroClear</b>
Metered aqueous nasal spray, 100 mcg per dose .....	2.87	200 dose OP	✓	<b>SteroClear</b>
<b>FLUTICASONE PROPIONATE</b>				
Metered aqueous nasal spray, 50 mcg per dose .....	1.98	120 dose OP	✓	<b>Flixonase Hayfever &amp; Allergy</b>
<b>IPRATROPIUM BROMIDE</b>				
Aqueous nasal spray, 0.03%.....	4.61	15 ml OP	✓	<b>Univent</b>

## Respiratory Devices

<b>MASK FOR SPACER DEVICE</b>				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
c) Only for children aged six years and under				
Small.....	2.20	1	✓	<b>e-chamber Mask</b>
<b>PEAK FLOW METER</b>				
a) Up to 25 dev available on a PSO				
b) Only on a PSO				
Low range.....	9.54	1	✓	<b>Mini-Wright AFS Low Range</b>
Normal range.....	9.54	1	✓	<b>Mini-Wright Standard</b>
<b>SPACER DEVICE</b>				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
220 ml (single patient) .....	2.95	1	✓	<b>e-chamber Turbo</b>
510 ml (single patient) .....	5.12	1	✓	<b>e-chamber La Grande</b>
800 ml.....	6.50	1	✓	<b>Volumatic</b>

## Respiratory Stimulants

<b>CAFFEINE CITRATE</b>				
Oral liq 20 mg per ml (10 mg base per ml) .....	15.10	25 ml OP	✓	<b>Biomed</b>
Biomed to be Sole Supply on 1 November 2019				

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

## Ear Preparations

### ACETIC ACID WITH 1, 2- PROPANEDIOL DIACETATE AND BENZETHONIUM

For Vosol ear drops with hydrocortisone powder refer Standard Formulae, [page 235](#)

Ear drops 2% with 1, 2-Propanediol diacetate 3% and

benzethonium chloride 0.02% .....6.97 35 ml OP ✓ **Vosol**

### FLUMETASONE PIVALATE

Ear drops 0.02% with clioquinol 1% .....4.46 7.5 ml OP ✓ **Locacorten-Viaform**

ED's

✓ **Locorten-Vioform**

### TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCIN AND NYSTATIN

Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate

2.5 mg and gramicidin 250 mcg per g .....5.16 7.5 ml OP ✓ **Kenacomb**

## Ear/Eye Preparations

### DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN

Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and

gramicidin 50 mcg per ml .....4.50 8 ml OP  
(9.27) Sofradex

### FRAMYCETIN SULPHATE

Ear/Eye drops 0.5%.....4.13 8 ml OP  
(8.65) Soframycin

## Eye Preparations

Eye preparations are only funded for use in the eye, unless explicitly stated otherwise.

### Anti-Infective Preparations

#### ACICLOVIR

\* Eye oint 3% .....14.92 4.5 g OP ✓ **ViruPOS**

#### CHLORAMPHENICOL

Eye oint 1% .....2.48 4 g OP ✓ **Chlorsig**

Eye drops 0.5% .....1.54 10 ml OP ✓ **Chlorafast**

a) Funded for use in the ear\*. Indications marked with \* are unapproved indications.

b) Chlorafast to be Sole Supply on 1 November 2019

#### CIPROFLOXACIN

Eye drops 0.3% – Subsidy by endorsement.....9.99 5 ml OP ✓ **Ciprofloxacin Teva**

When prescribed for the treatment of bacterial keratitis or severe bacterial conjunctivitis resistant to chloramphenicol; or for the second line treatment of chronic suppurative otitis media (CSOM)\*; and the prescription is endorsed accordingly.

Note: Indication marked with a \* is an unapproved indication.

#### GENTAMICIN SULPHATE

Eye drops 0.3% .....11.40 5 ml OP ✓ **Genoptic**

#### PROPAMIDINE ISETHIONATE

\* Eye drops 0.1% .....2.97 10 ml OP  
(14.55) Brolene

#### SODIUM FUSIDATE [FUSIDIC ACID]

Eye drops 1% .....5.29 5 g OP ✓ **Fucithalmic**

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>TOBRAMYCIN</b>				
Eye oint 0.3% .....	10.45	3.5 g OP	✓	<b>Tobrex</b>
Eye drops 0.3% .....	11.48	5 ml OP	✓	<b>Tobrex</b>

## Corticosteroids and Other Anti-Inflammatory Preparations

### DEXAMETHASONE

* Eye oint 0.1% .....	5.86	3.5 g OP	✓	<b>Maxidex</b>
* Eye drops 0.1% .....	4.50	5 ml OP	✓	<b>Maxidex</b>
Ocular implant 700 mcg – Special Authority see <a href="#">SA1680 below</a> – Retail pharmacy.....	1,444.50	1	✓	<b>Ozurdex</b>

#### ►SA1680 Special Authority for Subsidy

**Initial application — (Diabetic macular oedema)** only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 - 6/48 with functional awareness of reduction in vision; and
- 3 Either:
  - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
  - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

**Renewal — (Diabetic macular oedema)** only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

**Initial application — (Women of child bearing age with diabetic macular oedema)** only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 - 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

**Renewal — (Women of child bearing age with diabetic macular oedema)** only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

### DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

* Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per g.....	5.39	3.5 g OP	✓	<b>Maxitrol</b>
* Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml .....	4.50	5 ml OP	✓	<b>Maxitrol</b>

### DICLOFENAC SODIUM

Eye drops 0.1% .....	13.80	5 ml OP	✓	<b>Voltaren Ophtha</b>
----------------------	-------	---------	---	------------------------

## SENSORY ORGANS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>FLUOROMETHOLONE</b>				
* Eye drops 0.1% .....	3.09	5 ml OP	✓ <b>FML</b>	
	5.20		✓ <b>Flucon</b>	
<b>LEVOCABASTINE</b>				
Eye drops 0.5 mg per ml .....	8.71	4 ml OP		Livostin
	(10.34)			
<b>LODOXAMIDE</b>				
Eye drops 0.1% .....	8.71	10 ml OP	✓ <b>Lomide</b>	
<b>PREDNISOLONE ACETATE</b>				
Eye drops 1% .....	3.93	10 ml OP	✓ <b>Prednisolone-AFT</b>	
	7.00	5 ml OP	✓ <b>Pred Forte</b>	
<b>PREDNISOLONE SODIUM PHOSPHATE</b> – Special Authority see <a href="#">SA1715 below</a> – Retail pharmacy				
Eye drops 0.5%, single dose (preservative free) .....	38.50	20 dose	✓ <b>Minims</b>	
			<b>Prednisolone</b>	

### ► [SA1715](#) Special Authority for Subsidy

**Initial application** only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

**Renewal** from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

### **SODIUM CROMOGLICATE**

Eye drops 2% .....	1.79	5 ml OP	✓ <b>Cromal</b> <small>§29</small>
			✓ <b>Rexacrom</b>

Rexacrom to be Sole Supply on 1 January 2020

## Glaucoma Preparations - Beta Blockers

### **BETAXOLOL**

* Eye drops 0.25% .....	11.80	5 ml OP	✓ <b>Betoptic S</b>
* Eye drops 0.5% .....	7.50	5 ml OP	✓ <b>Betoptic</b>

### **TIMOLOL**

* Eye drops 0.25% .....	1.43	5 ml OP	✓ <b>Arrow-Timolol</b>
* Eye drops 0.25%, gel forming .....	3.30	2.5 ml OP	✓ <b>Timoptol XE</b>
* Eye drops 0.5% .....	1.43	5 ml OP	✓ <b>Arrow-Timolol</b>
* Eye drops 0.5%, gel forming .....	3.78	2.5 ml OP	✓ <b>Timoptol XE</b>

*(Timoptol XE Eye drops 0.25%, gel forming to be delisted 1 January 2020)*

## Glaucoma Preparations - Carbonic Anhydrase Inhibitors

### **ACETAZOLAMIDE**

* Tab 250 mg .....	17.03	100	✓ <b>Diamox</b>
--------------------	-------	-----	-----------------

### **BRINZOLAMIDE**

* Eye drops 1% .....	9.77	5 ml OP	✓ <b>Azopt</b>
----------------------	------	---------	----------------

### **DORZOLAMIDE HYDROCHLORIDE**

* Eye drops 2% .....	9.77	5 ml OP	
	(17.44)		Trusopt

### **DORZOLAMIDE WITH TIMOLOL**

* Eye drops 2% with timolol 0.5% .....	2.87	5 ml OP	✓ <b>Dortimopt</b>
--	------	---------	--------------------

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
<b>Glaucoma Preparations - Prostaglandin Analogues</b>			
<b>BIMATOPROST</b>			
* Eye drops 0.03% .....	3.30	3 ml OP	✓ <b>Bimatoprost</b> <b>Multichem</b>
<b>LATANOPROST</b>			
* Eye drops 0.005% .....	1.57	2.5 ml OP	✓ <b>Teva</b>
<b>TRAVOPROST</b>			
* Eye drops 0.004% .....	7.30	5 ml OP	✓ <b>Travopt</b>
	19.50	2.5 ml OP	✓ <b>Travatan</b>

<b>Glaucoma Preparations - Other</b>			
<b>BRIMONIDINE TARTRATE</b>			
* Eye drops 0.2% .....	4.29	5 ml OP	✓ <b>Arrow-Brimonidine</b>
<b>BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE</b>			
* Eye drops 0.2% with timolol maleate 0.5% .....	18.50	5 ml OP	✓ <b>Combigan</b>
<b>PILOCARPINE HYDROCHLORIDE</b>			
* Eye drops 1% .....	4.26	15 ml OP	✓ <b>Isopto Carpine</b>
* Eye drops 2% .....	5.35	15 ml OP	✓ <b>Isopto Carpine</b>
* Eye drops 4% .....	7.99	15 ml OP	✓ <b>Isopto Carpine</b>
Subsidised for oral use pursuant to the Standard Formulae.			
* Eye drops 2% single dose – Special Authority see <a href="#">SA0895</a> below – Retail pharmacy .....	31.95	20 dose	✓ <b>Minims Pilocarpine</b>

➔ [SA0895](#) **Special Authority for Subsidy**

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:  
Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be “tools of trade” and are not approved as special authority items.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

<b>Mydriatics and Cycloplegics</b>			
<b>ATROPINE SULPHATE</b>			
* Eye drops 1% .....	17.36	15 ml OP	✓ <b>Atropt</b>
<b>CYCLOPENTOLATE HYDROCHLORIDE</b>			
* Eye drops 1% .....	8.76	15 ml OP	✓ <b>Cyclogyl</b>
<b>TROPICAMIDE</b>			
* Eye drops 0.5% .....	7.15	15 ml OP	✓ <b>Mydriacyl</b>
* Eye drops 1% .....	8.66	15 ml OP	✓ <b>Mydriacyl</b>

**Preparations for Tear Deficiency**

For acetylcysteine eye drops refer Standard Formulae, [page 235](#)

<b>HYPROMELLOSE</b>			
* Eye drops 0.5% .....	2.00 (3.92)	15 ml OP	Methopt

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>HYPROMELLOSE WITH DEXTRAN</b>				
* Eye drops 0.3% with dextran 0.1%.....	2.30	15 ml OP	✓	<b>Poly-Tears</b>
<b>POLYVINYL ALCOHOL</b>				
* Eye drops 1.4% .....	2.62	15 ml OP	✓	<b>Vistil</b>
* Eye drops 3% .....	3.68	15 ml OP	✓	<b>Vistil Forte</b>
<i>(Vistil Eye drops 1.4% to be delisted 1 January 2020)</i>				
<i>(Vistil Forte Eye drops 3% to be delisted 1 March 2020)</i>				

## Preservative Free Ocular Lubricants

### ►SA1388 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:  
Both:

- 1 Confirmed diagnosis by slit lamp of severe secretory dry eye; and
- 2 Either:
  - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
  - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

**Renewal** from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

**CARBOMER** – Special Authority see [SA1388 above](#) – Retail pharmacy

Ophthalmic gel 0.3%, 0.5 g .....	8.25	30	✓	<b>Poly-Gel</b>
----------------------------------	------	----	---	-----------------

**MACROGOL 400 AND PROPYLENE GLYCOL** – Special Authority see [SA1388 above](#) – Retail pharmacy

Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml .....	4.30	24	✓	<b>Systane Unit Dose</b>
--	------	----	---	--------------------------

**SODIUM HYALURONATE [HYALURONIC ACID]** – Special Authority see [SA1388 above](#) – Retail pharmacy

Eye drops 1 mg per ml .....	22.00	10 ml OP	✓	<b>Hylo-Fresh</b>
-----------------------------	-------	----------	---	-------------------

Hylo-Fresh has a 6 month expiry after opening. The Pharmacy Procedures Manual restriction allowing one bottle per month is not relevant and therefore only the prescribed dosage to the nearest OP may be claimed.

## Other Eye Preparations

**NAPHAZOLINE HYDROCHLORIDE**

* Eye drops 0.1% .....	4.15	15 ml OP	✓	<b>Naphcon Forte</b>
------------------------	------	----------	---	----------------------

**OLOPATADINE**

Eye drops 0.1% .....	10.00	5 ml OP	✓	<b>Patanol</b>
----------------------	-------	---------	---	----------------

**PARAFFIN LIQUID WITH WOOL FAT**

* Eye oint 3% with wool fat 3% .....	3.63	3.5 g OP	✓	<b>Poly-Visc</b>
--------------------------------------	------	----------	---	------------------

**RETINOL PALMITATE**

Eye oint 138 mcg per g.....	3.80	5 g OP	✓	<b>VitA-POS</b>
-----------------------------	------	--------	---	-----------------



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

## Various

### PHARMACY SERVICES

May only be claimed once per patient.

* Brand switch fee .....	4.50	1 fee	<ul style="list-style-type: none"> <li>✓ BSF Logem</li> <li>✓ BSF Mylan <ul style="list-style-type: none"> <li>Efavirenz</li> <li>Emtricitabine</li> <li>Tenofovir</li> </ul> </li> <li>✓ BSF Teva <ul style="list-style-type: none"> <li>Atazanavir Sulphate</li> </ul> </li> <li>✓ BSF Teva <ul style="list-style-type: none"> <li>Emtricitabine</li> <li>Tenofovir Disoprox</li> </ul> </li> </ul>
a) The Pharmacode for BSF Teva Atazanavir Sulphate is 2573857 - see also <a href="#">page 107</a>			
b) The Pharmacode for BSF Teva Emtricitabine Tenofovir Disoprox is 2573865 - see also <a href="#">page 103</a>			
c) The Pharmacode for BSF Mylan Efavirenz Emtricitabine Tenofovir is 2573873 - see also <a href="#">page 106</a>			
d) The Pharmacode for BSF Logem is 2575949 - see also <a href="#">page 127</a>			

(BSF Logem Brand switch fee to be delisted 1 January 2020)

(BSF Mylan Efavirenz Emtricitabine Tenofovir Brand switch fee to be delisted 1 December 2019)

(BSF Teva Atazanavir Sulphate Brand switch fee to be delisted 1 December 2019)

(BSF Teva Emtricitabine Tenofovir Disoprox Brand switch fee to be delisted 1 December 2019)

## Agents Used in the Treatment of Poisonings

### Antidotes

ACETYL CYSTEINE – Retail pharmacy-Specialist			
Inj 200 mg per ml, 10 ml ampoule .....	58.76	10	✓ <u>DBL Acetylcysteine</u>
NALOXONE HYDROCHLORIDE			
a) Up to 5 inj available on a PSO			
b) Only on a PSO			
* Inj 400 mcg per ml, 1 ml ampoule .....	22.60	5	✓ <u>DBL Naloxone Hydrochloride</u>

### Removal and Elimination

CHARCOAL			
* Oral liq 50 g per 250 ml .....	43.50	250 ml OP	✓ <u>Carbosorb-X</u>
a) Up to 250 ml available on a PSO			
b) Only on a PSO			
DEFERASIROX – Special Authority see <a href="#">SA1492 on the next page</a> – Retail pharmacy			
Wastage claimable			
Tab 125 mg dispersible .....	276.00	28	✓ <u>Exjade</u>
Tab 250 mg dispersible .....	552.00	28	✓ <u>Exjade</u>
Tab 500 mg dispersible .....	1,105.00	28	✓ <u>Exjade</u>

►SA1492 Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria:  
All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
  - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2\*; or
  - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
  - 3.3 Treatment with deferiprone has resulted in arthritis; or
  - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per µL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 - 1.0 cells per µL).

**Renewal** only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria:  
Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels.

DEFERIPRONE – Special Authority see SA1480 below – Retail pharmacy

Tab 500 mg .....	533.17	100	✓ Feriprox
Oral liq 100 mg per 1 ml .....	266.59	250 ml OP	✓ Feriprox

►SA1480 Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DEFERRIOXAMINE MESILATE

* Inj 500 mg vial .....	84.53	10	✓ <b>DBL</b> <b>Desferrioxamine</b> <b>Mesylate for Inj</b> <b>BP</b>
-------------------------	-------	----	--

SODIUM CALCIUM EDETATE

* Inj 200 mg per ml, 5 ml .....	53.31 (156.71)	6	Calcium Disodium Versenate
---------------------------------	-------------------	---	-------------------------------

## Standard Formulae

### ACETYLCYSTEINE EYE DROPS

Acetylcysteine inj 200 mg per ml, 10 ml	qs
Suitable eye drop base	qs

### PHENOBARBITONE ORAL LIQUID

Phenobarbitone Sodium	1 g
Glycerol BP	70 ml
Water	to 100 ml

### ASPIRIN AND CHLOROFORM APPLICATION

Aspirin Soluble tabs 300 mg	12 tabs
Chloroform	to 100 ml

### PHENOBARBITONE SODIUM PAEDIATRIC ORAL LIQUID (10 mg per ml)

Phenobarbitone Sodium	400 mg
Glycerol BP	4 ml
Water	to 40 ml

### CODEINE LINCTUS (3 mg per 5 ml)

Codeine phosphate	60 mg
Glycerol	40 ml
Preservative	qs
Water	to 100 ml

### PILOCARPINE ORAL LIQUID

Pilocarpine 4% eye drops	qs
Preservative	qs
Water	to 500 ml

### CODEINE LINCTUS (15 mg per 5 ml)

Codeine phosphate	300 mg
Glycerol	40 ml
Preservative	qs
Water	to 100 ml

(Preservative should be used if quantity supplied is for more than 5 days.)

### FOLINIC MOUTHWASH

Calcium folinate 15 mg tab	1 tab
Preservative	qs
Water	to 500 ml

### SALIVA SUBSTITUTE FORMULA

Methylcellulose	5 g
Preservative	qs
Water	to 500 ml

(Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.)

(Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.)

### MAGNESIUM HYDROXIDE 8% MIXTURE

Magnesium hydroxide paste 29%	275 g
Methyl hydroxybenzoate	1.5 g
Water	to 1,000 ml

### SODIUM CHLORIDE ORAL LIQUID

Sodium chloride inj 23.4%, 20 ml	qs
Water	qs

(Only funded if prescribed for treatment of hyponatraemia)

### METHADONE MIXTURE

Methadone powder	qs
Glycerol	qs
Water	to 100 ml

### VANCOMYCIN ORAL SOLUTION (50 mg per ml)

Vancomycin 500 mg injection	10 vials
Glycerol BP	40 ml
Water	to 100 ml

(Only funded if prescribed for treatment of Clostridium difficile following metronidazole failure)

### METHYL HYDROXYBENZOATE 10% SOLUTION

Methyl hydroxybenzoate	10 g
Propylene glycol	to 100 ml

(Use 1 ml of the 10% solution per 100 ml of oral liquid mixture)

### VOSOL EAR DROPS

### WITH HYDROCORTISONE POWDER 1%

Hydrocortisone powder	1%
Vosol Ear Drops	to 35 ml

### OMEPRAZOLE SUSPENSION

Omeprazole capules or powder	qs
Sodium bicarbonate powder BP	8.4 g
Water	to 100 ml

# EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>Extemporaneously Compounded Preparations and Galenicals</b>				
<b>BENZOIN</b>				
Tincture compound BP .....	24.42 (39.90)	500 ml		Pharmacy Health
<i>(Pharmacy Health Tincture compound BP to be delisted 1 March 2020)</i>				
<b>CHLOROFORM</b>				
a) Only in combination				
b) Maximum of 100 ml per prescription				
c) Only in aspirin and chloroform application.				
d) Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined.				
Chloroform BP .....	25.50	500 ml	✓ PSM	
<b>CODEINE PHOSPHATE – Safety medicine; prescriber may determine dispensing frequency</b>				
Powder – Only in combination .....	63.09 (90.09)	25 g		Douglas
Only in extemporaneously compounded codeine linctus.				
<b>COLLODION FLEXIBLE</b>				
Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined.				
Collodion flexible .....	19.30	100 ml	✓ PSM	
<b>COMPOUND HYDROXYBENZOATE – Only in combination</b>				
Only in extemporaneously compounded oral mixtures.				
Soln .....	30.00	100 ml	✓ <u>Midwest</u>	
<b>GLYCERIN WITH SODIUM SACCHARIN – Only in combination</b>				
Only in combination with Ora-Plus.				
Suspension .....	30.95	473 ml	✓ <u>Ora-Sweet SF</u>	
<b>GLYCERIN WITH SUCROSE – Only in combination</b>				
Only in combination with Ora-Plus.				
Suspension .....	30.95	473 ml	✓ <u>Ora-Sweet</u>	
<b>GLYCEROL</b>				
* Liquid – Only in combination .....	3.28	500 ml	✓ <u>healthE Glycerol BP</u>	
Only in extemporaneously compounded oral liquid preparations.				
<b>MAGNESIUM HYDROXIDE</b>				
Paste 29% .....	22.61	500 g	✓ PSM	
<i>(PSM Paste 29% to be delisted 1 July 2020)</i>				
<b>METHADONE HYDROCHLORIDE</b>				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency				
d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets).				
Powder .....	7.84	1 g	✓ AFT	
<b>METHYL HYDROXYBENZOATE</b>				
Powder .....	8.98	25 g	✓ <u>Midwest</u>	
<b>METHYLCELLULOSE</b>				
Powder .....	36.95	100 g	✓ <u>MidWest</u>	
Suspension – Only in combination .....	30.95	473 ml	✓ <u>Ora-Plus</u>	

# EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN – Only in combination				
Suspension.....	30.95	473 ml	✓	<b><u>Ora-Blend SF</u></b>
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE – Only in combination				
Suspension.....	30.95	473 ml	✓	<b><u>Ora-Blend</u></b>
PHENOBARBITONE SODIUM				
Powder – Only in combination.....	52.50	10 g	✓	<b>MidWest</b>
	325.00	100 g	✓	<b>MidWest</b>
Only in children up to 12 years				
PROPYLENE GLYCOL				
Only in extemporaneously compounded methyl hydroxybenzoate 10% solution.				
Liq.....	11.25	500 ml	✓	<b>Midwest</b>
SODIUM BICARBONATE				
Powder BP – Only in combination.....	10.05	500 g	✓	<b>Midwest</b>
	9.80			
	(29.50)			David Craig
a) Only in extemporaneously compounded omeprazole and lansoprazole suspension.				
b) Midwest to be Sole Supply on 1 January 2020				
<i>(David Craig Powder BP to be delisted 1 January 2020)</i>				
SYRUP (PHARMACEUTICAL GRADE) – Only in combination				
Only in extemporaneously compounded oral liquid preparations.				
Liq.....	14.95	500 ml	✓	<b>Midwest</b>
Midwest to be Sole Supply on 1 January 2020				
WATER				
Tap – Only in combination.....	0.00	1 ml	✓	<b>Tap water</b>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

Nutrient Modules

Carbohydrate

»SA1522 Special Authority for Subsidy

**Initial application — (Cystic fibrosis or kidney disease)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- Either:
- 1 cystic fibrosis; or
  - 2 chronic kidney disease.

**Initial application — (Indications other than cystic fibrosis or renal failure)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Any of the following:
- 1 cancer in children; or
  - 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
  - 3 faltering growth in an infant/child; or
  - 4 bronchopulmonary dysplasia; or
  - 5 premature and post premature infant; or
  - 6 inborn errors of metabolism; or
  - 7 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

**Renewal — (Cystic fibrosis or renal failure)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- Both:
- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

**Renewal — (Indications other than cystic fibrosis or renal failure)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT – Special Authority see SA1522 above – Hospital pharmacy [HP3]  
Powder .....5.29 400 g OP ✓ Polycal

Carbohydrate And Fat

»SA1376 Special Authority for Subsidy

**Initial application — (Cystic fibrosis)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

**Initial application — (Indications other than cystic fibrosis)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
  - 2.1 cancer in children; or
  - 2.2 faltering growth; or
  - 2.3 bronchopulmonary dysplasia; or
  - 2.4 premature and post premature infants.

**Renewal — (Cystic fibrosis)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

**Renewal — (Indications other than cystic fibrosis)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE AND FAT SUPPLEMENT – Special Authority see [SA1376 on the previous page](#) – Hospital pharmacy [HP3]

Powder (neutral) .....	60.31	400 g OP	✓ Duocal Super Soluble Powder
------------------------	-------	----------	-------------------------------

Fat

»SA1523 Special Authority for Subsidy

**Initial application — (Inborn errors of metabolism)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

**Initial application — (Indications other than inborn errors of metabolism)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome; or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

- 10 ascites; or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

**Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner.

Approvals valid for 3 years for applications meeting the following criteria:

- Both:
- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

**Renewal — (Indications other than inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

**FAT SUPPLEMENT** – Special Authority see [SA1523 on the previous page](#) – Hospital pharmacy [HP3]

Emulsion (neutral) .....	12.30	200 ml OP	✓ Calogen
	30.75	500 ml OP	✓ Calogen
Emulsion (strawberry).....	12.30	200 ml OP	✓ Calogen
Oil .....	30.00	500 ml OP	✓ MCT oil (Nutricia)
Oil, 250 ml .....	114.92	4 OP	✓ Liquigen

## Protein

### ►SA1524 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

**PROTEIN SUPPLEMENT** – Special Authority see [SA1524 above](#) – Hospital pharmacy [HP3]

Powder .....	7.90	225 g OP	✓ Protifar
	8.95	227 g OP	✓ Resource Beneprotein



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

## Oral Supplements/Complete Diet (Nasogastric/Gastrostomy Tube Feed)

### Respiratory Products

#### ►SA1094 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has CORD and hypercapnia, defined as a CO<sub>2</sub> value exceeding 55 mmHg.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CORD ORAL FEED 1.5KCAL/ML – Special Authority see SA1094 above – Hospital pharmacy [HP3]

Liquid .....	1.66	237 ml OP	✓ Pulmocare
--------------	------	-----------	-------------

### Diabetic Products

#### ►SA1095 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML – Special Authority see SA1095 above – Hospital pharmacy [HP3]

Liquid .....	7.50	1,000 ml OP	✓ Diasion RTH
			✓ Glucerna Select RTH

DIABETIC ORAL FEED 1KCAL/ML – Special Authority see SA1095 above – Hospital pharmacy [HP3]

Liquid (strawberry) .....	1.50	200 ml OP	✓ Diasip
Liquid (vanilla) .....	1.50	200 ml OP	✓ Diasip
	1.88	250 ml OP	✓ Glucerna Select
	1.78	237 ml OP	
	(2.10)		Resource Diabetic
	(2.10)		Sustagen Diabetic

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## Fat Modified Products

### »SA1525 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED – Special Authority see [SA1525 above](#) – Hospital pharmacy [HP3]

Powder ..... 60.48      400 g OP      ✓ **Monogen**

## Paediatric Products For Children Awaiting Liver Transplant

### »SA1098 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML – Special Authority see [SA1098 above](#) – Hospital pharmacy [HP3]

Powder (unflavoured) ..... 78.97      400 g OP      ✓ **Heparon Junior**

## Paediatric Products For Children With Chronic Renal Failure

### »SA1099 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
ENTERAL/ORAL FEED 1KCAL/ML – Special Authority see <a href="#">SA1099 on the previous page</a> – Hospital pharmacy [HP3]			
Liquid .....	54.00	400 g OP	✓ <b>Kindergen</b>

## Paediatric Products

### ► [SA1379](#) Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
  - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
  - 2.2 any condition causing malabsorption; or
  - 2.3 faltering growth in an infant/child; or
  - 2.4 increased nutritional requirements; or
  - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML – Special Authority see <a href="#">SA1379 above</a> – Hospital pharmacy [HP3]			
Liquid .....	6.00	500 ml OP	✓ <b>Nutrini Energy RTH</b>
PAEDIATRIC ENTERAL FEED 1KCAL/ML – Special Authority see <a href="#">SA1379 above</a> – Hospital pharmacy [HP3]			
Liquid .....	2.68	500 ml OP	✓ <b>Nutrini RTH</b> ✓ <b>Pediasure RTH</b>
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority see <a href="#">SA1379 above</a> – Hospital pharmacy [HP3]			
Liquid .....	6.00	500 ml OP	✓ <b>Nutrini Energy Multi Fibre</b>
PAEDIATRIC ORAL FEED 1.5KCAL/ML – Special Authority see <a href="#">SA1379 above</a> – Hospital pharmacy [HP3]			
Liquid (strawberry) .....	1.60	200 ml OP	✓ <b>Fortini</b>
Liquid (vanilla) .....	1.60	200 ml OP	✓ <b>Fortini</b>
PAEDIATRIC ORAL FEED 1KCAL/ML – Special Authority see <a href="#">SA1379 above</a> – Hospital pharmacy [HP3]			
Liquid (chocolate) .....	1.07	200 ml OP	✓ <b>Pediasure</b>
Liquid (strawberry) .....	1.07	200 ml OP	✓ <b>Pediasure</b>
Liquid (vanilla) .....	1.07	200 ml OP	✓ <b>Pediasure</b>
	1.34	250 ml OP	✓ <b>Pediasure</b>
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority see <a href="#">SA1379 above</a> – Hospital pharmacy [HP3]			
Liquid (unflavoured) .....	1.60	200 ml OP	✓ <b>Fortini Multi Fibre</b>
Liquid (chocolate) .....	1.60	200 ml OP	✓ <b>Fortini Multi Fibre</b>
Liquid (strawberry) .....	1.60	200 ml OP	✓ <b>Fortini Multi Fibre</b>
Liquid (vanilla) .....	1.60	200 ml OP	✓ <b>Fortini Multi Fibre</b>
PEPTIDE-BASED ORAL FEED – Special Authority see <a href="#">SA1379 above</a> – Hospital pharmacy [HP3]			
Powder .....	43.60	400 g OP	✓ <b>Peptamen Junior</b>

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

## Renal Products

### ►SA1101 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- Both:
- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML – Special Authority see [SA1101 above](#) – Hospital pharmacy [HP3]

Liquid.....	6.08	500 ml OP	✓ <b>Nepro HP RTH</b>
-------------	------	-----------	-----------------------

RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see [SA1101 above](#) – Hospital pharmacy [HP3]

Liquid.....	2.67	220 ml OP	✓ <b>Nepro HP</b> <b>(strawberry)</b> ✓ <b>Nepro HP (vanilla)</b>
-------------	------	-----------	---

RENAL ORAL FEED 2 KCAL/ML – Special Authority see [SA1101 above](#) – Hospital pharmacy [HP3]

Liquid.....	2.88	237 ml OP	
	(3.31)		NovaSource Renal
Liquid (apricot) 125 ml.....	11.52	4 OP	✓ <b>Renilon 7.5</b>
Liquid (caramel) 125 ml.....	11.52	4 OP	✓ <b>Renilon 7.5</b>

## Specialised And Elemental Products

### ►SA1377 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML – Special Authority see <a href="#">SA1377 on the previous page</a> – Hospital pharmacy [HP3]			
Liquid .....	18.06	1,000 ml OP	✓ <b>Vital</b>
ORAL ELEMENTAL FEED 0.8KCAL/ML – Special Authority see <a href="#">SA1377 on the previous page</a> – Hospital pharmacy [HP3]			
Liquid (grapefruit), 250 ml carton.....	171.00	18 OP	✓ <b>Elemental 028 Extra</b>
Liquid (pineapple & orange), 250 ml carton.....	171.00	18 OP	✓ <b>Elemental 028 Extra</b>
Liquid (summer fruits), 250 ml carton.....	171.00	18 OP	✓ <b>Elemental 028 Extra</b>
ORAL ELEMENTAL FEED 1KCAL/ML – Special Authority see <a href="#">SA1377 on the previous page</a> – Hospital pharmacy [HP3]			
Powder (unflavoured) .....	4.50	80 g OP	✓ <b>Vivonex TEN</b>
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML – Special Authority see <a href="#">SA1377 on the previous page</a> – Hospital pharmacy [HP3]			
Liquid .....	12.04	1,000 ml OP	✓ <b>Peptisorb</b>

## Paediatric Products For Children With Low Energy Requirements

### ► [SA1196](#) Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML – Special Authority see [SA1196 above](#) – Hospital pharmacy [HP3]

Liquid .....	4.00	500 ml OP	✓ <b>Nutrini Low Energy Multi Fibre</b>
--------------	------	-----------	---

## Standard Supplements

### ► [SA1554](#) Special Authority for Subsidy

**Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
  - 2.1 The patient has a condition causing malabsorption; or
  - 2.2 The patient has failure to thrive; or
  - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

**Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease)** only from a dietitian, relevant

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:  
All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

**Initial application — (Children - exclusive enteral nutrition for Crohn's disease)** only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

**Renewal — (Children - exclusive enteral nutrition for Crohn's disease)** only from a gastroenterologist, dietitian on the recommendation of a gastroenterologist or vocationally registered general practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

**Initial application — (Adults)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:  
Patient is Malnourished
  - 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
  - 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
  - 1.3 Patient has a BMI of less than 20 kg/m<sup>2</sup> and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:  
Patient has not responded to first-line dietary measures over a 4 week period by:
  - 2.1 Increasing their food intake frequency (eg snacks between meals); or
  - 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
  - 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

**Renewal — (Adults)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:  
Patient is Malnourished
  - 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
  - 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
  - 2.3 Patient has a BMI of less than 20 kg/m<sup>2</sup> and unintentional weight loss greater than 5% within the last 3-6 months.

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

**Initial application — (Short-term medical condition)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
    - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
    - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

**Renewal — (Short-term medical condition)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner.

Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
    - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
    - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

**Initial application — (Long-term medical condition)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube - refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm<sup>3</sup>); or
- 12 Chronic pancreatitis.

**Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube - refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML – Special Authority see [SA1554 on page 245](#) – Hospital pharmacy [HP3]

Liquid.....	7.00	1,000 ml OP	✓ <b>Nutrison Energy</b>
-------------	------	-------------	--------------------------

ENTERAL FEED 1KCAL/ML – Special Authority see [SA1554 on page 245](#) – Hospital pharmacy [HP3]

Liquid.....	1.24	250 ml OP	✓ <b>Isosource Standard</b>
	5.29	1,000 ml OP	✓ <b>Nutrison Standard RTH</b>
			✓ <b>Osmolite RTH</b>

ENTERAL FEED WITH FIBRE 0.83 KCAL/ML – Special Authority see [SA1554 on page 245](#) – Hospital pharmacy [HP3]

Liquid.....	5.29	1,000 ml OP	✓ <b>Nutrison 800 Complete Multi Fibre</b>
-------------	------	-------------	--

ENTERAL FEED WITH FIBRE 1 KCAL/ML – Special Authority see [SA1554 on page 245](#) – Hospital pharmacy [HP3]

Liquid.....	5.29	1,000 ml OP	✓ <b>Jevity RTH</b>
			✓ <b>Nutrison Multi Fibre</b>

ENTERAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority see [SA1554 on page 245](#) – Hospital pharmacy [HP3]

Liquid.....	1.75	250 ml OP	✓ <b>Ensure Plus HN</b>
	7.00	1,000 ml OP	✓ <b>Ensure Plus RTH</b>
			✓ <b>Jevity HiCal RTH</b>
			✓ <b>Nutrison Energy Multi Fibre</b>



	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
<b>ORAL FEED (POWDER) – Special Authority see <a href="#">SA1554 on page 245</a> – Hospital pharmacy [HP3]</b>			
Note: Higher subsidy for Sustagen Hospital Formula will only be reimbursed for patients with both a valid Special Authority number and an appropriately endorsed prescription.			
Powder (chocolate) – Higher subsidy of up to \$26.00 per 850 g			
with Endorsement.....	26.00	850 g OP	✓ <b>Ensure</b>  Sustagen Hospital Formula Active
	9.54	840 g OP	
	(26.00)		
Additional subsidy by endorsement is available for patients with fat malabsorption, fat intolerance or chyle leak. The prescription must be endorsed accordingly.			
Powder (vanilla) – Higher subsidy of up to \$26.00 per 850 g			
with Endorsement.....	8.54	857 g OP	✓ <b>Fortisip</b> ✓ <b>Ensure</b>  Sustagen Hospital Formula Active
	26.00	850 g OP	
	9.54	840 g OP	
	(26.00)		
Additional subsidy by endorsement is available for patients with fat malabsorption, fat intolerance or chyle leak. The prescription must be endorsed accordingly.			
<b>ORAL FEED 1.5KCAL/ML – Special Authority see <a href="#">SA1554 on page 245</a> – Hospital pharmacy [HP3]</b>			
Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease. The prescription must be endorsed accordingly.			
Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement.....	0.72	200 ml OP	Ensure Plus Fortisip
	(1.26)		
	(1.26)		
Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement.....	0.72	200 ml OP	Ensure Plus Fortisip
	(1.26)		
	(1.26)		
Liquid (fruit of the forest) – Higher subsidy of \$1.26 per 200 ml			
with Endorsement.....	0.72	200 ml OP	Ensure Plus
	(1.26)		
Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement.....	0.72	200 ml OP	Ensure Plus Fortisip
	(1.26)		
	(1.26)		
Liquid (vanilla) – Higher subsidy of up to \$1.33 per 237 ml with			
Endorsement.....	0.85	237 ml OP	Ensure Plus
	(1.33)		
	0.72	200 ml OP	Ensure Plus Fortisip
	(1.26)		
	(1.26)		

## SPECIAL FOODS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
ORAL FEED WITH FIBRE 1.5 KCAL/ML – Special Authority see <a href="#">SA1554 on page 245</a> – Hospital pharmacy [HP3]				
Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.				
Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with				
Endorsement .....	0.72 (1.26)	200 ml OP		Fortisip Multi Fibre
Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with				
Endorsement .....	0.72 (1.26)	200 ml OP		Fortisip Multi Fibre
Liquid (vanilla) – Higher subsidy of \$1.26 per 200 ml with				
Endorsement .....	0.72 (1.26)	200 ml OP		Fortisip Multi Fibre

## High Calorie Products

### ►SA1195 Special Authority for Subsidy

**Initial application — (Cystic fibrosis)** only from a dietitian, relevant specialist or vocationally registered general practitioner.

Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

**Initial application — (Indications other than cystic fibrosis)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 any condition causing malabsorption; or
  - 1.2 faltering growth in an infant/child; or
  - 1.3 increased nutritional requirements; or
  - 1.4 fluid restricted; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

**Renewal — (Cystic fibrosis)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

**Renewal — (Indications other than cystic fibrosis)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML – Special Authority see [SA1195 above](#) – Hospital pharmacy [HP3]

Liquid .....	5.50	500 ml OP	✓ <b>Nutrison Concentrated</b>
	11.00	1,000 ml OP	✓ <b>Two Cal HN RTH</b>

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
ORAL FEED 2 KCAL/ML – Special Authority see <a href="#">SA1195 on the previous page</a> – Hospital pharmacy [HP3]			
Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.			
Liquid (vanilla) – Higher subsidy of \$1.90 per 200 ml with			
Endorsement.....	0.96 (1.90)	200 ml OP	Two Cal HN

## Food Thickeners

### ►SA1106 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FOOD THICKENER – Special Authority see [SA1106 above](#) – Hospital pharmacy [HP3]

Powder .....	6.53	300 g OP	✓ Nutilis
	7.25	380 g OP	✓ Feed Thickener Karicare Aptamil

## Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by PHARMAC from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

### ►SA1729 Special Authority for Subsidy

**Initial application — (all patients)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

**Initial application — (paediatric patients diagnosed by ESPGHAN criteria)** only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX – Special Authority see [SA1729 above](#) – Hospital pharmacy [HP3]

Powder .....	2.81	1,000 g OP	
	(5.15)		Healthies Simple Baking Mix

GLUTEN FREE BREAD MIX – Special Authority see [SA1729 above](#) – Hospital pharmacy [HP3]

Powder .....	3.93	1,000 g OP	
	(7.32)		NZB Low Gluten Bread Mix
	3.51		
	(10.87)		Horleys Bread Mix

## SPECIAL FOODS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
GLUTEN FREE FLOUR – Special Authority see <a href="#">SA1729 on the previous page</a> – Hospital pharmacy [HP3]				
Powder .....	5.62	2,000 g OP		
	(18.10)			Horleys Flour
GLUTEN FREE PASTA – Special Authority see <a href="#">SA1729 on the previous page</a> – Hospital pharmacy [HP3]				
Buckwheat Spirals .....	2.00	250 g OP		
	(3.11)			Orgran
Corn and Vegetable Shells .....	2.00	250 g OP		
	(2.92)			Orgran
Corn and Vegetable Spirals .....	2.00	250 g OP		
	(2.92)			Orgran
Rice and Corn Lasagne Sheets .....	1.60	200 g OP		
	(3.82)			Orgran
Rice and Corn Macaroni .....	2.00	250 g OP		
	(2.92)			Orgran
Rice and Corn Penne .....	2.00	250 g OP		
	(2.92)			Orgran
Rice and Maize Pasta Spirals .....	2.00	250 g OP		
	(2.92)			Orgran
Rice and Millet Spirals .....	2.00	250 g OP		
	(3.11)			Orgran
Rice and corn spaghetti noodles .....	2.00	375 g OP		
	(2.92)			Orgran
Vegetable and Rice Spirals .....	2.00	250 g OP		
	(2.92)			Orgran
Italian long style spaghetti .....	2.00	220 g OP		
	(3.11)			Orgran

## Foods And Supplements For Inborn Errors Of Metabolism

### ► [SA1108](#) Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

### Supplements For Homocystinuria

AMINOACID FORMULA WITHOUT METHIONINE – Special Authority see [SA1108 above](#) – Hospital pharmacy [HP3]  
Powder .....461.94 500 g OP ✓ **XMET Maxamum**

### Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE – Special Authority see [SA1108 above](#) – Hospital pharmacy [HP3]  
Powder .....437.22 500 g OP ✓ **MSUD Maxamum**

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE – Special Authority see [SA1108 on the previous page](#) – Hospital pharmacy [HP3]

Tabs.....	99.00	75 OP	✓ Phlexy 10
Powder (chocolate) 36 g sachet.....	393.00	30	✓ PKU Anamix Junior Chocolate
Powder (unflavoured) 27.8 g sachets.....	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 36 g sachets.....	393.00	30	✓ PKU Anamix Junior
Powder (vanilla) 36 g sachet.....	393.00	30	✓ PKU Anamix Junior Vanilla
Infant formula.....	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange).....	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured).....	320.00	500 g OP	✓ XP Maxamum
Liquid (berry).....	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange).....	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (unflavoured).....	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton.....	540.00	18 OP	✓ Easiphen Liquid
Liquid (juicy tropical) 125 ml.....	936.00	30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g.....	1,123.20	36 OP	✓ PKU Lophlex Sensation 20
Liquid (juicy berries) 62.5 ml.....	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml.....	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml.....	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml.....	936.00	30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml.....	936.00	30 OP	✓ PKU Lophlex LQ 20

## Foods

LOW PROTEIN BAKING MIX – Special Authority see [SA1108 on the previous page](#) – Hospital pharmacy [HP3]

Powder.....	8.22	500 g OP	✓ Loprofin Mix
-------------	------	----------	----------------

LOW PROTEIN PASTA – Special Authority see [SA1108 on the previous page](#) – Hospital pharmacy [HP3]

Animal shapes.....	11.91	500 g OP	✓ Loprofin
Lasagne.....	5.95	250 g OP	✓ Loprofin
Low protein rice pasta.....	11.91	500 g OP	✓ Loprofin
Macaroni.....	5.95	250 g OP	✓ Loprofin
Penne.....	11.91	500 g OP	✓ Loprofin
Spaghetti.....	11.91	500 g OP	✓ Loprofin
Spirals.....	11.91	500 g OP	✓ Loprofin

## Infant Formulae

### For Williams Syndrome

#### »SA1110 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA – Special Authority see [SA1110 above](#) – Hospital pharmacy [HP3]

Powder .....	44.40	400 g OP	✓ Locasol
--------------	-------	----------	-----------

### Gastrointestinal and Other Malabsorptive Problems

AMINO ACID FORMULA – Special Authority see [SA1219 below](#) – Hospital pharmacy [HP3]

Powder .....	43.60	400 g OP	✓ Alfamino Junior
Powder (unflavoured) .....	53.00	400 g OP	✓ Elecare
			✓ Elecare LCP
			✓ Neocate Gold
			✓ Neocate Junior Unflavoured
			✓ Neocate SYNEO
Powder (vanilla) .....	53.00	400 g OP	✓ Elecare
			✓ Neocate Junior Vanilla

#### »SA1219 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Extensively hydrolysed formula has been reasonably trialled and is inappropriate due to documented severe intolerance or allergy or malabsorption; or
- 2 History of anaphylaxis to cows milk protein formula or dairy products; or
- 3 Eosinophilic oesophagitis.

Note: A reasonable trial is defined as a 2-4 week trial.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
EXTENSIVELY HYDROLYSED FORMULA – Special Authority see <a href="#">SA1557 below</a> – Hospital pharmacy [HP3]			
Powder .....	15.21	450 g OP	✓ Aptamil Gold+ Pepti Junior

► **SA1557** Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
  - 1.2 Either:
    - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
    - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malabsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
  - 11.1 For step down from Amino Acid Formula; and
  - 11.2 The infant is currently receiving funded amino acid formula; and
  - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
  - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

## Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML – Special Authority see [SA1698 below](#) – Hospital pharmacy [HP3]

Liquid .....	2.35	125 ml OP	✓ Infatrini
--------------	------	-----------	-------------

► **SA1698** Special Authority for Subsidy

**Initial application** only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula;

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

and

- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

**Renewal** only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

## Ketogenic Diet

### ►SA1197 Special Authority for Subsidy

**Initial application** only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

**Renewal** only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA – Special Authority see [SA1197 above](#) – Retail pharmacy

Powder (unflavoured) .....	35.50	300 g OP	✓ <b>KetoCal 4:1</b>
			✓ <b>Ketocal 3:1</b>
Powder (vanilla).....	35.50	300 g OP	✓ <b>KetoCal 4:1</b>



Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

## Vaccinations

### ADULT DIPHTHERIA AND TETANUS VACCINE – [Xpharm]

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid in 0.5 ml.....0.00 5 ✓ **ADT Booster**

Any of the following:

- 1) For vaccination of patients aged 45 and 65 years old; or
- 2) For vaccination of previously unimmunised or partially immunised patients; or
- 3) For revaccination following immunosuppression; or
- 4) For boosting of patients with tetanus-prone wounds; or
- 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

### BACILLUS CALMETTE-GUERIN VACCINE – [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000

Note a list of countries with high rates of TB are available at [www.health.govt.nz/tuberculosis](http://www.health.govt.nz/tuberculosis) (search for downloads) or [www.bcgatlas.org/index.php](http://www.bcgatlas.org/index.php).

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),  
Danish strain 1331, live attenuated, vial with diluent.....0.00 10 ✓ **BCG Vaccine**

### DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE – [Xpharm]

Funded for any of the following criteria:

- 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
- 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
- 3) A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
- 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens.

Notes: Tdap is not registered for patients aged less than 10 years. Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg  
pertussis toxoid, 8 mcg pertussis filamentous  
haemagglutinin and 2.5 mcg pertactin in 0.5 ml syringe .....0.00 10  
1 ✓ **Boostrix**  
✓ **Boostrix**

# NATIONAL IMMUNISATION SCHEDULE

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
<b>DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE – [Xpharm]</b>				
Funded for any of the following:				
1) A single dose for children up to the age of 7 who have completed primary immunisation; or				
2) A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or				
3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or				
4) Five doses will be funded for children requiring solid organ transplantation.				
Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.				
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units poliomyelitis virus in 0.5ml syringe .....	0.00	10	✓	<b>Infanrix IPV</b>
<b>DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm]</b>				
Funded for patients meeting any of the following criteria:				
1) Up to four doses for children up to and under the age of 10 for primary immunisation; or				
2) An additional four doses (as appropriate) are funded for (re-)immunisation for children up to and under the age of 10 who are patients post haematopoietic stem cell transplantation, or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or				
3) Up to five doses for children up to and under the age of 10 receiving solid organ transplantation.				
Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.				
Inj 30IU diphtheriatoxoid with 40IU tetanustoxoid, 25mcg pertussistoxoid, 25mcg pertussisfilamentoushaemagglutinin, 8 mcgpertactin, 80 D-AgUpoliovirus, 10mcghepatitisBsurfaceantigen in 0.5ml syringe .....	0.00	10	✓	<b>Infanrix-hexa</b>
<b>HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm]</b>				
One dose for patients meeting any of the following:				
1) For primary vaccination in children; or				
2) An additional dose (as appropriate) is funded for (re-)immunisation for patients post haematopoietic stem cell transplantation, or chemotherapy; functional asplenic; pre or post splenectomy; pre- or post solid organ transplant, pre- or post cochlear implants, renal dialysis and other severely immunosuppressive regimens; or				
3) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.				
Haemophilus Influenzae type B polysaccharide 10 mcg conjugated to tetanus toxoid as carrier protein 20-40 mcg; prefilled syringe plus vial 0.5 ml.....	0.00	1	✓	<b>Hiberix</b>
<b>HEPATITIS A VACCINE – [Xpharm]</b>				
Funded for patients meeting any of the following criteria:				
1) Two vaccinations for use in transplant patients; or				
2) Two vaccinations for use in children with chronic liver disease; or				
3) One dose of vaccine for close contacts of known hepatitis A cases.				
Inj 1440 ELISA units in 1 ml syringe.....	0.00	1	✓	<b>Havrix</b>
Inj 720 ELISA units in 0.5 ml syringe.....	0.00	1	✓	<b>Havrix Junior</b>

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
<b>HEPATITIS B RECOMBINANT VACCINE – [Xpharm]</b>			
Inj 5 mcg per 0.5 ml vial.....	0.00	1	✓ <b>HBvaxPRO</b>
Funded for patients meeting any of the following criteria:			
1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or			
2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or			
3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or			
4) for HIV positive patients; or			
5) for hepatitis C positive patients; or			
6) for patients following non-consensual sexual intercourse; or			
7) for patients following immunosuppression; or			
8) for solid organ transplant patients; or			
9) for post-haematopoietic stem cell transplant (HSCT) patients; or			
10) following needle stick injury.			
Inj 10 mcg per 1 ml vial.....	0.00	1	✓ <b>HBvaxPRO</b>
Funded for patients meeting any of the following criteria:			
1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or			
2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or			
3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or			
4) for HIV positive patients; or			
5) for hepatitis C positive patients; or			
6) for patients following non-consensual sexual intercourse; or			
7) for patients following immunosuppression; or			
8) for solid organ transplant patients; or			
9) for post-haematopoietic stem cell transplant (HSCT) patients; or			
10) following needle stick injury.			
Inj 20 mcg per 1 ml prefilled syringe.....	0.00	1	✓ <b>Engerix-B</b>
Funded for patients meeting any of the following criteria:			
1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or			
2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or			
3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or			
4) for HIV positive patients; or			
5) for hepatitis C positive patients; or			
6) for patients following non-consensual sexual intercourse; or			
7) for patients following immunosuppression; or			
8) for solid organ transplant patients; or			
9) for post-haematopoietic stem cell transplant (HSCT) patients; or			
10) following needle stick injury; or			
11) for dialysis patients; or			
12) for liver or kidney transplant patients.			
Inj 40 mcg per 1 ml vial.....	0.00	1	✓ <b>HBvaxPRO</b>
Funded for any of the following criteria:			
1) for dialysis patients; or			
2) for liver or kidney transplant patient.			

# NATIONAL IMMUNISATION SCHEDULE

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV] – [Xpharm]			
Any of the following:			
1) Maximum of two doses for children aged 14 years and under; or			
2) Maximum of three doses for patients meeting any of the following criteria:			
1) People aged 15 to 26 years inclusive; or			
2) Either:			
People aged 9 to 26 years inclusive			
1) Confirmed HIV infection; or			
2) Transplant (including stem cell) patients: or			
3) Maximum of four doses for people aged 9 to 26 years inclusive post chemotherapy			
Inj 270 mcg in 0.5 ml syringe.....	0.00	10	✓ <b>Gardasil 9</b>

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
<b>INFLUENZA VACCINE</b>			
Inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine) – [Xpharm].....	9.00	1	✓ <b>Fluarix Tetra</b>
<b>A) INFLUENZA VACCINE – child aged 6 months to 35 months</b> is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by PHARMAC: <ul style="list-style-type: none"> <li>i) have any of the following cardiovascular diseases               <ul style="list-style-type: none"> <li>a) ischaemic heart disease, or</li> <li>b) congestive heart failure, or</li> <li>c) rheumatic heart disease, or</li> <li>d) congenital heart disease, or</li> <li>e) cerebo-vascular disease; or</li> </ul> </li> <li>ii) have either of the following chronic respiratory diseases:               <ul style="list-style-type: none"> <li>a) asthma, if on a regular preventative therapy, or</li> <li>b) other chronic respiratory disease with impaired lung function; or</li> </ul> </li> <li>iii) have diabetes; or</li> <li>iv) have chronic renal disease; or</li> <li>v) have any cancer, excluding basal and squamous skin cancers if not invasive; or</li> <li>vi) have any of the following other conditions:               <ul style="list-style-type: none"> <li>a) autoimmune disease, or</li> <li>b) immune suppression or immune deficiency, or</li> <li>c) HIV, or</li> <li>d) transplant recipients, or</li> <li>e) neuromuscular and CNS diseases/disorders, or</li> <li>f) haemoglobinopathies, or</li> <li>g) on long term aspirin, or</li> <li>h) have a cochlear implant, or</li> <li>i) errors of metabolism at risk of major metabolic decompensation, or</li> <li>j) pre and post splenectomy, or</li> <li>k) down syndrome, or</li> </ul> </li> <li>vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;</li> </ul> Unless meeting the criteria set out above, the following conditions are excluded from funding: <ul style="list-style-type: none"> <li>a) asthma not requiring regular preventative therapy,</li> <li>b) hypertension and/or dyslipidaemia without evidence of end-organ disease.</li> </ul>			
<b>B) INFLUENZA VACCINE – pregnant women</b> <ul style="list-style-type: none"> <li>a) are pregnant</li> </ul>			
<b>C) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.</b>			
Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine).....	45.00 90.00	5 10	✓ <b>FluQuadri</b> ✓ <b>Afluria Quad</b> ✓ <b>Influvac Tetra</b>

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
a) Only on a prescription			
b) No patient co-payment payable			
c)			
<b>A) INFLUENZA VACCINE – people 3 years and over</b>			
is available each year for patients aged 3 years and over who meet the following criteria, as set by PHARMAC:			
a) all people 65 years of age and over; or			
b) people under 65 years of age who:			
i) have any of the following cardiovascular diseases:			
a) ischaemic heart disease, or			
b) congestive heart failure, or			
c) rheumatic heart disease, or			
d) congenital heart disease, or			
e) cerebo-vascular disease; or			
ii) have either of the following chronic respiratory diseases:			
a) asthma, if on a regular preventative therapy, or			
b) other chronic respiratory disease with impaired lung function; or			
iii) have diabetes; or			
iv) have chronic renal disease; or			
v) have any cancer, excluding basal and squamous skin cancers if not invasive; or			
vi) have any of the following other conditions:			
a) autoimmune disease, or			
b) immune suppression or immune deficiency, or			
c) HIV, or			
d) transplant recipients, or			
e) neuromuscular and CNS diseases/disorders, or			
f) haemoglobinopathies, or			
g) are children on long term aspirin, or			
h) have a cochlear implant, or			
i) errors of metabolism at risk of major metabolic decompensation, or			
j) pre and post splenectomy, or			
k) down syndrome, or			
vii) are pregnant; or			
c) children aged four years or less (but over three years) who have been hospitalised for respiratory illness or have a history of significant respiratory illness;			
Unless meeting the criteria set out above, the following conditions are excluded from funding:			
a) asthma not requiring regular preventative therapy,			
b) hypertension and/or dyslipidaemia without evidence of end-organ disease.			
B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.			
C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.			
<b>MEASLES, MUMPS AND RUBELLA VACCINE – [Xpharm]</b>			
A maximum of two doses for any patient meeting the following criteria:			
1) For primary vaccination in children; or			
2) For revaccination following immunosuppression; or			
3) For any individual susceptible to measles, mumps or rubella; or			
4) A maximum of three doses for children who have had their first dose prior to 12 months.			
Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.			
Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,			
Rubella virus 1,000 CCID50; prefilled syringe/ampoule of			
diluent 0.5 ml .....	0.00	10	✓ <b>Priorix</b>

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
<b>MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE – [Xpharm]</b>			
Any of the following:			
1) Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or			
2) One dose for close contacts of meningococcal cases; or			
3) A maximum of two doses for bone marrow transplant patients; or			
4) A maximum of two doses for patients following immunosuppression*.			
Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.			
*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.			
Inj 4 mcg of each meningococcal polysaccharide conjugated to a total of approximately 48 mcg of diphtheria toxoid carrier per 0.5 ml vial .....	0.00	1	✓ <b>Menactra</b>
<b>MENINGOCOCCAL C CONJUGATE VACCINE – [Xpharm]</b>			
Any of the following:			
1) Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or			
2) One dose for close contacts of meningococcal cases; or			
3) A maximum of two doses for bone marrow transplant patients; or			
4) A maximum of two doses for patients following immunosuppression*.			
Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.			
*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.			
Inj 10 mcg in 0.5 ml syringe .....	0.00	1	✓ <b>Neisvac-C</b>
<b>PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE – [Xpharm]</b>			
Either:			
1) A primary course of four doses for previously unvaccinated individuals up to the age of 59 months inclusive; or			
2) Up to three doses as appropriate to complete the primary course of immunisation for individuals under the age of 59 months who have received one to three doses of PCV13.			
Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes			
Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B, 7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml prefilled syringe .....	0.00	10	✓ <b>Synflorix</b>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

## PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE – [Xpharm]

Any of the following:

- 1) One dose is funded for high risk children (over the age of 17 months and under 18 years) who have previously received four doses of PCV10; or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
  - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
  - b) with primary immune deficiencies; or
  - c) with HIV infection; or
  - d) with renal failure, or nephrotic syndrome; or
  - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
  - f) with cochlear implants or intracranial shunts; or
  - g) with cerebrospinal fluid leaks; or
  - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
  - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
  - j) pre term infants, born before 28 weeks gestation; or
  - k) with cardiac disease, with cyanosis or failure; or
  - l) with diabetes; or
  - m) with Down syndrome; or
  - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- 4) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml

syringe.....	0.00	10	✓ Prevenar 13
		1	✓ Prevenar 13



	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
<b>PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE – [Xpharm]</b>			
Either:			
1) Up to three doses (as appropriate) for patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy or with functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or 2) All of the following: <ul style="list-style-type: none"> <li>a) Patient is a child under 18 years for (re-)immunisation; and</li> <li>b) Treatment is for a maximum of two doses; and</li> <li>c) Any of the following:               <ul style="list-style-type: none"> <li>i) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or</li> <li>ii) with primary immune deficiencies; or</li> <li>iii) with HIV infection; or</li> <li>iv) with renal failure, or nephrotic syndrome; or</li> <li>v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or</li> <li>vi) with cochlear implants or intracranial shunts; or</li> <li>vii) with cerebrospinal fluid leaks; or</li> <li>viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or</li> <li>ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or</li> <li>x) pre term infants, born before 28 weeks gestation; or</li> <li>xi) with cardiac disease, with cyanosis or failure; or</li> <li>xii) with diabetes; or</li> <li>xiii) with Down syndrome; or</li> <li>xiv) who are pre- or post-splenectomy, or with functional asplenia.</li> </ul> </li> </ul>			
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype) .....	0.00	1	✓ <b><u>Pneumovax 23</u></b>
<b>POLIOMYELITIS VACCINE – [Xpharm]</b>			
Up to three doses for patients meeting either of the following:			
1) For partially vaccinated or previously unvaccinated individuals; or 2) For revaccination following immunosuppression.			
Note: Please refer to the Immunisation Handbook for appropriate schedule for catch-up programmes.			
Inj 80D antigen units in 0.5 ml syringe.....	0.00	1	✓ <b><u>IPOL</u></b>
<b>ROTAVIRUS ORAL VACCINE – [Xpharm]</b>			
Maximum of two doses for patients meeting the following:			
1) first dose to be administered in infants aged under 14 weeks of age; and 2) no vaccination being administered to children aged 24 weeks or over.			
Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator.....	0.00	10	✓ <b><u>Rotarix</u></b>

# NATIONAL IMMUNISATION SCHEDULE

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

## VARICELLA VACCINE [CHICKENPOX VACCINE] – [Xpharm]

Either:

- 1) Maximum of one dose for primary vaccination for either:
  - a) Any infant born on or after 1 April 2016; or
  - b) For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox), or
- 2) Maximum of two doses for any of the following:
  - a) Any of the following for non-immune patients:
    - i) with chronic liver disease who may in future be candidates for transplantation; or
    - ii) with deteriorating renal function before transplantation; or
    - iii) prior to solid organ transplant; or
    - iv) prior to any elective immunosuppression\*, or
    - v) for post exposure prophylaxis who are immune competent inpatients.; or
  - b) For patients at least 2 years after bone marrow transplantation, on advice of their specialist, or
  - c) For patients at least 6 months after completion of chemotherapy, on advice of their specialist, or
  - d) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist, or
  - e) For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella, or
  - f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella, or
  - g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.

\* immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

Inj 2000 PFU prefilled syringe plus vial .....	0.00	1	✓ <b>Varilrix</b>
		10	✓ <b>Varilrix</b>

## VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED VACCINE [SHINGLES VACCINE] – [Xpharm]

Funded for patients meeting either of the following criteria:

- 1) One dose for all people aged 65 years; or
- 2) One dose for all people aged between 66 and 80 years inclusive from 1 April 2018 and 31 March 2020.

Inj 19,400 PFU prefilled syringe plus vial .....	0.00	1	✓ <b>Zostavax</b>
		10	✓ <b>Zostavax</b>

## Diagnostic Agents

### TUBERCULIN PPD [MANTOUX] TEST – [Xpharm]

Inj 5 TU per 0.1 ml, 1 ml vial .....	0.00	1	✓ <b>Tubersol</b>
--------------------------------------	------	---	-------------------

<b>- Symbols -</b>		
3TC.....	106	
<b>- A -</b>		
A-Scabies.....	67	
Abacavir sulphate.....	106	
Abacavir sulphate with lamivudine.....	106	
Abiraterone acetate.....	171	
Acarbose.....	11	
Acarbose Mylan.....	11	
Accarb.....	11	
Accuretic 10.....	46	
Accuretic 20.....	46	
Acetazolamide.....	230	
Acetic acid with 1, 2- propanediol diacetate and benzethonium.....	228	
Acetic acid with hydroxyquinoline and ricinoleic acid.....	75	
Acetylcysteine.....	233	
Aci-Jel.....	75	
Aciclovir		
Infection.....	102	
Sensory.....	228	
Acidex.....	6	
Acipimox.....	52	
Acitretin.....	67	
Aclasta.....	114	
Aclin.....	110	
Actemra.....	211	
Actinomycin D.....	161	
Actrapid.....	10	
Actrapid Penfill.....	10	
Acupan.....	121	
Adalat 10.....	50	
Adalat Oros.....	50	
Adalimumab.....	180	
Adapalene.....	59	
Adefin.....	50	
Adefin XL.....	50	
Adefovir dipivoxil.....	101	
Adenuric.....	117	
ADR Cartridge 1.8.....	23	
Adrenaline.....	54	
ADT Booster.....	257	
Adult diphtheria and tetanus vaccine.....	257	
Advantan.....	62	
Advate.....	38	
Adynovate.....	39	
Afinitor.....	218	
Aflibercept.....	190	
Afluria Quad.....	261	
AFT Carbimazole.....	83	
AFT-Pyrazinamide.....	100	
Agents Affecting the Renin-Angiotensin System.....	45	
Agents for Parkinsonism and Related Disorders.....	118	
Agents Used in the Treatment of Poisonings.....	233	
Agrylin.....	160	
Alanase.....	227	
Albendazole.....	89	
Albey.....	220	
Albustix.....	77	
Alidurazyme.....	28	
Alendronate sodium.....	111	
Alendronate sodium with colecalciferol.....	111	
Alfacalcidol.....	32	
Alfamino Junior.....	254	
Alginic acid.....	6	
Alglucosidase alfa.....	26	
Alkeran.....	156	
Allersoothe.....	221	
Alimercap.....	159	
Allopurinol.....	116	
Alpha-Adrenoceptor Blockers.....	45	
Alpha-Keri Lotion.....	64	
Alphamox 125.....	92	
Alphamox 250.....	92	
Alprolix.....	36	
Alu-Tab.....	6	
Aluminium hydroxide.....	6	
Amantadine hydrochloride.....	118	
Ambrisentan.....	55	
Amiloride hydrochloride.....	51	
Amiloride hydrochloride with furosemide.....	51	
Amiloride hydrochloride with hydrochlorothiazide.....	52	
Aminophylline.....	226	
Amiodarone hydrochloride.....	47	
Amisulpride.....	131	
Amtripyline.....	124	
Amlodipine.....	49	
Amorolfine.....	60	
Amoxicillin.....	92	
Amoxicillin with clavulanic acid.....	92	
Amphotericin B.....	31	
Amsacrine.....	160	
AmsaLyo.....	160	
Amsidine.....	160	
Amzoate.....	28	
Anaesthetics.....	120	
Anagrelide hydrochloride.....	160	
Analgesics.....	121	
Anastrozole.....	173	
Andriol Testocaps.....	80	
Androderm.....	80	
Anoro Ellipta.....	224	
Antabuse.....	152	
Antacids and Antiflatulents.....	6	
Anten.....	124	
Anthelmintics.....	89	
Antiacne Preparations.....	59	
Antiallergy Preparations.....	220	
Antianaemics.....	35	
Antiandrogen Oral Contraceptives.....	75	
Antiarrhythmics.....	47	
Antibacterials.....	89	
Antibacterials Topical.....	59	
Anticholinergic Agents.....	223	
Anticholinesterases.....	110	
Antidepressants.....	124	
Antidiarrhoeals.....	6	
Antiepilepsy Drugs.....	126	
Antifibrinolytics, Haemostatics and Local Sclerosants.....	36	
Antifibrotics.....	224	
Antifungals.....	96	
Antifungals Topical.....	60	
Antihistamines.....	221	
Antihypotensives.....	48	
Antimalarials.....	99	
Antimigraine Preparations.....	129	
Antinausea and Vertigo Agents.....	130	
Antiparasitics.....	99	
Antipruritic Preparations.....	61	
Antipsychotics.....	131	
Antiretrovirals.....	105	
Antirheumatoid Agents.....	111	
Antispasmodics and Other Agents Altering Gut Motility.....	8	
Antithrombotic Agents.....	39	
Antithymocyte globulin (equine).....	180	
Antitrichomonal Agents.....	99	
Antituberculosics and Antileprotics.....	99	
Antulcerants.....	8	
Antivirals.....	101	
Anxiolytics.....	134	
Anzatax.....	163	
Apidra.....	10	
Apidra SoloStar.....	10	
Apo-Amlodipine.....	49	
Apo-Amoxi.....	92	
Apo-Azithromycin.....	90	
Apo-Bromocriptine.....	118	

Apo-Ciclopirox .....	60	Arrow-Losartan & Hydrochlorothiazide.....	46	Bacillus Calmette-Guerin (BCG) vaccine .....	180
Apo-Cilazapril .....	45	Arrow-Morphine LA.....	123	Bacillus Calmette-Guerin vaccine .....	257
Apo-Cilazapril/ Hydrochlorothiazide.....	46	Arrow-Norfloxacin.....	109	Baclofen.....	117
Apo-Clarithromycin Alimentary .....	8	Arrow-Ornidazole.....	99	Bactroban.....	60
Infection .....	91	Arrow-Quinapril 10.....	45	Barrier Creams and Emollients.....	64
Apo-Clomipramine.....	124	Arrow-Quinapril 20.....	45	BCG Vaccine.....	257
Apo-Diclo SR.....	110	Arrow-Quinapril 5.....	45	Beclazone 100.....	221
Apo-Diltiazem CD.....	50	Arrow-Roxithromycin .....	91	Beclazone 250.....	221
Apo-Doxazosin .....	45	Arrow-Sertraline.....	126	Beclazone 50.....	221
Apo-Folic Acid.....	36	Arrow-Timolol.....	230	Beclomethasone dipropionate.....	221, 227
Apo-Furosemide.....	51	Arrow-Tolterodine.....	77	Bee venom allergy treatment.....	220
Apo-Gabapentin .....	127	Arrow-Topiramate.....	128	Bendamustine hydrochloride.....	155
Apo-Leflunomide.....	111	Arrow-Tramadol.....	124	Bendrofluazide.....	52
Apo-Megestrol.....	172	Arsenic trioxide.....	160	Bendroflumethiazide [Bendrofluazide].....	52
Apo-Metoprolol.....	49	Asacol.....	7	BeneFIX.....	38
Apo-Mirtazapine.....	126	Asamax.....	7	Benzathine benzylpenicillin .....	92
Apo-Montelukast.....	226	Ascorbic acid.....	31	Benzatropine mesylate.....	119
Apo-Nadolol.....	49	Aspen Adrenaline .....	54	Benzbromaron AL 100.....	116
Apo-Nicotinic Acid.....	52	Aspirin		Benzoin.....	236
Apo-Ondansetron .....	131	Blood.....	39	Benzotrop.....	119
Apo-Oxybutynin.....	76	Nervous.....	121	Benzydamine hydrochloride.....	30
Apo-Paroxetine.....	125	Asthalin.....	223	Benzylpenicillin sodium [Penicillin G].....	92
Apo-Perindopril.....	45	Atazanavir sulphate.....	107	Beta Cream.....	62
Apo-Pindolol.....	49	Atenolol.....	48	Beta Ointment.....	62
Apo-Pravastatin.....	53	Atenolol AFT.....	48	Beta Scalp.....	68
Apo-Prazosin.....	45	ATGAM.....	180	Beta-Adrenoceptor Agonists.....	223
Apo-Prednisone.....	80	Ativan.....	135	Beta-Adrenoceptor Blockers.....	48
Apo-Primidone.....	128	Atomoxetine.....	147	Betadine.....	65
Apo-Propranolol.....	49	Atorvastatin.....	53	Betadine Skin Prep.....	65
Apo-Pyridoxine.....	31	Atropine sulphate		Betaferon.....	144
Apo-Ropinirole.....	118	Cardiovascular.....	47	Betahistine dihydrochloride .....	130
Apo-Selegiline S29.....	118	Sensory.....	231	Betaine.....	27
Apo-Sumatriptan.....	129	Atropt.....	231	Betaloc CR.....	49
Apo-Temozolomide.....	164	Atrovent.....	223	Betamethasone dipropionate.....	62
Apo-Terazosin.....	45	AU Synacthen.....	80	Betamethasone dipropionate with calcipotriol.....	67
Apo-Timol.....	49	Aubagio.....	139	Betamethasone sodium phosphate with betamethasone acetate.....	79
Apomorphine hydrochloride.....	118	Augmentin.....	92	Betamethasone valerate.....	62, 68
Aprepitant.....	130	Aurorix.....	125	Betamethasone valerate with clicloquinol.....	63
Apresoline.....	54	AutoSoft 30.....	20	Betamethasone valerate with sodium fusidate [fusidic acid].....	63
Aptamil Gold+ Pepti Junior.....	255	AutoSoft 90.....	22	Betaxolol.....	230
Aqueous cream.....	64	Avelox.....	94	Betnovate.....	62
Aratac.....	47	Avonex.....	142	Betnovate-C.....	63
Aripiprazole.....	131	Avonex Pen.....	142	Betoptic.....	230
Aripiprazole Sandoz.....	131	Azaciditidine.....	157	Betoptic S.....	230
Aristocort.....	63	Azaciditidine Dr Reddy's .....	157	Bezafibrate.....	52
Arrow - Clopid.....	39	Azamun.....	174	Bezalip.....	52
Arrow-Amitriptyline.....	124	Azathioprine.....	174		
Arrow-Bendrofluazide.....	52	Azithromycin.....	90		
Arrow-Brimonidine.....	231	Azol.....	88		
Arrow-Calcium.....	33	Azopt.....	230		
Arrow-Diazepam.....	134	AZT.....	106-107		
Arrow-Doxorubicin.....	162				
Arrow-Fluoxetine.....	125	<b>- B -</b>			
Arrow-Lamotrigine.....	127	B-D Micro-Fine.....	13		
		B-D Ultra Fine.....	14		
		B-D Ultra Fine II.....	14		

Bezalip Retard.....	52	Budesonide		Carmellose sodium with gelatin and pectin.....	30
Bicalutamide.....	172	Alimentary.....	6	Carmustine.....	156
Bicillin LA.....	92	Respiratory.....	221, 227	Carvedilol.....	48
BiCNU.....	156	Budesonide with eformoterol.....	222	Carvedilol Sandoz.....	48
Bicnu Heritage.....	156	Bumetanide.....	51	Catapres.....	50
Bile and Liver Therapy.....	9	Buprenorphine with naloxone.....	151	Cathejell.....	120
Biltricide.....	89	Bupropion hydrochloride.....	152	CeeNU.....	156
Bimatoprost.....	231	Burinex.....	51	Cefaclor monohydrate.....	89
Bimatoprost Multichem.....	231	Buscopan.....	8	Cefalexin.....	89
Binarex.....	172	Buspirone hydrochloride.....	134	Cefalexin Sandoz.....	89
Binocrit.....	36	Busulfan.....	156	Cefazolin.....	89
Biodone.....	122			Ceftriaxone.....	90
Biodone Extra Forte.....	122	- C -		Ceftriaxone-AFT.....	90
Biodone Forte.....	122	Cabergoline.....	88	Cefuroxime axetil.....	90
Bisacodyl.....	26	Cafergot.....	129	Celebrex.....	111
Bisoprolol fumarate.....	48	Cafergot S29.....	129	Celecoxib.....	111
BK Lotion.....	64	Caffeine citrate.....	227	Celecoxib Pfizer.....	111
Bleomycin sulphate.....	161	Calamine.....	61	Celestone Chronodose.....	79
Blood Colony-stimulating Factors.....	42	Calcipotriol.....	67	Celiprolol.....	48
Blood glucose diagnostic test meter.....	12	Calcitonin.....	78	Cellcept.....	174
Blood glucose diagnostic test strip.....	13	Calcitriol.....	32	Celol.....	48
Blood glucose test strips (visually impaired).....	13	Calcitriol-AFT.....	32	Centrally-Acting Agents.....	50
Blood Ketone Diagnostic Test Strip.....	11	Calcium carbonate.....	6, 33	Cephalexin ABM.....	89
Bonjela.....	30	Calcium Channel Blockers.....	49	Cetirizine hydrochloride.....	221
Boostrix.....	257	Calcium Disodium Versenate.....	234	Cetomacrogol.....	64
Bortezomib.....	161	Calcium folinate.....	158	Cetomacrogol with glycerol.....	64
Bosentan.....	55	Calcium Folate Ebewe.....	158	Cetuximab.....	191
Bosentan Dr Reddy's.....	55	Calcium Folate Sandoz.....	158	Charcoal.....	233
Bosvate.....	48	Calcium gluconate.....	33	Chemotherapeutic Agents.....	155
Bplex.....	31	Calcium Homeostasis.....	78	Chickenpox vaccine.....	266
Breo Ellipta.....	222	Calcium polystyrene sulphonate.....	44	Chlorafast.....	228
Brevinor 1/21.....	73	Calcium Resonium.....	44	Chlorambucil.....	156
Brevinor 1/28.....	73	Calcium Sandoz.....	33	Chloramphenicol.....	228
Brevinor 21.....	73	Calogen.....	240	Chlorhexidine gluconate	
Bricanyl Turbuhaler.....	223	Candesartan cilexetil.....	46	Alimentary.....	30
Brilinta.....	39	Candestar.....	46	Dermatological.....	63
Brimonidine tartrate.....	231	Canesten.....	60	Chloroform.....	236
Brimonidine tartrate with timolol maleate.....	231	Capecitabine.....	158	Chlorothiazide.....	52
Brinov.....	158	Capoten.....	45	Chlorpheniramine maleate.....	221
Brinzolamide.....	230	Capsaicin		Chlorpromazine hydrochloride.....	131
Brolene.....	228	Musculoskeletal.....	111	Chlorsig.....	228
Bromocriptine mesylate.....	118	Nervous.....	121	Chlortalidone [Chlorthalidone].....	52
Brufen SR.....	110	Captopril.....	45	Chlorthalidone.....	52
BSF Logem.....	233	Carafate.....	9	Chlorvescent.....	44
BSF Mylan Efavirenz Emtricitabine Tenofovir.....	233	Carbaccord.....	156	Choice Load 375.....	72
BSF Teva Atazanavir Sulphate.....	233	Carbamazepine.....	126	Choice TT380 Short.....	72
BSF Teva Emtricitabine Tenofovir Disoprox.....	233	Carbimazole.....	83	Choice TT380 Standard.....	72
Buccastem.....	131	Carbomer.....	232	Choline salicylate with cetalkonium chloride.....	30
		Carboplatin.....	156	Ciclopirox olamine.....	60
		Carboplatin Ebewe.....	156	Ciclosporin.....	218
		Carbosorb-X.....	233	Cilazapril.....	45
		Cardinol LA.....	49	Cilazapril with hydrochlorothiazide.....	46
		CareSens Dual.....	12	Cilicaine.....	93
		CareSens N.....	12-13		
		CareSens N POP.....	12		
		CareSens N Premier.....	12		
		CareSens PRO.....	13		

Cilicaine VK.....	93	Colestipol hydrochloride.....	52	Daivonex.....	67
Cinacalcet.....	78	Colgout.....	116	Daktarin.....	61
Cipflox.....	94	Colifoam.....	7	Dalacin C.....	94
Ciprofloxacin		Colistin sulphomethate.....	94	Dalteparin sodium.....	40
Infection.....	94	Colistin-Link.....	94	Danazol.....	88
Sensory.....	228	Collodion flexible.....	236	Dantrium.....	117
Ciprofloxacin Teva.....	228	Colloidal bismuth subcitrate.....	9	Dantrium S29.....	117
Circadin.....	146	Colofac.....	8	Dantrolene.....	117
Cisplatin.....	156	Coloxyl.....	25	Daonil.....	11
Cisplatin Ebewe.....	156	Combigan.....	231	Dapa-Tabs.....	52
Citalopram hydrobromide.....	125	Compound electrolytes.....	44	Dapsone.....	100
Cladribine.....	158	Compound electrolytes with glucose		Daraprim.....	95
Clarithromycin		[Dextrose].....	44	Darunavir.....	107
Alimentary.....	8	Compound hydroxybenzoate.....	236	Dasatinib.....	166
Infection.....	91	Concerta.....	149	Daunorubicin.....	162
Clexane.....	41	Condoms.....	71	DBL Acetylcysteine.....	233
Clindamycin.....	94	Condyline.....	69	DBL Aminophylline.....	226
Clindamycin ABM.....	94	Contraceptives - Hormonal.....	72	DBL Bleomycin Sulfate.....	161
Clinicians Renal Vit.....	32	Contraceptives - Non-hormonal.....	70	DBL Carboplatin.....	156
Clobazam.....	126	Copaxone.....	141	DBL Cisplatin.....	156
Clobetasol propionate.....	62, 68	Cordarone-X.....	47	DBL Dacarbazine.....	161
Clobetasone butyrate.....	62	Corticosteroids and Related Agents		DBL Desferrioxamine Mesylate for Inj	
Clofazimine.....	99	for Systemic Use.....	79	BP.....	234
Clomazol		Corticosteroids Topical.....	62	DBL Docetaxel.....	162
Dermatological.....	60	Cosentyx.....	209	DBL Ergometrine.....	75
Genito-Urinary.....	75	Cosmegen.....	161	DBL Gemcitabine.....	158
Clomifene citrate.....	88	Coumadin.....	42	DBL Gentamicin.....	94
Clomipramine hydrochloride.....	124	Creon 10000.....	23	DBL Leucovorin Calcium.....	158
Clonazepam.....	126, 134	Creon 25000.....	23	DBL Methotrexate Onco-Vial.....	159
Clonidine.....	50	Cromal.....	230	DBL Morphine Sulphate.....	123
Clonidine BNM.....	50	Crotamiton.....	61	DBL Morphine Tartrate.....	123
Clonidine hydrochloride.....	50	Crystaderm.....	59	DBL Naloxone Hydrochloride.....	233
Clopidogrel.....	39	Curam.....	92	DBL Octreotide.....	172
Clopine.....	132	Cvite.....	31	DBL Pethidine Hydrochloride.....	124
Clopiol.....	133-134	Cyclizine hydrochloride.....	130	DBL Vincristine Sulfate.....	166
Clotrimazole		Cyclizine lactate.....	130	De-Worm.....	89
Dermatological.....	60	Cyclogyl.....	231	Decozol.....	31
Genito-Urinary.....	75	Cyclopentolate hydrochloride.....	231	Deferasirox.....	233
Clozapine.....	132	Cyclophosphamide.....	156	Deferiprone.....	234
Clozaril.....	132	Cyclorin.....	99	Denosumab.....	111
Clustran.....	129	Cycloserine.....	99	Deolate.....	98
Co-trimoxazole.....	96	Cyklokapron.....	39	Deoxycoformycin.....	164
Coal tar.....	68	Cyproterone acetate.....	80	Depo-Medrol.....	79
Coal tar with allantoin, menthol,		Cyproterone acetate with		Depo-Provera.....	74
phenol and sulphur.....	68	ethinyloestradiol.....	75	Depo-Testosterone.....	80
Coal tar with salicylic acid and		Cystadane.....	27	Deprim.....	96
sulphur.....	68	Cytarabine.....	158	Derm Assist.....	62
Coco-Scalp.....	68	Cytotec.....	8	Dermol.....	62, 68
Codeine phosphate		Cytoxan.....	156	Desferrioxamine mesilate.....	234
Extemporaneous.....	236			Desmopressin acetate.....	87
Nervous.....	121			Desmopressin-PH&T.....	87
Cogentin.....	119			Detection of Substances in	
Colaspase [L-asparaginase].....	161			Urine.....	77
Colchicine.....	116			Dexamethasone	
Colecalciferol.....	32			Hormone.....	79
Colestid.....	52			Sensory.....	229

**- D -**

Dexamethasone phosphate.....	79	Diuretics.....	51	Eformoterol fumarate dihydrate.....	222
Dexamethasone with framycetin and gramicidin.....	228	Diurin 40.....	51	Eftrenonacog alfa [Recombinant factor IX].....	36
Dexamethasone with neomycin sulphate and polymyxin B sulphate.....	229	Docetaxel.....	162	Efudix.....	69
Dexamfetamine sulfate.....	148	Docetaxel Accord.....	162	Egopsoryl TA.....	68
Dexmethsone.....	79	Docetaxel Sandoz.....	162	Elaprase.....	27
Dextrochlorpheniramine maleate.....	221	Docusate sodium.....	25	Elecare.....	254
Dextrose.....	43-44	Docusate sodium with sennosides.....	25	Elecare LCP.....	254
DHC Continus.....	121	Dolutegravir.....	107	Elilyso.....	29
Diabetes.....	9	Domperidone.....	130	Elemental 028 Extra.....	245
Diabetes Management.....	11	Donepezil hydrochloride.....	150	Elocon.....	63
Diacomit.....	128	Donepezil-Rex.....	150	Elocon Alcohol Free.....	63
Diagnostic Agents.....	266	Dopress.....	124	Eltrombopag.....	36
Diamide Relief.....	6	Dornase alfa.....	226	Eltroxin.....	83
Diamox.....	230	Dortimopt.....	230	EMB Fatol.....	100
Diasip.....	241	Dorzolamide hydrochloride.....	230	Emend Tri-Pack.....	130
Diason RTH.....	241	Dorzolamide with timolol.....	230	EMLA.....	120
Diazepam.....	126, 134	Dostinex.....	88	Emtricitabine.....	106
Diazoxide.....	9	Dosulepin [Dothiepin] hydrochloride.....	124	Emtricitabine with tenofovir disoproxil.....	103
Dibenzyline.....	45	Dothiepin.....	124	Emtriva.....	106
Diclofenac Sandoz.....	110	Doxazosin.....	45	Emulsifying ointment.....	64
Diclofenac sodium Musculoskeletal.....	110	Doxepin hydrochloride.....	124	Enalapril maleate.....	45
Sensory.....	229	Doxine.....	93	Enbrel.....	174
Differin.....	59	Doxorubicin Ebewe.....	162	Endocrine Therapy.....	171
Diffiam.....	30	Doxorubicin hydrochloride.....	162	Endoxan.....	156
Diflucan.....	96	Doxy-50.....	93	Enerlyte.....	44
Diflucan S29.....	96	Doxycycline.....	93	Engerix-B.....	259
Diflucortolone valerate.....	62	DP Lotion.....	64	Enlax XR.....	126
Digestives Including Enzymes.....	23	DP Lotn HC.....	62	Enoxaparin sodium.....	41
Digoxin.....	47	DP-Allopurinol.....	116	Ensure.....	249
Dihydrocodeine tartrate.....	121	Dr Reddy's Omeprazole.....	9	Ensure Plus.....	249
Dilantin.....	128	Drugs Affecting Bone Metabolism.....	111	Ensure Plus HN.....	248
Dilantin Infatab.....	128	Dual blood glucose and blood ketone diagnostic test meter.....	12	Ensure Plus RTH.....	248
Diltiazem hydrochloride.....	50	Duocal Super Soluble Powder.....	239	Entacapone.....	118
Dilzem.....	50	Duolin.....	223	Entapone.....	118
Dimethicone.....	64-65	Duolin HFA.....	223	Entecavir.....	101
Dimethyl fumarate.....	135	Durex Confidence.....	72	Entecavir Sandoz.....	101
Dipentum.....	7	Durex Extra Safe.....	71	Entocort CIR.....	6
Diphtheria, tetanus and pertussis vaccine.....	257	Duride.....	54	Entresto 24/26.....	46
Diphtheria, tetanus, pertussis and polio vaccine.....	258	- E -		Entresto 49/51.....	46
Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine.....	258	e-chamber La Grande.....	227	Entresto 97/103.....	46
Diprosone.....	62	e-chamber Mask.....	227	Epilim.....	128
Diprosone OV.....	62	e-chamber Turbo.....	227	Epilim Crushable.....	128
Dipyridamole.....	39	E-Mycin.....	91	Epilim IV.....	128
Disinfecting and Cleansing Agents.....	63	Ear Preparations.....	228	Epilim S/F Liquid.....	128
Disopyramide phosphate.....	47	Ear/Eye Preparations.....	228	Epilim Syrup.....	128
Disulfiram.....	152	Easiphen Liquid.....	253	Epirubicin Ebewe.....	162
		Econazole nitrate.....	60	Epirubicin hydrochloride.....	162
		Efavirenz.....	106	Eplerenone.....	51
		Efavirenz with emtricitabine and tenofovir disoproxil.....	106	Epoetin alfa.....	36
		Effient.....	39	Epoprostenol.....	57
		Eformoterol fumarate.....	222	Eptacog alfa [Recombinant factor VIIa].....	37
				ERA.....	91
				Erbitux.....	191

Ergometrine maleate .....	75	Felo 5 ER.....	50	Fluorouracil sodium .....	69
Ergotamine tartrate with caffeine .....	129	Felodipine .....	50	Fluoxetine hydrochloride .....	125
Erlotinib.....	167	Femme-Tab ED .....	73	Flupenthixol decanoate .....	133
Erythrocin IV .....	91	Fentanyl.....	122	FluQuadri.....	261
Erythromycin (as lactobionate).....	91	Fentanyl Sandoz.....	122	Flutamide.....	172
Erythromycin ethyl succinate.....	91	Ferinject.....	33	Flutamide Mylan.....	172
Erythromycin stearate.....	91	Ferodan .....	34	Flutamin.....	172
Esbriet .....	225	Ferric carboxymaltose .....	33	Fluticasone .....	222
Escitalopram.....	125	Ferriprox .....	234	Fluticasone furoate with vilanterol .....	222
Escitalopram-Apotex .....	125	Ferro-F-Tabs .....	34	Fluticasone propionate .....	227
Eskazole .....	89	Ferro-tab.....	34	Fluticasone with salmeterol .....	222
Estradot .....	81	Ferrograd.....	34	FML .....	230
Estradot 50 mcg .....	81	Ferrosig .....	34	Foban .....	60
Estrofem .....	81	Ferrous fumarate.....	34	Folic acid .....	36
Etanercept .....	174	Ferrous fumarate with folic acid.....	34	Food Thickeners.....	251
Ethambutol hydrochloride.....	100	Ferrous sulfate.....	34	Foods And Supplements For Inborn Errors Of Metabolism.....	252
Ethics Aspirin.....	121	Ferrous sulphate.....	34	Foradil.....	222
Ethics Aspirin EC.....	39	Ferrum H .....	34	Forteo .....	113
Ethics Enalapril.....	45	Fexofenadine hydrochloride .....	221	Fortini.....	243
Ethics Lisinopril.....	45	Fibro-vein.....	39	Fortini Multi Fibre.....	243
Ethinylestradiol .....	81	Filgrastim.....	42	Fortisip .....	249
Ethinylestradiol with desogestrel .....	73	Finasteride.....	76	Fortisip Multi Fibre.....	250
Ethinylestradiol with levonorgestrel .....	73	Fingolimod.....	136	Fosamax.....	111
Ethinylestradiol with norethisterone.....	73	Firazyr.....	220	Fosamax Plus.....	111
Ethosuximide .....	127	Flagyl.....	99	Fragmin.....	40
Etopophos .....	162	Flagyl-S .....	99	Framycetin sulphate .....	228
Etoposide.....	162	Flamazine.....	60	Fresenius Kabi.....	43-44
Etoposide phosphate.....	162	Flecainide acetate .....	47	Frisium.....	126
Etravirine .....	106	Flecainide BNM .....	47	Frumil.....	51
Eumovate .....	62	Flecainide Controlled Release Teva.....	47	Frusemide.....	51
Everet .....	127	Fleet Phosphate Enema .....	25	Frusemide-Claris .....	51
Everolimus.....	218	Flixonase Hayfever & Allergy .....	227	Fucicort.....	63
Evista.....	112	Flixotide .....	222	Fucidin.....	95
Exelon.....	151	Flixotide Accuhaler.....	222	Fucithalmic .....	228
Exemestane.....	173	Floair.....	222	Fungilin.....	31
Exjade.....	233	Florinef.....	79	Furosemide [Frusemide].....	51
Extemporaneously Compounded Preparations and Galenicals .....	236	Fluanxol.....	133	fusidic acid Dermatological .....	60, 63
Eye Preparations.....	228	Fluarix Tetra .....	261	Infection .....	95
Eylea.....	190	Flucil.....	92	Sensory.....	228
Ezetimibe.....	53	Flucloxacillin .....	92		
Ezetimibe Sandoz.....	53	Flucloxin .....	92		
Ezetimibe with simvastatin .....	53	Flucon.....	230		
		Fluconazole .....	96		
		Fludara Oral.....	158		
		Fludarabine Ebewe.....	158		
		Fludarabine phosphate.....	158		
		Fludrocortisone acetate.....	79		
		Fluids and Electrolytes .....	43		
		Flumetasone pivalate .....	228		
		Fluocortolone caproate with fluocortolone pivalate and cinchocaine.....	7		
		Fluorometholone.....	230		
		Fluorouracil.....	158		
		Fluorouracil Ebewe.....	158		

**- F -**
**- G -**



Gemfibrozil .....	52	healthE Dimethicone 4% Lotion .....	65	Hydrogen peroxide	
Gemzar .....	158	healthE Dimethicone 5% .....	64	Alimentary .....	31
Genoptic .....	228	healthE Glycerol BP .....	236	Dermatological .....	59
Gentamicin sulphate		healthE Urea Cream .....	64	Hydroxocobalamin .....	31
Infection .....	94	Healtheries Simple Baking Mix .....	251	Hydroxychloroquine .....	111
Sensory .....	228	Hemastix .....	77	Hydroxyurea .....	162
Gilenya .....	136	Heparin sodium .....	41	Hygroton .....	52
Ginet .....	75	Heparinised saline .....	42	Hylo-Fresh .....	232
Glatiramer acetate .....	141	Heparon Junior .....	242	Hymenoptera .....	220
Glecaprevir with pibrentasvir .....	103	Hepatitis A vaccine .....	258	Hyoscine butylbromide .....	8
Glibenclamide .....	11	Hepatitis B recombinant		Hyoscine hydrobromide .....	130
Gliclazide .....	11	vaccine .....	259	Hypam .....	147
Glipizide .....	11	Hepsera .....	101	Hyperuricaemia and Antigout .....	116
Glivec .....	167	Herceptin .....	214	Hypromellose .....	231
Glizide .....	11	Hexamine hippurate .....	109	Hypromellose with dextran .....	232
Glucagen Hypokit .....	9	Hiberix .....	258		- I -
Glucagon hydrochloride .....	9	Hiprex .....	109	Ibiamox .....	92
Glucerna Select .....	241	Histaclear .....	221	Ibuprofen .....	110
Glucerna Select RTH .....	241	Histafen .....	221	Icatibant .....	220
Glucobay .....	11	Holoxan .....	156	Idarubicin hydrochloride .....	162
Glucose [Dextrose] .....	43	Horleys Bread Mix .....	251	Idursulfase .....	27
Gluten Free Foods .....	251	Horleys Flour .....	252	Ifosfamide .....	156
Glycerin with sodium saccharin .....	236	Hormone Replacement Therapy -		Ikorel .....	55
Glycerin with sucrose .....	236	Systemic .....	80	Iloprost .....	58
Glycerol		HPV .....	260	Imatinib mesilate .....	167
Alimentary .....	25	Humalog .....	10	Imatinib-AFT .....	167
Extemporaneous .....	236	Humalog Mix 25 .....	10	Imipramine hydrochloride .....	124
Glyceryl trinitrate		Humalog Mix 50 .....	10	Imiquimod .....	69
Alimentary .....	8	Human papillomavirus (6, 11, 16, 18,		Immune Modulators .....	107
Cardiovascular .....	54	31, 33, 45, 52 and 58) vaccine		Immunosuppressants .....	174
Glycopyrronium .....	223	[HPV] .....	260	Imuran .....	174
Glycopyrronium bromide .....	8	Humatin .....	95	Incruse Ellipta .....	224
Glycopyrronium with		Humira .....	180	Indacaterol .....	222
indacaterol .....	224	HumiraPen .....	180	Indapamide .....	52
Glytrin .....	54	Humulin 30/70 .....	10	Infanrix IPV .....	258
Gold Knight .....	71-72	Humulin NPH .....	10	Infanrix-hexa .....	258
Goserelin .....	87	Humulin R .....	10	Infant Formulae .....	254
Gutron .....	48	Hyaluronic acid .....	232	Infatrin .....	255
Gynaecological Anti-infectives .....	75	Hybloc .....	48	Infliximab .....	191
	- H -	Hydralazine .....	54	Influenza vaccine .....	261
Habitrol .....	153	Hydralazine hydrochloride .....	54	Influvac Tetra .....	261
Haemophilus influenzae type B		Hydrea .....	162	Inhaled Corticosteroids .....	221
vaccine .....	258	Hydrocortisone		Inhaled Long-acting	
Haldol .....	133	Dermatological .....	62	Beta-adrenoceptor Agonists .....	222
Haldol Concentrate .....	133	Hormone .....	79	Inspra .....	51
Haldol Decanoas .....	133	Hydrocortisone acetate .....	7	Insulin aspart .....	10
Haloperidol .....	132	Hydrocortisone acetate with		Insulin aspart with insulin aspart	
Haloperidol decanoate .....	133	pramoxine hydrochloride .....	7	protamine .....	10
Hamilton Sunscreen .....	69	Hydrocortisone and paraffin liquid		Insulin glargine .....	10
Harvoni .....	103	and lanolin .....	62	Insulin glulisine .....	10
Havrix .....	258	Hydrocortisone butyrate .....	62, 68	Insulin isophane .....	10
Havrix Junior .....	258	Hydrocortisone with cinchocaine .....	7	Insulin isophane with insulin	
HBvaxPRO .....	259	Hydrocortisone with miconazole .....	63	neutral .....	10
healthE Calamine Aqueous Cream		Hydrocortisone with natamycin and		Insulin lispro .....	10
BP .....	61	neomycin .....	63	Insulin lispro with insulin lispro	
healthE Dimethicone 10% .....	64				

protamine.....	10	Ivermectin.....	65	Latanoprost.....	231
Insulin neutral.....	10	- J -		Lax-Suppositories.....	26
Insulin pen needles.....	13	Jadelle.....	74	Lax-Tab.....	26
Insulin pump.....	14	Jakavi.....	170	Laxatives.....	25
Insulin pump cartridge.....	19	Jevity HiCal RTH.....	248	Laxsol.....	25
Insulin pump infusion set (steel cannula).....	19	Jevity RTH.....	248	Ledipasvir with sofosbuvir.....	103
Insulin pump infusion set (steel cannula, straight insertion).....	20	Juno Pemetrexed.....	159	Leflunomide.....	111
Insulin pump infusion set (teflon cannula, angle insertion with insertion device).....	20	- K -		Lenalidomide.....	162
Insulin pump infusion set (teflon cannula, angle insertion).....	21	Kaletra.....	107	Letrole.....	174
Insulin pump infusion set (teflon cannula, straight insertion with insertion device).....	22	Keflor.....	89	Letrozole.....	174
Insulin pump infusion set (teflon cannula, straight insertion).....	23	Kemadrin.....	119	Leukeran FC.....	156
Insulin pump reservoir.....	23	Kenacomb.....	228	Leukotriene Receptor Antagonists.....	226
Insulin syringes, disposable with attached needle.....	14	Kenacort-A 10.....	80	Leunase.....	161
Intal Forte CFC Free.....	226	Kenacort-A 40.....	80	Leuprorelin.....	87
Intelence.....	106	Kenalog in Orabase.....	30	Leustatin.....	158
Interferon alfa-2a.....	108	Ketocal 3:1.....	256	Levetiracetam.....	127
Interferon beta-1-alpha.....	142	KetoCal 4:1.....	256	Levetiracetam-AFT.....	127
Interferon beta-1-beta.....	144	Ketoconazole.....		Levlen ED.....	73
Intra-uterine device.....	72	Dermatological.....	68	Levocabastine.....	230
Invega Sustenna.....	133	Infection.....	97	Levodopa with benserazide.....	118
IPOL.....	265	Ketogenic Diet.....	256	Levodopa with carbidopa.....	118
Ipratropium bromide.....	223, 227	Ketoprofen.....	110	Levomepromazine hydrochloride.....	132
Iressa.....	167	KetoSens.....	11	Levomepromazine maleate.....	132
Irinotecan Accord.....	159	Keytruda.....	216	Levonorgestrel.....	
Irinotecan Actavis 100.....	159	Kindergen.....	243	Genito-Urinary.....	74
Irinotecan hydrochloride.....	159	Kivexa.....	106	Hormone.....	82
Irinotecan-Rex.....	159	Klacid.....	91	Levothyroxine.....	83
Iron polymaltose.....	34	Kliogest.....	81	Lidocaine [Lignocaine].....	120
ISENTRESS.....	107	Kliovance.....	81	Lidocaine [Lignocaine] hydrochloride.....	120
ISENTRESS HD.....	107	Kogenate FS.....	38	Lidocaine [Lignocaine] with chlorhexidine.....	120
Ismo 20.....	54	Konakion MM.....	39	Lidocaine [Lignocaine] with prilocaine.....	120
Ismo 40 Retard.....	54	Konsyl-D.....	25	Lidocaine-Claris.....	120
Isoniazid.....	100	Kuvan.....	28	Lignocaine.....	120
Isoniazid with rifampicin.....	100	- L -		Lioresal Intrathecal.....	117
Isoprenaline [Isoproterenol].....	54	L-asparaginase.....	161	Lipazil.....	52
Isoproterenol.....	54	Labetalol.....	48	Lipid-Modifying Agents.....	52
Isoptin.....	50	Lacosamide.....	127	Liquigen.....	240
Isoptin SR.....	50	Lactulose.....	25	Lisinopril.....	45
Isopto Carpine.....	231	Laevolac.....	25	Lithicarb FC.....	132
Isosorbide mononitrate.....	54	Lamictal.....	127	Lithium carbonate.....	132
Isosource Standard.....	248	Lamivudine.....	101, 106	Livostin.....	230
Isotretinoin.....	59	Lamivudine Alphapharm.....	106	LMX4.....	120
Ispaghula (psyllium) husk.....	25	Lamotrigine.....	127	Locacorten-Viaform ED's.....	228
Isuprel.....	54	Lamprene.....	99	Local preparations for Anal and Rectal Disorders.....	7
Itch-Soothe.....	61	Lanoxin.....	47	Locasol.....	254
Itraconazole.....	97	Lanoxin PG.....	47	Locoid.....	62, 68
Itrazole.....	97	Lanoxin S29.....	47	Locoid Crelo.....	62
		Lansoprazole.....	8	Locoid Lipocream.....	62
		Lantus.....	10	Locorten-Vioform.....	228
		Lantus SoloStar.....	10	Lodi.....	47
		Lanvis.....	160		
		Lanzol Relief.....	8		
		Lapatinib ditosylate.....	168		
		Largactil.....	131		
		Laronidase.....	28		
		Lasix.....	51		

Lodoxamide.....	230	Medrol.....	79	Metronidazole.....	99
Logem.....	127	Medroxyprogesterone acetate		Metoprolol IV Mylan.....	49
Lomide.....	230	Genito-Urinary.....	74	Metyrapone.....	88
Lomustine.....	156	Hormone.....	81-82	Mexiletine hydrochloride.....	48
Loniten.....	55	Mefenamic acid.....	110	Mexiletine Hydrochloride USP.....	48
Loperamide hydrochloride.....	6	Megestrol acetate.....	172	Miacalcic.....	78
Lopinavir with ritonavir.....	107	Melatonin.....	146	Micollette.....	26
Loprofin.....	253	Melphalan.....	156	Miconazole.....	31
Loprofin Mix.....	253	Menactra.....	263	Miconazole nitrate	
Lorafix.....	221	Meningococcal (groups A, C, Y and		Dermatological.....	61
Loratadine.....	221	W-135) conjugate vaccine.....	263	Genito-Urinary.....	75
Lorazepam.....	135	Meningococcal C conjugate		Micreme.....	75
Lorfast.....	221	vaccine.....	263	Micreme H.....	63
Lorstat.....	53	Menthhol.....	61	Microgynon 20 ED.....	73
Losartan Actavis.....	46	Mercaptopurine.....	159	Microgynon 30.....	73
Losartan potassium.....	46	Mercilon 28.....	73	Microgynon 50 ED.....	73
Losartan potassium with		Mesalazine.....	7	Microlut.....	74
hydrochlorothiazide.....	46	Mesna.....	163	Midazolam.....	146
Lovir.....	102	Mestinon.....	110	Midazolam-Claris.....	146
Loxamine.....	125	Metabolic Disorder Agents.....	26	Midodrine.....	48
Lucrin Depot 1-month.....	87	Meterol.....	222	Milan Laboratories.....	51
Lucrin Depot 3-month.....	87	Metformin hydrochloride.....	11	Minerals.....	33
Ludiomil.....	125	Methadone hydrochloride		Mini-Wright AFS Low Range.....	227
Lyderm.....	67	Extemporaneous.....	236	Mini-Wright Standard.....	227
- M -		Nervous.....	122	Minidiab.....	11
m-Eslon.....	123	Methatabs.....	122	MiniMed 640G.....	14
Mabthera.....	200	Methopt.....	231	Minims Pilocarpine.....	231
Macrogl 3350 with potassium		Methotrexate.....	159	Minims Prednisolone.....	230
chloride, sodium bicarbonate and		Methotrexate Ebewe.....	159	Minirin.....	87
sodium chloride.....	25	Methotrexate Sandoz.....	159	Mino-tabs.....	93
Macrogl 400 and propylene		Methyl hydroxybenzoate.....	236	Minocycline hydrochloride.....	93
glycol.....	232	Methylcellulose.....	236	Minomycin.....	93
Madopar 125.....	118	Methylcellulose with glycerin and		Minor Skin Infections.....	65
Madopar 250.....	118	sodium saccharin.....	237	Minoxidil.....	55
Madopar 62.5.....	118	Methylcellulose with glycerin and		Mirena.....	82
Madopar HBS.....	118	sucrose.....	237	Mirtazapine.....	126
Madopar Rapid.....	118	Methylidopa.....	50	Misoprostol.....	8
Magnesium hydroxide		Methylidopa Mylan.....	50	Mitomycin C.....	163
Alimentary.....	34	Methylidopa Mylan S29.....	50	Mitozantrone.....	163
Extemporaneous.....	236	Methylnaltrexone bromide.....	25	Mitozantrone Ebewe.....	163
Magnesium sulphate.....	34	Methylphenidate ER - Teva.....	149	Mixtard 30.....	10
Mantoux.....	266	Methylphenidate hydrochloride.....	148	Moclobemide.....	125
Maprotiline hydrochloride.....	125	Methylphenidate hydrochloride		Modafinil.....	150
Marevan.....	42	extended-release.....	149	Modavigil.....	150
Marine Blue Lotion SPF 50+.....	69	Methylprednisolone.....	79	Moduretic.....	52
Marvelon 28.....	73	Methylprednisolone (as sodium		Molaxole.....	25
Mask for spacer device.....	227	succinate).....	79	Moments.....	71
Mast Cell Stabilisers.....	226	Methylprednisolone aceponate.....	62	Mometasone furoate.....	63
Maviret.....	103	Methylprednisolone acetate.....	79	Monogen.....	242
Maxidex.....	229	Methylxanthines.....	226	Montelukast.....	226
Maxitrol.....	229	Metoclopramide Actavis 10.....	130	Montelukast Mylan.....	226
MCT oil (Nutricia).....	240	Metoclopramide hydrochloride.....	130	Moroctocog alfa [Recombinant factor	
Measles, mumps and rubella		Metolazone.....	51	VIII].....	38
vaccine.....	262	Metopirone.....	88	Morphine hydrochloride.....	122
Mebendazole.....	89	Metoprolol succinate.....	49	Morphine sulphate.....	123
Mebeverine hydrochloride.....	8	Metoprolol tartrate.....	49	Morphine tartrate.....	123

Motetis .....	119	Neocate Junior Vanilla.....	254	Novatrelin .....	67
Mouth and Throat .....	30	Neocate SYNEO.....	254	NovoMix 30 FlexPen .....	10
Movapo.....	118	Neoral.....	218	NovoRapid.....	10
Moxifloxacin.....	94	Neostigmine metilsulfate .....	110	NovoRapid FlexPen.....	10
MSUD Maxamum .....	252	Nepro HP (strawberry).....	244	NovoRapid Penfill .....	10
Mucilaginous laxatives with stimulants .....	25	Nepro HP (vanilla).....	244	NovoSeven RT .....	37
Mucolytics.....	226	Nepro HP RTH .....	244	Noxafil.....	97
Mucosoothe.....	120	Nerisone .....	62	Nozinan .....	132
Multiple Sclerosis Treatments .....	135	Neulactil.....	132	Nuelin .....	226
Multivitamin renal.....	32	Neulastim.....	42	Nuelin-SR .....	226
Multivitamins.....	32	NeuroTabs.....	33	Nutilis.....	251
Mupirocin.....	60	Nevirapine .....	106	Nutrient Modules .....	238
Muscle Relaxants .....	117	Nevirapine Alphapharm .....	106	Nutrini Energy Multi Fibre .....	243
Mvite.....	32	Nicorandil.....	55	Nutrini Energy RTH .....	243
Myambutol.....	100	Nicotine .....	153	Nutrini Low Energy Multi Fibre .....	245
Mycobutin.....	100	Nicotinic acid .....	52	Nutrini RTH.....	243
MycoNail .....	60	Nifedipine.....	50	Nutrison 800 Complete Multi Fibre .....	248
Mycophenolate mofetil.....	174	Nifuran.....	109	Nutrison Concentrated.....	250
Mycostatin .....	61	Nilotinib .....	168	Nutrison Energy.....	248
Mydriacyl .....	231	Nilstat .....		Nutrison Energy Multi Fibre .....	248
Mylan Atenolol.....	48	Alimentary.....	31	Nutrison Multi Fibre .....	248
Mylan Clomiphen.....	88	Genito-Urinary.....	75	Nutrison Standard RTH .....	248
Myleran.....	156	Infection .....	97	Nyefax Retard.....	50
Myocrisin .....	111	Nintedanib .....	224	Nystatin .....	
Myometrial and Vaginal Hormone Preparations .....	75	Nipent .....	164	Alimentary .....	31
Myozyme .....	26	Nitrados .....	146	Dermatological.....	61
Mysoline S29 .....	128	Nitrates .....	54	Genito-Urinary.....	75
<b>- N -</b>		Nitrazepam .....	146	Infection .....	97
Nadolol .....	49	Nitroderm TTS.....	54	NZB Low Gluten Bread Mix .....	251
Naglazyme.....	27	Nitrofurantoin.....	109	<b>- O -</b>	
Nalcrom.....	7	Nitrolingual Pump Spray.....	54	O/W Fatty Emulsion Cream.....	64
Naloxone hydrochloride.....	233	Nivestim.....	42	Obinutuzumab .....	198
Nalttraccord.....	152	Nivolumab.....	215	Octocog alfa [Recombinant factor VIII] (Advate).....	38
Naltrexone hydrochloride.....	152	Nizoral .....	97	Octocog alfa [Recombinant factor VIII] (Kogenate FS).....	38
Naphazoline hydrochloride.....	232	Nodia .....	6	Octreotide .....	172
Naphcon Forte.....	232	Noflam 250 .....	110	Octreotide LAR (somatostatin analogue).....	172
Naprosyn SR 1000 .....	110	Noflam 500 .....	110	Oestradiol .....	81
Naprosyn SR 750 .....	110	Non-Steroidal Anti-Inflammatory Drugs .....	110	Oestradiol valerate .....	81
Naproxen.....	110	Nonacog alfa [Recombinant factor IX] .....	38	Oestradiol with norethisterone.....	81
Nardil.....	125	Nonacog gamma, [Recombinant Factor IX].....	38	Oestriol .....	
Nardil S29.....	125	Norethisterone .....		Genito-Urinary.....	75
Nasal Preparations .....	227	Genito-Urinary.....	74	Hormone .....	82
Natalizumab.....	138	Hormone .....	82	Oestrogens.....	81
Natulan .....	164	Norflex .....	117	Ofev .....	224
Nausafix.....	131	Norflex .....	109	Oil in water emulsion .....	64
Nausicalm.....	130	Noriday 28 .....	74	Olanzapine .....	132-133
Navelbine.....	166	Norimin .....	73	Olbetam .....	52
Nedocromil .....	226	Normacol Plus .....	25	Olopatadine .....	232
Nefopam hydrochloride .....	121	Normison .....	147	Olsalazine .....	7
Neisvac-C.....	263	Norpress .....	125	Omalizumab .....	199
Neo-B12 .....	31	Nortriptyline hydrochloride.....	125	Omeprazole.....	9
Neo-Mercazole .....	83	Norvir.....	107	Omeprazole actavis 10.....	9
Neocate Gold.....	254	NovaSource Renal .....	244		
Neocate Junior Unflavoured .....	254				

Omeprazole actavis 20.....	9	Pacifen.....	117	Parnate S29.....	125
Omeprazole actavis 40.....	9	Paclitaxel.....	163	Paromomycin.....	95
Omnitrope.....	83	Paclitaxel Actavis.....	163	Paroxetine.....	125
Onbrez Breezhaler.....	222	Paclitaxel Ebewe.....	163	Paser.....	100
Oncaspar.....	163	Paediatric Seravit.....	32	Patanol.....	232
OncoTICE.....	180	Paliperidone.....	133	Paxam.....	134
Ondansetron.....	131	Pamidronate disodium.....	112	Pazopanib.....	169
Ondansetron ODT-DRLA.....	131	Pamisol.....	112	Peak flow meter.....	227
Ondansetron ODT-ORLA.....	131	Pancreatic enzyme.....	23	Pedialyte - Bubblegum.....	44
One-Alpha.....	32	Pantoprazole.....	9	Pediasure.....	243
Opdivo.....	215	Panzop Relief.....	9	Pediasure RTH.....	243
Ora-Blend.....	237	Panzytrat.....	23	Pegaspargase.....	163
Ora-Blend SF.....	237	Papaverine hydrochloride.....	55	Pegasys.....	108
Ora-Plus.....	236	Para-amino salicylic acid.....	100	Pegfilgrastim.....	42
Ora-Sweet.....	236	Paracare.....	121	Pegylated interferon alfa-2a.....	108
Ora-Sweet SF.....	236	Paracare Double Strength.....	121	Pembrolizumab.....	216
Orabase.....	30	Paracetamol.....	121	Pemetrexed.....	159
Oral Supplements/Complete Diet (Nasogastric/Gastrostomy Tube Feed).....	241	Paracetamol + Codeine (Relieve).....	123	Penicillamine.....	111
Oratane.....	59	Paracetamol Pharmacare.....	121	Penicillin G.....	92
Ordine.....	122	Paracetamol with codeine.....	123	PenMix 30.....	10
Orgran.....	252	Paradigm 1.8 Reservoir.....	23	PenMix 40.....	10
Orion Temozolomide.....	164	Paradigm 3.0 Reservoir.....	23	PenMix 50.....	10
Ornidazole.....	99	Paradigm Mio MMT-921.....	22	Pentasa.....	7
Orphenadrine citrate.....	117	Paradigm Mio MMT-923.....	22	Pentostatin [Deoxycoformycin].....	164
Ortho-tolidine.....	77	Paradigm Mio MMT-925.....	22	Pentoxifylline [Oxpentifylline].....	55
Oruvail SR.....	110	Paradigm Mio MMT-941.....	22	Peptamen Junior.....	243
Osmolite RTH.....	248	Paradigm Mio MMT-943.....	22	Peptisoothe.....	8
Other Endocrine Agents.....	88	Paradigm Mio MMT-945.....	22	Peptisorb.....	245
Other Oestrogen Preparations.....	81	Paradigm Mio MMT-965.....	22	Perhexiline maleate.....	50
Other Progestogen Preparations.....	82	Paradigm Mio MMT-975.....	22	Pericyazine.....	132
Other Skin Preparations.....	69	Paradigm Quick-Set MMT-386.....	23	Perindopril.....	45
Ovestin.....		Paradigm Quick-Set MMT-387.....	23	Perjeta.....	200
Genito-Urinary.....	75	Paradigm Quick-Set MMT-396.....	23	Permethrin.....	67
Hormone.....	82	Paradigm Quick-Set MMT-397.....	23	Perrigo.....	69
Ox-Pam.....	135	Paradigm Quick-Set MMT-398.....	23	Pertuzumab.....	200
Oxalliccord.....	157	Paradigm Quick-Set MMT-399.....	23	Peteha.....	100
Oxaliplatin.....	157	Paradigm Silhouette MMT-368.....	21	Pethidine hydrochloride.....	124
Oxaliplatin Accord.....	157	Paradigm Silhouette MMT-377.....	21	Pevaryl.....	60
Oxaliplatin Actavis 100.....	157	Paradigm Silhouette MMT-378.....	21	Pexsig.....	50
Oxaliplatin Ebewe.....	157	Paradigm Silhouette MMT-381.....	21	Pfizer Exemestane.....	173
Oxazepam.....	135	Paradigm Silhouette MMT-382.....	21	Pharmacy Health Sorbolene with Glycerin.....	64
Oxis Turbuhaler.....	222	Paradigm Silhouette MMT-383.....	21	Pharmacy Services.....	233
Oxpentifylline.....	55	Paradigm Silhouette MMT-384.....	21	Pheburane.....	29
Oxybutynin.....	76	Paradigm Sure-T MMT-864.....	19	Phenaseen.....	160
Oxycodone hydrochloride.....	123	Paradigm Sure-T MMT-866.....	19	Phenelzine sulphate.....	125
Oxycodone Sandoz.....	123	Paradigm Sure-T MMT-874.....	19	Phenobarbitone.....	127
OxyNorm.....	123	Paradigm Sure-T MMT-876.....	19	Phenobarbitone sodium Extemporaneous.....	237
Oxytocin.....	75	Paradigm Sure-T MMT-884.....	19	Nervous.....	147
Oxytocin BNM.....	75	Paradigm Sure-T MMT-886.....	19	Phenothrin.....	67
Oxytocin with ergometrine maleate.....	75	Paraffin.....	64-65	Phenoxybenzamine hydrochloride.....	45
Ozurdex.....	229	Paraffin liquid with wool fat.....	232	Phenoxymethylpenicillin (Penicillin V).....	93
		Paraldehyde.....	126	Phenytoin sodium.....	126, 128
		Parasidose.....	67		
		Parasitidal Preparations.....	65		
		Parnate.....	125		

Phlexy 10.....	253	Prednisolone sodium phosphate.....	230	Pytazen SR.....	39
Phosphate Phebra.....	44	Prednisolone-AFT.....	230	- Q -	
Phosphorus.....	44	Prednisone.....	80	Q 300.....	99
Phytomenadione.....	39	Pregabalin.....	128	Quetapel.....	132
Pilocarpine hydrochloride.....	231	Pregabalin Pfizer.....	128	Quetiapine.....	132
Pimafucort.....	63	Pregnancy Tests - hCG Urine.....	75	Quick-Set MMT-390.....	23
Pindolol.....	49	Premarin.....	81	Quick-Set MMT-391.....	23
Pine tar with trolamine laurilsulfate and fluorescein.....	68	Presolol.....	48	Quick-Set MMT-392.....	23
Pinetarsol.....	68	Prevenar 13.....	264	Quick-Set MMT-393.....	23
Pioglitazone.....	11	Prezista.....	107	Quinapril.....	45
Pirfenidone.....	225	Priadel.....	132	Quinapril with hydrochlorothiazide.....	46
Pizotifen.....	130	Primacin.....	99	Quinine sulphate.....	99
PKU Anamix Infant.....	253	Primaquine phosphate.....	99	Qvar.....	221
PKU Anamix Junior.....	253	Primidone.....	128	- R -	
PKU Anamix Junior Chocolate.....	253	Primolut N.....	82	RA-Morph.....	122
PKU Anamix Junior LQ.....	253	Priorix.....	262	Raloxifene hydrochloride.....	112
PKU Anamix Junior Vanilla.....	253	Probenecid.....	117	Raltegravir potassium.....	107
PKU Lophlex LQ 10.....	253	Probenecid-AFT.....	117	Ramipex.....	118
PKU Lophlex LQ 20.....	253	Procaine penicillin.....	93	Ranbaxy-Cefaclor.....	89
PKU Lophlex Powder.....	253	Procarbazine hydrochloride.....	164	Ranitidine.....	8
PKU Lophlex Sensation 20.....	253	Prochlorperazine.....	131	Ranitidine Relief.....	8
Plaquenil.....	111	Proctofoam.....	7	Rapamune.....	218
Plendil ER.....	50	Proctosedyl.....	7	Reandron 1000.....	80
Pneumococcal (PCV10) conjugate vaccine.....	263	Procyclidine hydrochloride.....	119	Recombinant factor IX.....	36, 38
Pneumococcal (PCV13) conjugate vaccine.....	264	Procytox.....	156	Recombinant factor VIIa.....	37
Pneumococcal (PPV23) polysaccharide vaccine.....	265	Progesterone.....	82	Recombinant factor VIII.....	38-39
Pneumovax 23.....	265	Proglidem.....	9	Rectogesic.....	8
Podophyllotoxin.....	69	Proglycem.....	9	Redipred.....	80
Polaramine.....	221	Progynova.....	81	Relieve.....	110
Poliomyelitis vaccine.....	265	Prolia.....	111	Relistor.....	25
Poloxamer.....	25	Promethazine hydrochloride.....	221	Remicade.....	191
Poly-Gel.....	232	Propafenone hydrochloride.....	48	Renilon 7.5.....	244
Poly-Tears.....	232	Propamidine isethionate.....	228	Resonium-A.....	44
Poly-Visc.....	232	Propranolol.....	49	Resource Beneprotein.....	240
Polycal.....	238	Propylene glycol.....	237	Resource Diabetic.....	241
Polyvinyl alcohol.....	232	Propylthiouracil.....	83	Respigen.....	223
Nonstan.....	110	Protaphane.....	10	Respiratory Devices.....	227
Posaconazole.....	97	Protaphane Penfill.....	10	Respiratory Stimulants.....	227
Postinor-1.....	74	Protifar.....	240	Retinol palmitate.....	232
Potassium chloride.....	43-44	Protionamide.....	100	ReTrieve.....	59
Potassium citrate.....	76	Provera.....	81	Retrovir.....	106
Potassium iodate.....	33	Provera HD.....	82	Revlimid.....	162
Povidone iodine.....	65	PSM Citalopram.....	125	Revolade.....	36
Pradaxa.....	42	Psoriasis and Eczema Preparations.....	67	Rexacrom.....	230
Pramipexole hydrochloride.....	118	PTU.....	83	RexAir.....	222
Prasugrel.....	39	Pulmicort Turbuhaler.....	221	Ribomustin.....	155
Pravastatin.....	53	Pulmocare.....	241	Ricit.....	76
Praziquantel.....	89	Pulmozyme.....	226	Rifabutin.....	100
Prazosin.....	45	Puri-nethol.....	159	Rifadin.....	101
Pred Forte.....	230	Puria.....	32	Rifampicin.....	101
Prednisolone.....	80	Pyrazinamide.....	100	Rifaximin.....	9
Prednisolone acetate.....	230	Pyridostigmine bromide.....	110	Rifinah.....	100
		Pyridoxine hydrochloride.....	31	Rilutek.....	119
		Pyrimethamine.....	95	Riluzole.....	119
				Riodine.....	65

Risedronate Sandoz.....	113	Senokot.....	26	acid].....	232
Risedronate sodium.....	113	Sensipar.....	78	Sodium phenylbutyrate.....	29
Risperdal Consta.....	134	SensoCard.....	13	Sodium polystyrene sulphonate.....	44
Risperidone.....	133-134	Serenace.....	132	Sodium tetradecyl sulphate.....	39
Risperon.....	133	Seretide.....	222	Sodium valproate.....	128
Ritalin.....	148	Seretide Accuhaler.....	222	Sofradex.....	228
Ritalin LA.....	149	Serevent.....	222	Soframycin.....	228
Ritalin SR.....	148	Serevent Accuhaler.....	222	Solian.....	131
Ritonavir.....	107	Sertraline.....	126	Solifenacin Mylan.....	76
Rituximab.....	200	Setrona.....	126	Solifenacin succinate.....	76
Rivaroxaban.....	42	Sevredol.....	123	Solu-Cortef.....	79
Rivastigmine.....	151	Sex Hormones Non		Solu-Medrol.....	79
Rivotril.....	126	Contraceptive.....	80	Solu-Medrol-Act-O-Vial.....	79
RIXUBIS.....	38	Shield 49.....	71	Somatropin (Omnitrope).....	83
Rizamelt.....	129	Shield Blue.....	71	Sotalol.....	49
Rizatriptan.....	129	Shield XL.....	72	Spacer device.....	227
Roferon-A.....	108	shingles vaccine.....	266	Span-K.....	44
Rolin.....	173	SII-Onco-BCG.....	180	Spilto Respimat.....	224
Ropin.....	118	Sildenafil.....	56	Spiractin.....	51
Ropinirole hydrochloride.....	118	Silhouette MMT-371.....	21	Spiriva.....	224
Rotarix.....	265	Silhouette MMT-373.....	21	Spiriva Respimat.....	224
Rotavirus oral vaccine.....	265	Siltuximab.....	210	Spiroolactone.....	51
Roxane		Simvastatin.....	53	Sporanox.....	97
Alimentary.....	6	Simvastatin Mylan.....	53	Sprycel.....	166
Cardiovascular.....	49	Sinemet.....	118	Staphlex.....	92
Roxithromycin.....	91	Sinemet CR.....	118	Stemetil.....	131
Rubifen.....	148	Sirolimus.....	218	SteroClear.....	227
Rubifen SR.....	148	Siterone.....	80	Stesolid.....	126
Rulide D.....	91	Slow-Lopresor.....	49	Stimulants/ADHD Treatments.....	147
Ruriotocog alfa pegol [Recombinant		Smith BioMed Rapid Pregnancy		Stiripentol.....	128
factor VIII].....	39	Test.....	75	Stocrin.....	106
Ruxolitinib.....	170	Sodibic.....	44	Stomahesive.....	30
Rythmodan.....	47	Sodium acid phosphate.....	25	Strattera.....	147
Rytmonorm.....	48	Sodium alginate.....	6	Stromectol.....	65
- S -		Sodium aurothiomalate.....	111	Suboxone.....	151
Sabril.....	128	Sodium benzoate.....	28	Sucalfate.....	9
Sacubitril with valsartan.....	46	Sodium bicarbonate		Sulfadiazine Silver.....	60
SalAir.....	223	Blood.....	43-44	Sulfadiazine sodium.....	95
Salazopyrin.....	7	Extemporaneous.....	237	Sulfasalazine.....	7
Salazopyrin EN.....	7	Sodium calcium edetate.....	234	Sulindac.....	110
Salbutamol.....	223	Sodium chloride		Sulphur.....	68
Salbutamol with ipratropium		Blood.....	43	Sulprix.....	131
bromide.....	223	Respiratory.....	227	Sumatriptan.....	129
Salicylic acid.....	68	Sodium citrate with sodium lauryl		Sunitinib.....	170
Salmeterol.....	222	sulphoacetate.....	26	Sunscreens.....	69
Sandomigran.....	130	Sodium citro-tartrate.....	76	Sunscreens, proprietary.....	69
Sandostatin LAR.....	172	Sodium cromoglicate		Sure-T MMT-863.....	19
Sapropterin dihydrochloride.....	28	Alimentary.....	7	Sure-T MMT-865.....	19
Scalp Preparations.....	68	Respiratory.....	226	Sure-T MMT-873.....	19
Scopoderm TTS.....	130	Sensory.....	230	Sure-T MMT-875.....	19
Sebizole.....	68	Sodium fluoride.....	33	Sure-T MMT-883.....	19
Secukinumab.....	209	Sodium Fusidate [fusidic acid]		Sure-T MMT-885.....	19
Sedatives and Hypnotics.....	146	Dermatological.....	60	Sustagen Diabetic.....	241
Seebri Breezhaler.....	223	Infection.....	95	Sustagen Hospital Formula	
Selegiline hydrochloride.....	118	Sensory.....	228	Active.....	249
Senna.....	26	Sodium hyaluronate [Hyaluronic		Sustanon Ampoules.....	80



Sutent.....	170	Tetracosactrin.....	80	Tretinoin	
Sylvant.....	210	Tetracyclin Wolff.....	93	Dermatological.....	59
Symbicort Turbuhaler 100/6.....	222	Tetracycline.....	93	Oncology.....	165
Symbicort Turbuhaler 200/6.....	222	Thalidomide.....	165	Trexate.....	159
Symbicort Turbuhaler 400/12.....	222	Thalomid.....	165	Triamcinolone acetonide	
Symmetrel.....	118	Theophylline.....	226	Alimentary.....	30
Sympathomimetics.....	54	Thiamine hydrochloride.....	31	Dermatological.....	63
Synacthen.....	80	THIO-TEPA.....	157	Hormone.....	80
Synacthen Depot.....	80	Thioguanine.....	160	Triamcinolone acetonide with	
Synacthen S29.....	80	Thiotepa.....	157	gramicidin, neomycin and nystatin	
Synacthene Retard.....	80	Thymol glycerin.....	31	Dermatological.....	63
Synflorix.....	263	Thyroid and Antithyroid Agents.....	83	Sensory.....	228
Synthroid.....	83	Ticagrelor.....	39	Triazolam.....	147
Syntometrine.....	75	Tilade.....	226	Trichazole.....	99
Syrup (pharmaceutical grade).....	237	Tilcotil.....	110	Triclosan.....	63
Systane Unit Dose.....	232	Timolol		Trimethoprim.....	96
<b>- T -</b>		Cardiovascular.....	49	Trimethoprim with	
T&R.....	34	Sensory.....	230	sulphamethoxazole	
Tacrolimus.....	218	Timoptol XE.....	230	[Co-trimoxazole].....	96
Tacrolimus Sandoz.....	218	Tiotropium bromide.....	224	Trisequens.....	81
Taliglucerase alfa.....	29	Tiotropium bromide with		Trisul.....	96
Tambocor.....	47	olodaterol.....	224	Trophic Hormones.....	83
Tambocor CR.....	47	Tivicay.....	107	Tropicamide.....	231
Tamoxifen citrate.....	173	TMP.....	96	Trusopt.....	230
Tamoxifen Sandoz.....	173	TOBI.....	96	TruSteel.....	20
Tamsulosin hydrochloride.....	76	Tobramycin		Tuberculin PPD [Mantoux] test.....	266
Tamsulosin-Rex.....	76	Infection.....	96	Tubersol.....	266
Tandem Cartridge.....	19	Sensory.....	229	Two Cal HN.....	251
Tandem t:slim X2.....	14	Tobramycin Mylan.....	96	Two Cal HN RTH.....	250
Tap water.....	237	Tobrex.....	229	Tykerb.....	168
Tarceva.....	167	Tocilizumab.....	211	Tysabri.....	138
Tasigna.....	168	Tofranil.....	124	<b>- U -</b>	
Tasmar.....	118	Tolcapone.....	118	Ultibro Breezhaler.....	224
Tecfidera.....	135	Tolterodine.....	77	Ultraproct.....	7
Tegretol.....	126	Topamax.....	128	Umeclidinium.....	224
Tegretol CR.....	126	Topical Products for Joint and		Umeclidinium with vilanterol.....	224
Telfast.....	221	Muscular Pain.....	111	Univent.....	223, 227
Temaccord.....	164	Topiramate.....	128	Ural.....	76
Temazepam.....	147	Topiramate Actavis.....	128	Urea.....	64
Temizole 20.....	164	Total parenteral nutrition (TPN).....	43	Urex Forte.....	51
Temozolomide.....	164	TPN.....	43	Urinary Agents.....	76
Tenofovir disoproxil.....	102	Tramadol hydrochloride.....	124	Urinary Tract Infections.....	109
Tenofovir Disoproxil Teva.....	102	Tramal SR 100.....	124	Uromitexan.....	163
Tenoxicam.....	110	Tramal SR 150.....	124	Ursodeoxycholic acid.....	24
Tepadina.....	157	Tramal SR 200.....	124	Ursosan.....	24
Terazosin.....	45	Trandate.....	48	Utrogestan.....	82
Terbinafine.....	98	Tranexamic acid.....	39	<b>- V -</b>	
Terbutaline sulphate.....	223	Tranylcypromine sulphate.....	125	Vaccinations.....	257
Teriflunomide.....	139	Trastuzumab.....	214	Vaclovir.....	102
Teriparatide.....	113	Travatan.....	231	Valaciclovir.....	102
Testosterone.....	80	Travoprost.....	231	Valganciclovir.....	102
Testosterone cypionate.....	80	Travopt.....	231	Valganciclovir Mylan.....	102
Testosterone esters.....	80	Treatments for Dementia.....	150	Vancomycin.....	96
Testosterone undecanoate.....	80	Treatments for Substance		Vannair.....	222
Tetrabenazine.....	119	Dependence.....	151	Varenicline Pfizer.....	153
Tetrabromophenol.....	77	Trental 400.....	55	Varenicline tartrate.....	153



Varicella vaccine [Chickenpox vaccine] .....	266	Votrient .....	169	Zytiga .....	171
Varicella zoster virus (Oka strain) live attenuated vaccine [shingles vaccine] .....	266	Vttack .....	98		
Varilrix .....	266	- W -			
Various .....	233	Warfarin sodium .....	42		
Vasodilators .....	54	Wart Preparations .....	69		
Vasopressin Agonists .....	87	Wasp venom allergy treatment .....	220		
Vedafil .....	56	Water			
Velcade .....	161	Blood .....	44		
Veletri .....	57	Extemporaneous .....	237		
Venlafaxine .....	126	Wool fat with mineral oil .....	64		
Venomil .....	220	- X -			
Ventavis .....	58	Xarelto .....	42		
Ventolin .....	223	Xifaxan .....	9		
Vepesid .....	162	XMET Maxamum .....	252		
Verapamil hydrochloride .....	50	Xolair .....	199		
Vergo 16 .....	130	XP Maxamum .....	253		
Vermox .....	89	Xylocaine .....	120		
Verpamil SR .....	50	Xylocaine 2% Jelly .....	120		
Vesanoid .....	165	Xyntha .....	38		
Vexazone .....	11	- Z -			
Vfend .....	98	Zantac .....	8		
Viaderm KC .....	63	Zapril .....	45		
Vidaza .....	157	Zarontin .....	127		
Vigabatrin .....	128	Zaroxolyn .....	51		
Vildagliptin .....	11	Zavedos .....	162		
Vildagliptin with metformin hydrochloride .....	11	Zeffix .....	101		
Vimpat .....	127	Zetlam .....	101		
Vinblastine sulphate .....	166	Ziagen .....	106		
Vincristine sulphate .....	166	Zidovudine [AZT] .....	106		
Vinorelbine .....	166	Zidovudine [AZT] with lamivudine .....	107		
Vinorelbine Ebewe .....	166	Zimybe .....	53		
Viramune Suspension .....	106	Zinc and castor oil .....	64		
ViruPOS .....	228	Zinc sulphate .....	34		
Vistil .....	232	Zincaps .....	34		
Vistil Forte .....	232	Zinnat .....	90		
Vit.D3 .....	32	Ziprasidone .....	133		
VitA-POS .....	232	Zista .....	221		
Vitabdeck .....	32	Zithromax .....	90		
Vitadol C .....	31	Zoladex .....	87		
Vital .....	245	Zoledronic acid			
Vitamin A with vitamins D and C .....	31	Hormone .....	78		
Vitamin B complex .....	31	Musculoskeletal .....	114		
Vitamin B6 25 .....	31	Zoledronic acid Mylan .....	78		
Vitamins .....	31-32	Zopiclone .....	147		
Vivonex TEN .....	245	Zopiclone Actavis .....	147		
Volibris .....	55	Zostavax .....	266		
Voltaren .....	110	Zostrix .....	111		
Voltaren D .....	110	Zostrix HP .....	121		
Voltaren Ophtha .....	229	Zuclopenthixol decanoate .....	134		
Volumatic .....	227	Zuclopenthixol hydrochloride .....	133		
Voriconazole .....	98	Zusdone .....	133		
Vosol .....	228	Zyban .....	152		
		Zypine .....	132		
		Zypine ODT .....	132		
		Zyprexa Relprev .....	133		













