November 2019 Volume 26 Number 2

Editors:

Kaye Wilson & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

Published each April, August and December. Changes to the contents are published in monthly updates.

Accessible in an electronic format at no cost from the PHARMAC website www.pharmac.govt.nz.

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at www.pharmac.govt.nz/subscriptions.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from www.pharmac.govt.nz/pub/schedule/archive/

Programmers

Anrik Drenth & John Geering email: texschedule@pharmac.govt.nz @Pharmaceutical Management Agency



ISSN 1179-3686 pdf ISSN 1172-9376 print

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to PHARMAC and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to PHARMAC should be in written form and not by reproduction of the PHARMAC logo. While care has been taken in compiling this Schedule, PHARMAC takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

	introducing i manimac	
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	35
	Cardiovascular System	45
	Dermatologicals	59
	Genito Urinary System	70
	Hormone Preparations – Systemic	78
	Infections – Agents For Systemic Use	89
	Musculoskeletal System	110
	Nervous System	118
	Oncology Agents & Immunosuppressants	155
	Respiratory System & Allergies	220
	Sensory Organs	228
	Various	233
Section C	Extemporaneous Compounds (ECPs)	235
Section D	Special Foods	238
Section I	National Immunisation Schedule	257

Index

Introducing PHARMAC

Introducing PHARMAC

The Pharmaceutical Management Agency (PHARMAC) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. PHARMAC negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

PHARMAC's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about PHARMAC and the way we make funding decisions can be found on the PHARMAC website at http://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by PHARMAC.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements PHARMAC has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example




Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS



SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg p sachet		30	✓	Gaviscon Infant
SODIUM ALGINATE				
* Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (8.60)	60		Gaviscon Double Strength
* Oral liq 500 mg with sodium bicarbonate 267 mg and calciun carbonate 160 mg per 10 ml		500 m		Acidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE * Tab 600 mg	12.56	100	•	Alu-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsementOnly when prescribed for patients unable to swallow cal inappropriate and the prescription is endorsed according	cium carbonate tablet	500 m s or v		Roxane um carbonate tablets are
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on * Tab 2 mg* * Cap 2 mg	10.75	400 400		Nodia Diamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE Cap 3 mg - Special Authority see SA1155 below - Retail pharmacy	166.50	90	,	Entocort CIR
■ SA1155 Special Authority for Subsidy Initial application — (Crohn's disease) from any relevant practithe following criteria: Both:	titioner. Approvals va	ılid fo	r 6 months	for applications meeting
Mild to moderate ileal, ileocaecal or proximal Crohn's disc	ease; and			

2.1 Diabetes; or2.2 Cushingoid habitus; or

2.3 Osteoporosis where there is significant risk of fracture; or

continued...

2 Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)	26.55	21.1 g OP	✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLO	RIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%	26.55	10 g OP	✓ Proctofoam S29
MESALAZINE			
Tab 400 mg	49.50	100	✓ Asacol
Tab EC 500 mg	49.50	100	✓ Asamax
Tab long-acting 500 mg	59.05	100	✓ Pentasa
Tab 800 mg	85.50	90	✓ Asacol
Modified release granules, 1 g		120 OP	✓ Pentasa
Enema 1 g per 100 ml	41.30	7	✓ Pentasa
Suppos 500 mg	22.80	20	✓ Asacol
Suppos 1 g	54.60	30	✓ Pentasa
OLSALAZINE			
Tab 500 mg	93.37	100	✓ Dipentum
Cap 250 mg		100	✓ Dipentum
SODIUM CROMOGLICATE			
Cap 100 mg	92.91	100	✓ Nalcrom
SULFASALAZINE			
* Tab 500 mg	14.00	100	✓ Salazopyrin
* Tab EC 500 mg		100	✓ Salazopyrin EN
Salazopyrin EN to be Sole Supply on 1 December 2019			.,

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cinchocaine hydrochloride 5 mg per g	30 g OP	✓ Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and cinchocaine hydrochloride 1 mg2.66	12	✓ Ultraproct
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	Proctosedyl

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Management of Anal Fissures

GLYCERYL TRINITRATE - Special Authority see SA1329 below - Retail pharmacy

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE

Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a		
PSO17.14	10	Max Health

HYOSCINE BUTYLBROMIDE

111	OCCURE DOTTEDITORIDE		
*	Tab 10 mg8.75	100	✓ Buscopan
*	Inj 20 mg, 1 ml - Up to 5 inj available on a PSO9.57	5	✓ Buscopan

MEBEVERINE HYDROCHLORIDE

*	Tab 135 mg18.0	00 90	✓ Colofac
---	----------------	-------	-----------

Antiulcerants

Antisecretory and Cytoprotective

MISOPROSTOL

*	Tab 200 mcg41.50	120	Cytote	С
---	------------------	-----	--------	---

Helicobacter Pylori Eradication

CLARITHROMYCIN

Tab 500 mg - Subsidy b	by endorsement	10.40	14	✓ A	po-Clarithromycin

- a) Maximum of 14 tab per prescription
- Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.
 Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

H2 Antagonists

RANITIDINE - Subsidy by endorsement

- a) Only on a prescription
- b) Subsidy by endorsement Subsidised for patients who were taking ranitidine prior to 1 November 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of ranitidine.

*	Tab 150 mg12.91	500	Ranitidine Relief
	Tab 300 mg18.21	500	✓ Ranitidine Relief
	Oral liq 150 mg per 10 ml5.14	300 ml	✓ Peptisoothe
*	Inj 25 mg per ml, 2 ml13.40	5	✓ Zantac

Proton Pump Inhibitors

ANICODDAZOLE

LA	NSOPRAZULE		
*	Cap 15 mg4.58	100	✓ Lanzol Relief
*	Cap 30 mg5.41	100	✓ Lanzol Relief

		Subsidy		Fully	
		(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
\ \	EPRAZOLE	Ψ	1 01		Wandactarci
JΙV	For omeprazole suspension refer Standard Formulae, page	235			
*	Cap 10 mg		90	/	Omeprazole actavis
	, ,				10
*	Cap 20 mg	1.96	90	✓	Omeprazole actavis
				_	<u>20</u>
*	Cap 40 mg	3.12	90	/	Omeprazole actavis
v	Dauder Only in combination	40.50	E ~	./	40 Midwest
*	Powder – Only in combination		5 g	•	Midwest
*	Only in extemporaneously compounded omeprazole su Inj 40 mg ampoule with diluent		5	J	Dr Reddy's
	ing to mg ampoute with undertt	00.30	J	•	Omeprazole
οΔ	NTOPRAZOLE				<u> </u>
	Tab EC 20 mg	2.02	100	1	Panzop Relief
	Tab EC 40 mg		100		Panzop Relief
	<u> </u>				
S	ite Protective Agents				
C	LLOIDAL BISMUTH SUBCITRATE				
	Tab 120 mg	14.51	50	1	Gastrodenol S29
21.1	CRALFATE				
,0	Tab 1 g	35.50	120		
		(48.28)	0		Carafate
_		(/			
B	ile and Liver Therapy				
RIF	AXIMIN - Special Authority see SA1461 below - Retail pha	rmacy			
	Tab 550 mg	625.00	56	1	Xifaxan
	SA1461 Special Authority for Subsidy				
	ial application only from a gastroenterologist, hepatologist of	r Practitioner on the re	ecomi	mendation	of a gastroenterologist o
	patologist. Approvals valid for 6 months where the patient ha				
	erated doses of lactulose.		, -		7
ìе	newal only from a gastroenterologist, hepatologist or Practition	oner on the recommen	datio	n of a gast	roenterologist or
ep	patologist. Approvals valid without further renewal unless not	ified where the treatme	ent re	emains app	propriate and the patient i
er	nefiting from treatment.				
	Shates				
D	iabetes				
Н	yperglycaemic Agents				
11/	ZOXIDE – Special Authority see SA1320 below – Retail pha	110 00	100	,	Proglicem \$29

DIAZONIDE – Special Authority see SA 1320 below –	Retail pharmacy		
Cap 25 mg	110.00	100	✓ Proglicem S29
Cap 100 mg	280.00	100	✓ Proglicem S29
Oral liq 50 mg per ml	620.00	30 ml OP	✓ Proglycem S29

⇒SA1320 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where used for the treatment of confirmed hypoglycaemia caused by hyperinsulinism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

GLUCAGON HYDROCHLORIDE

Inj 1 mg syringe kit − Up to 5 kit available on a PSO......32.00 1 ✓ Glucagen Hypokit

	Subsidy		Fully	Brand or
	(Manufacturer's F		sidised	Generic
	\$	Per		Manufacturer
Inculin Chart acting Drangrations				
Insulin - Short-acting Preparations				
NSULIN NEUTRAL				
Inj human 100 u per ml	25.26	10 ml OP	✓ A	Actrapid
,			✓ H	lumulin R
Inj human 100 u per ml, 3 ml	42.66	5	✓ A	Actrapid Penfill
,			✓ H	lumulin R
Insulin - Intermediate-acting Preparations				
NSULIN ASPART WITH INSULIN ASPART PROTAMINE				
Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	✓ N	lovoMix 30 FlexPen
NSULIN ISOPHANE				
	17.60	10 ml OP	./ L	lumulin NPH
Inj human 100 u per ml	17.00	10 1111 OF	_	
h Ini human 100	00.00	-		Protaphane
Inj human 100 u per ml, 3 ml	29.86	5		lumulin NPH
			• 1	Protaphane Penfill
NSULIN ISOPHANE WITH INSULIN NEUTRAL				
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP		lumulin 30/70
				Mixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5		lumulin 30/70
				PenMix 30
				PenMix 40
			✓ P	PenMix 50
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE				
Inj lispro 25% with insulin lispro protamine 75% 100 u per ml,				
3 ml		5	✓ H	lumalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml,		ū	-	
3 ml		5	✓ H	lumalog Mix 50
V 111		<u> </u>	• •	iumalog iinx oo
Insulin - Long-acting Preparations				
NSULIN GLARGINE				
Inj 100 u per ml, 10 ml	63.00	1	√ I	antus.
Inj 100 u per ml, 3 ml	94 50	5		antus
Inj 100 u per ml, 3 ml disposable pen		5	_	antus SoloStar
L my 100 a por mi, o mi aloposablo por minimi minimi				
Insulin - Rapid Acting Preparations				
NSULIN ASPART				
▲ Inj 100 u per ml, 10 ml	30.03	1	✓ N	lovoRapid
Inj 100 u per ml, 3 ml	51 10	5		lovoRapid Penfill
Inj 100 u per ml, 3 ml syringe		5		lovoRapid FlexPen
		3	• 1	iovonapiu riexreii
NSULIN GLULISINE		_		
Inj 100 u per ml, 10 ml		1	_	Apidra
Inj 100 u per ml, 3 ml		5		Apidra
Inj 100 u per ml, 3 ml disposable pen	46.07	5	✓ A	Apidra SoloStar
NSULIN LISPRO				
▲ Inj 100 u per ml, 10 ml	34.92	10 ml OP	✓ H	lumalog
▲ Inj 100 u per ml, 3 ml		5	_	lumalog
				•

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
Alpha Glucosidase Inhibitors				
ACARBOSE				
* Tab 50 mg	3.50	90	1	Glucobay
v	10.47		✓	Accarb
★ Tab 100 mg	6.40	90	✓	Glucobay
	11.24	50	✓	Acarbose Mylan S29
	20.23	90	✓	Accarb
Acarbose Mylan 👀 Tab 100 mg to be delisted 1 January	2020)			
Oral Hypoglycaemic Agents				
GLIBENCLAMIDE				
* Tab 5 mg	6.00	100	1	Daonil
GLICLAZIDE				
★ Tab 80 mg	10.29	500	1	Glizide
GLIPIZIDE		000		
₭ Tab 5 mg	3 27	100	1	Minidiab
· ·		100	•	Millidiab
METFORMIN HYDROCHLORIDE	0.60	1 000		Anatov
★ Tab immediate-release 500 mg ★ Tab immediate-release 850 mg		1,000 500		Apotex Apotex
Ç	7.04	300	•	Mhorex
PIOGLITAZONE	0.47	00	,	W
₭ Tab 15 mg		90		Vexazone
K Tab 45 mg		90		Vexazone Vexazone
≰ Tab 45 mg	7.10	90	•	Vexazone
'ILDAGLIPTIN			_	
Tab 50 mg	40.00	60	/	Galvus
ILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride		60		Galvumet
Tab 50 mg with 850 mg metformin hydrochloride	40.00	60	✓	Galvumet

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

10 strip OP ✓ KetoSens

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes: or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

1 OP CareSens Dual

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes; or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 diagnostic test

1 OP ✓ CareSens N

✓ CareSens N POP 20.00

✓ CareSens N Premier

Note: Only 1 meter available per PSO

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	1	CareSens N
		1	CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips	.20 50 te	st OP	SensoCard
---------------------------	-----------	-------	-----------

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

INSULIN PEN NEEDLES - Maximum of 200 dev per prescription

*	29 g × 12.7 mm10.50	100	✓ B-D Micro-Fine
*	31 g × 5 mm11.75	100	✓ B-D Micro-Fine
	31 g × 6 mm9.50		✓ Berpu
	31 g × 8 mm		✓ B-D Micro-Fine
	32 g × 4 mm		✓ B-D Micro-Fine

Subsidy Fully Bran	
(Manufacturer's Price) Subsidised Gene \$ Per ✔ Man	
\$ Per ♥ Man	ufacturer
INSULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE - Maximum of 200 dev per prescription	
★ Syringe 0.3 ml with 29 g x 12.7 mm needle13.00 100 ✓ B-D Ult	tra Fine
1.30 10	
(1.99) B-D Ult	ra Fine
★ Syringe 0.3 ml with 31 g × 8 mm needle13.00 100 ✓ B-D Ult	tra Fine II
1.30 10	
(1.99) B-D Ult	tra Fine II
※ Syringe 0.5 ml with 29 g × 12.7 mm needle13.00 100 ✓ B-D Ult	tra Fine
1.30 10	
(1.99) B-D Ult	ra Fine
※ Syringe 0.5 ml with 31 g × 8 mm needle13.00 100 ✓ B-D Ult	tra Fine II
1.30 10	
(1.99) B-D Ult	tra Fine II
※ Syringe 1 ml with 29 g × 12.7 mm needle13.00 100 ✓ B-D Ult	tra Fine
1.30 10	
(1.99) B-D Ult	ra Fine
※ Syringe 1 ml with 31 g × 8 mm needle	tra Fine II
1.30 10	
(1.99) B-D Ult	tra Fine II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription
- c) Maximum of 1 insulin pump per patient each four year period.

Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 640G
Min basal rate 0.1 U/h	4.500.00	1	✓ Tandem t:slim X2

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Fither:

continued...

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
(Waitulatule 31 Noe)	Per	oubsidised ✓	Manufacturer	

continued...

- 4.1 Applicant is a relevant specialist; or
- 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — **(HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Fither:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Insulin Pump Consumables

⇒SA1604 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Fither:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist: or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — **(severe unexplained hypoglycaemia)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 Applicant is a relevant specialist; or

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (**Previous use before 1 September 2012**) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less

continued...

	ALIMENTARY	INACIAN	ID WEI ADOLISH
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per 🗸	
continued			
than 80 mmol/mol; and			
2 The patient's HbA1c has not deteriorated more than 5 mm			
3 The patient has not had an increase in severe unexplaine 4 Either:	d hypoglycaemic epis	odes from base	eline; and
4.1 Applicant is a relevant specialist; or			
4.1 Applicant is a relevant specialist, of 4.2 Applicant is a nurse practitioner working within the	ir vocational scope		
11 1	•		
INSULIN PUMP CARTRIDGE – Special Authority see SA1604 c	on page 17 – Retail pr	armacy	
a) Maximum of 3 sets per prescription b) Only on a prescription			
c) Maximum of 13 packs of cartridge sets will be funded per	r vear.		
Cartridge 300 U, t:lock × 10		1 OP 🗸	Tandem Cartridge
INSULIN PUMP INFUSION SET (STEEL CANNULA) - Special		on page 17 – F	Retail pharmacy
a) Maximum of 3 sets per prescription	ridinolity 500 Ortioo i	on page 17	iotali priarritacy
b) Only on a prescription			
c) Maximum of 13 infusion sets will be funded per year.			
10 mm steel needle; 29 G; manual insertion; 60 cm tubing \times			
10 with 10 needles	130.00	1 OP 🗸	Paradigm Sure-T
40			MMT-884
10 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock	100.00	100 ./	Sure-T MMT-883
10 mm steel needle; 29 G; manual insertion; 80 cm tubing x		1 OP 🗸	Sure-1 WIWI1-003
10 with 10 needles		1 OP 🗸	Paradigm Sure-T
10 110 100 100 100 100 100 100 100 100			MMT-886
10 mm steel needle; 29 G; manual insertion; 80 cm tubing ×			
10 with 10 needles; luer lock	130.00	1 OP 🗸	Sure-T MMT-885
6 mm steel needle; 29 G; manual insertion; 60 cm tubing \times			
10 with 10 needles	130.00	1 OP 🗸	Paradigm Sure-T
			MMT-864
6 mm steel needle; 29 G; manual insertion; 60 cm tubing x	100.00	100 ./	Cure T MMT 000
10 with 10 needles; luer lock	130.00	1 OP ✓	Sure-T MMT-863
10 with 10 needles	130.00	1 OP 🗸	Paradigm Sure-T
TO WILL TO HOUGHOU		101	MMT-866
6 mm steel needle; 29 G; manual insertion; 80 cm tubing ×			
10 with 10 needles; luer lock	130.00	1 OP 🗸	Sure-T MMT-865
8 mm steel needle; 29 G; manual insertion; 60 cm tubing x			
10 with 10 needles	130.00	1 OP 🗸	Paradigm Sure-T
			MMT-874
8 mm steel needle; 29 G; manual insertion; 60 cm tubing x	100.00	1 OD - 1	C T MAT 070
10 with 10 needles; luer lock	130.00	1 OP 🗸	Sure-T MMT-873
8 mm steel needle; 29 G; manual insertion; 80 cm tubing \times			

✓ Paradigm Sure-T

MMT-876

✓ Sure-T MMT-875

1 OP

1 OP

8 mm steel needle; 29 G; manual insertion; 80 cm tubing \times

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
 \$	Per	✓	Manufacturer	

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) – Special Authority see SA1604 on page 17 – Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm steel cannula; straight insertion; 60 cm line × 10 with			
10 needles	130.00	1 OP	✓ TruSteel
6 mm steel cannula; straight insertion; 81 cm line × 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 81 cm line × 10 with	130.00	1 OP	✓ TruSteel

INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION WITH INSERTION DEVICE) - Special Authority see SA1604 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

1 OP

✓ AutoSoft 30

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	/	Manufacturer	

INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION) - Special Authority see SA1604 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription

c) Maximum of 13 infusion sets will be funded per year. 13 mm teflon cannula; angle insertion; 120 cm line × 10 with			
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-382
13 mm teflon cannula; angle insertion; 45 cm line × 10 with 10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-368
13 mm teflon cannula; angle insertion; 60 cm line x 10 with 10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-381
13 mm teflon cannula; angle insertion; 80 cm line x 10 with 10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-383
17 mm teflon cannula; angle insertion; 110 cm line x 10 with 10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-377
17 mm teflon cannula; angle insertion; 110 cm line x 10 with 10 needles; luer lock	130.00	1 OP	✓ Silhouette MMT-371
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-378
17 mm teflon cannula; angle insertion; 60 cm line × 10 with 10 needles; luer lock	130.00	1 OP	✓ Silhouette MMT-373
10 needles	130.00	1 OP	✓ Paradigm Silhouette

MMT-384

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
\$	Per	/	Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION WITH INSERTION DEVICE) - Special Authority see SA1604 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription

b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; insertion device; 45 cm		
blue tubing × 10 with 10 needles	1 OP	✓ Paradigm Mio MMT-941
6 mm teflon cannula; straight insertion; insertion device; 45 cm pink tubing × 10 with 10 needles	1 OP	✓ Paradigm Mio MMT-921
6 mm teflon cannula; straight insertion; insertion device; 60 cm blue tubing × 10 with 10 needles130.00	1 OP	✓ Paradigm Mio MMT-943
6 mm teflon cannula; straight insertion; insertion device; 60 cm pink tubing × 10 with 10 needles130.00	1 OP	✓ Paradigm Mio MMT-923
6 mm teflon cannula; straight insertion; insertion device; 80 cm blue tubing × 10 with 10 needles130.00	1 OP	✓ Paradigm Mio MMT-945
6 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing × 10 with 10 needles130.00	1 OP	✓ Paradigm Mio MMT-965
6 mm teflon cannula; straight insertion; insertion device; 80 cm pink tubing × 10 with 10 needles	1 OP	✓ Paradigm Mio MMT-925
9 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing × 10 with 10 needles130.00	1 OP	✓ Paradigm Mio MMT-975
6 mm teflon cannula; straight insertion; insertion device; 110 cm line × 10 with 10 needles140.00	1 OP	✓ AutoSoft 90
6 mm teflon cannula; straight insertion; insertion device; 60 cm line × 10 with 10 needles140.00	1 OP	✓ AutoSoft 90
9 mm teflon cannula; straight insertion; insertion device;		

110 cm line × 10 with 10 needles140.00

line × 10 with 10 needles......140.00

9 mm teflon cannula; straight insertion; insertion device; 60 cm

1 OP

1 OP

✓ AutoSoft 90

✓ AutoSoft 90

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Manufacturer INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION) - Special Authority see SA1604 on page 17 -Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula: straight insertion: 110 cm tubing × 10 with 1 OP ✓ Paradigm Quick-Set MMT-398 6 mm teflon cannula: straight insertion: 110 cm tubing × 10 with 1 OP ✓ Quick-Set MMT-391 6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 1 OP ✓ Paradigm Quick-Set MMT-399 6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 1 OP ✓ Quick-Set MMT-393 6 mm teflon cannula; straight insertion; 80 cm tubing × 10 with 1 OP ✓ Paradigm Quick-Set MMT-387 9 mm teflon cannula; straight insertion; 106 cm tubing × 10 with ✓ Paradigm Quick-Set 1 OP MMT-396 9 mm teflon cannula; straight insertion; 110 cm tubing × 10 with ✓ Quick-Set MMT-390 1 OP 9 mm teflon cannula: straight insertion: 60 cm tubing × 10 with 1 OP ✓ Paradigm Quick-Set MMT-397 9 mm teflon cannula: straight insertion: 60 cm tubing × 10 with 1 OP ✓ Quick-Set MMT-392 9 mm teflon cannula; straight insertion; 80 cm tubing × 10 with 1 OP ✓ Paradigm Quick-Set MMT-386 INSULIN PUMP RESERVOIR - Special Authority see SA1604 on page 17 - Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of reservoir sets will be funded per year. 10 × luer lock conversion cartridges 1.8 ml for Paradigm pumps......50.00 1 OP ✓ ADR Cartridge 1.8 Cartridge for 5 and 7 series pump; 1.8 ml × 1050.00 1 OP Paradigm 1.8 Reservoir Cartridge for 7 series pump; 3.0 ml × 1050.00 1 OP ✓ Paradigm 3.0 Reservoir **Digestives Including Enzymes** PANCREATIC ENZYME Cap pancreatin 150 mg (amylase 8.000 Ph Eur U. lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)34.93 100 Creon 10000 Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase, 100 **Panzytrat**

Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase

25.000 Ph Eur U, total protease 1.000 Ph Eur U)94.38

✓ Creon 25000

100

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic	
	\$	Per	✓	Manufacturer	
URSODEOXYCHOLIC ACID - Special Authority see SA1739 be	olow – Retail pharmac	у			
Cap 250 mg	37.95	100	√ U	rsosan	

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner.

Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults: and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Fully

Brand or

Subsidy

	(Manufacturer's P	rice) Subs	idised Generic Manufacturer
Laxatives			
Bulk-forming Agents			
ISPAGHULA (PSYLLIUM) HUSK – Only on a prescription * Powder for oral soln MUCILAGINOUS LAXATIVES WITH STIMULANTS * Dry		500 g OP 500 g OP	✓ Konsyl-D
	(17.32) 2.41 (8.72)	200 g OP	Normacol Plus Normacol Plus
Faecal Softeners			
DOCUSATE SODIUM - Only on a prescription * Tab 50 mg * Tab 120 mg DOCUSATE SODIUM WITH SENNOSIDES	3.13	100 100	✓ <u>Coloxyl</u> ✓ <u>Coloxyl</u>
Tab 50 mg with sennosides 8 mg POLOXAMER — Only on a prescription Not funded for use in the ear. Oral drops 10%		200 30 ml OP	✓ <u>Laxsol</u> ✓ Coloxyl
Opioid Receptor Antagonists - Peripheral			
METHYLNALTREXONE BROMIDE – Special Authority see SA1 Inj 12 mg per 0.6 ml vial		ail pharmacy 1 7	✓ Relistor ✓ Relistor
Initial application — (Opioid induced constipation) from any unless notified for applications meeting the following criteria: Both:	relevant practitio	ner. Approvals	s valid without further renewal
 The patient is receiving palliative care; and Either: Oral and rectal treatments for opioid induced cons Oral and rectal treatments for opioid induced cons 	•		ed.
Osmotic Laxatives			
GLYCEROL * Suppos 3.6 g – Only on a prescription LACTULOSE – Only on a prescription	9.25	20	✓ <u>PSM</u>
* Oral liq 10 g per 15 ml		500 ml	✓ <u>Laevolac</u>
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BI Powder for oral soln 13.125 g with potassium chloride 46.6 r sodium bicarbonate 178.5 mg and sodium chloride 350.	ng,	ND SODIUM C	Molaxole

SODIUM ACID PHOSPHATE - Only on a prescription

✓ Fleet Phosphate Enema

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml.	, , ,	otion		
5 ml	29.98	50	✓ <u>N</u>	<u>licolette</u>
Stimulant Laxatives				
BISACODYL - Only on a prescription				
* Tab 5 mg		200	_	ax-Tab
* Suppos 10 mg	3.74	10	✓ [ax-Suppositories
SENNA - Only on a prescription				
* Tab, standardised	2.17	100		
	(6.84)		S	Senokot
	0.43	20		
	(1.72)		S	Senokot

Metabolic Disorder Agents

ALGLUCOSIDASE ALFA — Special Authority se	ee SA1622 below – Retail pharmacy		
Inj 50 mg vial	1,142.60	1	Myozyme

⇒SA1622 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides: or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT): and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates: and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT: and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation: and
- 7 There is no evidence of new or progressive cardiomyopathy.

✓ Cystadane

180 g OP

	Subsidy (Manufacturer's Price)		Fully	Brand or Generic	
	\$	Per		Manufacturer	
BETAINE - Special Authority see SA1727 below - Retail pharma	ncy				

⇒SA1727 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism: and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE - Special Authority see SA1593 below - Retail pharmacy ✓ Naglazyme

⇒SA1593 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Fither:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to FRT.

IDURSULFASE - Special Authority see SA1623 below - Retail pharmacy ✓ Elaprase

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assav in cultured skin fibroblasts: or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT): and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer	
LARONIDASE – Special Authority see SA1695 below – Retail p	,	1	✓ A	ldurazvme	

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Fither:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

SAPROPTERIN DIHYDROCHLORIDE – Special Authority see SA1757 below – Retail pharmacy
Tab soluble 100 mg.......1,452.70 30 OP
✓ Kuvan

⇒SA1757 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy: or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Fully

Subsidy (Manufacturer's Price) \$ Price

Subsidised Per

Brand or Generic Manufacturer

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

174 g OP ✓ Pheburane

⇒SA1598 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA1734 below - Retail pharmacy

⇒SA1734 Special Authority for Subsidy

Special Authority approved by the Gaucher Treatment Panel

Notes: Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The Co-ordinator, Gaucher Treatment Panel Phone: 04 460 4990 PHARMAC PO Box 10 254 Facsimile: 04 916 7571

Wellington Email: gaucherpanel@pharmac.govt.nz

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Access Criteria

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2) Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and
- Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, serum glucosylsphingosine, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
 - Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia, thrombocytopenia; at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or massive symptomatic splenomegaly; or
 - 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g. hips or shoulder); spontaneous fractures or vertebral collapse; chronic bone pain not controlled by other pharmaceuticals; or
 - 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4) Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher
- 5) Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1) Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated; and
- 2) Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3) Radiological (MRI) signs of bone activity performed at one year and two years since initiation of treatment begins, and two to three yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4) Serum glucosylsphingosine levels taken at least 6 to 12 monthly show a decrease compared with baseline; and
- 5) Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 6) Patient has not developed another medical condition that might reasonably be expected to compromise a response to
- 7) Patient is compliant with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- 8) Supporting clinical information including test reports, MRI whole body STIR, serum glucosylsphingosine, haematological data, and other relevant investigations are submitted to the Gaucher Panel for assessment as required.

Mouth and Throat

Agents Used in Mouth Ulceration

BENZYDAMINE HYDROCHI ORIDE	

00 01.1070	g			
Endorser	nent9.0	0	500 ml	
	(20.3	1)		Difflam

Additional subsidy by endorsement for a patient who has oral mucositis as a result of treatment for cancer, and the prescription is endorsed accordingly.

CARMELLOSE SODIUM WITH GELATIN AND PECTIN

Soln 0.15% - Higher subsidy of \$20.31 per 500 ml with

Paste	17.20	56 g OP	Stomahesive
	4.55	15 g OP	
	(7.90)	-	Orabase
	1.52	5 g OP	
	(3.60)	_	Orabase
Powder	8.48	28 g OP	
	(10.95)	_	Stomahesive
CHLORHEXIDINE GLUCONATE			
Mouthwash 0.2%	2.57	200 ml OP	✓ healthE
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE			
* Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP	
-	(6.00)	_	Bonjela
TRIAMCINOLONE ACETONIDE			
Paste 0.1%	5.33	5 g OP	✓ Kenalog in Orabase

	Subsidy (Manufacturer's Pr	rice) Subs	Fully Brand or sidised Generic
	\$	Per	✓ Manufacturer
Oropharyngeal Anti-infectives			
AMPHOTERICIN B Lozenges 10 mg	5.86	20	✓ Fungilin
MICONAZOLE Oral gel 20 mg per g	4.74	40 g OP	✓ <u>Decozol</u>
NYSTATIN Oral liq 100,000 u per ml	1.95	24 ml OP	✓ <u>Nilstat</u>
Other Oral Agents			
For folinic mouthwash, pilocarpine oral liquid or saliva substitute HYDROGEN PEROXIDE	e formula refer Star	ndard Formula	e, page 235
★ Soln 3% (10 vol) – Maximum of 200 ml per prescription Pharmacy Health Soln 3% (10 vol) to be delisted 1 July 2020) THYMOL GLYCERIN	1.40	100 ml	✓ Pharmacy Health
* Compound, BPC	9.15	500 ml	✓ PSM
Vitamins			
Vitamin A			
/ITAMIN A WITH VITAMINS D AND C ★ Soln 1000 u with Vitamin D 400 u and ascorbic acid 30 mg 10 drops Vitadol C Soln 1000 u with Vitamin D 400 u and ascorbic acid	4.50	10 ml OP s to be delisted	✓ Vitadol C d 1 December 2019)
Vitamin B			
HYDROXOCOBALAMIN Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a l PYRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose	PSO1.89	3	✓ <u>Neo-B12</u>
b) Only on a prescription ★ Tab 25 mg – No patient co-payment payable ★ Tab 50 mg		90 500	✓ <u>Vitamin B6 25</u> ✓ <u>Apo-Pyridoxine</u>
THIAMINE HYDROCHLORIDE - Only on a prescription ★ Tab 50 mg	4.89	100	✓ Max Health
/ITAMIN B COMPLEX * Tab, strong, BPC		500	✓ Bplex
Vitamin C			•
ASCORBIC ACID			
a) No more than 100 mg per dose b) Only on a prescription * Tab 100 mg	8 10	500	✓ Cvite
- 145 177 IIIg		550	- 01110

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	(Manufacturer's Price) \$	Subsidi Per	sed •	Generic Manufacturer
Vitamin D				
ALFACALCIDOL * Cap 0.25 mcg * Cap 1 mcg * Oral drops 2 mcg per ml	87.98	100	✓ Or	ne-Alpha ne-Alpha ne-Alpha
CALCITRIOL * Cap 0.25 mcg Cap 0.5 mcg	7.95			alcitriol-AFT alcitriol-AFT
COLECALCIFEROL * Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescripti * Oral liq 188 mcg per ml (7,500 iu per ml)			✓ <u>Vi</u> ✓ Pu	<u></u>

Subsidy

Fully

Brand or

Multivitamin Preparations

MULTIVITAMIN RENAL - Special Authority see SA1546 belo	w – Retail pharmacy		
* Cap	6.49	30	 Clinicians Renal Vit
⇒SA1546 Special Authority for Subsidy			

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or
- 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73 m² body surface area (BSA).

MULTIVITAMINS - Special Authority see SA1036 below - Retail pharmacy 200 a OP ✓ Paediatric Seravit

⇒SA1036 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins.

VITAMINIC

VII	AIVIIVO		
*	Tab (BPC cap strength)10.50	1,000	✓ Mvite
	Cap (fat soluble vitamins A, D, E, K) - Special Authority see		
	SA1720 below – Retail pharmacy	60	✓ Vitabdeck

⇒SA1720 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency; or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Minerals				
Calcium				
CALCIUM CARBONATE * Tab eff 1.75 g (1 g elemental)	28.40	20	√ (Calcium Sandoz (\$29)
* Tab 1.25 g (500 mg elemental)		250		Arrow-Calcium
* Inj 10%, 10 ml ampoule	64.00	20	✓ N	Max Health S29
Fluoride				
SODIUM FLUORIDE * Tab 1.1 mg (0.5 mg elemental)	5.75	100	√ F	PSM
lodine				
POTASSIUM IODATE * Tab 253 mcg (150 mcg elemental iodine)	4.69	90	✓ <u>N</u>	leuroTabs
Iron				
FERRIC CARBOXYMALTOSE – Special Authority see SA1840 Inj 50 mg per ml, 10 ml		acy 1	√ F	Ferinject
■ SA1840 Special Authority for Subsidy Initial application — (serum ferritin less than or equal to 20 months for applications meeting the following criteria: Both:	ncg/L) from any rele	/ant p	oractitioner.	Approvals valid for 3

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease

continued...

				_
Subsidy		Fully	Brand or	_
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	/	Manufacturer	

continued...

and a trial of oral iron is unlikely to be effective; or

2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

FERROUS FUMARATE * Tab 200 mg (65 mg elemental)	3.09	100	✓ Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID * Tab 310 mg (100 mg elemental) with folic acid 350 mcg	4.68	60	✓ <u>Ferro-F-Tabs</u>
FERROUS SULFATE * Oral liq 30 mg (6 mg elemental) per 1 ml	12.08	500 ml	✓ Ferodan
FERROUS SULPHATE * Tab long-acting 325 mg (105 mg elemental)	2.06	30	✓ Ferrograd
IRON POLYMALTOSE			
* Inj 50 mg per ml, 2 ml ampoule	15.22	5	✓ Ferrum H
	34.50		✓ Ferrosig
(Ferrum H Inj 50 mg per ml, 2 ml ampoule to be delisted 1 February 202	20)		

Magnesium

For magnesium hydroxide mixture refer Standard Formulae, page 235

Zincaps to be Sole Supply on 1 December 2019

//AGNESIUM HYD	ROXIDE	
----------------	--------	--

Suspension 8%	72.20	500 ml	✓ T&R S29
MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule	10.21	10	✓ <u>DBL</u> ✓ DBI \$29 \$29

Zinc

ZIN	IC SULPHATE			
*	Cap 137.4 mg (50 mg elemental)11.00	100	✓ Zincaps

BLOOD AND BLOOD FORMING ORGANS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — **(chronic renal failure)** from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus: and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus: and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*: and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 ju per week.

Note: Indication marked with * is an unapproved indication

BLOOD AND BLOOD FORMING ORGANS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
EPOETIN ALFA – Special Authority see SA1775 on the previous	page - Retail pharm	асу		
Wastage claimable	250.00	6	1	Binocrit
Inj 1,000 iu in 0.5 ml, syringe Inj 2,000 iu in 1 ml, syringe		6	_	Binocrit
Inj 3,000 iu in 0.3 ml, syringe		6		Binocrit
Inj 4,000 iu in 0.4 ml, syringe		6		Binocrit
Inj 5,000 iu in 0.5 ml, syringe		6	✓	Binocrit
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	✓	Binocrit
Inj 8,000 iu in 0.8 ml, syringe	175.00	6	✓	Binocrit
Inj 10,000 iu in 1 ml, syringe	197.50	6	✓	Binocrit
Inj 40,000 iu in 1 ml, syringe	250.00	1	✓	Binocrit

Megaloblastic

	10	40	ī
FOL	JU.	ΑU	ıU

*	Tab 0.8 mg21.84	1,000	•	Apo-Folic Acid
*	Tab 5 mg12.12	500	•	Apo-Folic Acid
	Oral lig 50 mcg per ml26.00	25 ml OP	•	Biomed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial	2,450.00	1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	Alprolix
ELTROMBOPAG – Special Authority see SA1743 Wastage claimable	below - Retail pharmacy		

Tab 50 mg3,100.00 **➤ SA1743** Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab);
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding: or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (**idiopathic thrombocytopenic purpura - preparation for splenectomy**) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist.

continued...

✓ Revolade

✓ Revolade

28 28

Subsidy (Manufacturer's	Price)	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Fither:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe	2,356.60	1	✓ NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓ NovoSeven RT
Ini 8 ma syringe	9.426.40	1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	0 1	✓ FEIBA NF
Inj 1,000 U2,630.0	0 1	✓ FEIBA NF
Inj 2,500 U6,575.0	0 1	✓ FEIBA NF

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] – [X For patients with haemophilia. Rare Clinical Circumstan treatment is managed by the Haemophilia Treaters Grou	ices Brand of short half-life			
subject to criteria.	,			
Inj 250 iu prefilled syringe	287.50	1	✓	Xyntha
Inj 500 iu prefilled syringe	575.00	1	✓	Xyntha
Inj 1,000 iu prefilled syringe	1,150.00	1	✓	Xyntha
Inj 2,000 iu prefilled syringe	2,300.00	1	✓	Xyntha
Inj 3,000 iu prefilled syringe	3,450.00	1	✓	Xyntha
NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xph	arm]			
For patients with haemophilia. Access to funded treatme	•	emop	hilia Treat	ers Group in conjunction
with the National Haemophilia Management Group.	3.1.1, 1.7.1	- 6		, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
Inj 500 iu vial	435.00	1	1	RIXUBIS
Inj 1,000 iu vial	870.00	1	✓	RIXUBIS
Inj 2,000 iu vial	1,740.00	1	✓	RIXUBIS
Inj 3,000 iu vial	2,610.00	1	✓	RIXUBIS
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE	-) _ [Xnharm]			
For patients with haemophilia. Preferred Brand of short		or VIII	Access to	funded treatment is
managed by the Haemophilia Treaters Group in conjunc				
Inj 250 iu vial		1		Advate
Ini 500 iu vial.		1		Advate
Inj 1,000 iu vial		1		Advate
Inj 1,500 iu vial		1		Advate
Inj 1,300 iu vial	·	1		Advate
Inj 3,000 iu vial	*	1	_	Advate
•	•		•	Auvaic
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGEN				
For patients with haemophilia. Rare Clinical Circumstan				
treatment is managed by the Haemophilia Treaters Grou	ip in conjunction with the I	Natior	nal Haemo	philia Management Group,
subject to criteria.				
Inj 250 iu vial		1		Kogenate FS
Inj 500 iu vial		1		Kogenate FS
Inj 1,000 iu vial		1		Kogenate FS
Inj 2,000 iu vial	,	1		Kogenate FS
Inj 3,000 iu vial	2,850.00	1	•	Kogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR				
For patients with haemophilia A receiving prophylaxis tre	eatment. Access to funder	d trea	tment is m	nanaged by the Haemophilia
Treaters Group in conjunction with the National Haemop	hilia Management group.			
Inj 250 iu vial	300.00	1	✓	Adynovate
Inj 500 iu vial	600.00	1	✓	Adynovate
lnj 1,000 iu vial	1,200.00	1	1	Adynovate
Inj 2,000 iu vial	2,400.00	1	✓	Adynovate
SODIUM TETRADECYL SULPHATE				
	20 50	5		
	Z0.:0U			
		·		Fibro-vein
* Inj 3% 2 ml	(73.00)	ŭ		Fibro-vein
* Inj 3% 2 ml TRANEXAMIC ACID Tab 500 mg	(73.00)	100	.1	Fibro-vein Cyklokapron

✓ Effient

	Subsidy (Manufacturer's Price)	Sub Per	Fully sidised	Brand or Generic Manufacturer
Vitamin K				
PHYTOMENADIONE Inj 2 mg per 0.2 ml — Up to 5 inj available on a PSO Inj 10 mg per ml, 1 ml — Up to 5 inj available on a PSO		5 5	- 1	onakion MM onakion MM
Antithrombotic Agents				
Antiplatelet Agents				
ASPIRIN * Tab 100 mg	10.80	990	√ <u>E</u> 1	thics Aspirin EC
* Tab 75 mg	5.44	84	✓ A	rrow - Clopid
DIPYRIDAMOLE * Tab long-acting 150 mg		60	✓ <u>P</u> y	ytazen SR
PRASUGREL – Special Authority see SA1201 below – Retail ph Tab 5 mg		28	✓ Ef	ffient

⇒SA1201 Special Authority for Subsidy

Initial application — (coronary angioplasty and bare metal stent) from any relevant practitioner. Approvals valid for 6 months where the patient has undergone coronary angioplasty in the previous 4 weeks and is clopidogrel-allergic*.

Initial application — (drug eluting stent) from any relevant practitioner. Approvals valid for 12 months where the patient has had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Initial application — (stent thromobosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (coronary angioplasty and bare metal stent) from any relevant practitioner. Approvals valid for 6 months where the patient has undergone coronary angioplasty or had a bare metal cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Renewal — (drug eluting stent) from any relevant practitioner. Approvals valid for 12 months where had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Note: * Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

⇒SA1382 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

 Subsidy		Fully	Brand or	
(Manufacturer's Price)	;	Subsidised	Generic	
\$	Per	✓	Manufacturer	

Heparin and Antagonist Preparations

DALTEPARIN SODIUM - Special Authority see SA1270 bel	ow – Retail pharmacy		
Inj 2,500 iu per 0.2 ml prefilled syringe	19.97	10	✓ Fragmin
Inj 5,000 iu per 0.2 ml prefilled syringe	39.94	10	✓ Fragmin
Inj 7,500 iu per 0.75 ml graduated syringe		10	✓ Fragmin
Inj 10,000 iu per 1 ml graduated syringe		10	✓ Fragmin
Inj 12,500 iu per 0.5 ml prefilled syringe		10	✓ Fragmin
Inj 15,000 iu per 0.6 ml prefilled syringe	120.05	10	✓ Fragmin
Inj 18,000 iu per 0.72 ml prefilled syringe	158.47	10	✓ Fragmin

(Fragmin Inj 2,500 iu per 0.2 ml prefilled syringe to be delisted 1 April 2020)
(Fragmin Inj 5,000 iu per 0.2 ml prefilled syringe to be delisted 1 April 2020)
(Fragmin Inj 7,500 iu per 0.75 ml graduated syringe to be delisted 1 April 2020)
(Fragmin Inj 10,000 iu per 1 ml graduated syringe to be delisted 1 April 2020)
(Fragmin Inj 12,500 iu per 0.5 ml prefilled syringe to be delisted 1 January 2020)
(Fragmin Inj 15,000 iu per 0.6 ml prefilled syringe to be delisted 1 January 2020)
(Fragmin Inj 18,000 iu per 0.72 ml prefilled syringe to be delisted 1 January 2020)

⇒SA1270 Special Authority for Subsidy

Initial application — (Pregnancy or Malignancy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (**Pregnancy or Malignancy**) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Fither:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, Acute Coronary Syndrome, cardioversion, or prior to oral anti-coagulation).

ENOXAPARIN SODIUM - Special Authority see SA1646 on the next page - Retail pharmacy

Inj 20 mg in 0.2 ml syringe	.27.93	10	✓ Clexane
Inj 40 mg in 0.4 ml syringe		10	✓ Clexane
Inj 60 mg in 0.6 ml syringe	.56.18	10	✓ Clexane
Inj 80 mg in 0.8 ml syringe		10	✓ Clexane
Inj 100 mg in 1 ml syringe		10	✓ Clexane
Inj 120 mg in 0.8 ml syringe		10	✓ Clexane
Inj 150 mg in 1 ml syringe	133.20	10	✓ Clexane

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1646 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

HEPARIN SODIUM

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

Inj 1,000 iu per ml, 5 ml ampoule58.57	50	✓ Pfizer
Inj 5,000 iu per ml, 1 ml28.40	5	✓ Hospira
		✓ Pfizer
Inj 5,000 iu per ml, 5 ml ampoule203.68	50	✓ Pfizer
Inj 25,000 iu per ml, 0.2 ml19.00	5	✓ Hospira
122.00	10	✓ Wockhardt S29
190.00	50	✓ Pfizer \$29
HEPARINISED SALINE		
Inj 10 iu per ml, 5 ml56.94	50	✓ Pfizer
11) 10 to por 111, 0 111		- 1 11201
Oral Anticoagulants		
DABIGATRAN		
Cap 75 mg - No more than 2 cap per day76.36	60	✓ Pradaxa
Cap 110 mg76.36	60	✓ Pradaxa
Cap 150 mg76.36	60	✓ Pradaxa
RIVAROXABAN		
Tab 10 mg - No more than 1 tab per day83.10	30	✓ Xarelto
Tab 15 mg - Up to 14 tab available on a PSO77.56	28	✓ Xarelto
Tab 20 mg77.56	28	✓ Xarelto

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	✓	Manufacturer
VARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
★ Tab 1 mg	3.46	50	✓ (Coumadin
-	7.60	100	√	Marevan
★ Tab 2 mg	4.31	50	✓ (Coumadin
★ Tab 3 mg	11.80	100	✓ I	Marevan
★ Tab 5 mg	5.93	50	✓ (Coumadin
	13.50	100	✓	Marevan

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10⁹/L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1384 below - Retail pharmacy

⇒SA1384 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*). Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration

GLUCOSE [DEXTROSE]		
* Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO29.50	5	✓ Biomed
* Inj 50%, 90 ml bottle - Up to 5 inj available on a PSO14.50	1	✓ Biomed
POTASSIUM CHLORIDE		
* Inj 75 mg per ml, 10 ml55.00	50	✓ AstraZeneca

	Subsidy	, -	Fully	
	(Manufacturer's Pric \$	e) S Per	Subsidised •	
SODIUM BICARBONATE	-			
Inj 8.4%, 50 ml	19.95	1	/	Biomed
a) Up to 5 inj available on a PSO		•		2.00
b) Not in combination				
Inj 8.4%, 100 ml	20.50	1	1	Biomed
a) Up to 5 inj available on a PSO				
b) Not in combination				
SODIUM CHLORIDE				
Not funded for use as a nasal drop. Not funded for nebuliser for nebuliser use.	·		·	
Inj 0.9%, bag – Up to 2000 ml available on a PSO		500 ml	_	Baxter
Only if a second had an a successible of a second district make	1.26	1,000 m		Baxter
Only if prescribed on a prescription for renal dialysis, mat for emergency use. (500 ml and 1,000 ml packs)				
Inj 23.4% (4 mmol/ml), 20 ml ampoule		5	•	Biomed
For Sodium chloride oral liquid formulation refer Standard			./	Fresenius Kabi
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO	7.00	20 50		InterPharma
	7.00	30		Multichem
Fresenius Kabi to be Sole Supply on 1 December 2019				
Inj 0.9%, 10 ml ampoule - Up to 5 inj available on a PSO	5.40	50	1	Fresenius Kabi
	6.63		✓	Pfizer
Fresenius Kabi to be Sole Supply on 1 December 2019				
Inj 0.9%, 20 ml ampoule	5.00	20		Fresenius Kabi
	7.50	30		Multichem InterPharma
Fresenius Kabi to be Sole Supply on 1 December 2019	7.50	30	•	IIILEI FII AIIII A
(InterPharma Inj 0.9%, 5 ml ampoule to be delisted 1 December 2	019)			
(Multichem Inj 0.9%, 5 ml ampoule to be delisted 1 December 201				
(Pfizer Inj 0.9%, 10 ml ampoule to be delisted 1 December 2019)	,			
(Multichem Inj 0.9%, 20 ml ampoule to be delisted 1 December 20				
(InterPharma Inj 0.9%, 20 ml ampoule to be delisted 1 December	*			
TOTAL PARENTERAL NUTRITION (TPN) - Retail pharmacy-Sp			_	
Infusion	CBS	1 OP	•	TPN
WATER				
1) On a prescription or Practitioner's Supply Order only wh	en on the same fo	rm as an	injection	listed in the Pharmaceutica
Schedule requiring a solvent or diluent; or				
2) On a bulk supply order; or	dua			
3) When used in the extemporaneous compounding of eye4) When used for the dilution of sodium chloride soln 7% for		ationte on	ılv	
To vitigit used for the dilution of soutuin chilolide Soill 7 % ic	n cysuc iibiosis pa	au c iilo Ul	ııy.	
Inj 5 ml ampoule - Up to 5 inj available on a PSO	7.00	50	1	InterPharma
Inj 10 ml ampoule – Up to 5 inj available on a PSO		50		Pfizer
Inj 20 ml ampoule - Up to 5 inj available on a PSO		20		Fresenius Kabi
				Multichem
	7.50	30	_	InterPharma
Oral Administration				
CALCIUM POLYSTYRENE SULPHONATE				
Powder	169.85	300 g OF	•	Calcium Resonium

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Pr \$		Fully Brand or lised Generic Manufacturer
COMPOUND ELECTROLYTES Powder for oral soln — Up to 10 sach available on a PSO (Enerlyte Powder for oral soln to be delisted 1 April 2020)	2.30 9.77	10 50	✓ Enerlyte ✓ Electral
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml)	6.55	1,000 ml OP	✓ <u>Pedialyte -</u> <u>Bubblegum</u>
PHOSPHORUS Tab eff 500 mg (16 mmol) POTASSIUM CHLORIDE	82.50	100	✓ Phosphate Phebra
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	5.26 (11.85)	60	Chlorvescent
* Tab long-acting 600 mg (8 mmol)	8.90	200	✓ <u>Span-K</u>
Cap 840 mg	8.52	100	✓ Sodibic✓ Sodibic
SODIUM POLYSTYRENE SULPHONATE Powder	84.65	454 g OP	✓ Resonium-A

S29 Unapproved medicine supplied under Section 29

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Alpha-Adrenoceptor Blockers

Alpha Adrenoceptor Blockers

DOXAZOSIN * Tab 2 mg	500 500	✓ <u>Apo-Doxazosin</u> ✓ <u>Apo-Doxazosin</u>
* Cap 10 mg	30	✓ BNM S29
216.67	100	✓ Dibenzyline S29
PRAZOSIN		•
* Tab 1 mg5.53	100	✓ Apo-Prazosin
* Tab 2 mg7.00	100	✓ Apo-Prazosin
* Tab 5 mg11.70	100	✓ Apo-Prazosin
TERAZOSIN		
* Tab 1 mg	28	✓ Actavis
* Tab 2 mg	500	✓ Apo-Terazosin
* Tab 5 mg	500	✓ Apo-Terazosin

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

CAPTOPRIL * Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of age.	94.99	95 ml OP	✓ Capoten
CILAZAPRIL			
* Tab 0.5 mg	2.09	90	✓ Zapril
* Tab 2.5 mg	4.80	90	✓ Zapril
· ·	7.20	200	✓ Apo-Cilazapril
Zapril to be Sole Supply on 1 February 2020			
* Tab 5 mg	8.35	90	✓ Zapril
·	12.00	200	✓ Apo-Cilazapril
Zapril to be Sole Supply on 1 February 2020		_00	
(Apo-Cilazapril Tab 2.5 mg to be delisted 1 February 2020)			
(Apo-Cilazapril Tab 5 mg to be delisted 1 February 2020)			
ENALAPRIL MALEATE			
* Tab 5 mg	3.84	100	 Ethics Enalapril
* Tab 10 mg		100	 Ethics Enalapril
* Tab 20 mg		100	✓ Ethics Enalapril
-			
LISINOPRIL			4 -
* Tab 5 mg	2.07	90	✓ Ethics Lisinopril
* Tab 10 mg	2.36	90	 Ethics Lisinopril
* Tab 20 mg	3.17	90	 Ethics Lisinopril
PERINDOPRIL			
	3.75	30	✓ Apo-Perindopril
1 22 = 19			. — —
* Tab 4 mg	4.80	30	✓ Apo-Perindopril

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
QUINAPRIL				
₭ Tab 5 mg	6.01	90		Arrow-Quinapril 5
★ Tab 10 mg		90		Arrow-Quinapril 10
≰ Tab 20 mg	4.89	90	•	Arrow-Quinapril 20
ACE Inhibitors with Diuretics				
CILAZAPRIL WITH HYDROCHLOROTHIAZIDE				
* Tab 5 mg with hydrochlorothiazide 12.5 mg	10.18	100	✓	Apo-Cilazapril/ Hydrochlorothiazide
QUINAPRIL WITH HYDROCHLOROTHIAZIDE				
★ Tab 10 mg with hydrochlorothiazide 12.5 mg	3.83	30	1	Accuretic 10
Fab 20 mg with hydrochlorothiazide 12.5 mg	4.92	30	•	Accuretic 20
Angiotensin II Antagonists				
ANDESARTAN CILEXETIL				
★ Tab 4 mg	1.90	90	✓	Candestar
★ Tab 8 mg	2.28	90		Candestar
₭ Tab 16 mg	3.67	90		Candestar
₹ Tab 32 mg	6.39	90	✓	<u>Candestar</u>
OSARTAN POTASSIUM				
₭ Tab 12.5 mg	1.39	84		Losartan Actavis
₹ Tab 25 mg	1.63	84		Losartan Actavis
★ Tab 50 mg		84		Losartan Actavis
₭ Tab 100 mg	2.31	84	•	Losartan Actavis
Angiotensin II Antagonists with Diuretics				
OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE				
Tab 50 mg with hydrochlorothiazide 12.5 mg	1.88	30	/	Arrow-Losartan & Hydrochlorothiazide

Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN - Special Authority see SA1751 below - Retail pharmacy

Note: Due to the angiotensin II receptor blocking activity of sacubitril with valsartan it should not be co-administered with an ACE inhibitor or another ARB.

ACE ITTIBILOT OF AITOLITIC AT ID.			
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/103

⇒SA1751 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or
 - 2.2 Patient is in NYHA/WHO functional class III: or

continued...

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic
 \$	Per	•	Manufacturer

continued...

- 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local,	page 118	
AMIODARONE HYDROCHLORIDE		
▲ Tab 100 mg − Retail pharmacy-Specialist3.80	30	✓ Aratac
4.66		Cordarone-X
Aratac to be Sole Supply on 1 December 2019		
▲ Tab 200 mg - Retail pharmacy-Specialist5.25	30	✓ Aratac
7.63		✓ Cordarone-X
Aratac to be Sole Supply on 1 December 2019		
Inj 50 mg per ml, 3 ml ampoule – Up to 6 inj available on a PSO9.98	5	✓ Lodi
11.98	6	✓ Cordarone-X
16.37	10	✓ Max Health
(Cordarone-X Tab 200 mg to be delisted 1 December 2019) (Lodi Inj 50 mg per ml, 3 ml ampoule to be delisted 1 February 2020) (Cordarone-X Inj 50 mg per ml, 3 ml ampoule to be delisted 1 February 2020) ATROPINE SULPHATE * Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a		
PSO12.07	10	✓ <u>Martindale</u>
DIGOXIN		
* Tab 62.5 mcg - Up to 30 tab available on a PSO7.00	240	✓ Lanoxin PG
* Tab 250 mcg - Up to 30 tab available on a PSO	240	✓ Lanoxin
* Oral lig 50 mcg per ml	60 ml	✓ Lanoxin
		✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE		
▲ Cap 100 mg23.87	100	✓ Rythmodan

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
ECAINIDE ACETATE - Retail pharmacy-Specialist				
Tab 50 mg	19.95	60	✓	Flecainide BNM
	38.95		•	Tambocor
Flecainide BNM to be Sole Supply on 1 February 2020			_	
Cap long-acting 100 mg		30		Tambocor CR
	39.51	90	•	Flecainide
				Controlled
				Release Teva
Flecainide Controlled Release Teva to be Sole Supply				
Cap long-acting 200 mg	61.06	90	•	Flecainide
				Controlled
	22.72		,	Release Teva
Floorinide Controlled Delegas Toyo to be Cale Consult.	68.78	30	•	Tambocor CR
Flecainide Controlled Release Teva to be Sole Supply		_	./	Tambocor
Inj 10 mg per ml, 15 ml ampoule	52.45	5	•	rambocor
ambocor Pab 30 hig to be delisted 1 Pebruary 2020) Fambocor CR Cap long-acting 100 mg to be delisted 1 Decem. Fambocor CR Cap long-acting 200 mg to be delisted 1 Decem.				
EXILETINE HYDROCHLORIDE				
Cap 150 mg	162.00	100	1	Mexiletine
				Hydrochloride
				USP S29
Cap 250 mg	202.00	100	✓	Mexiletine
				Hydrochloride
				USP S29
ROPAFENONE HYDROCHLORIDE - Retail pharmacy-Speci	alist			
Tab 150 mg		50	/	Rytmonorm
				,
Antihypotensives				
IDODRINE – Special Authority see SA1474 below – Retail ph	•	100	./	Cutuan
Tab 2.5 mg		100		Gutron
Tab 5 mg	/ 9.00	100	•	Gutron

⇒SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Note: Treatment should be started with small doses and titrated upwards as necessary. Hypertension should be avoided, and the usual target is a standing systolic blood pressure of 90 mm Hg.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

ATI	ENOLOL		
*	Tab 50 mg4.26	500	Mylan Atenolol
	Tab 100 mg		✓ Mylan Atenolol
	Oral lig 25 mg per 5 ml		✓ Atenolol AFT
	Restricted to children under 12 years of age		

	Subsidy		Fully	Brand or	
	(Manufacturer's Price) \$	Per	Subsidised		
DICORDO OL FUNADATE	Ψ	rei		Manuacturer	
BISOPROLOL FUMARATE	0.50	00		Descripto	
* Tab 2.5 mg		90		Bosvate Bosvate	
* Tab 10 mg		90		Bosvate Bosvate	
* Tab 10 mg	9.40	90	•	Bosvate	
CARVEDILOL			_		
* Tab 6.25 mg		60		Carvedilol Sandoz	
* Tab 12.5 mg		60		Carvedilol Sandoz	
* Tab 25 mg	2.95	60	•	Carvedilol Sandoz	
CELIPROLOL					
* Tab 200 mg	21.40	180	✓	Celol	
LABETALOL					
Tab 100 mg	11.36	100	/	Hybloc	
145 100 mg				Presolol S29	
Tab 200 mg	20.74	100		Hybloc	
740 200 mg	20.7 ¬	100		Presolol S29	
* In F ma nor ml 20 ml amnoula	E0.06	5	•	Presolol 329	
* Inj 5 mg per ml, 20 ml ampoule		Э		Trandate	
(Highlas Tab 100 mg to be delicted 1 December 2010)	(88.60)			Tranuale	
(Hybloc Tab 100 mg to be delisted 1 December 2019)					
(Hybloc Tab 200 mg to be delisted 1 February 2020)					
METOPROLOL SUCCINATE			_		
* Tab long-acting 23.75 mg		30		Betaloc CR	
* Tab long-acting 47.5 mg		30		Betaloc CR	
* Tab long-acting 95 mg		30		Betaloc CR	
* Tab long-acting 190 mg	3.00	30	/	Betaloc CR	
METOPROLOL TARTRATE					
* Tab 50 mg	5.66	100	✓	Apo-Metoprolol	
* Tab 100 mg	7.55	60	✓	Apo-Metoprolol	
* Tab long-acting 200 mg	23.40	28	✓	Slow-Lopresor	
* Inj 1 mg per ml, 5 ml vial	29.50	5	✓	Metroprolol IV	
				Mylan	
NADOLOL					
* Tab 40 mg	16 69	100	1	Apo-Nadolol	
* Tab 80 mg		100		Apo-Nadolol	
ŭ	20.40	100	•	Apo Nadoloi	
PINDOLOL W. Tab 5 mm	10.00	100		Ana Dindalal	
* Tab 5 mg		100		Apo-Pindolol	
* Tab 10 mg		100		Apo-Pindolol	
* Tab 15 mg	33.31	100	•	Apo-Pindolol	
PROPRANOLOL					
* Tab 10 mg	4.64	100		Apo-Propranolol	
* Tab 40 mg		100		Apo-Propranolol	
* Cap long-acting 160 mg	18.17	100	•	Cardinol LA	
* Oral liq 4 mg per ml - Special Authority see SA1327 belo	w –				
Retail pharmacy	CBS	500 m	nl 🗸	Roxane S29	
m.CA1227 Special Authority for Subsidy					

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only): or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTALOL

VMI UDIDIVIE

	Tab 80 mg		500 100	✓ Mylan ✓ Mylan
TIN	MOLOL			
*	Tab 10 mg	10.55	100	✓ Apo-Timol

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

ΑIV	ILODIPINE		
*	Tab 2.5 mg	100	✓ Apo-Amlodipine
*	Tab 5 mg	250	✓ Apo-Amlodipine
*	Tab 10 mg4.40	250	✓ Apo-Amlodipine
FE	LODIPINE		
*	Tab long-acting 2.5 mg	30	✓ Plendil ER
*	Tab long-acting 5 mg	90	✓ Felo 5 ER
*	Tab long-acting 10 mg4.32	90	✓ Felo 10 ER
NIF	EDIPINE		
*	Tab long-acting 10 mg	60	Adalat 10
			✓ Adefin S29
*	Tab long-acting 20 mg9.59	100	✓ Nyefax Retard
*	Tab long-acting 30 mg3.14	30	✓ Adalat Oros
			Adefin XL
*	Tab long-acting 60 mg5.67	30	✓ Adalat Oros
			✓ Adefin XL

(Adefin XL Tab long-acting 30 mg to be delisted 1 March 2020)

Other Calcium Channel Blockers

DILTIAZEM HYDROCHLORIDE			
* Tab 30 mg	4.60	100	✓ Dilzem
* Tab 60 mg	8.50	100	✓ Dilzem
* Cap long-acting 120 mg	33.42	500	✓ Apo-Diltiazem CD
* Cap long-acting 180 mg	50.05	500	✓ Apo-Diltiazem CD
* Cap long-acting 240 mg	66.76	500	✓ Apo-Diltiazem CD
PERHEXILINE MALEATE			
* Tab 100 mg	62.90	100	✓ Pexsig

	Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	
/ERAPAMIL HYDROCHLORIDE				
* Tab 40 mg	7.01	100	1	Isoptin
* Tab 80 mg		100		Isoptin
* Tab long-acting 120 mg		250		Verpamil SR
Tables and a firm OAO man	36.02	100		Isoptin SR
* Tab long-acting 240 mg		250	•	Verpamil SR
Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO		5	./	loontin
Verpamil SR Tab long-acting 120 mg to be delisted 1 May 2020		5	•	Isoptin
Centrally-Acting Agents				
CLONIDINE				
★ Patch 2.5 mg, 100 mcg per day - Only on a prescription	7 40	4	1	Mylan
* Patch 5 mg, 200 mcg per day - Only on a prescription		4	_	Mylan Mylan
* Patch 7.5 mg, 300 mcg per day - Only on a prescription		4		Mylan
CLONIDINE HYDROCHLORIDE				
* Tab 25 mcg	8.75	112	1	Clonidine BNM
* Tab 150 mcg		100		Catapres
* Inj 150 mcg per ml, 1 ml ampoule	25.96	10	1	Medsurge
METHYLDOPA				-
				Made de la la la Mada la la
★ Tab 250 mg	15.10	100	•	Methyldopa Mylan
* Tab 250 mg	15.10 52.85	100 500		Methyldopa Mylan
* Tab 250 mg				
·				Methyldopa Mylan
Diuretics				Methyldopa Mylan
·				Methyldopa Mylan
Diuretics Loop Diuretics				Methyldopa Mylan
Diuretics Loop Diuretics BUMETANIDE	52.85	500		Methyldopa Mylan S29 S29
Diuretics Loop Diuretics BUMETANIDE * Tab 1 mg	52.85	100		Methyldopa Mylan S29 S29 Burinex
Diuretics Loop Diuretics BUMETANIDE * Tab 1 mg	52.85	500		Methyldopa Mylan S29 S29
Diuretics Loop Diuretics BUMETANIDE * Tab 1 mg * Inj 500 mcg per ml, 4 ml vial FUROSEMIDE [FRUSEMIDE]		100 5	<i>y</i>	Methyldopa Mylan S29 S29 Burinex Burinex
Diuretics Loop Diuretics BUMETANIDE * Tab 1 mg		100	<i>y y y y y</i>	Methyldopa Mylan S29 S29 Burinex Burinex Apo-Furosemide
Diuretics Loop Diuretics BUMETANIDE * Tab 1 mg * Inj 500 mcg per ml, 4 ml vial FUROSEMIDE [FRUSEMIDE] Tab 40 mg - Up to 30 tab available on a PSO		100 5	V V V V V	Methyldopa Mylan S29 S29 Burinex Burinex
Diuretics Loop Diuretics BUMETANIDE * Tab 1 mg * Inj 500 mcg per ml, 4 ml vial FUROSEMIDE [FRUSEMIDE] Tab 40 mg - Up to 30 tab available on a PSO		100 5 1,000	V V V V V	Methyldopa Mylan S29 S29 Burinex Burinex Apo-Furosemide Diurin 40
Diuretics Loop Diuretics BUMETANIDE * Tab 1 mg * Inj 500 mcg per ml, 4 ml vial FUROSEMIDE [FRUSEMIDE] Tab 40 mg - Up to 30 tab available on a PSO		100 5 1,000 50	V V V V V	Methyldopa Mylan S29 S29 Burinex Burinex Apo-Furosemide Diurin 40 Urex Forte
Diuretics Loop Diuretics BUMETANIDE * Tab 1 mg * Inj 500 mcg per ml, 4 ml vial FUROSEMIDE [FRUSEMIDE] Tab 40 mg - Up to 30 tab available on a PSO * Tab 500 mg Oral liq 10 mg per ml		100 5 1,000 50	/ / / / / / / / / /	Methyldopa Mylan S29 S29 Burinex Burinex Apo-Furosemide Diurin 40 Urex Forte
Diuretics Loop Diuretics BUMETANIDE * Tab 1 mg * Inj 500 mcg per ml, 4 ml vial FUROSEMIDE [FRUSEMIDE] Tab 40 mg - Up to 30 tab available on a PSO * Tab 500 mg * Tab 500 mg Lasix to be Sole Supply on 1 January 2020 * Inj 10 mg per ml, 25 ml ampoule		100 5 1,000 50 0 ml C		Methyldopa Mylan S29 S29 Burinex Burinex Apo-Furosemide Diurin 40 Urex Forte Lasix Lasix
Diuretics Loop Diuretics BUMETANIDE * Tab 1 mg * Inj 500 mcg per ml, 4 ml vial		100 5 1,000 50 80 ml C		Methyldopa Mylan S29 S29 Burinex Burinex Apo-Furosemide Diurin 40 Urex Forte Lasix
Diuretics Loop Diuretics BUMETANIDE * Tab 1 mg * Inj 500 mcg per ml, 4 ml vial FUROSEMIDE [FRUSEMIDE] Tab 40 mg - Up to 30 tab available on a PSO * Tab 500 mg * Tab 500 mg Lasix to be Sole Supply on 1 January 2020 * Inj 10 mg per ml, 25 ml ampoule		100 5 1,000 50 0 ml C		Methyldopa Mylan S29 S29 Burinex Burinex Apo-Furosemide Diurin 40 Urex Forte Lasix Lasix
Diuretics Loop Diuretics BUMETANIDE * Tab 1 mg		1000 5 1,0000 50 ml C 6 5		Methyldopa Mylan S29 S29 Burinex Burinex Apo-Furosemide Diurin 40 Urex Forte Lasix Lasix Frusemide-Claris
Diuretics Loop Diuretics BUMETANIDE * Tab 1 mg * Inj 500 mcg per ml, 4 ml vial FUROSEMIDE [FRUSEMIDE] Tab 40 mg — Up to 30 tab available on a PSO * Tab 500 mg * Oral liq 10 mg per ml		100 5 1,000 50 0 ml C		Methyldopa Mylan S29 S29 Burinex Burinex Apo-Furosemide Diurin 40 Urex Forte Lasix Lasix
Diuretics Loop Diuretics BUMETANIDE * Tab 1 mg		1000 5 1,0000 6 5 5 ml C		Methyldopa Mylan S29 S29 Burinex Burinex Apo-Furosemide Diurin 40 Urex Forte Lasix Lasix Frusemide-Claris
Diuretics Loop Diuretics BUMETANIDE * Tab 1 mg	52.85	1000 5 1,0000 6 5 5 ml C		Methyldopa Mylan S29 S29 Burinex Burinex Apo-Furosemide Diurin 40 Urex Forte Lasix Lasix Frusemide-Claris

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

- CARRIOVACCIU AR OVOTEM				
CARDIOVASCULAR SYSTEM				
	Subsidy (Manufacturer's P \$	rice) Subsi Per	Fully idised	Brand or Generic Manufacturer
■ SA1728 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valithe following criteria: Both: 1 Patient has heart failure with ejection fraction less than 40 2 Either:)%; and	renewal unless	notifie	d for applications meeting
2.1 Patient is intolerant to optimal dosing of spironolac2.2 Patient has experienced a clinically significant adv		on optimal dos	ing of s	pironolactone.
METOLAZONE Tab 5 mg	CBS	1 50		letolazone S29 aroxolyn S29
SPIRONOLACTONE * Tab 25 mg * Tab 100 mg Oral liq 5 mg per ml	11.80	100 100 25 ml OP	√ S	piractin piractin iomed
Potassium Sparing Combination Diuretics				
AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE * Tab 5 mg with furosemide 40 mg AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZ * Tab 5 mg with hydrochlorothiazide 50 mg	IDE	28 50		rumil
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE] * Tab 2.5 mg - Up to 150 tab available on a PSO	12.50	500	✓ <u>A</u>	rrow- Bendrofluazide
May be supplied on a PSO for reasons other than emerge * Tab 5 mg	• •	500	✓ <u>A</u>	rrow- Bendrofluazide
CHLOROTHIAZIDE Oral liq 50 mg per ml CHLORTALIDONE [CHLORTHALIDONE]	26.00	25 ml OP	✓ B	iomed
d. T. L. 05	0.50	50		

* Tab 25 mg6.50

Hygroton to be Sole Supply on 1 December 2019

F	ī	h	24	^	^	^

INDAPAMIDE

BEZAFIBRATE		
* Tab 200 mg19.01	90	✓ Bezalip
* Tab long-acting 400 mg12.89	30	✓ Bezalip Retard
GEMFIBROZIL		
* Tab 600 mg19.56	60	✓ Lipazil
-		-

✓ Hygroton

✓ Dapa-Tabs

50

90

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Other Lipid-Modifying Agents				
ACIPIMOX * Cap 250 mg NICOTINIC ACID	18.75	30	√ 0	Dibetam
* Tab 50 mg * Tab 500 mg	4.12 17.89	100 100	_	Apo-Nicotinic Acid Apo-Nicotinic Acid
Resins				
COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g	28.60	30	√ 0	Colestid
HMG CoA Reductase Inhibitors (Statins)				
Prescribing Guidelines				

Treatment with HMG CoA Reductase Inhibitors (statins) is recommended for patients with dyslipidaemia and an absolute 5 year cardiovascular risk of 15% or greater.

500

✓ I orstat

ATORVASTATIN	 See prescribing guideline above
¥ Tob 10 ma	

*	Tab 20 mg	9.99	500	✓ Lorstat
*	Tab 40 mg	15.93	500	✓ Lorstat
*	Tab 80 mg	27.19	500	✓ Lorstat
PR	AVASTATIN - See prescribing guideline above			
	Tab 20 mg	4.72	100	✓ Apo-Pravastatin
	Tab 40 mg		100	✓ Apo-Pravastatin
SIN	IVASTATIN - See prescribing guideline above			
	Tab 10 mg	0.95	90	Simvastatin Mylan
*	Tab 20 mg	1.52	90	✓ Simvastatin Mylan
*	Tab 40 mg	2.63	90	✓ Simvastatin Mylan
*	Tab 80 mg	6.00	90	✓ Simvastatin Mylan

Selective Cholesterol Absorption Inhibitors

ΕZI	ETIMIBE - Special Authority see SA1045 below - Retail pharmacy			
*	Tab 10 mg2.00	30	1	Ezetimibe Sandoz

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 x normal) when treated with one statin: or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
<u> </u>	Per	•	Manufacturer

continued...

a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy. If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA	1046 below – Retail	pharmacy	
Tab 10 mg with simvastatin 10 mg	5.15	30	✓ Zimybe
Tab 10 mg with simvastatin 20 mg		30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg		30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg		30	✓ Zimybe

⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy. If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

GLYCERYL TRINITRATE		
* Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO4.45	250 dose OP	✓ Nitrolingual Pump Spray
* Oral spray, 400 mcg per dose – Up to 200 dose available on a PSO	200 dose OP 30 30	✓ Glytrin ✓ Nitroderm TTS ✓ Nitroderm TTS
SOSORBIDE MONONITRATE	100 30 90	✓ Ismo 20 ✓ Ismo 40 Retard ✓ Duride
Sympathomimetics ADRENALINE Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PSO4.98 5.25 Inj 1 in 10,000, 10 ml ampoule – Up to 5 inj available on a PSO27.00	5	✓ Aspen Adrenaline ✓ DBL Adrenaline ✓ Hospira
49.00	10	✓ Aspen Adrenaline

	CARDIOVASCULAR SYSTEM				
	Subsidy (Manufacturer's Price)	Fu Subsidis Per	,		
ISOPRENALINE [ISOPROTERENOL]					
* Inj 200 mcg per ml, 1 ml ampoule	36.80 (164.20)	25	Isuprel		
V	, ,		•		
Vasodilators					
HYDRALAZINE HYDROCHLORIDE					
* Tab 25 mg - Special Authority see SA1321 below - Retail					
pharmacy	CBS		/ Hydralazine		
			Onelink S29		
			AMDIPHARM \$29		
			Onelink \$29		
* Inj 20 mg ampoule	25.90	5	✓ Apresoline		
⇒SA1321 Special Authority for Subsidy					
Initial application from any relevant practitioner. Approvals valid the following criteria: Either:	d without further renev	wal unless no	tified for applications meeting		
 For the treatment of refractory hypertension; or For the treatment of heart failure in combination with a nitr inhibitors and/or angiotensin receptor blockers. 	ate, in patients who a	re intolerant	or have not responded to ACE		
MINOXIDIL					
▲ Tab 10 mg	70.00	100	✓ Loniten		
NICORANDIL					
▲ Tab 10 mg	25.57	60	✓ Ikorel		
Ikorel to be Sole Supply on 1 December 2019					
▲ Tab 20 mg	32.28	60	✓ Ikorel		
Ikorel to be Sole Supply on 1 December 2019					
PAPAVERINE HYDROCHLORIDE					
* Inj 12 mg per ml, 10 ml ampoule	217.90	5	✓ Hospira		
PENTOXIFYLLINE [OXPENTIFYLLINE]					
Tab 400 mg	42.26	50	✓ Trental 400		
Endothelin Receptor Antagonists					
AMBRISENTAN - Special Authority see SA1702 below - Retail p	pharmacy				
Tab 5 mg		30	✓ Volibris		
Tab 10 mg		30	✓ Volibris		
⇒SA1702 Special Authority for Subsidy					
Special Authority approved by the Pulmonary Arterial Hypertensic	on Panel				
Notes: Application details may be obtained from PHARMAC's we	ebsite http://www.pha	rmac.govt.nz	or:		
The Coordinator, PAH Panel					
PHARMAC, PO Box 10-254, WELLINGTON					
Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac					
BOSENTAN – Special Authority see SA1712 on the next page –					
Tab 62.5 mg	141.00	60	Bosentan Dr Reddy's		
Tab 125 mg	141.00	60	✓ Bosentan Dr Reddy's		
			itoday 5		

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

⇒SA1712 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil; or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHAWHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised: or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	✓	Manufacturer	

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see SA1825 below – Retail pharmacy			
Tab 25 mg	0.64	4	✓ Vedafil
Tab 50 mg	0.64	4	✓ Vedafil
Tab 100 mg	6.60	12	✓ Vedafil

⇒SA1825 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II; or
 - 3.2 PAH is in NYHA/WHO functional class III; or
 - 3.3 PAH is in NYHA/WHO functional class IV; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Fither:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 4.5.0
 - 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
 - 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Prostacyclin Analogues

EPOPROSTENOL - Special Authority see SA1696 below - Retail pharmacy

1 ✓ Veletri ✓ Veletri Inj 1.5 mg vial73.21

⇒SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The Coordinator, PAH Panel

PHARMAC. PO Box 10-254. WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ILOPROST - Special Authority see SA1705 below - Retail pharmacy

Nebuliser soln 10 mcg per ml, 2 ml740.10 30 ✓ Ventavis Ventavis to be Sole Supply on 1 January 2020

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The Coordinator, PAH Panel

PHARMAC. PO Box 10-254. WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 89

ADAPALENE

- a) Maximum of 30 g per prescription
- b) Only on a prescription

b) Only on a prescription			
Crm 0.1%	22.89	30 g OP	Differin
Gel 0.1%	22.89	30 g OP	Differin
ISOTRETINOIN - Special Authority see SA1475 below - Retail	pharmacy		
Cap 5 mg	8.14	60	Oratane
Cap 10 mg	13.34	120	✓ Oratane
Cap 20 mg	20.49	120	✓ Oratane

⇒SA1475 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice: and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
 - 3.2 Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
- 2 Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

TRFTINOIN

Crm 0.5 mg per q − Maximum of 50 g per prescription......13.90 50 g OP ✓ ReTrieve

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 89

HYDROGEN PEROXIDE

* Crm 1%......8.56 10 g OP ✓ Crystaderm 15 g OP ✓ Crystaderm

	Subsidy		Fully	Brand or
	(Manufacturer's F \$	Price) Subs Per	sidised •	Generic Manufacturer
MUPIROCIN	*	-		-
Oint 2%	6.60	15 g OP		
	(9.26)		Ва	ctroban
a) Only on a prescription				
b) Not in combination				
SODIUM FUSIDATE [FUSIDIC ACID] Crm 2%	1.50	5 g OP	√ Fo	han
a) Maximum of 5 g per prescription	1.59	3 y Oi	v <u>10</u>	<u>ban</u>
b) Only on a prescription				
c) Not in combination				
Oint 2%	1.59	5 g OP	✓ Fo	<u>ban</u>
a) Maximum of 5 g per prescription				
b) Only on a prescription				
c) Not in combination				
SULFADIAZINE SILVER			م	
Crm 1%	10.80	50 g OP	✓ Fla	<u>imazine</u>
a) Up to 250 g available on a PSOb) Not in combination				
b) Not in combination				
Antifungals Topical				
·	nala mana 00			
For systemic antifungals, refer to INFECTIONS, Antifun	gais, page 96			
AMOROLFINE				
a) Only on a prescription b) Not in combination				
b) Not in combination Nail soln 5%	15 95	5 ml OP	✓ Mı	coNail
CICLOPIROX OLAMINE		0 1111 01	- 111)	
a) Only on a prescription				
b) Not in combination				
Nail-soln 8%	5.72	7 ml OP	✓ Ap	o-Ciclopirox
CLOTRIMAZOLE				-
* Crm 1%	0.70	20 g OP	✓ Cl	omazol
a) Only on a prescription		•		
b) Not in combination				
₭ Soln 1%		20 ml OP	0-	naatan
a) Only on a proportion	(7.55)		Ca	nesten
a) Only on a prescription b) Not in combination				
CONAZOLE NITRATE				
Crm 1%	1 00	20 g OP		
Jiii 1/0	(7.48)	20 y Oi	Pe	varyl
a) Only on a prescription	(3)		. •)-
, , , ,				
b) Not in combination		3		
Foaming soln 1%, 10 ml sachets	9.89	3		
Foaming soln 1%, 10 ml sachets	9.89 (17.23)	3	Pe	varyl
		3	Pe	varyl

✓ MidWest

	Subsidy (Manufacturer's F \$	Price) Subs	Fully Brand or sidised Generic Manufacturer
MICONAZOLE NITRATE	Ψ	1 01	manadadad
* Crm 2%	0.74	15 g OP	✓ Multichem
a) Only on a prescription		10 9 01	· <u>mantonem</u>
b) Not in combination			
* Lotn 2%	4.36	30 ml OP	
	(10.03)		Daktarin
 a) Only on a prescription 			
b) Not in combination			
* Tinct 2%		30 ml OP	Dalstanin
a) Only an a numerounisticus	(12.10)		Daktarin
a) Only on a prescriptionb) Not in combination			
NYSTATIN			
Crm 100,000 u per g		15 g OP	
	(7.90)		Mycostatin
a) Only on a prescription			
b) Not in combination			
Antipruritic Preparations			
CALAMINE			
a) Only on a prescription			
b) Not in combination			
Crm, aqueous, BP	1.26	100 g	✓ <u>healthE Calamine</u>
			Aqueous Cream
			<u>BP</u>
Lotn, BP	12.94	2,000 ml	✓ PSM
(PSM Lotn, BP to be delisted 1 July 2020)			
CROTAMITON			
a) Only on a prescription			
b) Not in combination	0.00	00 = 00	A Hab Caatha
Crm 10%	3.29	20 g OP	✓ <u>Itch-Soothe</u>
MENTHOL – Only in combination			
 Only in combination with a dermatological base of the combination with a dermatological galenicals. 	or proprietary Topical C	orticosteriod –	Plain
Crystals	6.92	25 g	✓ MidWest
	00.00	400	/ MI-DM1

29.60

100 g

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

Corticosteroids Topical

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 79

Corticosteroids - Plain

BETAMETHASONE DIPROPIONATE			
Crm 0.05%	2.96	15 g OP	✓ Diprosone
	8.97	50 g OP	✓ Diprosone
Crm 0.05% in propylene glycol base	4.33	30 g OP	Diprosone OV
Oint 0.05%	2.96	15 g OP	Diprosone
	8.97	50 g OP	Diprosone
Oint 0.05% in propylene glycol base		30 g OP	Diprosone OV
(Diprosone OV Crm 0.05% in propylene glycol base to be delisted	1 May 2020)		
BETAMETHASONE VALERATE			
* Crm 0.1%	3.45	50 g OP	✓ Beta Cream
* Oint 0.1%	3.45	50 g OP	✓ Beta Ointment
* Lotn 0.1%	18.00	50 ml OP	✓ Betnovate
CLOBETASOL PROPIONATE			
* Crm 0.05%	2.18	30 g OP	✓ Dermol
* Oint 0.05%	2.12	30 g OP	✓ Dermol
CLOBETASONE BUTYRATE		J	
Crm 0.05%	5.38	30 g OP	
CIII 0.00 / S	(7.09)	00 g 0.	Eumovate
DIFLUCORTOLONE VALERATE	(1100)		
Crm 0.1%	8 07	50 g OP	
OIIII 0.170	(15.86)	30 g Oi	Nerisone
Fatty oint 0.1%		50 g OP	Nonsone
Taky one of the second	(15.86)	00 g 0.	Nerisone
HYDROCORTISONE	(10100)		
* Crm 1% – Only on a prescription	1 11	30 g OP	✓ DermAssist
The contract of the contract o	16.25	500 g	✓ Pharmacy Health
* Powder – Only in combination		25 g	✓ ABM
Up to 5% in a dermatological base (not proprietary Topica			
galenicals		2 · · · · · · · · · · · · · · · · · · ·	
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN			
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only or			
a prescription	10.57	250 ml	✓ <u>DP Lotn HC</u>
HYDROCORTISONE BUTYRATE			
Lipocream 0.1%	3.42	30 g OP	✓ Locoid Lipocream
	6.85	100 g OP	Locoid Lipocream
Oint 0.1%	13.70	100 g OP	✓ <u>Locoid</u>
Milky emul 0.1%	13.70	100 ml OP	✓ Locoid Crelo
METHYLPREDNISOLONE ACEPONATE			
Crm 0.1%	4.95	15 g OP	✓ Advantan
Oint 0.1%	4.95	15 g OP	✓ Advantan

	Subsidy		Fully	Brand or
	(Manufacturer's Pr	ice) Sub Per	sidised •	Generic Manufacturer
MONETACONE ELIDOATE	Ψ	1 61		Wallulacturei
MOMETASONE FUROATE Crm 0.1%	1 51	15 a OD	√ EI	ocon Alcohol Free
OIII 0.1%	2.50	15 g OP 50 g OP	_	ocon Alcohol Free
Oint 0.1%		15 g OP	✓ EI	
Ont 0.170	2.90	50 g OP	✓ EI	
Lotn 0.1%		30 ml OP	_	ocon
TRIAMCINOLONE ACETONIDE				
Crm 0.02%	6.30	100 g OP	✓ Aı	ristocort
Oint 0.02%		100 g OP	_	ristocort
Corticosteroids - Combination				
BETAMETHASONE VALERATE WITH CLIOQUINOL - Only on	a procerintian			
Crm 0.1% with clioquinol 3%		15 g OP		
Offit 0.170 with Gloquinor 070	(4.90)	13 9 01	Re	etnovate-C
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE (FU	, ,		50	oniovato o
Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP		
Offit 0.170 with 30diditi lasidate (tasiale acia) 270	(10.45)	13 9 01	Fı	ucicort
a) Maximum of 15 g per prescription	(10.10)			1010011
b) Only on a prescription				
HYDROCORTISONE WITH MICONAZOLE - Only on a prescrip	ation			
* Crm 1% with miconazole nitrate 2%		15 g OP	✓ Mi	icreme H
		·	• 1111	ioromo m
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN — C Crm 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP	√ Di	mafucort
Oint 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP		mafucort
		·	•	maraoort
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC		N		
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 m		15 ~ OD		
and gramicidin 250 mcg per g - Only on a prescription		15 g OP	\/i	aderm KC
	(6.60)		VI	ademi KC
Disinfecting and Cleansing Agents				
CHLORHEXIDINE GLUCONATE - Subsidy by endorsement				
a) No more than 500 ml per month				
b) Only if prescribed for a dialysis patient and the prescription	on is endorsed acc	cordingly.		
* Handrub 1% with ethanol 70%		500 ml	✓ he	ealthE
* Soln 4% wash	3.98	500 ml	✓ he	ealthE
TRICLOSAN - Subsidy by endorsement				
a) Maximum of 500 ml per prescription				
b)				
a) Only if prescribed for a patient identified with Methic		ohylococcus a	aureus (N	MRSA) prior to elective
surgery in hospital and the prescription is endorsed				
b) Only if prescribed for a patient with recurrent Staph	ylococcus aureus	infection and	the pres	cription is endorsed
accordingly	5.00	F00 - 1 0 F		- Int- F
Soln 1%	5.90	500 ml OP	✓ he	ealthE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Barrier Creams and Emollients Barrier Creams** DIMETHICONE Crm 5% pump bottle......4.48 500 ml OP ✓ healthE Dimethicone 5% 500 ml OP ✓ healthE Dimethicone 10% ZINC AND CASTOR OIL ✓ Boucher 500 q **Emollients** AQUEOUS CREAM 500 q ✓ Boucher CETOMACROGOL * Crm BP......2.48 500 q ✓ healthE CETOMACROGOL WITH GLYCEROL Crm 90% with glycerol 10%......2.35 500 ml OP ✓ Boucher 2.82 ✓ Pharmacy Health Sorbolene with Glycerin ✓ Boucher 3.10 1.000 ml OP 3.87 ✓ Pharmacy Health Sorbolene with Glycerin (Pharmacy Health Sorbolene with Glycerin Crm 90% with glycerol 10% to be delisted 1 March 2020) (Pharmacy Health Sorbolene with Glycerin Crm 90% with glycerol 10% to be delisted 1 March 2020) **EMULSIFYING OINTMENT** 500 g ✓ AFT OIL IN WATER EMULSION 500 a ✓ O/W Fatty Emulsion Cream PARAFFIN Oint liquid paraffin 50% with white soft paraffin 50%......5.35 500 ml OP ✓ healthE URFA ✓ healthE Urea Cream 100 g OP WOOL FAT WITH MINERAL OIL - Only on a prescription

DP Lotion

DP Lotion

BK Lotion

BK Lotion

Alpha-Keri Lotion

1.000 ml

250 ml OP

1.000 ml

250 ml OP

(11.95)

1.40

(4.53)

5.60

1.40

(7.73)

(20.53) (23.91)

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	1	Manufacturer	

Other Dermatological Bases

PARAFFIN

White soft - Only in combination	4.99	450 g	✓ healthE
•	19.99	2,500 g	✓ healthE
	3.58	500 g	
	(7.78)	ŭ	IPW
	(8.69)		PSM

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid - Plain. (IPW White soft to be delisted 1 April 2020) (PSM White soft to be delisted 1 May 2020)

Minor Skin Infections

ᇚ	IODINE

Oint 10%......3.27 25 q OP a) Maximum of 100 g per prescription

b) Only on a prescription

Riodine to be Sole Supply on 1 February 2020 500 ml

5.40 6.20 1.28

> (13.27)0.19 (7.41)

> > (6.64)

Skin preparation, povidone iodine 10% with 30% alcohol......10.00 1.63 (3.48)

Skin preparation, povidone iodine 10% with 70% alcohol......1.63

100 ml 100 ml

100 ml

15 ml

100 ml

15 ml

500 ml

✓ Betadine

✓ Riodine

Riodine ✓ Riodine

✓ Betadine

Betadine

Retadine ✓ Betadine Skin Prep

Betadine Skin Prep

Pfizer

(Betadine Antiseptic soln 10% to be delisted 1 February 2020) (Betadine Antiseptic soln 10% to be delisted 1 February 2020)

(Betadine Antiseptic soln 10% to be delisted 1 February 2020)

Parasiticidal Preparations

DIMETHICONE

200 ml OP ✓ healthE

IVERMECTIN - Special Authority see SA1225 on the next page - Retail pharmacy

Dimethicone 4% Lotion

✓ Stromectol

1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.

- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- 3) For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.



Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

⇒SA1225 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:

continued...

Subsidy	;	Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
	Per	✓	Manufacturer

continued...

- 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
- 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — **(Other parasitic infections)** only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

PERMETHRIN

Crm 5% 4.95 Lotn 5% 3.69	30 g OP 30 ml OP	✓ <u>Lyderm</u> ✓ <u>A-Scabies</u>
PHENOTHRIN Shampoo 0.5%	200 ml OP	✓ Parasidose

Psoriasis and Eczema Preparations

		ACITRETIN – Special Authority see SA1476 below – Retail pharmacy	ACITRE
✓ Novatretin	60	Cap 10 mg17.86	Cap
✓ Novatretin	60	Cap 25 mg41.36	Cap

⇒SA1476 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is female and has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of two years after the completion of the treatment; or
 - 3.2 Patient is male.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is female and has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of two years after the completion of the treatment; or
- 2 Patient is male.

BETAMETHAS	ONF DIPRO	OPIONATE WITH (CAI CIPOTRIOI

Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g		60 g OP 30 g OP	✓ <u>Daivobet</u> ✓ <u>Daivobet</u>
CALCIPOTRIOL Oint 50 mcg per g	45.00	100 g OP	✓ Daivonex

	Subsidy (Manufacturer's P	rice) Subs	Fully	Brand or Generic
	\$	Per	✓	Manufacturer
COAL TAR				
Soln BP - Only in combination	36.25	200 ml	✓ N	lidwest
 Up to 10% only in combination with a dermatologic With or without other dermatological galenicals. 	cal base or propri	etary Topical C	Corticos	teriod – Plain
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SUL	PHUR			
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% an	d			
allantoin crm 2.5%	6.59	75 g OP		
	(8.00)		Е	gopsoryl TA
	3.43	30 g OP		
	(4.35)		Е	gopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR				
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97	25 g OP	✓ C	oco-Scalp
	7.95	40 g OP	✓ C	oco-Scalp
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE	SCEIN - Only or	n a prescription	1	
★ Soln 2.3% with trolamine laurilsulfate and fluorescein sodiun	າ3.86 [°]	500 ml	✓ P	inetarsol
SALICYLIC ACID			_	
Powder – Only in combination	18.88	250 g	✓ N	lidwest
· · · · · · · · · · · · · · · · · · ·			✓ P	SM
 Only in combination with a dermatological base or With or without other dermatological galenicals. 	proprietary Topic	al Corticostero	oid – Pla	ain or collodion flexible
SULPHUR				
Precipitated - Only in combination	6.35	100 g	✓ N	lidwest
 Only in combination with a dermatological base or With or without other dermatological galenicals. 		al Corticostero	oid – Pla	ain

Scal	. D	1010	242	0.00
-10% II	. 2	a =10 r	:	
		_		

BETAMETHASONE VALERATE * Scalp app 0.1%	7.75	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE * Scalp app 0.05%			✓ <u>Dermol</u>
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%	7.30	100 ml OP	✓ <u>Locoid</u>
KETOCONAZOLE Shampoo 2%		100 ml OP	✓ <u>Sebizole</u>

- a) Maximum of 100 ml per prescriptionb) Only on a prescription

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

Crm......3.30 100 g OP

(Hamilton Sunscreen Crm to be delisted 1 March 2020)

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 67

IMIQUIMOD

PODOPHYLLOTOXIN

a) Maximum of 3.5 ml per prescription

b) Only on a prescription

Other Skin Preparations

Antineoplastics

FLUOROURACIL SODIUM

GENITO-URINARY SYSTEM

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Contraceptives - Non-hormonal

Condoms

NDOMS 49 mm – Up to 144 dev available on a PSO	11.42	144	✓ Moments
•	13.36		✓ Shield 49
53 mm		10	✓ Moments
	1.11	12	✓ Gold Knight
	11.64	144	✓ Moments
	13.36	• • • •	✓ Shield Blue
a) Maximum of 60 dev per prescription			00.12 2.120
b) Up to 60 dev available on a PSO			
53 mm (chocolate)	13.36	144	✓ Gold Knight
a) Maximum of 60 dev per prescription	10.00	177	• dola kiligiti
b) Up to 60 dev available on a PSO			
	10.06	144	✓ Cold Knight
53 mm (strawberry)	13.30	144	Gold Knight
a) Maximum of 60 dev per prescription			
b) Up to 60 dev available on a PSO			
53 mm, 0.05 mm thickness		10	✓ Moments
	11.42	144	✓ Moments
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
53 mm, chocolate, brown	0.95	10	Moments
	11.64	144	✓ Moments
 a) Up to 60 dev available on a PSO 			
 b) Maximum of 60 dev per prescription 			
53 mm, strawberry, red	0.95	10	✓ Moments
	11.64	144	✓ Moments
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
56 mm	0.97	10	✓ Moments
	11.64	144	✓ Moments
	13.36		✓ Durex Extra Safe
			✓ Gold Knight
a) Maximum of 60 dev per prescription			J
b) Up to 60 dev available on a PSO			
56 mm, 0.05 mm thickness	1.30	12	✓ Gold Knight
,	15.57	144	✓ Gold Knight
a) Up to 60 dev available on a PSO	10.07		
b) Maximum of 60 dev per prescription			
56 mm, 0.08 mm thickness	0.07	10	✓ Moments
JO 11111, U.UO 111111 UIIUN11635	11.64	144	✓ Moments
a) Lin to 60 day available are a DOO	11.04	144	- MOHIGHE
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription	0.07	10	/ Namenta
56 mm, 0.08 mm thickness, red		10	✓ Moments
	11.64	144	✓ Moments
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
56 mm, chocolate		12	✓ Gold Knight
	15.57	144	Gold Knight
a) Up to 60 dev available on a PSO			

AThree Hon May COUNTY MAGE BE CASE PERSON ENGINE TIME if endorsed "certified exemption" by the prescriber or pharmacist.

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
* 56 mm, shaped	13.36 (16.08)	144		Durex Confidence
a) Maximum of 60 dev per prescription	(10.00)			Durex Corniderice
b) Up to 60 dev available on a PSO				
* 56 mm, strawberry	1.30	12	✓	Gold Knight
•	15.57	144	✓	Gold Knight
a) Up to 60 dev available on a PSO				-
b) Maximum of 60 dev per prescription				
* 60 mm - Up to 144 dev available on a PSO	13.36	144	✓	Shield XL
(Shield 49 49 mm to be delisted 1 March 2020)				
(Gold Knight 53 mm to be delisted 1 March 2020)				
(Shield Blue 53 mm to be delisted 1 March 2020)				
(Gold Knight 53 mm (chocolate) to be delisted 1 March 2020)				
(Gold Knight 53 mm (strawberry) to be delisted 1 March 2020)				
(Durex Extra Safe 56 mm to be delisted 1 March 2020)				
(Gold Knight 56 mm to be delisted 1 March 2020)				
(Durex Confidence 56 mm, shaped to be delisted 1 March 2020)				

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

- ✓ Choice TT380 Short
- ✓ Choice
- TT380 Standard ✓ Choice Load 375

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

continued...

			GENIT	ΓO-UR	NARY SYSTEM
		Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
	ntinued				
	ecial Authorities approved before 1 November 1999 remain val men are still either:	lid until the expiry da	te and ca	an be rer	newed providing that
WO	on a Social Welfare benefit; or				
	 have an income no greater than the benefit. 				
cor	e approval numbers of Special Authorities approved before 1 N nbined oral contraceptives and progestogen-only contraceptive				
	HINYLOESTRADIOL WITH DESOGESTREL				
*	Tab 20 mcg with desogestrel 150 mcg and 7 inert tab		84		Mercilon 28
	a) Higher subsider of \$12.00 per 04 tob with Crossic Auth	(19.80)	the pre	-	
	a) Higher subsidy of \$13.80 per 84 tab with Special Authb) Up to 84 tab available on a PSO	ionly see SA0500 or	i trie pre	vious pa	ye .
*	Tab 30 mcg with desogestrel 150 mcg and 7 inert tab	6.62	84		
		(19.80)		N	Marvelon 28
	a) Higher subsidy of \$13.80 per 84 tab with Special Authb) Up to 84 tab available on a PSO	nority see SA0500 or	the pre	vious pa	ge
ΕT	HINYLOESTRADIOL WITH LEVONORGESTREL				
*	Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets -	_			
	Up to 112 tab available on a PSO		84		Microgynon 20 ED
		6.45	112	✓ F	emme-Tab ED
*	Tab 50 mcg with levonorgestrel 125 mcg and 7 inert tab — Up		0.4		November 50 FD
*	to 84 tab available on a PSO		84 63	• 1	Microgynon 50 ED
~	Tab 50 flicg will revolidigestier 150 flicg	(16.50)	03	N	Microgynon 30
	a) Higher subsidy of \$15.00 per 63 tab with Special Auth	` ,	the pre		0,
	b) Up to 63 tab available on a PSO	ionly occ or lococ or	r and pro	riodo pa	90
*	Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets -	_			
	Up to 112 tab available on a PSO	1.77	84	√ <u>L</u>	<u>evlen ED</u>
		6.45	112	√ F	emme-Tab ED
ΕT	HINYLOESTRADIOL WITH NORETHISTERONE				
*	Tab 35 mcg with norethisterone 1 mg - Up to 63 tab available				
	on a PSO		63	✓ E	Brevinor 1/21
*	Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to				
	84 tab available on a PSO	6.95	84	✓ [Brevinor 1/28
*	Tab 35 mcg with norethisterone 500 mcg – Up to 63 tab	6 60	60	./ -	Brevinor 21
	available on a PSO	0.0∠	63	▼ [DIEVINOI ZI

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

* Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - Up

(Brevinor 1/21 Tab 35 mcg with norethisterone 1 mg to be delisted 1 July 2020) (Brevinor 21 Tab 35 mcg with norethisterone 500 mcg to be delisted 1 July 2020)

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

84

✓ Norimin

	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer
continued			

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit: and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

LEVONORGESTREL

*	Tab 30 mcg	84
	(16.50) Microlut

- a) Higher subsidy of \$13.80 per 84 tab with Special Authority see SA0500 on the previous page
- b) Up to 84 tab available on a PSO

Subdermal implant $(2 \times 75 \text{ mg rods})$ – Up to 3 pack available

on a PSO106.92	1	✓ <u>Jadelle</u>
MEDROXYPROGESTERONE ACETATE		
Inj 150 mg per ml, 1 ml syringe - Up to 5 inj available on a PSO7.98	1	✓ Depo-Provera
Depo-Provera to be Sole Supply on 1 December 2019		

NORETHISTERONE

*	Tab 350 mcg - Up to 84 tal	available on a PSO	6.25	84	✓ Noriday 28

Emergency Contraceptives

LEVONORGESTRE

*	Tab 1.5 mg4.9	95 1	✓ Postinor-1

- a) Maximum of 2 tab per prescription
- b) Up to 5 tab available on a PSO
- c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

★ Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO.......4.67 168 ✓ Ginet

Gynaecological Anti-infectives

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID			
Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate			
0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator	8.43	100 g OP	
(24	4.00)		Aci-Jel
CLOTRIMAZOLE			
* Vaginal crm 1% with applicators	2.50	35 g OP	✓ Clomazol
Clomazol to be Sole Supply on 1 January 2020			
* Vaginal crm 2% with applicators	3.00	20 g OP	Clomazol
Clomazol to be Sole Supply on 1 January 2020			
MICONAZOLE NITRATE			
* Vaginal crm 2% with applicator	3.88	40 g OP	✓ Micreme
NYSTATIN			
Vaginal crm 100,000 u per 5 g with applicator(s)	4.45	75 g OP	✓ Nilstat
3 , 1 3 11 ()		0	

Myometrial and Vaginal Hormone Preparations

,			3	
ERGO	METRINI	MALE	EATE	

Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO105.00	5	✓ DBL Ergometrine
OESTRIOL * Crm 1 mg per g with applicator	15 g OP 15	✓ Ovestin ✓ Ovestin
OXYTOCIN — Up to 5 inj available on a PSO Inj 5 iu per ml, 1 ml ampoule	5 5	✓ Oxytocin BNM ✓ Oxytocin BNM
OXYTOCIN WITH ERGOMETRINE MALEATE – Up to 5 inj available on a PSO Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml15.00	5	✓ Syntometrine

Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 107

5-Alpha Reductase Inhibitors

FINASTERIDE – Special Authority see SA0928 below – Retail pharmacy

* Tab 5 mg4.81 100

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE − Special Authority see SA1032 below − Retail pharmacy

* Cap 400 mcg17.73 100

✓ Tamsulosin-Rex

Tamsulosin-Rex to be Sole Supply on 1 January 2020

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

OXYBUTYNIN * Tab 5 mg	8.85	500	✓ Apo-Oxybutynin
* Oral liq 5 mg per 5 ml		473 ml	✓ Apo-Oxybutynin
POTASSIUM CITRATE			
Oral liq 3 mmol per ml - Special Authority see SA1083 below -			
Retail pharmacy	31.80	200 ml OP	✓ Biomed

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM	CITRO-	TARTRATE
--------	--------	----------

*	Grans eff 4 g sachets	2.34	28	•	<u>Ural</u>
SOI	LIFENACIN SUCCINATE				
	Tab 5 mg	3.00	30	1	Solifenacin Mylan
	Tab 10 mg	5.50	30	1	Solifenacin Mylan

	Subsidy		Fully	Brand or	
	(Manufacturer's Price)	Su	bsidised	Generic	
	\$	Per	1	Manufacturer	
TOLTERODINE - Special Authority see SA1272 below - Retail	pharmacy				
Tab 1 mg	14.56	56	✓ A	rrow-Tolterodine	
Tab 2 mg	14.56	56	✓ A	rrow-Tolterodine	
(Arrow-Tolterodine Tab 1 mg to be delisted 1 March 2020)					

⇒SA1272 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where patient has overactive bladder and a documented intolerance of, or is non-responsive to oxybutynin.

Detection of Substances in Urine

ORTHO-TOLIDINE			
* Compound diagnostic sticks	7.50	50 test OP	
, ,	(8.25)		Hemastix
TETRABROMOPHENOL			
* Blue diagnostic strips	7.02	100 test OP	
•	(13.92)		Albustix

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

Calcium Homeostasis

CAL	CITONIN		
*	Inj 100 iu per ml, 1 ml ampoule121.00	5	✓ Miacalcic
CIN	ACALCET – Special Authority see SA1618 below – Retail pharmacy		
	Tab 30 mg - Wastage claimable210.30	28	✓ Sensipar

⇒SA1618 Special Authority for Subsidy

Initial application only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial − Special Authority see SA1687 below −

Retail pharmacy......38.03 1

✓ Zoledronic acid

Mylan

⇒SA1687 Special Authority for Subsidy

Initial application — **(bone metastases)** only from an oncologist, haematologist or palliative care specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
 - 2.1 Patient has bone metastases or involvement; and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
 - 3.1 Patient has bone metastases or involvement; and
 - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer) only from an oncologist or medical practitioner on the recommendation of a oncologist. Approvals valid for 2 years for applications meeting the following criteria:
All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.

BE	TAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETA	ГЕ	
*	Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml19.20	5	
	(36.96)		Celestone
			Chronodose
DE	XAMETHASONE		
*	Tab 0.5 mg - Retail pharmacy-Specialist0.99	30	✓ Dexmethsone
	Up to 60 tab available on a PSO		
*	Tab 4 mg - Retail pharmacy-Specialist1.90	30	✓ <u>Dexmethsone</u>
	Up to 30 tab available on a PSO		
	Oral liq 1 mg per ml – Retail pharmacy-Specialist45.00	25 ml OP	✓ Biomed
	Oral liq prescriptions:		
	Must be written by a Paediatrician or Paediatric Cardiologist; or		
	2) On the recommendation of a Paediatrician or Paediatric Cardiologic	ıst.	
DE	XAMETHASONE PHOSPHATE		
	Dexamethasone phosphate injection will not be funded for oral use.		
	Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO14.19	10	✓ Max Health
	Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO25.18	10	✓ Max Health
	JDROCORTISONE ACETATE		
*	Tab 100 mcg14.32	100	✓ Florinef
ΗY	DROCORTISONE		
*	Tab 5 mg8.10	100	✓ Douglas
*	Tab 20 mg20.32	100	✓ Douglas
*	Inj 100 mg vial5.30	1	✓ Solu-Cortef
	a) Up to 5 inj available on a PSO		
	b) Only on a PSO		
ME	THYLPREDNISOLONE - Retail pharmacy-Specialist		
*	Tab 4 mg112.00	100	✓ <u>Medrol</u>
*	Tab 100 mg194.00	20	✓ <u>Medrol</u>
ME	THYLPREDNISOLONE (AS SODIUM SUCCINATE) - Retail pharmacy-Spec	cialist	
	Inj 40 mg vial18.90	1	✓ Solu-Medrol-Act-
			<u>O-Vial</u>
	11405		
	Inj 125 mg vial28.90	1	✓ Solu-Medrol-Act-
			<u>O-Vial</u>
	Inj 500 mg vial22.78	1	✓ Solu-Medrol-Act-
	11) 000 Hg Vid	•	0-Vial
			<u>*</u>
	Inj 1 g vial27.83	1	✓ Solu-Medrol
ME	THYLPREDNISOLONE ACETATE		
	Inj 40 mg per ml, 1 ml vial44.40	5	✓ Depo-Medrol

	Subsidy		Fully	Brand or
	(Manufacturer's Pr \$	rice) Subsi Per	idised •	Generic Manufacturer
PREDNISOLONE				
Yoral liq 5 mg per ml - Up to 30 ml available on a PSO Restricted to children under 12 years of age.	6.00	30 ml OP	√ <u>I</u>	Redipred
PREDNISONE				
* Tab 1 mg	10.68	500	1	Apo-Prednisone
* Tab 2.5 mg	12.09	500	1	Apo-Prednisone
* Tab 5 mg - Up to 30 tab available on a PSO	11.09	500	1	Apo-Prednisone
* Tab 20 mg		500	1	Apo-Prednisone
TETRACOSACTRIN				
* Inj 250 mcg per ml, 1 ml ampoule	75.00	1	1	AU Synacthen
, , , , , , , , , , , , , , , , , , , ,				Synacthen
				Synacthen S29 S29
* Inj 1 mg per ml, 1 ml ampoule	690.00	1		Synacthen Depot
and the porting for the compositor		•		Synacthene
				Retard \$29
(Synacthen S29 S29 Inj 250 mcg per ml, 1 ml ampoule to be de	elisted 1 January 2	2020)		
TRIAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule	20.80	5	1	Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule		5		Kenacort-A 40
, , , , , , , , , , , , , , , , , , , ,			-	

Sex Hormones Non Contraceptive

Androgen Agonists and Antagonists

CYPROTERONE ACETATE – Retail pharmacy-Specialist			
Tab 50 mg	13.17	50	✓ Siterone
Tab 100 mg	26.75	50	✓ Siterone
TESTOSTERONE Patch 5 mg per day	90.00	30	✓ Androderm
TESTOSTERONE CIPIONATE – Retail pharmacy-Specialist Inj 100 mg per ml, 10 ml vial	76.50	1	✓ <u>Depo-Testosterone</u>
TESTOSTERONE ESTERS – Retail pharmacy-Specialist Inj 250 mg per ml, 1 ml	12.98	1	✓ Sustanon Ampoules
TESTOSTERONE UNDECANOATE – Retail pharmacy-Specialist Cap 40 mg	21.00	60	✓ Andriol Testocaps
Inj 250 mg per ml, 4 ml vial	86.00	1	✓ Reandron 1000

Hormone Replacement Therapy - Systemic

Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

	(Ma	Subsidy nufacturer's P	Price) Subs	Fully sidised	
		\$	Per	1	Manufacturer
Destrogens					
ESTRADIOL – See prescribing guideline	on the previous page				
Tab 1 mg		4.12	28 OP		
		(11.10)			Estrofem
Tab 2 mg		4.12	28 OP		
		(11.10)			Estrofem
Patch 25 mcg per day		6.12	8	1	Estradot
 a) No more than 2 patch per wee 	(
b) Only on a prescription					
Patch 50 mcg per day		7.04	8	1	Estradot 50 mcg
a) No more than 2 patch per wee	(
b) Only on a prescription					
Patch 75 mcg per day		7.91	8	1	Estradot
a) No more than 2 patch per wee					
b) Only on a prescription					
Patch 100 mcg per day		7.91	8	1	Estradot
a) No more than 2 patch per wee		-	-		
b) Only on a prescription	•				
	na quidalina an tha necessir	10 0000			
ESTRADIOL VALERATE – See prescrib			0.4		D
Tab 1 mg			84	_	Progynova
Tab 2 mg		12.36	84	•	Progynova
ESTROGENS – See prescribing guidelir					
Conjugated, equine tab 300 mcg		3.01	28		
		(13.50)			Premarin
Conjugated, equine tab 625 mcg		4.12	28		
		(13.50)			Premarin
Progestogens					
EDROXYPROGESTERONE ACETATE	- See prescribing guideline	on the prev	vious page		
Tab 2.5 mg			30	1	Provera
Tab 5 mg		14.00	100	✓	Provera
Tab 10 mg		7.15	30	1	Provera
Progestogen and Oestrogen Co	ombined Preparation	ıs			
ESTRADIOL WITH NORETHISTERONE	- See prescribing guidelin	ne on the pre	evious page		
Tab 1 mg with 0.5 mg norethisterone a			28 OP		
		(18.10)	_0 0.		Kliovance
Tab 2 mg with 1 mg norethisterone acc	tate	` '	28 OP		
gg		(18.10)			Kliogest
Tab 2 mg with 1 mg norethisterone acc	state (10) and 2 mg	(10.10)			9001
oestradiol tab (12) and 1 mg oestr		5.40	28 OP		
destraction tab (12) and 1 mg destr	zuioi lau (u)	(18.10)	28 OP		Trisequens
		(10.10)			Посциено
Other Oestrogen Preparations					
THINYLOESTRADIOL					
Tab 10 mcg		17.60	100	1	NZ Medical and

 $[\]blacktriangle \textit{Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. }$

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
OESTRIOL	7.00	30	•	Ovestin
Other Progestogen Preparations				
LEVONORGESTREL				
* Intra-uterine device 52 mg	269.50	1	1	Mirena
* Intra-uterine device 13.5 mg	215.60	1	✓	<u>Jaydess</u>
MEDROXYPROGESTERONE ACETATE				
Tab 100 mg - Retail pharmacy-Specialist	101.00	100	✓	Provera HD
NORETHISTERONE				
* Tab 5 mg - Up to 30 tab available on a PSO Primolut N to be Sole Supply on 1 January 2020	18.29	100	✓	Primolut N
PROGESTERONE				
Cap 100 mg - Special Authority see SA1609 below - Retail				
pharmacy	16.50	30	1	Utrogestan
⇒SA1609 Special Authority for Subsidy				

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 For the prevention of pre-term labour*; and
- 2 Either:
 - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 For the prevention of pre-term labour*; and
- 2 Treatment is required for second or subsequent pregnancy; and
- 3 Either:
 - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are unapproved indications.

Thyroid and Antithyroid Agents		
CARBIMAZOLE		
* Tab 5 mg10.80	100	✓ AFT
		Carbimazole S29
		✓ Neo-Mercazole
LEVOTHYROXINE		
* Tab 25 mcg3.89	90	✓ Synthroid
* Tab 50 mcg1.71	28	✓ Mercury Pharma
4.05	90	✓ Synthroid
64.28	1,000	✓ Eltroxin
* Tab 100 mcg1.78	28	Mercury Pharma
4.21	90	✓ Synthroid
66.78	1,000	✓ Eltroxin

	(Manufacturer's Price)	Su	bsidised	Generic	
	\$	Per	✓	Manufacturer	
PROPYLTHIOURACIL – Special Authority see SA1199 below –	' '				

Propylthiouracil is not recommended for patients under the age of 18 years unless the patient is pregnant and other treatments are contraindicated.

100 ✓ PTU S29

⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) - Special Authority see SA1629 below	v – Retail ph	armacy	
*	Inj 5 mg cartridge	34.88	1	Omnitrope
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
*	Inj 15 mg cartridge	.104.63	1	✓ Omnitrope

⇒SA1629 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under: and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and</p>
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Pri	,	Subsidised	Generic	
\$	Per		Manufacturer	

continued...

- 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
- 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and

Subsidy)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
	Per	✓	Manufacturer

continued...

6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients.

GnRH Analogues

GOSERELIN			
Implant 3.6 mg, syringe	66.48	1	✓ Zoladex
Implant 10.8 mg, syringe		1	✓ Zoladex

Subsidy

Fully

Brand or

	(Manufacturer's Price)	Subs Per	idised	Generic Manufacturer
EUPRORELIN				
Additional subsidy by endorsement where the patient is a chil goserelin and the prescription is endorsed accordingly.	d or adolescent and	is unable t	to tolera	ate administration of
Inj 3.75 mg prefilled dual chamber syringe - Higher subsidy of				
\$221.60 per 1 inj with Endorsement	66.48	1		
	(221.60)		L	ucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe - Higher subsidy				
of \$591.68 per 1 inj with Endorsement	177.50	1		
	(591.68)		L	ucrin Depot 3-month

Vasopressin Agonists

DESMOPRESSIN ACETATE	
----------------------	--

ΙF

Tab 100 mcg - Special Authority see SA1401 below - Retail pharmacy	25.00	30	✓ Minirin
Tab 200 mcg - Special Authority see SA1401 below - Retail pharmacy	39.03	30 2.5 ml OP 6 ml OP	✓ Minirin ✓ Minirin ✓ <u>Desmopressin-PH&T</u>
Inj 4 mcg per ml, 1 ml – Special Authority see SA1401 below – Retail pharmacy	67.18	10	✓ Minirin

⇒SA1401 Special Authority for Subsidy

Initial application — (Desmopressin tablets for Nocturnal enuresis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has primary nocturnal enuresis; and
- 2 The nasal forms of desmopressin are contraindicated; and
- 3 An enuresis alarm is contraindicated.

Initial application — (Desmopressin tablets for Diabetes insipidus) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has cranial diabetes insipidus; and
- 2 The nasal forms of desmopressin are contraindicated.

Renewal — (Desmopressin tablets) from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from the treatment.

Initial application — (Desmopressin injection) only from a relevant specialist. Approvals valid for 2 years where the patient cannot use desmopressin nasal spray or nasal drops.

Renewal — (Desmopressin injection) only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Other Endocrine Agents

CABERGOLINE

		Tab 0.5 mg - Maximum of 2 tab per prescription; can be
✓ Dostinex	2	waived by Special Authority see SA1370 on the next page3.75
✓ Dostinex	8	15.20

Subsidy		Fully	Brand or
(Manufacturer's Pri	ice)	Subsidised	Generic
\$	Per		Manufacturer

⇒SA1370 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 pathological hyperprolactinemia; or
- 2 acromegaly*.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

1 :1 () [\]	IFEI	NI — 1	H	1 -

Tab 50 mg	10	✓ Mylan Clomiphen S29
DANAZOL		
Cap 100 mg	100	✓ Azol
Cap 200 mg	100	✓ Azol
METYRAPONE		
Cap 250 mg - Retail pharmacy-Specialist520.00	50	Metopirone

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Anthelmintics

ALBENDAZOLE - Special Authority see SA1318 below - Retail pharmacy

⇒SA1318 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

MEBENDAZOLE - Only on a prescription

Tab 100 mg	24.19	24	De-Worm
Oral lig 100 mg per 5 ml	2.18	15 ml	
, 31	(7.17)		Vermox
PRAZIQUANTEL			
Tab 600 mg	68.00	8	✓ Biltricide

Antibacterials

CEFACLOR MONOHYDRATE

- a) For topical antibacterials, refer to DERMATOLOGICALS, page 59
- b) For anti-infective eve preparations, refer to SENSORY ORGANS, page 228

Cephalosporins and Cephamycins

Cap 250 mg	24.70	100	✓ Ranbaxy-Cefactor
Grans for oral liq 125 mg per 5 ml - Wastage claimable	3.53	100 ml	✓ Ranbaxy-Cefactor
	4.33		✓ Keflor
CEFALEXIN			
Cap 250 mg	3.33	20	 Cephalexin ABM
Cap 500 mg	3.95	20	✓ Cephalexin ABM
Grans for oral liq 25 mg per ml - Wastage claimable	8.75	100 ml	 Cefalexin Sandoz
Grans for oral liq 50 mg per ml - Wastage claimable	11.75	100 ml	✓ Cefalexin Sandoz

CEFAZOLIN - Subsidy by endorsement

Only if prescribed for dialysis or cellulitis in accordance with a DHB approved protocol and the prescription is endorsed accordingly.

Inj 500 mg vial	3.39	5	✓ <u>AFT</u>
Inj 1 g vial	3.29	5	✓ AFT

CEFTRIAXONE - Subsidy by endorsement

- a) Up to 10 inj available on a PSO
- b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of pelvic inflammatory disease, or the treatment of suspected meningococcal disease, and the prescription or PSO is endorsed accordingly.

Inj 500 mg vial	0.89	1	✓ Ceftriaxone-AFT
,	1.20		✓ DEVA
Ceftriaxone-AFT to be Sole Supply on 1 January 2020			
Inj 1 g vial	0.84	1	✓ DEVA
, 0	3.99	5	✓ Ceftriaxone-AFT

Ceftriaxone-AFT to be Sole Supply on 1 January 2020

(DEVA Inj 500 mg vial to be delisted 1 January 2020)

(DEVA Inj 1 g vial to be delisted 1 January 2020)

	(Manufacturer's Price)		ubsidised	Generic
	\$	Per		Manufacturer
CEFUROXIME AXETIL — Subsidy by endorsement				
Only if prescribed for prophylaxis of endocarditis and the pre	scription is endorsed	accordi	ngly.	
Tab 250 mg	45.93	50	✓ Z	innat
Zinnat to be Sole Supply on 1 February 2020				

Cubaidu

Eully.

Drand or

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Authority. Tab 250 mg8.**	19 30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO		✓ Apo-Azithromycin
Grans for oral liq 200 mg per 5 ml (40 mg per ml) – Wastage		
claimable14.3	38 15 ml	✓ Zithromax

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

Grans for oral liq 250 mg per 5 ml – Wastage claimable......192.00 50 ml

	Subsidy	Fully	Brand or
(Man	ufacturer's Price)	Subsidised	Generic
	\$ P	er 🗸	Manufacturer

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN (AS LACTOBIONATE)	40.00		
Inj 1 g vial	10.00	1	Erythrocin IV
Erythrocin IV to be Sole Supply on 1 December 2019			
ERYTHROMYCIN ETHYL SUCCINATE			4
Tab 400 mg	16.95	100	E-Mycin
a) Up to 20 tab available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			4
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	E-Mycin
a) Up to 300 ml available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	E-Mycin
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
ERYTHROMYCIN STEARATE			
Tab 250 mg - Up to 30 tab available on a PSO	14.95	100	
	(22.29)		ERA
Tab 500 mg	29.90	100	
	(44.58)		ERA
ROXITHROMYCIN			
Tab disp 50 mg	8.29	10	✓ Rulide D
Restricted to children under 12 years of age.			
Tab 150 mg	8.28	50	✓ Arrow-
-			Roxithromycin
Tab 300 mg	16.33	50	✓ Arrow-
			Roxithromycin

Cap 250 mg	_	Subsidy (Manufacturer's Price	e) Subs	Fully Brand or idised Generic
AMOXICILLIN Cap 250 mg		\$	Per	✓ Manufacturer
Cap 250 mg	Penicillins			
a) Up to 30 cap available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP Cap 500 mg	AMOXICILLIN			
a) Up to 30 cap available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP Cap 500 mg	Cap 250 mg	14.97	500	✓ Apo-Amoxi
b) Up to 10 x the maximum PSO quantity for RFPP Cap 500 mg		22.50		✓ Alphamox
Cap 500 mg	a) Up to 30 cap available on a PSO			
a) Up to 30 cap available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml	b) Up to 10 x the maximum PSO quantity for RFPP			
a) Up to 30 cap available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml	Cap 500 mg	16.75	500	✓ Apo-Amoxi
b) Up to 10 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml		36.98		✓ Alphamox
Grans for oral liq 125 mg per 5 ml	a) Up to 30 cap available on a PSO			
a) Up to 200 ml available on a PSO b) Wastage claimable Grans for oral liq 250 mg per 5 ml	b) Up to 10 x the maximum PSO quantity for RFPP			
b) Wastage claimable Grans for oral liq 250 mg per 5 ml	Grans for oral liq 125 mg per 5 ml	1.20	100 ml	✓ Alphamox 125
Grans for oral liq 250 mg per 5 ml	a) Up to 200 ml available on a PSO			
a) Up to 300 ml available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP c) Wastage claimable lnj 250 mg vial	b) Wastage claimable			
b) Up to 10 x the maximum PSO quantity for RFPP c) Wastage claimable lnj 250 mg vial	Grans for oral liq 250 mg per 5 ml	1.31	100 ml	✓ Alphamox 250
c) Wastage claimable Inj 250 mg vial				
Inj 250 mg vial	, ,			
Inj 500 mg vial				
Inj 1 g vial — Up to 5 inj available on a PSO				
(Apo-Amoxi Cap 250 mg to be delisted 1 April 2020) (Apo-Amoxi Cap 500 mg to be delisted 1 April 2020) AMOXICILLIN WITH CLAVULANIC ACID Tab 500 mg with clavulanic acid 125 mg − Up to 30 tab available on a PSO	, ,		. •	
Apo-Amoxi Cap 500 mg to be delisted 1 April 2020) AMOXICILLIN WITH CLAVULANIC ACID Tab 500 mg with clavulanic acid 125 mg — Up to 30 tab available on a PSO		17.29	10	✓ <u>Ibiamox</u>
AMOXICILLIN WITH CLAVULANIC ACID Tab 500 mg with clavulanic acid 125 mg — Up to 30 tab available on a PSO				
Tab 500 mg with clavulanic acid 125 mg − Up to 30 tab available on a PSO	(Apo-Amoxi Cap 500 mg to be delisted 1 April 2020)			
available on a PSO	AMOXICILLIN WITH CLAVULANIC ACID			
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 mg per ml	Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab			
per ml	available on a PSO	1.88	20	✓ Augmentin
per ml	Grans for oral lig amoxicillin 25 mg with clavulanic acid 6.25	mg		_ -
b) Wastage claimable Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 mg per ml – Up to 200 ml available on a PSO	per ml	3.83	100 ml	✓ Augmentin
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 mg per ml – Up to 200 ml available on a PSO	a) Up to 200 ml available on a PSO			
per ml − Up to 200 ml available on a PSO	b) Wastage claimable			
per ml − Up to 200 ml available on a PSO	Grans for oral lig amoxicillin 50 mg with clavulanic acid 12.5	mg		
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj available on a PSO	per ml - Up to 200 ml available on a PSO	2.20 1	00 ml OP	✓ Curam
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj available on a PSO	BENZATHINE BENZYI PENICII I IN			
available on a PSO				
BENZYLPENICILLIN SODIUM [PENICILLIN G] Inj 600 mg (1 million units) vial – Up to 5 inj available on a PSO10.35 25.88 25 Sandoz Pan-Penicillin G		344.93	10	✓ Bicillin LA
Inj 600 mg (1 million units) vial – Up to 5 inj available on a PSO10.35 10 Sandoz 25.88 25 Pan-Penicillin G				
25.88 25 ✓ Pan-Penicillin G		SO 10.35	10	✓ Sandoz
	ing 500 mg (1 million units) viai – op to 5 mg available on a F			
		20.00	20	Sodium S29

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	I Generic Manufacturer
	Ψ	rei		Wallulacturei
LUCLOXACILLIN				
Cap 250 mg – Up to 30 cap available on a PSO		250		Staphlex
Cap 500 mg		500		Staphlex
Grans for oral liq 25 mg per ml	2.29	100 m	•	<u>AFT</u>
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 50 mg per ml	3.68	100 m	•	<u>AFT</u>
 a) Up to 200 ml available on a PSO 				
b) Wastage claimable			_	
Inj 250 mg vial		10		Flucloxin
Inj 500 mg vial		10		Flucloxin
Inj 1 g vial – Up to 5 inj available on a PSO	5.22	5	/	Flucil
HENOXYMETHYLPENICILLIN (PENICILLIN V)				
Cap 250 mg - Up to 30 cap available on a PSO	2.59	50	✓	Cilicaine VK
Cap 500 mg		50	✓	Cilicaine VK
a) Up to 20 cap available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	2.99	100 m	✓	AFT
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
c) AFT to be Sole Supply on 1 January 2020				
Grans for oral liq 250 mg per 5 ml	3.99	100 m	✓	AFT
a) Up to 300 ml available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
d) AFT to be Sole Supply on 1 January 2020				
ROCAINE PENICILLIN				
	102.50	5	1	Cilicaine
Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO	123.50	э	•	Cilicaine
Tetracyclines				
OXYCYCLINE				
Tab 50 mg - Up to 30 tab available on a PSO		30		
	(6.00)		_	Doxy-50
Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	•	Doxine
Doxy-50 Tab 50 mg to be delisted 1 January 2020)				
INOCYCLINE HYDROCHLORIDE				
Tab 50 mg - Additional subsidy by Special Authority see				
SA1355 below – Retail pharmacy	5.79	60		
,,	(12.05)			Mino-tabs
Cap 100 mg		100		
	(52.04)			Minomycin
SA1355 Special Authority for Manufacturers Price	(/			, .
itial application from any relevant practitioner. Approvals valid	d without further ren	ייי ובייי	aloce netit	ind where the nations ha
, , , , , , , , , , , , , , , , , ,	a williout luitile! Telle	wai ui	11011 66511	ieu wiiere irie palient na
Sacea.	Doto!!!			
ETRACYCLINE – Special Authority see SA1332 on the next pa	•	,		Tetroevelir
Cap 500 mg	46.00	30	•	Tetracyclin
				Wolff S29

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 59

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg - Up to 5 tab available on a PSO	1.45	28	✓ Cipflox
Tab 500 mg - Up to 5 tab available on a PSO	1.99	28	✓ Cipflox
Tab 750 mg	3.15	28	✓ Cipflox
CLINDAMYCIN			
Cap hydrochloride 150 mg – Maximum of 4 cap per			
prescription; can be waived by endorsement - Retail			
pharmacy - Specialist	4.10	16	Clindamycin ABM
	4.61	24	Dalacin C
Inj phosphate 150 mg per ml, 4 ml ampoule - Retail			
pharmacy-Specialist	39.00	10	✓ Dalacin C
(Clindamycin ABM Cap hydrochloride 150 mg to be delisted 1 A			
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist -	•	mont	
			ordingly
Only if prescribed for dialysis or cystic fibrosis patient and the			
Inj 150 mg		1	✓ Colistin-Link
GENTAMICIN SULPHATE			
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement.	25.00	5	DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient			ection and the prescription is
endorsed accordingly.		,	
Inj 40 mg per ml, 2 ml ampoule – Subsidy by endorsement.	17 50	10	✓ Pfizer
ing 40 mg por mi, 2 mi ampoule Gubbidy by endorsement.	30.00	50	✓ Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient	00.00	• •	
endorsed accordingly.	or complicated uni	iary tract iiii	ection and the prescription is
MOXIFLOXACIN - Special Authority see SA1740 below - Reta	il nharmacy		

⇒SA1740 Special Authority for Subsidy

Tab 400 mg52.00

No patient co-payment payable

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

1 Both:

continued...

5

✓ Avelox

Subsidy	Fu	lly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
\$	Per	/	Manufacturer

continued...

- 1.1 Active tuberculosis*; and
- 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Either:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy

⇒SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Fither:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharmacy

⇒SA1328 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy: or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

	Subsidy		Fully	Brand or	-
	(Manufacturer's Price) \$	Subs Per	idised	Generic Manufacturer	
ACRIMA ELIGIPATE (ELIGIPIO A CIP)	Ψ	FEI		Waltulacturei	=
SODIUM FUSIDATE [FUSIDIC ACID]	24.50	10	./ 5	مامانه	
Tab 250 mg - Retail pharmacy-Specialist Prescriptions must be written by, or on the recommendat		12 disease ph	✓ <u>Fu</u>		^
Frescriptions must be written by, or on the recommendat	ion or, an infectious	uisease pii	iysician c	or a cirrical microbiologis	>
SULFADIAZINE SODIUM - Special Authority see SA1331 below	Datail pharmany				
Tab 500 mg		56	✓ W	ockhardt \$29	
	343.20	50	• ***	OCKIIAI UL 329	
⇒SA1331 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals valid	l without further rene	wal unlace	notified	for applications meeting	7
he following criteria:	i williout furtifier refre	wai uilioss	riounca	ioi applications meeting	,
Any of the following:					
1 For the treatment of toxoplasmosis in patients with HIV for	a period of 3 month	s; or			
2 For pregnant patients for the term of the pregnancy; or	·				
3 For infants with congenital toxoplasmosis until 12 months of	of age.				
OBRAMYCIN					
Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement		5		bramycin Mylan	
Only if prescribed for dialysis or cystic fibrosis patient and	the prescription is	endorsed a	ccording	gly.	
Solution for inhalation 60 mg per ml, 5 ml – Subsidy by	0.000.00				
endorsement	2,200.00 5	66 dose	✓ T0)BI	
a) Wastage claimableb) Only if prescribed for a cystic fibrosis patient and the	aracariation is andar	and annor	lingly		
	prescription is endor	seu accord	inigiy.		
'RIMETHOPRIM 烙 Tab 300 mg – Up to 30 tab available on a PSO	16 50	50	✓ TN	ND	
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXA		00	• 111	<u></u>	
★ Tab trimethoprim 80 mg and sulphamethoxazole 400 mg — U	•				
to 30 tab available on a PSO		500	✓ Tri	isul	
♦ Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to 200 r	nl				
available on a PSO		100 ml	✓ De	<u>eprim</u>	
/ANCOMYCIN – Subsidy by endorsement					
Only if prescribed for a dialysis or cystic fibrosis patient or for			for treatn	nent of Clostridium	
difficile following metronidazole failure and the prescription is		•			
Inj 500 mg vial	2.37	1	✓ My	<u>/lan</u>	
Antifungals					ı
Antinangaio					ı
n) For topical antifungals refer to DERMATOLOGICALS, page 60)				
) For topical antifungals refer to GENITO URINARY, page 75					
FLUCONAZOLE			_		
Cap 50 mg - Retail pharmacy-Specialist		28	✓ <u>My</u>		
Cap 150 mg — Subsidy by endorsement		1	✓ <u>M</u> y		
 a) Maximum of 1 cap per prescription; can be waived by b) Patient has vaginal candida albicans and the practitio 					
not recommended and the prescription is endorsed at	ccordingly: can be w	aived by e	ndorsem	ent - Retail pharmacy -	
Specialist.				,,,	
Cap 200 mg - Retail pharmacy-Specialist	5.08	28	✓ My	<u>/lan</u>	
Powder for oral suspension 10 mg per ml - Special Authority			_		
see SA1359 on the next page - Retail pharmacy		35 ml		flucan S29 S29	
Wastage claimable	98.50		✓ Dii	flucan	
vvasiaye cialitable					

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOLE

by endorsement - Retail pharmacy - Specialist Specialist must be an infectious disease physician, clinical microbiologist,

clinical immunologist or dermatologist.

Oral liq 10 mg per ml — Special Authority see SA1322 below —
Retail pharmacy.......141.80

150 ml OP

Sporanox

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE

rab 200 mg - PCT - Retail pharmacy-Specialist - Subsid	уру		
endorsement	CBS	30	✓ Link Healthcare §29
			✓ Nizoral S29
Prescriptions must be written by, or on the recommend	dation of an oncolog	ist	

NYSTATIN

Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
·	(15.47)		Nilstat

	Subsidy (Manufacturer's Price	a) S	Fully	Brand or Generic	
	\$	Per	✓	Manufacturer	
POSACONAZOLE - Special Authority see SA1285 below - Reta	ail pharmacy				
Tab modified-release 100 mg	869.86	24	✓ N	oxafil	
Oral liq 40 mg per ml	761.13 1	105 ml O	P 🗸 N	oxafil	

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy: or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

IERBINAFINE			
* Tab 250 mg	1.33	14	✓ Deolate
VORICONAZOLE - Special Authority see SA1273 below - R	etail pharmacy		
Tab 50 mg	91.00	56	✓ Vttack
Tab 200 mg	350.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage			
claimable	1,437.00	70 ml	✓ Vfend

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis: or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	:	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
- 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
- 3.3 Patient has fluconazole resistant candidiasis; or
- 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

PRIMAQUINE PHOSPHATE - Special Authority see SA1684 below - Retail pharmacy

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Antiparasitics

Antiprotozoals

QUININE SUI PHATE

Antitrichomonal Agents

√	FD'	ノレニ	DA7	\sim	

C

Tab 200 mg - Up to 30 tab available on a PSO	10.45	100	Trichozole
Tab 400 mg - Up to 15 tab available on a PSO	18.15	100	Trichozole
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg	24.48	10	✓ Flagyl
ORNIDAZOLE			

Antituberculotics and Antileprotics

Tab 500 mg23.00

Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status.

10

✓ Arrow-Ornidazole

CLOFAZIMINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist.

		Subsidy (Manufacturer's Price)		Fully	
		(Manufacturer's Price) \$	Per	Subsidised	
CYCLOSERINE - Retail pharma	cv-Specialist				
a) No patient co-payment pa	, ,				
	ten by, or on the recommenda	ation of, an infectious d	iseas	e physicia	n, clinical microbiologist
Cap 250 mg		344.00	60	1	Cyclorin S29
APSONE - Retail pharmacy-Sp	pecialist				
a) No patient co-payment pa					
 b) Prescriptions must be writ dermatologist 	ten by, or on the recommenda	ation of, an infectious d	iseas	e physicia	n, clinical microbiologist
			100		Dapsone
Tab 100 mg		329.50	100	•	Dapsone
THAMBUTOL HYDROCHLORIE	DE - Retail pharmacy-Special	list			
 a) No patient co-payment pa 					
respiratory physician	ten by, or on the recommenda		iseas	se physicia	n, clinical microbiologist
Tab 100 mg		85.73	100	/	EMB Fatol \$29
Tab 400 mg		49.34	56	1	Myambutol \$29
SONIAZID - Retail pharmacy-Sp	pecialist				
 a) No patient co-payment pa 					
	ten by, or on the recommenda	ation of, an internal me	dicine	e physician	, paediatrician, clinical
	gist or public health physician	00.00	100	./	PSM
Tab 100 mg		22.00	100	•	PSIVI
ONIAZID WITH RIFAMPICIN -					
a) No patient co-payment pa b) Prescriptions must be write	yable ten by, or on the recommenda	ation of an internal me	dicina	nhveician	naediatrician clinical
	gist or public health physician	allon oi, an internal me	ulcille	priysician	, paediatriciari, cirricar
Tab 100 mg with rifampicin 1		85.54	100	1	Rifinah
Tab 150 mg with rifampicin 3			100	1	Rifinah
ARA-AMINO SALICYLIC ACID	- Retail pharmacy-Specialist				
a) No patient co-payment pa	yable				
	ten by, or on the recommenda	ation of, an infectious d	iseas	e specialis	t, clinical microbiologist
respiratory physician		000.00			D
· -		280.00	30	•	Paser S29
ROTIONAMIDE – Retail pharm					
a) No patient co-payment pa	,	stion of an infantiace of	io e = -	o one-!-!!	t aliaiaal misseleiste sist
b) Prescriptions must be writ respiratory physician	ten by, or on the recommenda	alion of, an intectious d	iseas	e specialis	si, cimicai microbiologist
		305.00	100	/	Peteha \$29
YRAZINAMIDE – Retail pharma			.00	•	
a) No patient co-payment pa	• •				
	ten by, or on the recommenda	ation of, an infectious d	iseas	e physicia	n, clinical microbiologist
€ Tab 500 mg		59.00	100	1	AFT-Pyrazinamide
IFABUTIN - Retail pharmacy-S					•
a) No patient co-payment pa					
	ten by, or on the recommenda	ation of, an infectious d	iseas	e physicia	n, respiratory physician o
gastroenterologist					
€ Cap 150 mg					Mycobutin

Subsidy	Fully Subsidised		Brand or	
(Manufacturer's Price)			Generic	
\$	Per	1	Manufacturer	

RIFAMPICIN - Subsidy by endorsement

- a) No patient co-payment payable
- b) For confirmed recurrent Staphylococcus aureus infection in combination with other effective anti-staphylococcal antimicrobial based on susceptibilities and the prescription is endorsed accordingly; can be waived by endorsement -Retail pharmacy - Specialist. Specialist must be an internal medicine physician, clinical microbiologist, dermatologist, paediatrician, or public health physician.

*	Cap 150 mg55.75	100	/	Rifadin
*	Cap 300 mg116.25	100	/	Rifadin
*	Oral liq 100 mg per 5 ml12.00	60 ml	✓	Rifadin

Antivirals

For eye preparations refer to Eye Preparations, Anti-Infective Preparations, page 228

Hepatitis B Treatment

⇒SA0829 Special Authority for Subsidy

Initial application only from a gastroenterologist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg+); and Documented resistance to lamivudine, defined as:
- 2 Patient has raised serum ALT (> 1 x ULN); and
- 3 Patient has HBV DNA greater than 100,000 copies per mL, or viral load 10 fold or higher over nadir; and
- 4 Detection of M204I or M204V mutation; and
- 5 Fither:
 - 5.1 Both
 - 5.1.1 Patient is cirrhotic; and
 - 5.1.2 adefovir dipivoxil to be used in combination with lamivudine: or
 - 5.2 Both:
 - 5.2.1 Patient is not cirrhotic; and
 - 5.2.2 adefovir dipivoxil to be used as monotherapy.

Renewal only from a gastroenterologist or infectious disease specialist. Approvals valid for 2 years where in the opinion of the treating physician, treatment remains appropriate and patient is benefiting from treatment.

Notes: Lamivudine should be added to adefovir dipivoxil if a patient develops documented resistance to adefovir dipivoxil, defined as:

- i) raised serum ALT (> 1 x ULN); and
- ii) HBV DNA greater than 100,000 copies per mL, or viral load 10 fold or higher over nadir; and
- iii) Detection of N236T or A181T/V mutation.

Adefovir dipivoxil should be stopped 6 months following HBeAg seroconversion for patients who were HBeAg+ prior to commencing adefovir dipivoxil.

The recommended dose of adefovir dipivoxil is no more than 10mg daily.

In patients with renal insufficiency adefovir dipivoxil dose should be reduced in accordance with the datasheet guidelines. Adefovir dipivoxil should be avoided in pregnant women and children.

ENTECAVIR

*	Tab 0.5 mg	52.00	30	 Entecavir Sandoz
LAN	MIVUDINE - Special Authority see SA1685 on the next page -	Retail pharma	су	
	Tab 100 mg	4.20	28	✓ Zetlam
	Oral liq 5 mg per ml	270.00	240 ml OP	✓ Zeffix

Subsidy	Fully	Brand or
	<u> </u>	
(Manutacturer's Price)	Subsidised	Generic
` ¢ ′	Por 🗸	Manufacturor

⇒SA1685 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B. TENOFOVIR DISOPROXII

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA1651., page 105

Herpesvirus Treatments

١.	\sim 1	\sim 1	OVIE	•
١	v	v	UVIE	١.

 * Tab dispersible 200 mg * Tab dispersible 400 mg * Tab dispersible 800 mg 	5.38	25 56 35	✓ <u>Lovir</u> ✓ <u>Lovir</u> ✓ <u>Lovir</u>
VALACICLOVIR Tab 500 mg Tab 1,000 mg		30 30	✓ <u>Vaclovir</u> ✓ <u>Vaclovir</u>
VALGANCICLOVIR – Special Authority see SA1404 below – Re Tab 450 mg		60	✓ <u>Valganciclovir</u> <u>Mylan</u>

⇒SA1404 Special Authority for Subsidy

Initial application — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
- 2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for

Subsidy	ı	Fully	Brand or
(Manufacturer's Price)			Generic
\$	Per	•	Manufacturer

continued...

3 months for applications meeting the following criteria:

Roth:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via PHARMAC's approved direct distribution supply. Further details can be found on

PHARMAC's website https://www.pharmac.govt.nz/hepatitis-c-treatments

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg......24,363.46 28 **✓ Harvoni**

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz/hepatitis-c-treatments or:

The Coordinator, Hepatitis C Treatment Panel

PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 460 4990.

Email: hepcpanel@pharmac.govt.nz

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA1842 on the next page

- a) Brand switch fee payable (Pharmacode 2573865) see page 233 for details
- b) Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 105 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the PHARMAC website.

Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a

_					
	Subsidy		Fully	Brand or	
	(Manufacturer's Price)	Sub	sidised	Generic	
	\$	Per	✓	Manufacturer	

⇒SA1842 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis, Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment: and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks: and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis, Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks: and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antiretrovirals

⇒SA1651 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — **(Confirmed HIV)** only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

continued...

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA1651 on the previous	page - Retail phar	macy	
Tab 50 mg	63.38	30	✓ Stocrin S29
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
Oral lig 30 mg per ml	145.79	180 ml OP	✓ Stocrin S29
(Stocrin S29 Tab 50 mg to be delisted 1 April 2020)			
(Stocrin S29 Oral liq 30 mg per ml to be delisted 1 August 20	20)		
ETRAVIRINE - Special Authority see SA1651 on the previous	s page – Retail pha	armacy	
Tab 200 mg	770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA1651 on the previous	s page – Retail pha	armacy	
Tab 200 mg	60.00	60	✓ <u>Nevirapine</u> <u>Alphapharm</u>
Oral suspension 10 mg per ml	203.55	240 ml	✓ Viramune Suspension

Nucleosides Reverse Transcriptase Inhibitors

ABACAVIR SULPHATE – Special Authority see SA1651 on the page 17 Tab 300 mg	180.00	Retail pharmad 60 240 ml OP	y ✓ <u>Ziagen</u> ✓ Ziagen	
ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority Note: abacavir with lamivudine (combination tablets) counts anti-retroviral Special Authority.	as two anti-retro	viral medicatio	ns for the purposes of the	
Tab 600 mg with lamivudine 300 mg	63.00	30	✓ <u>Kivexa</u>	
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPF	ROXIL - Special	Authority see	SA1651 on the previous page	e –
Retail pharmacy	000 () .	.,		
 a) Brand switch fee payable (Pharmacode 2573873) - see p 				
 Note: Efavirenz with emtricitabine and tenofovir disoprox the anti-retroviral Special Authority 		e anti-retroviral	medications for the purposes	s of
Tab 600 mg with emtricitabine 200 mg and tenofovir disoprox	cil			
245 mg (300 mg as a maleate)	106.88	30	✓ Mylan	
EMTRICITABINE - Special Authority see SA1651 on the previous	ıs page – Retail	pharmacy		
Con 000 ma	207.20	20	/ Emtrivo	

			<u>,</u>			
EMTRICITABINE – Special Authority see SA1651 on the Cap 200 mg		pharmacy 30	✓ Emtriva			

LAMIVUDINE – Special Authority see SA1651 on the previous page – Retail pharmacy						
Tab 150 mg	52.50	60	✓ Lamivudine Alphapharm			
Oral liq 10 mg per ml	102.50	240 ml OP	✓ 3TC			
ZIDOVUDINE [AZT] - Special Authority see SA1651 on the previous page - Retail pharmacy						
Cap 100 mg	152.25	100	✓ Retrovir			
Oral liq 10 mg per ml	30.45	200 ml OP	✓ Retrovir			

	Subsidy	-) 0:	Fully	Brand or		
	(Manufacturer's Pric \$	e) Subsi Per	aisea 🗸	Generic Manufacturer		
	Ф	rei		Manufacturer		
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority see SA1651 on page 105 – Retail pharmacy Note: zidovudine [AZT] with lamivudine (combination tablets) counts as two anti-retroviral medications for the purposes of the anti-retroviral Special Authority.						
Tab 300 mg with lamivudine 150 mg	33.00	60	✓ <u>A</u>	<u>phapharm</u>		
Protease Inhibitors						
ATAZANAVIR SULPHATE – Special Authority see SA1651 on pa Brand switch fee payable (Pharmacode 2573857) - see page	233 for details	,				
Cap 150 mg		60	√ <u>T</u> e			
Cap 200 mg	188.91	60	✓ Te	eva		
DARUNAVIR - Special Authority see SA1651 on page 105 - Reta	ail pharmacy					
Tab 400 mg	335.00	60	✓ Pı	rezista		
Tab 600 mg	476.00	60	✓ Pı	<u>rezista</u>		
LOPINAVIR WITH RITONAVIR - Special Authority see SA1651 of	n page 105 – Ret	ail pharmacy				
Tab 100 mg with ritonavir 25 mg		60	✓ Ka	aletra		
Tab 200 mg with ritonavir 50 mg		120	✓ K	aletra		
Oral lig 80 mg with ritonavir 20 mg per ml		300 ml OP	✓ K	aletra		
RITONAVIR - Special Authority see SA1651 on page 105 - Retai						
Tab 100 mg	, ,	30	✓ <u>N</u>	<u>orvir</u>		
Strand Transfer Inhibitors						
DOLUTEGRAVIR - Special Authority see SA1651 on page 105 - Tab 50 mg	, ,	30	✓ Ti	vicay		
RALTEGRAVIR POTASSIUM – Special Authority see SA1651 on page 105 – Retail pharmacy						
Tab 400 mg		60	√ ls	entress		
Tab 600 mg	,	60		entress HD		
1 ab 000 mg	1,030.00	00	- 15	CHUCOO HD		

Immune Modulators

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

- 1) Diagnosis
 - Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
 - PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
 - Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Exclusion Criteria

- Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia ($< 2.0 \times 10^9$) and/or thrombocytopenia.

Subsi	sidy Ful	ly Brand or
(Manufacture	rer's Price) Subsidise	d Generic
\$	Per	Manufacturer

continued...

4) Continuing alcohol abuse and/or continuing intravenous drug users.

Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

INTERFERON ALFA-2A - PCT - Retail pharmacy-Specialist

- a) See prescribing guideline on the previous page
- b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician or ophthalmologist
- - a) See prescribing guideline on the previous page
 - b) Note: PHARMAC will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at PHARMAC on 0800-023-588 option 4.

500.00 4 **✓ Pegasys**

✓ Roferon-A

⇒SA1400 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400.000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C. genotype 1: and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and

INFECTIONS - AGENTS FOR SYSTEMIC USE

S	Subsidy	Fully	Brand or
(Manufa	acturer's Price) Subsi	dised	Generic
	\$ Per	1	Manufacturer

continued...

- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
 - 4 Patient is to be treated in combination with boceprevir; and
 - 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use: and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Notes:

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

Urinary Tract Infections

HE	XAMINE HIPPURATE			
*	Tab 1 g	18.40	100	
	•	(40.01)		Hiprex
Nľ	TROFURANTOIN			
*	Tab 50 mg	22.20	100	✓ Nifuran
*	Tab 100 mg	37.50	100	✓ Nifuran
NC	PRFLOXACIN			
	Tab 400 mg - Subsidy by endorsement	135.00	100	✓ Arrow-Norfloxacin
	Only if prescribed for a patient with an uncomplicated uri	nary tract infection	n that is unre	esponsive to a first line agent or

with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy (Manufacturer's Price) \$) Per	Fully Subsidised	
Anticholinesterases				
NEOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	98.00	50	/	AstraZeneca
PYRIDOSTIGMINE BROMIDE				
▲ Tab 60 mg	45.79	100	•	Mestinon
Non-Steroidal Anti-Inflammatory Drugs				
DICLOFENAC SODIUM				
* Tab EC 25 mg	1.23	50	1	Diclofenac Sandoz
* Tab 50 mg dispersible	1.50	20	1	Voltaren D
* Tab EC 50 mg	1.23	50	1	Diclofenac Sandoz
* Tab long-acting 75 mg	22.80	500	•	Apo-Diclo SR
* Tab long-acting 100 mg	25.15	500	•	Apo-Diclo SR
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a l		5		Voltaren
* Suppos 12.5 mg		10	_	Voltaren
* Suppos 25 mg		10	_	Voltaren
* Suppos 50 mg – Up to 10 supp available on a PSO		10	_	Voltaren
* Suppos 100 mg	7.00	10	•	Voltaren
IBUPROFEN			_	
* Tab 200 mg		1,000		Relieve
* Tab long-acting 800 mg		30	_	Brufen SR
* Oral liq 20 mg per ml	1.88	200 m		<u>Ethics</u>
KETOPROFEN				
* Cap long-acting 200 mg	12.07	28	•	Oruvail SR
MEFENAMIC ACID				
* Cap 250 mg	1.25	50		
	(9.16)			Ponstan
	0.50	20		
	(5.60)			Ponstan
NAPROXEN				
* Tab 250 mg	32.69	500	✓	Noflam 250
* Tab 500 mg	22.19	250		Noflam 500
* Tab long-acting 750 mg	6.16	28		Naprosyn SR 750
* Tab long-acting 1 g	8.21	28	•	Naprosyn SR 1000
SULINDAC				
* Tab 100 mg	8.55	50	✓	Aclin
* Tab 200 mg	15.10	50	•	Aclin
TENOXICAM				
* Tab 20 mg	9.15	100	✓	<u>Tilcotil</u>
* Inj 20 mg vial		1	_	AFT
NSAIDs Other				
OFI FOOVID				
CELECOXIB	0.00	00		Oalahwasi
Cap 100 mg	3.63	60		Celebrex
Can 200 mg	0.00	20		Celebrar
Cap 200 mg	2.30	30		Celebrex Celecoxib Pfizer
(Celebrex Cap 100 mg to be delisted 1 September 2020)			•	OGIGCOXID FIIZEI

Subsidy		Fully	Brand or	Ī
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	1	Manufacturer	

Topical Products for Joint and Muscular Pain

CAPSAICIN

Crm 0.025% - Special Authority see SA1289 below - Retail 25 g OP ✓ Zostrix ✓ Zostrix 9.95 45 g OP

⇒SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

ntirhaumatoid Agente

7.98	100	✓ Plaquenil
		•
2.90	30	✓ Apo-Leflunomide
2.90	30	✓ Apo-Leflunomide
67.23	100	D-Penamine
110.12	100	✓ D-Penamine
76.87	10	✓ Myocrisin
113.17	10	✓ Myocrisin
217.23	10	Myocrisin
ed 1 March 2020)		
,		
ed 1 March 2020)		
		2.90 30 2.90 30

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

ALEINDI IOIVATE GODIOM		
* Tab 70 mg2.44	4	✓ Fosamax
ALENDRONATE SODIUM WITH COLECALCIFEROL		
* Tab 70 mg with colecalciferol 5,600 iu	4	✓ Fosamax Plus

Other Treatments

ALENDRONATE SODILIM

DENOSUMAB - Special Authority see SA1777 below - Retail pharm	асу		
Inj 60 mg prefilled syringe	326.00	1	✓ Prolia

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Fither:

continued...

✓ Fosamax Plus

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	sidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.1 The patient is female and postmenopausal; or
- 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

*	Tab 60 mg	53.76	28	Evista
RA	LOXIFENE HYDROCHLORIDE - Special Authority see SA1779	below - Retail	pharmacy	
	Inj 9 mg per ml, 10 ml vial	17.05	1	✓ Pamisol
	Inj 6 mg per ml, 10 ml vial	15.02	1	✓ Pamisol
	inj 3 mg per mi, 10 mi viai	5.98	1	✓ Pamisoi

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Subsidy (Manufacturer's Price)	Sul	Fully bsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

Subsidy (Manufacturer's	Fu Price) Subsidise	,	
\$	Per	Manufacturer	

continued...

- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

⇒SA1780 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy)

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
 \$	Per	•	Manufacturer

continued...

prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and

3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garyan) which incorporates BMD measurements (see Note); or
 - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene: and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a

ubsidy cturer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

continued...

fall from a standing height or less.

d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

ALLOPURINOL * Tab 100 mg	4.54	500	✓ DP-Allopurinol
* Tab 300 mg	10.35	500	✓ DP-Allopurinol
BENZBROMARONE - Special Authority see SA1537 below -	Retail pharmacy		
Tab 100 mg	45.00	100	✓ Benzbromaron AL
			100 \$29

⇒SA1537 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose: or
 - 2.3 Both:
 - 2.3.1 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Notes); and
 - 2.3.2 The patient has a rate of creatinine clearance greater than or equal to 20 ml/min; or
 - 2.4 All of the following:
 - 2.4.1 The patient is taking azathioprine and requires urate-lowering therapy; and
 - 2.4.2 Allopurinol is contraindicated; and
 - 2.4.3 Appropriate doses of probenecid are ineffective or probenecid cannot be used due to reduced renal function: and
- 3 The patient is receiving monthly liver function tests.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function

Notes: Benzbromarone has been associated with potentially fatal hepatotoxicity.

In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

The New Zealand Rheumatology Association has developed information for prescribers which can be accessed from its website at www.rheumatology.org.nz/home/resources-2/

\sim			NIE
\cup	LCH	IUI	IN⊏

* ∃	ab 500 mcg	9.58	100	✓ Colgout
------------	------------	------	-----	-----------

Per	•	Generic Manufacturer	
28	✓	Adenuric	
28	✓	Adenuric	
_	28	28	28 ✓ Adenuric

SA1538 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose: or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note).

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

PROBENECID

*	Tab 500 mg	55.00	100	✓ Probenecid-AFT
---	------------	-------	-----	------------------

Muscle Relaxants

$D \Lambda$	\sim	\sim	_	_	N I
BΑ	UЛ	()	ь.	_	IVI

*	Tab 10 mg	100	✓ Pacifen
	Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement11.55	1	✓ Lioresal Intrathecal
	Subsidised only for use in a programmable pump in patients where oral ant	tispastic a	gents have been ineffective or have
	caused intolerable side effects and the prescription is endorsed accordingly	/.	

Inj 2 mg per ml, 5 ml ampoule — Subsidy by endorsement............372.98 5 Medsurge
Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have caused intolerable side effects and the prescription is endorsed accordingly.

DANTROI ENF

Cap 25 mg	65.00	100	Dantrium
Cap 50 mg	77.00	100	✓ Dantrium S29 S29 ✓ Dantrium
ORPHENADRINE CITRATE			
Tab 100 mg	18.54	100	✓ Norflex

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

Agents for Parkinsonism and Related Disorders

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE			
▲ Cap 100 mg	38.24	60	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE			
▲ Inj 10 mg per ml, 2 ml ampoule	119.00	5	✓ Movapo
BROMOCRIPTINE MESYLATE			
* Tab 2.5 mg	32.08	100	✓ Apo-Bromocriptine
ENTACAPONE			
▲ Tab 200 mg	22.00	100	✓ Entapone
LEVODOPA WITH BENSERAZIDE			
* Tab dispersible 50 mg with benserazide 12.5 mg	13.25	100	Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg		100	Madopar 62.5
* Cap 100 mg with benserazide 25 mg		100	✓ Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg		100	✓ Madopar HBS
* Cap 200 mg with benserazide 50 mg	26.25	100	✓ Madopar 250
LEVODOPA WITH CARBIDOPA			
* Tab 100 mg with carbidopa 25 mg	17.97	100	✓ Kinson
			✓ Sinemet
* Tab long-acting 100 mg with carbidopa 25 mg		100	✓ Mylan S29
* Tab long-acting 200 mg with carbidopa 50 mg	37.15	100	✓ Sinemet CR
	46.73		✓ Mylan S29
* Tab 250 mg with carbidopa 25 mg	32.67	100	✓ Sinemet
PRAMIPEXOLE HYDROCHLORIDE			
▲ Tab 0.25 mg	6.12	100	✓ Ramipex
▲ Tab 1 mg	20.73	100	✓ Ramipex
ROPINIROLE HYDROCHLORIDE			
▲ Tab 0.25 mg	2.78	100	✓ Apo-Ropinirole
	2.85	84	✓ Ropin
▲ Tab 1 mg	3.95	84	✓ Ropin
	5.00	100	✓ Apo-Ropinirole
▲ Tab 2 mg		84	Ropin
A. Tab Face	7.72	100	✓ Apo-Ropinirole
▲ Tab 5 mg	12.50	84 100	✓ Ropin
(Ann Danining Tab 0.05 may to be delicted 1 Mayob 0000)	10.01	100	✓ Apo-Ropinirole
(Apo-Ropinirole Tab 0.25 mg to be delisted 1 March 2020) (Apo-Ropinirole Tab 1 mg to be delisted 1 March 2020)			
(Apo-Ropinirole Tab 2 mg to be delisted 1 March 2020)			
(Apo-Ropinirole Tab 5 mg to be delisted 1 March 2020)			
SELEGILINE HYDROCHLORIDE			
* Tab 5 mg	22.00	100	✓ Apo-Selegiline
Tab 5 mg	22.00	100	S29 S29
TOLOADONE			323 020
TOLCAPONE A Tol 100 mg	100 50	100	./ Taamar
▲ Tab 100 mg	13∠.30	100	✓ Tasmar

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
Anticholinergics	Ψ	rei	<u> </u>	Manuacturei
BENZATROPINE MESYLATE Tab 2 mg Inj 1 mg per ml, 2 ml	95.00	60	1	Benztrop Cogentin
a) Up to 10 inj available on a PSO b) Only on a PSO	190.00	10		Omega
PROCYCLIDINE HYDROCHLORIDE Tab 5 mg	7.40	100	•	Kemadrin
Agents for Essential Tremor, Chorea and Relate	d Disorders			
RILUZOLE – Special Authority see SA1403 below – Retail pharm Wastage claimable Tab 50 mg	•	56	✓	Rilutek
 ▶SA1403 Special Authority for Subsidy Initial application only from a neurologist or respiratory specialist following criteria: All of the following: 1 The patient has amyotrophic lateral sclerosis with disease 2 The patient has at least 60 percent of predicted forced vita 3 The patient has not undergone a tracheostomy; and 4 The patient has not experienced respiratory failure; and 5 Any of the following:	duration of 5 years o	or less onths	; and prior to the	initial application; and
Tab 25 mg	91.10	112	✓ !	<u>Motetis</u>
Anaesthetics Local				
LIDOCAINE [LIGNOCAINE] Gel 2%, tube — Subsidy by endorsement	dministration and the	25	cription is e	<u>Cathejell</u>

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

	Subsidy (Manufacturer's Price)	Fully	
	\$	Per	✓	
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%	38.00	200 m	ı 🗸	Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	8.75	25	/	Lidocaine-Claris
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	8.25	25	✓	Lidocaine-Claris
Inj 1%, 20 ml ampoule – Up to 5 inj available on a PSO	12.00	5		
	(20.00)			Xylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	6.20	5	/	Lidocaine-Claris
Inj 2%, 20 ml vial - Up to 5 inj available on a PSO	6.45	5	/	Lidocaine-Claris
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –				
Subsidy by endorsement	81.50	10	1	Pfizer
a) Up to 5 each available on a PSO				

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] - Special Authority see SA0906 abo	ve – Retail phar	macy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Author	ority see SA0906	above – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	✓ EMLA

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

Non-opioid Analgesics

For aspirin & chloroform application refer Standard Formulae, page 235

ASPIRIN

*	Tab dispersible 300 mg	- Up to 30 ta	b available on a PSO	4.50	100	✓ Ethics Aspirin
---	------------------------	---------------	----------------------	------	-----	------------------

CAPSAICIN - Subsidy by endorsement

Subsidised only if prescribed for post-herpetic neuralgia or diabetic peripheral neuropathy and the prescription is endorsed accordingly.

Crm 0.075%	12.50	45 a OP	✓ Zostrix HP
GIII 0.073 /6		40 u OF	▼ ∠USUIX FIF

NEFORAM HYDROCHLORIDE

ALI OI /IMITI DI IOONEONIDE			
Tab 30 mg	23.40	90	Acupan

	Subsidy	Duit 1 Out -	Fully Brand or
	(Manufacturer's F \$	Per Per	sidised Generic Manufacturer
RACETAMOL			
Tab 500 mg - blister pack - Up to 30 tab available on a PS	7.12	1,000	✓ Paracetamol Pharmacare
			✓ <u>Pharmacare</u>✓ Pharmacy Health
Tab 500 mg - bottle pack	6.32	1,000	✓ Pharmacare
Oral liq 120 mg per 5 ml		1,000 ml	✓ Paracare
a) Up to 200 ml available on a PSO			
b) Not in combination			_
Oral liq 250 mg per 5 ml	5.81	1,000 ml	✓ Paracare Double
a). Un to 100 ml available on a BCO			<u>Strength</u>
a) Up to 100 ml available on a PSO b) Not in combination			
Suppos 125 mg	3 29	10	✓ Gacet
Suppos 250 mg		10	✓ Gacet
Suppos 500 mg		50	✓ Gacet
harmacy Health Tab 500 mg - blister pack to be delisted 1 Ja			<u></u>
Opioid Analgesics			
DDEINE PHOSPHATE – Safety medicine; prescriber may de	tarmina disnansin	na frequency	
Tab 15 mg		100	✓ PSM
Tab 30 mg		100	✓ PSM
Tab 60 mg		100	✓ PSM
HYDROCODEINE TARTRATE			
Tab long-acting 60 mg	8.60	60	✓ <u>DHC Continus</u>
NTANYL			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing f			
Inj 50 mcg per ml, 2 ml ampoule		10	✓ Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule		10	Boucher and Muir
Patch 12.5 mcg per hour		5 5	✓ Fentanyl Sandoz
Patch 25 mcg per hour Patch 50 mcg per hour		5 5	 ✓ Fentanyl Sandoz ✓ Fentanyl Sandoz
Patch 75 mcg per hour		5	✓ Fentanyl Sandoz
Patch 100 mcg per hour		5	✓ Fentanyl Sandoz
THADONE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing f	requency		
 d) Extemporaneously compounded methadone will only be (methadone powder, not methadone tablets). 		e rate of the ch	neapest form available
e) For methadone hydrochloride oral liquid refer Standard	Formulae, page 2	235	
Tab 5 mg		10	✓ <u>Methatabs</u>
Tab 5 mg - bottle pack		10	Methatabs
Oral liq 2 mg per ml		200 ml	✓ <u>Biodone</u>
Oral liq 5 mg per ml		200 ml	✓ Biodone Forte
Oral liq 10 mg per ml		200 ml	✓ Biodone Extra Forte
Inj 10 mg per ml, 1 ml	61.00	10	✓ AFT

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Price	e) Su	bsidised	Generic
	\$	Per	✓	Manufacturer
MORPHINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing from		000 1		
Oral liq 1 mg per ml		200 ml		RA-Morph
Oral liq 2 mg per ml		200 ml	_	RA-Morph
Oral liq 5 mg per ml	19.44	200 ml	-	Ordine S29
			✓ <u>F</u>	RA-Morph
Oral lig 10 mg per ml	27.74	200 ml	✓ (Ordine S29
			√ F	RA-Morph
MORPHINE SULPHATE			_	
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing from				
Tab immediate-release 10 mg		10		Sevredol
Tab long-acting 10 mg		10		Arrow-Morphine LA
Tab immediate-release 20 mg		10		Sevredol
Tab long-acting 30 mg		10		Arrow-Morphine LA
Tab long-acting 60 mg		10		Arrow-Morphine LA
Tab long-acting 100 mg		10		Arrow-Morphine LA
Cap long-acting 10 mg	2.05	10	✓ n	n-Eslon
m-Eslon to be Sole Supply on 1 January 2020				
Cap long-acting 30 mg	3.00	10	✓ n	n-Eslon
m-Eslon to be Sole Supply on 1 January 2020				
Cap long-acting 60 mg	6.12	10	✓ n	n-Eslon
m-Eslon to be Sole Supply on 1 January 2020				
Cap long-acting 100 mg	7.13	10	✓ n	n-Eslon
m-Eslon to be Sole Supply on 1 January 2020				
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a Ps	SO6.27	5	✓ [BL Morphine
, , , , , , , , , , , , , , , , , , , ,			_	Sulphate
Inj 10 mg per ml, 1 ml ampoule - Up to 5 inj available on a F	PSO 4.47	5	✓ [BL Morphine
ing to mg per mily this amposite to pite only available on a .		ŭ	=	Sulphate
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a F	990 4.76	5	√ Γ	BL Morphine
ing 13 mg per mi, 1 mi ampodie – op to 3 mg available on a r	304.70	J	• •	Sulphate
Ini 20 mg normi 1 ml omnovio . Linto E ini quailable on a l	200 640	E	./ -	
Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on a F	-206.19	5	V L	BL Morphine
				<u>Sulphate</u>
MORPHINE TARTRATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fro	equency			
Inj 80 mg per ml, 1.5 ml ampoule		5	✓ [BL Morphine
, 01				Tartrate

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	S	ubsidised	Generic
	\$	Per	•	Manufacturer
OXYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	auency			
Tab controlled-release 5 mg		20	1	Oxycodone Sandoz
Tab controlled-release 10 mg		20		Oxycodone Sandoz
· ·				
Tab controlled-release 20 mg		20		Oxycodone Sandoz
Tab controlled-release 40 mg		20		Oxycodone Sandoz
Tab controlled-release 80 mg		20		Oxycodone Sandoz
Cap immediate-release 5 mg		20		OxyNorm
Cap immediate-release 10 mg		20		<u>OxyNorm</u>
Cap immediate-release 20 mg		20		OxyNorm
Oral liq 5 mg per 5 ml		250 ml		OxyNorm
Inj 10 mg per ml, 1 ml ampoule	7.28	5	1	<u>OxyNorm</u>
Inj 10 mg per ml, 2 ml ampoule	14.36	5	1	<u>OxyNorm</u>
Inj 50 mg per ml, 1 ml ampoule	30.60	5	1	OxyNorm
PARACETAMOL WITH CODEINE - Safety medicine; prescriber	may determine dispe	ensina t	requency	I
* Tab paracetamol 500 mg with codeine phosphate 8 mg		1.000		Paracetamol +
Tab paracetamor 500 mg with codeline phospitate o mg	10.21	1,000	•	Codeine (Relieve)
				Codellie (Helleve)
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	quency			
Tab 50 mg	4.46	10	1	PSM
Inj 50 mg per ml, 1 ml ampoule - Up to 5 inj available on a P	SO 4.98	5	1	DBL Pethidine
,				Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a P	SO 512	5	1	DBL Pethidine
ing so mg per mi, 2 mi ampedie — op to o mj avaliable on a r	000.12	J	-	Hydrochloride
				<u> </u>
TRAMADOL HYDROCHLORIDE				
Tab sustained-release 100 mg		20		Tramal SR 100
Tab sustained-release 150 mg	2.10	20		Tramal SR 150
Tab sustained-release 200 mg		20		Tramal SR 200
Cap 50 mg	2.25	100	✓.	Arrow-Tramadol
Antidepressants				
<u> </u>				
Cyclic and Related Agents				
AMITRIPTYLINE - Safety medicine; prescriber may determine di	isnensing frequency			
Tab 10 mg		100	1	Arrow-Amitriptyline
Tab 25 mg		100		Arrow-Amitriptyline
ŭ		100		Arrow-Amitriptyline
Tab 50 mg	2.31	100	•	Arrow-Ammurptynne

CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; prescriber may determine dispensing frequency

100

50

100

9.46

✓ Apo-Clomipramine✓ Apo-Clomipramine

✓ Apo-Clomipramine

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	S Per	Fully subsidised	Brand or Generic Manufacturer
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy by e	ndorsement			
a) Safety medicine; prescriber may determine dispensing f	requency			
b) Subsidy by endorsement - Subsidised for patients who	were taking dosulepin	[dothie	pin] hydro	ochloride prior to 1 June
2019 and the prescription is endorsed accordingly. Pha		the pr	escription	n as endorsed where there
exists a record of prior dispensing of dosulepin [dothiepi	n] hydrochloride.			
Tab 75 mg		100		Dopress
Cap 25 mg	6.45	100	•	Dopress
(Dopress Tab 75 mg to be delisted 1 August 2020)				
(Dopress Cap 25 mg to be delisted 1 January 2020)				
DOXEPIN HYDROCHLORIDE - Subsidy by endorsement				
a) Safety medicine; prescriber may determine dispensing f				
b) Subsidy by endorsement – Subsidised for patients who				
prescription is endorsed accordingly. Pharmacists may	annotate the prescript	ion as e	endorsed	where there exists a recor
of prior dispensing of doxepin hydrochloride.	0.00	400		
Cap 10 mg		100		Anten
Cap 25 mg		100 100		Anten Anten
Cap 50 mg	0.55	100	•	Anten
(Anten Cap 10 mg to be delisted 1 January 2020) (Anten Cap 25 mg to be delisted 1 April 2020)				
(Anten Cap 23 mg to be delisted 1 April 2020) (Anten Cap 50 mg to be delisted 1 May 2020)				
	and the second second second second			
IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescribe		-		
Tab 10 mg	10.96	50 100	_	Tofranil Tofranil
Tab 25 mg		50		Tofranil
MAPROTILINE HYDROCHLORIDE – Safety medicine; prescrit				
Tab 25 mg		30 30		cy Ludiomil
Tab 25 mg	12.53	50		Ludiomil
	25.06	100		Ludiomil
Tab 75 mg		20		Ludiomil
· · - · · · · · · · · · · · · · · ·	21.01	30		Ludiomil
NORTRIPTYLINE HYDROCHLORIDE - Safety medicine; presi	criher may determine o	lisnens	ina freau	ency
Tab 10 mg		100		Norpress
Tab 25 mg		180		Norpress
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	Selective			
PHENELZINE SULPHATE				
* Tab 15 mg	70.80	60	1	Nardil S29 S29
· ·	118.00	100	✓	Nardil
TRANYLCYPROMINE SULPHATE				
* Tab 10 mg	12 85	28	1	Parnate S29 S29
Tab to my	22.94	50		Parnate
	96.00	100	_	Parnate S29 S29
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE				
* Tab 150 mg	6.40	60	1	<u>Aurorix</u>
•		60		Aurorix
* Tab 300 mg				

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
Selective Serotonin Reuptake Inhibitors			
CITALOPRAM HYDROBROMIDE			
* Tab 20 mg	1.52	84	✓ PSM Citalopram
ESCITALOPRAM			
* Tab 10 mg	1.11	28	✓ Escitalopram- Apotex
* Tab 20 mg	1.90	28	✓ <u>Escitalopram-</u> <u>Apotex</u>
FLUOXETINE HYDROCHLORIDE			
* Tab dispersible 20 mg, scored - Subsidy by endorsement	1.98 2.47	30	✓ Fluox✓ Arrow-Fluoxetine
Subsidised by endorsement 1) When prescribed for a patient who cannot swallo accordingly; or 2) When prescribed in a daily dose that is not a mul endorsed. Note: Tablets should be combined w	tiple of 20 mg in which	case	the prescription is deemed to be
₩ Con 20 mg	1.00	90	✓ Arrow-Fluoxetine
* Cap 20 mg	2.91	90 84	✓ Fluox
'Arrow-Fluoxetine Cap 20 mg to be delisted 1 April 2020) PAROXETINE * Tab 20 mg	3.61 4.02	90	✓ Loxamine ✓ Apo-Paroxetine
Apo-Paroxetine Tab 20 mg to be delisted 1 March 2020)	1.02		- Apo Faroxoniio
	0.92	30	✓ Setrona
* Tab 50 mg	0.92 3.05	30 90	✓ Setrona ✓ Arrow-Sertraline
	3.05		***************************************
· ·	3.05	90	✓ Arrow-Sertraline
* Tab 100 mg	3.05 1.61	90 30	✓ Arrow-Sertraline✓ Setrona
•	3.05 1.61	90 30	✓ Arrow-Sertraline✓ Setrona
* Tab 100 mg	3.05 1.61	90 30	✓ Arrow-Sertraline✓ Setrona
* Tab 100 mg	3.05 1.61 5.25	90 30 90 30	✓ Arrow-Sertraline ✓ Setrona ✓ Arrow-Sertraline ✓ Apo-Mirtazapine
* Tab 100 mg	3.05 1.61 5.25	90 30 90	✓ Arrow-Sertraline ✓ Setrona ✓ Arrow-Sertraline
* Tab 100 mg	3.05 1.61 5.25 2.63 3.48	90 30 90 30 30 30	✓ Arrow-Sertraline ✓ Setrona ✓ Arrow-Sertraline ✓ Apo-Mirtazapine ✓ Apo-Mirtazapine
* Tab 100 mg	3.05 	90 30 90 30 30 30 84	✓ Arrow-Sertraline ✓ Setrona ✓ Arrow-Sertraline ✓ Apo-Mirtazapine ✓ Apo-Mirtazapine ✓ Enlafax XR
* Tab 100 mg 'Arrow-Sertraline Tab 50 mg to be delisted 1 March 2020) 'Arrow-Sertraline Tab 100 mg to be delisted 1 March 2020) Other Antidepressants MIRTAZAPINE Tab 30 mg	3.05 	90 30 90 30 30 30 84 84	✓ Arrow-Sertraline ✓ Setrona ✓ Arrow-Sertraline ✓ Apo-Mirtazapine ✓ Apo-Mirtazapine ✓ Enlafax XR ✓ Enlafax XR
* Tab 100 mg	3.05 	90 30 90 30 30 30 84	✓ Arrow-Sertraline ✓ Setrona ✓ Arrow-Sertraline ✓ Apo-Mirtazapine ✓ Apo-Mirtazapine ✓ Enlafax XR
* Tab 100 mg 'Arrow-Sertraline Tab 50 mg to be delisted 1 March 2020) 'Arrow-Sertraline Tab 100 mg to be delisted 1 March 2020) Other Antidepressants MIRTAZAPINE Tab 30 mg	3.05 	90 30 90 30 30 30 84 84	✓ Arrow-Sertraline ✓ Setrona ✓ Arrow-Sertraline ✓ Apo-Mirtazapine ✓ Apo-Mirtazapine ✓ Enlafax XR ✓ Enlafax XR
* Tab 100 mg	3.05 	90 30 90 30 30 30 84 84	✓ Arrow-Sertraline ✓ Setrona ✓ Arrow-Sertraline ✓ Apo-Mirtazapine ✓ Apo-Mirtazapine ✓ Enlafax XR ✓ Enlafax XR
* Tab 100 mg	3.05	90 30 90 30 30 30 84 84	✓ Arrow-Sertraline ✓ Setrona ✓ Arrow-Sertraline ✓ Apo-Mirtazapine ✓ Apo-Mirtazapine ✓ Enlafax XR ✓ Enlafax XR

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
DIAZEPAM - Safety medicine; prescriber may determine dispensing frequency			
Inj 5 mg per ml, 2 ml ampoule - Subsidy by endorsement11.83	5	1	Hospira
a) Up to 5 inj available on a PSO			
b) Only on a PSO			
c) PSO must be endorsed "not for anaesthetic procedures".		_	
Rectal tubes 5 mg - Up to 5 tube available on a PSO40.87	5		Stesolid
Rectal tubes 10 mg - Up to 5 tube available on a PSO40.87	5	/	Stesolid
PARALDEHYDE			
* Inj 5 ml	5	1	AFT S29
PHENYTOIN SODIUM			
* Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO 88.63	5	1	Hospira
* Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a			
PSO	5	1	Hospira
			r

Control of Epilepsy

CARBAMAZEPINE			
* Tab 200 mg	14.53	100	Tegretol
* Tab long-acting 200 mg	16.98	100	Tegretol CR
* Tab 400 mg	34.58	100	✓ Tegretol
* Tab long-acting 400 mg	39.17	100	✓ Tegretol CR
* Oral liq 20 mg per ml	26.37	250 ml	✓ Tegretol
CLOBAZAM - Safety medicine; prescriber may determine dispens	sing frequency		
Tab 10 mg		50	✓ Frisium
CLONAZEPAM - Safety medicine; prescriber may determine disp		nov	
Oral drops 2.5 mg per ml		10 ml OP	✓ Rivotril
	1.30	10 IIII OF	HIVOUII
ETHOSUXIMIDE			_
Cap 250 mg		100	Zarontin
Oral liq 250 mg per 5 ml	56.35	200 ml	Zarontin
GABAPENTIN			
Note: Not subsidised in combination with subsidised pregabal	in		
* Cap 100 mg		100	✓ Apo-Gabapentin
* Cap 300 mg		100	✓ Apo-Gabapentin
* Cap 400 mg		100	✓ Apo-Gabapentin
LACOSAMIDE – Special Authority see SA1125 below – Retail pha			
▲ Tab 50 mg	•	14	✓ Vimpat
▲ Tab 100 mg		14	✓ Vimpat
_ 100 100 mg	200.24	56	✓ Vimpat
▲ Tab 150 mg		14	✓ Vimpat
Tab 190 mg	300.40	56	✓ Vimpat ✓ Vimpat
A Tob 200 mg		56	•
▲ Tab 200 mg	400.55	96	✓ Vimpat

⇒SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

LAN	Ю.	TR	G	N	Е

LA	WICTRIGINE			
\blacktriangle	Tab dispersible 2 mg	6.74	30	✓ Lamictal
\blacktriangle	Tab dispersible 5 mg	9.64	30	✓ Lamictal
		15.00	56	✓ Arrow-Lamotrigine
\blacktriangle	Tab dispersible 25 mg - Brand switch fee payable			•
	(Pharmacode 2575949) - see page 233 for details	2.76	56	✓ Logem
\blacktriangle	Tab dispersible 50 mg - Brand switch fee payable			
_	(Pharmacode 2575949) - see page 233 for details	3.31	56	✓ Logem
•	Tab dispersible 100 mg – Brand switch fee payable			
_	(Pharmacode 2575949) - see page 233 for details	4 40	56	✓ Logem
	, , ,		00	Logom
LE	VETIRACETAM Table 250 mm	4.00	00	/ Franch
	Tab 250 mg		60	✓ Everet
	Tab 500 mg		60	✓ Everet
	Tab 750 mg		60	✓ <u>Everet</u>
	Tab 1,000 mg		60	✓ Everet
	Oral liq 100 mg per ml	44.78	300 ml OP	✓ Levetiracetam-AFT
PH	ENOBARBITONE			
	For phenobarbitone oral liquid refer Standard Formulae, p	age 235		
*	Tab 15 mg	40.00	500	✓ PSM
*	Tab 30 mg	40.00	500	✓ PSM
РН	ENYTOIN SODIUM			
	Tab 50 mg	75.00	200	✓ Dilantin Infatab
•	Cap 30 mg		200	✓ Dilantin
	Cap 100 mg		200	✓ Dilantin
*	Oral liq 30 mg per 5 ml		500 ml	✓ Dilantin
	EGABALIN		000	
ГП	Note: Not subsidised in combination with subsidised gaba	nontin		
×	•	•	56	✓ Drogobolin Dfizor
	Cap 25 mg		56	✓ Pregabalin Pfizer
*				✓ Pregabalin Pfizer
*	Cap 150 mg		56 50	✓ Pregabalin Pfizer
*	Cap 300 mg	/.38	56	✓ <u>Pregabalin Pfizer</u>
PR	IMIDONE			
*	Tab 250 mg	17.25	100	✓ Apo-Primidone
		62.00	200	✓ Mysoline S29 S29

	Subsidy		Fully	Brand or	
	(Manufacturer's Price	e)	Subsidised	Generic	
	\$	Per	1	Manufacturer	
SODIUM VALPROATE					
Tab 100 mg	13.65	100	✓	Epilim Crushable	
Tab 200 mg EC	27.44	100	✓	Epilim	
Tab 500 mg EC	52.24	100	✓	Epilim	
* Oral lig 200 mg per 5 ml	20.48	300 m	 	Epilim S/F Liquid	
, ,,			✓	Epilim Syrup	
* Inj 100 mg per ml, 4 ml	41.50	1	1	Epilim IV	
STIRIPENTOL - Special Authority see SA1330 below - Retail p	harmacy				
Cap 250 mg	509.29	60	1	Diacomit S29	
Powder for oral liq 250 mg sachet	509.29	60	✓	Diacomit S29	

⇒SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Patient has confirmed diagnosis of Dravet syndrome; and

▲ Tab 25 mg11.07

▲ Sprinkle cap 15 mg.......20.84

▲ Sprinkle cap 25 mg.......26.04

VIGABATRIN − Special Authority see SA1072 below − Retail pharmacy

▲ Tab 500 mg.......119.30

2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

60

60

100

✓ Arrow-Topiramate

✓ Topamax✓ Topamax

✓ Sabril

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

				Topiramate Actavis
		26.04		✓ Topamax
\blacktriangle	Tab 50 mg	18.81	60	✓ Arrow-Topiramate
				✓ Topiramate Actavis
		44.26		✓ Topamax
\blacktriangle	Tab 100 mg	31.99	60	✓ Arrow-Topiramate
	•			✓ Topiramate Actavis
		75.25		✓ Topamax
\blacktriangle	Tab 200 mg	55.19	60	✓ Arrow-Topiramate
	•			✓ Topiramate Actavis
		129.85		✓ Topamax

⇒SA1072 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

1 Fither:

TOPIRAMATE

- 1.1 Patient has infantile spasms; or
- 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Fither:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and

			NER	VOUS SYSTEM
	Subsidy (Manufacturer's Price) \$	Sub:	Fully sidised	Brand or Generic Manufacturer
continued				
2 Either:				
6-monthly basis thereafter); or 2.2 It is impractical or impossible (due to comorbid cond Notes: ``Optimal treatment with other antiepilepsy agents" is define indicated and clinically appropriate for the patient, given in adequating the pharmacokinetics of the drug with good evidence of a Vigabatrin is associated with a risk of irreversible visual field defeatenewal from any relevant practitioner. Approvals valid without following criteria: Both:	ned as treatment with the doses for the pati compliance. ts, which may be as	n other an ent's age ymptoma	tiepileps , weight tic in the	sy agents which are , and other features e early stages.
1 The patient has demonstrated a significant and sustained i2 Either:	mprovement in seizu	ire rate oi	severit	y and or quality of life; and
 Patient is receiving regular automated visual field to of treatment with vigabatrin; or 		,		
2.2 It is impractical or impossible (due to comorbid cond	,	•		
Notes: As a guideline, clinical trials have referred to a notional 50 anticonvulsant therapy and have assessed quality of life from the			ncy as a	n indicator of success with
Vigabatrin is associated with a risk of irreversible visual field defect	ts, which may be as	ymptoma	tic in the	e early stages.
Antimigraine Preparations				

Antimigraine Preparations			
For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page	ge 110		
Acute Migraine Treatment			
ERGOTAMINE TARTRATE WITH CAFFEINE Tab 1 mg with caffeine 100 mg	31.00	100	✓ Cafergot ✓ Cafergot S29 S29
RIZATRIPTAN Tab prodimersible 10 mg	E 06	30	✓ Rizamelt
Tab orodispersible 10 mg SUMATRIPTAN	3.20	30	<u> </u>
Tab 50 mg	24.44	100	✓ Apo-Sumatriptan
Tab 100 mg		100	✓ Apo-Sumatriptan
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj per			
prescription	42.67	2 OP	✓ Sun Pharma S29
	81.15		✓ Clustran
Prophylaxis of Migraine			
For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSPIZOTIFEN	STEM, page 48		
* Tab 500 mcg	23.21	100	✓ Sandomigran
Antinausea and Vertigo Agents			
For Antispasmodics refer to ALIMENTARY TRACT, page 8			

✓ Emend Tri-Pack

3 OP

APREPITANT – Special Authority see SA0987 on the next page – Retail pharmacy Cap 2 × 80 mg and 1 × 125 mg.......84.00



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA0987 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLOR	RIDE
--------------------------	------

* Tab 16 mg	2.89	84	✓ Vergo 16
CYCLIZINE HYDROCHLORIDE			
Tab 50 mg	0.55	10	✓ Nausicalm
CYCLIZINE LACTATE			
Inj 50 mg per ml, 1 ml	14.95	5	✓ Nausicalm
DOMPERIDONE			
* Tab 10 mg	2.25	100	✓ Pharmacy Health
HYOSCINE HYDROBROMIDE			
* Inj 400 mcg per ml, 1 ml ampoule	46.50	5	✓ Hospira
	93.00	10	✓ Martindale S29
Patch 1.5 mg - Special Authority see SA1387 below - Retail			
pharmacy	14.11	2	Scopoderm TTS

⇒SA1387 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

1 20

100

ME	FOCLOPRAMIDE HYDROCHLORIDE
N/e	Tab 40

* Tab 10 mg1.30) 100	Actavis 10
* Inj 5 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO9.50	10	✓ Pfizer
13.56	3	✓ Link Healthcare S29
Pfizer to be Sole Supply on 1 January 2020		
(Link Healthcare 29 Inj 5 mg per ml, 2 ml ampoule to be delisted 1 January	(2020)	
ONDANSETRON		
* Tab 4 mg	3 50	✓ Onrex
3.36	6	✓ Apo-Ondansetron
* Tab disp 4 mg - Up to 10 tab available on a PSO0.95	5 10	✓ Ondansetron
		ODT-ORLA
* Tab 8 mg4.57	7 50	Onrex
4.77	7	Apo-Ondansetron
* Tab disp 8 mg - Up to 10 tab available on a PSO1.43	3 10	✓ Ondansetron
		ODT-DRLA

(Apo-Ondansetron Tab 4 mg to be delisted 1 April 2020)

(Apo-Ondansetron Tab 8 mg to be delisted 1 April 2020)

	Subsidy (Manufacturer's Price)		Fully Brand or Subsidised Generic
	\$	Per	✓ Manufacturer
PROCHLORPERAZINE			
* Tab 3 mg buccal	5.97	50	
	(15.00)		Buccastem
* Tab 5 mg - Up to 30 tab available on a PSO	6.35 [′]	250	✓ Nausafix
* Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO	25.81	10	✓ Stemetil
Antipsychotics			
General			
AMICUL DDIDE Cofety medicines prescriber may determine a	liananaina fraguanas		
AMISULPRIDE – Safety medicine; prescriber may determine of Tab 100 mg		30	✓ Sulprix
Tab 100 mg		60	✓ <u>Sulprix</u> ✓ Sulprix
Tab 400 mg		60	✓ <u>Sulprix</u> ✓ Sulprix
Sulprix to be Sole Supply on 1 February 2020	29.70	00	Juiprix
Oral liq 100 mg per ml	65 53	60 ml	✓ Solian
(Solian Oral lig 100 mg per ml to be delisted 1 July 2020)		00 1111	• Johan
	diamanaina fua aa.		
ARIPIPRAZOLE – Safety medicine; prescriber may determine		20	Aviningonala Candan
Tab 5 mg Tab 10 mg		30 30	 ✓ <u>Aripiprazole Sandoz</u> ✓ Aripiprazole Sandoz
Tab 15 mg		30	✓ Aripiprazole Sandoz ✓ Aripiprazole Sandoz
Tab 20 mg		30	✓ Aripiprazole Sandoz ✓ Aripiprazole Sandoz
Tab 30 mg		30	✓ Aripiprazole Sandoz
· ·			
CHLORPROMAZINE HYDROCHLORIDE – Safety medicine; p	•		
Tab 10 mg – Up to 30 tab available on a PSO	14.83	100	✓ Largactil
Largactil to be Sole Supply on 1 January 2020 Tab 25 mg - Up to 30 tab available on a PSO	15.60	100	✓ Largactil
Largactil to be Sole Supply on 1 January 2020	13.02	100	Largaciii
Tab 100 mg – Up to 30 tab available on a PSO	36.73	100	✓ Largactil
Largactil to be Sole Supply on 1 January 2020		100	Largactii
Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO	30.79	10	✓ Largactil
Largactil to be Sole Supply on 1 January 2020			- Largaotti
CLOZAPINE – Hospital pharmacy [HP4]			
Safety medicine; prescriber may determine dispensing frequency	ulenov		
Tab 25 mg	' '	50	✓ Clozaril
1 db 25 mg	6.69	00	✓ Clopine
	11.36	100	✓ Clozaril
	13.37		✓ Clopine
Tab 50 mg		50	✓ Clopine
	17.33	100	✓ Clopine
Tab 100 mg	14.73	50	✓ Clozaril
· ·	17.33		✓ Clopine
	29.45	100	✓ Clozaril
	34.65		✓ Clopine
Tab 200 mg	34.65	50	✓ Clopine
	69.30	100	✓ Clopine
Suspension 50 mg per ml	17.33	100 m	✓ Clopine

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	0.4-14.		F. III	. D
	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	✓	Manufacturer
HALOPERIDOL – Safety medicine; prescriber may determine di	isnensing frequency			
Tab 500 mcg — Up to 30 tab available on a PSO		100	1	Serenace
Tab 1.5 mg – Up to 30 tab available on a PSO		100		Serenace
Tab 5 mg - Up to 30 tab available on a PSO		100		Serenace
Oral liq 2 mg per ml — Up to 200 ml available on a PSO		100 m	_	Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a P		10	•	Serenace
LEVOMEPROMAZINE HYDROCHLORIDE – Safety medicine; p		10		Nozinan
Inj 25 mg per ml, 1 ml ampoule	47.89	10		Wockhardt
(Wockhardt Inj 25 mg per ml, 1 ml ampoule to be delisted 1 April			•	WOCKHAIGE
	*			
LEVOMEPROMAZINE MALEATE – Safety medicine; prescriber				
Tab 25 mg		100		Nozinan
Tab 100 mg	41./5	100	•	<u>Nozinan</u>
LITHIUM CARBONATE - Safety medicine; prescriber may dete		luency		
Tab 250 mg	34.30	500		Lithicarb FC
Tab long-acting 400 mg		100		Priadel
Cap 250 mg	9.42	100	•	Douglas
OLANZAPINE - Safety medicine; prescriber may determine disp	pensing frequency			
Tab 2.5 mg		28	/	Zypine
Tab 5 mg	1.15	28	1	Zypine
Tab orodispersible 5 mg	1.25	28	1	Zypine ODT
Tab 10 mg	1.65	28		Zypine
Tab orodispersible 10 mg	2.05	28	1	Zypine ODT
PERICYAZINE - Safety medicine; prescriber may determine dis	spensing frequency			
Tab 2.5 mg		84	1	Neulactil
	12.49	100		Neulactil
Tab 10 mg	37.34	84	1	Neulactil
and a g	44.45	100	1	Neulactil
QUETIAPINE - Safety medicine; prescriber may determine disp	encina frequency			
Tab 25 mg		90	1	Quetapel
Tab 100 mg		90		Quetapel
Tab 200 mg		90		Quetapel
Tab 300 mg		90		Quetapel
· ·				<u> </u>
RISPERIDONE – Safety medicine; prescriber may determine dis Tab 0.5 mg		60	./	Actovio
Tab 1 mg		60 60		Actavis Actavis
Tab 2 mg		60	_	Actavis
Tab 3 mg		60	_	Actavis
Tab 4 mg		60		Actavis
Oral lig 1 mg per ml		30 ml		Risperon
		50 1111	•	- maperon
ZIPRASIDONE – Safety medicine; prescriber may determine dis		00		7
Cap 20 mg		60		Zusdone
Cap 40 mg		60		Zusdone
Cap 60 mg		60	_	Zusdone
Cap 80 mg		60		Zusdone
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pre	•			
Tab 10 mg	31.45	100	✓	Clopixol

Subsidy		Fully	Brand or	
(Manufacturer's Price)) Su	ıbsidised	Generic	
\$	Per	✓	Manufacturer	

Depot Injections

FLUPENTHIXOL DECANOATE – Safety medicine; presc Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSI Inj 20 mg per ml, 2 ml – Up to 5 inj available on a PSI Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSI	O13.14 O20.90	nsing freq 5 5 5	uency Fluanxol Fluanxol Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescri	ber may determine dispen	sing fregu	ency
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PS0		5	✓ Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PS	SO55.90	5	✓ Haldol Concentrate
			✓ Haldol
			Decanoas S29
OLANZAPINE - Special Authority see SA1428 below - R	etail pharmacy		
Safety medicine; prescriber may determine dispensing	g frequency		
Inj 210 mg vial	252.00	1	✓ Zyprexa Relprevv
Inj 300 mg vial	414.00	1	✓ Zyprexa Relprevv
Inj 405 mg vial	504.00	1	✓ Zyprexa Relprevv

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing	j frequency		
Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe	435.12	1	✓ Invega Sustenna

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the



Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
\$	Per	•	Manufacturer

continued...

initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE - Special Authority see SA1427 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequ	iency		
Inj 25 mg vial	135.98	1	 Risperdal Consta
Inj 37.5 mg vial	178.71	1	✓ Risperdal Consta
Inj 50 mg vial	217.56	1	✓ Risperdal Consta

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine: prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml − Up to 5 inj available on a PSO19.80 5 **Clopixol**

Anxiolytics

BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	20.23	100	✓ Orion
* Tab 10 mg	13.16	100	✓ Orion
CLONAZEPAM - Safety medicine; prescriber may determi	ne dispensing frequency		
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg		100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determine of	lispensing frequency		
Tab 2 mg	15.05	500	✓ Arrow-Diazepam
Tab 5 mg	16.18	500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine	e dispensing frequency		
Tab 1 mg	9.72	250	✓ Ativan
Tab 2.5 mg	12.50	100	✓ Ativan
OXAZEPAM - Safety medicine; prescriber may determine	dispensing frequency		
Tab 10 mg	6.17	100	✓ Ox-Pam
Tab 15 mg		100	✓ Ox-Pam

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Multiple Sclerosis Treatments

DIMETHYL FUMARATE - Special Authority see SA1559 below - Retail pharmacy

Wastage claimable

 Cap 120 mg
 520.00
 14
 ✓ Tecfidera

 Cap 240 mg
 2,000.00
 56
 ✓ Tecfidera

⇒SA1559 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- a) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- b) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- c) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- d) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria):
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse:
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point:
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- e) applications must be made by the patient's neurologist or general physician; and
- f) patients must have no previous history of lack of response to dimethyl fumarate; and
- g) patients must have not previously had intolerance to dimethyl fumarate; and



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

h) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to dimethyl fumarate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

FINGOLIMOD - Special Authority see SA1562 below - Retail pharmacy

Wastage claimable

Cap 0.5 mg.......2,200.00 28 **✓ Gilenya**

⇒SA1562 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- i) a gadolinium enhancing lesion; or
- ii) a Diffusion Weighted Imaging positive lesion; or
- iii) a T2 lesion with associated local swelling; or
- iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
- v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point:
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) patients must have no previous history of lack of response to fingolimod; and
- 7) patients must have not previously had intolerance to fingolimod; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0: or
 - c) 1.5 to 3.5: or
 - d) 2.0 to 4.0: or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to fingolimod; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

NATALIZUMAB - Special Authority see SA1563 below - Retail pharmacy

⇒SA1563 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) treatment must be initiated and supervised by a neurologist who is registered in the Tysabri Australasian Prescribing Programme operated by the supplier; and
- 7) patients must have no previous history of lack of response to natalizumab; and
- 8) patients must have not previously had intolerance to natalizumab; and
- 9) a) Patient is JC virus negative, or
 - Patient is JC virus positive and has given written informed consent acknowledging an understanding of the risk of progressive multifocal leucoencephalopathy (PML) associated with natalizumab
- 10) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	sidised	Generic	
\$	Per	1	Manufacturer	

continued...

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0: or
 - c) 1.5 to 3.5: or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to natalizumab; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier.

Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate.

Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

TERIFLUNOMIDE - Special Authority see SA1560 below - Retail pharmacy

Wastage claimable

⇒SA1560 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

continued...

- ii) a Diffusion Weighted Imaging positive lesion; or
- iii) a T2 lesion with associated local swelling; or
- iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
- v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) patients must have no previous history of lack of response to teriflunomide; and
- 7) patients must have not previously had intolerance to teriflunomide; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - a) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to teriflunomide: or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate and teriflunomide is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

Other Multiple Sclerosis Treatments

GLATIRAMER ACETATE − Special Authority see SA1808 on the next page − Retail pharmacy Inj 40 mg prefilled syringe − No patient co-payment payable......2,275.00 12 Copaxone

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

⇒SA1808 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. As a new Special Authority approval number is required the MSTAC coordinator should be notified of the change and a new prescription provided. **Entry Criteria**

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
•	Por 🗸	Manufacturor

continued...

- b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment.
 Progression of disability is defined as progress by any of the following EDDSS Points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0: or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

INTERFERON BETA-1-ALPHA - Special Authority see SA1809 below - Retail pharmacy

No patient co-payment payable

Inj 6 million iu prefilled syringe	1,170.00	4	✓ Avonex
Injection 6 million ju per 0.5 ml pen injector	1.170.00	4	✓ Avonex Pen

⇒SA1809 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Subsidy (Manufacturer's Price)	,		Brand or Generic	
\$	Per	Subsidised <	Manufacturer	

continued...

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. As a new Special Authority approval number is required the MSTAC coordinator should be notified of the change and a new prescription provided. **Entry Criteria**

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or
 - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment.
 Progression of disability is defined as progress by any of the following EDDSS Points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5: or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5: or
 - f) 3.0 to 4.5; or
 - a) 3.5 to 4.5; or
 - h) 4.0 to 4.5.



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

continued...

- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

INTERFERON BETA-1-BETA - Special Authority see SA1810 below - Retail pharmacy

No patient co-payment payable

⇒SA1810 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. As a new Special Authority approval number is required the MSTAC coordinator should be notified of the change and a new prescription provided. **Entry Criteria**

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or

NERVOUS SYSTEM

Subsidy		Fully	Brand or	
(Manufacturer's Price)		ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
- v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or
 - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment.
 Progression of disability is defined as progress by any of the following EDDSS Points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0: or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0: or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.



Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	/	Manufacturer

Sedatives and Hypnotics

MELATONIN - Special Authority see SA1666 below - Retail pharmacy

30 Circadin

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

IIDAZOLAM – Safety medicine; prescriber may determine o	dispensing frequency	
Inj 1 mg per ml, 5 ml ampoule	4.30 10	✓ Midazolam-Claris
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj avai	lable	
on a PSO	14.90 10	✓ Pfizer
On a PSO for status epilepticus use only. PSO mus	t be endorsed for status epile	epticus use only.
Inj 5 mg per ml, 3 ml ampoule	2.50 5	✓ Midazolam-Claris
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj avail	able on	
a PSO	11.90 5	✓ Pfizer
On a PSO for status epilepticus use only. PSO mus	t be endorsed for status epile	epticus use only.

NITRAZEPAM - Subsidy by endorsement

- a) Safety medicine; prescriber may determine dispensing frequency
- b) Subsidy by endorsement subsidised for patients who were taking nitrazepam prior to 1 August 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior

Tab 5 mg	5.22	100	✓ Nitrados
(Nitrados Tab 5 mg to be delisted 1 January 2021)			
PHENOBARBITONE SODIUM - Special Authority see SA1386 be	low – Retail ph	armacy	
Inj 200 mg per ml, 1 ml ampoule	30.00	5	✓ Aspen S29

⇒SA1386 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	✓	Manufacturer
TEMAZEPAM - Safety medicine; prescriber may determine disp	ensing frequency			
Tab 10 mg	1.27	25	✓ N	lormison_
TRIAZOLAM - Safety medicine; prescriber may determine dispe	ensing frequency			
Tab 125 mcg	5.10	100		
· ·	(9.85)		Н	lypam
Tab 250 mcg	4.10	100		
•	(11.20)		Н	lypam
ZOPICLONE - Safety medicine; prescriber may determine dispe	ensing frequency			
Tab 7.5 mg	9.56	500	√ <u>Z</u>	Copicione Actavis

Stimulants/ADHD Treatments

ATOMOXETINE - Special Authority see SA1416 below - Retail pharmacy		
Cap 10 mg107.03	28	✓ Strattera
Cap 18 mg107.03	28	✓ Strattera
Cap 25 mg107.03	28	✓ Strattera
Cap 40 mg107.03	28	✓ Strattera
Cap 60 mg107.03	28	✓ Strattera
Cap 80 mg139.11	28	✓ Strattera
Cap 100 mg139.11	28	✓ Strattera

⇒SA1416 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder) diagnosed according to DSM-IV or ICD 10 criteria; and
- 2 Once-daily dosing; and
- 3 Any of the following:
 - 3.1 Treatment with a subsidised formulation of a stimulant has resulted in the development or worsening of serious adverse reactions or where the combination of subsidised stimulant treatment with another agent would pose an unacceptable medical risk; or
 - 3.2 Treatment with a subsidised formulation of a stimulant has resulted in worsening of co-morbid substance abuse or there is a significant risk of diversion with subsidised stimulant therapy; or
 - 3.3 An effective dose of a subsidised formulation of a stimulant has been trialled and has been discontinued because of inadequate clinical response; or
 - 3.4 Treatment with a subsidised formulation of a stimulant is considered inappropriate because the patient has a history of psychoses or has a first-degree relative with schizophrenia; and
- 4 The patient will not be receiving treatment with atomoxetine in combination with a subsidised formulation of a stimulant, except for the purposes of transitioning from subsidised stimulant therapy to atomoxetine.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: A "subsidised formulation of a stimulant" refers to currently subsidised methylphenidate hydrochloride tablet formulations (immediate-release, sustained-release and extended-release) or dexamfetamine sulphate tablets.

DEXAMFETAMINE SULFATE - Special Authority see SA1149 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following



Su	ubsidy F	ully	Brand or
(Manufac	cturer's Price) Subsid	ised	Generic
	\$ Per	✓	Manufacturer

criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

- Both:
 - 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
 - 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1150 below - Retail pharmacy

- a) Only on a controlled drug form

b) Safety medicine; prescriber may determine dispensi	ng rrequency		
Tab immediate-release 5 mg	3.20	30	Rubifen
Tab immediate-release 10 mg	3.00	30	✓ Ritalin
•			Rubifen
Tab immediate-release 20 mg	7.85	30	Rubifen
Tab sustained-release 20 mg		30	Rubifen SR
·	50.00	100	Ritalin SR

⇒SA1150 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	sidised	Generic
\$	Per	•	Manufacturer

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Roth:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1151 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab extended-release 18 mg	18.20	30	Methylphenidate ERTeva
	58.96		✓ Concerta
Tab extended-release 27 mg	22.00	30	Methylphenidate ERTeva
	65.44		✓ Concerta
Tab extended-release 36 mg	22.40	30	Methylphenidate ERTeva
	71.93		✓ Concerta
Tab extended-release 54 mg	26.40	30	Methylphenidate ERTeva
	86.24		✓ Concerta
Cap modified-release 10 mg	15.60	30	✓ Ritalin LA
Cap modified-release 20 mg	20.40	30	✓ Ritalin LA
Cap modified-release 30 mg		30	✓ Ritalin LA
Cap modified-release 40 mg	30.60	30	✓ Ritalin LA

⇒SA1151 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or



Subsidy	Į	Fully	Brand or
(Manufacturer's Price)	Subsid	lised	Generic
\$	Per	✓	Manufacturer

- 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Fither:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

⇒SA1126 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	4.34	90	✓ Donepezil-Rex
* Tab 10 mg	6.64	90	✓ Donepezil-Rex
RIVASTIGMINE - Special Authority see SA1488 below - Retail	pharmacy		
Patch 4.6 mg per 24 hour	90.00	30	✓ Exelon
Patch 9.5 mg per 24 hour	90.00	30	✓ Exelon

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 The patient has been diagnosed with dementia; and

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic	
\$	Per	1	Manufacturer	

2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 2 mg with naloxone 0.5 mg	18.37	28
	57.40	
Tab sublingual 8 mg with naloxone 2 mg	53.12	28
	100.00	

✓ Buprenorphine Naloxone BNM

✓ Suboxone

✓ Buprenorphine Naloxone BNM

✓ Suboxone

(Suboxone Tab sublingual 2 mg with naloxone 0.5 mg to be delisted 1 April 2020) (Suboxone Tab sublingual 8 mg with naloxone 2 mg to be delisted 1 April 2020)

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and



Subsidy (Manufacturer's Price)	S	Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

Tab modified-release 150 mg	11.00	30	✓ Zyban
DISULFIRAM Tab 200 mg	153.00	100	✓ Antabuse
NALTREXONE HYDROCHLORIDE - Special Authority	see SA1408 below - Reta	il pharmacy	
Tab 50 mg			✓ Naltraccore

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic
` \$	Per	•	Manufacturer

NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.
- b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A. ✓ Habitrol Patch 7 mg - Up to 28 patch available on a PSO17.28 28

Patch 7 mg for direct distribution only - [Xpharm]	3.94	7	✓ <u>Habitrol</u>
Patch 14 mg - Up to 28 patch available on a PSO	19.00	28	✓ Habitrol
Patch 14 mg for direct distribution only - [Xpharm]	4.52	7	✓ <u>Habitrol</u>
Patch 21 mg - Up to 28 patch available on a PSO	21.77	28	✓ <u>Habitrol</u>
Patch 21 mg for direct distribution only - [Xpharm]	5.18	7	✓ <u>Habitrol</u>
Lozenge 1 mg - Up to 216 loz available on a PSO	18.27	216	✓ <u>Habitrol</u>
Lozenge 1 mg for direct distribution only - [Xpharm].	3.20	36	✓ <u>Habitrol</u>
Lozenge 2 mg - Up to 216 loz available on a PSO	20.02	216	✓ <u>Habitrol</u>
Lozenge 2 mg for direct distribution only - [Xpharm].	3.24	36	✓ <u>Habitrol</u>
Gum 2 mg (Fruit) - Up to 384 piece available on a PS	SO36.39	384	✓ <u>Habitrol</u>
Gum 2 mg (Fruit) for direct distribution only - [Xpharr	n]8.64	96	✓ <u>Habitrol</u>
Gum 2 mg (Mint) - Up to 384 piece available on a PS	O36.39	384	✓ <u>Habitrol</u>
Gum 2 mg (Mint) for direct distribution only - [Xpharn	1]8.64	96	✓ <u>Habitrol</u>
Gum 4 mg (Fruit) - Up to 384 piece available on a PS	SO42.07	384	✓ <u>Habitrol</u>
Gum 4 mg (Fruit) for direct distribution only - [Xpharn	n]10.01	96	✓ <u>Habitrol</u>
Gum 4 mg (Mint) - Up to 384 piece available on a PS	O42.07	384	✓ <u>Habitrol</u>
Gum 4 mg (Mint) for direct distribution only - [Xpharn	10.01	96	✓ <u>Habitrol</u>

VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 4225.64	53 OP	✓ Varenicline Pfizer
Tab 1 mg27.10	56	✓ Varenicline Pfizer

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking;
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme. which includes prescriber or nurse monitoring; and
- 3 Fither:
 - 3.1 The patient has tried but failed to guit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement
 - 3.2 The patient has tried but failed to guit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking:

NERVOUS SYSTEM

(Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

and

- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see SA1667 below

	271.35	1	_
, ,	1,085.38	1	✓ Ribomustin
Inj 1 mg for ECP	11.40	1 mg	✓ Baxter

⇒SA1667 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 All of the following:
 - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
 - 3.2.2 The patient has not received prior bendamustine therapy; and
 - 3.2.3 Fither:
 - 3.2.3.1 Both:
 - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more: or
 - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Fither:
 - 2.1 Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	1	Manufacturer

- 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.
 Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

BUSULFAN - PCT - Retail pharmacy-Specialist			
Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			
Inj 10 mg per ml, 45 ml vial	32.59	1	DBL Carboplatin
	45.20		Carboplatin Ebewe
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist			
Inj 100 mg vial	1,387.00	1	✓ BiCNU
			✓ Bicnu Heritage S29
Inj 100 mg for ECP	1,387.00	100 mg OP	✓ Baxter
CHLORAMBUCIL - PCT - Retail pharmacy-Specialist			
Tab 2 mg	29.06	25	Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	12.29	1	✓ DBL Cisplatin
	15.00		✓ Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial	19.70	1	 DBL Cisplatin
	21.00		Cisplatin Ebewe
Inj 1 mg for ECP	0.25	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE			
Tab 50 mg - PCT - Retail pharmacy-Specialist	79.00	50	✓ Endoxan S29
	158.00	100	✓ Procytox S29
Wastage claimable			
Inj 1 g vial - PCT - Retail pharmacy-Specialist	35.65	1	✓ Endoxan
	127.80	6	Cytoxan
Inj 2 g vial – PCT only – Specialist		1	✓ Endoxan
Inj 1 mg for ECP - PCT only - Specialist	0.04	1 mg	✓ Baxter
IFOSFAMIDE - PCT only - Specialist			
lnj 1 g	96.00	1	✓ Holoxan
lnj 2 g	180.00	1	✓ Holoxan
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
LOMUSTINE - PCT - Retail pharmacy-Specialist			
Cap 10 mg	132.59	20	✓ CeeNU
Cap 40 mg	399.15	20	✓ CeeNU
MELPHALAN			
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	✓ Alkeran
Inj 50 mg - PCT only - Specialist	67.80	1	✓ Alkeran

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
OXALIPLATIN - PCT only - Specialist	*			
Inj 100 mg vial	25.01	1	•	Oxaliplatin Actavis
	110.00		/	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	46.32	1	✓	Oxaliccord Oxaliplatin Accord
Inj 1 mg for ECP(Oxaliccord Inj 5 mg per ml, 20 ml vial to be delisted 1 February 2		1 mg	_	Baxter
THIOTEPA – PCT only – Specialist				
Inj 15 mg vial	CBS	1	1	Bedford \$29
.,		•	1	THIO-TEPA \$29
			✓	Tepadina S29
Inj 100 mg vial	CBS	1	✓	Tepadina \$29
Antimetabolites				
AZACITIDINE - PCT only - Specialist - Special Authority see S.	A1467 below			
Inj 100 mg vial	139.00	1	✓	Azacitidine Dr Reddy's
	605.00		1	Vidaza
Inj 1 mg for ECP	1.53	1 mg	✓	Baxter

⇒SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome: or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder);
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- 4 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

	0.1.1			B 1
	Subsidy (Manufacturer's Price	e) Subsi	Fully dised	Brand or Generic
	\$	Per	✓	Manufacturer
CALCIUM FOLINATE				
Tab 15 mg - PCT - Retail pharmacy-Specialist	104.26	10		BL Leucovorin Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17.10	5	✓ Ho	ospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Speciali	ist7.28	1		alcium Folinate Sandoz
Inj 50 mg - PCT - Retail pharmacy-Specialist	18.25	5		alcium Folinate Ebewe
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	9.49	1		alcium Folinate Sandoz
Inj 100 mg - PCT only - Specialist	7.33	1		alcium Folinate Ebewe
Inj 300 mg - PCT only - Specialist	22.51	1		alcium Folinate Ebewe
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	25.14	1	✓ Ca	alcium Folinate Sandoz
Inj 1 g - PCT only - Specialist	67.51	1	✓ Ca	alcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	72.00	1		alcium Folinate Sandoz
Inj 1 mg for ECP – PCT only – Specialist(Calcium Folinate Ebewe Inj 50 mg to be delisted 1 March 2020)	0.06	1 mg	✓ Ba	
CAPECITABINE - Retail pharmacy-Specialist				
Tab 150 mg		60	✓ Br	
Tab 500 mg	62.28	120	✓ Br	inov
CLADRIBINE – PCT only – Specialist Inj 1 mg per ml, 10 ml	E 040 70	7	./ La	eustatin
Inj 10 mg for ECP		10 mg OP	✓ Ba	
CYTARABINE		To mg Or		
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Speciali	ist400.00	5	✓ Pf	izer
Inj 100 mg per ml, 20 ml vial – PCT – Retail		•		
pharmacy-Specialist	41.36	1	✓ Pf	izer
Inj 1 mg for ECP - PCT only - Specialist		10 mg	✓ Ba	axter
Inj 100 mg intrathecal syringe for ECP - PCT only - Special	ist80.00	100 mg OP	✓ Ba	axter
FLUDARABINE PHOSPHATE				
Tab 10 mg - PCT - Retail pharmacy-Specialist		20	_	udara Oral
Inj 50 mg vial – PCT only – Specialist		5		udarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist	115.29	50 mg OP	✓ Ba	axter
FLUOROURACIL			_	
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist		1		uorouracil Ebewe
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist		1		uorouracil Ebewe
Inj 1 mg for ECP – PCT only – Specialist	0.66	100 mg	✓ Ba	axter
GEMCITABINE HYDROCHLORIDE – PCT only – Specialist	00			
Inj 1 g, 26.3 ml vial		1		BL Gemcitabine
Inj 1 g		1		emcitabine Ebewe
Inj 1 mg for ECP	349.20 0.02	1 mg	✓ Ge	emzar exter
,		·y		

	Subsidy (Manufacturer's Pric	e) Per	Fully Subsidised	d Generic
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist				
Inj 20 mg per ml, 5 ml vial	71.44	1	•	Irinotecan Accord \$29
			✓	Irinotecan Actavis
	100.00		✓	Irinotecan-Rex
Inj 1 mg for ECP	0.75	1 mg	✓	Baxter
MERCAPTOPURINE				
Tab 50 mg - PCT - Retail pharmacy-Specialist	37.00	25	1	Puri-nethol
Oral suspension 20 mg per ml - Retail pharmacy-Specialist	_			
Special Authority see SA1725 below		100 ml (OP 🗸	Allmercap

⇒SA1725 Special Authority for Subsidy

METHOTREXATE

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

IVI	INOTREMATE		
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist8.05	90	✓ Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist31.75	90	✓ Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist47.50	5	✓ Hospira
*	Inj 7.5 mg prefilled syringe14.61	1	✓ Methotrexate
			Sandoz
*	Inj 10 mg prefilled syringe14.66	1	✓ Methotrexate
			Sandoz
*	Inj 15 mg prefilled syringe14.77	1	✓ Methotrexate
			Sandoz
*	Inj 20 mg prefilled syringe14.88	1	✓ Methotrexate
			Sandoz
*	Inj 25 mg prefilled syringe14.99	1	✓ Methotrexate
			Sandoz
*	Inj 30 mg prefilled syringe15.09	1	✓ Methotrexate
			Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist30.00	5	✓ DBL Methotrexate
			Onco-Vial
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist45.00	1	✓ DBL Methotrexate
			Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist25.00	1	✓ Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail		
	pharmacy-Specialist79.99	1	✓ Methotrexate Ebewe
*	Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist4.73	5 mg OP	✓ Baxter
PE	METREXED - PCT only - Specialist - Special Authority see SA1679 on the r	next page	
	Inj 100 mg vial60.89	1	✓ Juno Pemetrexed
	Inj 500 mg vial217.77	1	✓ Juno Pemetrexed
	11		

✓ Baxter

Subs	sidy Fu	lly Brand or
(Manufactur	rer's Price) Subsidis	ed Generic
\$	Per	✓ Manufacturer

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:

Tab 40 ma

- 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Permetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
- 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

100.01

OF

/ 1 amida

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

1 No evidence of disease progression; and

THIOGUANINE - PCT - Retail pharmacy-Specialist

- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

1ab 40 mg126.31	25	Lanvis
Other Cytotoxic Agents		
AMSACRINE - PCT only - Specialist		
Inj 50 mg per ml, 1.5 ml ampoule	6	✓ Amsidine S29
Inj 75 mg1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-Specialist		
Cap 0.5 mgCBS	100	✓ Agrylin S29
		✓ Teva S29
ARSENIC TRIOXIDE - PCT only - Specialist		
Inj 1 mg per ml, 10 ml vial	10	✓ Phenasen
Inj 10 mg for ECP481.70	10 mg OP	✓ Baxter

	Subsidy (Manufacturer's Price	e) Sub	Fully	Brand or Generic	
	\$	Per	1	Manufacturer	
BLEOMYCIN SULPHATE - PCT only - Specialist					
Inj 15,000 iu, vial	161.01	1	✓ D	DBL Bleomycin Sulfate	
Inj 1,000 iu for ECP	12.45	1,000 iu	✓ B	Baxter	
BORTEZOMIB - PCT only - Specialist - Special Authority see S	A1576 below				
Inj 3.5 mg vial	1,892.50	1	✓ V	/elcade	
Inj 1 mg for ECP	594.77	1 mg	✓ B	Baxter	

⇒SA1576 Special Authority for Subsidy

Initial application — (Treatment naive multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient has treatment-naive symptomatic multiple myeloma; or
 - 1.2 The patient has treatment-naive symptomatic systemic AL amyloidosis *; and
- 2 Maximum of 9 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Relapsed/refractory multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has relapsed or refractory multiple myeloma; or
 - 1.2 The patient has relapsed or refractory systemic AL amyloidosis *; and
- 2 The patient has received only one prior front line chemotherapy for multiple myeloma or amyloidosis; and
- 3 The patient has not had prior publicly funded treatment with bortezomib; and
- 4 Maximum of 4 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Relapsed/refractory multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 The patient's disease obtained at least a partial response from treatment with bortezomib at the completion of cycle 4; and
- 2 Maximum of 4 further treatment cycles (making a total maximum of 8 consecutive treatment cycles).

Notes: Responding relapsed/refractory multiple myeloma patients should receive no more than 2 additional cycles of treatment beyond the cycle at which a confirmed complete response was first achieved. A line of therapy is considered to comprise either:

- a) a known therapeutic chemotherapy regimen and supportive treatments; or
- b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments.

Refer to datasheet for recommended dosage and number of doses of bortezomib per treatment cycle.

COLASPASE [L-ASPARAGINASE] - PCT only - Specialist ✓ Leunase 10.000 iu OP ✓ Baxter DACARBAZINE - PCT only - Specialist Inj 200 mg vial58.06 ✓ DBL Dacarbazine 580.60 10 ✓ Dacarbazine **ΔPP** \$29 Inj 200 mg for ECP58.06 200 ma OP ✓ Baxter DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist ✓ Cosmegen 1 0.5 mg OP ✓ Baxter

	Subsidy (Manufacturer's Price)			Full	,
	(Manufacturer's \$	Price)	Per	Subsidise	d Generic Manufacturer
AUNORUBICIN - PCT only - Specialist					
Inj 2 mg per ml, 10 ml	130.00		1	-	Pfizer
Inj 20 mg for ECP		20	mg (Baxter
, ,			9 \		
OCETAXEL - PCT only - Specialist Inj 10 mg per ml, 2 ml vial	10.40		1		DBL Docetaxel
Inj 20 mgInj 20 mg			1		Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial			1		Docetaxel Sandoz
Inj 20 mg per ml, 4 ml vial			1		Docetaxel
iiij 20 iiig poi iiii, 4 iiii viai	20.00		•	•	Accord S29
Inj 80 mg	105.00		1		Docetaxel Sandoz
Inj 1 mg for ECP			ı 1 mg		Baxter
	0.55		ı ıııy	•	Daxiei
OXORUBICIN HYDROCHLORIDE – PCT only – Specialist	40.00				. D
Inj 2 mg per ml, 5 ml vial			1		Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial			1		Doxorubicin Ebewe
lai O are are al. 50 artisist	17.00				Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial			1		Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial	65.00		1		´ Doxorubicin Ebewe ´ Arrow-Doxorubicin
Inj 1 mg for ECP			1 mg		Baxter
	0.29		ı ıııy	•	Daxiei
PIRUBICIN HYDROCHLORIDE - PCT only - Specialist	05.00				
Inj 2 mg per ml, 5 ml vial			1		Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial			1		Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial			1		Epirubicin Ebewe
Inj 1 mg for ECP	0.37		1 mg	•	Baxter
TOPOSIDE					
Cap 50 mg - PCT - Retail pharmacy-Specialist			20		Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist			10		<u>Vepesid</u>
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialis			1		Rex Medical
Inj 1 mg for ECP - PCT only - Specialist	0.09		1 mg	•	Baxter
ΓOPOSIDE PHOSPHATE − PCT only − Specialist					
Inj 100 mg (of etoposide base)	40.00		1		Étopophos
Inj 1 mg (of etoposide base) for ECP	0.47		1 mg	•	' Baxter
YDROXYUREA - PCT - Retail pharmacy-Specialist					
Cap 500 mg	31.76		100	•	Y Hydrea
ARUBICIN HYDROCHLORIDE					
Inj 5 mg vial - PCT only - Specialist	93.00		1	/	Zavedos
Inj 10 mg vial - PCT only - Specialist			1		Zavedos
Inj 1 mg for ECP - PCT only - Specialist			1 mg		Baxter
ENALIDOMIDE - Retail pharmacy-Specialist - Special Authority			9		
Wastage claimable	300 0/11700	SOIOW			
Cap 10 mg	6.207.00		21	•	Revlimid
Cap 15 mg			21		Revlimid
Cap 25 mg			21		Revlimid
SA1468 Special Authority for Subsidy	,0=1.00				

⇒SA1468 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	I Generic	
\$	Per 🗸	Manufacturer	

continued...

MESNA

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Either
 - 2.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 2.2 Both:
 - 2.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 2.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 3 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MEGNA		
Tab 400 mg - PCT - Retail pharmacy-Specialist314.00	50	✓ <u>Uromitexan</u>
Tab 600 mg - PCT - Retail pharmacy-Specialist448.50	50	✓ <u>Uromitexan</u>
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist177.45	15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist	15	✓ Uromitexan
Inj 1 mg for ECP - PCT only - Specialist2.96	100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist		
Inj 5 mg vial204.08	1	✓ Arrow
Inj 1 mg for ECP42.04	1 mg	✓ Baxter
MITOZANTRONE - PCT only - Specialist	•	
Inj 2 mg per ml, 10 ml vial	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP5.51	1 mg	✓ Baxter
PACLITAXEL - PCT only - Specialist	9	
, ·	_	4 B 19 1 E
Inj 30 mg47.30	5	Paclitaxel Ebewe
Inj 100 mg20.00	1	Paclitaxel Ebewe
91.67		Paclitaxel Actavis
Inj 150 mg26.69	1	Paclitaxel Ebewe
137.50		✓ Anzatax
		✓ Paclitaxel Actavis
Inj 300 mg35.35	1	✓ Paclitaxel Ebewe
275.00		✓ Anzatax
		✓ Paclitaxel Actavis
Inj 1 mg for ECP0.19	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see SA1325 below		
Inj 3,750 IU per 5 ml3,005.00	1	✓ Oncaspar S29
7 - 7 P		- · · · · · · · · · · · · · · · · · · ·

⇒SA1325 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and
- 3 Treatment is with curative intent.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and
- 3 Treatment is with curative intent.

PENTOSTATIN [DEOXYCOFORMYCIN] – PCT only – Specialist Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail pharmacy-S	pecialist		
Cap 50 mg	980.00	50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA1741 below - Retail	pharmacy		
Cap 5 mg	10.20	5	✓ Orion
			Temozolomide
Cap 20 mg	16.38	5	✓ Temaccord
	18.30		✓ Apo-Temozolomide
			Orion
			Temozolomide
			✓ Temizole 20 S29
Cap 100 mg	35.98	5	✓ Temaccord
	40.20		Apo-Temozolomide
			Orion
			Temozolomide
Cap 140 mg	50.12	5	Temaccord
	56.00		Orion
			Temozolomide
Cap 250 mg	86.34	5	✓ Temaccord
	96.80		✓ Orion
			Temozolomide

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	✓	Manufacturer

continued...

- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m2 per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special	Authority see SA1124 below		
Cap 50 mg	378.00	28	✓ Thalomid
Cap 100 mg	756.00	28	✓ Thalomid

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 12 months for applications meeting the following criteria: Fither:

- - 1 The patient has multiple myeloma; or
 - 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

٧

TILLINOIN		
Cap 10 mg - PCT - Retail pharmacy-Specialist479.50	100	Vesanoid
/INBLASTINE SULPHATE		
Inj 1 mg per ml, 10 ml vial - PCT - Retail pharmacy-Specialist 186.46	5	Hospira
Ini 1 mg for FCP - PCT only - Specialist 4 14	1 ma	✓ Baxter

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)		Fully Subsidised	Generic
	\$	Per		Manufacturer
VINCRISTINE SULPHATE				
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist	74.52	5	•	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist	85.61	5	✓	DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist	11.30	1 mg	✓	Baxter
VINORELBINE - PCT only - Specialist				
Inj 10 mg per ml, 1 ml vial	12.00	1	✓	Navelbine
, ,	42.00		1	Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial	56.00	1	1	Navelbine
	210.00		1	Vinorelbine Ebewe
Inj 1 mg for ECP	1.25	1 mg	✓	Baxter

Protein-tyrosine Kinase Inhibitors

DASATINIB - Special Authority see SA1805 below - Retail pharmacy

Wastage claimable

Tab 20 mg	3,774.06	60	✓ Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
Tab 70 mg	7,692.58	60	✓ Sprycel

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
 - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or

- 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
- 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB - Retail pharmacy-Specialist - Special Authorit	y see SA1653 on the n	ext page	
Tab 100 mg	764.00	30	✓ Tarceva
Tab 150 mg	1,146.00	30	✓ Tarceva

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1653 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Either:
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB - Retail pharmacy-Specialist - Special Authority see SA1654 below

⇒SA1654 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Fither:
 - 2.1 Patient is treatment naive: or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

IMATINIB MESILATE

Note: Imatinib-AFT is not a registered for the treatment of Gastro Intestinal Stromal Tumours (GIST). The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule.

Tab 100 mg - [Xpharm] - Special Authority see SA1460

	below	2,400.00	60	✓ Glivec
*	Cap 100 mg		60	✓ Imatinib-AFT
*	Cap 400 mg	197.50	30	✓ Imatinib-AFT

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 PHARMAC Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

	,	ully Brand or	
(Manufact	turer's Price) Subsidi	sed Generic	
	\$ Per	 Manufac 	turer

continued...

Special Authority criteria for GIST – access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE – Special Authority see SA1191 below – Retail pharmacy
Tab 250 mg

⇒SA1191 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient has not previously received trastuzumab treatment for HER 2 positive metastatic breast cancer; and
 - 1.3 Lapatinib not to be given in combination with trastuzumab; and
 - 1.4 Lapatinib to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2.2 The patient started trastuzumab for metastatic breast cancer but discontinued trastuzumab within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on trastuzumab; and
 - 2.4 Lapatinib not to be given in combination with trastuzumab; and
 - 2.5 Lapatinib to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology);
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

Wastage claimable

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Either
 - 2.1 Patient has documented CML treatment failure* with imatinib; or

Subsidy		Fully	Brand or
(Manufacturer's	Price) Subs	sidised	Generic
<u> </u>	Per	1	Manufacturer

continued...

- 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PAZOPANIB - Special Authority se	e SA1190 below – Retail pharmacy
----------------------------------	----------------------------------

Tab 200 mg	1,334.70	30	✓ Votrient
Tab 400 mg	2,669.40	30	✓ Votrient

⇒SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB - Special Authority see SA1753 on the next page - Retail pharmacy

wastage cialmable			
Tab 5 mg	2,500.00	56	Jakavi
Tab 15 mg	5.000.00	56	✓ Jakavi
Tab 20 mg	•	56	✓ Jakavi
1 ab 20 mg		50	• Oakavi

Subsidy (Manufacturer's Price)	Subsid	Fully dised	Brand or Generic
\$	Per	1	Manufacturer

⇒SA1753 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA1266 below - Retail pharmacy

Cap 12.5 mg	2,315.38	28	Sutent
Cap 25 mg	·	28	✓ Sutent
Cap 50 mg	·	28	✓ Sutent

⇒SA1266 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval: or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1 The patient's disease has progressed following treatment with imatinib; or
- 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 83

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA1767 below

Wastage claimable

⇒SA1767 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
- 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2: and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline: and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

BICALUTAMIDE Tab 50 mg	3.80	28	✓ Binarex
FLUTAMIDE - Retail pharmacy-Specialist			
Tab 250 mg	100.38	84	✓ Flutamide
	119.50	100	Mylan S29 ✓ Flutamin
MECECTROL ACETATE - Detail phormacy Capacialis		100	• Hutaniin
MEGESTROL ACETATE – Retail pharmacy-Specialist Tab 160 mg		30	✓ Apo-Megestrol
OCTREOTIDE	03.33	30	Apo-wegestron
Inj 50 mcg per ml, 1 ml vial	30 64	5	✓ DBL Octreotide
Inj 100 mcg per ml, 1 ml vial	18.69	5	✓ DBL Octreotide
Inj 500 mcg per ml, 1 ml vial		5	✓ DBL Octreotide
OCTREOTIDE LAR (SOMATOSTATIN ANALOGUE) -	- Special Authority see SA101	6 below -	Retail pharmacy
Inj LAR 10 mg prefilled syringe	1,772.50	1	Sandostatin LAR
Inj LAR 20 mg prefilled syringe		1	Sandostatin LAR
Inj LAR 30 mg prefilled syringe	2,951.25	1	Sandostatin LAR

⇒SA1016 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly: and
- 2 Any of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
 \$	Per	✓	Manufacturer

continued...

- 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
- 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed: or
- 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Fither:
 - 2.2.1 Patient has failed surgery: or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

* Tab 20 mg 5.60

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — **(Other Indications)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

TAMOXIFEN CITRATE

	•	
Aromatase Inhibitors		
ANASTROZOLE		
* Tab 1 mg5.04	30	✓ Rolin
EXEMESTANE		
* Tab 25 mg14.50	30	✓ Pfizer Exemestane
LETROZOLE		
* Tab 2.5 mg4.68	30	✓ <u>Letrole</u>

✓ Tamoxifen Sandoz

✓ Tamoxifen Sandoz

60

60

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
` • ′	Dor /	Manufacturar

Immunosuppressants

Cytotoxic Immunosuppressants

AZATHIOPRINE – Retail pharmacy-Specialist		
* Tab 25 mg	60	Azamun
9.66	100	Imuran
Azamun to be Sole Supply on 1 January 2020		
* Tab 50 mg	100	Azamun
10.58		Imuran
Azamun to be Sole Supply on 1 January 2020		
* Inj 50 mg vial	1	Imuran
(Imuran Tab 25 mg to be delisted 1 January 2020)		
(Imuran Tab 50 mg to be delisted 1 January 2020)		
, ,		
MYCOPHENOLATE MOFETIL		
Tab 500 mg25.00	50	Cellcept
Cap 250 mg25.00	100	 Cellcept
Powder for oral liq 1 g per 5 ml - Subsidy by endorsement	165 ml OP	✓ Cellcept
		•

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

Fusion Proteins

ETANERCEPT	 Special I 	Authority see	SA1812 below	 Retail pharmacy

Inj 25 mg	4	Enbrel
Inj 50 mg autoinjector	4	✓ Enbrel
Inj 50 mg prefilled syringe	4	✓ Enbrel

⇒SA1812 Special Authority for Subsidy

Initial application — (juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for juvenile idiopathic arthritis (JIA); and
 - 12 Fither
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for JIA: or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient diagnosed with Juvenile Idiopathic Arthritis (JIA); and
 - 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
 - 2.5 Both:
 - 2.5.1 Either:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

ioints: or

- 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
- 2.5.2 Physician's global assessment indicating severe disease.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis: or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
 - 2.6 Fither:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.7 Fither:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or

- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either: 1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
- 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or

2 All of the following:

- 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
- 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
- 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
- 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
- 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
- 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*: and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 4 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
- 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules;
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a named specialist or rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician: or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a dermatologist; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price) :	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment: and
- 3 A maximum of 4 doses.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

OLODULINI (COLUNE) DOT anh. Chasialist

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Spe	cialist					
Inj 50 mg per ml, 5 ml	2,351.25	5	✓ ATGAM			
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT on	ly – Specialist					
Subsidised only for bladder cancer.						
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE			
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG S29			
(SII-Onco-BCG \$29 Inj 40 mg per ml, vial to be delisted 1 Ap	oril 2021)					

Monoclonal Antibodies

		7 below – Retail pharmacy	ADALIMUMAB – Special Authority see SA1847 I
Humira	2	1,599.96	Inj 20 mg per 0.4 ml prefilled syringe
✓ HumiraPen	2	1,599.96	Inj 40 mg per 0.8 ml prefilled pen
Humira	2	1,599.96	Inj 40 mg per 0.8 ml prefilled syringe

⇒SA1847 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Fither:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 CDAI score is 150 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 PCDAI score is 15 or less: or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed;
 - 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Fither:

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
- 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
- 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD: or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Fither
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroillitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active</p>

Subsidy (Manufacturer's Price)	F Subsidi	ully	Brand or Generic
 \$	Per	√	Manufacturer

continued...

vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or

- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's disease.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for juvenile idiopathic arthritis; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient diagnosed with JIA; and
 - 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
 - 2.5 Both:
 - 2.5.1 Either:

Subsidy	ı	Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	•	Manufacturer

continued...

- 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender ioints: or
- 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
- 2.5.2 Physician's global assessment indicating severe disease.

Renewal — (juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a named specialist or rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Fither:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline: or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Fither:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — **(psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 4 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement: and
- 2 Patient continues to require treatment; and
- 3 A maximum of 4 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept: or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis: or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

- 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Fither:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease that is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has not responded adequately to treatment with infliximab (see Notes); or
 - 2.2 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has experienced intolerable side effects from treatment with infliximab; and
- 3 The patient is experiencing significant loss of quality of life; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: Behcet's disease diagnosed according to the International Study Group for Behcet's disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al, J Rheumatol. 2004;31:931-7.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plague psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline valuee; or

2.2 Both:

- 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
- 2.2.2 Fither:
 - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values: or
 - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value: and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for severe ocular inflammation: or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Initial application — (hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)	Suk	Fully	Brand or Generic
 \$	Per	√	Manufacturer

continued...

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 The patient has 3 or more active lesions (e.g. inflammatory nodules, abscesses, draining fistulae); and
- 4 The patient has a Dermatology Quality of Life Index of 10 or more and the assessment is no more than 1 month old at time of application; and
- 5 Following the initial loading doses, adalimumab is to be administered at doses no greater than 40mg every 7 days.

Renewal — (hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

AFLIBERCEPT – Special Authority see SA1772 below – Retail pharmacy

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab: or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
 - 2 Either:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	1	Manufacturer	

continued...

applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

CETUXIMAB – PCT only – Specialist – Special Authority see SA1697 below

Inj 5 mg per ml, 20 ml vial	364.00	1	Erbitux
Inj 5 mg per ml, 100 ml vial	1,820.00	1	Erbitux
Inj 1 mg for ECP	3.82	1 mg	Baxter

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB - PCT only - Special Authority see SA1831 below

Inj 100 mg	806.00	1	✓ Remicade
Inj 1 mg for ECP	8.29	1 mg	✓ Baxter

⇒SA1831 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less: or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (**Graft vs host disease**) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria: Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Fither:

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer Manufacturer	

continued...

- 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
- 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or

Subsidy (Manufacturer's Price)	Fu Subsidis	ılly	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Fither:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Fither:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 15, where lesions have been present for at least 6 months from the time of initial diagnosis; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 15, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Fither:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis: or
 - 2.4 Severe ocular inflammation: or
 - 2.5 Chronic ocular inflammation: or
 - 2.6 Crohn's disease (adults): or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease: or
 - 2.9 Severe fulminant ulcerative colitis; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	;	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.10 Severe ulcerative colitis; or
- 2.11 Plaque psoriasis; or
- 2.12 Neurosarcoidosis: or
- 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for psoriatic arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept: and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Fither:

continued...

- 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
- 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

1 The patient has had a good clinical response following 3 initial doses; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (severe ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
 - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids: and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (severe ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

OBINUTUZUMAB - PCT only - Specialist - Special Autl	hority see SA1627 below		
Inj 25 mg per ml, 40 ml vial	5,910.00	1	✓ Gazyva
Inj 1 mg for ECP	6.21	1 mg	✓ Baxter

⇒SA1627 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and</p>
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and

Subsidy	Fı	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
 \$	Per	•	Manufacturer

continued...

6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

OMALIZUMAB - Special Authority see SA1744 below - Retail phart	macy		
Inj 150 mg prefilled syringe	450.00	1	✓ Xolair
Inj 150 mg vial	450.00	1	✓ Xolair

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months. unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Fither:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
- 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PERTUZUMAB - PCT only - Specialist - Special Authority see SA1606 below

Inj 30 mg per ml, 14 ml vial	3,927.00	1	Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	Baxter

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology): and
- 2 Fither:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab: and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB - PCT only - Specialist - Special Authority see SA1818 on the next page

Inj 100 mg per 10 ml vial	1,075.50	2	Mabthera
Inj 500 mg per 50 ml vial	2,688.30	1	Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter

Subsidy (Manufacturer's Price) \$

Fully Subsidised Per

Brand or Generic Manufacturer

⇒SA1818 Special Authority for Subsidy

Initial application — (ABO-incompatible renal transplant) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible renal transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated renal transplant rejection) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated renal transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 The patient is rituximab treatment naive; and
- 3 Either:
 - 3.1 The patient is chemotherapy treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
 - 3.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; and
- 4 The patient has good performance status; and
- 5 The patient does not have chromosome 17p deletion CLL; and
- 6 Rituximab to be administered in combination with fludarabine and cyclophosphamide or bendamustine for a maximum of 6 treatment cycles; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

7 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
- 2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
- 3 The patient does not have chromosome 17p deletion CLL; and
- 4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide or bendamustine for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

Subsidy	y Full	/ Brand or
(Manufacturer's	s Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded: and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

Subsidy	Fu	,	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — **(haemophilia with inhibitors)** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre: or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy).

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Subsidy (Manufacturer's Price)	Fu Subsidis	ılly	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with rituximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 haemophilia with inhibitors; or
 - 2.2 rheumatoid arthritis: or
 - 2.3 severe cold haemagglutinin disease (CHAD); or
 - 2.4 warm autoimmune haemolytic anaemia (warm AIHA); or
 - 2.5 immune thrombocytopenic purpura (ITP); or
 - 2.6 thrombotic thrombocytopenic purpura (TTP); or
 - 2.7 pure red cell aplasia (PRCA); or
 - 2.8 ANCA associated vasculitis; or
 - 2.9 treatment refractory systemic lupus erythematosus (SLE); or
 - 2.10 steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS).

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of

Subsidy	Fı	ully Brand or	
(Manufacturer's Price)	Subsidis	sed Generic	
\$	Per	✓ Manufacturer	

continued...

leflunomide alone or in combination with oral or parenteral methotrexate; and

- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Fither:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Fither:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis: and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1.000 mg infusions of rituximab given two weeks apart.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physiciann; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria: Both:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms.

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria: Either:

- 1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
- 2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Patient has warm autoimmune haemolytic anaemia*: and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB - Special Authority see SA1754 below - Retail pharmacy

⇒SA1754 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and

4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

Note: Siltuximad is to be administered at doses no greater	rtnan i i mg/kg ever	y 3 weeks.	
Inj 100 mg vial	770.57	1	Sylvant
Inj 400 mg vial	3,082.33	1	Sylvant

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Generic
TOCILIZUMAB - PCT only - Special Authority see SA1858 belo	ow .			
Inj 20 mg per ml, 4 ml vial	220.00	1	✓	Actemra
Inj 20 mg per ml, 10 ml vial	550.00	1	✓	Actemra
Inj 20 mg per ml, 20 ml vial		1	1	Actemra
Inj 1 mg for ECP		1 mg	1	Baxter

⇒SA1858 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease: or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Fither:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules: and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Fither:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints:
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate: non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 Fither:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD): or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Bules of the Pharmaceutical Schedule: and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992:19:424-430); and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
- 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
 - 2.4 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.5 Both:
 - 2.5.1 Either:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status.

Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB - PCT only - Specialist - Special Authority see SA1632 below

Inj 150 mg vial	1,350.00	1	✓ Herceptin
Inj 440 mg vial	3,875.00	1	✓ Herceptin
Inj 1 mg for ECP	9.36	1 mg	✓ Baxter

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Fither:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:
All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

Programmed Cell Death-1 (PD-1) Inhibitors

NIVOLOMAB - POT only - Specialist - Special Authority see SA1656 on the next page			
Inj 10 mg per ml, 4 ml vial	1,051.98	1	Opdivo
Inj 10 mg per ml, 10 ml vial	2,629.96	1	Opdivo
Inj 1 mg for ECP	27.62	1 mg	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1656 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma stage III or IV; and
- 2 Patient has measurable disease as defined by the presence of at least one CT or MRI measurable lesion; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Nivolumab is to be used at a maximum dose of 3 mg/kg every 2 weeks for a maximum of 12 weeks (6 cycles); and
- 6 Baseline measurement of overall tumour burden is documented (see Note); and
- 7 Documentation confirming that the patient has been informed and acknowledges that the initial funded treatment period of nivolumab will not be continued beyond 12 weeks (6 cycles) if their disease progresses during this time.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note; or
 - 1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.3 Patient has stable disease according to RECIST criteria (see Note); and
- 2 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; and
- 3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; and
- 5 Nivolumab will be used at a maximum dose of 3 mg/kg every 2 weeks for a maximum of 12 weeks (6 cycles).

Notes: Disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Target lesion measurements should be assessed using CT or MRI imaging with the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

	IBROLIZUMAB - PCT only - Specialist - Special Authority see SA1657 on the next page		
✓ Keytruda	1	4,680.00	Inj 25 mg per ml, 4 ml vial
✓ Baxter	1 mg	49.14	Inj 1 mg for ECP

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1657 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma stage III or IV; and
- 2 Patient has measurable disease as defined by the presence of at least one CT or MRI measurable lesion; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Pembrolizumab is to be used at a maximum dose of 2 mg/kg every 3 weeks for a maximum of 12 weeks (4 cycles); and
- 6 Baseline measurement of overall tumour burden is documented (see Note); and
- 7 Documentation confirming that the patient has been informed and acknowledges that the initial funded treatment period of pembrolizumab will not be continued beyond 12 weeks (4 cycles) if their disease progresses during this time.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.3 Patient has stable disease according to RECIST criteria (see Note); and
- 2 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; and
- 3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; and
- 5 Pembrolizumab will be used at a maximum dose of 2 mg/kg every 3 weeks for a maximum of 12 weeks (4 cycles).

Notes: Disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Target lesion measurements should be assessed using CT or MRI imaging with the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Tab 5 mg4,555.76

	(Manufacturer's Price) Si	ubsidised	Generic	
	\$	Per	ıbsiuiseu ✓	Manufacturer	
Other Immunosuppressants					
CICLOSPORIN					
Cap 25 mg	44.63	50	✓	Neoral	
Cap 50 mg	88.91	50	✓	Neoral	
Cap 100 mg	177.81	50	✓	Neoral	
Oral liq 100 mg per ml	198.13 5	60 ml OP	1	Neoral	
EVEROLIMUS – Special Authority see SA1491 below – Retail ph Wastage claimable	armacy				
Tab 10 mg	6,512.29	30	1	Afinitor	

Subsidy

Fully

✓ Afinitor

Brand or

⇒SA1491 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

SIROLIMUS - Special Authority see SA0866 below - Retail pharmacy

Tab 1 mg	749.99	100	Rapamune
Tab 2 mg	1,499.99	100	Rapamune
Oral lig 1 mg per ml	449.99	60 ml OP	✓ Rapamune

⇒SA0866 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min; or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis: or
- . HUS or TTP; or
- Leukoencepthalopathy: or
- · Significant malignant disease

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

torio = initial operation of the state of th			
Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg	99.30	100	✓ Tacrolimus Sandoz
Cap 1 mg		100	✓ Tacrolimus Sandoz
Cap 5 mg		50	✓ Tacrolimus Sandoz

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified

continued...

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

Antiallergy Preparations

Allergic Emergencies

ICATIBANT - Special Authority see SA1558 below - Retail pharmacy

Inj 10 mg per ml, 3 ml prefilled syringe.......2,668.00 1 ✓ Firazyr

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Maintenance kit - 6 vials 120 mcg freeze dried venom, with

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above - Retail pharmacy

diluent	285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent	t		
9 ml, 3 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with dilu	ent305.00	1 OP	✓ Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority se	ee SA1367 above -	- Retail pharr	nacy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze	е		
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze	:		
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freez	e		
dried venom, with diluent	305.00	1 OP	✓ Venomil S29

	Subsidy		Fully Brand or
	(Manufacturer's Price		
	\$	Per	✓ Manufacturer
			Than a data or
Antihistamines			
CETIRIZINE HYDROCHLORIDE			
* Tab 10 mg	1 12	100	✓ Zista
* Oral lig 1 mg per ml		200 ml	✓ Histaclear
	2.00	200 1111	Tilstacical
CHLORPHENIRAMINE MALEATE			
* Oral liq 2 mg per 5 ml	9.37	500 ml	✓ Histafen
DEXTROCHLORPHENIRAMINE MALEATE			
* Tab 2 mg	2.02	40	
· · · · · · · · · · · · · · · · · · ·	(8.40)		Polaramine
	1.01	20	
	(5.99)		Polaramine
* Oral liq 2 mg per 5 ml		100 ml	1 olaramine
Talling 2 mg por 3 mi	(10.29)	100 1111	Polaramine
	(10.23)		Tolaramine
FEXOFENADINE HYDROCHLORIDE			
* Tab 60 mg	4.34	20	
	(8.23)		Telfast
* Tab 120 mg	4.74	10	
	(8.23)		Telfast
	14.22	30	
	(26.44)		Telfast
LORATADINE	, ,		
* Tab 10 mg	1.60	100	✓ Lorafix
	1.09	100	Loralix
Lorafix to be Sole Supply on 1 February 2020	0.45	100 1	. Laufaat
* Oral liq 1 mg per ml	2.15	120 ml	✓ Lorfast
PROMETHAZINE HYDROCHLORIDE			
* Tab 10 mg	1.68	50	✓ <u>Allersoothe</u>
* Tab 25 mg	1.89	50	✓ <u>Allersoothe</u>
* Oral liq 1 mg per 1 ml	2.69	100 ml	✓ Allersoothe
* Inj 25 mg per ml, 2 ml ampoule - Up to 5 inj available on a F	PSO 15.54	5	✓ Hospira
			•
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler, 50 mcg per dose	9.30 2	00 dose OP	✓ Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP	✓ Beclazone 50
Aerosol inhaler, 100 mcg per dose		200 dose OP	✓ Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP	✓ Beclazone 100
		200 dose OP	✓ Beclazone 250
Aerosol inhaler, 250 mcg per dose CFC-free	22.01 2	ou dose OP	▼ Deciazone 200
BUDESONIDE			_
Powder for inhalation, 100 mcg per dose	17.00 2	00 dose OP	✓ Pulmicort
			Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00 2	00 dose OP	✓ Pulmicort
			Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00 2	00 dose OP	✓ Pulmicort
i owder for initialation, 400 mby per dose	02.00 2	.ou dose or	Turbuhaler
			i ui builalei

	Subsidy		Fully	
	(Manufacturer's \$	Price) S	ubsidised	I Generic Manufacturer
LUTICACONE	Ψ	1 61		Manuacturei
LUTICASONE Aerosol inhaler, 50 mcg per dose	4.68	120 dose (ם בר	Floair
Aerosol inhaler, 50 mcg per dose CFC-free		120 dose (Flixotide
Powder for inhalation, 50 mcg per dose		60 dose C		Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose C	•	Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose (_	Floair
Aerosol inhaler, 125 mcg per dose CFC-free		120 dose (Flixotide
Aerosol inhaler, 250 mcg per dose		120 dose (Floair
Aerosol inhaler, 250 mcg per dose CFC-free		120 dose (DP 🗸	Flixotide
Powder for inhalation, 250 mcg per dose		60 dose C		Flixotide Accuhaler
Inhaled Long-acting Beta-adrenoceptor Ago	onists			
EFORMOTEROL FUMARATE				
Powder for inhalation, 12 mcg per dose, and monodose	e device20.64	60 dose		
	(35.80)			Foradil
EFORMOTEROL FUMARATE DIHYDRATE				
Powder for inhalation 4.5 mcg per dose, breath activate	ed.			
(equivalent to eformoterol furnarate 6 mcg metered		60 dose C	ıΡ	
(Squiralon to Sistinctorol lamarato o mog motoroc	(16.90)	00 0000 0	•	Oxis Turbuhaler
NDACATEROL	(10.00)			Calo i di ballaloi
	61.00	30 dose C	D ./	Onbrez Breezhaler
Powder for inhalation 150 mcgPowder for inhalation 300 mcg		30 dose C		Onbrez Breezhaler
•	01.00	ou dose C	, A	Olibiez Dieezlialer
SALMETEROL				
Aerosol inhaler CFC-free, 25 mcg per dose		120 dose (Serevent
Aerosol inhaler 25 mcg per dose		120 dose (Meterol
Powder for inhalation, 50 mcg per dose, breath activate	ed25.00	60 dose C)P /	Serevent Accuhaler
Inhaled Corticosteroids with Long-Acting B	eta-Adrenocept	or Agonis	sts	
BUDESONIDE WITH EFORMOTEROL				
Aerosol inhaler 100 mcg with eformoterol fumarate 6 m	ıca18.23	120 dose (OP 🗸	Vannair
Aerosor infraler 100 fficg with elofffoleror furnarate of ff		120 0036		
Powder for inhalation 100 mcg with eformoterol fumara		120 dose (Symbicort
Powder for inhalation 100 mcg with eformoterol fumara	te 6 mcg33.74		OP 🗸	Symbicort
Powder for inhalation 100 mcg with eformoterol fumara Aerosol inhaler 200 mcg with eformoterol fumarate 6 m	nte 6 mcg33.74	120 dose (OP ✓	Symbicort Turbuhaler 100/6
Powder for inhalation 100 mcg with eformoterol fumara	nte 6 mcg33.74	120 dose (OP ✓	Symbicort Turbuhaler 100/6 Vannair
Powder for inhalation 100 mcg with eformoterol fumara Aerosol inhaler 200 mcg with eformoterol fumarate 6 m Powder for inhalation 200 mcg with eformoterol fumara	acg21.40 tte 6 mcg44.08	120 dose (OP ✓	Symbicort Turbuhaler 100/6 Vannair Symbicort
Powder for inhalation 100 mcg with eformoterol fumarate 6 m Aerosol inhaler 200 mcg with eformoterol fumarate 6 m Powder for inhalation 200 mcg with eformoterol fumarate Powder for inhalation 400 mcg with eformoterol fumarate	ncg21.40 tte 6 mcg44.08 tte	120 dose (OP OP	Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6
Powder for inhalation 100 mcg with eformoterol fumara Aerosol inhaler 200 mcg with eformoterol fumarate 6 m Powder for inhalation 200 mcg with eformoterol fumara	ncg21.40 tte 6 mcg44.08 tte	120 dose (120 dose (120 dose (OP OP	Symbicort Turbuhaler 100/6 Vannair Symbicort
Powder for inhalation 100 mcg with eformoterol fumarate 6 m Powder for inhalation 200 mcg with eformoterol fumarate 6 m Powder for inhalation 200 mcg with eformoterol fumarate Powder for inhalation 400 mcg with eformoterol fumarate 12 mcg – No more than 2 dose per day	ncg21.40 tte 6 mcg44.08 tte	120 dose (120 dose (120 dose (OP OP	Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort
Powder for inhalation 100 mcg with eformoterol fumara Aerosol inhaler 200 mcg with eformoterol fumarate 6 m Powder for inhalation 200 mcg with eformoterol fumara Powder for inhalation 400 mcg with eformoterol fumara 12 mcg – No more than 2 dose per day	te 6 mcg33.74 lcg21.40 lte 6 mcg44.08 lte44.08	120 dose (120 dose (120 dose (60 dose C	OP VOP VOP VOP VOP VOP VOP VOP VOP VOP V	Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12
Powder for inhalation 100 mcg with eformoterol fumara Aerosol inhaler 200 mcg with eformoterol fumarate 6 m Powder for inhalation 200 mcg with eformoterol fumara Powder for inhalation 400 mcg with eformoterol fumara 12 mcg – No more than 2 dose per day	te 6 mcg33.74 lcg21.40 lte 6 mcg44.08 lte44.08	120 dose (120 dose (120 dose (OP VOP VOP VOP VOP VOP VOP VOP VOP VOP V	Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort
Powder for inhalation 100 mcg with eformoterol fumara Aerosol inhaler 200 mcg with eformoterol fumarate 6 m Powder for inhalation 200 mcg with eformoterol fumara Powder for inhalation 400 mcg with eformoterol fumara 12 mcg – No more than 2 dose per day	te 6 mcg33.74 lcg21.40 lte 6 mcg44.08 lte44.08	120 dose (120 do	OP VOP VOP VOP VOP VOP VOP VOP VOP VOP V	Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta
Powder for inhalation 100 mcg with eformoterol fumara Aerosol inhaler 200 mcg with eformoterol fumarate 6 m Powder for inhalation 200 mcg with eformoterol fumara Powder for inhalation 400 mcg with eformoterol fumara 12 mcg – No more than 2 dose per day	te 6 mcg33.74 lcg21.40 lte 6 mcg44.08 lte44.0844.0844.8	120 dose (120 dose (120 dose (60 dose C	OP / OP / OP / OP /	Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta
Powder for inhalation 100 mcg with eformoterol fumara Aerosol inhaler 200 mcg with eformoterol fumarate 6 m Powder for inhalation 200 mcg with eformoterol fumara Powder for inhalation 400 mcg with eformoterol fumara 12 mcg – No more than 2 dose per day	te 6 mcg33.74 tcg21.40 tte 6 mcg44.08 tte44.08 44.08 44.8 458 33.74	120 dose (120 do	OP / OP / OP / OP /	Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta RexAir Seretide
Powder for inhalation 100 mcg with eformoterol fumara Aerosol inhaler 200 mcg with eformoterol fumarate 6 m Powder for inhalation 200 mcg with eformoterol fumara Powder for inhalation 400 mcg with eformoterol fumara 12 mcg – No more than 2 dose per day	te 6 mcg33.74 tcg21.40 tte 6 mcg44.08 tte44.08 44.08 44.8 458 33.7416.83	120 dose (120 do	OP V OP V OP V OP V OP V	Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta RexAir Seretide RexAir
Powder for inhalation 100 mcg with eformoterol fumara Aerosol inhaler 200 mcg with eformoterol fumarate 6 m Powder for inhalation 200 mcg with eformoterol fumara Powder for inhalation 400 mcg with eformoterol fumara 12 mcg – No more than 2 dose per day	te 6 mcg33.74 tcg21.40 tte 6 mcg44.08 tte44.08 44.08 44.8 458 33.74 16.83 .44.08	120 dose (120 do	OP V OP V OP V OP V OP V	Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta RexAir Seretide
Powder for inhalation 100 mcg with eformoterol fumara Aerosol inhaler 200 mcg with eformoterol fumarate 6 m Powder for inhalation 200 mcg with eformoterol fumara Powder for inhalation 400 mcg with eformoterol fumara 12 mcg — No more than 2 dose per day	te 6 mcg33.74 log21.40 te 6 mcg44.08 te44.08 44.08 44.08 44.08 44.08 44.08 44.08 44.08 44.08 44.08 44.08 44.08 44.08 44.08	120 dose (120 do	OP / OP / OP / OP / OP / OP /	Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta RexAir Seretide RexAir Seretide
Powder for inhalation 100 mcg with eformoterol fumara Aerosol inhaler 200 mcg with eformoterol fumarate 6 m Powder for inhalation 200 mcg with eformoterol fumara 12 mcg - No more than 2 dose per day	te 6 mcg33.74 tcg21.40 tte 6 mcg44.08 tte44.08 44.08 44.08 458 33.7416.83 44.08 - No33.74	120 dose (120 do	OP / OP / OP / OP / OP / OP /	Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta RexAir Seretide RexAir
Powder for inhalation 100 mcg with eformoterol fumara Aerosol inhaler 200 mcg with eformoterol fumarate 6 m Powder for inhalation 200 mcg with eformoterol fumara Powder for inhalation 400 mcg with eformoterol fumara 12 mcg — No more than 2 dose per day	te 6 mcg33.74 tcg21.40 tte 6 mcg44.08 tte44.08 44.08 44.08 458 33.7416.83 44.08 - No33.74	120 dose (120 do	OP / OP / OP / OP / OP / OP /	Symbicort Turbuhaler 100/6 Vannair Symbicort Turbuhaler 200/6 Symbicort Turbuhaler 400/12 Breo Ellipta RexAir Seretide RexAir Seretide

				_
	Subsidy		Fully Brand or	
	(Manufacturer's			
	\$	Per	✓ Manufacturer	
Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Oral liq 400 mcg per ml	20.00	150 ml	✓ Ventolin	
Infusion 1 mg per ml, 5 ml		10	✓ Ventolin	
Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO		5	✓ Ventolin	
ing door may per mily i mile op to o my aramable on a roo mile				
Inhaled Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Aerosol inhaler, 100 mcg per dose CFC free - Up to 1000				
dose available on a PSO	3.80	200 dose OP	✓ Respigen	
			✓ SalAir	
	(6.00)		Ventolin	
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule - Up to 30 neb	, ,			
available on a PSO		20	✓ Asthalin	
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb				
available on a PSO		20	✓ Asthalin	
			<u>-141141111</u>	
TERBUTALINE SULPHATE	07.00	200 dose OP	A Briganul Turkubalar	
Powder for inhalation, 250 mcg per dose, breath activated	27.30	200 dose OP	Bricanyl Turbuhaler	
Anticholinergic Agents				
Antichonnergic Agents				
IPRATROPIUM BROMIDE				
Aerosol inhaler, 20 mcg per dose CFC-free - Up to 400 dos	е			
available on a PSO		200 dose OP	✓ Atrovent	
Nebuliser soln, 250 mcg per ml, 1 ml ampoule - Up to 40 ne	eb			
available on a PSO		20	✓ Univent	
Nebuliser soln, 250 mcg per ml, 2 ml ampoule - Up to 40 ne	eb			
available on a PSO		20	✓ Univent	
Univent to be Sole Supply on 1 January 2020				
Inhaled Beta-Adrenoceptor Agonists with Antic	holinergic A	Agents		
SALBUTAMOL WITH IPRATROPIUM BROMIDE				
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p	oer			
dose CFC-free	12.19	200 dose OP	✓ Duolin HFA	
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per				
vial, 2.5 ml ampoule - Up to 20 neb available on a PSO	5.20	20	✓ Duolin	
Long-Acting Muscarinic Antagonists				
GLYCOPYRRONIUM - Subsidy by endorsement				
a) Inhaled glycopyrronium treatment will not be subsidised if	f patient is also	receiving treatme	ent with subsidised tiotropium	ı or
umeclidinium.	-	-	'	
b) Glycopyrronium powder for inhalation 50 mcg per dose is	subsidised only	y for patients who	have been diagnosed as	
having COPD using spirometry, and the prescription is er	ndorsed accordi	ngly.	-	
Powder for inhalation 50 mcg per dose	61.00	30 dose OP	 Seebri Breezhaler 	

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

Powder for inhalation, 18 mcg per dose	50.37	30 dose	✓ Spiriva	
Soln for inhalation 2.5 mcg per dose	50.37	60 dose OP	✓ Spiriva F	Respimat

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA1584 at	ove – Retail ph	armacy
Powder for Inhalation 50 mcg with indacaterol 110 mcg81.00	30 dose OP	✓ Ultibro Breezhaler
TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584	above – Retail	pharmacy
Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg81.00	60 dose OP	✓ Spiolto Respimat

Antifibrotics

NINTEDANIB - Special Authority see SA1755 below - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

Cap 100 mg	2,554.00	60 OP	Ofev
Cap 150 mg	3,870.00	60 OP	Ofev

⇒SA1755 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and

continued...

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic	
	Per	1	Manufacturer	

continued...

- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA1748 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

⇒SA1748 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 80% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

	Subsidy	iaa\ (Fully	Brand or
	(Manufacturer's Pr \$	rice) ? Per	Subsidised ✓	Generic Manufacturer
Leukotriene Receptor Antagonists				
MONTELUKAST				
* Tab 4 mg		28	_	Montelukast Mylan
Montalulant Mulan to be Cale Cumply on 1 January 2000	5.25		•	Apo-Montelukast
Montelukast Mylan to be Sole Supply on 1 January 2020 * Tab 5 mg		28	1	Montelukast Mylan
1 Tub 0 Hig	5.50	20	_	Apo-Montelukast
Montelukast Mylan to be Sole Supply on 1 January 2020)			•
* Tab 10 mg	3.95	28	✓	Montelukast Mylan
	5.65			Accord \$29
Mantalula at Mulan ta ha Cala Cumhu an 1 January 2006	`		•	Apo-Montelukast
Montelukast Mylan to be Sole Supply on 1 January 2020)			
(Apo-Montelukast Tab 4 mg to be delisted 1 January 2020) (Apo-Montelukast Tab 5 mg to be delisted 1 January 2020)				
(Accord \$29 Tab 10 mg to be delisted 1 January 2020)				
(Apo-Montelukast Tab 10 mg to be delisted 1 January 2020)				
Mast Cell Stabilisers				
NEDOCROMIL				
Aerosol inhaler, 2 mg per dose CFC-free	28.07	112 dose	OP 🗸	Tilade
SODIUM CROMOGLICATE				
Aerosol inhaler, 5 mg per dose CFC-free	28.07	112 dose	OP 🗸	Intal Forte CFC Free
Methylxanthines				
Methylxantimies				
AMINOPHYLLINE				
* Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a		_		BBI 4 1 1 111
PSO	124.37	5	•	DBL Aminophylline
THEOPHYLLINE ** Tob long acting 250 mg	22.02	100	./	Nuelin-SR
* Tab long-acting 250 mg Nuelin-SR to be Sole Supply on 1 January 2020	23.02	100	•	Nueilli-on
* Oral liq 80 mg per 15 ml	16.60	500 ml	1	Nuelin
Nuelin to be Sole Supply on 1 January 2020				
Manadation				
Mucolytics				
DORNASE ALFA - Special Authority see SA0611 below - Retail	l pharmacy			
Nebuliser soln, 2.5 mg per 2.5 ml ampoule		6	✓	Pulmozyme
⇒SA0611 Special Authority for Subsidy				
Special Authority approved by the Cystic Fibrosis Advisory Panel				
Notes: Application details may be obtained from PHARMAC's w		.pharmac.	govt.nz or	:
,	04) 460 4990			
	: (04) 916 7571			
	FPanel@pharmac	-		
Prescriptions for patients approved for treatment must be written	by respiratory phy	ysicians o	r paediatri	cians who have experience
and expertise in treating cystic fibrosis.				
SODIUM CHLORIDE				
Not funded for use as a nasal drop. Soln 7%	24.50	90 ml O	p 🗸	Biomed
				Dioilieu

	Subsidy		Fully Brand or
	(Manufacturer's		idised Generic
	\$	Per	✓ Manufacturer
Nasal Preparations			
Allergy Prophylactics			
BECLOMETHASONE DIPROPIONATE			
Metered aqueous nasal spray, 50 mcg per dose	2.35 (5.26)	200 dose OP	Alanase
Metered aqueous nasal spray, 100 mcg per dose	2.46 (6.00)	200 dose OP	Alanase
(Alanase Metered aqueous nasal spray, 50 mcg per dose to be a (Alanase Metered aqueous nasal spray, 100 mcg per dose to be		•	
BUDESONIDE			
Metered aqueous nasal spray, 50 mcg per dose		200 dose OP	✓ <u>SteroClear</u>
Metered aqueous nasal spray, 100 mcg per dose FLUTICASONE PROPIONATE	2.87	200 dose OP	✓ <u>SteroClear</u>
Metered aqueous nasal spray, 50 mcg per dose	1.98	120 dose OP	✓ Flixonase Hayfever & Allergy
IPRATROPIUM BROMIDE			
Aqueous nasal spray, 0.03%	4.61	15 ml OP	✓ <u>Univent</u>
Respiratory Devices			
MASK FOR SPACER DEVICE			
a) Up to 50 dev available on a PSO			
b) Only on a PSO			
c) Only for children aged six years and under Small	2.20	1	✓ e-chamber Mask
	2.20	1	• e-chamber wask
PEAK FLOW METER			
a) Up to 25 dev available on a PSO			
b) Only on a PSO Low range	0.54	1	✓ Mini-Wright AFS
Low range		'	Low Range
Normal range	9.54	1	✓ Mini-Wright Standard
SPACER DEVICE			
a) Up to 50 dev available on a PSO			
b) Only on a PSO			
220 ml (single patient)	2.95	1	✓ e-chamber Turbo
510 ml (single patient)		1	✓ e-chamber La
			Grande
800 ml	6.50	1	✓ Volumatic
Respiratory Stimulants			
CAFFEINE CITRATE		05 100	4 B: 1

25 ml OP

✓ Biomed

	Subsidy		Fully	Brand or
	(Manufacturer's Pr	ica) Sul	osidised	Generic
	\$	Per	Joiuiseu	Manufacturer
	Ψ	FEI		Manuacturei
Ear Preparations				
ACETIC ACID MITH 4 O PROPANERIOL DIACETATE AND RE	NIZETUONIUM			
ACETIC ACID WITH 1, 2- PROPANEDIOL DIACETATE AND BE				
For Vosol ear drops with hydrocortisone powder refer Standa	ard Formulae, <mark>pag</mark>	je 235		
Ear drops 2% with 1, 2-Propanediol diacetate 3% and				
benzethonium chloride 0.02%	6 97	35 ml OP	✓ V	osol (
		00 1111 01	• •	0301
FLUMETASONE PIVALATE				
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	√ L	ocacorten-Viaform
				ED's
			. / I	ocorten-Vioform
			• -	ocorten-violoriii
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTATI	N		
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate				
	F 40	7.5 0.0		/aa.
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	• 1	Cenacomb
Ear/Eye Preparations				
, ,				
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN				
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and				
	4.50	0 1 OD		
gramicidin 50 mcg per ml		8 ml OP	_	
	(9.27)		S	Sofradex
FRAMYCETIN SULPHATE				
Ear/Eve drops 0.5%	4 12	8 ml OP		
Ear/Eye drops 0.5%		8 IIII OP	_	
	(8.65)		S	Soframycin
Eye Preparations				
Eye preparations are only funded for use in the eye, unless explicit	oitly stated athory	iico		
Lye preparations are only funded for use in the eye, unless expin	citiy stated otherw	nse.		
Anti-Infective Preparations				
Anti-infective r reparations				
ACICLOVIR				
	11.00	4.5 × OD		······
* Eye oint 3%	14.92	4.5 g OP	✓ V	'iruPOS
CHLORAMPHENICOL				
Eye oint 1%	2 48	4 g OP	✓ 0	Chlorsig
Eye drops 0.5%		10 ml OP		Chlorafast
			• <u>u</u>	illoralast
Funded for use in the ear*. Indications marked with * are	e unapproved ind	ications.		
CIPROFLOXACIN				
Eye drops 0.3% - Subsidy by endorsement	9 99	5 ml OP	√ 0	Ciprofloxacin Teva
			_	
When prescribed for the treatment of bacterial keratitis o				
for the second line treatment of chronic suppurative otitis		; and the pre	scription	is endorsed accordingly.
Note: Indication marked with a * is an unapproved indic	ation.			
GENTAMICIN SULPHATE				
	11 40	E ml OD		Conontio
Eye drops 0.3%	11.40	5 ml OP	• 6	Genoptic
PROPAMIDINE ISETHIONATE				
* Eye drops 0.1%	2 97	10 ml OP		
= -,0 3/000 0.1/0		10 1111 01	ь	Brolene
	(14.55)			NOIGHE
SODIUM FUSIDATE [FUSIDIC ACID]				
Eye drops 1%	5.29	5 g OP	√ F	ucithalmic
, ,		ŭ		

	Subsidy		Fully	Brand or
	(Manufacturer's P	rice) Subs	sidised	Generic
	\$	Per	✓	Manufacturer
TOBRAMYCIN				
Eye oint 0.3%	10.45	3.5 g OP	✓ T	obrex
Eye drops 0.3%		5 ml OP	✓ T	obrex
· ·				
Corticosteroids and Other Anti-Inflammatory Pr	eparations			
DEXAMETHASONE				
* Eye oint 0.1%	5.86	3.5 g OP	✓ N	Maxidex (
* Eye drops 0.1%		5 ml OP	✓ N	laxidex
Ocular implant 700 mcg - Special Authority see SA1680 bel				
- Retail pharmacy		1	✓ 0)zurdex

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not vet completed a family: and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

*	Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b			
	sulphate 6,000 u per g5.3	39	3.5 g OP	✓ Maxitrol
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml4.5	50	5 ml OP	✓ Maxitrol
DIC	CLOFENAC SODIUM			
	Eye drops 0.1%	80	5 ml OP	✓ Voltaren Ophtha

	Subsidy (Manufacturer's Pr	rian) Cuba	Fully	Brand or Generic
	(Manufacturer's Fr	Per Subs	siuiseu ✓	Manufacturer
FLUOROMETHOLONE				
* Eye drops 0.1%	3.09	5 ml OP	√ F	ML
	5.20		√ F	lucon
LEVOCABASTINE				
Eye drops 0.5 mg per ml	8.71	4 ml OP		
• •	(10.34)		L	ivostin
LODOXAMIDE				
Eye drops 0.1%	8.71	10 ml OP	√ L	.omide
PREDNISOLONE ACETATE				
Eye drops 1%	3.93	10 ml OP	✓ P	rednisolone-AFT
Year all a second	7.00	5 ml OP	✓ P	red Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority s	ee SA1715 below	- Retail pharr	nacv	
Eye drops 0.5%, single dose (preservative free)		20 dose		linims Prednisolone

⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM CROMOGLICATE

5 ml OP ✓ Cromal S29 ✓ Rexacrom

Rexacrom to be Sole Supply on 1 January 2020

Glaucoma Preparations - Beta Blockers

BETAXOLOL			
* Eye drops 0.25%	11.80	5 ml OP	✓ Betoptic S
* Eye drops 0.5%	7.50	5 ml OP	✓ Betoptic
TIMOLOL			
* Eye drops 0.25%	1.43	5 ml OP	✓ Arrow-Timolol
* Eye drops 0.25%, gel forming	3.30	2.5 ml OP	✓ Timoptol XE
* Eye drops 0.5%		5 ml OP	✓ Arrow-Timolol
* Eye drops 0.5%, gel forming	3.78	2.5 ml OP	✓ Timoptol XE
(Timoptol XE Eye drops 0.25%, gel forming to be deliste			•

Glaucoma Preparations - Carbonic Anhydrase Inhibitors

ACETAZOLAMIDE * Tab 250 mg	. 17.03	100	✓ <u>Diamox</u>
BRINZOLAMIDE	9.77	5 ml OP	✓ Azopt
DORZOLAMIDE HYDROCHLORIDE * Eye drops 2%	9.77	5 ml OP	
	(17.44)		Trusopt
DORZOLAMIDE WITH TIMOLOL * Eye drops 2% with timolol 0.5%	2.87	5 ml OP	✓ <u>Dortimopt</u>

	Subsidy (Manufacturer's Pr	rice) Subsi	Fully	Brand or Generic
	(Manufacturer's Fi	Per Per	uiseu ✓	Manufacturer
Glaucoma Preparations - Prostaglandin Analogu	ues			
BIMATOPROST				
* Eye drops 0.03%	3.30	3 ml OP		matoprost Multichem
LATANOPROST				
* Eye drops 0.005%	1.57	2.5 ml OP	✓ Te	eva
TRAVOPROST				
* Eye drops 0.004%		5 ml OP		avopt
	19.50	2.5 ml OP	✓ Tr	avatan
Glaucoma Preparations - Other				
BRIMONIDINE TARTRATE				
* Eye drops 0.2%	4.29	5 ml OP	✓ Ai	row-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE				
* Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ Co	ombigan
PILOCARPINE HYDROCHLORIDE				
* Eye drops 1%		15 ml OP		opto Carpine
* Eye drops 2% * Eve drops 4%		15 ml OP 15 ml OP		opto Carpine
* Eye drops 4%		13 1111 07	▼ 150	opto Carpine
* Eye drops 2% single dose – Special Authority see SA0895				
below – Retail pharmacy	31.95	20 dose	✓ Mi	inims Pilocarpine
				•

⇒SA0895 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics		
ATROPINE SULPHATE * Eye drops 1%	15 ml OP	✓ Atropt
CYCLOPENTOLATE HYDROCHLORIDE * Eye drops 1%	15 ml OP	✓ Cyclogyl
TROPICAMIDE * Eye drops 0.5%	15 ml OP 15 ml OP	✓ Mydriacyl✓ Mydriacyl
Preparations for Tear Deficiency		
For acetylcysteine eye drops refer Standard Formulae, page 235 HYPROMELLOSE * Eye drops 0.5%	15 ml OP	

(3.92)

Methopt

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

SENSORY ORGANS

	Subsidy (Manufacturer's P \$	rice) Subs	Fully sidised	Brand or Generic Manufacturer
HYPROMELLOSE WITH DEXTRAN * Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	✓ P	Poly-Tears
POLYVINYL ALCOHOL * Eye drops 1.4%	2.62	15 ml OP	√ V	/istil
* Eye drops 3%(Vistil Eye drops 1.4% to be delisted 1 January 2020) (Vistil Forte Eye drops 3% to be delisted 1 March 2020)	3.68	15 ml OP	✓ V	istil Forte

Preservative Free Ocular Lubricants

⇒SA1388 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER – Special Authority see SA1388 above – Retail pha	ırmacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL - Special Authori	ity see SA1388 ab	ove – Retai	il pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	 Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special Author	ority see SA1388	above – Re	tail pharmacy
Eye drops 1 mg per ml	22.00	10 ml OP	✓ Hylo-Fresh
Hylo-Fresh has a 6 month expiry after opening. The Pha	rmacy Procedures	s Manual re	striction allowing one bottle per
		. 00	1 1 1 1 1

month is not relevant and therefore only the prescribed dosage to the nearest OP may be claimed.

Other Eye Preparations

* Eye drops 0.1%4.15	15 ml OP	✓ Naphcon Forte
OLOPATADINE Eye drops 0.1%10.00	5 ml OP	✓ Patanol
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE Eye oint 138 mcg per g3.80	5 g OP	✓ VitA-POS

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

Various

PHARMACY SERVICES

May only be claimed once per patient.

1 fee **✓ BSF Logem**

✓ BSF Mylan Efavirenz Emtricitbane

Tenofov

- ✓ BSF Teva

 Atazanavir

 Sulphate
- ✓ BSF Teva Emtricitabine Tenofoir Disoprox
- a) The Pharmacode for BSF Teva Atazanavir Sulphate is 2573857 see also page 107
- b) The Pharmacode for BSF Teva Emtricitabine Tenofoir Disoprox is 2573865 see also page 103
- c) The Pharmacode for BSF Mylan Efavirenz Emtricitbane Tenofov is 2573873 see also page 106
- d) The Pharmacode for BSF Logem is 2575949 see also page 127

(BSF Logem Brand switch fee to be delisted 1 January 2020)

(BSF Mylan Efavirenz Emtricitbane Tenofov Brand switch fee to be delisted 1 December 2019)

(BSF Teva Atazanavir Sulphate Brand switch fee to be delisted 1 December 2019)

(BSF Teva Emtricitabine Tenofoir Disoprox Brand switch fee to be delisted 1 December 2019)

Agents Used in the Treatment of Poisonings

Antidotes

ACETYLCYSTEINE - Retail pharmacy-Specialist Inj 200 mg per ml, 10 ml ampoule58.76	10	✓ DBL Acetylcysteine
NALOXONE HYDROCHLORIDE		
a) Up to 5 inj available on a PSO		
b) Only on a PSO		
* Inj 400 mcg per ml, 1 ml ampoule22.60	5	✓ DBL Naloxone
		Hydrochloride

Removal and Elimination

വ				

*	Oral liq 50 g per 250 ml	43.50	250 ml OP	✓ Carbosorb-X
---	--------------------------	-------	-----------	---------------

a) Up to 250 ml available on a PSO

b) Only on a PSO

DEFERASIROX - Special Authority see SA1492 on the next page - Retail pharmacy

W	as	tage	cla	imat	ole
T	. L	105		ممناه	

Tab 125 mg dispersible	276.00	28	Exjade
Tab 250 mg dispersible	552.00	28	Exjade
Tab 500 mg dispersible	1,105.00	28	✓ Exjade

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.



Subsidy Fully (Manufacturer's Price) Subsidised		. ,	Brand or Generic
 \$	Per	/Siuiseu	Manufacturer

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE - Special Authority see SA1480 below -	Retail pharmacy		
Tab 500 mg	533.17	100	✓ Ferriprox
Oral lig 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox

⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESERBIOXAMINE MESILATE

* Inj 500 mg vial	84.53	10	✓ <u>DBL</u> Desferrioxamine Mesylate for Inj BP
SODIUM CALCIUM EDETATE * Inj 200 mg per ml, 5 ml	52 21	6	
* "1] 200 mg per mi, o mi	(156.71)	O	Calcium Disodium Versenate

OMEPRAZOLE SUSPENSION Omeprazole capules or powder

Water

Sodium bicarbonate powder BP

Standard Formulae ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml	qs	PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium	1 g
Suitable eye drop base	qs	Glycerol BP Water	70 ml to 100 ml
ASPIRIN AND CHLOROFORM APPLICATION Aspirin Soluble tabs 300 mg Chloroform	12 tabs to 100 ml	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml)	LIQUID (10
CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate Glycerol	60 mg 40 ml	Phenobarbitone Sodium Glycerol BP Water	400 mg 4 ml to 40 ml
Preservative Water	qs to 100 ml	PILOCARPINE ORAL LIQUID Pilocarpine 4% eye drops Preservative	qs qs
CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate Glycerol Preservative	300 mg 40 ml gs	Water (Preservative should be used if quantity supplied is than 5 days.)	to 500 ml
Water	to 100 ml	SALIVA SUBSTITUTE FORMULA	E a
FOLINIC MOUTHWASH Calcium folinate 15 mg tab Preservative Water	1 tab qs to 500 ml	Methylcellulose Preservative Water (Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	5 g qs to 500 ml for more
(Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)		SODIUM CHLORIDE ORAL LIQUID	
MAGNESIUM HYDROXIDE 8% MIXTURE Magnesium hydroxide paste 29%	275 g	Sodium chloride inj 23.4%, 20 ml Water (Only funded if prescribed for treatment of hyponatr	qs qs aemia)
Methyl hydroxybenzoate Water	1.5 g to 1,000 m	Il VANCOMYCIN ORAL SOLUTION (50 mg per ml) Vancomycin 500 mg injection	10 vials
METHADONE MIXTURE Methadone powder Glycerol Water	qs qs to 100 ml	Glycerol BP Water (Only funded if prescribed for treatment of Clostridiu following metronidazole failure)	40 ml to 100 ml um difficile
METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liqu	10 g to 100 ml uid mixture)	VOSOL EAR DROPS WITH HYDROCORTISONE POWDER 1% Hydrocortisone powder Vosol Ear Drops	1% to 35 ml

qs

8.4 g

to 100 ml

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy

(Manufacturer's Price)

Fully

Subsidised

Brand or

Generic

Per Manufacturer Extemporaneously Compounded Preparations and Galenicals BENZOIN Tincture compound BP......24.42 500 ml (39.90)Pharmacy Health (Pharmacy Health Tincture compound BP to be delisted 1 March 2020) CHI OROFORM a) Only in combination b) Maximum of 100 ml per prescription c) Only in aspirin and chloroform application. d) Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined. Chloroform BP......25.50 500 ml ✓ PSM CODEINE PHOSPHATE - Safety medicine: prescriber may determine dispensing frequency 25 q (90.09)Douglas Only in extemporaneously compounded codeine linctus. **COLLODION FLEXIBLE** Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined. 100 ml ✓ PSM COMPOUND HYDROXYBENZOATE - Only in combination Only in extemporaneously compounded oral mixtures. 100 ml Midwest GLYCERIN WITH SODIUM SACCHARIN - Only in combination Only in combination with Ora-Plus. 473 ml ✓ Ora-Sweet SF GLYCERIN WITH SUCROSE - Only in combination Only in combination with Ora-Plus. 473 ml ✓ Ora-Sweet **GLYCEROL** 500 ml ✓ healthE Glycerol BP Only in extemporaneously compounded oral liquid preparations. MAGNESIUM HYDROXIDE 500 q PSM (PSM Paste 29% to be delisted 1 July 2020) METHADONE HYDROCHLORIDE a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine: prescriber may determine dispensing frequency d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets). 1 q 🗸 AFT METHYL HYDROXYBENZOATE 25 q Midwest METHYLCELLULOSE ✓ MidWest 100 g 473 ml Ora-Plus

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Pric	e) Sub	Fully Brand or osidised Generic	
	\$	Per	✓ Manufacturer	
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHA	ARIN – Only in co	mbination		
Suspension		473 ml	✓ Ora-Blend SF	
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Only	in combination			
Suspension		473 ml	✓ Ora-Blend	
PHENOBARBITONE SODIUM				
Powder - Only in combination	52.50	10 g	✓ MidWest	
	325.00	100 g	✓ MidWest	
Only in children up to 12 years				
PROPYLENE GLYCOL				
Only in extemporaneously compounded methyl hydroxybenzo	ate 10% solution.			
Liq	11.25	500 ml	✓ Midwest	
SODIUM BICARBONATE				
Powder BP - Only in combination	10.05	500 g	✓ Midwest	
•	9.80	ŭ		
	(29.50)		David Craig	
a) Only in extemporaneously compounded omeprazole a	and lansoprazole s	suspension.		
b) Midwest to be Sole Supply on 1 January 2020	·	•		
(David Craig Powder BP to be delisted 1 January 2020)				
SYRUP (PHARMACEUTICAL GRADE) - Only in combination				
Only in extemporaneously compounded oral liquid preparation	ns.			
Liq		500 ml	✓ Midwest	
Midwest to be Sole Supply on 1 January 2020				
WATER				
Tap - Only in combination	0.00	1 ml	✓ Tap water	

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1522 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 inborn errors of metabolism: or
- 7 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

✓ fully subsidised 239

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per		Manufacturer

continued...

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT – Special Authority see SA1523 on the previous page – Hospital pharmacy [HP3]

Emulsion (neutral)	12.30 200	ml OP	Calogen
	30.75 500	ml OP 🗸	Calogen
Emulsion (strawberry)	12.30 200	ml OP 🗸	Calogen
Oil	30.00 500	ml OP 🗸	MCT oil (Nutricia)
Oil, 250 ml1	14.92 4	OP 🗸	Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT	 Special Authority see SA1524 above – Hospital p 	narmacy [HP3]	
Powder	7.90	225 g OP	✓ Protifar
	8.95	227 g OP	✓ Resource
		•	Beneprotein

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

Sustagen Diabetic

Oral Supplements/Complete Diet (Nasogastric/Gastrostomy Tube Feed)

Respiratory Products

⇒SA1094 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has CORD and hypercapnia, defined as a CO2 value exceeding 55 mmHg.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see SA1095 above	 Hospital pharm 	nacy [HP3]
Liquid7.50	1,000 ml OP	Diason RTH
		✓ Glucerna Select RTH
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 above - Ho	ospital pharmacy	[HP3]
Liquid (strawberry)1.50	200 ml OP	✓ Diasip
Liquid (vanilla)1.50	200 ml OP	✓ Diasip
1.88	250 ml OP	✓ Glucerna Select
1.78	237 ml OP	
(2.10)		Resource Diabetic

(2.10)



Subsidy (Manufacturer's Price) Fully Subsidised Brand or Generic Manufacturer

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism: or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED − Special Authority see SA1525 above − Hospital pharmacy [HP3]
Powder60.48 400 g OP
✓ Monogen

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	•	Manufacturer

ENTERAL/ORAL FEED 1KCAL/ML − Special Authority see SA1099 on the previous page − Hospital pharmacy [HP3] Liquid.......54.00 400 g OP ✓ Kindergen

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1379 a Liquid6.00	bove – Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini Energy RTH
PAEDIATRIC ENTERAL FEED 1KCAL/ML – Special Authority see SA1379 about Liquid	ove – Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini RTH ✓ Pediasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority s Liquid6.00	see SA1379 above − Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA1379 above	e – Hospital pharmacy [HP3]
Liquid (strawberry)1.60	200 ml OP 🗸 Fortini
Liquid (vanilla)	200 ml OP ✓ Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA1379 above -	- Hospital pharmacy [HP3]
Liquid (chocolate)1.07	200 ml OP ✓ Pediasure
Liquid (strawberry)1.07	200 ml OP ✓ Pediasure
Liquid (vanilla)	200 ml OP ✓ Pediasure
1.34	250 ml OP ✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority see Liquid (unflavoured)	
	200 ml OP ✓ Fortini Multi Fibre
Liquid (chocolate)	200 ml OP ✓ Fortini Multi Fibre
Liquid (strawberry)	200 ml OP ✓ Fortini Multi Fibre
Liquid (vanilla)1.60	
PEPTIDE-BASED ORAL FEED - Special Authority see SA1379 above - Hospi	, ,, ,
Powder43.60	400 g OP ✓ Peptamen Junior

Subsidy	Fully	Brand or
,	. ,	
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML – Special Authority see Liquid			,
RENAL ORAL FEED 1.8 KCAL/ML - Special Authority see SA	101 above – Hos	pital pharmacy	[HP3]
Liquid	2.67	220 ml OP	
			(strawberry)
			✓ Nepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA11	01 above - Hospi	ital pharmacy [H	HP3]
Liquid	2.88	237 ml OP	
	(3.31)		NovaSource Renal
Liquid (apricot) 125 ml	11.52	4 OP	✓ Renilon 7.5
Liquid (caramel) 125 ml	11.52	4 OP	Renilon 7.5

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Brand or

Fully

	(Manufacturer's F	Price) Subs Per	idised Generic ✓ Manufacturer
ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML — Spe pharmacy [HP3] Liquid		e SA1377 on th 1,000 ml OP	e previous page – Hospital Vital
ORAL ELEMENTAL FEED 0.8KCAL/ML — Special Authority see Liquid (grapefruit), 250 ml cartonLiquid (pineapple & orange), 250 ml cartonLiquid (summer fruits), 250 ml carton	171.00 171.00	previous page - 18 OP 18 OP 18 OP	- Hospital pharmacy [HP3] ✓ Elemental 028 Extra ✓ Elemental 028 Extra ✓ Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see S Powder (unflavoured)	4.50	80 g OP	✓ Vivonex TEN
[HP3] Liquid	·	1,000 ml OP	✓ Peptisorb

Subsidy

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML - Special Authority see SA1196 above - Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini Low Energy Liquid 4.00 Multi Fibre

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

continued...

✓ fully subsidised 245

bsidy	Fully	Brand or
turer's Price) Subsid	dised	Generic
 \$ Per	✓	

continued...

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or

continued...

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per ✓	Manufacturer	

continued...

- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 la baina fad v
 - 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
 - 2 Cystic Fibrosis; or
 - 3 Liver disease; or
 - 4 Chronic Renal failure; or
 - 5 Inflammatory bowel disease; or
 - 6 Chronic obstructive pulmonary disease with hypercapnia; or
 - 7 Short bowel syndrome: or
 - 8 Bowel fistula: or
 - 9 Severe chronic neurological conditions; or
 - 10 Epidermolysis bullosa: or
 - 11 AIDS (CD4 count < 200 cells/mm³); or
 - 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications

continued...

SPECIAL FOODS

Su		Fully	Brand or
(Manufact		dised	Generic
	\$ Per	•	Manufacturer

continued...

meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or

9 Severe chronic neurological conditions.		
ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 245 - F Liquid7.00		[HP3] ✓ Nutrison Energy
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on page 245 - Ho Liquid	250 ml OP	HP3] ✓ Isosource Standard ✓ Nutrison Standard RTH ✓ Osmolite RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML – Special Authority see SA1859 CLiquid		spital pharmacy [HP3] Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority see SA1859 on p Liquid5.29		al pharmacy [HP3] ✓ Jevity RTH ✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority see SA1859 on Liquid1.75 7.00	250 ml OP	ital pharmacy [HP3] ✓ Ensure Plus HN ✓ Ensure Plus RTH ✓ Jevity HiCal RTH ✓ Nutrison Energy

Multi Fibre

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

ORAL FEED (POWDER) - Special Authority see SA1859 on page 245 - Hospital pharmacy [HP3]

Note: Higher subsidy for Sustagen Hospital Formula will only be reimbursed for patients with both a valid Special Authority number and an appropriately endorsed prescription.

Powder (chocolate) - Higher subsidy of up to \$26.00 per 850 g			
with Endorsement	26.00	850 g OP	✓ Ensure
	9.54	840 g OP	
	(26.00)	•	Sustagen Hospital
			Formula Active

Additional subsidy by endorsement is available for patients with fat malabsorption, fat intolerance or chyle leak. The prescription must be endorsed accordingly.

Powder (vanilla) – Higher subsidy of up to \$26.00 per 850 g			
with Endorsement	8.54	857 g OP	✓ Fortisip
	26.00	850 g OP	✓ Ensure
	9.54	840 g OP	
	(26.00)	•	Sustagen Hospital
			Formula Active

Additional subsidy by endorsement is available for patients with fat malabsorption, fat intolerance or chyle leak. The prescription must be endorsed accordingly.

ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 245 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease. The prescription must be endorsed accordingly.

Liquid (banana) — Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with	, ,		·
Endorsement		200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
Liquid (fruit of the forest) - Higher subsidy of \$1.26 per 200 ml			
with Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.33 per 237 ml with			
Endorsement	0.85	237 ml OP	
	(1.33)		Ensure Plus
	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip

✓ fully subsidised

249

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsic	lised	Generic	
 \$	Per	1	Manufacturer	

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 245 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with

Endorsement	0.72	200 ml OP	
LIIU0156III6III	(1.26)	200 IIII OF	Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with	` ,		·
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 above -	Hospital p	harmacy [HP3]	
Liquid	5.50	500 ml OP	✓ Nutrison
			Concentrated
	11.00	1,000 ml OP	✓ Two Cal HN RTH

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$1.90 per 200 ml with

00) Two Cal HN

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by PHARMAC from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

Powder Powder		⁷ 3]
	(5.15)	Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see S	SA1729 above – Hospital pharmacy [HP:	3]
Powder	3.93 1,000 g OP	
	(7.32)	NZB Low Gluten Bread Mix
	3.51	
	(10.87)	Horleys Bread Mix

✓ fully subsidised 251

	Subsidy (Manufacturer's Price) \$	Sub:	Fully sidised	Brand or Generic Manufacturer
GLUTEN FREE FLOUR - Special Authority see SA17	'29 on the previous page – Hos	oital pharr	nacy [H	P3]
Powder		00 g OP		
	(18.10)		Н	lorleys Flour
GLUTEN FREE PASTA - Special Authority see SA17	29 on the previous page - Hosp	ital pharn	nacy [Hi	P3]
Buckwheat Spirals	2.00 25	60 g OP	, .	•
	(3.11)	•	С)rgran
Corn and Vegetable Shells	2.00 25	60 g OP		-
	(2.92)		C	Orgran
Corn and Vegetable Spirals	2.00 25	60 g OP		
	(2.92)		С	Orgran
Rice and Corn Lasagne Sheets	1.60 20	10 g OP		
	(3.82)		С)rgran
Rice and Corn Macaroni		60 g OP		
	(2.92)		С)rgran
Rice and Corn Penne		60 g OP	_	
	(2.92)		С	Orgran
Rice and Maize Pasta Spirals		60 g OP	_	
	(2.92)		C	Orgran
Rice and Millet Spirals		60 g OP	_	
	(3.11)	- 00	C	Orgran
Rice and com spaghetti noodles Vegetable and Rice Spirals		'5 g OP	_	
	(2.92)	·0 OP	C	Orgran
		60 g OP	_	\
Italian long style spaghetti	(2.92)	00 ~ OP	C	Orgran
		20 g OP	^	\u00e4an
	(3.11)		C)rgran

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 above - Hospital pharmacy [HP3]

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
•	Por	1	Manufacturer	

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

Tabs	99.00	75 OP	✓ Phlexy 10
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior Chocolate
Powder (unflavoured) 27.8 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior Vanilla
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange)	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured)	320.00	500 g OP	XP Maxamum
Liquid (berry)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	 Easiphen Liquid
Liquid (juicy tropical) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g	1,123.20	36 OP	✓ PKU Lophlex Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20

Foods

LOW PROTEIN BAKING WIX - Special Authority see	SATIOS on the previous pa	ge – Hospitai p	marmacy [HP3]
Powder	8.22	500 g OP	✓ Loprofin Mix
LOW PROTEIN PASTA - Special Authority see SA110	08 on the previous page – 1	Hospital pharm	acy [HP3]
Animal shapes	11.91	500 g OP	Loprofin
Lasagne	5.95	250 g OP	Loprofin
Low protein rice pasta	11.91	500 g OP	Loprofin
Macaroni	5.95	250 g OP	Loprofin

 Spaghetti
 11.91
 500 g OP
 ✓ Loprofin

 Spirals
 11.91
 500 g OP
 ✓ Loprofin

✓ fully subsidised 253

500 g OP

✓ Loprofin

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per Brand or Generic Manufacturer

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]
Powder44.40 400 g OP ✓ Locasol

Gastrointestinal and Other Malabsorptive Problems

AMINO ACID FORMULA – Special Authority see SA1219 below - Powder		y [HP3] 00 a OP	✓ Alfamino Junior
Powder (unflavoured)		00 g OP	✓ Elecare
· · ·		•	✓ Elecare LCP
			✓ Neocate Gold
			 Neocate Junior Unflavoured
			✓ Neocate SYNEO
Powder (vanilla)	53.00 4	00 g OP	✓ Elecare
			✓ Neocate Junior Vanilla

⇒SA1219 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Extensively hydrolysed formula has been reasonably trialled and is inappropriate due to documented severe intolerance or allergy or malabsorption; or
- 2 History of anaphylaxis to cows milk protein formula or dairy products; or
- 3 Eosinophilic oesophagitis.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's Prio \$	ce) Subs Per	Fully sidised	Brand or Generic Manufacturer
EXTENSIVELY HYDROLYSED FORMULA – Special Authority s Powder			✓ A	y [HP3] ptamil Gold+ Pepti Junior
	30.42	900 g OP		llerpro 1 llerpro 2

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption: or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea: or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption: or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption: or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure: or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML − Special Authority see SA1698 below − Hospital pharmacy [HP3] Liquid......2.35 125 ml OP ✓ Infatrini

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

continued...

✓ fully subsidised 255



Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Vaccinations

ADULT DIPHTHERIA AND TETANUS VACCINE - [Xpharm]

- 1) For vaccination of patients aged 45 and 65 years old; or
- 2) For vaccination of previously unimmunised or partially immunised patients; or
- 3) For revaccination following immunosuppression; or
- 4) For boosting of patients with tetanus-prone wounds; or
- 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or

www.bcgatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

10 **✓ BCG Vaccine**

DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE - [Xpharm]

Funded for any of the following criteria:

- 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
- 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
- 3) A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
- 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens.

Notes: Tdap is not registered for patients aged less than 10 years. Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid, 8 mcg pertussis filamentous

10

✓ Boostrix
✓ Boostrix

	Subsidy (Manufacturer's Price) \$	F Subsidi Per	Fully Brand or sed Generic Manufactur	er
DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE Funded for any of the following: 1) A single dose for children up to the age of 7 who have 2) A course of four vaccines is funded for catch up prograprimary immunisation; or 3) An additional four doses (as appropriate) are funded for pre- or post splenectomy; pre- or post solid organ trans regimens; or 4) Five doses will be funded for children requiring solid or Note: Please refer to the Immunisation Handbook for approing 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagluttinin, 8 mcg pertactin and 80 D-antigen units poliomyelitis virus in 0.5ml syringe	— [Xpharm] completed primary im ammes for children (to primary implementation for call and primary implementation. gan transplantation. priate schedule for call and primary implementation. MND HAEMOPHILUS of 10 for primary immular (re-)immunisation for primary immular (re-)immunisation for call and primary immular (re-)immunisation for primary immular (re-)immunisatio	munisation; the age of 1 r patients poind other seventh of the progration of the pro	or 0 years) to compost HSCT, or checerely immunosurammes. Infanrix IPV LE TYPE B VACCO	olete full motherapy; opressive
10 who are patients post haematopoietic stem cell tran post solid organ transplant, renal dialysis and other set 3) Up to five doses for children up to and under the age of Note: A course of up-to four vaccines is funded for catch up to complete full primary immunisation. Please refer to the Inprogrammes. Inj 30IU diphtheriatoxoid with 40IU tetanustoxoid, 25mcg pertussistoxoid, 25mcg pertussisfilamentoushaemagluttinin, 8 mcgpertactin, 80 D-AgUpoliovirus, 10mcghepatitisBsurfaceantigen in 0.5ml syringe	verely immunosuppres of 10 receiving solid ord programmes for child nmunisation Handboo	ssive regime gan transpla Iren (up to a	ns; or ntation. nd under the age	of 10 years) e for catch up
AEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm] One dose for patients meeting any of the following: 1) For primary vaccination in children; or 2) An additional dose (as appropriate) is funded for (re-)ir transplantation, or chemotherapy; functional asplenic; or post cochlear implants, renal dialysis and other seve 3) For use in testing for primary immunodeficiency diseas paediatrician. Haemophilus Influenzae type B polysaccharide 10 mcg	pre or post splenecton erely immunosuppress	ny; pre- or p	ost solid organ tra s; or	ansplant, pre-
conjugated to tetanus toxoid as carrier protein 20-40 morprefilled syringe plus vial 0.5 ml	0.00 disease; or	1	✓ <u>Hiberix</u>	
Inj 1440 ELISA units in 1 ml syringe Inj 720 ELISA units in 0.5 ml syringe		1	✓ <u>Havrix</u> ✓ <u>Havrix Junio</u>	<u>or</u>

		Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
		\$	Per	1	Manufacturer
HEPATITIS F	RECOMBINANT VACCINE - [Xpharm]				
	per 0.5 ml vial	0.00	1	1	HBvaxPRO
, ,	led for patients meeting any of the following criteria:		'	•	IIDVAXI IIO
	for household or sexual contacts of known acute h		onat	itic B carrie	ore: or
,	for children born to mothers who are hepatitis B su				513, 01
,	for children up to and under the age of 18 years inc	0 1			achieved a nositive
0)	serology and require additional vaccination or requ				
۵)	for HIV positive patients; or	ire a primary course o	n vac	omation, o	
,	for hepatitis C positive patients; or				
	for patients following non-consensual sexual interc	ourse: or			
	for patients following immunosuppression; or	ouroo, or			
	for solid organ transplant patients; or				
	for post-haematopoietic stem cell transplant (HSC	patients: or			
	following needle stick injury.	, panomo, o			
,					
Ini 10 mg	g per 1 ml vial	0.00	1	1	HBvaxPRO
•	led for patients meeting any of the following criteria:		•		
	for household or sexual contacts of known acute h		enat	itis B carrie	ers. Ur
	for children born to mothers who are hepatitis B su				310, 01
,	for children up to and under the age of 18 years inc	0 1			ve achieved a positive
٠,	serology and require additional vaccination or requ				
4)	for HIV positive patients; or			, -	
	for hepatitis C positive patients; or				
	for patients following non-consensual sexual interc	ourse; or			
	for patients following immunosuppression; or	,			
8)	for solid organ transplant patients; or				
9)	for post-haematopoietic stem cell transplant (HSC)	Γ) patients; or			
10)	following needle stick injury.				
				_	
	g per 1 ml prefilled syringe		1	/	Engerix-B
	led for patients meeting any of the following criteria:				
	for household or sexual contacts of known acute h				ers; or
	for children born to mothers who are hepatitis B su				
3)	for children up to and under the age of 18 years inc				
	serology and require additional vaccination or requ	ire a primary course of	f vac	cination; o	r
	for HIV positive patients; or				
	for hepatitis C positive patients; or				
,	for patients following non-consensual sexual interc	ourse; or			
,	for patients following immunosuppression; or				
,	for solid organ transplant patients; or	T\			
,	for post-haematopoietic stem cell transplant (HSC) patients; or			
,	following needle stick injury; or				
,	for dialysis patients; or				
12)	for liver or kidney transplant patients.				
Ini 40 mo	g per 1 ml vial	0.00	1	1	HBvaxPRO
	led for any of the following criteria:		'	•	IID VANE ITO
	for dialysis patients; or				
,	for liver or kidney transplant patient.				
2)	ioi iivoi oi kiuney iianopiani palleni.				

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
` \$	Per	✓	Manufacturer	

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV] - [Xpharm]

Any of the following:

- 1) Maximum of two doses for children aged 14 years and under; or
- 2) Maximum of three doses for patients meeting any of the following criteria:
 - 1) People aged 15 to 26 years inclusive; or
 - 2) Either:

People aged 9 to 26 years inclusive

- 1) Confirmed HIV infection; or
- 2) Transplant (including stem cell) patients: or
- 3) Maximum of four doses for people aged 9 to 26 years inclusive post chemotherapy

✓ Gardasil 9 10

Subsidy (Manufacturer's Price)	Subsid	Fully dised	Brand or Generic
 \$	Per	•	Manufacturer

INFLUENZA VACCINE

Inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine) -

[Xpharm]......9.00 1 ✓ Fluarix Tetra

A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by PHARMAC:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes: or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.

B) INFLUENZA VACCINE - pregnant women

- a) are pregnant
- C) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

5 ✓ FluQuadri	5	Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)45.00
10	10	90.00
✓ Influyac Tet		

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

- a) Only on a prescription
- b) No patient co-payment payable

С

A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by PHARMAC:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or
- c) children aged four years or less (but over three years) who have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

	Subsidy	Fully	Brand or
	acturer's Price)	Subsidised	Generic
(manuse	\$ Per	✓	Manufacturer

MEASLES. MUMPS AND RUBELLA VACCINE

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment from the Funder for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50, Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional
 or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
- 2) One dose for close contacts of meningococcal cases; or
- 3) A maximum of two doses for bone marrow transplant patients; or
- 4) A maximum of two doses for patients following immunosuppression*.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 4 mcg of each meningococcal polysaccharide conjugated to

a total of approximately 48 mcg of diphtheria toxoid carrier

MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm]

Any of the following:

- Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional
 or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
- 2) One dose for close contacts of meningococcal cases: or
- 3) A maximum of two doses for bone marrow transplant patients; or
- 4) A maximum of two doses for patients following immunosuppression*.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm]

Either:

- 1) A primary course of four doses for previously unvaccinated individuals up to the age of 59 months inclusive; or
- Up to three doses as appropriate to complete the primary course of immunisation for individuals under the age of 59 months who have received one to three doses of PCV13.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B,

7F, 9V, 14 and 23F; 3 mcg of pneumococcal

polysaccharide serotypes 4, 18C and 19F in 0.5 ml

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- One dose is funded for high risk children (over the age of 17 months and under 18 years) who have previously received four doses of PCV10; or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) with primary immune deficiencies: or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - n) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - i) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cyanosis or failure; or
 - I) with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency: or
- For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

	NATIONAL	IMMUNISA	TION SCHEDULE
	Subsidy (Manufacturer's Price) \$	Full Subsidise Per	,
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE	– [Xpharm]		
Either:			
 Up to three doses (as appropriate) for patients with I chemotherapy; pre- or post-splenectomy or with function complement deficiency (acquired or inherited), cochi All of the following: a) Patient is a child under 18 years for (re-)immur 	etional asplenia, pre- or ear implants, or primary	post-solid orga	n transplant, renal dialysis,
b) Treatment is for a maximum of two doses; and			
c) Any of the following: i) on immunosuppressive therapy or radiati immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; v) who are immune-suppressed following or or vi) with cochlear implants or intracranial shu vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more	or gan transplantation (inc nts; or than two weeks, and wh	luding haemate no are on an eq	opoietic stem cell transplant); uivalent daily dosage of
prednisone of 2 mg/kg per day or greater 20 mg or greater; or ix) with chronic pulmonary disease (including x) pre term infants, born before 28 weeks grail with cardiac disease, with cyanosis or fair xiii) with diabetes; or xiiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with	g asthma treated with hisestation; or ure; or		, ,
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	0.00	1	Pneumovax 23
POLIOMYELITIS VACCINE - [Xpharm] Up to three doses for patients meeting either of the followi 1) For partially vaccinated or previously unvaccinated in 2) For revaccination following immunosuppression.	ng: ndividuals; or		
Note: Please refer to the Immunisation Handbook for app Inj 80D antigen units in 0.5 ml syringe			nmes. ´ IPOL
ROTAVIRUS ORAL VACCINE – [Xpharm] Maximum of two doses for patients meeting the following: 1) first dose to be administered in infants aged under 1 2) no vaccination being administered to children aged 2	-		
Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator	0.00	10	Rotarix

	Subsidy (Manufacturer's Price) \$	Subsic Per	Fully lised	Brand or Generic Manufacturer
/ARICELLA VACCINE [CHICKENPOX VACCINE] – [Xpharm]				
Either:				
 Maximum of one dose for primary vaccination for either a) Any infant born on or after 1 April 2016; or 	er.			
 b) For previously unvaccinated children turning 11 y varicella infection (chickenpox), or 	ears old on or after 1	July 2017,	who ha	ave not previously had a
2) Maximum of two doses for any of the following:				
 a) Any of the following for non-immune patients: 				
 i) with chronic liver disease who may in future ii) with deteriorating renal function before tran iii) prior to solid organ transplant; or 		nsplantation	n; or	
iv) prior to solid organ transplant, or iv) prior to any elective immunosuppression*,	or			
v) for post exposure prophylaxis who are imm		nts.; or		
b) For patients at least 2 years after bone marrow to				
c) For patients at least 6 months after completion o				
 d) For HIV positive non immune to varicella with mi e) For patients with inborn errors of metabolism at r varicella, or 				
f) For household contacts of paediatric patients wh immune compromise where the household conta				ing a procedure leading to
g) For household contacts of adult patients who have immunocompromised, or undergoing a procedur has no clinical history of varicella.	ve no clinical history of e leading to immune co	varicella a ompromise	nd who where	are severely the household contact
* immunosuppression due to steroid or other immunosuppre 28 days	essive therapy must be	for a treatr	nent p	eriod of greater than
Inj 2000 PFU prefilled syringe plus vial	0.00	1 10		arilrix arilrix
	ED VACCINE ISHINGI	LES VACC	INE] -	[Xpharm]
/ARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUAT Funded for patients meeting either of the following criteria:				
	•	2018 and	31 Ma	rch 2020.
Funded for patients meeting either of the following criteria: 1) One dose for all people aged 65 years; or	s inclusive from 1 April	2018 and 3	✓ Z	rch 2020. ostavax ostavax
Funded for patients meeting either of the following criteria: 1) One dose for all people aged 65 years; or 2) One dose for all people aged between 66 and 80 year	s inclusive from 1 April	1	✓ Z	ostavax

- Symbols -		AFT-Pyrazinamide	100	Anaesthetics	119
3TC	106	Agents Affecting the		Anagrelide hydrochloride	160
- A -		Renin-Angiotensin System	45	Analgesics	120
A-Scabies	67	Agents for Parkinsonism and Rela	ited	Anastrozole	173
Abacavir sulphate	106	Disorders	118	Andriol Testocaps	80
Abacavir sulphate with		Agents Used in the Treatment of		Androderm	80
lamivudine	106	Poisonings	233	Anoro Ellipta	224
Abiraterone acetate	171	Agrylin	160	Antabuse	152
Acarbose	11	Alanase	227	Antacids and Antiflatulents	6
Acarbose Mylan	11	Albendazole	89	Anten	124
Accarb	11	Albey	220	Anthelmintics	89
Accuretic 10	46	Albustix	77	Antiacne Preparations	59
Accuretic 20	46	Aldurazyme	28	Antiallergy Preparations	220
Acetazolamide	230	Alendronate sodium	111	Antianaemics	35
Acetic acid with 1, 2- propane	diol	Alendronate sodium with		Antiandrogen Oral	
diacetate and		colecalciferol	111	Contraceptives	75
benzethonium	228	Alfacalcidol	32	Antiarrhythmics	
Acetic acid with hydroxyquino	line and	Alfamino Junior	254	Antibacterials	89
ricinoleic acid		Alginic acid	6	Antibacterials Topical	
Acetylcysteine		Alglucosidase alfa		Anticholinergic Agents	
Aci-Jel		Alkeran		Anticholinesterases	
Aciclovir		Allerpro 1		Antidepressants	
Infection	102	Allerpro 2		Antidiarrhoeals	
Sensory		Allersoothe		Antiepilepsy Drugs	
Acidex		Allmercap		Antifibrinolytics, Haemostatics and	
Acipimox		Allopurinol		Local Sclerosants	
Acitretin		Alpha-Adrenoceptor Blockers		Antifibrotics	
Aclasta		Alpha-Keri Lotion		Antifungals	
Aclin		Alphamox		Antifungals Topical	
Actemra		Alphamox 125		Antihistamines	
Actinomycin D		Alphamox 250		Antihypotensives	
Actrapid		Alprolix		Antimalarials	
Actrapid Penfill		Alu-Tab		Antimigraine Preparations	
Acupan		Aluminium hydroxide		Antinausea and Vertigo Agents	
Adalat 10		Amantadine hydrochloride		Antiparasitics	
Adalat Oros		Ambrisentan		Antipruritic Preparations	
Adalimumab		Amiloride hydrochloride		Antipsychotics	
Adapalene		Amiloride hydrochloride with		Antiretrovirals	
Adefin		furosemide	52	Antirheumatoid Agents	
Adefin XL		Amiloride hydrochloride with		Antispasmodics and Other Agents	
Adefovir dipivoxil		hydrochlorothiazide	52	Altering Gut Motility	
Adenuric		Aminophylline		Antithrombotic Agents	
ADR Cartridge 1.8		Amiodarone hydrochloride		Antithymocyte globulin	
Adrenaline		Amisulpride		(equine)	180
ADT Booster		Amitriptyline		Antitrichomonal Agents	
Adult diphtheria and tetanus		Amlodipine		Antituberculotics and	
vaccine	257	Amorolfine		Antileprotics	gg
Advantan		Amoxicillin		Antiucprotics	
Advate		Amoxicillin with clavulanic acid		Antivirals	
Adynovate		Amphotericin B		Anxiolytics	
Afinitor		Amsacrine		Anzatax	
Aflibercept		AmsaLyo		Apidra	
Afluria Quad		Amsidine		Apidra SoloStar	
AFT Carbimazole		Amzoate		Apo-Amlodipine	
Outomia_010		outo	20	p. /	

Apo-Amoxi	92	Arrow-Doxorubicin	162	B-D Micro-Fine	13
Apo-Azithromycin	90	Arrow-Fluoxetine	125	B-D Ultra Fine	14
Apo-Bromocriptine	118	Arrow-Lamotrigine	127	B-D Ultra Fine II	14
Apo-Ciclopirox	60	Arrow-Losartan &		Bacillus Calmette-Guerin (BCG)	
Apo-Cilazapril		Hydrochlorothiazide	46	vaccine	180
Apo-Cilazapril/		Arrow-Morphine LA	122	Bacillus Calmette-Guerin	
Hydrochlorothiazide	46	Arrow-Norfloxacin	109	vaccine	257
Apo-Clarithromycin		Arrow-Ornidazole	99	Baclofen	117
Alimentary	8	Arrow-Quinapril 10	46	Bactroban	6C
Infection	90	Arrow-Quinapril 20	46	Barrier Creams and Emollients	
Apo-Clomipramine	123	Arrow-Quinapril 5	46	BCG Vaccine	257
Apo-Diclo SR	110	Arrow-Roxithromycin	91	Beclazone 100	221
Apo-Diltiazem CD	<mark>50</mark>	Arrow-Sertraline	125	Beclazone 250	221
Apo-Doxazosin		Arrow-Timolol	230	Beclazone 50	221
Apo-Folic Acid	36	Arrow-Tolterodine	77	Beclomethasone	
Apo-Furosemide		Arrow-Topiramate		dipropionate22	21, <mark>22</mark> 7
Apo-Gabapentin	126	Arrow-Tramadol	123	Bee venom allergy treatment	220
Apo-Leflunomide		Arsenic trioxide	160	Bendamustine hydrochloride	155
Apo-Megestrol	172	Asacol	7	Bendrofluazide	52
Apo-Metoprolol	49	Asamax	7	Bendroflumethiazide	
Apo-Mirtazapine	125	Ascorbic acid	31	[Bendrofluazide]	
Apo-Montelukast	226	Aspen Adrenaline	54	Benzathine benzylpenicillin	92
Apo-Nadolol		Aspirin		Benzatropine mesylate	119
Apo-Nicotinic Acid	53	Blood	39	Benzbromaron AL 100	116
Apo-Ondansetron	130	Nervous	120	Benzbromarone	116
Apo-Oxybutynin		Asthalin	223	Benzoin	236
Apo-Paroxetine		Atazanavir sulphate	107	Benztrop	119
Apo-Perindopril	45	Atenolol	48	Benzydamine hydrochloride	30
Apo-Pindolol	49	Atenolol AFT	48	Benzylpenicillin sodium [Penicillin	
Apo-Pravastatin	53	ATGAM	180	G]	
Apo-Prazosin	45	Ativan	134	Beta Cream	
Apo-Prednisone	80	Atomoxetine	147	Beta Ointment	62
Apo-Primidone	127	Atorvastatin	53	Beta Scalp	68
Apo-Propranolol		Atropine sulphate		Beta-Adrenoceptor Agonists	223
Apo-Pyridoxine	31	Cardiovascular	47	Beta-Adrenoceptor Blockers	48
Apo-Ropinirole	118	Sensory		Betadine	65
Apo-Selegiline S29	118	Atropt	231	Betadine Skin Prep	65
Apo-Sumatriptan	129	Atrovent	223	Betaferon	144
Apo-Temozolomide	164	AU Synacthen		Betahistine dihydrochloride	130
Apo-Terazosin		Aubagio		Betaine	27
Apo-Timol	50	Augmentin	92	Betaloc CR	49
Apomorphine hydrochloride	118	Aurorix	124	Betamethasone dipropionate	
Aprepitant		AutoSoft 30	20	Betamethasone dipropionate with	
Apresoline		AutoSoft 90		calcipotriol	
Aptamil Gold+ Pepti Junior		Avelox		Betamethasone sodium phosphat	
Aqueous cream		Avonex		with betamethasone acetate	
Aratac		Avonex Pen		Betamethasone valerate	.62, 68
Aripiprazole	131	Azacitidine	157	Betamethasone valerate with	
Aripiprazole Sandoz	131	Azacitidine Dr Reddy's	157	clioquinol	
Aristocort	63	Azamun	174	Betamethasone valerate with sodi	
Arrow - Clopid		Azathioprine		fusidate [fusidic acid]	
Arrow-Amitriptyline	123	Azithromycin		Betaxolol	
Arrow-Bendrofluazide	52	Azol		Betnovate	
Arrow-Brimonidine		Azopt		Betnovate-C	
Arrow-Calcium		AZT	106–107	Betoptic	
Arrow-Diazepam	134	- B -		Betoptic S	230

Bezafibrate	52	Disoprox	233	CareSens N POP	
Bezalip	52	Buccastem	131	CareSens N Premier	
Bezalip Retard	52	Budesonide		CareSens PRO	13
Bicalutamide	172	Alimentary	6	Carmellose sodium with gelatin a	nd
Bicillin LA		Respiratory		pectin	
BiCNU	156	Budesonide with eformoterol.	222	Carmustine	
Bicnu Heritage	156	Bumetanide	51	Carvedilol	49
Bile and Liver Therapy	9	Buprenorphine Naloxone BNI	И 1 <u>51</u>	Carvedilol Sandoz	49
Biltricide		Buprenorphine with naloxone	151	Catapres	51
Bimatoprost	231	Bupropion hydrochloride		Cathejell	
Bimatoprost Multichem	231	Burinex	51	CeeNU	156
Binarex	172	Buscopan	8	Cefaclor monohydrate	
Binocrit	36	Buspirone hydrochloride	134	Cefalexin	89
Biodone	121	Busulfan	156	Cefalexin Sandoz	89
Biodone Extra Forte	121	- C -		Cefazolin	89
Biodone Forte	121	Cabergoline		Ceftriaxone	
Bisacodyl	26	Cafergot		Ceftriaxone-AFT	
Bisoprolol fumarate	49	Cafergot S29	129	Cefuroxime axetil	90
BK Lotion	64	Caffeine citrate	227	Celebrex	110
Bleomycin sulphate	161	Calamine	61	Celecoxib	
Blood Colony-stimulating		Calcipotriol		Celecoxib Pfizer	
Factors	42	Calcitonin		Celestone Chronodose	
Blood glucose diagnostic test		Calcitriol	32	Celiprolol	49
meter	12	Calcitriol-AFT	32	Cellcept	174
Blood glucose diagnostic test		Calcium carbonate	6, 33	Celol	49
strip	13	Calcium Channel Blockers	50	Centrally-Acting Agents	51
Blood glucose test strips (visua	ally	Calcium Disodium Versenate	234	Cephalexin ABM	89
impaired)		Calcium folinate	158	Cetirizine hydrochloride	
Blood Ketone Diagnostic Test		Calcium Folinate Ebewe	158	Cetomacrogol	64
Strip	11	Calcium Folinate Sandoz	158	Cetomacrogol with glycerol	64
Bonjela	30	Calcium gluconate	33	Cetuximab	191
Boostrix	257	Calcium Homeostasis	78	Charcoal	233
Bortezomib	161	Calcium polystyrene sulphona	ate43	Chemotherapeutic Agents	155
Bosentan	55	Calcium Resonium	43	Chickenpox vaccine	266
Bosentan Dr Reddy's	55	Calcium Sandoz	33	Chlorafast	228
Bosvate	49	Calogen	240	Chlorambucil	156
Bplex		Candesartan cilexetil	46	Chloramphenicol	228
Breo Ellipta	222	Candestar	46	Chlorhexidine gluconate	
Brevinor 1/21	73	Canesten	60	Alimentary	30
Brevinor 1/28	73	Capecitabine		Dermatological	63
Brevinor 21	73	Capoten	45	Chloroform	236
Bricanyl Turbuhaler	223	Capsaicin		Chlorothiazide	52
Brilinta	39	Musculoskeletal	111	Chlorpheniramine maleate	221
Brimonidine tartrate	231	Nervous		Chlorpromazine hydrochloride	131
Brimonidine tartrate with timolo	ol	Captopril		Chlorsig	228
maleate	231	Carafate	9	Chlortalidone [Chlorthalidone]	52
Brinov		Carbaccord		Chlorthalidone	
Brinzolamide	230	Carbamazepine	126	Chlorvescent	44
Brolene	228	Carbimazole		Choice Load 375	
Bromocriptine mesylate		Carbomer		Choice TT380 Short	
Brufen SR		Carboplatin		Choice TT380 Standard	
BSF Logem		Carboplatin Ebewe		Choline salicylate with cetalkoniur	m
BSF Mylan Efavirenz Emtricitba		Carbosorb-X		chloride	30
Tenofov	233	Cardinol LA	49	Ciclopirox olamine	60
BSF Teva Atazanavir Sulphate	233	CareSens Dual	12	Ciclosporin	218
BSF Teva Emtricitabine Tenofo	oir	CareSens N	12–13	Cilazapril	

Cilazapril with	Colchicine 116	Dacarbazine APP16
hydrochlorothiazide46	Colecalciferol32	Dactinomycin [Actinomycin D]16
Cilicaine93	Colestid53	Daivobet6
Cilicaine VK93	Colestipol hydrochloride53	Daivonex6
Cinacalcet78	Colgout	Daktarin6
Cipflox94	Colifoam7	Dalacin C9
Ciprofloxacin	Colistin sulphomethate94	Dalteparin sodium4
Infection94	Colistin-Link94	Danazol8
Sensory228	Collodion flexible236	Dantrium11
Ciprofloxacin Teva228	Colloidal bismuth subcitrate9	Dantrium S2911
Circadin146	Colofac8	Dantrolene11
Cisplatin	Coloxyl25	Daonil 1
Cisplatin Ebewe	Combigan231	Dapa-Tabs5
Citalopram hydrobromide 125	Compound electrolytes44	Dapsone10
Cladribine 158	Compound electrolytes with glucose	Daraprim9
Clarithromycin	[Dextrose]44	Darunavir10
Alimentary8	Compound hydroxybenzoate236	Dasatinib16
Infection90	Concerta149	Daunorubicin16
Clexane	Condoms71	DBL Acetylcysteine23
Clindamycin94	Condyline	DBL Adrenaline5
Clindamycin ABM94	Contraceptives - Hormonal	DBL Aminophylline22
Clinicians Renal Vit32	Contraceptives - Non-hormonal70	DBL Bleomycin Sulfate16
Clobazam126	Copaxone140	DBL Carboplatin15
	Cordarone-X47	DBL Carbopiatin
Clobetasol propionate		
Closeramine 62	Corticosteroids and Related Agents	DBL Dacarbazine16 DBL Desferrioxamine Mesylate for Inj
Clonazal	for Systemic Use	BP23
Clomazol	Corticosteroids Topical	
Dermatological 60	Cosentyx209	DBL Docetaxel
Genito-Urinary	Cosmegen	DBL Ergometrine
Clomifene citrate	Coumadin	DBL Gemcitabine
Clomipramine hydrochloride123	Creon 1000023	DBL Gentamicin9
Clonazepam 125–126, 134	Creon 2500023	DBL Leucovorin Calcium15
Clonidine51	Cromal230	DBL Methotrexate Onco-Vial
Clonidine BNM51	Crotamiton61	DBL Morphine Sulphate12
Clonidine hydrochloride51	Crystaderm59	DBL Morphine Tartrate12
Clopidogrel39	Curam92	DBL Naloxone Hydrochloride23
Clopine 131	Cvite31	DBL Octreotide17
Clopixol	Cyclizine hydrochloride130	DBL Pethidine Hydrochloride12
Clotrimazole	Cyclizine lactate130	DBL Vincristine Sulfate16
Dermatological60	Cyclogyl231	De-Worm8
Genito-Urinary75	Cyclopentolate hydrochloride231	Decozol3
Clozapine131	Cyclophosphamide156	Deferasirox23
Clozaril131	Cyclorin100	Deferiprone23
Clustran	Cycloserine100	Denosumab11
Co-trimoxazole96	Cyklokapron38	Deolate9
Coal tar68	Cyproterone acetate80	Deoxycoformycin16
Coal tar with allantoin, menthol,	Cyproterone acetate with	Depo-Medrol7
phenol and sulphur68	ethinyloestradiol75	Depo-Provera7
Coal tar with salicylic acid and	Cystadane27	Depo-Testosterone8
sulphur68	Cytarabine 158	Deprim9
Coco-Scalp68	Cytotec8	DermAssist6
Codeine phosphate	Cytoxan156	Dermol62, 6
Extemporaneous236	- D -	Desferrioxamine mesilate23
Nervous121	D-Penamine111	Desmopressin acetate8
Cogentin	Dabigatran41	Desmopressin-PH&T8
Colaspase [L-asparaginase]161	Dacarbazine161	Detection of Substances in
· · · · · · · · ·		

Urine	Disinfecting and Cleansing	Efavirenz with emtricitabine and	
Dexamethasone	Agents	tenofovir disoproxil	10
Hormone79	Disopyramide phosphate47	Effient	
Sensory229	Disulfiram152	Eformoterol fumarate	
Dexamethasone phosphate79	Diuretics51	Eformoterol fumarate dihydrate	
Dexamethasone with framycetin and	Diurin 4051	Eftrenonacog alfa [Recombinant	
gramicidin228	Docetaxel	factor IX]	3(
Dexamethasone with neomycin	Docetaxel Accord162	Efudix	
sulphate and polymyxin B	Docetaxel Sandoz162	Egopsoryl TA	
sulphate229	Docusate sodium25	Elaprase	
Dexamfetamine sulfate147	Docusate sodium with	Elecare	
Dexmethsone79	sennosides25	Elecare LCP	
Dextrochlorpheniramine	Dolutegravir107	Electral	
maleate221	Domperidone	Elelyso	
Dextrose42, 44	Donepezil hydrochloride150	Elemental 028 Extra	
DHC Continus121	Donepezil-Rex150	Elocon	
Diabetes9	Dopress	Elocon Alcohol Free	
Diabetes Management11	Dornase alfa226	Eltrombopag	
Diacomit	Dortimopt230	Eltroxin	
Diagnostic Agents	Dorzolamide hydrochloride230	EMB Fatol	
Diamide Relief6	Dorzolamide with timolol230	Emend Tri-Pack	
Diamox230	Dostinex87	EMLA	
Diasip241	Dosulepin [Dothiepin]	Emtricitabine	
Diason RTH241	hydrochloride124	Emtricitabilite with tenofovir	10
Diazepam	Dothiepin124	disoproxil	10
Diazoxide9	Doxazosin	Emtriva	
Dibenzyline45	Doxepin hydrochloride124	Emulsifying ointment	
Diclofenac Sandoz110	Doxine	Enalapril maleate	
Diclofenac sodium	Doxorubicin Ebewe162	Enbrel	
Musculoskeletal110	Doxorubicin hydrochloride	Endocrine Therapy	17
Sensory229	Doxy-5093	Endoxan	
Differin	Doxycycline93	Enerlyte	
Difflam30	DP Lotion	Engerix-B	
Diflucan	DP Lotn HC	Enlafax XR	
Diflucan S2996	DP-Allopurinol	Enoxaparin sodium	
Diflucortolone valerate	Dr Reddy's Omeprazole9	Ensure	
Digestives Including Enzymes23	Drugs Affecting Bone	Ensure Plus	24
	Metabolism111	Ensure Plus HN	
Digoxin	Dual blood glucose and blood ketone	Ensure Plus RTH	
Dilantin		Entacapone	
Dilantin Infatab	diagnostic test meter	Entapone	449
Diltiazem hydrochloride	Duolin223	Entecavir	10
Dilzem50	Duolin HFA	Entecavir Sandoz	10
	Durex Confidence	Entocort CIR	
Dimethicone	Durex Extra Safe71	Entresto 24/26	
Dimethyl fumarate	Duride54	Entresto 49/51	
Dipentum	- E -	Entresto 97/103	
Diphtheria, tetanus and pertussis	e-chamber La Grande227		
vaccine		Epilim Crushabla	
Diphtheria, tetanus, pertussis and	e-chamber Mask	Epilim Crushable	
polio vaccine	e-chamber Turbo	Epilim IV	
Diphtheria, tetanus, pertussis, polio,	E-Mycin	Epilim S/F Liquid	
hepatitis B and haemophilus	Ear Preparations	Epilim Syrup	12
influenzae type B vaccine	Ear/Eye Preparations	Epirubicin Ebewe	
Diprosone 62	Easiphen Liquid	Epirubicin hydrochloride	
Diprosone OV	Econazole nitrate 60	Eplerenone	
Dipyridamole39	Efavirenz106	Epoetin alfa	3

Epoprostenol	58	Febuxostat	117	fluocortolone pivalate and	
Eptacog alfa [Recombinant factor		Feed Thickener Karicare		cinchocaine	
VIIa]	37	Aptamil	251	Fluorometholone	23
ERA	91	FEIBA NF	37	Fluorouracil	15
Erbitux	191	Felo 10 ER	50	Fluorouracil Ebewe	15
Ergometrine maleate	75	Felo 5 ER		Fluorouracil sodium	6
Ergotamine tartrate with		Felodipine	50	Fluox	
caffeine	. 129	Femme-Tab ED	73	Fluoxetine hydrochloride	12
Erlotinib	166	Fentanyl	121	Flupenthixol decanoate	13
Erythrocin IV	91	Fentanyl Sandoz	121	FluQuadri	26
Erythromycin (as lactobionate)	91	Ferinject	33	Flutamide	17
Erythromycin ethyl succinate	91	Ferodan		Flutamide Mylan	17
Erythromycin stearate	91	Ferric carboxymaltose	33	Flutamin	17
Esbriet		Ferriprox	234	Fluticasone	22
Escitalopram	125	Ferro-F-Tabs	34	Fluticasone furoate with	
Escitalopram-Apotex	125	Ferro-tab	34	vilanterol	22
Eskazole		Ferrograd	34	Fluticasone propionate	22
Estradot	81	Ferrosig		Fluticasone with salmeterol	22
Estradot 50 mcg	81	Ferrous fumarate		FML	
Estrofem	81	Ferrous fumarate with folic acid	34	Foban	6
Etanercept	174	Ferrous sulfate	34	Folic acid	3
Ethambutol hydrochloride	100	Ferrous sulphate	34	Food Thickeners	25
Ethics Aspirin		Ferrum H		Foods And Supplements For Inbo	rn
Ethics Aspirin EC		Fexofenadine hydrochloride	221	Errors Of Metabolism	
Ethics Enalapril		Fibro-vein		Foradil	22
Ethics Lisinopril		Filgrastim		Forteo	11
Ethinyloestradiol		Finasteride	76	Fortini	24
Ethinyloestradiol with		Fingolimod	136	Fortini Multi Fibre	24
desogestrel	73	Firazyr	220	Fortisip	24
Ethinyloestradiol with		Flagyl	99	Fortisip Multi Fibre	25
levonorgestrel	73	Flagyl-S		Fosamax	
Ethinyloestradiol with		Flamazine		Fosamax Plus	
norethisterone	73	Flecainide acetate	48	Fragmin	4
Ethosuximide	126	Flecainide BNM		Framycetin sulphate	22
Etopophos		Flecainide Controlled Release		Fresenius Kabi	
Etoposide		Teva	48	Frisium	12
Etoposide phosphate		Fleet Phosphate Enema	25	Frumil	5
Etravirine		Flixonase Hayfever & Allergy		Frusemide	5
Eumovate		Flixotide		Frusemide-Claris	
Everet	127	Flixotide Accuhaler	222	Fucicort	6
Everolimus	218	Floair	222	Fucidin	9
Evista	112	Florinef	79	Fucithalmic	22
Exelon	150	Fluanxol	133	Fungilin	3
Exemestane	173	Fluarix Tetra	261	Furosemide [Frusemide]	
Exjade	233	Flucil	93	fusidic acid	
Extemporaneously Compounded		Flucloxacillin	93	Dermatological	60, 6
Preparations and		Flucloxin	93	Infection	
Galenicals	. 236	Flucon	230	Sensory	22
Eye Preparations		Fluconazole	96	´ - G -	
Eylea		Fludara Oral		Gabapentin	12
Ezetimibe		Fludarabine Ebewe		Gacet	
Ezetimibe Sandoz		Fludarabine phosphate		Galsulfase	
Ezetimibe with simvastatin		Fludrocortisone acetate	79	Galvumet	
- F -		Fluids and Electrolytes		Galvus	
Factor eight inhibitor bypassing		Flumetasone pivalate		Gardasil 9	
fraction	37	Fluocortolone caproate with		Gastrodenol	
		•			

Cavinaan Daubla Strangth 6	Louriy 050	Hydrocortisone butyrate62, 68
Gaviscon Double Strength		Hydrocortisone with cinchocaine
Gazyva198		Hydrocortisone with miconazole63
Gefitinib		
Gemcitabine Ebewe		Hydrocortisone with natamycin and
		neomycin
Gemcitabine hydrochloride		Hydrogen peroxide Alimentary31
Gemfibrozil		
General 158		Dermatological
Genoptic		
Gentamicin sulphate	healthE Urea Cream	Hydroxychloroquine111
Infection94		Hydroxyurea162
Sensory228		Hygroton
Gilenya		Hylo-Fresh
Ginet	· · · · · · · · · · · · · · · · · · ·	Hymenoptera220
Glatiramer acetate	•	Hyoscine butylbromide
Glecaprevir with pibrentasvir		Hyoscine hydrobromide
Glibenclamide11		Hypam147
Gliclazide11		Hyperuricaemia and Antigout
Glipizide11		Hypromellose231
Glivec		Hypromellose with dextran232
Glizide11		-1-
Glucagen Hypokit		Ibiamox92
Glucagon hydrochloride		lbuprofen110
Glucerna Select		Icatibant220
Glucerna Select RTH241		Idarubicin hydrochloride162
Glucobay11		Idursulfase27
Glucose [Dextrose]42		Ifosfamide156
Gluten Free Foods251	,	Ikorel55
Glycerin with sodium saccharin236		lloprost58
Glycerin with sucrose236	•	Imatinib mesilate167
Glycerol	HPV260	Imatinib-AFT167
Alimentary25		Imipramine hydrochloride124
Extemporaneous236	•	Imiquimod69
Glyceryl trinitrate	Humalog Mix 5010	Immune Modulators107
Alimentary		Immunosuppressants174
Cardiovascular54		Imuran174
Glycopyrronium223		Incruse Ellipta224
Glycopyrronium bromide		Indacaterol222
Glycopyrronium with	Humira180	Indapamide52
indacaterol224		Infanrix IPV258
Glytrin54		Infanrix-hexa258
Gold Knight71–72		Infant Formulae254
Goserelin86		Infatrini255
Gutron48	•	Infliximab191
Gynaecological Anti-infectives75		Influenza vaccine261
-H-	Hydralazine55	Influvac Tetra261
Habitrol153	, ,	Inhaled Corticosteroids221
Haemophilus influenzae type B	Hydrea162	Inhaled Long-acting
vaccine258	,	Beta-adrenoceptor Agonists 222
Haldol133	•	Inspra51
Haldol Concentrate133		Insulin aspart10
Haldol Decanoas133	Hydrocortisone acetate7	Insulin aspart with insulin aspart
Haloperidol132		protamine10
Haloperidol decanoate133		Insulin glargine10
Hamilton Sunscreen69		Insulin glulisine10
Harvoni103	and lanolin 62	Insulin isophane10

Insulin isophane with insulin	Isuprel	55	Lanzol Relief	
neutral10	Itch-Soothe	61	Lapatinib ditosylate	16
Insulin lispro10	Itraconazole	97	Largactil	13
Insulin lispro with insulin lispro	Itrazole	97	Laronidase	2
protamine10	Ivermectin	65	Lasix	5
Insulin neutral10	- J -		Latanoprost	23
Insulin pen needles13	Jadelle	74	Lax-Suppositories	2
Insulin pump14	Jakavi	169	Lax-Tab	
Insulin pump cartridge19	Jaydess	82	Laxatives	2
Insulin pump infusion set (steel	Jevity HiCal RTH	248	Laxsol	2
cannula)19	Jevity RTH	248	Ledipasvir with sofosbuvir	10
Insulin pump infusion set (steel	Juno Pemetrexed	159	Leflunomide	11
cannula, straight insertion) 20	- K -		Lenalidomide	16
Insulin pump infusion set (teflon	Kaletra	107	Letrole	
cannula, angle insertion with	Keflor	89	Letrozole	17
insertion device)20	Kemadrin	119	Leukeran FC	15
Insulin pump infusion set (teflon	Kenacomb	228	Leukotriene Receptor	
cannula, angle insertion)21	Kenacort-A 10	80	Antagonists	22
Insulin pump infusion set (teflon	Kenacort-A 40	80	Leunase	
cannula, straight insertion with	Kenalog in Orabase	30	Leuprorelin	8
insertion device)22	Ketocal 3:1		Leustatin	15
Insulin pump infusion set (teflon	KetoCal 4:1	256	Levetiracetam	12
cannula, straight insertion) 23	Ketoconazole		Levetiracetam-AFT	12
Insulin pump reservoir23	Dermatological	68	Levlen ED	7
Insulin syringes, disposable with	Infection		Levocabastine	23
attached needle14	Ketogenic Diet		Levodopa with benserazide	11
Intal Forte CFC Free226	Ketoprofen		Levodopa with carbidopa	11
Intelence106	KetoSens		Levomepromazine	
Interferon alfa-2a108	Keytruda	216	hydrochloride	13
Interferon beta-1-alpha142	Kindergen		Levomepromazine maleate	
Interferon beta-1-beta144	Kivexa		Levonorgestrel	
Intra-uterine device72	Klacid		Genito-Urinary	7
Invega Sustenna133	Kliogest		Hormone	
IPOL	Kliovance		Levothyroxine	
Ipratropium bromide223, 227	Kogenate FS		Lidocaine [Lignocaine]11	
Iressa	Konakion MM		Lidocaine [Lignocaine]	
Irinotecan Accord159	Konsyl-D		hydrochloride	12
Irinotecan Actavis 100	Kuvan		Lidocaine [Lignocaine] with	
Irinotecan hydrochloride	-L-		chlorhexidine	12
Irinotecan-Rex	L-asparaginase	161	Lidocaine [Lignocaine] with	
Iron polymaltose34	Labetalol		prilocaine	12
Isentress	Lacosamide		Lidocaine-Claris	
Isentress HD107	Lactulose		Lignocaine11	
Ismo 2054	Laevolac		Lioresal Intrathecal	
Ismo 40 Retard54	Lamictal		Lipazil	
Isoniazid100	Lamivudine10		Lipid-Modifying Agents	
Isoniazid with rifampicin100	Lamivudine Alphapharm		Liquigen	24
Isoprenaline [Isoproterenol]55	Lamotrigine		Lisinopril	
Isoproterenol55	Lamprene		Lithicarb FC	
Isoptin51	Lanoxin		Lithium carbonate	
Isoptin SR51	Lanoxin PG		Livostin	
Isopto Carpine231	Lanoxin S29		LMX4	
Isosorbide mononitrate54	Lansoprazole		Locacorten-Viaform ED's	
Isosource Standard248	Lantus		Local preparations for Anal and	
Isotretinoin59	Lantus SoloStar		Rectal Disorders	
Ispaghula (psyllium) husk25	Lanvis		Locasol	

Locoid	62, 68	MCT oil (Nutricia)	240	Metoclopramide hydrochloride	130
Locoid Crelo	62	Measles, mumps and rubella		Metolazone	5
Locoid Lipocream	62	vaccine	263	Metopirone	8
Locorten-Vioform	228	Mebendazole	89	Metoprolol succinate	49
Lodi	47	Mebeverine hydrochloride	8	Metoprolol tartrate	
Lodoxamide	230	Medrol	79	Metronidazole	9
Logem	127	Medroxyprogesterone acetate		Metroprolol IV Mylan	49
Lomide	230	Genito-Urinary	74	Metyrapone	
Lomustine	156	Hormone	81-82	Mexiletine hydrochloride	
Loniten	55	Mefenamic acid	110	Mexiletine Hydrochloride USP	4
Loperamide hydrochloride	6	Megestrol acetate	172	Miacalcic	
Lopinavir with ritonavir	107	Melatonin	146	Micolette	20
Loprofin		Melphalan	156	Miconazole	3
Loprofin Mix	253	Menactra		Miconazole nitrate	
Lorafix		Meningococcal (groups A, C, Y a	and	Dermatological	6
Loratadine	221	W-135) conjugate vaccine	263	Genito-Urinary	7
Lorazepam	134	Meningococcal C conjugate		Micreme	
Lorfast		vaccine	263	Micreme H	6
Lorstat	53	Menthol	61	Microgynon 20 ED	7
Losartan Actavis	46	Mercaptopurine	159	Microgynon 30	
Losartan potassium	46	Mercilon 28		Microgynon 50 ED	
Losartan potassium with		Mesalazine		Microlut	
hydrochlorothiazide	46	Mesna		Midazolam	
Lovir		Mestinon	110	Midazolam-Claris	140
Loxamine	125	Metabolic Disorder Agents	26	Midodrine	
Lucrin Depot 1-month	87	Meterol	222	Minerals	3
Lucrin Depot 3-month		Metformin hydrochloride		Mini-Wright AFS Low Range	
Ludiomil		Methadone hydrochloride		Mini-Wright Standard	
Lyderm	67	Extemporaneous	236	Minidiab	
- M -		Nervous		MiniMed 640G	
m-Eslon	122	Methatabs	121	Minims Pilocarpine	23
Mabthera	200	Methopt		Minims Prednisolone	
Macrogol 3350 with potassiun	n	Methotrexate		Minirin	8
chloride, sodium bicarbona		Methotrexate Ebewe		Mino-tabs	
sodium chloride		Methotrexate Sandoz		Minocycline hydrochloride	
Macrogol 400 and propylene		Methyl hydroxybenzoate		Minomycin	
glycol	232	Methylcellulose		Minor Śkin Infections	6
Madopar 125		Methylcellulose with glycerin and		Minoxidil	
Madopar 250		sodium saccharin		Mirena	
Madopar 62.5		Methylcellulose with glycerin and		Mirtazapine	
Madopar HBS		sucrose		Misoprostol	
Madopar Rapid		Methyldopa		Mitomycin C	
Magnesium hydroxide		Methyldopa Mylan		Mitozantrone	
Alimentary	34	Methyldopa Mylan S29		Mitozantrone Ebewe	
Extemporaneous		Methylnaltrexone bromide		Mixtard 30	1
Magnesium sulphate		Methylphenidate ER - Teva		MMR II	
Mantoux		Methylphenidate hydrochloride		Moclobemide	
Maprotiline hydrochloride		Methylphenidate hydrochloride		Modafinil	
Marevan		extended-release	149	Modavigil	
Marine Blue Lotion SPF 50+ .		Methylprednisolone		Moduretic	
Marvelon 28		Methylprednisolone (as sodium		Molaxole	
Mask for spacer device		succinate)	79	Moments	
Mast Cell Stabilisers		Methylprednisolone aceponate		Mometasone furoate	
Maviret		Methylprednisolone acetate		Monogen	
Maxidex		Methylxanthines		Montelukast	
Maxitrol		Metoclopramide Actavis 10		Montelukast Mylan	22

Moroctocog alfa [Recombinant fac	tor	Neisvac-C	263	Nortriptyline hydrochloride	124
VIII]	38	Neo-B12	31	Norvir	
Morphine hydrochloride		Neo-Mercazole	82	NovaSource Renal	244
Morphine sulphate		Neocate Gold	254	Novatretin	6
Morphine tartrate		Neocate Junior Unflavoured	254	NovoMix 30 FlexPen	10
Motetis		Neocate Junior Vanilla		NovoRapid	
Mouth and Throat		Neocate SYNEO		NovoRapid FlexPen	
Movapo		Neoral		NovoRapid Penfill	
Moxifloxacin		Neostigmine metilsulfate		NovoSeven RT	
MSUD Maxamum		Nepro HP (strawberry)		Noxafil	Q.
Mucilaginous laxatives with	202	Nepro HP (vanilla)		Nozinan	
stimulants	25	Nepro HP RTH		Nuelin	
Mucolytics		Nerisone		Nuelin-SR	
Mucosoothe		Neulactil		Nutilis	
Multiple Sclerosis Treatments		Neulastim		Nutrient Modules	
				Nutrini Energy Multi Fibre	
Multivitamin renal		NeuroTabs			
Multivitamins		Nevirapine		Nutrini Energy RTH	
Mupirocin		Nevirapine Alphapharm		Nutrini Low Energy Multi Fibre	
Muscle Relaxants		Nicorandil		Nutrini RTH	24
Mvite		Nicotine		Nutrison 800 Complete Multi	
Myambutol		Nicotinic acid		Fibre	
Mycobutin		Nifedipine		Nutrison Concentrated	
MycoNail		Nifuran	109	Nutrison Energy	
Mycophenolate mofetil	174	Nilotinib	168	Nutrison Energy Multi Fibre	
Mycostatin	61	Nilstat		Nutrison Multi Fibre	248
Mydriacyl	231	Alimentary	31	Nutrison Standard RTH	248
Mylan Atenolol	48	Genito-Urinary	75	Nyefax Retard	50
Mylan Clomiphen	88	Infection	97	Nystatin	
Myleran	156	Nintedanib	224	Alimentary	3
Myocrisin	111	Nipent	164	Dermatological	6
Myometrial and Vaginal Hormone		Nitrados	146	Genito-Urinary	
Preparations	75	Nitrates		Infection	
Myozyme		Nitrazepam	146	NZB Low Gluten Bread Mix	25
Mysoline S29		Nitroderm TTS		-0-	
- N -		Nitrofurantoin		O/W Fatty Emulsion Cream	64
Nadolol	49	Nitrolingual Pump Spray		Obinutuzumab	
Naglazyme		Nivestim		Octocog alfa [Recombinant factor	
Nalcrom		Nivolumab		VIII] (Advate)	38
Naloxone hydrochloride		Nizoral		Octocog alfa [Recombinant factor	
Naltraccord		Nodia		VIII] (Kogenate FS)	38
Naltrexone hydrochloride		Noflam 250		Octreotide	
Naphazoline hydrochloride		Noflam 500		Octreotide LAR (somatostatin	174
Naphcon Forte		Non-Steroidal Anti-Inflammatory	110	analogue)	17
		_	110	Oestradiol	
Naprosyn SR 1000		Drugs Nonacog gamma, [Recombinant	110	Oestradiol valerate	
Naprosyn SR 750		Factor IX]	20	Oestradiol with norethisterone	
Naproxen			30		0
Nardil		Norethisterone	7.4	Oestriol	
Nardil S29		Genito-Urinary		Genito-Urinary	
Nasal Preparations		Hormone		Hormone	
Natalizumab		Norflex		Oestrogens	
Natulan		Norfloxacin		Ofev	224
Nausafix		Noriday 28		Oil in water emulsion	
Nausicalm		Norimin		Olanzapine132	
Navelbine		Normacol Plus		Olbetam	
Nedocromil		Normison		Olopatadine	
Nefopam hydrochloride	120	Norpress	124	Olsalazine	

Omalizumab	. 199	Oxytocin with ergometrine	Paraffin liquid with wool fat	23
Omeprazole	9	maleate	Paraldehyde	120
Omeprazole actavis 10	9	Ozurdex229	Parasidose	
Omeprazole actavis 20	9	- P -	Parasiticidal Preparations	6
Omeprazole actavis 40	9	Pacifen117	Parnate	
Omnitrope		Paclitaxel163	Parnate S29	124
Onbrez Breezhaler	222	Paclitaxel Actavis163	Paromomycin	9
Oncaspar		Paclitaxel Ebewe163	Paroxetine	12
OncoTICE	.180	Paediatric Seravit32	Paser	100
Ondansetron	130	Paliperidone133	Patanol	23
Ondansetron ODT-DRLA	.130	Pamidronate disodium112	Paxam	13
Ondansetron ODT-ORLA		Pamisol112	Pazopanib	16
One-Alpha	32	Pan-Penicillin G Sodium92	Peak flow meter	22
Onrex		Pancreatic enzyme23	Pedialyte - Bubblegum	4
Opdivo	215	Pantoprazole9	Pediasure	
Ora-Blend		Panzop Relief9	Pediasure RTH	24
Ora-Blend SF	237	Panzytrat23	Pegaspargase	16
Ora-Plus	236	Papaverine hydrochloride55	Pegasys	
Ora-Sweet	236	Para-amino salicylic acid100	Pegfilgrastim	
Ora-Sweet SF	236	Paracare121	Pegylated interferon alfa-2a	
Orabase	30	Paracare Double Strength121	Pembrolizumab	
Oral Supplements/Complete Diet		Paracetamol121	Pemetrexed	159
(Nasogastric/Gastrostomy Tube		Paracetamol + Codeine	Penicillamine	11
Feed)	241	(Relieve) 123	Penicillin G	
Oratane		Paracetamol Pharmacare121	PenMix 30	
Ordine	122	Paracetamol with codeine123	PenMix 40	10
Orgran		Paradigm 1.8 Reservoir23	PenMix 50	
Orion Temozolomide		Paradigm 3.0 Reservoir23	Pentasa	
Ornidazole		Paradigm Mio MMT-92122	Pentostatin [Deoxycoformycin]	16
Orphenadrine citrate		Paradigm Mio MMT-92322	Pentoxifylline [Oxpentifylline]	
Ortho-tolidine		Paradigm Mio MMT-92522	Peptamen Junior	
Oruvail SR		Paradigm Mio MMT-94122	Peptisoothe	
Osmolite RTH		Paradigm Mio MMT-94322	Peptisorb	
Other Endocrine Agents		Paradigm Mio MMT-94522	Perhexiline maleate	50
Other Oestrogen Preparations		Paradigm Mio MMT-96522	Pericyazine	
Other Progestogen		Paradigm Mio MMT-97522	Perindopril	
Preparations	82	Paradigm Quick-Set MMT-38623	Perjeta	
Other Skin Preparations		Paradigm Quick-Set MMT-38723	Permethrin	
Ovestin		Paradigm Quick-Set MMT-39623	Perrigo	
Genito-Urinary	75	Paradigm Quick-Set MMT-39723	Pertuzumab	
Hormone		Paradigm Quick-Set MMT-39823	Peteha	
Ox-Pam		Paradigm Quick-Set MMT-39923	Pethidine hydrochloride	
Oxaliccord		Paradigm Silhouette MMT-36821	Pevaryl	
Oxaliplatin		Paradigm Silhouette MMT-37721	Pexsig	
Oxaliplatin Accord		Paradigm Silhouette MMT-37821	Pfizer Exemestane	
Oxaliplatin Actavis 100		Paradigm Silhouette MMT-38121	Pharmacy Health Sorbolene with	
Oxaliplatin Ebewe		Paradigm Silhouette MMT-38221	Glycerin	6
Oxazepam		Paradigm Silhouette MMT-38321	Pharmacy Services	
Oxis Turbuhaler		Paradigm Silhouette MMT-38421	Pheburane	2
Oxpentifylline		Paradigm Sure-T MMT-86419	Phenasen	
Oxybutynin		Paradigm Sure-T MMT-86619	Phenelzine sulphate	
Oxycodone hydrochloride		Paradigm Sure-T MMT-87419	Phenobarbitone	12
Oxycodone Sandoz	123	Paradigm Sure-T MMT-87619	Phenobarbitone sodium	
OxyNorm		Paradigm Sure-T MMT-88419	Extemporaneous	23
Oxytocin		Paradigm Sure-T MMT-88619	Nervous	
Oxytocin BNM		Paraffin	Phenothrin	6

Phenoxybenzamine	Praziquantel	89	Puria	3
hydrochloride4	5 Prazosin	45	Pyrazinamide	10
Phenoxymethylpenicillin (Penicillin	Pred Forte	230	Pyridostigmine bromide	11
V)9	3 Prednisolone	80	Pyridoxine hydrochloride	
Phenytoin sodium126–12		230	Pyrimethamine	
Phlexy 1025			Pytazen SR	3
Phosphate Phebra4	4 phosphate	230	_ Q -	
Phosphorus4	4 Prednisolone-AFT		Q 300	9
Phytomenadione3		80	Quetapel	
Pilocarpine hydrochloride23		127	Quetiapine	
Pimafucort6			Quick-Set MMT-390	
Pindolol4	•		Quick-Set MMT-391	2
Pine tar with trolamine laurilsulfate	Premarin		Quick-Set MMT-392	
and fluorescein6	8 Presolol	49	Quick-Set MMT-393	
Pinetarsol6	8 Prevenar 13	264	Quinapril	
Pioglitazone1		107	Quinapril with	
Pirfenidone22		132	hydrochlorothiazide	4
Pizotifen12		99	Quinine sulphate	
PKU Anamix Infant25			Qvar	
PKU Anamix Junior25			-R-	
PKU Anamix Junior Chocolate25			RA-Morph	12
PKU Anamix Junior LQ25		263	Raloxifene hydrochloride	
PKU Anamix Junior Vanilla25			Raltegravir potassium	
PKU Lophlex LQ 1025			Ramipex	
PKU Lophlex LQ 2025			Ranbaxy-Cefaclor	8
PKU Lophlex Powder25	3 Procarbazine hydrochloride		Ranitidine	
PKU Lophlex Sensation 2025	3 Prochlorperazine		Ranitidine Relief	
Plaquenil11	1 Proctofoam		Rapamune	
Plendil ER5		7	Reandron 1000	
Pneumococcal (PCV10) conjugate	Procyclidine hydrochloride		Recombinant factor IX	
vaccine			Recombinant factor VIIa	
Pneumococcal (PCV13) conjugate	Progesterone		Recombinant factor VIII	
vaccine26			Rectogesic	
Pneumococcal (PPV23)	Proglycem		Redipred	8
polysaccharide vaccine 26			Relieve	
Pneumovax 2326			Relistor	
Podophyllotoxin6			Remicade	
Polaramine22			Renilon 7.5	
Poliomyelitis vaccine26	. ,		Resonium-A	
Poloxamer2			Resource Beneprotein	
Poly-Gel23			Resource Diabetic	
Poly-Tears23			Respigen	
Poly-Visc23		10	Respiratory Devices	
Polycal23			Respiratory Stimulants	
Polyvinyl alcohol23			Retinol palmitate	
Ponstan11			ReTrieve	
Posaconazole 9		81	Retrovir	
Postinor-17			Revlimid	
Potassium chloride42, 4			Revolade	
Potassium citrate7			Rexacrom	
Potassium iodate3		67	RexAir	
Povidone iodine6			Ribomustin	
Pradaxa			Ricit	
Pramipexole hydrochloride11			Rifabutin	
Prasugrel3			Rifadin	
Pravastatin5			Rifampicin	

Rifaximin	9	Secukinumab	209	Sodium Fusidate [fusidic acid]	
Rifinah		Sedatives and Hypnotics		Dermatological	60
Rilutek		Seebri Breezhaler		Infection	
Riluzole		Selegiline hydrochloride		Sensory	
Riodine		Senna		Sodium hyaluronate [Hyaluronic	
Risedronate Sandoz		Senokot		acid]	232
Risedronate sodium		Sensipar		Sodium phenylbutyrate	
Risperdal Consta		SensoCard		Sodium polystyrene sulphonate	
Risperidone		Serenace		Sodium tetradecyl sulphate	
Risperon	132	Seretide		Sodium valproate	
Ritalin		Seretide Accuhaler		Sofradex	
Ritalin LA		Serevent		Soframycin	
Ritalin SR		Serevent Accuhaler		Solian	
Ritonavir		Sertraline		Solifenacin Mylan	
Rituximab		Setrona		Solifenacin succinate	
Rivaroxaban		Sevredol		Solu-Cortef	
Rivastigmine		Sex Hormones Non		Solu-Medrol	
Rivotril		Contraceptive	80	Solu-Medrol-Act-O-Vial	
RIXUBIS		Shield 49		Somatropin (Omnitrope)	
Rizamelt		Shield Blue		Sotalol	
Rizatriptan		Shield XL		Spacer device	
Roferon-A		shingles vaccine		Span-K	
Rolin		SII-Onco-BCG		Spiolto Respimat	
Ropin		Sildenafil		Spiractin	52
Ropinirole hydrochloride		Silhouette MMT-371		Spiriva	
Rotarix		Silhouette MMT-373		Spiriva Respimat	
Rotavirus oral vaccine		Siltuximab		Spironolactone	22-
Roxane	200	Simvastatin		Sporanox	07
Alimentary	6	Simvastatin Mylan		Sprycel	
Cardiovascular		Sinemet		Staphlex	100
Roxithromycin		Sinemet CR		Stemetil	
Rubifen		Sirolimus		SteroClear	
Rubifen SR		Siterone		Stesolid	
Rulide D		Slow-Lopresor		Stimulants/ADHD Treatments	
		•		Stiripentol	
Rurioctocog alfa pegol [Reco		Smith BioMed Rapid Pregnancy Test		Stocrin	
factor VIII]		Sodibic			
Ruxolitinib				Stomahesive Strattera	
Rythmodan		Sodium acid phosphate		Stromectol	
Rytmonorm	40	Sodium alginateSodium aurothiomalate		Suboxone	
Sabril	100	Sodium benzoate		Sucralfate	
Sacubitril with valsartan		Sodium bicarbonate	20	Sulfadiazine Silver	
SalAir			19 11	Sulfadiazine sodium	
Salazopyrin		Blood		Sulfasalazine	
Salazopyrin EN		Extemporaneous		Sulindac	
Salbutamol		Sodium calcium edetate Sodium chloride	234		
	223	Blood	49	Sulphur Sulprix	
Salbutamol with ipratropium	000			Sulpiix	10
bromide		Respiratory		Sumatriptan	120
Salicylic acid		Sodium citrate with sodium laury		Sunitinib	
Salmeterol		sulphoacetate		Sunscreens	
Sandomigran		Sodium citro-tartrate	/b	Sunscreens, proprietary	68
Sandostatin LAR		Sodium cromoglicate	7	Sure-T MMT-863 Sure-T MMT-865	
Sapropterin dihydrochloride.		Alimentary			
Scalp Preparations		Respiratory		Sure-T MMT-873	
Scopoderm TTS		Sensory		Sure-T MMT-875	
Sebizole	<u>68</u>	Sodium fluoride	33	Sure-T MMT-883	18

Sure-T MMT-885	19	Testosterone cipionate	80	Travopt	231
Sustagen Diabetic	241	Testosterone esters	80	Treatments for Dementia	
Sustagen Hospital Formula		Testosterone undecanoate	80	Treatments for Substance	
Active	249	Tetrabenazine	119	Dependence	151
Sustanon Ampoules	80	Tetrabromophenol	77	Trental 400	55
Sutent	170	Tetracosactrin		Tretinoin	
Sylvant		Tetracyclin Wolff	93	Dermatological	59
Symbicort Turbuhaler 100/6		Tetracycline		Oncology	
Symbicort Turbuhaler 200/6		Thalidomide		Trexate	
Symbicort Turbuhaler 400/12		Thalomid		Triamcinolone acetonide	
Symmetrel		Theophylline		Alimentary	30
Sympathomimetics		Thiamine hydrochloride		Dermatological	
Synacthen		THIO-TEPA		Hormone	
Synacthen Depot		Thioguanine		Triamcinolone acetonide with	
Synacthen S29		Thiotepa		gramicidin, neomycin and nys	statin
Synacthene Retard		Thymol glycerin		Dermatological	
Synflorix		Thyroid and Antithyroid Agents		Sensory	
Synthroid		Ticagrelor		Triazolam	
Syntometrine		Tilade		Trichozole	
Syrup (pharmaceutical grade)		Tilcotil	110	Triclosan	
Systane Unit Dose	232	Timolol		Trimethoprim	96
- T -		Cardiovascular		Trimethoprim with	
T&R		Sensory	230	sulphamethoxazole	
Tacrolimus	218	Timoptol XE		[Co-trimoxazole]	96
Tacrolimus Sandoz	218	Tiotropium bromide	224	Trisequens	
Taliglucerase alfa	29	Tiotropium bromide with		Trisul	96
Tambocor	48	olodaterol	224	Trophic Hormones	83
Tambocor CR	48	Tivicay	107	Tropicamide	231
Tamoxifen citrate	173	TMP	96	Trusopt	
Tamoxifen Sandoz	173	TOBI	96	TruSteel	
Tamsulosin hydrochloride		Tobramycin		Tuberculin PPD [Mantoux] test	
Tamsulosin-Rex		Infection	96	Tubersol	
Tandem Cartridge		Sensory		Two Cal HN	
Tandem t:slim X2		Tobramycin Mylan		Two Cal HN RTH	
Tap water		Tobrex		Tykerb	
Tarceva		Tocilizumab		Tysabri	
Tasigna		Tofranil		- U -	
Tasmar		Tolcapone		Ultibro Breezhaler	224
Tecfidera		Tolterodine			
Tegretol		Topamax		UltraproctUmeclidinium	
		Topical Products for Joint and	120	Umeclidinium with vilanterol	
Tegretol CR		•	444		
Telfast		Muscular Pain		Univent	
Temaccord		Topiramate		Ural	
Temazepam		Topiramate Actavis		Urea	
Temizole 20		Total parenteral nutrition (TPN)		Urex Forte	
Temozolomide		TPN		Urinary Agents	
Tenofovir disoproxil		Tramadol hydrochloride		Urinary Tract Infections	
Tenofovir Disoproxil Teva	102	Tramal SR 100		Uromitexan	
Tenoxicam		Tramal SR 150		Ursodeoxycholic acid	
Tepadina	157	Tramal SR 200	123	Ursosan	24
Terazosin	45	Trandate	49	Utrogestan	82
Terbinafine	98	Tranexamic acid	38	- V -	
Terbutaline sulphate	223	Tranylcypromine sulphate	124	Vaccinations	257
Teriflunomide		Trastuzumab		Vaclovir	
Teriparatide		Travatan		Valaciclovir	
Testosterone		Travoprost		Valganciclovir	

Valganciclovir Mylan102	Voltaren D
Vancomycin96	Voltaren Ophtha229
Vannair222	Volumatic227
Varenicline Pfizer153	Voriconazole98
Varenicline tartrate153	Vosol
Varicella vaccine [Chickenpox	Votrient
vaccine]266	Vttack98
Varicella zoster virus (Oka strain) live	- W -
attenuated vaccine [shingles	Warfarin sodium42
vaccine]	Wart Preparations
Varilrix	Wasp venom allergy treatment220
Various 233	Water
Vasodilators	Blood43
Vasopressin Agonists87	
	Extemporaneous
Vedafil	- X -
Velcade161	~
Veletri58	Xarelto41
Venlafaxine125	Xifaxan9
Venomil220	XMET Maxamum252
Ventavis58	Xolair
Ventolin223	XP Maxamum253
Vepesid	Xylocaine120
Verapamil hydrochloride51	Xylocaine 2% Jelly119
Vergo 16130	Xyntha38
Vermox89	- Z -
Verpamil SR51	Zantac8
Vesanoid165	Zapril45
Vexazone11	Zarontin126
Vfend98	Zaroxolyn52
Viaderm KC63	Zavedos
Vidaza157	Zeffix101
Vigabatrin128	Zetlam101
Vildagliptin11	Ziagen106
Vildagliptin with metformin	Zidovudine [AZT]106
hydrochloride11	Zidovudine [AZT] with
Vimpat126	lamivudine
Vinblastine sulphate165	Zimybe54
Vincristine sulphate166	Zinc and castor oil64
Vinorelbine166	Zinc sulphate34
Vinorelbine Ebewe	Zincaps
Viramune Suspension	Zinnat90
ViruPOS228	Ziprasidone 132
Vistil 232	Zista 221
Vistil Forte232	Zithromax90
Vit.D332	Zoladex 86
VitA-POS	Zoledronic acid
Vitabdeck32	Hormone78
	Musculoskeletal114
Vitadol C31	
Vital	Zoledronic acid Mylan78
Vitamin A with vitamins D and C31	Zopiclone
Vitamin B complex31	Zopiclone Actavis147
Vitamin B6 2531	Zostavax266
Vitamins31–32	Zostrix111
Vivonex TEN245	Zostrix HP120
Volibris55	Zuclopenthixol decanoate134
Voltaren 110	Zuclopenthixol hydrochloride 132

Zusdone	132
Zyban	
Zypine	
Zypine ODT	
Zyprexa Relprevv	
Zytina	