August 2020 Volume 27 Number 2

Editors:

Kaye Wilson & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

Published each April, August and December. Changes to the contents are published in monthly updates.

Accessible in an electronic format at no cost from the PHARMAC website www.pharmac.govt.nz/schedule.

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at www.pharmac.govt.nz/subscriptions.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth & John Geering email: texschedule@pharmac.govt.nz @Pharmaceutical Management Agency



ISSN 1179-3686 pdf ISSN 1172-9376 print

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to PHARMAC and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to PHARMAC should be in written form and not by reproduction of the PHARMAC logo. While care has been taken in compiling this Schedule, PHARMAC takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

	introducing Phanimac	
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	38
	Cardiovascular System	47
	Dermatologicals	61
	Genito Urinary System	71
	Hormone Preparations – Systemic	77
	Infections – Agents For Systemic Use	88
	Musculoskeletal System	110
	Nervous System	119
	Oncology Agents & Immunosuppressants	157
	Respiratory System & Allergies	232
	Sensory Organs	240
	Various	245
Section C	Extemporaneous Compounds (ECPs)	247
Section D	Special Foods	250
Sastian I	N	070
Section I	National Immunisation Schedule	270
	la desc	

Index

Introducing PHARMAC

Introducing PHARMAC

The Pharmaceutical Management Agency (PHARMAC) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. PHARMAC negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

PHARMAC's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about PHARMAC and the way we make funding decisions can be found on the PHARMAC website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by PHARMAC.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements PHARMAC has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg p sachet		30	•	Gaviscon Infant
SODIUM ALGINATE Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (8.60)	60		Gaviscon Double Strength
Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 m	-	Acidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE * Tab 600 mg	12.56	100	✓.	Alu-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsementOnly when prescribed for patients unable to swallow cal inappropriate and the prescription is endorsed according	cium carbonate tablet	500 m s or v		Roxane um carbonate tablets are
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on * Tab 2 mg* * Cap 2 mg	10.75	400 400		Nodia Diamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE Cap 3 mg - Special Authority see SA1886 below - Retail pharmacy	166.50	90	✓	Entocort CIR
⇒SA1886 Special Authority for Subsidy Initial application — (Crohn's disease) from any relevant practithe following criteria: Both:	titioner. Approvals va	ılid fo	r 6 months	for applications meeting
Mild to moderate ileal, ileocaecal or proximal Crohn's disc	ease; and			

2.3 Osteoporosis where there is significant risk of fracture; or

2 Any of the following: 2.1 Diabetes; or 2.2 Cushingoid habitus; or

Subs	sidy Full	/ Brand or
(Manufactur	rer's Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*: and
- 2 Patient does not have cirrhosis: and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

TITOTIO CONTINUINE ACETATE		
Rectal foam 10%, CFC-Free (14 applications)26.55	21.1 g OP	✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam S29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab EC 500 mg49.50	100	✓ Asamax
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
Modified release granules, 1 g141.72	120 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g54.60	30	✓ Pentasa
OLSALAZINE		
Tab 500 mg93.37	100	✓ Dipentum
Cap 250 mg53.00	100	✓ Dipentum

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
SODIUM CROMOGLICATE Cap 100 mg	92.91	100	✓ N	alcrom
SULFASALAZINE * Tab 500 mg Tab EC 500 mg		100 100		alazopyrin alazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CI	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cinchocaine hydrochloride 5 mg per g6.35	30 g OP	✓ Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and cinchocaine hydrochloride 1 mg2.66	12	✓ Ultraproct
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	 Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl

Management of Anal Fissures

GL	YCERYL TRINITRATE - Special Authority see SA1329 below - Retail pha	rmacy	
*	Oint 0.2%22.00	30 g OP	✓ Rectogesic

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule - Up to 10 inj available on	а		
PSO	17.14	10	Max Health
	34.32	5	✓ Robinul
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg	6.35	100	Buscopan
Buscopan to be Sole Supply on 1 October 2020			
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO	6.35	5	✓ Buscopan
MEBEVERINE HYDROCHLORIDE			
* Tab 135 mg	9.20	90	✓ Colofac
	9.20	90	✓ Colofac

Antiulcerants

Antisecretory and Cytoprotective

MISOPROSTOL

Subsidised on a PSO only if from a Family Planning New Zealand Clinic or an abortion service provider with a DHB contract and the PSO is endorsed with the name of the institution for which the PSO is required.

*	Tab 200 mcg - l	Up to 120 tab available on a F	PSO41.50	120	Cytotec
---	-----------------	--------------------------------	----------	-----	---------

Subsidy		Fully	Brand or	
(Manufacturer's Price) 5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

Helicobacter Pylori Eradication

CLARITHROMYCIN

- a) Maximum of 14 tab per prescription
- Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.
 Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

H2 Antagonists

FAI	MOTIDINE - Only on a prescription		
*	Tab 20 mg	100	✓ Famotidine
	·		Hovid S29
*	Tab 40 mg8.48	100	✓ Famotidine
			Hovid S29

RANITIDINE - Subsidy by endorsement

- a) Only on a prescription
- b) Subsidy by endorsement Subsidised for patients who were taking ranitidine prior to 1 November 2019 and the
 prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record
 of prior dispensing of ranitidine.

*	Tab 150 mg	12.91	500	Ranitidine Relief
	Tab 300 mg		500	Ranitidine Relief
	Oral liq 150 mg per 10 ml		300 ml	✓ Peptisoothe
	Inj 25 mg per ml, 2 ml		5	✓ Zantac
(Ra	nitidine Relief Tab 150 mg to be delisted 1 October 2020)			
(Ra	nitidine Relief Tab 300 mg to be delisted 1 October 2020)			
(Za	ntac Inj 25 mg per ml, 2 ml to be delisted 1 June 2021)			

Proton Pump Inhibitors

NSOPRAZOLE Cap 15 mg	100 100	✓ Lanzol Relief ✓ Lanzol Relief
IEPRAZOLE		
Cap 10 mg	90	Omeprazole actavis10
Cap 20 mg	90	 Omeprazole actavis 20
Cap 40 mg	90	✓ Omeprazole actavis 40
Powder – Only in combination42.50	5 g	✓ Midwest
Only in extemporaneously compounded omeprazole suspension.	•	
Inj 40 mg ampoule with diluent	5	✓ <u>Dr Reddy's</u> <u>Omeprazole</u>
		✓ Ocicure S29
NTOPRAZOI F		
	100	✓ Panzop Relief
Tab EC 40 mg	100	✓ Panzop Relief
,	Cap 30 mg 5.41 MEPRAZOLE For omeprazole suspension refer Standard Formulae, page 247 Cap 10 mg 1.98 Cap 20 mg 1.96 Cap 40 mg 3.12 Powder – Only in combination 42.50 Only in extemporaneously compounded omeprazole suspension. 1nj 40 mg ampoule with diluent 33.98 NTOPRAZOLE Tab EC 20 mg 2.02	Cap 15 mg 4.58 100 Cap 30 mg 5.41 100 MEPRAZOLE For omeprazole suspension refer Standard Formulae, page 247 90 Cap 10 mg 1.98 90 Cap 20 mg 1.96 90 Cap 40 mg 3.12 90 Powder – Only in combination 42.50 5 g Only in extemporaneously compounded omeprazole suspension. 1nj 40 mg ampoule with diluent 33.98 5 NTOPRAZOLE Tab EC 20 mg 2.02 100

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE Tab 120 mgSUCRALFATE	14.51	50	√ (Gastrodenol S29
Tab 1 g	35.50 (48.28)	120	(Carafate
Bile and Liver Therapy				
RIFAXIMIN - Special Authority see SA1461 below - Retail Tab 550 mg	,	56	✓)	(ifaxan
⇒SA1461 Special Authority for Subsidy nitial application only from a gastroenterologist, hepatolog nepatologist. Approvals valid for 6 months where the patient olerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or Practice approvals valid without further renewal unless penefiting from treatment.	t has hepatic encephalor	pathy d	lespite an a	dequate trial of maximum penterologist or
Diabetes				
Hyperglycaemic Agents				
DIAZOXIDE – Special Authority see SA1320 below – Retail Cap 25 mg Cap 100 mg Oral liq 50 mg per ml ➤ SA1320 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals mypoglycaemia caused by hyperinsulinism. Renewal from any relevant practitioner. Approvals valid with			P F	
appropriate and the patient is benefiting from treatment. GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit — Up to 5 kit available on a PSO	32.00	1	√ <u>(</u>	Glucagen Hypokit
Insulin - Short-acting Preparations				
NSULIN NEUTRAL ▲ Inj human 100 u per ml	25.26	10 ml C		Actrapid Humulin R
▲ Inj human 100 u per ml, 3 ml	42.66	5		Actrapid Penfill Iumulin R
Insulin - Intermediate-acting Preparations				
NSULIN ASPART WITH INSULIN ASPART PROTAMINE Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	✓ N	NovoMix 30 FlexPen

	Subsidy	F	ully Brand or
	(Manufacturer's Price	e) Subsidis	,
	\$	Per	✓ Manufacturer
NSULIN ISOPHANE			
Inj human 100 u per ml	17.68		✓ Humulin NPH✓ Protaphane
Inj human 100 u per ml, 3 ml	29.86	5	✓ Humulin NPH
- , ,			✓ Protaphane Penfill
NSULIN ISOPHANE WITH INSULIN NEUTRAL			
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70✓ Mixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70
•			✓ PenMix 30
			✓ PenMix 40
JOHN N. JORDO WITH INCH IN JORDO DROTAMINE			✓ PenMix 50
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE Inj lispro 25% with insulin lispro protamine 75% 100 u per ml.			
3 ml		5	✓ Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml			
3 ml		5	✓ Humalog Mix 50
Insulin - Long-acting Preparations			
SULIN GLARGINE			
Inj 100 u per ml, 10 ml	63.00	1	✓ Lantus
Inj 100 u per ml, 3 ml		5	✓ Lantus
Inj 100 u per ml, 3 ml disposable pen		5	✓ Lantus SoloStar
Insulin - Rapid Acting Preparations			
NSULIN ASPART			
Inj 100 u per ml, 10 ml			✓ NovoRapid
Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe	51.19	5	✓ NovoRapid FlexPen
NSULIN GLULISINE Inj 100 u per ml, 10 ml	27.02	1	✓ Apidra
Inj 100 u per ml, 3 ml			✓ Apidra
Inj 100 u per ml, 3 ml disposable pen			✓ Apidra SoloStar
NSULIN LISPRO			
Inj 100 u per ml, 10 ml		10 ml OP	✓ Humalog
Inj 100 u per ml, 3 ml	59.52	5	✓ Humalog
Alpha Glucosidase Inhibitors			
CARBOSE			
★ Tab 50 mg		90	✓ Glucobay
₭ Tab 100 mg	10.47 6.40	90	✓ Accarb✓ Glucobay
- 145 155 Hig	20.23	00	✓ Accarb
Oral Hypoglycaemic Agents			
ilibenclamide			
₭ Tab 5 mg	6.00	100	✓ <u>Daonil</u>
			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
GLICLAZIDE				
* Tab 80 mg	15.18	500	•	Glizide
GLIPIZIDE				
* Tab 5 mg	3.27	100	1	<u>Minidiab</u>
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg	8.63	1,000	/	Apotex
* Tab immediate-release 850 mg	7.04	500	•	Apotex
PIOGLITAZONE				
* Tab 15 mg	3.47	90	1	Vexazone
* Tab 30 mg		90	1	Vexazone
* Tab 45 mg	7.10	90	✓	<u>Vexazone</u>
VILDAGLIPTIN				
Tab 50 mg	40.00	60	✓	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride	40.00	60	1	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride		60	1	Galvumet

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes: or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	I Generic	
\$	Per 🗸	Manufacturer	

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	✓ CareSens N
		✓ CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood g	lucose test stri	os26.20	50 test OP	✓ SensoCard
---------	------------------	---------	------------	-------------

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	I Generic	
\$	Per 🗸	Manufacturer	

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

INSULIN PEN NEEDLES – Maximum	of 200 deviper prescription
-------------------------------	-----------------------------

*	29 g × 12.7 mm		100	✓ B-D Micro-Fi	
*	31 g × 5 mm		100	✓ B-D Micro-Fi	ne
*	31 g × 6 mm		100	✓ Berpu	
*	31 g × 8 mm		100	✓ B-D Micro-Fi	
*	32 g × 4 mm	10.50	100	B-D Micro-Fi	ne
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLI	E - Maximum of 2	00 dev per p	prescription	
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.00	100	✓ B-D Ultra Fir	ne
		1.30	10		
		(1.99)		B-D Ultra Fine	е
*	Syringe 0.3 ml with 31 g × 8 mm needle	` '	100	✓ B-D Ultra Fir	
	-, g g	1.30	10		
		(1.99)		B-D Ultra Fine	e II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	` '	100	✓ B-D Ultra Fir	ne.
•		1.30	10	2 2 0	
		(1.99)	. •	B-D Ultra Fin	e
*	Syringe 0.5 ml with 31 g × 8 mm needle	` '	100	✓ B-D Ultra Fir	
•••	Cynnigo dio nii Mar o'r g x o niin noodio	1.30	10	- 550 (11.01.11	
		(1.99)	10	B-D Ultra Fin	اا م
*	Syringe 1 ml with 29 g x 12.7 mm needle	` '	100	✓ B-D Ultra Fir	
~	Symige 1 mi with 25 g x 12.7 min needle	1.30	100	· D-D Oldarii	10
			10	B-D Ultra Fin	•
	Onderson Annal with Od an One or and the	(1.99)	400		
*	Syringe 1 ml with 31 g × 8 mm needle		100	B-D Ultra Fir	ie II
		1.30	10		
		(1.99)		B-D Ultra Fin	e II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

 c) Maximum of 1 insulin pump per patient each four year 	ear period.		
Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 640G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim X2

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and

Subsidy (Manufacturer's Pr	rice)	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist: or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — **(severe unexplained hypoglycaemia)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Eithei
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Drand or	_
Subsidy		. ,	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol: and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Fither:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Insulin Pump Consumables

⇒SA1906 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	1	Manufacturer	

continued...

8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline, according to the most recent result.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol. according to the most recent result: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline.

Initial application — (**Previous use before 1 September 2012**) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Subsic		Fully	Brand or
(Manufacture	r's Price) Subs	idised	Generic
\$	Per	1	Manufacturer

continued...

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol according to a recent laboratory result; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application, according to the most recent result; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline.

INSULIN PUMP CARTRIDGE - Special Authority see SA1906 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 packs of cartridge sets will be funded per year.

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA) - Special Authority see SA1906 on page 17 - Retail pharmacy

a)	Maximum	of 3 sets per	prescription

a) Maximum of 3 sets per prescriptionb) Only on a prescriptionc) Maximum of 13 infusion sets will be funded per year.			
10 mm steel needle; 60 cm tubing × 10	130.00	1 OP	MiniMed Sure-T MMT-884A
10 mm steel needle; 80 cm tubing × 10	130.00	1 OP	MiniMed Sure-T MMT-886A
6 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Sure-T MMT-864A
6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Sure-T MMT-866A
8 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Sure-T MMT-874A
8 mm steel needle; 80 cm tubing x 10	130.00	1 OP	✓ MiniMed Sure-T MMT-876A
10 mm steel needle; 29 G; manual insertion; 60 cm tubing \times			_
10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T MMT-884
10 mm steel needle; 29 G; manual insertion; 60 cm tubing x	100.00	1 OD	Come T MMT 000
10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-883
10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T MMT-886
10 mm steel needle; 29 G; manual insertion; 80 cm tubing x			
10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-885
6 mm steel needle; 29 G; manual insertion; 60 cm tubing x			
10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T MMT-864
6 mm steel needle; 29 G; manual insertion; 60 cm tubing ×			
10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-863
6 mm steel needle; 29 G; manual insertion; 80 cm tubing x	400.00	4.00	/ Dame diame Occurs T
10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T MMT-866
6 mm steel needle; 29 G; manual insertion; 80 cm tubing x			
10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-865
8 mm steel needle; 29 G; manual insertion; 60 cm tubing \times 10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T MMT-874
8 mm steel needle; 29 G; manual insertion; 60 cm tubing x			mini Vi T
10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-873
8 mm steel needle; 29 G; manual insertion; 80 cm tubing \times			
10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T

MMT-876

	Subsidy (Manufacturer's Price)		Fully lised	Brand or Generic Manufacturer
8 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles; luer lock	120.00	1 OP	./ 0	ure-T MMT-875
(Sure-T MMT-883 10 mm steel needle; 29 G; manual insertion; 60 September 2020)		-	-	
(Sure-T MMT-885 10 mm steel needle; 29 G; manual insertion; 80 September 2020)	0 cm tubing \times 10 with	n 10 needles	s; luer	lock to be delisted 1
(Sure-T MMT-865 6 mm steel needle; 29 G; manual insertion; 80 September 2020)	cm tubing × 10 with	10 needles;	luer lo	ock to be delisted 1
(Sure-T MMT-875 8 mm steel needle; 29 G; manual insertion; 80 September 2020)	cm tubing × 10 with	10 needles;	luer lo	ock to be delisted 1
INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT Retail pharmacy	INSERTION) - Spe	cial Authori	ty see	SA1906 on page 17 –
a) Maximum of 3 sets per prescription b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year. 6 mm steel cannula; straight insertion; 60 cm line x 10 with				
10 needles	130.00	1 OP	✓ T	ruSteel
10 needles	130.00	1 OP	✓ T	ruSteel
8 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles	130.00	1 OP	✓ T	ruSteel
8 mm steel cannula; straight insertion; 81 cm line × 10 with 10 needles	130.00	1 OP	✓ T	ruSteel

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1906 on page 17 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription

c) Maximum of 13 infusion sets will be funded per ye	ear.		
13 mm teflon needle, 110 cm tubing × 10		1 OP	✓ MiniMed Silhouette MMT-382A
13 mm teflon needle, 45 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-368A
13 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-381A
13 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-383A
17 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-377A
17 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-378A
17 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-384A
6 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-398A
6 mm teflon needle, 45 cm blue tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-941A
6 mm teflon needle, 45 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-921A
6 mm teflon needle, 60 cm blue tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-943A
6 mm teflon needle, 60 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-923A
6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-399A
6 mm teflon needle, 80 cm blue tubing	130.00	1 OP	✓ MiniMed Mio MMT-945A
6 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-965A
6 mm teflon needle, 80 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-925A
6 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-387A
9 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-396A
9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-397A
9 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-975A
9 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-386A

Fully

Brand or

Subsidy

	(Manufacturer's P	rice) Sub Per	osidised Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN SA1906 on page 17 – Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 13 mm teflon cannula; angle insertion; insertion device; 110 c		H INSERTION	N DEVICE) - Special Authority see
line x 10 with 10 needles	140.00	1 OP	✓ AutoSoft 30
line x 10 with 10 needles		1 OP	✓ AutoSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 13 mm teflon cannula; angle insertion; 120 cm line × 10 with	ISERTION) - S	pecial Author	ity see SA1906 on page 17 –
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-382
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-368
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-381
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-383
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-377
10 needles; luer lock	130.00	1 OP	✓ Silhouette MMT-371
17 mm teflon cannula; angle insertion; 60 cm line × 10 with 10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-378
17 mm teflon cannula; angle insertion; 60 cm line × 10 with 10 needles; luer lock	130.00	1 OP	✓ Silhouette MMT-373
10 needles10 needles	130.00	1 OP	✓ Paradigm Silhouette

(Silhouette MMT-371 17 mm teflon cannula; angle insertion; 110 cm line \times 10 with 10 needles; luer lock to be delisted 1 September 2020)

6 mm teflon cannula: straight insertion: insertion device: 60 cm

9 mm teflon cannula: straight insertion: insertion device: 60 cm

9 mm teflon cannula; straight insertion; insertion device;

line × 10 with 10 needles......140.00

110 cm line × 10 with 10 needles140.00

line × 10 with 10 needles......140.00

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Por	/	Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION WITH INSERTION DEVICE) - Special Authority see SA1906 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription

c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; insertion device; 45 cm			
blue tubing × 10 with 10 needles130.0	00 10	P ✓ Paradigm Mi MMT-941	io
6 mm teflon cannula; straight insertion; insertion device; 45 cm pink tubing × 10 with 10 needles	00 10	P Paradigm Mi	io
6 mm teflon cannula; straight insertion; insertion device; 60 cm blue tubing × 10 with 10 needles130.0	00 10		io
6 mm teflon cannula; straight insertion; insertion device; 60 cm pink tubing × 10 with 10 needles130.0	00 10	P ✓ Paradigm Mi MMT-923	io
6 mm teflon cannula; straight insertion; insertion device; 80 cm blue tubing × 10 with 10 needles130.0	00 10	P ✓ Paradigm Mi MMT-945	io
6 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing x 10 with 10 needles130.0	00 1 0	P ✓ Paradigm Mi MMT-965	io
6 mm teflon cannula; straight insertion; insertion device; 80 cm pink tubing × 10 with 10 needles130.0	00 1 0	P ✓ Paradigm Mi MMT-925	io
9 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing × 10 with 10 needles130.0	00 1 0	P Paradigm Mi MMT-975	io
6 mm teflon cannula; straight insertion; insertion device; 110 cm line × 10 with 10 needles140.0	00 10		

✓ AutoSoft 90

✓ AutoSoft 90

✓ AutoSoft 90

1 OP

1 OP

1 OP

Subsidy Fully Brand or (Manufacturer's Price) Generic Subsidised \$ Per Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION) - Special Authority see SA1906 on page 17 -Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription

c) Maximum of 13 infusion sets will be funded per year.		
6 mm teflon cannula; straight insertion; 110 cm tubing × 10 with 10 needles130.00	1 OP	✓ Paradigm Quick-Set MMT-398
6 mm teflon cannula; straight insertion; 110 cm tubing × 10 with 10 needles; luer lock	1 OP	✓ Quick-Set MMT-391
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with		/ Damadiana Oviale Cat
10 needles	1 OP	✓ Paradigm Quick-Set MMT-399
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 10 needles; luer lock130.00	1 OP	✓ Quick-Set MMT-393
6 mm teflon cannula; straight insertion; 80 cm tubing x 10 with 10 needles130.00	1 OP	✓ Paradigm Quick-Set
9 mm teflon cannula; straight insertion; 106 cm tubing × 10 with		MMT-387
10 needles130.00	1 OP	✓ Paradigm Quick-Set MMT-396
9 mm teflon cannula; straight insertion; 110 cm tubing x 10 with 10 needles; luer lock130.00	1 OP	✓ Quick-Set MMT-390
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 with		
10 needles130.00	1 OP	✓ Paradigm Quick-Set MMT-397
9 mm teflon cannula; straight insertion; 60 cm tubing x 10 with 10 needles; luer lock130.00	1 OP	✓ Quick-Set MMT-392
9 mm teflon cannula; straight insertion; 80 cm tubing × 10 with		
10 needles	1 OP	✓ Paradigm Quick-Set

(Quick-Set MMT-391 6 mm teflon cannula; straight insertion; 110 cm tubing x 10 with 10 needles; luer lock to be delisted 1 September 2020)

(Quick-Set MMT-390 9 mm teflon cannula; straight insertion; 110 cm tubing x 10 with 10 needles; luer lock to be delisted 1 September 2020)

INSULIN PUMP RESERVOIR - Special Authority see SA1906 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 packs of reservoir sets will be funded per year.

of maximum or to passes or recorrent code min so tantaca per years		
10 × luer lock conversion cartridges 1.8 ml for Paradigm pumps50.00	1 OP	✓ ADR Cartridge 1.8
Cartridge for 5 and 7 series pump; 1.8 ml × 1050.00	1 OP	✓ Paradigm
Cartridge for 7 series pump; 3.0 ml × 1050.00	1 OP	1.8 Reservoir ✓ Paradigm
California of the California o	101	3.0 Reservoir

Subsid	idy Fully	Brand or
(Manufacture	er's Price) Subsidised	Generic
\$	Per 🗸	Manufacturer

Digestives Including Enzymes

PANCREATIC FNZYME

FANOREATIC ENZINE			
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ <u>Creon 10000</u>
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase, 1,250 U protease))	94.40	100	✓ Panzytrat
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ Creon 25000
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph			_
Eur U)	34.93	20 g OP	Creon Micro
URSODEOXYCHOLIC ACID - Special Authority see SA1739 below	/ – Retail pha	rmacy	
Cap 250 mg Ursosan to be Sole Supply on 1 October 2020		100	✓ Ursosan

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (**Primary biliary cholangitis**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (**Pregnancy**) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

	ALIMENTAR	RY TRACT	AND	METABOLISM
	Subsidy (Manufacturer's Price \$		Fully dised	Brand or Generic Manufacturer
continued Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment. Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels. Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure — doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.				
Laxatives Bulk-forming Agents				
ISPAGHULA (PSYLLIUM) HUSK – Only on a prescription * Powder for oral soln Konsyl-D to be Sole Supply on 1 November 2020 MUCILAGINOUS LAXATIVES WITH STIMULANTS	12.20	500 g OP	✓ K	onsyl-D
* Dry	(17.32)	500 g OP 200 g OP		ormacol Plus
Faecal Softeners				
DOCUSATE SODIUM - Only on a prescription				

* Tab 50 mg	2.31	100	Coloxyl
Coloxyl to be Sole Supply on 1 October 2020 * Tab 120 mg	3.13	100	✓ Coloxyl
Coloxyl to be Sole Supply on 1 October 2020			•
DOCUSATE SODIUM WITH SENNOSIDES * Tab 50 mg with sennosides 8 mg	3.10	200	✓ <u>Laxsol</u>
POLOXAMER - Only on a prescription			
Not funded for use in the ear.			
* Oral drops 10%	3.98	30 ml OP	Coloxyl
Coloxyl to be Sole Supply on 1 November 2020			

Opioid Receptor Antagonists - Peripheral

METHYLNALTREXONE BROMIDE - Special Authority	see SA1691 below - Retail ph	armacy	1
Inj 12 mg per 0.6 ml vial	36.00	1	✓ Relistor
	246.00	7	✓ Relistor

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

continued...

Both:

- 1 The patient is receiving palliative care; and
- 2 Either:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

Osmotic Laxatives

GLYCEROL		
* Suppos 3.6 g - Only on a prescription	20	✓ <u>PSM</u>
LACTULOSE – Only on a prescription		
* Oral liq 10 g per 15 ml	500 ml	✓ Laevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE A Powder for oral soln 13.125 g with potassium chloride 46.6 mg,	ND SODIUM C	CHLORIDE
sodium bicarbonate 178.5 mg and sodium chloride 350.7 mg 6.70 Molaxole to be Sole Supply on 1 October 2020	30	✓ Molaxole
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%2.50	1	✓ Fleet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE - Only on a pro	escription	
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml,		
5 ml	50	✓ <u>Micolette</u>
Stimulant Laxatives		
BISACODYL - Only on a prescription		
* Tab 5 mg5.99	200	✓ <u>Lax-Tab</u>
* Suppos 10 mg	10	 Lax-Suppositories

Metabolic Disorder Agents

SENNA - Only on a prescription

ALGLUCOSIDASE ALFA - Special Authority see SA1920 below - Ret	tail pharmacy		
Inj 50 mg vial1,	,142.60 1	,	✓ Myozyme

* Tab, standardised......2.17

⇒SA1920 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease;
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or

continued...

100

20

(8.21)

(2.06)

Senokot

Senokot

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
- 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
- 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

⇒SA1921 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism: and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation. Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient

is benefiting from treatment.

GALSULFASE – Special Authority see SA1922 below – Retail pharmacy

Inj 1 mg per ml, 5 ml vial......2,234.00

......2,234.00 1 **V** Naglazyme

⇒SA1922 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

IDURSULFASE - Special Authority see SA1623 below - Retail pharmacy ✓ Elaprase

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT): and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

LARONIDASE - Special Authority see SA1695 below - Retail pharmacy ✓ Aldurazyme

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assav in cultured skin fibroblasts: or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome: and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT): and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

SAPROPTERIN DIHYDROCHLORIDE - Special Authority see SA1923 on the next page - Retail pharmacy 30 OP ✓ Kuvan

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

⇒SA1923 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE - Special Authority see SA1599 below	/ – Retail pharmacy		
Soln 100 mg per ml	CBS	100 ml	✓ Amzoate S29

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1924 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

TALIGLUCERASE ALFA - S	Special Authority see SA1880 on the next page -	Retail pharmacy	
Inj 200 unit vial		1	✓ Elelyso

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

⇒SA1880 Special Authority for Subsidy

Special Authority approved by the Gaucher Treatment Panel

Notes: Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The Co-ordinator, Gaucher Treatment Panel Phone: 04 460 4990 PHARMAC PO Box 10 254 Facsimile: 04 916 7571

Wellington Email: gaucherpanel@pharmac.govt.nz

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Access Criteria

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and
- Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
- Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia, thrombocytopenia; at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or massive symptomatic splenomegaly; or
 - 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g. hips or shoulder); spontaneous fractures or vertebral collapse; chronic bone pain not controlled by other pharmaceuticals; or
 - 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
 - 5) Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated; and
- 2) Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3) Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and three yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4) Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and

	Subsidy (Manufacturer's	Price) Subs	Fully	Brand or Generic
	\$	Per	1	Manufacturer
ontinued 6) Patient is compliant with regular treatment and taligluce every other week rounded to the nearest whole vial (2007) Supporting clinical information including test reports, MF investigations are submitted to the Gaucher Panel for as) units), unless of RI whole body STI	herwise agreed R, haematologi	by PH.	ARMAC; and
Mouth and Throat				
Agents Used in Mouth Ulceration				
BENZYDAMINE HYDROCHLORIDE				
Soln 0.15% - Higher subsidy of \$20.31 per 500 ml with				
Endorsement		500 ml	_	
A 100 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	(20.31)		_)ifflam
Additional subsidy by endorsement for a patient who h prescription is endorsed accordingly.	as oral mucositis	as a result of tre	eatmen	t for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN				
Paste	17 20	56 g OP	√ 0	Stomahesive
i dole	4.55	15 g OP	• 0	otomanesive
	(7.90)	10 g 01	C	Drabase
	1.52	5 g OP		
	(3.60)	Ü	C	Drabase
Powder	8.48	28 g OP		
	(10.95)		S	Stomahesive
CHLORHEXIDINE GLUCONATE				
Mouthwash 0.2%	2.57	200 ml OP	√ h	ealthE
healthE Mouthwash 0.2% to be delisted 1 November 2020)				
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE				
★ Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP		
	(6.00)		В	Bonjela
RIAMCINOLONE ACETONIDE				
Paste 0.1%		5 g OP	✓ K	Cenalog in Orabase
Kenalog in Orabase to be Sole Supply on 1 November	2020			
Oropharyngeal Anti-infectives				
MPHOTERICIN B				
Lozenges 10 mg	5.86	20	✓ F	ungilin
MICONAZOLE				=
Oral gel 20 mg per g	4.74	40 g OP	✓ D)ecozol
YSTATIN		3 -	_	
0.18.40000				

Other Oral Agents

For folinic mouthwash, pilocarpine oral liquid or saliva substitute formula refer Standard Formulae, page 247

24 ml OP

✓ Nilstat

THYMOL GLYCERIN

Oral liq 100,000 u per ml1.76

Nilstat to be Sole Supply on 1 October 2020

	Subsidy (Manufacturer's Price	a)	Fully Subsidised	Brand or Generic
	\$	Per		Manufacturer
Vitamins				
Vitamin B				
HYDROXOCOBALAMIN				
* Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a	PSO1.89	3	•	Neo-B12
PYRIDOXINE HYDROCHLORIDE				
a) No more than 100 mg per dose				
b) Only on a prescription	0.70	00		Vitamin DC 05
* Tab 25 mg - No patient co-payment payable	2.70	90	•	Vitamin B6 25
* Tab 50 mg	13.63	500	1	Apo-Pyridoxine
THIAMINE HYDROCHLORIDE – Only on a prescription				
* Tab 50 mg	4.89	100	1	Max Health
VITAMIN B COMPLEX				
* Tab, strong, BPC	7.15	500	1	Bplex
Vitamin C				
ASCORBIC ACID				
a) No more than 100 mg per dose				
b) Only on a prescription			_	
* Tab 100 mg	9.90	500	•	<u>Cvite</u>
Vitamin D				
ALFACALCIDOL				
* Cap 0.25 mcg		100	_	One-Alpha
* Cap 1 mcg * Oral drops 2 mcg per ml		100 20 ml O	_	One-Alpha One-Alpha
	00.08	20 1111 0	· ·	One-Aipha
CALCITRIOL * Cap 0.25 mcg	7 95	100	1	Calcitriol-AFT
* Cap 0.5 mcg		100	_	Calcitriol-AFT
COLECALCIFEROL				
* Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescri	ption2.50	12	1	Vit.D3
* Oral liq 188 mcg per ml (7,500 iu per ml)		4.8 ml C)P 🗸	Puria
Multivitamin Preparations				
MULTIVITAMIN RENAL - Special Authority see SA1546 below	v – Retail pharmacy			
* Cap	6.49	30	1	Clinicians Renal Vit
⇒SA1546 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals va	alid without further rer	newal ur	nless notif	ied for applications meeting
the following criteria:				
Either: 1 The patient has chronic kidney disease and is receiving	either peritoneal diel	ucic or h	aamadial	veie: or
 The patient has chronic kidney disease grade 5, defined The patient has chronic kidney disease grade 5, defined ml/min/1.73 m² body surface area (BSA). 				
MULTIVITAMINS - Special Authority see SA1036 on the next	page – Retail pharma	acv		
* Powder		200 g O	P 🗸	Paediatric Seravit
		-		

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer	

⇒SA1036 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins.

VITAMINS

*	Tab (BPC cap strength)1	1.45	1,000	✓ Mvite
*	Cap (fat soluble vitamins A, D, E, K) - Special Authority see			
	SA1720 below – Retail pharmacy2	3.40	60	✓ Vitabdeck

⇒SA1720 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency; or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

Minerals		
Calcium		
CALCIUM CARBONATE * Tab eff 1.75 g (1 g elemental)28.40	20	✓ Calcium Sandoz S29
* Tab 1.25 g (500 mg elemental)	250	✓ Arrow-Calcium
* Inj 10%, 10 ml ampoule32.00	10	✓ Max Health - HameIn S29
64.00	20	✓ Max Health S29
Fluoride		
SODIUM FLUORIDE * Tab 1.1 mg (0.5 mg elemental)	100	✓ PSM
lodine		
POTASSIUM IODATE Tab 253 mcg (150 mcg elemental iodine)	90	✓ NeuroTabs
Iron		
FERRIC CARBOXYMALTOSE – Special Authority see SA1840 below – Retail phar Inj 50 mg per ml, 10 ml150.00	rmacy 1	✓ Ferinject
To CA1940 Chaniel Authority for Cubaidy		•

⇒SA1840 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

100	✓ <u>Ferro-tab</u>
60	✓ Ferro-F-Tabs
500 ml	✓ Ferodan
30	✓ Ferrograd
5	✓ Ferrosig
	60 500 ml 30

Magnesium

For magnesium hydroxide mixture refer Standard Formulae, page 247

MAGNESIUM HYDROXIDE	
Suspension 8%	

Cappanoidi C/S	000 1111	
MAGNESIUM SULPHATE		
* Inj 2 mmol per ml, 5 ml ampoule10.21	10	✓ DBL
		✓ DBL S29 S29

✓ T&R S29

500 ml

72 20

ALIMENTARY TRACT AND METABOLISM

✓ Zincaps

100

Subsidy (Manufacturer's Price) Subsidised Per Subsidised Manufacturer

Subsidy (Manufacturer's Price) Subsidised Per Manufacturer

✓ Manufacturer

ZINC SULPHATE

* Cap 137.4 mg (50 mg elemental)......11.00

Subsidy (Manufacturer's Price) \$

Fully Subsidised Per

Brand or Generic Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus: and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*: and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 ju per week.

Note: Indication marked with * is an unapproved indication

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Subsidised	
	\$	Per	1	Manufacturer
EPOETIN ALFA - Special Authority see SA1775 on the previous	page - Retail phan	macy		
Wastage claimable		,		
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	✓	Binocrit
Inj 2,000 iu in 1 ml, syringe		6	✓	Binocrit
Inj 3,000 iu in 0.3 ml, syringe	150.00	6	✓	Binocrit
Inj 4,000 iu in 0.4 ml, syringe		6	1	Binocrit
Inj 5,000 iu in 0.5 ml, syringe	125.00	6	1	Binocrit
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	1	Binocrit
Inj 8,000 iu in 0.8 ml, syringe	175.00	6	1	Binocrit
Inj 10,000 iu in 1 ml, syringe	197.50	6	1	Binocrit
Inj 40,000 iu in 1 ml, syringe	250.00	1	✓	Binocrit
Megaloblastic				
FOLIC ACID				
* Tab 0.8 mg	21.84	1,000	1	Apo-Folic Acid
* Tab 5 mg		500		Apo-Folic Acid
Oral lig 50 mcg per ml		25 ml C)P 🗸	Biomed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemonhilia Management group

ricators aroup in conjunction with the National I	iacinopinna management giot	φ.	
Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial	2,450.00	1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	✓ Alprolix
ELTROMBOPAG - Special Authority see SA1743 be	elow – Retail pharmacy		
Wastage claimable			
Tab 25 mg	1,550.00	28	Revolade
Tab 50 mg	3,100.00	28	Revolade

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy: and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab);
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding: or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist.

Subsidy	Į	Fully	Brand or
(Manufacturer's Price)	Subsid	lised	Generic
\$	Per	✓	Manufacturer

continued...

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Fither:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe	2,356.60	1	✓ NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓ NovoSeven RT
Inj 8 mg syringe	9,426.40	1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	1,315.00	1	✓ FEIBA NF
Inj 1,000 U	2,630.00	1	✓ FEIBA NF
Inj 2,500 U	6,575.00	1	✓ FEIBA NF

	DECOD AND	DECOD! O	TIMING OTTAKIO
	Subsidy	Full	y Brand or
	(Manufacturer's Price)	Subsidise	
	\$	Per 🗸	Manufacturer
MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpha			
For patients with haemophilia. Rare Clinical Circumstances			
treatment is managed by the Haemophilia Treaters Group in	conjunction with the N	National Haem	ophilia Management Group,
subject to criteria.			
Inj 250 iu prefilled syringe			' Xyntha
Inj 500 iu prefilled syringe			Xyntha
Inj 1,000 iu prefilled syringe			Xyntha
Inj 2,000 iu prefilled syringe			Xyntha
Inj 3,000 iu prefilled syringe	*	1	' Xyntha
NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xpharm			
For patients with haemophilia. Access to funded treatment i	s managed by the Hae	emophilia Trea	ters Group in conjunction
with the National Haemophilia Management Group.			
Inj 500 iu vial			RIXUBIS
Inj 1,000 iu vial			RIXUBIS
Inj 2,000 iu vial	,		RIXUBIS
Inj 3,000 iu vial	2,610.00	1	RIXUBIS
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) -			
For patients with haemophilia. Preferred Brand of short half			
managed by the Haemophilia Treaters Group in conjunction	with the National Hae	mophilia Mana	igement Group.
Inj 250 iu vial	210.00	1	' Advate
Inj 500 iu vial			' Advate
Inj 1,000 iu vial			Advate
Inj 1,500 iu vial	,		Advate
Inj 2,000 iu vial	,		Advate
Inj 3,000 iu vial	•	1	' Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE	FS) - [Xpharm]		
For patients with haemophilia. Rare Clinical Circumstances			
treatment is managed by the Haemophilia Treaters Group in	conjunction with the N	National Haem	ophilia Management Group,
subject to criteria.			_
Inj 250 iu vial			Kogenate FS
Inj 500 iu vial			Kogenate FS
Inj 1,000 iu vial			Kogenate FS
Inj 2,000 iu vial	,		Kogenate FS
Inj 3,000 iu vial	•	1	Kogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]			
For patients with haemophilia A receiving prophylaxis treatm		d treatment is i	managed by the Haemophilia
Treaters Group in conjunction with the National Haemophilia			
Inj 250 iu vial			Adynovate
Inj 500 iu vial			Adynovate
Inj 1,000 iu vial	· ·		Adynovate
Inj 2,000 iu vial	2,400.00	1	' Adynovate
SODIUM TETRADECYL SULPHATE			
* Inj 3% 2 ml	28.50	5	
	(73.00)		Fibro-vein
TRANEXAMIC ACID			
Tab 500 mg	9.45	60	Mercury Pharma

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
Vitamin K				
PHYTOMENADIONE Inj 2 mg per 0.2 ml - Up to 5 inj available on a PSO Inj 10 mg per ml, 1 ml - Up to 5 inj available on a PSO		5 5		Konakion MM Konakion MM
Antithrombotic Agents				
Antiplatelet Agents				
ASPIRIN * Tab 100 mg CLOPIDOGREL	10.80	990	•	Ethics Aspirin EC
* Tab 75 mg	4.60	84	•	Clopidogrel Multichem
DIPYRIDAMOLE * Tab long-acting 150 mg PRASUGREL – Special Authority see SA1201 below – Retail ph		60	✓	Pytazen SR

⇒SA1201 Special Authority for Subsidy

Initial application — (coronary angioplasty and bare metal stent) from any relevant practitioner. Approvals valid for 6 months where the patient has undergone coronary angioplasty in the previous 4 weeks and is clopidogrel-allergic*.

Initial application — (drug eluting stent) from any relevant practitioner. Approvals valid for 12 months where the patient has had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Initial application — (stent thromobosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (coronary angioplasty and bare metal stent) from any relevant practitioner. Approvals valid for 6 months where the patient has undergone coronary angioplasty or had a bare metal cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Renewal — (drug eluting stent) from any relevant practitioner. Approvals valid for 12 months where had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Note: * Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

TICAGRELOR - Special Authority see SA1887 below - Retail pharmacy

56 ✓ Brilinta

⇒SA1887 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention post neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

1 Patient has had a neurological stenting procedure* in the last 60 days; and

continued...

✓ Effient

✓ Effient

28

(Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer		Subsidy	Fully	Brand or
\$ Per ✓ Manufacturer	(Manufa			Generic
		\$ Per	r 🗸	Manufacturer

continued...

- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention post neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Note: indications marked with * are unapproved indications.

Heparin and Antagonist Preparations

ENOXAPARIN SODIUM — Special Authority see SA1646 be	elow – Retail pharmacy	,	
Inj 20 mg in 0.2 ml syringe	27.93	10	Clexane
Inj 40 mg in 0.4 ml syringe		10	Clexane
Inj 60 mg in 0.6 ml syringe		10	Clexane
Inj 80 mg in 0.8 ml syringe		10	Clexane
Inj 100 mg in 1 ml syringe		10	Clexane
Inj 120 mg in 0.8 ml syringe		10	Clexane
			Clexane Forte
Inj 150 mg in 1 ml syringe	133.20	10	Clexane
			✓ Clexane Forte

(Clexane Inj 120 mg in 0.8 ml syringe to be delisted 1 January 2021) (Clexane Ini 150 mg in 1 ml syringe to be delisted 1 January 2021)

⇒SA1646 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
continued	1			
Renewal — (Pregnancy, Malignancy or Haemodialysis) fro	m any relevant practition	oner. App	rovals v	alid for 1 year for
pplications meeting the following criteria:				
ny of the following:				
Low molecular weight heparin treatment is required dur		•		
2 For the treatment of venous thromboembolism where the			dialuaia	
3 For the prevention of thrombus formation in the extra-co				
Renewal — (Venous thromboembolism other than in pregr				
ralid for 1 month where low molecular weight heparin treatmen	t or prophylaxis is requ	iired for a	secona	or subsequent event
surgery, ACS, cardioversion, or prior to oral anti-coagulation).				
HEPARIN SODIUM	F0 F7	F0		M:
Inj 1,000 iu per ml, 5 ml ampoule		50	_	<u>ffizer</u>
Inj 5,000 iu per ml, 1 ml		5	-	fizer
Ini F 000 in manual. Furthermorals	32.66	F0		lospira
Inj 5,000 iu per ml, 5 ml ampoule		50	_	<u>ffizer</u>
Inj 25,000 iu per ml, 0.2 ml		5		lospira
	42.40			leparin DBL S29
			✓ F	leparin
				Ratiopharm S29
	122.00	10	✓ V	Vockhardt S29
	190.00	50	✓ P	fizer S29
Heparin Ratiopharm 👀 Inj 25,000 iu per ml, 0.2 ml to be de	listed 1 January 2021)			
Wockhardt 🕯 Inj 25,000 iu per ml, 0.2 ml to be delisted 1 J	anuary 2021)			
HEPARINISED SALINE				
Inj 10 iu per ml, 5 ml	65.48	50	✓ P	fizer
			•	
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg - No more than 2 cap per day		60		radaxa
Cap 110 mg		60		radaxa
Cap 150 mg	/6.36	60	✓ F	radaxa
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day		30		(arelto
Tab 15 mg - Up to 14 tab available on a PSO		28		Carelto
Tab 20 mg	77.56	28	✓ X	arelto
VARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
★ Tab 1 mg	3.46	50		Coumadin
	6.46	100	✓ N	larevan
₭ Tab 2 mg	4.31	50	✓ (Coumadin
₭ Tab 3 mg	10.03	100	✓ N	larevan
•	10.03		✓ N	

Blood Colony-	stimulating Factors
---------------	---------------------

FILGRASTIM – Special Authority see SA1259 on the next page -	 Retail pharmacy 		
Inj 300 mcg per 0.5 ml prefilled syringe	96.22	10	✓ Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe	161.50	10	✓ Nivestim

✓ Marevan

100

11.48

Subsidy (Manufacturer's Price)	s	Fully	Brand or Generic	
\$	Per	✓	Manufacturer	

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×109/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10⁹/L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM – Special Authority see SA1912 below – Retail pharmacy
Inj 6 mg per 0.6 ml syringe1,080.00 1

✓ Neulastim

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration

b) Not in combination

GLUCOSE [DEXTROSE]			
* Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO	30.65	5	✓ Biomed
Biomed to be Sole Supply on 1 November 2020			
* Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO	15.00	1	✓ Biomed
Biomed to be Sole Supply on 1 November 2020			
POTASSIUM CHLORIDE			
* Inj 75 mg per ml, 10 ml	55.00	50	✓ AstraZeneca
			Potassium Chloride
			Aguettant S29
SODIUM BICARBONATE			
Inj 8.4%, 50 ml	19.95	1	✓ Biomed
a) Up to 5 inj available on a PSO			
b) Not in combination			
Inj 8.4%, 100 ml	20.50	1	✓ Biomed
a) Up to 5 inj available on a PSO			- 51011104
a) Up to 3 iiij avaliable on a F3O			

	Subsidy		Fully Brand or
	(Manufacturer's F	Price) Subs Per	idised Generic Manufacturer
SODIUM CHLORIDE	Ψ	1 01	- Marianacaror
Not funded for use as a nasal drop. Not funded for nebulise	r use except whe	en used in conju	unction with an antibiotic intended
for nebuliser use.	•	,	
Inj 0.9%, bag – Up to 2000 ml available on a PSO	1.23 1.26	500 ml 1,000 ml	✓ Baxter✓ Baxter
Only if prescribed on a prescription for renal dialysis, ma	aternity or post-n	,	home of the patient, or on a PSC
for emergency use. (500 ml and 1,000 ml packs)			
Inj 23.4% (4 mmol/ml), 20 ml ampoule		5	✓ Biomed
For Sodium chloride oral liquid formulation refer Standar			/ Formation Wald
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO		20	Fresenius Kabi
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO Inj 0.9%, 20 ml ampoule		50 20	 ✓ <u>Fresenius Kabi</u> ✓ Fresenius Kabi
• •	5.00	20	Fresenius Kabi
TOTAL PARENTERAL NUTRITION (TPN)	000		4
Infusion	CBS	1 OP	✓ TPN
VATER			
 On a prescription or Practitioner's Supply Order only w Schedule requiring a solvent or diluent; or On a bulk supply order; or When used in the extemporaneous compounding of ey When used for the dilution of sodium chloride soln 7% 	/e drops; or		ection issed in the Pharmaceutic
4) When used for the dilution of sociality chiloride soil 1 //s	ioi cysuc iibiosis	patients only.	
Inj 5 ml ampoule – Up to 5 inj available on a PSO	7.00	50	✓ InterPharma
Inj 10 ml ampoule – Up to 5 inj available on a PSO		50	✓ Pfizer
Inj 20 ml ampoule – Up to 5 inj available on a PSO		20	✓ Fresenius Kabi
, , ,			✓ Multichem
	7.50	30	✓ InterPharma
Oral Administration			
CALCIUM POLYSTYRENE SULPHONATE			
Powder	169.85	300 g OP	✓ Calcium Resonium
COMPOUND ELECTROLYTES		555 9 51	
Powder for oral soln — Up to 5 sach available on a PSO	9 77	50	✓ Electral
		30	Liectiai
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]		1 000 00	✓ Dadiabata
Soln with electrolytes (2 × 500 ml)	6.55	1,000 ml OP	✓ <u>Pedialyte -</u> Bubblegum
			Bubbleguiii
PHOSPHORUS			
Tab eff 500 mg (16 mmol)	82.50	100	Phosphate Phebra
POTASSIUM CHLORIDE			
★ Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	5.26	60	
	(11.85)		Chlorvescent
* Tab long-acting 600 mg (8 mmol)	8.90	200	✓ <u>Span-K</u>
SODIUM BICARBONATE			
Cap 840 mg	8.52	100	✓ Sodibic ✓ Sodibic

SODIUM POLYSTYRENE SULPHONATE

✓ Resonium-A

454 g OP

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Alpha-Adrenoceptor Blockers

Alpha Adrenoceptor Blockers

DOXAZOSIN * Tab 2 mg	500 500	✓ Apo-Doxazosin✓ Apo-Doxazosin
PHENOXYBENZAMINE HYDROCHLORIDE		
* Cap 10 mg65.00	30	✓ BNM S29
216.67	100	✓ Dibenzyline \$29
PRAZOSIN		
* Tab 1 mg5.53	100	✓ Apo-Prazosin
* Tab 2 mg7.00	100	✓ Apo-Prazosin
* Tab 5 mg11.70	100	✓ Apo-Prazosin
TERAZOSIN		
* Tab 1 mg	28	✓ Actavis
Tab 2 mg7.50	500	✓ Apo-Terazosin
Tab 5 mg10.90	500	✓ Apo-Terazosin
(Actavis Tab 1 mg to be delisted 1 October 2020)		•

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

CAPTOPRIL		_
* Oral liq 5 mg per ml94.99	95 ml OP	✓ Capoten
Oral liquid restricted to children under 12 years of age.		
CILAZAPRIL		
* Tab 0.5 mg2.09	90	✓ Zapril
* Tab 2.5 mg4.80	90	✓ Zapril
* Tab 5 mg	90	✓ Zapril
ENALAPRIL MALEATE		
* Tab 5 mg1.82	100	✓ Acetec
* Tab 10 mg	100	✓ Acetec
* Tab 20 mg	100	✓ Acetec
··· · · · · · · · · · · · · ·	100	Acetec
LISINOPRIL		
* Tab 5 mg	90	✓ Ethics Lisinopril
* Tab 10 mg2.36	90	Ethics Lisinopril
* Tab 20 mg3.17	90	Ethics Lisinopril
PERINDOPRIL		
* Tab 2 mg	30	Apo-Perindopril
* Tab 4 mg4.80	30	✓ Apo-Perindopril
QUINAPRIL		
* Tab 5 mg	90	✓ Arrow-Quinapril 5
* Tab 10 mg	90	✓ Arrow-Quinapril 10
	90	✓ Arrow-Quinapril 20
* Tab 20 mg4.89	90	MITOW-Quillapili 20

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)	Subsi Per	Fully dised	Brand or Generic Manufacturer	
ACE Inhibitors with Divrotios					

ACE Inhibitors with Diuretics

CILAZAPRIL WITH HYDROCHLOROTHIAZIDE - Subsidy by endorsement

Subsidy by endorsement - Subsidised for patients who were taking cilazapril with hydrochlorothiazide prior to 1 March 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril with hydrochlorothiazide.

Tab 5 mg with hydrochlorothiazide 12.5 mg......10.18 100 ✓ Apo-Cilazapril/ Hydrochlorothiazide

(Apo-Cilazapril/ Hydrochlorothiazide Tab 5 mg with hydrochlorothiazide 12.5 mg to be delisted 1 December 2020)

HINAPRII	WITH HYD	ROCHL	OROTHIAZIDE

	Tab 10 mg with hydrochlorothiazide 12.5 mg	3.57	28	✓ Accuretic
	• •	3.83	30	✓ Accuretic 10
*	Tab 20 mg with hydrochlorothiazide 12.5 mg	4.92	30	✓ Accuretic 20

Angiotensin II Antagonists

CANDESARTAN CILEXETII

0,	INDEONITY OF CHERE ITE			
*	Tab 4 mg	1.90	90	✓ Candestar
	Tab 8 mg		90	✓ Candestar
	Tab 16 mg		90	✓ Candestar
	Tab 32 mg		90	✓ Candestar
	SARTAN POTASSIUM			
*	Tab 12.5 mg	1.39	84	✓ Losartan Actavis
*	Tab 25 mg	1.63	84	Losartan Actavis
*	Tab 50 mg	2.00	84	Losartan Actavis
	Tab 100 mg		84	✓ Losartan Actavis

Angiotensin II Antagonists with Diuretics

LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE		
* Tab 50 mg with hydrochlorothiazide 12.5 mg	30	✓ Arrow-Losartan &
		Hydrochlorothiazide

Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN - Special Authority see SA1905 below - Retail pharmacy

Note: Due to the angiotensin II receptor blocking activity of sacubitril with valsartan it should not be co-administered with an ACE inhibitor or another ARR

AGE ITTIBITOR OF ALIGHRAP.			
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/103

⇒SA1905 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II: or

Subsidy	Fu	,	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
\$	Per	/	Manufacturer

continued...

- 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, page 119

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaestnetics, Local, p	age 119	
AMIODARONE HYDROCHLORIDE		
▲ Tab 100 mg	30	✓ Aratac
▲ Tab 200 mg5.25	30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule - Up to 10 inj available on a		
PSO16.37	10	✓ Max Health
ATROPINE SULPHATE		
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO12.07	10	✓ Martindale
	10	Martinuale
DIGOXIN		
* Tab 62.5 mcg - Up to 30 tab available on a PSO7.00	240	✓ Lanoxin PG
* Tab 250 mcg - Up to 30 tab available on a PSO15.20	240	✓ <u>Lanoxin</u>
* Oral liq 50 mcg per ml16.60	60 ml	Lanoxin
		✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE		
▲ Cap 100 mg23.87	100	✓ Rythmodan
FLECAINIDE ACETATE		,
▲ Tab 50 mg	60	✓ Flecainide BNM
▲ Cap long-acting 100 mg	90	✓ Flecainide Bivivi
Cap long-acting 100 mg9.51	90	Controlled
		Release Teva
A Con long acting 000 mg	00	✓ Flecainide
▲ Cap long-acting 200 mg61.06	90	
		<u>Controlled</u> Release Teva
1,10	_	
Inj 10 mg per ml, 15 ml ampoule100.00	5	✓ Tambocor
MEXILETINE HYDROCHLORIDE		
▲ Cap 150 mg162.00	100	Mexiletine
		Hydrochloride
		USP S29
▲ Cap 250 mg202.00	100	Mexiletine
		Hydrochloride
		USP S29
PROPAFENONE HYDROCHLORIDE		
▲ Tab 150 mg	50	✓ Rytmonorm
		,

CARDIOVASCULAR SYSTEM

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Por	_	Manufacturer

Antihypotensives

MIDODRINE - Special Authority see SA1474 below - Retail pharm	nacy		
Tab 2.5 mg	53.00	100	Gutron
Tab 5 mg	79.00	100	Gutron

⇒SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Note: Treatment should be started with small doses and titrated upwards as necessary. Hypertension should be avoided, and the usual target is a standing systolic blood pressure of 90 mm Hg.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

TENOLOL		
≰ Tab 50 mg4.26	500	Mylan Atenolol
≰ Tab 100 mg7.30	500	Mylan Atenolol
♦ Oral liq 25 mg per 5 ml21.25	300 ml OP	✓ Atenolol AFT
		✓ Atenolol AFT S29 S29
Restricted to children under 12 years of age.		023 023
SISOPROLOL FUMARATE		
₭ Tab 2.5 mg	90	✓ Bosvate
₭ Tab 5 mg5.15	90	✓ Bosvate
₭ Tab 10 mg9.40	90	✓ Bosvate
CARVEDILOL		
₹ Tab 6.25 mg2.24	60	✓ Carvedilol Sandoz
₭ Tab 12.5 mg2.30	60	✓ Carvedilol Sandoz
₭ Tab 25 mg2.95	60	✓ Carvedilol Sandoz
CELIPROLOL		
₹ Tab 200 mg21.40	180	✓ Celol
ABETALOL	100	5 60101
	100	(5)
₭ Tab 100 mg11.36	100	✓ Presolol S29
14.50		✓ Trandate
Trandate to be Sole Supply on 1 September 2020	400	/ Turn data
≰ Tab 200 mg27.00	100	✓ Trandate
29.74		✓ Presolol S29
Trandate to be Sole Supply on 1 September 2020	_	
k Inj 5 mg per ml, 20 ml ampoule59.06	5	
(88.60)		Trandate
≰ inj 5 mg per ml, 20 ml vial42.29	1	
(48.20)		Alvogen S29
Presolol S29 Tab 100 mg to be delisted 1 September 2020)		
Presolol S29 Tab 200 mg to be delisted 1 September 2020)		

	Subsidy		Fully	
	(Manufacturer's Price)		Subsidised	
	<u> </u>	Per		Manufacturer
METOPROLOL SUCCINATE				
* Tab long-acting 23.75 mg	1.03	30	✓	Betaloc CR
* Tab long-acting 47.5 mg		30	✓	Betaloc CR
* Tab long-acting 95 mg	1.99	30	✓	Betaloc CR
* Tab long-acting 190 mg		30	✓	Betaloc CR
METOPROLOL TARTRATE				
* Tab 50 mg	5.66	100	/	Apo-Metoprolol
* Tab 100 mg		60		Apo-Metoprolol
* Tab long-acting 200 mg		28		Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial		5	✓	Metroprolol IV
				Mylan
NADOLOL				
* Tab 40 mg	16.69	100	/	Apo-Nadolol
* Tab 80 mg		100		Apo-Nadolol
PINDOLOL				
* Tab 5 mg	13 22	100	1	Apo-Pindolol
* Tab 10 mg		100		Apo-Pindolol
* Tab 15 mg		100		Apo-Pindolol
		100	•	Apo i illuoloi
PROPRANOLOL	4.04	400	,	
* Tab 10 mg		100		Apo-Propranolol
* Tab 40 mg		100		Apo-Propranolol
* Cap long-acting 160 mg		100	•	Cardinol LA
* Oral liq 4 mg per ml - Special Authority see SA1327 below				
Retail pharmacy	CBS	500 m	nı 🗸	Roxane S29

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

ТΑ		

*	Tab 80 mg	32.58	500	Mylan
	Tab 160 mg		100	✓ Mylan
TIN	MOLOL			
*	Tab 10 mg	10.55	100	✓ Apo-Timol

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

AMLODIPINE			
Tab 2.5 mg	1.72	100	✓ Apo-Amlodipine
Tab 5 mg	3.33	250	✓ Apo-Amlodipine
Tab 10 mg	4.40	250	✓ Apo-Amlodipine

CARDIOVASCULAR SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
LODIPINE	•			
Tab long-acting 2.5 mg	1.45	30	✓	Plendil ER
Tab long-acting 5 mg		90	_	Felo 5 ER
Tab long-acting 10 mg		90	_	Felo 10 ER
FEDIPINE				
Tab long-acting 10 mg	10.62	60	1	Adalat 10
rab long-acting to mg	10.03	00		
Tab lane action 00 mm	17.70	100		Adefin S29
Tab long-acting 20 mg.		100		Nyefax Retard
Tab long-acting 30 mg		30	_	Adalat Oros
Tab long-acting 60 mg	5.67	30	_	Adalat Oros
			•	Adefin XL
Other Calcium Channel Blockers				
TIAZEM HYDROCHLORIDE			_	
Tab 30 mg		100		Dilzem
Tab 60 mg		100		Dilzem
Cap long-acting 120 mg		500		Apo-Diltiazem CD
Cap long-acting 180 mg		500		Apo-Diltiazem CD
Cap long-acting 240 mg	66.76	500	•	Apo-Diltiazem CD
RHEXILINE MALEATE				
Tab 100 mg	62.90	100	✓	Pexsig
RAPAMIL HYDROCHLORIDE				
Tab 40 mg	7.01	100	✓	Isoptin
Tab 80 mg		100		Isoptin
Tab long-acting 120 mg		100		Isoptin Retard \$29
. az 151.g azıg 1=0g				Isoptin SR
Tab long-acting 240 mg	15.12	30		Isoptin SR
. az 151.g azıg = 10g	25.00	250	_	Verpamil SR
Inj 2.5 mg per ml, 2 ml ampoule - Up to 5 inj available on a				
PSO		5	✓	Isoptin
erpamil SR Tab long-acting 240 mg to be delisted 1 Septemb	er 2020)			·
Centrally-Acting Agents				
ONIDINE				
Patch 2.5 mg, 100 mcg per day - Only on a prescription	10.34	4	•	Mylan
Mylan to be Sole Supply on 1 November 2020				
Patch 5 mg, 200 mcg per day – Only on a prescription	13.18	4	✓	Mylan
Mylan to be Sole Supply on 1 November 2020			_	
Patch 7.5 mg, 300 mcg per day - Only on a prescription	16.93	4	/	Mylan
Mylan to be Sole Supply on 1 November 2020				
ONIDINE HYDROCHLORIDE				
Tab 25 mcg	8.75	112	✓	Clonidine BNM
Tab 150 mcg		100		Catapres
Inj 150 mcg per ml, 1 ml ampoule		10	_	Medsurge
THYLDOPA				
Tab 250 mg	15 10	100	ſ	Methyldopa Mylan
		100	•	moniyidopa iviyiali
1 ab 250 mg	52.85	500		Methyldopa Mylan

28

50

✓ Frumil

✓ Moduretic

	Subsidy	Ful	ly Brand or
	(Manufacturer's Price		
	\$	Per •	Manufacturer
Diuretics			
Loop Diuretics			
BUMETANIDE			
* Tab 1 mg	4.91	30	Burinex S29 S29
•	16.36	100	Burinex
* Inj 500 mcg per ml, 4 ml vial	7.95	5	Burinex
FUROSEMIDE [FRUSEMIDE]			
* Tab 40 mg - Up to 30 tab available on a PSO	7.24	1,000	Apo-Furosemide
* Tab 500 mg	25.00	50	Urex Forte
* Oral liq 10 mg per ml	11.20	30 ml OP ✓	Lasix
* Inj 10 mg per ml, 25 ml ampoule	60.65	6 ✓	Lasix Lasix
* Inj 10 mg per ml, 2 ml ampoule - Up to 5 inj available on a	PSO 1.15	5	Frusemide-Claris
Potassium Sparing Diuretics			
AMILORIDE HYDROCHLORIDE			
Oral liq 1 mg per ml	30.00	25 ml OP ✔	/ Biomed
EPLERENONE – Special Authority see SA1728 below – Retail			
Tab 50 mg		30	/ Inspra
Tab 25 mg			Inspra
⇒SA1728 Special Authority for Subsidy			
Initial application from any relevant practitioner. Approvals va	lid without further rer	newal unless not	ified for applications meeting
the following criteria:			
Both:			
1 Patient has heart failure with ejection fraction less than 4	10%; and		
2 Either:	·		
2.1 Patient is intolerant to optimal dosing of spironola	ctone; or		
2.2 Patient has experienced a clinically significant ad	verse effect while on	optimal dosing	of spironolactone.
METOLAZONE			•
Tab 5 mg	CDC	1	Metolazone S29
1 ab 5 mg		-	
		50 ✔	Zaroxolyn S29
SPIRONOLACTONE	4.00	400	.
* Tab 25 mg			Spiractin
* Tab 100 mg			Spiractin
Oral liq 5 mg per ml	30.60	25 ml OP	Biomed

Potassium Sparing Combination Diuretics AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE

* Tab 5 mg with furosemide 40 mg8.63

* Tab 5 mg with hydrochlorothiazide 50 mg......5.00

AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE

	Subsidy (Manufacturer's Pr	ica) Su	Fully bsidised	Brand or Generic
	(Wandiacturer 3 i i	Per	D3IUI3EU ✓	Manufacturer
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]				
* Tab 2.5 mg - Up to 150 tab available on a PSO	12.50	500	✓ A	rrow- Bendrofluazide
May be supplied on a PSO for reasons other than emerg	ency.			
* Tab 5 mg	20.42	500	✓ A	rrow- Bendrofluazide
CHLOROTHIAZIDE				
Oral liq 50 mg per ml	26.00	25 ml OP	✓ B	liomed
CHLORTALIDONE [CHLORTHALIDONE]				
* Tab 25 mg	6.50	50	✓ H	lygroton
NDAPAMIDE	10.45	00	./ n	lone Tobe
* Tab 2.5 mg	10.45	90	• 1	apa-Tabs
Lipid-Modifying Agents				
Fibrates				
BEZAFIBRATE				
* Tab 200 mg		90	_	lezalip
* Tab long-acting 400 mg	12.89	30	✓ <u>B</u>	Sezalip Retard
GEMFIBROZIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who were endorsed accordingly. Pharmacists may annotate the prescr				
dispensing of gemfibrozil.	10.56	60	. / I	inazil
* Tab 600 mg(Lipazil Tab 600 mg to be delisted 1 January 2021)	19.50	60	• •	ipazil
Other Lipid-Modifying Agents				
ACIPIMOX				
* Cap 250 mg	21.56	30	✓ 0	lbetam
			√ 0	Olbetam S29 S29
NICOTINIC ACID	4.40	400		
Tab 50 mg Tab 500 mg		100 100		po-Nicotinic Acid
,	17.09	100	V A	po-Nicotinic Acid
Resins				
COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g	28.60	30	√ 0	olestid
HMG CoA Reductase Inhibitors (Statins)				
ATORVASTATIN				
* Tab 10 mg	6.96	500	√ L	orstat
* Tab 20 mg		500		orstat
* Tab 40 mg		500	_	orstat
* Tab 80 mg	27.19	500	✓ <u>L</u>	orstat

CARDIOVASCULAR SYSTEM

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	•	Manufacturer
PRAVASTATIN				
★ Tab 10 mg	3.55	28	✓	Pravastatin Mylan
★ Tab 20 mg	4.72	100	✓	Apo-Pravastatin
₭ Tab 40 mg		28	✓	Pravastatin Mylan
·	8.06	100	1	Apo-Pravastatin
SIMVASTATIN				
★ Tab 10 mg	1.23	90	✓	Simvastatin Mylan
Simvastatin Mylan to be Sole Supply on 1 November	2020			•
₭ Tab 20 mg	2.03	90	✓	Simvastatin Mylan
Simvastatin Mylan to be Sole Supply on 1 November				•
₭ Tab 40 mg	3.58	90	✓	Simvastatin Mylan
Simvastatin Mylan to be Sole Supply on 1 November				,
★ Tab 80 mg		90	1	Simvastatin Mylan
Simvastatin Mylan to be Sole Supply on 1 November				,

Selective Cholesterol Absorption Inhibitors

EZETIMIBE – Special Authority see SA1045 below – Retail pharmacy		
* Tab 10 mg1.95	30	Ezetimibe Sandoz
Ezetimibe Sandoz to be Sole Supply on 1 October 2020		

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than $10 \times \text{normal}$) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin: or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 below - Retail pharmacy

Tab 10 mg with simvastatin 10 mg	5.15	30	✓ Zimybe
Tab 10 mg with simvastatin 20 mg	6.15	30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg		30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg	8.15	30	✓ Zimybe

⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and

	Subsidy (Manufacturer's Price \$) Sul	Fully osidised	Brand or Generic Manufacturer
continued				
 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and 3 The patient has not reduced their LDL cholesterol to less atorvastatin. 	than 2.0 mmol/litre w	vith the us	e of the r	maximal tolerated dose o
Notes: A patient who has failed to reduce their LDL cholesterol t statin should use a more potent statin prior to consideration bein Other treatment options including fibrates, resins and nicotinic ac If a patient's LDL cholesterol cannot be calculated because the tr	g given to the use of cid should be conside riglyceride level is too	non-statir ered after o high the	n therapion failure of n a repea	es. f statin therapy. at test should be
performed and if the LDL cholesterol again cannot be calculated 2.0 mmol/litre.	then it can be consid	dered that	the LDL	cholesterol is greater that
Renewal from any relevant practitioner. Approvals valid for 2 ye benefiting from treatment.	ars where the treatm	nent remai	ins appro	opriate and the patient is
Nitrates				
GLYCERYL TRINITRATE				
* Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	4.4E 0E	0 dose OF	> √ N	litralingual Dump
available on a P50	4.45 25	u dose Of	• • N	litrolingual Pump Spray

不	Oral pump spray, 400 mcg per dose – Op to 250 dose			
	available on a PSO	4.45	250 dose OP	✓ Nitrolingual Pump Spray
*	Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
*	Patch 50 mg, 10 mg per day	18.62	30	✓ Nitroderm TTS
IS	OSORBIDE MONONITRATE			
*	Tab 20 mg	19.55	100	✓ Ismo 20
	Ismo 20 to be Sole Supply on 1 November 2020			
*	Tab long-acting 40 mg	8.20	30	✓ Ismo 40 Retard
	Ismo 40 Retard to be Sole Supply on 1 November 2020			
*	Tab long-acting 60 mg	9.25	90	Duride
	Duride to be Sole Supply on 1 November 2020			

Sympathomimetics		
ADRENALINE		
Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a PSO4.98	5	Aspen Adrenaline
10.76		✓ DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PSO27.00	5	✓ Hospira
49.00	10	✓ Aspen Adrenaline
ISOPRENALINE [ISOPROTERENOL]		
* Inj 200 mcg per ml, 1 ml ampoule	25	
(164.20)		Isuprel

Vasodilators		
HYDRALAZINE HYDROCHLORIDE		
* Tab 25 mg - Special Authority see SA1321 on the next page -		
Retail pharmacyCBS	1	✓ Hydralazine
	56	✓ Onelink S29
	84	✓ AMDIPHARM S29
	100	✓ Onelink S29
* Inj 20 mg ampoule25.90	5	✓ Apresoline

CARDIOVASCULAR SYSTEM

Subsidy (Manufacturer's Price)	Fully Subsidised		,		Brand or Generic
` \$	Per	•	Manufacturer		

⇒SA1321 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

MINIOVIDII

- 1 For the treatment of refractory hypertension; or
- 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

MINOXIDIL			
▲ Tab 10 mg	70.00	100	Loniten
NICORANDIL			
▲ Tab 10 mg	25.57	60	✓ Ikorel
▲ Tab 20 mg	32.28	60	✓ Ikorel
PAPAVERINE HYDROCHLORIDE			
* Inj 12 mg per ml, 10 ml ampoule	217.90	5	✓ Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]			
Tab 400 mg	42.26	50	✓ Trental 400

Endothelin Receptor Antagonists

		AMBRISENTAN – Special Authority see SA1702 below – Retail pharmacy
✓ Volibris	30	Tab 5 mg4,585.00
✓ Volibris	30	Tab 10 mg4,585.00

⇒SA1702 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

BOSENTAN - Special Authority see SA1908 below - Retail pharmacy

1ab 62.5 mg141.00	60	Bosentan Dr
		Reddy's
Tab 125 mg141.00	60	✓ Bosentan Dr
		Reddy's

⇒SA1908 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV: and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil: or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or

ARDIOVASCULAR SYSTEM

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	/	Manufacturer

continued...

- 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
- 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV: or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy: and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL - Special Authority see SA1909 below - Retail pharmacy			
Tab 25 mg	.0.64	4	✓ Vedafil
Tab 50 mg	.0.64	4	✓ Vedafil
Tab 100 mg	.6.60	12	✓ Vedafil

⇒SA1909 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital

CARDIOVASCULAR SYSTEM

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

ulceration; digital ulcers; or gangrene); and

- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II: or
 - 3.2 PAH is in NYHA/WHO functional class III: or
 - 3.3 PAH is in NYHA/WHO functional class IV; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age, or health system capacity constraints.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Prostacyclin Analogues

EPOPROSTENOL - Special Authority see SA1696 below - Retail pharmacy	/	
Inj 500 mcg vial36.6	1 1	✓ Veletri
Inj 1.5 mg vial73.21	1 1	✓ Veletri

⇒SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ILOPROST - Special Authority see SA1705 on the next page - Retail pharmacy

Nebuliser soln 10 mcg per ml, 2 ml740.10 30 ✓ Ventavis

CARDIOVASCULAR SYSTEM

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 88

ADAPALENE

- a) Maximum of 30 g per prescription
- b) Only on a prescription

b) Only on a prescription			
Crm 0.1%	22.89	30 g OP	Differin
Gel 0.1%	22.89	30 g OP	Differin
ISOTRETINOIN - Special Authority see SA1475 below - Rei	ail pharmacy		
Cap 5 mg	8.14	60	Oratane
Cap 10 mg	13.34	120	✓ Oratane
Cap 20 mg	20.49	120	✓ Oratane

⇒SA1475 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice: and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
 - 3.2 Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
- 2 Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

TRFTINOIN

Crm 0.5 mg per q − Maximum of 50 g per prescription13.90 50 g OP ✓ ReTrieve

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 88

HYDROGEN PEROXIDE

	Subsidy (Manufacturer's F \$	Price) Subs	sidised	Brand or Generic Manufacturer
MUPIROCIN Oint 2%	6.60 (9.26)	15 g OP	Bac	troban
a) Only on a prescriptionb) Not in combination	,			
SODIUM FUSIDATE [FUSIDIC ACID] Crm 2%	1.59	5 g OP	✓ Fok	an_
a) Maximum of 5 g per prescription b) Only on a prescription c) Not in combination Oint 2%	1 50	5 g OP	✓ Fol	an
a) Maximum of 5 g per prescription b) Only on a prescription c) Not in combination	1.00	3 y Oi	• <u>101</u>	<u>uun</u>
SULFADIAZINE SILVER Crm 1%	10.80	50 g OP	✓ Fla	mazine
For systemic antifungals, refer to INFECTIONS, Antifungals, AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		5 ml OP	✓ Myd	coNail
a) Only on a prescription b) Not in combination Nail-soln 8%	5.72	7 ml OP	✓ <u>Apo</u>	o-Ciclopirox
CLOTRIMAZOLE * Crm 1% a) Only on a prescription	0.70	20 g OP	✓ Clo	mazol
b) Not in combination * Soln 1% a) Only on a prescription	4.36 (7.55)	20 ml OP	Car	esten
b) Not in combination ECONAZOLE NITRATE				
Crm 1%	1.00 (7.48)	20 g OP	Pev	aryl
a) Only on a prescription b) Not in combination Foaming soln 1%, 10 ml sachets	9.89 (17.23)	3	Pev	aryl
a) Only on a prescriptionb) Not in combination	()		. 2.	•

				ATOLOGICALO
	Subsidy (Manufacturer's I	Price) Su Per	Fully ubsidised	Brand or Generic Manufacturer
MICONAZOLE NITRATE				_
* Crm 2%	0.74	15 g OP	✓ N	lultichem
* Lotn 2%	4.36 (10.03)	30 ml OP	D	aktarin
a) Only on a prescription b) Not in combination * Tinct 2%	4.36	30 ml OP		
a) Only on a prescription b) Not in combination	(12.10)	55 m 6 .	D	aktarin
Antipruritic Preparations				
CALAMINE a) Only on a prescription b) Not in combination Crm, aqueous, BP	1.26	100 g	√ h	ealthE Calamine

CROTAMITON

a) Only on a prescription

b) Not in combination

Crm 10%......3.29

29.60

✓ Itch-Soothe

ΒP

Aqueous Cream

MENTHOL - Only in combination

1) Only in combination with a dermatological base or proprietary Topical Corticosteriod - Plain

2) With or without other dermatological galenicals.

Crystals......6.92

25 g 100 g

20 g OP

✓ MidWest

✓ MidWest

Corticosteroids Topical

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 78

Corticosteroids - Plain

BE	TAMETHASONE DIPROPIONATE		
	Crm 0.05%2.96	15 g OP	Diprosone
	8.97	50 g OP	Diprosone
	Oint 0.05%2.96	15 g OP	Diprosone
	8.97	50 g OP	Diprosone
	Oint 0.05% in propylene glycol base4.33	30 g OP	Diprosone OV
BE	TAMETHASONE VALERATE		
*	Crm 0.1%3.45	50 g OP	✓ Beta Cream
	Oint 0.1%3.45		✓ Beta Ointment
*	Lotn 0.1%	50 ml OP	✓ Betnovate
CL	OBETASOL PROPIONATE		
*	Crm 0.05%2.18	30 g OP	✓ Dermol
*	Oint 0.05%2.12	30 g OP	✓ Dermol

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's P		sidised	Generic
	\$	Per		Manufacturer
CLOBETASONE BUTYRATE				
Crm 0.05%	5.38	30 g OP		
	(7.09)	-	1	Eumovate
DIFLUCORTOLONE VALERATE				
Crm 0.1%	8 97	50 g OP		
0111 0.170	(15.86)	00 g 01		Nerisone
Fatty oint 0.1%		50 g OP	'	INCHOOLIC
ratty office. 170	(15.86)	30 g Oi	ı	Nerisone
(Nerisone Crm 0.1% to be delisted 1 December 2020)	(10.00)			1101100110
Nerisone Fatty oint 0.1% to be delisted 1 December 2020)				
,				
HYDROCORTISONE	0.40	00 00		
★ Crm 1% – Only on a prescription		30 g OP		DermAssist
	3.70	100 g OP	✓ I	Hydrocortisone
				(PSM)
	17.15	500 g	✓ I	Hydrocortisone
				(PSM)
Hydrocortisone (PSM) to be Sole Supply on 1 September				
★ Powder – Only in combination		25 g		ABM
Up to 5% in a dermatological base (not proprietary Topical	al Corticosteriod	I – Plain) with o	or with	out other dermatological
galenicals				
DermAssist Crm 1% to be delisted 1 September 2020)				
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only or	n			
a prescription		250 ml	✓ 1	DP Lotn HC
DP Lotn HC to be Sole Supply on 1 October 2020		200 1111		D. 200
HYDROCORTISONE BUTYRATE				
Lipocream 0.1%	6 05	100 g OP	./ 1	Loosid Linearcom
•		•		Locoid Lipocream Locoid
Oint 0.1%		100 g OP 100 ml OP	_	Locoid Crelo
Milky emul 0.1%	13.70	100 IIII OP	•	Locold Creio
ACTUVU DDCDNUCOLONG ACCDONIATE				
Crm 0.1%	4.95	15 g OP		Advantan
		15 g OP 15 g OP		Advantan Advantan
Crm 0.1% Oint 0.1%		•		
Crm 0.1% Oint 0.1%	4.95	•	✓ /	
Crm 0.1% Oint 0.1% MOMETASONE FUROATE	4.95	15 g OP 15 g OP	√	Advantan
Crm 0.1% Oint 0.1% MOMETASONE FUROATE	4.95 1.51 2.50	15 g OP 15 g OP 50 g OP	✓ <u>/</u> ✓ <u> </u>	Advantan Elocon Alcohol Free Elocon Alcohol Free
Crm 0.1% Oint 0.1% MOMETASONE FUROATE Crm 0.1%	4.95 1.51 2.50	15 g OP 15 g OP 50 g OP 15 g OP	✓ / ✓ ! ✓ !	Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon
Crm 0.1% Oint 0.1% #IOMETASONE FUROATE Crm 0.1% Oint 0.1%	4.95 1.51 2.50 1.51 2.90	15 g OP 15 g OP 50 g OP	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon Elocon
Crm 0.1%	4.95 1.51 2.50 1.51 2.90	15 g OP 15 g OP 50 g OP 15 g OP 50 g OP	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon
Crm 0.1% Oint 0.1% MOMETASONE FUROATE Crm 0.1% Oint 0.1% Lotn 0.1% TRIAMCINOLONE ACETONIDE	4.95 1.51 2.50 1.51 2.90 6.30	15 g OP 15 g OP 50 g OP 15 g OP 50 g OP 30 ml OP	✓ / · · · · · · · · · · · · · · · · · ·	Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon Elocon Elocon
Crm 0.1% Oint 0.1% MOMETASONE FUROATE Crm 0.1% Oint 0.1% Lotn 0.1% TRIAMCINOLONE ACETONIDE Crm 0.02%	4.95 1.51 2.50 1.51 2.90 6.30	15 g OP 15 g OP 50 g OP 15 g OP 50 g OP	✓ / · · · · · · · · · · · · · · · · · ·	Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon Elocon
Oint 0.1%	4.95 1.51 2.50 1.51 2.90 6.30	15 g OP 15 g OP 50 g OP 15 g OP 50 g OP 30 ml OP	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Advantan Elocon Alcohol Free Elocon Elocon Elocon Elocon Aristocort
Crm 0.1% Oint 0.1% MOMETASONE FUROATE Crm 0.1% Oint 0.1% Lotn 0.1% TRIAMCINOLONE ACETONIDE Crm 0.02% Aristocort to be Sole Supply on 1 November 2020 Oint 0.02%	4.95 1.51 2.50 1.51 2.90 6.30	15 g OP 15 g OP 50 g OP 15 g OP 50 g OP 30 ml OP	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon Elocon Elocon
Crm 0.1% Oint 0.1% ### MOMETASONE FUROATE Crm 0.1% Oint 0.1% Lotn 0.1% FRIAMCINOLONE ACETONIDE Crm 0.02% Aristocort to be Sole Supply on 1 November 2020 Oint 0.02% Aristocort to be Sole Supply on 1 November 2020	4.95 1.51 2.50 1.51 2.90 6.30	15 g OP 15 g OP 50 g OP 15 g OP 50 g OP 30 ml OP	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon Elocon Elocon Aristocort
Crm 0.1%	4.95 1.51 2.50 1.51 2.90 6.30	15 g OP 15 g OP 50 g OP 15 g OP 50 g OP 30 ml OP	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Advantan Elocon Alcohol Free Elocon Elocon Elocon Elocon Aristocort
Crm 0.1%	4.95 1.51 2.50 1.51 2.90 6.30 6.30	15 g OP 15 g OP 50 g OP 15 g OP 50 g OP 30 ml OP	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon Elocon Elocon Aristocort
Crm 0.1%	4.951.51 2.501.51 2.906.306.35	15 g OP 15 g OP 50 g OP 15 g OP 50 g OP 30 ml OP 100 g OP	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Advantan Elocon Alcohol Free Elocon Elocon Elocon Elocon Aristocort
Crm 0.1%	4.951.51 2.501.51 2.906.306.35	15 g OP 15 g OP 50 g OP 15 g OP 50 g OP 30 ml OP	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Advantan Elocon Alcohol Free Elocon Alcohol Free Elocon Elocon Elocon Aristocort

	Subsidy (Manufacturer's P	rice) Subsi	Fully Brand or idised Generic ✓ Manufacturer
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FU Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP	Fucicort
a) Maximum of 15 g per prescriptionb) Only on a prescription			
HYDROCORTISONE WITH MICONAZOLE − Only on a prescri ★ Crm 1% with miconazole nitrate 2%		15 g OP	✓ Micreme H
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN — Crm 1% with natamycin 1% and neomycin sulphate 0.5% Oint 1% with natamycin 1% and neomycin sulphate 0.5% TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	3.35 3.35	15 g OP 15 g OP	✓ Pimafucort ✓ Pimafucort
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 n and gramicidin 250 mcg per g - Only on a prescription	ng	15 g OP	Viaderm KC
Disinfecting and Cleansing Agents			
CHLORHEXIDINE GLUCONATE – Subsidy by endorsement a) No more than 500 ml per month b) Only if prescribed for a dialysis patient and the prescript * Handrub 1% with ethanol 70%	4.29 3.98	cordingly. 500 ml 500 ml	✓ healthE ✓ healthE
a) Maximum of 500 ml per prescription b) a) Only if prescribed for a patient identified with Meth surgery in hospital and the prescription is endorsed b) Only if prescribed for a patient with recurrent Stapl accordingly Soln 1%	d accordingly; or nylococcus aureus	. ,	, ,,
(healthE Soln 1% to be delisted 1 November 2020)		300 1111 01	• Health
Barrier Creams and Emollients			
Barrier Creams			
DIMETHICONE * Crm 5% pump bottle	4.48	500 ml OP	✓ <u>healthE</u> Dimethicone 5%
* Crm 10% pump bottle	4.52	500 ml OP	✓ <u>healthE</u> <u>Dimethicone 10%</u>
ZINC AND CASTOR OIL * Oint	4.25	500 g	✓ Boucher
Emollients			
AQUEOUS CREAM * Crm	1.92	500 g	✓ Boucher

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's		sidised	Generic
	\$	Per		Manufacturer
CETOMACROGOL				
├ Crm BP	2.48	500 g	√ h	nealthE
CETOMACROGOL WITH GLYCEROL			_	
Crm 90% with glycerol 10%	2.35	500 ml OP	✓ p	\DF
Offit 30 /6 with grycolor 10 /6	2.00	300 1111 01	-	Boucher
	3.10	1,000 ml OP	_	ADE
	3.10	1,000 1111 01	-	Boucher
MULSIFYING OINTMENT			٠ -	<u>Jouonici</u>
* Oint BP	2.50	500 a	√	\CT
	3.59	500 g	•	AFI
DIL IN WATER EMULSION				
₭ Crm	2.19	500 g	√ <u>C</u>	D/W Fatty Emulsion
				<u>Cream</u>
PARAFFIN				
Oint liquid paraffin 50% with white soft paraffin 50%	5.35	500 ml OP	✓ <u>h</u>	<u>realthE</u>
JREA				
* Crm 10%	1.37	100 g OP	√ h	ealthE Urea Cream
NOOL FAT WITH MINERAL OIL - Only on a prescription			•	
Lotn hydrous 3% with mineral oil	E 60	1 000 ml		
K Lour nyurous 3% with mineral oil		1,000 ml	_	OP Lotion
	(11.95)	050 00	L	DP LOUON
	1.40	250 ml OP	-	ND Latina
	(4.53)	4 000	L	OP Lotion
	5.60	1,000 ml		Mala a Marit Latina
	(20.53)			Alpha-Keri Lotion
	(23.91)	050 OD		BK Lotion
	1.40	250 ml OP	_	NZ Latian
	(7.73)			3K Lotion
Other Dermatological Bases				
Other Berniatological Bases				
PARAFFIN				
White soft - Only in combination	4.99	450 g		<u>nealthE</u>
	19.99	2,500 g	✓ h	<u>realthE</u>
Only in combination with a dermatological galenical or	as a diluent for a	proprietary Top	ical Co	rticosteroid – Plain.
Minor Skin Infections				
POVIDONE IODINE				
Oint 10%	7.40	65 g OP	/ F	Betadine
a) Maximum of 130 g per prescription	7.40	03 g Oi	٠.	Detaume
,				
b) Only on a prescription				
c) Betadine to be Sole Supply on 1 October 2020	0.55	100 ml	./ -	Riodine
Antiseptic Solution 10%			_	
Antiseptic soln 10%		15 ml	_	Riodine
Chin proporation positions in the 400/ with 000/ -ttt	5.40	500 ml	A F	Riodine
Skin preparation, povidone iodine 10% with 30% alcohol		100 ml	_	Ostadina Cliin Duar
<u> </u>	(3.48)	100 ml	ь	Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	(7.78)	100 1111	_	Pfizer

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Parasiticidal Preparations

DIMETHICONE

* Lotn 4% 200 ml OP healthE Dimethicone 4% Lotion

IVERMECTIN - Special Authority see SA1225 below - Retail pharmacy

Tab 3 mg - Up to 100 tab available on a PSO.......17.20 ✓ Stromectol

- 1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- 3) For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA1225 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scables hyperinfestation (Crusted/ Norwegian scables); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy: or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation. Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria: Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

microbiologist; and

- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scables hyperinfestation (Crusted/ Norwegian scables); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy: or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

PERMETHRIN

Crm 5%	75 3	60gOP ∙	Lyderm
Lyderm to be Sole Supply on 1 November 2020	20 00	0 OD	/ A Cooking
Lotn 5%	99 30	J MI OP	✓ A-Scabies
PHENOTHRIN			
Shampoo 0.5% 11.3	36 20	0 ml OP	✓ Parasidose

Psoriasis and Eczema Preparations

ACITRETIN - Special Authority see SA1476 below - Retail pharmacy	1		
Cap 10 mg	17.86	60	✓ Novatretin
Novatretin to be Sole Supply on 1 October 2020			
Cap 25 mg	41.36	60	Novatretin
Novatretin to be Sole Supply on 1 October 2020			

⇒SA1476 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and

		DE	:RM/	ATOLOGICALS
	Subsidy (Manufacturer's Pric		ully ised	Brand or Generic Manufacturer
continued				
3 Either:				
3.1 Patient is female and has been counselled and und pregnancy and the applicant has ensured that the commencement of the treatment and that the patie treatment and for a period of two years after the countries. 3.2 Patient is male.	possibility of pregnent is informed that ompletion of the tre	ancy has beer she must not b atment; or	exclu pecom	uded prior to the ne pregnant during
Renewal from any relevant practitioner. Approvals valid for 1 year Either:	ar for applications i	neeting the fol	lowing	g criteria:
 Patient is female and has been counselled and understandand the applicant has ensured that the possibility of pregnot reatment and that the patient is informed that she must not years after the completion of the treatment; or Patient is male. 	ancy has been exc	luded prior to	the co	ommencement of the
BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL				
Foam spray 500 mcg with calcipotriol 50 mcg per g	59.95	60 g OP	√ E	nstilar
Gel 500 mcg with calcipotriol 50 mcg per g	52.24	60 g OP	✓ D	aivobet
Oint 500 mcg with calcipotriol 50 mcg per g	19.95	30 g OP	✓ D	aivobet
CALCIPOTRIOL		-		
Oint 50 mcg per g	40.00	120 g OP	✓ D	aivonex
		120 g O1	• •	aivolicx
COAL TAR	22.25	000 1		
Soln BP – Only in combination		200 ml	_	lidwest
 Up to 10% only in combination with a dermatologic With or without other dermatological galenicals. 	al base or proprieta	ary Topical Co	rticost	eriod – Plain
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULF Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and				
allantoin crm 2.5%		75 g OP		
	(8.00)		E	gopsoryl TA
	3.43	30 g OP	_	3-1
	(4.35)		E	gopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR	()		_	3-1
	4.07	25 a OB	./ c	ooo Cooln
Soln 12% with salicylic acid 2% and sulphur 4% oint	7.95	25 g OP		oco-Scalp
		40 g OP	• (oco-Scalp
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium Pinetarsol to be Sole Supply on 1 November 2020		a prescription 500 ml	✓ P	inetarsol
SALICYLIC ACID				
Powder - Only in combination	18.88	250 g	✓ M ✓ P	lidwest SM
 Only in combination with a dermatological base or p With or without other dermatological galenicals. 	proprietary Topical	Corticosteroid	i – Pla	uin or collodion flexible

1) Only in combination with a dermatological base or proprietary Topical Corticosteroid - Plain

100 g

✓ Midwest

Precipitated - Only in combination......6.35

2) With or without other dermatological galenicals.

SULPHUR

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Scalp Preparations

BETAMETHASONE VALERATE			
* Scalp app 0.1%	7.75	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE * Scalp app 0.05%	5.69	30 ml OP	✓ <u>Dermol</u>
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%	7.30	100 ml OP	✓ Locoid
KETOCONAZOLE Shampoo 2%	3.23	100 ml OP	✓ Sebizole

- a) Maximum of 100 ml per prescription
- b) Only on a prescription
- c) Sebizole to be Sole Supply on 1 November 2020

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

200 q OP Marine Blue Lotion

SPF 50+

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 68

IMIQUIMOD

Crm 5%, 250 mg sachet......21.72

24

3.5 ml OP

✓ Perrigo

PODOPHYLLOTOXIN

✓ Condyline

✓ Condyline S29 S29

- a) Maximum of 3.5 ml per prescription
- b) Only on a prescription

(Condyline S29 S29 Soln 0.5% to be delisted 1 September 2020)

Other Skin Preparations

Antineoplastics

FLUOROURACIL SODIUM

20 g OP ✓ Efudix

Subsidy (Manufacturer's Price) \$ Per

Fully Subsidised Brand or Generic Manufacturer

Contraceptives - Non-hormonal

Condoms

CO	NDOMS			
*	49 mm - Up to 144 dev available on a PSO	11.42	144	✓ Moments
	53 mm		10	✓ Moments
		11.64	144	✓ Moments
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
*	53 mm, 0.05 mm thickness	0.95	10	✓ Moments
-	, , , , , , , , , , , , , , , , , , , ,	11.42	144	✓ Moments
	a) Up to 60 dev available on a PSO			<u></u>
	b) Maximum of 60 dev per prescription			
*	53 mm, chocolate, brown	0.95	10	✓ Moments
-1-	oo min, onocolato, brown	11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO	11.04	1 1 1 1	Momento
	b) Maximum of 60 dev per prescription			
*	53 mm, strawberry, red	0.05	10	✓ Moments
*	55 min, snawberry, red	11.64	144	✓ Moments
	a) The to CO downwish has an a DCO	11.04	144	wioments
	a) Up to 60 dev available on a PSO			
*	b) Maximum of 60 dev per prescription	0.07	10	✓ Moments
不	56 mm		10	✓ Moments
	\	11.64	144	woments
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO	4.00	40	40 111/111
*	56 mm, 0.05 mm thickness		12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
*	56 mm, 0.08 mm thickness		10	✓ <u>Moments</u>
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
*	56 mm, 0.08 mm thickness, red		10	✓ Moments
		11.64	144	✓ Moments
	 a) Up to 60 dev available on a PSO 			
	 b) Maximum of 60 dev per prescription 			
*	56 mm, chocolate	1.30	12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
*	56 mm, strawberry	1.30	12	✓ Gold Knight
		15.57	144	Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
*	60 mm - Up to 144 dev available on a PSO	14.87	144	✓ Shield XL

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

GENITO-URINARY SYSTEM

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

*	IUD 29.1 mm length × 23.2 mm width	18.45	1	✓ Choice TT380 Short
---	------------------------------------	-------	---	----------------------

***** IUD 33.6 mm length × 29.9 mm width18.45

TT380 Standard

Choice

✓ Choice Load 375

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- · on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

a) Higher subsidy of \$13.80 per 84 tab with Special Authority see \$A0500 above

b) Up to 84 tab available on a PSO

a) Higher subsidy of \$13.80 per 84 tab with Special Authority see \$A0500 above

b) Up to 84 tab available on a PSO

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	I Generic
	\$	Per	•	Manufacturer
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO	2.18	84	1	Microgynon 20 ED
'	6.45	112		Femme-Tab ED
* Tab 50 mcg with levonorgestrel 125 mcg and 7 inert tab - Up	0			
to 84 tab available on a PSO	9.45	84	1	Microgynon 50 ED
* Tab 30 mcg with levonorgestrel 150 mcg	6.62	63		•
	(16.50)			Microgynon 30
a) Higher subsidy of \$15.00 per 63 tab with Special Auth	nority see SA0500 on	the i	previous p	age
b) Up to 63 tab available on a PSO	, ,			
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO		84	1	Levien ED
op to 112 tab available on a 1 co	6.45	112		Femme-Tab ED
ETHINIVI OFOTO ADIOL MITH MODETH HOTEDONE	0.10		-	· ciiiiio rub Eb
ETHINYLOESTRADIOL WITH NORETHISTERONE				
Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to				
84 tab available on a PSO	6.95	84	✓	Brevinor 1/28
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - U	р			
to 84 tab available on a PSO	6.62	84	✓	Necon
			1	Norimin

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Manyelon

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

CHORGEOTILE			
Tab 30 mcg - Up to 84 tab available on a PSO	.16.50	84	✓ Microlut
	22.00	112	✓ Microlut
Subdermal implant (2 × 75 mg rods) – Up to 3 pack available			
on a PSO1	106.92	1	Jadelle
	Subdermal implant (2 \times 75 mg rods) – Up to 3 pack available	Tab 30 mcg - Up to 84 tab available on a PSO16.50 22.00	Tab 30 mcg $-$ Up to 84 tab available on a PSO

GENITO-URINARY SYSTEM

	Subsidy		Fully	Brand or	
	(Manufacturer's Price)	S	ubsidised	Generic	
	\$	Per	✓	Manufacturer	
MEDROXYPROGESTERONE ACETATE Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a P NORETHISTERONE Tab 350 mcg – Up to 84 tab available on a PSO		1 84	_	epo-Provera oriday 28	
Emergency Contraceptives					
LEVONORGESTREL * Tab 1.5 mg	4.95	1	✓ P	ostinor-1	
a) Maximum of 2 tab per prescription					

- b) Up to 5 tab available on a PSO
- c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

* Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs - Up ✓ Ginet to 168 tab available on a PSO......4.67 168

Gynaecological Anti-infectives

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC AC Jelly with glacial acetic acid 0.94%, hydroxyguinoline sulphate	CID		
0.025%, glycerol 5% and ricinoleic acid 0.75% with applica	tor8.43 (24.00)	100 g OP	Aci-Jel
CLOTRIMAZOLE			
* Vaginal crm 1% with applicators	2.50	35 g OP	✓ Clomazol
* Vaginal crm 2% with applicators	3.00	20 g OP	✓ Clomazol
MICONAZOLE NITRATE			
* Vaginal crm 2% with applicator	6.89	40 a OP	✓ Micreme
Micreme to be Sole Supply on 1 November 2020		3 -	
NYSTATIN			
Vaginal crm 100,000 u per 5 g with applicator(s)	4.00	75 g OP	✓ Nilstat
Nilstat to be Sole Supply on 1 October 2020		3	

Myometrial and Vaginal Hormone Preparations

DOMETHINE MALLATE			
	105.00	5	✓ DBL Ergometrine
STRIOL			3
	6.62	15 g OP	✓ Ovestin
Ovestin to be Sole Supply on 1 October 2020			
Pessaries 500 mcg	6.86	15	Ovestin
Ovestin to be Sole Supply on 1 October 2020			
	STRIOL Crm 1 mg per g with applicator Ovestin to be Sole Supply on 1 October 2020 Pessaries 500 mcg	Inj 500 mcg per ml, 1 ml ampoule — Up to 5 inj available on a PSO	Inj 500 mcg per ml, 1 ml ampoule — Up to 5 inj available on a PSO

EDCOMETDINE MALEATE

✓ Tamsulosin-Rex

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	Brand or Generic Manufacturer
DXYTOCIN – Up to 5 inj available on a PSO				
Inj 5 iu per ml, 1 ml ampoule	3.98	5	✓	Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule	4.98	5	1	Oxytocin BNM
DXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj avail	able on a PSO			
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml		5	✓	<u>Syntometrine</u>

Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 106

5-Alpha Reductase Inhibitors

FINASTERIDE - Special Authority see SA0928 below - Retail pharmacy

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Fither:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 Patient has symptomatic benign prostatic hyperplasia; and

2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

OXYBUTYNIN		
* Tab 5 mg11.70	500	✓ Apo-Oxybutynin
* Oral liq 5 mg per 5 ml	473 ml	✓ Apo-Oxybutynin
POTASSIUM CITRATE		
Oral liq 3 mmol per ml - Special Authority see SA1083 on the		
next page – Retail pharmacy31.80	200 ml OP	✓ Biomed

GENITO-URINARY SYSTEM

Subsidy	F	ully	Brand or	
(Manufacturer's Pr	rice) Subsid	ised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SOL	DIUM	CITRO)-TART	RATE	
	O				

* Grans eff 4 g sachets	2.22	28	✓ Ural
Ural to be Sole Supply on 1 October 2020			
SOLIFENACIN SUCCINATE			
Tab 5 mg	3.00	30	 Solifenacin Mylan
Tab 10 mg	5.50	30	✓ Solifenacin Mylan

Detection of Substances in Urine

	TΗ				

٥.	THO TOLIBITE			
*	Compound diagnostic sticks	7.50	50 test OP	
		(8.25)		Hemastix
TE	TRABROMOPHENOL			
*	Blue diagnostic strips	7.02	100 test OP	

Obstetric Preparations

Antiprogesterones

MIFEPRISTONE

Subsidised on a PSO only if from a Family Planning New Zealand Clinic or an abortion service provider with a DHB contract and the PSO is endorsed with the name of the institution for which the PSO is required.

(13.92)

- a) Up to 15 tab available on a PSO
- b) Only on a PSO

Albustix

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	/	Manufacturer	

Calcium Homeostasis

CA	וי) ו	11(1	IN	IN

CINACALCET - Special Authority see SA1618 below - Retail pharmacy

Tab 30 mg − Wastage claimable......210.30 28 ✓ Sensipar

⇒SA1618 Special Authority for Subsidy

Initial application only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial − Special Authority see SA1687 below −
Retail pharmacy......38.03 1

✓ Zoledronic acid
Mylan

⇒SA1687 Special Authority for Subsidy

Initial application — **(bone metastases)** only from an oncologist, haematologist or palliative care specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
 - 2.1 Patient has bone metastases or involvement; and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
 - 3.1 Patient has bone metastases or involvement; and
 - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer) only from an oncologist or medical practitioner on the recommendation of a oncologist. Approvals valid for 2 years for applications meeting the following criteria:
All of the following:

Subsid	dy Full	/ Brand or
(Manufacture	r's Price) Subsidise	d Generic
\$	Per 💌	Manufacturer

continued...

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.

Corticosteroids and Related Agents for Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE

* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	5	Celestone
		Chronodose
BEXAMETHASONE ★ Tab 0.5 mg − Up to 60 tab available on a PSO	30 30 25 ml OP	✓ <u>Dexmethsone</u> ✓ <u>Dexmethsone</u> ✓ Biomed
DEXAMETHASONE PHOSPHATE		
Dexamethasone phosphate injection will not be funded for oral use.		
* Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO9.25	10	✓ <u>Dexamethasone</u> <u>Phosphate</u> <u>Panpharma</u>
* Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO16.37	10	✓ <u>Dexamethasone</u> <u>Phosphate</u> <u>Panpharma</u>
FLUDROCORTISONE ACETATE		
* Tab 100 mcg14.32	100	✓ Florinef
HYDROCORTISONE		
* Tab 5 mg8.10	100	✓ <u>Douglas</u>
* Tab 20 mg	100	✓ <u>Douglas</u>
 Inj 100 mg vial	1	✓ Solu-Cortef
METHYLPREDNISOLONE		
* Tab 4 mg112.00	100	✓ <u>Medrol</u>
* Tab 100 mg194.00	20	✓ <u>Medrol</u>
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)		
Inj 40 mg vial18.90	1	✓ <u>Solu-Medrol-Act-</u> <u>O-Vial</u>
Inj 125 mg vial28.90	1	✓ <u>Solu-Medrol-Act-O-Vial</u>
Inj 500 mg vial22.78	1	✓ <u>Solu-Medrol-Act-O-Vial</u>
Inj 1 g vial27.83	1	✓ Solu-Medrol
METHYLPREDNISOLONE ACETATE	'	- Join-Michiol
Inj 40 mg per ml, 1 ml vial44.40	5	✓ Depo-Medrol
PREDNISOLONE	3	- Depo-medioi
* Oral liq 5 mg per ml – Up to 30 ml available on a PSO	30 ml OP	✓ <u>Redipred</u>

	Subsidy		Fully	
	(Manufacturer's Price)	_	Subsidised	
	\$	Per		Manufacturer
PREDNISONE				
* Tab 1 mg	10.68	500	1	Apo-Prednisone
* Tab 2.5 mg		500	1	Apo-Prednisone
* Tab 5 mg - Up to 30 tab available on a PSO		500	1	Apo-Prednisone
* Tab 20 mg - Up to 30 tab available on a PSO	29.03	500	✓	Apo-Prednisone
TETRACOSACTRIN				
* Inj 250 mcg per ml, 1 ml ampoule	75.00	1	1	UK Synacthen S29
, , · · · · g				AU Synacthen
				Synacthen
* Inj 1 mg per ml, 1 ml ampoule	690.00	1	✓	Synacthen Depot
, 01			1	Synacthene
				Retard S29
TRIAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule	20.80	5	1	Kenacort-A 10
	26.62		1	Adcortyl S29
Kenacort-A 10 to be Sole Supply on 1 November 2020				,
Inj 40 mg per ml, 1 ml ampoule	11.30	1	1	Triaver \$29
, , , ,	51.10	5	1	Kenacort-A 40
	70.62		1	Kenalog \$29
Kenacort-A 40 to be Sole Supply on 1 November 2020				

Sex Hormones Non Contraceptive

Androgen Agonists and Antagonists

CVDDOTEDONE ACETATE

CYPROTERONE AGETATE			
Tab 50 mg	13.17	50	✓ Siterone
Tab 100 mg	26.75	50	✓ Siterone
TESTOSTERONE			
Patch 5 mg per day	90.00	30	✓ Androderm
TESTOSTERONE CIPIONATE			
Inj 100 mg per ml, 10 ml vial	76.50	1	✓ Depo-Testosterone
TESTOSTERONE ESTERS			•
Inj 250 mg per ml, 1 ml	12.98	1	✓ Sustanon Ampoules
TESTOSTERONE UNDECANOATE			,
Cap 40 mg	21.00	60	✓ Andriol Testocaps
Inj 250 mg per ml, 4 ml vial		1	✓ Reandron 1000
, ===,			

Hormone Replacement Therapy - Systemic

Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

		Subsidy (Manufacturer's Pr		Fully sidised	Brand or Generic
_		\$	Per		Manufacturer
0	estrogens				
	STRADIOL – See prescribing guideline on the previous page				
*	Tab 1 mg		28 OP	_	'atuafaua
*	Tab 2 mg	(11.10)	28 OP		strofem
~	Tab 2 mg	(11.10)	20 01	F	Strofem
*	Patch 100 mcg per 24 hours	, ,	4		limara
	a) No more than 1 patch per week				
	b) Only on a prescription				
*	Patch 50 mcg per 24 hours	7.04	4	✓ (Climara
	a) No more than 1 patch per week				
	b) Only on a prescription		_	, .	
	Patch 25 mcg per day	6.12	8	✓ E	stradot
	a) No more than 2 patch per week				
	b) Only on a prescription Patch 50 mcg per day	7.04	8	√	stradot 50 mcg
	a) No more than 2 patch per week	7.04	O	•	Strauot 50 mcg
	b) Only on a prescription				
	Patch 75 mcg per day	7.91	8	√ E	stradot
	a) No more than 2 patch per week				
	b) Only on a prescription				
	Patch 100 mcg per day	7.91	8	√ E	stradot
	a) No more than 2 patch per week				
	b) Only on a prescription				
	STRADIOL VALERATE - See prescribing guideline on the pr				
	Tab 1 mg		84		rogynova
*	Tab 2 mg	12.36	84	✓ <u>F</u>	Progynova
	STROGENS - See prescribing guideline on the previous page				
*	Conjugated, equine tab 300 mcg		28	_	
×	Conjugated aguing tab 605 mag	(13.50)	28	۲	remarin
*	Conjugated, equine tab 625 mcg	(13.50)	20	p	Premarin
		(10.50)		'	Temami
P	rogestogens				
ИE	DROXYPROGESTERONE ACETATE - See prescribing guid	eline on the prev	vious page		
*	Tab 2.5 mg	3.75	30		rovera
	Tab 5 mg		100	_	rovera
*	Tab 10 mg	7.15	30	✓ P	rovera
P	rogestogen and Oestrogen Combined Prepara	tions			
	STRADIOL WITH NORETHISTERONE - See prescribing gui		vious page		
	Tab 1 mg with 0.5 mg norethisterone acetate	5.40	28 OP		
٠		(18.10)		K	(liovance
*	Tab 2 mg with 1 mg norethisterone acetate		28 OP		Warrant.
	Tab O man with down manathiates are a sale to (40) as 10	(18.10)		K	liogest
*	Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	E 40	20 OD		
	oestradiol tab (12) and 1 mg oestradiol tab (6)	5.40	28 OP	т	risequens
		(10.10)			nooquono

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Other Oestrogen Preparations				
ETHINYLOESTRADIOL * Tab 10 mcg	17.60	100	✓ <u>N</u>	IZ Medical and Scientific
OESTRIOL * Tab 2 mg	7.00	30	√ <u>(</u>	Ovestin
Other Progestogen Preparations				
LEVONORGESTREL * Intra-uterine device 52 mg * Intra-uterine device 13.5 mg MEDROXYPROGESTERONE ACETATE		1		<u>Airena</u> laydess
Tab 100 mg	101.00	100	√ F	Provera HD
NORETHISTERONE * Tab 5 mg – Up to 30 tab available on a PSO PROGESTERONE	18.29	100	√ <u>F</u>	Primolut N
Cap 100 mg - Special Authority see SA1609 below - Retail pharmacy	16.50	30	√ (Jtrogestan

SA1609 Special Authority for Subsidy

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 For the prevention of pre-term labour*; and
- 2 Either:
 - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 For the prevention of pre-term labour*: and
- 2 Treatment is required for second or subsequent pregnancy; and
- 3 Fither:
 - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are unapproved indications.

Thyroid and Antithyroid Agents

CARBIMAZOLE		
* Tab 5 mg10.80	100	✓ AFT
		Carbimazole S29
		✓ Neo-Mercazole

	Subsidy		Fully	Brand or
	(Manufacturer's Price	e)	Subsidised	Generic
	\$	Per	•	Manufacturer
LEVOTHYROXINE				
* Tab 25 mcg	3.89	90	✓	Synthroid
* Tab 50 mcg		28	✓	Mercury Pharma
·	4.05	90	✓	Synthroid
	64.28	1,000	✓	Eltroxin
* Tab 100 mcg	1.78	28	✓	Mercury Pharma
•	4.21	90	✓	Synthroid
	66.78	1,000	✓	Eltroxin
PROPYLTHIOURACIL – Special Authority see SA1199 belo Propylthiouracil is not recommended for patients under the treatments are contraindicated.		ss the p	atient is p	regnant and other
Tab 50 mg	35.00	100	1	PTU S29

⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) - Special Authority see SA1629 below - Re	etail pharmacy	
*	Inj 5 mg cartridge34.8	38 1	Omnitrope
*	Inj 10 mg cartridge69.7	75 1	✓ Omnitrope
*	Inj 15 mg cartridge104.6	63 1	✓ Omnitrope

⇒SA1629 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and</p>
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — **(short stature without growth hormone deficiency)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	

continued...

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — **(Prader-Willi syndrome)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria;
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight eximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy (Manufacturer's Pri	ce)	Fully Subsidised	Brand or Generic	
\$	Per	•	Manufacturer	

continued...

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
GnRH Analogues				
GOSERELIN Implant 3.6 mg, syringe Implant 10.8 mg, syringe		1	-	Zoladex Zoladex
LEUPRORELIN Additional subsidy by endorsement where the patient is a chi goserelin and the prescription is endorsed accordingly. Inj 3.75 mg prefilled dual chamber syringe — Higher subsidy	ld or adolescent and i	s un	able to toler	ate administration of
\$221.60 per 1 inj with Endorsement	66.48 (221.60)	1	L	_ucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe – Higher subsidy of \$591.68 per 1 inj with Endorsement		1	l	Lucrin Depot 3-month

Vasopressin Agonists

DESMOPRESSIN	∧ ∩ ET ∧ T E
DESIMORRESSIN	AUFIAIF

	Tab 100 mcg - Special Authority see SA1401 below - Retail pharmacy	25.00	30	✓ Minirin
^	Tab 200 mcg - Special Authority see SA1401 below - Retail pharmacy	54.45 39.03	30 2.5 ml OP 6 ml OP	✓ Minirin ✓ Minirin ✓ Desmopressin- PH&T
	Desmopressin-PH&T to be Sole Supply on 1 November 2020 Inj 4 mcg per ml, 1 ml – Special Authority see SA1401 below – Retail pharmacy		10	✓ Minirin

⇒SA1401 Special Authority for Subsidy

Initial application — (Desmopressin tablets for Nocturnal enuresis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has primary nocturnal enuresis; and
- 2 The nasal forms of desmopressin are contraindicated; and
- 3 An enuresis alarm is contraindicated.

Initial application — (Desmopressin tablets for Diabetes insipidus) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has cranial diabetes insipidus; and
- 2 The nasal forms of desmopressin are contraindicated.

Renewal — (Desmopressin tablets) from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from the treatment.

Initial application — (Desmopressin injection) only from a relevant specialist. Approvals valid for 2 years where the patient cannot use desmopressin nasal spray or nasal drops.

Renewal — (Desmopressin injection) only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Other Endocrine Agents

CABERGOLINE

⇒SA1370 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 pathological hyperprolactinemia; or
- 2 acromegaly*.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

CLC	MIF	ENE	CITRA	ATE.

Tab 50 mg29).84	10	Mylan Clomiphen S29
DANAZOL			
Cap 100 mg19	.13 2	28	Mylan S29
Cap 200 mg97	'.83 1	00	Azol
METYRAPONE			
Cap 250 mg558	3.00	50	Metopirone
Metopirone to be Sole Supply on 1 November 2020			•

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

Anthelmintics

ALBENDAZOLE – Special Authority see SA131	8 below – Retail pharmacy		
Tab 400 mg	469.20	60	✓ Eskazole S29

⇒SA1318 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

MEBENDAZOLE - Only on a prescription

Tab 100 mg	24.19	24	De-Worm
Oral lig 100 mg per 5 ml	2.18	15 ml	
,	(7.17)		Vermox
PRAZIQUANTEL			
Tab 600 mg	68.00	8	✓ Biltricide

Antibacterials

- a) For topical antibacterials, refer to DERMATOLOGICALS, page 61
- b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 240

Cephalosporins and Cephamycins

CEFACLOR MONOHYDRATE			
Cap 250 mg	24.70	100	✓ Ranbaxy-Cefaclor
Grans for oral liq 125 mg per 5 ml - Wastage claimable	3.53	100 ml	✓ Ranbaxy-Cefaclor
	4.33		✓ Keflor
CEFALEXIN			
Cap 250 mg	3.33	20	✓ Cephalexin ABM
			✓ Ibilex S29
Cap 500 mg	3.95	20	✓ Cephalexin ABM
Grans for oral liq 25 mg per ml - Wastage claimable		100 ml	✓ Cefalexin Sandoz
Grans for oral lig 50 mg per ml - Wastage claimable		100 ml	✓ Cefalexin Sandoz
CEFAZOLIN — Subsidy by endorsement Only if prescribed for dialysis or cellulitis in accordance with accordingly. Inj 500 mg vial	3.39	protocol and to	the prescription is endorsed AFT AFT
AFT to be Sole Supply on 1 November 2020			
CEFTRIAXONE – Subsidy by endorsement a) Up to 10 inj available on a PSO b) Subsidised only if prescribed for a dialysis or cystic fibro pelvic inflammatory disease, or the treatment of suspect endorsed accordingly.			•
Inj 500 mg vial		1	✓ <u>Ceftriaxone-AFT</u>
Inj 1 g vial	3.99	5	✓ Ceftriaxone-AFT

Only if prescribed for prophylaxis of endocarditis and the prescription is endorsed accordingly.

Tab 250 mg45.93

CEFUROXIME AXETIL - Subsidy by endorsement

✓ Zinnat

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	sidised	Generic
\$	Per	1	Manufacturer

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO	2	✓ Apo-Azithromycin
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage		
claimable	15 ml	✓ Zithromax

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Fither:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

Subsidy		Fully	Brand or	_
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

continued...

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN (AS LACTOBIONATE) Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	E-Mycin
a) Up to 20 tab available on a PSOb) Up to 2 x the maximum PSO quantity for RFPP			4
Grans for oral liq 200 mg per 5 ml		100 ml	✓ E-Mycin
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	✓ E-Mycin
ERYTHROMYCIN STEARATE			
Tab 250 mg - Up to 30 tab available on a PSO		100	
	(22.29)		ERA
Tab 500 mg		100	
	(44.58)		ERA
ROXITHROMYCIN			
Tab disp 50 mgRestricted to children under 12 years of age.	8.29	10	✓ Rulide D
Tab 150 mg	8.28	50	✓ <u>Arrow-</u> <u>Roxithromycin</u>
Tab 300 mg	16.33	50	✓ <u>Arrow-</u> <u>Roxithromycin</u>

Tab 500 mg with clavulanic acid 125 mg − Up to 30 tab available on a PSO					_
Penicillins AMOXICILLIN Cap 250 mg			, .	,	
Penicillins AMOXICILLIN Cap 250 mg		· .			
AMOXICILLIN Cap 250 mg		Ψ	101	- Manadadaroi	_
Cap 250 mg	Penicillins				
a) Up to 30 cap available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP Cap 500 mg	AMOXICILLIN				
b) Up to 10 x the maximum PSO quantity for RFPP Cap 500 mg	Cap 250 mg	22.50	500	✓ Alphamox	
Cap 500 mg	a) Up to 30 cap available on a PSO				
a) Up to 30 cap available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml	b) Up to 10 x the maximum PSO quantity for RFPP				
b) Up to 10 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml	Cap 500 mg	36.98	500	✓ <u>Alphamox</u>	
Grans for oral liq 125 mg per 5 ml					
a) Up to 200 ml available on a PSO b) Wastage claimable c) Alphamox 125 to be Sole Supply on 1 November 2020 Grans for oral liq 250 mg per 5 ml	b) Up to 10 x the maximum PSO quantity for RFPP				
b) Wastage claimable c) Alphamox 125 to be Sole Supply on 1 November 2020 Grans for oral liq 250 mg per 5 ml	Grans for oral liq 125 mg per 5 ml	1.40	100 ml	Alphamox 125	
c) Alphamox 125 to be Sole Supply on 1 November 2020 Grans for oral liq 250 mg per 5 ml	a) Up to 200 ml available on a PSO				
Grans for oral liq 250 mg per 5 ml					
a) Up to 300 ml available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP c) Wastage claimable d) Alphamox 250 to be Sole Supply on 1 November 2020 lnj 250 mg vial	c) Alphamox 125 to be Sole Supply on 1 November 202	20			
b) Up to 10 x the maximum PSO quantity for RFPP c) Wastage claimable d) Alphamox 250 to be Sole Supply on 1 November 2020 Inj 250 mg vial	Grans for oral liq 250 mg per 5 ml	1.73	100 ml	Alphamox 250	
c) Wastage claimable d) Alphamox 250 to be Sole Supply on 1 November 2020 Inj 250 mg vial	 a) Up to 300 ml available on a PSO 				
d) Alphamox 250 to be Sole Supply on 1 November 2020 Inj 250 mg vial					
Inj 250 mg vial	, 0				
Inj 500 mg vial					
Inj 1 g vial − Up to 5 inj available on a PSO	, 0				
AMOXICILLIN WITH CLAVULANIC ACID Tab 500 mg with clavulanic acid 125 mg − Up to 30 tab available on a PSO	, ,				
Tab 500 mg with clavulanic acid 125 mg − Up to 30 tab available on a PSO	Inj 1 g vial – Up to 5 inj available on a PSO	17.29	10	✓ Ibiamox	
available on a PSO	AMOXICILLIN WITH CLAVULANIC ACID				
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 mg per ml	Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab				
per ml			20	Augmentin	
a) Up to 200 ml available on a PSO b) Wastage claimable Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 mg per ml − Up to 200 ml available on a PSO2.20 100 ml OP ✓ Curam BENZATHINE BENZYLPENICILLIN					
b) Wastage claimable Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 mg per ml − Up to 200 ml available on a PSO2.20 100 ml OP ✓ Curam BENZATHINE BENZYLPENICILLIN	per ml	5.00	100 ml	Augmentin	
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 mg per ml − Up to 200 ml available on a PSO2.20 100 ml OP ✓ Curam BENZATHINE BENZYLPENICILLIN	, .				
per ml − Up to 200 ml available on a PSO2.20 100 ml OP ✓ Curam BENZATHINE BENZYLPENICILLIN	b) Wastage claimable				
BENZATHINE BENZYLPENICILLIN	, ,	•			
	per ml – Up to 200 ml available on a PSO	2.20	100 ml OF	Curam	
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj	BENZATHINE BENZYLPENICILLIN				
	Inj 900 mg (1.2 million units) in 2.3 ml syringe - Up to 5 inj				
available on a PSO	available on a PSO	344.93	10	✓ Bicillin LA	
BENZYLPENICILLIN SODIUM [PENICILLIN G]	BENZYLPENICILLIN SODIUM [PENICILLIN G]				
Inj 600 mg (1 million units) vial − Up to 5 inj available on a PSO11.09		SO 11.09	10	✓ Sandoz	
25.88 25 ✓ Pan-Penicillin G	, 3(1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		25	✓ Pan-Penicillin G	
Sodium S29				Sodium S29	

Sandoz to be Sole Supply on 1 November 2020

(Pan-Penicillin G Sodium S29 Inj 600 mg (1 million units) vial to be delisted 1 November 2020)

	Subsidy (Manufacturer's Price \$		ully Brand or sed Generic Manufacturer
FLUCLOXACILLIN	Ψ	1 01	Waltalactarci
Cap 250 mg - Up to 30 cap available on a PSO	16.83	250	✓ Staphlex
Cap 500 mg - Up to 30 cap available on a PSO			✓ Staphlex
Grans for oral liq 25 mg per ml			✓ AFT
a) Up to 200 ml available on a PSO		100 1111	- <u>/</u>
b) Wastage claimable			
Grans for oral liq 50 mg per ml	3.68	100 ml	✓ AFT
a) Up to 200 ml available on a PSO		100 1111	- <u>/ /</u>
b) Wastage claimable			
Inj 250 mg vial	9.00	10	✓ Flucloxin
Inj 500 mg vial			✓ Flucioxin
Inj 1 g vial – Up to 5 inj available on a PSO	5.70	. •	✓ Flucil
Flucil to be Sole Supply on 1 November 2020		3	· I Iucii

PHENOXYMETHYLPENICILLIN (PENICILLIN V)	0.50	F0	✓ Ollinaina VIV
Cap 250 mg - Up to 30 cap available on a PSO			✓ <u>Cilicaine VK</u>
Cap 500 mg	4.26	50	✓ Cilicaine VK
a) Up to 20 cap available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 125 mg per 5 ml	2.99	100 ml	✓ <u>AFT</u>
a) Up to 200 ml available on a PSO			
b) Wastage claimable	0.00	400	.
Grans for oral liq 250 mg per 5 ml	3.99	100 ml	✓ <u>AFT</u>
a) Up to 300 ml available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
PROCAINE PENICILLIN			
Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO	123.50	5	✓ Cilicaine
Tetracyclines			
OOXYCYCLINE			
★ Tab 100 mg – Up to 30 tab available on a PSO	64.43	500	✓ Doxine
IINOCYCLINE HYDROCHLORIDE			
Tab 50 mg - Additional subsidy by Special Authority see			
SA1355 below – Retail pharmacy	5.79	60	
CATIOGO DOION TIOIGII PHAITHAU	(12.05)	00	Mino-tabs
		100	mino tabo
F Cup 100 mg	(52.04)	100	Minomycin
CA10EE Chasial Authority for Manufacturers Briss	(02.04)		······oiiiyoiii
⇒SA1355 Special Authority for Manufacturers Price	d without fouthouse	awal unlass =	stified where the noticet be
nitial application from any relevant practitioner. Approvals valid	u williout iurtrier ren	ewai uilless N	ninea where the patient ha
OSACEA.	Dotoil pho		
ETRACYCLINE – Special Authority see SA1332 on the next pa	•	•	/ A
Tab 250 mg			Accord \$29
Cap 500 mg	46.00	30	✓ Tetracyclin
			Wolff S29

28

✓ Cinflox

Subsidy	F	ully	Brand or
(Manufacturer's Price	e) Subsidis	sed	Generic
\$	Per	1	Manufacturer

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 61

CIPROFLOXACIN

Recommended for patients with any of the following:

i) microbiologically confirmed and clinically significant pseudomonas infection; or

- ii) prostatitis: or
- iii) pyelonephritis; or
- iv) gonorrhoea.

2.42	20	Cipliox
3.40	28	✓ Cipflox
5.95	28	✓ Cipflox
4.61	24	✓ Dalacin C
39.00	10	✓ Dalacin C
Subsidy by endorse	ment	
ne prescription is en	dorsed acc	ordingly.
65.00	1	✓ Colistin-Link
or complicated urin	ary tract inf	fection and the prescription is
182.00	10	✓ Teligent S29
or complicated urin	ary tract inf	fection and the prescription is
17.50	10	✓ Pfizer
87.50	50	✓ Pfizer
or complicated urin	ary tract inf	fection and the prescription is
il pharmacy		
•		
52.00	5	✓ Avelox

⇒SA1740 Special Authority for Subsidy

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

1 Both:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1.1 Active tuberculosis*; and
- 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Either:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy

⇒SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Fither:

- 1 Patient has confirmed cryptosporidium infection; or
 - 2 For the eradication of Entamoeba histolyica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharmacy

⇒SA1328 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

ıı,	NFECTIONS - A	JEN	13 701	1 SYSTEMIC USE
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
SODIUM FUSIDATE [FUSIDIC ACID]				
Tab 250 mg		12	•	Fucidin
SULFADIAZINE SODIUM – Special Authority see SA1331 below			_	
Tab 500 mg	543.20	56	/	Wockhardt S29
Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months	r a period of 3 months		nless notif	ied for applications meeting
	or ago.			
TOBRAMYCIN Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement	15.00	5	_	Tobramycin Mylan
Only if prescribed for dialysis or cystic fibrosis patient an Solution for inhalation 60 mg per ml, 5 ml – Subsidy by				
endorsement	2,200.00 5	6 dos	se 🗸	TOBI
a) Wastage claimable				
b) Only if prescribed for a cystic fibrosis patient and the	prescription is endors	sed a	ccordingly	
TRIMETHOPRIM				
* Tab 300 mg - Up to 30 tab available on a PSO	16.50	50	/	<u>TMP</u>
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX	AZOLE]			
* Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - U	Jp			
to 30 tab available on a PSO		500	/	Trisul
* Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to 200 u available on a PSO		100 m	ıl 🗸	Deprim
VANCOMYCIN – Subsidy by endorsement				
Only if prescribed for a dialysis or cystic fibrosis patient or for			is or for tre	eatment of Clostridium
difficile following metronidazole failure and the prescription is	•	•		
Inj 500 mg vial	2.35	1	•	Mylan
Mylan to be Sole Supply on 1 October 2020				
Antifungals				
a) For topical antifungals refer to DERMATOLOGICALS, page 67	2			
b) For topical antifungals refer to GENITO URINARY, page 74	-			
FLUCONAZOLE				
Cap 50 mg	2.75	28	1	Mylan
Mylan to be Sole Supply on 1 November 2020				•
Cap 150 mg	0.65	1	✓	Mylan
Mylan to be Sole Supply on 1 November 2020				
Cap 200 mg		28	•	Mylan
Powder for oral suspension 10 mg per ml — Special Authority		2E		Diffuson COC coc
see SA1359 on the next page – Retail pharmacy	34.56 98.50	35 m		Diflucan S29 S29 Diflucan
Wastage claimable	50.50		•	Dinavaii
. radiago dallilabio				

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOLE

Cap 100 mg	4.27	15	✓ <u>Itrazole</u>
Oral liq 10 mg per ml - Special Authority see SA1322 below -			
Retail pharmacy14	1.80	150 ml OP	✓ Sporanox

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE

Tab 200 mg - PCT	.CBS	30	✓ Link Healthcare S29
			✓ Nizoral S29
		100	✓ Strides Shasun S29
NYSTATIN			
Tab 500,000 u	.14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	.12.81	50	
	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 on the next page -	Retail phar	macy	
Tab modified-release 100 mg	369.86	24	✓ Noxafil
Oral liq 40 mg per ml7	761.13	105 ml OP	✓ Noxafil

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE

*	Tab 250 mg	1.33	14	✓ Deolate
۷O	RICONAZOLE - Special Authority see SA1273 below - Retail pharm	acy		
	Tab 50 mg9	1.00	56	✓ Vttack
	Tab 200 mg35	0.00	56	✓ Vttack
	Powder for oral suspension 40 mg per ml - Wastage			
	claimable1,43	7.00	70 ml	✓ Vfend

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis: or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

Antimalarials

PRIMAQUINE - Special Authority see SA1684 below - Retail pharmacy

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Antiparasitics

Antiprotozoals

QUININE SULP	PHATE
--------------	-------

Antitrichomonal Agents

METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	10.45	100	✓ Trichozole
• ,	36.35	250	✓ Metrogyl
Tab 400 mg - Up to 15 tab available on a PSO	5.55	21	✓ Metrogyl
, i	18.15	100	✓ Trichozole
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg		10	✓ Flagyl
(Trichozole Tab 200 mg to be delisted 1 September 2020) (Trichozole Tab 400 mg to be delisted 1 September 2020)			
ORNIDAZOLE			
Tab 500 mg	32.95	10	✓ Arrow-Ornidazole

Antituberculotics and Antileprotics

Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status.

CLOFAZIMINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist.

CYCLOSERINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician.

INFECTIONS - AGENTS FOR SYSTEMIC USE						
	Subsidy (Manufacturer's Price) \$	Su Per	Fully Brand or ubsidised Generic ✓ Manufacturer			
DAPSONE – Retail pharmacy-Specialist						
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat dermatologist Tab 25 mg		isease p	ohysician, clinical microbiologist or Dapsone			
Tab 100 mg		100	✓ Dapsone			
ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Specialis	st					
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician	ion of, an infectious di	isease p	physician, clinical microbiologist or			
Tab 100 mg	85.73	100	✓ EMB Fatol S29			
Tab 400 mg	49.34	56	✓ Myambutol S29			
ISONIAZID - Retail pharmacy-Specialist						
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat microbiologist, dermatologist or public health physician 	ion of, an internal med	dicine ph	hysician, paediatrician, clinical			
* Tab 100 mg	22.00	100	✓ <u>PSM</u>			
ISONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist						
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat microbiologist, dermatologist or public health physician 		dicine ph	hysician, paediatrician, clinical			
* Tab 100 mg with rifampicin 150 mg		100	Rifinah			
* Tab 150 mg with rifampicin 300 mg	170.60	100	✓ <u>Rifinah</u>			
PARA-AMINO SALICYLIC ACID - Retail pharmacy-Specialist						
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician 	ion of, an infectious di	isease s	specialist, clinical microbiologist or			
Grans for oral liq 4 g sachet	280.00	30	✓ Paser S29			
PROTIONAMIDE - Retail pharmacy-Specialist						
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician 		isease s	specialist, clinical microbiologist or			
Tab 250 mg	305.00	100	✓ Peteha S29			
PYRAZINAMIDE - Retail pharmacy-Specialist						
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician 		isease p	physician, clinical microbiologist or			
* Tab 500 mg	59.00	100	AFT-Pyrazinamide			
RIFABUTIN - Retail pharmacy-Specialist						
No patient co-payment payable	tour of an infant		ala antata a managanta a antanta d			
 b) Prescriptions must be written by, or on the recommendat gastroenterologist Cap 150 mg 		isease p				
本 Cap 130 IIIy	288.75	30	✓ Mycobutin			

	Subsidy (Manufacturer's Price) \$	Su Per	Fully ubsidised	Brand or Generic Manufacturer	
RIFAMPICIN – Subsidy by endorsement					

- a) No patient co-payment payable
- b) For confirmed recurrent Staphylococcus aureus infection in combination with other effective anti-staphylococcal antimicrobial based on susceptibilities and the prescription is endorsed accordingly; can be waived by endorsement -Retail pharmacy - Specialist. Specialist must be an internal medicine physician, clinical microbiologist, dermatologist, naediatrician, or nublic health physician

	paediatician, or public nealth physician.			
*	Cap 150 mg5	8.54	100	✓ Rifadin
	Rifadin to be Sole Supply on 1 November 2020			
*	Cap 300 mg12	2.06	100	Rifadin
	Rifadin to be Sole Supply on 1 November 2020			
*	Oral liq 100 mg per 5 ml1	2.60	60 ml	Rifadin
	Rifadin to be Sole Supply on 1 November 2020			

Antivirals

For eye preparations refer to Eye Preparations, Anti-Infective Preparations, page 240

Hepatitis B Treatment

ADEFOVIR DIPIVOXIL - Special Authority see \$A0829 below - Retail pharmacy 30 ✓ Hepsera Tab 10 mg670.00

⇒SA0829 Special Authority for Subsidy

Initial application only from a gastroenterologist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg+); and Documented resistance to lamivudine, defined as:
- 2 Patient has raised serum ALT (> 1 × ULN); and
- 3 Patient has HBV DNA greater than 100,000 copies per mL, or viral load 10 fold or higher over nadir; and
- 4 Detection of M204I or M204V mutation; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 Patient is cirrhotic: and
 - 5.1.2 adefovir dipivoxil to be used in combination with lamivudine; or
 - 5.2 Both:
 - 5.2.1 Patient is not cirrhotic; and
 - 5.2.2 adefovir dipivoxil to be used as monotherapy.

Renewal only from a gastroenterologist or infectious disease specialist. Approvals valid for 2 years where in the opinion of the treating physician, treatment remains appropriate and patient is benefiting from treatment.

Notes: Lamivudine should be added to adefovir dipivoxil if a patient develops documented resistance to adefovir dipivoxil. defined as:

- i) raised serum ALT (> 1 × ULN); and
- ii) HBV DNA greater than 100.000 copies per mL, or viral load 10 fold or higher over nadir; and
- iii) Detection of N236T or A181T/V mutation.

Adefovir dipivoxil should be stopped 6 months following HBeAg seroconversion for patients who were HBeAg+ prior to commencing adefovir dipivoxil.

The recommended dose of adefovir dipivoxil is no more than 10mg daily.

In patients with renal insufficiency adefovir dipivoxil dose should be reduced in accordance with the datasheet guidelines. Adefovir dipivoxil should be avoided in pregnant women and children.

FNTFCAVIR

30 ✓ Entecavir Sandoz

	Subsidy (Manufacturer's Prio \$	e) Per	Fully Subsidised	Generic
LAMIVUDINE – Special Authority see SA1685 below – Retail pha Tab 100 mg	•	28	1	Zetlam
Zetlam to be Sole Supply on 1 November 2020 Oral liq 5 mg per ml	270.00	240 ml	OP 🗸	Zeffix

⇒SA1685 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B. TENOFOVIR DISOPROXIL

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA1651., page 104

*	Tab 245 mg (300.6 mg as a succinate)		30	•	Tenofovir Disoproxil
					Teva

Herpesvirus Treatments			
ACICLOVIR			
* Tab dispersible 200 mg	1.60	25	✓ <u>Lovir</u>
* Tab dispersible 400 mg	5.38	56	✓ Lovir
* Tab dispersible 800 mg	5.98	35	✓ <u>Lovir</u>
VALACICLOVIR			
Tab 500 mg	5.75	30	✓ Vaclovir
Tab 1,000 mg	11.35	30	✓ Vaclovir
VALGANCICLOVIR - Special Authority see SA1404 below -	Retail pharmacy		
Tab 450 mg	225.00	60	✓ Valganciclovir
			Mylan

⇒SA1404 Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
- 2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months): and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Patient has undergone a lung transplant; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

- Both:
 - 1 Patient is immunocompromised; and
 - 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via PHARMAC's approved direct distribution supply. Further details can be found on

PHARMAC's website https://www.pharmac.govt.nz/hepatitis-c-treatments

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz/hepatitis-c-treatments or:

The Coordinator, Hepatitis C Treatment Panel

PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

Fully

Subsidy (Manufacturer's Price) Subsidised Per

Brand or Generic Manufacturer

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement: can be waived by Special Authority see SA1904

below Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabline with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 104 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the PHARMAC website.

Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a

Teva

⇒SA1904 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment: and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men: and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment: and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks: and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and

Subsidy	Full	/ Brand or
(Manufacturer's	s Price) Subsidise	d Generic
\$	Per 💌	Manufacturer

continued...

- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Antiretrovirals

⇒SA1651 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	ıbsidised	Generic
 \$	Per	/	Manufacturer

continued...

prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria: Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ – Special Authority see SA1651 o	n the previous page – Retail pharn	nacy	
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
ETRAVIRINE - Special Authority see SA1651	on the previous page - Retail phar	macy	
Tab 200 mg	770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA1651	on the previous page – Retail phar	macy	
Tab 200 mg	60.00	60	✓ <u>Nevirapine</u>
			<u>Alphapharm</u>
Oral suspension 10 mg per ml	203.55	240 ml	✓ Viramune
			Suspension

Nucleosides Reverse Transcriptase Inhibitors

ABACAVIR SULPHATE – Special Authority see SA1651 on the p Tab 300 mg Oral lig 20 mg per ml	180.00	Retail pharmad 60 240 ml OP	cy ✓ <u>Ziagen</u> ✓ Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE - Special Authority			• ,
Note: abacavir with lamivudine (combination tablets) counts a	as two anti-retro	oviral medicatio	ns for the purposes of the
anti-retroviral Special Authority.			
Tab 600 mg with lamivudine 300 mg	63.00	30	✓ Kivexa
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPR	OXII - Specia	I Authority see	SA1651 on the previous page
Retail pharmacy	Ortiz Opoolo		or moor on mo promode pay
Note: Efavirenz with emtricitabine and tenofovir disoproxil co	unte ae throp a	nti-ratroviral ma	dications for the nurnoses o
Note. Liaviferiz with emitherabile and teriologic disoproxil co	unio ao unice a	mi-renovitat me	dications for the purposes of

of the anti-retroviral Special Authority

Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil 245 mg (300 mg as a maleate)......106.88 30 Mvlan ige -

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Pric \$		Fully Brand or dised Generic Manufacturer
EMTRICITABINE – Special Authority see SA1651 on page 104 - Cap 200 mg		30	✓ <u>Emtriva</u>
LAMIVUDINE – Special Authority see SA1651 on page 104 – Re Tab 150 mg		60	✓ Lamivudine Alphapharm
Lamivudine Alphapharm to be Sole Supply on 1 Novembor Oral liq 10 mg per ml		240 ml OP	✓ 3TC
ZIDOVUDINE [AZT] – Special Authority see SA1651 on page 10 Cap 100 mg Oral liq 10 mg per ml	152.25	0y 100 200 ml OP	✓ Retrovir ✓ Retrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority see Note: zidovudine [AZT] with lamivudine (combination tablets the anti-retroviral Special Authority.			•
Tab 300 mg with lamivudine 150 mg	33.00	60	✓ Alphapharm
Protease Inhibitors			
ATAZANAVIR SULPHATE – Special Authority see SA1651 on p Cap 150 mg Cap 200 mg	141.68	harmacy 60 60	✓ <u>Teva</u> ✓ <u>Teva</u>
DARUNAVIR – Special Authority see SA1651 on page 104 – Re Tab 400 mg Tab 600 mg	335.00	60 60	✓ Prezista✓ Prezista
LOPINAVIR WITH RITONAVIR — Special Authority see SA1651 Tab 100 mg with ritonavir 25 mg Tab 200 mg with ritonavir 50 mg Oral liq 80 mg with ritonavir 20 mg per ml	183.75 463.00 735.00	tail pharmacy 60 120 300 ml OP	✓ Kaletra✓ Kaletra✓ Kaletra
RITONAVIR – Special Authority see SA1651 on page 104 – Ret Tab 100 mg		30	✓ <u>Norvir</u>
Strand Transfer Inhibitors			
DOLUTEGRAVIR – Special Authority see SA1651 on page 104 Tab 50 mg		30	✓ Tivicay
RALTEGRAVIR POTASSIUM - Special Authority see SA1651 o Tab 400 mg		il pharmacy 60	✓ Isentress

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

1) Diagnosis

bsidy	Fully	Brand or
turer's Price) Subsi	idised	Generic
 \$ Per	1	

continued...

- Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
- PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
- Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Exclusion Criteria

- Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia (< 2.0 × 10⁹) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

INTERFERON ALFA-2A - PCT

See prescribing guideline on the previous page

(Roferon-A Ini 3 m iu prefilled syringe to be delisted 1 December 2020)

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA1936 below - Retail pharmacy

- a) See prescribing guideline on the previous page
- b) Note: PHARMAC will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at PHARMAC on 0800-023-588 option 4.

⇒SA1936 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Fither:

Subsidy (Manufacturer's Price)	Ful Subsidise	,	
\$	Per •	Manufacturer	

continued...

- 3.1 Patient has responder relapsed; or
- 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease: and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegulated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

1 Patient has a cutaneous T cell lymphoma*; or

- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:

INFECTIONS - AGENTS FOR SYSTEMIC USE

Subsidy	Fu	ly Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per	 Manufacturer 	

continued...

- 3.1 Patient has a myeloproliferative disorder; and
- 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

Notes:

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet quidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

Urinary Tract Infections

METHENAMINE (HEXAMINE) HIPPURATE			
* Tab 1 g	40.01	100	✓ Hiprex
NITROFURANTOIN			
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
* Tab 100 mg	37.50	100	✓ Nifuran
NORFLOXACIN			
Tab 400 mg - Subsidy by endorsement	135.00	100	Arrow-Norfloxacin
Only if any and four and four and in the any constant of	all and a source to a state of a site of	a disart to consul-	and a section to a floor line and a second

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

	0		E. II.	Daniel au
	Subsidy (Manufacturaria Price	`	Fully	Brand or
	(Manufacturer's Price \$) Per	Subsidised	Generic Manufacturer
	ў	rei		Manuacturer
Autichalinastavassa				
Anticholinesterases				
NECOTIONINE METHOLILEATE				
NEOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	98.00	50	•	AstraZeneca
PYRIDOSTIGMINE BROMIDE				
▲ Tab 60 mg	45 79	100	1	Mestinon
		100		<u> </u>
Non-Steroidal Anti-Inflammatory Drugs				
Non-Steroidal Anti-Inflamiliatory Drugs				
DICLOFENAC SODIUM				
* Tab EC 25 mg	1 22	50	1	Diclofenac Sandoz
•		20		Voltaren D
· · · · · · · · · · · · · · · · · · ·				
* Tab EC 50 mg		50		Diclofenac Sandoz
* Tab long-acting 75 mg		500		Apo-Diclo SR
* Tab long-acting 100 mg		500		Apo-Diclo SR
* Inj 25 mg per ml, 3 ml ampoule - Up to 5 inj available on a	PSO 13.20	5	✓	Voltaren
* Suppos 12.5 mg		10	✓	Voltaren
* Suppos 25 mg		10		Voltaren
* Suppos 50 mg - Up to 10 supp available on a PSO		10		Voltaren
* Suppos 100 mg		10		Voltaren
	7.00	10	•	Voltaren
IBUPROFEN			_	
* Tab 200 mg	11.71	1,000		Relieve
Tab long-acting 800 mg	5.99	30	•	Ibuprofen SR BNM
	7.99		✓	Brufen SR
Ibuprofen SR BNM to be Sole Supply on 1 December 2	020			
* Oral liq 20 mg per ml		200 m	/	Ethics
(Brufen SR Tab long-acting 800 mg to be delisted 1 December 2			-	
	.020)			
KETOPROFEN				
* Cap long-acting 200 mg	12.07	28	•	Oruvail SR
MEFENAMIC ACID				
* Cap 250 mg	1 25	50		
σαρ 200 mg	(9.16)	00		Ponstan
	0.50	20		i Olistali
		20		Ponstan
	(5.60)			Ponsian
NAPROXEN				
* Tab 250 mg	32.69	500	✓	Noflam 250
* Tab 500 mg	22.19	250	✓	Noflam 500
* Tab long-acting 750 mg	6.16	28	/	Naprosyn SR 750
* Tab long-acting 1 g		28		Naprosyn SR 1000
5 5 5				
SULINDAC	0		_	A 11
* Tab 100 mg		50		Aclin
	9.57	56	✓	Mylan S29
* Tab 200 mg	15.10	50	✓	Aclin
Ť	16.91	56	1	Sulindac Mylan S29
(Aclin Tab 100 mg to be delisted 1 September 2020)	10.01	50	-	iwao myian
, , , , , , , , , , , , , , , , , , , ,				
TENOXICAM			_	
* Tab 20 mg	9.15	100	•	Tilcotil
* Inj 20 mg vial	9.95	1	✓	AFT

	Subsidy		Fully Brand or
	(Manufacturer's Pric	ce) Sub	sidised Generic Manufacturer
NSAIDs Other			
CELECOXIB	0.00	00	✓ Oolohuuu
Cap 100 mg	3.63	60	✓ Celebrex✓ Celecoxib Pfizer
Cap 200 mg	2.30	30	✓ Celebrex ✓ Celecoxib Pfizer
(Celebrex Cap 100 mg to be delisted 1 September 2020)			Celecoxib Plizer
Topical Products for Joint and Muscular Pain			
CAPSAICIN			
Crm 0.025% - Special Authority see SA1289 below - Retain			
pharmacy		25 g OP	✓ Zostrix
	9.95 13.27	45 g OP 60 g OP	✓ Zostrix✓ Rugby Capsaicin
	10.27	00 y O1	Topical Cream \$29
osteoarthritis that is not responsive to paracetamol and oral non- Antirheumatoid Agents	steroidal anti-inflan	nmatories ar	e contraindicated.
Subsidy by endorsement - Subsidised only if prescribed for malaria treatment or suppression, relevant dermatological creatment and mucosal ulceration)* and the prescription is prescription as endorsed where there exists a record of prior	onditions (cutaneou endorsed accordin	is forms of lungly. Pharma	pus and lichen planus, cutaned acists may annotate the
Subsidy by endorsement - Subsidised only if prescribed for malaria treatment or suppression, relevant dermatological creatment and mucosal ulceration)* and the prescription is prescription as endorsed where there exists a record of prio with a * is an unapproved indication.	onditions (cutaneou endorsed accordin r dispensing of hydi	is forms of lungly. Pharma	pus and lichen planus, cutaned acists may annotate the line. Note: Indication marked
malaria treatment or suppression, relevant dermatological or vasculitides and mucosal ulceration)* and the prescription is prescription as endorsed where there exists a record of prio with a * is an unapproved indication. * Tab 200 mg	onditions (cutaneou endorsed accordin r dispensing of hydi	is forms of lungly. Pharma	pus and lichen planus, cutaned acists may annotate the
Subsidy by endorsement - Subsidised only if prescribed for malaria treatment or suppression, relevant dermatological or vasculitides and mucosal ulceration)* and the prescription is prescription as endorsed where there exists a record of prio with a * is an unapproved indication. * Tab 200 mg	onditions (cutaneou endorsed accordin r dispensing of hydi 	is forms of lungly. Pharma	pus and lichen planus, cutaned acists may annotate the line. Note: Indication marked
Subsidy by endorsement - Subsidised only if prescribed for malaria treatment or suppression, relevant dermatological or vasculitides and mucosal ulceration)* and the prescription is prescription as endorsed where there exists a record of prio with a * is an unapproved indication. * Tab 200 mg	onditions (cutaneou endorsed accordin r dispensing of hydi 	is forms of lungly. Pharma roxychloroqu	pus and lichen planus, cutaned acists may annotate the line. Note: Indication marked Plaquenil
Subsidy by endorsement - Subsidised only if prescribed for malaria treatment or suppression, relevant dermatological or vasculitides and mucosal ulceration)* and the prescription is prescription as endorsed where there exists a record of prio with a * is an unapproved indication. * Tab 200 mg	onditions (cutaneous endorsed according dispensing of hydromasses) 2.90 6.00 2.90	is forms of lungly. Pharma roxychloroqu	pus and lichen planus, cutaner acists may annotate the ine. Note: Indication marked Plaquenil Apo-Leflunomide Arava Apo-Leflunomide
Subsidy by endorsement - Subsidised only if prescribed for malaria treatment or suppression, relevant dermatological or vasculitides and mucosal ulceration)* and the prescription is prescription as endorsed where there exists a record of prio with a * is an unapproved indication. * Tab 200 mg EFLUNOMIDE Tab 10 mg Tab 20 mg	enditions (cutaneous endorsed according dispensing of hydrometric from 1	us forms of luggly. Pharma roxychloroqu 100 30	pus and lichen planus, cutaned acists may annotate the line. Note: Indication marked Plaquenil Apo-Leflunomide Arava
Subsidy by endorsement - Subsidised only if prescribed for malaria treatment or suppression, relevant dermatological or vasculitides and mucosal ulceration)* and the prescription is prescription as endorsed where there exists a record of prio with a * is an unapproved indication. * Tab 200 mg LEFLUNOMIDE Tab 10 mg Tab 20 mg	onditions (cutaneous endorsed according dispensing of hydromasses) 2.90 6.00 2.90	us forms of luggly. Pharma roxychloroqu 100 30	pus and lichen planus, cutaned acists may annotate the line. Note: Indication marked Plaquenil Apo-Leflunomide Arava Apo-Leflunomide
Subsidy by endorsement - Subsidised only if prescribed for malaria treatment or suppression, relevant dermatological or vasculitides and mucosal ulceration)* and the prescription is prescription as endorsed where there exists a record of prio with a * is an unapproved indication. * Tab 200 mg	endorsed according dispensing of hydrometric dispensing of hydrometric dispensing of hydrometric dispensing of hydrometric dispension of hydrometric	us forms of luggly. Pharma roxychloroqu 100 30	pus and lichen planus, cutaned acists may annotate the line. Note: Indication marked Plaquenil Apo-Leflunomide Arava Apo-Leflunomide

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

AL	ENDRONATE SODIUM			
*	Tab 70 mg	4	4	✓ Fosamax
ALI	ENDRONATE SODIUM WITH COLECALCIFEROL			
*	Tab 70 mg with colecalciferol 5,600 iu	1	4	✓ Fosamax Plus

100

✓ D-Penamine

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy (Manufacturer's Price)

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

Other Treatments

DENOSUMAB – Special Authority see SA1777 below – Retail pharmacy Inj 60 mg prefilled syringe.......326.00

1 ✓ Prolia

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Either:
 - 2.1 The patient is female and postmenopausal; or
 - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	
PAMIDRONATE DISODIUM				
Inj 3 mg per ml, 10 ml vial	5.98	1	/	Pamisol
Inj 6 mg per ml, 10 ml vial	15.02	1	1	Pamisol
Inj 9 mg per ml, 10 ml vial		1	1	Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see S/ * Tab 60 mg		armacy 28	/	Evista

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM Tab 35 mg3.10	4	✓ Risedronate Sandoz
TERIPARATIDE – Special Authority see SA1139 below – Retail pharmacy		
Inj 250 mcg per ml, 2.4 ml490.00	1	✓ Forteo

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a

	Subsidy		Fully	Brand or
(M	anufacturer's Price)	Subs	idised	Generic
	\$	Per	✓	Manufacturer

continued...

funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, vial - Special Authority see 100 ml OP Aclasta

⇒SA1780 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1

Subsidy (Manufacturer's Price)	Sul	Fully	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note): or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined): and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Notes:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

ALL ODLIDINO

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

AL	LOPURINOL			
*	Tab 100 mg	11.47	500	✓ DP-Allopurinol
	DP-Allopurinol to be Sole Supply on 1 November 2020			•
*	Tab 300 mg	28.57	500	✓ DP-Allopurinol
	DP-Allopurinol to be Sole Supply on 1 November 2020			
BE	NZBROMARONE – Special Authority see SA1537 below – Reta	il pharmacy		
	Tab 50 mg	22.50	100	✓ Narcaricin mite S29
	Tab 100 mg	13.50	30	✓ Desuric S29
				✓ Urinorm S29
		45.00	100	Benzbromaron AL
				100 S29

⇒SA1537 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 Both:
 - 2.3.1 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Notes); and
 - 2.3.2 The patient has a rate of creatinine clearance greater than or equal to 20 ml/min; or
 - 2.4 All of the following:
 - 2.4.1 The patient is taking azathioprine and requires urate-lowering therapy; and
 - 2.4.2 Allopurinol is contraindicated; and
 - 2.4.3 Appropriate doses of probenecid are ineffective or probenecid cannot be used due to reduced renal function; and
- 3 The patient is receiving monthly liver function tests.

Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

Notes: Benzbromarone has been associated with potentially fatal hepatotoxicity.

In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

The New Zealand Rheumatology Association has developed information for prescribers which can be accessed from its website at www.rheumatology.org.nz/home/resources-2/

COLCHICINE

* Tab 500 mcg	9.58	100	✓ Colgout
FEBUXOSTAT - Special Authority see SA1931 below - Reta	ail pharmacy		
Tab 80 mg	39.50	28	✓ Adenuric
Tab 120 mg	39.50	28	Adenurio

⇒SA1931 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
 - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

PROBENECID

*	Tab 500 mg	55.00	100	1	Probenecid-AFT
---	------------	-------	-----	---	----------------

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Muscle Relaxants

masore relaxants			
BACLOFEN			
* Tab 10 mg		100	✓ Pacifen
Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement	11.55	1	Lioresal Intrathecal
Subsidised only for use in a programmable pump in patier			ents have been ineffective or have
caused intolerable side effects and the prescription is end-	orsed according	ly.	
Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement	372.98	5	✓ <u>Medsurge</u>
Subsidised only for use in a programmable pump in patier caused intolerable side effects and the prescription is end-			ents have been ineffective or have
DANTROLENE			
Cap 25 mg	97.50	100	✓ Dantrium
			✓ Dantrium S29 S29
Cap 50 mg	77.00	100	✓ Dantrium
ORPHENADRINE CITRATE			
Tab 100 mg	18 54	100	✓ Norflex

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Agents for Parkinsonism and Related Disorders

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE		
▲ Cap 100 mg38.24	60	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE		
▲ Inj 10 mg per ml, 2 ml ampoule59.50	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule121.84	5	✓ Movapo
BROMOCRIPTINE MESYLATE		
* Tab 2.5 mg32.08	100	✓ Apo-Bromocriptine
ENTACAPONE		
▲ Tab 200 mg22.00	100	✓ Entapone
LEVODOPA WITH BENSERAZIDE		
* Tab dispersible 50 mg with benserazide 12.5 mg	100	✓ Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg	100	✓ Madopar 62.5
* Cap 100 mg with benserazide 25 mg15.80	100	✓ Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg22.85	100	✓ Madopar HBS
* Cap 200 mg with benserazide 50 mg26.25	100	Madopar 250
LEVODOPA WITH CARBIDOPA		
* Tab 100 mg with carbidopa 25 mg17.97	100	✓ Kinson
		✓ Sinemet
* Tab long-acting 100 mg with carbidopa 25 mg23.84	100	✓ Mylan S29
* Tab long-acting 200 mg with carbidopa 50 mg37.15	100	✓ Sinemet CR
46.73		✓ Mylan S29
* Tab 250 mg with carbidopa 25 mg32.67	100	✓ Sinemet
PRAMIPEXOLE HYDROCHLORIDE		
▲ Tab 0.25 mg6.12	100	✓ Ramipex
▲ Tab 1 mg20.73	100	✓ Ramipex
ROPINIROLE HYDROCHLORIDE		
▲ Tab 0.25 mg2.85	84	✓ Ropin
3.39	100	✓ Mylan S29
▲ Tab 1 mg3.95	84	✓ Ropin
4.70	100	✓ Mylan S29
▲ Tab 2 mg5.48	84	Ropin
▲ Tab 5 mg12.50	84	✓ Ropin
SELEGILINE HYDROCHLORIDE		
* Tab 5 mg22.00	100	Apo-Selegiline
		S29 S29
TOLCAPONE		
▲ Tab 100 mg152.38	100	✓ Tasmar

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
Anticholinergics				
BENZATROPINE MESYLATE				
Tab 2 mg	7.99	60	1	Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	1	Cogentin
				Phebra
	190.00	10	•	Omega
 a) Up to 10 inj available on a PSO 				
b) Only on a PSO				
(Cogentin Inj 1 mg per ml, 2 ml to be delisted 1 December 2020)				
(Omega Inj 1 mg per ml, 2 ml to be delisted 1 December 2020)				
PROCYCLIDINE HYDROCHLORIDE				
Tab 5 mg	7.40	100	✓	Kemadrin
Agents for Essential Tremor, Chorea and Relate	ad Disorders			

Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE - Special Authority see SA1403 below - Retail pharmacy

Wastage claimable

⇒SA1403 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
 - 5.1 The patient is ambulatory; or
 - 5.2 The patient is able to use upper limbs; or
 - 5.3 The patient is able to swallow.

Renewal from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
 - 3.1 The patient is ambulatory; or
 - 3.2 The patient is able to use upper limbs; or
 - 3.3 The patient is able to swallow.

TETRABENAZINE

Tab 25 mg91.10 112 ✓ Motetis

Subsidy		Fully	Brand or	_
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

Anaesthetics

Local

LIDOCAINE [LIGNOCAINE]			
Gel 2%, tube - Subsidy by endorsement	14.50	30 ml	Xylocaine 2% Jelly
a) Up to 150 ml available on a PSO			
b) Subsidised only if prescribed for urethral or cervical a	administration and	the prescript	tion is endorsed accordingly.
Gel 2%, 11 ml urethral syringe - Subsidy by endorsement	42.00	10	✓ Instillagel Lido
a) Up to 5 each available on a PSO			_
b) Subsidised only if prescribed for urethral or cervical a	administration and	the prescript	tion is endorsed accordingly.
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE			
Oral (gel) soln 2%	38.00	200 ml	✓ Mucosoothe
Inj 1%, 5 ml ampoule – Up to 25 inj available on a PSO		25	✓ Lidocaine-Claris
, .,	17.50	50	
	(35.00)		Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	8.25 [°]	25	✓ Lidocaine-Claris
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5	
	(20.00)		Xylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	6.20	5	✓ Lidocaine-Claris
Inj 2%, 20 ml vial - Up to 5 inj available on a PSO	6.45	5	✓ <u>Lidocaine-Claris</u>
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE			
, , ,	81.50	10	✓ Pfizer
• •		. •	
Inj 1%, 20 ml ampoule – Up to 5 inj available on a PSO Inj 1%, 20 ml vial – Up to 5 inj available on a PSO	12.00 (20.00) 6.20 6.45	5	Xylocaine ✓ Lidocaine-Claris

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] - Special Authority see \$A0906 above	- Retail phar	macy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Authority	see SA0906	above – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	✓ EMLA

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

Non-opioid Analgesics

For aspirin & chloroform application refer Standard Formulae, page 247

ASPIRIN

★ Tab dispersible 300 mg - Up to 30 tab available on a PSO......4.50
100
✓ Ethics Aspirin

		Subsidy		Fully	Brand or
		(Manufacturer's Pri \$	ce) Subs Per	sidised •	Generic Manufacturer
CAF	SAICIN - Subsidy by endorsement				
	Subsidised only if prescribed for post-herpetic neuralgia or dia accordingly.	abetic peripheral	neuropathy a	nd the	prescription is endorsed
	Crm 0.075%	12.50	45 g OP	1	Zostrix HP
		15.83	57 g OP		Rugby Capsaicin Topical Cream S29
IEF	OPAM HYDROCHLORIDE				
	Tab 30 mg	23.40	90	1	Acupan
	ACETAMOL				·
	Tab 500 mg - blister pack	0.50	20		Medco Paracare
		1.12		✓	Ethics Paracetamol Classic
		2.48	100	✓	Paracare
		24.82	1,000	•	Paracetamol Pharmacare
				1	Pharmacare
	a) Maximum of 300 tab per prescription; can be waived b) Up to 30 tab available on a PSO c) 1) Subsidy by endorsement for higher quantities is	available for pati			
	b) Up to 30 tab available on a PSO c) 1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, are annotate the prescription as endorsed where disc. 2) Maximum of 100 tab per dispensing for non-endorsed patients), then dispense in re	available for patind the prescription spensing history slorsed patients.	n is annotated supports a lon f quantities p	d accor g-term rescrib	rdingly. Pharmacists may a condition. ed for more than 100 tabs
	b) Up to 30 tab available on a PSO c) 1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, are annotate the prescription as endorsed where disc. 2) Maximum of 100 tab per dispensing for non-endorsed.	available for patind the prescription spensing history solorsed patients. I peat dispensings	n is annotated supports a lon f quantities p	d accord g-term rescrib g 100	rdingly. Pharmacists may n condition. ed for more than 100 tabs
	b) Up to 30 tab available on a PSO c) 1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, are annotate the prescription as endorsed where disc. 2) Maximum of 100 tab per dispensing for non-endorsed patients), then dispense in retab 500 mg - bottle pack — Maximum of 300 tab per	available for patind the prescription spensing history solorsed patients. I peat dispensings 24.82 ailable for patients cription is annotal supports a long-teed patients. If queen a supports a long-teed patients. If queen and the supports a long-teed patients.	n is annotated supports a lor f quantities protection 1,000 s with long teled according term condition lantities prescription.	d accordance of	rdingly. Pharmacists may a condition. ed for more than 100 tabs tab per dispensing. Pharmacare ditions who require regular armacists may annotate the for more than 100 tabs (for more than 100 tabs)
	b) Up to 30 tab available on a PSO c) 1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, are annotate the prescription as endorsed where die 2) Maximum of 100 tab per dispensing for non-ence (for non-endorsed patients), then dispense in re Tab 500 mg - bottle pack — Maximum of 300 tab per prescription; can be waived by endorsement	available for pating the prescription of the prescription opensing history solorsed patients. I peat dispensings24.82 aliable for patients cription is annotated supports a long-teed patients. If quespensings not except	n is annotated supports a lor f quantities protection 1,000 s with long teled according term condition lantities prescription.	d accordance of	rdingly. Pharmacists may a condition. ed for more than 100 tabs tab per dispensing. Pharmacare ditions who require regular armacists may annotate the for more than 100 tabs (for more than 100 tabs).
	b) Up to 30 tab available on a PSO c) 1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, ar annotate the prescription as endorsed where die 2) Maximum of 100 tab per dispensing for non-end (for non-endorsed patients), then dispense in re Tab 500 mg - bottle pack — Maximum of 300 tab per prescription; can be waived by endorsement	available for pating the prescription of the prescription opensing history solorsed patients. I peat dispensings24.82 aliable for patients cription is annotated supports a long-teed patients. If quespensings not except	n is annotated supports a lor f quantities protection 1,000 s with long tested according term condition lantities prescueding 100 to the condition to the condition lantities prescueding 100 to the condition to the condition lantities prescueding 100 to the condition lantities prescu	d accordance of	rdingly. Pharmacists may a condition. ed for more than 100 tabs tab per dispensing. Pharmacare ditions who require regular armacists may annotate to for more than 100 tabs (for dispensing.
*	b) Up to 30 tab available on a PSO c) 1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, at annotate the prescription as endorsed where dis 2) Maximum of 100 tab per dispensing for non-endorsed patients), then dispense in re Tab 500 mg - bottle pack — Maximum of 300 tab per prescription; can be waived by endorsement	available for patind the prescription spensing history solorsed patients. I peat dispensings24.82 at a liable for patients cription is annotated supports a long-teed patients. If quespensings not exc5.45	n is annotated supports a lor f quantities protection 1,000 s with long tested according term condition lantities prescueding 100 to the condition to the condition lantities prescueding 100 to the condition to the condition lantities prescueding 100 to the condition lantities prescu	d according-term rescrib g 100 mm con ly. Ph cribed ab per	rdingly. Pharmacists may a condition. ed for more than 100 tabs tab per dispensing. Pharmacare ditions who require regula armacists may annotate to for more than 100 tabs (for dispensing.
K	b) Up to 30 tab available on a PSO c) 1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, are annotate the prescription as endorsed where dis 2) Maximum of 100 tab per dispensing for non-endorsed patients), then dispense in re Tab 500 mg - bottle pack — Maximum of 300 tab per prescription; can be waived by endorsement	available for patind the prescription spensing history storsed patients. I peat dispensings	n is annotated supports a lor f quantities protection of the second of t	d according-term rescrib g 100 mm con ly. Ph cribed ab per	rdingly. Pharmacists may a condition. ed for more than 100 tabs tab per dispensing. Pharmacare ditions who require regular armacists may annotate tifor more than 100 tabs (for dispensing. Paracare Paracare Double
*	b) Up to 30 tab available on a PSO c) 1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, are annotate the prescription as endorsed where disc. 2) Maximum of 100 tab per dispensing for non-endorsed patients), then dispense in refactor non-endorsed patients, then dispense in refactor non-endorsed patients, then dispense in represcription; can be waived by endorsement	available for patind the prescription spensing history storsed patients. I peat dispensings	n is annotated supports a lor f quantities protected according the second secon	d according term rescrib g 100 mm con ly. Ph	rdingly. Pharmacists may a condition. ed for more than 100 tabs tab per dispensing. Pharmacare ditions who require regular armacists may annotate the for more than 100 tabs (for dispensing.) Paracare Paracare Double Strength
*	b) Up to 30 tab available on a PSO c) 1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, are annotate the prescription as endorsed where dis 2) Maximum of 100 tab per dispensing for non-endorsed patients), then dispense in re Tab 500 mg - bottle pack — Maximum of 300 tab per prescription; can be waived by endorsement	available for patind the prescription spensing history storsed patients. I peat dispensings	n is annotated supports a lor f quantities protection of the second of t	d according decreased acco	rdingly. Pharmacists may a condition. ed for more than 100 tabs tab per dispensing. Pharmacare ditions who require regula armacists may annotate the for more than 100 tabs (for dispensing. Paracare Paracare Double

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
Opioid Analgesics				
CODEINE PHOSPHATE - Safety medicine; prescriber may dete	ermine dispensing fre	quen	СУ	
Tab 15 mg		100		PSM
PSM to be Sole Supply on 1 November 2020				
Tab 30 mg	7.45	100	1	PSM
PSM to be Sole Supply on 1 November 2020				
Tab 60 mg	14.25	100	•	PSM
PSM to be Sole Supply on 1 November 2020				
DIHYDROCODEINE TARTRATE				
Tab long-acting 60 mg	8.60	60	✓	DHC Continus
FENTANYL				
a) Only on a controlled drug form				
b) No patient co-payment payable				
 c) Safety medicine; prescriber may determine dispensing free 	eauencv			
Inj 50 mcg per ml, 2 ml ampoule		5	/	Fentanyl GH
,	3.56	10		Boucher and Muir
			/	Fentanyl IE S29
Inj 50 mcg per ml, 10 ml ampoule	9.41	10		Boucher and Muir
Patch 12.5 mcg per hour		5		Fentanyl Sandoz
Patch 25 mcg per hour		5	1	Fentanyl Sandoz
Patch 50 mcg per hour		5	✓	Fentanyl Sandoz
Patch 75 mcg per hour	9.25	5	1	Fentanyl Sandoz
Patch 100 mcg per hour	11.40	5	✓	Fentanyl Sandoz
METHADONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	equency			
d) Extemporaneously compounded methadone will only be	reimbursed at the rat	e of th	ne cheapes	st form available
(methadone powder, not methadone tablets).				
 e) For methadone hydrochloride oral liquid refer Standard F 				
Tab 5 mg		10		Methatabs
Oral liq 2 mg per ml		200 m		Biodone
Oral liq 5 mg per ml		200 m		Biodone Forte
Oral liq 10 mg per ml		200 m		Biodone Extra Forte
Inj 10 mg per ml, 1 ml	61.00	10	•	AFT
MORPHINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre				
Oral liq 1 mg per ml		200 m		RA-Morph
Oral liq 2 mg per ml		200 m		RA-Morph
Oral liq 5 mg per ml	19.44	200 m	_	Ordine \$29
0.15.40	07.74			RA-Morph
Oral liq 10 mg per ml	27.74	200 m		Ordine \$29
			•	RA-Morph

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per	•	Manufacturer
MORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
 Safety medicine; prescriber may determine dispensing fr 			_	
Tab immediate-release 10 mg	2.80	10	/	Sevredol
Sevredol to be Sole Supply on 1 November 2020				
Tab long-acting 10 mg		10		Arrow-Morphine LA
Tab immediate-release 20 mg	5.52	10	•	Sevredol
Sevredol to be Sole Supply on 1 November 2020	0.05	10	./	Aurau Marahina I A
Tab long-acting 30 mg		10		Arrow-Morphine LA Arrow-Morphine LA
Tab long-acting 60 mg Cap long-acting 10 mg		10		m-Eslon
Cap long-acting 10 mg		10		m-Esion
Cap long-acting 50 mg		10		m-Esion
Cap long-acting 100 mg		10		m-Esion
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a P		5		DBL Morphine
inj o mg per mi, i mi ampodie – op to o inj avaliable on a r	00	J	•	Sulphate
Inj 10 mg per ml, 1 ml ampoule - Up to 5 inj available on a	PSO 4.47	5	1	DBL Morphine
ing to mg per mi, i mi ampodie – op to 5 ing available on a	304.47	J	•	Sulphate
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a	PSO 4.76	5	1	DBL Morphine
ing 15 mg per mi, 1 mi ampodie – op to 5 ing available on a	304.70	J	•	Sulphate
Ini 20 mg nor ml. 1 ml amnoulo. Lin to 5 ini available on a	DSO 610	5	1	DBL Morphine
Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a	F3U 0. 19	5	•	Sulphate
(Arrow-Morphine LA Tab long-acting 10 mg to be delisted 1 Octo	phor 2020)			Julphate
, , ,	DDEI 2020)			
MORPHINE TARTRATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fr		_		DDI Marrahina
Inj 80 mg per ml, 1.5 ml ampoule	42.72	5	•	DBL Morphine Tartrate
(DDI Marahina Tartrata Ini 00 mg nay ml 1 E ml amnayla ta ha	dalistad 1 Cantambar	2020	١	rartrate
(DBL Morphine Tartrate Inj 80 mg per ml, 1.5 ml ampoule to be of	ielistea i September i	2020,	'	
OXYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fr			,	
Tab controlled-release 5 mg		20		Oxycodone Sandoz
Tab controlled-release 10 mg		20		Oxycodone Sandoz
Tab controlled-release 20 mg		20		Oxycodone Sandoz
Tab controlled-release 40 mg		20 20		Oxycodone Sandoz
Tab controlled-release 80 mg		20		Oxycodone Sandoz OxyNorm
Cap immediate-release 5 mg Cap immediate-release 10 mg		20		OxyNorm
Cap immediate-release 10 mg		20		OxyNorm
Oral liq 5 mg per 5 ml		20 250 m		OxyNorm
Inj 10 mg per ml, 1 ml ampoule		5		OxyNorm
Inj 10 mg per ml, 2 ml ampoule		5		OxyNorm
Inj 50 mg per ml, 1 ml ampoule		5		OxyNorm
PARACETAMOL WITH CODEINE – Safety medicine; prescribe		-		
* Tab paracetamol 500 mg with codeine phosphate 8 mg	,	1,000	•	paracetamol +
Tab paracetamor 500 mg with codeline phosphate 6 mg	10.21	1,000	,	Codeine (Relieve)
				Soucille (Helleve)

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre				
Tab 50 mg		10		<u>PSM</u>
Inj 50 mg per ml, 1 ml ampoule - Up to 5 inj available on a P	°SO4.98	5	•	DBL Pethidine Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a P	PSO5.12	5	✓	DBL Pethidine Hydrochloride
TRAMADOL HYDROCHLORIDE				
Tab sustained-release 100 mg	1.52	20	✓	Tramal SR 100
Tramal SR 100 to be Sole Supply on 1 November 2020				
Tab sustained-release 150 mg	2.10	20	✓	Tramal SR 150
Tramal SR 150 to be Sole Supply on 1 November 2020				
Tab sustained-release 200 mg	2.75	20	•	Tramal SR 200
Tramal SR 200 to be Sole Supply on 1 November 2020			_	
Cap 50 mg	2.25	100	/	Arrow-Tramadol
Antidepressants				
Cyclic and Related Agents				
AMITRIPTYLINE - Safety medicine; prescriber may determine d	ispensing frequency			
Tab 10 mg		100	1	Arrow-Amitriptyline
Tab 25 mg		100		Arrow-Amitriptyline
Tab 50 mg	2.51	100	✓	Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; prescri	iber may determine d	lispen	sing fregu	ency
Tab 10 mg	-	100		Anafranil S29
1 do 10 mg				Apo-Clomipramine
Tab 25 mg	4.73	50		Apo-Clomipramine
· · · · · · · · · · · · · · · · · · ·	9.46	100		Apo-Clomipramine
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy by en	dorsement			
a) Safety medicine; prescriber may determine dispensing fre				
b) Subsidy by endorsement – Subsidised for patients who w		[doth	ienin] hvdr	rochloride prior to 1 June
2019 and the prescription is endorsed accordingly. Pharr				
exists a record of prior dispensing of dosulepin [dothiepin				
Tab 75 mg		30	1	Dosulepin Mylan
Cap 25 mg		50		Dosulepin
				Mylan S29
IMIPRAMINE HYDROCHLORIDE - Safety medicine; prescriber	may determine diene	neine	fraguanc	•
Tab 10 mg		50		y Tofranil
rab to fig	10.96	100		Tofranil
Tab 25 mg		50		Tofranil
MAPROTILINE HYDROCHLORIDE – Safety medicine; prescribe				
Tab 25 mg		30		Ludiomil
100 20 mg	12.53	50		Ludiomil
	25.06	100		Ludiomil
Tab 75 mg		20		Ludiomil
100 / J mg	17.01	20		Ludioniii

✓ Ludiomil

30

21.01

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	Brand or Generic Manufacturer
NORTRIPTYLINE HYDROCHLORIDE - Safety medicine; pres				
Tab 10 mg Tab 25 mg		100		Norpress
Tab 25 mg	5.96	180	• !	Norpress
Monoamine-Oxidase Inhibitors (MAOIs) - Non	Selective			
PHENELZINE SULPHATE – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who were prescription is endorsed accordingly. Pharmacists may any prior dispensing of phenelzine sulphate.	notate the prescription	as end	orsed whe	ere there exists a record
* Tab 15 mg		60	√	Lupin S29 Nardil S29 S29
// unin coo Tob 15 mg to be delicted 1 October 2000)	118.00	100	•	Nardil
(Lupin S29 Tab 15 mg to be delisted 1 October 2020) (Nardil S29 S29 Tab 15 mg to be delisted 1 October 2020) (Nardil Tab 15 mg to be delisted 1 October 2020)				
TRANYLCYPROMINE SULPHATE				
Tab 10 mg	12.85 22.94	28 50		Parnate S29 S29 Parnate
	96.00	100		Parnate S29 S29
Monoamine-Oxidase Type A Inhibitors	33.33			
••				
MOCLOBEMIDE	0.40	00		·
* Tab 150 mg * Tab 300 mg		60 60	-	<u>Aurorix</u> Aurorix
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE				
* Tab 20 mg	1.52	84	✓ <u>I</u>	PSM Citalopram
ESCITALOPRAM				
* Tab 10 mg	1.11	28	√ I	Escitalopram- Apotex
* Tab 20 mg	1.90	28	√ I	Escitalopram-
				Apotex
FLUOXETINE HYDROCHLORIDE				
* Tab dispersible 20 mg, scored – Subsidy by endorsement.	1.98	30	√ I	Fluox
	9.93		✓ /	Arrow-Fluoxetine
Subsidised by endorsement 1) When prescribed for a patient who cannot swallor accordingly; or	w whole tablets or caps	sules a	nd the pre	escription is endorsed
 When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined with 				
Cap 20 mg	2.91	84	√	Fluox
, - 3	7.49	90		Arrow-Fluoxetine
PAROXETINE				
* Tab 20 mg	0.04	90		Loxamine

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	1	Manufacturer
SERTRALINE				
Tab 50 mg		30		Setrona
≮ Tab 100 mg	1.61	30		Setrona
Other Antidepressants				
IIRTAZAPINE				
Tab 30 mg		30		Apo-Mirtazapine
Tab 45 mg	3.48	30	•	Apo-Mirtazapine
ENLAFAXINE	2.22	0.4		- 1 (VD
€ Cap 37.5 mg		84	_	Enlafax XR
Cap 150 mg		84	_	Enlafax XR
€ Cap 150 mg	11.10	84		Enlafax XR
Antiepilepsy Drugs				
Agents for Control of Status Epilepticus				
LONAZEPAM - Safety medicine; prescriber may determ				
Inj 1 mg per ml, 1 ml	21.00	5	✓	Rivotril
IAZEPAM - Safety medicine; prescriber may determine of	dispensing frequency			
Inj 5 mg per ml, 2 ml ampoule - Subsidy by endorsem	ent23.66	5	•	Hospira
a) Up to 5 inj available on a PSO				
b) Only on a PSO				
c) PSO must be endorsed "not for anaesthetic pro		_		O
Rectal tubes 5 mg - Up to 5 tube available on a PSO.		5		Stesolid
Rectal tubes 10 mg - Up to 5 tube available on a PSC	740.87	5	•	Stesolid
ARALDEHYDE		_		
5 Inj 5 ml	1,500.00	5	•	AFT S29
HENYTOIN SODIUM			_	
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available		5	/	Hospira
Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available		_		
PSO	133.92	5	•	Hospira
Control of Epilepsy				
ARBAMAZEPINE			_	
Tab 200 mg		100		Tegretol CD
Tab long-acting 200 mg		100 100	_	Tegretol CR
Tab long-acting 400 mg		100		Tegretol CR
Oral lig 20 mg per ml		250 m		Tegretol
LOBAZAM – Safety medicine; prescriber may determine		_0011		9. 0.0.
Tab 10 mg		50	J	Frisium
•		50	•	i riolulli
LONAZEPAM – Safety medicine; prescriber may determ		0 ml 0	10 J	Rivotril
Oral drops 2.5 mg per ml	1.30 I	O IIII C	л. 🔻	myum
THOSUXIMIDE	140.00	100		Zavantin
Cap 250 mg		100 200 m		Zarontin Zarontin
Oral liq 250 mg per 5 ml		200 II		L ai Villii

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy		Fully	Brand or
(Manutacturer's Price) \$	Per		Generic Manufacturer
abalin			
2.65	100	✓ /	Apo-Gabapentin
	100	1	Apo-Gabapentin
	100	1	Apo-Gabapentin
pharmacy			
	14	✓ \	/impat
50.06	14	✓ \	/impat
200.24	56	✓ \	/impat
75.10	14	✓ \	/impat
300.40	56	✓ \	/impat
400.55	56		/impat
	(Manufacturer's Price) \$ abalin2.654.075.64 pharmacy25.0450.06 200.2475.10 300.40	(Manufacturer's Price) \$ Per abalin2.65 1004.07 1005.64 100 pharmacy25.04 1450.06 1450.06 1450.10 14300.40 56	(Manufacturer's Price) \$ Subsidised Per \$ Subsidised Per \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$

⇒SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

LAMOTRIGINE

▲ Tab dispe	ersible 2 mg	6.74	30	✓ La	ımictal
▲ Tab dispe	ersible 5 mg	9.64	30	✓ La	ımictal
		15.00	56	✓ Aı	row-Lamotrigine
* Tab dispe	ersible 25 mg	2.76	56	✓ Lo	gem
* Tab dispe	ersible 50 mg	3.31	56	✓ Lo	gem
* Tab dispe	ersible 100 mg	4.40	56	✓ Lo	ogem
	rigine Tab dispersible 5 mg to be delisted 1 October 2020				_
LEVETIRACE	TAM				
Tab 250 i	ng	4.99	60	✓ E\	<u>reret</u>
Tab 500 i	ng	8.79	60	✓ E\	veret .
Tab 750 i	ng	.14.39	60	✓ Ev	veret .
Tab 1,000	O mg	.18.59	60	✓ Ev	veret .
	00 mg per ml		300 ml OP	✓ Le	vetiracetam-AFT
PHENOBARE	BITONE				
For phen	obarbitone oral liquid refer Standard Formulae, page 247				
* Tab 15 m	g	.40.00	500	✓ PS	SM
	g		500	✓ PS	SM
PHENYTOIN	SODIUM				
* Tab 50 m	g	.75.00	200	✓ Di	lantin Infatab
	ig		200	✓ Di	lantin
	mg		200	✓ Di	lantin
	0 mg per 5 ml		500 ml	✓ Di	lantin

	Subsidy		Fully	
			rully	Brand or
	(Manufacturer's Price)		sidised	Generic
	\$	Per	•	Manufacturer
REGABALIN				
Note: Not subsidised in combination with subsidised gaba	apentin			
★ Cap 25 mg	2.25	56	✓ P	regabalin Pfizer
≮ Cap 75 mg		56	✓ P	regabalin Pfizer
K Cap 150 mg		56	✓ P	regabalin Pfizer
K Cap 300 mg		56	✓ P	regabalin Pfizer
PRIMIDONE				
F Tab 250 mg	17.25	100	✓ A	po-Primidone
•	62.00	200	✓ N	lysoline S29 S29
SODIUM VALPROATE				
Tab 100 mg	13.65	100	√ E	pilim Crushable
Tab 200 mg EC		100	✓ E	pilim
Tab 500 mg EC		100	✓ E	pilim
FOral liq 200 mg per 5 ml		300 ml	✓ E	pilim S/F Liquid
1 01				pilim Syrup
k Inj 100 mg per ml, 4 ml	41.50	1		pilim IV
STIRIPENTOL - Special Authority see SA1330 below - Retai				•
Cap 250 mg		60	✓ D	iacomit S29
Powder for oral liq 250 mg sachet		60	✓ D	iacomit S29

⇒SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

TO	PIRAMATE		
\blacktriangle	Tab 25 mg11.07	60	✓ Arrow-Topiramate
			✓ Topiramate Actavis
	26.04		✓ Topamax
\blacktriangle	Tab 50 mg18.81	60	✓ Arrow-Topiramate
			✓ Topiramate Actavis
	44.26		✓ Topamax
\blacktriangle	Tab 100 mg31.99	60	✓ Arrow-Topiramate
	-		✓ Topiramate Actavis
	75.25		✓ Topamax
\blacktriangle	Tab 200 mg55.19	60	✓ Arrow-Topiramate
	-		✓ Topiramate Actavis
	129.85		✓ Topamax
\blacktriangle	Sprinkle cap 15 mg	60	✓ Topamax
\blacktriangle	Sprinkle cap 25 mg	60	✓ Topamax
VIC	ABATRIN - Special Authority see SA1907 on the next page - Retail pharm	nacy	
	Tab 500 mg119.30	100	✓ Sabril



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

⇒SA1907 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
 - 2.2 It is impractical or impossible (due to comorbid conditions, or health system capacity constraints) to monitor the patient's visual fields..

Notes: ``Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions, or health system capacity constraints) to monitor the patient's visual fields..

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

RIZATRIPTAN

Tab orodispersible 10 mg......3.65 30 ✓ Rizamelt

Rizamelt to be Sole Supply on 1 October 2020

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
SUMATRIPTAN				
Tab 50 mg	24.44	100	/	Apo-Sumatriptan
Tab 100 mg		100		Apo-Sumatriptan
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj pe	er			-
prescription	34.00	2 OP	/	Imigran
	42.67			Sun Pharma S29
	81.15		/	Clustran
Imigran to be Sole Supply on 1 September 2020	"			
Sun Pharma 829 Inj 12 mg per ml, 0.5 ml prefilled pen to be de	,	020)		
Clustran Inj 12 mg per ml, 0.5 ml prefilled pen to be delisted 1 Se	eptember 2020)			
Prophylaxis of Migraine				
or Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SY	STEM, page 50			
PIZOTIFEN	71 0			
₹ Tab 500 mcg	23.21	100	/	Sandomigran
5				J
Antinausea and Vertigo Agents				
or Antispasmodics refer to ALIMENTARY TRACT, page 8				
PREPITANT – Special Authority see SA0987 below – Retail ph	armacy			
Cap 2 × 80 mg and 1 × 125 mg	84.00	3 OP		Emend Tri-Pack
⇒SA0987 Special Authority for Subsidy				
nitial application from any relevant practitioner. Approvals valid	d for 12 months wher	e the	patient is u	undergoing highly
metogenic chemotherapy and/or anthracycline-based chemothe				
tenewal from any relevant practitioner. Approvals valid for 12 m hemotherapy and/or anthracycline-based chemotherapy for the			undergoin	ig nigniy emetogenic
BETAHISTINE DIHYDROCHLORIDE	irealinent of mangna	icy.		
* Tab 16 mg	3.88	84	1	Vergo 16
Vergo 16 to be Sole Supply on 1 November 2020		04	•	vergo to
CYCLIZINE HYDROCHLORIDE				
Tab 50 mg	0.55	10	1	Nausicalm
CYCLIZINE LACTATE		. •		
Inj 50 mg per ml, 1 ml	14 95	5	/	Nausicalm
OMPERIDONE	14.00	Ü	•	Nuuoiouiiii
★ Tab 10 mg	2 25	100	1	Pharmacy Health
YOSCINE HYDROBROMIDE		. 50		
For Inj 400 mcg per ml, 1 ml ampoule	46 50	5	1	Hospira
r inj too mog por mi, i mi ampoulo	93.00	10		Martindale S29
	30.00	10	•	mai di luale des
Patch 1.5 mg Special Authority see SA1927 below - Retai	I			
Patch 1.5 mg - Special Authority see SA1927 below - Retai pharmacy		2	/	Scopoderm TTS

⇒SA1927 Special Authority for Subsidy

Initial application — (control of intractable nausea, vomiting or inability to swallow saliva or clozapine induced hypersalivation) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

METOCI OPRAMIDE HYDROCHI ORIDE

(Manu	Subsidy	Fully	Brand or
	facturer's Price)	Subsidised	Generic
	\$ P	Per 🗸	Manufacturer

continued...

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Initial application — (pandemic circumstances- symptomatic relief of respiratory secretions in palliative care) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Requires palliative care in the community setting; and
- 2 Requires symptomatic relief of respiratory secretions that is not possible with 'as required subcutaneous hyoscine injections' due to COVID-19 constraints on the health sector; and
- 3 Access to a syringe driver for administration of injectable hyoscine is not possible due to COVID-19 constraints on the health sector.

Renewal — (control of intractable nausea, vomiting or inability to swallow saliva or clozapine induced hypersalivation) from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

	Tab 10 mg - Up to 30 tab available on a PSO1.30	100	✓ Metoclopramide Actavis 10
	Metoclopramide Actavis 10 to be Sole Supply on 1 October 2020		
*	Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO9.50	10	✓ Pfizer
O١	IDANSETRON		
*	Tab 4 mg2.68	50	✓ Onrex
*	Tab disp 4 mg - Up to 10 tab available on a PSO0.76	10	Ondansetron ODT-DRLA
	Ondansetron ODT-DRLA to be Sole Supply on 1 October 2020		
*	Tab 8 mg4.57	50	✓ Onrex
*	Tab disp 8 mg - Up to 10 tab available on a PSO1.13	10	Ondansetron ODT-DRLA
	Ondansetron ODT-DRLA to be Sole Supply on 1 October 2020		
PR	OCHLORPERAZINE		
*	Tab 3 mg buccal	50	Buccastem
*	Tab 5 mg - Up to 30 tab available on a PSO	250	✓ Nausafix
*	Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO25.81	10	✓ Stemetil

Antipsychotics

General

AMISULPRIDE - Safety medicine; prescriber may de	termine dispensing frequency	y	
Tab 100 mg	5.15	30	✓ Sulprix
•	17.16	100	✓ Amisulpride
			Mylan S29
Tab 200 mg	14.96	60	✓ Sulprix
Tab 400 mg	29.78	60	✓ Sulprix

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
A DIDIDIDA ZOLE	-	1 01	<u> </u>	Wandlacturer
ARIPIPRAZOLE – Safety medicine; prescriber may determine		30	./	Arininrozolo Condoz
Tab 5 mg Tab 10 mg		30		Aripiprazole Sandoz Aripiprazole Sandoz
Tab 15 mg		30		Aripiprazole Sandoz
Tab 20 mg		30		Aripiprazole Sandoz
Tab 30 mg		30		Aripiprazole Sandoz
ŭ				
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; p				
Tab 10 mg - Up to 30 tab available on a PSO		100		<u>Largactil</u>
Tab 25 mg – Up to 30 tab available on a PSO		100		Largactil
Tab 100 mg – Up to 30 tab available on a PSO		100		Largactil
Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO	30.79	10	•	<u>Largactil</u>
CLOZAPINE – Hospital pharmacy [HP4]				
Safety medicine; prescriber may determine dispensing freq	uency			
Tab 25 mg	5.69	50	✓	Clozaril
	6.69		✓	Clopine
	11.36	100	✓	Clozaril
	13.37		✓	Clopine
Tab 50 mg	8.67	50		Clopine
	17.33	100	✓	Clopine
Tab 100 mg	14.73	50	✓	Clozaril
	17.33		✓	Clopine
	29.45	100	✓	Clozaril
	34.65		✓	Clopine
Tab 200 mg	34.65	50		Clopine
	69.30	100		Clopine
Suspension 50 mg per ml	17.33	100 m	nl 🗸	Clopine
HALOPERIDOL - Safety medicine; prescriber may determine of	dispensing frequency			
Tab 500 mcg - Up to 30 tab available on a PSO		100	1	Serenace
Tab 1.5 mg - Up to 30 tab available on a PSO		100		Serenace
Tab 5 mg - Up to 30 tab available on a PSO		100		Serenace
Oral lig 2 mg per ml - Up to 200 ml available on a PSO		100 m		Serenace
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a F		10		Serenace
LEVOMEPROMAZINE - Safety medicine; prescriber may dete	armina dienancina fraa	uana	,	
Tab 25 mg (33.8 mg as a maleate)		100		Nozinan (Swiss)
Tab 25 mg as a maleate		100		Nozinan (Swiss)
Tab 100 mg (135 mg as a maleate)		100		Nozinan (Swiss)
Tab 100 mg as a maleate		100		Nozinan
· ·				
LEVOMEPROMAZINE HYDROCHLORIDE - Safety medicine;				
Inj 25 mg per ml, 1 ml ampoule		10		<u>Nozinan</u>
LITHIUM CARBONATE - Safety medicine; prescriber may dete	ermine dispensing freq	uency	У	
Tab 250 mg - Subsidy by endorsement		500		Lithicarb FC
Subsidised for patients who were taking lithium carbon				
endorsed accordingly. Pharmacists may annotate the	prescription as endors	ed wh	nere there e	exists a record of prior
dispensing of lithium carbonate.				
Tab long-acting 400 mg	72.00	100	✓	Priadel
0 050	0.40	100		Douglas
Cap 250 mg(Lithicarb FC Tab 250 mg to be delisted 1 November 2020)	9.42	100	•	Douglas

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
ANZAPINE - Safety medicine; prescriber may determine disp	ensing frequency			
Tab 2.5 mg	1.35	28	✓	Zypine
Zypine to be Sole Supply on 1 November 2020				
Tab 5 mg	1.58	28	✓	Zypine
Zypine to be Sole Supply on 1 November 2020				
Tab orodispersible 5 mg	1.81	28	✓	Zypine ODT
Zypine ODT to be Sole Supply on 1 November 2020				
Tab 10 mg	2.01	28	1	Zypine
Zypine to be Sole Supply on 1 November 2020				••
Tab orodispersible 10 mg	2.38	28	1	Zypine ODT
Zypine ODT to be Sole Supply on 1 November 2020				,,
,	noncina froguency			
ERICYAZINE – Safety medicine; prescriber may determine dis		84	.1	Neulactil
Tab 2.5 mg	12.49	100		Neulactil
Toh 10 mg		84		Neulactil
Tab 10 mg				
	44.45	100	•	Neulactil
JETIAPINE – Safety medicine; prescriber may determine dispe				
Tab 25 mg	2.15	90	✓	Quetapel
Quetapel to be Sole Supply on 1 November 2020				
Tab 100 mg	5.06	90	1	Quetapel
Quetapel to be Sole Supply on 1 November 2020				
Tab 200 mg	8.90	90	✓	Quetapel
Quetapel to be Sole Supply on 1 November 2020				
Tab 300 mg	12.86	90	✓	Quetapel
Quetapel to be Sole Supply on 1 November 2020				
SPERIDONE - Safety medicine; prescriber may determine dis	snensing frequency			
Tab 0.5 mg		60	1	Actavis
Tab 1 mg		60		Actavis
Tab 2 mg		60		Actavis
Tab 3 mg		60		Actavis
Tab 4 mg		60		Actavis
Oral lig 1 mg per ml		30 m		Risperon
Risperon to be Sole Supply on 1 November 2020		JU 111		порегон
PRASIDONE – Safety medicine; prescriber may determine dis			_	
Cap 20 mg		60		Zusdone
Cap 40 mg		60		Zusdone
Cap 60 mg		60		Zusdone
Cap 80 mg	39.70	60	•	Zusdone
JCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pre-	scriber may determin	e dis	pensing fre	equency
Tab 10 mg		100		Clopixol

Depot Injections

uency	dispensing f	er may determine	FLUPENTHIXOL DECANOATE - Safety medicine; prescribe
✓ Fluanxol	. 5	13.14	Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PSO
✓ Fluanxol	5	20.90	Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PSO
✓ Fluanxol	5	40.87	Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO.

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
HALOPERIDOL DECANOATE - Safety medicine; prescriber ma	ay determine dispensir	ng frequ	ency	
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO	28.39	5	. ✓ H	laldol
Inj 100 mg per ml, 1 ml – Úp to 5 inj available on a PSO	55.90	5		laldol Concentrate laldol Decanoas S29
OLANZAPINE - Special Authority see SA1428 below - Retail pl	narmacy			
Safety medicine; prescriber may determine dispensing frequ	ency			
Inj 210 mg vial	252.00	1	√ <u>Z</u>	yprexa Relprevy
Inj 300 mg vial	414.00	1	✓ <u>Z</u>	'yprexa Relprevv
Inj 405 mg vial	504.00	1	√ <u>Z</u>	Zyprexa Relprevv

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing	ı frequency		
Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe		1	✓ Invega Sustenna
Inj 100 mg syringe		1	✓ Invega Sustenna
Inj 150 mg syringe		1	✓ Invega Sustenna

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE – Special Authority see SA1427 on the next page – Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency			
Inj 25 mg vial	135.98	1	✓ Risperdal Consta
Inj 37.5 mg vial	178.71	1	✓ Risperdal Consta
Inj 50 mg vial	217.56	1	✓ Risperdal Consta



Sub	osidy Fu	ly Brand or
(Manufactu	urer's Price) Subsidise	ed Generic
	\$ Per	/ Manufacturer

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml − Up to 5 inj available on a PSO......19.80 5 Clopixol

Anxiolytics

BUSPIRONE HYDROCHLORIDE * Tab 5 mg Tab 10 mg		100 100	✓ <u>Orion</u> ✓ <u>Orion</u>
CLONAZEPAM – Safety medicine; prescriber may determine dis Tab 500 mcg Tab 2 mg	5.64	100 100	✓ <u>Paxam</u> ✓ <u>Paxam</u>
DIAZEPAM – Safety medicine; prescriber may determine dispen Tab 2 mg Tab 5 mg	15.05	500 500	✓ Arrow-Diazepam✓ Arrow-Diazepam
LORAZEPAM – Safety medicine; prescriber may determine disp Tab 1 mg Tab 2.5 mg	9.72	250 100	✓ Ativan ✓ Ativan
OXAZEPAM — Safety medicine; prescriber may determine dispertable 10 mg	6.17	100 100	✓ Ox-Pam ✓ Ox-Pam

Multiple Sclerosis Treatments

DIMETHYL FUMARATE - Special Authority see SA	A1559 below – Retail pharmacy		
Wastage claimable			
Cap 120 mg	520.00	14	✓ Tecfidera
Cap 240 mg	2,000.00	56	✓ Tecfidera

⇒SA1559 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

NERVOUS SYSTEM

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per Manufacturer

(Manufacturer's \$

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

continued...
The coordinator

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Phone: 04 460 4990

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- a) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- b) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- c) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
- v) new T2 lesions compared with a previous MR scan; and
- d) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week:
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point:
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- e) applications must be made by the patient's neurologist or general physician; and
- f) patients must have no previous history of lack of response to dimethyl fumarate; and
- g) patients must have not previously had intolerance to dimethyl fumarate; and
- h) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5: or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5: or
 - f) 3.0 to 4.5; or



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

g) 3.5 to 4.5; or

h) 4.0 to 4.5.

- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to dimethyl fumarate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate, teriflunomide and ocrelizumab is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

FINGOLIMOD - Special Authority see SA1562 below - Retail pharmacy

Wastage claimable

Cap 0.5 mg......2,200.00 28 **✓ Gilenya**

⇒SA1562 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990 Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);

NERVOUS SYSTEM

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	osidised	Generic	
\$	Per	1	Manufacturer	

continued...

- c) last at least one week;
- d) start at least one month after the onset of a previous relapse;
- e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
- f) be distinguishable from the effects of general fatigue; and
- g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) patients must have no previous history of lack of response to fingolimod; and
- 7) patients must have not previously had intolerance to fingolimod; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to fingolimod; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate, teriflunomide and ocrelizumab is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

NATALIZUMAB - Special Authority see SA1563 below - Retail pharmacy

⇒SA1563 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria):
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least
 - f) be distinguishable from the effects of general fatigue; and
 - a) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) treatment must be initiated and supervised by a neurologist who is registered in the Tysabri Australasian Prescribing Programme operated by the supplier; and
- 7) patients must have no previous history of lack of response to natalizumab; and
- 8) patients must have not previously had intolerance to natalizumab; and
- a) Patient is JC virus negative, or
 - b) Patient is JC virus positive and has given written informed consent acknowledging an understanding of the risk of progressive multifocal leucoencephalopathy (PML) associated with natalizumab
- 10) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0: or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or

NERVOUS SYSTEM

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 3) intolerance to natalizumab; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier.

Switching between natalizumab, fingolimod, dimethyl fumarate, teriflunomide and ocrelizumab is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

OCRELIZUMAB - Special Authority see SA1867 below - Retail pharmacy

⇒SA1867 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- a) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- b) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- c) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- d) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;



Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least
- f) be distinguishable from the effects of general fatigue; and
- g) not be associated with a fever (T> 37.5°C); and
- e) applications must be made by the patient's neurologist or general physician; and
- f) patients must have no previous history of lack of response to ocrelizumab; and
- g) patients must have not previously had intolerance to ocrelizumab; and
- h) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0: or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0: or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to ocrelizumab: or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate, teriflunomide and ocrelizumab is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

TERIFLUNOMIDE - Special Authority see SA1560 below - Retail pharmacy

Wastage claimable

28 ✓ Aubagio

⇒SA1560 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990 Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

				=
Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week:
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) patients must have no previous history of lack of response to teriflunomide; and
- 7) patients must have not previously had intolerance to teriflunomide; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to teriflunomide; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate, teriflunomide and ocrelizumab is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria



Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

continued...

are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

Other Multiple Sclerosis Treatments

GLATIRAMER ACETATE - Special Authority see SA1808 below - Retail pharmacy

Inj 40 mg prefilled syringe......2,275.00

12

✓ Copaxone

⇒SA1808 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. As a new Special Authority approval number is required the MSTAC coordinator should be notified of the change and a new prescription provided. **Entry Criteria**

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;

Subsidy (Manufacturer's Price)	s	Fully	Brand or Generic	
\$	Per	✓	Manufacturer	

- f) be distinguishable from the effects of general fatigue; and
- g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or
 - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment.
 Progression of disability is defined as progress by any of the following EDDSS Points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0: or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

INTERFERON BETA-1-ALPHA - Special Authority see SA1809 below - Retail pharmacy

Inj 6 million iu prefilled syringe	1,170.00	4	✓ Avonex
Injection 6 million ju per 0.5 ml pen injector	1.170.00	4	✓ Avonex Pen

⇒SA1809 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. As a new Special Authority approval number is required the MSTAC coordinator should be notified of the change and a new prescription provided.

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria):
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or
 - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

Stopping Criteria

Any of the following:

1) Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment. Progression of disability is defined as progress by any of the following EDDSS Points:

Subsidy	Fu	illy Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

- a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
- b) 1.0 to 3.0; or
- c) 1.5 to 3.5; or
- d) 2.0 to 4.0: or
- e) 2.5 to 4.5; or
- f) 3.0 to 4.5; or
- g) 3.5 to 4.5; or h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

INTERFERON BETA-1-BETA – Special Authority see SA1810 below – Retail pharmacy

⇒SA1810 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. As a new Special Authority approval number is required the MSTAC coordinator should be notified of the change and a new prescription provided. **Entry Criteria**

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:



-				
	Subsidy	Fully	Brand or	
	(Manufacturer's Price)	Subsidised	Generic	
	•	Por 🗸	Manufacturor	

- a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week:
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or
 - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

Stopping Criteria

Any of the following:

- 1) Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment. Progression of disability is defined as progress by any of the following EDDSS Points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0: or
 - c) 1.5 to 3.5: or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the

NERVOUS SYSTE

Subsidy (Manufacturer's Pric	e)	Fully Subsidised	Brand or Generic	
\$	Per		Manufacturer	

continued...

beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

Sedatives and Hypnotics

MELATONIN - Special Authority see SA1666 below - Retail pharmacy ✓ Circadin Tab modified-release 2 mg - No more than 5 tab per day......28.22

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

. 9	g p	y ·
ing frequency	10	✓ Midazolam-Claris
4.30	10	• Wildazolaiii-Claris
14.90	10	✓ Pfizer
dorsed for statu	is epilepticu	s use only.
2.50	5	Midazolam-Claris
44.00	_	4 D (1)
	-	✓ Pfizer s use only.
		1 August 2019 and the prescription e there exists a record of prior
5.22	100	✓ Nitrados
	14.90 dorsed for statu2.5011.90 dorsed for statu ency taking nitrazep scription as ence	14.90 10 dorsed for status epilepticu2.50 511.90 5 dorsed for status epilepticu ency taking nitrazepam prior to scription as endorsed wher

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
HENOBARBITONE SODIUM - Special Authority see SA138	36 below - Retail pharm	acy		
Inj 200 mg per ml, 1 ml ampoule	30.00	5	✓	Aspen S29
	68.00	10	✓	Max Health S29
Aspen 229 Inj 200 mg per ml, 1 ml ampoule to be delisted 1 SA1386 Special Authority for Subsidy itial application from any relevant practitioner. Approvals v		wal u	nless notil	ied for applications med
ne following criteria:				
Soth: 1 For the treatment of terminal agitation that is unrespons 2 The applicant is part of a multidisciplinary team working		t		
EMAZEPAM - Safety medicine; prescriber may determine d	ispensing frequency			
Tab 10 mg Normison to be Sole Supply on 1 November 2020	1.33	25	•	Normison
RIAZOLAM – Safety medicine; prescriber may determine dis	spensing frequency			
Tab 125 mcg	4	100		
T 050	(9.85)	400		Hypam
Tab 250 mcg		100		Ulumana
	(11.20)			Hypam
OPICLONE – Safety medicine; prescriber may determine dis		500	./	Zamialama Astavia
Tab 7.5 mg	9.50	500		Zopiclone Actavis
Stimulants/ADHD Treatments				
TOMOXETINE Con 10 mg	10 /1	28	./	Generic Partners
Cap 10 mg	107.03	20		Strattera
Cap 18 mg		28		Generic Partners
Cap to mg	107.03	20		Strattera
Cap 25 mg		28		Generic Partners
οωρ = οg	107.03			Strattera
Cap 40 mg		28		Generic Partners
Cap 40 mg	107.03			Strattera
Cap 40 mg				Generic Partners
•		28	•	
Cap 60 mg		28		
Cap 60 mg	46.51 107.03	28 28	•	Strattera Generic Partners
•	46.51 107.03		√	Strattera
Cap 60 mg	46.51 107.03 56.45 139.11		√ ✓	Strattera Generic Partners
Cap 60 mg	46.51 107.03 56.45 139.11	28	\ \ \	Strattera Generic Partners Strattera
Cap 60 mg Cap 80 mg Cap 100 mg DEXAMFETAMINE SULFATE - Special Authority see SA114		28 28	\ \ \	Strattera Generic Partners Strattera Generic Partners
Cap 80 mg Cap 100 mg		28 28	\ \ \	Strattera Generic Partners Strattera Generic Partners

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or
 (Manufacturer's Price) \$	Per 🗸	Generic Manufacturer

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1150 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine: prescriber may determine dispensing frequency

b) Salety medicine, prescriber may determine dispensing	y nequency		
Tab immediate-release 5 mg	3.20	30	Rubifen
Tab immediate-release 10 mg		30	Ritalin
-			Rubifen
Tab immediate-release 20 mg	7.85	30	Rubifen
Tab sustained-release 20 mg	10.95	30	Rubifen SR
	50.00	100	✓ Ritalin SR

⇒SA1150 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for



Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	✓	Manufacturer	

applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Roth:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1151 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab extended-release 18 mg	18.20	30	Methylphenidate ERTeva
	58.96		✓ Concerta
Tab extended-release 27 mg	22.00	30	Methylphenidate ERTeva
	65.44		✓ Concerta
Tab extended-release 36 mg	22.40	30	Methylphenidate ERTeva
	71.93		✓ Concerta
Tab extended-release 54 mg	26.40	30	Methylphenidate ERTeva
	86.24		✓ Concerta
Cap modified-release 10 mg		30	✓ Ritalin LA
Cap modified-release 20 mg	20.40	30	Ritalin LA
Cap modified-release 30 mg	25.52	30	✓ Ritalin LA
Cap modified-release 40 mg	30.60	30	✓ Ritalin LA

⇒SA1151 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the

Subsidy (Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer	
*				

last 2 years and has recommended treatment for the patient in writing; and

- 4 Either:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

		MODAFINIL – Special Authority see SA1932 below – Retail pharmacy
✓ Modavigil	30	Tab 100 mg32.00
✓ Modavigil	60	64.00

⇒SA1932 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Any of the following:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 A multiple sleep latency test is not possible due to COVID-19 constraints on the health sectors; or
 - 2.3 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Fither:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	4.34	90	✓ Donepezil-Rex
* Tab 10 mg	6.64	90	✓ Donepezil-Rex
RIVASTIGMINE - Special Authority see SA1488 below -	Retail pharmacy		
Patch 4.6 mg per 24 hour	48.75	30	✓ Generic Partners
Patch 9.5 mg per 24 hour	48.75	30	✓ Generic Partners

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:



Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
\$	Per	✓	Manufacturer

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 2 mg with naloxone 0.5 mg.	18.37
---	-------

Tab sublingual 8 mg with naloxone 2 mg53.12

✓ <u>Buprenorphine</u>
Naloxone BNM

28

28

✓ Buprenorphine
Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone: and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone);
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE			
Tab modified-release 150 mg	11.00	30	Zyban
DISULFIRAM			
Tab 200 mg	153.00	100	✓ Antabuse
NALTREXONE HYDROCHLORIDE - Special Authority	see SA1408 below - Retai	l pharmacy	
Tab 50 mg	112.55	30	✓ Naltraccord

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment: or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.
- b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A. Patch 7 mg - Up to 28 patch available on a PSO17.28 28 ✓ Habitrol Patch 7 mg for direct distribution only - [Xpharm]......3.94 7 Habitrol ✓ Habitrol Patch 14 mg - Up to 28 patch available on a PSO19.00 28 7 ✓ Habitrol Patch 14 mg for direct distribution only - [Xpharm]......4.52 Patch 21 mg - Up to 28 patch available on a PSO21.77 ✓ Habitrol 28 Patch 21 mg for direct distribution only - [Xpharm]......5.18 ✓ Habitrol 7 Lozenge 1 mg - Up to 216 loz available on a PSO......18.27 ✓ Habitrol 216 ✓ Habitrol 36 Lozenge 2 mg - Up to 216 loz available on a PSO......20.02 216 ✓ Habitrol ✓ Habitrol 36 ✓ Habitrol Gum 2 mg (Fruit) - Up to 384 piece available on a PSO36.39 384 Gum 2 mg (Fruit) for direct distribution only - [Xpharm].......8.64 96 ✓ Habitrol Gum 2 mg (Mint) - Up to 384 piece available on a PSO......36.39 ✓ Habitrol 384 Gum 2 mg (Mint) for direct distribution only - [Xpharm].....8.64 96 ✓ Habitrol Gum 4 mg (Fruit) - Up to 384 piece available on a PSO42.07 384 ✓ Habitrol Gum 4 mg (Fruit) for direct distribution only - [Xpharm].................10.01 96 ✓ Habitrol Gum 4 mg (Mint) - Up to 384 piece available on a PSO......42.07 ✓ Habitrol 384 ✓ Habitrol Gum 4 mg (Mint) for direct distribution only - [Xpharm]......10.01 96



Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	25.64	53 OP	✓ Varenicline Pfizer
Tab 1 mg	27.10	56	✓ Varenicline Pfizer

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to guit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see SA1667 below

Inj 25 mg vial	 271.35 ´	1	✓ Ribomustin
Inj 100 mg vial	 1,085.38	1	✓ Ribomustin
Inj 1 mg for ECP	 11.40	1 mg	✓ Baxter

⇒SA1667 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 All of the following:
 - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
 - 3.2.2 The patient has not received prior bendamustine therapy; and
 - 3.2.3 Fither:
 - 3.2.3.1 Both:
 - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more: or
 - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Fither:
 - 2.1 Both:

_					
	Subsidy		Fully	Brand or	
	(Manufacturer's Price)	Sub	sidised	Generic	
	\$	Per	✓	Manufacturer	

- 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.
 Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

macrogiobalinacinia.			
BUSULFAN - PCT - Retail pharmacy-Specialist			
Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			
Inj 10 mg per ml, 45 ml vial	32.59	1	✓ DBL Carboplatin
, , , , , , , , , , , , , , , , , , , ,	45.20		✓ Carboplatin Ebewe
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist			
Inj 100 mg vial	1.387.00	1	✓ BiCNU
ing 100 mg var		•	✓ Bicnu Heritage \$29
Inj 100 mg for ECP	1 387 00	100 mg OP	✓ Baxter
		roo mg or	- Bunton
CHLORAMBUCIL – PCT – Retail pharmacy-Specialist	20.06	25	✓ Leukeran FC
Tab 2 mg	29.06	25	• Leukeran FC
CISPLATIN - PCT only - Specialist			4
Inj 1 mg per ml, 50 ml vial		1	✓ DBL Cisplatin
	15.00		✓ Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial		1	✓ DBL Cisplatin
lei 4 mm (m FOD	21.00	4	✓ Cisplatin Ebewe
Inj 1 mg for ECP	0.25	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE			
Tab 50 mg - PCT - Retail pharmacy-Specialist	79.00	50	✓ Endoxan S29
	158.00	100	✓ Procytox S29
Wastage claimable			•
Inj 1 g vial - PCT - Retail pharmacy-Specialist	35.65	1	✓ Endoxan
	127.80	6	Cytoxan
Inj 2 g vial – PCT only – Specialist		1	Endoxan
Inj 1 mg for ECP - PCT only - Specialist	0.04	1 mg	✓ Baxter
IFOSFAMIDE - PCT only - Specialist			
Inj 1 g	96.00	1	✓ Holoxan
Inj 2 g	180.00	1	✓ Holoxan
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
LOMUSTINE - PCT - Retail pharmacy-Specialist			
Cap 10 mg	132.59	20	✓ CeeNU
Cap 40 mg		20	✓ CeeNU
MELPHALAN			
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	✓ Alkeran
Inj 50 mg - PCT only - Specialist		1	✓ Alkeran
,	213.00	•	✓ Alkeran s29 S29
	420.00		✓ Tillomed \$29
	420.00		+ I IIIOIIIEU 323

	Subsidy		Fully	/ Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
OXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	/	Oxaliplatin Actavis
.,				100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	46.32	1	✓	Oxaliplatin Accord
Inj 1 mg for ECP		1 mg	✓	Baxter
THIOTEPA - PCT only - Specialist		•		
Inj 15 mg vial	CBS	1	✓	Bedford S29
, •		•	THIO-TEPA S29	
			•	Tepadina S29
Inj 100 mg vial	CBS	1	✓	Tepadina S29
Antimetabolites				
AZACITIDINE - PCT only - Specialist - Special Authority see SA	1467 below			
Inj 100 mg vial		1	1	Azacitidine Dr
, ,				Reddy's
	605.00		/	Vidaza
Inj 1 mg for ECP		1 mg		Baxter
", ' "g o Lo "		9	,	Bunto

⇒SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- 4 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

	Subsidy		Fully Brand or
(M	anufacturer's Price \$	e) S Per	Subsidised Generic Manufacturer
ALCIUM FOLINATE			
Tab 15 mg - PCT - Retail pharmacy-Specialist	114.69	10	✓ DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17.10	5	✓ Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specialist	7.28	1	✓ <u>Calcium Folinate</u> <u>Sandoz</u>
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist		1	Calcium Folinate Sandoz
Inj 100 mg - PCT only - Specialist	7.33	1	Calcium Folinate Ebewe
Inj 300 mg - PCT only - Specialist	22.51	1	Calcium Folinate Ebewe
Inj 10 mg per ml, 35 ml vial – PCT only – Specialist	25.14	1	Calcium Folinate Sandoz
Inj 1 g - PCT only - Specialist	67.51	1	Calcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	72.00	1	✓ Calcium Folinate Sandoz
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	✓ Baxter
APECITABINE - Retail pharmacy-Specialist			
Tab 150 mg	10.00	60	✓ Capercit
Tab 500 mg	49.00	120	✓ Capercit
LADRIBINE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml		1	✓ Leustatin
Inj 10 mg for ECP	749.96	10 mg Ol	P ✓ Baxter
/TARABINE			
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist Inj 100 mg per ml, 20 ml vial – PCT – Retail	400.00	5	✓ Pfizer
pharmacy-Specialist		1	✓ Pfizer
Inj 1 mg for ECP - PCT only - Specialist		10 mg	✓ Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Specialist	80.00 1	00 mg O	P ✓ Baxter
UDARABINE PHOSPHATE			
Tab 10 mg - PCT - Retail pharmacy-Specialist		20	✓ Fludara Oral
Inj 50 mg vial - PCT only - Specialist		5	✓ Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist	115.29	50 mg Ol	P Baxter
UOROURACIL			
Inj 50 mg per ml, 20 ml vial - PCT only - Specialist		1	✓ Fluorouracil Ebewe
Inj 50 mg per ml, 100 ml vial - PCT only - Specialist		1	✓ Fluorouracil Ebewe
Inj 1 mg for ECP - PCT only - Specialist	0.66	100 mg	✓ Baxter
EMCITABINE HYDROCHLORIDE - PCT only - Specialist			
Inj 1 g, 26.3 ml vial	62.50	1	DBL Gemcitabine
lnj 1 g	15.89	1	✓ Gemcitabine Ebewe
	349.20		✓ Gemzar
Inj 1 mg for ECP	0.02	1 mg	✓ Baxter
Gemzar Inj 1 g to be delisted 1 September 2020)			

	Subsidy		Fully	Brand or
	(Manufacturer's Price	e) Sub	sidised	Generic
	\$	Per	•	Manufacturer
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist				
Inj 20 mg per ml, 5 ml vial	71.44	1	✓ Ir	rinotecan
, - 3,				Accord \$29
			√ le	inotecan Actavis
			• "	100
	100.00			
	100.00			rinotecan-Rex
Inj 1 mg for ECP	0.75	1 mg	✓ B	Baxter
MERCAPTOPURINE				
Tab 50 mg - PCT - Retail pharmacy-Specialist	37.00	25	√ <u>P</u>	uri-nethol
Oral suspension 20 mg per ml - Retail pharmacy-Specialis	t –			
Special Authority see SA1725 below	428.00 1	00 ml OP	✓ A	Ilmercap

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

R A	_	ГΗ	\sim	т	-	۸ ٦	_

IVIL	MOTIENATE		
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist8.05	90	✓ Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist31.75	90	✓ Trexate
*		5	✓ Hospira
*	Inj 7.5 mg prefilled syringe14.61	1	✓ Methotrexate
			Sandoz
*	Inj 10 mg prefilled syringe14.66	1	✓ Methotrexate
	, , , ,		Sandoz
*	Inj 15 mg prefilled syringe14.77	1	✓ Methotrexate
	, 01		Sandoz
*	Inj 20 mg prefilled syringe14.88	1	✓ Methotrexate
	, , , , , , , , , , , , , , , , , , , ,		Sandoz
*	Inj 25 mg prefilled syringe14.99	1	✓ Methotrexate
	,g p		Sandoz
*	Inj 30 mg prefilled syringe15.09	1	✓ Methotrexate
	, , g.		Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist30.00	5	✓ DBL Methotrexate
	,g p,		Onco-Vial
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist45.00	1	✓ DBL Methotrexate
	,,	•	Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist25.00	1	✓ Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail	·	
-,-	pharmacy-Specialist	1	✓ Methotrexate Ebewe
	Methotrexate Ebewe to be Sole Supply on 1 October 2020	•	inothotioxato Ebono
*		1 mg	✓ Baxter
*	Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist4.73	5 mg OP	✓ Baxter
	METREXED – PCT only – Specialist – Special Authority see SA1679 on the r	•	
Г	Inj 100 mg vial60.89	1 text page	✓ Juno Pemetrexed
	Inj 500 mg vial	1	✓ Juno Pemetrexed
	Inj 300 fig Viai	1 mg	✓ Baxter
	iij i iig ioi Loi	ı iliy	- Payrei

Sub	sidy Fu	ully Brand or
(Manufactu	ırer's Price) Subsidis	sed Generic
	\$ Per	✓ Manufacturer

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: All of the following:

1 No evidence of disease progression; and

THIOGUANINE - PCT - Retail pharmacy-Specialist

- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

Tab 4	0 mg	126.31	25	✓ Lanvis	
Other	Cytotoxic Agents				
AMSACR	INE - PCT only - Specialist				
lnj 50	mg per ml, 1.5 ml ampoule	1,500.00	6	✓ Amsidine \$29	
		4,736.00		✓ Amsidine S29	
lnj 75	mg	1,250.00	5	✓ AmsaLyo S29	
ANAGREI	LIDE HYDROCHLORIDE - PCT - Retail pharma	acy-Specialist			
	0.5 mg		100	✓ Agrylin S29 S29	
	•			✓ Teva S29	
		1,175.87		✓ Agrylin	
ARSENIC	TRIOXIDE - PCT only - Specialist				
lnj 1 r	ng per ml, 10 ml vial	4,817.00	10	✓ Phenasen	
lnj 10	mg for ECP	481.70	10 mg OP	✓ Baxter	

	Subsidy		Fully	Brand or
	Manufacturer's Pr	ice) Su Per	bsidised	Generic Manufacturer
	Ψ	rei		iviariulaciurei
BLEOMYCIN SULPHATE - PCT only - Specialist				
Inj 15,000 iu, vial	161.01	1	✓ [OBL Bleomycin
				Sulfate
Inj 1,000 iu for ECP	12.45	1,000 iu	✓ E	Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority see SA	1889 below			
Inj 3.5 mg vial	105.00	1	✓ E	Bortezomib
,				Dr-Reddy's
Inj 1 mg for ECP	31.20	1 mg	✓ E	Baxter
SA1880 Special Authority for Subsidy		•		

SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.

Note: Indications marked with * are unapproved indications

Note: Indications marked with ^ are unapproved indications.			
COLASPASE [L-ASPARAGINASE] - PCT only - Specialist			
Inj 10,000 iu	102.32	1	✓ Leunase
Inj 10,000 iu for ECP		10,000 iu OP	✓ Baxter
(Leunase Inj 10,000 iu to be delisted 1 December 2020)		.,	
(Baxter Inj 10,000 iu for ECP to be delisted 1 December 2020)			
DACARBAZINE – PCT only – Specialist	CO 70	4	/ DDI Desemberine
Inj 200 mg vial		1	✓ DBL Dacarbazine
	580.60	10	Dacarbazine
			APP S29
Inj 200 mg for ECP	62.70	200 mg OP	✓ Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist			
Inj 0.5 mg vial	255.00	1	✓ Cosmegen
Inj 0.5 mg for ECP		0.5 mg OP	✓ Baxter
. •	200.00	o.o mg or	- Dunioi
DAUNORUBICIN – PCT only – Specialist	140.50	4	✓ Pfizer
Inj 2 mg per ml, 10 ml		1	
Inj 20 mg for ECP	149.50	20 mg OP	✓ Baxter
DOCETAXEL - PCT only - Specialist			
Inj 10 mg per ml, 2 ml vial	12.40	1	✓ DBL Docetaxel
Inj 20 mg	48.75	1	 Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial	26.95	1	✓ DBL Docetaxel
Inj 20 mg per ml, 4 ml vial	26.95	1	✓ Docetaxel
			Accord S29
Inj 80 mg	195.00	1	✓ Docetaxel Sandoz
Inj 1 mg for ECP		1 mg	✓ Baxter
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist		9	
Inj 2 mg per ml, 5 ml vial	10.00	1	✓ Doxorubicin Ebewe
, , ,		1	✓ Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial	17.00	1	✓ Arrow-Doxorubicin
lai O ann ann an FO anl sinl		4	
Inj 2 mg per ml, 50 ml vial		1	✓ Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	✓ Doxorubicin Ebewe
laid as a few FOR	65.00	4	✓ Arrow-Doxorubicin
Inj 1 mg for ECP	0.29	1 mg	✓ Baxter

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
PIRUBICIN HYDROCHLORIDE - PCT only - Specialist	Ψ	1 01		Manadada
Inj 2 mg per ml, 5 ml vial	25.00	1	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1		Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1		Epirubicin Ebewe
Inj 1 mg for ECP		1 mg	_	Baxter
TOPOSIDE		9		
Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	1	Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10		Vepesid
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Spec		1		Rex Medical
Inj 1 mg for ECP – PCT only – Specialist		1 mg	_	Baxter
TOPOSIDE PHOSPHATE – PCT only – Specialist		9		
Inj 100 mg (of etoposide base)	40.00	1	1	Etopophos
Inj 1 mg (of etoposide base) for ECP		1 mg	_	Baxter
, , ,		9	-	Duxtor
YDROXYUREA – PCT – Retail pharmacy-Specialist	21.76	100	./	Lludroo
Cap 500 mg	31.70	100	•	Hydrea
DARUBICIN HYDROCHLORIDE	20.00		,	
Inj 5 mg vial – PCT only – Specialist		1		Zavedos
Inj 10 mg vial – PCT only – Specialist		_ 1	_	Zavedos
Inj 1 mg for ECP - PCT only - Specialist	21.84	1 mg	•	Baxter
ENALIDOMIDE – Retail pharmacy-Specialist – Special Autho	ority see SA1897 below	1		
Wastage claimable			_	
Cap 5 mg		28		Revlimid
Cap 10 mg		21		Revlimid
0 45	6,207.00	28		Revlimid
Cap 15 mg		21		Revlimid
0 05	7,239.18	28		Revlimid
Cap 25 mg		21	•	Revlimid

⇒SA1897 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Either:
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and

Sub	bsidy F	ully Br	and or
(Manufactu	urer's Price) Subsidis	sed Ge	eneric
•	\$ Per	✓ Ma	anufacturer

continued...

- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 The patient has ECOG performance score of 0-1; and
- 5 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MESNA

Tab 400 mg - PCT - Retail pharmacy-Specialist	314.00	50	✓ Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist	448.50	50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist	177.45	15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist	407.40	15	✓ Uromitexan
Inj 1 mg for ECP - PCT only - Specialist	2.96	100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist			
Inj 5 mg vial	851.37	1	✓ Teva
Inj 20 mg vial	816.32	1	✓ Omegapharm \$29
Inj 1 mg for ECP		1 mg	✓ Baxter
(Omegapharm \$29 Inj 20 mg vial to be delisted 1 November 2020,)		
MITOZANTRONE - PCT only - Specialist			
Inj 2 mg per ml, 10 ml vial	97.50	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
OLAPARIB - Retail pharmacy-Specialist - Special Authority see S.	A1883 below		
Tab 100 mg	3,701.00	56	✓ Lynparza
Tab 150 mg		56	✓ Lynparza
Cap 50 mg - Wastage claimable	7,402.00	448	✓ Lynparza

⇒SA1883 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Patient has received at least two lines of previous treatment with platinum-based chemotherapy; and
- 4 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line of platinum-based chemotherapy; and
- 5 Patient's disease must have achieved partial or complete response to treatment with the immediately preceding platinum-based regimen; and

	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per	Manufacturer
continued			

- 6 Patient's disease has not progressed following prior treatment with olaparib; and
- 7 Treatment will be commenced within 8 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 8 Treatment to be administered as maintenance treatment; and
- 9 Treatment not to be administered in combination with other chemotherapy.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 No evidence of progressive disease; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy.

Note: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

PACLITAXEL - PCT only - Specialist			
Inj 30 mg	47.30	5	✓ Paclitaxel Ebewe
Inj 100 mg	24.00	1	✓ Paclitaxel Ebewe
•	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
•	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg	44.00	1	✓ Paclitaxel Ebewe
	275.00		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority se	ee SA1325 below		
Inj 750 iu per ml, 5 ml vial	3,455.00	1	✓ Oncaspar LYO S29

⇒SA1325 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and
- 3 Treatment is with curative intent.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and
- 3 Treatment is with curative intent.

		OXYCOFORMYCIN] - PCT only - Specialist	PENTOSTATIN [DEOX
✓ Nipent S29	1	CBS	Inj 10 mg
		YDROCHLORIDE - PCT - Retail pharmacy-Specialist	PROCARBAZINE HYD
✓ Natulan S29	50	980.00	Can 50 mg

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	✓	Manufacturer
TEMOZOLOMIDE - Special Authority see SA1741 below - Reta	ail pharmacy			
Cap 5 mg		5	✓	Temaccord
Cap 20 mg	16.38	5	✓	Temaccord
	18.30		✓	Apo-Temozolomide
	136.00	14	✓	Accord S29
Cap 100 mg	35.98	5	✓	Temaccord
	40.20		✓	Apo-Temozolomide
	532.00	14	✓	Accord S29
Cap 140 mg	50.12	5	✓	Temaccord
	400.00		✓	Amneal S29
Cap 180 mg	620.00	14	✓	Accord S29
Cap 250 mg		5	✓	Temaccord
•	688.00		1	Amneal \$29

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 No evidence of disease progression; and

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	√	

continued...

2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

Cap 10 mg - PCT - Retail pharmacy-Specialist479.50	100	✓ Vesanoid
VENETOCLAX - Retail pharmacy-Specialist - Special Authority see SA1868 belo	W	
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg1,771.86	42 OP	✓ Venclexta
Tab 10 mg95.78	14 OP	✓ Venclexta
Tab 50 mg239.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable8,209.41	120	✓ Venclexta

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Sub	bsidy F	ully Br	Brand or
(Manufactu	urer's Price) Subsidis	sed Ge	eneric
•	\$ Per	✓ Ma	anufacturer

continued...

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

VINBLASTINE SULPHATE

Inj 1 mg per mi, 10 mi viai – PC1 – Retail pharmacy-Specialist270.37	5	✓ DBL VInblastine S29
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Hospira✓ Baxter
Inj 1 mg per ml, 1 ml vial – PCT – Retail pharmacy-Specialist74.52	5	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist102.73	5	DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	✓ Baxter
VINORELBINE - PCT only - Specialist		
Inj 10 mg per ml, 1 ml vial	1	✓ Navelbine✓ Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial56.00 210.00	1	✓ Navelbine✓ Vinorelbine Ebewe
Inj 1 mg for ECP1.25	1 mg	✓ Baxter

Protein-tyrosine Kinase Inhibitors

ALECTINIB - Retail pharmacy-Specialist - Special Authority see SA1870 below Wastage claimable 224

Alecensa

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid

continued...

DRI Vinhlactina 920

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

continued...

for 6 months for applications meeting the following criteria:

Roth:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB – Special Authority see SA1805 below – Retail pharmacy

wastage ciaimable			
Tab 20 mg	3,774.06	60	Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
•	7,692.58	60	✓ Sprycel
	,		-1. 7

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
 - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
 - 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

⇒SA1915 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and

✓ Iressa

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	Manufacturer

continued...

- 3 Either:
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued defitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

GEFITINIB - Retail pharmacy-Specialist - Special Authority see SA1916 below	
Tab 250 mg1,700.00	30

⇒SA1916 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
 - 2.1 Patient is treatment naive: or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib: and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Gefitinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

IMATINIB MESII ATE

Note: Imatinib-AFT is not a registered for the treatment of Gastro Intestinal Stromal Tumours (GIST). The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule.

Tab 100 mg - [Xpharm] - Special Authority see SA1460 on the

	next page	2,400.00	60	✓ Glivec
*	Cap 100 mg		60	✓ Imatinib-AFT
*	Cap 400 mg	197.50	30	✓ Imatinib-AFT

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 PHARMAC Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

Special Authority criteria for GIST – access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA1191 below - Retail pharmacy

Tab 250 mg1,899.00 70 **✓ Tykerb**

⇒SA1191 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient has not previously received trastuzumab treatment for HER 2 positive metastatic breast cancer; and
 - 1.3 Lapatinib not to be given in combination with trastuzumab; and
 - 1.4 Lapatinib to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2.2 The patient started trastuzumab for metastatic breast cancer but discontinued trastuzumab within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on trastuzumab; and
 - 2.4 Lapatinib not to be given in combination with trastuzumab; and
 - 2.5 Lapatinib to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 on the next page - Retail pharmacy

Wastage claimable

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Fither:
 - 2.1 Patient has documented CML treatment failure* with imatinib; or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PALBOCICLIB - Retail pharmacy-Specialist - Special Authority see SA1894 below

wastage ciaimable			
Cap 75 mg	4,000.00	21	✓ Ibrance
Cap 100 mg	4,000.00	21	✓ Ibrance
Cap 125 mg	4.000.00	21	✓ Ibrance

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Either:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state: and
- 4.2.2 Fither:
 - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
 - 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PAZOPANIB – Special Authority see SA1190 below – Retail pha	ırmacy			
Tab 200 mg	1,334.70	30	✓ \	/otrient
Tab 400 mg	2,669.40	30	•	/otrient

SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

wastage cialmable			
Tab 5 mg	2,500.00	56	Jakavi
Tab 15 mg	5,000.00	56	Jakavi
Tab 20 mg	5.000.00	56	✓ Jakavi

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis: and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

- 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS: and
- 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

Cap 12.5 mg2,315.38	28	✓ Sutent
Cap 25 mg	28	✓ Sutent
Cap 50 mg	28	✓ Sutent

⇒SA1917 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib: or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 82

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA1914 below

Wastage claimable

Tab 250 mg4,276.19 120 ✓ Zytiga

⇒SA1914 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases: and
- 3 Patient's disease is castration resistant; and
- 4 Fither:

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	Subsi	idised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of clinical disease progression; and
- 2 No initiation of taxane chemotherapy with abiraterone; and
- 3 The treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease: and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

MEGESTROL ACETATE Tab 160 mg63.53	30	✓ Apo-Megestrol
OCTREOTIDE		
Inj 100 mcg per ml, 1 ml ampoule18.69	5	✓ Octreotide GH S29
Inj 50 mcg per ml, 1 ml vial30.64	5	✓ DBL Octreotide
		Octreotide
		MaxRx S29
Inj 100 mcg per ml, 1 ml vial18.69	5	✓ DBL Octreotide
Inj 500 mcg per ml, 1 ml vial72.50	5	✓ DBL Octreotide
222.00		Octreotide
		(Sun) S29

	\$	Per	✓ Manufacturer
OCTREOTIDE LAR (SOMATOSTATIN ANALOGUE)	- Special Authority see SA191	8 below	- Retail pharmacy
Inj LAR 10 mg prefilled syringe	1,772.50	1	✓ Sandostatin LAR
Inj LAR 20 mg prefilled syringe		1	Sandostatin LAR
Inj LAR 30 mg prefilled syringe		1	Sandostatin LAR

Subsidy

(Manufacturer's Price)

Fully

Subsidised

Brand or

Generic

⇒SA1918 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed: or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery: or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
- 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for

applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

TAMOXIFEN CITRATE

*	Tab 10 mg	15.00	60	✓ Tamoxifen Sandoz
	Tamoxifen Sandoz to be Sole Supply on 1 November 2020			
*	Tab 20 mg	6.65	60	✓ Tamoxifen Sandoz
	Tamoxifen Sandoz to be Sole Supply on 1 November 2020			

Aromatase Inhibitors

ANASTROZOLE	30	✓ Rolin
EXEMESTANE	30	✓ Pfizer Exemestane
LETROZOLE	30	✓ Letrole

Immunosuppressants

Cytotoxic Immunosuppressants

AΖ	ATHIOPRINE
*	Tah 25 mg

	1 ab 20 mg	7 .00	00	- /tEaman
*	Tab 50 mg	7.60	100	✓ Azamun
	Inj 50 mg vial		1	✓ Imuran
MY	COPHENOLATE MOFETIL			
	Tab 500 mg	35.90	50	✓ Cellcept
	Cap 250 mg		100	✓ Cellcept

Powder for oral liq 1 g per 5 ml - Subsidy by endorsement........... 187.25

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

Fusion Proteins

ETANERCEPT - Special Authority see SA1949 on the ne.	xt page – Retail pharmacy	,	
Inj 25 mg	690.00	4	✓ Enbrel
Inj 50 mg autoinjector	1,050.00	4	✓ Enbrel
Inj 50 mg prefilled syringe	1,050.00	4	✓ Enbrel

✓ ∆zamıın

✓ Cellcept

165 ml OP

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

⇒SA1949 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
- 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
- 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient diagnosed with Juvenile Idiopathic Arthritis (JIA); and
 - 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
 - 2.5 Both:
 - 2.5.1 Either:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender

continued...

ioints: or

- 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
- 2.5.2 Physician's global assessment indicating severe disease.

Renewal — (juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a named specialist or rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
\$	Per	1	Manufacturer

continued...

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
 - 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
 - 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis: or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Fither:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Fither:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and

Subsidy		Fully	Brand or	
(Manufacturer's Price	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Fither:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Fither:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
- 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Spec	cialist		
Inj 50 mg per ml, 5 ml	2,351.25	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT onl	y – Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG S29
(SII-Onco-BCG \$29 Ini 40 mg per ml. vial to be delisted 1 Ap	ril 2021)		

Monoclonal Antibodies

		below – Retail pharmacy	ADALIMUMAB – Special Authority see SA1950 be
Humira	2	1,599.96	Inj 20 mg per 0.4 ml prefilled syringe
✓ HumiraPen	2	1,599.96	Inj 40 mg per 0.8 ml prefilled pen
Humira	2	1,599.96	Inj 40 mg per 0.8 ml prefilled syringe

⇒SA1950 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD): or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ıbsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992:19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroillitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab;
 - 2.1.2 CDAI score is 150 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or

Subsidy	F	ılly	Brand or
(Manufacturer's	s Price) Subsidis	sed	Generic
\$	Per	✓	Manufacturer

continued...

- 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab: or
 - 2.1.2 PCDAI score is 15 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's disease.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage III or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 The patient has 3 or more active lesions (e.g. inflammatory nodules, abscesses, draining fistulae); and
- 4 The patient has a Dermatology Quality of Life Index of 10 or more and the assessment is no more than 1 month old at time of application; and
- 5 Following the initial loading doses, adalimumab is to be administered at doses no greater than 40mg every 7 days.

Renewal — (hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for juvenile idiopathic arthritis (JIA); and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for juvenile idiopathic arthritis; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient diagnosed with JIA; and
 - 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
 - 2.5 Both:
 - 2.5.1 Either:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

Renewal — (juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a named specialist or rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Fither:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for psoriatic arthritis; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following: 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
 - 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
 - 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g.
- prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement: and
- 2 Patient continues to require treatment: and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis: and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis: or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
 - 2.6 Fither:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Fither:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease that is significantly impacting the patient's quality of life (see Notes); and
- 2 Fither
 - 2.1 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has not responded adequately to treatment with infliximab (see Notes); or
 - 2.2 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has experienced intolerable side effects from treatment with infliximab; and
- 3 The patient is experiencing significant loss of quality of life; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: Behcet's disease diagnosed according to the International Study Group for Behcet's disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al. J Rheumatol. 2004;31:931-7.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept: or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline valuee; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for severe ocular inflammation; or
- 2 Both:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

- 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
- 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

- Both:
 - 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
 - 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

AFLIBERCEPT - Special Authority see SA1772 below - Retail pharmacy

1 ✓ Eylea

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab: or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Fither:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	

continued...

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

		- PCT only - Specialist - Special Authority see SA1697 below	CEI
Erbitux	1	per ml, 20 ml vial364.00	
Erbitux	1	per ml, 100 ml vial	
✓ Baxter	1 mg	or ECP	

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB – PCT only – Special Authority see SA1951 below	
Ini 100 ma	

Inj 100 mg	806.00	1	✓ Remicade
Inj 1 mg for ECF	98.29	1 mg	Baxter

⇒SA1951 Special Authority for Subsidy

Initial application — (Graft vs host disease) from any medical practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria

(Mar	Subsidy nufacturer's Price)	Sub	Fully sidised	Brand or Generic
	\$	Per	1	Manufacturer

continued...

for adalimumab and/or etanercept; and

3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for psoriatic arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for osoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
 - 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Initial application — (severe ocular inflammation) from any medical practitioner. Approvals valid for 3 doses for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation: or
- 2 Roth:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (chronic ocular inflammation) from any medical practitioner. Approvals valid for 3 doses for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to

-	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
=	<u> </u>				

continued...

achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (Pulmonary sarcoidosis) from any medical practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- ...
 1 Patient has life-threatening pulmonary sarcoidosis that is refractory to other treatments; and
- 2 Treatment is to be prescribed by, or has been recommended by, a physician with expertise in the treatment of pulmonary sarcoidosis.

Initial application — (Crohn's disease (adults)) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initial application — (**Crohn's disease (children)**) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Fither:

Subsidy	Fu	,	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
- 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Initial application — (severe ulcerative colitis) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Fither:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
 - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (severe ulcerative colitis) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Fither:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initial application — (plaque psoriasis) only from a dermatologist. Approvals valid for 3 doses for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plaque psoriasis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plague psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, cyclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) only from a dermatologist. Approvals valid for 3 doses for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (neurosarcoidosis) only from a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Biopsy consistent with diagnosis of neurosarcoidosis; and
- 2 Patient has CNS involvement: and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Fither:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Either:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (severe Behcet's disease) from any medical practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes:

- a) Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.
- b) Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any medical practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved guality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pvoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement: and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

MEPOLIZUMAB - Special Authority see SA1896 below - Retail pharmacy

⇒SA1896 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10 9 cells/L in the last 12 months; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- - 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
 - 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB - PCT only - Specialist - Special Author	ority see SA1627 below		
Inj 25 mg per ml, 40 ml vial	5,910.00	1	✓ Gazyva
Inj 1 mg for ECP	6.21	1 mg	✓ Baxter

⇒SA1627 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and</p>
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL: and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

OMALIZUMAB - Special Authority see SA1744 on the next pag	e – Retail pharmac	y	
Inj 150 mg prefilled syringe	450.00	1	✓ Xolair
Inj 150 mg vial		1	✓ Xolair

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Patient has previously adequately responded* to 6 doses of omalizumab; or

continued...

- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PERTUZUMAB - PCT only - Specialist - Special Authorit	y see SA1606 below		
Inj 30 mg per ml, 14 ml vial	3,927.00	1	✓ Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	✓ Baxter

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist	 Special Authority see SA190 	1 below	
Inj 100 mg per 10 ml vial	1,075.50	2	✓ Mabthera
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

⇒SA1901 Special Authority for Subsidy

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment;
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Fither:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Su	ubsidy	Fully	Brand or
(Manufac	cturer's Price) Subsid	lised	Generic
	\$ Per	✓	Manufacturer

continued...

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physiciann; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used: and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

Subsidy	y Full	/ Brand or
(Manufacturer's	s Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*: and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - S	pecial Authority see SA1937 be	elow	
Inj 100 mg per 10 ml vial	275.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Ini 1 mg for ECP	1 22	1 ma	✓ Rayter (Rivimyo)

⇒SA1937 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — **(ANCA associated vasculitis)** from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive: or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 222 Both
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Fither:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL;

Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
- 1.2.3 The patient does not have chromosome 17p deletion CLL; and
- 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sul	bsidised	Generic
\$	Per	1	Manufacturer

continued...

500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and

- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks. or two 1.000 mg doses given two weeks apart: and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Fither:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*: and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*: and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer
--

continued...

All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles: or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — **(haemophilia with inhibitors)** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre: or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*: and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AlHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or

Subsidy (Manufacturer's Price)	Su	Fully ibsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 x 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	sidised Generic
--	-----------------

continued...

- 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
- 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

⇒SA1754 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4

	Subsidy		Fully	Brand or
(Ma	anufacturer's Price)		ubsidised	Generic
	\$	Per		Manufacturer

continued...

months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB - PCT only - Special Authority see SA1858 on the next page

Inj 20 mg per m	nl, 4 ml vial	220.00	1	✓ Actemra
, , ,		550.00	1	✓ Actemra
		1,100.00	1	✓ Actemra
, , ,		2.85	1 mg	✓ Baxter
, ,			0	

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1858 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy

(Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses. **Initial application — (previous use)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis: or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	Fi	ılly	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	/	Manufacturer	

continued...

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Fither:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...
Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for juvenile idiopathic arthritis (JIA); and
- 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or

2 All of the following:

- 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
- 2.2 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
- 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
- 2.4 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
- 2.5 Both:
 - 2.5.1 Either:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender ioints: or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by

Subsidy	:)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
<u> </u>	Per	✓	Manufacturer

continued...

toxicity or intolerance; and

- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB - PCT only - Specialist - Special Authority see SA1632 below

Inj 150 mg vial	1,350.00	1	✓ Herceptin
Inj 440 mg vial	3,875.00	1	✓ Herceptin
Inj 1 mg for ECP	9.36	1 mg	✓ Baxter

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:
All of the following:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA1871 below

Inj 100 mg vial	 2,320.00	1	✓ Kadcyla
Inj 160 mg vial	 3,712.00	1	✓ Kadcyla
Inj 1 mg for ECP	 23.20	1 mg	✓ Baxter

⇒SA1871 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4 Patient has a good performance status (ECOG 0-1); and
 - 5 Fither
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
 - 6 Treatment to be discontinued at disease progression.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

Programmed Cell Death-1 (PD-1) Inhibitors

NIVOLUMAB - PCT only - Specialist - Special Authority see SA1911 below

Inj 10 mg per ml, 4 ml vial	1,051.98	1	Opdivo
Inj 10 mg per ml, 10 ml vial	2,629.96	1	Opdivo
Inj 1 mg for ECP	27.62	1 mg	✓ Baxter

⇒SA1911 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:

(Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer		Subsidy	F	ılly	Brand or
\$ Per ✓ Manufacturer	(N	fanufacturer's Price)	Subsidis	ed	
		\$	Per	<u> </u>	Manufacturer

continued...

- 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
- 2.2 Patient has signs of disease progression; and
- 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB – PCT	「only – Specialist – Special Authority see SA1910	0 below	
Inj 25 mg per ml, 4 ml v	vial4,680.00	00 1 ✓ Keytru	da
Inj 1 mg for ECP	49.14	4 1 mg ✓ Baxter	

⇒SA1910 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	✓	Manufacturer

continued...

- 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
- 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
- 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
- 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

CICLOSPORIN		
Cap 25 mg44.63	50	✓ Neoral
Cap 50 mg88.91	50	Neoral
Cap 100 mg177.81	50	✓ Neoral
Oral liq 100 mg per ml198.13	50 ml OP	Neoral
EVEROLIMUS - Special Authority see SA1913 below - Retail pharmacy		
Wastage claimable		
Tab 10 mg6,512.29	30	Afinitor
Tab 5 mg4,555.76	30	Afinitor

⇒SA1913 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Everolimus to be discontinued at progression of SEGAs; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Note: : MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

SIROLIMUS - Special Authority see SA0866 below - Retail pharmacy

Tab 1 mg	749.99	100	Rapamune
Tab 2 mg	1,499.99	100	✓ Rapamune
Oral lig 1 mg per ml	449.99	60 ml OP	✓ Rapamune

⇒SA0866 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- · Rapidly progressive transplant vasculopathy; or
- · Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP; or
- · Leukoencepthalopathy; or
- · Significant malignant disease

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg		100	✓ Tacrolimus Sandoz
Cap 1 mg		100	✓ Tacrolimus Sandoz
Cap 5 mg	248.20	50	✓ Tacrolimus Sandoz

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Antiallergy Preparations

Allergic Emergencies

ICATIBANT - Special Authority see SA1558 below - Retail pharmacy

Inj 10 mg per ml, 3 ml prefilled syringe......2,668.00 1 ✓ Firazyr

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Maintenance kit - 6 vials 120 mcg freeze dried venom, with

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above - Retail pharmacy

diluent	285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent			
9 ml, 3 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with dilue	nt305.00	1 OP	✓ Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority see	SA1367 above	- Retail pharr	macy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			•
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze			
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29

	Subsidy		Fully Brand or
	(Manufacturer's P		,
	\$	Per	✓ Manufacturer
Antihistamines			
Antimotalinios			
CETIRIZINE HYDROCHLORIDE			
* Tab 10 mg	1 12	100	✓ Zista
* Oral liq 1 mg per ml		200 ml	✓ Histaclear
	2.00	200 1111	- Instantai
CHLORPHENIRAMINE MALEATE			
* Oral liq 2 mg per 5 ml	9.37	500 ml	✓ Histafen
DEXTROCHLORPHENIRAMINE MALEATE			
* Tab 2 mg	2.02	40	
	(8.40)		Polaramine
	1.01	20	rolaranino
	(5.99)	20	Polaramine
₩ Orol lig 2 mg nor E ml		100 ml	Folarallille
* Oral liq 2 mg per 5 ml		100 1111	Deleventine
	(10.29)		Polaramine
FEXOFENADINE HYDROCHLORIDE			
* Tab 60 mg	4.34	20	
·	(8.23)		Telfast
* Tab 120 mg		10	
	(8.23)		Telfast
	14.22	30	Tondot
	(26.44)	00	Telfast
	(20.44)		Tollast
LORATADINE			
* Tab 10 mg		100	✓ Lorafix
* Oral liq 1 mg per ml	2.15	120 ml	✓ Lorfast
PROMETHAZINE HYDROCHLORIDE			
* Tab 10 mg	1 68	50	✓ Allersoothe
* Tab 25 mg		50	✓ Allersoothe
* Oral lig 1 mg per 1 ml		100 ml	✓ Allersoothe
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a		5	✓ Hospira
* Inj 25 mg per mi, 2 mi ampoule – op to 5 mj avaliable on a	F3U 17.07	5	• поѕрна
Inhaled Corticosteroids			
Illialed Corticosterolds			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler, 50 mcg per dose	0.30	200 dose OP	✓ Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP	✓ Beclazone 50
		200 dose OP	✓ Qvar
Aerosol inhaler, 100 mcg per dose			
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP	✓ Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free	22.67	200 dose OP	✓ Beclazone 250
BUDESONIDE			
Powder for inhalation, 100 mcg per dose	17.00	200 dose OP	✓ Pulmicort
,gr	- *		Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00	200 dose OP	✓ Pulmicort
1 Officer for initial attorn, 200 mby per dose	13.00	200 0036 OF	Turbuhaler
Develop for the letters 400 means and as	00.00	000 -1 05	
Powder for inhalation, 400 mcg per dose	32.00	200 dose OP	✓ Pulmicort
			Turbuhaler

	Subsidy		Fully	Brand or
	(Manufacturer's		idised	Generic
	\$	Per	1	Manufacturer
LUTICASONE				
Aerosol inhaler, 50 mcg per dose	4.68	120 dose OP	1	Floair
	7.19		1	Flixotide
Flixotide to be Sole Supply on 1 September 2020				
Powder for inhalation, 50 mcg per dose	7.50	60 dose OP	1	Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP	1	Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP	1	Floair
• • • • • • • • • • • • • • • • • • • •	13.60		1	Flixotide
Flixotide to be Sole Supply on 1 September 2020				
Aerosol inhaler, 250 mcg per dose	10 18	120 dose OP	1	Floair
7 to room whater, 200 mag par dood	24.62	120 0000 01		Flixotide
Flixotide to be Sole Supply on 1 September 2020	L-1.0L		-	. IIAGUAG
Powder for inhalation, 250 mcg per dose	13 60	60 dose OP	1	Flixotide Accuhaler
		JU GUSE OF	•	I IIAOUGE ACCUIIGICI
Floair Aerosol inhaler, 50 mcg per dose to be delisted 1 Septem				
Floair Aerosol inhaler, 125 mcg per dose to be delisted 1 Septe	mber 2020)			
Floair Aerosol inhaler, 250 mcg per dose to be delisted 1 Septe	mber 2020)			
Inhalad Lang action Data advancement wenter	40			
Inhaled Long-acting Beta-adrenoceptor Agonis	is			
EFORMOTEROL FUMARATE				
Powder for inhalation, 12 mcg per dose, and monodose dev	ioo 00.64	60 dose		
Powder for inhalation, 12 mcg per dose, and monodose dev		60 dose		Га на «III
	(35.80)			Foradil
FORMOTEROL FUMARATE DIHYDRATE				
Powder for inhalation 4.5 mcg per dose, breath activated				
(equivalent to eformoterol fumarate 6 mcg metered dose	e)10.32	60 dose OP		
-	(16.90)			Oxis Turbuhaler
NDACATEROL				
Powder for inhalation 150 mcg	61 00	30 dose OP	1	Onbrez Breezhaler
Powder for inhalation 300 mcg		30 dose OP		Onbrez Breezhaler
G	01.00	JU GUSE OF	•	טווטוכב טוככבוומופו
SALMETEROL			_	_
Aerosol inhaler CFC-free, 25 mcg per dose		120 dose OP		Serevent
Aerosol inhaler 25 mcg per dose		120 dose OP		Meterol
Powder for inhalation, 50 mcg per dose, breath activated	25.00	60 dose OP		Serevent Accuhaler
Inhaled Corticosteroids with Long-Acting Beta-	-Adrenocept	or Agonists		
DUDESONIDE WITH EFORMOTERO				
BUDESONIDE WITH EFORMOTEROL	40.00	100 dess 00		Vannair
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg		120 dose OP		Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6 r	ncg33.74	120 dose OP	•	Symbicort
				Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg		120 dose OP		Vannair
Powder for inhalation 200 mcg with eformoterol fumarate 6 r	ncg 44.08	120 dose OP		Symbicort
-	=			Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate				
12 mcg – No more than 2 dose per day	44 NR	60 dose OP	1	Symbicort
12 mbg 140 more man 2 0000 per day		JU GUSE OF	•	Turbuhaler 400/12
				runumater 400/12
LUTICASONE FUROATE WITH VILANTEROL				
Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose OP	1	Breo Ellipta
•				-

			ALLLIGIEU
	Subsidy	Duite a)	Fully Brand or
	(Manufacturer's \$	Price) Subsi	dised Generic ✓ Manufacturer
FLUTICASONE WITH SALMETEROL			
Aerosol inhaler 50 mcg with salmeterol 25 mcg		120 dose OP	✓ RexAir
Seretide to be Sole Supply on 1 September 2020	25.79		✓ Seretide
Aerosol inhaler 125 mcg with salmeterol 25 mcg	16.83	120 dose OP	✓ RexAir
Ç Ç	32.60		✓ Seretide
Seretide to be Sole Supply on 1 September 2020			
Powder for inhalation 100 mcg with salmeterol 50 mcg - No more than 2 dose per day		60 dose OP	✓ Seretide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg – No			
more than 2 dose per day		60 dose OP	✓ Seretide Accuhaler
(RexAir Aerosol inhaler 50 mcg with salmeterol 25 mcg to be deli (RexAir Aerosol inhaler 125 mcg with salmeterol 25 mcg to be de			
(1.10% iii 7.101030) iiiiitatoi 120 mag witii Saimeteloi 20 mag to be de	moteu i oepteli	1001 2020)	
Beta-Adrenoceptor Agonists			
SALBUTAMOL			
Oral liq 400 mcg per ml		150 ml	✓ <u>Ventolin</u>
Infusion 1 mg per ml, 5 ml Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO		10 5	✓ Ventolin✓ Ventolin
ing 500 mag per mil, i mil – op to 5 mg available on a 1 50		J	Ventoniii
Inhaled Beta-Adrenoceptor Agonists			
SALBUTAMOL			
Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000	2.22	000 4. 07	(December
dose available on a PSO	3.80	200 dose OP	✓ Respigen✓ SalAir
	(6.00)		Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb			
available on a PSO Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb		20	✓ <u>Asthalin</u>
available on a PSO		20	✓ Asthalin
TERBUTALINE SULPHATE			
Powder for inhalation, 200 mcg per dose (equivalent to			
250 mcg metered dose), breath activated		120 dose OP 200 dose OP	✓ Bricanyl Turbuhaler
Powder for inhalation, 250 mcg per dose, breath activated (Bricanyl Turbuhaler Powder for inhalation, 250 mcg per dose, br			✓ Bricanyl Turbuhaler October 2020)
Anticholinergic Agents			
IPRATROPIUM BROMIDE			
Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dose		000 daaa 00	Atmoscomt
available on a PSO Nebuliser soln, 250 mcg per ml, 1 ml ampoule – Up to 40 ne		200 dose OP	✓ Atrovent
available on a PSO		20	✓ Univent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule - Up to 40 ne			•
available on a PSO	11.73	20	✓ <u>Univent</u>

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	

Inhaled Beta-Adrenoceptor Agonists with Anticholinergic Agents

SALBUTAMOL WITH IPRATROPIUM BROMIDE

Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg per		
dose CFC-free	200 dose OP	Duolin HFA
Nabulian ada O F and with invetoration beautide O F and add		

Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule − Up to 20 neb available on a PSO5.20 20 ✓ Duolin

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium.
- Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD, and the prescription is endorsed accordingly.

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL — Special Authority see SA1584 above — Retail pharmacy Powder for Inhalation 50 mcg with indacaterol 110 mcg.....81.00 30 dose OP ✓ Ultibro Breezhaler TIOTROPIUM BROMIDE WITH OLODATEROL — Special Authority see SA1584 above — Retail pharmacy

Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg81.00 60 dose OP ✓ Spiolto Respimat

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antifibrotics

NINTEDANIB - Special Authority see SA1928 below - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

⇒SA1928 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA1929 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

 Tab 801 mg
 3,645.00
 90
 ✓ Esbriet

 Cap 267 mg
 − Wastage claimable
 3,645.00
 270
 ✓ Esbriet

⇒SA1929 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

	Subsidy	Full	/ Brand or
(Manu	ufacturer's Price)	Subsidise	d Generic
	\$ F	Per 🗸	Manufacturer

continued...

Renewal — (idiopathic pulmonary fibrosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Leukotriene Receptor Antagonists

MC	NTELUKAST			
*	Tab 4 mg	4.25	28	✓ Montelukast Mylan
*	Tab 5 mg	4.25	28	✓ Montelukast Mylan
*	Tab 10 mg	3.95	28	✓ Montelukast Mylan
	•			✓ Montelukast Mylan

Mast Cell Stabilisers

NEDOCROMIL - Subsidy by endorsement

Subsidy by endorsement - Subsidised for patients who were taking nedocromil prior to 1 July 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of nedocromil.

Aerosol inhaler, 2 mg per dose CFC-free......28.07 112 dose OP ✓ Tilade

(Tilade Aerosol inhaler, 2 mg per dose CFC-free to be delisted 1 February 2021)

SODIUM CROMOGLICATE - Subsidy by endorsement

Subsidy by endorsement - Subsidised for patients who were taking sodium cromoglicate prior to 1 July 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of sodium cromoglicate.

112 dose OP ✓ Intal Forte CFC Free

(Intal Forte CFC Free Aerosol inhaler, 5 mg per dose CFC-free to be delisted 1 May 2021)

Methylxanthines

AMINOPHYLLINE

*	Inj 25 mg per ml, 10 ml ampoule - Up to 5 inj available on a		
	PSO124.37	5	DBL Aminophylline
TH	EOPHYLLINE		
*	Tab long-acting 250 mg23.02	100	✓ Nuelin-SR
*	Oral liq 80 mg per 15 ml16.60	500 ml	✓ <u>Nuelin</u>

Mucolytics

DORNASE ALFA - Special Authority see SA0611 below - Ret	tail pharmacy		
Nebuliser soln, 2.5 mg per 2.5 ml ampoule	250.00	6	✓ Pulmozyme

⇒SA0611 Special Authority for Subsidy

Special Authority approved by the Cystic Fibrosis Advisory Panel

Notes: Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

			EM AND ALLERGIES
	Subsidy (Manufacturer's \$		Fully Brand or idised Generic Manufacturer
continued			
The Co-ordinator, Cystic Fibrosis Advisory Panel PHARMAC, PO Box 10 254 Wellington	Phone: (04) 460 4990 Facsimile: (04) 916 757 Email: <u>CFPanel@pharm</u>		
Prescriptions for patients approved for treatment must and expertise in treating cystic fibrosis.	be written by respiratory	physicians or pae	ediatricians who have experience
SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7%	24.50	90 ml OP	✓ Biomed
Nasal Preparations			
Allergy Prophylactics			
BUDESONIDE Metered aqueous nasal spray, 50 mcg per dose SteroClear to be Sole Supply on 1 October 20		200 dose OP	✓ SteroClear
Metered aqueous nasal spray, 100 mcg per dose SteroClear to be Sole Supply on 1 October 20	2.84	200 dose OP	✓ SteroClear
FLUTICASONE PROPIONATE Metered aqueous nasal spray, 50 mcg per dose	1.98	120 dose OP	✓ Flixonase Hayfever & Allergy
IPRATROPIUM BROMIDE Aqueous nasal spray, 0.03%	4.61	15 ml OP	✓ Univent
Respiratory Devices			
MASK FOR SPACER DEVICE a) Up to 50 dev available on a PSO b) Only on a PSO c) Only for children aged six years and under Small.	2.20	1	✓ e-chamber Mask
PEAK FLOW METER a) Up to 25 dev available on a PSO b) Only on a PSO		·	- Continues mass
Low range	9.54	1	Mini-Wright AFS Low Range
Normal range	9.54	1	Mini-Wright Standard
SPACER DEVICE a) Up to 50 dev available on a PSO b) Only on a PSO			
220 ml (single patient)		1 1	✓ e-chamber Turbo✓ e-chamber LaGrande
800 ml	6.50	1	✓ Volumatic
Respiratory Stimulants			
CAFFEINE CITRATE Oral liq 20 mg per ml (10 mg base per ml)	15.10	25 ml OP	✓ Biomed

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	(Manufacturer's P	rice) Subsi	idised Generic Manufacturer
Ear Preparations			
ACETIC ACID WITH 1, 2- PROPANEDIOL DIACETATE AND BE For Vosol ear drops with hydrocortisone powder refer Stand.		ge 247	
Ear drops 2% with 1, 2-Propanediol diacetate 3% and		ge 247	
benzethonium chloride 0.02% FLUMETASONE PIVALATE	6.97	35 ml OP	✓ Vosol
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform
			ED's ✓ Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTAT	IN	- Location Violonii
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g	5 16	7.5 ml OP	✓ Kenacomb
		7:51111 01	Rendeomb
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and			
gramicidin 50 mcg per ml		8 ml OP	
FRAMYCETIN SULPHATE	(9.27)		Sofradex
Ear/Eye drops 0.5%		8 ml OP	
	(8.65)		Soframycin
Eye Preparations			
Eye preparations are only funded for use in the eye, unless expli	citly stated otherv	vise.	
Anti-Infective Preparations			
ACICLOVIR	14.00	4.5 = OD	√ Vimpoo
CHLORAMPHENICOL	14.92	4.5 g OP	✓ ViruPOS
Eye oint 1%		5 g OP	✓ <u>Devatis</u>
Eye drops 0.5%		10 ml OP dications.	✓ <u>Chlorafast</u>
CIPROFLOXACIN	0.00	5 100	40: "
Eye drops 0.3% – Subsidy by endorsement		5 ml OP al conjunctivitis	✓ Ciprofloxacin Teva resistant to chloramphenicol; or
for the second line treatment of chronic suppurative otiti. Note: Indication marked with a * is an unapproved indic		; and the preso	cription is endorsed accordingly.
GENTAMICIN SULPHATE	alion.		
Eye drops 0.3%	11.40	5 ml OP	✓ Genoptic
PROPAMIDINE ISETHIONATE * Eye drops 0.1%	2.97	10 ml OP	
•	(14.55)		Brolene
SODIUM FUSIDATE [FUSIDIC ACID] Eye drops 1%	5 20	5 g OP	✓ Fucithalmic
Lyo diopo 1/0		3 g Oi	- i uoitiitainiit

Subsidy

Fully

Brand or

	Subsidy		Fully	Brand or	
	(Manufacturer's P	rice) Sul	osidised	Generic	
	\$	Per	1	Manufacturer	
TOBRAMYCIN					
Eye oint 0.3%	10.45	3.5 g OP	✓ T	obrex	
Eye drops 0.3%		5 ml OP	✓ T	obrex	
Corticosteroids and Other Anti-Inflammatory P	reparations				
DEXAMETHASONE					
* Eye oint 0.1%	5.86	3.5 g OP	✓ N	/laxidex	
* Eye drops 0.1%	4.50	5 ml OP	✓ N	/laxidex	
Ocular implant 700 mcg - Special Authority see SA1680 be					
- Retail pharmacy		1	✓ 0)zurdex	

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not vet completed a family: and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

*	Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b		
	sulphate 6,000 u per g5.39	3.5 g OP	✓ Maxitrol
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin		
	b sulphate 6,000 u per ml4.50	5 ml OP	✓ Maxitrol
DIC	CLOFENAC SODIUM		
	Eye drops 0.1%13.80	5 ml OP	✓ Voltaren Ophtha

SENSORY ORGANS

	Subsidy (Manufacturer's Prio \$	ce) Subs	Fully sidised	Brand or Generic Manufacturer
FLUOROMETHOLONE				
* Eye drops 0.1%	3.09	5 ml OP	√ F	ML
	5.20		√ F	lucon
LEVOCABASTINE				
Eye drops 0.5 mg per ml	8.71	4 ml OP		
• •	(10.34)		L	ivostin
LODOXAMIDE				
Eye drops 0.1%	8.71	10 ml OP	√ L	.omide
PREDNISOLONE ACETATE				
Eye drops 1%	5.93	10 ml OP	✓ P	rednisolone-AFT
) and the second	7.00	5 ml OP	✓ P	red Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority se	ee SA1715 below -	- Retail phari	macy	
Eye drops 0.5%, single dose (preservative free)		20 dose	•-	linims Prednisolone

⇒SA1715 Special Authority for Subsidy

SODIUM CROMOGLICATE

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Eye drops 2%	5 ml OP	✓ Rexacrom
Glaucoma Preparations - Beta Blockers		
BETAXOLOL * Eye drops 0.25%	5 ml OP 5 ml OP	✓ Betoptic S✓ Betoptic
* Eye drops 0.25% 1.43 * Eye drops 0.5% 1.43 * Eye drops 0.5%, gel forming 3.78	5 ml OP 5 ml OP 2.5 ml OP	✓ Arrow-Timolol ✓ Arrow-Timolol ✓ Timoptol XE
Glaucoma Preparations - Carbonic Anhydrase Inhibitors		
ACETAZOLAMIDE * Tab 250 mg17.03	100	✓ Diamox
BRINZOLAMIDE	100	· Diamox
* Eye drops 1%9.77	5 ml OP	✓ Azopt

	Subsidy (Manufacturer's I \$	Price) Subs Per	Fully Brand or idised Generic ✓ Manufacturer
Glaucoma Preparations - Prostaglandin Analog	jues		
BIMATOPROST			
* Eye drops 0.03%	3.30	3 ml OP	✓ Bimatoprost Multichem
LATANOPROST			
* Eye drops 0.005%	1.57	2.5 ml OP	✓ Teva
TRAVOPROST			
* Eye drops 0.004%	7.30	5 ml OP	✓ Travopt
	19.50	2.5 ml OP	✓ Travatan
Glaucoma Preparations - Other			
BRIMONIDINE TARTRATE			
* Eye drops 0.2%	4.29	5 ml OP	✓ Arrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE			
* Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ Combigan
PILOCARPINE HYDROCHLORIDE			· ·
* Eye drops 1%	4.26	15 ml OP	✓ Isopto Carpine
* Eye drops 2%	5.35	15 ml OP	✓ Isopto Carpine
* Eye drops 4%		15 ml OP	✓ Isopto Carpine
Subsidised for oral use pursuant to the Standard Formu	lae.		
* Eye drops 2% single dose – Special Authority see SA0895			4
below – Retail pharmacy	31.95	20 dose	Minims Pilocarpine
⇒SA0895 Special Authority for Subsidy			
Initial application from any relevant practitioner. Approvals val	id for 2 years for	applications me	eeting the following criteria:
Either:	vary to the present	nuctives or	
 Patient has to use an unpreserved solution due to an alle Patient wears soft contact lenses. 	rgy to the preser	valive, or	
Note: Minims for a general practice are considered to be "tools"	of trade" and are	not approved a	s special authority items
Renewal from any relevant practitioner. Approvals valid for 2 years			
benefiting from treatment.			ppp
Mydriatics and Cycloplegics			
ATROPINE SULPHATE			
* Eye drops 1%	17.36	15 ml OP	✓ Atropt
Atropt to be Sole Supply on 1 October 2020		10 1111 01	- 7110pt

15 ml OP

15 ml OP

15 ml OP

✓ Cyclogyl

✓ Mydriacyl✓ Mydriacyl

Preparations for Tear Deficiency

CYCLOPENTOLATE HYDROCHLORIDE

For acetylcysteine eye drops refer Standard Formulae, page 247

* Eye drops 0.5%......7.15

HYPROMELLOSE

TROPICAMIDE

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

SENSORY ORGANS

	Subsidy (Manufacturer's P	rice) Subs	Fully idised	Brand or Generic
	\$	Per	1	Manufacturer
HYPROMELLOSE WITH DEXTRAN * Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	✓ P	oly-Tears

Preservative Free Ocular Lubricants

⇒SA1388 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER - Special Authority see SA1388 above - Retail	pharmacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL - Special Au	thority see SA1388 above	– Retail	pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	 Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special A	Authority see SA1388 abo	ve – Reta	ail pharmacy
Eye drops 1 mg per ml	22.00 10	ml OP	✓ Hylo-Fresh
Hylo-Fresh has a 6 month expiry after opening. The	Pharmacy Procedures M	lanual res	triction allowing one bottle per
month is not relevant and therefore only the prescribe	ed dosage to the nearest	OP may b	pe claimed.

Other Eye Preparations

NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%4.15	15 ml OP	✓ Naphcon Forte
OLOPATADINE		
Eye drops 0.1%2.20	5 ml OP	✓ Olopatadine Teva✓ Patanol
Olopatadine Teva to be Sole Supply on 1 October 2020 (Patanol Eye drops 0.1% to be delisted 1 October 2020)		
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE Eve oint 138 mcg per g	5 a OP	✓ VitA-POS

Subsidy (Manufacturer's Price) S \$ Per

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

Agents Used in the Treatment of Poisonings

Antidotes

ACETYL CYSTEINE

NALOXONE HYDROCHLORIDE

- a) Up to 5 inj available on a PSO
- b) Only on a PSO
- * Inj 400 mcg per ml, 1 ml ampoule22.60

5 ✓ <u>DBL Naloxone</u>

Hydrochloride

Removal and Elimination

CHARCOAL

*	Oral liq 50 g per 250 ml	43.50	250 ml OP	✓ Carbosorb-X
---	--------------------------	-------	-----------	---------------

- a) Up to 250 ml available on a PSO
- b) Only on a PSO

DEFERASIROX – Special Authority see SA1492 below – Retail pharmacy

Tractage claimable			
Tab 125 mg dispersible	276.00	28	Exjade
Tab 250 mg dispersible	552.00	28	✓ Exjade
Tab 500 mg dispersible	1,105.00	28	✓ Exjade

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE	- Special Authority see SA	1480 on the next	page – Retail pharmacy

Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox



5	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	•	Manufacturer

⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

* Inj 500 mg vial	84.53	10	✓ <u>DBL</u> <u>Desferrioxamine</u> <u>Mesylate for Inj</u> <u>BP</u>
SODIUM CALCIUM EDETATE	50.04		

DESERBIOXAMINE MESILATE

SO	DIUM CALCIUM EDETATE			
*	Inj 200 mg per ml, 5 ml	53.31	6	
		(156.71)		Calcium Disodium
				Versenate

Standard Formulae ACETYLCYSTEINE EYE DROPS		PHENOBARBITONE ORAL LIQUID	
Acetylcysteine inj 200 mg per ml, 10 ml	qs	Phenobarbitone Sodium	1 g
Suitable eye drop base	qs	Glycerol BP Water	70 ml to 100 ml
ASPIRIN AND CHLOROFORM APPLICATION Aspirin Soluble tabs 300 mg Chloroform	12 tabs to 100 ml	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml)	LIQUID (10
CODEINE LINCTUS (3 mg per 5 ml)		Phenobarbitone Sodium	400 mg 4 ml
Codeine phosphate	60 mg	Glycerol BP Water	to 40 ml
Glycerol Preservative	40 ml	DIL OCADDINE ODAL LIQUID	
Water	qs to 100 ml	PILOCARPINE ORAL LIQUID Pilocarpine 4% eye drops	qs
	10 100 1111	Preservative	qs
CODEINE LINCTUS (15 mg per 5 ml)	000	Water	to 500 ml
Codeine phosphate Glycerol	300 mg 40 ml	(Preservative should be used if quantity supplied is	for more
Preservative	qs	than 5 days.)	
Water	to 100 ml	SALIVA SUBSTITUTE FORMULA	
FOLINIC MOUTHWASH		Methylcellulose Preservative	5 g
Calcium folinate 15 mg tab	1 tab	Water	qs to 500 ml
Preservative	qs	(Preservative should be used if quantity supplied is	
Water	to 500 ml	than 5 days. Maximum 500 ml per prescription.)	
(Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	for more	SODIUM CHLORIDE ORAL LIQUID	
, , , , , , , , , , , , , , , , , , , ,		Sodium chloride inj 23.4%, 20 ml	qs
MAGNESIUM HYDROXIDE 8% MIXTURE	075 -	Water	qs
Magnesium hydroxide paste 29% Methyl hydroxybenzoate	275 g 1.5 g	(Only funded if prescribed for treatment of hyponatr	aemia)
Water	to 1,000 m	NANCOMYCIN ORAL SOLUTION (50 mg per ml) Vancomycin 500 mg injection	10 vials
METHADONE MIXTURE		Glycerol BP	40 ml
Methadone powder	qs	Water	to 100 ml
Glycerol	qs	(Only funded if prescribed for treatment of Clostridia	um difficile
Water	to 100 ml	following metronidazole failure)	
METHYL HYDROXYBENZOATE 10% SOLUTION	40	VOSOL EAR DROPS	
Methyl hydroxybenzoate Propylene glycol	10 g to 100 ml	WITH HYDROCORTISONE POWDER 1% Hydrocortisone powder	1%
(Use 1 ml of the 10% solution per 100 ml of oral liqu		Vosol Ear Drops	to 35 ml

OMEPRAZOLE SUSPENSION Omeprazole capules or powder qs Sodium bicarbonate powder BP 8.4 g Water to 100 ml

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Douglas

Extemporaneously Compounded Preparations and Galenicals

OROFORM	

- a) Only in combination
- b) Maximum of 100 ml per prescription
- c) Only in aspirin and chloroform application.
- d) Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be

(90.09)

✓ PSM Chloroform BP......25.50 500 ml (PSM Chloroform BP to be delisted 1 November 2020) CODEINE PHOSPHATE - Safety medicine; prescriber may determine dispensing frequency Powder - Only in combination......63.09

Only in extemporaneously compounded codeine linctus.

COLLODION FLEXIBLE

Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined.

Collodion flexible19.30 100 ml ✓ PSM

COMPOUND HYDROXYBENZOATE - Only in combination

Only in extemporaneously compounded oral mixtures.

100 ml Midwest

GLYCERIN WITH SODIUM SACCHARIN - Only in combination

Only in combination with Ora-Plus.

473 ml ✓ Ora-Sweet SF

GLYCERIN WITH SUCROSE - Only in combination Only in combination with Ora-Plus.

473 ml

✓ Ora-Sweet

GLYCEROL

✓ healthE Glycerol BP 500 ml

a) Only in extemporaneously compounded oral liquid preparations.

b) healthE Glycerol BP to be Sole Supply on 1 October 2020

MAGNESIUM HYDROXIDE

✓ PSM 500 a

(PSM Paste 29% to be delisted 1 January 2021)

METHADONE HYDROCHLORIDE

- a) Only on a controlled drug form
- b) No patient co-payment payable
- c) Safety medicine; prescriber may determine dispensing frequency
- d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets).

✓ AFT 1 q METHYL HYDROXYBENZOATF ✓ Midwest 25 a METHYLCELLULOSE ✓ MidWest 100 g 473 ml ✓ Ora-Plus METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN - Only in combination 473 ml ✓ Ora-Blend SF

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Pri \$	ce) Subs	Fully sidised	Brand or Generic Manufacturer
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - On Suspension	•	473 ml	√ <u>C</u>	Ora-Blend
PHENOBARBITONE SODIUM Powder – Only in combination	52.50 325.00	10 g		/lidWest
Only in children up to 12 years PROPYLENE GLYCOL	325.00	100 g	V IV	niawest
Only in extemporaneously compounded methyl hydroxybenz		ı. 500 ml	✓ N	/lidwest
SODIUM BICARBONATE Powder BP - Only in combination	10.05	500 g	✓ <u>N</u>	<u>//lidwest</u>
Only in extemporaneously compounded omeprazole and SYRUP (PHARMACEUTICAL GRADE) – Only in combination		spension.		
Only in extemporaneously compounded oral liquid preparation		500 ml	✓ <u>N</u>	<u>/lidwest</u>
WATER Tap - Only in combination	0.00	1 ml	√ T	ap water

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

400 a OP ✓ Polycal

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

✓ fully subsidised 251

Subsidy		Fully	Brand or
(Manufacturer's Price)	_ 8	Subsidised	Generic
\$	Per		Manufacturer

continued...

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT – Special Authority see SA1523 on the previous page – Hospital pharmacy [HP3]

Emulsion (neutral)	12.30 200	ml OP	Calogen
	30.75 500	ml OP 🗸	Calogen
Emulsion (strawberry)	12.30 200	ml OP 🗸	Calogen
Oil	30.00 500	ml OP 🗸	MCT oil (Nutricia)
Oil, 250 ml1	14.92 4	OP 🗸	Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT – Special Authority see SA1524 above – Hospital pharmacy [HP3]
Powder7.90 225 g OP
8.95 227 g OP
✓ Resource
Beneprotein

Subsidy (Manufacturer's Price)

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

Oral Supplements/Complete Diet (Nasogastric/Gastrostomy Tube Feed)

Respiratory Products

⇒SA1094 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has CORD and hypercapnia, defined as a CO2 value exceeding 55 mmHg.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see SA109 Liquid		Hospital pharm 1,000 ml OP	acy [HP3] ✓ Diason RTH ✓ Glucerna Select RTH
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 ab	ove – Hos	pital pharmacy	[HP3]
Liquid (strawberry)	1.50	200 ml OP	✓ Diasip
Liquid (vanilla)	1.50	200 ml OP	✓ Diasip
	1.88	250 ml OP	✓ Glucerna Select
	1.78	237 ml OP	
	(2.10)		Resource Diabetic
	(2.10)		Sustagen Diabetic



Subsidy (Manufacturer's Price)

Fully Subsidised Brand or Generic Manufacturer

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism: or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults,

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED - Special Authority see SA1525 above - Hospital pharmacy [HP3] 400 g OP Monogen

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

400 a OP ✓ Heparon Junior

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	1	Manufacturer

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1379 ab Liquid6.00	oove – Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini Energy RTH
PAEDIATRIC ENTERAL FEED 1KCAL/ML – Special Authority see \$A1379 abov Liquid2.68	ve – Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini RTH ✓ Pediasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority st Liquid6.00	ee SA1379 above – Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA1379 above	- Hospital pharmacy [HP3]
Liquid (strawberry)1.60	200 ml OP ✓ Fortini
Liquid (vanilla)1.60	200 ml OP ✓ Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA1379 above -	Hospital pharmacy [HP3]
Liquid (chocolate)1.07	200 ml OP ✓ Pediasure
Liquid (strawberry)1.07	200 ml OP ✓ Pediasure
Liquid (vanilla)1.07	200 ml OP ✓ Pediasure
1.34	250 ml OP ✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority see S	A1379 above – Hospital pharmacy [HP3]
Liquid (unflavoured)	200 ml OP ✓ Fortini Multi Fibre
Liquid (chocolate)	200 ml OP ✓ Fortini Multi Fibre
Liquid (strawberry)1.60	200 ml OP ✓ Fortini Multi Fibre
Liquid (vanilla)	200 ml OP ✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED - Special Authority see SA1379 above - Hospita	al nharmacy [HP3]
Powder	400 g OP ✓ Peptamen Junior
70.00	100 g Ci - 1 optamen damen

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
· · · · ·	Dor		Manufacturor

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML - Special Authori	•	Hospital pharm 500 ml OP	,
RENAL ORAL FEED 1.8 KCAL/ML - Special Authority se	e SA1101 above – Hos	pital pharmacy	[HP3]
Liquid		220 ml OP	✓ Nepro HP (strawberry) ✓ Nepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML - Special Authority see	SA1101 above – Hospi	tal pharmacy [F	HP3]
Liquid	2.88	237 ml OP	
	(3.31)		NovaSource Renal
Liquid (apricot) 125 ml	11.52	4 OP	✓ Renilon 7.5
Liquid (caramel) 125 ml	11.52	4 OP	✓ Renilon 7.5

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's P \$	rice) Subs Per	Fully idised	Brand or Generic Manufacturer
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see Liquid (grapefruit), 250 ml carton Liquid (pineapple & orange), 250 ml carton Liquid (summer fruits), 250 ml carton	171.00 171.00	orevious page - 18 OP 18 OP 18 OP	✓ E	ital pharmacy [HP3] Elemental 028 Extra Elemental 028 Extra Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML – Special Authority see S Powder (unflavoured)	4.50	80 g OP	√ ∨	ivonex TÉN
[HP3] Liquid	•	1,000 ml OP		Peptisorb

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and

continued...

Subsidy (Manufacturer's Pr	rice)	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age, and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g., to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant: and

continued...

(Man	Subsidy ufacturer's Price)	Subsid	Fully lised	Brand or Generic
	\$	Per	1	Manufacturer

continued...

- 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum: or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease: or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia: or
- 7 Short bowel syndrome; or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm3); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

continued...

	Subsidy (Manufacturer's F		Subsid er	Fully lised	Brand or Generic Manufacturer
continued					
 Is being fed via a tube or a tube is to be inserted for the pucondition criteria); or Cystic Fibrosis; or Liver disease; or Chronic Renal failure; or Inflammatory bowel disease; or Chronic obstructive pulmonary disease with hypercapnia; Short bowel syndrome; or Bowel fistula; or Severe chronic neurological conditions. 	or				
ENTERAL FEED 1.5KCAL/ML – Special Authority see SA1859 c Liquid		lospital p 1,000		•] utrison Energy
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on Liquid		spital ph 250 n 1,000	nl OP ´	✓ İs	osource Standard utrison Standard RTH smolite RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authorit	v coo SA1850 o	n naga	257 _ Ho		
LiquidLiquid	•	1,000			utrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority so Liquid		1,000		√ Je	armacy [HP3] evity RTH utrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority		page 25 250 n 1,000	nl OP	✓ E	narmacy [HP3] nsure Plus HN nsure Plus RTH evity HiCal RTH utrison Energy Multi Fibre
ORAL FEED (POWDER) — Special Authority see SA1859 on page Note: Higher subsidy for Sustagen Hospital Formula will only number and an appropriately endorsed prescription. Powder (chocolate) — Higher subsidy of up to \$26.00 per 850	y be reimbursed				
with Endorsement		850 g 840 g	•		nsure ustagen Hospital
	(20.00)			3	Formula Active
Additional subsidy by endorsement is available for patier prescription must be endorsed accordingly. Paydor (varilla) Higher subsidy of up to \$00 per 950 g.	nts with fat mala	bsorptio	n, fat into	olerand	ce or chyle leak. The
Powder (vanilla) – Higher subsidy of up to \$26.00 per 850 g with Endorsement	8.54	857	1 OP	✓ F	ortisip
	26.00 9.54	850 (840 (OP	✓ E	nsure

Additional subsidy by endorsement is available for patients with fat malabsorption, fat intolerance or chyle leak. The prescription must be endorsed accordingly.

(26.00)

Sustagen Hospital Formula Active

	Subsidy	Fully	Brand or
(1)	Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 257 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
Liquid (fruit of the forest) - Higher subsidy of \$1.26 per 200 ml			
with Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.33 per 237 ml with			
Endorsement	0.85	237 ml OP	
	(1.33)		Ensure Plus
	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML – Special Authority see SA1859 on page 257 – Hospital pharmacy [HP3]
Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolata) – Higher subsidy of \$1.26 per 200 ml with

Endorsement	0.72	200 ml OP	
	(1.26)	200 1111 01	Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

continued...

Su	ubsidy F	ully	Brand or
(Manufac	cturer's Price) Subsid	ised	Generic
	\$ Per	•	Manufacturer

continued...

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$1.90 per 200 ml with

(1.90) Two Cal HN

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully osidised	Brand or Generic Manufacturer	
FOOD THICKENER - Special Authority see SA1106 on the p			,		
Powder	6.53 3	00 g OP	✓ N	lutilis	
	7.25 3	80 g OP		eed Thickener Karicare Aptamil	

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by PHARMAC from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

e – Hospital	pharmacy [HP3]	
2.81	1,000 g OP	
(5.15)	, 3	Healtheries Simple Baking Mix
– Hospital p	oharmacy [HP3]	
3.93	1,000 g OP	
(7.32)		NZB Low Gluten Bread Mix
3.51		
(10.87)		Horleys Bread Mix
ospital pharn	nacy [HP3]	
5.62	2,000 g OP	
(18.10)	, 3 -	Horleys Flour
	2.81 (5.15) 9 – Hospital µ 3.93 (7.32) 3.51 (10.87) lospital pharr5.62	(5.15) B – Hospital pharmacy [HP3]3.93 1,000 g OP (7.32) 3.51 (10.87) lospital pharmacy [HP3]5.62 2,000 g OP

	Subsidy (Manufacturer's Pri	ice) Su Per	Fully bsidised	Brand or Generic Manufacturer
	\$			
GLUTEN FREE PASTA – Special Authority see SA1729 on the		lospital pha	rmacy [H	P3]
Buckwheat Spirals	2.00	250 g OP		
	(3.11)		C	Orgran
Corn and Vegetable Shells	2.00	250 g OP		
	(2.92)		C	Orgran
Corn and Vegetable Spirals	2.00	250 g OP		
	(2.92)	-	C	Orgran
Rice and Corn Lasagne Sheets	1.60	200 g OP		•
·	(3.82)	ŭ	C	Orgran
Rice and Corn Macaroni	` ,	250 g OP		•
	(2.92)	3 -	C	Orgran
Rice and Corn Penne	` ,	250 g OP		
	(2.92)	5	(Orgran
Rice and Maize Pasta Spirals		250 g OP		g.u
. 100 a.i.a . 11a.20 . a.u.a opi alo	(2.92)	_00 g 0.	(Orgran
Rice and Millet Spirals	, ,	250 g OP	·	rigian
Those and Williot Ophialo	(3.11)	200 g O1		Orgran
Rice and corn spaghetti noodles	` ,	375 g OP		rigian
Thee and com spagnetti hoodies	(2.92)	073 g Oi	_	Orgran
Vegetable and Rice Spirals	` ,	250 g OP		rigian
Vogetable and ince opilals	(2.92)	200 g OF		Orgran
Italian long style speaketti	` ,	220 a OB	_	rigian
Italian long style spaghetti		220 g OP)raran
	(3.11)		C	Orgran

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 above - Hospital pharmacy [HP3]

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	osidised	Generic
\$	Per	/	Manufacturer

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

Tabs	99.00	75 OP	✓ Phlexy 10
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior Chocolate
Powder (unflavoured) 27.8 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior Vanilla
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange)	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured)	320.00	500 g OP	XP Maxamum
Liquid (berry)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	 Easiphen Liquid
Liquid (juicy tropical) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g	1,123.20	36 OP	✓ PKU Lophlex Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20

Foods

the previous p	age – Hospital p	harmacy [HP3]
8.22	500 g OP	Loprofin Mix
revious page -	Hospital pharm	acy [HP3]
11.91	500 g OP	Loprofin
5.95	250 g OP	Loprofin
11.91	500 g OP	Loprofin
5.95	250 g OP	Loprofin
11.91	500 g OP	Loprofin
11.91	500 g OP	Loprofin
11.91	500 g OP	Loprofin
		revious page – Hospital pharm



Subsidy (Manufacturer's Price) Fully Subsidised

Per

Brand or Generic Manufacturer

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]
Powder44.40 400 g OP ✓ Locasol

Gastrointestinal and Other Malabsorptive Problems

AMINO ACID FORMULA – Special Authority see SA1940 below Powder	,. ,.	✓ Alfamino Junior
Powder (unflavoured)	53.00 400 g OP	✓ Elecare
,	ū	✓ Elecare LCP
		✓ Neocate Gold
		 Neocate Junior Unflavoured
		✓ Neocate SYNEO
Powder (vanilla)	53.00 400 g OP	✓ Elecare
. ,	· ·	 Neocate Junior Vanilla

⇒SA1940 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short aut: or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Fither:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist.

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Approvals valid for 6 months for applications meeting the following criteria:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency: or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Fither:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
 - 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
 - 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Fither:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

continued...

SPECIAL FOODS

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products: or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see SA1557 below - Hospital pharmacy [HP3]

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome: or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia: or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued

recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula: and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)	35.50	300 g OP	✓ KetoCal 4:1 ✓ Ketocal 3:1
Powder (vanilla)	35.50	300 g OP	✓ KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Vaccinations

ADULT DIPHTHERIA AND TETANUS VACCINE - [Xpharm]

- 1) For vaccination of patients aged 45 and 65 years old; or
- 2) For vaccination of previously unimmunised or partially immunised patients; or
- 3) For revaccination following immunosuppression; or
- 4) For boosting of patients with tetanus-prone wounds; or
- 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

(ADT Booster Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid in 0.5 ml to be delisted 1 October 2020)

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent......0.00 10

10 **✓ BCG Vaccine**

BCG Vaccine to be Sole Supply on 1 October 2020

DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE - [Xpharm]

Funded for any of the following criteria:

- 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
- 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
- 3) A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
- 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens: or
- 5) A single dose for vaccination of patients aged 65 years old; or
- 6) A single dose for vaccination of patients aged 45 years old who have not had 4 previous tetanus doses; or
- 7) For vaccination of previously unimmunised or partially immunised patients; or
- 8) For revaccination following immunosuppression; or
- 9) For boosting of patients with tetanus-prone wounds.

Notes: Tdap is not registered for patients aged less than 10 years. Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid, 8 mcg pertussis filamentous

Boostrix to be Sole Supply on 1 October 2020

Subsidised

Subsidy

(Manufacturer's Price)

Fully

Brand or

Generic

	`	\$	Per	•	Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE -	- [Xphari	m]			
Funded for any of the following:					
 A single dose for children up to the age of 7 who have of 					
 A course of four vaccines is funded for catch up prograte primary immunisation; or 	mmes fo	r children (to	the age of 1	10 year	rs) to complete full
 An additional four doses (as appropriate) are funded fo pre- or post splenectomy; pre- or post solid organ trans 					
regimens; or		•		,	
4) Five doses will be funded for children requiring solid org	•	•			
Note: Please refer to the Immunisation Handbook for approp	oriate sci	nedule for cat	ch up progr	amme	S.
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg					
pertussis toxoid, 25 mcg pertussis filamentous					
haemagglutinin, 8 mcg pertactin and 80 D-antigen units			40		
poliomyelitis virus in 0.5ml syringe		0.00	10	✓ Int	fanrix IPV
Infanrix IPV to be Sole Supply on 1 October 2020					
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B A [Xpharm]	ND HAE	MOPHILUS I	NFLUENZA	E TYF	'E B VACCINE -
Funded for patients meeting any of the following criteria:					
1) Up to four doses for children up to and under the age o	f 10 for r	orimary immur	nisation: or		
2) An additional four doses (as appropriate) are funded for	r (re-)imr	munisation for	children ur	o to an	d under the age of
10 who are patients post haematopoietic stem cell trans					
post solid organ transplant, renal dialysis and other sev					
3) Up to five doses for children up to and under the age of					
Note: A course of up-to four vaccines is funded for catch up					
to complete full primary immunisation. Please refer to the Im					
programmes.					
Inj 30IU diphtheriatoxoid with 40IU tetanustoxoid, 25mcg					
pertussistoxoid, 25mcg					
pertussisfilamentoushaemagglutinin, 8 mcgpertactin,					
80 D-AgUpoliovirus, 10mcghepatitisBsurfaceantigen in					
0.5ml syringe		0.00	10	✓ Inf	fanrix-hexa
Infanrix-hexa to be Sole Supply on 1 October 2020					
HAEMOPHILUS INFLUENZAE TYPE B VACCINE - [Xpharm]					
One dose for patients meeting any of the following:					
, , ,					
For primary vaccination in children; or An additional data (see appropriate) in funded for (re.) in		ion for notion	a noot boo	matan.	niatio atam aall
 An additional dose (as appropriate) is funded for (re-)in transplantation, or chemotherapy; functional asplenic; p 					
or post cochlear implants, renal dialysis and other seve					iu organ transpiant, pre-
For use in testing for primary immunodeficiency disease					al modicino physician or
paediatrician.	55, OH III	e recommend	alion of an	IIII	ai medicine priysician oi
paeulatiiciaii.					
Harmond Startes for the Committee of the Administration of the Adm					
Haemophilus Influenzae type B polysaccharide 10 mcg					
conjugated to tetanus toxoid as carrier protein 20-40 mg		0.00		<i>-</i>	L. a. da.
prefilled syringe plus vial 0.5 ml		0.00	1	✓ Hil	Derix

	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer
HEPATITIS A VACCINE – [Xpharm] Funded for patients meeting any of the following criteria: 1) Two vaccinations for use in transplant patients; or 2) Two vaccinations for use in children with chronic liver 3) One dose of vaccine for close contacts of known hepatics.	,		

Inj 1440 ELISA units in 1 ml syringe	1	✓ Havrix
Havrix to be Sole Supply on 1 October 2020		
Inj 720 ELISA units in 0.5 ml syringe	1	Havrix Junior
Havrix Junior to be Sole Supply on 1 October 2020		

		Subsidy		Fu		and or
		(Manufacturer's Price)	Per	Subsidis		eneric anufacturer
IEDATITIC D	DECOMPINANT VACCINE [Valored]	Ψ	1 01		- 1410	and dotal of
	RECOMBINANT VACCINE – [Xpharm] per 0.5 ml vial	0.00	1		✓ HBva	vDDO
, ,	led for patients meeting any of the following criteria:	0.00	1		приа	XPNU
	for household or sexual contacts of known acute he	notitio P notionto or h	onot	itio D oo	rriara: ar	
,	for children born to mothers who are hepatitis B sui				mers, or	
	for children up to and under the age of 18 years inc				avo ach	ioved a nocitive
0)	serology and require additional vaccination or requi					icved a positive
4)	for HIV positive patients; or	re a primary occinc o	ı vuc	omation	, 01	
,	for hepatitis C positive patients; or					
,	for patients following non-consensual sexual interco	ourse; or				
,	for patients following immunosuppression; or	•				
	for solid organ transplant patients; or					
9)	for post-haematopoietic stem cell transplant (HSCT) patients; or				
	following needle stick injury.	,,				
					_	
	g per 1 ml vial	0.00	1	•	HBva	xPRO
	led for patients meeting any of the following criteria:					
,	for household or sexual contacts of known acute he		•		rriers; or	,
	for children born to mothers who are hepatitis B sur					
3)	for children up to and under the age of 18 years inc					leved a positive
4\	serology and require additional vaccination or requi	re a primary course o	t vac	cination	; or	
,	for HIV positive patients; or					
	for hepatitis C positive patients; or for patients following non-consensual sexual interco	NIFOO: OF				
,	for patients following immunosuppression; or	ourse, or				
,	for solid organ transplant patients; or					
	for post-haematopoietic stem cell transplant (HSCT) natients: or				
,	following needle stick injury.	, panomo, or				
-,	3					
Inj 20 mc	g per 1 ml prefilled syringe	0.00	1	,	Enge	rix-B
a) F	unded for patients meeting any of the following crite	ria:				
	1) for household or sexual contacts of known acute					s; or
	2) for children born to mothers who are hepatitis B	surface antigen (HBs	Ag)	positive;	or	
	3) for children up to and under the age of 18 years					achieved a positive
	serology and require additional vaccination or re	equire a primary cours	se of	vaccina	tion; or	
	4) for HIV positive patients; or					
	5) for hepatitis C positive patients; or					
	6) for patients following non-consensual sexual int	ercourse; or				
	7) for patients following immunosuppression; or					
	8) for solid organ transplant patients; or	CT) notionto, or				
	 9) for post-haematopoietic stem cell transplant (HSIO) following needle stick injury; or 	oci) patients; or				
	11) for dialysis patients; or					
	11) for dialysis patients, or 12) for liver or kidney transplant patients.					
	ngerix-B to be Sole Supply on 1 October 2020					
,	g per 1 ml vialg	0.00	1	,	✓ HBva	xPRO
,	J F · ···· · ·		•			· · ·

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
Funded for any of the following criteria:

- 1) for dialysis patients; or
- 2) for liver or kidney transplant patient.

(HBvaxPRO Inj 5 mcg per 0.5 ml vial to be delisted 1 October 2020) (HBvaxPRO Inj 10 mcg per 1 ml vial to be delisted 1 October 2020) (HBvaxPRO Inj 40 mcg per 1 ml vial to be delisted 1 October 2020)

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV] - [Xpharm]

Any of the following:

- 1) Maximum of two doses for children aged 14 years and under; or
- 2) Maximum of three doses for patients meeting any of the following criteria:
 - 1) People aged 15 to 26 years inclusive; or
 - 2) Either:

People aged 9 to 26 years inclusive

- 1) Confirmed HIV infection; or
- 2) Transplant (including stem cell) patients: or
- 3) Maximum of four doses for people aged 9 to 26 years inclusive post chemotherapy

	Subsidy (Manufacturer's Price)	Subs Per	Fully idised	Brand or Generic Manufacturer
INFLUENZA VACCINE	<u> </u>			

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine)

- [Xpharm]......9.00 1 ✓ Afluria Quad Junior (2020 Formulation)

A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by PHARMAC:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes: or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders. or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

Influvac Tetra	1	ccine)9.00	Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)
(2020 formulation)			
✓ Afluria Quad	10	90.00	
(2020 Formulation)			

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

- a) Only on a prescription
- b) No patient co-payment payable

С

A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by PHARMAC:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease: or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or
- c) children aged four years or less (but over three years) who have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Subsidy (Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer	
*				

MEASI ES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment from the Funder for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

✓ MMR II	5	diluent 0.5 ml
✓ Priorix	10	250.00

Priorix to be Sole Supply on 1 October 2020

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Either:

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
 - 2) One dose for close contacts of meningococcal cases; or
 - 3) A maximum of two doses for bone marrow transplant patients; or
 - 4) A maximum of two doses for patients following immunosuppression*; or
- B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) Either:
 - One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
 - One dose for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 December 2019 to 30 November 2020.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 4 mcg of each meningococcal polysaccharide conjugated to

a total of approximately 48 mcg of diphtheria toxoid carrier			
per 0.5 ml vial	0.00	1	✓ Menactra
Menactra to be Sole Supply on 1 October 2020			

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Subsidised Manufacturer

MENINGOCOCCAL C CONJUGATE VACCINE − [Xpharm]
Both:

- 1) The child is under 9 months of age; and
- 2) Any of the following:
 - Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
 - 2) Two doses for close contacts of meningococcal cases; or
 - 3) A maximum of two doses for bone marrow transplant patients; or
 - 4) A maximum of two doses for patients pre- and post-immunosuppression*.

Note: children under nine months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for booster schedules with meningococcal ACWY vaccine.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 10 mcg in 0.5 ml syringe	0.00	1	✓ Neisvac-C
PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm]			
1) A primary course of three doses for previously unvaccina	ated individuals (up to the age	e of 59 months inclusive
Note: please refer to the Immunisation Handbook for the app	ropriate schedule	e for catch u	p programmes
Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B	,		
7F, 9V, 14 and 23F; 3 mcg of pneumococcal			
polysaccharide serotypes 4, 18C and 19F in 0.5 ml			
prefilled syringe	0.00	10	Synflorix
Synflorix to be Sole Supply on 1 October 2020			

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- 1) Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) with primary immune deficiencies: or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cyanosis or failure; or
 - I) with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note:	please refer to the Im	munisation Han	dbook for the	appropriate	schedule for	catch up	orogrammes
Inj 30.	8 mcg of pneumococc	al polysacchario	de serotypes 1	1, 3, 4,			

NATIONAL IMMUNISATION SCHEDULE					
	Subsidy (Manufacturer's Price) \$	Per	F Subsidi	ully sed	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE - Either:	- [Xpharm]				
Up to three doses (as appropriate) for patients with I chemotherapy; pre- or post-splenectomy or with func complement deficiency (acquired or inherited), cochle All of the following: a) Patient is a child under 18 years for (re-)immun	tional asplenia, pre- or pear implants, or primary	oost-	solid or	gan tr	ansplant, renal dialysis,
b) Treatment is for a maximum of two doses; andc) Any of the following:					
 i) on immunosuppressive therapy or radiation immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; v) who are immune-suppressed following on 	or				
or vi) with cochlear implants or intracranial shur vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more to prednisone of 2 mg/kg per day or greater, 20 mg or greater; or ix) with chronic pulmonary disease (including x) pre term infants, born before 28 weeks ge xi) with cardiac disease, with cyanosis or fail xii) with diabetes; or	than two weeks, and who or children who weigh or children who weigh or asthma treated with high estation; or	more	than 1	0 kg c	on a total daily dosage of
xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with	functional asplenia.				
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	ng:	1		√ P	neumovax 23
Note: Please refer to the Immunisation Handbook for app Inj 80D antigen units in 0.5 ml syringeIPOL to be Sole Supply on 1 October 2020		tch-u 1	p progr	amme ✓ IF	
ROTAVIRUS ORAL VACCINE – [Xpharm] Maximum of two doses for patients meeting the following: 1) first dose to be administered in infants aged under 14 2) no vaccination being administered to children aged 2					
Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator Rotarix to be Sole Supply on 1 October 2020	0.00	10		✔ R	otarix

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] – [Xpharm] Either:				
Maximum of one dose for primary vaccination for either	r:			
a) Any infant born on or after 1 April 2016; or	•			
b) For previously unvaccinated children turning 11 y varicella infection (chickenpox), or	ears old on or after 1	July 2	017, who h	ave not previously had a
2) Maximum of two doses for any of the following:				
 a) Any of the following for non-immune patients: 				
 i) with chronic liver disease who may in future ii) with deteriorating renal function before trans iii) prior to solid organ transplant; or iv) prior to any elective immunosuppression*, ov v) for post exposure prophylaxis who are immunished b) For patients at least 2 years after bone marrow tr. c) For patients at least 6 months after completion of d) For HIV positive non immune to varicella with mile e) For patients with inborn errors of metabolism at rivaricella, or f) For household contacts of paediatric patients who immune compromise where the household contacts g) For household contacts of adult patients who have 	splantation; or or une competent inpatie ansplantation, on adv chemotherapy, on ac d or moderate immun- sk of major metabolic o are immunocompror ct has no clinical histo	ents.; of the divice of the osuppose decormised, ory of v	or their specia f their spec ression on mpensation or undergo varicella, or	ialist, or advice of HIV specialist, or , with no clinical history of sing a procedure leading to
immunocompromised, or undergoing a procedure has no clinical history of varicella.				
* immunosuppression due to steroid or other immunosuppre 28 days	ssive therapy must be	e for a	treatment p	period of greater than
Inj 1350 PFU prefilled syringe	0.00	1 10		arivax arivax
Varivax to be Sole Supply on 1 October 2020 Inj 2000 PFU prefilled syringe plus vial	0.00	1 10		arilrix arilrix
(Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 (Varil				
VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATE Funded for patients meeting either of the following criteria: 1) One dose for all people aged 65 years; or	•		•	., .
2) One dose for all people aged between 66 and 80 years	inclusive from 1 Apri	l 2018	and 31 De	cember 2020.
Inj 19,400 PFU prefilled syringe plus vial	0.00	1 10		ostavax ostavax
Diagnostic Agents				
TUBERCULIN PPD [MANTOUX] TEST – [Xpharm] Inj 5 TU per 0.1 ml, 1 ml vial Tubersol to be Sole Supply on 1 October 2020	0.00	1	✓ T	ubersol

- Symbols -		Aflibercept	196	Amisulpride Mylan	13
UK Synacthen	79	Afluria Quad		Amitriptyline	12
3TC		(2020 Formulation)	275	Amlodipine	
- A -		Afluria Quad Junior		Amneal	
A-Scabies	68	(2020 Formulation)	275	Amorolfine	6
Abacavir sulphate		AFT Carbimazole		Amoxicillin	
Abacavir sulphate with		AFT-Pyrazinamide		Amoxicillin with clavulanic acid	
lamivudine	105	Agents Affecting the		Amphotericin B	3
Abiraterone acetate		Renin-Angiotensin System	47	Amsacrine	
Acarbose	11	Agents for Parkinsonism and Rel		AmsaLyo	
Accarb	11	Disorders	119	Amsidine	
Accuretic	48	Agents Used in the Treatment of		Amzoate	
Accuretic 10	48	Poisonings	245	Anaesthetics	12
Accuretic 20	48	Agrylin		Anafranil	12
Acetazolamide	242	Agrylin S29		Anagrelide hydrochloride	16
Acetec	47	Albendazole		Analgesics	12
Acetic acid with 1, 2- propane	diol	Albey		Anastrozole	17
diacetate and		Albustix	76	Andriol Testocaps	
benzethonium	240	Aldurazyme	30	Androderm	7
Acetic acid with hydroxyquino	line and	Alecensa	169	Anoro Ellipta	
ricinoleic acid	74	Alectinib	169	Antabuse	
Acetylcysteine		Alendronate sodium	111	Antacids and Antiflatulents	
Aci-Jel	74	Alendronate sodium with		Anthelmintics	8
Aciclovir		colecalciferol	111	Antiacne Preparations	6
Infection	101	Alfacalcidol	34	Antiallergy Preparations	23
Sensory	240	Alfamino Junior	266	Antianaemics	3
Acidex	6	Alginic acid	6	Antiandrogen Oral	
Acipimox	54	Alglucosidase alfa	28	Contraceptives	
Acitretin	68	Alkeran	158	Antiarrhythmics	4
Aclasta	114	Alkeran s29	158	Antibacterials	
Aclin	110	Allerpro 1	268	Antibacterials Topical	
Actemra		Allerpro 2		Anticholinergic Agents	
Actinomycin D	163	Allersoothe	233	Anticholinesterases	
Actrapid		Allmercap		Antidepressants	
Actrapid Penfill		Allopurinol		Antidiarrhoeals	
Acupan		Alpha-Adrenoceptor Blockers		Antiepilepsy Drugs	
Adalat 10		Alpha-Keri Lotion		Antifibrinolytics, Haemostatics and	
Adalat Oros		Alphamox		Local Sclerosants	
Adalimumab		Alphamox 125		Antifibrotics	
Adapalene		Alphamox 250		Antifungals	
Adcortyl		Alprolix		Antifungals Topical	
Adefin		Alu-Tab		Antihistamines	
Adefin XL		Aluminium hydroxide		Antihypotensives	
Adefovir dipivoxil		Alvogen		Antimalarials	
Adenuric		Amantadine hydrochloride		Antimigraine Preparations	
ADR Cartridge 1.8		Ambrisentan		Antinausea and Vertigo Agents	
Adrenaline		Amiloride hydrochloride	53	Antiparasitics	
ADT Booster	270	Amiloride hydrochloride with		Antipruritic Preparations	
Adult diphtheria and tetanus	070	furosemide	53	Antipsychotics	
vaccine		Amiloride hydrochloride with	E0	Antiretrovirals	
Advata		hydrochlorothiazide		Antirheumatoid Agents	
AdvateAdynovate		Aminophylline Amiodarone hydrochloride		Antispasmodics and Other Agents	
Afinitor		Amisulpride		Altering Gut Motility Antithrombotic Agents	
/\!!!!!!\U!	200	Amauphuc	102	ATTURNOTION AGEIRS	4

Antithymocyte globulin		Aristocort	64	Azithromycin	8
(equine)		Arrow-Amitriptyline		Azol	
Antitrichomonal Agents	98	Arrow-Bendrofluazide		Azopt	
Antituberculotics and		Arrow-Brimonidine	243	AZT	10
Antileprotics		Arrow-Calcium		- B -	
Antiulcerants	8	Arrow-Diazepam	136	B-D Micro-Fine	
Antivirals		Arrow-Doxorubicin	163	B-D Ultra Fine	
Anxiolytics	136	Arrow-Fluoxetine		B-D Ultra Fine II	14
Anzatax		Arrow-Lamotrigine	128	Bacillus Calmette-Guerin (BCG)	
Apidra		Arrow-Losartan &		vaccine	180
Apidra SoloStar		Hydrochlorothiazide		Bacillus Calmette-Guerin	
Apo-Amlodipine		Arrow-Morphine LA		vaccine	
Apo-Azithromycin		Arrow-Norfloxacin		Baclofen	
Apo-Bromocriptine		Arrow-Ornidazole		Bactroban	
Apo-Ciclopirox	62	Arrow-Quinapril 10		Barrier Creams and Emollients	
Apo-Cilazapril/		Arrow-Quinapril 20		BCG Vaccine	
Hydrochlorothiazide	48	Arrow-Quinapril 5		Beclazone 100	
Apo-Clarithromycin		Arrow-Roxithromycin		Beclazone 250	
Alimentary	9	Arrow-Timolol	242	Beclazone 50	
Infection		Arrow-Topiramate		Beclomethasone dipropionate	
Apo-Clomipramine		Arrow-Tramadol	125	Bee venom allergy treatment	
Apo-Diclo SR		Arsenic trioxide		Bendamustine hydrochloride	15
Apo-Diltiazem CD		Asacol	7	Bendrofluazide	5
Apo-Doxazosin		Asamax		Bendroflumethiazide	
Apo-Folic Acid		Ascorbic acid		[Bendrofluazide]	54
Apo-Furosemide	53	Aspen Adrenaline	56	Benzathine benzylpenicillin	9
Apo-Gabapentin	128	Aspirin		Benzatropine mesylate	120
Apo-Leflunomide		Blood	42	Benzbromaron AL 100	
Apo-Megestrol		Nervous	121	Benzbromarone	110
Apo-Metoprolol		Asthalin	235	Benztrop	120
Apo-Mirtazapine	127	Atazanavir sulphate	106	Benzydamine hydrochloride	3
Apo-Nadolol	51	Atenolol	50	Benzylpenicillin sodium [Penicillin	
Apo-Nicotinic Acid	54	Atenolol AFT	50	G]	9 [.]
Apo-Oxybutynin	75	Atenolol AFT S29	50	Beta Cream	6
Apo-Perindopril	47	ATGAM	186	Beta Ointment	6
Apo-Pindolol	51	Ativan	136	Beta Scalp	70
Apo-Pravastatin	55	Atomoxetine	150	Beta-Adrenoceptor Agonists	23
Apo-Prazosin	47	Atorvastatin	54	Beta-Adrenoceptor Blockers	
Apo-Prednisone	79	Atropine sulphate		Betadine	6
Apo-Primidone	129	Cardiovascular	49	Betadine Skin Prep	6
Apo-Propranolol	51	Sensory	243	Betaferon	14
Apo-Pyridoxine	34	Atropt		Betahistine dihydrochloride	13
Apo-Selegiline S29		Atrovent	235	Betaine	
Apo-Sumatriptan	131	AU Synacthen	79	Betaloc CR	5
Apo-Temozolomide	167	Aubagio	142	Betamethasone dipropionate	6
Apo-Terazosin	47	Augmentin	91	Betamethasone dipropionate with	
Apo-Timol		Aurorix		calcipotriol	6
Apomorphine hydrochloride	119	AutoSoft 30	<mark>23</mark>	Betamethasone sodium phosphat	te
Aprepitant		AutoSoft 90		with betamethasone acetate	78
Apresoline	56	Avelox	<mark>93</mark>	Betamethasone valerate	.63, 70
Aptamil Gold+ Pepti Junior		Avonex		Betamethasone valerate with	
Aqueous cream		Avonex Pen	145	clioquinol	6
Aratac		Azacitidine		Betamethasone valerate with sod	
Arava	111	Azacitidine Dr Reddy's		fusidate [fusidic acid]	6
Aripiprazole	133	Azamun		Betaxolol	
Aripiprazole Sandoz		Azathioprine		Betnovate	

INDEX: Generic Chemicals and Brands

Betnovate-C	64	Budesonide with eformoterol	234	Carmustine	15
Betoptic	242	Bumetanide	53	Carvedilol	5
Betoptic S		Buprenorphine Naloxone BNM		Carvedilol Sandoz	
Bezafibrate		Buprenorphine with naloxone	154	Catapres	5
Bezalip	54	Bupropion hydrochloride		CeeNU	15
Bezalip Retard		Burinex		Cefaclor monohydrate	
Bicalutamide		Burinex S29	53	Cefalexin	8
Bicillin LA		Buscopan	8	Cefalexin Sandoz	8
BiCNU	158	Buspirone hydrochloride	136	Cefazolin	
Bicnu Heritage	158	Busulfan		Ceftriaxone	
Bile and Liver Therapy		- C -		Ceftriaxone-AFT	8
Biltricide		Cabergoline	87	Cefuroxime axetil	
Bimatoprost	243	Caffeine citrate	239	Celebrex	11
Bimatoprost Multichem	243	Calamine	63	Celecoxib	11
Binarex		Calcipotriol	69	Celecoxib Pfizer	11
Binocrit	39	Calcitonin	77	Celestone Chronodose	7
Biodone	123	Calcitriol		Celiprolol	5
Biodone Extra Forte	123	Calcitriol-AFT	34	Cellcept	
Biodone Forte	123	Calcium carbonate	6, 35	Celol	
Bisacodyl		Calcium Channel Blockers		Centrally-Acting Agents	5
Bisoprolol fumarate		Calcium Disodium Versenate	246	Cephalexin ABM	
BK Lotion		Calcium folinate	160	Cetirizine hydrochloride	
Bleomycin sulphate		Calcium Folinate Ebewe	160	Cetomacrogol	6
Blood Colony-stimulating		Calcium Folinate Sandoz	160	Cetomacrogol with glycerol	
Factors	44	Calcium gluconate		Cetuximab	
Blood glucose diagnostic test		Calcium Homeostasis	77	Charcoal	24
meter	13	Calcium polystyrene sulphonate		Chemotherapeutic Agents	
Blood glucose diagnostic test		Calcium Resonium		Chickenpox vaccine	
strip	13	Calcium Sandoz		Chlorafast	
Blood glucose test strips (visual		Calogen	252	Chlorambucil	
impaired)	•	Candesartan cilexetil		Chloramphenicol	
Blood Ketone Diagnostic Test		Candestar		Chlorhexidine gluconate	
Strip	12	Canesten	62	Alimentary	3
Bonjela		Capecitabine		Dermatological	
Boostrix		Capercit	160	Chloroform	
Bortezomib		Capoten	47	Chlorothiazide	5
Bortezomib Dr-Reddy's		Capsaicin		Chlorpheniramine maleate	
Bosentan		Musculoskeletal	111	Chlorpromazine hydrochloride	
Bosentan Dr Reddy's		Nervous	122	Chlortalidone [Chlorthalidone]	
Bosvate		Captopril	47	Chlorthalidone	
Bplex	34	Carafate	10	Chlorvescent	4
Breo Ellipta	234	Carbaccord	158	Choice Load 375	
Brevinor 1/28		Carbamazepine	127	Choice TT380 Short	7
Bricanyl Turbuhaler	235	Carbimazole		Choice TT380 Standard	7
Brilinta	42	Carbomer	244	Choline salicylate with cetalkonium	ı
Brimonidine tartrate	243	Carboplatin	158	chloride	3
Brimonidine tartrate with timolol		Carboplatin Ebewe	158	Ciclopirox olamine	
maleate	243	Carbosorb-X		Ciclosporin	
Brinzolamide	242	Cardinol LA		Cilazapril	
Brolene		CareSens Dual		Cilazapril with	
Bromocriptine mesylate		CareSens N	13	hydrochlorothiazide	4
Brufen SR		CareSens N POP		Cilicaine	9
Buccastem		CareSens N Premier		Cilicaine VK	9
Budesonide		CareSens PRO		Cinacalcet	
Alimentary	6	Carmellose sodium with gelatin a		Cipflox	
Respiratory		pectin			

Ciprofloxacin	Colestipol hydrochloride54	Daktarin6
Infection93	Colgout 117	Dalacin C9
Sensory240	Colifoam7	Danazol8
Ciprofloxacin Teva240	Colistin sulphomethate93	Dantrium11
Circadin149	Colistin-Link 93	Dantrium S2911
Cisplatin	Collodion flexible248	Dantrolene11
Cisplatin Ebewe158	Colloidal bismuth subcitrate10	Daonil1
Citalopram hydrobromide126	Colofac8	Dapa-Tabs5
Cladribine	Coloxyl27	Dapsone9
Clarithromycin	Combigan243	Daraprim9
Alimentary9	Compound electrolytes46	Darunavir10
Infection89	Compound electrolytes with glucose	Dasatinib17
Clexane43	[Dextrose]46	Daunorubicin16
Clexane Forte43	Compound hydroxybenzoate248	DBL Acetylcysteine24
Climara80	Concerta152	DBL Adrenaline5
Climara80	Condoms71	DBL Aminophylline23
Clindamycin93	Condyline70	DBL Bleomycin Sulfate16
Clinicians Renal Vit34	Condyline S2970	DBL Carboplatin15
Clobazam127	Contraceptives - Hormonal72	DBL Cisplatin15
Clobetasol propionate	Contraceptives - Non-hormonal71	DBL Dacarbazine16
Clobetasone butyrate64	Copaxone144	DBL Desferrioxamine Mesylate for Inj
Clofazimine98	Corticosteroids and Related Agents	BP24
Clomazol	for Systemic Use78	DBL Docetaxel16
Dermatological62	Corticosteroids Topical	DBL Ergometrine
Genito-Urinary74	Cosentyx221	DBL Gemcitabine16
Clomifene citrate87	Cosmegen 163	DBL Gentamicin
Clomipramine hydrochloride125	Coumadin44	DBL Leucovorin Calcium16
Clonazepam127, 136	Creon 1000026	DBL Methotrexate Onco-Vial16
Clonidine	Creon 2500026	DBL Morphine Sulphate12
Clonidine BNM	Creon Micro	DBL Morphine Tartrate12
Clonidine hydrochloride52	Crotamiton63	DBL Naloxone Hydrochloride24
Clopidogrel42	Crystaderm61	DBL Octreotide17
Clopidogrel Multichem42	Curam91	DBL Pethidine Hydrochloride 12
Clopine	Cvite34	DBL Vinblastine
Clopixol	Cyclizine hydrochloride131	DBL Vincristine Sulfate16
Clotrimazole	Cyclizine lactate131	De-Worm8
Dermatological62	Cyclogyl243	Decozol3
Genito-Urinary74	Cyclopentolate hydrochloride243	Deferasirox24
Clozapine	Cyclophosphamide158	Deferiprone24
Clozaril133	Cyclorin98	Denosumab11
Clustran	Cycloserine98	Deolate9
Co-trimoxazole95	Cyproterone acetate79	Deoxycoformycin16
Coal tar	Cyproterone acetate with	Depo-Medrol7
Coal tar with allantoin, menthol,	ethinyloestradiol74	Depo-Provera7
phenol and sulphur69	Cystadane29	Depo-Testosterone79
Coal tar with salicylic acid and	Cytarabine	Deprim9
sulphur69	Cytotec8	DermAssist6
Coco-Scalp69	Cytoxan158	Dermol63, 7
Codeine phosphate	- D -	Desferrioxamine mesilate24
Extemporaneous248	D-Penamine111	Desmopressin acetate8
Nervous123	Dabigatran44	Desmopressin-PH&T8
Cogentin	Dacarbazine	Desuric11
Colaspase [L-asparaginase]163	Dacarbazine APP163	Detection of Substances in
Colchicine	Dactinomycin [Actinomycin D]163	Urine7
Colecalciferol	Daivobet69	Dexamethasone
Colestid	Daivonex	Hormone7

INDEX: Generic Chemicals and Brands

Sensory241	Agents 65	factor IX]	3
Dexamethasone phosphate78	Disopyramide phosphate49	Efudix	7
Dexamethasone Phosphate	Disulfiram155	Egopsoryl TA	
Panpharma78	Diuretics53	Elaprase	3
Dexamethasone with framycetin and	Docetaxel163	Elecare	26
gramicidin240	Docetaxel Accord163	Elecare LCP	
Dexamethasone with neomycin	Docetaxel Sandoz163	Electral	4
sulphate and polymyxin B	Docusate sodium27	Elelyso	3
sulphate241	Docusate sodium with	Elemental 028 Extra	25
Dexamfetamine sulfate150	sennosides27	Elocon	6
Dexmethsone78	Dolutegravir106	Elocon Alcohol Free	
Dextrochlorpheniramine	Domperidone 131	Eltrombopag	3
maleate233	Donepezil hydrochloride153	Eltroxin	8
Dextrose45–46	Donepezil-Rex153	EMB Fatol	9
DHC Continus123	Dornase alfa238	Emend Tri-Pack	
Diabetes10	Dortimopt242	EMLA	12
Diabetes Management12	Dorzolamide hydrochloride242	Emtricitabine	
Diacomit	Dorzolamide with timolol242	Emtricitabine with tenofovir	
Diagnostic Agents281	Dostinex87	disoproxil	10
Diamide Relief6	Dosulepin [Dothiepin]	Emtriva	
Diamox242	hydrochloride125	Emulsifying ointment	6
Diasip253	Dosulepin Mylan125	Enalapril maleate	4
Diason RTH253	Dothiepin	Enbrel	
Diazepam127, 136	Doxazosin47	Endocrine Therapy	
Diazoxide10	Doxine92	Endoxan	
Dibenzyline47	Doxorubicin Ebewe163	Engerix-B	27
Diclofenac Sandoz110	Doxorubicin hydrochloride163	Enlafax XR	
Diclofenac sodium	Doxycycline92	Enoxaparin sodium	4
Musculoskeletal110	DP Lotion66	Enstilar	
Sensory241	DP Lotn HC64	Ensure	26
Differin61	DP-Allopurinol116	Ensure Plus	26
Difflam33	Dr Reddy's Omeprazole9	Ensure Plus HN	26
Diflucan95	Drugs Affecting Bone	Ensure Plus RTH	
Diflucan S2995	Metabolism 111	Entacapone	11
Diflucortolone valerate64	Dual blood glucose and blood ketone	Entapone	
Digestives Including Enzymes26	diagnostic test meter 12	Entecavir	
Digoxin49	Duocal Super Soluble Powder251	Entecavir Sandoz	
Dihydrocodeine tartrate123	Duolin	Entocort CIR	
Dilantin	Duolin HFA236	Entresto 24/26	
Dilantin Infatab128	Duride56	Entresto 49/51	4
Diltiazem hydrochloride52	-E-	Entresto 97/103	4
Dilzem52	e-chamber La Grande239	Epilim	
Dimethicone65, 67	e-chamber Mask239	Epilim Crushable	
Dimethyl fumarate	e-chamber Turbo239	Epilim IV	
Dipentum7	E-Mycin90	Epilim S/F Liquid	
Diphtheria, tetanus and pertussis	Ear Preparations240	Epilim Syrup	12
vaccine 270	Ear/Eye Preparations240	Epirubicin Ebewe	16
Diphtheria, tetanus, pertussis and	Easiphen Liquid265	Epirubicin hydrochloride	
polio vaccine271	Econazole nitrate62	Eplerenone	
Diphtheria, tetanus, pertussis, polio,	Efavirenz105	Epoetin alfa	
hepatitis B and haemophilus	Efavirenz with emtricitabine and	Epoprostenol	
influenzae type B vaccine 271	tenofovir disoproxil105	Eptacog alfa [Recombinant factor	_
Diprosone63	Effient	VIIa]	4
Diprosone OV63	Eformoterol fumarate234	ERA	
Dipyridamole42	Eformoterol fumarate dihydrate234	Erbitux	
Disinfecting and Cleansing	Eftrenonacog alfa [Recombinant	Ergometrine maleate	
g and croanong			

Erlotinib	170	Felodipine		Fluox	126
Erythrocin IV	90	Femme-Tab ED	73	Fluoxetine hydrochloride	126
Erythromycin (as lactobionate)	90	Fentanyl		Flupenthixol decanoate	134
Erythromycin ethyl succinate	90	Fentanyl GH	123	Flutamide	177
Erythromycin stearate	90	Fentanyl IE	123	Flutamin	177
Esbriet	237	Fentanyl Sandoz		Fluticasone	234
Escitalopram	126	Ferinject	35	Fluticasone furoate with	
Escitalopram-Apotex	126	Ferodan		vilanterol	234
Eskazole		Ferric carboxymaltose	35	Fluticasone propionate	239
Estradot	80	Ferriprox	245	Fluticasone with salmeterol	
Estradot 50 mcg	80	Ferro-F-Tabs	36	FML	242
Estrofem		Ferro-tab	36	Foban	62
Etanercept	179	Ferrograd	36	Folic acid	39
Ethambutol hydrochloride		Ferrosig		Food Thickeners	262
Ethics Aspirin		Ferrous fumarate		Foods And Supplements For Ir	nborn
Ethics Aspirin EC		Ferrous fumarate with folic acid.		Errors Of Metabolism	
Ethics Lisinopril		Ferrous sulfate	36	Foradil	
Ethics Paracetamol Classic		Ferrous sulphate		Forteo	
Ethinyloestradiol		Fexofenadine hydrochloride		Fortini	
Ethinyloestradiol with		Fibro-vein		Fortini Multi Fibre	
desogestrel	72	Filgrastim		Fortisip	
Ethinyloestradiol with		Finasteride		Fortisip Multi Fibre	
levonorgestrel	73	Fingolimod		Fosamax	111
Ethinyloestradiol with		Firazyr		Fosamax Plus	
norethisterone	73	Flagyl		Framycetin sulphate	
Ethosuximide		Flagyl-S		Frisium	
Etopophos		Flamazine		Frumil	
Etoposide		Flecainide acetate		Frusemide	
Etoposide phosphate		Flecainide BNM		Frusemide-Claris	
Etravirine		Flecainide Controlled Release		Fucicort	
Eumovate		Teva	49	Fucidin	
Everet		Fleet Phosphate Enema		Fucithalmic	
Everolimus		Flixonase Hayfever & Allergy		Fulvestrant	
Evista		Flixotide		Fungilin	
Exemestane		Flixotide Accuhaler		Furosemide [Frusemide]	
Exjade		Floair		fusidic acid	
Extemporaneously Compounded		Florinef		Dermatological	62 65
Preparations and		Fluanxol		Infection	
Galenicals	248	Flucil		Sensory	
Eye Preparations		Flucloxacillin		- G -	270
Eylea		Flucloxin		Gabapentin	109
Ezetimibe		Flucon		Gacet	
Ezetimibe Sandoz		Fluconazole		Galsulfase	
Ezetimibe sandoz		Fludara Oral		Galvumet	
- F -		Fludarabine Ebewe		Galvus	
				Gardasil 9	
Factor eight inhibitor bypassing	40	Fludragertisens agetate		Gastrodenol	
fraction		Fluids and Floatralities		Gaviscon Double Strength	
Famotidine		Fluids and Electrolytes		Caviscon Infant	
Famotidine Hovid		Flumetasone pivalate	240	Gaviscon Infant	
Faslodex		Fluocortolone caproate with		Gazyva	
Febuxostat	117	fluocortolone pivalate and	0	Gefitinib	
Feed Thickener Karicare	000	cinchocaine		Gemcitabine Ebewe	
Aptamil		Fluorometholone		Gemcitabine hydrochloride	
FEIBA NF		Fluorouracil		Gemfibrozil	
Felo 10 ER		Fluorouracil Ebewe		Gemzar	
Felo 5 ER	52	Fluorouracil sodium	/U	Generic Partners	. 150, 153

INDEX: Generic Chemicals and Brands

Genoptic240	Healtheries Simple Baking Mix263	Hylo-Fresh24
Gentamicin sulphate	Hemastix76	Hymenoptera23
Infection93	Heparin Ratiopharm44	Hyoscine butylbromide
Sensory240	Heparin sodium44	Hyoscine hydrobromide13
Gilenya 138	Heparinised saline44	Hypam15
Ginet74	Heparon Junior254	Hyperuricaemia and Antigout 11
Glatiramer acetate144	Hepatitis A vaccine272	Hypromellose24
Glecaprevir with pibrentasvir 102	Hepatitis B recombinant	Hypromellose with dextran24
Glibenclamide11	vaccine 273	-1-
Gliclazide12	Hepsera100	Ibiamox9
Glipizide12	Herceptin226	lbilex8
Glivec171	Hiberix271	Ibrance17
Glizide12	Hiprex109	lbuprofen11
Glucagen Hypokit10	Histaclear233	Ibuprofen SR BNM11
Glucagon hydrochloride10	Histafen233	lcatibant23
Glucerna Select253	Holoxan158	Idarubicin hydrochloride16
Glucerna Select RTH253	Horleys Bread Mix263	Idursulfase3
Glucobay11	Horleys Flour263	Ifosfamide15
Glucose [Dextrose]45	Hormone Replacement Therapy -	Ikorel5
Gluten Free Foods263	Systemic	lloprost5
Glycerin with sodium saccharin248	HPV274	Imatinib mesilate17
Glycerin with sucrose248	Humalog11	Imatinib-AFT17
Glycerol	Humalog Mix 2511	Imigran13
Alimentary28	Humalog Mix 5011	Imipramine hydrochloride 12
Extemporaneous248	Human papillomavirus (6, 11, 16, 18,	Imiguimod
Glyceryl trinitrate	31, 33, 45, 52 and 58) vaccine	Immune Modulators10
Alimentary8	[HPV]274	Immunosuppressants
Cardiovascular56	Humatin	Imuran
Glycopyrronium236	Humira186	Incruse Ellipta23
	HumiraPen186	Indacaterol23
Glycopyrronium bromide8	Humulin 30/7011	Indapamide5
Glycopyrronium with indacaterol236	Humulin NPH11	Infanrix IPV27
Gold Knight71	Humulin R	Infanrix-hexa27
Goserelin86	Hyaluronic acid244	Infant Formulae26
Gutron	Hydralazine56	Infatrini
Gynaecological Anti-infectives74	Hydralazine hydrochloride56	Infliximab19
-H-	Hydrea164	Influenza vaccine27
Habitrol	Hydrocortisone	Influvac Tetra
Haemophilus influenzae type B	Dermatological64	(2020 formulation)
vaccine271	Hormone	Inhaled Corticosteroids23
Haldol	Hydrocortisone (PSM)64	Inhaled Long-acting
Haldol Concentrate135	Hydrocortisone acetate7	Beta-adrenoceptor Agonists 23
Haldol Decanoas	Hydrocortisone acetate with	Inspra5
Haloperidol133	pramoxine hydrochloride	Instillagel Lido12
Haloperidol decanoate135	Hydrocortisone and paraffin liquid	Insulin aspart1
Harvoni102	and lanolin 64	Insulin aspart with insulin aspart
Havrix272	Hydrocortisone butyrate64, 70	protamine1
Havrix Junior272	Hydrocortisone with cinchocaine8	Insulin glargine1
HBvaxPRO273	Hydrocortisone with miconazole65	Insulin glulisine1
healthE Calamine Aqueous Cream	Hydrocortisone with natamycin and	Insulin isophane1
BP63	neomycin65	Insulin isophane with insulin
healthE Dimethicone 10%65	Hydrogen peroxide61	neutral1
healthE Dimethicone 4% Lotion67	Hydroxocobalamin34	Insulin lispro1
healthE Dimethicone 5%65	Hydroxychloroquine111	Insulin lispro with insulin lispro
healthE Glycerol BP248	Hydroxyurea164	protamine1
healthE Urea Cream66	Hygroton54	Insulin neutral1

Insulin pen needles	14	Itrazole	96	Lapatinib ditosylate	172
Insulin pump		Ivermectin		Largactil	133
Insulin pump cartridge		- J -		Laronidase	
Insulin pump infusion set (steel		Jadelle	73	Lasix	
cannula)	20	Jakavi		Latanoprost	
Insulin pump infusion set (steel	0	Jaydess		Lax-Suppositories	
cannula, straight insertion)	21	Jevity HiCal RTH		Lax-Tab	
Insulin pump infusion set (teflon	2 1	Jevity RTH	260	Laxatives	
cannula)	22	Juno Pemetrexed		Laxsol	
Insulin pump infusion set (teflon	22	- K -	101	Ledipasvir with sofosbuvir	
cannula, angle insertion with		Kadcyla	227	Leflunomide	
insertion device)	23	Kaletra		Lenalidomide	
Insulin pump infusion set (teflon	20	Keflor		Letrole	
cannula, angle insertion)	23	Kemadrin		Letrozole	
Insulin pump infusion set (teflon	20	Kenacomb		Leukeran FC	
		Kenacort-A 10			130
cannula, straight insertion with	04			Leukotriene Receptor	220
insertion device)	24	Kenacort-A 40		Antagonists	
Insulin pump infusion set (teflon	0.5	Kenalog		Leunase	
cannula, straight insertion)		Kenalog in Orabase		Leuprorelin	
Insulin pump reservoir	25	Ketocal 3:1		Leustatin	
Insulin syringes, disposable with		KetoCal 4:1	269	Levetiracetam	
attached needle		Ketoconazole		Levetiracetam-AFT	
Intal Forte CFC Free		Dermatological		Levlen ED	
Intelence		Infection		Levocabastine	
Interferon alfa-2a		Ketogenic Diet		Levodopa with benserazide	
Interferon beta-1-alpha		Ketoprofen		Levodopa with carbidopa	
Interferon beta-1-beta		KetoSens		Levomepromazine	133
Intra-uterine device		Keytruda		Levomepromazine	
Invega Sustenna		Kindergen		hydrochloride	133
IPOL		Kivexa	105	Levonorgestrel	
Ipratropium bromide235	, 239	Klacid		Genito-Urinary	
Iressa		Kliogest	80	Hormone	
Irinotecan Accord	.161	Kliovance	80	Levothyroxine	82
Irinotecan Actavis 100	.161	Kogenate FS	41	Lidocaine [Lignocaine]	121
Irinotecan hydrochloride	.161	Konakion MM	42	Lidocaine [Lignocaine]	
Irinotecan-Rex		Konsyl-D	27	hydrochloride	121
Iron polymaltose	36	Kuvan	30	Lidocaine [Lignocaine] with	
Isentress	.106	-L-		chlorhexidine	121
Isentress HD	.106	L-asparaginase	163	Lidocaine [Lignocaine] with	
Ismo 20	56	Labetalol	50	prilocaine	121
Ismo 40 Retard	56	Lacosamide	128	Lidocaine-Claris	
Isoniazid	99	Lactulose	28	Lignocaine	121
Isoniazid with rifampicin	99	Laevolac	28	Lioresal Intrathecal	
Isoprenaline [Isoproterenol]	56	Lamictal	128	Lipazil	
Isoproterenol	56	Lamivudine	101, 106	Lipid-Modifying Agents	54
Isoptin	52	Lamivudine Alphapharn	n106	Liquigen	
Isoptin Retard	52	Lamotrigine	128	Lisinopril	
Isoptin SR	52	Lamprene		Lithicarb FC	133
Isopto Carpine		Lanoxin		Lithium carbonate	133
Isosorbide mononitrate		Lanoxin PG		Livostin	242
Isosource Standard		Lanoxin S29	49	LMX4	121
Isotretinoin	61	Lansoprazole		Locacorten-Viaform ED's	240
Ispaghula (psyllium) husk		Lantus		Local preparations for Anal and	
Isuprel		Lantus SoloStar		Rectal Disorders	8
Itch-Soothe		Lanvis		Locasol	
Itraconazole		Lanzol Relief		Locoid	

Locoid Crelo	64	Maxitrol241	succinate)	7
Locoid Lipocream	64	MCT oil (Nutricia)252	Methylprednisolone aceponate	6
Locorten-Vioform	240	Measles, mumps and rubella	Methylprednisolone acetate	7
Lodoxamide	242	vaccine277	Methylxanthines	
Logem	128	Mebendazole88	Metoclopramide Actavis 10	13
Lomide		Mebeverine hydrochloride8	Metoclopramide hydrochloride	13
Lomustine	158	Medco122	Metolazone	5
Loniten	57	Medrol78	Metopirone	8
Loperamide hydrochloride	6	Medroxyprogesterone acetate	Metoprolol succinate	5
Lopinavir with ritonavir		Genito-Urinary74	Metoprolol tartrate	
Loprofin	265	Hormone80–81	Metrogyl	9
Loprofin Mix		Mefenamic acid110	Metronidazole	9
Lorafix	233	Megestrol acetate177	Metroprolol IV Mylan	
Loratadine	233	Melatonin149	Metyrapone	
Lorazepam	136	Melphalan158	Mexiletine hydrochloride	
Lorfast	233	Menactra277	Mexiletine Hydrochloride USP	
Lorstat		Meningococcal (groups A, C, Y and	Miacalcic	
Losartan Actavis	48	W-135) conjugate vaccine 277	Micolette	
Losartan potassium	48	Meningococcal C conjugate	Miconazole	
Losartan potassium with		vaccine278	Miconazole nitrate	
hydrochlorothiazide	48	Menthol63	Dermatological	6
Lovir		Mepolizumab204	Genito-Urinary	
Loxamine		Mercaptopurine161	Micreme	
Lucrin Depot 1-month	86	Mercilon 2872	Micreme H	
Lucrin Depot 3-month		Mesalazine7	Microgynon 20 ED	
Ludiomil		Mesna165	Microgynon 30	
Lupin		Mestinon110	Microgynon 50 ED	
Lyderm		Metabolic Disorder Agents28	Microlut	
Lynparza		Meterol234	Midazolam	14
- M -		Metformin hydrochloride12	Midazolam-Claris	14
m-Eslon	124	Methadone hydrochloride	Midodrine	5
Mabthera	207	Extemporaneous248	Mifegyne	
Macrogol 3350 with potassium		Nervous123	Mifepristone	
chloride, sodium bicarbonate	e and	Methatabs123	Minerals	
sodium chloride	28	Methenamine (hexamine)	Mini-Wright AFS Low Range	
Macrogol 400 and propylene		hippurate109	Mini-Wright Standard	23
glycol	244	Methopt243	Minidiab	1
Madopar 125		Methotrexate161	MiniMed 640G	1
Madopar 250		Methotrexate Ebewe161	MiniMed Mio MMT-921A	
Madopar 62.5		Methotrexate Sandoz161	MiniMed Mio MMT-923A	
Madopar HBS		Methyl hydroxybenzoate248	MiniMed Mio MMT-925A	2
Madopar Rapid		Methylcellulose248	MiniMed Mio MMT-941A	2
Magnesium hydroxide		Methylcellulose with glycerin and	MiniMed Mio MMT-943A	
Alimentary	36	sodium saccharin248	MiniMed Mio MMT-945A	2
Extemporaneous	248	Methylcellulose with glycerin and	MiniMed Mio MMT-965A	2
Magnesium sulphate		sucrose249	MiniMed Mio MMT-975A	2
Mantoux		Methyldopa52	MiniMed Quick-Set MMT-386A	2
Maprotiline hydrochloride	125	Methyldopa Mylan52	MiniMed Quick-Set MMT-387A	2
Marevan		Methyldopa Mylan S2952	MiniMed Quick-Set MMT-396A	2
Marine Blue Lotion SPF 50+	70	Methylnaltrexone bromide27	MiniMed Quick-Set MMT-397A	2
Martindale Pharma	245	Methylphenidate ER - Teva152	MiniMed Quick-Set MMT-398A	2
Marvelon 28		Methylphenidate hydrochloride 151	MiniMed Quick-Set MMT-399A	
Mask for spacer device	239	Methylphenidate hydrochloride	MiniMed Silhouette MMT-368A	
Mast Cell Stabilisers		extended-release152	MiniMed Silhouette MMT-377A	
Maviret		Methylprednisolone78	MiniMed Silhouette MMT-378A	
Maxidex	241	Methylprednisolone (as sodium	MiniMed Silhouette MMT-381A	

MiniMed Silhouette MMT-382A.	22	Mycobutin	99	Nifedipine	5
MiniMed Silhouette MMT-383A.	22	MycoNail	62	Nifuran	10
MiniMed Silhouette MMT-384A.	22	Mycophenolate mofetil	179	Nilotinib	17
MiniMed Sure-T MMT-864A	20	Mydriacyl	243	Nilstat	
MiniMed Sure-T MMT-866A	20	Mylan Atenolol	50	Alimentary	3
MiniMed Sure-T MMT-874A	20	Mylan Clomiphen		Genito-Urinary	74
MiniMed Sure-T MMT-876A	20	Myleran		Infection	9
MiniMed Sure-T MMT-884A	20	Myometrial and Vaginal Hormo		Nintedanib	
MiniMed Sure-T MMT-886A	20	Preparations		Nipent	16
Minims Pilocarpine	243	Myozyme	28	Nitrados	
Minims Prednisolone	242	Mysoline S29		Nitrates	5
Minirin		- N -		Nitrazepam	
Mino-tabs	92	Nadolol	51	Nitroderm TTS	
Minocycline hydrochloride	92	Naglazyme		Nitrofurantoin	
Minomycin		Nalcrom	8	Nitrolingual Pump Spray	5
Minor Śkin Infections		Naloxone hydrochloride	245	Nivestim	
Minoxidil		Naltraccord		Nivolumab	
Mirena		Naltrexone hydrochloride		Nizoral	
Mirtazapine		Naphazoline hydrochloride		Nodia	
Misoprostol		Naphcon Forte		Noflam 250	
Mitomycin C	165	Naprosyn SR 1000	110	Noflam 500	
Mitozantrone		Naprosyn SR 750		Non-Steroidal Anti-Inflammatory	
Mitozantrone Ebewe		Naproxen		Drugs	110
Mixtard 30		Narcaricin mite		Nonacog gamma, [Recombinant	
MMR II		Nardil		Factor IX]	4
Moclobemide		Nardil S29		Norethisterone	
Modafinil		Nasal Preparations		Genito-Urinary	7,
Modavigil		Natalizumab		Hormone	
Moduretic		Natulan		Norflex	
Molaxole		Nausafix		Norfloxacin	
Moments		Nausicalm		Noriday 28	
Mometasone furoate		Navelbine		Norimin	
Monogen		Necon		Normacol Plus	
Montelukast		Nedocromil		Normison	
Montelukast Mylan		Nefopam hydrochloride		Norpress	
Moroctocog alfa [Recombinant fa		Neisvac-C		Nortriptyline hydrochloride	
VIII]		Neo-B12		Norvir	
Morphine hydrochloride		Neo-Mercazole		NovaSource Renal	
Morphine sulphate		Neocate Gold		Novatretin	
Morphine tartrate		Neocate Junior Unflavoured		NovoMix 30 FlexPen	
Motetis		Neocate Junior Vanilla		NovoRapid	
Mouth and Throat		Neocate SYNEO		NovoRapid FlexPen	
Movapo		Neoral		NovoRapid Penfill	
Moxifloxacin		Neostigmine metilsulfate		NovoSeven RT	<u>4</u>
MSUD Maxamum		Nepro HP (strawberry)		Noxafil	
Mucilaginous laxatives with	204	Nepro HP (vanilla)		Nozinan	
stimulants	27	Nepro HP RTH		Nozinan (Swiss)	
Mucolytics		Nerisone		Nucala	
Mucosoothe		Neulactil		Nuelin	
Multiple Sclerosis Treatments		Neulastim		Nuelin-SR	201
Multivitamin renal		NeuroTabs		Nutilis	
Multivitamins		Nevirapine		Nutrient Modules	
Mupirocin		Nevirapine Alphapharm	105	Nutrini Energy Multi Fibre	
Muscle Relaxants	118	Nicorandil		Nutrini Energy RTH	
Mvite		Nicotine		Nutrini Low Energy Multi Fibre	
Myambutol		Nicotinic acid		Nutrini RTH	
IVI V UI I IDULUI					

			_
Nutrison 800 Complete Multi	Ondansetron ODT-DRLA132	Pan-Penicillin G Sodium	9
Fibre260	One-Alpha34	Pancreatic enzyme	
Nutrison Concentrated262	Onrex132	Pantoprazole	
Nutrison Energy260	Opdivo228	Panzop Relief	
Nutrison Energy Multi Fibre260	Ora-Blend249	Panzytrat	2
Nutrison Multi Fibre260	Ora-Blend SF248	Papaverine hydrochloride	5
Nutrison Standard RTH260	Ora-Plus248	Para-amino salicylic acid	9
Nyefax Retard52	Ora-Sweet248	Paracare	.12
Nystatin	Ora-Sweet SF248	Paracare Double Strength	.12
Alimentary33	Orabase33	Paracetamol	.12
Genito-Urinary74	Oral Supplements/Complete Diet	Paracetamol + Codeine	
Infection96	(Nasogastric/Gastrostomy Tube	(Relieve)	. 12
NZB Low Gluten Bread Mix263	Feed) 253	Paracetamol Pharmacare	. 12
- 0 -	Oratane61	Paracetamol with codeine	
O/W Fatty Emulsion Cream66	Ordine123	Paradigm 1.8 Reservoir	2
Obinutuzumab205	Orgran264	Paradigm 3.0 Reservoir	2
Obstetric Preparations76	Ornidazole98	Paradigm Mio MMT-921	
Ocicure9	Orphenadrine citrate118	Paradigm Mio MMT-923	2
Ocrelizumab141	Ortho-tolidine76	Paradigm Mio MMT-925	
Ocrevus141	Oruvail SR110	Paradigm Mio MMT-941	2
Octocog alfa [Recombinant factor	Osmolite RTH260	Paradigm Mio MMT-943	2
VIII] (Advate)41	Other Endocrine Agents87	Paradigm Mio MMT-945	2
Octocog alfa [Recombinant factor	Other Oestrogen Preparations81	Paradigm Mio MMT-965	2
VIII] (Kogenate FS)41	Other Progestogen	Paradigm Mio MMT-975	2
Octreotide177	Preparations81	Paradigm Quick-Set MMT-386	2
Octreotide (Sun) 177	Other Skin Preparations70	Paradigm Quick-Set MMT-387	2
Octreotide GH177	Ovestin	Paradigm Quick-Set MMT-396	2
Octreotide LAR (somatostatin	Genito-Urinary74	Paradigm Quick-Set MMT-397	2
analogue) 178	Hormone81	Paradigm Quick-Set MMT-398	2
Octreotide MaxRx177	Ox-Pam136	Paradigm Quick-Set MMT-399	2
Oestradiol80	Oxaliplatin159	Paradigm Silhouette MMT-368	2
Oestradiol valerate80	Oxaliplatin Accord159	Paradigm Silhouette MMT-377	2
Oestradiol with norethisterone80	Oxaliplatin Actavis 100159	Paradigm Silhouette MMT-378	2
Oestriol	Oxaliplatin Ebewe159	Paradigm Silhouette MMT-381	2
Genito-Urinary74	Oxazepam 136	Paradigm Silhouette MMT-382	2
Hormone81	Oxis Turbuhaler234	Paradigm Silhouette MMT-383	2
Oestrogens80	Oxpentifylline57	Paradigm Silhouette MMT-384	2
Ofev237	Oxybutynin	Paradigm Sure-T MMT-864	2
Oil in water emulsion66	Oxycodone hydrochloride124	Paradigm Sure-T MMT-866	2
Olanzapine 134–135	Oxycodone Sandoz124	Paradigm Sure-T MMT-874	2
Olaparib165	OxyNorm124	Paradigm Sure-T MMT-876	2
Olbetam54	Oxytocin75	Paradigm Sure-T MMT-884	
Olbetam S2954	Oxytocin BNM75	Paradigm Sure-T MMT-886	2
Olopatadine244	Oxytocin with ergometrine	Paraffin	
Olopatadine Teva244	maleate	Paraffin liquid with wool fat	.24
Olsalazine7	Ozurdex241	Paraldehyde	.12
Omalizumab205	-P-	Parasidose	6
Omeprazole9	Pacifen118	Parasiticidal Preparations	6
Omeprazole actavis 109	Paclitaxel166	Parnate	.12
Omeprazole actavis 209	Paclitaxel Actavis166	Parnate S29	.12
Omeprazole actavis 409	Paclitaxel Ebewe166	Paromomycin	9
Omnitrope82	Paediatric Seravit34	Paroxetine	.12
Onbrez Breezhaler234	Palbociclib173	Paser	9
Oncaspar LYO166	Paliperidone135	Patanol	.24
OncoTICE186	Pamidronate disodium113	Paxam	
Ondansetron132	Pamisol113	Pazopanib	. 17

Peak flow meter	239	Pioglitazone	12	Prevenar 13	279
Pedialyte - Bubblegum	46	Pirfenidone		Prezista	106
Pediasure	255	Pizotifen	131	Priadel	133
Pediasure RTH	255	PKU Anamix Infant	265	Primacin	
Pegaspargase		PKU Anamix Junior	265	Primaquine	
Pegasys	107	PKU Anamix Junior Chocolate	265	Primidone	129
Pegfilgrastim	45	PKU Anamix Junior LQ	265	Primolut N	81
Pegylated interferon alfa-2a	107	PKU Anamix Junior Vanilla	265	Priorix	
Pembrolizumab	229	PKU Lophlex LQ 10	265	Probenecid	117
Pemetrexed	161	PKU Lophlex LQ 20	265	Probenecid-AFT	
Penicillamine	111	PKU Lophlex Powder	265	Procaine penicillin	
Penicillin G	91	PKU Lophlex Sensation 20		Procarbazine hydrochloride	
PenMix 30		Plaquenil	111	Prochlorperazine	132
PenMix 40		Plendil ER		Proctofoam	
PenMix 50		Pneumococcal (PCV10) conjuga		Proctosedyl	
Pentasa		vaccine		Procyclidine hydrochloride	
Pentostatin [Deoxycoformycin]	166	Pneumococcal (PCV13) conjuga		Procytox	158
Pentoxifylline [Oxpentifylline]	57	vaccine	279	Progesterone	81
Peptamen Junior	255	Pneumococcal (PPV23)		Proglicem	
Peptisoothe		polysaccharide vaccine	280	Proglycem	10
Peptisorb	257	Pneumovax 23	280	Progynova	80
Perhexiline maleate		Podophyllotoxin	70	Prolia	112
Pericyazine		Polaramine	233	Promethazine hydrochloride	233
Perindopril		Poliomyelitis vaccine	280	Propafenone hydrochloride	49
Perjeta	207	Poloxamer	27	Propamidine isethionate	240
Permethrin		Poly-Gel		Propranolol	
Perrigo	70	Poly-Tears	244	Propylene glycol	
Pertuzumab	207	Poly-Visc	244	Propylthiouracil	82
Peteha	99	Polycal	250	Protaphane	11
Pethidine hydrochloride	125	Ponstan	110	Protaphane Penfill	11
Pevaryl	62	Posaconazole	96	Protifar	252
Pexsig	5 <mark>2</mark>	Postinor-1	74	Protionamide	
Pfizer Exemestane	179	Potassium chloride	45-46	Provera	80
Pheburane	31	Potassium Chloride Aguettant	45	Provera HD	81
Phenasen	162	Potassium citrate	75	PSM Citalopram	126
Phenelzine sulphate	126	Potassium iodate	35	Psoriasis and Eczema	
Phenobarbitone	128	Povidone iodine	66	Preparations	68
Phenobarbitone sodium		Pradaxa	44	PTU	
Extemporaneous	249	Pramipexole hydrochloride	119	Pulmicort Turbuhaler	233
Nervous	150	Prasugrel	42	Pulmocare	253
Phenothrin	68	Pravastatin	55	Pulmozyme	238
Phenoxybenzamine		Pravastatin Mylan	55	Puri-nethol	
hydrochloride	47	Praziquantel	88	Puria	34
Phenoxymethylpenicillin (Penicilli		Prazosin		Pyrazinamide	99
V)	92	Pred Forte		Pyridostigmine bromide	110
Phenytoin sodium1	27-128	Prednisolone	78	Pyridoxine hydrochloride	
Phlexy 10		Prednisolone acetate	242	Pyrimethamine	
Phosphate Phebra	46	Prednisolone sodium		Pytazen SR	42
Phosphorus		phosphate	242	- Q -	
Phytomenadione		Prednisolone-AFT		Q 300	98
Pilocarpine hydrochloride		Prednisone		Quetapel	
Pimafucort		Pregabalin	129	Quetiapine	
Pindolol		Pregabalin Pfizer	129	Quick-Set MMT-390	
Pine tar with trolamine laurilsulfat		Pregnancy Tests - hCG Urine		Quick-Set MMT-391	
and fluorescein		Premarin		Quick-Set MMT-392	
Pinetarsol	69	Presolol		Quick-Set MMT-393	2

Quinapril	47	Rituximab (Mabthera)	207	Serenace	13
Quinapril with		Rituximab (Riximyo)	212	Seretide	23
hydrochlorothiazide	48	Rivaroxaban	44	Seretide Accuhaler	23
Quinine sulphate		Rivastigmine	153	Serevent	23
Qvar		Rivotril		Serevent Accuhaler	
- R -		Riximyo	212	Sertraline	12
RA-Morph	123	RIXUBIS	41	Setrona	12
Raloxifene hydrochloride	113	Rizamelt	130	Sevredol	12
Raltegravir potassium		Rizatriptan	130	Sex Hormones Non	
Ramipex		Robinul		Contraceptive	<mark>7</mark>
Ranbaxy-Cefaclor		Roferon-A	107	Shield XL	
Ranitidine		Rolin	179	shingles vaccine	
Ranitidine Relief		Ropin	119	SII-Onco-BCG	18
Rapamune	231	Ropinirole hydrochloride	119	Sildenafil	5
Reandron 1000		Rotarix		Silhouette MMT-371	
Recombinant factor IX		Rotavirus oral vaccine		Silhouette MMT-373	
Recombinant factor VIIa	,	Roxane		Siltuximab	22
Recombinant factor VIII		Alimentary	6	Simvastatin	
Rectogesic		Cardiovascular		Simvastatin Mylan	5
Redipred		Roxithromycin		Sinemet	11
Relieve		Rubifen		Sinemet CR	
Relistor		Rubifen SR		Sirolimus	
Remicade		Rugby Capsaicin Topical Cre		Siterone	
Renilon 7.5		Musculoskeletal		Slow-Lopresor	
Resonium-A		Nervous		Smith BioMed Rapid Pregnancy	
Resource Beneprotein		Rulide D		Test	7
Resource Diabetic		Rurioctocog alfa pegol [Reco		Sodibic	
Respigen		factor VIII]		Sodium acid phosphate	
		•			
Respiratory Devices		Ruxolitinib		Sodium alginate Sodium benzoate	
Respiratory Stimulants		Rythmodan			3
Retinol palmitate		Rytmonorm	49	Sodium bicarbonate	4E 4
ReTrieve			400	Blood	
Retrovir		Sabril		Extemporaneous	
Revlimid		Sacubitril with valsartan		Sodium calcium edetate	24
Revolade		SalAir		Sodium chloride	
Rexacrom		Salazopyrin		Blood	
RexAir		Salazopyrin EN		Respiratory	23
Ribomustin		Salbutamol	235	Sodium citrate with sodium lauryl	
Ricit		Salbutamol with ipratropium		sulphoacetate	
Rifabutin		bromide		Sodium citro-tartrate	7
Rifadin		Salicylic acid		Sodium cromoglicate	
Rifampicin		Salmeterol		Alimentary	
Rifaximin	10	Sandomigran		Respiratory	
Rifinah		Sandostatin LAR		Sensory	
Rilutek		Sapropterin dihydrochloride.		Sodium fluoride	3
Riluzole	120	Scalp Preparations	70	Sodium Fusidate [fusidic acid]	
Riodine		Scopoderm TTS	131	Dermatological	6
Risedronate Sandoz	113	Sebizole	70	Infection	9
Risedronate sodium	113	Secukinumab	221	Sensory	24
Risperdal Consta	135	Sedatives and Hypnotics	149	Sodium hyaluronate [Hyaluronic	
Risperidone		Seebri Breezhaler	236	acid]	24
Risperon		Selegiline hydrochloride	119	Sodium phenylbutyrate	
Ritalin		Senna		Sodium polystyrene sulphonate	
Ritalin LA	152	Senokot	<mark>28</mark>	Sodium tetradecyl sulphate	
Ritalin SR		Sensipar		Sodium valproate	12
Ritonavir		SensoCard		Sofradex	

Soframycin	240	Sympathomimetics	56	Thiotepa	159
Solifenacin Mylan		Synacthen	79	Thymol glycerin	3
Solifenacin succinate	76	Synacthen Depot	79	Thyroid and Antithyroid Agents	8
Solu-Cortef	78	Synacthene Retard		Ticagrelor	4
Solu-Medrol	78	Synflorix		Tilade	
Solu-Medrol-Act-O-Vial	78	Synthroid		Tilcotil	
Somatropin (Omnitrope)	82	Syntometrine	75	Tillomed	15
Sotalol		Syrup (pharmaceutical grade)		Timolol	
Spacer device	239	Systane Unit Dose		Cardiovascular	5
Span-K		· -T-		Sensory	24
Spiolto Respimat		Tacrolimus	231	Timoptol XE	
Spiractin		Tacrolimus Sandoz	231	Tiotropium bromide	
Spiriva		Taliglucerase alfa	31	Tiotropium bromide with	
Spiriva Respimat		Tambocor		olodaterol	23
Spironolactone		Tamoxifen citrate	179	Tivicay	
Sporanox		Tamoxifen Sandoz	179	TMP	
Sprycel		Tamsulosin hydrochloride		TOBI	
Staphlex		Tamsulosin-Rex		Tobramycin	
Stemetil		Tandem Cartridge		Infection	9
SteroClear		Tandem t:slim X2		Sensory	
Stesolid		Tap water		Tobramycin Mylan	
Stimulants/ADHD Treatments .		Tarceva		Tobrex	
Stiripentol		Tasigna		Tocilizumab	
Stocrin		Tasmar		Tofranil	
Stomahesive		Tecfidera		Tolcapone	
Strattera	150	Tegretol		Topamax	
Strides Shasun		Tegretol CR		Topical Products for Joint and	
Stromectol		Telfast		Muscular Pain	11
Sucralfate		Teligent		Topiramate	
Sulfadiazine Silver		Temaccord		Topiramate Actavis	
Sulfadiazine sodium		Temazepam		Total parenteral nutrition (TPN)	
Sulfasalazine		Temozolomide		TPN	
Sulindac		Tenofovir disoproxil		Tramadol hydrochloride	
Sulindac Mylan		Tenofovir Disoproxil Teva		Tramal SR 100	
Sulphur		Tenoxicam		Tramal SR 150	
Sulprix		Tepadina		Tramal SR 200	
Sumatriptan		Terazosin		Trandate	
Sunitinib		Terbinafine		Tranexamic acid	
Sunscreens		Terbutaline sulphate		Tranylcypromine sulphate	
Sunscreens, proprietary		Teriflunomide		Trastuzumab	22
Sure-T MMT-863		Teriparatide		Trastuzumab emtansine	22
Sure-T MMT-865		Testosterone		Travatan	
Sure-T MMT-873		Testosterone cipionate		Travoprost	
Sure-T MMT-875		Testosterone esters		Travopt	
Sure-T MMT-883		Testosterone undecanoate		Treatments for Dementia	
Sure-T MMT-885		Tetrabenazine		Treatments for Substance	
Sustagen Diabetic		Tetrabromophenol		Dependence	15
Sustagen Hospital Formula		Tetracosactrin		Trental 400	
Active		Tetracyclin Wolff		Tretinoin	
Sustanon Ampoules		Tetracycline	92	Dermatological	6
Sutent		Thalidomide		Oncology	16
Sylvant		Thalomid		Trexate	
Symbicort Turbuhaler 100/6		Theophylline		Triamcinolone acetonide	
Symbicort Turbuhaler 200/6		Thiamine hydrochloride		Alimentary	3:
Symbicort Turbuhaler 400/12		THIO-TEPA	159	Dermatological	6
Symmetrel		Thioguanine		Hormone	70
~j					, ,

Triamcinolone acetonide with	Varivax	281	Wool fat with mineral oil	66
gramicidin, neomycin and nystatin	Vasodilators	<u>56</u>	- X -	
Dermatological65	Vasopressin Agonists		Xarelto	44
Sensory240	Vedafil		Xifaxan	10
Triaver	Veletri		XMET Maxamum	
Triazolam150	Venclexta		Xolair	
Trichozole98	Venetoclax		XP Maxamum	
Triclosan	Venlafaxine		Xylocaine	
Trimethoprim95	Venomil		Xylocaine 2% Jelly	
Trimethoprim with	Ventavis		Xyntha	
sulphamethoxazole	Ventolin		- Z -	41
[Co-trimoxazole]95	Ventolin		Zantac	
			Zapril	
Trisequens80	Verapamil hydrochloride			
Trisul	Vergo 16		Zarontin	
Trophic Hormones	Vermox		Zaroxolyn	
Tropicamide243	Verpamil SR		Zavedos	
Trusopt242	Vesanoid		Zeffix	
TruSteel21	Vexazone		Zetlam	
Tuberculin PPD [Mantoux] test281	Vfend		Ziagen	
Tubersol281	Viaderm KC		Zidovudine [AZT]	106
Two Cal HN262	Vidaza		Zidovudine [AZT] with	
Two Cal HN RTH262	Vigabatrin		lamivudine	
Tykerb172	Vildagliptin	12	Zimybe	55
Tysabri139	Vildagliptin with metformin		Zinc and castor oil	
- U -	hydrochloride	12	Zinc sulphate	
Ultibro Breezhaler236	Vimpat		Zincaps	37
Ultraproct8	Vinblastine sulphate	169	Zinnat	8
Umeclidinium236	Vincristine sulphate	169	Ziprasidone	134
Umeclidinium with vilanterol237	Vinorelbine		Zista	233
Univent235, 239	Vinorelbine Ebewe	169	Zithromax	
Ural	Viramune Suspension		Zoladex	86
Urea66	ViruPOS		Zoledronic acid	
Urex Forte53	Vit.D3		Hormone	77
Urinary Agents75	VitA-POS		Musculoskeletal	
Urinary Tract Infections109	Vitabdeck		Zoledronic acid Mylan	
Urinorm	Vital		Zopiclone	
Uromitexan	Vitamin B complex		Zopiclone Actavis	150
Ursodeoxycholic acid26	Vitamin B6 25		Zostavax	281
Ursosan	Vitamins		Zostrix	
Utrogestan81	Vivonex TEN		Zostrix HP	
- V -	Volibris		Zuclopenthixol decanoate	
Vaccinations270	Voltaren			
	Voltaren D		Zuclopenthixol hydrochloride Zusdone	
Vaclovir101				
	Voltaren Ophtha		Zyban	
Valganciclovir101	Volumatic		Zypine	134
Valganciclovir Mylan101	Voriconazole		Zypine ODT	134
Vancomycin95	Vosol		Zyprexa Relprevv	
Vannair234	Votrient		Zytiga	176
Varenicline Pfizer156	Vttack	97		
Varenicline tartrate156	- W -			
Varicella vaccine [Chickenpox	Warfarin sodium			
vaccine] 281	Wart Preparations			
Varicella zoster virus (Oka strain) live	Wasp venom allergy treatment	t <mark>232</mark>		
attenuated vaccine [shingles	Water			
vaccine] 281	Blood	46		
Varilrix281	Extemporaneous	249		