September 2020 Volume 27 Number 2

Editors:

Kaye Wilson & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

Published each April, August and December. Changes to the contents are published in monthly updates.

Accessible in an electronic format at no cost from the PHARMAC website www.pharmac.govt.nz/schedule.

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at www.pharmac.govt.nz/subscriptions.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth & John Geering email: texschedule@pharmac.govt.nz @Pharmaceutical Management Agency



ISSN 1179-3686 pdf ISSN 1172-9376 print

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to PHARMAC and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to PHARMAC should be in written form and not by reproduction of the PHARMAC logo. While care has been taken in compiling this Schedule, PHARMAC takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

	introducing i Harimao	_
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	37
	Cardiovascular System	47
	Dermatologicals	61
	Genito Urinary System	71
	Hormone Preparations – Systemic	77
	Infections – Agents For Systemic Use	88
	Musculoskeletal System	110
	Nervous System	119
	Oncology Agents & Immunosuppressants	158
	Respiratory System & Allergies	233
	Sensory Organs	241
	Various	246
Section C	Extemporaneous Compounds (ECPs)	248
Section D	Special Foods	251
Section I	National Immunisation Schedule	272

Introducing PHARMAC

Index

Introducing PHARMAC

The Pharmaceutical Management Agency (PHARMAC) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. PHARMAC negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

PHARMAC's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about PHARMAC and the way we make funding decisions can be found on the PHARMAC website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by PHARMAC.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements PHARMAC has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg p sachet		30	•	Gaviscon Infant
SODIUM ALGINATE Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (8.60)	60		Gaviscon Double Strength
Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 m	-	Acidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE * Tab 600 mg	12.56	100	✓.	Alu-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsementOnly when prescribed for patients unable to swallow cal inappropriate and the prescription is endorsed according	cium carbonate tablet	500 m s or v		Roxane um carbonate tablets are
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on * Tab 2 mg* * Cap 2 mg	10.75	400 400		Nodia Diamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE Cap 3 mg - Special Authority see SA1886 below - Retail pharmacy	166.50	90	✓	Entocort CIR
⇒SA1886 Special Authority for Subsidy Initial application — (Crohn's disease) from any relevant practithe following criteria: Both:	titioner. Approvals va	ılid fo	r 6 months	for applications meeting
Mild to moderate ileal, ileocaecal or proximal Crohn's disc	ease; and			

2.3 Osteoporosis where there is significant risk of fracture; or

2 Any of the following: 2.1 Diabetes; or 2.2 Cushingoid habitus; or

Subs	sidy Full	/ Brand or
(Manufactur	rer's Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*: and
- 2 Patient does not have cirrhosis: and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

TITOTIO CONTINUINE ACETATE		
Rectal foam 10%, CFC-Free (14 applications)26.55	21.1 g OP	✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam S29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab EC 500 mg49.50	100	✓ Asamax
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
Modified release granules, 1 g141.72	120 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g54.60	30	✓ Pentasa
OLSALAZINE		
Tab 500 mg93.37	100	✓ Dipentum
Cap 250 mg53.00	100	✓ Dipentum

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
SODIUM CROMOGLICATE Cap 100 mg	92.91	100	✓ N	alcrom
SULFASALAZINE * Tab 500 mg Tab EC 500 mg		100 100		alazopyrin alazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CI	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cinchocaine hydrochloride 5 mg per g6.35	30 g OP	✓ Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and cinchocaine hydrochloride 1 mg2.66	12	✓ Ultraproct
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	 Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl

Management of Anal Fissures

GL	YCERYL TRINITRATE - Special Authority see SA1329 below - Retail pha	rmacy	
*	Oint 0.2%22.00	30 g OP	✓ Rectogesic

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule - Up to 10 inj available on	а		
PSO	17.14	10	Max Health
	34.32	5	✓ Robinul
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg	6.35	100	Buscopan
Buscopan to be Sole Supply on 1 October 2020			
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO	6.35	5	✓ Buscopan
MEBEVERINE HYDROCHLORIDE			
* Tab 135 mg	9.20	90	✓ Colofac
	9.20	90	✓ Colofac

Antiulcerants

Antisecretory and Cytoprotective

MISOPROSTOL

Subsidised on a PSO only if from a Family Planning New Zealand Clinic or an abortion service provider with a DHB contract and the PSO is endorsed with the name of the institution for which the PSO is required.

*	Tab 200 mcg - l	Up to 120 tab available on a F	PSO41.50	120	Cytotec
---	-----------------	--------------------------------	----------	-----	---------

Subsidy		Fully	Brand or	Ī
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	1	Manufacturer	

Helicobacter Pylori Eradication

CLARITHROMYCIN

- a) Maximum of 14 tab per prescription
- Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.
 Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

H2 Antagonists

FAI	MOTIDINE - Only on a prescription		
*	Tab 20 mg4.91	100	✓ Famotidine
	•		Hovid S29
*	Tab 40 mg8.48	100	✓ Famotidine
			Hovid S20

RANITIDINE - Subsidy by endorsement

- a) Only on a prescription
- b) Subsidy by endorsement Subsidised for patients who were taking ranitidine prior to 1 November 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of ranitidine.

*	Tab 150 mg	12.91	500	✓ Ranitidine Relief
	Tab 300 mg		500	Ranitidine Relief
*	Oral liq 150 mg per 10 ml	5.14	300 ml	✓ Peptisoothe
*	Inj 25 mg per ml, 2 ml	13.40	5	✓ Zantac
(Ra	anitidine Relief Tab 150 mg to be delisted 1 October 2020)			
(Ra	anitidine Relief Tab 300 mg to be delisted 1 October 2020)			
172	antac Ini 25 mg per ml. 2 ml to be delisted 1 June 2021)			

Proton Pump Inhibitors

	•			
LA *	NSOPRAZOLE Cap 15 mg		100 100	✓ Lanzol Relief ✓ Lanzol Relief
ON	IEPRAZOLE			
	For omeprazole suspension refer Standard Formulae, page 248			
*	Cap 10 mg	.98	90	✓ Omeprazole actavis 10
*	Cap 20 mg	.96	90	✓ Omeprazole actavis 20
*	Cap 40 mg	.12	90	✓ Omeprazole actavis 40
*	Powder – Only in combination42	.50	5 g	✓ Midwest
	Only in extemporaneously compounded omeprazole suspension.		•	
*	Inj 40 mg ampoule with diluent	.98	5	✓ <u>Dr Reddy's</u> <u>Omeprazole</u> ✓ Ocicure S29
				Ocicule 529
PA	NTOPRAZOLE			_
*	Tab EC 20 mg2		100	✓ Panzop Relief
*	Tab EC 40 mg2	.85	100	✓ Panzop Relief

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE Tab 120 mgSUCRALFATE	14.51	50	√ (Gastrodenol S29
Tab 1 g	35.50 (48.28)	120	(Carafate
Bile and Liver Therapy				
RIFAXIMIN - Special Authority see SA1461 below - Retail Tab 550 mg	,	56	✓)	(ifaxan
⇒SA1461 Special Authority for Subsidy nitial application only from a gastroenterologist, hepatolog nepatologist. Approvals valid for 6 months where the patient olerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or Practice approvals valid without further renewal unless penefiting from treatment.	t has hepatic encephalor	pathy d	lespite an a	dequate trial of maximum penterologist or
Diabetes				
Hyperglycaemic Agents				
DIAZOXIDE – Special Authority see SA1320 below – Retail Cap 25 mg Cap 100 mg Oral liq 50 mg per ml ➤ SA1320 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals mypoglycaemia caused by hyperinsulinism. Renewal from any relevant practitioner. Approvals valid with			P F	
appropriate and the patient is benefiting from treatment. GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit — Up to 5 kit available on a PSO	32.00	1	√ <u>(</u>	Glucagen Hypokit
Insulin - Short-acting Preparations				
NSULIN NEUTRAL ▲ Inj human 100 u per ml	25.26	10 ml C		Actrapid Humulin R
▲ Inj human 100 u per ml, 3 ml	42.66	5		Actrapid Penfill Iumulin R
Insulin - Intermediate-acting Preparations				
NSULIN ASPART WITH INSULIN ASPART PROTAMINE Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	✓ N	NovoMix 30 FlexPen

	Subsidy	F	ully Brand or
	(Manufacturer's Price	e) Subsidis	,
	\$	Per	✓ Manufacturer
NSULIN ISOPHANE			
Inj human 100 u per ml	17.68		✓ Humulin NPH✓ Protaphane
Inj human 100 u per ml, 3 ml	29.86	5	✓ Humulin NPH
- , ,			✓ Protaphane Penfill
NSULIN ISOPHANE WITH INSULIN NEUTRAL			
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70✓ Mixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70
•			✓ PenMix 30
			✓ PenMix 40
JOHN N. JORDO WITH INCH IN JORDO DROTAMINE			✓ PenMix 50
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE Inj lispro 25% with insulin lispro protamine 75% 100 u per ml.			
3 ml		5	✓ Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml			
3 ml		5	✓ Humalog Mix 50
Insulin - Long-acting Preparations			
SULIN GLARGINE			
Inj 100 u per ml, 10 ml	63.00	1	✓ Lantus
Inj 100 u per ml, 3 ml		5	✓ Lantus
Inj 100 u per ml, 3 ml disposable pen		5	✓ Lantus SoloStar
Insulin - Rapid Acting Preparations			
NSULIN ASPART			
Inj 100 u per ml, 10 ml			✓ NovoRapid
Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe	51.19	5	✓ NovoRapid FlexPen
NSULIN GLULISINE Inj 100 u per ml, 10 ml	27.02	1	✓ Apidra
Inj 100 u per ml, 3 ml			✓ Apidra
Inj 100 u per ml, 3 ml disposable pen			✓ Apidra SoloStar
NSULIN LISPRO			
Inj 100 u per ml, 10 ml		10 ml OP	✓ Humalog
Inj 100 u per ml, 3 ml	59.52	5	✓ Humalog
Alpha Glucosidase Inhibitors			
CARBOSE			
★ Tab 50 mg		90	✓ Glucobay
₭ Tab 100 mg	10.47 6.40	90	✓ Accarb✓ Glucobay
- 145 155 Hig	20.23	00	✓ Accarb
Oral Hypoglycaemic Agents			
ilibenclamide			
₭ Tab 5 mg	6.00	100	✓ <u>Daonil</u>
			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
GLICLAZIDE				
* Tab 80 mg	15.18	500	•	Glizide
GLIPIZIDE				
* Tab 5 mg	3.27	100	✓	<u>Minidiab</u>
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg	8.63	1,000	/	Apotex
* Tab immediate-release 850 mg	7.04	500	•	Apotex
PIOGLITAZONE				
* Tab 15 mg	3.47	90	1	Vexazone
* Tab 30 mg		90	1	Vexazone
* Tab 45 mg	7.10	90	✓	<u>Vexazone</u>
VILDAGLIPTIN				
Tab 50 mg	40.00	60	✓	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride	40.00	60	1	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride		60	1	Galvumet

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes: or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	✓ CareSens N
		✓ CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood g	lucose test stri	os26.20	50 test OP	✓ SensoCard
---------	------------------	---------	------------	-------------

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

IIV	DOLIN I LIN NEEDELO — Maximum di 200 dev per prescrip	MON		
*	29 g × 12.7 mm	10.50	100	✓ B-D Micro-Fine
*	31 g × 5 mm	11.75	100	✓ B-D Micro-Fine
*	31 g × 6 mm	9.50	100	✓ Berpu
*	31 g × 8 mm		100	✓ B-D Micro-Fine
*	32 g × 4 mm		100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEED		200 dev per p	prescription
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.00	100	B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g × 8 mm needle	13.00	100	✓ B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.00	100	✓ B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	13.00	100	✓ B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.00	100	✓ B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 1 ml with 31 g x 8 mm needle	13.00	100	✓ B-D Ultra Fine II
	-	1.30	10	
		(1.99)		B-D Ultra Fine II
		, ,		

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

 c) Maximum of 1 insulin pump per patient each four year 	period.		
Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 640G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem Basal IQ
			✓ Tandem t:slim X2

(Tandem t:slim X2 Min basal rate 0.1 U/h to be delisted 1 December 2020)

⇒SA1603 Special Authority for Subsidy

Initial application — **(permanent neonatal diabetes)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and

	Subsidy	F	ully	Brand or
(Ma	nufacturer's Price)	Subsidi	sed	Generic
·	\$	Per	✓	Manufacturer

continued...

- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Fither:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	•	Manufacturer	

continued...

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Fither:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Either:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Insulin Pump Consumables

⇒SA1906 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:

			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per ✓	Manufacturer

continued...

- 8.1 Applicant is a relevant specialist; or
- 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline, according to the most recent result.

Initial application — **(HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol, according to the most recent result; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline.

Initial application — (**Previous use before 1 September 2012**) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) from any relevant practitioner. Approvals valid for 2 years for

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	
----------------------------------------------------------------------------------------	--

continued...

applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol according to a recent laboratory result; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application, according to the most recent result; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline.

INSULIN PUMP CARTRIDGE - Special Authority see SA1906 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 packs of cartridge sets will be funded per year.

Cartridge 300 U, t:lock × 10......50.00 1 OP ✓ Tandem Cartridge

	Subsidy	Fully	Brand or
(Man	ufacturer's Price)	Subsidised	Generic
·	\$ P	er 🗸	Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA) - Special Authority see SA1906 on page 17 - Retail pharmacy

a)	Maximum	of 3 sets	per pre	scription
----	---------	-----------	---------	-----------

a) Maximum of 3 sets per prescriptionb) Only on a prescriptionc) Maximum of 13 infusion sets will be funded per year.			
10 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Sure-T MMT-884A
10 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Sure-T MMT-886A
6 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Sure-T MMT-864A
6 mm steel needle; 80 cm tubing x 10	130.00	1 OP	✓ MiniMed Sure-T MMT-866A
8 mm steel needle; 60 cm tubing x 10	130.00	1 OP	✓ MiniMed Sure-T MMT-874A
8 mm steel needle; 80 cm tubing x 10	130.00	1 OP	✓ MiniMed Sure-T MMT-876A
10 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T
10 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T
6 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles	130.00	1 OP	MMT-886 ✓ Paradigm Sure-T
6 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock	120.00	1 OP	MMT-864 ✓ Sure-T MMT-863
6 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles		1 OP	✓ Paradigm Sure-T
8 mm steel needle; 29 G; manual insertion; 60 cm tubing x	400.00	4.00	MMT-866
10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T MMT-874
8 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-873
8 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T

MMT-876

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) – Special Authority see SA1906 on page 17 – Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles	1 OP	✓ TruSteel
6 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles130.00) 1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles130.00	0 1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles130.00	1 OP	✓ TruSteel

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1906 on page 17 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription

c) Maximum of 13 infusion sets will be funded per ye	ear.		
13 mm teflon needle, 110 cm tubing × 10		1 OP	✓ MiniMed Silhouette MMT-382A
13 mm teflon needle, 45 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-368A
13 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-381A
13 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-383A
17 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-377A
17 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-378A
17 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-384A
6 mm teflon needle, 110 cm tubing x 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-398A
6 mm teflon needle, 45 cm blue tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-941A
6 mm teflon needle, 45 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-921A
6 mm teflon needle, 60 cm blue tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-943A
6 mm teflon needle, 60 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-923A
6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-399A
6 mm teflon needle, 80 cm blue tubing	130.00	1 OP	✓ MiniMed Mio MMT-945A
6 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-965A
6 mm teflon needle, 80 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-925A
6 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-387A
9 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-396A
9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-397A
9 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-975A
9 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-386A

see

	Subsidy (Manufacturer's Pr	ice) Sub	Fully osidised	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE I SA1906 on page 17 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year.	NSERTION WITH	INSERTION	N DEVICE) - Special Authority
13 mm teflon cannula; angle insertion; insertion device; 110 line × 10 with 10 needles	140.00	1 OP	✓ Aι	utoSoft 30
line × 10 with 10 needles		1 OP	✓ Au	utoSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE I Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 13 mm teflon cannula; angle insertion; 120 cm line × 10 with 10 needles		pecial Author	✓ Pa	aradigm Silhouette
13 mm teflon cannula; angle insertion; 45 cm line × 10 with 10 needles	130.00	1 OP	✓ Pa	MMT-382 rradigm Silhouette MMT-368
13 mm teflon cannula; angle insertion; 60 cm line × 10 with 10 needles	130.00	1 OP		radigm Silhouette MMT-381
13 mm teflon cannula; angle insertion; 80 cm line × 10 with 10 needles	130.00	1 OP		radigm Silhouette MMT-383
17 mm teflon cannula; angle insertion; 110 cm line × 10 with 10 needles		1 OP		radigm Silhouette MMT-377
17 mm teflon cannula; angle insertion; 60 cm line x 10 with 10 needles	130.00	1 OP		radigm Silhouette MMT-378
17 mm teflon cannula; angle insertion; 60 cm line × 10 with 10 needles; luer lock	130.00	1 OP	✓ Si	Ihouette MMT-373
17 mm teflon cannula; angle insertion; 80 cm line × 10 with				

✓ Paradigm Silhouette MMT-384

1 OP

6 mm teflon cannula: straight insertion: insertion device: 60 cm

9 mm teflon cannula: straight insertion: insertion device: 60 cm

9 mm teflon cannula; straight insertion; insertion device;

line × 10 with 10 needles......140.00

110 cm line × 10 with 10 needles140.00

line × 10 with 10 needles......140.00

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Por	/	Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION WITH INSERTION DEVICE) - Special Authority see SA1906 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription

c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; insertion device; 45 cm			
blue tubing × 10 with 10 needles130.0	00 10	P ✓ Paradigm Mi MMT-941	io
6 mm teflon cannula; straight insertion; insertion device; 45 cm pink tubing × 10 with 10 needles	00 10	P Paradigm Mi	io
6 mm teflon cannula; straight insertion; insertion device; 60 cm blue tubing × 10 with 10 needles130.0	00 10		io
6 mm teflon cannula; straight insertion; insertion device; 60 cm pink tubing × 10 with 10 needles130.0	00 10	P ✓ Paradigm Mi MMT-923	io
6 mm teflon cannula; straight insertion; insertion device; 80 cm blue tubing × 10 with 10 needles130.0	00 10	P ✓ Paradigm Mi MMT-945	io
6 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing x 10 with 10 needles130.0	00 1 0	P ✓ Paradigm Mi MMT-965	io
6 mm teflon cannula; straight insertion; insertion device; 80 cm pink tubing × 10 with 10 needles130.0	00 1 0	P ✓ Paradigm Mi MMT-925	io
9 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing × 10 with 10 needles130.0	00 1 0	P Paradigm Mi MMT-975	io
6 mm teflon cannula; straight insertion; insertion device; 110 cm line × 10 with 10 needles140.0	00 10		

✓ AutoSoft 90

✓ AutoSoft 90

✓ AutoSoft 90

1 OP

1 OP

1 OP

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Manufacturer INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION) - Special Authority see SA1906 on page 17 -Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula: straight insertion: 110 cm tubing × 10 with 1 OP ✓ Paradigm Quick-Set MMT-398 6 mm teflon cannula: straight insertion: 60 cm tubing × 10 with 1 OP ✓ Paradigm Quick-Set MMT-399 6 mm teflon cannula: straight insertion: 60 cm tubing × 10 with 1 OP ✓ Quick-Set MMT-393 6 mm teflon cannula; straight insertion; 80 cm tubing x 10 with 1 OP ✓ Paradigm Quick-Set MMT-387 9 mm teflon cannula; straight insertion; 106 cm tubing × 10 with 1 OP ✓ Paradigm Quick-Set MMT-396 9 mm teflon cannula: straight insertion: 60 cm tubing × 10 with 1 OP ✓ Paradigm Quick-Set MMT-397 9 mm teflon cannula: straight insertion: 60 cm tubing × 10 with 1 OP ✓ Quick-Set MMT-392 9 mm teflon cannula; straight insertion; 80 cm tubing \times 10 with 1 OP ✓ Paradigm Quick-Set MMT-386 INSULIN PUMP RESERVOIR - Special Authority see SA1906 on page 17 - Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of reservoir sets will be funded per year. 10 × luer lock conversion cartridges 1.8 ml for Paradigm pumps......50.00 ✓ ADR Cartridge 1.8 1 OP Cartridge for 5 and 7 series pump; 1.8 ml × 1050.00 1 OP ✓ Paradigm 1.8 Reservoir Cartridge for 7 series pump: 3.0 ml × 1050.00 ✓ Paradigm 1 OP 3.0 Reservoir **Digestives Including Enzymes** PANCREATIC FNZYME Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)34.93 100 Creon 10000 Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase, 100 **Panzytrat** Cap pancreatin 300 mg (amylase 18,000 Ph Eur U. lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)94.38 100 Creon 25000 Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph 20 q OP ✓ Creon Micro

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
URSODEOXYCHOLIC ACID - Special Authority see SA1739 be	low – Retail pharmac	у			
Cap 250 mg	32.95	100	✓ (Jrsosan	

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Either:
 - 1 Patient has been diagnosed with Alagille syndrome; or

Ursosan to be Sole Supply on 1 October 2020

2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (**Pregnancy**) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (**Pregnancy/Primary biliary cholangitis**) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Fully

Brand or

Subsidy

	Subsidy (Manufacturer's F	Price) Subs	Fully Brand or idised Generic
Laxatives	\$	rei	✓ Manufacturer
Bulk-forming Agents			
SPAGHULA (PSYLLIUM) HUSK – Only on a prescription * Powder for oral soln Konsyl-D to be Sole Supply on 1 November 2020 MUCILAGINOUS LAXATIVES WITH STIMULANTS	12.20	500 g OP	✓ Konsyl-D
* Dry	6.02 (17.32) 2.41	500 g OP 200 g OP	Normacol Plus
	(8.72)	-	Normacol Plus
Faecal Softeners			
DOCUSATE SODIUM – Only on a prescription * Tab 50 mg Coloxyl to be Sole Supply on 1 October 2020	2.31	100	✓ Coloxyl
* Tab 120 mg Coloxyl to be Sole Supply on 1 October 2020 DOCUSATE SODIUM WITH SENNOSIDES	3.13	100	✓ Coloxyl
* Tab 50 mg with sennosides 8 mg POLOXAMER – Only on a prescription	3.10	200	✓ <u>Laxsol</u>
Not funded for use in the ear. ** Oral drops 10% Coloxyl to be Sole Supply on 1 November 2020	3.98	30 ml OP	✓ Coloxyl
Opioid Receptor Antagonists - Peripheral			
METHYLNALTREXONE BROMIDE - Special Authority see S Inj 12 mg per 0.6 ml vial		tail pharmacy 1 7	✓ Relistor ✓ Relistor
■ SA1691 Special Authority for Subsidy Initial application — (Opioid induced constipation) from ar unless notified for applications meeting the following criteria: Both: 1 The patient is receiving palliative care; and 2 Either: 2.1 Oral and rectal treatments for opioid induced con 2.2 Oral and rectal treatments for opioid induced con	nstipation are ineff	ective; or	
Osmotic Laxatives			
GLYCEROL * Suppos 3.6 g – Only on a prescription LACTULOSE – Only on a prescription	9.25	20	✓ <u>PSM</u>
* Oral liq 10 g per 15 ml	BICARBONATE A	500 ml ND SODIUM C	✓ <u>Laevolac</u> HLORIDE
Powder for oral soln 13.125 g with potassium chloride 46.6 sodium bicarbonate 178.5 mg and sodium chloride 35		30	✓ Molaxole

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Molaxole to be Sole Supply on 1 October 2020

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%	2.50	1	✓ Fleet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml	, , ,	otion 50	✓ <u>Micolette</u>
Stimulant Laxatives			
BISACODYL – Only on a prescription * Tab 5 mg * Suppos 10 mg		200 10	✓ <u>Lax-Tab</u> ✓ <u>Lax-Suppositories</u>
SENNA – Only on a prescription * Tab, standardised	2.17 (8.21) 0.43 (2.06)	100 20	Senokot Senokot

Metabolic Disorder Agents

ALGLUCOSIDASE ALFA - Special Authority see SA1920 be	elow - Retail pharmacy		
Inj 50 mg vial	1,142.60	1	✓ Myozyme

⇒SA1920 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidise	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to FRT: and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

⇒SA1921 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria: and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.
 Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE – Special Authority see SA1922 below – Retail pharmacy
Inj 1 mg per ml, 5 ml vial.......2,234.00 1 ✓ Naglazyme

⇒SA1922 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts: or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
- 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

SAPROPTERIN DIHYDROCHLORIDE – Special Authority see SA1923 below – Retail pharmacy
Tab soluble 100 mg.......1,452.70 30 OP
✓ Kuvan

⇒SA1923 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	•	Manutacturer	

continued...

- 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
- 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
- 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE - Special Authority see SA1599 below - Retail pharmacy

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1924 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

⇒SA1880 Special Authority for Subsidy

Special Authority approved by the Gaucher Treatment Panel

Notes: Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The Co-ordinator, Gaucher Treatment Panel Phone: 04 460 4990 PHARMAC PO Box 10 254 Facsimile: 04 916 7571

Wellington Email: gaucherpanel@pharmac.govt.nz

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Access Criteria

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific
 deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2) Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3) Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
- 1) Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia, 6) thrombocytopenia, at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or massive symptomatic splenomegaly; or
 - 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g., hips or shoulder); spontaneous fractures or vertebral collapse; chronic bone pain not controlled by other pharmaceuticals; or
 - 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 4) Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher
 - 5) Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1) Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated: and
- 2) Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3) Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and three yearly thereafter. demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4) Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 5) Patient has not developed another medical condition that might reasonably be expected to compromise a response to
- 6) Patient is compliant with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- 7) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations are submitted to the Gaucher Panel for assessment as required.

Mouth and Throat

Agents Used in Mouth Ulceration

BENZYDAMINE HYDROCHLORIDE

Soln 0.15% - Higher subsidy of \$20.31 per 500 ml with Endorsement9.00 500 ml (20.31)

Additional subsidy by endorsement for a patient who has oral mucositis as a result of treatment for cancer, and the prescription is endorsed accordingly.

Difflam

	Subsidy (Manufacturer's Pr	ice) Subs	Fully sidised	
CARMELLOSE SODIUM WITH GELATIN AND PECTIN				
Paste	17.20	56 g OP	1	Stomahesive
	4.55	15 g OP		
	(7.90)	10 9 01		Orabase
	1.52	5 g OP		Olababo
	(3.60)	3 g Oi		Orabase
Powder		28 g OP		Olabase
i owdei	(10.95)	20 g Oi		Stomahesive
	(10.55)			Otomanesive
CHLORHEXIDINE GLUCONATE			_	
Mouthwash 0.2%	2.57	200 ml OP	•	healthE
(healthE Mouthwash 0.2% to be delisted 1 November 2020)				
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE				
* Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP		
-	(6.00)	ŭ		Bonjela
TRIAMCINOLONE ACETONIDE	\/			•
Paste 0.1%	E 22	5 g OP	ſ	Kenalog in Orabase
Kenalog in Orabase to be Sole Supply on 1 November		o y Or	•	renalog III Olabase
Tronalog in Crabase to be sole supply on 1 November				
Oropharyngeal Anti-infectives				
AMPHOTERICIN B				
Lozenges 10 mg	5.86	20	1	Fungilin
MICONAZOLE				. 3
	4.74	40 = OD		Danasal
Oral gel 20 mg per g	4./4	40 g OP	•	<u>Decozol</u>
NYSTATIN				
Oral liq 100,000 u per ml	1.76	24 ml OP	/	Nilstat
Nilstat to be Sole Supply on 1 October 2020				
Other Oral Agents				
For folinic mouthwash, pilocarpine oral liquid or saliva substitute	formula refer Stan	idard Formula	ie. pai	ge 248
THYMOL GLYCERIN				
* Compound, BPC	0.15	500 ml	ſ	PSM
* Compound, Bro	9.10	300 IIII		r 3 Wi
Vitamins				
Vitamin B				
IVDDOVOCODAL AMINI				
HYDROXOCOBALAMIN ** Ini 1 mg nor ml 1 ml amnoula . I In to 6 ini gyailable on a R)CO 100	2	.1	Non P12
* Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a P	1.09	3	•	Neo-B12
PYRIDOXINE HYDROCHLORIDE				
a) No more than 100 mg per dose				
b) Only on a prescription				
* Tab 25 mg - No patient co-payment payable	2.70	90	1	Vitamin B6 25
Vitamin B6 25 to be Sole Supply on 1 October 2020				
* Tab 50 mg	13.63	500	1	Apo-Pyridoxine
				,
THIAMINE HYDROCHLORIDE – Only on a prescription	4.00	100	,	May Haald
en lan bul Ma	4 89	100	•	Max Health
* Tab 50 mg				
VITAMIN B COMPLEX * Tab, strong, BPC				

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

Vitamin C

ASCORBIC ACID

- a) No more than 100 mg per dose
- b) Only on a prescription

*	Tab 100 mg9.90	500	Cvite	е

Vitamin D

ALFACALCIDOL * Cap 0.25 mcg 26.32 * Cap 1 mcg 87.98 * Oral drops 2 mcg per ml 60.68	100 100 20 ml OP	✓ One-Alpha✓ One-Alpha✓ One-Alpha
CALCITRIOL * Cap 0.25 mcg	100 100	✓ Calcitriol-AFT ✓ Calcitriol-AFT
COLECALCIFEROL * Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescription2.50 * Oral liq 188 mcg per ml (7,500 iu per ml)	12 4.8 ml OP	✓ Vit.D3✓ Puria

Multivitamin Preparations

MU	LTIVITAMIN RENAL - Special Authority see SA1546 below - Retail pharmacy			
*	Cap	30	,	Clinicians Renal Vit

⇒SA1546 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or
- 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73 m² body surface area (BSA).

MULTIVITAMINS - Special Authority see SA1036 below - Retail pharmacy

*	Powder	72.00	200 a OP	✓ Paediatric Seravit

⇒SA1036 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins.

VITAMINIC

VI	AMINO		
*	Tab (BPC cap strength)11.45	1,000	✓ Mvite
	Cap (fat soluble vitamins A, D, E, K) - Special Authority see		
	SA1720 below – Retail pharmacy23.40	60	✓ Vitabdeck

⇒SA1720 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency; or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
Minerals				
Calcium				
CALCIUM CARBONATE * Tab eff 1.75 g (1 g elemental)	28.40	20	•	Calcium Sandoz S29
* Tab 1.25 g (500 mg elemental)	7.52	250	✓.	Arrow-Calcium
CALCIUM GLUCONATE * Inj 10%, 10 ml ampoule	32.00	10	✓	Max Health - Hameln 829
	64.00	20	✓	Max Health \$29
Fluoride				
SODIUM FLUORIDE * Tab 1.1 mg (0.5 mg elemental)	5.75	100	✓	PSM
lodine				
POTASSIUM IODATE Tab 253 mcg (150 mcg elemental iodine)	4.58	90	✓	NeuroTabs
Iron				
FERRIC CARBOXYMALTOSE – Special Authority see SA1840 Inj 50 mg per ml, 10 ml		acy 1	•	Ferinject

⇒SA1840 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:

	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer
continued			

FERROUS FUMARATE

- 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
- 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
- 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
- 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

* Tab 200 mg (65 mg elemental)	100	✓ Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID		
* Tab 310 mg (100 mg elemental) with folic acid 350 mcg4.68	60	✓ Ferro-F-Tabs
FERROUS SULFATE		
* Oral liq 30 mg (6 mg elemental) per 1 ml	500 ml	✓ Ferodan
FERROUS SULPHATE		
* Tab long-acting 325 mg (105 mg elemental)2.06	30	✓ Ferrograd
IRON POLYMALTOSE		
* Inj 50 mg per ml, 2 ml ampoule	5	✓ Ferrosig
Magnesium		
For magnesium hydroxide mixture refer Standard Formulae, page 248		
MAGNESIUM HYDROXIDE		
Suspension 8%72.20	500 ml	✓ T&R \$29
MAGNESIUM SULPHATE		
* Inj 2 mmol per ml, 5 ml ampoule10.21	10	✓ DBL
		✓ DBL S29 S29
Zinc		
ZINC SULPHATE		
* Cap 137.4 mg (50 mg elemental)11.00	100	✓ Zincaps

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus: and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus: and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*: and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per		Manufacturer
EPOETIN ALFA - Special Authority see SA1775 on the previous	page – Retail pharm	асу		
Wastage claimable				
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	✓	Binocrit
Inj 2,000 iu in 1 ml, syringe		6	✓	Binocrit
Inj 3,000 iu in 0.3 ml, syringe	150.00	6	✓	Binocrit
Inj 4,000 iu in 0.4 ml, syringe	96.50	6	✓	Binocrit
Inj 5,000 iu in 0.5 ml, syringe	125.00	6	✓	Binocrit
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	✓	Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	1	Binocrit
Inj 10,000 iu in 1 ml, syringe		6	✓	Binocrit
Inj 40,000 iu in 1 ml, syringe		1	•	Binocrit

Megaloblastic

-01	10	40	
-OL	_IC	AC	טו

*	Tab 0.8 mg21.84	1,000	1	Apo-Folic Acid
	Tab 5 mg	500	1	Apo-Folic Acid
	Oral lig 50 mcg per ml	25 ml OP	1	Biomed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
lnj 1,000 iu vial	2,450.00	1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	Alprolix
ELTROMBOPAG – Special Authority see SA1743 Wastage claimable	below - Retail pharmacy		

Tab 50 mg3,100.00 **➤ SA1743** Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab);
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding: or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (**idiopathic thrombocytopenic purpura - preparation for splenectomy**) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist.

continued...

✓ Revolade

✓ Revolade

28 28

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
 \$	Per	1	Manufacturer

continued...

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Fither:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe	2,356.60	1	✓ NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓ NovoSeven RT
Ini 8 ma svrinae	9.426.40	1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	1,315.00	1	✓ FEIBA NF
Inj 1,000 U	2,630.00	1	✓ FEIBA NF
Inj 2,500 U	6,575.00	1	✓ FEIBA NF

	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
	(Manufacturer's Price)	Per	Subsidised •	Manufacturer
MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [X	pharm]			
For patients with haemophilia. Rare Clinical Circumstance				
treatment is managed by the Haemophilia Treaters Group	ho in conjunction with the N	lation	al Haemop	hilia Management Group,
subject to criteria.				
Inj 250 iu prefilled syringe		1		(yntha
Inj 500 iu prefilled syringe		1		(yntha
Inj 1,000 iu prefilled syringe	·	1		(yntha
Inj 2,000 iu prefilled syringe	· ·	1		(yntha
Inj 3,000 iu prefilled syringe	•	1	• ,	(yntha
NONACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xpha				
For patients with haemophilia. Access to funded treatme	nt is managed by the Hae	emoph	nilia Treate	rs Group in conjunction
with the National Haemophilia Management Group.	405.00			NVIIDIO
Inj 500 iu vial		1		RIXUBIS
Inj 1,000 iu vial		1	_	RIXUBIS
Inj 2,000 iu vial Inj 3,000 iu vial	,	1		rixubis Rixubis
• •	•	1	V F	IIVODIO
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE)) – [Xpharm]			
For patients with haemophilia. Preferred Brand of short h	nalf-life recombinant facto	r VIII.	Access to	funded treatment is
managed by the Haemophilia Treaters Group in conjunct				
Inj 250 iu vial		1		Advate
Inj 500 iu vial		1	_	Advate
Inj 1,000 iu vial		1	_	Advate Advate
Inj 1,500 iu vial Inj 2,000 iu vial	*	1		Advate Advate
Inj 3,000 iu vial	,	1		Advate
DCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENA	•	'	• ,	wate
For patients with haemophilia. Rare Clinical Circumstand		****	mbinant fa	ator VIII Assocs to fundad
treatment is managed by the Haemophilia Treaters Group				
subject to criteria.	in conjunction with the r	valion	ai naeiliop	irilla Mariagement Group,
Inj 250 iu vial	237 50	1	√ k	Kogenate FS
		•		
•	475.00	1	K	
Inj 500 iu vial		1		(ogenate FS
lnj 500 iu vial Inj 1,000 iu vial	950.00	1 1 1	✓ k	Cogenate FS Cogenate FS
Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial	950.00 1,900.00	1	✓ k ✓ k	Kogenate FS Kogenate FS Kogenate FS
Inj 500 iu vial	950.00 1,900.00 2,850.00	1	✓ k ✓ k	Cogenate FS Cogenate FS
Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial Inj 3,000 iu vial RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR \	950.00 1,900.00 2,850.00 /III] – [Xpharm]	1 1 1	✓ h ✓ h	Cogenate FS Cogenate FS Cogenate FS Cogenate FS
Inj 500 iu vial	950.00 1,900.00 2,850.00 /III] – [Xpharm] atment. Access to funded	1 1 1	✓ h ✓ h	Cogenate FS Cogenate FS Cogenate FS Cogenate FS
Inj 500 iu vial		1 1 1 treat	✓ k ✓ k tment is ma	Kogenate FS Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophili
Inj 500 iu vial		1 1 1	Ment is ma	Kogenate FS Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophili
Inj 500 iu vial		1 1 1 treat	tment is ma	Kogenate FS Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophilic Adynovate Adynovate
Inj 500 iu vial		1 1 1 treat	iment is ma	Kogenate FS Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophili
Inj 500 iu vial		1 1 1 treat 1 1	iment is ma	Kogenate FS Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophilis Adynovate Adynovate Adynovate Adynovate
Inj 500 iu vial		1 1 1 treat 1 1 1	iment is ma	Kogenate FS Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophilia Adynovate Adynovate Adynovate Adynovate
Inj 500 iu vial		1 1 1 treat 1 1	tment is ma	Kogenate FS Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophilia Adynovate Adynovate Adynovate Adynovate Adynovate Adynovate Adynovate
Inj 500 iu vial		1 1 1 treat 1 1 1	tment is ma	Kogenate FS Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophilia Adynovate Adynovate Adynovate Adynovate
Inj 500 iu vial		1 1 1 treat 1 1 1	tment is ma	Kogenate FS Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophilia Adynovate Adynovate Adynovate Adynovate Adynovate Adynovate Adynovate

	Subsidy (Manufacturer's Price) \$	Sub:	Fully sidised	Brand or Generic Manufacturer	
Vitamin K					
PHYTOMENADIONE Inj 2 mg per 0.2 ml — Up to 5 inj available on a PSO Inj 10 mg per ml, 1 ml — Up to 5 inj available on a PSO		5 5	• • •	onakion MM onakion MM	
Antithrombotic Agents					
Antiplatelet Agents					
ASPIRIN * Tab 100 mg CLOPIDOGREL	10.80	990	✓ <u>E</u>	thics Aspirin EC	
* Tab 75 mg	4.60	84	_	lopidogrel Multichem	
DIPYRIDAMOLE					
* Tab long-acting 150 mg	10.90	60	✓ P	ytazen SR	
PRASUGREL - Special Authority see SA1954 below - Retail pha	armacy				
Tab 5 mg		28	_	ffient	
Tab 10 mg	120.00	28	✓ E	ffient	

(Effient Tab 5 mg to be delisted 1 February 2021) (Effient Tab 10 mg to be delisted 1 February 2021)

⇒SA1954 Special Authority for Subsidy

Renewal — (coronary angioplasty and bare metal stent) from any relevant practitioner. Approvals valid for 6 months where the patient has undergone coronary angioplasty or had a bare metal cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Renewal — (drug eluting stent) from any relevant practitioner. Approvals valid for 12 months where had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Note: * Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

TICAGRELOR - Special Authority see SA1955 below - Retail pharmacy

***** Tab 90 mg90.00 56 **✓ Brilinta**

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

function assay and requires antiplatelet treatment with ticagrelor; or

- 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal — (**Percutaneous coronary intervention with stent deployment**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Heparin and Antagonist Preparations

ENOXAPARIN SODIUM - Special Authority see SA1646 on the next page - Retail pharmacy

Inj 20 mg in 0.2 ml syringe	27.93	10	✓ Clexane
Inj 40 mg in 0.4 ml syringe	37.27	10	✓ Clexane
Inj 60 mg in 0.6 ml syringe		10	Clexane
Inj 80 mg in 0.8 ml syringe		10	Clexane
Inj 100 mg in 1 ml syringe		10	Clexane
Inj 120 mg in 0.8 ml syringe	116.55	10	Clexane
, , , ,			 Clexane Forte
Inj 150 mg in 1 ml syringe	133.20	10	Clexane
			 Clexane Forte

(Clexane Inj 120 mg in 0.8 ml syringe to be delisted 1 January 2021)

(Clexane Inj 150 mg in 1 ml syringe to be delisted 1 January 2021)

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised Per	Generic Manufacturer

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

HEPARIN SODIUM			
Inj 1,000 iu per ml, 5 ml ampoule	58.57	50	✓ Pfizer
Inj 5,000 iu per ml, 1 ml	28.40	5	✓ Pfizer
	32.66		✓ Hospira
Inj 5,000 iu per ml, 5 ml ampoule	203.68	50	✓ Pfizer
Inj 25,000 iu per ml, 0.2 ml		5	✓ Hospira
	42.40		✓ Heparin DBL S29
			✓ Heparin
			Ratiopharm S29
	122.00	10	✓ Wockhardt S29
	190.00	50	✓ Pfizer S29
(Heparin Ratiopharm \$29 Inj 25,000 iu per ml, 0.2 ml t	o be delisted 1 January 202	1)	
(Wockhardt \$29 Inj 25,000 iu per ml, 0.2 ml to be delis	ted 1 January 2021)	,	
(Pfizer \$29) Inj 25,000 iu per ml, 0.2 ml to be delisted 1	• '		
HEPARINISED SALINE			
	6E 40	50	✓ Pfizer
Inj 10 iu per ml, 5 ml	00.40	30	▼ Flizer
Oral Anticoagulants			
Oral Anticoagulants			
DARIGATRAN			

60

60 60 ✓ Pradaxa
✓ Pradaxa

✓ Pradaxa

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	I Generic
	\$	Per		Manufacturer
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day	83.10	30	✓	Xarelto
Tab 15 mg - Up to 14 tab available on a PSO	77.56	28	✓	Xarelto
Tab 20 mg	77.56	28	✓	Xarelto
VARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
★ Tab 1 mg	3.46	50	✓	Coumadin
v	6.46	100	✓	Marevan
★ Tab 2 mg	4.31	50	✓	Coumadin
★ Tab 3 mg		100	✓	Marevan
₭ Tab 5 mg		50	✓	Coumadin
ř	11.48	100	✓	Marevan
Blood Colony-stimulating Factors				
UI ODAOTIM — Ossaisi Asilysiikassa OA4050 kalaas — Datailys	l		·	
FILGRASTIM – Special Authority see SA1259 below – Retail p		40	,	NII Al
Inj 300 mcg per 0.5 ml prefilled syringe		10	_	Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe	161.50	10	•	<u>Nivestim</u>

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10⁹/L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1912 below - Retail pharmacy ✓ Neulastim

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration

GLUCOSE [DEXTROSE] * Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO.......30.65 ✓ Biomed Biomed to be Sole Supply on 1 November 2020 ✓ Riomed

Biomed to be Sole Supply on 1 November 2020

	DECOD AIV	D DEOOD	7 I STIMING STIGANS
	Subsidy (Manufacturer's Prio \$	ce) Subs Per	Fully Brand or sidised Generic ✓ Manufacturer
POTASSIUM CHLORIDE			
* Inj 75 mg per ml, 10 ml	55.00	50	 ✓ AstraZeneca ✓ Potassium Chloride Aguettant \$29
SODIUM BICARBONATE Inj 8.4%, 50 ml	10.05	1	✓ Biomed
a) Up to 5 inj available on a PSO b) Not in combination			
Inj 8.4%, 100 ml	20.50	1	✓ Biomed
a) Up to 5 inj available on a PSOb) Not in combination			
SODIUM CHLORIDE			
Not funded for use as a nasal drop. Not funded for nebuliser for nebuliser use.	ruse except when	used in conju	unction with an antibiotic intended
Inj 0.9%, bag - Up to 2000 ml available on a PSO	1.23	500 ml	✓ Baxter
	1.26	1,000 ml	✓ Baxter
Only if prescribed on a prescription for renal dialysis, ma for emergency use. (500 ml and 1,000 ml packs)	ternity or post-nata	al care in the	home of the patient, or on a PSC
Inj 23.4% (4 mmol/ml), 20 ml ampoule		5	✓ Biomed
For Sodium chloride oral liquid formulation refer Standar			. Cracanius Kahi
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO	2.80 E 40	20 50	 ✓ Fresenius Kabi ✓ Fresenius Kabi
Inj 0.9%, 20 ml ampoule — Op to 5 inj available on a PSO		20	✓ Fresenius Kabi
TOTAL PARENTERAL NUTRITION (TPN)		20	1 Toochias Itabi
Infusion	CBS	1 OP	✓ TPN
WATER 1) On a prescription or Practitioner's Supply Order only who Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of eyel, when used for the dilution of sodium chloride soln 7% for the supply of the solution of sodium chloride soln 7% for the supply order.	e drops; or		
Inj 5 ml ampoule - Up to 5 inj available on a PSO	7.00	50	✓ InterPharma
Inj 10 ml ampoule - Up to 5 inj available on a PSO		50	✓ Pfizer
Inj 20 ml ampoule – Up to 5 inj available on a PSO	5.00	20	✓ Fresenius Kabi✓ Multichem
	7.50	30	✓ InterPharma
Oral Administration			
CALCIUM POLYSTYRENE SULPHONATE Powder	169.85	300 g OP	✓ Calcium Resonium
COMPOUND ELECTROLYTES		-	
Powder for oral soln — Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]	9.//	50	✓ <u>Electral</u>
Soln with electrolytes (2 × 500 ml)	6.55	,000 ml OP	✓ <u>Pedialyte -</u> <u>Bubblegum</u>
PHOSPHORUS	00.75	400	/ Bi
Tab eff 500 mg (16 mmol)	82.50	100	✓ Phosphate Phebra

	Subsidy (Manufacturer's Price	e) Sı	Fully obsidised	Brand or Generic	
	\$	Per	✓	Manufacturer	
POTASSIUM CHLORIDE					
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	5.26	60			
	(11.85)		C	Chlorvescent	
* Tab long-acting 600 mg (8 mmol)	8.90	200	✓ S	pan-K	
SODIUM BICARBONATE					
Cap 840 mg	8.52	100	✓ S	odibic	
			√ S	odibic	
SODIUM POLYSTYRENE SULPHONATE					
Powder	84.65	454 a OP	✓ R	Resonium-A	

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Alpha-Adrenoceptor Blockers

Alpha Adrenoceptor Blockers

DOXAZOSIN			
Tab 2 mg		500	✓ Apo-Doxazosin
Tab 4 mg	10.80	500	✓ Apo-Doxazosin
PHENOXYBENZAMINE HYDROCHLORIDE			
* Cap 10 mg	65.00	30	✓ BNM S29
	216.67	100	✓ Dibenzyline S29
PRAZOSIN			
* Tab 1 mg	5.53	100	✓ Apo-Prazosin
* Tab 2 mg	7.00	100	✓ Apo-Prazosin
* Tab 5 mg	11.70	100	✓ Apo-Prazosin
TERAZOSIN			
* Tab 1 mg	0.59	28	✓ Actavis
Tab 2 mg	7.50	500	✓ Apo-Terazosin
Tab 5 mg		500	✓ Apo-Terazosin
(Actavis Tab 1 mg to be delisted 1 October 2020)			

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

CAPTOPRIL		_
* Oral liq 5 mg per ml94.99	95 ml OP	✓ Capoten
Oral liquid restricted to children under 12 years of age.		
CILAZAPRIL		
* Tab 0.5 mg2.09	90	✓ Zapril
* Tab 2.5 mg4.80	90	✓ Zapril
* Tab 5 mg	90	✓ Zapril
ENALAPRIL MALEATE		
* Tab 5 mg1.82	100	✓ Acetec
* Tab 10 mg	100	✓ Acetec
* Tab 20 mg	100	✓ Acetec
··· · · · · · · · · · · · · ·	100	Acetec
LISINOPRIL		
* Tab 5 mg2.07	90	✓ Ethics Lisinopril
* Tab 10 mg2.36	90	Ethics Lisinopril
* Tab 20 mg3.17	90	Ethics Lisinopril
PERINDOPRIL		
* Tab 2 mg	30	Apo-Perindopril
* Tab 4 mg4.80	30	✓ Apo-Perindopril
QUINAPRIL		
* Tab 5 mg	90	✓ Arrow-Quinapril 5
* Tab 10 mg	90	✓ Arrow-Quinapril 10
	90	✓ Arrow-Quinapril 20
* Tab 20 mg4.89	90	MITOW-Quillapili 20

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)	Subsi Per	Fully dised	Brand or Generic Manufacturer	
ACE Inhibitors with Divrotios					

ACE Inhibitors with Diuretics

CILAZAPRIL WITH HYDROCHLOROTHIAZIDE - Subsidy by endorsement

Subsidy by endorsement - Subsidised for patients who were taking cilazapril with hydrochlorothiazide prior to 1 March 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril with hydrochlorothiazide.

Tab 5 mg with hydrochlorothiazide 12.5 mg......10.18 100 ✓ Apo-Cilazapril/ Hydrochlorothiazide

(Apo-Cilazapril/ Hydrochlorothiazide Tab 5 mg with hydrochlorothiazide 12.5 mg to be delisted 1 December 2020)

HINAPRII	WITH HYD	ROCHL	OROTHIAZIDE

	Tab 10 mg with hydrochlorothiazide 12.5 mg	3.57	28	✓ Accuretic
	• •	3.83	30	✓ Accuretic 10
*	Tab 20 mg with hydrochlorothiazide 12.5 mg	4.92	30	✓ Accuretic 20

Angiotensin II Antagonists

CANDESARTAN CILEXETII

0,	INDEONITY OF CHERE ITE			
*	Tab 4 mg	1.90	90	✓ Candestar
	Tab 8 mg		90	✓ Candestar
	Tab 16 mg		90	✓ Candestar
	Tab 32 mg		90	✓ Candestar
	SARTAN POTASSIUM			
*	Tab 12.5 mg	1.39	84	✓ Losartan Actavis
*	Tab 25 mg	1.63	84	Losartan Actavis
*	Tab 50 mg	2.00	84	Losartan Actavis
	Tab 100 mg		84	✓ Losartan Actavis

Angiotensin II Antagonists with Diuretics

LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE		
* Tab 50 mg with hydrochlorothiazide 12.5 mg	30	✓ Arrow-Losartan &
		Hydrochlorothiazide

Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN - Special Authority see SA1905 below - Retail pharmacy

Note: Due to the angiotensin II receptor blocking activity of sacubitril with valsartan it should not be co-administered with an ACE inhibitor or another ARR

AGE ITTIBITOR OF ALIGHRIC AT ID.			
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/103

⇒SA1905 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II: or

continued...

Subsidy	Fu	,	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
\$	Per	/	Manufacturer

continued...

- 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, page 119

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaestnetics, Local, p	age 119	
AMIODARONE HYDROCHLORIDE		
▲ Tab 100 mg	30	✓ Aratac
▲ Tab 200 mg5.25	30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule - Up to 10 inj available on a		
PSO16.37	10	✓ Max Health
ATROPINE SULPHATE		
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO12.07	10	✓ Martindale
	10	Martinuale
DIGOXIN		
* Tab 62.5 mcg - Up to 30 tab available on a PSO7.00	240	✓ Lanoxin PG
* Tab 250 mcg - Up to 30 tab available on a PSO15.20	240	✓ <u>Lanoxin</u>
* Oral liq 50 mcg per ml16.60	60 ml	Lanoxin
		✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE		
▲ Cap 100 mg23.87	100	✓ Rythmodan
FLECAINIDE ACETATE		,
▲ Tab 50 mg	60	✓ Flecainide BNM
▲ Cap long-acting 100 mg	90	✓ Flecainide Bivivi
Cap long-acting 100 mg9.51	90	Controlled
		Release Teva
A Con long acting 000 mg	00	✓ Flecainide
▲ Cap long-acting 200 mg61.06	90	
		<u>Controlled</u> Release Teva
1,10	_	
Inj 10 mg per ml, 15 ml ampoule100.00	5	✓ Tambocor
MEXILETINE HYDROCHLORIDE		
▲ Cap 150 mg162.00	100	Mexiletine
		Hydrochloride
		USP S29
▲ Cap 250 mg202.00	100	Mexiletine
		Hydrochloride
		USP S29
PROPAFENONE HYDROCHLORIDE		
▲ Tab 150 mg	50	✓ Rytmonorm
		,

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
•	Por	1	Manufacturer

Antihypotensives

MIDODRINE - Special Authority see SA1474 below - Retail pharm	nacy		
Tab 2.5 mg	53.00	100	Gutron
Tab 5 mg	79.00	100	Gutron

⇒SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Note: Treatment should be started with small doses and titrated upwards as necessary. Hypertension should be avoided, and the usual target is a standing systolic blood pressure of 90 mm Hg.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

	ENOLOL		
*	Tab 50 mg		Mylan Atenolol
*	Tab 100 mg7.30		Mylan Atenolol
*	Oral liq 25 mg per 5 ml21.25		Atenolol AFT
		•	✓ Atenolol AFT
			S29 S29
	Restricted to children under 12 years of age.		
BIS	OPROLOL FUMARATE		
*	Tab 2.5 mg	90	✓ Bosvate
*	Tab 5 mg	90	✓ Bosvate
*	Tab 10 mg		✓ Bosvate
CA	RVEDILOL		
*	Tab 6.25 mg	60	✓ Carvedilol Sandoz
*	Tab 12.5 mg		✓ Carvedilol Sandoz
*	Tab 25 mg	60	 Carvedilol Sandoz
CF	LIPROLOL		
*	Tab 200 mg21.40	180	✓ Celol
LAE	BETALOL		
*	Tab 100 mg14.50	100	✓ Trandate
	Tab 200 mg27.00		✓ Trandate
*	Inj 5 mg per ml, 20 ml ampoule59.06		
	(88.60))	Trandate
*	inj 5 mg per ml, 20 ml vial	1	
	(48.20)		Alvogen S29
ME	TOPROLOL SUCCINATE		
*	Tab long-acting 23.75 mg	30	✓ Betaloc CR
*	Tab long-acting 47.5 mg		✓ Betaloc CR
*	Tab long-acting 95 mg1.99		✓ Betaloc CR
*	Tab long-acting 190 mg		✓ Betaloc CR
	• •		

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
	\$	Per		Manufacturer
METOPROLOL TARTRATE				
* Tab 50 mg	5.66	100	1	Apo-Metoprolol
* Tab 100 mg		60	✓	Apo-Metoprolol
* Tab long-acting 200 mg	23.40	28	1	Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial	29.50	5	✓	Metroprolol IV
				Mylan
NADOLOL				
* Tab 40 mg	16.69	100	1	Apo-Nadolol
* Tab 80 mg		100	1	Apo-Nadolol
PINDOLOL				
* Tab 5 mg	13.22	100	1	Apo-Pindolol
* Tab 10 mg		100		Apo-Pindolol
* Tab 15 mg		100	_	Apo-Pindolol
PROPRANOLOL				
* Tab 10 mg	4 64	100	1	Apo-Propranolol
* Tab 40 mg		100		Apo-Propranolol
* Cap long-acting 160 mg		100		Cardinol LA
Cap long acting 100 mg Special Authority see SA1327 below -		100	•	Variation En
, , ,		500 m	al ./	Roxane S29
Retail pharmacy		וו טטכ	II •	nuxaile 229

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTAL	JL

	Tab 80 mg Tab 160 mg		500 100	✓ Mylan✓ Mylan
TIN	MOLOL			
*	Tab 10 mg	10.55	100	✓ Apo-Timol

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

LODIPINE			
Tab 2.5 mg	1.72	100	✓ Apo-Amlodipine
Tab 5 mg	3.33	250	✓ Apo-Amlodipine
Tab 10 mg	4.40	250	✓ Apo-Amlodipine
ODIPINE			
Tab long-acting 2.5 mg	1.45	30	✓ Plendil ER
Tab long-acting 5 mg	3.93	90	✓ Felo 5 ER
Tab long-acting 10 mg	4.32	90	✓ Felo 10 ER
	Tab 2.5 mg Tab 5 mg Tab 10 mg ODIPINE Tab long-acting 2.5 mg. Tab long-acting 5 mg.	Tab 2.5 mg 1.72 Tab 5 mg 3.33 Tab 10 mg 4.40	Tab 2.5 mg 1.72 100 Tab 5 mg 3.33 250 Tab 10 mg 4.40 250 ODIPINE 250 Tab long-acting 2.5 mg 1.45 30 Tab long-acting 5 mg 3.93 90

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
FEDIPINE				
Tab long-acting 10 mg	10.63	60	1	Adalat 10
				Adefin S29
Tab long-acting 20 mg	17.72	100	•	Nyefax Retard
Tab long-acting 30 mg	3.14	30		Adalat Oros
Tab long-acting 60 mg	5.67	30		Adalat Oros
			•	Adefin XL
Other Calcium Channel Blockers				
LTIAZEM HYDROCHLORIDE				
Tab 30 mg	4.60	100		Dilzem
Tab 60 mg	8.50	100		Dilzem
Cap long-acting 120 mg	33.42	500	✓	Apo-Diltiazem CD
Cap long-acting 180 mg	50.05	500	✓	Apo-Diltiazem CD
Cap long-acting 240 mg	66.76	500	1	Apo-Diltiazem CD
RHEXILINE MALEATE				
Tab 100 mg	62.90	100	•	Pexsig
RAPAMIL HYDROCHLORIDE				
Tab 40 mg	7.01	100	1	Isoptin
Tab 80 mg	11.74	100	✓	Isoptin
Tab long-acting 120 mg	36.02	100	✓	Isoptin Retard \$29
			1	Isoptin SR
Tab long-acting 240 mg	15.12	30	✓	Isoptin SR
Inj 2.5 mg per ml, 2 ml ampoule - Up to 5 inj available of	na			
PSO	25.00	5	/	Isoptin
Centrally-Acting Agents				
ONIDINE				
Patch 2.5 mg, 100 mcg per day - Only on a prescription	10.34	4	1	Mylan
Mylan to be Sole Supply on 1 November 2020				
Patch 5 mg, 200 mcg per day - Only on a prescription	13.18	4	•	Mylan
Mylan to be Sole Supply on 1 November 2020			_	
Patch 7.5 mg, 300 mcg per day — Only on a prescription Mylan to be Sole Supply on 1 November 2020	16.93	4	•	Mylan
, , , , , , , , , , , , , , , , , , , ,				
ONIDINE HYDROCHLORIDE	0.75	440	,	Olamidina Dalla
Tab 25 mcg		112		Clonidine BNM
Tab 150 mcglpi 150 mcg.por ml 1 ml ampoulo		100		Catapres
Inj 150 mcg per ml, 1 ml ampoule	∠ე.५0	10	•	<u>Medsurge</u>
ETHYLDOPA	45.40	400	_	Markland day 184 1
Tab 250 mg	15.10 52.85	100 500		Methyldopa Mylan Methyldopa Mylan

Brand or

Fully

28

50

✓ Frumil

✓ Moduretic

Subsidy

Diuretics		(Manufacturer's P	Price) Subs Per		neric Inufacturer
# Tab 1 mg	Diuretics				
# Tab 1 mg	Loop Diuretics				
16.36	BUMETANIDE				
# Inj 500 mcg per ml, 4 ml vial	* Tab 1 mg	4.91	30	Burin	ex S29 S29
FUROSEMIDE [FRUSEMIDE] * Tab 40 mg - Up to 30 tab available on a PSO					
* Tab 40 mg - Up to 30 tab available on a PSO	* Inj 500 mcg per ml, 4 ml vial	7.95	5	Burin	ex
* Tab 500 mg					_
* Oral liq 10 mg per ml	• 1		,		
* Inj 10 mg per ml, 25 ml ampoule	•				
* Inj 10 mg per ml, 2 ml ampoule − Up to 5 inj available on a PSO1.15 5					
Potassium Sparing Diuretics AMILORIDE HYDROCHLORIDE Oral liq 1 mg per ml					
AMILORIDE HYDROCHLORIDE Oral liq 1 mg per ml					
Oral liq 1 mg per ml	Potassium Sparing Diuretics				
EPLERENONE — Special Authority see SA1728 below — Retail pharmacy Tab 50 mg	AMILORIDE HYDROCHLORIDE				
Tab 50 mg	Oral liq 1 mg per ml	30.00	25 ml OP	✓ Biome	ed
Tab 25 mg	EPLERENONE - Special Authority see SA1728 below - Retail p	pharmacy			
Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both: 1 Patient has heart failure with ejection fraction less than 40%; and 2 Either: 2.1 Patient is intolerant to optimal dosing of spironolactone; or 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone. METOLAZONE Tab 5 mg	Tab 50 mg	17.00			_
Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both: 1 Patient has heart failure with ejection fraction less than 40%; and 2 Either: 2.1 Patient is intolerant to optimal dosing of spironolactone; or 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone. METOLAZONE Tab 5 mg		11.87	30	✓ Inspra	<u>a</u>
the following criteria: Both: 1 Patient has heart failure with ejection fraction less than 40%; and 2 Either: 2.1 Patient is intolerant to optimal dosing of spironolactone; or 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone. METOLAZONE Tab 5 mg					
Both: 1 Patient has heart failure with ejection fraction less than 40%; and 2 Either: 2.1 Patient is intolerant to optimal dosing of spironolactone; or 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone. METOLAZONE Tab 5 mg		d without further	renewal unless	notified for	applications meeting
1 Patient has heart failure with ejection fraction less than 40%; and 2 Either: 2.1 Patient is intolerant to optimal dosing of spironolactone; or 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone. METOLAZONE Tab 5 mg	· · · · · · · · · · · · · · · · · · ·				
2 Either: 2.1 Patient is intolerant to optimal dosing of spironolactone; or 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone. METOLAZONE Tab 5 mg		00/ · and			
2.1 Patient is intolerant to optimal dosing of spironolactone; or 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone. METOLAZONE Tab 5 mg	•	7/0, and			
2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone. METOLAZONE Tab 5 mg		ctone: or			
METOLAZONE Tab 5 mg			on optimal dos	ing of spiror	nolactone.
Tab 5 mg CBS 1 ✓ Metolazone S29 50 ✓ Zaroxolyn S29 SPIRONOLACTONE ★ Tab 25 mg 4.38 100 ✓ Spiractin ★ Tab 100 mg 11.80 100 ✓ Spiractin	METOLAZONE		·	•	
50 ✓ Zaroxolyn 529 SPIRONOLACTONE * Tab 25 mg		CBS	1	✓ Metal	azone S29
SPIRONOLACTONE ★ Tab 25 mg 4.38 100 ✓ Spiractin ★ Tab 100 mg 11.80 100 ✓ Spiractin	Tab o mg		•		
★ Tab 25 mg 4.38 100 ✓ Spiractin ★ Tab 100 mg 11.80 100 ✓ Spiractin	SPIRONOL ACTONE		00	Luion	
* Tab 100 mg11.80 100 ✓ Spiractin		4 38	100	✓ Spira	ctin
·	3			•	
	•		25 ml OP		

Potassium Sparing Combination Diuretics

AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE

* Tab 5 mg with furosemide 40 mg8.63

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE] * Tab 2.5 mg – Up to 150 tab available on a PSO	20.00	500	•	Arrow- Bendrofluazide
a) May be supplied on a PSO for reasons other than em b) Arrow-Bendrofluazide to be Sole Supply on 1 Decem * Tab 5 mg	ber 2020	500	•	Arrow- Bendrofluazide
Arrow-Bendrofluazide to be Sole Supply on 1 December	2020			
CHLOROTHIAZIDE Oral liq 50 mg per ml	26.00 25	5 ml C	P 🗸	Biomed
CHLORTALIDONE [CHLORTHALIDONE] * Tab 25 mg	6.50	50	1	<u>Hygroton</u>
INDAPAMIDE * Tab 2.5 mg Dapa-Tabs to be Sole Supply on 1 November 2020	10.45	90	•	Dapa-Tabs
Lipid-Modifying Agents				
Fibrates				
BEZAFIBRATE * Tab 200 mg	12.89 taking gemfibrozil pr		1 August 2	
dispensing of gemfibrozil. * Tab 600 mg		60		Lipazil
Other Lipid-Modifying Agents				
ACIPIMOX * Cap 250 mg	21.56	30		Olbetam Olbetam S29 S29
NICOTINIC ACID Tab 50 mg Tab 500 mg		100 100		Apo-Nicotinic Acid Apo-Nicotinic Acid
Resins				
COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g	28.60	30	/	Colestid

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
HMG CoA Reductase Inhibitors (Statins)				
ATORVASTATIN				
* Tab 10 mg	6.96	500	√ <u>L</u>	<u>orstat</u>
* Tab 20 mg	9.99	500	✓ <u>L</u>	<u>orstat</u>
* Tab 40 mg	15.93	500	✓ <u>L</u>	<u>-orstat</u>
* Tab 80 mg	27.19	500	✓ <u>L</u>	<u>-orstat</u>
PRAVASTATIN				
* Tab 10 mg	3.55	28	✓ F	Pravastatin Mylan
* Tab 20 mg		100	√	Apo-Pravastatin
* Tab 40 mg		28	✓ F	ravastatin Mylan
·	8.06	100		Apo-Pravastatin
SIMVASTATIN				
* Tab 10 mg Simvastatin Mylan to be Sole Supply on 1 November 20		90	√ 9	Simvastatin Mylan
* Tab 20 mg		90	1 0	Simvastatin Mylan
Simvastatin Mylan to be Sole Supply on 1 November 20		30	• •	Jilivastatili Niylali
* Tab 40 mg		90	/ 9	Simvastatin Mylan
Simvastatin Mylan to be Sole Supply on 1 November 20		00		miraotatiii mytan
* Tab 80 mg		90	/ 9	Simvastatin Mylan
Simvastatin Mylan to be Sole Supply on 1 November 20		00		mylan
Cimiracia in injunito de Colo Cappi) on i interember 20	,			
Selective Cholesterol Absorption Inhibitors				
Control of the Contro				

ΕZ	ETIMIBE - Special Authority see SA1045 below - Retail pharmacy		
*	Tab 10 mg1.95	30	✓ Ezetimibe Sandoz
	Ezetimibe Sandoz to be Sole Supply on 1 October 2020		

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 × normal) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atoryastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy. If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 on the next page - Retail pharmacy

Tab 10 mg with simvastatin 10 mg	5.15	30	✓ Zimybe
Tab 10 mg with simvastatin 20 mg		30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg		30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg		30	✓ Zimybe

	Subsidy (Manufacturer's Price)	Sı	Fully ibsidised	Brand or Generic	
	\$	Per	1	Manufacturer	
⇒SA1046 Special Authority for Subsidy					

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and

3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy. If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

GLYCERYL TRI	NITRATE	
--------------	---------	--

*	available on a PSO	4.45	250 dose OP	✓ Nitrolingual Pump
				Spray
*	Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
*	Patch 50 mg, 10 mg per day	18.62	30	✓ Nitroderm TTS
ISC	DSORBIDE MONONITRATE			
*	Tab 20 mg	19.55	100	✓ Ismo 20
	Ismo 20 to be Sole Supply on 1 November 2020			
*	Tab long-acting 40 mg	8.20	30	✓ Ismo 40 Retard
	Ismo 40 Retard to be Sole Supply on 1 November 2020			
*	Tab long-acting 60 mg	9.25	90	Duride
	Duride to be Sole Supply on 1 November 2020			

Sympathomimetics

ADDENIAL INC

ADRENALINE			
Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a PSO	4.98	5	Aspen Adrenaline
	10.76	,	/ DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PSO	27.00	5	✓ Hospira
	49.00	10	Aspen Adrenaline
ISOPRENALINE [ISOPROTERENOL]			
* Inj 200 mcg per ml, 1 ml ampoule	36.80	25	
	(164.20)		Isuprel

Vasodilators

HYDRALAZINE HYDROCHLORIDE

Tab 25 mg - Special Authority see SA1321 on the next page -			
Retail pharmacy	CBS	1	Hydralazine
		56	✓ Onelink S29
		84	✓ AMDIPHARM \$29
		100	✓ Onelink S29
Inj 20 mg ampoule	25.90	5	✓ Apresoline
	' '	Tab 25 mg - Special Authority see SA1321 on the next page - Retail pharmacy	Retail pharmacy

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1321 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 For the treatment of refractory hypertension; or
- 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

MINOXIDIL

▲ Tab 10 mg	70.00	100	Loniten
NICORANDIL			
▲ Tab 10 mg	25.57	60	✓ Ikorel
▲ Tab 20 mg	32.28	60	✓ Ikorel
PAPAVERINE HYDROCHLORIDE			
* Inj 12 mg per ml, 10 ml ampoule	217.90	5	Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]			
Tab 400 mg	42.26	50	✓ Trental 400
•			

Endothelin Receptor Antagonists

AMBRISENTAN – Special Authority see SA1702 below – Retail pharmacy		
Tab 5 mg4,585.00	30	✓ Volibris
Tab 10 mg4,585.00	30	✓ Volibris

⇒SA1702 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

BOSENTAN	 Special / 	Authority see	SA1908 below -	- Retail pharmacy

Tab 62.5 mg	141.00	60	✓ Bosentan Dr
			Reddy's
Tab 125 mg	141.00	60	✓ Bosentan Dr
			Reddy's

⇒SA1908 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil: or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or

continued...

_				
	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Sı	ubsidised	Generic
	\$	Per	/	Manufacturer

continued...

- 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
- 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV: or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy: and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Phosphodiesterase Type 5 Inhibitors

ILDENAFIL - Special Authority see SA1909 below - Retail pharmacy		
Tab 25 mg	4	✓ Vedafil
Tab 50 mg	4	✓ Vedafil
Tab 100 mg	12	✓ Vedafil

⇒SA1909 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital

continued...

SI

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

ulceration; digital ulcers; or gangrene); and

- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II: or
 - 3.2 PAH is in NYHA/WHO functional class III: or
 - 3.3 PAH is in NYHA/WHO functional class IV; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age, or health system capacity constraints.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Prostacyclin Analogues

EPOPROSTENOL - Special Authority see SA1696 below - Retail pharmacy	/	
Inj 500 mcg vial36.6	1 1	✓ Veletri
Inj 1.5 mg vial73.21	1 1	✓ Veletri

⇒SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ILOPROST - Special Authority see SA1705 on the next page - Retail pharmacy

Nebuliser soln 10 mcg per ml, 2 ml740.10 30 ✓ Ventavis

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 88

ADAPALENE

- a) Maximum of 30 g per prescription
- b) Only on a prescription

b) Only on a prescription			
Crm 0.1%	22.89	30 g OP	Differin
Gel 0.1%	22.89	30 g OP	Differin
ISOTRETINOIN - Special Authority see SA1475 below - Retail pharm	nacy		
Cap 5 mg	8.14	60	Oratane
Cap 10 mg	13.34	120	✓ Oratane
Cap 20 mg	20.49	120	✓ Oratane

⇒SA1475 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice: and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
 - 3.2 Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
- 2 Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

TRFTINOIN

Crm 0.5 mg per q − Maximum of 50 g per prescription......13.90 50 g OP ✓ ReTrieve

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 88

HYDROGEN PEROXIDE

	Subsidy	Dai1 0 :	Fully	Brand or
	(Manufacturer's F \$	Price) Subs Per	sidised •	Generic Manufacturer
MUPIROCIN				
Oint 2%	6.60 (9.26)	15 g OP	В	actroban
a) Only on a prescriptionb) Not in combination				
SODIUM FUSIDATE [FUSIDIC ACID]				
Crm 2%	1.59	5 g OP	✓ <u>F</u>	<u>oban</u>
a) Maximum of 5 g per prescription				
b) Only on a prescription				
c) Not in combination	1.50	5 a OB	./ =	ohon
Oint 2%	1.59	5 g OP	• <u>F</u>	<u>oban</u>
a) Maximum of 5 g per prescriptionb) Only on a prescription				
c) Not in combination				
SULFADIAZINE SILVER				
Crm 1%	10.80	50 g OP	√ □	lamazine
a) Up to 250 g available on a PSO	10.00	30 g Oi	• 1	idilidžiii0
b) Not in combination				
For systemic antifungals, refer to INFECTIONS, Antifungals, p AMOROLFINE a) Only on a prescription b) Net in combination	•			
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		5 ml OP	✓ M	lycoNail
AMOROLFINE a) Only on a prescription b) Not in combination		5 ml OP	✓ N	lycoNail
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		5 ml OP	✓ N	lycoNail
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		5 ml OP	✓ N	lycoNail
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	14.93	5 ml OP		lycoNail .po-Ciclopirox
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	14.93			
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	14.93		✓ <u>A</u>	
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	14.93	7 ml OP	✓ <u>A</u>	po-Ciclopirox
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	5.72 0.70	7 ml OP 20 g OP	✓ <u>A</u>	po-Ciclopirox
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		7 ml OP	✓ <u>A</u>	.po-Ciclopirox Clomazol
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	5.72 0.70	7 ml OP 20 g OP	✓ <u>A</u>	po-Ciclopirox
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		7 ml OP 20 g OP	✓ <u>A</u>	.po-Ciclopirox Clomazol
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		7 ml OP 20 g OP	✓ <u>A</u>	.po-Ciclopirox Clomazol
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	5.72 0.70 4.36 (7.55)	7 ml OP 20 g OP 20 ml OP	✓ <u>A</u>	.po-Ciclopirox Clomazol
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	5.72 0.70 4.36 (7.55)	7 ml OP 20 g OP	✓ <u>A</u>	po-Ciclopirox Clomazol
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	5.72 0.70 4.36 (7.55)	7 ml OP 20 g OP 20 ml OP	✓ <u>A</u>	.po-Ciclopirox Clomazol
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	5.72 0.70 4.36 (7.55)	7 ml OP 20 g OP 20 ml OP	✓ <u>A</u>	po-Ciclopirox Clomazol
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		7 ml OP 20 g OP 20 ml OP 20 g OP	✓ <u>A</u>	po-Ciclopirox Clomazol
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		7 ml OP 20 g OP 20 ml OP	✓ <u>A</u> ✓ C	apo-Ciclopirox Clomazol Canesten
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		7 ml OP 20 g OP 20 ml OP 20 g OP	✓ <u>A</u> ✓ C	po-Ciclopirox Clomazol

	Subsidy (Manufacturer's Pri \$	ice) Sul Per	Fully bsidised	Brand or Generic Manufacturer
MICONAZOLE NITRATE				
* Crm 2%	0.74	15 g OP	✓ N	Nultichem
a) Only on a prescription				
b) Not in combination				
* Lotn 2%		30 ml OP		
	(10.03)			Daktarin
a) Only on a prescription				
b) Not in combination				
* Tinct 2%		30 ml OP	_	
	(12.10)		D	Daktarin
a) Only on a prescription				
b) Not in combination				
Antipruritic Preparations				
CALAMINE				
a) Only on a prescription				

a) Only on a prescription

b) Not in combination

✓ healthE Calamine **Aqueous Cream**

ΒP

CROTAMITON

a) Only on a prescription

b) Not in combination

20 g OP

100 a

✓ Itch-Soothe

MENTHOL - Only in combination

1) Only in combination with a dermatological base or proprietary Topical Corticosteriod - Plain

2) With or without other dermatological galenicals.

Crystals......6.92 29.60

25 g 100 g ✓ MidWest

✓ MidWest

Corticosteroids Topical

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 78

Corticosteroids - Plain

BE.	TAMETHASONE DIPROPIONATE		
	Crm 0.05%2.96	15 g OP	Diprosone
	36.00	50 g OP	Diprosone
	Oint 0.05%2.96	15 g OP	Diprosone
	36.00	50 g OP	✓ Diprosone
	Oint 0.05% in propylene glycol base4.33	30 g OP	Diprosone OV
BE	TAMETHASONE VALERATE		
*	Crm 0.1%3.45	50 g OP	✓ Beta Cream
	Oint 0.1%3.45	50 g OP	✓ Beta Ointment
*	Lotn 0.1%	50 ml OP	✓ Betnovate
CL	OBETASOL PROPIONATE		
*	Crm 0.05%2.18	30 g OP	✓ Dermol
*	Oint 0.05%2.12	30 g OP	✓ Dermol

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully Brand or
	(Manufacturer's I	Price) Subs Per	idised Generic ✓ Manufacturer
LODETAGONE BUTYDATE	Ψ	FEI	- Ivianuiaciurei
LOBETASONE BUTYRATE	5.00		
Crm 0.05%		30 g OP	
	(7.09)		Eumovate
IFLUCORTOLONE VALERATE			
Crm 0.1%	8 97	50 g OP	
0111 0.1 /0	(15.86)	00 g O1	Nerisone
Fatty oint 0.1%		50 ~ OD	Nellsolle
rally oint 0.1%		50 g OP	
	(15.86)		Nerisone
Nerisone Crm 0.1% to be delisted 1 December 2020)			
Nerisone Fatty oint 0.1% to be delisted 1 August 2021)			
YDROCORTISONE			
Crm 1% - Only on a prescription	3.70	100 g OP	✓ Hydrocortisone
Citi 1 % – Otily off a prescription		100 g OF	
	4= 4-	500	(<u>PSM)</u>
	17.15	500 g	✓ <u>Hydrocortisone</u>
			<u>(PSM)</u>
Powder – Only in combination	49.95	25 g	✓ ABM
Up to 5% in a dermatological base (not proprietary Top		d – Plain) with o	or without other dermatologica
galenicals			
· ·			
YDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN			
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only	y on		
a prescription	10.57	250 ml	✓ DP Lotn HC
DP Lotn HC to be Sole Supply on 1 October 2020			
,			
YDROCORTISONE BUTYRATE		100 00	
Lipocream 0.1%		100 g OP	 Locoid Lipocream
Oint 0.1%	13.70	100 g OP	✓ <u>Locoid</u>
Milky emul 0.1%	13.70	100 ml OP	✓ Locoid Crelo
IETHYLPREDNISOLONE ACEPONATE			
Crm 0.1%	1.16	15 g OP	✓ Advantan
	4.40	15 y OF	Auvanian
Advantan to be Sole Supply on 1 December 2020		05	
Oint 0.1%	4.46	15 g OP	✓ Advantan
Advantan to be Sole Supply on 1 December 2020			
IOMETASONE FUROATE			
Crm 0.1%	1 51	15 g OP	✓ Elocon Alcohol Free
OIII 0.1 /0	2.50	50 g OP	✓ Elocon Alcohol Free
Oint 0 10/		•	
Oint 0.1%		15 g OP	✓ <u>Elocon</u>
	2.90	50 g OP	✓ <u>Elocon</u>
Lotn 0.1%	6.30	30 ml OP	✓ Elocon
RIAMCINOLONE ACETONIDE			
Crm 0.02%	6 30	100 g OP	✓ Aristocort
	0.30	100 y OF	Alistocolt
Aristocort to be Sole Supply on 1 November 2020	2.05	100 05	/ Autotopo i
Oint 0.02%	6.35	100 g OP	✓ Aristocort
Aristocort to be Sole Supply on 1 November 2020			
Corticosteroids - Combination			
ETAMETHASONE VALEDATE WITH OLIOOLIINOL. Only	n a proceedation		
ETAMETHASONE VALERATE WITH CLIOQUINOL — Only of		45 . 05	
		15 ~ ()[)	
Crm 0.1% with clioquinol 3%	3.49 (4.90)	15 g OP	Betnovate-C

	Subsidy (Manufacturer's P	rice) Subsi	Fully Brand or idised Generic ✓ Manufacturer
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FU Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP	Fucicort
a) Maximum of 15 g per prescriptionb) Only on a prescription			
HYDROCORTISONE WITH MICONAZOLE - Only on a prescri * Crm 1% with miconazole nitrate 2%		15 g OP	✓ <u>Micreme H</u>
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN — Crm 1% with natamycin 1% and neomycin sulphate 0.5% Oint 1% with natamycin 1% and neomycin sulphate 0.5% TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	3.35 3.35	15 g OP 15 g OP	✓ Pimafucort ✓ Pimafucort
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 n and gramicidin 250 mcg per g - Only on a prescription	ng	15 g OP	Viaderm KC
Disinfecting and Cleansing Agents			
CHLORHEXIDINE GLUCONATE – Subsidy by endorsement a) No more than 500 ml per month b) Only if prescribed for a dialysis patient and the prescript Handrub 1% with ethanol 70%	4.29 3.98	cordingly. 500 ml 500 ml	✓ healthE ✓ healthE
a) Maximum of 500 ml per prescription b) a) Only if prescribed for a patient identified with Meth surgery in hospital and the prescription is endorser b) Only if prescribed for a patient with recurrent Stapl accordingly Soln 1% (health E Soln 1% to be delisted 1 November 2020)	d accordingly; or nylococcus aureus		, ,,
Barrier Creams and Emollients			
Barrier Creams			
DIMETHICONE			
* Crm 5% pump bottle	4.48	500 ml OP	✓ <u>healthE</u> Dimethicone 5%
* Crm 10% pump bottle	4.52	500 ml OP	✓ <u>healthE</u> Dimethicone 10%
ZINC AND CASTOR OIL * Oint	4.25	500 g	✓ Boucher
Emollients			
AQUEOUS CREAM * Crm	1.92	500 g	✓ Boucher

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully Brand or	
	(Manufacturer's	Price) Subs	sidised Generic Manufacti	
	<u> </u>	Per	Manuracti	urer
ETOMACROGOL	0.45	500		
€ Crm BP	2.48	500 g	✓ <u>healthE</u>	
ETOMACROGOL WITH GLYCEROL				
Crm 90% with glycerol 10%	2.35	500 ml OP	✓ ADE	
			✓ Boucher	
			✓ Kenkay So	rbolene
	3.10	1,000 ml OP	✓ ADE	
			✓ Boucher	
MULSIFYING OINTMENT				
♦ Oint BP	3.59	500 g	✓ AFT	
OIL IN WATER EMULSION		· ·		
Crm	2 10	500 g	✓ O/W Fatty I	Emuleion
		500 g	Cream	u.
ADAFFIN			Orcalli	
ARAFFIN Oint liquid paraffin E00/, with white eaft paraffin E00/	F 0F	E00 L OD	. hasker	
Oint liquid paraffin 50% with white soft paraffin 50%	5.35	500 ml OP	✓ <u>healthE</u>	
IREA				
← Crm 10%	1.37	100 g OP	✓ healthE Ure	ea Cream
VOOL FAT WITH MINERAL OIL - Only on a prescription				
Lotn hydrous 3% with mineral oil	5.60	1,000 ml		
•	(11.95)	,	DP Lotion	
	1.40	250 ml OP		
	(4.53)	-	DP Lotion	
	5.60	1,000 ml		
	(20.53)	,	Alpha-Keri I	_otion
	(23.91)		BK Lotion	
	1.40	250 ml OP		
	(7.73)	-	BK Lotion	
	` ,			
Other Dermatological Bases				
•				
ARAFFIN	4.00	450 ~	√ hoci4h⊏	
•		450 g	✓ healthE	
ARAFFIN White soft – Only in combination	19.99	2,500 g	✓ healthE	Dloin
ARAFFIN	19.99	2,500 g	✓ healthE	– Plain.
VARAFFIN White soft – Only in combination Only in combination with a dermatological galenical or	19.99	2,500 g	✓ healthE	– Plain.
ARAFFIN White soft – Only in combination	19.99	2,500 g	✓ healthE	– Plain.
VARAFFIN White soft – Only in combination Only in combination with a dermatological galenical or	19.99	2,500 g	✓ healthE	– Plain.
ARAFFIN White soft – Only in combination Only in combination with a dermatological galenical or Minor Skin Infections	19.99 r as a diluent for a	2,500 g	✓ healthE	– Plain.
ARAFFIN White soft – Only in combination Only in combination with a dermatological galenical or Minor Skin Infections OVIDONE IODINE	19.99 r as a diluent for a	2,500 g proprietary Top	✓ healthE ical Corticosteroid	– Plain.
ARAFFIN White soft – Only in combination Only in combination with a dermatological galenical or Minor Skin Infections OVIDONE IODINE Oint 10%	19.99 r as a diluent for a	2,500 g proprietary Top	✓ healthE ical Corticosteroid	– Plain.
ARAFFIN White soft – Only in combination Only in combination with a dermatological galenical or Minor Skin Infections OVIDONE IODINE Oint 10%	19.99 r as a diluent for a	2,500 g proprietary Top	✓ healthE ical Corticosteroid	– Plain.
ARAFFIN White soft – Only in combination Only in combination with a dermatological galenical or Minor Skin Infections OVIDONE IODINE Oint 10%	19.99 r as a diluent for a	2,500 g proprietary Top	✓ healthE ical Corticosteroid	– Plain.
ARAFFIN White soft – Only in combination Only in combination with a dermatological galenical or Minor Skin Infections OVIDONE IODINE Oint 10%	19.99 r as a diluent for a	2,500 g proprietary Top 65 g OP	 ✓ healthE ical Corticosteroid ✓ Betadine 	– Plain.
ARAFFIN White soft – Only in combination Only in combination with a dermatological galenical or Minor Skin Infections OVIDONE IODINE Oint 10%	19.99 r as a diluent for a	2,500 g proprietary Top 65 g OP	 ✓ healthE ical Corticosteroid ✓ Betadine ✓ Riodine 	– Plain.
ARAFFIN White soft – Only in combination Only in combination with a dermatological galenical or Minor Skin Infections OVIDONE IODINE Oint 10%	19.99 r as a diluent for a7.402.553.83 5.40	2,500 g proprietary Top 65 g OP 100 ml 15 ml	 ✓ healthE ical Corticosteroid ✓ Betadine ✓ Riodine ✓ Riodine 	– Plain.
ARAFFIN White soft – Only in combination Only in combination with a dermatological galenical or Minor Skin Infections OVIDONE IODINE Oint 10%	19.99 r as a diluent for a7.402.553.83 5.401.63	2,500 g proprietary Top 65 g OP 100 ml 15 ml 500 ml	 ✓ healthE ical Corticosteroid ✓ Betadine ✓ Riodine ✓ Riodine ✓ Riodine ✓ Riodine 	
ARAFFIN White soft – Only in combination Only in combination with a dermatological galenical or Minor Skin Infections OVIDONE IODINE Oint 10%	19.99 r as a diluent for a7.402.553.83 5.401.63 (3.48)	2,500 g proprietary Top 65 g OP 100 ml 15 ml 500 ml	 ✓ healthE ical Corticosteroid ✓ Betadine ✓ Riodine ✓ Riodine 	

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Parasiticidal Preparations

DIMETHICONE

* Lotn 4% 200 ml OP healthE Dimethicone 4% Lotion

IVERMECTIN - Special Authority see SA1225 below - Retail pharmacy

✓ Stromectol

- 1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- 3) For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA1225 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scables hyperinfestation (Crusted/ Norwegian scables); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy: or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation. Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria: Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

microbiologist; and

- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy: or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — **(Other parasitic infections)** only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

PERMETHRIN

Crm 5%	75	30 g OP	Lyderm
Lyderm to be Sole Supply on 1 November 2020 Lotn 5%	99	30 ml OP	✓ A-Scables
A-Scabies to be Sole Supply on 1 November 2020			71 000.000
PHENOTHRIN			
Shampoo 0.5% 11.5	36	200 ml OP	✓ Parasidose

Psoriasis and Eczema Preparations

ACITRETIN – Special Authority see SA1476 below – Retail pharmacy			
Cap 10 mg	.17.86	60	Novatretin
Novatretin to be Sole Supply on 1 October 2020			
Cap 25 mg	.41.36	60	Novatretin
Novatretin to be Sole Supply on 1 October 2020			

⇒SA1476 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and

continued...

		DERM	ATOLOGICALS
	Subsidy (Manufacturer's Price)	Fully Subsidised Per	Brand or Generic Manufacturer
continued			
3 Either:			
3.1 Patient is female and has been counselled and und pregnancy and the applicant has ensured that the pregnancement of the treatment and that the patient treatment and for a period of two years after the condition. 3.2 Patient is male.	possibility of pregnar nt is informed that sl mpletion of the treat	ncy has been excluse he must not becon ment; or	uded prior to the ne pregnant during
Renewal from any relevant practitioner. Approvals valid for 1 year Either:	ar for applications m	eeting the following	g criteria:
 Patient is female and has been counselled and understand and the applicant has ensured that the possibility of pregna treatment and that the patient is informed that she must no years after the completion of the treatment; or Patient is male. 	ancy has been exclu	ided prior to the co	ommencement of the
BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL			
Foam spray 500 mcg with calcipotriol 50 mcg per g			instilar
Gel 500 mcg with calcipotriol 50 mcg per g		_	aivobet
Oint 500 mcg with calcipotriol 50 mcg per g	19.95	30 g OP ✓ <u>D</u>	<u>aivobet</u>
CALCIPOTRIOL Oint 50 mcg per g	40.00 1	20 g OP ✓ D	aivonex
COAL TAR			
Soln BP - Only in combination	36.25	200 ml ✓ <u>N</u>	lidwest .
 Up to 10% only in combination with a dermatologica With or without other dermatological galenicals. 	al base or proprietar	y Topical Corticos	teriod – Plain
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULP Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and			
allantoin crm 2.5%		75 g OP	
	(8.00)		gopsoryl TA
	3.43	30 g OP	• • •
	(4.35)	Ē	gopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR			
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97 2	25 g OP ✓ C	coco-Scalp
·	7.95	40 g OP ✓ C	coco-Scalp
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES	SCEIN - Only on a	prescription	•
Soln 2.3% with trolamine laurilsulfate and fluorescein sodium Pinetarsol to be Sole Supply on 1 November 2020	,	' '	inetarsol
SALICYLIC ACID			
Powder - Only in combination	18.88	250 g	lidwest SM
 Only in combination with a dermatological base or p With or without other dermatological galenicals. 	proprietary Topical C	Corticosteroid – Pla	ain or collodion flexible

SULPHUR

Precipitated - Only in combination......6.35 100 g ✓ Midwest

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteroid Plain
- 2) With or without other dermatological galenicals.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Scalp Preparations

BETAMETHASONE VALERATE * Scalp app 0.1%	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE		✓ Dermol
* Scalp app 0.05%		
KETOCONAZOLE	100 ml OP	
Shampoo 2%3.23	100 ml OP	Sebizole

- a) Maximum of 100 ml per prescription
- b) Only on a prescription
- c) Sebizole to be Sole Supply on 1 November 2020

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

SPF 50+

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 68

IMIQUIMOD

Crm 5%, 250 mg sachet......21.72

24

✓ Perrigo

PODOPHYLLOTOXIN

3.60 3.5 ml OP

✓ Condyline

- a) Maximum of 3.5 ml per prescription
- b) Only on a prescription

Other Skin Preparations

Antineoplastics

FLUOROURACIL SODIUM

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Contraceptives - Non-hormonal

Condoms

CO	NDOMS			
*	49 mm - Up to 144 dev available on a PSO	11.42	144	✓ Moments
	53 mm		10	✓ Moments
		11.64	144	✓ Moments
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
*	53 mm, 0.05 mm thickness	0.95	10	✓ Moments
-	, , , , , , , , , , , , , , , , , , , ,	11.42	144	✓ Moments
	a) Up to 60 dev available on a PSO			<u></u>
	b) Maximum of 60 dev per prescription			
*	53 mm, chocolate, brown	0.95	10	✓ Moments
-1-	oo min, onocolato, brown	11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO	11.04	1 -1 -1	Momento
	b) Maximum of 60 dev per prescription			
æ	53 mm, strawberry, red	0.05	10	✓ Moments
*	55 min, snawberry, red	11.64	144	✓ Moments
	a) The to CO downwish to an a DCO	11.04	144	wioments
	a) Up to 60 dev available on a PSO			
*	b) Maximum of 60 dev per prescription	0.07	10	✓ Moments
不	56 mm		10	✓ Moments
	\	11.64	144	woments
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO	4.00	40	40 111/111
*	56 mm, 0.05 mm thickness		12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
*	56 mm, 0.08 mm thickness		10	✓ <u>Moments</u>
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
*	56 mm, 0.08 mm thickness, red		10	✓ Moments
		11.64	144	✓ Moments
	 a) Up to 60 dev available on a PSO 			
	 b) Maximum of 60 dev per prescription 			
*	56 mm, chocolate	1.30	12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
*	56 mm, strawberry	1.30	12	✓ Gold Knight
		15.57	144	Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
*	60 mm - Up to 144 dev available on a PSO	14.87	144	✓ Shield XL

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

GENITO-URINARY SYSTEM

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ubsidised	Generic	
\$	Per	✓	Manufacturer	

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO
- ✓ Choice TT380 Short

TT380 Standard

✓ Choice Load 375

✓ Choice

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

Mercilon 28

a) Higher subsidy of \$13.80 per 84 tab with Special Authority see \$A0500 above

b) Up to 84 tab available on a PSO

Marvelon 28 (19.80)

a) Higher subsidy of \$13.80 per 84 tab with Special Authority see \$A0500 above

b) Up to 84 tab available on a PSO

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	9	Subsidised	Generic
	\$	Per	1	Manufacturer
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO		84	1	Microgynon 20 ED
	6.45	112		Femme-Tab ED
* Tab 50 mcg with levonorgestrel 125 mcg and 7 inert tab - U	Jp .			
to 84 tab available on a PSO	9.45	84	✓	Microgynon 50 ED
* Tab 30 mcg with levonorgestrel 150 mcg	6.62	63		• •
ů ů	(16.50)			Microgynon 30
a) Higher subsidy of \$15.00 per 63 tab with Special Aut	hority see SA0500 or	the pr	revious pa	age
b) Up to 63 tab available on a PSO	,		•	
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO		84	1	Levien ED
op to 112 tab available on a 1 oo	6.45	112		Femme-Tab ED
ETHINNI OFOTRADIOL MUTHINODETHIOTEDONE	0.10			Tommo Tab Eb
ETHINYLOESTRADIOL WITH NORETHISTERONE				
Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to			_	
84 tab available on a PSO		84	•	Brevinor 1/28
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - L				
to 84 tab available on a PSO	6.62	84	✓	Necon
			✓	Norimin
	8.83	112	✓	Brevinor 28

(Brevinor 28 Tab 35 mcg with norethisterone 500 mcg and 7 inert tab to be delisted 1 January 2021)

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

GENITO-URINARY SYSTEM

0	Subsidy Manufacturer's Price)	Ç.	Fully ubsidised	Brand or Generic
(i	\$	Per	Jusiuiseu 🗸	Manufacturer
LEVONORGESTREL				
* Tab 30 mcg - Up to 84 tab available on a PSO	16.50	84	✓ N	licrolut
	22.00	112	✓ N	licrolut
Subdermal implant (2 x 75 mg rods) – Up to 3 pack available on a PSOJadelle to be Sole Supply on 1 December 2020	106.92	1	✓ J	adelle
MEDROXYPROGESTERONE ACETATE Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PSI NORETHISTERONE	O7.98	1	✓ <u>D</u>	epo-Provera
Tab 350 mcg - Up to 84 tab available on a PSO	6.25	84	✓ N	loriday 28
Emergency Contraceptives				
LEVONORGESTREL				

LE'	V	O	N	0	R	G	Е	Sī	ΓF	3	ΕΙ	L

- ✓ Postinor-1
 - a) Maximum of 2 tab per prescription
 - b) Up to 5 tab available on a PSO c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

* Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs - Up to 168 tab available on a PSO......4.67 ✓ Ginet 168

Gynaecological Anti-infectives

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID)		
Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate			
0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator	r8.43	100 g OP	
	(24.00)		Aci-Jel
CLOTRIMAZOLE			
* Vaginal crm 1% with applicators	2.50	35 g OP	✓ Clomazol
* Vaginal crm 2% with applicators	3.00	20 g OP	✓ Clomazol
MICONAZOLE NITRATE			
* Vaginal crm 2% with applicator	6.89	40 g OP	✓ Micreme
Micreme to be Sole Supply on 1 November 2020			
NYSTATIN			
Vaginal crm 100,000 u per 5 g with applicator(s)	4.00	75 g OP	✓ Nilstat
Nilstat to be Sole Supply on 1 October 2020		ŭ	

Myometrial and Vaginal Hormone Preparations

ERGON	/FTF	INF	MAI	FATE

Inj 500 mcg per ml, 1 ml ampoule - Up to 5 inj available on a ✓ DBL Ergometrine

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's P \$	Price) Subs Per	Fully sidised	Brand or Generic Manufacturer
OESTRIOL				
* Crm 1 mg per g with applicator Ovestin to be Sole Supply on 1 October 2020	6.62	15 g OP	√ 0	vestin
* Pessaries 500 mcg Ovestin to be Sole Supply on 1 October 2020	6.86	15	√ 0	vestin
OXYTOCIN - Up to 5 inj available on a PSO				
Inj 5 iu per ml, 1 ml ampoule	3.98	5	✓ <u>0</u>	xytocin BNM
Inj 10 iu per ml, 1 ml ampoule	4.98	5	√ <u>0</u>	xytocin BNM
OXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj availa Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml		5	√ <u>S</u>	syntometrine

Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

✓ Smith BioMed Rapid 40 test OP **Pregnancy Test**

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 106

5-Alpha Reductase Inhibitors

FINASTERIDE - Special Authority see \$A0928 below - Retail pharmacy 100 ✓ Ricit

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy ✓ Tamsulosin-Rex

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

GENITO-URINARY SYSTEM

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
\$	Per	1	Manufacturer

Other Urinary Agents

OYVRI ITVNIN

* Tab 5 mg11.70	500	✓ Apo-Oxybutynin
* Oral liq 5 mg per 5 ml	473 ml	✓ Apo-Oxybutynin
POTASSIUM CITRATE		
Oral lig 3 mmol per ml - Special Authority see SA1083 below -		

200 ml OP ✓ Biomed Retail pharmacy......31.80

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE

* Grans eff 4 g sachets	2.22	28	✓ Ural
Ural to be Sole Supply on 1 October 2020			
SOLIFENACIN SUCCINATE			
Tab 5 mg	3.00	30	 Solifenacin Mylan
Tab 10 mg	5.50	30	✓ Solifenacin Mylan

Detection of Substances in Urine

ORTHO-TOLIDINE	
* Compound diagnostic sticks	

~	Compound diagnostic sticks	30 1031 01	
	(8.25)		Hemastix

TETRABROMOPHENOL

*	Blue diagnostic strips7.02	100 test OP	
	(13.92)		Albustix

Obstetric Preparations

Antiprogesterones

MIFEPRISTONE

Subsidised on a PSO only if from a Family Planning New Zealand Clinic or an abortion service provider with a DHB contract and the PSO is endorsed with the name of the institution for which the PSO is required.

7 50

50 tost OP

✓ Mifegyne

- a) Up to 15 tab available on a PSO
- b) Only on a PSO

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	/	Manufacturer	

Calcium Homeostasis

CA	ויו	11(1	IN	IN

CINACALCET - Special Authority see SA1618 below - Retail pharmacy

Tab 30 mg − Wastage claimable......210.30 28 ✓ Sensipar

⇒SA1618 Special Authority for Subsidy

Initial application only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial − Special Authority see SA1687 below −
Retail pharmacy......38.03 1

✓ Zoledronic acid
Mylan

⇒SA1687 Special Authority for Subsidy

Initial application — **(bone metastases)** only from an oncologist, haematologist or palliative care specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
 - 2.1 Patient has bone metastases or involvement; and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
 - 3.1 Patient has bone metastases or involvement; and
 - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer) only from an oncologist or medical practitioner on the recommendation of a oncologist. Approvals valid for 2 years for applications meeting the following criteria:
All of the following:

Subsid	dy Full	/ Brand or
(Manufacture	r's Price) Subsidise	d Generic
\$	Per 💌	Manufacturer

continued...

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.

Corticosteroids and Related Agents for Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE

* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	5	Celestone
		Chronodose
BEXAMETHASONE ★ Tab 0.5 mg − Up to 60 tab available on a PSO	30 30 25 ml OP	✓ <u>Dexmethsone</u> ✓ <u>Dexmethsone</u> ✓ Biomed
DEXAMETHASONE PHOSPHATE		
Dexamethasone phosphate injection will not be funded for oral use.		
* Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO9.25	10	✓ <u>Dexamethasone</u> <u>Phosphate</u> <u>Panpharma</u>
* Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO16.37	10	✓ <u>Dexamethasone</u> <u>Phosphate</u> <u>Panpharma</u>
FLUDROCORTISONE ACETATE		
* Tab 100 mcg14.32	100	✓ Florinef
HYDROCORTISONE		
* Tab 5 mg8.10	100	✓ <u>Douglas</u>
* Tab 20 mg	100	✓ <u>Douglas</u>
 Inj 100 mg vial	1	✓ Solu-Cortef
METHYLPREDNISOLONE		
* Tab 4 mg112.00	100	✓ <u>Medrol</u>
* Tab 100 mg194.00	20	✓ <u>Medrol</u>
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)		
Inj 40 mg vial18.90	1	✓ <u>Solu-Medrol-Act-</u> <u>O-Vial</u>
Inj 125 mg vial28.90	1	✓ <u>Solu-Medrol-Act-O-Vial</u>
Inj 500 mg vial22.78	1	✓ <u>Solu-Medrol-Act-O-Vial</u>
Inj 1 g vial27.83	1	✓ Solu-Medrol
METHYLPREDNISOLONE ACETATE	'	- Join-Michiol
Inj 40 mg per ml, 1 ml vial44.40	5	✓ Depo-Medrol
PREDNISOLONE	3	- Depo-medioi
* Oral liq 5 mg per ml – Up to 30 ml available on a PSO	30 ml OP	✓ <u>Redipred</u>

	Subsidy		Fully	
	(Manufacturer's Price)	_	Subsidised	
	\$	Per		Manufacturer
PREDNISONE				
* Tab 1 mg	10.68	500	1	Apo-Prednisone
* Tab 2.5 mg		500	1	Apo-Prednisone
* Tab 5 mg - Up to 30 tab available on a PSO		500	1	Apo-Prednisone
* Tab 20 mg - Up to 30 tab available on a PSO	29.03	500	✓	Apo-Prednisone
TETRACOSACTRIN				
* Inj 250 mcg per ml, 1 ml ampoule	75.00	1	1	UK Synacthen S29
, , · · · · g				AU Synacthen
				Synacthen
* Inj 1 mg per ml, 1 ml ampoule	690.00	1	✓	Synacthen Depot
, 01			1	Synacthene
				Retard S29
TRIAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule	20.80	5	1	Kenacort-A 10
	26.62		1	Adcortyl S29
Kenacort-A 10 to be Sole Supply on 1 November 2020				,
Inj 40 mg per ml, 1 ml ampoule	11.30	1	1	Triaver \$29
, , , ,	51.10	5	1	Kenacort-A 40
	70.62		1	Kenalog \$29
Kenacort-A 40 to be Sole Supply on 1 November 2020				

Sex Hormones Non Contraceptive

Androgen Agonists and Antagonists

CVDDOTEDONE ACETATE

CYPROTERONE AGETATE			
Tab 50 mg	13.17	50	✓ Siterone
Tab 100 mg	26.75	50	✓ Siterone
TESTOSTERONE			
Patch 5 mg per day	90.00	30	✓ Androderm
TESTOSTERONE CIPIONATE			
Inj 100 mg per ml, 10 ml vial	76.50	1	✓ Depo-Testosterone
TESTOSTERONE ESTERS			•
Inj 250 mg per ml, 1 ml	12.98	1	✓ Sustanon Ampoules
TESTOSTERONE UNDECANOATE			,
Cap 40 mg	21.00	60	✓ Andriol Testocaps
Inj 250 mg per ml, 4 ml vial		1	✓ Reandron 1000
, ===,			

Hormone Replacement Therapy - Systemic

Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

		Subsidy (Manufacturer's Pr		Fully sidised	Brand or Generic
_		\$	Per		Manufacturer
0	estrogens				
	STRADIOL – See prescribing guideline on the previous page				
*	Tab 1 mg		28 OP	_	'atuafaua
*	Tab 2 mg	(11.10)	28 OP		strofem
~	Tab 2 mg	(11.10)	20 01	F	Strofem
*	Patch 100 mcg per 24 hours	, ,	4		limara
	a) No more than 1 patch per week				
	b) Only on a prescription				
*	Patch 50 mcg per 24 hours	7.04	4	✓ (Climara
	a) No more than 1 patch per week				
	b) Only on a prescription		_	, .	
	Patch 25 mcg per day	6.12	8	✓ E	stradot
	a) No more than 2 patch per week				
	b) Only on a prescription Patch 50 mcg per day	7.04	8	√	stradot 50 mcg
	a) No more than 2 patch per week	7.04	O	•	Strauot 50 mcg
	b) Only on a prescription				
	Patch 75 mcg per day	7.91	8	√ E	stradot
	a) No more than 2 patch per week				
	b) Only on a prescription				
	Patch 100 mcg per day	7.91	8	√ E	stradot
	a) No more than 2 patch per week				
	b) Only on a prescription				
	STRADIOL VALERATE - See prescribing guideline on the pr				
	Tab 1 mg		84		rogynova
*	Tab 2 mg	12.36	84	✓ <u>F</u>	Progynova
	STROGENS - See prescribing guideline on the previous page				
*	Conjugated, equine tab 300 mcg		28	_	
×	Conjugated aguing tab 605 mag	(13.50)	28	۲	remarin
*	Conjugated, equine tab 625 mcg	(13.50)	20	p	Premarin
		(10.50)		'	Temami
P	rogestogens				
ИE	DROXYPROGESTERONE ACETATE - See prescribing guid	eline on the prev	vious page		
*	Tab 2.5 mg	3.75	30		rovera
	Tab 5 mg		100	_	rovera
*	Tab 10 mg	7.15	30	✓ P	rovera
P	rogestogen and Oestrogen Combined Prepara	tions			
	STRADIOL WITH NORETHISTERONE - See prescribing gui		vious page		
	Tab 1 mg with 0.5 mg norethisterone acetate	5.40	28 OP		
٠		(18.10)		K	(liovance
*	Tab 2 mg with 1 mg norethisterone acetate		28 OP		Warrant.
	Tab O many with down manythiates are a sale to (40) and 10	(18.10)		K	liogest
*	Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	E 40	20 OD		
	oestradiol tab (12) and 1 mg oestradiol tab (6)	5.40	28 OP	т	risequens
		(10.10)			nooquono

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Other Oestrogen Preparations				
ETHINYLOESTRADIOL * Tab 10 mcg	17.60	100	✓ <u>N</u>	IZ Medical and Scientific
OESTRIOL * Tab 2 mg	7.00	30	√ <u>(</u>	Ovestin
Other Progestogen Preparations				
LEVONORGESTREL * Intra-uterine device 52 mg * Intra-uterine device 13.5 mg MEDROXYPROGESTERONE ACETATE		1		<u>Airena</u> laydess
Tab 100 mg	101.00	100	√ F	Provera HD
NORETHISTERONE * Tab 5 mg – Up to 30 tab available on a PSO PROGESTERONE	18.29	100	√ <u>F</u>	Primolut N
Cap 100 mg - Special Authority see SA1609 below - Retail pharmacy	16.50	30	√ (Jtrogestan

SA1609 Special Authority for Subsidy

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 For the prevention of pre-term labour*; and
- 2 Either:
 - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 For the prevention of pre-term labour*: and
- 2 Treatment is required for second or subsequent pregnancy; and
- 3 Fither:
 - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are unapproved indications.

Thyroid and Antithyroid Agents

CARBIMAZOLE		
* Tab 5 mg10.80	100	✓ AFT
		Carbimazole S29
		✓ Neo-Mercazole

	Subsidy		Fully	Brand or
	(Manufacturer's Price	e)	Subsidised	Generic
	\$	Per	•	Manufacturer
LEVOTHYROXINE				
* Tab 25 mcg	3.89	90	✓	Synthroid
* Tab 50 mcg		28	✓	Mercury Pharma
·	4.05	90	✓	Synthroid
	64.28	1,000	✓	Eltroxin
* Tab 100 mcg	1.78	28	✓	Mercury Pharma
•	4.21	90	✓	Synthroid
	66.78	1,000	✓	Eltroxin
PROPYLTHIOURACIL – Special Authority see SA1199 belo Propylthiouracil is not recommended for patients under the treatments are contraindicated.		ss the p	atient is p	regnant and other
Tab 50 mg	35.00	100	1	PTU S29

⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) - Special Authority see SA1629 below - Re	etail pharmacy	
*	Inj 5 mg cartridge34.8	38 1	Omnitrope
*	Inj 10 mg cartridge69.7	75 1	✓ Omnitrope
*	Inj 15 mg cartridge104.6	63 1	✓ Omnitrope

⇒SA1629 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and</p>
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — **(short stature without growth hormone deficiency)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	

continued...

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria;
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight eximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy (Manufacturer's Pri	ce)	Fully Subsidised	Brand or Generic	
\$	Per	•	Manufacturer	

continued...

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
GnRH Analogues				
GOSERELIN Implant 3.6 mg, syringeImplant 10.8 mg, syringe		1	-	Zoladex Zoladex
LEUPRORELIN Additional subsidy by endorsement where the patient is a chi goserelin and the prescription is endorsed accordingly. Inj 3.75 mg prefilled dual chamber syringe — Higher subsidy	ld or adolescent and i	s un	able to toler	ate administration of
\$221.60 per 1 inj with Endorsement	66.48 (221.60)	1	L	_ucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe – Higher subsidy of \$591.68 per 1 inj with Endorsement		1	l	Lucrin Depot 3-month

Vasopressin Agonists

DESMOPRESSIN	∧ ∩ ET ∧ T E
DESIMORRESSIN	AUFIAIF

	Tab 100 mcg - Special Authority see SA1401 below - Retail pharmacy	25.00	30	✓ Minirin
^	Tab 200 mcg - Special Authority see SA1401 below - Retail pharmacy	54.45 39.03	30 2.5 ml OP 6 ml OP	✓ Minirin ✓ Minirin ✓ Desmopressin- PH&T
	Desmopressin-PH&T to be Sole Supply on 1 November 2020 Inj 4 mcg per ml, 1 ml – Special Authority see SA1401 below – Retail pharmacy		10	✓ Minirin

⇒SA1401 Special Authority for Subsidy

Initial application — (Desmopressin tablets for Nocturnal enuresis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has primary nocturnal enuresis; and
- 2 The nasal forms of desmopressin are contraindicated; and
- 3 An enuresis alarm is contraindicated.

Initial application — (Desmopressin tablets for Diabetes insipidus) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has cranial diabetes insipidus; and
- 2 The nasal forms of desmopressin are contraindicated.

Renewal — (Desmopressin tablets) from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from the treatment.

Initial application — (Desmopressin injection) only from a relevant specialist. Approvals valid for 2 years where the patient cannot use desmopressin nasal spray or nasal drops.

Renewal — (Desmopressin injection) only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Other Endocrine Agents

CABERGOLINE

⇒SA1370 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 pathological hyperprolactinemia; or
- 2 acromegaly*.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

CLC	MIF	ENE	CITRA	ATE.

Tab 50 mg29).84	10	Mylan Clomiphen S29
DANAZOL			
Cap 100 mg19	.13 2	28	Mylan S29
Cap 200 mg97	'.83 1	00	Azol
METYRAPONE			
Cap 250 mg558	3.00	50	Metopirone
Metopirone to be Sole Supply on 1 November 2020			•

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	1	Manufacturer	

Anthelmintics

ALBENDAZOLE -	Special Authority see SA1318 below – Retail pharmacy		
Tab 400 mg	469.20	60	✓ Eskazole S29

⇒SA1318 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

MEBENDAZOLE - Only on a prescription

Tab 100 mg	24.19	24	De-Worm
Oral lig 100 mg per 5 ml		15 ml	
	(7.17)		Vermox
PRAZIQUANTEL			
Tah 600 mg	68.00	8	✓ Riltricide

Antibacterials

CEFACI OR MONOHYDRATE

- a) For topical antibacterials, refer to DERMATOLOGICALS, page 61
- b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 241

Cephalosporins and Cephamycins

CEFACEON MONORITEDRATE			
Cap 250 mg	24.70	100	✓ Ranbaxy-Cefaclor
Grans for oral lig 125 mg per 5 ml - Wastage claimable.	3.53	100 ml	✓ Ranbaxy-Cefaclor
	4.33		✓ Keflor
(Keflor Grans for oral liq 125 mg per 5 ml to be delisted 1 Dec	cember 2020)		
CEFALEXIN			
Cap 250 mg	3.33	20	 Cephalexin ABM
			✓ Ibilex S29
Cap 500 mg	3.95	20	Cephalexin ABM
Grans for oral liq 25 mg per ml - Wastage claimable		100 ml	 Cefalexin Sandoz
Grans for oral lig 50 mg per ml - Wastage claimable		100 ml	✓ Cefalexin Sandoz
(Ibilex S29 Cap 250 mg to be delisted 1 February 2021)			
CEFAZOLIN - Subsidy by endorsement			
Only if prescribed for dialysis or cellulitis in accordance waccordingly.	rith a DHB approved p	protocol and t	he prescription is endorsed
Inj 500 mg vial	3.39	5	✓ AFT
AFT to be Sole Supply on 1 November 2020		3	
Inj 1 g vial	3 49	5	✓ AFT
AFT to be Sole Supply on 1 November 2020		J	• • •
711 1 to be cold capply on 1 Hoveliber 2020			

CEFTRIAXONE - Subsidy by endorsement

- a) Up to 10 inj available on a PSO
- b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of pelvic inflammatory disease, or the treatment of suspected meningococcal disease, and the prescription or PSO is endorsed accordingly.

Inj 500 mg vial	0.89	1	✓ <u>Ceftriaxone-AFT</u>
lnj 1 g vial	3.99	5	✓ Ceftriaxone-AFT

	Subsidy (Manufacturer's Price)	Subsic	Fully lised	Brand or Generic
	\$	Per	1	Manufacturer
CEFUROXIME AXETIL – Subsidy by endorsement				
Only if prescribed for prophylaxis of endocarditis and the pres	scription is endorsed	accordingly		
Tab 250 mg	45.93	50	✓ Zi	innat

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special

Authority.		
Tab 250 mg8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO	2	✓ Apo-Azithromycin
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage		
claimable	15 ml	✓ Zithromax

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN – Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 on the next page Tab 250 mg3.98 14 Apo-Clarithromycin

Grans for oral liq 250 mg per 5 ml − Wastage claimable......192.00 50 ml ✓ Klacid

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated.

Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician.

Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN (AS LACTOBIONATE)

Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	E-Mycin
a) Up to 20 tab available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	E-Mycin
 a) Up to 300 ml available on a PSO 			
b) Up to 2 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	E-Mycin
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
ERYTHROMYCIN STEARATE			
Tab 250 mg - Up to 30 tab available on a PSO	14.95	100	
	(22.29)		ERA
Tab 500 mg		100	
	(44.58)		ERA
ROXITHROMYCIN			
Tab disp 50 mg	8.29	10	✓ Rulide D
Restricted to children under 12 years of age.			
Tab 150 mg	8.28	50	✓ Arrow-
			Roxithromycin
Tab 300 mg	16.33	50	✓ Arrow-
			Roxithromycin

	Subsidy (Manufacturer's Pr \$	ice) Subs	Fully Brand or sidised Generic Manufacturer
Penicillins			
AMOXICILLIN			
Cap 250 mg	22.50	500	✓ <u>Alphamox</u>
a) Up to 30 cap available on a PSO			
b) Up to 10 x the maximum PSO quantity for RFPP			
Cap 500 mg	36.98	500	✓ <u>Alphamox</u>
a) Up to 30 cap available on a PSO			
b) Up to 10 x the maximum PSO quantity for RFPP	4.40	400 1	4 4 1 1 1 1 1 1 1 1 1 1
Grans for oral liq 125 mg per 5 ml	1.40	100 ml	✓ Alphamox 125
a) Up to 200 ml available on a PSO			
b) Wastage claimablec) Alphamox 125 to be Sole Supply on 1 November 202	0		
Grans for oral lig 250 mg per 5 ml		100 ml	✓ Alphamox 250
a) Up to 300 ml available on a PSO	1.73	100 1111	Alphaniox 250
b) Up to 10 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
d) Alphamox 250 to be Sole Supply on 1 November 202	0		
Inj 250 mg vial		10	✓ Ibiamox
Inj 500 mg vial	12.41	10	✓ Ibiamox
Inj 1 g vial - Up to 5 inj available on a PSO	17.29	10	✓ Ibiamox
AMOXICILLIN WITH CLAVULANIC ACID			
Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab			
available on a PSO	1.88	20	✓ Augmentin
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 r	ng		•
per ml	5.00	100 ml	✓ Augmentin
a) Up to 200 ml available on a PSO			-
b) Wastage claimable			
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 r	ng		
per ml - Up to 200 ml available on a PSO	2.20	100 ml OP	✓ Curam
BENZATHINE BENZYLPENICILLIN			
Inj 900 mg (1.2 million units) in 2.3 ml syringe - Up to 5 inj			
available on a PSO	344.93	10	✓ Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]			
Inj 600 mg (1 million units) vial – Up to 5 inj available on a PS	SO 11.09	10	✓ Sandoz
, ,	25.88	25	✓ Pan-Penicillin G
			Sodium S29

Sandoz to be Sole Supply on 1 November 2020

(Pan-Penicillin G Sodium S29 Inj 600 mg (1 million units) vial to be delisted 1 November 2020)

	Subsidy (Manufacturer's Price \$		ully Brand or sed Generic Manufacturer
FLUCLOXACILLIN	Ψ	1 01	Waltalactarci
Cap 250 mg - Up to 30 cap available on a PSO	16.83	250	✓ Staphlex
Cap 500 mg - Up to 30 cap available on a PSO			✓ Staphlex
Grans for oral liq 25 mg per ml			✓ AFT
a) Up to 200 ml available on a PSO		100 1111	- <u>/</u>
b) Wastage claimable			
Grans for oral liq 50 mg per ml	3.68	100 ml	✓ AFT
a) Up to 200 ml available on a PSO		100 1111	- <u>/ /</u>
b) Wastage claimable			
Inj 250 mg vial	9.00	10	✓ Flucloxin
Inj 500 mg vial			✓ Flucioxin
Inj 1 g vial – Up to 5 inj available on a PSO	5.70	. •	✓ Flucil
Flucil to be Sole Supply on 1 November 2020		3	· I Iucii

PHENOXYMETHYLPENICILLIN (PENICILLIN V)	0.50	F0	✓ Ollinaina VIV
Cap 250 mg - Up to 30 cap available on a PSO			✓ <u>Cilicaine VK</u>
Cap 500 mg	4.26	50	✓ Cilicaine VK
a) Up to 20 cap available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 125 mg per 5 ml	2.99	100 ml	✓ <u>AFT</u>
a) Up to 200 ml available on a PSO			
b) Wastage claimable	0.00	400	.
Grans for oral liq 250 mg per 5 ml	3.99	100 ml	✓ <u>AFT</u>
a) Up to 300 ml available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
PROCAINE PENICILLIN			
Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO	123.50	5	✓ Cilicaine
Tetracyclines			
OOXYCYCLINE			
★ Tab 100 mg – Up to 30 tab available on a PSO	64.43	500	✓ Doxine
IINOCYCLINE HYDROCHLORIDE			
Tab 50 mg - Additional subsidy by Special Authority see			
SA1355 below – Retail pharmacy	5.79	60	
CATIOGO DOION TIOIGII PHAITHAU	(12.05)	00	Mino-tabs
		100	mino tabo
F Cup 100 mg	(52.04)	100	Minomycin
CA10EE Chasial Authority for Manufacturers Briss	(02.04)		······oiiiyoiii
⇒SA1355 Special Authority for Manufacturers Price	d without fouthouse	awal unlass =	stified where the noticet be
nitial application from any relevant practitioner. Approvals valid	u williout iurtrier ren	ewai uilless N	ninea where the patient ha
OSACEA.	Dotoil pho		
ETRACYCLINE – Special Authority see SA1332 on the next pa	•	•	/ A
Tab 250 mg			Accord \$29
Cap 500 mg	46.00	30	✓ Tetracyclin
			Wolff S29

28

✓ Cinflox

Subsidy (Manufacturer's Price)	Ç,	Fully ubsidised	Brand or Generic	
(Manuacturer's Frice)	O.	ibsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 61

CIPROFLOXACIN

Recommended for patients with any of the following:

i) microbiologically confirmed and clinically significant pseudomonas infection; or

- ii) prostatitis: or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Cipflox to be Sole Supply on 1 November 2020	20	Cipliox	
Tab 500 mg - Up to 5 tab available on a PSO	28	✓ Cipflox	
Tab 750 mg	28	✓ Cipflox	
Cipflox to be Sole Supply on 1 November 2020			
CLINDAMYCIN			
Cap hydrochloride 150 mg4.61	24	✓ Dalacin C	
Inj phosphate 150 mg per ml, 4 ml ampoule39.00	10	✓ Dalacin C	
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist - Subsidy by endors	sement		
Only if prescribed for dialysis or cystic fibrosis patient and the prescription is	endorsed acc	ordingly.	
Inj 150 mg65.00	1	✓ Colistin-Link	
GENTAMICIN SULPHATE			
Inj 10 mg per ml, 1 ml ampoule – Subsidy by endorsement25.00			
Only if prescribed for a dialysis or cystic fibrosis patient or complicated un endorsed accordingly.	inary tract inf	ection and the prescription is	;
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement182.00	10	✓ Teligent S29	
Only if prescribed for a dialysis or cystic fibrosis patient or complicated un endorsed accordingly.	inary tract inf	ection and the prescription is	;
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement17.50	10	✓ Pfizer	
87.50	50	✓ Pfizer	
Only if prescribed for a dialysis or cystic fibrosis patient or complicated un endorsed accordingly.	inary tract inf	ection and the prescription is	;
MOXIFLOXACIN – Special Authority see SA1740 below – Retail pharmacy No patient co-payment payable			
Tab 400 mg42.00	5	✓ Avelox	
Avelox to be Sole Supply on 1 December 2020			

⇒SA1740 Special Authority for Subsidy

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1 Both:
 - 1.1 Active tuberculosis*; and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Either:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy

⇒SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Fither:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharmacy

⇒SA1328 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

ıı,	NFECTIONS - A	JEN	13 701	1 SYSTEMIC USE
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
SODIUM FUSIDATE [FUSIDIC ACID]				
Tab 250 mg		12	•	Fucidin
SULFADIAZINE SODIUM – Special Authority see SA1331 below			_	
Tab 500 mg	543.20	56	/	Wockhardt S29
Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months	r a period of 3 months		nless notif	ied for applications meeting
	or ago.			
TOBRAMYCIN Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement	15.00	5	_	Tobramycin Mylan
Only if prescribed for dialysis or cystic fibrosis patient an Solution for inhalation 60 mg per ml, 5 ml – Subsidy by				
endorsement	2,200.00 5	6 dos	se 🗸	TOBI
a) Wastage claimable				
b) Only if prescribed for a cystic fibrosis patient and the	prescription is endors	sed a	ccordingly	
TRIMETHOPRIM				
* Tab 300 mg - Up to 30 tab available on a PSO	16.50	50	/	<u>TMP</u>
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX	AZOLE]			
* Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - U	Jp			
to 30 tab available on a PSO		500	/	Trisul
* Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to 200 i available on a PSO		100 m	ıl 🗸	Deprim
VANCOMYCIN – Subsidy by endorsement				
Only if prescribed for a dialysis or cystic fibrosis patient or for			is or for tre	eatment of Clostridium
difficile following metronidazole failure and the prescription is	•	•		
Inj 500 mg vial	2.35	1	•	Mylan
Mylan to be Sole Supply on 1 October 2020				
Antifungals				
a) For topical antifungals refer to DERMATOLOGICALS, page 67	2			
b) For topical antifungals refer to GENITO URINARY, page 74	-			
FLUCONAZOLE				
Cap 50 mg	2.75	28	1	Mylan
Mylan to be Sole Supply on 1 November 2020				•
Cap 150 mg	0.65	1	✓	Mylan
Mylan to be Sole Supply on 1 November 2020				
Cap 200 mg		28	•	Mylan
Powder for oral suspension 10 mg per ml — Special Authority		2E		Diffuson COC coc
see SA1359 on the next page – Retail pharmacy	34.56 98.50	35 m		Diflucan S29 S29 Diflucan
Wastage claimable	50.50		•	Dinavaii
. radiago dallilabio				

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOLE

Cap 100 mg	4.27	15	✓ <u>Itrazole</u>
Oral liq 10 mg per ml - Special Authority see SA1322 below -			
Retail pharmacy14	1.80	150 ml OP	✓ Sporanox

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE

Tab 200 mg - PCT	.CBS	30	✓ Link Healthcare S29
			✓ Nizoral S29
		100	✓ Strides Shasun S29
NYSTATIN			
Tab 500,000 u	.14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	.12.81	50	
	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 on the next page -	Retail phar	macy	
Tab modified-release 100 mg	369.86	24	✓ Noxafil
Oral liq 40 mg per ml7	761.13	105 ml OP	✓ Noxafil

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE

*	Tab 250 mg	1.33	14	✓ Deolate
۷O	RICONAZOLE - Special Authority see SA1273 below - Retail pharm	acy		
	Tab 50 mg9	1.00	56	✓ Vttack
	Tab 200 mg35	0.00	56	✓ Vttack
	Powder for oral suspension 40 mg per ml - Wastage			
	claimable1,43	7.00	70 ml	✓ Vfend

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis: or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Subsidy	Ful	ly Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	 Manufacturer 	

Antimalarials

PRIMAQUINE - Special Authority see SA1684 below - Retail pharmacy

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Antiparasitics

Antiprotozoals

\cap	IININE	SHI	DH7.	ΓF

Antitrichomonal Agents

METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl
Metrogyl to be Sole Supply on 1 December 2020			
Tab 400 mg - Up to 15 tab available on a PSO	5.23	21	✓ Metrogyl
Metrogyl to be Sole Supply on 1 December 2020			
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg	24.48	10	✓ Flagyl
ORNIDAZOLE			
Tab 500 mg	32.95	10	Arrow-Ornidazole

Antituberculotics and Antileprotics

Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status.

CLOFAZIMINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist.

CYCLOSERINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician.
- Cap 250 mg......344.00 60 **✓ Cyclorin** ©29

'	NFECTIONS - AC	JENIS	S FUR SYSTEMIC USE
	Subsidy (Manufacturer's Price) \$	Su Per	Fully Brand or ubsidised Generic ✓ Manufacturer
DAPSONE – Retail pharmacy-Specialist			
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat dermatologist Tab 25 mg		isease p	ohysician, clinical microbiologist or Dapsone
Tab 100 mg		100	✓ Dapsone
ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Specialis	st		
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician	ion of, an infectious di	isease p	physician, clinical microbiologist or
Tab 100 mg	85.73	100	✓ EMB Fatol S29
Tab 400 mg	49.34	56	✓ Myambutol S29
ISONIAZID - Retail pharmacy-Specialist			
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat microbiologist, dermatologist or public health physician 	ion of, an internal med	dicine ph	hysician, paediatrician, clinical
* Tab 100 mg	22.00	100	✓ <u>PSM</u>
ISONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist			
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat microbiologist, dermatologist or public health physician 		dicine ph	hysician, paediatrician, clinical
* Tab 100 mg with rifampicin 150 mg		100	Rifinah
* Tab 150 mg with rifampicin 300 mg	170.60	100	✓ <u>Rifinah</u>
PARA-AMINO SALICYLIC ACID – Retail pharmacy-Specialist			
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician 	ion of, an infectious di	isease s	specialist, clinical microbiologist or
Grans for oral liq 4 g sachet	280.00	30	✓ Paser S29
PROTIONAMIDE - Retail pharmacy-Specialist			
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician 		isease s	specialist, clinical microbiologist or
Tab 250 mg	305.00	100	✓ Peteha S29
PYRAZINAMIDE - Retail pharmacy-Specialist			
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician 		isease p	physician, clinical microbiologist or
* Tab 500 mg	59.00	100	AFT-Pyrazinamide
RIFABUTIN - Retail pharmacy-Specialist			
No patient co-payment payable	to a strong to to all the		ala antata a managanta a antanta d
 b) Prescriptions must be written by, or on the recommendat gastroenterologist Cap 150 mg 		isease p	
本 Cap 130 IIIy	288.75	30	✓ Mycobutin

	· · · · · · · · · · · · · · · · · · ·				
	Sub (Manufactu	,	,	Fully Brand or Subsidised Generic	Generic
		3	Per	/	Manufacturer
RII	RIFAMPICIN – Subsidy by endorsement				
	a) No patient co-payment payable				
	b) For confirmed recurrent Staphylococcus aureus infection in combina	tion with	other effe	ctive anti	-staphylococcal
	antimicrobial based on susceptibilities and the prescription is endors	ed acco	rdingly: car	be waiv	red by endorsement -
	Retail pharmacy - Specialist. Specialist must be an internal medicin				
	paediatrician, or public health physician.		,		
*	* Can 150 mg	.	100	√ B	ifadin

*	Cap 150 mg	58.54	100	Kıtadın
	Rifadin to be Sole Supply on 1 November 2020			
*	Cap 300 mg	122.06	100	Rifadin
	Rifadin to be Sole Supply on 1 November 2020			
*	Oral liq 100 mg per 5 ml	12.60	60 ml	Rifadin
	Rifadin to be Sole Supply on 1 November 2020			

Antivirals

For eye preparations refer to Eye Preparations, Anti-Infective Preparations, page 241

Hepatitis B Treatment

⇒SA0829 Special Authority for Subsidy

Initial application only from a gastroenterologist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg+); and Documented resistance to lamivudine, defined as:
- 2 Patient has raised serum ALT (> 1 x ULN); and
- 3 Patient has HBV DNA greater than 100,000 copies per mL, or viral load 10 fold or higher over nadir; and
- 4 Detection of M204I or M204V mutation; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 Patient is cirrhotic: and
 - 5.1.2 adefovir dipivoxil to be used in combination with lamivudine; or
 - 5.2 Both:
 - 5.2.1 Patient is not cirrhotic; and
 - 5.2.2 adefovir dipivoxil to be used as monotherapy.

Renewal only from a gastroenterologist or infectious disease specialist. Approvals valid for 2 years where in the opinion of the treating physician, treatment remains appropriate and patient is benefiting from treatment.

Notes: Lamivudine should be added to adefovir dipivoxil if a patient develops documented resistance to adefovir dipivoxil, defined as:

- i) raised serum ALT (> 1 × ULN); and
- ii) HBV DNA greater than 100,000 copies per mL, or viral load 10 fold or higher over nadir; and
- iii) Detection of N236T or A181T/V mutation.

Adefovir dipivoxil should be stopped 6 months following HBeAg seroconversion for patients who were HBeAg+ prior to commencing adefovir dipivoxil.

The recommended dose of adefovir dipivoxil is no more than 10mg daily.

In patients with renal insufficiency adefovir dipivoxil dose should be reduced in accordance with the datasheet guidelines. Adefovir dipivoxil should be avoided in pregnant women and children.

ENTECAVIR

	Subsidy (Manufacturer's Prio \$	e) Per	Fully Subsidised	Generic
LAMIVUDINE – Special Authority see SA1685 below – Retail phr Tab 100 mg	•	28	1	Zetlam
Zetlam to be Sole Supply on 1 November 2020 Oral liq 5 mg per ml	270.00	240 ml	OP 🗸	Zeffix

⇒SA1685 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B. TENOFOVIR DISOPROXIL

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA1651., page 104

*	Tab 245 mg (300.6 mg as a succinate)		30	•	Tenofovir Disoproxil
					Teva

Herpesvirus Treatments			
ACICLOVIR			
* Tab dispersible 200 mg	1.60	25	✓ <u>Lovir</u>
* Tab dispersible 400 mg	5.38	56	✓ Lovir
* Tab dispersible 800 mg	5.98	35	✓ <u>Lovir</u>
VALACICLOVIR			
Tab 500 mg	5.75	30	✓ Vaclovir
Tab 1,000 mg	11.35	30	✓ Vaclovir
VALGANCICLOVIR - Special Authority see SA1404 below -	Retail pharmacy		
Tab 450 mg	225.00	60	✓ Valganciclovir
			Mylan

⇒SA1404 Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
- 2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months): and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Patient has undergone a lung transplant; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

- Both:
 - 1 Patient is immunocompromised; and
 - 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via PHARMAC's approved direct distribution supply. Further details can be found on

PHARMAC's website https://www.pharmac.govt.nz/hepatitis-c-treatments

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz/hepatitis-c-treatments or:

The Coordinator, Hepatitis C Treatment Panel

PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

Fully

Subsidy (Manufacturer's Price) Subsidised Per

Brand or Generic Manufacturer

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement: can be waived by Special Authority see SA1904

below Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabline with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 104 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the PHARMAC website.

Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a

Teva

⇒SA1904 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment: and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men: and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment: and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks: and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and

Subsidy	Full	/ Brand or
(Manufacturer's	s Price) Subsidise	d Generic
\$	Per 💌	Manufacturer

continued...

- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Antiretrovirals

⇒SA1651 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	ıbsidised	Generic
 \$	Per	✓	Manufacturer

continued...

prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria: Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ – Special Authority see SA1651 o	n the previous page – Retail pharn	nacy	
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
ETRAVIRINE - Special Authority see SA1651	on the previous page - Retail phar	macy	
Tab 200 mg	770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA1651	on the previous page – Retail phar	macy	
Tab 200 mg	60.00	60	✓ <u>Nevirapine</u>
			<u>Alphapharm</u>
Oral suspension 10 mg per ml	203.55	240 ml	✓ Viramune
			Suspension

Nucleosides Reverse Transcriptase Inhibitors

ABACAVIR SULPHATE – Special Authority see SA1651 on the p Tab 300 mg Oral lig 20 mg per ml	180.00	Retail pharma 60 240 ml OP	cy ✓ <u>Ziagen</u> ✓ Ziagen
			. 3
ABACAVIR SULPHATE WITH LAMIVUDINE - Special Authority			• , ,
Note: abacavir with lamivudine (combination tablets) counts a	as two anti-retro	oviral medication	ons for the purposes of the
anti-retroviral Special Authority.			
Tab 600 mg with lamivudine 300 mg	63.00	30	✓ Kivexa
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPR	OXII - Specia	I Authority see	SA1651 on the previous page
Retail pharmacy	O72 Op 00.0		orridor dirano providuo pas
Note: Efavirenz with emtricitabine and tenofovir disoproxil co	unte ae throp a	nti_ratroviral me	adications for the nurnoses o
Note. Liaviferiz with emitherabile and teriolovii disoproxii co	unio ao imee a	iiu-ieuoviiai iiie	fulcations for the purposes of

of the anti-retroviral Special Authority

Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil 245 mg (300 mg as a maleate)......106.88 30 Mvlan ige -

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Prio \$		Fully dised	
EMTRICITABINE – Special Authority see SA1651 on page 104 - Cap 200 mg		30	/	<u>Emtriva</u>
LAMIVUDINE – Special Authority see SA1651 on page 104 – Re Tab 150 mg		60	•	Lamivudine Alphapharm
Lamivudine Alphapharm to be Sole Supply on 1 Novembor Oral liq 10 mg per ml		240 ml OP	•	зтс
ZIDOVUDINE [AZT] – Special Authority see SA1651 on page 10 Cap 100 mg Oral liq 10 mg per ml	152.25	cy 100 200 ml OP		Retrovir Retrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority see Note: zidovudine [AZT] with lamivudine (combination tablets the anti-retroviral Special Authority.				•
Tab 300 mg with lamivudine 150 mg	33.00	60	•	Alphapharm
Protease Inhibitors				
ATAZANAVIR SULPHATE – Special Authority see SA1651 on p Cap 150 mg Cap 200 mg	141.68	harmacy 60 60		<u>Teva</u> Teva
DARUNAVIR – Special Authority see SA1651 on page 104 – Re Tab 400 mg Tab 600 mg	335.00	60 60		Prezista Prezista
LOPINAVIR WITH RITONAVIR — Special Authority see SA1651 Tab 100 mg with ritonavir 25 mg Tab 200 mg with ritonavir 50 mg Oral liq 80 mg with ritonavir 20 mg per ml	183.75 463.00 735.00	tail pharmacy 60 120 300 ml OP	1	Kaletra Kaletra Kaletra
RITONAVIR – Special Authority see SA1651 on page 104 – Ret Tab 100 mg		30	•	Norvir
Strand Transfer Inhibitors				
DOLUTEGRAVIR – Special Authority see SA1651 on page 104 Tab 50 mg		30	1	Tivicay
RALTEGRAVIR POTASSIUM - Special Authority see SA1651 o Tab 400 mg		ail pharmacy 60	•	Isentress Isentress HD

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

1) Diagnosis

	ubsidy	Fully	Brand or
	cturer's Price) Su	ubsidised	Generic
(wich funds	\$ Per	ubsidised ✓	Manufacturer

continued...

- Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
- PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
- Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Exclusion Criteria

- Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia (< 2.0 × 10⁹) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

INTERFERON ALFA-2A - PCT

See prescribing guideline on the previous page

(Roferon-A Inj 3 m iu prefilled syringe to be delisted 1 December 2020)

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA1936 below - Retail pharmacy

- a) See prescribing guideline on the previous page
- b) Note: PHARMAC will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at PHARMAC on 0800-023-588 option 4.

⇒SA1936 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's	Price)	Subsidised	Generic	
\$	Per	· /	Manufacturer	

continued...

- 3.1 Patient has responder relapsed; or
- 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis): and
- 6 Compensated liver disease: and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*: and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Roth:

INFECTIONS - AGENTS FOR SYSTEMIC USE

Subsidy	Fu	ly Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per	 Manufacturer 	

continued...

- 3.1 Patient has a myeloproliferative disorder; and
- 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

Notes:

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet quidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

Urinary Tract Infections

METHENAMINE (HEXAMINE) HIPPURATE			
* Tab 1 g	40.01	100	✓ Hiprex
NITROFURANTOIN			
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
* Tab 100 mg	37.50	100	✓ Nifuran
NORFLOXACIN			
Tab 400 mg - Subsidy by endorsement	135.00	100	Arrow-Norfloxacin
Only if any and four and four and in the any constant of	all and a source to a set to the other	a disart to consul-	and a section to a floor line and a second

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy		Fully	Brand or
	(Manufacturer's Price) 5	Subsidised	I Generic
	` \$	Per	1	Manufacturer
Anticholinesterases				
Antionomicotorasco				
NEOSTIGMINE METILSULFATE				
	10.00	40		l
Inj 2.5 mg per ml, 1 ml ampoule		10		Juno S29
	98.00	50	•	AstraZeneca
PYRIDOSTIGMINE BROMIDE				
▲ Tab 60 mg	45 79	100	1	Mestinon
		100		<u> </u>
Non-Steroidal Anti-Inflammatory Drugs				
Non-Steroidal Anti-illianimatory Drugs				
DICLOFENAC SODIUM				
	1.00	ΕO	./	Dielefense Candan
* Tab EC 25 mg		50		Diclofenac Sandoz
* Tab 50 mg dispersible		20		Voltaren D
* Tab EC 50 mg		50		Diclofenac Sandoz
* Tab long-acting 75 mg	22.80	500	✓	Apo-Diclo SR
* Tab long-acting 100 mg	25.15	500	✓	Apo-Diclo SR
* Inj 25 mg per ml, 3 ml ampoule - Up to 5 inj available on a F	SO 13.20	5	/	Voltaren
* Suppos 12.5 mg		10	1	Voltaren
* Suppos 25 mg		10		Voltaren
1.1		10		Voltaren
* Suppos 50 mg – Up to 10 supp available on a PSO				
* Suppos 100 mg	7.00	10	•	Voltaren
IBUPROFEN				
* Tab 200 mg	11.71	1,000	/	Relieve
Tab long-acting 800 mg		30		Ibuprofen SR BNM
Tab long doing ood mg	7.99	00		Brufen SR
Ibunyafan CD DNM ta ba Cala Cumhy an 1 Dagambay 00			•	Diuleii Sh
Ibuprofen SR BNM to be Sole Supply on 1 December 20		000	,	Fal. I
* Oral liq 20 mg per ml		200 ml	•	<u>Ethics</u>
(Brufen SR Tab long-acting 800 mg to be delisted 1 December 20	020)			
KETOPROFEN				
* Cap long-acting 200 mg	12.07	28	/	Oruvail SR
, , ,		_0		
MEFENAMIC ACID				
* Cap 250 mg	1.25	50		
	(9.16)			Ponstan
	0.50	20		
	(5.60)			Ponstan
NAPROXEN	, ,			
	20.60	E00	.,	Notion 250
* Tab 250 mg		500		Noflam 250
* Tab 500 mg		250		Noflam 500
* Tab long-acting 750 mg		28		Naprosyn SR 750
* Tab long-acting 1 g	8.21	28	1	Naprosyn SR 1000
SULINDAC				
* Tab 100 mg	0.57	56	ı	Mylan S29
· · · · · · · · · · · · · · · · · · ·				•
* Tab 200 mg		50		Aclin
	16.91	56	•	Sulindac Mylan S29
TENOXICAM				
* Tab 20 mg	9 15	100	1	Tilcotil
* Inj 20 mg vial		1		AFT
* III 20 IIIY viai	3.30	1	•	OL I

	N			
	Subsidy (Manufacturer's Pric	e) Sub Per	sidised (Brand or Generic Manufacturer
NSAIDs Other				
CELECOXIB			_	
Cap 100 mg		60		ecoxib Pfizer
Cap 200 mg	2.30	30	✓ Celo ✓ Celo	eprex ecoxib Pfizer
Topical Products for Joint and Muscular Pain				
Topical Floducts for Joint and Muscular Fain				
CAPSAICIN				
Crm 0.025% - Special Authority see SA1289 below - Retain the service of the servi		05 × 0D		Aud
pharmacy	6.95 9.95	25 g OP 45 g OP	✓ Zos ✓ Zos	
	13.27	45 g OP 60 g OP		by Capsaicin
	10.21	00 g Oi	To	opical
			С	ream S29
⇒SA1289 Special Authority for Subsidy				
nitial application from any relevant practitioner. Approvals val osteoarthritis that is not responsive to paracetamol and oral non-	id without further re -steroidal anti-inflan	newal unles nmatories ar	s notified v e contraind	vhere the patient has dicated.
Antirheumatoid Agents HYDROXYCHLOROQUINE – Subsidy by endorsement				
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, system suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmonary and non-pulmonary and annotate the prescription as endorsed who hydroxychloroquine. Note: Indication marked with a * is an	forms of lupus and onary)*, and the pre ere there exists a re unapproved indica	lichen planu escription is ecord of prior tion.	us, cutaned endorsed a r dispensin	ous vasculitides and accordingly.
HYDROXYCHLOROQUINE – Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, systen suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulm- Pharmacists may annotate the prescription as endorsed who hydroxychloroquine. Note: Indication marked with a * is an Tab 200 mg	forms of lupus and onary)*, and the pre ere there exists a re unapproved indica	lichen planu escription is ecord of prior	ıs, cutaned endorsed a	ous vasculitides and accordingly.
HYDROXYCHLOROQUINE – Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, system suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulm- Pharmacists may annotate the prescription as endorsed who hydroxychloroquine. Note: Indication marked with a * is an * Tab 200 mg	forms of lupus and onary)*, and the pre- ere there exists a re- unapproved indica 	lichen planuescription is ecord of priorition. 100	us, cutaned endorsed a r dispensin	ous vasculitides and accordingly. g of quenil
HYDROXYCHLOROQUINE – Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, systen suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulm- Pharmacists may annotate the prescription as endorsed who hydroxychloroquine. Note: Indication marked with a * is an Tab 200 mg	forms of lupus and onary)*, and the preere there exists a re unapproved indica	lichen planu escription is ecord of prior tion.	us, cutaned endorsed a r dispensin Place	ous vasculitides and accordingly. Ig of quenil -Leflunomide
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, systen suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulm Pharmacists may annotate the prescription as endorsed wh hydroxychloroquine. Note: Indication marked with a * is an * Tab 200 mg	forms of lupus and onary)*, and the pre- ere there exists a re- unapproved indica 	lichen planuescription is ecord of priorition. 100	us, cutaned endorsed a r dispensin	ous vasculitides and accordingly. Ig of quenil -Leflunomide
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, system suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulme Pharmacists may annotate the prescription as endorsed whole drown the sendorsed whole substitution of the substitution of the sendorsed whole substitution of the sendorsed whole substitution of the sendorsed whole substitution of the substitution of the sendorsed whole sub	forms of lupus and onary)*, and the preere there exists a re unapproved indicated	lichen planuescription is ecord of priorition. 100	us, cutaned endorsed a r dispensin Plac Apo Ara	ous vasculitides and accordingly. Ig of quenil -Leflunomide va
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, systen suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulm Pharmacists may annotate the prescription as endorsed wh hydroxychloroquine. Note: Indication marked with a * is an * Tab 200 mg	forms of lupus and onary)*, and the preere there exists a re unapproved indicated	lichen planu escription is ecord of prior tion. 100	us, cutaned endorsed a r dispensin Plac Apo Ara	ous vasculitides and accordingly. Ig of quenil -Leflunomide va -Leflunomide
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, systen suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulme Pharmacists may annotate the prescription as endorsed whole the prescri	forms of lupus and onary)*, and the preere there exists a re unapproved indicated	lichen planu escription is ecord of prior tion. 100	is, cutanecendorsed a r dispensin Place Apo Ara	ous vasculitides and accordingly. Ig of quenil -Leflunomide va -Leflunomide
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, system suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulm- Pharmacists may annotate the prescription as endorsed wh- hydroxychloroquine. Note: Indication marked with a * is an * Tab 200 mg	forms of lupus and onary)*, and the preere there exists a re unapproved indicated	lichen planu escription is ecord of prior tion. 100	is, cutanecendorsed a r dispensin Place Apo Ara	ous vasculitides and accordingly. Ig of quenil -Leflunomide va -Leflunomide
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, system suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulm- Pharmacists may annotate the prescription as endorsed wh- hydroxychloroquine. Note: Indication marked with a * is an * Tab 200 mg	forms of lupus and onary)*, and the preere there exists a re unapproved indicated	lichen planu escription is ecord of prior tion. 100	is, cutanecendorsed a r dispensin Place Apo Ara	ous vasculitides and accordingly. Ig of quenil -Leflunomide va -Leflunomide
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, systen suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulm Pharmacists may annotate the prescription as endorsed wh hydroxychloroquine. Note: Indication marked with a * is an * Tab 200 mg	forms of lupus and onary)*, and the preere there exists a re unapproved indicated	lichen planu escription is ecord of prior tion. 100	is, cutanecendorsed a r dispensin Place Apo Ara	ous vasculitides and accordingly. Ig of quenil -Leflunomide va -Leflunomide
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, system suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulme Pharmacists may annotate the prescription as endorsed whe hydroxychloroquine. Note: Indication marked with a * is an * Tab 200 mg ** Tab 200 mg	forms of lupus and onary)*, and the preere there exists a re unapproved indicated	lichen planuscription is ecord of priorition. 100 30 30	is, cutanecendorsed a r dispensin Place Apo Ara Apo Ara	ous vasculitides and accordingly. Ig of quenil D-Leflunomide va D-Leflunomide va enamine
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, system suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulm Pharmacists may annotate the prescription as endorsed wh hydroxychloroquine. Note: Indication marked with a * is an * Tab 200 mg	forms of lupus and onary)*, and the preere there exists a re unapproved indicated	lichen planuscription is ecord of priorition. 100 30	is, cutanecendorsed a r dispensin Place Apo Ara Apo Ara	ous vasculitides and accordingly. Ig of quenil D-Leflunomide va D-Leflunomide va
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, system suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulme Pharmacists may annotate the prescription as endorsed whe hydroxychloroquine. Note: Indication marked with a * is an * Tab 200 mg ** Tab 200 mg	forms of lupus and onary)*, and the preere there exists a re unapproved indicated	lichen planuscription is ecord of priorition. 100 30 30	is, cutanecendorsed a r dispensin Place Apo Ara Apo Ara	ous vasculitides and accordingly. Ig of quenil D-Leflunomide va D-Leflunomide va enamine
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, systen suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulm- Pharmacists may annotate the prescription as endorsed wh- hydroxychloroquine. Note: Indication marked with a * is an Tab 200 mg	forms of lupus and onary)*, and the preere there exists a re unapproved indicated	lichen planuscription is ecord of priorition. 100 30 30	is, cutanecendorsed a r dispensin Place Apo Ara Apo Ara	ous vasculitides and accordingly. Ig of quenil D-Leflunomide va D-Leflunomide va enamine
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, systen suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulm Pharmacists may annotate the prescription as endorsed wh hydroxychloroquine. Note: Indication marked with a * is an * Tab 200 mg LEFLUNOMIDE Tab 10 mg Arava to be Sole Supply on 1 December 2020 Tab 20 mg Arava to be Sole Supply on 1 December 2020 (Apo-Leflunomide Tab 10 mg to be delisted 1 December 2020) (Apo-Leflunomide Tab 20 mg to be delisted 1 December 2020) PENICILLAMINE Tab 125 mg Tab 250 mg Drugs Affecting Bone Metabolism Alendronate for Osteoporosis ALENDRONATE SODIUM	forms of lupus and onary)*, and the preere there exists a reunapproved indication	lichen planuscription is ecord of priorition. 100 30 30	is, cutanecendorsed a r dispensin Plat Apc Ara Apc Ara	ous vasculitides and accordingly. Ig of aquenil D-Leflunomide va D-Leflunomide va Penamine eenamine
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, systen suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulm Pharmacists may annotate the prescription as endorsed wh hydroxychloroquine. Note: Indication marked with a * is an * Tab 200 mg LEFLUNOMIDE Tab 10 mg Arava to be Sole Supply on 1 December 2020 Tab 20 mg Arava to be Sole Supply on 1 December 2020 (Apo-Leflunomide Tab 10 mg to be delisted 1 December 2020) (Apo-Leflunomide Tab 20 mg to be delisted 1 December 2020) PENICILLAMINE Tab 125 mg Tab 250 mg Drugs Affecting Bone Metabolism Alendronate for Osteoporosis ALENDRONATE SODIUM	forms of lupus and onary)*, and the preere there exists a reunapproved indication	lichen planuscription is ecord of priorition. 100 30 30	is, cutanecendorsed a r dispensin Place Apo Ara Apo Ara	ous vasculitides and accordingly. Ig of aquenil D-Leflunomide va D-Leflunomide va Penamine eenamine
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, systen suppression, relevant dermatological conditions (cutaneous mucosal ulceration)*, sarcoidosis (pulmonary and non-pulm- Pharmacists may annotate the prescription as endorsed wh- hydroxychloroquine. Note: Indication marked with a * is an * Tab 200 mg	forms of lupus and onary)*, and the preere there exists a reunapproved indication	lichen planuscription is ecord of priorition. 100 30 30	is, cutanecendorsed a r dispensin Plat Apc Ara Apc Ara	ous vasculitides and accordingly. Ig of aquenil D-Leflunomide va D-Leflunomide va Penamine eenamine

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy (Manufacturer's Price)

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

Other Treatments

DENOSUMAB – Special Authority see SA1777 below – Retail pharmacy Inj 60 mg prefilled syringe.......326.00

1 ✓ Prolia

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Either:
 - 2.1 The patient is female and postmenopausal; or
 - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	
PAMIDRONATE DISODIUM				
Inj 3 mg per ml, 10 ml vial	5.98	1	/	Pamisol
Inj 6 mg per ml, 10 ml vial	15.02	1	1	Pamisol
Inj 9 mg per ml, 10 ml vial		1	1	Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see S/ * Tab 60 mg		armacy 28	/	Evista

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM Tab 35 mg3.10	4	✓ Risedronate Sandoz
TERIPARATIDE – Special Authority see SA1139 below – Retail pharmacy		
Inj 250 mcg per ml, 2.4 ml490.00	1	✓ Forteo

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a

	Subsidy		Fully	Brand or
(M	anufacturer's Price)	Subs	idised	Generic
	\$	Per	✓	Manufacturer

continued...

funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, vial - Special Authority see 100 ml OP Aclasta

⇒SA1780 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain: or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1

Subsidy (Manufacturer's Price)	Sul	Fully	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note): or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Notes:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

ALL ODLIDINO

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

AL	LOPURINOL			
*	Tab 100 mg	11.47	500	✓ DP-Allopurinol
	DP-Allopurinol to be Sole Supply on 1 November 2020			•
*	Tab 300 mg	28.57	500	✓ DP-Allopurinol
	DP-Allopurinol to be Sole Supply on 1 November 2020			
BE	NZBROMARONE – Special Authority see SA1537 below – Reta	il pharmacy		
	Tab 50 mg	22.50	100	✓ Narcaricin mite S29
	Tab 100 mg	13.50	30	✓ Desuric S29
				✓ Urinorm S29
		45.00	100	Benzbromaron AL
				100 S29

⇒SA1537 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 Both:
 - 2.3.1 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Notes); and
 - 2.3.2 The patient has a rate of creatinine clearance greater than or equal to 20 ml/min; or
 - 2.4 All of the following:
 - 2.4.1 The patient is taking azathioprine and requires urate-lowering therapy; and
 - 2.4.2 Allopurinol is contraindicated; and
 - 2.4.3 Appropriate doses of probenecid are ineffective or probenecid cannot be used due to reduced renal function; and
- 3 The patient is receiving monthly liver function tests.

Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

Notes: Benzbromarone has been associated with potentially fatal hepatotoxicity.

In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

The New Zealand Rheumatology Association has developed information for prescribers which can be accessed from its website at www.rheumatology.org.nz/home/resources-2/

COLCHICINE

* Tab 500 mcg	9.58	100	✓ Colgout
FEBUXOSTAT - Special Authority see SA1931 below - Reta	ail pharmacy		
Tab 80 mg	39.50	28	✓ Adenuric
Tab 120 mg	39.50	28	Adenurio

⇒SA1931 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
 - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

PROBENECID

*	Tab 500 mg	55.00	100	1	Probenecid-AFT
---	------------	-------	-----	---	----------------

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Muscle Relaxants

masore relaxants			
BACLOFEN			
* Tab 10 mg		100	✓ Pacifen
Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement	11.55	1	Lioresal Intrathecal
Subsidised only for use in a programmable pump in patier			ents have been ineffective or have
caused intolerable side effects and the prescription is end-	orsed according	ly.	
Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement	372.98	5	✓ <u>Medsurge</u>
Subsidised only for use in a programmable pump in patier caused intolerable side effects and the prescription is end-			ents have been ineffective or have
DANTROLENE			
Cap 25 mg	97.50	100	✓ Dantrium
			✓ Dantrium S29 S29
Cap 50 mg	77.00	100	✓ Dantrium
ORPHENADRINE CITRATE			
Tab 100 mg	18 54	100	✓ Norflex

Subsidy (Manufacturer's Price) Su

Fully Subsidised Per

Brand or Generic Manufacturer

Agents for Parkinsonism and Related Disorders

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE			
▲ Cap 100 mg	38.24	60	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE		_	4
▲ Inj 10 mg per ml, 2 ml ampoule		5	✓ <u>Movapo</u>
▲ Inj 10 mg per ml, 5 ml ampoule	121.84	5	✓ Movapo
BROMOCRIPTINE MESYLATE			
* Tab 2.5 mg	32.08	100	✓ Apo-Bromocriptine
ENTACAPONE			
▲ Tab 200 mg	22.00	100	✓ Entapone
LEVODOPA WITH BENSERAZIDE			
* Tab dispersible 50 mg with benserazide 12.5 mg		100	Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg		100	✓ Madopar 62.5
* Cap 100 mg with benserazide 25 mg		100	✓ Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg		100	✓ Madopar HBS
* Cap 200 mg with benserazide 50 mg	26.25	100	✓ Madopar 250
LEVODOPA WITH CARBIDOPA			<i>a</i>
* Tab 100 mg with carbidopa 25 mg		100	✓ Kinson
Cinemat to be Cale Cumply on 1 December 2000	21.11		✓ Sinemet
Sinemet to be Sole Supply on 1 December 2020	00.04	100	/ Mulan 000
* Tab long-acting 100 mg with carbidopa 25 mg Tab long-acting 200 mg with carbidopa 50 mg		100 100	✓ Mylan S29 ✓ Sinemet CR
* Tab long-acting 200 mg with carbidopa 50 mg		100	
* Tab 250 mg with carbidopa 25 mg	46.73	100	✓ Mylan S29 ✓ Sinemet
Sinemet to be Sole Supply on 1 December 2020		100	▼ Sillelliet
(Kinson Tab 100 mg with carbidopa 25 mg to be delisted 1 Dec	remher 2020)		
(Mylan \$29) Tab long-acting 100 mg with carbidopa 25 mg to b		nhar 2020)	
	de delisted i Deceli	1061 2020)	
PRAMIPEXOLE HYDROCHLORIDE Tab 0.25 mg	6.10	100	✓ Ramipex
▲ Tab 0.25 filg		100	✓ Ramipex
-	20.70	100	· <u>mamipex</u>
ROPINIROLE HYDROCHLORIDE	0.05	84	✓ Ropin
▲ Tab 0.25 mg	3.39	100	✓ <u>Hopin</u> ✓ Mylan \$29
▲ Tab 1 mg		84	✓ Ropin
Tab ing	4.70	100	✓ Mylan \$29
▲ Tab 2 mg	•	84	✓ Ropin
▲ Tab 5 mg		84	✓ Ropin
SELEGILINE HYDROCHLORIDE	12.00	0-1	- IIOPIII
* Tab 5 mg	22.00	100	✓ Apo-Selegiline
* Tab only	22.00	100	S29 S29
TOLOADONE			323 323
TOLCAPONE A Tob 100 mg	150.00	100	✓ Tasmar
▲ Tab 100 mg	152.38	100	₹ rasmar



NERVOUS STSTEM				
	Subsidy (Manufacturer's Price)	Subs Per	Fully idised	Brand or Generic Manufacturer
Anticholinergics				
BENZATROPINE MESYLATE Tab 2 mg		60 5 10	✓ C	Benztrop Cogentin Phebra Omega
a) Up to 10 inj available on a PSO b) Only on a PSO c) Phebra to be Sole Supply on 1 December 2020 (Cogentin Inj 1 mg per ml, 2 ml to be delisted 1 December 2020) (Omega Inj 1 mg per ml, 2 ml to be delisted 1 December 2020) PROCYCLIDINE HYDROCHLORIDE Tab 5 mg	7.40	100	√ k	Kemadrin
Agents for Essential Tremor, Chorea and Relate	ed Disorders			
RILUZOLE – Special Authority see SA1403 below – Retail pharr Wastage claimable Tab 50 mg	•	56	√ <u>F</u>	Rilutek
■ SA1403 Special Authority for Subsidy Initial application only from a neurologist or respiratory specialis following criteria: All of the following: 1 The patient has amyotrophic lateral sclerosis with disease 2 The patient has at least 60 percent of predicted forced vital contents.	duration of 5 years o	or less; and	i	
3 The patient has not undergone a tracheostomy; and 4 The patient has not experienced respiratory failure; and				

- 5 Any of the following:
 - 5.1 The patient is ambulatory; or
 - 5.2 The patient is able to use upper limbs; or
 - 5.3 The patient is able to swallow.

Renewal from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
 - 3.1 The patient is ambulatory; or
 - 3.2 The patient is able to use upper limbs; or
 - 3.3 The patient is able to swallow.

Т	E٦	ΓF	₹/	٩В	El	N٨	٩Z	ΙN	Ε

Tab 25 mg91.10 112 ✓ <u>Motetis</u>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
Anaesthetics				
Local				
LIDOCAINE [LIGNOCAINE] Gel 2%, tube - Subsidy by endorsement		30 m		Xylocaine 2% Jelly
 b) Subsidised only if prescribed for urethral or cervical a Gel 2%, 11 ml urethral syringe – Subsidy by endorsement a) Up to 5 each available on a PSO b) Subsidised only if prescribed for urethral, cervical or accordingly. 	42.00	10	•	Instillagel Lido
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE Oral (gel) soln 2%	8.75 17.50 (35.00) 8.25 12.00 (20.00) 6.20	200 m 25 50 25 5 5	•	Mucosoothe Lidocaine-Claris Xylocaine Lidocaine-Claris Xylocaine Lidocaine-Claris Lidocaine-Claris
Subsidy by endorsement		10 e pres		Pfizer endorsed accordingly.
Topical Local Anaesthetics				
■ SA0906 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid condition requiring frequent injections or venepuncture. Renewal from any relevant practitioner. Approvals valid for 2 years from the traction of the provided in the provided i				

benefiting from treatment.

LIDOCAINE [LIGNOCAINE] - Special Authority see SA0906 above	- Retail phai	rmacy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Authorit	y see SA090	6 above – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	EMLA

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

Non-opioid Analgesics

For aspirin & chloroform application refer Standard Formulae, page 248

ASPIRIN

* Tab dispersible 300 mg - Up to 30 tab available on a PSO......4.50 100 ✓ Ethics Aspirin

_		Subsidy (Manufacturer's Pr \$	ice) Sub Per	Fully sidised	Brand or Generic Manufacturer
CAI	PSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or d accordingly.	iabetic peripheral	neuropathy a	ınd the	prescription is endorsed
	Crm 0.075%	12.50 15.83	45 g OP 57 g OP		Zostrix HP Rugby Capsaicin Topical Cream ®29
NEI	FOPAM HYDROCHLORIDE				• • • • • • • • • • • • • • • • • • • •
	Tab 30 mg	23.40	90	1	Acupan
PAF	RACETAMOL				
	Tab 500 mg - blister pack	0.50	20	✓ I	Medco Paracare Pharmacy Health
		1.12			Ethics Paracetamol Classic
		2.48	100		Paracare Pharmacy Health
		11.75 24.82	96 1,000		Panadol Mini Caps Paracetamol Pharmacare
	c)				
	Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, a annotate the prescription as endorsed where d Maximum of 100 tab per dispensing for non-en (for non-endorsed patients), then dispense in real tab 500 mg - bottle pack — Maximum of 300 tab per	nd the prescriptions spensing history states of the desired patients.	n is annotate supports a loi If quantities p	d accor ng-term rescrib ng 100 t	dingly. Pharmacists may condition. ed for more than 100 tabs tab per dispensing.
	1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, a annotate the prescription as endorsed where d 2) Maximum of 100 tab per dispensing for non-en (for non-endorsed patients), then dispense in refab 500 mg - bottle pack — Maximum of 300 tab per prescription; can be waived by endorsement	and the prescription spensing history statements of the desired patients. Seperat dispensings	n is annotate supports a lor If quantities p not exceedir 1,000	d accor ng-term rescribe ng 100 t	dingly. Pharmacists may condition. ed for more than 100 tabs tab per dispensing. Pharmacare
	1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, a annotate the prescription as endorsed where d 2) Maximum of 100 tab per dispensing for non-en (for non-endorsed patients), then dispense in real tab 500 mg - bottle pack — Maximum of 300 tab per	nd the prescription is pensing history statements. Seperat dispensings 24.82 railable for patient scription is annotal supports a long-tised patients. If questions is supports a long-tised patients.	n is annotate supports a loi If quantities p not exceedir 1,000 s with long te ted according erm conditior uantities pres	d accorning-term rescribing 100 to the control of t	dingly. Pharmacists may condition. ed for more than 100 tabs tab per dispensing. Pharmacare ditions who require regula armacists may annotate the or more than 100 tabs (for the conditions).
*	1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, a annotate the prescription as endorsed where d 2) Maximum of 100 tab per dispensing for non-en (for non-endorsed patients), then dispense in reference to the pack — Maximum of 300 tab per prescription; can be waived by endorsement	nd the prescription is pensing history statements. Seperat dispensings 24.82 railable for patient scription is annotal supports a long-tied patients. If qui spensings not except the properties of patients and patients and patients are patients.	n is annotate supports a loi If quantities p not exceedir 1,000 s with long te ted according erm conditior uantities pres	d accorng-term rescribing 100 to the rescribing 100 to the rescribing 100 to the rescribing to the rescribed fab per second above the rescribed fab per second to the rescribed fab per second to the rescribed fab per second to the rescribing to th	dingly. Pharmacists may condition. ed for more than 100 tabs tab per dispensing. Pharmacare ditions who require regula armacists may annotate the or more than 100 tabs (for the conditions).
	1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, a annotate the prescription as endorsed where d 2) Maximum of 100 tab per dispensing for non-en (for non-endorsed patients), then dispense in reference for non-endorsed patients), then dispense in reference for non-endorsed patients). Tab 500 mg - bottle pack — Maximum of 300 tab per prescription; can be waived by endorsement	nd the prescription is pensing history addressed patients. It is pensing the peat dispensings and the peat dispensing the peat dispension is annotated patients. If question is an ongstone pensings not except patients. If question is spensings not except patients. If question is spensings not except patients. If question is spensings not except patients.	n is annotate supports a loi f quantities p not exceedir 1,000 s with long te ted according erm conditior uantities pres ceeding 100 t	d accorng-term rescribing 100 for a congly. Phana cribed for ab per decided as a congression of the congress	dingly. Pharmacists may condition. ed for more than 100 tabs tab per dispensing. Pharmacare ditions who require regularmacists may annotate the or more than 100 tabs (for dispensing.
*	1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, a annotate the prescription as endorsed where d 2) Maximum of 100 tab per dispensing for non-en (for non-endorsed patients), then dispense in respectively. Tab 500 mg - bottle pack — Maximum of 300 tab per prescription; can be waived by endorsement	nd the prescription is pensing history addressed patients. In peneat dispensings and a large and a lar	n is annotate supports a lor f quantities protected according erm condition untities preseding 100 the 1,000 ml	d accorng-term rescribing 100 to 100	dingly. Pharmacists may condition. ed for more than 100 tabs tab per dispensing. Pharmacare ditions who require regula armacists may annotate the or more than 100 tabs (for dispensing. Paracare Paracare Double Strength
*	1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, a annotate the prescription as endorsed where d 2) Maximum of 100 tab per dispensing for non-en (for non-endorsed patients), then dispense in reference of the pack — Maximum of 300 tab per prescription; can be waived by endorsement	nd the prescription is pensing history addressed patients. It is pensing be peat dispensings along to the peat dispensing anota supports a long-ties of patients. If question is anota supports a long-ties of patients. If question is seen a long-ties of patients. If question is seen a long-ties of patients. If question is seen a long-ties of patients. It question is seen a long-ti	n is annotate supports a lor f quantities protected according error annities presceeding 1,000 to 1,000 ml	d accorng-term rescribing 100 ft	dingly. Pharmacists may condition. ed for more than 100 tabs tab per dispensing. Pharmacare ditions who require regula armacists may annotate the or more than 100 tabs (for dispensing. Paracare Paracare Double

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
Opioid Analgesics				
CODEINE PHOSPHATE - Safety medicine; prescriber may dete	ermine dispensing fre	quenc	٧	
Tab 15 mg		100		PSM
PSM to be Sole Supply on 1 November 2020				
Tab 30 mg	7.45	100	1	PSM
PSM to be Sole Supply on 1 November 2020				
Tab 60 mg	14.25	100	•	PSM
PSM to be Sole Supply on 1 November 2020				
DIHYDROCODEINE TARTRATE				
Tab long-acting 60 mg	8.60	60	1	DHC Continus
FENTANYL				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	equency			
Inj 50 mcg per ml, 2 ml ampoule		5	1	Fentanyl GH
, , , ,	3.56	10	✓	Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule	9.41	10	✓	Boucher and Muir
Patch 12.5 mcg per hour	2.95	5	1	Fentanyl Sandoz
Patch 25 mcg per hour	3.66	5	1	Fentanyl Sandoz
Patch 50 mcg per hour		5		Fentanyl Sandoz
Patch 75 mcg per hour		5		Fentanyl Sandoz
Patch 100 mcg per hour		5	•	Fentanyl Sandoz
(Fentanyl GH Inj 50 mcg per ml, 2 ml ampoule to be delisted 1 Ja	anuary 2021)			
METHADONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre				
d) Extemporaneously compounded methadone will only be	reimbursed at the rate	e of th	e cheapes	st form available
(methadone powder, not methadone tablets).				
e) For methadone hydrochloride oral liquid refer Standard F	71 0		_	
Tab 5 mg		10		<u>Methatabs</u>
Oral liq 2 mg per ml		200 m		Biodone Biodone
Oral liq 5 mg per ml		200 m		Biodone Forte
Oral liq 10 mg per ml		200 ml 10		Biodone Extra Forte AFT
Inj 10 mg per ml, 1 ml	01.00	10	•	AFI
MORPHINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre				
Oral liq 1 mg per ml		200 m		RA-Morph
Oral liq 2 mg per ml		200 m		RA-Morph
Oral liq 5 mg per ml	19.44 2	200 m		Ordine \$29
Out line 40 and a sound	07.74	200		RA-Morph
Oral liq 10 mg per ml	27.74 2	200 m		Ordine \$29
			•	RA-Morph

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
MORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
 Safety medicine; prescriber may determine dispensing free 				
Tab immediate-release 10 mg	2.80	10	•	Sevredol
Sevredol to be Sole Supply on 1 November 2020			_	
Tab long-acting 10 mg		10		Arrow-Morphine LA
Tab immediate-release 20 mg	5.52	10	•	Sevredol
Sevredol to be Sole Supply on 1 November 2020	0.05	40		Americ Manakina I A
Tab long-acting 30 mg		10		Arrow-Morphine LA
Tab long-acting 60 mgCap long-acting 10 mg		10 10		Arrow-Morphine LA m-Eslon
Cap long-acting 10 mg		10		m-Esion
		10		m-Eslon
Cap long-acting 60 mg Cap long-acting 100 mg		10		m-Esion
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS		5		DBL Morphine
ing 5 mg per mi, 1 mi ampoule - Op to 5 ing available on a 1 c	0.27	J	•	Sulphate
Inj 10 mg per ml, 1 ml ampoule - Up to 5 inj available on a P	SO 1.17	5	1	DBL Morphine
ing to mg per mi, i mi ampoule — op to 5 ing available on a r	504.47	J	•	Sulphate
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a P	SO 476	5	1	DBL Morphine
ing 13 mg per mi, 1 mi ampoule – op to 3 mg available on a F	304.70	5	•	Sulphate
Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on a P	SO 610	5	1	DBL Morphine
ing so mg per mi, i mi ampoule – op to s ing available on a F	300.19	5	•	Sulphate
Arrow-Morphine LA Tab long-acting 10 mg to be delisted 1 Octo	har 2020)			Julphate
,	Del 2020)			
DXYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre		00	,	0
Tab controlled-release 5 mg		20		Oxycodone Sandoz
Tab controlled-release 10 mg		20		Oxycodone Sandoz Oxycodone Sandoz
Tab controlled release 20 mg		20 20		
Tab controlled-release 40 mg Tab controlled-release 80 mg		20		Oxycodone Sandoz Oxycodone Sandoz
Cap immediate-release 5 mg		20		OxyNorm
Cap immediate-release 5 mg		20		OxyNorm
Cap immediate-release 10 mg		20		OxyNorm
Oral lig 5 mg per 5 ml		250 n		OxyNorm
Inj 10 mg per ml, 1 ml ampoule		5		OxyNorm
Inj 10 mg per ml, 2 ml ampoule		5		OxyNorm
Inj 50 mg per ml, 1 ml ampoule		5		OxyNorm
PARACETAMOL WITH CODEINE – Safety medicine; prescriber		-		
* Tab paracetamol 500 mg with codeine phosphate 8 mg		1,000		y Paracetamol +
Tab paracetation booting with codellie phosphate offig	10.21	1,000	,	Codeine (Relieve)

Codeine (Relieve)

			NLN	VOUS STSTEW
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing f				
Tab 50 mg		10	✓ [
Inj 50 mg per ml, 1 ml ampoule - Up to 5 inj available on a	PSO4.98	5	✓ [BL Pethidine Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a	PSO 5.12	5	√ Γ	BL Pethidine
ing 50 mg per mi, 2 mi ampoule – op to 5 mg available on a	F305.12	5	v L	Hydrochloride
TRAMADOL HYDROCHLORIDE				•
Tab sustained-release 100 mg	1.52	20	√ T	ramal SR 100
Tramal SR 100 to be Sole Supply on 1 November 2020				
Tab sustained-release 150 mg	2.10	20	√ T	ramal SR 150
Tramal SR 150 to be Sole Supply on 1 November 2020	1			
Tab sustained-release 200 mg		20	√ T	ramal SR 200
Tramal SR 200 to be Sole Supply on 1 November 2020				
Cap 50 mg		100	✓ A	Arrow-Tramadol
Arrow-Tramadol to be Sole Supply on 1 December 202	0			
Antidepressants				
Antidepressants				
Cyclic and Related Agents				
AMITRIPTYLINE - Safety medicine; prescriber may determine	dispensina frequency			
Tab 10 mg		100	✓ A	Arrow-Amitriptyline
Arrow-Amitriptyline to be Sole Supply on 1 December 2	2020			
Tab 25 mg		100	✓ A	Arrow-Amitriptyline
Arrow-Amitriptyline to be Sole Supply on 1 December 2				
Tab 50 mg		100	✓ p	Arrow-Amitriptyline
Arrow-Amitriptyline to be Sole Supply on 1 December 2	020			
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; presc	riber may determine d	ispen	sing freque	псу
Tab 10 mg	13.99	100	✓ A	Anafranil S29
				po-Clomipramine
Tab 25 mg		50		po-Clomipramine
	9.46	100	✓ <u>P</u>	po-Clomipramine
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy by e	ndorsement			
 a) Safety medicine; prescriber may determine dispensing f 				
b) Subsidy by endorsement – Subsidised for patients who				
2019 and the prescription is endorsed accordingly. Pha		the p	prescription	as endorsed where the
exists a record of prior dispensing of dosulepin [dothiepi				
Tab 75 mg		30		Oosulepin Mylan
Cap 25 mg	1.83	50	•	Oosulepin
				Mylan S29
IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescribe	, ,	_		
Tab 10 mg		50		ofranil
T 05	10.96	100	✓ T	ofranil

50

✓ Tofranil

Tab 25 mg8.80

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
MAPROTILINE HYDROCHLORIDE - Subsidy by endorsemen	t			
 a) Safety medicine; prescriber may determine dispensing b) Subsidy by endorsement – Subsidised for patients who 2020 and the prescription is endorsed accordingly. Pha exists a record of prior dispensing of maprotiline hydroc 	were taking maprotilin armacists may annotat hloride.	e the	prescriptio	n as endorsed where ther
Tab 25 mg		30		Ludiomil
	12.53	50		Ludiomil
Tob 75 mg	25.06	100		Ludiomil
Tab 75 mg	14.01	20 30		Ludiomil Ludiomil
(Ludiomil Tab 25 mg to be delisted 1 February 2021) (Ludiomil Tab 25 mg to be delisted 1 February 2021) (Ludiomil Tab 25 mg to be delisted 1 February 2021) (Ludiomil Tab 75 mg to be delisted 1 August 2021) (Ludiomil Tab 75 mg to be delisted 1 August 2021)				
NORTRIPTYLINE HYDROCHLORIDE - Safety medicine; pres				
Tab 10 mg		100		Norpress
Tab 25 mg	5.98	180	•	Norpress
Subsidy by endorsement – Subsidised for patients who we prescription is endorsed accordingly. Pharmacists may an prior dispensing of phenelzine sulphate. * Tab 15 mg	notate the prescription		ndorsed wh	
prescription is endorsed accordingly. Pharmacists may an	notate the prescription	as er	ndorsed wh	Lupin S29 Nardil S29 S29
Subsidy by endorsement – Subsidised for patients who we prescription is endorsed accordingly. Pharmacists may an prior dispensing of phenelzine sulphate. * Tab 15 mg	notate the prescription70.80 118.00	as er	ndorsed wh	Lupin S29 Nardil S29 S29
Subsidy by endorsement – Subsidised for patients who we prescription is endorsed accordingly. Pharmacists may an prior dispensing of phenelzine sulphate. * Tab 15 mg	notate the prescription70.80 118.00	60 100	ndorsed wh	nere there exists a record Lupin \$29 Nardil \$29 \$29 Nardil
Subsidy by endorsement – Subsidised for patients who we prescription is endorsed accordingly. Pharmacists may an prior dispensing of phenelzine sulphate. * Tab 15 mg	notate the prescription70.80 118.00	60 100 28	ndorsed wh	Lupin \$29 Nardil \$29 Nardil Parnate \$29 \$29
Subsidy by endorsement – Subsidised for patients who we prescription is endorsed accordingly. Pharmacists may an prior dispensing of phenelzine sulphate. * Tab 15 mg	notate the prescription70.80 118.0012.85 22.94	60 100 28 50	ndorsed wh	Lupin \$29 Nardil \$29 Nardil Parnate \$29 \$29 Parnate
Subsidy by endorsement – Subsidised for patients who we prescription is endorsed accordingly. Pharmacists may an prior dispensing of phenelzine sulphate. * Tab 15 mg		60 100 28 50	ndorsed what when the second s	Lupin \$29 Nardil \$29 Nardil Parnate \$29 \$29 Parnate
Subsidy by endorsement – Subsidised for patients who we prescription is endorsed accordingly. Pharmacists may an prior dispensing of phenelzine sulphate. * Tab 15 mg		28 50 100 60	ndorsed what when the second s	Lupin \$29 Nardil \$29 Nardil \$29 Nardil Parnate \$29 Parnate Parnate \$29 Parnate \$29 Parnate \$29 Parnate \$29

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	I Generic
ESCITALOPRAM	*	. 01		. managaran Ol
* Tab 10 mg	1.11	28	•	Escitalopram- Apotex
* Tab 20 mg	1.90	28	•	Escitalopram- Apotex
FLUOXETINE HYDROCHLORIDE				
* Tab dispersible 20 mg, scored – Subsidy by endorsement	1.98	30	•	Fluox
, G.	9.93		✓	Arrow-Fluoxetine
Subsidised by endorsement 1) When prescribed for a patient who cannot swallow accordingly; or 2) When prescribed in a daily dose that is not a multip	•		·	
endorsed. Note: Tablets should be combined with				
Cap 20 mg	2.91	84	1	Fluox
оар <u>20 mg</u>	7.49	90		Arrow-Fluoxetine
PAROXETINE				
* Tab 20 mg	3.61	90	/	Loxamine
SERTRALINE				
Tab 50 mg	0.92	30	✓	Setrona
* Tab 100 mg	1.61	30		Setrona
Other Antidepressants				
MIRTAZAPINE				
Tab 30 mg	2.63	30	1	Apo-Mirtazapine
Tab 45 mg	3.48	30		Apo-Mirtazapine
VENLAFAXINE				
* Cap 37.5 mg	6.38	84	•	Enlafax XR
* Cap 75 mg	8.11	84	✓	Enlafax XR
* Cap 150 mg	11.16	84	•	Enlafax XR
Antiepilepsy Drugs				
Agents for Control of Status Epilepticus				
CLONAZEPAM – Safety medicine; prescriber may determine displining 1 mg per ml, 1 ml		5	/	Rivotril
DIAZEPAM – Safety medicine; prescriber may determine dispens				
Inj 5 mg per ml, 2 ml ampoule — Subsidy by endorsement a) Up to 5 inj available on a PSO b) Only on a PSO c) PSO must be endorsed "not for anaesthetic procedure	23.66	5	•	Hospira
Rectal tubes 5 mg - Up to 5 tube available on a PSO		5	1	Stesolid
Rectal tubes 10 mg – Up to 5 tube available on a PSO		5		Stesolid
(Stesolid Rectal tubes 10 mg to be delisted 1 December 2020)	····· · · · · · · ·	-		
PARALDEHYDE				
* Inj 5 ml	1,500.00	5	•	AFT S29

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

NERVOUS SYSTEM

PHENYTOIN SODIUM * Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO88.63 5 Hospira * Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a PSO88.63 5 Hospira * Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a PSO88.63 5 Hospira * Control of Epilepsy * CARBAMAZEPINE * Tab 200 mg	
★ Inj 50 mg per ml, 2 ml ampoule — Up to 5 inj available on a PSO88.63 5 ✓ Hospira ★ Inj 50 mg per ml, 5 ml ampoule — Up to 5 inj available on a PSO	
★ Inj 50 mg per ml, 5 ml ampoule — Up to 5 inj available on a PSO	
Control of Epilepsy CARBAMAZEPINE * Tab 200 mg	
Control of Epilepsy CARBAMAZEPINE ★ Tab 200 mg 14.53 100 ✓ Tegretol ★ Tab long-acting 200 mg 16.98 100 ✓ Tegretol CR ★ Tab 400 mg 34.58 100 ✓ Tegretol CR ★ Tab long-acting 400 mg 39.17 100 ✓ Tegretol CR ★ Oral liq 20 mg per ml 26.37 250 ml ✓ Tegretol CLOBAZAM – Safety medicine; prescriber may determine dispensing frequency Tab 10 mg 9.12 50 ✓ Frisium CLONAZEPAM – Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml 7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE Cap 250 mg 140.88 100 ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin	
CARBAMAZEPINE # Tab 200 mg 14.53 100 ✓ Tegretol # Tab long-acting 200 mg 16.98 100 ✓ Tegretol CR # Tab 400 mg 34.58 100 ✓ Tegretol CR # Tab long-acting 400 mg 39.17 100 ✓ Tegretol CR # Oral liq 20 mg per ml 26.37 250 ml ✓ Tegretol CLOBAZAM – Safety medicine; prescriber may determine dispensing frequency Tab 10 mg 9.12 50 ✓ Frisium CLONAZEPAM – Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml 7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE Cap 250 mg 140.88 100 ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin	
★ Tab 200 mg 14.53 100 ✓ Tegretol ★ Tab long-acting 200 mg 16.98 100 ✓ Tegretol CR ★ Tab 400 mg 34.58 100 ✓ Tegretol ★ Tab long-acting 400 mg 39.17 100 ✓ Tegretol CR ★ Oral liq 20 mg per ml 26.37 250 ml ✓ Tegretol CLOBAZAM - Safety medicine; prescriber may determine dispensing frequency Tab 10 mg 9.12 50 ✓ Frisium CLONAZEPAM - Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml 7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE Cap 250 mg 140.88 100 ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin	
★ Tab long-acting 200 mg 16.98 100 ✓ Tegretol CR ★ Tab 400 mg 34.58 100 ✓ Tegretol ★ Tab long-acting 400 mg 39.17 100 ✓ Tegretol CR ★ Oral liq 20 mg per ml 26.37 250 ml ✓ Tegretol CLOBAZAM - Safety medicine; prescriber may determine dispensing frequency Tab 10 mg 9.12 50 ✓ Frisium CLONAZEPAM - Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml 7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE Cap 250 mg 140.88 100 ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin	
★ Tab 400 mg 34.58 100 ✓ Tegretol ★ Tab long-acting 400 mg 39.17 100 ✓ Tegretol CR ★ Oral liq 20 mg per ml 26.37 250 ml ✓ Tegretol CLOBAZAM – Safety medicine; prescriber may determine dispensing frequency Tab 10 mg 9.12 50 ✓ Frisium CLONAZEPAM – Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml 7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE Cap 250 mg 140.88 100 ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin	
★ Tab long-acting 400 mg 39.17 100 ✓ Tegretol CR ★ Oral liq 20 mg per ml 26.37 250 ml ✓ Tegretol CLOBAZAM — Safety medicine; prescriber may determine dispensing frequency Tab 10 mg 9.12 50 ✓ Frisium CLONAZEPAM — Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml 7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE Cap 250 mg 140.88 100 ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin	
★ Oral liq 20 mg per ml 26.37 250 ml ✓ Tegretol CLOBAZAM - Safety medicine; prescriber may determine dispensing frequency Tab 10 mg 9.12 50 ✓ Frisium CLONAZEPAM - Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml 7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE Cap 250 mg 140.88 100 ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin	
CLOBAZAM – Safety medicine; prescriber may determine dispensing frequency Tab 10 mg	
Tab 10 mg	
CLONAZEPAM – Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml	
Oral drops 2.5 mg per ml .7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE	
ETHOSUXIMIDE Cap 250 mg	
Cap 250 mg 140.88 100 ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin	
Oral liq 250 mg per 5 ml	
3,11	
CADADENTIN	
Note: Not subsidised in combination with subsidised pregabalin	
★ Cap 100 mg	
※ Cap 300 mg	
★ Cap 400 mg	
LACOSAMIDE - Special Authority see SA1125 below - Retail pharmacy	
▲ Tab 50 mg25.04 14 ✓ Vimpat	
▲ Tab 100 mg50.06 14 ✓ Vimpat	
200.24 56 ✓ Vimpat	
▲ Tab 150 mg	
300.40 56 ✓ Vimpat	
▲ Tab 200 mg	

⇒SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

	Subsidy (Manufactured Dri	(aa)	Fully	
	(Manufacturer's Pri \$	ice) Sub: Per	sidised •	Generic Manufacturer
AMOTRIGINE	· · · · · · · · · · · · · · · · · · ·	-		
Tab dispersible 2 mg	55.00	30	1	Lamictal
Tab dispersible 5 mg		56		Arrow-Lamotrigine
	50.00	30		Lamictal
Fab dispersible 25 mg	2.76	56	1	Logem
Tab dispersible 50 mg		56	1	Logem
Tab dispersible 100 mg		56		Logem
Arrow-Lamotrigine Tab dispersible 5 mg to be delisted 1 Oct	ober 2020)			
EVETIRACETAM				
Tab 250 mg	4.99	60	1	Everet
Tab 500 mg	8.79	60	1	Everet
Tab 750 mg	14.39	60	1	Everet
Tab 1,000 mg	18.59	60	1	Everet
Oral liq 100 mg per ml	44.78	300 ml OP	1	Levetiracetam-AFT
HENOBARBITONE				
For phenobarbitone oral liquid refer Standard Formulae,	page 248			
F Tab 15 mg		500	1	PSM
€ Tab 30 mg		500	_	PSM
HENYTOIN SODIUM				
* Tab 50 mg	75.00	200	1	Dilantin Infatab
Cap 30 mg		200		Dilantin
Cap 100 mg		200		Dilantin
Oral liq 30 mg per 5 ml		500 ml		Dilantin
		000 1111	•	Dilantin
REGABALIN Note: Not subsidised in combination with subsidised gat	onontin			
•	•	56	./	Pregabalin Pfizer
· · · · · · · · · · · · · · · · ·		56		Pregabalin Pfizer
		56		Pregabalin Pfizer
		56		Pregabalin Pfizer
	7.30	50	٠	riegaballii riizei
RIMIDONE				
Fab 250 mg		100		Apo-Primidone
	62.00	200	/	Mysoline S29 S29
ODIUM VALPROATE				
Tab 100 mg	13.65	100	1	Epilim Crushable
Tab 200 mg EC	27.44	100		Epilim
Tab 500 mg EC	52.24	100	1	Epilim
Oral liq 200 mg per 5 ml	20.48	300 ml	1	Epilim S/F Liquid
			1	Epilim Syrup
f Inj 100 mg per ml, 4 ml	41.50	1	1	Epilim IV
TIRIPENTOL – Special Authority see SA1330 below – Reta	ail pharmacy			
• • • • • • • • • • • • • • • • • • • •		00		Di
Cap 250 mg	509.29	60	•	Diacomit S29

⇒SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Patient has confirmed diagnosis of Dravet syndrome; and



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

TOPIRAMATE

\blacktriangle	Tab 25 mg	11.07	60	Arrow-Topiramate
	-			✓ Topiramate Actavis
		26.04		✓ Topamax
\blacktriangle	Tab 50 mg	18.81	60	✓ Arrow-Topiramate
				✓ Topiramate Actavis
		44.26		✓ Topamax
\blacktriangle	Tab 100 mg	31.99	60	✓ Arrow-Topiramate
	-			✓ Topiramate Actavis
		75.25		✓ Topamax
\blacktriangle	Tab 200 mg	55.19	60	✓ Arrow-Topiramate
	•			✓ Topiramate Actavis
		129.85		✓ Topamax
\blacktriangle	Sprinkle cap 15 mg	20.84	60	✓ Topamax
\blacktriangle	Sprinkle cap 25 mg	26.04	60	✓ Topamax
VIG	GABATRIN - Special Authority see SA1907 below - Retail	pharmacy		
	Tab 500 mg	, ,	100	✓ Sabril

SA1907 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and
- 2 Fither:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
 - 2.2 It is impractical or impossible (due to comorbid conditions, or health system capacity constraints) to monitor the patient's visual fields..

Notes: ``Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages. **Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
- 2.2 It is impractical or impossible (due to comorbid conditions, or health system capacity constraints) to monitor the patient's visual fields..

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

Acute Migraine Treatment

RIZATRIPTAN			
Tab orodispersible 10 mg	3.65	30	✓ Rizamelt
Rizamelt to be Sole Supply on 1 October 2020			
SUMATRIPTAN			
Tab 50 mg	24.44	100	✓ Apo-Sumatriptan
Tab 100 mg	46.23	100	✓ Apo-Sumatriptan
Inj 12 mg per ml, 0.5 ml prefilled pen	34.00	2 OP	✓ Imigran
a) Brand switch fee payable (Pharmacode 2597330)	- see page 246 for	details	

b) Maximum of 10 inj per prescription

Prophylaxis of Migraine

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 50

PIZOTIFEN

Mr. T. I. 40

※ Tab 500 mcg......23.21 100 **✓ Sandomigran**

Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT – Special Authority see SA0987 below – Retail pharmacy

Can 2 x 80 mg and 1 x 125 mg

84 00 3 OP

 Cap 2 × 80 mg and 1 × 125 mg.......84.00
 3 OP
 ✓ Emend Tri-Pack

⇒SA0987 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetodenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

* Tab 16 mg3.88 Vergo 16 to be Sole Supply on 1 November 2020	84	♥ vergo 16
CYCLIZINE HYDROCHLORIDE Tab 50 mg0.55	10	✓ Nausicalm
CYCLIZINE LACTATE Inj 50 mg per ml, 1 ml14.95	5	✓ Nausicalm
DOMPERIDONE * Tab 10 mg2.25	100	✓ Pharmacy Health

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
HYOSCINE HYDROBROMIDE					
* Inj 400 mcg per ml, 1 ml ampoule	93.00	10	✓	Martindale S29	
Patch 1.5 mg - Special Authority see SA1927 below - Retai	ļ				
pharmacy	14.11	2	/	Scopoderm TTS	

⇒SA1927 Special Authority for Subsidy

Initial application — (control of intractable nausea, vomiting or inability to swallow saliva or clozapine induced hypersalivation) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Initial application — (pandemic circumstances- symptomatic relief of respiratory secretions in palliative care) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Requires palliative care in the community setting; and
- 2 Requires symptomatic relief of respiratory secretions that is not possible with 'as required subcutaneous hyoscine injections' due to COVID-19 constraints on the health sector; and
- 3 Access to a syringe driver for administration of injectable hyoscine is not possible due to COVID-19 constraints on the health sector.

Renewal — (control of intractable nausea, vomiting or inability to swallow saliva or clozapine induced hypersalivation) from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE		
* Tab 10 mg - Up to 30 tab available on a PSO	100	Metoclopramide Actavis 10
Metoclopramide Actavis 10 to be Sole Supply on 1 October 2020		
* Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO9.50	10	✓ Pfizer
ONDANSETRON		
* Tab 4 mg	50	✓ Onrex
* Tab disp 4 mg - Up to 10 tab available on a PSO0.76	10	 ✓ Ondansetron ODT-DRLA
Ondansetron ODT-DRLA to be Sole Supply on 1 October 2020		
* Tab 8 mg4.57	50	✓ Onrex
* Tab disp 8 mg - Up to 10 tab available on a PSO1.13	10	✓ Ondansetron ODT-DRLA
Ondansetron ODT-DRLA to be Sole Supply on 1 October 2020		
PROCHLORPERAZINE		
* Tab 3 mg buccal5.97	50	
(30.00)		Buccastem
* Tab 5 mg – Up to 30 tab available on a PSO	250	✓ Nausafix
* Inj 12.5 mg per ml, 1 ml – Up to 5 inj available on a PSO25.81	10	✓ Stemetil

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antipsychotics

General

AMISULPRIDE - Safety medicine; prescriber may determine d	lispensing frequen	су	
Tab 100 mg	5.15	30	✓ Sulprix
	17.16	100	✓ Amisulpride
			Mylan S29
Tab 200 mg	14.96	60	✓ Sulprix
Tab 400 mg		60	✓ Sulprix
ARIPIPRAZOLE – Safety medicine; prescriber may determine		nev	
Tab 5 mg		30	✓ Aripiprazole Sandoz
Tab o mg	28.58	49	✓ Aripiprazole 341402
	20.50	40	Pharma S29
Tab 40 mm	17.50	00	
Tab 10 mg		30	✓ <u>Aripiprazole Sandoz</u>
Tab 15 mg		30	✓ Aripiprazole Sandoz
Tab 20 mg		30	✓ Aripiprazole Sandoz
Tab 30 mg		30	✓ <u>Aripiprazole Sandoz</u>
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; p			
Tab 10 mg - Up to 30 tab available on a PSO	14.83	100	✓ Largactil
Tab 25 mg - Up to 30 tab available on a PSO		100	✓ Largactil
Tab 100 mg - Up to 30 tab available on a PSO		100	✓ Largactil
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO	30.79	10	✓ Largactil
CLOZAPINE - Hospital pharmacy [HP4]			
Safety medicine; prescriber may determine dispensing freq	uency		
Tab 25 mg	5.69	50	✓ Clozaril
•	6.69		✓ Clopine
	11.36	100	✓ Clozaril
	13.37		✓ Clopine
Tab 50 mg	8.67	50	✓ Clopine
	17.33	100	✓ Clopine
Tab 100 mg	14.73	50	✓ Clozaril
	17.33		✓ Clopine
	29.45	100	✓ Clozaril
	34.65		✓ Clopine
Tab 200 mg	34.65	50	✓ Clopine
	69.30	100	✓ Clopine
Suspension 50 mg per ml	17.33	100 ml	✓ Clopine
HALOPERIDOL - Safety medicine; prescriber may determine	dispensina freauer	ncv	
Tab 500 mcg - Up to 30 tab available on a PSO		100	✓ Serenace
Tab 1.5 mg - Up to 30 tab available on a PSO		100	✓ Serenace
Tab 5 mg - Up to 30 tab available on a PSO		100	✓ Serenace
Oral lig 2 mg per ml – Up to 200 ml available on a PSO		100 ml	✓ Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a I		10	✓ Serenace
LEVOMEPROMAZINE - Safety medicine; prescriber may dete			
Tab 25 mg (33.8 mg as a maleate)		100	✓ Nozinan (Swiss)
Tab 25 mg as a maleate		100	✓ Nozinan
Tab 100 mg (135 mg as a maleate)		100	✓ Nozinan (Swiss)
Tab 100 mg as a maleate		100	✓ Nozinan (Swiss)
rab rob mg ab a maioato		100	- ITOLITIUII

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price)	Subsid	Fully Brand or dised Generic
	\$	Per	✓ Manufacturer
LEVOMEPROMAZINE HYDROCHLORIDE - Safety medicine; p	rescriber may determ	nine dispens	sing frequency
Inj 25 mg per ml, 1 ml ampoule		10	✓ <u>Nozinan</u>
LITHIUM CARBONATE - Safety medicine; prescriber may deter	mine dispensing freq	uency	
Tab 250 mg - Subsidy by endorsement		500	✓ Lithicarb FC
Subsidised for patients who were taking lithium carbonat	te tab 250 mg prior to	1 January	2020 and the prescription is
endorsed accordingly. Pharmacists may annotate the pr	rescription as endors	ed where th	ere exists a record of prior
dispensing of lithium carbonate.			
Tab long-acting 400 mg		100	✓ Priadel
Cap 250 mg	9.42	100	Douglas
(Lithicarb FC Tab 250 mg to be delisted 1 November 2020)			
OLANZAPINE - Safety medicine; prescriber may determine disp	ensing frequency		
Tab 2.5 mg	1.35	28	✓ Zypine
Zypine to be Sole Supply on 1 November 2020			
Tab 5 mg	1.58	28	✓ Zypine
Zypine to be Sole Supply on 1 November 2020			<i>4</i>
Tab orodispersible 5 mg	1.81	28	Zypine ODT
Zypine ODT to be Sole Supply on 1 November 2020	0.04	00	/ 7
Tab 10 mg	2.01	28	✓ Zypine
Zypine to be Sole Supply on 1 November 2020 Tab orodispersible 10 mg	0.00	28	./ Zunina ODT
Zypine ODT to be Sole Supply on 1 November 2020	2.30	20	✓ Zypine ODT
,			
PERICYAZINE – Safety medicine; prescriber may determine dis		0.4	4 N. J. W.
Tab 2.5 mg		84	✓ Neulactil
Tab 10 mm	12.49	100	✓ Neulactil
Tab 10 mg	37.34 44.45	84 100	✓ Neulactil ✓ Neulactil
		100	Neulacui
QUETIAPINE – Safety medicine; prescriber may determine disposition			40
Tab 25 mg	2.15	90	✓ Quetapel
Quetapel to be Sole Supply on 1 November 2020	F 00	00	/ Overhead
Tab 100 mg	5.06	90	Quetapel
Quetapel to be Sole Supply on 1 November 2020 Tab 200 mg	9.00	90	✓ Quetapel
Quetapel to be Sole Supply on 1 November 2020	0.90	30	• Quetapei
Tab 300 mg	12.86	90	✓ Quetapel
Quetapel to be Sole Supply on 1 November 2020		00	- audiapoi
RISPERIDONE – Safety medicine; prescriber may determine dis	enoncina froquency		
Tab 0.5 mg		60	✓ Risperidone (Teva)
Risperidone (Teva) to be Sole Supply on 1 December 20		00	- moperiuone (revu)
Tab 1 mg		60	✓ Risperidone (Teva)
Risperidone (Teva) to be Sole Supply on 1 December 20		00	- moponaono (rova)
Tab 2 mg		60	✓ Risperidone (Teva)
Risperidone (Teva) to be Sole Supply on 1 December 20			.,
Tab 3 mg		60	✓ Risperidone (Teva)
Risperidone (Teva) to be Sole Supply on 1 December 20)20		. , ,
Tab 4 mg	3.42	60	✓ Risperidone (Teva)
Risperidone (Teva) to be Sole Supply on 1 December 20	020		
Oral liq 1 mg per ml	8.90	30 ml	✓ Risperon
Risperon to be Sole Supply on 1 November 2020			

✓ Zyprexa Relprevy

✓ Zyprexa Relprevv

	Subsidy		Fully Brand or
	(Manufacturer's Price)	_	Subsidised Generic
	\$	Per	✓ Manufacturer
ZIPRASIDONE - Safety medicine; prescriber may determine di	spensing frequency		
Cap 20 mg	14.50	60	✓ Zusdone
Cap 40 mg	24.70	60	✓ Zusdone
Cap 60 mg	33.80	60	✓ Zusdone
Cap 80 mg	39.70	60	✓ Zusdone
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pre	escriber may determin	e dis	pensing frequency
Tab 10 mg	•	100	
Depot Injections			
FLUPENTHIXOL DECANOATE - Safety medicine; prescriber n	nay determine dispens	ing f	requency
Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PSO		5	✓ Fluanxol
Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PSO	20.90	5	✓ Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	40.87	5	✓ Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescriber ma	av determine dispensi	na fre	eauencv
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO		5	✓ Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO		5	✓ Haldol Concentrate
, ,,,			✓ Haldol
			Decanoas S29
OLANZAPINE - Special Authority see SA1428 below - Retail p	harmacy		
Safety medicine; prescriber may determine dispensing frequ			
Inj 210 mg vial		1	✓ Zyprexa Relprevv

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents: and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

 Safety medicine; prescriber may determine dispensing frequency
 194.25
 1
 Invega Sustenna

 Inj 25 mg syringe
 271.95
 1
 Invega Sustenna

 Inj 75 mg syringe
 357.42
 1
 Invega Sustenna

 Inj 100 mg syringe
 435.12
 1
 Invega Sustenna

 Inj 150 mg syringe
 435.12
 1
 Invega Sustenna

 Inj 150 mg syringe
 435.12
 1
 Invega Sustenna

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE - Special Authority see SA1427 below - Retail pharmacy

Inj 25 mg vial 135.98 1 ✓ Risperdal Consta Inj 37.5 mg vial 178.71 1 ✓ Risperdal Consta Inj 50 mg vial 217.56 1 ✓ Risperdal Consta	Safety medicine; prescriber may determine dispensin	ig frequency		
,	Inj 25 mg vial	135.98	1	Risperdal Consta
Ini 50 mg vial 217 56 1 Risperdal Consta	Inj 37.5 mg vial	178.71	1	✓ Risperdal Consta
This period consultation is a second consultation of the consultat	lnj 50 mg vial	217.56	1	✓ Risperdal Consta

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

✓ Clopixol

Anxiolytics

BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	20.23	100	✓ Orion
* Tab 10 mg	13.16	100	✓ Orion
CLONAZEPAM - Safety medicine; prescriber may determine di	spensing frequency	,	
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determine dispe	nsing frequency		
Tab 2 mg	61.07	500	✓ Arrow-Diazepam
Arrow-Diazepam to be Sole Supply on 1 December 202	20		-
Tab 5 mg	73.60	500	✓ Arrow-Diazepam
Arrow-Diazepam to be Sole Supply on 1 December 202	20		-

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
LORAZEPAM – Safety medicine; prescriber may determine disp	ensing frequency			
Tab 1 mg	9.72	250	✓	<u>Ativan</u>
Tab 2.5 mg	12.50	100	1	<u>Ativan</u>
OXAZEPAM - Safety medicine; prescriber may determine disper	nsing frequency			
Tab 10 mg	6.17	100	✓ (Ox-Pam
Tab 15 mg		100	✓ (Ox-Pam

Multiple Sclerosis Treatments

⇒SA1559 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- a) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- b) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- c) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- d) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week:
 - d) start at least one month after the onset of a previous relapse;



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

continued...

- be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point:
- f) be distinguishable from the effects of general fatigue; and
- g) not be associated with a fever (T> 37.5°C); and
- e) applications must be made by the patient's neurologist or general physician; and
- f) patients must have no previous history of lack of response to dimethyl fumarate; and
- g) patients must have not previously had intolerance to dimethyl fumarate; and
- h) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to dimethyl fumarate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate, teriflunomide and ocrelizumab is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

FINGOLIMOD - Special Authority see SA1562 below - Retail pharmacy

Wastage claimable

Cap 0.5 mg.......2,200.00 28 ✓ Gilenya

⇒SA1562 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

Subsidy	Fu	lly Drand or	
Subsidy		,	
(Manufacturer's P	rice) Subsidise	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week:
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) patients must have no previous history of lack of response to fingolimod; and
- 7) patients must have not previously had intolerance to fingolimod; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to fingolimod; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate, teriflunomide and ocrelizumab is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria



Subsidy (Manufacturer's Price) Per

Fully Subsidised

Brand or Generic Manufacturer

continued...

are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

NATALIZUMAB - Special Authority see SA1563 below - Retail pharmacy

✓ Tysabri

⇒SA1563 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990 Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week:
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) treatment must be initiated and supervised by a neurologist who is registered in the Tysabri Australasian Prescribing Programme operated by the supplier: and
- 7) patients must have no previous history of lack of response to natalizumab; and

NERVOUS SYSTEM

Subsidy	F	ılly	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
\$	Per	/	Manufacturer

continued...

- 8) patients must have not previously had intolerance to natalizumab; and
 - a) Patient is JC virus negative, or
 - Patient is JC virus positive and has given written informed consent acknowledging an understanding of the risk of progressive multifocal leucoencephalopathy (PML) associated with natalizumab
- 10) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5: or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to natalizumab: or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier.

Switching between natalizumab, fingolimod, dimethyl fumarate, teriflunomide and ocrelizumab is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

OCRELIZUMAB - Special Authority see SA1867 below - Retail pharmacy

⇒SA1867 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- a) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- b) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and

_				
	Subsidy	Fully	Brand or	
	(Manufacturer's Price)	Subsidised	Generic	
	\$	Por 🗸	Manufacturer	

continued...

- c) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
 - d) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria):
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week:
 - d) start at least one month after the onset of a previous relapse;
 - be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point:
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
 - e) applications must be made by the patient's neurologist or general physician; and
 - f) patients must have no previous history of lack of response to ocrelizumab; and
 - g) patients must have not previously had intolerance to ocrelizumab; and
 - h) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5: or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to ocrelizumab: or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate, teriflunomide and ocrelizumab is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

TERIFLUNOMIDE - Special Authority see SA1560 on the next page - Retail pharmacy

Wastage claimable

NERVOUS SYSTEM

Subsidy (Manufacturer's Price) \$

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

⇒SA1560 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) patients must have no previous history of lack of response to teriflunomide; and
- 7) patients must have not previously had intolerance to teriflunomide; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

b) 1.0 to 3.0; or

c) 1.5 to 3.5; or

d) 2.0 to 4.0; or

e) 2.5 to 4.5; or

f) 3.0 to 4.5; or

g) 3.5 to 4.5; or

h) 4.0 to 4.5.

- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to teriflunomide; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate, teriflunomide and ocrelizumab is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

Other Multiple Sclerosis Treatments

GLATIRAMER ACETATE - Special Authority see SA1808 below - Retail pharmacy

⇒SA1808 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. As a new Special Authority approval number is required the MSTAC coordinator should be notified of the change and a new prescription provided. **Entry Criteria**

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- i) a gadolinium enhancing lesion; or
- ii) a Diffusion Weighted Imaging positive lesion; or
- iii) a T2 lesion with associated local swelling; or
- iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
- v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point:
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or
 - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment.
 Progression of disability is defined as progress by any of the following EDDSS Points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0: or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

⇒SA1809 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990 Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. As a new Special Authority approval number is required the MSTAC coordinator should be notified of the change and a new prescription provided. **Entry Criteria**

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- f) be distinguishable from the effects of general fatigue; and
- g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or
 - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment.
 Progression of disability is defined as progress by any of the following EDDSS Points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
- h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

⇒SA1810 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. As a new Special Authority approval number is required the MSTAC coordinator should be notified of the change and a new prescription provided.

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria):
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or
 - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

Stopping Criteria

Any of the following:

Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment.
 Progression of disability is defined as progress by any of the following EDDSS Points:

Subsidy (Manufacturer's Price)	Sı	Fully ibsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
- b) 1.0 to 3.0; or
- c) 1.5 to 3.5; or
- d) 2.0 to 4.0: or
- e) 2.5 to 4.5; or
- f) 3.0 to 4.5; or
- g) 3.5 to 4.5; or h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

Sedatives and Hypnotics

MELATONIN - Special Authority see SA1666 below - Retail pharmacy

Tab modified-release 2 mg - No more than 5 tab per day28.22

30

✓ Circadin

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	(Manufacturer's Price)	Per	Subsidised	
MIDAZOLAM - Safety medicine; prescriber may determine disper	nsing frequency			
Inj 1 mg per ml, 5 ml ampoule	4.30	10	1	Midazolam-Claris
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available			_	
on a PSO		10		Pfizer
On a PSO for status epilepticus use only. PSO must be e		epilep 5		only. Midazolam-Claris
Inj 5 mg per ml, 3 ml ampoule		5	•	WIIUaZUIaIII-Ciai iS
a PSOa		5	1	Pfizer
On a PSO for status epilepticus use only. PSO must be e		pilep		
NITRAZEPAM – Subsidy by endorsement				•
Safety medicine; prescriber may determine dispensing free	guency			
b) Subsidy by endorsement – subsidised for patients who we		prior	to 1 Augu	ust 2019 and the prescription
is endorsed accordingly. Pharmacists may annotate the p				
dispensing of nitrazepam in the preceding 12 months.			_	
Tab 5 mg	5.22	100	/	Nitrados
Nitrados Tab 5 mg to be delisted 1 January 2021)				
PHENOBARBITONE SODIUM – Special Authority see SA1386 b	•	•	_	
Inj 200 mg per ml, 1 ml ampoule	68.00	10	•	Max Health S29
SA1386 Special Authority for Subsidy				
nitial application from any relevant practitioner. Approvals valid	without further rene	wal ui	nless notif	ied for applications meeting
ne following criteria: Hoth:				
For the treatment of terminal agitation that is unresponsive	to other agents: and	1		
2 The applicant is part of a multidisciplinary team working in		•		
EMAZEPAM – Safety medicine; prescriber may determine dispe	•			
Tab 10 mg	0 1 1	25	1	Normison
Normison to be Sole Supply on 1 November 2020	1.00	20	•	TTOTTIII OOTI
TRIAZOLAM – Safety medicine; prescriber may determine disper	nsina fraguancy			
Tab 125 mcg	•	100		
	(9.85)			Hypam
Tab 250 mcg	4.10 [′]	100		••
	(11.20)			Hypam
OPICLONE - Safety medicine; prescriber may determine disper	nsing frequency			
T-1-75	0.50	E00	,	Tandalana Askasila

500

✓ Zopiclone Actavis

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Generic
Stimulants/ADHD Treatments				
ATOMOXETINE				
Cap 10 mg	18.41	28	/	Generic Partners
	(107.03)			Strattera
Generic Partners to be Sole Supply on 1 December 2020)			
Cap 18 mg	27.06	28	✓	Generic Partners
	(107.03)			Strattera
Generic Partners to be Sole Supply on 1 December 2020)			
Cap 25 mg	29.22	28	✓	Generic Partners
	(107.03)			Strattera
Generic Partners to be Sole Supply on 1 December 2020			_	
Cap 40 mg		28	/	Generic Partners
	(107.03)			Strattera
Generic Partners to be Sole Supply on 1 December 2020			_	
Cap 60 mg		28	•	Generic Partners
	(107.03)			Strattera
Generic Partners to be Sole Supply on 1 December 2020				
Cap 80 mg		28	•	Generic Partners
Occasio Bertanas te ha Octa Occasio and December 2006	(139.11)			Strattera
Generic Partners to be Sole Supply on 1 December 2020		28		Generic Partners
Cap 100 mg		28	•	Strattera
Generic Partners to be Sole Supply on 1 December 2020	(139.11)			Strattera
,)			
(Strattera Cap 10 mg to be delisted 1 December 2020)				
(Strattera Cap 18 mg to be delisted 1 December 2020) (Strattera Cap 25 mg to be delisted 1 December 2020)				
(Strattera Cap 40 mg to be delisted 1 December 2020)				
(Strattera Cap 40 mg to be delisted 1 December 2020)				
(Strattera Cap 80 mg to be delisted 1 December 2020)				
(Strattera Cap 100 mg to be delisted 1 December 2020)				
,	alow Datail nharma			
DEXAMFETAMINE SULFATE - Special Authority see SA1149 b	elow – netali priarma	СУ		
a) Only on a controlled drug formb) Safety medicine; prescriber may determine dispensing fre	allopov			
		100	.,	DCM
Tab 5 mg	20.00	100	•	<u>PSM</u>

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:



Subsidy (Manufacturer's Price)	Fully Subsidised	
 \$	Per 🗸	Manufacturer

continued...

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1150 below - Retail pharmacy

- a) Only on a controlled drug form

50.00 (Ritalin SR Tab sustained-release 20 mg to be delisted 1 June 2021)

⇒SA1150 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — **(ADHD in patients 5 or over)** only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

continued...

✓ Ritalin SR

100

Subsid	dy Fi	ılly Brand or	
(Manufacture	r's Price) Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1151 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine: prescriber may determine dispensing frequency

Tab extended-release 18 mg	18.20	30	✓ Methylphenidate ER - Teva
	58.96		✓ Concerta
Tab extended-release 27 mg	22.00	30	✓ Methylphenidate ER - Teva
	65.44		✓ Concerta
Tab extended-release 36 mg	22.40	30	Methylphenidate ERTeva
	71.93		✓ Concerta
Tab extended-release 54 mg	26.40	30	Methylphenidate ERTeva
	86.24		✓ Concerta
Cap modified-release 10 mg	15.60	30	✓ Ritalin LA
Cap modified-release 20 mg	20.40	30	✓ Ritalin LA
Cap modified-release 30 mg	25.52	30	✓ Ritalin LA
Cap modified-release 40 mg		30	Ritalin LA

⇒SA1151 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

Subsidy	Fi	ılly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
\$	Per	/	Manufacturer

continued...

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

./ Madaviali

MODAFINIL - Special Authority see SA1932 below - Retail pharmacy

♥ iwodavigii	30	1ab 100 mg32.00
✓ Modaviqil	60	64.00

⇒SA1932 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Any of the following:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 A multiple sleep latency test is not possible due to COVID-19 constraints on the health sectors; or
 - 2.3 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE		
* Tab 5 mg4.34	90	✓ Donepezil-Rex
Donepezil-Rex to be Sole Supply on 1 December 2020		
* Tab 10 mg6.64	90	Donepezil-Rex
Donepezil-Rex to be Sole Supply on 1 December 2020		
RIVASTIGMINE - Special Authority see SA1488 below - Retail pharmacy		
Patch 4.6 mg per 24 hour48.75	30	✓ Generic Partners
Patch 9.5 mg per 24 hour48.75	30	✓ Generic Partners

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate: and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

28

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 8 mg with naloxone 2 mg53.12

✓ <u>Buprenorphine</u> Naloxone BNM

✓ Buprenorphine Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (**Detoxification**) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

Tab modified-release 150 mg......11.00 30 ✓ Zyban

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
DISULFIRAM Tab 200 mg	153.00	100	✓ A	ntabuse
NALTREXONE HYDROCHLORIDE – Special Authority see SA1 Tab 50 mg		harmacy 30		altraccord

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

a) Nicotine will not be funded in amounts less than 4 weeks of treatment.

 b) Note: Direct Provision by a pharmacist permitted under the p 	rovisions in	Part I of Section	n A.
Patch 7 mg - Up to 28 patch available on a PSO	18.14	28	Habitrol
Patch 7 mg for direct distribution only - [Xpharm]	3.94	7	Habitrol
Patch 14 mg - Up to 28 patch available on a PSO	19.95	28	Habitrol
Patch 14 mg for direct distribution only - [Xpharm]	4.52	7	Habitrol
Patch 21 mg - Up to 28 patch available on a PSO	22.86	28	Habitrol
Patch 21 mg for direct distribution only - [Xpharm]	5.18	7	Habitrol
Lozenge 1 mg - Up to 216 loz available on a PSO	19.18	216	Habitrol
Lozenge 1 mg for direct distribution only - [Xpharm]	3.20	36	Habitrol
Lozenge 2 mg - Up to 216 loz available on a PSO	21.02	216	Habitrol
Lozenge 2 mg for direct distribution only - [Xpharm]	3.24	36	Habitrol
Gum 2 mg (Fruit) - Up to 384 piece available on a PSO	38.21	384	Habitrol
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]	8.64	96	Habitrol
Gum 2 mg (Mint) - Up to 384 piece available on a PSO	38.21	384	Habitrol
Gum 2 mg (Mint) for direct distribution only - [Xpharm]	8.64	96	Habitrol
Gum 4 mg (Fruit) - Up to 384 piece available on a PSO	44.17	384	Habitrol
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]	10.01	96	Habitrol
Gum 4 mg (Mint) - Up to 384 piece available on a PSO	44.17	384	Habitrol
Gum 4 mg (Mint) for direct distribution only - [Xpharm]	10.01	96	Habitrol

VARENICLINE TARTRATE - Special Authority see SA1845 on the next page - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	25.64	53 OP	✓ Varenicline Pfizer
Tab 1 mg	27.10	56	✓ Varenicline Pfizer

Subsidy		Fully	Brand or	
(Manufacturer's Price)			Generic	
\$	Per	•	Manufacturer	

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see SA1667 below

Inj 25 mg vial	 	271.35	1	✓ Ribomustin
Inj 100 mg vial	 	1,085.38	1	✓ Ribomustin
Inj 1 mg for ECP	 	11.40	1 mg	✓ Baxter

⇒SA1667 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 All of the following:
 - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
 - 3.2.2 The patient has not received prior bendamustine therapy; and
 - 3.2.3 Fither:
 - 3.2.3.1 Both:
 - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more: or
 - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Either:
 - 2.1 Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

- 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.
 Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

BUSULFAN – PCT – Retail pharmacy-Specialist Tab 2 mg	90.05	100	✓ Myleran
•	09.20	100	♥ Wyleran
CARBOPLATIN – PCT only – Specialist	20.50	1	✓ DDI Carbanistin
Inj 10 mg per ml, 45 ml vial	32.59 45.20	ı	✓ DBL Carboplatin✓ Carboplatin Ebewe
	48.50		✓ Carbopiatiii Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist		9	
Inj 100 mg vial	1 387 00	1	✓ BiCNU
ing 100 mg viai			✓ Bicnu Heritage S29
Inj 100 mg for ECP	1,387.00	100 mg OP	✓ Baxter
CHLORAMBUCIL - PCT - Retail pharmacy-Specialist			
Tab 2 mg	29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	12.29	1	✓ DBL Cisplatin
, , ,	15.00		✓ Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial	19.70	1	✓ DBL Cisplatin
, .,	21.00		✓ Cisplatin Ebewe
Inj 1 mg for ECP	0.25	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE			
Tab 50 mg - PCT - Retail pharmacy-Specialist	79.00	50	✓ Endoxan S29
	158.00	100	✓ Procytox S29
Wastage claimable			•
Inj 1 g vial - PCT - Retail pharmacy-Specialist	35.65	1	✓ Endoxan
	127.80	6	✓ Cytoxan
Inj 2 g vial – PCT only – Specialist		1	✓ Endoxan
Inj 1 mg for ECP - PCT only - Specialist	0.04	1 mg	✓ Baxter
IFOSFAMIDE - PCT only - Specialist			
lnj 1 g		1	✓ Holoxan
lnj 2 g		1	✓ Holoxan
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
LOMUSTINE - PCT - Retail pharmacy-Specialist			
Cap 10 mg		20	✓ CeeNU
Cap 40 mg	399.15	20	✓ CeeNU
MELPHALAN			
Tab 2 mg - PCT - Retail pharmacy-Specialist		25	✓ Alkeran
Inj 50 mg - PCT only - Specialist	67.80	1	✓ Alkeran
	213.00		✓ Alkeran s29 S29
	420.00		✓ Tillomed S29

	Subsidy		Fully	Brand or
	(Manufacturer's Price	e) (e	Subsidised	
	\$	Per	✓	Manufacturer
OXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	✓	Oxaliplatin Actavis 100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	46.32	1		Oxaliplatin Accord
Inj 1 mg for ECP	0.48	1 mg	1	Baxter
THIOTEPA - PCT only - Specialist		-		
Inj 15 mg vial	CBS	1	1	Bedford \$29
, - 3			1	THIO-TEPA S29
			_	Tepadina S29
Inj 100 mg vial	CBS	1		Tepadina S29
inj 100 mg vidi				repudina
Antimetabolites				
AZACITIDINE - PCT only - Specialist - Special Authority see	SA1467 below			
Inj 100 mg vial		1	1	Azacitidine Dr
, · · · · · · · · · · · · · · · · · ·		•		Reddy's
	605.00		1	Vidaza

⇒SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or

1 ma

✓ Baxter

- 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and

- 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- 4 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

	Subsidy (Manufacturer's Price)	20) C··	Fully Brand or
	(Manufacturer's Prior	ce) Su Per	bsidised Generic Manufacturer
CALCIUM FOLINATE			
Tab 15 mg – PCT – Retail pharmacy-Specialist	114.69	10	✓ DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17.10	5	✓ Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specia		1	✓ <u>Calcium Folinate</u> Sandoz
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	9.49	1	✓ Calcium Folinate Sandoz
Inj 100 mg - PCT only - Specialist	7.33	1	✓ Calcium Folinate Ebewe
Inj 300 mg - PCT only - Specialist	22.51	1	✓ Calcium Folinate Ebewe
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	25.14	1	✓ Calcium Folinate Sandoz
Inj 1 g - PCT only - Specialist	67.51	1	✓ Calcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	72.00	1	✓ Calcium Folinate Sandoz
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	✓ Baxter
CAPECITABINE – Retail pharmacy-Specialist	40.00	00	()
Tab 500 mg		60 120	✓ <u>Capercit</u> ✓ Capercit
Tab 500 mg	49.00	120	Capercit
CLADRIBINE - PCT only - Specialist Inj 1 mg per ml, 10 ml	7/0.06	1	✓ Leustatin
Inj 10 mg for ECP		10 mg OP	✓ Baxter
CYTARABINE			
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia	alist400.00	5	✓ Pfizer
Inj 100 mg per ml, 20 ml vial – PCT – Retail		•	
pharmacy-Specialist	41.36	1	✓ Pfizer
Inj 1 mg for ECP - PCT only - Specialist		10 mg	✓ Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Specia	alist80.00	100 mg OP	✓ Baxter
FLUDARABINE PHOSPHATE			
Tab 10 mg - PCT - Retail pharmacy-Specialist		20	✓ Fludara Oral
Inj 50 mg vial – PCT only – Specialist		5	✓ Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist	115.29	50 mg OP	✓ Baxter
FLUOROURACIL			4
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist		1	✓ Fluorouracil Ebewe
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist		1 100 mg	✓ Fluorouracil Ebewe✓ Baxter
Inj 1 mg for ECP – PCT only – Specialist	0.00	100 mg	Daxiei
GEMCITABINE HYDROCHLORIDE – PCT only – Specialist	60.50	1	✓ DPI Comeitabine
Inj 1 g, 26.3 ml vial		1	✓ DBL Gemcitabine ✓ Gemcitabine Flower
Inj 1 mg for ECP		1 mg	 ✓ Gemcitabine Ebewe ✓ Baxter
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist	0.02	illy	· DUALGI
Inj 20 mg per ml, 5 ml vial	71 44	1	✓ Irinotecan
ing 20 mg por mi, o mi viai		I	Accord \$29
			✓ Irinotecan Actavis
			100
lei 4 mm for FOR	100.00	4.	✓ Irinotecan-Rex
Inj 1 mg for ECP	0.75	1 mg	✓ Baxter

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Prio \$	e) S Per	Fully Subsidised	Brand or Generic Manufacturer
MERCAPTOPURINE Tab 50 mg - PCT - Retail pharmacy-Specialist	37.00	25	√ P	uri-nethol
Oral suspension 20 mg per ml – Retail pharmacy-Specialist Special Authority see SA1725 below		100 ml O	P ✓ A	llmercap

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

METHOTREXATE

MF	THOTREXATE		
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist8.05	90	✓ Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist31.75	90	✓ Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist47.50	5	✓ Hospira
*	Inj 7.5 mg prefilled syringe14.61	1	✓ Methotrexate
			Sandoz
*	Inj 10 mg prefilled syringe14.66	1	✓ Methotrexate
	, 0, , 0		Sandoz
*	Inj 15 mg prefilled syringe14.77	1	✓ Methotrexate
•	, 10g p. 6 9	·	Sandoz
*	Inj 20 mg prefilled syringe14.88	1	✓ Methotrexate
•••	11) 20 119 promos oyringo	•	Sandoz
*	Inj 25 mg prefilled syringe14.99	1	✓ Methotrexate
-,-	This 20 mg promited dynnigo	•	Sandoz
*	Inj 30 mg prefilled syringe	1	✓ Methotrexate
~	ing 50 mg premied syringe13.09	'	Sandoz
*	Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist30.00	5	✓ DBL Methotrexate
不	ing 25 mg per mi, 2 mi viai – PCT – netali pharmacy-specialist50.00	5	Onco-Vial
	Ini OF man and OO milaid ROT Detail abanessas Considiet 45 00		✓ DBL Methotrexate
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist45.00	1	Onco-Vial
.1.	lei 400 m a manuel 40 ml - BOT - Batallahama an Occasiolist - OF 00	_	*****
*	Inj 100 mg per ml, 10 ml – PCT – Retail pharmacy-Specialist25.00	1	Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail		
	pharmacy-Specialist	1	Methotrexate Ebewe
	Methotrexate Ebewe to be Sole Supply on 1 October 2020		4
*	Inj 1 mg for ECP – PCT only – Specialist	1 mg	✓ Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist4.73	5 mg OP	✓ Baxter
PΕ	METREXED - PCT only - Specialist - Special Authority see SA1679 below		
	Inj 100 mg vial60.89	1	 Juno Pemetrexed
	Inj 500 mg vial217.77	1	 Juno Pemetrexed
	Inj 1 mg for ECP	1 mg	✓ Baxter

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant

Subsidy (Manufacturer's Price)	Fu Subsidis	,	
\$	Per	 Manufacturer 	

continued...

specialist. Approvals valid for 8 months for applications meeting the following criteria:

- All of the following:
 - 1 No evidence of disease progression; and
 - 2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

Tab 40 mg	126.31	25	✓ Lanvis	
Other Cytotoxic Agents				

AMSACRINE - PCT only - Specialist			
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	✓ Amsidine S29
	4,736.00		✓ Amsidine S29
Inj 75 mg	1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-S	Specialist		
Cap 0.5 mg	CBS	100	✓ Agrylin S29 S29
, ,			✓ Teva S29
	1,175.87		✓ Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml vial	4,817.00	10	✓ Phenasen
Inj 10 mg for ECP	481.70	10 mg OP	✓ Baxter
BLEOMYCIN SULPHATE - PCT only - Specialist			
Inj 15,000 iu, vial	161.01	1	✓ DBL Bleomycin Sulfate
Inj 1,000 iu for ECP	12.45	1,000 iu	✓ Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority see		next page	
Inj 3.5 mg vial		1	✓ Bortezomib
, ,			Dr-Reddy's
Ini 1 ma for ECP	31.20	1 ma	✓ Baxter

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	/	Manufacturer	

⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

1 The patient has symptomatic multiple myeloma; or			
2 The patient has symptomatic systemic AL amyloidosis *.			
Note: Indications marked with * are unapproved indications.			
COLASPASE [L-ASPARAGINASE] – PCT only – Specialist			
Inj 10,000 iu		1	Leunase
Inj 10,000 iu for ECP	102.32	10,000 iu OP	✓ Baxter
(Leunase Inj 10,000 iu to be delisted 1 December 2020)			
(Baxter Inj 10,000 iu for ECP to be delisted 1 December 2020)			
DACARBAZINE - PCT only - Specialist			
Inj 200 mg vial	62.70	1	✓ DBL Dacarbazine
, •	580.60	10	 Dacarbazine
			APP S29
Inj 200 mg for ECP	62.70	200 mg OP	✓ Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist		_00g 0.	
Inj 0.5 mg vial	255.00	1	✓ Cosmegen
Inj 0.5 mg for ECP		0.5 mg OP	✓ Baxter
	255.00	0.5 mg OF	Daxiei
DAUNORUBICIN – PCT only – Specialist	4 40 50		4 D#
Inj 2 mg per ml, 10 ml		1	✓ Pfizer
Inj 20 mg for ECP	149.50	20 mg OP	✓ Baxter
DOCETAXEL - PCT only - Specialist			
Inj 10 mg per ml, 2 ml vial	12.40	1	✓ DBL Docetaxel
Inj 20 mg		1	 Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial		1	✓ DBL Docetaxel
Inj 20 mg per ml, 4 ml vial	26.95	1	Docetaxel
			Accord \$29
Inj 80 mg	195.00	1	✓ Docetaxel Sandoz
Inj 1 mg for ECP		1 mg	✓ Baxter
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist		· ·	
Inj 2 mg per ml, 5 ml vial	10.00	1	✓ Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	✓ Doxorubicin Ebewe
11) 2 11g por 111, 20 111 val	17.00	•	✓ Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial		1	✓ Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	✓ Doxorubicin Ebewe
.,g p,	65.00	•	✓ Arrow-Doxorubicin
Inj 1 mg for ECP	0.29	1 mg	✓ Baxter
EPIRUBICIN HYDROCHLORIDE – PCT only – Specialist		9	
Inj 2 mg per ml, 5 ml vial	25.00	1	✓ Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	✓ Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	✓ Epirubicin Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
		i ilig	Duxter
ETOPOSIDE	040.70	00	/ Vanasid
Cap 50 mg - PCT - Retail pharmacy-Specialist		20	✓ <u>Vepesid</u>
Cap 100 mg - PCT - Retail pharmacy-Specialist		10	✓ <u>Vepesid</u>
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialis		1	✓ Rex Medical
Inj 1 mg for ECP - PCT only - Specialist	0.09	1 mg	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	d Generic
ETOPOSIDE PHOSPHATE – PCT only – Specialist Inj 100 mg (of etoposide base) Inj 1 mg (of etoposide base) for ECP		1 1 mg	_	Etopophos Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] – PCT – Retail pharm Cap 500 mg	, ,	100		Devatis Hydrea
(Hydrea Cap 500 mg to be delisted 1 February 2021)				•
IDARUBICIN HYDROCHLORIDE Inj 5 mg vial – PCT only – Specialist Inj 10 mg vial – PCT only – Specialist Inj 1 mg for ECP – PCT only – Specialist	198.00	1 1 1 mg	✓	Zavedos Zavedos Baxter
LENALIDOMIDE – Retail pharmacy-Specialist – Special Authority Wastage claimable	see SA1897 below	1		
Cap 5 mg	5,122.76	28	✓	Revlimid
Cap 10 mg		21		Revlimid
	6,207.00	28		Revlimid
Cap 15 mg		21	_	Revlimid
	7,239.18	28		Revlimid
Cap 25 mg	7,627.00	21	/	Revlimid

⇒SA1897 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Fither:
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 The patient has ECOG performance score of 0-1; and
- 5 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
\$	Per	1	Manufacturer

continued...

recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MFSNA

Tab 400 mg - PCT - Retail pharmacy-Specialist314.00	50	✓ Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist448.50	50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist177.45	5 15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist407.40) 15	✓ Uromitexan
Inj 1 mg for ECP - PCT only - Specialist2.96	100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist		
Inj 5 mg vial851.37	' 1	✓ Teva
Inj 20 mg vial816.32	2 1	✓ Omegapharm S29
Inj 1 mg for ECP175.38	3 1 mg	✓ Baxter
(Omegapharm S29 Inj 20 mg vial to be delisted 1 November 2020)		
MITOZANTRONE - PCT only - Specialist		
Inj 2 mg per ml, 10 ml vial97.50) 1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP5.51	1 mg	✓ Baxter
OLAPARIB - Retail pharmacy-Specialist - Special Authority see SA1883 bel	low	
Tab 100 mg3,701.00	56	✓ Lynparza
Tab 150 mg3,701.00	56	✓ Lynparza
Cap 50 mg - Wastage claimable7,402.00) 448	✓ Lynparza

⇒SA1883 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Patient has received at least two lines of previous treatment with platinum-based chemotherapy; and
- 4 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line of platinum-based chemotherapy; and
- 5 Patient's disease must have achieved partial or complete response to treatment with the immediately preceding platinum-based regimen; and
- 6 Patient's disease has not progressed following prior treatment with olaparib; and
- 7 Treatment will be commenced within 8 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 8 Treatment to be administered as maintenance treatment; and
- 9 Treatment not to be administered in combination with other chemotherapy.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Treatment remains clinically appropriate and patient is benefitting from treatment; and

Subsidy		Fully	Brand or	Т
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2 No evidence of progressive disease; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy.

Note: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

PACLITAXEL - PCT only - Specialist

PACLITAXEL - PCT only - Specialist			
Inj 30 mg	47.30	5	✓ Paclitaxel Ebewe
Inj 100 mg	24.00	1	Paclitaxel Ebewe
•	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
•	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg	44.00	1	✓ Paclitaxel Ebewe
•	275.00		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see SA13	325 below		
Inj 750 iu per ml, 5 ml vial	3,455.00	1	✓ Oncaspar LYO S29

⇒SA1325 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and
- 3 Treatment is with curative intent.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol; and
- 3 Treatment is with curative intent.

PENTOSTATIN [DEOXYCOFORMYCIN]	- PCT	only - 9	Spacialist
PENTOSTATIN [DEOXTGOPONINTGIN]	- FUI	Offiny — 3	opecialist

Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - R	etail pharmacy-Specialist		
Cap 50 mg	980.00	50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA174	1 on the next page – Retail phar	rmacy	
Cap 5 mg	9.13	5	✓ <u>Temaccord</u>
Cap 20 mg	16.38	5	✓ <u>Temaccord</u>
	18.30		✓ Apo-Temozolomide
	136.00	14	✓ Accord S29
Cap 100 mg	35.98	5	✓ Temaccord
	40.20		✓ Apo-Temozolomide
	532.00	14	✓ Accord S29
Cap 140 mg	50.12	5	✓ <u>Temaccord</u>
	400.00		✓ Amneal S29
Cap 180 mg	620.00	14	✓ Accord S29
Cap 250 mg	86.34	5	✓ Temaccord
	688.00		✓ Amneal S29

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

- Either:
 - 1 The patient has multiple myeloma; or
 - 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

✓ Vesanoid	100	Cap 10 mg - PCT - Retail pharmacy-Specialist479.50
	elow	VENETOCLAX – Retail pharmacy-Specialist – Special Authority see SA1868 bel
✓ Venclexta	42 OP	Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg
✓ Venclexta	14 OP	Tab 10 mg95.78
✓ Venclexta	7 OP	Tab 50 mg239.44
✓ Venclexta	120	Tab 100 mg - Wastage claimable

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

	Subsidy acturer's Price) \$	Per	Fully Subsidised	
VINBLASTINE SULPHATE				
Inj 1 mg per ml, 10 ml vial - PCT - Retail pharmacy-Specialist27	70.37	5	1	DBL Vinblastine S29
			1	Hospira
Inj 1 mg for ECP - PCT only - Specialist	.6.00	1 mg	1	Baxter
VINCRISTINE SULPHATE				
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist	74.52	5	/	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist10)2.73	5	•	DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist	12.60	1 mg	1	Baxter
VINORELBINE - PCT only - Specialist				
Inj 10 mg per ml, 1 ml vial	12.00	1	1	Navelbine
	42.00		1	Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial	56.00	1	✓	Navelbine
2.	10.00		•	Vinorelbine Ebewe
Inj 1 mg for ECP	.1.25	1 mg	•	Baxter

Protein-tyrosine Kinase Inhibitors

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA1805 below - Retail pharmacy

Wastage claimable			
Tab 20 mg	3,774.06	60	✓ Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
Tab 70 mg	7,692.58	60	✓ Sprycel

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and

(Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	Subsidy	Fully	Brand or
\$ Per ✓ Manufacturer	(Manufacturer's Price)		
	 \$	Per 🗸	Manufacturer

continued...

- 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day: or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
 - 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB	 Retail pharmacy-Specialist 	 Special Authority se 	e SA1915 below

Tab 100 mg	764.00	30	Tarceva
Tab 150 mg	1,146.00	30	✓ Tarceva

⇒SA1915 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Either:
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

GEEITINIR - Rotail ph	armacy-Specialist _	Special Authority con	SA1916 on the next page	αn
Carrillinio – Reiali on	iarmacy-Soecialisi -		SATATO OF THE DEXLOR	ııe.

Tab 2	0 mg	1,700.00	30	✓ Iressa

Subsid	y Full	/ Brand or
(Manufacturer	's Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

⇒SA1916 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Fither:
 - 2.1 Patient is treatment naive: or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Gefitinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

IMATINIB MESILATE

Note: Imatinib-AFT is not a registered for the treatment of Gastro Intestinal Stromal Tumours (GIST). The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule.

Tab 100 mg - [Xpharm] - Special Authority see SA1460

	below	2,400.00	60	Glivec
*	Cap 100 mg		60	✓ Imatinib-AFT
*	Cap 400 mg	197.50	30	✓ Imatinib-AFT

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 PHARMAC Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

Special Authority criteria for GIST - access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA1191	on the next page - Ret	ail pharmacy		
Tab 250 mg	1,899.00	70	1	Tykerb
(Tykerh Tah 250 mg to be delicted 1 June 2021)				-

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

⇒SA1191 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient has not previously received trastuzumab treatment for HER 2 positive metastatic breast cancer; and
 - 1.3 Lapatinib not to be given in combination with trastuzumab; and
 - 1.4 Lapatinib to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2.2 The patient started trastuzumab for metastatic breast cancer but discontinued trastuzumab within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on trastuzumab; and
 - 2.4 Lapatinib not to be given in combination with trastuzumab; and
 - 2.5 Lapatinib to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

Wastage claimable

Cap 150 mg	4,680.00	120	Tasigna
Cap 200 mg	6,532.00	120	Tasigna

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Either:
 - 2.1 Patient has documented CML treatment failure* with imatinib; or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PALBOCICLIB – Retail pharmacy-Specialist – Special Authority Wastage claimable	see SA1894 below			
Cap 75 mg	4 000 00	21	√ 1	brance
Cap 100 mg	•	21	-	brance
Cap 125 mg	•	21	-	brance

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Fither:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state: and
- 4.2.2 Either:
 - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
 - 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

PAZOPANIB - Special Authority see SA1190 below - Retail pharmacy

Tab 200 mg	1,334.70	30	✓ Votrient
Tab 400 mg	2,669.40	30	✓ Votrient

⇒SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and

(Mani	Subsidy	Fully	Brand or
	ufacturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB – Special Authority see SA1890 below – Retail pharmacy

wasiaye ciaimable			
Tab 5 mg	2,500.00	56	Jakavi
Tab 15 mg	5,000.00	56	Jakavi
Tab 20 mg	·	56	Jakavi

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS: and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB – Special Authority see SA1917 on the next page – Retail pharmacy	
Cap 12.5 mg2,315.38	28

Cap 12.5 mg	2,315.38	28	Sutent
Cap 25 mg	4,630.77	28	Sutent
Cap 50 mg	9.261.54	28	Sutent

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

⇒SA1917 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
- 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 82

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA1914 below

Wastage claimable

⇒SA1914 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of clinical disease progression; and
- 2 No initiation of taxane chemotherapy with abiraterone; and
- 3 The treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
BICALUTAMIDE Tab 50 mg	3.80	28	✓ Bi	inarex
FLUTAMIDE Tab 250 mg		100	./ EI	lutamin
FULVESTRANT - Retail pharmacy-Specialist - Special Authority	see SA1895 below	100	▼ F1	lutaniiii
Inj 50 mg per ml, 5 ml prefilled syringe	1,068.00	2	✓ Fa	aslodex

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

MEGESTROL ACETATE			
Tab 160 mg	63.53	30	✓ Apo-Megestrol
OCTREOTIDE			
Inj 100 mcg per ml, 1 ml ampoule	18.69	5	✓ Octreotide GH S29
Inj 50 mcg per ml, 1 ml vial		5	✓ DBL Octreotide
, ,			✓ Octreotide MaxRx \$29
Inj 100 mcg per ml, 1 ml vial	18.69	5	✓ DBL Octreotide
Inj 500 mcg per ml, 1 ml ampoule		5	✓ Octreotide GH S29
Inj 500 mcg per ml, 1 ml vial		5	✓ DBL Octreotide
, ,	222.00		✓ Octreotide (Sun) S29
Inj 50 mcg per ml, 1 ml ampoule	30.64	5	✓ Octreotide GH S29
OCTREOTIDE LAR (SOMATOSTATIN ANALOGUE) - Spec	cial Authority see SA1	918 below -	- Retail pharmacy
Inj LAR 10 mg prefilled syringe		1	✓ Sandostatin LAR
Inj LAR 20 mg prefilled syringe		1	✓ Sandostatin LAR
Inj LAR 30 mg prefilled syringe		1	✓ Sandostatin LAR

⇒SA1918 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery: or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly: and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and

Tamoxifen Sandoz to be Sole Supply on 1 November 2020

Tamoxifen Sandoz to be Sole Supply on 1 November 2020

	Subsidy (Manufactured Price)	0.	Fully	Brand or
	(Manufacturer's Price) \$	Per	ibsidised •	Generic Manufacturer
continued 3 The regular Special Authority renewal requirements canno	t be met due to COV	'ID-19 c	onstraints	on the health sector.
TAMOXIFEN CITRATE * Tab 10 mg	15.00	60	√ T	amoxifen Sandoz

Aromatase Inhibitors

ANASTROZOLE		_
* Tab 1 mg5.04	30	✓ Rolin
EXEMESTANE		
* Tab 25 mg	30	✓ Pfizer Exemestane
LETROZOLE		
* Tab 2.5 mg	30	✓ <u>Letrole</u>

Immunosuppressants

A Z A TUI O D D I N I E

Cytotoxic Immunosuppressants

AZATRIOPRINE			
* Tab 25 mg	7.35	60	Azamun
* Tab 50 mg		100	✓ Azamun
* Inj 50 mg vial1		1	✓ Imuran
MYCOPHENOLATE MOFETIL			
Tab 500 mg	35.90	50	Cellcept
Cap 250 mg	35.90	100	✓ Cellcept
Powder for oral liq 1 g per 5 ml — Subsidy by endorsement1	87.25	165 ml OP	 Cellcept
Mysenhanolate neurolar for availiantid to subsidiated only for noti-	المصور والمصاد	to ourollour to	blota and cancul

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

Fusion Proteins

ETANERCEPT - Special Authority see SA1949 below - Retail	pharmacy		
Inj 25 mg	690.00	4	Enbrel
Inj 50 mg autoinjector	1,050.00	4	✓ Enbrel
Ini 50 ma prefilled syringe	1.050.00	4	✓ Enbrel

⇒SA1949 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and

continued...

✓ Tamoxifen Sandoz

60

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	•	Manufacturer

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest: and
 - 2.3 Patient has bilateral sacroillitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient diagnosed with Juvenile Idiopathic Arthritis (JIA); and
 - 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
 - 2.5 Both:
 - 251 Fither
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender ioints: or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

Renewal — (juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Applicant is a named specialist or rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
\$	Per	1	Manufacturer

continued...

- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for psoriatic arthritis; and
 - 12 Fither
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for psoriatic arthritis: or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Fither:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Subsidy	Fu	ılly	Brand or	
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold: or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints;
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Fither:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

All of the following:

- 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
 - 2.1.2 Fither:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Fither

Subsidy	Fu	ılly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
 \$	Per	•	

continued...

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Specia	alist				
Inj 50 mg per ml, 5 ml	2,351.25	5	✓ ATGAM		
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT only					
Subsidised only for bladder cancer.					
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE		
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG S29		
(SII-Onco-BCG S29 Inj 40 mg per ml, vial to be delisted 1 April 2021)					

Monoclonal Antibodies

ADALIMUMAB - Special Authority see SA1950 below -	Retail pharmacy		
Inj 20 mg per 0.4 ml prefilled syringe	1,599.96	2	Humira
Inj 40 mg per 0.8 ml prefilled pen	1,599.96	2	✓ HumiraPen
Inj 40 mg per 0.8 ml prefilled syringe	1,599.96	2	Humira

⇒SA1950 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm

25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm

35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm

55-64 years - Male: 5.5 cm; Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	
\$	Per 🗸	Manufacturer

continued...

- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or

Subsidy (Manufacturer's Price)	Fully Subsidised		
\$	Per 🗸	Manufacturer	

continued...

- 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Fither:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 CDAI score is 150 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — **(Crohn's disease - children)** only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 PCDAI score is 15 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's disease.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 The patient has 3 or more active lesions (e.g. inflammatory nodules, abscesses, draining fistulae); and
- 4 The patient has a Dermatology Quality of Life Index of 10 or more and the assessment is no more than 1 month old at time of application; and
- 5 Following the initial loading doses, adalimumab is to be administered at doses no greater than 40mg every 7 days.

Renewal — (hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for juvenile idiopathic arthritis (JIA); and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept: or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for juvenile idiopathic arthritis; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
 - 2.2 Patient diagnosed with JIA; and
 - 2.3 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
 - 2.5 Both:
 - 2.5.1 Fither:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender joints; or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

Renewal — (juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a named specialist or rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a named specialist or rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Fither:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Fither:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
 \$	Per	•	

continued...

- 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
- 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis; or
- 2 All of the following:

Subsidy (Manufacturer's Price)	F Subsidis	ully	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Fither:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Fither:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 The patient has severe Behcet's disease that is significantly impacting the patient's quality of life (see Notes); and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Either:
 - 2.1 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has not responded adequately to treatment with infliximab (see Notes); or
 - 2.2 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has experienced intolerable side effects from treatment with infliximab; and
- 3 The patient is experiencing significant loss of quality of life; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: Behcet's disease diagnosed according to the International Study Group for Behcet's disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al, J Rheumatol. 2004;31:931-7.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Fither

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1.1 Applicant is a dermatologist; or
- 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 212 Fither
 - 2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline values; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values: or
 - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value: and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

iti ici.

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

AFLIBERCEPT - Special Authority see SA1772 below - Retail pharmacy

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Fither:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting

	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	✓	Manufacturer

continued...

the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 below

Inj 5 mg per ml, 20 ml vial		1	✓ Erbitux
Inj 5 mg per ml, 100 ml vial		1	✓ Erbitux
Inj 1 mg for ECP	3.82	1 mg	✓ Baxter

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB - PCT only - Special Authority see SA1951 below

Inj 100 mg	806.00	1	✓ Remicade
Inj 1 mg for ECP	8.29	1 mg	✓ Baxter

⇒SA1951 Special Authority for Subsidy

Initial application — (**Graft vs host disease**) from any medical practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the qut.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

significant response to treatment in the opinion of the physician; and

3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for psoriatic arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (severe ocular inflammation) from any medical practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
- 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
- 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (chronic ocular inflammation) from any medical practitioner. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	l Generic	
\$	Per 🗸	Manufacturer	

continued...

Initial application — (Pulmonary sarcoidosis) from any medical practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has life-threatening pulmonary sarcoidosis that is refractory to other treatments; and
- 2 Treatment is to be prescribed by, or has been recommended by, a physician with expertise in the treatment of pulmonary sarcoidosis.

Initial application — (Crohn's disease (adults)) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection;
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initial application — **(Crohn's disease (children))** only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
 - 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain: and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initial application — (severe ulcerative colitis) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
 - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65: and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids: and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Renewal — (severe ulcerative colitis) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initial application — (plaque psoriasis) only from a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, cyclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) only from a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and

Subsidy		Fully	Brand or	Т
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	/	Manufacturer	

continued...

- 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
- 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (neurosarcoidosis) only from a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Biopsy consistent with diagnosis of neurosarcoidosis; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Fither:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Either:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (severe Behcet's disease) from any medical practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes:

a) Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	•	Manufacturer	

continued...

b) Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any medical practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment: and
- 3 A maximum of 8 doses.

⇒SA1896 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

the first dose to assess response to treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB - PCT only - Specialist - Special Authority see SA1627 below

lnj 25 mg per ml	, 40 ml vial5,910.00	1	✓ Gazyva
Inj 1 mg for ECP	6.21	1 mg	✓ Baxter

⇒SA1627 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

Inj 150 mg prefilled syringe	450.00	1	Xolair
Inj 150 mg vial	450.00	1	✓ Xolair

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	✓ Manufacturer	

continued...

12 months, unless contraindicated or not tolerated; or

- 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PERTUZUMAB - PCT only - Specialist - Special Authority see SA1606 on the next page

Inj 30 mg per ml, 14 ml vial	3,927.00	1	Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	✓ Baxter

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist - Special Authority see SA1901 below

Inj 100 mg per 10 ml vial	1,075.50 2	Mabthera
Inj 500 mg per 50 ml vial	2,688.30 1	✓ Mabthera
Inj 1 mg for ECP	5.64 1 mg	✓ Baxter (Mabthera)

⇒SA1901 Special Authority for Subsidy

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment;
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and

Subsidy	Fu	,	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months,

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

but the condition has relapsed and the patient now requires repeat treatment; and

3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — **(haemophilia with inhibitors)** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and

Subsidy	Fully		Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

continued...

- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Fither:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept: or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

	Subsidy		Fully	Brand or
(M	anufacturer's Price)	Sub	sidised	Generic
	\$	Per	✓	Manufacturer

continued...

- 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physiciann; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	Fı	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
 \$	Per	•	

continued...

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA1937 below

inj 100 mg per 10 mi viai	2/5.33	2	✔ HIXIMYO
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

⇒SA1937 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*: and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	√	

continued...

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
 - 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Fither
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks. or two 1.000 mg doses given two weeks apart: and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*: and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects: and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective;
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

Subsidy	F	ully	Brand or	_
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre: or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or

Subs (Manufactur		
\$	Per	Manufacturer

continued...

- 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
- 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles: or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy;
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

continued...

recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms: and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks: and
- 2 Fither
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1.000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 x 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease: and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

SECUKINUMAB - Special Authority see SA1754 below - Retail pharmacy

⇒SA1754 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

prior treatment course: and

4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

Note: Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Inj 100 mg vial770.57

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB - PCT only - Special Authority see SA1858 below

Inj 20 mg per ml, 4 ml vial	220.00	1	✓ Actemra
Inj 20 mg per ml, 10 ml vial	550.00	1	✓ Actemra
Inj 20 mg per ml, 20 ml vial		1	✓ Actemra
Inj 1 mg for ECP	2.85	1 mg	✓ Baxter

⇒SA1858 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis: or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease: or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
 - 3.2.2 Fither:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Fither:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints;
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD): or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992:19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had severe active polyarticular course JIA for 6 months duration or longer; and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	/	Manufacturer	

continued...

- 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate (at a dose of 10-20 mg/m² weekly or at the maximum tolerated dose) in combination with either oral corticosteroids (prednisone 0.25 mg/kg or at the maximum tolerated dose) or a full trial of serial intra-articular corticosteroid injections; and
- 2.4 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.5 Both:
 - 2.5.1 Fither:
 - 2.5.1.1 Patient has persistent symptoms of poorly-controlled and active disease in at least 20 swollen, tender ioints; or
 - 2.5.1.2 Patient has persistent symptoms of poorly-controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, shoulder, cervical spine, hip; and
 - 2.5.2 Physician's global assessment indicating severe disease.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status.

Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
TRASTUZUMAB - PCT only - Specialist - Special Authority see	SA1632 below			
Inj 150 mg vial	1,350.00	1	✓	Herceptin
Inj 440 mg vial	3,875.00	1	✓	Herceptin
Inj 1 mg for ECP	9.36	1 mg	1	Baxter

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology);
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE - PCT only - Specialist -	- Special Authority see SA1871	below	
Inj 100 mg vial	2,320.00	1	✓ Kadcyla

 Inj 160 mg vial
 3,712.00
 1
 ✓ Kadcyla

 Inj 1 mg for ECP
 23.20
 1 mg
 ✓ Baxter

⇒SA1871 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Fither:
 - 3.1 The patient has received prior therapy for metastatic disease*: or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Fither:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Treatment to be discontinued at disease progression.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

Programmed Cell Death-1 (PD-1) Inhibitors

NIVOLUMAB - PCT only - Specialist - Special Authority see SA1911 below		
Inj 10 mg per ml, 4 ml vial	1	Opdivo
Inj 10 mg per ml, 10 ml vial2,629.96	1	✓ Opdivo
Inj 1 mg for ECP27.62	1 mg	✓ Baxter

⇒SA1911 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Fither:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes: and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10. mm.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

⇒SA1910 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

continued...

per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

CICLOSPORIN		
Cap 25 mg	44.63 50 ✓ Ne	oral
Cap 50 mg	88.91 50 ✓ Ne	oral
Cap 100 mg17	77.81 50 V Ne	oral
Oral liq 100 mg per ml19	98.13 50 ml OP ✓ Ne	oral
EVEROLIMUS - Special Authority see SA1913 below - Retail pharmac	;y	
Wastage claimable		
Tab 10 mg6,5	12.29 30 ✓ Afi	nitor
Tab 5 mg4,55	55.76 30 ✓ Afi	nitor

⇒SA1913 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Roth:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Everolimus to be discontinued at progression of SEGAs; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Note: : MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

	Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	Generic
SIROLIMUS – Special Authority see SA0866 below – Retail phar	,	100	./	Rapamune
Tab 1 mg Tab 2 mg Oral liq 1 mg per ml	1,499.99	100 100 60 ml C	1	Rapamune Rapamune

⇒SA0866 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min; or
- · Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP; or
- · Leukoencepthalopathy; or
- · Significant malignant disease

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg		100	✓ Tacrolimus Sandoz
Cap 1 mg	84.30	100	✓ Tacrolimus Sandoz
Cap 5 mg	248.20	50	✓ Tacrolimus Sandoz

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

✓ Firazyr

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antiallergy Preparations

Allergic Emergencies

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Maintenance kit - 6 vials 120 mon freeze dried venom with

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above - Retail pharmacy

diluent	285.00	1 OP	✓ Venomil \$29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent			
9 ml, 3 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority see \$	SA1367 above	- Retail pharr	nacy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze			
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze			-
dried venom, with diluent	305.00	1 OP	✓ Venomil S29

	Subsidy		Fully Brand or
	(Manufacturer's Price	e) Subs	idised Generic
	\$	Per	✓ Manufacturer
Authistonius			
Antihistamines			
CETIRIZINE HYDROCHLORIDE			
* Tab 10 mg	1 12	100	✓ Zista
* Oral liq 1 mg per ml		200 ml	✓ Histaclear
CHLORPHENIRAMINE MALEATE	2.00		
	0.07	500 ml	✓ Histafen
* Oral liq 2 mg per 5 ml	9.37	300 1111	▼ nistaleli
DEXTROCHLORPHENIRAMINE MALEATE			
* Tab 2 mg		40	
	(8.40)		Polaramine
	1.01	20	5
W 0 11 0 5 1	(5.99)	400 1	Polaramine
* Oral liq 2 mg per 5 ml		100 ml	D
	(10.29)		Polaramine
FEXOFENADINE HYDROCHLORIDE			
* Tab 60 mg	4.34	20	
	(8.23)		Telfast
* Tab 120 mg	4.74	10	
	(8.23)		Telfast
	14.22	30	
	(26.44)		Telfast
LORATADINE			
* Tab 10 mg	1.69	100	✓ Lorafix
* Oral liq 1 mg per ml	2.15	120 ml	✓ Lorfast
PROMETHAZINE HYDROCHLORIDE			
* Tab 10 mg	1.68	50	✓ Allersoothe
* Tab 25 mg		50	✓ Allersoothe
* Oral lig 1 mg per 1 ml		100 ml	✓ Allersoothe
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a		5	✓ Hospira
		•	
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler, 50 mcg per dose		00 dose OP	✓ Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		00 dose OP	✓ Beclazone 50
Aerosol inhaler, 100 mcg per dose		00 dose OP	✓ Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		00 dose OP	✓ Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free	22.67 2	00 dose OP	✓ Beclazone 250
BUDESONIDE			
Powder for inhalation, 100 mcg per dose	17.00 2	00 dose OP	✓ Pulmicort
			Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00 2	00 dose OP	✓ Pulmicort
- ·			Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00 2	00 dose OP	✓ Pulmicort
			Turbuhaler

	Subsidy (Manufacturer's l	Price) Sub Per	Fully sidised		
FLUTICASONE					
Aerosol inhaler, 50 mcg per dose	7.19	120 dose OF		Flixotide	
Powder for inhalation, 50 mcg per dose	7.50	60 dose OP	1	Flixotide Accuhaler	
Powder for inhalation, 100 mcg per dose		60 dose OP		Flixotide Accuhaler	
Aerosol inhaler, 125 mcg per dose		120 dose OF		Flixotide	
Aerosol inhaler, 250 mcg per dose		120 dose OF		Flixotide	
Powder for inhalation, 250 mcg per dose	13.60	60 dose OP	•	Flixotide Accuhaler	
Inhaled Long-acting Beta-adrenoceptor Agonisi	ts				
EFORMOTEROL FUMARATE					
Powder for inhalation, 12 mcg per dose, and monodose device	ce20.64	60 dose			
	(35.80)			Foradil	
EFORMOTEROL FUMARATE DIHYDRATE					
Powder for inhalation 4.5 mcg per dose, breath activated					
(equivalent to eformoterol fumarate 6 mcg metered dose) 10.32	60 dose OP			
, ,	(16.90)			Oxis Turbuhaler	
INDACATEROL					
Powder for inhalation 150 mcg	61.00	30 dose OP	1	Onbrez Breezhaler	
Powder for inhalation 300 mcg	61.00	30 dose OP	1	Onbrez Breezhaler	
SALMETEROL					
Aerosol inhaler CFC-free, 25 mcg per dose	25.00	120 dose OF	/	Serevent	
Aerosol inhaler 25 mcg per dose		120 dose OF	1	Meterol	
Powder for inhalation, 50 mcg per dose, breath activated	25.00	60 dose OP	1	Serevent Accuhaler	
(Meterol Aerosol inhaler 25 mcg per dose to be delisted 1 Januar	y 2021)				
Inhaled Corticosteroids with Long-Acting Beta-	Adrenocept	or Agonist	3		
BUDESONIDE WITH EFORMOTEROL					
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg	18 23	120 dose OF	/	Vannair	
Powder for inhalation 100 mcg with eformoterol fumarate 6 m		120 dose OF		Symbicort	
. chack for annualist rooming that connected running and	.09	0 0000 0.		Turbuhaler 100/6	
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg	21.40	120 dose OF	/	Vannair	
Powder for inhalation 200 mcg with eformoterol fumarate 6 m		120 dose OF	1	Symbicort	
G	· ·			Turbuhaler 200/6	
Powder for inhalation 400 mcg with eformoterol fumarate					
12 mcg - No more than 2 dose per day	44.08	60 dose OP	1	Symbicort	
				Turbuhaler 400/12	
FLUTICASONE FUROATE WITH VILANTEROL					
Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose OP	✓	Breo Ellipta	
FLUTICASONE WITH SALMETEROL					
Aerosol inhaler 50 mcg with salmeterol 25 mcg	25.79	120 dose OF	1	Seretide	
Aerosol inhaler 125 mcg with salmeterol 25 mcg		120 dose OF		Seretide	
Powder for inhalation 100 mcg with salmeterol 50 mcg - No					
more than 2 dose per day	33.74	60 dose OP	✓	Seretide Accuhaler	
Powder for inhalation 250 mcg with salmeterol 50 mcg - No					
more than 2 dose per day	44.08	60 dose OP	✓	Seretide Accuhaler	

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's F	Price) Subsid	dised	Generic
	\$	Per	•	Manufacturer
Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Oral liq 400 mcg per ml	20.00	150 ml	✓ Vei	ntolin
Infusion 1 mg per ml, 5 ml		10	✓ Vei	ntolin
Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO		5	✓ Vei	
, ooog po, op to o, avanasie on a . oo				
Inhaled Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Aerosol inhaler, 100 mcg per dose CFC free - Up to 1000				
dose available on a PSO	3.80	200 dose OP		spigen
			✓ Sal	Air
	(6.00)		Ver	ntolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule - Up to 30 neb				
available on a PSO	3 93	20	✓ Ast	halin
		20	· <u>A</u>	
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb	4.00	00		
available on a PSO	4.03	20	✓ Ast	<u>inalin</u>
TERBUTALINE SULPHATE				
Powder for inhalation, 200 mcg per dose (equivalent to				
250 mcg metered dose), breath activated	22.20	120 dose OP	√ Bri	canyl Turbuhaler
				•
Powder for inhalation, 250 mcg per dose, breath activated		200 dose OP		canyl Turbuhaler
(Bricanyl Turbuhaler Powder for inhalation, 250 mcg per dose, bro	eath activated to	o be delisted 1 O	ctober 2	!020)
Anticholinergic Agents				
IDDATEORIUM PROMINE				
IPRATROPIUM BROMIDE				
Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dose	9			
available on a PSO	16.20	200 dose OP	✓ Atr	ovent
Nebuliser soln, 250 mcg per ml, 1 ml ampoule - Up to 40 ne	b			
available on a PSO		20	✓ Uni	ivent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 ne		20	- 0	····
		00	. I II.	
available on a PSO		20	✓ <u>Un</u>	vent
(Univent Nebuliser soln, 250 mcg per ml, 1 ml ampoule to be delis	sted 1 January 2	2021)		
Inhaled Beta-Adrenoceptor Agonists with Anticl	nolinergic A	gents		
SALBUTAMOL WITH IPRATROPIUM BROMIDE				
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p	er			
dose CFC-free	12.19	200 dose OP	✓ Du	olin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per				
vial, 2.5 ml ampoule – Up to 20 neb available on a PSO	5.20	20	✓ Du	olin
That, 2.5 th ampould op to 20 hob available of a 1 00		20	- 50	<u> </u>
Long-Acting Muscarinic Antagonists				
Long-Acting Muscarinic Antagonists				
GLYCOPYRRONIUM - Subsidy by endorsement				
• •				and a fall and the top of the con-
a) Inhaled glycopyrronium treatment will not be subsidised if	patient is also r	receiving treatme	ent with s	subsidised tiotropium or
umeclidinium.				
b) Glycopyrronium powder for inhalation 50 mcg per dose is		for patients who	have b	een diagnosed as
having COPD, and the prescription is endorsed according	μly.			
Powder for inhalation 50 mcg per dose		30 dose OP	✓ See	ebri Breezhaler
31				

Subsidy	Fully	Brand or
(Manufacturer's Pi	rice) Subsidised	Generic
\$	Per 🗸	Manufacturer

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed

accinica chacicoa.			
Powder for inhalation, 18 mcg per dose	50.37	30 dose	✓ Spiriva
Soln for inhalation 2.5 mcg per dose	50.37	60 dose OP	✓ Spiriva Respimat

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD, and the prescription is endorsed accordingly.

Powder for inhalation 62.5 mcg per dose	61.50	30 dose OP	✓ Incruse Ellipta
-----------------------------------------	-------	------------	-------------------

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA1584 above	ve – Retail phar	macy
Powder for Inhalation 50 mcg with indacaterol 110 mcg81.00	30 dose OP	✓ Ultibro Breezhaler
TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 a	ıbove – Retail p	harmacy
Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg81.00	60 dose OP	✓ Spiolto Respimat
UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Re	etail pharmacy	

OMEOLIBITION WITH VILLANDE	opoolar riamonty coo or th	00 . abovo	riotan priarmacy		
Douglar for inhalation 60 E mag with	vilantaral OE mag	77.00	20 doos OB	Anara I	Illinto

Powder for inhalation 62.5 mcg with vilanterol 25 mcg	✓ Ar	noro Ellipta
-------------------------------------------------------	------	--------------

Antifibrotics

NINTEDANIB - Special Authority see SA1928 below - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone. Cap 100 mg......2.554.00 60 OP ✓ Ofev ✓ Ofev 60 OP

⇒SA1928 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient has been diagnosed with idiopathic pulmonary fibrosis; and

Subsidy	Fully		Brand or	
(Manufacturer's Price	,	Subsidised	Generic	
\$	Per		Manufacturer	

continued...

- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA1929 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Tab 801 mg	3,645.00	90	Esbriet
Cap 267 mg - Wastage claimable	3,645.00	270	Esbriet

⇒SA1929 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Leukotriene Receptor Antagonists

MC	NTELUKAST		
*	Tab 4 mg4.25	28	✓ Montelukast Mylan
*	Tab 5 mg4.25	28	✓ Montelukast Mylan
*	Tab 10 mg	28	✓ Montelukast Mylan
	v		✓ Montelukast Mylan

Fully

Subsidy (Manufacturer's Price)

Subsidised Per

Brand or Generic Manufacturer

Mast Cell Stabilisers

NEDOCROMIL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking nedocromil prior to 1 July 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of nedocromil.

Aerosol inhaler, 2 mg per dose CFC-free.......28.07 112 dose OP ✓ Tilade

(Tilade Aerosol inhaler, 2 mg per dose CFC-free to be delisted 1 February 2021)

SODIUM CROMOGLICATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking sodium cromoglicate prior to 1 July 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of sodium cromoglicate.

Methylxanthines

AMINOPHYLLINE

Mucolytics

DORNASE ALFA – Special Authority see SA0611 below – Retail pharmacy Nebuliser soln, 2.5 mg per 2.5 ml ampoule......250.00

6 ✓ Pulmozyme

⇒SA0611 Special Authority for Subsidy

Special Authority approved by the Cystic Fibrosis Advisory Panel

Notes: Application details may be obtained from PHARMAC's website www.pharmac.govt.nz/SAForms or:

The Co-ordinator, Cystic Fibrosis Advisory Panel Phone: (04) 460 4990 PHARMAC, PO Box 10 254 Facsimile: (04) 916 7571 Wellington Email: CFPanel@pharmac.govt.nz

Prescriptions for patients approved for treatment must be written by respiratory physicians or paediatricians who have experience and expertise in treating cystic fibrosis.

SODIUM CHI ORIDE

Not funded for use as a nasal drop.

Nasal Preparations

Allergy Prophylactics

BUDESONIDE

SteroClear to be Sole Supply on 1 October 2020

	ıbsidy		Fully	Brand or
(Manufac	turer's Price)		sidised	Generic
	\$	Per		Manufacturer
FLUTICASONE PROPIONATE				
Metered aqueous nasal spray, 50 mcg per dose1	.98 120	dose OP	✓ [Flixonase Hayfever
				& Allergy
PRATROPIUM BROMIDE				
Aqueous nasal spray, 0.03%4	.61 15	ml OP	√ (Jnivent
Respiratory Devices				
MASK FOR SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
c) Only for children aged six years and under				
Small	20	1	1	e-chamber Mask
PEAK FLOW METER	0	•	•	onambor maon
a) Up to 25 dev available on a PSO				
b) Only on a PSO Low range9	54	1	1 1	Mini-Wright AFS
Low range	.04	'	• 1	Low Range
Normal range9	54	1	✓ I	Mini-Wright
Troma range	.01	•	٠,	Standard
SPACER DEVICE				oturida d
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
220 ml (single patient)	95	1	1	e-chamber Turbo
510 ml (single patient)		i	•	e-chamber La
- · · · · · · · · · · · · · · · · · · ·		•		Grande
800 ml	.50	1	1	/olumatic
		•		

CAFFEINE CITRATE			
Oral liq 20 mg per ml (10 mg base per ml)	15.10	25 ml OP	✓ Biomed

			SENSORY UNGANS
	Subsidy (Manufacturer's Pri \$	ice) Subs	Fully Brand or sidised Generic Manufacturer
Ear Preparations			
ACETIC ACID WITH 1, 2- PROPANEDIOL DIACETATE AND BE			
For Vosol ear drops with hydrocortisone powder refer Stand	ard Formulae, pag	e 248	
Ear drops 2% with 1, 2-Propanediol diacetate 3% and benzethonium chloride 0.02%	6.07	35 ml OP	✓ Vosol
FLUMETASONE PIVALATE	0.97	33 IIII OF	▼ V0501
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform
			ED's
			✓ Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTATI	N	
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate	E 40	7.5 OD	/ Vanasamb
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	✓ Kenacomb
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and			
gramicidin 50 mcg per ml	4.50 (9.27)	8 ml OP	Sofradex
FRAMYCETIN SULPHATE	(9.27)		Soliauex
Ear/Eye drops 0.5%	4.13	8 ml OP	
, ,	(8.65)		Soframycin
F Duanamakia na			
Eye Preparations			
Eye preparations are only funded for use in the eye, unless expli	citly stated otherw	ise.	
Anti-Infective Preparations			
A CICL OVID			

ACICLOVIR		
* Eye oint 3%	4.5 g OP	✓ ViruPOS
CHLORAMPHENICOL		
Eye oint 1%1.55	5 g OP	✓ Devatis
Eye drops 0.5%1.54	10 ml OP	✓ Chlorafast
Funded for use in the ear*. Indications marked with * are unapproved in	dications.	
CIPROFLOXACIN		
Eye drops 0.3% – Subsidy by endorsement	5 ml OP	✓ Ciprofloxacin Teva
When prescribed for the treatment of bacterial keratitis or severe bacteria		•
for the second line treatment of chronic suppurative otitis media (CSOM))*; and the pres	cription is endorsed accordingly.
Note: Indication marked with a * is an unapproved indication.	•	
GENTAMICIN SULPHATE		
Eve drops 0.3%	5 ml OP	✓ Genoptic
PROPAMIDINE ISETHIONATE		
* Eye drops 0.1%2.97	10 ml OP	
(14.55)	10 1111 01	Brolene
,		Diolette
SODIUM FUSIDATE [FUSIDIC ACID]	- 00	4 =
Eye drops 1%5.29	5 g OP	Fucithalmic

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Pr	rice) Sub	Fully	Brand or Generic
	\$	Per	1	Manufacturer
TOBRAMYCIN				
Eye oint 0.3%	10.45	3.5 g OP	✓ T	obrex
Eye drops 0.3%	11.48	5 ml OP	√ T	obrex
Corticosteroids and Other Anti-Inflammatory P	reparations			
DEXAMETHASONE				
* Eye oint 0.1%	5.86	3.5 g OP	✓ N	/laxidex
* Eye drops 0.1%	4.50	5 ml OP	✓ N	// Axidex
Ocular implant 700 mcg - Special Authority see SA1680 be				

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient has diabetic macular oedema with pseudophakic lens; and

- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not vet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

*	Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b	5.00	0.5 0.0	/ Mandana I	
	sulphate 6,000 u per g	5.39	3.5 g OP	✓ Maxitrol	
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin				
	b sulphate 6,000 u per ml	4.50	5 ml OP	✓ Maxitrol	
DIC	CLOFENAC SODIUM				
	Eye drops 0.1%	13.80	5 ml OP	✓ Voltaren Ophtha	

✓ Ozurdex

	Subsidy		Fully	Brand or
	(Manufacturer's F	Price) Subs	sidised	Generic
	\$	Per	✓	Manufacturer
FLUOROMETHOLONE				
* Eye drops 0.1%	3.09	5 ml OP	√ F	:ML
,	5.20		√ F	lucon
LEVOCABASTINE				
Eye drops 0.5 mg per ml	8.71	4 ml OP		
, , , , , , , , , , , , , , , , , , , ,	(10.34)		L	ivostin
LODOXAMIDE	, ,			
Eye drops 0.1%	8.71	10 ml OP	√ L	.omide
PREDNISOLONE ACETATE				
Eye drops 1%	5.93	10 ml OP	√ F	rednisolone-AFT
	7.00	5 ml OP	-	red Forte
DDEDNICOLONE CODILIM DUOCDUATE Consist A		· · ·		104 1 0110
PREDNISOLONE SODIUM PHOSPHATE - Special A			•-	
Eye drops 0.5%, single dose (preservative free)	38.50	20 dose	✓ I/	linims
				Prednisolone

⇒SA1715 Special Authority for Subsidy

SODIUM CROMOGLICATE

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Eye drops 2%1.79	5 ml OP	✓ Rexacrom					
Glaucoma Preparations - Beta Blockers							
BETAXOLOL # Eye drops 0.25%	5 ml OP 5 ml OP	✓ Betoptic S✓ Betoptic					
* Eye drops 0.25%	5 ml OP	✓ Arrow-Timolol					
September 2020 ** Eye drops 0.5%	5 ml OP	✓ Arrow-Timolol					
* Eye drops 0.5%, gel forming3.78	2.5 ml OP	✓ Timoptol XE					
Glaucoma Preparations - Carbonic Anhydrase Inhibitors							
ACETAZOLAMIDE * Tab 250 mg17.03 BRINZOLAMIDE	100	✓ Diamox					
* Eye drops 1%	5 ml OP	✓ Azopt					
DORZOLAMIDE HYDROCHLORIDE	5 ml OD						
* Eye drops 2%	5 ml OP	Trusopt					
DORZOLAMIDE WITH TIMOLOL							
* Eye drops 2% with timolol 0.5%	5 ml OP	✓ Dortimopt					

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully Brand or idised Generic ✓ Manufacturer
Glaucoma Preparations - Prostaglandin Analogo	ues		
BIMATOPROST * Eye drops 0.03%	3.30	3 ml OP	✓ Bimatoprost Multichem
LATANOPROST * Eye drops 0.005% TRAVOPROST	1.57	2.5 ml OP	✓ <u>Teva</u>
* Eye drops 0.004%	7.30 19.50	5 ml OP 2.5 ml OP	✓ Travopt✓ Travatan
Glaucoma Preparations - Other			
BRIMONIDINE TARTRATE * Eye drops 0.2% BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE	4.29	5 ml OP	✓ Arrow-Brimonidine
* Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ Combigan
PILOCARPINE HYDROCHLORIDE * Eye drops 1%	5.35 7.99	15 ml OP 15 ml OP 15 ml OP	✓ Isopto Carpine✓ Isopto Carpine✓ Isopto Carpine
 ★ Eye drops 2% single dose - Special Authority see SA0895 below - Retail pharmacy	I for 2 years for gy to the preser f trade" and are	vative; or not approved a	s special authority items.
Mydriatics and Cycloplegics			
* Eye drops 1%	17.36	15 ml OP	✓ Atropt
CYCLOPENTOLATE HYDROCHLORIDE * Eye drops 1%	8.76	15 ml OP	✓ Cyclogyl
TROPICAMIDE * Eye drops 0.5% * Eye drops 1%		15 ml OP 15 ml OP	✓ Mydriacyl✓ Mydriacyl
Preparations for Tear Deficiency			
For acetylcysteine eye drops refer Standard Formulae, page 248 HYPROMELLOSE * Eye drops 0.5%	2.00 (3.92)	15 ml OP	Methopt

	Subsidy (Manufacturer's Price)		Fully idised	Brand or Generic
	\$	Per		Manufacturer
HYPROMELLOSE WITH DEXTRAN * Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	✓ P	oly-Tears

Preservative Free Ocular Lubricants

⇒SA1388 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER - Special Authority see SA1388 above - Retail ph	armacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL - Special Autho	rity see SA1388 at	ove – Retai	il pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special Autl	hority see SA1388	above – Re	tail pharmacy
Eye drops 1 mg per ml	22.00	10 ml OP	✓ Hylo-Fresh
Hylo-Fresh has a 6 month expiry after opening. The Ph	armacy Procedure	s Manual re	striction allowing one bottle n

Hylo-Fresh has a 6 month expiry after opening. The Pharmacy Procedures Manual restriction allowing one bottle per month is not relevant and therefore only the prescribed dosage to the nearest OP may be claimed.

Other Ey	e Preparations
----------	----------------

NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%	15 ml OP	✓ Naphcon Forte
OLOPATADINE		•
Eye drops 0.1%2.20	5 ml OP	Olopatadine Teva
10.00		✓ Patanol
Olopatadine Teva to be Sole Supply on 1 October 2020		
(Patanol Eye drops 0.1% to be delisted 1 October 2020)		
PARAFFIN LIQUID WITH WOOL FAT		
* Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE		
Eve oint 138 mcg per a	5 a OP	✓ VitA-POS



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

✓ BSF Imigran

Various

PHARMACY SERVICES

May only be claimed once per patient.

The Pharmacode for BSF Imigran is 2597330 - see also page 131

(BSF Imigran Brand switch fee to be delisted 1 December 2020)

Agents Used in the Treatment of Poisonings

Antidotes

ACFTYL	CVST	FINE

NALOXONE HYDROCHLORIDE

- a) Up to 5 inj available on a PSO
- b) Only on a PSO

Removal and Elimination

CHARCOAL

•				
*	Oral lig 50 g per 250 ml	43.50	250 ml OP	✓ Carbosorb-X

- a) Up to 250 ml available on a PSO
- b) Only on a PSO

DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy

Wastage claimable		
Tab 125 mg dispersible276.00	28	Exjade
Tab 250 mg dispersible552.00	28	✓ Exjade
Tab 500 mg dispersible	28	Exjade

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Either:



	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Subs	dised	Generic
	\$	Per	✓	Manufacturer
continued				
 For the first renewal following 2 years of therapy, the treatment improvement in all three parameters namely serum ferritin, For subsequent renewals, the treatment has been tolerated in all three parameters namely serum ferritin, cardiac MRI 1 	cardiac MRI T2* ar I and has resulted in	nd liver MRI n clinical sta	T2* le	vels; or
DEFERIPRONE - Special Authority see SA1480 below - Retail p	harmacy			
Tab 500 mg	533.17	100	√ F	erriprox
Oral liq 100 mg per 1 ml	266.59 25	60 ml OP	√ F	erriprox
⇒SA1480 Special Authority for Subsidy				
Initial application only from a haematologist. Approvals valid with	nout further renewa	l unless not	ified fo	r applications meeting the
following criteria:				
Either:				
The patient has been diagnosed with chronic iron overload The patient has been diagnosed with chronic iron overload	U		,	or

* inj 500 mg via	ll	84.53	10	◆ DRF
				<u>Desferrioxamine</u> Mesylate for Inj
				
				<u>BP</u>
SODIUM CALCIUM				

DESFERRIOXAMINE MESILATE



Standard Formulae			
ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base	qs qs	PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water	1 g 70 ml to 100 ml
ASPIRIN AND CHLOROFORM APPLICATION Aspirin Soluble tabs 300 mg Chloroform	12 tabs to 100 ml	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml)	LIQUID (10
CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate Glycerol	60 mg 40 ml	Phenobarbitone Sodium Glycerol BP Water	400 mg 4 ml to 40 ml
Preservative Water	qs to 100 ml	PILOCARPINE ORAL LIQUID Pilocarpine 4% eye drops Preservative	qs qs
CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate Glycerol Preservative	300 mg 40 ml qs	Water (Preservative should be used if quantity supplied is than 5 days.)	to 500 ml for more
Water	to 100 ml	SALIVA SUBSTITUTE FORMULA Methylcellulose	5 g
FOLINIC MOUTHWASH Calcium folinate 15 mg tab Preservative Water	1 tab qs to 500 ml	Preservative Water (Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	qs to 500 ml for more
(Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	for more	SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml	qs
MAGNESIUM HYDROXIDE 8% MIXTURE Magnesium hydroxide paste 29% Methyl hydroxybenzoate Water	275 g 1.5 g	Water (Only funded if prescribed for treatment of hyponatrolly VANCOMYCIN ORAL SOLUTION (50 mg per ml)	qs
METHADONE MIXTURE Methadone powder Glycerol Water	qs qs to 100 ml	Vancomycin 500 mg injection Glycerol BP Water (Only funded if prescribed for treatment of Clostridiu following metronidazole failure)	10 vials 40 ml to 100 ml ım difficile
METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liqu	10 g to 100 ml	VOSOL EAR DROPS WITH HYDROCORTISONE POWDER 1% Hydrocortisone powder Vosol Ear Drops	1% to 35 ml
OMEPRAZOLE SUSPENSION Omeprazole capules or powder Sodium bicarbonate powder BP	qs 8.4 g		

8.4 g to 100 ml

Water

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy
(Manufacturer's Price)
Subsi

Fully Subsidised Brand or Generic Manufacturer

Extemporaneously Compounded Preparations and Galenicals

CHLOROFORM

- a) Only in combination
- b) Maximum of 100 ml per prescription
- c) Only in aspirin and chloroform application.
- d) Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined.

(90.09) Douglas

473 ml

Only in extemporaneously compounded codeine linctus.

COLLODION FLEXIBLE

Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined.

Collodion flexible19.30 100 ml ✓ PSM

COMPOUND HYDROXYBENZOATE - Only in combination

Only in extemporaneously compounded oral mixtures.

GLYCERIN WITH SODIUM SACCHARIN - Only in combination

Only in combination with Ora-Plus.

GLYCERIN WITH SUCROSE - Only in combination

Only in combination with Ora-Plus.

TOERIN WITH SUCHOSE - Only III COMBINATION

GLYCEROL

500 ml ✓ healthE Glycerol BP

✓ Ora-Sweet

- a) Only in extemporaneously compounded oral liquid preparations.
- b) healthE Glycerol BP to be Sole Supply on 1 October 2020

MAGNESIUM HYDROXIDE

(PSM Paste 29% to be delisted 1 January 2021)

METHADONE HYDROCHLORIDE

- a) Only on a controlled drug form
- b) No patient co-payment payable
- c) Safety medicine; prescriber may determine dispensing frequency
- d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets).

Powder 7.84 ✓ AFT 1 q METHYL HYDROXYBENZOATE ✓ Midwest 25 a METHYLCELI UI OSF 100 g ✓ MidWest ✓ Ora-Plus 473 ml METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN - Only in combination Suspension.......30.95 473 ml ✓ Ora-Blend SF

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy		Fully Brand or		
	(Manufacturer's Price	ce) Subsi	sidised Generic		
	\$	Per	✓ Manufacturer		
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - On	•				
Suspension	30.95	473 ml	✓ Ora-Blend		
PHENOBARBITONE SODIUM					
Powder – Only in combination	52.50	10 g	✓ MidWest		
•	325.00	100 g	✓ MidWest		
Only in children up to 12 years		ŭ			
PROPYLENE GLYCOL					
Only in extemporaneously compounded methyl hydroxybenz	coate 10% solution.				
Lig		500 ml	✓ Midwest		
SODIUM BICARBONATE					
Powder BP - Only in combination	10.05	500 g	✓ Midwest		
Only in extemporaneously compounded omeprazole and lansoprazole suspension.					
SYRUP (PHARMACEUTICAL GRADE) - Only in combination					
Only in extemporaneously compounded oral liquid preparation	ons.				
Liq		500 ml	✓ <u>Midwest</u>		
WATER					
Tap - Only in combination	0.00	1 ml	✓ Tap water		

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

continued...

✓ fully subsidised 251



Subsidy	Fu	ly Brand or	
(Manufacturer's Price)	Subsidise	ed Generic	
\$	Per	Manufacturer	

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome; or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Authority see SA1523 on the previous page - Hospital pharmacy [HP3]

Emulsion (neutral)	12.30	200 ml OP	✓ Calogen
	30.75	500 ml OP	✓ Calogen
Emulsion (strawberry)	12.30	200 ml OP	✓ Calogen
Oil	30.00	500 ml OP	✓ MCT oil (Nutricia)
Oil, 250 ml	114.92	4 OP	✓ Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	rmacy [HP3]	PROTEIN SUPPLEMENT — Special Authority see SA1524 above — Hospital pha
✓ Protifar	225 g OP	Powder7.90
✓ Resource	227 g OP	8.95
Beneprotein		

Subsidy (Manufacturer's Price) Fully Subsidised

Brand or Generic Manufacturer

Oral Supplements/Complete Diet (Nasogastric/Gastrostomy Tube Feed)

Respiratory Products

⇒SA1094 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has CORD and hypercapnia, defined as a CO2 value exceeding 55 mmHg.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Liquid	- Hospital Pharm 1,000 ml OP	acy [HP3] ✓ Diason RTH ✓ Glucerna Select RTH
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 above - Ho	ospital pharmacy	[HP3]
Liquid (strawberry)1.50	200 ml OP	✓ Diasip
Liquid (vanilla)	200 ml OP	✓ Diasip
1.88	250 ml OP	✓ Glucerna Select
1.78	237 ml OP	
(2.10)		Resource Diabetic
(2.10)		Sustagen Diabetic

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism: or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED - Special Authority see SA1525 above - Hospital pharmacy [HP3]

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

r	
PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1379 ab Liquid6.00	ove – Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini Energy RTH
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see SA1379 about Liquid	ve – Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini RTH ✓ Pediasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML — Special Authority se Liquid	ee SA1379 above – Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML – Special Authority see SA1379 above Liquid (strawberry)	 Hospital pharmacy [HP3] 200 ml OP ✓ Fortini 200 ml OP ✓ Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA1379 above - Liquid (chocolate)	Hospital pharmacy [HP3] 200 ml OP
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority see S Liquid (unflavoured)	200 ml OP 200 m
PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 above – Hospita	ai pnarmacy [HP3]

✓ Peptamen Junior

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML - Special Authorit Liquid	•	Hospital pharm 500 ml OP	,
RENAL ORAL FEED 1.8 KCAL/ML - Special Authority se	e SA1101 above – Hos	pital pharmacy	[HP3]
Liquid		220 ml OP	
RENAL ORAL FEED 2 KCAL/ML - Special Authority see	SA1101 above – Hospi	tal pharmacy [H	IP3]
Liquid	2.88	237 ml OP	
	(3.31)		NovaSource Renal
Liquid (apricot) 125 ml	11.52	4 OP	✓ Renilon 7.5
Liquid (caramel) 125 ml	11.52	4 OP	✓ Renilon 7.5

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's Pri \$	ice) Subsi Per	Fully idised	Brand or Generic Manufacturer
ORAL ELEMENTAL FEED 0.8KCAL/ML — Special Authority see Liquid (grapefruit), 250 ml carton Liquid (pineapple & orange), 250 ml carton Liquid (summer fruits), 250 ml carton	171.00 171.00	revious page - 18 OP 18 OP 18 OP	✓ E	tal pharmacy [HP3] lemental 028 Extra lemental 028 Extra lemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see S Powder (unflavoured)	•	vious page – I 80 g OP		l pharmacy [HP3] ivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Auth [HP3] Liquid	•	on the previou 1,000 ml OP		- Hospital pharmacy

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML	 Special Authority 	see SA1196 ab	ove -	– Hospitai pharmacy [HP3]
Liquid	4.00	500 ml OP	1	Nutrini Low Energy
				Multi Fibre

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	sed	Generic
\$	Per	•	Manufacturer

continued...

3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g., to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (**Long-term medical condition**) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia: or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

	Subsidy Manufacturaria Pr	ioo)	Fully Subsidised	
(Manufacturer's Pr \$	Per	Subsidised •	Manufacturer
ontinued				
 Is being fed via a tube or a tube is to be inserted for the purcondition criteria); or Cystic Fibrosis; or Liver disease; or Chronic Renal failure; or Inflammatory bowel disease; or Chronic obstructive pulmonary disease with hypercapnia; or 		(not naso	ogastric tul	oe - refer to specific medic
7 Short bowel syndrome; or8 Bowel fistula; or9 Severe chronic neurological conditions.				
ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on	nage 258 – Ho	snital nha	armacy [HI	23]
Liquid		1,000 ml	, .	Nutrison Energy
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on p Liquid	1.24	oital pharr 250 ml (1,000 ml	OP ✓	Isosource Standard Nutrison Standard RTH
ENTERAL FEED MITH FIRRE O CO MON MAIL. Conside Authority	044050	05		Osmolite RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML — Special Authority Liquid		1,000 ml		Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority see Liquid		ge 258 – 1,000 ml	OP 🗸	harmacy [HP3] Jevity RTH Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.5KCAL/ML — Special Authority se Liquid	1.75	age 258 - 250 ml (1,000 ml	OP ✓	pharmacy [HP3] Ensure Plus HN Ensure Plus RTH Jevity HiCal RTH Nutrison Energy Multi Fibre
ORAL FEED (POWDER) — Special Authority see SA1859 on page Note: Higher subsidy for Sustagen Hospital Formula will only number and an appropriately endorsed prescription. Powder (chocolate) — Higher subsidy of up to \$26.00 per 850	be reimbursed f			n a valid Special Authority
with Endorsement	•	850 g C 840 g C		Ensure Sustagen Hospital Formula Active
Additional subsidy by endorsement is available for patients prescription must be endorsed accordingly.	s with fat malabs	sorption,	fat intolera	

(26.00) Sustagen Hospital Formula Active
Additional subsidy by endorsement is available for patients with fat malabsorption, fat intolerance or chyle leak. The

26.00

9.54

857 g OP

850 g OP

840 g OP

✓ Fortisip✓ Ensure

Powder (vanilla) - Higher subsidy of up to \$26.00 per 850 g

prescription must be endorsed accordingly.

with Endorsement......8.54

Subsidy		Fully	Brand or	
(Manufacturer's Pric	ce)	Subsidised	Generic	
\$	Per	/	Manufacturer	

ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 258 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
Lituoisement	(1.26) (1.26)	200 IIII OF	Ensure Plus Fortisip
Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (fruit of the forest) - Higher subsidy of \$1.26 per 200 ml			
with Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	
	(1.26)		Fortisip
Liquid (vanilla) – Higher subsidy of up to \$1.33 per 237 ml with			
Endorsement		237 ml OP	
	(1.33)		Ensure Plus
	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 258 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisin Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

0.1.1		F "	D 1	
Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous	page - Hospital p	harmacy [HP3]
Liquid5.50	500 ml OP	✓ Nutrison
		Concentrated
11.00	1,000 ml OP	Two Cal HN RTH

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$1.90 per 200 ml with

Two Cal HN

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

SPECIAL FOODS

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
FOOD THICKENER – Special Authority see SA1106 on the pre	6.53 30	pharmac 00 g OP 80 g OP	✓ N ✓ F	lutilis Feed Thickener Karicare Aptamil

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by PHARMAC from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX – Special Authority see SA1729 abov		pharmacy [HP3] 1,000 g OP	
	(5.15)	.,000 g 0.	Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA1729 abov	e – Hospital	pharmacy [HP3]	
Powder	3.93	1,000 g OP	
	(7.32)	-	NZB Low Gluten Bread Mix
	3.51		
	(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1729 above - F	lospital phari	macy [HP3]	
Powder	5.62	2,000 g OP	
	(18.10)		Horleys Flour

	Subsidy		Fully	Brand or			
	(Manufacturer's Pri		idised	Generic			
	\$	Per	•	Manufacturer			
GLUTEN FREE PASTA – Special Authority see SA1729 on the previous page – Hospital pharmacy [HP3]							
Buckwheat Spirals		250 g OP					
	(3.11)	•	0)rgran			
Corn and Vegetable Shells	2.00 [°]	250 g OP		·			
·	(2.92)	· ·	0)rgran			
Corn and Vegetable Spirals	2.00 [°]	250 g OP		·			
	(2.92)	· ·	0)rgran			
Rice and Corn Lasagne Sheets	1.60 [°]	200 g OP		·			
·	(3.82)	· ·	0)rgran			
Rice and Corn Macaroni		250 g OP		ŭ			
	(2.92)	Ü	0)rgran			
Rice and Corn Penne	, ,	250 g OP		ŭ			
	(2.92)	· ·	0)rgran			
Rice and Maize Pasta Spirals	2.00 [°]	250 g OP		·			
·	(2.92)	· ·	0)rgran			
Rice and Millet Spirals	2.00 [°]	250 g OP		·			
•	(3.11)	· ·	0)rgran			
Rice and corn spaghetti noodles	2.00 [°]	375 g OP		·			
. •	(2.92)	· ·	0)rgran			
Vegetable and Rice Spirals	2.00 [°]	250 g OP		·			
	(2.92)	· ·	0)rgran			
Italian long style spaghetti	2.00	220 g OP		•			
	(3.11)	ŭ	0)rgran			
	. ,			-			

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 above - Hospital pharmacy [HP3]

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

Tabs	99.00	75 OP	✓ Phlexy 10
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior Chocolate
Powder (unflavoured) 27.8 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior Vanilla
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange)	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured)	320.00	500 g OP	✓ XP Maxamum
Liquid (berry)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	Easiphen Liquid
Liquid (juicy tropical) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g	1,123.20	36 OP	✓ PKU Lophlex Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
		1 0001	

(PKU Lophlex Powder Powder (unflavoured) 27.8 g sachets to be delisted 1 March 2021)

Foods

LOW PROTEIN BAKING MIX – Special Authority see SA Powder			oharmacy [HP3] ✓ Loprofin Mix
LOW PROTEIN PASTA - Special Authority see SA1108	on the previous page – I	Hospital pharm	acy [HP3]
Animal shapes	11.91	500 g OP	✓ Loprofin
Lasagne	5.95	250 g OP	✓ Loprofin
Low protein rice pasta	11.91	500 g OP	✓ Loprofin
Macaroni	5.95	250 g OP	✓ Loprofin
Penne	11.91	500 g OP	✓ Loprofin
Spaghetti	11.91	500 g OP	✓ Loprofin
Spirals	11.91	500 g OP	✓ Loprofin

Subsidy (Manufacturer's Price) \$

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

Vanilla

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Gastrointestinal and Other Malabsorptive Problems

AMINO ACID FORMULA - Special Authority see SA1940		,	
Powder	43.60	400 g OP	 Alfamino Junior
Powder (unflavoured)	53.00	400 g OP	✓ Elecare
,		ŭ	✓ Elecare LCP
			✓ Neocate Gold
			✓ Neocate Junior Unflavoured
			✓ Neocate SYNEO
Powder (vanilla)	53.00	400 g OP	✓ Elecare
,		Ü	✓ Neocate Junior

⇒SA1940 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut: or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Fither:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — **(Children 12 months of age and over)** only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist.

continued...

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Approvals valid for 6 months for applications meeting the following criteria:

. -...

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency: or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Fither:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
 - 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
 - 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Fither:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

RAL LIQUID PEPTIDE FORMULA – Special	Authority see SA1953 below –	· Hospital pharma	cy [HP3]
quid 1 kcal/ml	10.45	500 ml OP	✓ Nutrini Peptisorb
quid 1.5 kcal/ml	15.68	500 ml OP	✓ Nutrini Peptisorb
			Energy

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea: or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure; or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and
 - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

continued...

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see SA1557 below - Hospital pharmacy [HP3]

Powder	15.21	450 g OP	✓ Aptamil Gold+ Pepti Junior
	30.42	900 g OP	✓ Allerpro 1 ✓ Allerpro 2

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption: or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea: or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula: and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)35.50	300 g OP	✓ KetoCal 4:1
		✓ Ketocal 3:1
Powder (vanilla)35.50	300 g OP	✓ KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Vaccinations

ADULT DIPHTHERIA AND TETANUS VACCINE - [Xpharm]

- 1) For vaccination of patients aged 45 and 65 years old: or
- 2) For vaccination of previously unimmunised or partially immunised patients; or
- 3) For revaccination following immunosuppression; or
- 4) For boosting of patients with tetanus-prone wounds; or
- 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

(ADT Booster Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid in 0.5 ml to be delisted 1 October 2020)

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent......0.00

10 ✓ BCG Vaccine

BCG Vaccine to be Sole Supply on 1 October 2020

DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE - [Xpharm]

Funded for any of the following criteria:

- 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
- 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
- A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
- 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens: or
- 5) A single dose for vaccination of patients aged 65 years old; or
- 6) A single dose for vaccination of patients aged 45 years old who have not had 4 previous tetanus doses; or
- 7) For vaccination of previously unimmunised or partially immunised patients; or
- 8) For revaccination following immunosuppression; or
- 9) For boosting of patients with tetanus-prone wounds.

Notes: Tdap is not registered for patients aged less than 10 years. Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid, 8 mcg pertussis filamentous

Boostrix to be Sole Supply on 1 October 2020

Subsidised

Per

Fully

Brand or

Generic

Manufacturer

Subsidy

(Manufacturer's Price)

\$

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE – [Xpharm]	
Funded for any of the following:	
 A single dose for children up to the age of 7 who have completed primary immunisation; or A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full 	
primary immunisation; or	
3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemothers	mv.
pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressiv	
regimens; or	C
Five doses will be funded for children requiring solid organ transplantation.	
Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.	
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg	
pertussis toxoid, 25 mcg pertussis filamentous	
haemagglutinin, 8 mcg pertactin and 80 D-antigen units	
poliomyelitis virus in 0.5ml syringe	
Infanrix IPV to be Sole Supply on 1 October 2020	
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE -	
[Xpharm]	
Funded for patients meeting any of the following criteria:	
1) Up to four doses for children up to and under the age of 10 for primary immunisation; or	
2) An additional four doses (as appropriate) are funded for (re-)immunisation for children up to and under the age of	
10 who are patients post haematopoietic stem cell transplantation, or chemotherapy; pre or post splenectomy; pr	e- or
post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or	
Up to five doses for children up to and under the age of 10 receiving solid organ transplantation.	
Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 y	ears)
to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for car	ch up
programmes.	
Inj 30IU diphtheriatoxoid with 40IU tetanustoxoid, 25mcg	
pertussistoxoid, 25mcg	
pertussisfilamentoushaemagglutinin, 8 mcgpertactin,	
80 D-AgUpoliovirus, 10mcghepatitisBsurfaceantigen in	
0.5ml syringe	
Infanrix-hexa to be Sole Supply on 1 October 2020	
HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm]	
One dose for patients meeting any of the following:	
For primary vaccination in children; or	
2) An additional dose (as appropriate) is funded for (re-)immunisation for patients post haematopoietic stem cell	
transplantation, or chemotherapy; functional asplenic; pre or post splenectomy; pre- or post solid organ transplan	t, pre-
or post cochlear implants, renal dialysis and other severely immunosuppressive regimens; or	
 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physic paediatrician. 	an or
paeulatician.	
Haemophilus Influenzae type B polysaccharide 10 mcg	
conjugated to tetanus toxoid as carrier protein 20-40 mcg;	
prefilled syringe plus vial 0.5 ml	
promote opinings produced that the minimum minimum minimum model is a minimum minimum model in the minimum model i	

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
HEPATITIS A VACCINE – [Xpharm] Funded for patients meeting any of the following criteria: 1) Two vaccinations for use in transplant patients; or 2) Two vaccinations for use in children with chronic liver of 3) One dose of vaccine for close contacts of known hepatimes.	*				

Inj 1440 ELISA units in 1 ml syringe Havrix to be Sole Supply on 1 October 2020	0.00	1	✓ Havrix
Inj 720 ELISA units in 0.5 ml syringe Havrix Junior to be Sole Supply on 1 October 2020	0.00	1	Havrix Junior

		Subsidy (Manufacturer's Price)	Per	F Subsidi	ully sed	Brand or Generic Manufacturer
IFPATITIS F	RECOMBINANT VACCINE - [Xpharm]					
	per 0.5 ml vial	0.00	1		✓ HI	BvaxPRO
	led for patients meeting any of the following criteria:		•		•	S Vaxi 110
	for household or sexual contacts of known acute he	natitie B nationte or h	anat	itic B o	arriare	·· or
,	for children born to mothers who are hepatitis B sur					s, OI
	for children up to and under the age of 18 years inc					achieved a nocitive
3)	serology and require additional vaccination or requi					acilieved a positive
4)	for HIV positive patients; or	ic a primary course o	ı vac	onianoi	1, 01	
,	for hepatitis C positive patients; or					
,	for patients following non-consensual sexual interco	nurse: or				
,	for patients following immunosuppression; or	70150, OI				
	for solid organ transplant patients; or					
,	for post-haematopoietic stem cell transplant (HSCT) nationts: or				
,	following needle stick injury.) patients, or				
10)	following fleedle stick injury.					
Ini 10 ma	g per 1 ml vial	0.00	1		✓ µ	BvaxPRO
•	ded for patients meeting any of the following criteria:	0.00	1		• 111	DVAXEITO
		notitio P notionto or h	anat	itio D o	orrioro	N. O.F.
	for household or sexual contacts of known acute he					5; Of
	for children born to mothers who are hepatitis B sur for children up to and under the age of 18 years inc					achieved a positive
٥)	serology and require additional vaccination or requi					achieved a positive
4)	for HIV positive patients; or	ie a primary course o	ı vac	Ullaliu	1, 01	
,	for hepatitis C positive patients; or					
	for patients following non-consensual sexual interco	ourco: or				
,	for patients following immunosuppression; or	Juise, oi				
,	for solid organ transplant patients; or					
,	for post-haematopoietic stem cell transplant (HSCT) nationts: or				
	following needle stick injury.) patients, or				
10)	tonowing needle stick injury.					
Ini 20 mc	g per 1 ml prefilled syringe	0.00	1		√ Fr	ngerix-B
	unded for patients meeting any of the following crite		•			Igerix B
a) i	for household or sexual contacts of known acute		or ho	natitic	B oarr	iore: or
	2) for children born to mothers who are hepatitis B					1615, 01
	3) for children up to and under the age of 18 years					ve achieved a nocitive
	serology and require additional vaccination or re					
	4) for HIV positive patients; or	quite a primary cours	oc UI	vaccini	auon,	OI
	5) for hepatitis C positive patients; or					
	6) for patients following non-consensual sexual into	arcourse. or				
	7) for patients following immunosuppression; or	Sicouisc, oi				
	8) for solid organ transplant patients; or					
	9) for post-haematopoietic stem cell transplant (HS	SCT) natients: or				
	10) following needle stick injury; or	or / patiente, or				
	11) for dialysis patients; or					
	12) for liver or kidney transplant patients.					
	ingerix-B to be Sole Supply on 1 October 2020					
	g per 1 ml vial	0.00	1		✓ HI	BvaxPRO
, 40 1110	3 50		•			

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer Funded for any of the following criteria:

- 1) for dialysis patients; or
- 2) for liver or kidney transplant patient.

(HBvaxPRO Inj 5 mcg per 0.5 ml vial to be delisted 1 October 2020) (HBvaxPRO Inj 10 mcg per 1 ml vial to be delisted 1 October 2020) (HBvaxPRO Inj 40 mcg per 1 ml vial to be delisted 1 October 2020)

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV] - [Xpharm]

Any of the following:

- 1) Maximum of two doses for children aged 14 years and under; or
- 2) Maximum of three doses for patients meeting any of the following criteria:
 - 1) People aged 15 to 26 years inclusive; or
 - 2) Either:

People aged 9 to 26 years inclusive

- 1) Confirmed HIV infection; or
- 2) Transplant (including stem cell) patients: or
- 3) Maximum of four doses for people aged 9 to 26 years inclusive post chemotherapy

✓ Gardasil 9 10 Gardasil 9 to be Sole Supply on 1 October 2020

	Subsidy (Manufacturer's Price) \$	F Subsidi Per	ully sed	Brand or Generic Manufacturer
INFLUENZA VACCINE				

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) ✓ Afluria Quad Junior (2020 Formulation)

A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by PHARMAC:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes: or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV. or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders. or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

Influvac Tetra	1	ccine)9.00	Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)
(2020 formulation)			
✓ Afluria Quad	10	90.00	
(2020 Formulation)			

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	/	Manufacturer	

- a) Only on a prescription
- b) No patient co-payment payable

С

A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by PHARMAC:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - a) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or
- c) children aged four years or less (but over three years) who have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
*				

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment from the Funder for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

✓ MMR II	5	diluent 0.5 ml
✓ Priorix	10	250.00

Priorix to be Sole Supply on 1 October 2020

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Either:

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
 - 2) One dose for close contacts of meningococcal cases; or
 - 3) A maximum of two doses for bone marrow transplant patients; or
 - 4) A maximum of two doses for patients following immunosuppression*; or
- B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) Either:
 - One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
 - ii) One dose for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 December 2019 to 30 November 2020.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 4 mcg of each meningococcal polysaccharide conjugated to

a total of approximately 48 mcg of diphtheria toxoid carrier			
per 0.5 ml vial	0.00	1	✓ Menactra
Menactra to be Sole Supply on 1 October 2020			

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm]					

- 1) The child is under 9 months of age; and
- 2) Any of the following:

PΝ

- 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
- 2) Two doses for close contacts of meningococcal cases; or
- 3) A maximum of two doses for bone marrow transplant patients; or
- 4) A maximum of two doses for patients pre- and post-immunosuppression*.

Note: children under nine months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for booster schedules with meningococcal ACWY vaccine.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 10 mcg in 0.5 ml syringe	0.00	1	✓ Neisvac-C
EUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xphar	m]		
1) A primary course of three doses for previously unvacc	cinated individuals u	p to the age	e of 59 months inclusive
Note: please refer to the Immunisation Handbook for the a	ppropriate schedule	for catch u	p programmes
Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6	6B,		
7F, 9V, 14 and 23F; 3 mcg of pneumococcal			
polysaccharide serotypes 4, 18C and 19F in 0.5 ml			
prefilled syringe	0.00	10	Synflorix
Synflorix to be Sole Supply on 1 October 2020			

		- ::	-	
Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- 1) Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response: or
 - b) with primary immune deficiencies: or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - i) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cvanosis or failure; or
 - I) with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV. for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- 4) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note:	please refer to the Immunisation Handbook for the app	propriate schedule for catch up programmes
Inj 30.	8 mcg of pneumococcal polysaccharide serotypes 1, 3,	, 4,

5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml		
syringe	10	✓ Prevenar 13
	1	✓ Prevenar 13

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE - [Xpharm] Either: 1) Up to three doses (as appropriate) for patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy or with functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or 2) All of the following: a) Patient is a child under 18 years for (re-)immunisation; and b) Treatment is for a maximum of two doses; and c) Any of the following: i) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; or v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); vi) with cochlear implants or intracranial shunts; or vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or x) pre term infants, born before 28 weeks gestation; or xi) with cardiac disease, with cyanosis or failure; or xii) with diabetes: or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with functional asplenia. Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 1 ✓ Pneumovax 23 Pneumovax 23 to be Sole Supply on 1 October 2020 POLIOMYELITIS VACCINE - [Xpharm] Up to three doses for patients meeting either of the following: 1) For partially vaccinated or previously unvaccinated individuals; or 2) For revaccination following immunosuppression. Note: Please refer to the Immunisation Handbook for appropriate schedule for catch-up programmes. ✓ IPOI IPOL to be Sole Supply on 1 October 2020 ROTAVIRUS ORAL VACCINE - [Xpharm] Maximum of two doses for patients meeting the following: 1) first dose to be administered in infants aged under 14 weeks of age; and 2) no vaccination being administered to children aged 24 weeks or over. Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator...............................0.00 10 Rotarix Rotarix to be Sole Supply on 1 October 2020

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully idised	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] – [Xpharm] Either: 1) Maximum of one dose for primary vaccination for either a) Any infant born on or after 1 April 2016; or b) For previously unvaccinated children turning 11 ye varicella infection (chickenpox), or 2) Maximum of two doses for any of the following: a) Any of the following for non-immune patients: i) with chronic liver disease who may in future ii) with deteriorating renal function before trans iii) prior to solid organ transplant; or iv) prior to any elective immunosuppression*, o v) for post exposure prophylaxis who are immu b) For patients at least 2 years after bone marrow tra c) For patients at least 6 months after completion of d) For HIV positive non immune to varicella with mile e) For patients with inborn errors of metabolism at ris varicella, or f) For household contacts of paediatric patients who	be candidates for train plantation; or rune competent inpatie ansplantation, on advictemotherapy, on addict or moderate immunosk of major metabolic pare immunocompron	July 2017, nsplantation ents.; or ice of their lvice of the osuppress decomper nised, or u	on; or special sir specion on ansation	list, or ialist, or advice of HIV specialist, or, with no clinical history of
 immune compromise where the household contact g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure has no clinical history of varicella. 	e no clinical history of	varicella a	and wh	
* immunosuppression due to steroid or other immunosuppres	ssive therapy must be	for a treat	tment p	eriod of greater than
28 days Inj 1350 PFU prefilled syringe	0.00	1 10	_	arivax arivax
Varivax to be Sole Supply on 1 October 2020 Inj 2000 PFU prefilled syringe plus vial	0.00	1 10		arilrix arilrix
(Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 C (Varilrix Inj 2000 PFU prefilled syringe plus vial to be delisted 1 C	,			
VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATE Funded for patients meeting either of the following criteria: 1) One dose for all people aged 65 years; or 2) One dose for all people aged between 66 and 80 years				
Inj 19,400 PFU prefilled syringe plus vial	0.00	1 10		ostavax ostavax
Diagnostic Agents				
TUBERCULIN PPD [MANTOUX] TEST - [Xpharm] Inj 5 TU per 0.1 ml, 1 ml vial Tubersol to be Sole Supply on 1 October 2020	0.00	1	√ T	ubersol

- Symbols -		Aflibercept	197	Amisulpride Mylan	13
UK Synacthen	79	Afluria Quad		Amitriptyline	12
3TC		(2020 Formulation)	277	Amlodipine	
- A -		Afluria Quad Junior		Amneal	
A-Scabies	68	(2020 Formulation)	277	Amorolfine	6
Abacavir sulphate		AFT Carbimazole		Amoxicillin	
Abacavir sulphate with		AFT-Pyrazinamide		Amoxicillin with clavulanic acid	
lamivudine	105	Agents Affecting the		Amphotericin B	3
Abiraterone acetate		Renin-Angiotensin System	47	Amsacrine	
Acarbose	11	Agents for Parkinsonism and Rela		AmsaLyo	
Accarb	11	Disorders	119	Amsidine	
Accuretic	48	Agents Used in the Treatment of		Amzoate	
Accuretic 10	48	Poisonings	246	Anaesthetics	12
Accuretic 20	48	Agrylin		Anafranil	12
Acetazolamide	243	Agrylin S29		Anagrelide hydrochloride	16
Acetec	47	Albendazole		Analgesics	12
Acetic acid with 1, 2- propaned	diol	Albey	233	Anastrozole	18
diacetate and		Albustix	76	Andriol Testocaps	
benzethonium	241	Aldurazyme	30	Androderm	7
Acetic acid with hydroxyquinol	line and	Alecensa	170	Anoro Ellipta	
ricinoleic acid	74	Alectinib	170	Antabuse	
Acetylcysteine		Alendronate sodium	111	Antacids and Antiflatulents	
Aci-Jel	74	Alendronate sodium with		Anthelmintics	8
Aciclovir		colecalciferol	111	Antiacne Preparations	6
Infection	101	Alfacalcidol	34	Antiallergy Preparations	23
Sensory	241	Alfamino Junior	267	Antianaemics	3
Acidex	6	Alginic acid	6	Antiandrogen Oral	
Acipimox	54	Alglucosidase alfa	28	Contraceptives	
Acitretin	68	Alkeran	159	Antiarrhythmics	4
Aclasta	114	Alkeran s29	159	Antibacterials	
Aclin	110	Allerpro 1	270	Antibacterials Topical	
Actemra		Allerpro 2		Anticholinergic Agents	
Actinomycin D	164	Allersoothe	234	Anticholinesterases	
Actrapid		Allmercap		Antidepressants	
Actrapid Penfill		Allopurinol		Antidiarrhoeals	
Acupan		Alpha-Adrenoceptor Blockers		Antiepilepsy Drugs	
Adalat 10		Alpha-Keri Lotion		Antifibrinolytics, Haemostatics and	
Adalat Oros		Alphamox		Local Sclerosants	
Adalimumab		Alphamox 125		Antifibrotics	
Adapalene		Alphamox 250		Antifungals	
Adcortyl		Alprolix		Antifungals Topical	
Adefin		Alu-Tab		Antihistamines	
Adefin XL		Aluminium hydroxide		Antihypotensives	
Adefovir dipivoxil		Alvogen		Antimalarials	
Adenuric		Amantadine hydrochloride		Antimigraine Preparations	
ADR Cartridge 1.8		Ambrisentan		Antinausea and Vertigo Agents	
Adrenaline		Amiloride hydrochloride	53	Antiparasitics	
ADT Booster	272	Amiloride hydrochloride with	50	Antipruritic Preparations	
Adult diphtheria and tetanus	070	furosemide	53	Antipsychotics	
vaccine		Amiloride hydrochloride with		Antiretrovirals	
Advata		hydrochlorothiazide		Antirheumatoid Agents	
Advate		Aminophylline		Antispasmodics and Other Agents	
Adynovate		Amiodarone hydrochloride		Altering Gut Motility	
Afinitor	∠31	Amisulpride	133	Antithrombotic Agents	4

					=
Antithymocyte globulin		Aripiprazole Sandoz	133	Azathioprine	
(equine)	187	Aristocort	64	Azithromycin	8
Antitrichomonal Agents	98	Arrow-Amitriptyline		Azol	8
Antituberculotics and		Arrow-Bendrofluazide	54	Azopt	24
Antileprotics	98	Arrow-Brimonidine	<mark>244</mark>	AZT	10
Antiulcerants	8	Arrow-Calcium	35	- B -	
Antivirals	100	Arrow-Diazepam	136	B-D Micro-Fine	
Anxiolytics	136	Arrow-Doxorubicin	164	B-D Ultra Fine	
Anzatax		Arrow-Fluoxetine		B-D Ultra Fine II	14
Apidra		Arrow-Lamotrigine	129	Bacillus Calmette-Guerin (BCG)	
Apidra SoloStar	11	Arrow-Losartan &		vaccine	18
Apo-Amlodipine		Hydrochlorothiazide		Bacillus Calmette-Guerin	
Apo-Azithromycin		Arrow-Morphine LA	124	vaccine	
Apo-Bromocriptine		Arrow-Norfloxacin	109	Baclofen	
Apo-Ciclopirox	62	Arrow-Ornidazole		Bactroban	6
Apo-Cilazapril/		Arrow-Quinapril 10	47	Barrier Creams and Emollients	
Hydrochlorothiazide	48	Arrow-Quinapril 20	47	BCG Vaccine	27
Apo-Clarithromycin		Arrow-Quinapril 5	47	Beclazone 100	23
Alimentary	9	Arrow-Roxithromycin	90	Beclazone 250	23
Infection	89	Arrow-Timolol	243	Beclazone 50	23
Apo-Clomipramine	125	Arrow-Topiramate	130	Beclomethasone dipropionate	23
Apo-Diclo SR	110	Arrow-Tramadol	125	Bee venom allergy treatment	23
Apo-Diltiazem CD		Arsenic trioxide		Bendamustine hydrochloride	
Apo-Doxazosin		Asacol	7	Bendrofluazide	5
Apo-Folic Acid	38	Asamax	7	Bendroflumethiazide	
Apo-Furosemide		Ascorbic acid	34	[Bendrofluazide]	54
Apo-Gabapentin	128	Aspen Adrenaline	<mark>56</mark>	Benzathine benzylpenicillin	9
Apo-Leflunomide		Aspirin		Benzatropine mesylate	
Apo-Megestrol	178	Blood	41	Benzbromaron AL 100	
Apo-Metoprolol		Nervous	121	Benzbromarone	110
Apo-Mirtazapine		Asthalin	236	Benztrop	120
Apo-Nadolol		Atazanavir sulphate	106	Benzydamine hydrochloride	3
Apo-Nicotinic Acid		Atenolol	50	Benzylpenicillin sodium [Penicillin	
Apo-Oxybutynin		Atenolol AFT		Gj	9
Apo-Perindopril		Atenolol AFT S29	50	Beta Cream	6
Apo-Pindolol		ATGAM	187	Beta Ointment	6
Apo-Pravastatin		Ativan	137	Beta Scalp	
Apo-Prazosin		Atomoxetine	151	Beta-Adrenoceptor Agonists	
Apo-Prednisone		Atorvastatin	<u>55</u>	Beta-Adrenoceptor Blockers	
Apo-Primidone		Atropine sulphate		Betadine	
Apo-Propranolol		Cardiovascular	49	Betadine Skin Prep	
Apo-Pyridoxine		Sensory		Betaferon	
Apo-Selegiline S29		Atropt		Betahistine dihydrochloride	
Apo-Sumatriptan		Atrovent		Betaine	
Apo-Temozolomide		AU Synacthen		Betaloc CR	
Apo-Terazosin		Aubagio		Betamethasone dipropionate	
Apo-Timol		Augmentin		Betamethasone dipropionate with	
Apomorphine hydrochloride	119	Aurorix		calcipotriol	69
Aprepitant		AutoSoft 30		Betamethasone sodium phosphate	
Apresoline		AutoSoft 90		with betamethasone acetate	
Aptamil Gold+ Pepti Junior		Avelox		Betamethasone valerate	
Aqueous cream		Avonex		Betamethasone valerate with	
Aratac		Avonex Pen		clioquinol	6
Arava		Azacitidine		Betamethasone valerate with sodii	
Aripiprazole		Azacitidine Dr Reddy's		fusidate [fusidic acid]	
Aripiprazole 1A Pharma		Azamun		Betaxolol	
r r					

INDEX: Generic Chemicals and Brands

Betnovate	63	Budesonide	CareSens PRO	1
Betnovate-C	64	Alimentary6	Carmellose sodium with gelatin and	
Betoptic	243	Respiratory234, 239	pectin	3
Betoptic S	243	Budesonide with eformoterol235	Carmustine	
Bezafibrate		Bumetanide53	Carvedilol	
Bezalip		Buprenorphine Naloxone BNM 155	Carvedilol Sandoz	
Bezalip Retard	54	Buprenorphine with naloxone155	Catapres	
Bicalutamide		Bupropion hydrochloride155	CeeNU	.15
Bicillin LA		Burinex53	Cefaclor monohydrate	8
BiCNU	159	Burinex S2953	Cefalexin	8
Bicnu Heritage	159	Buscopan8	Cefalexin Sandoz	8
Bile and Liver Therapy		Buspirone hydrochloride136	Cefazolin	
Biltricide		Busulfan159	Ceftriaxone	8
Bimatoprost		- C -	Ceftriaxone-AFT	8
Bimatoprost Multichem		Cabergoline87	Cefuroxime axetil	
Binarex		Caffeine citrate240	Celebrex	
Binocrit		Calamine63	Celecoxib	
Biodone		Calcipotriol69	Celecoxib Pfizer	
Biodone Extra Forte		Calcitonin77	Celestone Chronodose	
Biodone Forte		Calcitriol34	Celiprolol	
Bisacodyl		Calcitriol-AFT34	Cellcept	
Bisoprolol fumarate		Calcium carbonate	Celol	
BK Lotion		Calcium Channel Blockers51	Centrally-Acting Agents	
Bleomycin sulphate		Calcium Disodium Versenate247	Cephalexin ABM	
Blood Colony-stimulating		Calcium folinate161	Cetirizine hydrochloride	
Factors	44	Calcium Folinate Ebewe161	Cetomacrogol	
Blood glucose diagnostic test		Calcium Folinate Sandoz161	Cetomacrogol with glycerol	6
meter	13	Calcium gluconate35	Cetuximab	
Blood glucose diagnostic test		Calcium Homeostasis	Charcoal	
strip	13	Calcium polystyrene sulphonate45	Chemotherapeutic Agents	
Blood glucose test strips (visuall		Calcium Resonium45	Chickenpox vaccine	
impaired)		Calcium Sandoz35	Chlorafast	
Blood Ketone Diagnostic Test		Calogen	Chlorambucil	
Strip	12	Candesartan cilexetil48	Chloramphenicol	
Bonjela		Candestar	Chlorhexidine gluconate	
Boostrix		Canesten 62	Alimentary	3
Bortezomib		Capecitabine161	Dermatological	
Bortezomib Dr-Reddy's		Capercit161	Chloroform	
Bosentan		Capoten47	Chlorothiazide	
Bosentan Dr Reddy's		Capsaicin	Chlorpheniramine maleate	
Bosvate		Musculoskeletal111	Chlorpromazine hydrochloride	
Bplex		Nervous122	Chlortalidone [Chlorthalidone]	
Breo Ellipta		Captopril47	Chlorthalidone	
Brevinor 1/28		Carafate	Chlorvescent	
Brevinor 28		Carbaccord	Choice Load 375	
Bricanyl Turbuhaler		Carbamazepine128	Choice TT380 Short	
Brilinta		Carbimazole81	Choice TT380 Standard	
Brimonidine tartrate		Carbomer245	Choline salicylate with cetalkonium	
Brimonidine tartrate with timolol	277	Carboplatin	chloride	3
maleate	244	Carboplatin Ebewe	Ciclopirox olamine	
Brinzolamide		Carbosorb-X	Ciclosporin	
Brolene		Cardinol LA51	Cilazapril	
Bromocriptine mesylate		CareSens Dual	Cilazapril with	4
Brufen SR		CareSens N	hydrochlorothiazide	4
BSF Imigran		CareSens N POP	Cilicaine	ტ
Buccastem		CareSens N Premier	Cilicaine VK	
DUUUQ315111	104	Out 6000110 14 1 10111101	UIIIU AII IE VIX	📆

	21.111	5.11.1
Cinacalcet77	Colestid54	Daktarin6
Cipflox93	Colestipol hydrochloride54	Dalacin C9
Ciprofloxacin	Colgout	Danazol8
Infection93	Colifoam7	Dantrium11
Sensory241	Colistin sulphomethate93	Dantrium S2911
Ciprofloxacin Teva241	Colistin-Link93	Dantrolene11
Circadin149	Collodion flexible249	Daonil1
Cisplatin	Colloidal bismuth subcitrate10	Dapa-Tabs5
Cisplatin Ebewe	Colofac8	Dapsone9
Citalopram hydrobromide 126	Coloxyl27	Daraprim9
Cladribine161	Combigan244	Darunavir10
Clarithromycin	Compound electrolytes45	Dasatinib17
Alimentary9	Compound electrolytes with glucose	Daunorubicin
Infection89	[Dextrose] 45	DBL Acetylcysteine24
Clexane42	Compound hydroxybenzoate249	DBL Adrenaline5
Clexane Forte42	Concerta153	DBL Aminophylline23
Climara80	Condoms71	DBL Bleomycin Sulfate16
Climara80	Condyline70	DBL Carboplatin15
Clindamycin93	Contraceptives - Hormonal72	DBL Cisplatin15
Clinicians Renal Vit34	Contraceptives - Non-hormonal71	DBL Dacarbazine16
Clobazam128	Copaxone144	DBL Desferrioxamine Mesylate for Inj
Clobetasol propionate63, 70	Corticosteroids and Related Agents	BP24
Clobetasone butyrate64	for Systemic Use78	DBL Docetaxel16
Clofazimine98	Corticosteroids Topical63	DBL Ergometrine7
Clomazol	Cosentyx222	DBL Gemcitabine16
Dermatological62	Cosmegen 164	DBL Gentamicin9
Genito-Urinary74	Coumadin44	DBL Leucovorin Calcium16
Clomifene citrate87	Creon 1000025	DBL Methotrexate Onco-Vial16
Clomipramine hydrochloride125	Creon 2500025	DBL Morphine Sulphate12
Clonazepam127–128, 136	Creon Micro25	DBL Naloxone Hydrochloride24
Clonidine52	Crotamiton63	DBL Octreotide17
Clonidine BNM52	Crystaderm61	DBL Pethidine Hydrochloride12
Clonidine hydrochloride52	Curam91	DBL Vinblastine17
Clopidogrel41	Cvite34	DBL Vincristine Sulfate17
Clopidogrel Multichem41	Cyclizine hydrochloride131	De-Worm8
Clopine	Cyclizine lactate131	Decozol3
Clopixol135–136	Cyclogyl244	Deferasirox24
Clotrimazole	Cyclopentolate hydrochloride244	Deferiprone24
Dermatological62	Cyclophosphamide159	Denosumab11
Genito-Urinary74	Cyclorin98	Deolate9
Clozapine133	Cycloserine98	Deoxycoformycin16
Clozaril	Cyproterone acetate79	Depo-Medrol7
Co-trimoxazole	Cyproterone acetate with	Depo-Provera7
Coal tar	ethinyloestradiol74	Depo-Testosterone7
Coal tar with allantoin, menthol,	Cystadane29	Deprim9
phenol and sulphur69	Cytarabine161	Dermol
Coal tar with salicylic acid and	Cytotec8	Desferrioxamine mesilate24
sulphur	Cytoxan159	Desmopressin acetate8
Coco-Scalp 69	- D -	Desmopressin-PH&T8
Codeine phosphate	D-Penamine	Desuric
Extemporaneous	Dabigatran	Detection of Substances in
Nervous 123	Dacarbazine ARR 164	Urine
Coleman II and Colema	Dacarbazine APP	Dexamethasone
Colaspase [L-asparaginase]164	Dactinomycin [Actinomycin D]164	Hormone
Colchicine	Daivobet	Sensory24
Colecalciferol34	Daivonex69	Dexamethasone phosphate7

INDEX: Generic Chemicals and Brands

Dexamethasone Phosphate	Disulfiram	156	Egopsoryl TA	69
Panpharma78	Diuretics	53	Elaprase	29
Dexamethasone with framycetin and	Docetaxel	164	Elecare	
gramicidin 241	Docetaxel Accord	164	Elecare LCP	267
Dexamethasone with neomycin	Docetaxel Sandoz		Electral	45
sulphate and polymyxin B	Docusate sodium	27	Elelyso	31
sulphate242	Docusate sodium with		Elemental 028 Extra	258
Dexamfetamine sulfate151	sennosides	27	Elocon	64
Dexmethsone78	Dolutegravir	106	Elocon Alcohol Free	64
Dextrochlorpheniramine	Domperidone	131	Eltrombopag	38
maleate234	Donepezil hydrochloride	154	Eltroxin	
Dextrose44–45	Donepezil-Rex	154	EMB Fatol	99
DHC Continus123	Dornase alfa	239	Emend Tri-Pack	131
Diabetes10	Dortimopt	243	EMLA	121
Diabetes Management12	Dorzolamide hydrochloride	243	Emtricitabine	106
Diacomit129	Dorzolamide with timolol	243	Emtricitabine with tenofovir	
Diagnostic Agents283	Dostinex	87	disoproxil	103
Diamide Relief6	Dosulepin [Dothiepin]		Emtriva	106
Diamox243	hydrochloride	125	Emulsifying ointment	66
Diasip254	Dosulepin Mylan	125	Enalapril maleate	47
Diason RTH254	Dothiepin		Enbrel	180
Diazepam127, 136	Doxazosin	47	Endocrine Therapy	177
Diazoxide10	Doxine	92	Endoxan	159
Dibenzyline47	Doxorubicin Ebewe	164	Engerix-B	275
Diclofenac Sandoz110	Doxorubicin hydrochloride	164	Enlafax XR	127
Diclofenac sodium	Doxycycline	92	Enoxaparin sodium	42
Musculoskeletal110	DP Lotion	66	Enstilar	69
Sensory242	DP Lotn HC	64	Ensure	261
Differin61	DP-Allopurinol	116	Ensure Plus	262
Difflam32	Dr Reddy's Omeprazole	9	Ensure Plus HN	261
Diflucan95	Drugs Affecting Bone		Ensure Plus RTH	261
Diflucan S2995	Metabolism	111	Entacapone	119
Diflucortolone valerate64	Dual blood glucose and blood ke	etone	Entapone	119
Digestives Including Enzymes25	diagnostic test meter	12	Entecavir	100
Digoxin49	Duocal Super Soluble Powder		Entecavir Sandoz	100
Dihydrocodeine tartrate123	Duolin	236	Entocort CIR	6
Dilantin129	Duolin HFA	236	Entresto 24/26	48
Dilantin Infatab129	Duride	56	Entresto 49/51	48
Diltiazem hydrochloride52	- E -		Entresto 97/103	48
Dilzem52	e-chamber La Grande	240	Epilim	
Dimethicone65, 67	e-chamber Mask		Epilim Crushable	129
Dimethyl fumarate137	e-chamber Turbo	240	Epilim IV	129
Dipentum7	E-Mycin	90	Epilim S/F Liquid	129
Diphtheria, tetanus and pertussis	Ear Preparations	241	Epilim Syrup	129
vaccine272	Ear/Eye Preparations		Epirubicin Ebewe	164
Diphtheria, tetanus, pertussis and	Easiphen Liquid	266	Epirubicin hydrochloride	164
polio vaccine273	Econazole nitrate	62	Eplerenone	5 <mark>3</mark>
Diphtheria, tetanus, pertussis, polio,	Efavirenz	105	Epoetin alfa	38
hepatitis B and haemophilus	Efavirenz with emtricitabine and		Epoprostenol	
influenzae type B vaccine 273	tenofovir disoproxil	105	Eptacog alfa [Recombinant facto	
Diprosone63	Effient		VIIa]	
Diprosone OV63	Eformoterol fumarate		ERA	90
Dipyridamole41	Eformoterol fumarate dihydrate	235	Erbitux	
Disinfecting and Cleansing	Eftrenonacog alfa [Recombinant		Ergometrine maleate	74
Agents 65	factor IX]		Erlotinib	171
Disopyramide phosphate49	Efudix	70	Erythrocin IV	90

Enthromusin (as lastabianata)	00	Fontonyl	100	Elutomin	170
Erythromycin (as lactobionate)		Fentanyl		FlutaminFluticasone	
Erythromycin ethyl succinate Erythromycin stearate		Fentanyl GHFentanyl Sandoz	100	Fluticasone furoate with	230
Esbriet		Ferinject	120	vilanterol	220
				Fluticasone propionate	
Escitalopram		Ferodan			
Escitalopram-Apotex		Ferric carboxymaltose		Fluticasone with salmeterol	
Eskazole		Ferriprox		FML	
Estradot		Ferro-F-Tabs		Foban	
Estradot 50 mcg		Ferro-tab		Folic acid	
Estrofem		Ferrograd		Food Thickeners	
Etanercept		Ferrosig		Foods And Supplements For In	
Ethambutol hydrochloride		Ferrous fumarate		Errors Of Metabolism	
Ethics Aspirin		Ferrous fumarate with folic acid.		Foradil	
Ethics Aspirin EC		Ferrous sulfate		Forteo	
Ethics Lisinopril		Ferrous sulphate		Fortini	
Ethics Paracetamol Classic		Fexofenadine hydrochloride		Fortini Multi Fibre	
Ethinyloestradiol	81	Fibro-vein		Fortisip	
Ethinyloestradiol with		Filgrastim	44	Fortisip Multi Fibre	262
desogestrel	72	Finasteride		Fosamax	<mark>11</mark> 1
Ethinyloestradiol with		Fingolimod	138	Fosamax Plus	
levonorgestrel	73	Firazyr	233	Framycetin sulphate	241
Ethinyloestradiol with		Flagyl	98	Frisium	
norethisterone	73	Flagyl-S	98	Frumil	53
Ethosuximide	128	Flamazine		Frusemide	53
Etopophos	165	Flecainide acetate	49	Frusemide-Claris	53
Etoposide		Flecainide BNM	49	Fucicort	65
Etoposide phosphate		Flecainide Controlled Release		Fucidin	95
Etravirine		Teva	49	Fucithalmic	
Eumovate		Fleet Phosphate Enema		Fulvestrant	
Everet		Flixonase Hayfever & Allergy		Fungilin	
Everolimus		Flixotide		Furosemide [Frusemide]	
Evista		Flixotide Accuhaler		fusidic acid	
Exemestane		Florinef		Dermatological	62 65
Exjade		Fluanxol		Infection	
Extemporaneously Compounded	= 10	Flucil		Sensory	
Preparations and		Flucloxacillin		- G -	
Galenicals	249	Flucloxin		Gabapentin	128
Eye Preparations		Flucon		Gacet	
Eylea		Fluconazole		Galsulfase	
Ezetimibe		Fludara Oral		Galvumet	
Ezetimibe Sandoz		Fludarabine Ebewe		Galvus	
Ezetimibe with simvastatin		Fludarabine phosphate		Gardasil 9	
- F -	33			Gastrodenol	
		Fluido and Floatrolyton		Gaviscon Double Strength	
Factor eight inhibitor bypassing fraction	20	Fluids and Electrolytes		•	
		Flumetasone pivalate	241	Gaviscon Infant	
Famotidine		Fluocortolone caproate with		Gazyva	
Famotidine Hovid		fluocortolone pivalate and		Gefitinib	
Faslodex		cinchocaine		Gemcitabine Ebewe	
Febuxostat	11/	Fluorometholone		Gemcitabine hydrochloride	
Feed Thickener Karicare	004	Fluorouracil		Gemfibrozil	
Aptamil		Fluorouracil Ebewe		Generic Partners	
FEIBA NF		Fluorouracil sodium		Genoptic	241
Felo 10 ER		Fluox		Gentamicin sulphate	
Felo 5 ER		Fluoxetine hydrochloride		Infection	93
Felodipine	51	Flupenthixol decanoate		Sensory	
Femme-Tab ED	73	Flutamide	178	Gilenya	138

Ginet74	Heparon Junior255	Hyoscine hydrobromide13
Glatiramer acetate144	Hepatitis A vaccine274	Hypam 15
Glecaprevir with pibrentasvir 102	Hepatitis B recombinant	Hyperuricaemia and Antigout11
Glibenclamide11	vaccine 275	Hypromellose24
Gliclazide12	Hepsera100	Hypromellose with dextran24
Glipizide12	Herceptin227	-1-
Glivec172	Hiberix273	Ibiamox9
Glizide12	Hiprex109	Ibilex8
Glucagen Hypokit10	Histaclear234	Ibrance17
Glucagon hydrochloride10	Histafen234	Ibuprofen11
Glucerna Select254	Holoxan159	Ibuprofen SR BNM11
Glucerna Select RTH254	Horleys Bread Mix264	Icatibant23
Glucobay11	Horleys Flour264	Idarubicin hydrochloride16
Glucose [Dextrose]44	Hormone Replacement Therapy -	Idursulfase2
Gluten Free Foods264	Systemic	Ifosfamide15
Glycerin with sodium saccharin249	HPV276	Ikorel5
Glycerin with sucrose249	Humalog11	Iloprost5
Glycerol	Humalog Mix 2511	Imatinib mesilate17
Alimentary27	Humalog Mix 5011	Imatinib-AFT17
Extemporaneous249	Human papillomavirus (6, 11, 16, 18,	Imigran13
Glyceryl trinitrate	31, 33, 45, 52 and 58) vaccine	Imipramine hydrochloride12
Alimentary8	[HPV]276	Imiquimod7
Cardiovascular56	Humatin94	Immune Modulators10
Glycopyrronium236	Humira187	Immunosuppressants18
Glycopyrronium bromide8	HumiraPen187	Imuran18
Glycopyrronium with	Humulin 30/7011	Incruse Ellipta23
indacaterol237	Humulin NPH11	Indacaterol23
Gold Knight71	Humulin R10	Indapamide5
Goserelin86	Hyaluronic acid245	Infanrix IPV27
Gutron50	Hydralazine56	Infanrix-hexa27
Gynaecological Anti-infectives74	Hydralazine hydrochloride56	Infant Formulae26
- H -	Hydrea165	Infatrini27
Habitrol156	Hydrocortisone	Infliximab19
Haemophilus influenzae type B	Dermatological64	Influenza vaccine27
vaccine273	Hormone78	Influvac Tetra
Haldol135	Hydrocortisone (PSM)64	(2020 formulation)27
Haldol Concentrate135	Hydrocortisone acetate7	Inhaled Corticosteroids23
Haldol Decanoas135	Hydrocortisone acetate with	Inhaled Long-acting
Haloperidol133	pramoxine hydrochloride7	Beta-adrenoceptor Agonists 23
Haloperidol decanoate135	Hydrocortisone and paraffin liquid	Inspra5
Harvoni102	and lanolin64	Instillagel Lido12
Havrix274	Hydrocortisone butyrate64, 70	Insulin aspart1
Havrix Junior274	Hydrocortisone with cinchocaine8	Insulin aspart with insulin aspart
HBvaxPRO275	Hydrocortisone with miconazole65	protamine1
healthE Calamine Aqueous Cream	Hydrocortisone with natamycin and	Insulin glargine1
BP63	neomycin 65	Insulin glulisine1
healthE Dimethicone 10%65	Hydrogen peroxide61	Insulin isophane1
healthE Dimethicone 4% Lotion67	Hydroxocobalamin33	Insulin isophane with insulin
healthE Dimethicone 5%65	hydroxycarbamide165	neutral1
healthE Glycerol BP249	Hydroxychloroquine111	Insulin lispro1
healthE Urea Cream66	Hydroxyurea	Insulin lispro with insulin lispro
Healtheries Simple Baking Mix264	[hydroxycarbamide]165	protamine1
Hemastix76	Hygroton54	Insulin neutral1
Heparin Ratiopharm43	Hylo-Fresh245	Insulin pen needles1
Heparin sodium43	Hymenoptera233	Insulin pump1
Heparinised saline43	Hyoscine butylbromide8	Insulin pump cartridge1

Insulin pump infusion set (steel	Jadelle	74	Largactil	133
cannula)20	Jakavi	175	Laronidase	30
Insulin pump infusion set (steel	Jaydess	81	Lasix	53
cannula, straight insertion) 21	Jevity HiCal RTH	261	Latanoprost	244
Insulin pump infusion set (teflon	Jevity RTH		Lax-Suppositories	
cannula)22	Juno		Lax-Tab	
Insulin pump infusion set (teflon	Juno Pemetrexed		Laxatives	
cannula, angle insertion with	- K -		Laxsol	27
insertion device)23	Kadcyla	228	Ledipasvir with sofosbuvir	
Insulin pump infusion set (teflon	Kaletra		Leflunomide	
cannula, angle insertion)	Keflor		Lenalidomide	
Insulin pump infusion set (teflon	Kemadrin		Letrole	
cannula, straight insertion with	Kenacomb		Letrozole	
insertion device)24	Kenacort-A 10		Leukeran FC	
Insulin pump infusion set (teflon	Kenacort-A 40		Leukotriene Receptor	
cannula, straight insertion) 25	Kenalog		Antagonists	238
Insulin pump reservoir25	Kenalog in Orabase		Leunase	
Insulin syringes, disposable with	Kenkay Sorbolene		Leuprorelin	
attached needle14	Ketocal 3:1		Leustatin	
Intal Forte CFC Free239	KetoCal 4:1		Levetiracetam	
Intelence	Ketoconazole	271	Levetiracetam-AFT	
Interferon alfa-2a	Dermatological	70	Levlen ED	
Interferon beta-1-alpha146	Infection		Levocabastine	
•	Ketogenic Diet		Levodopa with benserazide	
Interferon beta-1-beta147				
Intra-uterine device	Ketoprofen		Levodopa with carbidopa Levomepromazine	100
Invega Sustenna	KetoSens		Levomepromazine	100
IPOL	Keytruda		•	10
Ipratropium bromide236, 240	Kindergen		hydrochloride	134
Iressa	Kivexa		Levonorgestrel	7.
Irinotecan Accord	Klacid		Genito-Urinary Hormone	/4
Irinotecan Actavis 100	Kliogest			
Irinotecan hydrochloride161	Kliovance		Levothyroxine	
Irinotecan-Rex161	Kogenate FS		Lidocaine [Lignocaine]	121
Iron polymaltose	Konakion MM		Lidocaine [Lignocaine]	40
Isentress	Konsyl-D		hydrochloride	121
Isentress HD106	Kuvan	30	Lidocaine [Lignocaine] with	40
Ismo 20	-L-	404	chlorhexidine	121
Ismo 40 Retard56	L-asparaginase		Lidocaine [Lignocaine] with	40
Isoniazid99	Labetalol		prilocaine	
Isoniazid with rifampicin99	Lacosamide		Lidocaine-Claris	
Isoprenaline [Isoproterenol]56	Lactulose		Lignocaine	
Isoproterenol56	Laevolac		Lioresal Intrathecal	
Isoptin	Lamictal		Lipazil	
Isoptin Retard52	Lamivudine		Lipid-Modifying Agents	
Isoptin SR52	Lamivudine Alphapharm		Liquigen	
Isopto Carpine244	Lamotrigine		Lisinopril	47
Isosorbide mononitrate56	Lamprene		Lithicarb FC	
Isosource Standard261	Lanoxin		Lithium carbonate	
Isotretinoin61	Lanoxin PG		Livostin	
Ispaghula (psyllium) husk27	Lanoxin S29		LMX4	
Isuprel56	Lansoprazole		Locacorten-Viaform ED's	241
Itch-Soothe63	Lantus		Local preparations for Anal and	
Itraconazole96	Lantus SoloStar		Rectal Disorders	
Itrazole96	Lanvis		Locasol	
Ivermectin67	Lanzol Relief		Locoid	
- J -	Lapatinib ditosylate	172	Locoid Crelo	64

Locoid Lipocream	64	MCT oil (Nutricia)2	253	Methylprednisolone aceponate	6
Locorten-Vioform	241	Measles, mumps and rubella		Methylprednisolone acetate	
Lodoxamide	243	vaccine2		Methylxanthines	
Logem	129	Mebendazole	88	Metoclopramide Actavis 10	13
Lomide		Mebeverine hydrochloride		Metoclopramide hydrochloride	
Lomustine	159	Medco1		Metolazone	
Loniten	57	Medrol	78	Metopirone	8
Loperamide hydrochloride	6	Medroxyprogesterone acetate		Metoprolol succinate	
Lopinavir with ritonavir		Genito-Urinary		Metoprolol tartrate	
Loprofin		Hormone80-		Metrogyl	
Loprofin Mix		Mefenamic acid1		Metronidazole	
Lorafix		Megestrol acetate1		Metroprolol IV Mylan	
Loratadine	234	Melatonin1		Metyrapone	
Lorazepam	137	Melphalan1		Mexiletine hydrochloride	
Lorfast		Menactra2		Mexiletine Hydrochloride USP	
Lorstat		Meningococcal (groups A, C, Y and		Miacalcic	
Losartan Actavis		W-135) conjugate vaccine 2		Micolette	
Losartan potassium		Meningococcal C conjugate		Miconazole	
Losartan potassium with		vaccine2		Miconazole nitrate	•
hydrochlorothiazide	48	Menthol		Dermatological	6
Lovir		Mepolizumab2		Genito-Urinary	
Loxamine		Mercaptopurine1		Micreme	
Lucrin Depot 1-month		Mercilon 28		Micreme H	
Lucrin Depot 3-month		Mesalazine		Microgynon 20 ED	
Ludiomil		Mesna1		Microgynon 30	
Lupin		Mestinon1		Microgynon 50 ED	
Lyderm		Metabolic Disorder Agents		Microlut	
Lynparza		Meterol2		Midazolam	
- M -		Metformin hydrochloride		Midazolam-Claris	
m-Eslon	124	Methadone hydrochloride		Midodrine	
Mabthera		Extemporaneous2		Mifegyne	
Macrogol 3350 with potassium		Nervous1		Mifepristone	
chloride, sodium bicarbonate	and	Methatabs1		Minerals	
sodium chloride		Methenamine (hexamine)		Mini-Wright AFS Low Range	
Macrogol 400 and propylene		hippurate1		Mini-Wright Standard	
glycol	245	Methopt2		Minidiab	
Madopar 125		Methotrexate1		MiniMed 640G	
Madopar 250		Methotrexate Ebewe1		MiniMed Mio MMT-921A	
Madopar 62.5		Methotrexate Sandoz1		MiniMed Mio MMT-923A	
Madopar HBS		Methyl hydroxybenzoate2		MiniMed Mio MMT-925A	
Madopar Rapid		Methylcellulose2		MiniMed Mio MMT-941A	
Magnesium hydroxide		Methylcellulose with glycerin and		MiniMed Mio MMT-943A	
Alimentary	36	sodium saccharin2		MiniMed Mio MMT-945A	
Extemporaneous		Methylcellulose with glycerin and		MiniMed Mio MMT-965A	
Magnesium sulphate		sucrose2		MiniMed Mio MMT-975A	
Mantoux		Methyldopa		MiniMed Quick-Set MMT-386A	
Maprotiline hydrochloride		Methyldopa Mylan		MiniMed Quick-Set MMT-387A	
Marevan		Methyldopa Mylan S29		MiniMed Quick-Set MMT-396A	
Marine Blue Lotion SPF 50+	70	Methylnaltrexone bromide		MiniMed Quick-Set MMT-397A	
Martindale Pharma		Methylphenidate ER - Teva1		MiniMed Quick-Set MMT-398A	
Marvelon 28		Methylphenidate hydrochloride 1		MiniMed Quick-Set MMT-399A	2
Mask for spacer device		Methylphenidate hydrochloride		MiniMed Silhouette MMT-368A	
Mast Cell Stabilisers		extended-release1		MiniMed Silhouette MMT-377A	
Maviret		Methylprednisolone		MiniMed Silhouette MMT-378A	
Maxidex		Methylprednisolone (as sodium		MiniMed Silhouette MMT-381A	
Maxitrol		succinate)		MiniMed Silhouette MMT-382A	

MiniMed Silhouette MMT-383A	22	Mycophenolate mofetil	180	Nilotinib	17
MiniMed Silhouette MMT-384A	22	Mydriacyl	244	Nilstat	
MiniMed Sure-T MMT-864A	20	Mylan Atenolol		Alimentary	3
MiniMed Sure-T MMT-866A	20	Mylan Clomiphen		Genito-Urinary	
MiniMed Sure-T MMT-874A		Myleran		Infection	
MiniMed Sure-T MMT-876A	20	Myometrial and Vaginal Hormone		Nintedanib	23
MiniMed Sure-T MMT-884A		Preparations	74	Nipent	
MiniMed Sure-T MMT-886A	20	Myozyme		Nitrados	
Minims Pilocarpine	244	Mysoline S29		Nitrates	
Minims Prednisolone		- N -		Nitrazepam	
Minirin	86	Nadolol	51	Nitroderm TTS	
Mino-tabs	92	Naglazyme	29	Nitrofurantoin	10
Minocycline hydrochloride	92	Nalcrom		Nitrolingual Pump Spray	50
Minomycin		Naloxone hydrochloride	246	Nivestim	4
Minor Skin Infections		Naltraccord		Nivolumab	22
Minoxidil	57	Naltrexone hydrochloride	156	Nizoral	
Mirena	81	Naphazoline hydrochloride	245	Nodia	
Mirtazapine	127	Naphcon Forte		Noflam 250	110
Misoprostol		Naprosyn SR 1000		Noflam 500	110
Mitomycin C	166	Naprosyn SR 750		Non-Steroidal Anti-Inflammatory	
Mitozantrone	166	Naproxen		Drugs	110
Mitozantrone Ebewe		Narcaricin mite		Nonacog gamma, [Recombinant	
Mixtard 30	11	Nardil	126	Factor IX]	40
MMR II	279	Nardil S29	126	Norethisterone	
Moclobemide	126	Nasal Preparations	239	Genito-Urinary	74
Modafinil	154	Natalizumab	140	Hormone	
Modavigil	154	Natulan	167	Norflex	118
Moduretic		Nausafix	132	Norfloxacin	10
Molaxole	27	Nausicalm	131	Noriday 28	
Moments	71	Navelbine	170	Norimin	
Mometasone furoate	64	Necon	73	Normacol Plus	2
Monogen	255	Nedocromil	239	Normison	150
Montelukast	238	Nefopam hydrochloride	122	Norpress	120
Montelukast Mylan	238	Neisvac-C	280	Nortriptyline hydrochloride	120
Moroctocog alfa [Recombinant fa	actor	Neo-B12	33	Norvir	10
VIII]	40	Neo-Mercazole	81	NovaSource Renal	25
Morphine hydrochloride	123	Neocate Gold	267	Novatretin	6
Morphine sulphate		Neocate Junior Unflavoured	267	NovoMix 30 FlexPen	
Motetis		Neocate Junior Vanilla	267	NovoRapid	1
Mouth and Throat		Neocate SYNEO	267	NovoRapid FlexPen	1
Movapo	119	Neoral	231	NovoRapid Penfill	1
Moxifloxacin	93	Neostigmine metilsulfate	110	NovoSeven RT	
MSUD Maxamum	265	Nepro HP (strawberry)	257	Noxafil	9
Mucilaginous laxatives with		Nepro HP (vanilla)	257	Nozinan13	
stimulants		Nepro HP RTH	257	Nozinan (Swiss)	13
Mucolytics	239	Nerisone	64	Nucala	20
Mucosoothe		Neulactil		Nuelin	
Multiple Sclerosis Treatments	137	Neulastim	44	Nuelin-SR	23
Multivitamin renal		NeuroTabs		Nutilis	
Multivitamins		Nevirapine		Nutrient Modules	
Mupirocin	62	Nevirapine Alphapharm		Nutrini Energy Multi Fibre	
Muscle Relaxants	118	Nicorandil	57	Nutrini Energy RTH	
Mvite		Nicotine		Nutrini Low Energy Multi Fibre	
Myambutol		Nicotinic acid		Nutrini Peptisorb	
Mycobutin		Nifedipine		Nutrini Peptisorb Energy	
MycoNail	62	Nifuran	109	Nutrini RTH	25

				_
Nutrison 800 Complete Multi	Ondansetron ODT-DRLA1	32	Pan-Penicillin G Sodium	91
Fibre261	One-Alpha	34	Panadol Mini Caps	122
Nutrison Concentrated263	Onrex1	32	Pancreatic enzyme	25
Nutrison Energy261	Opdivo22	29	Pantoprazole	9
Nutrison Energy Multi Fibre261	Ora-Blend2		Panzop Relief	9
Nutrison Multi Fibre261	Ora-Blend SF24	49	Panzytrat	
Nutrison Standard RTH261	Ora-Plus24	49	Papaverine hydrochloride	
Nyefax Retard52	Ora-Sweet24		Para-amino salicylic acid	
Nystatin	Ora-Sweet SF24		Paracare	
Alimentary33	Orabase		Paracare Double Strength	
Genito-Urinary74	Oral Supplements/Complete Diet	•	Paracetamol	
Infection96	(Nasogastric/Gastrostomy Tube		Paracetamol + Codeine	122
NZB Low Gluten Bread Mix264	Feed) 2	54	(Relieve)	124
- O -	Oratane		Paracetamol Pharmacare	
O/W Fatty Emulsion Cream66	Ordine1		Paracetamol with codeine	
Obinutuzumab	Orgran		Paradigm 1.8 Reservoir	
Obstetric Preparations76	Ornidazole		Paradigm 3.0 Reservoir	
Ocicure9	Orphenadrine citrate1		Paradigm Mio MMT-921	
Ocrelizumab141	Ortho-tolidine		Paradigm Mio MMT-923	
Ocrevus141	Oruvail SR1		Paradigm Mio MMT-925	
Octocog alfa [Recombinant factor	Osmolite RTH20		Paradigm Mio MMT-941	
VIII] (Advate)40	Other Endocrine Agents		Paradigm Mio MMT-943	
Octocog alfa [Recombinant factor	Other Oestrogen Preparations	81	Paradigm Mio MMT-945	
VIII] (Kogenate FS)40	Other Progestogen		Paradigm Mio MMT-965	
Octreotide	Preparations		Paradigm Mio MMT-975	24
Octreotide (Sun) 178	Other Skin Preparations	70	Paradigm Quick-Set MMT-386	25
Octreotide GH178	Ovestin		Paradigm Quick-Set MMT-387	25
Octreotide LAR (somatostatin	Genito-Urinary	75	Paradigm Quick-Set MMT-396	25
analogue) 178	Hormone	81	Paradigm Quick-Set MMT-397	25
Octreotide MaxRx178	Ox-Pam1	37	Paradigm Quick-Set MMT-398	25
Oestradiol80	Oxaliplatin1		Paradigm Quick-Set MMT-399	
Oestradiol valerate80	Oxaliplatin Accord1	60	Paradigm Silhouette MMT-368	
Oestradiol with norethisterone80	Oxaliplatin Actavis 1001		Paradigm Silhouette MMT-377	
Oestriol	Oxaliplatin Ebewe1		Paradigm Silhouette MMT-378	
Genito-Urinary75	Oxazepam1		Paradigm Silhouette MMT-381	
Hormone81	Oxis Turbuhaler2		Paradigm Silhouette MMT-382	
Oestrogens 80	Oxpentifylline		Paradigm Silhouette MMT-383	
Ofev237	Oxybutynin		Paradigm Silhouette MMT-384	
Oil in water emulsion	Oxycodone hydrochloride12		Paradigm Sure-T MMT-864	
Olanzapine	Oxycodone Sandoz12		Paradigm Sure-T MMT-866	
Olaparib	OxyNorm12		Paradigm Sure-T MMT-874	
Olbetam54	Oxytocin		Paradigm Sure-T MMT-876	
			Paradigm Sure-T MMT-884	
Olbetam S2954	Oxytocin BNM	/5		
Olopatadine	Oxytocin with ergometrine	75	Paradigm Sure-T MMT-886	
Olopatadine Teva245	maleate		Paraffin	
Olsalazine7	Ozurdex24	42	Paraffin liquid with wool fat	
Omalizumab206	-P-		Paraldehyde	
Omeprazole9	Pacifen1		Parasidose	
Omeprazole actavis 109	Paclitaxel10		Parasiticidal Preparations	
Omeprazole actavis 209	Paclitaxel Actavis1		Parnate	
Omeprazole actavis 409	Paclitaxel Ebewe1		Parnate S29	
Omnitrope82	Paediatric Seravit		Paromomycin	94
Onbrez Breezhaler235	Palbociclib1		Paroxetine	127
Oncaspar LYO167	Paliperidone1	35	Paser	99
OncoTICE187	Pamidronate disodium1		Patanol	245
Ondansetron132	Pamisol1	13	Paxam	136

Pazopanib	174	and fluorescein	69	Premarin	8
Peak flow meter	240	Pinetarsol	69	Prevenar 13	28
Pedialyte - Bubblegum	45	Pioglitazone	12	Prezista	100
Pediasure	256	Pirfenidone	238	Priadel	13
Pediasure RTH	256	Pizotifen	131	Primacin	9
Pegaspargase	167	PKU Anamix Infant	266	Primaquine	9
Pegasys	107	PKU Anamix Junior	266	Primidone	12
Pegfilgrastim		PKU Anamix Junior Chocolate	266	Primolut N	8
Pegylated interferon alfa-2a		PKU Anamix Junior LQ		Priorix	27
Pembrolizumab		PKU Anamix Junior Vanilla	266	Probenecid	11
Pemetrexed	162	PKU Lophlex LQ 10	266	Probenecid-AFT	11
Penicillamine		PKU Lophlex LQ 20		Procaine penicillin	9
Penicillin G	91	PKU Lophlex Powder	266	Procarbazine hydrochloride	16
PenMix 30	11	PKU Lophlex Sensation 20		Prochlorperazine	
PenMix 40	11	Plaquenil		Proctofoam	
PenMix 50		Plendil ER		Proctosedyl	
Pentasa	7	Pneumococcal (PCV10) conjugate		Procyclidine hydrochloride	120
Pentostatin [Deoxycoformycin]	167	vaccine	280	Procytox	
Pentoxifylline [Oxpentifylline]		Pneumococcal (PCV13) conjugate		Progesterone	
Peptamen Junior		vaccine	281	Proglicem	
Peptisoothe		Pneumococcal (PPV23)		Proglycem	
Peptisorb		polysaccharide vaccine	282	Progynova	
Perhexiline maleate		Pneumovax 23		Prolia	
Pericyazine		Podophyllotoxin		Promethazine hydrochloride	
Perindopril		Polaramine		Propafenone hydrochloride	
Perjeta		Poliomyelitis vaccine		Propamidine isethionate	
Permethrin		Poloxamer		Propranolol	
Perrigo		Poly-Gel		Propylene glycol	
Pertuzumab		Poly-Tears		Propylthiouracil	
Peteha	99	Poly-Visc		Protaphane	1
Pethidine hydrochloride	125	Polycal	251	Protaphane Penfill	1
Pevaryl		Ponstan	110	Protifar	25
Pexsig		Posaconazole		Protionamide	
Pfizer Exemestane		Postinor-1		Provera	8
Pharmacy Services		Potassium chloride		Provera HD	
Pheburane		Potassium Chloride Aguettant		PSM Citalopram	
Phenasen		Potassium citrate		Psoriasis and Eczema	
Phenelzine sulphate	126	Potassium iodate		Preparations	6
Phenobarbitone		Povidone iodine		PTU	
Phenobarbitone sodium		Pradaxa	43	Pulmicort Turbuhaler	23
Extemporaneous	250	Pramipexole hydrochloride	119	Pulmocare	25
Nervous		Prasugrel		Pulmozyme	
Phenothrin	68	Pravastatin		Puri-nethol	
Phenoxybenzamine		Pravastatin Mylan	55	Puria	3
hydrochloride	47	Praziquantel		Pyrazinamide	
Phenoxymethylpenicillin (Penicillin		Prazosin		Pyridostigmine bromide	
V)		Pred Forte		Pyridoxine hydrochloride	
Phenytoin sodium12	8-129	Prednisolone	78	Pyrimethamine	
Phlexy 10		Prednisolone acetate	243	Pytazen SR	4
Phosphate Phebra		Prednisolone sodium		- Q -	
Phosphorus		phosphate	243	Q 300	9
Phytomenadione		Prednisolone-AFT	243	Quetapel	
Pilocarpine hydrochloride		Prednisone		Quetiapine	13
Pimafucort	65	Pregabalin		Quick-Set MMT-392	2
Pindolol		Pregabalin Pfizer		Quick-Set MMT-393	
Pine tar with trolamine laurilsulfate		Pregnancy Tests - hCG Urine		Quinapril	
		· ·			

Quinapril with		Rituximab (Riximyo)	213	Seretide	23
hydrochlorothiazide	48	Rivaroxaban		Seretide Accuhaler	23
Quinine sulphate	98	Rivastigmine	154	Serevent	23
Qvar	234	Rivotril		Serevent Accuhaler	23
- R -		Riximyo		Sertraline	12
RA-Morph	123	RIXUBIS	40	Setrona	12
Raloxifene hydrochloride	113	Rizamelt	131	Sevredol	12
Raltegravir potassium		Rizatriptan	131	Sex Hormones Non	
Ramipex		Robinul	8	Contraceptive	7
Ranbaxy-Cefaclor		Roferon-A	107	Shield XL	
Ranitidine		Rolin	180	shingles vaccine	28
Ranitidine Relief	9	Ropin	119	SII-Onco-BCG	18
Rapamune	232	Ropinirole hydrochloride	119	Sildenafil	5
Reandron 1000		Rotarix		Silhouette MMT-373	2
Recombinant factor IX	38, 40	Rotavirus oral vaccine	282	Siltuximab	22
Recombinant factor VIIa		Roxane		Simvastatin	
Recombinant factor VIII	40	Alimentary	6	Simvastatin Mylan	
Rectogesic	8	Cardiovascular	51	Sinemet	11
Redipred		Roxithromycin	90	Sinemet CR	
Relieve		Rubifen		Sirolimus	23
Relistor	27	Rubifen SR		Siterone	7
Remicade		Rugby Capsaicin Topical C	ream	Slow-Lopresor	5
Renilon 7.5		Musculoskeletal		Smith BioMed Rapid Pregnancy	
Resonium-A		Nervous		Test	7
Resource Beneprotein	253	Rulide D	90	Sodibic	
Resource Diabetic		Rurioctocog alfa pegol [Red	combinant	Sodium acid phosphate	
Respigen		factor VIII]		Sodium alginate	
Respiratory Devices		Ruxolitinib		Sodium benzoate	
Respiratory Stimulants		Rythmodan	49	Sodium bicarbonate	
Retinol palmitate		Rytmonorm		Blood	.45-4
ReTrieve		· - \$ -		Extemporaneous	
Retrovir		Sabril	130	Sodium calcium edetate	
Revlimid	165	Sacubitril with valsartan	48	Sodium chloride	
Revolade		SalAir	236	Blood	4
Rexacrom		Salazopyrin		Respiratory	
Ribomustin		Salazopyrin EN		Sodium citrate with sodium lauryl	
Ricit		Salbutamol		sulphoacetate	
Rifabutin		Salbutamol with ipratropiun		Sodium citro-tartrate	
Rifadin		bromide		Sodium cromoglicate	
Rifampicin		Salicylic acid		Alimentary	
Rifaximin		Salmeterol		Respiratory	
Rifinah		Sandomigran		Sensory	
Rilutek		Sandostatin LAR		Sodium fluoride	
Riluzole		Sapropterin dihydrochloride		Sodium Fusidate [fusidic acid]	
Riodine		Scalp Preparations		Dermatological	6
Risedronate Sandoz		Scopoderm TTS	132	Infection	
Risedronate sodium		Sebizole		Sensory	
Risperdal Consta		Secukinumab		Sodium hyaluronate [Hyaluronic	
Risperidone		Sedatives and Hypnotics		acid]	24
Risperidone (Teva)		Seebri Breezhaler		Sodium phenylbutyrate	3
Risperon		Selegiline hydrochloride		Sodium polystyrene sulphonate	
Ritalin		Senna		Sodium tetradecyl sulphate	
Ritalin LA		Senokot		Sodium valproate	
Ritalin SR		Sensipar		Sofradex	
Ritonavir		SensoCard		Soframycin	
Rituximab (Mabthera)		Serenace		Solifenacin Mylan	
·					/ '

Solifenacin succinate	76	Syntometrine	75	Tilcotil	110
Solu-Cortef	78	Syrup (pharmaceutical grade).	250	Tillomed	159
Solu-Medrol	78	Systane Unit Dose	245	Timolol	
Solu-Medrol-Act-O-Vial	78	- T -		Cardiovascular	51
Somatropin (Omnitrope)	82	Tacrolimus	232	Sensory	243
Sotalol	51	Tacrolimus Sandoz	232	Timoptol XE	
Spacer device	240	Taliglucerase alfa	31	Tiotropium bromide	
Span-K	46	Tambocor		Tiotropium bromide with	
Spiolto Respimat		Tamoxifen citrate		olodaterol	237
Spiractin		Tamoxifen Sandoz	180	Tivicay	106
Spiriva	237	Tamsulosin hydrochloride	75	TMP	
Spiriva Respimat		Tamsulosin-Rex	75	TOBI	
Spironolactone		Tandem Basal IQ		Tobramycin	
Sporanox		Tandem Cartridge	19	Infection	95
Sprycel		Tandem t:slim X2		Sensory	242
Staphlex		Tap water		Tobramycin Mylan	95
Stemetil		Tarceva		Tobrex	242
SteroClear		Tasigna		Tocilizumab	
Stesolid		Tasmar		Tofranil	
Stimulants/ADHD Treatments		Tecfidera		Tolcapone	
Stiripentol		Tegretol		Topamax	
Stocrin		Tegretol CR		Topical Products for Joint and	
Stomahesive		Telfast		Muscular Pain	111
Strattera		Teligent		Topiramate	
Strides Shasun		Temaccord		Topiramate Actavis	
Stromectol		Temazepam		Total parenteral nutrition (TPN)	
Sucralfate		Temozolomide		TPN	
Sulfadiazine Silver		Tenofovir disoproxil		Tramadol hydrochloride	
Sulfadiazine sodium				Tramal SR 100	
Sulfasalazine		Tenofovir Disoproxil Teva		Tramal SR 150	
		Tenoxicam			
Sulindac		Tepadina		Tramal SR 200	
Sulindac Mylan		Terazosin			
Sulphur		Terbinafine		Tranexamic acid	
Sulprix		Terbutaline sulphate		Tranylcypromine sulphate	
Sumatriptan		Teriflunomide		Trastuzumab	
Sunitinib		Teriparatide	113	Trastuzumab emtansine	
Sunscreens		Testosterone		Travatan	
Sunscreens, proprietary		Testosterone cipionate		Travoprost	244
Sure-T MMT-863		Testosterone esters		Travopt	
Sure-T MMT-873		Testosterone undecanoate		Treatments for Dementia	154
Sustagen Diabetic	254	Tetrabenazine		Treatments for Substance	
Sustagen Hospital Formula		Tetrabromophenol		Dependence	
Active		Tetracosactrin		Trental 400	57
Sustanon Ampoules		Tetracyclin Wolff		Tretinoin	
Sutent		Tetracycline		Dermatological	
Sylvant		Thalidomide	168	Oncology	169
Symbicort Turbuhaler 100/6	235	Thalomid		Trexate	162
Symbicort Turbuhaler 200/6	235	Theophylline	239	Triamcinolone acetonide	
Symbicort Turbuhaler 400/12		Thiamine hydrochloride		Alimentary	33
Symmetrel	119	THIO-TEPA	160	Dermatological	64
Sympathomimetics		Thioguanine		Hormone	
Synacthen	79	Thiotepa	160	Triamcinolone acetonide with	
Synacthen Depot		Thymol glycerin	33	gramicidin, neomycin and nyst	atin
Synacthene Retard		Thyroid and Antithyroid Agents	81	Dermatological	
Synflorix		Ticagrelor	41	Sensory	
Synthroid		Tilade		Triaver	

Triazolam	150	Venclexta	169	XP Maxamum	26
Triclosan	65	Venetoclax	169	Xylocaine	12
Trimethoprim	95	Venlafaxine	127	Xylocaine 2% Jelly	12
Trimethoprim with		Venomil	233	Xyntha	4
sulphamethoxazole		Ventavis	59	- Z -	
[Co-trimoxazole]	95	Ventolin	236	Zantac	
Trisequens	80	Vepesid	164	Zapril	
Trisul	95	Verapamil hydrochloride	52	Zarontin	
Trophic Hormones	82	Vergo 16		Zaroxolyn	
Tropicamide	244	Vermox	88	Zavedos	16
Trusopt	243	Vesanoid	169	Zeffix	10
TruSteel	<mark>21</mark>	Vexazone	12	Zetlam	10
Tuberculin PPD [Mantoux] test	283	Vfend	97	Ziagen	10
Tubersol	283	Viaderm KC	65	Zidovudine [AZT]	10
Two Cal HN	263	Vidaza	160	Zidovudine [AZT] with	
Two Cal HN RTH		Vigabatrin	130	lamivudine	
Tykerb	172	Vildagliptin	12	Zimybe	
Tysabri	140	Vildagliptin with metformin		Zinc and castor oil	6
- U -		hydrochloride	12	Zinc sulphate	3
Ultibro Breezhaler	237	Vimpat	128	Zincaps	3
Ultraproct	8	Vinblastine sulphate	170	Zinnat	
Umeclidinium	237	Vincristine sulphate	170	Ziprasidone	13
Umeclidinium with vilanterol	237	Vinorelbine		Zista	23
Univent	.236, 240	Vinorelbine Ebewe	170	Zithromax	8
Ural	76	Viramune Suspension	105	Zoladex	8
Urea	66	ViruPOS	241	Zoledronic acid	
Urex Forte	53	Vit.D3	34	Hormone	<mark>7</mark>
Urinary Agents	75	VitA-POS	245	Musculoskeletal	11
Urinary Tract Infections		Vitabdeck	34	Zoledronic acid Mylan	<mark>7</mark>
Urinorm		Vital	257	Zopiclone	
Uromitexan	166	Vitamin B complex	33	Zopiclone Actavis	15
Ursodeoxycholic acid	26	Vitamin B6 25	33	Zostavax	28
Ursosan		Vitamins	33–34	Zostrix	
Utrogestan		Vivonex TEN		Zostrix HP	12
- V -		Volibris		Zuclopenthixol decanoate	
Vaccinations	272	Voltaren		Zuclopenthixol hydrochloride	
Vaclovir		Voltaren D		Zusdone	
Valaciclovir		Voltaren Ophtha		Zyban	
Valganciclovir		Volumatic		Zypine	
Valganciclovir Mylan		Voriconazole		Zypine ODT	
Vancomycin		Vosol	241	Zyprexa Relprevv	13
Vannair		Votrient		Zytiga	17
Varenicline Pfizer		Vttack	97	, 0	
Varenicline tartrate		- W -			
Varicella vaccine [Chickenpox		Warfarin sodium	44		
vaccine]	283	Wart Preparations			
Varicella zoster virus (Oka stra		Wasp venom allergy treatme			
attenuated vaccine [shingles		Water			
vaccine]		Blood	45		
Varilrix		Extemporaneous			
Various		Wool fat with mineral oil			
Varivax		- X -			
Vasodilators		Xarelto	44		
Vasopressin Agonists		Xifaxan			
Vedafil		XMET Maxamum			
Veletri		Xolair			