

## December 2020

### Volume 8 Number 4

#### Editor:

Kaye Wilson & Doris Chong  
email: [enquiry@pharmac.govt.nz](mailto:enquiry@pharmac.govt.nz)  
Telephone +64 4 460 4990  
Facsimile +64 4 460 4995  
Level 9, 40 Mercer Street  
PO Box 10 254 Wellington 6143

#### Freephone Information Line

0800 66 00 50 (9am – 5pm weekdays)

#### Circulation

Published each March, July and November.  
Changes to the contents are published in  
monthly updates.

Accessible in an electronic format at  
no cost from the PHARMAC website  
[www.pharmac.govt.nz/schedule](http://www.pharmac.govt.nz/schedule).

You can register to have an electronic  
version of the Pharmaceutical Schedule,  
Section H for Hospital Pharmaceuticals (link  
to PDF copy) emailed to your nominated  
email address each month by subscribing at  
[schedule.pharmac.govt.nz/subscribe](http://schedule.pharmac.govt.nz/subscribe).

#### Production

Typeset automatically from XML and  $\text{\TeX}$ .  
XML version of the Schedule available from  
[schedule.pharmac.govt.nz/pub/HML](http://schedule.pharmac.govt.nz/pub/HML)

#### Programmers

Anrik Drenth & John Geering

email: [texschedule@pharmac.govt.nz](mailto:texschedule@pharmac.govt.nz)

©Pharmaceutical Management Agency



ISSN 1179-3708 pdf

ISSN 1172-9694 print

This work is licensed under the Creative  
Commons Attribution 4.0 International licence.

In essence, you are free to copy, distribute  
and adapt it, as long as you attribute the work  
to PHARMAC and abide by the other licence  
terms. To view a copy of this licence, visit:  
[creativecommons.org/licenses/by/4.0/](http://creativecommons.org/licenses/by/4.0/).

Attribution to PHARMAC should be in  
written form and not by reproduction of the  
PHARMAC logo. While care has been taken  
in compiling this Schedule, PHARMAC takes  
no responsibility for any errors or omissions,  
and shall not be liable for any consequences  
arising there from.

Part I	General Rules	4
--------	---------------	---

Part II	Alimentary Tract and Metabolism	5
	Blood and Blood Forming Organs	24
	Cardiovascular System	39
	Dermatologicals	54
	Genito-Urinary System	60
	Hormone Preparations	64
	Infections	74
	Musculoskeletal System	97
	Nervous System	106
	Oncology Agents and Immunosuppressants	130
	Respiratory System and Allergies	204
	Sensory Organs	212
	Various	219
	Extemporaneous Compounds (ECPs)	227
	Special Foods	230
	Vaccines	246

Part III	Optional Pharmaceuticals	257
----------	--------------------------	-----

	Index	258
--	-------	-----

# Introducing PHARMAC

The Pharmaceutical Management Agency (PHARMAC) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. PHARMAC negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

## PHARMAC's role:

*“Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided.”*

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about PHARMAC and the way we make funding decisions can be found on the PHARMAC website at <https://www.pharmac.govt.nz/about>.

## Glossary

### Units of Measure

gram .....	g	microgram.....	mcg	millimole.....	mmol
kilogram.....	kg	milligram.....	mg	unit.....	u
international unit.....	iu	millilitre.....	ml		

### Abbreviations

application .....	app	enteric coated .....	EC	solution .....	soln
capsule .....	cap	granules.....	grans	suppository.....	suppos
cream.....	crm	injection .....	inj	tablet.....	tab
dispersible .....	disp	liquid .....	liq	tincture.....	tinc
effervescent.....	eff	lotion .....	lotn		
emulsion .....	emul	ointment.....	oint		

HSS    Hospital Supply Status

# Guide to Section H listings

## Example

ANATOMICAL HEADING			
	Price (ex man. Excl. GST) \$	Per	Brand or Generic Manufacturer
THERAPEUTIC HEADING			
Generic name listed by therapeutic group and subgroup	CHEMICAL A - <b>Restricted</b> see terms below ⚡ Presentation A.....10.00	100	Brand A
	➡ <b>Restricted</b> Only for use in children under 12 years of age		Brand or manufacturer's name
Indicates only presentation B1 is Restricted	CHEMICAL B - <b>Some items restricted</b> see terms below ⚡ Presentation B1.....1,589,00 Presentation B2 ➡ <b>Restricted</b> Oncologist or haematologist	1	Brand B1 e.g. Brand B2
From 1 January 2012 to 30 June 2014, at least 99% of the total volume of this item purchased must be Brand C	CHEMICAL C Presentation C - <b>-1% DV Limit Jan-12 to 2014</b> .....15.00	28	Brand C
	CHEMICAL D - <b>Restricted</b> see terms below ⚡ Presentation D - <b>-1% DV Limit Mar-13 to 2014</b> .....38.65	500	Brand D
Standard national price excluding GST	➡ <b>Restricted</b> <i>Limited to five weeks' treatment</i> Either: 1 For the prophylaxis of venous thromboembolism following a total hip replacement; or 2 For the prophylaxis of venous thromboembolism following a total knee replacement.		Quantity the Price applies to
Form and strength	CHEMICAL E Presentation E		e.g. Brand E
			Not a contracted product
⚡ Item restricted (see above); ⚡ Item restricted (see below) Products with Hospital Supply Status (HSS) are in <b>bold</b>			

## PART I: GENERAL RULES

---

General Rules for Section H of the Pharmaceutical Schedule are included in Section A.

Read the [General Rules](https://www.pharmac.govt.nz/section-a) : <https://www.pharmac.govt.nz/section-a>.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## Antacids and Antiflatulents

### Antacids and Reflux Barrier Agents

#### ALUMINIUM HYDROXIDE WITH MAGNESIUM HYDROXIDE AND SIMETICONE

Tab 200 mg with magnesium hydroxide 200 mg and simeticone 20 mg

*e.g. Mylanta*

Oral liq 400 mg with magnesium hydroxide 400 mg and simeticone

*e.g. Mylanta Double Strength*

30 mg per 5 ml

#### SIMETICONE

Oral drops 100 mg per ml

Oral drops 20 mg per 0.3 ml

Oral drops 40 mg per ml

#### SODIUM ALGINATE WITH MAGNESIUM ALGINATE

Powder for oral soln 225 mg with magnesium alginate 87.5 mg, sachet

*e.g. Gaviscon Infant*

#### SODIUM ALGINATE WITH SODIUM BICARBONATE AND CALCIUM CARBONATE

Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate

160 mg

*e.g. Gaviscon Double Strength*

Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate

160 mg per 10 ml.....4.95

500 ml

Acidex

#### SODIUM CITRATE

Oral liq 8.8% (300 mmol/l)

## Phosphate Binding Agents

#### ALUMINIUM HYDROXIDE

Tab 600 mg

#### CALCIUM CARBONATE – **Restricted** see terms [below](#)

↓ Oral liq 250 mg per ml (100 mg elemental per ml) .....39.00

500 ml

Roxane

→ **Restricted (RS1698)**

#### Initiation

Only when prescribed for patients unable to swallow calcium carbonate tablets or where calcium carbonate tablets are inappropriate..

## Antidiarrhoeals and Intestinal Anti-Inflammatory Agents

### Antipropulsives

#### DIPHENOXYLATE HYDROCHLORIDE WITH ATROPINE SULPHATE

Tab 2.5 mg with atropine sulphate 25 mcg

#### LOPERAMIDE HYDROCHLORIDE

Tab 2 mg .....10.75

400

Nodia

Cap 2 mg – **1% DV Oct-19 to 2022** .....6.25

400

**Diamide Relief**

## Rectal and Colonic Anti-Inflammatories

#### BUDESONIDE – **Restricted** see terms [on the next page](#)

↓ Cap 3 mg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➔ Restricted (RS1723)</b>			
<b>Initiation – Crohn's disease</b>			
Both:			
1	Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and		
2	Any of the following:		
2.1	Diabetes; or		
2.2	Cushingoid habitus; or		
2.3	Osteoporosis where there is significant risk of fracture; or		
2.4	Severe acne following treatment with conventional corticosteroid therapy; or		
2.5	History of severe psychiatric problems associated with corticosteroid treatment; or		
2.6	History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or		
2.7	Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).		
<b>Initiation – Collagenous and lymphocytic colitis (microscopic colitis)</b>			
Patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.			
<b>Initiation – Gut Graft versus Host disease</b>			
Patient has gut Graft versus Host disease following allogenic bone marrow transplantation.			
<b>Initiation – non-cirrhotic autoimmune hepatitis</b>			
<i>Re-assessment required after 6 months</i>			
All of the following:			
1	Patient has autoimmune hepatitis*; and		
2	Patient does not have cirrhosis; and		
3	Any of the following:		
3.1	Diabetes; or		
3.2	Cushingoid habitus; or		
3.3	Osteoporosis where there is significant risk of fracture; or		
3.4	Severe acne following treatment with conventional corticosteroid therapy; or		
3.5	History of severe psychiatric problems associated with corticosteroid treatment; or		
3.6	History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or		
3.7	Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or		
3.8	Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).		
Note: Indications marked with * are unapproved indications.			
<b>Continuation – non-cirrhotic autoimmune hepatitis</b>			
<i>Re-assessment required after 6 months</i>			
Treatment remains appropriate and the patient is benefitting from the treatment.			
<b>HYDROCORTISONE ACETATE</b>			
Rectal foam 10%, CFC free (14 applications) .....	26.55	21.1 g	Colifoam
<b>HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE</b>			
Topical Aerosol foam, 1% with pramoxine hydrochloride 1%			
<b>MESALAZINE</b>			
Tab EC 400 mg .....	49.50	100	Asacol
Tab EC 500 mg .....	49.50	100	Asamax
Tab long-acting 500 mg – 1% DV Jul-20 to 2023 .....	56.10	100	<b>Pentasa</b>
Tab 800 mg .....	85.50	90	Asacol
Modified release granules 1 g .....	141.72	120 g	Pentasa
Suppos 500 mg .....	22.80	20	Asacol
Suppos 1 g .....	54.60	30	Pentasa
Enema 1 g per 100 ml .....	41.30	7	Pentasa

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>OLSALAZINE</b>			
Tab 500 mg .....	93.37	100	Dipentum
Cap 250 mg .....	53.00	100	Dipentum
<b>PREDNISOLONE SODIUM</b>			
Rectal foam 20 mg per dose (14 applications) .....	74.10	1	Essential Prednisolone
<b>SODIUM CROMOGLICATE</b>			
Cap 100 mg			
<b>SULFASALAZINE</b>			
Tab 500 mg .....	14.00	100	Salazopyrin
Tab EC 500 mg – <b>1% DV Dec-19 to 2022</b> .....	15.53	100	<b>Salazopyrin EN</b>

## Local Preparations for Anal and Rectal Disorders

### Antihaemorrhoidal Preparations

<b>CINCHOCAINE HYDROCHLORIDE WITH HYDROCORTISONE</b>			
Oint 5 mg with hydrocortisone 5 mg per g .....	15.00	30 g	Proctosedyl
Suppos 5 mg with hydrocortisone 5 mg per g .....	9.90	12	Proctosedyl
<b>FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE</b>			
Oint 950 mcg with fluocortolone pivalate 920 mcg and cinchocaine hydrochloride 5 mg per g .....	6.35	30 g	Ultraproct
Suppos 630 mcg with fluocortolone pivalate 610 mcg and cinchocaine hydrochloride 1 mg .....	2.66	12	Ultraproct

### Management of Anal Fissures

<b>GLYCERYL TRINITRATE</b>			
Oint 0.2% .....	22.00	30 g	Rectogesic

### Rectal Sclerosants

<b>OILY PHENOL [PHENOL OILY]</b>			
Inj 5%, 5 ml vial			

## Antispasmodics and Other Agents Altering Gut Motility

<b>GLYCOPYRRONIUM BROMIDE</b>			
Inj 200 mcg per ml, 1 ml ampoule .....	17.14	10	Max Health
<b>HYOSCINE BUTYLBROMIDE</b>			
Tab 10 mg – <b>1% DV Oct-20 to 2023</b> .....	6.35	100	<b>Buscopan</b>
Inj 20 mg, 1 ml ampoule – <b>1% DV Jul-20 to 2023</b> .....	6.35	5	<b>Buscopan</b>
<b>MEBEVERINE HYDROCHLORIDE</b>			
Tab 135 mg – <b>1% DV Jul-20 to 2023</b> .....	9.20	90	<b>Colofac</b>

## Antulcerants

### Antisecretory and Cytoprotective

<b>MISOPROSTOL</b>			
Tab 200 mcg .....	41.50	120	Cytotec

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>H2 Antagonists</b>			
CIMETIDINE			
Tab 200 mg			
Tab 400 mg			
FAMOTIDINE			
Tab 20 mg			
Tab 40 mg			
Inj 10 mg per ml, 2 ml vial			
Inj 10 mg per ml, 4 ml vial			
RANITIDINE – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Tab 150 mg			
↓ Tab 300 mg			
↓ Oral liq 150 mg per 10 ml .....	5.14	300 ml	Peptisoothe
↓ Inj 25 mg per ml, 2 ml ampoule .....	13.40	5	Zantac
<i>(Peptisoothe Oral liq 150 mg per 10 ml to be delisted 1 September 2021)</i>			
<i>(Zantac Inj 25 mg per ml, 2 ml ampoule to be delisted 1 March 2021)</i>			
➔ <b>Restricted</b> ( <a href="#">RS1703</a> )			
<b>Initiation</b>			
Either:			
1 For continuation use; or			
2 Routine prevention of allergic reactions..			
<b>Proton Pump Inhibitors</b>			
LANSOPRAZOLE			
Cap 15 mg – <b>1% DV Sep-18 to 2021</b> .....	4.58	100	Lanzol Relief
Cap 30 mg – <b>1% DV Sep-18 to 2021</b> .....	5.41	100	Lanzol Relief
OMEPRAZOLE			
↓ Tab dispersible 20 mg			
➔ <b>Restricted</b> ( <a href="#">RS1027</a> )			
<b>Initiation</b>			
Only for use in tube-fed patients.			
Cap 10 mg .....	1.98	90	Omeprazole actavis 10
Cap 20 mg .....	1.96	90	Omeprazole actavis 20
Cap 40 mg .....	3.12	90	Omeprazole actavis 40
Powder for oral liq .....	42.50	5 g	Midwest
Inj 40 mg ampoule with diluent – <b>1% DV Oct-19 to 2022</b> .....	33.98	5	Dr Reddy's Omeprazole
Inj 40 mg vial – <b>1% DV Oct-19 to 2022</b> .....	11.46	5	Omezol IV
PANTOPRAZOLE			
Tab EC 20 mg – <b>1% DV Oct-19 to 2022</b> .....	2.02	100	Panzop Relief
Tab EC 40 mg – <b>1% DV Oct-19 to 2022</b> .....	2.85	100	Panzop Relief
Inj 40 mg vial			
<b>Site Protective Agents</b>			
COLLOIDAL BISMUTH SUBCITRATE			
Tab 120 mg .....	14.51	50	Gastrodenol
SUCRALFATE			
Tab 1 g			

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

## Bile and Liver Therapy

L-ORNITHINE L-ASPARTATE – **Restricted** see terms [below](#)

↓ Grans for oral liquid 3 g

→ **Restricted (RS1261)**

### Initiation

For patients with chronic hepatic encephalopathy who have not responded to treatment with, or are intolerant to lactulose, or where lactulose is contraindicated.

RIFAXIMIN – **Restricted** see terms [below](#)

↓ Tab 550 mg – 1% DV Mar-21 to 2023 ..... 625.00 56 **Xifaxan**

→ **Restricted (RS1416)**

### Initiation

For patients with hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

## Diabetes

### Alpha Glucosidase Inhibitors

ACARBOSE

Tab 50 mg – 1% DV Sep-18 to 2021 ..... 3.50 90 **Glucobay**

Tab 100 mg – 1% DV Sep-18 to 2021 ..... 6.40 90 **Glucobay**

### Hyperglycaemic Agents

DIAZOXIDE – **Restricted** see terms [below](#)

↓ Cap 25 mg ..... 110.00 100 **Proglidem**

↓ Cap 100 mg ..... 280.00 100 **Proglidem**

↓ Oral liq 50 mg per ml ..... 620.00 30 ml **Proglycem**

→ **Restricted (RS1028)**

### Initiation

For patients with confirmed hypoglycaemia caused by hyperinsulinism.

GLUCAGON HYDROCHLORIDE

Inj 1 mg syringe kit – 1% DV Jul-20 to 2023 ..... 32.00 1 **Glucagen Hypokit**

GLUCOSE [DEXTROSE]

Tab 1.5 g

Tab 3.1 g

Tab 4 g

Gel 40%

GLUCOSE WITH SUCROSE AND FRUCTOSE

Gel 19.7% with sucrose 35% and fructose 19.7%, 18 g sachet

### Insulin - Intermediate-Acting Preparations

INSULIN ASPART WITH INSULIN ASPART PROTAMINE

Inj insulin aspart 30% with insulin aspart protamine 70%, 100 u per ml,  
3 ml prefilled pen ..... 52.15 5 **NovoMix 30 FlexPen**

INSULIN ISOPHANE

Inj insulin human 100 u per ml, 10 ml vial

Inj insulin human 100 u per ml, 3 ml cartridge

## ALIMENTARY TRACT AND METABOLISM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>INSULIN LISPRO WITH INSULIN LISPRO PROTAMINE</b>			
Inj insulin lispro 25% with insulin lispro protamine 75%, 100 u per ml, 3 ml cartridge.....	42.66	5	Humalog Mix 25
Inj insulin lispro 50% with insulin lispro protamine 50%, 100 u per ml, 3 ml cartridge.....	42.66	5	Humalog Mix 50
<b>INSULIN NEUTRAL WITH INSULIN ISOPHANE</b>			
Inj insulin neutral 30% with insulin isophane 70%, 100 u per ml, 10 ml vial			
Inj insulin neutral 30% with insulin isophane 70%, 100 u per ml, 3 ml cartridge			
Inj insulin neutral 40% with insulin isophane 60%, 100 u per ml, 3 ml cartridge			
Inj insulin neutral 50% with insulin isophane 50%, 100 u per ml, 3 ml cartridge			
<b>Insulin - Long-Acting Preparations</b>			
<b>INSULIN GLARGINE</b>			
Inj 100 u per ml, 3 ml disposable pen.....	94.50	5	Lantus SoloStar
Inj 100 u per ml, 3 ml cartridge.....	94.50	5	Lantus
Inj 100 u per ml, 10 ml vial.....	63.00	1	Lantus
<b>Insulin - Rapid-Acting Preparations</b>			
<b>INSULIN ASPART</b>			
Inj 100 u per ml, 10 ml vial			
Inj 100 u per ml, 3 ml cartridge			
Inj 100 u per ml, 3 ml syringe .....	51.19	5	NovoRapid FlexPen
<b>INSULIN GLULISINE</b>			
Inj 100 u per ml, 10 ml vial.....	27.03	1	Apidra
Inj 100 u per ml, 3 ml cartridge.....	46.07	5	Apidra
Inj 100 u per ml, 3 ml disposable pen.....	46.07	5	Apidra Solostar
<b>INSULIN LISPRO</b>			
Inj 100 u per ml, 10 ml vial			
Inj 100 u per ml, 3 ml cartridge			
<b>Insulin - Short-Acting Preparations</b>			
<b>INSULIN NEUTRAL</b>			
Inj human 100 u per ml, 10 ml vial			
Inj human 100 u per ml, 3 ml cartridge			
<b>Oral Hypoglycaemic Agents</b>			
<b>GLIBENCLAMIDE</b>			
Tab 5 mg – 1% DV Oct-18 to 2021.....	6.00	100	Daonil
<b>GLICLAZIDE</b>			
Tab 80 mg – 1% DV Nov-20 to 2023.....	15.18	500	Glizide
<b>GLIPIZIDE</b>			
Tab 5 mg – 1% DV Dec-18 to 2021 .....	3.27	100	Minidiab

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>METFORMIN HYDROCHLORIDE</b>			
Tab immediate-release 500 mg – <b>1% DV Feb-19 to 2021</b> .....	8.63	1,000	<b>Apotex</b>
Tab immediate-release 850 mg – <b>1% DV Feb-19 to 2021</b> .....	7.04	500	<b>Apotex</b>
<b>PIOGLITAZONE</b>			
Tab 15 mg – <b>1% DV Oct-18 to 2021</b> .....	3.47	90	<b>Vexazone</b>
Tab 30 mg – <b>1% DV Oct-18 to 2021</b> .....	5.06	90	<b>Vexazone</b>
Tab 45 mg – <b>1% DV Oct-18 to 2021</b> .....	7.10	90	<b>Vexazone</b>
<b>VILDAGLIPTIN</b>			
Tab 50 mg .....	40.00	60	Galvus
<b>VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE</b>			
Tab 50 mg with 1,000 mg metformin hydrochloride .....	40.00	60	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride .....	40.00	60	Galvumet

## Digestives Including Enzymes

### PANCREATIC ENZYME

Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase, 1,250 U protease))

Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U) – **1% DV Sep-18 to 2021** ..... 34.93 100 **Creon 10000**

Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U) – **1% DV Sep-18 to 2021** ..... 94.38 100 **Creon 25000**

Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph Eur U) ..... 34.93 20 g Creon Micro

Powder pancreatin 60.12 mg (3,600 Ph. Eur. u/amylase, 5,000 Ph. Eur. u/lipase and 200 Ph. Eur. u/protease)

### URSODEOXYCHOLIC ACID – **Restricted** see terms [below](#)

↓ Cap 250 mg – **1% DV Oct-20 to 2023** ..... 32.95 100 **Ursosan**

### → **Restricted (RS1647)**

#### **Initiation – Alagille syndrome or progressive familial intrahepatic cholestasis**

Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

#### **Initiation – Chronic severe drug induced cholestatic liver injury**

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

#### **Initiation – Primary biliary cholangitis**

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

#### **Initiation – Pregnancy**

Patient diagnosed with cholestasis of pregnancy.

#### **Initiation – Haematological transplant**

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to

continued...

## ALIMENTARY TRACT AND METABOLISM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

allogenic stem cell or bone marrow transplantation; and

2 Treatment for up to 13 weeks.

### Initiation – Total parenteral nutrition induced cholestasis

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by TPN; and
- 2 Liver function has not improved with modifying the TPN composition.

## Laxatives

### Bowel-Cleansing Preparations

#### CITRIC ACID WITH MAGNESIUM OXIDE AND SODIUM PICOSULFATE

Powder for oral soln 12 g with magnesium oxide 3.5 g and sodium picosulfate 10 mg per sachet

*e.g. PicoPrep*

#### MACROGOL 3350 WITH ASCORBIC ACID, POTASSIUM CHLORIDE AND SODIUM CHLORIDE

Powder for oral soln 755.68 mg with ascorbic acid 85.16 mg, potassium chloride 10.55 mg, sodium chloride 37.33 mg and sodium sulphate 80.62 mg per g, 210 g sachet

*e.g. Glycoprep-C*

Powder for oral soln 755.68 mg with ascorbic acid 85.16 mg, potassium chloride 10.55 mg, sodium chloride 37.33 mg and sodium sulphate 80.62 mg per g, 70 g sachet

*e.g. Glycoprep-C*

#### MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE, SODIUM CHLORIDE AND SODIUM SULPHATE

Powder for oral soln 59 g with potassium chloride 0.7425 g, sodium bicarbonate 1.685 g, sodium chloride 1.465 g and sodium sulphate 5.685 g per sachet – **1% DV Aug-19 to 2022** ..... 14.31

4

**Klean Prep**

### Bulk-Forming Agents

#### ISPAGHULA (PSYLLIUM) HUSK

Powder for oral soln – **1% DV Nov-20 to 2023** ..... 12.20

500 g

**Konsyl-D**

#### STERCULIA WITH FRANGULA – **Restricted:** For continuation only

➡ Powder for oral soln

### Faecal Softeners

#### DOCUSATE SODIUM

Tab 50 mg – **1% DV Oct-20 to 2023** ..... 2.31

100

**Coloxyl**

Tab 120 mg – **1% DV Oct-20 to 2023** ..... 3.13

100

**Coloxyl**

#### DOCUSATE SODIUM WITH SENNOSIDES

Tab 50 mg with sennosides 8 mg – **1% DV Jun-18 to 2021** ..... 3.10

200

**Laxsol**

#### PARAFFIN

Oral liquid 1 mg per ml

Enema 133 ml

#### POLOXAMER

Oral drops 10% – **1% DV Nov-20 to 2023** ..... 3.98

30 ml

**Coloxyl**

### Opioid Receptor Antagonists - Peripheral

#### METHYLNALTREXONE BROMIDE – **Restricted** see terms [on the next page](#)

⚡ Inj 12 mg per 0.6 ml vial ..... 36.00

246.00

1

Relistor

7

Relistor

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

### ➔ Restricted (RS1601)

#### Initiation – Opioid induced constipation

Both:

- 1 The patient is receiving palliative care; and
- 2 Either:
  - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
  - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

### Osmotic Laxatives

#### GLYCEROL

Suppos 1.27 g			
Suppos 2.55 g			
Suppos 3.6 g – 1% DV Oct-18 to 2021	9.25	20	<b>PSM</b>

#### LACTULOSE

Oral liq 10 g per 15 ml – 1% DV Nov-19 to 2022	3.33	500 ml	<b>Laevolac</b>
--	------	--------	-----------------

#### MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE AND SODIUM CHLORIDE

Powder for oral soln 6.563 g with potassium chloride 23.3 mg, sodium bicarbonate 89.3 mg and sodium chloride 175.4 mg			
Powder for oral soln 13.125 g with potassium chloride 46.6 mg, sodium bicarbonate 178.5 mg and sodium chloride 350.7 mg – 1% DV Oct-20 to 2023	6.70	30	<b>Molaxole</b>

#### SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE

Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml – 1% DV Nov-19 to 2022	29.98	50	<b>Micolette</b>
---	-------	----	------------------

#### SODIUM PHOSPHATE WITH PHOSPHORIC ACID

Oral liq 16.4% with phosphoric acid 25.14%			
Enema 10% with phosphoric acid 6.58%	2.50	1	<b>Fleet Phosphate Enema</b>

### Stimulant Laxatives

#### BISACODYL

Tab 5 mg – 1% DV Sep-18 to 2021	5.99	200	<b>Lax-Tabs</b>
Suppos 10 mg – 1% DV Sep-18 to 2021	3.74	10	<b>Lax-Suppositories</b>

#### SENNOSIDES

Tab 7.5 mg			
------------	--	--	--

### Metabolic Disorder Agents

#### ALGLUCOSIDASE ALFA – Restricted see terms [below](#)

⚡ Inj 50 mg vial	1,142.60	1	<b>Myozyme</b>
------------------	----------	---	----------------

### ➔ Restricted (RS1750)

#### Initiation

Metabolic physician

*Re-assessment required after 12 months*

All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2 Any of the following:

- 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
- 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
- 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
- 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and

- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

## Continuation

*Re-assessment required after 12 months*

All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

## ARGININE

Powder

Inj 600 mg per ml, 25 ml vial

BETAINE – **Restricted** see terms [below](#)

↓ Powder for oral soln.....575.00 180 g Cystadane

→ **Restricted (RS1751)**

## Initiation

Metabolic physician

*Re-assessment required after 12 months*

All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
  - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
  - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
  - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

## Continuation

*Re-assessment required after 12 months*

The treatment remains appropriate and the patient is benefiting from treatment.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

BIOTIN – **Restricted** see terms [below](#)

↓ Cap 50 mg

↓ Cap 100 mg

↓ Inj 10 mg per ml, 5 ml vial

→ **Restricted** (RS1330)

Metabolic physician or metabolic disorders dietitian

GALSULFASE – **Restricted** see terms [below](#)

↓ Inj 1 mg per ml, 5 ml vial.....2,234.00      1      Naglazyme

→ **Restricted** (RS1752)

#### Initiation

Metabolic physician

*Re-assessment required after 12 months*

Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency confirmed by either enzyme activity assay in leukocytes or skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI.

#### Continuation

*Re-assessment required after 12 months*

All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

HAEM ARGINATE

Inj 25 mg per ml, 10 ml ampoule

IDURSULFASE – **Restricted** see terms [below](#)

↓ Inj 2 mg per ml, 3 ml vial.....4,608.30      1      Elaprase

→ **Restricted** (RS1546)

#### Initiation

Metabolic physician

*Limited to 24 weeks treatment*

All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
  - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

LARONIDASE – **Restricted** see terms [below](#)

⚡ Inj 100 U per ml, 5 ml vial ..... 1,335.16 1 Aldurazyme

➡ **Restricted (RS1607)**

## Initiation

Metabolic physician

*Limited to 24 weeks treatment*

All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysaccharidosis I-H); and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE – **Restricted** see terms [below](#)

⚡ Cap 500 mg

⚡ Oral soln 1,000 mg per 10 ml

⚡ Oral soln 1,100 mg per 15 ml

⚡ Inj 200 mg per ml, 5 ml vial

➡ **Restricted (RS1035)**

Neurologist, metabolic physician or metabolic disorders dietitian

PYRIDOXAL-5-PHOSPHATE – **Restricted** see terms [below](#)

⚡ Tab 50 mg

➡ **Restricted (RS1331)**

Neurologist, metabolic physician or metabolic disorders dietitian

SAPROPTERIN DIHYDROCHLORIDE – **Restricted** see terms [below](#)

⚡ Tab soluble 100 mg ..... 1,452.70 30 Kuvan

➡ **Restricted (RS1753)**

## Initiation

Metabolic physician

*Re-assessment required after 1 month*

All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

## Continuation

*Re-assessment required after 12 months*

All of the following:

- 1 Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
- 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
  - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
  - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
  - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

#### SODIUM BENZOATE

Cap 500 mg  
Powder  
Soln 100 mg per ml  
Inj 20%, 10 ml ampoule

#### SODIUM PHENYLBUTYRATE – **Some items restricted** see terms [below](#)

Tab 500 mg			
↓ Grans 483 mg per g.....	1,920.00	174 g	Pheburane
Oral liq 250 mg per ml			
Inj 200 mg per ml, 10 ml ampoule			

➔ **Restricted (RS1754)**

#### Initiation

Metabolic physician

*Re-assessment required after 12 months*

For the chronic management of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

#### Continuation

*Re-assessment required after 12 months*

The treatment remains appropriate and the patient is benefiting from treatment.

#### TALIGLUCERASE ALFA – **Restricted** see terms [below](#)

↓ Inj 200 unit vial.....	1,072.00	1	Elelyso
--------------------------	----------	---	---------

➔ **Restricted (RS1034)**

#### Initiation

Only for use in patients with approval by the Gaucher Treatment Panel.

#### TRIENTINE DIHYDROCHLORIDE

Cap 300 mg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Minerals</b>			
<b>Calcium</b>			
CALCIUM CARBONATE			
Tab 1.25 g (500 mg elemental) – <b>1% DV May-21 to 2023</b> .....	7.52	250	Arrow-Calcium
	6.69		<b>Calci-Tab 500</b>
Tab eff 1.25 g (500 mg elemental)			
Tab eff 1.75 g (1 g elemental)			
<i>(Arrow-Calcium Tab 1.25 g (500 mg elemental) to be delisted 1 May 2021)</i>			
<b>Fluoride</b>			
SODIUM FLUORIDE			
Tab 1.1 mg (0.5 mg elemental)			
<b>Iodine</b>			
POTASSIUM IODATE			
Tab 253 mcg (150 mcg elemental iodine) – <b>1% DV Oct-20 to 2023</b> .....	4.58	90	NeuroTabs
POTASSIUM IODATE WITH IODINE			
Oral liq 10% with iodine 5%			
<b>Iron</b>			
FERRIC CARBOXYMALTOSE – <b>Restricted</b> see terms <a href="#">below</a>			
⚡ Inj 50 mg per ml, 10 ml vial.....	150.00	1	Ferinject
➡ <b>Restricted (RS1417)</b>			
<b>Initiation</b>			
Treatment with oral iron has proven ineffective or is clinically inappropriate.			
FERROUS FUMARATE			
Tab 200 mg (65 mg elemental) – <b>1% DV Jan-19 to 2021</b> .....	3.09	100	Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID			
Tab 310 mg (100 mg elemental) with folic acid 350 mcg – <b>1% DV</b>			
<b>Jun-18 to 2021</b> .....	4.68	60	Ferro-F-Tabs
FERROUS GLUCONATE WITH ASCORBIC ACID			
Tab 170 mg (20 mg elemental) with ascorbic acid 40 mg			
FERROUS SULFATE			
Oral liq 30 mg (6 mg elemental) per ml – <b>1% DV Nov-19 to 2022</b> .....	12.08	500 ml	Ferodan
FERROUS SULPHATE			
Tab long-acting 325 mg (105 mg elemental) – <b>1% DV Jun-18 to 2021</b> .....	2.06	30	Ferrograd
FERROUS SULPHATE WITH ASCORBIC ACID			
Tab long-acting 325 mg (105 mg elemental) with ascorbic acid 500 mg			
IRON POLYMALTOSE			
Inj 50 mg per ml, 2 ml ampoule .....	34.50	5	Ferrosig
IRON SUCROSE			
Inj 20 mg per ml, 5 ml ampoule .....	100.00	5	Venofer

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Magnesium</b>			
MAGNESIUM AMINO ACID CHELATE Cap 750 mg (150 mg elemental)			
MAGNESIUM CHLORIDE Inj 1 mmol per 1 ml, 100 ml bag			
MAGNESIUM HYDROXIDE Tab 311 mg (130 mg elemental)			
MAGNESIUM OXIDE Cap 663 mg (400 mg elemental) Cap 696 mg (420 mg elemental)			
MAGNESIUM OXIDE WITH MAGNESIUM ASPARTATE, MAGNESIUM AMINO ACID CHELATE AND MAGNESIUM CITRATE Cap 500 mg with magnesium aspartate 100 mg, magnesium amino acid chelate 100 mg and magnesium citrate 100 mg (360 mg elemental magnesium)			
MAGNESIUM SULPHATE Inj 0.4 mmol per ml, 250 ml bag Inj 2 mmol per ml, 5 ml ampoule ..... Inj 100 mg per ml, 50 ml bag	10.21	10	DBL

**Zinc**

ZINC Oral liq 5 mg per 5 drops			
ZINC CHLORIDE Inj 5.3 mg per ml (5.1 mg per ml elemental), 2 ml ampoule			
ZINC SULPHATE Cap 137.4 mg (50 mg elemental) – <b>1% DV Dec-19 to 2022</b> .....	11.00	100	<b>Zincaps</b>

**Mouth and Throat****Agents Used in Mouth Ulceration**

BENZYLAMINE HYDROCHLORIDE Soln 0.15% Spray 0.15% Spray 0.3%			
BENZYLAMINE HYDROCHLORIDE WITH CETYLPYRIDINIUM CHLORIDE Lozenge 3 mg with cetylpyridinium chloride			
CARBOXYMETHYLCELLULOSE Oral spray			
CARMELLOSE SODIUM WITH PECTIN AND GELATINE Paste Powder			
CHLORHEXIDINE GLUCONATE Mouthwash 0.2%			
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE Adhesive gel 8.7% with cetalkonium chloride 0.01%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## DICHLOROBENZYL ALCOHOL WITH AMYLMETACRESOL

Lozenge 1.2 mg with amylmetacresol 0.6 mg

## TRIAMCINOLONE ACETONIDE

Paste 0.1% – 1% DV Nov-20 to 2023.....5.33 5 g **Kenalog in Orabase**

### Oropharyngeal Anti-Infectives

#### AMPHOTERICIN B

Lozenge 10 mg.....5.86 20 Fungilin

#### MICONAZOLE

Oral gel 20 mg per g – 1% DV Sep-18 to 2021 .....4.74 40 g **Decozol**

#### NYSTATIN

Oral liquid 100,000 u per ml – 1% DV Oct-20 to 2023.....1.76 24 ml **Nilstat**

### Other Oral Agents

#### HYALURONIC ACID WITH LIDOCAINE [LIGNOCAINE]

Inj 20 mg per ml

#### SODIUM HYALURONATE [HYALURONIC ACID] – **Restricted** see terms [below](#)

⚠ Inj 20 mg per ml, 1 ml syringe

➡ **Restricted (RS1175)**

Otolaryngologist

#### THYMOL GLYCERIN

Compound, BPC.....9.15 500 ml **PSM**

### Vitamins

#### Multivitamin Preparations

#### MULTIVITAMIN AND MINERAL SUPPLEMENT – **Restricted** see terms [below](#)

⚠ Cap.....23.35 180 Clinicians Multivit & Mineral Boost

➡ **Restricted (RS1498)**

#### Initiation

Limited to 3 months treatment

Both:

- 1 Patient was admitted to hospital with burns; and
- 2 Any of the following:
  - 2.1 Burn size is greater than 15% of total body surface area (BSA) for all types of burns; or
  - 2.2 Burn size is greater than 10% of BSA for mid-dermal or deep dermal burns; or
  - 2.3 Nutritional status prior to admission or dietary intake is poor.

#### MULTIVITAMIN RENAL – **Restricted** see terms [below](#)

⚠ Cap.....6.49 30 Clinicians Renal Vit

➡ **Restricted (RS1499)**

#### Initiation

Either:

- 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or
- 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73m<sup>2</sup> body surface area (BSA).

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>MULTIVITAMINS</b>			
Tab (BPC cap strength) – <b>1% DV Mar-20 to 2022</b> .....	11.45	1,000	<b>Mvite</b>
↓ cap vitamin A 2500 u, betacarotene 3 mg, cholecalciferol 11 mcg, alpha tocopherol 150 u, phytomenadione 150 mcg, folic acid 0.2 mg, ascorbic acid 100 mg, thiamine 1.5 mg, pantothenic acid 12 mg, riboflavin 1.7 mg, niacin 20 mg, pyridoxine hydrochloride 1.9 mg, cyanocobalamin 3 mcg, zinc 7.5 mg and biotin 100 mcg			<i>e.g. Vitabdeck</i>
→ <b>Restricted (RS1620)</b>			
<b>Initiation</b>			
Any of the following:			
1 Patient has cystic fibrosis with pancreatic insufficiency; or			
2 Patient is an infant or child with liver disease or short gut syndrome; or			
3 Patient has severe malabsorption syndrome.			
↓ Powder vitamin A 4200 mcg with vitamin D 155.5 mcg, vitamin E 21.4 mg, vitamin C 400 mg, vitamin K1 166 mcg thiamine 3.2 mg, riboflavin 4.4 mg, niacin 35 mg, vitamin B6 3.4 mg, folic acid 303 mcg, vitamin B12 8.6 mcg, biotin 214 mcg, pantothenic acid 17 mg, choline 350 mg and inositol 700 mg			<i>e.g. Paediatric Seravit</i>
→ <b>Restricted (RS1178)</b>			
<b>Initiation</b>			
Patient has inborn errors of metabolism.			
Inj thiamine hydrochloride 250 mg with riboflavin 4 mg and pyridoxine hydrochloride 50 mg, 5 ml ampoule (1) and inj ascorbic acid 500 mg with nicotinamide 160 mg and glucose 1000 mg, 5 ml ampoule (1)			<i>e.g. Pabrinex IV</i>
Inj thiamine hydrochloride 250 mg with riboflavin 4 mg and pyridoxine hydrochloride 50 mg, 5 ml ampoule (1) and inj ascorbic acid 500 mg with nicotinamide 160 mg, 2 ml ampoule (1)			<i>e.g. Pabrinex IM</i>
Inj thiamine hydrochloride 500 mg with riboflavin 8 mg and pyridoxine hydrochloride 100 mg, 10 ml ampoule (1) and inj ascorbic acid 1000 mg with nicotinamide 320 mg and glucose 2000 mg, 10 ml ampoule (1)			<i>e.g. Pabrinex IV</i>
<b>Vitamin A</b>			
<b>RETINOL</b>			
Tab 10,000 iu			
Cap 25,000 iu			
Oral liq 150,000 iu per ml			
Oral liq 666.7 mcg per 2 drops, 10 ml			
Oral liq 5,000 iu per drop, 30 ml			
<b>Vitamin B</b>			
<b>HYDROXOCOBALAMIN</b>			
Inj 1 mg per ml, 1 ml ampoule – <b>1% DV Sep-18 to 2021</b> .....	1.89	3	<b>Neo-B12</b>
<b>PYRIDOXINE HYDROCHLORIDE</b>			
Tab 25 mg – <b>1% DV Oct-20 to 2023</b> .....	2.70	90	<b>Vitamin B6 25</b>
Tab 50 mg .....	13.63	500	<b>Apo-Pyridoxine</b>
Inj 100 mg per ml, 2 ml vial			
Inj 100 mg per ml, 1 ml ampoule			
Inj 100 mg per ml, 30 ml vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>THIAMINE HYDROCHLORIDE</b>			
Tab 50 mg .....	4.89	100	Max Health
Tab 100 mg .....			
Inj 100 mg per ml, 1 ml vial .....			<i>e.g. Benerva</i>
Inj 100 mg per ml, 2 ml vial .....			
<b>VITAMIN B COMPLEX</b>			
Tab strong, BPC.....	7.15	500	Bplex

## Vitamin C

<b>ASCORBIC ACID</b>			
Tab 100 mg – <b>1% DV Mar-20 to 2022</b> .....	9.90	500	<b>Cvite</b>
Tab chewable 250 mg .....			

## Vitamin D

<b>ALFACALCIDOL</b>			
Cap 0.25 mcg .....	26.32	100	One-Alpha
Cap 1 mcg .....	87.98	100	One-Alpha
Oral drops 2 mcg per ml .....	60.68	20 ml	One-Alpha
<b>CALCITRIOL</b>			
Cap 0.25 mcg – <b>1% DV Oct-19 to 2022</b> .....	7.95	100	<b>Calcitriol-AFT</b>
Cap 0.5 mcg – <b>1% DV Oct-19 to 2022</b> .....	13.75	100	<b>Calcitriol-AFT</b>
Oral liq 1 mcg per ml .....			
Inj 1 mcg per ml, 1 ml ampoule .....			
<b>COLECALCIFEROL</b>			
Cap 1.25 mg (50,000 iu) – <b>1% DV Feb-21 to 2023</b> .....	2.95	12	<b>Vit.D3</b>
Oral liq 188 mcg per ml (7,500 iu per ml) .....	9.00	4.8 ml	Puria

## Vitamin E

ALPHA TOCOPHERYL – **Restricted** see terms [below](#)

↓ Oral liq 156 u per ml

➔ **Restricted (RS1632)**

**Initiation – Cystic fibrosis**

Both:

1 Cystic fibrosis patient; and

2 Either:

- 2.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck); or
- 2.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for the patient.

**Initiation – Osteoradionecrosis**

For the treatment of osteoradionecrosis.

**Initiation – Other indications**

All of the following:

- 1 Infant or child with liver disease or short gut syndrome; and
- 2 Requires vitamin supplementation; and
- 3 Either:

- 3.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplements (Vitabdeck); or
- 3.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for patient.

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

ALPHA TOCOPHERYL ACETATE – **Restricted** see terms [below](#)

- ⚡ Cap 100 u
- ⚡ Cap 500 u
- ⚡ Oral liq 156 u per ml

➡ **Restricted** ([RS1176](#))

**Initiation – Cystic fibrosis**

- Both:
- 1 Cystic fibrosis patient; and
  - 2 Either:
    - 2.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck); or
    - 2.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for the patient.

**Initiation – Osteoradionecrosis**

For the treatment of osteoradionecrosis.

**Initiation – Other indications**

- All of the following:
- 1 Infant or child with liver disease or short gut syndrome; and
  - 2 Requires vitamin supplementation; and
  - 3 Either:
    - 3.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplements (Vitabdeck); or
    - 3.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for patient.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## Antianaemics

### Hypoplastic and Haemolytic

EPOETIN ALFA – **Restricted** see terms [below](#)

⚡ Inj 1,000 iu in 0.5 ml syringe – 1% DV Apr-19 to 2022.....	250.00	6	<b>Binocrit</b>
⚡ Inj 2,000 iu in 1 ml syringe – 1% DV Apr-19 to 2022.....	100.00	6	<b>Binocrit</b>
⚡ Inj 3,000 iu in 0.3 ml syringe – 1% DV Apr-19 to 2022.....	150.00	6	<b>Binocrit</b>
⚡ Inj 4,000 iu in 0.4 ml syringe – 1% DV Apr-19 to 2022.....	96.50	6	<b>Binocrit</b>
⚡ Inj 5,000 iu in 0.5 ml syringe – 1% DV Apr-19 to 2022.....	125.00	6	<b>Binocrit</b>
⚡ Inj 6,000 iu in 0.6 ml syringe – 1% DV Apr-19 to 2022.....	145.00	6	<b>Binocrit</b>
⚡ Inj 8,000 iu in 0.8 ml syringe – 1% DV Apr-19 to 2022.....	175.00	6	<b>Binocrit</b>
⚡ Inj 10,000 iu in 1 ml syringe – 1% DV Apr-19 to 2022.....	197.50	6	<b>Binocrit</b>
⚡ Inj 40,000 iu in 1 ml syringe – 1% DV Apr-19 to 2022.....	250.00	1	<b>Binocrit</b>

➡ **Restricted (RS1660)**

#### Initiation – chronic renal failure

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Either:
  - 3.1 Both:
    - 3.1.1 Patient does not have diabetes mellitus; and
    - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
  - 3.2 Both:
    - 3.2.1 Patient has diabetes mellitus; and
    - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; and
- 4 Patient is on haemodialysis or peritoneal dialysis.

#### Initiation – myelodysplasia\*

*Re-assessment required after 2 months*

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS); and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification-based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

#### Continuation – myelodysplasia\*

*Re-assessment required after 12 months*

All of the following:

- 1 The patient's transfusion requirement continues to be reduced with epoetin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

#### Initiation – all other indications

Haematologist

For use in patients where blood transfusion is not a viable treatment alternative.

Note: Indications marked with \* are unapproved indications

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

**EPOETIN BETA – Restricted** see terms [below](#)

Note: Epoetin beta is considered a Discretionary Variance Pharmaceutical for epoetin alfa.

- ↓ Inj 2,000 iu in 0.3 ml syringe
- ↓ Inj 3,000 iu in 0.3 ml syringe
- ↓ Inj 4,000 iu in 0.3 ml syringe
- ↓ Inj 5,000 iu in 0.3 ml syringe
- ↓ Inj 6,000 iu in 0.3 ml syringe
- ↓ Inj 10,000 iu in 0.6 ml syringe

→ **Restricted (RS1661)**

**Initiation – chronic renal failure**

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Either:
  - 3.1 Both:
    - 3.1.1 Patient does not have diabetes mellitus; and
    - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
  - 3.2 Both:
    - 3.2.1 Patient has diabetes mellitus; and
    - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; and
- 4 Patient is on haemodialysis or peritoneal dialysis.

**Initiation – myelodysplasia\***

*Re-assessment required after 12 months*

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS); and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification-based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

**Continuation – myelodysplasia\***

*Re-assessment required after 2 months*

All of the following:

- 1 The patient's transfusion requirement continues to be reduced with epoetin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

**Initiation – all other indications**

Haematologist.

For use in patients where blood transfusion is not a viable treatment alternative.

\*Note: Indications marked with \* are unapproved indications.

## Megaloblastic

**FOLIC ACID**

Tab 0.8 mg – <b>1% DV Oct-18 to 2021</b> .....	21.84	1,000	<b>Apo-Folic Acid</b>
Tab 5 mg – <b>1% DV Oct-18 to 2021</b> .....	12.12	500	<b>Apo-Folic Acid</b>
Oral liq 50 mcg per ml .....	26.00	25 ml	Biomed
Inj 5 mg per ml, 10 ml vial			

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

## Antifibrinolytics, Haemostatics and Local Sclerosants

ALUMINIUM CHLORIDE – **Restricted** see terms [below](#)

↓ Topical soln 20% w/v

*e.g. Driclor*

→ **Restricted** (RS1500)

### Initiation

For use as a haemostatis agent.

APROTININ – **Restricted** see terms [below](#)

↓ Inj 10,000 kIU per ml (equivalent to 200 mg per ml), 50 ml vial

→ **Restricted** (RS1332)

### Initiation

Cardiac anaesthetist

Either:

- 1 Paediatric patient undergoing cardiopulmonary bypass procedure; or
- 2 Adult patient undergoing cardiac surgical procedure where the significant risk of massive bleeding outweighs the potential adverse effects of the drug.

ELTROMBOPAG – **Restricted** see terms [below](#)

↓ Tab 25 mg ..... 1,550.00

28 Revolade

↓ Tab 50 mg ..... 3,100.00

28 Revolade

→ **Restricted** (RS1648)

### Initiation – idiopathic thrombocytopenic purpura - post-splenectomy

Haematologist

*Re-assessment required after 6 weeks*

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
  - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
  - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
  - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

### Initiation – idiopathic thrombocytopenic purpura - preparation for splenectomy

Haematologist

*Limited to 6 weeks treatment*

The patient requires eltrombopag treatment as preparation for splenectomy.

### Continuation – idiopathic thrombocytopenic purpura - post-splenectomy

Haematologist

*Re-assessment required after 12 months*

The patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre

### Initiation – idiopathic thrombocytopenic purpura contraindicated to splenectomy

Haematologist

*Re-assessment required after 3 months*

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
  - 3.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per microliter; or
  - 3.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

**Continuation – idiopathic thrombocytopenic purpura contraindicated to splenectomy**

Haematologist

*Re-assessment required after 12 months*

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

**Initiation – severe aplastic anaemia**

Haematologist

*Re-assessment required after 3 months*

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
  - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
  - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

**Continuation – severe aplastic anaemia**

Haematologist

*Re-assessment required after 12 months*

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

**EMICIZUMAB – Restricted see terms below**

↓ Inj 30 mg in 1 ml vial.....	3,570.00	1	Hemlibra
↓ Inj 60 mg in 0.4 ml vial.....	7,138.00	1	Hemlibra
↓ Inj 105 mg in 0.7 ml vial.....	12,492.00	1	Hemlibra
↓ Inj 150 mg in 1 ml vial.....	17,846.00	1	Hemlibra

➔ **Restricted (RS1780)**

**Initiation**

Haematologist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and
- 2 Either:
  - 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or

continued...

## BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more; and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:
  - 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
  - 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
- 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

### Continuation

Haematologist

*Re-assessment required after 6 months*

Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

### FERRIC SUBSULFATE

Gel 25.9%

Soln 500 ml

### POLIDOCANOL

Inj 0.5%, 30 ml vial

### SODIUM TETRADECYL SULPHATE

Inj 3%, 2 ml ampoule

### THROMBIN

Powder

### TRANEXAMIC ACID

Tab 500 mg – 1% DV May-20 to 2022.....9.45

Inj 100 mg per ml, 5 ml ampoule – 1% DV Sep-18 to 2021.....6.95

Inj 100 mg per ml, 10 ml ampoule – 1% DV Sep-18 to 2021.....10.95

60

5

5

**Mercury Pharma**

**Tranexamic-AFT**

**Tranexamic-AFT**

## Anticoagulant Reversal Agents

IDARUCIZUMAB – **Restricted** see terms [below](#)

⚡ Inj 50 mg per ml, 50 ml vial.....4,250.00

2

Praxbind

➡ **Restricted (RS1535)**

### Initiation

For the reversal of the anticoagulant effects of dabigatran when required in situations of life-threatening or uncontrolled bleeding, or for emergency surgery or urgent procedures.

## Blood Factors

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – **Restricted** see terms [on the next page](#)

⚡ Inj 250 iu vial.....612.50

1

Alprolix

⚡ Inj 500 iu vial.....1,225.00

1

Alprolix

⚡ Inj 1,000 iu vial.....2,450.00

1

Alprolix

⚡ Inj 2,000 iu vial.....4,900.00

1

Alprolix

⚡ Inj 3,000 iu vial.....7,350.00

1

Alprolix

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>→ Restricted (RS1684)</b>			
<b>Initiation</b>			
For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group			
EPTACOG ALFA [RECOMBINANT FACTOR VIIA] – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Inj 1 mg syringe .....	1,178.30	1	NovoSeven RT
↓ Inj 2 mg syringe .....	2,356.60	1	NovoSeven RT
↓ Inj 5 mg syringe .....	5,891.50	1	NovoSeven RT
↓ Inj 8 mg syringe .....	9,426.40	1	NovoSeven RT
<b>→ Restricted (RS1704)</b>			
<b>Initiation</b>			
For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria			
FACTOR EIGHT INHIBITOR BYPASSING FRACTION – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Inj 500 U .....	1,315.00	1	FEIBA NF
↓ Inj 1,000 U .....	2,630.00	1	FEIBA NF
↓ Inj 2,500 U .....	6,575.00	1	FEIBA NF
<b>→ Restricted (RS1705)</b>			
<b>Initiation</b>			
For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group			
MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Inj 250 iu prefilled syringe .....	287.50	1	Xyntha
↓ Inj 500 iu prefilled syringe .....	575.00	1	Xyntha
↓ Inj 1,000 iu prefilled syringe .....	1,150.00	1	Xyntha
↓ Inj 2,000 iu prefilled syringe .....	2,300.00	1	Xyntha
↓ Inj 3,000 iu prefilled syringe .....	3,450.00	1	Xyntha
<b>→ Restricted (RS1706)</b>			
<b>Initiation</b>			
For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria			
NONACOG GAMMA, [RECOMBINANT FACTOR IX] – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Inj 500 iu vial .....	435.00	1	RIXUBIS
↓ Inj 1,000 iu vial .....	870.00	1	RIXUBIS
↓ Inj 2,000 iu vial .....	1,740.00	1	RIXUBIS
↓ Inj 3,000 iu vial .....	2,610.00	1	RIXUBIS
<b>→ Restricted (RS1679)</b>			
<b>Initiation</b>			
For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group			
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) – <b>Restricted</b> see terms <a href="#">on the next page</a>			
↓ Inj 250 iu vial .....	210.00	1	Advate
↓ Inj 500 iu vial .....	420.00	1	Advate
↓ Inj 1,000 iu vial .....	840.00	1	Advate
↓ Inj 1,500 iu vial .....	1,260.00	1	Advate
↓ Inj 2,000 iu vial .....	1,680.00	1	Advate
↓ Inj 3,000 iu vial .....	2,520.00	1	Advate

## BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

### ➔ Restricted (RS1707)

#### Initiation

For patients with haemophilia. Preferred Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group

OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE FS) – **Restricted** see terms [below](#)

⚡ Inj 250 iu vial.....	237.50	1	Kogenate FS
⚡ Inj 500 iu vial.....	475.00	1	Kogenate FS
⚡ Inj 1,000 iu vial.....	950.00	1	Kogenate FS
⚡ Inj 2,000 iu vial.....	1,900.00	1	Kogenate FS
⚡ Inj 3,000 iu vial.....	2,850.00	1	Kogenate FS

### ➔ Restricted (RS1708)

#### Initiation

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria

RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII] – **Restricted** see terms [below](#)

⚡ Inj 250 iu vial.....	300.00	1	Adynovate
⚡ Inj 500 iu vial.....	600.00	1	Adynovate
⚡ Inj 1,000 iu vial.....	1,200.00	1	Adynovate
⚡ Inj 2,000 iu vial.....	2,400.00	1	Adynovate

### ➔ Restricted (RS1682)

#### Initiation

For patients with haemophilia A receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group

## Vitamin K

#### PHYTOMENADIONE

Inj 2 mg in 0.2 ml ampoule .....	8.00	5	Konakion MM
Inj 10 mg per ml, 1 ml ampoule .....	9.21	5	Konakion MM

## Antithrombotics

### Anticoagulants

BIVALIRUDIN – **Restricted** see terms [below](#)

⚡ Inj 250 mg vial

### ➔ Restricted (RS1181)

#### Initiation

Either:

- 1 For use in heparin-induced thrombocytopenia, heparin resistance or heparin intolerance; or
- 2 For use in patients undergoing endovascular procedures.

#### CITRATE SODIUM

Inj 4% (200 mg per 5 ml), 5 ml ampoule  
Inj 46.7% (1.4 g per 3 ml), 3 ml syringe  
Inj 46.7% (2.36 g per 5 ml), 5 ml ampoule

#### DABIGATRAN

Cap 75 mg.....	76.36	60	Pradaxa
Cap 110 mg.....	76.36	60	Pradaxa
Cap 150 mg.....	76.36	60	Pradaxa

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>DANAPAROID – Restricted</b> see terms <a href="#">below</a>			
↓ Inj 750 u in 0.6 ml ampoule			
➔ <b>Restricted (RS1182)</b>			
<b>Initiation</b>			
For use in heparin-induced thrombocytopenia, heparin resistance or heparin intolerance.			
<b>DEFIBROTIDE – Restricted</b> see terms <a href="#">below</a>			
↓ Inj 80 mg per ml, 2.5 ml ampoule			
➔ <b>Restricted (RS1183)</b>			
<b>Initiation</b>			
Haematologist			
Patient has moderate or severe sinusoidal obstruction syndrome as a result of chemotherapy or regimen-related toxicities.			
<b>DEXTROSE WITH SODIUM CITRATE AND CITRIC ACID [ACID CITRATE DEXTROSE A]</b>			
Inj 24.5 mg with sodium citrate 22 mg and citric acid 7.3 mg per ml, 100 ml bag			
<b>ENOXAPARIN SODIUM</b>			
Inj 20 mg in 0.2 ml syringe.....	27.93	10	Clexane
Inj 40 mg in 0.4 ml ampoule			
Inj 40 mg in 0.4 ml syringe.....	37.27	10	Clexane
Inj 60 mg in 0.6 ml syringe.....	56.18	10	Clexane
Inj 80 mg in 0.8 ml syringe.....	74.90	10	Clexane
Inj 100 mg in 1 ml syringe.....	93.80	10	Clexane
Inj 120 mg in 0.8 ml syringe.....	116.55	10	Clexane
			Clexane Forte
Inj 150 mg in 1 ml syringe.....	133.20	10	Clexane
			Clexane Forte
<i>(Clexane Inj 120 mg in 0.8 ml syringe to be delisted 1 January 2021)</i>			
<i>(Clexane Inj 150 mg in 1 ml syringe to be delisted 1 January 2021)</i>			
<b>FONDAPARINUX SODIUM – Restricted</b> see terms <a href="#">below</a>			
↓ Inj 2.5 mg in 0.5 ml syringe			
↓ Inj 7.5 mg in 0.6 ml syringe			
➔ <b>Restricted (RS1184)</b>			
<b>Initiation</b>			
For use in heparin-induced thrombocytopenia, heparin resistance or heparin intolerance.			
<b>HEPARIN SODIUM</b>			
Inj 100 iu per ml, 250 ml bag			
Inj 1,000 iu per ml, 1 ml ampoule .....	197.06	50	Hospira
Inj 1,000 iu per ml, 5 ml ampoule – <b>1% DV Nov-18 to 2021</b> .....	58.57	50	<b>Pfizer</b>
Inj 5,000 iu in 0.2 ml ampoule			
Inj 5,000 iu per ml, 1 ml ampoule .....	32.66	5	Hospira
Inj 5,000 iu per ml, 5 ml ampoule – <b>1% DV Nov-18 to 2021</b> .....	203.68	50	<b>Pfizer</b>
<b>HEPARINISED SALINE</b>			
Inj 10 iu per ml, 5 ml ampoule .....	65.48	50	Pfizer
Inj 100 iu per ml, 2 ml ampoule			
Inj 100 iu per ml, 5 ml ampoule			
<b>PHENINDIONE</b>			
Tab 10 mg			
Tab 25 mg			
Tab 50 mg			
<b>PROTAMINE SULPHATE</b>			
Inj 10 mg per ml, 5 ml ampoule			

## BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>RIVAROXABAN</b>			
Tab 10 mg .....	83.10	30	Xarelto
Tab 15 mg .....	77.56	28	Xarelto
Tab 20 mg .....	77.56	28	Xarelto
<b>SODIUM CITRATE WITH SODIUM CHLORIDE AND POTASSIUM CHLORIDE</b>			
Inj 4.2 mg with sodium chloride 5.7 mg and potassium chloride 74.6 mcg per ml, 5,000 ml bag			
<b>WARFARIN SODIUM</b>			
Tab 1 mg .....	6.46	100	Marevan
Tab 2 mg .....			
Tab 3 mg .....	10.03	100	Marevan
Tab 5 mg .....	11.48	100	Marevan
<b>Antiplatelets</b>			
<b>ASPIRIN</b>			
Tab 100 mg – <b>10% DV Nov-19 to 2022</b> .....	1.95	90	<b>Ethics Aspirin EC</b>
	10.80	990	<b>Ethics Aspirin EC</b>
Suppos 300 mg			
<b>CLOPIDOGREL</b>			
Tab 75 mg – <b>1% DV May-20 to 2022</b> .....	4.60	84	<b>Clopidogrel Multichem</b>
<b>DIPYRIDAMOLE</b>			
Tab 25 mg			
Tab long-acting 150 mg – <b>1% DV Oct-19 to 2022</b> .....	10.90	60	<b>Pytazen SR</b>
Inj 5 mg per ml, 2 ml ampoule			
<b>EPTIFIBATIDE</b> – <b>Restricted</b> see terms <a href="#">below</a>			
⚡ Inj 2 mg per ml, 10 ml vial – <b>1% DV Nov-18 to 2021</b> .....	138.75	1	<b>Integrilin</b>
⚡ Inj 750 mcg per ml, 100 ml vial – <b>1% DV Nov-18 to 2021</b> .....	405.00	1	<b>Integrilin</b>
➡ <b>Restricted (RS1759)</b>			
<b>Initiation</b>			
Any of the following:			
1 For use in patients with acute coronary syndromes undergoing percutaneous coronary intervention; or			
2 For use in patients with definite or strongly suspected intra-coronary thrombus on coronary angiography; or			
3 For use in patients undergoing intra-cranial intervention.			
<b>LYSINE ACETYSALICYLATE [LYSINE ASPIRIN]</b> – <b>Restricted</b> see terms <a href="#">below</a>			
⚡ Inj 500 mg			<i>e.g. Aspegic</i>
➡ <b>Restricted (RS1689)</b>			
<b>Initiation</b>			
Both:			
1 For use when an immediate antiplatelet effect is required prior to an urgent interventional neuro-radiology or interventional cardiology procedure; and			
2 Administration of oral aspirin would delay the procedure.			
<b>PRASUGREL</b> – <b>Restricted:</b> For continuation only			
➡ Tab 5 mg .....	108.00	28	Effient
➡ Tab 10 mg .....	120.00	28	Effient
(Effient Tab 5 mg to be delisted 1 February 2021)			
(Effient Tab 10 mg to be delisted 1 February 2021)			
<b>TICAGRELOR</b> – <b>Restricted</b> see terms <a href="#">on the next page</a>			
⚡ Tab 90 mg .....	90.00	56	Brilinta

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)

*e.g. Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

➔ **Restricted (RS1774)**

**Initiation**

Restricted to treatment of acute coronary syndromes specifically for patients who have recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome, and in whom fibrinolytic therapy has not been given in the last 24 hours and is not planned.

**Initiation – thrombosis prevention neurological stenting**

*Re-assessment required after 12 months*

- Both:
- 1 Either:
    - 1.1 Patient has had a neurological stenting procedure\* in the last 60 days; or
    - 1.2 Patient is about to have a neurological stenting procedure performed\*; and
  - 2 Either:
    - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
    - 2.2 Either:
      - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
      - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent..

**Continuation – thrombosis prevention neurological stenting**

*Re-assessment required after 12 months*

- Both:
- 1 Patient is continuing to benefit from treatment; and
  - 2 Treatment continues to be clinically appropriate.

**Initiation – Percutaneous coronary intervention with stent deployment**

*Limited to 12 months treatment*

- All of the following:
- 1 Patient has undergone percutaneous coronary intervention; and
  - 2 Patient has had a stent deployed in the previous 4 weeks; and
  - 3 Patient is clopidogrel-allergic\*\*.

**Initiation – Stent thrombosis**

Patient has experienced cardiac stent thrombosis whilst on clopidogrel.

**Initiation – Myocardial infarction**

*Limited to 1 week treatment*

For short term use while in hospital following ST-elevated myocardial infarction.

Notes: Indications marked with \* are unapproved indications.

Note: \*\* Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment

TICLOPIDINE

Tab 250 mg

**Fibrinolytic Agents**

ALTEPLASE

- Inj 2 mg vial
- Inj 10 mg vial
- Inj 50 mg vial

TENECTEPLASE

- Inj 50 mg vial

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>UROKINASE</b>			
Inj 5,000 iu vial			
Inj 10,000 iu vial			
Inj 50,000 iu vial			
Inj 100,000 iu vial			
Inj 500,000 iu vial			

## Colony-Stimulating Factors

### Drugs Used to Mobilise Stem Cells

**PLERIXAFOR – Restricted** see terms [below](#)

⚡ Inj 20 mg per ml, 1.2 ml vial.....8,740.00 1 Mozobil

➔ **Restricted (RS1536)**

**Initiation – Autologous stem cell transplant**

Haematologist

Limited to 3 days treatment

All of the following:

- 1 Patient is to undergo stem cell transplantation; and
- 2 Patient has not had a previous unsuccessful mobilisation attempt with plerixafor; and
- 3 Any of the following:
  - 3.1 Both:
    - 3.1.1 Patient is undergoing G-CSF mobilisation; and
    - 3.1.2 Either:
      - 3.1.2.1 Has a suboptimal peripheral blood CD34 count of less than or equal to  $10 \times 10^6/L$  on day 5 after 4 days of G-CSF treatment; or
      - 3.1.2.2 Efforts to collect  $> 1 \times 10^6$  CD34 cells/kg have failed after one apheresis procedure; or
  - 3.2 Both:
    - 3.2.1 Patient is undergoing chemotherapy and G-CSF mobilisation; and
    - 3.2.2 Any of the following:
      - 3.2.2.1 Both:
        - 3.2.2.1.1 Has rising white blood cell counts of  $> 5 \times 10^9/L$ ; and
        - 3.2.2.1.2 Has a suboptimal peripheral blood CD34 count of less than or equal to  $10 \times 10^6/L$ ; or
      - 3.2.2.2 Efforts to collect  $> 1 \times 10^6$  CD34 cells/kg have failed after one apheresis procedure; or
      - 3.2.2.3 The peripheral blood CD34 cell counts are decreasing before the target has been received; or
  - 3.3 A previous mobilisation attempt with G-CSF or G-CSF plus chemotherapy has failed.

### Granulocyte Colony-Stimulating Factors

**FILGRASTIM – Restricted** see terms [below](#)

⚡ Inj 300 mcg in 0.5 ml prefilled syringe – **1% DV May-19 to 2021** .....96.22 10 **Nivestim**

⚡ Inj 300 mcg in 1 ml vial.....520.00 4 **Neupogen**

⚡ Inj 480 mcg in 0.5 ml prefilled syringe – **1% DV Mar-19 to 2021** .....161.50 10 **Nivestim**

➔ **Restricted (RS1188)**

Haematologist or oncologist

**PEGFILGRASTIM – Restricted** see terms [below](#)

⚡ Inj 6 mg per 0.6 ml syringe .....1,080.00 1 Neulastim

➔ **Restricted (RS1743)**

**Initiation**

For prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...  
equal to 5%\*).

Note: \*Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines

## Fluids and Electrolytes

### Intravenous Administration

#### CALCIUM CHLORIDE

Inj 100 mg per ml, 10 ml vial

Inj 100 mg per ml, 50 ml syringe

*e.g. Baxter*

#### CALCIUM GLUCONATE

Inj 10%, 10 ml ampoule

*e.g. Max Health*

#### COMPOUND ELECTROLYTES

Inj sodium 140 mmol/l, potassium 5 mmol/l, magnesium 1.5 mmol/l,  
chloride 98 mmol/l, acetate 27 mmol/l, gluconate 23 mmol/l, 500 ml  
bag – **1% DV Jun-18 to 2021** .....

44.10

18

**Plasma-Lyte 148**

Inj sodium 140 mmol/l, potassium 5 mmol/l, magnesium 1.5 mmol/l,  
chloride 98 mmol/l, acetate 27 mmol/l, gluconate 23 mmol/l,  
1,000 ml bag – **1% DV Jun-18 to 2021** .....

27.24

12

**Plasma-Lyte 148**

#### COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]

Inj sodium 140 mmol/l, 5 mmol/l potassium, 1.5 mmol/l magnesium,  
98 mmol/l chloride, 27 mmol/l acetate and 23 mmol/l gluconate,  
glucose 23 mmol/l (5%), 1,000 ml bag – **1% DV Jun-18 to 2021** .....

211.92

12

**Plasma-Lyte 148 & 5%  
Glucose**

#### COMPOUND SODIUM LACTATE [HARTMANN'S SOLUTION]

Inj sodium 131 mmol/l with potassium 5 mmol/l, calcium 2 mmol/l,  
bicarbonate 29 mmol/l, chloride 111 mmol/l, 500 ml bag – **1% DV  
Jun-18 to 2021** .....

23.40

18

**Baxter**

Inj sodium 131 mmol/l with potassium 5 mmol/l, calcium 2 mmol/l,  
bicarbonate 29 mmol/l, chloride 111 mmol/l, 1,000 ml bag – **1% DV  
Jun-18 to 2021** .....

15.72

12

**Baxter**

#### GLUCOSE [DEXTROSE]

Inj 5%, 1,000 ml bag – **1% DV Aug-18 to 2021** .....

16.80

10

**Fresenius Kabi**

Inj 5%, 100 ml bag – **1% DV Aug-18 to 2021** .....

77.50

50

**Fresenius Kabi**

Inj 5%, 250 ml bag – **1% DV Aug-18 to 2021** .....

52.50

30

**Fresenius Kabi**

Inj 5%, 50 ml bag – **1% DV Jun-18 to 2021** .....

143.40

60

**Baxter Glucose 5%**

Inj 5%, 500 ml bag – **1% DV Aug-18 to 2021** .....

24.00

20

**Fresenius Kabi**

Inj 10%, 1,000 ml bag – **1% DV Jun-18 to 2021** .....

111.96

12

**Baxter Glucose 10%**

Inj 10%, 500 ml bag – **1% DV Jun-18 to 2021** .....

109.98

18

**Baxter Glucose 10%**

Inj 50%, 10 ml ampoule – **1% DV Nov-20 to 2023** .....

30.65

5

**Biomed**

Inj 50%, 500 ml bag – **1% DV Jun-18 to 2021** .....

337.32

18

**Baxter Glucose 50%**

Inj 50%, 90 ml bottle – **1% DV Nov-20 to 2023** .....

15.00

1

**Biomed**

#### GLUCOSE WITH POTASSIUM CHLORIDE

Inj 10% glucose with 20 mmol/l potassium chloride, 500 ml bag

## BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>GLUCOSE WITH POTASSIUM CHLORIDE AND SODIUM CHLORIDE</b>			
Inj 2.5% glucose with potassium chloride 20 mmol/l and sodium chloride 0.45%, 3,000 ml bag			
Inj 10% glucose with potassium chloride 10 mmol/l and sodium chloride 15 mmol/l, 500 ml bag			
Inj 4% glucose with potassium chloride 20 mmol/l and sodium chloride 0.18%, 1,000 ml bag – <b>1% DV Jun-18 to 2021</b>	203.40	12	<b>Baxter</b>
Inj 5% glucose with potassium chloride 20 mmol/l and sodium chloride 0.45%, 1,000 ml bag – <b>1% DV Jun-18 to 2021</b>	159.96	12	<b>Baxter</b>
Inj 5% glucose with potassium chloride 20 mmol/l and sodium chloride 0.9%, 1,000 ml bag – <b>1% DV Jun-18 to 2021</b>	282.72	12	<b>Baxter</b>
<b>GLUCOSE WITH SODIUM CHLORIDE</b>			
Inj glucose 2.5% with sodium chloride 0.45%, 500 ml bag			
Inj 4% glucose and sodium chloride 0.18%, 1,000 ml bag – <b>1% DV Jun-18 to 2021</b>	163.32	12	<b>Baxter</b>
Inj 5% glucose and sodium chloride 0.45%, 1,000 ml bag – <b>1% DV Jun-18 to 2021</b>	163.20	12	<b>Baxter</b>
Inj 5% glucose and sodium chloride 0.9%, 1,000 ml bag – <b>1% DV Jun-18 to 2021</b>	173.40	12	<b>Baxter</b>
<b>POTASSIUM CHLORIDE</b>			
Inj 75 mg (1 mmol) per ml, 10 ml ampoule			
Inj 225 mg (3 mmol) per ml, 20 ml ampoule			
<b>POTASSIUM CHLORIDE WITH SODIUM CHLORIDE</b>			
Inj 10 mmol potassium chloride with 0.29% sodium chloride, 100 ml bag – <b>1% DV Jun-18 to 2021</b>	476.64	48	<b>Baxter</b>
Inj 20 mmol potassium chloride with 0.9% sodium chloride, 1,000 ml bag – <b>1% DV Jun-18 to 2021</b>	163.08	12	<b>Baxter</b>
Inj 40 mmol potassium chloride with 0.9% sodium chloride, 1,000 ml bag – <b>1% DV Jun-18 to 2021</b>	253.32	12	<b>Baxter</b>
Inj 40 mmol potassium chloride with 0.9% sodium chloride, 100 ml bag – <b>1% DV Jun-18 to 2021</b>	772.32	48	<b>Baxter</b>
<b>POTASSIUM DIHYDROGEN PHOSPHATE</b>			
Inj 1 mmol per ml, 10 ml ampoule	151.80	10	Hospira
<b>RINGER'S SOLUTION</b>			
Inj sodium 147 mmol/l with potassium 4 mmol/l, calcium 2.2 mmol/l, chloride 156 mmol/l, 1,000 ml bag			
<b>SODIUM ACETATE</b>			
Inj 4 mmol per ml, 20 ml ampoule			
<b>SODIUM BICARBONATE</b>			
Inj 8.4%, 10 ml vial			
Inj 8.4%, 50 ml vial	19.95	1	Biomed
Inj 8.4%, 100 ml vial	20.50	1	Biomed

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>SODIUM CHLORIDE</b>			
Inj 0.9%, 5 ml ampoule – <b>1% DV Dec-19 to 2022</b> .....	2.80	20	<b>Fresenius Kabi</b>
Inj 0.9%, 10 ml ampoule – <b>1% DV Dec-19 to 2022</b> .....	5.40	50	<b>Fresenius Kabi</b>
↓ Inj 0.9%, 3 ml syringe, non-sterile pack – <b>1% DV Sep-18 to 2021</b> .....	160.90	480	<b>BD PosiFlush</b>
→ <b>Restricted (RS1297)</b>			
<b>Initiation</b>			
For use in flushing of in-situ vascular access devices only.			
↓ Inj 0.9%, 5 ml syringe, non-sterile pack – <b>1% DV Sep-18 to 2021</b> .....	162.91	480	<b>BD PosiFlush</b>
→ <b>Restricted (RS1297)</b>			
<b>Initiation</b>			
For use in flushing of in-situ vascular access devices only.			
↓ Inj 0.9%, 10 ml syringe, non-sterile pack – <b>1% DV Sep-18 to 2021</b> .....	170.35	480	<b>BD PosiFlush</b>
→ <b>Restricted (RS1297)</b>			
<b>Initiation</b>			
For use in flushing of in-situ vascular access devices only.			
Inj 0.9%, 20 ml ampoule – <b>1% DV Dec-19 to 2022</b> .....	5.00	20	<b>Fresenius Kabi</b>
Inj 23.4% (4 mmol/ml), 20 ml ampoule.....	33.00	5	Biomed
Inj 0.45%, 500 ml bag.....	71.28	18	Baxter
Inj 3%, 1,000 ml bag.....	91.20	12	Baxter
Inj 0.9%, 50 ml bag.....	109.80	60	Baxter
Inj 0.9%, 100 ml bag.....	78.24	48	Baxter
Inj 0.9%, 250 ml bag.....	44.64	24	Baxter
Inj 0.9%, 500 ml bag.....	22.14	18	Baxter
Inj 0.9%, 1,000 ml bag.....	15.12	12	Baxter
Inj 1.8%, 500 ml bottle			
<b>SODIUM DIHYDROGEN PHOSPHATE [SODIUM ACID PHOSPHATE]</b>			
Inj 1 mmol per ml, 20 ml ampoule – <b>1% DV Oct-18 to 2021</b> .....	48.70	5	<b>Biomed</b>
<b>WATER</b>			
Inj 5 ml ampoule .....	7.00	50	InterPharma
Inj 10 ml ampoule.....	6.63	50	Pfizer
Inj 20 ml ampoule .....	5.00	20	Fresenius Kabi
	7.50	30	InterPharma
	5.00	20	Multichem
Inj 250 ml bag			
Inj 500 ml bag			
Inj, 1,000 ml bag.....	19.08	12	Baxter
<i>(InterPharma Inj 5 ml ampoule to be delisted 1 June 2021)</i>			
<i>(InterPharma Inj 20 ml ampoule to be delisted 1 June 2021)</i>			
<b>Oral Administration</b>			
<b>CALCIUM POLYSTYRENE SULPHONATE</b>			
Powder .....	169.85	300 g	Calcium Resonium
<b>COMPOUND ELECTROLYTES</b>			
Powder for oral soln – <b>1% DV Apr-20 to 2022</b> .....	9.77	50	<b>Electral</b>
<b>COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]</b>			
Soln with electrolytes (2 x 500 ml) – <b>1% DV Nov-18 to 2021</b> .....	6.55	1,000 ml	<b>Pedialyte - Bubblegum</b>
<b>PHOSPHORUS</b>			
Tab eff 500 mg (16 mmol)			

## BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>POTASSIUM CHLORIDE</b>			
Tab eff 548 mg (14 mmol) with chloride 285 mg (8 mmol)			
Tab long-acting 600 mg (8 mmol) – <b>1% DV Oct-18 to 2021</b> .....	8.90	200	<b>Span-K</b>
Oral liq 2 mmol per ml			
<b>SODIUM BICARBONATE</b>			
Cap 840 mg .....	8.52	100	Sodibic
<b>SODIUM CHLORIDE</b>			
Tab 600 mg			
Oral liq 2 mmol/ml			
<b>SODIUM POLYSTYRENE SULPHONATE</b>			
Powder – <b>1% DV Sep-18 to 2021</b> .....	84.65	454 g	<b>Resonium A</b>
<b>Plasma Volume Expanders</b>			
<b>GELATINE, SUCCINYLATED</b>			
Inj 4%, 500 ml bag – <b>1% DV Jun-18 to 2021</b> .....	120.00	10	<b>Gelofusine</b>

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

## Agents Affecting the Renin-Angiotensin System

### ACE Inhibitors

#### CAPTOPRIL

↓ Oral liq 5 mg per ml .....	94.99	95 ml	Capoten
------------------------------	-------	-------	---------

→ **Restricted (RS1263)**

#### Initiation

Any of the following:

- 1 For use in children under 12 years of age; or
- 2 For use in tube-fed patients; or
- 3 For management of rebound transient hypertension following cardiac surgery.

#### CILAZAPRIL

Tab 0.5 mg – 1% DV Sep-19 to 2022 .....	2.09	90	<b>Zapril</b>
Tab 2.5 mg – 1% DV Feb-20 to 2022 .....	4.80	90	<b>Zapril</b>
Tab 5 mg – 1% DV Feb-20 to 2022 .....	8.35	90	<b>Zapril</b>

#### ENALAPRIL MALEATE

Tab 5 mg – 1% DV Jun-20 to 2022 .....	1.82	100	<b>Acetec</b>
Tab 10 mg – 1% DV Jun-20 to 2022 .....	2.02	100	<b>Acetec</b>
Tab 20 mg – 1% DV Jun-20 to 2022 .....	2.42	100	<b>Acetec</b>

#### LISINOPRIL

Tab 5 mg – 1% DV Dec-18 to 2021 .....	2.07	90	<b>Ethics Lisinopril</b>
Tab 10 mg – 1% DV Dec-18 to 2021 .....	2.36	90	<b>Ethics Lisinopril</b>
Tab 20 mg – 1% DV Dec-18 to 2021 .....	3.17	90	<b>Ethics Lisinopril</b>

#### PERINDOPRIL

Tab 2 mg .....	3.75	30	Apo-Perindopril
Tab 4 mg .....	4.80	30	Apo-Perindopril

#### QUINAPRIL

Tab 5 mg – 1% DV Nov-18 to 2021 .....	6.01	90	<b>Arrow-Quinapril 5</b>
Tab 10 mg – 1% DV Nov-18 to 2021 .....	3.16	90	<b>Arrow-Quinapril 10</b>
Tab 20 mg – 1% DV Nov-18 to 2021 .....	4.89	90	<b>Arrow-Quinapril 20</b>

### ACE Inhibitors with Diuretics

#### CILAZAPRIL WITH HYDROCHLOROTHIAZIDE – **Restricted:** For continuation only

→ Tab 5 mg with hydrochlorothiazide 12.5 mg .....	10.18	100	Apo-Cilazapril/ Hydrochlorothiazide
---	-------	-----	--

*(Apo-Cilazapril/ Hydrochlorothiazide Tab 5 mg with hydrochlorothiazide 12.5 mg to be delisted 1 May 2021)*

#### QUINAPRIL WITH HYDROCHLOROTHIAZIDE

Tab 10 mg with hydrochlorothiazide 12.5 mg – 1% DV Dec-18 to 2021 .....	3.83	30	<b>Accuretic 10</b>
Tab 20 mg with hydrochlorothiazide 12.5 mg – 1% DV Dec-18 to 2021 .....	4.92	30	<b>Accuretic 20</b>

### Angiotensin II Antagonists

#### CANDESARTAN CILEXETIL

Tab 4 mg – 1% DV Sep-18 to 2021 .....	1.90	90	<b>Candestar</b>
Tab 8 mg – 1% DV Sep-18 to 2021 .....	2.28	90	<b>Candestar</b>
Tab 16 mg – 1% DV Sep-18 to 2021 .....	3.67	90	<b>Candestar</b>
Tab 32 mg – 1% DV Sep-18 to 2021 .....	6.39	90	<b>Candestar</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>LOSARTAN POTASSIUM</b>			
Tab 12.5 mg – 1% DV Jan-21 to 2023 .....	1.56	84	<b>Losartan Actavis</b>
Tab 25 mg – 1% DV Jan-21 to 2023 .....	1.84	84	<b>Losartan Actavis</b>
Tab 50 mg – 1% DV Jan-21 to 2023 .....	2.25	84	<b>Losartan Actavis</b>
Tab 100 mg – 1% DV Jan-21 to 2023 .....	3.50	84	<b>Losartan Actavis</b>

## Angiotensin II Antagonists with Diuretics

<b>LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE</b>			
Tab 50 mg with hydrochlorothiazide 12.5 mg – 1% DV Jan-19 to 2021 .....	1.88	30	<b>Arrow-Losartan &amp; Hydrochlorothiazide</b>

## Angiotensin II Antagonists with Neprilysin Inhibitors

<b>SACUBITRIL WITH VALSARTAN – Restricted</b> see terms <a href="#">below</a>			
⚡ Tab 24.3 mg with valsartan 25.7 mg .....	190.00	56	Entresto 24/26
⚡ Tab 48.6 mg with valsartan 51.4 mg .....	190.00	56	Entresto 49/51
⚡ Tab 97.2 mg with valsartan 102.8 mg .....	190.00	56	Entresto 97/103

➡ **Restricted (RS1738)**

### Initiation

*Re-assessment required after 12 months*

All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
  - 2.1 Patient is in NYHA/WHO functional class II; or
  - 2.2 Patient is in NYHA/WHO functional class III; or
  - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
  - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
  - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

### Continuation

*Re-assessment required after 12 months*

The treatment remains appropriate and the patient is benefiting from treatment.

Note: Due to the angiotensin II receptor blocking activity of sacubitril with valsartan it should not be co-administered with an ACE inhibitor or another ARB.

## Alpha-Adrenoceptor Blockers

<b>DOXAZOSIN</b>			
Tab 2 mg .....	8.95	500	Apo-Doxazosin
Tab 4 mg .....	10.80	500	Apo-Doxazosin
<b>PHENOXYBENZAMINE HYDROCHLORIDE</b>			
Cap 10 mg			
Inj 50 mg per ml, 1 ml ampoule			
Inj 50 mg per ml, 2 ml ampoule			
<b>PHENTOLAMINE MESYLATE</b>			
Inj 5 mg per ml, 1 ml ampoule			
Inj 10 mg per ml, 1 ml ampoule			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PRazosin</b>			
Tab 1 mg .....	5.53	100	Apo-Prazosin
Tab 2 mg .....	7.00	100	Apo-Prazosin
Tab 5 mg .....	11.70	100	Apo-Prazosin
<b>TERazosin</b> – <b>Restricted:</b> For continuation only			
➔ Tab 1 mg .....			
➔ Tab 2 mg .....	7.50	500	Apo-Terazosin
➔ Tab 5 mg .....	10.90	500	Apo-Terazosin

## Antiarrhythmics

<b>ADENOSINE</b>			
Inj 3 mg per ml, 2 ml vial – <b>1% DV Feb-20 to 2022</b> .....	62.73	6	<b>Adenocor</b>
↓ Inj 3 mg per ml, 10 ml vial			
➔ <b>Restricted (RS1266)</b>			
<b>Initiation</b>			
For use in cardiac catheterisation, electrophysiology and MRI.			
<b>AJMALINE</b> – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Inj 5 mg per ml, 10 ml ampoule			
➔ <b>Restricted (RS1001)</b>			
Cardiologist			
<b>AMIODARONE HYDROCHLORIDE</b>			
Tab 100 mg – <b>1% DV Dec-19 to 2022</b> .....	3.80	30	<b>Aratac</b>
Tab 200 mg – <b>1% DV Dec-19 to 2022</b> .....	5.25	30	<b>Aratac</b>
Inj 50 mg per ml, 3 ml ampoule – <b>1% DV Feb-20 to 2022</b> .....	16.37	10	<b>Max Health</b>
<b>ATROPINE SULPHATE</b>			
Inj 600 mcg per ml, 1 ml ampoule – <b>1% DV Oct-18 to 2021</b> .....	12.07	10	<b>Martindale</b>
<b>DIGOXIN</b>			
Tab 62.5 mcg – <b>1% DV Nov-19 to 2022</b> .....	7.00	240	<b>Lanoxin PG</b>
Tab 250 mcg – <b>1% DV Nov-19 to 2022</b> .....	15.20	240	<b>Lanoxin</b>
Oral liq 50 mcg per ml			
Inj 250 mcg per ml, 2 ml vial			
<b>DISOPYRAMIDE PHOSPHATE</b>			
Cap 100 mg			
<b>FLECAINIDE ACETATE</b>			
Tab 50 mg – <b>1% DV Feb-20 to 2022</b> .....	19.95	60	<b>Flecainide BNM</b>
Cap long-acting 100 mg – <b>1% DV Dec-19 to 2022</b> .....	39.51	90	<b>Flecainide Controlled Release Teva</b>
Cap long-acting 200 mg – <b>1% DV Dec-19 to 2022</b> .....	61.06	90	<b>Flecainide Controlled Release Teva</b>
Inj 10 mg per ml, 15 ml ampoule .....	100.00	5	<b>Tambocor</b>
<b>IVABRADINE</b> – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Tab 5 mg			
➔ <b>Restricted (RS1566)</b>			
<b>Initiation</b>			
Both:			
1 Patient is indicated for computed tomography coronary angiography; and			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2 Either:

2.1 Patient has a heart rate of greater than 70 beats per minute while taking a maximally tolerated dose of beta blocker;  
or

2.2 Patient is unable to tolerate beta blockers.

## MEXILETINE HYDROCHLORIDE

Cap 150 mg .....	162.00	100	Mexiletine Hydrochloride USP
Cap 250 mg .....	202.00	100	Mexiletine Hydrochloride USP

## PROPAFENONE HYDROCHLORIDE

Tab 150 mg

## Antihypensives

MIDODRINE – **Restricted** see terms [below](#)

↓ Tab 2.5 mg

↓ Tab 5 mg

→ **Restricted** (RS1427)

### Initiation

Patient has disabling orthostatic hypotension not due to drugs.

## Beta-Adrenoceptor Blockers

### ATENOLOL

Tab 50 mg – 1% DV Sep-18 to 2021 .....	4.26	500	<b>Mylan Atenolol</b>
Tab 100 mg – 1% DV Sep-18 to 2021 .....	7.30	500	<b>Mylan Atenolol</b>
Oral liq 5 mg per ml .....	21.25	300 ml	Atenolol-AFT

### BISOPROLOL FUMARATE

Tab 2.5 mg – 1% DV Apr-21 to 2023 .....	1.84	90	<b>Bisoprolol Mylan</b>
	3.53		Bosvate
Tab 5 mg – 1% DV Apr-21 to 2023 .....	2.55	90	<b>Bisoprolol Mylan</b>
	5.15		Bosvate
Tab 10 mg – 1% DV Apr-21 to 2023 .....	3.62	90	<b>Bisoprolol Mylan</b>
	9.40		Bosvate

(Bosvate Tab 2.5 mg to be delisted 1 April 2021)

(Bosvate Tab 5 mg to be delisted 1 April 2021)

(Bosvate Tab 10 mg to be delisted 1 April 2021)

### CARVEDILOL

Tab 6.25 mg .....	2.24	60	Carvedilol Sandoz
Tab 12.5 mg .....	2.30	60	Carvedilol Sandoz
Tab 25 mg .....	2.95	60	Carvedilol Sandoz

CELIPROLOL – **Restricted:** For continuation only

→ Tab 200 mg ..... 21.40 | 180 | Celol |

(Celol Tab 200 mg to be delisted 1 April 2021)

### ESMOLOL HYDROCHLORIDE

Inj 10 mg per ml, 10 ml vial

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>LABETALOL</b>			
Tab 50 mg			
Tab 100 mg – 1% DV Sep-20 to 2024	14.50	100	<b>Trandate</b>
Tab 200 mg – 1% DV Sep-20 to 2024	27.00	100	<b>Trandate</b>
Inj 5 mg per ml, 20 ml ampoule			
<b>METOPROLOL SUCCINATE</b>			
Tab long-acting 23.75 mg	1.45	30	Betaloc CR
Tab long-acting 47.5 mg	1.43	30	Betaloc CR
Tab long-acting 95 mg	2.15	30	Betaloc CR
Tab long-acting 190 mg	4.27	30	Betaloc CR
<b>METOPROLOL TARTRATE</b>			
Tab 50 mg – 1% DV Oct-18 to 2021	5.66	100	<b>Apo-Metoprolol</b>
Tab 100 mg – 1% DV Oct-18 to 2021	7.55	60	<b>Apo-Metoprolol</b>
Tab long-acting 200 mg	23.40	28	Slow-Lopresor
Inj 1 mg per ml, 5 ml vial – 1% DV Feb-19 to 31 Jan 2022	29.50	5	<b>Metoprolol IV Mylan</b>
<b>NADOLOL</b>			
Tab 40 mg – 1% DV Oct-18 to 2021	16.69	100	<b>Apo-Nadolol</b>
Tab 80 mg – 1% DV Oct-18 to 2021	26.43	100	<b>Apo-Nadolol</b>
<b>PINDOLOL</b>			
Tab 5 mg – 1% DV Oct-18 to 2021	13.22	100	<b>Apo-Pindolol</b>
Tab 10 mg – 1% DV Oct-18 to 2021	23.12	100	<b>Apo-Pindolol</b>
Tab 15 mg – 1% DV Oct-18 to 2021	33.31	100	<b>Apo-Pindolol</b>
<b>PROPRANOLOL</b>			
Tab 10 mg – 1% DV Oct-18 to 2021	4.64	100	<b>Apo-Propranolol</b>
Tab 40 mg – 1% DV Oct-18 to 2021	5.72	100	<b>Apo-Propranolol</b>
Cap long-acting 160 mg	18.17	100	Cardinol LA
Oral liq 4 mg per ml			
Inj 1 mg per ml, 1 ml ampoule			
<b>SOTALOL</b>			
Tab 80 mg – 1% DV Oct-19 to 2022	32.58	500	<b>Mylan</b>
Tab 160 mg – 1% DV Oct-19 to 2022	10.98	100	<b>Mylan</b>
<b>TIMOLOL MALEATE</b>			
Tab 10 mg			

## Calcium Channel Blockers

### Dihydropyridine Calcium Channel Blockers

<b>AMLODIPINE</b>			
Tab 2.5 mg – 1% DV Jun-21 to 2023	1.72	100	Apo-Amlodipine
	1.08	90	<b>Vasorex</b>
Tab 5 mg – 1% DV Jun-21 to 2023	3.33	250	Apo-Amlodipine
	0.96	90	<b>Vasorex</b>
Tab 10 mg – 1% DV Jun-21 to 2023	4.40	250	Apo-Amlodipine
	1.19	90	<b>Vasorex</b>

(Apo-Amlodipine Tab 2.5 mg to be delisted 1 June 2021)

(Apo-Amlodipine Tab 5 mg to be delisted 1 June 2021)

(Apo-Amlodipine Tab 10 mg to be delisted 1 June 2021)

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>FELODIPINE</b>			
Tab long-acting 2.5 mg – 1% DV Sep-18 to 2021 .....	1.45	30	<b>Plendil ER</b>
Tab long-acting 5 mg – 1% DV Dec-18 to 2021 .....	3.93	90	<b>Felo 5 ER</b>
Tab long-acting 10 mg – 1% DV Dec-18 to 2021 .....	4.32	90	<b>Felo 10 ER</b>
<b>ISRADIPINE</b>			
Tab 2.5 mg			
Cap 2.5 mg			
<b>NICARDIPINE HYDROCHLORIDE – Restricted</b> see terms <a href="#">below</a>			
↓ Inj 2.5 mg per ml, 10 ml vial			
➔ <b>Restricted (RS1699)</b>			
<b>Initiation</b>			
Anaesthetist, intensivist, cardiologist or paediatric cardiologist			
Any of the following:			
1 Patient has hypertension requiring urgent treatment with an intravenous agent; or			
2 Patient has excessive ventricular afterload; or			
3 Patient is awaiting or undergoing cardiac surgery using cardiopulmonary bypass.			
<b>NIFEDIPINE</b>			
Tab long-acting 10 mg.....	10.63	60	Adalat 10
Tab long-acting 20 mg.....	17.72	100	Nyefax Retard
Tab long-acting 30 mg.....	3.14	30	Adalat Oros
Tab long-acting 60 mg.....	5.67	30	Adalat Oros
Cap 5 mg			
<b>NIMODIPINE</b>			
Tab 30 mg – 1% DV Jul-20 to 2022 .....	350.00	100	<b>Nimotop</b>
Inj 200 mcg per ml, 50 ml vial – 1% DV Jul-20 to 2022.....	67.50	1	<b>Nimotop</b>
<b>Other Calcium Channel Blockers</b>			
<b>DILTIAZEM HYDROCHLORIDE</b>			
Tab 30 mg .....	4.60	100	Dilzem
Tab 60 mg .....	8.50	100	Dilzem
Cap long-acting 120 mg – 1% DV Oct-18 to 2021.....	33.42	500	<b>Apo-Diltiazem CD</b>
Cap long-acting 180 mg – 1% DV Oct-18 to 2021.....	50.05	500	<b>Apo-Diltiazem CD</b>
Cap long-acting 240 mg – 1% DV Oct-18 to 2021.....	66.76	500	<b>Apo-Diltiazem CD</b>
Inj 5 mg per ml, 5 ml vial			
<b>PERHEXILINE MALEATE</b>			
Tab 100 mg – 1% DV Oct-19 to 2022.....	62.90	100	<b>Pexsig</b>
<b>VERAPAMIL HYDROCHLORIDE</b>			
Tab 40 mg .....	7.01	100	Isoptin
Tab 80 mg .....	11.74	100	Isoptin
Tab long-acting 120 mg.....	36.02	100	Isoptin SR
Tab long-acting 240 mg.....	15.12	30	Isoptin SR
Inj 2.5 mg per ml, 2 ml ampoule .....	25.00	5	Isoptin
<b>Centrally-Acting Agents</b>			
<b>CLONIDINE</b>			
Patch 2.5 mg, 100 mcg per day – 1% DV Nov-20 to 2023 .....	10.34	4	<b>Mylan</b>
Patch 5 mg, 200 mcg per day – 1% DV Nov-20 to 2023 .....	13.18	4	<b>Mylan</b>
Patch 7.5 mg, 300 mcg per day – 1% DV Nov-20 to 2023 .....	16.93	4	<b>Mylan</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>CLONIDINE HYDROCHLORIDE</b>			
Tab 25 mcg – <b>1% DV Oct-18 to 2021</b> .....	8.75	112	<b>Clonidine BNM</b>
Tab 150 mcg.....	34.32	100	Catapres
Inj 150 mcg per ml, 1 ml ampoule – <b>1% DV Oct-18 to 2021</b> .....	25.96	10	<b>Medsurge</b>
<b>METHYLDOPA</b>			
Tab 250 mg .....	15.10	100	Methyldopa Mylan
<b>Diuretics</b>			
<b>Loop Diuretics</b>			
<b>BUMETANIDE</b>			
Tab 1 mg .....	16.36	100	Burinex
Inj 500 mcg per ml, 4 ml vial			
<b>FUROSEMIDE [FRUSEMIDE]</b>			
Tab 40 mg – <b>1% DV Dec-19 to 2021</b> .....	7.24	1,000	<b>Apo-Furosemide</b>
Tab 500 mg – <b>1% DV Mar-19 to 2021</b> .....	25.00	50	<b>Urex Forte</b>
Oral liq 10 mg per ml – <b>1% DV Jan-20 to 2022</b> .....	11.20	30 ml	<b>Lasix</b>
Inj 10 mg per ml, 2 ml ampoule – <b>1% DV Oct-19 to 2022</b> .....	1.15	5	<b>Frusemide-Clarix</b>
			Furosemide-Baxter
Inj 10 mg per ml, 25 ml ampoule – <b>1% DV Jan-20 to 2022</b> .....	60.65	6	<b>Lasix</b>
<i>(Frusemide-Clarix Inj 10 mg per ml, 2 ml ampoule to be delisted 1 March 2021)</i>			
<b>Osmotic Diuretics</b>			
<b>MANNITOL</b>			
Inj 10%, 1,000 ml bag – <b>1% DV Jun-18 to 2021</b> .....	747.24	12	<b>Baxter</b>
Inj 20%, 500 ml bag – <b>1% DV Jun-18 to 2021</b> .....	1,096.92	18	<b>Baxter</b>
<b>Potassium Sparing Combination Diuretics</b>			
<b>AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE</b>			
Tab 5 mg with furosemide 40 mg			
<b>AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE</b>			
Tab 5 mg with hydrochlorothiazide 50 mg			
<b>Potassium Sparing Diuretics</b>			
<b>AMILORIDE HYDROCHLORIDE</b>			
Tab 5 mg			
Oral liq 1 mg per ml .....	30.00	25 ml	Biomed
<b>EPLERENONE – Restricted see terms below</b>			
↓ Tab 25 mg – <b>1% DV Sep-18 to 2021</b> .....	11.87	30	<b>Inspira</b>
↓ Tab 50 mg – <b>1% DV Dec-18 to 2021</b> .....	17.00	30	<b>Inspira</b>
<b>→ Restricted (RS1640)</b>			
<b>Initiation</b>			
Both:			
1 Patient has heart failure with ejection fraction less than 40%; and			
2 Either:			
2.1 Patient is intolerant to optimal dosing of spironolactone; or			
2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>SPIRONOLACTONE</b>			
Tab 25 mg .....	4.38	100	Spiractin
Tab 100 mg .....	11.80	100	Spiractin
Oral liq 50 mg per ml – 1% DV Nov-19 to 2022 .....	30.60	25 ml	Biomed

## Thiazide and Related Diuretics

<b>BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]</b>			
Tab 2.5 mg – 1% DV Dec-20 to 2023 .....	20.00	500	Arrow-Bendrofluazide
Tab 5 mg – 1% DV Dec-20 to 2023 .....	34.55	500	Arrow-Bendrofluazide
<b>CHLOROTHIAZIDE</b>			
Oral liq 50 mg per ml .....	26.00	25 ml	Biomed
<b>CHLORTALIDONE [CHLORTHALIDONE]</b>			
Tab 25 mg – 1% DV Dec-19 to 2022 .....	6.50	50	Hygroton
<b>INDAPAMIDE</b>			
Tab 2.5 mg – 1% DV Nov-20 to 2023 .....	10.45	90	Dapa-Tabs
<b>METOLAZONE</b>			
Tab 5 mg .....			

## Lipid-Modifying Agents

### Fibrates

<b>BEZAFIBRATE</b>			
Tab 200 mg – 1% DV Dec-18 to 2021 .....	19.01	90	Bezalip
Tab long-acting 400 mg – 1% DV Dec-18 to 2021 .....	12.89	30	Bezalip Retard
<b>GEMFIBROZIL – Restricted:</b> For continuation only			
➔ Tab 600 mg .....	19.56	60	Lipazil
(Lipazil Tab 600 mg to be delisted 1 January 2021)			

### HMG CoA Reductase Inhibitors (Statins)

<b>ATORVASTATIN</b>			
Tab 10 mg – 1% DV Sep-18 to 2021 .....	6.96	500	Lorstat
Tab 20 mg – 1% DV Sep-18 to 2021 .....	9.99	500	Lorstat
Tab 40 mg – 1% DV Sep-18 to 2021 .....	15.93	500	Lorstat
Tab 80 mg – 1% DV Sep-18 to 2021 .....	27.19	500	Lorstat
<b>PRAVASTATIN</b>			
Tab 10 mg .....			
Tab 20 mg – 1% DV Apr-21 to 2023 .....	4.72	100	Apo-Pravastatin
	2.11	28	Pravastatin Mylan
Tab 40 mg – 1% DV Apr-21 to 2023 .....	8.06	100	Apo-Pravastatin
	3.61	28	Pravastatin Mylan
(Apo-Pravastatin Tab 20 mg to be delisted 1 April 2021)			
(Apo-Pravastatin Tab 40 mg to be delisted 1 April 2021)			
<b>SIMVASTATIN</b>			
Tab 10 mg – 1% DV Nov-20 to 2023 .....	1.23	90	Simvastatin Mylan
Tab 20 mg – 1% DV Nov-20 to 2023 .....	2.03	90	Simvastatin Mylan
Tab 40 mg – 1% DV Nov-20 to 2023 .....	3.58	90	Simvastatin Mylan
Tab 80 mg – 1% DV Nov-20 to 2023 .....	7.12	90	Simvastatin Mylan

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Resins

CHOLESTYRAMINE  
Powder for oral liq 4 g

COLESTIPOL HYDROCHLORIDE  
Grans for oral liq 5 g

Selective Cholesterol Absorption Inhibitors

EZETIMIBE – **Restricted** see terms [below](#)

↓ Tab 10 mg – 1% DV Oct-20 to 2023.....	1.95	30	Ezetimibe Sandoz
---	------	----	------------------

➔ **Restricted** (RS1005)

**Initiation**

All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
  - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 x normal) when treated with one statin; or
  - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
  - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

EZETIMIBE WITH SIMVASTATIN – **Restricted** see terms [below](#)

↓ Tab 10 mg with simvastatin 10 mg.....	5.15	30	Zimybe
↓ Tab 10 mg with simvastatin 20 mg.....	6.15	30	Zimybe
↓ Tab 10 mg with simvastatin 40 mg.....	7.15	30	Zimybe
↓ Tab 10 mg with simvastatin 80 mg.....	8.15	30	Zimybe

➔ **Restricted** (RS1006)

**Initiation**

All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Other Lipid-Modifying Agents

ACIPIMOX  
Cap 250 mg

NICOTINIC ACID

Tab 50 mg .....	4.12	100	Apo-Nicotinic Acid
Tab 500 mg .....	17.89	100	Apo-Nicotinic Acid

(Apo-Nicotinic Acid Tab 50 mg to be delisted 1 May 2021)  
(Apo-Nicotinic Acid Tab 500 mg to be delisted 1 May 2021)

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## Nitrates

### GLYCERYL TRINITRATE

Inj 1 mg per ml, 5 ml ampoule			
Inj 1 mg per ml, 10 ml ampoule			
Inj 1 mg per ml, 50 ml vial			
Inj 5 mg per ml, 10 ml ampoule .....	100.00	5	Hospira
Oral pump spray, 400 mcg per dose .....	4.45	250 dose	Nitrolingual Pump Spray
Patch 25 mg, 5 mg per day .....	15.73	30	Nitroderm TTS 5
Patch 50 mg, 10 mg per day .....	18.62	30	Nitroderm TTS 10

### ISOSORBIDE MONONITRATE

Tab 20 mg – 1% DV Nov-20 to 2023 .....	19.55	100	<b>Ismo 20</b>
Tab long-acting 40 mg – 1% DV Nov-20 to 2023 .....	8.20	30	<b>Ismo 40 Retard</b>
Tab long-acting 60 mg – 1% DV Nov-20 to 2023 .....	9.25	90	<b>Duride</b>

## Other Cardiac Agents

LEVOSIMENDAN – **Restricted** see terms [below](#)

- ⚡ Inj 2.5 mg per ml, 5 ml vial
- ⚡ Inj 2.5 mg per ml, 10 ml vial
- ➡ **Restricted (RS1007)**

### Initiation – Heart transplant

Either:

- 1 For use as a bridge to heart transplant, in patients who have been accepted for transplant; or
- 2 For the treatment of heart failure following heart transplant.

### Initiation – Heart failure

Cardiologist or intensivist

For the treatment of severe acute decompensated heart failure that is non-responsive to dobutamine.

## Sympathomimetics

### ADRENALINE

Inj 1 in 1,000, 1 ml ampoule .....	4.98	5	Aspen Adrenaline
	10.76		DBL Adrenaline
Inj 1 in 1,000, 30 ml vial			
Inj 1 in 10,000, 10 ml ampoule .....	49.00	10	Aspen Adrenaline
	27.00	5	Hospira
Inj 1 in 10,000, 10 ml syringe			

### DOBUTAMINE

Inj 12.5 mg per ml, 20 ml ampoule – 1% DV Jan-19 to 2021 .....	61.13	5	<b>Dobutamine-hameln</b>
--	-------	---	--------------------------

### DOPAMINE HYDROCHLORIDE

Inj 40 mg per ml, 5 ml ampoule – 1% DV Sep-18 to 2021 .....	29.73	10	<b>Max Health Ltd</b>
---	-------	----	-----------------------

### EPHEDRINE

Inj 3 mg per ml, 10 ml syringe			
Inj 30 mg per ml, 1 ml ampoule – 1% DV Oct-20 to 2023 .....	30.63	10	<b>Max Health</b>

### ISOPRENALINE [ISOPROTERENOL]

Inj 200 mcg per ml, 1 ml ampoule			
Inj 200 mcg per ml, 5 ml ampoule			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>METARAMINOL</b>			
Inj 0.5 mg per ml, 10 ml syringe			
Inj 0.5 mg per ml, 20 ml syringe			
Inj 0.5 mg per ml, 5 ml syringe			
Inj 1 mg per ml, 1 ml ampoule			
Inj 1 mg per ml, 10 ml syringe			
Inj 10 mg per ml, 1 ml ampoule – 1% DV Jan-21 to 2023 .....	55.20	10	Torbay
<b>NORADRENALINE</b>			
Inj 0.06 mg per ml, 100 ml bag			
Inj 0.06 mg per ml, 50 ml syringe			
Inj 0.1 mg per ml, 100 ml bag			
Inj 0.1 mg per ml, 50 ml syringe			
Inj 0.12 mg per ml, 100 ml bag			
Inj 0.12 mg per ml, 50 ml syringe			
Inj 0.16 mg per ml, 50 ml syringe			
Inj 1 mg per ml, 100 ml bag			
Inj 1 mg per ml, 4 ml ampoule – 1% DV Oct-19 to 2022 .....	45.00	10	Noradrenaline BNM
<b>PHENYLEPHRINE HYDROCHLORIDE</b>			
Inj 10 mg per ml, 1 ml ampoule .....	142.07	25	Neosynephrine HCL
<b>Vasodilators</b>			
<b>ALPROSTADIL HYDROCHLORIDE</b>			
Inj 500 mcg per ml, 1 ml ampoule – 1% DV Dec-18 to 2021 .....	1,765.50	5	Prostin VR
<b>DIAZOXIDE</b>			
Inj 15 mg per ml, 20 ml ampoule			
<b>HYDRALAZINE HYDROCHLORIDE</b>			
↓ Tab 25 mg			
➔ <b>Restricted (RS1008)</b>			
<b>Initiation</b>			
Either:			
1 For the treatment of refractory hypertension; or			
2 For the treatment of heart failure, in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.			
Inj 20 mg ampoule .....	25.90	5	Apresoline
<b>MILRINONE</b>			
Inj 1 mg per ml, 10 ml ampoule – 1% DV Sep-18 to 2021 .....	99.00	10	Primacor
<b>MINOXIDIL</b>			
Tab 10 mg .....	70.00	100	Loniten
<b>NICORANDIL</b>			
Tab 10 mg – 1% DV Dec-19 to 2022 .....	25.57	60	Ikorel
Tab 20 mg – 1% DV Dec-19 to 2022 .....	32.28	60	Ikorel
<b>PAPAVERINE HYDROCHLORIDE</b>			
Inj 30 mg per ml, 1 ml vial			
Inj 12 mg per ml, 10 ml ampoule .....	217.90	5	Hospira
<b>PENTOXIFYLLINE [OXPENTIFYLLINE]</b>			
Tab 400 mg			
<b>SODIUM NITROPRUSSIDE</b>			
Inj 50 mg vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Endothelin Receptor Antagonists</b>			
AMBRISENTAN – <b>Restricted</b> see terms <a href="#">below</a>			
⚡ Tab 5 mg – 1% DV Mar-21 to 2023 .....	1,550.00	30	<b>Ambrisentan Mylan</b>
	4,585.00		Volibris
⚡ Tab 10 mg – 1% DV Mar-21 to 2023 .....	1,550.00	30	<b>Ambrisentan Mylan</b>
	4,585.00		Volibris

(Volibris Tab 5 mg to be delisted 1 March 2021)

(Volibris Tab 10 mg to be delisted 1 March 2021)

➡ **Restricted** ([RS1621](#))

#### Initiation

Either:

- 1 For use in patients with a valid Special Authority approval for ambrisentan by the Pulmonary Arterial Hypertension Panel; or
- 2 In-hospital stabilisations in emergency situations.

BOSENTAN – **Restricted** see terms [below](#)

⚡ Tab 62.5 mg – 1% DV Dec-18 to 2021 .....	141.00	60	<b>Bosentan Dr Reddy's</b>
⚡ Tab 125 mg – 1% DV Dec-18 to 2021 .....	141.00	60	<b>Bosentan Dr Reddy's</b>

➡ **Restricted** ([RS1622](#))

#### Initiation – Pulmonary arterial hypertension

*Re-assessment required after 6 months*

Either:

- 1 All of the following:
  - 1.1 Patient has pulmonary arterial hypertension (PAH); and
  - 1.2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
  - 1.3 PAH is at NYHA/WHO functional class II, III, or IV; and
  - 1.4 Any of the following:
    - 1.4.1 Both:
      - 1.4.1.1 Bosentan is to be used as PAH monotherapy; and
      - 1.4.1.2 Either:
        - 1.4.1.2.1 Patient is intolerant or contraindicated to sildenafil; or
        - 1.4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
    - 1.4.2 Both:
      - 1.4.2.1 Bosentan is to be used as PAH dual therapy; and
      - 1.4.2.2 Either:
        - 1.4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
        - 1.4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
    - 1.4.3 Both:
      - 1.4.3.1 Bosentan is to be used as PAH triple therapy; and
      - 1.4.3.2 Any of the following:
        - 1.4.3.2.1 Patient is on the lung transplant list; or
        - 1.4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
        - 1.4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
        - 1.4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2 In-hospital stabilisation in emergency situations.

**Continuation – Pulmonary arterial hypertension**

*Re-assessment required after 6 months*

Any of the following:

- 1 Both:
  - 1.1 Bosentan is to be used as PAH monotherapy; and
  - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
  - 2.1 Bosentan is to be used as PAH dual therapy; and
  - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
  - 3.1 Bosentan is to be used as PAH triple therapy; and
  - 3.2 Any of the following:
    - 3.2.1 Patient is on the lung transplant list; or
    - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
    - 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
    - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

**Phosphodiesterase Type 5 Inhibitors**

SILDENAFIL – **Restricted** see terms [below](#)

⬇ Tab 25 mg – <b>1% DV Sep-18 to 2021</b> .....	0.64	4	<b>Vedafil</b>
⬇ Tab 50 mg – <b>1% DV Sep-18 to 2021</b> .....	0.64	4	<b>Vedafil</b>
⬇ Tab 100 mg – <b>1% DV Sep-18 to 2021</b> .....	6.60	12	<b>Vedafil</b>
⬇ Inj 0.8 mg per ml, 12.5 ml vial			

➡ **Restricted (RS1740)**

**Initiation – tablets Raynaud’s Phenomenon**

All of the following:

- 1 Patient has Raynaud's phenomenon; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (proper body insulation, avoidance of cold exposure, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient has persisting severe symptoms despite treatment with calcium channel blockers and nitrates (unless contraindicated or not tolerated).

**Initiation – tablets Pulmonary arterial hypertension**

Any of the following:

- 1 All of the following:
  - 1.1 Patient has pulmonary arterial hypertension (PAH); and
  - 1.2 Any of the following:
    - 1.2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
    - 1.2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
    - 1.2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
  - 1.3 Any of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.3.1 PAH is in NYHA/WHO functional class II; or
- 1.3.2 PAH is in NYHA/WHO functional class III; or
- 1.3.3 PAH is in NYHA/WHO functional class IV; and
- 1.4 Either:
  - 1.4.1 All of the following:
    - 1.4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 1.4.1.2 Either:
      - 1.4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
      - 1.4.1.2.2 Patient is peri Fontan repair; and
    - 1.4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm<sup>-5</sup>); or
  - 1.4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age, or health system capacity constraints; or
- 2 For use in neonatal units for persistent pulmonary hypertension of the newborn (PPHN); or
- 3 In-hospital stabilisation in emergency situations.

## Initiation – tablets other conditions

Any of the following:

- 1 For use in weaning patients from inhaled nitric oxide; or
- 2 For perioperative use in cardiac surgery patients; or
- 3 For use in intensive care as an alternative to nitric oxide; or
- 4 For use in the treatment of erectile dysfunction secondary to spinal cord injury in patients being treated in a spinal unit.

## Initiation – injection

Both:

- 1 For use in the treatment of pulmonary hypertension in infants or children being treated in paediatric intensive care units and neonatal intensive care units when the enteral route is not accessible; and
- 2 Any of the following:
  - 2.1 For perioperative use following cardiac surgery; or
  - 2.2 For use in persistent pulmonary hypertension of the newborn (PPHN); or
  - 2.3 For use in congenital diaphragmatic hernia.

## Prostacyclin Analogues

EPOPROSTENOL – **Restricted** see terms [below](#)

⚡ Inj 500 mcg vial.....	36.61	1	Veletri
⚡ Inj 1.5 mg vial .....	73.21	1	Veletri

➡ **Restricted** (RS1624)

### Initiation

Either:

- 1 For use in patients with a valid Special Authority approval for epoprostenol by the Pulmonary Arterial Hypertension Panel; or
- 2 In-hospital stabilisation in emergency situations.

### ILOPROST

Inj 50 mcg in 0.5 ml ampoule – <b>1% DV Jan-20 to 2022</b> .....	305.00	5	Clinect
⚡ Nebuliser soln 10 mcg per ml, 2 ml – <b>1% DV Jan-20 to 2022</b> .....	740.10	30	Ventavis

➡ **Restricted** (RS1625)

### Initiation

Any of the following:

continued...

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

continued...

- 1 For use in patients with a valid Special Authority approval for iloprost by the Pulmonary Arterial Hypertension Panel; or
- 2 For diagnostic use in catheter laboratories; or
- 3 For use following mitral or tricuspid valve surgery; or
- 4 In-hospital stabilisation in emergency situations.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Anti-Infective Preparations</b>			
<b>Antibacterials</b>			
HYDROGEN PEROXIDE			
Crm 1%.....	8.56	15 g	Crystaderm
Soln 3% (10 vol)			
MAFENIDE ACETATE – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Powder 50 g sachet			
➔ <b>Restricted</b> ( <a href="#">RS1299</a> )			
<b>Initiation</b>			
For the treatment of burns patients.			
MUPIROCIN			
Oint 2%			
SODIUM FUSIDATE [FUSIDIC ACID]			
Crm 2% – <b>1% DV May-19 to 2021</b> .....	1.59	5 g	<b>Foban</b>
Oint 2% – <b>1% DV May-19 to 2021</b> .....	1.59	5 g	<b>Foban</b>
SULFADIAZINE SILVER			
Crm 1%.....	10.80	50 g	Flamazine
<b>Antifungals</b>			
AMOROLFINE			
Nail soln 5% – <b>1% DV Oct-20 to 2023</b> .....	14.93	5 ml	<b>MycoNail</b>
CICLOPIROX OLAMINE			
Nail soln 8% – <b>1% DV Sep-18 to 2021</b> .....	5.72	7 ml	<b>Apo-Ciclopirox</b>
➔ Soln 1% – <b>Restricted:</b> For continuation only			
CLOTRIMAZOLE			
Crm 1%.....	0.70	20 g	Clomazol
➔ Soln 1% – <b>Restricted:</b> For continuation only			
ECONAZOLE NITRATE			
➔ Crm 1% – <b>Restricted:</b> For continuation only			
Foaming soln 1%			
KETOCONAZOLE			
Shampoo 2% – <b>1% DV Nov-20 to 2023</b> .....	3.23	100 ml	<b>Sebizole</b>
METRONIDAZOLE			
Gel 0.75%			
MICONAZOLE NITRATE			
Crm 2% – <b>1% DV Feb-21 to 2023</b> .....	0.81	15 g	<b>Multichem</b>
➔ Lotn 2% – <b>Restricted:</b> For continuation only			
Tinc 2%			
NYSTATIN			
Crm 100,000 u per g			
<b>Antiparasitics</b>			
DIMETHICONE			
Lotn 4% – <b>1% DV Oct-19 to 2022</b> .....	4.98	200 ml	<b>healthE Dimethicone 4% Lotion</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MALATHION [MALDISON] Lotn 0.5% Shampoo 1%			
PERMETHRIN Crm 5% – 1% DV Nov-20 to 2023 .....	5.75	30 g	Lyderm
Lotn 5% – 1% DV Nov-20 to 2023 .....	3.99	30 ml	A-Scabies
PHENOTHRLIN Shampoo 0.5%			

### Antiacne Preparations

ADAPALENE Crm 0.1% Gel 0.1%			
BENZOYL PEROXIDE Soln 5%			
ISOTRETINOIN Cap 5 mg – 1% DV Oct-18 to 2021 .....	8.14	60	Oratane
Cap 10 mg – 1% DV Oct-18 to 2021 .....	13.34	120	Oratane
Cap 20 mg – 1% DV Oct-18 to 2021 .....	20.49	120	Oratane
TRETINOIN Crm 0.05% – 1% DV Jun-18 to 2021 .....	13.90	50 g	ReTrieve

### Antipruritic Preparations

CALAMINE Crm, aqueous, BP – 1% DV Nov-18 to 2021 .....	1.26	100 g	healthE Calamine Aqueous Cream BP
CROTAMITON Crm 10% – 1% DV Sep-18 to 2021 .....	3.29	20 g	Itch-Soothe

### Barrier Creams and Emollients

#### Barrier Creams

DIMETHICONE Crm 5% tube – 1% DV Oct-19 to 2022 .....	1.53	100 g	healthE Dimethicone 5%
Crm 5% pump bottle .....	4.48	500 ml	healthE Dimethicone 5%
Crm 10% pump bottle – 1% DV Sep-18 to 2021 .....	4.52	500 ml	healthE Dimethicone 10%
ZINC Crm			e.g. Zinc Cream (Orion-) ;Zinc Cream (PSM)
Oint Paste			e.g. Zinc oxide (PSM)

## DERMATOLOGICALS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>ZINC AND CASTOR OIL</b>			
Crm.....	1.63	20 g	Orion
Oint.....	4.25	500 g	Boucher
Note: DV limit applies to the pack sizes of greater than 30 g.			
Oint, BP.....	1.26	20 g	healthE
Note: DV limit applies to the pack sizes of 30 g or less.			
<b>ZINC WITH WOOL FAT</b>			
Crm zinc 15.25% with wool fat 4%			<i>e.g. Sudocrem</i>
<b>Emollients</b>			
<b>AQUEOUS CREAM</b>			
Crm 100 g – <b>1% DV Oct-18 to 2021</b> .....	1.05	100 g	<b>Pharmacy Health SLS-free</b>
Note: DV limit applies to the pack sizes of 100 g or less.			
Crm 500 g – <b>1% DV Dec-18 to 2021</b> .....	1.92	500 g	<b>Boucher</b>
Note: DV limit applies to the pack sizes of greater than 100 g.			
<b>CETOMACROGOL</b>			
Crm BP, 500 g – <b>1% DV Sep-18 to 2021</b> .....	2.48	500 g	<b>healthE</b>
Crm BP, 100 g – <b>1% DV Sep-18 to 2021</b> .....	1.42	1	<b>healthE</b>
<b>CETOMACROGOL WITH GLYCEROL</b>			
Crm 90% with glycerol 10%, – <b>1% DV Dec-19 to 2022</b> .....	1.65	100 g	<b>healthE</b>
Note: DV limit applies to the pack sizes of 100 g or less.			
Crm 90% with glycerol 10% – <b>1% DV Mar-20 to 2022</b> .....	2.35	500 ml	ADE
	3.10	1,000 ml	ADE
	2.35	500 ml	<b>Boucher</b>
	3.10	1,000 ml	<b>Boucher</b>
Note: DV limit applies to the pack sizes of greater than 100 g.			
<b>EMULSIFYING OINTMENT</b>			
Oint BP – <b>1% DV Oct-20 to 2023</b> .....	1.84	100 g	<b>Jaychem</b>
Note: DV limit applies to pack sizes of less than 200 g.			
Oint BP, 500 g – <b>1% DV Mar-21 to 2023</b> .....	3.59	500 g	AFT
	3.40		<b>Emulsifying Ointment ADE</b>
Note: DV limit applies to pack sizes of greater than 200 g.			
<i>(AFT Oint BP, 500 g to be delisted 1 March 2021)</i>			
<b>GLYCEROL WITH PARAFFIN</b>			
Crm glycerol 10% with white soft paraffin 5% and liquid paraffin 10%			<i>e.g. QV cream</i>
<b>OIL IN WATER EMULSION</b>			
Crm, 500 g – <b>1% DV Jan-19 to 2021</b> .....	2.19	500 g	<b>O/W Fatty Emulsion Cream</b>
Note: DV limit applies to the pack sizes of greater than 100 g.			
Crm, 100 g – <b>1% DV Dec-18 to 2021</b> .....	1.44	1	<b>healthE Fatty Cream</b>
<b>PARAFFIN</b>			
Oint liquid paraffin 50% with white soft paraffin 50% – <b>1% DV Jan-19 to 2021</b> .....	1.97	100 g	<b>healthE</b>
Note: DV limit applies to the pack sizes of 100 g or greater.			
White soft – <b>1% DV Sep-18 to 2021</b> .....	0.79	10 g	<b>healthE</b>
Note: DV limit applies to pack sizes of 30 g or less, and to both white soft paraffin and yellow soft paraffin.			
White soft, – <b>1% DV Apr-20 to 2022</b> .....	4.99	450 g	<b>healthE</b>
Yellow soft			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PARAFFIN WITH WOOL FAT</b>			
Lotn liquid paraffin 15.9% with wool fat 0.6%			<i>e.g. AlphaKeri;BK ;DP; Hydroderm Lotn</i>
Lotn liquid paraffin 91.7% with wool fat 3%			<i>e.g. Alpha Keri Bath Oil</i>
<b>UREA</b>			
Crm 10%.....	1.37	100 g	healthE Urea Cream
<b>WOOL FAT</b>			
Crm			
<b>Corticosteroids</b>			
<b>BETAMETHASONE DIPROPIONATE</b>			
Crm 0.05% – <b>1% DV Feb-21 to 2023</b> .....	36.00	50 g	<b>Diprosone</b>
Note: DV limit applies to the pack sizes of greater than 30 g.			
Oint 0.05% – <b>1% DV Feb-21 to 2023</b> .....	36.00	50 g	<b>Diprosone</b>
Note: DV limit applies to the pack sizes of greater than 30 g.			
<b>BETAMETHASONE VALERATE</b>			
Crm 0.1% – <b>1% DV Oct-18 to 2021</b> .....	3.45	50 g	<b>Beta Cream</b>
Oint 0.1% – <b>1% DV Oct-18 to 2021</b> .....	3.45	50 g	<b>Beta Ointment</b>
Lotn 0.1% – <b>1% DV Dec-18 to 2021</b> .....	18.00	50 ml	<b>Betnovate</b>
<b>CLOBETASOL PROPIONATE</b>			
Crm 0.05% – <b>1% DV Nov-19 to 2022</b> .....	2.18	30 g	<b>Dermol</b>
Oint 0.05% – <b>1% DV Nov-19 to 2022</b> .....	2.12	30 g	<b>Dermol</b>
<b>CLOBETASONE BUTYRATE</b>			
Crm 0.05%			
<b>DIFLUCORTOLONE VALERATE – Restricted:</b> For continuation only			
➔ Crm 0.1%			
➔ Fatty oint 0.1%			
<b>HYDROCORTISONE</b>			
Crm 1%, 100 g – <b>1% DV Sep-20 to 2022</b> .....	3.70	100 g	<b>Hydrocortisone (PSM)</b>
Note: DV limit applies to the pack sizes of less than or equal to 100 g.			
Crm 1%, 500 g – <b>1% DV Dec-20 to 2023</b> .....	17.15	500 g	<b>Hydrocortisone (PSM)</b>
<b>HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN</b>			
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – <b>1% DV Oct-20 to 2023</b> .....	10.57	250 ml	<b>DP Lotn HC</b>
<b>HYDROCORTISONE BUTYRATE</b>			
Crm 0.1%.....	6.85	100 g	Locoid Lipocream
Oint 0.1% – <b>1% DV Mar-19 to 2021</b> .....	13.70	100 g	<b>Locoid</b>
Milky emul 0.1% – <b>1% DV Mar-19 to 2021</b> .....	13.70	100 ml	<b>Locoid Crelo</b>
<b>METHYLPREDNISOLONE ACEPONATE</b>			
Crm 0.1% – <b>1% DV Dec-20 to 2023</b> .....	4.46	15 g	<b>Advantan</b>
Oint 0.1% – <b>1% DV Dec-20 to 2023</b> .....	4.46	15 g	<b>Advantan</b>
<b>MOMETASONE FUROATE</b>			
Crm 0.1% – <b>1% DV Nov-18 to 2021</b> .....	1.51	15 g	<b>Elocon Alcohol Free</b>
	2.50	50 g	<b>Elocon Alcohol Free</b>
Oint 0.1% – <b>1% DV Nov-18 to 2021</b> .....	1.51	15 g	<b>Elocon</b>
	2.90	50 g	<b>Elocon</b>
Lotn 0.1% – <b>1% DV Nov-18 to 2021</b> .....	6.30	30 ml	<b>Elocon</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>TRIAMCINOLONE ACETONIDE</b>			
Crm 0.02% – 1% DV Nov-20 to 2023 .....	6.30	100 g	<b>Aristocort</b>
Oint 0.02% – 1% DV Nov-20 to 2023 .....	6.35	100 g	<b>Aristocort</b>

### Corticosteroids with Anti-Infective Agents

BETAMETHASONE VALERATE WITH CLIOQUINOL – **Restricted** see terms [below](#)

↓ Crm 0.1% with clioquinol 3%

→ **Restricted (RS1125)**

#### Initiation

Either:

- 1 For the treatment of intertrigo; or
- 2 For continuation use.

BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUSIDIC ACID]

Crm 0.1% with sodium fusidate (fusidic acid) 2%

HYDROCORTISONE WITH MICONAZOLE

Crm 1% with miconazole nitrate 2% – 1% DV Sep-18 to 2021 ..... 2.00 15 g **Micreme H**

HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN

Crm 1% with natamycin 1% and neomycin sulphate 0.5% ..... 3.35 15 g Pimafucort

Oint 1% with natamycin 1% and neomycin sulphate 0.5% ..... 3.35 15 g Pimafucort

TRIAMCINOLONE ACETONIDE WITH NEOMYCIN SULPHATE, GRAMICIDIN AND NYSTATIN

Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and  
gramicidin 250 mcg per g

### Psoriasis and Eczema Preparations

ACITRETIN

Cap 10 mg – 1% DV Oct-20 to 2023 ..... 17.86 60 **Novatrelin**

Cap 25 mg – 1% DV Oct-20 to 2023 ..... 41.36 60 **Novatrelin**

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

Foam spray 500 mcg with calcipotriol 50 mcg per g ..... 59.95 60 g Enstilar

Gel 500 mcg with calcipotriol 50 mcg per g – 1% DV Dec-18 to 2021 ..... 52.24 60 g **Daivobet**

Oint 500 mcg with calcipotriol 50 mcg per g – 1% DV Dec-18 to 2021 ..... 19.95 30 g **Daivobet**

CALCIPOTRIOL

Oint 50 mcg per g ..... 40.00 120 g Daivonex

COAL TAR WITH SALICYLIC ACID AND SULPHUR

Oint 12% with salicylic acid 2% and sulphur 4%

METHOXSALEN [8-METHOXYPSORALEN]

Tab 10 mg

Lotn 1.2%

PIMECROLIMUS – **Restricted** see terms [below](#)

↓ Crm 1% – 1% DV Mar-21 to 2023 ..... 28.50 15 g **Elidel**

→ **Restricted (RS1781)**

#### Initiation

Dermatologist, paediatrician or ophthalmologist

Both:

- 1 Patient has atopic dermatitis on the eyelid; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy, documented allergy to topical corticosteroids, cataracts, glaucoma, or raised intraocular pressure.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN			
Soln 2.3% with trolamine laurilsulfate and fluorescein sodium – <b>1% DV</b>			
<b>Nov-20 to 2023</b> .....	4.44	500 ml	<b>Pinetarsol</b>
POTASSIUM PERMANGANATE			
Tab 400 mg			
Crystals			

Scalp Preparations

BETAMETHASONE VALERATE			
Scalp app 0.1% – <b>1% DV Oct-18 to 2021</b> .....	7.75	100 ml	<b>Beta Scalp</b>
CLOBETASOL PROPIONATE			
Scalp app 0.05% – <b>1% DV Nov-19 to 2022</b> .....	5.69	30 ml	<b>Dermol</b>
HYDROCORTISONE BUTYRATE			
Scalp lotn 0.1% – <b>1% DV Mar-19 to 2021</b> .....	7.30	100 ml	<b>Locoid</b>

Wart Preparations

IMIQUIMOD			
Crm 5%, 250 mg sachet .....	21.72	24	Perrigo
PODOPHYLLOTOXIN			
Soln 0.5% .....	33.60	3.5 ml	Condyline
SILVER NITRATE			
Sticks with applicator			

Other Skin Preparations

DIPHEMANIL METILSULFATE			
Powder 2%			
SUNSCREEN, PROPRIETARY			
Lotn – <b>1% DV Mar-20 to 2022</b> .....	5.10	200 g	<b>Marine Blue Lotion SPF 50+</b>

Antineoplastics

FLUOROURACIL SODIUM			
Crm 5% – <b>1% DV Sep-18 to 2021</b> .....	7.95	20 g	<b>Efudix</b>
METHYL AMINOLEVULINATE HYDROCHLORIDE – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Crm 16%			
➡ <b>Restricted (RS1127)</b>			
Dermatologist or plastic surgeon			

Wound Management Products

CALCIUM GLUCONATE			
Gel 2.5%			<i>e.g. Orion</i>

## GENITO-URINARY SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

### Anti-Infective Agents

#### ACETIC ACID

Soln 3%

Soln 5%

#### ACETIC ACID WITH HYDROXYQUINOLINE, GLYCEROL AND RICINOLEIC ACID

Jelly 0.94% with hydroxyquinoline sulphate 0.025%, glycerol 5% and  
ricinoleic acid 0.75% with applicator

#### CHLORHEXIDINE GLUCONATE

Crm 1%

Lotn 1%

#### CLOTRIMAZOLE

Vaginal crm 1% with applicator – 1% DV Jan-20 to 2022 .....2.50

35 g

**Clomazol**

Vaginal crm 2% with applicator – 1% DV Jan-20 to 2022 .....3.00

20 g

**Clomazol**

#### MICONAZOLE NITRATE

Vaginal crm 2% with applicator – 1% DV Nov-20 to 2023 .....6.89

40 g

**Micreme**

#### NIYSTATIN

Vaginal crm 100,000 u per 5 g with applicator(s) – 1% DV Oct-20 to 2023 ....4.00

75 g

**Nilstat**

### Contraceptives

#### Antiandrogen Oral Contraceptives

#### CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

Tab 2 mcg with ethinyloestradiol 35 mcg and 7 inert tablets.....4.67

168

Ginet

#### Combined Oral Contraceptives

#### ETHINYLOESTRADIOL WITH DESOGESTREL

Tab 20 mcg with desogestrel 150 mcg

Tab 30 mcg with desogestrel 150 mcg

#### ETHINYLOESTRADIOL WITH LEVONORGESTREL

Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets .....2.18

84

Microgynon 20 ED

Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets .....1.77

84

Leven ED

Tab 20 mcg with levonorgestrel 100 mcg

Tab 30 mcg with levonorgestrel 150 mcg

Tab 50 mcg with levonorgestrel 125 mcg.....9.45

84

Microgynon 50 ED

#### ETHINYLOESTRADIOL WITH NORETHISTERONE

Tab 35 mcg with norethisterone 1 mg

Tab 35 mcg with norethisterone 1 mg and 7 inert tab – 1% DV Mar-20  
to 2022 .....6.95

84

**Brevinor 1/28**

Tab 35 mcg with norethisterone 500 mcg

#### NORETHISTERONE WITH MESTRANOL

Tab 1 mg with mestranol 50 mcg

### Contraceptive Devices

#### INTRA-UTERINE DEVICE

IUD 29.1 mm length x 23.2 mm width – 1% DV Nov-19 to 2022 .....18.45

1

**Choice TT380 Short**

IUD 33.6 mm length x 29.9 mm width – 1% DV Nov-19 to 2022 .....18.45

1

**Choice TT380 Standard**

IUD 35.5 mm length x 19.6 mm width – 1% DV Nov-19 to 2022 .....15.50

1

**Choice Load 375**

Price  
(ex man. excl. GST)  
\$ Per Brand or  
Generic  
Manufacturer

## Emergency Contraception

LEVONORGESTREL

Tab 1.5 mg ..... 4.95 1 Postinor-1

## Progestogen-Only Contraceptives

LEVONORGESTREL

Tab 30 mcg – 1% DV May-20 to 2022 ..... 16.50 84 **Microlut**

Subdermal implant (2 x 75 mg rods) – 1% DV Dec-20 to 2023 ..... 106.92 1 **Jadelle**

Intra-uterine device 52 mg – 1% DV Nov-19 to 31 Oct 2022 ..... 269.50 1 **Mirena**

Intra-uterine device 13.5 mg – 1% DV Nov-19 to 31 Oct 2022 ..... 215.60 1 **Jaydess**

MEDROXYPROGESTERONE ACETATE

Inj 150 mg per ml, 1 ml syringe – 1% DV Dec-19 to 2022 ..... 7.98 1 **Depo-Provera**

NORETHISTERONE

Tab 350 mcg – 1% DV Sep-18 to 2021 ..... 6.25 84 **Noriday 28**

## Obstetric Preparations

### Antiprogestogens

MIFEPRISTONE

Tab 200 mg

### Oxytocics

CARBOPROST TROMETAMOL

Inj 250 mcg per ml, 1 ml ampoule

DINOPROSTONE

Pessaries 10 mg

Vaginal gel 1 mg in 3 g ..... 56.86 1 Prostin E2

Vaginal gel 2 mg in 3 g ..... 69.77 1 Prostin E2

ERGOMETRINE MALEATE

Inj 500 mcg per ml, 1 ml ampoule ..... 105.00 5 DBL Ergometrine

OXYTOCIN

Inj 5 iu per ml, 1 ml ampoule – 1% DV Nov-18 to 2021 ..... 3.98 5 **Oxytocin BNM**

Inj 10 iu per ml, 1 ml ampoule – 1% DV Nov-18 to 2021 ..... 4.98 5 **Oxytocin BNM**

OXYTOCIN WITH ERGOMETRINE MALEATE

Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampoule – 1%  
DV Oct-18 to 2021 ..... 15.00 5 **Syntometrine**

### Tocolytics

PROGESTERONE – **Restricted** see terms [below](#)

⚡ Cap 100 mg ..... 16.50 30 Utrogestan

➡ **Restricted (RS1533)**

**Initiation**

Gynaecologist or obstetrician

Re-assessment required after 12 months

Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 For the prevention of pre-term labour\*; and
- 2 Either:
  - 2.1 The patient has a short cervix on ultrasound (defined as < 25mm at 16 to 28 weeks); or
  - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

## Continuation

Gynaecologist or obstetrician

*Re-assessment required after 12 months*

All of the following:

- 1 For the prevention of pre-term labour\*; and
- 2 Treatment is required for second or subsequent pregnancy; and
- 3 Either:
  - 3.1 The patient has a short cervix on ultrasound (defined as < 25mm at 16 to 28 weeks); or
  - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with \* are unapproved indications.

TERBUTALINE – **Restricted** see terms [below](#)

↓ Inj 500 mcg ampoule

→ **Restricted** (RS1130)

Obstetrician

## Oestrogens

OESTRIOL

Crm 1 mg per g with applicator – 1% DV Oct-20 to 2023 .....	6.62	15 g	<b>Ovestin</b>
Pessaries 500 mcg – 1% DV Oct-20 to 2023 .....	6.86	15	<b>Ovestin</b>

## Urologicals

### 5-Alpha Reductase Inhibitors

FINASTERIDE – **Restricted** see terms [below](#)

↓ Tab 5 mg ..... 4.81 100 Ricit

→ **Restricted** (RS1131)

Initiation

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
  - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
  - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

### Alpha-1A Adrenoceptor Blockers

TAMSULOSIN HYDROCHLORIDE – **Restricted** see terms [below](#)

↓ Cap 400 mcg – 1% DV Jan-20 to 2022 ..... 17.73 100 Tamsulosin-Rex

→ **Restricted** (RS1132)

Initiation

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## Urinary Alkalisers

POTASSIUM CITRATE – **Restricted** see terms [below](#)

↓ Oral liq 3 mmol per ml – **1% DV Oct-18 to 2021** ..... 31.80      200 ml      **Biomed**

➔ **Restricted (RS1133)**

### Initiation

Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

SODIUM CITRO-TARTRATE

Grans eff 4 g sachets – **1% DV Oct-20 to 2023** ..... 2.22      28      **Ural**

## Urinary Antispasmodics

OXYBUTYNIN

Tab 5 mg ..... 11.70      500      Apo-Oxybutynin

Oral liq 5 mg per 5 ml ..... 60.40      473 ml      Apo-Oxybutynin

SOLIFENACIN SUCCINATE – **Some items restricted** see terms [below](#)

Tab 5 mg – **1% DV Dec-18 to 2021** ..... 3.00      30      **Solifenacin Mylan**

Tab 10 mg – **1% DV Dec-18 to 2021** ..... 5.50      30      **Solifenacin Mylan**

➔ **Restricted (RS1274)**

### Initiation

Patient has overactive bladder and a documented intolerance of, or is non-responsive to, oxybutynin.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## Anabolic Agents

### OXANDROLONE

↓ Tab 2.5 mg

→ **Restricted (RS1302)**

#### Initiation

For the treatment of burns patients.

## Androgen Agonists and Antagonists

### CYPROTERONE ACETATE

Tab 50 mg – **1% DV Dec-18 to 2021** ..... 13.17 50 **Siterone**

Tab 100 mg – **1% DV Dec-18 to 2021** ..... 26.75 50 **Siterone**

### TESTOSTERONE

Patch 5 mg per day ..... 90.00 30 **Androderm**

### TESTOSTERONE CIPIONATE

Inj 100 mg per ml, 10 ml vial..... 76.50 1 **Depo-Testosterone**

### TESTOSTERONE ESTERS

Inj testosterone decanoate 100 mg, testosterone isocarproate 60 mg,  
testosterone phenylpropionate 60 mg and testosterone propionate  
30 mg per ml, 1 ml ampoule

### TESTOSTERONE UNDECANOATE

Cap 40 mg – **1% DV Nov-18 to 2021** ..... 21.00 60 **Andriol Testocaps**

Inj 250 mg per ml, 4 ml vial..... 86.00 1 **Reandron 1000**

## Calcium Homeostasis

### CALCITONIN

Inj 100 iu per ml, 1 ml ampoule ..... 121.00 5 **Miacalcic**

### CINACALCET – **Restricted** see terms [below](#)

↓ Tab 30 mg – **1% DV Sep-18 to 2021** ..... 210.30 28 **Sensipar**

→ **Restricted (RS1540)**

#### Initiation

Nephrologist or endocrinologist

*Re-assessment required after 6 months*

Either:

- 1 All of the following:
  - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
  - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
  - 1.3 The patient is symptomatic; or
- 2 All of the following:
  - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriopathy); and
  - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
  - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
<b>Continuation</b>			
Nephrologist or endocrinologist			
Both:			
1 The patient's serum calcium level has fallen to < 3mmol/L; and			
2 The patient has experienced clinically significant symptom improvement.			
Note: This does not include parathyroid adenomas unless these have become malignant.			
<b>ZOLEDRONIC ACID</b>			
↓ Inj 4 mg per 5 ml, vial – <b>1% DV May-19 to 2021</b> .....	38.03	1	<b>Zoledronic acid Mylan</b>
➔ <b>Restricted (RS1602)</b>			
<b>Initiation – bone metastases</b>			
Oncologist, haematologist or palliative care specialist			
Any of the following:			
1 Patient has hypercalcaemia of malignancy; or			
2 Both:			
2.1 Patient has bone metastases or involvement; and			
2.2 Patient has severe bone pain resistant to standard first-line treatments; or			
3 Both:			
3.1 Patient has bone metastases or involvement; and			
3.2 Patient is at risk of skeletal-related events (pathological fracture, spinal cord compression, radiation to bone or surgery to bone).			
<b>Initiation – early breast cancer</b>			
Oncologist			
All of the following:			
1 Treatment to be used as adjuvant therapy for early breast cancer; and			
2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and			
3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.			
<b>Corticosteroids</b>			
<b>BETAMETHASONE</b>			
Tab 500 mcg			
Inj 4 mg per ml, 1 ml ampoule			
<b>BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE</b>			
Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml ampoule			
<b>DEXAMETHASONE</b>			
Tab 0.5 mg – <b>1% DV Oct-18 to 2021</b> .....	0.99	30	<b>Dexamethasone</b>
Tab 4 mg – <b>1% DV Oct-18 to 2021</b> .....	1.90	30	<b>Dexamethasone</b>
Oral liq 1 mg per ml .....	45.00	25 ml	<b>Biomed</b>
<b>DEXAMETHASONE PHOSPHATE</b>			
Inj 4 mg per ml, 1 ml ampoule – <b>1% DV Jul-20 to 2022</b> .....	9.25	10	<b>Dexamethasone Phosphate Panpharma</b>
Inj 4 mg per ml, 2 ml ampoule – <b>1% DV Jul-20 to 2022</b> .....	16.37	10	<b>Dexamethasone Phosphate Panpharma</b>
<b>FLUDROCORTISONE ACETATE</b>			
Tab 100 mcg .....	14.32	100	<b>Florinef</b>

## HORMONE PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>HYDROCORTISONE</b>			
Tab 5 mg – 1% DV Sep-18 to 2021 .....	8.10	100	<b>Douglas</b>
Tab 20 mg – 1% DV Sep-18 to 2021 .....	20.32	100	<b>Douglas</b>
Inj 100 mg vial .....	5.30	1	<b>Solu-Cortef</b>
<b>METHYLPREDNISOLONE (AS SODIUM SUCCINATE)</b>			
Tab 4 mg – 1% DV Dec-18 to 2021 .....	112.00	100	<b>Medrol</b>
Tab 100 mg – 1% DV Dec-18 to 2021 .....	194.00	20	<b>Medrol</b>
Inj 40 mg vial – 1% DV Dec-18 to 2021 .....	18.90	1	<b>Solu-Medrol Act-O-Vial</b>
Inj 125 mg vial – 1% DV Dec-18 to 2021 .....	28.90	1	<b>Solu-Medrol Act-O-Vial</b>
Inj 500 mg vial – 1% DV Dec-18 to 2021 .....	22.78	1	<b>Solu-Medrol Act-O-Vial</b>
Inj 1 g vial – 1% DV Dec-18 to 2021 .....	27.83	1	<b>Solu-Medrol</b>
<b>METHYLPREDNISOLONE ACETATE</b>			
Inj 40 mg per ml, 1 ml vial – 1% DV Dec-18 to 2021 .....	44.40	5	<b>Depo-Medrol</b>
<b>PREDNISOLONE</b>			
Oral liq 5 mg per ml – 1% DV Jun-18 to 2021 .....	6.00	30 ml	<b>Redipred</b>
Enema 200 mcg per ml, 100 ml			
<b>PREDNISONE</b>			
Tab 1 mg .....	10.68	500	<b>Apo-Prednisone</b>
Tab 2.5 mg .....	12.09	500	<b>Apo-Prednisone</b>
Tab 5 mg .....	11.09	500	<b>Apo-Prednisone</b>
Tab 20 mg .....	29.03	500	<b>Apo-Prednisone</b>
<b>TRIAMCINOLONE ACETONIDE</b>			
Inj 10 mg per ml, 1 ml ampoule – 5% DV Apr-21 to 2023 .....	20.80	5	<b>Kenacort-A 10</b>
Inj 40 mg per ml, 1 ml ampoule – 1% DV Apr-21 to 2023 .....	51.10	5	<b>Kenacort-A 40</b>
<b>TRIAMCINOLONE HEXACETONIDE</b>			
Inj 20 mg per ml, 1 ml vial			

## Hormone Replacement Therapy

### Oestrogens

<b>OESTRADIOL</b>			
Tab 1 mg			
Patch 25 mcg per day.....	6.12	8	<b>Estradot</b>
Patch 50 mcg per day.....	7.04	8	<b>Estradot</b>
Patch 75 mcg per day.....	7.91	8	<b>Estradot</b>
Patch 100 mcg per day.....	7.91	8	<b>Estradot</b>
<b>OESTRADIOL VALERATE</b>			
Tab 1 mg – 1% DV Sep-18 to 2021 .....	12.36	84	<b>Progynova</b>
Tab 2 mg – 1% DV Sep-18 to 2021 .....	12.36	84	<b>Progynova</b>
<b>OESTROGENS (CONJUGATED EQUINE)</b>			
Tab 300 mcg			
Tab 625 mcg			

### Progestogen and Oestrogen Combined Preparations

<b>OESTRADIOL WITH NORETHISTERONE ACETATE</b>			
Tab 1 mg with 0.5 mg norethisterone acetate			
Tab 2 mg with 1 mg norethisterone acetate			
Tab 2 mg with 1 mg norethisterone acetate (10), and tab 2 mg oestradiol (12) and tab 1 mg oestradiol (6)			

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)  
e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## OESTROGENS WITH MEDROXYPROGESTERONE ACETATE

- Tab 625 mcg conjugated equine with 2.5 mg medroxyprogesterone acetate  
 Tab 625 mcg conjugated equine with 5 mg medroxyprogesterone acetate

## Progestogens

## MEDROXYPROGESTERONE ACETATE

Tab 2.5 mg .....	3.75	30	Provera
Tab 5 mg .....	14.00	100	Provera
Tab 10 mg .....	7.15	30	Provera

## Other Endocrine Agents

CABERGOLINE – **Restricted** see terms [below](#)

↓ Tab 0.5 mg – 1% DV Sep-18 to 2021 .....	3.75	2	Dostinex
	15.20	8	Dostinex

→ **Restricted (RS1319)**

## Initiation

Any of the following:

- 1 Inhibition of lactation; or
- 2 Patient has pathological hyperprolactinemia; or
- 3 Patient has acromegaly.

## CLOMIFENE CITRATE

Tab 50 mg .....	29.84	10	Mylan Clomiphene
-----------------	-------	----	------------------

## DANAZOL

Cap 100 mg .....	19.13	28	Mylan
Cap 200 mg .....	97.83	100	Azol

(Mylan Cap 100 mg to be delisted 1 April 2021)

(Azol Cap 200 mg to be delisted 1 April 2021)

## GESTRINONE

Cap 2.5 mg

## METYRAPONE

Cap 250 mg

## PENTAGASTRIN

Inj 250 mcg per ml, 2 ml ampoule

## Other Oestrogen Preparations

## ETHINYLOESTRADIOL

Tab 10 mcg – 1% DV Sep-18 to 2021 .....	17.60	100	NZ Medical and Scientific
---	-------	-----	---------------------------

## OESTRADIOL

Implant 50 mg

## OESTRIOL

Tab 2 mg – 1% DV Sep-20 to 2023 .....	7.00	30	Ovestin
---------------------------------------	------	----	---------

## Other Progestogen Preparations

## MEDROXYPROGESTERONE

Tab 100 mg .....	101.00	100	Provera HD
------------------	--------	-----	------------

## HORMONE PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
NORETHISTERONE			
Tab 5 mg – 1% DV Dec-19 to 2021 .....	18.29	100	Primolut N

### Pituitary and Hypothalamic Hormones and Analogues

#### CORTICOTRORELIN (OVINE)

Inj 100 mcg vial

#### THYROTROPIN ALFA

Inj 900 mcg vial

### Adrenocorticotrophic Hormones

#### TETRACOSACTIDE [TETRACOSACTRIN]

Inj 250 mcg per ml, 1 ml ampoule ..... 75.00

Inj 1 mg per ml, 1 ml ampoule ..... 690.00

1 Synacthen  
1 Synacthen Depot

### GnRH Agonists and Antagonists

#### BUSERELIN

Inj 1 mg per ml, 5.5 ml vial

#### GONADORELIN

Inj 100 mcg vial

#### GOSERELIN

Implant 3.6 mg, syringe – 1% DV May-21 to 2023 ..... 65.68

66.48

Implant 10.8 mg, syringe – 1% DV May-21 to 2023 ..... 122.37

177.50

1 Teva  
Zoladex  
1 Teva  
Zoladex

*(Zoladex Implant 3.6 mg, syringe to be delisted 1 May 2021)*

*(Zoladex Implant 10.8 mg, syringe to be delisted 1 May 2021)*

#### LEUPRORELIN ACETATE

Inj 3.75 mg prefilled dual chamber syringe ..... 221.60

Inj 11.25 mg prefilled dual chamber syringe ..... 591.68

1 Lucrin Depot 1-month  
1 Lucrin Depot 3-month

### Gonadotrophins

#### CHORIOGONADOTROPIN ALFA

Inj 250 mcg in 0.5 ml syringe

### Growth Hormone

#### SOMATROPIN – Restricted see terms [below](#)

⚡ Inj 5 mg cartridge – 1% DV Oct-18 to 2021 ..... 34.88

⚡ Inj 10 mg cartridge – 1% DV Oct-18 to 2021 ..... 69.75

⚡ Inj 15 mg cartridge – 1% DV Oct-18 to 2021 ..... 104.63

1 Omnitrope  
1 Omnitrope  
1 Omnitrope

➡ Restricted (RS1549)

#### Initiation – growth hormone deficiency in children

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

Either:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
- 2 All of the following:
  - 2.1 Height velocity < 25th percentile for age; and adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
  - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
  - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
  - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
  - 2.5 Appropriate imaging of the pituitary gland has been obtained.

#### **Continuation – growth hormone deficiency in children**

Endocrinologist or paediatric endocrinologist

*Re-assessment required after 12 months*

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

#### **Initiation – Turner syndrome**

Endocrinologist or paediatric endocrinologist

*Re-assessment required after 12 months*

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

#### **Continuation – Turner syndrome**

Endocrinologist or paediatric endocrinologist

*Re-assessment required after 12 months*

All of the following:

- 1 Height velocity greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

#### **Initiation – short stature without growth hormone deficiency**

Endocrinologist or paediatric endocrinologist

*Re-assessment required after 12 months*

All of the following:

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

## Continuation – short stature without growth hormone deficiency

Endocrinologist or paediatric endocrinologist

*Re-assessment required after 12 months*

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 Current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

## Initiation – short stature due to chronic renal insufficiency

Endocrinologist, paediatric endocrinologist or renal physician on the recommendation of an endocrinologist or paediatric endocrinologist

*Re-assessment required after 12 months*

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
  - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73 m<sup>2</sup> as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l × 40 = corrected GFR (ml/min/1.73 m<sup>2</sup> ) in a child who may or may not be receiving dialysis; or
  - 6.2 The patient has received a renal transplant and has received < 5mg/ m<sup>2</sup> /day of prednisone or equivalent for at least 6 months.

## Continuation – short stature due to chronic renal insufficiency

Endocrinologist, paediatric endocrinologist or renal physician on the recommendation of an endocrinologist or paediatric endocrinologist

*Re-assessment required after 12 months*

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

**Initiation – Prader-Willi syndrome**

Endocrinologist or paediatric endocrinologist

*Re-assessment required after 12 months*

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
  - 5.1 Both:
    - 5.1.1 The patient is aged two years or older; and
    - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
  - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

**Continuation – Prader-Willi syndrome**

Endocrinologist or paediatric endocrinologist

*Re-assessment required after 12 months*

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient’s specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

**Initiation – adults and adolescents**

Endocrinologist or paediatric endocrinologist

*Re-assessment required after 12 months*

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient’s serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until it is within 1 standard deviation of the mean normal value for age and sex; and

The dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

## Continuation – adults and adolescents

Endocrinologist or paediatric endocrinologist

*Re-assessment required after 12 months*

Either:

- 1 All of the following:
  - 1.1 The patient has been treated with somatropin for < 12 months; and
  - 1.2 There has been an improvement in the Quality of Life Assessment defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
  - 1.3 Serum IGF-I levels have increased to within  $\pm 1$ SD of the mean of the normal range for age and sex; and
  - 1.4 The dose of somatropin does not exceed 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
  - 2.1 The patient has been treated with somatropin for more than 12 months; and
  - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
  - 2.3 Serum IGF-I levels have continued to be maintained within  $\pm 1$ SD of the mean of the normal range for age and sex (other than for obvious external factors); and
  - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients.

## Thyroid and Antithyroid Preparations

CARBIMAZOLE

Tab 5 mg

IODINE

Soln BP 50 mg per ml

LEVOTHYROXINE

Tab 25 mcg

Tab 50 mcg

Tab 100 mcg

LIOTHYRONINE SODIUM

↓ Tab 20 mcg

➔ **Restricted (RS1301)**

### Initiation

For a maximum of 14 days' treatment in patients with thyroid cancer who are due to receive radioiodine therapy.

Inj 20 mcg vial

Inj 100 mcg vial

POTASSIUM IODATE

Tab 170 mg

POTASSIUM PERCHLORATE

Cap 200 mg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

PROPYLTHIOURACIL – **Restricted** see terms [below](#)

↓ Tab 50 mg .....35.00 100 PTU

➔ **Restricted (RS1276)**

**Initiation**

Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Note: Propylthiouracil is not recommended for patients under the age of 18 years unless the patient is pregnant and other treatments are contraindicated.

**PROTIRELIN**

Inj 100 mcg per ml, 2 ml ampoule

Vasopressin Agents

**ARGIPRESSIN [VASOPRESSIN]**

Inj 20 u per ml, 1 ml ampoule

DESMOPRESSIN ACETATE – **Some items restricted** see terms [below](#)

↓ Tab 100 mcg.....25.00 30 Minirin

↓ Tab 200 mcg.....54.45 30 Minirin

Nasal spray 10 mcg per dose – **1% DV Nov-20 to 2023** .....27.95 6 ml **Desmopressin-PH&T**

Inj 4 mcg per ml, 1 ml ampoule

Inj 15 mcg per ml, 1 ml ampoule

Nasal drops 100 mcg per ml

➔ **Restricted (RS1339)**

**Initiation – Nocturnal enuresis**

Either:

- 1 The nasal forms of desmopressin are contraindicated; or
- 2 An enuresis alarm is contraindicated.

Note: Cranial diabetes insipidus and the nasal forms of desmopressin are contraindicated.

**TERLIPRESSIN**

Inj 0.1 mg per ml, 8.5 ml ampoule .....450.00 5 Glypressin

Inj 1 mg per 8.5 ml ampoule .....215.00 5 Glypressin

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Antibacterials</b>			
<b>Aminoglycosides</b>			
AMIKACIN – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Inj 5 mg per ml, 10 ml syringe			
↓ Inj 5 mg per ml, 5 ml syringe .....	18.50	1	Biomed
↓ Inj 15 mg per ml, 5 ml syringe			
↓ Inj 250 mg per ml, 2 ml vial – <b>1% DV Aug-18 to 2021</b> .....	265.00	5	<b>DBL Amikacin</b>
➔ <b>Restricted (RS1041)</b>			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
GENTAMICIN SULPHATE			
↓ Inj 10 mg per ml, 1 ml ampoule .....	25.00	5	DBL Gentamicin
↓ Inj 40 mg per ml, 2 ml ampoule .....	17.50	10	Pfizer
PAROMOMYCIN – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Cap 250 mg .....	126.00	16	Humatin
➔ <b>Restricted (RS1603)</b>			
Clinical microbiologist, infectious disease specialist or gastroenterologist			
STREPTOMYCIN SULPHATE – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Inj 400 mg per ml, 2.5 ml ampoule			
➔ <b>Restricted (RS1043)</b>			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
TOBRAMYCIN			
↓ Powder			
➔ <b>Restricted (RS1475)</b>			
<b>Initiation</b>			
For addition to orthopaedic bone cement.			
↓ Inj 40 mg per ml, 2 ml vial – <b>1% DV Sep-18 to 2021</b> .....	15.00	5	<b>Tobramycin Mylan</b>
➔ <b>Restricted (RS1044)</b>			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
↓ Inj 100 mg per ml, 5 ml vial			
➔ <b>Restricted (RS1044)</b>			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
↓ Solution for inhalation 60 mg per ml, 5 ml – <b>1% DV May-21 to 2023</b> .....	2,200.00 395.00	56 dose	TOBI <b>Tobramycin BNM</b>
➔ <b>Restricted (RS1435)</b>			
<b>Initiation</b>			
Patient has cystic fibrosis.			
<i>(TOBI Solution for inhalation 60 mg per ml, 5 ml to be delisted 1 May 2021)</i>			
<b>Carbapenems</b>			
ERTAPENEM – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Inj 1 g vial – <b>1% DV Aug-19 to 2022</b> .....	70.00	1	<b>Invanz</b>
➔ <b>Restricted (RS1045)</b>			
Clinical microbiologist or infectious disease specialist			
IMIPENEM WITH CILASTATIN – <b>Restricted</b> see terms <a href="#">on the next page</a>			
↓ Inj 500 mg with 500 mg cilastatin vial – <b>1% DV Jul-19 to 2022</b> .....	60.00	1	<b>Imipenem+Cilastatin RBX</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➔ Restricted (RS1046)</b>			
Clinical microbiologist or infectious disease specialist			
MEROPENEM – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Inj 500 mg vial – <b>1% DV Apr-21 to 2023</b> .....	4.00	1	Meropenem Ranbaxy
	33.92	10	<b>Meropenem-AFT</b>
↓ Inj 1 g vial – <b>1% DV Apr-21 to 2023</b> .....	8.00	1	Meropenem Ranbaxy
	45.04	10	<b>Meropenem-AFT</b>
<i>(Meropenem Ranbaxy Inj 500 mg vial to be delisted 1 April 2021)</i>			
<i>(Meropenem Ranbaxy Inj 1 g vial to be delisted 1 April 2021)</i>			
<b>➔ Restricted (RS1047)</b>			
Clinical microbiologist or infectious disease specialist			

### Cephalosporins and Cephamycins - 1st Generation

CEFALEXIN			
Cap 250 mg – <b>1% DV Nov-19 to 2022</b> .....	3.33	20	<b>Cephalexin ABM</b>
Cap 500 mg .....	3.95	20	Cephalexin ABM
Grans for oral liq 25 mg per ml – <b>1% DV Oct-18 to 2021</b> .....	8.75	100 ml	<b>Cefalexin Sandoz</b>
Grans for oral liq 50 mg per ml – <b>1% DV Oct-18 to 2021</b> .....	11.75	100 ml	<b>Cefalexin Sandoz</b>
CEFAZOLIN			
Inj 500 mg vial – <b>1% DV Nov-20 to 2023</b> .....	3.39	5	<b>AFT</b>
Inj 1 g vial – <b>1% DV Nov-20 to 2023</b> .....	3.49	5	<b>AFT</b>

### Cephalosporins and Cephamycins - 2nd Generation

CEFACTOR			
Cap 250 mg – <b>1% DV Oct-19 to 2022</b> .....	24.70	100	<b>Ranbaxy-Cefactor</b>
Grans for oral liq 25 mg per ml – <b>1% DV Oct-19 to 2022</b> .....	3.53	100 ml	<b>Ranbaxy-Cefactor</b>
CEFOXITIN			
Inj 1 g vial .....	58.00	10	Cefoxitin Actavis
<i>(Cefoxitin Actavis Inj 1 g vial to be delisted 1 January 2021)</i>			
CEFUROXIME			
Tab 250 mg – <b>1% DV Feb-20 to 2022</b> .....	45.93	50	<b>Zinnat</b>
Inj 750 mg vial .....	9.85	10	Cefuroxime Actavis
Inj 1.5 g vial .....	14.36	10	Cefuroxime Actavis

### Cephalosporins and Cephamycins - 3rd Generation

CEFOTAXIME			
Inj 500 mg vial .....	1.90	1	Cefotaxime Sandoz
Inj 1 g vial – <b>1% DV Nov-20 to 2023</b> .....	45.00	10	<b>DBL Cefotaxime</b>
CEFTAZIDIME – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Inj 1 g vial – <b>1% DV Dec-20 to 2023</b> .....	2.69	1	<b>Ceftazidime-AFT</b>
<b>➔ Restricted (RS1048)</b>			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
CEFTRIAOXONE			
Inj 500 mg vial – <b>1% DV Jan-20 to 2022</b> .....	0.89	1	<b>Ceftriaxone-AFT</b>
Inj 1 g vial – <b>1% DV Jan-20 to 2022</b> .....	3.99	5	<b>Ceftriaxone-AFT</b>
Inj 2 g vial – <b>1% DV Jan-20 to 2022</b> .....	1.98	1	<b>Ceftriaxone-AFT</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## Cephalosporins and Cephamycins - 4th Generation

CEFEPIME – **Restricted** see terms [below](#)

⚡ Inj 1 g vial – 1% DV Sep-18 to 2021 .....	3.75	1	Cefepime-AFT
⚡ Inj 2 g vial – 1% DV Sep-18 to 2021 .....	5.69	1	Cefepime-AFT

➡ **Restricted** (RS1049)

Clinical microbiologist or infectious disease specialist

## Cephalosporins and Cephamycins - 5th Generation

CEFTAROLINE FOSAMIL – **Restricted** see terms [below](#)

⚡ Inj 600 mg vial .....	1,595.00	10	Zinforo
-------------------------	----------	----	---------

➡ **Restricted** (RS1446)

**Initiation – multi-resistant organism salvage therapy**

Clinical microbiologist or infectious disease specialist

Either:

- 1 for patients where alternative therapies have failed; or
- 2 for patients who have a contraindication or hypersensitivity to standard current therapies.

## Macrolides

AZITHROMYCIN – **Restricted** see terms [below](#)

⚡ Tab 250 mg – 1% DV Sep-18 to 2021 .....	8.19	30	Apo-Azithromycin
⚡ Tab 500 mg – 1% DV Sep-18 to 2021 .....	0.93	2	Apo-Azithromycin
⚡ Grans for oral liq 200 mg per 5 ml (40 mg per ml) – 1% DV Dec-18 to 2021 .....	14.38	15 ml	Zithromax

➡ **Restricted** (RS1598)

**Initiation – bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections**

Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome\*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome\*; or
- 3 Patient has cystic fibrosis and has chronic infection with *Pseudomonas aeruginosa* or *Pseudomonas* related gram negative organisms\*; or
- 4 Patient has an atypical *Mycobacterium* infection.

Note: Indications marked with \* are unapproved indications

**Initiation – non-cystic fibrosis bronchiectasis\***

Respiratory specialist or paediatrician

*Re-assessment required after 12 months*

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis\*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
  - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
  - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with \* are unapproved indications. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis will be subsidised in the community.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

#### **Continuation – non-cystic fibrosis bronchiectasis\***

Respiratory specialist or paediatrician

*Re-assessment required after 12 months*

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

Note: Indications marked with \* are unapproved indications. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis will be subsidised in the community.

#### **Initiation – other indications**

*Re-assessment required after 5 days*

For any other condition.

#### **Continuation – other indications**

*Re-assessment required after 5 days*

For any other condition.

#### **CLARITHROMYCIN – Restricted see terms [below](#)**

↓ Tab 250 mg .....	3.98	14	Apo-Clarithromycin
↓ Tab 500 mg .....	10.40	14	Apo-Clarithromycin
↓ Grans for oral liq 50 mg per ml .....	192.00	50 ml	Klacid
↓ Inj 500 mg vial – <b>1% DV Dec-20 to 2023</b> .....	9.87	1	<b>Martindale</b>

#### **→ Restricted (RS1709)**

#### **Initiation – Tab 250 mg and oral liquid**

Any of the following:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug resistance or intolerance to standard pharmaceutical agents; or
- 3 Helicobacter pylori eradication; or
- 4 Prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated.

#### **Initiation – Tab 500 mg**

Helicobacter pylori eradication.

#### **Initiation – Infusion**

Any of the following:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug resistance or intolerance to standard pharmaceutical agents; or
- 3 Community-acquired pneumonia.

#### **ERYTHROMYCIN (AS ETHYLSUCCINATE)**

Tab 400 mg .....	16.95	100	E-Mycin
Grans for oral liq 200 mg per 5 ml .....	5.00	100 ml	E-Mycin
Grans for oral liq 400 mg per 5 ml .....	6.77	100 ml	E-Mycin

#### **ERYTHROMYCIN (AS LACTOBIONATE)**

Inj 1 g vial – <b>1% DV Dec-19 to 2022</b> .....	10.00	1	<b>Erythrocin IV</b>
--	-------	---	----------------------

#### **ERYTHROMYCIN (AS STEARATE) – Restricted: For continuation only**

→ Tab 250 mg

→ Tab 500 mg

#### **ROXITHROMYCIN – Some items restricted see terms [on the next page](#)**

↓ Tab dispersible 50 mg .....	8.29	10	Rulide D
Tab 150 mg – <b>1% DV Sep-19 to 2022</b> .....	8.28	50	<b>Arrow-Roxithromycin</b>
Tab 300 mg – <b>1% DV Sep-19 to 2022</b> .....	16.33	50	<b>Arrow-Roxithromycin</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➔ Restricted (RS1569)</b>			
<b>Initiation</b>			
Only for use in patients under 12 years of age.			
<b>Penicillins</b>			
<b>AMOXICILLIN</b>			
Cap 250 mg – 1% DV Apr-20 to 2022 .....	22.50	500	<b>Alphamox</b>
Cap 500 mg – 1% DV Apr-20 to 2022 .....	36.98	500	<b>Alphamox</b>
Grans for oral liq 125 mg per 5 ml – 1% DV Nov-20 to 2023 .....	1.40	100 ml	<b>Alphamox 125</b>
Grans for oral liq 250 mg per 5 ml – 1% DV Nov-20 to 2023 .....	1.73	100 ml	<b>Alphamox 250</b>
Inj 250 mg vial .....	10.67	10	Ibiamox
Inj 500 mg vial .....	12.41	10	Ibiamox
Inj 1 g vial .....	17.29	10	Ibiamox
<b>AMOXICILLIN WITH CLAVULANIC ACID</b>			
Tab 500 mg with clavulanic acid 125 mg .....	1.88	20	Augmentin
Grans for oral liq 25 mg with clavulanic acid 6.25 mg per ml .....	5.00	100 ml	Augmentin
Grans for oral liq 50 mg with clavulanic acid 12.5 mg per ml .....	2.20	100 ml	Curam
Inj 500 mg with clavulanic acid 100 mg vial .....	28.18	10	m-Amoxiclav
Inj 1,000 mg with clavulanic acid 200 mg vial .....	43.30	10	m-Amoxiclav
<b>BENZATHINE BENZYL PENICILLIN</b>			
Inj 900 mg (1.2 million units) in 2.3 ml syringe – 1% DV Dec-18 to 2021 .....	344.93	10	<b>Bicillin LA</b>
<b>BENZYL PENICILLIN SODIUM [PENICILLIN G]</b>			
Inj 600 mg (1 million units) vial – 1% DV Nov-20 to 2023 .....	11.09	10	<b>Sandoz</b>
<b>FLUCLOXACILLIN</b>			
Cap 250 mg – 1% DV Sep-18 to 2021 .....	16.83	250	<b>Staphlex</b>
Cap 500 mg – 1% DV Sep-18 to 2021 .....	56.61	500	<b>Staphlex</b>
Grans for oral liq 25 mg per ml – 1% DV Oct-18 to 2021 .....	2.29	100 ml	<b>AFT</b>
Grans for oral liq 50 mg per ml – 1% DV Oct-18 to 2021 .....	3.68	100 ml	<b>AFT</b>
Inj 250 mg vial .....	9.00	10	Flucloxin
Inj 500 mg vial .....	9.40	10	Flucloxin
Inj 1 g vial – 1% DV Nov-20 to 2023 .....	5.70	5	<b>Flucil</b>
<b>PHENOXYMETHYLPENICILLIN [PENICILLIN V]</b>			
Cap 250 mg – 1% DV Sep-18 to 2021 .....	2.59	50	<b>Cilicaine VK</b>
Cap 500 mg – 1% DV Sep-18 to 2021 .....	4.26	50	<b>Cilicaine VK</b>
Grans for oral liq 125 mg per 5 ml – 1% DV Jan-20 to 2022 .....	2.99	100 ml	<b>AFT</b>
Grans for oral liq 250 mg per 5 ml – 1% DV Jan-20 to 2022 .....	3.99	100 ml	<b>AFT</b>
<b>PIPERACILLIN WITH TAZOBACTAM – Restricted see terms <a href="#">below</a></b>			
⚡ Inj 4 g with tazobactam 0.5 g vial .....	38.00	10	PipTaz Sandoz PiperTaz Sandoz
<b>➔ Restricted (RS1053)</b>			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
<b>PROCAINE PENICILLIN</b>			
Inj 1.5 g in 3.4 ml syringe .....	123.50	5	Cilicaine
<b>TICARCILLIN WITH CLAVULANIC ACID – Restricted see terms <a href="#">below</a></b>			
⚡ Inj 3 g with clavulanic acid 0.1 mg vial			
<b>➔ Restricted (RS1054)</b>			
Clinical microbiologist, infectious disease specialist or respiratory specialist			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Quinolones</b>			
CIPROFLOXACIN – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Tab 250 mg – <b>1% DV Nov-20 to 2023</b> .....	2.42	28	<b>Cipflox</b>
↓ Tab 500 mg – <b>1% DV Nov-20 to 2023</b> .....	3.40	28	<b>Cipflox</b>
↓ Tab 750 mg – <b>1% DV Nov-20 to 2023</b> .....	5.95	28	<b>Cipflox</b>
↓ Oral liq 50 mg per ml			
↓ Oral liq 100 mg per ml			
↓ Inj 2 mg per ml, 100 ml bag – <b>1% DV Oct-18 to 2021</b> .....	68.20	10	<b>Cipflox</b>
→ <b>Restricted (RS1055)</b>			
Clinical microbiologist or infectious disease specialist			
MOXIFLOXACIN – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Tab 400 mg – <b>1% DV Dec-20 to 2023</b> .....	42.00	5	<b>Avelox</b>
↓ Inj 1.6 mg per ml, 250 ml bottle – <b>1% DV Apr-20 to 2022</b> .....	39.00	1	<b>Moxifloxacin Kabi</b>
→ <b>Restricted (RS1644)</b>			
<b>Initiation – Mycobacterium infection</b>			
Infectious disease specialist, clinical microbiologist or respiratory specialist			
Any of the following:			
1 Both:			
1.1 Active tuberculosis; and			
1.2 Any of the following:			
1.2.1 Documented resistance to one or more first-line medications; or			
1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or			
1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or			
1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or			
1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or			
2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated; or			
3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.			
<b>Initiation – Pneumonia</b>			
Infectious disease specialist or clinical microbiologist			
Either:			
1 Immunocompromised patient with pneumonia that is unresponsive to first-line treatment; or			
2 Pneumococcal pneumonia or other invasive pneumococcal disease highly resistant to other antibiotics.			
<b>Initiation – Penetrating eye injury</b>			
Ophthalmologist			
Five days treatment for patients requiring prophylaxis following a penetrating eye injury.			
<b>Initiation – Mycoplasma genitalium</b>			
All of the following:			
1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium and is symptomatic; and			
2 Either:			
2.1 Has tried and failed to clear infection using azithromycin; or			
2.2 Has laboratory confirmed azithromycin resistance; and			
3 Treatment is only for 7 days.			
NORFLOXACIN			
Tab 400 mg .....	135.00	100	Arrow-Norfloxacin

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Tetracyclines</b>			
DEMECLOCYCLINE HYDROCHLORIDE			
Tab 150 mg			
Cap 150 mg			
Cap 300 mg			
DOXYCYCLINE			
➔ Tab 50 mg – <b>Restricted:</b> For continuation only			
Tab 100 mg .....	64.43	500	Doxine
Inj 5 mg per ml, 20 ml vial			
MINOCYCLINE			
Tab 50 mg			
➔ Cap 100 mg – <b>Restricted:</b> For continuation only			
TETRACYCLINE			
Tab 250 mg .....	21.42	28	Accord
Cap 500 mg			
TIGECYCLINE – <b>Restricted</b> see terms <a href="#">below</a>			
⚡ Inj 50 mg vial			
➔ <b>Restricted</b> (RS1059)			
Clinical microbiologist or infectious disease specialist			
<b>Other Antibacterials</b>			
AZTREONAM – <b>Restricted</b> see terms <a href="#">below</a>			
⚡ Inj 1 g vial .....	364.92	10	Azactam
➔ <b>Restricted</b> (RS1277)			
Clinical microbiologist or infectious disease specialist			
CHLORAMPHENICOL – <b>Restricted</b> see terms <a href="#">below</a>			
⚡ Inj 1 g vial			
➔ <b>Restricted</b> (RS1277)			
Clinical microbiologist or infectious disease specialist			
CLINDAMYCIN – <b>Restricted</b> see terms <a href="#">below</a>			
⚡ Cap 150 mg – <b>1% DV Apr-20 to 2022</b> .....	4.61	24	<b>Dalacin C</b>
⚡ Oral liq 15 mg per ml			
⚡ Inj 150 mg per ml, 4 ml ampoule – <b>1% DV Oct-19 to 2022</b> .....	39.00	10	<b>Dalacin C</b>
➔ <b>Restricted</b> (RS1061)			
Clinical microbiologist or infectious disease specialist			
COLISTIN SULPHOMETHATE [COLESTIMETHATE] – <b>Restricted</b> see terms <a href="#">below</a>			
⚡ Inj 150 mg per ml, 1 ml vial.....	65.00	1	Colistin-Link
➔ <b>Restricted</b> (RS1062)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
DAPTOMYCIN – <b>Restricted</b> see terms <a href="#">below</a>			
⚡ Inj 500 mg vial .....	243.52	1	Cubicin
➔ <b>Restricted</b> (RS1063)			
Clinical microbiologist or infectious disease specialist			
FOSFOMYCIN – <b>Restricted</b> see terms <a href="#">below</a>			
⚡ Powder for oral solution, 3 g sachet			<i>e.g. UroFos</i>
➔ <b>Restricted</b> (RS1315)			
Clinical microbiologist or infectious disease specialist			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>LINCOMYCIN – Restricted</b> see terms <a href="#">below</a>			
↓ Inj 300 mg per ml, 2 ml vial			
➔ <b>Restricted (RS1065)</b>			
Clinical microbiologist or infectious disease specialist			
<b>LINEZOLID – Restricted</b> see terms <a href="#">below</a>			
↓ Tab 600 mg – <b>1% DV Oct-18 to 2021</b> .....	553.77	10	<b>Zyvox</b>
↓ Oral liq 20 mg per ml – <b>1% DV Dec-18 to 2021</b> .....	1,879.00	150 ml	<b>Zyvox</b>
↓ Inj 2 mg per ml, 300 ml bottle – <b>1% DV Feb-19 to 2021</b> .....	18.50	1	<b>Linezolid Kabi</b>
➔ <b>Restricted (RS1066)</b>			
Clinical microbiologist or infectious disease specialist			
<b>METHENAMINE (HEXAMINE) HIPPURATE</b>			
Tab 1 g .....	40.01	100	Hiprex
<b>NITROFURANTOIN</b>			
Tab 50 mg – <b>1% DV Apr-19 to 2021</b> .....	22.20	100	<b>Nifuran</b>
Tab 100 mg – <b>1% DV Apr-19 to 2021</b> .....	37.50	100	<b>Nifuran</b>
<b>PIVMECILLINAM – Restricted</b> see terms <a href="#">below</a>			
↓ Tab 200 mg			
➔ <b>Restricted (RS1322)</b>			
Clinical microbiologist or infectious disease specialist			
<b>SODIUM FUSIDATE [FUSIDIC ACID] – Restricted</b> see terms <a href="#">below</a>			
↓ Tab 250 mg .....	34.50	12	Fucidin
➔ <b>Restricted (RS1064)</b>			
Clinical microbiologist or infectious disease specialist			
<b>SULPHADIAZINE – Restricted</b> see terms <a href="#">below</a>			
↓ Tab 500 mg			
➔ <b>Restricted (RS1067)</b>			
Clinical microbiologist, infectious disease specialist or maternal-foetal medicine specialist			
<b>TEICOPLANIN – Restricted</b> see terms <a href="#">below</a>			
↓ Inj 400 mg vial – <b>1% DV Jul-20 to 2021</b> .....	56.50	1	<b>Teicoplanin Mylan</b>
➔ <b>Restricted (RS1068)</b>			
Clinical microbiologist or infectious disease specialist			
<b>TRIMETHOPRIM</b>			
Tab 100 mg			
Tab 300 mg – <b>1% DV Oct-18 to 2021</b> .....	16.50	50	<b>TMP</b>
<b>TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXAZOLE]</b>			
Tab 80 mg with sulphamethoxazole 400 mg			
Oral liq 8 mg with sulphamethoxazole 40 mg per ml.....	2.97	100 ml	Deprim
Inj 16 mg with sulphamethoxazole 80 mg per ml, 5 ml ampoule			
<b>VANCOMYCIN – Restricted</b> see terms <a href="#">below</a>			
↓ Inj 500 mg vial – <b>1% DV Oct-20 to 2023</b> .....	2.35	1	<b>Mylan</b>
➔ <b>Restricted (RS1069)</b>			
Clinical microbiologist or infectious disease specialist			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## Antifungals

### Imidazoles

#### KETOCONAZOLE

⚡ Tab 200 mg

➡ **Restricted (RS1410)**

Oncologist

### Polyene Antimycotics

#### AMPHOTERICIN B

⚡ Inj (liposomal) 50 mg vial.....	3,450.00	10	AmBisome
-----------------------------------	----------	----	----------

➡ **Restricted (RS1071)**

#### Initiation

Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist

Either:

- 1 Proven or probable invasive fungal infection, to be prescribed under an established protocol; or
- 2 Both:
  - 2.1 Possible invasive fungal infection; and
  - 2.2 A multidisciplinary team (including an infectious disease physician or a clinical microbiologist) considers the treatment to be appropriate.

⚡ Inj 50 mg vial

➡ **Restricted (RS1316)**

Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist

#### NYSTATIN

Tab 500,000 u .....	17.09	50	Nilstat
Cap 500,000 u .....	15.47	50	Nilstat

### Triazoles

#### FLUCONAZOLE – **Restricted** see terms [below](#)

⚡ Cap 50 mg – 1% DV Nov-20 to 2023 .....	2.75	28	<b>Mylan</b>
⚡ Cap 150 mg – 1% DV Nov-20 to 2023 .....	0.65	1	<b>Mylan</b>
⚡ Cap 200 mg – 1% DV Nov-20 to 2023 .....	12.89	28	<b>Mylan</b>
⚡ Oral liquid 50 mg per 5 ml .....	109.34	35 ml	Diflucan
⚡ Inj 2 mg per ml, 50 ml vial – 1% DV Oct-19 to 2022 .....	2.80	1	<b>Fluconazole-Clarix</b>
⚡ Inj 2 mg per ml, 100 ml vial – 1% DV Oct-19 to 2022 .....	3.45	1	<b>Fluconazole-Clarix</b>

➡ **Restricted (RS1072)**

Consultant

#### ITRACONAZOLE – **Restricted** see terms [below](#)

⚡ Cap 100 mg – 1% DV Nov-19 to 2022 .....	4.27	15	<b>Itrazole</b>
⚡ Oral liquid 10 mg per ml			

➡ **Restricted (RS1073)**

Clinical immunologist, clinical microbiologist, dermatologist or infectious disease specialist

#### POSACONAZOLE – **Restricted** see terms [on the next page](#)

⚡ Tab modified-release 100 mg .....	869.86	24	Noxafil
⚡ Oral liq 40 mg per ml .....	761.13	105 ml	Noxafil

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

### ➔ Restricted (RS1074)

#### Initiation

Haematologist or infectious disease specialist

*Re-assessment required after 6 weeks*

Both:

- 1 Either:
  - 1.1 Patient has acute myeloid leukaemia; or
  - 1.2 Patient is planned to receive a stem cell transplant and is at high risk for aspergillus infection; and
- 2 Patient is to be treated with high dose remission induction therapy or re-induction therapy.

#### Continuation

Haematologist or infectious disease specialist

*Re-assessment required after 6 weeks*

Both:

- 1 Patient has previously received posaconazole prophylaxis during remission induction therapy; and
- 2 Any of the following:
  - 2.1 Patient is to be treated with high dose remission re-induction therapy; or
  - 2.2 Patient is to be treated with high dose consolidation therapy; or
  - 2.3 Patient is receiving a high risk stem cell transplant.

VORICONAZOLE – **Restricted** see terms [below](#)

↓ Tab 50 mg – 1% DV Sep-18 to 2021 .....	91.00	56	<b>Vttack</b>
↓ Tab 200 mg – 1% DV Sep-18 to 2021 .....	350.00	56	<b>Vttack</b>
↓ Powder for oral suspension 40 mg per ml – 1% DV Dec-18 to 2021.....	1,437.00	70 ml	<b>Vfend</b>
↓ Inj 200 mg vial – 1% DV Oct-19 to 2022.....	44.00	1	<b>Neo Health</b>

### ➔ Restricted (RS1075)

#### Initiation – Proven or probable aspergillus infection

Clinical microbiologist, haematologist or infectious disease specialist

Both:

- 1 Patient is immunocompromised; and
- 2 Patient has proven or probable invasive aspergillus infection.

#### Initiation – Possible aspergillus infection

Clinical microbiologist, haematologist or infectious disease specialist

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient has possible invasive aspergillus infection; and
- 3 A multidisciplinary team (including an infectious disease physician) considers the treatment to be appropriate.

#### Initiation – Resistant candidiasis infections and other moulds

Clinical microbiologist, haematologist or infectious disease specialist

All of the following:

- 1 Patient is immunocompromised; and
- 2 Either:
  - 2.1 Patient has fluconazole resistant candidiasis; or
  - 2.2 Patient has mould strain such as *Fusarium* spp. and *Scedosporium* spp; and
- 3 A multidisciplinary team (including an infectious disease physician or clinical microbiologist) considers the treatment to be appropriate.

## Other Antifungals

CASPOFUNGIN – **Restricted** see terms [on the next page](#)

↓ Inj 50 mg vial – 1% DV Dec-19 to 2022 .....	220.28	1	<b>Max Health</b>
↓ Inj 70 mg vial – 1% DV Dec-19 to 2022.....	284.63	1	<b>Max Health</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## ➔ **Restricted (RS1076)**

### Initiation

Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist  
Either:

- 1 Proven or probable invasive fungal infection, to be prescribed under an established protocol; or
- 2 Both:
  - 2.1 Possible invasive fungal infection; and
  - 2.2 A multidisciplinary team (including an infectious disease physician or a clinical microbiologist) considers the treatment to be appropriate.

FLUCYTOSINE – **Restricted** see terms [below](#)

⚡ Cap 500 mg

## ➔ **Restricted (RS1279)**

Clinical microbiologist or infectious disease specialist

TERBINAFINE

Tab 250 mg .....	1.33	14	Deolate
------------------	------	----	---------

## Antimycobacterials

### Antileprotics

CLOFAZIMINE – **Restricted** see terms [below](#)

⚡ Cap 50 mg

## ➔ **Restricted (RS1077)**

Clinical microbiologist, dermatologist or infectious disease specialist

DAPSONE – **Restricted** see terms [below](#)

⚡ Tab 25 mg .....	268.50	100	Dapsone
⚡ Tab 100 mg .....	329.50	100	Dapsone

## ➔ **Restricted (RS1078)**

Clinical microbiologist, dermatologist or infectious disease specialist

### Antituberculotics

CYCLOSERINE – **Restricted** see terms [below](#)

⚡ Cap 250 mg

## ➔ **Restricted (RS1079)**

Clinical microbiologist, infectious disease specialist or respiratory specialist

ETHAMBUTOL HYDROCHLORIDE – **Restricted** see terms [below](#)

⚡ Tab 100 mg .....			
⚡ Tab 400 mg .....	49.34	56	Myambutol

## ➔ **Restricted (RS1080)**

Clinical microbiologist, infectious disease specialist or respiratory specialist

ISONIAZID – **Restricted** see terms [below](#)

⚡ Tab 100 mg – 1% DV Oct-18 to 2021 .....	22.00	100	PSM
---	-------	-----	-----

## ➔ **Restricted (RS1281)**

Clinical microbiologist, dermatologist, paediatrician, public health physician or internal medicine physician

ISONIAZID WITH RIFAMPICIN – **Restricted** see terms [below](#)

⚡ Tab 100 mg with rifampicin 150 mg – 1% DV Sep-18 to 2021 .....	85.54	100	Rifinah
⚡ Tab 150 mg with rifampicin 300 mg – 1% DV Sep-18 to 2021 .....	170.60	100	Rifinah

## ➔ **Restricted (RS1282)**

Clinical microbiologist, dermatologist, paediatrician, public health physician or internal medicine physician

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PARA-AMINOSALICYLIC ACID – Restricted</b> see terms <a href="#">below</a>			
↓ Grans for oral liq 4 g.....	280.00	30	Paser
→ <b>Restricted (RS1083)</b>			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
<b>PROTIONAMIDE – Restricted</b> see terms <a href="#">below</a>			
↓ Tab 250 mg.....	305.00	100	Peteha
→ <b>Restricted (RS1084)</b>			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
<b>PYRAZINAMIDE – Restricted</b> see terms <a href="#">below</a>			
↓ Tab 500 mg.....			
→ <b>Restricted (RS1085)</b>			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
<b>RIFABUTIN – Restricted</b> see terms <a href="#">below</a>			
↓ Cap 150 mg.....	299.75	30	Mycobutin
→ <b>Restricted (RS1086)</b>			
Clinical microbiologist, gastroenterologist, infectious disease specialist or respiratory specialist			
<b>RIFAMPICIN – Restricted</b> see terms <a href="#">below</a>			
↓ Cap 150 mg – 1% DV Nov-20 to 2023.....	58.54	100	<b>Rifadin</b>
↓ Cap 300 mg – 1% DV Nov-20 to 2023.....	122.06	100	<b>Rifadin</b>
↓ Oral liq 100 mg per 5 ml – 1% DV Nov-20 to 2023.....	12.60	60 ml	<b>Rifadin</b>
↓ Inj 600 mg vial – 1% DV Nov-20 to 2023.....	134.98	1	<b>Rifadin</b>
→ <b>Restricted (RS1087)</b>			
Clinical microbiologist, dermatologist, internal medicine physician, paediatrician or public health physician			

## Antiparasitics

### Anthelmintics

<b>ALBENDAZOLE – Restricted</b> see terms <a href="#">below</a>			
↓ Tab 200 mg.....			
↓ Tab 400 mg.....			
→ <b>Restricted (RS1088)</b>			
Clinical microbiologist or infectious disease specialist			
<b>IVERMECTIN – Restricted</b> see terms <a href="#">below</a>			
↓ Tab 3 mg.....	17.20	4	Stromectol
→ <b>Restricted (RS1283)</b>			
Clinical microbiologist, dermatologist or infectious disease specialist			
<b>MEBENDAZOLE</b>			
Tab 100 mg.....	24.19	24	De-Worm
	7.97	6	Vermox
Oral liq 100 mg per 5 ml			
<i>(De-Worm Tab 100 mg to be delisted 1 March 2021)</i>			
<b>PRAZIQUANTEL</b>			
Tab 600 mg.....			

### Antiprotozoals

<b>ARTEMETHER WITH LUMEFANTRINE – Restricted</b> see terms <a href="#">on the next page</a>			
↓ Tab 20 mg with lumefantrine 120 mg.....			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔ <b>Restricted (RS1090)</b> Clinical microbiologist or infectious disease specialist ARTESUNATE – <b>Restricted</b> see terms <a href="#">below</a> ⚡ Inj 60 mg vial			
➔ <b>Restricted (RS1091)</b> Clinical microbiologist or infectious disease specialist ATOVAQUONE WITH PROGUANIL HYDROCHLORIDE – <b>Restricted</b> see terms <a href="#">below</a> ⚡ Tab 62.5 mg with proguanil hydrochloride 25 mg.....	25.00	12	Malarone Junior
⚡ Tab 250 mg with proguanil hydrochloride 100 mg.....	64.00	12	Malarone
➔ <b>Restricted (RS1092)</b> Clinical microbiologist or infectious disease specialist CHLOROQUINE PHOSPHATE – <b>Restricted</b> see terms <a href="#">below</a> ⚡ Tab 250 mg			
➔ <b>Restricted (RS1093)</b> Clinical microbiologist, dermatologist, infectious disease specialist or rheumatologist MEFLOQUINE – <b>Restricted</b> see terms <a href="#">below</a> ⚡ Tab 250 mg			
➔ <b>Restricted (RS1094)</b> Clinical microbiologist, dermatologist, infectious disease specialist or rheumatologist METRONIDAZOLE Tab 200 mg – 1% DV Dec-20 to 2023 .....	33.15	250	<b>Metrogyl</b>
Tab 400 mg – 1% DV Dec-20 to 2023 .....	5.23	21	<b>Metrogyl</b>
Oral liq benzoate 200 mg per 5 ml .....	25.00	100 ml	Flagyl-S
Injection 5 mg per ml, 100 ml bottle.....	1.39	100 ml	AFT
Inj 5 mg per ml, 100 ml bottle .....	34.80	20	Colpocin-T
Inj 5 mg per ml, 100 ml bag – 1% DV Feb-21 to 2023.....	27.50	10	<b>Baxter</b>
Suppos 500 mg .....	24.48	10	Flagyl
<i>(AFT Injection 5 mg per ml, 100 ml bottle to be delisted 1 February 2021)</i> <i>(Colpocin-T Inj 5 mg per ml, 100 ml bottle to be delisted 1 February 2021)</i>			
NITAZOXANIDE – <b>Restricted</b> see terms <a href="#">below</a> ⚡ Tab 500 mg .....	1,680.00	30	Alinia
⚡ Oral liq 100 mg per 5 ml			
➔ <b>Restricted (RS1095)</b> Clinical microbiologist or infectious disease specialist ORNIDAZOLE Tab 500 mg .....	32.95	10	Arrow-Ornidazole
PENTAMIDINE ISETHIONATE – <b>Restricted</b> see terms <a href="#">below</a> ⚡ Inj 300 mg vial – 1% DV Nov-19 to 2022.....	216.00	5	<b>Pentacarinat</b>
➔ <b>Restricted (RS1096)</b> Clinical microbiologist or infectious disease specialist PRIMAQUINE – <b>Restricted</b> see terms <a href="#">below</a> ⚡ Tab 15 mg ⚡ Tab 7.5 mg			
➔ <b>Restricted (RS1097)</b> Clinical microbiologist or infectious disease specialist PYRIMETHAMINE – <b>Restricted</b> see terms <a href="#">below</a> ⚡ Tab 25 mg			
➔ <b>Restricted (RS1098)</b> Clinical microbiologist, infectious disease specialist or maternal-foetal medicine specialist			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>QUININE DIHYDROCHLORIDE – Restricted</b> see terms <a href="#">below</a>			
↓ Inj 60 mg per ml, 10 ml ampoule			
↓ Inj 300 mg per ml, 2 ml vial			
→ <b>Restricted (RS1099)</b>			
Clinical microbiologist or infectious disease specialist			
<b>QUININE SULPHATE</b>			
Tab 300 mg	61.91	500	Q 300
<b>SODIUM STIBOGLUCONATE – Restricted</b> see terms <a href="#">below</a>			
↓ Inj 100 mg per ml, 1 ml vial			
→ <b>Restricted (RS1100)</b>			
Clinical microbiologist or infectious disease specialist			
<b>SPIRAMYCIN – Restricted</b> see terms <a href="#">below</a>			
↓ Tab 500 mg			
→ <b>Restricted (RS1101)</b>			
Maternal-foetal medicine specialist			
<b>Antiretrovirals</b>			
<b>Non-Nucleoside Reverse Transcriptase Inhibitors</b>			
→ <b>Restricted (RS1571)</b>			
<b>Initiation – Confirmed HIV</b>			
Patient has confirmed HIV infection.			
<b>Initiation – Prevention of maternal transmission</b>			
Either:			
1 Prevention of maternal foetal transmission; or			
2 Treatment of the newborn for up to eight weeks.			
<b>Initiation – Post-exposure prophylaxis following non-occupational exposure to HIV</b>			
Both:			
1 Treatment course to be initiated within 72 hours post exposure; and			
2 Any of the following:			
2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or			
2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or			
2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.			
<b>Initiation – Percutaneous exposure</b>			
Patient has percutaneous exposure to blood known to be HIV positive.			
<b>EFAVIRENZ – Restricted</b> see terms <a href="#">above</a>			
† Tab 200 mg	190.15	90	Stocrin
† Tab 600 mg	63.38	30	Stocrin
† Oral liq 30 mg per ml			
<b>ETRAVIRINE – Restricted</b> see terms <a href="#">above</a>			
† Tab 200 mg	770.00	60	Intelence
<b>NEVIRAPINE – Restricted</b> see terms <a href="#">above</a>			
† Tab 200 mg – <b>1% DV Sep-18 to 2021</b>	60.00	60	<b>Nevirapine Alphapharm</b>
† Oral suspension 10 mg per ml	203.55	240 ml	Viramune Suspension

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

## Nucleoside Reverse Transcriptase Inhibitors

### ➔ Restricted (RS1572)

#### Initiation – Confirmed HIV

Patient has confirmed HIV infection.

#### Initiation – Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

#### Initiation – Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

#### Initiation – Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

#### ABACAVIR SULPHATE – Restricted see terms [above](#)

† Tab 300 mg – 1% DV Jul-19 to 2022 .....	180.00	60	<b>Ziagen</b>
† Oral liq 20 mg per ml .....	256.31	240 ml	Ziagen

#### ABACAVIR SULPHATE WITH LAMIVUDINE – Restricted see terms [above](#)

† Tab 600 mg with lamivudine 300 mg – 1% DV Jul-19 to 2022 .....	63.00	30	<b>Kivexa</b>
--	-------	----	---------------

#### EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPROXIL – Restricted see terms [above](#)

† Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil 245 mg (300 mg as a maleate) – 1% DV Jun-19 to 2022 .....	106.88	30	<b>Mylan</b>
--	--------	----	--------------

#### EMTRICITABINE – Restricted see terms [above](#)

† Cap 200 mg – 1% DV Jul-19 to 2022 .....	307.20	30	<b>Emtriva</b>
---	--------	----	----------------

#### LAMIVUDINE – Restricted see terms [above](#)

† Tab 150 mg – 1% DV Nov-20 to 2023 .....	84.50	60	<b>Lamivudine Alphapharm</b>
---	-------	----	----------------------------------

† Oral liq 10 mg per ml

#### STAVUDINE – Restricted see terms [above](#)

† Cap 30 mg

† Cap 40 mg

† Powder for oral soln 1 mg per ml

#### ZIDOVUDINE [AZT] – Restricted see terms [above](#)

† Cap 100 mg .....	152.25	100	<b>Retrovir</b>
† Oral liq 10 mg per ml .....	30.45	200 ml	<b>Retrovir</b>
† Inj 10 mg per ml, 20 ml vial .....	750.00	5	<b>Retrovir IV</b>

#### ZIDOVUDINE [AZT] WITH LAMIVUDINE – Restricted see terms [above](#)

† Tab 300 mg with lamivudine 150 mg .....	33.00	60	<b>Alphapharm</b>
---	-------	----	-------------------

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

## Protease Inhibitors

### ➔ Restricted (RS1573)

#### Initiation – Confirmed HIV

Patient has confirmed HIV infection.

#### Initiation – Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

#### Initiation – Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

#### Initiation – Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

#### ATAZANAVIR SULPHATE – Restricted see terms [above](#)

† Cap 150 mg – 1% DV Jun-19 to 2022	141.68	60	Teva
† Cap 200 mg – 1% DV Jun-19 to 2022	188.91	60	Teva

#### DARUNAVIR – Restricted see terms [above](#)

† Tab 400 mg – 1% DV Apr-21 to 2023	132.00	60	Darunavir Mylan
	335.00		Prezista
† Tab 600 mg – 1% DV Apr-21 to 2023	196.65	60	Darunavir Mylan
	476.00		Prezista

(Prezista Tab 400 mg to be delisted 1 April 2021)

(Prezista Tab 600 mg to be delisted 1 April 2021)

#### INDINAVIR – Restricted see terms [above](#)

- † Cap 200 mg
- † Cap 400 mg

#### LOPINAVIR WITH RITONAVIR – Restricted see terms [above](#)

† Tab 100 mg with ritonavir 25 mg	183.75	60	Kaletra
† Tab 200 mg with ritonavir 50 mg	463.00	120	Kaletra
† Oral liq 80 mg with ritonavir 20 mg per ml	735.00	300 ml	Kaletra

#### RITONAVIR – Restricted see terms [above](#)

† Tab 100 mg – 1% DV Jul-19 to 2022	43.31	30	Norvir
-------------------------------------	-------	----	--------

## Strand Transfer Inhibitors

### ➔ Restricted (RS1574)

#### Initiation – Confirmed HIV

Patient has confirmed HIV infection.

#### Initiation – Prevention of maternal transmission

Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

## Initiation – Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

## Initiation – Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

**DOLUTEGRAVIR – Restricted** see terms [on the previous page](#)

⚡ Tab 50 mg ..... 1,090.00 30 Tivicay

**RALTEGRAVIR POTASSIUM – Restricted** see terms [on the previous page](#)

⚡ Tab 400 mg ..... 1,090.00 60 Isentress  
 ⚡ Tab 600 mg ..... 1,090.00 60 Isentress HD

## Antivirals

### Hepatitis B

**ADEFOVIR DIPIVOXIL – Restricted** see terms [below](#)

⚡ Tab 10 mg ..... 670.00 30 Hepsera  
 (Hepsera Tab 10 mg to be delisted 1 March 2021)

➡ **Restricted (RS1104)**

#### Initiation

Gastroenterologist or infectious disease specialist

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg+); and Documented resistance to lamivudine defined as:
- 2 Patient has raised serum ALT ( $> 1 \times \text{ULN}$ ); and
- 3 Patient has HBV DNA greater than 100,000 copies per mL, or viral load greater than or equal to 10-fold over nadir; and
- 4 Detection of M204I or M204V mutation; and
- 5 Either:
  - 5.1 Both:
    - 5.1.1 Patient is cirrhotic; and
    - 5.1.2 Adefovir dipivoxil to be used in combination with lamivudine; or
  - 5.2 Both:
    - 5.2.1 Patient is not cirrhotic; and
    - 5.2.2 Adefovir dipivoxil to be used as monotherapy.

**ENTECAVIR**

Tab 0.5 mg – **1% DV Nov-18 to 2021** ..... 52.00 30 **Entecavir Sandoz**

**LAMIVUDINE**

Tab 100 mg – **1% DV Nov-20 to 2023** ..... 6.95 28 **Zetlam**  
 Oral liq 5 mg per ml ..... 270.00 240 ml Zeffix

**TENOFOVIR DISOPROXIL**

Tab 245 mg (300.6 mg as a succinate) – **1% DV Sep-18 to 2021** ..... 38.10 30 **Tenofovir Disoproxil Teva**

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Hepatitis C</b>			
GLECAPREVIR WITH PIBRENTASVIR			
Note: the supply of treatment is via PHARMAC's approved direct distribution supply. Further details can be found on PHARMAC's website <a href="https://www.pharmac.govt.nz/maviret">https://www.pharmac.govt.nz/maviret</a> .			
Tab 100 mg with pibrentasvir 40 mg .....	24,750.00	84	Maviret
LEDIPASVIR WITH SOFOSBUVIR – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Tab 90 mg with sofosbuvir 400 mg.....	24,363.46	28	Harvoni
→ <b>Restricted (RS1528)</b>			
<b>Initiation</b>			
Note: Only for use in patients with approval by the Hepatitis C Treatment Panel (HepCTP). Applications will be considered by HepCTP at its regular meetings and approved subject to eligibility according to the Access Criteria (set out in Section B of the Pharmaceutical Schedule).			
<b>Herpesviridae</b>			
ACICLOVIR			
Tab dispersible 200 mg – 1% DV Oct-19 to 2022 .....	1.60	25	Lovir
Tab dispersible 400 mg – 1% DV Oct-19 to 2022 .....	5.38	56	Lovir
Tab dispersible 800 mg – 1% DV Oct-19 to 2022 .....	5.98	35	Lovir
Inj 250 mg vial – 1% DV Sep-18 to 2021 .....	9.60	5	Aciclovir-Baxter Aciclovir-Claris
<i>(Aciclovir-Claris Inj 250 mg vial to be delisted 1 March 2021)</i>			
CIDOFOVIR – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Inj 75 mg per ml, 5 ml vial			
→ <b>Restricted (RS1108)</b>			
Clinical microbiologist, infectious disease specialist, otolaryngologist or oral surgeon			
FOSCARNET SODIUM – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Inj 24 mg per ml, 250 ml bottle			
→ <b>Restricted (RS1109)</b>			
Clinical microbiologist or infectious disease specialist			
GANCICLOVIR – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Inj 500 mg vial .....	380.00	5	Cymevene
→ <b>Restricted (RS1110)</b>			
Clinical microbiologist or infectious disease specialist			
VALACICLOVIR			
Tab 500 mg – 1% DV Sep-18 to 2021 .....	5.75	30	Vaclovir
Tab 1,000 mg – 1% DV Sep-18 to 2021 .....	11.35	30	Vaclovir
VALGANCICLOVIR – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Tab 450 mg – 1% DV May-19 to 2021.....	225.00	60	Valganciclovir Mylan
→ <b>Restricted (RS1112)</b>			
<b>Initiation – Transplant cytomegalovirus prophylaxis</b>			
<i>Limited to 3 months treatment</i>			
Patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.			
<b>Initiation – Lung transplant cytomegalovirus prophylaxis</b>			
<i>Limited to 6 months treatment</i>			
Both:			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient has undergone a lung transplant; and
- 2 Either:
  - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
  - 2.2 The recipient is cytomegalovirus positive.

#### Initiation – Cytomegalovirus in immunocompromised patients

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

### HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL – **Restricted** see terms [below](#)

↓ Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a succinate) – 1% DV Jun-19 to 2022 .....	61.15	30	Teva
→ <b>Restricted (RS1737)</b>			

#### Initiation – Confirmed HIV

Patient has confirmed HIV infection.

#### Initiation – Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

#### Initiation – Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

#### Initiation – Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

#### Initiation – Pre-exposure prophylaxis

*Re-assessment required after 3 months*

All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or <https://ashm.org.au/HIV/PrEP/> for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
  - 6.1 All of the following:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 6.1.1 Patient is male or transgender; and
- 6.1.2 Patient has sex with men; and
- 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
- 6.1.4 Any of the following:
  - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
  - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
  - 6.1.4.3 Patient has used methamphetamine in the last three months; or
- 6.2 All of the following:
  - 6.2.1 Patient has a regular partner who has HIV infection; and
  - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
  - 6.2.3 Condoms have not been consistently used.

### Continuation – Pre-exposure prophylaxis

*Re-assessment required after 3 months*

All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or <https://ashm.org.au/HIV/PrEP/> for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
  - 6.1 All of the following:
    - 6.1.1 Patient is male or transgender; and
    - 6.1.2 Patient has sex with men; and
    - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
    - 6.1.4 Any of the following:
      - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
      - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
      - 6.1.4.3 Patient has used methamphetamine in the last three months; or
  - 6.2 All of the following:
    - 6.2.1 Patient has a regular partner who has HIV infection; and
    - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
    - 6.2.3 Condoms have not been consistently used.

## Influenza

OSELTAMIVIR – **Restricted** see terms [below](#)

Note: The restriction on the use of oseltamivir to hospitalised patients means that supply into the community for a new course is not permitted. Supply of a part original pack on discharge where initiated as a hospital inpatient is permitted.

↓ Tab 75 mg

↓ Powder for oral suspension 6 mg per ml

→ **Restricted (RS1307)**

### Initiation

Either:

- 1 Only for hospitalised patient with known or suspected influenza; or
- 2 For prophylaxis of influenza in hospitalised patients as part of a DHB hospital approved infections control plan.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## ZANAMIVIR

Note: The restriction on the use of zanamivir to hospitalised patients means that supply into the community for a new course is not permitted. Supply of a part original pack on discharge where initiated as a hospital inpatient is permitted.

↓ Powder for inhalation 5 mg.....37.38 20 dose Relenza Rotadisk

→ **Restricted (RS1369)**

### Initiation

Either:

- 1 Only for hospitalised patient with known or suspected influenza; or
- 2 For prophylaxis of influenza in hospitalised patients as part of a DHB hospital approved infections control plan.

## Immune Modulators

### INTERFERON ALFA-2B

Inj 18 m iu, 1.2 ml multidose pen

Inj 30 m iu, 1.2 ml multidose pen

Inj 60 m iu, 1.2 ml multidose pen

INTERFERON GAMMA – **Restricted** see terms [below](#)

↓ Inj 100 mcg in 0.5 ml vial

→ **Restricted (RS1113)**

### Initiation

Patient has chronic granulomatous disease and requires interferon gamma.

PEGYLATED INTERFERON ALFA-2A – **Restricted** see terms [below](#)

↓ Inj 180 mcg prefilled syringe.....500.00 4 Pegasys

→ **Restricted (RS1782)**

**Initiation – Chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant**

*Limited to 48 weeks treatment*

Any of the following:

- 1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
- 2 Patient has chronic hepatitis C and is co-infected with HIV; or
- 3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant.

Notes: Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.

Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml.

**Continuation – Chronic hepatitis C - genotype 1 infection**

Gastroenterologist, infectious disease specialist or general physician

*Re-assessment required after 48 weeks*

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

**Initiation – Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior**

Gastroenterologist, infectious disease specialist or general physician

*Limited to 48 weeks treatment*

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; or
  - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir.

**Initiation – Chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV**

*Limited to 6 months treatment*

Patient has chronic hepatitis C, genotype 2 or 3 infection.

**Initiation – Hepatitis B**

Gastroenterologist, infectious disease specialist or general physician

*Limited to 48 weeks treatment*

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naïve; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log<sub>10</sub> IU/ml; and
- 5 Either:
  - 5.1 HBeAg positive; or
  - 5.2 Serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (greater than or equal to Metavir Stage F2 or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon.

Notes: Approved dose is 180 mcg once weekly.

The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.

In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon alfa-2a dose should be reduced to 135 mcg once weekly.

In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.

Pegylated Interferon alfa-2a is not approved for use in children.

**Initiation – myeloproliferative disorder or cutaneous T cell lymphoma**

*Re-assessment required after 12 months*

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma\*; or
- 2 All of the following:
  - 2.1 Patient has a myeloproliferative disorder\*; and
  - 2.2 Patient is intolerant of hydroxyurea; and
  - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
  - 3.1 Patient has a myeloproliferative disorder; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

3.2 Patient is pregnant, planning pregnancy or lactating.

**Continuation – myeloproliferative disorder or cutaneous T cell lymphoma**

*Re-assessment required after 12 months*

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:

3.1 Patient has a cutaneous T cell lymphoma\*; or

3.2 Both:

3.2.1 Patient has a myeloproliferative disorder\*; and

3.2.2 Either:

3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or

3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with \* are unapproved indications

**Initiation – ocular surface squamous neoplasia**

Ophthalmologist

*Re-assessment required after 12 months*

Patient has ocular surface squamous neoplasia\*.

**Continuation – ocular surface squamous neoplasia**

Ophthalmologist

*Re-assessment required after 12 months*

The treatment remains appropriate and patient is benefitting from treatment.

Note: Indications marked with \* are unapproved indications

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## Anticholinesterases

EDROPHONIUM CHLORIDE – **Restricted** see terms [below](#)

↓ Inj 10 mg per ml, 15 ml vial

↓ Inj 10 mg per ml, 1 ml ampoule

→ **Restricted** (RS1015)

### Initiation

For the diagnosis of myasthenia gravis.

NEOSTIGMINE METILSULFATE

Inj 2.5 mg per ml, 1 ml ampoule .....	98.00	50	AstraZeneca
---------------------------------------	-------	----	-------------

NEOSTIGMINE METILSULFATE WITH GLYCOPYRRONIUM BROMIDE

Inj 2.5 mg with glycopyrronium bromide 0.5 mg per ml, 1 ml ampoule .....	20.90	10	Max Health
--	-------	----	------------

PYRIDOSTIGMINE BROMIDE

Tab 60 mg – <b>1% DV Nov-19 to 2022</b> .....	45.79	100	<b>Mestinon</b>
---	-------	-----	-----------------

## Antirheumatoid Agents

HYDROXYCHLOROQUINE – **Restricted** see terms [below](#)

↓ Tab 200 mg – <b>1% DV Sep-18 to 2021</b> .....	7.98	100	<b>Plaquenil</b>
--	------	-----	------------------

→ **Restricted** (RS1776)

### Initiation

Any of the following:

- 1 Rheumatoid arthritis; or
- 2 Systemic or discoid lupus erythematosus; or
- 3 Malaria treatment or suppression; or
- 4 Relevant dermatological conditions (cutaneous forms of lupus and lichen planus, cutaneous vasculitides and mucosal ulceration); or
- 5 Sarcoidosis (pulmonary and non-pulmonary).

LEFLUNOMIDE

Tab 10 mg – <b>1% DV Dec-20 to 2023</b> .....	6.00	30	<b>Arava</b>
---	------	----	--------------

Tab 20 mg – <b>1% DV Dec-20 to 2023</b> .....	6.00	30	<b>Arava</b>
---	------	----	--------------

PENICILLAMINE

Tab 125 mg .....	67.23	100	D-Penamine
------------------	-------	-----	------------

Tab 250 mg .....	110.12	100	D-Penamine
------------------	--------	-----	------------

SODIUM AUROTHIOMALATE

Inj 10 mg in 0.5 ml ampoule

Inj 20 mg in 0.5 ml ampoule

Inj 50 mg in 0.5 ml ampoule

## Drugs Affecting Bone Metabolism

### Bisphosphonates

ALENDRONATE SODIUM

Tab 70 mg – <b>1% DV Apr-19 to 2022</b> .....	2.44	4	<b>Fosamax</b>
---	------	---	----------------

ALENDRONATE SODIUM WITH COLECALCIFEROL

Tab 70 mg with colecalciferol 5,600 iu – <b>1% DV Apr-19 to 2022</b> .....	1.51	4	<b>Fosamax Plus</b>
--	------	---	---------------------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PAMIDRONATE DISODIUM</b>			
Inj 3 mg per ml, 10 ml vial.....	5.98	1	Pamisol
Inj 6 mg per ml, 10 ml vial.....	15.02	1	Pamisol
Inj 9 mg per ml, 10 ml vial.....	17.05	1	Pamisol

<b>RISEDRONATE SODIUM</b>			
Tab 35 mg – 1% DV Oct-19 to 2022.....	3.10	4	<b>Risedronate Sandoz</b>

<b>ZOLEDRONIC ACID</b>			
⚡ Inj 5 mg per 100 ml, vial – 1% DV Oct-19 to 2022.....	60.00	100 ml	<b>Aclasta</b>

➡ **Restricted (RS1663)**

## Initiation – Inherited bone fragility disorders

Any specialist

Patient has been diagnosed with an inherited bone fragility disorder (e.g. osteogenesis imperfecta).

## Initiation – Osteoporosis

Any specialist

*Therapy limited to 3 doses*

Both:

- Any of the following:
  - History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
  - History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
  - History of two significant osteoporotic fractures demonstrated radiologically; or
  - Documented T-Score greater than or equal to -3.0 (see Note); or
  - A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
  - Patient has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

## Initiation – glucocorticosteroid therapy

Any specialist

*Re-assessment required after 12 months*

All of the following:

- The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- Any of the following:
  - The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or
  - The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
  - The patient has had a Special Authority approval for alendronate (Underlying cause - glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

## Continuation – glucocorticosteroid therapy

Any specialist

*Re-assessment required after 12 months*

Both:

- The patient is continuing systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone

continued...

	Price	Brand or
	(ex man. excl. GST)	Generic
	\$	Manufacturer
	Per	

continued...  
equivalents); and  
2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

**Initiation – Paget's disease**

Any specialist  
*Re-assessment required after 12 months*  
All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
  - 2.1 Bone or articular pain; or
  - 2.2 Bone deformity; or
  - 2.3 Bone, articular or neurological complications; or
  - 2.4 Asymptomatic disease, but risk of complications; or
  - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

**Continuation – Paget's disease**

Any specialist  
*Re-assessment required after 12 months*  
Both:

- 1 Any of the following:
  - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
  - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
  - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Notes:

- 1 BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- 2 Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- 3 Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- 4 A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

**Other Drugs Affecting Bone Metabolism**

DENOSUMAB – **Restricted** see terms [below](#)

⬇ Inj 60 mg prefilled syringe..... 326.00      1      Prolia  
➡ **Restricted (RS1665)**

**Initiation**

All of the following:

- 1 The patient has severe, established osteoporosis; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2 Either:
- 2.1 The patient is female and postmenopausal; or
  - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
- 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
  - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
  - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
  - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
  - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
  - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- 1 BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- 2 Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab.
- 3 Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- 4 A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- 5 Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialed so that the patient achieves the minimum requirement of 12 months' continuous therapy.

RALOXIFENE – **Restricted** see terms [below](#)

⚡ Tab 60 mg .....	53.76	28	Evista
-------------------	-------	----	--------

➡ **Restricted (RS1666)**

**Initiation**

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- History of two significant osteoporotic fractures demonstrated radiologically; or
- Documented T-Score greater than or equal to -3.0 (see Notes); or
- A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- Patient has had a Special Authority approval for zoledronic acid (Underlying cause - Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

#### Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

TERIPARATIDE – **Restricted** see terms [below](#)

↓ Inj 250 mcg per ml, 2.4 ml cartridge ..... 490.00      1      Forteo

→ **Restricted (RS1143)**

#### Initiation

Limited to 18 months treatment

All of the following:

- The patient has severe, established osteoporosis; and
- The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- The patient has had two or more fractures due to minimal trauma; and
- The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

#### Notes:

- The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- Antiresorptive agents and their adequate doses for the purposes of this restriction are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

## Enzymes

### HYALURONIDASE

Inj 1,500 iu ampoule

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Hyperuricaemia and Antigout</b>			
<b>ALLOPURINOL</b>			
Tab 100 mg – 1% DV Nov-20 to 2023 .....	11.47	500	<b>DP-Allopurinol</b>
Tab 300 mg – 1% DV Nov-20 to 2023 .....	28.57	500	<b>DP-Allopurinol</b>
<b>BENZBROMARONE – Restricted:</b> For continuation only			
➔ Tab 50 mg			
➔ Tab 100 mg .....	45.00	100	Benzbromaron AL 100
<b>COLCHICINE</b>			
Tab 500 mcg – 1% DV Jan-19 to 2021 .....	9.58	100	<b>Colgout</b>
<b>FEBUXOSTAT – Restricted</b> see terms <a href="#">below</a>			
⚡ Tab 80 mg .....	39.50	28	Adenuric
⚡ Tab 120 mg .....	39.50	28	Adenuric
➔ <b>Restricted (RS1760)</b>			
<b>Initiation</b>			
Any specialist			
Both:			
1 Patient has been diagnosed with gout; and			
2 Any of the following:			
2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or			
2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or			
2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or			
2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout.			
Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.			
<b>PROBENECID</b>			
Tab 500 mg			
<b>RASBURICASE – Restricted</b> see terms <a href="#">below</a>			
⚡ Inj 1.5 mg vial			
➔ <b>Restricted (RS1016)</b>			
Haematologist			
<b>Muscle Relaxants and Related Agents</b>			
<b>ATRAURIUM BESYLATE</b>			
Inj 10 mg per ml, 2.5 ml ampoule – 1% DV Jun-18 to 2021 .....	10.00	5	<b>Tracrium</b>
Inj 10 mg per ml, 5 ml ampoule – 1% DV Jun-18 to 2021 .....	12.50	5	<b>Tracrium</b>
<b>BACLOFEN</b>			
Tab 10 mg – 1% DV Oct-18 to 2021 .....	4.20	100	<b>Pacifen</b>
Oral liq 1 mg per ml			
Inj 0.05 mg per ml, 1 ml ampoule .....	11.55	1	Lioresal Intrathecal
Inj 2 mg per ml, 5 ml ampoule – 1% DV Apr-19 to 2021 .....	372.98	5	<b>Medsurge</b>

⚡ Item restricted (see ➔ above); ⚡ Item restricted (see ➔ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>CLOSTRIDIUM BOTULINUM TYPE A TOXIN</b>			
Inj 100 u vial .....	467.50	1	Botox
Inj 300 u vial .....	388.50	1	Dysport
Inj 500 u vial .....	1,295.00	2	Dysport
<b>DANTROLENE</b>			
Cap 25 mg .....	97.50	100	Dantrium
Cap 50 mg .....	77.00	100	Dantrium
Inj 20 mg vial .....	888.00	6	Dantrium IV
<b>MIVACURIUM CHLORIDE</b>			
Inj 2 mg per ml, 5 ml ampoule .....	33.92	5	Mivacron
Inj 2 mg per ml, 10 ml ampoule .....	67.17	5	Mivacron
<b>ORPHENADRINE CITRATE</b>			
Tab 100 mg – <b>1% DV Jun-18 to 2021</b> .....	18.54	100	<b>Norflex</b>
<b>PANCURONIUM BROMIDE</b>			
Inj 2 mg per ml, 2 ml ampoule			
<b>ROCURONIUM BROMIDE</b>			
Inj 10 mg per ml, 5 ml ampoule – <b>1% DV Aug-20 to 2022</b> .....	31.14	10	<b>Hameln</b>
<b>SUXAMETHONIUM CHLORIDE</b>			
Inj 50 mg per ml, 2 ml ampoule – <b>1% DV Feb-21 to 2023</b> .....	78.00	50	AstraZeneca
	23.40	10	<b>Martindale</b>
<i>(AstraZeneca Inj 50 mg per ml, 2 ml ampoule to be delisted 1 February 2021)</i>			
<b>VECURIONIUM BROMIDE</b>			
Inj 10 mg vial			

### Reversers of Neuromuscular Blockade

**SUGAMMADEX – Restricted** see terms [below](#)

⚡ Inj 100 mg per ml, 2 ml vial .....	1,200.00	10	Bridion
⚡ Inj 100 mg per ml, 5 ml vial .....	3,000.00	10	Bridion

➡ **Restricted (RS1370)**

#### Initiation

Any of the following:

- 1 Patient requires reversal of profound neuromuscular blockade following rapid sequence induction that has been undertaken using rocuronium (i.e. suxamethonium is contraindicated or undesirable); or
- 2 Severe neuromuscular degenerative disease where the use of neuromuscular blockade is required; or
- 3 Patient has an unexpectedly difficult airway that cannot be intubated and requires a rapid reversal of anaesthesia and neuromuscular blockade; or
- 4 The duration of the patient's surgery is unexpectedly short; or
- 5 Neostigmine or a neostigmine/anticholinergic combination is contraindicated (for example the patient has ischaemic heart disease, morbid obesity or COPD); or
- 6 Patient has a partial residual block after conventional reversal.

### Non-Steroidal Anti-Inflammatory Drugs

<b>CELECOXIB</b>			
Cap 100 mg .....	3.63	60	Celecoxib Pfizer
Cap 200 mg .....	2.30	30	Celecoxib Pfizer

# MUSCULOSKELETAL SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>DICLOFENAC SODIUM</b>			
Tab EC 25 mg – 1% DV Oct-18 to 2021 .....	1.23	50	<b>Diclofenac Sandoz</b>
Tab 50 mg dispersible .....	1.50	20	Voltaren D
Tab EC 50 mg – 1% DV Oct-18 to 2021 .....	1.23	50	<b>Diclofenac Sandoz</b>
Tab long-acting 75 mg – 1% DV Oct-18 to 2021 .....	22.80	500	<b>Apo-Diclo SR</b>
Tab long-acting 100 mg – 1% DV Oct-18 to 2021 .....	25.15	500	<b>Apo-Diclo SR</b>
Inj 25 mg per ml, 3 ml ampoule .....	13.20	5	Voltaren
Suppos 12.5 mg .....	2.04	10	Voltaren
Suppos 25 mg .....	2.44	10	Voltaren
Suppos 50 mg .....	4.22	10	Voltaren
Suppos 100 mg .....	7.00	10	Voltaren
<b>ETORICOXIB – Restricted</b> see terms <a href="#">below</a>			
⚡ Tab 30 mg			
⚡ Tab 60 mg			
⚡ Tab 90 mg			
⚡ Tab 120 mg			
➡ <b>Restricted (RS1290)</b>			
<b>Initiation</b>			
For in-vivo investigation of allergy only.			
<b>IBUPROFEN</b>			
Tab 200 mg .....	11.71	1,000	Relieve
➡ Tab 400 mg – <b>Restricted:</b> For continuation only			
➡ Tab 600 mg – <b>Restricted:</b> For continuation only			
Tab long-acting 800 mg – 1% DV Apr-20 to 2021 .....	5.99	30	<b>Ibuprofen SR BNM</b>
Oral liq 20 mg per ml – 1% DV May-19 to 2021 .....	1.88	200 ml	<b>Ethics</b>
Inj 5 mg per ml, 2 ml ampoule			
Inj 10 mg per ml, 2 ml vial			
<b>INDOMETHACIN</b>			
Cap 25 mg			
Cap 50 mg			
Cap long-acting 75 mg			
Inj 1 mg vial			
Suppos 100 mg			
<b>KETOPROFEN</b>			
Cap long-acting 200 mg .....	12.07	28	Oruvail SR
<b>MEFENAMIC ACID – Restricted:</b> For continuation only			
➡ Cap 250 mg			
<b>NAPROXEN</b>			
Tab 250 mg – 1% DV Dec-18 to 2021 .....	32.69	500	<b>Noflam 250</b>
Tab 500 mg – 1% DV Dec-18 to 2021 .....	22.19	250	<b>Noflam 500</b>
Tab long-acting 750 mg – 1% DV Oct-18 to 2021 .....	6.16	28	<b>Naprosyn SR 750</b>
Tab long-acting 1 g – 1% DV Oct-18 to 2021 .....	8.21	28	<b>Naprosyn SR 1000</b>
<b>PARECOXIB</b>			
Inj 40 mg vial .....	100.00	10	Dynastat
<b>SULINDAC</b>			
Tab 100 mg			
Tab 200 mg			
<b>TENOXICAM</b>			
Tab 20 mg – 1% DV Oct-19 to 2022 .....	9.15	100	<b>Tilcotil</b>
Inj 20 mg vial .....	9.95	1	<b>AFT</b>

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

Topical Products for Joint and Muscular Pain

CAPSAICIN – **Restricted** see terms [below](#)

↓ Crm 0.025%.....9.95 45 g Zostrix

➡ **Restricted** ([RS1309](#))

Initiation

Patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## Agents for Parkinsonism and Related Disorders

### Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE – **Restricted** see terms [below](#)

↓ Tab 50 mg – **1% DV Aug-18 to 2021**..... 130.00 56 **Rilutek**  
 → **Restricted (RS1351)**

#### Initiation

Neurologist or respiratory specialist

*Re-assessment required after 6 months*

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
  - 5.1 The patient is ambulatory; or
  - 5.2 The patient is able to use upper limbs; or
  - 5.3 The patient is able to swallow.

#### Continuation

*Re-assessment required after 18 months*

All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
  - 3.1 The patient is ambulatory; or
  - 3.2 The patient is able to use upper limbs; or
  - 3.3 The patient is able to swallow.

TETRABENAZINE

Tab 25 mg – **1% DV Oct-19 to 2022**.....91.10 112 **Motetis**

### Anticholinergics

BENZATROPINE MESYLATE

Tab 2 mg ..... 7.99 60 **Benztrop**  
 Inj 1 mg per ml, 2 ml ampoule – **1% DV Dec-20 to 2023**.....95.00 5 **Phebra**

PROCYCLIDINE HYDROCHLORIDE

Tab 5 mg

### Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE

Cap 100 mg .....38.24 60 **Symmetrel**

APOMORPHINE HYDROCHLORIDE

Inj 10 mg per ml, 2 ml ampoule – **1% DV Jan-20 to 2023** .....59.50 5 **Movapo**  
 Inj 10 mg per ml, 5 ml ampoule – **1% DV Feb-20 to 2023** ..... 121.84 5 **Movapo**

BROMOCRIPTINE

Tab 2.5 mg  
 Cap 5 mg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>ENTACAPONE</b>			
Tab 200 mg – 1% DV Sep-18 to 2021 .....	22.00	100	<b>Entapone</b>
<b>LEVODOPA WITH BENSERAZIDE</b>			
Tab dispersible 50 mg with benserazide 12.5 mg .....	13.25	100	Madopar Rapid
Cap 50 mg with benserazide 12.5 mg .....	13.75	100	Madopar 62.5
Cap 100 mg with benserazide 25 mg .....	15.80	100	Madopar 125
Cap long-acting 100 mg with benserazide 25 mg .....	22.85	100	Madopar HBS
Cap 200 mg with benserazide 50 mg .....	26.25	100	Madopar 250
<b>LEVODOPA WITH CARBIDOPA</b>			
Tab 100 mg with carbidopa 25 mg – 1% DV Dec-20 to 2023 .....	21.11	100	<b>Sinemet</b>
Tab long-acting 100 mg with carbidopa 25 mg .....			
Tab long-acting 200 mg with carbidopa 50 mg – 1% DV Feb-21 to 2023 .....	43.65	100	<b>Sinemet CR</b>
Tab 250 mg with carbidopa 25 mg – 1% DV Dec-20 to 2023 .....	38.39	100	<b>Sinemet</b>
<b>PRAMIPEXOLE HYDROCHLORIDE</b>			
Tab 0.25 mg – 1% DV Oct-19 to 2022 .....	6.12	100	<b>Ramipex</b>
Tab 1 mg – 1% DV Oct-19 to 2022 .....	20.73	100	<b>Ramipex</b>
<b>ROPINIROLE HYDROCHLORIDE</b>			
Tab 0.25 mg – 1% DV Mar-20 to 2022 .....	2.85	84	<b>Ropin</b>
Tab 1 mg – 1% DV Mar-20 to 2022 .....	3.95	84	<b>Ropin</b>
Tab 2 mg – 1% DV Mar-20 to 2022 .....	5.48	84	<b>Ropin</b>
Tab 5 mg – 1% DV Mar-20 to 2022 .....	12.50	84	<b>Ropin</b>
<b>SELEGILINE HYDROCHLORIDE</b>			
Tab 5 mg .....			
<b>TOLCAPONE</b>			
Tab 100 mg .....	152.38	100	Tasmar

## Anaesthetics

### General Anaesthetics

<b>DESFLURANE</b>			
Soln for inhalation 100%, 240 ml bottle .....	1,350.00	6	Suprane
<b>DEXMEDETOMIDINE</b>			
Inj 100 mcg per ml, 2 ml vial – 1% DV Mar-21 to 2023 .....	97.88	5	<b>Dexmedetomidine-Teva</b>
	357.00		Precedex
<i>(Precedex Inj 100 mcg per ml, 2 ml vial to be delisted 1 March 2021)</i>			
<b>ETOMIDATE</b>			
Inj 2 mg per ml, 10 ml ampoule .....			
<b>ISOFLURANE</b>			
Soln for inhalation 100%, 250 ml bottle .....	1,020.00	6	Aerrane
<b>KETAMINE</b>			
Inj 1 mg per ml, 100 ml bag – 1% DV Feb-20 to 2022 .....	135.00	5	<b>Biomed</b>
Inj 10 mg per ml, 10 ml syringe – 1% DV Feb-20 to 2022 .....	70.00	5	<b>Biomed</b>
Inj 100 mg per ml, 2 ml ampoule .....	155.60	5	Ketamine-Baxter
Inj 100 mg per ml, 2 ml vial – 1% DV Jan-19 to 2021 .....	31.50	5	<b>Ketalar</b>
	155.60		Ketamine-Claris
<i>(Ketamine-Claris Inj 100 mg per ml, 2 ml vial to be delisted 1 March 2021)</i>			
<b>METHOHEXITAL SODIUM</b>			
Inj 10 mg per ml, 50 ml vial .....			

## NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PROPOFOL</b>			
Inj 10 mg per ml, 20 ml ampoule – <b>10% DV Dec-19 to 2022</b> .....	4.35	5	<b>Fresofol 1% MCT/LCT</b>
Inj 10 mg per ml, 50 ml vial – <b>10% DV Oct-19 to 2022</b> .....	19.50	10	<b>Fresofol 1% MCT/LCT</b>
Inj 10 mg per ml, 100 ml vial – <b>10% DV Oct-19 to 2022</b> .....	39.00	10	<b>Fresofol 1% MCT/LCT</b>
<b>SEVOFLURANE</b>			
Soln for inhalation 100%, 250 ml bottle.....	840.00	6	Baxter
<b>THIOPENTAL [THIOPENTONE] SODIUM</b>			
Inj 500 mg ampoule			
<b>Local Anaesthetics</b>			
<b>ARTICAINE HYDROCHLORIDE</b>			
Inj 1%			
<b>ARTICAINE HYDROCHLORIDE WITH ADRENALINE</b>			
Inj 4% with adrenaline 1:100,000, 1.7 ml dental cartridge			
Inj 4% with adrenaline 1:100,000, 2.2 ml dental cartridge			
Inj 4% with adrenaline 1:200,000, 1.7 ml dental cartridge			
Inj 4% with adrenaline 1:200,000, 2.2 ml dental cartridge			
<b>BENZOCAINE</b>			
Gel 20%			
<b>BENZOCAINE WITH TETRACAINE HYDROCHLORIDE</b>			
Gel 18% with tetracaine hydrochloride 2%			<i>e.g. ZAP Topical Anaesthetic Gel</i>
<b>BUPIVACAINE HYDROCHLORIDE</b>			
Inj 5 mg per ml, 4 ml ampoule – <b>1% DV Oct-20 to 2023</b> .....	50.00	5	<b>Marcaïn Isobaric</b>
Inj 2.5 mg per ml, 20 ml ampoule			
Inj 2.5 mg per ml, 20 ml ampoule sterile pack – <b>1% DV Aug-20 to 2023</b> .....	23.36	5	<b>Marcaïn</b>
Inj 5 mg per ml, 10 ml ampoule sterile pack – <b>1% DV Aug-20 to 2023</b> .....	16.20	5	<b>Marcaïn</b>
Inj 5 mg per ml, 20 ml ampoule			
Inj 5 mg per ml, 20 ml ampoule sterile pack – <b>1% DV Aug-20 to 2023</b> .....	16.56	5	<b>Marcaïn</b>
Inj 1.25 mg per ml, 100 ml bag			
Inj 1.25 mg per ml, 200 ml bag			
Inj 2.5 mg per ml, 100 ml bag – <b>1% DV Oct-20 to 2023</b> .....	150.00	5	<b>Marcaïn</b>
Inj 2.5 mg per ml, 200 ml bag			
Inj 1.25 mg per ml, 500 ml bag			
<b>BUPIVACAINE HYDROCHLORIDE WITH ADRENALINE</b>			
Inj 2.5 mg per ml with adrenaline 1:400,000, 20 ml vial – <b>1% DV Aug-19 to 2022</b> .....	94.50	5	<b>Marcaïn with Adrenaline</b>
Inj 5 mg per ml with adrenaline 1:200,000, 20 ml vial – <b>1% DV Aug-19 to 2022</b> .....	80.50	5	<b>Marcaïn with Adrenaline</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>BUPIVACAINE HYDROCHLORIDE WITH FENTANYL</b>			
Inj 0.625 mg with fentanyl 2 mcg per ml, 100 ml bag			
Inj 0.625 mg with fentanyl 2 mcg per ml, 200 ml bag – <b>1% DV Apr-20 to 2022</b>	152.50	5	<b>Biomed</b>
Inj 1.25 mg with fentanyl 2 mcg per ml, 100 ml syringe			
Inj 1.25 mg with fentanyl 2 mcg per ml, 100 ml bag – <b>1% DV Nov-19 to 2022</b>	112.50	5	<b>Bupafen</b>
Inj 1.25 mg with fentanyl 2 mcg per ml, 200 ml bag – <b>1% DV Nov-19 to 2022</b>	117.50	5	<b>Bupafen</b>
Inj 1.25 mg with fentanyl 2 mcg per ml, 50 ml syringe			
Inj 1.25 mg with fentanyl 2 mcg per ml, 15 ml syringe	36.00	5	Biomed
Inj 1.25 mg with fentanyl 2 mcg per ml, 20 ml syringe	46.00	5	Biomed
<b>BUPIVACAINE HYDROCHLORIDE WITH GLUCOSE</b>			
Inj 0.5% with glucose 8%, 4 ml ampoule	38.00	5	Marcaïn Heavy
<b>COCAINE HYDROCHLORIDE</b>			
Paste 5%			
Soln 15%, 2 ml syringe			
Soln 4%, 2 ml syringe	25.46	1	Biomed
<b>COCAINE HYDROCHLORIDE WITH ADRENALINE</b>			
Paste 15% with adrenaline 0.06%			
Paste 25% with adrenaline 0.06%			
<b>ETHYL CHLORIDE</b>			
Spray 100%			
<b>LIDOCAINE [LIGNOCAINE]</b>			
Crm 4%	5.40	5 g	LMX4
	27.00	30 g	LMX4
<b>LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE</b>			
Gel 2% – <b>1% DV Nov-18 to 2021</b>	4.87	20 g	<b>Orion</b>
Soln 4%			
Spray 10% – <b>1% DV Jul-19 to 2022</b>	75.00	50 ml	<b>Xylocaine</b>
Oral (gel) soln 2%	38.00	200 ml	Mucosoothie
Inj 1%, 20 ml ampoule, sterile pack			
Inj 2%, 20 ml ampoule, sterile pack			
Inj 1%, 5 ml ampoule	8.75	25	Lidocaine-Clarís
Inj 1%, 20 ml vial – <b>1% DV Jul-19 to 2022</b>	6.20	5	<b>Lidocaine-Clarís</b>
Inj 2%, 5 ml ampoule – <b>1% DV Nov-19 to 2022</b>	8.25	25	<b>Lidocaine-Clarís</b>
Inj 2%, 20 ml vial – <b>1% DV Jul-19 to 2022</b>	6.45	5	<b>Lidocaine-Clarís</b>
Gel 2%, 11 ml urethral syringe – <b>1% DV Apr-20 to 2022</b>	42.00	10	<b>Instillagel Lido</b>
<b>LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH ADRENALINE</b>			
Inj 1% with adrenaline 1:100,000, 5 ml ampoule – <b>1% DV Nov-19 to 2022</b>	29.00	10	<b>Xylocaine</b>
Inj 1% with adrenaline 1:200,000, 20 ml vial	50.00	5	Xylocaine
Inj 2% with adrenaline 1:80,000, 1.7 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 1.8 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 2.2 ml dental cartridge			
Inj 2% with adrenaline 1:200,000, 20 ml vial	60.00	5	Xylocaine
<b>LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH ADRENALINE AND TETRACAINE HYDROCHLORIDE</b>			
Soln 4% with adrenaline 0.1% and tetracaine hydrochloride 0.5%, 5 ml syringe	17.50	1	Topicaïne

## NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH CHLORHEXIDINE			
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringe .....	81.50	10	Pfizer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH PHENYLEPHRINE HYDROCHLORIDE			
Nasal spray 5% with phenylephrine hydrochloride 0.5%			
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE			
Crm 2.5% with prilocaine 2.5%.....	45.00	30 g	EMLA
Patch 25 mcg with prilocaine 25 mcg.....	115.00	20	EMLA
Crm 2.5% with prilocaine 2.5%, 5 g.....	45.00	5	EMLA
MEPIVACAINE HYDROCHLORIDE			
Inj 3%, 1.8 ml dental cartridge .....	43.60	50	Scandonest 3%
Inj 3%, 2.2 ml dental cartridge .....	43.60	50	Scandonest 3%
PRILOCAINE HYDROCHLORIDE			
Inj 0.5%, 50 ml vial .....	100.00	5	Citanest
Inj 2%, 5 ml ampoule			
PRILOCAINE HYDROCHLORIDE WITH FELYPRESSIN			
Inj 3% with felypressin 0.03 iu per ml, 1.8 ml dental cartridge			
Inj 3% with felypressin 0.03 iu per ml, 2.2 ml dental cartridge			
ROPIVACAINE HYDROCHLORIDE			
Inj 2 mg per ml, 10 ml ampoule – 1% DV Nov-20 to 2023.....	9.25	5	Ropivacaine Kabi
Inj 2 mg per ml, 20 ml ampoule – 1% DV Nov-20 to 2023.....	9.65	5	Ropivacaine Kabi
Inj 2 mg per ml, 100 ml bag – 1% DV Nov-20 to 2023 .....	31.00	5	Ropivacaine Kabi
Inj 2 mg per ml, 200 ml bag – 1% DV Nov-20 to 2023 .....	40.95	5	Ropivacaine Kabi
Inj 7.5 mg per ml, 10 ml ampoule – 1% DV Nov-20 to 2023.....	10.40	5	Ropivacaine Kabi
Inj 7.5 mg per ml, 20 ml ampoule – 1% DV Nov-20 to 2023.....	12.75	5	Ropivacaine Kabi
Inj 10 mg per ml, 10 ml ampoule – 1% DV Nov-20 to 2023.....	11.10	5	Ropivacaine Kabi
Inj 10 mg per ml, 20 ml ampoule – 1% DV Nov-20 to 2023.....	16.60	5	Ropivacaine Kabi
ROPIVACAINE HYDROCHLORIDE WITH FENTANYL			
Inj 2 mg with fentanyl 2 mcg per ml, 100 ml bag.....	198.50	5	Naropin
Inj 2 mg with fentanyl 2 mcg per ml, 200 ml bag.....	270.00	5	Naropin
TETRACAINE [AMETHOCAINE] HYDROCHLORIDE			
Gel 4%			

## Analgesics

### Non-Opioid Analgesics

ASPIRIN			
Tab dispersible 300 mg – 1% DV Oct-19 to 2022 .....	4.50	100	Ethics Aspirin
CAPSAICIN – <b>Restricted</b> see terms <a href="#">below</a>			
⚡ Crm 0.075%.....	12.50	45 g	Zostrix HP
➡ <b>Restricted (RS1145)</b>			
<b>Initiation</b>			
For post-herpetic neuralgia or diabetic peripheral neuropathy.			
METHOXYFLURANE – <b>Restricted</b> see terms <a href="#">below</a>			
⚡ Soln for inhalation 99.9%, 3 ml bottle			
➡ <b>Restricted (RS1292)</b>			
<b>Initiation</b>			
Both:			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient is undergoing a painful procedure with an expected duration of less than one hour; and
- 2 Only to be used under supervision by a medical practitioner or nurse who is trained in the use of methoxyflurane.

#### NEFOPAM HYDROCHLORIDE

Tab 30 mg

#### PARACETAMOL – Some items restricted see terms [below](#)

Tab soluble 500 mg

Tab 500 mg

Oral liq 120 mg per 5 ml – **20% DV Nov-20 to 2023**.....5.45

1,000 ml

**Paracare**

Oral liq 250 mg per 5 ml – **20% DV Nov-20 to 2023**.....6.25

1,000 ml

**Paracare Double  
Strength**

↓ Inj 10 mg per ml, 100 ml vial – **1% DV Nov-20 to 2023**.....8.90

10

**Paracetamol Kabi**

Suppos 25 mg – **1% DV Nov-19 to 2022**.....58.50

20

**Biomed**

Suppos 50 mg – **1% DV Nov-19 to 2022**.....58.50

20

**Biomed**

Suppos 125 mg – **1% DV Nov-18 to 2021**.....3.29

10

**Gacet**

Suppos 250 mg – **1% DV Nov-18 to 2021**.....3.79

10

**Gacet**

Suppos 500 mg – **1% DV Feb-19 to 2021**.....12.40

50

**Gacet**

➔ **Restricted (RS1146)**

#### Initiation

Intravenous paracetamol is only to be used where other routes are unavailable or impractical, or where there is reduced absorption. The need for IV paracetamol must be re-assessed every 24 hours.

#### SUCROSE

Oral liq 25% – **1% DV Feb-20 to 2022**.....13.00

25 ml

**Biomed**

↓ Oral liq 66.7% (preservative free)

➔ **Restricted (RS1763)**

#### Initiation

For use in neonatal patients only.

## Opioid Analgesics

#### ALFENTANIL

Inj 0.5 mg per ml, 2 ml ampoule – **1% DV Nov-20 to 2023**.....24.75

10

**Hameln**

#### CODEINE PHOSPHATE

Tab 15 mg – **1% DV Nov-20 to 2023**.....6.25

100

**PSM**

Tab 30 mg – **1% DV Nov-20 to 2023**.....7.45

100

**PSM**

Tab 60 mg – **1% DV Nov-20 to 2023**.....14.25

100

**PSM**

#### DIHYDROCODEINE TARTRATE

Tab long-acting 60 mg – **1% DV Oct-19 to 2022**.....8.60

60

**DHC Continus**

## NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>FENTANYL</b>			
Inj 10 mcg per ml, 10 ml syringe			
Inj 50 mcg per ml, 2 ml ampoule – 1% DV Nov-18 to 2021	3.56	10	<b>Boucher and Muir</b>
Inj 10 mcg per ml, 50 ml bag	210.00	10	Biomed
Inj 10 mcg per ml, 50 ml syringe	165.00	10	Biomed
Inj 50 mcg per ml, 10 ml ampoule – 1% DV Nov-18 to 2021	9.41	10	<b>Boucher and Muir</b>
Inj 10 mcg per ml, 100 ml bag – 1% DV Nov-19 to 2022	110.00	5	<b>Biomed</b>
Inj 20 mcg per ml, 50 ml syringe – 1% DV Oct-18 to 2021	18.74	1	<b>Biomed</b>
Inj 20 mcg per ml, 100 ml bag			
Patch 12.5 mcg per hour	2.95	5	Fentanyl Sandoz
Patch 25 mcg per hour	3.66	5	Fentanyl Sandoz
Patch 50 mcg per hour	6.65	5	Fentanyl Sandoz
Patch 75 mcg per hour	9.25	5	Fentanyl Sandoz
Patch 100 mcg per hour	11.40	5	Fentanyl Sandoz
<b>METHADONE HYDROCHLORIDE</b>			
Tab 5 mg – 1% DV Sep-19 to 2022	1.40	10	<b>Methatabs</b>
Oral liq 2 mg per ml – 1% DV Oct-18 to 2021	5.79	200 ml	<b>Biodone</b>
Oral liq 5 mg per ml – 1% DV Oct-18 to 2021	5.79	200 ml	<b>Biodone Forte</b>
Oral liq 10 mg per ml – 1% DV Oct-18 to 2021	6.79	200 ml	<b>Biodone Extra Forte</b>
Inj 10 mg per ml, 1 ml vial	61.00	10	AFT
<b>MORPHINE HYDROCHLORIDE</b>			
Oral liq 1 mg per ml – 1% DV Dec-18 to 2021	9.28	200 ml	<b>RA-Morph</b>
Oral liq 2 mg per ml – 1% DV Dec-18 to 2021	16.24	200 ml	<b>RA-Morph</b>
Oral liq 5 mg per ml – 1% DV Dec-18 to 2021	19.44	200 ml	<b>RA-Morph</b>
Oral liq 10 mg per ml – 1% DV Dec-18 to 2021	27.74	200 ml	<b>RA-Morph</b>
<b>MORPHINE SULPHATE</b>			
Tab immediate-release 10 mg – 1% DV Nov-20 to 2023	2.80	10	<b>Sevredol</b>
Tab immediate-release 20 mg – 1% DV Nov-20 to 2023	5.52	10	<b>Sevredol</b>
Tab long-acting 30 mg	2.85	10	Arrow-Morphine LA
Tab long-acting 60 mg	5.60	10	Arrow-Morphine LA
Cap long-acting 10 mg – 1% DV Jan-20 to 2022	2.05	10	<b>m-Eslon</b>
Cap long-acting 30 mg – 1% DV Jan-20 to 2022	3.00	10	<b>m-Eslon</b>
Cap long-acting 60 mg – 1% DV Jan-20 to 2022	6.12	10	<b>m-Eslon</b>
Cap long-acting 100 mg – 1% DV Jan-20 to 2022	7.13	10	<b>m-Eslon</b>
Inj 1 mg per ml, 100 ml bag – 1% DV Nov-20 to 2023	102.25	5	<b>Biomed</b>
Inj 1 mg per ml, 10 ml syringe – 1% DV Nov-20 to 2023	24.50	5	<b>Biomed</b>
Inj 1 mg per ml, 50 ml syringe – 1% DV Nov-20 to 2023	52.00	5	<b>Biomed</b>
Inj 1 mg per ml, 2 ml syringe			
Inj 2 mg per ml, 30 ml syringe	135.00	10	Biomed
Inj 5 mg per ml, 1 ml ampoule	6.27	5	DBL Morphine Sulphate
Inj 10 mg per ml, 1 ml ampoule	4.47	5	DBL Morphine Sulphate
Inj 10 mg per ml, 100 mg cassette			
Inj 10 mg per ml, 100 ml bag			
Inj 15 mg per ml, 1 ml ampoule	4.76	5	DBL Morphine Sulphate
Inj 30 mg per ml, 1 ml ampoule	6.19	5	DBL Morphine Sulphate
Inj 200 mcg in 0.4 ml syringe			
Inj 300 mcg in 0.3 ml syringe			
<i>(Arrow-Morphine LA Tab long-acting 60 mg to be delisted 1 April 2021)</i>			
<b>MORPHINE TARTRATE</b>			
Inj 80 mg per ml, 1.5 ml ampoule			

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>OXYCODONE HYDROCHLORIDE</b>			
Tab controlled-release 5 mg – <b>1% DV May-19 to 2021</b> .....	2.15	20	<b>Oxycodone Sandoz</b>
Tab controlled-release 10 mg – <b>1% DV May-19 to 2021</b> .....	2.15	20	<b>Oxycodone Sandoz</b>
Tab controlled-release 20 mg – <b>1% DV May-19 to 2021</b> .....	2.15	20	<b>Oxycodone Sandoz</b>
Tab controlled-release 40 mg – <b>1% DV May-19 to 2021</b> .....	3.20	20	<b>Oxycodone Sandoz</b>
Tab controlled-release 80 mg – <b>1% DV May-19 to 2021</b> .....	10.98	20	<b>Oxycodone Sandoz</b>
Cap immediate-release 5 mg – <b>1% DV Sep-18 to 2021</b> .....	1.88	20	<b>OxyNorm</b>
Cap immediate-release 10 mg – <b>1% DV Sep-18 to 2021</b> .....	3.32	20	<b>OxyNorm</b>
Cap immediate-release 20 mg – <b>1% DV Sep-18 to 2021</b> .....	5.81	20	<b>OxyNorm</b>
Oral liq 5 mg per 5 ml .....	11.20	250 ml	<b>OxyNorm</b>
Inj 1 mg per ml, 100 ml bag			
Inj 10 mg per ml, 1 ml ampoule – <b>1% DV Sep-18 to 2021</b> .....	7.28	5	<b>OxyNorm</b>
Inj 10 mg per ml, 2 ml ampoule – <b>1% DV Sep-18 to 2021</b> .....	14.36	5	<b>OxyNorm</b>
Inj 50 mg per ml, 1 ml ampoule – <b>1% DV Sep-18 to 2021</b> .....	30.60	5	<b>OxyNorm</b>
<b>PARACETAMOL WITH CODEINE</b>			
Tab paracetamol 500 mg with codeine phosphate 8 mg .....	18.21	1,000	Paracetamol + Codeine (Relieve)
<b>PETHIDINE HYDROCHLORIDE</b>			
Tab 50 mg – <b>1% DV Sep-18 to 2021</b> .....	4.46	10	<b>PSM</b>
Inj 5 mg per ml, 10 ml syringe			
Inj 5 mg per ml, 100 ml bag			
Inj 10 mg per ml, 100 ml bag			
Inj 10 mg per ml, 50 ml syringe			
Inj 50 mg per ml, 1 ml ampoule .....	4.98	5	DBL Pethidine Hydrochloride
Inj 50 mg per ml, 2 ml ampoule .....	5.12	5	DBL Pethidine Hydrochloride
<b>REMIFENTANIL</b>			
Inj 1 mg vial – <b>1% DV Oct-20 to 2023</b> .....	13.95	5	<b>Remifentanil-AFT</b>
Inj 2 mg vial – <b>1% DV Oct-20 to 2023</b> .....	19.95	5	<b>Remifentanil-AFT</b>
<b>TRAMADOL HYDROCHLORIDE</b>			
Tab sustained-release 100 mg – <b>1% DV Nov-20 to 2023</b> .....	1.52	20	<b>Tramal SR 100</b>
Tab sustained-release 150 mg – <b>1% DV Nov-20 to 2023</b> .....	2.10	20	<b>Tramal SR 150</b>
Tab sustained-release 200 mg – <b>1% DV Nov-20 to 2023</b> .....	2.75	20	<b>Tramal SR 200</b>
Cap 50 mg – <b>1% DV Dec-20 to 2023</b> .....	2.80	100	<b>Arrow-Tramadol</b>
Oral soln 10 mg per ml			
Inj 10 mg per ml, 100 ml bag			
Inj 50 mg per ml, 1 ml ampoule – <b>1% DV Oct-20 to 2023</b> .....	4.50	5	<b>Tramal 50</b>
Inj 50 mg per ml, 2 ml ampoule – <b>1% DV Oct-20 to 2023</b> .....	3.83	5	<b>Tramal 100</b>

## Antidepressants

### Cyclic and Related Agents

<b>AMITRIPTYLINE</b>			
Tab 10 mg – <b>1% DV Dec-20 to 2023</b> .....	2.49	100	<b>Arrow-Amitriptyline</b>
Tab 25 mg – <b>1% DV Dec-20 to 2023</b> .....	1.51	100	<b>Arrow-Amitriptyline</b>
Tab 50 mg – <b>1% DV Dec-20 to 2023</b> .....	2.51	100	<b>Arrow-Amitriptyline</b>
<b>CLOMIPRAMINE HYDROCHLORIDE</b>			
Tab 10 mg – <b>1% DV Oct-18 to 2021</b> .....	13.99	100	<b>Apo-Clomipramine</b>
Tab 25 mg – <b>1% DV Oct-18 to 2021</b> .....	9.46	100	<b>Apo-Clomipramine</b>

## NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – <b>Restricted:</b> For continuation only			
➡ Cap 25 mg .....	7.83	50	Dosulepin Mylan
DOXEPIN HYDROCHLORIDE – <b>Restricted:</b> For continuation only			
➡ Cap 10 mg			
➡ Cap 25 mg			
➡ Cap 50 mg			
IMIPRAMINE HYDROCHLORIDE			
Tab 10 mg .....	5.48	50	Tofranil
	6.58	60	Tofranil
Tab 25 mg .....	8.80	50	Tofranil
MAPROTILINE HYDROCHLORIDE – <b>Restricted:</b> For continuation only			
➡ Tab 25 mg			
➡ Tab 75 mg			
MIANSERIN HYDROCHLORIDE – <b>Restricted:</b> For continuation only			
➡ Tab 30 mg			
NORTRIPTYLINE HYDROCHLORIDE			
Tab 10 mg – <b>1% DV Oct-19 to 2022</b> .....	2.44	100	Norpress
Tab 25 mg – <b>1% DV Oct-19 to 2022</b> .....	5.98	180	Norpress

### Monoamine-Oxidase Inhibitors - Non-Selective

PHENELZINE SULPHATE			
Tab 15 mg			
TRANLYCYPROMINE SULPHATE			
Tab 10 mg			

### Monoamine-Oxidase Type A Inhibitors

MOCLOBEMIDE			
Tab 150 mg – <b>1% DV Apr-19 to 2021</b> .....	6.40	60	Aurorix
Tab 300 mg – <b>1% DV Apr-19 to 2021</b> .....	9.80	60	Aurorix

### Other Antidepressants

MIRTAZAPINE			
Tab 30 mg – <b>1% DV Oct-18 to 2021</b> .....	2.63	30	Apo-Mirtazapine
Tab 45 mg – <b>1% DV Oct-18 to 2021</b> .....	3.48	30	Apo-Mirtazapine
VENLAFAXINE			
Cap 37.5 mg .....	6.38	84	Enlafax XR
Cap 75 mg .....	8.11	84	Enlafax XR
Cap 150 mg .....	11.16	84	Enlafax XR

### Selective Serotonin Reuptake Inhibitors

CITALOPRAM HYDROBROMIDE			
Tab 20 mg – <b>1% DV Sep-18 to 2021</b> .....	1.52	84	PSM Citalopram
ESCITALOPRAM			
Tab 10 mg .....	1.11	28	Escitalopram-Apotex
Tab 20 mg .....	1.90	28	Escitalopram-Apotex

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>FLUOXETINE HYDROCHLORIDE</b>			
Tab dispersible 20 mg, scored – <b>1% DV Feb-21 to 2022</b> .....	9.93	30	Arrow-Fluoxetine
	1.98		<b>Fluox</b>
Cap 20 mg – <b>1% DV Feb-21 to 2022</b> .....	7.49	90	Arrow-Fluoxetine
	2.91	84	<b>Fluox</b>
<i>(Arrow-Fluoxetine Tab dispersible 20 mg, scored to be delisted 1 February 2021)</i>			
<i>(Arrow-Fluoxetine Cap 20 mg to be delisted 1 February 2021)</i>			
<b>PAROXETINE</b>			
Tab 20 mg – <b>1% DV Mar-20 to 2022</b> .....	3.61	90	<b>Loxamine</b>
<b>SERTRALINE</b>			
Tab 50 mg – <b>1% DV Mar-20 to 2022</b> .....	0.92	30	<b>Setrona</b>
Tab 100 mg – <b>1% DV Mar-20 to 2022</b> .....	1.61	30	<b>Setrona</b>

## Antiepilepsy Drugs

### Agents for the Control of Status Epilepticus

<b>CLONAZEPAM</b>			
Inj 1 mg per ml, 1 ml ampoule .....	21.00	5	Rivotril
<b>DIAZEPAM</b>			
Inj 5 mg per ml, 2 ml ampoule .....	23.66	5	Hospira
Rectal tubes 5 mg.....	43.50	5	Stesolid
Rectal tubes 10 mg			
<b>LORAZEPAM</b>			
Inj 2 mg vial			
Inj 4 mg per ml, 1 ml vial			
<b>PARALDEHYDE</b>			
Inj 5 ml ampoule			
Soln 97%			
<b>PHENYTOIN SODIUM</b>			
Inj 50 mg per ml, 2 ml ampoule .....	88.63	5	Hospira
Inj 50 mg per ml, 5 ml ampoule .....	133.92	5	Hospira

### Control of Epilepsy

<b>CARBAMAZEPINE</b>			
Tab 200 mg .....	14.53	100	Tegretol
Tab long-acting 200 mg.....	16.98	100	Tegretol CR
Tab 400 mg .....	34.58	100	Tegretol
Tab long-acting 400 mg.....	39.17	100	Tegretol CR
Oral liq 20 mg per ml .....	26.37	250 ml	Tegretol
<b>CLOBAZAM</b>			
Tab 10 mg			
<b>CLONAZEPAM</b>			
Oral drops 2.5 mg per ml			
<b>ETHOSUXIMIDE</b>			
Cap 250 mg .....	140.88	100	Zarontin
Oral liq 50 mg per ml .....	56.35	200 ml	Zarontin

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>GABAPENTIN</b>			
Note: Gabapentin not to be given in combination with pregabalin			
Cap 100 mg – 1% DV Aug-18 to 2021 .....	2.65	100	<b>Apo-Gabapentin</b>
Cap 300 mg – 1% DV Aug-18 to 2021 .....	4.07	100	<b>Apo-Gabapentin</b>
Cap 400 mg – 1% DV Aug-18 to 2021 .....	5.64	100	<b>Apo-Gabapentin</b>
<b>LACOSAMIDE – Restricted</b> see terms <a href="#">below</a>			
↓ Tab 50 mg .....	25.04	14	Vimpat
↓ Tab 100 mg .....	50.06	14	Vimpat
	200.24	56	Vimpat
↓ Tab 150 mg .....	75.10	14	Vimpat
	300.40	56	Vimpat
↓ Tab 200 mg .....	400.55	56	Vimpat
↓ Inj 10 mg per ml, 20 ml vial			
➔ <b>Restricted (RS1151)</b>			
<b>Initiation</b>			
<i>Re-assessment required after 15 months</i>			
Both:			
1 Patient has partial-onset epilepsy; and			
2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).			
Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.			
<b>Continuation</b>			
Patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).			
Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective			
<b>LAMOTRIGINE</b>			
Tab dispersible 2 mg .....	55.00	30	Lamictal
Tab dispersible 5 mg .....	50.00	30	Lamictal
Tab dispersible 25 mg – 5% DV Oct-19 to 2022 .....	2.76	56	<b>Logem</b>
Tab dispersible 50 mg – 5% DV Oct-19 to 2022 .....	3.31	56	<b>Logem</b>
Tab dispersible 100 mg – 5% DV Oct-19 to 2022 .....	4.40	56	<b>Logem</b>
<b>LEVETIRACETAM</b>			
Tab 250 mg – 1% DV Aug-19 to 2022 .....	4.99	60	<b>Everet</b>
Tab 500 mg – 1% DV Aug-19 to 2022 .....	8.79	60	<b>Everet</b>
Tab 750 mg – 1% DV Aug-19 to 2022 .....	14.39	60	<b>Everet</b>
Tab 1,000 mg – 1% DV Aug-19 to 2022 .....	18.59	60	<b>Everet</b>
Oral liq 100 mg per ml .....	44.78	300 ml	Levetiracetam-AFT
Inj 100 mg per ml, 5 ml vial – 1% DV Oct-19 to 2022 .....	38.95	10	<b>Levetiracetam-AFT</b>
<b>PHENOBARBITONE</b>			
Tab 15 mg – 1% DV Oct-18 to 2021 .....	40.00	500	<b>PSM</b>
Tab 30 mg – 1% DV Oct-18 to 2021 .....	40.00	500	<b>PSM</b>
<b>PHENYTOIN</b>			
Tab 50 mg			
<b>PHENYTOIN SODIUM</b>			
Cap 30 mg			
Cap 100 mg			
Oral liq 6 mg per ml			

↑ Item restricted (see ➔ above); ↓ Item restricted (see ➔ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PREGABALIN</b>			
Note: Pregabalin not to be given in combination with gabapentin			
Cap 25 mg – 1% DV Jul-18 to 2021 .....	2.25	56	<b>Pregabalin Pfizer</b>
Cap 75 mg – 1% DV Jul-18 to 2021 .....	2.65	56	<b>Pregabalin Pfizer</b>
Cap 150 mg – 1% DV Jul-18 to 2021 .....	4.01	56	<b>Pregabalin Pfizer</b>
Cap 300 mg – 1% DV Jul-18 to 2021 .....	7.38	56	<b>Pregabalin Pfizer</b>
<b>PRIMIDONE</b>			
Tab 250 mg			
<b>SODIUM VALPROATE</b>			
Tab 100 mg			
Tab EC 200 mg			
Tab EC 500 mg			
Oral liq 40 mg per ml			
Inj 100 mg per ml, 4 ml vial – 1% DV Sep-18 to 2021 .....	9.98	1	<b>Epilim IV</b>
<b>STIRIPENTOL – Restricted</b> see terms <a href="#">below</a>			
⚡ Cap 250 mg .....	509.29	60	Diacomit
⚡ Powder for oral liq 250 mg sachet .....	509.29	60	Diacomit
➔ <b>Restricted (RS1152)</b>			
<b>Initiation</b>			
Paediatric neurologist			
<i>Re-assessment required after 6 months</i>			
Both:			
1 Patient has confirmed diagnosis of Dravet syndrome; and			
2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.			
<b>Continuation</b>			
Paediatric neurologist			
Patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.			
<b>TOPIRAMATE</b>			
Tab 25 mg .....	11.07	60	Arrow-Topiramate
	26.04		Topamax
	11.07		Topiramate Actavis
Tab 50 mg .....	18.81	60	Arrow-Topiramate
	44.26		Topamax
	18.81		Topiramate Actavis
Tab 100 mg .....	31.99	60	Arrow-Topiramate
	75.25		Topamax
	31.99		Topiramate Actavis
Tab 200 mg .....	55.19	60	Arrow-Topiramate
	129.85		Topamax
	55.19		Topiramate Actavis
Cap sprinkle 15 mg .....	20.84	60	Topamax
Cap sprinkle 25 mg .....	26.04	60	Topamax

**VIGABATRIN – Restricted** see terms [below](#)

⚡ Tab 500 mg

➔ **Restricted (RS1739)**

#### Initiation

*Re-assessment required after 15 months*

Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Either:
  - 1.1 Patient has infantile spasms; or
  - 1.2 Both:
    - 1.2.1 Patient has epilepsy; and
    - 1.2.2 Either:
      - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
      - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and
- 2 Either:
  - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
  - 2.2 It is impractical or impossible (due to comorbid conditions, or health system capacity constraints) to monitor the patient's visual fields.

Notes: "Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

## Continuation

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life; and
- 2 Either:
  - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
  - 2.2 It is impractical or impossible (due to comorbid conditions, or health system capacity constraints) to monitor the patient's visual fields.

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

## Antimigraine Preparations

### Acute Migraine Treatment

DIHYDROERGOTAMINE MESYLATE

Inj 1 mg per ml, 1 ml ampoule

METOCLOPRAMIDE HYDROCHLORIDE WITH PARACETAMOL

Tab 5 mg with paracetamol 500 mg

RIZATRIPTAN

Tab orodispersible 10 mg – 1% DV Oct-20 to 2023 ..... 3.65      30      **Rizamelt**

SUMATRIPTAN

Tab 50 mg – 1% DV Oct-19 to 2022 ..... 24.44      100      **Apo-Sumatriptan**  
 Tab 100 mg – 1% DV Oct-19 to 2022 ..... 46.23      100      **Apo-Sumatriptan**  
 Inj 12 mg per ml, 0.5 ml prefilled pen – 1% DV Sep-20 to 2022 ..... 34.00      2      **Imigran**

### Prophylaxis of Migraine

PIZOTIFEN

Tab 500 mcg ..... 23.21      100      **Sandomigran**

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Antinausea and Vertigo Agents</b>			
APREPITANT – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Cap 2 × 80 mg and 1 × 125 mg – <b>1% DV Jul-18 to 2021</b> .....	84.00	3	<b>Emend Tri-Pack</b>
→ <b>Restricted (RS1154)</b>			
<b>Initiation</b>			
Patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.			
BETAHISTINE DIHYDROCHLORIDE			
Tab 16 mg – <b>1% DV Nov-20 to 2023</b> .....	3.88	84	<b>Vergo 16</b>
CYCLIZINE HYDROCHLORIDE			
Tab 50 mg – <b>1% DV Jan-19 to 2021</b> .....	0.55	10	<b>Nausicalm</b>
CYCLIZINE LACTATE			
Inj 50 mg per ml, 1 ml ampoule – <b>1% DV May-21 to 2022</b> .....	21.53	10	<b>Hameln</b>
	14.95	5	<b>Nausicalm</b>
<i>(Nausicalm Inj 50 mg per ml, 1 ml ampoule to be delisted 1 May 2021)</i>			
DOMPERIDONE			
Tab 10 mg – <b>1% DV Mar-19 to 2021</b> .....	2.25	100	<b>Pharmacy Health</b>
DROPERIDOL			
Inj 2.5 mg per ml, 1 ml ampoule – <b>1% DV May-20 to 2022</b> .....	30.95	10	<b>Droleptan</b>
GRANISETRON			
Inj 1 mg per ml, 3 ml ampoule – <b>1% DV Jan-21 to 2023</b> .....	1.20	1	<b>Deva</b>
HYOSCINE HYDROBROMIDE			
Inj 400 mcg per ml, 1 ml ampoule			
↓ Patch 1.5 mg .....	14.11	2	<b>Scopoderm TTS</b>
→ <b>Restricted (RS1155)</b>			
<b>Initiation</b>			
Any of the following:			
1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or			
2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective; or			
3 For treatment of post-operative nausea and vomiting where cyclizine, droperidol and a 5HT3 antagonist have proven ineffective, are not tolerated or are contraindicated.			
METOCLOPRAMIDE HYDROCHLORIDE			
Tab 10 mg – <b>1% DV Oct-20 to 2023</b> .....	1.30	100	<b>Metoclopramide Actavis 10</b>
Oral liq 5 mg per 5 ml			
Inj 5 mg per ml, 2 ml ampoule – <b>1% DV Jan-20 to 2022</b> .....	9.50	10	<b>Pfizer</b>
ONDANSETRON			
Tab 4 mg – <b>1% DV Apr-20 to 2022</b> .....	2.68	50	<b>Onrex</b>
Tab dispersible 4 mg – <b>1% DV Oct-20 to 2023</b> .....	0.76	10	<b>Ondansetron ODT-DRLA</b>
Tab 8 mg – <b>1% DV Apr-20 to 2022</b> .....	4.57	50	<b>Onrex</b>
Tab dispersible 8 mg – <b>1% DV Oct-20 to 2023</b> .....	1.13	10	<b>Ondansetron ODT-DRLA</b>
Inj 2 mg per ml, 2 ml ampoule .....	1.50	5	<b>Ondansetron-Claris</b>
Inj 2 mg per ml, 4 ml ampoule .....	2.20	5	<b>Ondansetron Kabi</b>

## NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PROCHLORPERAZINE</b>			
Tab buccal 3 mg			
Tab 5 mg – 1% DV Dec-20 to 2023 .....	8.00	250	<b>Nausafix</b>
Inj 12.5 mg per ml, 1 ml ampoule			
Suppos 25 mg			
<b>TROPISETRON</b>			
Inj 1 mg per ml, 2 ml ampoule – 1% DV Sep-18 to 2021 .....	8.95	1	<b>Tropisetron-AFT</b>
Inj 1 mg per ml, 5 ml ampoule .....	13.95	1	Tropisetron-AFT
<b>Antipsychotic Agents</b>			
<b>General</b>			
<b>AMISULPRIDE</b>			
Tab 100 mg – 1% DV Nov-19 to 2022 .....	5.15	30	<b>Sulprix</b>
Tab 200 mg – 1% DV Nov-19 to 2022 .....	14.96	60	<b>Sulprix</b>
Tab 400 mg – 1% DV Feb-20 to 2022 .....	29.78	60	<b>Sulprix</b>
Oral liq 100 mg per ml			
<b>ARIPIPRAZOLE</b>			
Tab 5 mg – 1% DV Aug-18 to 2021 .....	17.50	30	<b>Aripiprazole Sandoz</b>
Tab 10 mg – 1% DV Aug-18 to 2021 .....	17.50	30	<b>Aripiprazole Sandoz</b>
Tab 15 mg – 1% DV Aug-18 to 2021 .....	17.50	30	<b>Aripiprazole Sandoz</b>
Tab 20 mg – 1% DV Aug-18 to 2021 .....	17.50	30	<b>Aripiprazole Sandoz</b>
Tab 30 mg – 1% DV Aug-18 to 2021 .....	17.50	30	<b>Aripiprazole Sandoz</b>
<b>CHLORPROMAZINE HYDROCHLORIDE</b>			
Tab 10 mg – 1% DV Jan-20 to 2022 .....	14.83	100	<b>Largactil</b>
Tab 25 mg – 1% DV Jan-20 to 2022 .....	15.62	100	<b>Largactil</b>
Tab 100 mg – 1% DV Jan-20 to 2022 .....	36.73	100	<b>Largactil</b>
Oral liq 10 mg per ml			
Oral liq 20 mg per ml			
Inj 25 mg per ml, 2 ml ampoule – 1% DV Jan-20 to 2022 .....	30.79	10	<b>Largactil</b>
<b>CLOZAPINE</b>			
Tab 25 mg .....	6.69	50	Clopine
	13.37	100	Clopine
	5.69	50	Clozaril
	11.36	100	Clozaril
Tab 50 mg .....	8.67	50	Clopine
	17.33	100	Clopine
Tab 100 mg .....	17.33	50	Clopine
	34.65	100	Clopine
	14.73	50	Clozaril
	29.45	100	Clozaril
Tab 200 mg .....	34.65	50	Clopine
	69.30	100	Clopine
Oral liq 50 mg per ml .....	17.33	100 ml	Clopine
<b>HALOPERIDOL</b>			
Tab 500 mcg – 1% DV Oct-19 to 2022 .....	6.23	100	<b>Serenace</b>
Tab 1.5 mg – 1% DV Oct-19 to 2022 .....	9.43	100	<b>Serenace</b>
Tab 5 mg – 1% DV Oct-19 to 2022 .....	29.72	100	<b>Serenace</b>
Oral liq 2 mg per ml – 1% DV Oct-19 to 2022 .....	23.84	100 ml	<b>Serenace</b>
Inj 5 mg per ml, 1ml ampoule – 1% DV Oct-19 to 2022 .....	21.55	10	<b>Serenace</b>

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>LEVOMEPROMAZINE</b>			
Tab 25 mg – 1% DV Sep-19 to 2022	16.10	100	Nozinan
Tab 100 mg – 1% DV Sep-19 to 2022	41.75	100	Nozinan
<b>LEVOMEPROMAZINE HYDROCHLORIDE</b>			
Inj 25 mg per ml, 1 ml ampoule – 1% DV Apr-20 to 2022	33.50	10	Nozinan
<b>LITHIUM CARBONATE</b>			
Tab long-acting 400 mg			
Cap 250 mg	9.42	100	Douglas
<b>OLANZAPINE</b>			
Tab 2.5 mg – 1% DV Nov-20 to 2023	1.35	28	Zypine
Tab 5 mg – 1% DV Nov-20 to 2023	1.58	28	Zypine
Tab orodispersible 5 mg – 1% DV Nov-20 to 2023	1.81	28	Zypine ODT
Tab 10 mg – 1% DV Nov-20 to 2023	2.01	28	Zypine
Tab orodispersible 10 mg – 1% DV Nov-20 to 2023	2.38	28	Zypine ODT
Inj 10 mg vial			
<b>PERICYAZINE</b>			
Tab 2.5 mg			
Tab 10 mg			
<b>QUETIAPINE</b>			
Tab 25 mg – 1% DV Nov-20 to 2023	2.15	90	Quetapel
Tab 100 mg – 1% DV Nov-20 to 2023	5.06	90	Quetapel
Tab 200 mg – 1% DV Nov-20 to 2023	8.90	90	Quetapel
Tab 300 mg – 1% DV Nov-20 to 2023	12.86	90	Quetapel
<b>RISPERIDONE</b>			
Tab 0.5 mg – 1% DV Dec-20 to 2023	1.86	60	Risperidone (Teva)
Tab 1 mg – 1% DV Dec-20 to 2023	2.06	60	Risperidone (Teva)
Tab 2 mg – 1% DV Dec-20 to 2023	2.29	60	Risperidone (Teva)
Tab 3 mg – 1% DV Dec-20 to 2023	2.50	60	Risperidone (Teva)
Tab 4 mg – 1% DV Dec-20 to 2023	3.42	60	Risperidone (Teva)
Oral liq 1 mg per ml – 1% DV Nov-20 to 2023	8.90	30 ml	Risperon
<b>ZIPRASIDONE</b>			
Cap 20 mg – 1% DV Dec-18 to 2021	14.50	60	Zusdone
Cap 40 mg – 1% DV Sep-18 to 2021	24.70	60	Zusdone
Cap 60 mg – 1% DV Sep-18 to 2021	33.80	60	Zusdone
Cap 80 mg – 1% DV Sep-18 to 2021	39.70	60	Zusdone
<b>ZUCLOPENTHIXOL ACETATE</b>			
Inj 50 mg per ml, 1 ml ampoule			
Inj 50 mg per ml, 2 ml ampoule			
<b>ZUCLOPENTHIXOL HYDROCHLORIDE</b>			
Tab 10 mg	31.45	100	Clopixol

### Depot Injections

<b>FLUPENTHIXOL DECANOATE</b>			
Inj 20 mg per ml, 1 ml ampoule	13.14	5	Fluanxol
Inj 20 mg per ml, 2 ml ampoule	20.90	5	Fluanxol
Inj 100 mg per ml, 1 ml ampoule	40.87	5	Fluanxol
<b>HALOPERIDOL DECANOATE</b>			
Inj 50 mg per ml, 1 ml ampoule	28.39	5	Haldol
Inj 100 mg per ml, 1 ml ampoule	55.90	5	Haldol Concentrate

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>OLANZAPINE – Restricted</b> see terms <a href="#">below</a>			
‡ Inj 210 mg vial – 1% DV Oct-18 to 2021.....	252.00	1	<b>Zyprexa Relprevv</b>
‡ Inj 300 mg vial – 1% DV Oct-18 to 2021.....	414.00	1	<b>Zyprexa Relprevv</b>
‡ Inj 405 mg vial – 1% DV Oct-18 to 2021.....	504.00	1	<b>Zyprexa Relprevv</b>

➔ **Restricted (RS1379)**

## Initiation

*Re-assessment required after 12 months*

Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or paliperidone depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia; and
  - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

## Continuation

*Re-assessment required after 12 months*

The initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

**PALIPERIDONE – Restricted** see terms [below](#)

‡ Inj 25 mg syringe .....	194.25	1	Invega Sustenna
‡ Inj 50 mg syringe .....	271.95	1	Invega Sustenna
‡ Inj 75 mg syringe .....	357.42	1	Invega Sustenna
‡ Inj 100 mg syringe .....	435.12	1	Invega Sustenna
‡ Inj 150 mg syringe .....	435.12	1	Invega Sustenna

➔ **Restricted (RS1381)**

## Initiation

*Re-assessment required after 12 months*

Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

## Continuation

*Re-assessment required after 12 months*

The initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

**PIPTHIAZINE PALMITATE – Restricted:** For continuation only

- ➔ Inj 50 mg per ml, 1 ml ampoule
- ➔ Inj 50 mg per ml, 2 ml ampoule

**RISPERIDONE – Restricted** see terms [below](#)

‡ Inj 25 mg vial .....	135.98	1	Risperdal Consta
‡ Inj 37.5 mg vial .....	178.71	1	Risperdal Consta
‡ Inj 50 mg vial .....	217.56	1	Risperdal Consta

➔ **Restricted (RS1380)**

## Initiation

*Re-assessment required after 12 months*

Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

### Continuation

*Re-assessment required after 12 months*

The initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

### ZUCLOPENTHIXOL DECANOATE

Inj 200 mg per ml, 1 ml ampoule .....	19.80	5	Clopixol
Inj 500 mg per ml, 1 ml ampoule .....			e.g. Clopixol Conc

## Anxiolytics

### BUSPIRONE HYDROCHLORIDE

Tab 5 mg – 1% DV Sep-18 to 2021 .....	20.23	100	Orion
Tab 10 mg – 1% DV Sep-18 to 2021 .....	13.16	100	Orion

### CLONAZEPAM

Tab 500 mcg – 1% DV Jun-18 to 2021 .....	5.64	100	Paxam
Tab 2 mg – 1% DV Jun-18 to 2021 .....	10.78	100	Paxam

### DIAZEPAM

Tab 2 mg – 1% DV Dec-20 to 2023 .....	61.07	500	Arrow-Diazepam
Tab 5 mg – 1% DV Dec-20 to 2023 .....	73.60	500	Arrow-Diazepam

### LORAZEPAM

Tab 1 mg – 1% DV Sep-18 to 2021 .....	9.72	250	Ativan
Tab 2.5 mg – 1% DV Sep-18 to 2021 .....	12.50	100	Ativan

### OXAZEPAM

Tab 10 mg .....	6.17	100	Ox-Pam
Tab 15 mg .....	8.53	100	Ox-Pam

## Multiple Sclerosis Treatments

### DIMETHYL FUMARATE – **Restricted** see terms [below](#)

↓ Cap 120 mg .....	520.00	14	Tecfidera
↓ Cap 240 mg .....	2,000.00	56	Tecfidera
→ <b>Restricted (RS1504)</b>			

### Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

### FINGOLIMOD – **Restricted** see terms [below](#)

↓ Cap 0.5 mg .....	2,200.00	28	Gilenya
→ <b>Restricted (RS1433)</b>			

### Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

NATALIZUMAB – **Restricted** see terms [below](#)

⚡ Inj 20 mg per ml, 15 ml vial..... 1,750.00 1 Tysabri

➡ **Restricted (RS1447)**

## Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

OCRELIZUMAB – **Restricted** see terms [below](#)

⚡ Inj 30 mg per ml, 10 ml vial..... 9,346.00 1 Ocrevus

➡ **Restricted (RS1711)**

## Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

TERIFLUNOMIDE – **Restricted** see terms [below](#)

⚡ Tab 14 mg ..... 1,582.62 28 Aubagio

➡ **Restricted (RS1505)**

## Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

## Other Multiple Sclerosis Treatments

➡ **Restricted (RS1434)**

## Initiation

Only for use in patients with approval by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (set out in Section B of the Pharmaceutical Schedule).

GLATIRAMER ACETATE – **Restricted** see terms [above](#)

⚡ Inj 40 mg prefilled syringe..... 2,275.00 12 Copaxone

INTERFERON BETA-1-ALPHA – **Restricted** see terms [above](#)

⚡ Inj 6 million iu in 0.5 ml pen injector..... 1,170.00 4 Avonex Pen

⚡ Inj 6 million iu in 0.5 ml syringe..... 1,170.00 4 Avonex

INTERFERON BETA-1-BETA – **Restricted** see terms [above](#)

⚡ Inj 8 million iu per ml, 1 ml vial

## Sedatives and Hypnotics

CHLORAL HYDRATE

Oral liq 100 mg per ml

Oral liq 200 mg per ml

LORMETAZEPAM – **Restricted:** For continuation only

➡ Tab 1 mg

MELATONIN – **Restricted** see terms [on the next page](#)

⚡ Tab modified-release 2 mg..... 28.22 30 Circadin

⚡ Tab 3 mg

Note: Only for use in compounding an oral liquid formulation, for in-hospital use only.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>➔ Restricted (RS1576)</b>			
<b>Initiation – insomnia secondary to neurodevelopmental disorder</b>			
Psychiatrist, paediatrician, neurologist or respiratory specialist			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder); and			
2 Behavioural and environmental approaches have been tried or are inappropriate; and			
3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and			
4 Patient is aged 18 years or under.			
<b>Continuation – insomnia secondary to neurodevelopmental disorder</b>			
Psychiatrist, paediatrician, neurologist or respiratory specialist			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 Patient is aged 18 years or under; and			
2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and			
3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and			
4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.			
<b>Initiation – insomnia where benzodiazepines and zopiclone are contraindicated</b>			
Both:			
1 Patient has insomnia and benzodiazepines and zopiclone are contraindicated; and			
2 For in-hospital use only.			
<b>MIDAZOLAM</b>			
Tab 7.5 mg			
Oral liq 2 mg per ml			
Inj 1 mg per ml, 5 ml ampoule – 1% DV Jan-19 to 2021 .....	2.98	10	<b>Mylan Midazolam</b>
Inj 5 mg per ml, 3 ml ampoule – 1% DV Jan-19 to 2021 .....	2.36	5	<b>Mylan Midazolam</b>
<b>PHENOBARBITONE</b>			
Inj 200 mg per ml, 1 ml ampoule			
<b>TEMAZEPAM</b>			
Tab 10 mg – 1% DV Nov-20 to 2023 .....	1.33	25	<b>Normison</b>
<b>TRIAZOLAM – Restricted:</b> For continuation only			
➔ Tab 125 mcg			
➔ Tab 250 mcg			
<b>ZOPICLONE</b>			
Tab 7.5 mg			
<b>Stimulants / ADHD Treatments</b>			
<b>ATOMOXETINE</b>			
Cap 10 mg – 1% DV Sep-20 to 2022 .....	18.41	28	<b>Generic Partners</b>
Cap 18 mg – 1% DV Sep-20 to 2022 .....	27.06	28	<b>Generic Partners</b>
Cap 25 mg – 1% DV Sep-20 to 2022 .....	29.22	28	<b>Generic Partners</b>
Cap 40 mg – 1% DV Sep-20 to 2022 .....	29.22	28	<b>Generic Partners</b>
Cap 60 mg – 1% DV Sep-20 to 2022 .....	46.51	28	<b>Generic Partners</b>
Cap 80 mg – 1% DV Sep-20 to 2022 .....	56.45	28	<b>Generic Partners</b>
Cap 100 mg – 1% DV Sep-20 to 2022 .....	58.48	28	<b>Generic Partners</b>
<b>CAFFEINE</b>			
Tab 100 mg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>DEXAMFETAMINE SULFATE – Restricted</b> see terms <a href="#">below</a>			
↓ Tab 5 mg – 1% DV Oct-18 to 2021.....	20.00	100	<b>PSM</b>
➔ <b>Restricted (RS1169)</b>			
<b>Initiation – ADHD</b>			
Paediatrician or psychiatrist			
Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria.			
<b>Initiation – Narcolepsy</b>			
Neurologist or respiratory specialist			
<i>Re-assessment required after 24 months</i>			
Patient suffers from narcolepsy.			
<b>Continuation – Narcolepsy</b>			
Neurologist or respiratory specialist			
<i>Re-assessment required after 24 months</i>			
The treatment remains appropriate and the patient is benefiting from treatment.			
<b>METHYLPHENIDATE HYDROCHLORIDE – Restricted</b> see terms <a href="#">below</a>			
↓ Tab extended-release 18 mg.....	58.96 7.75	30	Concerta Methylphenidate ER - Teva
↓ Tab extended-release 27 mg.....	65.44 11.45	30	Concerta Methylphenidate ER - Teva
↓ Tab extended-release 36 mg.....	71.93 15.50	30	Concerta Methylphenidate ER - Teva
↓ Tab extended-release 54 mg.....	86.24 22.25	30	Concerta Methylphenidate ER - Teva
↓ Tab immediate-release 5 mg.....	3.20	30	Rubifen
↓ Tab immediate-release 10 mg.....	3.00	30	Ritalin Rubifen
↓ Tab immediate-release 20 mg.....	7.85	30	Rubifen
↓ Tab sustained-release 20 mg.....	50.00 10.95	100	Ritalin SR Rubifen SR
↓ Cap modified-release 10 mg.....	15.60	30	Ritalin LA
↓ Cap modified-release 20 mg.....	20.40	30	Ritalin LA
↓ Cap modified-release 30 mg.....	25.52	30	Ritalin LA
↓ Cap modified-release 40 mg.....	30.60	30	Ritalin LA
<i>(Ritalin SR Tab sustained-release 20 mg to be delisted 1 June 2021)</i>			
➔ <b>Restricted (RS1294)</b>			
<b>Initiation – ADHD (immediate-release and sustained-release formulations)</b>			
Paediatrician or psychiatrist			
Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria.			
<b>Initiation – Narcolepsy (immediate-release and sustained-release formulations)</b>			
Neurologist or respiratory specialist			
<i>Re-assessment required after 24 months</i>			
Patient suffers from narcolepsy.			
<b>Continuation – Narcolepsy (immediate-release and sustained-release formulations)</b>			
Neurologist or respiratory specialist			
<i>Re-assessment required after 24 months</i>			
The treatment remains appropriate and the patient is benefiting from treatment.			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

#### Initiation – Extended-release and modified-release formulations

Paediatrician or psychiatrist

Both:

- 1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria; and
- 2 Either:
  - 2.1 Patient is taking a currently listed formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
  - 2.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

MODAFINIL – **Restricted** see terms [below](#)

↓ Tab 100 mg .....	64.00	60	Modavigil
--------------------	-------	----	-----------

→ **Restricted (RS1761)**

#### Initiation – Narcolepsy

Neurologist or respiratory specialist

*Re-assessment required after 24 months*

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Any of the following:
  - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
  - 2.2 A multiple sleep latency test is not possible due to COVID-19 constraints on the health sector; or
  - 2.3 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
  - 3.1 An effective dose of a listed formulation of methylphenidate or dexamphetamine has been trialled and discontinued because of intolerable side effects; or
  - 3.2 Methylphenidate and dexamphetamine are contraindicated.

#### Continuation – Narcolepsy

Neurologist or respiratory specialist

*Re-assessment required after 24 months*

The treatment remains appropriate and the patient is benefiting from treatment.

## Treatments for Dementia

DONEPEZIL HYDROCHLORIDE

Tab 5 mg – 1% DV Dec-20 to 2023 .....	4.34	90	Donepezil-Rex
Tab 10 mg – 1% DV Dec-20 to 2023 .....	6.64	90	Donepezil-Rex

RIVASTIGMINE – **Restricted** see terms [below](#)

↓ Patch 4.6 mg per 24 hour – 1% DV Apr-20 to 2021 .....	48.75	30	Generic Partners
↓ Patch 9.5 mg per 24 hour – 1% DV Apr-20 to 2021 .....	48.75	30	Generic Partners

→ **Restricted (RS1436)**

#### Initiation

*Re-assessment required after 6 months*

Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

## Continuation

*Re-assessment required after 12 months*

Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

## Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE – **Restricted** see terms [below](#)

⚡ Tab 2 mg with naloxone 0.5 mg – 1% DV Apr-20 to 2022 .....	18.37	28	Buprenorphine Naloxone BNM
⚡ Tab 8 mg with naloxone 2 mg – 1% DV Apr-20 to 2022 .....	53.12	28	Buprenorphine Naloxone BNM

➡ **Restricted (RS1172)**

### Initiation – Detoxification

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Prescriber works in an opioid treatment service approved by the Ministry of Health.

### Initiation – Maintenance treatment

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Prescriber works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

Tab modified-release 150 mg – 1% DV Mar-21 to 2023.....	11.00	30	Zyban
---	-------	----	-------

DISULFIRAM

Tab 200 mg .....	153.00	100	Antabuse
------------------	--------	-----	----------

NALTREXONE HYDROCHLORIDE – **Restricted** see terms [below](#)

⚡ Tab 50 mg – 1% DV Jan-21 to 2023 .....	133.33	30	Naltrexone
--	--------	----	------------

➡ **Restricted (RS1173)**

### Initiation – Alcohol dependence

Both:

- 1 Patient is currently enrolled, or is planned to be enrolled, in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Naltrexone is to be prescribed by, or on the recommendation of, a physician working in an Alcohol and Drug Service.

### Initiation – Constipation

For the treatment of opioid-induced constipation.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>NICOTINE – Some items restricted</b> see terms <a href="#">below</a>			
Patch 7 mg per 24 hours .....	18.14	28	Habitrol
Patch 14 mg per 24 hours .....	19.95	28	Habitrol
Patch 21 mg per 24 hours .....	22.86	28	Habitrol
↓ Oral spray 1 mg per dose .....			<i>e.g. Nicorette QuickMist Mouth Spray</i>
Lozenge 1 mg .....	19.18	216	Habitrol
Lozenge 2 mg .....	21.02	216	Habitrol
↓ Soln for inhalation 15 mg cartridge .....			<i>e.g. Nicorette Inhalator</i>
Gum 2 mg .....	38.21	384	Habitrol (Fruit) Habitrol (Mint)
Gum 4 mg .....	44.17	384	Habitrol (Fruit) Habitrol (Mint)

➔ **Restricted (RS1310)**

**Initiation**

Any of the following:

- 1 For perioperative use in patients who have a 'nil by mouth' instruction; or
- 2 For use within mental health inpatient units; or
- 3 For acute use in agitated patients who are unable to leave the hospital facilities.

**VARENICLINE – Restricted** see terms [below](#)

↓ Tab 0.5 mg × 11 and 1 mg × 42 – <b>1% DV Mar-19 to 2021</b> .....	25.64	53	<b>Varenicline Pfizer</b>
↓ Tab 1 mg – <b>1% DV Mar-19 to 2021</b> .....	27.10	56	<b>Varenicline Pfizer</b>

➔ **Restricted (RS1702)**

**Initiation**

All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
  - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
  - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline in a 12 month period.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## Chemotherapeutic Agents

### Alkylating Agents

**BENDAMUSTINE HYDROCHLORIDE** – **Restricted** see terms [below](#)

↓ Inj 25 mg vial .....	271.35	1	Ribomustin
↓ inj 100 mg vial.....	1,085.38	1	Ribomustin

➡ **Restricted (RS1578)**

#### Initiation – treatment naive CLL

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m<sup>2</sup> on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

#### Initiation – Indolent, Low-grade lymphomas

*Re-assessment required after 9 months*

All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
  - 3.1 Both:
    - 3.1.1 Patient is treatment naive; and
    - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
  - 3.2 All of the following:
    - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
    - 3.2.2 The patient has not received prior bendamustine therapy; and
    - 3.2.3 Either:
      - 3.2.3.1 Both:
        - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
        - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
      - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

#### Continuation – Indolent, Low-grade lymphomas

*Re-assessment required after 9 months*

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
    - 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients. Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenström's macroglobulinaemia.			
<b>BUSULFAN</b>			
Tab 2 mg .....	89.25	100	Myleran
Inj 6 mg per ml, 10 ml ampoule			
<b>CARMUSTINE</b>			
Inj 100 mg vial .....	1,387.00	1	BiCNU Bicnu Heritage
<b>CHLORAMBUCIL</b>			
Tab 2 mg			
<b>CYCLOPHOSPHAMIDE</b>			
Tab 50 mg .....	79.00	50	Endoxan
	158.00	100	Procytox
Inj 1 g vial – <b>1% DV Oct-18 to 2021</b> .....	35.65	1	<b>Endoxan</b>
Inj 2 g vial – <b>1% DV Oct-18 to 2021</b> .....	71.25	1	<b>Endoxan</b>
<b>IFOSFAMIDE</b>			
Inj 1 g vial .....	96.00	1	Holoxan
Inj 2 g vial .....	180.00	1	Holoxan
<b>LOMUSTINE</b>			
Cap 10 mg .....	132.59	20	Ceenu
Cap 40 mg .....	399.15	20	Ceenu
<b>MELPHALAN</b>			
Tab 2 mg			
Inj 50 mg vial			
<b>THIOTEPA</b>			
Inj 15 mg vial			
Inj 100 mg vial			

### Anthracyclines and Other Cytotoxic Antibiotics

<b>BLEOMYCIN SULPHATE</b>			
Inj 15,000 iu vial – <b>1% DV Dec-18 to 2021</b> .....	161.01	1	<b>DBL Bleomycin Sulfate</b>
<b>DACTINOMYCIN [ACTINOMYCIN D]</b>			
Inj 0.5 mg vial .....	255.00	1	Cosmegen
<b>DAUNORUBICIN</b>			
Inj 2 mg per ml, 10 ml vial.....	149.50	1	Pfizer
<b>DOXORUBICIN HYDROCHLORIDE</b>			
Inj 2 mg per ml, 5 ml vial			
Inj 2 mg per ml, 25 ml vial.....	11.50	1	Doxorubicin Ebewe
Note: DV limit applies to all 50 mg presentations of doxorubicin hydrochloride.			
Inj 50 mg vial			
Inj 2 mg per ml, 50 ml vial.....	23.00	1	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial – <b>1% DV Jan-19 to 2021</b> .....	56.15	1	<b>Doxorubicin Ebewe</b>
<b>EPIRUBICIN HYDROCHLORIDE</b>			
Inj 2 mg per ml, 5 ml vial.....	25.00	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial.....	30.00	1	Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial – <b>1% DV Apr-19 to 2021</b> .....	85.00	1	<b>Epirubicin Ebewe</b>

## ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>IDARUBICIN HYDROCHLORIDE</b>			
Inj 5 mg vial – <b>1% DV Sep-18 to 2021</b> .....	93.00	1	<b>Zavedos</b>
Inj 10 mg vial – <b>1% DV Sep-18 to 2021</b> .....	198.00	1	<b>Zavedos</b>
<b>MITOMYCIN C</b>			
Inj 5 mg vial .....	851.37	1	Teva
Inj 20 mg vial .....	3,275.00	1	Teva
<b>MITOZANTRONE</b>			
Inj 2 mg per ml, 10 ml vial.....	97.50	1	Mitozantrone Ebewe
<b>Antimetabolites</b>			
<b>AZACITIDINE – Restricted</b> see terms <a href="#">below</a>			
⚡ Inj 100 mg vial – <b>1% DV Dec-18 to 2021</b> .....	139.00	1	<b>Azacitidine Dr Reddy's</b>
➡ <b>Restricted (RS1418)</b>			
<b>Initiation</b>			
Haematologist			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 Any of the following:			
1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or			
1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or			
1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and			
2 The patient has performance status (WHO/ECOG) grade 0-2; and			
3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and			
4 The patient has an estimated life expectancy of at least 3 months.			
<b>Continuation</b>			
Haematologist			
<i>Re-assessment required after 12 months</i>			
Both:			
1 No evidence of disease progression, and; and			
2 The treatment remains appropriate and patient is benefitting from treatment.			
<b>CAPECITABINE</b>			
Tab 150 mg – <b>1% DV Jul-20 to 2022</b> .....	10.00	60	<b>Capercit</b>
Tab 500 mg – <b>1% DV Jul-20 to 2022</b> .....	49.00	120	<b>Capercit</b>
<b>CLADRIBINE</b>			
Inj 2 mg per ml, 5 ml vial			
Inj 1 mg per ml, 10 ml vial.....	749.96	1	Leustatin
<b>CYTARABINE</b>			
Inj 20 mg per ml, 5 ml vial.....	400.00	5	Pfizer
Inj 100 mg per ml, 20 ml vial – <b>1% DV Dec-18 to 2021</b> .....	41.36	1	<b>Pfizer</b>
<b>FLUDARABINE PHOSPHATE</b>			
Tab 10 mg – <b>1% DV Sep-18 to 2021</b> .....	412.00	20	<b>Fludara Oral</b>
Inj 50 mg vial – <b>1% DV Nov-19 to 2022</b> .....	576.45	5	<b>Fludarabine Ebewe</b>
<b>FLUOROURACIL</b>			
Inj 50 mg per ml, 20 ml vial – <b>1% DV Oct-18 to 2021</b> .....	12.00	1	<b>Fluorouracil Ebewe</b>
Inj 50 mg per ml, 100 ml vial – <b>1% DV Oct-18 to 2021</b> .....	30.00	1	<b>Fluorouracil Ebewe</b>

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)  
e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>GEMCITABINE</b>			
Inj 10 mg per ml, 100 ml vial – <b>1% DV Jul-20 to 2023</b> .....	15.89	1	<b>Gemcitabine Ebewe</b>
<b>MERCAPTOPURINE</b>			
Tab 50 mg – <b>1% DV Jul-19 to 2022</b> .....	37.00	25	<b>Puri-nethol</b>
↓ Oral suspension 20 mg per ml.....	428.00	100 ml	Allmercap
➔ <b>Restricted (RS1635)</b>			
<b>Initiation</b>			
Paediatric haematologist or paediatric oncologist			
<i>Re-assessment required after 12 months</i>			
The patient requires a total dose of less than one full 50 mg tablet per day.			
<b>Continuation</b>			
Paediatric haematologist or paediatric oncologist			
<i>Re-assessment required after 12 months</i>			
The patient requires a total dose of less than one full 50 mg tablet per day.			
<b>METHOTREXATE</b>			
Tab 2.5 mg – <b>1% DV Jan-19 to 2021</b> .....	8.05	90	<b>Trexate</b>
Tab 10 mg – <b>1% DV Jan-19 to 2021</b> .....	31.75	90	<b>Trexate</b>
Inj 2.5 mg per ml, 2 ml vial			
Inj 7.5 mg prefilled syringe.....	14.61	1	Methotrexate Sandoz
Inj 10 mg prefilled syringe.....	14.66	1	Methotrexate Sandoz
Inj 15 mg prefilled syringe.....	14.77	1	Methotrexate Sandoz
Inj 20 mg prefilled syringe.....	14.88	1	Methotrexate Sandoz
Inj 25 mg prefilled syringe.....	14.99	1	Methotrexate Sandoz
Inj 30 mg prefilled syringe.....	15.09	1	Methotrexate Sandoz
Inj 25 mg per ml, 2 ml vial.....	30.00	5	DBL Methotrexate
			Onco-Vial
			Methotrexate DBL
			Onco-Vial
Inj 25 mg per ml, 20 ml vial.....	45.00	1	DBL Methotrexate
			Onco-Vial
Inj 100 mg per ml, 10 ml vial.....	25.00	1	Methotrexate Ebewe
Inj 100 mg per ml, 50 ml vial – <b>1% DV Oct-20 to 2023</b> .....	79.99	1	<b>Methotrexate Ebewe</b>
<i>(DBL Methotrexate Onco-Vial Inj 25 mg per ml, 2 ml vial to be delisted 1 May 2021)</i>			
<b>PEMETREXED – Restricted</b> see terms <a href="#">below</a>			
↓ Inj 100 mg vial .....	60.89	1	Juno Pemetrexed
↓ Inj 500 mg vial .....	217.77	1	Juno Pemetrexed
➔ <b>Restricted (RS1596)</b>			
<b>Initiation – Mesothelioma</b>			
<i>Re-assessment required after 8 months</i>			
Both:			
1 Patient has been diagnosed with mesothelioma; and			
2 Pemetrexed to be administered at a dose of 500 mg/m <sup>2</sup> every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.			
<b>Continuation – Mesothelioma</b>			
<i>Re-assessment required after 8 months</i>			
All of the following:			
1 No evidence of disease progression; and			
2 The treatment remains appropriate and the patient is benefitting from treatment; and			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

3 Pemetrexed to be administered at a dose of 500mg/m<sup>2</sup> every 21 days for a maximum of 6 cycles.

## Initiation – Non small cell lung cancer

*Re-assessment required after 8 months*

Both:

1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and

2 Either:

2.1 Both:

2.1.1 Patient has chemotherapy-naïve disease; and

2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m<sup>2</sup> every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or

2.2 All of the following:

2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and

2.2.2 Patient has not received prior funded treatment with pemetrexed; and

2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m<sup>2</sup> every 21 days for a maximum of 6 cycles.

## Continuation – Non small cell lung cancer

*Re-assessment required after 8 months*

All of the following:

1 No evidence of disease progression; and

2 The treatment remains appropriate and the patient is benefitting from treatment; and

3 Pemetrexed is to be administered at a dose of 500mg/m<sup>2</sup> every 21 days.

## THIOGUANINE

Tab 40 mg

## Other Cytotoxic Agents

### AMSACRINE

Inj 50 mg per ml, 1.5 ml ampoule

Inj 75 mg

### ANAGRELIDE HYDROCHLORIDE

Cap 0.5 mg

### ARSENIC TRIOXIDE

Inj 1 mg per ml, 10 ml vial.....4,817.00

10

Phenasen

### BORTEZOMIB – **Restricted** see terms [below](#)

⚡ Inj 3.5 mg vial – **1% DV Aug-20 to 2022**.....105.00

1

**Bortezomib Dr-Reddy's**

➡ **Restricted (RS1725)**

### Initiation – multiple myeloma/amyloidosis

Either:

1 The patient has symptomatic multiple myeloma; or

2 The patient has symptomatic systemic AL amyloidosis.

### DACARBAZINE

Inj 200 mg vial .....62.70

1

DBL Dacarbazine

### ETOPOSIDE

Cap 50 mg – **1% DV Jul-19 to 2022**.....340.73

20

**Vepesid**

Cap 100 mg – **1% DV Jul-19 to 2022**.....340.73

10

**Vepesid**

Inj 20 mg per ml, 5 ml vial.....7.90

1

Rex Medical

### ETOPOSIDE (AS PHOSPHATE)

Inj 100 mg vial .....40.00

1

Etopophos

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
HYDROXYUREA [HYDROXYCARBAMIDE]			
Cap 500 mg – 1% DV Feb-21 to 2023.....	23.82	100	<b>Devatis</b>
	31.76		Hydrea
<i>(Hydrea Cap 500 mg to be delisted 1 February 2021)</i>			
IRINOTECAN HYDROCHLORIDE			
Inj 20 mg per ml, 5 ml vial – 1% DV Apr-19 to 2021.....	71.44	1	<b>Irinotecan Actavis 100</b>
LENALIDOMIDE – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Cap 5 mg.....	5,122.76	28	Revlimid
↓ Cap 10 mg.....	4,655.25	21	Revlimid
	6,207.00	28	Revlimid
↓ Cap 15 mg.....	5,429.39	21	Revlimid
	7,239.18	28	Revlimid
↓ Cap 25 mg.....	7,627.00	21	Revlimid

→ **Restricted (RS1730)**

**Initiation – Relapsed/refractory disease**

Haematologist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Either:
  - 3.1 Lenalidomide to be used as third line\* treatment for multiple myeloma; or
  - 3.2 Both:
    - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
    - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

**Continuation – Relapsed/refractory disease**

Haematologist

*Re-assessment required after 6 months*

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

**Initiation – Maintenance following first-line autologous stem cell transplant (SCT)**

Haematologist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 The patient has ECOG performance score of 0-1; and
- 5 Lenalidomide to be administered at a maximum dose of 15 mg/day.

**Continuation – Maintenance following first-line autologous stem cell transplant (SCT)**

Haematologist

*Re-assessment required after 6 months*

Both:

- 1 No evidence of disease progression; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with \* is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

OLAPARIB – **Restricted** see terms [below](#)

⚡ Tab 100 mg .....	3,701.00	56	Lynparza
⚡ Tab 150 mg .....	3,701.00	56	Lynparza
⚡ Cap 50 mg .....	7,402.00	448	Lynparza

➡ **Restricted (RS1722)**

## Initiation

Medical oncologist

*Re-assessment required after 12 months*

All of the following:

- 1 Patient has a high-grade serous\* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Patient has received at least two lines of previous treatment with platinum-based chemotherapy; and
- 4 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line of platinum-based chemotherapy; and
- 5 Patient's disease must have achieved partial or complete response to treatment with the immediately preceding platinum-based regimen; and
- 6 Patient's disease has not progressed following prior treatment with olaparib; and
- 7 Treatment will be commenced within 8 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 8 Treatment to be administered as maintenance treatment; and
- 9 Treatment not to be administered in combination with other chemotherapy.

## Continuation

Medical oncologist

*Re-assessment required after 12 months*

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 No evidence of progressive disease; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy.

Note: \*Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

PEGASPARGASE – **Restricted** see terms [below](#)

⚡ Inj 750 iu per ml, 5 ml vial .....	3,455.00	1	Oncaspar LYO
--------------------------------------	----------	---	--------------

➡ **Restricted (RS1788)**

## Initiation – Newly diagnosed ALL

*Limited to 12 months treatment*

Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

## Initiation – Relapsed ALL

*Limited to 12 months treatment*

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

### Initiation – Lymphoma

Limited to 12 months treatment

Patient has lymphoma requiring L-asparaginase containing protocol (e.g. SMILE).

### PENTOSTATIN [DEOXYCOFORMYCIN]

Inj 10 mg vial

### PROCARBAZINE HYDROCHLORIDE

Cap 50 mg .....	980.00	50	Natulan
-----------------	--------	----	---------

### TEMOZOLOMIDE – **Restricted** see terms [below](#)

↓ Cap 5 mg – 1% DV May-20 to 2022 .....	9.13	5	<b>Temaccord</b>
↓ Cap 20 mg – 1% DV May-20 to 2022 .....	16.38	5	<b>Temaccord</b>
↓ Cap 100 mg – 1% DV May-20 to 2022 .....	35.98	5	<b>Temaccord</b>
↓ Cap 140 mg – 1% DV May-20 to 2022 .....	50.12	5	<b>Temaccord</b>
↓ Cap 250 mg – 1% DV May-20 to 2022 .....	86.34	5	<b>Temaccord</b>

### → **Restricted (RS1645)**

### Initiation – High grade gliomas

Re-assessment required after 12 months

All of the following:

- 1 Either:
  - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
  - 1.2 Patient has newly diagnosed anaplastic astrocytoma\*;
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m<sup>2</sup> per day.

### Continuation – High grade gliomas

Re-assessment required after 12 months

Either:

- 1 Both:
  - 1.1 Patient has glioblastoma multiforme; and
  - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
  - 2.1 Patient has anaplastic astrocytoma\*;
  - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
  - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

### Initiation – Neuroendocrine tumours

Re-assessment required after 9 months

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour\*;
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m<sup>2</sup> per day; and
- 4 Temozolomide to be discontinued at disease progression.

### Continuation – Neuroendocrine tumours

Re-assessment required after 6 months

Both:

- 1 No evidence of disease progression; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- The treatment remains appropriate and the patient is benefitting from treatment.

## Initiation – ewing's sarcoma

*Re-assessment required after 9 months*

Patient has relapse or refractory Ewing's sarcoma.

## Continuation – ewing's sarcoma

*Re-assessment required after 6 months*

Both:

- No evidence of disease progression; and
- The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a \* is an unapproved indication. Temozolomide is not funded for the treatment of relapsed high grade glioma.

THALIDOMIDE – **Restricted** see terms [below](#)

⚡ Cap 50 mg .....	378.00	28	Thalomid
⚡ Cap 100 mg .....	756.00	28	Thalomid

➡ **Restricted (RS1192)**

## Initiation

*Re-assessment required after 12 months*

Any of the following:

- The patient has multiple myeloma; or
- The patient has systemic AL amyloidosis\*; or
- The patient has erythema nodosum leprosum.

## Continuation

Patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen

Indication marked with \* is an unapproved indication

## TRETINOIN

Cap 10 mg .....	479.50	100	Vesanoid
-----------------	--------	-----	----------

VENETOCLAX – **Restricted** see terms [below](#)

⚡ Tab 14 x 10 mg, 7 x 50 mg, 21 x 100 mg .....	1,771.86	42	Venclexta
⚡ Tab 10 mg .....	95.78	14	Venclexta
⚡ Tab 50 mg .....	239.44	7	Venclexta
⚡ Tab 100 mg .....	8,209.41	120	Venclexta

➡ **Restricted (RS1713)**

## Initiation – relapsed/refractory chronic lymphocytic leukaemia

Haematologist

*Re-assessment required after 7 months*

All of the following:

- Patient has chronic lymphocytic leukaemia requiring treatment; and
- Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- Patient has not previously received funded venetoclax; and
- The patient's disease has relapsed within 36 months of previous treatment; and
- Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- Patient has an ECOG performance status of 0-2.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

### Continuation – relapsed/refractory chronic lymphocytic leukaemia

Haematologist

*Re-assessment required after 6 months*

Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

### Initiation – previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*

Haematologist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

### Continuation – previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*

Haematologist

*Re-assessment required after 6 months*

The treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)\* and B-cell prolymphocytic leukaemia (B-PLL)\*. Indications marked with \* are unapproved indications.

## Platinum Compounds

### CARBOPLATIN

Inj 10 mg per ml, 45 ml vial – **1% DV Jun-19 to 2021** .....45.20      1      **Carboplatin Ebewe**

### CISPLATIN

Inj 1 mg per ml, 50 ml vial..... 12.29      1      **DBL Cisplatin**

Inj 1 mg per ml, 100 ml vial – **1% DV Sep-18 to 2021** ..... 19.70      1      **DBL Cisplatin**

*(DBL Cisplatin Inj 1 mg per ml, 50 ml vial to be delisted 1 April 2021)*

### OXALIPLATIN

Inj 5 mg per ml, 20 ml vial – **1% DV Feb-20 to 2021** .....46.32      1      **Oxaliplatin Accord**

## Protein-Tyrosine Kinase Inhibitors

### ALECTINIB – Restricted see terms [below](#)

↓ Cap 150 mg .....7,935.00      224      **Alecensa**

➔ **Restricted (RS1712)**

#### Initiation

*Re-assessment required after 6 months*

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test; and
- 3 Patient has an ECOG performance score of 0-2.

#### Continuation

*Re-assessment required after 6 months*

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>DASATINIB – Restricted</b> see terms <a href="#">below</a>			
⚡ Tab 20 mg .....	3,774.06	60	Sprycel
⚡ Tab 50 mg .....	6,214.20	60	Sprycel
⚡ Tab 70 mg .....	7,692.58	60	Sprycel

➡ **Restricted (RS1685)**

## Initiation

Haematologist or any relevant practitioner on the recommendation of a haematologist

*Re-assessment required after 6 months*

Any of the following:

- 1 Both:
  - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
  - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
  - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
  - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
  - 3.1 The patient has a diagnosis of CML in chronic phase; and
  - 3.2 Maximum dose of 100 mg/day; and
  - 3.3 Any of the following:
    - 3.3.1 Patient has documented treatment failure\* with imatinib; or
    - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
    - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
    - 3.3.4 Patients is enrolled in the KISS study\*\* and requires dasatinib treatment according to the study protocol.

## Continuation

Haematologist or any relevant practitioner on the recommendation of a haematologist

*Re-assessment required after 6 months*

All of the following:

- 1 Lack of treatment failure while on dasatinib\*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: \*treatment failure for CML as defined by Leukaemia Net Guidelines. \*\*Kinase-Inhibition Study with Sprycel Start-up

[https://www.cancertrialsnz.ac.nz/kiss/](https://www.cancertrials.nz.ac.nz/kiss/)

**ERLOTINIB – Restricted** see terms [below](#)

⚡ Tab 100 mg .....	764.00	30	Tarceva
⚡ Tab 150 mg .....	1,146.00	30	Tarceva

➡ **Restricted (RS1747)**

## Initiation

*Re-assessment required after 4 months*

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Either:
  - 3.1 Patient is treatment naive; or
  - 3.2 Both:
    - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
    - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

### Continuation

*Re-assessment required after 6 months*

Both:

- 1 Radiological assessment (preferably including CT scan) indicates NSCLC has not progressed; and
- 2 Erlotinib is to be given for a maximum of 3 months.

### Continuation – pandemic circumstances

*Re-assessment required after 6 months*

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular renewal requirements cannot be met due to COVID-19 constraints on the health sector.

GEFITINIB – **Restricted** see terms [below](#)

↓ Tab 250 mg ..... 1,700.00 30 Iressa

→ **Restricted (RS1748)**

### Initiation

*Re-assessment required after 4 months*

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
  - 2.1 Patient is treatment naive; or
  - 2.2 Both:
    - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
    - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

### Continuation

*Re-assessment required after 6 months*

Both:

- 1 Radiological assessment (preferably including CT scan) indicates NSCLC has not progressed; and
- 2 Gefitinib is to be given for a maximum of 3 months.

### Continuation – pandemic circumstances

*Re-assessment required after 6 months*

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains; and
- 2 Gefitinib to be discontinued at progression; and
- 3 The regular renewal requirements cannot be met due to COVID-19 constraints on the health sector.

### IMATINIB MESILATE

Imatinib-AFT is not a registered for the treatment of Gastro Intestinal Stromal Tumours (GIST). The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule

↓ Tab 100 mg ..... 2,400.00 60 Glivec

→ **Restricted (RS1402)**

### Initiation

*Re-assessment required after 12 months*

Both:

- 1 Patient has diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- tumour (GIST); and
- 2 Maximum dose of 400 mg/day.

## Continuation

*Re-assessment required after 12 months*

Adequate clinical response to treatment with imatinib (prescriber determined).

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule.

Cap 100 mg.....	98.00	60	Imatinib-AFT
Cap 400 mg – 1% DV Jun-21 to 2023.....	197.50	30	Imatinib-AFT
	84.79		<b>Imatinib-Rex</b>

(Imatinib-AFT Cap 400 mg to be delisted 1 June 2021)

LAPATINIB – **Restricted** see terms [below](#)

⚡ Tab 250 mg .....	1,899.00	70	Tykerb
--------------------	----------	----	--------

(Tykerb Tab 250 mg to be delisted 1 June 2021)

➡ **Restricted (RS1197)**

## Initiation

*Re-assessment required after 12 months*

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The patient has not previously received trastuzumab treatment for HER 2 positive metastatic breast cancer; and
  - 1.3 Lapatinib not to be given in combination with trastuzumab; and
  - 1.4 Lapatinib to be discontinued at disease progression; or
- 2 All of the following:
  - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 2.2 The patient started trastuzumab for metastatic breast cancer but discontinued trastuzumab within 3 months of starting treatment due to intolerance; and
  - 2.3 The cancer did not progress whilst on trastuzumab; and
  - 2.4 Lapatinib not to be given in combination with trastuzumab; and
  - 2.5 Lapatinib to be discontinued at disease progression.

## Continuation

*Re-assessment required after 12 months*

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB – **Restricted** see terms [below](#)

⚡ Cap 150 mg.....	4,680.00	120	Tasigna
⚡ Cap 200 mg.....	6,532.00	120	Tasigna

➡ **Restricted (RS1437)**

## Initiation

Haematologist

*Re-assessment required after 6 months*

All of the following:

continued...

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

continued...

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Either:
  - 2.1 Patient has documented CML treatment failure\* with imatinib; or
  - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: \*treatment failure as defined by Leukaemia Net Guidelines.

### Continuation

Haematologist

*Re-assessment required after 6 months*

All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PALBOCICLIB – **Restricted** see terms [below](#)

↓ Cap 75 mg.....	4,000.00	21	Ibrance
↓ Cap 100 mg.....	4,000.00	21	Ibrance
↓ Cap 125 mg.....	4,000.00	21	Ibrance

→ **Restricted (RS1731)**

### Initiation

Medical oncologist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Either:
  - second or subsequent line setting
  - 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
  - 4.2 Both:
    - first line setting
    - 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
    - 4.2.2 Either:
      - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
      - 4.2.2.2 All of the following:
        - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
        - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
        - 4.2.2.2.3 There is no evidence of progressive disease; and
      - 4.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

### Continuation

Medical oncologist

*Re-assessment required after 12 months*

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefiting from treatment.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PAZOPANIB – Restricted</b> see terms <a href="#">below</a>			
⚡ Tab 200 mg .....	1,334.70	30	Votrient
⚡ Tab 400 mg .....	2,669.40	30	Votrient

➡ **Restricted (RS1198)**

## Initiation

*Re-assessment required after 3 months*

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 Both:
    - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
    - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
- 5 All of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; and
  - 5.2 Haemoglobin level < lower limit of normal; and
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); and
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; and
  - 5.5 Karnofsky performance score of less than or equal to 70; and
  - 5.6 2 or more sites of organ metastasis.

## Continuation

*Re-assessment required after 3 months*

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

**RUXOLITINIB – Restricted** see terms [below](#)

⚡ Tab 5 mg .....	2,500.00	56	Jakavi
⚡ Tab 15 mg .....	5,000.00	56	Jakavi
⚡ Tab 20 mg .....	5,000.00	56	Jakavi

➡ **Restricted (RS1726)**

## Initiation

Haematologist

*Re-assessment required after 12 months*

All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
  - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
  - 2.2 Both:
    - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

DIPSS; and

2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and

3 A maximum dose of 20 mg twice daily is to be given.

#### Continuation

Relevant specialist or medical practitioner on the recommendation of a Relevant specialist

*Re-assessment required after 12 months*

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB – **Restricted** see terms [below](#)

↓ Cap 12.5 mg .....	2,315.38	28	Sutent
↓ Cap 25 mg .....	4,630.77	28	Sutent
↓ Cap 50 mg .....	9,261.54	28	Sutent

→ **Restricted (RS1749)**

#### Initiation – RCC

*Re-assessment required after 3 months*

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
  - 2.4 Both:
    - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
    - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
- 5 All of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; and
  - 5.2 Haemoglobin level < lower limit of normal; and
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); and
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; and
  - 5.5 Karnofsky performance score of less than or equal to 70; and
  - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Notes: RCC - Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

#### Continuation – RCC

*Re-assessment required after 3 months*

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

## Initiation – GIST

*Re-assessment required after 3 months*

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
  - 2.1 The patient's disease has progressed following treatment with imatinib; or
  - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

## Continuation – GIST

*Re-assessment required after 6 months*

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
  - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
  - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non-measurable disease); or
  - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

## Continuation – GIST pandemic circumstances

*Re-assessment required after 6 months*

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Note: GIST - It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

## Taxanes

### DOCETAXEL

Inj 10 mg per ml, 2 ml vial.....	12.40	1	DBL Docetaxel
Inj 10 mg per ml, 8 ml vial.....	26.95	1	DBL Docetaxel

*(DBL Docetaxel Inj 10 mg per ml, 2 ml vial to be delisted 1 June 2021)*

### PACLITAXEL

Inj 6 mg per ml, 5 ml vial.....	47.30	5	Paclitaxel Ebewe
Inj 6 mg per ml, 16.7 ml vial – 1% DV Nov-20 to 2023 .....	24.00	1	<b>Paclitaxel Ebewe</b>
Inj 6 mg per ml, 25 ml vial.....	26.69	1	Paclitaxel Ebewe
Inj 6 mg per ml, 50 ml vial – 1% DV Nov-20 to 2023 .....	44.00	1	<b>Paclitaxel Ebewe</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## Treatment of Cytotoxic-Induced Side Effects

### CALCIUM FOLINATE

Tab 15 mg .....	114.69	10	DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml ampoule .....			
Inj 10 mg per ml, 5 ml ampoule .....	18.25	5	Calcium Folate Ebewe
Inj 10 mg per ml, 5 ml vial – 1% DV Jan-20 to 2022 .....	7.28	1	<b>Calcium Folate</b> <b>Sandoz</b>
Inj 10 mg per ml, 10 ml vial – 1% DV Jan-20 to 2022 .....	9.49	1	<b>Calcium Folate</b> <b>Sandoz</b>
Inj 10 mg per ml, 30 ml vial .....	22.51	1	Calcium Folate Ebewe
Inj 10 mg per ml, 35 ml vial – 1% DV Nov-19 to 2022 .....	25.14	1	<b>Calcium Folate</b> <b>Sandoz</b>
Inj 10 mg per ml, 100 ml vial – 1% DV Mar-20 to 2022 .....	72.00	1	<b>Calcium Folate</b> <b>Sandoz</b>

DEXRAZOXANE – **Restricted** see terms [below](#)

↓ Inj 500 mg

→ **Restricted (RS1695)**

e.g. *Cardioxane*

### Initiation

Medical oncologist, paediatric oncologist, haematologist or paediatric haematologist

All of the following:

- 1 Patient is to receive treatment with high dose anthracycline given with curative intent; and
- 2 Based on current treatment plan, patient's cumulative lifetime dose of anthracycline will exceed 250mg/m<sup>2</sup> doxorubicin equivalent or greater; and
- 3 Dextrazoxane to be administered only whilst on anthracycline treatment; and
- 4 Either:
  - 4.1 Treatment to be used as a cardioprotectant for a child or young adult; or
  - 4.2 Treatment to be used as a cardioprotectant for secondary malignancy.

### MESNA

Tab 400 mg – 1% DV Nov-19 to 2022 .....	314.00	50	<b>Uromitexan</b>
Tab 600 mg – 1% DV Nov-19 to 2022 .....	448.50	50	<b>Uromitexan</b>
Inj 100 mg per ml, 4 ml ampoule – 1% DV Nov-19 to 2022 .....	177.45	15	<b>Uromitexan</b>
Inj 100 mg per ml, 10 ml ampoule – 1% DV Nov-19 to 2022 .....	407.40	15	<b>Uromitexan</b>

## Vinca Alkaloids

### VINBLASTINE SULPHATE

Inj 1 mg per ml, 10 ml vial .....	270.37	5	Hospira
-----------------------------------	--------	---	---------

### VINCISTINE SULPHATE

Inj 1 mg per ml, 1 ml vial .....	74.52	5	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial .....	102.73	5	DBL Vincristine Sulfate

### VINOIRELBINE

Inj 10 mg per ml, 1 ml vial .....	12.00	1	Navelbine
Inj 10 mg per ml, 5 ml vial .....	56.00	1	Navelbine

## Endocrine Therapy

ABIRATERONE ACETATE – **Restricted** see terms [on the next page](#)

↓ Tab 250 mg ..... 4,276.19 120 Zytiga

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## ➔ Restricted (RS1746)

### Initiation

Medical oncologist, radiation oncologist or urologist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
  - 4.1 All of the following:
    - 4.1.1 Patient is symptomatic; and
    - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
    - 4.1.3 Patient has ECOG performance score of 0-1; and
    - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
  - 4.2 All of the following:
    - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
    - 4.2.2 Patient has ECOG performance score of 0-2; and
    - 4.2.3 Patient has not had prior treatment with abiraterone.

### Continuation

Medical oncologist, radiation oncologist or urologist

*Re-assessment required after 6 months*

All of the following:

- 1 No evidence of clinical disease progression; and
- 2 No initiation of taxane chemotherapy with abiraterone; and
- 3 The treatment remains appropriate and the patient is benefiting from treatment.

### BICALUTAMIDE

Tab 50 mg .....	3.80	28	Binarex
-----------------	------	----	---------

### FLUTAMIDE

Tab 250 mg .....	119.50	100	Flutamin
------------------	--------	-----	----------

### FULVESTRANT – Restricted see terms below

⚡ Inj 50 mg per ml, 5 ml prefilled syringe .....	1,068.00	2	Faslodex
--	----------	---	----------

## ➔ Restricted (RS1732)

### Initiation

Medical oncologist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

### Continuation

Medical oncologist

*Re-assessment required after 6 months*

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 No evidence of disease progression.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MEGESTROL ACETATE			
Tab 160 mg – 1% DV Oct-18 to 2021.....	63.53	30	<b>Apo-Megestrol</b>
OCTREOTIDE – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Inj 50 mcg per ml, 1 ml ampoule .....	30.64	5	DBL Octreotide
↓ Inj 100 mcg per ml, 1 ml ampoule .....	18.69	5	DBL Octreotide
↓ Inj 500 mcg per ml, 1 ml ampoule .....	72.50	5	DBL Octreotide
↓ Inj 10 mg vial .....	1,772.50	1	Sandostatin LAR
↓ Inj 20 mg vial .....	2,358.75	1	Sandostatin LAR
↓ Inj 30 mg vial .....	2,951.25	1	Sandostatin LAR

→ **Restricted (RS1744)**

#### Initiation – Malignant bowel obstruction

All of the following:

- 1 The patient has nausea\* and vomiting\* due to malignant bowel obstruction\*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with \* are unapproved indications

#### Initiation – acromegaly

*Re-assessment required after 3 months*

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
  - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
  - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
  - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

#### Continuation – acromegaly

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks.

#### Initiation – Other indications

Any of the following:

- 1 VIPomas and glucagonomas - for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
  - 2.1 Gastrinoma; and
  - 2.2 Either:
    - 2.2.1 Patient has failed surgery; or
    - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
  - 3.1 Insulinomas; and
  - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:

continued...

## ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
- 5.2 Disabling symptoms not controlled by maximal medical therapy.

### Continuation – Acromegaly - pandemic circumstances

*Re-assessment required after 6 months*

All of the following:

- 1 Patient has acromegaly; and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Note: restriction applies only to the long-acting formulations of octreotide

### TAMOXIFEN CITRATE

Tab 10 mg – 1% DV Nov-20 to 2023	15.00	60	Tamoxifen Sandoz
Tab 20 mg – 1% DV Nov-20 to 2023	6.65	60	Tamoxifen Sandoz

## Aromatase Inhibitors

### ANASTROZOLE

Tab 1 mg – 1% DV Apr-21 to 2023	4.55	30	Anastrole
	5.04		Rolin

*(Rolin Tab 1 mg to be delisted 1 April 2021)*

### EXEMESTANE

Tab 25 mg	14.50	30	Pfizer Exemestane
-----------	-------	----	-------------------

### LETROZOLE

Tab 2.5 mg – 1% DV Nov-18 to 2021	4.68	30	Letrole
-----------------------------------	------	----	---------

## Imaging Agents

### AMINOLEVULINIC ACID HYDROCHLORIDE – Restricted see terms below

⚡ Powder for oral soln, 30 mg per ml, 1.5 g vial	4,400.00	1	Gliolan
	44,000.00	10	Gliolan

➡ **Restricted (RS1565)**

### Initiation – high grade malignant glioma

All of the following:

- 1 Patient has newly diagnosed, untreated, glioblastoma multiforme; and
- 2 Treatment to be used as adjuvant to fluorescence-guided resection; and
- 3 Patient's tumour is amenable to complete resection.

## Immunosuppressants

### Calcineurin Inhibitors

### CICLOSPORIN

Cap 25 mg	44.63	50	Neoral
Cap 50 mg	88.91	50	Neoral
Cap 100 mg	177.81	50	Neoral
Oral liq 100 mg per ml	198.13	50 ml	Neoral
Inj 50 mg per ml, 5 ml ampoule	276.30	10	Sandimmun

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>TACROLIMUS – Restricted</b> see terms <a href="#">below</a>			
↓ Cap 0.5 mg.....	49.60	100	Tacrolimus Sandoz
↓ Cap 0.75 mg.....	99.30	100	Tacrolimus Sandoz
↓ Cap 1 mg.....	84.30	100	Tacrolimus Sandoz
↓ Cap 5 mg.....	248.20	50	Tacrolimus Sandoz
↓ Inj 5 mg per ml, 1 ml ampoule			

➔ **Restricted (RS1651)**

**Initiation – organ transplant recipients**

Any specialist

For use in organ transplant recipients.

**Initiation – non-transplant indications\***

Any specialist

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with \* are unapproved indications

## Fusion Proteins

**ETANERCEPT – Restricted** see terms [below](#)

↓ Inj 25 mg vial – 5% DV Sep-19 to 2024.....	690.00	4	<b>Enbrel</b>
↓ Inj 50 mg autoinjector – 5% DV Sep-19 to 2024.....	1,050.00	4	<b>Enbrel</b>
↓ Inj 50 mg syringe – 5% DV Sep-19 to 2024.....	1,050.00	4	<b>Enbrel</b>

➔ **Restricted (RS1783)**

**Initiation – polyarticular course juvenile idiopathic arthritis**

Rheumatologist or named specialist

*Re-assessment required after 6 months*

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
  - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or

2 All of the following:

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
  - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

## **Continuation – polyarticular course juvenile idiopathic arthritis**

Rheumatologist or named specialist

*Re-assessment required after 6 months*

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

## **Initiation – oligoarticular course juvenile idiopathic arthritis**

Rheumatologist or named specialist

*Re-assessment required after 6 months*

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

## **Continuation – oligoarticular course juvenile idiopathic arthritis**

Rheumatologist or named specialist

*Re-assessment required after 6 months*

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

## **Initiation – rheumatoid arthritis**

Rheumatologist

*Re-assessment required after 6 months*

Either:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
  - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis; or

2 All of the following:

- 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 2.5 Any of the following:
  - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
  - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
  - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
  - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
  - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Continuation – rheumatoid arthritis**

Rheumatologist

*Re-assessment required after 6 months*

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

## Initiation – ankylosing spondylitis

Rheumatologist

*Re-assessment required after 6 months*

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
  - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
  - 2.6 Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of starting treatment.

Average normal chest expansion corrected for age and gender:

Age	Male	Female
18-24	7.0 cm	5.5 cm
25-34	7.5 cm	5.5 cm
35-44	6.5 cm	4.5 cm
45-54	6.0 cm	5.0 cm
55-64	5.5 cm	4.0 cm
65-74	4.0 cm	4.0 cm
75+	3.0 cm	2.5 cm

## Continuation – ankylosing spondylitis

Rheumatologist

*Re-assessment required after 6 months*

All of the following:

- 1 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

**Initiation – psoriatic arthritis**

Rheumatologist

*Re-assessment required after 6 months*

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
  - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
  - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Continuation – psoriatic arthritis**

Rheumatologist

*Re-assessment required after 6 months*

Both:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initiation – severe chronic plaque psoriasis, prior TNF use**

Dermatologist

*Limited to 4 months treatment*

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab; or
  - 2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; and
- 3 Patient must be reassessed for continuation after 3 doses.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

## **Initiation – severe chronic plaque psoriasis, treatment-naïve**

Dermatologist

*Limited to 4 months treatment*

All of the following:

- 1 Either:
  - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

## **Continuation – severe chronic plaque psoriasis**

Dermatologist

*Re-assessment required after 6 months*

Both:

- 1 Either:
  - 1.1 Both:
    - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 1.1.2 Either:
      - 1.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-etanercept treatment baseline value; or
      - 1.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
  - 1.2 Both:
    - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
    - 1.2.2 Either:
      - 1.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
      - 1.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-etanercept treatment baseline value; and
- 2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

continued...

	Price		Brand or
(ex man. excl. GST)			Generic
\$	Per		Manufacturer

continued...

**Initiation – pyoderma gangrenosum**

Dermatologist

All of the following:

- 1 Patient has pyoderma gangrenosum\*;
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with \* are unapproved indications.

**Continuation – pyoderma gangrenosum**

Dermatologist

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

**Initiation – adult-onset Still's disease**

Rheumatologist

*Re-assessment required after 6 months*

Either:

- 1 Both:
  - 1.1 Either:
    - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
    - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the Section H rules; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
    - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

**Continuation – adult-onset Still's disease**

Rheumatologist

*Re-assessment required after 6 months*

The patient has a sustained improvement in inflammatory markers and functional status.

**Initiation – undifferentiated spondyloarthritis**

Rheumatologist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis\* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
  - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with \* are unapproved indications.

## Continuation – undifferentiated spondyloarthritis

Rheumatologist or medical practitioner on the recommendation of a Rheumatologist

*Re-assessment required after 6 months*

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

## Monoclonal Antibodies

ABCIXIMAB – **Restricted** see terms [below](#)

⚡ Inj 2 mg per ml, 5 ml vial.....579.53 1 ReoPro

*(ReoPro Inj 2 mg per ml, 5 ml vial to be delisted 1 January 2021)*

➡ **Restricted** ([RS1202](#))

### Initiation

Either:

- 1 For use in patients with acute coronary syndromes undergoing percutaneous coronary intervention; or
- 2 For use in patients undergoing intra-cranial intervention.

ADALIMUMAB – **Restricted** see terms [below](#)

⚡ Inj 20 mg per 0.4 ml syringe ..... 1,599.96 2 Humira  
 ⚡ Inj 40 mg per 0.8 ml pen..... 1,599.96 2 HumiraPen  
 ⚡ Inj 40 mg per 0.8 ml syringe ..... 1,599.96 2 Humira

➡ **Restricted** ([RS1784](#))

### Initiation – polyarticular course juvenile idiopathic arthritis

Rheumatologist or named specialist

*Re-assessment required after 6 months*

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- arthritis (JIA); and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
  - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for polyarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

**Continuation – polyarticular course juvenile idiopathic arthritis**

Rheumatologist or named specialist

*Re-assessment required after 6 months*

- Both:
- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2 Either:
    - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
    - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

**Initiation – oligoarticular course juvenile idiopathic arthritis**

Rheumatologist or named specialist

*Re-assessment required after 6 months*

- Either:
- 1 Both:
    - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
    - 1.2 Either:
      - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
      - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for oligoarticular course JIA; or
  - 2 All of the following:
    - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
    - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
    - 2.3 Any of the following:
      - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
      - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
      - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

## **Continuation – oligoarticular course juvenile idiopathic arthritis**

Rheumatologist or named specialist

*Re-assessment required after 6 months*

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

## **Initiation – fistulising Crohn's disease**

Gastroenterologist

*Re-assessment required after 4 months*

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Either:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

## **Continuation – fistulising Crohn's disease**

Gastroenterologist

*Re-assessment required after 6 months*

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

## **Initiation – Crohn's disease - adults**

Gastroenterologist

*Re-assessment required after 3 months*

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

## **Continuation – Crohn's disease - adults**

Gastroenterologist

*Re-assessment required after 3 months*

Both:

- 1 Either:
  - 1.1 Either:

continued...

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

continued...

- 1.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
- 1.1.2 CDAI score is 150 or less; or
- 1.2 Both:
  - 1.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
  - 1.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

## **Initiation – Crohn's disease - children**

Gastroenterologist

*Re-assessment required after 3 months*

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

## **Continuation – Crohn's disease - children**

Gastroenterologist

*Re-assessment required after 3 months*

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 100 points from the PCDAI score when the patient was initiated on adalimumab; or
  - 1.2 PCDAI score is 150 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

## **Initiation – rheumatoid arthritis**

Rheumatologist

*Re-assessment required after 6 months*

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
  - 2.5 Any of the following:

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
- 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
  - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
  - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

## Continuation – rheumatoid arthritis

Rheumatologist

*Re-assessment required after 6 months*

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

## Initiation – ankylosing spondylitis

Rheumatologist

*Re-assessment required after 6 months*

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
  - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
- 2.5 Either:
  - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by

continued...

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

continued...

the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or

- 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and

2.6 Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of starting treatment.

Average normal chest expansion corrected for age and gender:

Age	Male	Female
18-24	7.0 cm	5.5 cm
25-34	7.5 cm	5.5 cm
35-44	6.5 cm	4.5 cm
45-54	6.0 cm	5.0 cm
55-64	5.5 cm	4.0 cm
65-74	4.0 cm	4.0 cm
75+	3.0 cm	2.5 cm

### Continuation – ankylosing spondylitis

Rheumatologist

*Re-assessment required after 6 months*

All of the following:

- Following 12 weeks' initial treatment and subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- Adalimumab to be administered at doses no greater than 40 mg every 14 days.

### Initiation – psoriatic arthritis

Rheumatologist

*Re-assessment required after 6 months*

Either:

- Both:
  - The patient has had an initial Special Authority approval for etanercept for psoriatic arthritis; and
  - Either:
    - The patient has experienced intolerable side effects from etanercept; or
    - The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for psoriatic arthritis; or
- All of the following:
  - Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - Either:
    - Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
  - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
  - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

## Continuation – psoriatic arthritis

Rheumatologist

*Re-assessment required after 6 months*

Both:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

## Initiation – plaque psoriasis, prior TNF use

Dermatologist

*Limited to 4 months treatment*

Both:

- 1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from etanercept; or
  - 2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis.

## Initiation – plaque psoriasis, treatment-naïve

Dermatologist

*Limited to 4 months treatment*

All of the following:

- 1 Either:
  - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot,

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

**Continuation – plaque psoriasis**

Dermatologist

*Re-assessment required after 6 months*

Both:

1 Either:

1.1 Both:

1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and

1.1.2 Either:

1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or

1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or

1.2 Both:

1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and

1.2.2 Either:

1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or

1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-etanercept treatment baseline value; and

2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initiation – pyoderma gangrenosum**

Dermatologist

All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with \* are unapproved indications.

**Continuation – pyoderma gangrenosum**

Dermatologist

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

**Initiation – adult-onset Still's disease**

Rheumatologist

*Re-assessment required after 6 months*

Either:

1 Both:

1.1 Either:

1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the Section H rules; and

1.2 Either:

1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or

1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or

2 All of the following:

2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and

2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and

2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

## Continuation – adult-onset Still's disease

Rheumatologist

*Re-assessment required after 6 months*

The patient has a sustained improvement in inflammatory markers and functional status.

## Initiation – severe Behcet's disease

Any relevant practitioner

*Re-assessment required after 3 months*

All of the following:

1 The patient has severe Behcet's disease that is significantly impacting the patient's quality of life (see Notes); and

2 Either:

2.1 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has not responded adequately to treatment with infliximab (see Notes); or

2.2 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has experienced intolerable side effects from treatment with infliximab; and

3 The patient is experiencing significant loss of quality of life; and

4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: Behcet's disease diagnosed according to the International Study Group for Behcet's disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al, J Rheumatol. 2004;31:931-7.

## Continuation – severe Behcet's disease

Any relevant practitioner

*Re-assessment required after 6 months*

Both:

1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and

2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

## Initiation – severe ocular inflammation

*Re-assessment required after 4 months*

Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; and

1.2 Either:

1.2.1 The patient has experienced intolerable side effects from infliximab; or

1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for severe ocular inflammation; or

2 Both:

2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

2.2 Any of the following:

- 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
- 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
- 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

**Continuation – severe ocular inflammation**

*Re-assessment required after 12 months*

Both:

1 Any of the following:

- 1.1 The patient has had a good clinical response following 3 initial doses; or
- 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and

2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

**Initiation – chronic ocular inflammation**

*Re-assessment required after 4 months*

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
  - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
  - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
  - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
  - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

**Continuation – chronic ocular inflammation**

*Re-assessment required after 12 months*

Both:

1 Any of the following:

- 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and

- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

## Initiation – hidradenitis suppurativa

Dermatologist

*Re-assessment required after 4 months*

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 The patient has 3 or more active lesions (e.g. inflammatory nodules, abscesses, draining fistulae); and
- 4 The patient has a Dermatology Quality of Life Index of 10 or more and the assessment is no more than 1 month old at time of application; and
- 5 Following the initial loading doses, adalimumab is to be administered at doses no greater than 40mg every 7 days.

## Continuation – hidradenitis suppurativa

Dermatologist

*Re-assessment required after 6 months*

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

AFLIBERCEPT – **Restricted** see terms [below](#)

‡ Inj 40 mg per ml, 0.1 ml vial..... 1,250.00 1 Eylea

➔ **Restricted (RS1659)**

## Initiation – Wet Age Related Macular Degeneration

Ophthalmologist

*Re-assessment required after 3 months*

Either:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Wet age-related macular degeneration (wet AMD); or
    - 1.1.2 Polypoidal choroidal vasculopathy; or
    - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
  - 1.2 Either:
    - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
    - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
  - 1.3 There is no structural damage to the central fovea of the treated eye; and
  - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
  - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.2 Patient has previously\* (\*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

### Continuation – Wet Age Related Macular Degeneration

Ophthalmologist

*Re-assessment required after 12 months*

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

### Initiation – Diabetic Macular Oedema

Ophthalmologist

*Re-assessment required after 4 months*

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 – 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

### Continuation – Diabetic Macular Oedema

Ophthalmologist

*Re-assessment required after 12 months*

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with aflibercept, patient has retrialled with at least one injection of bevacizumab and had no response.

**BASILIXIMAB – Restricted** see terms [below](#)

↓ Inj 20 mg vial .....2,560.00 1 Simulect

➔ **Restricted (RS1203)**

### Initiation

For use in solid organ transplants.

**BEVACIZUMAB – Restricted** see terms [below](#)

↓ Inj 25 mg per ml, 4 ml vial

↓ Inj 25 mg per ml, 16 ml vial

➔ **Restricted (RS1691)**

### Initiation – Recurrent Respiratory Papillomatosis

Otolaryngologist

*Re-assessment required after 12 months*

All of the following:

- 1 Maximum of 6 doses; and
- 2 The patient has recurrent respiratory papillomatosis; and
- 3 The treatment is for intra-lesional administration.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

## Continuation – Recurrent Respiratory Papillomatosis

Otolaryngologist

*Re-assessment required after 12 months*

All of the following:

- 1 Maximum of 6 doses; and
- 2 The treatment is for intra-lesional administration; and
- 3 There has been a reduction in surgical treatments or disease regrowth as a result of treatment.

## Initiation – ocular conditions

Either:

- 1 Ocular neovascularisation; or
- 2 Exudative ocular angiopathy.

CETUXIMAB – **Restricted** see terms [below](#)

⚡ Inj 5 mg per ml, 20 ml vial.....	364.00	1	Erbitux
⚡ Inj 5 mg per ml, 100 ml vial.....	1,820.00	1	Erbitux

➡ **Restricted (RS1613)**

## Initiation

Medical oncologist

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB – **Restricted** see terms [below](#)

⚡ Inj 100 mg.....	806.00	1	Remicade
-------------------	--------	---	----------

➡ **Restricted (RS1772)**

## Initiation – Graft vs host disease

Patient has steroid-refractory acute graft vs. host disease of the gut.

## Initiation – rheumatoid arthritis

Rheumatologist

*Re-assessment required after 4 months*

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

## Continuation – rheumatoid arthritis

Rheumatologist

*Re-assessment required after 6 months*

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline

continued...

	Price	Brand or
(ex man. excl. GST)		Generic
\$	Per	Manufacturer

continued...

- and a clinically significant response to treatment in the opinion of the physician; or
- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

#### **Initiation – ankylosing spondylitis**

Rheumatologist

*Re-assessment required after 3 months*

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

#### **Continuation – ankylosing spondylitis**

Rheumatologist

*Re-assessment required after 6 months*

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

#### **Initiation – psoriatic arthritis**

Rheumatologist

*Re-assessment required after 4 months*

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for psoriatic arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for psoriatic arthritis.

#### **Continuation – psoriatic arthritis**

Rheumatologist

*Re-assessment required after 6 months*

Both:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

#### **Initiation – severe ocular inflammation**

*Re-assessment required after 4 months*

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or

2 Both:

- 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
- 2.2 Any of the following:
  - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
  - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
  - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

## Continuation – severe ocular inflammation

*Re-assessment required after 12 months*

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

## Initiation – chronic ocular inflammation

*Re-assessment required after 4 months*

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
  - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
  - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
  - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at therapeutic dose; or
  - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

## Continuation – chronic ocular inflammation

*Re-assessment required after 12 months*

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions,

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

or resolution of uveitic cystoid macular oedema); or

- 3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

#### **Initiation – Pulmonary sarcoidosis**

Both:

- 1 Patient has life-threatening pulmonary sarcoidosis that is refractory to other treatments; and
- 2 Treatment is to be prescribed by, or has been recommended by, a physician with expertise in the treatment of pulmonary sarcoidosis.

#### **Initiation – Crohn's disease (adults)**

Gastroenterologist

*Re-assessment required after 3 months*

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

#### **Continuation – Crohn's disease (adults)**

Gastroenterologist

*Re-assessment required after 6 months*

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
  - 1.2 CDAI score is 150 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

#### **Initiation – Crohn's disease (children)**

Gastroenterologist

*Re-assessment required after 3 months*

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 5 Patient must be reassessed for continuation after 3 months of therapy.

## **Continuation – Crohn's disease (children)**

Gastroenterologist

*Re-assessment required after 6 months*

Both:

- 1 Any of the following:
  - 1.1 PDAI score has reduced by 10 points from the PDAI score when the patient was initiated on infliximab; or
  - 1.2 PDAI score is 15 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but PDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

## **Initiation – fistulising Crohn's disease**

Gastroenterologist

*Re-assessment required after 4 months*

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Either:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e).

## **Continuation – fistulising Crohn's disease**

Gastroenterologist

*Re-assessment required after 6 months*

Both:

- 1 Either:
  - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

## **Initiation – acute severe fulminant ulcerative colitis**

Gastroenterologist

*Limited to 6 weeks treatment*

Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

## **Continuation – severe fulminant ulcerative colitis**

Gastroenterologist

*Re-assessment required after 6 months*

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

**Initiation – severe ulcerative colitis**

Gastroenterologist

*Re-assessment required after 3 months*

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
  - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
  - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

**Continuation – severe ulcerative colitis**

Gastroenterologist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
  - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
  - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

**Initiation – plaque psoriasis**

Dermatologist

*Re-assessment required after 3 doses*

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
  - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, cyclosporin, or acitretin; and

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and

- 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

## Continuation – plaque psoriasis

Dermatologist

*Re-assessment required after 3 doses*

Both:

- 1 Either:

- 1.1 Both:

- 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
- 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or

- 1.2 Both:

- 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and

- 1.2.2 Either:

- 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
- 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value; and

- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

## Initiation – neurosarcoidosis

Neurologist

*Re-assessment required after 18 months*

All of the following:

- 1 Biopsy consistent with diagnosis of neurosarcoidosis; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
  - 4.1 IV cyclophosphamide has been tried; or
  - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

## Continuation – neurosarcoidosis

Neurologist

*Re-assessment required after 18 months*

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
  - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
  - 2.2 There has been a marked reduction in prednisone dose; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

### 2.3 Either:

2.3.1 There has been an improvement in MRI appearances; or

2.3.2 Marked improvement in other symptomology.

### Initiation – severe Behcet's disease

*Re-assessment required after 4 months*

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
  - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
  - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes:

- 1 Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.
- 2 Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

### Continuation – severe Behcet's disease

*Re-assessment required after 6 months*

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

### Initiation – pyoderma gangrenosum

Dermatologist

All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with \* are unapproved indications.

### Continuation – pyoderma gangrenosum

Dermatologist

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

MEPOLIZUMAB – **Restricted** see terms [below](#)

↓ Inj 100 mg vial ..... 1,638.00 1 Nucala

→ **Restricted (RS1733)**

### Initiation – Severe eosinophilic asthma

Respiratory physician or clinical immunologist

*Re-assessment required after 12 months*

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than  $0.5 \times 10^9$  cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
  - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
  - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment.

## Continuation – Severe eosinophilic asthma

Respiratory physician or clinical immunologist

*Re-assessment required after 2 years*

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
  - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
  - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB – **Restricted** see terms [below](#)

⚡ Inj 25 mg per ml, 40 ml vial.....5,910.00      1      Gazyva

➡ **Restricted (RS1550)**

## Initiation

Haematologist

*Limited to 6 months treatment*

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts\* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

\* greater than or equal to  $1.5 \times 10^9$ /L and platelets greater than or equal to  $75 \times 10^9$ /L

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
OMALIZUMAB – <b>Restricted</b> see terms <a href="#">below</a>			
↓ Inj 150 mg prefilled syringe.....	450.00	1	Xolair
↓ Inj 150 mg vial .....	450.00	1	Xolair

➔ **Restricted (RS1652)**

**Initiation – severe asthma**

Clinical immunologist or respiratory specialist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient must be aged 6 years or older ; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
  - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
  - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

**Continuation – severe asthma**

Respiratory specialist

*Re-assessment required after 6 months*

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

**Initiation – severe chronic spontaneous urticaria**

Clinical immunologist or dermatologist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
    - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; and
- 3 Any of the following:
  - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
  - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
  - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
  - 4.1 Treatment to be stopped if inadequate response\* following 4 doses; or
  - 4.2 Complete response\* to 6 doses of omalizumab.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

## Continuation – severe chronic spontaneous urticaria

Clinical immunologist or dermatologist

*Re-assessment required after 6 months*

Either:

- 1 Patient has previously had a complete response\* to 6 doses of omalizumab; or
- 2 Both:
  - 2.1 Patient has previously had a complete response\* to 6 doses of omalizumab; and
  - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: \*Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

**PERTUZUMAB – Restricted** see terms [below](#)

⚡ Inj 30 mg per ml, 14 ml vial.....3,927.00 1 Perjeta

➡ **Restricted (RS1551)**

## Initiation

*Re-assessment required after 12 months*

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
  - 2.1 Patient is chemotherapy treatment naive; or
  - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

## Continuation

*Re-assessment required after 12 months*

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

**RANIBIZUMAB – Restricted** see terms [below](#)

⚡ Inj 10 mg per ml, 0.23 ml vial

⚡ Inj 10 mg per ml, 0.3 ml vial

➡ **Restricted (RS1637)**

## Initiation – Wet Age Related Macular Degeneration

Ophthalmologist

*Re-assessment required after 3 months*

Either:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Wet age-related macular degeneration (wet AMD); or
    - 1.1.2 Polypoidal choroidal vasculopathy; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
- 1.2 Either:
  - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
  - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
- 1.3 There is no structural damage to the central fovea of the treated eye; and
- 1.4 Patient has not previously been treated with aflibercept for longer than 3 months; or
- 2 Patient has current approval to use aflibercept for treatment of wAMD and was found to be intolerant to aflibercept within 3 months.

## Continuation – Wet Age Related Macular Degeneration

Ophthalmologist

*Re-assessment required after 12 months*

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

RITUXIMAB (MABTHERA) – **Restricted** see terms [below](#)

⬇ Inj 10 mg per ml, 10 ml vial.....	1,075.50	2	Mabthera
⬇ Inj 10 mg per ml, 50 ml vial.....	2,688.30	1	Mabthera

➔ **Restricted (RS1785)**

## Initiation – rheumatoid arthritis - prior TNF inhibitor use

Rheumatologist

*Limited to 4 months treatment*

All of the following:

- 1 Both:
  - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
    - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

## Initiation – rheumatoid arthritis - TNF inhibitors contraindicated

Rheumatologist

*Limited to 4 months treatment*

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
  - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of cyclosporin; or
  - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
  - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
  - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
  - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
  - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

## Continuation – rheumatoid arthritis - re-treatment in 'partial responders' to rituximab

Rheumatologist

*Re-assessment required after 4 months*

All of the following:

- 1 Any of the following:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

## Continuation – rheumatoid arthritis - re-treatment in 'responders' to rituximab

Rheumatologist

*Re-assessment required after 4 months*

All of the following:

- 1 Either:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) – **Restricted** see terms [below](#)

↓ Inj 10 mg per ml, 10 ml vial.....	275.33	2	Riximyo
↓ Inj 10 mg per ml, 50 ml vial.....	688.20	1	Riximyo

→ **Restricted (RS1764)**

#### Initiation – haemophilia with inhibitors

Haematologist

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

#### Continuation – haemophilia with inhibitors

Haematologist

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

#### Initiation – post-transplant

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with \* are unapproved indications.

#### Continuation – post-transplant

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with \* are unapproved indications.

#### Initiation – indolent, low-grade lymphomas or hairy cell leukaemia\*

*Re-assessment required after 9 months*

Either:

- 1 Both:
  - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
  - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia\* requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

## **Continuation – indolent, low-grade lymphomas or hairy cell leukaemia\***

*Re-assessment required after 12 months*

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

## **Initiation – aggressive CD20 positive NHL**

Either:

- 1 All of the following:
  - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
  - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
  - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
  - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia.

## **Continuation – aggressive CD20 positive NHL**

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia.

## **Initiation – Chronic lymphocytic leukaemia**

*Re-assessment required after 12 months*

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
  - 2.1 The patient is rituximab treatment naive; or
  - 2.2 Either:
    - 2.2.1 The patient is chemotherapy treatment naive; or
    - 2.2.2 Both:
      - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
      - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
  - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
  - 4.1 The patient does not have chromosome 17p deletion CLL; or
  - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

**Continuation – Chronic lymphocytic leukaemia**

*Re-assessment required after 12 months*

Both:

- 1 Either:
  - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
  - 1.2 All of the following:
    - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
    - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
    - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
    - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

**Initiation – severe cold haemagglutinin disease (CHAD)**

Haematologist

*Re-assessment required after 8 weeks*

All of the following:

- 1 Patient has cold haemagglutinin disease\*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Continuation – severe cold haemagglutinin disease (CHAD)**

Haematologist

*Re-assessment required after 8 weeks*

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m<sup>2</sup> weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

## **Initiation – warm autoimmune haemolytic anaemia (warm AIHA)**

Haematologist

*Re-assessment required after 8 weeks*

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia\*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

## **Continuation – warm autoimmune haemolytic anaemia (warm AIHA)**

Haematologist

*Re-assessment required after 8 weeks*

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m<sup>2</sup> weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

## **Initiation – immune thrombocytopenic purpura (ITP)**

Haematologist

*Re-assessment required after 8 weeks*

All of the following:

- 1 Either:
  - 1.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per microlitre; or
  - 1.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
  - 2.1 Treatment with steroids and splenectomy have been ineffective; or
  - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
  - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

## **Continuation – immune thrombocytopenic purpura (ITP)**

Haematologist

*Re-assessment required after 8 weeks*

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m<sup>2</sup> weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura\*; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2.2 An initial response lasting at least 12 months was demonstrated; and
- 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

**Initiation – thrombotic thrombocytopenic purpura (TTP)**

Haematologist

*Re-assessment required after 8 weeks*

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks; and
- 2 Either:
  - 2.1 Patient has thrombotic thrombocytopenic purpura\* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
  - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura\* with neurological or cardiovascular pathology.

Note: Indications marked with \* are unapproved indications.

**Continuation – thrombotic thrombocytopenic purpura (TTP)**

Haematologist

*Re-assessment required after 8 weeks*

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura\*;
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Initiation – pure red cell aplasia (PRCA)**

Haematologist

*Re-assessment required after 6 weeks*

Patient has autoimmune pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with \* are unapproved indications.

**Continuation – pure red cell aplasia (PRCA)**

Haematologist

*Re-assessment required after 6 weeks*

Patient was previously treated with rituximab for pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with \* are unapproved indications.

**Initiation – ANCA associated vasculitis**

*Re-assessment required after 8 weeks*

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*;
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
  - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
  - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
  - 3.3 Cyclophosphamide and methotrexate are contraindicated; or

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

3.4 Patient is a female of child-bearing potential; or

3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with \* are unapproved indications.

## **Continuation – ANCA associated vasculitis**

*Re-assessment required after 8 weeks*

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*;
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

## **Initiation – treatment refractory systemic lupus erythematosus (SLE)**

Rheumatologist or nephrologist

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE\*;
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

## **Continuation – treatment refractory systemic lupus erythematosus (SLE)**

Rheumatologist or nephrologist

All of the following:

- 1 Patient's SLE\* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

## **Initiation – Antibody-mediated organ transplant rejection**

Patient has been diagnosed with antibody-mediated organ transplant rejection\*.

Note: Indications marked with \* are unapproved indications.

## **Initiation – ABO-incompatible organ transplant**

Patient is to undergo an ABO-incompatible solid organ transplant\*.

Note: Indications marked with \* are unapproved indications.

## **Initiation – Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)**

Nephrologist

*Re-assessment required after 8 weeks*

All of the following:

- 1 Patient is a child with SDNS\* or FRNS\*;
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with a \* are unapproved indications.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per

continued...

**Continuation – Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)**

Nephrologist

*Re-assessment required after 8 weeks*

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for > 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with a \* are unapproved indications.

**Initiation – Steroid resistant nephrotic syndrome (SRNS)**

Nephrologist

*Re-assessment required after 8 weeks*

All of the following:

- 1 Patient is a child with SRNS\* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with a \* are unapproved indications.

**Continuation – Steroid resistant nephrotic syndrome (SRNS)**

Nephrologist

*Re-assessment required after 8 weeks*

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with a \* are unapproved indications.

**Initiation – Neuromyelitis Optica Spectrum Disorder (NMOSD)**

*Re-assessment required after 6 months*

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m<sup>2</sup> administered weekly for four weeks; and
- 2 Either:
  - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
  - 2.2 All of the following:
    - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
    - 2.2.2 The patient is receiving treatment with mycophenolate; and
    - 2.2.3 The patients is receiving treatment with corticosteroids.

**Continuation – Neuromyelitis Optica Spectrum Disorder (NMOSD)**

*Re-assessment required after 2 years*

All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 375 mg/m<sup>2</sup> administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

## Initiation – Severe Refractory Myasthenia Gravis

Neurologist

*Re-assessment required after 2 years*

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m<sup>2</sup> of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
  - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
  - 2.2 Both:
    - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
    - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

## Continuation – Severe Refractory Myasthenia Gravis

Neurologist

*Re-assessment required after 2 years*

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m<sup>2</sup> of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
  - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
  - 3.2 Both:
    - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
    - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

## Initiation – Severe antisynthetase syndrome

*Re-assessment required after 12 months*

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
  - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
  - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000 mg infusions of rituximab.

## Continuation – Severe antisynthetase syndrome

*Re-assessment required after 12 months*

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 3 Maximum of two cycles of  $2 \times 1,000$  mg infusions of rituximab given two weeks apart.

#### Initiation – graft versus host disease

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not been effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of  $375 \text{ mg/m}^2$  of body surface area per week for a total of 4 weeks.

#### Initiation – severe chronic inflammatory demyelinating polyneuropathy

Neurologist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used:  $375 \text{ mg/m}^2$  of body surface area per week for a total of four weeks, or  $500 \text{ mg}$  once weekly for four weeks, or two  $1,000 \text{ mg}$  doses given two weeks apart.

#### Continuation – severe chronic inflammatory demyelinating polyneuropathy

Neurologist or medical practitioner on the recommendation of a Neurologist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used:  $375 \text{ mg/m}^2$  of body surface area per week for a total of four weeks, or  $500 \text{ mg}$  once weekly for four weeks, or two  $1,000 \text{ mg}$  doses given two weeks apart.

#### Initiation – anti-NMDA receptor autoimmune encephalitis

Neurologist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used:  $375 \text{ mg/m}^2$  of body surface area per week for a total of four weeks, or  $500 \text{ mg}$  once weekly for four weeks, or two  $1,000 \text{ mg}$  doses given two weeks apart.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

## **Continuation – anti-NMDA receptor autoimmune encephalitis**

Neurologist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m<sup>2</sup> of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

## **Initiation – CD20+ low grade or follicular B-cell NHL**

*Re-assessment required after 9 months*

Either:

- 1 Both:
  - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
  - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

## **Continuation – CD20+ low grade or follicular B-cell NHL**

*Re-assessment required after 24 months*

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m<sup>2</sup> every 8 weeks (maximum of 12 cycles).

## **SECUKINUMAB – Restricted see terms below**

↓ Inj 150 mg per ml, 1 ml prefilled syringe..... 1,599.00      2      Cosentyx

➡ **Restricted (RS1653)**

## **Initiation – severe chronic plaque psoriasis, second-line biologic**

Dermatologist

*Re-assessment required after 4 months*

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialed infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
  - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQI assessment is no more than 1 month old at the time of application.

## **Continuation – severe chronic plaque psoriasis, second-line biologic**

Dermatologist

*Re-assessment required after 6 months*

Both:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1 Either:
  - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
  - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

**Initiation – severe chronic plaque psoriasis, first-line biologic**

Dermatologist

*Re-assessment required after 4 months*

All of the following:

- 1 Either:
  - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

**Continuation – severe chronic plaque psoriasis, first-line biologic**

Dermatologist

*Re-assessment required after 6 months*

Both:

- 1 Either:
  - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
  - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

SILTUXIMAB – **Restricted** see terms [below](#)

↓ Inj 100 mg vial .....	770.57	1	Sylvant
↓ Inj 400 mg vial .....	3,082.33	1	Sylvant

➔ **Restricted (RS1525)**

**Initiation**

Haematologist or rheumatologist

*Re-assessment required after 6 months*

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

## Continuation

Haematologist or rheumatologist

*Re-assessment required after 12 months*

The treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB – **Restricted** see terms [below](#)

⚡ Inj 20 mg per ml, 4 ml vial.....	220.00	1	Actemra
⚡ Inj 20 mg per ml, 10 ml vial.....	550.00	1	Actemra
⚡ Inj 20 mg per ml, 20 ml vial.....	1,100.00	1	Actemra

➡ **Restricted (RS1786)**

## Initiation – cytokine release syndrome

*Therapy limited to 3 doses*

Either:

- 1 All of the following:
  - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
  - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
  - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
  - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
  - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
  - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

## Initiation – previous use

Any relevant practitioner

*Limited to 6 months treatment*

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 rheumatoid arthritis; or
  - 2.2 systemic juvenile idiopathic arthritis; or
  - 2.3 adult-onset Still's disease; or
  - 2.4 polyarticular juvenile idiopathic arthritis; or
  - 2.5 idiopathic multicentric Castleman's disease.

## Initiation – Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)

Rheumatologist or Practitioner on the recommendation of a rheumatologist

*Limited to 6 months treatment*

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and

continued...

	Price	Brand or
(ex man. excl. GST)		Generic
\$	Per	Manufacturer

continued...

3 Either:

3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or

3.2 Both:

3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and

3.2.2 Either:

3.2.2.1 The patient has experienced intolerable side effects from rituximab; or

3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

### Initiation – Rheumatoid Arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

*Re-assessment required after 6 months*

All of the following:

1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and

2 Tocilizumab is to be used as monotherapy; and

3 Either:

3.1 Treatment with methotrexate is contraindicated; or

3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and

4 Either:

4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or

4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and

5 Either:

5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or

5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

6 Either:

6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or

6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

### Initiation – systemic juvenile idiopathic arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

*Re-assessment required after 6 months*

Both:

1 Patient diagnosed with systemic juvenile idiopathic arthritis; and

2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

### Initiation – adult-onset Still's disease

Rheumatologist or Practitioner on the recommendation of a rheumatologist

*Re-assessment required after 6 months*

Either:

1 Both:

1.1 Either:

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
- 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule; and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

## Initiation – polyarticular juvenile idiopathic arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

*Re-assessment required after 4 months*

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
  - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.4 Any of the following:
    - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

## Initiation – idiopathic multicentric Castleman's disease

Haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist

*Re-assessment required after 6 months*

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

## Continuation – Rheumatoid Arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

*Re-assessment required after 6 months*

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

**Continuation – systemic juvenile idiopathic arthritis**

Rheumatologist or Practitioner on the recommendation of a rheumatologist

*Re-assessment required after 6 months*

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

**Continuation – adult-onset Still's disease**

Rheumatologist or Practitioner on the recommendation of a rheumatologist

*Re-assessment required after 6 months*

the patient has a sustained improvement in inflammatory markers and functional status.

**Continuation – polyarticular juvenile idiopathic arthritis**

Rheumatologist or Practitioner on the recommendation of a rheumatologist

*Re-assessment required after 6 months*

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

**Continuation – idiopathic multicentric Castleman's disease**

Haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist

*Re-assessment required after 12 months*

the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

**TRASTUZUMAB – Restricted** see terms [below](#)

↓ Inj 150 mg vial .....	1,350.00	1	Herceptin
↓ Inj 440 mg vial .....	3,875.00	1	Herceptin

→ **Restricted (RS1554)**

**Initiation – Early breast cancer**

*Limited to 12 months treatment*

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
  - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
  - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
  - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
  - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
  - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

**Initiation – metastatic breast cancer (trastuzumab-naïve patients)**

*Limited to 12 months treatment*

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology);

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

and

2 Either:

2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or

2.2 Both:

2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and

2.2.2 The cancer did not progress whilst on lapatinib; and

3 Either:

3.1 Trastuzumab will not be given in combination with pertuzumab; or

3.2 All of the following:

3.2.1 Trastuzumab to be administered in combination with pertuzumab; and

3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and

3.2.3 The patient has good performance status (ECOG grade 0-1); and

4 Trastuzumab not to be given in combination with lapatinib; and

5 Trastuzumab to be discontinued at disease progression.

## Initiation – metastatic breast cancer (patients previously treated with trastuzumab)

Limited to 12 months treatment

All of the following:

1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and

2 Either:

2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or

2.2 Both:

2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and

2.2.2 The cancer did not progress whilst on lapatinib; and

3 Either:

3.1 Trastuzumab will not be given in combination with pertuzumab; or

3.2 All of the following:

3.2.1 Trastuzumab to be administered in combination with pertuzumab; and

3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and

3.2.3 The patient has good performance status (ECOG grade 0-1); and

4 Trastuzumab not to be given in combination with lapatinib; and

5 Trastuzumab to be discontinued at disease progression.

## Continuation – metastatic breast cancer

Re-assessment required after 12 months

All of the following:

1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and

2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and

3 Trastuzumab not to be given in combination with lapatinib; and

4 Trastuzumab to be discontinued at disease progression.

TRASTUZUMAB EMTANSINE – **Restricted** see terms [on the next page](#)

⚡ Inj 100 mg vial .....	2,320.00	1	Kadcyla
⚡ Inj 160 mg vial .....	3,712.00	1	Kadcyla

	Price		Brand or
(ex man. excl. GST)	\$	Per	Generic
			Manufacturer

➔ **Restricted (RS1715)**

**Initiation**

*Re-assessment required after 6 months*

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
  - 3.1 The patient has received prior therapy for metastatic disease\*; or
  - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy\*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:
  - 5.1 Patient does not have symptomatic brain metastases; or
  - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Treatment to be discontinued at disease progression.

**Continuation**

*Re-assessment required after 6 months*

- Both:
- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
  - 2 Treatment to be discontinued at disease progression.

Note: \*Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

**Programmed Cell Death-1 (PD-1) Inhibitors**

**NIVOLUMAB – Restricted** see terms [below](#)

↓ Inj 10 mg per ml, 4 ml vial.....	1,051.98	1	Opdivo
↓ Inj 10 mg per ml, 10 ml vial.....	2,629.96	1	Opdivo

➔ **Restricted (RS1742)**

**Initiation**

Medical oncologist

*Re-assessment required after 4 months*

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
  - 4.1 Patient has not received funded pembrolizumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

**Continuation**

Medical oncologist

*Re-assessment required after 4 months*

Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
    - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
    - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
  - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
  - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
  - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks.

Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB – **Restricted** see terms [below](#)

⚡ Inj 25 mg per ml, 4 ml vial.....4,680.00      1      Keytruda

➡ **Restricted (RS1741)**

## Initiation

Medical oncologist

*Re-assessment required after 4 months*

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
  - 4.1 Patient has not received funded nivolumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

4.2.2 The cancer did not progress while the patient was on nivolumab; and

5 Baseline measurement of overall tumour burden is documented (see Note); and

6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

## Continuation

Medical oncologist

*Re-assessment required after 4 months*

Either:

1 All of the following:

1.1 Any of the following:

1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or

1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or

1.1.3 Patient has stable disease according to RECIST criteria (see Note); and

1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and

1.3 No evidence of progressive disease according to RECIST criteria (see Note); and

1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or

2 All of the following:

2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and

2.2 Patient has signs of disease progression; and

2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks.

Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

## Other Immunosuppressants

ANTITHYMOCYTE GLOBULIN (EQUINE)

Inj 50 mg per ml, 5 ml ampoule .....	2,351.25	5	ATGAM
--------------------------------------	----------	---	-------

ANTITHYMOCYTE GLOBULIN (RABBIT)

Inj 25 mg vial

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>AZATHIOPRINE</b>			
Tab 25 mg – 1% DV Jan-20 to 2022 .....	7.35	60	<b>Azamun</b>
Tab 50 mg – 1% DV Jan-20 to 2022 .....	7.60	100	<b>Azamun</b>
Inj 50 mg vial – 1% DV Nov-19 to 2022 .....	199.00	1	<b>Imuran</b>
<b>BACILLUS CALMETTE-GUERIN (BCG) – Restricted</b> see terms <a href="#">below</a>			
⚡ Inj 2-8 x 10 <sup>8</sup> CFU vial .....	149.37	1	<b>OncoTICE</b>
➡ <b>Restricted (RS1206)</b>			
<b>Initiation</b>			
For use in bladder cancer.			
<b>EVEROLIMUS – Restricted</b> see terms <a href="#">below</a>			
⚡ Tab 5 mg .....	4,555.76	30	<b>Afinitor</b>
⚡ Tab 10 mg .....	6,512.29	30	<b>Afinitor</b>
➡ <b>Restricted (RS1745)</b>			
<b>Initiation</b>			
Neurologist or oncologist			
<i>Re-assessment required after 3 months</i>			
Both:			
1 Patient has tuberous sclerosis; and			
2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.			
<b>Continuation – pandemic circumstances</b>			
<i>Re-assessment required after 6 months</i>			
All of the following:			
1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and			
2 Everolimus to be discontinued at progression of SEGAs; and			
3 The regular renewal requirements cannot be met due to COVID-19 constraints on the health sector.			
Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.			
<b>Continuation</b>			
Neurologist or oncologist			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and			
2 The treatment remains appropriate and the patient is benefiting from treatment; and			
3 Everolimus to be discontinued at progression of SEGAs.			
Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.			
<b>MYCOPHENOLATE MOFETIL</b>			
Tab 500 mg .....	35.90	50	<b>CellCept</b>
Cap 250 mg .....	35.90	100	<b>CellCept</b>
Powder for oral liq 1 g per 5 ml .....	187.25	165 ml	<b>CellCept</b>
Inj 500 mg vial .....	133.33	4	<b>CellCept</b>
<b>PICIBANIL</b>			
Inj 100 mg vial			
<b>SIROLIMUS – Restricted</b> see terms <a href="#">on the next page</a>			
⚡ Tab 1 mg .....	749.99	100	<b>Rapamune</b>
⚡ Tab 2 mg .....	1,499.99	100	<b>Rapamune</b>
⚡ Oral liq 1 mg per ml .....	449.99	60 ml	<b>Rapamune</b>

	Price	Brand or
	(ex man. excl. GST)	Generic
	\$	Per
		Manufacturer

➔ **Restricted** ([RS1208](#))

**Initiation**

For rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR < 30 ml/min; or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- HUS or TTP; or
- Leukoencephalopathy; or
- Significant malignant disease

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## Antiallergy Preparations

### Allergic Emergencies

ICATIBANT – **Restricted** see terms [below](#)

⚡ Inj 10 mg per ml, 3 ml prefilled syringe .....	2,668.00	1	Firazyr
--	----------	---	---------

➡ **Restricted** (RS1501)

#### Initiation

Clinical immunologist or relevant specialist

*Re-assessment required after 12 months*

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

#### Continuation

*Re-assessment required after 12 months*

The treatment remains appropriate and the patient is benefiting from treatment.

### Allergy Desensitisation

BEE VENOM – **Restricted** see terms [below](#)

⚡ Maintenance kit - 6 vials 120 mcg freeze dried venom, with diluent			
⚡ Inj 550 mcg vial with diluent			
⚡ Initiation Kit - 5 vials freeze dried venom with diluent .....	305.00	1	VENOX
⚡ Maintenance Kit - 1 vial freeze dried venom with diluent .....	305.00	1	VENOX

➡ **Restricted** (RS1117)

#### Initiation

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

PAPER WASP VENOM – **Restricted** see terms [below](#)

⚡ Treatment kit - 6 vials 120 mcg freeze dried venom, with diluent			
⚡ Inj 550 mcg vial with diluent			

➡ **Restricted** (RS1118)

#### Initiation

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

YELLOW JACKET WASP VENOM – **Restricted** see terms [below](#)

⚡ Treatment kit - 6 vials 120 mcg freeze dried venom, with diluent			
⚡ Inj 550 mcg vial with diluent			

➡ **Restricted** (RS1119)

#### Initiation

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Allergy Prophylactics</b>			
<b>BUDESONIDE</b>			
Nasal spray 50 mcg per dose – <b>1% DV Oct-20 to 2023</b> .....	2.54	200 dose	<b>SteroClear</b>
Nasal spray 100 mcg per dose – <b>1% DV Oct-20 to 2023</b> .....	2.84	200 dose	<b>SteroClear</b>
<b>FLUTICASONE PROPIONATE</b>			
Nasal spray 50 mcg per dose – <b>1% DV Nov-18 to 2021</b> .....	1.98	120 dose	<b>Flixonase Hayfever &amp; Allergy</b>
<b>IPRATROPIUM BROMIDE</b>			
Aqueous nasal spray 0.03% .....	4.61	15 ml	Univent
<b>SODIUM CROMOGLICATE</b>			
Nasal spray 4%			
<b>Antihistamines</b>			
<b>CETIRIZINE HYDROCHLORIDE</b>			
Tab 10 mg – <b>1% DV Nov-19 to 2022</b> .....	1.12	100	<b>Zista</b>
Oral liq 1 mg per ml .....	2.99	200 ml	Histaclear
<b>CHLORPHENIRAMINE MALEATE</b>			
Oral liq 0.4 mg per ml			
Inj 10 mg per ml, 1 ml ampoule			
<b>CYPROHEPTADINE HYDROCHLORIDE</b>			
Tab 4 mg			
<b>FEXOFENADINE HYDROCHLORIDE</b>			
Tab 60 mg			
Tab 120 mg			
Tab 180 mg			
<b>LORATADINE</b>			
Tab 10 mg – <b>1% DV Feb-20 to 2022</b> .....	1.69	100	<b>Lorafix</b>
Oral liq 1 mg per ml .....	2.95	120 ml	Lorfast
<b>PROMETHAZINE HYDROCHLORIDE</b>			
Tab 10 mg – <b>1% DV Sep-18 to 2021</b> .....	1.68	50	<b>Allersoothe</b>
Tab 25 mg – <b>1% DV Sep-18 to 2021</b> .....	1.89	50	<b>Allersoothe</b>
Oral liq 1 mg per ml – <b>1% DV Sep-18 to 2021</b> .....	2.69	100 ml	<b>Allersoothe</b>
Inj 25 mg per ml, 2 ml ampoule .....	17.87	5	Hospira
<b>Anticholinergic Agents</b>			
<b>IPRATROPIUM BROMIDE</b>			
Aerosol inhaler 20 mcg per dose			
Nebuliser soln 250 mcg per ml, 1 ml ampoule .....	3.35	20	Univent
Nebuliser soln 250 mcg per ml, 2 ml ampoule – <b>1% DV Jan-20 to 2022</b> .....	11.73	20	<b>Univent</b>
<i>(Univent Nebuliser soln 250 mcg per ml, 1 ml ampoule to be delisted 1 January 2021)</i>			
<b>Anticholinergic Agents with Beta-Adrenoceptor Agonists</b>			
<b>SALBUTAMOL WITH IPRATROPIUM BROMIDE</b>			
Aerosol inhaler 100 mcg with ipratropium bromide 20 mcg per dose			
Nebuliser soln 2.5 mg with ipratropium bromide 0.5 mg per 2.5 ml ampoule – <b>1% DV Oct-18 to 2021</b> .....	5.20	20	<b>Duolin</b>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## Long-Acting Muscarinic Agents

### GLYCOPYRRONIUM

Note: inhaled glycopyrronium treatment must not be used if the patient is also receiving treatment with subsidised tiotropium or umeclidinium.

Powder for inhalation 50 mcg per dose ..... 61.00      30 dose      Seebri Breezhaler

### TIOTROPIUM BROMIDE

Note: tiotropium treatment must not be used if the patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.

Soln for inhalation 2.5 mcg per dose ..... 50.37      60 dose      Spiriva Respimat

Powder for inhalation 18 mcg per dose ..... 50.37      30 dose      Spiriva

### UMECLIDINIUM

Note: Umeclidinium must not be used if the patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.

Powder for inhalation 62.5 mcg per dose ..... 61.50      30 dose      Incruse Ellipta

## Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

### ➡ Restricted (RS1518)

#### Initiation

*Re-assessment required after 2 years*

- Both:
- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
  - 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

#### Continuation

*Re-assessment required after 2 years*

- Both:
- 1 Patient is compliant with the medication; and
  - 2 Patient has experienced improved COPD symptom control (prescriber determined).

Note: Combination long acting muscarinic antagonist and long acting beta-2 agonist must not be used if the patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

### GLYCOPYRRONIUM WITH INDACATEROL – Restricted see terms [above](#)

† Powder for Inhalation 50 mcg with indacaterol 110 mcg ..... 81.00      30 dose      Ultibro Breezhaler

### TIOTROPIUM BROMIDE WITH OLODATEROL – Restricted see terms [above](#)

† Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg ..... 81.00      60 dose      Spiolto Respimat

### UMECLIDINIUM WITH VILANTEROL – Restricted see terms [above](#)

† Powder for inhalation 62.5 mcg with vilanterol 25 mcg ..... 77.00      30 dose      Anoro Ellipta

## Antifibrotics

### NINTEDANIB – Restricted see terms [below](#)

‡ Cap 100 mg ..... 2,554.00      60      Ofev

‡ Cap 150 mg ..... 3,870.00      60      Ofev

### ➡ Restricted (RS1756)

#### Initiation – idiopathic pulmonary fibrosis

Respiratory specialist

*Re-assessment required after 12 months*

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with pirfenidone; or
  - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Continuation – idiopathic pulmonary fibrosis

Re-assessment required after 12 months

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE – **Restricted** see terms [below](#)

⬇ Tab 801 mg .....	3,645.00	90	Esbriet
⬇ Cap 267 mg .....	3,645.00	270	Esbriet

➔ **Restricted (RS1757)**

Initiation – idiopathic pulmonary fibrosis

Respiratory specialist

Re-assessment required after 12 months

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Notes); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with nintedanib; or
  - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Continuation – idiopathic pulmonary fibrosis

Re-assessment required after 12 months

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

## RESPIRATORY SYSTEM AND ALLERGIES

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Beta-Adrenoceptor Agonists</b>			
<b>SALBUTAMOL</b>			
Oral liq 400 mcg per ml – <b>1% DV Nov-18 to 2021</b> .....	20.00	150 ml	<b>Ventolin</b>
Inj 500 mcg per ml, 1 ml ampoule			
Inj 1 mg per ml, 5 ml ampoule			
Aerosol inhaler, 100 mcg per dose .....	3.80	200 dose	SalAir
	6.00		Ventolin
Nebuliser soln 1 mg per ml, 2.5 ml ampoule – <b>1% DV Oct-18 to 2021</b> .....	3.93	20	<b>Asthalin</b>
Nebuliser soln 2 mg per ml, 2.5 ml ampoule – <b>1% DV Oct-18 to 2021</b> .....	4.03	20	<b>Asthalin</b>
<b>TERBUTALINE SULPHATE</b>			
Powder for inhalation 250 mcg per dose			
Inj 0.5 mg per ml, 1 ml ampoule			
Powder for inhalation, 200 mcg per dose (equivalent to 250 mcg metered dose), breath activated .....	22.20	120 dose	Bricanyl Turbuhaler
<b>Cough Suppressants</b>			
<b>PHOLCODINE</b>			
Oral liq 1 mg per ml – <b>1% DV Jun-20 to 2022</b> .....	3.09	200 ml	<b>AFT Pholcodine Linctus BP</b>
<b>Decongestants</b>			
<b>OXYMETAZOLINE HYDROCHLORIDE</b>			
Aqueous nasal spray 0.25 mg per ml			
Aqueous nasal spray 0.5 mg per ml			
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b>			
Tab 60 mg			
<b>SODIUM CHLORIDE</b>			
Aqueous nasal spray isotonic			
<b>SODIUM CHLORIDE WITH SODIUM BICARBONATE</b>			
Soln for nasal irrigation			
<b>XYLOMETAZOLINE HYDROCHLORIDE</b>			
Aqueous nasal spray 0.05%			
Aqueous nasal spray 0.1%			
Nasal drops 0.05%			
Nasal drops 0.1%			
<b>Inhaled Corticosteroids</b>			
<b>BECLOMETHASONE DIPROPIONATE</b>			
Aerosol inhaler 50 mcg per dose .....	8.54	200 dose	Beclazone 50
	9.30		Qvar
Aerosol inhaler 100 mcg per dose .....	12.50	200 dose	Beclazone 100
	15.50		Qvar
Aerosol inhaler 250 mcg per dose .....	22.67	200 dose	Beclazone 250

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>BUDESONIDE</b>			
Nebuliser soln 250 mcg per ml, 2 ml ampoule			
Nebuliser soln 500 mcg per ml, 2 ml ampoule			
Powder for inhalation 100 mcg per dose			
Powder for inhalation 200 mcg per dose			
Powder for inhalation 400 mcg per dose			
<b>FLUTICASONE</b>			
Aerosol inhaler 50 mcg per dose – <b>1% DV Sep-20 to 2023</b>	7.19	120 dose	<b>Flixotide</b>
Powder for inhalation 50 mcg per dose	8.67	60 dose	Flixotide Accuhaler
Powder for inhalation 100 mcg per dose	13.87	60 dose	Flixotide Accuhaler
Aerosol inhaler 125 mcg per dose – <b>1% DV Sep-20 to 2023</b>	13.60	120 dose	<b>Flixotide</b>
Aerosol inhaler 250 mcg per dose – <b>1% DV Sep-20 to 2023</b>	24.62	120 dose	<b>Flixotide</b>
Powder for inhalation 250 mcg per dose	24.51	60 dose	Flixotide Accuhaler

### Leukotriene Receptor Antagonists

<b>MONTELUKAST</b>			
Tab 4 mg – <b>1% DV Jan-20 to 2022</b>	4.25	28	<b>Montelukast Mylan</b>
Tab 5 mg – <b>1% DV Jan-20 to 2022</b>	4.25	28	<b>Montelukast Mylan</b>
Tab 10 mg – <b>1% DV Jan-20 to 2022</b>	3.95	28	<b>Montelukast Mylan</b>

### Long-Acting Beta-Adrenoceptor Agonists

<b>EFORMOTEROL FUMARATE</b>			
Powder for inhalation 12 mcg per dose			
<b>EFORMOTEROL FUMARATE DIHYDRATE</b>			
Powder for inhalation 4.5 mcg per dose, breath activated (equivalent to eformoterol fumarate 6 mcg metered dose)			
<b>INDACATEROL</b>			
Powder for inhalation 150 mcg per dose	61.00	30 dose	Onbrez Breezhaler
Powder for inhalation 300 mcg per dose	61.00	30 dose	Onbrez Breezhaler
<b>SALMETEROL</b>			
Aerosol inhaler 25 mcg per dose	9.90	120 dose	Meterol
	25.00		Serevent
Powder for inhalation 50 mcg per dose	25.00	60 dose	Serevent Accuhaler
<i>(Meterol Aerosol inhaler 25 mcg per dose to be delisted 1 January 2021)</i>			

### Inhaled Corticosteroids with Long-Acting Beta-Adrenoceptor Agonists

<b>BUDESONIDE WITH EFORMOTEROL</b>			
Powder for inhalation 100 mcg with eformoterol fumarate 6 mcg			
Powder for inhalation 200 mcg with eformoterol fumarate 6 mcg			
Powder for inhalation 400 mcg with eformoterol fumarate 12 mcg			
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg			
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg			
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide with 6 mcg eformoterol fumarate metered dose)	41.50	120 dose	DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fumarate per dose (equivalent to 400 mcg budesonide with 12 mcg eformoterol fumarate metered dose)	82.50	120 dose	DuoResp Spiromax

## RESPIRATORY SYSTEM AND ALLERGIES

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>FLUTICASONE FUROATE WITH VILANTEROL</b>			
Powder for inhalation 100 mcg with vilanterol 25 mcg .....	44.08	30 dose	Breo Ellipta
<b>FLUTICASONE WITH SALMETEROL</b>			
Aerosol inhaler 50 mcg with salmeterol 25 mcg – 1% DV Sep-20 to 2023.....	25.79	120 dose	<b>Seretide</b>
Powder for inhalation 100 mcg with salmeterol 50 mcg .....	33.74	60 dose	Seretide Accuhaler
Aerosol inhaler 125 mcg with salmeterol 25 mcg – 1% DV Sep-20 to 2023 .....	32.60	120 dose	<b>Seretide</b>
Powder for inhalation 250 mcg with salmeterol 50 mcg .....	44.08	60 dose	Seretide Accuhaler

### Mast Cell Stabilisers

#### NEDOCROMIL

Aerosol inhaler 2 mg per dose

(Any Aerosol inhaler 2 mg per dose to be delisted 1 February 2021)

#### SODIUM CROMOGLICATE

Aerosol inhaler 5 mg per dose

(Any Aerosol inhaler 5 mg per dose to be delisted 1 May 2021)

### Methylxanthines

#### AMINOPHYLLINE

Inj 25 mg per ml, 10 ml ampoule ..... 124.37

5

DBL Aminophylline

#### CAFFEINE CITRATE

Oral liq 20 mg per ml (caffeine 10 mg per ml) – 1% DV Nov-19 to 2022..... 15.10

25 ml

**Biomed**

Inj 20 mg per ml (caffeine 10 mg per ml), 2.5 ml ampoule – 1% DV  
Nov-19 to 2022 .....

63.25

5

**Biomed**

#### THEOPHYLLINE

Tab long-acting 250 mg – 1% DV Jan-20 to 2022..... 23.02

100

**Nuelin-SR**

Oral liq 80 mg per 15 ml – 1% DV Jan-20 to 2022 ..... 16.60

500 ml

**Nuelin**

### Mucolytics and Expectorants

#### DORNASE ALFA – **Restricted** see terms [below](#)

⚡ Nebuliser soln 2.5 mg per 2.5 ml ampoule ..... 250.00

6

Pulmozyme

➔ **Restricted (RS1787)**

#### Initiation – cystic fibrosis

Respiratory physician or paediatrician

Re-assessment required after 12 months

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
  - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
  - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
  - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or
  - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

continued...

	Price		Brand or
(ex man. excl. GST)			Generic
\$	Per		Manufacturer

continued...

## Continuation – cystic fibrosis

Respiratory physician or paediatrician

The treatment remains appropriate and the patient continues to benefit from treatment.

## Initiation – significant mucus production

Limited to 4 weeks treatment

Both:

- 1 Patient is an in-patient; and
- 2 The mucus production cannot be cleared by first line chest techniques.

## Initiation – pleural emphyema

Limited to 3 days treatment

Both:

- 1 Patient is an in-patient; and
- 2 Patient diagnoses with pleural emphyema.

## SODIUM CHLORIDE

Nebuliser soln 7%, 90 ml bottle – **1% DV Nov-19 to 2022** .....24.50      90 ml      **Biomed**

## Pulmonary Surfactants

### BERACTANT

Soln 200 mg per 8 ml vial

### PORACTANT ALFA

Soln 120 mg per 1.5 ml vial .....425.00      1      Curosurf

Soln 240 mg per 3 ml vial .....695.00      1      Curosurf

## Respiratory Stimulants

### DOXAPRAM

Inj 20 mg per ml, 5 ml vial

## Sclerosing Agents

### TALC

Powder

Soln (slurry) 100 mg per ml, 50 ml

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Anti-Infective Preparations</b>			
<b>Antibacterials</b>			
CHLORAMPHENICOL			
Eye oint 1% – 1% DV May-20 to 2022.....	1.55	5 g	Devatis
Ear drops 0.5%			
Eye drops 0.5% – 1% DV Nov-19 to 2022.....	1.54	10 ml	Chlorafast
Eye drops 0.5%, single dose			
CIPROFLOXACIN			
Eye drops 0.3% .....	9.99	5 ml	Ciprofloxacin Teva
FRAMYCETIN SULPHATE			
Ear/eye drops 0.5%			
GENTAMICIN SULPHATE			
Eye drops 0.3% .....	11.40	5 ml	Genoptic
PROPAMIDINE ISETHIONATE			
Eye drops 0.1%			
SODIUM FUSIDATE [FUSIDIC ACID]			
Eye drops 1% .....	5.29	5 g	Fucithalmic
SULPHACETAMIDE SODIUM			
Eye drops 10%			
TOBRAMYCIN			
Eye oint 0.3% .....	10.45	3.5 g	Tobrex
Eye drops 0.3% .....	11.48	5 ml	Tobrex
<b>Antifungals</b>			
NATAMYCIN			
Eye drops 5%			
<b>Antivirals</b>			
ACICLOVIR			
Eye oint 3% .....	14.92	4.5 g	ViruPOS
<b>Combination Preparations</b>			
CIPROFLOXACIN WITH HYDROCORTISONE			
Ear drops ciprofloxacin 0.2% with 1% hydrocortisone.....	16.30	10 ml	Ciproxin HC Otic
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/eye drops 500 mcg with framycetin sulphate 5 mg and gramicidin 50 mcg per ml			
DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE			
Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per g .....	5.39	3.5 g	Maxitrol
Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml .....	4.50	5 ml	Maxitrol
DEXAMETHASONE WITH TOBRAMYCIN			
Eye drops 0.1% with tobramycin 0.3% .....	12.64	5 ml	Tobradex

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FLUMETASONE PIVALATE WITH CLIQUINOL Ear drops 0.02% with cliquinol 1%			
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCIN AND NYSTATIN Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g .....	5.16	7.5 ml	Kenacomb

## Anti-Inflammatory Preparations

### Corticosteroids

DEXAMETHASONE			
Eye oint 0.1% .....	5.86	3.5 g	Maxidex
Eye drops 0.1% .....	4.50	5 ml	Maxidex
↓ Ocular implant 700 mcg.....	1,444.50	1	Ozurdex

#### → Restricted (RS1606)

#### Initiation – Diabetic macular oedema

Ophthalmologist

*Re-assessment required after 12 months*

All of the following:

- 1 Patients have diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 – 6/48 with functional awareness of reduction in vision; and
- 3 Either:
  - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
  - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

#### Continuation – Diabetic macular oedema

Ophthalmologist

*Re-assessment required after 12 months*

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

#### Initiation – Women of child bearing age with diabetic macular oedema

Ophthalmologist

*Re-assessment required after 12 months*

All of the following:

- 1 Patients have diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 – 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

#### Continuation – Women of child bearing age with diabetic macular oedema

Ophthalmologist

*Re-assessment required after 12 months*

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

## SENSORY ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FLUOROMETHOLONE			
Eye drops 0.1% .....	3.09	5 ml	FML
PREDNISOLONE ACETATE			
Eye drops 0.12% .....			
Eye drops 1% .....	7.00	5 ml	Pred Forte
	5.93	10 ml	Prednisolone- AFT
PREDNISOLONE SODIUM PHOSPHATE			
Eye drops 0.5%, single dose (preservative free).....	38.50	20 dose	Minims Prednisolone

### Non-Steroidal Anti-Inflammatory Drugs

DICLOFENAC SODIUM			
Eye drops 0.1% .....	13.80	5 ml	Voltaren Ophtha
KETOROLAC TROMETAMOL			
Eye drops 0.5% .....			

### Decongestants and Antiallergics

#### Antiallergic Preparations

LEVOCABASTINE			
Eye drops 0.05% .....			
LODOXAMIDE			
Eye drops 0.1% .....	8.71	10 ml	Lomide
OLOPATADINE			
Eye drops 0.1% – 1% DV Oct-20 to 2022 .....	2.20	5 ml	Olopatadine Teva
SODIUM CROMOGLICATE			
Eye drops 2% – 1% DV Jan-20 to 2022 .....	1.79	5 ml	Rexacrom

#### Decongestants

NAPHAZOLINE HYDROCHLORIDE			
Eye drops 0.1% .....	4.15	15 ml	Naphcon Forte

### Diagnostic and Surgical Preparations

#### Diagnostic Dyes

FLUORESCEIN SODIUM			
Eye drops 2%, single dose			
Inj 10%, 5 ml vial .....	125.00	12	Fluorescite
Ophthalmic strips 1 mg			
FLUORESCEIN SODIUM WITH LIGNOCAINE HYDROCHLORIDE			
Eye drops 0.25% with lignocaine hydrochloride 4%, single dose			
LISSAMINE GREEN			
Ophthalmic strips 1.5 mg			
ROSE BENGAL SODIUM			
Ophthalmic strips 1%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Irrigation Solutions</b>			
<b>MIXED SALT SOLUTION FOR EYE IRRIGATION</b>			
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 15 ml dropper bottle .....	5.00	15 ml	Balanced Salt Solution
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 250 ml			<i>e.g. Balanced Salt Solution</i>
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 500 ml bottle.....	10.50	500 ml	Balanced Salt Solution
<b>Ocular Anaesthetics</b>			
<b>OXYBUPROCAINE HYDROCHLORIDE</b>			
Eye drops 0.4%, single dose			
<b>PROXYMETACAINE HYDROCHLORIDE</b>			
Eye drops 0.5%			
<b>TETRACAINE [AMETHOCAINE] HYDROCHLORIDE</b>			
Eye drops 0.5%, single dose			
Eye drops 1%, single dose			
<b>Viscoelastic Substances</b>			
<b>HYPROMELLOSE</b>			
Inj 2%, 1 ml syringe			
Inj 2%, 2 ml syringe			
<b>SODIUM HYALURONATE [HYALURONIC ACID]</b>			
Inj 14 mg per ml, 0.85 ml syringe – <b>1% DV Oct-19 to 2022</b> .....	50.00	1	<b>Healon GV</b>
Inj 14 mg per ml, 0.55 ml syringe – <b>1% DV Oct-19 to 2022</b> .....	50.00	1	<b>Healon GV</b>
Inj 23 mg per ml, 0.6 ml syringe – <b>1% DV Oct-19 to 2022</b> .....	60.00	1	<b>Healon 5</b>
Inj 10 mg per ml, 0.85 ml syringe – <b>1% DV Oct-19 to 2022</b> .....	28.50	1	<b>Healon</b>
<b>SODIUM HYALURONATE [HYALURONIC ACID] WITH CHONDROITIN SULPHATE</b>			
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.35 ml syringe and inj 10 mg sodium hyaluronate [hyaluronic acid] per ml, 0.4 ml syringe .....	64.00	1	Duovisc
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.5 ml syringe and inj 10 mg sodium hyaluronate [hyaluronic acid] per ml, 0.55 ml syringe .....	74.00	1	Duovisc
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.75 ml syringe.....	67.00	1	Viscoat
<b>Other</b>			
<b>DISODIUM EDETATE</b>			
Inj 150 mg per ml, 20 ml ampoule			
Inj 150 mg per ml, 20 ml vial			
Inj 150 mg per ml, 100 ml vial			

## SENSORY ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

RIBOFLAVIN 5-PHOSPHATE  
Soln trans epithelial riboflavin  
Inj 0.1%  
Inj 0.1% plus 20% dextran T500

### Glaucoma Preparations

#### Beta Blockers

BETAXOLOL			
Eye drops 0.25% .....	11.80	5 ml	Betoptic S
Eye drops 0.5% .....	7.50	5 ml	Betoptic
TIMOLOL			
Eye drops 0.25% – 1% DV Dec-20 to 2023 .....	1.81	5 ml	Arrow-Timolol
Eye drops 0.5% – 1% DV Dec-20 to 2023 .....	2.04	5 ml	Arrow-Timolol
Eye drops 0.5%, gel forming .....	3.78	2.5 ml	Timoptol XE

#### Carbonic Anhydrase Inhibitors

ACETAZOLAMIDE			
Tab 250 mg .....	17.03	100	Diamox
Inj 500 mg			
BRINZOLAMIDE			
Eye drops 1%			
DORZOLAMIDE			
Eye drops 2%			
DORZOLAMIDE WITH TIMOLOL			
Eye drops 2% with timolol 0.5% – 1% DV Jan-19 to 2021 .....	2.87	5 ml	Dortimopt

#### Miotics

ACETYLCHOLINE CHLORIDE			
Inj 20 mg vial with diluent			
CARBACHOL			
Inj 150 mcg vial			
PILOCARPINE HYDROCHLORIDE			
Eye drops 1% .....	4.26	15 ml	Isopto Carpine
Eye drops 2% .....	5.35	15 ml	Isopto Carpine
Eye drops 2%, single dose			
Eye drops 4% .....	7.99	15 ml	Isopto Carpine

#### Prostaglandin Analogues

BIMATOPROST			
Eye drops 0.03% – 1% DV Feb-19 to 2021 .....	3.30	3 ml	Bimatoprost Multichem
LATANOPROST			
Eye drops 0.005% – 1% DV Apr-19 to 2021 .....	1.57	2.5 ml	Teva
TRAVOPROST			
Eye drops 0.004% .....	7.30	5 ml	Travopt

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Sympathomimetics</b>			
APRACLOPIDINE			
Eye drops 0.5% .....	19.77	5 ml	Iopidine
BRIMONIDINE TARTRATE			
Eye drops 0.2% .....	4.29	5 ml	Arrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL			
Eye drops 0.2% with timolol 0.5%			
<b>Mydriatics and Cycloplegics</b>			
<b>Anticholinergic Agents</b>			
ATROPINE SULPHATE			
Eye drops 0.5%			
Eye drops 1%, single dose			
Eye drops 1% – <b>1% DV Oct-20 to 2023</b> .....	17.36	15 ml	<b>Atropit</b>
CYCLOPENTOLATE HYDROCHLORIDE			
Eye drops 0.5%, single dose			
Eye drops 1% .....	8.76	15 ml	Cyclogyl
Eye drops 1%, single dose			
TROPICAMIDE			
Eye drops 0.5% .....	7.15	15 ml	Mydriacyl
Eye drops 0.5%, single dose			
Eye drops 1% .....	8.66	15 ml	Mydriacyl
Eye drops 1%, single dose			
<b>Sympathomimetics</b>			
PHENYLEPHRINE HYDROCHLORIDE			
Eye drops 2.5%, single dose			
Eye drops 10%, single dose			
<b>Ocular Lubricants</b>			
CARBOMER			
Ophthalmic gel 0.3%, single dose .....	8.25	30	Poly Gel
Ophthalmic gel 0.2%			
CARMELOSE SODIUM WITH PECTIN AND GELATINE			
Eye drops 0.5%			
Eye drops 0.5%, single dose			
Eye drops 1%			
Eye drops 1%, single dose			
HYPROMELLOSE			
Eye drops 0.5% .....	3.92	15 ml	Methopt
HYPROMELLOSE WITH DEXTRAN			
Eye drops 0.3% with dextran 0.1%.....	2.30	15 ml	Poly-Tears
Eye drops 0.3% with dextran 0.1%, single dose			
MACROGOL 400 AND PROPYLENE GLYCOL			
Eye drops 0.4% with propylene glycol 0.3% preservative free, single dose.....	4.30	24	Systane Unit Dose

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

PARAFFIN LIQUID WITH SOFT WHITE PARAFFIN			
Eye oint 42.5% with soft white paraffin 57.3%			
PARAFFIN LIQUID WITH WOOL FAT			
Eye oint 3% with wool fat 3% .....	3.63	3.5 g	Poly-Visc
POLYVINYL ALCOHOL WITH POVIDONE			
Eye drops 1.4% with povidone 0.6%, single dose			
RETINOL PALMITATE			
Oint 138 mcg per g .....	3.80	5 g	VitA-POS
SODIUM HYALURONATE [HYALURONIC ACID]			
Eye drops 1 mg per ml .....	22.00	10 ml	Hylo-Fresh

Other Otological Preparations

ACETIC ACID WITH PROPYLENE GLYCOL
Ear drops 2.3% with propylene glycol 2.8%
DOCUSATE SODIUM
Ear drops 0.5%

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Agents Used in the Treatment of Poisonings</b>			
<b>Antidotes</b>			
<b>ACETYL CYSTEINE</b>			
Tab eff 200 mg			
Inj 200 mg per ml, 10 ml ampoule – <b>1% DV Sep-18 to 2021</b> .....	58.76	10	<b>DBL Acetylcysteine</b>
<b>AMYL NITRITE</b>			
Liq 98% in 3 ml capsule			
<b>DIGOXIN IMMUNE FAB</b>			
Inj 38 mg vial			
Inj 40 mg vial			
<b>ETHANOL</b>			
Liq 96%			
<b>ETHANOL WITH GLUCOSE</b>			
Inj 10% with glucose 5%, 500 ml bottle			
<b>ETHANOL, DEHYDRATED</b>			
Inj 100%, 5 ml ampoule			
Inj 96%			
<b>FLUMAZENIL</b>			
Inj 0.1 mg per ml, 5 ml ampoule – <b>1% DV Dec-18 to 2021</b> .....	132.68	10	<b>Hameln</b>
<b>HYDROXOCOBALAMIN</b>			
Inj 5 g vial			
Inj 2.5 g vial			
<b>NALOXONE HYDROCHLORIDE</b>			
Inj 400 mcg per ml, 1 ml ampoule – <b>1% DV Aug-18 to 2021</b> .....	22.60	5	<b>DBL Naloxone Hydrochloride</b>
<b>PRALIDOXIME IODIDE</b>			
Inj 25 mg per ml, 20 ml ampoule			
<b>SODIUM NITRITE</b>			
Inj 30 mg per ml, 10 ml ampoule			
<b>SODIUM THIOSULFATE</b>			
Inj 250 mg per ml, 10 ml vial			
Inj 250 mg per ml. 50 ml vial			
Inj 500 mg per ml, 10 ml vial			
Inj 500 mg per ml, 20 ml ampoule			
<b>SOYA OIL</b>			
Inj 20%, 500 ml bag			
Inj 20%, 500 ml bottle			

**Antitoxins**

<b>BOTULISM ANTITOXIN</b>			
Inj 250 ml vial			
<b>DIPHTHERIA ANTITOXIN</b>			
Inj 10,000 iu vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## Antivenoms

RED BACK SPIDER ANTIVENOM

Inj 500 u vial

SNAKE ANTIVENOM

Inj 50 ml vial

## Removal and Elimination

CHARCOAL

Oral liq 200 mg per ml ..... 43.50      250 ml      Carbasorb-X

DEFERASIROX – **Restricted** see terms [below](#)

↓ Tab 125 mg dispersible .....	276.00	28	Exjade
↓ Tab 250 mg dispersible .....	552.00	28	Exjade
↓ Tab 500 mg dispersible .....	1,105.00	28	Exjade

➔ **Restricted (RS1444)**

### Initiation

Haematologist

*Re-assessment required after 2 years*

All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
  - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2\*; or
  - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
  - 3.3 Treatment with deferiprone has resulted in arthritis; or
  - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per µL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 - 1.0 cells per µL).

### Continuation

Haematologist

*Re-assessment required after 2 years*

Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels. .

DEFERIPRONE – **Restricted** see terms [below](#)

↓ Tab 500 mg .....	533.17	100	Ferriprox
↓ Oral liq 100 mg per ml .....	266.59	250 ml	Ferriprox

➔ **Restricted (RS1445)**

### Initiation

Patient has been diagnosed with chronic iron overload due to congenital inherited anaemia or acquired red cell aplasia.

DESFERIOXAMINE MESILATE

Inj 500 mg vial – 1% DV Mar-19 to 2021 .....	84.53	10	<b>DBL Desferrioxamine Mesylate for Inj BP</b>
--	-------	----	--

DICOBALT EDETATE

Inj 15 mg per ml, 20 ml ampoule

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DIMERCAPROL			
Inj 50 mg per ml, 2 ml ampoule			
DIMERCAPTOSUCCINIC ACID			
Cap 100 mg			e.g. PCNZ, Optimus Healthcare, Chemet
Cap 200 mg			e.g. PCNZ, Optimus Healthcare, Chemet
SODIUM CALCIUM EDETATE			
Inj 200 mg per ml, 2.5 ml ampoule			
Inj 200 mg per ml, 5 ml ampoule			

## Antiseptics and Disinfectants

CHLORHEXIDINE			
Soln 4%			
Soln 5% .....	15.50	500 ml	healthE
CHLORHEXIDINE WITH CETRIMIDE			
Crm 0.1% with cetrimide 0.5%			
Foaming soln 0.5% with cetrimide 0.5%			
CHLORHEXIDINE WITH ETHANOL			
Soln 0.5% with ethanol 70%			
Soln 2% with ethanol 70%			
Soln 0.5% with ethanol 70%, non-staining (pink) 25 ml .....	1.55	1	healthE
IODINE WITH ETHANOL			
Soln 1% with ethanol 70%			
ISOPROPYL ALCOHOL			
Soln 70%, 500 ml .....	5.65	1	healthE
POVIDONE-IODINE			
↓ Vaginal tab 200 mg			
➔ <b>Restricted (RS1354)</b>			
<b>Initiation</b>			
Rectal administration pre-prostate biopsy.			
Oint 10% – <b>1% DV Oct-20 to 2023</b> .....	7.40	65 g	<b>Betadine</b>
Soln 10% – <b>1% DV Nov-19 to 2021</b> .....	2.55	100 ml	<b>Riodine</b>
Soln 5%			
Soln 7.5%			
Soln 10%, – <b>1% DV Dec-19 to 2022</b> .....	3.83	15 ml	<b>Riodine</b>
	5.40	500 ml	<b>Riodine</b>
Pad 10%			
Swab set 10%			
POVIDONE-IODINE WITH ETHANOL			
Soln 10% with ethanol 30%			
Soln 10% with ethanol 70%			
SODIUM HYPOCHLORITE			
Soln			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Contrast Media</b>			
<b>Iodinated X-ray Contrast Media</b>			
<b>DIATRIZOATE MEGLUMINE WITH SODIUM AMIDOTRIZOATE</b>			
Oral liq 660 mg per ml with sodium amidotrizoate 100 mg per ml, 100 ml bottle.....	22.50	100 ml	Gastrografin
Inj 260 mg with sodium amidotrizoate 40 mg per ml, 250 ml bottle.....	80.00	1	Urografin
<b>DIATRIZOATE SODIUM</b>			
Oral liq 370 mg per ml, 10 ml sachet.....	156.12	50	Ioscan
<b>IODISED OIL</b>			
Inj 38% w/w (480 mg per ml), 10 ml ampoule .....	410.00	1	Lipiodol Ultra Fluid
<b>IODIXANOL</b>			
Inj 270 mg per ml (iodine equivalent), 50 ml bottle.....	220.00	10	Visipaque
Inj 270 mg per ml (iodine equivalent), 100 ml bottle.....	430.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 50 ml bottle.....	220.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 100 ml bottle.....	430.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 200 ml bottle.....	850.00	10	Visipaque
<b>IOHEXOL</b>			
Inj 240 mg per ml (iodine equivalent), 50 ml bottle.....	77.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 20 ml bottle.....	59.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 50 ml bottle.....	77.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 100 ml bottle.....	154.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 20 ml bottle.....	61.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 50 ml bottle.....	77.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 75 ml bottle.....	117.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 100 ml bottle.....	154.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 200 ml bottle.....	298.00	10	Omnipaque
<b>Non-iodinated X-ray Contrast Media</b>			
<b>BARIUM SULPHATE</b>			
Powder for oral liq 20 mg per g (2% w/w), 22.1 g sachet.....	507.50	50	E-Z-Cat Dry
Oral liq 400 mg per ml (40% w/v, 30% w/w), bottle .....	17.39	148 g	Varibar - Thin Liquid
Oral liq 600 mg per g (60% w/v, 30% w/w), tube .....	36.51	454 g	E-Z-Paste
Oral liq 400 mg per ml (40% w/v), bottle .....	155.35	250 ml	Varibar - Honey
	38.40	240 ml	Varibar - Nectar
	145.04	230 ml	Varibar - Pudding
Enema 1,250 mg per ml (125% w/v), 500 ml bag .....	282.30	12	Liquibar
Oral liq 22 mg per g (2.2% w/w), 250 ml bottle.....	175.00	24	CT Plus+
Oral liq 22 mg per g (2.2% w/w), 450 ml bottle.....	220.00	24	CT Plus+
Oral liq 1 mg per ml (0.1% w/v, 0.1% w/w), 450 ml bottle .....	441.12	24	Volumen
Oral liq 20.9 mg per ml (2.1% w/v, 2% w/w), 250 ml bottle .....	140.94	24	Readi-CAT 2
Powder for oral soln 97.65% w/w, 300 g bottle .....	237.76	24	X-Opaque-HD
Oral liq 400 mg per ml (40% w/v, 30% w/w), 20 ml bottle .....	52.35	3	Tagitol V
Oral liq 1,250 mg per ml (125% w/v), 2,000 ml bottle.....	91.77	1	Liquibar
<b>BARIUM SULPHATE WITH SODIUM BICARBONATE</b>			
Grans eff 382.2 mg per g with sodium bicarbonate 551.3 mg per g, 4 g sachet.....	102.93	50	E-Z-Gas II

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>CITRIC ACID WITH SODIUM BICARBONATE</b>			
Powder 382.2 mg per g with sodium bicarbonate 551.3 mg per g, 4 g sachet			<i>e.g. E-Z-GAS II</i>

### Paramagnetic Contrast Media

<b>GADOBENIC ACID</b>			
Inj 334 mg per ml, 10 ml vial.....	324.74	10	Multihance
Inj 334 mg per ml, 20 ml vial.....	636.28	10	Multihance
<b>GADOBUTROL</b>			
Inj 1 mmol per ml, 15 ml vial			
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 5 ml prefilled syringe.....	120.00	5	Gadovist 1.0
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 7.5 ml prefilled syringe.....	180.00	5	Gadovist 1.0
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 15 ml prefilled syringe.....	700.00	10	Gadovist 1.0
<b>GADODIAMIDE</b>			
Inj 287 mg per ml, 10 ml prefilled syringe.....	200.00	10	Omniscan
Inj 287 mg per ml, 10 ml vial.....	170.00	10	Omniscan
Inj 287 mg per ml, 5 ml vial.....	120.00	10	Omniscan
Inj 287 mg per ml, 15 ml prefilled syringe.....	320.00	10	Omniscan
<b>GADOTERIC ACID</b>			
Inj 279.32 mg per ml (0.5 mmol per ml), 10 ml prefilled syringe.....	24.50	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 15 ml bottle .....	34.50	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 15 ml prefilled syringe.....	41.00	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 20 ml prefilled syringe.....	55.00	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 10 ml bottle .....	23.20	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 20 ml bottle .....	46.30	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 5 ml bottle .....	12.30	1	Dotarem
<b>GADOXETATE DISODIUM</b>			
Inj 181.43 mg per ml (equivalent to 0.25 mmol per ml), 10 ml prefilled syringe.....	300.00	1	Primovist
<b>MEGLUMINE GADOPENTETATE</b>			
Inj 469 mg per ml, 10 ml prefilled syringe.....	95.00	5	Magnevist
Inj 469 mg per ml, 10 ml vial.....	185.00	10	Magnevist
<b>MEGLUMINE IOTROXATE</b>			
Inj 105 mg per ml, 100 ml bottle .....	150.00	100 ml	Biliscopin

### Ultrasound Contrast Media

<b>PERFLUTREN</b>			
Inj 1.1 mg per ml, 1.5 ml vial.....	180.00	1	Definity
	720.00	4	Definity

### Diagnostic Agents

<b>ARGININE</b>			
Inj 50 mg per ml, 500 ml bottle			
Inj 100 mg per ml, 300 ml bottle			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>HISTAMINE ACID PHOSPHATE</b>			
Nebuliser soln 0.6%, 10 ml vial			
Nebuliser soln 2.5%, 10 ml vial			
Nebuliser soln 5%, 10 ml vial			
<b>MANNITOL</b>			
Powder for inhalation			<i>e.g. Aridol</i>
<b>METHACHOLINE CHLORIDE</b>			
Powder 100 mg			
<b>SECRETIN PENTAHYDROCHLORIDE</b>			
Inj 100 u ampoule			
<b>SINCALIDE</b>			
Inj 5 mcg per vial			

## Diagnostic Dyes

<b>BONNEY'S BLUE DYE</b>			
Soln			
<b>INDIGO CARMINE</b>			
Inj 4 mg per ml, 5 ml ampoule			
Inj 8 mg per ml, 5 ml ampoule			
<b>INDOCYANINE GREEN</b>			
Inj 25 mg vial			
<b>METHYLTHIONINIUM CHLORIDE [METHYLENE BLUE]</b>			
Inj 5 mg per ml, 10 ml ampoule .....	240.35	5	Proveblue
<b>PATENT BLUE V</b>			
Inj 2.5%, 2 ml ampoule .....	440.00	5	Obex Medical
Inj 2.5%, 5 ml prefilled syringe .....	420.00	5	InterPharma

## Irrigation Solutions

**CHLORHEXIDINE WITH CETRIMIDE**  
 ‡ Irrigation soln 0.015% with cetrimide 0.15%, 500 ml bottle

➔ **Restricted (RS1683)**

### Initiation

*Re-assessment required after 3 months*

All of the following:

- 1 Patient has burns that are greater than 30% of total body surface area (BSA); and
- 2 For use in the perioperative preparation and cleansing of large burn areas requiring debridement/skin grafting; and
- 3 The use of 30 ml ampoules is impractical due to the size of the area to be covered.

### Continuation

*Re-assessment required after 3 months*

The treatment remains appropriate for the patient and the patient is benefiting from the treatment.

Irrigation soln 0.015% with cetrimide 0.15%, 30 ml ampoule – **1% DV**

**Aug-18 to 2021** ..... 29.76 30 **Pfizer**

### GLYCINE

Irrigation soln 1.5%, 3,000 ml bag – **1% DV Sep-18 to 2021** ..... 31.20 4 **B Braun**

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>SODIUM CHLORIDE</b>			
Irrigation soln 0.9%, 3,000 ml bag – <b>1% DV Sep-18 to 2021</b> .....	26.80	4	<b>B Braun</b>
Irrigation soln 0.9%, 30 ml ampoule – <b>1% DV Sep-18 to 2021</b> .....	7.00	20	<b>Interpharma</b>
Irrigation soln 0.9%, 1,000 ml bottle – <b>1% DV Jun-18 to 2021</b> .....	14.90	10	<b>Baxter Sodium</b>
			<b>Chloride 0.9%</b>
Irrigation soln 0.9%, 250 ml bottle – <b>1% DV Aug-18 to 2021</b> .....	17.64	12	<b>Fresenius Kabi</b>
<b>WATER</b>			
Irrigation soln, 3,000 ml bag – <b>1% DV Sep-18 to 2021</b> .....	28.80	4	<b>B Braun</b>
Irrigation soln, 1,000 ml bottle – <b>1% DV Jun-18 to 2021</b> .....	17.30	10	<b>Baxter Water for</b>
			<b>Irrigation</b>
Irrigation soln, 250 ml bottle – <b>1% DV Aug-18 to 2021</b> .....	17.64	12	<b>Fresenius Kabi</b>

## Surgical Preparations

### BISMUTH SUBNITRATE AND IODOFORM PARAFFIN

Paste

### DIMETHYL SULFOXIDE

Soln 50%

Soln 99%

### PHENOL

Inj 6%, 10 ml ampoule

### PHENOL WITH IOXAGLIC ACID

Inj 12%, 10 ml ampoule

### TROMETAMOL

Inj 36 mg per ml, 500 ml bottle

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Cardioplegia Solutions</b>			
<b>ELECTROLYTES</b>			
Inj 15 mmol/l sodium chloride, 9 mmol/l potassium chloride, 1 mmol/l potassium hydrogen 2-ketoglutarate, 4 mmol/l magnesium chloride, 18 mmol/l histidine hydrochloride, 180 mmol/l histidine, 2 mmol/l tryptophan, 30 mmol/l mannitol, 0.015 mmol/l calcium chloride, 1,000 ml bag			<i>e.g. Custodiol-HTK</i>
Inj aspartic acid 10.43 mg per ml, citric acid 0.22476 mg per ml, glutamic acid 11.53 mg per ml, sodium phosphate 0.1725 mg per ml, potassium chloride 2.15211 mg per ml, sodium citrate 1.80768 mg per ml, sodium hydroxide 6.31 mg per ml and trometamol 11.2369 mg per ml, 364 ml bag			<i>e.g. Cardioplegia Enriched Paed. Soln.</i>
Inj aspartic acid 8.481 mg per ml, citric acid 0.8188 mg per ml, glutamic acid 9.375 mg per ml, sodium phosphate 0.6285 mg per ml, potassium chloride 2.5 mg per ml, sodium citrate 6.585 mg per ml, sodium hydroxide 5.133 mg per ml and trometamol 9.097 mg per ml, 527 ml bag			<i>e.g. Cardioplegia Enriched Solution</i>
Inj citric acid 0.07973 mg per ml, sodium phosphate 0.06119 mg per ml, potassium chloride 2.181 mg per ml, sodium chloride 1.788 mg ml, sodium citrate 0.6412 mg per ml and trometamol 5.9 mg per ml, 523 ml bag			<i>e.g. Cardioplegia Base Solution</i>
Inj 110 mmol/l sodium, 16 mmol/l potassium, 1.2 mmol/l calcium, 16 mmol/l magnesium and 160 mmol/l chloride, 1,000 ml bag			<i>e.g. Cardioplegia Solution AHB7832</i>
Inj 143 mmol/l sodium, 16 mmol/l potassium, 16 mmol/l magnesium and 1.2 mmol/l calcium, 1,000 ml bag			<i>e.g. Cardioplegia Electrolyte Solution</i>
<b>MONOSODIUM GLUTAMATE WITH SODIUM ASPARTATE</b>			
Inj 42.68 mg with sodium aspartate 39.48 mg per ml, 250 ml bottle			
<b>MONOSODIUM L-ASPARTATE</b>			
Inj 14 mmol per 10 ml, 10 ml			

## Cold Storage Solutions

### SODIUM WITH POTASSIUM

Inj 29 mmol/l with potassium 125 mmol/l, 1,000 ml bag

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Extemporaneously Compounded Preparations</b>			
ACETIC ACID Liq			
ALUM Powder BP			
ARACHIS OIL [PEANUT OIL] Liq			
ASCORBIC ACID Powder			
BENZOIN Tincture compound BP			
BISMUTH SUBGALLATE Powder			
BORIC ACID Powder			
CARBOXYMETHYLCELLULOSE Soln 1.5%			
CETRIMIDE Soln 40%			
CHLORHEXIDINE GLUCONATE Soln 20 %			
CHLOROFORM Liq BP			
CITRIC ACID Powder BP			
CLOVE OIL Liq			
COAL TAR Soln BP – <b>1% DV Nov-19 to 2022</b> .....	36.25	200 ml	<b>Midwest</b>
CODEINE PHOSPHATE Powder			
COLLODION FLEXIBLE Liq			
COMPOUND HYDROXYBENZOATE Soln – <b>1% DV Aug-19 to 2022</b> .....	30.00	100 ml	<b>Midwest</b>
CYSTEAMINE HYDROCHLORIDE Powder			
DISODIUM HYDROGEN PHOSPHATE WITH SODIUM DIHYDROGEN PHOSPHATE Inj 37.46 mg with sodium dihydrogen phosphate 47.7 mg in 1.5 ml ampoule			
DITHRANOL Powder			
GLUCOSE [DEXTROSE] Powder			

## EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GLYCERIN WITH SODIUM SACCHARIN			
Suspension – 1% DV Jul-19 to 2022.....	30.95	473 ml	<b>Ora-Sweet SF</b>
GLYCERIN WITH SUCROSE			
Suspension – 1% DV Jul-19 to 2022.....	30.95	473 ml	<b>Ora-Sweet</b>
GLYCEROL			
Liq – 1% DV Oct-20 to 2023.....	3.23	500 ml	<b>healthE Glycerol BP Liquid</b>
HYDROCORTISONE			
Powder .....	49.95	25 g	ABM
LACTOSE			
Powder			
MAGNESIUM HYDROXIDE			
Paste			
Suspension			
MENTHOL			
Crystals			
METHADONE HYDROCHLORIDE			
Powder			
METHYL HYDROXYBENZOATE			
Powder – 1% DV Jul-19 to 2022.....	8.98	25 g	<b>Midwest</b>
METHYLCELLULOSE			
Powder – 1% DV Jul-19 to 2022.....	36.95	100 g	<b>Midwest</b>
Suspension – 1% DV Jul-19 to 2022.....	30.95	473 ml	<b>Ora-Plus</b>
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN			
Suspension – 1% DV Jul-19 to 2022.....	30.95	473 ml	<b>Ora-Blend SF</b>
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE			
Suspension – 1% DV Jul-19 to 2022.....	30.95	473 ml	<b>Ora-Blend</b>
OLIVE OIL			
Liq			
PARAFFIN			
Liq			
PHENOBARBITONE SODIUM			
Powder			
PHENOL			
Liq			
PILOCARPINE NITRATE			
Powder			
POLYHEXAMETHYLENE BIGUANIDE			
Liq			
POVIDONE K30			
Powder			
SALICYLIC ACID			
Powder			
SILVER NITRATE			
Crystals			
SODIUM BICARBONATE			
Powder BP – 1% DV Jan-20 to 2022.....	10.05	500 g	<b>Midwest</b>

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

# EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SODIUM CITRATE			
Powder			
SODIUM METABISULFITE			
Powder			
STARCH			
Powder			
SULPHUR			
Precipitated			
Sublimed			
SYRUP			
Liq (pharmaceutical grade) – 1% DV Jan-20 to 2022.....	14.95	500 ml	<b>Midwest</b>
THEOBROMA OIL			
Oint			
TRI-SODIUM CITRATE			
Crystals			
TRICHLORACETIC ACID			
Grans			
UREA			
Powder BP			
WOOL FAT			
Oint, anhydrous			
XANTHAN			
Gum 1%			
ZINC OXIDE			
Powder			

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

Food Modules

Carbohydrate

➡ **Restricted (RS1467)**

**Initiation – Use as an additive**

Any of the following:

- 1 Cystic fibrosis; or
- 2 Chronic kidney disease; or
- 3 Cancer in children; or
- 4 Cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 5 Faltering growth in an infant/child; or
- 6 Bronchopulmonary dysplasia; or
- 7 Premature and post premature infant; or
- 8 Inborn errors of metabolism.

**Initiation – Use as a module**

For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

**CARBOHYDRATE SUPPLEMENT – Restricted** see terms [above](#)

† Powder 95 g carbohydrate per 100 g, 368 g can

† Powder 96 g carbohydrate per 100 g, 400 g can

*e.g. Polycal*

Fat

➡ **Restricted (RS1468)**

**Initiation – Use as an additive**

Any of the following:

- 1 Patient has inborn errors of metabolism; or
- 2 Faltering growth in an infant/child; or
- 3 Bronchopulmonary dysplasia; or
- 4 Fat malabsorption; or
- 5 Lymphangiectasia; or
- 6 Short bowel syndrome; or
- 7 Infants with necrotising enterocolitis; or
- 8 Biliary atresia; or
- 9 For use in a ketogenic diet; or
- 10 Chyle leak; or
- 11 Ascites; or
- 12 Patient has increased energy requirements, and for whom dietary measures have not been successful.

**Initiation – Use as a module**

For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

**LONG-CHAIN TRIGLYCERIDE SUPPLEMENT – Restricted** see terms [above](#)

† Liquid 50 g fat per 100 ml, 200 ml bottle

*e.g. Calogen*

† Liquid 50 g fat per 100 ml, 500 ml bottle

*e.g. Calogen*

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>MEDIUM-CHAIN TRIGLYCERIDE SUPPLEMENT – Restricted</b> see terms <a href="#">on the previous page</a>			
† Liquid 50 g fat per 100 ml, 250 ml bottle			e.g. <i>Liquigen</i>
† Liquid 95 g fat per 100 ml, 500 ml bottle			e.g. <i>MCT Oil</i>

**WALNUT OIL – Restricted** see terms [on the previous page](#)

† Liq

## Protein

➔ **Restricted (RS1469)**

**Initiation – Use as an additive**

Either:

- 1 Protein losing enteropathy; or
- 2 High protein needs.

**Initiation – Use as a module**

For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

**PROTEIN SUPPLEMENT – Restricted** see terms [above](#)

† Powder 5 g protein, 0.67 g carbohydrate and 0.6 g fat per 6.6 g, 275 g can			
† Powder 6 g protein per 7 g, can .....	8.95	227 g	Resource Beneprotein
† Powder 89 g protein, < 1.5 g carbohydrate and 2 g fat per 100 g, 225 g can			e.g. <i>Protifar</i>

## Other Supplements

**BREAST MILK FORTIFIER**

Powder 0.2 g protein, 0.7 g carbohydrate and 0.02 g fat per 1 g sachet

Powder 0.5 g protein, 1.2 g carbohydrate and 0.08 g fat per 2 g sachet

Powder 0.6 g protein and 1.4 g carbohydrate per 2.2 g sachet

e.g. *FM 85*

e.g. *S26 Human Milk Fortifier*

e.g. *Nutricia Breast Milk Fortifier*

**CARBOHYDRATE AND FAT SUPPLEMENT – Restricted** see terms [below](#)

† Powder 72.7 g carbohydrate and 22.3 g fat per 100 g, 400 g can

e.g. *Super Soluble Duocal*

➔ **Restricted (RS1212)**

**Initiation**

Both:

- 1 Infant or child aged four years or under; and
- 2 Any of the following:
  - 2.1 Cystic fibrosis; or
  - 2.2 Cancer in children; or
  - 2.3 Faltering growth; or
  - 2.4 Bronchopulmonary dysplasia; or
  - 2.5 Premature and post premature infants.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## Food/Fluid Thickeners

### NOTE:

While pre-thickened drinks and supplements have not been included in Section H, DHB hospitals may continue to use such products for patients with dysphagia, provided that:

- use was established prior to 1 July 2013; and
- the product has not been specifically considered and excluded by PHARMAC; and
- use of the product conforms to any applicable indication restrictions for similar products that are listed in Section H (for example, use of thickened high protein products should be in line with the restriction for high protein oral feed in Section H).

PHARMAC intends to make a further decision in relation to pre-thickened drinks and supplements in the future, and will notify of any change to this situation.

#### CAROB BEAN GUM WITH MAIZE STARCH AND MALTODEXTRIN

Powder

*e.g. Feed Thickener  
Karicare Aptamil*

#### GUAR GUM

Powder

*e.g. Guarcol*

#### MAIZE STARCH

Powder

*e.g. Resource Thicken  
Up; Nuttilis*

#### MALTODEXTRIN WITH XANTHAN GUM

Powder

*e.g. Instant Thick*

#### MALTODEXTRIN WITH XANTHAN GUM AND ASCORBIC ACID

Powder

*e.g. Easy Thick*

## Metabolic Products

### ➡ Restricted (RS1232)

#### Initiation

Any of the following:

- 1 For the dietary management of homocystinuria, maple syrup urine disease, phenylketonuria (PKU), glutaric aciduria, isovaleric acidemia, propionic acidemia, methylmalonic acidemia, tyrosinaemia or urea cycle disorders; or
- 2 Patient has adrenoleukodystrophy; or
- 3 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

## Glutaric Aciduria Type 1 Products

AMINO ACID FORMULA (WITHOUT LYSINE AND LOW TRYPTOPHAN) – **Restricted** see terms [above](#)

⚡ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can

*e.g. GA1 Anamix Infant*

⚡ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can

*e.g. XLYS Low TRY  
Maxamaid*

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Homocystinuria Products

AMINO ACID FORMULA (WITHOUT METHIONINE) – **Restricted** see terms [on the previous page](#)

† Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can	e.g. HCU Anamix Infant
† Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can	e.g. XMET Maxamaid
† Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can	e.g. XMET Maxamum
† Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle	e.g. HCU Anamix Junior LQ

Isovaleric Acidaemia Products

AMINO ACID FORMULA (WITHOUT LEUCINE) – **Restricted** see terms [on the previous page](#)

† Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can	e.g. IVA Anamix Infant
† Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can	e.g. XLEU Maxamaid
† Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can	e.g. XLEU Maxamum

Maple Syrup Urine Disease Products

AMINO ACID FORMULA (WITHOUT ISOLEUCINE, LEUCINE AND VALINE) – **Restricted** see terms [on the previous page](#)

† Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can	e.g. MSUD Anamix Infant
† Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can	e.g. MSUD Maxamum
† Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle	e.g. MSUD Anamix Junior LQ

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Phenylketonuria Products</b>			
AMINO ACID FORMULA (WITHOUT PHENYLALANINE) – <b>Restricted</b> see terms <a href="#">on page 232</a>			
† Tab 8.33 mg			e.g. Phlexy-10
† Powder 20 g protein, 2.5 g carbohydrate and 0.22 g fibre per 27.8 g sachet			e.g. PKU Lophlex Powder (unflavoured)
† Powder 20 g protein, 3.8 g carbohydrate and 0.23 g fibre per 28 g sachet			e.g. PKU Lophlex Powder (unflavoured)
† Powder 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet			e.g. PKU Anamix Junior (van/choc/unfl)
† Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can			e.g. PKU Anamix Infant
† Powder 13.1 g protein, 50.1 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can			e.g. PKU Anamix Infant
† Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can			e.g. XP Maxamum
† Powder 8.33 g protein and 8.8 g carbohydrate per 20 g sachet			e.g. Phlexy-10
† Liquid 10 g protein, 4.4 g carbohydrate and 0.25 g fibre per 100 ml, 62.5 ml bottle			e.g. PKU Lophlex LQ 10
† Liquid 20 g protein, 8.8 g carbohydrate and 0.34 g fibre per 100 ml, 125 ml bottle			e.g. PKU Lophlex LQ 20
† Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, bottle	13.10	125 ml	PKU Anamix Junior LQ (Berry) PKU Anamix Junior LQ (Orange) PKU Anamix Junior LQ (Unflavoured)
† Liquid 16 g protein, 7 g carbohydrate and 0.27 g fibre per 100 ml, 125 ml bottle			e.g. PKU Lophlex LQ 20
† Liquid 16 g protein, 7 g carbohydrate and 0.27 g fibre per 100 ml, 62.5 ml bottle			e.g. PKU Lophlex LQ 10
† Liquid 16 g protein, 7 g carbohydrate and 0.4 g fibre per 100 ml, 125 ml bottle			e.g. PKU Lophlex LQ 20
† Liquid 16 g protein, 7 g carbohydrate and 0.4 g fibre per 100 ml, 62.5 ml bottle			e.g. PKU Lophlex LQ 10
† Liquid 6.7 g protein, 5.1 g carbohydrate and 2 g fat per 100 ml, 250 ml carton			e.g. Easiphen
† Semi-solid 18.3 g protein, 18.5 g carbohydrate and 0.92 g fibre per 100 g, 109 g pot			e.g. PKU Lophlex Sensations 20 (berries)
(e.g. PKU Lophlex Powder (unflavoured) Powder 20 g protein, 2.5 g carbohydrate and 0.22 g fibre per 27.8 g sachet to be delisted 1 March 2021)			
(e.g. PKU Anamix Infant Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can to be delisted 1 June 2021)			

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

## Propionic Acidaemia and Methylmalonic Acidaemia Products

AMINO ACID FORMULA (WITHOUT Isoleucine, Methionine, Threonine and Valine) – **Restricted** see terms [on page 232](#)

- † Powder 13.1 g protein, 50.1 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can e.g. MMA/PA Anamix Infant
  - † Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can e.g. MMA/PA Anamix Infant
  - † Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can e.g. XMTVI Maxamaid
  - † Powder 39 g protein and 34 g carbohydrate per 100 g, 500 g can e.g. XMTVI Maxamum
- (e.g. MMA/PA Anamix Infant Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can to be delisted 1 March 2021)

## Protein Free Supplements

PROTEIN FREE SUPPLEMENT – **Restricted** see terms [on page 232](#)

- † Powder nil added protein and 67 g carbohydrate per 100 g, 400 g can e.g. Energivit

## Tyrosinaemia Products

AMINO ACID FORMULA (WITHOUT PHENYLALANINE AND TYROSINE) – **Restricted** see terms [on page 232](#)

- † Powder 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet e.g. TYR Anamix Junior
- † Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can e.g. TYR Anamix Infant
- † Powder 25 g protein and 51 g carbohydrate per 100 g, 400 g can e.g. XPHEN, TYR Maxamaid
- † Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle e.g. TYR Anamix Junior LQ

## Urea Cycle Disorders Products

AMINO ACID SUPPLEMENT – **Restricted** see terms [on page 232](#)

- † Powder 25 g protein and 65 g carbohydrate per 100 g, 200 g can e.g. Dialamine
- † Powder 79 g protein per 100 g, 200 g can e.g. Essential Amino Acid Mix

## X-Linked Adrenoleukodystrophy Products

GLYCEROL TRIERUCATE – **Restricted** see terms [on page 232](#)

- † Liquid, 1,000 ml bottle

GLYCEROL TRIOLEATE – **Restricted** see terms [on page 232](#)

- † Liquid, 500 ml bottle

## Specialised Formulas

## Diabetic Products

➔ **Restricted (RS1215)****Initiation**

Any of the following:

- 1 For patients with type I or type II diabetes suffering weight loss and malnutrition that requires nutritional support; or
- 2 For patients with pancreatic insufficiency; or
- 3 For patients who have, or are expected to, eat little or nothing for 5 days; or
- 4 For patients who have a poor absorptive capacity and/or high nutrient losses and/or increased nutritional needs from causes such as catabolism; or
- 5 For use pre- and post-surgery; or
- 6 For patients being tube-fed; or
- 7 For tube-feeding as a transition from intravenous nutrition.

LOW-GI ENTERAL FEED 1 KCAL/ML – **Restricted** see terms [above](#)

† Liquid 5 g protein, 9.6 g carbohydrate and 5.4 g fat per 100 ml, 1,000 ml bottle.....	7.50	1,000 ml	Glucerna Select RTH (Vanilla)
† Liquid 4.3 g protein, 11.3 g carbohydrate and 4.2 g fat per 100 ml, 1,000 ml bag			<i>e.g. Nutrison Advanced Diason</i>

LOW-GI ORAL FEED 1 KCAL/ML – **Restricted** see terms [above](#)

† Liquid 4.5 g protein, 9.8 g carbohydrate, 4.4 g fat and 1.9 g fibre per 100 ml, can.....	2.10	237 ml	Sustagen Diabetic (Vanilla)
† Liquid 5 g protein, 9.6 g carbohydrate and 5.4 g fat per 100 ml, 250 ml bottle.....	1.88	250 ml	Glucerna Select (Vanilla)
† Liquid 6 g protein, 9.5 g carbohydrate, 4.7 g fat and 2.6 g fibre per 100 ml, can.....	2.10	237 ml	Resource Diabetic (Vanilla)
† Liquid 4.9 g protein, 11.7 g carbohydrate, 3.8 g fat and 2 g fibre per 100 ml, 200 ml bottle			<i>e.g. Diasip</i>

## Elemental and Semi-Elemental Products

➔ **Restricted (RS1216)****Initiation**

Any of the following:

- 1 Malabsorption; or
- 2 Short bowel syndrome; or
- 3 Enterocutaneous fistulas; or
- 4 Eosinophilic enteritis (including oesophagitis); or
- 5 Inflammatory bowel disease; or
- 6 Acute pancreatitis where standard feeds are not tolerated; or
- 7 Patients with multiple food allergies requiring enteral feeding.

AMINO ACID ORAL FEED – **Restricted** see terms [above](#)

† Powder 11 g protein, 62 g carbohydrate and 1 g fat per sachet.....	4.50	80 g	Vivonex TEN
--	------	------	-------------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>AMINO ACID ORAL FEED 0.8 KCAL/ML – <b>Restricted</b> see terms <a href="#">on the previous page</a></b>			
⬆ Liquid 2.5 g protein, 11 g carbohydrate and 3.5 g fat per 100 ml, 250 ml carton			<i>e.g. Elemental 028 Extra</i>
<b>PEPTIDE-BASED ENTERAL FEED 1 KCAL/ML – <b>Restricted</b> see terms <a href="#">on the previous page</a></b>			
⬆ Liquid 4 g protein, 17.6 g carbohydrate and 1.7 g fat per 100 ml, 1,000 ml bag			<i>e.g. Nutrison Advanced Peptisorb</i>
⬆ Liquid 4 g protein, 17.7 g carbohydrate and 1.7 g fat per 100 ml, 1,000 ml bag			<i>e.g. Nutrison Advanced Peptisorb</i>
<i>(e.g. Nutrison Advanced Peptisorb Liquid 4 g protein, 17.6 g carbohydrate and 1.7 g fat per 100 ml, 1,000 ml bag to be delisted 1 February 2021)</i>			
<b>PEPTIDE-BASED ENTERAL FEED 1.5 KCAL/ML – <b>Restricted</b> see terms <a href="#">on the previous page</a></b>			
⬆ Liquid 6.75 g protein, 18.4 g carbohydrate and 5.5 g fat per 100 ml, bottle..... 18.06	1,000 ml	Vital	
<b>PEPTIDE-BASED ORAL FEED – <b>Restricted</b> see terms <a href="#">on the previous page</a></b>			
⬆ Powder 13.7 g protein, 62.9 g carbohydrate and 17.5 g fat per 100 g, 400 g can			<i>e.g. Peptamen Junior</i>
⬆ Powder 13.8 g protein, 59 g carbohydrate and 18 g fat per 100 g, 400 g can			<i>e.g. MCT Pepdite; MCT Pepdite 1+</i>
<b>PEPTIDE-BASED ORAL FEED 1 KCAL/ML – <b>Restricted</b> see terms <a href="#">on the previous page</a></b>			
⬆ Liquid 5 g protein, 16 g carbohydrate and 1.69 g fat per 100 ml, carton..... 4.95	237 ml	Peptamen OS 1.0 (Vanilla)	

## Fat Modified Products

**FAT-MODIFIED FEED – **Restricted** see terms [below](#)**

⬆ Powder 12.9 g protein, 69.1 g carbohydrate and 12.9 g fat per 100 g, 400 g can

*e.g. Monogen*

➔ **Restricted (RS1470)**

### Initiation

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

## Hepatic Products

➔ **Restricted (RS1217)**

### Initiation

For children (up to 18 years) who require a liver transplant.

**HEPATIC ORAL FEED – **Restricted** see terms [above](#)**

⬆ Powder 11 g protein, 64 g carbohydrate and 20 g fat per 100 g, can ..... 78.97

400 g Heparon Junior

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

## High Calorie Products

### ➔ Restricted (RS1317)

#### Initiation

Any of the following:

- 1 Patient is fluid volume or rate restricted; or
- 2 Patient requires low electrolyte; or
- 3 Both:
  - 3.1 Any of the following:
    - 3.1.1 Cystic fibrosis; or
    - 3.1.2 Any condition causing malabsorption; or
    - 3.1.3 Faltering growth in an infant/child; or
    - 3.1.4 Increased nutritional requirements; and
  - 3.2 Patient has substantially increased metabolic requirements.

ENTERAL FEED 2 KCAL/ML – **Restricted** see terms [above](#)

↑ Liquid 7.5 g protein, 20 g carbohydrate and 10 g fat per 100 ml, bottle .....	5.50	500 ml	Nutrison Concentrated
↑ Liquid 8.4 g protein, 21.9 g carbohydrate, 9.1 g fat and 0.5 g fibre per 100 ml, bottle .....	11.00	1,000 ml	TwoCal HN RTH (Vanilla)

ORAL FEED 2 KCAL/ML – **Restricted** see terms [above](#)

↑ Liquid 8.4 g protein, 22.4 g carbohydrate, 8.9 g fat and 0.8 g fibre per 100 ml, bottle .....	1.90	200 ml	Two Cal HN
---	------	--------	------------

## High Protein Products

HIGH PROTEIN ENTERAL FEED 1.25 KCAL/ML – **Restricted** see terms [below](#)

↓ Liquid 6.3 g protein, 14.2 g carbohydrate and 4.9 g fat per 100 ml, 1,000 ml bottle			<i>e.g. Nutrison Protein Plus</i>
---	--	--	-----------------------------------

### ➔ Restricted (RS1327)

#### Initiation

Both:

- 1 The patient has a high protein requirement; and
- 2 Any of the following:
  - 2.1 Patient has liver disease; or
  - 2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
  - 2.3 Patient is fluid restricted; or
  - 2.4 Patient's needs cannot be more appropriately met using high calorie product.

HIGH PROTEIN ENTERAL FEED 1.26 KCAL/ML – **Restricted** see terms [below](#)

↓ Liquid 10 g protein, 10.4 g carbohydrate and 4.9 g fat per 100 ml, bottle .....	5.78	500 ml	Nutrison Protein Intense
---	------	--------	--------------------------

### ➔ Restricted (RS1327)

#### Initiation

Both:

- 1 The patient has a high protein requirement; and
- 2 Any of the following:
  - 2.1 Patient has liver disease; or
  - 2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
  - 2.3 Patient is fluid restricted; or
  - 2.4 Patient's needs cannot be more appropriately met using high calorie product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

#### HIGH PROTEIN ENTERAL FEED 1.28 KCAL/ML – **Restricted** see terms [below](#)

↓ Liquid 6.3 g protein, 14.1 g carbohydrate, 4.9 g fat and 1.5 g fibre per 100 ml, 1,000 ml bag

*e.g. Nutrison Protein Plus Multi Fibre*

→ **Restricted (RS1327)**

#### Initiation

Both:

- 1 The patient has a high protein requirement; and
- 2 Any of the following:
  - 2.1 Patient has liver disease; or
  - 2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
  - 2.3 Patient is fluid restricted; or
  - 2.4 Patient's needs cannot be more appropriately met using high calorie product.

### Infant Formulas

#### AMINO ACID FORMULA – **Restricted** see terms [below](#)

↓ Powder 1.95 g protein, 8.1 g carbohydrate and 3.5 g fat per 100 ml, 400 g can

*e.g. Neocate*

↓ Powder 13 g protein, 49 g carbohydrate and 23 g fat per 100 g, 400 g can

*e.g. Neocate SYNEO unflavoured*

↓ Powder 13.3 g protein, 56 g carbohydrate and 22 g fat per 100 g, 400 g can

*e.g. Neocate Junior Unflavoured*

↓ Powder 13.5 g protein, 52 g carbohydrate and 24.5 g fat per 100 g, can ..... 53.00 400 g

*Neocate Gold (Unflavoured)*

↓ Powder 14.8 g protein, 51.4 g carbohydrate and 23 g fat per 100 g, can ..... 53.00 400 g

*Neocate Junior Vanilla*

↓ Powder 15 g protein, 56 g carbohydrate and 20 g fat per 100 g, can ..... 43.60 400 g

*Alfamino Junior*

↓ Powder 2.2 g protein, 7.8 g carbohydrate and 3.4 g fat per 100 ml, can ..... 53.00 400 g

*Elecare LCP*

*(Unflavoured)*

↓ Powder 2.2 g protein, 7.8 g carbohydrate and 3.4 g fat per 100 ml, can ..... 53.00 400 g

*Elecare (Unflavoured)*

*Elecare (Vanilla)*

→ **Restricted (RS1765)**

#### Initiation

Any of the following:

- 1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; or
- 2 History of anaphylaxis to cows' milk protein formula or dairy products; or
- 3 Eosinophilic oesophagitis; or
- 4 Ultra-short gut; or
- 5 Severe Immune deficiency.

#### Continuation

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows' milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
- 3 Amino acid formula is required for a nutritional deficit.

#### ENTERAL LIQUID PEPTIDE FORMULA – **Restricted** see terms [on the next page](#)

↓ Liquid 2.75 g protein, 13.7 g carbohydrate and 3.89 g fat per 100 ml ..... 10.45 500 ml

*Nutrini Peptisorb*

↓ Liquid 4.2 g protein, 18.6 g carbohydrate and 6.58 g fat per 100 ml ..... 15.68 500 ml

*Nutrini Peptisorb Energy*

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ **Restricted (RS1775)**

**Initiation**

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
  - 2.1 Severe malabsorption; or
  - 2.2 Short bowel syndrome; or
  - 2.3 Intractable diarrhoea; or
  - 2.4 Biliary atresia; or
  - 2.5 Cholestatic liver diseases causing malabsorption; or
  - 2.6 Cystic fibrosis; or
  - 2.7 Proven fat malabsorption; or
  - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
  - 2.9 Intestinal failure; or
  - 2.10 Both:
    - 2.10.1 The patient is currently receiving funded amino acid formula; and
    - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
  - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
  - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

**Continuation**

Both:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula.

EXTENSIVELY HYDROLYSED FORMULA – **Restricted** see terms [below](#)

↓ Powder 1.6 g protein, 7.5 g carbohydrate and 3.1 g fat per 100 ml, 900 g can.....	30.42	900 g	Aptamil AllerPro SYNEO 1
↓ Powder 1.6 g protein, 7.8 g carbohydrate and 3.2 g fat per 100 ml, 900 g can.....	30.42	900 g	Aptamil AllerPro SYNEO 2
↓ Powder 14 g protein, 53.4 g carbohydrate and 27.3 g fat per 100 g, 450 g can			<i>e.g. Aptamil Gold+ Pepti Junior</i>

➔ **Restricted (RS1502)**

**Initiation**

Any of the following:

- 1 Both:
  - 1.1 Cows' milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
  - 1.2 Either:
    - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
    - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 For step down from Amino Acid Formula.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

### Continuation

Both:

- 1 An assessment as to whether the infant can be transitioned to a cows' milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula.

### FRUCTOSE-BASED FORMULA

Powder 14.6 g protein, 49.7 g carbohydrate and 30.8 g fat per 100 g,  
400 g can

*e.g. Galactomin 19*

### LACTOSE-FREE FORMULA

Powder 1.3 g protein, 7.3 g carbohydrate and 3.5 g fat per 100 ml, 900 g  
can

*e.g. Karicare Aptamil  
Gold De-Lact*

Powder 1.5 g protein, 7.2 g carbohydrate and 3.6 g fat per 100 ml, 900 g  
can

*e.g. S26 Lactose Free*

### LOW-CALCIUM FORMULA

Powder 14.6 g protein, 53.7 g carbohydrate and 26.1 g fat per 100 g,  
400 g can

*e.g. Locasol*

### PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML – **Restricted** see terms [below](#)

↓ Liquid 2.6 g protein, 10.3 g carbohydrate, 5.4 g fat and 0.6 g fibre per  
100 ml, bottle.....2.35 125 ml Infatrini

### → **Restricted (RS1614)**

#### Initiation – Fluid restricted or volume intolerance with faltering growth

Both:

- 1 Either:
  - 1.1 The patient is fluid restricted or volume intolerant; or
  - 1.2 The patient has increased nutritional requirements due to faltering growth; and
- 2 Patient is under 18 months old and weighs less than 8kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialed appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

### PRETERM FORMULA – **Restricted** see terms [below](#)

↓ Liquid 2.2 g protein, 8.4 g carbohydrate and 4.4 g fat per 100 ml, bottle .....0.75 100 ml S26 LBW Gold RTF

↓ Liquid 2.3 g protein, 8.6 g carbohydrate and 4.2 g fat per 100 ml, 90 ml  
bottle *e.g. Pre Nan Gold RTF*

↓ Liquid 2.6 g protein, 8.4 g carbohydrate and 3.9 g fat per 100 ml, 70 ml  
bottle *e.g. Karicare Aptamil  
Gold+Preterm*

### → **Restricted (RS1224)**

#### Initiation

For infants born before 33 weeks' gestation or weighing less than 1.5 kg at birth.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>THICKENED FORMULA</b>			
Powder 1.8 g protein, 8.1 g carbohydrate and 3.3 g fat per 100 ml, 900 g can			<i>e.g. Karicare Aptamil Thickened AR</i>

## Ketogenic Diet Products

**HIGH FAT FORMULA – Restricted** see terms [below](#)

⚡ Powder 14.4 g protein, 2.9 g carbohydrate and 69.2 g fat per 100 g, can .....35.50	300 g	Ketocal 4:1 (Unflavoured)
⚡ Powder 15.3 g protein, 7.2 g carbohydrate and 67.7 g fat per 100 g, can .....35.50	300 g	Ketocal 4:1 (Vanilla)
		Ketocal 3:1 (Unflavoured)

➡ **Restricted (RS1225)**

### Initiation

For patients with intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

## Paediatric Products

➡ **Restricted (RS1473)**

### Initiation

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
  - 2.1 The child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
  - 2.2 Any condition causing malabsorption; or
  - 2.3 Faltering growth in an infant/child; or
  - 2.4 Increased nutritional requirements; or
  - 2.5 The child is being transitioned from TPN or tube feeding to oral feeding; or
  - 2.6 The child has eaten, or is expected to eat, little or nothing for 3 days.

**PAEDIATRIC ENTERAL FEED 0.76 KCAL/ML – Restricted** see terms [above](#)

⚡ Liquid 2.5 g protein, 12.5 g carbohydrate, 3.3 g fat and 0.7 g fibre per 100 ml, bag.....4.00	500 ml	Nutrini Low Energy Multifibre RTH
---	--------	--------------------------------------

**PAEDIATRIC ENTERAL FEED 1 KCAL/ML – Restricted** see terms [above](#)

⚡ Liquid 2.8 g protein, 11.2 g carbohydrate and 5 g fat per 100 ml, bag.....2.68	500 ml	Pediasure RTH
⚡ Liquid 2.8 g protein, 12.3 g carbohydrate and 4.4 g fat per 100 ml, 500 ml bag		<i>e.g. Nutrini RTH</i>
<b>PAEDIATRIC ENTERAL FEED 1.5 KCAL/ML – Restricted</b> see terms <a href="#">above</a>		
⚡ Liquid 4.1 g protein, 18.5 g carbohydrate, 6.7 g fat and 0.8 g fibre per 100 ml, bag.....6.00	500 ml	Nutrini Energy Multi Fibre
⚡ Liquid 4.1 g protein, 18.5 g carbohydrate and 6.7 g fat per 100 ml, 500 ml bag		<i>e.g. Nutrini Energy RTH</i>

**PAEDIATRIC ORAL FEED 1 KCAL/ML – Restricted** see terms [above](#)

⚡ Liquid 4.2 g protein, 16.7 g carbohydrate and 7.5 g fat per 100 ml, bottle .....1.07	200 ml	Pediasure (Chocolate) Pediasure (Strawberry) Pediasure (Vanilla)
⚡ Liquid 4.2 g protein, 16.7 g carbohydrate and 7.5 g fat per 100 ml, can .....1.34	250 ml	Pediasure (Vanilla)

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>PAEDIATRIC ORAL FEED 1.5 KCAL/ML – Restricted</b> see terms <a href="#">on the previous page</a>			
⬇ Liquid 3.4 g protein, 18.8 g carbohydrate and 6.8 g fat per 100 ml, 200 ml bottle			<i>e.g. Fortini</i>
⬇ Liquid 4.0 g protein, 18.8 g carbohydrate, 6.8 g fat and 1.5 g fibre per 100 ml, 200 ml bottle			<i>e.g. Fortini Multifibre</i>

## Renal Products

**LOW ELECTROLYTE ENTERAL FEED 1.8 KCAL/ML – Restricted** see terms [below](#)

⬇ Liquid 8.1 g protein, 14.74 g carbohydrate, 9.77 g fat and 1.26 g fibre per 100 ml, bottle.....	6.08	500 ml	Nepro HP RTH
--	------	--------	--------------

➔ **Restricted (RS1229)**

### Initiation

For patients with acute or chronic kidney disease.

**LOW ELECTROLYTE ORAL FEED – Restricted** see terms [below](#)

⬇ Powder 7.5 g protein, 59 g carbohydrate and 26.3 g fat per 100 g, 400 g can			<i>e.g. Kindergen</i>
--	--	--	-----------------------

➔ **Restricted (RS1227)**

### Initiation

For children (up to 18 years) with acute or chronic kidney disease.

**LOW ELECTROLYTE ORAL FEED 1.8 KCAL/ML**

⬇ Liquid 8 g protein, 14.74 g carbohydrate, 9.77 g fat and 1.26 g fibre per 100 ml, carton.....	2.67	220 ml	Nepro HP (Strawberry) Nepro HP (Vanilla)
--	------	--------	---

➔ **Restricted (RS1228)**

### Initiation

For patients with acute or chronic kidney disease.

**LOW ELECTROLYTE ORAL FEED 2 KCAL/ML – Restricted** see terms [below](#)

⬇ Liquid 9.1 g protein, 19 g carbohydrate and 10 g fat per 100 ml, carton.....	3.31	237 ml	Novasource Renal (Vanilla)
⬇ Liquid 3 g protein, 25.5 g carbohydrate and 9.6 g fat per 100 ml, 237 ml bottle			
⬇ Liquid 7.5 g protein, 20 g carbohydrate and 10 g fat per 100 ml, 125 ml carton			<i>e.g. Renilon 7.5</i>

➔ **Restricted (RS1228)**

### Initiation

For patients with acute or chronic kidney disease.

## Surgical Products

**HIGH ARGININE ORAL FEED 1.4 KCAL/ML – Restricted** see terms [below](#)

⬇ Liquid 10.1 g protein, 15 g carbohydrate, 4.5 g fat and 0 g fibre per 100 ml, carton.....	4.00	178 ml	Impact Advanced Recovery
--	------	--------	-----------------------------

➔ **Restricted (RS1231)**

### Initiation

Three packs per day for 5 to 7 days prior to major gastrointestinal, head or neck surgery.

**PREOPERATIVE CARBOHYDRATE FEED 0.5 KCAL/ML – Restricted** see terms [on the next page](#)

⬇ Oral liq 0 g protein, 12.6 g carbohydrate and 0 g fat per 100 ml, 200 ml bottle.....	6.80	4	preOp
---	------	---	-------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

### ➔ Restricted (RS1415)

#### Initiation

Maximum of 400 ml as part of an Enhanced Recovery After Surgery (ERAS) protocol 2 to 3 hours before major abdominal surgery.

### Standard Foods

### ➔ Restricted (RS1214)

#### Initiation

Any of the following:

For patients with malnutrition, defined as any of the following:

1 Any of the following:

1.1 BMI < 18.5; or

1.2 Greater than 10% weight loss in the last 3-6 months; or

1.3 BMI < 20 with greater than 5% weight loss in the last 3-6 months; or

2 For patients who have, or are expected to, eat little or nothing for 5 days; or

3 For patients who have a poor absorptive capacity and/or high nutrient losses and/or increased nutritional needs from causes such as catabolism; or

4 For use pre- and post-surgery; or

5 For patients being tube-fed; or

6 For tube-feeding as a transition from intravenous nutrition; or

7 For any other condition that meets the community Special Authority criteria.

ENTERAL FEED 1.5 KCAL/ML – **Restricted** see terms [above](#)

† Liquid 6 g protein, 18.3 g carbohydrate and 5.8 g fat per 100 ml, bag.....	7.00	1,000 ml	Nutrison Energy
† Liquid 6 g protein, 18.4 g carbohydrate, 5.8 g fat and 1.5 g fibre per 100 ml, 1,000 ml bag			<i>e.g. Nutrison Energy Multi Fibre</i>
† Liquid 6.25 g protein, 20 g carbohydrate and 5 g fat per 100 ml, can .....	1.75	250 ml	Ensure Plus HN
† Liquid 6.27 g protein, 20.4 g carbohydrate and 4.9 g fat per 100 ml, bag .....	7.00	1,000 ml	Ensure Plus HN RTH
† Liquid 6.38 g protein, 21.1 g carbohydrate, 4.9 g fat and 1.2 g fibre per 100 ml, bag.....	7.00	1,000 ml	Jevity HiCal RTH

ENTERAL FEED 1 KCAL/ML – **Restricted** see terms [above](#)

† Liquid 4 g protein, 13.6 g carbohydrate and 3.4 g fat per 100 ml, bottle .....	5.29	1,000 ml	Osmolite RTH
† Liquid 4 g protein, 14.1 g carbohydrate, 3.47 g fat and 1.76 g fibre per 100 ml, bottle .....	5.29	1,000 ml	Jevity RTH
† Liquid 4 g protein, 12.3 g carbohydrate and 3.9 g fat per 100 ml, 1,000 ml bag			<i>e.g. NutrisonStdRTH; NutrisonLowSodium</i>
† Liquid 4 g protein, 12.3 g carbohydrate and 3.9 g fat per 100 ml, 1,000 ml bottle			<i>e.g. Nutrison Low Sodium</i>
† Liquid 4 g protein, 12.3 g carbohydrate, 3.9 g fat and 1.5 g fibre per 100 ml, 1000 ml bag			<i>e.g. Nutrison Multi Fibre</i>

ENTERAL FEED 1.2 KCAL/ML – **Restricted** see terms [above](#)

† Liquid 5.55 g protein, 15.1 g carbohydrate, 3.93 g fat and 2 g fibre per 100 ml, 1,000 ml bag			<i>e.g. Jevity Plus RTH</i>
---	--	--	-----------------------------

ENTERAL FEED WITH FIBRE 0.83 KCAL/ML – **Restricted** see terms [above](#)

† Liquid 5.5 g protein, 8.8 g carbohydrate, 2.5 g fat and 1.5 g fibre per 100 ml, bottle .....	5.29	1,000 ml	Nutrison 800 Complete Multi Fibre
--	------	----------	-----------------------------------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>ORAL FEED – Restricted</b> see terms <a href="#">on the previous page</a>			
† Powder 15.9 g protein, 57.4 g carbohydrate and 14 g fat per 100 g, can .....	26.00	850 g	Ensure (Chocolate) Ensure (Vanilla)
† Powder 20.8 g protein, 61 g carbohydrate and 9.4 g fat per 100 g, can .....	8.54	857 g	Fortisip (Vanilla)
† Powder 23 g protein, 65 g carbohydrate and 2.5 g fat per 100 g, can .....	26.00	840 g	Sustagen Hospital Formula Active (Choc) Sustagen Hospital Formula Active (Van)
Note: Community subsidy of Sustagen Hospital Formula is subject to both Special Authority criteria and a manufacturer's surcharge. Higher subsidy by endorsement is available for patients meeting the following endorsement criteria; fat malabsorption, fat intolerance or chyle leak.			
<i>(Fortisip (Vanilla) Powder 20.8 g protein, 61 g carbohydrate and 9.4 g fat per 100 g, can to be delisted 1 August 2021)</i>			
<b>ORAL FEED 1 KCAL/ML – Restricted</b> see terms <a href="#">on the previous page</a>			
† Liquid 3.8 g protein, 23 g carbohydrate and 12.7 g fibre per 100 ml, 237 ml carton			<i>e.g. Resource Fruit Beverage</i>
<b>ORAL FEED 1.5 KCAL/ML – Restricted</b> see terms <a href="#">on the previous page</a>			
† Liquid 5.5 g protein, 21.1 g carbohydrate and 4.81 g fat per 100 ml, can .....	1.33	237 ml	Ensure Plus (Vanilla)
† Liquid 6.25 g protein, 20.2 g carbohydrate and 4.92 g fat per 100 ml, carton.....	1.26	200 ml	Ensure Plus (Banana) Ensure Plus (Chocolate) Ensure Plus (Fruit of the Forest) Ensure Plus (Vanilla) <i>e.g. Fortijuice</i>
† Liquid 4 g protein and 33.5 g carbohydrate per 100 ml, 200 ml bottle			<i>e.g. Fortisip</i>
† Liquid 6 g protein, 18.4 g carbohydrate and 5.8 g fat per 100 ml, 200 ml bottle			<i>e.g. Fortisip</i>
† Liquid 6 g protein, 18.4 g carbohydrate, 5.8 g fat and 2.3 g fibre per 100 ml, 200 ml bottle			<i>e.g. Fortisip Multi Fibre</i>

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

## Bacterial and Viral Vaccines

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE – **Restricted** see terms [below](#)

† Inj 30 IU diphtheria toxoid with 30IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units poliomyelitis virus in 0.5 ml syringe  
– 0% DV Oct-20 to 2024..... 0.00 10 **Infanrix IPV**

➔ **Restricted** ([RS1387](#))

### Initiation

Any of the following:

- 1 A single dose for children up to the age of 7 who have completed primary immunisation; or
- 2 A course of up to four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
- 3 An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 4 Five doses will be funded for children requiring solid organ transplantation.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes

DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE –

**Restricted** see terms [below](#)

† Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin, 80 D-antigen units poliomyelitis virus, 10 mcg hepatitis B  
– 0% DV Oct-20 to 2024..... 0.00 10 **Infanrix-hexa**

➔ **Restricted** ([RS1478](#))

### Initiation

Any of the following:

- 1 Up to four doses for children up to and under the age of 10 for primary immunisation; or
- 2 An additional four doses (as appropriate) are funded for (re-)immunisation for children up to and under the age of 10 who are patients post haematopoietic stem cell transplantation, or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 3 Up to five doses for children up to and under the age of 10 receiving solid organ transplantation.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

## Bacterial Vaccines

BACILLUS CALMETTE-GUERIN VACCINE – **Restricted** see terms [below](#)

† Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin), Danish strain 1331, live attenuated, vial Danish strain 1331, live attenuated, vial with diluent – 0% DV Oct-20 to 2024..... 0.00 10 **BCG Vaccine**

➔ **Restricted** ([RS1233](#))

### Initiation

All of the following:

For infants at increased risk of tuberculosis defined as:

- 1 Living in a house or family with a person with current or past history of TB; and
- 2 Having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; and
- 3 During their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000.

Note: A list of countries with high rates of TB are available at <http://www.health.govt.nz/tuberculosis> (Search for Downloads) or [www.bcgatlas.org/index.php](http://www.bcgatlas.org/index.php)

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

**DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE – Restricted** see terms [below](#)

<p>↓ Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis toxoid, 8 mcg pertussis filamentous haemagglutinin and 2.5 mcg pertactin in 0.5 ml syringe – <b>0% DV Oct-20 to 2024</b>.....</p>	0.00	1 10	<b>Boostrix</b> <b>Boostrix</b>
--	------	---------	------------------------------------

→ **Restricted (RS1766)**

**Initiation**

Any of the following:

- 1 A single dose for pregnant women in the second or third trimester of each pregnancy; or;
- 2 A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or;
- 3 A course of up to four doses is funded for children from age 7 up the age of 18 years inclusive to complete full primary immunisation; or
- 4 An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 5 A single dose for vaccination of patients aged 65 years old; or
- 6 A single dose for vaccination of patients aged 45 years old who have not had 4 previous tetanus doses; or
- 7 For vaccination of previously unimmunised or partially immunised patients; or
- 8 For revaccination following immunosuppression; or
- 9 For boosting of patients with tetanus-prone wounds.

Note: Tdap is not registered for patients aged less than 10 years. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

**HAEMOPHILUS INFLUENZAE TYPE B VACCINE – Restricted** see terms [below](#)

<p>↓ Haemophilus Influenzae type B polysaccharide 10 mcg conjugated to tetanus toxoid as carrier protein 20-40 mcg; prefilled syringe plus vial 0.5 ml .....</p>	0.00	1	Hiberix
--	------	---	---------

→ **Restricted (RS1520)**

**Initiation**

*Therapy limited to 1 dose*

Any of the following:

- 1 For primary vaccination in children; or
- 2 An additional dose (as appropriate) is funded for (re-)immunisation for patients post haematopoietic stem cell transplantation, or chemotherapy; functional asplenic; pre or post splenectomy; pre- or post solid organ transplant, pre- or post cochlear implants, renal dialysis and other severely immunosuppressive regimens; or
- 3 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

**MENINGOCOCCAL (A, C, Y AND W-135) CONJUGATE VACCINE – Restricted** see terms [below](#)

<p>↓ Inj 4 mcg of each meningococcal polysaccharide conjugated to a total of approximately 48 mcg of diphtheria toxoid carrier per 0.5 ml vial – <b>0% DV Oct-20 to 2024</b>.....</p>	0.00	1	<b>Menactra</b>
---	------	---	-----------------

→ **Restricted (RS1778)**

**Initiation**

Either:

- 1 Any of the following:
  - 1.1 Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with HIV, complement deficiency (acquired or inherited), functional or anatomic asplenia or pre or post solid organ transplant; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.2 One dose for close contacts of meningococcal cases; or
- 1.3 A maximum of two doses for bone marrow transplant patients; or
- 1.4 A maximum of two doses for patients following immunosuppression\*; or

## 2 Both:

- 2.1 Person is aged between 13 and 25 years, inclusive; and
- 2.2 Either:

- 2.2.1 One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
- 2.2.2 One dose for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 December 2019 to 30 November 2021.

Notes: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

\*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

**MENINGOCOCCAL C CONJUGATE VACCINE – Restricted** see terms [below](#)

⚡ Inj 10 mcg in 0.5 ml syringe ..... 0.00 1 Neisvac-C

➡ **Restricted (RS1767)**

**Initiation – Children under 9 months of age**

Any of the following:

- 1 Up to three doses for patients pre- and post splenectomy and for patients with HIV, complement deficiency (acquired or inherited), functional or anatomic asplenia or pre or post solid organ transplant; or
- 2 Two doses for close contacts of meningococcal cases; or
- 3 A maximum of two doses for bone marrow transplant patients; or
- 4 A maximum of two doses for patients pre- and post-immunosuppression\*.

Notes: children under nine months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for booster schedules with meningococcal ACWY vaccine.

\*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

**PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE – Restricted** see terms [below](#)

⚡ mcg of pneumococcal polysaccharide serotypes 1, 5, 6B, 7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml prefilled syringe – **0% DV Oct-20 to 2024** ..... 0.00 10 **Synflorix**

➡ **Restricted (RS1768)**

**Initiation**

A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive.

Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

**PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE – Restricted** see terms [below](#)

⚡ Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5 ml syringe ..... 0.00 1 Prevenar 13  
10 Prevenar 13

➡ **Restricted (RS1769)**

**Initiation – High risk children who have received PCV10**

*Therapy limited to 1 dose*

Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10.

**Initiation – High risk children aged under 5 years**

*Therapy limited to 4 doses*

Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Up to an additional four doses (as appropriate) are funded for children aged under 5 years for (re-)immunisation; and
- 2 Any of the following:
  - 2.1 On immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
  - 2.2 With primary immune deficiencies; or
  - 2.3 With HIV infection; or
  - 2.4 With renal failure, or nephrotic syndrome; or
  - 2.5 Who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
  - 2.6 With cochlear implants or intracranial shunts; or
  - 2.7 With cerebrospinal fluid leaks; or
  - 2.8 Receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
  - 2.9 With chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
  - 2.10 Pre term infants, born before 28 weeks gestation; or
  - 2.11 With cardiac disease, with cyanosis or failure; or
  - 2.12 With diabetes; or
  - 2.13 With Down syndrome; or
  - 2.14 Who are pre-or post-splenectomy, or with functional asplenia.

#### Initiation – High risk adults and children 5 years and over

*Therapy limited to 4 doses*

Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency.

#### Initiation – Testing for primary immunodeficiency diseases

For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

**PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE – Restricted** see terms [below](#)

↓ Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype) – **0% DV Oct-20 to 2024**.....0.00 1 **Pneumovax 23**

→ **Restricted (RS1587)**

#### Initiation – High risk patients

*Therapy limited to 3 doses*

For patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy; or with functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency.

#### Initiation – High risk children

*Therapy limited to 2 doses*

Both:

- 1 Patient is a child under 18 years for (re-)immunisation; and
- 2 Any of the following:
  - 2.1 On immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
  - 2.2 With primary immune deficiencies; or
  - 2.3 With HIV infection; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.4 With renal failure, or nephrotic syndrome; or
- 2.5 Who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
- 2.6 With cochlear implants or intracranial shunts; or
- 2.7 With cerebrospinal fluid leaks; or
- 2.8 Receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
- 2.9 With chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
- 2.10 Pre term infants, born before 28 weeks gestation; or
- 2.11 With cardiac disease, with cyanosis or failure; or
- 2.12 With diabetes; or
- 2.13 With Down syndrome; or
- 2.14 Who are pre-or post-splenectomy, or with functional asplenia.

**Initiation – Testing for primary immunodeficiency diseases**

For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

SALMONELLA TYPHI VACCINE – **Restricted** see terms [below](#)

⬇ Inj 25 mcg in 0.5 ml syringe

➡ **Restricted (RS1243)**

**Initiation**

For use during typhoid fever outbreaks.

Viral Vaccines

HEPATITIS A VACCINE – **Restricted** see terms [below](#)

⬇ Inj 720 ELISA units in 0.5 ml syringe – <b>0% DV Oct-20 to 2024</b> .....	0.00	1	Havrix Junior
⬇ Inj 1440 ELISA units in 1 ml syringe – <b>0% DV Oct-20 to 2024</b> .....	0.00	1	Havrix

➡ **Restricted (RS1638)**

**Initiation**

Any of the following:

- 1 Two vaccinations for use in transplant patients; or
- 2 Two vaccinations for use in children with chronic liver disease; or
- 3 One dose of vaccine for close contacts of known hepatitis A cases.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

## HEPATITIS B RECOMBINANT VACCINE

↓ Inj 10 mcg per 0.5 ml prefilled syringe ..... 0.00 1 Engerix-B

→ **Restricted (RS1588)**

**Initiation**

Any of the following:

- 1 For household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2 For children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3 For children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4 For HIV positive patients; or
- 5 For hepatitis C positive patients; or
- 6 for patients following non-consensual sexual intercourse; or
- 7 For patients following immunosuppression; or
- 8 For solid organ transplant patients; or
- 9 For post-haematopoietic stem cell transplant (HSCT) patients; or
- 10 Following needle stick injury.

↓ Inj 20 mcg per 1 ml prefilled syringe – **0% DV Oct-20 to 2024** ..... 0.00 1 Engerix-B

→ **Restricted (RS1671)**

**Initiation**

Any of the following:

- 1 For household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2 For children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3 For children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4 For HIV positive patients; or
- 5 For hepatitis C positive patients; or
- 6 for patients following non-consensual sexual intercourse; or
- 7 For patients following immunosuppression; or
- 8 For solid organ transplant patients; or
- 9 For post-haematopoietic stem cell transplant (HSCT) patients; or
- 10 Following needle stick injury; or
- 11 For dialysis patients; or
- 12 For liver or kidney transplant patients.

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV] – **Restricted** see terms [below](#)

↓ Inj 270 mcg in 0.5 ml syringe – **0% DV Oct-20 to 2024** ..... 0.00 10 Gardasil 9

→ **Restricted (RS1693)**

**Initiation – Children aged 14 years and under**

*Therapy limited to 2 doses*

Children aged 14 years and under.

**Initiation – other conditions**

Either:

- 1 Up to 3 doses for people aged 15 to 26 years inclusive; or
- 2 Both:
  - 2.1 People aged 9 to 26 years inclusive; and
  - 2.2 Any of the following:
    - 2.2.1 Up to 3 doses for confirmed HIV infection; or
    - 2.2.2 Up to 3 doses for transplant (including stem cell) patients; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2.2.3 Up to 4 doses for Post chemotherapy.

## Initiation – Recurrent Respiratory Papillomatosis

All of the following:

- 1 Either:
  - 1.1 Maximum of two doses for children aged 14 years and under; or
  - 1.2 Maximum of three doses for people aged 15 years and over; and
- 2 The patient has recurrent respiratory papillomatosis; and
- 3 The patient has not previously had an HPV vaccine.

## INFLUENZA VACCINE

⚡ Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine).....	9.00	1	Afluria Quad Junior (2020 Formulation)
--	------	---	---

### ➡ Restricted (RS1675)

## Initiation – cardiovascular disease for patients aged 6 months to 35 months

Any of the following:

- 1 Ischaemic heart disease; or
- 2 Congestive heart failure; or
- 3 Rheumatic heart disease; or
- 4 Congenital heart disease; or
- 5 Cerebro-vascular disease.

Note: hypertension and/or dyslipidaemia without evidence of end-organ disease is excluded from funding.

## Initiation – chronic respiratory disease for patients aged 6 months to 35 months

Either:

- 1 Asthma, if on a regular preventative therapy; or
- 2 Other chronic respiratory disease with impaired lung function.

Note: asthma not requiring regular preventative therapy is excluded from funding.

## Initiation – Other conditions for patients aged 6 months to 35 months

Any of the following:

- 1 Diabetes; or
- 2 Chronic renal disease; or
- 3 Any cancer, excluding basal and squamous skin cancers if not invasive; or
- 4 Autoimmune disease; or
- 5 Immune suppression or immune deficiency; or
- 6 HIV; or
- 7 Transplant recipient; or
- 8 Neuromuscular and CNS diseases/ disorders; or
- 9 Haemoglobinopathies; or
- 10 Is a child on long term aspirin; or
- 11 Has a cochlear implant; or
- 12 Errors of metabolism at risk of major metabolic decompensation; or
- 13 Pre and post splenectomy; or
- 14 Down syndrome; or
- 15 Child who has been hospitalised for respiratory illness or has a history of significant respiratory illness.

⚡ Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine).....	90.00	10	Afluria Quad (2020 Formulation)
	9.00	1	Influvac Tetra (2020 formulation)

### ➡ Restricted (RS1674)

## Initiation – People over 65

The patient is 65 years of age or over.

continued...

Price (ex man. excl. GST)	Brand or Generic Manufacturer
\$ Per	

continued...

**Initiation – cardiovascular disease for patients 3 years and over**

Any of the following:

- 1 Ischaemic heart disease; or
- 2 Congestive heart failure; or
- 3 Rheumatic heart disease; or
- 4 Congenital heart disease; or
- 5 Cerebro-vascular disease.

Note: hypertension and/or dyslipidaemia without evidence of end-organ disease is excluded from funding.

**Initiation – chronic respiratory disease for patients 3 years and over**

Either:

- 1 Asthma, if on a regular preventative therapy; or
- 2 Other chronic respiratory disease with impaired lung function.

Note: asthma not requiring regular preventative therapy is excluded from funding.

**Initiation – Other conditions for patients 3 years and over**

Either:

- 1 Any of the following:
  - 1.1 Diabetes; or
  - 1.2 chronic renal disease; or
  - 1.3 Any cancer, excluding basal and squamous skin cancers if not invasive; or
  - 1.4 Autoimmune disease; or
  - 1.5 Immune suppression or immune deficiency; or
  - 1.6 HIV; or
  - 1.7 Transplant recipient; or
  - 1.8 Neuromuscular and CNS diseases/ disorders; or
  - 1.9 Haemoglobinopathies; or
  - 1.10 Is a child on long term aspirin; or
  - 1.11 Has a cochlear implant; or
  - 1.12 Errors of metabolism at risk of major metabolic decompensation; or
  - 1.13 Pre and post splenectomy; or
  - 1.14 Down syndrome; or
  - 1.15 Is pregnant; or
  - 1.16 Is a child aged four and under who has been hospitalised for respiratory illness or has a history of significant respiratory illness; or
- 2 Patients in a long-stay inpatient mental health care unit or who are compulsorily detained long-term in a forensic unit within a DHB hospital.

MEASLES, MUMPS AND RUBELLA VACCINE – **Restricted** see terms [below](#)

↓ Injection, measles virus 1,000 CCID <sub>50</sub> , mumps virus 5,012 CCID <sub>50</sub> , Rubella virus 1,000 CCID <sub>50</sub> ; prefilled syringe/ampoule of diluent 0.5 ml – <b>0% DV Oct-20 to 2024</b>	0.00	10	<b>Priorix</b>
---	------	----	----------------

→ **Restricted (RS1487)**

**Initiation – first dose prior to 12 months**

*Therapy limited to 3 doses*

Any of the following:

- 1 For primary vaccination in children; or
- 2 For revaccination following immunosuppression; or
- 3 For any individual susceptible to measles, mumps or rubella.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

## Initiation – first dose after 12 months

*Therapy limited to 2 doses*

Any of the following:

- 1 For primary vaccination in children; or
- 2 For revaccination following immunosuppression; or
- 3 For any individual susceptible to measles, mumps or rubella.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

POLIOMYELITIS VACCINE – **Restricted** see terms [below](#)

⚡ Inj 80 D-antigen units in 0.5 ml syringe – **0% DV Oct-20 to 2024**..... 0.00 1 **IPOL**

➡ **Restricted (RS1398)**

## Initiation

*Therapy limited to 3 doses*

Either:

- 1 For partially vaccinated or previously unvaccinated individuals; or
- 2 For revaccination following immunosuppression.

Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

## RABIES VACCINE

Inj 2.5 IU vial with diluent

ROTA VIRUS ORAL VACCINE – **Restricted** see terms [below](#)

⚡ Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose,  
prefilled oral applicator – **0% DV Oct-20 to 2024**..... 0.00 10 **Rotarix**

➡ **Restricted (RS1590)**

## Initiation

*Therapy limited to 2 doses*

Both:

- 1 First dose to be administered in infants aged under 14 weeks of age; and
- 2 No vaccination being administered to children aged 24 weeks or over.

## VARICELLA VACCINE [CHICKENPOX VACCINE]

⚡ Inj 1350 PFU prefilled syringe – **0% DV Oct-20 to 2024**..... 0.00 1 **Varivax**  
10 **Varivax**

➡ **Restricted (RS1591)**

## Initiation – primary vaccinations

*Therapy limited to 1 dose*

Either:

- 1 Any infant born on or after 1 April 2016; or
- 2 For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox).

## Initiation – other conditions

*Therapy limited to 2 doses*

Any of the following:

- 1 Any of the following:  
for non-immune patients:
  - 1.1 With chronic liver disease who may in future be candidates for transplantation; or
  - 1.2 With deteriorating renal function before transplantation; or
  - 1.3 Prior to solid organ transplant; or
  - 1.4 Prior to any elective immunosuppression\*; or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1.5 For post exposure prophylaxis who are immune competent inpatients; or
- 2 For patients at least 2 years after bone marrow transplantation, on advice of their specialist; or
- 3 For patients at least 6 months after completion of chemotherapy, on advice of their specialist; or
- 4 For HIV positive patients non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
- 5 For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or
- 6 For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or
- 7 For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.

Note: \* immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

↓ Inj 2000 PFU prefilled syringe plus vial

➔ **Restricted (RS1777)**

**Initiation – infants between 9 and 12 months of age**

*Therapy limited to 2 doses*

Any of the following:

- 1 Any of the following:
  - for non-immune patients:
    - 1.1 With chronic liver disease who may in future be candidates for transplantation; or
    - 1.2 With deteriorating renal function before transplantation; or
    - 1.3 Prior to solid organ transplant; or
    - 1.4 Prior to any elective immunosuppression\*; or
    - 1.5 For post exposure prophylaxis who are immune competent inpatients; or
  - 2 For patients at least 2 years after bone marrow transplantation, on advice of their specialist; or
  - 3 For patients at least 6 months after completion of chemotherapy, on advice of their specialist; or
  - 4 For HIV positive patients non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
  - 5 For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or
  - 6 For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or
  - 7 For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.

Note: \* immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

VARICELLA ZOSTER VACCINE [SHINGLES VACCINE] – **Restricted** see terms [below](#)

↓ Varicella zoster virus (Oka strain) live attenuated vaccine [shingles

vaccine] .....	0.00	1	Zostavax
		10	Zostavax

➔ **Restricted (RS1779)**

**Initiation – people aged 65 years**

*Therapy limited to 1 dose*

One dose for all people aged 65 years.

**Initiation – people aged between 66 and 80 years**

*Therapy limited to 1 dose*

One dose for all people aged between 66 and 80 years inclusive from 1 April 2018 and 31 December 2021.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Diagnostic Agents

TUBERCULIN PPD [MANTOUX] TEST			
Inj 5 TU per 0.1 ml, 1 ml vial – 0% DV Oct-20 to 2024 .....	0.00	1	Tubersol

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
<b>Optional Pharmaceuticals</b>			
<b>NOTE:</b>			
In addition to the products expressly listed here in Part III: Optional Pharmaceuticals, a range of hospital medical devices are listed in an addendum to Part III which is available at <a href="http://schedule.pharmac.govt.nz">schedule.pharmac.govt.nz</a> . The Optional Pharmaceuticals listed in the addendum are deemed to be listed in Part III, and the Rules of the Pharmaceutical Schedule applying to products listed in Part III apply to them.			
<b>BLOOD GLUCOSE DIAGNOSTIC TEST METER</b>			
1 meter with 50 lancets, a lancing device, and 10 diagnostic test strips .....	20.00	1	CareSens N Premier
	10.00		Caresens N
			Caresens N POP
<b>BLOOD GLUCOSE DIAGNOSTIC TEST STRIP</b>			
Blood glucose test strips.....	10.56	50 test	CareSens N
Test strips .....	10.56	50 test	CareSens PRO
<b>BLOOD KETONE DIAGNOSTIC TEST STRIP</b>			
Test strips .....	15.50	10 strip	KetoSens
<b>DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER</b>			
Meter with 50 lancets, a lancing device, and 10 blood glucose diagnostic test strips .....	20.00	1	CareSens Dual
<b>MASK FOR SPACER DEVICE</b>			
Small.....	2.20	1	e-chamber Mask
<b>PEAK FLOW METER</b>			
Low Range .....	9.54	1	Mini-Wright AFS Low Range
Normal Range .....	9.54	1	Mini-Wright Standard
<b>PREGNANCY TEST - HCG URINE</b>			
Cassette .....	12.00	40 test	Smith BioMed Rapid Pregnancy Test
<b>SODIUM NITROPRUSSIDE</b>			
Test strip.....	22.00	50 strip	Ketostix
<b>SPACER DEVICE</b>			
220 ml (single patient) .....	2.95	1	e-chamber Turbo
510 ml (single patient) .....	5.12	1	e-chamber La Grande
800 ml.....	6.50	1	Volumatic

<b>- Symbols -</b>		
8-methoxypsoralen.....	58	(2020 Formulation).....252
<b>- A -</b>		
A-Scabies.....	55	AFT Pholcodine Linctus BP.....208
Abacavir sulphate.....	88	Agents Affecting the
Abacavir sulphate with		Renin-Angiotensin System.....39
lamivudine.....	88	Agents for Parkinsonism and Related
Abciximab.....	158	Disorders.....106
Abiraterone acetate.....	147	Agents Used in the Treatment of
Acarbose.....	9	Poisonings.....219
Accuretic 10.....	39	Ajmaline.....41
Accuretic 20.....	39	Albendazole.....85
Acetazolamide.....	216	Aldurazyme.....16
Acetec.....	39	Alecensa.....139
Acetic acid		Alectinib.....139
Extemporaneously Compounded		Alendronate sodium.....97
Preparations.....	227	Alendronate sodium with
Genito-Urinary.....	60	colecalfiferol.....97
Acetic acid with hydroxyquinoline,		Alfacalcidol.....22
glycerol and ricinoleic acid.....	60	Alfamino Junior.....239
Acetic acid with propylene		Alfentanil.....111
glycol.....	218	Alglucosidase alfa.....13
Acetylcholine chloride.....	216	Alinia.....86
Acetylcysteine.....	219	Allersoothe.....205
Aciclovir		Allmercap.....133
Infections.....	91	Allopurinol.....102
Sensory.....	212	Alpha tocopheryl.....22
Aciclovir-Baxter.....	91	Alpha tocopheryl acetate.....23
Aciclovir-Claris.....	91	Alpha-Adrenoceptor Blockers.....40
Acid Citrate Dextrose A.....	31	Alphamox.....78
Acidex.....	5	Alphamox 125.....78
Acipimox.....	47	Alphamox 250.....78
Acitretin.....	58	Alprolix.....28
Aclasta.....	98	Alprostadii hydrochloride.....49
Actemra.....	194	Alteplase.....33
Actinomycin D.....	131	Alum.....227
Adalat 10.....	44	Aluminium chloride.....26
Adalat Oros.....	44	Aluminium hydroxide.....5
Adalimumab.....	158	Aluminium hydroxide with
Adapalene.....	55	magnesium hydroxide and
Adefovir dipivoxil.....	90	simeticone.....5
Adenocor.....	41	Amantadine hydrochloride.....106
Adenosine.....	41	AmBisome.....82
Adenuric.....	102	Ambrisentan.....50
Adrenaline.....	48	Ambrisentan Mylan.....50
Advantan.....	57	Amethocaine
Advate.....	29	Nervous.....110
Adynovate.....	30	Sensory.....215
Aerrane.....	107	Amikacin.....74
Afinitor.....	202	Amiloride hydrochloride.....45
Aflibercept.....	168	Amiloride hydrochloride with
Afluria Quad		furosemide.....45
(2020 Formulation).....	252	Amiloride hydrochloride with
Afluria Quad Junior		hydrochlorothiazide.....45
		Aminolevulinic acid
		hydrochloride.....150
		Aminophylline.....210
		Amiodarone hydrochloride.....41
		Amisulpride.....120
		Amitriptyline.....113
		Amlodipine.....43
		Amorolfine.....54
		Amoxicillin.....78
		Amoxicillin with clavulanic acid.....78
		Amphotericin B
		Alimentary.....20
		Infections.....82
		Amsacrine.....134
		Amyl nitrite.....219
		Anabolic Agents.....64
		Anaesthetics.....107
		Anagrelide hydrochloride.....134
		Analgesics.....110
		Anastrozole.....150
		Anatrole.....150
		Andriol Testocaps.....64
		Androderm.....64
		Androgen Agonists and
		Antagonists.....64
		Anoro Ellipta.....206
		Antabuse.....128
		Antacids and Antiflatulents.....5
		Anti-Infective Agents.....60
		Anti-Infective Preparations
		Dermatological.....54
		Sensory.....212
		Anti-Inflammatory Preparations.....213
		Antiacne Preparations.....55
		Antiallergy Preparations.....204
		Antianaemics.....24
		Antiarrhythmics.....41
		Antibacterials.....74
		Anticholinergic Agents.....205
		Anticholinesterases.....97
		Antidepressants.....113
		Antidiarrhoeals and Intestinal
		Anti-Inflammatory Agents.....5
		Antiepilepsy Drugs.....115
		Antifibrinolytics, Haemostatics and
		Local Sclerosants.....26
		Antifibrotics.....206
		Antifungals.....82
		Antihypotensives.....42
		Antimigraine Preparations.....118
		Antimycobacterials.....84
		Antinausea and Vertigo Agents.....119
		Antiparasitics.....85
		Antipruritic Preparations.....55
		Antipsychotic Agents.....120
		Antiretrovirals.....87

Antirheumatoid Agents .....	97	Aripiprazole.....	120	Avonex.....	124
Antiseptics and Disinfectants.....	221	Aripiprazole Sandoz .....	120	Avonex Pen .....	124
Antispasmodics and Other Agents		Aristocort.....	58	Azacididine .....	132
Altering Gut Motility .....	7	Arrow-Amitriptyline .....	113	Azacididine Dr Reddy's .....	132
Antithrombotics.....	30	Arrow-Bendrofluazide .....	46	Azactam.....	80
Antithymocyte globulin		Arrow-Brimonidine .....	217	Azamun.....	202
(equine) .....	201	Arrow-Calcium .....	18	Azathioprine.....	202
Antithymocyte globulin (rabbit) .....	201	Arrow-Diazepam.....	123	Azithromycin.....	76
Antilucerants.....	7	Arrow-Fluoxetine .....	115	Azol.....	67
Antivirals.....	90	Arrow-Losartan &		AZT.....	88
Anxiolytics.....	123	Hydrochlorothiazide.....	40	Aztreonam .....	80
Apidra .....	10	Arrow-Morphine LA.....	112		
Apidra Solostar .....	10	Arrow-Norfloxacin .....	79	<b>- B -</b>	
Apo-Amlodipine .....	43	Arrow-Ornidazole.....	86	Bacillus calmette-guerin (BCG) .....	202
Apo-Azithromycin .....	76	Arrow-Quinapril 10.....	39	Bacillus calmette-guerin	
Apo-Ciclopirox .....	54	Arrow-Quinapril 20.....	39	vaccine .....	246
Apo-Cilazapril/		Arrow-Quinapril 5.....	39	Baclofen.....	102
Hydrochlorothiazide .....	39	Arrow-Roxithromycin .....	77	Bacterial and Viral Vaccines.....	246
Apo-Clarithromycin .....	77	Arrow-Timolol .....	216	Bacterial Vaccines .....	246
Apo-Clomipramine.....	113	Arrow-Topiramate .....	117	Balanced Salt Solution .....	215
Apo-Diclo SR .....	104	Arrow-Tramadol .....	113	Barium sulphate.....	222
Apo-Diltiazem CD .....	44	Arsenic trioxide .....	134	Barium sulphate with sodium	
Apo-Doxazosin .....	40	Artemether with lumefantrine.....	85	bicarbonate.....	222
Apo-Folic Acid .....	25	Artesunate .....	86	Barrier Creams and Emollients.....	55
Apo-Furosemide .....	45	Articaine hydrochloride .....	108	Basiliximab .....	169
Apo-Gabapentin .....	116	Articaine hydrochloride with		BCG Vaccine .....	246
Apo-Megestrol .....	149	adrenaline.....	108	BD PosiFlush.....	37
Apo-Metoprolol .....	43	Asacol.....	6	Beclazone 100.....	208
Apo-Mirtazapine .....	114	Asamax.....	6	Beclazone 250.....	208
Apo-Nadolol.....	43	Ascorbic acid		Beclazone 50.....	208
Apo-Nicotinic Acid .....	47	Alimentary .....	22	Beclomethasone dipropionate .....	208
Apo-Oxybutynin .....	63	Extemporaneously Compounded		Bee venom .....	204
Apo-Perindopril.....	39	Preparations .....	227	Bendamustine hydrochloride .....	130
Apo-Pindolol .....	43	Aspen Adrenaline .....	48	Bendrofluazide .....	46
Apo-Pravastatin.....	46	Aspirin		Bendroflumethiazide	
Apo-Prazosin .....	41	Blood.....	32	[Bendrofluazide].....	46
Apo-Prednisone.....	66	Nervous.....	110	Benzathine benzylpenicillin .....	78
Apo-Propranolol.....	43	Asthalin .....	208	Benzatropine mesylate .....	106
Apo-Pyridoxine .....	21	Atazanavir sulphate .....	89	Benzbromaron AL 100.....	102
Apo-Sumatriptan.....	118	Atenolol.....	42	Benzbromarone.....	102
Apo-Terazosin .....	41	Atenolol-AFT.....	42	Benzocaine.....	108
Apomorphine hydrochloride.....	106	ATGAM.....	201	Benzocaine with tetracaine	
Apraclonidine.....	217	Ativan.....	123	hydrochloride.....	108
Aprepitant .....	119	Atomoxetine.....	125	Benzoin.....	227
Apresoline.....	49	Atorvastatin .....	46	Benzoyl peroxide.....	55
Aprotinin .....	26	Atovaquone with proguanil		Benzotrop .....	106
Aptamil AllerPro SYNEO 1 .....	240	hydrochloride .....	86	Benzylamine hydrochloride .....	19
Aptamil AllerPro SYNEO 2 .....	240	Atracurium besylate .....	102	Benzylamine hydrochloride with	
Aqueous cream .....	56	Atropine sulphate		cetylpyridinium chloride .....	19
Arachis oil [Peanut oil].....	227	Cardiovascular .....	41	Benzylpenicillin sodium [Penicillin	
Aratac .....	41	Sensory.....	217	G].....	78
Arava .....	97	Atropt .....	217	Beractant .....	211
Arginine		Aubagio .....	124	Beta Cream .....	57
Alimentary .....	14	Augmentin .....	78	Beta Ointment.....	57
Various .....	223	Aurorix .....	114	Beta Scalp .....	59
Argipressin [Vasopressin].....	73	Avelox.....	79	Beta-Adrenoceptor Agonists.....	208
				Beta-Adrenoceptor Blockers.....	42

Betadine .....	221	Bosentan .....	50	Calcium Homeostasis .....	64	
Betahistine dihydrochloride .....	119	Bosentan Dr Reddy's .....	50	Calcium polystyrene sulphonate .....	37	
Betaine .....	14	Bosvate .....	42	Calcium Resonium .....	37	
Betaloc CR .....	43	Botox .....	103	Candesartan cilexetil .....	39	
Betamethasone .....	65	Botulism antitoxin .....	219	Candestar .....	39	
Betamethasone dipropionate .....	57	Bplex .....	22	Capecitabine .....	132	
Betamethasone dipropionate with calcipotriol .....	58	Breo Ellipta .....	210	Capercit .....	132	
Betamethasone sodium phosphate with betamethasone acetate .....	65	Brevinor 1/28 .....	60	Capoten .....	39	
Betamethasone valerate .....	57, 59	Bricanyl Turbuhaler .....	208	Capsaicin Musculoskeletal .....	105	
Betamethasone valerate with clioquinol .....	58	Bridion .....	103	Nervous .....	110	
Betamethasone valerate with sodium fusidate [Fusidic acid] .....	58	Brilinta .....	32	Captopril .....	39	
Betaxolol .....	216	Brimonidine tartrate .....	217	Carbachol .....	216	
Betnovate .....	57	Brimonidine tartrate with timolol .....	217	Carbamazepine .....	115	
Betoptic .....	216	Brinzolamide .....	216	Carbasorb-X .....	220	
Betoptic S .....	216	Bromocriptine .....	106	Carbimazole .....	72	
Bevacizumab .....	169	Budesonide .....		Carbomer .....	217	
Bezafibrate .....	46	Alimentary .....	5	Carboplatin .....	139	
Bezalip .....	46	Respiratory .....	205, 209	Carboplatin Ebewe .....	139	
Bezalip Retard .....	46	Budesonide with eformoterol .....	209	Carboprost trometamol .....	61	
Bicalutamide .....	148	Bumetanide .....	45	Carboxymethylcellulose Alimentary .....	19	
Bicillin LA .....	78	Bupafen .....	109	Extemporaneously Compounded Preparations .....	227	
BiCNU .....	131	Bupivacaine hydrochloride .....	108	Cardinol LA .....	43	
Bicnu Heritage .....	131	Bupivacaine hydrochloride with adrenaline .....	108	CareSens Dual .....	257	
Bile and Liver Therapy .....	9	Bupivacaine hydrochloride with fentanyl .....	109	Caresens N .....	257	
Biliscopin .....	223	Bupivacaine hydrochloride with glucose .....	109	Caresens N POP .....	257	
Bimatoprost .....	216	Buprenorphine Naloxone BNM .....	128	CareSens N Premier .....	257	
Bimatoprost Multichem .....	216	Buprenorphine with naloxone .....	128	CareSens PRO .....	257	
Binarex .....	148	Bupropion hydrochloride .....	128	Carmellose sodium with pectin and gelatine .....		
Binocrit .....	24	Burinex .....	45	Alimentary .....	19	
Biodone .....	112	Buscopan .....	7	Sensory .....	217	
Biodone Extra Forte .....	112	Buserelin .....	68	Carmustine .....	131	
Biodone Forte .....	112	Buspirone hydrochloride .....	123	Carvedilol .....	42	
Biotin .....	15	Busulfan .....	131	Carvedilol Sandoz .....	42	
Bisacodyl .....	13	<b>- C -</b>			Caspofungin .....	83
Bismuth subgallate .....	227	Cabergoline .....	67	Catapres .....	45	
Bismuth subnitrate and iodoform paraffin .....	225	Caffeine .....	125	Ceenu .....	131	
Bisoprolol fumarate .....	42	Caffeine citrate .....	210	Cefaclor .....	75	
Bisoprolol Mylan .....	42	Calamine .....	55	Cefalexin .....	75	
Bivalirudin .....	30	Calci-Tab 500 .....	18	Cefalexin Sandoz .....	75	
Bleomycin sulphate .....	131	Calcipotriol .....	58	Cefazolin .....	75	
Blood glucose diagnostic test meter .....	257	Calcitonin .....	64	Cefepime .....	76	
Blood glucose diagnostic test strip .....	257	Calcitriol .....	22	Cefepime-AFT .....	76	
Blood ketone diagnostic test strip .....	257	Calcitriol-AFT .....	22	Cefotaxime .....	75	
Bonney's blue dye .....	224	Calcium carbonate .....	5, 18	Cefotaxime Sandoz .....	75	
Boostrix .....	247	Calcium Channel Blockers .....	43	Cefoxitin .....	75	
Boric acid .....	227	Calcium chloride .....	35	Cefoxitin Actavis .....	75	
Bortezomib .....	134	Calcium folinate .....	147	Ceftaroline fosamil .....	76	
Bortezomib Dr-Reddy's .....	134	Calcium Folate Ebewe .....	147	Ceftazidime .....	75	
		Calcium Folate Sandoz .....	147	Ceftazidime-AFT .....	75	
		Calcium gluconate Blood .....	35	Ceftriaxone .....	75	
		Dermatological .....	59	Ceftriaxone-AFT .....	75	
				Cefuroxime .....	75	

Cefuroxime Actavis.....	75	hydrocortisone.....	7	sulphur.....	58
Celecoxib.....	103	Cipfloxx.....	79	Cocaine hydrochloride.....	109
Celecoxib Pfizer.....	103	Ciprofloxacin.....		Cocaine hydrochloride with	
Celiprolol.....	42	Infections.....	79	adrenaline.....	109
CellCept.....	202	Sensory.....	212	Codeine phosphate	
Celol.....	42	Ciprofloxacin Teva.....	212	Extemporaneously Compounded	
Centrally-Acting Agents.....	44	Ciprofloxacin with		Preparations.....	227
Cephalexin ABM.....	75	hydrocortisone.....	212	Nervous.....	111
Cetirizine hydrochloride.....	205	Ciproxin HC Otic.....	212	Colchicine.....	102
Cetomacrogol.....	56	Circadin.....	124	Colecalciferol.....	22
Cetomacrogol with glycerol.....	56	Cisplatin.....	139	Colestimethate.....	80
Cetrimide.....	227	Citalopram hydrobromide.....	114	Colestipol hydrochloride.....	47
Cetuximab.....	170	Citanest.....	110	Colgout.....	102
Charcoal.....	220	Citrate sodium.....	30	Colifoam.....	6
Chemotherapeutic Agents.....	130	Citric acid.....	227	Colistin sulphomethate	
Chickenpox vaccine.....	254	Citric acid with magnesium oxide and		[Coolestimethate].....	80
Chlorafast.....	212	sodium picosulfate.....	12	Colistin-Link.....	80
Chloral hydrate.....	124	Citric acid with sodium		Colloidon flexible.....	227
Chlorambucil.....	131	bicarbonate.....	223	Colloidal bismuth subcitrate.....	8
Chloramphenicol		Cladribine.....	132	Colofac.....	7
Infections.....	80	Clarithromycin.....	77	Colony-Stimulating Factors.....	34
Sensory.....	212	Clexane.....	31	Coloxyl.....	12
Chlorhexidine.....	221	Clexane Forte.....	31	Colpocin-T.....	86
Chlorhexidine gluconate		Clindamycin.....	80	Compound electrolytes.....	35, 37
Alimentary.....	19	Clinect.....	52	Compound electrolytes with glucose	
Extemporaneously Compounded		Clinicians Multivit & Mineral		[Dextrose].....	35, 37
Preparations.....	227	Boost.....	20	Compound hydroxybenzoate.....	227
Genito-Urinary.....	60	Clinicians Renal Vit.....	20	Compound sodium lactate	
Chlorhexidine with		Clobazam.....	115	[Hartmann's solution].....	35
cetrimide.....	221, 224	Clobetasol propionate.....	57, 59	Concerta.....	126
Chlorhexidine with ethanol.....	221	Clobetasone butyrate.....	57	Condyline.....	59
Chloroform.....	227	Clofazimine.....	84	Contraceptives.....	60
Chloroquine phosphate.....	86	Clomazol		Contrast Media.....	222
Chlorothiazide.....	46	Dermatological.....	54	Copaxone.....	124
Chlorpheniramine maleate.....	205	Genito-Urinary.....	60	Corticosteroids	
Chlorpromazine hydrochloride.....	120	Clomifene citrate.....	67	Dermatological.....	57
Chlortalidone [Chlorthalidone].....	46	Clomipramine hydrochloride.....	113	Hormone Preparations.....	65
Chlorthalidone.....	46	Clonazepam.....	115, 123	Corticotorelin (ovine).....	68
Choice Load 375.....	60	Clonidine.....	44	Cosentyx.....	192
Choice TT380 Short.....	60	Clonidine BNM.....	45	Cosmegen.....	131
Choice TT380 Standard.....	60	Clonidine hydrochloride.....	45	Cough Suppressants.....	208
Cholestyramine.....	47	Clopidogrel.....	32	Creon 10000.....	11
Choline salicylate with cetalkonium		Clopidogrel Multichem.....	32	Creon 25000.....	11
chloride.....	19	Clopine.....	120	Creon Micro.....	11
Chorogonadotropin alfa.....	68	Clopixol.....	121, 123	Crotamiton.....	55
Ciclopirox olamine.....	54	Clostridium botulinum type A		Crystaderm.....	54
Ciclosporin.....	150	toxin.....	103	CT Plus+.....	222
Cidofovir.....	91	Clotrimazole		Cubicin.....	80
Cilazapril.....	39	Dermatological.....	54	Curam.....	78
Cilazapril with		Genito-Urinary.....	60	Curosurf.....	211
hydrochlorothiazide.....	39	Clove oil.....	227	Cvite.....	22
Cilicaine.....	78	Clozapine.....	120	Cyclizine hydrochloride.....	119
Cilicaine VK.....	78	Clozaril.....	120	Cyclizine lactate.....	119
Cimetidine.....	8	Co-trimoxazole.....	81	Cyclogyl.....	217
Cinacalcet.....	64	Coal tar.....	227	Cyclopentolate hydrochloride.....	217
Cinchocaine hydrochloride with		Coal tar with salicylic acid and		Cyclophosphamide.....	131

Cycloserine.....	84	Deferasirox.....	220	Diazepam.....	115, 123
Cymevene.....	91	Deferiprone.....	220	Diazoxide.....	
Cyproheptadine hydrochloride.....	205	Defibrotide.....	31	Alimentary.....	9
Cyproterone acetate.....	64	Definity.....	223	Cardiovascular.....	49
Cyproterone acetate with ethinyloestradiol.....	60	Demeclocycline hydrochloride.....	80	Dichlorobenzyl alcohol with amylmetacresol.....	20
Cystadane.....	14	Denosumab.....	99	Diclofenac Sandoz.....	104
Cysteamine hydrochloride.....	227	Deolate.....	84	Diclofenac sodium.....	
Cytarabine.....	132	Deoxycoformycin.....	137	Musculoskeletal.....	104
Cytotec.....	7	Depo-Medrol.....	66	Sensory.....	214
<b>- D -</b>					
D-Penamine.....	97	Depo-Provera.....	61	Dicobalt edetate.....	220
Dabigatran.....	30	Depo-Testosterone.....	64	Diffucan.....	82
Dacarbazine.....	134	Deprim.....	81	Diflucortolone valerate.....	57
Dactinomycin [Actinomycin D].....	131	Dermol.....	57, 59	Digestives Including Enzymes.....	11
Daivobet.....	58	Desferrioxamine mesilate.....	220	Digoxin.....	41
Daivonex.....	58	Desflurane.....	107	Digoxin immune Fab.....	219
Dalacin C.....	80	Desmopressin acetate.....	73	Dihydrocodeine tartrate.....	111
Danaparoid.....	31	Desmopressin-PH&T.....	73	Dihydroergotamine mesylate.....	118
Danazol.....	67	Dexamethasone.....		Diltiazem hydrochloride.....	44
Dantrium.....	103	Hormone Preparations.....	65	Dilzem.....	44
Dantrium IV.....	103	Sensory.....	213	Dimercaprol.....	221
Dantrolene.....	103	Dexamethasone phosphate.....	65	Dimercaptosuccinic acid.....	221
Daonil.....	10	Dexamethasone Phosphate Panpharma.....	65	Dimethicone.....	54-55
Dapa-Tabs.....	46	Dexamethasone with framycetin and gramicidin.....	212	Dimethyl fumarate.....	123
Dapsone.....	84	Dexamethasone with neomycin sulphate and polymyxin B.....		Dimethyl sulfoxide.....	225
Daptomycin.....	80	sulphate.....	212	Dinoprostone.....	61
Darunavir.....	89	Dexamethasone with tobramycin.....	212	Dipentum.....	7
Darunavir Mylan.....	89	Dexamfetamine sulfate.....	126	Diphemamil metilsulfate.....	59
Dasatinib.....	140	Dexmedetomidine.....	107	Diphenoxylate hydrochloride with atropine sulphate.....	5
Daunorubicin.....	131	Dexmedetomidine-Teva.....	107	Diphtheria antitoxin.....	219
DBL Acetylcysteine.....	219	Dexmethsone.....	65	Diphtheria, tetanus and pertussis vaccine.....	247
DBL Adrenaline.....	48	Dextrazoxane.....	147	Diphtheria, tetanus, pertussis and polio vaccine.....	246
DBL Amikacin.....	74	Dextrose.....		Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine.....	246
DBL Aminophylline.....	210	Alimentary.....	9	Diprosone.....	57
DBL Bleomycin Sulfate.....	131	Blood.....	35, 37	Dipyridamole.....	32
DBL Cefotaxime.....	75	Extemporaneously Compounded Preparations.....	227	Disodium edetate.....	215
DBL Cisplatin.....	139	Dextrose with sodium citrate and citric acid [Acid Citrate Dextrose A].....	31	Disodium hydrogen phosphate with sodium dihydrogen phosphate.....	227
DBL Dacarbazine.....	134	DHC Continus.....	111	Disopyramide phosphate.....	41
DBL Desferrioxamine Mesylate for Inj BP.....	220	Diabetes.....	9	Disulfiram.....	128
DBL Docetaxel.....	146	Diacomit.....	117	Dithranol.....	227
DBL Ergometrine.....	61	Diagnostic Agents.....		Diuretics.....	45
DBL Gentamicin.....	74	Vaccines.....	256	Dobutamine.....	48
DBL Leucovorin Calcium.....	147	Various.....	223	Dobutamine-hameln.....	48
DBL Methotrexate Onco-Vial.....	133	Diagnostic and Surgical Preparations.....	214	Docetaxel.....	146
DBL Morphine Sulphate.....	112	Diamide Relief.....	5	Docusate sodium.....	
DBL Naloxone Hydrochloride.....	219	Diamox.....	216	Alimentary.....	12
DBL Octreotide.....	149	Diatrizoate meglumine with sodium amidotrizoate.....	222	Sensory.....	218
DBL Pethidine Hydrochloride.....	113	Diatrizoate sodium.....	222	Docusate sodium with sennosides.....	12
DBL Vincristine Sulfate.....	147				
De-Worm.....	85				
Decongestants.....	208				
Decongestants and Antiallergics.....	214				
Decozol.....	20				

Dolutegravir .....	90	Elaprase .....	15	Erbitux .....	170
Domperidone .....	119	Elecare (Unflavoured).....	239	Ergometrine maleate .....	61
Donepezil hydrochloride.....	127	Elecare (Vanilla).....	239	Erlotinib .....	140
Donepezil-Rex .....	127	Elecare LCP (Unflavoured).....	239	Ertapenem .....	74
Dopamine hydrochloride.....	48	Electral .....	37	Erythrocin IV .....	77
Dornase alfa .....	210	Electrolytes .....	226	Erythromycin (as	
Dortimopt .....	216	Elelyso .....	17	ethylsuccinate).....	77
Dorzolamide .....	216	Elidel .....	58	Erythromycin (as lactobionate).....	77
Dorzolamide with timolol.....	216	Elocon.....	57	Erythromycin (as stearate).....	77
Dostinex .....	67	Elocon Alcohol Free .....	57	Esbriet .....	207
Dosulepin [Dothiepin]		Eltrombopag.....	26	Escitalopram .....	114
hydrochloride .....	114	Emend Tri-Pack .....	119	Escitalopram-Apotex .....	114
Dosulepin Mylan .....	114	Emicizumab .....	27	Esmolol hydrochloride .....	42
Dotarem .....	223	EMLA .....	110	Essential Prednisolone .....	7
Dothiepin .....	114	Emtricitabine.....	88	Estradot .....	66
Doxapram .....	211	Emtricitabine with tenofovir		Etanercept .....	151
Doxazosin .....	40	disoproxil .....	92	Ethambutol hydrochloride.....	84
Doxepin hydrochloride.....	114	Emtriva .....	88	Ethanol .....	219
Doxine .....	80	Emulsifying ointment .....	56	Ethanol with glucose.....	219
Doxorubicin Ebewe.....	131	Emulsifying Ointment ADE .....	56	Ethanol, dehydrated .....	219
Doxorubicin hydrochloride.....	131	Enalapril maleate.....	39	Ethics Aspirin .....	110
Doxycycline .....	80	Enbrel .....	151	Ethics Aspirin EC.....	32
DP Lotn HC .....	57	Endocrine Therapy .....	147	Ethics Lisinopril.....	39
DP-Allopurinol.....	102	Endoxan .....	131	Ethinylloestradiol .....	67
Dr Reddy's Omeprazole .....	8	Engerix-B.....	251	Ethinylloestradiol with	
Droletan .....	119	Enlifax XR.....	114	desogestrel .....	60
Droperidol .....	119	Enoxaparin sodium.....	31	Ethinylloestradiol with	
Drugs Affecting Bone		Enstilar .....	58	levonorgestrel .....	60
Metabolism .....	97	Ensure (Chocolate).....	245	Ethinylloestradiol with	
Dual blood glucose and blood ketone		Ensure (Vanilla).....	245	norethisterone.....	60
diagnostic test meter .....	257	Ensure Plus (Banana).....	245	Ethosuximide .....	115
Duolin .....	205	Ensure Plus (Chocolate).....	245	Ethyl chloride .....	109
DuoResp Spiromax .....	209	Ensure Plus (Fruit of the		Etomidate .....	107
Duovisc .....	215	Forest).....	245	Etopophos .....	134
Duride .....	48	Ensure Plus (Vanilla).....	245	Etoposide.....	134
Dynastat .....	104	Ensure Plus HN .....	244	Etoposide (as phosphate).....	134
Dysport .....	103	Ensure Plus HN RTH.....	244	Etoricoxib .....	104
- E -		Entacapone .....	107	Etravirine .....	87
e-chamber La Grande .....	257	Entapone .....	107	Everet .....	116
e-chamber Mask .....	257	Entecavir.....	90	Everolimus.....	202
e-chamber Turbo .....	257	Entecavir Sandoz .....	90	Evista .....	100
E-Mycin.....	77	Entresto 24/26 .....	40	Exemestane.....	150
E-Z-Cat Dry .....	222	Entresto 49/51 .....	40	Exjade.....	220
E-Z-Gas II .....	222	Entresto 97/103 .....	40	Extemporaneously Compounded	
E-Z-Paste .....	222	Enzymes .....	101	Preparations .....	227
Econazole nitrate .....	54	Ephedrine .....	48	Eylea .....	168
Edrophonium chloride.....	97	Epilim IV .....	117	Ezetimibe.....	47
Efavirenz .....	87	Epirubicin Ebewe.....	131	Ezetimibe Sandoz.....	47
Efavirenz with emtricitabine and		Epirubicin hydrochloride .....	131	Ezetimibe with simvastatin .....	47
tenofovir disoproxil.....	88	Eplerenone .....	45	- F -	
Effient .....	32	Poetin alfa .....	24	Factor eight inhibitor bypassing	
Eformoterol fumarate.....	209	Epoetin beta .....	25	fraction .....	29
Eformoterol fumarate dihydrate .....	209	Epoprostenol .....	52	Famotidine.....	8
Eftrenonacog alfa [Recombinant		Eptacog alfa [Recombinant factor		Faslodex .....	148
factor IX] .....	28	Vlla] .....	29	Febuxostat.....	102
Efudix.....	59	Eptifibatide .....	32	FEIBA NF .....	29

Felo 10 ER.....	44	cinchocaine.....	7	Gadoxetate disodium.....	223
Felo 5 ER.....	44	Fluorescein sodium.....	214	Galsulfase.....	15
Felodipine.....	44	Fluorescein sodium with lignocaine hydrochloride.....	214	Galvumet.....	11
Fentanyl.....	112	Fluorescein.....	214	Galvus.....	11
Fentanyl Sandoz.....	112	Fluorometholone.....	214	Ganciclovir.....	91
Ferinject.....	18	Fluorouracil.....	132	Gardasil 9.....	251
Ferodan.....	18	Fluorouracil Ebewe.....	132	Gastrodenol.....	8
Ferric carboxymaltose.....	18	Fluorouracil sodium.....	59	Gastrografin.....	222
Ferric subsulfate.....	28	Fluox.....	115	Gazyva.....	178
Ferriprox.....	220	Fluoxetine hydrochloride.....	115	Gefitinib.....	141
Ferro-F-Tabs.....	18	Flupenthixol decanoate.....	121	Gelatine, succinylated.....	38
Ferro-tab.....	18	Flutamide.....	148	Gelofusine.....	38
Ferrogad.....	18	Flutamin.....	148	Gemcitabine.....	133
Ferrosig.....	18	Fluticasone.....	209	Gemcitabine Ebewe.....	133
Ferrous fumarate.....	18	Fluticasone furoate with vilanterol.....	210	Gemfibrozil.....	46
Ferrous fumarate with folic acid.....	18	Fluticasone propionate.....	205	Genoptic.....	212
Ferrous gluconate with ascorbic acid.....	18	Fluticasone with salmeterol.....	210	Gentamicin sulphate	
Ferrous sulfate.....	18	FML.....	214	Infections.....	74
Ferrous sulphate.....	18	Foban.....	54	Sensory.....	212
Ferrous sulphate with ascorbic acid.....	18	Folic acid.....	25	Gestrinone.....	67
Fexofenadine hydrochloride.....	205	Fondaparinux sodium.....	31	Gilenya.....	123
Filgrastim.....	34	Food Modules.....	230	Ginet.....	60
Finasteride.....	62	Food/Fluid Thickeners.....	232	Glatiramer acetate.....	124
Fingolimod.....	123	Forteo.....	101	Glaucoma Preparations.....	216
Firazyr.....	204	Fortisip (Vanilla).....	245	Glecaprevir with pibrentasvir.....	91
Flagyl.....	86	Fosamax.....	97	Glibenclamide.....	10
Flagyl-S.....	86	Fosamax Plus.....	97	Gliclazide.....	10
Flamazine.....	54	Foscarnet sodium.....	91	Gliolan.....	150
Flecainide acetate.....	41	Fosfomycin.....	80	Glipizide.....	10
Flecainide BNM.....	41	Framycetin sulphate.....	212	Glivec.....	141
Flecainide Controlled Release		Fresenius Kabi		Glizide.....	10
Teva.....	41	Blood.....	35, 37	Glucagen Hypokit.....	9
Fleet Phosphate Enema.....	13	Various.....	225	Glucagon hydrochloride.....	9
Flixonase Hayfever & Allergy.....	205	Fresofol 1% MCT/LCT.....	108	Glucerna Select (Vanilla).....	236
Flixotide.....	209	Frusemide.....	45	Glucerna Select RTH (Vanilla).....	236
Flixotide Accuhaler.....	209	Frusemide-Claris.....	45	Glucobay.....	9
Florinef.....	65	Fucidin.....	81	Glucose [Dextrose]	
Fluanxol.....	121	Fucithalmic.....	212	Alimentary.....	9
Flucil.....	78	Fulvestrant.....	148	Blood.....	35
Flucloxacillin.....	78	Fungilin.....	20	Extemporaneously Compounded	
Flucloxin.....	78	Furosemide [Frusemide].....	45	Preparations.....	227
Fluconazole.....	82	Furosemide-Baxter.....	45	Glucose with potassium chloride.....	35
Fluconazole-Claris.....	82	Fusidic acid		Glucose with potassium chloride and sodium chloride.....	36
Flucytosine.....	84	Dermatological.....	54, 58	Glucose with sodium chloride.....	36
Fludara Oral.....	132	Infections.....	81	Glucose with sucrose and fructose.....	9
Fludarabine Ebewe.....	132	Sensory.....	212	Glycerin with sodium saccharin.....	228
Fludarabine phosphate.....	132			Glycerin with sucrose.....	228
Fludrocortisone acetate.....	65			Glycerol	
Fluids and Electrolytes.....	35			Alimentary.....	13
Flumazenil.....	219			Extemporaneously Compounded	
Flumetasone pivalate with clioquinol.....	213			Preparations.....	228
Fluocortolone caproate with fluocortolone pivalate and				Glycerol with paraffin.....	56
				Glyceryl trinitrate	
				Alimentary.....	7

**- G -**

Cardiovascular.....	48	[HPV].....	251	Imatinib-AFT.....	142
Glycine.....	224	Humatin.....	74	Imatinib-Rex.....	142
Glycopyrronium.....	206	Humira.....	158	Imigran.....	118
Glycopyrronium bromide.....	7	HumiraPen.....	158	Imipenem with cilastatin.....	74
Glycopyrronium with		Hyaluronic acid		Imipenem-Cilastatin RBX.....	74
indacaterol.....	206	Alimentary.....	20	Imipramine hydrochloride.....	114
Glypressin.....	73	Sensory.....	215, 218	Imiquimod.....	59
Gonadorelin.....	68	Hyaluronic acid with lidocaine		Immune Modulators.....	94
Goserelin.....	68	[lignocaine].....	20	Immunosuppressants.....	150
Granisetron.....	119	Hyaluronidase.....	101	Impact Advanced Recovery.....	243
<b>- H -</b>		Hydralazine hydrochloride.....	49	Imuran.....	202
Habitrol.....	129	Hydrea.....	135	Incruse Ellipta.....	206
Habitrol (Fruit).....	129	Hydrocortisone		Indacaterol.....	209
Habitrol (Mint).....	129	Dermatological.....	57	Indapamide.....	46
Haem arginate.....	15	Extemporaneously Compounded		Indigo carmine.....	224
Haemophilus influenzae type B		Preparations.....	228	Indinavir.....	89
vaccine.....	247	Hormone Preparations.....	66	Indocyanine green.....	224
Haldol.....	121	Hydrocortisone (PSM).....	57	Indomethacin.....	104
Haldol Concentrate.....	121	Hydrocortisone acetate.....	6	Infanrix IPV.....	246
Haloperidol.....	120	Hydrocortisone acetate with		Infanrix-hexa.....	246
Haloperidol decanoate.....	121	pramoxine hydrochloride.....	6	Infatrin.....	241
Hartmann's solution.....	35	Hydrocortisone and paraffin liquid		Infliximab.....	170
Harvoni.....	91	and lanolin.....	57	Influenza vaccine.....	252
Havrix.....	250	Hydrocortisone butyrate.....	57, 59	Influvac Tetra	
Havrix Junior.....	250	Hydrocortisone with miconazole.....	58	(2020 formulation).....	252
Healon.....	215	Hydrocortisone with natamycin and		Inhaled Corticosteroids.....	208
Healon 5.....	215	neomycin.....	58	Inspra.....	45
Healon GV.....	215	Hydrogen peroxide.....	54	Instillagel Lido.....	109
healthE Calamine Aqueous Cream		Hydroxocobalamin		Insulin aspart.....	10
BP.....	55	Alimentary.....	21	Insulin aspart with insulin aspart	
healthE Dimethicone 10%.....	55	Various.....	219	protamine.....	9
healthE Dimethicone 4% Lotion.....	54	hydroxycarbamide.....	135	Insulin glargine.....	10
healthE Dimethicone 5%.....	55	Hydroxychloroquine.....	97	Insulin glulisine.....	10
healthE Fatty Cream.....	56	Hydroxyurea		Insulin isophane.....	9
healthE Glycerol BP Liquid.....	228	[hydroxycarbamide].....	135	Insulin lispro.....	10
healthE Urea Cream.....	57	Hygroton.....	46	Insulin lispro with insulin lispro	
Hemlibra.....	27	Hylo-Fresh.....	218	protamine.....	10
Heparin sodium.....	31	Hyoscine butylbromide.....	7	Insulin neutral.....	10
Heparinised saline.....	31	Hyoscine hydrobromide.....	119	Insulin neutral with insulin	
Heparin Junior.....	237	Hyperuricaemia and Antigout.....	102	isophane.....	10
Hepatitis A vaccine.....	250	Hypromellose.....	215, 217	Integrilin.....	32
Hepatitis B recombinant		Hypromellose with dextran.....	217	Intelence.....	87
vaccine.....	251	<b>- I -</b>		Interferon alfa-2b.....	94
Hepsera.....	90	Ibiamox.....	78	Interferon beta-1-alpha.....	124
Herceptin.....	197	Ibrance.....	143	Interferon beta-1-beta.....	124
Hiberix.....	247	Ibuprofen.....	104	Interferon gamma.....	94
Hiprex.....	81	Ibuprofen SR BNM.....	104	Intra-uterine device.....	60
Histaclear.....	205	Icatibant.....	204	Invanz.....	74
Histamine acid phosphate.....	224	Idarubicin hydrochloride.....	132	Invega Sustenna.....	122
Holoxan.....	131	Idarucizumab.....	28	Iodine.....	72
Hormone Replacement Therapy.....	66	Idursulfase.....	15	Iodine with ethanol.....	221
HPV.....	251	Ifosfamide.....	131	Iodised oil.....	222
Humalog Mix 25.....	10	Ikorel.....	49	Iodixanol.....	222
Humalog Mix 50.....	10	Iloprost.....	52	Iohexol.....	222
Human papillomavirus (6, 11, 16, 18,		Imaging Agents.....	150	Iopidine.....	217
31, 33, 45, 52 and 58) vaccine		Imatinib mesilate.....	141	Ioscan.....	222

IPOL.....	254	KetoSens.....	257	Levosimendan.....	48
Ipratropium bromide.....	205	Ketostix.....	257	Levothyroxine.....	72
Iressa.....	141	Keytruda.....	200	Lidocaine [Lignocaine].....	109
Irinotecan Actavis 100.....	135	Kivexa.....	88	Lidocaine [Lignocaine] hydrochloride.....	109
Irinotecan hydrochloride.....	135	Klacid.....	77	Lidocaine [Lignocaine] hydrochloride with adrenaline.....	109
Iron polymaltose.....	18	Klean Prep.....	12	Lidocaine [Lignocaine] hydrochloride with adrenaline and tetracaine hydrochloride.....	109
Iron sucrose.....	18	Kogenate FS.....	30	Lidocaine [Lignocaine] hydrochloride with chlorhexidine.....	110
Irrigation Solutions.....	224	Konakion MM.....	30	Lidocaine [Lignocaine] hydrochloride with phenylephrine hydrochloride.....	110
Isentress.....	90	Konsyl-D.....	12	Lidocaine [Lignocaine] with prilocaine.....	110
Isentress HD.....	90	Kuvan.....	16	Lidocaine-Claris.....	109
Ismo 20.....	48	<b>- L -</b>			lignocaine
Ismo 40 Retard.....	48	L-ornithine L-aspartate.....	9	Alimentary.....	20
Isoflurane.....	107	Labeltalol.....	43	Nervous.....	109-110
Isoniazid.....	84	Lacosamide.....	116	Lincomycin.....	81
Isoniazid with rifampicin.....	84	Lactose.....	228	Linezolid.....	81
Isoprenaline [Soprotrenol].....	48	Lactulose.....	13	Linezolid Kabi.....	81
Isopropyl alcohol.....	221	Laevolac.....	13	Lioresal Intrathecal.....	102
Isoproterenol.....	48	Lamictal.....	116	Liothyronine sodium.....	72
Isoptin.....	44	Lamivudine.....	88, 90	Lipazil.....	46
Isoptin SR.....	44	Lamivudine Alphapharm.....	88	Lipid-Modifying Agents.....	46
Isopto Carpine.....	216	Lamotrigine.....	116	Lipiodol Ultra Fluid.....	222
Isosorbide mononitrate.....	48	Lanoxin.....	41	Liquibar.....	222
Isotretinoin.....	55	Lanoxin PG.....	41	Lisinopril.....	39
Ispaghula (psyllium) husk.....	12	Lansoprazole.....	8	Lissamine green.....	214
Isradipine.....	44	Lantus.....	10	Lithium carbonate.....	121
Itch-Soothe.....	55	Lantus SoloStar.....	10	LMX4.....	109
Itraconazole.....	82	Lanzol Relief.....	8	Local Preparations for Anal and Rectal Disorders.....	7
Itrazole.....	82	Lapatinib.....	142	Locoid.....	57, 59
Ivabradine.....	41	Largactil.....	120	Locoid Crelo.....	57
Ivermectin.....	85	Laronidase.....	16	Locoid Lipocream.....	57
<b>- J -</b>		Lasix.....	45	Lodoxamide.....	214
Jadelle.....	61	Latanoprost.....	216	Logem.....	116
Jakavi.....	144	Lax-Suppositories.....	13	Lomide.....	214
Jaydess.....	61	Lax-Tabs.....	13	Lomustine.....	131
Jevity HiCal RTH.....	244	Laxatives.....	12	Long-Acting Beta-Adrenoceptor Agonists.....	209
Jevity RTH.....	244	Laxsol.....	12	Loniten.....	49
Juno Pemetrexed.....	133	Ledipasvir with sofosbuvir.....	91	Loperamide hydrochloride.....	5
<b>- K -</b>		Leflunomide.....	97	Lopinavir with ritonavir.....	89
Kadcyla.....	198	Lenalidomide.....	135	Lorafix.....	205
Kaletra.....	89	Letrole.....	150	Loratadine.....	205
Kenacomb.....	213	Letrozole.....	150	Lorazepam.....	115, 123
Kenacort-A 10.....	66	Leukotriene Receptor Antagonists.....	209	Lorfast.....	205
Kenacort-A 40.....	66	Leuporelin acetate.....	68	Lormetazepam.....	124
Kenalog in Orabase.....	20	Leustatin.....	132	Lorstat.....	46
Ketalar.....	107	Levetiracetam.....	116	Losartan Actavis.....	40
Ketamine.....	107	Levetiracetam-AFT.....	116	Losartan potassium.....	40
Ketamine-Baxter.....	107	Levlen ED.....	60		
Ketamine-Claris.....	107	Levocabastine.....	214		
Ketocal 3:1 (Unflavoured).....	242	Levocarnitine.....	16		
Ketocal 4:1 (Unflavoured).....	242	Levodopa with benserazide.....	107		
Ketocal 4:1 (Vanilla).....	242	Levodopa with carbidopa.....	107		
Ketoconazole.....		Levomepromazine.....	121		
Dermatological.....	54	Levodopa hydrochloride.....	121		
Infections.....	82	Levonorgestrel.....	61		
Ketoprofen.....	104				
Ketorolac trometamol.....	214				

Losartan potassium with hydrochlorothiazide .....	40	Marcaïn Heavy.....	109	Methotrexate.....	133
Lovir.....	91	Marcaïn Isobaric .....	108	Methotrexate DBL Onco-Vial.....	133
Loxamine.....	115	Marcaïn with Adrenaline .....	108	Methotrexate Ebewe.....	133
Lucrin Depot 1-month.....	68	Marevan.....	32	Methotrexate Sandoz .....	133
Lucrin Depot 3-month.....	68	Marine Blue Lotion SPF 50+ .....	59	Methoxsalen	
Lyderm.....	55	Mask for spacer device.....	257	[8-methoxy psoralen].....	58
Lynparza.....	136	Mast Cell Stabilisers .....	210	Methoxyflurane .....	110
Lysine acetylsalicylate [Lysine aspirin].....	32	Maviret.....	91	Methyl aminolevulinate hydrochloride.....	59
Lysine aspirin .....	32	Maxidex.....	213	Methyl hydroxybenzoate.....	228
<b>- M -</b>		Maxitrol.....	212	Methylcellulose .....	228
m-Amoxiclav.....	78	Measles, mumps and rubella vaccine.....	253	Methylcellulose with glycerin and sodium saccharin.....	228
m-Eslon.....	112	Mebendazole.....	85	Methylcellulose with glycerin and sucrose.....	228
Mabthera.....	181	Mebeverine hydrochloride .....	7	Methyldopa.....	45
Macrogl 3350 with ascorbic acid, potassium chloride and sodium chloride.....	12	Medrol.....	66	Methyldopa Mylan .....	45
Macrogl 3350 with potassium chloride, sodium bicarbonate and sodium chloride .....	13	Medroxyprogesterone.....	67	Methylene blue .....	224
Macrogl 3350 with potassium chloride, sodium bicarbonate, sodium chloride and sodium sulphate.....	12	Medroxyprogesterone acetate		Methylnaltrexone bromide .....	12
Macrogl 400 and propylene glycol.....	217	Genito-Urinary.....	61	Methylphenidate ER - Teva.....	126
Madopar 125 .....	107	Hormone Preparations.....	67	Methylphenidate hydrochloride.....	126
Madopar 250 .....	107	Mefenamic acid .....	104	Methylprednisolone (as sodium succinate) .....	66
Madopar 62.5 .....	107	Mefloquine.....	86	Methylprednisolone aceponate.....	57
Madopar HBS.....	107	Megestrol acetate.....	149	Methylprednisolone acetate.....	66
Madopar Rapid.....	107	Meglumine gadopentetate .....	223	Methylthioninium chloride [Methylene blue].....	224
Mafenide acetate.....	54	Meglumine iotroxate .....	223	Methylxanthines.....	210
Magnesium amino acid chelate .....	19	Melatonin.....	124	Metoclopramide Actavis 10 .....	119
Magnesium chloride .....	19	Melphalan.....	131	Metoclopramide hydrochloride .....	119
Magnesium hydroxide		Menactra.....	247	Metoclopramide hydrochloride with paracetamol.....	118
Alimentary.....	19	Meningococcal (A, C, Y and W-135) conjugate vaccine.....	247	Metolazone.....	46
Extemporaneously Compounded Preparations .....	228	Meningococcal C conjugate vaccine .....	248	Metoprolol succinate.....	43
Magnesium oxide .....	19	Menthol.....	228	Metoprolol tartrate .....	43
Magnesium oxide with magnesium aspartate, magnesium amino acid chelate and magnesium citrate.....	19	Mepivacaine hydrochloride.....	110	Metrogyl.....	86
Magnesium sulphate .....	19	Mepolizumab .....	177	Metronidazole	
Magnevist.....	223	Mercaptopurine.....	133	Dermatological.....	54
Malarone.....	86	Meropenem.....	75	Infections.....	86
Malarone Junior.....	86	Meropenem Ranbaxy .....	75	Metoprolol IV Mylan.....	43
Malathion [Maldison].....	55	Meropenem-AFT .....	75	Metyrapone.....	67
Maldison .....	55	Mesalazine .....	6	Mexiletine hydrochloride.....	42
Mannitol		Mesna.....	147	Mexiletine Hydrochloride USP.....	42
Cardiovascular.....	45	Mestinon.....	97	Miacalcic.....	64
Various.....	224	Metabolic Disorder Agents .....	13	Mianserin hydrochloride .....	114
Mantoux.....	256	Metabolic Products .....	232	Micolette.....	13
Maprotiline hydrochloride .....	114	Metaraminol.....	49	Miconazole .....	20
Marcaïn.....	108	Meterol.....	209	Miconazole nitrate	
		Metformin hydrochloride .....	11	Dermatological.....	54
		Methacholine chloride .....	224	Genito-Urinary.....	60
		Methadone hydrochloride		Micreme.....	60
		Extemporaneously Compounded Preparations .....	228	Micreme H .....	58
		Nervous.....	112	Microgynon 20 ED.....	60
		Methatabs.....	112	Microgynon 50 ED.....	60
		Methenamine (Hexamine) hippurate.....	81	Microlut.....	61
		Methohexital sodium.....	107	Midazolam .....	125
		Methopt.....	217		

Midodrine.....	42	Mycosail.....	54	Genito-Urinary.....	60
Mifepristone.....	61	Mycophenolate mofetil.....	202	Infections.....	82
Milrinone.....	49	Mydriacyl.....	217	Nimodipine.....	44
Minerals.....	18	Mydriatics and Cycloplegics.....	217	Nimotop.....	44
Mini-Wright AFS Low Range.....	257	Mylan Atenolol.....	42	Nintedanib.....	206
Mini-Wright Standard.....	257	Mylan Clomiphen.....	67	Nitazoxanide.....	86
Minidiab.....	10	Mylan Midazolam.....	125	Nitrates.....	48
Minims Prednisolone.....	214	Myleran.....	131	Nitroderm TTS 10.....	48
Minirin.....	73	Myozyme.....	13	Nitroderm TTS 5.....	48
Minocycline.....	80	- N -		Nitrofurantoin.....	81
Minoxidil.....	49	Nadolol.....	43	Nitrolingual Pump Spray.....	48
Mirena.....	61	Naglazyme.....	15	Nivestim.....	34
Mirtazapine.....	114	Naloxone hydrochloride.....	219	Nivolumab.....	199
Misoprostol.....	7	Naltracord.....	128	Nodia.....	5
Mitomycin C.....	132	Naltrexone hydrochloride.....	128	Noflam 250.....	104
Mitozantrone.....	132	Naphazoline hydrochloride.....	214	Noflam 500.....	104
Mitozantrone Ebewe.....	132	Naphcon Forte.....	214	Non-Steroidal Anti-Inflammatory Drugs.....	103
Mivacron.....	103	Naprosyn SR 1000.....	104	Nonacog gamma, [Recombinant factor IX].....	29
Mivacurium chloride.....	103	Naprosyn SR 750.....	104	Noradrenaline.....	49
Mixed salt solution for eye irrigation.....	215	Naproxen.....	104	Noradrenaline BNM.....	49
Moclobemide.....	114	Naropin.....	110	Norethisterone Genito-Urinary.....	61
Modafinil.....	127	Natalizumab.....	124	Hormone Preparations.....	68
Modavigil.....	127	Natamycin.....	212	Norethisterone with mestranol.....	60
Molaxole.....	13	Natulan.....	137	Norflex.....	103
Mometasone furoate.....	57	Nausafix.....	120	Norflexacin.....	79
Monosodium glutamate with sodium aspartate.....	226	Nausicalm.....	119	Noriday 28.....	61
Monosodium l-aspartate.....	226	Navelbine.....	147	Normison.....	125
Montelukast.....	209	Nedocromil.....	210	Norpress.....	114
Montelukast Mylan.....	209	Nefopam hydrochloride.....	111	Nortriptyline hydrochloride.....	114
Morotcog alfa [Recombinant factor VIII].....	29	Neisvac-C.....	248	Norvir.....	89
Morphine hydrochloride.....	112	Neo-B12.....	21	Novasource Renal (Vanilla).....	243
Morphine sulphate.....	112	Neocate Gold (Unflavoured).....	239	Novatrein.....	58
Morphine tartrate.....	112	Neocate Junior Vanilla.....	239	NovoMix 30 FlexPen.....	9
Motetis.....	106	Neoral.....	150	NovoRapid FlexPen.....	10
Mouth and Throat.....	19	Neostigmine metilsulfate.....	97	NovoSeven RT.....	29
Movapo.....	106	Neostigmine metilsulfate with glycopyrronium bromide.....	97	Noxafil.....	82
Moxifloxacin.....	79	Neosynephrine HCL.....	49	Nozinan.....	121
Moxifloxacin Kabi.....	79	Nepro HP (Strawberry).....	243	Nucala.....	177
Mozobil.....	34	Nepro HP (Vanilla).....	243	Nuelin.....	210
Mucolytics and Expectorants.....	210	Nepro HP RTH.....	243	Nuelin-SR.....	210
Mucosoothe.....	109	Neulastim.....	34	Nutrini Energy Multi Fibre.....	242
Multihance.....	223	Neupogen.....	34	Nutrini Low Energy Multifibre RTH.....	242
Multiple Sclerosis Treatments.....	123	NeuroTabs.....	18	Nutrini Peptisorb.....	239
Multivitamin and mineral supplement.....	20	Nevirapine.....	87	Nutrini Peptisorb Energy.....	239
Multivitamin renal.....	20	Nevirapine Alphapharm.....	87	Nutrison 800 Complete Multi Fibre.....	244
Multivitamins.....	21	Nicardipine hydrochloride.....	44	Nutrison Concentrated.....	238
Mupirocin.....	54	Nicorandil.....	49	Nutrison Energy.....	244
Muscle Relaxants and Related Agents.....	102	Nicotine.....	129	Nutrison Protein Intense.....	238
Mvite.....	21	Nicotinic acid.....	47	Nyefax Retard.....	44
Myambutol.....	84	Nifedipine.....	44		
Mycobutin.....	85	Nifuran.....	81		
		Nilotinib.....	142		
		Nilstat.....			
		Alimentary.....	20		

Nystatin		Opdivo	199	Paracare Double Strength	111
Alimentary	20	Optional Pharmaceuticals	257	Paracetamol	111
Dermatological	54	Ora-Blend	228	Paracetamol Kabi	111
Genito-Urinary	60	Ora-Blend SF	228	Paracetamol with codeine	113
Infections	82	Ora-Plus	228	Paraffin	
- O -		Ora-Sweet	228	Alimentary	12
O/W Fatty Emulsion Cream	56	Ora-Sweet SF	228	Dermatological	56
Obinutuzumab	178	Oratane	55	Extemporaneously Compounded	
Obstetric Preparations	61	Ornidazole	86	Preparations	228
Ocrelizumab	124	Orphenadrine citrate	103	Paraffin liquid with soft white	
Ocrevus	124	Oruvail SR	104	paraffin	218
Octocog alfa [Recombinant factor		Oseltamivir	93	Paraffin liquid with wool fat	218
VIII] (Advate)	29	Osmolite RTH	244	Paraffin with wool fat	57
Octocog alfa [Recombinant factor		Other Cardiac Agents	48	Paraldehyde	115
VIII] (Kogenate FS)	30	Other Endocrine Agents	67	Parecoxib	104
Octreotide	149	Other Oestrogen Preparations	67	Paromomycin	74
Ocular Lubricants	217	Other Otological Preparations	218	Paroxetine	115
Oestradiol	66-67	Other Progestogen		Paser	85
Oestradiol valerate	66	Preparations	67	Patent blue V	224
Oestradiol with norethisterone		Other Skin Preparations	59	Paxam	123
acetate	66	Ovestin		Pazopanib	144
Oestriol		Genito-Urinary	62	Peak flow meter	257
Genito-Urinary	62	Hormone Preparations	67	Peanut oil	227
Hormone Preparations	67	Ox-Pam	123	Pedialyte - Bubblegum	37
Oestrogens	62	Oxaliplatin	139	Pediasure (Chocolate)	242
Oestrogens (conjugated equine)	66	Oxaliplatin Accord	139	Pediasure (Strawberry)	242
Oestrogens with		Oxandrolone	64	Pediasure (Vanilla)	242
medroxyprogesterone		Oxazepam	123	Pediasure RTH	242
acetate	67	Oxpentifylline	49	Pegaspargase	136
Ofev	206	Oxybuprocaine hydrochloride	215	Pegasys	94
Oil in water emulsion	56	Oxybutynin	63	Pegfilgrastim	34
Oily phenol [Phenol oily]	7	Oxycodone hydrochloride	113	Pegylated interferon alfa-2a	94
Olanzapine	121-122	Oxycodone Sandoz	113	Pembrolizumab	200
Olaparib	136	Oxymetazoline hydrochloride	208	Pemetrexed	133
Olive oil	228	OxyNorm	113	Penicillamine	97
Olopatadine	214	Oxytocin	61	Penicillin G	78
Olopatadine Teva	214	Oxytocin BNM	61	Penicillin V	78
Olsalazine	7	Oxytocin with ergometrine		Pentacarinat	86
Omaliuzumab	179	maleate	61	Pentagastrin	67
Omeprazole	8	Ozurdex	213	Pentamidine isethionate	86
Omeprazole actavis 10	8	- P -		Pentasa	6
Omeprazole actavis 20	8	Pacifen	102	Pentostatin [Deoxycoryformycin]	137
Omeprazole actavis 40	8	Paclitaxel	146	Pentoxifylline [Oxpentifylline]	49
Omezol IV	8	Paclitaxel Ebewe	146	Peptamen OS 1.0 (Vanilla)	237
Omnipaque	222	Palbociclib	143	Peptisoothe	8
Omniscan	223	Paliperidone	122	Perflutren	223
Omnitrope	68	Pamidronate disodium	98	Perhexiline maleate	44
Onbrez Breezhaler	209	Pamisol	98	Pericyazine	121
Oncaspar LYO	136	Pancreatic enzyme	11	Perindopril	39
OncoTICE	202	Pancuronium bromide	103	Perjeta	180
Ondansetron	119	Pantoprazole	8	Permethrin	55
Ondansetron Kabi	119	Panzop Relief	8	Perrigo	59
Ondansetron ODT-DRLA	119	Papaverine hydrochloride	49	Pertuzumab	180
Ondansetron-Clarix	119	Paper wasp venom	204	Peteha	85
One-Alpha	22	Para-aminosalicylic Acid	85	Pethidine hydrochloride	113
Onrex	119	Paracare	111	Pexsig	44

Pfizer Exemestane .....	150	Pneumococcal (PCV10) conjugate vaccine .....	248	Prevenar 13.....	248
Pheburane .....	17	Pneumococcal (PCV13) conjugate vaccine .....	248	Prezista.....	89
Phenasen .....	134	Pneumococcal (PPV23) polysaccharide vaccine .....	249	Prilocaine hydrochloride .....	110
Phenelzine sulphate .....	114	Pneumovax 23.....	249	Prilocaine hydrochloride with felypressin .....	110
Phenindione.....	31	Podophyllotoxin .....	59	Primacor .....	49
Phenobarbitone .....	116, 125	Polidocanol .....	28	Primaquine .....	86
Phenobarbitone sodium.....	228	Poliomyelitis vaccine .....	254	Primidone .....	117
Phenol		Poloxamer .....	12	Primolut N.....	68
Extemporaneously Compounded Preparations .....	228	Poly Gel .....	217	Primovist .....	223
Various.....	225	Poly-Tears .....	217	Priorix .....	253
Phenol oily .....	7	Poly-Visc.....	218	Probenecid .....	102
Phenol with ioxaglic acid .....	225	Polyhexamethylene biguanide.....	228	Procaine penicillin.....	78
Phenothrin .....	55	Polyvinyl alcohol with povidone .....	218	Procarbazine hydrochloride .....	137
Phenoxybenzamine hydrochloride .....	40	Poractant alfa .....	211	Prochlorperazine .....	120
Phenoxyethylpenicillin [Penicillin V] .....	78	Posaconazole .....	82	Proctosedyl.....	7
Phentolamine mesylate .....	40	Postinor-1 .....	61	Procyclidine hydrochloride.....	106
Phenylephrine hydrochloride		Potassium chloride .....	36, 38	Procytox.....	131
Cardiovascular .....	49	Potassium chloride with sodium chloride .....	36	Progesterone .....	61
Sensory .....	217	Potassium citrate.....	63	Proglycem.....	9
Phenytin .....	116	Potassium dihydrogen phosphate.....	36	Progynova .....	66
Phenytoin sodium .....	115-116	Potassium iodate		Prolia .....	99
Pholcodine .....	208	Alimentary .....	18	Promethazine hydrochloride.....	205
Phosphorus .....	37	Hormone Preparations.....	72	Propafenone hydrochloride .....	42
Phytomenadione.....	30	Potassium iodate with iodine .....	18	Propamidine isethionate .....	212
Picibanil .....	202	Potassium perchlorate.....	72	Propofol .....	108
Pilocarpine hydrochloride .....	216	Potassium permanganate.....	59	Propranolol .....	43
Pilocarpine nitrate.....	228	Povidone K30 .....	228	Propylthiouracil .....	73
Pimafucort .....	58	Povidone-iodine .....	221	Prostin E2 .....	61
Pimecrolimus .....	58	Povidone-iodine with ethanol.....	221	Prostin VR .....	49
Pindolol.....	43	Pradaxa .....	30	Protamine sulphate.....	31
Pine tar with trolamine laurilsulfate and fluorescein .....	59	Pralidoxime iodide .....	219	Protionamide .....	85
Pinetarsol.....	59	Pramipexole hydrochloride .....	107	Protirelin .....	73
Pioglitazone.....	11	Prasugrel .....	32	Proveblue .....	224
Piperacillin with tazobactam .....	78	Pravastatin.....	46	Provera .....	67
PiperTaz Sandoz .....	78	Pravastatin Mylan.....	46	Provera HD.....	67
Pipothiazine palmitate .....	122	Praxbind .....	28	Proxymetacaine hydrochloride .....	215
PipTaz Sandoz .....	78	Praziquantel.....	85	Pseudoephedrine hydrochloride.....	208
Pirfenidone .....	207	Prazosin.....	41	PSM Citalopram .....	114
Pituitary and Hypothalamic Hormones and Analogues.....	68	Precedex .....	107	Psoriasis and Eczema Preparations .....	58
Pivmecillinam.....	81	Pred Forte .....	214	PTU .....	73
Pizotifen.....	118	Prednisolone.....	66	Pulmonary Surfactants .....	211
PKU Anamix Junior LQ (Berry).....	234	Prednisolone acetate.....	214	Pulmozyme .....	210
PKU Anamix Junior LQ (Orange) .....	234	Prednisolone sodium .....	7	Puri-nethol .....	133
PKU Anamix Junior LQ (Unflavoured).....	234	Prednisolone sodium phosphate.....	214	Puria .....	22
Plaquenil.....	97	Prednisolone- AFT.....	214	Pyrazinamide.....	85
Plasma-Lyte 148.....	35	Prednisone .....	66	Pyridoxal-5-phosphate.....	16
Plasma-Lyte 148 & 5% Glucose.....	35	Pregabalin .....	117	Pyridoxine hydrochloride .....	21
Plendil ER.....	44	Pregabalin Pfizer .....	117	Pyrimethamine .....	86
Plerixafor .....	34	Pregnancy test - hCG urine .....	257	Pytazen SR.....	32
		preOp.....	243		

Quetapel.....	121	Riodine.....	221	Scandonest 3%.....	110
Quetiapine.....	121	Risedronate Sandoz.....	98	Sclerosing Agents.....	211
Quinapril.....	39	Risedronate sodium.....	98	Scopoderm TTS.....	119
Quinapril with hydrochlorothiazide.....	39	Risperdal Consta.....	122	Sebizole.....	54
Quinine dihydrochloride.....	87	Risperidone.....	121–122	Secretin pentahydrochloride.....	224
Quinine sulphate.....	87	Risperidone (Teva).....	121	Secukinumab.....	192
Qvar.....	208	Risperon.....	121	Sedatives and Hypnotics.....	124
<b>- R -</b>					
RA-Morph.....	112	Ritalin.....	126	Seebri Breezhaler.....	206
Rabies vaccine.....	254	Ritalin LA.....	126	Selegiline hydrochloride.....	107
Raloxifene.....	100	Ritalin SR.....	126	Sennosides.....	13
Raltegravir potassium.....	90	Ritonavir.....	89	Sensipar.....	64
Ramipex.....	107	Rituximab (mabthera).....	181	Serenace.....	120
Ranbaxy-Cefaclor.....	75	Rituximab (riximyo).....	183	Seretide.....	210
Ranibizumab.....	180	Rivaroxaban.....	32	Seretide Accuhaler.....	210
Ranitidine.....	8	Rivastigmine.....	127	Serevent.....	209
Rapamune.....	202	Rivotril.....	115	Serevent Accuhaler.....	209
Rasburicase.....	102	Riximyo.....	183	Sertraline.....	115
Readi-CAT 2.....	222	RIXUBIS.....	29	Setrona.....	115
Reandron 1000.....	64	Rizamelt.....	118	Sevoflurane.....	108
Recombinant factor IX.....	28–29	Rizatriptan.....	118	Sevredol.....	112
Recombinant factor VIIa.....	29	Rocuronium bromide.....	103	Shingles vaccine.....	255
Recombinant factor VIII.....	29–30	Rolin.....	150	Sildenafil.....	51
Rectogesic.....	7	Ropin.....	107	Siltuximab.....	193
Red back spider antivenom.....	220	Ropinirole hydrochloride.....	107	Silver nitrate	
Redipred.....	66	Ropivacaine hydrochloride.....	110	Dermatological.....	59
Relenza Rotadisk.....	94	Ropivacaine hydrochloride with fentanyl.....	110	Extemporaneously Compounded Preparations.....	228
Relistor.....	12	Ropivacaine Kabi.....	110	Simeticone.....	5
Remicade.....	170	Rose bengal sodium.....	214	Simulect.....	169
Remifentanil.....	113	Rotarix.....	254	Simvastatin.....	46
Remifentanil-AFT.....	113	Rotavirus oral vaccine.....	254	Simvastatin Mylan.....	46
ReoPro.....	158	Roxane.....	5	Sinacalide.....	224
Resonium A.....	38	Roxithromycin.....	77	Sinemet.....	107
Resource Beneprotein.....	231	Rubifen.....	126	Sinemet CR.....	107
Resource Diabetic (Vanilla).....	236	Rubifen SR.....	126	Sirolimus.....	202
Respiratory Stimulants.....	211	Rulide D.....	77	Siterone.....	64
Retinol.....	21	Rurioctocog alfa pegol [Recombinant factor VIII].....	30	Slow-Lopresor.....	43
Retinol Palmitate.....	218	Ruxolitinib.....	144	Smith BioMed Rapid Pregnancy Test.....	257
ReTrieve.....	55	<b>- S -</b>			
Retrovir.....	88	S26 LBW Gold RTF.....	241	Snake antivenom.....	220
Retrovir IV.....	88	Sacubitril with valsartan.....	40	Sodibic.....	38
Revlimid.....	135	SalAir.....	208	Sodium acetate.....	36
Revolade.....	26	Salazopyrin.....	7	Sodium acid phosphate.....	37
Rexacrom.....	214	Salazopyrin EN.....	7	Sodium alginate with magnesium alginate.....	5
Riboflavin 5-phosphate.....	216	Salbutamol.....	208	Sodium alginate with sodium bicarbonate and calcium carbonate.....	5
Ribomustin.....	130	Salbutamol with ipratropium bromide.....	205	Sodium aurothiomalate.....	97
Ricit.....	62	Salicylic acid.....	228	Sodium benzoate.....	17
Rifabutin.....	85	Salmeterol.....	209	Sodium bicarbonate	
Rifadin.....	85	Salmonella typhi vaccine.....	250	Blood.....	36, 38
Rifampicin.....	85	Sandimmun.....	150	Extemporaneously Compounded Preparations.....	228
Rifaximin.....	9	Sandomigran.....	118	Sodium calcium edetate.....	221
Rifinah.....	84	Sandostatin LAR.....	149		
Rilutek.....	106	Sapropterin Dihydrochloride.....	16		
Riluzole.....	106	Scalp Preparations.....	59		
Ringer's solution.....	36				

Sodium chloride		Specialised Formulas	236	Taliglucerase alfa	17
Blood	37–38	Spilto Respimat	206	Tambocor	41
Respiratory	208, 211	Spiractin	46	Tamoxifen citrate	150
Various	225	Spiramycin	87	Tamoxifen Sandoz	150
Sodium chloride with sodium bicarbonate	208	Spiriva	206	Tamsulosin hydrochloride	62
Sodium citrate		Spiriva Respimat	206	Tamsulosin-Rex	62
Alimentary	5	Spiroolactone	46	Tarceva	140
Extemporaneously Compounded Preparations	229	Sprycel	140	Tasigna	142
Sodium citrate with sodium chloride and potassium chloride	32	Standard Feeds	244	Tasmar	107
Sodium citrate with sodium lauryl sulphoacetate	13	Staphlex	78	Tecfidera	123
Sodium citro-tartrate	63	Starch	229	Tegretol	115
Sodium cromoglicate		Stavudine	88	Tegretol CR	115
Alimentary	7	Sterculia with frangula	12	Teicoplanin	81
Respiratory	205, 210	SteroClear	205	Teicoplanin Mylan	81
Sensory	214	Stesolid	115	Temaccord	137
Sodium dihydrogen phosphate [Sodium acid phosphate]	37	Stimulants / ADHD Treatments	125	Temazepam	125
Sodium fluoride	18	Stiripentol	117	Temozolomide	137
Sodium fusidate [Fusidic acid]		Stocrin	87	Tenecteplase	33
Dermatological	54	Streptomycin sulphate	74	Tenofovir disoproxil	90
Infections	81	Stromectol	85	Tenofovir Disoproxil Teva	90
Sensory	212	Sucalfate	8	Tenoxicam	104
Sodium hyaluronate [Hyaluronic acid]		Sucrose	111	Terazosin	41
Alimentary	20	Sugammadex	103	Terbinafine	84
Sensory	215, 218	Sulfadiazine silver	54	Terbutaline	62
Sodium hyaluronate [Hyaluronic acid] with chondroitin sulphate	215	Sulfasalazine	7	Terbutaline sulphate	208
Sodium hypochlorite	221	Sulindac	104	Teriflunomide	124
Sodium metabisulfite	229	Sulphacetamide sodium	212	Teriparatide	101
Sodium nitrite	219	Sulphadiazine	81	Terlipressin	73
Sodium nitroprusside		Sulphur	229	Testosterone	64
Cardiovascular	49	Sulprix	120	Testosterone cipionate	64
Optional Pharmaceuticals	257	Sumatriptan	118	Testosterone esters	64
Sodium phenylbutyrate	17	Sunitinib	145	Testosterone undecanoate	64
Sodium phosphate with phosphoric acid	13	Sunscren, proprietary	59	Tetrabenazine	106
Sodium polystyrene sulphonate	38	Suprane	107	Tetracaine [Amethocaine] hydrochloride	
Sodium stibogluconate	87	Surgical Preparations	225	Nervous	110
Sodium tetradecyl sulphate	28	Sustagen Diabetic (Vanilla)	236	Sensory	215
Sodium thiosulfate	219	Sustagen Hospital Formula Active (Choc)	245	Tetracosactide [Tetracosactrin]	68
Sodium valproate	117	Sustagen Hospital Formula Active (Van)	245	Tetracosactrin	68
Sodium with potassium	226	Sutent	145	Tetracycline	80
Solifenacin Mylan	63	Suxamethonium chloride	103	Thalidomide	138
Solifenacin succinate	63	Sylvant	193	Thalomid	138
Solu-Cortef	66	Symmetrel	106	Thiobroma oil	229
Solu-Medrol	66	Sympathomimetics	48	Theophylline	210
Solu-Medrol Act-O-Vial	66	Synacthen	68	Thiamine hydrochloride	22
Somatropin	68	Synacthen Depot	68	Thioguanine	134
Sotalol	43	Synflorix	248	Thiopental [Thiopentone] sodium	108
Soya oil	219	Syntometrine	61	Thiopentone	108
Spacer device	257	Syrup	229	Thiotepa	131
Span-K	38	Systane Unit Dose	217	Thrombin	28
		- T -		Thymol glycerin	20
		Tacrolimus	151	Thyroid and Antithyroid Preparations	72
		Tacrolimus Sandoz	151	Thyrotropin alfa	68
		Tagitol V	222	Ticagrelor	32
		Talc	211	Ticarillin with clavulanic acid	78

Ticlopidine .....	33	Triamcinolone acetonide with gramicidin, neomycin and nystatin .....	213	vaccine] .....	254
Tigecycline.....	80	Triamcinolone acetonide with neomycin sulphate, gramicidin and nystatin .....	58	Varicella zoster vaccine [Shingles vaccine] .....	255
Tilcotil .....	104	Triamcinolone hexacetonide.....	66	Varivax.....	254
Timolol.....	216	Triazolam .....	125	Vasodilators.....	49
Timolol maleate.....	43	Trichloracetic acid.....	229	Vasopressin.....	73
Timoptol XE .....	216	Trientine dihydrochloride .....	17	Vasopressin Agents.....	73
Tiotropium bromide.....	206	Trimethoprim .....	81	Vasorex .....	43
Tiotropium bromide with olodaterol.....	206	Trimethoprim with sulphamethoxazole [Co-trimoxazole] .....	81	Vecuronium bromide .....	103
Tivicay .....	90	Trometamol .....	225	Vedafil.....	51
TMP .....	81	Tropicamide.....	217	Veletri .....	52
TOBI .....	74	Tropisetron .....	120	Venclexta.....	138
Tobradex .....	212	Tropisetron-AFT .....	120	Venetoclax.....	138
Tobramycin		Tuberculin PPD [Mantoux] test.....	256	Venlafaxine.....	114
Infections.....	74	Tubersol.....	256	Venofer.....	18
Sensory.....	212	Two Cal HN .....	238	VENOX.....	204
Tobramycin BNM.....	74	TwoCal HN RTH (Vanilla).....	238	Ventavis.....	52
Tobramycin Mylan .....	74	Tykerb.....	142	Ventolin.....	208
Tobrex .....	212	Tysabri.....	124	Vepesid.....	134
Tocilizumab .....	194	- U -		Verapamil hydrochloride.....	44
Tofranil.....	114	Ultibro Breezhaler.....	206	Vergo 16.....	119
Tolcapone.....	107	Ultraproct .....	7	Vermox.....	85
Topamax.....	117	Umeclidinium.....	206	Vesanoid.....	138
Topicaïne.....	109	Umeclidinium with vilanterol .....	206	Vexazone.....	11
Topical Products for Joint and Muscular Pain.....	105	Univent .....	205	Vfend .....	83
Topiramate .....	117	Ural.....	63	Vigabatrin .....	117
Topiramate Actavis.....	117	Urea		Vildagliptin .....	11
Torbay .....	49	Dermatological.....	57	Vildagliptin with metformin hydrochloride .....	11
Tracrium .....	102	Extemporaneously Compounded Preparations .....	229	Vimpat .....	116
Tramadol hydrochloride.....	113	Urex Forte.....	45	Vinblastine sulphate .....	147
Tramal 100 .....	113	Urografin.....	222	Vincristine sulphate .....	147
Tramal 50 .....	113	Urokinase .....	34	Vinorelbine.....	147
Tramal SR 100 .....	113	Urologicals.....	62	Viral Vaccines.....	250
Tramal SR 150 .....	113	Uromitexan .....	147	Viramune Suspension .....	87
Tramal SR 200 .....	113	Ursodeoxycholic acid.....	11	ViruPOS.....	212
Trandate.....	43	Ursosan .....	11	Viscoat.....	215
Tranexamic acid .....	28	Utrogestan .....	61	Visipaque.....	222
Tranexamic-AFT .....	28	- V -		Vit.D3.....	22
Tranylcypromine sulphate .....	114	Vaclovir .....	91	VitA-POS.....	218
Trastuzumab.....	197	Valaciclovir .....	91	Vital.....	237
Trastuzumab emtansine.....	198	Valganciclovir .....	91	Vitamin B complex.....	22
Travoprost .....	216	Valganciclovir Mylan.....	91	Vitamin B6 25 .....	21
Travopt .....	216	Vancomycin.....	81	Vitamins.....	20
Treatments for Dementia.....	127	Varenicline.....	129	Vivonex TEN.....	236
Treatments for Substance Dependence .....	128	Varenicline Pfizer.....	129	Volibris.....	50
Tretinoin		Varibar - Honey .....	222	Voltaren .....	104
Dermatological.....	55	Varibar - Nectar .....	222	Voltaren D.....	104
Oncology.....	138	Varibar - Pudding.....	222	Voltaren Ophtha .....	214
Trexate .....	133	Varibar - Thin Liquid .....	222	Volumatic.....	257
Tri-sodium citrate.....	229	Varicella vaccine [Chickenpox vaccine] .....	254	VoLumen .....	222
Triamcinolone acetonide				Voriconazole.....	83
Alimentary.....	20			Votrient .....	144
Dermatological.....	58			Vttack.....	83
Hormone Preparations.....	66			- W -	
				Warfarin sodium .....	32

Wart Preparations.....	59	Zuclopenthixol decanoate.....	123
Water		Zuclopenthixol hydrochloride.....	121
Blood.....	37	Zusdone.....	121
Various.....	225	Zyban.....	128
Wool fat		Zypine.....	121
Dermatological.....	57	Zypine ODT .....	121
Extemporaneously Compounded		Zyprexa Relprevv .....	122
Preparations.....	229	Zytiga.....	147
- X -		Zyvox.....	81
X-Opaque-HD.....	222		
Xanthan.....	229		
Xarelto.....	32		
Xifaxan.....	9		
Xolair.....	179		
Xylocaine.....	109		
Xylometazoline hydrochloride.....	208		
Xyntha.....	29		
- Y -			
Yellow jacket wasp venom .....	204		
- Z -			
Zanamivir.....	94		
Zantac.....	8		
Zapril.....	39		
Zarontin.....	115		
Zavedos.....	132		
Zeffix.....	90		
Zetlam.....	90		
Ziagen.....	88		
Zidovudine [AZT] .....	88		
Zidovudine [AZT] with			
lamivudine .....	88		
Zimybe.....	47		
Zinc			
Alimentary.....	19		
Dermatological.....	55		
Zinc and castor oil.....	56		
Zinc chloride .....	19		
Zinc oxide.....	229		
Zinc sulphate.....	19		
Zinc with wool fat.....	56		
Zincaps.....	19		
Zinfo.....	76		
Zinnat.....	75		
Ziprasidone.....	121		
Zista.....	205		
Zithromax.....	76		
Zoladex.....	68		
Zoledronic acid			
Hormone Preparations.....	65		
Musculoskeletal .....	98		
Zoledronic acid Mylan.....	65		
Zopiclone.....	125		
Zostavax.....	255		
Zostrix.....	105		
Zostrix HP.....	110		
Zuclopenthixol acetate.....	121		



























