January 2021 Volume 28 Number 0

Editors:

Kaye Wilson & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

Published each April, August and December. Changes to the contents are published in monthly updates.

Accessible in an electronic format at no cost from the PHARMAC website www.pharmac.govt.nz/schedule.

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth & John Geering email: texschedule@pharmac.govt.nz @Pharmaceutical Management Agency



ISSN 1179-3686 pdf ISSN 1172-9376 print

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to PHARMAC and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to PHARMAC should be in written form and not by reproduction of the PHARMAC logo. While care has been taken in compiling this Schedule, PHARMAC takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

2	Introducing PHARMAC	
5	General Rules	Section A
6	Alimentary Tract & Metabolism	Section B
38	Blood & Blood Forming Organs	
48	Cardiovascular System	
62	Dermatologicals	
72	Genito Urinary System	
79	Hormone Preparations – Systemic	
90	Infections – Agents For Systemic Use	
112	Musculoskeletal System	
120	Nervous System	
157	Oncology Agents & Immunosuppressants	
231	Respiratory System & Allergies	
239	Sensory Organs	
244	Various	
246	Extemporaneous Compounds (ECPs)	Section C
249	Special Foods	Section D
270	National Immunisation Schedule	Section I
280	Index	

Introducing PHARMAC

Introducing PHARMAC

The Pharmaceutical Management Agency (PHARMAC) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. PHARMAC negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

PHARMAC's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about PHARMAC and the way we make funding decisions can be found on the PHARMAC website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by PHARMAC.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements PHARMAC has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg p sachet		30	•	Gaviscon Infant
SODIUM ALGINATE Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (8.60)	60		Gaviscon Double Strength
Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 m	-	Acidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE * Tab 600 mg	12.56	100	✓.	Alu-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsementOnly when prescribed for patients unable to swallow cal inappropriate and the prescription is endorsed according	cium carbonate tablet	500 m s or v		Roxane um carbonate tablets are
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on * Tab 2 mg* * Cap 2 mg	10.75	400 400		Nodia Diamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE Cap 3 mg - Special Authority see SA1886 below - Retail pharmacy	166.50	90	✓	Entocort CIR
⇒SA1886 Special Authority for Subsidy Initial application — (Crohn's disease) from any relevant practithe following criteria: Both:	titioner. Approvals va	ılid fo	r 6 months	for applications meeting
Mild to moderate ileal, ileocaecal or proximal Crohn's disc	ease; and			

2.3 Osteoporosis where there is significant risk of fracture; or

2 Any of the following: 2.1 Diabetes; or 2.2 Cushingoid habitus; or

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)	26.55	21.1 g OP	✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLO	RIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%	26.55	10 g OP	✓ Proctofoam S29
MESALAZINE			
Tab 400 mg	49.50	100	✓ Asacol
Tab EC 500 mg	49.50	100	✓ Asamax
Tab long-acting 500 mg	56.10	100	✓ Pentasa
Tab 800 mg	85.50	90	✓ Asacol
Modified release granules, 1 g	118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml		7	✓ Pentasa
Suppos 500 mg	22.80	20	✓ Asacol
Suppos 1 g	54.60	30	✓ Pentasa
OLSALAZINE			
Tab 500 mg	93.37	100	Dipentum
Cap 250 mg		100	✓ Dipentum

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PREDNISOLONE SODIUM				
Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	•	Essential Prednisolone S29
SODIUM CROMOGLICATE Cap 100 mgSULFASALAZINE	92.91	100	✓	Nalcrom
* Tab 500 mg		100 100		Salazopyrin Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CI	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and		
cinchocaine hydrochloride 5 mg per g6.35	30 g OP	Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and		
cinchocaine hydrochloride 1 mg2.66	12	Ultraproct
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl

Management of Anal Fissures

GLYCERYL TRINITRATE − Special Authority see SA1329 below − Retail pharmacy

★ Oint 0.2%......22.00 30 g OP

✓ Rectogesic

⇒SA1329 Special Authority for Subsidy

CL VCODVDDONII IM DDOMIDE

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule - Up to 10 inj available on a PSO	17.14	10	✓ Max Health
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg	6.35	100	Buscopan
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO		5	✓ Buscopan
MEBEVERINE HYDROCHLORIDE			
* Tab 135 mg	9.20	90	✓ Colofac

Antiulcerants

Antisecretory and Cytoprotective

MISOPROSTOL

Subsidised on a PSO only if from a Family Planning New Zealand Clinic or an abortion service provider with a DHB contract and the PSO is endorsed with the name of the institution for which the PSO is required.

★ Tab 200 mcg - Up to 120 tab available on a PSO41.50
120 ✓ Cytotec

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

Helicobacter Pylori Eradication

CLARITHROMYCIN

Tab 500 mg − Subsidy by endorsement......10.40 14 ✓ Apo-Clarithromycin

- a) Maximum of 14 tab per prescription
- Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.
 Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

H2 Antagonists

	MOTIDINE – Only on a prescription Tab 20 mg	<i>1</i> Q1	100	✓ Famotidine
~	1 ab 20 mg	4.01	100	Hovid \$29
*	Tab 40 mg	8.48	100	✓ Famotidine
				Hovid S29
*	Inj 10 mg per ml, 4 ml - Subsidy by endorsement		. •	✓ Mylan S29
	Subsidy by endorsement – Subsidised for patients receiving tre	eatment as pa	rt of palliativ	e care.

RANITIDINE - Subsidy by endorsement

- a) Only on a prescription
- b) Subsidy by endorsement Subsidised for patients who were taking ranitidine prior to 1 November 2019 and the
 prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record
 of prior dispensing of ranitidine.

*	Oral liq 150 mg per 10 ml5	.14	300 ml	✓ Peptisoothe
*	Inj 25 mg per ml, 2 ml13	.40	5	✓ Zantac
/ D	" " 0 1" 150			

(Peptisoothe Oral liq 150 mg per 10 ml to be delisted 1 September 2021) (Zantac Inj 25 mg per ml, 2 ml to be delisted 1 June 2021)

Proton Pump Inhibitors

Proton Pump inimpitors		
LANSOPRAZOLE		
* Cap 15 mg4.58	100	✓ Lanzol Relief
* Cap 30 mg5.41	100	✓ Lanzol Relief
OMEPRAZOLE		
For omeprazole suspension refer Standard Formulae, page 246		
* Cap 10 mg1.98	90	Omeprazole actavis10
* Cap 20 mg1.96	90	Omeprazole actavis20
* Cap 40 mg3.12	90	 Omeprazole actavis 40
* Powder – Only in combination42.50	5 g	✓ Midwest
Only in extemporaneously compounded omeprazole suspension.	•	
* Inj 40 mg ampoule with diluent33.98	5	✓ Dr Reddy's
		<u>Omeprazole</u>
		✓ Ocicure S29
PANTOPRAZOLE		
* Tab EC 20 mg2.02	100	✓ Panzop Relief
* Tab EC 40 mg2.85	100	✓ Panzop Relief

	Subsidy (Manufacturer's Price \$) Per	Fully Brand or Subsidised Generic Manufacturer
Site Protective Agents			
COLLOIDAL BISMUTH SUBCITRATE Tab 120 mgSUCRALFATE	14.51	50	✓ Gastrodenol \$29
Tab 1 g	35.50 (48.28)	120	Carafate
Bile and Liver Therapy			
RIFAXIMIN – Special Authority see SA1461 below Tab 550 mg	625.00	56	✓ Xifaxan
■ SA1461 Special Authority for Subsidy nitial application only from a gastroenterologist nepatologist. Approvals valid for 6 months where olerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist. Approvals valid without further ren- penefiting from treatment.	e the patient has hepatic encephalop ogist or Practitioner on the recomme	oathy d	lespite an adequate trial of maximum n of a gastroenterologist or
Diabetes			
Hyperglycaemic Agents			
DIAZOXIDE - Special Authority see SA1320 be	low – Retail pharmacy		
Cap 25 mg		100	
Cap 100 mg		100	✓ Proglicem S29
Oral liq 50 mg per ml	620.00	80 ml C	OP ✓ Proglycem S29
➤SA1320 Special Authority for Subsidy nitial application from any relevant practitioner	. Approvals valid for 12 months whe	re use	ed for the treatment of confirmed
nypoglycaemia caused by hyperinsulinism. Renewal from any relevant practitioner. Approva appropriate and the patient is benefiting from trea		ss noti	ified where the treatment remains
GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit - Up to 5 kit available or	n a PSO32.00	1	✓ Glucagen Hypokit
Insulin - Short-acting Preparations			
NSULIN NEUTRAL			
Inj human 100 u per ml		0 ml C	✓ Humulin R
▲ Inj human 100 u per ml, 3 ml	42.66	5	✓ Actrapid Penfill✓ Humulin R
Insulin - Intermediate-acting Prepar	ations		
NSULIN ASPART WITH INSULIN ASPART PRO			
▲ Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	✓ NovoMix 30 FlexPen

	Subsidy	F	ully Brand or
	(Manufacturer's Price	e) Subsidis	,
	\$	Per	✓ Manufacturer
NSULIN ISOPHANE			
Inj human 100 u per ml	17.68		✓ Humulin NPH✓ Protaphane
Inj human 100 u per ml, 3 ml	29.86	5	✓ Humulin NPH
- , ,			✓ Protaphane Penfill
NSULIN ISOPHANE WITH INSULIN NEUTRAL			
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70✓ Mixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70
•			✓ PenMix 30
			✓ PenMix 40
JOHN N. JORDO WITH INCH IN LIARDO PROTAMINE			✓ PenMix 50
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE Inj lispro 25% with insulin lispro protamine 75% 100 u per ml.			
3 ml		5	✓ Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml			
3 ml		5	✓ Humalog Mix 50
Insulin - Long-acting Preparations			
SULIN GLARGINE			
Inj 100 u per ml, 10 ml	63.00	1	✓ Lantus
Inj 100 u per ml, 3 ml		5	✓ Lantus
Inj 100 u per ml, 3 ml disposable pen		5	✓ Lantus SoloStar
Insulin - Rapid Acting Preparations			
NSULIN ASPART			
Inj 100 u per ml, 10 ml			✓ NovoRapid
Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe	51.19	5	✓ NovoRapid FlexPen
NSULIN GLULISINE Inj 100 u per ml, 10 ml	27.02	1	✓ Apidra
Inj 100 u per ml, 3 ml			✓ Apidra
Inj 100 u per ml, 3 ml disposable pen			✓ Apidra SoloStar
NSULIN LISPRO			
Inj 100 u per ml, 10 ml		10 ml OP	✓ Humalog
Inj 100 u per ml, 3 ml	59.52	5	✓ Humalog
Alpha Glucosidase Inhibitors			
CARBOSE			
★ Tab 50 mg		90	✓ Glucobay
₭ Tab 100 mg	10.47 6.40	90	✓ Accarb✓ Glucobay
- 145 155 Hig	20.23	00	✓ Accarb
Oral Hypoglycaemic Agents			
ilibenclamide			
₭ Tab 5 mg	6.00	100	✓ <u>Daonil</u>
			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
GLICLAZIDE * Tab 80 mg	15.18	500	•	Glizide
GLIPIZIDE * Tab 5 mg	3.27	100	•	<u>Minidiab</u>
METFORMIN HYDROCHLORIDE * Tab immediate-release 500 mg Tab immediate-release 850 mg	8.63 7.04	1,000 500	_ '	Apotex Apotex
PIOGLITAZONE * Tab 15 mg * Tab 30 mg * Tab 45 mg	3.47 5.06	90 90 90	1	<u>Vexazone</u> <u>Vexazone</u> Vexazone
VILDAGLIPTIN Tab 50 mg		60		Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE Tab 50 mg with 1,000 mg metformin hydrochloride Tab 50 mg with 850 mg metformin hydrochloride		60 60	_	Galvumet Galvumet

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes: or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

10 strip OP ✓ KetoSens

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

✓ CareSens Dual diagnostic test strips.......20.00 1 OP

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	I Generic	
\$	Per 🗸	Manufacturer	

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	✓ CareSens N
		✓ CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood g	lucose test stri	os26.20	50 test OP	✓ SensoCard
---------	------------------	---------	------------	-------------

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

INSULIN PEN NEEDLES	 Maximum of 200 	dev per	prescription
---------------------	------------------------------------	---------	--------------

IIVC	oblini bin nebelo inaximam of 200 dev per prescripti	OH		
*	29 g × 12.7 mm	10.50	100	✓ B-D Micro-Fine
*	31 g × 5 mm		100	✓ B-D Micro-Fine
*	31 g × 6 mm		100	✓ Berpu
*	31 g × 8 mm		100	✓ B-D Micro-Fine
*	32 g × 4 mm		100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEED	LE - Maximum of 2	00 dev per p	prescription
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.00	100	✓ B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g x 8 mm needle	13.00	100	✓ B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.00	100	✓ B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	13.00	100	B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.00	100	B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.00	100	B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

c) Maximum of 1 insulin pump per patient each four	year period.		
Min basal rate 0.025 U/h		1	✓ MiniMed 640G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			X2 with Basal-IQ

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
\$	Per	✓	Manufacturer

continued...

education from an appropriate health professional); and

- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist: or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Eithei
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Drand or	_
Subsidy		. ,	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol: and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Fither:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Insulin Pump Consumables

⇒SA1906 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	1	Manufacturer	

continued...

8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline, according to the most recent result.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol. according to the most recent result: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline.

Initial application — (**Previous use before 1 September 2012**) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Subsic		Fully	Brand or
(Manufacture	r's Price) Subs	idised	Generic
\$	Per	1	Manufacturer

continued...

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol according to a recent laboratory result; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application, according to the most recent result; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline.

INSULIN PUMP CARTRIDGE - Special Authority see SA1906 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 packs of cartridge sets will be funded per year.

Subsidy Fully Brand or (Manufacturer's Price) Generic Subsidised \$ Per Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA) - Special Authority see SA1906 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription

c) Maximum of 13 infusion sets will be funded per year.			
10 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Sure-T MMT-884A
10 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Sure-T MMT-886A
6 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Sure-T MMT-864A
6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Sure-T MMT-866A
8 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Sure-T MMT-874A
8 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Sure-T MMT-876A
10 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T
10 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles	130.00	1 OP	MMT-884 ✓ Paradigm Sure-T
6 mm steel needle; 29 G; manual insertion; 60 cm tubing ×			MMT-886
10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T MMT-864
6 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-863
6 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T
8 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T
8 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock	130.00	1 OP	✓ Sure-T MMT-873
8 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Sure-T

(Paradigm Sure-T MMT-884 10 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Sure-T MMT-886 10 mm steel needle; 29 G; manual insertion; 80 cm tubing x 10 with 10 needles to be delisted 1 April

(Paradigm Sure-T MMT-864 6 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Sure-T MMT-866 6 mm steel needle; 29 G; manual insertion; 80 cm tubing x 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Sure-T MMT-874 8 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles to be delisted 1 April

(Paradigm Sure-T MMT-876 8 mm steel needle; 29 G; manual insertion; 80 cm tubing x 10 with 10 needles to be delisted 1 April 2021)

MMT-876

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) – Special Authority see SA1906 on page 17 – Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles	1 OP	✓ TruSteel
6 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles130.00) 1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles130.00	0 1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles130.00	1 OP	✓ TruSteel

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1906 on page 17 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription

c) Maximum of 13 infusion sets will be funded per ye	ear.		
13 mm teflon needle, 110 cm tubing × 10		1 OP	✓ MiniMed Silhouette MMT-382A
13 mm teflon needle, 45 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-368A
13 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-381A
13 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-383A
17 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-377A
17 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-378A
17 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-384A
6 mm teflon needle, 110 cm tubing x 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-398A
6 mm teflon needle, 45 cm blue tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-941A
6 mm teflon needle, 45 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-921A
6 mm teflon needle, 60 cm blue tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-943A
6 mm teflon needle, 60 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-923A
6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-399A
6 mm teflon needle, 80 cm blue tubing	130.00	1 OP	✓ MiniMed Mio MMT-945A
6 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-965A
6 mm teflon needle, 80 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-925A
6 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-387A
9 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-396A
9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-397A
9 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-975A
9 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-386A

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION WITH INSERTION DEVICE) - Special Authority see SA1906 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION) - Special Authority see SA1906 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

130.00	1 OP	✓ Paradigm Silhouette MMT-382
130.00	1 OP	✓ Paradigm Silhouette MMT-368
130.00	1 OP	✓ Paradigm Silhouette MMT-381
130.00	1 OP	✓ Paradigm Silhouette MMT-383
130.00	1 OP	✓ Paradigm Silhouette MMT-377
130.00	1 OP	✓ Paradigm Silhouette MMT-378
130.00	1 OP	✓ Silhouette MMT-373
130.00	1 OP	✓ Paradigm Silhouette
	130.00130.00130.00130.00130.00130.00130.00	130.00 1 OP130.00 1 OP130.00 1 OP130.00 1 OP130.00 1 OP

(Paradigm Silhouette MMT-382 13 mm teflon cannula; angle insertion; 120 cm line x 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Silhouette MMT-368 13 mm teflon cannula; angle insertion; 45 cm line × 10 with 10 needles to be delisted 1 April 2021) (Paradigm Silhouette MMT-381 13 mm teflon cannula; angle insertion; 60 cm line × 10 with 10 needles to be delisted 1 April 2021) (Paradigm Silhouette MMT-383 13 mm teflon cannula; angle insertion; 80 cm line × 10 with 10 needles to be delisted 1 April 2021) (Paradigm Silhouette MMT-377 17 mm teflon cannula; angle insertion; 110 cm line × 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Silhouette MMT-378 17 mm teflon cannula; angle insertion; 60 cm line \times 10 with 10 needles to be delisted 1 April 2021) (Paradigm Silhouette MMT-384 17 mm teflon cannula; angle insertion; 80 cm line \times 10 with 10 needles to be delisted 1 April 2021)

MMT-384

6 mm teflon cannula: straight insertion: insertion device: 60 cm

9 mm teflon cannula; straight insertion; insertion device;

line × 10 with 10 needles......140.00

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	/	Manufacturer	

INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION WITH INSERTION DEVICE) - Special Authority see SA1906 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- h) Only on a prescription

c) Maximum of 13 infusion sets will be funded per year.			
6 mm teflon cannula; straight insertion; insertion device; 45 cm blue tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Mio MMT-941
6 mm teflon cannula; straight insertion; insertion device; 45 cm pink tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Mio MMT-921
6 mm teflon cannula; straight insertion; insertion device; 60 cm blue tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Mio MMT-943
6 mm teflon cannula; straight insertion; insertion device; 60 cm pink tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Mio MMT-923
6 mm teflon cannula; straight insertion; insertion device; 80 cm blue tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Mio MMT-945
6 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Mio
6 mm teflon cannula; straight insertion; insertion device; 80 cm pink tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Mio
9 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing \times 10 with 10 needles	130.00	1 OP	✓ Paradigm Mio MMT-975
6 mm teflon cannula; straight insertion; insertion device; 110 cm line x 10 with 10 needles	140.00	1 OP	✓ AutoSoft 90

1 OP

1 OP

✓ AutoSoft 90

✓ AutoSoft 90

Subsidy		Fully	Brand or	
		i uliy	Dialiu di	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	1	Manufacturer	

9 mm teflon cannula; straight insertion; insertion device; 60 cm

(Paradigm Mio MMT-941 6 mm teflon cannula; straight insertion; insertion device; 45 cm blue tubing x 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Mio MMT-921 6 mm teflon cannula; straight insertion; insertion device; 45 cm pink tubing \times 10 with 10 needles to be delisted 1 April 2021)

(Paradigm $\dot{\text{Mio}}$ MMT-943 6 mm teflon cannula; straight insertion; insertion device; 60 cm blue tubing \times 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Mio MMT-923 6 mm teflon cannula; straight insertion; insertion device; 60 cm pink tubing \times 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Mio MMT-945 6 mm teflon cannula; straight insertion; insertion device; 80 cm blue tubing x 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Mio MMT-965 6 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing \times 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Mio MMT-925 6 mm teflon cannula; straight insertion; insertion device; 80 cm pink tubing \times 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Mio MMT-975 9 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing \times 10 with 10 needles to be delisted 1 April 2021)

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION) - Special Authority see SA1906 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm teflon cannula; straight insertion; 110 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Quick-Set
6 mm toflen connula: atraight insertion; 60 cm tubing v 10 with			MMT-398
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-399
6 mm teflon cannula; straight insertion; 60 cm tubing \times 10 with			
10 needles; luer lock	130.00	1 OP	✓ Quick-Set MMT-393
6 mm teflon cannula; straight insertion; 80 cm tubing × 10 with 10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-387
9 mm teflon cannula; straight insertion; 106 cm tubing \times 10 with			
10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-396
9 mm teflon cannula; straight insertion; 60 cm tubing \times 10 with			_
10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-397
9 mm teflon cannula; straight insertion; 60 cm tubing \times 10 with			
10 needles; luer lock	130.00	1 OP	✓ Quick-Set MMT-392
9 mm teflon cannula; straight insertion; 80 cm tubing × 10 with	100.00	1 OD	. Dougalisms Ovials Cat
10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-386

(Paradigm Quick-Set MMT-398 6 mm teflon cannula; straight insertion; 110 cm tubing × 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Quick-Set MMT-399 6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Quick-Set MMT-387 6 mm teflon cannula; straight insertion; 80 cm tubing x 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Quick-Set MMT-396 9 mm teflon cannula; straight insertion; 106 cm tubing × 10 with 10 needles to be delisted 1 April 2021)

Paradigm Quick-Set MMT-397 9 mm teflon cannula; straight insertion; 60 cm tubing x 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Quick-Set MMT-386 9 mm teflon cannula; straight insertion; 80 cm tubing x 10 with 10 needles to be delisted 1 April 2021)

INSULIN PUMP RESERVOIR - Special Authority see SA1906 on page 17 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 packs of reservoir sets will be funded per year

10 × luer lock conversion cartridges 1.8 ml for Paradigm pumps50.00 Cartridge for 5 and 7 series pump; 1.8 ml × 10	1 OP 1 OP	✓ ADR Cartridge 1.8✓ Paradigm
out in ago 10. o unu r conso pump, no mir comministration		1.8 Reservoir
Cartridge for 7 series pump; 3.0 ml × 1050.00	1 OP	✓ Paradigm 3.0 Reservoir

Fully

Brand or

	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer	
Digestives Including Enzymes					
PANCREATIC ENZYME					
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ <u>Cı</u>	reon 10000	
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylas 1,250 U protease))	*	100	√ Pa	anzytrat	
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ <u>Cı</u>	reon 25000	
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 F	Ph				
Eur U)	34.93	20 g OF	Cı	reon Micro	
URSODEOXYCHOLIC ACID – Special Authority see SA1739 b	'	cy 100	✓ <u>Ur</u>	rsosan	

Subsidy

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner.

Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Patient has been diagnosed with Alagille syndrome: or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6

Su	Subsidy	Fully	Brand or
(Manufac	cturer's Price) Subsid	dised	Generic
	\$ Per	•	Manufacturer

continued...

months where the patient continues to benefit from treatment.

ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Laxatives

Bulk-forming Agents

* Powder for oral soln	12.20	500 g OP	✓ Konsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS		·	
* Dry	6.02	500 g OP	
·	(17.32)	•	Normacol Plus
	2.41	200 g OP	
	(8.72)		Normacol Plus

Faecal Softeners

* Tab 50 mg		✓ Coloxyl
* Tab 120 mg3.13	100	✓ Coloxyl
DOCUSATE SODIUM WITH SENNOSIDES		
* Tab 50 mg with sennosides 8 mg	200	✓ Laxsol
POLOXAMER – Only on a prescription		
Not funded for use in the ear.		
* Oral drops 10%	30 ml OP	✓ Coloxyl

Opioid Receptor Antagonists - Peripheral

DOCUSATE SODIUM - Only on a prescription

MET	HYLNALTREXONE BROMIDE - Special Authorit	ty see SA1691 below – Retail pl	narmacy		
	nj 12 mg per 0.6 ml vial	36.00	1	1	Relistor
		246.00	7	1	Relistor

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Fither:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	S Per	ubsidised 🗸	Generic Manufacturer
Osmotic Laxatives	•			
GLYCEROL * Suppos 3.6 g – Only on a prescription LACTULOSE – Only on a prescription	9.25	20	✓ <u>P</u>	<u>SM</u>
* Oral liq 10 g per 15 ml	3.33	500 ml	√ L	aevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BI Powder for oral soln 13.125 g with potassium chloride 46.6 n	CARBONATE AND	SODIUN	I CHLORII	DE
sodium bicarbonate 178.5 mg and sodium chloride 350.	7 mg 6.70	30	✓ N	<u>lolaxole</u>
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%	2.50	1	√ F	leet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml,	, ,	iption		
5 ml		50	✓ <u>N</u>	<u>licolette</u>
Stimulant Laxatives				
BISACODYL – Only on a prescription * Tab 5 mg * Suppos 10 mg		200 10		ax-Tab ax-Suppositories
SENNA – Only on a prescription * Tab, standardised	2.17 (8.21)	100	S	Senokot
	0.43 (2.06)	20	S	Senokot

Metabolic Disorder Agents

⇒SA1920 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

⇒SA1921 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency: or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE – Special Authority see SA1922 below – Retail pharmacy

✓ Naglazyme

⇒SA1922 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
	\$	Per	✓	Manufacturer	
IDURSULFASE - Special Authority see SA1623 below - Retail p	harmacy				
Inj 2 mg per ml, 3 ml vial	4,608.30	1	✓ [Elaprase	

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Fither:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

⇒SA1923 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 Fither

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
- 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE - Special Authority see SA1599 below - Retail pharmacy

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE - Special Authority see SA1924 below - Retail pharmacy
Grans 483 mg per g.......2.016.00 174 g OP

✓ Pheburane

⇒SA1924 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

⇒SA1880 Special Authority for Subsidy

Special Authority approved by the Gaucher Treatment Panel

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The Co-ordinator, Gaucher Treatment Panel Phone: 04 460 4990 PHARMAC PO Box 10 254 Facsimile: 04 916 7571

Wellington Email: gaucherpanel@pharmac.govt.nz

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Access Criteria

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	✓	Manufacturer

continued...

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2) Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and
- 3) Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
- Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia, thrombocytopenia; at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or massive symptomatic splenomegaly; or
 - 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g. hips or shoulder); spontaneous fractures or vertebral collapse; chronic bone pain not controlled by other pharmaceuticals; or
 - 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
 - Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated: and
- 2) Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and three yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4) Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 5) Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT: and
- 6) Patient is compliant with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations are submitted to the Gaucher Panel for assessment as required.

Mouth and Throat

Agents Used in Mouth Ulceration

BENZYDAMINE HYDROCHLORIDE

Soln 0.15% – Higher subsidy of \$20.31 per 500 ml with

Additional subsidy by endorsement for a patient who has oral mucositis as a result of treatment for cancer, and the prescription is endorsed accordingly.

	Subsidy (Manufacturer's Pric \$	ce) Sub	Fully Brand or bsidised Generic Manufacturer	
CARMELLOSE SODIUM WITH GELATIN AND PECTIN	3	Per	Manuacturer	
Paste	17.20	56 a OP	✓ Stomahesive	
rasie	4.55	0	▼ Stomanesive	
		15 g OP	Orahaaa	
	(7.90)	r - OD	Orabase	
	1.52	5 g OP	Oughana	
December	(3.60)	00 - OD	Orabase	
Powder		28 g OP	01	
	(10.95)		Stomahesive	
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE				
★ Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP		
	(6.00)		Bonjela	
TRIAMCINOLONE ACETONIDE				
Paste 0.1%	5.33	5 g OP	✓ Kenalog in Ora	abase
1 4010 0.170		3 y Oi	- Kenalog III Ola	asusc
Oropharyngeal Anti-infectives				
AMPHOTERICIN B				
Lozenges 10 mg	5.86	20	✓ Fungilin	
/ICONAZOLE			•	
Oral gel 20 mg per g	171	40 g OP	✓ Decozol	
0 01 0	4.74	40 y OF	Decozoi	
NYSTATIN				
Oral liq 100,000 u per ml	1.76	24 ml OP	✓ Nilstat	
Other Oral Agents				
For folinic mouthwash, pilocarpine oral liquid or saliva substitu	ita formula rafar Stano	lard Formul	lae nage 246	
	ale ioinidia lelei Stant	iaiu i oiiiiui	iae, page 240	
THYMOL GLYCERIN			4	
★ Compound, BPC	9.15	500 ml	✓ PSM	
Vitamins				
When he B				
Vitamin B				
HYDROXOCOBALAMIN	- DOO 400		A Nove B40	
★ Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a	a PSO1.89	3	✓ <u>Neo-B12</u>	
			✓ Vita-B12	
	3.15	5	 Hydroxocobala 	
			Mercury Pha	ırma
PYRIDOXINE HYDROCHLORIDE				
a) No more than 100 mg per dose				
b) Only on a prescription				
Tab 25 mg - No patient co-payment payable	2 70	90	✓ Vitamin B6 25	
K Tab 50 mg		500	✓ Apo-Pyridoxin	۵
<u> </u>	13.03	500	• Apo-rynuoxin	C
THIAMINE HYDROCHLORIDE – Only on a prescription				
★ Tab 50 mg	7.09	100	Max Health	
/ITAMIN B COMPLEX				
★ Tab, strong, BPC	7.15	500	✓ Bplex	
			PIVA	

	Subsidy (Manufacturer's Pri	ce) Subsi Per	dised G	rand or eneric lanufacturer
Vitamin C				
ASCORBIC ACID a) No more than 100 mg per dose b) Only on a prescription				
* Tab 100 mg	9.90	500	✓ Cvit	<u>e</u>
Vitamin D				
ALFACALCIDOL * Cap 0.25 mcg * Cap 1 mcg * Oral drops 2 mcg per ml	87.98	100 100 20 ml OP	✓ One ✓ One ✓ One	-Alpha
* Cap 0.25 mcg * Cap 0.5 mcg		100 100		itriol-AFT itriol-AFT
COLECALCIFEROL * Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescrip Vit.D3 to be Sole Supply on 1 February 2021 * Oral liq 188 mcg per ml (7,500 iu per ml)		12 4.8 ml OP	✓ Vit.E	
	9.00	4.0 1111 01	· run	a
Multivitamin Preparations				
MULTIVITAMIN RENAL – Special Authority see SA1546 below * Cap	, ,	30	✓ Clini	icians Renal Vit
■ SA1546 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals value following criteria: Either: 1 The patient has chronic kidney disease and is receiving 2 The patient has chronic kidney disease grade 5, defined 15 ml/min/1.73 m² body surface area (BSA).	either peritoneal dia	llysis or haem	odialysis;	or
MULTIVITAMINS – Special Authority see SA1036 below – Ret ** Powder		200 g OP	✓ Paed	diatric Seravit
■ SA1036 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valinborn errors of metabolism. Renewal from any relevant practitioner. Approvals valid without approval for multivitamins. VITAMINS				
* Tab (BPC can strength)	11 45	1 000	✓ Mvit	e

• • •	7			
*	Tab (BPC cap strength)11	1.45	1,000	✓ Mvite
*	Cap (fat soluble vitamins A, D, E, K) - Special Authority see			
	SA1720 below – Retail pharmacy 23	3.40	60	✓ Vitabdecl

⇒SA1720 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency; or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

	<u> </u>	1 01	- Mananataron
Minerals			
Calcium			
CALCIUM CARBONATE			
* Tab eff 1.75 g (1 g elemental)	28.40	20	✓ Calcium Sandoz S29
* Tab 1.25 g (500 mg elemental)	6.69 7.52	250	✓ Calci-Tab 500 ✓ Arrow-Calcium
* Tab eff 1.25 g (500 mg elemental) - Subsidy by endorsement		76	✓ Cacit (\$29)
Subsidy by endorsement – Only when prescribed for paed considered unsuitable.			
Calcium Sandoz 329 Tab eff 1.75 g (1 g elemental) to be deliste Arrow-Calcium Tab 1.25 g (500 mg elemental) to be delisted 1 Ma	, ,		
CALCIUM GLUCONATE			
* Inj 10%, 10 ml ampoule	32.00	10	✓ Max Health - HameIn S29
	64.00	20	✓ Max Health S29
Fluoride			
SODIUM FLUORIDE * Tab 1.1 mg (0.5 mg elemental)	5.75	100	✓ PSM
lodine			
POTASSIUM IODATE Tab 253 mcg (150 mcg elemental iodine)	4.58	90	✓ <u>NeuroTabs</u>
Iron			
FERRIC CARBOXYMALTOSE – Special Authority see SA1840 bi		armacy 1	✓ Ferinject
⇒SA1840 Special Authority for Subsidy nitial application — (serum ferritin less than or equal to 20 mononths for applications meeting the following criteria:	cg/L) from any	relevant prac	titioner. Approvals valid for 3

Subsidy

(Manufacturer's Price)

Fully

Subsidised

Per

Brand or Generic

Manufacturer

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:

 - - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist,

	ALIMENTAR	Y TRAC	T AND	METABOLISM
	Subsidy (Manufacturer's Price \$	e) Sub Per	Fully sidised	Brand or Generic Manufacturer
continued				
anaesthetist or medical practitioner on the recommendation of a anaesthetist. Approvals valid for 3 months for applications mee			stetricia	n, gynaecologist or
Both:	ung the following chi	ciia.		
1 Patient has been diagnosed with iron-deficiency anaemia	a; and			
2 Any of the following:	nt and treatment has	nravan ina	ffa ativa v	
2.1 Patient has been compliant with oral iron treatme2.2 Treatment with oral iron has resulted in dose-limit		proven me	necuve;	OI
2.3 Patient has symptomatic heart failure, chronic kid	ney disease stage 3	or more or	active ir	nflammatory bowel disease
and a trial of oral iron is unlikely to be effective; or 2.4 Rapid correction of anaemia is required.	ſ			
Renewal — (iron deficiency anaemia) only from an internal m	nedicine physician, ol	ostetrician.	avnaec	ologist, anaesthetist or
medical practitioner on the recommendation of a internal medici	ne physician, obstetr			
Approvals valid for 3 months for applications meeting the following Both:	ng criteria:			
1 Patient continues to have iron-deficiency anaemia; and				
2 A re-trial with oral iron is clinically inappropriate.				
FERROUS FUMARATE				
* Tab 200 mg (65 mg elemental)	3.09	100	✓ <u>F</u>	erro-tab
FERROUS FUMARATE WITH FOLIC ACID * Tab 310 mg (100 mg elemental) with folic acid 350 mcg	4 68	60	√ F	erro-F-Tabs
FERROUS SULFATE		00		<u> </u>
* Oral liq 30 mg (6 mg elemental) per 1 ml	12.08	500 ml	√ <u>F</u>	erodan
FERROUS SULPHATE				
* Tab long-acting 325 mg (105 mg elemental)	2.06	30	✓ <u>F</u>	errograd
IRON POLYMALTOSE * Inj 50 mg per ml, 2 ml ampoule	34 50	5	√ F	errosig
				511-55.ig
Magnesium				
For magnesium hydroxide mixture refer Standard Formulae, pag	ge 246			
MAGNESIUM HYDROXIDE	00.00	055 1		
Suspension 8%	33.60	355 ml	✓ P	Phillips Milk of Magnesia S29
	72.20	500 ml	✓ T	'&R S29

'	72.20	500 ml	Magnesia S29 ✓ T&R S29
(T&R S29 Suspension 8% to be delisted 1 February 2021)			
MAGNESIUM SULPHATE			
* Inj 2 mmol per ml, 5 ml ampoule	10.21	10	✓ DBL
			✓ DBL S29 S29

Zinc

ZINIC	SI II	PHATE
ZIIVO	JUL	

★ Cap 137.4 mg (50 mg elemental)......11.00 100 ✓ Zincaps

Subsidy (Manufacturer's Price) \$

Fully Subsidised Per

Brand or Generic Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus: and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*: and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 ju per week.

Note: Indication marked with * is an unapproved indication

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Subsidised	
	\$	Per	1	Manufacturer
EPOETIN ALFA - Special Authority see SA1775 on the previous	page - Retail phan	macy		
Wastage claimable		,		
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	✓	Binocrit
Inj 2,000 iu in 1 ml, syringe		6	✓	Binocrit
Inj 3,000 iu in 0.3 ml, syringe	150.00	6	✓	Binocrit
Inj 4,000 iu in 0.4 ml, syringe		6	1	Binocrit
Inj 5,000 iu in 0.5 ml, syringe	125.00	6	1	Binocrit
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	1	Binocrit
Inj 8,000 iu in 0.8 ml, syringe	175.00	6	1	Binocrit
Inj 10,000 iu in 1 ml, syringe	197.50	6	1	Binocrit
Inj 40,000 iu in 1 ml, syringe	250.00	1	✓	<u>Binocrit</u>
Megaloblastic				
FOLIC ACID				
* Tab 0.8 mg	21.84	1,000	1	Apo-Folic Acid
* Tab 5 mg		500		Apo-Folic Acid
Oral lig 50 mcg per ml		25 ml C)P 🗸	Biomed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemonhilia Management group

ricators aroup in conjunction with the National I	iacinopinna management giot	φ.	
Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial	2,450.00	1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	✓ Alprolix
ELTROMBOPAG - Special Authority see SA1743 be	elow – Retail pharmacy		
Wastage claimable			
Tab 25 mg	1,550.00	28	Revolade
Tab 50 mg	3,100.00	28	Revolade

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy: and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab);
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding: or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Fither:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see SA1969 below

Inj 30 mg in 1 ml vial	3,570.00	1	✓ Hemlibra
Inj 60 mg in 0.4 ml vial	7,138.00	1	✓ Hemlibra
, ,	12,492.00	1	✓ Hemlibra
lni 150 mg in 1 ml vial	17.846.00	1	✓ Hemlibra

⇒SA1969 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and
- 2 Fither:

	AA	0 !- !	0
(Mar	ufacturer's Price)	Subsidised	Generic
	\$ P	Per 🗸	Manufacturer

continued...

- 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
- 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more; and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:
 - 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
 - 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
- 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe	2,356.60	1	✓ NovoSeven RT
Inj 5 mg syringe	·	1	✓ NovoSeven RT
Ini 8 mg syringe	9.426.40	1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	1,315.00	1	✓ FEIBA NF
Inj 1,000 U	2,630.00	1	✓ FEIBA NF
Inj 2,500 U	6,575.00	1	✓ FEIBA NF

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

287.50	1	Xyntha
	1	Xyntha
	575.00 1,150.00	575.00 1 1,150.00 1 2,300.00 1

NONACOG GAMMA. [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 iu vial	435.00	1	✓ RIXUBIS
Inj 1,000 iu vial	870.00	1	✓ RIXUBIS
Inj 2,000 iu vial		1	✓ RIXUBIS
Inj 3,000 iu vial	·	1	✓ RIXUBIS
, -,	,		

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) -[
For patients with haemophilia. Preferred Brand of short half-				
managed by the Haemophilia Treaters Group in conjunction v				
Inj 250 iu vial		1		Advate
Inj 500 iu vial		1		Advate
Inj 1,000 iu vial		1		Advate
Inj 1,500 iu vial Inj 2,000 iu vial	,	1		Advate Advate
Inj 3,000 iu vial	,	1		Advate
• •	,	'	•	Auvale
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE I				
For patients with haemophilia. Rare Clinical Circumstances E				
treatment is managed by the Haemophilia Treaters Group in	conjunction with the r	valior	iai naemi	pprillia Management Group,
subject to criteria. Inj 250 iu vial	227 50	1	1	Kogenate FS
Inj 500 iu vial		1		Kogenate FS
Inj 1,000 iu vial		1		Kogenate FS
Inj 2,000 iu vial		1		Kogenate FS
Inj 3,000 iu vial	,	1		Kogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]	*	-		
For patients with haemophilia A receiving prophylaxis treatme		d traa	tmont ic n	nanaged by the Haemonhilia
Treaters Group in conjunction with the National Haemophilia		u ii ca	unciii is ii	nanaged by the Haemophila
Inj 250 iu vial	0 0 1	1	1	Adynovate
Inj 500 iu vial		1		Adynovate
Inj 1,000 iu vial		i		Adynovate
Inj 2,000 iu vial	,	1		Adynovate
SODIUM TETRADECYL SULPHATE	,			.,
* Inj 3% 2 ml	28 50	5		
7 III 0/0 Z III	(73.00)	J		Fibro-vein
TDANEVAMIC ACID	(10.00)			TIDIO TOILI
TRANEXAMIC ACID Tab 500 mg	0.45	60	./	Mercury Pharma
Tab 500 flig	9.40	00	•	wercury Pharma
Vitamin K				
Vitaliili K				
PHYTOMENADIONE				
Inj 2 mg per 0.2 ml - Up to 5 inj available on a PSO	8.00	5	1	Konakion MM
Inj 10 mg per ml, 1 ml - Up to 5 inj available on a PSO	9.21	5	1	Konakion MM
Antithrombotic Agents				
Antiplatelet Agents				
ASPIRIN				
* Tab 100 mg	10.80	990	1	Ethics Aspirin EC
•	10.00	990	•	Ettiles Aspiriti EC
CLOPIDOGREL	4.00	0.4	,	Olevel de souel
* Tab 75 mg	4.60	84	•	Clopidogrel
				<u>Multichem</u>
DIPYRIDAMOLE			_	
* Tab long-acting 150 mg	10.90	60	/	Pytazen SR

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
PRASUGREL - Special Authority see SA1954 below - Retail pl	narmacy			
Tab 5 mg	108.00	28	✓	Effient
Tab 10 mg	120.00	28	1	Effient
(Effient Tab 5 mg to be delisted 1 February 2021) (Effient Tab 10 mg to be delisted 1 February 2021)				

⇒SA1954 Special Authority for Subsidy

Renewal — (coronary angioplasty and bare metal stent) from any relevant practitioner. Approvals valid for 6 months where the patient has undergone coronary angioplasty or had a bare metal cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Renewal — (drug eluting stent) from any relevant practitioner. Approvals valid for 12 months where had a drug-eluting cardiac stent inserted in the previous 4 weeks and is clopidogrel-allergic*.

Note: * Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Eithe
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Fither:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal — (**Percutaneous coronary intervention with stent deployment**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Heparin and Antagonist Preparations

ENOXAPARIN SODIUM - Special Authority see S	SA1646 below – Retail pharmacy
---	--------------------------------

Inj 20 mg in 0.2 ml syringe	27.93	10	Clexane
Inj 40 mg in 0.4 ml syringe	37.27	10	Clexane
Inj 60 mg in 0.6 ml syringe		10	Clexane
Inj 80 mg in 0.8 ml syringe		10	Clexane
Inj 100 mg in 1 ml syringe		10	Clexane
Inj 120 mg in 0.8 ml syringe		10	 Clexane Forte
Ini 150 mg in 1 ml svringe		10	✓ Clexane Forte

⇒SA1646 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or

	BLOOD AND	BLOOD I	FORI	MING ORGANS
	Subsidy (Manufacturer's Price)		Fully ised	Brand or Generic Manufacturer
continued				
3 For the prevention of thrombus formation in the extra-cor	poreal circulation duri	ng haemodia	alysis.	
Renewal — (Venous thromboembolism other than in pregna	ncy or malignancy)	from any re	levant	t practitioner. Approvals
valid for 1 month where low molecular weight heparin treatment				
(surgery, ACS, cardioversion, or prior to oral anti-coagulation).				
HEPARIN SODIUM				
Inj 1,000 iu per ml, 5 ml ampoule	58.57	50	✓ P	fizer
Inj 5,000 iu per ml, 1 ml	28.40	5		fizer
	32.66		✓ D	BL Heparin
				Sodium S29
				ospira
Inj 5,000 iu per ml, 5 ml ampoule		50	✓ <u>P</u>	
Inj 25,000 iu per ml, 0.2 ml		5		ospira
	42.40		✓ H	eparin DBL S29
(Pfizer Inj 5,000 iu per ml, 1 ml to be delisted 1 March 2021)				
HEPARINISED SALINE				
Inj 10 iu per ml, 5 ml	65.48	50	✓ P	fizer
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg - No more than 2 cap per day	76.36	60	✓ P	radaxa
Cap 110 mg	76.36	60	✓ P	radaxa
Cap 150 mg	76.36	60	✓ P	radaxa
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day	83.10	30	✓ X	arelto
Tab 15 mg - Up to 14 tab available on a PSO		28	✓ X	arelto
Tab 20 mg	77.56	28	✓ X	arelto
WARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg		50	-	oumadin
	6.46	100		larevan
* Tab 2 mg		50	-	oumadin
* Tab 5 mg		100		larevan
* Tab 5 mg	5.93 11.48	50 100	-	oumadin Iarevan
Plant Order of the Information	11.70	100	- 141	ui (vull

Blood Co	lony-stimu	lating Factors
-----------------	------------	----------------

FILGRASTIM - Special Authority see SA1259 below - Retail pharma	асу		
Inj 300 mcg per 0.5 ml prefilled syringe	96.22	10	✓ Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe	161.50	10	✓ Nivestim

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

Subsidy	I	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	/	Manufacturer

continued...

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10⁹/L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1912 below - Retail pharmacy

✓ Neulastim

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration

CI	LICOSE	[DEYTROSE]	

*	Inj 50%, 10 ml ampoule - Up to 5 inj available on a PSO30.65	5	✓ Biomed
*	Inj 50%, 90 ml bottle - Up to 5 inj available on a PSO15.00	1	Biomed

POTASSIUM CHI ORIDE

*	Inj 75 mg per ml,	10 ml	55.00	50	✓ AstraZeneca
---	-------------------	-------	-------	----	---------------

✓ Juno S29 ✓ Potassium Chloride

Aquettant S29

SODIUM BICARBONATE

Inj 8.4%, 50 ml	 1	9.95	1	Biomed
\ 11 · = · ·	 200			

a) Up to 5 inj available on a PSO b) Not in combination

Inj 8.4%, 100 ml......20.50

✓ Biomed

a) Up to 5 inj available on a PSO

b) Not in combination

SODIUM CHI ORIDE

Not funded for use as a nasal drop. Not funded for nebuliser use except when used in conjunction with an antibiotic intended for nebuliser use.

Inj 0.9%, bag - Up to 2000 ml available on a PSO	1.23	500 ml	Baxter
	1 26	1 000 ml	✓ Baxter

Only if prescribed on a prescription for renal dialysis, maternity or post-natal care in the home of the patient, or on a PSO for emergency use (500 ml and 1 000 ml nacks)

Inj 23.4% (4 mmol/ml), 20 ml ampoule	33.00	5	
For Sodium chloride oral liquid formulation refer Standar	rd Formulae, page 2	246	
Inj 0.9%, 5 ml ampoule - Up to 5 inj available on a PSO	2.80	20	
Inj 0.9%, 10 ml ampoule - Up to 5 inj available on a PSO	5.40	50	
Ini 0.9%. 20 ml ampoule	5.00	20	

1	Biomed	

✓ Fresenius Kabi

✓ Fresenius Kabi ✓ Fresenius Kabi

	Subsidy (Manufacturer's Price)	Sub Per	Fully osidised	Brand or Generic Manufacturer
TOTAL PARENTERAL NUTRITION (TPN) Infusion	CBS	1 OP	✓ T	

WATER

- On a prescription or Practitioner's Supply Order only when on the same form as an injection listed in the Pharmaceutical Schedule requiring a solvent or diluent; or
- 2) On a bulk supply order; or
- 3) When used in the extemporaneous compounding of eye drops; or
- 4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.

Inj 5 ml ampoule – Up to 5 inj available on a PSO7	7.00	50	✓ InterPharma
Inj 10 ml ampoule - Up to 5 inj available on a PSO6	6.63	50	✓ Pfizer
Inj 20 ml ampoule - Up to 5 inj available on a PSO	5.00	20	Fresenius Kabi
			✓ Multichem
7	7.50	30	✓ InterPharma

(InterPharma Inj 5 ml ampoule to be delisted 1 June 2021) (InterPharma Inj 20 ml ampoule to be delisted 1 June 2021)

Oral Administration		
CALCIUM POLYSTYRENE SULPHONATE Powder169.85	300 g OP	✓ Calcium Resonium
COMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO9.77	50	✓ <u>Electral</u>
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml)6.55	1,000 ml OP	✓ <u>Pedialyte -</u> Bubblegum
PHOSPHORUS		<u>Dabbiegum</u>
Tab eff 500 mg (16 mmol)82.50	100	Phosphate Phebra
POTASSIUM CHLORIDE		
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	60	Chlorvescent
* Tab long-acting 600 mg (8 mmol)8.90	200	✓ Span-K
SODIUM BICARBONATE		
Cap 840 mg8.52	100	✓ Sodibic✓ Sodibic
SODIUM POLYSTYRENE SULPHONATE		
Powder84.65	454 g OP	✓ Resonium-A

	CARDIOVASCULAR SYSTEM				
		Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	
A	lpha-Adrenoceptor Blockers				
Α	lpha Adrenoceptor Blockers				
DO	XAZOSIN				
*	Tab 2 mg	8.95	500	1	Apo-Doxazosin
*	Tab 4 mg	10.80	500	1	Apo-Doxazosin
РΗ	ENOXYBENZAMINE HYDROCHLORIDE				
*	Cap 10 mg	65.00	30	1	BNM S29
•	- Cap 10 11g	216.67	100		Dibenzyline S29
חח	AZOCINI	210.07	100	-	Disconzy into
	AZOSIN Tab 1 mg	5.52	100	1	Apo-Prazosin
	Tab 2 mg		100		Apo-Prazosin
	Tab 5 mg		100	_	Apo-Prazosin
	-		100	•	Apo i iuzoomi
	RAZOSIN - Subsidy by endorsement Subsidy by endorsement - Subsidised for patients who were	takina tarazasin pri	orto 1	Ootobor 3	1000 and the prescription is
	endorsed accordingly. Pharmacists may annotate the presc				
	dispensing of terazosin.	ription as chaolsca	WIICICI	ITICIC CAIS	is a record or prior
	Tab 2 mg	7.50	500	1	Apo-Terazosin
		14.20	28		Teva S29
	Tab 5 mg		500		Apo-Terazosin
		24.80	28		Teva S29
		00			
Α	gents Affecting the Renin-Angiotensin Systen	n			
	· · · · · · · · · · · · · · · · · · ·				
Α	CE Inhibitors				
0.4	DTODDII				
	PTOPRIL	04.00	95 ml C	n ./	Canatan
木	Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of age.	94.99	15 IIII C	P V	Capoten
	, ,				
	AZAPRIL Table 6.5 mm	0.00	00	,	7
	Tab 0.5 mg		90		Zapril
*	Tab 2.5 mg		90 90		Zapril Zapril
	Tab 5 mg	0.33	90	•	<u>Zaprii</u>
	ALAPRIL MALEATE	4.00	400		
	Tab 5 mg		100		Acetec
	Tab 10 mg		100		Acetec
	Tab 20 mg	2.42	100	•	Acetec
_	INOPRIL	0.5-			-
	Tab 5 mg		90		Ethics Lisinopril
	Tab 10 mg		90		Ethics Lisinopril
	Tab 20 mg	3.1/	90	•	Ethics Lisinopril
PE	RINDOPRIL	_	_	_	
	Tab 2 mg		30		Apo-Perindopril
	Tab 4 mg	4.80	30	/	Apo-Perindopril

* Tab 5 mg6.01

90

90

90

✓ Arrow-Quinapril 5

✓ Arrow-Quinapril 10

✓ Arrow-Quinapril 20

QUINAPRIL

	(CARDIO	VASC	ULAR SYSTEM
	Subsidy (Manufacturer's Price) \$	Subs	Fully sidised	Brand or Generic Manufacturer
ACE Inhibitors with Diuretics				
CILAZAPRIL WITH HYDROCHLOROTHIAZIDE — Subsidy b Subsidy by endorsement — Subsidised for patients who w 2020 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of cilazapril with hydro	vere taking cilazapril with macists may annotate th chlorothiazide.		tion as	
Table ing warmydroonloreanazide 12.0 mg		100		Hydrochlorothiazide
(Apo-Cilazapril/ Hydrochlorothiazide Tab 5 mg with hydrochlo QUINAPRIL WITH HYDROCHLOROTHIAZIDE	rothiazide 12.5 mg to be	e delisted 1	May 2	021)
Tab 10 mg with hydrochlorothiazide 12.5 mg		28	-	Accuretic
* Tab 20 mg with hydrochlorothiazide 12.5 mg	3.83 4.92	30 30		Accuretic 10 Accuretic 20
Angiotensin II Antagonists				
CANDESARTAN CILEXETIL				
* Tab 4 mg		90	_	Candestar
* Tab 8 mg		90	_	Candestar Candestar
* Tab 16 mg Tab 32 mg		90 90	_	Candestar
LOSARTAN POTASSIUM		00	• •	<u> </u>
* Tab 12.5 mg	1.56	84	√ L	osartan Actavis
* Tab 25 mg		84	_	osartan Actavis
* Tab 50 mg	2.25	84	✓ <u>L</u>	osartan Actavis
* Tab 100 mg	3.50	84	✓ L	osartan Actavis
Angiotensin II Antagonists with Diuretics				
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE Tab 50 mg with hydrochlorothiazide 12.5 mg		30	√ <u>Þ</u>	Arrow-Losartan & Hydrochlorothiazide
Angiotensin II Antagonists with Neprilysin In	hibitors			
SACUBITRIL WITH VALSARTAN – Special Authority see SA Note: Due to the angiotensin II receptor blocking activity ACE inhibitor or another ARB.	of sacubitril with valsart		d not be	e co-administered with an
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	√ E	intresto 24/26

ACE IIIIIDIO O GIOTICI ATID.			
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/103

⇒SA1905 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or

	Subsidy	Fu	,	Brand or
(Manuf	facturer's Price)	Subsidis	ed	Generic
	\$	Per	/	Manufacturer

continued...

- 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment: and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, page 120

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

AMIODARONE HYDROCHLORIDE	page .=e	
▲ Tab 100 mg	30	✓ Aratac
▲ Tab 200 mg	30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule – Up to 10 inj available on a PSO16.37	10	✓ Max Health
ATROPINE SULPHATE		
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a		
PSO12.07	10	✓ Martindale
DIGOXIN		
* Tab 62.5 mcg – Up to 30 tab available on a PSO	240	✓ Lanoxin PG
* Tab 250 mcg – Up to 30 tab available on a PSO	240	✓ Lanoxin
* Oral lig 50 mcg per ml	60 ml	✓ Lanoxin
		✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE		
▲ Cap 100 mg23.87	100	✓ Rythmodan
FLECAINIDE ACETATE	100	- rryumioudii
Tab 50 mg	60	✓ Flecainide BNM
▲ Cap long-acting 100 mg	90	✓ Flecainide Bivini
Out long dealing 100 mg	30	Controlled
		Release Teva
▲ Cap long-acting 200 mg61.06	90	✓ Flecainide
		Controlled
		Release Teva
Inj 10 mg per ml, 15 ml ampoule100.00	5	✓ Tambocor
MEXILETINE HYDROCHLORIDE		
▲ Cap 150 mg	100	✓ ANI S29
		✓ Mexiletine
		Hydrochloride
		USP S29
▲ Cap 250 mg202.00	100	✓ Mexiletine
		Hydrochloride
		USP S29
PROPAFENONE HYDROCHLORIDE		
▲ Tab 150 mg40.90	50	✓ Rytmonorm

		CARDIO	ASCL	JLAR SYSTEM
	Subsidy (Manufacturer's Pri	ce) Subs Per	Fully idised	Brand or Generic Manufacturer
Antihypotensives				
MIDODRINE – Special Authority see SA1474 below – Retail pl Tab 2.5 mg Tab 5 mg Tab 5 mg Initial application from any relevant practitioner. Approvals va not due to drugs. Note: Treatment should be started with small doses and titrate the usual target is a standing systolic blood pressure of 90 mm Renewal from any relevant practitioner. Approvals valid for 2 y benefiting from treatment.	53.00 79.00 alid for 2 years where d upwards as neces Hg.	sary. Hypert	ension s	ntron g orthostatic hypotension hould be avoided, and
Beta-Adrenoceptor Blockers				
Beta Adrenoceptor Blockers				
ATENOLOL * Tab 50 mg * Tab 100 mg * Oral liq 25 mg per 5 ml	7.30	500 500 300 ml OP	✓ My ✓ Ato ✓ Ato	rlan Atenolol rlan Atenolol enolol AFT enolol AFT S29 S29
Restricted to children under 12 years of age.				
BISOPROLOL FUMARATE * Tab 2.5 mg Bisoprolol Mylan to be Sole Supply on 1 April 2021	1.84 3.53	90		soprolol Mylan osvate
* Tab 5 mg Bisoprolol Mylan to be Sole Supply on 1 April 2021	2.55 5.15	90		soprolol Mylan svate
* Tab 10 mg	3.62 9.40	90		soprolol Mylan osvate

CELIPROLOL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking celiprolol prior to 1 October 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of celiprolol.

	3			
*	Tab 200 mg	21.40	180	Celol

(Celol Tab 200 mg to be delisted 1 April 2021)

✓ Carvedilol Sandoz

✓ Carvedilol Sandoz✓ Carvedilol Sandoz

60

60

		Subsidy		Fully	
		(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
	BETALOL	<u> </u>		-	Marialactaror
*	Tab 100 mg	14 50	100	1	Trandate
*	Tab 200 mg		100		Trandate
*	Inj 5 mg per ml, 20 ml ampoule		5	•	<u>Trandate</u>
-1-	inj o mg por mi, 20 mi ampodio	(88.60)	Ü		Trandate
*	inj 5 mg per ml, 20 ml vial	` ,	1		Trandato
•••	", o mg por m, 20 m na	(48.20)	•		Alvogen \$29
ME	TOPROLOL SUCCINATE	, ,			•
*	Tab long-acting 23.75 mg	1.45	30	1	Betaloc CR
*	Tab long-acting 47.5 mg	1.43	30	1	Betaloc CR
*	Tab long-acting 95 mg		30	1	Betaloc CR
*	Tab long-acting 190 mg		30	✓	Betaloc CR
ME	TOPROLOL TARTRATE				
*	Tab 50 mg	5.66	100	✓	Apo-Metoprolol
*	Tab 100 mg	7.55	60	✓	Apo-Metoprolol
*	Tab long-acting 200 mg	23.40	28	✓	Slow-Lopresor
*	Inj 1 mg per ml, 5 ml vial	29.50	5	•	Metroprolol IV
					<u>Mylan</u>
NA	DOLOL				
*	Tab 40 mg	16.69	100	✓	Apo-Nadolol
*	Tab 80 mg	26.43	100	•	Apo-Nadolol
PIN	IDOLOL				
*	Tab 5 mg	13.22	100	✓	Apo-Pindolol
*	Tab 10 mg	23.12	100	✓	Apo-Pindolol
*	Tab 15 mg	33.31	100	✓	Apo-Pindolol
PR	OPRANOLOL				
*	Tab 10 mg	4.64	100	1	Apo-Propranolol
*	Tab 40 mg		100		Apo-Propranolol
*	Cap long-acting 160 mg		100		Cardinol LA
*	Oral liq 4 mg per ml - Special Authority see SA1327 below -				
	Retail pharmacy		500 m	nl 🗸	Roxane S29
L.,	CA1207 Chariel Authority for Cubaidy				

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

S	O	ΓΑ	LC)L

*	Tab 80 mg	32.58	500	✓ Mylan
	Tab 160 mg		100	✓ Mylan
TIN	MOLOL			
*	Tab 10 mg	10.55	100	✓ Apo-Timol

Subsidy (Manufacturer's Price) \$ Per

Fully Subsidised Per Brand or Generic Manufacturer

Calcium Channel Blockers

AMI ODIDINIE

Dihydropyridine Calcium Channel Blockers

MLODIPINE Tab 2.5 mg	90	✓ Vasorex
1.72	100	✓ Apo-Amlodipine
16.20	28	✓ Bristol \$29
Tab 5 mg0.96	90	✓ Vasorex
1.56	28	✓ Sandoz S29
		✓ Teva S29
3.33	250	✓ Apo-Amlodipine
Tab 10 mg1.19	90	✓ Vasorex
1.66	28	✓ Sandoz S29
4.40	250	✓ Apo-Amlodipine
Apo-Amlodipine Tab 2.5 mg to be delisted 1 June 2021)		. ,
Apo-Amlodipine Tab 5 mg to be delisted 1 June 2021)		
Apo-Amlodipine Tab 10 mg to be delisted 1 June 2021)		
ELODIPINE		
₭ Tab long-acting 2.5 mg1.45	30	✓ Plendil ER
★ Tab long-acting 5 mg3.93	90	✓ Felo 5 ER
★ Tab long-acting 10 mg4.32	90	✓ Felo 10 ER
IIFEDIPINE		
★ Tab long-acting 10 mg10.63	60	✓ Adalat 10
		✓ Adefin S29
18.80	56	✓ Tensipine MR10 S2
₭ Tab long-acting 20 mg17.72	100	✓ Nyefax Retard
★ Tab long-acting 30 mg3.14	30	✓ Adalat Oros
34.10	100	✓ Mylan S29
Fab long-acting 60 mg5.67	30	✓ Adalat Oros
		Adefin XL
52.81	100	✓ Mylan (\$29)
Adalat 10 Tab long-acting 10 mg to be delisted 1 August 2021)		-
Adefin S29 Tab long-acting 10 mg to be delisted 1 August 2021)		
Adalat Oros Tab long-acting 30 mg to be delisted 1 August 2021)		

Other Calcium Channel Blockers

(Adalat Oros Tab long-acting 60 mg to be delisted 1 August 2021) (Adefin XL Tab long-acting 60 mg to be delisted 1 August 2021)

DIL	TIAZEM HYDROCHLORIDE			
*	Tab 30 mg	4.60	100	✓ Dilzem
*	Tab 60 mg	8.50	100	✓ Dilzem
*	Cap long-acting 120 mg	33.42	500	✓ Apo-Diltiazem CD
	Cap long-acting 180 mg		500	✓ Apo-Diltiazem CD
	Cap long-acting 240 mg		500	✓ Apo-Diltiazem CD
PΕ	RHEXILINE MALEATE			
*	Tab 100 mg	62.90	100	✓ Pexsig

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	
	(Manufacturer's Price)) Per	Subsidised	Generic Manufacturer
ERAPAMIL HYDROCHLORIDE				
₭ Tab 40 mg	7.01	100		Isoptin
├ Tab 80 mg	11.74	100	•	Isoptin
Fab long-acting 120 mg	36.02	100		Isoptin Retard \$29
K. Talalana astina 040 mm	45.40	00		Isoptin SR
★ Tab long-acting 240 mg★ Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a	15.12	30	•	Isoptin SR
PSOPSO	25.00	5	1	Isoptin
Controlly Acting Agents				•
Centrally-Acting Agents				
CLONIDINE	40.04			
Patch 2.5 mg, 100 mcg per day — Only on a prescription		4		Mylan Mylan
Patch 5 mg, 200 mcg per day — Only on a prescription		4 4		Mylan Mylan
Patch 7.5 mg, 300 mcg per day — Only on a prescription	16.93	4	•	<u>Mylan</u>
CLONIDINE HYDROCHLORIDE ├ Tab 25 mcg	Q 75	112	1	Clonidine BNM
₭ Tab 150 mcg		100		Catapres
light is the light in the light is the light is the light is the light is the light in the light is the light is the light is the light is the light in the ligh		100	_	Medsurge
METHYLDOPA	25.50	10	•	<u>weasurge</u>
К Таb 250 mg	15 10	100	1	Methyldopa Mylan
140 200 mg	52.85	500		Methyldopa Mylan
	02.00	000	•	S29 S29
				329 329
51				329 329
Diuretics				323 323
Diuretics Loop Diuretics				323 929
Loop Diuretics				323 929
Loop Diuretics BUMETANIDE	4.91	30		Burinex S29 S29
Loop Diuretics BUMETANIDE K Tab 1 mg	16.36	100	1	Burinex S29 S29 Burinex
Loop Diuretics BUMETANIDE * Tab 1 mg	16.36		1	Burinex S29 S29
Loop Diuretics BUMETANIDE * Tab 1 mg	16.36	100	1	Burinex S29 S29 Burinex
Loop Diuretics BUMETANIDE * Tab 1 mg * Inj 500 mcg per ml, 4 ml vial BUROSEMIDE [FRUSEMIDE] * Tab 40 mg - Up to 30 tab available on a PSO	16.36 7.95	100 5 1,000	<i>'</i>	Burinex S29 S29 Burinex Burinex Apo-Furosemide
Loop Diuretics BUMETANIDE * Tab 1 mg * Inj 500 mcg per ml, 4 ml vial BUROSEMIDE [FRUSEMIDE] * Tab 40 mg — Up to 30 tab available on a PSO * Tab 500 mg	16.36 7.95 7.24 25.00	100 5 1,000 50	<i>y y y</i>	Burinex S29 S29 Burinex Burinex Apo-Furosemide Urex Forte
Loop Diuretics BUMETANIDE * Tab 1 mg * Inj 500 mcg per ml, 4 ml vial FUROSEMIDE [FRUSEMIDE] * Tab 40 mg — Up to 30 tab available on a PSO * Tab 500 mg * Oral liq 10 mg per ml	16.36 7.95 7.24 25.00 11.20 3	100 5 1,000 50 50 ml C		Burinex S29 S29 Burinex Burinex Apo-Furosemide Urex Forte Lasix
Loop Diuretics BUMETANIDE Tab 1 mg Inj 500 mcg per ml, 4 ml vial BUROSEMIDE [FRUSEMIDE] Tab 40 mg - Up to 30 tab available on a PSO Tab 500 mg Oral liq 10 mg per ml Inj 10 mg per ml, 25 ml ampoule	16.36 7.95 7.24 25.00 11.20 3	100 5 1,000 50 50 ml C		Burinex S29 S29 Burinex Burinex Apo-Furosemide Urex Forte Lasix Lasix
Loop Diuretics BUMETANIDE K Tab 1 mg K Inj 500 mcg per ml, 4 ml vial BUROSEMIDE [FRUSEMIDE] K Tab 40 mg — Up to 30 tab available on a PSO K Tab 500 mg K Oral liq 10 mg per ml K Inj 10 mg per ml, 25 ml ampoule	16.36 7.95 7.24 25.00 11.20 3	100 5 1,000 50 50 ml C		Burinex S29 S29 Burinex Burinex Apo-Furosemide Urex Forte Lasix Lasix Frusemide-Claris
Loop Diuretics FUMETANIDE For Tab 1 mg	16.36 7.95 7.24 25.00 11.20 3 60.65 PSO1.15	100 5 1,000 50 50 ml C		Burinex S29 S29 Burinex Burinex Apo-Furosemide Urex Forte Lasix Lasix
Loop Diuretics BUMETANIDE K Tab 1 mg	16.36 7.95 7.24 25.00 11.20 3 60.65 PSO1.15	100 5 1,000 50 50 ml C		Burinex S29 S29 Burinex Burinex Apo-Furosemide Urex Forte Lasix Lasix Frusemide-Claris
Loop Diuretics BUMETANIDE * Tab 1 mg	16.36 7.95 7.24 25.00 11.20 3 60.65 PSO1.15	100 5 1,000 50 50 ml C		Burinex S29 S29 Burinex Burinex Apo-Furosemide Urex Forte Lasix Lasix Frusemide-Claris
Loop Diuretics BUMETANIDE * Tab 1 mg	16.36 7.95 7.24 25.00 11.20 60.65 'SO1.15	100 5 1,000 50 50 ml C		Burinex S29 S29 Burinex Burinex Apo-Furosemide Urex Forte Lasix Lasix Frusemide-Claris
Loop Diuretics BUMETANIDE K Tab 1 mg	16.36 7.95 24 25.00 11.20 3 60.65 PSO1.15 15 	100 5 1,000 50 80 ml C 6 5		Burinex S29 S29 Burinex Burinex Apo-Furosemide Urex Forte Lasix Lasix Frusemide-Claris Furosemide-Baxter
Loop Diuretics BUMETANIDE * Tab 1 mg * Inj 500 mcg per ml, 4 ml vial FUROSEMIDE [FRUSEMIDE] * Tab 40 mg - Up to 30 tab available on a PSO * Tab 500 mg * Oral liq 10 mg per ml	16.36 7.95 7.24 25.00 11.20 3 60.65 PSO1.15 1 March 2021)	100 5 1,000 50 80 ml C 6 5		Burinex S29 S29 Burinex Burinex Apo-Furosemide Urex Forte Lasix Lasix Frusemide-Claris Furosemide-Baxter

	Subsidy (Manufacturer's Pri	ice) Sub	Fully	Brand or Generic Manufacturer
	\$	Per		Manutacturer
⇒SA1728 Special Authority for Subsidy				
nitial application from any relevant practitioner. Approx	vals valid without further re	enewal unles	s notifie	d for applications meet
ne following criteria:		511011ai ai ii 00		a 101 appiloanono moon
Both:				
1 Patient has heart failure with ejection fraction less2 Either:	than 40%; and			
2.1 Patient is intolerant to optimal dosing of spi2.2 Patient has experienced a clinically signific		on optimal do	sing of s	spironolactone.
METOLAZONE				
Tab 5 mg	CBS	1	✓ N	letolazone S29
		50	✓ Z	aroxolyn S29
PIRONOLACTONE				
€ Tab 25 mg		100		piractin
← Tab 100 mg		100 25 ml OP		piractin iomed
Oral liq o mg per mi		23 1111 01	· <u>-</u>	nomeu
Potassium Sparing Combination Diuretics	3			
MILORIDE HYDROCHLORIDE WITH FUROSEMIDE				
* Tab 5 mg with furosemide 40 mg	8.63	28	√ F	rumil
MILORIDE HYDROCHLORIDE WITH HYDROCHLORO	THIAZIDE			
Fab 5 mg with hydrochlorothiazide 50 mg	5.00	50	✓ N	loduretic
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]				
Tab 2.5 mg - Up to 150 tab available on a PSO	20.00	500	✓ <u>A</u>	rrow- Bendrofluazide
May be supplied on a PSO for reasons other tha				

* Tab 2.5 mg – Up to 150 tab available on a PSO20.00	0 500	✓ <u>Arrow-</u> <u>Bendrofluazide</u>
May be supplied on a PSO for reasons other than emergency. * Tab 5 mg34.59	5 500	✓ <u>Arrow-</u> Bendrofluazide
CHLOROTHIAZIDE Oral liq 50 mg per ml	0 25 ml OP	✓ Biomed
Tab 25 mg3.90		✓ Igroton S29 ✓ <u>Hygroton</u>

Lipid-Modif	ying .	Agents
-------------	--------	--------

	ra	

INDAPAMIDE

BE	ZAFIBRATE			
*	Tab 200 mg19	.01	90 🗸	['] Bezalip
	Tab long-acting 400 mg12		30	Bezalip Retard

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

✓ Dapa-Tabs

90

(Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	
Other Lipid-Modifying Agents				
CIPIMOX € Cap 250 mg	21.56	30		Olbetam Olbetam S29 S29
ICOTINIC ACID				
Tab 50 mg		100		Apo-Nicotinic Acid
Tab 500 mg Apo-Nicotinic Acid Tab 50 mg to be delisted 1 May 2021) Apo-Nicotinic Acid Tab 500 mg to be delisted 1 May 2021)	17.89	100	•	Apo-Nicotinic Acid
Resins				
OLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g	28.60	30	•	Colestid
HMG CoA Reductase Inhibitors (Statins)				
TORVASTATIN				
← Tab 10 mg	6.96	500	1	Lorstat
€ Tab 20 mg		500	✓	Lorstat
€ Tab 40 mg		500		Lorstat
€ Tab 80 mg	27.19	500	•	Lorstat
← Tab 10 mg	3 55	28	/	Pravastatin Mylan
€ Tab 20 mg		28		Pravastatin Mylan
	4.72	100		Apo-Pravastatin
Pravastatin Mylan to be Sole Supply on 1 April 2021 Tab 40 mg	2.61	28		Pravastatin Mylan
1 ab 40 mg	8.06	100		Apo-Pravastatin
Pravastatin Mylan to be Sole Supply on 1 April 2021	0.00	100	•	Apo-Fiavasiaiiii
Pravastatin Mylan Tab 10 mg to be delisted 1 April 2021)				
Apo-Pravastatin Tab 20 mg to be delisted 1 April 2021)				
Apo-Pravastatin Tab 40 mg to be delisted 1 April 2021)				
IMVASTATIN				
€ Tab 10 mg	1.23	90	1	Simvastatin Mylan
€ Tab 20 mg		90		Simvastatin Mylan
F Tab 40 mg		90		Simvastatin Mylan
F Tab 80 mg	7.12	90	•	Simvastatin Mylan
Selective Cholesterol Absorption Inhibitors				
ZETIMIBE - Special Authority see SA1045 below - Retail pharm	•			
← Tab 10 mg	1.95	30	/	Ezetimibe Sandoz

1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and

continued...

All of the following:

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
 \$	Per	•	Manufacturer

continued...

- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 x normal) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atoryastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 below - Retail pharmacy

Tab 10 mg with simvastatin 10 mg	5.15	30	✓ Zimybe
Tab 10 mg with simvastatin 20 mg	6.15	30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg		30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg	8.15	30	✓ Zimybe

⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

GLYCERYL TRINITRATE

*	Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	4.45	250 dose OP	✓ Nitrolingual Pump
				Spray
*	Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
	Patch 50 mg, 10 mg per day		30	✓ Nitroderm TTS
ISC	SORBIDE MONONITRATE			
*	Tab 20 mg	19.55	100	✓ Ismo 20
*	Tab long-acting 40 mg	8.20	30	✓ Ismo 40 Retard
*	Tab long-acting 60 mg	9.25	90	✓ <u>Duride</u>

OATIBIO VACCCEATI CTOTEM			
	Subsidy (Manufacturer's Price) \$	Subsidis Per	ully Brand or sed Generic Manufacturer
Sympathomimetics			
ADRENALINE			
Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PS	O4.98	5	✓ Aspen Adrenaline
, ,, , ,	10.76		✓ DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a F	PSO27.00	5	✓ Hospira
	49.00	10	Aspen Adrenaline
ISOPRENALINE [ISOPROTERENOL]			
* Inj 200 mcg per ml, 1 ml ampoule	36.80	25	
	(164.20)		Isuprel
(Isuprel Inj 200 mcg per ml, 1 ml ampoule to be delisted 1 Febru	ary 2021)		
Vasodilators			
HYDRALAZINE HYDROCHLORIDE			
* Tab 25 mg - Special Authority see SA1321 below - Retail			
pharmacy	CBS	1	✓ Hydralazine
priamacy			✓ Onelink \$29
			✓ AMDIPHARM \$29
			✓ Onelink \$29
* Inj 20 mg ampoule	25 90		✓ Apresoline
⇒SA1321 Special Authority for Subsidy	20.00	J	Apresonne
Initial application from any relevant practitioner. Approvals val	id without further renev	wal unlace no	atified for applications meeting
the following criteria:	ia without further refle	wai uiliess iid	nilled for applications meeting
Either:			
1 For the treatment of refractory hypertension; or			
2 For the treatment of heart failure in combination with a ni	trate, in patients who a	re intolerant	or have not responded to ACE
inhibitors and/or angiotensin receptor blockers.	•		•
MINOXIDIL			
▲ Tab 10 mg	70.00	100	✓ Loniten
NICORANDIL			
▲ Tab 10 mg	25.57	60	✓ <u>lkorel</u>
▲ Tab 20 mg			✓ <u>lkorel</u>
PAPAVERINE HYDROCHLORIDE			
* Inj 12 mg per ml, 10 ml ampoule	217.90	5	✓ Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]			
Tab 400 mg	42 26	50	✓ Trental 400
145 455 frig		00	Tremai 400
Endothelin Receptor Antagonists			
AMBRISENTAN - Special Authority see SA1702 on the next pa	ge - Retail pharmacy		
Tab 5 mg	1,550.00	30	✓ Ambrisentan Mylan
-	4,585.00		✓ Volibris
Ambrisentan Mylan to be Sole Supply on 1 March 2021			
Tab 10 mg			✓ Ambrisentan Mylan
Ancheiranten Mulanta ha Cala Cumplu an 1 Mayala 0001	4,585.00	,	✓ Volibris

(Volibris Tab 5 mg to be delisted 1 March 2021) (Volibris Tab 10 mg to be delisted 1 March 2021)

Ambrisentan Mylan to be Sole Supply on 1 March 2021

Reddy's

Subsidy (Manufacturer's	Full s Price) Subsidise	
\$	Per •	Manufacturer

⇒SA1702 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

BOSENTAN - Special Authority see SA1908 below - Retail pharmacy

 Tab 62.5 mg
 141.00
 60
 ✓ Bosentan Dr Reddy's

 Tab 125 mg
 141.00
 60
 ✓ Bosentan Dr

⇒SA1908 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Fither:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil; or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV: or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV: or
 - 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see SA1909 below – Retail p	oharmacy		
Tab 25 mg	0.64	4	✓ Vedafil
Tab 50 mg	0.64	4	✓ Vedafil
Tab 100 mg	6.60	12	✓ Vedafil

⇒SA1909 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II: or
 - 3.2 PAH is in NYHA/WHO functional class III: or
 - 3.3 PAH is in NYHA/WHO functional class IV: and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and

Subsid		Fully	Brand or
(Manufacturer		ubsidised	Generic
\$	Per	•	Manufacturer

continued...

- 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
- 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age, or health system capacity constraints.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Prostacyclin Analogues

EPOPROSTENOL - Special Authority see SA1696 below - Retail pharmacy		
Inj 500 mcg vial36.61	1	✓ Veletri
Inj 1.5 mg vial73.21	1	✓ Veletri

⇒SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz ILOPROST - Special Authority see SA1705 below - Retail pharmacv Nebuliser soln 10 mcg per ml, 2 ml740.10

30 Ventavis

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz



Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 90

ADAPAI FNF

IS

a) Maximum of 30 g per prescription

h'	Only	i on a	prescri	ntion
v	, OHIII	v un a	DIESCII	DUUII

b) Only on a procomption			
Crm 0.1%	22.89	30 g OP	Differin
Gel 0.1%	22.89		✓ Differin
SOTRETINOIN - Special Authority see SA1475 below - F	Retail pharmacy		
Cap 5 mg	8.14	60	Oratane
Cap 10 mg	13.34	120	✓ Oratane
Can 20 mg	20 49	120	✓ Oratane

⇒SA1475 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
 - 3.2 Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is female and has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of one month after the completion of the treatment; or
- 2 Patient is male.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

TRFTINOIN

Crm 0.5 mg per g − Maximum of 50 g per prescription13.90 50 g OP ✓ ReTrieve

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 90

HYDROGEN PEROXIDE

111	DITOGEN I ETIONIDE		
*	Crm 1%8.56	10 g OP	Crystaderm
		15 g OP	 Crystaderm

		_	
	Subsidy (Manufacturer's I \$	Price) Subs	Fully Brand or sidised Generic ✓ Manufacturer
MUPIROCIN Oint 2%	6.60	15 g OP	Bactroban
a) Only on a prescriptionb) Not in combination	(/		
SODIUM FUSIDATE [FUSIDIC ACID] Crm 2%	1.59	5 g OP	✓ <u>Foban</u>
 a) Maximum of 5 g per prescription b) Only on a prescription c) Not in combination 			
Oint 2%a) Maximum of 5 g per prescription	1.59	5 g OP	✓ <u>Foban</u>
b) Only on a prescription c) Not in combination			
SULFADIAZINE SILVER Crm 1%	10.80	50 g OP	✓ Flamazine
a) Up to 250 g available on a PSOb) Not in combination			
Antifungals Topical			
For systemic antifungals, refer to INFECTIONS, Antifung AMOROLFINE a) Only on a prescription b) Not in combination	gals, page 97		
Nail soln 5%	14.93	5 ml OP	✓ MycoNail
CICLOPIROX OLAMINE a) Only on a prescription b) Not in combination			
Nail-soln 8% CLOTRIMAZOLE	5.72	7 ml OP	✓ Apo-Ciclopirox
** Crm 1% a) Only on a prescription b) Not in combination	0.70	20 g OP	✓ Clomazol
* Soln 1%	4.36 (7.55)	20 ml OP	Canesten
a) Only on a prescriptionb) Not in combination			
ECONAZOLE NITRATE Crm 1%	1.00 (7.48)	20 g OP	Pevaryl
a) Only on a prescriptionb) Not in combination	, ,		•
Foaming soln 1%, 10 ml sachets	9.89 (17.23)	3	Pevaryl
a) Only on a prescriptionb) Not in combination			

DERMATOLOGICALS

	Subsidy (Manufacturer's Pr \$	rice) Subs Per	Fully sidised	Brand or Generic Manufacturer
MICONAZOLE NITRATE * Crm 2% a) Only on a prescription	0.81	15 g OP	✓ M	ultichem
b) Not in combination c) Multichem to be Sole Supply on 1 February 2021 * Lotn 2%	4.36 (10.03)	30 ml OP	D	aktarin
b) Not in combination * Tinct 2%	4.36 (12.10)	30 ml OP	D	aktarin
a) Only on a prescriptionb) Not in combination				
Antipruritic Preparations				
CALAMINE a) Only on a prescription b) Not in combination Crm, aqueous, BP	1.26	100 g	√ <u>h</u>	ealthE Calamine Aqueous Cream BP
CROTAMITON a) Only on a prescription b) Not in combination Crm 10%	3.29	20 g OP	√ <u>lt</u>	ch-Soothe
MENTHOL - Only in combination 1) Only in combination with a dermatological base or pro 2) With or without other dermatological galenicals.	prietary Topical Co	orticosteriod –	Plain	
Crystals	6.92 29.60	25 g 100 g		idWest idWest
Corticosteroids Topical				
For systemic corticosteroids, refer to CORTICOSTEROIDS AND	RELATED AGEN	NTS, page 80		
Corticosteroids - Plain				
BETAMETHASONE DIPROPIONATE Crm 0.05%	2.96	15 g OP	✓ D	iprosone

BETAMETHASONE DIPROPIONATE			
Crm 0.05%	2.96	15 g OP	✓ Diprosone
	36.00	50 g OP	Diprosone
Diprosone to be Sole Supply on 1 February 2021			
Oint 0.05%	2.96	15 g OP	Diprosone
	36.00	50 g OP	Diprosone
Diprosone to be Sole Supply on 1 February 2021			
Oint 0.05% in propylene glycol base	4.33	30 g OP	Diprosone OV
BETAMETHASONE VALERATE			
* Crm 0.1%	3.45	50 g OP	✓ Beta Cream
* Oint 0.1%	3.45	50 g OP	✓ Beta Ointment
* Lotn 0.1%	18.00	50 ml OP	✓ Betnovate

	Subsidy		Fully Brand or
	(Manufacturer's P		sidised Generic
	\$	Per	✓ Manufacturer
CLOBETASOL PROPIONATE			
* Crm 0.05%	2.18	30 g OP	✓ Dermol
* Oint 0.05%	2.12	30 g OP	✓ Dermol
		9	
CLOBETASONE BUTYRATE	Г 00	00 = OD	
Crm 0.05%		30 g OP	F
	(10.00)		Eumovate
DIFLUCORTOLONE VALERATE			
Fatty oint 0.1%	8.97	50 g OP	
	(15.86)		Nerisone
(Nerisone Fatty oint 0.1% to be delisted 1 August 2021)			
HYDROCORTISONE			
* Crm 1% – Only on a prescription	3.70	100 g OP	✓ Hydrocortisone
The Office of a proportion in the proportion in		100 g O1	(PSM)
	17.15	E00 a	✓ Hydrocortisone
	17.15	500 g	
# D O ' ' '	40.05	0.5	(PSM)
* Powder – Only in combination		25 g	✓ ABM
Up to 5% in a dermatological base (not proprietary Topic	cal Corticosteriod	d – Plain) with o	or without other dermatological
galenicals			
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN			
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only of	on		
a prescription		250 ml	✓ DP Lotn HC
		200	<u> </u>
HYDROCORTISONE BUTYRATE	0.05	100 = OD	/ Lassid Lineauseus
Lipocream 0.1%		100 g OP	✓ Locoid Lipocream
Oint 0.1%		100 g OP	Locoid
Milky emul 0.1%	13.70	100 ml OP	✓ Locoid Crelo
METHYLPREDNISOLONE ACEPONATE			
Crm 0.1%	4.46	15 g OP	✓ Advantan
Oint 0.1%	4.46	15 g OP	✓ Advantan
MOMETASONE FUROATE			
Crm 0.1%	1 51	15 g OP	✓ Elocon Alcohol Free
G111 G17 /	2.50	50 g OP	✓ Elocon Alcohol Free
Oint 0.1%		15 g OP	✓ Elocon
Olit 0.170	2.90	50 g OP	✓ Elocon
Lotn 0.1%		30 ml OP	✓ Elocon
	0.00	00 1111 01	Liocom
TRIAMCINOLONE ACETONIDE		400 00	
Crm 0.02%		100 g OP	✓ <u>Aristocort</u>
Oint 0.02%	6.35	100 g OP	✓ Aristocort
A			
Corticosteroids - Combination			
DETAMETUAÇONE VALEDATE MITU OLIOOUINOL			
BETAMETHASONE VALERATE WITH CLIOQUINOL — Only on		45 = OD	
Crm 0.1% with clioquinol 3%		15 g OP	Data susta O
(Data and a O Oma O 40) with all and 1000 to 1 life in the	(4.90)		Betnovate-C
(Betnovate-C Crm 0.1% with clioquinol 3% to be delisted 1 June	2021)		
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FU	SIDIC ACID]		
Crm 0.1% with sodium fusidate (fusidic acid) 2%	3.49	15 g OP	
, ,	(10.45)	<u> </u>	Fucicort
a) Maximum of 15 g per prescription	. ,		
b) Only on a prescription			
A = A = - a b = - a			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

DERMATOLOGICALS

	Subsidy (Manufacturer's F \$	Price) Subsi Per	Fully Brand or dised Generic ✓ Manufacturer
HYDROCORTISONE WITH MICONAZOLE - Only on a presci ★ Crm 1% with miconazole nitrate 2%		15 g OP	✓ Micreme H
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN – Crm 1% with natamycin 1% and neomycin sulphate 0.5%. Oint 1% with natamycin 1% and neomycin sulphate 0.5%	3.35	-	✓ Pimafucort ✓ Pimafucort
FRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMY Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 u and gramicidin 250 mcg per g – Only on a prescription	mg	ГIN 15 g ОР	Viaderm KC
Barrier Creams and Emollients			
Barrier Creams			
DIMETHICONE * Crm 5% pump bottle	4.48	500 ml OP	✓ <u>healthE</u> Dimethicone 5%
* Crm 10% pump bottle	4.52	500 ml OP	✓ healthE Dimethicone 10%
ZINC AND CASTOR OIL * Oint	4.25	500 g	✓ Boucher
Emollients			
AQUEOUS CREAM * Crm CETOMACROGOL	1.92	500 g	✓ <u>Boucher</u>
* Crm BPCETOMACROGOL WITH GLYCEROL	2.48	500 g	✓ <u>healthE</u>
Crm 90% with glycerol 10%	2.35	500 ml OP	✓ ADE ✓ Boucher
	3.10	1,000 ml OP	✓ Kenkay Sorbolene✓ ADE✓ Boucher
EMULSIFYING OINTMENT * Oint BP	3.40	500 g	✓ Emulsifying
Fundation Circumst ADE to be Oak Ourselver A Mar	3.59		Ointment ADE ✓ AFT
Emulsifying Ointment ADE to be Sole Supply on 1 Mar (AFT Oint BP to be delisted 1 March 2021)	rcn 2021		
DIL IN WATER EMULSION * Crm	2.19	500 g	✓ O/W Fatty Emulsion Cream
PARAFFIN Oint liquid paraffin 50% with white soft paraffin 50%	5.35	500 ml OP	✓ <u>healthE</u>
JREA * Crm 10%	1.37	100 g OP	✓ healthE Urea Cream

				▝
	Subsidy		Fully Brand or	
	(Manufacturer's F		sidised Generic	
	\$	Per	✓ Manufacturer	
WOOL FAT WITH MINERAL OIL – Only on a prescription				
* Lotn hydrous 3% with mineral oil	5.60	1,000 ml		
·	(11.95)		DP Lotion	
	1.40	250 ml OP		
	(4.53)		DP Lotion	
	5.60	1,000 ml		
	(20.53)		Alpha-Keri Lotion	
	(23.91)		BK Lotion	
	1.40	250 ml OP		
	(7.73)		BK Lotion	
Other Dermatological Bases				
PARAFFIN				
White soft - Only in combination	4.99	450 g	✓ healthE	
•	19.99	2,500 g	✓ healthE	
Only in combination with a dermatological galenical or	as a diluent for a p	, ,		

Minor Skin Infections

POVIDONE IODINE			
Oint 10%	7.40	65 g OP	✓ <u>Betadine</u>
a) Maximum of 130 g per prescription			
b) Only on a prescription			
Antiseptic Solution 10%	2.55	100 ml	✓ Riodine
Antiseptic soln 10%	3.83	15 ml	✓ Riodine
	5.40	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
	(3.48)		Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	1.63	100 ml	
	(7.78)		Pfizer

Parasiticidal Preparations

DIMETHICONE

* Lotn 4%	4.98	200 ml OP	✓ <u>healthE</u> <u>Dimethicone 4%</u> <u>Lotion</u>
IVERMECTIN - Special Authority see SA1225 below - F	letail pharmacy		
Tab 3 mg - Up to 100 tab available on a PSO	17.20	4	✓ Stromectol

1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.

- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA1225 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy: or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria: Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	,	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — **(Other parasitic infections)** only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

PERMETHRIN

Ρ

Crm 5% 5.75 Lotn 5% 3.99		✓ <u>Lyderm</u> ✓ <u>A-Scabies</u>
PHENOTHRIN		
Shampoo 0.5%	200 ml OP	✓ Parasidose

Psoriasis and Eczema Preparations

ACITRETIN - Special Authority see SA1476 below - Retail pl	narmacy		
Cap 10 mg	17.86	60	✓ Novatretin
Cap 25 mg	41.36	60	✓ Novatretin

⇒SA1476 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Fither:
 - 3.1 Patient is female and has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of two years after the completion of the treatment; or
 - 3.2 Patient is male.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is female and has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and the applicant has ensured that the possibility of pregnancy has been excluded prior to the commencement of the treatment and that the patient is informed that she must not become pregnant during treatment and for a period of two years after the completion of the treatment: or
- 2 Patient is male.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIO

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL			
Foam spray 500 mcg with calcipotriol 50 mcg per g	59.95	60 g OP	Enstilar
Gel 500 mcg with calcipotriol 50 mcg per g	52.24	60 g OP	✓ Daivobet
Oint 500 mcg with calcipotriol 50 mcg per g	19.95	30 g OP	Daivobet
CALCIPOTRIOL			
Oint 50 mcg per g	40.00	120 g OP	Daivonex
COAL TAR			
Soln BP - Only in combination	36.25	200 ml	✓ <u>Midwest</u>

- 1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

DERMATOLOGICALS

	Subsidy (Manufacturerla Dri	aa\ Cuba	Fully Brand or sidised Generic
	(Manufacturer's Pri \$	Per Subs	sidised Generic Manufacturer
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SUL	_PHUR		
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% at			
allantoin crm 2.5%		75 g OP	
	(8.00) 3.43	30 g OP	Egopsoryl TA
	(4.35)	00 g O1	Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR			
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97 7.95	25 g OP 40 g OP	✓ Coco-Scalp✓ Coco-Scalp
PIMECROLIMUS - Special Authority see SA1970 below - Reta	ail pharmacy		
a) Maximum of 15 g per prescription			
b) Note: a maximum of 15 g per prescription and no more Cream 1%		on per 12 we 15 g OP	eks. ✓ Elidel
Elidel to be Sole Supply on 1 March 2021	20.50	15 y OF	Eliuei
⇒SA1970 Special Authority for Subsidy			
Initial application only from a dermatologist, paediatrician, oph	thalmologist or any	relevant prac	ctitioner on the recommendation
of a dermatologist, paediatrician or ophthalmologist. Approvals	valid without furthe	r renewal unl	less notified for applications
meeting the following criteria:			
Both: 1 Patient has atopic dermatitis on the eyelid; and			
Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure.			
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUOR	ESCEIN – Only on	a prescription	n
* Soln 2.3% with trolamine laurilsulfate and fluorescein sodium		500 ml	✓ Pinetarsol
SALICYLIC ACID			
Powder - Only in combination	18.88	250 g	✓ Midwest✓ PSM
 Only in combination with a dermatological base of With or without other dermatological galenicals. 	r proprietary Topica	l Corticostero	oid – Plain or collodion flexible
2) Will of Willout office defination global galerinodio.			
SULPHUR			
Precipitated – Only in combination	6.35	100 g	✓ Midwest
1) Only in combination with a dermatological base of	r proprietary Topica	l Corticostero	oid – Plain
2) With or without other dermatological galenicals.			
Scalp Preparations			
BETAMETHASONE VALERATE			

BETAMETHASONE VALERATE * Scalp app 0.1%	7.75	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE * Scalp app 0.05%	5.69	30 ml OP	✓ <u>Dermol</u>
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%	7.30	100 ml OP	✓ Locoid
KETOCONAZOLE Shampoo 2%	3.23	100 ml OP	✓ Sebizole
a) Maximum of 100 ml per prescription			

b) Only on a prescription

DERMATOLOGICALS

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 69

IMIQUIMOD

PODOPHYLLOTOXIN

a) Maximum of 3.5 ml per prescription

b) Only on a prescription

Other Skin Preparations

Antineoplastics
FLUOROURACIL SODIUM

GENITO-URINARY SYSTEM

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Brand or

Generic

Manufacturer

Subsidy Fully
(Manufacturer's Price) Subsidised
\$ Per

Contraceptives - Non-hormonal

Condoms

-	IDOMS			
	49 mm - Up to 144 dev available on a PSO		144	✓ <u>Moments</u>
	53 mm		10	✓ <u>Moments</u>
	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	11.64	144	✓ <u>Moments</u>
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO		, .	
	53 mm, 0.05 mm thickness		10	✓ <u>Moments</u>
	\	11.42	144	✓ <u>Moments</u>
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	2.05	40	/ 11
	53 mm, chocolate, brown		10	✓ Moments
	-\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	11.64	144	✓ <u>Moments</u>
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	0.05	40	/ Name : :- te
	53 mm, strawberry, red		10	✓ Moments
	.) 11-1-00 1	11.64	144	✓ <u>Moments</u>
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	2.27	40	/ 11
	56 mm		10	✓ Moments
		11.64	144	✓ <u>Moments</u>
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
	56 mm, 0.05 mm thickness		12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	56 mm, 0.05mm thickness (bulk pack)	14.61	144	✓ Gold Knight
	 a) Maximum of 60 dev per prescription 			
	b) Up to 60 dev available on a PSO			_
	56 mm, 0.08 mm thickness		10	✓ Moments
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	56 mm, 0.08 mm thickness, red		10	✓ Moments
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	56 mm, chocolate		12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	56 mm, strawberry	1.30	12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	60 mm	1.42	12	✓ Gold Knight XL
		14.87	144	✓ Shield XL
		17.02		Gold Knight XL

GENITO-URINARY SYSTEM

b) Up to 60 dev available on a PSO

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
*	60 mm (bulk pack)	14.87	144	√ <u>G</u>	Gold Knight XL
	a) Maximum of 60 dev per prescription				

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

*	IUD 29.1 mm length × 23.2 mm width	1	✓ Choice TT380 Short
	IUD 33.6 mm length × 29.9 mm width	1	✓ Choice
	•		TT380 Standard
*	IUD 35.5 mm length × 19.6 mm width15.50	1	✓ Choice Load 375

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

*	Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to	0		
	84 tab available on a PSO	19.80	84	✓ Mercilon 28
*	Tab 30 mcg with desogestrel 150 mcg and 7 inert tab	6.62	84	
		(19.80)		Marvelon 28

- a) Higher subsidy of \$13.80 per 84 tab with Special Authority see \$A0500 above
- b) Up to 84 tab available on a PSO

				_
	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Su	bsidised	Generic
	\$	Per	✓	Manufacturer
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO	2.18	84	✓ M	licrogynon 20 ED
•	6.45	112		emme-Tab ED
* Tab 50 mcg with levonorgestrel 125 mcg and 7 inert tab -	Up			
to 84 tab available on a PSO	9.45	84	✓ M	licrogynon 50 ED
* Tab 30 mcg with levonorgestrel 150 mcg	6.62	63		•
	(16.50)		M	licrogynon 30
a) Higher subsidy of \$15.00 per 63 tab with Special Au	thority see SA0500 on	the pre	vious pag	е
b) Up to 63 tab available on a PSO	,	•		
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO		84	✓ L	evlen ED
	6.45	112	✓ F	emme-Tab ED
ETHINYLOESTRADIOL WITH NORETHISTERONE				
Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up	to			
84 tab available on a PSO		84	✓ B	revinor 1/28
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab -	Un		_	
to 84 tab available on a PSO		84		econ orimin

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

	ONONGEOTHEE			
*	Tab 30 mcg - Up to 84 tab available on a PSO	16.50	84	✓ Microlut
		22.00	112	✓ Microlut
*	Subdermal implant (2 × 75 mg rods) - Up to 3 pack available			
	on a PSO1	06.92	1	✓ <u>Jadelle</u>

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
MEDROXYPROGESTERONE ACETATE Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PS	SO7.98	1	✓ [Depo-Provera
NORETHISTERONE Tab 350 mcg - Up to 84 tab available on a PSO	6.25	84	√ V	loriday 28
Emergency Contraceptives				

Į	_E	۷	OI	١	Ю	R	G	E٥	ST	R	Е	L

- Tab 1.5 mg4.95 ✓ Postinor-1
 - a) Maximum of 2 tab per prescription
 - b) Up to 5 tab available on a PSO
 - c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

*	Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up			
	to 168 tab available on a PSO	4.98	168	Ginet
	Ginet to be Sole Supply on 1 April 2021			

Gynaecological Anti-infectives

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID
Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate

0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43 (24.00)	100 g OP	Aci-Jel
CLOTRIMAZOLE * Vaginal crm 1% with applicators2.50	35 g OP	✓ Clomazol
* Vaginal crm 2% with applicators3.00 MICONAZOLE NITRATE	20 g OP	✓ <u>Clomazol</u>
* Vaginal crm 2% with applicator	40 g OP	✓ <u>Micreme</u>
Vaginal crm 100,000 u per 5 g with applicator(s)4.00	75 g OP	✓ Nilstat

Myometrial and Vaginal Hormone Preparations

ERGOMETRINE MALEATE			
Inj 500 mcg per ml, 1 ml ampoule - Up to 5 inj available on a			
PSO	105.00	5	 DBL Ergometrine
OESTRIOL			·
* Crm 1 mg per g with applicator	6.62	15 g OP	✓ Ovestin
* Pessaries 500 mcg	6.86	15	✓ Ovestin
OXYTOCIN – Up to 5 inj available on a PSO			
Inj 5 iu per ml, 1 ml ampoule	3.98	5	Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule	4.98	5	✓ Oxytocin BNM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
OXYTOCIN WITH ERGOMETRINE MALEATE — Up to 5 inj ava Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml		5	√ <u>s</u>	yntometrine	
Pregnancy Tests - hCG Urine					
PREGNANCY TESTS - HCG URINE					

- a) Up to 200 test available on a PSO
- b) Only on a PSO

40 test OP

✓ David One Step Cassette **Pregnancy Test**

✓ Smith BioMed Rapid **Pregnancy Test**

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 108

5-Alpha Reductase Inhibitors

FINASTERIDE - Special Authority see SA0928 below - Retail pharmacy ✓ Ricit 100 Ricit to be Sole Supply on 1 April 2021

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Fither:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy ✓ Tamsulosin-Rex

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

OXYBUTYNIN * Tab 5 mg	500 473 ml	✓ Apo-Oxybutynin ✓ Apo-Oxybutynin
POTASSIUM CITRATE Oral liq 3 mmol per ml — Special Authority see SA1083 on the next page — Retail pharmacy	200 ml OP	✓ <u>Biomed</u>

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

GENITO-URINARY SYSTEM

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ubsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE * Grans eff 4 g sachets	2.22	28	✓ <u>Ural</u>
SOLIFENACIN SUCCINATE			
Tab 5 mg	3.00	30	✓ Solifenacin Mylan
Tab 10 mg	5.50	30	✓ Solifenacin Mylan

Detection of Substances in Urine

ORTHO-TOLIDINE			
* Compound diagnostic sticks	7.50	50 test OP	
	(8.25)		Hemastix
TETRABROMOPHENOL			
* Blue diagnostic strips	7.02	100 test OP	
-	(13.92)		Albustix

Obstetric Preparations

Antiprogesterones

MIFEPRISTONE

Subsidised on a PSO only if from a Family Planning New Zealand Clinic or an abortion service provider with a DHB contract and the PSO is endorsed with the name of the institution for which the PSO is required.

Mifegyne

- a) Up to 15 tab available on a PSO
- b) Only on a PSO

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Por	_	Manufacturer	

Calcium Homeostasis

$C\Delta I$		

★ Inj 100 iu per ml, 1 ml ampoule121.00
5
✓ Miacalcic

CINACALCET - Special Authority see SA1618 below - Retail pharmacy

Tab 30 mg − Wastage claimable210.30 28 ✓ Sensipar

⇒SA1618 Special Authority for Subsidy

Initial application only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial − Special Authority see SA1687 below −
Retail pharmacy......38.03 1

✓ Zoledronic acid
Mylan

⇒SA1687 Special Authority for Subsidy

Initial application — **(bone metastases)** only from an oncologist, haematologist or palliative care specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
 - 2.1 Patient has bone metastases or involvement; and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
 - 3.1 Patient has bone metastases or involvement; and
 - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer) only from an oncologist or medical practitioner on the recommendation of a oncologist. Approvals valid for 2 years for applications meeting the following criteria:
All of the following:

Subsid	dy Full	/ Brand or
(Manufacture	r's Price) Subsidise	d Generic
\$	Per 💌	Manufacturer

continued...

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.

Corticosteroids and Related Agents for Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETA	ATE	
* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml19.20	5	
(36.96)		Celestone
,		Chronodose
DEXAMETHASONE		
* Tab 0.5 mg - Up to 60 tab available on a PSO	30	✓ Dexmethsone
5 1	30	
* Tab 4 mg – Up to 30 tab available on a PSO		✓ <u>Dexmethsone</u>
Oral liq 1 mg per ml45.00	25 ml OP	✓ Biomed
DEXAMETHASONE PHOSPHATE		
Dexamethasone phosphate injection will not be funded for oral use.		
* Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO9.25	10	✓ <u>Dexamethasone</u>
		<u>Phosphate</u>
		<u>Panpharma</u>
* Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO16.37	10	✓ <u>Dexamethasone</u>
		<u>Phosphate</u>
		<u>Panpharma</u>
FLUDROCORTISONE ACETATE		
* Tab 100 mcg	100	✓ Florinef
HYDROCORTISONE	100	✓ Davidas
* Tab 5 mg	100	✓ <u>Douglas</u>
* Tab 20 mg	100	✓ <u>Douglas</u>
* Inj 100 mg vial5.30	1	✓ Solu-Cortef
a) Up to 5 inj available on a PSO		
b) Only on a PSO		
METHYLPREDNISOLONE		
* Tab 4 mg112.00	100	✓ Medrol
* Tab 100 mg	20	✓ Medrol
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)		
Inj 40 mg vial18.90	1	✓ Solu-Medrol-Act-
IIIJ 40 IIIY Vidi10.90	ı	
		<u>O-Vial</u>
Inj 125 mg vial28.90	1	✓ Solu-Medrol-Act-
iiij 120 iiig vidi20.90	'	0-Vial
		U-VIAI
Inj 500 mg vial22.78	1	✓ Solu-Medrol-Act-
111 500 111g viai	'	0-Vial
		<u>O-Viai</u>
Inj 1 g vial27.83	1	✓ Solu-Medrol
. •	•	o doid illiculor
METHYLPREDNISOLONE ACETATE	_	4
Inj 40 mg per ml, 1 ml vial44.40	5	✓ Depo-Medrol
PREDNISOLONE		
* Oral liq 5 mg per ml - Up to 30 ml available on a PSO	30 ml OP	✓ Redipred
Restricted to children under 12 years of age.		_
, ,		

				_
	Subsidy		Fully Brand or	
	(Manufacturer's Price)	_	Subsidised Generic	
	\$	Per	r 🗸 Manufacturer	
PREDNISONE				
* Tab 1 mg	10.68	500	✓ Apo-Prednisone	
* Tab 2.5 mg		500	✓ Apo-Prednisone	
* Tab 5 mg - Up to 30 tab available on a PSO	11.09	500	✓ Apo-Prednisone	
* Tab 20 mg - Up to 30 tab available on a PSO	29.03	500	✓ Apo-Prednisone	
TETRACOSACTRIN			·	
* Inj 250 mcg per ml, 1 ml ampoule	75.00	1	✓ UK Synacthen S29	
, , ,			✓ AU Synacthen	
			✓ Synacthen	
* Inj 1 mg per ml, 1 ml ampoule	690.00	1	✓ Synacthen Depot	
, , ,			✓ Synacthene	
			Retard \$29	
TRIAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule	20.80	5	✓ Kenacort-A 10	
ing to mg per mi, i mi ampoule		J	-	
Kanasant A 40 ta ba Oala Oanabaan 4 Annil 2004	26.62		✓ Adcortyl S29	
Kenacort-A 10 to be Sole Supply on 1 April 2021			_	
Inj 40 mg per ml, 1 ml ampoule		1	✓ Triaver S29	
	51.10	5	Kenacort-A 40	
	70.62		✓ Kenalog S29	
Kenacort-A 40 to be Sole Supply on 1 April 2021			-	

Sex Hormones Non Contraceptive

Androgen Agonists and Antagonists

CVDDOTEDONE ACETATE

CYPROTERONE ACETATE			
Tab 50 mg	13.17	50	✓ Siterone
Tab 100 mg	26.75	50	✓ Siterone
TESTOSTERONE			
Patch 5 mg per day	90.00	30	✓ Androderm
TESTOSTERONE CIPIONATE			
Inj 100 mg per ml, 10 ml vial	76.50	1	✓ Depo-Testosterone
TESTOSTERONE ESTERS			
Inj 250 mg per ml, 1 ml	12.98	1	 Sustanon Ampoules
TESTOSTERONE UNDECANOATE			
Cap 40 mg	21.00	60	Andriol Testocaps
Inj 250 mg per ml, 4 ml vial		1	✓ Reandron 1000

Hormone Replacement Therapy - Systemic

Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

	Subsidy (Manufacturer's Price	e) Sub	Fully	Brand or Generic
	\$	Per	✓	Manufacturer
Oestrogens				
OESTRADIOL - See prescribing guideline on the previous page	ge			
* Tab 1 mg	4.12	28 OP		
	(11.10)		Е	Estrofem
* Tab 2 mg	4.12	28 OP		
	(11.10)		Е	Estrofem
* Patch 100 mcg per 24 hours	7.91	4	✓ (Climara
a) No more than 1 patch per week				
b) Only on a prescription				
* Patch 50 mcg per 24 hours	7.04	4	✓ (Climara
a) No more than 1 patch per week				
b) Only on a prescription				
Patch 25 mcg per day	6.12	8	√ E	stradot
	7.85			stradiol TDP
				Mylan S29
a) No more than 2 patch per week				my an -
b) Only on a prescription				
Patch 50 mcg per day	7.04	8	./ 5	stradot 50 mcg
r atcir 50 meg per day	9.22	U		stradiol TDP
	3.22		٠.	
) N				Mylan S29
a) No more than 2 patch per week				
b) Only on a prescription	7.04	0		
Patch 75 mcg per day	7.91	8	•	stradot
a) No more than 2 patch per week				
b) Only on a prescription		_		
Patch 100 mcg per day	7.91	8	✓ E	stradot
 a) No more than 2 patch per week 				
b) Only on a prescription				
OESTRADIOL VALERATE - See prescribing guideline on the	previous page			
* Tab 1 mg		84	✓ P	Progynova
* Tab 2 mg	12.36	84	_	Progynova
OESTROGENS - See prescribing guideline on the previous pa			_	
* Conjugated, equine tab 300 mcg		28		
* Conjugated, equine tab 500 meg	(13.50)	20	Р	remarin
* Conjugated, equine tab 625 mcg		28		Tomami
* Conjugated, equilie tab 025 meg	(13.50)	20	Р	remarin
	(13.30)		'	Terriariii
Progestogens				
MEDROXYPROGESTERONE ACETATE - See prescribing gu	uideline on the previou	us page		
* Tab 2.5 mg		30	✓ P	Provera
* Tab 5 mg	14.00	100	✓ P	Provera
* Tab 10 mg	7.15	30	√ P	Provera

	Subsidy (Manufacturer's Price) Su Per	Fully Brand or ibsidised Generic Manufacturer
Progestogen and Oestrogen Combined Prepa	rations		
OESTRADIOL WITH NORETHISTERONE - See prescribing	guideline on page 81		
* Tab 1 mg with 0.5 mg norethisterone acetate		28 OP	LCP:
* Tab 2 mg with 1 mg norethisterone acetate	(18.10)	28 OP	Kliovance
* Tab 2 mg with 1 mg norethisterone acetate	(18.10)	20 OF	Kliogest
* Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	` ,		· ·····ges
oestradiol tab (12) and 1 mg oestradiol tab (6)		28 OP	
	(18.10)		Trisequens
Other Oestrogen Preparations			
ETHINYLOESTRADIOL * Tab 10 mcg	17.60	100	✓ NZ Medical and
Tab to mog	17.00	100	Scientific
OESTRIOL			
* Tab 2 mg	7.00	30	✓ Ovestin
Other Progestogen Preparations			
LEVONORGESTREL			
* Intra-uterine device 52 mg	269 50	1	✓ Mirena
* Intra-uterine device 13.5 mg		1	✓ Jaydess
MEDROXYPROGESTERONE ACETATE			
Tab 100 mg	101.00	100	✓ Provera HD
NORETHISTERONE			
* Tab 5 mg - Up to 30 tab available on a PSO	18.29	100	✓ <u>Primolut N</u>
PROGESTERONE			
Cap 100 mg - Special Authority see SA1609 below - Reta			4
pharmacy	16.50	30	Utrogestan

⇒SA1609 Special Authority for Subsidy

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 For the prevention of pre-term labour*; and
- 2 Either:
 - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 For the prevention of pre-term labour*; and
- 2 Treatment is required for second or subsequent pregnancy; and
- 3 Either:
 - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are unapproved indications.

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
Thyroid and Antithyroid Agents			
CARBIMAZOLE * Tab 5 mg	10.80	100	✓ AFT Carbimazole \$29 ✓ Neo-Mercazole ✓ Neo-Mercazole S29 \$29
(AFT Carbimazole S29 Tab 5 mg to be delisted 1 March 2021)			
LEVOTHYROXINE * Tab 25 mcg		90	✓ Synthroid
* Tab 50 mcg	5.79	28 90 1.000	✓ Mercury Pharma✓ Synthroid✓ Eltroxin
* Tab 100 mcg	1.78 6.01	28 90 1.000	Mercury PharmaSynthroid
PROPYLTHIOURACIL – Special Authority see SA1199 below – Propylthiouracil is not recommended for patients under the treatments are contraindicated.	- Retail pharmacy	,	
Tab 50 mg	35.00	100	✓ PTU S29
⇒SA1199 Special Authority for Subsidy			

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SC	MATROPIN (OMNITROPE) - Special Authority see SA	1629 below – Retail pharm	acy	
*	Inj 5 mg cartridge	34.88	i	Omnitrope
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
	Inj 15 mg cartridge		1	✓ Omnitrope

⇒SA1629 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months

Subsidy	/ Fully	y Brand or
(Manufacturer's		
\$	Per ✓	Manufacturer

continued...

using the standards of Tanner and Davies (1985); and

- 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
- 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
- 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

ibsidy turer's Price) Subs	Fully	Brand or Generic
 \$ Per	✓	Manufacturer

continued...

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — (**short stature due to chronic renal insufficiency**) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Fither:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months...

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
 - 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
 - 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
 - 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
 - 5 No malignancy has developed after growth hormone therapy was commenced; and
 - 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per	✓	Manufacturer	
				_

continued...

Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and

- 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
- 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients.

GnRH Analogues

\sim	Q.	= [- 1	N	

Implant 3.6 mg, syringe	65.68	1	✓ Teva
1 0, 7 0	66.48		✓ Zoladex
Implant 10.8 mg, syringe	122.37	1	✓ Teva
	177.50		✓ Zoladex

(Zoladex Implant 3.6 mg, syringe to be delisted 1 May 2021) (Zoladex Implant 10.8 mg, syringe to be delisted 1 May 2021)

LEUPRORELIN

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

Inj 3.75 mg prefilled dual chamber syringe - Higher subsider	dy of		
\$221.60 per 1 inj with Endorsement	66.48	1	
	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe - Higher subs	sidy		

of \$591.68 per 1 inj with Endorsement.......177.50 (591.68)Lucrin Depot 3-month

Vasopressin Agonists

DE	SMOPRESSIN		
	Wafer 120 mcg47.00	30	Minirin Melt
DE	SMOPRESSIN ACETATE		
	Tab 100 mcg25.00	30	✓ Minirin
	Tab 200 mcg54.45	30	✓ Minirin
\blacktriangle	Nasal drops 100 mcg per ml39.03	2.5 ml OP	✓ Minirin
\blacktriangle	Nasal spray 10 mcg per dose27.95	6 ml OP	✓ Desmopressin-
			PH&T
	Inj 4 mcg per ml, 1 ml67.18	10	✓ Minirin

Other Endocrine Agents

CABERGOLINE

		Tab 0.5 mg – Maximum of 2 tab per prescription; can be
✓ Dostines	2	waived by Special Authority see SA1370 on the next page3.75
✓ Dostine:	8	15.20

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	1	Manufacturer	

⇒SA1370 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 pathological hyperprolactinemia; or
- 2 acromegaly*.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

CLOMIFENE CITRATE Tab 50 mg29.84	10	✓ Mylan Clomiphen \$29
DANAZOL		
Cap 100 mg19.13	3 28	✓ Mylan S29
Cap 200 mg97.83	3 100	✓ Azol
(Mylan S29 Cap 100 mg to be delisted 1 April 2021) (Azol Cap 200 mg to be delisted 1 April 2021)		
, ,		
METYRAPONE Cap 250 mg558.00	50	✓ <u>Metopirone</u>

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	oubsidised ✓	Manufacturer	

Anthelmintics

		E – Special Authority see SA1318 below – Retail pharmacy	ALBENDAZOLE – S
✓ Eskazole S29	60	J469.20	Tab 400 mg

⇒SA1318 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

Tab 100 mg	7.97	6	✓ Vermox
	24.19	24	✓ De-Worm
Oral lig 100 mg per 5 ml	2.18	15 ml	
	(7.53)		Vermox
(De-Worm Tab 100 mg to be delisted 1 March 2021)	,		
PRAZIQUANTEL			
Tab 600 mg	68.00	8	 Biltricide

Antibacterials

- a) For topical antibacterials, refer to DERMATOLOGICALS, page 62
- b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 239

Cephalosporins and Cephamycins

CEFACLOR MONOHYDRATE			
Cap 250 mg	24.70	100	✓ Ranbaxy-Cefaclor
Grans for oral liq 125 mg per 5 ml - Wastage claimable	3.53	100 ml	✓ Ranbaxy-Cefaclor
CEFALEXIN			
Cap 250 mg	3.33	20	✓ Cephalexin ABM
			✓ Ibilex S29
Cap 500 mg	3.95	20	 Cephalexin ABM
Grans for oral liq 25 mg per ml - Wastage claimable	8.75	100 ml	✓ Cefalexin Sandoz
Grans for oral liq 50 mg per ml - Wastage claimable	11.75	100 ml	✓ Cefalexin Sandoz
(Ibilex S29 Cap 250 mg to be delisted 1 February 2021)			
CEFAZOLIN - Subsidy by endorsement			
Only if prescribed for dialysis or cellulitis in accordance with a accordingly.	DHB approved	protocol and t	the prescription is endorsed
Inj 500 mg vial	3 30	5	✓ AFT
Inj 1 g vial		5	✓ AFT
IIIJ I Y VIAI	3.49	J	* <u>ALI</u>

CEFTRIAXONE - Subsidy by endorsement

- a) Up to 10 inj available on a PSO
- b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of pelvic inflammatory disease, or the treatment of suspected meningococcal disease, and the prescription or PSO is endorsed accordingly.

Inj 500 mg vial	0.89	1	✓ Ceftriaxone-AFT
Ini 1 g vial	3 00	5	✓ Ceftriaxone-AFT

CEFUROXIME AXETIL - Subsidy by endorsement

Only if prescribed for prophylaxis of endocarditis and the prescription is endorsed accordingly. 50 Zinnat Tab 250 mg45.93

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidise	ed Generic	
\$	Per	 Manufacturer 	

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO		2	✓ Apo-Azithromycin
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage			
claimable	14.38	15 ml	✓ Zithromax

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Atypical mycobacterial infection; or
 - 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN (AS LACTOBIONATE) Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	✓ E-Mycin
a) Up to 20 tab available on a PSO b) Up to 2 x the maximum PSO quantity for RFPP			·
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	E-Mycin
a) Up to 300 ml available on a PSOb) Up to 2 x the maximum PSO quantity for RFPPc) Wastage claimable			
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	✓ E-Mycin
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
ERYTHROMYCIN STEARATE			
Tab 250 mg - Up to 30 tab available on a PSO		100	
	(22.29)		ERA
Tab 500 mg		100	
	(44.58)		ERA
ROXITHROMYCIN			
Tab disp 50 mg	8.29	10	✓ Rulide D
Restricted to children under 12 years of age.			
Tab 150 mg	8.28	50	✓ <u>Arrow-</u> <u>Roxithromycin</u>
Tab 300 mg	16.33	50	✓ <u>Arrow-</u> Roxithromycin

	Subsidy (Manufacturer's Prio \$	ce) Subs Per	Fully Brand or sidised Generic Manufacturer
Penicillins			
AMOXICILLIN			
Cap 250 mg	22.50	500	✓ <u>Alphamox</u>
a) Up to 30 cap available on a PSO			
b) Up to 10 x the maximum PSO quantity for RFPP			
Cap 500 mg	36.98	500	✓ <u>Alphamox</u>
a) Up to 30 cap available on a PSO			
b) Up to 10 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml	1.40	100 ml	✓ Alphamox 125
a) Up to 200 ml available on a PSO	1.40	100 1111	Alphamox 125
b) Wastage claimable			
Grans for oral liq 250 mg per 5 ml	1.73	100 ml	✓ Alphamox 250
a) Up to 300 ml available on a PSO	1.70	100 1111	Alphaniox 200
b) Up to 10 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
Inj 250 mg vial	10.67	10	✓ Ibiamox
Inj 500 mg vial	12.41	10	✓ Ibiamox
Inj 1 g vial – Up to 5 inj available on a PSO	17.29	10	✓ Ibiamox
AMOXICILLIN WITH CLAVULANIC ACID			
Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab			
available on a PSO	1.88	20	✓ Augmentin
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25	mg		•
per ml		100 ml	Augmentin
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5			
per ml – Up to 200 ml available on a PSO	2.20	100 ml OP	✓ Curam
BENZATHINE BENZYLPENICILLIN			
Inj 900 mg (1.2 million units) in 2.3 ml syringe - Up to 5 inj			
available on a PSO	344.93	10	✓ Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]			
Inj 600 mg (1 million units) vial - Up to 5 inj available on a P	SO 11.09	10	✓ Sandoz
FLUCLOXACILLIN			
Cap 250 mg - Up to 30 cap available on a PSO	16.83	250	✓ Staphlex
Cap 500 mg - Up to 30 cap available on a PSO		500	✓ Staphlex
Grans for oral liq 25 mg per ml	2.29	100 ml	✓ <u>AFT</u>
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
Grans for oral liq 50 mg per ml	3.68	100 ml	✓ <u>AFT</u>
a) Up to 200 ml available on a PSO			
b) Wastage claimable	0.00	40	/ Floring
Inj 250 mg vial		10	✓ Fluctoxin
Inj 500 mg vialInj 1 g vial – Up to 5 inj available on a PSO		10 5	✓ Flucloxin✓ Flucil
inj i g viai – up to 5 inj avaliable on a PSO	3./0	b	▼ <u>FIUCII</u>

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

PHENOXYMETHYLPENICILLIN (PENICILLIN V) Cap 250 mg — Up to 30 cap available on a PSO		Subsidy (Manufacturer's Price) \$	Sub:	Fully sidised	Generic
Cap 500 mg	PHENOXYMETHYLPENICILLIN (PENICILLIN V)				
a) Up to 20 cap available on a PSO b) Up to 2 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml	Cap 250 mg - Up to 30 cap available on a PSO	2.59	50		
b) Up to 2 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml		4.26	50	✓	Cilicaine VK
b) Up to 2 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml	a) Up to 20 cap available on a PSO				
a) Up to 200 ml available on a PSO b) Wastage claimable Grans for oral liq 250 mg per 5 ml					
b) Wastage claimable Grans for oral liq 250 mg per 5 ml	Grans for oral liq 125 mg per 5 ml	2.99	100 ml	1	<u>AFT</u>
Grans for oral liq 250 mg per 5 ml	a) Up to 200 ml available on a PSO				
a) Up to 300 ml available on a PSO b) Up to 2 x the maximum PSO quantity for RFPP c) Wastage claimable PROCAINE PENICILLIN Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO	b) Wastage claimable				
a) Up to 300 ml available on a PSO b) Up to 2 x the maximum PSO quantity for RFPP c) Wastage claimable PROCAINE PENICILLIN Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO	Grans for oral liq 250 mg per 5 ml	3.99	100 ml	1	<u>AFT</u>
c) Wastage claimable PROCAINE PENICILLIN Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO					
c) Wastage claimable PROCAINE PENICILLIN Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO	b) Up to 2 x the maximum PSO quantity for RFPP				
Inj 1.5 g in 3.4 ml syringe − Up to 5 inj available on a PSO					
Inj 1.5 g in 3.4 ml syringe − Up to 5 inj available on a PSO	PROCAINE PENICII I IN				
Tetracyclines DOXYCYCLINE * Tab 100 mg − Up to 30 tab available on a PSO		123.50	5	1	Cilicaine
* Tab 100 mg − Up to 30 tab available on a PSO	Tetracyclines				
MINOCYCLINE HYDROCHLORIDE * Tab 50 mg − Additional subsidy by Special Authority see SA1355 below − Retail pharmacy	DOXYCYCLINE				
 * Tab 50 mg - Additional subsidy by Special Authority see SA1355 below - Retail pharmacy	* Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	/	Doxine
* Tab 50 mg - Additional subsidy by Special Authority see SA1355 below - Retail pharmacy 5.79 60 (12.05) Mino-tabs * Cap 100 mg 19.32 100 (52.04) Minomycin SA1355 Special Authority for Manufacturers Price Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea. TETRACYCLINE - Special Authority see SA1332 below - Retail pharmacy Tab 250 mg 28 ✓ Accord \$29	MINOCYCLINE HYDROCHLORIDE				
SA1355 below – Retail pharmacy					
** Cap 100 mg		5 70	60		
** Cap 100 mg	OA 1000 below Tietali pharmacy		00		Mino-tahs
(52.04) Minomycin SA1355 Special Authority for Manufacturers Price Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea. TETRACYCLINE – Special Authority see SA1332 below – Retail pharmacy Tab 250 mg	* Can 100 mg		100		Willio tabo
▶SA1355 Special Authority for Manufacturers Price Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea. TETRACYCLINE – Special Authority see SA1332 below – Retail pharmacy Tab 250 mg			100		Minomycin
Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea. TETRACYCLINE - Special Authority see SA1332 below - Retail pharmacy Tab 250 mg	- CA10EE Charles Authority for Manufacturers Dries	(02.01)			Will Corry Corr
rosacea. TETRACYCLINE - Special Authority see SA1332 below - Retail pharmacy Tab 250 mg21.42 28 ✓ Accord \$29		id without further ren	owel unles	n notif	iad whara the nationt had
TETRACYCLINE − Special Authority see SA1332 below − Retail pharmacy Tab 250 mg21.42 28 ✓ Accord \$29		ia williout furtifier ferit	ewai uilles	5 HOUI	ieu where the patient has
Tab 250 mg21.42 28 ✓ Accord 529		il pharmaoy			
	, ,		00	./	Accord Coo
	•	21.42	28	•	ACCORD 829
		id for 3 months for ap	plications	meetir	ng the following criteria:
Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:	Both:				

Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 62

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg - Up to 5 tab available on a PSO	2.42	28	✓ Cipflox
Tab 500 mg - Up to 5 tab available on a PSO	3.40	28	✓ Cipflox
Tab 750 mg	5.95	28	✓ Cipflox

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
CLINDAMYCIN				
Cap hydrochloride 150 mg	4.61	24	1	Dalacin C
Inj phosphate 150 mg per ml, 4 ml ampoule		10	1	Dalacin C
COLISTIN SULPHOMETHATE – Retail pharmacy-Specialist – S Only if prescribed for dialysis or cystic fibrosis patient and the Inj 150 mg	e prescription is endo			∕. Colistin-Link
GENTAMICIN SULPHATE				
Inj 10 mg per ml, 1 ml ampoule – Subsidy by endorsement	25.00	5	1	DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.			infection a	and the prescription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement	182.00	10	1	Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.		tract	infection a	and the prescription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement	17.50	10	1	Pfizer
	87.50	50	✓	Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient or endorsed accordingly.	or complicated urinary	tract	infection a	and the prescription is
MOXIFLOXACIN – Special Authority see SA1740 below – Retail No patient co-payment payable	pharmacy			
Tab 400 mg	42.00	5	•	<u>Avelox</u>
⇒SA1740 Special Authority for Subsidy				

⇒SA1740 Special Authority for Subsidy

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Active tuberculosis*; and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications;
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 1 Has nucleio
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

	Subsidy (Manufacturer's Price \$) Sub Per	Fully sidised	Brand or Generic Manufacturer
PAROMOMYCIN – Special Authority see SA1689 below – Retail Cap 250 mg		16	✓ H	umatin S29
■ SA1689 Special Authority for Subsidy Initial application only from an infectious disease specialist, clini month for applications meeting the following criteria: Either:	cal microbiologist o	r gastroen	terologist	. Approvals valid for 1
Patient has confirmed cryptosporidium infection; or For the eradication of Entamoeba histolyica carriage. Renewal only from an infectious disease specialist, clinical microl applications meeting the following criteria:	biologist or gastroer	nterologist.	Approv	als valid for 1 month for
Either: 1 Patient has confirmed cryptosporidium infection; or 2 For the eradication of Entamoeba histolyica carriage.				
PYRIMETHAMINE – Special Authority see SA1328 below – Reta Tab 25 mg		30	✓ D	araprim S29
 SA1328 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following: For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months. 	a period of 3 montl		ss notified	d for applications meeting
SODIUM FUSIDATE [FUSIDIC ACID] Tab 250 mg	34.50	12	✓ F	ucidin
SULFADIAZINE SODIUM - Special Authority see SA1331 below Tab 500 mg		56	✓ W	ockhardt §29
 SA1331 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following: For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months or the second content of the pregnancy. 	a period of 3 montl		ss notified	d for applications meeting
TOBRAMYCIN Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement	15 00	5	√ T	obramycin Mylan
Only if prescribed for dialysis or cystic fibrosis patient and Solution for inhalation 60 mg per ml, 5 ml – Subsidy by		-		
endorsement	395.00 2,200.00	56 dose	✓ T	obramycin BNM OBI
 a) Wastage claimable b) Only if prescribed for a cystic fibrosis patient and the (TOBI Solution for inhalation 60 mg per ml, 5 ml to be delisted 1 M TRIMETHOPRIM 		rsed accor	rdingly.	
* Tab 300 mg – Up to 30 tab available on a PSO	16.50	50	✓ <u>T</u>	MP

	Subsidy (Manufacturer's Pric \$	e) Subs	Fully sidised	Brand or Generic Manufacturer
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX	AZOLE]			
* Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - L to 30 tab available on a PSO		500	✓ T	risul
* Oral liq 8 mg sulphamethoxazole 40 mg per ml - Up to 200 i available on a PSO		100 ml	✓ D	eprim
VANCOMYCIN – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient or for	r prophylaxis of end	locarditis or	for treat	ment of Clostridium
difficile following metronidazole failure and the prescription is Inj 500 mg vial	endorsed according		✓ <u>M</u>	lylan_

Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, page 63
- b) For topical antifungals refer to GENITO URINARY, page 76

FLUCONAZOLE

2.75	28	Mylan
0.65	1	✓ Mylan
12.89	28	✓ Mylan
ty.		-
109.34	35 ml	Diflucan
	0.65 12.89 y	

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOLE

Cap 100 mg4.27	15	✓ <u>Itrazole</u>
Oral liq 10 mg per ml - Special Authority see SA1322 on the		
next page – Retail pharmacy141.80	150 ml OP	✓ Sporanox

Subsidy (Manufacturer's Price)	Subo	Fully	Brand or Generic
(Manufacturer's Price)	Per	√ sidised	Manufacturer

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOI E

Tab 200 mg - PCT	CBS	30	✓ Link Healthcare S29
			✓ Nizoral S29
		100	✓ Strides Shasun S29
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 below - Reta	il pharmacy		
Tab modified-release 100 mg	869.86	24	✓ Noxafil
Oral liq 40 mg per ml	761.13	105 ml OP	✓ Noxafil

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Fither:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERRINAFINE

* Tab 250 mg	1.33	14	✓ Deolate
VORICONAZOLE - Special Authority see SA1273 on the next	page – Retail phari	macy	
Tab 50 mg	91.00	56	✓ Vttack
Tab 200 mg	350.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage			
claimable	1,437.00	70 ml	✓ Vfend

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis: or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

PRIMAQUINE – Special Authority see SA1684 below – Retail pharmacy
Tab 7.5 mg117.00 56

CA4COA Consist Authority for Cubaids

✓ Primacin S29

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Antiparasitics

Antiprotozoals

QUININE SULPHATE

99

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
Antitrichomonal Agents			
METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl
Tab 400 mg - Up to 15 tab available on a PSO		21	✓ <u>Metrogyl</u>
Oral liq benzoate 200 mg per 5 ml		100 m	. 3,
Suppos 500 mg	24.46	10	✓ Flagyl
ORNIDAZOLE Tab 500 mg	32.95	10	✓ Arrow-Ornidazole
Antituberculotics and Antileprotics			
Note: There is no co-payment charge for all pharmaceuticals list	ed in the Antitubercul	otics a	and Antileprotics group regardless o
immigration status.	od iii tilo 7 tilitaborodi	Olloo (and Anthopronos group regardioss c
CLOFAZIMINE - Retail pharmacy-Specialist			
a) No patient co-payment payable			
b) Prescriptions must be written by, or on the recommendati	ion of, an infectious d	isease	e physician, clinical microbiologist o
dermatologist.			
* Cap 50 mg	442.00	100	✓ Lamprene S29
CYCLOSERINE - Retail pharmacy-Specialist			
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician. 	ion of, an infectious d	isease	e physician, clinical microbiologist o
Cap 250 mg	344.00	60	✓ Cyclorin S29
DAPSONE - Retail pharmacy-Specialist			-
a) No patient co-payment payable			
b) Prescriptions must be written by, or on the recommendat dermatologist	ion of, an infectious d	isease	e physician, clinical microbiologist of
Tab 25 mg	268.50	100	✓ Dapsone
Tab 100 mg	329.50	100	✓ Dapsone
ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Specialis	st		
a) No patient co-payment payable			
b) Prescriptions must be written by, or on the recommendat	ion of, an infectious d	isease	e physician, clinical microbiologist o
respiratory physician Tab 100 mg	85 73	100	✓ EMB Fatol S29
Tab 400 mg		56	✓ Myambutol \$29
ISONIAZID – Retail pharmacy-Specialist		00	inyumbutor
a) No patient co-payment payable			
b) Prescriptions must be written by, or on the recommendat	ion of, an internal med	dicine	physician, paediatrician, clinical
microbiologist, dermatologist or public health physician		u.oo	prijetetari, padarametari, dirindar
* Tab 100 mg	22.00	100	✓ <u>PSM</u>
ISONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist			
a) No patient co-payment payable			
b) Prescriptions must be written by, or on the recommendat	ion of, an internal med	dicine	physician, paediatrician, clinical
microbiologist, dermatologist or public health physician	0E F4	100	√ Difinah
* Tab 100 mg with rifampicin 150 mg * Tab 150 mg with rifampicin 300 mg		100 100	✓ <u>Rifinah</u>✓ Rifinah
Tab 100 mg with mampion 300 mg	170.00	100	<u> ⊓iiiiaii</u>

		NFECTIONS - AC	GENTS	S FOR S	SYSTEMIC USE
		Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
PARA	-AMINO SALICYLIC ACID - Retail pharmacy-Specialist				
b	No patient co-payment payable Prescriptions must be written by, or on the recommendative respiratory physician				•
	rans for oral liq 4 g sachet	280.00	30	✓ P	aser S29
	TONAMIDE – Retail pharmacy-Specialist				
	 No patient co-payment payable Prescriptions must be written by, or on the recommendated respiratory physician 	tion of, an infectious di	isease s	pecialist,	clinical microbiologist or
Т	ab 250 mg	305.00	100	✓ P	eteha S29
PYRA	ZINAMIDE - Retail pharmacy-Specialist				
	 No patient co-payment payable Prescriptions must be written by, or on the recommendal respiratory physician 	tion of, an infectious di	isease p	hysician,	clinical microbiologist or
* T	ab 500 mg	59.00	100	✓ A	FT-Pyrazinamide
RIFA	BUTIN - Retail pharmacy-Specialist				-
	No patient co-payment payable Prescriptions must be written by, or on the recommendat gastroenterologist	tion of, an infectious di	isease p	hysician,	respiratory physician or
* C	ap 150 mg	299.75	30	✓ N	lycobutin
RIFA	MPICIN - Subsidy by endorsement				
	No patient co-payment payable For confirmed recurrent Staphylococcus aureus infection antimicrobial based on susceptibilities and the prescriptic Retail pharmacy - Specialist. Specialist must be an interpaediatrician, or public health physician.	on is endorsed accordi	ingly; ca	n be waiv al microbi	ved by endorsement - ologist, dermatologist,
	ap 150 mg		100	_	tifadin
	ap 300 mg		100	_	lifadin
* (ral liq 100 mg per 5 ml	12.60	60 ml	▼ <u>H</u>	lifadin
Ani	ivirals				
For e	re preparations refer to Eye Preparations, Anti-Infective Pro	eparations, page 239			
Hep	patitis B Treatment				

ADEFOVIR DIPIVOXIL - Special Authority see SA0829 below - Retail pharmacy Tab 10 mg670.00

(Hepsera Tab 10 mg to be delisted 1 March 2021)

⇒SA0829 Special Authority for Subsidy

Initial application only from a gastroenterologist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

30

✓ Hepsera

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg+); and Documented resistance to lamivudine, defined as:
- 2 Patient has raised serum ALT (> 1 x ULN); and
- 3 Patient has HBV DNA greater than 100,000 copies per mL, or viral load 10 fold or higher over nadir; and
- 4 Detection of M204I or M204V mutation: and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
` e ´	Por 🗸	Manufacturor

continued...

- 5 Either:
 - 5.1 Both:
 - 5.1.1 Patient is cirrhotic; and
 - 5.1.2 adefovir dipivoxil to be used in combination with lamivudine: or
 - 5.2 Roth
 - 5.2.1 Patient is not cirrhotic; and
 - 5.2.2 adefovir dipivoxil to be used as monotherapy.

Renewal only from a gastroenterologist or infectious disease specialist. Approvals valid for 2 years where in the opinion of the treating physician, treatment remains appropriate and patient is benefiting from treatment.

Notes: Lamivudine should be added to adefovir dipivoxil if a patient develops documented resistance to adefovir dipivoxil, defined as:

- i) raised serum ALT (> 1 x ULN); and
- ii) HBV DNA greater than 100,000 copies per mL, or viral load 10 fold or higher over nadir; and
- iii) Detection of N236T or A181T/V mutation.

Adefovir dipivoxil should be stopped 6 months following HBeAg seroconversion for patients who were HBeAg+ prior to commencing adefovir dipivoxil.

The recommended dose of adefovir dipivoxil is no more than 10mg daily.

In patients with renal insufficiency adefovir dipivoxil dose should be reduced in accordance with the datasheet guidelines. Adefovir dipivoxil should be avoided in pregnant women and children.

ENTECAVIR

*	Tab 0.5 mg	52.00	30	✓ Entecavir Sandoz
	MIVUDINE - Special Authority see SA1685 below - Retail pharma			
	Tab 100 mg	6.95	28	✓ Zetlam
	Oral liq 5 mg per ml	.270.00 2	240 ml OP	✓ Zeffix
_				

⇒SA1685 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B. TENOFOVIR DISOPROXIL

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA1651... page 105

Herpesvirus Treatments

ACICLOVIR			
* Tab dispersible 200 mg	1.60	25	✓ Lovir
* Tab dispersible 400 mg	5.38	56	✓ Lovir
* Tab dispersible 800 mg	5.98	35	✓ <u>Lovir</u>
VALACICLOVIR			
Tab 500 mg	5.75	30	✓ Vaclovir
Tab 1,000 mg	11.35	30	✓ Vaclovir
VALGANCICLOVIR - Special Authority see SA1404 on the next p	age – Retail pha	armacy	
Tab 450 mg	225.00	60	✓ Valganciclovir
-			Mylan

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

⇒SA1404 Special Authority for Subsidy

Initial application — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
- 2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via PHARMAC's approved direct distribution supply. Further details can be found on PHARMAC's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see \$A1605 below

No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg......24,363.46 28 **✓ Harvoni**

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP) Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz/mayiret or:

The Coordinator, Hepatitis C Treatment Panel

PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA1904 below

Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 105 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the PHARMAC website.

Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a

⇒SA1904 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and

Subsidy	Fully	Brand or	Т
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

6.2.3 Condoms have not been consistently used.

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Antiretrovirals

⇒SA1651 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — **(Confirmed HIV)** only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initial application — (**Percutaneous exposure**) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA1651 on the previous	ous page – Retail pharr	nacy	
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
ETRAVIRINE - Special Authority see SA1651 on the prev	ous page – Retail phar	macy	
Tab 200 mg	770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA1651 on the prev	ious page – Retail phar	macy	
Tab 200 mg	60.00	60	✓ Nevirapine
			<u>Alphapharm</u>
Oral suspension 10 mg per ml	203.55	240 ml	✓ Viramune
			Suspension

Fully

Brand or

Subsidy

	(Manufacturer's F	Price) Subsi	dised Generic Manufacturer		
Nucleosides Reverse Transcriptase Inhibitors					
ABACAVIR SULPHATE – Special Authority see SA1651 on pag Tab 300 mg Oral liq 20 mg per ml ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority Note: abacavir with lamivudine (combination tablets) counts	180.00 256.31 see SA1651 on	60 240 ml OP page 105 – Rei	, ,		
anti-retroviral Special Authority. Tab 600 mg with lamivudine 300 mg	63.00	30	✓ <u>Kivexa</u>		
pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil counti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil	ounts as three ar	•	, ,		
245 mg (300 mg as a maleate) EMTRICITABINE – Special Authority see SA1651 on page 105 -		30	✓ <u>Mylan</u>		
Cap 200 mg	•	30	✓ Emtriva		
LAMIVUDINE – Special Authority see SA1651 on page 105 – Re Tab 150 mg		60	✓ <u>Lamivudine</u> Alphapharm		
Oral liq 10 mg per ml		240 ml OP	✓ 3TC		
ZIDOVUDINE [AZT] – Special Authority see SA1651 on page 10 Cap 100 mg		100 200 ml OP ge 105 – Retail p			
Protease Inhibitors					
ATAZANAVIR SULPHATE – Special Authority see SA1651 on p Cap 150 mg Cap 200 mg DARUNAVIR – Special Authority see SA1651 on page 105 – Re	141.68 188.91 tail pharmacy	60 60	✓ <u>Teva</u> ✓ <u>Teva</u>		
Tab 400 mg Darunavir Mylan to be Sole Supply on 1 April 2021	335.00	60	✓ Darunavir Mylan✓ Prezista		
Tab 600 mg Darunavir Mylan to be Sole Supply on 1 April 2021	196.65 476.00	60	✓ Darunavir Mylan✓ Prezista		
(Prezista Tab 400 mg to be delisted 1 April 2021) (Prezista Tab 600 mg to be delisted 1 April 2021)					
LOPINAVIR WITH RITONAVIR – Special Authority see SA1651 Tab 100 mg with ritonavir 25 mg Tab 200 mg with ritonavir 50 mg Oral lig 80 mg with ritonavir 20 mg per ml	183.75 463.00	Retail pharmacy 60 120 300 ml OP	✓ Kaletra ✓ Kaletra ✓ Kaletra		
RITONAVIR – Special Authority see SA1651 on page 105 – Ret Tab 100 mg	ail pharmacy	30	✓ <u>Norvir</u>		

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Strand Transfer Inhibitors

DOLUTEGRAVIR - Special Authority see SA1651 or	n page 105 – Retail pharmacy		
Tab 50 mg	1,090.00	30	Tivicay
RALTEGRAVIR POTASSIUM - Special Authority se	e SA1651 on page 105 – Retail	pharmacy	
Tab 400 mg	1,090.00	60	✓ Isentress
Tab 600 mg	1.090.00	60	✓ Isentress HD

Immune Modulators

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

- 1) Diagnosis
 - Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
 - PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
 - Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Exclusion Criteria

- Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia (< 2.0 × 10⁹) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA1972 below - Retail pharmacy

- a) See prescribing guideline above
- b) Note: PHARMAC will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at PHARMAC on 0800-023-588 option 4.

⇒SA1972 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and

INFECTIONS - AGENTS FOR SYSTEMIC USE

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	1	Manufacturer

continued...

2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400.000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and

INFECTIONS - AGENTS FOR SYSTEMIC USE

Subsidy	Fully	Brand or
(Manufacturer's	Price) Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Initial application — (ocular surface squamous neoplasia) only from an ophthalmologist. Approvals valid for 12 months where patient has ocular surface squamous neoplasia *.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

3 Either:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
 - - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate: or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

Renewal — (ocular surface squamous neoplasia) only from an ophthalmologist. Approvals valid for 12 months where the treatment remains appropriate and patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications.

Notes:

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet auidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

Urinary			_
	/ Irant	II a trata)	
	/ Haut	11111	114116

METHENAMINE (HEXAMINE) HIPPURATE			
* Tab 1 g40.01	100	✓ Hiprex	
NITROFURANTOIN			
* Tab 50 mg - Up to 30 tab available on a PSO22.20	100	✓ Nifuran	
* Tab 100 mg37.50	100	✓ Nifuran	

INFECTIONS - AGENTS FOR SYSTEMIC USE

Subsidy (Manufacturer's Price)	Ful Subsidise	,
 \$	Per •	Manufacturer

NORFLOXACIN

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	` \$ ´	Per	1	Manufacturer
Anticholinesterases				
Alluonomicotorasco				
NEOSTIGMINE METILSULFATE				
	10.00	40		l 000
Inj 2.5 mg per ml, 1 ml ampoule		10		Juno S29
	29.40			Max Health
	98.00	50	•	AstraZeneca
PYRIDOSTIGMINE BROMIDE				
▲ Tab 60 mg	45.70	100	./ I	Mestinon
_ 1au 60 mg	43.79	100	▼ <u>I</u>	Westilloll
N. A. 1114 M				
Non-Steroidal Anti-Inflammatory Drugs				
DICLOFENAC SODIUM				
* Tab EC 25 mg	1.23	50	√ [Diclofenac Sandoz
* Tab 50 mg dispersible	1.50	20	✓ \	/oltaren D
* Tab EC 50 mg	1.23	50	√ [Diclofenac Sandoz
* Tab long-acting 75 mg		500	_	Apo-Diclo SR
* Tab long-acting 100 mg		500		Apo-Diclo SR
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a		5		/oltaren
* Suppos 12.5 mg		10		/oltaren
* Suppos 25 mg		10		/oltaren
* Suppos 50 mg - Up to 10 supp available on a PSO	4.22	10	✓ \	/oltaren
* Suppos 100 mg	7.00	10	✓ \	/oltaren
IBUPROFEN				
	44.74	4 000		N. II
* Tab 200 mg		1,000		Relieve
* Tab long-acting 800 mg		30	_	buprofen SR BNM
* Oral liq 20 mg per ml	1.88	200 m	ı 🗸 <u>E</u>	<u>Ethics</u>
KETOPROFEN				
* Cap long-acting 200 mg	12.07	28	10	Oruvail SR
	12.07	20	• (Jiuvali Sii
MEFENAMIC ACID				
* Cap 250 mg	1.25	50		
, ,	(9.16)		F	Ponstan
	0.50	20	•	
	(5.60)	20		Ponstan
	(3.00)		'	Ulstall
NAPROXEN				
* Tab 250 mg	32.69	500	✓ 1	Noflam 250
* Tab 500 mg		250		Noflam 500
* Tab long-acting 750 mg		28	_	Naprosyn SR 750
* Tab long-acting 1 g		28		Naprosyn SR 1000
5 5 5	0.21	20	▼ Ī	tapiosyli on 1000
SULINDAC				
* Tab 100 mg	9.57	56	✓ I	Mylan S29
* Tab 200 mg		50		Aclin
	16.91	56		
	10.91	90	₩ :	Sulindac Mylan S29
TENOXICAM				
* Tab 20 mg	9.15	100	✓ 1	Filcotil
* Inj 20 mg vial		1	√ /	
, == 111g 11cl			- 7	

	MUSCULOSKELETAL SYSTEM				
	Subsidy (Manufacturer's Prio \$	ce) Subs Per	Fully Brand or sidised Generic ✓ Manufacturer		
NSAIDs Other					
CELECOXIB Cap 100 mg Cap 200 mg		60 30	✓ Celecoxib Pfizer✓ Celebrex✓ Celecoxib Pfizer		
Topical Products for Joint and Muscular Pain CAPSAICIN Crm 0.025% - Special Authority see SA1289 below - Retail pharmacy	9.75 13.27	45 g OP 60 g OP	✓ Zostrix ✓ Rugby Capsaicin Topical Cream S29		
Zostrix to be Sole Supply on 1 April 2021 (Rugby Capsaicin Topical Cream \$29 Crm 0.025% to be deliste >SA1289 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid osteoarthritis that is not responsive to paracetamol and oral non-s Antirheumatoid Agents	I without further re				
HYDROXYCHLOROQUINE – Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, systemi suppression, relevant dermatological conditions (cutaneous f mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmo Pharmacists may annotate the prescription as endorsed whe hydroxychloroquine. Note: Indication marked with a * is an u	orms of lupus and nary)*, and the pre re there exists a re	lichen planuescription is ecord of prior	s, cutaneous vasculitides and endorsed accordingly.		
* Tab 200 mg		100	✓ <u>Plaquenil</u>		
LEFLUNOMIDE Tab 10 mg Tab 20 mg		30 30	✓ <u>Arava</u> ✓ <u>Arava</u>		
PENICILLAMINE Tab 125 mg Tab 250 mg		100 100	✓ D-Penamine ✓ D-Penamine		
Drugs Affecting Bone Metabolism					
Alendronate for Osteoporosis					
ALENDRONATE SODIUM * Tab 70 mg	2.44	4	✓ Fosamax		
ALENDRONATE SODIUM WITH COLECALCIFEROL * Tab 70 mg with colecalciferol 5,600 iu	1.51	4	✓ Fosamax Plus		
Other Treatments					

DENOSUMAB - Special Authority see SA1777 on the next page - Retail pharmacy

Inj 60 mg prefilled syringe......326.00

✓ Prolia

1

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Fither:
 - 2.1 The patient is female and postmenopausal; or
 - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

DVVVI	ואססח	ISODIUM

Inj 3 mg per ml, 10 ml vial	5.98	1	✓ Pamisol
Inj 6 mg per ml, 10 ml vial	15.02	1	✓ Pamisol
, , ,		1	✓ Pamisol
DALOVIEENE LIVERDOOLII ORIBE	On a sight Audio site and OA4770 and the most many	D	I I

Si	Subsidy	Fully	Brand or
(Manufac	cturer's Price) Subsi	dised	Generic
	\$ Per	1	Manufacturer

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically: or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM

Tab 35 mg3.10	0 4	✓ Risedronate Sandoz
TERIPARATIDE - Special Authority see SA1139 below - Retail pharmacy		
Inj 250 mcg per ml, 2.4 ml490.00	0 1	✓ Forteo

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops

Si	ubsidy	Fully	Brand or
(Manufac	cturer's Price) Subsi	dised	Generic
`	\$ Per	1	Manufacturer

continued...

during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

⇒SA1780 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or

Subsidy	ce)	Fully	Brand or
(Manufacturer's Pric		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
- 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 Any of the following:

- 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
- 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
- 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below

	,	ully Brand or	
(Manufact	turer's Price) Subsidi	sed Generic	
	\$ Per	 Manufac 	turer

continued...

- -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

ALLOPURINOL			
* Tab 100 mg	11.47	500	✓ DP-Allopurinol
* Tab 300 mg	28.57	500	✓ DP-Allopurinol
BENZBROMARONE - Special Authority see SA1963	below - Retail pharmacy		
Tab 50 mg	22.50	100	✓ Narcaricin mite \$29
Tab 100 mg	13.50	30	✓ Desuric S29
			✓ Urinorm S29
	45.00	100	Benzbromaron AL
			100 \$29

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

0 50

100

✓ Colgout

COLCHICINE * Tab 500 mcg

* Tab 500 mog		100	• Ooigout
FEBUXOSTAT - Special Authority see SA1931 below - Re	etail pharmacy		
Tab 80 mg	39.50	28	Adenuric
Tab 120 mg	39.50	28	✓ Adenuric

⇒SA1931 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
 - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine

✓ Dantrium

✓ Dantrium

✓ Norflex

✓ Dantrium S29 S29

100

100

100

		,000_0	, L L	LIALOIOILM
	Subsidy (Manufacturer's Price)	Sub Per	Fully sidised	Brand or Generic Manufacturer
continued clearance less than 30 ml/minute). No dosage adjustment of febrimpairment. Optimal treatment with allopurinol in patients with reclearance-adjusted dose of allopurinol then, if serum urate remain allopurinol to 600 mg or the maximum tolerated dose.	nal impairment is det	ined as tr	eatment	to the creatinine
PROBENECID * Tab 500 mg	55.00	100	√ P	robenecid-AFT
Muscle Relaxants				
BACLOFEN				
* Tab 10 mg	4.20	100	✓ P	acifen
Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement		1	✓ L	ioresal Intrathecal
Subsidised only for use in a programmable pump in patie caused intolerable side effects and the prescription is en-		pastic age	ents have	e been ineffective or have
Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement	372.98	5	✓ N	ledsurge
Subsidised only for use in a programmable pump in patie caused intolerable side effects and the prescription is en		pastic age	ents have	e been ineffective or have
DANTROLENE				

Cap 50 mg......77.00

ORPHENADRINE CITRATE

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

Agents for Parkinsonism and Related Disorders

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE			
▲ Cap 100 mg	38.24	60	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE			
▲ Inj 10 mg per ml, 2 ml ampoule	59.50	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule	121.84	5	✓ Movapo
BROMOCRIPTINE MESYLATE			
* Tab 2.5 mg	32.08	100	✓ Apo-Bromocriptine
ENTACAPONE			·
▲ Tab 200 mg	22.00	100	✓ Entapone
LEVODOPA WITH BENSERAZIDE			
* Tab dispersible 50 mg with benserazide 12.5 mg	13.25	100	✓ Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg		100	✓ Madopar 62.5
* Cap 100 mg with benserazide 25 mg		100	✓ Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg		100	✓ Madopar HBS
* Cap 200 mg with benserazide 50 mg	26.25	100	✓ Madopar 250
LEVODOPA WITH CARBIDOPA			
* Tab 100 mg with carbidopa 25 mg	21.11	100	✓ Sinemet
* Tab long-acting 200 mg with carbidopa 50 mg	43.65	100	✓ Sinemet CR
	46.73		✓ Mylan S29
Sinemet CR to be Sole Supply on 1 February 2021			
* Tab 250 mg with carbidopa 25 mg		100	✓ Sinemet
(Mylan S29) Tab long-acting 200 mg with carbidopa 50 mg to b	e delisted 1 Febru	ary 2021)	
PRAMIPEXOLE HYDROCHLORIDE			
▲ Tab 0.25 mg	6.12	100	✓ Ramipex
▲ Tab 1 mg	20.73	100	✓ Ramipex
ROPINIROLE HYDROCHLORIDE			
▲ Tab 0.25 mg	2.85	84	✓ Ropin
	3.39	100	✓ Mylan S29
▲ Tab 1 mg	3.95	84	✓ Ropin
	4.70	100	✓ Mylan S29
▲ Tab 2 mg	5.48	84	✓ Ropin
▲ Tab 5 mg	12.50	84	✓ <u>Ropin</u>
SELEGILINE HYDROCHLORIDE			
* Tab 5 mg	22.00	100	✓ Apo-Selegiline
			S29 S29
TOLCAPONE			
▲ Tab 100 mg	152.38	100	✓ Tasmar
•			

Anticholinergics

RENI7	JUINE	MESVI	

Tab 2 mg	7.99	60	✓ Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	✓ Phebra

a) Up to 10 inj available on a PSO

b) Only on a PSO

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	d Generic
PROCYCLIDINE HYDROCHLORIDE Tab 5 mg	7.40	100		Kemadrin
Agents for Essential Tremor, Chorea and Relate	d Disorders			
RILUZOLE – Special Authority see SA1403 below – Retail pharm Wastage claimable Tab 50 mg	•	56	√	Rilutek
■ SA1403 Special Authority for Subsidy Initial application only from a neurologist or respiratory specialis following criteria: All of the following:	t. Approvals valid for	r 6 m	onths for a	applications meeting the
The patient has amyotrophic lateral sclerosis with disease The patient has at least 60 percent of predicted forced vita The patient has not undergone a tracheostomy; and The patient has not experienced respiratory failure; and Any of the following: 5.1 The patient is ambulatory; or 5.2 The patient is able to use upper limbs; or 5.3 The patient is able to swallow.				ie initial application; and
Renewal from any relevant practitioner. Approvals valid for 18 m All of the following: 1 The patient has not undergone a tracheostomy; and 2 The patient has not experienced respiratory failure; and 3 Any of the following: 3.1 The patient is ambulatory; or 3.2 The patient is able to use upper limbs; or 3.3 The patient is able to swallow.	onths for applications	s mee	iting the fo	ollowing criteria:
TETRABENAZINE Tab 25 mg	91.10	112	/	Motetis
Anaesthetics				
Local				
LIDOCAINE [LIGNOCAINE] Gel 2%, tube - Subsidy by endorsement	dministration and the	30 ml pres 10	cription is	Xylocaine 2% Jelly endorsed accordingly. Instillagel Lido

b) Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed

a) Up to 5 each available on a PSO

accordingly.

	Subsidy (Manufacture de Brise	٠	Fully	Brand or Generic
	(Manufacturer's Price \$	Per	ubsidised •	Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%	38.00	200 ml	✓ I	Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	8.75	25	√ I	Lidocaine-Claris
	17.50	50		
	(35.00))	Kylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	8.25	25	✓ <u>I</u>	Lidocaine-Claris
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5		
	(20.00))	Kylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	6.20	5	✓ [Lidocaine-Claris
Inj 2%, 20 ml vial - Up to 5 inj available on a PSO	6.45	5	✓ [Lidocaine-Claris
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –				
Subsidy by endorsement	81.50	10	√ [Pfizer
a) Up to 5 each available on a PSO		. •	-	
b) On heiding a health if a man with a different transfer deal and a minimum transfer				

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] – Special Authority see	SA0906 above – Retail pharn	nacy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - S	Special Authority see SA0906	above – Reta	ail pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	EMLA

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 112

Non-opioid Analgesics

•			
ASPIRIN			
* Tab dispersible 300 mg - Up to 30 tab available on a PS	O4.50	100	✓ Ethics Aspirin
CAPSAICIN - Subsidy by endorsement			
Subsidised only if prescribed for post-herpetic neuralgia of	r diabetic peripheral	neuropathy a	nd the prescription is endorsed
accordingly.			
Crm 0.075%	11.95	45 g OP	Zostrix HP
	15.83	57 g OP	Rugby Capsaicin
			Topical
			Cream S29
Zostrix HP to be Sole Supply on 1 April 2021			
NEFOPAM HYDROCHLORIDE			
Tab 30 mg	23.40	90	✓ Acupan

	Subsidy (Manufacturer's Price) \$	Si Per	Fully ubsidised	Brand or Generic Manufacturer
PARACETAMOL				
Tab 500 mg - blister pack	0.50	20	1	Medco Paracare Pharmacy Health
	1.12		•	Ethics Paracetamol Classic
	2.48	100		Paracare Pharmacy Health
	11.75	96	1	Panadol Mini Caps
	24.82	1,000	•	Paracetamol Pharmacare
			1	Pharmacare

- a) Maximum of 300 tab per prescription; can be waived by endorsement
- b) Up to 30 tab available on a PSO

c)

- Subsidy by endorsement for higher quantities is available for patients with long term conditions who require
 regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may
 annotate the prescription as endorsed where dispensing history supports a long-term condition.
- 2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.

Tab 500 mg - bottle pack - Maximum of 300 tab per			
prescription; can be waived by endorsement	24.82	1,000	Paracetamol Pharmacare
			Pharmacare

- Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.
- Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.

*	Oral liq 120 mg per 5 ml	5.45	1,000 ml	✓ Paracare
	 a) Up to 200 ml available on a PSO 			
	b) Not in combination			
*	Oral liq 250 mg per 5 ml	6.25	1,000 ml	✓ Paracare Double
				<u>Strength</u>
	a) Up to 100 ml available on a PSO			
	b) Not in combination			
*	Suppos 125 mg	3.29	10	✓ Gacet
*	Suppos 250 mg	3.79	10	✓ Gacet
*	Suppos 500 mg	12.40	50	✓ Gacet
(Pr	armacare Tab 500 mg - bottle pack to be delisted 1 March 2021)			

Opioid Analgesics

CODEINE PHOSPHATE - Safety medicine; prescriber may of	determine dispensing	frequency	
Tab 15 mg		100	✓ PSM
Tab 30 mg	7.45	100	✓ PSM
Tab 60 mg	14.25	100	✓ PSM
DIHYDROCODEINE TARTRATE			
Tab long-acting 60 mg	8.60	60	✓ DHC Continus

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
	Ψ	rei		Manuacturei
FENTANYL				
a) Only on a controlled drug form				
b) No patient co-payment payable				
 c) Safety medicine; prescriber may determine dispensing for 				
Inj 50 mcg per ml, 2 ml ampoule		10	✓	Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule	9.41	10	✓	Boucher and Muir
Patch 12.5 mcg per hour	2.95	5	✓	Fentanyl Sandoz
Patch 25 mcg per hour	3.66	5	✓	Fentanyl Sandoz
Patch 50 mcg per hour	6.65	5		Fentanyl Sandoz
Patch 75 mcg per hour	9.25	5		Fentanyl Sandoz
Patch 100 mcg per hour	11.40	5	✓	Fentanyl Sandoz
 c) Safety medicine; prescriber may determine dispensing fid) d) Extemporaneously compounded methadone will only be (methadone powder, not methadone tablets). e) For methadone hydrochloride oral liquid refer Standard I 	reimbursed at the rat	e of th	ne cheapes	st form available
Tab 5 mg		10	1	Methatabs
Oral lig 2 mg per ml		200 n		Biodone
Oral lig 5 mg per ml		200 n		Biodone Forte
Oral liq 10 mg per ml		200 n	nl 🗸	Biodone Extra Forte
Inj 10 mg per ml, 1 ml		10	1	AFT
IORPHINE HYDROCHLORIDE				
a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing fi	requency			
Oral liq 1 mg per ml		200 n	nl 🗸	RA-Morph
Oral liq 2 mg per ml		200 n	nl 🗸	RA-Morph
Oral lig 5 mg per ml		200 n	nl 🗸	Ordine \$29
-·-·				DA 14

Oral liq 10 mg per ml27.74

✓ <u>RA-Morph</u>
✓ Ordine S29

✓ RA-Morph

200 ml

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	✓	Manufacturer
MORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing for	requency			
Tab immediate-release 10 mg	2.80	10	✓	<u>Sevredol</u>
Tab immediate-release 20 mg	5.52	10	✓	Sevredol
Tab long-acting 30 mg	2.85	10	✓	Arrow-Morphine LA
Tab long-acting 60 mg		10	1	Arrow-Morphine LA
Cap long-acting 10 mg		10		m-Eslon
Cap long-acting 30 mg		10		m-Eslon
Cap long-acting 60 mg		10		m-Eslon
Cap long-acting 100 mg		10		m-Esion
1 0 0		5		
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a P	500.27	Э	V	DBL Morphine
				Sulphate
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a	PSO 4.47	5	/	DBL Morphine
				Sulphate
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a	PSO 4.76	5	1	DBL Morphine
,g p, ap		-		Sulphate
Ini 20 ma nor ml. 1 ml amnoula. Lin to E ini available on a	DCO 6 10	5	./	DBL Morphine
Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on a	F300.19	5	•	
				Sulphate
(Arrow-Morphine LA Tab long-acting 30 mg to be delisted 1 June				
(Arrow-Morphine LA Tab long-acting 60 mg to be delisted 1 April	il 2021)			
OXYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
, ,				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing for			,	
Tab controlled-release 5 mg		20		Oxycodone Sandoz
Tab controlled-release 10 mg		20		Oxycodone Sandoz
Tab controlled-release 20 mg	2.15	20		Oxycodone Sandoz
Tab controlled-release 40 mg	3.20	20	•	Oxycodone Sandoz
Tab controlled-release 80 mg	10.98	20	✓	Oxycodone Sandoz
Cap immediate-release 5 mg	1.88	20	✓	OxyNorm
Cap immediate-release 10 mg		20	1	OxyNorm
Cap immediate-release 20 mg		20		OxyNorm
Oral lig 5 mg per 5 ml		250 m		OxyNorm
		5		OxyNorm OxyNorm
Inj 10 mg per ml, 1 ml ampoule				
Inj 10 mg per ml, 2 ml ampoule		5		OxyNorm .
Inj 50 mg per ml, 1 ml ampoule	30.60	5	•	<u>OxyNorm</u>
PARACETAMOL WITH CODEINE - Safety medicine; prescribe	er may determine dispe	ensing	g frequenc	V
* Tab paracetamol 500 mg with codeine phosphate 8 mg		1,000		Paracetamol +
3 · · · · · · · · · · · · · · · · · · ·		,		Codeine (Relieve)
DETUIDING UNADDOCUM ODIDE				Coucino (ricino co)
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fi	requency			
Tab 50 mg		10	✓	PSM
Inj 50 mg per ml, 1 ml ampoule - Up to 5 inj available on a		5		DBL Pethidine
,	******	-		Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a	DSO 5.10	5	J	DBL Pethidine
ing 50 mg per mi, 2 mi ampoule – Op to 5 mg available on a	1 000.12	J	•	
				Hydrochloride

	Subsidy (Manufacturer's Price)	р-	Fully Subsidised	I Generic
	\$	Per		Manufacturer
RAMADOL HYDROCHLORIDE	4.50			T 100.400
Tab sustained-release 100 mg		20		Tramal SR 100
Tab sustained-release 150 mg		20		Tramal SR 150
Tab sustained-release 200 mg		20		Tramal SR 200
Cap 50 mg	2.80	100	•	Arrow-Tramadol
Antidepressants				
Cyclic and Related Agents				
MITRIPTYLINE - Safety medicine; prescriber may deter	, , ,		_	
Tab 10 mg		100		Arrow-Amitriptyline
Tab 25 mg		100		Arrow-Amitriptyline
Tab 50 mg		100		Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine;	prescriber may determine d	isper	ising frequ	ency
Tab 10 mg	13.99	100		Anafranil S29
-			1	Apo-Clomipramine
Tab 25 mg	9.46	100		Apo-Clomipramine
Anafranil S29 Tab 10 mg to be delisted 1 May 2021)				
OOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy	hy endorsement			
,	•			
 a) Safety medicine; prescriber may determine dispens b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. 	who were taking dosulepin			
	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride4.93		prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	30 50	prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin Mylan (\$29)
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	the 30 50 nsing	prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 y
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	30 50 nsing	prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 y Tofranil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	30 50 nsing 50 100	prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 y Tofranil Tofranil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	30 50 nsing	prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 y Tofranil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	30 50 nsing 50 100	prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 y Tofranil Tofranil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	30 50 nsing 50 100 50	prescription	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 y Tofranil Tofranil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	30 50 nsing 50 100 50	prescription graphic frequency rochloride	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 y Tofranil Tofranil Tofranil prior to 1 September
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	30 50 nsing 50 100 50	prescription graphic frequency rochloride	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 y Tofranil Tofranil Tofranil prior to 1 September
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	30 50 nsing 50 100 50	prescription frequence rochloride prescription	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 Y Tofranil Tofranil Tofranil prior to 1 September n as endorsed where the
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	30 50 nsing 50 100 50	prescription frequency rochloride prescription	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 Y Tofranil Tofranil Tofranil prior to 1 September n as endorsed where the Ludiomil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	30 50 nsing 50 100 50 50 4 hyd 30 50	prescription frequency rochloride prescription	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 Y Tofranil Tofranil Tofranil prior to 1 September n as endorsed where the Ludiomil Ludiomil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	s the 30 50 nsing 50 100 50 50 50 100 100	rochloride prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 Y Tofranil Tofranil Tofranil prior to 1 September n as endorsed where the Ludiomil Ludiomil Ludiomil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	s the 30 50 100 50 50 50 100 50 100 20	rochloride prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 Y Tofranil Tofranil Tofranil prior to 1 September n as endorsed where the Ludiomil Ludiomil Ludiomil Ludiomil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	s the 30 50 nsing 50 100 50 50 50 100 100	rochloride prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 Y Tofranil Tofranil Tofranil prior to 1 September n as endorsed where the Ludiomil Ludiomil Ludiomil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	s the 30 50 100 50 50 50 100 50 100 20	rochloride prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 Y Tofranil Tofranil Tofranil prior to 1 September n as endorsed where the Ludiomil Ludiomil Ludiomil Ludiomil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	s the 30 50 100 50 50 50 100 50 100 20	rochloride prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 Y Tofranil Tofranil Tofranil prior to 1 September n as endorsed where the Ludiomil Ludiomil Ludiomil Ludiomil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	s the 30 50 100 50 50 50 100 50 100 20	rochloride prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 Y Tofranil Tofranil Tofranil prior to 1 September n as endorsed where the Ludiomil Ludiomil Ludiomil Ludiomil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	s the 30 50 100 50 50 50 100 50 100 20	rochloride prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 Y Tofranil Tofranil Tofranil prior to 1 September n as endorsed where the Ludiomil Ludiomil Ludiomil Ludiomil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	30 50 nsing 50 100 50 50 4 hyd 4 the 30 50 100 20 30	prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 y Tofranil Tofranil Tofranil prior to 1 September n as endorsed where the Ludiomil Ludiomil Ludiomil Ludiomil Ludiomil Ludiomil Ludiomil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	30 50 nsing 50 100 50 50 4 hyd 4 the 30 50 100 20 30	prescription frequence rochloride prescription	n as endorsed where the Dosulepin Mylan Dosulepin Mylan 529 y Tofranil Tofranil Tofranil prior to 1 September n as endorsed where the Ludiomil Ludiomil Ludiomil Ludiomil Ludiomil Ludiomil
b) Subsidy by endorsement – Subsidised for patients 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dot Tab 75 mg	who were taking dosulepin Pharmacists may annotate hiepin] hydrochloride	30 50 nsing 50 100 50 50 4 hyd 4 the 30 50 100 20 30	prescription frequence rochloride prescription	n as endorsed where the Dosulepin Mylan Dosulepin Mylan \$29 y Tofranil Tofranil Tofranil prior to 1 September n as endorsed where the Ludiomil Ludiomil Ludiomil Ludiomil Ludiomil Ludiomil Ludiomil

				VOOD OTOTEM
	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Su Per	bsidised •	Generic Manufacturer
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	Selective			
TRANYLCYPROMINE SULPHATE				
Tab 10 mg	12.85	28	√ P	arnate S29 S29
Č	22.94	50	✓ P	arnate
	45.88	100	✓ P	arnate S29 S29
	96.00		✓ P	arnate S29 S29
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE				
* Tab 150 mg	6.40	60		urorix
Tab 300 mg	9.80	60	✓ <u>A</u>	urorix
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE				
* Tab 20 mg	1.52	84	✓ <u>P</u>	SM Citalopram
ESCITALOPRAM				
* Tab 10 mg	1.40	28	√ E	scitalopram- Apotex
* Tab 20 mg	2.49	28	√ E	scitalopram- Apotex
FLUOXETINE HYDROCHLORIDE	4.00	00		1
* Tab dispersible 20 mg, scored – Subsidy by endorsement	1.98 9.93	30		luox rrow-Fluoxetine
Subsidised by endorsement	9.93		▼ A	irow-riuoxetine
When prescribed for a patient who cannot swallov accordingly; or	v whole tablets or caps	sules an	d the pres	scription is endorsed
2) When prescribed in a daily dose that is not a multi	iple of 20 mg in which	case the	e prescrip	tion is deemed to be
endorsed. Note: Tablets should be combined wit	th capsules to facilitate	ncrem	ental 10 n	ng doses.
Cap 20 mg	2.91	84		luox
	7.49	90	✓ A	rrow-Fluoxetine
(Arrow-Fluoxetine Tab dispersible 20 mg, scored to be delisted (Arrow-Fluoxetine Cap 20 mg to be delisted 1 February 2021)	1 February 2021)			
PAROXETINE				
* Tab 20 mg	3.61	90	✓ L	<u>oxamine</u>
SERTRALINE				
Tab 50 mg	0.92	30	√ S	etrona
-				etrona AU
	3.05	90	✓ A	rrow-Sertraline
Tab 100 mg	1.61	30	_	<u>etrona</u>
	5.05	00	✓ S	etrona AU

5.25

90

✓ Arrow-Sertraline

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

MIRTAZAPINE		Subsidy (Manufacturer's Price)	Per	Fully Brand or Subsidised Generic r Manufacturer
Tab 45 mg	Other Antidepressants			
# Cap 37.5 mg	Tab 30 mg			
Antiepilepsy Drugs Agents for Control of Status Epilepticus CLONAZEPAM — Safety medicine; prescriber may determine dispensing frequency Inj 1 mg per ml, 1 ml. — 21.00	* Cap 37.5 mg * Cap 75 mg	8.11	84	✓ Enlafax XR
CLONAZEPAM - Safety medicine; prescriber may determine dispensing frequency				
Inj 1 mg per ml, 1 ml	Agents for Control of Status Epilepticus			
Inj 5 mg per ml, 2 ml ampoule — Subsidy by endorsement	· · · · · · · · · · · · · · · · · · ·		5	✓ Rivotril
Rectal tubes 5 mg - Up to 5 tube available on a PSO	Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement a) Up to 5 inj available on a PSO		5	✓ Hospira
★ Inj 5 ml	Rectal tubes 5 mg - Up to 5 tube available on a PSO		5	✓ Stesolid
★ Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a PSO	* Inj 5 ml	1,500.00	5	✓ AFT S29
CARBAMAZEPINE * Tab 200 mg	* Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a			·
CARBAMAZEPINE		133.92	5	✓ Hospira
★ Tab 200 mg 14.53 100 ✓ Tegretol ★ Tab long-acting 200 mg 16.98 100 ✓ Tegretol CR ★ Tab 400 mg 34.58 100 ✓ Tegretol ★ Tab long-acting 400 mg 39.17 100 ✓ Tegretol CR ★ Oral liq 20 mg per ml 26.37 250 ml ✓ Tegretol CLOBAZAM - Safety medicine; prescriber may determine dispensing frequency Tab 10 mg 9.12 50 ✓ Frisium CLONAZEPAM - Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml 7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE Cap 250 mg 140.88 100 ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin GABAPENTIN Note: Not subsidised in combination with subsidised pregabalin ★ Cap 100 mg 2.65 100 ✓ Apo-Gabapentin ★ Cap 300 mg 4.07 100 ✓ Apo-Gabapentin				
Tab 10 mg	* Tab 200 mg	16.98 34.58 39.17 26.37 2	100 100 100	✓ Tegretol CR ✓ Tegretol ✓ Tegretol CR
Oral drops 2.5 mg per ml 7.38 10 ml OP ✔ Rivotril ETHOSUXIMIDE Cap 250 mg 140.88 100 ✔ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✔ Zarontin GABAPENTIN Note: Not subsidised in combination with subsidised pregabalin ★ Cap 100 mg 2.65 100 ✔ Apo-Gabapentin ★ Cap 300 mg 4.07 100 ✔ Apo-Gabapentin	Tab 10 mg	9.12	50	✓ Frisium
Cap 250 mg 140.88 100 ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin GABAPENTIN Note: Not subsidised in combination with subsidised pregabalin * Cap 100 mg 2.65 100 ✓ Apo-Gabapentin * Cap 300 mg 4.07 100 ✓ Apo-Gabapentin	Oral drops 2.5 mg per ml		ml C	OP
Note: Not subsidised in combination with subsidised pregabalin ★ Cap 100 mg	Cap 250 mg			
★ Cap 300 mg		alin		
	* Cap 300 mg	4.07	100	✓ Apo-Gabapentin

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer	
LACOSAMIDE - Special Authority see SA1125 below - Retail p	harmacy				
▲ Tab 50 mg	25.04	14	✓ V	impat	
▲ Tab 100 mg	50.06	14	✓ V	impat	
•	200.24	56	✓ V	impat	
▲ Tab 150 mg	75.10	14	✓ V	impat	
ŭ	300.40	56	✓ V	impat	
▲ Tab 200 mg	400.55	56		impat	

⇒SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

LAMOTRIGINE

LA	VIOTRIGINE			
\blacktriangle	Tab dispersible 2 mg	55.00	30	✓ Lamictal
\blacktriangle	Tab dispersible 5 mg	50.00	30	✓ Lamictal
*	Tab dispersible 25 mg	2.76	56	✓ Logem
*	Tab dispersible 50 mg	3.31	56	Logem
*	Tab dispersible 100 mg		56	✓ Logem
۱F	VETIRACETAM			-
	Tab 250 mg	4.99	60	✓ Everet
	Tab 500 mg		60	✓ Everet
	Tab 750 mg		60	✓ Everet
	Tab 1,000 mg		60	✓ Everet
	Oral liq 100 mg per ml		300 ml OP	✓ Levetiracetam-AFT
РН	ENOBARBITONE			
	For phenobarbitone oral liquid refer Standard Formula	e nage 246		
*	Tab 15 mg		500	✓ PSM
*	Tab 30 mg		500	✓ PSM
	ENYTOIN SODIUM		000	<u> </u>
*		75.00	200	✓ Dilantin Infatab
不	Tab 50 mg		200	✓ Dilantin
	Cap 100 mg		200	✓ Dilantin
*	Cap 100 mg Oral liq 30 mg per 5 ml		500 ml	✓ Dilantin
-	. •	22.03	300 1111	Dilantin
PR	EGABALIN			
	Note: Not subsidised in combination with subsidised g	, ,		4.5. I !! D!!
*	Cap 25 mg		56	✓ Pregabalin Pfizer
*	Cap 75 mg		56	✓ Pregabalin Pfizer
*	Cap 150 mg	4.01	56	✓ Lyrica
	0 000	7.00		✓ <u>Pregabalin Pfizer</u>
*	Cap 300 mg	7.38	56	Pregabalin Pfizer

NERVOUS SYSTEM

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Subsidised	I Generic
	\$	Per		Manufacturer
PRIMIDONE				
* Tab 250 mg	17.25	100	✓	Apo-Primidone
	62.00	200	•	Mysoline S29 S29
SODIUM VALPROATE				
Tab 100 mg	13.65	100	✓	Epilim Crushable
Tab 200 mg EC	27.44	100	✓	Epilim
Tab 500 mg EC		100	✓	Epilim
* Oral liq 200 mg per 5 ml		300 m	nl 🗸	Epilim S/F Liquid
			✓	Epilim Syrup
* Inj 100 mg per ml, 4 ml	41.50	1	•	Epilim IV
STIRIPENTOL - Special Authority see SA1330 below - Retail p	harmacy			
Cap 250 mg	509.29	60	✓	Diacomit S29
Powder for oral liq 250 mg sachet	509.29	60	✓	Diacomit S29

⇒SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

Tab 25 mg	11.07	60	✓ Arrow-Topiramate
· ·			✓ Topiramate Actavis
	26.04		✓ Topamax
Tab 50 mg	18.81	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
	44.26		✓ Topamax
Tab 100 mg	31.99	60	✓ Arrow-Topiramate
			✓ Topiramate Actavis
	75.25		✓ Topamax
Tab 200 mg	55.19	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
	129.85		✓ Topamax
Sprinkle cap 15 mg	20.84	60	✓ Topamax
Sprinkle cap 25 mg	26.04	60	✓ Topamax
SABATRIN - Special Authority see SA1907 below			
Tab 500 mg		100	✓ Sabril

⇒SA1907 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

1 Either:

TOPIRAMATE

- 1.1 Patient has infantile spasms; or
- 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

- 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
- 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter): or
 - 2.2 It is impractical or impossible (due to comorbid conditions, or health system capacity constraints) to monitor the patient's visual fields..

Notes: ``Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages. **Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions, or health system capacity constraints) to monitor the patient's visual fields..

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 112

Acute Migraine Treatment

Tab orodispersible 10 mg3.65	30	✓ Rizamelt
SUMATRIPTAN Tab 50 mg24.44	100	✓ Apo-Sumatriptan
Tab 100 mg46.23	100	✓ Apo-Sumatriptan
Inj 12 mg per ml, 0.5 ml prefilled pen – Maximum of 10 inj per prescription34.00	2 OP	✓ <u>Imigran</u>

Prophylaxis of Migraine

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 51

PIZOTIFEN

※ Tab 500 mcg.......23.21 100 **✓ Sandomigran**

Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT - Special Authority see SA0987 on the next page - Retail pharmacy



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA0987 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLOR	RIDE
--------------------------	------

* Tab 16 mg	84	✓ Vergo 16
CYCLIZINE HYDROCHLORIDE		
Tab 50 mg0.55	10	✓ <u>Nausicalm</u>
CYCLIZINE LACTATE		
Inj 50 mg per ml, 1 ml14.95	5	✓ Nausicalm
21.53	10	✓ Hameln
(Nausicalm Inj 50 mg per ml, 1 ml to be delisted 1 May 2021)		
DOMPERIDONE		
* Tab 10 mg	100	✓ Pharmacy Health
HYOSCINE HYDROBROMIDE		
* Inj 400 mcg per ml, 1 ml ampoule	10	✓ Martindale S29
Patch 1.5 mg - Special Authority see SA1927 below - Retail		
pharmacy14.11	2	✓ Scopoderm TTS

⇒SA1927 Special Authority for Subsidy

Initial application — (control of intractable nausea, vomiting or inability to swallow saliva or clozapine induced hypersalivation) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Initial application — (pandemic circumstances- symptomatic relief of respiratory secretions in palliative care) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Requires palliative care in the community setting; and
- 2 Requires symptomatic relief of respiratory secretions that is not possible with 'as required subcutaneous hyoscine injections' due to COVID-19 constraints on the health sector; and
- 3 Access to a syringe driver for administration of injectable hyoscine is not possible due to COVID-19 constraints on the health sector.

Renewal — (control of intractable nausea, vomiting or inability to swallow saliva or clozapine induced hypersalivation) from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE

*	Tab 10 mg - Up to 30 tab available on a PSO1.30	100	✓ <u>Metoclopramide</u> <u>Actavis 10</u>
*	Inj 5 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO9.50	10	✓ <u>Pfizer</u>
O١	IDANSETRON		
*	Tab 4 mg2.68	50	✓ Onrex
*	Tab disp 4 mg - Up to 10 tab available on a PSO	10	✓ Ondansetron ODT-DRLA
*	Tab 8 mg4.57	50	✓ Onrex
*	Tab disp 8 mg – Up to 10 tab available on a PSO1.13	10	✓ Ondansetron ODT-DRLA

				INE	NVOUS STSTEIN
		Subsidy		Fully	Brand or
		(Manufacturer's Price)		Subsidised	
_		\$	Per		Manufacturer
PR	OCHLORPERAZINE				
*	Tab 3 mg buccal		50		
	T	(30.00)	050	,	Buccastem
*	Tab 5 mg – Up to 30 tab available on a PSO		250		Nausafix Stamatil
*	Inj 12.5 mg per ml, 1 ml – Up to 5 inj available on a PSO	25.61	10		Stemetil
A	ntipsychotics				
G	eneral				
AM	ISULPRIDE - Safety medicine; prescriber may determine di	spensing frequency			
	Tab 100 mg		30	1	Sulprix
	· ·	17.16	100		Amisulpride
					Mylan S29
	Tab 200 mg	14.96	60	1	Sulprix
	Tab 400 mg		60		Sulprix
ΔR	IPIPRAZOLE – Safety medicine; prescriber may determine of				
ALL	Tab 5 mg		30	1	Aripiprazole Sandoz
		28.58	49		Aripiprazole 1A
		20.00			Pharma S29
	Tab 10 mg	17 50	30	1	Aripiprazole Sandoz
	Tab 15 mg		30		Aripiprazole Sandoz
	Tab 20 mg		30		Aripiprazole Sandoz
	Tab 30 mg		30		Aripiprazole Sandoz
\cap L	LORPROMAZINE HYDROCHLORIDE - Safety medicine; pi				
OH	Tab 10 mg - Up to 30 tab available on a PSO		100		<u>Largactil</u>
	Tab 25 mg - Up to 30 tab available on a PSO	15.62	100		Largactil
	Tab 100 mg - Up to 30 tab available on a PSO		100		Largactil
	Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO		100		Largactil
CI	OZAPINE – Hospital pharmacy [HP4]				<u></u>
CL		lonov.			
	Safety medicine; prescriber may determine dispensing frequency and 25 mg		50		Clozaril
	Tab 25 mg	6.69	50		Clopine
		11.36	100		Clozaril
		13.37	100		Clopine
	Tab 50 mg		50		Clopine
	Tab 50 mg	17.33	100		Clopine
	Tab 100 mg		50		Clozaril
	740 700 mg	17.33	00		Clopine
		29.45	100		Clozaril
		34.65			Clopine
	Tab 200 mg	34.65	50		Clopine
	3	69.30	100	1	Clopine
	Suspension 50 mg per ml		100 m		Clopine
НΔ	LOPERIDOL - Safety medicine; prescriber may determine d	lisnensina frequency			•
, \	Tab 500 mcg - Up to 30 tab available on a PSO		100	1	Serenace
	Tab 1.5 mg - Up to 30 tab available on a PSO		100		Serenace
	Tab 5 mg - Up to 30 tab available on a PSO		50		Serenace
	J	29.72	100		Serenace
	Oral liq 2 mg per ml - Up to 200 ml available on a PSO		100 m		Serenace
	Ini 5 mg nor ml. 1 ml ampaula. Un to 5 ini available on a 5	21.55	10		Soronaco

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO21.55

✓ Serenace

10

	Subsidy		Fully	Brand or
	(Manufacturer's Price))	Subsidised	
	` \$	Per	•	Manufacturer
LEVOMEPROMAZINE - Safety medicine; prescriber may deter	rmine dispensing fred	illenc\	,	
Tab 25 mg (33.8 mg as a maleate)	, ,	100	_	Nozinan (Swiss)
Tab 25 mg as a maleate		100		Nozinan
Tab 100 mg (135 mg as a maleate)		100		Nozinan (Swiss)
Tab 100 mg as a maleate		100		Nozinan
LEVOMEPROMAZINE HYDROCHLORIDE - Safety medicine; p				
Inj 25 mg per ml, 1 ml ampoule		10		Nozinan
LITHIUM CARBONATE – Safety medicine; prescriber may dete	, ,			
Tab long-acting 400 mg		100		Priadel
Cap 250 mg	9.42	100	•	Douglas
OLANZAPINE - Safety medicine; prescriber may determine disp	pensing frequency			
Tab 2.5 mg	1.35	28	✓	Zypine
Tab 5 mg	1.58	28	✓	Zypine
Tab orodispersible 5 mg	1.81	28	✓	Zypine ODT
Tab 10 mg	2.01	28	✓	Zypine
Tab orodispersible 10 mg		28	✓	Zypine ODT
PERICYAZINE - Safety medicine; prescriber may determine dis	spensing frequency			
Tab 2.5 mg		84	/	Neulactil
1 db 2.0 mg	12.49	100		Neulactil
Tab 10 mg		84		Neulactil
145 10 mg	44.45	100		Neulactil
OLICTIADING Cofety medicines prescriber may determine dian		100		Touldotti
QUETIAPINE – Safety medicine; prescriber may determine disp		90	./	Oustand
Tab 25 mg		90		Quetapel Quetapel
Tab 100 mg Tab 200 mg		90		Quetapel
Tab 300 mg		90		Quetapel
•		90	•	<u>Quetaper</u>
RISPERIDONE – Safety medicine; prescriber may determine dis				
Tab 0.5 mg		60		Risperidone (Teva)
Tab 1 mg		60		Risperidone (Teva)
Tab 2 mg		60		Risperidone (Teva)
Tab 3 mg		60		Risperidone (Teva)
Tab 4 mg		60	_	Risperidone (Teva)
Oral liq 1 mg per ml	8.90	30 ml	•	Risperon
ZIPRASIDONE - Safety medicine; prescriber may determine dis	spensing frequency			
Cap 20 mg	14.50	60	✓	Zusdone
Cap 40 mg	24.70	60	✓	Zusdone
Cap 60 mg	33.80	60	✓	Zusdone
Cap 80 mg	39.70	60	✓	Zusdone
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pre	scriber may determin	ne dist	ensina fre	eauencv
Tab 10 mg		100		Clopixol
				'
Depot Injections				
FLUPENTHIXOL DECANOATE - Safety medicine; prescriber m		sing fr		
Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PSO		5		Fluanxol
Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PSO		5		Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	40.87	5	•	Fluanxol

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
HALOPERIDOL DECANOATE - Safety medicine; prescriber ma	ay determine dispensir	ng frequ	ency	
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO	28.39	5	. ✓ H	laldol
Inj 100 mg per ml, 1 ml – Úp to 5 inj available on a PSO	55.90	5		laldol Concentrate laldol Decanoas S29
OLANZAPINE - Special Authority see SA1428 below - Retail pl	narmacy			
Safety medicine; prescriber may determine dispensing frequ	ency			
Inj 210 mg vial	252.00	1	√ <u>Z</u>	yprexa Relprevy
Inj 300 mg vial	414.00	1	✓ <u>Z</u>	'yprexa Relprevv
Inj 405 mg vial	504.00	1	√ <u>Z</u>	Zyprexa Relprevv

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing	ı frequency		
Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe		1	✓ Invega Sustenna
Inj 100 mg syringe		1	✓ Invega Sustenna
Inj 150 mg syringe		1	✓ Invega Sustenna

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE – Special Authority see SA1427 on the next page – Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency			
Inj 25 mg vial	135.98	1	✓ Risperdal Consta
Inj 37.5 mg vial	178.71	1	✓ Risperdal Consta
Inj 50 mg vial	217.56	1	✓ Risperdal Consta



Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	idised	Generic
\$	Per	•	Manufacturer

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml − Up to 5 inj available on a PSO.......19.80 5 ✓ Clopixol

Anxiolytics

BUSPIRONE HYDROCHI ORIDE

* Tab 5 mg * Tab 10 mg		100 100	✓ <u>Orion</u> ✓ Orion
CLONAZEPAM – Safety medicine; prescriber may determine Tab 500 mcg		100	✓ Paxam
Tab 2 mg		100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determine dis	pensing frequency		
Tab 2 mg	61.07	500	✓ Arrow-Diazepam
Tab 5 mg	73.60	500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine of	dispensing frequency		
Tab 1 mg	9.72	250	✓ Ativan
Tab 2.5 mg	12.50	100	✓ <u>Ativan</u>
OXAZEPAM - Safety medicine; prescriber may determine di	spensing frequency		

Multiple Sclerosis Treatments

DIMETHYL FUMARATE - Special Authority see SA1	559 below – Retail pharmacy		
Wastage claimable			
Cap 120 mg	520.00	14	Tecfidera
Cap 240 mg	2,000.00	56	Tecfidera

⇒SA1559 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

continued...

100

100

✓ Ox-Pam✓ Ox-Pam

NERVOUS SYSTEM

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per Manufacturer

(Manufacturer's \$

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

continued...
The coordinator

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Phone: 04 460 4990

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- a) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- b) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- c) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
- v) new T2 lesions compared with a previous MR scan; and
- d) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week:
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point:
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- e) applications must be made by the patient's neurologist or general physician; and
- f) patients must have no previous history of lack of response to dimethyl fumarate; and
- g) patients must have not previously had intolerance to dimethyl fumarate; and
- h) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5: or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5: or
 - f) 3.0 to 4.5; or



	Subsidy	Fully	Brand or
(Manu	ufacturer's Price)	Subsidised	Generic
	\$ P	er 🗸	Manufacturer

- g) 3.5 to 4.5; or
- h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to dimethyl fumarate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate, teriflunomide and ocrelizumab is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

FINGOLIMOD - Special Authority see SA1562 below - Retail pharmacy

Wastage claimable

Cap 0.5 mg......2,200.00 28 **✓ Gilenya**

⇒SA1562 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic	
\$	Per	1	Manufacturer	

- c) last at least one week:
- d) start at least one month after the onset of a previous relapse;
- e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
- f) be distinguishable from the effects of general fatigue; and
- g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) patients must have no previous history of lack of response to fingolimod; and
- 7) patients must have not previously had intolerance to fingolimod; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to fingolimod; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate, teriflunomide and ocrelizumab is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

NATALIZUMAB - Special Authority see SA1563 below - Retail pharmacy

⇒SA1563 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria):
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least
 - f) be distinguishable from the effects of general fatigue; and
 - a) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) treatment must be initiated and supervised by a neurologist who is registered in the Tysabri Australasian Prescribing Programme operated by the supplier; and
- 7) patients must have no previous history of lack of response to natalizumab; and
- 8) patients must have not previously had intolerance to natalizumab; and
- a) Patient is JC virus negative, or
 - b) Patient is JC virus positive and has given written informed consent acknowledging an understanding of the risk of progressive multifocal leucoencephalopathy (PML) associated with natalizumab
- 10) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0: or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or

NERVOUS SYSTEM

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	1	Manufacturer	

continued...

- 3) intolerance to natalizumab; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier.

Switching between natalizumab, fingolimod, dimethyl fumarate, teriflunomide and ocrelizumab is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

OCRELIZUMAB - Special Authority see SA1867 below - Retail pharmacy

⇒SA1867 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

- a) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- b) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- c) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- d) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;



Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

- e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least
- f) be distinguishable from the effects of general fatigue; and
- g) not be associated with a fever (T> 37.5°C); and
- e) applications must be made by the patient's neurologist or general physician; and
- f) patients must have no previous history of lack of response to ocrelizumab; and
- g) patients must have not previously had intolerance to ocrelizumab; and
- h) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- 1) Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0: or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0: or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to ocrelizumab: or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate, teriflunomide and ocrelizumab is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

TERIFLUNOMIDE - Special Authority see SA1560 below - Retail pharmacy

Wastage claimable

28 ✓ Aubagio

⇒SA1560 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990 Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Entry Criteria

Subsidy (Manufacturer's Price)	Fully Price) Subsidised		Brand or Generic	
\$	Per	✓	Manufacturer	

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week:
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist or general physician; and
- 6) patients must have no previous history of lack of response to teriflunomide; and
- 7) patients must have not previously had intolerance to teriflunomide; and
- 8) patient must not be co-prescribed beta interferon or glatiramer acetate.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months. Progression of disability is defined as progress by any
 of the following EDDSS points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to teriflunomide; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Switching between natalizumab, fingolimod, dimethyl fumarate, teriflunomide and ocrelizumab is permitted provided the EDSS stopping criteria are not met. Switching to interferon or glatiramer acetate is only permitted provided the EDSS stopping criteria are not met and both fingolimod and natalizumab are either not tolerated or treatment with both agents would be clinically inappropriate. Continued relapses on treatment would be expected to lead to a switch of treatment provided the stopping criteria



Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

continued...

are not met. If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur.

Other Multiple Sclerosis Treatments

GLATIRAMER ACETATE - Special Authority see SA1808 below - Retail pharmacy

Inj 40 mg prefilled syringe......2,275.00

12 Copaxone

⇒SA1808 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. As a new Special Authority approval number is required the MSTAC coordinator should be notified of the change and a new prescription provided. **Entry Criteria**

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- f) be distinguishable from the effects of general fatigue; and
- g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or
 - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

Stopping Criteria

Any of the following:

- Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment.
 Progression of disability is defined as progress by any of the following EDDSS Points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0; or
 - c) 1.5 to 3.5; or
 - d) 2.0 to 4.0: or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
- h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

INTERFERON BETA-1-ALPHA - Special Authority see SA1809 below - Retail pharmacy

Inj 6 million iu prefilled syringe	1,170.00	4	Avonex
Injection 6 million iu per 0.5 ml pen injector	1,170.00	4	Avonex Pen

⇒SA1809 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. As a new Special Authority approval number is required the MSTAC coordinator should be notified of the change and a new prescription provided.

Entry Criteria

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:
 - a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the
 past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria):
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week;
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or
 - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

Stopping Criteria

Any of the following:

1) Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment. Progression of disability is defined as progress by any of the following EDDSS Points:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
- b) 1.0 to 3.0; or
- c) 1.5 to 3.5; or
- d) 2.0 to 4.0; or
- e) 2.5 to 4.5; or
- f) 3.0 to 4.5; or g) 3.5 to 4.5; or
- h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the beta-interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

INTERFERON BETA-1-BETA – Special Authority see SA1810 below – Retail pharmacy

SA1810 Special Authority for Subsidy

Special Authority approved by the Multiple Sclerosis Treatment Committee

Notes: Special Authority approved by the Multiple Sclerosis Treatment Assessment Committee (MSTAC). Applications will be considered by MSTAC at its regular meetings and approved subject to eligibility according to the Entry and Stopping criteria (below).

Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The coordinator Phone: 04 460 4990

Multiple Sclerosis Treatment Assessment Committee Facsimile: 04 916 7571

PHARMAC PO Box 10 254 Email: mstaccoordinator@pharmac.govt.nz

Wellington

Completed application forms must be sent to the coordinator for MSTAC and will be considered by MSTAC at the next practicable opportunity.

Notification of MSTAC's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Only prescriptions for 6 million iu of interferon beta-1-alpha per week, or 8 million iu of interferon beta-1-beta every other day, 40 mg glatiramer acetate 3 times weekly will be subsidised.

Switching between treatments is permitted within the 12 month approval period without reapproval by MSTAC. As a new Special Authority approval number is required the MSTAC coordinator should be notified of the change and a new prescription provided. **Entry Criteria**

- 1) Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2) patients must have Clinically Definite Relapsing Remitting MS with or without underlying progression; and
- 3) patients must have:



-				
	Subsidy	Fully	Brand or	
	(Manufacturer's Price)	Subsidised	Generic	
	•	Por 🗸	Manufacturor	

continued...

- a) EDSS score 0 4.0 and:
 - Experienced at least 1 significant relapse of MS in the previous 12 months or 2 significant relapses in the past 24 months; and
 - Evidence of new inflammatory activity on an MR scan within the past 24 months, any of the following:
 - i) a gadolinium enhancing lesion; or
 - ii) a Diffusion Weighted Imaging positive lesion; or
 - iii) a T2 lesion with associated local swelling; or
 - iv) a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse; or
 - v) new T2 lesions compared with a previous MR scan; and
- 4) A significant relapse must:
 - a) be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse but the neurologist/physician must be satisfied that the clinical features were characteristic and met the specified criteria);
 - b) be associated with characteristic new symptom(s)/sign(s) or substantial worsening of previously experienced symptom(s)/sign(s);
 - c) last at least one week:
 - d) start at least one month after the onset of a previous relapse;
 - e) be severe enough to change either the EDSS or at least one of the Kurtzke Functional System scores by at least 1 point;
 - f) be distinguishable from the effects of general fatigue; and
 - g) not be associated with a fever (T> 37.5°C); and
- 5) applications must be made by the patient's neurologist; and
- 6) patients must have no previous history of lack of response to beta-interferon or glatiramer acetate; and
- 7) patients must have either:
 - a) intolerance to both natalizumab and fingolimod; or
 - b) treatment with both natalizumab and fingolimod is considered clinically inappropriate; and
- 8) patient will not be co-prescribed natalizumab or fingolimod.

Stopping Criteria

Any of the following:

- 1) Confirmed progression of disability that is sustained for six months during a minimum of one year of treatment. Progression of disability is defined as progress by any of the following EDDSS Points:
 - a) from starting at EDSS 0 increasing to (i.e. stopping on reaching) EDSS 3.0; or
 - b) 1.0 to 3.0: or
 - c) 1.5 to 3.5: or
 - d) 2.0 to 4.0; or
 - e) 2.5 to 4.5; or
 - f) 3.0 to 4.5; or
 - g) 3.5 to 4.5; or
 - h) 4.0 to 4.5.
- 2) increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment)(see note); or
- 3) intolerance to interferon beta-1-alpha, and/or interferon beta-1-beta and/or glatiramer acetate; or
- 4) non-compliance with treatment, including refusal to undergo annual assessment.

Note: Treatment with interferon beta -1-beta, interferon beta-1-alpha and glatiramer acetate, is permitted only if treatment with both natalizumab and fingolimod is not tolerated or treatment with both would be clinically inappropriate. Beta-interferon or glatiramer acetate will not be funded as second line treatments if EDSS progression has occurred on treatment with natalizumab or fingolimod. Patients who have an increasing relapse rate over 12 months of treatment (compared with the relapse rate on starting treatment) and who do not meet the EDSS Stopping Criteria at annual review may switch from either of the

NERVOUS SYSTEM

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa. Patients may switch from either of the beta-interferon's [interferon beta-1-beta or interferon beta-1-alpha] to glatiramer acetate or vice versa for increased relapses only once, after which they will be required to stop funded treatment if they meet any of the Stopping Criteria at annual review (including the criterion relating to increasing relapse rate over 12 months of treatment). If a relapse has resulted in an increased EDSS score that potentially may lead to discontinuation of treatment according to stopping criteria, a period of 6 months is allowed from the start of the relapse for recovery to occur. In this setting anti-JCV antibody positive status may be accepted as a clinically inappropriate reason for treatment with natalizumab.

Sedatives and Hypnotics

MELATONIN – Special Authority see SA1666 below – Retail pharmacy
Tab modified-release 2 mg − No more than 5 tab per day28.22 30 ✓ Circadin

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

MIDAZOLAM - Safety medicine; prescriber may determine dispensir	g frequency		
Inj 1 mg per ml, 5 ml ampoule	4.30	10	✓ Midazolam-Baxter
, , , , ,			✓ Midazolam-Claris
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available			
on a PSO	14.90	10	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be end	orsed for status e	pilepticus us	se only.
Inj 5 mg per ml, 3 ml ampoule	2.50	5	✓ Midazolam-Baxter
			✓ Midazolam-Claris
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj available on			
a PSO	11.90	5	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be end	orsed for status e	pilepticus us	se only.
(Midazolam-Claris Inj 1 mg per ml, 5 ml ampoule to be delisted 1 Mar	ch 2021)		
(Midazolam-Claris Inj 5 mg per ml, 3 ml ampoule to be delisted 1 Ma	rch 2021)		
PHENOBARBITONE SODIUM - Special Authority see SA1386 on the	e next page - Re	tail pharmad	су
Inj 200 mg per ml, 1 ml ampoule	78.20	10	✓ Max Health S29



Subsidy (Manufacturer's Price)	Subi	Fully sidised	Brand or Generic
(Manufacturer's Price)	Per	siuiseu •	Manufacturer

⇒SA1386 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

TEMAZEPAM – Safety medicine; prescriber may determine of Tab 10 mg		25	✓ <u>Normison</u>
TRIAZOLAM - Safety medicine; prescriber may determine dis	spensing frequency		
Tab 125 mcg	5.10	100	
•	(9.85)		Hypam
Tab 250 mcg	4.10 [′]	100	,,
· ·	(11.20)		Hypam
ZOPICLONE - Safety medicine; prescriber may determine di	spensing frequency		
Tab 7.5 mg		500	✓ Zopiclone Actavis

Stimulants/ADHD Treatments

ATOMOXETINE - Brand switch fee payable (Pharmaco	de 2576996) - see page 24	4 for details	3
Cap 10 mg	18.41	28	✓ Generic Partners
Cap 18 mg	27.06	28	✓ Generic Partners
Cap 25 mg	29.22	28	✓ Generic Partners
Cap 40 mg	29.22	28	✓ Generic Partners
Cap 60 mg	46.51	28	✓ Generic Partners
Cap 80 mg	56.45	28	✓ Generic Partners
Cap 100 mg	58.48	28	✓ Generic Partners
DEXAMFETAMINE SULFATE - Special Authority see S	A1149 below – Retail phar	macy	
a) Only on a controlled drug form			
b) Safety medicine; prescriber may determine dispe	nsing frequency		

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

100

✓ PSM

All of the following:

1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and

- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Subsidy	Fu	ully	Brand or
(Manufacturer's Price)	Subsidis		Generic
\$	Per	✓	Manufacturer

continued

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

ig iroquorioy		
3.20	30	✓ Rubifen
	30	✓ Ritalin
		✓ Rubifen
7.75	30	✓ Methylphenidate ER
		- Teva
7.85	30	✓ Rubifen
	30	✓ Rubifen SR
50.00	100	✓ Ritalin SR
11.45	30	✓ Methylphenidate ER
		- Teva
15.50	30	 Methylphenidate ER
		- Teva
22.25	30	✓ Methylphenidate ER
	30	- Teva
		3.20 30 3.00 30

(Ritalin SR Tab sustained-release 20 mg to be delisted 1 June 2021)

⇒SA1964 Special Authority for Subsidy


Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:



Subsidy (Manufacturer's Price)	Subsid	Fully dised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine: prescriber may determine dispensing frequency

Tab extended-release 18 mg	58.96	30	✓ Concerta
Tab extended-release 27 mg		30	✓ Concerta
Tab extended-release 36 mg		30	✓ Concerta
Tab extended-release 54 mg		30	✓ Concerta
Cap modified-release 10 mg		30	✓ Ritalin LA
Cap modified-release 20 mg		30	✓ Ritalin LA
Cap modified-release 30 mg		30	✓ Ritalin LA
Cap modified-release 40 mg		30	Ritalin LA

⇒SA1965 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in

NERVOUS SYSTEM

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic
` \$	Per	•	Manufacturer

continued...

writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL - Special Authority see SA1932 below - Retail pharma	су		
Tab 100 mg	64.00	60	Modavigil

⇒SA1932 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Any of the following:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 A multiple sleep latency test is not possible due to COVID-19 constraints on the health sectors; or
 - 2.3 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	4.34	90	✓ Donepezil-Rex
* Tab 10 mg	6.64	90	✓ Donepezil-Rex
RIVASTIGMINE - Special Authority see SA1488 below -	Retail pharmacy		
Patch 4.6 mg per 24 hour	48.75	30	✓ Generic Partners
Patch 9.5 mg per 24 hour	48.75	30	✓ Generic Partners

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate: and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

......18.37 28

Tab sublingual 8 mg with naloxone 2 mg53.12

✓ Buprenorphine Naloxone BNM

✓ Buprenorphine Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

	Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic	
	\$	Per	✓	Manufacturer	
DISULFIRAM					
Tab 200 mg	250.00	100	✓ A	Antabuse	
NALTREXONE HYDROCHLORIDE - Special Authority see S.	A1408 below – Retail p	harmac	y		
Tab 50 mg	133.33	30	✓ <u>N</u>	laltraccord	

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

a) Nicotine will not be funded in amounts less than 4 weeks of treatment.

 b) Note: Direct Provision by a pharmacist permitted under the provision 	ovisions in Part	I of Section	A.
Patch 7 mg - Up to 28 patch available on a PSO	18.14	28	Habitrol
Patch 7 mg for direct distribution only - [Xpharm]	3.94	7	Habitrol
Patch 14 mg - Up to 28 patch available on a PSO	19.95	28	Habitrol
Patch 14 mg for direct distribution only - [Xpharm]	4.52	7	✓ Habitrol
Patch 21 mg - Up to 28 patch available on a PSO	22.86	28	✓ Habitrol
Patch 21 mg for direct distribution only - [Xpharm]	5.18	7	✓ Habitrol
Lozenge 1 mg - Up to 216 loz available on a PSO	19.18	216	Habitrol
Lozenge 1 mg for direct distribution only - [Xpharm]	3.20	36	✓ Habitrol
Lozenge 2 mg - Up to 216 loz available on a PSO	21.02	216	✓ Habitrol
Lozenge 2 mg for direct distribution only - [Xpharm]	3.24	36	✓ Habitrol
Gum 2 mg (Fruit) - Up to 384 piece available on a PSO	38.21	384	✓ Habitrol
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]	8.64	96	✓ Habitrol
Gum 2 mg (Mint) - Up to 384 piece available on a PSO	38.21	384	✓ Habitrol
Gum 2 mg (Mint) for direct distribution only - [Xpharm]	8.64	96	✓ Habitrol
Gum 4 mg (Fruit) - Up to 384 piece available on a PSO	44.17	384	✓ Habitrol
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]	10.01	96	✓ Habitrol
Gum 4 mg (Mint) - Up to 384 piece available on a PSO	44.17	384	✓ Habitrol
Gum 4 mg (Mint) for direct distribution only - [Xpharm]	10.01	96	✓ Habitrol

VARENICLINE TARTRATE - Special Authority see SA1845 on the next page - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	25.64	53 OP	✓ Varenicline Pfizer
Tab 1 mg		56	✓ Varenicline Pfizer



Subsidy (Manufacturer's Price)	F Subsidi	ully	Brand or Generic
 \$	Per	√	Manufacturer

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking;
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to guit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy: or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking:
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see SA1667 below

Inj 25 mg vial	 271.35 ´	1	✓ Ribomustin
Inj 100 mg vial	 1,085.38	1	✓ Ribomustin
Inj 1 mg for ECP	 11.40	1 mg	✓ Baxter

⇒SA1667 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 All of the following:
 - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
 - 3.2.2 The patient has not received prior bendamustine therapy; and
 - 3.2.3 Fither:
 - 3.2.3.1 Both:
 - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more: or
 - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Fither:
 - 2.1 Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

- 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.
 Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

BUSULFAN – PCT – Retail pharmacy-Specialist Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN – PCT only – Specialist			,
Inj 10 mg per ml, 45 ml vial	32.59	1	✓ DBL Carboplatin
, , , , , , , , , , , , , , , , , , , ,	45.20		✓ Carboplatin Ebewe
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist			
Inj 100 mg vial	1,387.00	1	✓ BiCNU
			✓ Bicnu Heritage S29
Inj 100 mg for ECP	1,387.00	100 mg OP	✓ Baxter
CHLORAMBUCIL - PCT - Retail pharmacy-Specialist			
Tab 2 mg	29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	12.29	1	✓ DBL Cisplatin
.,	15.00		✓ Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial	19.70	1	✓ DBL Cisplatin
, •	21.00		✓ Cisplatin Ebewe
Inj 1 mg for ECP	0.25	1 mg	✓ Baxter
(DBL Cisplatin Inj 1 mg per ml, 50 ml vial to be delisted 1 April	il 2021)		
CYCLOPHOSPHAMIDE			
Tab 50 mg - PCT - Retail pharmacy-Specialist	79.00	50	✓ Endoxan S29
	158.00	100	✓ Procytox S29
Wastage claimable			•
Inj 1 g vial - PCT - Retail pharmacy-Specialist	35.65	1	✓ Endoxan
	127.80	6	Cytoxan
Inj 2 g vial – PCT only – Specialist		1	Endoxan
Inj 1 mg for ECP - PCT only - Specialist	0.04	1 mg	✓ Baxter
IFOSFAMIDE - PCT only - Specialist			
lnj 1 g	96.00	1	✓ Holoxan
lnj 2 g	180.00	1	✓ Holoxan
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
LOMUSTINE - PCT - Retail pharmacy-Specialist			
Cap 10 mg	132.59	20	✓ CeeNU
Cap 40 mg	399.15	20	✓ CeeNU
MELPHALAN			
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	✓ Alkeran
Inj 50 mg - PCT only - Specialist	67.80	1	✓ Alkeran
			✓ Alkeran S29 S29
	420.00		✓ Tillomed S29

	Subsidy		Fully	/ Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
OXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	/	Oxaliplatin Actavis
.,				100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	46.32	1	✓	Oxaliplatin Accord
Inj 1 mg for ECP		1 mg	· •	Baxter
THIOTEPA - PCT only - Specialist		•		
Inj 15 mg vial	CBS	1	✓	Bedford S29
, •			1	THIO-TEPA S29
			•	Tepadina S29
Inj 100 mg vial	CBS	1	✓	Tepadina S29
Antimetabolites				
AZACITIDINE - PCT only - Specialist - Special Authority see SA	1467 below			
Inj 100 mg vial		1	1	Azacitidine Dr
, ,				Reddy's
	605.00		/	Vidaza
Inj 1 mg for ECP		1 mg		Baxter
", ' "g o Lo "		9	,	Bunto

⇒SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- 4 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

	Subsidy		Fully Brand or
((Manufacturer's Pri	ce) S Per	ubsidised Generic ✓ Manufacturer
ALCIUM FOLINATE			
Tab 15 mg - PCT - Retail pharmacy-Specialist	114.69	10	✓ DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specialis		5 1	 ✓ Hospira ✓ Calcium Folinate
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	9.49	1	Sandoz ✓ Calcium Folinate Sandoz
Inj 100 mg - PCT only - Specialist	7.33	1	✓ Calcium Folinate Ebewe
Inj 300 mg - PCT only - Specialist	22.51	1	Calcium Folinate Ebewe
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	25.14	1	Calcium Folinate Sandoz
Inj 1 g - PCT only - Specialist	67.51	1	✓ Calcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial – PCT only – Specialist		1	✓ Calcium Folinate Sandoz
Inj 1 mg for ECP – PCT only – Specialist APECITABINE – Retail pharmacy-Specialist	0.06	1 mg	✓ Baxter
Tab 150 mg		60 120	✓ <u>Capercit</u> ✓ <u>Capercit</u>
ADRIBINE - PCT only - Specialist Inj 1 mg per ml, 10 ml	749 96	1	✓ Leustatin
Inj 10 mg for ECP		10 mg OF	
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialis Inj 100 mg per ml, 20 ml vial – PCT – Retail	t400.00	5	✓ Pfizer
pharmacy-Specialist	41.36	1	✓ Pfizer
Inj 1 mg for ECP - PCT only - Specialist		10 mg	✓ Baxter
Inj 100 mg intrathecal syringe for ECP – PCT only – Specialis JDARABINE PHOSPHATE		100 mg Ol	P Baxter
Tab 10 mg - PCT - Retail pharmacy-Specialist		20	✓ Fludara Oral
Inj 50 mg vial – PCT only – Specialist		5	✓ Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist	115.29	50 mg OF	○ ✓ Baxter
UOROURACIL	10.00		/ Fluoresmanii Firema
Inj 50 mg per ml, 20 ml vial - PCT only - Specialist		1 1	✓ Fluorouracil Ebewe✓ Fluorouracil Ebewe
Inj 1 mg for ECP – PCT only – Specialist		100 mg	✓ Fluorouracii Ebewe ✓ Baxter
EMCITABINE HYDROCHLORIDE - PCT only - Specialist		100 mg	DUNIGI
Inj 1 g, 26.3 ml vial	62 50	1	✓ DBL Gemcitabine
Inj 1 g		1	✓ Gemcitabine Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
NOTECAN HYDROCHLORIDE - PCT only - Specialist		3	
Inj 20 mg per ml, 5 ml vial	71.44	1	✓ Irinotecan Accord \$29
			✓ Irinotecan Actavis 100
	100.00		✓ Irinotecan-Rex
Inj 1 mg for ECP	0.75	1 mg	✓ Baxter

	Subsidy (Manufacturer's Pric	e) Per	Fully Subsidised	Brand or Generic Manufacturer	
MERCAPTOPURINE Tab 50 mg - PCT - Retail pharmacy-Specialist	37.00	25	√ <u>P</u>	uri-nethol	
Oral suspension 20 mg per ml – Retail pharmacy-Specialist - Special Authority see SA1725 below		100 ml (OP ✓ A	Ilmercap	

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

ME	THOTREXATE		
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist8.05	90	✓ Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist31.75	90	✓ Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist47.50	5	✓ Hospira
			 Methotrexate DBL
*	Inj 7.5 mg prefilled syringe14.61	1	✓ Methotrexate
			Sandoz
*	Inj 10 mg prefilled syringe14.66	1	✓ Methotrexate
			Sandoz
*	Inj 15 mg prefilled syringe14.77	1	✓ Methotrexate
			Sandoz
*	Inj 20 mg prefilled syringe14.88	1	✓ Methotrexate
			Sandoz
*	Inj 25 mg prefilled syringe14.99	1	✓ Methotrexate
			Sandoz
*	Inj 30 mg prefilled syringe15.09	1	✓ Methotrexate
			Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist30.00	5	✓ DBL Methotrexate
			Onco-Vial
			✓ Methotrexate DBL
			Onco-Vial
*	Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Specialist45.00	1	 DBL Methotrexate
			Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist25.00	1	Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail		
	pharmacy-Specialist79.99	1	✓ Methotrexate Ebewe
	Inj 1 mg for ECP - PCT only - Specialist0.06	1 mg	✓ Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist4.73	5 mg OP	✓ Baxter
,	ospira Inj 2.5 mg per ml, 2 ml to be delisted 1 May 2021)		
(DI	BL Methotrexate Onco-Vial Inj 25 mg per ml, 2 ml vial to be delisted 1 May 202	1)	
PΕ	METREXED - PCT only - Specialist - Special Authority see SA1679 below		
	Inj 100 mg vial60.89	1	Juno Pemetrexed
	Inj 500 mg vial217.77	1	Juno Pemetrexed
	Inj 1 mg for ECP	1 mg	✓ Baxter

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce) Per	Subsidised	Generic Manufacturer	
Ψ	1 61		Manuacturer	

continued...

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

Tab 40 mg	126.31	25	✓ Lanvis
Other Cutotoxic Agents			

Other Cytotoxic Agents		
AMSACRINE - PCT only - Specialist		
Inj 50 mg per ml, 1.5 ml ampoule	6	✓ Amsidine S29
4,736.00		✓ Amsidine S29
Inj 75 mg1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-Specialist		
Cap 0.5 mgCBS	100	✓ Agrylin S29 S29
		✓ Teva S29
1,175.87		✓ Agrylin
(Agrylin S29 S29 Cap 0.5 mg to be delisted 1 April 2021)		
(Teva S29 Cap 0.5 mg to be delisted 1 April 2021)		
ARSENIC TRIOXIDE - PCT only - Specialist		
Inj 1 mg per ml, 10 ml vial4,817.00	10	✓ Phenasen
Inj 10 mg for ECP481.70	10 mg OP	✓ Baxter

	(Fully sidised	Brand or Generic
	\$	Per		Manufacturer
BLEOMYCIN SULPHATE - PCT only - Specialist				
Inj 15,000 iu, vial	161.01	1	✓ [BL Bleomycin
				Sulfate
Inj 1,000 iu for ECP	12.45	1,000 iu	√ E	Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority see \$	SA1889 below			
Inj 3.5 mg vial	105.00	1	✓ E	Bortezomib
, 0				Dr-Reddy's
Inj 1 mg for ECP	31.20	1 mg	✓ E	Baxter

⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.

Note: Indications marked with * are unapproved indications.

DACARBAZINE - PCT only - Specialist			
Inj 200 mg vial	62.70	1	✓ DBL Dacarbazine
	580.60	10	✓ Dacarbazine
			APP S29
Inj 200 mg for ECP	62.70	200 mg OP	✓ Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist			
Inj 0.5 mg vial	255.00	1	✓ Cosmegen
Inj 0.5 mg for ECP	255.00	0.5 mg OP	✓ Baxter
DAUNORUBICIN - PCT only - Specialist			
Inj 2 mg per ml, 10 ml	149.50	1	✓ Pfizer
Inj 20 mg for ECP		20 mg OP	✓ Baxter
DOCETAXEL - PCT only - Specialist		_	
Inj 10 mg per ml, 2 ml vial	12.40	1	✓ DBL Docetaxel
Inj 20 mg		1	✓ Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial		1	✓ DBL Docetaxel
Inj 20 mg per ml, 4 ml vial	26.95	1	✓ Docetaxel
			Accord S29
Inj 80 mg	195.00	1	✓ Docetaxel Sandoz
Inj 1 mg for ECP		1 mg	✓ Baxter
(DBL Docetaxel Inj 10 mg per ml, 2 ml vial to be delisted 1 June 2			
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial	10.00	1	✓ Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial	11.50	1	Doxorubicin Ebewe
	17.00		✓ Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial		1	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial	56.15	1	Doxorubicin Ebewe
	65.00		✓ Arrow-Doxorubicin
Inj 1 mg for ECP	0.29	1 mg	✓ Baxter
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial	25.00	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		. 1	✓ Epirubicin Ebewe
Inj 1 mg for ECP	0.43	1 mg	✓ Baxter

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
FTODOOIDE	Ψ	1 61		Manufacturer
ETOPOSIDE Pote il plant de la constaliat	040.70	00	,	Vld
Cap 50 mg – PCT – Retail pharmacy-Specialist		20		Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10		<u>Vepesid</u>
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia		1	_	Rex Medical
Inj 1 mg for ECP - PCT only - Specialist	0.09	1 mg	•	Baxter
ETOPOSIDE PHOSPHATE - PCT only - Specialist				
Inj 100 mg (of etoposide base)	40.00	1	✓	Etopophos
Inj 1 mg (of etoposide base) for ECP	0.47	1 mg	✓	Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail pha	rmacy-Specialist			
Cap 500 mg	, ,	100	/	Devatis
	31.76		/	Hydrea
Devatis to be Sole Supply on 1 February 2021				,
(Hydrea Cap 500 mg to be delisted 1 February 2021)				
IDARUBICIN HYDROCHLORIDE				
Inj 5 mg vial – PCT only – Specialist	03.00	1	1	Zavedos
Inj 10 mg vial – PCT only – Specialist		1		Zavedos
Inj 1 mg for ECP - PCT only - Specialist		1 mg	_	Baxter
		•	•	Daxiei
LENALIDOMIDE - Retail pharmacy-Specialist - Special Author	ity see SA1897 below			
Wastage claimable			_	
Cap 5 mg		28		Revlimid
Cap 10 mg		21		Revlimid
	6,207.00	28		Revlimid
Cap 15 mg	5,429.39	21		Revlimid
	7,239.18	28	✓	Revlimid
Cap 25 mg	7,627.00	21	✓	Revlimid

⇒SA1897 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Either:
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 The patient has ECOG performance score of 0-1; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	bsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

5 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MFSNA

Tab 400 mg - PCT - Retail pharmacy-Specialist	314.00	50	✓ Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist	448.50	50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist.	177.45	15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist	407.40	15	✓ Uromitexan
Inj 1 mg for ECP - PCT only - Specialist	2.96	100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist			
Inj 5 mg vial	851.37	1	✓ Teva
Inj 20 mg vial	3,275.00	1	✓ Omegapharm S29
, ,	•		✓ Teva
Inj 1 mg for ECP	288.09	1 mg	✓ Baxter
(Teva Inj 5 mg vial to be delisted 1 June 2021)		•	
MITOZANTRONE - PCT only - Specialist			
Inj 2 mg per ml, 10 ml vial	97.50	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP	5.51	1 mg	✓ Baxter
OLAPARIB - Retail pharmacy-Specialist - Special Authority s	ee SA1883 below		
Tab 100 mg	3,701.00	56	✓ Lynparza
Tab 150 mg		56	✓ Lynparza
Cap 50 mg - Wastage claimable	7,402.00	448	✓ Lynparza

⇒SA1883 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Patient has received at least two lines of previous treatment with platinum-based chemotherapy; and
- 4 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line of platinum-based chemotherapy; and
- 5 Patient's disease must have achieved partial or complete response to treatment with the immediately preceding platinum-based regimen; and
- 6 Patient's disease has not progressed following prior treatment with olaparib; and

Subsidy	F	ully	Brand or	
(Manufacturer's Pr	rice) Subsid	ised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 7 Treatment will be commenced within 8 weeks of the patient's last dose of the immediately preceding platinum-based regimen: and
- 8 Treatment to be administered as maintenance treatment; and
- 9 Treatment not to be administered in combination with other chemotherapy.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 No evidence of progressive disease; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy.

Note: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

Inj 30 mg	47.30	5	✓ Paclitaxel Ebewe
Inj 100 mg		1	✓ Paclitaxel Ebewe
, ,	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
,	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg	44.00	1	✓ Paclitaxel Ebewe
, .	275.00		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	✓ Baxter
EGASPARGASE - PCT only - Special Authority s	ee SA1979 below		
Inj 750 iu per ml, 5 ml vial	3.455.00	1	✓ Oncaspar LYO S29

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only -	Specialist		
Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail	pharmacy-Specialist		
Cap 50 mg	980.00	50	✓ Natulan S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
TEMOZOLOMIDE - Special Authority see SA1741 below - Reta	ail pharmacy			
Cap 5 mgCap 20 mg	9.13 16.38	5 5	1	Temaccord Temaccord
Cap 100 mg	18.30 136.00 35.98 40.20	14 5	✓ <u>/</u>	Apo-Temozolomide Accord ⁸²⁹ <u>Temaccord</u> Apo-Temozolomide
Cap 140 mg	532.00	14 5	✓ [Accord S29 Temaccord Amneal S29
Cap 180 mg		14 5	1	Accord §29 Temaccord Amneal §29

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 No evidence of disease progression; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRFTINOIN

Cap 10 mg - PCT - Retail pharmacy-Specialist	100	✓ Vesanoid
VENETOCLAX - Retail pharmacy-Specialist - Special Authority see SA1868 be	elow	
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	42 OP	Venclexta
Tab 10 mg95.78	14 OP	✓ Venclexta
Tab 50 mg239.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable8,209.41	120	✓ Venclexta

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

continued...

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

VINBLASTINE SULPHATE

Inj 1 mg per mi, 10 mi viai – PCT – Retail pharmacy-Specialist2/0.3/	5	✓ DBL Vinblastine \$29
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Hospira✓ Baxter
Inj 1 mg per ml, 1 ml vial – PCT – Retail pharmacy-Specialist74.52	5	✓ DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist 102.73	5	✓ DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	✓ Baxter
VINORELBINE - PCT only - Specialist		
Inj 10 mg per ml, 1 ml vial	1	✓ Navelbine✓ Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial56.00 210.00	1	✓ Navelbine✓ Vinorelbine Ebewe
Inj 1 mg for ECP1.25	1 mg	✓ Baxter

Protein-tyrosine Kinase Inhibitors

ALECTINIB - Retail pharmacy-Specialist - Special Authority see SA1870 below Wastage claimable

224 Alecensa

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid

Subsidy		Fully	Brand or	
(Manufacturer's Pr	ice) S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA1805 below - Retail pharmacy

Wastage claimable	•		
Tab 20 mg	3,774.06	60	✓ Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
Tab 70 mg	7,692.58	60	✓ Sprycel

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
 - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
 - 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*: and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

⇒SA1915 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and

Subsidy (Manufacturer's Price)	Fully Subsidised	
 \$	Per 🗸	Manufacturer

continued...

- 3 Either:
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued defitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

GEFITINIB – Retail pharmacy-Specialist – Special Authority see S	A1916 below		
Tab 250 mg	1,700.00	30	✓ Iressa

⇒SA1916 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
 - 2.1 Patient is treatment naive: or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib: and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Gefitinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

IMATINIB MESII ATE

Note: Imatinib-AFT is not a registered for the treatment of Gastro Intestinal Stromal Tumours (GIST). The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule.

Tab 100 mg - [Xpharm] - Special Authority see SA1460 on the

	next page2,4	00.00	0 •	✓ Glivec
*	Cap 100 mg		0 •	/ Imatinib-Rex
	•	98.00	•	/ Imatinib-AFT
*	Cap 400 mg	84.79 3	80	/ Imatinib-Rex
		97.50	•	/ Imatinib-AFT

(Imatinib-AFT Cap 100 mg to be delisted 1 June 2021) (Imatinib-AFT Cap 400 mg to be delisted 1 June 2021)

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from PHARMAC's website <u>schedule.pharmac.govt.nz/SAForms</u>, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 PHARMAC Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

Special Authority criteria for GIST - access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA1191 below - Retail pharmacy

(Tykerb Tab 250 mg to be delisted 1 June 2021)

SA1191 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient has not previously received trastuzumab treatment for HER 2 positive metastatic breast cancer; and
 - 1.3 Lapatinib not to be given in combination with trastuzumab; and
 - 1.4 Lapatinib to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2.2 The patient started trastuzumab for metastatic breast cancer but discontinued trastuzumab within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on trastuzumab; and
 - 2.4 Lapatinib not to be given in combination with trastuzumab; and
 - 2.5 Lapatinib to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 on the next page - Retail pharmacy

Wastage claimable

Cap 150 mg4,680.00	120	Tasigna
Cap 200 mg6,532.00	120	Tasigna

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Fither:
 - 2.1 Patient has documented CML treatment failure* with imatinib; or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PALBOCICLIB - Retail pharmacy-Specialist - Special Authority see SA1894 below

wastage ciaimable			
Cap 75 mg	4,000.00	21	✓ Ibrance
Cap 100 mg	4,000.00	21	✓ Ibrance
Cap 125 mg	4.000.00	21	✓ Ibrance

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Either:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state: and
- 4.2.2 Fither:
 - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
 - 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PAZOPANIB – Special Authority see SA1190 below – Retail pha	ırmacy			
Tab 200 mg	1,334.70	30	✓ \	/otrient
Tab 400 mg	2,669.40	30	•	/otrient

SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

wastage cialmable			
Tab 5 mg	2,500.00	56	Jakavi
Tab 15 mg	5,000.00	56	Jakavi
Tab 20 mg	5.000.00	56	✓ Jakavi

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis: and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

- 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS: and
- 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

Cap 12.5 mg2,315.38	28	✓ Sutent
Cap 25 mg	28	✓ Sutent
Cap 50 mg	28	✓ Sutent

⇒SA1917 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib: or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 84

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA1914 below

Wastage claimable

⇒SA1914 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases: and
- 3 Patient's disease is castration resistant; and
- 4 Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)		sidised	Generic
 5	Per		Manutacturer

continued...

- 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
- 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of clinical disease progression; and
- 2 No initiation of taxane chemotherapy with abiraterone; and
- 3 The treatment remains appropriate and the patient is benefiting from treatment.

BICALUTAMIDE

Tab 50 mg	4.07 4.21	30 28	✓ Binarex✓ Binarex
Binarex to be Sole Supply on 1 April 2021			
FLUTAMIDE			
Tab 250 mg	119.50	100	Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Special Author	ity see SA1895 belo	OW	
Inj 50 mg per ml, 5 ml prefilled syringe	1,068.00	2	✓ Faslodex

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

MEGESTROL ACETATE

Tab 160 mg	63.53	30	✓ Apo-Megestrol
------------	-------	----	-----------------

	Subsidy (Manufacturer's Price) \$	S Per	Fully Brand or Subsidised Generic Manufacturer
OCTREOTIDE			
Inj 100 mcg per ml, 1 ml ampoule	18.69	5	✓ Octreotide GH S29
Inj 50 mcg per ml, 1 ml ampoule	30.64	5	✓ Octreotide GH S29
Inj 50 mcg per ml, 1 ml vial		5	✓ DBL Octreotide ✓ Octreotide MaxRx (\$29)
Inj 100 mcg per ml, 1 ml vial	18.69	5	✓ DBL Octreotide
Inj 500 mcg per ml, 1 ml ampoule		5	✓ Octreotide GH S29
Inj 500 mcg per ml, 1 ml vial		5	✓ DBL Octreotide ✓ Octreotide (Sun) \$29
OCTREOTIDE LAR (SOMATOSTATIN ANALOGUE) - Special A	Authority see SA1918	below	– Retail pharmacy
Inj LAR 10 mg prefilled syringe	1,772.50	1	✓ Sandostatin LAR
Inj LAR 20 mg prefilled syringe		1	Sandostatin LAR
Inj LAR 30 mg prefilled syringe	2,951.25	1	Sandostatin LAR

⇒SA1918 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed: or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Initial application — **(Other Indications)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Any of the following:

60

60

✓ <u>Tamoxifen Sandoz</u>
 ✓ Tamoxifen Sandoz

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Fither:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

TAMOXIFEN CITRATE

Aromatase Inhibitors		
ANASTROZOLE * Tab 1 mg	30	✓ Anatrole ✓ Rolin
EXEMESTANE * Tab 25 mg14.50 LETROZOLE	30	✓ Pfizer Exemestane
* Tab 2.5 mg4.68	30	✓ <u>Letrole</u>

Immunosuppressants

Cytotoxic Immunosuppressants

AZATHIOPRINE

*	Tab 25 mg	7.35	60	Azamun
	Tab 50 mg		100	✓ Azamun
	Inj 50 mg vial199		1	✓ Imuran

	(Fully	Brand or Generic	
	\$	Per		Manufacturer	
MYCOPHENOLATE MOFETIL					
Tab 500 mg	35.90	50	✓ (Cellcept	
Cap 250 mg	35.90	100	✓ (Cellcept	
Powder for oral lig 1 g per 5 ml - Subsidy by endorsement	187.25	165 ml OP	✓ (Cellcept	
Mycophenolate powder for oral liquid is subsidised only	for patients unab	le to swallow to	ablets a	and capsules, and when	
the prescription is endorsed accordingly.	•			•	

Fusion Proteins

ETANERCEPT - Special Authority see SA1974 below - Retail pharmacy		
Inj 25 mg690.00	4	Enbrel
Inj 50 mg autoinjector	4	✓ Enbrel
Inj 50 mg prefilled syringe1,050.00	4	✓ Enbrel

⇒SA1974 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Roth:
 - 1.1 Fither:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

```
18-24 years - Male: 7.0 cm; Female: 5.5 cm
25-34 years - Male: 7.5 cm; Female: 5.5 cm
35-44 years - Male: 6.5 cm; Female: 4.5 cm
45-54 years - Male: 6.0 cm; Female: 5.0 cm
55-64 years - Male: 5.5 cm; Female: 4.0 cm
65-74 years - Male: 4.0 cm; Female: 4.0 cm
75+ years - Male: 3.0 cm; Female: 2.5 cm
```

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or

Subsidy	Fully	Brand or
(Manufacturer's		Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
\$	Per	1	Manufacturer

continued...

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for psoriatic arthritis; and
 - 1.2 Fither
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for psoriatic arthritis: or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Fither:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Subsidy	Fu	ılly	Brand or	
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold: or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints;
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Fither:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

All of the following:

- 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Fither:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose): and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose): and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Fither

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
\$	Per	•	Manufacturer

continued...

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Specia	list		
Inj 50 mg per ml, 5 ml	2,351.25	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT only	Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG S29
(SII-Onco-BCG S29 Inj 40 mg per ml, vial to be delisted 1 April	2022)		

Monoclonal Antibodies

		pelow – Retail pharmacy	ADALIMUMAB – Special Authority see SA1975 bel
Humira	2	1,599.96	Inj 20 mg per 0.4 ml prefilled syringe
✓ HumiraPen	2	1,599.96	Inj 40 mg per 0.8 ml prefilled pen
✓ Humira	2	1.599.96	Ini 40 mg per 0.8 ml prefilled syringe

⇒SA1975 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — **(ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm

25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm

35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm

55-64 years - Male: 5.5 cm; Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — **(ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	
\$	Per 🗸	Manufacturer

continued...

- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or

Subsidy (Manufacturer's Price)	Fully Subsidised		
\$	Per 🗸	Manufacturer	

continued...

- 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Fither:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 CDAI score is 150 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — **(Crohn's disease - children)** only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 PCDAI score is 15 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 The patient has 3 or more active lesions (e.g. inflammatory nodules, abscesses, draining fistulae); and
- 4 The patient has a Dermatology Quality of Life Index of 10 or more and the assessment is no more than 1 month old at time of application: and
- 5 Following the initial loading doses, adalimumab is to be administered at doses no greater than 40mg every 7 days.

Renewal — (hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline: and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Fither:

Subsidy (Manufactureria Price)		Fully	Brand or
 (Manufacturer's Price) \$	Subside Per	JISEO 🗸	Generic Manufacturer

continued...

- 1.2.1 The patient has experienced intolerable side effects from etanercept; or
- 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline: or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA): and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Fither:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist.

Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold: or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

continued...

2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease that is significantly impacting the patient's quality of life (see Notes); and
- 2 Fither:
 - 2.1 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has not responded adequately to treatment with infliximab (see Notes); or
 - 2.2 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has experienced intolerable side effects from treatment with infliximab; and
- 3 The patient is experiencing significant loss of quality of life; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: Behcet's disease diagnosed according to the International Study Group for Behcet's disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al, J Rheumatol. 2004;31:931-7.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plague psoriasis; and
 - 12 Fither
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

continued...

1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or

- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plague psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis: or
 - 2.1.2 Patient has severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot, where the plague or plagues have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline valuee; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
 - 2.2.2 Fither:
 - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values: or
 - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value: and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Any of the following:

- 1.1 The patient has had a good clinical response following 3 initial doses; or
- 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

✓ Eylea

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
- 1.3 There is no structural damage to the central fovea of the treated eye; and
- 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB – PCT only – Special Authority see SA1982 of	on the next page		
Inj 100 mg	806.00	1	✓ Remicade
Inj 1 mg for ECP	8.29	1 mg	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1982 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

continued...

considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Graft vs host disease) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria: Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation: or

2 Roth:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss: and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose: or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Roth:

- 1 Patient has confirmed Crohn's disease; and
- 2 Fither:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Fither:

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

- 2.3.1 There has been an improvement in MRI appearances; or
- 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria: Fither:

- 1 Roth:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plaque psoriasis: and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plague psoriasis; or
- 2 All of the following:
 - 2.1 Fither:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis: or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plague or plagues have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

1 Fither

Both:

- 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values: or

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis: or
 - 2.4 Severe ocular inflammation; or
 - 2.5 Chronic ocular inflammation; or
 - 2.6 Crohn's disease (adults): or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease: or
 - 2.9 Severe fulminant ulcerative colitis: or
 - 2.10 Severe ulcerative colitis; or
 - 2.11 Plaque psoriasis: or
 - 2.12 Neurosarcoidosis; or
 - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for psoriatic arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- - 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
 - - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and

Subsidy (Manufacturer's Price)	Fully Subsidised	
 \$	Per 🗸	Manufacturer

continued...

3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis: and
- 2 Either:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
 - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement: and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

⇒SA1896 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded: and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10⁹ cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids: or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
OBINUTUZUMAB - PCT only - Specialist - Special Authority se	ee SA1627 helow	rei		Manufacturer
Inj 25 mg per ml, 40 ml vial	5,910.00	1 1 mg		Gazyva Baxter
CA1627 Capaigl Authority for Subaidy	0.21	ı ıııg	•	Daxiei

SA1627 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and</p>
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

OMALIZUMAB - Special Authority see SA1744 below - Reta	ail pharmacy		
Inj 150 mg prefilled syringe	450.00	1	✓ Xolair
Inj 150 mg vial	450.00	1	✓ Xolair

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Fither:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient must be aged 12 years or older; and

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PERTUZUMAB - PCT only - Specialist - Special Authority see SA1606 below

Inj 30 mg per ml, 14 ml vial	3,927.00	1	Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	Baxter

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and

Subsidy)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist -	- Special Authority see SA1976	6 below		
Inj 100 mg per 10 ml vial	1,075.50	2	Mab	hera
Inj 500 mg per 50 ml vial	2,688.30	1	Mab	hera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxt	er (Mabthera)

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Fither:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

1 Both:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis: and
- 2 Fither:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician: or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA1937 on the next page

inj 100 mg per 10 mi viai	2/5.33	2	✓ <u>RIXIMYO</u>
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

⇒SA1937 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g or a further repeat 3 month induction
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4.1 The patient does not have chromosome 17p deletion CLL; or
- 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Roth:

1 Either:

- 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
- 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and

Subsidy	Fully	Brand or	_
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

1 Patient is a child with SDNS* or FRNS*; and

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	1	

continued...

- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Subsidy	-	ully	Brand or	
(Manufacturer's Price)	Subsid	ised	Generic	
\$	Per	✓	Manufacturer	

continued...

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
 - 2 An initial response lasting at least 12 months was demonstrated; and
 - 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre: or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	•	Manufacturer	

continued...

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks: and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Subsidy (Manufacturer's Price)	Suk	Fully	Brand or Generic
 \$	Per	√	Manufacturer

continued...

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or 2 All of the following:
- - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*: and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 × 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Fither:
 - 2.1 Both
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

SECUKINUMAB - Special Authority see SA1754 below - Retail pharmacy

⇒SA1754 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab: or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

	Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic	
	\$	Per	1	Manufacturer	
SILTUXIMAB - Special Authority see SA1596 below - Retail ph	armacy				
Note: Siltuximab is to be administered at doses no greater t	han 11 mg/kg every 3	weeks.			
Inj 100 mg vial	770.57	1	✓ S	ylvant	
Inj 400 mg vial	3,082.33	1	✓ S	ylvant	

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB - PCT only - Special Authority see SA1977 below

Inj 20 mg per ml, 4 ml vial2	20.00 1	✓ Actemra
Inj 20 mg per ml, 10 ml vial5		✓ Actemra
Inj 20 mg per ml, 20 ml vial	00.00 1	✓ Actemra
Inj 1 mg for ECP	2.85 1 m	g Saxter

⇒SA1977 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial: and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis: or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease: or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis: and
- 3 Fither:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules: and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Fither:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Fither:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints:
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Bules of the Pharmaceutical Schedule: and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

Sub: (Manufactu	,	,	
\$	Per	✓ Manufacturer	

continued...

2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB - PCT only - Specialist - Special Authority see SA1632 below

Inj 150 mg vial	1,350.00	1	Herceptin
Inj 440 mg vial	3,875.00	1	✓ Herceptin
Inj 1 mg for ECP		1 mg	✓ Baxter

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
	Per 🗸	Manufacturer

continued...

- 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully idised	I Generic
TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special	Authority see SA187	'1 below		
Inj 100 mg vial	2,320.00	1	1	Kadcyla
Inj 160 mg vial	3,712.00	1	1	Kadcyla
Inj 1 mg for ECP	23.20	1 mg	1	Baxter

⇒SA1871 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease*: or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Fither:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Treatment to be discontinued at disease progression.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

Programmed Cell Death-1 (PD-1) Inhibitors

NIVOLUMAB – PCT only – Specialist – Special Autho	rity see SA1911 below		
Inj 10 mg per ml, 4 ml vial	1,051.98	1	Opdivo
Inj 10 mg per ml, 10 ml vial	2,629.96	1	✓ Opdivo
Inj 1 mg for ECP	27.62	1 mg	✓ Baxter

⇒SA1911 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1: and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Fither:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB - PCT only - Specialist - Special Authority see SA1910 below

⇒SA1910 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV: and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab

(Manu	Subsidy	Fully	Brand or
	facturer's Price)	Subsidised	Generic
	\$ P	Per 🗸	Manufacturer

continued...

within 12 weeks of starting treatment due to intolerance; and

- 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

44.63	50	Neoral
88.91	50	Neoral
	50	✓ Neoral
	50 ml OP	✓ Neoral
	44.63 88.91 177.81 198.13	88.91 50 177.81 50

	Subsidy (Manufacturer's Price) \$		Fully ubsidised	Generic
EVEROLIMUS – Special Authority see SA1913 below – Retail ph Wastage claimable	armacy			
Tab 10 mg	6,512.29	30	1	Afinitor
Tab 5 mg	4,555.76	30	1	Afinitor

⇒SA1913 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Everolimus to be discontinued at progression of SEGAs; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Note: : MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

SIROLIMUS - Special Authority see SA0866 below - Retail pharmacy

Tab 1 mg	749.99	100	Rapamune
Tab 2 mg	1,499.99	100	✓ Rapamune
Oral liq 1 mg per ml	449.99	60 ml OP	Rapamune

⇒SA0866 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- · Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis: or
- . HUS or TTP: or
- · Leukoencepthalopathy; or
- Significant malignant disease

TACROLIMIIS	- Special Authority see	SA1745 on the next page -	- Retail nharmacy

Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg	99.30	100	✓ Tacrolimus Sandoz
Cap 1 mg	84.30	100	✓ Tacrolimus Sandoz
Cap 5 mg	248.20	50	✓ Tacrolimus Sandoz

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	Manufacturer

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antiallergy Preparations

Allergic Emergencies

ICATIBANT – Special Authority see SA1558 below – Retail pharmacy
Inj 10 mg per ml, 3 ml prefilled syringe.......2,668.00 1 ✓ Firazyr

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above -	Retail pharma	су
Initiation kit - 5 vials freeze dried venom with diluent305.00	1 OP	✓ VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent305.00	1 OP	✓ VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with		
diluent285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent		
9 ml, 3 diluent 1.8 ml305.00	1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent 305.00	1 OP	✓ Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority see SA1367 above	- Retail pharn	nacy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze		·
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze		
dried venom, with diluent305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze		
dried venom, with diluent305.00	1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze		
dried venom, with diluent305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze		
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml305.00	1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze		
dried venom, with diluent305.00	1 OP	✓ Venomil S29

	Subsidy		Fully Brand or
	(Manufacturer's Pr		
	\$	Per	✓ Manufacturer
Authistonius			
Antihistamines			
CETIRIZINE HYDROCHLORIDE			
* Tab 10 mg	1.12	100	✓ Zista
* Oral lig 1 mg per ml		200 ml	✓ Histaclear
CHLORPHENIRAMINE MALEATE			
* Oral liq 2 mg per 5 ml	9 37	500 ml	✓ Histafen
		300 1111	Tilotalen
DEXTROCHLORPHENIRAMINE MALEATE	0.00	40	
* Tab 2 mg		40	Delevenine
	(8.40)	20	Polaramine
	1.01	20	Polaramine
* Oral liq 2 mg per 5 ml	(5.99)	100 ml	Polaramine
* Oral liq 2 mg per 5 ml		100 1111	Polaramine
	(10.29)		Polarattille
FEXOFENADINE HYDROCHLORIDE			
* Tab 60 mg		20	-
	(8.23)		Telfast
* Tab 120 mg		10	T 16 .
	(8.23)	00	Telfast
	14.22	30	Talfast
	(26.44)		Telfast
LORATADINE			
* Tab 10 mg		100	✓ <u>Lorafix</u>
* Oral liq 1 mg per ml	2.95	120 ml	✓ Lorfast
PROMETHAZINE HYDROCHLORIDE			
* Tab 10 mg	1.68	50	✓ Allersoothe
* Tab 25 mg		50	✓ Allersoothe
* Oral liq 1 mg per 1 ml	2.69	100 ml	✓ Allersoothe
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a	PSO 17.87	5	✓ Hospira
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler, 50 mcg per dose	0.20	200 dose OP	✓ Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OF	✓ Beclazone 50
Aerosol inhaler, 30 mcg per dose or o-nee		200 dose OP	✓ Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP	✓ Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free		200 dose OP	✓ Beclazone 250
		200 0000 01	5 Deciazone 200
BUDESONIDE Boundar for inhelation, 100 mag par door	17.00	200 docs OD	√ Dulminort
Powder for inhalation, 100 mcg per dose	17.00	200 dose OP	✓ Pulmicort Turbuhaler
Douglas for inhalation 200	10.00	000 das- OD	
Powder for inhalation, 200 mcg per dose	19.00	200 dose OP	✓ Pulmicort
D 1 ('11 1 " (00)	00.00	200 1 65	Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00	200 dose OP	✓ Pulmicort
			Turbuhaler

	Subsidy		Fully B	rand or
	(Manufacturer's	Price) Subs	idised G	eneric
	\$	Per	✓ M	anufacturer
FLUTICASONE				
Aerosol inhaler, 50 mcg per dose	7.19	120 dose OP	✓ Flixo	tide
Powder for inhalation, 50 mcg per dose		60 dose OP		tide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP		tide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP	✓ Flixo	
Aerosol inhaler, 250 mcg per dose	24.62	120 dose OP	✓ Flixo	
Powder for inhalation, 250 mcg per dose		60 dose OP		tide Accuhaler
,,				
Inhaled Long-acting Beta-adrenoceptor Agonis	ts			
EFORMOTEROL FUMARATE				
	ioo 20.64	60 dose		
Powder for inhalation, 12 mcg per dose, and monodose dev		ou dose	Fora	انام
	(35.80)		rora	JII
EFORMOTEROL FUMARATE DIHYDRATE				
Powder for inhalation 4.5 mcg per dose, breath activated				
(equivalent to eformoterol fumarate 6 mcg metered dose	e) 10.32	60 dose OP		
	(16.90)		Oxis	Turbuhaler
INDACATEROL				
Powder for inhalation 150 mcg	61.00	30 dose OP	Onbi	rez Breezhaler
Powder for inhalation 300 mcg	61.00	30 dose OP	Onbi	rez Breezhaler
SALMETEROL				
Aerosol inhaler CFC-free, 25 mcg per dose	25.00	120 dose OP	✓ Sere	vent
Powder for inhalation, 50 mcg per dose, breath activated		60 dose OP		vent Accuhaler
Torraci is initiation, so may per uses, stouth astronous				
Inhaled Corticosteroids with Long-Acting Beta-	-Adrenocept	or Agonists		
BUDESONIDE WITH EFORMOTEROL				
Powder for inhalation 160 mcg with 4.5 mcg eformoterol				
fumarate per dose (equivalent to 200 mcg budesonide v	with			
6 mcg eformoterol fumarate metered dose)		120 dose OP	✓ Duol	Resp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fuma		120 0000 01	· Duoi	toop opnomax
per dose (equivalent to 400 mcg budesonide with 12 mc				
eformoterol fumarate metered dose) — No more than 2	<i>-</i> y			
dose per day	82 50	120 dose OP	✓ Duol	Resp Spiromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg		120 dose OP	✓ Vanr	
Powder for inhalation 100 mcg with eformoterol furnarate 6 r		120 dose OP	✓ Sym	
1 owder for initial attorn 100 they with elormoteror furnarate of	11cg55.74	120 0036 01		rbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg	21 //0	120 dose OP	✓ Vanr	
Powder for inhalation 200 mcg with eformoterol furnarate 6 mcg		120 dose OP	✓ Sym	
1 owder for initial auton 200 integ with elothioteror idilial ate 6 i	110g 14 .00	120 0086 OF		rbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate			ıu	. Salialoi 200/0
12 mcg - No more than 2 dose per day	// NO	60 dose OP	✓ Sym	hicart
12 mg - No more man 2 dose per day	44.00	ou dose OF		rbuhaler 400/12
			ıu	Dulidici 400/12
FLUTICASONE FUROATE WITH VILANTEROL				
Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose OP	✓ Breo	Ellipta

	Subsidy (Manufacturer's \$		Fully sidised	Brand or Generic Manufacturer
LUTICASONE WITH SALMETEROL Aerosol inhaler 50 mcg with salmeterol 25 mcg	25 79	120 dose OP	_	Seretide
Aerosol inhaler 125 mcg with salmeterol 25 mcg	32.60	120 dose OP		Seretide Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg - 1 more than 2 dose per day		60 dose OP	•	Seretide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg - 1 more than 2 dose per day		60 dose OP	•	Seretide Accuhaler
Beta-Adrenoceptor Agonists				
ALBUTAMOL				
Oral liq 400 mcg per ml	20.00	150 ml		Ventolin
Infusion 1 mg per ml, 5 ml	53.00	10 5		Ventolin Ventolin
Inhaled Beta-Adrenoceptor Agonists				
ALBUTAMOL				
Aerosol inhaler, 100 mcg per dose CFC free - Up to 1000				
dose available on a PSO	3.80	200 dose OP		Respigen SalAir
	(6.00)			Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 n- available on a PSO		20	,	Asthalin
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 n		20	•	Astriaini
available on a PSO	4.03	20	✓	<u>Asthalin</u>
ERBUTALINE SULPHATE				
Powder for inhalation, 200 mcg per dose (equivalent to 250 mcg metered dose), breath activated	22.20	120 dose OP	•	Bricanyl Turbuhaler
Anticholinergic Agents				
PRATROPIUM BROMIDE				
Aerosol inhaler, 20 mcg per dose CFC-free	16.20	200 dose OP	1	Atrovent
a) Up to 400 dose available on a PSO				
 b) No patient co-payment payable Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 	nah			
available on a PSO		20	•	<u>Univent</u>
Inhaled Beta-Adrenoceptor Agonists with Ant	icholinergic	Agents		
ALBUTAMOL WITH IPRATROPIUM BROMIDE				
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mc dose CFC-free		200 dose OP	/	Duolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg pe				
vial, 2.5 ml ampoule - Up to 20 neb available on a PS	SO5.20	20	1	<u>Duolin</u>

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

\$ Per ✔ Manufacturer

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium
- Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD, and the prescription is endorsed accordingly.

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL — Special Authority see SA1584 above — Retail pharmacy
Powder for Inhalation 50 mcg with indacaterol 110 mcg......81.00 30 dose OP ✓ Ultibro Breezhaler

TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 above - Retail pharmacy

Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg81.00 60 dose OP ✓ Spiolto Respimat

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

Antifibrotics

NINTEDANIB - Special Authority see SA1928 on the next page - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

 Cap 100 mg
 2,554.00
 60 OP
 ✓ Ofev

 Cap 150 mg
 3,870.00
 60 OP
 ✓ Ofev

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

				
	Subsidy	Fully	Brand or	
	(Manufacturer's Price)	Subsidised	Generic	
	\$	Per 🗸	Manufacturer	

⇒SA1928 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA1929 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Tab 801 mg	3,645.00	90	Esbriet
Cap 267 mg - Wastage claimable	3,645.00	270	Esbriet

⇒SA1929 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

Leukotriene Receptor Antagonists

МО	NTELUKAST		
*	Tab 4 mg4.25	28	✓ Montelukast Mylan
	Tab 5 mg4.25	28	✓ Montelukast Mylan
*	Tab 10 mg3.95	28	✓ Montelukast Mylan

Mast Cell Stabilisers

NEDOCROMIL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking nedocromil prior to 1 July 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of nedocromil.

Aerosol inhaler, 2 mg per dose CFC-free.......28.07 112 dose OP ✓ Tilade

(Tilade Aerosol inhaler, 2 mg per dose CFC-free to be delisted 1 February 2021)

SODIUM CROMOGLICATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking sodium cromoglicate prior to 1 July 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of sodium cromoglicate.

Methylxanthines

AMINOPHYLLINE

*	Inj 25 mg per ml, 10 ml ampoule - Up to 5 inj available on a		
	PSO124.37	5	✓ DBL Aminophylline
TH	EOPHYLLINE		
*	Tab long-acting 250 mg23.02	100	✓ Nuelin-SR
*	Oral lig 80 mg per 15 ml 16.60	500 ml	✓ Nuelin

Mucolytics

DORNASE ALFA – Special Authority see SA1978 below – Reta	ail pharmacy		
Nebuliser soln, 2.5 mg per 2.5 ml ampoule	250.00	6	✓ Pulmozyme

⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

	Subsidy	D: \	Fully Brand or
	(Manufacturer's \$	Price) Subsi	idised Generic Manufacturer
ODIUM CHLORIDE			
Not funded for use as a nasal drop.			
Soln 7%	24.50	90 ml OP	✓ Biomed
Nasal Preparations			
Allergy Prophylactics			
UDESONIDE			
Metered aqueous nasal spray, 50 mcg per dose	2.54	200 dose OP	✓ SteroClear
Metered aqueous nasal spray, 100 mcg per dose		200 dose OP	✓ SteroClear
LUTICASONE PROPIONATE			
Metered aqueous nasal spray, 50 mcg per dose	1.98	120 dose OP	✓ Flixonase Hayfever
			<u>& Allergy</u>
PRATROPIUM BROMIDE	5.00	45 1 0 D	∠ Habaant
Aqueous nasal spray, 0.03%	5.23	15 ml OP	✓ Univent
Univent to be Sole Supply on 1 April 2021			
Respiratory Devices			
ASK FOR SPACER DEVICE			
a) Up to 50 dev available on a PSO			
b) Only on a PSO			
 c) Only for children aged six years and under 			
Small	2.20	1	e-chamber Mask
EAK FLOW METER			
a) Up to 25 dev available on a PSO			
b) Only on a PSO			_
Low range	9.54	1	✓ Mini-Wright AFS
	0.54		Low Range
Normal range	9.54	1	✓ Mini-Wright Standard
DA OED DEVIOE			Standard
PACER DEVICE			
a) Up to 50 dev available on a PSO			
b) Only on a PSO 220 ml (single patient)	2 95	1	✓ e-chamber Turbo
510 ml (single patient)		1	✓ e-chamber La
o to the following patients)		•	Grande
800 ml	6.50	1	✓ Volumatic
Deceminatory Stimulants			
Respiratory Stimulants			
AFFEINE CITRATE			
Oral liq 20 mg per ml (10 mg base per ml)		25 ml OP	

✓ Locacorten-Viaform ED's✓ Locorten-Vioform

	Subsidy (Manufacturer's Price \$	e) Sub Per	Fully esidised	Brand or Generic Manufacturer	
Ear Preparations					
ACETIC ACID WITH 1, 2- PROPANEDIOL DIACETATE AND BE For Vosol ear drops with hydrocortisone powder refer Standa Ear drops 2% with 1, 2-Propanediol diacetate 3% and benzethonium chloride 0.02%	ard Formulae, page	246 35 ml OP	√ Vo	osol	
FLUMETASONE PIVALATE			•		

Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate

Ear/Eye Preparations

DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN

Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and gramicidin 50 mcg per ml4.50

..4.50 8 ml OP (9.27) Sofradex

7.5 ml OP

FRAMYCETIN SULPHATE

8.65) Soframycin

Eye Preparations

Eye preparations are only funded for use in the eye, unless explicitly stated otherwise.

Anti-Infective Preparations

Tita inioonivo i roparationo			
ACICLOVIR * Eye oint 3%	14.92	4.5 g OP	✓ ViruPOS
CHLORAMPHENICOL		3 -	
Eye oint 1%	1.55	5 g OP	✓ Devatis
Eye drops 0.5%	1.54	10 ml OP	✓ Chlorafast
Funded for use in the ear*. Indications marked with '	are unapproved in	dications.	
CIPROFLOXACIN			
Eye drops 0.3% - Subsidy by endorsement	12.15	5 ml OP	✓ Ciprofloxacin Teva
When prescribed for the treatment of bacterial keratit for the second line treatment of chronic suppurative on Note: Indication marked with a * is an unapproved in	titis media (CSOM)	,	
GENTAMICIN SULPHATE			
F 0.00/	44.40	5 I OD	/ A

Eye drops 0.3%	11.40	5 ml OP	✓ Genoptic
PROPAMIDINE ISETHIONATE			

(14.55) Brolene

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Pr	rice) Sub	Fully	Brand or Generic
	\$	Per	1	Manufacturer
TOBRAMYCIN				
Eye oint 0.3%	10.45	3.5 g OP	✓ T	obrex
Eye drops 0.3%	11.48	5 ml OP	√ T	obrex
Corticosteroids and Other Anti-Inflammatory P	reparations			
DEXAMETHASONE				
* Eye oint 0.1%	5.86	3.5 g OP	✓ N	/laxidex
* Eye drops 0.1%	4.50	5 ml OP	✓ N	// Axidex
Ocular implant 700 mcg - Special Authority see SA1680 be				

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient has diabetic macular oedema with pseudophakic lens; and

- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not vet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

*	Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per g	5.39	3.5 a OP	✓ Maxitrol
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml		5 ml OP	✓ Maxitrol
	LOFENAC SODIUM Eye drops 0.1%	13.80	5 ml OP	✓ Voltaren Ophtha

✓ Ozurdex

	Subsidy (Manufacturer's Pri	ce) Su Per	Fully bsidised	Brand or Generic Manufacturer	
FLUOROMETHOLONE					
* Eye drops 0.1%	3.09	5 ml OP	√ F	:ML	
,	5.20		√ F	lucon	
KETOROLAC TROMETAMOL - Special Authority see SA1981	below - Retail pha	rmacy			
Eye drops 0.5%		5 ml OP	✓ A	cular	
On cold Analysis for Only day					

SA1981 Special Authority for Subsidy

Initial application — (macular oedema) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 The patient has established post-operative or inflammatory (uveitic) cystoid macular oedema; or
- 2 Both:
 - 2.1 The patient is at risk of postoperative macular oedema; and
 - 2.2 The patient has had, or is scheduled to have imminent cataract surgery.

LEVOCABASTINE

Eye drops 0.5 mg per ml	8.71	4 ml OP	
	(10.34)		Livostin
LODOXAMIDE			
Eye drops 0.1%	8.71	10 ml OP	✓ Lomide
NEPAFENAC			
Eye drops 0.3%	13.80	3 ml OP	✓ Ilevro
PREDNISOLONE ACETATE			
Eye drops 1%	5.93	10 ml OP	✓ Prednisolone-AFT
	7.00	5 ml OP	✓ Pred Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority s	see SA1715 below	- Retail pharn	nacy
Eye drops 0.5%, single dose (preservative free)	38.50	20 dose	Minims
			Prednisolone

⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

5 ml OP

2.5 ml OP

Rexacrom

✓ Timoptol XE

SODIUM CROMOGLICATE

Glaucoma Preparations - Beta Blockers		
BETAXOLOL * Eye drops 0.25%	5 ml OP 5 ml OP	✓ Betoptic S✓ Betoptic
TIMOLOL ** Eve drapa 0.259/	5 ml OP	✓ Arrow-Timolol
* Eye drops 0.25%	5 ml OP	✓ Arrow-Timolol

	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully Brand didised Generical Manufa	0
Glaucoma Preparations - Carbonic Anhydrase	nhibitors			
ACETAZOLAMIDE * Tab 250 mg	17.03	100	✓ Diamox	
BRINZOLAMIDE	17.00	100	Diamox	
* Eye drops 1%	9.77	5 ml OP	✓ Azopt	
DORZOLAMIDE HYDROCHLORIDE				
* Eye drops 2%		5 ml OP	Trucant	
DORZOLAMIDE WITH TIMOLOL	(17.44)		Trusopt	
* Eye drops 2% with timolol 0.5%	2.87	5 ml OP	✓ Dortimor	ot
Glaucoma Preparations - Prostaglandin Analog				_
	400			
BIMATOPROST * Eve drops 0.03%	3.30	3 ml OP	✓ Bimatop	nst
		0 1111 01	Multich	
ATANOPROST				
* Eye drops 0.005%	1.57	2.5 ml OP	✓ Teva	
FRAVOPROST ★ Eye drops 0.004%	7.00	E ml OD	-/ Tuoyont	
* Eye drops 0.004%	10.50	5 ml OP	✓ Travopt ✓ Mylan S29	
	19.50	2.5 ml OP	✓ Travatan	
Glaucoma Preparations - Other				
BRIMONIDINE TARTRATE				
* Eye drops 0.2%	12.25	5 ml OP	✓ Arrow-Bi	imonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE				
* Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	Combiga	n
PILOCARPINE HYDROCHLORIDE	4.00	45 100	41	
★ Eye drops 1%★ Eye drops 2%		15 ml OP 15 ml OP	✓ Isopto Call ✓ Isopto Call	
* Eye drops 2% ***********************************		15 ml OP	✓ Isopto Ca	•
Subsidised for oral use pursuant to the Standard Formu				
* Eye drops 2% single dose - Special Authority see SA0895				
below – Retail pharmacy	31.95	20 dose	Minims F	ilocarpine
⇒SA0895 Special Authority for Subsidy	160	P P	and an about the Harri	to a substant a
nitial application from any relevant practitioner. Approvals vali Either:	d for 2 years for	applications me	eting the follow	ing criteria:
Patient has to use an unpreserved solution due to an alle Patient wears soft contact lenses.	rgy to the preser	vative; or		
Note: Minims for a general practice are considered to be "tools of	of trade" and are	not approved a	is special autho	rity items
Renewal from any relevant practitioner. Approvals valid for 2 years penefiting from treatment.				
Mydriatics and Cycloplegics				
ATROPINE SULPHATE				
W Fue drame 10/	17.00	15 ml OD	./ Atront	

* Eye drops 1%......17.36

✓ Atropt

15 ml OP

	Subsidy		Fully Brand or
	(Manufacturer's Pr	rice) Subsi	idised Generic
	\$	Per	✓ Manufacturer
CYCLOPENTOLATE HYDROCHLORIDE			
	0.70	45 LOD	A Overland
* Eye drops 1%	8./6	15 ml OP	Cyclogyl
* Eye drops 1%, single dose (preservative free) - Only on a			
prescription	52.86	20 dose	✓ Minims
F F			Cyclopentolate
TD001044405			Cyclopolitolato
TROPICAMIDE			
* Eye drops 0.5%	7.15	15 ml OP	✓ Mydriacyl
* Eye drops 1%	8.66	15 ml OP	✓ Mydriacyl
, ,			
Preparations for Tear Deficiency			
r reparations for real benciency			
For contributation and draps refer Standard Formulae page 246	,		
For acetylcysteine eye drops refer Standard Formulae, page 246)		
HYPROMELLOSE			
* Eye drops 0.5%	2.00	15 ml OP	
	(3.92)		Methopt
	(0.32)		Methopt
HYPROMELLOSE WITH DEXTRAN			
* Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	✓ Poly-Tears
• •			•

Preservative Free Ocular Lubricants

⇒SA1388 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER - Special Authority see SA1388 above - Retail pharmac Ophthalmic gel 0.3%, 0.5 g		30	✓ Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL – Special Authority see Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml			harmacy Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] — Special Authority Eye drops 1 mg per ml Hylo-Fresh has a 6 month expiry after opening. The Pharmac month is not relevant and therefore only the prescribed dosage	22.00 1 by Procedures	I0 ml OP Manual restr	✓ Hylo-Fresh iction allowing one bottle per

Other Eye Preparations

15 ml OP	Naphcon Forte
5 ml OP	✓ Olopatadine Teva
3.5 g OP	✓ Poly-Visc
5 g OP	✓ VitA-POS
	5 ml OP 3.5 g OP



Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	✓	Manufacturer	

Various

PHARMACY SERVICES

May only be claimed once per patient.

The Pharmacode for BSF Atomoxetine Generic Partners is 2576996 - see also page 150 (BSF Atomoxetine Generic Partners Brand switch fee to be delisted 1 March 2021)

Agents Used in the Treatment of Poisonings

Antidotes

ACETYLCYSTEINE

✓ Martindale
Pharma \$29

NAI OXONE HYDROCHI ORIDE

a) Up to 5 inj available on a PSO

b) Only on a PSO

Removal and Elimination

CHARCOAL

a) Up to 250 ml available on a PSO

b) Only on a PSO

DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy

Wastage claimable

Tractage of	airidalo			
Tab 125 mg	dispersible	276.00	28	Exjade
Tab 250 mg	dispersible	552.00	28	✓ Exiade
	dispersible		28	✓ Exjade
	, ,	*		•

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria:



Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
---	-----	---------------------	-------------------------------------	--

continued...

Fither:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE - Special Authority see	SA1480 below – Retail pharmacy
-------------------------------------	--------------------------------

Tab 500 mg	533.17	100	Ferriprox
Oral lig 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox

■ SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE

* Inj 500 mg vial	84.53	10	✓ <u>DBL</u> <u>Desferrioxamine</u> <u>Mesylate for Inj</u> <u>BP</u>
SODIUM CALCIUM EDETATE			
* Inj 200 mg per ml, 5 ml	53.31	6	
	(156.71)		Calcium Disodium
			Versenate

Standard Formulae

Standard Formulae			
ACETYLCYSTEINE EYE DROPS		PHENOBARBITONE ORAL LIQUID	
Acetylcysteine inj 200 mg per ml, 10 ml	qs	Phenobarbitone Sodium	1 g
Suitable eye drop base	qs	Glycerol BP	70 ml
•		Water	to 100 ml
CODEINE LINCTUS (3 mg per 5 ml)			
Codeine phosphate	60 mg	PHENOBARBITONE SODIUM PAEDIATRIC ORAL	LIQUID (10
Glycerol	40 ml	mg per ml)	
Preservative	qs	Phenobarbitone Sodium	400 mg
Water	to 100 ml	Glycerol BP	4 ml
CODEINE LINCTUS (15 mg per 5 ml)		Water	to 40 ml
Codeine phosphate	300 mg	PILOCARPINE ORAL LIQUID	
Glycerol	40 ml	Pilocarpine 4% eye drops	qs
Preservative	qs	Preservative	qs
Water	to 100 ml	Water	to 500 ml
		(Preservative should be used if quantity supplied is f	
FOLINIC MOUTHWASH		than 5 days.)	
Calcium folinate 15 mg tab	1 tab	• •	
Preservative	qs	SALIVA SUBSTITUTE FORMULA	
Water	to 500 ml	Methylcellulose	5 g
(Preservative should be used if quantity supplied is f	or more	Preservative	qs
than 5 days. Maximum 500 ml per prescription.)		Water	to 500 ml
MAGNESIUM HYDROXIDE 8% MIXTURE		(Preservative should be used if quantity supplied is f	or more
Magnesium hydroxide paste 29%	275 g	than 5 days. Maximum 500 ml per prescription.)	
Methyl hydroxybenzoate	1.5 g	SODIUM CHLORIDE ORAL LIQUID	
Water	to 1,000 m		qs
	,	Water	qs
METHADONE MIXTURE		(Only funded if prescribed for treatment of hyponatra	iemia)
Methadone powder	qs	VANCOLIVOIN ORAL COLUTION (50	
Glycerol	qs	VANCOMYCIN ORAL SOLUTION (50 mg per ml)	40
Water	to 100 ml	Vancomycin 500 mg injection	10 vials
METHYL HYDROXYBENZOATE 10% SOLUTION		Glycerol BP	40 ml
Methyl hydroxybenzoate	10 g	Water (Only funded if prescribed for treatment of Clostridium	to 100 ml
Propylene glycol	to 100 ml	following metronidazole failure)	III UIIIICII C
(Use 1 ml of the 10% solution per 100 ml of oral liqui		following metroritidazole failure)	
	ia mixtaro)	VOSOL EAR DROPS	
OMEPRAZOLE SUSPENSION		WITH HYDROCORTISONE POWDER 1%	
Omeprazole capules or powder	qs	Hydrocortisone powder	1%
Sodium bicarbonate powder BP	8.4 g	Vosol Ear Drops	to 35 ml
Water	to 100 ml	·	

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

	(Manutacturer's Pi \$	rice) Subs Per	sidised Generic Manufacturer
Extemporaneously Compounded Preparations	and Galenica	ls	
CODEINE PHOSPHATE - Safety medicine; prescriber may det	ermine dispensing	g frequency	
Powder - Only in combination		25 g	Davidas
Only in extemporaneously compounded codeine linctus	(90.09)		Douglas
COLLODION FLEXIBLE			
Note: This product is no longer being manufactured by the determined.	supplier and will b	e delisted fror	n the Schedule at a date to be
Collodion flexible	19.30	100 ml	✓ PSM
COMPOUND HYDROXYBENZOATE – Only in combination Only in extemporaneously compounded oral mixtures.			
Soln		100 ml	✓ <u>Midwest</u>
GLYCERIN WITH SODIUM SACCHARIN — Only in combination Only in combination with Ora-Plus.			
Suspension	30.95	473 ml	✓ Ora-Sweet SF
GLYCERIN WITH SUCROSE – Only in combination			
Only in combination with Ora-Plus. Suspension	30.95	473 ml	✓ Ora-Sweet
GLYCEROL			
* Liquid – Only in combination		500 ml	✓ healthE Glycerol BP
Only in extemporaneously compounded oral liquid prepare	arations.		
METHADONE HYDROCHLORIDE			
a) Only on a controlled drug formb) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing fr	requency		
d) Extemporaneously compounded methadone will only be		rate of the ch	eapest form available
(methadone powder, not methadone tablets).			4
Powder	7.84	1 g	✓ AFT
METHYL HYDROXYBENZOATE Powder	0.00	0E a	✓ Midwoot
	0.90	25 g	✓ <u>Midwest</u>
METHYLCELLULOSE Powder	36.95	100 g	✓ MidWest
Suspension – Only in combination		473 ml	✓ Ora-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCH		ombination	
Suspension	•	473 ml	✓ Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - On	ly in combination		
Suspension	30.95	473 ml	✓ Ora-Blend
PHENOBARBITONE SODIUM			4
Powder - Only in combination	52.50 325.00	10 g 100 g	✓ MidWest ✓ MidWest
Only in children up to 12 years	323.00	100 g	• Iviiuvest
PROPYLENE GLYCOL			
Only in extemporaneously compounded methyl hydroxyben:		n.	
Liq	11.25	500 ml	✓ Midwest
SODIUM BICARBONATE	_		
Powder BP — Only in combination		500 g	✓ <u>Midwest</u>

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Only in extemporaneously compounded omeprazole and lansoprazole suspension.

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully osidised	Brand or Generic Manufacturer
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparation Liq		500 ml	✓ <u>M</u>	lidwest
WATER Tap - Only in combination	0.00	1 ml	✓ Ta	ap water

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

1 cancer in children: or

Both:

- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

continued...

✓ fully subsidised 249



Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	/	Manufacturer	

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

Subsidy		Fully	Brand or
(Manufacturer's	Price) Sub	sidised	Generic
<u> </u>	Per	1	Manufacturer

continued...

- 10 ascites: or
 - 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Authority see SA1523 on the previous page - Hospital pharmacy [HP3]

Emulsion (neutral)	12.30 200 1	ml OP	Calogen
	30.75 500 i	ml OP	Calogen
Emulsion (strawberry)	12.30 200 1	ml OP	Calogen
Oil	30.00 500 1	ml OP	MCT oil (Nutricia)
Oil, 250 ml	114.92 4	OP 🗸	′ Liquigen (

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT - Special Authority	y see SA1524 above – Hospital pha	rmacy [HP3]	
Powder	7.90	225 g OP	✓ Protifar
	8.95	227 g OP	✓ Resource
		-	Beneprotein

Subsidy Fully (Manufacturer's Price) Subsidised \$ Per ✓

Brand or Generic Manufacturer

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3] 1.000 ml OP ✓ Diason RTH ✓ Glucerna Select DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3] ✓ Diasip 200 ml OP 200 ml OP ✓ Diasip 1,88 250 ml OP ✓ Glucerna Select 237 ml OP 1.78 (2.10)Resource Diabetic (2.10)Sustagen Diabetic

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism: or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED − Special Authority see SA1525 above − Hospital pharmacy [HP3]
Powder60.48 400 g OP ✓ Monogen

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

continued...

 Subsidy Manufacturer's Price)	S	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1379 o Liquid		
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see SA1379 on Liquid	the previous page 500 ml OP	 Hospital pharmacy [HP3] Nutrini RTH Pediasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority spharmacy [HP3]	see SA1379 on the	e previous page – Hospital
Liquid	500 ml OP	✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML – Special Authority see SA1379 on the Liquid (strawberry)	e previous page – 200 ml OP 200 ml OP	Hospital pharmacy [HP3] ✓ Fortini ✓ Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA1379 on the Liquid (chocolate)	orevious page – H 200 ml OP 200 ml OP 200 ml OP 250 ml OP	lospital pharmacy [HP3] ✓ Pediasure ✓ Pediasure ✓ Pediasure ✓ Pediasure ✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority see Spharmacy [HP3]	SA1379 on the pre	evious page – Hospital
Liquid (unflavoured). 1.60 Liquid (chocolate) 1.60 Liquid (strawberry) 1.60 Liquid (vanilla) 1.60	200 ml OP 200 ml OP 200 ml OP 200 ml OP	 ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 on the previou Powder	s page – Hospital 400 g OP	pharmacy [HP3] Peptamen Junior

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML - Special Authority :	see SA1101 above –	Hospital pharma	cy [HP3]
Liquid	6.08	500 ml OP	✓ Nepro HP RTH

	Subsidy (Manufacturer's Pri		Fully dised	Brand or Generic Manufacturer
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see SA1 Liquid		is page – Hosp 220 ml OP	✓ N	narmacy [HP3] lepro HP (strawberry) lepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA110 Liquid		page – Hospit 237 ml OP	al pha	rmacy [HP3]
Liquid (apricot) 125 ml Liquid (caramel) 125 ml		4 OP 4 OP	✓ R	lovaSource Renal Renilon 7.5 Renilon 7.5

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LiquidLiquid	•		, , , ,
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see	e SA1377 above	- Hospital phar	macy [HP3]
Liquid (grapefruit), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
Liquid (summer fruits), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see	SA1377 above -	Hospital pharm	acy [HP3]
Powder (unflavoured)	4.50	80 g OP	✓ Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Aut	hority see SA137	7 above – Hosp	pital pharmacy [HP3]
Liquid	12.04	1,000 ml OP	✓ Peptisorb

Subsidy (Manufacturer's Price) Fully Subsidised Per Brand or Generic Manufacturer

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML –	 Special Authority 	see SA1196 ab	ove -	 Hospital pharmacy [HP3]
Liquid	4.00	500 ml OP	1	Nutrini Low Energy
				Multi Fibre

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

continued...

Subsidy (Manufacturer's Price)		Fully	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

continued...

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease: or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions: or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML − Special Authority see SA1859 on page 256 − Hospital pharmacy [HP3]

Liquid7.00 1,000 ml OP ✓ Nutrison Energy

	Subsidy		Fully Brand or
	(Manufacturer's I	Price) Subsi	•
	\$	Per	 Manufacturer
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on Liquid		spital pharmacy 250 ml OP 1,000 ml OP	[HP3] ✓ Isosource Standard ✓ Nutrison Standard RTH
			✓ Osmolite RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authorit Liquid	•	on page 256 – H 1,000 ml OP	ospital pharmacy [HP3] Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML – Special Authority se Liquid		page 256 – Hosp 1,000 ml OP	oital pharmacy [HP3] ✓ Jevity RTH ✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority s Liquid		page 256 – Hos 250 ml OP 1,000 ml OP	spital pharmacy [HP3] Ensure Plus HN Ensure Plus RTH Jevity HiCal RTH Nutrison Energy Multi Fibre
ORAL FEED (POWDER) – Special Authority see SA1859 on pag Note: Higher subsidy for Sustagen Hospital Formula will only number and an appropriately endorsed prescription. Powder (chocolate) – Higher subsidy of up to \$26.00 per 850	be reimbursed		
with Endorsement	0	850 g OP 840 g OP	✓ Ensure Sustagen Hospital
Additional subsidy by endorsement is available for patien prescription must be endorsed accordingly. Powder (vanilla) – Higher subsidy of up to \$26.00 per 850 g		, .	,
with Endorsement	26.00 9.54	857 g OP 850 g OP 840 g OP	✓ Fortisip ✓ Ensure
	(26.00)		Sustagen Hospital Formula Active

Additional subsidy by endorsement is available for patients with fat malabsorption, fat intolerance or chyle leak. The prescription must be endorsed accordingly. (Fortisip Powder (vanilla) to be delisted 1 August 2021)

Subsidy	1	Fully	Brand or	
(Manufacturer's Price)	Subsid	ised	Generic	
\$	Per	1	Manufacturer	

ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 256 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) — Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
	(1.26) (1.26)	200 0.	Ensure Plus Fortisip
Liquid (fruit of the forest) — Higher subsidy of \$1.26 per 200 ml with Endorsement		200 ml OP	
Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with	(1.26)		Ensure Plus
Endorsement	0.72 (1.26)	200 ml OP	Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.33 per 237 ml with Endorsement	0.85	237 ml OP	
	(1.33) 0.72	200 ml OP	Ensure Plus
	(1.26) (1.26)		Ensure Plus Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 256 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisin Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

continued...

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements: or
 - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/	ИL	– Spe	cial A	Autho	rity see SA11	95 on tl	ne previous p	age – Hospital p	harmacy [HP3]
Liquid							5.50	500 ml OP	✓ Nutrison
									Concentrated
							11.00	1,000 ml OP	Two Cal HN RTH
0041 5550 01/041 04	_								71 ID 41

ORAL FEED 2 KCAL/ML – Special Authority see SA1195 on the previous page – Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe

epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$1.90 per 200 ml with

90) Two Cal HN

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

SPECIAL FOODS

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully osidised	Brand or Generic Manufacturer
FOOD THICKENER – Special Authority see SA1106 on the pre-	6.53 30	pharmad 00 g OP 80 g OP	✓ N ✓ F	lutilis leed Thickener Karicare Aptamil

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by PHARMAC from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX – Special Authority see SA1729 Powder		,	
rowdei	(5.15)	1,000 g OP	Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA1729	above - Hospital pha	rmacy [HP3]	
Powder	3.93 1	1,000 g OP	
	(7.32)	-	NZB Low Gluten Bread Mix
	3.51		
	(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1729 above	<mark>re</mark> – Hospital pharmac	cy [HP3]	
Powder	5.62 2	2,000 g OP	
	(18.10)		Horleys Flour

	Subsidy		Fully	Brand or
	(Manufacturer's Pri \$	ce) Sub Per	sidised •	Generic Manufacturer
GLUTEN FREE PASTA - Special Authority see SA1729 on the	previous page – H	lospital pharr	nacy [HF	P3]
Buckwheat Spirals	2.00	250 g OP		
	(3.11)		0)rgran
Corn and Vegetable Shells	2.00	250 g OP		
	(2.92)		0)rgran
Corn and Vegetable Spirals	2.00	250 g OP		
	(2.92)		0)rgran
Rice and Corn Lasagne Sheets	1.60	200 g OP		
	(3.82)		0)rgran
Rice and Corn Macaroni	2.00	250 g OP		
	(2.92)	_	0)rgran
Rice and Corn Penne	2.00	250 g OP		
	(2.92)	_	0)rgran
Rice and Maize Pasta Spirals	2.00	250 g OP		•
	(2.92)	•	0)rgran
Rice and Millet Spirals	2.00	250 g OP		•
	(3.11)	•	0)rgran
Rice and corn spaghetti noodles	2.00	375 g OP		•
	(2.92)	•	0)rgran
Vegetable and Rice Spirals	2.00	250 g OP		•
-	(2.92)	,	0)rgran
Italian long style spaghetti	2.00	220 g OP		-
- · · ·	(3.11)	-	0)rgran

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 above - Hospital pharmacy [HP3]

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

Tabs	99.00	75 OP	✓ Phlexy 10
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior
			Chocolate
Powder (unflavoured) 27.8 g sachets	936.00	30	✓ PKU Lophlex
			Powder
Powder (unflavoured) 28 g sachets	936.00	30	✓ PKU Lophlex
			Powder
Powder (unflavoured) 36 g sachets		30	PKU Anamix Junior
Powder (vanilla) 36 g sachet	393.00	30	PKU Anamix Junior
			Vanilla
Infant formula	174.72	400 g OP	PKU Anamix Infant
Powder (orange)	320.00	500 g OP	XP Maxamum
Powder (unflavoured)	320.00	500 g OP	XP Maxamum
Liquid (berry)	13.10	125 ml OP	PKU Anamix Junior
			LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior
			LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior
,			LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	✓ Easiphen Liquid
Liquid (juicy tropical) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g		36 OP	✓ PKU Lophlex
	,,,		Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
		-1- 00041	*

(PKU Lophlex Powder Powder (unflavoured) 27.8 g sachets to be delisted 1 March 2021)

Foods

LOW PROTEIN BAKING MIX - Special Authority see SA1 Powder			oharmacy [HP3] ✓ Loprofin Mix
LOW PROTEIN PASTA - Special Authority see SA1108 of	n the previous page – I	Hospital pharm	acy [HP3]
Animal shapes	11.91	500 g OP	Loprofin
Lasagne	5.95	250 g OP	Loprofin
Low protein rice pasta	11.91	500 g OP	Loprofin
Macaroni	5.95	250 g OP	✓ Loprofin
Penne	11.91	500 g OP	✓ Loprofin
Spaghetti	11.91	500 g OP	✓ Loprofin
Spirals	11.91	500 g OP	✓ Loprofin

Subsidy (Manufacturer's Price) \$

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

Vanilla

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]
Powder44.40 400 g OP ✓ Locasol

Gastrointestinal and Other Malabsorptive Problems

AMINO ACID FORMULA - Special Authority see SA19	40 below – Hospital pharı	macy [HP3]	
Powder	43.60	400 g OP	Alfamino Junior
Powder (unflavoured)	53.00	400 g OP	✓ Elecare
,		Ü	✓ Elecare LCP
			✓ Neocate Gold
			✓ Neocate Junior
			Unflavoured
			✓ Neocate SYNEO
Powder (vanilla)	53.00	400 g OP	✓ Elecare
,		Ü	✓ Neocate Junior

⇒SA1940 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut: or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Fither:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — **(Children 12 months of age and over)** only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist.

continued...

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

Approvals valid for 6 months for applications meeting the following criteria:

1 Either:

- 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency: or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Fither:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 0.4 D-1

2 Both:

- 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
- 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Fither:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

continued...

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

ENTERAL LIQUID PEPTIDE FORMULA	 Special Authority see SA1953 below 	– Hospital phar	macy [HP3]
Liquid 1 kcal/ml	10.45	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml	15.68	500 ml OP	✓ Nutrini Peptisorb
			Energy

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea; or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure; or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and
 - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

continued...



Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
 \$	Per	•	Manufacturer

continued...

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see SA1557 below - Hospital pharmacy [HP3]

D 1	15.21		✓ Aptamil Gold+ Pepti Junior
	30.42	900 g OP	 Aptamil AllerPro SYNEO 1
			 Aptamil AllerPro SYNEO 2

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Subsidy Fully (Manufacturer's Price) Subsidised \$ Per

sed Generic

Manufacturer

Brand or

Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML - Special Authority see SA1698 below - Hospital pharmacy [HP3] Liquid.......2.35 125 ml OP ✓ Infatrini

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula: and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

 Powder (unflavoured)
 35.50
 300 g OP
 KetoCal 4:1

 V Ketocal 3:1
 KetoCal 4:1

 Powder (vanilla)
 35.50
 300 g OP
 KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Vaccinations** BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm] For infants at increased risk of tuberculosis. Increased risk is defined as: 1) living in a house or family with a person with current or past history of TB; or 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or egual to 40 per 100,000 for 6 months or longer; or 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php. Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin), Danish strain 1331, live attenuated, vial with diluent.................0.00 10 ✓ BCG Vaccine DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE - [Xpharm] Funded for any of the following criteria: 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or 3) A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or 5) A single dose for vaccination of patients aged from 65 years old; or 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or 7) For vaccination of previously unimmunised or partially immunised patients: or 8) For revaccination following immunosuppression; or 9) For boosting of patients with tetanus-prone wounds. Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis toxoid, 8 mcg pertussis filamentous 10 **Boostrix Boostrix** DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE - [Xpharm] Funded for any of the following: 1) A single dose for children up to the age of 7 who have completed primary immunisation; or 2) A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or 3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive 4) Five doses will be funded for children requiring solid organ transplantation. Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Ini 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units

Infanrix IPV

10

	NATIONAL I	IMMUNISAT	TION SCHEDULE
(Subsidy Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AN	D HAEMOPHILUS I	NFLUENZAE T	YPE B VACCINE -
[Xpharm] Funded for patients meeting any of the following criteria: 1) Up to four doses for children up to and under the age of 1 2) An additional four doses (as appropriate) are funded for (10 who are patients post haematopoietic stem cell transp post solid organ transplant, renal dialysis and other sevel 3) Up to five doses for children up to and under the age of 1 Note: A course of up-to four vaccines is funded for catch up pr to complete full primary immunisation. Please refer to the Imm programmes. Inj 30IU diphtheriatoxoid with 40IU tetanustoxoid, 25mcg pertussistoxoid, 25mcg pertussisfilamentoushaemagglutinin, 8 mcgpertactin, 80 D-AgUpoliovirus, 10mcghepatitisBsurfaceantigen in 0.5ml syringe	re-)immunisation for plantation, or chemot rely immunosuppres 0 receiving solid org rogrammes for childi nunisation Handbook	r children up to therapy; pre or sive regimens; gan transplanta ren (up to and u s for the approp	oost splenectomy; pre- or or ion. under the age of 10 years)
HAEMOPHILUS INFLUENZAE TYPE B VACCINE — [Xpharm] One dose for patients meeting any of the following: 1) For primary vaccination in children; or 2) An additional dose (as appropriate) is funded for (re-)imm transplantation, or chemotherapy; functional asplenic; pre or post cochlear implants, renal dialysis and other severe 3) For use in testing for primary immunodeficiency diseases paediatrician.	nunisation for patient e or post splenectom lly immunosuppressi	ts post haemate ny; pre- or post ive regimens; o	opoietic stem cell solid organ transplant, pre- r
Haemophilus Influenzae type B polysaccharide 10 mcg conjugated to tetanus toxoid as carrier protein 20-40 mcg; prefilled syringe plus vial 0.5 ml HEPATITIS A VACCINE — [Xpharm] Funded for patients meeting any of the following criteria: 1) Two vaccinations for use in transplant patients; or 2) Two vaccinations for use in children with chronic liver dis 3) One dose of vaccine for close contacts of known hepatitis	0.00 ease; or	1 🗸	Hiberix
Inj 1440 ELISA units in 1 ml syringe Inj 720 ELISA units in 0.5 ml syringe			Havrix Havrix Junior

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Inj 10 mo	RECOMBINANT VACCINE – [Xpharm] g per 0.5 ml prefilled syringeled for patients meeting any of the following criteria	0.00 a:	1	√ E	ingerix-B
2) 3) 4) 5) 6) 7) 8) 9)	for household or sexual contacts of known acute for children born to mothers who are hepatitis B s for children up to and under the age of 18 years in serology and require additional vaccination or req for HIV positive patients; or for hepatitis C positive patients; or for patients following non-consensual sexual interfor patients following immunosuppression; or for solid organ transplant patients; or for post-haematopoietic stem cell transplant (HSC)	urface antigen (HBsAg nclusive who are consic uire a primary course o course; or	, pos derec	itive; or I not to have	
Inj 20 mc Func 1) 2) 3) 4) 5) 6) 7) 8) 9) 10)	g per 1 ml prefilled syringe	a: nepatitis B patients or h urface antigen (HBsAg nclusive who are consid uire a primary course o course; or) pos derec	itis B carrier itive; or I not to have	
Any of th 1) Mai 2) Mai 1 2	ILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND e following: ximum of two doses for children aged 14 years and ximum of three doses for patients meeting any of the people aged 15 to 26 years inclusive; or Either: People aged 9 to 26 years inclusive 1) Confirmed HIV infection; or 2) Transplant (including stem cell) patients: oximum of four doses for people aged 9 to 26 years	d under; or ne following criteria: or			
Inj 270 m	cg in 0.5 ml syringe	0.00	10	√ <u>G</u>	Gardasil 9

	Subsidy (Manufacturer's Price) \$	F Subsidi Per	ully sed	Brand or Generic Manufacturer
INFLUENZA VACCINE				

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) ✓ Afluria Quad Junior (2020 Formulation)

A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by PHARMAC:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes: or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV. or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders. or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

Influvac Tetra	1	ccine)9.00	Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)
(2020 formulation)			
✓ Afluria Quad	10	90.00	
(2020 Formulation)			

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

- a) Only on a prescription
- b) No patient co-payment payable

С

A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by PHARMAC:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease: or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or
- c) children aged four years or less (but over three years) who have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	
--	--

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression: or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment from the Funder for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Fither:

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
 - 2) One dose for close contacts of meningococcal cases; or
 - 3) A maximum of two doses for bone marrow transplant patients: or
 - 4) A maximum of two doses for patients following immunosuppression*; or
- B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) Either:
 - i) One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
 - ii) One dose for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 December 2019 to 30 November 2021.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 4 mcg of each meningococcal polysaccharide conjugated to

a total of approximately 48 mcg of diphtheria toxoid carrier

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm] Both: 1) The child is under 9 months of age; and 2) Any of the following: 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or 2) Two doses for close contacts of meningococcal cases; or 3) A maximum of two doses for bone marrow transplant patients; or 4) A maximum of two doses for patients pre- and post-immunosuppression*. Note: children under nine months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for booster schedules with meningococcal ACWY vaccine. *Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days. 1 ✓ Neisvac-C

PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm]

1) A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B,

7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml

10

Synflorix

		- ::	-	
Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously
 received two doses of the primary course of PCV10: or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) with primary immune deficiencies: or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cyanosis or failure; or
 - I) with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note:	please refer to the Immunisation Handbook for the appropria	ate schedule for catch up programme	S
Inj 30.	8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,		

	Subsidy		Full	
	(Manufacturer's Price) \$	Per	Subsidise	d Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCIN Either:	IE - [Xpharm]			
Up to three doses (as appropriate) for patients wi	th HIV for nationts nost ha	omot	onoiotio et	om call transplant or
chemotherapy; pre- or post-splenectomy or with f				
complement deficiency (acquired or inherited), co	, , ,		•	
2) All of the following:	omean implante, or primary			, 0.
a) Patient is a child under 18 years for (re-)imr	nunisation; and			
b) Treatment is for a maximum of two doses; a	and			
c) Any of the following:				
 i) on immunosuppressive therapy or rad 	liation therapy, vaccinate wl	hen t	here is ex	pected to be a sufficient
immune response; or				
ii) with primary immune deficiencies; or				
iii) with HIV infection; oriv) with renal failure, or nephrotic syndror	me: or			
v) who are immune-suppressed following		ludin	n haemato	nnoietic stem cell transplant).
or	g organ transplantation (mo	i a a ii i	griadinak	polotio otom con transplanty,
vi) with cochlear implants or intracranial s	shunts; or			
vii) with cerebrospinal fluid leaks; or				
viii) receiving corticosteroid therapy for mo				
prednisone of 2 mg/kg per day or grea	ater, or children who weigh	more	than 10 k	g on a total daily dosage of
20 mg or greater; or	ding anthma tracted with his	ah da	oo oortio	otoroid thorony); or
ix) with chronic pulmonary disease (inclu x) pre term infants, born before 28 week		gri-uc	ose conicc	isteroid trierapy), or
xi) with cardiac disease, with cyanosis or	•			
xii) with diabetes; or				
xiii) with Down syndrome; or				
xiv) who are pre-or post-splenectomy, or v	vith functional asplenia.			
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each				
23 pneumococcal serotype)	0.00	1	•	Pneumovax 23
POLIOMYELITIS VACCINE - [Xpharm]				
Up to three doses for patients meeting either of the foll				
For partially vaccinated or previously unvaccinate	ed individuals; or			
2) For revaccination following immunosuppression.				
Note: Please refer to the Immunisation Handbook for a		tcn-u 1		imes. IPOL
Inj 80D antigen units in 0.5 ml syringe	0.00	ı	•	<u>IPOL</u>
ROTAVIRUS ORAL VACCINE – [Xpharm] Maximum of two decay for patients meeting the following the follo	na:			
Maximum of two doses for patients meeting the followi 1) first dose to be administered in infants aged under	•			
no vaccination being administered to children age				
Oral susp live attenuated human rotavirus				
1,000,000 CCID50 per dose, prefilled oral applicat	or0.00	10	•	Rotarix

	NATIONAL	IMMUNIS	SATIO	ON SCHEDULE
	Subsidy (Manufacturer's Price)	Subsid Per	Fully lised	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] – [Xpharm] Either:				
1) Maximum of one dose for primary vaccination for either	r:			
 a) Any infant born on or after 1 April 2016; or b) For previously unvaccinated children turning 11 y varicella infection (chickenpox), or 	ears old on or after 1	July 2017, v	who ha	ave not previously had a
2) Maximum of two doses for any of the following:				
 a) Any of the following for non-immune patients: 				
i) with chronic liver disease who may in future ii) with deteriorating renal function before tran iii) prior to called a reach transplant or iii)		nsplantation	ı; or	
iii) prior to solid organ transplant; oriv) prior to any elective immunosuppression*,	or			
v) for poet exposure prophylaxic who are imm		nto · or		

- v) for post exposure prophylaxis who are immune competent inpatients.; or
- b) For patients at least 2 years after bone marrow transplantation, on advice of their specialist, or
- c) For patients at least 6 months after completion of chemotherapy, on advice of their specialist, or
- d) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist, or
- For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella, or
- f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella, or
- g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.

* imm	nosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater tha
28 da	S

Inj 1350 PFU prefilled syringe	0.00 1	✓ Varivax
	10	✓ Varivax

VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED VACCINE [SHINGLES VACCINE] – [Xpharm] Funded for patients meeting either of the following criteria:

1) One dose for all people aged 65 years; or

2) One dose for all people aged between 66 and 80 years inclusive from 1 April 2018 and 31 December 2021.

Inj 19,400 PFU prefilled syringe plus vial	0.00 1	•	Zostavax
	10	•	7ostavax

Diagnostic Agents

TUBERCULIN PPD [MANTOUX] TEST - [Xpharm]			
Ini 5 TU per 0.1 ml. 1 ml vial	0.00	1	✓ Tuberso

- Symbols -		(2020 Formulation)	273	Amneal	16
UK Synacthen	81	Afluria Quad Junior		Amorolfine	
3TC		(2020 Formulation)	273	Amoxicillin	9
- A -		AFT Carbimazole		Amoxicillin with clavulanic acid	
A-Scabies	69	AFT-Pyrazinamide	101	Amphotericin B	
Abacavir sulphate	107	Agents Affecting the		Amsacrine	
Abacavir sulphate with		Renin-Angiotensin System	48	AmsaLyo	
lamivudine	107	Agents for Parkinsonism and Re		Amsidine	16
Abiraterone acetate		Disorders	120	Amzoate	3
Acarbose	11	Agents Used in the Treatment of		Anaesthetics	12
Accarb	11	Poisonings	244	Anafranil	12
Accuretic	49	Agrylin	162	Anagrelide hydrochloride	
Accuretic 10	49	Agrylin S29	162	Analgesics	12
Accuretic 20	49	Albendazole		Anastrozole	
Acetazolamide	242	Albey	231	Anatrole	17
Acetec	48	Albustix	78	Andriol Testocaps	8
Acetic acid with 1, 2- propanedi	ol	Aldurazyme	31	Androderm	8
diacetate and		Alecensa	169	ANI	
benzethonium	239	Alectinib	169	Anoro Ellipta	23
Acetic acid with hydroxyquinolir	ne and	Alendronate sodium	113	Antabuse	15
ricinoleic acid	<mark>76</mark>	Alendronate sodium with		Antacids and Antiflatulents	
Acetylcysteine		colecalciferol	113	Anthelmintics	
Aci-Jel		Alfacalcidol	35	Antiacne Preparations	
Aciclovir		Alfamino Junior	265	Antiallergy Preparations	
Infection	102	Alginic acid		Antianaemics	
Sensory	239	Alglucosidase alfa	29	Antiandrogen Oral	
Acidex	6	Alkeran		Contraceptives	7
Acipimox	56	Alkeran S29		Antiarrhythmics	
Acitretin		Allersoothe	232	Antibacterials	9
Aclasta	116	Allmercap		Antibacterials Topical	6
Aclin	112	Allopurinol	118	Anticholinergic Agents	
Actemra	221	Alpha-Adrenoceptor Blockers	48	Anticholinesterases	
Actinomycin D	163	Alpha-Keri Lotion	67	Antidepressants	
Actrapid	10	Alphamox	93	Antidiarrhoeals	
Actrapid Penfill	10	Alphamox 125	93	Antiepilepsy Drugs	12
Acular	241	Alphamox 250	93	Antifibrinolytics, Haemostatics and	
Acupan	122	Alprolix	39	Local Sclerosants	3
Adalat 10	53	Alu-Tab	6	Antifibrotics	
Adalat Oros	53	Aluminium hydroxide	6	Antifungals	9
Adalimumab	187	Alvogen	52	Antifungals Topical	
Adapalene	62	Amantadine hydrochloride	120	Antihistamines	
Adcortyl	81	Ambrisentan	58	Antihypotensives	
Adefin	53	Ambrisentan Mylan	58	Antimalarials	9
Adefin XL	53	Amiloride hydrochloride	54	Antimigraine Preparations	
Adefovir dipivoxil	101	Amiloride hydrochloride with		Antinausea and Vertigo Agents	
Adenuric		furosemide	55	Antiparasitics	
ADR Cartridge 1.8		Amiloride hydrochloride with		Antipruritic Preparations	
Adrenaline		hydrochlorothiazide		Antipsychotics	
Advantan		Aminophylline		Antiretrovirals	
Advate		Amiodarone hydrochloride		Antirheumatoid Agents	11
Adynovate		Amisulpride		Antispasmodics and Other Agents	
Afinitor		Amisulpride Mylan		Altering Gut Motility	
Aflibercept	197	Amitriptyline		Antithrombotic Agents	4
Afluria Quad		Amlodipine	53	Antithymocyte globulin	

(equine)	187	Aripiprazole Sandoz	133	Azathioprine	
Antitrichomonal Agents	100	Aristocort	65	Azithromycin	9
Antituberculotics and		Arrow-Amitriptyline	126	Azol	89
Antileprotics	100	Arrow-Bendrofluazide	55	Azopt	242
Antiulcerants		Arrow-Brimonidine	242	AZT	107
Antivirals	101	Arrow-Calcium	36	- B -	
Anxiolytics	136	Arrow-Diazepam	136	B-D Micro-Fine	
Anzatax	166	Arrow-Doxorubicin	163	B-D Ultra Fine	
Apidra		Arrow-Fluoxetine	127	B-D Ultra Fine II	14
Apidra SoloStar	11	Arrow-Losartan &		Bacillus Calmette-Guerin (BCG)	
Apo-Amlodipine		Hydrochlorothiazide	49	vaccine	187
Apo-Azithromycin		Arrow-Morphine LA		Bacillus Calmette-Guerin	
Apo-Bromocriptine		Arrow-Norfloxacin	111	vaccine	270
Apo-Ciclopirox	63	Arrow-Ornidazole		Baclofen	119
Apo-Cilazapril/		Arrow-Quinapril 10		Bactroban	
Hydrochlorothiazide	49	Arrow-Quinapril 20		Barrier Creams and Emollients	66
Apo-Clarithromycin		Arrow-Quinapril 5		BCG Vaccine	
Alimentary	9	Arrow-Roxithromycin	92	Beclazone 100	
Infection	91	Arrow-Sertraline	127	Beclazone 250	
Apo-Clomipramine	126	Arrow-Timolol		Beclazone 50	232
Apo-Diclo SR		Arrow-Topiramate		Beclomethasone dipropionate	232
Apo-Diltiazem CD	53	Arrow-Tramadol	126	Bee venom allergy treatment	23
Apo-Doxazosin	48	Arsenic trioxide	162	Bendamustine hydrochloride	157
Apo-Folic Acid	39	Asacol	7	Bendrofluazide	5
Apo-Furosemide		Asamax	7	Bendroflumethiazide	
Apo-Gabapentin	128	Ascorbic acid	35	[Bendrofluazide]	5
Apo-Megestrol	177	Aspen Adrenaline	58	Benzathine benzylpenicillin	93
Apo-Metoprolol	52	Aspirin		Benzatropine mesylate	120
Apo-Mirtazapine	128	Blood	42	Benzbromaron AL 100	118
Apo-Nadolol	52	Nervous	122	Benzbromarone	118
Apo-Nicotinic Acid	56	Asthalin	<mark>234</mark>	Benztrop	120
Apo-Oxybutynin	77	Atazanavir sulphate	107	Benzydamine hydrochloride	3
Apo-Perindopril	48	Atenolol	51	Benzylpenicillin sodium [Penicillin	
Apo-Pindolol	52	Atenolol AFT	51	G]	93
Apo-Pravastatin	56	Atenolol AFT S29	51	Beta Cream	
Apo-Prazosin	48	ATGAM	187	Beta Ointment	64
Apo-Prednisone	81	Ativan	136	Beta Scalp	
Apo-Primidone	130	Atomoxetine	150	Beta-Adrenoceptor Agonists	234
Apo-Propranolol	52	Atorvastatin	<mark>56</mark>	Beta-Adrenoceptor Blockers	
Apo-Pyridoxine		Atropine sulphate		Betadine	6
Apo-Selegiline S29		Cardiovascular	50	Betadine Skin Prep	6
Apo-Sumatriptan		Sensory	242	Betaferon	147
Apo-Temozolomide		Atropt	242	Betahistine dihydrochloride	132
Apo-Terazosin	48	Atrovent		Betaine	30
Apo-Timol		AU Synacthen	81	Betaloc CR	52
Apomorphine hydrochloride		Aubagio	142	Betamethasone dipropionate	64
Aprepitant		Augmentin	93	Betamethasone dipropionate with	
Apresoline	58	Aurorix	127	calcipotriol	
Aptamil AllerPro SYNEO 1		AutoSoft 30		Betamethasone sodium phosphat	
Aptamil AllerPro SYNEO 2		AutoSoft 90	24–25	with betamethasone acetate	
Aptamil Gold+ Pepti Junior		Avelox		Betamethasone valerate	
Aqueous cream		Avonex		Betamethasone valerate with	•
Aratac		Avonex Pen		clioquinol	65
Arava		Azacitidine		Betamethasone valerate with sodi	
Aripiprazole		Azacitidine Dr Reddy's		fusidate [fusidic acid]	
Aripiprazole 1A Pharma		Azamun		Betaxolol	

INDEX: Generic Chemicals and Brands

Betnovate	64	Buccastem133	CareSens N1
Betnovate-C	65	Budesonide	CareSens N POP1
Betoptic	241	Alimentary6	CareSens N Premier1
Betoptic S		Respiratory232, 238	CareSens PRO1
Bezafibrate		Budesonide with eformoterol233	Carmellose sodium with gelatin and
Bezalip	55	Bumetanide54	pectin3
Bezalip Retard		Buprenorphine Naloxone BNM 154	Carmustine15
Bicalutamide		Buprenorphine with naloxone154	Carvedilol
Bicillin LA		Bupropion hydrochloride154	Carvedilol Sandoz
BiCNU		Burinex54	Catapres
Bicnu Heritage		Burinex S2954	CeeNU15
Bile and Liver Therapy		Buscopan8	Cefaclor monohydrate
Biltricide		Buspirone hydrochloride136	Cefalexin
Bimatoprost		Busulfan158	Cefalexin Sandoz
Bimatoprost Multichem		- C -	Cefazolin
Binarex		Cabergoline88	Ceftriaxone
Binocrit		Cacit	Ceftriaxone-AFT
Biodone		Caffeine citrate238	Cefuroxime axetil
Biodone Extra Forte		Calamine	Celebrex11
Biodone Forte		Calci-Tab 500	Celecoxib11
Bisacodyl		Calcipotriol69	Celecoxib Pfizer11
		Calcitonin	Celestone Chronodose
Bisoprolol fumarate		Calcitriol	
Bisoprolol Mylan		Calcitriol-AFT35	Celliprolol
BK Lotion			Cellcept18
Bleomycin sulphate	103	Calcium carbonate	Celol
Blood Colony-stimulating	45	Calcium Channel Blockers	Centrally-Acting Agents
Factors	45	Calcium Disodium Versenate245	Cephalexin ABM
Blood glucose diagnostic test	40	Calcium folinate	Cetirizine hydrochloride23
meter	13	Calcium Folinate Ebewe	•
Blood glucose diagnostic test	40	Calcium Folinate Sandoz160	Cetomacrogol with glycerol
strip		Calcium gluconate	Cetuximab19
Blood glucose test strips (visually		Calcium Homeostasis79	Charcoal24
impaired)	13	Calcium polystyrene sulphonate47	Chemotherapeutic Agents15
Blood Ketone Diagnostic Test		Calcium Resonium47	Chickenpox vaccine27
Strip		Calcium Sandoz36	Chlorafast23
Bonjela		Calogen251	Chlorambucil15
Boostrix		Candesartan cilexetil49	Chloramphenicol23
Bortezomib		Candestar49	Chlorothiazide5
Bortezomib Dr-Reddy's		Canesten63	•
Bosentan		Capecitabine160	Chlorpromazine hydrochloride13
Bosentan Dr Reddy's		Capercit160	Chlortalidone [Chlorthalidone]5
Bosvate		Capoten48	Chlorthalidone
Bplex	34	Capsaicin	Chlorvescent4
Breo Ellipta		Musculoskeletal113	
Brevinor 1/28		Nervous122	Choice TT380 Short7
Bricanyl Turbuhaler	234	Captopril48	Choice TT380 Standard
Brilinta		Carafate10	Choline salicylate with cetalkonium
Brimonidine tartrate	242	Carbaccord	chloride3
Brimonidine tartrate with timolol		Carbamazepine128	Ciclopirox olamine6
maleate	242	Carbimazole84	
Brinzolamide	242	Carbomer243	Cilazapril4
Bristol	53	Carboplatin158	Cilazapril with
Brolene	239	Carboplatin Ebewe158	
Bromocriptine mesylate	120	Carbosorb-X244	
BSF Atomoxetine Generic		Cardinol LA52	
Partners	244	CareSens Dual12	

Cipflox	94	Colifoam	<mark>7</mark>	Dantrium	119
Ciprofloxacin		Colistin sulphomethate		Dantrium S29	119
Infection	94	Colistin-Link	95	Dantrolene	119
Sensory	239	Collodion flexible	247	Daonil	1 1
Ciprofloxacin Teva		Colloidal bismuth subcitrate	10	Dapa-Tabs	55
Circadin	149	Colofac	8	Dapsone	100
Cisplatin		Coloxyl	28	Daraprim	96
Cisplatin Ebewe	158	Combigan	242	Darunavir	
Citalopram hydrobromide		Compound electrolytes		Darunavir Mylan	107
Cladribine		Compound electrolytes with gluc	cose	Dasatinib	170
Clarithromycin		[Dextrose]		Daunorubicin	
Alimentary	9	Compound hydroxybenzoate		David One Step Cassette Pregna	ncy
Infection		Concerta		Test	
Clexane	44	Condoms	73	DBL Acetylcysteine	
Clexane Forte		Condyline	71	DBL Adrenaline	
Climara	82	Contraceptives - Hormonal		DBL Aminophylline	
Climara		Contraceptives - Non-hormonal.		DBL Bleomycin Sulfate	
Clindamycin		Copaxone		DBL Carboplatin	
Clinicians Renal Vit		Corticosteroids and Related Age		DBL Cisplatin	
Clobazam		for Systemic Use		DBL Dacarbazine	
Clobetasol propionate		Corticosteroids Topical		DBL Desferrioxamine Mesylate fo	
Clobetasone butyrate		Cosentyx		BP	
Clofazimine		Cosmegen		DBL Docetaxel	
Clomazol	100	Coumadin		DBL Ergometrine	
Dermatological	63	Creon 10000		DBL Gemcitabine	
Genito-Urinary		Creon 25000		DBL Gentamicin	
Clomifene citrate		Creon Micro		DBL Heparin Sodium	
Clomipramine hydrochloride		Crotamiton		DBL Leucovorin Calcium	
Clonazepam		Crystaderm		DBL Methotrexate Onco-Vial	
Clonidine		Curam		DBL Morphine Sulphate	
Clonidine BNM		Cvite		DBL Naloxone Hydrochloride	24/
				DBL Natioxoffe Trydrochionae DBL Octreotide	
Clonidarel		Cyclizine hydrochloride			
Clopidogrel		Cyclizine lactate		DBL Pethidine Hydrochloride	
Clopidogrel Multichem		Cyclogyl		DBL Vinblastine	
Clopine		Cyclopentolate hydrochloride		DBL Vincristine Sulfate	
Clopixol	134, 136	Cyclophosphamide		De-Worm	
Clotrimazole	00	Cyclorin		Decozol	
Dermatological		Cycloserine		Deferasirox	
Genito-Urinary		Cyproterone acetate	81	Deferiprone	
Clozapine		Cyproterone acetate with		Denosumab	
Clozaril		ethinyloestradiol		Deolate	
Co-trimoxazole		Cystadane		Deoxycoformycin	
Coal tar		Cytarabine		Depo-Medrol	
Coal tar with allantoin, mentho		Cytotec	8	Depo-Provera	
phenol and sulphur		Cytoxan	158	Depo-Testosterone	
Coal tar with salicylic acid and		- D -		Deprim	97
sulphur		D-Penamine	113	Dermol	.65, 70
Coco-Scalp	70	Dabigatran	45	Desferrioxamine mesilate	245
Codeine phosphate		Dacarbazine		Desmopressin	8
Extemporaneous	247	Dacarbazine APP	163	Desmopressin acetate	
Nervous	123	Dactinomycin [Actinomycin D]	163	Desmopressin-PH&T	
Colchicine	118	Daivobet		Desuric	
Colecalciferol		Daivonex		Detection of Substances in	
Colestid	56	Daktarin	64	Urine	78
Colestipol hydrochloride		Dalacin C		Dexamethasone	
Colgout		Danazol		Hormone	80

INDEX: Generic Chemicals and Brands

Sensory240	Diuretics54	Egopsoryl TA	
Dexamethasone phosphate80	Docetaxel163	Elaprase	3
Dexamethasone Phosphate	Docetaxel Accord163	Elecare	.26
Panpharma80	Docetaxel Sandoz163	Elecare LCP	.26
Dexamethasone with framycetin and	Docusate sodium28	Electral	
gramicidin239	Docusate sodium with	Elelyso	3
Dexamethasone with neomycin	sennosides28	Elemental 028 Extra	.25
sulphate and polymyxin B	Dolutegravir108	Elidel	
sulphate240	Domperidone 132	Elocon	6
Dexamfetamine sulfate150	Donepezil hydrochloride153	Elocon Alcohol Free	
Dexmethsone80	Donepezil-Rex153	Eltrombopag	3
Dextrochlorpheniramine	Dornase alfa237	Eltroxin	8
maleate232	Dortimopt242	EMB Fatol	
Dextrose46–47	Dorzolamide hydrochloride242	Emend Tri-Pack	
DHC Continus123	Dorzolamide with timolol242	Emicizumab	4
Diabetes10	Dostinex88	EMLA	
Diabetes Management12	Dosulepin [Dothiepin]	Emtricitabine	. 10
Diacomit130	hydrochloride126	Emtricitabine with tenofovir	
Diagnostic Agents279	Dosulepin Mylan126	disoproxil	
Diamide Relief6	Dothiepin	Emtriva	. 10
Diamox242	Doxazosin48	Emulsifying ointment	
Diasip252	Doxine94	Emulsifying Ointment ADE	6
Diason RTH252	Doxorubicin Ebewe163	Enalapril maleate	4
Diazepam128, 136	Doxorubicin hydrochloride163	Enbrel	
Diazoxide10	Doxycycline94	Endocrine Therapy	. 17
Dibenzyline48	DP Lotion67	Endoxan	
Diclofenac Sandoz112	DP Lotn HC65	Engerix-B	.27
Diclofenac sodium	DP-Allopurinol118	Enlafax XR	.12
Musculoskeletal112	Dr Reddy's Omeprazole9	Enoxaparin sodium	4
Sensory240	Drugs Affecting Bone	Enstilar	6
Differin62	Metabolism 113	Ensure	
Difflam33	Dual blood glucose and blood ketone	Ensure Plus	.26
Diflucan97	diagnostic test meter 12	Ensure Plus HN	.25
Diflucortolone valerate65	Duocal Super Soluble Powder250	Ensure Plus RTH	.25
Digestives Including Enzymes27	Duolin234	Entacapone	.12
Digoxin50	Duolin HFA234	Entapone	
Dihydrocodeine tartrate123	DuoResp Spiromax233	Entecavir	. 10
Dilantin129	Duride57	Entecavir Sandoz	
Dilantin Infatab129	-E-	Entocort CIR	
Diltiazem hydrochloride53	e-chamber La Grande238	Entresto 24/26	
Dilzem53	e-chamber Mask238	Entresto 49/51	4
Dimethicone66–67	e-chamber Turbo238	Entresto 97/103	
Dimethyl fumarate136	E-Mycin92	Epilim	.13
Dipentum7	Ear Preparations239	Epilim Crushable	.13
Diphtheria, tetanus and pertussis	Ear/Eye Preparations239	Epilim IV	.13
vaccine270	Easiphen Liquid264	Epilim S/F Liquid	
Diphtheria, tetanus, pertussis and	Econazole nitrate63	Epilim Syrup	.13
polio vaccine270	Efavirenz106	Epirubicin Ebewe	.16
Diphtheria, tetanus, pertussis, polio,	Efavirenz with emtricitabine and	Epirubicin hydrochloride	
hepatitis B and haemophilus	tenofovir disoproxil 107	Eplerenone	5
influenzae type B vaccine 271	Effient43	Epoetin alfa	
Diprosone64	Eformoterol fumarate233	Epoprostenol	
Diprosone OV64	Eformoterol fumarate dihydrate233	Eptacog alfa [Recombinant factor	
Dipyridamole42	Eftrenonacog alfa [Recombinant	VIIa]	4
Disopyramide phosphate50	factor IX]39	ERA	
Disulfiram155	Efudix71	Erbitux	.19

Ergometrine maleate	76	FEIBA NF	41	Fluox	127
Erlotinib	170	Felo 10 ER	53	Fluoxetine hydrochloride	127
Erythrocin IV	92	Felo 5 ER	53	Flupenthixol decanoate	134
Erythromycin (as lactobionate)		Felodipine	53	Flutamide	
Erythromycin ethyl succinate	92	Femme-Tab ED		Flutamin	177
Erythromycin stearate		Fentanyl	124	Fluticasone	233
Esbriet	236	Fentanyl Sandoz		Fluticasone furoate with	
Escitalopram		Ferinject	36	vilanterol	233
Escitalopram-Apotex		Ferodan		Fluticasone propionate	238
Eskazole		Ferric carboxymaltose	36	Fluticasone with salmeterol	
Essential Prednisolone	8	Ferriprox		FML	241
Estradiol TDP Mylan	82	Ferro-F-Tabs		Foban	
Estradot		Ferro-tab	37	Folic acid	39
Estradot 50 mcg		Ferrograd	37	Food Thickeners	261
Estrofem		Ferrosig		Foods And Supplements For In	ıborn
Etanercept		Ferrous fumarate		Errors Of Metabolism	
Ethambutol hydrochloride		Ferrous fumarate with folic acid.		Foradil	
Ethics Aspirin		Ferrous sulfate		Forteo	
Ethics Aspirin EC		Ferrous sulphate		Fortini	
Ethics Lisinopril		Fexofenadine hydrochloride		Fortini Multi Fibre	254
Ethics Paracetamol Classic		Fibro-vein		Fortisip	
Ethinyloestradiol		Filgrastim		Fortisip Multi Fibre	260
Ethinyloestradiol with		Finasteride		Fosamax	
desogestrel	74	Fingolimod		Fosamax Plus	
Ethinyloestradiol with		Firazyr		Framycetin sulphate	
levonorgestrel	75	Flagyl		Frisium	
Ethinyloestradiol with		Flagyl-S		Frumil	
norethisterone	75	Flamazine		Frusemide	
Ethosuximide		Flecainide acetate		Frusemide-Claris	
Etopophos		Flecainide BNM		Fucicort	
Etoposide		Flecainide Controlled Release		Fucidin	
Etoposide phosphate		Teva	50	Fucithalmic	
Etravirine		Fleet Phosphate Enema		Fulvestrant	
Eumovate		Flixonase Hayfever & Allergy		Fungilin	
Everet		Flixotide		Furosemide [Frusemide]	
Everolimus		Flixotide Accuhaler		Furosemide-Baxter	
Evista		Florinef		fusidic acid	
Exemestane		Fluanxol		Dermatological	63 69
Exjade		Flucil		Infection	
Extemporaneously Compounded		Flucloxacillin		Sensory	
Preparations and		Flucloxin		- G -	200
Galenicals	247	Flucon		Gabapentin	129
Eye Preparations		Fluconazole		Gacet	
Eylea		Fludara Oral		Galsulfase	
Ezetimibe		Fludarabine Ebewe		Galvumet	
Ezetimibe Sandoz		Fludarabine phosphate		Galvus	
Ezetimibe with simvastatin		Fludrocortisone acetate		Gardasil 9	
- E -	37	Fluids and Electrolytes		Gastrodenol	
Factor eight inhibitor hypocoing				Gaviscon Double Strength	
Factor eight inhibitor bypassing fraction	41	Fluncartologo capraeto with	208		
Famotidine		Fluocortolone caproate with		Gaviscon Infant	
Famotidine Hovid	•	fluocortolone pivalate and	0	GazyvaGefitinib	
		cinchocaine			
Faslodex		Fluorometholone		Gemcitabine Ebewe	
Febuxostat Feed Thickener Karicare	110	FluorouracilFluorouracil Ebewe		Gemcitabine hydrochloride	
	060	Fluorouracii Edewe		Genoptic	239
Aptamil	262	FIUOTOUTACII SODIUM	/ 1		

INDEX: Generic Chemicals and Brands

Gentamicin sulphate	Hemastix78	Hydroxyurea	
Infection95	Hemlibra40	[hydroxycarbamide]	16
Sensory239	Heparin sodium45	Hygroton	5
Gilenya138	Heparinised saline45	Hylo-Fresh	
Ginet76	Heparon Junior253	Hymenoptera	
Glatiramer acetate	Hepatitis A vaccine271	Hyoscine butylbromide	
Glecaprevir with pibrentasvir103	Hepatitis B recombinant	Hyoscine hydrobromide	
Glibenclamide11	vaccine272	Hypam	
Gliclazide12	Hepsera101	Hyperuricaemia and Antigout	
Glipizide12	Herceptin224	Hypromellose	
Glivec171	Hiberix271	Hypromellose with dextran	
Glizide12	Hiprex110	-1-	
Glucagen Hypokit10	Histaclear232	Ibiamox	9
Glucagon hydrochloride10	Histafen232	Ibilex	9
Glucerna Select252	Holoxan158	Ibrance	17
Glucerna Select RTH252	Horleys Bread Mix262	Ibuprofen	11
Glucobay11	Horleys Flour262	Ibuprofen SR BNM	11
Glucose [Dextrose]46	Hormone Replacement Therapy -	lcatibant	
Gluten Free Foods262	Systemic 81	Idarubicin hydrochloride	16
Glycerin with sodium saccharin247	HPV272	Idursulfase	
Glycerin with sucrose247	Humalog11	Ifosfamide	15
Glycerol	Humalog Mix 2511	Igroton	5
Alimentary29	Humalog Mix 5011	Ikorel	
Extemporaneous247	Human papillomavirus (6, 11, 16, 18,	llevro	
Glyceryl trinitrate	31, 33, 45, 52 and 58) vaccine	lloprost	6
Alimentary8	[HPV]272	Imatinib mesilate	
Cardiovascular57	Humatin96	Imatinib-AFT	17
Glycopyrronium235	Humira187	Imatinib-Rex	17
Glycopyrronium bromide8	HumiraPen187	Imigran	13
Glycopyrronium with	Humulin 30/7011	Imipramine hydrochloride	
indacaterol235	Humulin NPH11	Imiquimod	
Gold Knight73	Humulin R10	Immune Modulators	
Gold Knight XL73–74	Hyaluronic acid243	Immunosuppressants	
Goserelin	Hydralazine58	Imuran	
Gutron51	Hydralazine hydrochloride58	Incruse Ellipta	23
Gynaecological Anti-infectives76	Hydrea164	Indacaterol	
´ -H-	Hydrocortisone	Indapamide	
Habitrol	Dermatological65	Infanrix IPV	
Haemophilus influenzae type B	Hormone80	Infanrix-hexa	27
vaccine271	Hydrocortisone (PSM)65	Infant Formulae	26
Haldol135	Hydrocortisone acetate7	Infatrini	26
Haldol Concentrate135	Hydrocortisone acetate with	Infliximab	19
Haldol Decanoas135	pramoxine hydrochloride 7	Influenza vaccine	27
Haloperidol 133	Hydrocortisone and paraffin liquid	Influvac Tetra	
Haloperidol decanoate135	and lanolin65	(2020 formulation)	27
Harvoni104	Hydrocortisone butyrate65, 70	Inhaled Corticosteroids	23
Havrix271	Hydrocortisone with cinchocaine8	Inhaled Long-acting	
Havrix Junior271	Hydrocortisone with miconazole66	Beta-adrenoceptor Agonists	23
healthE Calamine Aqueous Cream	Hydrocortisone with natamycin and	Inspra	
BP64	neomycin 66	Instillagel Lido	
healthE Dimethicone 10%66	Hydrogen peroxide62	Insulin aspart	
healthE Dimethicone 4% Lotion 67	Hydroxocobalamin34	Insulin aspart with insulin aspart	
healthE Dimethicone 5%	Hydroxocobalamin Mercury	protamine	1
healthE Glycerol BP247	Pharma34	Insulin glargine	
healthE Urea Cream66	hydroxycarbamide164	Insulin glulisine	
Healtheries Simple Baking Mix262	Hydroxychloroquine113	Insulin isophane	

Insulin isophane with insulin	Isotretinoin	62	Lansoprazole	9
neutral			Lantus	
Insulin lispro			Lantus SoloStar	
Insulin lispro with insulin lispro	Itch-Soothe		Lanvis	162
protamine	11 Itraconazole	97	Lanzol Relief	
Insulin neutral			Lapatinib ditosylate	
Insulin pen needles			Largactil	133
Insulin pump			Laronidase	
Insulin pump cartridge		75	Lasix	
Insulin pump infusion set (steel	Jakavi		Latanoprost	
cannula)			Lax-Suppositories	29
Insulin pump infusion set (steel	Jevity HiCal RTH		Lax-Tab	
cannula, straight insertion)			Laxatives	
Insulin pump infusion set (teflon	Juno Pemetrexed		Laxsol	28
cannula)			Ledipasvir with sofosbuvir	
Insulin pump infusion set (teflon	Kadcyla	226	Leflunomide	
cannula, angle insertion with	Kaletra		Lenalidomide	
insertion device)			Letrole	
Insulin pump infusion set (teflon	Kenacomb		Letrozole	
cannula, angle insertion)			Leukeran FC	
Insulin pump infusion set (teflon	Kenacort-A 40		Leukotriene Receptor	
cannula, straight insertion with	Kenalog		Antagonists	237
insertion device)	•		Leuprorelin	
Insulin pump infusion set (teflon	Kenkay Sorbolene		Leustatin	
cannula, straight insertion)	•		Levetiracetam	
Insulin pump reservoir			Levetiracetam-AFT	
Insulin syringes, disposable with	Ketoconazole		Levlen ED	
attached needle		70	Levocabastine	
Intal Forte CFC Free2			Levodopa with benserazide	
Intelence1			Levodopa with carbidopa	
Interferon beta-1-alpha1			Levomepromazine	
Interferon beta-1-beta1			Levomepromazine	
Intra-uterine device			hydrochloride	134
Invega Sustenna1			Levonorgestrel	
IPOL2			Genito-Urinary	75-76
Ipratropium bromide234, 2			Hormone	
Iressa1			Levothyroxine	
Irinotecan Accord1			Lidocaine [Lignocaine]	
Irinotecan Actavis 1001	•		Lidocaine [Lignocaine]	
Irinotecan hydrochloride			hydrochloride	122
Irinotecan-Rex1			Lidocaine [Lignocaine] with	
Iron polymaltose			chlorhexidine	122
Isentress1			Lidocaine [Lignocaine] with	
Isentress HD1			prilocaine	122
Ismo 20		52	Lidocaine-Claris	122
Ismo 40 Retard			Lignocaine	
Isoniazid1			Lioresal Intrathecal	
Isoniazid with rifampicin1			Lipid-Modifying Agents	
Isoprenaline [Isoproterenol]			Liquigen	251
Isoproterenol			Lisinopril	23
Isoptin		m 102, 107	Lithium carbonate	
Isoptin Retard			Livostin	
Isoptin SR			LMX4	
Isopto Carpine2			Locacorten-Viaform ED's	
Isosorbide mononitrate	57 Lanoxin PG		Local preparations for Anal and	
Isosource Standard2	259 Lanoxin S29		Rectal Disorders	,
13030u100 3ta1lualu	Landani Jej		1 1561a1 DISUIUEIS	

Locasol	265	Maxitrol240	succinate)	8
Locoid	65, 70	MCT oil (Nutricia)251	,	
Locoid Crelo	65	Measles, mumps and rubella	Methylprednisolone acetate	8
Locoid Lipocream	65	vaccine275	Methylxanthines	23
Locorten-Vioform		Mebendazole90	•	
Lodoxamide	241	Mebeverine hydrochloride		
Logem	129	Medco123	Metolazone	5
Lomide		Medrol80		
Lomustine		Medroxyprogesterone acetate	Metoprolol succinate	5
Loniten	58	Genito-Urinary76		
Loperamide hydrochloride	6	Hormone82–83		10
Lopinavir with ritonavir	107	Mefenamic acid112		
Loprofin		Megestrol acetate177		
Loprofin Mix		Melatonin149		
Lorafix	232	Melphalan 158	Mexiletine hydrochloride	5
Loratadine	232	Menactra275	Mexiletine Hydrochloride USP	5
Lorazepam	136	Meningococcal (groups A, C, Y and	Miacalcic	
Lorfast	232	W-135) conjugate vaccine 275	Micolette	2
Lorstat	<u>56</u>	Meningococcal C conjugate	Miconazole	3
Losartan Actavis	49	vaccine276	Miconazole nitrate	
Losartan potassium	49	Menthol64	Dermatological	6
Losartan potassium with		Mepolizumab206	Genito-Urinary	7
hydrochlorothiazide	49	Mercaptopurine161		
Lovir		Mercilon 2874		6
Loxamine	127	Mesalazine		
Lucrin Depot 1-month	88	Mesna165		
Lucrin Depot 3-month	88	Mestinon112	Microgynon 50 ED	
Ludiomil	126	Metabolic Disorder Agents29		
Lyderm		Metformin hydrochloride12		
Lynparza		Methadone hydrochloride	Midazolam-Baxter	14
Lyrica		Extemporaneous247	Midazolam-Claris	14
- M -		Nervous124		5
m-Eslon	125	Methatabs124	Mifegyne	7
Mabthera	209	Methenamine (hexamine)	Mifepristone	7
Macrogol 3350 with potassium	1	hippurate110		
chloride, sodium bicarbona		Methopt243		23
sodium chloride	29	Methotrexate161		
Macrogol 400 and propylene		Methotrexate DBL Onco-Vial161		
glycol	243	Methotrexate Ebewe161	MiniMed 640G	1
Madopar 125		Methotrexate Sandoz161		
Madopar 250		Methyl hydroxybenzoate247	MiniMed Mio MMT-923A	2
Madopar 62.5		Methylcellulose247		2
Madopar HBS		Methylcellulose with glycerin and	MiniMed Mio MMT-941A	
Madopar Rapid	120	sodium saccharin247	MiniMed Mio MMT-943A	2
Magnesium hydroxide		Methylcellulose with glycerin and	MiniMed Mio MMT-945A	2
Magnesium sulphate		sucrose247		
Mantoux	279	Methyldopa54	MiniMed Mio MMT-975A	2
Maprotiline hydrochloride	126	Methyldopa Mylan54		2
Marevan		Methyldopa Mylan S2954	MiniMed Quick-Set MMT-387A	2
Marine Blue Lotion SPF 50+	71	Methylnaltrexone bromide28	MiniMed Quick-Set MMT-396A	2
Martindale Pharma		Methylphenidate ER - Teva151	MiniMed Quick-Set MMT-397A	2
Marvelon 28	74	Methylphenidate hydrochloride 151		
Mask for spacer device		Methylphenidate hydrochloride	MiniMed Quick-Set MMT-399A	
Mast Cell Stabilisers		extended-release152	MiniMed Silhouette MMT-368A	
Maviret		Methylprednisolone80	MiniMed Silhouette MMT-377A	2
Maxidex	240	Methylprednisolone (as sodium	MiniMed Silhouette MMT-378A	

MiniMed Silhouette MMT-381A	22	Mvite	35	Nicotine	15
MiniMed Silhouette MMT-382A	22	Myambutol	100	Nicotinic acid	5
MiniMed Silhouette MMT-383A	22	Mycobutin	101	Nifedipine	5
MiniMed Silhouette MMT-384A	22	MycoNail		Nifuran	
MiniMed Sure-T MMT-864A	20	Mycophenolate mofetil	180	Nilotinib	17
MiniMed Sure-T MMT-866A	20	Mydriacyl		Nilstat	
MiniMed Sure-T MMT-874A	20	Mylan Atenolol		Alimentary	3
MiniMed Sure-T MMT-876A	20	Mylan Clomiphen		Genito-Urinary	
MiniMed Sure-T MMT-884A	20	Myleran		Infection	9
MiniMed Sure-T MMT-886A	20	Myometrial and Vaginal Hormo		Nintedanib	
Minims Cyclopentolate	243	Preparations		Nipent	16
Minims Pilocarpine		Myozyme		Nitrates	
Minims Prednisolone	241	Mysoline S29		Nitroderm TTS	5
Minirin		- N -		Nitrofurantoin	110
Minirin Melt	88	Nadolol	52	Nitrolingual Pump Spray	
Mino-tabs	94	Naglazyme		Nivestim	
Minocycline hydrochloride		Nalcrom		Nivolumab	
Minomycin		Naloxone hydrochloride		Nizoral	
Minor Skin Infections		Naltraccord		Nodia	
Minoxidil		Naltrexone hydrochloride		Noflam 250	
Mirena		Naphazoline hydrochloride		Noflam 500	
Mirtazapine		Naphcon Forte		Non-Steroidal Anti-Inflammatory	
Misoprostol		Naprosyn SR 1000		Drugs	112
Mitomycin C		Naprosyn SR 750		Nonacog gamma, [Recombinant	
Mitozantrone	165	Naproxen		Factor IX]	4 ⁻
Mitozantrone Ebewe		Narcaricin mite		Norethisterone	
Mixtard 30		Nasal Preparations		Genito-Urinary	7
MMR II		Natalizumab		Hormone	
Moclobemide		Natulan		Norflex	
Modafinil		Nausafix		Norfloxacin	
Modavigil		Nausicalm		Noriday 28	
Moduretic		Navelbine		Norimin	
Molaxole		Necon		Normacol Plus	
Moments		Nedocromil		Normison	
Mometasone furoate		Nefopam hydrochloride		Norpress	
Monogen		Neisvac-C		Nortriptyline hydrochloride	
Montelukast		Neo-B12		Norvir	
Montelukast Mylan		Neo-Mercazole		NovaSource Renal	
Moroctocog alfa [Recombinant f		Neo-Mercazole S29		Novatretin	
VIII]		Neocate Gold		NovoMix 30 FlexPen	
Morphine hydrochloride		Neocate Junior Unflavoured		NovoRapid	
Morphine sulphate		Neocate Junior Vanilla		NovoRapid FlexPen	
Motetis		Neocate SYNEO		NovoRapid Penfill	
Mouth and Throat		Neoral		NovoSeven RT	
Movapo		Neostigmine metilsulfate		Noxafil	
Moxifloxacin		Nepafenac		Nozinan	
MSUD Maxamum		Nepro HP (strawberry)		Nozinan (Swiss)	
Mucilaginous laxatives with	200	Nepro HP (vanilla)	255	Nucala	
stimulants	28	Nepro HP RTH	254	Nuelin	
Mucolytics		Nerisone		Nuelin-SR	
Mucosoothe		Neulactil		Nutilis	
Multiple Sclerosis Treatments		Neulastim		Nutrient Modules	
Multivitamin renal		NeuroTabs		Nutrini Energy Multi Fibre	
Multivitamins		Nevirapine			
Mupirocin		Nevirapine Alphapharm		Nutrini Energy RTH Nutrini Low Energy Multi Fibre	
Muscle Relaxants		Nicorandil		Nutrini Peptisorb	
IVIUOUIC I ICIANAI ILO	7	1 VIOOI AI IUII			40

INDEX: Generic Chemicals and Brands

Nutrini Peptisorb Energy267	OncoTICE187	Panadol Mini Caps	12
Nutrini RTH254	Ondansetron132	Pancreatic enzyme	2
Nutrison 800 Complete Multi	Ondansetron ODT-DRLA132	Pantoprazole	
Fibre259	One-Alpha35	Panzop Relief	
Nutrison Concentrated261	Onrex132	Panzytrat	
Nutrison Energy258	Opdivo226	Papaverine hydrochloride	5
Nutrison Energy Multi Fibre259	Ora-Blend247	Para-amino salicylic acid	10
Nutrison Multi Fibre259	Ora-Blend SF247	Paracare	
Nutrison Standard RTH259	Ora-Plus247	Paracare Double Strength	12
Nyefax Retard53	Ora-Sweet247	Paracetamol	12
Nystatin	Ora-Sweet SF247	Paracetamol + Codeine	
Alimentary34	Orabase34	(Relieve)	12
Genito-Urinary76	Oral and Enteral Feeds252	Paracetamol Pharmacare	12
Infection98	Oratane62	Paracetamol with codeine	
NZB Low Gluten Bread Mix262	Ordine124	Paradigm 1.8 Reservoir	2
-0-	Orgran263	Paradigm 3.0 Reservoir	2
O/W Fatty Emulsion Cream66	Ornidazole100	Paradigm Mio MMT-921	
Obinutuzumab207	Orphenadrine citrate119	Paradigm Mio MMT-923	2
Obstetric Preparations78	Ortho-tolidine78	Paradigm Mio MMT-925	2
Ocicure9	Oruvail SR112	Paradigm Mio MMT-941	2
Ocrelizumab141	Osmolite RTH259	Paradigm Mio MMT-943	2
Ocrevus141	Other Endocrine Agents88	Paradigm Mio MMT-945	2
Octocog alfa [Recombinant factor	Other Oestrogen Preparations83	Paradigm Mio MMT-965	2
VIII] (Advate)42	Other Progestogen	Paradigm Mio MMT-975	2
Octocog alfa [Recombinant factor	Preparations83	Paradigm Quick-Set MMT-386	2
VIII] (Kogenate FS)42	Other Skin Preparations71	Paradigm Quick-Set MMT-387	2
Octreotide178	Ovestin	Paradigm Quick-Set MMT-396	2
Octreotide (Sun) 178	Genito-Urinary76	Paradigm Quick-Set MMT-397	2
Octreotide GH178	Hormone83	Paradigm Quick-Set MMT-398	2
Octreotide LAR (somatostatin	Ox-Pam136	Paradigm Quick-Set MMT-399	2
analogue) 178	Oxaliplatin159	Paradigm Silhouette MMT-368	2
Octreotide MaxRx178	Oxaliplatin Accord159	Paradigm Silhouette MMT-377	2
Oestradiol82	Oxaliplatin Actavis 100 159	Paradigm Silhouette MMT-378	2
Oestradiol valerate82	Oxaliplatin Ebewe159	Paradigm Silhouette MMT-381	2
Oestradiol with norethisterone83	Oxazepam 136	Paradigm Silhouette MMT-382	
Oestriol	Oxis Turbuhaler233	Paradigm Silhouette MMT-383	
Genito-Urinary76	Oxpentifylline58	Paradigm Silhouette MMT-384	2
Hormone83	Oxybutynin77	Paradigm Sure-T MMT-864	
Oestrogens82	Oxycodone hydrochloride125	Paradigm Sure-T MMT-866	2
Ofev235	Oxycodone Sandoz125	Paradigm Sure-T MMT-874	2
Oil in water emulsion66	OxyNorm125	Paradigm Sure-T MMT-876	2
Olanzapine 134–135	Oxytocin76	Paradigm Sure-T MMT-884	2
Olaparib	Oxytocin BNM76	Paradigm Sure-T MMT-886	2
Olbetam56	Oxytocin with ergometrine	Paraffin6	6-6
Olbetam S2956	maleate77	Paraffin liquid with wool fat	24
Olopatadine243	Ozurdex240	Paraldehyde	12
Olopatadine Teva243	- P -	Parasidose	6
Olsalazine7	Pacifen119	Parasiticidal Preparations	6
Omalizumab207	Paclitaxel 166	Parnate	12
Omeprazole9	Paclitaxel Actavis166	Parnate S29	12
Omeprazole actavis 109	Paclitaxel Ebewe166	Paromomycin	
Omeprazole actavis 209	Paediatric Seravit35	Paroxetine	
Omeprazole actavis 409	Palbociclib 173	Paser	
Omnitrope84	Paliperidone135	Paxam	
Onbrez Breezhaler233	Pamidronate disodium114	Pazopanib	17
Oncaspar LYO166	Pamisol114	Peak flow meter	23
· · · · · · · · · · · · · · · · · · ·			

Pedialyte - Bubblegum	47	Pinetarsol		Premarin	8
Pediasure	254	Pioglitazone	12	Prevenar 13	
Pediasure RTH	254	Pirfenidone	236	Prezista	10
Pegaspargase	166	Pizotifen	131	Priadel	13
Pegasys	108	PKU Anamix Infant	264	Primacin	
Pegfilgrastim	46	PKU Anamix Junior	264	Primaquine	9
Pegylated interferon alfa-2a	108	PKU Anamix Junior Chocolate	264	Primidone	130
Pembrolizumab		PKU Anamix Junior LQ	264	Primolut N	
Pemetrexed	161	PKU Anamix Junior Vanilla	264	Priorix	27
Penicillamine	113	PKU Lophlex LQ 10		Probenecid	119
Penicillin G	93	PKU Lophlex LQ 20	264	Probenecid-AFT	119
PenMix 30	11	PKU Lophlex Powder	264	Procaine penicillin	9
PenMix 40		PKU Lophlex Sensation 20	264	Procarbazine hydrochloride	16
PenMix 50	11	Plaquenil	113	Prochlorperazine	13
Pentasa	7	Plendil ER	53	Proctofoam	
Pentostatin [Deoxycoformycin]	166	Pneumococcal (PCV10) conjugate		Proctosedyl	
Pentoxifylline [Oxpentifylline]	58	vaccine	. 276	Procyclidine hydrochloride	12
Peptamen Junior		Pneumococcal (PCV13) conjugate		Procytox	
Peptisoothe		vaccine	. 277	Progesterone	
Peptisorb		Pneumococcal (PPV23)		Proglicem	10
Perhexiline maleate		polysaccharide vaccine	. 278	Proglycem	
Pericyazine		Pneumovax 23		Progynova	
Perindopril		Podophyllotoxin	71	Prolia	
Perjeta		Polaramine		Promethazine hydrochloride	23
Permethrin		Poliomyelitis vaccine		Propafenone hydrochloride	5
Perrigo		Poloxamer		Propamidine isethionate	
Pertuzumab		Poly-Gel		Propranolol	
Peteha		Poly-Tears		Propylene glycol	
Pethidine hydrochloride		Poly-Visc		Propylthiouracil	
Pevaryl		Polycal	. 249	Protaphane	
Pexsig		Ponstan		Protaphane Penfill	
Pfizer Exemestane		Posaconazole		Protifar	
Pharmacy Services		Postinor-1		Protionamide	
Pheburane		Potassium chloride4		Provera	
Phenasen		Potassium Chloride Aguettant		Provera HD	
Phenobarbitone		Potassium citrate		PSM Citalopram	
Phenobarbitone sodium		Potassium iodate		Psoriasis and Eczema	
Extemporaneous	247	Povidone iodine		Preparations	69
Nervous		Pradaxa		PTU	
Phenothrin		Pramipexole hydrochloride		Pulmicort Turbuhaler	
Phenoxybenzamine		Prasugrel		Pulmozyme	
hydrochloride	48	Pravastatin		Puri-nethol	
Phenoxymethylpenicillin (Penicillin		Pravastatin Mylan		Puria	
V)		Praziquantel		Pyrazinamide	
Phenytoin sodium12		Prazosin		Pyridostigmine bromide	
Phillips Milk of Magnesia		Pred Forte		Pyridoxine hydrochloride	
Phlexy 10		Prednisolone		Pyrimethamine	91
Phosphate Phebra		Prednisolone acetate		Pytazen SR	
Phosphorus		Prednisolone sodium		- Q -	
Phytomenadione		Prednisolone sodium		Q 300	Q
Pilocarpine hydrochloride	242	phosphate	241	Quetapel	
Pimafucort		Prednisolone-AFT	241	Quetiapine	12
Pimecrolimus		Prednisone		Quick-Set MMT-392	ان ان
Pindolol		Pregabalin		Quick-Set MMT-393	ا2
Pine tar with trolamine laurilsulfate		Pregabalin Pfizer		Quinapril	
and fluorescein		Pregnancy Tests - hCG Urine		Quinapril with	40
and iidorooodiii	/ 🗸	Trogramor roots Tiou office	I I	Ganapin mui	

INDEX: Generic Chemicals and Brands

hydrochlorothiazide	49	Rivastigmine	153	Sertraline	12
Quinine sulphate		Rivotril		Setrona	
Qvar	232	Riximyo	210	Setrona AU	12
- R -		RIXUBIS	41	Sevredol	12
RA-Morph	124	Rizamelt	131	Sex Hormones Non	
Raloxifene hydrochloride		Rizatriptan		Contraceptive	8
Raltegravir potassium		Rolin	179	Shield XL	
Ramipex		Ropin		shingles vaccine	
Ranbaxy-Cefaclor		Ropinirole hydrochloride		SII-Onco-BCG	18
Ranitidine		Rotarix		Sildenafil	
Rapamune		Rotavirus oral vaccine		Silhouette MMT-373	
Reandron 1000		Roxane		Siltuximab	
Recombinant factor IX		Alimentary	6	Simvastatin	
Recombinant factor VIIa		Cardiovascular		Simvastatin Mylan	
Recombinant factor VIII		Roxithromycin		Sinemet	
Rectogesic		Rubifen		Sinemet CR	
Redipred		Rubifen SR		Sirolimus	
Relieve		Rugby Capsaicin Topical Cr		Siterone	
Relistor		Musculoskeletal		Slow-Lopresor	
Remicade				Smith BioMed Rapid Pregnancy	
Renilon 7.5		Nervous Rulide D		Test	7
Resonium-A				Sodibic	
		Rurioctocog alfa pegol [Reco			
Resource Beneprotein		factor VIII]		Sodium acid phosphate	
Resource Diabetic		Ruxolitinib		Sodium alginate	
Respigen		Rythmodan		Sodium benzoate	3
Respiratory Devices		Rytmonorm	50	Sodium bicarbonate	40.4
Respiratory Stimulants		•	400	Blood	
Retinol palmitate		Sabril		Extemporaneous	
ReTrieve		Sacubitril with valsartan		Sodium calcium edetate	24
Retrovir		SalAir		Sodium chloride	
Revlimid		Salazopyrin		Blood	
Revolade		Salazopyrin EN		Respiratory	23
Rexacrom		Salbutamol		Sodium citrate with sodium lauryl	
Ribomustin		Salbutamol with ipratropium		sulphoacetate	
Ricit		bromide		Sodium citro-tartrate	7
Rifabutin		Salicylic acid		Sodium cromoglicate	
Rifadin		Salmeterol		Alimentary	
Rifampicin		Sandomigran		Respiratory	
Rifaximin		Sandostatin LAR		Sensory	
Rifinah		Sapropterin dihydrochloride.		Sodium fluoride	3
Rilutek		Scalp Preparations		Sodium Fusidate [fusidic acid]	
Riluzole		Scopoderm TTS	132	Dermatological	
Riodine	67	Sebizole		Infection	
Risedronate Sandoz	115	Secukinumab		Sensory	23
Risedronate sodium	115	Sedatives and Hypnotics	149	Sodium hyaluronate [Hyaluronic	
Risperdal Consta	135	Seebri Breezhaler	235	acid]	24
Risperidone	134–135	Selegiline hydrochloride	120	Sodium phenylbutyrate	3
Risperidone (Teva)	134	Senna	29	Sodium polystyrene sulphonate	4
Risperon	134	Senokot		Sodium tetradecyl sulphate	4
Ritalin		Sensipar	79	Sodium valproate	13
Ritalin LA	152	SensoCard	13	Sofradex	
Ritalin SR	151	Serenace	133	Soframycin	23
Ritonavir	107	Seretide	234	Solifenacin Mylan	7
Rituximab (Mabthera)		Seretide Accuhaler	234	Solifenacin succinate	7
Rituximab (Riximyo)		Serevent		Solu-Cortef	8
Rivaroxaban		Serevent Accuhaler		Solu-Medrol	

Solu-Medrol-Act-O-Vial	80	Tacrolimus	229	Timoptol XE	241
Somatropin (Omnitrope)		Tacrolimus Sandoz		Tiotropium bromide	
Sotalol		Taliglucerase alfa		Tiotropium bromide with	
Spacer device		Tambocor		olodaterol	235
Span-K		Tamoxifen citrate		Tivicay	
Spiolto Respimat		Tamoxifen Sandoz		TMP	
Spiractin		Tamsulosin hydrochloride		TOBI	
Spiriva		Tamsulosin-Rex		Tobramycin	
Spiriva Respimat		Tandem Cartridge		Infection	96
Spironolactone		Tandem t:slim X2 with Basal-IC		Sensory	
Sporanox		Tap water		Tobramycin BNM	
Sprycel		Tarceva		Tobramycin Mylan	
Staphlex		Tasigna		Tobrex	
Stemetil		Tasmar		Tocilizumab	
SteroClear		Tecfidera		Tofranil	
Stesolid		Tegretol		Tolcapone	
Stimulants/ADHD Treatments		Tegretol CR		Topamax	
Stiripentol		Telfast		Topical Products for Joint and	
Stocrin		Teligent		Muscular Pain	119
Stomahesive		Temaccord	167	Topiramate	
Strides Shasun		Temazepam		Topiramate Actavis	
Stromectol		Temozolomide		Total parenteral nutrition (TPN)	
Sucralfate		Tenofovir disoproxil		TPN	
Sulfadiazine Silver		Tenofovir Disoproxil Teva		Tramadol hydrochloride	
Sulfadiazine sodium		Tenoxicam		Tramal SR 100	
Sulfasalazine		Tensipine MR10		Tramal SR 150	
Sulindac		Tepadina		Tramal SR 200	
Sulindac Mylan		Terazosin	139	Trandate	
Sulphur		Terbinafine		Tranexamic acid	
Sulprix		Terbutaline sulphate		Tranylcypromine sulphate Trastuzumab	
Sumatriptan		Teriflunomide			
Sunitinib		Teriparatide		Trastuzumab emtansine	
Sunscreens		Testosterone		Travatan	
Sunscreens, proprietary		Testosterone cipionate		Travoprost	242
Sure-T MMT-863		Testosterone esters		Travopt	242
Sure-T MMT-873		Testosterone undecanoate		Treatments for Dementia	150
Sustagen Diabetic	252	Tetrabenazine		Treatments for Substance	45
Sustagen Hospital Formula	050	Tetrabromophenol		Dependence	
Active		Tetracosactrin		Trental 400	58
Sustanon Ampoules		Tetracycline		Tretinoin	0.0
Sutent		Thalidomide		Dermatological	
Sylvant		Thalomid		Oncology	168
Symbicort Turbuhaler 100/6		Theophylline	237	Trexate	161
Symbicort Turbuhaler 200/6		Thiamine hydrochloride		Triamcinolone acetonide	
Symbicort Turbuhaler 400/12		THIO-TEPA		Alimentary	34
Symmetrel		Thioguanine		Dermatological	
Sympathomimetics		Thiotepa		Hormone	
Synacthen		Thymol glycerin		Triamcinolone acetonide with	
Synacthen Depot	81	Thyroid and Antithyroid Agents		gramicidin, neomycin and nyst	
Synacthene Retard		Ticagrelor		Dermatological	
Synflorix		Tilade		Sensory	
Synthroid		Tilcotil		Triaver	
Syntometrine		Tillomed	158	Triazolam	
Syrup (pharmaceutical grade)		Timolol		Trimethoprim	96
Systane Unit Dose	243	Cardiovascular		Trimethoprim with	
- T -		Sensory	241	sulphamethoxazole	

INDEX: Generic Chemicals and Brands

[Co-trimoxazole]	97	Ventavis	61	Xyntha	4
Trisequens	83	Ventolin	234	- Z -	
Trisul	97	Vepesid	164	Zantac	
Trophic Hormones	84	Verapamil hydrochloride	54	Zapril	4
Tropicamide		Vergo 16		Zarontin	12
Trusopt		Vermox		Zaroxolyn	
TruSteel		Vesanoid	168	Zavedos	
Tuberculin PPD [Mantoux] test	279	Vexazone	12	Zeffix	10
Tubersol		Vfend	98	Zetlam	10
Two Cal HN	261	Viaderm KC	66	Ziagen	10
Two Cal HN RTH	261	Vidaza	159	Zidovudine [AZT]	
Tykerb		Vigabatrin		Zidovudine AZT with	
Tysabri		Vildagliptin		lamivudine	10
. U -		Vildagliptin with metformin		Zimybe	5
Ultibro Breezhaler	235	hydrochloride	12	Zinc and castor oil	
Ultraproct	8	Vimpat		Zinc sulphate	
Umeclidinium		Vinblastine sulphate		Zincaps	3
Umeclidinium with vilanterol		Vincristine sulphate		Zinnat	9
Univent		Vinorelbine		Ziprasidone	
Ural		Vinorelbine Ebewe		Zista	
Urea		Viramune Suspension		Zithromax	
Urex Forte		ViruPOS		Zoladex	
Urinary Agents		Vit.D3		Zoledronic acid	
Urinary Tract Infections		Vita-B12		Hormone	7
Urinorm		VitA-POS		Musculoskeletal	11
Uromitexan		Vitabdeck		Zoledronic acid Mylan	
Ursodeoxycholic acid		Vital		Zopiclone	
Ursosan		Vitamin B complex		Zopiclone Actavis	15
Utrogestan		Vitamin B6 25		Zostavax	
- V -		Vitamins		Zostrix	
Vaccinations	270	Vivonex TEN		Zostrix HP	
Vaclovir		Volibris		Zuclopenthixol decanoate	
Valaciclovir		Voltaren		Zuclopenthixol hydrochloride	
Valganciclovir		Voltaren D		Zusdone	
Valganciclovir Mylan		Voltaren Ophtha		Zyban	
Vancomycin		Volumatic		Zypine	
Vannair		Voriconazole		Zypine ODT	
Varenicline Pfizer		Vosol		Zyprexa Relprevv	
Varenicline tartrate		Votrient		Zytiga	
Varicella vaccine [Chickenpox		Vttack		2y tiga	
vaccine]	270	- W -			
Varicella zoster virus (Oka strair		Warfarin sodium	45		
attenuated vaccine [shingles	1) 1140	Wart Preparations			
vaccine]	270	Wasp venom allergy treatmer			
Various		Water	201		
Varivax		Blood	47		
Vasodilators		Extemporaneous			
Vasopressin Agonists		Wool fat with mineral oil	67		
Vasorex		• X -	0/		
Vedafil		Xarelto	15		
Veletri		Xifaxan			
Venclexta		XMET Maxamum			
Venetoclax		Xolair			
Venlafaxine		XP Maxamum			
		Xylocaine			
Venomil					
VENOX	231	Xylocaine 2% Jelly	121		